

# Computer Algebra Independent Integration Tests

Summer 2023 edition with Rubi V 4.17.3

7-Inverse-hyperbolic-functions/7.2-Inverse-hyperbolic-cosine/190-  
7.2.4-f-x-<sup>m</sup>-d+e-x<sup>2</sup>-<sup>p</sup>-a+b-arccosh-c-x-<sup>n</sup>

Nasser M. Abbasi

December 9, 2023

Compiled on December 9, 2023 at 7:19am

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>detailed summary tables of results</b>	<b>20</b>
<b>3</b>	<b>Listing of integrals</b>	<b>189</b>
<b>4</b>	<b>Appendix</b>	<b>4102</b>

# CHAPTER 1

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	3
1.2	Results . . . . .	4
1.3	Time and leaf size Performance . . . . .	7
1.4	Performance based on number of rules Rubi used . . . . .	9
1.5	Performance based on number of steps Rubi used . . . . .	10
1.6	Solved integrals histogram based on leaf size of result . . . . .	11
1.7	Solved integrals histogram based on CPU time used . . . . .	12
1.8	Leaf size vs. CPU time used . . . . .	13
1.9	list of integrals with no known antiderivative . . . . .	14
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	14
1.11	list of integrals solved by CAS but failed verification . . . . .	14
1.12	Timing . . . . .	15
1.13	Verification . . . . .	15
1.14	Important notes about some of the results . . . . .	16
1.15	Design of the test system . . . . .	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 569 ]. This is test number [ 190 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 13.3.1 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	99.12 ( 564 )	0.88 ( 5 )
Mathematica	98.24 ( 559 )	1.76 ( 10 )
Maple	83.66 ( 476 )	16.34 ( 93 )
Maxima	42.00 ( 239 )	58.00 ( 330 )
Fricas	41.65 ( 237 )	58.35 ( 332 )
Mupad	25.31 ( 144 )	74.69 ( 425 )
Giac	18.63 ( 106 )	81.37 ( 463 )
Sympy	16.34 ( 93 )	83.66 ( 476 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Mathematica	68.717	1.933	3.866	25.483
Rubi	58.875	0.000	15.290	25.835
Maple	37.258	17.223	4.218	41.301
Fricas	18.278	2.460	0.000	79.262
Maxima	15.114	1.230	1.933	81.722
Giac	0.176	0.176	0.703	98.946
Mupad	0.000	0.351	0.000	99.649
Sympy	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	10	30.00	70.00	0.00
Rubi	5	100.00	0.00	0.00
Maple	93	100.00	0.00	0.00
Fricas	332	80.12	0.00	19.88
Maxima	330	90.30	0.00	9.70
Giac	463	52.48	0.65	46.87
Mupad	425	0.00	100.00	0.00
Sympy	476	69.33	30.67	0.00

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.



System	Mean time (sec)
Fricas	0.27
Maxima	0.51
Rubi	0.97
Maple	1.74
Giac	2.96
Mupad	3.16
Mathematica	3.33
Sympy	34.99

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	26.68	1.01	28.00	1.00
Sympy	28.12	1.09	27.00	1.00
Giac	29.36	1.02	26.50	1.00
Fricas	175.00	1.92	103.00	1.51
Rubi	198.42	0.93	154.50	1.00
Maxima	200.69	4.56	134.00	1.00
Mathematica	256.98	1.07	144.00	1.01
Maple	429.84	1.78	192.00	1.05

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following gives the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{148, 149, 150, 233, 234, 235, 236, 237, 238, 239, 261, 265, 266, 272, 273, 274, 275, 280, 281, 282, 283, 288, 289, 290, 291, 297, 298, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 318, 319, 324, 325, 326, 327, 331, 332, 333, 334, 338, 339, 340, 341, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 373, 378, 382, 383, 387, 391, 395, 396, 400, 406, 407, 412, 413, 417, 418, 422, 423, 427, 428, 432, 433, 438, 439, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 514, 515, 522, 523, 524, 530, 531, 532, 533, 537, 538, 539, 540, 541, 542, 546, 547, 548, 549, 550, 551, 555, 556, 559, 560, 564, 565, 568, 569}

## 1.10 List of integrals solved by CAS but has no known antiderivative

**Rubi** {}

**Mathematica** {}

**Maple** {}

**Maxima** {}

**Fricas** {}

**Sympy** {}

**Giac** {}

**Mupad** {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

**Rubi** {6, 8, 15, 16, 17, 18, 24, 26, 27, 175, 177, 183, 185, 191, 192, 201, 203, 476, 478, 487}

**Mathematica** {15, 22, 24, 26, 29, 30, 37, 38, 39, 41, 42, 44, 46, 48, 50, 51, 52, 53, 54, 56, 57, 58, 60, 61, 68, 69, 70, 71, 72, 73, 74, 75, 84, 85, 86, 87, 88, 89, 90, 91, 92, 100, 101, 102, 112, 115, 117, 120, 122, 125, 130, 132, 142, 169, 171, 173, 174, 175, 176, 177, 179, 181, 182, 183, 184, 185, 187, 189, 190, 191, 192, 193, 202, 203, 204, 205, 206, 207, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 232, 245, 246, 247, 251, 252, 260, 267, 268, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 315, 316, 369, 370, 372, 374, 375, 377, 392, 397, 414, 415, 462, 464, 471, 473, 475, 480, 482, 484, 486, 489, 490, 502, 506, 509, 510, 511, 512, 513, 543, 544, 557, 558, 566}

**Maple** {490, 492, 494, 495, 496, 497, 498, 500, 501, 502, 503, 504, 505, 506, 509, 510, 511, 512, 513}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals



were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### 1.14.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

### 1.14.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

### 1.14.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

#### 1.14.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

# 1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1. Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
June 27, 2013  
Design v0.01

# CHAPTER 2

## DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	21
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	28
2.3	Detailed conclusion table specific for Rubi results . . . . .	171

## 2.1 List of integrals sorted by grade for each CAS

2.1.1	Rubi . . . . .	21
2.1.2	Mma . . . . .	22
2.1.3	Maple . . . . .	23
2.1.4	Fricas . . . . .	23
2.1.5	Maxima . . . . .	24
2.1.6	Giac . . . . .	25
2.1.7	Mupad . . . . .	26
2.1.8	Sympy . . . . .	26

### 2.1.1 Rubi

**A grade** { 1, 2, 3, 4, 5, 7, 9, 10, 11, 12, 13, 14, 16, 18, 19, 20, 21, 22, 23, 25, 27, 40, 47, 49, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 170, 171, 172, 173, 174, 176, 178, 179, 180, 181, 182, 184, 188, 189, 190, 192, 194, 195, 196, 197, 198, 199, 200, 202, 210, 212, 220, 225, 226, 227, 228, 229, 230, 232, 240, 241, 242, 246, 247, 248, 249, 253, 254, 255, 256, 257, 258, 260, 267, 268, 269, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 292, 293, 294, 295, 296, 299, 300, 301, 302, 315, 316, 317, 320, 328, 330, 335, 337, 342, 343, 344, 347, 368, 369, 370, 371, 372, 374, 375, 376, 377, 381, 384, 385, 386, 390, 394, 397, 398, 399, 401, 402, 403, 404, 405, 408, 409, 411, 414, 415, 416, 419, 420, 421, 424, 425, 426, 429, 430, 431, 434, 435, 436, 437, 440, 441, 442, 443, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 516, 517, 518, 519, 520, 521, 525, 526, 527, 528, 529, 534, 535, 543, 544, 545, 552, 553, 554, 557, 561, 562, 566, 567 }  
}

**B grade** { }

**C grade** { 6, 8, 15, 17, 24, 26, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 48, 50, 51, 52, 53, 54, 55, 56, 57, 167, 168, 169, 175, 177, 183, 185, 191, 201, 203, 204, 205, 206, 207, 208, 209, 211, 213, 214, 215, 216, 217, 218, 219, 221, 223, 224, 231, 243, 244, 245, 250, 251, 252, 259, 262, 263, 264, 321, 322, 323, 329, 336, 345, 346, 379, 380, 389, 392, 393, 410, 536, 558, 563 }  
}

**F normal fail** { 186, 187, 193, 222, 388 }

**F(-1) timeout fail { }**

**F(-2) exception fail { }**

## 2.1.2 Mma

**A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 177, 178, 179, 180, 181, 182, 183, 185, 186, 187, 188, 189, 190, 191, 193, 194, 195, 196, 197, 198, 199, 200, 201, 203, 204, 205, 206, 207, 208, 209, 210, 211, 213, 214, 215, 216, 217, 218, 219, 220, 221, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 253, 254, 255, 256, 257, 258, 259, 262, 263, 264, 267, 268, 269, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 292, 293, 294, 295, 296, 299, 300, 301, 302, 317, 320, 321, 322, 323, 328, 329, 330, 335, 336, 337, 342, 343, 344, 345, 346, 347, 368, 369, 370, 372, 374, 375, 377, 379, 380, 381, 384, 385, 386, 388, 389, 390, 392, 393, 394, 397, 398, 399, 401, 402, 403, 404, 405, 408, 409, 410, 411, 414, 415, 416, 419, 420, 421, 424, 425, 426, 429, 430, 431, 434, 435, 436, 437, 440, 441, 442, 443, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 490, 492, 493, 495, 497, 499, 507, 508, 519, 520, 521, 525, 526, 527, 528, 529, 534, 535, 536, 543, 544, 545, 552, 553, 554, 557, 558, 561, 562, 563, 566 }**

**B grade { 33, 42, 176, 184, 192, 202, 212, 222, 260, 315, 316 }**

**C grade { 251, 252, 489, 491, 494, 496, 498, 500, 501, 502, 503, 504, 505, 506, 509, 510, 511, 512, 513, 516, 517, 518 }**

**F normal fail { 371, 376, 567 }**

**F(-1) timeout fail { 333, 334, 341, 555, 556, 559, 560 }**

**F(-2) exception fail { }**

### 2.1.3 Maple

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 56, 57, 62, 68, 69, 70, 74, 75, 84, 85, 86, 90, 91, 100, 101, 102, 109, 110, 112, 113, 114, 115, 116, 118, 121, 122, 123, 124, 125, 126, 130, 131, 132, 133, 134, 139, 142, 164, 165, 166, 167, 168, 169, 183, 185, 191, 193, 204, 205, 206, 208, 214, 216, 218, 227, 228, 229, 231, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 254, 255, 256, 257, 259, 262, 263, 264, 267, 268, 269, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 292, 293, 294, 296, 299, 300, 301, 302, 315, 316, 317, 320, 322, 328, 329, 335, 336, 342, 343, 344, 345, 346, 347, 368, 381, 386, 390, 394, 399, 405, 411, 416, 437, 443, 461, 462, 463, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 525, 526, 527, 528, 534, 535, 536, 544, 545 }

**B grade** { 58, 59, 60, 61, 63, 64, 65, 66, 67, 71, 72, 73, 76, 77, 78, 79, 80, 81, 82, 83, 87, 88, 89, 92, 93, 94, 95, 96, 97, 98, 99, 103, 104, 105, 106, 107, 108, 111, 117, 119, 120, 127, 128, 129, 135, 136, 137, 138, 140, 141, 170, 171, 172, 173, 175, 177, 178, 179, 180, 181, 186, 187, 188, 189, 194, 195, 196, 197, 198, 199, 201, 203, 207, 209, 211, 213, 215, 217, 219, 221, 223, 224, 225, 226, 250, 251, 252, 253, 295, 321, 323, 330, 337, 464, 499, 507, 508, 543 }

**C grade** { 32, 55, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 500, 501, 502, 503, 504, 505, 506, 509, 510, 511, 512, 513 }

**F normal fail** { 143, 144, 145, 146, 147, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 174, 176, 182, 184, 190, 192, 200, 202, 210, 212, 220, 222, 230, 232, 258, 260, 369, 370, 371, 372, 374, 375, 376, 377, 379, 380, 384, 385, 388, 389, 392, 393, 397, 398, 401, 402, 403, 404, 408, 409, 410, 414, 415, 419, 420, 421, 424, 425, 426, 429, 430, 431, 434, 435, 436, 440, 441, 442, 516, 517, 518, 519, 520, 521, 529, 552, 553, 554, 557, 558, 561, 562, 563, 566, 567 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### 2.1.4 Fricas

**A grade** { 1, 2, 3, 4, 5, 7, 9, 10, 11, 12, 13, 14, 16, 18, 19, 20, 21, 22, 23, 25, 27, 40, 47, 49, 63, 64, 65, 66, 67, 76, 77, 78, 79, 80, 81, 82, 83, 93, 94, 95, 96, 97, 98, 99, 104, 106, 108, 111, 113, 114, 116, 118, 124, 126, 128, 136, 138, 164, 165, 166, 170, 172, 178, 180, 186, 188, 194, 196, 198, 226, 228, 240, 241, 242, 254, 256, 411, 416, 461, 462, 463, 464, 465, 467, 469, 470, 471, 472, 473, 474, 476, 478, 479, 480, 481, 482, 483, 485, 487, 488, 516, 525, 526, 527 }

**B grade** { 62, 141, 296, 302, 347, 368, 437, 443, 499, 507, 508, 517, 518, 528 }

**C grade** { }

**F normal fail** { 6, 8, 15, 17, 24, 26, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 48, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 68, 69, 70, 71, 72, 73, 74, 75, 84, 85, 86, 87,



88, 89, 90, 91, 92, 100, 101, 102, 103, 105, 107, 109, 110, 112, 115, 117, 119, 120, 121, 122, 123, 125, 127, 129, 130, 131, 132, 133, 134, 135, 137, 139, 140, 142, 143, 144, 145, 146, 147, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 167, 168, 169, 171, 173, 174, 175, 176, 177, 179, 181, 182, 183, 184, 185, 187, 189, 190, 191, 192, 193, 195, 197, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 227, 229, 230, 231, 232, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 255, 257, 258, 259, 260, 262, 263, 264, 267, 268, 269, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 292, 293, 294, 295, 299, 300, 301, 315, 316, 317, 320, 321, 322, 323, 328, 329, 330, 335, 336, 337, 342, 343, 344, 345, 346, 419, 420, 421, 424, 425, 426, 429, 430, 431, 434, 435, 436, 440, 441, 442, 466, 468, 475, 477, 484, 486, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 500, 501, 502, 503, 504, 505, 506, 509, 510, 511, 512, 513, 519, 520, 521, 529, 534, 535, 536, 543, 544, 545 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 412, 413, 414, 415, 417, 418, 552, 553, 554, 555, 556, 557, 558, 559, 560, 561, 562, 563, 564, 565, 566, 567, 568, 569 }

## 2.1.5 Maxima

**A grade** { 1, 2, 3, 4, 5, 7, 9, 10, 12, 14, 16, 18, 23, 25, 27, 63, 64, 65, 66, 67, 77, 78, 79, 80, 81, 82, 83, 94, 95, 96, 97, 98, 99, 104, 106, 108, 113, 116, 119, 121, 126, 127, 129, 133, 134, 164, 165, 166, 170, 172, 178, 180, 186, 188, 194, 196, 198, 240, 241, 242, 461, 462, 463, 464, 465, 467, 469, 470, 471, 472, 473, 474, 476, 478, 479, 480, 481, 482, 483, 485, 487, 488, 525, 526, 527, 528 }

**B grade** { 11, 13, 19, 20, 21, 22, 40 }

**C grade** { 62, 76, 93, 111, 136, 138, 141, 226, 228, 254, 256 }

**F normal fail** { 6, 8, 15, 17, 24, 26, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 68, 69, 70, 71, 72, 73, 74, 75, 84, 85, 86, 87, 88, 89, 90, 91, 92, 100, 101, 102, 105, 107, 109, 110, 112, 114, 115, 117, 118, 120, 122, 123, 124, 125, 128, 130, 131, 132, 139, 140, 142, 143, 144, 145, 146, 147, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 167, 168, 169, 171, 173, 174, 175, 176, 177, 179, 181, 182, 183, 184, 185, 187, 189, 190, 191, 192, 193, 195, 197, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 229, 230, 231, 232, 243, 244, 245, 249, 250, 251, 252, 257, 258, 259, 260, 262, 263, 264, 267, 268, 269, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 292, 293, 294, 295, 296, 299, 300, 301, 302, 315, 316, 317, 320, 321, 322, 323, 328, 329, 330, 335, 336, 337, 342, 343, 344, 345, 346, 347, 368, 369, 370, 371, 372, 374, 375, 376, 377, 379, 380, 381, 384, 385, 386, 388, 389, 390, 392, 393, 394, 397, 398, 399, 401, 402, 403, 404, 405, 408, 409, 410, 411, 414, 415, 416, 419, 420, 421, 424, 425, 426, 429, 430, 431, 434, 435, 436, 437, 440, 441, 442, 443, 466, 468, 475, 477, 484, 486, 490, 492, 494, 496, 498, 500,

501, 506, 507, 508, 509, 510, 517, 518, 519, 520, 521, 534, 535, 536, 543, 544, 545, 552, 553, 554, 557, 558, 561, 562, 563, 566, 567 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { 103, 135, 137, 225, 227, 246, 247, 248, 253, 255, 489, 491, 493, 495, 497, 499, 502, 503, 504, 505, 511, 512, 513, 514, 515, 516, 529, 530, 531, 532, 555, 559 }

## 2.1.6 Giac

**A grade** { 134 }

**B grade** { 528 }

**C grade** { 138, 141, 228, 256 }

**F normal fail** { 30, 31, 32, 33, 34, 35, 36, 39, 40, 41, 42, 43, 44, 45, 46, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 71, 72, 87, 88, 103, 105, 107, 108, 109, 110, 111, 112, 113, 117, 118, 119, 120, 121, 122, 123, 125, 127, 128, 129, 130, 131, 132, 133, 135, 137, 139, 140, 142, 143, 144, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 167, 168, 169, 171, 179, 187, 195, 197, 198, 199, 200, 201, 202, 203, 207, 208, 209, 210, 211, 212, 213, 215, 217, 218, 219, 220, 221, 222, 223, 225, 227, 229, 230, 231, 232, 243, 244, 245, 249, 250, 253, 255, 257, 258, 259, 260, 262, 263, 264, 267, 269, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 292, 294, 295, 296, 300, 301, 302, 315, 316, 317, 321, 322, 323, 328, 329, 330, 335, 336, 337, 343, 345, 346, 347, 368, 370, 372, 375, 377, 381, 386, 390, 392, 393, 394, 397, 398, 399, 401, 402, 403, 404, 405, 408, 409, 410, 411, 414, 415, 416, 435, 436, 437, 441, 442, 443, 466, 467, 468, 469, 475, 476, 477, 478, 484, 485, 486, 487, 489, 491, 492, 493, 494, 495, 496, 497, 499, 500, 501, 502, 503, 504, 505, 508, 509, 510, 511, 512, 513, 516, 517, 518, 519, 520, 521, 529, 534, 535, 536, 543, 544, 545, 552, 553, 554, 558, 561, 562, 563, 566, 567 }

**F(-1) timeout fail** { 454, 457, 458 }

**F(-2) exception fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 37, 38, 47, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 104, 106, 114, 115, 116, 124, 126, 136, 145, 146, 147, 151, 152, 153, 164, 165, 166, 170, 172, 173, 174, 175, 176, 177, 178, 180, 181, 182, 183, 184, 185, 186, 188, 189, 190, 191, 192, 193, 194, 196, 204, 205, 206, 214, 216, 224, 226, 233, 234, 235, 240, 241, 242, 246, 247, 248, 251, 252, 254, 268, 272, 274, 280, 282, 288, 290, 293, 299, 303, 306, 308, 310, 311, 320, 324, 326, 331, 333, 338, 340, 342, 344, 348, 350, 352, 354, 357, 359, 361, 363, 364, 369, 371, 373, 374, 376, 378, 379, 380, 384, 385, 388, 389, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 440, 452, 453, 461, 462, 463, 464, 465, 470, 471, 472, 473, 474, 479, 480, 481, 482, 483, 488, 490, 498, 506, 507, 525, 526, 527, 557 }

### 2.1.7 Mupad

**A grade** { }

**B grade** { 347, 368 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 262, 263, 264, 267, 268, 269, 270, 271, 276, 277, 278, 279, 284, 285, 286, 287, 292, 293, 294, 295, 296, 299, 300, 301, 302, 315, 316, 317, 320, 321, 322, 323, 328, 329, 330, 335, 336, 337, 342, 343, 344, 345, 346, 369, 370, 371, 372, 374, 375, 376, 377, 379, 380, 381, 384, 385, 386, 388, 389, 390, 392, 393, 394, 397, 398, 399, 401, 402, 403, 404, 405, 408, 409, 410, 411, 414, 415, 416, 419, 420, 421, 424, 425, 426, 429, 430, 431, 434, 435, 436, 437, 440, 441, 442, 443, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 516, 517, 518, 519, 520, 521, 525, 526, 527, 528, 529, 534, 535, 536, 543, 544, 545, 552, 553, 554, 557, 558, 561, 562, 563, 566, 567 }

**F(-2) exception fail** { }

### 2.1.8 Sympy

**A grade** { }

**B grade** { }

**C grade** { }

**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 73, 74, 75, 83, 84, 85, 86, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126,

127, 128, 129, 130, 135, 136, 137, 138, 139, 140, 141, 142, 146, 147, 153, 154, 155, 160, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 180, 181, 182, 183, 184, 185, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 215, 216, 217, 218, 219, 220, 225, 226, 227, 228, 229, 230, 231, 232, 240, 241, 242, 243, 244, 245, 247, 248, 249, 250, 251, 253, 254, 255, 256, 257, 258, 259, 260, 262, 263, 264, 267, 268, 269, 270, 271, 276, 277, 278, 279, 292, 293, 294, 295, 296, 299, 300, 301, 302, 315, 316, 317, 320, 321, 322, 323, 328, 329, 330, 342, 343, 344, 345, 346, 347, 368, 369, 370, 371, 372, 374, 375, 376, 377, 379, 380, 381, 385, 386, 392, 393, 394, 398, 399, 401, 403, 404, 405, 409, 410, 411, 415, 419, 420, 421, 434, 435, 436, 437, 440, 441, 442, 443, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 502, 503, 504, 516, 517, 520, 521, 525, 526, 527, 528, 529, 534, 535, 536, 543, 544, 545, 552, 553, 554, 557, 558, 561, 562, 563, 566, 567 }

**F(-1) timeout fail** { 19, 52, 53, 54, 64, 71, 72, 76, 77, 78, 79, 80, 81, 82, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 131, 132, 133, 134, 143, 144, 145, 150, 151, 152, 156, 157, 158, 159, 161, 162, 178, 179, 186, 187, 188, 189, 190, 191, 192, 193, 214, 221, 222, 223, 224, 233, 234, 238, 246, 252, 284, 285, 286, 287, 288, 289, 290, 291, 310, 314, 334, 335, 336, 337, 338, 339, 340, 341, 355, 356, 357, 358, 359, 360, 361, 362, 363, 366, 367, 384, 388, 389, 390, 391, 397, 402, 407, 408, 413, 414, 416, 417, 418, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 449, 450, 452, 453, 456, 457, 458, 460, 479, 501, 505, 506, 507, 508, 509, 510, 511, 512, 513, 518, 519, 523, 524, 538, 547, 560, 569 }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column N.S. means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	163	91	94	184	113	0	0	0
N.S.	1	1.08	0.60	0.62	1.22	0.75	0.00	0.00	0.00
time (sec)	N/A	0.354	0.118	0.319	0.219	0.263	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	147	166	151	202	108	0	0	0
N.S.	1	1.09	1.23	1.12	1.50	0.80	0.00	0.00	0.00
time (sec)	N/A	0.341	0.073	0.161	0.212	0.266	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	127	89	86	145	103	0	0	0
N.S.	1	1.05	0.74	0.71	1.20	0.85	0.00	0.00	0.00
time (sec)	N/A	0.330	0.081	0.173	0.281	0.265	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	101	100	136	162	98	0	0	0
N.S.	1	1.03	1.02	1.39	1.65	1.00	0.00	0.00	0.00
time (sec)	N/A	0.254	0.110	0.322	0.219	0.263	0.000	0.000	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	86	89	71	71	97	83	0	0	0
N.S.	1	1.03	0.83	0.83	1.13	0.97	0.00	0.00	0.00
time (sec)	N/A	0.279	0.066	0.027	0.202	0.271	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	140	130	131	0	0	0	0	0
N.S.	1	1.20	1.11	1.12	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.658	0.095	0.878	0.000	0.000	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	74	110	92	66	127	0	0	0
N.S.	1	0.97	1.45	1.21	0.87	1.67	0.00	0.00	0.00
time (sec)	N/A	0.325	0.125	0.049	0.331	0.282	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	F	F	F	F(-2)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	141	107	128	0	0	0	0	0
N.S.	1	1.04	0.79	0.95	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.677	0.067	0.303	0.000	0.000	0.000	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	93	127	117	89	146	0	0	0
N.S.	1	1.03	1.41	1.30	0.99	1.62	0.00	0.00	0.00
time (sec)	N/A	0.336	0.182	0.045	0.361	0.283	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	206	200	124	124	319	165	0	0	0
N.S.	1	0.97	0.60	0.60	1.55	0.80	0.00	0.00	0.00
time (sec)	N/A	0.538	0.136	0.595	0.221	0.263	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	231	194	192	346	161	0	0	0
N.S.	1	1.16	0.97	0.96	1.73	0.80	0.00	0.00	0.00
time (sec)	N/A	0.524	0.142	0.512	0.252	0.285	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	177	180	116	116	261	153	0	0	0
N.S.	1	1.02	0.66	0.66	1.47	0.86	0.00	0.00	0.00
time (sec)	N/A	0.511	0.117	0.486	0.213	0.249	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	133	126	170	287	149	0	0	0
N.S.	1	0.98	0.93	1.25	2.11	1.10	0.00	0.00	0.00
time (sec)	N/A	0.289	0.141	0.532	0.252	0.263	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	143	155	99	99	194	133	0	0	0
N.S.	1	1.08	0.69	0.69	1.36	0.93	0.00	0.00	0.00
time (sec)	N/A	0.433	0.104	0.493	0.227	0.255	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	184	241	208	192	0	0	0	0	0
N.S.	1	1.31	1.13	1.04	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.977	0.491	0.952	0.000	0.000	0.000	0.000	0.000



Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	135	145	131	139	143	201	0	0	0
N.S.	1	1.07	0.97	1.03	1.06	1.49	0.00	0.00	0.00
time (sec)	N/A	0.531	0.133	0.490	0.302	0.287	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	F	F	F	F(-2)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	249	182	206	0	0	0	0	0
N.S.	1	1.24	0.91	1.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.028	0.231	0.895	0.000	0.000	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	142	163	135	150	137	213	0	0	0
N.S.	1	1.15	0.95	1.06	0.96	1.50	0.00	0.00	0.00
time (sec)	N/A	0.523	0.128	0.478	0.288	0.288	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	256	241	147	154	465	201	0	0	0
N.S.	1	0.94	0.57	0.60	1.82	0.79	0.00	0.00	0.00
time (sec)	N/A	0.795	0.172	0.585	0.239	0.272	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	230	219	162	233	501	197	0	0	0
N.S.	1	0.95	0.70	1.01	2.18	0.86	0.00	0.00	0.00
time (sec)	N/A	0.486	0.216	0.546	0.221	0.256	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	227	221	139	146	388	189	0	0	0
N.S.	1	0.97	0.61	0.64	1.71	0.83	0.00	0.00	0.00
time (sec)	N/A	0.748	0.180	0.555	0.275	0.250	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	166	161	150	202	423	185	0	0	0
N.S.	1	0.97	0.90	1.22	2.55	1.11	0.00	0.00	0.00
time (sec)	N/A	0.311	0.244	0.543	0.231	0.268	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	191	194	123	130	302	169	0	0	0
N.S.	1	1.02	0.64	0.68	1.58	0.88	0.00	0.00	0.00
time (sec)	N/A	0.647	0.176	0.468	0.243	0.267	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	F	F	F	F(-2)	F(-1)
verified	N/A	No	No	Yes	TBD	TBD	TBD	TBD	TBD
size	239	362	305	245	0	0	0	0	0
N.S.	1	1.51	1.28	1.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.355	0.488	0.948	0.000	0.000	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	180	177	136	178	231	249	0	0	0
N.S.	1	0.98	0.76	0.99	1.28	1.38	0.00	0.00	0.00
time (sec)	N/A	0.743	0.258	0.452	0.329	0.272	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	F	F	F	F(-2)	F(-1)
verified	N/A	No	No	Yes	TBD	TBD	TBD	TBD	TBD
size	267	370	303	258	0	0	0	0	0
N.S.	1	1.39	1.13	0.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.431	0.462	0.997	0.000	0.000	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	195	187	142	195	208	253	0	0	0
N.S.	1	0.96	0.73	1.00	1.07	1.30	0.00	0.00	0.00
time (sec)	N/A	0.792	0.256	0.461	0.500	0.278	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	158	187	227	232	0	0	0	0	0
N.S.	1	1.18	1.44	1.47	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.863	0.367	0.704	0.000	0.000	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	140	146	151	217	0	0	0	0	0
N.S.	1	1.04	1.08	1.55	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.683	0.197	0.614	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	102	102	155	181	0	0	0	0	0
N.S.	1	1.00	1.52	1.77	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.559	0.204	0.616	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	75	85	137	0	0	0	0	0
N.S.	1	1.01	1.15	1.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.439	0.066	0.546	0.000	0.000	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	58	64	180	0	0	0	0	0
N.S.	1	0.98	1.08	3.05	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.350	0.052	2.332	0.000	0.000	0.000	0.000	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	65	129	191	0	0	0	0	0
N.S.	1	1.07	2.11	3.13	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.442	0.267	0.855	0.000	0.000	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	99	132	143	0	0	0	0	0
N.S.	1	1.04	1.39	1.51	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.585	0.192	0.711	0.000	0.000	0.000	0.000	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	118	117	212	243	0	0	0	0	0
N.S.	1	0.99	1.80	2.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.679	0.313	0.699	0.000	0.000	0.000	0.000	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	178	223	190	0	0	0	0	0
N.S.	1	1.13	1.42	1.21	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.866	0.238	0.786	0.000	0.000	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	177	198	244	254	0	0	0	0	0
N.S.	1	1.12	1.38	1.44	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.868	0.775	0.757	0.000	0.000	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	179	180	209	202	0	0	0	0	0
N.S.	1	1.01	1.17	1.13	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.770	0.518	0.658	0.000	0.000	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	124	121	206	192	0	0	0	0	0
N.S.	1	0.98	1.66	1.55	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.568	0.609	0.731	0.000	0.000	0.000	0.000	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	53	64	134	65	0	0	0
N.S.	1	1.00	0.87	1.05	2.20	1.07	0.00	0.00	0.00
time (sec)	N/A	0.238	0.173	0.495	0.273	0.264	0.000	0.000	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	120	118	189	192	0	0	0	0	0
N.S.	1	0.98	1.58	1.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.531	1.048	0.658	0.000	0.000	0.000	0.000	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	116	121	244	254	0	0	0	0	0
N.S.	1	1.04	2.10	2.19	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.658	0.613	0.736	0.000	0.000	0.000	0.000	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	196	283	207	0	0	0	0	0
N.S.	1	1.15	1.66	1.22	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.846	0.545	0.822	0.000	0.000	0.000	0.000	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	152	207	289	280	0	0	0	0	0
N.S.	1	1.36	1.90	1.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.959	1.157	0.763	0.000	0.000	0.000	0.000	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	248	313	377	251	0	0	0	0	0
N.S.	1	1.26	1.52	1.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.289	1.304	0.838	0.000	0.000	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	249	249	287	256	0	0	0	0	0
N.S.	1	1.00	1.15	1.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.962	1.292	0.717	0.000	0.000	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	133	83	136	0	101	0	0	0
N.S.	1	0.98	0.61	1.00	0.00	0.74	0.00	0.00	0.00
time (sec)	N/A	0.329	0.264	0.506	0.000	0.257	0.000	0.000	0.000



Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	186	180	287	256	0	0	0	0	0
N.S.	1	0.97	1.54	1.38	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.803	1.150	0.662	0.000	0.000	0.000	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	89	64	86	0	98	0	0	0
N.S.	1	0.98	0.70	0.95	0.00	1.08	0.00	0.00	0.00
time (sec)	N/A	0.258	0.234	0.480	0.000	0.253	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	180	174	316	256	0	0	0	0	0
N.S.	1	0.97	1.76	1.42	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.725	0.795	0.690	0.000	0.000	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	171	200	328	324	0	0	0	0	0
N.S.	1	1.17	1.92	1.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.933	0.966	0.789	0.000	0.000	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	230	270	362	269	0	0	0	0	0
N.S.	1	1.17	1.57	1.17	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.133	1.193	0.829	0.000	0.000	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	250	312	402	391	0	0	0	0	0
N.S.	1	1.25	1.61	1.56	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.375	1.079	0.832	0.000	0.000	0.000	0.000	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	310	406	471	314	0	0	0	0	0
N.S.	1	1.31	1.52	1.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.659	1.325	0.881	0.000	0.000	0.000	0.000	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	52	77	169	0	0	0	0	0
N.S.	1	0.98	1.45	3.19	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.336	0.037	1.960	0.000	0.000	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	109	107	120	161	0	0	0	0	0
N.S.	1	0.98	1.10	1.48	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.519	0.692	0.479	0.000	0.000	0.000	0.000	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	164	158	223	205	0	0	0	0	0
N.S.	1	0.96	1.36	1.25	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.696	1.793	0.545	0.000	0.000	0.000	0.000	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	278	231	198	878	0	0	0	0	0
N.S.	1	0.83	0.71	3.16	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.149	0.956	0.835	0.000	0.000	0.000	0.000	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	201	176	151	367	0	0	0	0	0
N.S.	1	0.88	0.75	1.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.803	0.849	0.578	0.000	0.000	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	124	124	144	278	0	0	0	0	0
N.S.	1	1.00	1.16	2.24	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.422	0.472	0.750	0.000	0.000	0.000	0.000	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	118	118	137	286	0	0	0	0	0
N.S.	1	1.00	1.16	2.42	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.481	0.419	0.944	0.000	0.000	0.000	0.000	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	B	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	88	88	153	150	462	0	0	0
N.S.	1	0.74	0.74	1.29	1.26	3.88	0.00	0.00	0.00
time (sec)	N/A	0.333	0.091	1.177	0.308	0.290	0.000	0.000	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	199	134	128	1742	146	549	0	0	0
N.S.	1	0.67	0.64	8.75	0.73	2.76	0.00	0.00	0.00
time (sec)	N/A	0.414	0.153	1.144	0.324	0.301	0.000	0.000	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	279	178	146	2537	207	615	0	0	0
N.S.	1	0.64	0.52	9.09	0.74	2.20	0.00	0.00	0.00
time (sec)	N/A	0.477	0.174	1.166	0.296	0.329	0.000	0.000	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	272	175	169	988	205	203	0	0	0
N.S.	1	0.64	0.62	3.63	0.75	0.75	0.00	0.00	0.00
time (sec)	N/A	0.463	0.173	0.687	0.297	0.272	0.000	0.000	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	195	132	126	640	144	176	0	0	0
N.S.	1	0.68	0.65	3.28	0.74	0.90	0.00	0.00	0.00
time (sec)	N/A	0.402	0.122	1.320	0.308	0.267	0.000	0.000	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	118	87	98	356	81	142	0	0	0
N.S.	1	0.74	0.83	3.02	0.69	1.20	0.00	0.00	0.00
time (sec)	N/A	0.301	0.124	0.513	0.247	0.263	0.000	0.000	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	213	150	233	394	0	0	0	0	0
N.S.	1	0.70	1.09	1.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.822	0.654	0.858	0.000	0.000	0.000	0.000	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	235	165	307	436	0	0	0	0	0
N.S.	1	0.70	1.31	1.86	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.824	0.890	1.221	0.000	0.000	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	315	215	290	541	0	0	0	0	0
N.S.	1	0.68	0.92	1.72	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.177	0.852	1.113	0.000	0.000	0.000	0.000	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	360	325	337	783	0	0	0	0	0
N.S.	1	0.90	0.94	2.18	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.507	3.468	0.748	0.000	0.000	0.000	0.000	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	281	270	270	883	0	0	0	0	0
N.S.	1	0.96	0.96	3.14	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.161	1.467	0.714	0.000	0.000	0.000	0.000	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	200	216	235	546	0	0	0	0	0
N.S.	1	1.08	1.18	2.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.659	0.962	0.919	0.000	0.000	0.000	0.000	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	197	209	223	244	0	0	0	0	0
N.S.	1	1.06	1.13	1.24	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.698	0.971	1.020	0.000	0.000	0.000	0.000	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	203	210	259	270	0	0	0	0	0
N.S.	1	1.03	1.28	1.33	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.769	0.835	1.121	0.000	0.000	0.000	0.000	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	98	94	2171	189	572	0	0	0
N.S.	1	0.59	0.57	13.08	1.14	3.45	0.00	0.00	0.00
time (sec)	N/A	0.342	0.066	1.223	0.299	0.310	0.000	0.000	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	247	143	136	3145	163	648	0	0	0
N.S.	1	0.58	0.55	12.73	0.66	2.62	0.00	0.00	0.00
time (sec)	N/A	0.414	0.170	1.173	0.300	0.305	0.000	0.000	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	328	191	154	4262	225	720	0	0	0
N.S.	1	0.58	0.47	12.99	0.69	2.20	0.00	0.00	0.00
time (sec)	N/A	0.526	0.210	1.294	0.293	0.307	0.000	0.000	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	409	231	170	5523	287	792	0	0	0
N.S.	1	0.56	0.42	13.50	0.70	1.94	0.00	0.00	0.00
time (sec)	N/A	0.722	0.282	1.386	0.446	0.330	0.000	0.000	0.000



Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	399	229	209	1846	285	275	0	0	0
N.S.	1	0.57	0.52	4.63	0.71	0.69	0.00	0.00	0.00
time (sec)	N/A	0.669	0.181	0.887	0.407	0.255	0.000	0.000	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	321	186	153	1376	223	245	0	0	0
N.S.	1	0.58	0.48	4.29	0.69	0.76	0.00	0.00	0.00
time (sec)	N/A	0.515	0.212	0.945	0.396	0.267	0.000	0.000	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	143	135	966	161	215	0	0	0
N.S.	1	0.59	0.56	3.98	0.66	0.88	0.00	0.00	0.00
time (sec)	N/A	0.444	0.095	0.528	0.383	0.264	0.000	0.000	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	165	98	107	620	102	185	0	0	0
N.S.	1	0.59	0.65	3.76	0.62	1.12	0.00	0.00	0.00
time (sec)	N/A	0.322	0.186	1.011	0.243	0.261	0.000	0.000	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	292	232	336	499	0	0	0	0	0
N.S.	1	0.79	1.15	1.71	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.166	0.962	2.029	0.000	0.000	0.000	0.000	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	311	241	500	538	0	0	0	0	0
N.S.	1	0.77	1.61	1.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.205	1.475	0.970	0.000	0.000	0.000	0.000	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	321	259	574	570	0	0	0	0	0
N.S.	1	0.81	1.79	1.78	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.199	1.169	1.167	0.000	0.000	0.000	0.000	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	454	431	500	1743	0	0	0	0	0
N.S.	1	0.95	1.10	3.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.020	6.035	0.904	0.000	0.000	0.000	0.000	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	371	376	415	1289	0	0	0	0	0
N.S.	1	1.01	1.12	3.47	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.572	3.698	0.744	0.000	0.000	0.000	0.000	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	293	306	347	885	0	0	0	0	0
N.S.	1	1.04	1.18	3.02	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.881	1.934	0.987	0.000	0.000	0.000	0.000	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	284	316	305	303	0	0	0	0	0
N.S.	1	1.11	1.07	1.07	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.027	1.566	1.094	0.000	0.000	0.000	0.000	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	293	312	319	340	0	0	0	0	0
N.S.	1	1.06	1.09	1.16	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.122	1.314	1.139	0.000	0.000	0.000	0.000	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	293	313	400	2429	0	0	0	0	0
N.S.	1	1.07	1.37	8.29	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.221	2.940	1.221	0.000	0.000	0.000	0.000	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	A	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	219	111	105	3775	224	703	0	0	0
N.S.	1	0.51	0.48	17.24	1.02	3.21	0.00	0.00	0.00
time (sec)	N/A	0.368	0.101	1.331	0.311	0.308	0.000	0.000	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	314	172	147	5007	187	795	0	0	0
N.S.	1	0.55	0.47	15.95	0.60	2.53	0.00	0.00	0.00
time (sec)	N/A	0.439	0.106	1.315	0.343	0.311	0.000	0.000	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	385	203	165	6382	251	879	0	0	0
N.S.	1	0.53	0.43	16.58	0.65	2.28	0.00	0.00	0.00
time (sec)	N/A	0.524	0.125	1.469	0.312	0.335	0.000	0.000	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	458	241	179	2374	313	353	0	0	0
N.S.	1	0.53	0.39	5.18	0.68	0.77	0.00	0.00	0.00
time (sec)	N/A	0.707	0.164	0.970	0.315	0.276	0.000	0.000	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	378	198	163	1840	249	317	0	0	0
N.S.	1	0.52	0.43	4.87	0.66	0.84	0.00	0.00	0.00
time (sec)	N/A	0.537	0.118	0.763	0.284	0.260	0.000	0.000	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	298	153	145	1102	185	281	0	0	0
N.S.	1	0.51	0.49	3.70	0.62	0.94	0.00	0.00	0.00
time (sec)	N/A	0.460	0.087	0.605	0.289	0.253	0.000	0.000	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	218	108	117	956	118	241	0	0	0
N.S.	1	0.50	0.54	4.39	0.54	1.11	0.00	0.00	0.00
time (sec)	N/A	0.332	0.202	0.556	0.203	0.263	0.000	0.000	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	379	326	471	620	0	0	0	0	0
N.S.	1	0.86	1.24	1.64	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.563	2.683	1.179	0.000	0.000	0.000	0.000	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	404	335	596	390	0	0	0	0	0
N.S.	1	0.83	1.48	0.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.631	3.711	1.994	0.000	0.000	0.000	0.000	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	407	343	660	691	0	0	0	0	0
N.S.	1	0.84	1.62	1.70	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.722	1.466	1.080	0.000	0.000	0.000	0.000	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	54	152	0	0	0	0	0
N.S.	1	1.00	0.82	2.30	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.312	0.117	0.937	0.000	0.000	0.000	0.000	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	248	140	670	195	176	0	0	0
N.S.	1	1.05	0.59	2.84	0.83	0.75	0.00	0.00	0.00
time (sec)	N/A	0.722	0.293	0.791	0.303	0.253	0.000	0.000	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	212	220	171	568	0	0	0	0	0
N.S.	1	1.04	0.81	2.68	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.713	0.944	0.938	0.000	0.000	0.000	0.000	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	160	113	382	131	146	0	0	0
N.S.	1	1.03	0.72	2.45	0.84	0.94	0.00	0.00	0.00
time (sec)	N/A	0.488	0.239	0.958	0.322	0.266	0.000	0.000	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	132	141	300	0	0	0	0	0
N.S.	1	1.00	1.07	2.27	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.429	0.635	0.866	0.000	0.000	0.000	0.000	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	85	158	63	117	0	0	0
N.S.	1	1.00	1.18	2.19	0.88	1.62	0.00	0.00	0.00
time (sec)	N/A	0.271	0.182	0.624	0.207	0.258	0.000	0.000	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	53	89	0	0	0	0	0
N.S.	1	1.00	1.00	1.68	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.224	0.028	0.558	0.000	0.000	0.000	0.000	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	86	153	326	0	0	0	0	0
N.S.	1	0.57	1.01	2.16	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.444	0.283	0.895	0.000	0.000	0.000	0.000	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	71	71	208	116	265	0	0	0
N.S.	1	1.00	1.00	2.93	1.63	3.73	0.00	0.00	0.00
time (sec)	N/A	0.295	0.049	1.003	0.321	0.296	0.000	0.000	0.000



Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	238	168	309	431	0	0	0	0	0
N.S.	1	0.71	1.30	1.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.720	0.893	1.114	0.000	0.000	0.000	0.000	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	154	174	192	134	479	0	0	0
N.S.	1	0.99	1.12	1.24	0.86	3.09	0.00	0.00	0.00
time (sec)	N/A	0.485	0.324	1.070	0.322	0.307	0.000	0.000	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	233	162	145	325	0	489	0	0	0
N.S.	1	0.70	0.62	1.39	0.00	2.10	0.00	0.00	0.00
time (sec)	N/A	0.494	0.079	1.343	0.000	0.300	0.000	0.000	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	226	243	192	301	0	0	0	0	0
N.S.	1	1.08	0.85	1.33	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.833	1.238	1.339	0.000	0.000	0.000	0.000	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	113	97	255	157	429	0	0	0
N.S.	1	0.75	0.65	1.70	1.05	2.86	0.00	0.00	0.00
time (sec)	N/A	0.388	0.056	1.076	0.308	0.284	0.000	0.000	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	143	143	159	279	0	0	0	0	0
N.S.	1	1.00	1.11	1.95	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.479	0.615	1.069	0.000	0.000	0.000	0.000	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	136	135	0	327	0	0	0
N.S.	1	1.00	1.79	1.78	0.00	4.30	0.00	0.00	0.00
time (sec)	N/A	0.292	0.389	0.874	0.000	0.296	0.000	0.000	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	72	180	70	0	0	0	0
N.S.	1	1.00	0.86	2.14	0.83	0.00	0.00	0.00	0.00
time (sec)	N/A	0.254	0.027	0.929	0.214	0.000	0.000	0.000	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	229	159	337	483	0	0	0	0	0
N.S.	1	0.69	1.47	2.11	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.730	2.140	1.220	0.000	0.000	0.000	0.000	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	158	122	114	266	144	0	0	0	0
N.S.	1	0.77	0.72	1.68	0.91	0.00	0.00	0.00	0.00
time (sec)	N/A	0.398	0.068	1.217	0.221	0.000	0.000	0.000	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	329	254	439	588	0	0	0	0	0
N.S.	1	0.77	1.33	1.79	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.111	3.677	1.292	0.000	0.000	0.000	0.000	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	250	177	161	367	0	0	0	0	0
N.S.	1	0.71	0.64	1.47	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.513	0.106	1.296	0.000	0.000	0.000	0.000	0.000

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	175	167	398	0	529	0	0	0
N.S.	1	0.72	0.69	1.64	0.00	2.18	0.00	0.00	0.00
time (sec)	N/A	0.488	0.127	1.299	0.000	0.317	0.000	0.000	0.000

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	224	264	225	366	0	0	0	0	0
N.S.	1	1.18	1.00	1.63	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.877	0.715	1.197	0.000	0.000	0.000	0.000	0.000

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	158	138	122	257	175	469	0	0	0
N.S.	1	0.87	0.77	1.63	1.11	2.97	0.00	0.00	0.00
time (sec)	N/A	0.396	0.126	0.941	0.253	0.299	0.000	0.000	0.000

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	108	101	458	169	0	0	0	0
N.S.	1	0.81	0.76	3.44	1.27	0.00	0.00	0.00	0.00
time (sec)	N/A	0.368	0.160	1.186	0.239	0.000	0.000	0.000	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	107	162	220	0	421	0	0	0
N.S.	1	0.84	1.28	1.73	0.00	3.31	0.00	0.00	0.00
time (sec)	N/A	0.316	0.446	1.139	0.000	0.299	0.000	0.000	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	162	179	132	477	157	0	0	0	0
N.S.	1	1.10	0.81	2.94	0.97	0.00	0.00	0.00	0.00
time (sec)	N/A	0.434	0.069	1.010	0.252	0.000	0.000	0.000	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	317	265	389	560	0	0	0	0	0
N.S.	1	0.84	1.23	1.77	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.141	7.067	1.331	0.000	0.000	0.000	0.000	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	248	173	173	434	0	0	0	0	0
N.S.	1	0.70	0.70	1.75	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.512	0.260	1.261	0.000	0.000	0.000	0.000	0.000

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	479	384	534	655	0	0	0	0	0
N.S.	1	0.80	1.11	1.37	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.674	7.208	1.221	0.000	0.000	0.000	0.000	0.000

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	338	226	237	403	276	0	0	0	0
N.S.	1	0.67	0.70	1.19	0.82	0.00	0.00	0.00	0.00
time (sec)	N/A	0.673	0.164	1.150	0.257	0.000	0.000	0.000	0.000

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	<b>F</b>	<b>F(-1)</b>	A	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	259	116	419	191	0	0	142	0
N.S.	1	1.05	0.47	1.70	0.78	0.00	0.00	0.58	0.00
time (sec)	N/A	0.636	0.074	1.372	0.252	0.000	0.000	0.334	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	145	153	93	456	0	0	0	0	0
N.S.	1	1.06	0.64	3.14	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.555	0.359	0.904	0.000	0.000	0.000	0.000	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	114	74	311	62	101	0	0	0
N.S.	1	1.04	0.67	2.83	0.56	0.92	0.00	0.00	0.00
time (sec)	N/A	0.381	0.094	1.181	0.925	0.259	0.000	0.000	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	75	223	0	0	0	0	0
N.S.	1	1.00	0.85	2.53	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.351	0.158	0.675	0.000	0.000	0.000	0.000	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	A	<b>F</b>	C	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	55	123	28	72	0	40	0
N.S.	1	1.00	1.12	2.51	0.57	1.47	0.00	0.82	0.00
time (sec)	N/A	0.230	0.069	0.851	0.760	0.264	0.000	0.289	0.000

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	45	51	0	0	0	0	0
N.S.	1	1.00	1.41	1.59	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.199	0.017	0.464	0.000	0.000	0.000	0.000	0.000

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	66	113	270	0	0	0	0	0
N.S.	1	0.64	1.10	2.62	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.388	0.148	0.871	0.000	0.000	0.000	0.000	0.000

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	B	<b>F</b>	C	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	57	168	73	89	0	80	0
N.S.	1	1.00	1.19	3.50	1.52	1.85	0.00	1.67	0.00
time (sec)	N/A	0.252	0.024	0.918	0.473	0.276	0.000	0.297	0.000

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	167	125	234	349	0	0	0	0	0
N.S.	1	0.75	1.40	2.09	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.576	0.276	0.986	0.000	0.000	0.000	0.000	0.000

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	98	98	100	0	0	0	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.309	0.082	0.000	0.000	0.000	0.000	0.000	0.000



Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	141	141	115	0	0	0	0	0	0
N.S.	1	1.00	0.82	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.359	0.031	0.000	0.000	0.000	0.000	0.000	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	429	423	387	0	0	0	0	0	0
N.S.	1	0.99	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.388	0.788	0.000	0.000	0.000	0.000	0.000	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	307	306	290	0	0	0	0	0	0
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.719	0.310	0.000	0.000	0.000	0.000	0.000	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	184	195	191	0	0	0	0	0	0
N.S.	1	1.06	1.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.476	0.171	0.000	0.000	0.000	0.000	0.000	0.000

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	27	27	29	27	32	31	41	31	29
N.S.	1	1.00	1.07	1.00	1.19	1.15	1.52	1.15	1.07
time (sec)	N/A	0.270	3.757	1.338	0.809	0.261	4.546	0.286	3.038

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	27	27	29	27	30	43	58	30	29
N.S.	1	1.00	1.07	1.00	1.11	1.59	2.15	1.11	1.07
time (sec)	N/A	0.556	9.494	1.354	0.758	0.261	64.463	0.289	3.051

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	27	27	29	27	32	57	0	31	29
N.S.	1	1.00	1.07	1.00	1.19	2.11	0.00	1.15	1.07
time (sec)	N/A	0.842	12.265	1.210	0.721	0.275	0.000	0.295	3.060

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	723	548	350	0	0	0	0	0	0
N.S.	1	0.76	0.48	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.704	0.927	0.000	0.000	0.000	0.000	0.000	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	455	396	274	0	0	0	0	0	0
N.S.	1	0.87	0.60	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.102	0.550	0.000	0.000	0.000	0.000	0.000	0.000

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	278	265	223	0	0	0	0	0	0
N.S.	1	0.95	0.80	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.714	0.210	0.000	0.000	0.000	0.000	0.000	0.000

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	176	176	147	0	0	0	0	0	0
N.S.	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.349	0.059	0.000	0.000	0.000	0.000	0.000	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	300	297	216	0	0	0	0	0	0
N.S.	1	0.99	0.72	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.637	0.178	0.000	0.000	0.000	0.000	0.000	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	450	430	319	0	0	0	0	0	0
N.S.	1	0.96	0.71	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.975	0.499	0.000	0.000	0.000	0.000	0.000	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	817	596	387	0	0	0	0	0	0
N.S.	1	0.73	0.47	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.600	1.648	0.000	0.000	0.000	0.000	0.000	0.000

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	503	428	288	0	0	0	0	0	0
N.S.	1	0.85	0.57	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.653	0.765	0.000	0.000	0.000	0.000	0.000	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	302	283	229	0	0	0	0	0	0
N.S.	1	0.94	0.76	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.906	0.155	0.000	0.000	0.000	0.000	0.000	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	188	188	153	0	0	0	0	0	0
N.S.	1	1.00	0.81	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.573	0.316	0.000	0.000	0.000	0.000	0.000	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	336	330	225	0	0	0	0	0	0
N.S.	1	0.98	0.67	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.195	0.324	0.000	0.000	0.000	0.000	0.000	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	504	484	413	0	0	0	0	0	0
N.S.	1	0.96	0.82	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.836	0.742	0.000	0.000	0.000	0.000	0.000	0.000

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	128	128	124	0	0	0	0	0	0
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.306	0.064	0.000	0.000	0.000	0.000	0.000	0.000

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	266	326	125	188	178	175	0	0	0
N.S.	1	1.23	0.47	0.71	0.67	0.66	0.00	0.00	0.00
time (sec)	N/A	2.139	0.190	0.496	0.295	0.257	0.000	0.000	0.000

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	195	226	101	140	134	142	0	0	0
N.S.	1	1.16	0.52	0.72	0.69	0.73	0.00	0.00	0.00
time (sec)	N/A	1.323	0.251	0.498	0.250	0.260	0.000	0.000	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	112	128	73	90	76	95	0	0	0
N.S.	1	1.14	0.65	0.80	0.68	0.85	0.00	0.00	0.00
time (sec)	N/A	0.734	0.159	0.140	0.272	0.261	0.000	0.000	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	86	95	187	0	0	0	0	0
N.S.	1	0.88	0.97	1.91	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.463	0.059	0.458	0.000	0.000	0.000	0.000	0.000

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	163	157	191	255	0	0	0	0	0
N.S.	1	0.96	1.17	1.56	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.628	0.891	0.484	0.000	0.000	0.000	0.000	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	258	256	319	320	0	0	0	0	0
N.S.	1	0.99	1.24	1.24	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.993	3.303	0.498	0.000	0.000	0.000	0.000	0.000

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	371	425	237	1284	326	349	0	0	0
N.S.	1	1.15	0.64	3.46	0.88	0.94	0.00	0.00	0.00
time (sec)	N/A	2.174	0.428	1.492	0.450	0.281	0.000	0.000	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	319	328	241	678	0	0	0	0	0
N.S.	1	1.03	0.76	2.13	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.872	2.125	0.654	0.000	0.000	0.000	0.000	0.000

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	163	181	726	204	280	0	0	0
N.S.	1	0.88	0.97	3.90	1.10	1.51	0.00	0.00	0.00
time (sec)	N/A	0.570	0.578	0.561	0.337	0.264	0.000	0.000	0.000

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	204	181	235	527	0	0	0	0	0
N.S.	1	0.89	1.15	2.58	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.751	1.164	0.770	0.000	0.000	0.000	0.000	0.000

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	402	238	449	0	0	0	0	0	0
N.S.	1	0.59	1.12	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.326	1.234	0.000	0.000	0.000	0.000	0.000	0.000

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	234	186	270	582	0	0	0	0	0
N.S.	1	0.79	1.15	2.49	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.847	1.695	0.856	0.000	0.000	0.000	0.000	0.000



Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	427	250	5035	0	0	0	0	0	0
N.S.	1	0.59	11.79	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.860	66.458	0.000	0.000	0.000	0.000	0.000	0.000

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	336	213	304	1751	0	0	0	0	0
N.S.	1	0.63	0.90	5.21	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.181	0.898	1.220	0.000	0.000	0.000	0.000	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	495	663	262	1952	388	432	0	0	0
N.S.	1	1.34	0.53	3.94	0.78	0.87	0.00	0.00	0.00
time (sec)	N/A	3.756	0.513	0.831	0.309	0.277	0.000	0.000	0.000

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	441	550	485	1721	0	0	0	0	0
N.S.	1	1.25	1.10	3.90	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.177	4.827	0.846	0.000	0.000	0.000	0.000	0.000

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	348	221	208	1270	278	367	0	0	0
N.S.	1	0.64	0.60	3.65	0.80	1.05	0.00	0.00	0.00
time (sec)	N/A	0.806	1.481	1.181	0.235	0.283	0.000	0.000	0.000

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	336	356	374	1061	0	0	0	0	0
N.S.	1	1.06	1.11	3.16	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.535	3.346	1.049	0.000	0.000	0.000	0.000	0.000

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	573	396	650	0	0	0	0	0	0
N.S.	1	0.69	1.13	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.499	2.181	0.000	0.000	0.000	0.000	0.000	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	453	393	433	464	0	0	0	0	0
N.S.	1	0.87	0.96	1.02	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.362	4.189	1.078	0.000	0.000	0.000	0.000	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	630	387	5444	0	0	0	0	0	0
N.S.	1	0.61	8.64	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.650	63.651	0.000	0.000	0.000	0.000	0.000	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	426	403	583	500	0	0	0	0	0
N.S.	1	0.95	1.37	1.17	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.237	1.873	1.231	0.000	0.000	0.000	0.000	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	<b>F</b>	A	B	A	A	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	880	0	288	2224	471	558	0	0	0
N.S.	1	0.00	0.33	2.53	0.54	0.63	0.00	0.00	0.00
time (sec)	N/A	0.000	0.603	0.775	0.308	0.276	0.000	0.000	0.000

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	<b>F</b>	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	N/A	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	841	0	910	2528	0	0	0	0	0
N.S.	1	0.00	1.08	3.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	5.916	0.878	0.000	0.000	0.000	0.000	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	470	259	234	1958	337	477	0	0	0
N.S.	1	0.55	0.50	4.17	0.72	1.01	0.00	0.00	0.00
time (sec)	N/A	1.017	1.512	0.742	0.219	0.278	0.000	0.000	0.000

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	486	561	740	1737	0	0	0	0	0
N.S.	1	1.15	1.52	3.57	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.372	3.647	1.054	0.000	0.000	0.000	0.000	0.000

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	836	613	963	0	0	0	0	0	0
N.S.	1	0.73	1.15	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.455	5.837	0.000	0.000	0.000	0.000	0.000	0.000

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	No	No	Yes	TBD	TBD	TBD	TBD	TBD
size	607	663	554	589	0	0	0	0	0
N.S.	1	1.09	0.91	0.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.332	5.260	1.243	0.000	0.000	0.000	0.000	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	890	609	5694	0	0	0	0	0	0
N.S.	1	0.68	6.40	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.564	65.179	0.000	0.000	0.000	0.000	0.000	0.000

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	<b>F</b>	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	N/A	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	638	0	803	652	0	0	0	0	0
N.S.	1	0.00	1.26	1.02	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	2.728	1.290	0.000	0.000	0.000	0.000	0.000

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	421	470	255	1314	407	348	0	0	0
N.S.	1	1.12	0.61	3.12	0.97	0.83	0.00	0.00	0.00
time (sec)	N/A	1.663	0.478	0.911	0.304	0.270	0.000	0.000	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	355	372	295	1092	0	0	0	0	0
N.S.	1	1.05	0.83	3.08	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.405	1.545	0.931	0.000	0.000	0.000	0.000	0.000

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	292	271	201	752	279	282	0	0	0
N.S.	1	0.93	0.69	2.58	0.96	0.97	0.00	0.00	0.00
time (sec)	N/A	0.880	0.428	0.944	0.298	0.275	0.000	0.000	0.000

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	226	189	228	563	0	0	0	0	0
N.S.	1	0.84	1.01	2.49	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.687	1.303	0.838	0.000	0.000	0.000	0.000	0.000

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	108	149	314	145	218	0	0	0
N.S.	1	0.70	0.96	2.03	0.94	1.41	0.00	0.00	0.00
time (sec)	N/A	0.352	0.736	0.593	0.224	0.263	0.000	0.000	0.000

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	53	149	0	0	0	0	0
N.S.	1	1.00	1.00	2.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.247	0.038	0.483	0.000	0.000	0.000	0.000	0.000

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	273	138	315	0	0	0	0	0	0
N.S.	1	0.51	1.15	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.659	0.672	0.000	0.000	0.000	0.000	0.000	0.000

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	186	138	237	476	0	0	0	0	0
N.S.	1	0.74	1.27	2.56	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.739	0.909	1.152	0.000	0.000	0.000	0.000	0.000

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	430	254	5036	0	0	0	0	0	0
N.S.	1	0.59	11.71	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.405	63.656	0.000	0.000	0.000	0.000	0.000	0.000

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	328	263	370	1521	0	0	0	0	0
N.S.	1	0.80	1.13	4.64	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.455	0.991	1.208	0.000	0.000	0.000	0.000	0.000

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	556	535	392	781	0	0	0	0	0
N.S.	1	0.96	0.71	1.40	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.094	2.995	1.456	0.000	0.000	0.000	0.000	0.000

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	440	412	343	756	0	0	0	0	0
N.S.	1	0.94	0.78	1.72	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.469	1.693	1.458	0.000	0.000	0.000	0.000	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	413	287	319	648	0	0	0	0	0
N.S.	1	0.69	0.77	1.57	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.737	1.424	1.201	0.000	0.000	0.000	0.000	0.000

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	257	199	270	738	0	0	0	0	0
N.S.	1	0.77	1.05	2.87	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.269	2.119	1.086	0.000	0.000	0.000	0.000	0.000



Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	125	219	341	0	0	0	0	0
N.S.	1	0.64	1.12	1.74	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.743	1.426	0.987	0.000	0.000	0.000	0.000	0.000

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	198	140	126	578	0	0	0	0	0
N.S.	1	0.71	0.64	2.92	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.656	0.317	0.975	0.000	0.000	0.000	0.000	0.000

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	471	260	613	0	0	0	0	0	0
N.S.	1	0.55	1.30	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.330	3.077	0.000	0.000	0.000	0.000	0.000	0.000

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	341	279	127	1271	0	0	0	0	0
N.S.	1	0.82	0.37	3.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.465	0.516	1.475	0.000	0.000	0.000	0.000	0.000

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	650	433	5362	0	0	0	0	0	0
N.S.	1	0.67	8.25	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.981	63.672	0.000	0.000	0.000	0.000	0.000	0.000

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	496	472	529	2163	0	0	0	0	0
N.S.	1	0.95	1.07	4.36	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.181	1.968	1.402	0.000	0.000	0.000	0.000	0.000

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	568	569	490	922	0	0	0	0	0
N.S.	1	1.00	0.86	1.62	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.474	4.390	1.457	0.000	0.000	0.000	0.000	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	482	461	382	2901	0	0	0	0	0
N.S.	1	0.96	0.79	6.02	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.708	1.962	1.354	0.000	0.000	0.000	0.000	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	336	330	356	635	0	0	0	0	0
N.S.	1	0.98	1.06	1.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.666	3.675	1.003	0.000	0.000	0.000	0.000	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	389	249	264	2657	0	0	0	0	0
N.S.	1	0.64	0.68	6.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.294	1.836	1.295	0.000	0.000	0.000	0.000	0.000

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	298	189	388	582	0	0	0	0	0
N.S.	1	0.63	1.30	1.95	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.940	2.495	1.191	0.000	0.000	0.000	0.000	0.000

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	331	278	289	2435	0	0	0	0	0
N.S.	1	0.84	0.87	7.36	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.709	1.643	1.161	0.000	0.000	0.000	0.000	0.000

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	597	448	818	0	0	0	0	0	0
N.S.	1	0.75	1.37	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.986	9.527	0.000	0.000	0.000	0.000	0.000	0.000

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	476	469	457	2857	0	0	0	0	0
N.S.	1	0.99	0.96	6.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.142	2.851	1.500	0.000	0.000	0.000	0.000	0.000

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	B	F	F	F	F(-1)	F	F(-1)
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD
size	796	0	5532	0	0	0	0	0	0
N.S.	1	0.00	6.95	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	65.919	0.000	0.000	0.000	0.000	0.000	0.000

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	562	750	534	3745	0	0	0	0	0
N.S.	1	1.33	0.95	6.66	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	6.305	2.796	1.387	0.000	0.000	0.000	0.000	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	429	407	220	794	0	0	0	0	0
N.S.	1	0.95	0.51	1.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.371	1.073	1.227	0.000	0.000	0.000	0.000	0.000

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	295	116	488	0	0	0	0	0
N.S.	1	1.21	0.48	2.01	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.128	0.358	0.832	0.000	0.000	0.000	0.000	0.000

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	177	215	123	343	105	150	0	0	0
N.S.	1	1.21	0.69	1.94	0.59	0.85	0.00	0.00	0.00
time (sec)	N/A	0.755	0.108	1.120	0.301	0.274	0.000	0.000	0.000

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	140	87	239	0	0	0	0	0
N.S.	1	0.93	0.58	1.58	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.536	0.166	0.624	0.000	0.000	0.000	0.000	0.000

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	<b>F</b>	C	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	80	54	139	50	114	0	76	0
N.S.	1	1.01	0.68	1.76	0.63	1.44	0.00	0.96	0.00
time (sec)	N/A	0.317	0.078	0.520	0.226	0.272	0.000	0.306	0.000

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	45	51	0	0	0	0	0
N.S.	1	1.00	1.41	1.59	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.211	0.018	0.530	0.000	0.000	0.000	0.000	0.000

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	183	106	151	0	0	0	0	0	0
N.S.	1	0.58	0.83	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.537	0.171	0.000	0.000	0.000	0.000	0.000	0.000

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	124	103	111	241	0	0	0	0	0
N.S.	1	0.83	0.90	1.94	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.574	0.371	0.898	0.000	0.000	0.000	0.000	0.000

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	296	194	233	0	0	0	0	0	0
N.S.	1	0.66	0.79	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.024	0.776	0.000	0.000	0.000	0.000	0.000	0.000

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	136	0	0	31
N.S.	1	1.00	1.06	0.94	1.00	4.39	0.00	0.00	1.00
time (sec)	N/A	5.211	2.228	2.790	0.423	0.265	0.000	0.000	3.222

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	87	0	0	31
N.S.	1	1.00	1.06	0.94	1.00	2.81	0.00	0.00	1.00
time (sec)	N/A	2.412	1.100	2.766	0.385	0.290	0.000	0.000	3.442

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	43	32	0	31
N.S.	1	1.00	1.06	0.94	1.00	1.39	1.03	0.00	1.00
time (sec)	N/A	1.013	0.765	3.006	0.347	0.274	72.012	0.000	3.053

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	58	32	31	31
N.S.	1	1.00	1.06	0.94	1.00	1.87	1.03	1.00	1.00
time (sec)	N/A	0.334	3.776	1.649	0.364	0.270	22.635	0.359	3.155

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	70	32	31	31
N.S.	1	1.00	1.06	0.94	1.00	2.26	1.03	1.00	1.00
time (sec)	N/A	0.340	4.457	3.079	0.435	0.259	26.558	0.365	3.383

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	84	0	31	31
N.S.	1	1.00	1.06	0.94	1.00	2.71	0.00	1.00	1.00
time (sec)	N/A	0.341	4.765	3.251	0.425	0.261	0.000	0.375	3.440

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	28	24	26	38	27	26	26
N.S.	1	1.00	1.08	0.92	1.00	1.46	1.04	1.00	1.00
time (sec)	N/A	0.271	0.736	1.497	0.327	0.262	11.446	0.358	3.240



Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	505	665	179	294	276	248	0	0	0
N.S.	1	1.32	0.35	0.58	0.55	0.49	0.00	0.00	0.00
time (sec)	N/A	4.798	0.355	0.549	0.207	0.263	0.000	0.000	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	388	431	147	218	210	204	0	0	0
N.S.	1	1.11	0.38	0.56	0.54	0.53	0.00	0.00	0.00
time (sec)	N/A	2.577	0.128	0.503	0.219	0.258	0.000	0.000	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	223	109	140	124	140	0	0	0
N.S.	1	1.27	0.62	0.80	0.71	0.80	0.00	0.00	0.00
time (sec)	N/A	1.262	0.151	0.146	0.195	0.256	0.000	0.000	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	144	128	129	253	0	0	0	0	0
N.S.	1	0.89	0.90	1.76	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.602	0.055	0.612	0.000	0.000	0.000	0.000	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	260	244	276	416	0	0	0	0	0
N.S.	1	0.94	1.06	1.60	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.556	1.406	0.542	0.000	0.000	0.000	0.000	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	387	414	455	527	0	0	0	0	0
N.S.	1	1.07	1.18	1.36	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.784	6.826	0.549	0.000	0.000	0.000	0.000	0.000

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	605	651	189	887	0	0	0	0	0
N.S.	1	1.08	0.31	1.47	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	6.940	0.920	1.311	0.000	0.000	0.000	0.000	0.000

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	402	390	148	536	0	0	0	0	0
N.S.	1	0.97	0.37	1.33	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.881	0.408	1.088	0.000	0.000	0.000	0.000	0.000

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	231	182	98	256	0	0	0	0	0
N.S.	1	0.79	0.42	1.11	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.278	0.207	0.727	0.000	0.000	0.000	0.000	0.000

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	55	0	0	0	0	0
N.S.	1	1.00	1.00	1.20	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.226	0.023	0.562	0.000	0.000	0.000	0.000	0.000

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	241	143	162	548	0	0	0	0	0
N.S.	1	0.59	0.67	2.27	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.770	0.225	1.143	0.000	0.000	0.000	0.000	0.000

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	C	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	413	292	270	955	0	0	0	0	0
N.S.	1	0.71	0.65	2.31	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.119	0.900	1.320	0.000	0.000	0.000	0.000	0.000

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	C	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	607	498	375	1319	0	0	0	0	0
N.S.	1	0.82	0.62	2.17	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.001	1.638	1.339	0.000	0.000	0.000	0.000	0.000

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	315	345	136	520	0	0	0	0	0
N.S.	1	1.10	0.43	1.65	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.173	0.567	0.888	0.000	0.000	0.000	0.000	0.000

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	259	140	375	131	205	0	0	0
N.S.	1	1.07	0.58	1.54	0.54	0.84	0.00	0.00	0.00
time (sec)	N/A	2.023	0.134	1.377	0.301	0.255	0.000	0.000	0.000

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	188	158	98	255	0	0	0	0	0
N.S.	1	0.84	0.52	1.36	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.151	0.216	0.696	0.000	0.000	0.000	0.000	0.000

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	<b>F</b>	C	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	95	101	155	65	159	0	103	0
N.S.	1	0.86	0.92	1.41	0.59	1.45	0.00	0.94	0.00
time (sec)	N/A	0.501	0.082	0.597	0.219	0.257	0.000	0.345	0.000

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	45	51	0	0	0	0	0
N.S.	1	1.00	1.41	1.59	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.213	0.020	0.522	0.000	0.000	0.000	0.000	0.000

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	265	154	488	0	0	0	0	0	0
N.S.	1	0.58	1.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.681	0.508	0.000	0.000	0.000	0.000	0.000	0.000

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	166	125	137	313	0	0	0	0	0
N.S.	1	0.75	0.83	1.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.705	0.436	1.050	0.000	0.000	0.000	0.000	0.000

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	460	273	1051	0	0	0	0	0	0
N.S.	1	0.59	2.28	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.496	4.225	0.000	0.000	0.000	0.000	0.000	0.000

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	30	30	32	28	30	68	31	30	30
N.S.	1	1.00	1.07	0.93	1.00	2.27	1.03	1.00	1.00
time (sec)	N/A	0.313	3.943	1.145	0.373	0.260	70.004	0.366	3.170

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	61	45	44	0	0	0	0	0
N.S.	1	0.91	0.67	0.66	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.364	0.374	0.540	0.000	0.000	0.000	0.000	0.000

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	48	34	33	0	0	0	0	0
N.S.	1	0.96	0.68	0.66	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.327	0.151	0.488	0.000	0.000	0.000	0.000	0.000

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	33	25	24	0	0	0	0	0
N.S.	1	1.14	0.86	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.299	0.094	0.100	0.000	0.000	0.000	0.000	0.000

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	25	24	22	24	22
N.S.	1	1.00	1.10	1.00	1.25	1.20	1.10	1.20	1.10
time (sec)	N/A	0.208	1.582	0.569	0.281	0.236	1.310	0.286	2.772

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	23	36	36	23	22
N.S.	1	1.00	1.10	1.00	1.15	1.80	1.80	1.15	1.10
time (sec)	N/A	0.210	5.058	0.282	0.286	0.247	4.231	0.286	3.043

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	339	190	188	297	0	0	0	0	0
N.S.	1	0.56	0.55	0.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.664	0.382	0.925	0.000	0.000	0.000	0.000	0.000

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	297	173	171	250	0	0	0	0	0
N.S.	1	0.58	0.58	0.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.642	0.344	0.882	0.000	0.000	0.000	0.000	0.000

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	139	90	103	165	0	0	0	0	0
N.S.	1	0.65	0.74	1.19	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.492	0.307	0.604	0.000	0.000	0.000	0.000	0.000

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	197	123	127	186	0	0	0	0	0
N.S.	1	0.62	0.64	0.94	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.501	0.293	0.513	0.000	0.000	0.000	0.000	0.000

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	139	90	105	165	0	0	0	0	0
N.S.	1	0.65	0.76	1.19	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.402	0.312	0.664	0.000	0.000	0.000	0.000	0.000



Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	28	26	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.00	0.93	0.00	1.00
time (sec)	N/A	0.628	1.534	0.971	0.306	0.256	1.205	0.000	3.052

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	32	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.14	0.96	1.00	1.00
time (sec)	N/A	0.502	1.796	0.917	0.289	0.243	1.657	0.310	2.699

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	32	27	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.14	0.96	0.00	1.00
time (sec)	N/A	0.299	7.707	1.078	0.286	0.236	3.077	0.000	3.050

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	32	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.14	0.96	1.00	1.00
time (sec)	N/A	0.300	1.248	1.109	0.277	0.253	8.026	0.303	2.906

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	397	224	215	318	0	0	0	0	0
N.S.	1	0.56	0.54	0.80	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.710	0.712	0.530	0.000	0.000	0.000	0.000	0.000

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	339	191	188	297	0	0	0	0	0
N.S.	1	0.56	0.55	0.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.616	0.624	0.711	0.000	0.000	0.000	0.000	0.000

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	297	174	172	252	0	0	0	0	0
N.S.	1	0.59	0.58	0.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.579	0.600	0.602	0.000	0.000	0.000	0.000	0.000

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	239	141	147	231	0	0	0	0	0
N.S.	1	0.59	0.62	0.97	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.475	0.549	0.859	0.000	0.000	0.000	0.000	0.000

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	28	26	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.00	0.93	0.00	1.00
time (sec)	N/A	0.982	1.628	1.149	0.286	0.254	10.390	0.000	2.908

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	32	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.14	0.96	1.00	1.00
time (sec)	N/A	0.812	2.406	0.905	0.282	0.249	11.347	0.316	3.082

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	32	27	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.14	0.96	0.00	1.00
time (sec)	N/A	0.324	8.156	1.054	0.272	0.249	27.240	0.000	3.146

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	32	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.14	0.96	1.00	1.00
time (sec)	N/A	0.316	1.336	1.253	0.279	0.248	71.907	0.311	2.964

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	397	223	216	320	0	0	0	0	0
N.S.	1	0.56	0.54	0.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.731	1.054	0.596	0.000	0.000	0.000	0.000	0.000

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	439	240	233	365	0	0	0	0	0
N.S.	1	0.55	0.53	0.83	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.748	1.040	0.802	0.000	0.000	0.000	0.000	0.000

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	397	223	216	318	0	0	0	0	0
N.S.	1	0.56	0.54	0.80	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.638	0.978	0.513	0.000	0.000	0.000	0.000	0.000

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	339	190	191	297	0	0	0	0	0
N.S.	1	0.56	0.56	0.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.570	0.756	0.949	0.000	0.000	0.000	0.000	0.000

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	45	0	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.61	0.00	0.00	1.00
time (sec)	N/A	1.381	1.536	1.017	0.306	0.246	0.000	0.000	2.724

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	49	0	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.75	0.00	1.00	1.00
time (sec)	N/A	1.200	2.591	0.980	0.309	0.250	0.000	0.307	2.761

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	49	0	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.75	0.00	0.00	1.00
time (sec)	N/A	0.315	7.868	1.283	0.318	0.255	0.000	0.000	2.707

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	49	0	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.75	0.00	1.00	1.00
time (sec)	N/A	0.321	1.528	1.304	0.314	0.261	0.000	0.318	2.709

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	55	69	91	0	0	0	0	0
N.S.	1	0.56	0.70	0.93	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.418	0.097	0.802	0.000	0.000	0.000	0.000	0.000

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	44	60	78	0	0	0	0	0
N.S.	1	0.68	0.92	1.20	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.403	0.079	1.030	0.000	0.000	0.000	0.000	0.000

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	44	60	67	0	0	0	0	0
N.S.	1	0.68	0.92	1.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.431	0.086	0.586	0.000	0.000	0.000	0.000	0.000

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	50	58	0	0	0	0	0
N.S.	1	1.00	1.79	2.07	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.317	0.071	0.778	0.000	0.000	0.000	0.000	0.000

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	47	48	0	55	0	0	0
N.S.	1	1.00	1.68	1.71	0.00	1.96	0.00	0.00	0.00
time (sec)	N/A	0.232	0.055	0.664	0.000	0.249	0.000	0.000	0.000

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	22	24	35	24	24	24
N.S.	1	1.00	1.08	0.92	1.00	1.46	1.00	1.00	1.00
time (sec)	N/A	0.262	0.687	1.542	0.281	0.240	1.372	0.308	2.736

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	22	24	37	26	24	24
N.S.	1	1.00	1.08	0.92	1.00	1.54	1.08	1.00	1.00
time (sec)	N/A	0.277	0.891	2.220	0.279	0.245	4.129	0.292	2.718

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	197	123	130	182	0	0	0	0	0
N.S.	1	0.62	0.66	0.92	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.553	0.468	1.236	0.000	0.000	0.000	0.000	0.000

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	139	90	99	161	0	0	0	0	0
N.S.	1	0.65	0.71	1.16	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.481	0.383	0.764	0.000	0.000	0.000	0.000	0.000

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	68	81	116	0	0	0	0	0
N.S.	1	0.74	0.88	1.26	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.511	0.277	0.506	0.000	0.000	0.000	0.000	0.000

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	54	55	0	65	0	0	0
N.S.	1	1.00	1.54	1.57	0.00	1.86	0.00	0.00	0.00
time (sec)	N/A	0.227	0.140	0.692	0.000	0.252	0.000	0.000	0.000

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	49	27	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.75	0.96	0.00	1.00
time (sec)	N/A	0.294	1.663	1.174	0.283	0.252	2.059	0.000	2.754



Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	53	29	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.89	1.04	1.00	1.00
time (sec)	N/A	0.304	2.373	1.110	0.280	0.243	3.676	0.313	2.862

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	63	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	2.25	0.96	1.00	1.00
time (sec)	N/A	0.313	4.836	0.516	0.322	0.254	8.695	0.344	3.170

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	28	24	26	61	26	0	26
N.S.	1	1.00	1.08	0.92	1.00	2.35	1.00	0.00	1.00
time (sec)	N/A	0.275	6.911	1.050	0.295	0.268	8.073	0.000	2.814

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	25	25	27	23	25	60	26	25	25
N.S.	1	1.00	1.08	0.92	1.00	2.40	1.04	1.00	1.00
time (sec)	N/A	0.232	0.193	0.925	0.297	0.255	6.765	0.325	2.724

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	64	27	0	28
N.S.	1	1.00	1.07	0.93	1.00	2.29	0.96	0.00	1.00
time (sec)	N/A	0.317	8.682	1.000	0.307	0.238	21.220	0.000	3.192

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	68	29	28	28
N.S.	1	1.00	1.07	0.93	1.00	2.43	1.04	1.00	1.00
time (sec)	N/A	0.312	9.981	1.092	0.309	0.255	48.171	0.336	2.861

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	38	0	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.36	0.00	0.00	1.00
time (sec)	N/A	0.300	1.452	1.651	0.285	0.259	0.000	0.000	2.741

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	28	27	0	28
N.S.	1	1.00	1.07	0.93	1.00	1.00	0.96	0.00	1.00
time (sec)	N/A	0.289	0.280	1.353	0.287	0.249	2.672	0.000	2.713

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	50	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.79	0.96	1.00	1.00
time (sec)	N/A	0.297	1.020	1.207	0.288	0.265	2.979	0.325	3.349

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	63	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	2.25	0.96	1.00	1.00
time (sec)	N/A	0.303	1.487	1.457	0.324	0.262	83.053	0.333	3.471

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	86	0	28	28
N.S.	1	1.00	1.07	0.93	1.00	3.07	0.00	1.00	1.00
time (sec)	N/A	0.301	1.912	1.987	0.319	0.257	0.000	0.346	3.827

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	98	83	257	107	0	0	0	0	0
N.S.	1	0.85	2.62	1.09	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.638	0.528	0.849	0.000	0.000	0.000	0.000	0.000

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	82	73	194	87	0	0	0	0	0
N.S.	1	0.89	2.37	1.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.604	0.344	0.490	0.000	0.000	0.000	0.000	0.000

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	57	65	61	0	0	0	0	0
N.S.	1	0.98	1.12	1.05	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.541	0.280	0.210	0.000	0.000	0.000	0.000	0.000

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	259	24	26	24	22
N.S.	1	1.00	1.10	1.00	12.95	1.20	1.30	1.20	1.10
time (sec)	N/A	0.514	2.897	0.661	0.415	0.248	2.633	0.300	2.703

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	351	36	41	23	22
N.S.	1	1.00	1.10	1.00	17.55	1.80	2.05	1.15	1.10
time (sec)	N/A	0.497	8.902	0.266	0.430	0.243	12.413	0.301	2.686

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	350	352	322	555	0	0	0	0	0
N.S.	1	1.01	0.92	1.59	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.051	0.559	1.308	0.000	0.000	0.000	0.000	0.000

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	262	130	279	0	0	0	0	0
N.S.	1	1.70	0.84	1.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.267	0.435	0.583	0.000	0.000	0.000	0.000	0.000

Problem 322	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	248	252	217	405	0	0	0	0	0
N.S.	1	1.02	0.88	1.63	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.298	0.381	0.636	0.000	0.000	0.000	0.000	0.000

Problem 323	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	146	131	121	269	0	0	0	0	0
N.S.	1	0.90	0.83	1.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.746	0.205	0.765	0.000	0.000	0.000	0.000	0.000

Problem 324	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	442	43	27	0	28
N.S.	1	1.00	1.07	0.93	15.79	1.54	0.96	0.00	1.00
time (sec)	N/A	1.024	9.975	1.041	0.692	0.253	2.860	0.000	3.000

Problem 325	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	427	49	29	28	28
N.S.	1	1.00	1.07	0.93	15.25	1.75	1.04	1.00	1.00
time (sec)	N/A	0.422	3.957	0.967	0.571	0.258	7.052	0.324	2.860

Problem 326	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	453	49	29	0	28
N.S.	1	1.00	1.07	0.93	16.18	1.75	1.04	0.00	1.00
time (sec)	N/A	0.290	97.628	1.207	0.772	0.271	21.382	0.000	3.451

Problem 327	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	456	49	29	28	28
N.S.	1	1.00	1.07	0.93	16.29	1.75	1.04	1.00	1.00
time (sec)	N/A	0.295	5.621	1.125	0.757	0.264	48.470	0.313	3.161

Problem 328	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	354	310	338	599	0	0	0	0	0
N.S.	1	0.88	0.95	1.69	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.304	0.849	0.792	0.000	0.000	0.000	0.000	0.000

Problem 329	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	348	360	327	583	0	0	0	0	0
N.S.	1	1.03	0.94	1.68	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.047	0.751	0.963	0.000	0.000	0.000	0.000	0.000

Problem 330	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	179	232	445	0	0	0	0	0
N.S.	1	0.73	0.94	1.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.755	0.490	0.920	0.000	0.000	0.000	0.000	0.000

Problem 331	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	476	43	27	0	28
N.S.	1	1.00	1.07	0.93	17.00	1.54	0.96	0.00	1.00
time (sec)	N/A	1.674	18.830	1.638	0.848	0.259	20.421	0.000	3.266

Problem 332	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	484	49	29	28	28
N.S.	1	1.00	1.07	0.93	17.29	1.75	1.04	1.00	1.00
time (sec)	N/A	0.835	20.941	1.378	0.941	0.248	65.610	0.312	2.925

Problem 333	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	<b>F(-1)</b>	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	0	26	483	49	29	0	28
N.S.	1	1.00	0.00	0.93	17.25	1.75	1.04	0.00	1.00
time (sec)	N/A	0.302	0.000	1.139	0.921	0.257	176.193	0.000	3.388

Problem 334	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	<b>F(-1)</b>	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	0	26	469	49	0	28	28
N.S.	1	1.00	0.00	0.93	16.75	1.75	0.00	1.00	1.00
time (sec)	N/A	0.650	0.000	1.151	0.767	0.262	0.000	0.322	3.520

Problem 335	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	454	460	446	773	0	0	0	0	0
N.S.	1	1.01	0.98	1.70	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.735	1.333	0.913	0.000	0.000	0.000	0.000	0.000



Problem 336	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	448	464	436	759	0	0	0	0	0
N.S.	1	1.04	0.97	1.69	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.845	1.136	0.845	0.000	0.000	0.000	0.000	0.000

Problem 337	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	351	229	343	623	0	0	0	0	0
N.S.	1	0.65	0.98	1.77	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.919	0.828	0.954	0.000	0.000	0.000	0.000	0.000

Problem 338	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	<b>F(-2)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	524	60	0	0	28
N.S.	1	1.00	1.07	0.93	18.71	2.14	0.00	0.00	1.00
time (sec)	N/A	2.372	16.723	0.958	1.095	0.257	0.000	0.000	3.457

Problem 339	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	534	66	0	28	28
N.S.	1	1.00	1.07	0.93	19.07	2.36	0.00	1.00	1.00
time (sec)	N/A	1.066	23.018	0.851	1.161	0.251	0.000	0.328	3.326

Problem 340	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	534	66	0	0	28
N.S.	1	1.00	1.07	0.93	19.07	2.36	0.00	0.00	1.00
time (sec)	N/A	0.308	165.753	1.724	1.150	0.254	0.000	0.000	3.367

Problem 341	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	F(-1)	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	0	26	533	66	0	28	28
N.S.	1	1.00	0.00	0.93	19.04	2.36	0.00	1.00	1.00
time (sec)	N/A	0.308	0.000	1.434	1.153	0.259	0.000	0.334	3.219

Problem 342	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	337	215	190	519	0	0	0	0	0
N.S.	1	0.64	0.56	1.54	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.762	0.498	0.974	0.000	0.000	0.000	0.000	0.000

Problem 343	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	169	149	387	0	0	0	0	0
N.S.	1	0.72	0.63	1.64	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.693	0.366	0.868	0.000	0.000	0.000	0.000	0.000

Problem 344	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	165	144	375	0	0	0	0	0
N.S.	1	0.70	0.61	1.58	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.679	0.335	1.167	0.000	0.000	0.000	0.000	0.000

Problem 345	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	121	117	243	0	0	0	0	0
N.S.	1	0.89	0.86	1.79	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.814	0.283	0.767	0.000	0.000	0.000	0.000	0.000

Problem 346	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	130	115	107	225	0	0	0	0	0
N.S.	1	0.88	0.82	1.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.652	0.240	0.605	0.000	0.000	0.000	0.000	0.000

Problem 347	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	50	57	0	75	0	0	59
N.S.	1	1.00	1.35	1.54	0.00	2.03	0.00	0.00	1.59
time (sec)	N/A	0.226	0.026	0.520	0.000	0.258	0.000	0.000	3.349

Problem 348	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	481	80	29	0	28
N.S.	1	1.00	1.07	0.93	17.18	2.86	1.04	0.00	1.00
time (sec)	N/A	0.420	4.229	1.116	0.826	0.270	7.062	0.000	3.153

Problem 349	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	491	86	31	28	28
N.S.	1	1.00	1.07	0.93	17.54	3.07	1.11	1.00	1.00
time (sec)	N/A	0.416	1.419	0.988	0.831	0.252	21.614	0.317	3.429

Problem 350	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	531	106	29	0	28
N.S.	1	1.00	1.07	0.93	18.96	3.79	1.04	0.00	1.00
time (sec)	N/A	0.314	24.453	1.570	1.280	0.250	64.100	0.000	3.455

Problem 351	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	503	106	29	28	28
N.S.	1	1.00	1.07	0.93	17.96	3.79	1.04	1.00	1.00
time (sec)	N/A	0.707	5.619	0.536	0.951	0.271	71.187	0.339	3.263

Problem 352	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	28	24	511	104	27	0	26
N.S.	1	1.00	1.08	0.92	19.65	4.00	1.04	0.00	1.00
time (sec)	N/A	0.270	19.571	1.121	0.928	0.256	59.906	0.000	3.589

Problem 353	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	25	25	27	23	502	103	27	25	25
N.S.	1	1.00	1.08	0.92	20.08	4.12	1.08	1.00	1.00
time (sec)	N/A	0.613	3.877	1.096	0.928	0.270	47.153	0.333	3.268

Problem 354	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	530	108	29	0	28
N.S.	1	1.00	1.07	0.93	18.93	3.86	1.04	0.00	1.00
time (sec)	N/A	0.303	21.384	1.404	1.268	0.257	123.725	0.000	3.567

Problem 355	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	538	114	0	28	28
N.S.	1	1.00	1.07	0.93	19.21	4.07	0.00	1.00	1.00
time (sec)	N/A	0.309	17.592	1.297	1.292	0.258	0.000	0.352	3.570

Problem 356	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	614	144	0	28	28
N.S.	1	1.00	1.07	0.93	21.93	5.14	0.00	1.00	1.00
time (sec)	N/A	0.826	5.065	1.457	1.354	0.262	0.000	0.354	3.265

Problem 357	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	637	144	0	0	28
N.S.	1	1.00	1.07	0.93	22.75	5.14	0.00	0.00	1.00
time (sec)	N/A	0.314	39.633	1.362	1.959	0.261	0.000	0.000	3.267

Problem 358	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	635	144	0	28	28
N.S.	1	1.00	1.07	0.93	22.68	5.14	0.00	1.00	1.00
time (sec)	N/A	0.318	8.992	1.431	1.401	0.260	0.000	0.339	3.261

Problem 359	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	28	24	623	142	0	0	26
N.S.	1	1.00	1.08	0.92	23.96	5.46	0.00	0.00	1.00
time (sec)	N/A	0.273	35.325	1.327	1.685	0.261	0.000	0.000	3.282

Problem 360	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	25	25	27	23	609	141	0	25	25
N.S.	1	1.00	1.08	0.92	24.36	5.64	0.00	1.00	1.00
time (sec)	N/A	0.626	4.647	1.270	1.328	0.262	0.000	0.353	3.507

Problem 361	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	638	144	0	0	28
N.S.	1	1.00	1.07	0.93	22.79	5.14	0.00	0.00	1.00
time (sec)	N/A	0.297	30.699	1.463	1.956	0.264	0.000	0.000	3.277

Problem 362	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	647	150	0	28	28
N.S.	1	1.00	1.07	0.93	23.11	5.36	0.00	1.00	1.00
time (sec)	N/A	0.304	12.774	1.605	1.853	0.269	0.000	0.349	3.356

Problem 363	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	30	30	32	28	577	54	0	0	30
N.S.	1	1.00	1.07	0.93	19.23	1.80	0.00	0.00	1.00
time (sec)	N/A	0.311	1.607	2.604	1.950	0.261	0.000	0.000	3.176

Problem 364	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	30	30	32	28	514	44	31	0	30
N.S.	1	1.00	1.07	0.93	17.13	1.47	1.03	0.00	1.00
time (sec)	N/A	0.293	0.369	2.286	1.429	0.265	6.264	0.000	3.375

Problem 365	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	30	30	32	28	544	82	31	30	30
N.S.	1	1.00	1.07	0.93	18.13	2.73	1.03	1.00	1.00
time (sec)	N/A	0.441	1.139	1.357	0.714	0.278	24.707	0.325	3.711

Problem 366	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	30	30	32	28	596	108	0	30	30
N.S.	1	1.00	1.07	0.93	19.87	3.60	0.00	1.00	1.00
time (sec)	N/A	0.325	1.672	1.627	1.264	0.273	0.000	0.362	4.078

Problem 367	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	30	30	32	28	700	146	0	30	30
N.S.	1	1.00	1.07	0.93	23.33	4.87	0.00	1.00	1.00
time (sec)	N/A	0.319	2.065	2.462	1.284	0.267	0.000	0.379	3.892



Problem 368	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	45	51	0	56	0	0	48
N.S.	1	1.00	1.41	1.59	0.00	1.75	0.00	0.00	1.50
time (sec)	N/A	0.211	0.022	0.703	0.000	0.251	0.000	0.000	3.339

Problem 369	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	259	463	300	0	0	0	0	0	0
N.S.	1	1.79	1.16	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.992	1.882	0.000	0.000	0.000	0.000	0.000	0.000

Problem 370	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	340	525	384	0	0	0	0	0	0
N.S.	1	1.54	1.13	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.941	1.286	0.000	0.000	0.000	0.000	0.000	0.000

Problem 371	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	241	275	0	0	0	0	0	0	0
N.S.	1	1.14	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.933	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 372	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	233	231	246	0	0	0	0	0	0
N.S.	1	0.99	1.06	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.921	1.098	0.000	0.000	0.000	0.000	0.000	0.000

Problem 373	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	<b>F(-2)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	27	27	29	25	30	0	87	0	27
N.S.	1	1.00	1.07	0.93	1.11	0.00	3.22	0.00	1.00
time (sec)	N/A	3.297	1.274	1.232	1.057	0.000	8.802	0.000	3.922

Problem 374	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	479	571	527	0	0	0	0	0	0
N.S.	1	1.19	1.10	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.274	2.771	0.000	0.000	0.000	0.000	0.000	0.000

Problem 375	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	462	723	498	0	0	0	0	0	0
N.S.	1	1.56	1.08	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.397	2.479	0.000	0.000	0.000	0.000	0.000	0.000

Problem 376	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	363	567	0	0	0	0	0	0	0
N.S.	1	1.56	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.680	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 377	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	351	331	387	0	0	0	0	0	0
N.S.	1	0.94	1.10	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.121	1.590	0.000	0.000	0.000	0.000	0.000	0.000

Problem 378	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	<b>F(-2)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	30	0	133	0	29
N.S.	1	1.00	1.07	0.93	1.03	0.00	4.59	0.00	1.00
time (sec)	N/A	4.178	1.191	1.226	1.122	0.000	9.341	0.000	3.252

Problem 379	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	351	352	154	0	0	0	0	0	0
N.S.	1	1.00	0.44	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.847	0.234	0.000	0.000	0.000	0.000	0.000	0.000

Problem 380	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	F	F	F(-2)	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	205	180	117	0	0	0	0	0	0
N.S.	1	0.88	0.57	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.637	0.137	0.000	0.000	0.000	0.000	0.000	0.000

Problem 381	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	48	41	0	0	0	0	0
N.S.	1	1.00	1.00	0.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.229	0.027	0.580	0.000	0.000	0.000	0.000	0.000

Problem 382	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	26	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	1.08	0.92	0.92
time (sec)	N/A	0.368	5.540	1.451	0.557	0.000	4.852	1.957	2.789

Problem 383	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	26	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	1.08	0.92	0.92
time (sec)	N/A	0.960	6.064	1.335	0.507	0.000	98.580	1.951	2.754

Problem 384	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	511	420	198	0	0	0	0	0	0
N.S.	1	0.82	0.39	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.113	0.454	0.000	0.000	0.000	0.000	0.000	0.000

Problem 385	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	302	203	136	0	0	0	0	0	0
N.S.	1	0.67	0.45	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.452	0.354	0.000	0.000	0.000	0.000	0.000	0.000

Problem 386	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	48	41	0	0	0	0	0
N.S.	1	1.00	1.00	0.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.230	0.028	0.550	0.000	0.000	0.000	0.000	0.000

Problem 387	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	26	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	1.08	0.92	0.92
time (sec)	N/A	0.364	5.251	1.287	0.504	0.000	46.948	3.616	2.708

Problem 388	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	580	0	213	0	0	0	0	0	0
N.S.	1	0.00	0.37	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	0.483	0.000	0.000	0.000	0.000	0.000	0.000

Problem 389	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	330	256	148	0	0	0	0	0	0
N.S.	1	0.78	0.45	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.604	0.442	0.000	0.000	0.000	0.000	0.000	0.000

Problem 390	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	48	41	0	0	0	0	0
N.S.	1	1.00	1.00	0.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.221	0.028	0.576	0.000	0.000	0.000	0.000	0.000

Problem 391	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	0	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.00	0.92	0.92
time (sec)	N/A	0.356	4.785	1.200	0.527	0.000	0.000	3.565	2.706

Problem 392	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	368	369	165	0	0	0	0	0	0
N.S.	1	1.00	0.45	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.276	0.347	0.000	0.000	0.000	0.000	0.000	0.000

Problem 393	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	211	186	121	0	0	0	0	0	0
N.S.	1	0.88	0.57	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.912	0.219	0.000	0.000	0.000	0.000	0.000	0.000

Problem 394	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	50	50	44	0	0	0	0	0
N.S.	1	1.00	1.00	0.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.219	0.046	0.570	0.000	0.000	0.000	0.000	0.000

Problem 395	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	20	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.83	0.92	0.92
time (sec)	N/A	0.361	7.046	1.848	0.535	0.000	4.751	1.354	3.021

Problem 396	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	20	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.83	0.92	0.92
time (sec)	N/A	0.954	3.538	1.684	0.530	0.000	97.627	1.374	2.763

Problem 397	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	525	447	219	0	0	0	0	0	0
N.S.	1	0.85	0.42	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	5.055	0.595	0.000	0.000	0.000	0.000	0.000	0.000

Problem 398	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	316	217	144	0	0	0	0	0	0
N.S.	1	0.69	0.46	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.754	0.427	0.000	0.000	0.000	0.000	0.000	0.000

Problem 399	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	50	50	50	44	0	0	0	0	0
N.S.	1	1.00	1.00	0.88	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.214	0.044	0.556	0.000	0.000	0.000	0.000	0.000



Problem 400	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	20	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.83	0.92	0.92
time (sec)	N/A	0.357	7.197	1.434	0.516	0.000	48.711	4.402	2.868

Problem 401	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	65	50	72	0	0	0	0	0	0
N.S.	1	0.77	1.11	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.371	0.078	0.000	0.000	0.000	0.000	0.000	0.000

Problem 402	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	438	205	209	0	0	0	0	0	0
N.S.	1	0.47	0.48	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.541	0.373	0.000	0.000	0.000	0.000	0.000	0.000

Problem 403	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	294	148	153	0	0	0	0	0	0
N.S.	1	0.50	0.52	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.468	0.245	0.000	0.000	0.000	0.000	0.000	0.000

Problem 404	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	175	104	114	0	0	0	0	0	0
N.S.	1	0.59	0.65	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.395	0.136	0.000	0.000	0.000	0.000	0.000	0.000

Problem 405	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	41	0	0	0	0	0
N.S.	1	1.00	1.00	0.89	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.232	0.029	0.543	0.000	0.000	0.000	0.000	0.000

Problem 406	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	27	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	1.12	0.92	0.92
time (sec)	N/A	0.216	5.500	1.173	0.523	0.000	15.144	2.223	2.788

Problem 407	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	0	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.00	0.92	0.92
time (sec)	N/A	0.215	5.838	1.234	0.506	0.000	0.000	2.280	2.876

Problem 408	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	433	241	411	0	0	0	0	0	0
N.S.	1	0.56	0.95	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.837	0.934	0.000	0.000	0.000	0.000	0.000	0.000

Problem 409	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	286	183	239	0	0	0	0	0	0
N.S.	1	0.64	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.703	0.447	0.000	0.000	0.000	0.000	0.000	0.000

Problem 410	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	170	148	127	0	0	0	0	0	0
N.S.	1	0.87	0.75	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.646	0.285	0.000	0.000	0.000	0.000	0.000	0.000

Problem 411	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	46	41	0	59	0	0	0
N.S.	1	1.00	1.00	0.89	0.00	1.28	0.00	0.00	0.00
time (sec)	N/A	0.223	0.029	0.559	0.000	0.259	0.000	0.000	0.000

Problem 412	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	27	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	1.12	0.92	0.92
time (sec)	N/A	0.559	4.758	1.405	0.532	0.000	170.159	0.366	3.103

Problem 413	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	0	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.00	0.92	0.92
time (sec)	N/A	0.569	5.766	1.578	0.506	0.000	0.000	0.388	2.858

Problem 414	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	329	251	317	0	0	0	0	0	0
N.S.	1	0.76	0.96	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.197	0.454	0.000	0.000	0.000	0.000	0.000	0.000

Problem 415	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	201	182	141	0	0	0	0	0	0
N.S.	1	0.91	0.70	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.615	0.263	0.000	0.000	0.000	0.000	0.000	0.000

Problem 416	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	A	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	48	41	0	59	0	0	0
N.S.	1	1.00	1.00	0.85	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.226	0.031	0.576	0.000	0.250	0.000	0.000	0.000

Problem 417	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	0	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.00	0.92	0.92
time (sec)	N/A	0.548	4.746	1.523	0.496	0.000	0.000	0.380	2.762

Problem 418	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	24	24	26	20	22	0	0	22	22
N.S.	1	1.00	1.08	0.83	0.92	0.00	0.00	0.92	0.92
time (sec)	N/A	0.557	4.584	1.591	0.520	0.000	0.000	0.390	2.746

Problem 419	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	253	184	182	0	0	0	0	0	0
N.S.	1	0.73	0.72	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.582	0.753	0.000	0.000	0.000	0.000	0.000	0.000

Problem 420	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	379	279	241	0	0	0	0	0	0
N.S.	1	0.74	0.64	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.637	0.970	0.000	0.000	0.000	0.000	0.000	0.000

Problem 421	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	253	184	214	0	0	0	0	0	0
N.S.	1	0.73	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.511	0.530	0.000	0.000	0.000	0.000	0.000	0.000

Problem 422	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	29	29	0	29
N.S.	1	1.00	1.07	0.93	1.00	1.00	1.00	0.00	1.00
time (sec)	N/A	0.784	0.338	1.281	0.520	0.265	4.595	0.000	3.082

Problem 423	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	29	31	0	29
N.S.	1	1.00	1.07	0.93	1.00	1.00	1.07	0.00	1.00
time (sec)	N/A	0.572	0.319	1.411	0.532	0.278	6.431	0.000	3.232

Problem 424	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	658	444	438	0	0	0	0	0	0
N.S.	1	0.67	0.67	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.900	2.314	0.000	0.000	0.000	0.000	0.000	0.000

Problem 425	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	578	404	500	0	0	0	0	0	0
N.S.	1	0.70	0.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.782	1.648	0.000	0.000	0.000	0.000	0.000	0.000

Problem 426	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	450	308	384	0	0	0	0	0	0
N.S.	1	0.68	0.85	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.633	1.598	0.000	0.000	0.000	0.000	0.000	0.000

Problem 427	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	<b>F(-2)</b>	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	29	0	0	29
N.S.	1	1.00	1.07	0.93	1.00	1.00	0.00	0.00	1.00
time (sec)	N/A	1.351	0.391	2.466	0.622	0.268	0.000	0.000	3.161

Problem 428	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	29	0	0	29
N.S.	1	1.00	1.07	0.93	1.00	1.00	0.00	0.00	1.00
time (sec)	N/A	1.045	0.829	1.238	0.522	0.283	0.000	0.000	3.416

Problem 429	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	870	567	677	0	0	0	0	0	0
N.S.	1	0.65	0.78	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.052	5.624	0.000	0.000	0.000	0.000	0.000	0.000

Problem 430	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	793	532	633	0	0	0	0	0	0
N.S.	1	0.67	0.80	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.885	3.069	0.000	0.000	0.000	0.000	0.000	0.000

Problem 431	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	674	447	538	0	0	0	0	0	0
N.S.	1	0.66	0.80	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.773	3.947	0.000	0.000	0.000	0.000	0.000	0.000



Problem 432	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	54	0	0	29
N.S.	1	1.00	1.07	0.93	1.00	1.86	0.00	0.00	1.00
time (sec)	N/A	2.076	0.424	1.655	0.546	0.272	0.000	0.000	3.201

Problem 433	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	54	0	0	29
N.S.	1	1.00	1.07	0.93	1.00	1.86	0.00	0.00	1.00
time (sec)	N/A	1.600	0.618	1.402	0.519	0.279	0.000	0.000	3.487

Problem 434	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	323	265	292	0	0	0	0	0	0
N.S.	1	0.82	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.642	0.974	0.000	0.000	0.000	0.000	0.000	0.000

Problem 435	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	211	170	212	0	0	0	0	0	0
N.S.	1	0.81	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.534	0.608	0.000	0.000	0.000	0.000	0.000	0.000

Problem 436	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	154	138	154	0	0	0	0	0	0
N.S.	1	0.90	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.449	0.212	0.000	0.000	0.000	0.000	0.000	0.000

Problem 437	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	56	53	0	213	0	0	0
N.S.	1	1.00	1.30	1.23	0.00	4.95	0.00	0.00	0.00
time (sec)	N/A	0.232	0.033	0.614	0.000	0.278	0.000	0.000	0.000

Problem 438	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	39	27	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.39	0.96	1.00	1.00
time (sec)	N/A	0.292	2.094	1.431	0.534	0.274	6.481	12.444	3.260

Problem 439	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	30	26	28	41	29	28	28
N.S.	1	1.00	1.07	0.93	1.00	1.46	1.04	1.00	1.00
time (sec)	N/A	0.292	1.184	1.285	0.499	0.286	31.447	12.451	3.264

Problem 440	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	379	279	291	0	0	0	0	0	0
N.S.	1	0.74	0.77	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.661	0.899	0.000	0.000	0.000	0.000	0.000	0.000

Problem 441	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	253	184	213	0	0	0	0	0	0
N.S.	1	0.73	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.572	0.630	0.000	0.000	0.000	0.000	0.000	0.000

Problem 442	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	182	152	153	0	0	0	0	0	0
N.S.	1	0.84	0.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.479	0.237	0.000	0.000	0.000	0.000	0.000	0.000

Problem 443	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	57	54	0	221	0	0	0
N.S.	1	1.00	1.00	0.95	0.00	3.88	0.00	0.00	0.00
time (sec)	N/A	0.244	0.037	0.562	0.000	0.281	0.000	0.000	0.000

Problem 444	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	42	29	29	29
N.S.	1	1.00	1.07	0.93	1.00	1.45	1.00	1.00	1.00
time (sec)	N/A	0.313	0.479	1.318	0.512	0.275	6.587	12.453	3.246

Problem 445	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	44	31	29	29
N.S.	1	1.00	1.07	0.93	1.00	1.52	1.07	1.00	1.00
time (sec)	N/A	0.316	0.535	1.469	0.534	0.274	31.692	12.594	3.256

Problem 446	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	56	31	29	29
N.S.	1	1.00	1.07	0.93	1.00	1.93	1.07	1.00	1.00
time (sec)	N/A	0.325	0.937	1.451	0.579	0.265	67.557	12.736	3.316

Problem 447	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	27	27	29	25	27	54	29	27	27
N.S.	1	1.00	1.07	0.93	1.00	2.00	1.07	1.00	1.00
time (sec)	N/A	0.279	0.634	1.188	0.520	0.289	67.704	12.572	3.147

Problem 448	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	28	24	26	53	27	26	26
N.S.	1	1.00	1.08	0.92	1.00	2.04	1.04	1.00	1.00
time (sec)	N/A	0.231	0.126	1.385	0.546	0.281	49.660	12.618	3.190

Problem 449	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	55	0	29	29
N.S.	1	1.00	1.07	0.93	1.00	1.90	0.00	1.00	1.00
time (sec)	N/A	0.317	0.669	1.518	0.541	0.280	0.000	12.581	3.696

Problem 450	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	27	29	57	0	29	29
N.S.	1	1.00	1.07	0.93	1.00	1.97	0.00	1.00	1.00
time (sec)	N/A	0.322	0.687	1.550	0.528	0.270	0.000	12.517	3.381

Problem 451	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	30	30	32	28	30	42	31	30	30
N.S.	1	1.00	1.07	0.93	1.00	1.40	1.03	1.00	1.00
time (sec)	N/A	0.292	0.572	1.294	0.509	0.276	89.725	12.612	3.440

Problem 452	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	29	32	43	0	0	31
N.S.	1	1.00	1.07	1.00	1.10	1.48	0.00	0.00	1.07
time (sec)	N/A	0.265	0.623	0.523	1.091	0.279	0.000	0.000	3.095

Problem 453	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	27	27	29	27	32	31	0	0	29
N.S.	1	1.00	1.07	1.00	1.19	1.15	0.00	0.00	1.07
time (sec)	N/A	0.231	0.396	0.314	1.052	0.273	0.000	0.000	3.152

Problem 454	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	18	18	15	0	18
N.S.	1	1.00	1.12	1.00	1.12	1.12	0.94	0.00	1.12
time (sec)	N/A	0.193	0.033	0.296	0.527	0.272	31.178	0.000	2.936

Problem 455	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	29	34	33	27	33	31
N.S.	1	1.00	1.07	1.00	1.17	1.14	0.93	1.14	1.07
time (sec)	N/A	0.278	1.058	0.546	0.553	0.275	79.713	13.112	3.331

Problem 456	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	29	29	31	29	32	45	0	32	31
N.S.	1	1.00	1.07	1.00	1.10	1.55	0.00	1.10	1.07
time (sec)	N/A	0.277	1.280	0.661	0.573	0.270	0.000	12.328	3.420

Problem 457	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-1)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	44	0	0	31
N.S.	1	1.00	1.06	0.94	1.00	1.42	0.00	0.00	1.00
time (sec)	N/A	0.319	1.014	1.575	0.526	0.277	0.000	0.000	3.310

Problem 458	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	F(-1)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	31	0	0	31
N.S.	1	1.00	1.06	0.94	1.00	1.00	0.00	0.00	1.00
time (sec)	N/A	0.304	0.275	1.562	0.538	0.280	0.000	0.000	3.039

Problem 459	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	46	32	31	31
N.S.	1	1.00	1.06	0.94	1.00	1.48	1.03	1.00	1.00
time (sec)	N/A	0.307	0.685	1.602	0.536	0.269	92.138	13.104	3.497

Problem 460	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	31	31	33	29	31	58	0	31	31
N.S.	1	1.00	1.06	0.94	1.00	1.87	0.00	1.00	1.00
time (sec)	N/A	0.323	1.027	2.093	0.565	0.297	0.000	12.809	3.532

Problem 461	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	177	173	122	123	178	140	0	0	0
N.S.	1	0.98	0.69	0.69	1.01	0.79	0.00	0.00	0.00
time (sec)	N/A	0.358	0.086	0.580	0.214	0.258	0.000	0.000	0.000

Problem 462	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	161	157	140	215	196	136	0	0	0
N.S.	1	0.98	0.87	1.34	1.22	0.84	0.00	0.00	0.00
time (sec)	N/A	0.352	0.130	0.379	0.200	0.299	0.000	0.000	0.000

Problem 463	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	138	137	101	105	139	119	0	0	0
N.S.	1	0.99	0.73	0.76	1.01	0.86	0.00	0.00	0.00
time (sec)	N/A	0.331	0.073	0.313	0.182	0.254	0.000	0.000	0.000



Problem 464	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	122	121	120	225	156	114	0	0	0
N.S.	1	0.99	0.98	1.84	1.28	0.93	0.00	0.00	0.00
time (sec)	N/A	0.314	0.092	0.343	0.190	0.262	0.000	0.000	0.000

Problem 465	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	99	76	76	91	94	0	0	0
N.S.	1	1.05	0.81	0.81	0.97	1.00	0.00	0.00	0.00
time (sec)	N/A	0.282	0.057	0.029	0.200	0.269	0.000	0.000	0.000

Problem 466	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	264	274	129	128	0	0	0	0	0
N.S.	1	1.04	0.49	0.48	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.920	0.135	1.244	0.000	0.000	0.000	0.000	0.000

Problem 467	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	76	105	102	63	132	0	0	0
N.S.	1	1.01	1.40	1.36	0.84	1.76	0.00	0.00	0.00
time (sec)	N/A	0.294	0.084	0.046	0.302	0.274	0.000	0.000	0.000

Problem 468	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	251	258	100	135	0	0	0	0	0
N.S.	1	1.03	0.40	0.54	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.878	0.082	0.603	0.000	0.000	0.000	0.000	0.000

Problem 469	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	97	128	141	85	139	0	0	0
N.S.	1	1.03	1.36	1.50	0.90	1.48	0.00	0.00	0.00
time (sec)	N/A	0.317	0.177	0.044	0.294	0.300	0.000	0.000	0.000

Problem 470	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	319	267	192	214	305	231	0	0	0
N.S.	1	0.84	0.60	0.67	0.96	0.72	0.00	0.00	0.00
time (sec)	N/A	0.600	0.184	0.835	0.216	0.265	0.000	0.000	0.000

Problem 471	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	341	266	214	364	332	227	0	0	0
N.S.	1	0.78	0.63	1.07	0.97	0.67	0.00	0.00	0.00
time (sec)	N/A	0.557	0.225	0.763	0.215	0.269	0.000	0.000	0.000

Problem 472	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	260	226	163	182	247	198	0	0	0
N.S.	1	0.87	0.63	0.70	0.95	0.76	0.00	0.00	0.00
time (sec)	N/A	0.564	0.150	0.814	0.201	0.269	0.000	0.000	0.000

Problem 473	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	269	246	183	338	273	195	0	0	0
N.S.	1	0.91	0.68	1.26	1.01	0.72	0.00	0.00	0.00
time (sec)	N/A	0.449	0.190	0.749	0.233	0.259	0.000	0.000	0.000

Problem 474	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	180	130	138	180	163	0	0	0
N.S.	1	0.92	0.66	0.70	0.92	0.83	0.00	0.00	0.00
time (sec)	N/A	0.466	0.121	0.630	0.204	0.267	0.000	0.000	0.000

Problem 475	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	342	372	228	223	0	0	0	0	0
N.S.	1	1.09	0.67	0.65	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.085	0.385	0.990	0.000	0.000	0.000	0.000	0.000

Problem 476	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	144	128	178	134	236	0	0	0
N.S.	1	0.90	0.80	1.11	0.84	1.48	0.00	0.00	0.00
time (sec)	N/A	0.522	0.169	0.614	0.278	0.286	0.000	0.000	0.000

Problem 477	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	321	329	173	196	0	0	0	0	0
N.S.	1	1.02	0.54	0.61	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.104	0.301	1.161	0.000	0.000	0.000	0.000	0.000

Problem 478	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	184	168	133	191	126	216	0	0	0
N.S.	1	0.91	0.72	1.04	0.68	1.17	0.00	0.00	0.00
time (sec)	N/A	0.515	0.161	0.576	0.310	0.313	0.000	0.000	0.000

Problem 479	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	435	365	276	319	451	334	0	0	0
N.S.	1	0.84	0.63	0.73	1.04	0.77	0.00	0.00	0.00
time (sec)	N/A	0.961	0.276	0.740	0.216	0.260	0.000	0.000	0.000

Problem 480	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	494	442	294	538	487	330	0	0	0
N.S.	1	0.89	0.60	1.09	0.99	0.67	0.00	0.00	0.00
time (sec)	N/A	0.864	0.361	0.766	0.226	0.272	0.000	0.000	0.000

Problem 481	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	365	313	236	273	374	289	0	0	0
N.S.	1	0.86	0.65	0.75	1.02	0.79	0.00	0.00	0.00
time (sec)	N/A	0.879	0.233	0.758	0.215	0.263	0.000	0.000	0.000

Problem 482	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	358	328	256	476	409	286	0	0	0
N.S.	1	0.92	0.72	1.33	1.14	0.80	0.00	0.00	0.00
time (sec)	N/A	0.552	0.277	0.763	0.194	0.264	0.000	0.000	0.000

Problem 483	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	287	253	193	211	287	241	0	0	0
N.S.	1	0.88	0.67	0.74	1.00	0.84	0.00	0.00	0.00
time (sec)	N/A	0.720	0.162	0.677	0.194	0.262	0.000	0.000	0.000

Problem 484	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	509	508	344	348	0	0	0	0	0
N.S.	1	1.00	0.68	0.68	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.328	0.791	1.166	0.000	0.000	0.000	0.000	0.000

Problem 485	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	265	217	182	273	222	327	0	0	0
N.S.	1	0.82	0.69	1.03	0.84	1.23	0.00	0.00	0.00
time (sec)	N/A	0.794	0.219	0.673	0.276	0.303	0.000	0.000	0.000

Problem 486	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	476	480	278	293	0	0	0	0	0
N.S.	1	1.01	0.58	0.62	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.005	0.556	1.520	0.000	0.000	0.000	0.000	0.000

Problem 487	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	<b>No</b>	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	260	198	184	270	197	322	0	0	0
N.S.	1	0.76	0.71	1.04	0.76	1.24	0.00	0.00	0.00
time (sec)	N/A	0.851	0.237	0.681	0.293	0.330	0.000	0.000	0.000

Problem 488	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	395	343	265	302	415	333	0	0	0
N.S.	1	0.87	0.67	0.76	1.05	0.84	0.00	0.00	0.00
time (sec)	N/A	0.871	0.230	0.633	0.203	0.267	0.000	0.000	0.000

Problem 489	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	627	627	524	364	0	0	0	0	0
N.S.	1	1.00	0.84	0.58	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.448	0.859	45.913	0.000	0.000	0.000	0.000	0.000

Problem 490	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	521	521	512	2111	0	0	0	0	0
N.S.	1	1.00	0.98	4.05	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.280	0.424	3.168	0.000	0.000	0.000	0.000	0.000

Problem 491	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	544	544	457	284	0	0	0	0	0
N.S.	1	1.00	0.84	0.52	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.231	0.438	23.725	0.000	0.000	0.000	0.000	0.000

Problem 492	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	449	449	447	1987	0	0	0	0	0
N.S.	1	1.00	1.00	4.43	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.140	0.109	0.881	0.000	0.000	0.000	0.000	0.000

Problem 493	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	501	501	397	232	0	0	0	0	0
N.S.	1	1.00	0.79	0.46	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.160	0.252	1.279	0.000	0.000	0.000	0.000	0.000

Problem 494	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	489	489	410	381	0	0	0	0	0
N.S.	1	1.00	0.84	0.78	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.342	0.220	1.603	0.000	0.000	0.000	0.000	0.000

Problem 495	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	543	543	549	331	0	0	0	0	0
N.S.	1	1.00	1.01	0.61	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.319	0.985	26.268	0.000	0.000	0.000	0.000	0.000



Problem 496	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	550	550	518	466	0	0	0	0	0
N.S.	1	1.00	0.94	0.85	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.447	0.393	2.088	0.000	0.000	0.000	0.000	0.000

Problem 497	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	624	624	657	424	0	0	0	0	0
N.S.	1	1.00	1.05	0.68	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.387	1.200	32.835	0.000	0.000	0.000	0.000	0.000

Problem 498	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	562	562	694	2132	0	0	0	0	0
N.S.	1	1.00	1.23	3.79	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.389	1.376	3.007	0.000	0.000	0.000	0.000	0.000

Problem 499	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F(-2)</b>	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	113	123	399	0	537	0	0	0
N.S.	1	1.00	1.09	3.53	0.00	4.75	0.00	0.00	0.00
time (sec)	N/A	0.304	0.258	7.562	0.000	0.283	0.000	0.000	0.000

Problem 500	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	598	598	754	505	0	0	0	0	0
N.S.	1	1.00	1.26	0.84	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.521	1.157	1.493	0.000	0.000	0.000	0.000	0.000

Problem 501	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	634	634	792	635	0	0	0	0	0
N.S.	1	1.00	1.25	1.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.612	1.555	1.609	0.000	0.000	0.000	0.000	0.000

Problem 502	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	839	839	777	897	0	0	0	0	0
N.S.	1	1.00	0.93	1.07	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.634	1.319	31.119	0.000	0.000	0.000	0.000	0.000

Problem 503	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	792	792	719	813	0	0	0	0	0
N.S.	1	1.00	0.91	1.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.380	6.079	24.162	0.000	0.000	0.000	0.000	0.000

Problem 504	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	804	804	734	828	0	0	0	0	0
N.S.	1	1.00	0.91	1.03	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.521	6.391	20.609	0.000	0.000	0.000	0.000	0.000

Problem 505	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	846	846	821	900	0	0	0	0	0
N.S.	1	1.00	0.97	1.06	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.468	6.719	37.202	0.000	0.000	0.000	0.000	0.000

Problem 506	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	737	737	1097	3551	0	0	0	0	0
N.S.	1	1.00	1.49	4.82	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.653	6.634	5.517	0.000	0.000	0.000	0.000	0.000

Problem 507	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F(-1)</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	231	220	192	1169	0	1217	0	0	0
N.S.	1	0.95	0.83	5.06	0.00	5.27	0.00	0.00	0.00
time (sec)	N/A	0.590	0.667	0.551	0.000	0.364	0.000	0.000	0.000

Problem 508	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	177	166	183	1126	0	1233	0	0	0
N.S.	1	0.94	1.03	6.36	0.00	6.97	0.00	0.00	0.00
time (sec)	N/A	0.360	0.834	0.625	0.000	0.375	0.000	0.000	0.000

Problem 509	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	772	772	1204	1225	0	0	0	0	0
N.S.	1	1.00	1.56	1.59	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.715	6.075	2.445	0.000	0.000	0.000	0.000	0.000

Problem 510	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	834	834	1261	1455	0	0	0	0	0
N.S.	1	1.00	1.51	1.74	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.833	6.086	2.692	0.000	0.000	0.000	0.000	0.000

Problem 511	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	<b>F(-2)</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	<b>No</b>	TBD	TBD	TBD	TBD	TBD
size	1224	1224	1185	1752	0	0	0	0	0
N.S.	1	1.00	0.97	1.43	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	4.136	11.031	49.457	0.000	0.000	0.000	0.000	0.000

Problem 512	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F(-2)	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	1234	1234	1193	1222	0	0	0	0	0
N.S.	1	1.00	0.97	0.99	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	3.106	11.030	23.198	0.000	0.000	0.000	0.000	0.000

Problem 513	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F(-2)	F	F(-1)	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	1234	1234	1161	1778	0	0	0	0	0
N.S.	1	1.00	0.94	1.44	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.986	10.354	29.030	0.000	0.000	0.000	0.000	0.000

Problem 514	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	0	20	19	20	20
N.S.	1	1.00	1.10	0.90	0.00	1.00	0.95	1.00	1.00
time (sec)	N/A	0.191	6.366	1.063	0.000	0.256	1.717	0.332	3.232

Problem 515	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	0	20	19	20	20
N.S.	1	1.00	1.10	0.90	0.00	1.00	0.95	1.00	1.00
time (sec)	N/A	0.199	2.906	1.158	0.000	0.268	1.065	0.307	3.317

Problem 516	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	<b>F(-2)</b>	A	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	101	101	556	0	0	332	0	0	0
N.S.	1	1.00	5.50	0.00	0.00	3.29	0.00	0.00	0.00
time (sec)	N/A	0.402	13.469	0.000	0.000	0.286	0.000	0.000	0.000

Problem 517	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	B	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	182	176	633	0	0	724	0	0	0
N.S.	1	0.97	3.48	0.00	0.00	3.98	0.00	0.00	0.00
time (sec)	N/A	0.375	1.869	0.000	0.000	0.326	0.000	0.000	0.000

Problem 518	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	<b>F</b>	<b>F</b>	B	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	284	270	685	0	0	1360	0	0	0
N.S.	1	0.95	2.41	0.00	0.00	4.79	0.00	0.00	0.00
time (sec)	N/A	1.155	3.004	0.000	0.000	0.386	0.000	0.000	0.000

Problem 519	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	558	544	397	0	0	0	0	0	0
N.S.	1	0.97	0.71	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.474	0.867	0.000	0.000	0.000	0.000	0.000	0.000

Problem 520	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	353	350	293	0	0	0	0	0	0
N.S.	1	0.99	0.83	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.740	0.323	0.000	0.000	0.000	0.000	0.000	0.000

Problem 521	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	198	205	186	0	0	0	0	0	0
N.S.	1	1.04	0.94	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.422	0.427	0.000	0.000	0.000	0.000	0.000	0.000

Problem 522	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	23	23	25	23	25	25	20	25	25
N.S.	1	1.00	1.09	1.00	1.09	1.09	0.87	1.09	1.09
time (sec)	N/A	0.241	2.856	2.470	0.284	0.269	16.244	0.278	3.551

Problem 523	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	23	23	25	23	25	36	0	25	25
N.S.	1	1.00	1.09	1.00	1.09	1.57	0.00	1.09	1.09
time (sec)	N/A	0.235	2.293	1.375	0.318	0.270	0.000	0.280	3.194

Problem 524	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	23	23	25	23	25	47	0	25	25
N.S.	1	1.00	1.09	1.00	1.09	2.04	0.00	1.09	1.09
time (sec)	N/A	0.237	3.602	2.325	0.286	0.277	0.000	0.279	3.308

Problem 525	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	609	609	453	632	684	586	0	0	0
N.S.	1	1.00	0.74	1.04	1.12	0.96	0.00	0.00	0.00
time (sec)	N/A	2.395	0.370	0.809	0.217	0.271	0.000	0.000	0.000

Problem 526	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	359	359	299	402	429	380	0	0	0
N.S.	1	1.00	0.83	1.12	1.19	1.06	0.00	0.00	0.00
time (sec)	N/A	1.471	0.276	0.752	0.211	0.275	0.000	0.000	0.000

Problem 527	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	168	174	203	218	209	0	0	0
N.S.	1	1.00	1.04	1.21	1.30	1.24	0.00	0.00	0.00
time (sec)	N/A	0.792	0.156	0.347	0.205	0.269	0.000	0.000	0.000



Problem 528	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	B	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	55	84	78	72	96	0	111	0
N.S.	1	1.08	1.65	1.53	1.41	1.88	0.00	2.18	0.00
time (sec)	N/A	0.390	0.094	0.348	0.202	0.256	0.000	0.386	0.000

Problem 529	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	763	763	623	0	0	0	0	0	0
N.S.	1	1.00	0.82	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.762	0.434	0.000	0.000	0.000	0.000	0.000	0.000

Problem 530	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	0	34	20	22	22
N.S.	1	1.00	1.09	0.91	0.00	1.55	0.91	1.00	1.00
time (sec)	N/A	0.208	15.372	0.669	0.000	0.260	1.637	0.399	3.075

Problem 531	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	0	34	20	22	22
N.S.	1	1.00	1.09	0.91	0.00	1.55	0.91	1.00	1.00
time (sec)	N/A	0.206	8.229	0.783	0.000	0.263	1.097	0.332	3.106

Problem 532	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	0	54	20	22	22
N.S.	1	1.00	1.09	0.91	0.00	2.45	0.91	1.00	1.00
time (sec)	N/A	0.213	13.235	1.615	0.000	0.264	7.622	0.326	3.172

Problem 533	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	107	65	20	22	22
N.S.	1	1.00	1.09	0.91	4.86	2.95	0.91	1.00	1.00
time (sec)	N/A	0.210	26.794	1.655	0.719	0.271	167.634	0.335	3.356

Problem 534	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	388	388	254	380	0	0	0	0	0
N.S.	1	1.00	0.65	0.98	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.001	0.417	2.646	0.000	0.000	0.000	0.000	0.000

Problem 535	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	139	180	125	178	0	0	0	0	0
N.S.	1	1.29	0.90	1.28	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.560	0.174	0.944	0.000	0.000	0.000	0.000	0.000

Problem 536	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	57	46	56	0	0	0	0	0
N.S.	1	1.06	0.85	1.04	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.412	0.057	0.054	0.000	0.000	0.000	0.000	0.000

Problem 537	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	22	29	17	22	22
N.S.	1	1.00	1.10	1.00	1.10	1.45	0.85	1.10	1.10
time (sec)	N/A	0.211	0.567	0.825	0.270	0.258	9.359	0.281	2.866

Problem 538	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	22	53	0	22	22
N.S.	1	1.00	1.10	1.00	1.10	2.65	0.00	1.10	1.10
time (sec)	N/A	0.216	2.607	0.738	0.269	0.242	0.000	0.283	2.836

Problem 539	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	22	19	22	22
N.S.	1	1.00	1.09	0.91	1.00	1.00	0.86	1.00	1.00
time (sec)	N/A	0.221	0.982	1.175	0.263	0.246	0.470	0.313	2.792

Problem 540	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	39	20	22	22
N.S.	1	1.00	1.09	0.91	1.00	1.77	0.91	1.00	1.00
time (sec)	N/A	0.216	0.812	1.336	0.267	0.248	0.729	0.298	2.819

Problem 541	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	63	20	22	22
N.S.	1	1.00	1.09	0.91	1.00	2.86	0.91	1.00	1.00
time (sec)	N/A	0.223	1.422	1.342	0.287	0.298	3.052	0.301	2.964

Problem 542	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	87	20	22	22
N.S.	1	1.00	1.09	0.91	1.00	3.95	0.91	1.00	1.00
time (sec)	N/A	0.227	2.805	1.339	0.268	0.308	34.561	0.304	2.958

Problem 543	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	510	510	663	1102	0	0	0	0	0
N.S.	1	1.00	1.30	2.16	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.190	2.580	1.985	0.000	0.000	0.000	0.000	0.000

Problem 544	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	Yes	TBD	TBD	TBD	TBD	TBD
size	257	257	338	465	0	0	0	0	0
N.S.	1	1.00	1.32	1.81	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.802	1.001	1.117	0.000	0.000	0.000	0.000	0.000

Problem 545	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	86	80	125	0	0	0	0	0
N.S.	1	0.96	0.89	1.39	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.767	0.334	0.073	0.000	0.000	0.000	0.000	0.000

Problem 546	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	816	57	19	22	22
N.S.	1	1.00	1.10	1.00	40.80	2.85	0.95	1.10	1.10
time (sec)	N/A	0.210	17.947	0.895	1.318	0.251	89.470	0.314	2.964

Problem 547	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1078	98	0	22	22
N.S.	1	1.00	1.10	1.00	53.90	4.90	0.00	1.10	1.10
time (sec)	N/A	0.210	29.873	0.662	1.766	0.274	0.000	0.304	2.901

Problem 548	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	596	36	20	22	22
N.S.	1	1.00	1.09	0.91	27.09	1.64	0.91	1.00	1.00
time (sec)	N/A	0.223	10.789	1.214	0.808	0.251	0.872	0.302	3.170

Problem 549	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	580	67	22	22	22
N.S.	1	1.00	1.09	0.91	26.36	3.05	1.00	1.00	1.00
time (sec)	N/A	0.215	10.972	1.308	1.099	0.296	1.978	0.312	3.178

Problem 550	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	857	108	22	22	22
N.S.	1	1.00	1.09	0.91	38.95	4.91	1.00	1.00	1.00
time (sec)	N/A	0.227	16.339	1.521	1.988	0.262	15.244	0.327	3.268

Problem 551	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	1117	149	22	22	22
N.S.	1	1.00	1.09	0.91	50.77	6.77	1.00	1.00	1.00
time (sec)	N/A	0.222	24.975	1.393	2.985	0.271	170.436	0.334	3.409

Problem 552	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	672	672	536	0	0	0	0	0	0
N.S.	1	1.00	0.80	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.673	4.906	0.000	0.000	0.000	0.000	0.000	0.000

Problem 553	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	322	322	317	0	0	0	0	0	0
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.457	1.973	0.000	0.000	0.000	0.000	0.000	0.000

Problem 554	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	102	104	100	0	0	0	0	0	0
N.S.	1	1.02	0.98	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.756	0.142	0.000	0.000	0.000	0.000	0.000	0.000

Problem 555	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	<b>F(-1)</b>	N/A	<b>F(-2)</b>	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	0	20	0	0	19	22	22
N.S.	1	1.00	0.00	0.91	0.00	0.00	0.86	1.00	1.00
time (sec)	N/A	0.219	0.000	1.408	0.000	0.000	1.588	12.583	3.091

Problem 556	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	<b>F(-1)</b>	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	0	20	22	0	20	22	22
N.S.	1	1.00	0.00	0.91	1.00	0.00	0.91	1.00	1.00
time (sec)	N/A	0.226	0.000	0.926	0.544	0.000	41.816	12.374	3.183

Problem 557	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F(-2)</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	442	442	812	0	0	0	0	0	0
N.S.	1	1.00	1.84	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	2.016	1.994	0.000	0.000	0.000	0.000	0.000	0.000

Problem 558	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	140	151	269	0	0	0	0	0	0
N.S.	1	1.08	1.92	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.871	0.398	0.000	0.000	0.000	0.000	0.000	0.000

Problem 559	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	<b>F(-1)</b>	N/A	<b>F(-2)</b>	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	0	20	0	0	19	22	22
N.S.	1	1.00	0.00	0.91	0.00	0.00	0.86	1.00	1.00
time (sec)	N/A	0.225	0.000	1.372	0.000	0.000	37.428	19.146	3.354



Problem 560	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	<b>F(-1)</b>	N/A	N/A	<b>F(-2)</b>	<b>F(-1)</b>	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	0	20	22	0	0	22	22
N.S.	1	1.00	0.00	0.91	1.00	0.00	0.00	1.00	1.00
time (sec)	N/A	0.226	0.000	1.216	0.708	0.000	0.000	19.263	3.176

Problem 561	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	608	608	530	0	0	0	0	0	0
N.S.	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.373	0.815	0.000	0.000	0.000	0.000	0.000	0.000

Problem 562	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	287	287	213	0	0	0	0	0	0
N.S.	1	1.00	0.74	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.765	0.418	0.000	0.000	0.000	0.000	0.000	0.000

Problem 563	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	88	96	100	0	0	0	0	0	0
N.S.	1	1.09	1.14	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.416	0.072	0.000	0.000	0.000	0.000	0.000	0.000

Problem 564	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	0	20	22	22
N.S.	1	1.00	1.09	0.91	1.00	0.00	0.91	1.00	1.00
time (sec)	N/A	0.220	0.182	1.359	0.500	0.000	5.045	12.922	3.076

Problem 565	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	<b>F(-2)</b>	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	0	22	22	22
N.S.	1	1.00	1.09	0.91	1.00	0.00	1.00	1.00	1.00
time (sec)	N/A	0.222	0.227	1.068	0.533	0.000	151.692	12.693	3.113

Problem 566	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	<b>No</b>	N/A	TBD	TBD	TBD	TBD	TBD
size	358	358	268	0	0	0	0	0	0
N.S.	1	1.00	0.75	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.051	1.260	0.000	0.000	0.000	0.000	0.000	0.000

Problem 567	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-2)</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	120	129	0	0	0	0	0	0	0
N.S.	1	1.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.812	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 568	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	0	20	22	22
N.S.	1	1.00	1.09	0.91	1.00	0.00	0.91	1.00	1.00
time (sec)	N/A	0.227	0.196	1.406	0.509	0.000	36.247	0.419	3.985

Problem 569	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	F(-1)	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	24	20	22	0	0	22	22
N.S.	1	1.00	1.09	0.91	1.00	0.00	0.00	1.00	1.00
time (sec)	N/A	0.239	0.250	1.210	0.559	0.000	0.000	0.421	3.784

## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [246] had the largest ratio of [1.27272999999999992]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	8	1.08	23	0.348
2	A	7	7	1.09	23	0.304
3	A	6	6	1.05	23	0.261
4	A	4	4	1.03	21	0.190
5	A	4	4	1.03	20	0.200
6	C	12	11	1.20	23	0.478
7	A	7	6	0.97	23	0.261
8	C	13	12	1.04	23	0.522
9	A	6	5	1.03	23	0.217
10	A	7	6	0.97	25	0.240
11	A	11	10	1.16	25	0.400
12	A	7	6	1.02	25	0.240
13	A	5	5	0.98	23	0.217
14	A	7	6	1.08	22	0.273
15	C	17	16	1.31	25	0.640
16	A	8	7	1.07	25	0.280
17	C	18	17	1.24	25	0.680
18	A	12	11	1.15	25	0.440
19	A	7	6	0.94	25	0.240
20	A	10	10	0.95	25	0.400
21	A	7	6	0.97	25	0.240
22	A	6	6	0.97	23	0.261

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
23	A	7	6	1.02	22	0.273
24	C	22	21	1.51	25	0.840
25	A	7	6	0.98	25	0.240
26	C	23	22	1.39	25	0.880
27	A	10	9	0.96	25	0.360
28	C	14	13	1.18	25	0.520
29	C	13	12	1.04	25	0.480
30	C	10	9	1.00	25	0.360
31	C	9	8	1.01	23	0.348
32	C	7	6	0.98	22	0.273
33	C	8	7	1.07	25	0.280
34	C	11	10	1.04	25	0.400
35	C	11	10	0.99	25	0.400
36	C	16	15	1.13	25	0.600
37	C	14	13	1.12	25	0.520
38	C	15	14	1.01	25	0.560
39	C	10	9	0.98	25	0.360
40	A	2	2	1.00	23	0.087
41	C	10	9	0.98	22	0.409
42	C	11	10	1.04	25	0.400
43	C	15	14	1.15	25	0.560
44	C	15	14	1.36	25	0.560
45	C	22	21	1.26	25	0.840
46	C	15	14	1.00	25	0.560
47	A	7	7	0.98	25	0.280
48	C	12	11	0.97	25	0.440
49	A	3	3	0.98	23	0.130
50	C	12	11	0.97	22	0.500
51	C	14	13	1.17	25	0.520
52	C	19	18	1.17	25	0.720
53	C	19	18	1.25	25	0.720
54	C	28	27	1.31	25	1.080
55	C	7	6	0.98	18	0.333
56	C	10	9	0.98	18	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	C	12	11	0.96	18	0.611
58	A	7	7	0.83	27	0.259
59	A	5	5	0.88	27	0.185
60	A	3	3	1.00	24	0.125
61	A	3	3	1.00	27	0.111
62	A	5	5	0.74	27	0.185
63	A	4	4	0.67	27	0.148
64	A	4	4	0.64	27	0.148
65	A	3	3	0.64	27	0.111
66	A	3	3	0.68	27	0.111
67	A	4	4	0.74	25	0.160
68	A	8	7	0.70	27	0.259
69	A	8	7	0.70	27	0.259
70	A	10	9	0.68	27	0.333
71	A	12	12	0.90	27	0.444
72	A	10	10	0.96	27	0.370
73	A	8	8	1.08	24	0.333
74	A	8	8	1.06	27	0.296
75	A	8	8	1.03	27	0.296
76	A	6	5	0.59	27	0.185
77	A	6	5	0.58	27	0.185
78	A	6	5	0.58	27	0.185
79	A	6	5	0.56	27	0.185
80	A	4	4	0.57	27	0.148
81	A	4	4	0.58	27	0.148
82	A	4	4	0.59	27	0.148
83	A	4	4	0.59	25	0.160
84	A	12	11	0.79	27	0.407
85	A	13	12	0.77	27	0.444
86	A	13	12	0.81	27	0.444
87	A	18	17	0.95	27	0.630
88	A	16	15	1.01	27	0.556
89	A	11	11	1.04	24	0.458
90	A	14	13	1.11	27	0.481

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
91	A	14	13	1.06	27	0.481
92	A	14	13	1.07	27	0.481
93	A	7	6	0.51	27	0.222
94	A	7	6	0.55	27	0.222
95	A	6	5	0.53	27	0.185
96	A	4	4	0.53	27	0.148
97	A	4	4	0.52	27	0.148
98	A	4	4	0.51	27	0.148
99	A	5	5	0.50	25	0.200
100	A	16	15	0.86	27	0.556
101	A	16	15	0.83	27	0.556
102	A	17	16	0.84	27	0.593
103	A	3	3	1.00	14	0.214
104	A	6	6	1.05	27	0.222
105	A	5	5	1.04	27	0.185
106	A	4	4	1.03	27	0.148
107	A	3	3	1.00	27	0.111
108	A	2	2	1.00	25	0.080
109	A	1	1	1.00	24	0.042
110	A	6	5	0.57	27	0.185
111	A	2	2	1.00	27	0.074
112	A	8	7	0.71	27	0.259
113	A	4	4	0.99	27	0.148
114	A	4	4	0.70	27	0.148
115	A	10	9	1.08	27	0.333
116	A	4	4	0.75	27	0.148
117	A	5	5	1.00	27	0.185
118	A	4	4	1.00	25	0.160
119	A	2	2	1.00	24	0.083
120	A	10	9	0.69	27	0.333
121	A	7	6	0.77	27	0.222
122	A	15	14	0.77	27	0.519
123	A	6	5	0.71	27	0.185
124	A	6	6	0.72	27	0.222

Continued on next page

## 2.3. Detailed conclusion table specific for Rubi results

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
125	A	11	10	1.18	27	0.370
126	A	4	4	0.87	27	0.148
127	A	6	5	0.81	27	0.185
128	A	4	4	0.84	25	0.160
129	A	5	5	1.10	24	0.208
130	A	14	13	0.84	27	0.481
131	A	6	5	0.70	27	0.185
132	A	19	18	0.80	27	0.667
133	A	6	5	0.67	27	0.185
134	A	9	9	1.05	20	0.450
135	A	5	5	1.06	22	0.227
136	A	4	4	1.04	22	0.182
137	A	3	3	1.00	22	0.136
138	A	2	2	1.00	20	0.100
139	A	1	1	1.00	19	0.053
140	A	6	5	0.64	22	0.227
141	A	2	2	1.00	22	0.091
142	A	8	7	0.75	22	0.318
143	A	1	1	1.00	30	0.033
144	A	1	1	1.00	31	0.032
145	A	9	9	0.99	27	0.333
146	A	8	8	1.00	27	0.296
147	A	6	6	1.06	25	0.240
148	N/A	1	0	1.00	27	0.000
149	N/A	6	0	1.00	27	0.000
150	N/A	10	0	1.00	27	0.000
151	A	12	12	0.76	29	0.414
152	A	8	8	0.87	29	0.276
153	A	3	3	0.95	29	0.103
154	A	1	1	1.00	29	0.034
155	A	5	5	0.99	29	0.172
156	A	8	8	0.96	29	0.276
157	A	12	12	0.73	35	0.343

Continued on next page



Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
158	A	8	8	0.85	35	0.229
159	A	3	3	0.94	35	0.086
160	A	1	1	1.00	35	0.029
161	A	5	5	0.98	35	0.143
162	A	8	8	0.96	35	0.229
163	A	1	1	1.00	24	0.042
164	A	11	11	1.23	20	0.550
165	A	10	10	1.16	20	0.500
166	A	7	7	1.14	18	0.389
167	C	8	7	0.88	20	0.350
168	C	14	13	0.96	20	0.650
169	C	16	15	0.99	20	0.750
170	A	14	14	1.15	29	0.483
171	A	11	11	1.03	29	0.379
172	A	7	7	0.88	27	0.259
173	A	5	5	0.89	26	0.192
174	A	9	8	0.59	29	0.276
175	C	11	10	0.79	29	0.345
176	A	11	10	0.59	29	0.345
177	C	16	15	0.63	29	0.517
178	A	25	25	1.34	29	0.862
179	A	21	21	1.25	29	0.724
180	A	9	8	0.64	27	0.296
181	A	12	12	1.06	26	0.462
182	A	16	15	0.69	29	0.517
183	C	20	19	0.87	29	0.655
184	A	17	16	0.61	29	0.552
185	C	26	25	0.95	29	0.862
186	F	0	0	N/A	0.000	N/A
187	F	0	0	N/A	0.000	N/A
188	A	10	9	0.55	27	0.333
189	A	14	14	1.15	26	0.538
190	A	24	23	0.73	29	0.793

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
191	C	29	28	1.09	29	0.966
192	A	25	24	0.68	29	0.828
193	F	0	0	N/A	0.000	N/A
194	A	14	14	1.12	29	0.483
195	A	11	11	1.05	29	0.379
196	A	7	7	0.93	29	0.241
197	A	5	5	0.84	29	0.172
198	A	2	2	0.70	27	0.074
199	A	1	1	1.00	26	0.038
200	A	7	6	0.51	29	0.207
201	C	10	9	0.74	29	0.310
202	A	11	10	0.59	29	0.345
203	C	13	12	0.80	29	0.414
204	C	22	21	0.96	29	0.724
205	C	19	18	0.94	29	0.621
206	C	14	13	0.69	29	0.448
207	C	13	12	0.77	29	0.414
208	C	10	9	0.64	27	0.333
209	C	10	9	0.71	26	0.346
210	A	16	15	0.55	29	0.517
211	C	20	19	0.82	29	0.655
212	A	28	27	0.67	29	0.931
213	C	24	23	0.95	29	0.793
214	C	19	18	1.00	29	0.621
215	C	19	18	0.96	29	0.621
216	C	20	19	0.98	29	0.655
217	C	16	15	0.64	29	0.517
218	C	11	10	0.63	27	0.370
219	C	14	13	0.84	26	0.500
220	A	26	25	0.75	29	0.862
221	C	24	23	0.99	29	0.793
222	F	0	0	N/A	0.000	N/A
223	C	29	28	1.33	29	0.966

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
224	C	18	17	0.95	22	0.773
225	A	11	11	1.21	24	0.458
226	A	8	8	1.21	24	0.333
227	A	5	5	0.93	24	0.208
228	A	3	3	1.01	22	0.136
229	A	1	1	1.00	21	0.048
230	A	7	6	0.58	24	0.250
231	C	9	8	0.83	24	0.333
232	A	11	10	0.66	24	0.417
233	N/A	25	0	1.00	31	0.000
234	N/A	15	0	1.00	31	0.000
235	N/A	6	0	1.00	31	0.000
236	N/A	1	0	1.00	31	0.000
237	N/A	1	0	1.00	31	0.000
238	N/A	1	0	1.00	31	0.000
239	N/A	1	0	1.00	26	0.000
240	A	23	22	1.32	20	1.100
241	A	18	17	1.11	20	0.850
242	A	11	11	1.27	18	0.611
243	C	9	8	0.89	20	0.400
244	C	20	19	0.94	20	0.950
245	C	23	22	1.07	20	1.100
246	A	28	28	1.08	22	1.273
247	A	18	18	0.97	22	0.818
248	A	6	6	0.79	22	0.273
249	A	1	1	1.00	22	0.045
250	C	11	10	0.59	22	0.455
251	C	16	15	0.71	22	0.682
252	C	23	22	0.82	22	1.000
253	A	11	11	1.10	24	0.458
254	A	10	10	1.07	24	0.417
255	A	6	6	0.84	24	0.250
256	A	4	4	0.86	22	0.182

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
257	A	1	1	1.00	21	0.048
258	A	8	7	0.58	24	0.292
259	C	10	9	0.75	24	0.375
260	A	15	14	0.59	24	0.583
261	N/A	1	0	1.00	30	0.000
262	C	6	5	0.91	20	0.250
263	C	6	5	0.96	20	0.250
264	C	6	5	1.14	18	0.278
265	N/A	1	0	1.00	20	0.000
266	N/A	1	0	1.00	20	0.000
267	A	4	3	0.56	28	0.107
268	A	4	3	0.58	28	0.107
269	A	4	3	0.65	28	0.107
270	A	4	3	0.62	26	0.115
271	A	6	5	0.65	25	0.200
272	N/A	2	0	1.00	28	0.000
273	N/A	2	0	1.00	28	0.000
274	N/A	1	0	1.00	28	0.000
275	N/A	1	0	1.00	28	0.000
276	A	4	3	0.56	28	0.107
277	A	4	3	0.56	28	0.107
278	A	4	3	0.59	26	0.115
279	A	5	4	0.59	25	0.160
280	N/A	2	0	1.00	28	0.000
281	N/A	2	0	1.00	28	0.000
282	N/A	1	0	1.00	28	0.000
283	N/A	1	0	1.00	28	0.000
284	A	4	3	0.56	28	0.107
285	A	4	3	0.55	28	0.107
286	A	4	3	0.56	26	0.115
287	A	6	5	0.56	25	0.200
288	N/A	2	0	1.00	28	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
289	N/A	2	0	1.00	28	0.000
290	N/A	1	0	1.00	28	0.000
291	N/A	1	0	1.00	28	0.000
292	A	5	4	0.56	24	0.167
293	A	5	4	0.68	24	0.167
294	A	5	4	0.68	24	0.167
295	A	4	3	1.00	22	0.136
296	A	1	1	1.00	21	0.048
297	N/A	1	0	1.00	24	0.000
298	N/A	1	0	1.00	24	0.000
299	A	5	4	0.62	28	0.143
300	A	5	4	0.65	28	0.143
301	A	9	8	0.74	26	0.308
302	A	1	1	1.00	25	0.040
303	N/A	1	0	1.00	28	0.000
304	N/A	1	0	1.00	28	0.000
305	N/A	1	0	1.00	28	0.000
306	N/A	1	0	1.00	26	0.000
307	N/A	1	0	1.00	25	0.000
308	N/A	1	0	1.00	28	0.000
309	N/A	1	0	1.00	28	0.000
310	N/A	1	0	1.00	28	0.000
311	N/A	1	0	1.00	28	0.000
312	N/A	1	0	1.00	28	0.000
313	N/A	1	0	1.00	28	0.000
314	N/A	1	0	1.00	28	0.000
315	A	5	4	0.85	20	0.200
316	A	5	4	0.89	20	0.200
317	A	5	4	0.98	18	0.222
318	N/A	2	0	1.00	20	0.000
319	N/A	2	0	1.00	20	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
320	A	6	5	1.01	28	0.179
321	C	15	14	1.70	28	0.500
322	C	16	15	1.02	26	0.577
323	C	14	13	0.90	25	0.520
324	N/A	13	0	1.00	28	0.000
325	N/A	2	0	1.00	28	0.000
326	N/A	1	0	1.00	28	0.000
327	N/A	1	0	1.00	28	0.000
328	A	8	7	0.88	28	0.250
329	C	15	14	1.03	26	0.538
330	A	8	7	0.73	25	0.280
331	N/A	12	0	1.00	28	0.000
332	N/A	4	0	1.00	28	0.000
333	N/A	1	0	1.00	28	0.000
334	N/A	4	0	1.00	28	0.000
335	A	7	6	1.01	28	0.214
336	C	14	13	1.04	26	0.500
337	A	7	6	0.65	25	0.240
338	N/A	11	0	1.00	28	0.000
339	N/A	3	0	1.00	28	0.000
340	N/A	1	0	1.00	28	0.000
341	N/A	1	0	1.00	28	0.000
342	A	6	5	0.64	28	0.179
343	A	6	5	0.72	28	0.179
344	A	6	5	0.70	28	0.179
345	C	14	13	0.89	28	0.464
346	C	12	11	0.88	26	0.423
347	A	1	1	1.00	25	0.040
348	N/A	2	0	1.00	28	0.000
349	N/A	2	0	1.00	28	0.000
350	N/A	1	0	1.00	28	0.000
351	N/A	3	0	1.00	28	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
352	N/A	1	0	1.00	26	0.000
353	N/A	3	0	1.00	25	0.000
354	N/A	1	0	1.00	28	0.000
355	N/A	1	0	1.00	28	0.000
356	N/A	4	0	1.00	28	0.000
357	N/A	1	0	1.00	28	0.000
358	N/A	1	0	1.00	28	0.000
359	N/A	1	0	1.00	26	0.000
360	N/A	4	0	1.00	25	0.000
361	N/A	1	0	1.00	28	0.000
362	N/A	1	0	1.00	28	0.000
363	N/A	1	0	1.00	30	0.000
364	N/A	1	0	1.00	30	0.000
365	N/A	2	0	1.00	30	0.000
366	N/A	1	0	1.00	30	0.000
367	N/A	1	0	1.00	30	0.000
368	A	1	1	1.00	21	0.048
369	A	5	4	1.79	27	0.148
370	A	5	4	1.54	27	0.148
371	A	10	9	1.14	25	0.360
372	A	5	4	0.99	24	0.167
373	N/A	9	0	1.00	27	0.000
374	A	5	4	1.19	29	0.138
375	A	5	4	1.56	29	0.138
376	A	9	8	1.56	27	0.296
377	A	5	4	0.94	26	0.154
378	N/A	8	0	1.00	29	0.000
379	C	18	17	1.00	24	0.708
380	C	12	11	0.88	24	0.458
381	A	1	1	1.00	24	0.042
382	N/A	2	0	1.00	24	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
383	N/A	4	0	1.00	24	0.000
384	A	16	15	0.82	24	0.625
385	A	8	7	0.67	24	0.292
386	A	1	1	1.00	24	0.042
387	N/A	2	0	1.00	24	0.000
388	F	0	0	N/A	0.000	N/A
389	C	15	14	0.78	24	0.583
390	A	1	1	1.00	24	0.042
391	N/A	2	0	1.00	24	0.000
392	C	19	18	1.00	24	0.750
393	C	12	11	0.88	24	0.458
394	A	1	1	1.00	24	0.042
395	N/A	3	0	1.00	24	0.000
396	N/A	6	0	1.00	24	0.000
397	A	17	16	0.85	24	0.667
398	A	8	7	0.69	24	0.292
399	A	1	1	1.00	24	0.042
400	N/A	3	0	1.00	24	0.000
401	A	8	7	0.77	19	0.368
402	A	6	5	0.47	24	0.208
403	A	5	4	0.50	24	0.167
404	A	6	5	0.59	24	0.208
405	A	1	1	1.00	24	0.042
406	N/A	1	0	1.00	24	0.000
407	N/A	1	0	1.00	24	0.000
408	A	6	5	0.56	24	0.208
409	A	7	6	0.64	24	0.250
410	C	11	10	0.87	24	0.417
411	A	1	1	1.00	24	0.042
412	N/A	3	0	1.00	24	0.000
413	N/A	4	0	1.00	24	0.000
414	A	13	12	0.76	24	0.500

Continued on next page



Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
415	A	10	9	0.91	24	0.375
416	A	1	1	1.00	24	0.042
417	N/A	3	0	1.00	24	0.000
418	N/A	4	0	1.00	24	0.000
419	A	4	3	0.73	29	0.103
420	A	4	3	0.74	27	0.111
421	A	6	5	0.73	26	0.192
422	N/A	2	0	1.00	29	0.000
423	N/A	2	0	1.00	29	0.000
424	A	4	3	0.67	29	0.103
425	A	4	3	0.70	27	0.111
426	A	5	4	0.68	26	0.154
427	N/A	2	0	1.00	29	0.000
428	N/A	2	0	1.00	29	0.000
429	A	4	3	0.65	29	0.103
430	A	4	3	0.67	27	0.111
431	A	6	5	0.66	26	0.192
432	N/A	2	0	1.00	29	0.000
433	N/A	2	0	1.00	29	0.000
434	A	5	4	0.82	28	0.143
435	A	5	4	0.81	28	0.143
436	A	6	5	0.90	26	0.192
437	A	1	1	1.00	25	0.040
438	N/A	1	0	1.00	28	0.000
439	N/A	1	0	1.00	28	0.000
440	A	5	4	0.74	29	0.138
441	A	5	4	0.73	29	0.138
442	A	6	5	0.84	27	0.185
443	A	1	1	1.00	26	0.038
444	N/A	1	0	1.00	29	0.000
445	N/A	1	0	1.00	29	0.000
446	N/A	1	0	1.00	29	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
447	N/A	1	0	1.00	27	0.000
448	N/A	1	0	1.00	26	0.000
449	N/A	1	0	1.00	29	0.000
450	N/A	1	0	1.00	29	0.000
451	N/A	1	0	1.00	30	0.000
452	N/A	1	0	1.00	29	0.000
453	N/A	1	0	1.00	27	0.000
454	N/A	1	0	1.00	16	0.000
455	N/A	1	0	1.00	29	0.000
456	N/A	1	0	1.00	29	0.000
457	N/A	1	0	1.00	31	0.000
458	N/A	1	0	1.00	31	0.000
459	N/A	1	0	1.00	31	0.000
460	N/A	1	0	1.00	31	0.000
461	A	7	7	0.98	19	0.368
462	A	7	7	0.98	19	0.368
463	A	5	5	0.99	19	0.263
464	A	5	5	0.99	17	0.294
465	A	4	4	1.05	16	0.250
466	A	4	4	1.04	19	0.211
467	A	5	4	1.01	19	0.211
468	A	4	4	1.03	19	0.211
469	A	6	5	1.03	19	0.263
470	A	7	6	0.84	21	0.286
471	A	10	9	0.78	21	0.429
472	A	7	6	0.87	21	0.286
473	A	8	7	0.91	19	0.368
474	A	7	6	0.92	18	0.333
475	A	4	4	1.09	21	0.190
476	A	8	7	0.90	21	0.333
477	A	4	4	1.02	21	0.190
478	A	10	9	0.91	21	0.429

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
479	A	7	6	0.84	21	0.286
480	A	12	11	0.89	21	0.524
481	A	7	6	0.86	21	0.286
482	A	9	8	0.92	19	0.421
483	A	7	6	0.88	18	0.333
484	A	4	4	1.00	21	0.190
485	A	7	6	0.82	21	0.286
486	A	4	4	1.01	21	0.190
487	A	10	9	0.76	21	0.429
488	A	7	6	0.87	18	0.333
489	A	2	2	1.00	21	0.095
490	A	2	2	1.00	21	0.095
491	A	2	2	1.00	21	0.095
492	A	2	2	1.00	19	0.105
493	A	2	2	1.00	18	0.111
494	A	2	2	1.00	21	0.095
495	A	2	2	1.00	21	0.095
496	A	2	2	1.00	21	0.095
497	A	2	2	1.00	21	0.095
498	A	2	2	1.00	21	0.095
499	A	5	4	1.00	19	0.211
500	A	2	2	1.00	21	0.095
501	A	2	2	1.00	21	0.095
502	A	2	2	1.00	21	0.095
503	A	2	2	1.00	21	0.095
504	A	2	2	1.00	18	0.111
505	A	2	2	1.00	21	0.095
506	A	2	2	1.00	21	0.095
507	A	11	10	0.95	21	0.476
508	A	6	5	0.94	19	0.263
509	A	2	2	1.00	21	0.095
510	A	2	2	1.00	21	0.095
511	A	2	2	1.00	21	0.095
512	A	2	2	1.00	21	0.095

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
513	A	2	2	1.00	18	0.111
514	N/A	1	0	1.00	20	0.000
515	N/A	1	0	1.00	20	0.000
516	A	7	6	1.00	20	0.300
517	A	8	7	0.97	20	0.350
518	A	10	9	0.95	20	0.450
519	A	8	8	0.97	23	0.348
520	A	7	7	0.99	23	0.304
521	A	5	5	1.04	21	0.238
522	N/A	1	0	1.00	23	0.000
523	N/A	1	0	1.00	23	0.000
524	N/A	1	0	1.00	23	0.000
525	A	2	2	1.00	20	0.100
526	A	2	2	1.00	20	0.100
527	A	2	2	1.00	18	0.111
528	A	3	3	1.08	10	0.300
529	A	2	2	1.00	20	0.100
530	N/A	1	0	1.00	22	0.000
531	N/A	1	0	1.00	22	0.000
532	N/A	1	0	1.00	22	0.000
533	N/A	1	0	1.00	22	0.000
534	A	2	2	1.00	20	0.100
535	A	2	2	1.29	18	0.111
536	C	11	10	1.06	10	1.000
537	N/A	1	0	1.00	20	0.000
538	N/A	1	0	1.00	20	0.000
539	N/A	1	0	1.00	22	0.000
540	N/A	1	0	1.00	22	0.000
541	N/A	1	0	1.00	22	0.000
542	N/A	1	0	1.00	22	0.000
543	A	2	2	1.00	20	0.100
544	A	2	2	1.00	18	0.111

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
545	A	10	9	0.96	10	0.900
546	N/A	1	0	1.00	20	0.000
547	N/A	1	0	1.00	20	0.000
548	N/A	1	0	1.00	22	0.000
549	N/A	1	0	1.00	22	0.000
550	N/A	1	0	1.00	22	0.000
551	N/A	1	0	1.00	22	0.000
552	A	2	2	1.00	22	0.091
553	A	2	2	1.00	20	0.100
554	A	9	8	1.02	12	0.667
555	N/A	1	0	1.00	22	0.000
556	N/A	1	0	1.00	22	0.000
557	A	2	2	1.00	20	0.100
558	C	11	10	1.08	12	0.833
559	N/A	1	0	1.00	22	0.000
560	N/A	1	0	1.00	22	0.000
561	A	2	2	1.00	22	0.091
562	A	2	2	1.00	20	0.100
563	C	9	8	1.09	12	0.667
564	N/A	1	0	1.00	22	0.000
565	N/A	1	0	1.00	22	0.000
566	A	2	2	1.00	20	0.100
567	A	9	8	1.08	12	0.667
568	N/A	1	0	1.00	22	0.000
569	N/A	1	0	1.00	22	0.000

# CHAPTER 3

## LISTING OF INTEGRALS

3.1	$\int x^4(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$ . . . . .	208
3.2	$\int x^3(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$ . . . . .	215
3.3	$\int x^2(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$ . . . . .	222
3.4	$\int x(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$ . . . . .	229
3.5	$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$ . . . . .	235
3.6	$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x} dx$ . . . . .	240
3.7	$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^2} dx$ . . . . .	248
3.8	$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx$ . . . . .	254
3.9	$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^4} dx$ . . . . .	262
3.10	$\int x^4(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	268
3.11	$\int x^3(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	275
3.12	$\int x^2(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	283
3.13	$\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	290
3.14	$\int (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	296
3.15	$\int \frac{(d - c^2 dx^2)^2(a + \operatorname{barccosh}(cx))}{x} dx$ . . . . .	302
3.16	$\int \frac{(d - c^2 dx^2)^2(a + \operatorname{barccosh}(cx))}{x^2} dx$ . . . . .	311
3.17	$\int \frac{(d - c^2 dx^2)^2(a + \operatorname{barccosh}(cx))}{x^3} dx$ . . . . .	318
3.18	$\int \frac{(d - c^2 dx^2)^2(a + \operatorname{barccosh}(cx))}{x^4} dx$ . . . . .	328
3.19	$\int x^4(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	336
3.20	$\int x^3(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	343
3.21	$\int x^2(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	351
3.22	$\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	358
3.23	$\int (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$ . . . . .	365
3.24	$\int \frac{(d - c^2 dx^2)^3(a + \operatorname{barccosh}(cx))}{x} dx$ . . . . .	372
3.25	$\int \frac{(d - c^2 dx^2)^3(a + \operatorname{barccosh}(cx))}{x^2} dx$ . . . . .	383
3.26	$\int \frac{(d - c^2 dx^2)^3(a + \operatorname{barccosh}(cx))}{x^3} dx$ . . . . .	390

3.27	$\int \frac{(d-c^2 dx^2)^3 (a+b \operatorname{arccosh}(cx))}{x^4} dx$	402
3.28	$\int \frac{x^4 (a+b \operatorname{arccosh}(cx))}{d-c^2 dx^2} dx$	410
3.29	$\int \frac{x^3 (a+b \operatorname{arccosh}(cx))}{d-c^2 dx^2} dx$	418
3.30	$\int \frac{x^2 (a+b \operatorname{arccosh}(cx))}{d-c^2 dx^2} dx$	426
3.31	$\int \frac{x (a+b \operatorname{arccosh}(cx))}{d-c^2 dx^2} dx$	433
3.32	$\int \frac{a+b \operatorname{arccosh}(cx)}{d-c^2 dx^2} dx$	439
3.33	$\int \frac{a+b \operatorname{arccosh}(cx)}{x(d-c^2 dx^2)} dx$	445
3.34	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^2(d-c^2 dx^2)} dx$	451
3.35	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^3(d-c^2 dx^2)} dx$	458
3.36	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^4(d-c^2 dx^2)} dx$	465
3.37	$\int \frac{x^4 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^2} dx$	473
3.38	$\int \frac{x^3 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^2} dx$	482
3.39	$\int \frac{x^2 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^2} dx$	491
3.40	$\int \frac{x (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^2} dx$	498
3.41	$\int \frac{a+b \operatorname{arccosh}(cx)}{(d-c^2 dx^2)^2} dx$	503
3.42	$\int \frac{a+b \operatorname{arccosh}(cx)}{x(d-c^2 dx^2)^2} dx$	510
3.43	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^2(d-c^2 dx^2)^2} dx$	517
3.44	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^3(d-c^2 dx^2)^2} dx$	526
3.45	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^4(d-c^2 dx^2)^2} dx$	535
3.46	$\int \frac{x^4 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^3} dx$	546
3.47	$\int \frac{x^3 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^3} dx$	556
3.48	$\int \frac{x^2 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^3} dx$	562
3.49	$\int \frac{x (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^3} dx$	571
3.50	$\int \frac{a+b \operatorname{arccosh}(cx)}{(d-c^2 dx^2)^3} dx$	576
3.51	$\int \frac{a+b \operatorname{arccosh}(cx)}{x(d-c^2 dx^2)^3} dx$	584
3.52	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^2(d-c^2 dx^2)^3} dx$	592
3.53	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^3(d-c^2 dx^2)^3} dx$	602
3.54	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^4(d-c^2 dx^2)^3} dx$	612
3.55	$\int \frac{\operatorname{arccosh}(ax)}{c-a^2 cx^2} dx$	626
3.56	$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2 cx^2)^2} dx$	631

3.57	$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^3} dx$	637
3.58	$\int x^4 \sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx)) dx$	645
3.59	$\int x^2 \sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx)) dx$	653
3.60	$\int \sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx)) dx$	659
3.61	$\int \frac{\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))}{x^2} dx$	664
3.62	$\int \frac{\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))}{x^4} dx$	669
3.63	$\int \frac{\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))}{x^6} dx$	675
3.64	$\int \frac{\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))}{x^8} dx$	682
3.65	$\int x^5 \sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx)) dx$	689
3.66	$\int x^3 \sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx)) dx$	695
3.67	$\int x \sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx)) dx$	701
3.68	$\int \frac{\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))}{x} dx$	706
3.69	$\int \frac{\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))}{x^3} dx$	713
3.70	$\int \frac{\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))}{x^5} dx$	720
3.71	$\int x^4 (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	728
3.72	$\int x^2 (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	738
3.73	$\int (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	747
3.74	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx$	754
3.75	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx$	761
3.76	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^6} dx$	768
3.77	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx$	775
3.78	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx$	782
3.79	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx$	789
3.80	$\int x^7 (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	796
3.81	$\int x^5 (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	803
3.82	$\int x^3 (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	810
3.83	$\int x (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	816
3.84	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx$	822
3.85	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx$	830
3.86	$\int \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx$	839
3.87	$\int x^4 (d-c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$	848
3.88	$\int x^2 (d-c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$	859
3.89	$\int (d-c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$	869
3.90	$\int \frac{(d-c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx$	878
3.91	$\int \frac{(d-c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^4} dx$	887



3.92	$\int \frac{(d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx))}{x^6} dx$	896
3.93	$\int \frac{(d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx))}{x^8} dx$	905
3.94	$\int \frac{(d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx))}{x^{10}} dx$	912
3.95	$\int \frac{(d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx))}{x^{12}} dx$	920
3.96	$\int x^7 (d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx)) dx$	927
3.97	$\int x^5 (d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx)) dx$	934
3.98	$\int x^3 (d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx)) dx$	941
3.99	$\int x (d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx)) dx$	948
3.100	$\int \frac{(d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx))}{x} dx$	954
3.101	$\int \frac{(d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx))}{x^3} dx$	963
3.102	$\int \frac{(d-c^2 dx^2)^{5/2} (a+b \operatorname{arccosh}(cx))}{x^5} dx$	973
3.103	$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx$	983
3.104	$\int \frac{x^5 (a+b \operatorname{arccosh}(cx))}{\sqrt{d-c^2 dx^2}} dx$	988
3.105	$\int \frac{x^4 (a+b \operatorname{arccosh}(cx))}{\sqrt{d-c^2 dx^2}} dx$	995
3.106	$\int \frac{x^3 (a+b \operatorname{arccosh}(cx))}{\sqrt{d-c^2 dx^2}} dx$	1001
3.107	$\int \frac{x^2 (a+b \operatorname{arccosh}(cx))}{\sqrt{d-c^2 dx^2}} dx$	1007
3.108	$\int \frac{x (a+b \operatorname{arccosh}(cx))}{\sqrt{d-c^2 dx^2}} dx$	1012
3.109	$\int \frac{a+b \operatorname{arccosh}(cx)}{\sqrt{d-c^2 dx^2}} dx$	1017
3.110	$\int \frac{a+b \operatorname{arccosh}(cx)}{x \sqrt{d-c^2 dx^2}} dx$	1021
3.111	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^2 \sqrt{d-c^2 dx^2}} dx$	1027
3.112	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^3 \sqrt{d-c^2 dx^2}} dx$	1032
3.113	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^4 \sqrt{d-c^2 dx^2}} dx$	1039
3.114	$\int \frac{x^5 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^{3/2}} dx$	1045
3.115	$\int \frac{x^4 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^{3/2}} dx$	1051
3.116	$\int \frac{x^3 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^{3/2}} dx$	1058
3.117	$\int \frac{x^2 (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^{3/2}} dx$	1064
3.118	$\int \frac{x (a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^{3/2}} dx$	1070
3.119	$\int \frac{a+b \operatorname{arccosh}(cx)}{(d-c^2 dx^2)^{3/2}} dx$	1075
3.120	$\int \frac{a+b \operatorname{arccosh}(cx)}{x (d-c^2 dx^2)^{3/2}} dx$	1079
3.121	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^2 (d-c^2 dx^2)^{3/2}} dx$	1086
3.122	$\int \frac{a+b \operatorname{arccosh}(cx)}{x^3 (d-c^2 dx^2)^{3/2}} dx$	1092

3.123	$\int \frac{a+\operatorname{barccosh}(cx)}{x^4(d-c^2dx^2)^{3/2}} dx$	1101
3.124	$\int \frac{x^5(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$	1107
3.125	$\int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$	1114
3.126	$\int \frac{x^3(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$	1121
3.127	$\int \frac{x^2(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$	1127
3.128	$\int \frac{x(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$	1133
3.129	$\int \frac{a+\operatorname{barccosh}(cx)}{(d-c^2dx^2)^{5/2}} dx$	1138
3.130	$\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^{5/2}} dx$	1144
3.131	$\int \frac{a+\operatorname{barccosh}(cx)}{x^2(d-c^2dx^2)^{5/2}} dx$	1153
3.132	$\int \frac{a+\operatorname{barccosh}(cx)}{x^3(d-c^2dx^2)^{5/2}} dx$	1159
3.133	$\int \frac{a+\operatorname{barccosh}(cx)}{x^4(d-c^2dx^2)^{5/2}} dx$	1170
3.134	$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{7/2}} dx$	1176
3.135	$\int \frac{x^4\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$	1183
3.136	$\int \frac{x^3\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$	1189
3.137	$\int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$	1194
3.138	$\int \frac{x\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$	1199
3.139	$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$	1203
3.140	$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx$	1207
3.141	$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx$	1212
3.142	$\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx$	1217
3.143	$\int \frac{(fx)^{3/2}(a+\operatorname{barccosh}(cx))}{\sqrt{1-c^2x^2}} dx$	1223
3.144	$\int \frac{(fx)^{3/2}(a+\operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$	1227
3.145	$\int (fx)^m (d-c^2dx^2)^3 (a+\operatorname{barccosh}(cx)) dx$	1231
3.146	$\int (fx)^m (d-c^2dx^2)^2 (a+\operatorname{barccosh}(cx)) dx$	1240
3.147	$\int (fx)^m (d-c^2dx^2) (a+\operatorname{barccosh}(cx)) dx$	1248
3.148	$\int \frac{(fx)^m (a+\operatorname{barccosh}(cx))}{d-c^2dx^2} dx$	1254
3.149	$\int \frac{(fx)^m (a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^2} dx$	1258
3.150	$\int \frac{(fx)^m (a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^3} dx$	1264
3.151	$\int (fx)^m (d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx)) dx$	1271
3.152	$\int (fx)^m (d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx)) dx$	1280
3.153	$\int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx)) dx$	1287

3.154	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$	1293
3.155	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$	1298
3.156	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$	1304
3.157	$\int (fx)^m (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + \operatorname{barccosh}(cx)) dx$	1312
3.158	$\int (fx)^m (d1 + cd1x)^{3/2} (d2 - cd2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx$	1322
3.159	$\int (fx)^m \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx)) dx$	1330
3.160	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d1 + cd1x} \sqrt{d2 - cd2x}} dx$	1336
3.161	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{3/2} (d2 - cd2x)^{3/2}} dx$	1341
3.162	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2}} dx$	1347
3.163	$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1 - a^2 x^2}} dx$	1355
3.164	$\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^2 dx$	1359
3.165	$\int (c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^2 dx$	1368
3.166	$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^2 dx$	1376
3.167	$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2 cx^2} dx$	1382
3.168	$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2 cx^2)^2} dx$	1388
3.169	$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2 cx^2)^3} dx$	1397
3.170	$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$	1407
3.171	$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$	1418
3.172	$\int x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$	1427
3.173	$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$	1434
3.174	$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx$	1441
3.175	$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx$	1449
3.176	$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$	1458
3.177	$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$	1466
3.178	$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$	1476
3.179	$\int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$	1492
3.180	$\int x (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$	1506
3.181	$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$	1514
3.182	$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$	1524
3.183	$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx$	1535
3.184	$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$	1547
3.185	$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$	1558
3.186	$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$	1571
3.187	$\int x^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$	1591

3.188	$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$	1613
3.189	$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$	1621
3.190	$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$	1633
3.191	$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx$	1648
3.192	$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$	1665
3.193	$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$	1681
3.194	$\int \frac{x^5 (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$	1697
3.195	$\int \frac{x^4 (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$	1708
3.196	$\int \frac{x^3 (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$	1718
3.197	$\int \frac{x^2 (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$	1726
3.198	$\int \frac{x (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$	1733
3.199	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$	1738
3.200	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x \sqrt{d - c^2 dx^2}} dx$	1742
3.201	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx$	1748
3.202	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx$	1755
3.203	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx$	1763
3.204	$\int \frac{x^5 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$	1773
3.205	$\int \frac{x^4 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$	1788
3.206	$\int \frac{x^3 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$	1802
3.207	$\int \frac{x^2 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$	1812
3.208	$\int \frac{x (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$	1820
3.209	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$	1827
3.210	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{3/2}} dx$	1834
3.211	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx$	1844
3.212	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx$	1855
3.213	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx$	1868
3.214	$\int \frac{x^5 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$	1882
3.215	$\int \frac{x^4 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$	1897
3.216	$\int \frac{x^3 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$	1911
3.217	$\int \frac{x^2 (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$	1922

3.218	$\int \frac{x(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$	1933
3.219	$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$	1941
3.220	$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$	1951
3.221	$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$	1965
3.222	$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$	1981
3.223	$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$	1997
3.224	$\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{7/2}} dx$	2015
3.225	$\int \frac{x^4\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$	2028
3.226	$\int \frac{x^3\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$	2036
3.227	$\int \frac{x^2\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$	2043
3.228	$\int \frac{x\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$	2049
3.229	$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$	2054
3.230	$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx$	2058
3.231	$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx$	2064
3.232	$\int \frac{\operatorname{arccosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx$	2070
3.233	$\int (fx)^m (d-c^2dx^2)^{5/2} (a+b\operatorname{arccosh}(cx))^2 dx$	2078
3.234	$\int (fx)^m (d-c^2dx^2)^{3/2} (a+b\operatorname{arccosh}(cx))^2 dx$	2096
3.235	$\int (fx)^m \sqrt{d-c^2dx^2} (a+b\operatorname{arccosh}(cx))^2 dx$	2106
3.236	$\int \frac{(fx)^m (a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$	2112
3.237	$\int \frac{(fx)^m (a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$	2116
3.238	$\int \frac{(fx)^m (a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$	2120
3.239	$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx$	2124
3.240	$\int (c-a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx$	2128
3.241	$\int (c-a^2cx^2)^2 \operatorname{arccosh}(ax)^3 dx$	2142
3.242	$\int (c-a^2cx^2) \operatorname{arccosh}(ax)^3 dx$	2153
3.243	$\int \frac{\operatorname{arccosh}(ax)^3}{c-a^2cx^2} dx$	2161
3.244	$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^2} dx$	2168
3.245	$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$	2179
3.246	$\int (c-a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$	2192
3.247	$\int (c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx$	2209
3.248	$\int \sqrt{c-a^2cx^2} \operatorname{arccosh}(ax)^3 dx$	2221
3.249	$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c-a^2cx^2}} dx$	2227

3.250	$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx$	2231
3.251	$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx$	2238
3.252	$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$	2249
3.253	$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$	2265
3.254	$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$	2274
3.255	$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$	2283
3.256	$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$	2289
3.257	$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$	2294
3.258	$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx$	2298
3.259	$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx$	2305
3.260	$\int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx$	2312
3.261	$\int \frac{(fx)^m (a+b\operatorname{arccosh}(cx))^3}{\sqrt{1-c^2x^2}} dx$	2322
3.262	$\int \frac{(c-a^2cx^2)^3}{\operatorname{arccosh}(ax)} dx$	2326
3.263	$\int \frac{(c-a^2cx^2)^2}{\operatorname{arccosh}(ax)} dx$	2331
3.264	$\int \frac{c-a^2cx^2}{\operatorname{arccosh}(ax)} dx$	2336
3.265	$\int \frac{1}{(c-a^2cx^2)\operatorname{arccosh}(ax)} dx$	2341
3.266	$\int \frac{1}{(c-a^2cx^2)^2\operatorname{arccosh}(ax)} dx$	2345
3.267	$\int \frac{x^4\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$	2349
3.268	$\int \frac{x^3\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$	2355
3.269	$\int \frac{x^2\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$	2361
3.270	$\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$	2366
3.271	$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$	2371
3.272	$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx$	2376
3.273	$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$	2381
3.274	$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$	2385
3.275	$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$	2389
3.276	$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$	2393
3.277	$\int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$	2399
3.278	$\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$	2405

3.279	$\int \frac{(1-c^2x^2)^{3/2}}{a+\operatorname{barccosh}(cx)} dx$	2410
3.280	$\int \frac{(1-c^2x^2)^{3/2}}{x(a+\operatorname{barccosh}(cx))} dx$	2415
3.281	$\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+\operatorname{barccosh}(cx))} dx$	2420
3.282	$\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+\operatorname{barccosh}(cx))} dx$	2425
3.283	$\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+\operatorname{barccosh}(cx))} dx$	2429
3.284	$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+\operatorname{barccosh}(cx)} dx$	2433
3.285	$\int \frac{x^2(1-c^2x^2)^{5/2}}{a+\operatorname{barccosh}(cx)} dx$	2439
3.286	$\int \frac{x(1-c^2x^2)^{5/2}}{a+\operatorname{barccosh}(cx)} dx$	2445
3.287	$\int \frac{(1-c^2x^2)^{5/2}}{a+\operatorname{barccosh}(cx)} dx$	2451
3.288	$\int \frac{(1-c^2x^2)^{5/2}}{x(a+\operatorname{barccosh}(cx))} dx$	2457
3.289	$\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+\operatorname{barccosh}(cx))} dx$	2462
3.290	$\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+\operatorname{barccosh}(cx))} dx$	2467
3.291	$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+\operatorname{barccosh}(cx))} dx$	2471
3.292	$\int \frac{x^4}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$	2475
3.293	$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$	2480
3.294	$\int \frac{x^2}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$	2485
3.295	$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$	2490
3.296	$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$	2495
3.297	$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$	2499
3.298	$\int \frac{1}{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$	2503
3.299	$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx$	2507
3.300	$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx$	2512
3.301	$\int \frac{x}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx$	2517
3.302	$\int \frac{1}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx$	2523
3.303	$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx$	2527
3.304	$\int \frac{1}{x^2\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx$	2531
3.305	$\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx$	2535
3.306	$\int \frac{x}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx$	2539
3.307	$\int \frac{1}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx$	2543
3.308	$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx$	2547

3.309	$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$	2551
3.310	$\int \frac{x^m(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$	2555
3.311	$\int \frac{x^m\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$	2559
3.312	$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$	2563
3.313	$\int \frac{x^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$	2567
3.314	$\int \frac{x^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))} dx$	2571
3.315	$\int \frac{(c-a^2cx^2)^3}{\operatorname{arccosh}(ax)^2} dx$	2575
3.316	$\int \frac{(c-a^2cx^2)^2}{\operatorname{arccosh}(ax)^2} dx$	2581
3.317	$\int \frac{c-a^2cx^2}{\operatorname{arccosh}(ax)^2} dx$	2586
3.318	$\int \frac{1}{(c-a^2cx^2)\operatorname{arccosh}(ax)^2} dx$	2591
3.319	$\int \frac{1}{(c-a^2cx^2)^2\operatorname{arccosh}(ax)^2} dx$	2596
3.320	$\int \frac{x^3\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2601
3.321	$\int \frac{x^2\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2609
3.322	$\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2619
3.323	$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2628
3.324	$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$	2636
3.325	$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$	2644
3.326	$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$	2649
3.327	$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$	2654
3.328	$\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2658
3.329	$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2666
3.330	$\int \frac{(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2676
3.331	$\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$	2683
3.332	$\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$	2691
3.333	$\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$	2697
3.334	$\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$	2702
3.335	$\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2707
3.336	$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2715
3.337	$\int \frac{(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$	2725



3.338	$\int \frac{(1-c^2x^2)^{5/2}}{x(a+\operatorname{barccosh}(cx))^2} dx$	2732
3.339	$\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+\operatorname{barccosh}(cx))^2} dx$	2740
3.340	$\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+\operatorname{barccosh}(cx))^2} dx$	2745
3.341	$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+\operatorname{barccosh}(cx))^2} dx$	2749
3.342	$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2753
3.343	$\int \frac{x^4}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2760
3.344	$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2766
3.345	$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2772
3.346	$\int \frac{x}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2780
3.347	$\int \frac{1}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2787
3.348	$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2791
3.349	$\int \frac{1}{x^2\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2796
3.350	$\int \frac{x^3}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$	2801
3.351	$\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$	2806
3.352	$\int \frac{x}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$	2811
3.353	$\int \frac{1}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$	2816
3.354	$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$	2821
3.355	$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$	2826
3.356	$\int \frac{x^4}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2830
3.357	$\int \frac{x^3}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2835
3.358	$\int \frac{x^2}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2840
3.359	$\int \frac{x}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2844
3.360	$\int \frac{1}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2848
3.361	$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2853
3.362	$\int \frac{1}{x^2(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2858
3.363	$\int \frac{(fx)^m(1-c^2x^2)^{3/2}}{(a+\operatorname{barccosh}(cx))^2} dx$	2863
3.364	$\int \frac{(fx)^m\sqrt{1-c^2x^2}}{(a+\operatorname{barccosh}(cx))^2} dx$	2868
3.365	$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$	2873
3.366	$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$	2878
3.367	$\int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$	2883
3.368	$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx$	2888

3.369	$\int \frac{x^3(d-c^2dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2893
3.370	$\int \frac{x^2(d-c^2dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2900
3.371	$\int \frac{x(d-c^2dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2907
3.372	$\int \frac{d-c^2dx^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2914
3.373	$\int \frac{d-c^2dx^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2919
3.374	$\int \frac{x^3(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2926
3.375	$\int \frac{x^2(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2933
3.376	$\int \frac{x(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2940
3.377	$\int \frac{(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2947
3.378	$\int \frac{(d-c^2dx^2)^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	2953
3.379	$\int (c-a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx$	2960
3.380	$\int \sqrt{c-a^2cx^2} \sqrt{\operatorname{arccosh}(ax)} dx$	2971
3.381	$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c-a^2cx^2}} dx$	2978
3.382	$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{3/2}} dx$	2982
3.383	$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{5/2}} dx$	2986
3.384	$\int (c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx$	2992
3.385	$\int \sqrt{c-a^2cx^2} \operatorname{arccosh}(ax)^{3/2} dx$	3003
3.386	$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c-a^2cx^2}} dx$	3009
3.387	$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx$	3013
3.388	$\int (c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx$	3017
3.389	$\int \sqrt{c-a^2cx^2} \operatorname{arccosh}(ax)^{5/2} dx$	3036
3.390	$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c-a^2cx^2}} dx$	3045
3.391	$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c-a^2cx^2)^{3/2}} dx$	3049
3.392	$\int (a^2-x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$	3053
3.393	$\int \sqrt{a^2-x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$	3064
3.394	$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2-x^2}} dx$	3072
3.395	$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{3/2}} dx$	3076
3.396	$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{5/2}} dx$	3081
3.397	$\int (a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$	3087

3.398	$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$	3098
3.399	$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx$	3104
3.400	$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx$	3108
3.401	$\int \frac{x}{\sqrt{1-x^2} \sqrt{\operatorname{arccosh}(x)}} dx$	3113
3.402	$\int \frac{(c-a^2cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$	3119
3.403	$\int \frac{(c-a^2cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$	3125
3.404	$\int \frac{\sqrt{c-a^2cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$	3130
3.405	$\int \frac{1}{\sqrt{c-a^2cx^2} \sqrt{\operatorname{arccosh}(ax)}} dx$	3135
3.406	$\int \frac{1}{(c-a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx$	3139
3.407	$\int \frac{1}{(c-a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx$	3143
3.408	$\int \frac{(c-a^2cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$	3147
3.409	$\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$	3153
3.410	$\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx$	3159
3.411	$\int \frac{1}{\sqrt{c-a^2cx^2} \operatorname{arccosh}(ax)^{3/2}} dx$	3166
3.412	$\int \frac{1}{(c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx$	3170
3.413	$\int \frac{1}{(c-a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx$	3175
3.414	$\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$	3180
3.415	$\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx$	3188
3.416	$\int \frac{1}{\sqrt{c-a^2cx^2} \operatorname{arccosh}(ax)^{5/2}} dx$	3195
3.417	$\int \frac{1}{(c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx$	3199
3.418	$\int \frac{1}{(c-a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx$	3204
3.419	$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^n dx$	3209
3.420	$\int x \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^n dx$	3214
3.421	$\int \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^n dx$	3219
3.422	$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^n}{x} dx$	3224
3.423	$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx$	3229
3.424	$\int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^n dx$	3233
3.425	$\int x (d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^n dx$	3239
3.426	$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^n dx$	3245
3.427	$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^n}{x} dx$	3250
3.428	$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx$	3255

3.429	$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$	3260
3.430	$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$	3266
3.431	$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$	3272
3.432	$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$	3278
3.433	$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx$	3285
3.434	$\int \frac{x^3 (a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$	3290
3.435	$\int \frac{x^2 (a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$	3295
3.436	$\int \frac{x (a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$	3300
3.437	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$	3305
3.438	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x \sqrt{1 - c^2 x^2}} dx$	3309
3.439	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx$	3313
3.440	$\int \frac{x^3 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$	3317
3.441	$\int \frac{x^2 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$	3322
3.442	$\int \frac{x (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$	3327
3.443	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$	3332
3.444	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} dx$	3336
3.445	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx$	3340
3.446	$\int \frac{x^2 (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$	3344
3.447	$\int \frac{x (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$	3348
3.448	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$	3352
3.449	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x (d - c^2 dx^2)^{3/2}} dx$	3356
3.450	$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx$	3360
3.451	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$	3364
3.452	$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx$	3368
3.453	$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx$	3372
3.454	$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx$	3376
3.455	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{d - c^2 dx^2} dx$	3380
3.456	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^2} dx$	3384
3.457	$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$	3388
3.458	$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$	3392
3.459	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$	3396
3.460	$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$	3400

3.461	$\int x^4(d+ex^2)(a+\operatorname{barccosh}(cx)) dx$	3404
3.462	$\int x^3(d+ex^2)(a+\operatorname{barccosh}(cx)) dx$	3411
3.463	$\int x^2(d+ex^2)(a+\operatorname{barccosh}(cx)) dx$	3418
3.464	$\int x(d+ex^2)(a+\operatorname{barccosh}(cx)) dx$	3424
3.465	$\int (d+ex^2)(a+\operatorname{barccosh}(cx)) dx$	3430
3.466	$\int \frac{(d+ex^2)(a+\operatorname{barccosh}(cx))}{x} dx$	3435
3.467	$\int \frac{(d+ex^2)(a+\operatorname{barccosh}(cx))}{x^2} dx$	3440
3.468	$\int \frac{(d+ex^2)(a+\operatorname{barccosh}(cx))}{x^3} dx$	3446
3.469	$\int \frac{(d+ex^2)(a+\operatorname{barccosh}(cx))}{x^4} dx$	3452
3.470	$\int x^4(d+ex^2)^2(a+\operatorname{barccosh}(cx)) dx$	3458
3.471	$\int x^3(d+ex^2)^2(a+\operatorname{barccosh}(cx)) dx$	3465
3.472	$\int x^2(d+ex^2)^2(a+\operatorname{barccosh}(cx)) dx$	3474
3.473	$\int x(d+ex^2)^2(a+\operatorname{barccosh}(cx)) dx$	3481
3.474	$\int (d+ex^2)^2(a+\operatorname{barccosh}(cx)) dx$	3489
3.475	$\int \frac{(d+ex^2)^2(a+\operatorname{barccosh}(cx))}{x} dx$	3496
3.476	$\int \frac{(d+ex^2)^2(a+\operatorname{barccosh}(cx))}{x^2} dx$	3503
3.477	$\int \frac{(d+ex^2)^2(a+\operatorname{barccosh}(cx))}{x^3} dx$	3510
3.478	$\int \frac{(d+ex^2)^2(a+\operatorname{barccosh}(cx))}{x^4} dx$	3517
3.479	$\int x^4(d+ex^2)^3(a+\operatorname{barccosh}(cx)) dx$	3525
3.480	$\int x^3(d+ex^2)^3(a+\operatorname{barccosh}(cx)) dx$	3533
3.481	$\int x^2(d+ex^2)^3(a+\operatorname{barccosh}(cx)) dx$	3542
3.482	$\int x(d+ex^2)^3(a+\operatorname{barccosh}(cx)) dx$	3549
3.483	$\int (d+ex^2)^3(a+\operatorname{barccosh}(cx)) dx$	3557
3.484	$\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x} dx$	3564
3.485	$\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x^2} dx$	3572
3.486	$\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x^3} dx$	3579
3.487	$\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x^4} dx$	3586
3.488	$\int (d+ex^2)^4(a+\operatorname{barccosh}(cx)) dx$	3594
3.489	$\int \frac{x^4(a+\operatorname{barccosh}(cx))}{d+ex^2} dx$	3601
3.490	$\int \frac{x^3(a+\operatorname{barccosh}(cx))}{d+ex^2} dx$	3609
3.491	$\int \frac{x^2(a+\operatorname{barccosh}(cx))}{d+ex^2} dx$	3616
3.492	$\int \frac{x(a+\operatorname{barccosh}(cx))}{d+ex^2} dx$	3623
3.493	$\int \frac{a+\operatorname{barccosh}(cx)}{d+ex^2} dx$	3630
3.494	$\int \frac{a+\operatorname{barccosh}(cx)}{x(d+ex^2)} dx$	3637
3.495	$\int \frac{a+\operatorname{barccosh}(cx)}{x^2(d+ex^2)} dx$	3644

3.496	$\int \frac{a+\operatorname{barccosh}(cx)}{x^3(d+ex^2)} dx$	3652
3.497	$\int \frac{a+\operatorname{barccosh}(cx)}{x^4(d+ex^2)} dx$	3660
3.498	$\int \frac{x^3(a+\operatorname{barccosh}(cx))}{(d+ex^2)^2} dx$	3669
3.499	$\int \frac{x(a+\operatorname{barccosh}(cx))}{(d+ex^2)^2} dx$	3676
3.500	$\int \frac{a+\operatorname{barccosh}(cx)}{x(d+ex^2)^2} dx$	3682
3.501	$\int \frac{a+\operatorname{barccosh}(cx)}{x^3(d+ex^2)^2} dx$	3689
3.502	$\int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d+ex^2)^2} dx$	3697
3.503	$\int \frac{x^2(a+\operatorname{barccosh}(cx))}{(d+ex^2)^2} dx$	3706
3.504	$\int \frac{a+\operatorname{barccosh}(cx)}{(d+ex^2)^2} dx$	3714
3.505	$\int \frac{a+\operatorname{barccosh}(cx)}{x^2(d+ex^2)^2} dx$	3722
3.506	$\int \frac{x^5(a+\operatorname{barccosh}(cx))}{(d+ex^2)^3} dx$	3731
3.507	$\int \frac{x^3(a+\operatorname{barccosh}(cx))}{(d+ex^2)^3} dx$	3740
3.508	$\int \frac{x(a+\operatorname{barccosh}(cx))}{(d+ex^2)^3} dx$	3749
3.509	$\int \frac{a+\operatorname{barccosh}(cx)}{x(d+ex^2)^3} dx$	3755
3.510	$\int \frac{a+\operatorname{barccosh}(cx)}{x^3(d+ex^2)^3} dx$	3765
3.511	$\int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d+ex^2)^3} dx$	3775
3.512	$\int \frac{x^2(a+\operatorname{barccosh}(cx))}{(d+ex^2)^3} dx$	3785
3.513	$\int \frac{a+\operatorname{barccosh}(cx)}{(d+ex^2)^3} dx$	3794
3.514	$\int \sqrt{d+ex^2}(a+\operatorname{barccosh}(cx)) dx$	3803
3.515	$\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{d+ex^2}} dx$	3807
3.516	$\int \frac{a+\operatorname{barccosh}(cx)}{(d+ex^2)^{3/2}} dx$	3811
3.517	$\int \frac{a+\operatorname{barccosh}(cx)}{(d+ex^2)^{5/2}} dx$	3817
3.518	$\int \frac{a+\operatorname{barccosh}(cx)}{(d+ex^2)^{7/2}} dx$	3824
3.519	$\int (fx)^m (d+ex^2)^3 (a+\operatorname{barccosh}(cx)) dx$	3832
3.520	$\int (fx)^m (d+ex^2)^2 (a+\operatorname{barccosh}(cx)) dx$	3841
3.521	$\int (fx)^m (d+ex^2) (a+\operatorname{barccosh}(cx)) dx$	3848
3.522	$\int \frac{(fx)^m (a+\operatorname{barccosh}(cx))}{d+ex^2} dx$	3854
3.523	$\int \frac{(fx)^m (a+\operatorname{barccosh}(cx))}{(d+ex^2)^2} dx$	3858
3.524	$\int \frac{(fx)^m (a+\operatorname{barccosh}(cx))}{(d+ex^2)^3} dx$	3862
3.525	$\int (d+ex^2)^3 (a+\operatorname{barccosh}(cx))^2 dx$	3866

3.526	$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx$	3875
3.527	$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx$	3881
3.528	$\int (a + \operatorname{barccosh}(cx))^2 dx$	3887
3.529	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{d + ex^2} dx$	3892
3.530	$\int \sqrt{d + ex^2} (a + \operatorname{barccosh}(cx))^2 dx$	3900
3.531	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d + ex^2}} dx$	3904
3.532	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d + ex^2)^{3/2}} dx$	3908
3.533	$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d + ex^2)^{5/2}} dx$	3912
3.534	$\int \frac{(d + ex^2)^2}{a + \operatorname{barccosh}(cx)} dx$	3916
3.535	$\int \frac{d + ex^2}{a + \operatorname{barccosh}(cx)} dx$	3923
3.536	$\int \frac{1}{a + \operatorname{barccosh}(cx)} dx$	3928
3.537	$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))} dx$	3934
3.538	$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))} dx$	3938
3.539	$\int \frac{\sqrt{d + ex^2}}{a + \operatorname{barccosh}(cx)} dx$	3942
3.540	$\int \frac{1}{\sqrt{d + ex^2} (a + \operatorname{barccosh}(cx))} dx$	3946
3.541	$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$	3950
3.542	$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx$	3954
3.543	$\int \frac{(d + ex^2)^2}{(a + \operatorname{barccosh}(cx))^2} dx$	3958
3.544	$\int \frac{d + ex^2}{(a + \operatorname{barccosh}(cx))^2} dx$	3967
3.545	$\int \frac{1}{(a + \operatorname{barccosh}(cx))^2} dx$	3973
3.546	$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^2} dx$	3979
3.547	$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2} dx$	3984
3.548	$\int \frac{\sqrt{d + ex^2}}{(a + \operatorname{barccosh}(cx))^2} dx$	3989
3.549	$\int \frac{1}{\sqrt{d + ex^2} (a + \operatorname{barccosh}(cx))^2} dx$	3993
3.550	$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$	3997
3.551	$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$	4002
3.552	$\int (d + ex^2)^2 \sqrt{a + \operatorname{barccosh}(cx)} dx$	4007
3.553	$\int (d + ex^2) \sqrt{a + \operatorname{barccosh}(cx)} dx$	4015
3.554	$\int \sqrt{a + \operatorname{barccosh}(cx)} dx$	4021
3.555	$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{d + ex^2} dx$	4027
3.556	$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{(d + ex^2)^2} dx$	4031
3.557	$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx$	4035
3.558	$\int (a + \operatorname{barccosh}(cx))^{3/2} dx$	4041

3.559	$\int \frac{(a+b\operatorname{arccosh}(cx))^{3/2}}{d+ex^2} dx$	4048
3.560	$\int \frac{(a+b\operatorname{arccosh}(cx))^{3/2}}{(d+ex^2)^2} dx$	4052
3.561	$\int \frac{(d+ex^2)^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx$	4056
3.562	$\int \frac{d+ex^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx$	4064
3.563	$\int \frac{1}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx$	4069
3.564	$\int \frac{1}{(d+ex^2)\sqrt{a+b\operatorname{arccosh}(cx)}} dx$	4075
3.565	$\int \frac{1}{(d+ex^2)^2\sqrt{a+b\operatorname{arccosh}(cx)}} dx$	4079
3.566	$\int \frac{d+ex^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	4083
3.567	$\int \frac{1}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	4088
3.568	$\int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	4094
3.569	$\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))^{3/2}} dx$	4098



### 3.1 $\int x^4(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$

3.1.1	Optimal result . . . . .	208
3.1.2	Mathematica [A] (verified) . . . . .	209
3.1.3	Rubi [A] (verified) . . . . .	209
3.1.4	Maple [A] (verified) . . . . .	212
3.1.5	Fricas [A] (verification not implemented) . . . . .	212
3.1.6	Sympy [F] . . . . .	213
3.1.7	Maxima [A] (verification not implemented) . . . . .	213
3.1.8	Giac [F(-2)] . . . . .	214
3.1.9	Mupad [F(-1)] . . . . .	214

#### 3.1.1 Optimal result

Integrand size = 23, antiderivative size = 151

$$\int x^4(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{152bd\sqrt{-1 + cx}\sqrt{1 + cx}}{3675c^5} - \frac{76bdx^2\sqrt{-1 + cx}\sqrt{1 + cx}}{3675c^3} - \frac{19bdx^4\sqrt{-1 + cx}\sqrt{1 + cx}}{1225c} + \frac{1}{49}bcdx^6\sqrt{-1 + cx}\sqrt{1 + cx} + \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) - \frac{1}{7}c^2dx^7(a + \operatorname{barccosh}(cx))$$

```
output 1/5*d*x^5*(a+b*arccosh(c*x))-1/7*c^2*d*x^7*(a+b*arccosh(c*x))-152/3675*b*d
*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5-76/3675*b*d*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/
2)/c^3-19/1225*b*d*x^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c+1/49*b*c*d*x^6*(c*x-1
)^(1/2)*(c*x+1)^(1/2)
```

### 3.1.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.60

$$\int x^4(d - c^2 dx^2)(a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{d(-105ax^5(-7 + 5c^2x^2) + \frac{b\sqrt{-1+cx}\sqrt{1+cx}(-152-76c^2x^2-57c^4x^4+75c^6x^6)}{c^5} - 105bx^5(-7 + 5c^2x^2) \operatorname{arccosh}(cx))}{3675}$$

input `Integrate[x^4*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `(d*(-105*a*x^5*(-7 + 5*c^2*x^2) + (b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(-152 - 76*c^2*x^2 - 57*c^4*x^4 + 75*c^6*x^6))/c^5 - 105*b*x^5*(-7 + 5*c^2*x^2)*ArcCosh[c*x]))/3675`

### 3.1.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$ , Rules used = {6336, 27, 960, 111, 27, 111, 27, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(d - c^2 dx^2)(a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6336$$

$$-bc \int \frac{dx^5(7 - 5c^2x^2)}{35\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^2 dx^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx))$$

$$\downarrow 27$$

$$-\frac{1}{35}bcd \int \frac{x^5(7 - 5c^2x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^2 dx^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx))$$

$$\downarrow 960$$

$$-\frac{1}{35}bcd \left( \frac{19}{7} \int \frac{x^5}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{5}{7}x^6\sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c^2 dx^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx))$$

$$\downarrow 111$$

---

3.1.  $\int x^4(d - c^2 dx^2)(a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& -\frac{1}{35}bcd \left( \frac{19}{7} \left( \frac{\int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6\sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c^2dx^7(a + \\
& \quad \text{barccosh}(cx)) + \frac{1}{5}dx^5(a + \text{barccosh}(cx)) \\
& \quad \downarrow 27 \\
& -\frac{1}{35}bcd \left( \frac{19}{7} \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6\sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c^2dx^7(a + \\
& \quad \text{barccosh}(cx)) + \frac{1}{5}dx^5(a + \text{barccosh}(cx)) \\
& \quad \downarrow 111 \\
& -\frac{1}{35}bcd \left( \frac{19}{7} \left( \frac{4 \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6\sqrt{cx-1}\sqrt{cx+1} \right) - \\
& \quad \frac{1}{7}c^2dx^7(a + \text{barccosh}(cx)) + \frac{1}{5}dx^5(a + \text{barccosh}(cx)) \\
& \quad \downarrow 27 \\
& -\frac{1}{35}bcd \left( \frac{19}{7} \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6\sqrt{cx-1}\sqrt{cx+1} \right) - \\
& \quad \frac{1}{7}c^2dx^7(a + \text{barccosh}(cx)) + \frac{1}{5}dx^5(a + \text{barccosh}(cx)) \\
& \quad \downarrow 83 \\
& -\frac{1}{7}c^2dx^7(a + \text{barccosh}(cx)) + \frac{1}{5}dx^5(a + \text{barccosh}(cx)) - \\
& \quad \frac{1}{35}bcd \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) - \frac{5}{7}x^6\sqrt{cx-1}\sqrt{cx+1} \right)
\end{aligned}$$

input `Int[x^4*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `-1/35*(b*c*d*((-5*x^6*sqrt[-1 + c*x]*sqrt[1 + c*x])/7 + (19*((x^4*sqrt[-1 + c*x]*sqrt[1 + c*x])/(5*c^2) + (4*((2*sqrt[-1 + c*x]*sqrt[1 + c*x])/(3*c^4) + (x^2*sqrt[-1 + c*x]*sqrt[1 + c*x])/(3*c^2)))/(5*c^2)))/7)) + (d*x^5*(a + b*ArcCosh[c*x]))/5 - (c^2*d*x^7*(a + b*ArcCosh[c*x]))/7`

## 3.1.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 111 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 960 `Int[((e_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.1.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.62

method	result	S
parts	$-da\left(\frac{1}{7}c^2x^7 - \frac{1}{5}x^5\right) - \frac{db\left(\frac{\operatorname{arccosh}(cx)c^7x^7}{7} - \frac{\operatorname{arccosh}(cx)c^5x^5}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(75c^6x^6 - 57c^4x^4 - 76c^2x^2 - 152)}{3675}\right)}{c^5}$	9
derivativedivides	$\frac{-da\left(\frac{1}{7}c^7x^7 - \frac{1}{5}c^5x^5\right) - db\left(\frac{\operatorname{arccosh}(cx)c^7x^7}{7} - \frac{\operatorname{arccosh}(cx)c^5x^5}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(75c^6x^6 - 57c^4x^4 - 76c^2x^2 - 152)}{3675}\right)}{c^5}$	9
default	$\frac{-da\left(\frac{1}{7}c^7x^7 - \frac{1}{5}c^5x^5\right) - db\left(\frac{\operatorname{arccosh}(cx)c^7x^7}{7} - \frac{\operatorname{arccosh}(cx)c^5x^5}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(75c^6x^6 - 57c^4x^4 - 76c^2x^2 - 152)}{3675}\right)}{c^5}$	9

input `int(x^4*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-d*a*(1/7*c^2*x^7-1/5*x^5)-d*b/c^5*(1/7*arccosh(c*x)*c^7*x^7-1/5*arccosh(c*x)*c^5*x^5-1/3675*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(75*c^6*x^6-57*c^4*x^4-76*c^2*x^2-152))`

### 3.1.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.75

$$\int x^4(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx)) dx = \frac{525 ac^7 dx^7 - 735 ac^5 dx^5 + 105(5 bc^7 dx^7 - 7 bc^5 dx^5) \log(cx + \sqrt{c^2 x^2 - 1}) - (75 bc^6 dx^6 - 57 bc^4 dx^4 - 76 b^2 c^2 dx^2 - 152 b^2 d) \sqrt{c^2 x^2 - 1}}{3675 c^5}$$

input `integrate(x^4*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `-1/3675*(525*a*c^7*d*x^7 - 735*a*c^5*d*x^5 + 105*(5*b*c^7*d*x^7 - 7*b*c^5*d*x^5)*log(c*x + sqrt(c^2*x^2 - 1)) - (75*b*c^6*d*x^6 - 57*b*c^4*d*x^4 - 76*b*c^2*d*x^2 - 152*b*d)*sqrt(c^2*x^2 - 1))/c^5`

---

3.1.  $\int x^4(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx)) dx$

### 3.1.6 Sympy [F]

$$\int x^4(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -d \left( \int (-ax^4) dx + \int ac^2 x^6 dx + \int (-bx^4 \operatorname{acosh}(cx)) dx + \int bc^2 x^6 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x**4*(-c**2*d*x**2+d)*(a+b*acosh(c*x)),x)`

output `-d*(Integral(-a*x**4, x) + Integral(a*c**2*x**6, x) + Integral(-b*x**4*acosh(c*x), x) + Integral(b*c**2*x**6*acosh(c*x), x))`

### 3.1.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.22

$$\int x^4(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{1}{7} ac^2 dx^7 + \frac{1}{5} adx^5 - \frac{1}{245} \left( 35x^7 \operatorname{arcosh}(cx) - \left( \frac{5\sqrt{c^2x^2 - 1}x^6}{c^2} + \frac{6\sqrt{c^2x^2 - 1}x^4}{c^4} + \frac{8\sqrt{c^2x^2 - 1}x^2}{c^6} + \frac{16\sqrt{c^2x^2 - 1}}{c^8} \right) c \right) bc^2 d + \frac{1}{75} \left( 15x^5 \operatorname{arcosh}(cx) - \left( \frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6} \right) c \right) bd$$

input `integrate(x^4*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/7*a*c^2*d*x^7 + 1/5*a*d*x^5 - 1/245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*c^2*d + 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d`

### 3.1.8 Giac [F(-2)]

Exception generated.

$$\int x^4 (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

### 3.1.9 Mupad [F(-1)]

Timed out.

$$\int x^4 (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2) dx$$

input `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2),x)`

output `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2), x)`

### 3.2 $\int x^3(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$

3.2.1	Optimal result . . . . .	215
3.2.2	Mathematica [A] (verified) . . . . .	215
3.2.3	Rubi [A] (verified) . . . . .	216
3.2.4	Maple [A] (verified) . . . . .	219
3.2.5	Fricas [A] (verification not implemented) . . . . .	219
3.2.6	Sympy [F] . . . . .	220
3.2.7	Maxima [A] (verification not implemented) . . . . .	220
3.2.8	Giac [F(-2)] . . . . .	221
3.2.9	Mupad [F(-1)] . . . . .	221

#### 3.2.1 Optimal result

Integrand size = 23, antiderivative size = 135

$$\int x^3(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{bdx\sqrt{-1+cx}\sqrt{1+cx}}{24c^3} - \frac{bdx^3\sqrt{-1+cx}\sqrt{1+cx}}{36c} + \frac{1}{36}bcdx^5\sqrt{-1+cx}\sqrt{1+cx} - \frac{bd\operatorname{arccosh}(cx)}{24c^4} + \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) - \frac{1}{6}c^2dx^6(a + \operatorname{barccosh}(cx))$$

output

```
-1/24*b*d*arccosh(c*x)/c^4+1/4*d*x^4*(a+b*arccosh(c*x))-1/6*c^2*d*x^6*(a+b*arccosh(c*x))-1/24*b*d*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-1/36*b*d*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c+1/36*b*c*d*x^5*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

#### 3.2.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.23

$$\int x^3(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \frac{1}{4}adx^4 - \frac{1}{6}ac^2dx^6 - \frac{bdx\sqrt{-1+cx}\sqrt{1+cx}}{24c^3} - \frac{bdx^3\sqrt{-1+cx}\sqrt{1+cx}}{36c} + \frac{1}{36}bcdx^5\sqrt{-1+cx}\sqrt{1+cx} + \frac{1}{4}bdx^4\operatorname{arccosh}(cx) - \frac{1}{6}bc^2dx^6\operatorname{arccosh}(cx) - \frac{bd\operatorname{arctanh}\left(\frac{\sqrt{-1+cx}}{\sqrt{1+cx}}\right)}{12c^4}$$



input `Integrate[x^3*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output  $(a*d*x^4)/4 - (a*c^2*d*x^6)/6 - (b*d*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/(24*c^3) - (b*d*x^3*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/(36*c) + (b*c*d*x^5*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/36 + (b*d*x^4*\text{ArcCosh}[c*x])/4 - (b*c^2*d*x^6*\text{ArcCosh}[c*x])/6 - (b*d*\text{ArcTanh}[\text{Sqrt}[-1 + c*x]/\text{Sqrt}[1 + c*x]])/(12*c^4)$

### 3.2.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {6336, 27, 960, 111, 27, 101, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3(d - c^2 dx^2)(a + \text{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6336} \\
 & -bc \int \frac{dx^4(3 - 2c^2 x^2)}{12\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{6}c^2 dx^6(a + \text{barccosh}(cx)) + \frac{1}{4}dx^4(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{12}bcd \int \frac{x^4(3 - 2c^2 x^2)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{6}c^2 dx^6(a + \text{barccosh}(cx)) + \frac{1}{4}dx^4(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{960} \\
 & -\frac{1}{12}bcd \left( \frac{4}{3} \int \frac{x^4}{\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{3}x^5\sqrt{cx - 1}\sqrt{cx + 1} \right) - \frac{1}{6}c^2 dx^6(a + \text{barccosh}(cx)) + \\
 & \quad \frac{1}{4}dx^4(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{111} \\
 & -\frac{1}{12}bcd \left( \frac{4}{3} \left( \frac{\int \frac{3x^2}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{4c^2} + \frac{x^3\sqrt{cx - 1}\sqrt{cx + 1}}{4c^2} \right) - \frac{1}{3}x^5\sqrt{cx - 1}\sqrt{cx + 1} \right) - \frac{1}{6}c^2 dx^6(a + \\
 & \quad \text{barccosh}(cx)) + \frac{1}{4}dx^4(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{12}bcd \left( \frac{4}{3} \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{6}c^2 dx^6 (a + \\
& \quad \text{barccosh}(cx)) + \frac{1}{4}dx^4 (a + \text{barccosh}(cx)) \\
& \quad \downarrow 101 \\
& -\frac{1}{12}bcd \left( \frac{4}{3} \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right) - \\
& \quad \frac{1}{6}c^2 dx^6 (a + \text{barccosh}(cx)) + \frac{1}{4}dx^4 (a + \text{barccosh}(cx)) \\
& \quad \downarrow 43 \\
& \quad -\frac{1}{6}c^2 dx^6 (a + \text{barccosh}(cx)) + \frac{1}{4}dx^4 (a + \text{barccosh}(cx)) - \\
& \frac{1}{12}bcd \left( \frac{4}{3} \left( \frac{3 \left( \frac{\text{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right)
\end{aligned}$$

input `Int[x^3*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `(d*x^4*(a + b*ArcCosh[c*x]))/4 - (c^2*d*x^6*(a + b*ArcCosh[c*x]))/6 - (b*c*d*(-1/3*(x^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (4*((x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(4*c^2) + (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCos[h[c*x]/(2*c^3)))/(4*c^2))))/3)/12`

### 3.2.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 101 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)*(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)n(e + f*x)pSimp[a2d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 111 `Int[((a_.) + (b_.)*(x_))(m_)((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)(m - 1)(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)(m - 2)(c + d*x)n(e + f*x)pSimp[a2d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`

rule 960 `Int[((e_.)*(x_))(m_)((a1_ + (b1_.)*(x_)(non2_))(p_)((a2_ + (b2_.)*(x_)(non2_))(p_)((c_) + (d_.)*(x_)(n_)), x_Symbol] := Simp[d*(e*x)(m + 1)(a1 + b1*x(n/2))(p + 1)((a2 + b2*x(n/2))(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)m(a1 + b1*x(n/2))p(a2 + b2*x(n/2))p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))(m_)((d_) + (e_.)*(x_)2)(p_), x_Symbol] := With[{u = IntHide[(f*x)m(d + e*x2)p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c2*d + e, 0] && IGtQ[p, 0]`

### 3.2.4 Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.12

method	result
parts	$-da\left(\frac{1}{6}c^2x^6 - \frac{1}{4}x^4\right) - \frac{db\left(\frac{\operatorname{arccosh}(cx)c^6x^6}{6} - \frac{c^4x^4\operatorname{arccosh}(cx)}{4} + \frac{\sqrt{cx-1}\sqrt{cx+1}\left(-2c^5x^5\sqrt{c^2x^2-1} + 2\sqrt{c^2x^2-1}c^3x^3 + 3cx\sqrt{c^2x^2-1}\right)}{72\sqrt{c^2x^2-1}}\right)}{c^4}$
derivativedivides	$-da\left(\frac{1}{6}c^6x^6 - \frac{1}{4}c^4x^4\right) - db\left(\frac{\operatorname{arccosh}(cx)c^6x^6}{6} - \frac{c^4x^4\operatorname{arccosh}(cx)}{4} + \frac{\sqrt{cx-1}\sqrt{cx+1}\left(-2c^5x^5\sqrt{c^2x^2-1} + 2\sqrt{c^2x^2-1}c^3x^3 + 3cx\sqrt{c^2x^2-1}\right)}{72\sqrt{c^2x^2-1}}\right)$
default	$-da\left(\frac{1}{6}c^6x^6 - \frac{1}{4}c^4x^4\right) - db\left(\frac{\operatorname{arccosh}(cx)c^6x^6}{6} - \frac{c^4x^4\operatorname{arccosh}(cx)}{4} + \frac{\sqrt{cx-1}\sqrt{cx+1}\left(-2c^5x^5\sqrt{c^2x^2-1} + 2\sqrt{c^2x^2-1}c^3x^3 + 3cx\sqrt{c^2x^2-1}\right)}{72\sqrt{c^2x^2-1}}\right)$

input `int(x^3*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-d*a*(1/6*c^2*x^6-1/4*x^4)-d*b/c^4*(1/6*arccosh(c*x)*c^6*x^6-1/4*c^4*x^4*a  
rccosh(c*x)+1/72*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-2*c^5*x^5*(c^2*x^2-1)^(1/2)  
+2*(c^2*x^2-1)^(1/2)*c^3*x^3+3*c*x*(c^2*x^2-1)^(1/2)+3*ln(c*x+(c^2*x^2-1)^(  
1/2)))/(c^2*x^2-1)^(1/2))`

### 3.2.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.80

$$\int x^3(d - c^2dx^2)(a + b\operatorname{arccosh}(cx)) dx = \frac{12ac^6dx^6 - 18ac^4dx^4 + 3(4bc^6dx^6 - 6bc^4dx^4 + bd)\log(cx + \sqrt{c^2x^2 - 1}) - (2bc^5dx^5 - 2bc^3dx^3 - 3b^2c^2dx^2)}{72c^4}$$

input `integrate(x^3*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `-1/72*(12*a*c^6*d*x^6 - 18*a*c^4*d*x^4 + 3*(4*b*c^6*d*x^6 - 6*b*c^4*d*x^4  
+ b*d)*log(c*x + sqrt(c^2*x^2 - 1)) - (2*b*c^5*d*x^5 - 2*b*c^3*d*x^3 - 3*b  
*c*d*x)*sqrt(c^2*x^2 - 1))/c^4`

### 3.2.6 Sympy [F]

$$\int x^3(d - c^2 dx^2)(a + \operatorname{barccosh}(cx)) dx = -d \left( \int (-ax^3) dx + \int ac^2 x^5 dx + \int (-bx^3 \operatorname{acosh}(cx)) dx + \int bc^2 x^5 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x**3*(-c**2*d*x**2+d)*(a+b*acosh(c*x)),x)`

output `-d*(Integral(-a*x**3, x) + Integral(a*c**2*x**5, x) + Integral(-b*x**3*acosh(c*x), x) + Integral(b*c**2*x**5*acosh(c*x), x))`

### 3.2.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.50

$$\int x^3(d - c^2 dx^2)(a + \operatorname{barccosh}(cx)) dx = -\frac{1}{6} ac^2 dx^6 + \frac{1}{4} adx^4 - \frac{1}{288} \left( 48x^6 \operatorname{arccosh}(cx) - \left( \frac{8\sqrt{c^2x^2 - 1}x^5}{c^2} + \frac{10\sqrt{c^2x^2 - 1}x^3}{c^4} + \frac{15\sqrt{c^2x^2 - 1}x}{c^6} + \frac{15 \log(2c^2x + 2\sqrt{c^2x^2 - 1})}{c^7} \right) c \right) b d + \frac{1}{32} \left( 8x^4 \operatorname{arccosh}(cx) - \left( \frac{2\sqrt{c^2x^2 - 1}x^3}{c^2} + \frac{3\sqrt{c^2x^2 - 1}x}{c^4} + \frac{3 \log(2c^2x + 2\sqrt{c^2x^2 - 1})}{c^5} \right) c \right) b d$$

input `integrate(x^3*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/6*a*c^2*d*x^6 + 1/4*a*d*x^4 - 1/288*(48*x^6*arccosh(c*x) - (8*sqrt(c^2*x^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c^4 + 15*sqrt(c^2*x^2 - 1)*x/c^6 + 15*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^7)*c)*b*c^2*d + 1/32*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*d`

### 3.2.8 Giac [F(-2)]

Exception generated.

$$\int x^3(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.2.9 Mupad [F(-1)]

Timed out.

$$\int x^3(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \int x^3 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2) dx$$

input `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2),x)`

output `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2), x)`

### 3.3 $\int x^2(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$

3.3.1	Optimal result . . . . .	222
3.3.2	Mathematica [A] (verified) . . . . .	223
3.3.3	Rubi [A] (verified) . . . . .	223
3.3.4	Maple [A] (verified) . . . . .	225
3.3.5	Fricas [A] (verification not implemented) . . . . .	226
3.3.6	Sympy [F] . . . . .	226
3.3.7	Maxima [A] (verification not implemented) . . . . .	227
3.3.8	Giac [F(-2)] . . . . .	227
3.3.9	Mupad [F(-1)] . . . . .	228

#### 3.3.1 Optimal result

Integrand size = 23, antiderivative size = 121

$$\int x^2(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{26bd\sqrt{-1 + cx}\sqrt{1 + cx}}{225c^3} - \frac{13bdx^2\sqrt{-1 + cx}\sqrt{1 + cx}}{225c} + \frac{1}{25}bcdx^4\sqrt{-1 + cx}\sqrt{1 + cx} + \frac{1}{3}dx^3(a + \operatorname{barccosh}(cx)) - \frac{1}{5}c^2dx^5(a + \operatorname{barccosh}(cx))$$

```
output 1/3*d*x^3*(a+b*arccosh(c*x))-1/5*c^2*d*x^5*(a+b*arccosh(c*x))-26/225*b*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-13/225*b*d*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c+1/25*b*c*d*x^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

### 3.3.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.74

$$\int x^2(d - c^2 dx^2)(a + \operatorname{arccosh}(cx)) dx = \frac{d(15ac^3x^3(-5 + 3c^2x^2) + b\sqrt{-1 + cx}\sqrt{1 + cx}(26 + 13c^2x^2 - 9c^4x^4) + 15bc^3x^3(-5 + 3c^2x^2) \operatorname{arccosh}(cx))}{225c^3}$$

input `Integrate[x^2*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `-1/225*(d*(15*a*c^3*x^3*(-5 + 3*c^2*x^2) + b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(26 + 13*c^2*x^2 - 9*c^4*x^4) + 15*b*c^3*x^3*(-5 + 3*c^2*x^2)*ArcCosh[c*x]))/c^3`

### 3.3.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {6336, 27, 960, 111, 27, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2(d - c^2 dx^2)(a + \operatorname{arccosh}(cx)) dx \\ & \quad \downarrow \text{6336} \\ & -bc \int \frac{dx^3(5 - 3c^2x^2)}{15\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{5}c^2 dx^5(a + \operatorname{arccosh}(cx)) + \frac{1}{3}dx^3(a + \operatorname{arccosh}(cx)) \\ & \quad \downarrow \text{27} \\ & -\frac{1}{15}bcd \int \frac{x^3(5 - 3c^2x^2)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{5}c^2 dx^5(a + \operatorname{arccosh}(cx)) + \frac{1}{3}dx^3(a + \operatorname{arccosh}(cx)) \\ & \quad \downarrow \text{960} \\ & -\frac{1}{15}bcd \left( \frac{13}{5} \int \frac{x^3}{\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{3}{5}x^4\sqrt{cx - 1}\sqrt{cx + 1} \right) - \frac{1}{5}c^2 dx^5(a + \operatorname{arccosh}(cx)) + \\ & \quad \quad \quad \frac{1}{3}dx^3(a + \operatorname{arccosh}(cx)) \\ & \quad \downarrow \text{111} \end{aligned}$$

---

3.3.  $\int x^2(d - c^2 dx^2)(a + \operatorname{arccosh}(cx)) dx$



$$\begin{aligned}
& -\frac{1}{15}bcd \left( \frac{13}{5} \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) - \frac{3}{5}x^4\sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{5}c^2dx^5(a + \\
& \quad \text{barccosh}(cx)) + \frac{1}{3}dx^3(a + \text{barccosh}(cx)) \\
& \quad \downarrow 27 \\
& -\frac{1}{15}bcd \left( \frac{13}{5} \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) - \frac{3}{5}x^4\sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{5}c^2dx^5(a + \\
& \quad \text{barccosh}(cx)) + \frac{1}{3}dx^3(a + \text{barccosh}(cx)) \\
& \quad \downarrow 83 \\
& -\frac{1}{5}c^2dx^5(a + \text{barccosh}(cx)) + \frac{1}{3}dx^3(a + \text{barccosh}(cx)) - \\
& \quad \frac{1}{15}bcd \left( \frac{13}{5} \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) - \frac{3}{5}x^4\sqrt{cx-1}\sqrt{cx+1} \right)
\end{aligned}$$

input `Int[x^2*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `-1/15*(b*c*d*((-3*x^4*sqrt[-1 + c*x]*sqrt[1 + c*x])/5 + (13*((2*sqrt[-1 + c*x]*sqrt[1 + c*x])/(3*c^4) + (x^2*sqrt[-1 + c*x]*sqrt[1 + c*x])/(3*c^2)))/5)) + (d*x^3*(a + b*ArcCosh[c*x]))/3 - (c^2*d*x^5*(a + b*ArcCosh[c*x]))/5`

### 3.3.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

```
rule 111 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

```
rule 960 Int[((e_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

```
rule 6336 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.3.4 Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.71

method	result	size
parts	$-da\left(\frac{1}{5}c^2x^5 - \frac{1}{3}x^3\right) - \frac{db\left(\frac{\operatorname{arccosh}(cx)c^5x^5}{5} - \frac{c^3x^3\operatorname{arccosh}(cx)}{3} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4x^4-13c^2x^2-26)}{225}\right)}{c^3}$	86
derivativedivides	$\frac{-da\left(\frac{1}{5}c^5x^5 - \frac{1}{3}c^3x^3\right) - db\left(\frac{\operatorname{arccosh}(cx)c^5x^5}{5} - \frac{c^3x^3\operatorname{arccosh}(cx)}{3} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4x^4-13c^2x^2-26)}{225}\right)}{c^3}$	90
default	$\frac{-da\left(\frac{1}{5}c^5x^5 - \frac{1}{3}c^3x^3\right) - db\left(\frac{\operatorname{arccosh}(cx)c^5x^5}{5} - \frac{c^3x^3\operatorname{arccosh}(cx)}{3} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4x^4-13c^2x^2-26)}{225}\right)}{c^3}$	90

```
input int(x^2*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

$$3.3. \int x^2(d - c^2dx^2)(a + b\operatorname{arccosh}(cx)) dx$$

output  $-d*a*(1/5*c^2*x^5-1/3*x^3)-d*b/c^3*(1/5*arccosh(c*x)*c^5*x^5-1/3*c^3*x^3*arccosh(c*x)-1/225*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(9*c^4*x^4-13*c^2*x^2-26))$

### 3.3.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.85

$$\int x^2(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx)) dx = \frac{45 ac^5 dx^5 - 75 ac^3 dx^3 + 15(3 bc^5 dx^5 - 5 bc^3 dx^3) \log(cx + \sqrt{c^2 x^2 - 1}) - (9 bc^4 dx^4 - 13 bc^2 dx^2 - 26 bd)}{225 c^3}$$

input `integrate(x^2*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $-1/225*(45*a*c^5*d*x^5 - 75*a*c^3*d*x^3 + 15*(3*b*c^5*d*x^5 - 5*b*c^3*d*x^3)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (9*b*c^4*d*x^4 - 13*b*c^2*d*x^2 - 26*b*d)*\sqrt{c^2*x^2 - 1})/c^3$

### 3.3.6 Sympy [F]

$$\int x^2(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx)) dx = -d \left( \int (-ax^2) dx + \int ac^2 x^4 dx + \int (-bx^2 \operatorname{acosh}(cx)) dx + \int bc^2 x^4 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x**2*(-c**2*d*x**2+d)*(a+b*acosh(c*x)),x)`

output  $-d*(\operatorname{Integral}(-a*x**2, x) + \operatorname{Integral}(a*c**2*x**4, x) + \operatorname{Integral}(-b*x**2*\operatorname{acosh}(c*x), x) + \operatorname{Integral}(b*c**2*x**4*\operatorname{acosh}(c*x), x))$

### 3.3.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.20

$$\begin{aligned} & \int x^2(d - c^2 dx^2)(a + \operatorname{arccosh}(cx)) dx \\ &= -\frac{1}{5} ac^2 dx^5 \\ & \quad - \frac{1}{75} \left( 15x^5 \operatorname{arccosh}(cx) - \left( \frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6} \right) c \right) bc^2d \\ & \quad + \frac{1}{3} adx^3 + \frac{1}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \right) bd \end{aligned}$$

input `integrate(x^2*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/5*a*c^2*d*x^5 - 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*c^2*d + 1/3*a*d*x^3 + 1/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*d`

### 3.3.8 Giac [F(-2)]

Exception generated.

$$\int x^2(d - c^2 dx^2)(a + \operatorname{arccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^2*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vector & l) Error: Bad Argument Value`

**3.3.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2) dx$$

input `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2),x)`output `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2), x)`

### 3.4 $\int x(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$

3.4.1	Optimal result . . . . .	229
3.4.2	Mathematica [A] (verified) . . . . .	229
3.4.3	Rubi [A] (verified) . . . . .	230
3.4.4	Maple [A] (verified) . . . . .	231
3.4.5	Fricas [A] (verification not implemented) . . . . .	232
3.4.6	Sympy [F] . . . . .	232
3.4.7	Maxima [A] (verification not implemented) . . . . .	233
3.4.8	Giac [F(-2)] . . . . .	233
3.4.9	Mupad [F(-1)] . . . . .	234

#### 3.4.1 Optimal result

Integrand size = 21, antiderivative size = 98

$$\int x(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{3bdx\sqrt{-1 + cx}\sqrt{1 + cx}}{32c} + \frac{bdx(-1 + cx)^{3/2}(1 + cx)^{3/2}}{16c} + \frac{3bd\operatorname{arccosh}(cx)}{32c^2} - \frac{d(1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{4c^2}$$

```
output 1/16*b*d*x*(c*x-1)^(3/2)*(c*x+1)^(3/2)/c+3/32*b*d*arccosh(c*x)/c^2-1/4*d*(-c^2*x^2+1)^2*(a+b*arccosh(c*x))/c^2-3/32*b*d*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c
```

#### 3.4.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.02

$$\int x(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \frac{d\left(cx(b\sqrt{-1 + cx}\sqrt{1 + cx}(5 - 2c^2 x^2) + 8acx(-2 + c^2 x^2)) + 8bc^2 x^2(-2 + c^2 x^2) \operatorname{arccosh}(cx) + 10b\operatorname{arctanh}\left(\frac{cx-1}{cx+1}\right)\right)}{32c^2}$$

```
input Integrate[x*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]
```

output 
$$\frac{-1/32*(d*(c*x*(b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(5 - 2*c^2*x^2) + 8*a*c*x*(-2 + c^2*x^2)) + 8*b*c^2*x^2*(-2 + c^2*x^2)*\text{ArcCosh}[c*x] + 10*b*\text{ArcTanh}[\text{Sqrt}[(-1 + c*x)/(1 + c*x)]])}{c^2}$$

### 3.4.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6329, 40, 40, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d - c^2 dx^2) (a + \text{barccosh}(cx)) dx$$

$$\downarrow 6329$$

$$\frac{bd \int (cx - 1)^{3/2} (cx + 1)^{3/2} dx}{4c} - \frac{d(1 - c^2 x^2)^2 (a + \text{barccosh}(cx))}{4c^2}$$

$$\downarrow 40$$

$$\frac{bd(\frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4} \int \sqrt{cx - 1}\sqrt{cx + 1} dx)}{4c} - \frac{d(1 - c^2 x^2)^2 (a + \text{barccosh}(cx))}{4c^2}$$

$$\downarrow 40$$

$$\frac{bd\left(\frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{1}{2} \int \frac{1}{\sqrt{cx - 1}\sqrt{cx + 1}} dx\right)\right)}{4c} - \frac{d(1 - c^2 x^2)^2 (a + \text{barccosh}(cx))}{4c^2}$$

$$\downarrow 43$$

$$\frac{bd\left(\frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\text{arccosh}(cx)}{2c}\right)\right)}{4c} - \frac{d(1 - c^2 x^2)^2 (a + \text{barccosh}(cx))}{4c^2}$$

input  $\text{Int}[x*(d - c^2*d*x^2)*(a + b*\text{ArcCosh}[c*x]), x]$

output 
$$-1/4*(d*(1 - c^2*x^2)^2*(a + b*\text{ArcCosh}[c*x]))/c^2 + (b*d*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/2 - \text{ArcCosh}[c*x]/(2*c))))/4)/(4*c)$$

### 3.4.3.1 Defintions of rubi rules used

rule 40 
$$\text{Int}[(a + b*x)^m * (c + d*x)^m, x\_Symbol] \rightarrow \text{Simp}[x * (a + b*x)^m * (c + d*x)^m / (2*m + 1), x] + \text{Simp}[2*a*c * (m / (2*m + 1)) \text{Int}[(a + b*x)^{m-1} * (c + d*x)^{m-1}, x], x] /;$$
 FreeQ[{a, b, c, d}, x] && EqQ[b\*c + a\*d, 0] && IGtQ[m + 1/2, 0]

rule 43 
$$\text{Int}[1 / (\text{Sqrt}[a + b*x] * \text{Sqrt}[c + d*x]), x\_Symbol] \rightarrow \text{Simp}[\text{ArcCosh}[b*(x/a) / (b*\text{Sqrt}[d/b])], x] /;$$
 FreeQ[{a, b, c, d}, x] && EqQ[b\*c + a\*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]

rule 6329 
$$\text{Int}[(a + \text{ArcCosh}[c*x])^n * (d + e*x^2)^p, x\_Symbol] \rightarrow \text{Simp}[(d + e*x^2)^{p+1} * (a + b*\text{ArcCosh}[c*x])^n / (2*e*(p+1)), x] - \text{Simp}[b*(n / (2*c*(p+1))) * \text{Simp}[(d + e*x^2)^p / ((1 + c*x)^{p+1} * (1 - c*x)^p) \text{Int}[(1 + c*x)^{p+1/2} * (-1 + c*x)^{p+1/2} * (a + b*\text{ArcCosh}[c*x])^{n-1}, x], x] /;$$
 FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2\*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]

### 3.4.4 Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.39

method	result
derivativedivides	$-\frac{da(c^2x^2-1)^2}{4} - db \left( \frac{c^4x^4 \operatorname{arccosh}(cx)}{4} - \frac{c^2x^2 \operatorname{arccosh}(cx)}{2} + \frac{\operatorname{arccosh}(cx)}{4} - \frac{\sqrt{cx-1}\sqrt{cx+1}(2\sqrt{c^2x^2-1}c^3x^3 - 5cx\sqrt{c^2x^2-1} + 3\ln(c^2x^2-1))}{32\sqrt{c^2x^2-1}} \right) / c^2$
default	$-\frac{da(c^2x^2-1)^2}{4} - db \left( \frac{c^4x^4 \operatorname{arccosh}(cx)}{4} - \frac{c^2x^2 \operatorname{arccosh}(cx)}{2} + \frac{\operatorname{arccosh}(cx)}{4} - \frac{\sqrt{cx-1}\sqrt{cx+1}(2\sqrt{c^2x^2-1}c^3x^3 - 5cx\sqrt{c^2x^2-1} + 3\ln(c^2x^2-1))}{32\sqrt{c^2x^2-1}} \right) / c^2$
parts	$-\frac{da(c^2x^2-1)^2}{4c^2} - \frac{db \left( \frac{c^4x^4 \operatorname{arccosh}(cx)}{4} - \frac{c^2x^2 \operatorname{arccosh}(cx)}{2} + \frac{\operatorname{arccosh}(cx)}{4} - \frac{\sqrt{cx-1}\sqrt{cx+1}(2\sqrt{c^2x^2-1}c^3x^3 - 5cx\sqrt{c^2x^2-1} + 3\ln(c^2x^2-1))}{32\sqrt{c^2x^2-1}} \right)}{c^2}$

input `int(x*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

3.4. 
$$\int x(d - c^2dx^2)(a + b\operatorname{arccosh}(cx)) dx$$



output  $1/c^2*(-1/4*d*a*(c^2*x^2-1)^2-d*b*(1/4*c^4*x^4*\operatorname{arccosh}(c*x)-1/2*c^2*x^2*\operatorname{arccosh}(c*x)+1/4*\operatorname{arccosh}(c*x)-1/32*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(2*(c^2*x^2-1)^{(1/2)}*c^3*x^3-5*c*x*(c^2*x^2-1)^{(1/2)}+3*\ln(c*x+(c^2*x^2-1)^{(1/2)}))/c^2*x^2-1)^{(1/2)})$

### 3.4.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00

$$\int x(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \frac{8ac^4 dx^4 - 16ac^2 dx^2 + (8bc^4 dx^4 - 16bc^2 dx^2 + 5bd) \log(cx + \sqrt{c^2 x^2 - 1}) - (2bc^3 dx^3 - 5bcdx)\sqrt{c^2 x^2 - 1}}{32c^2}$$

input `integrate(x*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $-1/32*(8*a*c^4*d*x^4 - 16*a*c^2*d*x^2 + (8*b*c^4*d*x^4 - 16*b*c^2*d*x^2 + 5*b*d)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (2*b*c^3*d*x^3 - 5*b*c*d*x)*\sqrt{c^2*x^2 - 1})/c^2$

### 3.4.6 Sympy [F]

$$\int x(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -d \left( \int (-ax) dx + \int ac^2 x^3 dx + \int (-bx \operatorname{acosh}(cx)) dx + \int bc^2 x^3 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x*(-c**2*d*x**2+d)*(a+b*acosh(c*x)),x)`

output  $-d*(\operatorname{Integral}(-a*x, x) + \operatorname{Integral}(a*c**2*x**3, x) + \operatorname{Integral}(-b*x*\operatorname{acosh}(c*x), x) + \operatorname{Integral}(b*c**2*x**3*\operatorname{acosh}(c*x), x))$

### 3.4.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.65

$$\int x(d - c^2 dx^2) (a + \operatorname{arccosh}(cx)) dx = -\frac{1}{4} ac^2 dx^4 - \frac{1}{32} \left( 8x^4 \operatorname{arccosh}(cx) - \left( \frac{2\sqrt{c^2x^2 - 1}x^3}{c^2} + \frac{3\sqrt{c^2x^2 - 1}x}{c^4} + \frac{3 \log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^5} \right) c \right) bc^2d + \frac{1}{2} adx^2 + \frac{1}{4} \left( 2x^2 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x}{c^2} + \frac{\log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^3} \right) \right) bd$$

input `integrate(x*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/4*a*c^2*d*x^4 - 1/32*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*c^2*d + 1/2*a*d*x^2 + 1/4*(2*x^2*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x/c^2 + log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^3))*b*d`

### 3.4.8 Giac [F(-2)]

Exception generated.

$$\int x(d - c^2 dx^2) (a + \operatorname{arccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.4.9 Mupad [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2) dx$$

input `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2),x)`output `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2), x)`

### 3.5 $\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$

3.5.1	Optimal result . . . . .	235
3.5.2	Mathematica [A] (verified) . . . . .	235
3.5.3	Rubi [A] (verified) . . . . .	236
3.5.4	Maple [A] (verified) . . . . .	237
3.5.5	Fricas [A] (verification not implemented) . . . . .	238
3.5.6	Sympy [F] . . . . .	238
3.5.7	Maxima [A] (verification not implemented) . . . . .	239
3.5.8	Giac [F(-2)] . . . . .	239
3.5.9	Mupad [F(-1)] . . . . .	239

#### 3.5.1 Optimal result

Integrand size = 20, antiderivative size = 86

$$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{7bd\sqrt{-1 + cx}\sqrt{1 + cx}}{9c} + \frac{1}{9}bcdx^2\sqrt{-1 + cx}\sqrt{1 + cx} + dx(a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 dx^3(a + \operatorname{barccosh}(cx))$$

```
output d*x*(a+b*arccosh(c*x))-1/3*c^2*d*x^3*(a+b*arccosh(c*x))-7/9*b*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c+1/9*b*c*d*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

#### 3.5.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.83

$$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \frac{d(b\sqrt{-1 + cx}\sqrt{1 + cx}(-7 + c^2x^2) + a(9cx - 3c^3x^3) - 3bcx(-3 + c^2x^2) \operatorname{arccosh}(cx))}{9c}$$

```
input Integrate[(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]
```

```
output (d*(b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-7 + c^2*x^2) + a*(9*c*x - 3*c^3*x^3) - 3*b*c*x*(-3 + c^2*x^2)*ArcCosh[c*x]))/(9*c)
```

### 3.5.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6309, 27, 960, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6309} \\
 & -bc \int \frac{dx(3 - c^2 x^2)}{3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2 dx^3(a + \operatorname{barccosh}(cx)) + dx(a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{3}bcd \int \frac{x(3 - c^2 x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2 dx^3(a + \operatorname{barccosh}(cx)) + dx(a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{960} \\
 & -\frac{1}{3}bcd \left( \frac{7}{3} \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{3}c^2 dx^3(a + \operatorname{barccosh}(cx)) + dx(a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{83} \\
 & -\frac{1}{3}c^2 dx^3(a + \operatorname{barccosh}(cx)) + dx(a + \operatorname{barccosh}(cx)) - \\
 & \quad \frac{1}{3}bcd \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right)
 \end{aligned}$$

input `Int[(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `-1/3*(b*c*d*((7*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2) - (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/3)) + d*x*(a + b*ArcCosh[c*x]) - (c^2*d*x^3*(a + b*ArcCosh[c*x]))/3`

## 3.5.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 960 `Int[((e_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

## 3.5.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.83

method	result	size
parts	$-da\left(\frac{1}{3}x^3c^2 - x\right) - \frac{db\left(\frac{c^3x^3 \operatorname{arccosh}(cx) - cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{9}(c^2x^2-7)}{c}\right)}{c}$	71
derivativedivides	$\frac{-da\left(\frac{1}{3}c^3x^3 - cx\right) - db\left(\frac{c^3x^3 \operatorname{arccosh}(cx) - cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{9}(c^2x^2-7)}{c}\right)}{c}$	73
default	$\frac{-da\left(\frac{1}{3}c^3x^3 - cx\right) - db\left(\frac{c^3x^3 \operatorname{arccosh}(cx) - cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{9}(c^2x^2-7)}{c}\right)}{c}$	73

input `int((-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

---

3.5.  $\int (d - c^2dx^2)(a + b\operatorname{arccosh}(cx)) dx$

output  $-d*a*(1/3*x^3*c^2-x)-d*b/c*(1/3*c^3*x^3*\operatorname{arccosh}(c*x)-c*x*\operatorname{arccosh}(c*x)-1/9*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(c^2*x^2-7))$

### 3.5.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.97

$$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \frac{3ac^3 dx^3 - 9acdx + 3(bc^3 dx^3 - 3bcdx) \log(cx + \sqrt{c^2 x^2 - 1}) - (bc^2 dx^2 - 7bd)\sqrt{c^2 x^2 - 1}}{9c}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $-1/9*(3*a*c^3*d*x^3 - 9*a*c*d*x + 3*(b*c^3*d*x^3 - 3*b*c*d*x)*\log(c*x + \operatorname{sqrt}(c^2*x^2 - 1)) - (b*c^2*d*x^2 - 7*b*d)*\operatorname{sqrt}(c^2*x^2 - 1))/c$

### 3.5.6 Sympy [F]

$$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = -d \left( \int (-a) dx + \int (-b \operatorname{acosh}(cx)) dx \right) + \int ac^2 x^2 dx + \int bc^2 x^2 \operatorname{acosh}(cx) dx$$

input `integrate((-c**2*d*x**2+d)*(a+b*acosh(c*x)),x)`

output  $-d*(\operatorname{Integral}(-a, x) + \operatorname{Integral}(-b*\operatorname{acosh}(c*x), x) + \operatorname{Integral}(a*c**2*x**2, x) + \operatorname{Integral}(b*c**2*x**2*\operatorname{acosh}(c*x), x))$

### 3.5.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.13

$$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{1}{3} ac^2 dx^3 - \frac{1}{9} \left( 3x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bc^2 d$$

$$+ adx + \frac{(cx \operatorname{arcosh}(cx) - \sqrt{c^2 x^2 - 1})bd}{c}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/3*a*c^2*d*x^3 - 1/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*c^2*d + a*d*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*d/c`

### 3.5.8 Giac [F(-2)]

Exception generated.

$$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.5.9 Mupad [F(-1)]

Timed out.

$$\int (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2) dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2), x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2), x)`



### 3.6 $\int \frac{(d-c^2 dx^2)(a+b \operatorname{arccosh}(cx))}{x} dx$

3.6.1	Optimal result	240
3.6.2	Mathematica [A] (verified)	241
3.6.3	Rubi [C] (warning: unable to verify)	241
3.6.4	Maple [A] (verified)	245
3.6.5	Fricas [F]	245
3.6.6	Sympy [F]	245
3.6.7	Maxima [F]	246
3.6.8	Giac [F(-2)]	246
3.6.9	Mupad [F(-1)]	247

#### 3.6.1 Optimal result

Integrand size = 23, antiderivative size = 117

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x} dx = \frac{1}{4}bcdx\sqrt{-1 + cx}\sqrt{1 + cx} - \frac{1}{4}b \operatorname{arccosh}(cx) + \frac{1}{2}d(1 - c^2 x^2)(a + b \operatorname{arccosh}(cx)) + \frac{d(a + b \operatorname{arccosh}(cx))^2}{2b} + d(a + b \operatorname{arccosh}(cx)) \log(1 + e^{-2 \operatorname{arccosh}(cx)}) - \frac{1}{2}bd \operatorname{PolyLog}(2, -e^{-2 \operatorname{arccosh}(cx)})$$

output

```
-1/4*b*d*arccosh(c*x)+1/2*d*(-c^2*x^2+1)*(a+b*arccosh(c*x))+1/2*d*(a+b*arccosh(c*x))^2/b+d*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)-1/2*b*d*polylog(2,-1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)+1/4*b*c*d*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

### 3.6.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.11

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{arccosh}(cx))}{x} dx = -\frac{1}{2}ac^2 dx^2 + \frac{1}{4}bcdx\sqrt{-1 + cx}\sqrt{1 + cx}$$

$$- \frac{1}{2}bc^2 dx^2 \operatorname{arccosh}(cx) + \frac{1}{2}bd \operatorname{arctanh}\left(\frac{\sqrt{-1 + cx}}{\sqrt{1 + cx}}\right)$$

$$+ ad \log(x) + \frac{1}{2}bd(\operatorname{arccosh}(cx) (\operatorname{arccosh}(cx)$$

$$+ 2 \log(1 + e^{-2\operatorname{arccosh}(cx)}))$$

$$- \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)}))$$

input `Integrate[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x,x]`

output `-1/2*(a*c^2*d*x^2) + (b*c*d*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/4 - (b*c^2*d*x^2*ArcCosh[c*x])/2 + (b*d*ArcTanh[Sqrt[-1 + c*x]/Sqrt[1 + c*x]])/2 + a*d*Log[x] + (b*d*(ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])])) - PolyLog[2, -E^(-2*ArcCosh[c*x])]))/2`

### 3.6.3 Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.66 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.20, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.478$ , Rules used = {6334, 40, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{arccosh}(cx))}{x} dx$$

$$\downarrow 6334$$

$$d \int \frac{a + \operatorname{arccosh}(cx)}{x} dx + \frac{1}{2}bcd \int \sqrt{cx - 1}\sqrt{cx + 1} dx + \frac{1}{2}d(1 - c^2 x^2)(a + \operatorname{arccosh}(cx))$$

$$\downarrow 40$$

$$\begin{aligned}
& d \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}bcd \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) + \\
& \quad \frac{1}{2}d(1-c^2x^2)(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow 43 \\
& d \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}d(1-c^2x^2)(a + \operatorname{barccosh}(cx)) + \\
& \quad \frac{1}{2}bcd \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \\
& \quad \downarrow 6297 \\
& \frac{d \int - \left( (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2}d(1-c^2x^2)(a + \\
& \quad \operatorname{barccosh}(cx)) + \frac{1}{2}bcd \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \\
& \quad \downarrow 25 \\
& - \frac{d \int (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2}d(1-c^2x^2)(a + \\
& \quad \operatorname{barccosh}(cx)) + \frac{1}{2}bcd \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \\
& \quad \downarrow 3042 \\
& \frac{d \int -i(a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2}d(1-c^2x^2)(a + \\
& \quad \operatorname{barccosh}(cx)) + \frac{1}{2}bcd \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \\
& \quad \downarrow 26 \\
& \frac{id \int (a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2}d(1-c^2x^2)(a + \\
& \quad \operatorname{barccosh}(cx)) + \frac{1}{2}bcd \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \\
& \quad \downarrow 4201 \\
& \frac{id \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 + e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2 \right)}{b} + \\
& \quad \frac{1}{2}d(1-c^2x^2)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}bcd \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \\
& \quad \downarrow 2620
\end{aligned}$$

---

3.6.  $\int \frac{(d-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx$

$$\frac{id(2i(\frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + b$$

$$\frac{1}{2}d(1 - c^2x^2)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}bcd\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)}{b}$$

↓ 2715

$$\frac{id(2i(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) -$$

$$\frac{1}{2}d(1 - c^2x^2)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}bcd\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)}{b}$$

↓ 2838

$$\frac{id(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx)))$$

$$\frac{1}{2}d(1 - c^2x^2)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}bcd\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)}{b}$$

input `Int[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x,x]`

output `(d*(1 - c^2*x^2)*(a + b*ArcCosh[c*x]))/2 + (b*c*d*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/2 + (I*d*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x]))]) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4))/b`

### 3.6.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4201 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6297 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`
- rule 6334 `Int[(((a_) + ArcCosh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_)^2)^(p_))/(x_), x_Symbol] := Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d)^p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.6.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.12

method	result
parts	$-da\left(\frac{c^2x^2}{2} - \ln(x)\right) - \frac{db \operatorname{arccosh}(cx)^2}{2} - \frac{db \operatorname{arccosh}(cx)c^2x^2}{2} + \frac{bcdx\sqrt{cx-1}\sqrt{cx+1}}{4} + \frac{bd \operatorname{arccosh}(cx)}{4} +$
derivativedivides	$-da\left(\frac{c^2x^2}{2} - \ln(cx)\right) - \frac{db \operatorname{arccosh}(cx)^2}{2} + \frac{bcdx\sqrt{cx-1}\sqrt{cx+1}}{4} - \frac{db \operatorname{arccosh}(cx)c^2x^2}{2} + \frac{bd \operatorname{arccosh}(cx)}{4} +$
default	$-da\left(\frac{c^2x^2}{2} - \ln(cx)\right) - \frac{db \operatorname{arccosh}(cx)^2}{2} + \frac{bcdx\sqrt{cx-1}\sqrt{cx+1}}{4} - \frac{db \operatorname{arccosh}(cx)c^2x^2}{2} + \frac{bd \operatorname{arccosh}(cx)}{4} +$

input `int((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`

output  $-d*a*(1/2*c^2*x^2-\ln(x))-1/2*d*b*\operatorname{arccosh}(c*x)^2-1/2*d*b*\operatorname{arccosh}(c*x)*c^2*x^2+1/4*b*c*d*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+1/4*b*d*\operatorname{arccosh}(c*x)+d*b*\operatorname{arccosh}(c*x)*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2)+1/2*d*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2)$

### 3.6.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x} dx = \int -\frac{(c^2 dx^2 - d)(b \operatorname{arccosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x,x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))/x, x)`

### 3.6.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x} dx = -d \left( \int \left( -\frac{a}{x} \right) dx + \int ac^2 x dx \right) + \int \left( -\frac{b \operatorname{arccosh}(cx)}{x} \right) dx + \int bc^2 x \operatorname{arccosh}(cx) dx$$

---

3.6.  $\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x} dx$

input `integrate((-c**2*d*x**2+d)*(a+b*acosh(c*x))/x,x)`

output `-d*(Integral(-a/x, x) + Integral(a*c**2*x, x) + Integral(-b*acosh(c*x)/x, x) + Integral(b*c**2*x*acosh(c*x), x))`

### 3.6.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x} dx = \int -\frac{(c^2 dx^2 - d)(b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `-1/2*a*c^2*d*x^2 + a*d*log(x) - integrate(b*c^2*d*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) - b*d*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

### 3.6.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.6.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d - c^2 dx^2)}{x} dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2))/x,x)`output `int((a + b*acosh(c*x))*(d - c^2*d*x^2))/x, x)`



### 3.7 $\int \frac{(d-c^2 dx^2)(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.7.1	Optimal result . . . . .	248
3.7.2	Mathematica [A] (verified) . . . . .	248
3.7.3	Rubi [A] (verified) . . . . .	249
3.7.4	Maple [A] (verified) . . . . .	251
3.7.5	Fricas [A] (verification not implemented) . . . . .	251
3.7.6	Sympy [F] . . . . .	252
3.7.7	Maxima [A] (verification not implemented) . . . . .	252
3.7.8	Giac [F(-2)] . . . . .	253
3.7.9	Mupad [F(-1)] . . . . .	253

#### 3.7.1 Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \frac{(d - c^2 dx^2)(a + b\operatorname{arccosh}(cx))}{x^2} dx = bcd\sqrt{-1 + cx}\sqrt{1 + cx} - \frac{d(a + b\operatorname{arccosh}(cx))}{x} - c^2 dx(a + b\operatorname{arccosh}(cx)) + bcd \arctan\left(\sqrt{-1 + cx}\sqrt{1 + cx}\right)$$

output `-d*(a+b*arccosh(c*x))/x-c^2*d*x*(a+b*arccosh(c*x))+b*c*d*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))+b*c*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)`

#### 3.7.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.45

$$\int \frac{(d - c^2 dx^2)(a + b\operatorname{arccosh}(cx))}{x^2} dx = -\frac{ad}{x} - ac^2 dx + bcd\sqrt{-1 + cx}\sqrt{1 + cx} - \frac{bd\operatorname{arccosh}(cx)}{x} - bc^2 dx\operatorname{arccosh}(cx) + \frac{bcd\sqrt{-1 + c^2 x^2} \arctan\left(\sqrt{-1 + c^2 x^2}\right)}{\sqrt{-1 + cx}\sqrt{1 + cx}}$$

input `Integrate[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x^2,x]`

output  $-\left(\frac{a*d}{x}\right) - a*c^2*d*x + b*c*d*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x] - (b*d*\text{ArcCosh}[c*x])/x - b*c^2*d*x*\text{ArcCosh}[c*x] + (b*c*d*\text{Sqrt}[-1 + c^2*x^2]*\text{ArcTan}[\text{Sqrt}[-1 + c^2*x^2]])/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])$

### 3.7.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {6336, 25, 27, 960, 103, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)(a + \text{barccosh}(cx))}{x^2} dx \\
 & \quad \downarrow \text{6336} \\
 & -bc \int -\frac{d(c^2 x^2 + 1)}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx + c^2(-d)x(a + \text{barccosh}(cx)) - \frac{d(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{25} \\
 & bc \int \frac{d(c^2 x^2 + 1)}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx + c^2(-d)x(a + \text{barccosh}(cx)) - \frac{d(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{27} \\
 & bcd \int \frac{c^2 x^2 + 1}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx + c^2(-d)x(a + \text{barccosh}(cx)) - \frac{d(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{960} \\
 & bcd \left( \int \frac{1}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx + \sqrt{cx - 1}\sqrt{cx + 1} \right) + c^2(-d)x(a + \text{barccosh}(cx)) - \frac{d(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{103} \\
 & bcd \left( c \int \frac{1}{(cx - 1)(cx + 1)c + c} d(\sqrt{cx - 1}\sqrt{cx + 1}) + \sqrt{cx - 1}\sqrt{cx + 1} \right) + c^2(-d)x(a + \text{barccosh}(cx)) - \frac{d(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

$$c^2(-d)x(a + \operatorname{barccosh}(cx)) - \frac{d(a + \operatorname{barccosh}(cx))}{x} + bcd\left(\arctan\left(\frac{\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{cx-1}\sqrt{cx+1}}\right) + \sqrt{cx-1}\sqrt{cx+1}\right)$$

input `Int[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x^2,x]`

output `-((d*(a + b*ArcCosh[c*x]))/x) - c^2*d*x*(a + b*ArcCosh[c*x]) + b*c*d*(Sqrt[-1 + c*x]*Sqrt[1 + c*x] + ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])`

### 3.7.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 103 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 960 `Int[((e_)*(x_)^(m_))*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

```
rule 6336 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_
)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.7.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.21

method	result
parts	$-da\left(c^2x + \frac{1}{x}\right) - dbc\left(cx \operatorname{arccosh}(cx) + \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}\left(\sqrt{c^2x^2-1} - \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right)\right)}{\sqrt{c^2x^2-1}}\right)$
derivativedivides	$c\left(-da\left(cx + \frac{1}{cx}\right) - db\left(cx \operatorname{arccosh}(cx) + \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}\left(\sqrt{c^2x^2-1} - \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right)\right)}{\sqrt{c^2x^2-1}}\right)\right)$
default	$c\left(-da\left(cx + \frac{1}{cx}\right) - db\left(cx \operatorname{arccosh}(cx) + \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}\left(\sqrt{c^2x^2-1} - \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right)\right)}{\sqrt{c^2x^2-1}}\right)\right)$

```
input int((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)
```

```
output -d*a*(c^2*x+1/x)-d*b*c*(c*x*arccosh(c*x)+arccosh(c*x)/c/x-(c*x-1)^(1/2)*(c
*x+1)^(1/2)/(c^2*x^2-1)^(1/2)*((c^2*x^2-1)^(1/2)-arctan(1/(c^2*x^2-1)^(1/2
))))
```

### 3.7.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.67

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^2} dx = \frac{ac^2 dx^2 - 2bcdx \arctan(-cx + \sqrt{c^2 x^2 - 1}) - \sqrt{c^2 x^2 - 1}bcdx - (bc^2 + b)dx \log(-cx + \sqrt{c^2 x^2 - 1}) + a}{x}$$

```
input integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^2,x, algorithm="fricas")
```

3.7.  $\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^2} dx$

output  $-(a*c^2*d*x^2 - 2*b*c*d*x*\arctan(-c*x + \sqrt{c^2*x^2 - 1}) - \sqrt{c^2*x^2 - 1}*b*c*d*x - (b*c^2 + b)*d*x*\log(-c*x + \sqrt{c^2*x^2 - 1}) + a*d + (b*c^2*d*x^2 - (b*c^2 + b)*d*x + b*d)*\log(c*x + \sqrt{c^2*x^2 - 1}))/x$

### 3.7.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^2} dx = -d \left( \int ac^2 dx + \int \left(-\frac{a}{x^2}\right) dx + \int bc^2 \operatorname{acosh}(cx) dx + \int \left(-\frac{b \operatorname{acosh}(cx)}{x^2}\right) dx \right)$$

input `integrate((-c**2*d*x**2+d)*(a+b*acosh(c*x))/x**2,x)`

output `-d*(Integral(a*c**2, x) + Integral(-a/x**2, x) + Integral(b*c**2*acosh(c*x), x) + Integral(-b*acosh(c*x)/x**2, x))`

### 3.7.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.87

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^2} dx = -ac^2 dx - \left( cx \operatorname{arcosh}(cx) - \sqrt{c^2 x^2 - 1} \right) bcd - \left( c \arcsin\left(\frac{1}{c|x|}\right) + \frac{\operatorname{arcosh}(cx)}{x} \right) bd - \frac{ad}{x}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output `-a*c^2*d*x - (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*c*d - (c*arcsin(1/(c*abs(x))) + arccosh(c*x)/x)*b*d - a*d/x`

### 3.7.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.7.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d - c^2 dx^2)}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2))/x^2,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2))/x^2, x)`

### 3.8 $\int \frac{(d-c^2dx^2)(a+b\operatorname{arccosh}(cx))}{x^3} dx$

3.8.1	Optimal result . . . . .	254
3.8.2	Mathematica [A] (verified) . . . . .	255
3.8.3	Rubi [C] (warning: unable to verify) . . . . .	255
3.8.4	Maple [A] (verified) . . . . .	259
3.8.5	Fricas [F] . . . . .	259
3.8.6	Sympy [F] . . . . .	260
3.8.7	Maxima [F] . . . . .	260
3.8.8	Giac [F(-2)] . . . . .	260
3.8.9	Mupad [F(-1)] . . . . .	261

#### 3.8.1 Optimal result

Integrand size = 23, antiderivative size = 135

$$\int \frac{(d - c^2dx^2)(a + b\operatorname{arccosh}(cx))}{x^3} dx = \frac{bcd\sqrt{-1 + cx}\sqrt{1 + cx}}{2x} - \frac{1}{2}bc^2d\operatorname{arccosh}(cx) - \frac{d(1 - c^2x^2)(a + b\operatorname{arccosh}(cx))}{2x^2} - \frac{c^2d(a + b\operatorname{arccosh}(cx))^2}{2b} - c^2d(a + b\operatorname{arccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)}) + \frac{1}{2}bc^2d \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})$$

output

```
-1/2*b*c^2*d*arccosh(c*x)-1/2*d*(-c^2*x^2+1)*(a+b*arccosh(c*x))/x^2-1/2*c^2*d*(a+b*arccosh(c*x))^2/b-c^2*d*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2))*(c*x+1)^(1/2))^2)+1/2*b*c^2*d*polylog(2,-1/(c*x+(c*x-1)^(1/2))*(c*x+1)^(1/2))^2)+1/2*b*c*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x
```

### 3.8.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.79

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx = -\frac{ad}{2x^2} + \frac{bcd\sqrt{-1+cx}\sqrt{1+cx}}{2x} - \frac{bd\operatorname{arccosh}(cx)}{2x^2} - ac^2 d \log(x) - \frac{1}{2}bc^2 d(\operatorname{arccosh}(cx) (\operatorname{arccosh}(cx) + 2 \log(1 + e^{-2\operatorname{arccosh}(cx)})) - \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)}))$$

input `Integrate[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x^3,x]`

output `-1/2*(a*d)/x^2 + (b*c*d*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*x) - (b*d*ArcCosh[c*x])/(2*x^2) - a*c^2*d*Log[x] - (b*c^2*d*(ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLog[2, -E^(-2*ArcCosh[c*x])]))/2`

### 3.8.3 Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.522$ , Rules used = {6335, 108, 27, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx \\ & \quad \downarrow \text{6335} \\ & c^2(-d) \int \frac{a + \operatorname{barccosh}(cx)}{x} dx - \frac{1}{2}bcd \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2} dx - \frac{d(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))}{2x^2} \\ & \quad \downarrow \text{108} \\ & c^2(-d) \int \frac{a + \operatorname{barccosh}(cx)}{x} dx - \frac{1}{2}bcd \left( \int \frac{c^2}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) - \\ & \quad \frac{d(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))}{2x^2} \\ & \quad \downarrow \text{27} \end{aligned}$$

---

3.8.  $\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx$



$$\begin{aligned}
& c^2(-d) \int \frac{a + \operatorname{barccosh}(cx)}{x} dx - \frac{1}{2}bcd \left( c^2 \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) - \\
& \quad \frac{d(1-c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow 43 \\
& c^2(-d) \int \frac{a + \operatorname{barccosh}(cx)}{x} dx - \frac{d(1-c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \\
& \quad \frac{1}{2}bcd \left( \operatorname{carccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) \\
& \quad \downarrow 6297 \\
& \frac{c^2d \int - \left( (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) \right) d(a + \operatorname{barccosh}(cx))}{\frac{d(1-c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{carccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)} \\
& \quad \downarrow 25 \\
& \frac{c^2d \int (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{\frac{d(1-c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{carccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)} \\
& \quad \downarrow 3042 \\
& \frac{c^2d \int -i(a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{\frac{d(1-c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{carccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)} \\
& \quad \downarrow 26 \\
& \frac{ic^2d \int (a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{\frac{d(1-c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{carccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)} \\
& \quad \downarrow 4201 \\
& \frac{ic^2d \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 + e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2 \right)}{\frac{d(1-c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{carccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)} \\
& \quad \downarrow 2620
\end{aligned}$$

---

3.8.  $\int \frac{(d-c^2x^2)(a + \operatorname{barccosh}(cx))}{x^3} dx$

$$\frac{ic^2d(2i(\frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx)) \frac{d(1 - c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)}{b}}{\downarrow 2715}$$

$$\frac{ic^2d(2i(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx)) \frac{d(1 - c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)}{b}}{\downarrow 2838}$$

$$\frac{ic^2d(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx)) \frac{d(1 - c^2x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd \left( \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right)}{b}}$$

input `Int[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x^3,x]`

output `-1/2*(d*(1 - c^2*x^2)*(a + b*ArcCosh[c*x]))/x^2 - (b*c*d*(-((Sqrt[-1 + c*x]*Sqrt[1 + c*x])/x) + c*ArcCosh[c*x]))/2 - (I*c^2*d*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4))/b`

### 3.8.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 108 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(m + 1))), x] - Simp[1/(b*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4201 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6297 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6335 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])/(f*(m + 1)), x] + (-Simp[b*c*((-d)^p/(f*(m + 1))) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x] - Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x]), x], x]) / ; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0] && ILtQ[(m + 1)/2, 0]`

### 3.8.4 Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.95

method	result
parts	$-\frac{da}{2x^2} - da c^2 \ln(x) - db c^2 \left( -\frac{\operatorname{arccosh}(cx)^2}{2} + \frac{-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx)}{2c^2x^2} + \operatorname{arccosh}(cx) \right)$
derivativedivides	$c^2 \left( -da \left( \ln(cx) + \frac{1}{2c^2x^2} \right) - db \left( -\frac{\operatorname{arccosh}(cx)^2}{2} + \frac{-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx)}{2c^2x^2} + \operatorname{arccosh}(cx) \right) \right)$
default	$c^2 \left( -da \left( \ln(cx) + \frac{1}{2c^2x^2} \right) - db \left( -\frac{\operatorname{arccosh}(cx)^2}{2} + \frac{-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx)}{2c^2x^2} + \operatorname{arccosh}(cx) \right) \right)$

input `int((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)`

output  $-1/2*d*a/x^2-d*a*c^2*\ln(x)-d*b*c^2*(-1/2*\operatorname{arccosh}(c*x)^2+1/2*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+c^2*x^2+\operatorname{arccosh}(c*x))/c^2/x^2+\operatorname{arccosh}(c*x)*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)+1/2*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2))$

### 3.8.5 Fracas [F]

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int -\frac{(c^2 dx^2 - d)(b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^3,x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))/x^3, x)`

---

3.8.  $\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^3} dx$

### 3.8.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx = -d \left( \int \left( -\frac{a}{x^3} \right) dx + \int \frac{ac^2}{x} dx \right. \\ \left. + \int \left( -\frac{b \operatorname{acosh}(cx)}{x^3} \right) dx + \int \frac{bc^2 \operatorname{acosh}(cx)}{x} dx \right)$$

input `integrate((-c**2*d*x**2+d)*(a+b*acosh(c*x))/x**3,x)`

output `-d*(Integral(-a/x**3, x) + Integral(a*c**2/x, x) + Integral(-b*acosh(c*x)/x**3, x) + Integral(b*c**2*acosh(c*x)/x, x))`

### 3.8.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx = \int -\frac{(c^2 dx^2 - d)(b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `-b*c^2*d*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x) - a*c^2*d*log(x) + 1/2*b*d*(sqrt(c^2*x^2 - 1)*c/x - arccosh(c*x)/x^2) - 1/2*a*d/x^2`

### 3.8.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.8.  $\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^3} dx$

**3.8.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d - c^2 dx^2)}{x^3} dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2))/x^3,x)`output `int((a + b*acosh(c*x))*(d - c^2*d*x^2))/x^3, x)`

### 3.9 $\int \frac{(d-c^2dx^2)(a+b\operatorname{arccosh}(cx))}{x^4} dx$

3.9.1	Optimal result . . . . .	262
3.9.2	Mathematica [A] (verified) . . . . .	262
3.9.3	Rubi [A] (verified) . . . . .	263
3.9.4	Maple [A] (verified) . . . . .	265
3.9.5	Fricas [A] (verification not implemented) . . . . .	265
3.9.6	Sympy [F] . . . . .	266
3.9.7	Maxima [A] (verification not implemented) . . . . .	266
3.9.8	Giac [F(-2)] . . . . .	267
3.9.9	Mupad [F(-1)] . . . . .	267

#### 3.9.1 Optimal result

Integrand size = 23, antiderivative size = 90

$$\int \frac{(d - c^2dx^2)(a + b\operatorname{arccosh}(cx))}{x^4} dx = \frac{bcd\sqrt{-1 + cx}\sqrt{1 + cx}}{6x^2} - \frac{d(a + b\operatorname{arccosh}(cx))}{3x^3} + \frac{c^2d(a + b\operatorname{arccosh}(cx))}{x} - \frac{5}{6}bc^3d \arctan\left(\sqrt{-1 + cx}\sqrt{1 + cx}\right)$$

output `-1/3*d*(a+b*arccosh(c*x))/x^3+c^2*d*(a+b*arccosh(c*x))/x-5/6*b*c^3*d*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))+1/6*b*c*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x^2`

#### 3.9.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.41

$$\int \frac{(d - c^2dx^2)(a + b\operatorname{arccosh}(cx))}{x^4} dx = -\frac{ad}{3x^3} + \frac{ac^2d}{x} + \frac{bcd\sqrt{-1 + cx}\sqrt{1 + cx}}{6x^2} - \frac{bd\operatorname{arccosh}(cx)}{3x^3} + \frac{bc^2d\operatorname{arccosh}(cx)}{x} - \frac{5bc^3d\sqrt{-1 + c^2x^2} \arctan\left(\sqrt{-1 + c^2x^2}\right)}{6\sqrt{-1 + cx}\sqrt{1 + cx}}$$

input `Integrate[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*(a*d)/x^3 + (a*c^2*d)/x + (b*c*d*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(6*x^2) - (b*d*ArcCosh[c*x])/(3*x^3) + (b*c^2*d*ArcCosh[c*x])/x - (5*b*c^3*d*Sqrt[-1 + c^2*x^2]*ArcTan[Sqrt[-1 + c^2*x^2]])/(6*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.9.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {6336, 27, 956, 103, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)(a + \text{barccosh}(cx))}{x^4} dx \\
 & \quad \downarrow \text{6336} \\
 & -bc \int -\frac{d(1 - 3c^2 x^2)}{3x^3 \sqrt{cx - 1} \sqrt{cx + 1}} dx + \frac{c^2 d(a + \text{barccosh}(cx))}{x} - \frac{d(a + \text{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3}bcd \int \frac{1 - 3c^2 x^2}{x^3 \sqrt{cx - 1} \sqrt{cx + 1}} dx + \frac{c^2 d(a + \text{barccosh}(cx))}{x} - \frac{d(a + \text{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{956} \\
 & \frac{1}{3}bcd \left( \frac{\sqrt{cx - 1} \sqrt{cx + 1}}{2x^2} - \frac{5}{2}c^2 \int \frac{1}{x \sqrt{cx - 1} \sqrt{cx + 1}} dx \right) + \frac{c^2 d(a + \text{barccosh}(cx))}{x} - \frac{d(a + \text{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{103} \\
 & \frac{1}{3}bcd \left( \frac{\sqrt{cx - 1} \sqrt{cx + 1}}{2x^2} - \frac{5}{2}c^3 \int \frac{1}{(cx - 1)(cx + 1)c + c} d(\sqrt{cx - 1} \sqrt{cx + 1}) \right) + \frac{c^2 d(a + \text{barccosh}(cx))}{x} - \frac{d(a + \text{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{218}
 \end{aligned}$$

---

3.9.  $\int \frac{(d - c^2 dx^2)(a + \text{barccosh}(cx))}{x^4} dx$



$$\frac{c^2 d(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{d(a + \operatorname{barccosh}(cx))}{3x^3} + \frac{1}{3}bcd \left( \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} - \frac{5}{2}c^2 \arctan\left(\sqrt{cx-1}\sqrt{cx+1}\right) \right)$$

input `Int[((d - c^2*d*x^2)*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*(d*(a + b*ArcCosh[c*x]))/x^3 + (c^2*d*(a + b*ArcCosh[c*x]))/x + (b*c*d*((Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*x^2) - (5*c^2*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/2))/3`

### 3.9.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 103 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 956 `Int[((e_)*(x_)^(m_))*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[c*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(a1*a2*e^(m + 1))), x] + Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(a1*a2*e^n*(m + 1)) Int[(e*x)^(m + n)*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]`

```
rule 6336 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_
)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.9.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.30

method	result
parts	$-da\left(-\frac{c^2}{x} + \frac{1}{3x^3}\right) - db c^3 \left(\frac{\operatorname{arccosh}(cx)}{3c^3x^3} - \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{6c^2x^2\sqrt{c^2x^2-1}} \left(5 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) c^2x^2 + \sqrt{c^2x^2-1}\right)\right)$
derivativedivides	$c^3 \left(-da\left(\frac{1}{3c^3x^3} - \frac{1}{cx}\right) - db \left(\frac{\operatorname{arccosh}(cx)}{3c^3x^3} - \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{6c^2x^2\sqrt{c^2x^2-1}} \left(5 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) c^2x^2 + \sqrt{c^2x^2-1}\right)\right)\right)$
default	$c^3 \left(-da\left(\frac{1}{3c^3x^3} - \frac{1}{cx}\right) - db \left(\frac{\operatorname{arccosh}(cx)}{3c^3x^3} - \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{6c^2x^2\sqrt{c^2x^2-1}} \left(5 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) c^2x^2 + \sqrt{c^2x^2-1}\right)\right)\right)$

```
input int((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)
```

```
output -d*a*(-c^2/x+1/3/x^3)-d*b*c^3*(1/3/c^3/x^3*arccosh(c*x)-arccosh(c*x)/c/x-1
/6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(5*arctan(1/(c^2*x^2-1)^(1/2))*c^2*x^2+(c^2
*x^2-1)^(1/2))/c^2/x^2/(c^2*x^2-1)^(1/2))
```

### 3.9.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.62

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^4} dx = \frac{10bc^3 dx^3 \arctan(-cx + \sqrt{c^2x^2 - 1}) - 6ac^2 dx^2 + 2(3bc^2 - b) dx^3 \log(-cx + \sqrt{c^2x^2 - 1}) - \sqrt{c^2x^2 - 1}}{6x^3}$$

```
input integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")
```

3.9.  $\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^4} dx$

output 
$$\frac{-1/6*(10*b*c^3*d*x^3*\arctan(-c*x + \sqrt{c^2*x^2 - 1}) - 6*a*c^2*d*x^2 + 2*(3*b*c^2 - b)*d*x^3*\log(-c*x + \sqrt{c^2*x^2 - 1}) - \sqrt{c^2*x^2 - 1}*b*c*d*x + 2*a*d - 2*(3*b*c^2*d*x^2 - (3*b*c^2 - b)*d*x^3 - b*d)*\log(c*x + \sqrt{c^2*x^2 - 1}))}{x^3}$$

### 3.9.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^4} dx = -d \left( \int \left( -\frac{a}{x^4} \right) dx + \int \frac{ac^2}{x^2} dx \right. \\ \left. + \int \left( -\frac{b \operatorname{acosh}(cx)}{x^4} \right) dx + \int \frac{bc^2 \operatorname{acosh}(cx)}{x^2} dx \right)$$

input `integrate((-c**2*d*x**2+d)*(a+b*acosh(c*x))/x**4,x)`

output `-d*(Integral(-a/x**4, x) + Integral(a*c**2/x**2, x) + Integral(-b*acosh(c*x)/x**4, x) + Integral(b*c**2*acosh(c*x)/x**2, x))`

### 3.9.7 Maxima [A] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.99

$$\int \frac{(d - c^2 dx^2)(a + b \operatorname{arccosh}(cx))}{x^4} dx \\ = \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arcosh}(cx)}{x} \right) bc^2 d \\ - \frac{1}{6} \left( \left( c^2 \arcsin \left( \frac{1}{c|x|} \right) - \frac{\sqrt{c^2 x^2 - 1}}{x^2} \right) c + \frac{2 \operatorname{arcosh}(cx)}{x^3} \right) bd + \frac{ac^2 d}{x} - \frac{ad}{3x^3}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")`

output `(c*arcsin(1/(c*abs(x))) + arccosh(c*x)/x)*b*c^2*d - 1/6*((c^2*arcsin(1/(c*abs(x))) - sqrt(c^2*x^2 - 1)/x^2)*c + 2*arccosh(c*x)/x^3)*b*d + a*c^2*d/x - 1/3*a*d/x^3`

### 3.9.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.9.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)(a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d - c^2 dx^2)}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2))/x^4,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2))/x^4, x)`

### 3.10 $\int x^4(d - c^2dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.10.1	Optimal result . . . . .	268
3.10.2	Mathematica [A] (verified) . . . . .	269
3.10.3	Rubi [A] (verified) . . . . .	269
3.10.4	Maple [A] (verified) . . . . .	272
3.10.5	Fricas [A] (verification not implemented) . . . . .	272
3.10.6	Sympy [F] . . . . .	273
3.10.7	Maxima [A] (verification not implemented) . . . . .	273
3.10.8	Giac [F(-2)] . . . . .	274
3.10.9	Mupad [F(-1)] . . . . .	274

#### 3.10.1 Optimal result

Integrand size = 25, antiderivative size = 206

$$\int x^4(d - c^2dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{8bd^2\sqrt{-1+cx}\sqrt{1+cx}}{315c^5} + \frac{4bd^2(-1+cx)^{3/2}(1+cx)^{3/2}}{945c^5} - \frac{bd^2(-1+cx)^{5/2}(1+cx)^{5/2}}{525c^5}$$

$$- \frac{10bd^2(-1+cx)^{7/2}(1+cx)^{7/2}}{441c^5} - \frac{bd^2(-1+cx)^{9/2}(1+cx)^{9/2}}{81c^5}$$

$$+ \frac{1}{5}d^2x^5(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2d^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}c^4d^2x^9(a + \operatorname{barccosh}(cx))$$

```
output 4/945*b*d^2*(c*x-1)^(3/2)*(c*x+1)^(3/2)/c^5-1/525*b*d^2*(c*x-1)^(5/2)*(c*x
+1)^(5/2)/c^5-10/441*b*d^2*(c*x-1)^(7/2)*(c*x+1)^(7/2)/c^5-1/81*b*d^2*(c*x
-1)^(9/2)*(c*x+1)^(9/2)/c^5+1/5*d^2*x^5*(a+b*arccosh(c*x))-2/7*c^2*d^2*x^7
*(a+b*arccosh(c*x))+1/9*c^4*d^2*x^9*(a+b*arccosh(c*x))-8/315*b*d^2*(c*x-1)
^(1/2)*(c*x+1)^(1/2)/c^5
```

### 3.10.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.60

$$\int x^4 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{d^2 (315ac^5 x^5 (63 - 90c^2 x^2 + 35c^4 x^4) - b\sqrt{-1 + cx}\sqrt{1 + cx} (2104 + 1052c^2 x^2 + 789c^4 x^4 - 2650c^6 x^6 + 1225c^8 x^8)) + 315b^2 c^5 x^5 (63 - 90c^2 x^2 + 35c^4 x^4) \operatorname{ArcCosh}[cx]}{99225c^5}$$

input `Integrate[x^4*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `(d^2*(315*a*c^5*x^5*(63 - 90*c^2*x^2 + 35*c^4*x^4) - b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2104 + 1052*c^2*x^2 + 789*c^4*x^4 - 2650*c^6*x^6 + 1225*c^8*x^8) + 315*b*c^5*x^5*(63 - 90*c^2*x^2 + 35*c^4*x^4)*ArcCosh[c*x])/(99225*c^5)`

### 3.10.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 200, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6336, 27, 1905, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6336}$$

$$-bc \int \frac{d^2 x^5 (35c^4 x^4 - 90c^2 x^2 + 63)}{315\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{9} c^4 d^2 x^9 (a + \operatorname{barccosh}(cx)) - \frac{2}{7} c^2 d^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5} d^2 x^5 (a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{27}$$

$$-\frac{1}{315} bcd^2 \int \frac{x^5 (35c^4 x^4 - 90c^2 x^2 + 63)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{9} c^4 d^2 x^9 (a + \operatorname{barccosh}(cx)) - \frac{2}{7} c^2 d^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5} d^2 x^5 (a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{1905}$$

---

3.10.  $\int x^4 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& -\frac{bcd^2\sqrt{c^2x^2-1}\int\frac{x^5(35c^4x^4-90c^2x^2+63)}{\sqrt{c^2x^2-1}}dx}{315\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{9}c^4d^2x^9(a+\operatorname{barccosh}(cx))-\frac{2}{7}c^2d^2x^7(a+\operatorname{barccosh}(cx))+\frac{1}{5}d^2x^5(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 1578 \\
& -\frac{bcd^2\sqrt{c^2x^2-1}\int\frac{x^4(35c^4x^4-90c^2x^2+63)}{\sqrt{c^2x^2-1}}dx^2}{630\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{9}c^4d^2x^9(a+\operatorname{barccosh}(cx))-\frac{2}{7}c^2d^2x^7(a+\operatorname{barccosh}(cx))+\frac{1}{5}d^2x^5(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 1195 \\
& -\frac{bcd^2\sqrt{c^2x^2-1}\int\left(\frac{35(c^2x^2-1)^{7/2}}{c^4}+\frac{50(c^2x^2-1)^{5/2}}{c^4}+\frac{3(c^2x^2-1)^{3/2}}{c^4}-\frac{4\sqrt{c^2x^2-1}}{c^4}+\frac{8}{c^4\sqrt{c^2x^2-1}}\right)dx^2}{630\sqrt{cx-1}\sqrt{cx+1}}+ \\
& \quad \frac{1}{9}c^4d^2x^9(a+\operatorname{barccosh}(cx))-\frac{2}{7}c^2d^2x^7(a+\operatorname{barccosh}(cx))+\frac{1}{5}d^2x^5(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 2009 \\
& \frac{\frac{1}{9}c^4d^2x^9(a+\operatorname{barccosh}(cx))-\frac{2}{7}c^2d^2x^7(a+\operatorname{barccosh}(cx))+\frac{1}{5}d^2x^5(a+\operatorname{barccosh}(cx))-bcd^2\sqrt{c^2x^2-1}\left(\frac{70(c^2x^2-1)^{9/2}}{9c^6}+\frac{100(c^2x^2-1)^{7/2}}{7c^6}+\frac{6(c^2x^2-1)^{5/2}}{5c^6}-\frac{8(c^2x^2-1)^{3/2}}{3c^6}+\frac{16\sqrt{c^2x^2-1}}{c^6}\right)}{630\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^4*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `-1/630*(b*c*d^2*sqrt[-1 + c^2*x^2]*((16*sqrt[-1 + c^2*x^2])/c^6 - (8*(-1 + c^2*x^2)^(3/2))/(3*c^6) + (6*(-1 + c^2*x^2)^(5/2))/(5*c^6) + (100*(-1 + c^2*x^2)^(7/2))/(7*c^6) + (70*(-1 + c^2*x^2)^(9/2))/(9*c^6)))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (d^2*x^5*(a + b*ArcCosh[c*x]))/5 - (2*c^2*d^2*x^7*(a + b*ArcCosh[c*x]))/7 + (c^4*d^2*x^9*(a + b*ArcCosh[c*x]))/9`

## 3.10.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 1195 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`
- rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`
- rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`



### 3.10.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.60

method	result
parts	$d^2 a \left( \frac{1}{9} c^4 x^9 - \frac{2}{7} c^2 x^7 + \frac{1}{5} x^5 \right) + \frac{d^2 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{2 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{\sqrt{cx-1} \sqrt{cx+1} (1225 c^8 x^8 - 2650 c^6 x^6 + 789 c^4 x^4 + 1052 c^2 x^2 + 2104)}{99225 c^5} \right)}{c^5}$
derivativedivides	$\frac{d^2 a \left( \frac{1}{9} c^9 x^9 - \frac{2}{7} c^7 x^7 + \frac{1}{5} c^5 x^5 \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{2 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{\sqrt{cx-1} \sqrt{cx+1} (1225 c^8 x^8 - 2650 c^6 x^6 + 789 c^4 x^4 + 1052 c^2 x^2 + 2104)}{99225 c^5} \right)}{c^5}$
default	$\frac{d^2 a \left( \frac{1}{9} c^9 x^9 - \frac{2}{7} c^7 x^7 + \frac{1}{5} c^5 x^5 \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{2 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{\sqrt{cx-1} \sqrt{cx+1} (1225 c^8 x^8 - 2650 c^6 x^6 + 789 c^4 x^4 + 1052 c^2 x^2 + 2104)}{99225 c^5} \right)}{c^5}$

input `int(x^4*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output  $d^2 a \left( \frac{1}{9} c^4 x^9 - \frac{2}{7} c^2 x^7 + \frac{1}{5} x^5 \right) + d^2 b / c^5 \left( \frac{1}{9} \operatorname{arccosh}(cx) c^9 x^9 - \frac{2}{7} \operatorname{arccosh}(cx) c^7 x^7 + \frac{1}{5} \operatorname{arccosh}(cx) c^5 x^5 - \frac{1}{99225} (cx-1)^{1/2} (cx+1)^{1/2} (1225 c^8 x^8 - 2650 c^6 x^6 + 789 c^4 x^4 + 1052 c^2 x^2 + 2104) \right)$

### 3.10.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.80

$$\int x^4 (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx$$

$$= \frac{11025 a c^9 d^2 x^9 - 28350 a c^7 d^2 x^7 + 19845 a c^5 d^2 x^5 + 315 (35 b c^9 d^2 x^9 - 90 b c^7 d^2 x^7 + 63 b c^5 d^2 x^5) \log(cx + \sqrt{c^2 x^2 - 1}) - (1225 b c^8 d^2 x^8 - 2650 b c^6 d^2 x^6 + 789 b c^4 d^2 x^4 + 1052 b c^2 d^2 x^2 + 2104 b d^2) \sqrt{c^2 x^2 - 1}}{99225 c^5}$$

input `integrate(x^4*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $\frac{1}{99225} (11025 a c^9 d^2 x^9 - 28350 a c^7 d^2 x^7 + 19845 a c^5 d^2 x^5 + 315 (35 b c^9 d^2 x^9 - 90 b c^7 d^2 x^7 + 63 b c^5 d^2 x^5) \log(cx + \sqrt{c^2 x^2 - 1}) - (1225 b c^8 d^2 x^8 - 2650 b c^6 d^2 x^6 + 789 b c^4 d^2 x^4 + 1052 b c^2 d^2 x^2 + 2104 b d^2) \sqrt{c^2 x^2 - 1}) / c^5$

### 3.10.6 Sympy [F]

$$\int x^4 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = d^2 \left( \int ax^4 dx + \int (-2ac^2 x^6) dx + \int ac^4 x^8 dx \right. \\ \left. + \int bx^4 \operatorname{acosh}(cx) dx + \int (-2bc^2 x^6 \operatorname{acosh}(cx)) dx + \int bc^4 x^8 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x**4*(-c**2*d*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `d**2*(Integral(a*x**4, x) + Integral(-2*a*c**2*x**6, x) + Integral(a*c**4*x**8, x) + Integral(b*x**4*acosh(c*x), x) + Integral(-2*b*c**2*x**6*acosh(c*x), x) + Integral(b*c**4*x**8*acosh(c*x), x))`

### 3.10.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.55

$$\int x^4 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{1}{9} ac^4 d^2 x^9 - \frac{2}{7} ac^2 d^2 x^7 \\ + \frac{1}{2835} \left( 315 x^9 \operatorname{arcosh}(cx) - \left( \frac{35 \sqrt{c^2 x^2 - 1} x^8}{c^2} + \frac{40 \sqrt{c^2 x^2 - 1} x^6}{c^4} + \frac{48 \sqrt{c^2 x^2 - 1} x^4}{c^6} + \frac{64 \sqrt{c^2 x^2 - 1} x^2}{c^8} \right) c \right) bc^2 d \\ + \frac{1}{5} ad^2 x^5 \\ - \frac{2}{245} \left( 35 x^7 \operatorname{arcosh}(cx) - \left( \frac{5 \sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6 \sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16 \sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) bc^2 d \\ + \frac{1}{75} \left( 15 x^5 \operatorname{arcosh}(cx) - \left( \frac{3 \sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4 \sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bd^2$$

input `integrate(x^4*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output  $1/9*a*c^4*d^2*x^9 - 2/7*a*c^2*d^2*x^7 + 1/2835*(315*x^9*\operatorname{arccosh}(c*x) - (35*\sqrt{c^2*x^2 - 1})*x^8/c^2 + 40*\sqrt{c^2*x^2 - 1})*x^6/c^4 + 48*\sqrt{c^2*x^2 - 1})*x^4/c^6 + 64*\sqrt{c^2*x^2 - 1})*x^2/c^8 + 128*\sqrt{c^2*x^2 - 1})/c^{10})*c)*b*c^4*d^2 + 1/5*a*d^2*x^5 - 2/245*(35*x^7*\operatorname{arccosh}(c*x) - (5*\sqrt{c^2*x^2 - 1})*x^6/c^2 + 6*\sqrt{c^2*x^2 - 1})*x^4/c^4 + 8*\sqrt{c^2*x^2 - 1})*x^2/c^6 + 16*\sqrt{c^2*x^2 - 1})/c^8)*c)*b*c^2*d^2 + 1/75*(15*x^5*\operatorname{arccosh}(c*x) - (3*\sqrt{c^2*x^2 - 1})*x^4/c^2 + 4*\sqrt{c^2*x^2 - 1})*x^2/c^4 + 8*\sqrt{c^2*x^2 - 1})/c^6)*c)*b*d^2$

### 3.10.8 Giac [F(-2)]

Exception generated.

$$\int x^4(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const ve cteur & l) Error: Bad Argument Value

### 3.10.9 Mupad [F(-1)]

Timed out.

$$\int x^4(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2 dx$$

input `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2,x)`

output `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2, x)`

### 3.11 $\int x^3(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.11.1	Optimal result . . . . .	275
3.11.2	Mathematica [A] (verified) . . . . .	276
3.11.3	Rubi [A] (verified) . . . . .	276
3.11.4	Maple [A] (verified) . . . . .	280
3.11.5	Fricas [A] (verification not implemented) . . . . .	280
3.11.6	Sympy [F] . . . . .	281
3.11.7	Maxima [B] (verification not implemented) . . . . .	281
3.11.8	Giac [F(-2)] . . . . .	282
3.11.9	Mupad [F(-1)] . . . . .	282

#### 3.11.1 Optimal result

Integrand size = 25, antiderivative size = 200

$$\int x^3(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{73bd^2x\sqrt{-1+cx}\sqrt{1+cx}}{3072c^3} - \frac{73bd^2x^3\sqrt{-1+cx}\sqrt{1+cx}}{4608c} + \frac{43bcd^2x^5\sqrt{-1+cx}\sqrt{1+cx}}{1152}$$

$$- \frac{1}{64}bc^3d^2x^7\sqrt{-1+cx}\sqrt{1+cx} - \frac{73bd^2\operatorname{arccosh}(cx)}{3072c^4} + \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx))$$

$$- \frac{1}{3}c^2d^2x^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}c^4d^2x^8(a + \operatorname{barccosh}(cx))$$

```
output -73/3072*b*d^2*arccosh(c*x)/c^4+1/4*d^2*x^4*(a+b*arccosh(c*x))-1/3*c^2*d^2
*x^6*(a+b*arccosh(c*x))+1/8*c^4*d^2*x^8*(a+b*arccosh(c*x))-73/3072*b*d^2*x
*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-73/4608*b*d^2*x^3*(c*x-1)^(1/2)*(c*x+1)^(
1/2)/c+43/1152*b*c*d^2*x^5*(c*x-1)^(1/2)*(c*x+1)^(1/2)-1/64*b*c^3*d^2*x^7*
(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

### 3.11.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.97

$$\int x^3 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{d^2 (2304ac^4x^4 - 3072ac^6x^6 + 1152ac^8x^8 - 219bcx\sqrt{-1+cx}\sqrt{1+cx} - 146bc^3x^3\sqrt{-1+cx}\sqrt{1+cx} + 344b^2c^5x^5\sqrt{-1+cx}\sqrt{1+cx} - 144b^2c^7x^7\sqrt{-1+cx}\sqrt{1+cx} + 384b^2c^4x^4(6 - 8c^2x^2 + 3c^4x^4)\operatorname{ArcCosh}[cx] - 438b^2\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1+cx)/(1+cx)])])}{9216c^4}$$

input `Integrate[x^3*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `(d^2*(2304*a*c^4*x^4 - 3072*a*c^6*x^6 + 1152*a*c^8*x^8 - 219*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 146*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 344*b^2*c^5*x^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 144*b*c^7*x^7*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 384*b*c^4*x^4*(6 - 8*c^2*x^2 + 3*c^4*x^4)*ArcCosh[c*x] - 438*b^2*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]])/(9216*c^4)`

### 3.11.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.16, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6336, 27, 1905, 1590, 27, 363, 262, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6336}$$

$$-bc \int \frac{d^2 x^4 (3c^4 x^4 - 8c^2 x^2 + 6)}{24\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{8} c^4 d^2 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} c^2 d^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 x^4 (a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{27}$$

$$-\frac{1}{24} bcd^2 \int \frac{x^4 (3c^4 x^4 - 8c^2 x^2 + 6)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{8} c^4 d^2 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} c^2 d^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 x^4 (a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{1905}$$

---

3.11.  $\int x^3 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& -\frac{bcd^2\sqrt{c^2x^2-1}\int\frac{x^4(3c^4x^4-8c^2x^2+6)}{\sqrt{c^2x^2-1}}dx}{24\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{8}c^4d^2x^8(a+\operatorname{barccosh}(cx))-\frac{1}{3}c^2d^2x^6(a+\operatorname{barccosh}(cx)) \\
& \quad +\frac{1}{4}d^2x^4(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 1590 \\
& -\frac{bcd^2\sqrt{c^2x^2-1}\left(\frac{\int\frac{c^2x^4(48-43c^2x^2)}{\sqrt{c^2x^2-1}}dx}{8c^2}+\frac{3}{8}c^2x^7\sqrt{c^2x^2-1}\right)}{24\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{8}c^4d^2x^8(a+\operatorname{barccosh}(cx))- \\
& \quad \frac{1}{3}c^2d^2x^6(a+\operatorname{barccosh}(cx))+\frac{1}{4}d^2x^4(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 27 \\
& -\frac{bcd^2\sqrt{c^2x^2-1}\left(\frac{1}{8}\int\frac{x^4(48-43c^2x^2)}{\sqrt{c^2x^2-1}}dx+\frac{3}{8}c^2x^7\sqrt{c^2x^2-1}\right)}{24\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{8}c^4d^2x^8(a+\operatorname{barccosh}(cx))- \\
& \quad \frac{1}{3}c^2d^2x^6(a+\operatorname{barccosh}(cx))+\frac{1}{4}d^2x^4(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 363 \\
& -\frac{bcd^2\sqrt{c^2x^2-1}\left(\frac{1}{8}\left(\frac{73}{6}\int\frac{x^4}{\sqrt{c^2x^2-1}}dx-\frac{43}{6}x^5\sqrt{c^2x^2-1}\right)+\frac{3}{8}c^2x^7\sqrt{c^2x^2-1}\right)}{24\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{8}c^4d^2x^8(a+\operatorname{barccosh}(cx))-\frac{1}{3}c^2d^2x^6(a+\operatorname{barccosh}(cx))+\frac{1}{4}d^2x^4(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 262 \\
& -\frac{bcd^2\sqrt{c^2x^2-1}\left(\frac{1}{8}\left(\frac{73}{6}\left(\frac{3\int\frac{x^2}{\sqrt{c^2x^2-1}}dx}{4c^2}+\frac{x^3\sqrt{c^2x^2-1}}{4c^2}\right)-\frac{43}{6}x^5\sqrt{c^2x^2-1}\right)+\frac{3}{8}c^2x^7\sqrt{c^2x^2-1}\right)}{24\sqrt{cx-1}\sqrt{cx+1}}+ \\
& \quad \frac{1}{8}c^4d^2x^8(a+\operatorname{barccosh}(cx))-\frac{1}{3}c^2d^2x^6(a+\operatorname{barccosh}(cx))+\frac{1}{4}d^2x^4(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 262 \\
& -\frac{bcd^2\sqrt{c^2x^2-1}\left(\frac{1}{8}\left(\frac{73}{6}\left(\frac{3\left(\frac{\int\frac{1}{\sqrt{c^2x^2-1}}dx}{2c^2}+\frac{x\sqrt{c^2x^2-1}}{2c^2}\right)}{4c^2}+\frac{x^3\sqrt{c^2x^2-1}}{4c^2}\right)-\frac{43}{6}x^5\sqrt{c^2x^2-1}\right)+\frac{3}{8}c^2x^7\sqrt{c^2x^2-1}\right)}{24\sqrt{cx-1}\sqrt{cx+1}}+ \\
& \quad \frac{1}{8}c^4d^2x^8(a+\operatorname{barccosh}(cx))-\frac{1}{3}c^2d^2x^6(a+\operatorname{barccosh}(cx))+\frac{1}{4}d^2x^4(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow 224
\end{aligned}$$

---

3.11.  $\int x^3(d-c^2dx^2)^2(a+\operatorname{barccosh}(cx))dx$

$$bcd^2\sqrt{c^2x^2-1} \left( \frac{1}{8} \left( \frac{73}{6} \left( \frac{3 \left( \frac{\int \frac{1-\frac{c^2x^2-d\sqrt{c^2x^2-1}}{1-\frac{c^2x^2-1}{2c^2}} + \frac{x\sqrt{c^2x^2-1}}{2c^2}}{4c^2} + \frac{x^3\sqrt{c^2x^2-1}}{4c^2} \right) - \frac{43}{6}x^5\sqrt{c^2x^2-1} + \frac{3}{8}c^2x^7\sqrt{c^2x^2-1}}{24\sqrt{cx-1}\sqrt{cx+1}} \right) \right) \right) - \frac{1}{3}c^2d^2x^6(a + \operatorname{barccosh}(cx)) + \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx))$$

↓ 219

$$\frac{1}{8}c^4d^2x^8(a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2d^2x^6(a + \operatorname{barccosh}(cx)) + \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) - bcd^2\sqrt{c^2x^2-1} \left( \frac{1}{8} \left( \frac{73}{6} \left( \frac{3 \left( \frac{\operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2-1}}\right) + \frac{x\sqrt{c^2x^2-1}}{2c^2}}{4c^2} + \frac{x^3\sqrt{c^2x^2-1}}{4c^2} \right) - \frac{43}{6}x^5\sqrt{c^2x^2-1} + \frac{3}{8}c^2x^7\sqrt{c^2x^2-1}}{24\sqrt{cx-1}\sqrt{cx+1}} \right) \right) \right)$$

input `Int[x^3*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `(d^2*x^4*(a + b*ArcCosh[c*x]))/4 - (c^2*d^2*x^6*(a + b*ArcCosh[c*x]))/3 + (c^4*d^2*x^8*(a + b*ArcCosh[c*x]))/8 - (b*c*d^2*Sqrt[-1 + c^2*x^2]*((3*c^2*x^7*Sqrt[-1 + c^2*x^2])/8 + ((-43*x^5*Sqrt[-1 + c^2*x^2])/6 + (73*((x^3*Sqrt[-1 + c^2*x^2])/(4*c^2) + (3*((x*Sqrt[-1 + c^2*x^2])/(2*c^2) + ArcTanh[(c*x)/Sqrt[-1 + c^2*x^2]]/(2*c^3)))/(4*c^2)))/6)/8))/(24*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.11.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

---

3.11.  $\int x^3(d - c^2dx^2)^2(a + \operatorname{barccosh}(cx)) dx$

- rule 224 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 262 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 363 `Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^(m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`
- rule 1590 `Int[((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]`
- rule 1905 `Int[((f_)*(x_)^(m_))*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(p_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_))*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`



### 3.11.4 Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.96

method	result
parts	$d^2 a \left( \frac{1}{8} c^4 x^8 - \frac{1}{3} c^2 x^6 + \frac{1}{4} x^4 \right) + \frac{d^2 b \left( \frac{\operatorname{arccosh}(cx) c^8 x^8}{8} - \frac{\operatorname{arccosh}(cx) c^6 x^6}{3} + \frac{c^4 x^4 \operatorname{arccosh}(cx)}{4} - \frac{\sqrt{cx-1} \sqrt{cx+1} (144 c^7 x^7 \sqrt{c^2 x^2 - 1}}{c^4} \right)}{c^4}$
derivativedivides	$\frac{d^2 a \left( \frac{1}{8} c^8 x^8 - \frac{1}{3} c^6 x^6 + \frac{1}{4} c^4 x^4 \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^8 x^8}{8} - \frac{\operatorname{arccosh}(cx) c^6 x^6}{3} + \frac{c^4 x^4 \operatorname{arccosh}(cx)}{4} - \frac{\sqrt{cx-1} \sqrt{cx+1} (144 c^7 x^7 \sqrt{c^2 x^2 - 1}}{c^4} \right)}{c^4}$
default	$\frac{d^2 a \left( \frac{1}{8} c^8 x^8 - \frac{1}{3} c^6 x^6 + \frac{1}{4} c^4 x^4 \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^8 x^8}{8} - \frac{\operatorname{arccosh}(cx) c^6 x^6}{3} + \frac{c^4 x^4 \operatorname{arccosh}(cx)}{4} - \frac{\sqrt{cx-1} \sqrt{cx+1} (144 c^7 x^7 \sqrt{c^2 x^2 - 1}}{c^4} \right)}{c^4}$

input `int(x^3*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{d^2 a \left( \frac{1}{8} c^4 x^8 - \frac{1}{3} c^2 x^6 + \frac{1}{4} x^4 \right) + d^2 b \left( \frac{1}{8} \operatorname{arccosh}(cx) c^8 x^8 - \frac{1}{3} \operatorname{arccosh}(cx) c^6 x^6 + \frac{1}{4} c^4 x^4 \operatorname{arccosh}(cx) - \frac{1}{9216} (cx-1)^{1/2} (cx+1)^{1/2} (144 c^7 x^7 (c^2 x^2 - 1)^{1/2} - 344 c^5 x^5 (c^2 x^2 - 1)^{1/2} + 146 (c^2 x^2 - 1)^{1/2} c^3 x^3 + 219 c x (c^2 x^2 - 1)^{1/2} + 219 \ln(cx + (c^2 x^2 - 1)^{1/2}))}{c^4} \right)}{(c^2 x^2 - 1)^{1/2}}$$

### 3.11.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.80

$$\int x^3 (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \frac{1152 ac^8 d^2 x^8 - 3072 ac^6 d^2 x^6 + 2304 ac^4 d^2 x^4 + 3(384 bc^8 d^2 x^8 - 1024 bc^6 d^2 x^6 + 768 bc^4 d^2 x^4 - 73 bd^2) \log(cx + \sqrt{c^2 x^2 - 1})}{9216 c^4}$$

input `integrate(x^3*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output 
$$\frac{1}{9216} (1152 a c^8 d^2 x^8 - 3072 a c^6 d^2 x^6 + 2304 a c^4 d^2 x^4 + 3(384 b c^8 d^2 x^8 - 1024 b c^6 d^2 x^6 + 768 b c^4 d^2 x^4 - 73 b d^2) \log(cx + \sqrt{c^2 x^2 - 1}) - (144 b c^7 d^2 x^7 - 344 b c^5 d^2 x^5 + 146 b c^3 d^2 x^3 + 219 b c d^2 x) \sqrt{c^2 x^2 - 1}) / c^4$$

### 3.11.6 Sympy [F]

$$\int x^3 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = d^2 \left( \int ax^3 dx + \int (-2ac^2 x^5) dx + \int ac^4 x^7 dx \right. \\ \left. + \int bx^3 \operatorname{acosh}(cx) dx + \int (-2bc^2 x^5 \operatorname{acosh}(cx)) dx + \int bc^4 x^7 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x**3*(-c**2*d*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `d**2*(Integral(a*x**3, x) + Integral(-2*a*c**2*x**5, x) + Integral(a*c**4*x**7, x) + Integral(b*x**3*acosh(c*x), x) + Integral(-2*b*c**2*x**5*acosh(c*x), x) + Integral(b*c**4*x**7*acosh(c*x), x))`

### 3.11.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 346 vs. 2(168) = 336.

Time = 0.25 (sec) , antiderivative size = 346, normalized size of antiderivative = 1.73

$$\int x^3 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{1}{8} ac^4 d^2 x^8 - \frac{1}{3} ac^2 d^2 x^6 \\ + \frac{1}{3072} \left( 384 x^8 \operatorname{arcosh}(cx) - \left( \frac{48 \sqrt{c^2 x^2 - 1} x^7}{c^2} + \frac{56 \sqrt{c^2 x^2 - 1} x^5}{c^4} + \frac{70 \sqrt{c^2 x^2 - 1} x^3}{c^6} + \frac{105 \sqrt{c^2 x^2 - 1} x}{c^8} \right) \right. \\ \left. + \frac{1}{4} ad^2 x^4 - \frac{1}{144} \left( 48 x^6 \operatorname{arcosh}(cx) - \left( \frac{8 \sqrt{c^2 x^2 - 1} x^5}{c^2} + \frac{10 \sqrt{c^2 x^2 - 1} x^3}{c^4} + \frac{15 \sqrt{c^2 x^2 - 1} x}{c^6} + \frac{15 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1})}{c^7} \right) \right) \right. \\ \left. + \frac{1}{32} \left( 8 x^4 \operatorname{arcosh}(cx) - \left( \frac{2 \sqrt{c^2 x^2 - 1} x^3}{c^2} + \frac{3 \sqrt{c^2 x^2 - 1} x}{c^4} + \frac{3 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1})}{c^5} \right) \right) c \right) bd^2$$

input `integrate(x^3*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

```
output 1/8*a*c^4*d^2*x^8 - 1/3*a*c^2*d^2*x^6 + 1/3072*(384*x^8*arccosh(c*x) - (48
*sqrt(c^2*x^2 - 1)*x^7/c^2 + 56*sqrt(c^2*x^2 - 1)*x^5/c^4 + 70*sqrt(c^2*x^
2 - 1)*x^3/c^6 + 105*sqrt(c^2*x^2 - 1)*x/c^8 + 105*log(2*c^2*x + 2*sqrt(c^
2*x^2 - 1)*c)/c^9)*c)*b*c^4*d^2 + 1/4*a*d^2*x^4 - 1/144*(48*x^6*arccosh(c*
x) - (8*sqrt(c^2*x^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c^4 + 15*sqrt
(c^2*x^2 - 1)*x/c^6 + 15*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^7)*c)*b*c^
2*d^2 + 1/32*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c
^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*d^2
```

### 3.11.8 Giac [F(-2)]

Exception generated.

$$\int x^3 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.11.9 Mupad [F(-1)]

Timed out.

$$\int x^3 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x^3 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2 dx$$

```
input int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2,x)
```

```
output int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2, x)
```

### 3.12 $\int x^2(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.12.1	Optimal result . . . . .	283
3.12.2	Mathematica [A] (verified) . . . . .	283
3.12.3	Rubi [A] (verified) . . . . .	284
3.12.4	Maple [A] (verified) . . . . .	286
3.12.5	Fricas [A] (verification not implemented) . . . . .	287
3.12.6	Sympy [F] . . . . .	287
3.12.7	Maxima [A] (verification not implemented) . . . . .	288
3.12.8	Giac [F(-2)] . . . . .	288
3.12.9	Mupad [F(-1)] . . . . .	289

#### 3.12.1 Optimal result

Integrand size = 25, antiderivative size = 177

$$\int x^2(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{8bd^2\sqrt{-1+cx}\sqrt{1+cx}}{105c^3} + \frac{4bd^2(-1+cx)^{3/2}(1+cx)^{3/2}}{315c^3}$$

$$- \frac{bd^2(-1+cx)^{5/2}(1+cx)^{5/2}}{175c^3} - \frac{bd^2(-1+cx)^{7/2}(1+cx)^{7/2}}{49c^3}$$

$$+ \frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) - \frac{2}{5}c^2d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}c^4d^2x^7(a + \operatorname{barccosh}(cx))$$

output  $4/315*b*d^2*(c*x-1)^{(3/2)}*(c*x+1)^{(3/2)}/c^3-1/175*b*d^2*(c*x-1)^{(5/2)}*(c*x+1)^{(5/2)}/c^3-1/49*b*d^2*(c*x-1)^{(7/2)}*(c*x+1)^{(7/2)}/c^3+1/3*d^2*x^3*(a+b*\operatorname{arccosh}(c*x))-2/5*c^2*d^2*x^5*(a+b*\operatorname{arccosh}(c*x))+1/7*c^4*d^2*x^7*(a+b*\operatorname{arccosh}(c*x))-8/105*b*d^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3$

#### 3.12.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.66

$$\int x^2(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{d^2(105ac^3x^3(35 - 42c^2x^2 + 15c^4x^4) - b\sqrt{-1+cx}\sqrt{1+cx}(818 + 409c^2x^2 - 612c^4x^4 + 225c^6x^6) + 105bc^3}{11025c^3}$$

input `Integrate[x^2*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output  $(d^2*(105*a*c^3*x^3*(35 - 42*c^2*x^2 + 15*c^4*x^4) - b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(818 + 409*c^2*x^2 - 612*c^4*x^4 + 225*c^6*x^6) + 105*b*c^3*x^3*(35 - 42*c^2*x^2 + 15*c^4*x^4)*\text{ArcCosh}[c*x]))/(11025*c^3)$

### 3.12.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6336, 27, 1905, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2(d - c^2dx^2)^2(a + \text{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6336} \\
 & -bc \int \frac{d^2x^3(15c^4x^4 - 42c^2x^2 + 35)}{105\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{7}c^4d^2x^7(a + \text{barccosh}(cx)) - \frac{2}{5}c^2d^2x^5(a + \text{barccosh}(cx)) + \\
 & \quad \frac{1}{3}d^2x^3(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{105}bcd^2 \int \frac{x^3(15c^4x^4 - 42c^2x^2 + 35)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{7}c^4d^2x^7(a + \text{barccosh}(cx)) - \frac{2}{5}c^2d^2x^5(a + \\
 & \quad \text{barccosh}(cx)) + \frac{1}{3}d^2x^3(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{1905} \\
 & -\frac{bcd^2\sqrt{c^2x^2 - 1} \int \frac{x^3(15c^4x^4 - 42c^2x^2 + 35)}{\sqrt{c^2x^2 - 1}} dx}{105\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{7}c^4d^2x^7(a + \text{barccosh}(cx)) - \frac{2}{5}c^2d^2x^5(a + \\
 & \quad \text{barccosh}(cx)) + \frac{1}{3}d^2x^3(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{1578} \\
 & -\frac{bcd^2\sqrt{c^2x^2 - 1} \int \frac{x^2(15c^4x^4 - 42c^2x^2 + 35)}{\sqrt{c^2x^2 - 1}} dx^2}{210\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{7}c^4d^2x^7(a + \text{barccosh}(cx)) - \frac{2}{5}c^2d^2x^5(a + \\
 & \quad \text{barccosh}(cx)) + \frac{1}{3}d^2x^3(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{1195}
 \end{aligned}$$

---

3.12.  $\int x^2(d - c^2dx^2)^2(a + \text{barccosh}(cx)) dx$

$$\begin{aligned}
& \frac{bcd^2\sqrt{c^2x^2-1} \int \left( \frac{15(c^2x^2-1)^{5/2}}{c^2} + \frac{3(c^2x^2-1)^{3/2}}{c^2} - \frac{4\sqrt{c^2x^2-1}}{c^2} + \frac{8}{c^2\sqrt{c^2x^2-1}} \right) dx^2}{210\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{7}c^4d^2x^7(a + \\
& \operatorname{barccosh}(cx)) - \frac{2}{5}c^2d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2009} \\
& \frac{\frac{1}{7}c^4d^2x^7(a + \operatorname{barccosh}(cx)) - \frac{2}{5}c^2d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) -}{210\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad bcd^2\sqrt{c^2x^2-1} \left( \frac{30(c^2x^2-1)^{7/2}}{7c^4} + \frac{6(c^2x^2-1)^{5/2}}{5c^4} - \frac{8(c^2x^2-1)^{3/2}}{3c^4} + \frac{16\sqrt{c^2x^2-1}}{c^4} \right)
\end{aligned}$$

input `Int[x^2*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `-1/210*(b*c*d^2*Sqrt[-1 + c^2*x^2]*((16*Sqrt[-1 + c^2*x^2])/c^4 - (8*(-1 + c^2*x^2)^(3/2))/(3*c^4) + (6*(-1 + c^2*x^2)^(5/2))/(5*c^4) + (30*(-1 + c^2*x^2)^(7/2))/(7*c^4)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (d^2*x^3*(a + b*ArcCosh[c*x]))/3 - (2*c^2*d^2*x^5*(a + b*ArcCosh[c*x]))/5 + (c^4*d^2*x^7*(a + b*ArcCosh[c*x]))/7`

### 3.12.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 1578 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m-1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m-1)/2]`

```
rule 1905 Int[((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_)^(non2_.))^(q_.)*((d2_) + (e2_.)
*(x_)^(non2_.))^(q_.)*((a_.) + (b_.)*(x_)^(n_) + (c_.)*(x_)^(n2_))^(p_.), x
_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[
q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a
+ b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p,
q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6336 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_
)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.12.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.66

method	result
parts	$d^2 a \left( \frac{1}{7} c^4 x^7 - \frac{2}{5} c^2 x^5 + \frac{1}{3} x^3 \right) + \frac{d^2 b \left( \frac{\operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{2 \operatorname{arccosh}(cx) c^5 x^5}{5} + \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - \frac{\sqrt{cx-1} \sqrt{cx+1} (225c^6 - 612c^4 x^6 + 409c^2 x^4 + 818)}{11025} \right)}{c^3}$
derivativedivides	$\frac{d^2 a \left( \frac{1}{7} c^7 x^7 - \frac{2}{5} c^5 x^5 + \frac{1}{3} c^3 x^3 \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{2 \operatorname{arccosh}(cx) c^5 x^5}{5} + \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - \frac{\sqrt{cx-1} \sqrt{cx+1} (225c^6 x^6 - 612c^4 x^4 + 409c^2 x^2 + 818)}{11025} \right)}{c^3}$
default	$\frac{d^2 a \left( \frac{1}{7} c^7 x^7 - \frac{2}{5} c^5 x^5 + \frac{1}{3} c^3 x^3 \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{2 \operatorname{arccosh}(cx) c^5 x^5}{5} + \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - \frac{\sqrt{cx-1} \sqrt{cx+1} (225c^6 x^6 - 612c^4 x^4 + 409c^2 x^2 + 818)}{11025} \right)}{c^3}$

```
input int(x^2*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output d^2*a*(1/7*c^4*x^7-2/5*c^2*x^5+1/3*x^3)+d^2*b/c^3*(1/7*arccosh(c*x)*c^7*x^
7-2/5*arccosh(c*x)*c^5*x^5+1/3*c^3*x^3*arccosh(c*x)-1/11025*(c*x-1)^(1/2)*
(c*x+1)^(1/2)*(225*c^6*x^6-612*c^4*x^4+409*c^2*x^2+818))
```

### 3.12.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.86

$$\int x^2 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{1575 ac^7 d^2 x^7 - 4410 ac^5 d^2 x^5 + 3675 ac^3 d^2 x^3 + 105 (15 bc^7 d^2 x^7 - 42 bc^5 d^2 x^5 + 35 bc^3 d^2 x^3) \log(cx + \sqrt{c^2 x^2 - 1})}{11025 c^3}$$

input `integrate(x^2*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `1/11025*(1575*a*c^7*d^2*x^7 - 4410*a*c^5*d^2*x^5 + 3675*a*c^3*d^2*x^3 + 105*(15*b*c^7*d^2*x^7 - 42*b*c^5*d^2*x^5 + 35*b*c^3*d^2*x^3)*log(c*x + sqrt(c^2*x^2 - 1)) - (225*b*c^6*d^2*x^6 - 612*b*c^4*d^2*x^4 + 409*b*c^2*d^2*x^2 + 818*b*d^2)*sqrt(c^2*x^2 - 1))/c^3`

### 3.12.6 Sympy [F]

$$\int x^2 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = d^2 \left( \int ax^2 dx + \int (-2ac^2 x^4) dx + \int ac^4 x^6 dx \right. \\ \left. + \int bx^2 \operatorname{acosh}(cx) dx \right. \\ \left. + \int (-2bc^2 x^4 \operatorname{acosh}(cx)) dx \right. \\ \left. + \int bc^4 x^6 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x**2*(-c**2*d*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `d**2*(Integral(a*x**2, x) + Integral(-2*a*c**2*x**4, x) + Integral(a*c**4*x**6, x) + Integral(b*x**2*acosh(c*x), x) + Integral(-2*b*c**2*x**4*acosh(c*x), x) + Integral(b*c**4*x**6*acosh(c*x), x))`



### 3.12.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.47

$$\int x^2(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{1}{7} ac^4 d^2 x^7 - \frac{2}{5} ac^2 d^2 x^5 + \frac{1}{245} \left( 35 x^7 \operatorname{arcosh}(cx) - \left( \frac{5 \sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6 \sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16 \sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) bc^4 d^2 - \frac{2}{75} \left( 15 x^5 \operatorname{arcosh}(cx) - \left( \frac{3 \sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4 \sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bc^2 d^2 + \frac{1}{3} ad^2 x^3 + \frac{1}{9} \left( 3 x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2 \sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bd^2$$

input `integrate(x^2*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/7*a*c^4*d^2*x^7 - 2/5*a*c^2*d^2*x^5 + 1/245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*c^4*d^2 - 2/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*c^2*d^2 + 1/3*a*d^2*x^3 + 1/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*d^2`

### 3.12.8 Giac [F(-2)]

Exception generated.

$$\int x^2(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^2*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vcteur & l) Error: Bad Argument Value`

**3.12.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2 dx$$

input `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2,x)`output `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2, x)`

### 3.13 $\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.13.1	Optimal result . . . . .	290
3.13.2	Mathematica [A] (verified) . . . . .	290
3.13.3	Rubi [A] (verified) . . . . .	291
3.13.4	Maple [A] (verified) . . . . .	293
3.13.5	Fricas [A] (verification not implemented) . . . . .	293
3.13.6	Sympy [F] . . . . .	294
3.13.7	Maxima [B] (verification not implemented) . . . . .	294
3.13.8	Giac [F(-2)] . . . . .	295
3.13.9	Mupad [F(-1)] . . . . .	295

#### 3.13.1 Optimal result

Integrand size = 23, antiderivative size = 136

$$\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = -\frac{5bd^2 x \sqrt{-1 + cx} \sqrt{1 + cx}}{96c} + \frac{5bd^2 x (-1 + cx)^{3/2} (1 + cx)^{3/2}}{144c} - \frac{bd^2 x (-1 + cx)^{5/2} (1 + cx)^{5/2}}{36c} + \frac{5bd^2 \operatorname{arccosh}(cx)}{96c^2} - \frac{d^2 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx))}{6c^2}$$

output  $5/144*b*d^2*x*(c*x-1)^{(3/2)}*(c*x+1)^{(3/2)}/c-1/36*b*d^2*x*(c*x-1)^{(5/2)}*(c*x+1)^{(5/2)}/c+5/96*b*d^2*\operatorname{arccosh}(c*x)/c^2-1/6*d^2*(-c^2*x^2+1)^3*(a+b*\operatorname{arccosh}(c*x))/c^2-5/96*b*d^2*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c$

#### 3.13.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.93

$$\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{d^2 \left( cx(b\sqrt{-1 + cx}\sqrt{1 + cx}(-33 + 26c^2x^2 - 8c^4x^4) + 48acx(3 - 3c^2x^2 + c^4x^4)) + 48bc^2x^2(3 - 3c^2x^2 + c^4x^4) \right)}{288c^2}$$

input `Integrate[x*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output  $(d^2*(c*x*(b*\sqrt{-1 + c*x})*\sqrt{1 + c*x}*(-33 + 26*c^2*x^2 - 8*c^4*x^4) + 48*a*c*x*(3 - 3*c^2*x^2 + c^4*x^4)) + 48*b*c^2*x^2*(3 - 3*c^2*x^2 + c^4*x^4)*\text{ArcCosh}[c*x] - 66*b*\text{ArcTanh}[\sqrt{(-1 + c*x)/(1 + c*x)}])/(288*c^2)$

### 3.13.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$ , Rules used = {6329, 40, 40, 40, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x(d - c^2 dx^2)^2 (a + \text{barccosh}(cx)) dx \\
 & \quad \downarrow 6329 \\
 & \frac{bd^2 \int (cx - 1)^{5/2} (cx + 1)^{5/2} dx}{6c} - \frac{d^2 (1 - c^2 x^2)^3 (a + \text{barccosh}(cx))}{6c^2} \\
 & \quad \downarrow 40 \\
 & \frac{bd^2 (\frac{1}{6}x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \int (cx - 1)^{3/2} (cx + 1)^{3/2} dx)}{6c} - \frac{d^2 (1 - c^2 x^2)^3 (a + \text{barccosh}(cx))}{6c^2} \\
 & \quad \downarrow 40 \\
 & \frac{bd^2 (\frac{1}{6}x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} (\frac{1}{4}x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \int \sqrt{cx - 1} \sqrt{cx + 1} dx))}{6c} - \frac{d^2 (1 - c^2 x^2)^3 (a + \text{barccosh}(cx))}{6c^2} \\
 & \quad \downarrow 40 \\
 & \frac{bd^2 (\frac{1}{6}x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} (\frac{1}{4}x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} (\frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{1}{2} \int \frac{1}{\sqrt{cx - 1}\sqrt{cx + 1}} dx)))}{6c} - \frac{d^2 (1 - c^2 x^2)^3 (a + \text{barccosh}(cx))}{6c^2} \\
 & \quad \downarrow 43
 \end{aligned}$$

$$\frac{d^2(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))}{6c^2} - \frac{bd^2\left(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6}\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)\right)}{6c}$$

input `Int[x*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `-1/6*(d^2*(1 - c^2*x^2)^3*(a + b*ArcCosh[c*x]))/c^2 - (b*d^2*((x*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/6 - (5*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/6))/(6*c)`

### 3.13.3.1 Defintions of rubi rules used

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n/(2*e*(p + 1)), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.13.4 Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.25

method	result
derivativedivides	$\frac{\frac{d^2 a (c^2 x^2 - 1)^3}{6} + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^6 x^6}{6} - \frac{c^4 x^4 \operatorname{arccosh}(cx)}{2} + \frac{c^2 x^2 \operatorname{arccosh}(cx)}{2} - \frac{\operatorname{arccosh}(cx)}{6} + \frac{\sqrt{cx-1} \sqrt{cx+1} (-8c^5 x^5 \sqrt{c^2 x^2 - 1} + \dots)}{c^2} \right)}{c^2}$
default	$\frac{\frac{d^2 a (c^2 x^2 - 1)^3}{6} + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^6 x^6}{6} - \frac{c^4 x^4 \operatorname{arccosh}(cx)}{2} + \frac{c^2 x^2 \operatorname{arccosh}(cx)}{2} - \frac{\operatorname{arccosh}(cx)}{6} + \frac{\sqrt{cx-1} \sqrt{cx+1} (-8c^5 x^5 \sqrt{c^2 x^2 - 1} + \dots)}{c^2} \right)}{c^2}$
parts	$\frac{d^2 a (c^2 x^2 - 1)^3}{6c^2} + \frac{d^2 b \left( \frac{\operatorname{arccosh}(cx) c^6 x^6}{6} - \frac{c^4 x^4 \operatorname{arccosh}(cx)}{2} + \frac{c^2 x^2 \operatorname{arccosh}(cx)}{2} - \frac{\operatorname{arccosh}(cx)}{6} + \frac{\sqrt{cx-1} \sqrt{cx+1} (-8c^5 x^5 \sqrt{c^2 x^2 - 1} + \dots)}{c^2} \right)}{c^2}$

input `int(x*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{c^2} \left( \frac{1}{6} d^2 a (c^2 x^2 - 1)^3 + d^2 b \left( \frac{1}{6} \operatorname{arccosh}(cx) c^6 x^6 - \frac{1}{2} c^4 x^4 \operatorname{arccosh}(cx) + \frac{1}{2} c^2 x^2 \operatorname{arccosh}(cx) - \frac{1}{6} \operatorname{arccosh}(cx) + \frac{1}{288} (cx-1)^{1/2} (cx+1)^{1/2} (-8c^5 x^5 \sqrt{c^2 x^2 - 1} + 26(c^2 x^2 - 1)^{1/2} c^3 x^3 - 33cx \sqrt{c^2 x^2 - 1} + 15 \ln(cx + \sqrt{c^2 x^2 - 1})) \right) \right) / (c^2 x^2 - 1)^{1/2} \right)$$

### 3.13.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.10

$$\int x (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \frac{48 ac^6 d^2 x^6 - 144 ac^4 d^2 x^4 + 144 ac^2 d^2 x^2 + 3(16 bc^6 d^2 x^6 - 48 bc^4 d^2 x^4 + 48 bc^2 d^2 x^2 - 11 bd^2) \log(cx + \sqrt{c^2 x^2 - 1})}{288 c^2}$$

input `integrate(x*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output 
$$\frac{1}{288} (48 a c^6 d^2 x^6 - 144 a c^4 d^2 x^4 + 144 a c^2 d^2 x^2 + 3(16 b c^6 d^2 x^6 - 48 b c^4 d^2 x^4 + 48 b c^2 d^2 x^2 - 11 b d^2) \log(cx + \sqrt{c^2 x^2 - 1}) - (8 b c^5 d^2 x^5 - 26 b c^3 d^2 x^3 + 33 b c d^2 x) \sqrt{c^2 x^2 - 1}) / c^2$$

### 3.13.6 Sympy [F]

$$\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = d^2 \left( \int ax dx + \int (-2ac^2 x^3) dx + \int ac^4 x^5 dx \right. \\ \left. + \int bx \operatorname{acosh}(cx) dx + \int (-2bc^2 x^3 \operatorname{acosh}(cx)) dx \right. \\ \left. + \int bc^4 x^5 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x*(-c**2*d*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `d**2*(Integral(a*x, x) + Integral(-2*a*c**2*x**3, x) + Integral(a*c**4*x**5, x) + Integral(b*x*acosh(c*x), x) + Integral(-2*b*c**2*x**3*acosh(c*x), x) + Integral(b*c**4*x**5*acosh(c*x), x))`

### 3.13.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs.  $2(113) = 226$ .

Time = 0.25 (sec) , antiderivative size = 287, normalized size of antiderivative = 2.11

$$\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{1}{6} ac^4 d^2 x^6 - \frac{1}{2} ac^2 d^2 x^4 \\ + \frac{1}{288} \left( 48 x^6 \operatorname{arcosh}(cx) - \left( \frac{8 \sqrt{c^2 x^2 - 1} x^5}{c^2} + \frac{10 \sqrt{c^2 x^2 - 1} x^3}{c^4} + \frac{15 \sqrt{c^2 x^2 - 1} x}{c^6} + \frac{15 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1})}{c^7} \right) \right) \\ - \frac{1}{16} \left( 8 x^4 \operatorname{arcosh}(cx) - \left( \frac{2 \sqrt{c^2 x^2 - 1} x^3}{c^2} + \frac{3 \sqrt{c^2 x^2 - 1} x}{c^4} + \frac{3 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1})}{c^5} \right) \right) c \Big) bc^2 d^2 \\ + \frac{1}{2} ad^2 x^2 + \frac{1}{4} \left( 2 x^2 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x}{c^2} + \frac{\log(2c^2 x + 2\sqrt{c^2 x^2 - 1})}{c^3} \right) \right) bd^2$$

input `integrate(x*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/6*a*c^4*d^2*x^6 - 1/2*a*c^2*d^2*x^4 + 1/288*(48*x^6*arccosh(c*x) - (8*sqrt(c^2*x^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c^4 + 15*sqrt(c^2*x^2 - 1)*x/c^6 + 15*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^7)*c)*b*c^4*d^2 - 1/16*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*c^2*d^2 + 1/2*a*d^2*x^2 + 1/4*(2*x^2*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x/c^2 + log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^3))*b*d^2`

**3.13.8 Giac [F(-2)]**

Exception generated.

$$\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.13.9 Mupad [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2 dx$$

input `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2,x)`

output `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^2, x)`



### 3.14 $\int (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.14.1	Optimal result . . . . .	296
3.14.2	Mathematica [A] (verified) . . . . .	296
3.14.3	Rubi [A] (verified) . . . . .	297
3.14.4	Maple [A] (verified) . . . . .	299
3.14.5	Fricas [A] (verification not implemented) . . . . .	299
3.14.6	Sympy [F] . . . . .	300
3.14.7	Maxima [A] (verification not implemented) . . . . .	300
3.14.8	Giac [F(-2)] . . . . .	301
3.14.9	Mupad [F(-1)] . . . . .	301

#### 3.14.1 Optimal result

Integrand size = 22, antiderivative size = 143

$$\int (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{8bd^2\sqrt{-1+cx}\sqrt{1+cx}}{15c} + \frac{4bd^2(-1+cx)^{3/2}(1+cx)^{3/2}}{45c} - \frac{bd^2(-1+cx)^{5/2}(1+cx)^{5/2}}{25c}$$

$$+ d^2x(a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2d^2x^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}c^4d^2x^5(a + \operatorname{barccosh}(cx))$$

output  $4/45*b*d^2*(c*x-1)^{(3/2)}*(c*x+1)^{(3/2)}/c-1/25*b*d^2*(c*x-1)^{(5/2)}*(c*x+1)^{(5/2)}/c+d^2*x*(a+b*\operatorname{arccosh}(c*x))-2/3*c^2*d^2*x^3*(a+b*\operatorname{arccosh}(c*x))+1/5*c^4*d^2*x^5*(a+b*\operatorname{arccosh}(c*x))-8/15*b*d^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c$

#### 3.14.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.69

$$\int (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{d^2(b\sqrt{-1+cx}\sqrt{1+cx}(-149 + 38c^2x^2 - 9c^4x^4) + 15acx(15 - 10c^2x^2 + 3c^4x^4) + 15bcx(15 - 10c^2x^2 + 3c^4x^4) + 15d^2x^5(a + \operatorname{barccosh}(cx)) - 2c^2d^2x^3(a + \operatorname{barccosh}(cx)))}{225c}$$

input `Integrate[(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output  $(d^2*(b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(-149 + 38*c^2*x^2 - 9*c^4*x^4) + 15*a*c*x*(15 - 10*c^2*x^2 + 3*c^4*x^4) + 15*b*c*x*(15 - 10*c^2*x^2 + 3*c^4*x^4)*\text{ArcCosh}[c*x]))/(225*c)$

### 3.14.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.08, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {6309, 27, 1905, 1576, 1140, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d - c^2 dx^2)^2 (a + \text{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6309} \\
 & -bc \int \frac{d^2 x (3c^4 x^4 - 10c^2 x^2 + 15)}{15\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{5} c^4 d^2 x^5 (a + \text{barccosh}(cx)) - \frac{2}{3} c^2 d^2 x^3 (a + \text{barccosh}(cx)) + \\
 & \quad d^2 x (a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{15} bcd^2 \int \frac{x(3c^4 x^4 - 10c^2 x^2 + 15)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{5} c^4 d^2 x^5 (a + \text{barccosh}(cx)) - \frac{2}{3} c^2 d^2 x^3 (a + \\
 & \quad \text{barccosh}(cx)) + d^2 x (a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{1905} \\
 & -\frac{bcd^2 \sqrt{c^2 x^2 - 1} \int \frac{x(3c^4 x^4 - 10c^2 x^2 + 15)}{\sqrt{c^2 x^2 - 1}} dx}{15\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5} c^4 d^2 x^5 (a + \text{barccosh}(cx)) - \frac{2}{3} c^2 d^2 x^3 (a + \\
 & \quad \text{barccosh}(cx)) + d^2 x (a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{1576} \\
 & -\frac{bcd^2 \sqrt{c^2 x^2 - 1} \int \frac{3c^4 x^4 - 10c^2 x^2 + 15}{\sqrt{c^2 x^2 - 1}} dx^2}{30\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5} c^4 d^2 x^5 (a + \text{barccosh}(cx)) - \frac{2}{3} c^2 d^2 x^3 (a + \text{barccosh}(cx)) + \\
 & \quad d^2 x (a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{1140} \\
 & -\frac{bcd^2 \sqrt{c^2 x^2 - 1} \int \left( 3(c^2 x^2 - 1)^{3/2} - 4\sqrt{c^2 x^2 - 1} + \frac{8}{\sqrt{c^2 x^2 - 1}} \right) dx^2}{30\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5} c^4 d^2 x^5 (a + \text{barccosh}(cx)) - \\
 & \quad \frac{2}{3} c^2 d^2 x^3 (a + \text{barccosh}(cx)) + d^2 x (a + \text{barccosh}(cx))
 \end{aligned}$$

---

3.14.  $\int (d - c^2 dx^2)^2 (a + \text{barccosh}(cx)) dx$

$$\frac{1}{5}c^4d^2x^5(a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2d^2x^3(a + \operatorname{barccosh}(cx)) + d^2x(a + \operatorname{barccosh}(cx)) - \frac{bcd^2\sqrt{c^2x^2-1}\left(\frac{6(c^2x^2-1)^{5/2}}{5c^2} - \frac{8(c^2x^2-1)^{3/2}}{3c^2} + \frac{16\sqrt{c^2x^2-1}}{c^2}\right)}{30\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `-1/30*(b*c*d^2*Sqrt[-1 + c^2*x^2]*((16*Sqrt[-1 + c^2*x^2])/c^2 - (8*(-1 + c^2*x^2)^(3/2))/(3*c^2) + (6*(-1 + c^2*x^2)^(5/2))/(5*c^2)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + d^2*x*(a + b*ArcCosh[c*x]) - (2*c^2*d^2*x^3*(a + b*ArcCosh[c*x]))/3 + (c^4*d^2*x^5*(a + b*ArcCosh[c*x]))/5`

### 3.14.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1140 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 0]`

rule 1576 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`

rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 6309 Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol]
  := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.14.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.69

method	result
parts	$d^2 a \left( \frac{1}{5} c^4 x^5 - \frac{2}{3} x^3 c^2 + x \right) + \frac{d^2 b \left( \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{2c^3 x^3 \operatorname{arccosh}(cx)}{3} + cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1} \sqrt{cx+1} (9c^4 x^4 - 38c^2 x^2 + 149)}{225} \right)}{c}$
derivativedivides	$\frac{d^2 a \left( \frac{1}{5} c^5 x^5 - \frac{2}{3} c^3 x^3 + cx \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{2c^3 x^3 \operatorname{arccosh}(cx)}{3} + cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1} \sqrt{cx+1} (9c^4 x^4 - 38c^2 x^2 + 149)}{225} \right)}{c}$
default	$\frac{d^2 a \left( \frac{1}{5} c^5 x^5 - \frac{2}{3} c^3 x^3 + cx \right) + d^2 b \left( \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{2c^3 x^3 \operatorname{arccosh}(cx)}{3} + cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1} \sqrt{cx+1} (9c^4 x^4 - 38c^2 x^2 + 149)}{225} \right)}{c}$

```
input int((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output d^2*a*(1/5*c^4*x^5-2/3*x^3*c^2+x)+d^2*b/c*(1/5*arccosh(c*x)*c^5*x^5-2/3*c^3*x^3*arccosh(c*x)+c*x*arccosh(c*x)-1/225*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(9*c^4*x^4-38*c^2*x^2+149))
```

### 3.14.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.93

$$\int (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \frac{45 ac^5 d^2 x^5 - 150 ac^3 d^2 x^3 + 225 acd^2 x + 15 (3 bc^5 d^2 x^5 - 10 bc^3 d^2 x^3 + 15 bcd^2 x) \log(cx + \sqrt{c^2 x^2 - 1}) - (9 b^2 c^4 d^2 x^4 - 38 b^2 c^2 d^2 x^2 + 149 b d^2) \sqrt{c^2 x^2 - 1}}{225 c}$$

```
input integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")
```

```
output 1/225*(45*a*c^5*d^2*x^5 - 150*a*c^3*d^2*x^3 + 225*a*c*d^2*x + 15*(3*b*c^5*d^2*x^5 - 10*b*c^3*d^2*x^3 + 15*b*c*d^2*x)*log(c*x + sqrt(c^2*x^2 - 1)) - (9*b*c^4*d^2*x^4 - 38*b*c^2*d^2*x^2 + 149*b*d^2)*sqrt(c^2*x^2 - 1))/c
```

---

3.14.  $\int (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx$

### 3.14.6 Sympy [F]

$$\int (d - c^2 dx^2)^2 (a + \operatorname{arccosh}(cx)) dx = d^2 \left( \int a dx + \int b \operatorname{acosh}(cx) dx + \int (-2ac^2 x^2) dx \right. \\ \left. + \int ac^4 x^4 dx + \int (-2bc^2 x^2 \operatorname{acosh}(cx)) dx \right. \\ \left. + \int bc^4 x^4 \operatorname{acosh}(cx) dx \right)$$

input `integrate((-c**2*d*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `d**2*(Integral(a, x) + Integral(b*acosh(c*x), x) + Integral(-2*a*c**2*x**2, x) + Integral(a*c**4*x**4, x) + Integral(-2*b*c**2*x**2*acosh(c*x), x) + Integral(b*c**4*x**4*acosh(c*x), x))`

### 3.14.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.36

$$\int (d - c^2 dx^2)^2 (a + \operatorname{arccosh}(cx)) dx \\ = \frac{1}{5} ac^4 d^2 x^5 \\ + \frac{1}{75} \left( 15x^5 \operatorname{arccosh}(cx) - \left( \frac{3\sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4\sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8\sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bc^4 d^2 \\ - \frac{2}{3} ac^2 d^2 x^3 - \frac{2}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bc^2 d^2 \\ + ad^2 x + \frac{(cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) bd^2}{c}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/5*a*c^4*d^2*x^5 + 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*c^4*d^2 - 2/3*a*c^2*d^2*x^3 - 2/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*c^2*d^2 + a*d^2*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*d^2/c`

**3.14.8 Giac [F(-2)]**

Exception generated.

$$\int (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.14.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2 dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^2,x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^2, x)`

### 3.15 $\int \frac{(d-c^2dx^2)^2(a+b\operatorname{arccosh}(cx))}{x} dx$

3.15.1	Optimal result	302
3.15.2	Mathematica [A] (warning: unable to verify)	303
3.15.3	Rubi [C] (warning: unable to verify)	303
3.15.4	Maple [A] (verified)	308
3.15.5	Fricas [F]	309
3.15.6	Sympy [F]	309
3.15.7	Maxima [F]	310
3.15.8	Giac [F(-2)]	310
3.15.9	Mupad [F(-1)]	310

#### 3.15.1 Optimal result

Integrand size = 25, antiderivative size = 184

$$\int \frac{(d - c^2dx^2)^2 (a + b\operatorname{arccosh}(cx))}{x} dx = \frac{11}{32}bcd^2x\sqrt{-1 + cx}\sqrt{1 + cx} - \frac{1}{16}bcd^2x(-1 + cx)^{3/2}(1 + cx)^{3/2} - \frac{11}{32}bd^2\operatorname{arccosh}(cx) + \frac{1}{2}d^2(1 - c^2x^2)(a + b\operatorname{arccosh}(cx)) + \frac{1}{4}d^2(1 - c^2x^2)^2(a + b\operatorname{arccosh}(cx)) + \frac{d^2(a + b\operatorname{arccosh}(cx))^2}{2b} + d^2(a + b\operatorname{arccosh}(cx))\log(1 + e^{-2\operatorname{arccosh}(cx)}) - \frac{1}{2}bd^2\operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})$$

output `-1/16*b*c*d^2*x*(c*x-1)^(3/2)*(c*x+1)^(3/2)-11/32*b*d^2*arccosh(c*x)+1/2*d^2*(-c^2*x^2+1)*(a+b*arccosh(c*x))+1/4*d^2*(-c^2*x^2+1)^2*(a+b*arccosh(c*x))+1/2*d^2*(a+b*arccosh(c*x))^2/b+d^2*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)-1/2*b*d^2*polylog(2,-1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)+11/32*b*c*d^2*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)`

### 3.15.2 Mathematica [A] (warning: unable to verify)

Time = 0.49 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.13

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x} dx = \frac{1}{4} d^2 \left( -4ac^2 x^2 + ac^4 x^4 - 4bc^2 x^2 \operatorname{arccosh}(cx) \right. \\ \left. + bc^4 x^4 \operatorname{arccosh}(cx) + 2b \left( cx \sqrt{-1 + cx} \sqrt{1 + cx} \right. \right. \\ \left. \left. + 2 \operatorname{arctanh} \left( \sqrt{\frac{-1 + cx}{1 + cx}} \right) \right) \right. \\ \left. - \frac{1}{8} b \left( cx \sqrt{\frac{-1 + cx}{1 + cx}} (3 + 3cx + 2c^2 x^2 + 2c^3 x^3) \right. \right. \\ \left. \left. + 6 \operatorname{arctanh} \left( \sqrt{\frac{-1 + cx}{1 + cx}} \right) \right) \right. \\ \left. + 2b \operatorname{arccosh}(cx) (\operatorname{arccosh}(cx) \right. \\ \left. + 2 \log(1 + e^{-2 \operatorname{arccosh}(cx)}) \right) + 4a \log(x) \\ \left. - 2b \operatorname{PolyLog}(2, -e^{-2 \operatorname{arccosh}(cx)}) \right)$$

input `Integrate[((d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]))/x,x]`

output `(d^2*(-4*a*c^2*x^2 + a*c^4*x^4 - 4*b*c^2*x^2*ArcCosh[c*x] + b*c^4*x^4*ArcCosh[c*x] + 2*b*(c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x]])) - (b*(c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*(3 + 3*c*x + 2*c^2*x^2 + 2*c^3*x^3) + 6*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x]])))/8 + 2*b*ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) + 4*a*Log[x] - 2*b*PolyLog[2, -E^(-2*ArcCosh[c*x])])/4`

### 3.15.3 Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.98 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.31, number of steps used = 17, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.640$ , Rules used = {6334, 27, 40, 40, 43, 6334, 40, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.15.  $\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x} dx$



$$\begin{aligned}
& \int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx \\
& \quad \downarrow \text{6334} \\
& d \int \frac{d(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4} bcd^2 \int (cx - 1)^{3/2} (cx + 1)^{3/2} dx + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \\
& \quad \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{27} \\
& d^2 \int \frac{(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4} bcd^2 \int (cx - 1)^{3/2} (cx + 1)^{3/2} dx + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \\
& \quad \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{40} \\
& d^2 \int \frac{(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \\
& \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \int \sqrt{cx - 1} \sqrt{cx + 1} dx \right) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{40} \\
& d^2 \int \frac{(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \\
& \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{1}{2} \int \frac{1}{\sqrt{cx - 1} \sqrt{cx + 1}} dx \right) \right) + \\
& \quad \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{43} \\
& d^2 \int \frac{(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \\
& \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \\
& \quad \downarrow \text{6334} \\
& d^2 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2} bc \int \sqrt{cx - 1} \sqrt{cx + 1} dx + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) \right) + \\
& \quad \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \\
& \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \\
& \quad \downarrow \text{40}
\end{aligned}$$

$$d^2 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2} bc \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1} \sqrt{cx+1}} dx \right) + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) - \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx-1)^{3/2} (cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 43

$$d^2 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{2} bc \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx-1)^{3/2} (cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 6297

$$d^2 \left( \frac{\int - \left( (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx-1)^{3/2} (cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 25

$$d^2 \left( - \frac{\int (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx-1)^{3/2} (cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 3042

$$d^2 \left( - \frac{\int -i(a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x (cx-1)^{3/2} (cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 26

---

3.15.  $\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx$

$$d^2 \left( \frac{i \int (a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \downarrow 4201$$

$$d^2 \left( \frac{i \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)} (a + \operatorname{barccosh}(cx))}{1 + e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2} i (a + \operatorname{barccosh}(cx))^2 \right)}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \downarrow 2620$$

$$d^2 \left( \frac{i \left( 2i \left( \frac{1}{2} b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{2} i (a + \operatorname{barccosh}(cx))^2 \right)}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \downarrow 2715$$

$$d^2 \left( \frac{i \left( 2i \left( -\frac{1}{4} b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{2} i (a + \operatorname{barccosh}(cx))^2 \right)}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \downarrow 2838$$

$$d^2 \left( \frac{i \left( 2i \left( \frac{1}{4} b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{2} i (a + \operatorname{barccosh}(cx))^2 \right)}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4} bcd^2 \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

input `Int[((d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]))/x,x]`

output `(d^2*(1 - c^2*x^2)^2*(a + b*ArcCosh[c*x]))/4 - (b*c*d^2*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c))))/4) + d^2*((1 - c^2*x^2)*(a + b*ArcCosh[c*x])/2 + (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c))))/2 + (I*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x]))] + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4)))/b`

### 3.15.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*(c + d*x)^(m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

```
rule 2715 Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
  :-> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
  ))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :-> Simp[-PolyLog[2
  , (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 3042 Int[u_, x_Symbol] :-> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
  Q[u, x]
```

```
rule 4201 Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
  _Symbol] :-> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
  (c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x))))], x], x]
  /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

```
rule 6297 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/(x_), x_Symbol] :-> Simp[1/b
  Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a
  , b, c}, x] && IGtQ[n, 0]
```

```
rule 6334 Int((((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.)/(x_),
  x_Symbol] :-> Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d
  Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d
  )^p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x]) /; FreeQ
  [{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.15.4 Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.04

method	result
parts	$d^2 a \left( \frac{c^4 x^4}{4} - c^2 x^2 + \ln(x) \right) + \frac{d^2 b \operatorname{arccosh}(cx) c^4 x^4}{4} - d^2 b \operatorname{arccosh}(cx) c^2 x^2 + \frac{d^2 b \operatorname{polylog}\left(2, -\frac{cx}{2}\right)}{2}$
derivativedivides	$d^2 a \left( \frac{c^4 x^4}{4} - c^2 x^2 + \ln(cx) \right) + \frac{13b d^2 \operatorname{arccosh}(cx)}{32} + d^2 b \operatorname{arccosh}(cx) \ln\left(1 + (cx + \sqrt{cx - 1})\right)$
default	$d^2 a \left( \frac{c^4 x^4}{4} - c^2 x^2 + \ln(cx) \right) + \frac{13b d^2 \operatorname{arccosh}(cx)}{32} + d^2 b \operatorname{arccosh}(cx) \ln\left(1 + (cx + \sqrt{cx - 1})\right)$

$$3.15. \quad \int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x} dx$$

input `int((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`

output `d^2*a*(1/4*c^4*x^4-c^2*x^2+ln(x))+1/4*d^2*b*arccosh(c*x)*c^4*x^4-d^2*b*arc  
cosh(c*x)*c^2*x^2+1/2*d^2*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2  
)-1/2*d^2*b*arccosh(c*x)^2+13/32*b*d^2*arccosh(c*x)-1/16*d^2*b*(c*x+1)^(1/  
2)*(c*x-1)^(1/2)*c^3*x^3+13/32*b*c*d^2*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)+d^2*b  
*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)`

### 3.15.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(c^2 dx^2 - d)^2 (b \operatorname{arccosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x,x, algorithm="fricas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c  
^2*d^2*x^2 + b*d^2)*arccosh(c*x))/x, x)`

### 3.15.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x} dx = d^2 \left( \int \frac{a}{x} dx + \int (-2ac^2 x) dx + \int ac^4 x^3 dx \right. \\ \left. + \int \frac{b \operatorname{arccosh}(cx)}{x} dx + \int (-2bc^2 x \operatorname{arccosh}(cx)) dx \right. \\ \left. + \int bc^4 x^3 \operatorname{arccosh}(cx) dx \right)$$

input `integrate((-c**2*d*x**2+d)**2*(a+b*acosh(c*x))/x,x)`

output `d**2*(Integral(a/x, x) + Integral(-2*a*c**2*x, x) + Integral(a*c**4*x**3,  
x) + Integral(b*acosh(c*x)/x, x) + Integral(-2*b*c**2*x*acosh(c*x), x) + I  
ntegral(b*c**4*x**3*acosh(c*x), x))`

### 3.15.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(c^2 dx^2 - d)^2 (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `1/4*a*c^4*d^2*x^4 - a*c^2*d^2*x^2 + a*d^2*log(x) + integrate(b*c^4*d^2*x^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) - 2*b*c^2*d^2*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + b*d^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

### 3.15.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.15.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2}{x} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x, x)`

**3.16**  $\int \frac{(d-c^2dx^2)^2(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.16.1 Optimal result . . . . . 311  
 3.16.2 Mathematica [A] (verified) . . . . . 312  
 3.16.3 Rubi [A] (warning: unable to verify) . . . . . 312  
 3.16.4 Maple [A] (verified) . . . . . 315  
 3.16.5 Fricas [A] (verification not implemented) . . . . . 315  
 3.16.6 Sympy [F] . . . . . 316  
 3.16.7 Maxima [A] (verification not implemented) . . . . . 316  
 3.16.8 Giac [F(-2)] . . . . . 317  
 3.16.9 Mupad [F(-1)] . . . . . 317

**3.16.1 Optimal result**

Integrand size = 25, antiderivative size = 135

$$\int \frac{(d - c^2dx^2)^2 (a + \operatorname{arccosh}(cx))}{x^2} dx = \frac{5}{3}bcd^2\sqrt{-1 + cx}\sqrt{1 + cx} - \frac{1}{9}bcd^2(-1 + cx)^{3/2}(1 + cx)^{3/2} - \frac{d^2(a + \operatorname{arccosh}(cx))}{x} - 2c^2d^2x(a + \operatorname{arccosh}(cx)) + \frac{1}{3}c^4d^2x^3(a + \operatorname{arccosh}(cx)) + bcd^2 \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})$$

output

```
-1/9*b*c*d^2*(c*x-1)^(3/2)*(c*x+1)^(3/2)-d^2*(a+b*arccosh(c*x))/x-2*c^2*d^2*x*(a+b*arccosh(c*x))+1/3*c^4*d^2*x^3*(a+b*arccosh(c*x))+b*c*d^2*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))+5/3*b*c*d^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```



### 3.16.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.97

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx = \frac{d^2 \left( -9a - 18ac^2x^2 + 3ac^4x^4 + 16bcx\sqrt{-1+cx}\sqrt{1+cx} - bc^3x^3\sqrt{-1+cx}\sqrt{1+cx} + 3b(-3 - 6c^2x^2 + c^4x^4) \operatorname{ArcCosh}[cx] - 9bcx \operatorname{ArcTan}\left[\frac{1}{\sqrt{-1+cx}\sqrt{1+cx}}\right] \right)}{9x}$$

input `Integrate[((d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]))/x^2,x]`

output `(d^2*(-9*a - 18*a*c^2*x^2 + 3*a*c^4*x^4 + 16*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 3*b*(-3 - 6*c^2*x^2 + c^4*x^4)*ArcCosh[c*x] - 9*b*c*x*ArcTan[1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])]))/(9*x)`

### 3.16.3 Rubi [A] (warning: unable to verify)

Time = 0.53 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$ , Rules used = {6336, 27, 1905, 1578, 1192, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx \\ & \quad \downarrow \text{6336} \\ & -bc \int -\frac{d^2(-c^4x^4 + 6c^2x^2 + 3)}{3x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{3}c^4d^2x^3(a + \operatorname{barccosh}(cx)) - 2c^2d^2x(a + \operatorname{barccosh}(cx)) - \frac{d^2(a + \operatorname{barccosh}(cx))}{x} \\ & \quad \downarrow \text{27} \\ & \frac{1}{3}bcd^2 \int \frac{-c^4x^4 + 6c^2x^2 + 3}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{3}c^4d^2x^3(a + \operatorname{barccosh}(cx)) - 2c^2d^2x(a + \operatorname{barccosh}(cx)) - \frac{d^2(a + \operatorname{barccosh}(cx))}{x} \\ & \quad \downarrow \text{1905} \end{aligned}$$

---

3.16.  $\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx$

$$\frac{bcd^2\sqrt{c^2x^2-1}\int\frac{-c^4x^4+6c^2x^2+3}{x\sqrt{c^2x^2-1}}dx}{3\sqrt{cx-1}\sqrt{cx+1}}+\frac{\frac{1}{3}c^4d^2x^3(a+\operatorname{barccosh}(cx))-2c^2d^2x(a+\operatorname{barccosh}(cx))-\frac{d^2(a+\operatorname{barccosh}(cx))}{x}}{x}}{1578}$$

$$\frac{bcd^2\sqrt{c^2x^2-1}\int\frac{-c^4x^4+6c^2x^2+3}{x^2\sqrt{c^2x^2-1}}dx^2}{6\sqrt{cx-1}\sqrt{cx+1}}+\frac{\frac{1}{3}c^4d^2x^3(a+\operatorname{barccosh}(cx))-2c^2d^2x(a+\operatorname{barccosh}(cx))-\frac{d^2(a+\operatorname{barccosh}(cx))}{x}}{x}}{1192}$$

$$\frac{bd^2\sqrt{c^2x^2-1}\int\frac{-c^4x^8+4c^4x^4+8c^4}{x^4+1}d\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}\sqrt{cx+1}}+\frac{\frac{1}{3}c^4d^2x^3(a+\operatorname{barccosh}(cx))-2c^2d^2x(a+\operatorname{barccosh}(cx))-\frac{d^2(a+\operatorname{barccosh}(cx))}{x}}{x}}{1467}$$

$$\frac{bd^2\sqrt{c^2x^2-1}\int\left(-x^4c^4+\frac{3c^4}{x^4+1}+5c^4\right)d\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}\sqrt{cx+1}}+\frac{\frac{1}{3}c^4d^2x^3(a+\operatorname{barccosh}(cx))-2c^2d^2x(a+\operatorname{barccosh}(cx))-\frac{d^2(a+\operatorname{barccosh}(cx))}{x}}{x}}{2009}$$

$$\frac{\frac{1}{3}c^4d^2x^3(a+\operatorname{barccosh}(cx))-2c^2d^2x(a+\operatorname{barccosh}(cx))-\frac{d^2(a+\operatorname{barccosh}(cx))}{x}}{x}+\frac{bd^2\sqrt{c^2x^2-1}\left(3c^4\arctan\left(\sqrt{c^2x^2-1}\right)-\frac{1}{3}c^4x^6+5c^4\sqrt{c^2x^2-1}\right)}{3c^3\sqrt{cx-1}\sqrt{cx+1}}}{x}$$

input `Int[((d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]))/x^2,x]`

output `-((d^2*(a + b*ArcCosh[c*x]))/x) - 2*c^2*d^2*x*(a + b*ArcCosh[c*x]) + (c^4*d^2*x^3*(a + b*ArcCosh[c*x]))/3 + (b*d^2*Sqrt[-1 + c^2*x^2]*(-1/3*(c^4*x^6) + 5*c^4*Sqrt[-1 + c^2*x^2] + 3*c^4*ArcTan[Sqrt[-1 + c^2*x^2]]))/(3*c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.16.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 1192 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`
- rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`
- rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`
- rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.16.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.03

method	result
parts	$d^2 a \left( \frac{c^4 x^3}{3} - 2cx - \frac{1}{x} \right) + d^2 bc \left( \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - 2cx \operatorname{arccosh}(cx) - \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{cx} \right)$
derivativedivides	$c \left( d^2 a \left( \frac{c^3 x^3}{3} - 2cx - \frac{1}{cx} \right) + d^2 b \left( \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - 2cx \operatorname{arccosh}(cx) - \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{cx} \right) \right)$
default	$c \left( d^2 a \left( \frac{c^3 x^3}{3} - 2cx - \frac{1}{cx} \right) + d^2 b \left( \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - 2cx \operatorname{arccosh}(cx) - \frac{\operatorname{arccosh}(cx)}{cx} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{cx} \right) \right)$

input `int((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)`

output  $d^2 a \left( \frac{1}{3} c^4 x^3 - 2c^2 x - \frac{1}{x} \right) + d^2 b c \left( \frac{1}{3} c^3 x^3 \operatorname{arccosh}(c x) - 2c x \operatorname{arccosh}(c x) - \frac{\operatorname{arccosh}(c x)}{c x} - \frac{\sqrt{c x - 1} \sqrt{c x + 1}}{c x} \right) + 9 \arctan \left( \frac{1}{(c^2 x^2 - 1)^{1/2}} \right) - 16 \frac{(c^2 x^2 - 1)^{1/2}}{(c^2 x^2 - 1)^{1/2}}$

### 3.16.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 201, normalized size of antiderivative = 1.49

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx$$

$$= \frac{3 a c^4 d^2 x^4 - 18 a c^2 d^2 x^2 + 18 b c d^2 x \arctan(-cx + \sqrt{c^2 x^2 - 1}) - 3 (b c^4 - 6 b c^2 - 3 b) d^2 x \log(-cx + \sqrt{c^2 x^2 - 1})}{x^2}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x, algorithm="fricas")`

output  $\frac{1}{9} (3 a c^4 d^2 x^4 - 18 a c^2 d^2 x^2 + 18 b c d^2 x \arctan(-c x + \sqrt{c^2 x^2 - 1})) - 3 (b c^4 - 6 b c^2 - 3 b) d^2 x \log(-c x + \sqrt{c^2 x^2 - 1}) - 9 a d^2 + 3 (b c^4 d^2 x^4 - 6 b c^2 d^2 x^2 - (b c^4 - 6 b c^2 - 3 b) d^2 x - 3 b d^2) \log(c x + \sqrt{c^2 x^2 - 1}) - (b c^3 d^2 x^3 - 16 b c d^2 x) \sqrt{c^2 x^2 - 1} / x$

---

3.16.  $\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx$

### 3.16.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx = d^2 \left( \int (-2ac^2) dx + \int \frac{a}{x^2} dx + \int ac^4 x^2 dx \right. \\ \left. + \int (-2bc^2 \operatorname{acosh}(cx)) dx + \int \frac{b \operatorname{acosh}(cx)}{x^2} dx \right. \\ \left. + \int bc^4 x^2 \operatorname{acosh}(cx) dx \right)$$

input `integrate((-c**2*d*x**2+d)**2*(a+b*acosh(c*x))/x**2,x)`

output `d**2*(Integral(-2*a*c**2, x) + Integral(a/x**2, x) + Integral(a*c**4*x**2, x) + Integral(-2*b*c**2*acosh(c*x), x) + Integral(b*acosh(c*x)/x**2, x) + Integral(b*c**4*x**2*acosh(c*x), x))`

### 3.16.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.06

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx \\ = \frac{1}{3} ac^4 d^2 x^3 + \frac{1}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bc^4 d^2 - 2ac^2 d^2 x \\ - 2 \left( cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1} \right) bcd^2 - \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) bd^2 - \frac{ad^2}{x}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output `1/3*a*c^4*d^2*x^3 + 1/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*c^4*d^2 - 2*a*c^2*d^2*x - 2*(c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*c*d^2 - (c*arcsin(1/(c*abs(x))) + arccosh(c*x)/x)*b*d^2 - a*d^2/x`

**3.16.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.16.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x^2,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x^2, x)`

### 3.17 $\int \frac{(d-c^2dx^2)^2(a+b\operatorname{arccosh}(cx))}{x^3} dx$

3.17.1	Optimal result	318
3.17.2	Mathematica [A] (verified)	319
3.17.3	Rubi [C] (warning: unable to verify)	319
3.17.4	Maple [A] (verified)	325
3.17.5	Fricas [F]	326
3.17.6	Sympy [F]	326
3.17.7	Maxima [F]	327
3.17.8	Giac [F(-2)]	327
3.17.9	Mupad [F(-1)]	327

#### 3.17.1 Optimal result

Integrand size = 25, antiderivative size = 200

$$\int \frac{(d - c^2dx^2)^2 (a + b\operatorname{arccosh}(cx))}{x^3} dx = \frac{1}{4}bc^3d^2x\sqrt{-1 + cx}\sqrt{1 + cx} - \frac{bcd^2(-1 + cx)^{3/2}(1 + cx)^{3/2}}{2x} - \frac{1}{4}bc^2d^2\operatorname{arccosh}(cx) - c^2d^2(1 - c^2x^2)(a + b\operatorname{arccosh}(cx)) - \frac{d^2(1 - c^2x^2)^2(a + b\operatorname{arccosh}(cx))}{2x^2} - \frac{c^2d^2(a + b\operatorname{arccosh}(cx))^2}{b} - 2c^2d^2(a + b\operatorname{arccosh}(cx))\log(1 + e^{-2\operatorname{arccosh}(cx)}) + bc^2d^2\operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})$$

output

```
-1/2*b*c*d^2*(c*x-1)^(3/2)*(c*x+1)^(3/2)/x-1/4*b*c^2*d^2*arccosh(c*x)-c^2*d^2*(-c^2*x^2+1)*(a+b*arccosh(c*x))-1/2*d^2*(-c^2*x^2+1)^2*(a+b*arccosh(c*x))/x^2-c^2*d^2*(a+b*arccosh(c*x))^2/b-2*c^2*d^2*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)+b*c^2*d^2*polylog(2,-1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)+1/4*b*c^3*d^2*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

### 3.17.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.91

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx$$

$$= \frac{d^2 \left( -2a + 2ac^4 x^4 + 2bcx\sqrt{-1 + cx}\sqrt{1 + cx} - bc^3 x^3\sqrt{-1 + cx}\sqrt{1 + cx} - 4bc^2 x^2 \operatorname{arccosh}(cx)^2 - 2bc^2 x^2 \operatorname{arccosh}(cx) \right)}{4x^2}$$

input `Integrate[((d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]))/x^3,x]`

output `(d^2*(-2*a + 2*a*c^4*x^4 + 2*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 4*b*c^2*x^2*ArcCosh[c*x]^2 - 2*b*c^2*x^2*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]] + 2*b*ArcCosh[c*x]*(-1 + c^4*x^4 - 4*c^2*x^2*Log[1 + E^(-2*ArcCosh[c*x])]) - 8*a*c^2*x^2*Log[x] + 4*b*c^2*x^2*PolyLog[2, -E^(-2*ArcCosh[c*x])]))/(4*x^2)`

### 3.17.3 Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.03 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.24, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.680$ , Rules used = {6335, 27, 108, 27, 40, 43, 6334, 40, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx$$

$$\downarrow 6335$$

$$-2c^2 d \int \frac{d(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{2} bcd^2 \int \frac{(cx - 1)^{3/2} (cx + 1)^{3/2}}{x^2} dx -$$

$$\frac{d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{2x^2}$$

$$\downarrow 27$$



$$\begin{aligned}
& -2c^2d^2 \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx + \frac{1}{2}bcd^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x^2} dx - \\
& \quad \frac{d^2(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow 108 \\
& -2c^2d^2 \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx + \\
& \frac{1}{2}bcd^2 \left( \int 3c^2\sqrt{cx-1}\sqrt{cx+1} dx - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) - \frac{d^2(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow 27 \\
& -2c^2d^2 \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx + \\
& \frac{1}{2}bcd^2 \left( 3c^2 \int \sqrt{cx-1}\sqrt{cx+1} dx - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) - \frac{d^2(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow 40 \\
& -2c^2d^2 \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx + \\
& \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) - \\
& \quad \frac{d^2(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow 43 \\
& -2c^2d^2 \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx - \frac{d^2(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2} + \\
& \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \\
& \quad \downarrow 6334 \\
& -2c^2d^2 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}bc \int \sqrt{cx-1}\sqrt{cx+1} dx + \frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx)) \right) - \\
& \quad \frac{d^2(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2} + \\
& \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \\
& \quad \downarrow 40
\end{aligned}$$

$$-2c^2 d^2 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2} bc \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1} \sqrt{cx+1}} dx \right) + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) \right. \\ \left. + \frac{d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2} bcd^2 \left( 3c^2 \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2} (cx+1)^{3/2}}{x} \right) \right)$$

↓ 43

$$-2c^2 d^2 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{2} bc \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \\ \left. + \frac{d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2} bcd^2 \left( 3c^2 \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2} (cx+1)^{3/2}}{x} \right) \right)$$

↓ 6297

$$-2c^2 d^2 \left( \frac{\int - \left( (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) \right) \\ \left. + \frac{d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2} bcd^2 \left( 3c^2 \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2} (cx+1)^{3/2}}{x} \right) \right)$$

↓ 25

$$-2c^2 d^2 \left( - \frac{\int (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2} (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) \right) \\ \left. + \frac{d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2} bcd^2 \left( 3c^2 \left( \frac{1}{2} x \sqrt{cx-1} \sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2} (cx+1)^{3/2}}{x} \right) \right)$$

↓ 3042

$$-2c^2 d^2 \left( \frac{\int -i(a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2}(1 - c^2 x^2)(a + \operatorname{barccosh}(cx)) \right. \\ \left. + \frac{d^2(1 - c^2 x^2)^2(a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \right)$$

↓ 26

$$-2c^2 d^2 \left( \frac{i \int (a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{2}(1 - c^2 x^2)(a + \operatorname{barccosh}(cx)) \right. \\ \left. + \frac{d^2(1 - c^2 x^2)^2(a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \right)$$

↓ 4201

$$-2c^2 d^2 \left( \frac{i \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 + e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2 \right)}{b} + \frac{1}{2}(1 - c^2 x^2)(a + \operatorname{barccosh}(cx)) \right. \\ \left. + \frac{d^2(1 - c^2 x^2)^2(a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \right)$$

↓ 2620

$$-2c^2 d^2 \left( \frac{i \left( 2i \left( \frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)) \right) \right)}{b} + \frac{1}{2}(1 - c^2 x^2)(a + \operatorname{barccosh}(cx)) \right. \\ \left. + \frac{d^2(1 - c^2 x^2)^2(a + \operatorname{barccosh}(cx))}{2x^2} + \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \right)$$

↓ 2715

$$\begin{aligned}
& -2c^2 d^2 \left( \frac{i(2i(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{arccosh}(cx)))}{b} \right. \\
& \quad \left. + \frac{d^2(1 - c^2x^2)^2(a + \operatorname{arccosh}(cx))}{2x^2} + \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \right) \\
& \quad \downarrow \text{2838} \\
& -2c^2 d^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{arccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{arccosh}(cx))) - \frac{1}{2}i(a + \operatorname{arccosh}(cx)))}{b} \right. \\
& \quad \left. + \frac{d^2(1 - c^2x^2)^2(a + \operatorname{arccosh}(cx))}{2x^2} + \frac{1}{2}bcd^2 \left( 3c^2 \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) - \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x} \right) \right)
\end{aligned}$$

input `Int[(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x])/x^3,x]`

output `-1/2*(d^2*(1 - c^2*x^2)^2*(a + b*ArcCosh[c*x])/x^2 + (b*c*d^2*(-((( -1 + c*x)^(3/2)*(1 + c*x)^(3/2))/x) + 3*c^2*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c))))/2 - 2*c^2*d^2(((1 - c^2*x^2)*(a + b*ArcCosh[c*x])/2 + (b*c*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/2 + (I*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x]])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]]/4)))/b)`

### 3.17.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

---

3.17.  $\int \frac{(d-c^2dx^2)^2(a+b\operatorname{arccosh}(cx))}{x^3} dx$

- rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*(c + d*x)^m/(2*m + 1), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 108 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(m + 1))), x] - Simp[1/(b*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6334 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.)/(x_), x_Symbol] := Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d)^(p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6335 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_)^(m_))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (-Simp[b*c*((-d)^p/(f*(m + 1))) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x] - Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0] && ILtQ[(m + 1)/2, 0]`

### 3.17.4 Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.03

method	result
derivativedivides	$c^2 \left( d^2 a \left( \frac{c^2 x^2}{2} - 2 \ln(cx) - \frac{1}{2c^2 x^2} \right) + d^2 b \operatorname{arccosh}(cx)^2 - \frac{bc d^2 x \sqrt{cx-1} \sqrt{cx+1}}{4} + \frac{d^2 b \operatorname{arccosh}(cx) c^2}{2} \right)$
default	$c^2 \left( d^2 a \left( \frac{c^2 x^2}{2} - 2 \ln(cx) - \frac{1}{2c^2 x^2} \right) + d^2 b \operatorname{arccosh}(cx)^2 - \frac{bc d^2 x \sqrt{cx-1} \sqrt{cx+1}}{4} + \frac{d^2 b \operatorname{arccosh}(cx) c^2}{2} \right)$
parts	$d^2 a \left( \frac{c^4 x^2}{2} - \frac{1}{2x^2} - 2c^2 \ln(x) \right) + d^2 b c^2 \operatorname{arccosh}(cx)^2 + \frac{d^2 b c^4 \operatorname{arccosh}(cx) x^2}{2} - \frac{b c^3 d^2 x \sqrt{cx-1} \sqrt{cx+1}}{4}$

input `int((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)`

$$3.17. \int \frac{(d-c^2 dx^2)^2 (a+b \operatorname{arccosh}(cx))}{x^3} dx$$

output  $c^2(d^2a*(1/2*c^2*x^2-2*\ln(cx)-1/2/c^2/x^2)+d^2*b*\operatorname{arccosh}(cx)^2-1/4*b*c*d^2*x*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}+1/2*d^2*b*\operatorname{arccosh}(cx)*c^2*x^2-1/4*b*d^2*\operatorname{arccosh}(cx)-1/2*d^2*b+1/2*d^2*b/c/x*(cx+1)^{(1/2)}*(cx-1)^{(1/2)}-1/2*d^2*b*\operatorname{arccosh}(cx)/c^2/x^2-2*d^2*b*\operatorname{arccosh}(cx)*\ln(1+(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2}))^2)-d^2*b*\operatorname{polylog}(2,-(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2}))^2))$

### 3.17.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(c^2 dx^2 - d)^2 (b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(cx))/x^3,x, algorithm="fricas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(cx))/x^3, x)`

### 3.17.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^3} dx = d^2 \left( \int \frac{a}{x^3} dx + \int \left( -\frac{2ac^2}{x} \right) dx + \int ac^4 x dx + \int \frac{b \operatorname{arccosh}(cx)}{x^3} dx + \int \left( -\frac{2bc^2 \operatorname{arccosh}(cx)}{x} \right) dx + \int bc^4 x \operatorname{arccosh}(cx) dx \right)$$

input `integrate((-c**2*d*x**2+d)**2*(a+b*acosh(cx))/x**3,x)`

output `d**2*(Integral(a/x**3, x) + Integral(-2*a*c**2/x, x) + Integral(a*c**4*x, x) + Integral(b*acosh(cx)/x**3, x) + Integral(-2*b*c**2*acosh(cx)/x, x) + Integral(b*c**4*x*acosh(cx), x))`

**3.17.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(c^2 dx^2 - d)^2 (b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `1/2*a*c^4*d^2*x^2 - 2*a*c^2*d^2*log(x) + 1/2*b*d^2*(sqrt(c^2*x^2 - 1)*c/x - arccosh(c*x)/x^2) - 1/2*a*d^2/x^2 + integrate(b*c^4*d^2*x*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) - 2*b*c^2*d^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x, x)`

**3.17.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.17.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x^3,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x^3, x)`

---

3.17.  $\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx$



**3.18**  $\int \frac{(d-c^2dx^2)^2(a+b\operatorname{arccosh}(cx))}{x^4} dx$

3.18.1	Optimal result . . . . .	328
3.18.2	Mathematica [A] (verified) . . . . .	328
3.18.3	Rubi [A] (warning: unable to verify) . . . . .	329
3.18.4	Maple [A] (verified) . . . . .	332
3.18.5	Fricas [A] (verification not implemented) . . . . .	333
3.18.6	Sympy [F] . . . . .	334
3.18.7	Maxima [A] (verification not implemented) . . . . .	334
3.18.8	Giac [F(-2)] . . . . .	335
3.18.9	Mupad [F(-1)] . . . . .	335

**3.18.1 Optimal result**

Integrand size = 25, antiderivative size = 142

$$\int \frac{(d - c^2dx^2)^2(a + \operatorname{arccosh}(cx))}{x^4} dx = -bc^3d^2\sqrt{-1 + cx}\sqrt{1 + cx} + \frac{bcd^2\sqrt{-1 + cx}\sqrt{1 + cx}}{6x^2} - \frac{d^2(a + \operatorname{arccosh}(cx))}{3x^3} + \frac{2c^2d^2(a + \operatorname{arccosh}(cx))}{x} + c^4d^2x(a + \operatorname{arccosh}(cx)) - \frac{11}{6}bc^3d^2 \arctan\left(\sqrt{-1 + cx}\sqrt{1 + cx}\right)$$

```
output -1/3*d^2*(a+b*arccosh(c*x))/x^3+2*c^2*d^2*(a+b*arccosh(c*x))/x+c^4*d^2*x*(
a+b*arccosh(c*x))-11/6*b*c^3*d^2*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))-b*c^3
*d^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)+1/6*b*c*d^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x
^2
```

**3.18.2 Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.95

$$\int \frac{(d - c^2dx^2)^2(a + \operatorname{arccosh}(cx))}{x^4} dx = \frac{d^2\left(-2a + 12ac^2x^2 + 6ac^4x^4 + bcx\sqrt{-1 + cx}\sqrt{1 + cx} - 6bc^3x^3\sqrt{-1 + cx}\sqrt{1 + cx} + 2b(-1 + 6c^2x^2 + 3c^4x^4)\right)}{6x^3}$$

input `Integrate[((d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]))/x^4,x]`

output `(d^2*(-2*a + 12*a*c^2*x^2 + 6*a*c^4*x^4 + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 6*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*b*(-1 + 6*c^2*x^2 + 3*c^4*x^4)*ArcCosh[c*x] + 11*b*c^3*x^3*ArcTan[1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])])/(6*x^3)`

### 3.18.3 Rubi [A] (warning: unable to verify)

Time = 0.52 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.15, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$ , Rules used = {6336, 27, 1905, 1578, 1192, 25, 1471, 25, 27, 299, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx \\
 & \quad \downarrow \text{6336} \\
 & -bc \int -\frac{d^2(-3c^4x^4 - 6c^2x^2 + 1)}{3x^3\sqrt{cx - 1}\sqrt{cx + 1}} dx + c^4 d^2 x(a + \operatorname{barccosh}(cx)) + \frac{2c^2 d^2 (a + \operatorname{barccosh}(cx))}{x} - \\
 & \quad \frac{d^2(a + \operatorname{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{3}bcd^2 \int \frac{-3c^4x^4 - 6c^2x^2 + 1}{x^3\sqrt{cx - 1}\sqrt{cx + 1}} dx + c^4 d^2 x(a + \operatorname{barccosh}(cx)) + \frac{2c^2 d^2 (a + \operatorname{barccosh}(cx))}{x} - \\
 & \quad \frac{d^2(a + \operatorname{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{1905} \\
 & \frac{bcd^2\sqrt{c^2x^2 - 1} \int \frac{-3c^4x^4 - 6c^2x^2 + 1}{x^3\sqrt{c^2x^2 - 1}} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + c^4 d^2 x(a + \operatorname{barccosh}(cx)) + \frac{2c^2 d^2 (a + \operatorname{barccosh}(cx))}{x} - \\
 & \quad \frac{d^2(a + \operatorname{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{1578} \\
 & \frac{bcd^2\sqrt{c^2x^2 - 1} \int \frac{-3c^4x^4 - 6c^2x^2 + 1}{x^4\sqrt{c^2x^2 - 1}} dx^2}{6\sqrt{cx - 1}\sqrt{cx + 1}} + c^4 d^2 x(a + \operatorname{barccosh}(cx)) + \frac{2c^2 d^2 (a + \operatorname{barccosh}(cx))}{x} - \\
 & \quad \frac{d^2(a + \operatorname{barccosh}(cx))}{3x^3}
 \end{aligned}$$

---

3.18.  $\int \frac{(d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx$

$$\begin{aligned}
& \downarrow 1192 \\
& \frac{bd^2\sqrt{c^2x^2-1} \int -\frac{3c^4x^8+12c^4x^4+8c^4}{(x^4+1)^2} d\sqrt{c^2x^2-1}}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{2c^2d^2(a+\operatorname{barccosh}(cx))} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3}} + c^4d^2x(a+\operatorname{barccosh}(cx)) + \\
& \downarrow 25 \\
& -\frac{bd^2\sqrt{c^2x^2-1} \int \frac{3c^4x^8+12c^4x^4+8c^4}{(x^4+1)^2} d\sqrt{c^2x^2-1}}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{2c^2d^2(a+\operatorname{barccosh}(cx))} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3}} + c^4d^2x(a+\operatorname{barccosh}(cx)) + \\
& \downarrow 1471 \\
& \frac{bd^2\sqrt{c^2x^2-1} \left( \frac{1}{2} \int -\frac{c^4(6x^4+17)}{x^4+1} d\sqrt{c^2x^2-1} + \frac{c^4\sqrt{c^2x^2-1}}{2(x^4+1)} \right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{2c^2d^2(a+\operatorname{barccosh}(cx))} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3}} + c^4d^2x(a+\operatorname{barccosh}(cx)) + \\
& \downarrow 25 \\
& \frac{bd^2\sqrt{c^2x^2-1} \left( \frac{c^4\sqrt{c^2x^2-1}}{2(x^4+1)} - \frac{1}{2} \int \frac{c^4(6x^4+17)}{x^4+1} d\sqrt{c^2x^2-1} \right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{2c^2d^2(a+\operatorname{barccosh}(cx))} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3}} + c^4d^2x(a+\operatorname{barccosh}(cx)) + \\
& \downarrow 27 \\
& \frac{bd^2\sqrt{c^2x^2-1} \left( \frac{c^4\sqrt{c^2x^2-1}}{2(x^4+1)} - \frac{1}{2}c^4 \int \frac{6x^4+17}{x^4+1} d\sqrt{c^2x^2-1} \right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{2c^2d^2(a+\operatorname{barccosh}(cx))} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3}} + c^4d^2x(a+\operatorname{barccosh}(cx)) + \\
& \downarrow 299 \\
& \frac{bd^2\sqrt{c^2x^2-1} \left( \frac{c^4\sqrt{c^2x^2-1}}{2(x^4+1)} - \frac{1}{2}c^4 \left( 11 \int \frac{1}{x^4+1} d\sqrt{c^2x^2-1} + 6\sqrt{c^2x^2-1} \right) \right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{2c^2d^2(a+\operatorname{barccosh}(cx))} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3}} + c^4d^2x(a+\operatorname{barccosh}(cx)) + \\
& \downarrow 216 \\
& c^4d^2x(a+\operatorname{barccosh}(cx)) + \frac{2c^2d^2(a+\operatorname{barccosh}(cx))}{x} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} + \\
& \frac{bd^2\sqrt{c^2x^2-1} \left( \frac{c^4\sqrt{c^2x^2-1}}{2(x^4+1)} - \frac{1}{2}c^4 \left( 11 \arctan \left( \sqrt{c^2x^2-1} \right) + 6\sqrt{c^2x^2-1} \right) \right)}{3c\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

---

3.18.  $\int \frac{(d-c^2dx^2)^2(a+\operatorname{barccosh}(cx))}{x^4} dx$

input `Int[((d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*(d^2*(a + b*ArcCosh[c*x]))/x^3 + (2*c^2*d^2*(a + b*ArcCosh[c*x]))/x + c^4*d^2*x*(a + b*ArcCosh[c*x]) + (b*d^2*Sqrt[-1 + c^2*x^2]*((c^4*Sqrt[-1 + c^2*x^2]))/(2*(1 + x^4)) - (c^4*(6*Sqrt[-1 + c^2*x^2] + 11*ArcTan[Sqrt[-1 + c^2*x^2]]))/2)/(3*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.18.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 1192 `Int[((d_.) + (e_.)*(x_)^(m_))*((f_.) + (g_.)*(x_)^(n_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1471 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Simp[1/(2*d*(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]`

rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 1905 `Int[((f_)*(x_)^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`

rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_)), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.18.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.06

---

3.18.  $\int \frac{(d-c^2dx^2)^2(a+b\operatorname{arccosh}(cx))}{x^4} dx$

method	result
parts	$d^2a\left(c^4x + \frac{2c^2}{x} - \frac{1}{3x^3}\right) + d^2b c^3\left(cx \operatorname{arccosh}(cx) - \frac{\operatorname{arccosh}(cx)}{3c^3x^3} + \frac{2 \operatorname{arccosh}(cx)}{cx} + \frac{\sqrt{cx-1}\sqrt{cx+1}}{cx}\right)$
derivativedivides	$c^3\left(d^2a\left(cx - \frac{1}{3c^3x^3} + \frac{2}{cx}\right) + d^2b\left(cx \operatorname{arccosh}(cx) - \frac{\operatorname{arccosh}(cx)}{3c^3x^3} + \frac{2 \operatorname{arccosh}(cx)}{cx} + \frac{\sqrt{cx-1}\sqrt{cx+1}}{cx}\right)\right)$
default	$c^3\left(d^2a\left(cx - \frac{1}{3c^3x^3} + \frac{2}{cx}\right) + d^2b\left(cx \operatorname{arccosh}(cx) - \frac{\operatorname{arccosh}(cx)}{3c^3x^3} + \frac{2 \operatorname{arccosh}(cx)}{cx} + \frac{\sqrt{cx-1}\sqrt{cx+1}}{cx}\right)\right)$

input `int((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)`

output  $d^2a*(c^4*x+2*c^2/x-1/3/x^3)+d^2*b*c^3*(c*x*\operatorname{arccosh}(c*x)-1/3/c^3/x^3*\operatorname{arccosh}(c*x)+2*\operatorname{arccosh}(c*x)/c/x+1/6*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(11*\arctan(1/(c^2*x^2-1)^{(1/2)})*c^2*x^2-6*c^2*x^2*(c^2*x^2-1)^{(1/2)}+(c^2*x^2-1)^{(1/2)})/c^2/x^2/(c^2*x^2-1)^{(1/2)})$

### 3.18.5 Fricas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.50

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^4} dx$$

$$= \frac{6ac^4d^2x^4 - 22bc^3d^2x^3 \arctan(-cx + \sqrt{c^2x^2 - 1}) + 12ac^2d^2x^2 - 2(3bc^4 + 6bc^2 - b)d^2x^3 \log(-cx + \sqrt{c^2x^2 - 1})}{x^4}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")`

output  $1/6*(6*a*c^4*d^2*x^4 - 22*b*c^3*d^2*x^3*\arctan(-c*x + \sqrt{c^2*x^2 - 1}) + 12*a*c^2*d^2*x^2 - 2*(3*b*c^4 + 6*b*c^2 - b)*d^2*x^3*\log(-c*x + \sqrt{c^2*x^2 - 1}) - 2*a*d^2 + 2*(3*b*c^4*d^2*x^4 + 6*b*c^2*d^2*x^2 - (3*b*c^4 + 6*b*c^2 - b)*d^2*x^3 - b*d^2)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (6*b*c^3*d^2*x^3 - b*c*d^2*x)*\sqrt{c^2*x^2 - 1})/x^3$

---

3.18.  $\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^4} dx$

### 3.18.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^4} dx = d^2 \left( \int ac^4 dx + \int \frac{a}{x^4} dx + \int \left( -\frac{2ac^2}{x^2} \right) dx \right. \\ \left. + \int bc^4 \operatorname{acosh}(cx) dx + \int \frac{b \operatorname{acosh}(cx)}{x^4} dx \right. \\ \left. + \int \left( -\frac{2bc^2 \operatorname{acosh}(cx)}{x^2} \right) dx \right)$$

input `integrate((-c**2*d*x**2+d)**2*(a+b*acosh(c*x))/x**4,x)`

output `d**2*(Integral(a*c**4, x) + Integral(a/x**4, x) + Integral(-2*a*c**2/x**2, x) + Integral(b*c**4*acosh(c*x), x) + Integral(b*acosh(c*x)/x**4, x) + Integral(-2*b*c**2*acosh(c*x)/x**2, x))`

### 3.18.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.96

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^4} dx \\ = ac^4 d^2 x + (cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) bc^3 d^2 + 2 \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) bc^2 d^2 \\ - \frac{1}{6} \left( \left( c^2 \arcsin \left( \frac{1}{c|x|} \right) - \frac{\sqrt{c^2 x^2 - 1}}{x^2} \right) c + \frac{2 \operatorname{arccosh}(cx)}{x^3} \right) bd^2 + \frac{2ac^2 d^2}{x} - \frac{ad^2}{3x^3}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")`

output `a*c^4*d^2*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*c^3*d^2 + 2*(c*arcsin(1/(c*abs(x))) + arccosh(c*x)/x)*b*c^2*d^2 - 1/6*((c^2*arcsin(1/(c*abs(x)))) - sqrt(c^2*x^2 - 1)/x^2)*c + 2*arccosh(c*x)/x^3)*b*d^2 + 2*a*c^2*d^2/x - 1/3*a*d^2/x^3`

**3.18.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.18.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x^4,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^2)/x^4, x)`



### 3.19 $\int x^4(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.19.1	Optimal result . . . . .	336
3.19.2	Mathematica [A] (verified) . . . . .	337
3.19.3	Rubi [A] (verified) . . . . .	337
3.19.4	Maple [A] (verified) . . . . .	340
3.19.5	Fricas [A] (verification not implemented) . . . . .	340
3.19.6	Sympy [F(-1)] . . . . .	341
3.19.7	Maxima [B] (verification not implemented) . . . . .	341
3.19.8	Giac [F(-2)] . . . . .	342
3.19.9	Mupad [F(-1)] . . . . .	342

#### 3.19.1 Optimal result

Integrand size = 25, antiderivative size = 256

$$\int x^4(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{16bd^3\sqrt{-1+cx}\sqrt{1+cx}}{1155c^5} + \frac{8bd^3(-1+cx)^{3/2}(1+cx)^{3/2}}{3465c^5}$$

$$- \frac{2bd^3(-1+cx)^{5/2}(1+cx)^{5/2}}{1925c^5} + \frac{bd^3(-1+cx)^{7/2}(1+cx)^{7/2}}{1617c^5}$$

$$+ \frac{4bd^3(-1+cx)^{9/2}(1+cx)^{9/2}}{297c^5} + \frac{bd^3(-1+cx)^{11/2}(1+cx)^{11/2}}{121c^5}$$

$$+ \frac{1}{5}d^3x^5(a + \operatorname{barccosh}(cx)) - \frac{3}{7}c^2d^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{3}c^4d^3x^9(a + \operatorname{barccosh}(cx)) - \frac{1}{11}c^6d^3x^{11}(a + \operatorname{barccosh}(cx))$$

```
output 8/3465*b*d^3*(c*x-1)^(3/2)*(c*x+1)^(3/2)/c^5-2/1925*b*d^3*(c*x-1)^(5/2)*(c
*x+1)^(5/2)/c^5+1/1617*b*d^3*(c*x-1)^(7/2)*(c*x+1)^(7/2)/c^5+4/297*b*d^3*(
c*x-1)^(9/2)*(c*x+1)^(9/2)/c^5+1/121*b*d^3*(c*x-1)^(11/2)*(c*x+1)^(11/2)/c
^5+1/5*d^3*x^5*(a+b*arccosh(c*x))-3/7*c^2*d^3*x^7*(a+b*arccosh(c*x))+1/3*c
^4*d^3*x^9*(a+b*arccosh(c*x))-1/11*c^6*d^3*x^11*(a+b*arccosh(c*x))-16/1155
*b*d^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5
```

### 3.19.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.57

$$\int x^4 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \frac{d^3 (3465ac^5 x^5 (-231 + 495c^2 x^2 - 385c^4 x^4 + 105c^6 x^6) + b\sqrt{-1 + cx}\sqrt{1 + cx}(50488 + 25244c^2 x^2 + 18933c^4 x^4 - 117625c^6 x^6 + 111475c^8 x^8 - 33075c^{10} x^{10}) + 3465b^2 c^5 x^5 (-231 + 495c^2 x^2 - 385c^4 x^4 + 105c^6 x^6) \operatorname{ArcCosh}[cx])}{c^5}$$

input `Integrate[x^4*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/4002075*(d^3*(3465*a*c^5*x^5*(-231 + 495*c^2*x^2 - 385*c^4*x^4 + 105*c^6*x^6) + b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(50488 + 25244*c^2*x^2 + 18933*c^4*x^4 - 117625*c^6*x^6 + 111475*c^8*x^8 - 33075*c^10*x^10) + 3465*b*c^5*x^5*(-231 + 495*c^2*x^2 - 385*c^4*x^4 + 105*c^6*x^6)*ArcCosh[c*x]))/c^5`

### 3.19.3 Rubi [A] (verified)

Time = 0.79 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6336, 27, 2113, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^4 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6336} \\ & -bc \int \frac{d^3 x^5 (-105c^6 x^6 + 385c^4 x^4 - 495c^2 x^2 + 231)}{1155\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{11} c^6 d^3 x^{11} (a + \operatorname{barccosh}(cx)) + \\ & \quad \frac{1}{3} c^4 d^3 x^9 (a + \operatorname{barccosh}(cx)) - \frac{3}{7} c^2 d^3 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5} d^3 x^5 (a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{27} \\ & -\frac{bcd^3 \int \frac{x^5 (-105c^6 x^6 + 385c^4 x^4 - 495c^2 x^2 + 231)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{1155} - \frac{1}{11} c^6 d^3 x^{11} (a + \operatorname{barccosh}(cx)) + \frac{1}{3} c^4 d^3 x^9 (a + \\ & \quad \operatorname{barccosh}(cx)) - \frac{3}{7} c^2 d^3 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5} d^3 x^5 (a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{2113} \end{aligned}$$

---

3.19.  $\int x^4 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& -\frac{bcd^3\sqrt{c^2x^2-1}\int\frac{x^5(-105c^6x^6+385c^4x^4-495c^2x^2+231)}{\sqrt{c^2x^2-1}}dx}{1155\sqrt{cx-1}\sqrt{cx+1}}-\frac{1}{11}c^6d^3x^{11}(a+\operatorname{barccosh}(cx))+ \\
& \frac{1}{3}c^4d^3x^9(a+\operatorname{barccosh}(cx))-\frac{3}{7}c^2d^3x^7(a+\operatorname{barccosh}(cx))+\frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2331} \\
& -\frac{bcd^3\sqrt{c^2x^2-1}\int\frac{x^4(-105c^6x^6+385c^4x^4-495c^2x^2+231)}{\sqrt{c^2x^2-1}}dx^2}{2310\sqrt{cx-1}\sqrt{cx+1}}-\frac{1}{11}c^6d^3x^{11}(a+\operatorname{barccosh}(cx))+ \\
& \frac{1}{3}c^4d^3x^9(a+\operatorname{barccosh}(cx))-\frac{3}{7}c^2d^3x^7(a+\operatorname{barccosh}(cx))+\frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2123} \\
& \frac{bcd^3\sqrt{c^2x^2-1}\int\left(-\frac{105(c^2x^2-1)^{9/2}}{c^4}-\frac{140(c^2x^2-1)^{7/2}}{c^4}-\frac{5(c^2x^2-1)^{5/2}}{c^4}+\frac{6(c^2x^2-1)^{3/2}}{c^4}-\frac{8\sqrt{c^2x^2-1}}{c^4}+\frac{16}{c^4\sqrt{c^2x^2-1}}\right)dx^2}{2310\sqrt{cx-1}\sqrt{cx+1}} \\
& \frac{1}{11}c^6d^3x^{11}(a+\operatorname{barccosh}(cx))+\frac{1}{3}c^4d^3x^9(a+\operatorname{barccosh}(cx))-\frac{3}{7}c^2d^3x^7(a+\operatorname{barccosh}(cx))+ \\
& \frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2009} \\
& -\frac{1}{11}c^6d^3x^{11}(a+ \\
& \operatorname{barccosh}(cx))+\frac{1}{3}c^4d^3x^9(a+\operatorname{barccosh}(cx))-\frac{3}{7}c^2d^3x^7(a+\operatorname{barccosh}(cx))+\frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx))- \\
& \frac{bcd^3\sqrt{c^2x^2-1}\left(-\frac{210(c^2x^2-1)^{11/2}}{11c^6}-\frac{280(c^2x^2-1)^{9/2}}{9c^6}-\frac{10(c^2x^2-1)^{7/2}}{7c^6}+\frac{12(c^2x^2-1)^{5/2}}{5c^6}-\frac{16(c^2x^2-1)^{3/2}}{3c^6}+\frac{32\sqrt{c^2x^2-1}}{c^6}\right)}{2310\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^4*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/2310*(b*c*d^3*sqrt[-1 + c^2*x^2]*((32*sqrt[-1 + c^2*x^2])/c^6 - (16*(-1 + c^2*x^2)^(3/2))/(3*c^6) + (12*(-1 + c^2*x^2)^(5/2))/(5*c^6) - (10*(-1 + c^2*x^2)^(7/2))/(7*c^6) - (280*(-1 + c^2*x^2)^(9/2))/(9*c^6) - (210*(-1 + c^2*x^2)^(11/2))/(11*c^6)))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (d^3*x^5*(a + b*ArcCosh[c*x]))/5 - (3*c^2*d^3*x^7*(a + b*ArcCosh[c*x]))/7 + (c^4*d^3*x^9*(a + b*ArcCosh[c*x]))/3 - (c^6*d^3*x^11*(a + b*ArcCosh[c*x]))/11`

## 3.19.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2113 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[P_x*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`
- rule 2123 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[P_x*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[P_x, x] && (IntegersQ[m, n] || IGtQ[m, -2])`
- rule 2331 `Int[(P_q)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, P_q, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[P_q, x^2] && IntegerQ[(m - 1)/2]`
- rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.19.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.60

method	result
parts	$-d^3 a \left( \frac{1}{11} c^6 x^{11} - \frac{1}{3} c^4 x^9 + \frac{3}{7} c^2 x^7 - \frac{1}{5} x^5 \right) - \frac{d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{11} x^{11}}{11} - \frac{\operatorname{arccosh}(cx) c^9 x^9}{3} + \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} \right)}{c^5}$
derivativedivides	$-d^3 a \left( \frac{1}{11} c^{11} x^{11} - \frac{1}{3} c^9 x^9 + \frac{3}{7} c^7 x^7 - \frac{1}{5} c^5 x^5 \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{11} x^{11}}{11} - \frac{\operatorname{arccosh}(cx) c^9 x^9}{3} + \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} \right)$
default	$-d^3 a \left( \frac{1}{11} c^{11} x^{11} - \frac{1}{3} c^9 x^9 + \frac{3}{7} c^7 x^7 - \frac{1}{5} c^5 x^5 \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{11} x^{11}}{11} - \frac{\operatorname{arccosh}(cx) c^9 x^9}{3} + \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} \right)$

input `int(x^4*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$-d^3 a \left( \frac{1}{11} c^6 x^{11} - \frac{1}{3} c^4 x^9 + \frac{3}{7} c^2 x^7 - \frac{1}{5} x^5 \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{11} x^{11}}{11} - \frac{\operatorname{arccosh}(cx) c^9 x^9}{3} + \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} \right)$$

### 3.19.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 201, normalized size of antiderivative = 0.79

$$\int x^4 (d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx)) dx = \frac{363825 ac^{11} d^3 x^{11} - 1334025 ac^9 d^3 x^9 + 1715175 ac^7 d^3 x^7 - 800415 ac^5 d^3 x^5 + 3465 (105 bc^{11} d^3 x^{11} - 385 bc^9 d^3 x^9 + 495 bc^7 d^3 x^7 - 231 bc^5 d^3 x^5) \log(cx + \sqrt{c^2 x^2 - 1}) - (33075 bc^{10} d^3 x^{10} - 111475 bc^8 d^3 x^8 + 117625 bc^6 d^3 x^6 - 18933 bc^4 d^3 x^4 - 25244 bc^2 d^3 x^2 - 50488 b d^3) \sqrt{c^2 x^2 - 1}}{c^5}$$

input `integrate(x^4*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output 
$$-1/4002075 * (363825 * a * c^{11} * d^3 * x^{11} - 1334025 * a * c^9 * d^3 * x^9 + 1715175 * a * c^7 * d^3 * x^7 - 800415 * a * c^5 * d^3 * x^5 + 3465 * (105 * b * c^{11} * d^3 * x^{11} - 385 * b * c^9 * d^3 * x^9 + 495 * b * c^7 * d^3 * x^7 - 231 * b * c^5 * d^3 * x^5) * \log(cx + \sqrt{c^2 * x^2 - 1}) - (33075 * b * c^{10} * d^3 * x^{10} - 111475 * b * c^8 * d^3 * x^8 + 117625 * b * c^6 * d^3 * x^6 - 18933 * b * c^4 * d^3 * x^4 - 25244 * b * c^2 * d^3 * x^2 - 50488 * b * d^3) * \sqrt{c^2 * x^2 - 1}) / c^5$$

### 3.19.6 Sympy [F(-1)]

Timed out.

$$\int x^4(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**4*(-c**2*d*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `Timed out`

### 3.19.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs.  $2(212) = 424$ .

Time = 0.24 (sec) , antiderivative size = 465, normalized size of antiderivative = 1.82

$$\begin{aligned} \int x^4(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = & -\frac{1}{11} ac^6 d^3 x^{11} + \frac{1}{3} ac^4 d^3 x^9 - \frac{3}{7} ac^2 d^3 x^7 \\ & - \frac{1}{7623} \left( 693 x^{11} \operatorname{arcosh}(cx) - \left( \frac{63 \sqrt{c^2 x^2 - 1} x^{10}}{c^2} + \frac{70 \sqrt{c^2 x^2 - 1} x^8}{c^4} + \frac{80 \sqrt{c^2 x^2 - 1} x^6}{c^6} + \frac{96 \sqrt{c^2 x^2 - 1} x^4}{c^8} \right) \right) \\ & + \frac{1}{945} \left( 315 x^9 \operatorname{arcosh}(cx) - \left( \frac{35 \sqrt{c^2 x^2 - 1} x^8}{c^2} + \frac{40 \sqrt{c^2 x^2 - 1} x^6}{c^4} + \frac{48 \sqrt{c^2 x^2 - 1} x^4}{c^6} + \frac{64 \sqrt{c^2 x^2 - 1} x^2}{c^8} \right) \right) \\ & + \frac{1}{5} ad^3 x^5 \\ & - \frac{3}{245} \left( 35 x^7 \operatorname{arcosh}(cx) - \left( \frac{5 \sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6 \sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16 \sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) bc^2 c \\ & + \frac{1}{75} \left( 15 x^5 \operatorname{arcosh}(cx) - \left( \frac{3 \sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4 \sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bd^3 \end{aligned}$$

input `integrate(x^4*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

```
output -1/11*a*c^6*d^3*x^11 + 1/3*a*c^4*d^3*x^9 - 3/7*a*c^2*d^3*x^7 - 1/7623*(693
*x^11*arccosh(c*x) - (63*sqrt(c^2*x^2 - 1)*x^10/c^2 + 70*sqrt(c^2*x^2 - 1)
*x^8/c^4 + 80*sqrt(c^2*x^2 - 1)*x^6/c^6 + 96*sqrt(c^2*x^2 - 1)*x^4/c^8 + 1
28*sqrt(c^2*x^2 - 1)*x^2/c^10 + 256*sqrt(c^2*x^2 - 1)/c^12)*c)*b*c^6*d^3 +
1/945*(315*x^9*arccosh(c*x) - (35*sqrt(c^2*x^2 - 1)*x^8/c^2 + 40*sqrt(c^2
*x^2 - 1)*x^6/c^4 + 48*sqrt(c^2*x^2 - 1)*x^4/c^6 + 64*sqrt(c^2*x^2 - 1)*x
^2/c^8 + 128*sqrt(c^2*x^2 - 1)/c^10)*c)*b*c^4*d^3 + 1/5*a*d^3*x^5 - 3/245*(
35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x
^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*c^2*
d^3 + 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^
2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d^3
```

### 3.19.8 Giac [F(-2)]

Exception generated.

$$\int x^4 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

```
input integrate(x^4*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve
cteur & l) Error: Bad Argument Value
```

### 3.19.9 Mupad [F(-1)]

Timed out.

$$\int x^4 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3 dx$$

```
input int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3,x)
```

```
output int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3, x)
```

### 3.20 $\int x^3(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.20.1	Optimal result . . . . .	343
3.20.2	Mathematica [A] (verified) . . . . .	344
3.20.3	Rubi [A] (verified) . . . . .	344
3.20.4	Maple [A] (verified) . . . . .	347
3.20.5	Fricas [A] (verification not implemented) . . . . .	348
3.20.6	Sympy [F] . . . . .	348
3.20.7	Maxima [B] (verification not implemented) . . . . .	349
3.20.8	Giac [F(-2)] . . . . .	350
3.20.9	Mupad [F(-1)] . . . . .	350

#### 3.20.1 Optimal result

Integrand size = 25, antiderivative size = 230

$$\int x^3(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{49bd^3x\sqrt{-1+cx}\sqrt{1+cx}}{5120c^3} + \frac{49bd^3x(-1+cx)^{3/2}(1+cx)^{3/2}}{7680c^3}$$

$$- \frac{49bd^3x(-1+cx)^{5/2}(1+cx)^{5/2}}{9600c^3} + \frac{7bd^3x(-1+cx)^{7/2}(1+cx)^{7/2}}{1600c^3}$$

$$+ \frac{bd^3x(-1+cx)^{9/2}(1+cx)^{9/2}}{100c^3} + \frac{49bd^3\operatorname{arccosh}(cx)}{5120c^4}$$

$$- \frac{d^3(-1+cx)^4(1+cx)^4(a + \operatorname{barccosh}(cx))}{8c^4} - \frac{d^3(-1+cx)^5(1+cx)^5(a + \operatorname{barccosh}(cx))}{10c^4}$$

output  $49/7680*b*d^3*x*(c*x-1)^{(3/2)}*(c*x+1)^{(3/2)}/c^3-49/9600*b*d^3*x*(c*x-1)^{(5/2)}*(c*x+1)^{(5/2)}/c^3+7/1600*b*d^3*x*(c*x-1)^{(7/2)}*(c*x+1)^{(7/2)}/c^3+1/100*b*d^3*x*(c*x-1)^{(9/2)}*(c*x+1)^{(9/2)}/c^3+49/5120*b*d^3*\operatorname{arccosh}(c*x)/c^4-1/8*d^3*(c*x-1)^4*(c*x+1)^4*(a+b*\operatorname{arccosh}(c*x))/c^4-1/10*d^3*(c*x-1)^5*(c*x+1)^5*(a+b*\operatorname{arccosh}(c*x))/c^4-49/5120*b*d^3*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3$



### 3.20.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.70

$$\int x^3 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx =$$

$$\frac{d^3 \left( 1920ac^4 x^4 (-10 + 20c^2 x^2 - 15c^4 x^4 + 4c^6 x^6) + bcx \sqrt{-1 + cx} \sqrt{1 + cx} (1185 + 790c^2 x^2 - 3208c^4 x^4 + \dots) \right)}{c^4}$$

input `Integrate[x^3*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output 
$$\frac{-1/76800*(d^3*(1920*a*c^4*x^4*(-10 + 20*c^2*x^2 - 15*c^4*x^4 + 4*c^6*x^6) + b*c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(1185 + 790*c^2*x^2 - 3208*c^4*x^4 + 2736*c^6*x^6 - 768*c^8*x^8) + 1920*b*c^4*x^4*(-10 + 20*c^2*x^2 - 15*c^4*x^4 + 4*c^6*x^6)*\operatorname{ArcCosh}[c*x] + 2370*b*\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]])}{c^4}$$

### 3.20.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6336, 27, 2003, 35, 646, 40, 40, 40, 40, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6336}$$

$$-bc \int -\frac{d^3(1 - c^2 x^2)^4 (4c^2 x^2 + 1)}{40c^4 \sqrt{cx - 1} \sqrt{cx + 1}} dx + \frac{d^3(1 - c^2 x^2)^5 (a + \operatorname{barccosh}(cx))}{10c^4} -$$

$$\frac{d^3(1 - c^2 x^2)^4 (a + \operatorname{barccosh}(cx))}{8c^4}$$

$$\downarrow \text{27}$$

$$\frac{bd^3 \int \frac{(1 - c^2 x^2)^4 (4c^2 x^2 + 1)}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{40c^3} + \frac{d^3(1 - c^2 x^2)^5 (a + \operatorname{barccosh}(cx))}{10c^4} - \frac{d^3(1 - c^2 x^2)^4 (a + \operatorname{barccosh}(cx))}{8c^4}$$

$$\downarrow \text{2003}$$

---

3.20.  $\int x^3 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& \frac{bd^3 \int \frac{(-cx-1)^4(cx-1)^{7/2}(4c^2x^2+1)}{\sqrt{cx+1}} dx}{40c^3} + \frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \\
& \quad \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} \\
& \quad \downarrow 35 \\
& \frac{bd^3 \int (cx-1)^{7/2}(cx+1)^{7/2}(4c^2x^2+1) dx}{40c^3} + \frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \\
& \quad \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} \\
& \quad \downarrow 646 \\
& \frac{bd^3 \left( \frac{7}{5} \int (cx-1)^{7/2}(cx+1)^{7/2} dx + \frac{2}{5}x(cx-1)^{9/2}(cx+1)^{9/2} \right)}{40c^3} + \\
& \frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} \\
& \quad \downarrow 40 \\
& \frac{bd^3 \left( \frac{7}{5} \left( \frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8} \int (cx-1)^{5/2}(cx+1)^{5/2} dx \right) + \frac{2}{5}x(cx-1)^{9/2}(cx+1)^{9/2} \right)}{40c^3} + \\
& \frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} \\
& \quad \downarrow 40 \\
& \frac{bd^3 \left( \frac{7}{5} \left( \frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8} \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \int (cx-1)^{3/2}(cx+1)^{3/2} dx \right) \right) + \frac{2}{5}x(cx-1)^{9/2}(cx+1)^{9/2} \right)}{40c^3} + \\
& \frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} \\
& \quad \downarrow 40 \\
& \frac{bd^3 \left( \frac{7}{5} \left( \frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8} \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \int \sqrt{cx-1}\sqrt{cx+1} dx \right) \right) \right) + \frac{2}{5}x(cx-1)^{9/2}(cx+1)^{9/2} \right)}{40c^3} + \\
& \frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} \\
& \quad \downarrow 40 \\
& \frac{bd^3 \left( \frac{7}{5} \left( \frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8} \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) \right) \right) \right) + \frac{2}{5}x(cx-1)^{9/2}(cx+1)^{9/2} \right)}{40c^3} + \\
& \frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} \\
& \quad \downarrow 43
\end{aligned}$$

---

3.20.  $\int x^3(d-c^2dx^2)^3(a+\operatorname{barccosh}(cx)) dx$

$$\frac{d^3(1-c^2x^2)^5(a+\operatorname{barccosh}(cx))}{10c^4} - \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^4} + \frac{bd^3\left(\frac{7}{5}\left(\frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8}\left(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6}\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}\right)\right)\right)\right)}{40c^3}$$

input `Int[x^3*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/8*(d^3*(1 - c^2*x^2)^4*(a + b*ArcCosh[c*x]))/c^4 + (d^3*(1 - c^2*x^2)^5*(a + b*ArcCosh[c*x]))/(10*c^4) + (b*d^3*((2*x*(-1 + c*x)^(9/2)*(1 + c*x)^(9/2))/5 + (7*((x*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2))/8 - (7*((x*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/6 - (5*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/6))/8))/5)/(40*c^3)`

### 3.20.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 35 `Int[(u_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(b/d)^m Int[u*(c + d*x)^(m + n), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && !(IntegerQ[n] && SimplerQ[a + b*x, c + d*x])`

rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

```
rule 646 Int[((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_)*(x_)
^2), x_Symbol] := Simp[b*x*(c + d*x)^(m + 1)*((e + f*x)^(n + 1)/(d*f*(2*m +
3))), x] - Simp[(b*c*e - a*d*f*(2*m + 3))/(d*f*(2*m + 3)) Int[(c + d*x)^
m*(e + f*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m, n] &&
EqQ[d*e + c*f, 0] && !LtQ[m, -1]
```

```
rule 2003 Int[(u_)*((c_) + (d_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] :
> Int[u*(c + d*x)^(n + p)*(a/c + (b/d)*x)^p, x] /; FreeQ[{a, b, c, d, n, p}
, x] && EqQ[b*c^2 + a*d^2, 0] && (IntegerQ[p] || (GtQ[a, 0] && GtQ[c, 0] &&
!IntegerQ[n]))
```

```
rule 6336 Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)
^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.20.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.01

method	result
parts	$-d^3 a \left( \frac{1}{10} c^6 x^{10} - \frac{3}{8} c^4 x^8 + \frac{1}{2} c^2 x^6 - \frac{1}{4} x^4 \right) - \frac{d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{10} x^{10}}{10} - \frac{3 \operatorname{arccosh}(cx) c^8 x^8}{8} + \frac{\operatorname{arccosh}(cx) c^6 x^6}{2} - \frac{c^4 x^4 \operatorname{arccosh}(cx)}{4} \right)}{\frac{1}{10} c^{10} x^{10} - \frac{3}{8} c^8 x^8 + \frac{1}{2} c^6 x^6 - \frac{1}{4} c^4 x^4} - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{10} x^{10}}{10} - \frac{3 \operatorname{arccosh}(cx) c^8 x^8}{8} + \frac{\operatorname{arccosh}(cx) c^6 x^6}{2} - \frac{c^4 x^4 \operatorname{arccosh}(cx)}{4} \right)$
derivativedivides	
default	$\frac{-d^3 a \left( \frac{1}{10} c^{10} x^{10} - \frac{3}{8} c^8 x^8 + \frac{1}{2} c^6 x^6 - \frac{1}{4} c^4 x^4 \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{10} x^{10}}{10} - \frac{3 \operatorname{arccosh}(cx) c^8 x^8}{8} + \frac{\operatorname{arccosh}(cx) c^6 x^6}{2} - \frac{c^4 x^4 \operatorname{arccosh}(cx)}{4} \right)}{\frac{1}{10} c^{10} x^{10} - \frac{3}{8} c^8 x^8 + \frac{1}{2} c^6 x^6 - \frac{1}{4} c^4 x^4} - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^{10} x^{10}}{10} - \frac{3 \operatorname{arccosh}(cx) c^8 x^8}{8} + \frac{\operatorname{arccosh}(cx) c^6 x^6}{2} - \frac{c^4 x^4 \operatorname{arccosh}(cx)}{4} \right)$

```
input int(x^3*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

output  $-d^3 a * (1/10 * c^6 * x^{10} - 3/8 * c^4 * x^8 + 1/2 * c^2 * x^6 - 1/4 * x^4) - d^3 b / c^4 * (1/10 * \operatorname{arc} \cosh(c * x) * c^{10} * x^{10} - 3/8 * \operatorname{arccosh}(c * x) * c^8 * x^8 + 1/2 * \operatorname{arccosh}(c * x) * c^6 * x^6 - 1/4 * c^4 * x^4 * \operatorname{arccosh}(c * x) + 1/76800 * (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * (-768 * c^9 * x^9 * (c^2 * x^2 - 1)^{(1/2)} + 2736 * c^7 * x^7 * (c^2 * x^2 - 1)^{(1/2)} - 3208 * c^5 * x^5 * (c^2 * x^2 - 1)^{(1/2)} + 790 * (c^2 * x^2 - 1)^{(1/2)} * c^3 * x^3 + 1185 * c * x * (c^2 * x^2 - 1)^{(1/2)} + 1185 * \ln(c * x + (c^2 * x^2 - 1)^{(1/2)})) / (c^2 * x^2 - 1)^{(1/2)})$

### 3.20.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.86

$$\int x^3 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \frac{7680 ac^{10} d^3 x^{10} - 28800 ac^8 d^3 x^8 + 38400 ac^6 d^3 x^6 - 19200 ac^4 d^3 x^4 + 15 (512 bc^{10} d^3 x^{10} - 1920 bc^8 d^3 x^8 +$$

input `integrate(x^3*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $-1/76800 * (7680 * a * c^{10} * d^3 * x^{10} - 28800 * a * c^8 * d^3 * x^8 + 38400 * a * c^6 * d^3 * x^6 - 19200 * a * c^4 * d^3 * x^4 + 15 * (512 * b * c^{10} * d^3 * x^{10} - 1920 * b * c^8 * d^3 * x^8 + 2560 * b * c^6 * d^3 * x^6 - 1280 * b * c^4 * d^3 * x^4 + 79 * b * d^3) * \log(c * x + \sqrt{c^2 * x^2 - 1}) - (768 * b * c^9 * d^3 * x^9 - 2736 * b * c^7 * d^3 * x^7 + 3208 * b * c^5 * d^3 * x^5 - 790 * b * c^3 * d^3 * x^3 - 1185 * b * c * d^3 * x) * \sqrt{c^2 * x^2 - 1}) / c^4$

### 3.20.6 Sympy [F]

$$\begin{aligned} \int x^3 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx &= -d^3 \left( \int (-ax^3) dx + \int 3ac^2 x^5 dx \right. \\ &\quad + \int (-3ac^4 x^7) dx + \int ac^6 x^9 dx \\ &\quad + \int (-bx^3 \operatorname{acosh}(cx)) dx \\ &\quad + \int 3bc^2 x^5 \operatorname{acosh}(cx) dx \\ &\quad + \int (-3bc^4 x^7 \operatorname{acosh}(cx)) dx \\ &\quad \left. + \int bc^6 x^9 \operatorname{acosh}(cx) dx \right) \end{aligned}$$

input `integrate(x**3*(-c**2*d*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `-d**3*(Integral(-a*x**3, x) + Integral(3*a*c**2*x**5, x) + Integral(-3*a*c**4*x**7, x) + Integral(a*c**6*x**9, x) + Integral(-b*x**3*acosh(c*x), x) + Integral(3*b*c**2*x**5*acosh(c*x), x) + Integral(-3*b*c**4*x**7*acosh(c*x), x) + Integral(b*c**6*x**9*acosh(c*x), x))`

### 3.20.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 501 vs.  $2(194) = 388$ .

Time = 0.22 (sec) , antiderivative size = 501, normalized size of antiderivative = 2.18

$$\int x^3 (d - c^2 dx^2)^3 (a + \operatorname{arccosh}(cx)) dx = -\frac{1}{10} ac^6 d^3 x^{10} + \frac{3}{8} ac^4 d^3 x^8 - \frac{1}{2} ac^2 d^3 x^6 - \frac{1}{12800} \left( 1280 x^{10} \operatorname{arccosh}(cx) - \left( \frac{128 \sqrt{c^2 x^2 - 1} x^9}{c^2} + \frac{144 \sqrt{c^2 x^2 - 1} x^7}{c^4} + \frac{168 \sqrt{c^2 x^2 - 1} x^5}{c^6} + \frac{210 \sqrt{c^2 x^2 - 1} x^3}{c^8} \right) + \frac{1}{1024} \left( 384 x^8 \operatorname{arccosh}(cx) - \left( \frac{48 \sqrt{c^2 x^2 - 1} x^7}{c^2} + \frac{56 \sqrt{c^2 x^2 - 1} x^5}{c^4} + \frac{70 \sqrt{c^2 x^2 - 1} x^3}{c^6} + \frac{105 \sqrt{c^2 x^2 - 1} x}{c^8} \right) + \frac{1}{4} ad^3 x^4 - \frac{1}{96} \left( 48 x^6 \operatorname{arccosh}(cx) - \left( \frac{8 \sqrt{c^2 x^2 - 1} x^5}{c^2} + \frac{10 \sqrt{c^2 x^2 - 1} x^3}{c^4} + \frac{15 \sqrt{c^2 x^2 - 1} x}{c^6} + \frac{15 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1})}{c^7} \right) + \frac{1}{32} \left( 8 x^4 \operatorname{arccosh}(cx) - \left( \frac{2 \sqrt{c^2 x^2 - 1} x^3}{c^2} + \frac{3 \sqrt{c^2 x^2 - 1} x}{c^4} + \frac{3 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1}c)}{c^5} \right) \right) c \right) bd^3$$

input `integrate(x^3*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

```
output -1/10*a*c^6*d^3*x^10 + 3/8*a*c^4*d^3*x^8 - 1/2*a*c^2*d^3*x^6 - 1/12800*(12
80*x^10*arccosh(c*x) - (128*sqrt(c^2*x^2 - 1)*x^9/c^2 + 144*sqrt(c^2*x^2 -
1)*x^7/c^4 + 168*sqrt(c^2*x^2 - 1)*x^5/c^6 + 210*sqrt(c^2*x^2 - 1)*x^3/c^
8 + 315*sqrt(c^2*x^2 - 1)*x/c^10 + 315*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c
)/c^11)*c)*b*c^6*d^3 + 1/1024*(384*x^8*arccosh(c*x) - (48*sqrt(c^2*x^2 - 1
)*x^7/c^2 + 56*sqrt(c^2*x^2 - 1)*x^5/c^4 + 70*sqrt(c^2*x^2 - 1)*x^3/c^6 +
105*sqrt(c^2*x^2 - 1)*x/c^8 + 105*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^9
)*c)*b*c^4*d^3 + 1/4*a*d^3*x^4 - 1/96*(48*x^6*arccosh(c*x) - (8*sqrt(c^2*x
^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c^4 + 15*sqrt(c^2*x^2 - 1)*x/c^
6 + 15*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^7)*c)*b*c^2*d^3 + 1/32*(8*x^
4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4
+ 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*d^3
```

### 3.20.8 Giac [F(-2)]

Exception generated.

$$\int x^3 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.20.9 Mupad [F(-1)]

Timed out.

$$\int x^3 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x^3 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3 dx$$

```
input int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3,x)
```

```
output int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3, x)
```

### 3.21 $\int x^2(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.21.1	Optimal result . . . . .	351
3.21.2	Mathematica [A] (verified) . . . . .	352
3.21.3	Rubi [A] (verified) . . . . .	352
3.21.4	Maple [A] (verified) . . . . .	355
3.21.5	Fricas [A] (verification not implemented) . . . . .	355
3.21.6	Sympy [F] . . . . .	356
3.21.7	Maxima [B] (verification not implemented) . . . . .	356
3.21.8	Giac [F(-2)] . . . . .	357
3.21.9	Mupad [F(-1)] . . . . .	357

#### 3.21.1 Optimal result

Integrand size = 25, antiderivative size = 227

$$\int x^2(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{16bd^3\sqrt{-1+cx}\sqrt{1+cx}}{315c^3} + \frac{8bd^3(-1+cx)^{3/2}(1+cx)^{3/2}}{945c^3} - \frac{2bd^3(-1+cx)^{5/2}(1+cx)^{5/2}}{525c^3}$$

$$+ \frac{bd^3(-1+cx)^{7/2}(1+cx)^{7/2}}{441c^3} + \frac{bd^3(-1+cx)^{9/2}(1+cx)^{9/2}}{81c^3}$$

$$+ \frac{1}{3}d^3x^3(a + \operatorname{barccosh}(cx)) - \frac{3}{5}c^2d^3x^5(a + \operatorname{barccosh}(cx)) + \frac{3}{7}c^4d^3x^7(a + \operatorname{barccosh}(cx)) - \frac{1}{9}c^6d^3x^9(a + \operatorname{barccosh}(cx))$$

```
output 8/945*b*d^3*(c*x-1)^(3/2)*(c*x+1)^(3/2)/c^3-2/525*b*d^3*(c*x-1)^(5/2)*(c*x
+1)^(5/2)/c^3+1/441*b*d^3*(c*x-1)^(7/2)*(c*x+1)^(7/2)/c^3+1/81*b*d^3*(c*x-
1)^(9/2)*(c*x+1)^(9/2)/c^3+1/3*d^3*x^3*(a+b*arccosh(c*x))-3/5*c^2*d^3*x^5*
(a+b*arccosh(c*x))+3/7*c^4*d^3*x^7*(a+b*arccosh(c*x))-1/9*c^6*d^3*x^9*(a+b
*arccosh(c*x))-16/315*b*d^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3
```



### 3.21.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.61

$$\int x^2 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \frac{d^3 (315ac^3 x^3 (-105 + 189c^2 x^2 - 135c^4 x^4 + 35c^6 x^6) + b\sqrt{-1 + cx}\sqrt{1 + cx}(5258 + 2629c^2 x^2 - 6297c^4 x^4 + 4675c^6 x^6 - 1225c^8 x^8) + 315b*c^3*x^3*(-105 + 189*c^2*x^2 - 135*c^4*x^4 + 35*c^6*x^6)*\operatorname{ArcCosh}[c*x])}{99225c^3}$$

input `Integrate[x^2*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/99225*(d^3*(315*a*c^3*x^3*(-105 + 189*c^2*x^2 - 135*c^4*x^4 + 35*c^6*x^6) + b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(5258 + 2629*c^2*x^2 - 6297*c^4*x^4 + 4675*c^6*x^6 - 1225*c^8*x^8) + 315*b*c^3*x^3*(-105 + 189*c^2*x^2 - 135*c^4*x^4 + 35*c^6*x^6)*ArcCosh[c*x]))/c^3`

### 3.21.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.97, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6336, 27, 2113, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6336} \\ & -bc \int \frac{d^3 x^3 (-35c^6 x^6 + 135c^4 x^4 - 189c^2 x^2 + 105)}{315\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{9}c^6 d^3 x^9 (a + \operatorname{barccosh}(cx)) + \frac{3}{7}c^4 d^3 x^7 (a + \operatorname{barccosh}(cx)) - \frac{3}{5}c^2 d^3 x^5 (a + \operatorname{barccosh}(cx)) + \frac{1}{3}d^3 x^3 (a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{27} \\ & -\frac{1}{315}bcd^3 \int \frac{x^3 (-35c^6 x^6 + 135c^4 x^4 - 189c^2 x^2 + 105)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{9}c^6 d^3 x^9 (a + \operatorname{barccosh}(cx)) + \frac{3}{7}c^4 d^3 x^7 (a + \operatorname{barccosh}(cx)) - \frac{3}{5}c^2 d^3 x^5 (a + \operatorname{barccosh}(cx)) + \frac{1}{3}d^3 x^3 (a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{2113} \end{aligned}$$

---

3.21.  $\int x^2 (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& -\frac{bcd^3\sqrt{c^2x^2-1}\int\frac{x^3(-35c^6x^6+135c^4x^4-189c^2x^2+105)}{\sqrt{c^2x^2-1}}dx}{315\sqrt{cx-1}\sqrt{cx+1}}-\frac{1}{9}c^6d^3x^9(a+\operatorname{barccosh}(cx))+\frac{3}{7}c^4d^3x^7(a+\operatorname{barccosh}(cx))-\frac{3}{5}c^2d^3x^5(a+\operatorname{barccosh}(cx))+\frac{1}{3}d^3x^3(a+\operatorname{barccosh}(cx))) \\
& \quad \downarrow \text{2331} \\
& -\frac{bcd^3\sqrt{c^2x^2-1}\int\frac{x^2(-35c^6x^6+135c^4x^4-189c^2x^2+105)}{\sqrt{c^2x^2-1}}dx^2}{630\sqrt{cx-1}\sqrt{cx+1}}-\frac{1}{9}c^6d^3x^9(a+\operatorname{barccosh}(cx))+\frac{3}{7}c^4d^3x^7(a+\operatorname{barccosh}(cx))-\frac{3}{5}c^2d^3x^5(a+\operatorname{barccosh}(cx))+\frac{1}{3}d^3x^3(a+\operatorname{barccosh}(cx))) \\
& \quad \downarrow \text{2123} \\
& \frac{bcd^3\sqrt{c^2x^2-1}\int\left(-\frac{35(c^2x^2-1)^{7/2}}{c^2}-\frac{5(c^2x^2-1)^{5/2}}{c^2}+\frac{6(c^2x^2-1)^{3/2}}{c^2}-\frac{8\sqrt{c^2x^2-1}}{c^2}+\frac{16}{c^2\sqrt{c^2x^2-1}}\right)dx^2}{630\sqrt{cx-1}\sqrt{cx+1}}-\frac{1}{9}c^6d^3x^9(a+\operatorname{barccosh}(cx))+\frac{3}{7}c^4d^3x^7(a+\operatorname{barccosh}(cx))-\frac{3}{5}c^2d^3x^5(a+\operatorname{barccosh}(cx))+\frac{1}{3}d^3x^3(a+\operatorname{barccosh}(cx))) \\
& \quad \downarrow \text{2009} \\
& -\frac{1}{9}c^6d^3x^9(a+\operatorname{barccosh}(cx))+\frac{3}{7}c^4d^3x^7(a+\operatorname{barccosh}(cx))-\frac{3}{5}c^2d^3x^5(a+\operatorname{barccosh}(cx))+\frac{1}{3}d^3x^3(a+\operatorname{barccosh}(cx))- \\
& \quad \frac{bcd^3\sqrt{c^2x^2-1}\left(-\frac{70(c^2x^2-1)^{9/2}}{9c^4}-\frac{10(c^2x^2-1)^{7/2}}{7c^4}+\frac{12(c^2x^2-1)^{5/2}}{5c^4}-\frac{16(c^2x^2-1)^{3/2}}{3c^4}+\frac{32\sqrt{c^2x^2-1}}{c^4}\right)}{630\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^2*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/630*(b*c*d^3*sqrt[-1 + c^2*x^2]*((32*sqrt[-1 + c^2*x^2])/c^4 - (16*(-1 + c^2*x^2)^(3/2))/(3*c^4) + (12*(-1 + c^2*x^2)^(5/2))/(5*c^4) - (10*(-1 + c^2*x^2)^(7/2))/(7*c^4) - (70*(-1 + c^2*x^2)^(9/2))/(9*c^4)))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (d^3*x^3*(a + b*ArcCosh[c*x]))/3 - (3*c^2*d^3*x^5*(a + b*ArcCosh[c*x]))/5 + (3*c^4*d^3*x^7*(a + b*ArcCosh[c*x]))/7 - (c^6*d^3*x^9*(a + b*ArcCosh[c*x]))/9`

## 3.21.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2113 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`
- rule 2123 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`
- rule 2331 `Int[(Pq_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`
- rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.21.4 Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.64

method	result
parts	$-d^3 a \left( \frac{1}{9} c^6 x^9 - \frac{3}{7} c^4 x^7 + \frac{3}{5} c^2 x^5 - \frac{1}{3} x^3 \right) - \frac{d^3 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{c^3 x^3}{3} \right)}{c^3}$
derivativedivides	$-d^3 a \left( \frac{1}{9} c^9 x^9 - \frac{3}{7} c^7 x^7 + \frac{3}{5} c^5 x^5 - \frac{1}{3} c^3 x^3 \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} \right) - \frac{d^3 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} \right)}{c^3}$
default	$-d^3 a \left( \frac{1}{9} c^9 x^9 - \frac{3}{7} c^7 x^7 + \frac{3}{5} c^5 x^5 - \frac{1}{3} c^3 x^3 \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} \right) - \frac{d^3 b \left( \frac{\operatorname{arccosh}(cx) c^9 x^9}{9} - \frac{3 \operatorname{arccosh}(cx) c^7 x^7}{7} + \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} - \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} \right)}{c^3}$

input `int(x^2*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$-d^3 a \left( \frac{1}{9} c^6 x^9 - \frac{3}{7} c^4 x^7 + \frac{3}{5} c^2 x^5 - \frac{1}{3} x^3 \right) - d^3 b / c^3 \left( \frac{1}{9} \operatorname{arccosh}(cx) c^9 x^9 - \frac{3}{7} \operatorname{arccosh}(cx) c^7 x^7 + \frac{3}{5} \operatorname{arccosh}(cx) c^5 x^5 - \frac{1}{3} c^3 x^3 \operatorname{arccosh}(cx) - \frac{1}{99225} (cx-1)^{1/2} (cx+1)^{1/2} (1225 c^8 x^8 - 4675 c^6 x^6 + 6297 c^4 x^4 - 2629 c^2 x^2 - 5258) \right)$$

### 3.21.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.83

$$\int x^2 (d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx)) dx = \frac{11025 ac^9 d^3 x^9 - 42525 ac^7 d^3 x^7 + 59535 ac^5 d^3 x^5 - 33075 ac^3 d^3 x^3 + 315 (35 bc^9 d^3 x^9 - 135 bc^7 d^3 x^7 + 189 bc^5 d^3 x^5 - 105 bc^3 d^3 x^3) \log(cx + \sqrt{c^2 x^2 - 1}) - (1225 b c^8 d^3 x^8 - 4675 b c^6 d^3 x^6 + 6297 b c^4 d^3 x^4 - 2629 b c^2 d^3 x^2 - 5258 b d^3) \sqrt{c^2 x^2 - 1}}{c^3}$$

input `integrate(x^2*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output 
$$-1/99225 * (11025 * a * c^9 * d^3 * x^9 - 42525 * a * c^7 * d^3 * x^7 + 59535 * a * c^5 * d^3 * x^5 - 33075 * a * c^3 * d^3 * x^3 + 315 * (35 * b * c^9 * d^3 * x^9 - 135 * b * c^7 * d^3 * x^7 + 189 * b * c^5 * d^3 * x^5 - 105 * b * c^3 * d^3 * x^3) * \log(cx + \sqrt{c^2 * x^2 - 1}) - (1225 * b * c^8 * d^3 * x^8 - 4675 * b * c^6 * d^3 * x^6 + 6297 * b * c^4 * d^3 * x^4 - 2629 * b * c^2 * d^3 * x^2 - 5258 * b * d^3) * \sqrt{c^2 * x^2 - 1}) / c^3$$

### 3.21.6 Sympy [F]

$$\int x^2(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = -d^3 \left( \int (-ax^2) dx + \int 3ac^2 x^4 dx \right. \\ \left. + \int (-3ac^4 x^6) dx + \int ac^6 x^8 dx \right. \\ \left. + \int (-bx^2 \operatorname{acosh}(cx)) dx \right. \\ \left. + \int 3bc^2 x^4 \operatorname{acosh}(cx) dx \right. \\ \left. + \int (-3bc^4 x^6 \operatorname{acosh}(cx)) dx \right. \\ \left. + \int bc^6 x^8 \operatorname{acosh}(cx) dx \right)$$

input `integrate(x**2*(-c**2*d*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `-d**3*(Integral(-a*x**2, x) + Integral(3*a*c**2*x**4, x) + Integral(-3*a*c**4*x**6, x) + Integral(a*c**6*x**8, x) + Integral(-b*x**2*acosh(c*x), x) + Integral(3*b*c**2*x**4*acosh(c*x), x) + Integral(-3*b*c**4*x**6*acosh(c*x), x) + Integral(b*c**6*x**8*acosh(c*x), x))`

### 3.21.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 388 vs.  $2(189) = 378$ .

Time = 0.27 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.71

$$\int x^2(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = -\frac{1}{9} ac^6 d^3 x^9 + \frac{3}{7} ac^4 d^3 x^7 \\ - \frac{1}{2835} \left( 315 x^9 \operatorname{arcosh}(cx) - \left( \frac{35 \sqrt{c^2 x^2 - 1} x^8}{c^2} + \frac{40 \sqrt{c^2 x^2 - 1} x^6}{c^4} + \frac{48 \sqrt{c^2 x^2 - 1} x^4}{c^6} + \frac{64 \sqrt{c^2 x^2 - 1} x^2}{c^8} \right) \right) bc^4 d^3 \\ - \frac{3}{5} ac^2 d^3 x^5 \\ + \frac{3}{245} \left( 35 x^7 \operatorname{arcosh}(cx) - \left( \frac{5 \sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6 \sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16 \sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) bc^4 d^3 \\ - \frac{1}{25} \left( 15 x^5 \operatorname{arcosh}(cx) - \left( \frac{3 \sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4 \sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bc^2 d^3 \\ + \frac{1}{3} ad^3 x^3 + \frac{1}{9} \left( 3 x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2 \sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bd^3$$

input `integrate(x^2*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/9*a*c^6*d^3*x^9 + 3/7*a*c^4*d^3*x^7 - 1/2835*(315*x^9*arccosh(c*x) - (3  
5*sqrt(c^2*x^2 - 1)*x^8/c^2 + 40*sqrt(c^2*x^2 - 1)*x^6/c^4 + 48*sqrt(c^2*x  
^2 - 1)*x^4/c^6 + 64*sqrt(c^2*x^2 - 1)*x^2/c^8 + 128*sqrt(c^2*x^2 - 1)/c^1  
0)*c)*b*c^6*d^3 - 3/5*a*c^2*d^3*x^5 + 3/245*(35*x^7*arccosh(c*x) - (5*sqrt  
(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*  
x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*c^4*d^3 - 1/25*(15*x^5*arccosh(c*  
x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c  
^2*x^2 - 1)/c^6)*c)*b*c^2*d^3 + 1/3*a*d^3*x^3 + 1/9*(3*x^3*arccosh(c*x) -  
c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*d^3`

### 3.21.8 Giac [F(-2)]

Exception generated.

$$\int x^2(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^2*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

### 3.21.9 Mupad [F(-1)]

Timed out.

$$\int x^2(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3 dx$$

input `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3,x)`

output `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3, x)`

## 3.22 $\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.22.1	Optimal result	358
3.22.2	Mathematica [A] (warning: unable to verify)	359
3.22.3	Rubi [A] (verified)	359
3.22.4	Maple [A] (verified)	361
3.22.5	Fricas [A] (verification not implemented)	361
3.22.6	Sympy [F]	362
3.22.7	Maxima [B] (verification not implemented)	363
3.22.8	Giac [F(-2)]	364
3.22.9	Mupad [F(-1)]	364

### 3.22.1 Optimal result

Integrand size = 23, antiderivative size = 166

$$\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = -\frac{35bd^3 x \sqrt{-1 + cx} \sqrt{1 + cx}}{1024c} + \frac{35bd^3 x (-1 + cx)^{3/2} (1 + cx)^{3/2}}{1536c} - \frac{7bd^3 x (-1 + cx)^{5/2} (1 + cx)^{5/2}}{384c} + \frac{bd^3 x (-1 + cx)^{7/2} (1 + cx)^{7/2}}{64c} + \frac{35bd^3 \operatorname{arccosh}(cx)}{1024c^2} - \frac{d^3 (1 - c^2 x^2)^4 (a + \operatorname{barccosh}(cx))}{8c^2}$$

output `35/1536*b*d^3*x*(c*x-1)^(3/2)*(c*x+1)^(3/2)/c-7/384*b*d^3*x*(c*x-1)^(5/2)*(c*x+1)^(5/2)/c+1/64*b*d^3*x*(c*x-1)^(7/2)*(c*x+1)^(7/2)/c+35/1024*b*d^3*arccosh(c*x)/c^2-1/8*d^3*(-c^2*x^2+1)^4*(a+b*arccosh(c*x))/c^2-35/1024*b*d^3*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c`

### 3.22.2 Mathematica [A] (warning: unable to verify)

Time = 0.24 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.90

$$\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \frac{d^3 \left( cx(b\sqrt{-1 + cx}\sqrt{1 + cx}(279 - 326c^2x^2 + 200c^4x^4 - 48c^6x^6) + 384acx(-4 + 6c^2x^2 - 4c^4x^4 + c^6x^6)) \right)}{3072c^2}$$

input `Integrate[x*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/3072*(d^3*(c*x*(b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(279 - 326*c^2*x^2 + 200*c^4*x^4 - 48*c^6*x^6) + 384*a*c*x*(-4 + 6*c^2*x^2 - 4*c^4*x^4 + c^6*x^6)) + 384*b*c^2*x^2*(-4 + 6*c^2*x^2 - 4*c^4*x^4 + c^6*x^6)*ArcCosh[c*x] + 558*b*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]])/c^2`

### 3.22.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.97, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.261$ , Rules used = {6329, 40, 40, 40, 40, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow 6329 \\ & \frac{bd^3 \int (cx - 1)^{7/2} (cx + 1)^{7/2} dx}{8c} - \frac{d^3 (1 - c^2 x^2)^4 (a + \operatorname{barccosh}(cx))}{8c^2} \\ & \quad \downarrow 40 \\ & \frac{bd^3 \left( \frac{1}{8} x (cx - 1)^{7/2} (cx + 1)^{7/2} - \frac{7}{8} \int (cx - 1)^{5/2} (cx + 1)^{5/2} dx \right)}{8c} - \frac{d^3 (1 - c^2 x^2)^4 (a + \operatorname{barccosh}(cx))}{8c^2} \\ & \quad \downarrow 40 \\ & \frac{bd^3 \left( \frac{1}{8} x (cx - 1)^{7/2} (cx + 1)^{7/2} - \frac{7}{8} \left( \frac{1}{6} x (cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \int (cx - 1)^{3/2} (cx + 1)^{3/2} dx \right) \right)}{8c} - \frac{d^3 (1 - c^2 x^2)^4 (a + \operatorname{barccosh}(cx))}{8c^2} \end{aligned}$$

---

3.22.  $\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$



↓ 40

$$\frac{bd^3\left(\frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8}\left(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6}\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\int\sqrt{cx-1}\sqrt{cx+1}\right)\right)\right)}{8c} \\ \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^2}$$

↓ 40

$$\frac{bd^3\left(\frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8}\left(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6}\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}\right)\right)\right)\right)}{8c} \\ \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^2}$$

↓ 43

$$\frac{bd^3\left(\frac{1}{8}x(cx-1)^{7/2}(cx+1)^{7/2} - \frac{7}{8}\left(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6}\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}\right)\right)\right)\right)}{8c} \\ \frac{d^3(1-c^2x^2)^4(a+\operatorname{barccosh}(cx))}{8c^2}$$

input `Int[x*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/8*(d^3*(1 - c^2*x^2)^4*(a + b*ArcCosh[c*x]))/c^2 + (b*d^3*((x*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2))/8 - (7*((x*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/6 - (5*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c))))/4))/6))/8)/(8*c)`

### 3.22.3.1 Defintions of rubi rules used

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

### 3.22.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.22

method	result
derivativedivides	$-\frac{d^3 a (c^2 x^2 - 1)^4}{8} - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^8 x^8}{8} - \frac{\operatorname{arccosh}(cx) c^6 x^6}{2} + \frac{3c^4 x^4 \operatorname{arccosh}(cx)}{4} - \frac{c^2 x^2 \operatorname{arccosh}(cx)}{2} + \frac{\operatorname{arccosh}(cx)}{8} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{c^2} \right)$
default	$-\frac{d^3 a (c^2 x^2 - 1)^4}{8} - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^8 x^8}{8} - \frac{\operatorname{arccosh}(cx) c^6 x^6}{2} + \frac{3c^4 x^4 \operatorname{arccosh}(cx)}{4} - \frac{c^2 x^2 \operatorname{arccosh}(cx)}{2} + \frac{\operatorname{arccosh}(cx)}{8} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{c^2} \right)$
parts	$-\frac{d^3 a (c^2 x^2 - 1)^4}{8c^2} - \frac{d^3 b \left( \frac{\operatorname{arccosh}(cx) c^8 x^8}{8} - \frac{\operatorname{arccosh}(cx) c^6 x^6}{2} + \frac{3c^4 x^4 \operatorname{arccosh}(cx)}{4} - \frac{c^2 x^2 \operatorname{arccosh}(cx)}{2} + \frac{\operatorname{arccosh}(cx)}{8} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{c^2} \right)}{c^2}$

```
input int(x*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output 1/c^2*(-1/8*d^3*a*(c^2*x^2-1)^4-d^3*b*(1/8*arccosh(c*x)*c^8*x^8-1/2*arccos
h(c*x)*c^6*x^6+3/4*c^4*x^4*arccosh(c*x)-1/2*c^2*x^2*arccosh(c*x)+1/8*arcco
sh(c*x)-1/3072*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(48*c^7*x^7*(c^2*x^2-1)^(1/2)-2
00*c^5*x^5*(c^2*x^2-1)^(1/2)+326*(c^2*x^2-1)^(1/2)*c^3*x^3-279*c*x*(c^2*x^
2-1)^(1/2)+105*ln(c*x+(c^2*x^2-1)^(1/2)))/(c^2*x^2-1)^(1/2))
```

### 3.22.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.11

$$\int x(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx)) dx = \frac{384 ac^8 d^3 x^8 - 1536 ac^6 d^3 x^6 + 2304 ac^4 d^3 x^4 - 1536 ac^2 d^3 x^2 + 3(128 bc^8 d^3 x^8 - 512 bc^6 d^3 x^6 + 768 bc^4 d^3 x^4 - 512 bc^2 d^3 x^2 + 256 bd^3)}{c^2(d - c^2 dx^2)^{3/2}}$$

```
input integrate(x*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")
```

3.22.  $\int x(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx)) dx$

output 
$$\frac{-1/3072*(384*a*c^8*d^3*x^8 - 1536*a*c^6*d^3*x^6 + 2304*a*c^4*d^3*x^4 - 1536*a*c^2*d^3*x^2 + 3*(128*b*c^8*d^3*x^8 - 512*b*c^6*d^3*x^6 + 768*b*c^4*d^3*x^4 - 512*b*c^2*d^3*x^2 + 93*b*d^3)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (48*b*c^7*d^3*x^7 - 200*b*c^5*d^3*x^5 + 326*b*c^3*d^3*x^3 - 279*b*c*d^3*x)*\sqrt{c^2*x^2 - 1})}{c^2}$$

### 3.22.6 Sympy [F]

$$\begin{aligned} \int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = & -d^3 \left( \int (-ax) dx + \int 3ac^2 x^3 dx \right. \\ & + \int (-3ac^4 x^5) dx + \int ac^6 x^7 dx \\ & + \int (-bx \operatorname{acosh}(cx)) dx + \int 3bc^2 x^3 \operatorname{acosh}(cx) dx \\ & + \int (-3bc^4 x^5 \operatorname{acosh}(cx)) dx \\ & \left. + \int bc^6 x^7 \operatorname{acosh}(cx) dx \right) \end{aligned}$$

input `integrate(x*(-c**2*d*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `-d**3*(Integral(-a*x, x) + Integral(3*a*c**2*x**3, x) + Integral(-3*a*c**4*x**5, x) + Integral(a*c**6*x**7, x) + Integral(-b*x*acosh(c*x), x) + Integral(3*b*c**2*x**3*acosh(c*x), x) + Integral(-3*b*c**4*x**5*acosh(c*x), x) + Integral(b*c**6*x**7*acosh(c*x), x))`

**3.22.7 Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 423 vs.  $2(137) = 274$ .

Time = 0.23 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.55

$$\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = -\frac{1}{8} ac^6 d^3 x^8 + \frac{1}{2} ac^4 d^3 x^6 - \frac{1}{3072} \left( 384 x^8 \operatorname{arcosh}(cx) - \left( \frac{48 \sqrt{c^2 x^2 - 1} x^7}{c^2} + \frac{56 \sqrt{c^2 x^2 - 1} x^5}{c^4} + \frac{70 \sqrt{c^2 x^2 - 1} x^3}{c^6} + \frac{105 \sqrt{c^2 x^2 - 1} x}{c^8} - \frac{3}{4} ac^2 d^3 x^4 + \frac{1}{96} \left( 48 x^6 \operatorname{arcosh}(cx) - \left( \frac{8 \sqrt{c^2 x^2 - 1} x^5}{c^2} + \frac{10 \sqrt{c^2 x^2 - 1} x^3}{c^4} + \frac{15 \sqrt{c^2 x^2 - 1} x}{c^6} + \frac{15 \log(2 c^2 x + 2 \sqrt{c^2 x^2 - 1} c)}{c^7} - \frac{3}{32} \left( 8 x^4 \operatorname{arcosh}(cx) - \left( \frac{2 \sqrt{c^2 x^2 - 1} x^3}{c^2} + \frac{3 \sqrt{c^2 x^2 - 1} x}{c^4} + \frac{3 \log(2 c^2 x + 2 \sqrt{c^2 x^2 - 1} c)}{c^5} \right) c \right) bc^2 d^3 + \frac{1}{2} ad^3 x^2 + \frac{1}{4} \left( 2 x^2 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x}{c^2} + \frac{\log(2 c^2 x + 2 \sqrt{c^2 x^2 - 1} c)}{c^3} \right) \right) bd^3 \right)$$

input `integrate(x*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/8*a*c^6*d^3*x^8 + 1/2*a*c^4*d^3*x^6 - 1/3072*(384*x^8*arccosh(c*x) - (48*sqrt(c^2*x^2 - 1)*x^7/c^2 + 56*sqrt(c^2*x^2 - 1)*x^5/c^4 + 70*sqrt(c^2*x^2 - 1)*x^3/c^6 + 105*sqrt(c^2*x^2 - 1)*x/c^8 + 105*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^9)*c)*b*c^6*d^3 - 3/4*a*c^2*d^3*x^4 + 1/96*(48*x^6*arccosh(c*x) - (8*sqrt(c^2*x^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c^4 + 15*sqrt(c^2*x^2 - 1)*x/c^6 + 15*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^7)*c)*b*c^4*d^3 - 3/32*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*c^2*d^3 + 1/2*a*d^3*x^2 + 1/4*(2*x^2*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x/c^2 + log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^3))*b*d^3`

**3.22.8 Giac [F(-2)]**

Exception generated.

$$\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.22.9 Mupad [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3 dx$$

input `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3,x)`

output `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^3, x)`

### 3.23 $\int (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.23.1	Optimal result . . . . .	365
3.23.2	Mathematica [A] (verified) . . . . .	365
3.23.3	Rubi [A] (verified) . . . . .	366
3.23.4	Maple [A] (verified) . . . . .	368
3.23.5	Fricas [A] (verification not implemented) . . . . .	368
3.23.6	Sympy [F] . . . . .	369
3.23.7	Maxima [A] (verification not implemented) . . . . .	370
3.23.8	Giac [F(-2)] . . . . .	370
3.23.9	Mupad [F(-1)] . . . . .	371

#### 3.23.1 Optimal result

Integrand size = 22, antiderivative size = 191

$$\int (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{16bd^3\sqrt{-1+cx}\sqrt{1+cx}}{35c} + \frac{8bd^3(-1+cx)^{3/2}(1+cx)^{3/2}}{105c}$$

$$- \frac{6bd^3(-1+cx)^{5/2}(1+cx)^{5/2}}{175c} + \frac{bd^3(-1+cx)^{7/2}(1+cx)^{7/2}}{49c}$$

$$+ d^3x(a + \operatorname{barccosh}(cx)) - c^2d^3x^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4d^3x^5(a + \operatorname{barccosh}(cx)) - \frac{1}{7}c^6d^3x^7(a + \operatorname{barccosh}(cx))$$

```
output 8/105*b*d^3*(c*x-1)^(3/2)*(c*x+1)^(3/2)/c-6/175*b*d^3*(c*x-1)^(5/2)*(c*x+1)^(5/2)/c+1/49*b*d^3*(c*x-1)^(7/2)*(c*x+1)^(7/2)/c+d^3*x*(a+b*arccosh(c*x))-c^2*d^3*x^3*(a+b*arccosh(c*x))+3/5*c^4*d^3*x^5*(a+b*arccosh(c*x))-1/7*c^6*d^3*x^7*(a+b*arccosh(c*x))-16/35*b*d^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c
```

#### 3.23.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.64

$$\int (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{d^3(b\sqrt{-1+cx}\sqrt{1+cx}(2161 - 757c^2x^2 + 351c^4x^4 - 75c^6x^6) + 105acx(-35 + 35c^2x^2 - 21c^4x^4 + 5c^6x^6))}{3675c}$$

input `Integrate[(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/3675*(d^3*(b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2161 - 757*c^2*x^2 + 351*c^4*x^4 - 75*c^6*x^6) + 105*a*c*x*(-35 + 35*c^2*x^2 - 21*c^4*x^4 + 5*c^6*x^6) + 105*b*c*x*(-35 + 35*c^2*x^2 - 21*c^4*x^4 + 5*c^6*x^6)*ArcCosh[c*x]))/c`

### 3.23.3 Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.02, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {6309, 27, 2113, 2331, 2389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6309} \\
 & -bc \int \frac{d^3 x (-5c^6 x^6 + 21c^4 x^4 - 35c^2 x^2 + 35)}{35\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^6 d^3 x^7 (a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4 d^3 x^5 (a + \operatorname{barccosh}(cx)) - c^2 d^3 x^3 (a + \operatorname{barccosh}(cx)) + d^3 x (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{35}bcd^3 \int \frac{x(-5c^6 x^6 + 21c^4 x^4 - 35c^2 x^2 + 35)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^6 d^3 x^7 (a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4 d^3 x^5 (a + \operatorname{barccosh}(cx)) - c^2 d^3 x^3 (a + \operatorname{barccosh}(cx)) + d^3 x (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{2113} \\
 & -\frac{bcd^3 \sqrt{c^2 x^2 - 1} \int \frac{x(-5c^6 x^6 + 21c^4 x^4 - 35c^2 x^2 + 35)}{\sqrt{c^2 x^2 - 1}} dx}{35\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}c^6 d^3 x^7 (a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4 d^3 x^5 (a + \operatorname{barccosh}(cx)) - c^2 d^3 x^3 (a + \operatorname{barccosh}(cx)) + d^3 x (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{2331} \\
 & -\frac{bcd^3 \sqrt{c^2 x^2 - 1} \int \frac{-5c^6 x^6 + 21c^4 x^4 - 35c^2 x^2 + 35}{\sqrt{c^2 x^2 - 1}} dx^2}{70\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}c^6 d^3 x^7 (a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4 d^3 x^5 (a + \operatorname{barccosh}(cx)) - c^2 d^3 x^3 (a + \operatorname{barccosh}(cx)) + d^3 x (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{2389}
 \end{aligned}$$

$$\frac{bcd^3\sqrt{c^2x^2-1}\int\left(-5(c^2x^2-1)^{5/2}+6(c^2x^2-1)^{3/2}-8\sqrt{c^2x^2-1}+\frac{16}{\sqrt{c^2x^2-1}}\right)dx^2}{70\sqrt{cx-1}\sqrt{cx+1}}-\frac{1}{7}c^6d^3x^7(a+\text{barccosh}(cx))+\frac{3}{5}c^4d^3x^5(a+\text{barccosh}(cx))-c^2d^3x^3(a+\text{barccosh}(cx))+d^3x(a+\text{barccosh}(cx)))$$

↓ 2009

$$-\frac{1}{7}c^6d^3x^7(a+\text{barccosh}(cx))+\frac{3}{5}c^4d^3x^5(a+\text{barccosh}(cx))-c^2d^3x^3(a+\text{barccosh}(cx))+d^3x(a+\text{barccosh}(cx))-\frac{bcd^3\sqrt{c^2x^2-1}\left(-\frac{10(c^2x^2-1)^{7/2}}{7c^2}+\frac{12(c^2x^2-1)^{5/2}}{5c^2}-\frac{16(c^2x^2-1)^{3/2}}{3c^2}+\frac{32\sqrt{c^2x^2-1}}{c^2}\right)}{70\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/70*(b*c*d^3*sqrt[-1 + c^2*x^2]*((32*sqrt[-1 + c^2*x^2])/c^2 - (16*(-1 + c^2*x^2)^(3/2))/(3*c^2) + (12*(-1 + c^2*x^2)^(5/2))/(5*c^2) - (10*(-1 + c^2*x^2)^(7/2))/(7*c^2)))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + d^3*x*(a + b*ArcCosh[c*x]) - c^2*d^3*x^3*(a + b*ArcCosh[c*x]) + (3*c^4*d^3*x^5*(a + b*ArcCosh[c*x]))/5 - (c^6*d^3*x^7*(a + b*ArcCosh[c*x]))/7`

### 3.23.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2113 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m])/(a*c + b*d*x^2)^FracPart[m] Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

rule 2331 `Int[(Pq_)*(x_)^((m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`



rule 2389 `Int[(Pq_)*((a_) + (b_)*(x_)^(n_.))^(p_.), x_Symbol] := Int[ExpandIntegrand [Pq*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, n}, x] && PolyQ[Pq, x] && (IGtQ[p, 0] || EqQ[n, 1])`

rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.23.4 Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.68

method	result
parts	$-d^3 a \left( \frac{1}{7} c^6 x^7 - \frac{3}{5} c^4 x^5 + x^3 c^2 - x \right) - \frac{d^3 b \left( \frac{\operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} + c^3 x^3 \operatorname{arccosh}(cx) - cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c} \right)}{c}$
derivativedivides	$-d^3 a \left( \frac{1}{7} c^7 x^7 - \frac{3}{5} c^5 x^5 + c^3 x^3 - cx \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} + c^3 x^3 \operatorname{arccosh}(cx) - cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c} \right)$
default	$-d^3 a \left( \frac{1}{7} c^7 x^7 - \frac{3}{5} c^5 x^5 + c^3 x^3 - cx \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^7 x^7}{7} - \frac{3 \operatorname{arccosh}(cx) c^5 x^5}{5} + c^3 x^3 \operatorname{arccosh}(cx) - cx \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c} \right)$

input `int((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-d^3*a*(1/7*c^6*x^7-3/5*c^4*x^5+x^3*c^2-x)-d^3*b/c*(1/7*arccosh(c*x)*c^7*x^7-3/5*arccosh(c*x)*c^5*x^5+c^3*x^3*arccosh(c*x)-c*x*arccosh(c*x)-1/3675*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(75*c^6*x^6-351*c^4*x^4+757*c^2*x^2-2161))`

### 3.23.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.88

$$\int (d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx)) dx = \frac{525 ac^7 d^3 x^7 - 2205 ac^5 d^3 x^5 + 3675 ac^3 d^3 x^3 - 3675 acd^3 x + 105 (5 bc^7 d^3 x^7 - 21 bc^5 d^3 x^5 + 35 bc^3 d^3 x^3 - 3675 c^2 d^3 x^2 - 2161 d^3 x)}{3675 c^2}$$

3.23.  $\int (d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx)) dx$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `-1/3675*(525*a*c^7*d^3*x^7 - 2205*a*c^5*d^3*x^5 + 3675*a*c^3*d^3*x^3 - 3675*a*c*d^3*x + 105*(5*b*c^7*d^3*x^7 - 21*b*c^5*d^3*x^5 + 35*b*c^3*d^3*x^3 - 35*b*c*d^3*x)*log(c*x + sqrt(c^2*x^2 - 1)) - (75*b*c^6*d^3*x^6 - 351*b*c^4*d^3*x^4 + 757*b*c^2*d^3*x^2 - 2161*b*d^3)*sqrt(c^2*x^2 - 1))/c`

### 3.23.6 Sympy [F]

$$\begin{aligned} \int (d - c^2 dx^2)^3 (a + \operatorname{arccosh}(cx)) dx &= -d^3 \left( \int (-a) dx + \int (-b \operatorname{acosh}(cx)) dx \right. \\ &\quad + \int 3ac^2 x^2 dx + \int (-3ac^4 x^4) dx + \int ac^6 x^6 dx \\ &\quad + \int 3bc^2 x^2 \operatorname{acosh}(cx) dx \\ &\quad + \int (-3bc^4 x^4 \operatorname{acosh}(cx)) dx \\ &\quad \left. + \int bc^6 x^6 \operatorname{acosh}(cx) dx \right) \end{aligned}$$

input `integrate((-c**2*d*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `-d**3*(Integral(-a, x) + Integral(-b*acosh(c*x), x) + Integral(3*a*c**2*x**2, x) + Integral(-3*a*c**4*x**4, x) + Integral(a*c**6*x**6, x) + Integral(3*b*c**2*x**2*acosh(c*x), x) + Integral(-3*b*c**4*x**4*acosh(c*x), x) + Integral(b*c**6*x**6*acosh(c*x), x))`

### 3.23.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 302, normalized size of antiderivative = 1.58

$$\int (d - c^2 dx^2)^3 (a + \operatorname{arccosh}(cx)) dx = -\frac{1}{7} ac^6 d^3 x^7 + \frac{3}{5} ac^4 d^3 x^5 - \frac{1}{245} \left( 35 x^7 \operatorname{arccosh}(cx) - \left( \frac{5 \sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6 \sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16 \sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) bc^6 d^3 + \frac{1}{25} \left( 15 x^5 \operatorname{arccosh}(cx) - \left( \frac{3 \sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4 \sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bc^4 d^3 - ac^2 d^3 x^3 - \frac{1}{3} \left( 3 x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2 \sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bc^2 d^3 + ad^3 x + \frac{(cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) bd^3}{c}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/7*a*c^6*d^3*x^7 + 3/5*a*c^4*d^3*x^5 - 1/245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*c^6*d^3 + 1/25*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*c^4*d^3 - a*c^2*d^3*x^3 - 1/3*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*c^2*d^3 + a*d^3*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*d^3/c`

### 3.23.8 Giac [F(-2)]

Exception generated.

$$\int (d - c^2 dx^2)^3 (a + \operatorname{arccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.23.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3 dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^3,x)`output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^3, x)`

**3.24**  $\int \frac{(d-c^2dx^2)^3(a+b\operatorname{arccosh}(cx))}{x} dx$

3.24.1 Optimal result . . . . . 372  
 3.24.2 Mathematica [A] (warning: unable to verify) . . . . . 373  
 3.24.3 Rubi [C] (warning: unable to verify) . . . . . 374  
 3.24.4 Maple [A] (verified) . . . . . 380  
 3.24.5 Fricas [F] . . . . . 381  
 3.24.6 Sympy [F] . . . . . 381  
 3.24.7 Maxima [F] . . . . . 382  
 3.24.8 Giac [F(-2)] . . . . . 382  
 3.24.9 Mupad [F(-1)] . . . . . 382

**3.24.1 Optimal result**

Integrand size = 25, antiderivative size = 239

$$\int \frac{(d - c^2dx^2)^3 (a + b\operatorname{arccosh}(cx))}{x} dx$$

$$= \frac{19}{48}bcd^3x\sqrt{-1+cx}\sqrt{1+cx} - \frac{7}{72}bcd^3x(-1+cx)^{3/2}(1+cx)^{3/2}$$

$$+ \frac{1}{36}bcd^3x(-1+cx)^{5/2}(1+cx)^{5/2} - \frac{19}{48}bd^3\operatorname{arccosh}(cx) + \frac{1}{2}d^3(1-c^2x^2)(a+b\operatorname{arccosh}(cx)) + \frac{1}{4}d^3(1-c^2x^2)^2$$

output

```
-7/72*b*c*d^3*x*(c*x-1)^(3/2)*(c*x+1)^(3/2)+1/36*b*c*d^3*x*(c*x-1)^(5/2)*(
c*x+1)^(5/2)-19/48*b*d^3*arccosh(c*x)+1/2*d^3*(-c^2*x^2+1)*(a+b*arccosh(c*
x))+1/4*d^3*(-c^2*x^2+1)^2*(a+b*arccosh(c*x))+1/6*d^3*(-c^2*x^2+1)^3*(a+b*
arccosh(c*x))+1/2*d^3*(a+b*arccosh(c*x))^2/b+d^3*(a+b*arccosh(c*x))*ln(1+1
/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)-1/2*b*d^3*polylog(2,-1/(c*x+(c*x-1)^(
1/2)*(c*x+1)^(1/2)))^2)+19/48*b*c*d^3*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

### 3.24.2 Mathematica [A] (warning: unable to verify)

Time = 0.49 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.28

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x} dx = -\frac{1}{144} d^3 \left( 216ac^2x^2 - 108ac^4x^4 + 24ac^6x^6 \right. \\ \left. + 33bcx \sqrt{\frac{-1+cx}{1+cx}} + 33bc^2x^2 \sqrt{\frac{-1+cx}{1+cx}} \right. \\ \left. + 22bc^3x^3 \sqrt{\frac{-1+cx}{1+cx}} + 22bc^4x^4 \sqrt{\frac{-1+cx}{1+cx}} \right. \\ \left. - 4bc^5x^5 \sqrt{\frac{-1+cx}{1+cx}} - 4bc^6x^6 \sqrt{\frac{-1+cx}{1+cx}} \right. \\ \left. - 108bcx \sqrt{-1+cx} \sqrt{1+cx} - 72\operatorname{barccosh}(cx)^2 \right. \\ \left. - 150\operatorname{barctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right) \right. \\ \left. + 12\operatorname{barccosh}(cx) (18c^2x^2 - 9c^4x^4 + 2c^6x^6 \right. \\ \left. - 12 \log(1 + e^{-2\operatorname{arccosh}(cx)})) - 144a \log(x) \right. \\ \left. + 72b \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)}) \right)$$

input `Integrate[((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x,x]`

output `-1/144*(d^3*(216*a*c^2*x^2 - 108*a*c^4*x^4 + 24*a*c^6*x^6 + 33*b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)] + 33*b*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)] + 22*b*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)] + 22*b*c^4*x^4*Sqrt[(-1 + c*x)/(1 + c*x)] - 4*b*c^5*x^5*Sqrt[(-1 + c*x)/(1 + c*x)] - 4*b*c^6*x^6*Sqrt[(-1 + c*x)/(1 + c*x)] - 108*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 72*b*ArcCosh[c*x]^2 - 150*b*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]] + 12*b*ArcCosh[c*x]*(18*c^2*x^2 - 9*c^4*x^4 + 2*c^6*x^6 - 12*Log[1 + E^(-2*ArcCosh[c*x])]) - 144*a*Log[x] + 72*b*PolyLog[2, -E^(-2*ArcCosh[c*x])]))`

### 3.24.3 Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.35 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.51, number of steps used = 22, number of rules used = 21,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.840$ , Rules used = {6334, 27, 40, 40, 40, 43, 6334, 40, 40, 43, 6334, 40, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x} dx \\
 & \quad \downarrow \text{6334} \\
 & d \int \frac{d^2 (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{6} bcd^3 \int (cx - 1)^{5/2} (cx + 1)^{5/2} dx + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & d^3 \int \frac{(1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{6} bcd^3 \int (cx - 1)^{5/2} (cx + 1)^{5/2} dx + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{40} \\
 & d^3 \int \frac{(1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{6} bcd^3 \left( \frac{1}{6} x (cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \int (cx - 1)^{3/2} (cx + 1)^{3/2} dx \right) + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{40} \\
 & d^3 \int \frac{(1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{6} bcd^3 \left( \frac{1}{6} x (cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \int \sqrt{cx - 1} \sqrt{cx + 1} dx \right) \right) + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{40}
 \end{aligned}$$

$$d^3 \int \frac{(1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{1}{2} \int \frac{1}{\sqrt{cx - 1}\sqrt{cx + 1}} \right) \right) + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) \right)$$

↓ 43

$$d^3 \int \frac{(1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 6334

$$d^3 \left( \int \frac{(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4} bc \int (cx - 1)^{3/2} (cx + 1)^{3/2} dx + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 40

$$d^3 \left( \int \frac{(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4} bc \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \int \sqrt{cx - 1}\sqrt{cx + 1} dx \right) + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 40

$$d^3 \left( \int \frac{(1 - c^2 x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4} bc \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{1}{2} \int \frac{1}{\sqrt{cx - 1}\sqrt{cx + 1}} \right) \right) + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6} d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right)$$

↓ 43



$$d^3 \left( \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx + \frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \right. \right. \\ \left. \left. \frac{1}{6}d^3(1-c^2x^2)^3(a+\operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 6334

$$d^3 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}bc \int \sqrt{cx-1}\sqrt{cx+1} dx + \frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx)) + \frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx)) \right. \\ \left. \frac{1}{6}d^3(1-c^2x^2)^3(a+\operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 40

$$d^3 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}bc \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) + \frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx)) \right. \\ \left. \frac{1}{6}d^3(1-c^2x^2)^3(a+\operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 43

$$d^3 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx)) + \frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx)) + \frac{1}{2}bc \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} \right. \right. \\ \left. \left. \frac{1}{6}d^3(1-c^2x^2)^3(a+\operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 6297

$$d^3 \left( \frac{\int -\left( (a+\operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b} \right) \right) d(a+\operatorname{barccosh}(cx))}{b} + \frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx)) \right. \\ \left. \frac{1}{6}d^3(1-c^2x^2)^3(a+\operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 25

$$d^3 \left( -\frac{\int (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 3042

$$d^3 \left( -\frac{\int -i(a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 26

$$d^3 \left( \frac{i \int (a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 4201

$$d^3 \left( \frac{i \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)} (a + \operatorname{barccosh}(cx))}{1 + e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2} i (a + \operatorname{barccosh}(cx))^2 \right)}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{6} bcd^3 \left( \frac{1}{6} x(cx - 1)^{5/2} (cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4} x(cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 2620

$$d^3 \left( \frac{i(2i(\frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx)))}{b} \right. \\ \left. + \frac{1}{6}d^3(1 - c^2x^2)^3(a + \operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx - 1)^{5/2}(cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 2715

$$d^3 \left( \frac{i(2i(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)))}{b} \right. \\ \left. + \frac{1}{6}d^3(1 - c^2x^2)^3(a + \operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx - 1)^{5/2}(cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

↓ 2838

$$d^3 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx)))}{b} \right. \\ \left. + \frac{1}{6}d^3(1 - c^2x^2)^3(a + \operatorname{barccosh}(cx)) + \frac{1}{6}bcd^3 \left( \frac{1}{6}x(cx - 1)^{5/2}(cx + 1)^{5/2} - \frac{5}{6} \left( \frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) \right) \right)$$

input `Int(((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x,x]`

output `(d^3*(1 - c^2*x^2)^3*(a + b*ArcCosh[c*x])/6 + (b*c*d^3*((x*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/6 - (5*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/6) + d^3*((1 - c^2*x^2)*(a + b*ArcCosh[c*x])/2 + ((1 - c^2*x^2)^2*(a + b*ArcCosh[c*x])/4 + (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/2 - (b*c*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/4 + (I*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])]) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]]/4)))/b)`

## 3.24.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*(c + d*x)^m/(2*m + 1), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6334 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_.) + (e_.)*(x_)^2)^(p_.)/(x_), x_Symbol] := Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d)^p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.24.4 Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.03

method	result
parts	$-d^3 a \left( \frac{c^6 x^6}{6} - \frac{3c^4 x^4}{4} + \frac{3c^2 x^2}{2} - \ln(x) \right) - \frac{d^3 b \operatorname{arccosh}(cx) c^6 x^6}{6} + \frac{3d^3 b \operatorname{arccosh}(cx) c^4 x^4}{4} - \frac{3d^3 b \operatorname{arccosh}(cx) c^2 x^2}{2}$
derivativedivides	$-d^3 a \left( \frac{c^6 x^6}{6} - \frac{3c^4 x^4}{4} + \frac{3c^2 x^2}{2} - \ln(cx) \right) + \frac{25b d^3 \operatorname{arccosh}(cx)}{48} + d^3 b \operatorname{arccosh}(cx) \ln \left( 1 + (cx + \sqrt{c^2 x^2 + d}) \right)$
default	$-d^3 a \left( \frac{c^6 x^6}{6} - \frac{3c^4 x^4}{4} + \frac{3c^2 x^2}{2} - \ln(cx) \right) + \frac{25b d^3 \operatorname{arccosh}(cx)}{48} + d^3 b \operatorname{arccosh}(cx) \ln \left( 1 + (cx + \sqrt{c^2 x^2 + d}) \right)$

input `int((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`

output  $-d^3 a \left( \frac{1}{6} c^6 x^6 - \frac{3}{4} c^4 x^4 + \frac{3}{2} c^2 x^2 - \ln(x) \right) - \frac{1}{6} d^3 b \operatorname{arccosh}(c x) c^6 x^6 + \frac{3}{4} d^3 b \operatorname{arccosh}(c x) c^4 x^4 - \frac{3}{2} d^3 b \operatorname{arccosh}(c x) c^2 x^2 + \frac{1}{2} d^3 b \operatorname{polylog}(2, -(c x + (c x - 1)^{1/2}) (c x + 1)^{1/2})^2 + d^3 b \operatorname{arccosh}(c x) \ln(1 + (c x + (c x - 1)^{1/2}) (c x + 1)^{1/2})^2 - \frac{1}{2} d^3 b \operatorname{arccosh}(c x)^2 + \frac{25}{48} b d^3 \operatorname{arccosh}(c x) + \frac{1}{36} d^3 b (c x + 1)^{1/2} (c x - 1)^{1/2} c^5 x^5 - \frac{11}{72} d^3 b (c x + 1)^{1/2} (c x - 1)^{1/2} c^3 x^3 + \frac{25}{48} b c d^3 x (c x - 1)^{1/2} (c x + 1)^{1/2}$

$$3.24. \quad \int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x} dx$$

## 3.24.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x} dx = \int -\frac{(c^2 dx^2 - d)^3 (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x,x, algorithm="fricas")`

output `integral(-(a*c^6*d^3*x^6 - 3*a*c^4*d^3*x^4 + 3*a*c^2*d^3*x^2 - a*d^3 + (b*c^6*d^3*x^6 - 3*b*c^4*d^3*x^4 + 3*b*c^2*d^3*x^2 - b*d^3)*arccosh(c*x))/x, x)`

## 3.24.6 Sympy [F]

$$\begin{aligned} \int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x} dx = & -d^3 \left( \int \left( -\frac{a}{x} \right) dx + \int 3ac^2 x dx + \int (-3ac^4 x^3) dx \right. \\ & + \int ac^6 x^5 dx + \int \left( -\frac{b \operatorname{acosh}(cx)}{x} \right) dx \\ & + \int 3bc^2 x \operatorname{acosh}(cx) dx \\ & + \int (-3bc^4 x^3 \operatorname{acosh}(cx)) dx \\ & \left. + \int bc^6 x^5 \operatorname{acosh}(cx) dx \right) \end{aligned}$$

input `integrate((-c**2*d*x**2+d)**3*(a+b*acosh(c*x))/x,x)`

output `-d**3*(Integral(-a/x, x) + Integral(3*a*c**2*x, x) + Integral(-3*a*c**4*x**3, x) + Integral(a*c**6*x**5, x) + Integral(-b*acosh(c*x)/x, x) + Integral(3*b*c**2*x*acosh(c*x), x) + Integral(-3*b*c**4*x**3*acosh(c*x), x) + Integral(b*c**6*x**5*acosh(c*x), x))`

**3.24.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x} dx = \int -\frac{(c^2 dx^2 - d)^3 (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `-1/6*a*c^6*d^3*x^6 + 3/4*a*c^4*d^3*x^4 - 3/2*a*c^2*d^3*x^2 + a*d^3*log(x) - integrate(b*c^6*d^3*x^5*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) - 3*b*c^4*d^3*x^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 3*b*c^2*d^3*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) - b*d^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/x, x)`

**3.24.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.24.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3}{x} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x, x)`

---

3.24.  $\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x} dx$

**3.25** 
$$\int \frac{(d-c^2dx^2)^3(a+b\operatorname{arccosh}(cx))}{x^2} dx$$

3.25.1	Optimal result	383
3.25.2	Mathematica [A] (verified)	383
3.25.3	Rubi [A] (verified)	384
3.25.4	Maple [A] (verified)	386
3.25.5	Fricas [A] (verification not implemented)	387
3.25.6	Sympy [F]	387
3.25.7	Maxima [A] (verification not implemented)	388
3.25.8	Giac [F(-2)]	388
3.25.9	Mupad [F(-1)]	389

**3.25.1 Optimal result**

Integrand size = 25, antiderivative size = 180

$$\int \frac{(d - c^2dx^2)^3 (a + b\operatorname{arccosh}(cx))}{x^2} dx$$

$$= \frac{11}{5}bcd^3\sqrt{-1+cx}\sqrt{1+cx} - \frac{1}{5}bcd^3(-1+cx)^{3/2}(1+cx)^{3/2}$$

$$+ \frac{1}{25}bcd^3(-1+cx)^{5/2}(1+cx)^{5/2} - \frac{d^3(a+b\operatorname{arccosh}(cx))}{x} - 3c^2d^3x(a+b\operatorname{arccosh}(cx)) + c^4d^3x^3(a+b\operatorname{arccosh}(cx))$$

output

```
-1/5*b*c*d^3*(c*x-1)^(3/2)*(c*x+1)^(3/2)+1/25*b*c*d^3*(c*x-1)^(5/2)*(c*x+1)^(5/2)-d^3*(a+b*arccosh(c*x))/x-3*c^2*d^3*x*(a+b*arccosh(c*x))+c^4*d^3*x^3*(a+b*arccosh(c*x))-1/5*c^6*d^3*x^5*(a+b*arccosh(c*x))+b*c*d^3*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))+11/5*b*c*d^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

**3.25.2 Mathematica [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.76

$$\int \frac{(d - c^2dx^2)^3 (a + b\operatorname{arccosh}(cx))}{x^2} dx = \frac{1}{25}d^3 \left( -\frac{25a}{x} - 75ac^2x + 25ac^4x^3 - 5ac^6x^5 \right.$$

$$+ bc\sqrt{-1+cx}\sqrt{1+cx}(61 - 7c^2x^2 + c^4x^4)$$

$$- \frac{5b(5 + 15c^2x^2 - 5c^4x^4 + c^6x^6) \operatorname{arccosh}(cx)}{x}$$

$$\left. - 25bc \operatorname{arctan} \left( \frac{1}{\sqrt{-1+cx}\sqrt{1+cx}} \right) \right)$$

3.25. 
$$\int \frac{(d-c^2dx^2)^3(a+b\operatorname{arccosh}(cx))}{x^2} dx$$



input `Integrate[((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x^2,x]`

output `(d^3*((-25*a)/x - 75*a*c^2*x + 25*a*c^4*x^3 - 5*a*c^6*x^5 + b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(61 - 7*c^2*x^2 + c^4*x^4) - (5*b*(5 + 15*c^2*x^2 - 5*c^4*x^4 + c^6*x^6)*ArcCosh[c*x])/x - 25*b*c*ArcTan[1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])]))/25`

### 3.25.3 Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6336, 27, 2113, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^3 (a + \text{barccosh}(cx))}{x^2} dx \\
 & \quad \downarrow \text{6336} \\
 & -bc \int -\frac{d^3(c^6 x^6 - 5c^4 x^4 + 15c^2 x^2 + 5)}{5x\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{5}c^6 d^3 x^5 (a + \text{barccosh}(cx)) + c^4 d^3 x^3 (a + \\
 & \quad \text{barccosh}(cx)) - 3c^2 d^3 x (a + \text{barccosh}(cx)) - \frac{d^3(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{5}bcd^3 \int \frac{c^6 x^6 - 5c^4 x^4 + 15c^2 x^2 + 5}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{5}c^6 d^3 x^5 (a + \text{barccosh}(cx)) + c^4 d^3 x^3 (a + \text{barccosh}(cx)) - \\
 & \quad 3c^2 d^3 x (a + \text{barccosh}(cx)) - \frac{d^3(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{2113} \\
 & \frac{bcd^3 \sqrt{c^2 x^2 - 1} \int \frac{c^6 x^6 - 5c^4 x^4 + 15c^2 x^2 + 5}{x\sqrt{c^2 x^2 - 1}} dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{1}{5}c^6 d^3 x^5 (a + \text{barccosh}(cx)) + c^4 d^3 x^3 (a + \text{barccosh}(cx)) - \\
 & \quad 3c^2 d^3 x (a + \text{barccosh}(cx)) - \frac{d^3(a + \text{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{2331} \\
 & \frac{bcd^3 \sqrt{c^2 x^2 - 1} \int \frac{c^6 x^6 - 5c^4 x^4 + 15c^2 x^2 + 5}{x^2 \sqrt{c^2 x^2 - 1}} dx^2}{10\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{1}{5}c^6 d^3 x^5 (a + \text{barccosh}(cx)) + c^4 d^3 x^3 (a + \\
 & \quad \text{barccosh}(cx)) - 3c^2 d^3 x (a + \text{barccosh}(cx)) - \frac{d^3(a + \text{barccosh}(cx))}{x}
 \end{aligned}$$

---

3.25.  $\int \frac{(d - c^2 dx^2)^3 (a + \text{barccosh}(cx))}{x^2} dx$

$$\begin{aligned}
 & \downarrow \text{2123} \\
 & \frac{bcd^3\sqrt{c^2x^2-1} \int \left( (c^2x^2-1)^{3/2} c^2 - 3\sqrt{c^2x^2-1}c^2 + \frac{11c^2}{\sqrt{c^2x^2-1}} + \frac{5}{x^2\sqrt{c^2x^2-1}} \right) dx^2}{10\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{5}c^6d^3x^5(a + \\
 & \text{barccosh}(cx)) + c^4d^3x^3(a + \text{barccosh}(cx)) - 3c^2d^3x(a + \text{barccosh}(cx)) - \frac{d^3(a + \text{barccosh}(cx))}{x} \\
 & \downarrow \text{2009} \\
 & -\frac{1}{5}c^6d^3x^5(a + \text{barccosh}(cx)) + c^4d^3x^3(a + \text{barccosh}(cx)) - 3c^2d^3x(a + \text{barccosh}(cx)) - \\
 & \frac{d^3(a + \text{barccosh}(cx))}{x} + \\
 & \frac{bcd^3\sqrt{c^2x^2-1} \left( 10 \arctan \left( \sqrt{c^2x^2-1} \right) + \frac{2}{5}(c^2x^2-1)^{5/2} - 2(c^2x^2-1)^{3/2} + 22\sqrt{c^2x^2-1} \right)}{10\sqrt{cx-1}\sqrt{cx+1}}
 \end{aligned}$$

input `Int[((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x^2,x]`

output `-((d^3*(a + b*ArcCosh[c*x]))/x) - 3*c^2*d^3*x*(a + b*ArcCosh[c*x]) + c^4*d^3*x^3*(a + b*ArcCosh[c*x]) - (c^6*d^3*x^5*(a + b*ArcCosh[c*x]))/5 + (b*c*d^3*Sqrt[-1 + c^2*x^2]*(22*Sqrt[-1 + c^2*x^2] - 2*(-1 + c^2*x^2)^(3/2) + (2*(-1 + c^2*x^2)^(5/2))/5 + 10*ArcTan[Sqrt[-1 + c^2*x^2]]))/(10*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.25.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2113 `Int[(P_x)*((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m])/(a*c + b*d*x^2)^FracPart[m] Int[P_x*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

rule 2123 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol]
:> Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c,
d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

rule 2331 `Int[(Pq_)*(x_)^((m_) + (b_)*(x_)^2)^(p_), x_Symbol] :> Simp[1/2 S
ubst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /;
FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`

rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_
)^2)^(p_), x_Symbol] :> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.25.4 Maple [A] (verified)

Time = 0.45 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.99

method	result
parts	$-d^3 a \left( \frac{c^6 x^5}{5} - c^4 x^3 + 3c^2 x + \frac{1}{x} \right) - d^3 b c \left( \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - c^3 x^3 \operatorname{arccosh}(cx) + 3cx \operatorname{arccosh}(cx) \right)$
derivativedivides	$c \left( -d^3 a \left( \frac{c^5 x^5}{5} - c^3 x^3 + 3cx + \frac{1}{cx} \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - c^3 x^3 \operatorname{arccosh}(cx) + 3cx \operatorname{arccosh}(cx) \right) \right)$
default	$c \left( -d^3 a \left( \frac{c^5 x^5}{5} - c^3 x^3 + 3cx + \frac{1}{cx} \right) - d^3 b \left( \frac{\operatorname{arccosh}(cx) c^5 x^5}{5} - c^3 x^3 \operatorname{arccosh}(cx) + 3cx \operatorname{arccosh}(cx) \right) \right)$

input `int((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)`

output  $-d^3 a \left( \frac{1}{5} c^6 x^5 - c^4 x^3 + 3c^2 x + \frac{1}{x} \right) - d^3 b c \left( \frac{1}{5} \operatorname{arccosh}(cx) c^5 x^5 - c^3 x^3 \operatorname{arccosh}(cx) + 3cx \operatorname{arccosh}(cx) \right) + \frac{1}{25} (cx-1)^{1/2} (cx+1)^{1/2} (-c^4 x^4 (c^2 x^2 - 1)^{1/2} + 7c^2 x^2 (c^2 x^2 - 1)^{1/2} + 5 \arctan(1/(c^2 x^2 - 1)^{1/2}) - 61 (c^2 x^2 - 1)^{1/2}) / (c^2 x^2 - 1)^{1/2}$

---

3.25.  $\int \frac{(d-c^2 dx^2)^3 (a+b \operatorname{arccosh}(cx))}{x^2} dx$

### 3.25.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.38

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^2} dx = \frac{5ac^6 d^3 x^6 - 25ac^4 d^3 x^4 + 75ac^2 d^3 x^2 - 50bcd^3 x \arctan(-cx + \sqrt{c^2 x^2 - 1}) - 5(bc^6 - 5bc^4 + 15bc^2 + 5b) d^3 x \log(-cx + \sqrt{c^2 x^2 - 1}) + 25a^2 d^3 + 5(b^2 c^6 d^3 x^6 - 5b^2 c^4 d^3 x^4 + 15b^2 c^2 d^3 x^2 - (b^2 c^6 - 5b^2 c^4 + 15b^2 c^2 + 5b^2) d^3 x + 5b^2 d^3) \log(cx + \sqrt{c^2 x^2 - 1}) - (b^2 c^5 d^3 x^5 - 7b^2 c^3 d^3 x^3 + 61b^2 c d^3 x) \sqrt{c^2 x^2 - 1}}{x}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x, algorithm="fricas")`

output `-1/25*(5*a*c^6*d^3*x^6 - 25*a*c^4*d^3*x^4 + 75*a*c^2*d^3*x^2 - 50*b*c*d^3*x*arctan(-c*x + sqrt(c^2*x^2 - 1)) - 5*(b*c^6 - 5*b*c^4 + 15*b*c^2 + 5*b)*d^3*x*log(-c*x + sqrt(c^2*x^2 - 1)) + 25*a*d^3 + 5*(b*c^6*d^3*x^6 - 5*b*c^4*d^3*x^4 + 15*b*c^2*d^3*x^2 - (b*c^6 - 5*b*c^4 + 15*b*c^2 + 5*b)*d^3*x + 5*b*d^3)*log(c*x + sqrt(c^2*x^2 - 1)) - (b*c^5*d^3*x^5 - 7*b*c^3*d^3*x^3 + 61*b*c*d^3*x)*sqrt(c^2*x^2 - 1))/x`

### 3.25.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^2} dx = -d^3 \left( \int 3ac^2 dx + \int \left(-\frac{a}{x^2}\right) dx + \int (-3ac^4 x^2) dx + \int ac^6 x^4 dx + \int 3bc^2 \operatorname{acosh}(cx) dx + \int \left(-\frac{b \operatorname{acosh}(cx)}{x^2}\right) dx + \int (-3bc^4 x^2 \operatorname{acosh}(cx)) dx + \int bc^6 x^4 \operatorname{acosh}(cx) dx \right)$$

input `integrate((-c**2*d*x**2+d)**3*(a+b*acosh(c*x))/x**2,x)`

output `-d**3*(Integral(3*a*c**2, x) + Integral(-a/x**2, x) + Integral(-3*a*c**4*x**2, x) + Integral(a*c**6*x**4, x) + Integral(3*b*c**2*acosh(c*x), x) + Integral(-b*acosh(c*x)/x**2, x) + Integral(-3*b*c**4*x**2*acosh(c*x), x) + Integral(b*c**6*x**4*acosh(c*x), x))`

---

3.25.  $\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^2} dx$

### 3.25.7 Maxima [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 231, normalized size of antiderivative = 1.28

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^2} dx$$

$$= -\frac{1}{5} ac^6 d^3 x^5$$

$$- \frac{1}{75} \left( 15 x^5 \operatorname{arccosh}(cx) - \left( \frac{3 \sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4 \sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bc^6 d^3$$

$$+ ac^4 d^3 x^3 + \frac{1}{3} \left( 3 x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2 \sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bc^4 d^3 - 3 ac^2 d^3 x$$

$$- 3 \left( cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1} \right) bcd^3 - \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) bd^3 - \frac{ad^3}{x}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output `-1/5*a*c^6*d^3*x^5 - 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*c^6*d^3 + a*c^4*d^3*x^3 + 1/3*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*c^4*d^3 - 3*a*c^2*d^3*x - 3*(c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*c*d^3 - (c*arcsin(1/(c*abs(x))) + arccosh(c*x)/x)*b*d^3 - a*d^3/x`

### 3.25.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.25.  $\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^2} dx$

**3.25.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x^2,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x^2, x)`

**3.26**  $\int \frac{(d-c^2dx^2)^3(a+b\operatorname{arccosh}(cx))}{x^3} dx$

3.26.1 Optimal result . . . . . 390  
 3.26.2 Mathematica [A] (warning: unable to verify) . . . . . 391  
 3.26.3 Rubi [C] (warning: unable to verify) . . . . . 391  
 3.26.4 Maple [A] (verified) . . . . . 399  
 3.26.5 Fricas [F] . . . . . 399  
 3.26.6 Sympy [F] . . . . . 400  
 3.26.7 Maxima [F] . . . . . 400  
 3.26.8 Giac [F(-2)] . . . . . 401  
 3.26.9 Mupad [F(-1)] . . . . . 401

**3.26.1 Optimal result**

Integrand size = 25, antiderivative size = 267

$$\int \frac{(d - c^2dx^2)^3 (a + b\operatorname{arccosh}(cx))}{x^3} dx = -\frac{3}{32}bc^3d^3x\sqrt{-1 + cx}\sqrt{1 + cx} - \frac{7}{16}bc^3d^3x(-1 + cx)^{3/2}(1 + cx)^{3/2} + \frac{bcd^3(-1 + cx)^{5/2}(1 + cx)^{5/2}}{2x} + \frac{3}{32}bc^2d^3\operatorname{arccosh}(cx) - \frac{3}{2}c^2d^3(1 - c^2x^2)(a + b\operatorname{arccosh}(cx)) - \frac{3}{4}c^2d^3(1 - c^2x^2)^2(a + b\operatorname{arccosh}(cx)) - \frac{d^3(1 - c^2x^2)^2}{4}$$

output

```
-7/16*b*c^3*d^3*x*(c*x-1)^(3/2)*(c*x+1)^(3/2)+1/2*b*c*d^3*(c*x-1)^(5/2)*(c*x+1)^(5/2)/x+3/32*b*c^2*d^3*arccosh(c*x)-3/2*c^2*d^3*(-c^2*x^2+1)*(a+b*arccosh(c*x))-3/4*c^2*d^3*(-c^2*x^2+1)^2*(a+b*arccosh(c*x))-1/2*d^3*(-c^2*x^2+1)^3*(a+b*arccosh(c*x))/x^2-3/2*c^2*d^3*(a+b*arccosh(c*x))^2/b-3*c^2*d^3*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)+3/2*b*c^2*d^3*polylog(2,-1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)-3/32*b*c^3*d^3*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)
```

### 3.26.2 Mathematica [A] (warning: unable to verify)

Time = 0.46 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.13

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx$$

$$= \frac{d^3 \left( -16a + 48ac^4 x^4 - 8ac^6 x^6 + 3bc^3 x^3 \sqrt{\frac{-1+cx}{1+cx}} + 3bc^4 x^4 \sqrt{\frac{-1+cx}{1+cx}} + 2bc^5 x^5 \sqrt{\frac{-1+cx}{1+cx}} + 2bc^6 x^6 \sqrt{\frac{-1+cx}{1+cx}} + 16a^2 \right)}{32x^2}$$

input `Integrate[((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x^3,x]`

output `(d^3*(-16*a + 48*a*c^4*x^4 - 8*a*c^6*x^6 + 3*b*c^3*x^3*sqrt[(-1 + c*x)/(1 + c*x)] + 3*b*c^4*x^4*sqrt[(-1 + c*x)/(1 + c*x)] + 2*b*c^5*x^5*sqrt[(-1 + c*x)/(1 + c*x)] + 2*b*c^6*x^6*sqrt[(-1 + c*x)/(1 + c*x)] + 16*b*c*x*sqrt[-1 + c*x]*sqrt[1 + c*x] - 24*b*c^3*x^3*sqrt[-1 + c*x]*sqrt[1 + c*x] - 48*b*c^2*x^2*ArcCosh[c*x]^2 - 42*b*c^2*x^2*ArcTanh[sqrt[(-1 + c*x)/(1 + c*x)]] - 8*b*ArcCosh[c*x]*(2 - 6*c^4*x^4 + c^6*x^6 + 12*c^2*x^2*Log[1 + E^(-2*ArcCosh[c*x])]) - 96*a*c^2*x^2*Log[x] + 48*b*c^2*x^2*PolyLog[2, -E^(-2*ArcCosh[c*x])]))/(32*x^2)`

### 3.26.3 Rubi [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.43 (sec) , antiderivative size = 370, normalized size of antiderivative = 1.39, number of steps used = 23, number of rules used = 22,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.880$ , Rules used = {6335, 27, 108, 27, 40, 40, 43, 6334, 40, 40, 43, 6334, 40, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx$$

$$\downarrow \text{6335}$$

$$-3c^2 d \int \frac{d^2(1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{2} bcd^3 \int \frac{(cx - 1)^{5/2} (cx + 1)^{5/2}}{x^2} dx - \frac{d^3(1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2}$$

---

3.26.  $\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx$



$$\begin{aligned}
& \downarrow 27 \\
& -3c^2 d^3 \int \frac{(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{2}bcd^3 \int \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x^2} dx - \\
& \quad \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} \\
& \downarrow 108 \\
& -3c^2 d^3 \int \frac{(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx - \\
& \frac{1}{2}bcd^3 \left( \int 5c^2(cx-1)^{3/2}(cx+1)^{3/2} dx - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) - \\
& \quad \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} \\
& \downarrow 27 \\
& -3c^2 d^3 \int \frac{(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx - \\
& \frac{1}{2}bcd^3 \left( 5c^2 \int (cx-1)^{3/2}(cx+1)^{3/2} dx - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) - \\
& \quad \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} \\
& \downarrow 40 \\
& -3c^2 d^3 \int \frac{(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx - \\
& \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \int \sqrt{cx-1}\sqrt{cx+1} dx \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) - \\
& \quad \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} \\
& \downarrow 40 \\
& -3c^2 d^3 \int \frac{(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx - \\
& \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) - \\
& \quad \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} \\
& \downarrow 43
\end{aligned}$$

$$\begin{aligned}
& -3c^2 d^3 \int \frac{(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx - \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \\
& \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) \\
& \quad \downarrow \text{6334} \\
& -3c^2 d^3 \left( \int \frac{(1-c^2x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4}bc \int (cx-1)^{3/2}(cx+1)^{3/2} dx + \frac{1}{4}(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx)) \right. \\
& \quad \left. \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \right. \\
& \left. \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) \right) \\
& \quad \downarrow \text{40} \\
& -3c^2 d^3 \left( \int \frac{(1-c^2x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4}bc \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \int \sqrt{cx-1}\sqrt{cx+1} dx \right) + \frac{1}{4}(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx)) \right. \\
& \quad \left. \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \right. \\
& \left. \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) \right) \\
& \quad \downarrow \text{40} \\
& -3c^2 d^3 \left( \int \frac{(1-c^2x^2) (a + \operatorname{barccosh}(cx))}{x} dx - \frac{1}{4}bc \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) \right) \right. \\
& \quad \left. \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \right. \\
& \left. \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) \right) \\
& \quad \downarrow \text{43} \\
& -3c^2 d^3 \left( \int \frac{(1-c^2x^2) (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{4}(1-c^2x^2)^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} \right. \right. \\
& \quad \left. \left. \frac{d^3(1-c^2x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \right. \right. \\
& \left. \left. \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right) \right) \right) \\
& \quad \downarrow \text{6334}
\end{aligned}$$

$$-3c^2 d^3 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2} bc \int \sqrt{cx-1} \sqrt{cx+1} dx + \frac{1}{4} (1-c^2x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{2} (1-c^2x^2) (a + \operatorname{barccosh}(cx)) \right) - \frac{d^3(1-c^2x^2)^3(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right)$$

↓ 40

$$-3c^2 d^3 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{2} bc \left( \frac{1}{2} x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) + \frac{1}{4} (1-c^2x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{2} (1-c^2x^2) (a + \operatorname{barccosh}(cx)) \right) - \frac{d^3(1-c^2x^2)^3(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right)$$

↓ 43

$$-3c^2 d^3 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x} dx + \frac{1}{4} (1-c^2x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{2} (1-c^2x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{2} bc \left( \frac{1}{2} x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) \right) - \frac{d^3(1-c^2x^2)^3(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right)$$

↓ 6297

$$-3c^2 d^3 \left( \frac{\int -\left( (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{4} (1-c^2x^2)^2 (a + \operatorname{barccosh}(cx)) + \frac{1}{2} (1-c^2x^2) (a + \operatorname{barccosh}(cx)) + \frac{1}{2} bc \left( \frac{1}{2} x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) \right) - \frac{d^3(1-c^2x^2)^3(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx-1)^{5/2}(cx+1)^{5/2}}{x} \right)$$

↓ 25

$$-3c^2 d^3 \left( -\frac{\int (a + \operatorname{barccosh}(cx)) \tanh \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \right. \\ \left. - \frac{d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx - 1)^{5/2} (cx + 1)^{5/2}}{x} \right) \right)$$

↓ 3042

$$-3c^2 d^3 \left( -\frac{\int -i(a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \right. \\ \left. - \frac{d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx - 1)^{5/2} (cx + 1)^{5/2}}{x} \right) \right)$$

↓ 26

$$-3c^2 d^3 \left( \frac{i \int (a + \operatorname{barccosh}(cx)) \tan \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right) d(a + \operatorname{barccosh}(cx))}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \right. \\ \left. - \frac{d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx - 1)^{5/2} (cx + 1)^{5/2}}{x} \right) \right)$$

↓ 4201

$$-3c^2 d^3 \left( \frac{i \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)} (a + \operatorname{barccosh}(cx))}{1 + e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2} i (a + \operatorname{barccosh}(cx))^2 \right)}{b} + \frac{1}{4} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) \right. \\ \left. - \frac{d^3 (1 - c^2 x^2)^3 (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2} bcd^3 \left( 5c^2 \left( \frac{1}{4} x (cx - 1)^{3/2} (cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2} x \sqrt{cx - 1} \sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx - 1)^{5/2} (cx + 1)^{5/2}}{x} \right) \right)$$

↓ 2620

$$-3c^2d^3 \left( \frac{i(2i(\frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)))}{b} - \frac{d^3(1 - c^2x^2)^3(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx - 1)^{5/2}(cx + 1)^{5/2}}{x} \right) \right)$$

↓ 2715

$$-3c^2d^3 \left( \frac{i(2i(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)))}{b} - \frac{d^3(1 - c^2x^2)^3(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx - 1)^{5/2}(cx + 1)^{5/2}}{x} \right) \right)$$

↓ 2838

$$-3c^2d^3 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx)))}{b} - \frac{d^3(1 - c^2x^2)^3(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bcd^3 \left( 5c^2 \left( \frac{1}{4}x(cx - 1)^{3/2}(cx + 1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx - 1}\sqrt{cx + 1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right) - \frac{(cx - 1)^{5/2}(cx + 1)^{5/2}}{x} \right) \right)$$

input `Int(((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x^3,x]`

output `-1/2*(d^3*(1 - c^2*x^2)^3*(a + b*ArcCosh[c*x]))/x^2 - (b*c*d^3*(-((( -1 + c*x)^(5/2)*(1 + c*x)^(5/2))/x) + 5*c^2*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4)))/2 - 3*c^2*d^3((((1 - c^2*x^2)*(a + b*ArcCosh[c*x]))/2 + ((1 - c^2*x^2)^2*(a + b*ArcCosh[c*x]))/4 + (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/2 - (b*c*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/4 + (I*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4)))/b)`

## 3.26.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*(c + d*x)^m/(2*m + 1), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 108 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(m + 1))), x] - Simp[1/(b*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

$$3.26. \int \frac{(d-c^2dx^2)^3(a+b\operatorname{arccosh}(cx))}{x^3} dx$$

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6334 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((d_) + (e_.)*(x_)^2)^(p_.)/(x_), x_Symbol] := Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d)^(p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6335 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (-Simp[b*c*((-d)^(p/(f*(m + 1)))) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x] - Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0] && ILtQ[(m + 1)/2, 0]`

### 3.26.4 Maple [A] (verified)

Time = 1.00 (sec) , antiderivative size = 258, normalized size of antiderivative = 0.97

method	result
derivativedivides	$c^2 \left( -d^3 a \left( \frac{c^4 x^4}{4} - \frac{3c^2 x^2}{2} + 3 \ln(cx) + \frac{1}{2c^2 x^2} \right) - \frac{d^3 b}{2} - \frac{21b d^3 \operatorname{arccosh}(cx)}{32} - 3d^3 b \operatorname{arccosh}(cx) \ln \left( 1 + (cx + \sqrt{c^2 x^2 + d}) \right) \right)$
default	$c^2 \left( -d^3 a \left( \frac{c^4 x^4}{4} - \frac{3c^2 x^2}{2} + 3 \ln(cx) + \frac{1}{2c^2 x^2} \right) - \frac{d^3 b}{2} - \frac{21b d^3 \operatorname{arccosh}(cx)}{32} - 3d^3 b \operatorname{arccosh}(cx) \ln \left( 1 + (cx + \sqrt{c^2 x^2 + d}) \right) \right)$
parts	$-d^3 a \left( \frac{c^6 x^4}{4} - \frac{3c^4 x^2}{2} + \frac{1}{2x^2} + 3c^2 \ln(x) \right) - \frac{d^3 b c^2}{2} - 3d^3 b c^2 \operatorname{arccosh}(cx) \ln \left( 1 + (cx + \sqrt{c^2 x^2 + d}) \right)$

input `int((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)`

output `c^2*(-d^3*a*(1/4*c^4*x^4-3/2*c^2*x^2+3*ln(c*x)+1/2/c^2/x^2)-1/2*d^3*b-21/32*b*d^3*arccosh(c*x)-3*d^3*b*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2))*(c*x+1)^(1/2))^2)+1/2*d^3*b/c/x*(c*x+1)^(1/2)*(c*x-1)^(1/2)-1/4*d^3*b*arccosh(c*x)*c^4*x^4+3/2*d^3*b*arccosh(c*x)*c^2*x^2-1/2*d^3*b*arccosh(c*x)/c^2/x^2+3/2*d^3*b*arccosh(c*x)^2+1/16*d^3*b*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^3*x^3-21/32*b*c*d^3*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)-3/2*d^3*b*polylog(2,-(c*x+(c*x-1)^(1/2))*(c*x+1)^(1/2))^2)`

### 3.26.5 Fracas [F]

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^3} dx = \int -\frac{(c^2 dx^2 - d)^3 (b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x, algorithm="fricas")`

output `integral(-(a*c^6*d^3*x^6 - 3*a*c^4*d^3*x^4 + 3*a*c^2*d^3*x^2 - a*d^3 + (b*c^6*d^3*x^6 - 3*b*c^4*d^3*x^4 + 3*b*c^2*d^3*x^2 - b*d^3)*arccosh(c*x))/x^3, x)`



## 3.26.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx = -d^3 \left( \int \left( -\frac{a}{x^3} \right) dx + \int \frac{3ac^2}{x} dx + \int (-3ac^4 x) dx \right. \\ \left. + \int ac^6 x^3 dx + \int \left( -\frac{b \operatorname{acosh}(cx)}{x^3} \right) dx \right. \\ \left. + \int \frac{3bc^2 \operatorname{acosh}(cx)}{x} dx \right. \\ \left. + \int (-3bc^4 x \operatorname{acosh}(cx)) dx \right. \\ \left. + \int bc^6 x^3 \operatorname{acosh}(cx) dx \right)$$

input `integrate((-c**2*d*x**2+d)**3*(a+b*acosh(c*x))/x**3,x)`

output `-d**3*(Integral(-a/x**3, x) + Integral(3*a*c**2/x, x) + Integral(-3*a*c**4*x, x) + Integral(a*c**6*x**3, x) + Integral(-b*acosh(c*x)/x**3, x) + Integral(3*b*c**2*acosh(c*x)/x, x) + Integral(-3*b*c**4*x*acosh(c*x), x) + Integral(b*c**6*x**3*acosh(c*x), x))`

## 3.26.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int -\frac{(c^2 dx^2 - d)^3 (b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `-1/4*a*c^6*d^3*x^4 + 3/2*a*c^4*d^3*x^2 - 3*a*c^2*d^3*log(x) + 1/2*b*d^3*(sqrt(c^2*x^2 - 1)*c/x - arccosh(c*x)/x^2) - 1/2*a*d^3/x^2 - integrate(b*c^6*d^3*x^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) - 3*b*c^4*d^3*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 3*b*c^2*d^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

**3.26.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.26.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x^3,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x^3, x)`

**3.27**  $\int \frac{(d-c^2dx^2)^3(a+b\operatorname{arccosh}(cx))}{x^4} dx$

3.27.1 Optimal result . . . . . 402  
 3.27.2 Mathematica [A] (verified) . . . . . 403  
 3.27.3 Rubi [A] (warning: unable to verify) . . . . . 403  
 3.27.4 Maple [A] (verified) . . . . . 406  
 3.27.5 Fricas [A] (verification not implemented) . . . . . 407  
 3.27.6 Sympy [F] . . . . . 407  
 3.27.7 Maxima [A] (verification not implemented) . . . . . 408  
 3.27.8 Giac [F(-2)] . . . . . 408  
 3.27.9 Mupad [F(-1)] . . . . . 409

**3.27.1 Optimal result**

Integrand size = 25, antiderivative size = 195

$$\int \frac{(d - c^2dx^2)^3 (a + \operatorname{arccosh}(cx))}{x^4} dx = -\frac{8}{3}bc^3d^3\sqrt{-1 + cx}\sqrt{1 + cx} + \frac{bcd^3\sqrt{-1 + cx}\sqrt{1 + cx}}{6x^2} + \frac{1}{9}bc^3d^3(-1 + cx)^{3/2}(1 + cx)^{3/2} - \frac{d^3(a + \operatorname{arccosh}(cx))}{3x^3} + \frac{3c^2d^3(a + \operatorname{arccosh}(cx))}{x} + 3c^4d^3x(a + \operatorname{arccosh}(cx)) - \frac{1}{3}c^6d^3x^3(a + \operatorname{arccosh}(cx)) - \frac{17}{6}bc^3d^3 \arctan\left(\sqrt{-1 + cx}\sqrt{1 + cx}\right)$$

```
output 1/9*b*c^3*d^3*(c*x-1)^(3/2)*(c*x+1)^(3/2)-1/3*d^3*(a+b*arccosh(c*x))/x^3+3
*c^2*d^3*(a+b*arccosh(c*x))/x+3*c^4*d^3*x*(a+b*arccosh(c*x))-1/3*c^6*d^3*x
^3*(a+b*arccosh(c*x))-17/6*b*c^3*d^3*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))-8
/3*b*c^3*d^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)+1/6*b*c*d^3*(c*x-1)^(1/2)*(c*x+1)
^(1/2)/x^2
```

### 3.27.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.73

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx$$

$$= \frac{d^3 \left( -6a + 54ac^2x^2 + 54ac^4x^4 - 6ac^6x^6 + bcx\sqrt{-1 + cx}\sqrt{1 + cx}(3 - 50c^2x^2 + 2c^4x^4) - 6b(1 - 9c^2x^2 - 9c^4x^4 + c^6x^6) \operatorname{ArcCosh}[cx] + 51b^2c^3x^3 \operatorname{ArcTan}\left[\frac{1}{\sqrt{-1 + cx}\sqrt{1 + cx}}\right] \right)}{18x^3}$$

input `Integrate[((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x^4,x]`

output `(d^3*(-6*a + 54*a*c^2*x^2 + 54*a*c^4*x^4 - 6*a*c^6*x^6 + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(3 - 50*c^2*x^2 + 2*c^4*x^4) - 6*b*(1 - 9*c^2*x^2 - 9*c^4*x^4 + c^6*x^6)*ArcCosh[c*x] + 51*b*c^3*x^3*ArcTan[1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])]))/(18*x^3)`

### 3.27.3 Rubi [A] (warning: unable to verify)

Time = 0.79 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.96, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$ , Rules used = {6336, 27, 2113, 2331, 2124, 27, 1192, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx$$

$$\downarrow \text{6336}$$

$$-bc \int -\frac{d^3(c^6x^6 - 9c^4x^4 - 9c^2x^2 + 1)}{3x^3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^6d^3x^3(a + \operatorname{barccosh}(cx)) + 3c^4d^3x(a + \operatorname{barccosh}(cx)) + \frac{3c^2d^3(a + \operatorname{barccosh}(cx))}{x} - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3}$$

$$\downarrow \text{27}$$

$$\frac{1}{3}bcd^3 \int \frac{c^6x^6 - 9c^4x^4 - 9c^2x^2 + 1}{x^3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^6d^3x^3(a + \operatorname{barccosh}(cx)) + 3c^4d^3x(a + \operatorname{barccosh}(cx)) + \frac{3c^2d^3(a + \operatorname{barccosh}(cx))}{x} - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3}$$

$$\downarrow \text{2113}$$

---

3.27.  $\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx$

$$\begin{aligned}
& \frac{bcd^3\sqrt{c^2x^2-1} \int \frac{c^6x^6-9c^4x^4-9c^2x^2+1}{x^3\sqrt{c^2x^2-1}} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3}c^6d^3x^3(a+\operatorname{barccosh}(cx)) + 3c^4d^3x(a+\operatorname{barccosh}(cx)) + \\
& \quad \frac{3c^2d^3(a+\operatorname{barccosh}(cx))}{x} - \frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{2331} \\
& \frac{bcd^3\sqrt{c^2x^2-1} \int \frac{c^6x^6-9c^4x^4-9c^2x^2+1}{x^4\sqrt{c^2x^2-1}} dx^2}{6\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3}c^6d^3x^3(a+\operatorname{barccosh}(cx)) + 3c^4d^3x(a+\operatorname{barccosh}(cx)) + \\
& \quad \frac{3c^2d^3(a+\operatorname{barccosh}(cx))}{x} - \frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{2124} \\
& \frac{bcd^3\sqrt{c^2x^2-1} \left( \int -\frac{2x^4c^6+18x^2c^4+17c^2}{2x^2\sqrt{c^2x^2-1}} dx^2 + \frac{\sqrt{c^2x^2-1}}{x^2} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3}c^6d^3x^3(a+\operatorname{barccosh}(cx)) + 3c^4d^3x(a + \\
& \quad \operatorname{barccosh}(cx)) + \frac{3c^2d^3(a+\operatorname{barccosh}(cx))}{x} - \frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{27} \\
& \frac{bcd^3\sqrt{c^2x^2-1} \left( \frac{\sqrt{c^2x^2-1}}{x^2} - \frac{1}{2} \int \frac{-2x^4c^6+18x^2c^4+17c^2}{x^2\sqrt{c^2x^2-1}} dx^2 \right)}{6\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3}c^6d^3x^3(a+\operatorname{barccosh}(cx)) + 3c^4d^3x(a + \\
& \quad \operatorname{barccosh}(cx)) + \frac{3c^2d^3(a+\operatorname{barccosh}(cx))}{x} - \frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{1192} \\
& \frac{bcd^3\sqrt{c^2x^2-1} \left( \frac{\sqrt{c^2x^2-1}}{x^2} - \frac{\int \frac{-2c^6x^8+14c^6x^4+33c^6}{x^4+1} d\sqrt{c^2x^2-1}}{c^4} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3}c^6d^3x^3(a+\operatorname{barccosh}(cx)) + \\
& \quad 3c^4d^3x(a+\operatorname{barccosh}(cx)) + \frac{3c^2d^3(a+\operatorname{barccosh}(cx))}{x} - \frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{1467} \\
& \frac{bcd^3\sqrt{c^2x^2-1} \left( \frac{\sqrt{c^2x^2-1}}{x^2} - \frac{\int (-2x^4c^6+\frac{17c^6}{x^4+1}+16c^6) d\sqrt{c^2x^2-1}}{c^4} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3}c^6d^3x^3(a+\operatorname{barccosh}(cx)) + \\
& \quad 3c^4d^3x(a+\operatorname{barccosh}(cx)) + \frac{3c^2d^3(a+\operatorname{barccosh}(cx))}{x} - \frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{2009} \\
& -\frac{1}{3}c^6d^3x^3(a+\operatorname{barccosh}(cx)) + 3c^4d^3x(a+\operatorname{barccosh}(cx)) + \frac{3c^2d^3(a+\operatorname{barccosh}(cx))}{x} - \\
& \quad \frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^3\sqrt{c^2x^2-1} \left( \frac{\sqrt{c^2x^2-1}}{x^2} - \frac{17c^6 \arctan(\sqrt{c^2x^2-1}) - \frac{2}{3}c^6x^6 + 16c^6\sqrt{c^2x^2-1}}{c^4} \right)}{6\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

---

3.27.  $\int \frac{(d-c^2dx^2)^3(a+\operatorname{barccosh}(cx))}{x^4} dx$

input `Int[((d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*(d^3*(a + b*ArcCosh[c*x]))/x^3 + (3*c^2*d^3*(a + b*ArcCosh[c*x]))/x + 3*c^4*d^3*x*(a + b*ArcCosh[c*x]) - (c^6*d^3*x^3*(a + b*ArcCosh[c*x]))/3 + (b*c*d^3*sqrt[-1 + c^2*x^2]*(sqrt[-1 + c^2*x^2]/x^2 - ((-2*c^6*x^6)/3 + 16*c^6*sqrt[-1 + c^2*x^2] + 17*c^6*ArcTan[sqrt[-1 + c^2*x^2]]/c^4))/(6*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.27.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1192 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4]^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2113 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m])/(a*c + b*d*x^2)^FracPart[m] Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

```
rule 2124 Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :
> With[{Qx = PolynomialQuotient[Px, a + b*x, x], R = PolynomialRemainder[Px,
a + b*x, x]}, Simp[R*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((m + 1)*(b*c -
a*d))), x] + Simp[1/((m + 1)*(b*c - a*d)) Int[(a + b*x)^(m + 1)*(c + d*x)
^n*ExpandToSum[(m + 1)*(b*c - a*d)*Qx - d*R*(m + n + 2), x], x], x]] /; Fr
eeQ[{a, b, c, d, n}, x] && PolyQ[Px, x] && LtQ[m, -1] && (IntegerQ[m] || !
ILtQ[n, -1])
```

```
rule 2331 Int[(Pq_)*(x_)^((m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 S
ubst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /;
FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]
```

```
rule 6336 Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(x_
)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.27.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.00

method	result
parts	$-d^3 a \left( \frac{x^3 c^6}{3} - 3c^4 x - \frac{3c^2}{x} + \frac{1}{3x^3} \right) - d^3 b c^3 \left( \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - 3cx \operatorname{arccosh}(cx) + \frac{\operatorname{arccosh}(cx)}{3c^3 x^3} \right)$
derivativedivides	$c^3 \left( -d^3 a \left( \frac{c^3 x^3}{3} - 3cx + \frac{1}{3c^3 x^3} - \frac{3}{cx} \right) - d^3 b \left( \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - 3cx \operatorname{arccosh}(cx) + \frac{\operatorname{arccosh}(cx)}{3c^3 x^3} \right) \right)$
default	$c^3 \left( -d^3 a \left( \frac{c^3 x^3}{3} - 3cx + \frac{1}{3c^3 x^3} - \frac{3}{cx} \right) - d^3 b \left( \frac{c^3 x^3 \operatorname{arccosh}(cx)}{3} - 3cx \operatorname{arccosh}(cx) + \frac{\operatorname{arccosh}(cx)}{3c^3 x^3} \right) \right)$

```
input int((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)
```

$$3.27. \int \frac{(d-c^2 dx^2)^3 (a+b \operatorname{arccosh}(cx))}{x^4} dx$$

output 
$$-d^3 a (1/3 x^3 c^6 - 3 c^4 x - 3 c^2/x + 1/3/x^3) - d^3 b c^3 (1/3 c^3 x^3 \operatorname{arccosh}(cx) - 3 c x \operatorname{arccosh}(cx) + 1/3 c^3/x^3 \operatorname{arccosh}(cx) - 3 \operatorname{arccosh}(cx)/c/x - 1/18 (cx-1)^{1/2} (cx+1)^{1/2} (2 c^4 x^4 (c^2 x^2 - 1)^{1/2} + 51 \operatorname{arctan}(1/(c^2 x^2 - 1)^{1/2}) c^2 x^2 - 50 c^2 x^2 (c^2 x^2 - 1)^{1/2} + 3 (c^2 x^2 - 1)^{1/2})/c^2 x^2 / (c^2 x^2 - 1)^{1/2})$$

### 3.27.5 Fricas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.30

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx = \frac{6 ac^6 d^3 x^6 - 54 ac^4 d^3 x^4 + 102 bc^3 d^3 x^3 \operatorname{arctan}(-cx + \sqrt{c^2 x^2 - 1}) - 54 ac^2 d^3 x^2 - 6 (bc^6 - 9 bc^4 - 9 bc^2 + \dots)}{x^4}$$

input `integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")`

output 
$$-1/18*(6*a*c^6*d^3*x^6 - 54*a*c^4*d^3*x^4 + 102*b*c^3*d^3*x^3*\operatorname{arctan}(-c*x + \operatorname{sqrt}(c^2*x^2 - 1)) - 54*a*c^2*d^3*x^2 - 6*(b*c^6 - 9*b*c^4 - 9*b*c^2 + b)*d^3*x^3*\log(-c*x + \operatorname{sqrt}(c^2*x^2 - 1)) + 6*a*d^3 + 6*(b*c^6*d^3*x^6 - 9*b*c^4*d^3*x^4 - 9*b*c^2*d^3*x^2 - (b*c^6 - 9*b*c^4 - 9*b*c^2 + b)*d^3*x^3 + b*d^3)*\log(c*x + \operatorname{sqrt}(c^2*x^2 - 1)) - (2*b*c^5*d^3*x^5 - 50*b*c^3*d^3*x^3 + 3*b*c*d^3*x)*\operatorname{sqrt}(c^2*x^2 - 1))/x^3$$

### 3.27.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx = -d^3 \left( \int (-3ac^4) dx + \int \left(-\frac{a}{x^4}\right) dx + \int \frac{3ac^2}{x^2} dx + \int ac^6 x^2 dx + \int (-3bc^4 \operatorname{acosh}(cx)) dx + \int \left(-\frac{b \operatorname{acosh}(cx)}{x^4}\right) dx + \int \frac{3bc^2 \operatorname{acosh}(cx)}{x^2} dx + \int bc^6 x^2 \operatorname{acosh}(cx) dx \right)$$

input `integrate((-c**2*d*x**2+d)**3*(a+b*acosh(c*x))/x**4,x)`

3.27. 
$$\int \frac{(d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx$$



```
output -d**3*(Integral(-3*a*c**4, x) + Integral(-a/x**4, x) + Integral(3*a*c**2/x
**2, x) + Integral(a*c**6*x**2, x) + Integral(-3*b*c**4*acosh(c*x), x) + I
ntegral(-b*acosh(c*x)/x**4, x) + Integral(3*b*c**2*acosh(c*x)/x**2, x) + I
ntegral(b*c**6*x**2*acosh(c*x), x))
```

### 3.27.7 Maxima [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 208, normalized size of antiderivative = 1.07

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx$$

$$= -\frac{1}{3} ac^6 d^3 x^3 - \frac{1}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bc^6 d^3 + 3ac^4 d^3 x$$

$$+ 3 \left( cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1} \right) bc^3 d^3 + 3 \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) bc^2 d^3$$

$$- \frac{1}{6} \left( \left( c^2 \arcsin \left( \frac{1}{c|x|} \right) - \frac{\sqrt{c^2 x^2 - 1}}{x^2} \right) c + \frac{2 \operatorname{arccosh}(cx)}{x^3} \right) bd^3 + \frac{3ac^2 d^3}{x} - \frac{ad^3}{3x^3}$$

```
input integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")
```

```
output -1/3*a*c^6*d^3*x^3 - 1/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^
2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*c^6*d^3 + 3*a*c^4*d^3*x + 3*(c*x*arccosh(c
*x) - sqrt(c^2*x^2 - 1))*b*c^3*d^3 + 3*(c*arcsin(1/(c*abs(x))) + arccosh(c
*x)/x)*b*c^2*d^3 - 1/6*((c^2*arcsin(1/(c*abs(x))) - sqrt(c^2*x^2 - 1)/x^2)
*c + 2*arccosh(c*x)/x^3)*b*d^3 + 3*a*c^2*d^3/x - 1/3*a*d^3/x^3
```

### 3.27.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c^2*d*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

---

3.27.  $\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx$

**3.27.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x^4,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^3)/x^4, x)`

### 3.28 $\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx$

3.28.1	Optimal result	410
3.28.2	Mathematica [A] (verified)	411
3.28.3	Rubi [C] (verified)	411
3.28.4	Maple [A] (verified)	415
3.28.5	Fricas [F]	416
3.28.6	Sympy [F]	416
3.28.7	Maxima [F]	416
3.28.8	Giac [F(-2)]	417
3.28.9	Mupad [F(-1)]	417

#### 3.28.1 Optimal result

Integrand size = 25, antiderivative size = 158

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{d - c^2dx^2} dx = \frac{11b\sqrt{-1 + cx}\sqrt{1 + cx}}{9c^5d} + \frac{bx^2\sqrt{-1 + cx}\sqrt{1 + cx}}{9c^3d} - \frac{x(a + \operatorname{arccosh}(cx))}{c^4d} - \frac{x^3(a + \operatorname{arccosh}(cx))}{3c^2d} + \frac{2(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^5d} + \frac{b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{c^5d} - \frac{b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{c^5d}$$

output

```
-x*(a+b*arccosh(c*x))/c^4/d-1/3*x^3*(a+b*arccosh(c*x))/c^2/d+2*(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^5/d+b*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^5/d-b*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^5/d+11/9*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5/d+1/9*b*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d
```

### 3.28.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.44

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \frac{18acx + 6ac^3x^3 - 18b\sqrt{\frac{-1+cx}{1+cx}} - 18bcx\sqrt{\frac{-1+cx}{1+cx}} - 4b\sqrt{-1+cx}\sqrt{1+cx} - 2bc^2x^2\sqrt{-1+cx}\sqrt{1+cx} + 1}{d - c^2 dx^2}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2), x]`

output `-1/18*(18*a*c*x + 6*a*c^3*x^3 - 18*b*Sqrt[(-1 + c*x)/(1 + c*x)] - 18*b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)] - 4*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 2*b*c^2*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 18*b*c*x*ArcCosh[c*x] + 6*b*c^3*x^3*ArcCosh[c*x] - 9*b*ArcCosh[c*x]^2 - 18*b*ArcCosh[c*x]*Log[1 + E^(-ArcCosh[c*x])] + 18*b*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] + 9*a*Log[1 - c*x] - 9*a*Log[1 + c*x] + 18*b*PolyLog[2, -E^(-ArcCosh[c*x])] + 18*b*PolyLog[2, E^ArcCosh[c*x]])/(c^5*d)`

### 3.28.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.86 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.18, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.520$ , Rules used = {6353, 27, 111, 27, 83, 6353, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx \\ & \quad \downarrow \text{6353} \\ & \frac{\int \frac{x^2(a + \operatorname{barccosh}(cx))}{d(1 - c^2 x^2)} dx}{c^2} + \frac{b \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3cd} - \frac{x^3(a + \operatorname{barccosh}(cx))}{3c^2 d} \\ & \quad \downarrow \text{27} \\ & \frac{\int \frac{x^2(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d} + \frac{b \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3cd} - \frac{x^3(a + \operatorname{barccosh}(cx))}{3c^2 d} \end{aligned}$$

---

3.28.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx$

$$\begin{aligned}
& \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx + \frac{b\left(\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)}{3cd} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} \\
& \quad \downarrow 111 \\
& \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx + \frac{b\left(\frac{2\int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)}{3cd} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} \\
& \quad \downarrow 27 \\
& \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} + \frac{b\left(\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)}{3cd} \\
& \quad \downarrow 83 \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx}{c^2} + \frac{b\int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{x(a+\operatorname{barccosh}(cx))}{c^2} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} + \\
& \quad \frac{c^2d}{b\left(\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)} \\
& \quad \downarrow 6353 \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{x(a+\operatorname{barccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} + \\
& \quad \frac{c^2d}{b\left(\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)} \\
& \quad \downarrow 83 \\
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx+1}} \operatorname{darccosh}(cx)}{c^3} - \frac{x(a+\operatorname{barccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} + \\
& \quad \frac{c^2d}{b\left(\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)} \\
& \quad \downarrow 6318 \\
& -\frac{\int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{c^3} - \frac{x(a+\operatorname{barccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} - \\
& \quad \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} + \frac{c^2d}{b\left(\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)} \\
& \quad \downarrow 3042 \\
& \quad \downarrow 26
\end{aligned}$$

---

3.28.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))}{d-c^2dx^2} dx$

$$\frac{-\frac{i \int (a + \operatorname{barccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{c^3} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{\frac{x^3(a + \operatorname{barccosh}(cx))}{3c^2d} + \frac{b\left(\frac{c^2d}{3c^4}\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)}{3cd}}$$

↓ 4670

$$\frac{-\frac{i \int \log(1 - e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) - i b \int \log(1 + e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))}{c^3} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2}}{\frac{x^3(a + \operatorname{barccosh}(cx))}{3c^2d} + \frac{b\left(\frac{c^2d}{3c^4}\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)}{3cd}}$$

↓ 2715

$$\frac{-\frac{i \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) d e^{\operatorname{arccosh}(cx)} - i b \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) d e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))}{c^3} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2}}{\frac{x^3(a + \operatorname{barccosh}(cx))}{3c^2d} + \frac{b\left(\frac{c^2d}{3c^4}\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)}{3cd}}$$

↓ 2838

$$\frac{-\frac{i \left(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx)) + i b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - i b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})\right)}{c^3} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{\frac{x^3(a + \operatorname{barccosh}(cx))}{3c^2d} + \frac{b\left(\frac{c^2d}{3c^4}\frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2}\right)}{3cd}}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2),x]`

output `(b*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2)))/(3*c*d) - (x^3*(a + b*ArcCosh[c*x]))/(3*c^2*d) + ((b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^3 - (x*(a + b*ArcCosh[c*x]))/c^2 - (I*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c^3)/(c^2*d)`

## 3.28.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx)] /; FreeQ[b, x]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 111 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^(p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.28.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.47

method	result
derivativedivides	$\frac{a\left(\frac{c^3x^3}{3} + cx + \frac{\ln(cx-1)}{2} - \frac{\ln(cx+1)}{2}\right)}{d} - \frac{b \operatorname{arccosh}(cx)c^3x^3}{3d} - \frac{b \operatorname{arccosh}(cx)cx}{d} - \frac{b \operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1}\sqrt{cx+1})}{d} + \frac{b \operatorname{arccosh}(cx) \ln(1-cx+\sqrt{cx-1}\sqrt{cx+1})}{d}$
default	$-\frac{a\left(\frac{c^3x^3}{3} + cx + \frac{\ln(cx-1)}{2} - \frac{\ln(cx+1)}{2}\right)}{d} - \frac{b \operatorname{arccosh}(cx)c^3x^3}{3d} - \frac{b \operatorname{arccosh}(cx)cx}{d} - \frac{b \operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1}\sqrt{cx+1})}{d} + \frac{b \operatorname{arccosh}(cx) \ln(1-cx+\sqrt{cx-1}\sqrt{cx+1})}{d}$
parts	$-\frac{a\left(\frac{1}{3}\frac{x^3c^2+x}{c^4} - \frac{\ln(cx+1)}{2c^5} + \frac{\ln(cx-1)}{2c^5}\right)}{d} + \frac{b \operatorname{arccosh}(cx) \ln(1+cx+\sqrt{cx-1}\sqrt{cx+1})}{dc^5} - \frac{b \operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1}\sqrt{cx+1})}{dc^5}$

input `int(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x,method=_RETURNVERBOSE)`

output `1/c^5*(-a/d*(1/3*c^3*x^3+c*x+1/2*ln(c*x-1)-1/2*ln(c*x+1))-1/3*b/d*arccosh(c*x)*c^3*x^3-b/d*arccosh(c*x)*c*x-b/d*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+b/d*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/9*b/d*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^2*x^2-b/d*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+b/d*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+11/9*b/d*(c*x-1)^(1/2)*(c*x+1)^(1/2))`

3.28. 
$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx$$



## 3.28.5 Fracas [F]

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^4}{c^2 dx^2 - d} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="fracas")`

output `integral(-(b*x^4*arccosh(c*x) + a*x^4)/(c^2*d*x^2 - d), x)`

## 3.28.6 Sympy [F]

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = -\int \frac{ax^4}{c^2 x^2 - 1} dx + \int \frac{bx^4 \operatorname{acosh}(cx)}{c^2 x^2 - 1} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(-c**2*d*x**2+d),x)`

output `-(Integral(a*x**4/(c**2*x**2 - 1), x) + Integral(b*x**4*acosh(c*x)/(c**2*x**2 - 1), x))/d`

## 3.28.7 Maxima [F]

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^4}{c^2 dx^2 - d} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `1/72*(4*c^4*(2*(c^2*x^3 + 3*x)/(c^8*d) - 3*log(c*x + 1)/(c^9*d) + 3*log(c*x - 1)/(c^9*d)) + 36*c^2*(2*x/(c^6*d) - log(c*x + 1)/(c^7*d) + log(c*x - 1)/(c^7*d)) + 648*c*integrate(1/12*x*log(c*x - 1)/(c^6*d*x^2 - c^4*d), x) - 3*(4*(2*c^3*x^3 + 6*c*x - 3*log(c*x + 1) + 3*log(c*x - 1))*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 3*log(c*x + 1)^2 + 6*log(c*x + 1)*log(c*x - 1))/(c^5*d) + 72*integrate(-1/6*(2*c^3*x^3 + 6*c*x - 3*log(c*x + 1) + 3*log(c*x - 1))/(c^7*d*x^3 - c^5*d*x + (c^6*d*x^2 - c^4*d)*sqrt(c*x + 1))*sqrt(c*x - 1), x) - 216*integrate(1/12*log(c*x - 1)/(c^6*d*x^2 - c^4*d), x))*b - 1/6*a*(2*(c^2*x^3 + 3*x)/(c^4*d) - 3*log(c*x + 1)/(c^5*d) + 3*log(c*x - 1)/(c^5*d))`

**3.28.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.28.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{d - c^2 dx^2} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2),x)`

output `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2), x)`

### 3.29 $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx$

3.29.1	Optimal result	418
3.29.2	Mathematica [A] (warning: unable to verify)	418
3.29.3	Rubi [C] (verified)	419
3.29.4	Maple [A] (verified)	423
3.29.5	Fricas [F]	423
3.29.6	Sympy [F]	424
3.29.7	Maxima [F]	424
3.29.8	Giac [F(-2)]	424
3.29.9	Mupad [F(-1)]	425

#### 3.29.1 Optimal result

Integrand size = 25, antiderivative size = 140

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{d - c^2dx^2} dx = \frac{bx\sqrt{-1 + cx}\sqrt{1 + cx}}{4c^3d} + \frac{\operatorname{arccosh}(cx)}{4c^4d} - \frac{x^2(a + \operatorname{arccosh}(cx))}{2c^2d} + \frac{(a + \operatorname{arccosh}(cx))^2}{2bc^4d} - \frac{(a + \operatorname{arccosh}(cx)) \log(1 - e^{2\operatorname{arccosh}(cx)})}{c^4d} - \frac{b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2c^4d}$$

output  $1/4*b*\operatorname{arccosh}(c*x)/c^4/d-1/2*x^2*(a+b*\operatorname{arccosh}(c*x))/c^2/d+1/2*(a+b*\operatorname{arccosh}(c*x))^2/b/c^4/d-(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/c^4/d-1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/c^4/d+1/4*b*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/d$

#### 3.29.2 Mathematica [A] (warning: unable to verify)

Time = 0.20 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.08

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{d - c^2dx^2} dx = \frac{2c^2x^2(a + \operatorname{arccosh}(cx)) - \frac{2(a+b\operatorname{arccosh}(cx))^2}{b} - b\left(cx\sqrt{-1 + cx}\sqrt{1 + cx} + 2\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right) + 4(a -$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2),x]`

output `-1/4*(2*c^2*x^2*(a + b*ArcCosh[c*x]) - (2*(a + b*ArcCosh[c*x])^2)/b - b*(c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]]) + 4*(a + b*ArcCosh[c*x])*Log[1 - E^ArcCosh[c*x]] + 4*(a + b*ArcCosh[c*x])*Log[1 + E^ArcCosh[c*x]] + 4*b*PolyLog[2, -E^ArcCosh[c*x]] + 4*b*PolyLog[2, E^ArcCosh[c*x]])/(c^4*d)`

### 3.29.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.04, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.480$ , Rules used = {6353, 27, 101, 43, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{d(1 - c^2 x^2)} dx}{c^2} + \frac{b \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2cd} - \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d} + \frac{b \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2cd} - \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d} \\
 & \quad \downarrow \text{101} \\
 & \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d} + \frac{b \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2cd} - \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d} \\
 & \quad \downarrow \text{43} \\
 & \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d} - \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2cd} \\
 & \quad \downarrow \text{6328}
 \end{aligned}$$

---

3.29.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx$

$$\begin{aligned}
& \frac{\int \frac{cx(a+\operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{c^4d} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2d} + \frac{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}{2cd} \\
& \quad \downarrow \text{3042} \\
& \frac{\int -i(a+\operatorname{barccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{\frac{c^4d}{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2d} + \\
& \quad \downarrow \text{26} \\
& \frac{i \int (a+\operatorname{barccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{\frac{c^4d}{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2d} + \\
& \quad \downarrow \text{4199} \\
& \frac{i\left(2i \int -\frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b}\right)}{\frac{c^4d}{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2d} + \\
& \quad \downarrow \text{25} \\
& \frac{i\left(-2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b}\right)}{\frac{c^4d}{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2d} + \\
& \quad \downarrow \text{2620} \\
& \frac{i\left(-2i\left(\frac{1}{2}b \int \log(1-e^{2\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - \frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx))\right) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b}\right)}{\frac{c^4d}{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2d} + \\
& \quad \downarrow \text{2715} \\
& \frac{i\left(-2i\left(\frac{1}{4}b \int e^{-2\operatorname{arccosh}(cx)} \log(1-e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx))\right) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b}\right)}{\frac{c^4d}{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2d} +
\end{aligned}$$


---

3.29.  $\int \frac{x^3(a+\operatorname{barccosh}(cx))}{d-c^2dx^2} dx$

↓ 2838

$$\frac{i\left(-2i\left(-\frac{1}{2}\log\left(1-e^{2\operatorname{arccosh}(cx)}\right)\right)(a+\operatorname{arccosh}(cx))-\frac{1}{4}b\operatorname{PolyLog}\left(2,e^{2\operatorname{arccosh}(cx)}\right)\right)-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}}{\frac{x^2(a+\operatorname{arccosh}(cx))}{2c^2d}+\frac{b\left(\frac{c^4d}{2c^3}+\frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}{2cd}}$$

input `Int[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2),x]`

output `-1/2*(x^2*(a + b*ArcCosh[c*x]))/(c^2*d) + (b*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/(2*c*d) + (I*(((1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])])) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/(c^4*d)`

### 3.29.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 43 `Int[1/(sqrt[(a_) + (b_.)*(x_)]*sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 101 `Int[((a_.) + (b_.)*(x_))^(n_)*((c_.) + (d_.)*(x_))^(m_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6328 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6353 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.29.4 Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.55

method	result
derivativedivides	$-\frac{a\left(\frac{c^2x^2}{2} + \frac{\ln(cx-1)}{2} + \frac{\ln(cx+1)}{2}\right)}{d} + \frac{b \operatorname{arccosh}(cx)^2}{2d} + \frac{b\sqrt{cx+1}\sqrt{cx-1}cx}{4d} - \frac{b \operatorname{arccosh}(cx)c^2x^2}{2d} + \frac{b \operatorname{arccosh}(cx)}{4d} - \frac{b \operatorname{arccosh}(cx)\ln(1-cx)}{4d}$
default	$-\frac{a\left(\frac{c^2x^2}{2} + \frac{\ln(cx-1)}{2} + \frac{\ln(cx+1)}{2}\right)}{d} + \frac{b \operatorname{arccosh}(cx)^2}{2d} + \frac{b\sqrt{cx+1}\sqrt{cx-1}cx}{4d} - \frac{b \operatorname{arccosh}(cx)c^2x^2}{2d} + \frac{b \operatorname{arccosh}(cx)}{4d} - \frac{b \operatorname{arccosh}(cx)\ln(1-cx)}{4d}$
parts	$-\frac{ax^2}{2dc^2} - \frac{a \ln(c^2x^2-1)}{2dc^4} + \frac{b \operatorname{arccosh}(cx)^2}{2dc^4} - \frac{b \operatorname{arccosh}(cx)x^2}{2dc^2} + \frac{bx\sqrt{cx-1}\sqrt{cx+1}}{4c^3d} + \frac{b \operatorname{arccosh}(cx)}{4c^4d} - \frac{b \operatorname{arccosh}(cx)\ln(1-cx)}{4c^4d}$

```
input int(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x,method=_RETURNVERBOSE)
```

```
output 1/c^4*(-a/d*(1/2*c^2*x^2+1/2*ln(c*x-1)+1/2*ln(c*x+1))+1/2*b/d*arccosh(c*x)^2+1/4*b/d*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c*x-1/2*b/d*arccosh(c*x)*c^2*x^2+1/4*b/d*arccosh(c*x)-b/d*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-b/d*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-b/d*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-b/d*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)))
```

### 3.29.5 Fricas [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arccosh}(cx) + a)x^3}{c^2 dx^2 - d} dx$$

```
input integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="fricas")
```

```
output integral(-(b*x^3*arccosh(c*x) + a*x^3)/(c^2*d*x^2 - d), x)
```



### 3.29.6 Sympy [F]

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = -\int \frac{\frac{ax^3}{c^2 x^2 - 1}}{d} dx + \int \frac{bx^3 \operatorname{acosh}(cx)}{c^2 x^2 - 1} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(-c**2*d*x**2+d),x)`

output `-(Integral(a*x**3/(c**2*x**2 - 1), x) + Integral(b*x**3*acosh(c*x)/(c**2*x**2 - 1), x))/d`

### 3.29.7 Maxima [F]

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^3}{c^2 dx^2 - d} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `-1/2*a*(x^2/(c^2*d) + log(c^2*x^2 - 1)/(c^4*d)) + 1/8*b*((2*c^2*x^2 - 4*(c^2*x^2 + log(c*x + 1) + log(c*x - 1))*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) + 2*(log(c*x - 1) + 1)*log(c*x + 1) + log(c*x + 1)^2 + log(c*x - 1)^2 + 2*log(c*x - 1))/(c^4*d) - 8*integrate(1/2*(c^2*x^2 + log(c*x + 1) + log(c*x - 1))/(c^6*d*x^3 - c^4*d*x + (c^5*d*x^2 - c^3*d)*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1))), x)`

### 3.29.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.29.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx$

**3.29.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{d - c^2 dx^2} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2),x)`output `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2), x)`

### 3.30 $\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx$

3.30.1	Optimal result	426
3.30.2	Mathematica [A] (warning: unable to verify)	426
3.30.3	Rubi [C] (verified)	427
3.30.4	Maple [A] (verified)	430
3.30.5	Fricas [F]	430
3.30.6	Sympy [F]	430
3.30.7	Maxima [F]	431
3.30.8	Giac [F]	431
3.30.9	Mupad [F(-1)]	432

#### 3.30.1 Optimal result

Integrand size = 25, antiderivative size = 102

$$\int \frac{x^2(a + b\operatorname{arccosh}(cx))}{d - c^2dx^2} dx = \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}}{c^3d} - \frac{x(a + b\operatorname{arccosh}(cx))}{c^2d} + \frac{2(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^3d} + \frac{b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{c^3d} - \frac{b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{c^3d}$$

output `-x*(a+b*arccosh(c*x))/c^2/d+2*(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d+b*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d-b*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d+b*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d`

#### 3.30.2 Mathematica [A] (warning: unable to verify)

Time = 0.20 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.52

$$\int \frac{x^2(a + b\operatorname{arccosh}(cx))}{d - c^2dx^2} dx = \frac{-2acx + 2b\sqrt{\frac{-1+cx}{1+cx}} + 2bcx\sqrt{\frac{-1+cx}{1+cx}} - 2bcx\operatorname{arccosh}(cx) + b\operatorname{arccosh}(cx)^2 + 2b\operatorname{arccosh}(cx)\log(1 + e^{-\operatorname{arccosh}(cx)})}{d - c^2dx^2}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2),x]`

output  $(-2*a*c*x + 2*b*\sqrt{(-1 + c*x)/(1 + c*x)} + 2*b*c*x*\sqrt{(-1 + c*x)/(1 + c*x)} - 2*b*c*x*ArcCosh[c*x] + b*ArcCosh[c*x]^2 + 2*b*ArcCosh[c*x]*Log[1 + E^{-ArcCosh[c*x]}] - 2*b*ArcCosh[c*x]*Log[1 - E^{-ArcCosh[c*x]}] - a*Log[1 - c*x] + a*Log[1 + c*x] - 2*b*PolyLog[2, -E^{-ArcCosh[c*x]}] - 2*b*PolyLog[2, E^{-ArcCosh[c*x]}])/(2*c^3*d)$

### 3.30.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.56 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$ , Rules used = {6353, 27, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{d(1 - c^2 x^2)} dx}{c^2} + \frac{b \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{cd} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2 x^2} dx}{c^2 d} + \frac{b \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{cd} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} \\
 & \quad \downarrow \text{83} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2 x^2} dx}{c^2 d} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d} \\
 & \quad \downarrow \text{6318} \\
 & - \frac{\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{c^3 d} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{\int i(a + \operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{c^3 d} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d}
 \end{aligned}$$

---

3.30.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx$

$$\begin{aligned}
& \downarrow 26 \\
& \frac{i \int (a + \operatorname{barccosh}(cx)) \csc(i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{c^3 d} - \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d} \\
& \downarrow 4670 \\
& \frac{i(ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d} \\
& \downarrow 2715 \\
& \frac{i(ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d} \\
& \downarrow 2838 \\
& \frac{i(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{c^3 d} \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2),x]`

output `(b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c^3*d) - (x*(a + b*ArcCosh[c*x]))/(c^2*d) - (I*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/(c^3*d)`

### 3.30.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`
- rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.30.4 Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.77

method	result
derivativedivides	$\frac{a \left( cx + \frac{\ln(cx-1)}{2} - \frac{\ln(cx+1)}{2} \right)}{d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{d} + \frac{b \operatorname{arccosh}(cx) \ln(1+cx+\sqrt{cx-1}\sqrt{cx+1})}{d} - \frac{b \operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1}\sqrt{cx+1})}{c^3 d}$
default	$-\frac{a \left( cx + \frac{\ln(cx-1)}{2} - \frac{\ln(cx+1)}{2} \right)}{d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{d} + \frac{b \operatorname{arccosh}(cx) \ln(1+cx+\sqrt{cx-1}\sqrt{cx+1})}{d} - \frac{b \operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1}\sqrt{cx+1})}{c^3 d}$
parts	$-\frac{a \left( \frac{x}{c^2} - \frac{\ln(cx+1)}{2c^3} + \frac{\ln(cx-1)}{2c^3} \right)}{d} - \frac{b \operatorname{arccosh}(cx)x}{dc^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3 d} - \frac{b \operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1}\sqrt{cx+1})}{dc^3} +$

input `int(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x,method=_RETURNVERBOSE)`

output `1/c^3*(-a/d*(c*x+1/2*ln(c*x-1)-1/2*ln(c*x+1))+b/d*(c*x-1)^(1/2)*(c*x+1)^(1/2)+b/d*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-b/d*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-b/d*arccosh(c*x)*c*x-b/d*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+b/d*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)))`

### 3.30.5 Fricas [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arccosh}(cx) + a)x^2}{c^2 dx^2 - d} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="fricas")`

output `integral(-(b*x^2*arccosh(c*x) + a*x^2)/(c^2*d*x^2 - d), x)`

### 3.30.6 Sympy [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = -\int \frac{ax^2}{c^2 x^2 - 1} dx + \int \frac{bx^2 \operatorname{acosh}(cx)}{c^2 x^2 - 1} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(-c**2*d*x**2+d),x)`

---

3.30.  $\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx$

output `-(Integral(a*x**2/(c**2*x**2 - 1), x) + Integral(b*x**2*acosh(c*x)/(c**2*x**2 - 1), x))/d`

### 3.30.7 Maxima [F]

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^2}{c^2 dx^2 - d} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `1/8*(4*c^2*(2*x/(c^4*d) - log(c*x + 1)/(c^5*d) + log(c*x - 1)/(c^5*d)) + 2*4*c*integrate(1/4*x*log(c*x - 1)/(c^4*d*x^2 - c^2*d), x) - (4*(2*c*x - log(c*x + 1) + log(c*x - 1))*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + log(c*x + 1)^2 + 2*log(c*x + 1)*log(c*x - 1))/(c^3*d) + 8*integrate(-1/2*(2*c*x - log(c*x + 1) + log(c*x - 1))/(c^5*d*x^3 - c^3*d*x + (c^4*d*x^2 - c^2*d)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) - 8*integrate(1/4*log(c*x - 1)/(c^4*d*x^2 - c^2*d), x))*b - 1/2*a*(2*x/(c^2*d) - log(c*x + 1)/(c^3*d) + log(c*x - 1)/(c^3*d))`

### 3.30.8 Giac [F]

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^2}{c^2 dx^2 - d} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)*x^2/(c^2*d*x^2 - d), x)`



**3.30.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{d - c^2 dx^2} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2),x)`output `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2), x)`

### 3.31 $\int \frac{x(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx$

3.31.1	Optimal result	433
3.31.2	Mathematica [A] (verified)	433
3.31.3	Rubi [C] (verified)	434
3.31.4	Maple [A] (verified)	436
3.31.5	Fricas [F]	437
3.31.6	Sympy [F]	437
3.31.7	Maxima [F]	437
3.31.8	Giac [F]	438
3.31.9	Mupad [F(-1)]	438

#### 3.31.1 Optimal result

Integrand size = 23, antiderivative size = 74

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{d - c^2dx^2} dx = \frac{(a + \operatorname{arccosh}(cx))^2}{2bc^2d} - \frac{(a + \operatorname{arccosh}(cx)) \log(1 - e^{2\operatorname{arccosh}(cx)})}{c^2d} - \frac{b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2c^2d}$$

output  $\frac{1}{2}*(a+b*\operatorname{arccosh}(c*x))^2/b/c^2/d - (a+b*\operatorname{arccosh}(c*x))*\ln(1 - (c*x + (c*x - 1)^{1/2})*(c*x + 1)^{1/2})/c^2/d - 1/2*b*\operatorname{polylog}(2, (c*x + (c*x - 1)^{1/2})*(c*x + 1)^{1/2})/c^2/d$

#### 3.31.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.15

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{d - c^2dx^2} dx = \frac{(a + \operatorname{arccosh}(cx))(a + \operatorname{arccosh}(cx) - 2b \log(1 - e^{\operatorname{arccosh}(cx)}) - 2b \log(1 + e^{\operatorname{arccosh}(cx)})) - 2b^2 \operatorname{PolyLog}(2, -E^{\operatorname{arccosh}(cx)}) - 2b^2 \operatorname{PolyLog}(2, E^{\operatorname{arccosh}(cx)})}{2bc^2d}$$

input `Integrate[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2), x]`

output  $((a + b*\operatorname{ArcCosh}[c*x])*(a + b*\operatorname{ArcCosh}[c*x] - 2*b*\operatorname{Log}[1 - E^{\operatorname{ArcCosh}[c*x]}] - 2*b*\operatorname{Log}[1 + E^{\operatorname{ArcCosh}[c*x]}]) - 2*b^2*\operatorname{PolyLog}[2, -E^{\operatorname{ArcCosh}[c*x]}] - 2*b^2*\operatorname{PolyLog}[2, E^{\operatorname{ArcCosh}[c*x]}])/(2*b*c^2*d)$

### 3.31.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.01, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$ , Rules used = {6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx \\
 & \quad \downarrow \text{6328} \\
 & \frac{\int \frac{cx(a + \operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{c^2 d} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -i(a + \operatorname{barccosh}(cx)) \tan\left(i \operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{c^2 d} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \int (a + \operatorname{barccosh}(cx)) \tan\left(i \operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{c^2 d} \\
 & \quad \downarrow \text{4199} \\
 & \frac{i \left( 2i \int -\frac{e^{2 \operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 - e^{2 \operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{c^2 d} \\
 & \quad \downarrow \text{25} \\
 & \frac{i \left( -2i \int \frac{e^{2 \operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 - e^{2 \operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{c^2 d} \\
 & \quad \downarrow \text{2620} \\
 & \frac{i \left( -2i \left( \frac{1}{2} b \int \log(1 - e^{2 \operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} \log(1 - e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) \right) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{c^2 d} \\
 & \quad \downarrow \text{2715}
 \end{aligned}$$

---

3.31.  $\int \frac{x(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx$

$$i \left( \frac{-2i \left( \frac{1}{4} b \int e^{-2 \operatorname{arccosh}(cx)} \log(1 - e^{2 \operatorname{arccosh}(cx)}) d e^{2 \operatorname{arccosh}(cx)} - \frac{1}{2} \log(1 - e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{arccosh}(cx)) \right) - \frac{i(a + b \operatorname{arccosh}(cx))^2}{2b}}{c^2 d} \right)$$

↓ 2838

$$i \left( \frac{-2i \left( -\frac{1}{2} \log(1 - e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{arccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog}(2, e^{2 \operatorname{arccosh}(cx)}) \right) - \frac{i(a + b \operatorname{arccosh}(cx))^2}{2b}}{c^2 d} \right)$$

input `Int[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2),x]`

output `(I*(((−1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(−1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x]]) - (b*PolyLog[2, E^(2*ArcCosh[c*x]]))/4)))/(c^2*d)`

### 3.31.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

### 3.31.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.85

method	result
parts	$-\frac{a \ln(c^2 x^2 - 1)}{2d c^2} - \frac{b \left( -\frac{\operatorname{arccosh}(cx)^2}{2} + \operatorname{arccosh}(cx) \ln(1 - cx - \sqrt{cx-1} \sqrt{cx+1}) + \operatorname{polylog}(2, cx + \sqrt{cx-1} \sqrt{cx+1}) + \operatorname{arccosh}(cx) \right)}{d c^2}$
derivativedivides	$\frac{a \left( \frac{\ln(cx-1)}{2} + \frac{\ln(cx+1)}{2} \right)}{d} - \frac{b \left( -\frac{\operatorname{arccosh}(cx)^2}{2} + \operatorname{arccosh}(cx) \ln(1 - cx - \sqrt{cx-1} \sqrt{cx+1}) + \operatorname{polylog}(2, cx + \sqrt{cx-1} \sqrt{cx+1}) + \operatorname{arccosh}(cx) \right)}{c^2 d}$
default	$-\frac{a \left( \frac{\ln(cx-1)}{2} + \frac{\ln(cx+1)}{2} \right)}{d} - \frac{b \left( -\frac{\operatorname{arccosh}(cx)^2}{2} + \operatorname{arccosh}(cx) \ln(1 - cx - \sqrt{cx-1} \sqrt{cx+1}) + \operatorname{polylog}(2, cx + \sqrt{cx-1} \sqrt{cx+1}) + \operatorname{arccosh}(cx) \right)}{c^2 d}$

input `int(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d), x, method=_RETURNVERBOSE)`

output `-1/2*a/d/c^2*ln(c^2*x^2-1)-b/d/c^2*(-1/2*arccosh(c*x)^2+arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)))`

---

3.31.  $\int \frac{x(a+b\operatorname{arccosh}(cx))}{d-c^2 dx^2} dx$

**3.31.5 Fracas [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x}{c^2 dx^2 - d} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="fricas")`

output `integral(-(b*x*arccosh(c*x) + a*x)/(c^2*d*x^2 - d), x)`

**3.31.6 Sympy [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = -\frac{\int \frac{ax}{c^2 x^2 - 1} dx + \int \frac{bx \operatorname{acosh}(cx)}{c^2 x^2 - 1} dx}{d}$$

input `integrate(x*(a+b*acosh(c*x))/(-c**2*d*x**2+d),x)`

output `-(Integral(a*x/(c**2*x**2 - 1), x) + Integral(b*x*acosh(c*x)/(c**2*x**2 - 1), x))/d`

**3.31.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x}{c^2 dx^2 - d} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `-1/8*b*((4*(log(c*x + 1) + log(c*x - 1))*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) - log(c*x + 1)^2 - 2*log(c*x + 1)*log(c*x - 1) - log(c*x - 1)^2)/(c^2*d) + 8*integrate(1/2*(log(c*x + 1) + log(c*x - 1))/(c^4*d*x^3 - c^2*d*x + (c^3*d*x^2 - c*d)*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1))), x) - 1/2*a*log(c^2*d*x^2 - d)/(c^2*d)`

**3.31.8 Giac [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arccosh}(cx) + a)x}{c^2 dx^2 - d} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)*x/(c^2*d*x^2 - d), x)`

**3.31.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{d - c^2 dx^2} dx$$

input `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2),x)`

output `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2), x)`

### 3.32 $\int \frac{a+b\operatorname{arccosh}(cx)}{d-c^2dx^2} dx$

3.32.1	Optimal result	439
3.32.2	Mathematica [A] (verified)	439
3.32.3	Rubi [C] (verified)	440
3.32.4	Maple [C] (verified)	442
3.32.5	Fricas [F]	442
3.32.6	Sympy [F]	443
3.32.7	Maxima [F]	443
3.32.8	Giac [F]	443
3.32.9	Mupad [F(-1)]	444

#### 3.32.1 Optimal result

Integrand size = 22, antiderivative size = 59

$$\int \frac{a + b\operatorname{arccosh}(cx)}{d - c^2dx^2} dx = \frac{2(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{cd} + \frac{b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{cd} - \frac{b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{cd}$$

output `2*(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c/d+b*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/c/d-b*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c/d`

#### 3.32.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.08

$$\int \frac{a + b\operatorname{arccosh}(cx)}{d - c^2dx^2} dx = \frac{-((a + b\operatorname{arccosh}(cx)) (\log(1 - e^{\operatorname{arccosh}(cx)}) - \log(1 + e^{\operatorname{arccosh}(cx)}))) + b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{cd}$$

input `Integrate[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2),x]`

output `((-(a + b*ArcCosh[c*x]))*(Log[1 - E^ArcCosh[c*x]] - Log[1 + E^ArcCosh[c*x]])) + b*PolyLog[2, -E^ArcCosh[c*x]] - b*PolyLog[2, E^ArcCosh[c*x]]/(c*d)`



### 3.32.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + b \operatorname{arccosh}(cx)}{d - c^2 dx^2} dx \\
 & \quad \downarrow \text{6318} \\
 & \frac{\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d \operatorname{arccosh}(cx)}{cd} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int i(a + b \operatorname{arccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{cd} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \int (a + b \operatorname{arccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{cd} \\
 & \quad \downarrow \text{4670} \\
 & \frac{i(ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)))}{cd} \\
 & \quad \downarrow \text{2715} \\
 & \frac{i(ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)))}{cd} \\
 & \quad \downarrow \text{2838} \\
 & \frac{i(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{cd}
 \end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2), x]`

output  $((-I)*((2*I)*(a + b*\text{ArcCosh}[c*x])*\text{ArcTanh}[E^{\text{ArcCosh}[c*x]}] + I*b*\text{PolyLog}[2, -E^{\text{ArcCosh}[c*x]}] - I*b*\text{PolyLog}[2, E^{\text{ArcCosh}[c*x]}]))/(c*d)$

### 3.32.3.1 Defintions of rubi rules used

rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F_{x_}), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 2715  $\text{Int}[\text{Log}[(a_ + (b_)*((F_)^{((e_)*((c_ + (d_)*(x_)))^n_))}] , x\_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n)], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x\} \ \&\& \ \text{GtQ}[a, 0]$

rule 2838  $\text{Int}[\text{Log}[(c_)*((d_ + (e_)*(x_)^{n_})]/(x_)), x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n/n, x] /; \text{FreeQ}\{c, d, e, n\}, x\} \ \&\& \ \text{EqQ}[c*d, 1]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4670  $\text{Int}[\text{csc}[(e_ + (\text{Complex}[0, fz_])*(f_)*(x_))*((c_ + (d_)*(x_))^m_)], x\_Symbol] \rightarrow \text{Simp}[-2*(c + d*x)^m*(\text{ArcTanh}[E^{((-I)*e + f*fz*x)}]/(f*fz*I)), x] + (-\text{Simp}[d*(m/(f*fz*I)) \text{Int}[(c + d*x)^{m-1}*\text{Log}[1 - E^{((-I)*e + f*fz*x)}]], x], x] + \text{Simp}[d*(m/(f*fz*I)) \text{Int}[(c + d*x)^{m-1}*\text{Log}[1 + E^{((-I)*e + f*fz*x)}]], x], x) /; \text{FreeQ}\{c, d, e, f, fz\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$

rule 6318  $\text{Int}[(a_ + \text{ArcCosh}[(c_)*(x_)]*(b_))^n_]/((d_ + (e_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[-(c*d)^{-1} \text{Subst}[\text{Int}[(a + b*x)^n*\text{Csch}[x], x], x, \text{ArcCosh}[c*x]], x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[n, 0]$

### 3.32.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.33 (sec) , antiderivative size = 180, normalized size of antiderivative = 3.05

method	result
derivativedivides	$\frac{\frac{a \operatorname{arctanh}(cx)}{d} - \frac{b \left( -\operatorname{arctanh}(cx) \operatorname{arccosh}(cx) - \frac{2i \left( \operatorname{arctanh}(cx) \ln \left( 1 + \frac{i(cx+1)}{\sqrt{-c^2x^2+1}} \right) - \operatorname{arctanh}(cx) \ln \left( 1 - \frac{i(cx+1)}{\sqrt{-c^2x^2+1}} \right) + \operatorname{dilog} \left( 1 + \frac{i}{\sqrt{-c^2x^2-1}} \right) \right)}{c^2x^2-1}}{c}}{d}$
default	$\frac{\frac{a \operatorname{arctanh}(cx)}{d} - \frac{b \left( -\operatorname{arctanh}(cx) \operatorname{arccosh}(cx) - \frac{2i \left( \operatorname{arctanh}(cx) \ln \left( 1 + \frac{i(cx+1)}{\sqrt{-c^2x^2+1}} \right) - \operatorname{arctanh}(cx) \ln \left( 1 - \frac{i(cx+1)}{\sqrt{-c^2x^2+1}} \right) + \operatorname{dilog} \left( 1 + \frac{i}{\sqrt{-c^2x^2-1}} \right) \right)}{c^2x^2-1}}{c}}{d}$
parts	$\frac{a \ln(cx+1)}{2dc} - \frac{a \ln(cx-1)}{2dc} - \frac{b \left( -\operatorname{arctanh}(cx) \operatorname{arccosh}(cx) - \frac{2i \left( \operatorname{arctanh}(cx) \ln \left( 1 + \frac{i(cx+1)}{\sqrt{-c^2x^2+1}} \right) - \operatorname{arctanh}(cx) \ln \left( 1 - \frac{i(cx+1)}{\sqrt{-c^2x^2+1}} \right) + \operatorname{dilog} \left( 1 + \frac{i}{\sqrt{-c^2x^2-1}} \right) \right)}{c}}{dc}$

input `int((a+b*arccosh(c*x))/(-c^2*d*x^2+d), x, method=_RETURNVERBOSE)`

output `1/c*(a/d*arctanh(c*x)-b/d*(-arctanh(c*x)*arccosh(c*x)-2*I*(arctanh(c*x)*ln(1+I*(c*x+1)/(-c^2*x^2+1)^(1/2))-arctanh(c*x)*ln(1-I*(c*x+1)/(-c^2*x^2+1)^(1/2))+dilog(1+I*(c*x+1)/(-c^2*x^2+1)^(1/2))-dilog(1-I*(c*x+1)/(-c^2*x^2+1)^(1/2)))*(-c^2*x^2+1)^(1/2)*(1/2*c*x+1/2)^(1/2)*(1/2*c*x-1/2)^(1/2)/(c^2*x^2-1))`

### 3.32.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{d - c^2 dx^2} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{c^2 dx^2 - d} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d), x, algorithm="fricas")`

output `integral(-(b*arccosh(c*x) + a)/(c^2*d*x^2 - d), x)`

### 3.32.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{d - c^2 dx^2} dx = -\int \frac{a}{c^2 x^2 - 1} dx + \int \frac{b \operatorname{arcosh}(cx)}{c^2 x^2 - 1} dx$$

input `integrate((a+b*acosh(c*x))/(-c**2*d*x**2+d),x)`

output `-(Integral(a/(c**2*x**2 - 1), x) + Integral(b*acosh(c*x)/(c**2*x**2 - 1), x))/d`

### 3.32.7 Maxima [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{d - c^2 dx^2} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{c^2 dx^2 - d} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `1/8*b*((4*(log(c*x + 1) - log(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) - log(c*x + 1)^2 - 2*log(c*x + 1)*log(c*x - 1))/(c*d) + 8*integrate(1/4*(3*c*x - 1)*log(c*x - 1)/(c^2*d*x^2 - d), x) + 8*integrate(1/2*(log(c*x + 1) - log(c*x - 1))/(c^3*d*x^3 - c*d*x + (c^2*d*x^2 - d)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) + 1/2*a*(log(c*x + 1)/(c*d) - log(c*x - 1)/(c*d))`

### 3.32.8 Giac [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{d - c^2 dx^2} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{c^2 dx^2 - d} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/(c^2*d*x^2 - d), x)`

**3.32.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{d - c^2 dx^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{d - c^2 dx^2} dx$$

input `int((a + b*acosh(c*x))/(d - c^2*d*x^2), x)`output `int((a + b*acosh(c*x))/(d - c^2*d*x^2), x)`

### 3.33 $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)} dx$

3.33.1	Optimal result . . . . .	445
3.33.2	Mathematica [B] (verified) . . . . .	445
3.33.3	Rubi [C] (verified) . . . . .	446
3.33.4	Maple [A] (verified) . . . . .	448
3.33.5	Fricas [F] . . . . .	449
3.33.6	Sympy [F] . . . . .	449
3.33.7	Maxima [F] . . . . .	449
3.33.8	Giac [F] . . . . .	450
3.33.9	Mupad [F(-1)] . . . . .	450

#### 3.33.1 Optimal result

Integrand size = 25, antiderivative size = 61

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x(d - c^2dx^2)} dx = \frac{2(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d} + \frac{b \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{2d} - \frac{b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2d}$$

```
output 2*(a+b*arccosh(c*x))*arctanh((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/d+1/2*b*
polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/d-1/2*b*polylog(2,(c*x+(c*
x-1)^(1/2)*(c*x+1)^(1/2))^2)/d
```

#### 3.33.2 Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 129 vs. 2(61) = 122.

Time = 0.27 (sec) , antiderivative size = 129, normalized size of antiderivative = 2.11

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x(d - c^2dx^2)} dx = \frac{-b\operatorname{arccosh}(cx)^2 - 2b\operatorname{arccosh}(cx) \log(1 + e^{-2\operatorname{arccosh}(cx)}) + 2b\operatorname{arccosh}(cx) \log(1 + e^{-\operatorname{arccosh}(cx)}) + 2b\operatorname{arccosh}(cx) \log(1 + e^{\operatorname{arccosh}(cx)})}{d}$$

```
input Integrate[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)),x]
```

output 
$$\frac{-1/2*(-(b*\text{ArcCosh}[c*x]^2) - 2*b*\text{ArcCosh}[c*x]*\text{Log}[1 + E^{(-2*\text{ArcCosh}[c*x])}] + 2*b*\text{ArcCosh}[c*x]*\text{Log}[1 + E^{(-\text{ArcCosh}[c*x])}] + 2*b*\text{ArcCosh}[c*x]*\text{Log}[1 - E^{\text{ArcCosh}[c*x]}] - 2*a*\text{Log}[x] + a*\text{Log}[1 - c^2*x^2] + b*\text{PolyLog}[2, -E^{(-2*\text{ArcCosh}[c*x])}] - 2*b*\text{PolyLog}[2, -E^{(-\text{ArcCosh}[c*x])}] + 2*b*\text{PolyLog}[2, E^{\text{ArcCosh}[c*x]}])}{d}$$

### 3.33.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.07, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$ , Rules used = {6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)} dx \\ & \quad \downarrow \text{6331} \\ & \frac{\int \frac{a + b \operatorname{arccosh}(cx)}{cx \sqrt{\frac{cx-1}{cx+1}}(cx+1)} d \operatorname{arccosh}(cx)}{d} \\ & \quad \downarrow \text{5984} \\ & \frac{2 \int (a + b \operatorname{arccosh}(cx)) \operatorname{csch}(2 \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{d} \\ & \quad \downarrow \text{3042} \\ & \frac{2 \int i(a + b \operatorname{arccosh}(cx)) \operatorname{csc}(2i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{d} \\ & \quad \downarrow \text{26} \\ & \frac{2i \int (a + b \operatorname{arccosh}(cx)) \operatorname{csc}(2i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{d} \\ & \quad \downarrow \text{4670} \\ & \frac{2i \left( \frac{1}{2} i b \int \log(1 - e^{2 \operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) - \frac{1}{2} i b \int \log(1 + e^{2 \operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + i \operatorname{arctanh}(e^{2 \operatorname{arccosh}(cx)}) \right)}{d} \\ & \quad \downarrow \text{2715} \end{aligned}$$

---

3.33. 
$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)} dx$$

$$\frac{2i\left(\frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1 - e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1 + e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)}\right)}{d}$$

↓ 2838

$$\frac{2i\left(i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) (a + b\operatorname{arccosh}(cx)) + \frac{1}{4}ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4}ib \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})\right)}{d}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)),x]`

output `((-2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])]) + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])])/d`

### 3.33.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`



rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^(n, x), x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[n, 0]`

### 3.33.4 Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 191, normalized size of antiderivative = 3.13

method	result
parts	$-\frac{a\left(\frac{\ln(cx+1)}{2}-\ln(x)+\frac{\ln(cx-1)}{2}\right)}{d} - \frac{b\left(-\operatorname{arccosh}(cx)\ln\left(1+(cx+\sqrt{cx-1}\sqrt{cx+1})^2\right)-\frac{\operatorname{polylog}\left(2,-(cx+\sqrt{cx-1}\sqrt{cx+1})^2\right)}{2}\right)}{d}$
derivativedivides	$-\frac{a\left(-\ln(cx)+\frac{\ln(cx+1)}{2}+\frac{\ln(cx-1)}{2}\right)}{d} - \frac{b\left(-\operatorname{arccosh}(cx)\ln\left(1+(cx+\sqrt{cx-1}\sqrt{cx+1})^2\right)-\frac{\operatorname{polylog}\left(2,-(cx+\sqrt{cx-1}\sqrt{cx+1})^2\right)}{2}\right)}{d}$
default	$-\frac{a\left(-\ln(cx)+\frac{\ln(cx+1)}{2}+\frac{\ln(cx-1)}{2}\right)}{d} - \frac{b\left(-\operatorname{arccosh}(cx)\ln\left(1+(cx+\sqrt{cx-1}\sqrt{cx+1})^2\right)-\frac{\operatorname{polylog}\left(2,-(cx+\sqrt{cx-1}\sqrt{cx+1})^2\right)}{2}\right)}{d}$

input `int((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d),x,method=_RETURNVERBOSE)`

output `-a/d*(1/2*ln(c*x+1)-ln(x)+1/2*ln(c*x-1))-b/d*(-arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)-1/2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)+arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)))`

**3.33.5 Fracas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d),x, algorithm="fricas")`

output `integral(-(b*arccosh(c*x) + a)/(c^2*d*x^3 - d*x), x)`

**3.33.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)} dx = -\int \frac{a}{c^2 x^3 - x} dx + \int \frac{b \operatorname{acosh}(cx)}{c^2 x^3 - x} dx$$

input `integrate((a+b*acosh(c*x))/x/(-c**2*d*x**2+d),x)`

output `-(Integral(a/(c**2*x**3 - x), x) + Integral(b*acosh(c*x)/(c**2*x**3 - x), x))/d`

**3.33.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `-1/2*a*(log(c*x + 1)/d + log(c*x - 1)/d - 2*log(x)/d) - b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(c^2*d*x^3 - d*x), x)`

**3.33.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d),x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)*x), x)`

**3.33.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(d - c^2 dx^2)} dx$$

input `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)),x)`

output `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)), x)`

### 3.34 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d-c^2dx^2)} dx$

3.34.1	Optimal result	451
3.34.2	Mathematica [A] (verified)	451
3.34.3	Rubi [C] (verified)	452
3.34.4	Maple [A] (verified)	455
3.34.5	Fricas [F]	455
3.34.6	Sympy [F]	456
3.34.7	Maxima [F]	456
3.34.8	Giac [F]	456
3.34.9	Mupad [F(-1)]	457

#### 3.34.1 Optimal result

Integrand size = 25, antiderivative size = 95

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2(d - c^2dx^2)} dx = -\frac{a + \operatorname{arccosh}(cx)}{dx} + \frac{bc \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d} + \frac{2c(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{d} + \frac{bc \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{d} - \frac{bc \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{d}$$

output  $(-a-b*\operatorname{arccosh}(c*x))/d/x+b*c*\arctan((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d+2*c*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d+b*c*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d-b*c*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d$

#### 3.34.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.39

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2(d - c^2dx^2)} dx = \frac{-\frac{a+b\operatorname{arccosh}(cx)}{x} + \frac{bc\sqrt{-1+c^2x^2} \arctan(\frac{\sqrt{-1+c^2x^2}}{\sqrt{-1+cx}\sqrt{1+cx}})}{\sqrt{-1+cx}\sqrt{1+cx}} - c(a + \operatorname{arccosh}(cx)) \log(1 - e^{\operatorname{arccosh}(cx)}) + c(a + \operatorname{arccosh}(cx)) \log(1 + e^{\operatorname{arccosh}(cx)})}{d}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)),x]`

output `((-(a + b*ArcCosh[c*x])/x) + (b*c*Sqrt[-1 + c^2*x^2]*ArcTan[Sqrt[-1 + c^2*x^2]])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - c*(a + b*ArcCosh[c*x])*Log[1 - E^ArcCosh[c*x]] + c*(a + b*ArcCosh[c*x])*Log[1 + E^ArcCosh[c*x]] + b*c*PolyLog[2, -E^ArcCosh[c*x]] - b*c*PolyLog[2, E^ArcCosh[c*x]])/d`

### 3.34.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.58 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6347, 27, 103, 218, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{x^2(d - c^2 dx^2)} dx \\
 & \quad \downarrow \text{6347} \\
 & c^2 \int \frac{a + \operatorname{barccosh}(cx)}{d(1 - c^2 x^2)} dx + \frac{bc \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{d} - \frac{a + \operatorname{barccosh}(cx)}{dx} \\
 & \quad \downarrow \text{27} \\
 & \frac{c^2 \int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2 x^2} dx}{d} + \frac{bc \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{d} - \frac{a + \operatorname{barccosh}(cx)}{dx} \\
 & \quad \downarrow \text{103} \\
 & \frac{c^2 \int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2 x^2} dx}{d} + \frac{bc^2 \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1})}{d} - \frac{a + \operatorname{barccosh}(cx)}{dx} \\
 & \quad \downarrow \text{218} \\
 & \frac{c^2 \int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2 x^2} dx}{d} - \frac{a + \operatorname{barccosh}(cx)}{dx} + \frac{bc \arctan(\sqrt{cx-1}\sqrt{cx+1})}{d} \\
 & \quad \downarrow \text{6318} \\
 & -\frac{c \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{d} - \frac{a + \operatorname{barccosh}(cx)}{dx} + \frac{bc \arctan(\sqrt{cx-1}\sqrt{cx+1})}{d}
 \end{aligned}$$

$$\begin{aligned}
& \downarrow 3042 \\
& \frac{c \int i(a + \operatorname{barccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d} - \frac{a + \operatorname{barccosh}(cx)}{dx} + \\
& \frac{bc \operatorname{arctan}(\sqrt{cx-1}\sqrt{cx+1})}{d} \\
& \downarrow 26 \\
& \frac{ic \int (a + \operatorname{barccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d} - \frac{a + \operatorname{barccosh}(cx)}{dx} + \\
& \frac{bc \operatorname{arctan}(\sqrt{cx-1}\sqrt{cx+1})}{d} \\
& \downarrow 4670 \\
& \frac{ic(ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \\
& \frac{a + \operatorname{barccosh}(cx)}{dx} + \frac{bc \operatorname{arctan}(\sqrt{cx-1}\sqrt{cx+1})}{d} \\
& \downarrow 2715 \\
& \frac{ic(ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \\
& \frac{a + \operatorname{barccosh}(cx)}{dx} + \frac{bc \operatorname{arctan}(\sqrt{cx-1}\sqrt{cx+1})}{d} \\
& \downarrow 2838 \\
& \frac{ic(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d} - \\
& \frac{a + \operatorname{barccosh}(cx)}{dx} + \frac{bc \operatorname{arctan}(\sqrt{cx-1}\sqrt{cx+1})}{d}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)),x]`

output `-((a + b*ArcCosh[c*x])/(d*x)) + (b*c*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/d - (I*c*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/d`

## 3.34.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 103 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/(f*fz*I)], x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6318 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

```
rule 6347 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]
```

### 3.34.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.51

method	result
parts	$-\frac{a\left(\frac{1}{x} - \frac{c \ln(cx+1)}{2} + \frac{c \ln(cx-1)}{2}\right)}{d} - \frac{bc\left(\frac{\operatorname{arccosh}(cx)}{cx} - 2 \arctan(cx + \sqrt{cx-1} \sqrt{cx+1}) - \operatorname{dilog}(cx + \sqrt{cx-1} \sqrt{cx+1}) - \operatorname{dilog}(cx - \sqrt{cx-1} \sqrt{cx+1})\right)}{d}$
derivativedivides	$c\left(-\frac{a\left(\frac{1}{cx} - \frac{\ln(cx+1)}{2} + \frac{\ln(cx-1)}{2}\right)}{d} - \frac{b\left(\frac{\operatorname{arccosh}(cx)}{cx} - 2 \arctan(cx + \sqrt{cx-1} \sqrt{cx+1}) - \operatorname{dilog}(cx + \sqrt{cx-1} \sqrt{cx+1}) - \operatorname{dilog}(cx - \sqrt{cx-1} \sqrt{cx+1})\right)}{d}\right)$
default	$c\left(-\frac{a\left(\frac{1}{cx} - \frac{\ln(cx+1)}{2} + \frac{\ln(cx-1)}{2}\right)}{d} - \frac{b\left(\frac{\operatorname{arccosh}(cx)}{cx} - 2 \arctan(cx + \sqrt{cx-1} \sqrt{cx+1}) - \operatorname{dilog}(cx + \sqrt{cx-1} \sqrt{cx+1}) - \operatorname{dilog}(cx - \sqrt{cx-1} \sqrt{cx+1})\right)}{d}\right)$

```
input int((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d),x,method=_RETURNVERBOSE)
```

```
output -a/d*(1/x-1/2*c*ln(c*x+1)+1/2*c*ln(c*x-1))-b/d*c*(arccosh(c*x)/c/x-2*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-dilog(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-dilog(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))
```

### 3.34.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)x^2} dx$$

```
input integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d),x, algorithm="fracas")
```

```
output integral(-(b*arccosh(c*x) + a)/(c^2*d*x^4 - d*x^2), x)
```

---

3.34.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)} dx$



**3.34.6 Sympy [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)} dx = -\frac{\int \frac{a}{c^2 x^4 - x^2} dx + \int \frac{b \operatorname{arccosh}(cx)}{c^2 x^4 - x^2} dx}{d}$$

input `integrate((a+b*acosh(c*x))/x**2/(-c**2*d*x**2+d),x)`

output `-(Integral(a/(c**2*x**4 - x**2), x) + Integral(b*acosh(c*x)/(c**2*x**4 - x**2), x))/d`

**3.34.7 Maxima [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `1/8*(24*c^3*integrate(1/4*x*log(c*x - 1)/(c^2*d*x^2 - d), x) - 4*c^2*(log(c*x + 1)/(c*d) - log(c*x - 1)/(c*d)) - 8*c^2*integrate(1/4*log(c*x - 1)/(c^2*d*x^2 - d), x) - (c*x*log(c*x + 1)^2 + 2*c*x*log(c*x + 1)*log(c*x - 1) - 4*(c*x*log(c*x + 1) - c*x*log(c*x - 1) - 2)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)))/(d*x) + 8*integrate(1/2*(c^2*x*log(c*x + 1) - c^2*x*log(c*x - 1) - 2*c)/(c^3*d*x^4 - c*d*x^2 + (c^2*d*x^3 - d*x)*sqrt(c*x + 1))*sqrt(c*x - 1)), x)*b + 1/2*a*(c*log(c*x + 1)/d - c*log(c*x - 1)/d - 2/(d*x))`

**3.34.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d),x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)*x^2), x)`

---

3.34.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)} dx$

**3.34.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2 (d - c^2 dx^2)} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)),x)`output `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)), x)`

### 3.35 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)} dx$

3.35.1	Optimal result	458
3.35.2	Mathematica [A] (verified)	458
3.35.3	Rubi [C] (verified)	459
3.35.4	Maple [A] (verified)	462
3.35.5	Fricas [F]	463
3.35.6	Sympy [F]	463
3.35.7	Maxima [F]	463
3.35.8	Giac [F]	464
3.35.9	Mupad [F(-1)]	464

#### 3.35.1 Optimal result

Integrand size = 25, antiderivative size = 118

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^3(d - c^2dx^2)} dx = \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{2dx} - \frac{a + b\operatorname{arccosh}(cx)}{2dx^2} + \frac{2c^2(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d} + \frac{bc^2 \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{2d} - \frac{bc^2 \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2d}$$

output `1/2*(-a-b*arccosh(c*x))/d/x^2+2*c^2*(a+b*arccosh(c*x))*arctanh((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d+1/2*b*c^2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d-1/2*b*c^2*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d+1/2*b*c*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/x`

#### 3.35.2 Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.80

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^3(d - c^2dx^2)} dx = \frac{a - bcx\sqrt{-1 + cx}\sqrt{1 + cx} + b\operatorname{arccosh}(cx) - 2bc^2x^2\operatorname{arccosh}(cx)^2 - 2bc^2x^2\operatorname{arccosh}(cx)\log(1 + e^{-2\operatorname{arccosh}(cx)})}{d}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)),x]`

output `-1/2*(a - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + b*ArcCosh[c*x] - 2*b*c^2*x^2*ArcCosh[c*x]^2 - 2*b*c^2*x^2*ArcCosh[c*x]*Log[1 + E^(-2*ArcCosh[c*x])] + 2*b*c^2*x^2*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] + 2*b*c^2*x^2*ArcCosh[c*x]*Log[1 + E^ArcCosh[c*x]] - 2*a*c^2*x^2*Log[x] + a*c^2*x^2*Log[1 - c^2*x^2] + b*c^2*x^2*PolyLog[2, -E^(-2*ArcCosh[c*x])] + 2*b*c^2*x^2*PolyLog[2, -E^ArcCosh[c*x]] + 2*b*c^2*x^2*PolyLog[2, E^ArcCosh[c*x]])/(d*x^2)`

### 3.35.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.99, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6347, 27, 106, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{x^3(d - c^2 dx^2)} dx \\
 & \quad \downarrow 6347 \\
 & c^2 \int \frac{a + \operatorname{barccosh}(cx)}{dx(1 - c^2 x^2)} dx + \frac{bc \int \frac{1}{x^2 \sqrt{cx-1} \sqrt{cx+1}} dx}{2d} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} \\
 & \quad \downarrow 27 \\
 & \frac{c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2 x^2)} dx}{d} + \frac{bc \int \frac{1}{x^2 \sqrt{cx-1} \sqrt{cx+1}} dx}{2d} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} \\
 & \quad \downarrow 106 \\
 & \frac{c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2 x^2)} dx}{d} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \frac{bc \sqrt{cx-1} \sqrt{cx+1}}{2dx} \\
 & \quad \downarrow 6331 \\
 & - \frac{c^2 \int \frac{a + \operatorname{barccosh}(cx)}{cx \sqrt{\frac{cx-1}{cx+1}} (cx+1)} d \operatorname{arccosh}(cx)}{d} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \frac{bc \sqrt{cx-1} \sqrt{cx+1}}{2dx} \\
 & \quad \downarrow 5984
 \end{aligned}$$

---

3.35.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d - c^2 dx^2)} dx$

$$\begin{aligned}
& \frac{2c^2 \int (a + \operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2dx} \\
& \quad \downarrow \text{3042} \\
& \frac{2c^2 \int i(a + \operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \\
& \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2dx} \\
& \quad \downarrow \text{26} \\
& \frac{2ic^2 \int (a + \operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \\
& \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2dx} \\
& \quad \downarrow \text{4670} \\
& \frac{2ic^2 \left( \frac{1}{2}ib \int \log(1 - e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2}ib \int \log(1 + e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) \right)}{d} \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2dx} \\
& \quad \downarrow \text{2715} \\
& \frac{2ic^2 \left( \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1 - e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1 + e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} \right)}{d} \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2dx} \\
& \quad \downarrow \text{2838} \\
& \frac{2ic^2 \left( i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4}ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4}ib \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right)}{d} \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2dx}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)),x]`

output `(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*d*x) - (a + b*ArcCosh[c*x])/(2*d*x^2) - ((2*I)*c^2*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])] + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])]))/d`

## 3.35.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 106 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((c_.)*(d_.)*(x_)))]^(n_.), x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/(f*fz*I)], x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

### 3.35.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.06

method	result
derivativedivides	$c^2 \left( -\frac{a \left( \frac{1}{2c^2x^2} - \ln(cx) + \frac{\ln(cx+1)}{2} + \frac{\ln(cx-1)}{2} \right)}{d} - \frac{b \left( \frac{-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx)}{2c^2x^2} - \operatorname{arccosh}(cx) \ln \left( 1+(cx+\sqrt{cx^2-1}) \right) \right)}{d} \right)$
default	$c^2 \left( -\frac{a \left( \frac{1}{2c^2x^2} - \ln(cx) + \frac{\ln(cx+1)}{2} + \frac{\ln(cx-1)}{2} \right)}{d} - \frac{b \left( \frac{-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx)}{2c^2x^2} - \operatorname{arccosh}(cx) \ln \left( 1+(cx+\sqrt{cx^2-1}) \right) \right)}{d} \right)$
parts	$-\frac{a \left( \frac{c^2 \ln(cx+1)}{2} + \frac{1}{2x^2} - c^2 \ln(x) + \frac{c^2 \ln(cx-1)}{2} \right)}{d} - \frac{b c^2 \left( \frac{-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx)}{2c^2x^2} - \operatorname{arccosh}(cx) \ln \left( 1+(cx+\sqrt{cx^2-1}) \right) \right)}{d}$

input `int((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d),x,method=_RETURNVERBOSE)`

output `c^2*(-a/d*(1/2/c^2/x^2-ln(c*x)+1/2*ln(c*x+1)+1/2*ln(c*x-1))-b/d*(1/2*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2+arccosh(c*x))/c^2/x^2-arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)-1/2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)+arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))))`

$$3.35. \int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)} dx$$

**3.35.5 Fricas [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d),x, algorithm="fricas")`

output `integral(-(b*arccosh(c*x) + a)/(c^2*d*x^5 - d*x^3), x)`

**3.35.6 Sympy [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)} dx = -\frac{\int \frac{a}{c^2 x^5 - x^3} dx + \int \frac{b \operatorname{acosh}(cx)}{c^2 x^5 - x^3} dx}{d}$$

input `integrate((a+b*acosh(c*x))/x**3/(-c**2*d*x**2+d),x)`

output `-(Integral(a/(c**2*x**5 - x**3), x) + Integral(b*acosh(c*x)/(c**2*x**5 - x**3), x))/d`

**3.35.7 Maxima [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `-1/2*(c^2*log(c*x + 1)/d + c^2*log(c*x - 1)/d - 2*c^2*log(x)/d + 1/(d*x^2))*a - b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(c^2*d*x^5 - d*x^3), x)`



**3.35.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d),x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)*x^3), x)`

**3.35.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (d - c^2 dx^2)} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)),x)`

output `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)), x)`

### 3.36 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^4(d-c^2dx^2)} dx$

3.36.1	Optimal result	465
3.36.2	Mathematica [A] (verified)	466
3.36.3	Rubi [C] (verified)	466
3.36.4	Maple [A] (verified)	470
3.36.5	Fricas [F]	471
3.36.6	Sympy [F]	471
3.36.7	Maxima [F]	472
3.36.8	Giac [F]	472
3.36.9	Mupad [F(-1)]	472

#### 3.36.1 Optimal result

Integrand size = 25, antiderivative size = 157

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^4(d - c^2dx^2)} dx = \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{6dx^2} - \frac{a + b\operatorname{arccosh}(cx)}{3dx^3} - \frac{c^2(a + b\operatorname{arccosh}(cx))}{dx} + \frac{7bc^3 \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{6d} + \frac{2c^3(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{d} + \frac{bc^3 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{d} - \frac{bc^3 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{d}$$

output  $\frac{1}{3}*(-a-b*\operatorname{arccosh}(c*x))/d/x^3-c^2*(a+b*\operatorname{arccosh}(c*x))/d/x+7/6*b*c^3*\arctan((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d+2*c^3*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d+b*c^3*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d-b*c^3*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d+1/6*b*c*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/x^2$

### 3.36.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.42

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)} dx$$

$$= \frac{-\frac{2a}{x^3} - \frac{6ac^2}{x} + \frac{bc\sqrt{-1+cx}\sqrt{1+cx}}{x^2} - \frac{2\operatorname{barccosh}(cx)}{x^3} - \frac{6bc^2\operatorname{arccosh}(cx)}{x} + \frac{7bc^3\sqrt{-1+c^2x^2}\arctan\left(\frac{\sqrt{-1+c^2x^2}}{\sqrt{-1+cx}\sqrt{1+cx}}\right)}{\sqrt{-1+cx}\sqrt{1+cx}} - 6ac^3 \log(1$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)),x]`

output  $((-2*a)/x^3 - (6*a*c^2)/x + (b*c*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x])/x^2 - (2*b*\operatorname{ArcCosh}[c*x])/x^3 - (6*b*c^2*\operatorname{ArcCosh}[c*x])/x + (7*b*c^3*\operatorname{Sqrt}[-1 + c^2*x^2]*\operatorname{ArcTan}[\operatorname{Sqrt}[-1 + c^2*x^2]])/(\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]) - 6*a*c^3*\operatorname{Log}[1 - E^{\operatorname{ArcCosh}[c*x]}] - 6*b*c^3*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - E^{\operatorname{ArcCosh}[c*x]}] + 6*a*c^3*\operatorname{Log}[1 + E^{\operatorname{ArcCosh}[c*x]}] + 6*b*c^3*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + E^{\operatorname{ArcCosh}[c*x]}] + 6*b*c^3*\operatorname{PolyLog}[2, -E^{\operatorname{ArcCosh}[c*x]}] - 6*b*c^3*\operatorname{PolyLog}[2, E^{\operatorname{ArcCosh}[c*x]}])/ (6*d)$

### 3.36.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.87 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.13, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$ , Rules used = {6347, 27, 114, 27, 103, 218, 6347, 103, 218, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)} dx$$

$$\downarrow 6347$$

$$c^2 \int \frac{a + \operatorname{barccosh}(cx)}{dx^2 (1 - c^2 x^2)} dx + \frac{bc \int \frac{1}{x^3 \sqrt{cx-1} \sqrt{cx+1}} dx}{3d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3}$$

$$\downarrow 27$$

$$\frac{c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (1 - c^2 x^2)} dx}{d} + \frac{bc \int \frac{1}{x^3 \sqrt{cx-1} \sqrt{cx+1}} dx}{3d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3}$$

---

3.36.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)} dx$

$$\begin{aligned}
& \downarrow 114 \\
& \frac{c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)} dx}{d} + \frac{bc \left( \frac{1}{2} \int \frac{c^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} \\
& \downarrow 27 \\
& \frac{c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)} dx}{d} + \frac{bc \left( \frac{1}{2} c^2 \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} \\
& \downarrow 103 \\
& \frac{c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)} dx}{d} + \frac{bc \left( \frac{1}{2} c^3 \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} \\
& \downarrow 218 \\
& \frac{c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)} dx}{d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2} c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d} \\
& \downarrow 6347 \\
& \frac{c^2 \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + bc \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2} c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d} \\
& \downarrow 103 \\
& \frac{c^2 \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + bc^2 \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2} c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d} \\
& \downarrow 218 \\
& \frac{c^2 \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx - \frac{a+\operatorname{barccosh}(cx)}{x} + bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) \right)}{d} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2} c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d} \\
& \downarrow 6318
\end{aligned}$$

$$\frac{c^2 \left( -c \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx) - \frac{a + \operatorname{barccosh}(cx)}{x} + bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) \right)}{d} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2}c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d}$$

↓ 3042

$$\frac{c^2 \left( -c \int i(a + \operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{a + \operatorname{barccosh}(cx)}{x} + bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) \right)}{d} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2}c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d}$$

↓ 26

$$\frac{c^2 \left( -ic \int (a + \operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{a + \operatorname{barccosh}(cx)}{x} + bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) \right)}{d} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2}c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d}$$

↓ 4670

$$\frac{c^2 \left( -ic \int (ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx))) \right)}{d} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2}c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d}$$

↓ 2715

$$\frac{c^2 \left( -ic \int (ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2ia \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx))) \right)}{d} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2}c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d}$$

↓ 2838

$$\frac{c^2 \left( -ic \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right) \right)}{d} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3dx^3} + \frac{bc \left( \frac{1}{2}c^2 \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)}{3d}$$

input `Int[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)),x]`

output `-1/3*(a + b*ArcCosh[c*x])/(d*x^3) + (b*c*((Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*x^2) + (c^2*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/2))/(3*d) + (c^2*(-((a + b*ArcCosh[c*x])/x) + b*c*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]] - I*c*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]])))/d`

### 3.36.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 103 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

rule 114 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`

rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^(e_)*((c_) + (d_)*(x_)))]^(n_), x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

### 3.36.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.21

method	result
derivativedivides	$c^3 \left( -\frac{a \left( \frac{1}{3c^3x^3} + \frac{1}{cx} - \frac{\ln(cx+1)}{2} + \frac{\ln(cx-1)}{2} \right)}{d} - \frac{b \left( \frac{6c^2x^2 \operatorname{arccosh}(cx) - \sqrt{cx-1}\sqrt{cx+1}cx + 2 \operatorname{arccosh}(cx) - 7 \arctan\left(\frac{cx+\sqrt{cx-1}}{3}\right)}{6c^3x^3} \right)}{d} \right)$
default	$c^3 \left( -\frac{a \left( \frac{1}{3c^3x^3} + \frac{1}{cx} - \frac{\ln(cx+1)}{2} + \frac{\ln(cx-1)}{2} \right)}{d} - \frac{b \left( \frac{6c^2x^2 \operatorname{arccosh}(cx) - \sqrt{cx-1}\sqrt{cx+1}cx + 2 \operatorname{arccosh}(cx) - 7 \arctan\left(\frac{cx+\sqrt{cx-1}}{3}\right)}{6c^3x^3} \right)}{d} \right)$
parts	$-\frac{a \left( -\frac{c^3 \ln(cx+1)}{2} + \frac{1}{3x^3} + \frac{c^2}{x} + \frac{c^3 \ln(cx-1)}{2} \right)}{d} - \frac{b c^3 \left( \frac{6c^2x^2 \operatorname{arccosh}(cx) - \sqrt{cx-1}\sqrt{cx+1}cx + 2 \operatorname{arccosh}(cx) - 7 \arctan\left(\frac{cx+\sqrt{cx-1}}{3}\right)}{6c^3x^3} \right)}{d}$

```
input int((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d),x,method=_RETURNVERBOSE)
```

```
output c^3*(-a/d*(1/3/c^3/x^3+1/c/x-1/2*ln(c*x+1)+1/2*ln(c*x-1))-b/d*(1/6*(6*c^2*x^2*arccosh(c*x)-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+2*arccosh(c*x))/c^3/x^3-7/3*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-dilog(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-dilog(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))
```

### 3.36.5 Fricas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)x^4} dx$$

```
input integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d),x, algorithm="fricas")
```

```
output integral(-(b*arccosh(c*x) + a)/(c^2*d*x^6 - d*x^4), x)
```

### 3.36.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)} dx = -\frac{\int \frac{a}{c^2 x^6 - x^4} dx + \int \frac{b \operatorname{arccosh}(cx)}{c^2 x^6 - x^4} dx}{d}$$

```
input integrate((a+b*acosh(c*x))/x**4/(-c**2*d*x**2+d),x)
```

```
output -(Integral(a/(c**2*x**6 - x**4), x) + Integral(b*acosh(c*x)/(c**2*x**6 - x**4), x))/d
```

---

3.36.  $\int \frac{a+b \operatorname{arccosh}(cx)}{x^4(d-c^2 dx^2)} dx$



**3.36.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `1/6*(3*c^3*log(c*x + 1)/d - 3*c^3*log(c*x - 1)/d - 2*(3*c^2*x^2 + 1)/(d*x^3))*a + 1/24*(216*c^5*integrate(1/12*x^3*log(c*x - 1)/(c^2*d*x^4 - d*x^2), x) - 12*c^4*(log(c*x + 1)/(c*d) - log(c*x - 1)/(c*d)) - 72*c^4*integrate(1/12*x^2*log(c*x - 1)/(c^2*d*x^4 - d*x^2), x) - 4*c^2*(c*log(c*x + 1)/d - c*log(c*x - 1)/d - 2/(d*x)) - (3*c^3*x^3*log(c*x + 1)^2 + 6*c^3*x^3*log(c*x + 1)*log(c*x - 1) - 4*(3*c^3*x^3*log(c*x + 1) - 3*c^3*x^3*log(c*x - 1) - 6*c^2*x^2 - 2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)))/(d*x^3) + 24*integrate(1/6*(3*c^4*x^3*log(c*x + 1) - 3*c^4*x^3*log(c*x - 1) - 6*c^3*x^2 - 2*c)/(c^3*d*x^6 - c*d*x^4 + (c^2*d*x^5 - d*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1)), x))*b`

**3.36.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d - c^2 dx^2)} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d),x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)*x^4), x)`

**3.36.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d - c^2 dx^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4(d - c^2 dx^2)} dx$$

input `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)),x)`

output `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)), x)`

---

3.36.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d - c^2 dx^2)} dx$

### 3.37 $\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$

3.37.1	Optimal result	473
3.37.2	Mathematica [A] (warning: unable to verify)	474
3.37.3	Rubi [C] (verified)	474
3.37.4	Maple [A] (verified)	479
3.37.5	Fricas [F]	479
3.37.6	Sympy [F]	480
3.37.7	Maxima [F]	480
3.37.8	Giac [F(-2)]	481
3.37.9	Mupad [F(-1)]	481

#### 3.37.1 Optimal result

Integrand size = 25, antiderivative size = 177

$$\int \frac{x^4(a + b\operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = -\frac{bx^2}{2c^3d^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{b\sqrt{-1+cx}\sqrt{1+cx}}{2c^5d^2} + \frac{3x(a + b\operatorname{arccosh}(cx))}{2c^4d^2} + \frac{x^3(a + b\operatorname{arccosh}(cx))}{2c^2d^2(1 - c^2x^2)} - \frac{3(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^5d^2} - \frac{3b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{2c^5d^2} + \frac{3b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{2c^5d^2}$$

output  $\frac{3}{2}x(a+b\operatorname{arccosh}(cx))/c^4/d^2+1/2x^3(a+b\operatorname{arccosh}(cx))/c^2/d^2/(-c^2x^2+1)-3(a+b\operatorname{arccosh}(cx))*\operatorname{arctanh}(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)})/c^5/d^2-3/2b*\operatorname{polylog}(2,-cx-(cx-1)^{(1/2)}*(cx+1)^{(1/2)})/c^5/d^2+3/2b*\operatorname{polylog}(2,cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)})/c^5/d^2-1/2b*x^2/c^3/d^2/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-1/2b*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/c^5/d^2$

### 3.37.2 Mathematica [A] (warning: unable to verify)

Time = 0.77 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.38

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^2} dx$$

$$= \frac{4acx - 3b\sqrt{\frac{-1+cx}{1+cx}} - 4bcx\sqrt{\frac{-1+cx}{1+cx}} + b\sqrt{\frac{-1+cx}{1+cx}} + \frac{bcx\sqrt{\frac{-1+cx}{1+cx}}}{1-cx} - \frac{2acx}{-1+c^2x^2} + 4bcx \operatorname{arccosh}(cx) + \frac{b \operatorname{arccosh}(cx)}{1-cx} - \frac{b \operatorname{arccosh}(cx)}{1+cx}}{d - c^2 dx^2}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `(4*a*c*x - 3*b*Sqrt[(-1 + c*x)/(1 + c*x)] - 4*b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)] + (b*Sqrt[(-1 + c*x)/(1 + c*x)]/(1 - c*x) + (b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) - (2*a*c*x)/(-1 + c^2*x^2) + 4*b*c*x*ArcCosh[c*x] + (b*ArcCosh[c*x])/(1 - c*x) - (b*ArcCosh[c*x])/(1 + c*x) + 6*b*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] - 6*b*ArcCosh[c*x]*Log[1 + E^ArcCosh[c*x]] + 3*a*Log[1 - c*x] - 3*a*Log[1 + c*x] - 6*b*PolyLog[2, -E^ArcCosh[c*x]] + 6*b*PolyLog[2, E^ArcCosh[c*x]])/(4*c^5*d^2)`

### 3.37.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.87 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.520$ , Rules used = {6349, 27, 109, 27, 83, 6353, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^2} dx$$

$$\downarrow \text{6349}$$

$$-\frac{3 \int \frac{x^2(a+b \operatorname{arccosh}(cx))}{d(1-c^2x^2)} dx}{2c^2d} + \frac{b \int \frac{x^3}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2cd^2} + \frac{x^3(a + b \operatorname{arccosh}(cx))}{2c^2d^2(1 - c^2x^2)}$$

$$\downarrow \text{27}$$

$$-\frac{3 \int \frac{x^2(a+b \operatorname{arccosh}(cx))}{1-c^2x^2} dx}{2c^2d^2} + \frac{b \int \frac{x^3}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2cd^2} + \frac{x^3(a + b \operatorname{arccosh}(cx))}{2c^2d^2(1 - c^2x^2)}$$

---

3.37.  $\int \frac{x^4(a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^2} dx$

$$\begin{aligned}
 & \downarrow 109 \\
 & -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{2c^2d^2} + \frac{b \left( -\frac{\int -\frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c^2} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2cd^2} + \frac{x^3(a + \operatorname{barccosh}(cx))}{2c^2d^2(1-c^2x^2)} \\
 & \downarrow 27 \\
 & -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{2c^2d^2} + \frac{b \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c^2} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2cd^2} + \frac{x^3(a + \operatorname{barccosh}(cx))}{2c^2d^2(1-c^2x^2)} \\
 & \downarrow 83 \\
 & -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{2c^2d^2} + \frac{x^3(a + \operatorname{barccosh}(cx))}{2c^2d^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2cd^2} \\
 & \downarrow 6353 \\
 & -\frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} + \frac{b \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} \right)}{2c^2d^2} + \frac{x^3(a + \operatorname{barccosh}(cx))}{2c^2d^2(1-c^2x^2)} + \\
 & \quad \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2cd^2} \\
 & \downarrow 83 \\
 & -\frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{2c^2d^2} + \frac{x^3(a + \operatorname{barccosh}(cx))}{2c^2d^2(1-c^2x^2)} + \\
 & \quad \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2cd^2} \\
 & \downarrow 6318 \\
 & -\frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{2c^2d^2} + \frac{x^3(a + \operatorname{barccosh}(cx))}{2c^2d^2(1-c^2x^2)} + \\
 & \quad \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2cd^2} \\
 & \downarrow 3042
 \end{aligned}$$

---

3.37.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$

$$\begin{aligned}
& \frac{3 \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{c^3} - \frac{x(a+\operatorname{barccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{2c^2d^2(1-c^2x^2)} + \frac{2c^2d^2}{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)} \\
& \quad \downarrow 26 \\
& \frac{3 \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{c^3} - \frac{x(a+\operatorname{barccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{2c^2d^2(1-c^2x^2)} + \frac{2c^2d^2}{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)} \\
& \quad \downarrow 4670 \\
& \frac{3 \left( -\frac{i \left( \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) \right)}{c^3} \right)}{2c^2d^2(1-c^2x^2)} + \frac{2c^2d^2}{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)} \\
& \quad \downarrow 2715 \\
& \frac{3 \left( -\frac{i \left( \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) \right)}{c^3} \right)}{2c^2d^2(1-c^2x^2)} + \frac{2c^2d^2}{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)} \\
& \quad \downarrow 2838 \\
& \frac{3 \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + \int \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - \int \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{c^3} \right) - \frac{x(a+\operatorname{barccosh}(cx))}{c^2}}{2c^2d^2(1-c^2x^2)} + \frac{2c^2d^2}{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

$$3.37. \quad \int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^2} dx$$

```
output (b*(-(x^2/(c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (2*Sqrt[-1 + c*x]*Sqrt[1 +
c*x])/c^4))/(2*c*d^2) + (x^3*(a + b*ArcCosh[c*x]))/(2*c^2*d^2*(1 - c^2*x^
2)) - (3*((b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^3 - (x*(a + b*ArcCosh[c*x]))/
c^2 - (I*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog
[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c^3))/(2*c^2*d^2)
```

### 3.37.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

```
rule 83 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f
*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]
```

```
rule 109 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f
*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1)
+ c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p)
+ b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c,
d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] ||
IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

```
rule 2715 Int[Log[(a_ + (b_.)*((F_)^(e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6349 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]`

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.37.4 Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.44

method	result
derivativedivides	$\frac{a\left(cx - \frac{1}{4(cx+1)} - \frac{3\ln(cx+1)}{4} - \frac{1}{4(cx-1)} + \frac{3\ln(cx-1)}{4}\right)}{d^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{d^2} + \frac{b\operatorname{arccosh}(cx)cx}{d^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{2d^2(c^2x^2-1)} - \frac{b\operatorname{arccosh}(cx)cx}{2d^2(c^2x^2-1)} + \frac{3b}{2d^2(c^2x^2-1)}$
default	$\frac{a\left(cx - \frac{1}{4(cx+1)} - \frac{3\ln(cx+1)}{4} - \frac{1}{4(cx-1)} + \frac{3\ln(cx-1)}{4}\right)}{d^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{d^2} + \frac{b\operatorname{arccosh}(cx)cx}{d^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{2d^2(c^2x^2-1)} - \frac{b\operatorname{arccosh}(cx)cx}{2d^2(c^2x^2-1)} + \frac{3b}{2d^2(c^2x^2-1)}$
parts	$\frac{a\left(\frac{x}{c^4} - \frac{1}{4c^5(cx+1)} - \frac{3\ln(cx+1)}{4c^5} - \frac{1}{4c^5(cx-1)} + \frac{3\ln(cx-1)}{4c^5}\right)}{d^2} + \frac{b\operatorname{arccosh}(cx)x}{d^2c^4} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^5d^2} - \frac{b\operatorname{arccosh}(cx)x}{2d^2c^4(c^2x^2-1)}$

input `int(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)`

output `1/c^5*(a/d^2*(c*x-1/4/(c*x+1)-3/4*ln(c*x+1)-1/4/(c*x-1)+3/4*ln(c*x-1))-b/d^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)+b/d^2*arccosh(c*x)*c*x-1/2*b/d^2/(c^2*x^2-1)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-1/2*b/d^2/(c^2*x^2-1)*arccosh(c*x)*c*x+3/2*b/d^2*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+3/2*b/d^2*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-3/2*b/d^2*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-3/2*b/d^2*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)))`

### 3.37.5 Fracas [F]

$$\int \frac{x^4(a + b\operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = \int \frac{(b\operatorname{arccosh}(cx) + a)x^4}{(c^2dx^2 - d)^2} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*x^4*arccosh(c*x) + a*x^4)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`



## 3.37.6 Sympy [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{ax^4}{c^4 x^4 - 2c^2 x^2 + 1} dx + \int \frac{bx^4 \operatorname{acosh}(cx)}{c^4 x^4 - 2c^2 x^2 + 1} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a*x**4/(c**4*x**4 - 2*c**2*x**2 + 1), x) + Integral(b*x**4*acosh(c*x)/(c**4*x**4 - 2*c**2*x**2 + 1), x))/d**2`

## 3.37.7 Maxima [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(c^2 dx^2 - d)^2} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `1/64*(16*c^4*(2*x/(c^10*d^2*x^2 - c^8*d^2) - 4*x/(c^8*d^2) + 3*log(c*x + 1)/(c^9*d^2) - 3*log(c*x - 1)/(c^9*d^2)) - 576*c^3*integrate(1/8*x^3*log(c*x - 1)/(c^8*d^2*x^4 - 2*c^6*d^2*x^2 + c^4*d^2), x) - 24*c^2*(2*x/(c^8*d^2*x^2 - c^6*d^2) + log(c*x + 1)/(c^7*d^2) - log(c*x - 1)/(c^7*d^2)) + 192*c^2*integrate(1/8*x^2*log(c*x - 1)/(c^8*d^2*x^4 - 2*c^6*d^2*x^2 + c^4*d^2), x) - 9*(c*(2/(c^8*d^2*x - c^7*d^2) - log(c*x + 1)/(c^7*d^2) + log(c*x - 1)/(c^7*d^2)) + 4*log(c*x - 1)/(c^8*d^2*x^2 - c^6*d^2))*c + 4*(3*(c^2*x^2 - 1)*log(c*x + 1)^2 + 6*(c^2*x^2 - 1)*log(c*x + 1)*log(c*x - 1) + 4*(4*c^3*x^3 - 6*c*x - 3*(c^2*x^2 - 1)*log(c*x + 1) + 3*(c^2*x^2 - 1)*log(c*x - 1))*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(c^7*d^2*x^2 - c^5*d^2) - 64*integrate(-1/4*(4*c^3*x^3 - 6*c*x - 3*(c^2*x^2 - 1)*log(c*x + 1) + 3*(c^2*x^2 - 1)*log(c*x - 1))/(c^9*d^2*x^5 - 2*c^7*d^2*x^3 + c^5*d^2*x + (c^8*d^2*x^4 - 2*c^6*d^2*x^2 + c^4*d^2))*sqrt(c*x + 1)*sqrt(c*x - 1), x) - 192*integrate(1/8*log(c*x - 1)/(c^8*d^2*x^4 - 2*c^6*d^2*x^2 + c^4*d^2), x))*b - 1/4*a*(2*x/(c^6*d^2*x^2 - c^4*d^2) - 4*x/(c^4*d^2) + 3*log(c*x + 1)/(c^5*d^2) - 3*log(c*x - 1)/(c^5*d^2))`

**3.37.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.37.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^2} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2,x)`

output `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2, x)`

### 3.38 $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$

3.38.1	Optimal result	482
3.38.2	Mathematica [A] (warning: unable to verify)	483
3.38.3	Rubi [C] (verified)	483
3.38.4	Maple [A] (verified)	488
3.38.5	Fricas [F]	489
3.38.6	Sympy [F]	489
3.38.7	Maxima [F]	489
3.38.8	Giac [F(-2)]	490
3.38.9	Mupad [F(-1)]	490

#### 3.38.1 Optimal result

Integrand size = 25, antiderivative size = 179

$$\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx = -\frac{b}{2c^4d^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{b\sqrt{-1+cx}}{2c^4d^2\sqrt{1+cx}} + \frac{b\operatorname{arccosh}(cx)}{2c^4d^2}$$

$$+ \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2d^2(1-c^2x^2)} - \frac{(a+b\operatorname{arccosh}(cx))^2}{2bc^4d^2}$$

$$+ \frac{(a+b\operatorname{arccosh}(cx))\log(1-e^{2\operatorname{arccosh}(cx)})}{c^4d^2}$$

$$+ \frac{b\operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2c^4d^2}$$

output  $1/2*b*\operatorname{arccosh}(c*x)/c^4/d^2+1/2*x^2*(a+b*\operatorname{arccosh}(c*x))/c^2/d^2/(-c^2*x^2+1)$   
 $-1/2*(a+b*\operatorname{arccosh}(c*x))^2/b/c^4/d^2+(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/c^4/d^2+1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/c^4/d^2-1/2*b/c^4/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/2*b*(c*x-1)^(1/2)/c^4/d^2/(c*x+1)^(1/2)$

### 3.38.2 Mathematica [A] (warning: unable to verify)

Time = 0.52 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.17

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx$$

$$= -b\sqrt{\frac{-1+cx}{1+cx}} + \frac{b\sqrt{\frac{-1+cx}{1+cx}}}{1-cx} + \frac{bcx\sqrt{\frac{-1+cx}{1+cx}}}{1-cx} - \frac{2a}{-1+c^2x^2} + \frac{\operatorname{barccosh}(cx)}{1-cx} + \frac{\operatorname{barccosh}(cx)}{1+cx} - 2\operatorname{barccosh}(cx)^2 + 4\operatorname{barccosh}(cx)$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `(- (b*sqrt[(-1 + c*x)/(1 + c*x)]) + (b*sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) + (b*c*x*sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) - (2*a)/(-1 + c^2*x^2) + (b*ArcCosh[c*x])/(1 - c*x) + (b*ArcCosh[c*x])/(1 + c*x) - 2*b*ArcCosh[c*x]^2 + 4*b*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] + 4*b*ArcCosh[c*x]*Log[1 + E^ArcCosh[c*x]] + 2*a*Log[1 - c^2*x^2] + 4*b*PolyLog[2, -E^ArcCosh[c*x]] + 4*b*PolyLog[2, E^ArcCosh[c*x]])/(4*c^4*d^2)`

### 3.38.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.77 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.01, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.560$ , Rules used = {6349, 27, 100, 27, 87, 43, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx$$

$$\downarrow \text{6349}$$

$$-\frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{d(1 - c^2 x^2)} dx}{c^2 d} + \frac{b \int \frac{x^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2cd^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)}$$

$$\downarrow \text{27}$$

$$-\frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d^2} + \frac{b \int \frac{x^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2cd^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)}$$

---

3.38.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx$

$$\begin{aligned}
 & \int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx + \frac{b \left( \frac{\int \frac{c^2 x}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c^3} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} \\
 & \quad \downarrow 100 \\
 & - \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d^2} + \frac{b \left( \frac{\int \frac{x}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} \\
 & \quad \downarrow 27 \\
 & - \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d^2} + \frac{b \left( \frac{\int \frac{1}{\sqrt{cx-1} \sqrt{cx+1}} dx}{c} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} \\
 & \quad \downarrow 87 \\
 & - \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2 d^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \quad \downarrow 43 \\
 & \frac{\int \frac{cx(a + \operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{c^4 d^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \quad \downarrow 6328 \\
 & \frac{\int -i(a + \operatorname{barccosh}(cx)) \tan \left( i \operatorname{arccosh}(cx) + \frac{\pi}{2} \right) d \operatorname{arccosh}(cx)}{c^4 d^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \quad \downarrow 3042 \\
 & \frac{\int -i(a + \operatorname{barccosh}(cx)) \tan \left( i \operatorname{arccosh}(cx) + \frac{\pi}{2} \right) d \operatorname{arccosh}(cx)}{c^4 d^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \quad \downarrow 26
 \end{aligned}$$

---

3.38.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx$

$$\begin{aligned}
 & - \frac{i \int (a + \operatorname{barccosh}(cx)) \tan \left( i \operatorname{arccosh}(cx) + \frac{\pi}{2} \right) d \operatorname{arccosh}(cx)}{c^4 d^2} + \frac{x^2 (a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \\
 & \quad \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \qquad \qquad \qquad \downarrow \text{4199} \\
 & - \frac{i \left( 2i \int - \frac{e^{2 \operatorname{arccosh}(cx)} (a + \operatorname{barccosh}(cx))}{1 - e^{2 \operatorname{arccosh}(cx)}} d \operatorname{arccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{c^4 d^2} + \frac{x^2 (a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \\
 & \quad \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & - \frac{i \left( -2i \int \frac{e^{2 \operatorname{arccosh}(cx)} (a + \operatorname{barccosh}(cx))}{1 - e^{2 \operatorname{arccosh}(cx)}} d \operatorname{arccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{c^4 d^2} + \frac{x^2 (a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \\
 & \quad \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \qquad \qquad \qquad \downarrow \text{2620} \\
 & - \frac{i \left( -2i \left( \frac{1}{2} b \int \log (1 - e^{2 \operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) - \frac{1}{2} \log (1 - e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) \right) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{c^4 d^2} + \\
 & \quad \frac{x^2 (a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \qquad \qquad \qquad \downarrow \text{2715} \\
 & - \frac{i \left( -2i \left( \frac{1}{4} b \int e^{-2 \operatorname{arccosh}(cx)} \log (1 - e^{2 \operatorname{arccosh}(cx)}) d e^{2 \operatorname{arccosh}(cx)} - \frac{1}{2} \log (1 - e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) \right) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{c^4 d^2} + \\
 & \quad \frac{x^2 (a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} \right)}{2cd^2} \\
 & \qquad \qquad \qquad \downarrow \text{2838}
 \end{aligned}$$

---

3.38.  $\int \frac{x^3 (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx$

$$i\left(-2i\left(-\frac{1}{2}\log(1 - e^{2\operatorname{arccosh}(cx)})\right)(a + \operatorname{arccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})\right) - \frac{i(a + \operatorname{arccosh}(cx))^2}{2b}\right) +$$

$$\frac{x^2(a + \operatorname{arccosh}(cx))}{2c^2d^2(1 - c^2x^2)} + \frac{b\left(\frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}}\right)}{2cd^2}$$

input `Int[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `(x^2*(a + b*ArcCosh[c*x]))/(2*c^2*d^2*(1 - c^2*x^2)) + (b*(-(1/(c^3*sqrt[-1 + c*x]*sqrt[1 + c*x])) + (-sqrt[-1 + c*x]/(c^2*sqrt[1 + c*x])) + ArcCosh[c*x]/c^2)/c)/(2*c*d^2) - (I*(((1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/(c^4*d^2)`

### 3.38.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 43 `Int[1/(sqrt[(a_) + (b_.)*(x_)]*sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

---

3.38.  $\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^2} dx$

- rule 100 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[(b*c - a*d)2(c + d*x)(n + 1)((e + f*x)(p + 1)/(d2(d*e - c*f)(n + 1))), x] - Simp[1/(d2(d*e - c*f)(n + 1)) Int[(c + d*x)(n + 1)(e + f*x)pSimp[a2d2f*(n + p + 2) + b2*c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b2*d*(d*e - c*f)(n + 1)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))`
- rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))(n_.)((c_.) + (d_.)*(x_))(m_.))/((a_.) + (b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))(n_.), x_Symbol] := Simp[((c + d*x)m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))n/a), x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)(m - 1)*Log[1 + b*((F^(g*(e + f*x)))n/a), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_))(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*xn/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4199 `Int[((c_.) + (d_.)*(x_))(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)m(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`
- rule 6328 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)(x_))/((d_) + (e_.)*(x_)2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c2*d + e, 0] && IGtQ[n, 0]`



```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1)))
Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - S
imp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)]
Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c
*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0]
&& GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.38.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.13

method	result
derivativedivides	$\frac{a \left( \frac{1}{4cx+4} + \frac{\ln(cx+1)}{2} - \frac{1}{4(cx-1)} + \frac{\ln(cx-1)}{2} \right) + b \left( -\frac{\operatorname{arccosh}(cx)^2}{2} - \frac{\sqrt{cx-1}\sqrt{cx+1}cx - c^2x^2 + \operatorname{arccosh}(cx)+1}{2(c^2x^2-1)} + \operatorname{arccosh}(cx) \ln(1-cx-1) \right)}{d^2} + \frac{\dots}{c^4}$
default	$\frac{a \left( \frac{1}{4cx+4} + \frac{\ln(cx+1)}{2} - \frac{1}{4(cx-1)} + \frac{\ln(cx-1)}{2} \right) + b \left( -\frac{\operatorname{arccosh}(cx)^2}{2} - \frac{\sqrt{cx-1}\sqrt{cx+1}cx - c^2x^2 + \operatorname{arccosh}(cx)+1}{2(c^2x^2-1)} + \operatorname{arccosh}(cx) \ln(1-cx-1) \right)}{d^2} + \frac{\dots}{c^4}$
parts	$\frac{a \left( \frac{1}{4c^4(cx+1)} + \frac{\ln(cx+1)}{2c^4} - \frac{1}{4c^4(cx-1)} + \frac{\ln(cx-1)}{2c^4} \right) + b \left( -\frac{\operatorname{arccosh}(cx)^2}{2} - \frac{\sqrt{cx-1}\sqrt{cx+1}cx - c^2x^2 + \operatorname{arccosh}(cx)+1}{2(c^2x^2-1)} + \operatorname{arccosh}(cx) \ln(1-cx-1) \right)}{d^2} + \frac{\dots}{c^4}$

```
input int(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)
```

```
output 1/c^4*(a/d^2*(1/4/(c*x+1)+1/2*ln(c*x+1)-1/4/(c*x-1)+1/2*ln(c*x-1))+b/d^2*(
-1/2*arccosh(c*x)^2-1/2*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-c^2*x^2+arccosh(c
*x)+1)/(c^2*x^2-1)+arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+poly
log(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)
*(c*x+1)^(1/2))+polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))))
```

$$3.38. \int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$$

**3.38.5 Fracas [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^3}{(c^2 dx^2 - d)^2} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*x^3*arccosh(c*x) + a*x^3)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.38.6 Sympy [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{ax^3}{c^4 x^4 - 2c^2 x^2 + 1} dx + \int \frac{bx^3 \operatorname{acosh}(cx)}{c^4 x^4 - 2c^2 x^2 + 1} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a*x**3/(c**4*x**4 - 2*c**2*x**2 + 1), x) + Integral(b*x**3*acosh(c*x)/(c**4*x**4 - 2*c**2*x**2 + 1), x))/d**2`

**3.38.7 Maxima [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^3}{(c^2 dx^2 - d)^2} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `-1/8*b*(((c^2*x^2 - 1)*log(c*x + 1)^2 + 2*(c^2*x^2 - 1)*log(c*x + 1)*log(c*x - 1) + (c^2*x^2 - 1)*log(c*x - 1)^2 - 4*(((c^2*x^2 - 1)*log(c*x + 1) + (c^2*x^2 - 1)*log(c*x - 1) - 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + 2)/(c^6*d^2*x^2 - c^4*d^2) - 8*integrate(1/2*(((c^2*x^2 - 1)*log(c*x + 1) + (c^2*x^2 - 1)*log(c*x - 1) - 1)/(c^8*d^2*x^5 - 2*c^6*d^2*x^3 + c^4*d^2*x + (c^7*d^2*x^4 - 2*c^5*d^2*x^2 + c^3*d^2))*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1))), x)) - 1/2*a*(1/(c^6*d^2*x^2 - c^4*d^2) - log(c^2*x^2 - 1)/(c^4*d^2))`

---

3.38.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx$

**3.38.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.38.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^2} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2,x)`

output `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2, x)`

**3.39** 
$$\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$$

3.39.1 Optimal result . . . . . 491  
 3.39.2 Mathematica [A] (warning: unable to verify) . . . . . 491  
 3.39.3 Rubi [C] (verified) . . . . . 492  
 3.39.4 Maple [A] (verified) . . . . . 495  
 3.39.5 Fricas [F] . . . . . 496  
 3.39.6 Sympy [F] . . . . . 496  
 3.39.7 Maxima [F] . . . . . 496  
 3.39.8 Giac [F] . . . . . 497  
 3.39.9 Mupad [F(-1)] . . . . . 497

**3.39.1 Optimal result**

Integrand size = 25, antiderivative size = 124

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = -\frac{b}{2c^3d^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{x(a + \operatorname{arccosh}(cx))}{2c^2d^2(1 - c^2x^2)} - \frac{(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^3d^2} - \frac{b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{2c^3d^2} + \frac{b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{2c^3d^2}$$

```
output 1/2*x*(a+b*arccosh(c*x))/c^2/d^2/(-c^2*x^2+1)-(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d^2-1/2*b*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d^2+1/2*b*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d^2-1/2*b/c^3/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.39.2 Mathematica [A] (warning: unable to verify)**

Time = 0.61 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.66

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = b\sqrt{\frac{-1+cx}{1+cx}} + \frac{b\sqrt{\frac{-1+cx}{1+cx}}}{1-cx} + \frac{bcx\sqrt{\frac{-1+cx}{1+cx}}}{1-cx} - \frac{2acx}{-1+c^2x^2} + \frac{\operatorname{arccosh}(cx)}{1-cx} - \frac{\operatorname{arccosh}(cx)}{1+cx} + 2\operatorname{arccosh}(cx) \log(1 - e^{\operatorname{arccosh}(cx)})$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `(b*Sqrt[(-1 + c*x)/(1 + c*x)] + (b*Sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) + (b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) - (2*a*c*x)/(-1 + c^2*x^2) + (b*ArcCosh[c*x])/(1 - c*x) - (b*ArcCosh[c*x])/(1 + c*x) + 2*b*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] - 2*b*ArcCosh[c*x]*Log[1 + E^ArcCosh[c*x]] + a*Log[1 - c*x] - a*Log[1 + c*x] - 2*b*PolyLog[2, -E^ArcCosh[c*x]] + 2*b*PolyLog[2, E^ArcCosh[c*x]])/(4*c^3*d^2)`

### 3.39.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.98, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$ , Rules used = {6349, 27, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx \\
 & \quad \downarrow \text{6349} \\
 & -\frac{\int \frac{a + \operatorname{barccosh}(cx)}{d(1 - c^2x^2)} dx}{2c^2d} + \frac{b \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2cd^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2c^2d^2(1 - c^2x^2)} \\
 & \quad \downarrow \text{27} \\
 & -\frac{\int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2x^2} dx}{2c^2d^2} + \frac{b \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2cd^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2c^2d^2(1 - c^2x^2)} \\
 & \quad \downarrow \text{83} \\
 & -\frac{\int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2x^2} dx}{2c^2d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2c^2d^2(1 - c^2x^2)} - \frac{b}{2c^3d^2\sqrt{cx-1}\sqrt{cx+1}} \\
 & \quad \downarrow \text{6318} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{2c^3d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2c^2d^2(1 - c^2x^2)} - \frac{b}{2c^3d^2\sqrt{cx-1}\sqrt{cx+1}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.39.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx$

$$\frac{\int i(a + \operatorname{barccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{2c^3 d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} - \frac{b}{2c^3 d^2 \sqrt{cx - 1} \sqrt{cx + 1}}$$

↓ 26

$$\frac{i \int (a + \operatorname{barccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{2c^3 d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} - \frac{b}{2c^3 d^2 \sqrt{cx - 1} \sqrt{cx + 1}}$$

↓ 4670

$$\frac{i(ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)))}{2c^3 d^2} - \frac{x(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} - \frac{b}{2c^3 d^2 \sqrt{cx - 1} \sqrt{cx + 1}}$$

↓ 2715

$$\frac{i(ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) d e^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) d e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)))}{2c^3 d^2} - \frac{x(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} - \frac{b}{2c^3 d^2 \sqrt{cx - 1} \sqrt{cx + 1}}$$

↓ 2838

$$\frac{i(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{2c^3 d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2c^2 d^2 (1 - c^2 x^2)} - \frac{b}{2c^3 d^2 \sqrt{cx - 1} \sqrt{cx + 1}}$$

input `Int[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `-1/2*b/(c^3*d^2*sqrt[-1 + c*x]*sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*c^2*d^2*(1 - c^2*x^2)) + ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/(c^3*d^2)`

## 3.39.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(d*f*(n + p + 2)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.39.4 Maple [A] (verified)

Time = 0.73 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.55

method	result
derivativedivides	$\frac{a \left( -\frac{1}{4(cx+1)} - \frac{\ln(cx+1)}{4} - \frac{1}{4(cx-1)} + \frac{\ln(cx-1)}{4} \right) + b \left( -\frac{cx \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1}}{2(c^2x^2-1)} + \frac{\operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1} \sqrt{cx+1})}{2} + \operatorname{polylog}(2, cx + (cx-1)^{1/2}) \right)}{d^2}$
default	$\frac{a \left( -\frac{1}{4(cx+1)} - \frac{\ln(cx+1)}{4} - \frac{1}{4(cx-1)} + \frac{\ln(cx-1)}{4} \right) + b \left( -\frac{cx \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1}}{2(c^2x^2-1)} + \frac{\operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1} \sqrt{cx+1})}{2} + \operatorname{polylog}(2, cx + (cx-1)^{1/2}) \right)}{c^3}$
parts	$\frac{a \left( -\frac{1}{4c^3(cx+1)} - \frac{\ln(cx+1)}{4c^3} - \frac{1}{4c^3(cx-1)} + \frac{\ln(cx-1)}{4c^3} \right) + b \left( -\frac{cx \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1}}{2(c^2x^2-1)} + \frac{\operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1} \sqrt{cx+1})}{2} + \operatorname{polylog}(2, cx + (cx-1)^{1/2}) \right)}{d^2}$

```
input int(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)
```

```
output 1/c^3*(a/d^2*(-1/4/(c*x+1)-1/4*ln(c*x+1)-1/4/(c*x-1)+1/4*ln(c*x-1))+b/d^2*(-1/2*(c*x*arccosh(c*x)+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(c^2*x^2-1)+1/2*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/2*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-1/2*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-1/2*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))))
```

---

3.39.  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$



**3.39.5 Fracas [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(c^2 dx^2 - d)^2} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*x^2*arccosh(c*x) + a*x^2)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.39.6 Sympy [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{ax^2}{c^4 x^4 - 2c^2 x^2 + 1} dx + \int \frac{bx^2 \operatorname{acosh}(cx)}{c^4 x^4 - 2c^2 x^2 + 1} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a*x**2/(c**4*x**4 - 2*c**2*x**2 + 1), x) + Integral(b*x**2*acosh(c*x)/(c**4*x**4 - 2*c**2*x**2 + 1), x))/d**2`

**3.39.7 Maxima [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(c^2 dx^2 - d)^2} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

```
output -1/64*(192*c^3*integrate(1/8*x^3*log(c*x - 1)/(c^6*d^2*x^4 - 2*c^4*d^2*x^2
+ c^2*d^2), x) + 8*c^2*(2*x/(c^6*d^2*x^2 - c^4*d^2) + log(c*x + 1)/(c^5*d
^2) - log(c*x - 1)/(c^5*d^2)) - 64*c^2*integrate(1/8*x^2*log(c*x - 1)/(c^6
*d^2*x^4 - 2*c^4*d^2*x^2 + c^2*d^2), x) + 3*(c*(2/(c^6*d^2*x - c^5*d^2) -
log(c*x + 1)/(c^5*d^2) + log(c*x - 1)/(c^5*d^2)) + 4*log(c*x - 1)/(c^6*d^2
*x^2 - c^4*d^2))*c - 4*((c^2*x^2 - 1)*log(c*x + 1)^2 + 2*(c^2*x^2 - 1)*log
(c*x + 1)*log(c*x - 1) - 4*(2*c*x + (c^2*x^2 - 1)*log(c*x + 1) - (c^2*x^2
- 1)*log(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)))/(c^5*d^2*x^2 -
c^3*d^2) + 64*integrate(1/4*(2*c*x + (c^2*x^2 - 1)*log(c*x + 1) - (c^2*x^2
- 1)*log(c*x - 1))/(c^7*d^2*x^5 - 2*c^5*d^2*x^3 + c^3*d^2*x + (c^6*d^2*x^
4 - 2*c^4*d^2*x^2 + c^2*d^2)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) + 64*integra
te(1/8*log(c*x - 1)/(c^6*d^2*x^4 - 2*c^4*d^2*x^2 + c^2*d^2), x))*b - 1/4*a
*(2*x/(c^4*d^2*x^2 - c^2*d^2) + log(c*x + 1)/(c^3*d^2) - log(c*x - 1)/(c^3
*d^2))
```

### 3.39.8 Giac [F]

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(c^2 dx^2 - d)^2} dx$$

```
input integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="giac")
```

```
output integrate((b*arccosh(c*x) + a)*x^2/(c^2*d*x^2 - d)^2, x)
```

### 3.39.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^2} dx$$

```
input int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2,x)
```

```
output int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2, x)
```

### 3.40 $\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$

3.40.1	Optimal result	498
3.40.2	Mathematica [A] (verified)	498
3.40.3	Rubi [A] (verified)	499
3.40.4	Maple [A] (verified)	500
3.40.5	Fricas [A] (verification not implemented)	500
3.40.6	Sympy [F]	501
3.40.7	Maxima [B] (verification not implemented)	501
3.40.8	Giac [F]	502
3.40.9	Mupad [F(-1)]	502

#### 3.40.1 Optimal result

Integrand size = 23, antiderivative size = 61

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = -\frac{bx}{2cd^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{a + \operatorname{arccosh}(cx)}{2c^2d^2(1 - c^2x^2)}$$

output `1/2*(a+b*arccosh(c*x))/c^2/d^2/(-c^2*x^2+1)-1/2*b*x/c/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)`

#### 3.40.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.87

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = \frac{a + bcx\sqrt{-1 + cx}\sqrt{1 + cx} + \operatorname{arccosh}(cx)}{2c^2d^2 - 2c^4d^2x^2}$$

input `Integrate[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `(a + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + b*ArcCosh[c*x])/(2*c^2*d^2 - 2*c^4*d^2*x^2)`

### 3.40.3 Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {6329, 41}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^2} dx$$

↓ 6329

$$\frac{b \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2cd^2} + \frac{a + \operatorname{arccosh}(cx)}{2c^2 d^2 (1 - c^2 x^2)}$$

↓ 41

$$\frac{a + \operatorname{arccosh}(cx)}{2c^2 d^2 (1 - c^2 x^2)} - \frac{bx}{2cd^2 \sqrt{cx-1} \sqrt{cx+1}}$$

input `Int[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `-1/2*(b*x)/(c*d^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*ArcCosh[c*x])/(2*c^2*d^2*(1 - c^2*x^2))`

#### 3.40.3.1 Defintions of rubi rules used

rule 41 `Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] :> Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`

rule 6329 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.40.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.05

method	result	size
derivativedivides	$\frac{-\frac{a}{2d^2(c^2x^2-1)} + \frac{b\left(-\frac{\operatorname{arccosh}(cx)}{2(c^2x^2-1)} - \frac{cx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2}}{c^2}$	64
default	$\frac{-\frac{a}{2d^2(c^2x^2-1)} + \frac{b\left(-\frac{\operatorname{arccosh}(cx)}{2(c^2x^2-1)} - \frac{cx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2}}{c^2}$	64
parts	$-\frac{a}{2d^2c^2(c^2x^2-1)} + \frac{b\left(-\frac{\operatorname{arccosh}(cx)}{2(c^2x^2-1)} - \frac{cx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2c^2}$	66

input `int(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)`

output `1/c^2*(-1/2*a/d^2/(c^2*x^2-1)+b/d^2*(-1/2/(c^2*x^2-1)*arccosh(c*x)-1/2/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c*x))`

### 3.40.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.07

$$\int \frac{x(a + b\operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = -\frac{ac^2x^2 + \sqrt{c^2x^2 - 1}bcx + b \log(cx + \sqrt{c^2x^2 - 1})}{2(c^4d^2x^2 - c^2d^2)}$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `-1/2*(a*c^2*x^2 + sqrt(c^2*x^2 - 1)*b*c*x + b*log(c*x + sqrt(c^2*x^2 - 1)))/(c^4*d^2*x^2 - c^2*d^2)`

### 3.40.6 Sympy [F]

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = \frac{\int \frac{ax}{c^4 x^4 - 2c^2 x^2 + 1} dx + \int \frac{bx \operatorname{acosh}(cx)}{c^4 x^4 - 2c^2 x^2 + 1} dx}{d^2}$$

input `integrate(x*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a*x/(c**4*x**4 - 2*c**2*x**2 + 1), x) + Integral(b*x*acosh(c*x)/(c**4*x**4 - 2*c**2*x**2 + 1), x))/d**2`

### 3.40.7 Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs.  $2(52) = 104$ .

Time = 0.27 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.20

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^2} dx = -\frac{1}{4} \left( \left( \frac{\sqrt{c^2 x^2 - 1} c^2 d^2}{c^7 d^4 x + c^6 d^4} + \frac{\sqrt{c^2 x^2 - 1} c^2 d^2}{c^7 d^4 x - c^6 d^4} \right) c^2 + \frac{2 \operatorname{arcosh}(cx)}{c^4 d^2 x^2 - c^2 d^2} \right) b - \frac{a}{2(c^4 d^2 x^2 - c^2 d^2)}$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `-1/4*((sqrt(c^2*x^2 - 1)*c^2*d^2/(c^7*d^4*x + c^6*d^4) + sqrt(c^2*x^2 - 1)*c^2*d^2/(c^7*d^4*x - c^6*d^4))*c^2 + 2*arccosh(c*x)/(c^4*d^2*x^2 - c^2*d^2))*b - 1/2*a/(c^4*d^2*x^2 - c^2*d^2)`

**3.40.8 Giac [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(c^2 dx^2 - d)^2} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x/(c^2*d*x^2 - d)^2, x)`

**3.40.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^2} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^2} dx$$

input `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2,x)`

output `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^2, x)`

### 3.41 $\int \frac{a+b\operatorname{arccosh}(cx)}{(d-c^2dx^2)^2} dx$

3.41.1	Optimal result	503
3.41.2	Mathematica [A] (warning: unable to verify)	503
3.41.3	Rubi [C] (verified)	504
3.41.4	Maple [A] (verified)	507
3.41.5	Fricas [F]	507
3.41.6	Sympy [F]	508
3.41.7	Maxima [F]	508
3.41.8	Giac [F]	509
3.41.9	Mupad [F(-1)]	509

#### 3.41.1 Optimal result

Integrand size = 22, antiderivative size = 120

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d - c^2dx^2)^2} dx = -\frac{b}{2cd^2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{x(a + \operatorname{arccosh}(cx))}{2d^2(1 - c^2x^2)} + \frac{(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{cd^2} + \frac{b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{2cd^2} - \frac{b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{2cd^2}$$

output  $1/2*x*(a+b*\operatorname{arccosh}(c*x))/d^2/(-c^2*x^2+1)+(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/c/d^2+1/2*b*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/c/d^2-1/2*b*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/c/d^2-1/2*b/c/d^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

#### 3.41.2 Mathematica [A] (warning: unable to verify)

Time = 1.05 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.58

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d - c^2dx^2)^2} dx = \frac{-2acx - 2b\sqrt{\frac{-1+cx}{1+cx}} - 2bcx\sqrt{\frac{-1+cx}{1+cx}} - 2b\operatorname{arccosh}(cx)\left(cx + (-1+c^2x^2)\log\left(1 - e^{\operatorname{arccosh}(cx)}\right) + (1-c^2x^2)\log\left(1 + e^{\operatorname{arccosh}(cx)}\right)\right) + (a - ac^2x^2)\log\left(\frac{cx + (-1+c^2x^2)\log\left(1 - e^{\operatorname{arccosh}(cx)}\right) + (1-c^2x^2)\log\left(1 + e^{\operatorname{arccosh}(cx)}\right)}{-1+c^2x^2}\right)}{4cd^2}$$



input `Integrate[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^2,x]`

output `((-2*a*c*x - 2*b*Sqrt[(-1 + c*x)/(1 + c*x)] - 2*b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)] - 2*b*ArcCosh[c*x]*(c*x + (-1 + c^2*x^2)*Log[1 - E^ArcCosh[c*x]] + (1 - c^2*x^2)*Log[1 + E^ArcCosh[c*x]]) + (a - a*c^2*x^2)*Log[1 - c*x] - a*Log[1 + c*x] + a*c^2*x^2*Log[1 + c*x])/(-1 + c^2*x^2) + 2*b*PolyLog[2, -E^ArcCosh[c*x]] - 2*b*PolyLog[2, E^ArcCosh[c*x]])/(4*c*d^2)`

### 3.41.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.98, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.409$ , Rules used = {6316, 27, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^2} dx \\
 & \quad \downarrow \text{6316} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{d(1 - c^2 x^2)} dx}{2d} + \frac{bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2 x^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2 x^2} dx}{2d^2} + \frac{bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2 x^2)} \\
 & \quad \downarrow \text{83} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{1 - c^2 x^2} dx}{2d^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2 x^2)} - \frac{b}{2cd^2 \sqrt{cx-1} \sqrt{cx+1}} \\
 & \quad \downarrow \text{6318} \\
 & -\frac{\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}} \operatorname{darccosh}(cx)} dx}{2cd^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2 x^2)} - \frac{b}{2cd^2 \sqrt{cx-1} \sqrt{cx+1}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.41.  $\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^2} dx$

$$\begin{aligned}
& -\frac{\int i(a + \operatorname{barccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{2cd^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2x^2)} - \\
& \quad \frac{b}{2cd^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow 26 \\
& -\frac{i \int (a + \operatorname{barccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{2cd^2} + \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2x^2)} - \\
& \quad \frac{b}{2cd^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow 4670 \\
& -\frac{i(ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2x^2)} - \frac{2cd^2}{2cd^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow 2715 \\
& -\frac{i(ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctan} \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2x^2)} - \frac{b}{2cd^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow 2838 \\
& -\frac{i(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{2cd^2} + \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{2d^2(1 - c^2x^2)} - \frac{b}{2cd^2\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^2,x]`

output `-1/2*b/(c*d^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*d^2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]])) / (c*d^2)`

## 3.41.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/(f*fz*I)], x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)/((d_.) + (e_.)*(x_)^2), x_Symbol]
:> Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

### 3.41.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.60

method	result
derivativedivides	$\frac{a \left( -\frac{1}{4(cx+1)} + \frac{\ln(cx+1)}{4} - \frac{1}{4(cx-1)} - \frac{\ln(cx-1)}{4} \right)}{d^2} + \frac{b \left( -\frac{cx \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1}}{2(c^2x^2-1)} - \frac{\operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1} \sqrt{cx+1})}{2} \right)}{c}$
default	$\frac{a \left( -\frac{1}{4(cx+1)} + \frac{\ln(cx+1)}{4} - \frac{1}{4(cx-1)} - \frac{\ln(cx-1)}{4} \right)}{d^2} + \frac{b \left( -\frac{cx \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1}}{2(c^2x^2-1)} - \frac{\operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1} \sqrt{cx+1})}{2} \right)}{c}$
parts	$\frac{a \left( -\frac{1}{4c(cx+1)} + \frac{\ln(cx+1)}{4c} - \frac{1}{4c(cx-1)} - \frac{\ln(cx-1)}{4c} \right)}{d^2} + \frac{b \left( -\frac{cx \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1}}{2(c^2x^2-1)} - \frac{\operatorname{arccosh}(cx) \ln(1-cx-\sqrt{cx-1} \sqrt{cx+1})}{2} \right)}{c}$

input `int((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)`

output `1/c*(a/d^2*(-1/4/(c*x+1)+1/4*ln(c*x+1)-1/4/(c*x-1)-1/4*ln(c*x-1))+b/d^2*(-1/2*(c*x*arccosh(c*x)+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(c^2*x^2-1)-1/2*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-1/2*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/2*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/2*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))))`

### 3.41.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^2} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="fracas")`

output `integral((b*arccosh(c*x) + a)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

---

3.41.  $\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^2} dx$

## 3.41.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^2} dx = \int \frac{a}{c^4 x^4 - 2c^2 x^2 + 1} dx + \int \frac{b \operatorname{acosh}(cx)}{c^4 x^4 - 2c^2 x^2 + 1} dx$$

input `integrate((a+b*acosh(c*x))/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a/(c**4*x**4 - 2*c**2*x**2 + 1), x) + Integral(b*acosh(c*x)/(c**4*x**4 - 2*c**2*x**2 + 1), x))/d**2`

## 3.41.7 Maxima [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `1/64*(192*c^3*integrate(1/8*x^3*log(c*x - 1)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x) - 8*c^2*(2*x/(c^4*d^2*x^2 - c^2*d^2) + log(c*x + 1)/(c^3*d^2) - log(c*x - 1)/(c^3*d^2)) - 64*c^2*integrate(1/8*x^2*log(c*x - 1)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x) + 3*(c*(2/(c^4*d^2*x - c^3*d^2) - log(c*x + 1)/(c^3*d^2) + log(c*x - 1)/(c^3*d^2)) + 4*log(c*x - 1)/(c^4*d^2*x^2 - c^2*d^2))*c - 4*((c^2*x^2 - 1)*log(c*x + 1)^2 + 2*(c^2*x^2 - 1)*log(c*x + 1)*log(c*x - 1) + 4*(2*c*x - (c^2*x^2 - 1)*log(c*x + 1) + (c^2*x^2 - 1)*log(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)))/(c^3*d^2*x^2 - c*d^2) + 64*integrate(-1/4*(2*c*x - (c^2*x^2 - 1)*log(c*x + 1) + (c^2*x^2 - 1)*log(c*x - 1))/(c^5*d^2*x^5 - 2*c^3*d^2*x^3 + c*d^2*x + (c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) + 64*integrate(1/8*log(c*x - 1)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x))*b - 1/4*a*(2*x/(c^2*d^2*x^2 - d^2) - log(c*x + 1)/(c*d^2) + log(c*x - 1)/(c*d^2))`

**3.41.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(c^2*d*x^2 - d)^2, x)`

**3.41.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(d - c^2 dx^2)^2} dx$$

input `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^2,x)`

output `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^2, x)`

### 3.42 $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)^2} dx$

3.42.1	Optimal result	510
3.42.2	Mathematica [B] (warning: unable to verify)	510
3.42.3	Rubi [C] (verified)	511
3.42.4	Maple [A] (verified)	514
3.42.5	Fricas [F]	515
3.42.6	Sympy [F]	515
3.42.7	Maxima [F]	515
3.42.8	Giac [F]	516
3.42.9	Mupad [F(-1)]	516

#### 3.42.1 Optimal result

Integrand size = 25, antiderivative size = 116

$$\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2dx^2)^2} dx = -\frac{bcx}{2d^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{a + \operatorname{arccosh}(cx)}{2d^2(1 - c^2x^2)} + \frac{2(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d^2} + \frac{b \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{2d^2} - \frac{b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2d^2}$$

output  $1/2*(a+b*\operatorname{arccosh}(c*x))/d^2/(-c^2*x^2+1)+2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/d^2+1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/d^2-1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/d^2-1/2*b*c*x/d^2/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}$

#### 3.42.2 Mathematica [B] (warning: unable to verify)

Leaf count is larger than twice the leaf count of optimal. 244 vs. 2(116) = 232.

Time = 0.61 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.10

$$\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2dx^2)^2} dx = -b\sqrt{\frac{-1+cx}{1+cx}} + \frac{b\sqrt{\frac{-1+cx}{1+cx}}}{1-cx} + \frac{bcx\sqrt{\frac{-1+cx}{1+cx}}}{1-cx} - \frac{2a}{-1+c^2x^2} + \frac{b\operatorname{arccosh}(cx)}{1-cx} + \frac{b\operatorname{arccosh}(cx)}{1+cx} + 4b\operatorname{arccosh}(cx)^2 + 4b\operatorname{arccosh}(cx)$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^2),x]`

output `(-(b*Sqrt[(-1 + c*x)/(1 + c*x)]) + (b*Sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) + (b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) - (2*a)/(-1 + c^2*x^2) + (b*ArcCosh[c*x])/(1 - c*x) + (b*ArcCosh[c*x])/(1 + c*x) + 4*b*ArcCosh[c*x]^2 + 4*b*ArcCosh[c*x]*Log[1 + E^(-2*ArcCosh[c*x])] - 4*b*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] - 4*b*ArcCosh[c*x]*Log[1 + E^ArcCosh[c*x]] + 4*a*Log[x] - 2*a*Log[1 - c^2*x^2] - 2*b*PolyLog[2, -E^(-2*ArcCosh[c*x])] - 4*b*PolyLog[2, -E^ArcCosh[c*x]] - 4*b*PolyLog[2, E^ArcCosh[c*x]])/(4*d^2)`

### 3.42.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.66 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6351, 27, 41, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^2} dx \\
 & \quad \downarrow \text{6351} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{dx(1 - c^2 x^2)} dx}{d} + \frac{bc \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2 x^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2 x^2)} dx}{d^2} + \frac{bc \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2 x^2)} \\
 & \quad \downarrow \text{41} \\
 & \frac{\int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2 x^2)} dx}{d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2 x^2)} - \frac{bcx}{2d^2 \sqrt{cx - 1} \sqrt{cx + 1}} \\
 & \quad \downarrow \text{6331} \\
 & -\frac{\int \frac{a + \operatorname{barccosh}(cx)}{cx \sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2 x^2)} - \frac{bcx}{2d^2 \sqrt{cx - 1} \sqrt{cx + 1}}
 \end{aligned}$$

---

3.42.  $\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^2} dx$



$$\begin{aligned}
& \downarrow 5984 \\
& \frac{2 \int (a + \operatorname{barccosh}(cx)) \operatorname{csch}(2 \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2x^2)} - \frac{bcx}{2d^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 3042 \\
& \frac{2 \int i(a + \operatorname{barccosh}(cx)) \operatorname{csc}(2i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2x^2)} - \frac{bcx}{2d^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 26 \\
& \frac{2i \int (a + \operatorname{barccosh}(cx)) \operatorname{csc}(2i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2x^2)} - \frac{bcx}{2d^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 4670 \\
& \frac{2i \left( \frac{1}{2} ib \int \log(1 - e^{2 \operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} ib \int \log(1 + e^{2 \operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + i \operatorname{arctanh}(e^{2 \operatorname{arccosh}(cx)}) \right)}{d^2} \\
& \quad + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2x^2)} - \frac{bcx}{2d^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 2715 \\
& \frac{2i \left( \frac{1}{4} ib \int e^{-2 \operatorname{arccosh}(cx)} \log(1 - e^{2 \operatorname{arccosh}(cx)}) de^{2 \operatorname{arccosh}(cx)} - \frac{1}{4} ib \int e^{-2 \operatorname{arccosh}(cx)} \log(1 + e^{2 \operatorname{arccosh}(cx)}) de^{2 \operatorname{arccosh}(cx)} \right)}{d^2} \\
& \quad + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2x^2)} - \frac{bcx}{2d^2\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 2838 \\
& \frac{2i \left( i \operatorname{arctanh}(e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} ib \operatorname{PolyLog}(2, -e^{2 \operatorname{arccosh}(cx)}) - \frac{1}{4} ib \operatorname{PolyLog}(2, e^{2 \operatorname{arccosh}(cx)}) \right)}{d^2} \\
& \quad + \frac{a + \operatorname{barccosh}(cx)}{2d^2(1 - c^2x^2)} - \frac{bcx}{2d^2\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^2), x]`

output `-1/2*(b*c*x)/(d^2*sqrt[-1 + c*x]*sqrt[1 + c*x]) + (a + b*ArcCosh[c*x])/(2*d^2*(1 - c^2*x^2)) - ((2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanH[E^(2*ArcCosh[c*x])]) + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])]) - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])])/d^2`

## 3.42.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 41 `Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/(f*fz*I)], x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 5984 `Int[Csch[(a_) + (b_)*(x_)]^(n_)*((c_) + (d_)*(x_))^(m_)*Sech[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)/((x_.)*((d_.) + (e_.)*(x_.)^2)), x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^m)*((d_.) + (e_.)*(x_.)^2)^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

### 3.42.4 Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 254, normalized size of antiderivative = 2.19

method	result
parts	$\frac{a\left(\frac{1}{4cx+4} - \frac{\ln(cx+1)}{2} + \ln(x) - \frac{1}{4(cx-1)} - \frac{\ln(cx-1)}{2}\right)}{d^2} + \frac{b\left(-\frac{\sqrt{cx-1}\sqrt{cx+1}cx-c^2x^2+\operatorname{arccosh}(cx)+1}{2(c^2x^2-1)} + \operatorname{arccosh}(cx) \ln\left(1+(cx-1)^2\right)\right)}{d^2}$
derivativedivides	$\frac{a\left(\ln(cx) + \frac{1}{4cx+4} - \frac{\ln(cx+1)}{2} - \frac{1}{4(cx-1)} - \frac{\ln(cx-1)}{2}\right)}{d^2} + \frac{b\left(-\frac{\sqrt{cx-1}\sqrt{cx+1}cx-c^2x^2+\operatorname{arccosh}(cx)+1}{2(c^2x^2-1)} + \operatorname{arccosh}(cx) \ln\left(1+(cx-1)^2\right)\right)}{d^2}$
default	$\frac{a\left(\ln(cx) + \frac{1}{4cx+4} - \frac{\ln(cx+1)}{2} - \frac{1}{4(cx-1)} - \frac{\ln(cx-1)}{2}\right)}{d^2} + \frac{b\left(-\frac{\sqrt{cx-1}\sqrt{cx+1}cx-c^2x^2+\operatorname{arccosh}(cx)+1}{2(c^2x^2-1)} + \operatorname{arccosh}(cx) \ln\left(1+(cx-1)^2\right)\right)}{d^2}$

input `int((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)`

output `a/d^2*(1/4/(c*x+1)-1/2*ln(c*x+1)+ln(x)-1/4/(c*x-1)-1/2*ln(c*x-1))+b/d^2*(-1/2*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-c^2*x^2+arccosh(c*x)+1)/(c^2*x^2-1)+arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)+1/2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)-arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)))`

$$3.42. \int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)^2} dx$$

## 3.42.5 Fricas [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(c^4*d^2*x^5 - 2*c^2*d^2*x^3 + d^2*x), x)`

## 3.42.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^2} dx = \int \frac{a}{c^4 x^5 - 2c^2 x^3 + x} dx + \int \frac{b \operatorname{acosh}(cx)}{c^4 x^5 - 2c^2 x^3 + x} dx$$

input `integrate((a+b*acosh(c*x))/x/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a/(c**4*x**5 - 2*c**2*x**3 + x), x) + Integral(b*acosh(c*x)/(c**4*x**5 - 2*c**2*x**3 + x), x))/d**2`

## 3.42.7 Maxima [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `-1/2*a*(1/(c^2*d^2*x^2 - d^2) + log(c*x + 1)/d^2 + log(c*x - 1)/d^2 - 2*log(x)/d^2) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(c^4*d^2*x^5 - 2*c^2*d^2*x^3 + d^2*x), x)`

**3.42.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^2*x), x)`

**3.42.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(d - c^2 dx^2)^2} dx$$

input `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^2),x)`

output `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^2), x)`

### 3.43 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d-c^2dx^2)^2} dx$

3.43.1	Optimal result	517
3.43.2	Mathematica [A] (verified)	518
3.43.3	Rubi [C] (verified)	518
3.43.4	Maple [A] (verified)	523
3.43.5	Fricas [F]	524
3.43.6	Sympy [F]	524
3.43.7	Maxima [F]	524
3.43.8	Giac [F]	525
3.43.9	Mupad [F(-1)]	525

#### 3.43.1 Optimal result

Integrand size = 25, antiderivative size = 170

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^2(d - c^2dx^2)^2} dx = -\frac{bc}{2d^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{a + b\operatorname{arccosh}(cx)}{d^2x(1 - c^2x^2)} + \frac{3c^2x(a + b\operatorname{arccosh}(cx))}{2d^2(1 - c^2x^2)} + \frac{bc \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d^2} + \frac{3c(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{d^2} + \frac{3bc \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{2d^2} - \frac{3bc \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{2d^2}$$

output

```
(-a-b*arccosh(c*x))/d^2/x/(-c^2*x^2+1)+3/2*c^2*x*(a+b*arccosh(c*x))/d^2/(-c^2*x^2+1)+b*c*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))/d^2+3*c*(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/d^2+3/2*b*c*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/d^2-3/2*b*c*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/d^2-1/2*b*c/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.43.2 Mathematica [A] (verified)

Time = 0.54 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.66

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx$$

$$= -\frac{4a}{x} + bc \sqrt{\frac{-1+cx}{1+cx}} + \frac{bc \sqrt{\frac{-1+cx}{1+cx}}}{1-cx} + \frac{bc^2 x \sqrt{\frac{-1+cx}{1+cx}}}{1-cx} - \frac{2ac^2 x}{-1+c^2 x^2} - \frac{4b \operatorname{arccosh}(cx)}{x} + \frac{b \operatorname{arccosh}(cx)}{1-cx} - \frac{b \operatorname{arccosh}(cx)}{1+cx} + \frac{4bc}{1+cx}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^2), x]`

output `((-4*a)/x + b*c*Sqrt[(-1 + c*x)/(1 + c*x)] + (b*c*Sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) + (b*c^2*x*Sqrt[(-1 + c*x)/(1 + c*x)]/(1 - c*x) - (2*a*c^2*x)/(-1 + c^2*x^2) - (4*b*ArcCosh[c*x])/x + (b*c*ArcCosh[c*x])/(1 - c*x) - (b*c*ArcCosh[c*x])/(1 + c*x) + (4*b*c*Sqrt[-1 + c^2*x^2]*ArcTan[Sqrt[-1 + c^2*x^2]])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - 6*b*c*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] + 6*b*c*ArcCosh[c*x]*Log[1 + E^ArcCosh[c*x]] - 3*a*c*Log[1 - c*x] + 3*a*c*Log[1 + c*x] + 6*b*c*PolyLog[2, -E^ArcCosh[c*x]] - 6*b*c*PolyLog[2, E^ArcCosh[c*x]])/(4*d^2)`

### 3.43.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.85 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.15, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.560$ , Rules used = {6347, 27, 115, 27, 103, 218, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx$$

$$\downarrow 6347$$

$$3c^2 \int \frac{a + b \operatorname{arccosh}(cx)}{d^2 (1 - c^2 x^2)^2} dx - \frac{bc \int \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx}{d^2} - \frac{a + b \operatorname{arccosh}(cx)}{d^2 x (1 - c^2 x^2)}$$

$$\downarrow 27$$

$$\begin{aligned}
& \frac{3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{d^2} - \frac{bc \int \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{d^2x(1-c^2x^2)} \\
& \quad \downarrow 115 \\
& \frac{3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{d^2} - \frac{bc \left( -\frac{\int \frac{c}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{d^2x(1-c^2x^2)} \\
& \quad \downarrow 27 \\
& \frac{3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{d^2} - \frac{bc \left( -\int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{d^2x(1-c^2x^2)} \\
& \quad \downarrow 103 \\
& \frac{3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{d^2} - \frac{bc \left( -c \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{d^2x(1-c^2x^2)} \\
& \quad \downarrow 218 \\
& \frac{3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{d^2x(1-c^2x^2)} - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
& \quad \downarrow 6316 \\
& \frac{3c^2 \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2} bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{d^2x(1-c^2x^2)} - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
& \quad \downarrow 83 \\
& \frac{3c^2 \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{d^2x(1-c^2x^2)} - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
& \quad \downarrow 6318
\end{aligned}$$



$$\begin{aligned}
 & \frac{3c^2 \left( -\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{2c} + \frac{x(a+b\operatorname{arccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{arccosh}(cx)}{d^2x(1-c^2x^2)} \\
 & \quad - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3c^2 \left( -\frac{\int i(a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+b\operatorname{arccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \\
 & \quad \frac{a + \operatorname{arccosh}(cx)}{d^2x(1-c^2x^2)} - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
 & \quad \downarrow \text{26} \\
 & \frac{3c^2 \left( -\frac{i \int (a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+b\operatorname{arccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \\
 & \quad \frac{a + \operatorname{arccosh}(cx)}{d^2x(1-c^2x^2)} - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
 & \quad \downarrow \text{4670} \\
 & \frac{3c^2 \left( -\frac{i \left( ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) \right)}{2c} \right)}{d^2} - \\
 & \quad \frac{a + \operatorname{arccosh}(cx)}{d^2x(1-c^2x^2)} - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
 & \quad \downarrow \text{2715} \\
 & \frac{3c^2 \left( -\frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right)}{d^2} - \\
 & \quad \frac{a + \operatorname{arccosh}(cx)}{d^2x(1-c^2x^2)} - \frac{bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} \\
 & \quad \downarrow \text{2838}
 \end{aligned}$$

---

3.43.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d-c^2dx^2)^2} dx$

$$3c^2 \left( -\frac{i(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a + b \operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{2c} + \frac{x(a + b \operatorname{arccosh}(cx))}{2(1 - c^2 x^2)} \right)$$


---


$$\frac{a + b \operatorname{arccosh}(cx)}{d^2 x (1 - c^2 x^2)} - \frac{bc \left( -\arctan \left( \frac{d^2}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2}$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^2), x]`

output `-(a + b*ArcCosh[c*x])/(d^2*x*(1 - c^2*x^2)) - (b*c*(-(1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) - ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]]))/d^2 + (3*c^2*(-1/2*b/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c))/d^2`

### 3.43.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

- rule 115 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

### 3.43.4 Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.22

method	result
derivativedivides	$c \left( \frac{a \left( -\frac{1}{cx} - \frac{1}{4(cx+1)} + \frac{3 \ln(cx+1)}{4} - \frac{1}{4(cx-1)} - \frac{3 \ln(cx-1)}{4} \right)}{d^2} + \frac{b \left( -\frac{3c^2 x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - 2 \operatorname{arccosh}(cx)}{2cx(c^2 x^2 - 1)} + 2 \operatorname{arctan} \left( \frac{\sqrt{cx-1} \sqrt{cx+1}}{c} \right)}{c} \right)}{d^2} \right)$
default	$c \left( \frac{a \left( -\frac{1}{cx} - \frac{1}{4(cx+1)} + \frac{3 \ln(cx+1)}{4} - \frac{1}{4(cx-1)} - \frac{3 \ln(cx-1)}{4} \right)}{d^2} + \frac{b \left( -\frac{3c^2 x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - 2 \operatorname{arccosh}(cx)}{2cx(c^2 x^2 - 1)} + 2 \operatorname{arctan} \left( \frac{\sqrt{cx-1} \sqrt{cx+1}}{c} \right)}{c} \right)}{d^2} \right)$
parts	$\frac{a \left( -\frac{c}{4(cx+1)} + \frac{3c \ln(cx+1)}{4} - \frac{1}{x} - \frac{c}{4(cx-1)} - \frac{3c \ln(cx-1)}{4} \right)}{d^2} + \frac{bc \left( -\frac{3c^2 x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - 2 \operatorname{arccosh}(cx)}{2cx(c^2 x^2 - 1)} + 2 \operatorname{arctan} \left( \frac{\sqrt{cx-1} \sqrt{cx+1}}{c} \right) \right)}{d^2}$

input `int((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)`

output `c*(a/d^2*(-1/c/x-1/4/(c*x+1)+3/4*ln(c*x+1)-1/4/(c*x-1)-3/4*ln(c*x-1))+b/d^2*(-1/2*(3*c^2*x^2*arccosh(c*x)+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-2*arccosh(c*x))/c/x/(c^2*x^2-1)+2*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+3/2*dilog(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+3/2*dilog(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+3/2*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))`

$$3.43. \int \frac{a+b \operatorname{arccosh}(cx)}{x^2(d-c^2 dx^2)^2} dx$$

**3.43.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2), x)`

**3.43.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx = \int \frac{\frac{a}{c^4 x^6 - 2c^2 x^4 + x^2} dx}{d^2} + \int \frac{b \operatorname{acosh}(cx)}{c^4 x^6 - 2c^2 x^4 + x^2} dx$$

input `integrate((a+b*acosh(c*x))/x**2/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a/(c**4*x**6 - 2*c**2*x**4 + x**2), x) + Integral(b*acosh(c*x)/(c**4*x**6 - 2*c**2*x**4 + x**2), x))/d**2`

**3.43.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output  $\frac{1}{64}(576c^5 \int \frac{1}{8x^3 \log(cx-1)} / (c^4 d^2 x^4 - 2c^2 d^2 x^2 + d^2), x) - 24c^4 (2x / (c^4 d^2 x^2 - c^2 d^2) + \log(cx+1) / (c^3 d^2) - \log(cx-1) / (c^3 d^2)) - 192c^4 \int \frac{1}{8x^2 \log(cx-1)} / (c^4 d^2 x^4 - 2c^2 d^2 x^2 + d^2), x) + 9(c(2 / (c^4 d^2 x - c^3 d^2) - \log(cx+1) / (c^3 d^2) + \log(cx-1) / (c^3 d^2)) + 4 \log(cx-1) / (c^4 d^2 x^2 - c^2 d^2)) c^3 + 16c^2 (2x / (c^2 d^2 x^2 - d^2) - \log(cx+1) / (c d^2) + \log(cx-1) / (c d^2)) + 192c^2 \int \frac{1}{8 \log(cx-1)} / (c^4 d^2 x^4 - 2c^2 d^2 x^2 + d^2), x) - 4(3(c^3 x^3 - cx) \log(cx+1)^2 + 6(c^3 x^3 - cx) \log(cx+1) \log(cx-1) + 4(6c^2 x^2 - 3(c^3 x^3 - cx) \log(cx+1) + 3(c^3 x^3 - cx) \log(cx-1) - 4) \log(cx + \sqrt{cx+1}) \sqrt{cx-1})) / (c^2 d^2 x^3 - d^2 x) + 64 \int (-1/4(6c^3 x^2 - 3(c^4 x^3 - c^2 x) \log(cx+1) + 3(c^4 x^3 - c^2 x) \log(cx-1) - 4c) / (c^5 d^2 x^6 - 2c^3 d^2 x^4 + c d^2 x^2 + (c^4 d^2 x^5 - 2c^2 d^2 x^3 + d^2 x) \sqrt{cx+1} \sqrt{cx-1})), x) * b - 1/4 a (2(3c^2 x^2 - 2) / (c^2 d^2 x^3 - d^2 x) - 3c \log(cx+1) / d^2 + 3c \log(cx-1) / d^2)$

### 3.43.8 Giac [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^2*x^2), x)`

### 3.43.9 Mupad [F(-1)]

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2 (d - c^2 dx^2)^2} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^2),x)`

output `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^2), x)`

### 3.44 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^2} dx$

3.44.1	Optimal result	526
3.44.2	Mathematica [A] (warning: unable to verify)	526
3.44.3	Rubi [C] (verified)	527
3.44.4	Maple [A] (verified)	532
3.44.5	Fricas [F]	532
3.44.6	Sympy [F]	533
3.44.7	Maxima [F]	533
3.44.8	Giac [F]	533
3.44.9	Mupad [F(-1)]	534

#### 3.44.1 Optimal result

Integrand size = 25, antiderivative size = 152

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^3(d - c^2dx^2)^2} dx = -\frac{bc}{2d^2x\sqrt{-1+cx}\sqrt{1+cx}} + \frac{c^2(a + \operatorname{arccosh}(cx))}{d^2(1 - c^2x^2)} - \frac{a + \operatorname{arccosh}(cx)}{2d^2x^2(1 - c^2x^2)} + \frac{4c^2(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d^2} + \frac{bc^2 \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{d^2} - \frac{bc^2 \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{d^2}$$

output  $c^2*(a+b*\operatorname{arccosh}(c*x))/d^2/(-c^2*x^2+1)+1/2*(-a-b*\operatorname{arccosh}(c*x))/d^2/x^2/(-c^2*x^2+1)+4*c^2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/d^2+b*c^2*\operatorname{polylog}(2, -(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/d^2-b*c^2*\operatorname{polylog}(2, (c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)/d^2-1/2*b*c/d^2/x/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}$

#### 3.44.2 Mathematica [A] (warning: unable to verify)

Time = 1.16 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.90

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^3(d - c^2dx^2)^2} dx = -\frac{a}{x^2} + \frac{ac^2}{1-c^2x^2} + 4ac^2 \log(x) - 2ac^2 \log(1 - c^2x^2) + \frac{1}{2}b \left( \frac{2cx\sqrt{-1+cx}\sqrt{1+cx}-2\operatorname{arccosh}(cx)}{x^2} + c^2 \left( -\frac{1}{\sqrt{-1+cx}} + \operatorname{arccosh}(cx) \right) \right)$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^2), x]`

output 
$$\begin{aligned} & \left( -\frac{a}{x^2} + \frac{a*c^2}{(1 - c^2*x^2)} + 4*a*c^2*\text{Log}[x] - 2*a*c^2*\text{Log}[1 - c^2*x^2] \right. \\ & + \frac{b*((2*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x] - 2*\text{ArcCosh}[c*x])}{x^2} + c^2* \\ & \left. \left( -\frac{1}{\text{Sqrt}[(-1 + c*x)/(1 + c*x)]} + \text{ArcCosh}[c*x]/(1 - c*x) \right) - c^2*(\text{Sqrt}[(-1 + c*x)/(1 + c*x)] \right. \right. \\ & \left. - \text{ArcCosh}[c*x]/(1 + c*x)) + 4*c^2*(\text{ArcCosh}[c*x]*(\text{ArcCos} \right. \\ & \left. \text{h}[c*x] + 2*\text{Log}[1 + E^{(-2*\text{ArcCosh}[c*x])}] - \text{PolyLog}[2, -E^{(-2*\text{ArcCosh}[c*x])}] \right. \\ & \left. \left. \right) + 2*c^2*(\text{ArcCosh}[c*x]*(\text{ArcCosh}[c*x] - 4*\text{Log}[1 + E^{\text{ArcCosh}[c*x]}]) - 4*\text{PolyLog}[2, -E^{\text{ArcCosh}[c*x]}]) \right. \\ & \left. \left. + 2*c^2*(\text{ArcCosh}[c*x]*(\text{ArcCosh}[c*x] - 4*\text{Log}[1 - E^{\text{ArcCosh}[c*x]}]) - 4*\text{PolyLog}[2, E^{\text{ArcCosh}[c*x]}]) \right) \right) / (2*d^2) \end{aligned}$$

### 3.44.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.96 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.36, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.560$ , Rules used = {6347, 27, 114, 27, 41, 6351, 41, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \text{barccosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx \\ & \quad \downarrow \text{6347} \\ & 2c^2 \int \frac{a + \text{barccosh}(cx)}{d^2 x (1 - c^2 x^2)^2} dx - \frac{bc \int \frac{1}{x^2 (cx-1)^{3/2} (cx+1)^{3/2}} dx}{2d^2} - \frac{a + \text{barccosh}(cx)}{2d^2 x^2 (1 - c^2 x^2)} \\ & \quad \downarrow \text{27} \\ & \frac{2c^2 \int \frac{a + \text{barccosh}(cx)}{x(1 - c^2 x^2)^2} dx}{d^2} - \frac{bc \int \frac{1}{x^2 (cx-1)^{3/2} (cx+1)^{3/2}} dx}{2d^2} - \frac{a + \text{barccosh}(cx)}{2d^2 x^2 (1 - c^2 x^2)} \\ & \quad \downarrow \text{114} \\ & \frac{2c^2 \int \frac{a + \text{barccosh}(cx)}{x(1 - c^2 x^2)^2} dx}{d^2} - \frac{bc \left( \int \frac{2c^2}{(cx-1)^{3/2} (cx+1)^{3/2}} dx + \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} - \frac{a + \text{barccosh}(cx)}{2d^2 x^2 (1 - c^2 x^2)} \\ & \quad \downarrow \text{27} \\ & \frac{2c^2 \int \frac{a + \text{barccosh}(cx)}{x(1 - c^2 x^2)^2} dx}{d^2} - \frac{bc \left( 2c^2 \int \frac{1}{(cx-1)^{3/2} (cx+1)^{3/2}} dx + \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} - \frac{a + \text{barccosh}(cx)}{2d^2 x^2 (1 - c^2 x^2)} \end{aligned}$$

---

3.44.  $\int \frac{a + \text{barccosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx$



$$\begin{aligned}
& \downarrow 41 \\
& \frac{2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} \\
& \downarrow 6351 \\
& \frac{2c^2 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2}bc \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \\
& \quad \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} \\
& \downarrow 41 \\
& \frac{2c^2 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \\
& \quad \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} \\
& \downarrow 6331 \\
& \frac{2c^2 \left( - \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \\
& \quad \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} \\
& \downarrow 5984 \\
& \frac{2c^2 \left( -2 \int (a + \operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} \\
& \downarrow 3042 \\
& \frac{2c^2 \left( -2 \int i(a + \operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2} \\
& \downarrow 26
\end{aligned}$$

$$\frac{2c^2 \left( -2i \int (a + \operatorname{barccosh}(cx)) \csc(2i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a + \operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2}$$

↓ 4670

$$\frac{2c^2 \left( -2i \left( \frac{1}{2} ib \int \log(1 - e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} ib \int \log(1 + e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + i \operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) \right) \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2}$$

↓ 2715

$$\frac{2c^2 \left( -2i \left( \frac{1}{4} ib \int e^{-2\operatorname{arccosh}(cx)} \log(1 - e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{4} ib \int e^{-2\operatorname{arccosh}(cx)} \log(1 + e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} \right) \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2}$$

↓ 2838

$$\frac{2c^2 \left( -2i \left( i \operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4} ib \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right) \right)}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2(1-c^2x^2)} - \frac{bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{2d^2}$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^2), x]`

output `-1/2*(b*c*(1/(x*sqrt[-1 + c*x]*sqrt[1 + c*x]) - (2*c^2*x)/(sqrt[-1 + c*x]*sqrt[1 + c*x]))) / d^2 - (a + b*ArcCosh[c*x]) / (2*d^2*x^2*(1 - c^2*x^2)) + (2*c^2*(-1/2*(b*c*x)/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (a + b*ArcCosh[c*x]) / (2*(1 - c^2*x^2)) - (2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])]) + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])])) / d^2`

## 3.44.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 41 `Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`
- rule 114 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[-(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

### 3.44.4 Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.84

method	result
derivativedivides	$c^2 \left( \frac{a \left( -\frac{1}{2c^2x^2} + 2 \ln(cx) + \frac{1}{4cx+4} - \ln(cx+1) - \frac{1}{4(cx-1)} - \ln(cx-1) \right)}{d^2} + \frac{b \left( -\frac{2c^2x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - \operatorname{arccosh}(cx)}{2(c^2x^2-1)c^2x^2} \right)}{d^2} \right)$
default	$c^2 \left( \frac{a \left( -\frac{1}{2c^2x^2} + 2 \ln(cx) + \frac{1}{4cx+4} - \ln(cx+1) - \frac{1}{4(cx-1)} - \ln(cx-1) \right)}{d^2} + \frac{b \left( -\frac{2c^2x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - \operatorname{arccosh}(cx)}{2(c^2x^2-1)c^2x^2} \right)}{d^2} \right)$
parts	$\frac{a \left( \frac{c^2}{4cx+4} - c^2 \ln(cx+1) - \frac{1}{2x^2} + 2c^2 \ln(x) - \frac{c^2}{4(cx-1)} - c^2 \ln(cx-1) \right)}{d^2} + \frac{b c^2 \left( -\frac{2c^2x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - \operatorname{arccosh}(cx)}{2(c^2x^2-1)c^2x^2} \right)}{d^2}$

input `int((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)`

output 
$$c^2 \cdot \left( \frac{a}{d^2} \cdot \left( -\frac{1}{2} \cdot \frac{1}{c^2 x^2} + 2 \ln(cx) + \frac{1}{4} \cdot \frac{1}{(cx+1)} - \ln(cx+1) - \frac{1}{4} \cdot \frac{1}{(cx-1)} - \ln(cx-1) \right) + \frac{b}{d^2} \cdot \left( -\frac{1}{2} \cdot \frac{2c^2x^2 \operatorname{arccosh}(cx) + (cx-1)^{1/2} (cx+1)^{1/2} cx - \operatorname{arccosh}(cx)}{(c^2x^2-1)c^2x^2} + \frac{1}{x^2} \cdot \frac{2c^2x^2 \operatorname{arccosh}(cx) * \ln(1+(cx+(cx-1)^{1/2} * (cx+1)^{1/2}))^2 + \operatorname{polylog}(2, -(cx+(cx-1)^{1/2} * (cx+1)^{1/2}))^2 - 2 \operatorname{arccosh}(cx) * \ln(1-cx-(cx-1)^{1/2} * (cx+1)^{1/2}) - 2 \operatorname{polylog}(2, cx+(cx-1)^{1/2} * (cx+1)^{1/2}) - 2 \operatorname{arccosh}(cx) * \ln(1+cx+(cx-1)^{1/2} * (cx+1)^{1/2}) - 2 \operatorname{polylog}(2, -cx-(cx-1)^{1/2} * (cx+1)^{1/2}))}{d^2} \right) \right)$$

### 3.44.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^2 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(c^4*d^2*x^7 - 2*c^2*d^2*x^5 + d^2*x^3), x)`

## 3.44.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx = \int \frac{a}{c^4 x^7 - 2c^2 x^5 + x^3} dx + \int \frac{b \operatorname{arcosh}(cx)}{c^4 x^7 - 2c^2 x^5 + x^3} dx$$

input `integrate((a+b*acosh(c*x))/x**3/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a/(c**4*x**7 - 2*c**2*x**5 + x**3), x) + Integral(b*acosh(c*x)/(c**4*x**7 - 2*c**2*x**5 + x**3), x))/d**2`

## 3.44.7 Maxima [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `-1/2*a*(2*c^2*log(c*x + 1)/d^2 + 2*c^2*log(c*x - 1)/d^2 - 4*c^2*log(x)/d^2 + (2*c^2*x^2 - 1)/(c^2*d^2*x^4 - d^2*x^2)) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(c^4*d^2*x^7 - 2*c^2*d^2*x^5 + d^2*x^3), x)`

## 3.44.8 Giac [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^2*x^3), x)`

**3.44.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (d - c^2 dx^2)^2} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^2), x)`output `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^2), x)`

### 3.45 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^4(d-c^2dx^2)^2} dx$

3.45.1	Optimal result	535
3.45.2	Mathematica [A] (verified)	536
3.45.3	Rubi [C] (verified)	536
3.45.4	Maple [A] (verified)	543
3.45.5	Fricas [F]	543
3.45.6	Sympy [F]	544
3.45.7	Maxima [F]	544
3.45.8	Giac [F]	545
3.45.9	Mupad [F(-1)]	545

#### 3.45.1 Optimal result

Integrand size = 25, antiderivative size = 248

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^4(d - c^2dx^2)^2} dx = -\frac{bc^3}{3d^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc}{6d^2x^2\sqrt{-1+cx}\sqrt{1+cx}}$$

$$-\frac{a + b\operatorname{arccosh}(cx)}{3d^2x^3(1 - c^2x^2)} - \frac{5c^2(a + b\operatorname{arccosh}(cx))}{3d^2x(1 - c^2x^2)}$$

$$+\frac{5c^4x(a + b\operatorname{arccosh}(cx))}{2d^2(1 - c^2x^2)} + \frac{13bc^3 \arctan(\sqrt{-1+cx}\sqrt{1+cx})}{6d^2}$$

$$+\frac{5c^3(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{d^2}$$

$$+\frac{5bc^3 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{2d^2} - \frac{5bc^3 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{2d^2}$$

output

```
1/3*(-a-b*arccosh(c*x))/d^2/x^3/(-c^2*x^2+1)-5/3*c^2*(a+b*arccosh(c*x))/d^
2/x/(-c^2*x^2+1)+5/2*c^4*x*(a+b*arccosh(c*x))/d^2/(-c^2*x^2+1)+13/6*b*c^3*
arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))/d^2+5*c^3*(a+b*arccosh(c*x))*arctanh(c
*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/d^2+5/2*b*c^3*polylog(2,-c*x-(c*x-1)^(1/2)
*(c*x+1)^(1/2))/d^2-5/2*b*c^3*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/d
^2-1/3*b*c^3/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/6*b*c/d^2/x^2/(c*x-1)^(1/2)
/(c*x+1)^(1/2)
```



### 3.45.2 Mathematica [A] (verified)

Time = 1.30 (sec) , antiderivative size = 377, normalized size of antiderivative = 1.52

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx =$$

$$-\frac{4a}{x^3} + \frac{24ac^2}{x} - 3bc^3 \sqrt{\frac{-1+cx}{1+cx}} + \frac{3bc^3 \sqrt{\frac{-1+cx}{1+cx}}}{-1+cx} + \frac{3bc^4 x \sqrt{\frac{-1+cx}{1+cx}}}{-1+cx} - \frac{2bc^3}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{2bc}{x^2 \sqrt{-1+cx}\sqrt{1+cx}} + \frac{6ac^4 x}{-1+c^2 x^2} + \frac{4ba}{d}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^2), x]`

output

$$-1/12*((4*a)/x^3 + (24*a*c^2)/x - 3*b*c^3*\sqrt{(-1 + c*x)/(1 + c*x)} + (3*b*c^3*\sqrt{(-1 + c*x)/(1 + c*x)})/(-1 + c*x) + (3*b*c^4*x*\sqrt{(-1 + c*x)/(1 + c*x)})/(-1 + c*x) - (2*b*c^3)/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (2*b*c)/(x^2*sqrt[-1 + c*x]*sqrt[1 + c*x]) + (6*a*c^4*x)/(-1 + c^2*x^2) + (4*b*ArcCosh[c*x])/x^3 + (24*b*c^2*ArcCosh[c*x])/x + (3*b*c^3*ArcCosh[c*x])/(-1 + c*x) + (3*b*c^3*ArcCosh[c*x])/(1 + c*x) - (26*b*c^3*sqrt[-1 + c^2*x^2]*ArcTan[sqrt[-1 + c^2*x^2]])/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + 30*b*c^3*ArcCosh[c*x]*Log[1 - E^ArcCosh[c*x]] - 30*b*c^3*ArcCosh[c*x]*Log[1 + E^ArcCosh[c*x]] + 15*a*c^3*Log[1 - c*x] - 15*a*c^3*Log[1 + c*x] - 30*b*c^3*PolyLog[2, -E^ArcCosh[c*x]] + 30*b*c^3*PolyLog[2, E^ArcCosh[c*x]])/d^2$$

### 3.45.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.29 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.26, number of steps used = 22, number of rules used = 21,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.840$ , Rules used = {6347, 27, 114, 27, 115, 27, 103, 218, 6347, 115, 27, 103, 218, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx$$

$$\downarrow \text{6347}$$

$$\frac{5}{3}c^2 \int \frac{a + b \operatorname{arccosh}(cx)}{d^2 x^2 (1 - c^2 x^2)^2} dx - \frac{bc \int \frac{1}{x^3 (cx-1)^{3/2} (cx+1)^{3/2}} dx}{3d^2} - \frac{a + b \operatorname{arccosh}(cx)}{3d^2 x^3 (1 - c^2 x^2)}$$

---

3.45.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{3d^2} - \frac{bc \int \frac{1}{x^3(cx-1)^{3/2}(cx+1)^{3/2}} dx}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} \\
& \downarrow 114 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{3d^2} - \frac{bc \left( \frac{1}{2} \int \frac{3c^2}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} \\
& \downarrow 27 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \int \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} \\
& \downarrow 115 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -\int \frac{\frac{c}{x\sqrt{cx-1}\sqrt{cx+1}}}{c} dx - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} \\
& \downarrow 27 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -\int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} \\
& \downarrow 103 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -c \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} \\
& \downarrow 218 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \\
& \downarrow 6347
\end{aligned}$$

---

3.45.  $\int \frac{a+\operatorname{barccosh}(cx)}{x^4(d-c^2dx^2)^2} dx$

$$\frac{5c^2 \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - bc \int \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} \right)}{3d^2} - \frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} -$$

$$\frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2}$$

↓ 115

$$\frac{5c^2 \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - bc \left( -\int \frac{\frac{c}{x\sqrt{cx-1}\sqrt{cx+1}}}{c} dx - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} \right)}{3d^2} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2}$$

↓ 27

$$\frac{5c^2 \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - bc \left( -\int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} \right)}{3d^2} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2}$$

↓ 103

$$\frac{5c^2 \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - bc \left( -c \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} \right)}{3d^2} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2}$$

↓ 218

$$\frac{5c^2 \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2}$$

↓ 6316

$$\frac{5c^2 \left( 3c^2 \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2} bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2} -$$

$$\frac{a + \operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2}$$

↓ 83

---

3.45.  $\int \frac{a+\operatorname{barccosh}(cx)}{x^4(d-c^2dx^2)^2} dx$

$$\frac{5c^2 \left( 3c^2 \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) \right) \right)}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \frac{a+\operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)}$$

↓ 6318

$$\frac{5c^2 \left( 3c^2 \left( -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) \right) \right)}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \frac{a+\operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)}$$

↓ 3042

$$\frac{5c^2 \left( 3c^2 \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) \right) \right)}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \frac{a+\operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)}$$

↓ 26

$$\frac{5c^2 \left( 3c^2 \left( -\frac{i \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) \right) \right)}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \frac{a+\operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)}$$

↓ 4670

$$\frac{5c^2 \left( 3c^2 \left( -\frac{i \left( ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx))}{2c} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) \right) \right)}{3d^2} - \frac{bc \left( \frac{3}{2}c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \frac{a+\operatorname{barccosh}(cx)}{3d^2x^3(1-c^2x^2)}$$

↓ 2715

---

3.45.  $\int \frac{a+\operatorname{barccosh}(cx)}{x^4(d-c^2dx^2)^2} dx$

$$5c^2 \left( 3c^2 \left( -\frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) dx + e^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) dx + e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right) \right.$$


---


$$\left. \frac{a + \operatorname{barccosh}(cx)}{3d^2 x^3 (1 - c^2 x^2)} - \frac{bc \left( \frac{3}{2} c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \right)$$

↓ 2838

---


$$5c^2 \left( 3c^2 \left( -\frac{i \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right) \right) + \frac{x(a + \operatorname{barccosh}(cx))}{2(1 - c^2 x^2)}$$


---


$$\frac{a + \operatorname{barccosh}(cx)}{3d^2 x^3 (1 - c^2 x^2)} - \frac{bc \left( \frac{3}{2} c^2 \left( -\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{2x^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \quad 3d^2$$

input `Int[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^2), x]`

output `-1/3*(a + b*ArcCosh[c*x])/(d^2*x^3*(1 - c^2*x^2)) - (b*c*(1/(2*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (3*c^2*(-(1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) - ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]]))/2))/(3*d^2) + (5*c^2*(-((a + b*ArcCosh[c*x])/(x*(1 - c^2*x^2))) - b*c*(-(1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) - ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]]) + 3*c^2*(-1/2*b/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c)))/(3*d^2)`

### 3.45.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x_, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x_, x], x] /; FreeQ[a, x] && !MatchQ[F_x_, (b_)*(G_x_)] /; FreeQ[b, x]`

- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`
- rule 114 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`
- rule 115 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^(e_.)*((c_.) + (d_.)*(x_)))]^(n_.), x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

### 3.45.4 Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.01

method	result
derivativedivides	$c^3 \left( \frac{a \left( -\frac{1}{3c^3x^3} - \frac{2}{cx} - \frac{1}{4(cx+1)} + \frac{5\ln(cx+1)}{4} - \frac{1}{4(cx-1)} - \frac{5\ln(cx-1)}{4} \right)}{d^2} + \frac{b \left( -\frac{15c^4x^4 \operatorname{arccosh}(cx) + 2\sqrt{cx-1}\sqrt{cx+1}c^3x^3 - 10c^2x}{6c^3x^3} \right)}{d^2} \right)$
default	$c^3 \left( \frac{a \left( -\frac{1}{3c^3x^3} - \frac{2}{cx} - \frac{1}{4(cx+1)} + \frac{5\ln(cx+1)}{4} - \frac{1}{4(cx-1)} - \frac{5\ln(cx-1)}{4} \right)}{d^2} + \frac{b \left( -\frac{15c^4x^4 \operatorname{arccosh}(cx) + 2\sqrt{cx-1}\sqrt{cx+1}c^3x^3 - 10c^2x}{6c^3x^3} \right)}{d^2} \right)$
parts	$\frac{a \left( -\frac{c^3}{4(cx+1)} + \frac{5c^3 \ln(cx+1)}{4} - \frac{1}{3x^3} - \frac{2c^2}{x} - \frac{c^3}{4(cx-1)} - \frac{5c^3 \ln(cx-1)}{4} \right)}{d^2} + \frac{b c^3 \left( -\frac{15c^4x^4 \operatorname{arccosh}(cx) + 2\sqrt{cx-1}\sqrt{cx+1}c^3x^3 - 10c^2x}{6c^3x^3} \right)}{d^2}$

input `int((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^2,x,method=_RETURNVERBOSE)`

output `c^3*(a/d^2*(-1/3/c^3/x^3-2/c/x-1/4/(c*x+1)+5/4*ln(c*x+1)-1/4/(c*x-1)-5/4*ln(c*x-1))+b/d^2*(-1/6*(15*c^4*x^4*arccosh(c*x)+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-10*c^2*x^2*arccosh(c*x)+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-2*arccosh(c*x))/c^3/x^3/(c^2*x^2-1)+13/3*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+5/2*dilog(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+5/2*dilog(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+5/2*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))`

### 3.45.5 Fricas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^2 x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(c^4*d^2*x^8 - 2*c^2*d^2*x^6 + d^2*x^4), x)`



### 3.45.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx = \int \frac{a}{c^4 x^8 - 2c^2 x^6 + x^4} dx + \int \frac{b \operatorname{arcosh}(cx)}{c^4 x^8 - 2c^2 x^6 + x^4} dx$$

input `integrate((a+b*acosh(c*x))/x**4/(-c**2*d*x**2+d)**2,x)`

output `(Integral(a/(c**4*x**8 - 2*c**2*x**6 + x**4), x) + Integral(b*acosh(c*x)/(c**4*x**8 - 2*c**2*x**6 + x**4), x))/d**2`

### 3.45.7 Maxima [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `1/12*(15*c^3*log(c*x + 1)/d^2 - 15*c^3*log(c*x - 1)/d^2 - 2*(15*c^4*x^4 - 10*c^2*x^2 - 2)/(c^2*d^2*x^5 - d^2*x^3))*a + 1/192*(8640*c^7*integrate(1/24*x^5*log(c*x - 1)/(c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2), x) - 120*c^6*(2*x/(c^4*d^2*x^2 - c^2*d^2) + log(c*x + 1)/(c^3*d^2) - log(c*x - 1)/(c^3*d^2)) - 2880*c^6*integrate(1/24*x^4*log(c*x - 1)/(c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2), x) + 45*(c*(2/(c^4*d^2*x - c^3*d^2) - log(c*x + 1)/(c^3*d^2) + log(c*x - 1)/(c^3*d^2)) + 4*log(c*x - 1)/(c^4*d^2*x^2 - c^2*d^2))*c^5 + 80*c^4*(2*x/(c^2*d^2*x^2 - d^2) - log(c*x + 1)/(c*d^2) + log(c*x - 1)/(c*d^2)) + 2880*c^4*integrate(1/24*x^2*log(c*x - 1)/(c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2), x) + 16*c^2*(2*(3*c^2*x^2 - 2)/(c^2*d^2*x^3 - d^2*x) - 3*c*log(c*x + 1)/d^2 + 3*c*log(c*x - 1)/d^2) - 4*(15*(c^5*x^5 - c^3*x^3)*log(c*x + 1)^2 + 30*(c^5*x^5 - c^3*x^3)*log(c*x + 1)*log(c*x - 1) + 4*(30*c^4*x^4 - 20*c^2*x^2 - 15*(c^5*x^5 - c^3*x^3)*log(c*x + 1) + 15*(c^5*x^5 - c^3*x^3)*log(c*x - 1) - 4)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(c^2*d^2*x^5 - d^2*x^3) + 192*integrate(-1/12*(30*c^5*x^4 - 20*c^3*x^2 - 15*(c^6*x^5 - c^4*x^3)*log(c*x + 1) + 15*(c^6*x^5 - c^4*x^3)*log(c*x - 1) - 4*c)/(c^5*d^2*x^8 - 2*c^3*d^2*x^6 + c*d^2*x^4 + (c^4*d^2*x^7 - 2*c^2*d^2*x^5 + d^2*x^3)*sqrt(c*x + 1))*sqrt(c*x - 1), x))*b`

**3.45.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^2 x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^2*x^4), x)`

**3.45.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4 (d - c^2 dx^2)^2} dx$$

input `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^2),x)`

output `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^2), x)`

### 3.46 $\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$

3.46.1	Optimal result	546
3.46.2	Mathematica [A] (warning: unable to verify)	547
3.46.3	Rubi [C] (verified)	547
3.46.4	Maple [A] (verified)	552
3.46.5	Fricas [F]	553
3.46.6	Sympy [F]	553
3.46.7	Maxima [F]	554
3.46.8	Giac [F]	554
3.46.9	Mupad [F(-1)]	555

#### 3.46.1 Optimal result

Integrand size = 25, antiderivative size = 249

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \frac{bx^3}{12c^2d^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} + \frac{b}{4c^5d^3\sqrt{-1 + cx}(1 + cx)^{3/2}}$$

$$- \frac{b(-1 + cx)^{3/2}}{12c^5d^3(1 + cx)^{3/2}} + \frac{3b}{8c^5d^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{x^3(a + \operatorname{arccosh}(cx))}{4c^2d^3(1 - c^2x^2)^2} - \frac{3x(a + \operatorname{arccosh}(cx))}{8c^4d^3(1 - c^2x^2)}$$

$$+ \frac{3(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{4c^5d^3}$$

$$+ \frac{3b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{8c^5d^3} - \frac{3b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{8c^5d^3}$$

output  $1/12*b*x^3/c^2/d^3/(c*x-1)^(3/2)/(c*x+1)^(3/2)-1/12*b*(c*x-1)^(3/2)/c^5/d^3/(c*x+1)^(3/2)+1/4*x^3*(a+b*\operatorname{arccosh}(c*x))/c^2/d^3/(-c^2*x^2+1)^2-3/8*x*(a+b*\operatorname{arccosh}(c*x))/c^4/d^3/(-c^2*x^2+1)+3/4*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^5/d^3+3/8*b*\operatorname{polylog}(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^5/d^3-3/8*b*\operatorname{polylog}(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^5/d^3+1/4*b/c^5/d^3/(c*x+1)^(3/2)/(c*x-1)^(1/2)+3/8*b/c^5/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)$

### 3.46.2 Mathematica [A] (warning: unable to verify)

Time = 1.29 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.15

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

$$= \frac{-\frac{b(-2+cx)\sqrt{1+cx}}{(-1+cx)^{3/2}} + \frac{b\sqrt{-1+cx}(2+cx)}{(1+cx)^{3/2}} + \frac{12acx}{(-1+c^2x^2)^2} + \frac{30acx}{-1+c^2x^2} + \frac{3\operatorname{barccosh}(cx)}{(-1+cx)^2} - \frac{3\operatorname{barccosh}(cx)}{(1+cx)^2} - 15b\left(-\frac{1}{\sqrt{\frac{-1+cx}{1+cx}}}\right)}{c^5 d^3}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output `(-((b*(-2 + c*x)*Sqrt[1 + c*x])/(-1 + c*x)^(3/2)) + (b*Sqrt[-1 + c*x]*(2 + c*x))/(1 + c*x)^(3/2) + (12*a*c*x)/(-1 + c^2*x^2)^2 + (30*a*c*x)/(-1 + c^2*x^2) + (3*b*ArcCosh[c*x])/(-1 + c*x)^2 - (3*b*ArcCosh[c*x])/(1 + c*x)^2 - 15*b*(-(1/Sqrt[(-1 + c*x)/(1 + c*x)]) + ArcCosh[c*x]/(1 - c*x)) - 15*b*(Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x]/(1 + c*x)) + (9*b*ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 - E^ArcCosh[c*x]]))/2 - (9*b*ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 + E^ArcCosh[c*x]]))/2 - 9*a*Log[1 - c*x] + 9*a*Log[1 + c*x] + 18*b*PolyLog[2, -E^ArcCosh[c*x]] - 18*b*PolyLog[2, E^ArcCosh[c*x]])/(48*c^5*d^3)`

### 3.46.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.96 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.560$ , Rules used = {6349, 27, 105, 100, 27, 48, 6349, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

$$\downarrow \text{6349}$$

$$-\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))}{d^2(1 - c^2 x^2)^2} dx}{4c^2 d} - \frac{b \int \frac{x^3}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4cd^3} + \frac{x^3(a + \operatorname{barccosh}(cx))}{4c^2 d^3 (1 - c^2 x^2)^2}$$

$$\downarrow \text{27}$$

---

3.46.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$

$$\begin{aligned}
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{4c^2d^3} - \frac{b \int \frac{x^3}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4cd^3} + \frac{x^3(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} \\
& \quad \downarrow 105 \\
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{4c^2d^3} - \frac{b \left( \frac{\int \frac{x^2}{(cx-1)^{3/2}(cx+1)^{5/2}} dx}{c} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} + \frac{x^3(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} \\
& \quad \downarrow 100 \\
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{4c^2d^3} - \frac{b \left( \frac{\int \frac{c\sqrt{cx-1}}{(cx+1)^{5/2}} dx}{c^3} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} + \\
& \quad \frac{x^3(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} \\
& \quad \downarrow 27 \\
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{4c^2d^3} - \frac{b \left( \frac{\int \frac{\sqrt{cx-1}}{(cx+1)^{5/2}} dx}{c^2} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} + \\
& \quad \frac{x^3(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} \\
& \quad \downarrow 48 \\
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{4c^2d^3} + \frac{x^3(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} - \\
& \quad \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3(cx+1)^{3/2}} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} \\
& \quad \downarrow 6349 \\
& -\frac{3 \left( -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx}{2c^2} + \frac{b \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{4c^2d^3} + \frac{x^3(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} - \\
& \quad \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3(cx+1)^{3/2}} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} \\
& \quad \downarrow 83
\end{aligned}$$

---

3.46.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^3} dx$

$$\begin{aligned}
& - \frac{3 \left( - \frac{\int \frac{a+b \operatorname{arccosh}(cx)}{1-c^2x^2} dx}{2c^2} + \frac{x(a+b \operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{4c^2d^3} + \frac{x^3(a+b \operatorname{arccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} - \\
& \quad \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3(cx+1)^{3/2}} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} \\
& \qquad \qquad \qquad \downarrow \text{6318} \\
& - \frac{3 \left( \frac{\int \frac{a+b \operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d \operatorname{arccosh}(cx)}{2c^3} + \frac{x(a+b \operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{4c^2d^3} + \frac{x^3(a+b \operatorname{arccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} - \\
& \quad \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3(cx+1)^{3/2}} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& - \frac{3 \left( \frac{\int i(a+b \operatorname{arccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{2c^3} + \frac{x(a+b \operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{4c^2d^3} + \\
& \quad \frac{x^3(a+b \operatorname{arccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} - \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3(cx+1)^{3/2}} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} \\
& \qquad \qquad \qquad \downarrow \text{26} \\
& - \frac{3 \left( \frac{i \int (a+b \operatorname{arccosh}(cx)) \operatorname{csc}(i \operatorname{arccosh}(cx)) d \operatorname{arccosh}(cx)}{2c^3} + \frac{x(a+b \operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{4c^2d^3} + \\
& \quad \frac{x^3(a+b \operatorname{arccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} - \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3(cx+1)^{3/2}} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} \\
& \qquad \qquad \qquad \downarrow \text{4670} \\
& - \frac{3 \left( \frac{i \left( b \int \log(1-e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) - b \int \log(1+e^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b \operatorname{arccosh}(cx)) \right)}{2c^3} \right)}{4c^2d^3} + \\
& \quad \frac{x^3(a+b \operatorname{arccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} - \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3(cx+1)^{3/2}} - \frac{1}{c^3\sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3}
\end{aligned}$$

---

3.46.  $\int \frac{x^4(a+b \operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$

↓ 2715

$$3 \left( \frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) dx + e^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) dx + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{2c^3} \right)$$

$$\frac{x^3(a + b \operatorname{arccosh}(cx))}{4c^2 d^3 (1 - c^2 x^2)^2} - \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3 (cx+1)^{3/2}} - \frac{1}{c^3 \sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3}$$

↓ 2838

$$3 \left( \frac{i \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c^3} \right) + \frac{x(a + b \operatorname{arccosh}(cx))}{2c^2(1 - c^2 x^2)}$$

$$\frac{x^3(a + b \operatorname{arccosh}(cx))}{4c^2 d^3 (1 - c^2 x^2)^2} - \frac{b \left( \frac{(cx-1)^{3/2}}{3c^3 (cx+1)^{3/2}} - \frac{1}{c^3 \sqrt{cx-1}(cx+1)^{3/2}} - \frac{x^3}{3c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output `-1/4*(b*(-1/3*x^3/(c*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (-1/(c^3*Sqrt[-1 + c*x]*(1 + c*x)^(3/2))) + (-1 + c*x)^(3/2)/(3*c^3*(1 + c*x)^(3/2)))/c)/(c*d^3) + (x^3*(a + b*ArcCosh[c*x]))/(4*c^2*d^3*(1 - c^2*x^2)^2) - (3*(-1/2*b/(c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*c^2*(1 - c^2*x^2)) + ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]])))/c^3)/(4*c^2*d^3)`

### 3.46.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

$$3.46. \int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

- rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp  
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{  
a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p  
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),  
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f  
*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 100 `Int[((a_.) + (b_.)*(x_))^(2)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(  
p_), x_] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d^2*(d  
*e - c*f)*(n + 1))), x] - Simp[1/(d^2*(d*e - c*f)*(n + 1)) Int[(c + d*x)^(  
n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1) + c*f*(  
p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n + 1)*x  
, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n  
+ p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))`
- rule 105 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_  
_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1)/((m +  
1)*(b*e - a*f))), x] - Simp[n*((d*e - c*f)/((m + 1)*(b*e - a*f))) Int[(a  
+ b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d,  
e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1]  
|| !SumSimplerQ[p, 1]) && NeQ[m, -1]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((c_.) + (d_.)*(x_)))]^(n_.), x_Symbol]  
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)  
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2  
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`



```
rule 4670 Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

```
rule 6318 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symb
ol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x
]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]
```

```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_
.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*(m - 1)/(2*e*(p + 1))
Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - S
imp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)]
Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c
*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0]
&& GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.46.4 Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.03

method	result
derivativedivides	$\frac{a \left( \frac{1}{16(cx+1)^2} - \frac{5}{16(cx+1)} - \frac{3 \ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} - \frac{5}{16(cx-1)} + \frac{3 \ln(cx-1)}{16} \right) b \left( -\frac{15c^3 x^3 \operatorname{arccosh}(cx) + 15\sqrt{cx-1}\sqrt{cx+1} c^2 x^2 - 24(c^4 x^4 - 2c^2 x^2)}{24(c^4 x^4 - 2c^2 x^2)} \right)}{d^3}$
default	$\frac{a \left( \frac{1}{16(cx+1)^2} - \frac{5}{16(cx+1)} - \frac{3 \ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} - \frac{5}{16(cx-1)} + \frac{3 \ln(cx-1)}{16} \right) b \left( -\frac{15c^3 x^3 \operatorname{arccosh}(cx) + 15\sqrt{cx-1}\sqrt{cx+1} c^2 x^2 - 24(c^4 x^4 - 2c^2 x^2)}{24(c^4 x^4 - 2c^2 x^2)} \right)}{d^3}$
parts	$\frac{a \left( \frac{1}{16c^5(cx+1)^2} - \frac{5}{16c^5(cx+1)} - \frac{3 \ln(cx+1)}{16c^5} - \frac{1}{16c^5(cx-1)^2} - \frac{5}{16c^5(cx-1)} + \frac{3 \ln(cx-1)}{16c^5} \right) b \left( -\frac{15c^3 x^3 \operatorname{arccosh}(cx) + 15\sqrt{cx-1}\sqrt{cx+1} c^2 x^2 - 24(c^4 x^4 - 2c^2 x^2)}{24(c^4 x^4 - 2c^2 x^2)} \right)}{d^3}$

```
input int(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)
```

$$3.46. \int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$$

output  $1/c^5*(-a/d^3*(1/16/(c*x+1)^2-5/16/(c*x+1)-3/16*\ln(c*x+1)-1/16/(c*x-1)^2-5/16/(c*x-1)+3/16*\ln(c*x-1))-b/d^3*(-1/24*(15*c^3*x^3*\operatorname{arccosh}(c*x)+15*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-9*c*x*\operatorname{arccosh}(c*x)-13*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))/(c^4*x^4-2*c^2*x^2+1)+3/8*\operatorname{arccosh}(c*x)*\ln(1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2))+3/8*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2))-3/8*\operatorname{arccosh}(c*x)*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2))-3/8*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2))}))$

### 3.46.5 Fricas [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^4}{(c^2 dx^2 - d)^3} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`

output `integral(-(b*x^4*arccosh(c*x) + a*x^4)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

### 3.46.6 Sympy [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = -\int \frac{ax^4}{c^6 x^6 - 3c^4 x^4 + 3c^2 x^2 - 1} dx + \int \frac{bx^4 \operatorname{acosh}(cx)}{c^6 x^6 - 3c^4 x^4 + 3c^2 x^2 - 1} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**3,x)`

output `-(Integral(a*x**4/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x) + Integral(b*x**4*acosh(c*x)/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x))/d**3`

## 3.46.7 Maxima [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^4}{(c^2dx^2 - d)^3} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

output `1/2048*(18432*c^5*integrate(1/32*x^5*log(c*x - 1)/(c^10*d^3*x^6 - 3*c^8*d^3*x^4 + 3*c^6*d^3*x^2 - c^4*d^3), x) + 80*c^4*(2*(5*c^2*x^3 - 3*x)/(c^12*d^3*x^4 - 2*c^10*d^3*x^2 + c^8*d^3) + 3*log(c*x + 1)/(c^9*d^3) - 3*log(c*x - 1)/(c^9*d^3)) - 6144*c^4*integrate(1/32*x^4*log(c*x - 1)/(c^10*d^3*x^6 - 3*c^8*d^3*x^4 + 3*c^6*d^3*x^2 - c^4*d^3), x) + 18*(c*(2*(5*c^2*x^2 + 3*c*x - 6)/(c^12*d^3*x^3 - c^11*d^3*x^2 - c^10*d^3*x + c^9*d^3) - 5*log(c*x + 1)/(c^9*d^3) + 5*log(c*x - 1)/(c^9*d^3)) + 16*(2*c^2*x^2 - 1)*log(c*x - 1)/(c^12*d^3*x^4 - 2*c^10*d^3*x^2 + c^8*d^3))*c^3 - 48*c^2*(2*(c^2*x^3 + x)/(c^10*d^3*x^4 - 2*c^8*d^3*x^2 + c^6*d^3) - log(c*x + 1)/(c^7*d^3) + log(c*x - 1)/(c^7*d^3)) + 12288*c^2*integrate(1/32*x^2*log(c*x - 1)/(c^10*d^3*x^6 - 3*c^8*d^3*x^4 + 3*c^6*d^3*x^2 - c^4*d^3), x) + 9*(c*(2*(3*c^2*x^2 - 3*c*x - 2)/(c^10*d^3*x^3 - c^9*d^3*x^2 - c^8*d^3*x + c^7*d^3) - 3*log(c*x + 1)/(c^7*d^3) + 3*log(c*x - 1)/(c^7*d^3)) - 16*log(c*x - 1)/(c^10*d^3*x^4 - 2*c^8*d^3*x^2 + c^6*d^3))*c - 32*(3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1)^2 + 6*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1)*log(c*x - 1) - 4*(10*c^3*x^3 - 6*c*x + 3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1) - 3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x - 1))*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(c^9*d^3*x^4 - 2*c^7*d^3*x^2 + c^5*d^3) + 2048*integrate(1/16*(10*c^3*x^3 - 6*c*x + 3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1) - 3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x - 1))/(c^11*d^3*x^7 - 3*c^9*d^3*x^5 + 3*c^7*d^3*x^3 - c^5*d^3*x + (...`

## 3.46.8 Giac [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^4}{(c^2dx^2 - d)^3} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)*x^4/(c^2*d*x^2 - d)^3, x)`

**3.46.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^3} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3,x)`output `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3, x)`

**3.47**  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$

3.47.1 Optimal result . . . . . 556  
 3.47.2 Mathematica [A] (verified) . . . . . 556  
 3.47.3 Rubi [A] (verified) . . . . . 557  
 3.47.4 Maple [A] (verified) . . . . . 559  
 3.47.5 Fricas [A] (verification not implemented) . . . . . 560  
 3.47.6 Sympy [F] . . . . . 560  
 3.47.7 Maxima [F] . . . . . 561  
 3.47.8 Giac [F(-2)] . . . . . 561  
 3.47.9 Mupad [F(-1)] . . . . . 561

**3.47.1 Optimal result**

Integrand size = 25, antiderivative size = 136

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \frac{bx^3}{12cd^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} + \frac{b}{4c^4d^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{b\sqrt{-1 + cx}}{4c^4d^3\sqrt{1 + cx}} - \frac{\operatorname{arccosh}(cx)}{4c^4d^3} + \frac{x^4(a + \operatorname{arccosh}(cx))}{4d^3(1 - c^2x^2)^2}$$

output `1/12*b*x^3/c/d^3/(c*x-1)^(3/2)/(c*x+1)^(3/2)-1/4*b*arccosh(c*x)/c^4/d^3+1/4*x^4*(a+b*arccosh(c*x))/d^3/(-c^2*x^2+1)^2+1/4*b/c^4/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/4*b*(c*x-1)^(1/2)/c^4/d^3/(c*x+1)^(1/2)`

**3.47.2 Mathematica [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.61

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \frac{bcx\sqrt{-1 + cx}\sqrt{1 + cx}(-3 + 4c^2x^2) + a(-3 + 6c^2x^2) + 3b(-1 + 2c^2x^2) \operatorname{arccosh}(cx)}{12c^4d^3(-1 + c^2x^2)^2}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output  $(b*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(-3 + 4*c^2*x^2) + a*(-3 + 6*c^2*x^2) + 3*b*(-1 + 2*c^2*x^2)*\text{ArcCosh}[c*x])/(12*c^4*d^3*(-1 + c^2*x^2)^2)$

### 3.47.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$ , Rules used = {6332, 109, 27, 100, 27, 87, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

$$\downarrow 6332$$

$$\frac{x^4(a + \text{barccosh}(cx))}{4d^3(1 - c^2 x^2)^2} - \frac{bc \int \frac{x^4}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4d^3}$$

$$\downarrow 109$$

$$\frac{x^4(a + \text{barccosh}(cx))}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( -\frac{\int -\frac{3x^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{3c^2} - \frac{x^3}{3c^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

$$\downarrow 27$$

$$\frac{x^4(a + \text{barccosh}(cx))}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{\int \frac{x^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{c^2} - \frac{x^3}{3c^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

$$\downarrow 100$$

$$\frac{x^4(a + \text{barccosh}(cx))}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{\int \frac{c^2 x}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c^3} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} - \frac{x^3}{3c^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

$$\downarrow 27$$

$$\frac{x^4(a + \text{barccosh}(cx))}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{\int \frac{x}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c} - \frac{1}{c^3 \sqrt{cx-1} \sqrt{cx+1}} - \frac{x^3}{3c^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

---

3.47.  $\int \frac{x^3(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$

$$\frac{x^4(a + \operatorname{arccosh}(cx))}{4d^3(1 - c^2x^2)^2} - \frac{bc \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x^3}{3c^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

↓ 87

$$\frac{x^4(a + \operatorname{arccosh}(cx))}{4d^3(1 - c^2x^2)^2} - \frac{bc \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x^3}{3c^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

↓ 43

input `Int[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output `(x^4*(a + b*ArcCosh[c*x]))/(4*d^3*(1 - c^2*x^2)^2) - (b*c*(-1/3*x^3/(c^2*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (-1/(c^3*sqrt[-1 + c*x]*sqrt[1 + c*x])) + (-sqrt[-1 + c*x]/(c^2*sqrt[1 + c*x])) + ArcCosh[c*x]/c^2)/c)/(4*d^3)`

### 3.47.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 43 `Int[1/(sqrt[(a_) + (b_)*(x_)]*sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`

3.47.  $\int \frac{x^3(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx$

rule 100 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))n((e_.) + (f_.)*(x_))p, x_] := Simp[(b*c - a*d)2(c + d*x)(n + 1)((e + f*x)(p + 1)/(d2(d*e - c*f)*(n + 1))), x] - Simp[1/(d2(d*e - c*f)*(n + 1)) Int[(c + d*x)(n + 1)(e + f*x)pSimp[a2d2f*(n + p + 2) + b2*c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b2*d*(d*e - c*f)*(n + 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))`

rule 109 `Int[((a_.) + (b_.)*(x_))m((c_.) + (d_.)*(x_))n((e_.) + (f_.)*(x_))p, x_] := Simp[(b*c - a*d)*(a + b*x)(m + 1)(c + d*x)(n - 1)((e + f*x)(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)(m + 1)(c + d*x)(n - 2)(e + f*x)pSimp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))n((f_.)*(x_))m((d_.) + (e_.)*(x_)2)p, x_Symbol] := Simp[(f*x)(m + 1)(d + e*x2)(p + 1)((a + b*ArcCosh[c*x])n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x2)p/((1 + c*x)p(-1 + c*x)p) Int[(f*x)(m + 1)(1 + c*x)(p + 1/2)(-1 + c*x)(p + 1/2)(a + b*ArcCosh[c*x])(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

### 3.47.4 Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.00

method	result
derivativedivides	$\frac{a \left( -\frac{1}{16(cx+1)^2} + \frac{3}{16(cx+1)} - \frac{1}{16(cx-1)^2} - \frac{3}{16(cx-1)} \right)}{d^3} - \frac{b \left( -\frac{\operatorname{arccosh}(cx)}{16(cx+1)^2} + \frac{3 \operatorname{arccosh}(cx)}{16(cx+1)} - \frac{\operatorname{arccosh}(cx)}{16(cx-1)^2} - \frac{3 \operatorname{arccosh}(cx)}{16(cx-1)} - \frac{cx}{12(cx+1)} \right)}{c^4 d^3}$
default	$\frac{a \left( -\frac{1}{16(cx+1)^2} + \frac{3}{16(cx+1)} - \frac{1}{16(cx-1)^2} - \frac{3}{16(cx-1)} \right)}{d^3} - \frac{b \left( -\frac{\operatorname{arccosh}(cx)}{16(cx+1)^2} + \frac{3 \operatorname{arccosh}(cx)}{16(cx+1)} - \frac{\operatorname{arccosh}(cx)}{16(cx-1)^2} - \frac{3 \operatorname{arccosh}(cx)}{16(cx-1)} - \frac{cx}{12(cx+1)} \right)}{c^4 d^3}$
parts	$-\frac{a \left( -\frac{1}{16c^4(cx+1)^2} + \frac{3}{16c^4(cx+1)} - \frac{1}{16c^4(cx-1)^2} - \frac{3}{16c^4(cx-1)} \right)}{d^3} - \frac{b \left( -\frac{\operatorname{arccosh}(cx)}{16(cx+1)^2} + \frac{3 \operatorname{arccosh}(cx)}{16(cx+1)} - \frac{\operatorname{arccosh}(cx)}{16(cx-1)^2} - \frac{3 \operatorname{arccosh}(cx)}{16(cx-1)} - \frac{cx}{12(cx+1)} \right)}{d^3 c^4}$

3.47.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$



input `int(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)`

output  $\frac{1}{c^4}(-a/d^3(-1/16/(c*x+1)^2+3/16/(c*x+1)-1/16/(c*x-1)^2-3/16/(c*x-1))-b/d^3(-1/16*\operatorname{arccosh}(c*x)/(c*x+1)^2+3/16*\operatorname{arccosh}(c*x)/(c*x+1)-1/16*\operatorname{arccosh}(c*x)/(c*x-1)^2-3/16*\operatorname{arccosh}(c*x)/(c*x-1)-1/12*c*x*(4*c^2*x^2-3)/(c*x+1)^(3/2)/(c*x-1)^(3/2)))$

### 3.47.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.74

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \frac{3ac^4x^4 + 3(2bc^2x^2 - b)\log(cx + \sqrt{c^2x^2 - 1}) + (4bc^3x^3 - 3bcx)\sqrt{c^2x^2 - 1}}{12(c^8d^3x^4 - 2c^6d^3x^2 + c^4d^3)}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`

output  $\frac{1}{12}*(3*a*c^4*x^4 + 3*(2*b*c^2*x^2 - b)*\log(c*x + \sqrt{c^2*x^2 - 1}) + (4*b*c^3*x^3 - 3*b*c*x)*\sqrt{c^2*x^2 - 1})/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3)$

### 3.47.6 Sympy [F]

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = -\int \frac{ax^3}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1} dx + \int \frac{bx^3 \operatorname{acosh}(cx)}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**3,x)`

output  $-(\operatorname{Integral}(a*x**3/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x) + \operatorname{Integral}(b*x**3*\operatorname{acosh}(c*x)/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x))/d**3$

**3.47.7 Maxima [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^3}{(c^2 dx^2 - d)^3} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

output `1/16*b*((4*c^2*x^2 + 4*(2*c^2*x^2 - 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) - 3)/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3) + 16*integrate(1/4*(2*c^2*x^2 - 1)/(c^10*d^3*x^7 - 3*c^8*d^3*x^5 + 3*c^6*d^3*x^3 - c^4*d^3*x + (c^9*d^3*x^6 - 3*c^7*d^3*x^4 + 3*c^5*d^3*x^2 - c^3*d^3)*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1))), x)) + 1/4*(2*c^2*x^2 - 1)*a/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3)`

**3.47.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.47.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^3} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3,x)`

output `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3, x)`

---

3.47.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$

**3.48**  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$

3.48.1	Optimal result . . . . .	562
3.48.2	Mathematica [A] (warning: unable to verify) . . . . .	563
3.48.3	Rubi [C] (verified) . . . . .	563
3.48.4	Maple [A] (verified) . . . . .	567
3.48.5	Fricas [F] . . . . .	568
3.48.6	Sympy [F] . . . . .	568
3.48.7	Maxima [F] . . . . .	568
3.48.8	Giac [F] . . . . .	569
3.48.9	Mupad [F(-1)] . . . . .	570

**3.48.1 Optimal result**

Integrand size = 25, antiderivative size = 186

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \frac{b}{12c^3d^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} + \frac{b}{8c^3d^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{x(a + \operatorname{arccosh}(cx))}{4c^2d^3(1 - c^2x^2)^2} - \frac{x(a + \operatorname{arccosh}(cx))}{8c^2d^3(1 - c^2x^2)}$$

$$- \frac{(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{4c^3d^3}$$

$$- \frac{b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{8c^3d^3} + \frac{b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{8c^3d^3}$$

output

```
1/12*b/c^3/d^3/(c*x-1)^(3/2)/(c*x+1)^(3/2)+1/4*x*(a+b*arccosh(c*x))/c^2/d^3/(-c^2*x^2+1)^2-1/8*x*(a+b*arccosh(c*x))/c^2/d^3/(-c^2*x^2+1)-1/4*(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d^3-1/8*b*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d^3+1/8*b*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c^3/d^3+1/8*b/c^3/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.48.2 Mathematica [A] (warning: unable to verify)

Time = 1.15 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.54

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

$$= \frac{-\frac{b(-2+cx)\sqrt{1+cx}}{(-1+cx)^{3/2}} + \frac{b\sqrt{-1+cx}(2+cx)}{(1+cx)^{3/2}} + \frac{12acx}{(-1+c^2x^2)^2} + \frac{6acx}{-1+c^2x^2} + \frac{3b \operatorname{arccosh}(cx)}{(-1+cx)^2} - \frac{3b \operatorname{arccosh}(cx)}{(1+cx)^2} - 3b \left( -\frac{1}{\sqrt{\frac{-1+cx}{1+cx}}} + a \right)}{48c^3d^3}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output `(-((b*(-2 + c*x)*Sqrt[1 + c*x])/(-1 + c*x)^(3/2)) + (b*Sqrt[-1 + c*x]*(2 + c*x))/(1 + c*x)^(3/2) + (12*a*c*x)/(-1 + c^2*x^2)^2 + (6*a*c*x)/(-1 + c^2*x^2) + (3*b*ArcCosh[c*x])/(-1 + c*x)^2 - (3*b*ArcCosh[c*x])/(1 + c*x)^2 - 3*b*(-(1/Sqrt[(-1 + c*x)/(1 + c*x)]) + ArcCosh[c*x]/(1 - c*x)) - 3*b*(Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x]/(1 + c*x)) - (3*b*ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 - E^ArcCosh[c*x]]))/2 + (3*b*ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 + E^ArcCosh[c*x]]))/2 + 3*a*Log[1 - c*x] - 3*a*Log[1 + c*x] - 6*b*PolyLog[2, -E^ArcCosh[c*x]] + 6*b*PolyLog[2, E^ArcCosh[c*x]])/(48*c^3*d^3)`

### 3.48.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.80 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.97, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$ , Rules used = {6349, 27, 83, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

$$\downarrow \text{6349}$$

$$-\frac{\int \frac{a+b \operatorname{arccosh}(cx)}{d^2(1-c^2x^2)^2} dx}{4c^2d} - \frac{b \int \frac{x}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4cd^3} + \frac{x(a + b \operatorname{arccosh}(cx))}{4c^2d^3(1 - c^2x^2)^2}$$

$$\downarrow \text{27}$$

---

3.48.  $\int \frac{x^2(a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^3} dx$

$$\begin{aligned}
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{4c^2d^3} - \frac{b \int \frac{x}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4cd^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} \\
& \quad \downarrow \text{83} \\
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{4c^2d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} + \frac{b}{12c^3d^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \quad \downarrow \text{6316} \\
& -\frac{\frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)}}{4c^2d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} + \\
& \quad \frac{b}{12c^3d^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \quad \downarrow \text{83} \\
& -\frac{\frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}}{4c^2d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} + \\
& \quad \frac{b}{12c^3d^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \quad \downarrow \text{6318} \\
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} + \frac{x(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} + \\
& \quad \frac{b}{12c^3d^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& -\frac{\int \frac{i(a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}}{4c^2d^3} + \\
& \quad \frac{x(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} + \frac{b}{12c^3d^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \quad \downarrow \text{26} \\
& -\frac{i \int \frac{(a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}}{4c^2d^3} + \\
& \quad \frac{x(a+\operatorname{barccosh}(cx))}{4c^2d^3(1-c^2x^2)^2} + \frac{b}{12c^3d^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \quad \downarrow \text{4670}
\end{aligned}$$

---

3.48.  $\int \frac{x^2(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^3} dx$

$$\begin{aligned}
 & - \frac{i \left( ib \int \log \left( 1 - e^{\operatorname{arccosh}(cx)} \right) d \operatorname{arccosh}(cx) - ib \int \log \left( 1 + e^{\operatorname{arccosh}(cx)} \right) d \operatorname{arccosh}(cx) + 2i \operatorname{arctanh} \left( e^{\operatorname{arccosh}(cx)} \right) (a + \operatorname{barccosh}(cx)) \right)}{2c} + \dots \\
 & \frac{4c^2 d^3}{4c^2 d^3 (1 - c^2 x^2)^2} + \frac{b}{12c^3 d^3 (cx - 1)^{3/2} (cx + 1)^{3/2}} \\
 & \quad \downarrow \text{2715} \\
 & - \frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log \left( 1 - e^{\operatorname{arccosh}(cx)} \right) d e^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log \left( 1 + e^{\operatorname{arccosh}(cx)} \right) d e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh} \left( e^{\operatorname{arccosh}(cx)} \right) \right)}{2c} \\
 & \frac{4c^2 d^3}{4c^2 d^3 (1 - c^2 x^2)^2} + \frac{b}{12c^3 d^3 (cx - 1)^{3/2} (cx + 1)^{3/2}} \\
 & \quad \downarrow \text{2838} \\
 & - \frac{i \left( 2i \operatorname{arctanh} \left( e^{\operatorname{arccosh}(cx)} \right) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog} \left( 2, -e^{\operatorname{arccosh}(cx)} \right) - ib \operatorname{PolyLog} \left( 2, e^{\operatorname{arccosh}(cx)} \right) \right)}{2c} + \frac{x(a + \operatorname{barccosh}(cx))}{2(1 - c^2 x^2)} \\
 & \frac{4c^2 d^3}{4c^2 d^3 (1 - c^2 x^2)^2} + \frac{b}{12c^3 d^3 (cx - 1)^{3/2} (cx + 1)^{3/2}}
 \end{aligned}$$

input `Int[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output `b/(12*c^3*d^3*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (x*(a + b*ArcCosh[c*x]))/(4*c^2*d^3*(1 - c^2*x^2)^2) - (-1/2*b/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c)/(4*c^2*d^3)`

### 3.48.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

---

3.48.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$

- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`
- rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.48.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.38

method	result
derivativedivides	$\frac{a \left( \frac{1}{16(cx+1)^2} - \frac{1}{16(cx+1)} + \frac{\ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} - \frac{1}{16(cx-1)} - \frac{\ln(cx-1)}{16} \right)}{d^3} - \frac{b \left( -\frac{3c^3x^3 \operatorname{arccosh}(cx) + 3\sqrt{cx-1}\sqrt{cx+1}c^2x^2 + 3cx}{24(c^4x^4 - 2c^2x^2 + 1)} \right)}{d^3}$
default	$\frac{a \left( \frac{1}{16(cx+1)^2} - \frac{1}{16(cx+1)} + \frac{\ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} - \frac{1}{16(cx-1)} - \frac{\ln(cx-1)}{16} \right)}{d^3} - \frac{b \left( -\frac{3c^3x^3 \operatorname{arccosh}(cx) + 3\sqrt{cx-1}\sqrt{cx+1}c^2x^2 + 3cx}{24(c^4x^4 - 2c^2x^2 + 1)} \right)}{d^3}$
parts	$-\frac{a \left( \frac{1}{16c^3(cx+1)^2} - \frac{1}{16c^3(cx+1)} + \frac{\ln(cx+1)}{16c^3} - \frac{1}{16c^3(cx-1)^2} - \frac{1}{16c^3(cx-1)} - \frac{\ln(cx-1)}{16c^3} \right)}{d^3} - \frac{b \left( -\frac{3c^3x^3 \operatorname{arccosh}(cx) + 3\sqrt{cx-1}\sqrt{cx+1}c^2x^2 + 3cx}{24(c^4x^4 - 2c^2x^2 + 1)} \right)}{d^3}$

```
input int(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)
```

```
output 1/c^3*(-a/d^3*(1/16/(c*x+1)^2-1/16/(c*x+1)+1/16*ln(c*x+1)-1/16/(c*x-1)^2-1/16/(c*x-1)-1/16*ln(c*x-1))-b/d^3*(-1/24*(3*c^3*x^3*arccosh(c*x)+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+3*c*x*arccosh(c*x)-(c*x-1)^(1/2)*(c*x+1)^(1/2)))/(c^4*x^4-2*c^2*x^2+1)-1/8*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-1/8*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/8*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/8*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))))
```

$$3.48. \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$$



## 3.48.5 Fricas [F]

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^2}{(c^2dx^2 - d)^3} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`

output `integral(-(b*x^2*arccosh(c*x) + a*x^2)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

## 3.48.6 Sympy [F]

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = -\int \frac{ax^2}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1} dx + \int \frac{bx^2 \operatorname{acosh}(cx)}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**3,x)`

output `-(Integral(a*x**2/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x) + Integral(b*x**2*acosh(c*x)/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x))/d**3`

## 3.48.7 Maxima [F]

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x^2}{(c^2dx^2 - d)^3} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

```

output -1/2048*(6144*c^5*integrate(1/32*x^5*log(c*x - 1)/(c^8*d^3*x^6 - 3*c^6*d^3
*x^4 + 3*c^4*d^3*x^2 - c^2*d^3), x) - 16*c^4*(2*(5*c^2*x^3 - 3*x)/(c^10*d^
3*x^4 - 2*c^8*d^3*x^2 + c^6*d^3) + 3*log(c*x + 1)/(c^7*d^3) - 3*log(c*x -
1)/(c^7*d^3)) - 2048*c^4*integrate(1/32*x^4*log(c*x - 1)/(c^8*d^3*x^6 - 3*
c^6*d^3*x^4 + 3*c^4*d^3*x^2 - c^2*d^3), x) + 6*(c*(2*(5*c^2*x^2 + 3*c*x -
6)/(c^10*d^3*x^3 - c^9*d^3*x^2 - c^8*d^3*x + c^7*d^3) - 5*log(c*x + 1)/(c^
7*d^3) + 5*log(c*x - 1)/(c^7*d^3)) + 16*(2*c^2*x^2 - 1)*log(c*x - 1)/(c^10
*d^3*x^4 - 2*c^8*d^3*x^2 + c^6*d^3))*c^3 - 16*c^2*(2*(c^2*x^3 + x)/(c^8*d^
3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3) - log(c*x + 1)/(c^5*d^3) + log(c*x - 1)/(
c^5*d^3)) + 4096*c^2*integrate(1/32*x^2*log(c*x - 1)/(c^8*d^3*x^6 - 3*c^6*
d^3*x^4 + 3*c^4*d^3*x^2 - c^2*d^3), x) + 3*(c*(2*(3*c^2*x^2 - 3*c*x - 2)/(
c^8*d^3*x^3 - c^7*d^3*x^2 - c^6*d^3*x + c^5*d^3) - 3*log(c*x + 1)/(c^5*d^3
) + 3*log(c*x - 1)/(c^5*d^3)) - 16*log(c*x - 1)/(c^8*d^3*x^4 - 2*c^6*d^3*x
^2 + c^4*d^3))*c - 32*((c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1)^2 + 2*(c^4*x
^4 - 2*c^2*x^2 + 1)*log(c*x + 1)*log(c*x - 1) + 4*(2*c^3*x^3 + 2*c*x - (c^
4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1) + (c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x -
1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)))/(c^7*d^3*x^4 - 2*c^5*d^3*x^2 +
c^3*d^3) + 2048*integrate(-1/16*(2*c^3*x^3 + 2*c*x - (c^4*x^4 - 2*c^2*x^2
+ 1)*log(c*x + 1) + (c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x - 1))/(c^9*d^3*x^7
- 3*c^7*d^3*x^5 + 3*c^5*d^3*x^3 - c^3*d^3*x + (c^8*d^3*x^6 - 3*c^6*d^3*...

```

### 3.48.8 Giac [F]

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arccosh}(cx) + a)x^2}{(c^2 dx^2 - d)^3} dx$$

```

input integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="giac")

```

```

output integrate(-(b*arccosh(c*x) + a)*x^2/(c^2*d*x^2 - d)^3, x)

```

**3.48.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^3} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3,x)`output `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3, x)`

### 3.49 $\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$

3.49.1	Optimal result	571
3.49.2	Mathematica [A] (verified)	571
3.49.3	Rubi [A] (verified)	572
3.49.4	Maple [A] (verified)	573
3.49.5	Fricas [A] (verification not implemented)	574
3.49.6	Sympy [F]	574
3.49.7	Maxima [F]	574
3.49.8	Giac [F]	575
3.49.9	Mupad [F(-1)]	575

#### 3.49.1 Optimal result

Integrand size = 23, antiderivative size = 91

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \frac{bx}{12cd^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} - \frac{bx}{6cd^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{a + \operatorname{arccosh}(cx)}{4c^2d^3(1 - c^2x^2)^2}$$

output  $1/12*b*x/c/d^3/(c*x-1)^{(3/2)}/(c*x+1)^{(3/2)}+1/4*(a+b*\operatorname{arccosh}(c*x))/c^2/d^3/(-c^2*x^2+1)^2-1/6*b*x/c/d^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

#### 3.49.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.70

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^3} dx = \frac{3a + bcx\sqrt{-1 + cx}\sqrt{1 + cx}(3 - 2c^2x^2) + 3\operatorname{arccosh}(cx)}{12c^2d^3(-1 + c^2x^2)^2}$$

input `Integrate[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output  $(3*a + b*c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(3 - 2*c^2*x^2) + 3*b*\operatorname{ArcCosh}[c*x])/((12*c^2*d^3*(-1 + c^2*x^2)^2)$

### 3.49.3 Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$ , Rules used = {6329, 42, 41}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx \\
 & \quad \downarrow \text{6329} \\
 & \frac{a + \operatorname{barccosh}(cx)}{4c^2 d^3 (1 - c^2 x^2)^2} - \frac{b \int \frac{1}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4cd^3} \\
 & \quad \downarrow \text{42} \\
 & \frac{a + \operatorname{barccosh}(cx)}{4c^2 d^3 (1 - c^2 x^2)^2} - \frac{b \left( -\frac{2}{3} \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3} \\
 & \quad \downarrow \text{41} \\
 & \frac{a + \operatorname{barccosh}(cx)}{4c^2 d^3 (1 - c^2 x^2)^2} - \frac{b \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4cd^3}
 \end{aligned}$$

input `Int[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output `-1/4*(b*(-1/3*x/((-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (2*x)/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/(c*d^3) + (a + b*ArcCosh[c*x])/(4*c^2*d^3*(1 - c^2*x^2)^2)`

#### 3.49.3.1 Defintions of rubi rules used

rule 41 `Int[1/(((a_) + (b_.)*(x_))^(3/2)*((c_) + (d_.)*(x_))^(3/2)), x_Symbol] :> Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`

---

3.49.  $\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$

rule 42 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[(-x)*(a + b*x)^(m + 1)*((c + d*x)^(m + 1)/(2*a*c*(m + 1))), x] + Simp[(2*m + 3)/(2*a*c*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(m + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && ILtQ[m + 3/2, 0]`

rule 6329 `Int[((a_) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.49.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$\frac{\frac{a}{4d^3(c^2x^2-1)^2} - \frac{b\left(-\frac{\operatorname{arccosh}(cx)}{4(c^2x^2-1)^2} + \frac{cx(2c^2x^2-3)}{12\sqrt{cx-1}\sqrt{cx+1}(c^2x^2-1)}\right)}{d^3}}{c^2}$	86
default	$\frac{\frac{a}{4d^3(c^2x^2-1)^2} - \frac{b\left(-\frac{\operatorname{arccosh}(cx)}{4(c^2x^2-1)^2} + \frac{cx(2c^2x^2-3)}{12\sqrt{cx-1}\sqrt{cx+1}(c^2x^2-1)}\right)}{d^3}}{c^2}$	86
parts	$\frac{a}{4d^3c^2(c^2x^2-1)^2} - \frac{b\left(-\frac{\operatorname{arccosh}(cx)}{4(c^2x^2-1)^2} + \frac{cx(2c^2x^2-3)}{12\sqrt{cx-1}\sqrt{cx+1}(c^2x^2-1)}\right)}{d^3c^2}$	88

input `int(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)`

output  $\frac{1}{c^2} \left( \frac{1}{4} \frac{a}{d^3} (c^2x^2-1)^{-2} - \frac{b}{d^3} \left( -\frac{1}{4} (c^2x^2-1)^{-2} \operatorname{arccosh}(cx) + \frac{1}{12} \frac{(cx-1)^{1/2}}{(cx+1)^{1/2}} cx (2c^2x^2-3) (c^2x^2-1)^{-1} \right) \right)$

**3.49.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.08

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

$$= -\frac{3ac^4x^4 - 6ac^2x^2 - 3b \log(cx + \sqrt{c^2x^2 - 1}) + (2bc^3x^3 - 3bcx)\sqrt{c^2x^2 - 1}}{12(c^6d^3x^4 - 2c^4d^3x^2 + c^2d^3)}$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`output `-1/12*(3*a*c^4*x^4 - 6*a*c^2*x^2 - 3*b*log(c*x + sqrt(c^2*x^2 - 1)) + (2*b*c^3*x^3 - 3*b*c*x)*sqrt(c^2*x^2 - 1))/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3)`**3.49.6 Sympy [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = -\int \frac{ax}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1} dx + \int \frac{bx \operatorname{acosh}(cx)}{c^6x^6 - 3c^4x^4 + 3c^2x^2 - 1} dx$$

input `integrate(x*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**3,x)`output `-(Integral(a*x/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x) + Integral(b*x*acosh(c*x)/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x))/d**3`**3.49.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x}{(c^2 dx^2 - d)^3} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`output `1/16*b*((4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) + 1)/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3) + 16*integrate(1/4/(c^8*d^3*x^7 - 3*c^6*d^3*x^5 + 3*c^4*d^3*x^3 - c^2*d^3*x + (c^7*d^3*x^6 - 3*c^5*d^3*x^4 + 3*c^3*d^3*x^2 - c*d^3)*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1))), x) + 1/4*a/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3)`

**3.49.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)x}{(c^2 dx^2 - d)^3} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)*x/(c^2*d*x^2 - d)^3, x)`

**3.49.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^3} dx$$

input `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3,x)`

output `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^3, x)`



### 3.50 $\int \frac{a+b\operatorname{arccosh}(cx)}{(d-c^2dx^2)^3} dx$

3.50.1	Optimal result	576
3.50.2	Mathematica [A] (warning: unable to verify)	577
3.50.3	Rubi [C] (verified)	577
3.50.4	Maple [A] (verified)	581
3.50.5	Fricas [F]	581
3.50.6	Sympy [F]	582
3.50.7	Maxima [F]	582
3.50.8	Giac [F]	583
3.50.9	Mupad [F(-1)]	583

#### 3.50.1 Optimal result

Integrand size = 22, antiderivative size = 180

$$\int \frac{a + b\operatorname{arccosh}(cx)}{(d - c^2dx^2)^3} dx = \frac{b}{12cd^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} - \frac{3b}{8cd^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{x(a + b\operatorname{arccosh}(cx))}{4d^3(1 - c^2x^2)^2} + \frac{3x(a + b\operatorname{arccosh}(cx))}{8d^3(1 - c^2x^2)}$$

$$+ \frac{3(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{4cd^3}$$

$$+ \frac{3b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{8cd^3} - \frac{3b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{8cd^3}$$

output `1/12*b/c/d^3/(c*x-1)^(3/2)/(c*x+1)^(3/2)+1/4*x*(a+b*arccosh(c*x))/d^3/(-c^2*x^2+1)^2+3/8*x*(a+b*arccosh(c*x))/d^3/(-c^2*x^2+1)+3/4*(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c/d^3+3/8*b*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/c/d^3-3/8*b*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/c/d^3-3/8*b/c/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)`

### 3.50.2 Mathematica [A] (warning: unable to verify)

Time = 0.80 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.76

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^3} dx$$

$$= \frac{4ax}{(-1+c^2x^2)^2} - \frac{6ax}{-1+c^2x^2} + \frac{b(\sqrt{-1+cx}\sqrt{1+cx}(2+cx)-3\operatorname{arccosh}(cx))}{3c(1+cx)^2} + \frac{b((2-cx)\sqrt{-1+cx}\sqrt{1+cx}+3\operatorname{arccosh}(cx))}{3c(-1+cx)^2} + \frac{3b}{\sqrt{\frac{-1+cx}{1+cx}}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^3,x]`

output `((4*a*x)/(-1 + c^2*x^2)^2 - (6*a*x)/(-1 + c^2*x^2) + (b*(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2 + c*x) - 3*ArcCosh[c*x]))/(3*c*(1 + c*x)^2) + (b*((2 - c*x)*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 3*ArcCosh[c*x]))/(3*c*(-1 + c*x)^2) + (3*b*(-1/Sqrt[(-1 + c*x)/(1 + c*x)]) + ArcCosh[c*x]/(1 - c*x))/c + (3*b*(Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x]/(1 + c*x))/c - (3*a*Log[1 - c*x])/c + (3*a*Log[1 + c*x])/c - (3*b*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 + E^ArcCosh[c*x]]) - 4*PolyLog[2, -E^ArcCosh[c*x]]))/(2*c) + (3*b*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 - E^ArcCosh[c*x]]) - 4*PolyLog[2, E^ArcCosh[c*x]]))/(2*c))/(16*d^3)`

### 3.50.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.72 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.97, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6316, 27, 83, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^3} dx$$

$$\downarrow \text{6316}$$

$$\frac{3 \int \frac{a + b \operatorname{arccosh}(cx)}{d^2(1 - c^2 x^2)^2} dx}{4d} - \frac{bc \int \frac{x}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4d^3} + \frac{x(a + b \operatorname{arccosh}(cx))}{4d^3(1 - c^2 x^2)^2}$$

---

3.50.  $\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^3} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{3 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{4d^3} - \frac{bc \int \frac{x}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4d^3(1-c^2x^2)^2} \\
& \downarrow 83 \\
& \frac{3 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{4d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4d^3(1-c^2x^2)^2} + \frac{b}{12cd^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \downarrow 6316 \\
& \frac{3\left(\frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)}\right)}{4d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4d^3(1-c^2x^2)^2} + \\
& \quad \frac{b}{12cd^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \downarrow 83 \\
& \frac{3\left(\frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{4d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4d^3(1-c^2x^2)^2} + \\
& \quad \frac{b}{12cd^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \downarrow 6318 \\
& \frac{3\left(\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{4d^3} + \frac{x(a+\operatorname{barccosh}(cx))}{4d^3(1-c^2x^2)^2} + \\
& \quad \frac{b}{12cd^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \downarrow 3042 \\
& \frac{3\left(-\frac{\int i(a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{4d^3} + \\
& \quad \frac{x(a+\operatorname{barccosh}(cx))}{4d^3(1-c^2x^2)^2} + \frac{b}{12cd^3(cx-1)^{3/2}(cx+1)^{3/2}} \\
& \downarrow 26 \\
& \frac{3\left(-\frac{i \int (a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{4d^3} + \\
& \quad \frac{x(a+\operatorname{barccosh}(cx))}{4d^3(1-c^2x^2)^2} + \frac{b}{12cd^3(cx-1)^{3/2}(cx+1)^{3/2}}
\end{aligned}$$

---

3.50.  $\int \frac{a+\operatorname{barccosh}(cx)}{(d-c^2dx^2)^3} dx$

↓ 4670

$$3 \left( -\frac{i \int \log(1 - e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - i b \int \log(1 + e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx))}{2c} \right) + \frac{4d^3}{4d^3 (1 - c^2 x^2)^2} + \frac{b}{12cd^3 (cx - 1)^{3/2} (cx + 1)^{3/2}}$$

↓ 2715

$$3 \left( -\frac{i \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) d e^{\operatorname{arccosh}(cx)} - i b \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) d e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{2c} \right) + \frac{4d^3}{4d^3 (1 - c^2 x^2)^2} + \frac{b}{12cd^3 (cx - 1)^{3/2} (cx + 1)^{3/2}}$$

↓ 2838

$$3 \left( -\frac{i (2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)) + i b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - i b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{2c} \right) + \frac{x(a + b \operatorname{arccosh}(cx))}{2(1 - c^2 x^2)} + \frac{4d^3}{4d^3 (1 - c^2 x^2)^2} + \frac{b}{12cd^3 (cx - 1)^{3/2} (cx + 1)^{3/2}}$$

input `Int[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^3,x]`

output `b/(12*c*d^3*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (x*(a + b*ArcCosh[c*x]))/(4*d^3*(1 - c^2*x^2)^2) + (3*(-1/2*b/(c*sqrt[-1 + c*x]*sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*ArcCosh[c*x]))*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c)/(4*d^3)`

## 3.50.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/(f*fz*I)], x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

```
rule 6318 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol]
  := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]
```

### 3.50.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.42

method	result
derivativedivides	$-\frac{a\left(\frac{1}{16(cx+1)^2} + \frac{3}{16(cx+1)} - \frac{3\ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} + \frac{3}{16(cx-1)} + \frac{3\ln(cx-1)}{16}\right)}{d^3} - \frac{b\left(\frac{9c^3x^3 \operatorname{arccosh}(cx) + 9\sqrt{cx-1}\sqrt{cx+1}c^2x^2 - 15cx}{24c^4x^4 - 48c^2x^2 + 24}\right)}{d^3}$
default	$-\frac{a\left(\frac{1}{16(cx+1)^2} + \frac{3}{16(cx+1)} - \frac{3\ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} + \frac{3}{16(cx-1)} + \frac{3\ln(cx-1)}{16}\right)}{d^3} - \frac{b\left(\frac{9c^3x^3 \operatorname{arccosh}(cx) + 9\sqrt{cx-1}\sqrt{cx+1}c^2x^2 - 15cx}{24c^4x^4 - 48c^2x^2 + 24}\right)}{d^3}$
parts	$-\frac{a\left(\frac{1}{16c(cx+1)^2} + \frac{3}{16c(cx+1)} - \frac{3\ln(cx+1)}{16c} - \frac{1}{16c(cx-1)^2} + \frac{3}{16c(cx-1)} + \frac{3\ln(cx-1)}{16c}\right)}{d^3} - \frac{b\left(\frac{9c^3x^3 \operatorname{arccosh}(cx) + 9\sqrt{cx-1}\sqrt{cx+1}c^2x^2 - 15cx}{24c^4x^4 - 48c^2x^2 + 24}\right)}{d^3}$

```
input int((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)
```

```
output 1/c*(-a/d^3*(1/16/(c*x+1)^2+3/16/(c*x+1)-3/16*ln(c*x+1)-1/16/(c*x-1)^2+3/16/(c*x-1)+3/16*ln(c*x-1))-b/d^3*(1/24*(9*c^3*x^3*arccosh(c*x)+9*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-15*c*x*arccosh(c*x)-11*(c*x-1)^(1/2)*(c*x+1)^(1/2)))/(c^4*x^4-2*c^2*x^2+1)+3/8*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+3/8*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-3/8*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-3/8*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))))
```

### 3.50.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^3} dx$$

```
input integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="fricas")
```

```
output integral(-(b*arccosh(c*x) + a)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)
```

---

3.50.  $\int \frac{a+b \operatorname{arccosh}(cx)}{(d-c^2 dx^2)^3} dx$

## 3.50.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^3} dx = -\int \frac{\frac{a}{c^6 x^6 - 3c^4 x^4 + 3c^2 x^2 - 1} dx}{d^3} + \int \frac{\frac{b \operatorname{arcosh}(cx)}{c^6 x^6 - 3c^4 x^4 + 3c^2 x^2 - 1} dx}{d^3}$$

input `integrate((a+b*acosh(c*x))/(-c**2*d*x**2+d)**3,x)`

output `-(Integral(a/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x) + Integral(b*acosh(c*x)/(c**6*x**6 - 3*c**4*x**4 + 3*c**2*x**2 - 1), x))/d**3`

## 3.50.7 Maxima [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

output `1/2048*(18432*c^5*integrate(1/32*x^5*log(c*x - 1)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x) - 48*c^4*(2*(5*c^2*x^3 - 3*x)/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3) + 3*log(c*x + 1)/(c^5*d^3) - 3*log(c*x - 1)/(c^5*d^3)) - 6144*c^4*integrate(1/32*x^4*log(c*x - 1)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x) + 18*(c*(2*(5*c^2*x^2 + 3*c*x - 6)/(c^8*d^3*x^3 - c^7*d^3*x^2 - c^6*d^3*x + c^5*d^3) - 5*log(c*x + 1)/(c^5*d^3) + 5*log(c*x - 1)/(c^5*d^3)) + 16*(2*c^2*x^2 - 1)*log(c*x - 1)/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3))*c^3 + 80*c^2*(2*(c^2*x^3 + x)/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3) - log(c*x + 1)/(c^3*d^3) + log(c*x - 1)/(c^3*d^3)) + 12288*c^2*integrate(1/32*x^2*log(c*x - 1)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x) + 9*(c*(2*(3*c^2*x^2 - 3*c*x - 2)/(c^6*d^3*x^3 - c^5*d^3*x^2 - c^4*d^3*x + c^3*d^3) - 3*log(c*x + 1)/(c^3*d^3) + 3*log(c*x - 1)/(c^3*d^3)) - 16*log(c*x - 1)/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3))*c - 32*(3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1)^2 + 6*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x + 1)*log(c*x - 1) + 4*(6*c^3*x^3 - 10*c*x - 3*(c^4*x^4 - 2*c^2*x^2 + 1))*log(c*x + 1) + 3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x - 1))*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(c^5*d^3*x^4 - 2*c^3*d^3*x^2 + c*d^3) + 2048*integrate(-1/16*(6*c^3*x^3 - 10*c*x - 3*(c^4*x^4 - 2*c^2*x^2 + 1))*log(c*x + 1) + 3*(c^4*x^4 - 2*c^2*x^2 + 1)*log(c*x - 1))/(c^7*d^3*x^7 - 3*c^5*d^3*x^5 + 3*c^3*d^3*x^3 - c*d^3*x + (c^6*d^3*x^6 - 3*c^4*d^3*x^4 + ...`

3.50.  $\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^3} dx$

**3.50.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/(c^2*d*x^2 - d)^3, x)`

**3.50.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(d - c^2 dx^2)^3} dx$$

input `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^3,x)`

output `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^3, x)`



### 3.51 $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)^3} dx$

3.51.1	Optimal result	584
3.51.2	Mathematica [A] (warning: unable to verify)	585
3.51.3	Rubi [C] (verified)	585
3.51.4	Maple [A] (verified)	589
3.51.5	Fricas [F]	590
3.51.6	Sympy [F]	590
3.51.7	Maxima [F]	591
3.51.8	Giac [F]	591
3.51.9	Mupad [F(-1)]	591

#### 3.51.1 Optimal result

Integrand size = 25, antiderivative size = 171

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x(d - c^2dx^2)^3} dx = \frac{bcx}{12d^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} - \frac{2bcx}{3d^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{a + b\operatorname{arccosh}(cx)}{4d^3(1 - c^2x^2)^2} + \frac{a + b\operatorname{arccosh}(cx)}{2d^3(1 - c^2x^2)}$$

$$+ \frac{2(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d^3}$$

$$+ \frac{b \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{2d^3} - \frac{b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2d^3}$$

output  $1/12*b*c*x/d^3/(c*x-1)^{(3/2)}/(c*x+1)^{(3/2)}+1/4*(a+b*\operatorname{arccosh}(c*x))/d^3/(-c^2*x^2+1)^2+1/2*(a+b*\operatorname{arccosh}(c*x))/d^3/(-c^2*x^2+1)+2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}((c*x+(c*x-1)^{(1/2})*(c*x+1)^{(1/2}))^2)/d^3+1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2})*(c*x+1)^{(1/2}))^2)/d^3-1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2})*(c*x+1)^{(1/2}))^2)/d^3-2/3*b*c*x/d^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.51.2 Mathematica [A] (warning: unable to verify)

Time = 0.97 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.92

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx$$

$$= \frac{a}{(-1+c^2x^2)^2} - \frac{2a}{-1+c^2x^2} - \frac{b(\sqrt{-1+cx}\sqrt{1+cx}(2+cx)-3\operatorname{arccosh}(cx))}{12(1+cx)^2} + \frac{b((2-cx)\sqrt{-1+cx}\sqrt{1+cx}+3\operatorname{arccosh}(cx))}{12(-1+cx)^2} + \frac{5}{4}b \left( -\frac{1}{\sqrt{-1+cx}} \right)$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^3), x]`

output `(a/(-1 + c^2*x^2)^2 - (2*a)/(-1 + c^2*x^2) - (b*(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2 + c*x) - 3*ArcCosh[c*x]))/(12*(1 + c*x)^2) + (b*((2 - c*x)*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 3*ArcCosh[c*x]))/(12*(-1 + c*x)^2) + (5*b*(-(1/Sqrt[(-1 + c*x)/(1 + c*x)]) + ArcCosh[c*x]/(1 - c*x)))/4 - (5*b*(Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x]/(1 + c*x)))/4 + 4*a*Log[x] - 2*a*Log[1 - c^2*x^2] + 2*b*(ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLog[2, -E^(-2*ArcCosh[c*x])]) + b*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 + E^ArcCosh[c*x]]) - 4*PolyLog[2, -E^ArcCosh[c*x]]) + b*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 - E^ArcCosh[c*x]]) - 4*PolyLog[2, E^ArcCosh[c*x]])))/(4*d^3)`

### 3.51.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.17, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.520$ , Rules used = {6351, 27, 42, 41, 6351, 41, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx$$

$$\downarrow \text{6351}$$

$$\frac{\int \frac{a + b \operatorname{arccosh}(cx)}{d^2 x(1 - c^2 x^2)^2} dx}{d} - \frac{bc \int \frac{1}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4d^3} + \frac{a + b \operatorname{arccosh}(cx)}{4d^3(1 - c^2 x^2)^2}$$

---

3.51.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^3} - \frac{bc \int \frac{1}{(cx-1)^{5/2}(cx+1)^{5/2}} dx}{4d^3} + \frac{a + \operatorname{barccosh}(cx)}{4d^3(1-c^2x^2)^2} \\
& \downarrow 42 \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^3} - \frac{bc \left( -\frac{2}{3} \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3} + \frac{a + \operatorname{barccosh}(cx)}{4d^3(1-c^2x^2)^2} \\
& \downarrow 41 \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{4d^3(1-c^2x^2)^2} - \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3} \\
& \downarrow 6351 \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2} bc \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)}}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{4d^3(1-c^2x^2)^2} - \\
& \quad \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3} \\
& \downarrow 41 \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{a + \operatorname{barccosh}(cx)}{4d^3(1-c^2x^2)^2} - \\
& \quad \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}}{d^3} \\
& \downarrow 6331 \\
& \frac{- \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{4d^3(1-c^2x^2)^2} - \\
& \quad \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3} \\
& \downarrow 5984 \\
& \frac{-2 \int (a + \operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}}{d^3} + \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{4d^3(1-c^2x^2)^2} - \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3} \\
& \downarrow 3042
\end{aligned}$$

---

3.51.  $\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^3} dx$

$$\frac{-2 \int i(a + \operatorname{barccosh}(cx)) \csc(2i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a + \operatorname{barccosh}(cx)}{2(1 - c^2 x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}}{d^3} +$$

$$\frac{a + \operatorname{barccosh}(cx)}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

↓ 26

$$\frac{-2i \int (a + \operatorname{barccosh}(cx)) \csc(2i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a + \operatorname{barccosh}(cx)}{2(1 - c^2 x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}}{d^3} +$$

$$\frac{a + \operatorname{barccosh}(cx)}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

↓ 4670

$$\frac{-2i \left( \frac{1}{2} ib \int \log(1 - e^{2 \operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} ib \int \log(1 + e^{2 \operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + i \operatorname{arctanh}(e^{2 \operatorname{arccosh}(cx)}) \right)}{d^3} +$$

$$\frac{a + \operatorname{barccosh}(cx)}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

↓ 2715

$$\frac{-2i \left( \frac{1}{4} ib \int e^{-2 \operatorname{arccosh}(cx)} \log(1 - e^{2 \operatorname{arccosh}(cx)}) de^{2 \operatorname{arccosh}(cx)} - \frac{1}{4} ib \int e^{-2 \operatorname{arccosh}(cx)} \log(1 + e^{2 \operatorname{arccosh}(cx)}) de^{2 \operatorname{arccosh}(cx)} \right)}{d^3} +$$

$$\frac{a + \operatorname{barccosh}(cx)}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

↓ 2838

$$\frac{-2i \left( i \operatorname{arctanh}(e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} ib \operatorname{PolyLog}(2, -e^{2 \operatorname{arccosh}(cx)}) - \frac{1}{4} ib \operatorname{PolyLog}(2, e^{2 \operatorname{arccosh}(cx)}) \right)}{d^3} +$$

$$\frac{a + \operatorname{barccosh}(cx)}{4d^3(1 - c^2 x^2)^2} - \frac{bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{4d^3}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^3), x]`

```
output -1/4*(b*c*(-1/3*x/((-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (2*x)/(3*Sqrt[-1 +
c*x]*Sqrt[1 + c*x])))/d^3 + (a + b*ArcCosh[c*x])/(4*d^3*(1 - c^2*x^2)^2) +
(-1/2*(b*c*x)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*ArcCosh[c*x])/(2*(1
- c^2*x^2)) - (2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])]) +
(I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh
[c*x])])))/d^3
```

### 3.51.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 41 Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := S
imp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && Eq
Q[b*c + a*d, 0]
```

```
rule 42 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[(-
x)*(a + b*x)^(m + 1)*((c + d*x)^(m + 1)/(2*a*c*(m + 1))), x] + Simp[(2*m +
3)/(2*a*c*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(m + 1), x], x] /; Fre
eQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && ILtQ[m + 3/2, 0]
```

```
rule 2715 Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_.) + (e_.)*(x_)^2)), x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

### 3.51.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 324, normalized size of antiderivative = 1.89

method	result
parts	$-\frac{a\left(-\frac{1}{16(cx+1)^2}-\frac{5}{16(cx+1)}+\frac{\ln(cx+1)}{2}-\ln(x)-\frac{1}{16(cx-1)^2}+\frac{5}{16(cx-1)}+\frac{\ln(cx-1)}{2}\right)}{d^3}-\frac{b\left(\frac{8\sqrt{cx-1}\sqrt{cx+1}c^3x^3-8c^4x^4+6c^2}{d^3}\right)}{d^3}$
derivativedivides	$-\frac{a\left(-\ln(cx)-\frac{1}{16(cx+1)^2}-\frac{5}{16(cx+1)}+\frac{\ln(cx+1)}{2}-\frac{1}{16(cx-1)^2}+\frac{5}{16(cx-1)}+\frac{\ln(cx-1)}{2}\right)}{d^3}-\frac{b\left(\frac{8\sqrt{cx-1}\sqrt{cx+1}c^3x^3-8c^4x^4+6c^2}{d^3}\right)}{d^3}$
default	$-\frac{a\left(-\ln(cx)-\frac{1}{16(cx+1)^2}-\frac{5}{16(cx+1)}+\frac{\ln(cx+1)}{2}-\frac{1}{16(cx-1)^2}+\frac{5}{16(cx-1)}+\frac{\ln(cx-1)}{2}\right)}{d^3}-\frac{b\left(\frac{8\sqrt{cx-1}\sqrt{cx+1}c^3x^3-8c^4x^4+6c^2}{d^3}\right)}{d^3}$

3.51.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)^3} dx$

input `int((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)`

output 
$$-a/d^3*(-1/16/(c*x+1)^2-5/16/(c*x+1)+1/2*\ln(c*x+1)-\ln(x)-1/16/(c*x-1)^2+5/16/(c*x-1)+1/2*\ln(c*x-1))-b/d^3*(1/12*(8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^3*x^3-8*c^4*x^4+6*c^2*x^2*arccosh(c*x)-9*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+16*c^2*x^2-9*arccosh(c*x)-8)/(c^4*x^4-2*c^2*x^2+1)-arccosh(c*x)*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2)-1/2*polylog(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2)+arccosh(c*x)*\ln(1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))+polylog(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))+arccosh(c*x)*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))+polylog(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))$$

### 3.51.5 Fricas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^3 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`

output `integral(-(b*arccosh(c*x) + a)/(c^6*d^3*x^7 - 3*c^4*d^3*x^5 + 3*c^2*d^3*x^3 - d^3*x), x)`

### 3.51.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx = -\int \frac{a}{c^6 x^7 - 3c^4 x^5 + 3c^2 x^3 - x} dx + \int \frac{b \operatorname{arccosh}(cx)}{c^6 x^7 - 3c^4 x^5 + 3c^2 x^3 - x} dx$$

input `integrate((a+b*acosh(c*x))/x/(-c**2*d*x**2+d)**3,x)`

output `-(Integral(a/(c**6*x**7 - 3*c**4*x**5 + 3*c**2*x**3 - x), x) + Integral(b*acosh(c*x)/(c**6*x**7 - 3*c**4*x**5 + 3*c**2*x**3 - x), x))/d**3`

**3.51.7 Maxima [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^3 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

output `-1/4*a*((2*c^2*x^2 - 3)/(c^4*d^3*x^4 - 2*c^2*d^3*x^2 + d^3) + 2*log(c*x + 1)/d^3 + 2*log(c*x - 1)/d^3 - 4*log(x)/d^3) - b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(c^6*d^3*x^7 - 3*c^4*d^3*x^5 + 3*c^2*d^3*x^3 - d^3*x), x)`

**3.51.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^3 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^3*x), x)`

**3.51.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(d - c^2 dx^2)^3} dx$$

input `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^3), x)`

output `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^3), x)`



### 3.52 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d-c^2dx^2)^3} dx$

3.52.1	Optimal result	592
3.52.2	Mathematica [A] (warning: unable to verify)	593
3.52.3	Rubi [C] (verified)	593
3.52.4	Maple [A] (verified)	599
3.52.5	Fricas [F]	599
3.52.6	Sympy [F(-1)]	600
3.52.7	Maxima [F]	600
3.52.8	Giac [F]	601
3.52.9	Mupad [F(-1)]	601

#### 3.52.1 Optimal result

Integrand size = 25, antiderivative size = 230

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2(d - c^2dx^2)^3} dx = \frac{bc}{12d^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} - \frac{7bc}{8d^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{a + \operatorname{arccosh}(cx)}{d^3x(1 - c^2x^2)^2} + \frac{5c^2x(a + \operatorname{arccosh}(cx))}{4d^3(1 - c^2x^2)^2}$$

$$+ \frac{15c^2x(a + \operatorname{arccosh}(cx))}{8d^3(1 - c^2x^2)} + \frac{bc \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d^3}$$

$$+ \frac{15c(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{4d^3}$$

$$+ \frac{15bc \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{8d^3} - \frac{15bc \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{8d^3}$$

output  $1/12*b*c/d^3/(c*x-1)^{(3/2)}/(c*x+1)^{(3/2)}+(-a-b*\operatorname{arccosh}(c*x))/d^3/x/(-c^2*x^2+1)^2+5/4*c^2*x*(a+b*\operatorname{arccosh}(c*x))/d^3/(-c^2*x^2+1)^2+15/8*c^2*x*(a+b*\operatorname{arccosh}(c*x))/d^3/(-c^2*x^2+1)+b*c*\operatorname{arctan}((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d^3+15/4*c*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d^3+15/8*b*c*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d^3-15/8*b*c*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d^3-7/8*b*c/d^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.52.2 Mathematica [A] (warning: unable to verify)

Time = 1.19 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.57

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx$$

$$= \frac{-\frac{96a}{x} + \frac{24ac^2x}{(-1+c^2x^2)^2} - \frac{84ac^2x}{-1+c^2x^2} - \frac{2bc((-2+cx)\sqrt{-1+cx}\sqrt{1+cx}-3\operatorname{arccosh}(cx))}{(-1+cx)^2} + \frac{2bc(\sqrt{-1+cx}\sqrt{1+cx}(2+cx)-3\operatorname{arccosh}(cx))}{(1+cx)^2}}{}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^3), x]`

output `((-96*a)/x + (24*a*c^2*x)/(-1 + c^2*x^2)^2 - (84*a*c^2*x)/(-1 + c^2*x^2) - (2*b*c*((-2 + c*x)*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 3*ArcCosh[c*x]))/(-1 + c*x)^2 + (2*b*c*(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2 + c*x) - 3*ArcCosh[c*x]))/(1 + c*x)^2 - (96*b*ArcCosh[c*x])/x + 42*b*c*(-(1/Sqrt[(-1 + c*x)/(1 + c*x)]) + ArcCosh[c*x]/(1 - c*x)) + 42*b*c*(Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x]/(1 + c*x)) + (96*b*c*Sqrt[-1 + c^2*x^2]*ArcTan[Sqrt[-1 + c^2*x^2]])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - 90*a*c*Log[1 - c*x] + 90*a*c*Log[1 + c*x] - 45*b*c*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 + E^ArcCosh[c*x]]) - 4*PolyLog[2, -E^ArcCosh[c*x]]) + 45*b*c*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 - E^ArcCosh[c*x]]) - 4*PolyLog[2, E^ArcCosh[c*x]]))/(96*d^3)`

### 3.52.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.13 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.17, number of steps used = 19, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.720$ , Rules used = {6347, 27, 115, 27, 115, 27, 103, 218, 6316, 83, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx$$

$$\downarrow 6347$$

$$5c^2 \int \frac{a + \operatorname{barccosh}(cx)}{d^3 (1 - c^2 x^2)^3} dx + \frac{bc \int \frac{1}{x(cx-1)^{5/2}(cx+1)^{5/2}} dx}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2 x^2)^2}$$

---

3.52.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx}{d^3} + \frac{bc \int \frac{1}{x(cx-1)^{5/2}(cx+1)^{5/2}} dx}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2x^2)^2} \\
& \downarrow 115 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx}{d^3} + \frac{bc \left( -\frac{\int \frac{3c}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx}{3c} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2x^2)^2} \\
& \downarrow 27 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx}{d^3} + \frac{bc \left( -\int \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2x^2)^2} \\
& \downarrow 115 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx}{d^3} + \frac{bc \left( \frac{\int \frac{c}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{c} + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2x^2)^2} \\
& \downarrow 27 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx}{d^3} + \frac{bc \left( \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2x^2)^2} \\
& \downarrow 103 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx}{d^3} + \\
& \frac{bc \left( c \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2x^2)^2} \\
& \downarrow 218 \\
& \frac{5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2x^2)^2} + \\
& \frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} \\
& \downarrow 6316
\end{aligned}$$

$$\frac{5c^2 \left( \frac{3}{4} \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - \frac{1}{4}bc \int \frac{x}{(cx-1)^{5/2}(cx+1)^{5/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} +$$

$$\frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3}$$

83

$$\frac{5c^2 \left( \frac{3}{4} \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} +$$

$$\frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3}$$

6316

$$\frac{5c^2 \left( \frac{3}{4} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} +$$

$$\frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3}$$

83

$$\frac{5c^2 \left( \frac{3}{4} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} +$$

$$\frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3}$$

6318

$$\frac{5c^2 \left( \frac{3}{4} \left( -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} +$$

$$\frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3}$$

3042

---

3.52.  $\int \frac{a+\operatorname{barccosh}(cx)}{x^2(d-c^2dx^2)^3} dx$

$$5c^2 \left( \frac{\frac{3}{4} \left( -\frac{i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2}}{d^3} \right. \\ \left. + \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} + \frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} \right)$$

↓ 26

$$5c^2 \left( \frac{\frac{3}{4} \left( -\frac{i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2}}{d^3} \right. \\ \left. + \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} + \frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} \right)$$

↓ 4670

$$5c^2 \left( \frac{\frac{3}{4} \left( -\frac{i(ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx))}{2c}}{d^3} \right. \right. \\ \left. \left. + \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} + \frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} \right) \right)$$

↓ 2715

$$5c^2 \left( \frac{\frac{3}{4} \left( -\frac{i(ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{2c}}{d^3} \right. \right. \\ \left. \left. + \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} + \frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} \right) \right)$$

↓ 2838

$$5c^2 \left( \frac{\frac{3}{4} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{2c}}{d^3} \right. \right. \\ \left. \left. + \frac{a + \operatorname{barccosh}(cx)}{d^3x(1-c^2x^2)^2} + \frac{bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{d^3} \right) \right)$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^3), x]`

3.52.  $\int \frac{a+\operatorname{barccosh}(cx)}{x^2(d-c^2dx^2)^3} dx$

output  $-\frac{(a + b \operatorname{ArcCosh}[c x])}{(d^3 x (1 - c^2 x^2)^2)} + \frac{b c (-1/3 + 1/((-1 + c x)^{3/2} (1 + c x)^{3/2})) + 1/(\operatorname{Sqrt}[-1 + c x] \operatorname{Sqrt}[1 + c x]) + \operatorname{ArcTan}[\operatorname{Sqrt}[-1 + c x] \operatorname{Sqrt}[1 + c x]])}{d^3} + \frac{5 c^2 (b/(12 c (-1 + c x)^{3/2} (1 + c x)^{3/2})) + (x (a + b \operatorname{ArcCosh}[c x]))}{4 (1 - c^2 x^2)^2} + \frac{3 (-1/2 b / (c \operatorname{Sqrt}[-1 + c x] \operatorname{Sqrt}[1 + c x]) + (x (a + b \operatorname{ArcCosh}[c x]))}{2 (1 - c^2 x^2)} - \frac{((I/2) ((2 I) (a + b \operatorname{ArcCosh}[c x]) \operatorname{ArcTanh}[E^{\operatorname{ArcCosh}[c x]}] + I b \operatorname{PolyLog}[2, -E^{\operatorname{ArcCosh}[c x]}] - I b \operatorname{PolyLog}[2, E^{\operatorname{ArcCosh}[c x]}]) / c)}{4}}{d^3}$

### 3.52.3.1 Defintions of rubi rules used

rule 26  $\operatorname{Int}[(\operatorname{Complex}[0, a]) (F x), x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Complex}[\operatorname{Identity}[0], a]) \operatorname{Int}[F x, x], x] /;$   $\operatorname{FreeQ}[a, x] \ \&\& \ \operatorname{EqQ}[a^2, 1]$

rule 27  $\operatorname{Int}[(a) (F x), x\_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F x, x], x] /;$   $\operatorname{FreeQ}[a, x] \ \&\& \ \operatorname{MatchQ}[F x, (b) (G x)] /;$   $\operatorname{FreeQ}[b, x]$

rule 83  $\operatorname{Int}[(a) + (b) (x) ((c) + (d) (x))^{(n)} ((e) + (f) (x))^{(p)}, x] \rightarrow \operatorname{Simp}[b (c + d x)^{(n+1)} ((e + f x)^{(p+1)} / (d f (n + p + 2))), x] /;$   $\operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \operatorname{NeQ}[n + p + 2, 0] \ \&\& \ \operatorname{EqQ}[a d f (n + p + 2) - b (d e (n + 1) + c f (p + 1)), 0]$

rule 103  $\operatorname{Int}[1/(\operatorname{Sqrt}[(a) + (b) (x)] \operatorname{Sqrt}[(c) + (d) (x)] ((e) + (f) (x))), x] \rightarrow \operatorname{Simp}[b f \operatorname{Subst}[\operatorname{Int}[1/(d (b e - a f)^2 + b f^2 x^2), x], x, \operatorname{Sqrt}[a + b x] \operatorname{Sqrt}[c + d x]], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{EqQ}[2 b d * e - f (b c + a d), 0]$

rule 115  $\operatorname{Int}[(a) + (b) (x) ((c) + (d) (x))^{(n)} ((e) + (f) (x))^{(p)}, x] \rightarrow \operatorname{Simp}[b (a + b x)^{(m+1)} (c + d x)^{(n+1)} ((e + f x)^{(p+1)} / ((m + 1) (b c - a d) (b e - a f))), x] + \operatorname{Simp}[1/((m + 1) (b c - a d) (b e - a f)) \operatorname{Int}[(a + b x)^{(m+1)} (c + d x)^n (e + f x)^p \operatorname{Simp}[a d f (m + 1) - b (d e (m + n + 2) + c f (m + p + 2)) - b d f (m + n + p + 3) x, x], x], x] /;$   $\operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \operatorname{LtQ}[m, -1] \ \&\& \ \operatorname{IntegersQ}[2 m, 2 * n, 2 * p]$

rule 218  $\operatorname{Int}[(a) + (b) (x)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2]/a) \operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /;$   $\operatorname{FreeQ}\{a, b\}, x] \ \&\& \ \operatorname{PosQ}[a/b]$

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))]^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x
_Symbol] :> Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p +
1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*
ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 +
c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a
+ b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*
d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symb
ol] :> Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x
]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1
))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp
[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(
f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^
(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && ILtQ[m, -1]`

### 3.52.4 Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.17

method	result
derivativedivides	$c \left( -\frac{a \left( \frac{1}{cx} + \frac{1}{16(cx+1)^2} + \frac{7}{16(cx+1)} - \frac{15 \ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} + \frac{7}{16(cx-1)} + \frac{15 \ln(cx-1)}{16} \right)}{d^3} - \frac{b \left( \frac{45c^4 x^4 \operatorname{arccosh}(cx) + 21\sqrt{cx-1}}{\dots} \right)}{\dots} \right)$
default	$c \left( -\frac{a \left( \frac{1}{cx} + \frac{1}{16(cx+1)^2} + \frac{7}{16(cx+1)} - \frac{15 \ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} + \frac{7}{16(cx-1)} + \frac{15 \ln(cx-1)}{16} \right)}{d^3} - \frac{b \left( \frac{45c^4 x^4 \operatorname{arccosh}(cx) + 21\sqrt{cx-1}}{\dots} \right)}{\dots} \right)$
parts	$-\frac{a \left( \frac{c}{16(cx+1)^2} + \frac{7c}{16(cx+1)} - \frac{15c \ln(cx+1)}{16} + \frac{1}{x} - \frac{c}{16(cx-1)^2} + \frac{7c}{16(cx-1)} + \frac{15c \ln(cx-1)}{16} \right)}{d^3} - \frac{bc \left( \frac{45c^4 x^4 \operatorname{arccosh}(cx) + 21\sqrt{cx-1}}{\dots} \right)}{\dots}$

input `int((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)`

output `c*(-a/d^3*(1/c/x+1/16/(c*x+1)^2+7/16/(c*x+1)-15/16*ln(c*x+1)-1/16/(c*x-1)^2+7/16/(c*x-1)+15/16*ln(c*x-1))-b/d^3*(1/24*(45*c^4*x^4*arccosh(c*x)+21*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-75*c^2*x^2*arccosh(c*x)-23*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+24*arccosh(c*x))/(c^4*x^4-2*c^2*x^2+1)/c/x-2*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-15/8*dilog(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-15/8*dilog(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-15/8*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))`

### 3.52.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^3 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`

output `integral(-(b*arccosh(c*x) + a)/(c^6*d^3*x^8 - 3*c^4*d^3*x^6 + 3*c^2*d^3*x^4 - d^3*x^2), x)`



### 3.52.6 Sympy [F(-1)]

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**2/(-c**2*d*x**2+d)**3,x)`

output `Timed out`

### 3.52.7 Maxima [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

output `1/2048*(92160*c^7*integrate(1/32*x^5*log(c*x - 1)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x) - 240*c^6*(2*(5*c^2*x^3 - 3*x)/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3) + 3*log(c*x + 1)/(c^5*d^3) - 3*log(c*x - 1)/(c^5*d^3)) - 30720*c^6*integrate(1/32*x^4*log(c*x - 1)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x) + 90*(c*(2*(5*c^2*x^2 + 3*c*x - 6)/(c^8*d^3*x^3 - c^7*d^3*x^2 - c^6*d^3*x + c^5*d^3) - 5*log(c*x + 1)/(c^5*d^3) + 5*log(c*x - 1)/(c^5*d^3)) + 16*(2*c^2*x^2 - 1)*log(c*x - 1)/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3))*c^5 + 400*c^4*(2*(c^2*x^3 + x)/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3) - log(c*x + 1)/(c^3*d^3) + log(c*x - 1)/(c^3*d^3)) + 61440*c^4*integrate(1/32*x^2*log(c*x - 1)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x) + 45*(c*(2*(3*c^2*x^2 - 3*c*x - 2)/(c^6*d^3*x^3 - c^5*d^3*x^2 - c^4*d^3*x + c^3*d^3) - 3*log(c*x + 1)/(c^3*d^3) + 3*log(c*x - 1)/(c^3*d^3)) - 16*log(c*x - 1)/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3))*c^3 + 128*c^2*(2*(3*c^2*x^3 - 5*x)/(c^4*d^3*x^4 - 2*c^2*d^3*x^2 + d^3) - 3*log(c*x + 1)/(c*d^3) + 3*log(c*x - 1)/(c*d^3)) - 30720*c^2*integrate(1/32*log(c*x - 1)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x) - 32*(15*(c^5*x^5 - 2*c^3*x^3 + c*x)*log(c*x + 1)^2 + 30*(c^5*x^5 - 2*c^3*x^3 + c*x)*log(c*x + 1)*log(c*x - 1) + 4*(30*c^4*x^4 - 50*c^2*x^2 - 15*(c^5*x^5 - 2*c^3*x^3 + c*x)*log(c*x + 1) + 15*(c^5*x^5 - 2*c^3*x^3 + c*x)*log(c*x - 1) + 16)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/c^4*d^3*x^5 ...`

**3.52.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^3*x^2), x)`

**3.52.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2 (d - c^2 dx^2)^3} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^3),x)`

output `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^3), x)`

### 3.53 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^3} dx$

3.53.1	Optimal result	602
3.53.2	Mathematica [A] (warning: unable to verify)	603
3.53.3	Rubi [C] (verified)	603
3.53.4	Maple [A] (verified)	609
3.53.5	Fricas [F]	610
3.53.6	Sympy [F(-1)]	610
3.53.7	Maxima [F]	610
3.53.8	Giac [F]	611
3.53.9	Mupad [F(-1)]	611

#### 3.53.1 Optimal result

Integrand size = 25, antiderivative size = 250

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^3(d - c^2dx^2)^3} dx = \frac{bc}{2d^3x(-1 + cx)^{3/2}(1 + cx)^{3/2}} - \frac{5bc^3x}{12d^3(-1 + cx)^{3/2}(1 + cx)^{3/2}}$$

$$- \frac{2bc^3x}{3d^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{3c^2(a + \operatorname{arccosh}(cx))}{4d^3(1 - c^2x^2)^2}$$

$$- \frac{a + \operatorname{arccosh}(cx)}{2d^3x^2(1 - c^2x^2)^2} + \frac{3c^2(a + \operatorname{arccosh}(cx))}{2d^3(1 - c^2x^2)}$$

$$+ \frac{6c^2(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d^3}$$

$$+ \frac{3bc^2 \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{2d^3} - \frac{3bc^2 \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{2d^3}$$

output  $1/2*b*c/d^3/x/(c*x-1)^{(3/2)}/(c*x+1)^{(3/2)}-5/12*b*c^3*x/d^3/(c*x-1)^{(3/2)}/(c*x+1)^{(3/2)}+3/4*c^2*(a+b*\operatorname{arccosh}(c*x))/d^3/(-c^2*x^2+1)^2+1/2*(-a-b*\operatorname{arccosh}(c*x))/d^3/x^2/(-c^2*x^2+1)^2+3/2*c^2*(a+b*\operatorname{arccosh}(c*x))/d^3/(-c^2*x^2+1)+6*c^2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2)/d^3+3+3/2*b*c^2*\operatorname{polylog}(2, -(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2)/d^3-3/2*b*c^2*\operatorname{polylog}(2, (c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2)/d^3-2/3*b*c^3*x/d^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.53.2 Mathematica [A] (warning: unable to verify)

Time = 1.08 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.61

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx$$

$$= \frac{-\frac{2a}{x^2} + \frac{ac^2}{(-1+c^2x^2)^2} - \frac{4ac^2}{-1+c^2x^2} - \frac{bc^2((-2+cx)\sqrt{-1+cx}\sqrt{1+cx}-3\operatorname{arccosh}(cx))}{12(-1+cx)^2} - \frac{bc^2(\sqrt{-1+cx}\sqrt{1+cx}(2+cx)-3\operatorname{arccosh}(cx))}{12(1+cx)^2}}{1}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^3), x]`

output

```
((-2*a)/x^2 + (a*c^2)/(-1 + c^2*x^2)^2 - (4*a*c^2)/(-1 + c^2*x^2) - (b*c^2
*((-2 + c*x)*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 3*ArcCosh[c*x]))/(12*(-1 + c*x
)^2) - (b*c^2*(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2 + c*x) - 3*ArcCosh[c*x]))/(
12*(1 + c*x)^2) + (2*b*(c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - ArcCosh[c*x]))/
x^2 + (9*b*c^2*(-(1/Sqrt[(-1 + c*x)/(1 + c*x)]) + ArcCosh[c*x]/(1 - c*x))
/4 - (9*b*c^2*(Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x]/(1 + c*x)))/4 + 1
2*a*c^2*Log[x] - 6*a*c^2*Log[1 - c^2*x^2] + 6*b*c^2*(ArcCosh[c*x]*(ArcCosh
[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLog[2, -E^(-2*ArcCosh[c*x])])
) + 3*b*c^2*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 + E^ArcCosh[c*x]]) - 4*P
olyLog[2, -E^ArcCosh[c*x]]) + 3*b*c^2*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[
1 - E^ArcCosh[c*x]]) - 4*PolyLog[2, E^ArcCosh[c*x]]))/(4*d^3)
```

### 3.53.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.37 (sec) , antiderivative size = 312, normalized size of antiderivative = 1.25, number of steps used = 19, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.720$ , Rules used = {6347, 27, 114, 27, 42, 41, 6351, 42, 41, 6351, 41, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx$$

↓ 6347

$$\begin{aligned}
& 3c^2 \int \frac{a + \operatorname{barccosh}(cx)}{d^3 x (1 - c^2 x^2)^3} dx + \frac{bc \int \frac{1}{x^2 (cx-1)^{5/2} (cx+1)^{5/2}} dx}{2d^3} - \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} \\
& \quad \downarrow 27 \\
& \frac{3c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(1-c^2 x^2)^3} dx}{d^3} + \frac{bc \int \frac{1}{x^2 (cx-1)^{5/2} (cx+1)^{5/2}} dx}{2d^3} - \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} \\
& \quad \downarrow 114 \\
& \frac{3c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(1-c^2 x^2)^3} dx}{d^3} + \frac{bc \left( \int \frac{4c^2}{(cx-1)^{5/2} (cx+1)^{5/2}} dx + \frac{1}{x(cx-1)^{3/2} (cx+1)^{3/2}} \right)}{2d^3} - \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} \\
& \quad \downarrow 27 \\
& \frac{3c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(1-c^2 x^2)^3} dx}{d^3} + \frac{bc \left( 4c^2 \int \frac{1}{(cx-1)^{5/2} (cx+1)^{5/2}} dx + \frac{1}{x(cx-1)^{3/2} (cx+1)^{3/2}} \right)}{2d^3} - \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} \\
& \quad \downarrow 42 \\
& \frac{3c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(1-c^2 x^2)^3} dx}{d^3} + \\
& \frac{bc \left( 4c^2 \left( -\frac{2}{3} \int \frac{1}{(cx-1)^{3/2} (cx+1)^{3/2}} dx - \frac{x}{3(cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2} (cx+1)^{3/2}} \right)}{2d^3} - \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} \\
& \quad \downarrow 41 \\
& \frac{3c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(1-c^2 x^2)^3} dx}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} + \\
& \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2} (cx+1)^{3/2}} \right)}{2d^3} \\
& \quad \downarrow 6351 \\
& \frac{3c^2 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x(1-c^2 x^2)^2} dx - \frac{1}{4} bc \int \frac{1}{(cx-1)^{5/2} (cx+1)^{5/2}} dx + \frac{a + \operatorname{barccosh}(cx)}{4(1-c^2 x^2)^2} \right)}{d^3} - \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} + \\
& \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2} (cx+1)^{3/2}} \right)}{2d^3} \\
& \quad \downarrow 42 \\
& \frac{3c^2 \left( \int \frac{a + \operatorname{barccosh}(cx)}{x(1-c^2 x^2)^2} dx - \frac{1}{4} bc \left( -\frac{2}{3} \int \frac{1}{(cx-1)^{3/2} (cx+1)^{3/2}} dx - \frac{x}{3(cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{a + \operatorname{barccosh}(cx)}{4(1-c^2 x^2)^2} \right)}{d^3} \\
& \frac{a + \operatorname{barccosh}(cx)}{2d^3 x^2 (1 - c^2 x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2} (cx+1)^{3/2}} \right)}{2d^3}
\end{aligned}$$

---

3.53.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx$

$$\begin{aligned} & \downarrow 41 \\ & \frac{3c^2 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx + \frac{a+\operatorname{barccosh}(cx)}{4(1-c^2x^2)^2} - \frac{1}{4}bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{d^3} - \\ & \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3} \\ & \downarrow 6351 \\ & \frac{3c^2 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2}bc \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} + \frac{a+\operatorname{barccosh}(cx)}{4(1-c^2x^2)^2} - \frac{1}{4}bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{d^3} - \\ & \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3} \\ & \downarrow 41 \\ & \frac{3c^2 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} + \frac{a+\operatorname{barccosh}(cx)}{4(1-c^2x^2)^2} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{4}bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{d^3} - \\ & \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3} \\ & \downarrow 6331 \\ & \frac{3c^2 \left( - \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} + \frac{a+\operatorname{barccosh}(cx)}{4(1-c^2x^2)^2} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{4}bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{d^3} - \\ & \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3} \\ & \downarrow 5984 \\ & \frac{3c^2 \left( -2 \int (a + \operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} + \frac{a+\operatorname{barccosh}(cx)}{4(1-c^2x^2)^2} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{4}bc \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{d^3} - \\ & \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3} \\ & \downarrow 3042 \end{aligned}$$

---

3.53.  $\int \frac{a+\operatorname{barccosh}(cx)}{x^3(d-c^2dx^2)^3} dx$

$$\frac{3c^2 \left( -2 \int i(a + \operatorname{barccosh}(cx)) \csc(2i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a + \operatorname{barccosh}(cx)}{2(1-c^2x^2)} + \frac{a + \operatorname{barccosh}(cx)}{4(1-c^2x^2)^2} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3}$$

↓ 26

$$\frac{3c^2 \left( -2i \int (a + \operatorname{barccosh}(cx)) \csc(2i \operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a + \operatorname{barccosh}(cx)}{2(1-c^2x^2)} + \frac{a + \operatorname{barccosh}(cx)}{4(1-c^2x^2)^2} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3}$$

↓ 4670

$$\frac{3c^2 \left( -2i \left( \frac{1}{2} ib \int \log(1 - e^{2 \operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} ib \int \log(1 + e^{2 \operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + i \operatorname{arctanh}(e^{2 \operatorname{arccosh}(cx)}) \right) \right)}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3}$$

↓ 2715

$$\frac{3c^2 \left( -2i \left( \frac{1}{4} ib \int e^{-2 \operatorname{arccosh}(cx)} \log(1 - e^{2 \operatorname{arccosh}(cx)}) de^{2 \operatorname{arccosh}(cx)} - \frac{1}{4} ib \int e^{-2 \operatorname{arccosh}(cx)} \log(1 + e^{2 \operatorname{arccosh}(cx)}) de^{2 \operatorname{arccosh}(cx)} \right) \right)}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3}$$

↓ 2838

$$\frac{3c^2 \left( -2i \left( i \operatorname{arctanh}(e^{2 \operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4} ib \operatorname{PolyLog}(2, -e^{2 \operatorname{arccosh}(cx)}) - \frac{1}{4} ib \operatorname{PolyLog}(2, e^{2 \operatorname{arccosh}(cx)}) \right) \right)}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2(1-c^2x^2)^2} + \frac{bc \left( 4c^2 \left( \frac{2x}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{x}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{2d^3}$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^3), x]`

```
output (b*c*(1/(x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + 4*c^2*(-1/3*x/((-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (2*x)/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/(2*d^3)
- (a + b*ArcCosh[c*x])/(2*d^3*x^2*(1 - c^2*x^2)^2) + (3*c^2*(-1/2*(b*c*x)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (b*c*(-1/3*x/((-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (2*x)/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/4 + (a + b*ArcCosh[c*x])/(4*(1 - c^2*x^2)^2) + (a + b*ArcCosh[c*x])/(2*(1 - c^2*x^2)) - (2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])]) + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])]))/d^3
```

### 3.53.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 41 Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]
```

```
rule 42 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[(-x)*(a + b*x)^(m + 1)*((c + d*x)^(m + 1)/(2*a*c*(m + 1))), x] + Simp[(2*m + 3)/(2*a*c*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(m + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && ILtQ[m + 3/2, 0]
```

```
rule 114 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])
```



rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] :> Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x
]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_) + (e_.)*(x_)^2)),
x_Symbol] :> Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x
, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IG
tQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1
))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp
[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(
f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^
(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && ILtQ[m, -1]`

```
rule 6351 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])
```

### 3.53.4 Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 391, normalized size of antiderivative = 1.56

method	result
derivativedivides	$c^2 \left( -\frac{a \left( \frac{1}{2c^2x^2} - 3\ln(cx) - \frac{1}{16(cx+1)^2} - \frac{9}{16(cx+1)} + \frac{3\ln(cx+1)}{2} - \frac{1}{16(cx-1)^2} + \frac{9}{16(cx-1)} + \frac{3\ln(cx-1)}{2} \right)}{d^3} - \frac{b \left( \frac{8\sqrt{cx+1}\sqrt{cx-1}}{\dots} \right)}{\dots} \right)$
default	$c^2 \left( -\frac{a \left( \frac{1}{2c^2x^2} - 3\ln(cx) - \frac{1}{16(cx+1)^2} - \frac{9}{16(cx+1)} + \frac{3\ln(cx+1)}{2} - \frac{1}{16(cx-1)^2} + \frac{9}{16(cx-1)} + \frac{3\ln(cx-1)}{2} \right)}{d^3} - \frac{b \left( \frac{8\sqrt{cx+1}\sqrt{cx-1}}{\dots} \right)}{\dots} \right)$
parts	$-\frac{a \left( -\frac{c^2}{16(cx+1)^2} - \frac{9c^2}{16(cx+1)} + \frac{3c^2\ln(cx+1)}{2} + \frac{1}{2x^2} - 3c^2\ln(x) - \frac{c^2}{16(cx-1)^2} + \frac{9c^2}{16(cx-1)} + \frac{3c^2\ln(cx-1)}{2} \right)}{d^3} - \frac{bc^2 \left( \frac{8\sqrt{cx+1}\sqrt{cx-1}}{\dots} \right)}{\dots}$

```
input int((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)
```

```
output c^2*(-a/d^3*(1/2/c^2/x^2-3*ln(c*x)-1/16/(c*x+1)^2-9/16/(c*x+1)+3/2*ln(c*x+1)-1/16/(c*x-1)^2+9/16/(c*x-1)+3/2*ln(c*x-1))-b/d^3*(1/12/(c^4*x^4-2*c^2*x^2+1)/c^2/x^2*(8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-8*c^6*x^6+18*c^4*x^4*arccosh(c*x)-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+16*c^4*x^4-27*c^2*x^2*arccosh(c*x)-6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-8*c^2*x^2+6*arccosh(c*x))-3*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)-3/2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)+3*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+3*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+3*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+3*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))))
```

$$3.53. \int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^3} dx$$

**3.53.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`

output `integral(-(b*arccosh(c*x) + a)/(c^6*d^3*x^9 - 3*c^4*d^3*x^7 + 3*c^2*d^3*x^5 - d^3*x^3), x)`

**3.53.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**3/(-c**2*d*x**2+d)**3,x)`

output `Timed out`

**3.53.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

output `-1/4*a*((6*c^4*x^4 - 9*c^2*x^2 + 2)/(c^4*d^3*x^6 - 2*c^2*d^3*x^4 + d^3*x^2) + 6*c^2*log(c*x + 1)/d^3 + 6*c^2*log(c*x - 1)/d^3 - 12*c^2*log(x)/d^3) - b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(c^6*d^3*x^9 - 3*c^4*d^3*x^7 + 3*c^2*d^3*x^5 - d^3*x^3), x)`

**3.53.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^3*x^3), x)`

**3.53.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (d - c^2 dx^2)^3} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^3),x)`

output `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^3), x)`

### 3.54 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^4(d-c^2dx^2)^3} dx$

3.54.1	Optimal result	612
3.54.2	Mathematica [A] (warning: unable to verify)	613
3.54.3	Rubi [C] (verified)	613
3.54.4	Maple [A] (verified)	622
3.54.5	Fricas [F]	623
3.54.6	Sympy [F(-1)]	623
3.54.7	Maxima [F]	623
3.54.8	Giac [F]	624
3.54.9	Mupad [F(-1)]	625

#### 3.54.1 Optimal result

Integrand size = 25, antiderivative size = 310

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx = -\frac{bc^3}{12d^3(-1 + cx)^{3/2}(1 + cx)^{3/2}} + \frac{bc}{6d^3x^2(-1 + cx)^{3/2}(1 + cx)^{3/2}}$$

$$-\frac{29bc^3}{24d^3\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{a + \operatorname{arccosh}(cx)}{3d^3x^3(1 - c^2x^2)^2}$$

$$-\frac{7c^2(a + \operatorname{arccosh}(cx))}{3d^3x(1 - c^2x^2)^2} + \frac{35c^4x(a + \operatorname{arccosh}(cx))}{12d^3(1 - c^2x^2)^2}$$

$$+ \frac{35c^4x(a + \operatorname{arccosh}(cx))}{8d^3(1 - c^2x^2)} + \frac{19bc^3 \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{6d^3}$$

$$+ \frac{35c^3(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{4d^3}$$

$$+ \frac{35bc^3 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{8d^3} - \frac{35bc^3 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{8d^3}$$

output

```
-1/12*b*c^3/d^3/(c*x-1)^(3/2)/(c*x+1)^(3/2)+1/6*b*c/d^3/x^2/(c*x-1)^(3/2)/
(c*x+1)^(3/2)+1/3*(-a-b*arccosh(c*x))/d^3/x^3/(-c^2*x^2+1)^2-7/3*c^2*(a+b*
arccosh(c*x))/d^3/x/(-c^2*x^2+1)^2+35/12*c^4*x*(a+b*arccosh(c*x))/d^3/(-c^
2*x^2+1)^2+35/8*c^4*x*(a+b*arccosh(c*x))/d^3/(-c^2*x^2+1)+19/6*b*c^3*arcta
n((c*x-1)^(1/2)*(c*x+1)^(1/2))/d^3+35/4*c^3*(a+b*arccosh(c*x))*arctanh(c*x
+(c*x-1)^(1/2)*(c*x+1)^(1/2))/d^3+35/8*b*c^3*polylog(2,-c*x-(c*x-1)^(1/2)*
(c*x+1)^(1/2))/d^3-35/8*b*c^3*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/d
^3-29/24*b*c^3/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.54.2 Mathematica [A] (warning: unable to verify)

Time = 1.33 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.52

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx$$

$$= \frac{-\frac{16a}{x^3} - \frac{144ac^2}{x} + \frac{12ac^4x}{(-1+c^2x^2)^2} - \frac{66ac^4x}{-1+c^2x^2} - \frac{bc^3((-2+cx)\sqrt{-1+cx}\sqrt{1+cx}-3\operatorname{arccosh}(cx))}{(-1+cx)^2} + \frac{bc^3(\sqrt{-1+cx}\sqrt{1+cx}(2+cx)-3\operatorname{arccosh}(cx))}{(1+cx)^2}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^3), x]`

output `((-16*a)/x^3 - (144*a*c^2)/x + (12*a*c^4*x)/(-1 + c^2*x^2)^2 - (66*a*c^4*x)/(-1 + c^2*x^2) - (b*c^3*((-2 + c*x)*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 3*ArcCosh[c*x]))/(-1 + c*x)^2 + (b*c^3*(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2 + c*x) - 3*ArcCosh[c*x]))/(1 + c*x)^2 + 33*b*c^3*(-(1/Sqrt[(-1 + c*x)/(1 + c*x)]) + ArcCosh[c*x]/(1 - c*x)) + 33*b*c^3*(Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x]/(1 + c*x)) + 144*b*c^2*(-(ArcCosh[c*x]/x) + (c*Sqrt[-1 + c^2*x^2]*ArcTan[Sqrt[-1 + c^2*x^2]]/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (8*b*(-2*ArcCosh[c*x] + (c*x*(-1 + c^2*x^2 + c^2*x^2*Sqrt[-1 + c^2*x^2]*ArcTan[Sqrt[-1 + c^2*x^2]])))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/x^3 - 105*a*c^3*Log[1 - c*x] + 105*a*c^3*Log[1 + c*x] - (105*b*c^3*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 + E^ArcCosh[c*x]]) - 4*PolyLog[2, -E^ArcCosh[c*x]]))/2 + (105*b*c^3*(ArcCosh[c*x]*(ArcCosh[c*x] - 4*Log[1 - E^ArcCosh[c*x]]) - 4*PolyLog[2, E^ArcCosh[c*x]]))/2)/(48*d^3)`

### 3.54.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.66 (sec) , antiderivative size = 406, normalized size of antiderivative = 1.31, number of steps used = 28, number of rules used = 27,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.080$ , Rules used = {6347, 27, 114, 27, 115, 27, 115, 27, 103, 218, 6347, 115, 27, 115, 27, 103, 218, 6316, 83, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx$$

---

3.54.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx$

$$\begin{aligned}
& \downarrow 6347 \\
& \frac{7}{3}c^2 \int \frac{a + \operatorname{barccosh}(cx)}{d^3 x^2 (1 - c^2 x^2)^3} dx + \frac{bc \int \frac{1}{x^3 (cx-1)^{5/2} (cx+1)^{5/2}} dx}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \downarrow 27 \\
& \frac{7c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (1 - c^2 x^2)^3} dx}{3d^3} + \frac{bc \int \frac{1}{x^3 (cx-1)^{5/2} (cx+1)^{5/2}} dx}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \downarrow 114 \\
& \frac{7c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (1 - c^2 x^2)^3} dx}{3d^3} + \frac{bc \left( \frac{1}{2} \int \frac{5c^2}{x (cx-1)^{5/2} (cx+1)^{5/2}} dx + \frac{1}{2x^2 (cx-1)^{3/2} (cx+1)^{3/2}} \right)}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \downarrow 27 \\
& \frac{7c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (1 - c^2 x^2)^3} dx}{3d^3} + \frac{bc \left( \frac{5}{2} c^2 \int \frac{1}{x (cx-1)^{5/2} (cx+1)^{5/2}} dx + \frac{1}{2x^2 (cx-1)^{3/2} (cx+1)^{3/2}} \right)}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \downarrow 115 \\
& \frac{7c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (1 - c^2 x^2)^3} dx}{3d^3} + \\
& \frac{bc \left( \frac{5}{2} c^2 \left( - \frac{\int \frac{3c}{x (cx-1)^{3/2} (cx+1)^{3/2}} dx}{3c} - \frac{1}{3 (cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{1}{2x^2 (cx-1)^{3/2} (cx+1)^{3/2}} \right)}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \downarrow 27 \\
& \frac{7c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (1 - c^2 x^2)^3} dx}{3d^3} + \\
& \frac{bc \left( \frac{5}{2} c^2 \left( - \int \frac{1}{x (cx-1)^{3/2} (cx+1)^{3/2}} dx - \frac{1}{3 (cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{1}{2x^2 (cx-1)^{3/2} (cx+1)^{3/2}} \right)}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \downarrow 115 \\
& \frac{7c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (1 - c^2 x^2)^3} dx}{3d^3} + \\
& \frac{bc \left( \frac{5}{2} c^2 \left( \frac{\int \frac{c}{x \sqrt{cx-1} \sqrt{cx+1}} dx}{c} + \frac{1}{\sqrt{cx-1} \sqrt{cx+1}} - \frac{1}{3 (cx-1)^{3/2} (cx+1)^{3/2}} \right) + \frac{1}{2x^2 (cx-1)^{3/2} (cx+1)^{3/2}} \right)}{3d^3} - \\
& \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \downarrow 27
\end{aligned}$$

---

3.54.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx$

$$\begin{aligned}
& \frac{7c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^3} dx}{3d^3} + \\
& \frac{bc \left( \frac{5}{2}c^2 \left( \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3} \\
& \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \quad \downarrow \text{103} \\
& \frac{7c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^3} dx}{3d^3} + \\
& \frac{bc \left( \frac{5}{2}c^2 \left( c \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3} \\
& \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} \\
& \quad \downarrow \text{218} \\
& \frac{7c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^3} dx}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} + \\
& \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3} \\
& \quad \downarrow \text{6347} \\
& \frac{7c^2 \left( 5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx + bc \int \frac{1}{x(cx-1)^{5/2}(cx+1)^{5/2}} dx - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} \right)}{3d^3} - \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} + \\
& \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3} \\
& \quad \downarrow \text{115} \\
& \frac{7c^2 \left( 5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx + bc \left( -\frac{\int \frac{3c}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx}{3c} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} \right)}{3d^3} \\
& \frac{a + \operatorname{barccosh}(cx)}{3d^3 x^3 (1 - c^2 x^2)^2} + \\
& \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3} \\
& \quad \downarrow \text{27}
\end{aligned}$$



$$\frac{7c^2 \left( 5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx + bc \left( - \int \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 115

$$\frac{7c^2 \left( 5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx + bc \left( \int \frac{\frac{c}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{c} + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 27

$$\frac{7c^2 \left( 5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx + bc \left( \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 103

$$\frac{7c^2 \left( 5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx + bc \left( c \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 218

$$\frac{7c^2 \left( 5c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^3} dx - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} + bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{3d^3} + \frac{bc \left( \frac{5}{2} c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 6316

$$\frac{7c^2 \left( 5c^2 \left( \frac{3}{4} \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - \frac{1}{4} bc \int \frac{x}{(cx-1)^{5/2}(cx+1)^{5/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} + bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 83

$$\frac{7c^2 \left( 5c^2 \left( \frac{3}{4} \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} + bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 6316

$$\frac{7c^2 \left( 5c^2 \left( \frac{3}{4} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2} bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{1}{12c(cx-1)} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} + bc \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 83

$$\frac{7c^2 \left( 5c^2 \left( \frac{3}{4} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 6318

$$\frac{7c^2 \left( 5c^2 \left( \frac{3}{4} \left( -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 3042

$$\frac{7c^2 \left( 5c^2 \left( \frac{3}{4} \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 26

$$\frac{7c^2 \left( 5c^2 \left( \frac{3}{4} \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x(a+\operatorname{barccosh}(cx))}{4(1-c^2x^2)^2} + \frac{b}{12c(cx-1)^{3/2}(cx+1)^{3/2}} \right) \right)}{3d^3} + \frac{a + \operatorname{barccosh}(cx)}{3d^3x^3(1-c^2x^2)^2} + \frac{bc \left( \frac{5}{2}c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 4670

$$7c^2 \left( 5c^2 \left( \frac{3}{4} \left( -\frac{i(ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)) dx}{2c} \right) \right) \right)$$

$$\frac{bc \left( \frac{5}{2} c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 2715

$$7c^2 \left( 5c^2 \left( \frac{3}{4} \left( -\frac{i(ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)) dx}{2c} \right) \right) \right)$$

$$\frac{bc \left( \frac{5}{2} c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

↓ 2838

$$7c^2 \left( 5c^2 \left( \frac{3}{4} \left( -\frac{i(2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{2c} \right) \right) + \frac{x(a + b \operatorname{arccosh}(cx))}{2(1 - c^2 x^2)} \right)$$

$$\frac{bc \left( \frac{5}{2} c^2 \left( \arctan(\sqrt{cx-1}\sqrt{cx+1}) + \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{3(cx-1)^{3/2}(cx+1)^{3/2}} \right) + \frac{1}{2x^2(cx-1)^{3/2}(cx+1)^{3/2}} \right)}{3d^3}$$

input `Int[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^3), x]`

output 
$$\begin{aligned} & -1/3*(a + b*\text{ArcCosh}[c*x])/(d^3*x^3*(1 - c^2*x^2)^2) + (b*c*(1/(2*x^2*(-1 + \\ & c*x)^{3/2}*(1 + c*x)^{3/2})) + (5*c^2*(-1/3*1/((-1 + c*x)^{3/2}*(1 + c*x)^{3/2})) \\ & + 1/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) + \text{ArcTan}[\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + \\ & c*x]]))/2)/(3*d^3) + (7*c^2*(-((a + b*\text{ArcCosh}[c*x])/(x*(1 - c^2*x^2)^2)) \\ & + b*c*(-1/3*1/((-1 + c*x)^{3/2}*(1 + c*x)^{3/2})) + 1/(\text{Sqrt}[-1 + c*x]*\text{Sqrt} \\ & [1 + c*x]) + \text{ArcTan}[\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]]) + 5*c^2*(b/(12*c*(-1 + \\ & c*x)^{3/2}*(1 + c*x)^{3/2})) + (x*(a + b*\text{ArcCosh}[c*x]))/(4*(1 - c^2*x^2)^2) \\ & + (3*(-1/2*b/(c*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) + (x*(a + b*\text{ArcCosh}[c*x]))/ \\ & (2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*\text{ArcCosh}[c*x])* \text{ArcTanh}[E^{\text{ArcCosh}[c \\ & *x]] + I*b*\text{PolyLog}[2, -E^{\text{ArcCosh}[c*x]}] - I*b*\text{PolyLog}[2, E^{\text{ArcCosh}[c*x]})]/ \\ & c))/4))/(3*d^3) \end{aligned}$$

### 3.54.3.1 Defintions of rubi rules used

rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F_x_), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /;$   $\text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 27  $\text{Int}[(a_)*(F_x_), x\_Symbol] \rightarrow \text{Simp}[a \text{Int}[F_x, x], x] /;$   $\text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x_)] /;$   $\text{FreeQ}[b, x]$

rule 83  $\text{Int}[(a_. + (b_.)*(x_))*((c_. + (d_.)*(x_))^{(n_.)*((e_. + (f_.)*(x_))^{(p_.)}, x_)] \rightarrow \text{Simp}[b*(c + d*x)^{(n + 1)}*(e + f*x)^{(p + 1)}/(d*f*(n + p + 2)), x] /;$   $\text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 2, 0] \ \&\& \ \text{EqQ}[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]$

rule 103  $\text{Int}[1/(\text{Sqrt}[(a_. + (b_.)*(x_)]*\text{Sqrt}[(c_. + (d_.)*(x_)]*((e_. + (f_.)*(x_)))), x_] \rightarrow \text{Simp}[b*f \text{Subst}[\text{Int}[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, \text{Sqrt}[a + b*x]*\text{Sqrt}[c + d*x]], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[2*b*d*e - f*(b*c + a*d), 0]$

rule 114  $\text{Int}[(a_. + (b_.)*(x_))^{(m_)*((c_. + (d_.)*(x_))^{(n_)*((e_. + (f_.)*(x_))^{(p_)}, x_)] \rightarrow \text{Simp}[b*(a + b*x)^{(m + 1)}*(c + d*x)^{(n + 1)}*(e + f*x)^{(p + 1)}/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + \text{Simp}[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n*(e + f*x)^p*\text{Simp}[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /;$   $\text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ (\text{IntegerQ}[n] \ || \ \text{IntegersQ}[2*n, 2*p] \ || \ \text{ILtQ}[m + n + p + 3, 0])$

rule 115 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

### 3.54.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 314, normalized size of antiderivative = 1.01

method	result
derivativedivides	$c^3 \left( -\frac{a \left( \frac{1}{3c^3x^3} + \frac{3}{cx} + \frac{1}{16(cx+1)^2} + \frac{11}{16(cx+1)} - \frac{35 \ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} + \frac{11}{16(cx-1)} + \frac{35 \ln(cx-1)}{16} \right)}{d^3} - \frac{b \left( \frac{105 \operatorname{arccosh}(cx)c^6}{16} \right)}{d^3} \right)$
default	$c^3 \left( -\frac{a \left( \frac{1}{3c^3x^3} + \frac{3}{cx} + \frac{1}{16(cx+1)^2} + \frac{11}{16(cx+1)} - \frac{35 \ln(cx+1)}{16} - \frac{1}{16(cx-1)^2} + \frac{11}{16(cx-1)} + \frac{35 \ln(cx-1)}{16} \right)}{d^3} - \frac{b \left( \frac{105 \operatorname{arccosh}(cx)c^6}{16} \right)}{d^3} \right)$
parts	$-\frac{a \left( \frac{c^3}{16(cx+1)^2} + \frac{11c^3}{16(cx+1)} - \frac{35c^3 \ln(cx+1)}{16} + \frac{1}{3x^3} + \frac{3c^2}{x} - \frac{c^3}{16(cx-1)^2} + \frac{11c^3}{16(cx-1)} + \frac{35c^3 \ln(cx-1)}{16} \right)}{d^3} - \frac{bc^3 \left( \frac{105 \operatorname{arccosh}(cx)c^6}{16} \right)}{d^3}$

input `int((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^3,x,method=_RETURNVERBOSE)`

output `c^3*(-a/d^3*(1/3/c^3/x^3+3/c/x+1/16/(c*x+1)^2+11/16/(c*x+1)-35/16*ln(c*x+1)-1/16/(c*x-1)^2+11/16/(c*x-1)+35/16*ln(c*x-1))-b/d^3*(1/24*(105*arccosh(c*x)*c^6*x^6+29*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-175*c^4*x^4*arccosh(c*x)-27*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+56*c^2*x^2*arccosh(c*x)-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+8*arccosh(c*x))/(c^4*x^4-2*c^2*x^2+1)/c^3/x^3-19/3*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-35/8*dilog(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-35/8*dilog(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-35/8*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))`

$$3.54. \int \frac{a+b \operatorname{arccosh}(cx)}{x^4(d-c^2dx^2)^3} dx$$

**3.54.5 Fracas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3 x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^3,x, algorithm="fricas")`

output `integral(-(b*arccosh(c*x) + a)/(c^6*d^3*x^10 - 3*c^4*d^3*x^8 + 3*c^2*d^3*x^6 - d^3*x^4), x)`

**3.54.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**4/(-c**2*d*x**2+d)**3,x)`

output `Timed out`

**3.54.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arcosh}(cx) + a}{(c^2 dx^2 - d)^3 x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`



```

output 1/6144*(1935360*c^9*integrate(1/96*x^7*log(c*x - 1)/(c^6*d^3*x^8 - 3*c^4*d
^3*x^6 + 3*c^2*d^3*x^4 - d^3*x^2), x) - 1680*c^8*(2*(5*c^2*x^3 - 3*x)/(c^8
*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3) + 3*log(c*x + 1)/(c^5*d^3) - 3*log(c*x
- 1)/(c^5*d^3)) - 645120*c^8*integrate(1/96*x^6*log(c*x - 1)/(c^6*d^3*x^8
- 3*c^4*d^3*x^6 + 3*c^2*d^3*x^4 - d^3*x^2), x) + 630*(c*(2*(5*c^2*x^2 + 3
*c*x - 6)/(c^8*d^3*x^3 - c^7*d^3*x^2 - c^6*d^3*x + c^5*d^3) - 5*log(c*x +
1)/(c^5*d^3) + 5*log(c*x - 1)/(c^5*d^3)) + 16*(2*c^2*x^2 - 1)*log(c*x - 1)
/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3))*c^7 + 2800*c^6*(2*(c^2*x^3 + x)/
(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3) - log(c*x + 1)/(c^3*d^3) + log(c*x
- 1)/(c^3*d^3)) + 1290240*c^6*integrate(1/96*x^4*log(c*x - 1)/(c^6*d^3*x^
8 - 3*c^4*d^3*x^6 + 3*c^2*d^3*x^4 - d^3*x^2), x) + 315*(c*(2*(3*c^2*x^2 -
3*c*x - 2)/(c^6*d^3*x^3 - c^5*d^3*x^2 - c^4*d^3*x + c^3*d^3) - 3*log(c*x +
1)/(c^3*d^3) + 3*log(c*x - 1)/(c^3*d^3)) - 16*log(c*x - 1)/(c^6*d^3*x^4 -
2*c^4*d^3*x^2 + c^2*d^3))*c^5 + 896*c^4*(2*(3*c^2*x^3 - 5*x)/(c^4*d^3*x^4
- 2*c^2*d^3*x^2 + d^3) - 3*log(c*x + 1)/(c*d^3) + 3*log(c*x - 1)/(c*d^3))
- 645120*c^4*integrate(1/96*x^2*log(c*x - 1)/(c^6*d^3*x^8 - 3*c^4*d^3*x^6
+ 3*c^2*d^3*x^4 - d^3*x^2), x) + 128*c^2*(2*(15*c^4*x^4 - 25*c^2*x^2 + 8)
/(c^4*d^3*x^5 - 2*c^2*d^3*x^3 + d^3*x) - 15*c*log(c*x + 1)/d^3 + 15*c*log(
c*x - 1)/d^3) - 32*(105*(c^7*x^7 - 2*c^5*x^5 + c^3*x^3)*log(c*x + 1)^2 + 2
10*(c^7*x^7 - 2*c^5*x^5 + c^3*x^3)*log(c*x + 1)*log(c*x - 1) + 4*(210*c...

```

### 3.54.8 Giac [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx = \int -\frac{b \operatorname{arccosh}(cx) + a}{(c^2 dx^2 - d)^3 x^4} dx$$

```

input integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^3,x, algorithm="giac")

```

```

output integrate(-(b*arccosh(c*x) + a)/((c^2*d*x^2 - d)^3*x^4), x)

```

**3.54.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4 (d - c^2 dx^2)^3} dx$$

input `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^3), x)`output `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^3), x)`

### 3.55 $\int \frac{\operatorname{arccosh}(ax)}{c-a^2cx^2} dx$

3.55.1	Optimal result	626
3.55.2	Mathematica [A] (verified)	626
3.55.3	Rubi [C] (verified)	627
3.55.4	Maple [C] (verified)	629
3.55.5	Fricas [F]	629
3.55.6	Sympy [F]	629
3.55.7	Maxima [F]	630
3.55.8	Giac [F]	630
3.55.9	Mupad [F(-1)]	630

#### 3.55.1 Optimal result

Integrand size = 18, antiderivative size = 53

$$\int \frac{\operatorname{arccosh}(ax)}{c-a^2cx^2} dx = \frac{2\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac} + \frac{\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{ac} - \frac{\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{ac}$$

output `2*arccosh(a*x)*arctanh(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c+polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c-polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c`

#### 3.55.2 Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.45

$$\int \frac{\operatorname{arccosh}(ax)}{c-a^2cx^2} dx = -\frac{\operatorname{arccosh}(ax)\log(1-e^{\operatorname{arccosh}(ax)})}{ac} + \frac{\operatorname{arccosh}(ax)\log(1+e^{\operatorname{arccosh}(ax)})}{ac} + \frac{\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{ac} - \frac{\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{ac}$$

input `Integrate[ArcCosh[a*x]/(c - a^2*c*x^2), x]`

output `-((ArcCosh[a*x]*Log[1 - E^ArcCosh[a*x]])/(a*c)) + (ArcCosh[a*x]*Log[1 + E^ArcCosh[a*x]])/(a*c) + PolyLog[2, -E^ArcCosh[a*x]]/(a*c) - PolyLog[2, E^ArcCosh[a*x]]/(a*c)`

### 3.55.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)}{c - a^2 cx^2} dx \\
 & \quad \downarrow \text{6318} \\
 & \int \frac{\operatorname{arccosh}(ax)}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax) \\
 & \quad \downarrow \text{3042} \\
 & \int i\operatorname{arccosh}(ax) \csc(i\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax) \\
 & \quad \downarrow \text{26} \\
 & i \int \operatorname{arccosh}(ax) \csc(i\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax) \\
 & \quad \downarrow \text{4670} \\
 & \frac{i \int \log(1 - e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - i \int \log(1 + e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2i\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac} \\
 & \quad \downarrow \text{2715} \\
 & \frac{i \int e^{-\operatorname{arccosh}(ax)} \log(1 - e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - i \int e^{-\operatorname{arccosh}(ax)} \log(1 + e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} + 2i\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac} \\
 & \quad \downarrow \text{2838} \\
 & \frac{i(2\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{ac}
 \end{aligned}$$

input `Int[ArcCosh[a*x]/(c - a^2*c*x^2),x]`

---

3.55.  $\int \frac{\operatorname{arccosh}(ax)}{c - a^2 cx^2} dx$

output  $((-I)*((2*I)*\text{ArcCosh}[a*x]*\text{ArcTanh}[E^{\text{ArcCosh}[a*x]}] + I*\text{PolyLog}[2, -E^{\text{ArcCosh}[a*x]}] - I*\text{PolyLog}[2, E^{\text{ArcCosh}[a*x]}]))/(a*c)$

### 3.55.3.1 Defintions of rubi rules used

rule 26  $\text{Int}[(\text{Complex}[0, a_])*(F_x), x\_Symbol] \rightarrow \text{Simp}[(\text{Complex}[\text{Identity}[0], a]) \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ \text{EqQ}[a^2, 1]$

rule 2715  $\text{Int}[\text{Log}[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^{(n_)}], x\_Symbol] \rightarrow \text{Simp}[1/(d*e*n*\text{Log}[F]) \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x))})^n], x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x \ \&\& \ \text{GtQ}[a, 0]$

rule 2838  $\text{Int}[\text{Log}[(c_)*((d_) + (e_)*(x_)^{(n_)})]/(x_), x\_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n/n, x] /; \text{FreeQ}\{c, d, e, n\}, x \ \&\& \ \text{EqQ}[c*d, 1]$

rule 3042  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$

rule 4670  $\text{Int}[\text{csc}[(e_) + (\text{Complex}[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^{(m_)}, x\_Symbol] \rightarrow \text{Simp}[-2*(c + d*x)^m*(\text{ArcTanh}[E^{((-I)*e + f*fz*x)}]/(f*fz*I)), x] + (-\text{Simp}[d*(m/(f*fz*I)) \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 - E^{((-I)*e + f*fz*x)}], x], x] + \text{Simp}[d*(m/(f*fz*I)) \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + E^{((-I)*e + f*fz*x)}], x], x]) /; \text{FreeQ}\{c, d, e, f, fz\}, x \ \&\& \ \text{IGtQ}[m, 0]$

rule 6318  $\text{Int}[(a_) + \text{ArcCosh}[(c_)*(x_)]*(b_)]^{(n_)} / ((d_) + (e_)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[-(c*d)^{-1} \text{Subst}[\text{Int}[(a + b*x)^n*\text{Csch}[x], x], x, \text{ArcCosh}[c*x]], x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[n, 0]$

### 3.55.4 Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.96 (sec) , antiderivative size = 169, normalized size of antiderivative = 3.19

method	result
derivativedivides	$\frac{\operatorname{arctanh}(ax) \operatorname{arccosh}(ax)}{c} + \frac{2i \left( \operatorname{arctanh}(ax) \ln \left( 1 + \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) - \operatorname{arctanh}(ax) \ln \left( 1 - \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) + \operatorname{dilog} \left( 1 + \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) - \operatorname{dilog} \left( 1 - \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) \right)}{c(a^2x^2-1)}$
default	$\frac{\operatorname{arctanh}(ax) \operatorname{arccosh}(ax)}{c} + \frac{2i \left( \operatorname{arctanh}(ax) \ln \left( 1 + \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) - \operatorname{arctanh}(ax) \ln \left( 1 - \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) + \operatorname{dilog} \left( 1 + \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) - \operatorname{dilog} \left( 1 - \frac{i(ax+1)}{\sqrt{-a^2x^2+1}} \right) \right)}{a c(a^2x^2-1)}$

input `int(arccosh(a*x)/(-a^2*c*x^2+c),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{a} \left( \frac{1}{c} \operatorname{arctanh}(a*x) \operatorname{arccosh}(a*x) + 2 \frac{I}{c} \left( \operatorname{arctanh}(a*x) \ln \left( 1 + \frac{I(a*x+1)}{\sqrt{-a^2*x^2+1}} \right) - \operatorname{arctanh}(a*x) \ln \left( 1 - \frac{I(a*x+1)}{\sqrt{-a^2*x^2+1}} \right) + \operatorname{dilog} \left( 1 + \frac{I(a*x+1)}{\sqrt{-a^2*x^2+1}} \right) - \operatorname{dilog} \left( 1 - \frac{I(a*x+1)}{\sqrt{-a^2*x^2+1}} \right) \right) \right) \frac{1}{c(a^2*x^2-1)}$$

### 3.55.5 Fricas [F]

$$\int \frac{\operatorname{arccosh}(ax)}{c - a^2cx^2} dx = \int -\frac{\operatorname{arccosh}(ax)}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c),x, algorithm="fricas")`

output `integral(-arccosh(a*x)/(a^2*c*x^2 - c), x)`

### 3.55.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)}{c - a^2cx^2} dx = -\frac{\int \frac{\operatorname{acosh}(ax)}{a^2x^2-1} dx}{c}$$

input `integrate(acosh(a*x)/(-a**2*c*x**2+c),x)`

output `-Integral(acosh(a*x)/(a**2*x**2 - 1), x)/c`

---

3.55. 
$$\int \frac{\operatorname{arccosh}(ax)}{c - a^2cx^2} dx$$

**3.55.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{c - a^2cx^2} dx = \int -\frac{\operatorname{arcosh}(ax)}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c),x, algorithm="maxima")`

output `1/8*(4*(log(a*x + 1) - log(a*x - 1))*log(a*x + sqrt(a*x + 1))*sqrt(a*x - 1) - log(a*x + 1)^2 - 2*log(a*x + 1)*log(a*x - 1) + log(a*x - 1)^2)/(a*c) + 1/2*(log(a*x - 1)*log(1/2*a*x + 1/2) + dilog(-1/2*a*x + 1/2))/(a*c) + integrate(1/2*(log(a*x + 1) - log(a*x - 1))/(a^3*c*x^3 - a*c*x + (a^2*c*x^2 - c)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

**3.55.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{c - a^2cx^2} dx = \int -\frac{\operatorname{arcosh}(ax)}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c),x, algorithm="giac")`

output `integrate(-arccosh(a*x)/(a^2*c*x^2 - c), x)`

**3.55.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{c - a^2cx^2} dx = \int \frac{\operatorname{acosh}(ax)}{c - a^2cx^2} dx$$

input `int(acosh(a*x)/(c - a^2*c*x^2),x)`

output `int(acosh(a*x)/(c - a^2*c*x^2), x)`

### 3.56 $\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^2} dx$

3.56.1	Optimal result	631
3.56.2	Mathematica [A] (warning: unable to verify)	631
3.56.3	Rubi [C] (verified)	632
3.56.4	Maple [A] (verified)	635
3.56.5	Fricas [F]	635
3.56.6	Sympy [F]	635
3.56.7	Maxima [F]	636
3.56.8	Giac [F]	636
3.56.9	Mupad [F(-1)]	636

#### 3.56.1 Optimal result

Integrand size = 18, antiderivative size = 109

$$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^2} dx = -\frac{1}{2ac^2\sqrt{-1+ax}\sqrt{1+ax}} + \frac{x\operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} + \frac{\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac^2} + \frac{\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{2ac^2} - \frac{\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{2ac^2}$$

output  $\frac{1}{2}x\operatorname{arccosh}(ax)/c^2/(-a^2x^2+1)+\operatorname{arccosh}(ax)*\operatorname{arctanh}(ax+(ax-1)^{1/2}*(ax+1)^{1/2})/a/c^2+1/2*\operatorname{polylog}(2,-ax-(ax-1)^{1/2}*(ax+1)^{1/2})/a/c^2-1/2*\operatorname{polylog}(2,ax+(ax-1)^{1/2}*(ax+1)^{1/2})/a/c^2-1/2/a/c^2/(ax-1)^{1/2}/(ax+1)^{1/2}$

#### 3.56.2 Mathematica [A] (warning: unable to verify)

Time = 0.69 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^2} dx = \frac{2\left(\sqrt{\frac{-1+ax}{1+ax}}(1+ax)+\operatorname{arccosh}(ax)\left(ax+(-1+a^2x^2)\log\left(1-e^{\operatorname{arccosh}(ax)}\right)+(1-a^2x^2)\log\left(1+e^{\operatorname{arccosh}(ax)}\right)\right)\right)}{-1+a^2x^2} + 2\operatorname{PolyLog}\left(2, -e^{\operatorname{arccosh}(ax)}\right) + 2\operatorname{PolyLog}\left(2, e^{\operatorname{arccosh}(ax)}\right)$$



input `Integrate[ArcCosh[a*x]/(c - a^2*c*x^2)^2,x]`

output `((-2*(Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x) + ArcCosh[a*x]*(a*x + (-1 + a^2*x^2)*Log[1 - E^ArcCosh[a*x]] + (1 - a^2*x^2)*Log[1 + E^ArcCosh[a*x]])))/(-1 + a^2*x^2) + 2*PolyLog[2, -E^ArcCosh[a*x]] - 2*PolyLog[2, E^ArcCosh[a*x]])/(4*a*c^2)`

### 3.56.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.98, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6316, 27, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^2} dx \\
 & \quad \downarrow \text{6316} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)}{c(1-a^2x^2)} dx}{2c} + \frac{a \int \frac{x}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} + \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx}{2c^2} + \frac{a \int \frac{x}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} + \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} \\
 & \quad \downarrow \text{83} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx}{2c^2} + \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} - \frac{1}{2ac^2\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{6318} \\
 & -\frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} - \frac{1}{2ac^2\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\int i \operatorname{arccosh}(ax) \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} - \frac{1}{2ac^2\sqrt{ax-1}\sqrt{ax+1}}
 \end{aligned}$$

---

3.56.  $\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^2} dx$

$$\begin{aligned}
& \downarrow 26 \\
& \frac{i \int \operatorname{arccosh}(ax) \operatorname{csc}(i \operatorname{arccosh}(ax)) d \operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} - \frac{1}{2ac^2\sqrt{ax-1}\sqrt{ax+1}} \\
& \downarrow 4670 \\
& \frac{i(i \int \log(1 - e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - i \int \log(1 + e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}))}{2ac^2} \\
& \quad \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} - \frac{1}{2ac^2\sqrt{ax-1}\sqrt{ax+1}} \\
& \downarrow 2715 \\
& \frac{i(i \int e^{-\operatorname{arccosh}(ax)} \log(1 - e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - i \int e^{-\operatorname{arccosh}(ax)} \log(1 + e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} + 2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}))}{2ac^2} \\
& \quad \frac{x \operatorname{arccosh}(ax)}{2c^2(1-a^2x^2)} - \frac{1}{2ac^2\sqrt{ax-1}\sqrt{ax+1}} \\
& \downarrow 2838 \\
& \frac{i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{2ac^2} - \\
& \quad \frac{1}{2ac^2\sqrt{ax-1}\sqrt{ax+1}}
\end{aligned}$$

input `Int[ArcCosh[a*x]/(c - a^2*c*x^2)^2,x]`

output `-1/2*1/(a*c^2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + (x*ArcCosh[a*x])/(2*c^2*(1 - a^2*x^2)) - ((I/2)*((2*I)*ArcCosh[a*x]*ArcTanh[E^ArcCosh[a*x]] + I*PolyLog[2, -E^ArcCosh[a*x]] - I*PolyLog[2, E^ArcCosh[a*x]]))/(a*c^2)`

### 3.56.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

---

3.56.  $\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^2} dx$

- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`
- rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

### 3.56.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.48

method	result
derivativedivides	$\frac{-\frac{ax \operatorname{arccosh}(ax) + \sqrt{ax-1} \sqrt{ax+1}}{2(a^2x^2-1)c^2} - \frac{\operatorname{arccosh}(ax) \ln(1-ax-\sqrt{ax-1} \sqrt{ax+1})}{2c^2} - \frac{\operatorname{polylog}(2, ax + \sqrt{ax-1} \sqrt{ax+1})}{2c^2} + \frac{\operatorname{arccosh}(ax) \ln(1+ax)}{2c^2}}{a}$
default	$\frac{-\frac{ax \operatorname{arccosh}(ax) + \sqrt{ax-1} \sqrt{ax+1}}{2(a^2x^2-1)c^2} - \frac{\operatorname{arccosh}(ax) \ln(1-ax-\sqrt{ax-1} \sqrt{ax+1})}{2c^2} - \frac{\operatorname{polylog}(2, ax + \sqrt{ax-1} \sqrt{ax+1})}{2c^2} + \frac{\operatorname{arccosh}(ax) \ln(1+ax)}{2c^2}}{a}$

input `int(arccosh(a*x)/(-a^2*c*x^2+c)^2,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{a} \frac{-\frac{1}{2}(ax \operatorname{arccosh}(ax) + (ax-1)^{1/2}(ax+1)^{1/2})}{(a^2x^2-1)/c^2 - 1/2c^2 \operatorname{arccosh}(ax) \ln(1-ax-(ax-1)^{1/2}(ax+1)^{1/2}) - 1/2c^2 \operatorname{polylog}(2, ax + (ax-1)^{1/2}(ax+1)^{1/2}) + 1/2c^2 \operatorname{arccosh}(ax) \ln(1+ax) + (ax-1)^{1/2}(ax+1)^{1/2}} + 1/2c^2 \operatorname{polylog}(2, -ax - (ax-1)^{1/2}(ax+1)^{1/2})}$$

### 3.56.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arccosh}(ax)}{(a^2cx^2 - c)^2} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^2,x, algorithm="fricas")`

output `integral(arccosh(a*x)/(a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2), x)`

### 3.56.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{acosh}(ax)}{c^2(a^4x^4 - 2a^2x^2 + 1)} dx$$

input `integrate(acosh(a*x)/(-a**2*c*x**2+c)**2,x)`

output `Integral(acosh(a*x)/(a**4*x**4 - 2*a**2*x**2 + 1), x)/c**2`

---

3.56. 
$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^2} dx$$

**3.56.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arcosh}(ax)}{(a^2cx^2 - c)^2} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^2,x, algorithm="maxima")`

output `-1/16*((a^2*x^2 - 1)*log(a*x + 1)^2 + 2*(a^2*x^2 - 1)*log(a*x + 1)*log(a*x - 1) - (a^2*x^2 - 1)*log(a*x - 1)^2 + 4*a*x + 4*(2*a*x - (a^2*x^2 - 1)*log(a*x + 1) + (a^2*x^2 - 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1)) - 2*(a^2*x^2 - 1)*log(a*x - 1)/(a^3*c^2*x^2 - a*c^2) + 1/4*(log(a*x - 1)*log(1/2*a*x + 1/2) + dilog(-1/2*a*x + 1/2))/(a*c^2) - 1/8*log(a*x + 1)/(a*c^2) + integrate(-1/4*(2*a*x - (a^2*x^2 - 1)*log(a*x + 1) + (a^2*x^2 - 1)*log(a*x - 1))/(a^5*c^2*x^5 - 2*a^3*c^2*x^3 + a*c^2*x + (a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

**3.56.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arcosh}(ax)}{(a^2cx^2 - c)^2} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^2,x, algorithm="giac")`

output `integrate(arccosh(a*x)/(a^2*c*x^2 - c)^2, x)`

**3.56.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{acosh}(ax)}{(c - a^2cx^2)^2} dx$$

input `int(acosh(a*x)/(c - a^2*c*x^2)^2,x)`

output `int(acosh(a*x)/(c - a^2*c*x^2)^2, x)`

### 3.57 $\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^3} dx$

3.57.1	Optimal result	637
3.57.2	Mathematica [A] (warning: unable to verify)	637
3.57.3	Rubi [C] (verified)	638
3.57.4	Maple [A] (verified)	642
3.57.5	Fricas [F]	642
3.57.6	Sympy [F]	643
3.57.7	Maxima [F]	643
3.57.8	Giac [F]	644
3.57.9	Mupad [F(-1)]	644

#### 3.57.1 Optimal result

Integrand size = 18, antiderivative size = 164

$$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^3} dx = \frac{1}{12ac^3(-1+ax)^{3/2}(1+ax)^{3/2}} - \frac{3}{8ac^3\sqrt{-1+ax}\sqrt{1+ax}}$$

$$+ \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \frac{3x\operatorname{arccosh}(ax)}{8c^3(1-a^2x^2)} + \frac{3\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{4ac^3}$$

$$+ \frac{3\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{8ac^3} - \frac{3\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{8ac^3}$$

output `1/12/a/c^3/(a*x-1)^(3/2)/(a*x+1)^(3/2)+1/4*x*arccosh(a*x)/c^3/(-a^2*x^2+1)^(2+3/8*x*arccosh(a*x)/c^3/(-a^2*x^2+1)+3/4*arccosh(a*x)*arctanh(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^3+3/8*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^3-3/8*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^3-3/8/a/c^3/(a*x-1)^(1/2)/(a*x+1)^(1/2)`

#### 3.57.2 Mathematica [A] (warning: unable to verify)

Time = 1.79 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.36

$$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^3} dx$$

$$= \frac{-2(-2+ax)\sqrt{1+ax}}{(-1+ax)^{3/2}} + \frac{2\sqrt{-1+ax}(2+ax)}{(1+ax)^{3/2}} + \frac{6\operatorname{arccosh}(ax)}{(-1+ax)^2} - \frac{6\operatorname{arccosh}(ax)}{(1+ax)^2} + 18\left(-\frac{1}{\sqrt{\frac{-1+ax}{1+ax}}} + \frac{\operatorname{arccosh}(ax)}{1-ax}\right) + 18\left(\sqrt{\frac{-1+ax}{1+ax}}\right)$$

input `Integrate[ArcCosh[a*x]/(c - a^2*c*x^2)^3,x]`

output `((-2*(-2 + a*x)*Sqrt[1 + a*x])/(-1 + a*x)^(3/2) + (2*Sqrt[-1 + a*x]*(2 + a*x))/(1 + a*x)^(3/2) + (6*ArcCosh[a*x])/(-1 + a*x)^2 - (6*ArcCosh[a*x])/(1 + a*x)^2 + 18*(-(1/Sqrt[(-1 + a*x)/(1 + a*x)]) + ArcCosh[a*x]/(1 - a*x)) + 18*(Sqrt[(-1 + a*x)/(1 + a*x)] - ArcCosh[a*x]/(1 + a*x)) + 9*ArcCosh[a*x]*(ArcCosh[a*x] - 4*Log[1 - E^ArcCosh[a*x]]) - 9*ArcCosh[a*x]*(ArcCosh[a*x] - 4*Log[1 + E^ArcCosh[a*x]]) + 36*PolyLog[2, -E^ArcCosh[a*x]] - 36*PolyLog[2, E^ArcCosh[a*x]])/(96*a*c^3)`

### 3.57.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.70 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.96, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$ , Rules used = {6316, 27, 83, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^3} dx \\
 & \quad \downarrow \text{6316} \\
 & \frac{3 \int \frac{\operatorname{arccosh}(ax)}{c^2(1-a^2x^2)^2} dx}{4c} - \frac{a \int \frac{x}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4c^3} + \frac{x \operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{3 \int \frac{\operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{4c^3} - \frac{a \int \frac{x}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4c^3} + \frac{x \operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} \\
 & \quad \downarrow \text{83} \\
 & \frac{3 \int \frac{\operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{4c^3} + \frac{x \operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \frac{1}{12ac^3(ax-1)^{3/2}(ax+1)^{3/2}} \\
 & \quad \downarrow \text{6316}
 \end{aligned}$$

---

3.57.  $\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^3} dx$

$$\begin{aligned}
& \frac{3\left(\frac{1}{2} \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx + \frac{1}{2}a \int \frac{x}{(ax-1)^{3/2}(ax+1)^{3/2}} dx + \frac{x\operatorname{arccosh}(ax)}{2(1-a^2x^2)}\right)}{4c^3} + \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \\
& \qquad \frac{1}{12ac^3(ax-1)^{3/2}(ax+1)^{3/2}} \\
& \qquad \downarrow \text{83} \\
& \frac{3\left(\frac{1}{2} \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx + \frac{x\operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}}\right)}{4c^3} + \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \\
& \qquad \frac{1}{12ac^3(ax-1)^{3/2}(ax+1)^{3/2}} \\
& \qquad \downarrow \text{6318} \\
& \frac{3\left(\frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)}{2a} + \frac{x\operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}}\right)}{4c^3} + \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \\
& \qquad \frac{1}{12ac^3(ax-1)^{3/2}(ax+1)^{3/2}} \\
& \qquad \downarrow \text{3042} \\
& \frac{3\left(-\frac{\int i\operatorname{arccosh}(ax) \csc(i\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2a} + \frac{x\operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}}\right)}{4c^3} + \\
& \qquad \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \frac{1}{12ac^3(ax-1)^{3/2}(ax+1)^{3/2}} \\
& \qquad \downarrow \text{26} \\
& \frac{3\left(-\frac{i \int \operatorname{arccosh}(ax) \csc(i\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2a} + \frac{x\operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}}\right)}{4c^3} + \\
& \qquad \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \frac{1}{12ac^3(ax-1)^{3/2}(ax+1)^{3/2}} \\
& \qquad \downarrow \text{4670} \\
& \frac{3\left(-\frac{i\left(i \int \log(1-e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - i \int \log(1+e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2i\operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})\right)}{2a}\right)}{4c^3} + \frac{x\operatorname{arccosh}(ax)}{2(1-a^2x^2)^2} + \\
& \qquad \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)^2} + \frac{1}{12ac^3(ax-1)^{3/2}(ax+1)^{3/2}} \\
& \qquad \downarrow \text{2715}
\end{aligned}$$



$$3 \left( -\frac{i \int e^{-\operatorname{arccosh}(ax)} \log(1 - e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - i \int e^{-\operatorname{arccosh}(ax)} \log(1 + e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} + 2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{2a} \right)$$


---


$$\frac{x \operatorname{arccosh}(ax)}{4c^3 (1 - a^2 x^2)^2} + \frac{1}{12ac^3 (ax - 1)^{3/2} (ax + 1)^{3/2}}$$

↓ 2838

$$3 \left( \frac{x \operatorname{arccosh}(ax)}{2(1 - a^2 x^2)} - \frac{i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{2a} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}} \right)$$


---


$$\frac{x \operatorname{arccosh}(ax)}{4c^3 (1 - a^2 x^2)^2} + \frac{4c^3}{12ac^3 (ax - 1)^{3/2} (ax + 1)^{3/2}}$$

input `Int[ArcCosh[a*x]/(c - a^2*c*x^2)^3,x]`

output `1/(12*a*c^3*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2)) + (x*ArcCosh[a*x])/(4*c^3*(1 - a^2*x^2)^2) + (3*(-1/2*1/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + (x*ArcCosh[a*x])/(2*(1 - a^2*x^2)) - ((I/2)*((2*I)*ArcCosh[a*x]*ArcTanh[E^ArcCosh[a*x]]) + I*PolyLog[2, -E^ArcCosh[a*x]] - I*PolyLog[2, E^ArcCosh[a*x]]))/a)/(4*c^3)`

### 3.57.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x
_Symbol] :> Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p +
1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*
ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 +
c*x)^p*(-1 + c*x)^p) Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a
+ b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*
d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symb
ol] :> Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x
]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

### 3.57.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.25

method	result
derivativedivides	$\frac{-9a^3x^3 \operatorname{arccosh}(ax) + 9a^2x^2\sqrt{ax-1}\sqrt{ax+1} - 15ax \operatorname{arccosh}(ax) - 11\sqrt{ax-1}\sqrt{ax+1}}{24(a^4x^4 - 2a^2x^2 + 1)c^3} - \frac{3 \operatorname{arccosh}(ax) \ln(1-ax-\sqrt{ax-1}\sqrt{ax+1})}{8c^3} - \frac{3 \operatorname{polylog}(2, a*x + (a*x-1)^{1/2}*(a*x+1)^{1/2})}{a}$
default	$\frac{-9a^3x^3 \operatorname{arccosh}(ax) + 9a^2x^2\sqrt{ax-1}\sqrt{ax+1} - 15ax \operatorname{arccosh}(ax) - 11\sqrt{ax-1}\sqrt{ax+1}}{24(a^4x^4 - 2a^2x^2 + 1)c^3} - \frac{3 \operatorname{arccosh}(ax) \ln(1-ax-\sqrt{ax-1}\sqrt{ax+1})}{8c^3} - \frac{3 \operatorname{polylog}(2, -a*x - (a*x-1)^{1/2}*(a*x+1)^{1/2})}{a}$

input `int(arccosh(a*x)/(-a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)`

output `1/a*(-1/24*(9*a^3*x^3*arccosh(a*x)+9*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-15*a*x*arccosh(a*x)-11*(a*x-1)^(1/2)*(a*x+1)^(1/2))/(a^4*x^4-2*a^2*x^2+1)/c^3-3/8/c^3*arccosh(a*x)*ln(1-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-3/8/c^3*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/8/c^3*arccosh(a*x)*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/8/c^3*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2)))`

### 3.57.5 Fricas [F]

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arccosh}(ax)}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^3,x, algorithm="fricas")`

output `integral(-arccosh(a*x)/(a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3), x)`

## 3.57.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^3} dx = -\int \frac{\operatorname{acosh}(ax)}{a^6x^6 - 3a^4x^4 + 3a^2x^2 - 1} \frac{dx}{c^3}$$

input `integrate(acosh(a*x)/(-a**2*c*x**2+c)**3,x)`

output `-Integral(acosh(a*x)/(a**6*x**6 - 3*a**4*x**4 + 3*a**2*x**2 - 1), x)/c**3`

## 3.57.7 Maxima [F]

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arcosh}(ax)}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^3,x, algorithm="maxima")`

output `-1/64*(10*a^3*x^3 + 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1)^2 + 6*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1)*log(a*x - 1) - 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1)^2 - 14*a*x + 4*(6*a^3*x^3 - 10*a*x - 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1) + 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1)) - 7*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1))/(a^5*c^3*x^4 - 2*a^3*c^3*x^2 + a*c^3) + 3/16*(log(a*x - 1)*log(1/2*a*x + 1/2) + dilog(-1/2*a*x + 1/2))/(a*c^3) - 7/64*log(a*x + 1)/(a*c^3) + integrate(-1/16*(6*a^3*x^3 - 10*a*x - 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1) + 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1))/(a^7*c^3*x^7 - 3*a^5*c^3*x^5 + 3*a^3*c^3*x^3 - a*c^3*x + (a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

**3.57.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arcosh}(ax)}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^3,x, algorithm="giac")`

output `integrate(-arccosh(a*x)/(a^2*c*x^2 - c)^3, x)`

**3.57.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^3} dx = \int \frac{\operatorname{acosh}(ax)}{(c - a^2cx^2)^3} dx$$

input `int(acosh(a*x)/(c - a^2*c*x^2)^3,x)`

output `int(acosh(a*x)/(c - a^2*c*x^2)^3, x)`

### 3.58 $\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

3.58.1	Optimal result	645
3.58.2	Mathematica [A] (warning: unable to verify)	646
3.58.3	Rubi [A] (verified)	646
3.58.4	Maple [B] (verified)	649
3.58.5	Fricas [F]	650
3.58.6	Sympy [F]	651
3.58.7	Maxima [F]	651
3.58.8	Giac [F]	651
3.58.9	Mupad [F(-1)]	652

#### 3.58.1 Optimal result

Integrand size = 27, antiderivative size = 278

$$\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \frac{bx^2 \sqrt{d - c^2 dx^2}}{32c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{bx^4 \sqrt{d - c^2 dx^2}}{96c \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{bcx^6 \sqrt{d - c^2 dx^2}}{36 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{16c^4} - \frac{x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{24c^2} + \frac{1}{6} x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{32bc^5 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

output 
$$-1/16*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/c^4-1/24*x^3*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/c^2+1/6*x^5*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}+1/32*b*x^2*(-c^2*d*x^2+d)^{(1/2)}/c^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/96*b*x^4*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/36*b*c*x^6*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/32*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

### 3.58.2 Mathematica [A] (warning: unable to verify)

Time = 0.96 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.71

$$\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{48acx\sqrt{d - c^2 dx^2}(-3 - 2c^2 x^2 + 8c^4 x^4) - 144a\sqrt{d} \arctan\left(\frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right) + \frac{b\sqrt{d - c^2 dx^2}(-72\operatorname{arccosh}(cx)^2 + 18\cosh(2\operatorname{arccosh}(cx)) - 9\cosh(4\operatorname{arccosh}(cx)) - 2\cosh(6\operatorname{arccosh}(cx)) + 12\operatorname{ArcCosh}[cx](-3\sinh(2\operatorname{arccosh}(cx)) + 3\sinh(4\operatorname{arccosh}(cx)) + \sinh(6\operatorname{arccosh}(cx))))}{\sqrt{d}(-1 + c^2 x^2)}}}{(2304c^5)}$$

input `Integrate[x^4*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `(48*a*c*x*Sqrt[d - c^2*d*x^2]*(-3 - 2*c^2*x^2 + 8*c^4*x^4) - 144*a*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + (b*Sqrt[d - c^2*d*x^2]*(-72*ArcCosh[c*x]^2 + 18*Cosh[2*ArcCosh[c*x]] - 9*Cosh[4*ArcCosh[c*x]] - 2*Cosh[6*ArcCosh[c*x]] + 12*ArcCosh[c*x]*(-3*Sinh[2*ArcCosh[c*x]] + 3*Sinh[4*ArcCosh[c*x]] + Sinh[6*ArcCosh[c*x]])))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))/(2304*c^5)`

### 3.58.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.83, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$ , Rules used = {6341, 15, 6354, 15, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6341$$

$$-\frac{\sqrt{d - c^2 dx^2} \int \frac{x^4 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{6\sqrt{cx - 1} \sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x^5 dx}{6\sqrt{cx - 1} \sqrt{cx + 1}} + \frac{1}{6} x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))$$

$$\downarrow 15$$

$$-\frac{\sqrt{d - c^2 dx^2} \int \frac{x^4 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{6\sqrt{cx - 1} \sqrt{cx + 1}} + \frac{1}{6} x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^6 \sqrt{d - c^2 dx^2}}{36\sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow 6354$$

---

3.58.  $\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& \frac{\sqrt{d-c^2dx^2} \left( \frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} - \frac{b \int x^3 dx}{4c} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{\frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{bcx^6\sqrt{d-c^2dx^2}}{36\sqrt{cx-1}\sqrt{cx+1}}}{\phantom{6\sqrt{cx-1}\sqrt{cx+1}}} \\
& \quad \downarrow 15 \\
& \frac{\sqrt{d-c^2dx^2} \left( \frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} - \frac{bx^4}{16c} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{\frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{bcx^6\sqrt{d-c^2dx^2}}{36\sqrt{cx-1}\sqrt{cx+1}}}{\phantom{6\sqrt{cx-1}\sqrt{cx+1}}} \\
& \quad \downarrow 6354 \\
& \frac{\sqrt{d-c^2dx^2} \left( \frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \int x dx}{2c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} - \frac{bx^4}{16c} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{\frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{bcx^6\sqrt{d-c^2dx^2}}{36\sqrt{cx-1}\sqrt{cx+1}}}{\phantom{6\sqrt{cx-1}\sqrt{cx+1}}} \\
& \quad \downarrow 15 \\
& \frac{\sqrt{d-c^2dx^2} \left( \frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} - \frac{bx^2}{4c} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} - \frac{bx^4}{16c} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{\frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{bcx^6\sqrt{d-c^2dx^2}}{36\sqrt{cx-1}\sqrt{cx+1}}}{\phantom{6\sqrt{cx-1}\sqrt{cx+1}}} \\
& \quad \downarrow 6308 \\
& \frac{\frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \sqrt{d-c^2dx^2} \left( \frac{x^3\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} + \frac{3 \left( \frac{(a+b\operatorname{arccosh}(cx))^2}{4bc^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} - \frac{bx^2}{4c} \right)}{4c^2} - \frac{bx^4}{16c} \right)}{6\sqrt{cx-1}\sqrt{cx+1}}}{\frac{bcx^6\sqrt{d-c^2dx^2}}{36\sqrt{cx-1}\sqrt{cx+1}}}
\end{aligned}$$

---

3.58.  $\int x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) dx$



input `Int[x^4*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `-1/36*(b*c*x^6*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^5*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/6 - (Sqrt[d - c^2*d*x^2]*(-1/16*(b*x^4)/c + (x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(4*c^2) + (3*(-1/4*(b*x^2)/c + (x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])))/(2*c^2) + (a + b*ArcCosh[c*x])^2/(4*b*c^3)))/(4*c^2)))/(6*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.58.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)])*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_)*Sqrt[(d_) + (e_.)*(x_)^2]), x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.58.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 877 vs.  $2(234) = 468$ .

Time = 0.84 (sec) , antiderivative size = 878, normalized size of antiderivative = 3.16

method	result
default	$-\frac{ax^3(-c^2dx^2+d)^{\frac{3}{2}}}{6c^2d} - \frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{8c^4d} + \frac{ax\sqrt{-c^2dx^2+d}}{16c^4} + \frac{ad \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{16c^4\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)}{32\sqrt{cx-1}\sqrt{cx+1}c^5}\right)$
parts	$-\frac{ax^3(-c^2dx^2+d)^{\frac{3}{2}}}{6c^2d} - \frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{8c^4d} + \frac{ax\sqrt{-c^2dx^2+d}}{16c^4} + \frac{ad \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{16c^4\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)}{32\sqrt{cx-1}\sqrt{cx+1}c^5}\right)$

```
input int(x^4*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output 
$$\begin{aligned} & -1/6*a*x^3*(-c^2*d*x^2+d)^{(3/2)}/c^2/d-1/8*a/c^4*x*(-c^2*d*x^2+d)^{(3/2)}/d+1 \\ & /16*a/c^4*x*(-c^2*d*x^2+d)^{(1/2)}+1/16*a/c^4*d/(c^2*d)^{(1/2)}*\arctan((c^2*d) \\ & ^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)})+b*(-1/32*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)} \\ & /((c*x+1)^{(1/2)}/c^5*\operatorname{arccosh}(c*x)^2+1/2304*(-d*(c^2*x^2-1))^{(1/2)}*(32*c^7*x \\ & ^7-64*c^5*x^5+32*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^6*x^6+38*c^3*x^3-48*(c*x+1) \\ & )^{(1/2)}*(c*x-1)^{(1/2)}*c^4*x^4-6*c*x+18*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2 \\ & -(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(-1+6*\operatorname{arccosh}(c*x)))/(c*x+1)/c^5/(c*x-1)+1/51 \\ & 2*(-d*(c^2*x^2-1))^{(1/2)}*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)} \\ & )^{(1/2)}*c^4*x^4+4*c*x-8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+(c*x-1)^{(1/2)}*(c*x+ \\ & 1)^{(1/2)}*(-1+4*\operatorname{arccosh}(c*x)))/(c*x+1)/c^5/(c*x-1)-1/256*(-d*(c^2*x^2-1))^{(1/2)} \\ & *(2*c^3*x^3-2*c*x+2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-(c*x-1)^{(1/2)}* \\ & (c*x+1)^{(1/2)}*(-1+2*\operatorname{arccosh}(c*x)))/(c*x+1)/c^5/(c*x-1)-1/256*(-d*(c^2*x^2- \\ & 1))^{(1/2)}*(-2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+2*c^3*x^3+(c*x-1)^{(1/2)}* \\ & (c*x+1)^{(1/2)}-2*c*x)*(1+2*\operatorname{arccosh}(c*x)))/(c*x+1)/c^5/(c*x-1)+1/512*(-d*(c^2 \\ & *x^2-1))^{(1/2)}*(-8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^4*x^4+8*c^5*x^5+8*(c*x-1) \\ & )^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-12*c^3*x^3-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+4*c*x) \\ & *(1+4*\operatorname{arccosh}(c*x)))/(c*x+1)/c^5/(c*x-1)+1/2304*(-d*(c^2*x^2-1))^{(1/2)}*(-32 \\ & *(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^6*x^6+32*c^7*x^7+48*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)} \\ & )^{(1/2)}*c^4*x^4-64*c^5*x^5-18*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+38*c^3*x^3+ \\ & (c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}-6*c*x)*(1+6*\operatorname{arccosh}(c*x)))/(c*x+1)/c^5/(c*x-1) \dots \end{aligned}$$

### 3.58.5 Fracas [F]

$$\int x^4 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output `integral((b*x^4*arccosh(c*x) + a*x^4)*sqrt(-c^2*d*x^2 + d), x)`

**3.58.6 Sympy [F]**

$$\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int x^4 \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate(x**4*(a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**4*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x)), x)`

**3.58.7 Maxima [F]**

$$\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/48*(8*(-c^2*d*x^2 + d)^(3/2)*x^3/(c^2*d) - 3*sqrt(-c^2*d*x^2 + d)*x/c^4 + 6*(-c^2*d*x^2 + d)^(3/2)*x/(c^4*d) - 3*sqrt(d)*arcsin(c*x)/c^5)*a + b*integrate(sqrt(-c^2*d*x^2 + d)*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1), x)`

**3.58.8 Giac [F]**

$$\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)*x^4, x)`

**3.58.9 Mupad [F(-1)]**

Timed out.

$$\int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2} dx$$

input `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2),x)`output `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2), x)`

### 3.59 $\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

3.59.1	Optimal result	653
3.59.2	Mathematica [A] (verified)	654
3.59.3	Rubi [A] (verified)	654
3.59.4	Maple [B] (verified)	656
3.59.5	Fricas [F]	657
3.59.6	Sympy [F]	657
3.59.7	Maxima [F]	658
3.59.8	Giac [F]	658
3.59.9	Mupad [F(-1)]	658

#### 3.59.1 Optimal result

Integrand size = 27, antiderivative size = 201

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \frac{bx^2 \sqrt{d - c^2 dx^2}}{16c \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{bcx^4 \sqrt{d - c^2 dx^2}}{16 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{8c^2} + \frac{1}{4} x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{16bc^3 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

output

```
-1/8*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^2+1/4*x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)+1/16*b*x^2*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/16*b*c*x^4*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/16*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.59.2 Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.75

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \frac{-16acx(-1 + 2c^2x^2) \sqrt{d - c^2dx^2} + 16a\sqrt{d} \arctan\left(\frac{cx\sqrt{d-c^2dx^2}}{\sqrt{d}(-1+c^2x^2)}\right) + \frac{b\sqrt{d-c^2dx^2}(\operatorname{sarccosh}(cx)^2 + \cosh(4\operatorname{arccosh}(cx)))}{\sqrt{\frac{-1+cx}{1+cx}}}}{128c^3}$$

input `Integrate[x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `-1/128*(-16*a*c*x*(-1 + 2*c^2*x^2)*Sqrt[d - c^2*d*x^2] + 16*a*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + (b*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))/c^3`

### 3.59.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.88, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6341, 15, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6341} \\ & -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x^3 dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4} x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{15} \\ & -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4} x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^4 \sqrt{d - c^2 dx^2}}{16\sqrt{cx-1}\sqrt{cx+1}} \\ & \quad \downarrow \text{6354} \end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{d-c^2dx^2} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \int x dx}{2c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a + \\
& \quad \operatorname{arccosh}(cx)) - \frac{bcx^4\sqrt{d-c^2dx^2}}{16\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow 15 \\
& \frac{\sqrt{d-c^2dx^2} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} - \frac{bx^2}{4c} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a + \\
& \quad \operatorname{arccosh}(cx)) - \frac{bcx^4\sqrt{d-c^2dx^2}}{16\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow 6308 \\
& \frac{\frac{1}{4}x^3\sqrt{d-c^2dx^2}(a + \operatorname{arccosh}(cx)) - \sqrt{d-c^2dx^2} \left( \frac{(a+b\operatorname{arccosh}(cx))^2}{4bc^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} - \frac{bx^2}{4c} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bcx^4\sqrt{d-c^2dx^2}}{16\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `-1/16*(b*c*x^4*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/4 - (Sqrt[d - c^2*d*x^2]*(-1/4*(b*x^2)/c + (x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(2*c^2) + (a + b*ArcCosh[c*x])^2/(4*b*c^3)))/(4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.59.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)])*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`



```
rule 6341 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) +
(e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqr
t[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0]
&& IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])
```

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.59.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 366 vs. 2(169) = 338.

Time = 0.58 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.83

method	result
default	$-\frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{4c^2d} + \frac{ax\sqrt{-c^2dx^2+d}}{8c^2} + \frac{ad \arctan\left(\frac{\sqrt{c^2d}x}{\sqrt{-c^2dx^2+d}}\right)}{8c^2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2}{16\sqrt{cx-1}\sqrt{cx+1}c^3} + \frac{\sqrt{-d(c^2x^2-1)}}{16\sqrt{cx-1}\sqrt{cx+1}c^3}\right)$
parts	$-\frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{4c^2d} + \frac{ax\sqrt{-c^2dx^2+d}}{8c^2} + \frac{ad \arctan\left(\frac{\sqrt{c^2d}x}{\sqrt{-c^2dx^2+d}}\right)}{8c^2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2}{16\sqrt{cx-1}\sqrt{cx+1}c^3} + \frac{\sqrt{-d(c^2x^2-1)}}{16\sqrt{cx-1}\sqrt{cx+1}c^3}\right)$

```
input int(x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output 
$$\begin{aligned} & -1/4*a*x*(-c^2*d*x^2+d)^{(3/2)}/c^2/d+1/8*a/c^2*x*(-c^2*d*x^2+d)^{(1/2)}+1/8*a \\ & /c^2*d/(c^2*d)^{(1/2)}*\arctan((c^2*d)^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)})+b*(-1/16 \\ & *(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}/c^3*\operatorname{arccosh}(c*x)^2+1/2 \\ & 56*(-d*(c^2*x^2-1))^{(1/2)}*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)} \\ & *c^4*x^4+4*c*x-8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+(c*x-1)^{(1/2)}*(c*x \\ & +1)^{(1/2)}*(-1+4*\operatorname{arccosh}(c*x)))/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c^2*x^2-1))^{(1/2)} \\ & *(-8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^4*x^4+8*c^5*x^5+8*(c*x-1)^{(1/2)}*(c \\ & *x+1)^{(1/2)}*c^2*x^2-12*c^3*x^3-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+4*c*x)*(1+4*\operatorname{arc} \\ & \operatorname{cosh}(c*x))/(c*x+1)/c^3/(c*x-1) \end{aligned}$$

### 3.59.5 Fricas [F]

$$\int x^2 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a) x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*x^2*arccosh(c*x) + a*x^2), x)`

### 3.59.6 Sympy [F]

$$\int x^2 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int x^2 \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate(x**2*(a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**2*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x)), x)`

**3.59.7 Maxima [F]**

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a) x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `1/8*a*(sqrt(-c^2*d*x^2 + d)*x/c^2 - 2*(-c^2*d*x^2 + d)^(3/2)*x/(c^2*d) + sqrt(d)*arcsin(c*x)/c^3) + b*integrate(sqrt(-c^2*d*x^2 + d)*x^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)), x)`

**3.59.8 Giac [F]**

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a) x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)*x^2, x)`

**3.59.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2} dx$$

input `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2),x)`

output `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2), x)`

### 3.60 $\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

3.60.1	Optimal result	659
3.60.2	Mathematica [A] (warning: unable to verify)	659
3.60.3	Rubi [A] (verified)	660
3.60.4	Maple [B] (verified)	661
3.60.5	Fricas [F]	662
3.60.6	Sympy [F]	662
3.60.7	Maxima [F]	662
3.60.8	Giac [F(-2)]	663
3.60.9	Mupad [F(-1)]	663

#### 3.60.1 Optimal result

Integrand size = 24, antiderivative size = 124

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = -\frac{bcx^2 \sqrt{d - c^2 dx^2}}{4\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{4bc\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output  $\frac{1}{2}x(a+b\operatorname{arccosh}(cx))(-c^2d^2x^2+d)^{(1/2)} - \frac{1}{4}b^2cx^2(-c^2d^2x^2+d)^{(1/2)} / (cx-1)^{(1/2)} / (cx+1)^{(1/2)} - \frac{1}{4}(a+b\operatorname{arccosh}(cx))^2(-c^2d^2x^2+d)^{(1/2)} / b/c / (cx-1)^{(1/2)} / (cx+1)^{(1/2)}$

#### 3.60.2 Mathematica [A] (warning: unable to verify)

Time = 0.47 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.16

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \frac{1}{8} \left( 4ax\sqrt{d - c^2 dx^2} - \frac{4a\sqrt{d} \arctan\left(\frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right)}{c} - \frac{b\sqrt{d - c^2 dx^2} (2\operatorname{arccosh}(cx))^2 + \cosh(2\operatorname{arccosh}(cx)) - 2\operatorname{arccosh}(cx) \sinh(2\operatorname{arccosh}(cx))}{c\sqrt{\frac{-1+cx}{1+cx}}(1 + cx)} \right)$$

input `Integrate[Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output 
$$\frac{(4*a*x*\sqrt{d - c^2*d*x^2} - (4*a*\sqrt{d}*\text{ArcTan}[(c*x*\sqrt{d - c^2*d*x^2})/(\sqrt{d}*(-1 + c^2*x^2))])/c - (b*\sqrt{d - c^2*d*x^2}*(2*\text{ArcCosh}[c*x]^2 + \text{Cosh}[2*\text{ArcCosh}[c*x]] - 2*\text{ArcCosh}[c*x]*\text{Sinh}[2*\text{ArcCosh}[c*x])))/(c*\sqrt{(-1 + c*x)/(1 + c*x)}*(1 + c*x)))/8$$

### 3.60.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {6310, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx)) dx \\ & \quad \downarrow \text{6310} \\ & -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \text{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx)) \\ & \quad \downarrow \text{15} \\ & -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \text{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx)) - \frac{bcx^2\sqrt{d - c^2 dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \\ & \quad \downarrow \text{6308} \\ & \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx)) - \frac{\sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx))^2}{4bc\sqrt{cx-1}\sqrt{cx+1}} - \frac{bcx^2\sqrt{d - c^2 dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \end{aligned}$$

input `Int[Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output 
$$-1/4*(b*c*x^2*\sqrt{d - c^2*d*x^2})/(\sqrt{-1 + c*x}*\sqrt{1 + c*x}) + (x*\text{Sqrt}[d - c^2*d*x^2]*(a + b*\text{ArcCosh}[c*x])/2 - (\text{Sqrt}[d - c^2*d*x^2]*(a + b*\text{ArcCosh}[c*x])^2)/(4*b*c*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]))$$

## 3.60.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

## 3.60.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 277 vs.  $2(104) = 208$ .

Time = 0.75 (sec) , antiderivative size = 278, normalized size of antiderivative = 2.24

method	result
default	$\frac{ax\sqrt{-c^2dx^2+d}}{2} + \frac{ad \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2}{4\sqrt{cx-1}\sqrt{cx+1}c} + \frac{\sqrt{-d(c^2x^2-1)}(2c^3x^3-2cx+2\sqrt{cx-1}\sqrt{cx+1})}{16(cx-1)c}\right)$
parts	$\frac{ax\sqrt{-c^2dx^2+d}}{2} + \frac{ad \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2}{4\sqrt{cx-1}\sqrt{cx+1}c} + \frac{\sqrt{-d(c^2x^2-1)}(2c^3x^3-2cx+2\sqrt{cx-1}\sqrt{cx+1})}{16(cx-1)c}\right)$

input `int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output  $\frac{1}{2}ax(-c^2dx^2+d)^{1/2} + \frac{1}{2}ad(c^2d)^{1/2} \arctan\left(\frac{(c^2d)^{1/2}x}{(-c^2dx^2+d)^{1/2}}\right) + b\left(-\frac{1}{4}(-d(c^2x^2-1))^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}/c \operatorname{arccosh}(cx)^2 + \frac{1}{16}(-d(c^2x^2-1))^{1/2}(2c^3x^3-2cx+2)(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2 - (cx-1)^{1/2}(cx+1)^{1/2}\right) \cdot (-1+2\operatorname{arccosh}(cx))/(cx-1)/(cx+1)/c + \frac{1}{16}(-d(c^2x^2-1))^{1/2}(-2(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2 + 2c^3x^3 + (cx-1)^{1/2}(cx+1)^{1/2} - 2cx) \cdot (1+2\operatorname{arccosh}(cx))/(cx-1)/(cx+1)/c$

### 3.60.5 Fricas [F]

$$\int \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a) dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a), x)`

### 3.60.6 Sympy [F]

$$\int \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int \sqrt{-d(cx-1)(cx+1)} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x)), x)`

### 3.60.7 Maxima [F]

$$\int \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a) dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output  $\frac{1}{2}(\sqrt{-c^2dx^2+d}x + \sqrt{d}) \arcsin(cx)/c + a + b \int \sqrt{-c^2dx^2+d} \log(cx + \sqrt{cx+1}\sqrt{cx-1}), x$

**3.60.8 Giac [F(-2)]**

Exception generated.

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.60.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2} dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2),x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2), x)`



### 3.61 $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.61.1	Optimal result	664
3.61.2	Mathematica [A] (warning: unable to verify)	664
3.61.3	Rubi [A] (verified)	665
3.61.4	Maple [B] (verified)	666
3.61.5	Fricas [F]	667
3.61.6	Sympy [F]	667
3.61.7	Maxima [F]	668
3.61.8	Giac [F(-2)]	668
3.61.9	Mupad [F(-1)]	668

#### 3.61.1 Optimal result

Integrand size = 27, antiderivative size = 118

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^2} dx = -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x} + \frac{c\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2b\sqrt{-1+cx}\sqrt{1+cx}} + \frac{bc\sqrt{d-c^2dx^2}\log(x)}{\sqrt{-1+cx}\sqrt{1+cx}}$$

output `-(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x+1/2*c*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*c*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)`

#### 3.61.2 Mathematica [A] (warning: unable to verify)

Time = 0.42 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.16

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^2} dx = -\frac{a\sqrt{d-c^2dx^2}}{x} + ac\sqrt{d}\arctan\left(\frac{cx\sqrt{d-c^2dx^2}}{\sqrt{d}(-1+c^2x^2)}\right) + \frac{1}{2}bc\sqrt{d-c^2dx^2}\left(-\frac{2\operatorname{arccosh}(cx)}{cx} + \frac{\operatorname{arccosh}(cx)^2 + 2\log(cx)}{\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}\right)$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^2,x]`

output `-((a*Sqrt[d - c^2*d*x^2])/x) + a*c*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + (b*c*Sqrt[d - c^2*d*x^2]*((-2*ArcCosh[c*x])/(c*x) + (ArcCosh[c*x]^2 + 2*Log[c*x])/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))))/2`

### 3.61.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6339, 14, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^2} dx$$

↓ 6339

$$\frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{\sqrt{cx - 1} \sqrt{cx + 1}} + \frac{bc \sqrt{d - c^2 dx^2} \int \frac{1}{x} dx}{\sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x}$$

↓ 14

$$\frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{\sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x} + \frac{bc \log(x) \sqrt{d - c^2 dx^2}}{\sqrt{cx - 1} \sqrt{cx + 1}}$$

↓ 6308

$$\frac{c \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{2b \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x} + \frac{bc \log(x) \sqrt{d - c^2 dx^2}}{\sqrt{cx - 1} \sqrt{cx + 1}}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^2,x]`

output `-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x) + (c*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (b*c*Sqrt[d - c^2*d*x^2]*Log[x])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.61.3.1 Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6339 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`

## 3.61.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 285 vs.  $2(102) = 204$ .

Time = 0.94 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.42

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{3}{2}}}{dx} - ac^2x\sqrt{-c^2dx^2+d} - \frac{ac^2d \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{\sqrt{c^2d}} + \frac{b\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2c}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{b\sqrt{-d(c^2x^2-1)}}{\sqrt{c}}$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{3}{2}}}{dx} - ac^2x\sqrt{-c^2dx^2+d} - \frac{ac^2d \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{\sqrt{c^2d}} + \frac{b\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2c}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{b\sqrt{-d(c^2x^2-1)}}{\sqrt{c}}$

input `int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

---

3.61. 
$$\int \frac{\sqrt{d-c^2x^2}(a+b\operatorname{arccosh}(cx))}{x^2} dx$$

output 
$$-a/d/x*(-c^2*d*x^2+d)^{(3/2)}-a*c^2*x*(-c^2*d*x^2+d)^{(1/2)}-a*c^2*d/(c^2*d)^{(1/2)}*\arctan((c^2*d)^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)})+1/2*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*\operatorname{arccosh}(c*x)^2*c-b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*\operatorname{arccosh}(c*x)*c-b*(-d*(c^2*x^2-1))^{(1/2)}*\operatorname{arccosh}(c*x)*x/(c*x-1)/(c*x+1)*c^2+b*(-d*(c^2*x^2-1))^{(1/2)}*\operatorname{arccosh}(c*x)/x/(c*x-1)/(c*x+1)+b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*c$$

### 3.61.5 Fricas [F]

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^2} dx = \int \frac{\sqrt{-c^2dx^2+d}(b\operatorname{arccosh}(cx)+a)}{x^2} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/x^2, x)`

### 3.61.6 Sympy [F]

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^2} dx = \int \frac{\sqrt{-d(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))}{x^2} dx$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2)/x**2,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))/x**2, x)`

**3.61.7 Maxima [F]**

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{\sqrt{-c^2 dx^2 + d}(b \operatorname{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="maxima")`

output `-(c*sqrt(d)*arcsin(c*x) + sqrt(-c^2*d*x^2 + d)/x)*a + b*integrate(sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x^2, x)`

**3.61.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.61.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^2,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^2, x)`

### 3.62 $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^4} dx$

3.62.1	Optimal result	669
3.62.2	Mathematica [A] (verified)	669
3.62.3	Rubi [A] (verified)	670
3.62.4	Maple [A] (verified)	671
3.62.5	Fricas [B] (verification not implemented)	672
3.62.6	Sympy [F]	673
3.62.7	Maxima [C] (verification not implemented)	673
3.62.8	Giac [F(-2)]	674
3.62.9	Mupad [F(-1)]	674

#### 3.62.1 Optimal result

Integrand size = 27, antiderivative size = 119

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^4} dx = -\frac{bc\sqrt{d-c^2dx^2}}{6x^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{3dx^3} - \frac{bc^3\sqrt{d-c^2dx^2}\log(x)}{3\sqrt{-1+cx}\sqrt{1+cx}}$$

output 
$$-1/3*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))/d/x^3-1/6*b*c*(-c^2*d*x^2+d)^{(1/2)}/x^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/3*b*c^3*\ln(x)*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

#### 3.62.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.74

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^4} dx = \frac{\sqrt{d-c^2dx^2}\left(\frac{(-1+cx)^{3/2}(1+cx)^{3/2}(a+b\operatorname{arccosh}(cx))}{3x^3} - \frac{1}{3}bc\left(\frac{1}{2x^2} + c^2\log(x)\right)\right)}{\sqrt{-1+cx}\sqrt{1+cx}}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^4,x]`

output `(Sqrt[d - c^2*d*x^2]*((( -1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(a + b*ArcCosh[c*x])))/(3*x^3) - (b*c*(1/(2*x^2) + c^2*Log[x]))/3)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.62.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.74, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6332, 25, 82, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^4} dx \\
 & \quad \downarrow \text{6332} \\
 & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{(1-cx)(cx+1)}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3dx^3} \\
 & \quad \downarrow \text{25} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3dx^3} \\
 & \quad \downarrow \text{82} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \frac{1-c^2 x^2}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3dx^3} \\
 & \quad \downarrow \text{244} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \left(\frac{1}{x^3} - \frac{c^2}{x}\right) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3dx^3} \\
 & \quad \downarrow \text{2009} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \left(c^2(-\log(x)) - \frac{1}{2x^2}\right)}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3dx^3}
 \end{aligned}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^4,x]`

---

3.62.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^4} dx$

output 
$$-1/3*((d - c^2*d*x^2)^{(3/2)}*(a + b*ArcCosh[c*x]))/(d*x^3) + (b*c*Sqrt[d - c^2*d*x^2]*(-1/2*1/x^2 - c^2*Log[x]))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])$$

### 3.62.3.1 Defintions of rubi rules used

rule 25 
$$\text{Int}[-(F_x), x\_Symbol] \text{ :> } \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$$

rule 82 
$$\text{Int}[(a + (b \cdot x)^m) \cdot ((c + (d \cdot x)^n) \cdot (e + (f \cdot x)^p))]^p, x] \text{ :> } \text{Int}[(a \cdot c + b \cdot d \cdot x^{2m})^m \cdot (e + f \cdot x)^p, x] \text{ ; FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{EqQ}[b \cdot c + a \cdot d, 0] \ \&\& \ \text{EqQ}[n, m] \ \&\& \ \text{IntegerQ}[m]$$

rule 244 
$$\text{Int}[(c \cdot x)^m \cdot (a + (b \cdot x)^2)^p, x\_Symbol] \text{ :> } \text{Int}[\text{Expand} \text{Integrand}[(c \cdot x)^m \cdot (a + b \cdot x^2)^p, x], x] \text{ ; FreeQ}\{a, b, c, m\}, x] \ \&\& \ \text{IGtQ}[p, 0]$$

rule 2009 
$$\text{Int}[u, x\_Symbol] \text{ :> } \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

rule 6332 
$$\text{Int}[(a + \text{ArcCosh}[c \cdot x]) \cdot (b \cdot x)^n \cdot (f \cdot x)^m \cdot (d + (e \cdot x)^2)^p, x\_Symbol] \text{ :> } \text{Simp}[(f \cdot x)^{m+1} \cdot (d + e \cdot x^2)^{p+1} \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^n / (d \cdot f \cdot (m+1)), x] + \text{Simp}[b \cdot c \cdot (n / (f \cdot (m+1))) \cdot \text{Simp}[(d + e \cdot x^2)^p / ((1 + c \cdot x)^p \cdot (-1 + c \cdot x)^p)] \quad \text{Int}[(f \cdot x)^{m+1} \cdot (1 + c \cdot x)^{p+1/2} \cdot (-1 + c \cdot x)^{p+1/2} \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^{n-1}, x], x] \text{ ; FreeQ}\{a, b, c, d, e, f, m, p\}, x] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{EqQ}[m + 2 \cdot p + 3, 0] \ \&\& \ \text{NeQ}[m, -1]$$

### 3.62.4 Maple [A] (verified)

Time = 1.18 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.29

method	result
default	$-\frac{a(-c^2 d x^2 + d)^{\frac{3}{2}}}{3 d x^3} + \frac{b \sqrt{-d(c^2 x^2 - 1)} \left( 2 \sqrt{c x + 1} \operatorname{arccosh}(c x) \sqrt{c x - 1} c^2 x^2 + 2 c^3 x^3 \operatorname{arccosh}(c x) - 2 \ln \left( 1 + (c x + \sqrt{c x - 1} \sqrt{c x + 1})^2 \right) \right) x^3 c}{6 \sqrt{c x - 1} \sqrt{c x + 1} x^3}$
parts	$-\frac{a(-c^2 d x^2 + d)^{\frac{3}{2}}}{3 d x^3} + \frac{b \sqrt{-d(c^2 x^2 - 1)} \left( 2 \sqrt{c x + 1} \operatorname{arccosh}(c x) \sqrt{c x - 1} c^2 x^2 + 2 c^3 x^3 \operatorname{arccosh}(c x) - 2 \ln \left( 1 + (c x + \sqrt{c x - 1} \sqrt{c x + 1})^2 \right) \right) x^3 c}{6 \sqrt{c x - 1} \sqrt{c x + 1} x^3}$

3.62. 
$$\int \frac{\sqrt{d - c^2 x^2} (a + b \operatorname{arccosh}(c x))}{x^4} dx$$



input `int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^4,x,method=_RETURNVERBOSE)`

output `-1/3*a/d/x^3*(-c^2*d*x^2+d)^(3/2)+1/6*b*(-d*(c^2*x^2-1))^(1/2)*(2*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2+2*c^3*x^3*arccosh(c*x)-2*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3-2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-c*x)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^3`

### 3.62.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 220 vs. 2(99) = 198.

Time = 0.29 (sec) , antiderivative size = 462, normalized size of antiderivative = 3.88

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^4} dx$$

$$= \frac{\begin{aligned} & 2(bc^4 x^4 - 2bc^2 x^2 + b)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1}) + (bc^5 x^5 - bc^3 x^3)\sqrt{-d} \log\left(\frac{c^2 dx^6 + c^2 dx^2 - dx^4 + \sqrt{d}}{6(c^2 x^5 - x^5)}\right) \\ & - 2(bc^5 x^5 - bc^3 x^3)\sqrt{d} \arctan\left(\frac{\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1}(x^2 + 1)\sqrt{d}}{c^2 dx^4 - (c^2 + 1)dx^2 + d}\right) - 2(bc^4 x^4 - 2bc^2 x^2 + b)\sqrt{-c^2 dx^2 + d} \log(cx - \sqrt{c^2 x^2 - 1}) \end{aligned}}{6(c^2 x^5 - x^5)}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^4,x,algorithm="fracas")`

output `[1/6*(2*(b*c^4*x^4 - 2*b*c^2*x^2 + b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + (b*c^5*x^5 - b*c^3*x^3)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + sqrt(-c^2*d*x^2 + d)*(b*c*x^3 - b*c*x)*sqrt(c^2*x^2 - 1) + 2*(a*c^4*x^4 - 2*a*c^2*x^2 + a)*sqrt(-c^2*d*x^2 + d))/(c^2*x^5 - x^3), -1/6*(2*(b*c^5*x^5 - b*c^3*x^3)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 2*(b*c^4*x^4 - 2*b*c^2*x^2 + b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - sqrt(-c^2*d*x^2 + d)*(b*c*x^3 - b*c*x)*sqrt(c^2*x^2 - 1) - 2*(a*c^4*x^4 - 2*a*c^2*x^2 + a)*sqrt(-c^2*d*x^2 + d))/(c^2*x^5 - x^3)]`

### 3.62.6 Sympy [F]

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)}(a + b \operatorname{acosh}(cx))}{x^4} dx$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2)/x**4,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))/x**4, x)`

### 3.62.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.26

$$\begin{aligned} & \int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^4} dx \\ &= \frac{\left( c^4 d^2 \sqrt{-\frac{1}{c^4 d}} \log\left(x^2 - \frac{1}{c^2}\right) + i(-1)^{-2c^2 dx^2 + 2d} c^2 d^{\frac{3}{2}} \log\left(-2c^2 d + \frac{2d}{x^2}\right) + \frac{\sqrt{-c^4 dx^4 + 2c^2 dx^2 - dd}}{x^2} \right) bc}{6d} \\ & \quad - \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} b \operatorname{arccosh}(cx)}{3 dx^3} - \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} a}{3 dx^3} \end{aligned}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^4,x, algorithm="maxima")`

output `1/6*(c^4*d^2*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2) + I*(-1)^(-2*c^2*d*x^2 + 2*d)*c^2*d^(3/2)*log(-2*c^2*d + 2*d/x^2) + sqrt(-c^4*d*x^4 + 2*c^2*d*x^2 - d)*d/x^2)*b*c/d - 1/3*(-c^2*d*x^2 + d)^(3/2)*b*arccosh(c*x)/(d*x^3) - 1/3*(-c^2*d*x^2 + d)^(3/2)*a/(d*x^3)`

**3.62.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^4,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.62.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2}}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^4,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^4, x)`

### 3.63 $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^6} dx$

3.63.1	Optimal result	675
3.63.2	Mathematica [A] (verified)	676
3.63.3	Rubi [A] (verified)	676
3.63.4	Maple [B] (verified)	678
3.63.5	Fricas [A] (verification not implemented)	679
3.63.6	Sympy [F]	679
3.63.7	Maxima [A] (verification not implemented)	680
3.63.8	Giac [F(-2)]	680
3.63.9	Mupad [F(-1)]	681

#### 3.63.1 Optimal result

Integrand size = 27, antiderivative size = 199

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^6} dx = -\frac{bc\sqrt{d-c^2dx^2}}{20x^4\sqrt{-1+cx}\sqrt{1+cx}} + \frac{bc^3\sqrt{d-c^2dx^2}}{30x^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{5dx^5} - \frac{2c^2(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{15dx^3} - \frac{2bc^5\sqrt{d-c^2dx^2}\log(x)}{15\sqrt{-1+cx}\sqrt{1+cx}}$$

output `-1/5*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/d/x^5-2/15*c^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/d/x^3-1/20*b*c*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/30*b*c^3*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/15*b*c^5*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)`

### 3.63.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.64

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^6} dx$$

$$= \frac{\sqrt{d - c^2 dx^2} (12(-1 + cx)^{3/2} (1 + cx)^{3/2} (a + \operatorname{barccosh}(cx)) + 8c^2 x^2 (-1 + cx)^{3/2} (1 + cx)^{3/2} (a + \operatorname{barccosh}(cx)) - 8c^2 x^2 (3 - 2c^2 x^2 + 8c^4 x^4 \operatorname{Log}[x]))}{60x^5 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^6,x]`

output `(Sqrt[d - c^2*d*x^2]*(12*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(a + b*ArcCosh[c*x]) + 8*c^2*x^2*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(a + b*ArcCosh[c*x]) - b*c*x*(3 - 2*c^2*x^2 + 8*c^4*x^4*Log[x]))/(60*x^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.63.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.67, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 1433, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^6} dx$$

$$\downarrow \text{6337}$$

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{-2c^4 x^4 - c^2 x^2 + 3}{15x^5} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{5dx^5} - \frac{2c^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{15dx^3}$$

$$\downarrow \text{27}$$

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{-2c^4 x^4 - c^2 x^2 + 3}{x^5} dx}{15\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{5dx^5} - \frac{2c^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{15dx^3}$$

$$\downarrow \text{1433}$$

---

3.63.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^6} dx$

$$\frac{bc\sqrt{d-c^2dx^2} \int \left(-\frac{2c^4}{x} - \frac{c^2}{x^3} + \frac{3}{x^5}\right) dx}{15\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{5dx^5} - \frac{2c^2(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{15dx^3}$$

↓ 2009

$$-\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{5dx^5} - \frac{2c^2(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{15dx^3} + \frac{bc\sqrt{d-c^2dx^2}\left(-2c^4\log(x) + \frac{c^2}{2x^2} - \frac{3}{4x^4}\right)}{15\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^6,x]`

output `-1/5*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(d*x^5) - (2*c^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(15*d*x^3) + (b*c*Sqrt[d - c^2*d*x^2]*(-3/(4*x^4) + c^2/(2*x^2) - 2*c^4*Log[x]))/(15*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.63.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1433 `Int[((d_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d*x)^m*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[p, 0] && (EqQ[p, 1] || !IntegerQ[(m + 1)/2])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.63.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1741 vs.  $2(167) = 334$ .

Time = 1.14 (sec) , antiderivative size = 1742, normalized size of antiderivative = 8.75

method	result	size
default	Expression too large to display	1742
parts	Expression too large to display	1742

```
input int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^6,x,method=_RETURNVERBOSE)
```

```
output -9/20*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c+1/2*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)*x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c^9-11/12*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)*x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c^7+21/20*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c^3+1/6*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)*x^5/(c*x-1)/(c*x+1)*c^10-3/5*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)*x^3/(c*x-1)/(c*x+1)*c^8+3/10*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)*x/(c*x-1)/(c*x+1)*c^6+9/5*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)/x^5/(c*x-1)/(c*x+1)*arccosh(c*x)-2/15*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)*x^9/(c*x-1)/(c*x+1)*c^14+4/15*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*c^5-2/15*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*c^5-1/4*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c^5+a*(-1/5/d/x^5*(-c^2*d*x^2+d)^(3/2)-2/15*c^2/d/x^3*(-c^2*d*x^2+d)^(3/2))+4/15*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)*x^7/(c*x-1)/(c*x+1)*c^12-6/5*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*c^5-3/10*b*(-d*(c^2*x^2-1))^(1/2)/(15*c^6*x^6-5*c^4*x^4-15*c^2*x^2+9...
```

### 3.63.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 549, normalized size of antiderivative = 2.76

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^6} dx$$

$$= \frac{4(2bc^6x^6 - bc^4x^4 - 4bc^2x^2 + 3b)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) + 4(bc^7x^7 - bc^5x^5)\sqrt{-d} \log\left(\frac{c^2dx^6}{\sqrt{-c^2dx^2 + d} - d}\right) - 8(bc^7x^7 - bc^5x^5)\sqrt{d} \arctan\left(\frac{\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}(x^2 + 1)\sqrt{d}}{c^2dx^4 - (c^2 + 1)dx^2 + d}\right) - 4(2bc^6x^6 - bc^4x^4 - 4bc^2x^2 + 3b)\sqrt{-c^2dx^2 + d}}{c^2dx^4 - (c^2 + 1)dx^2 + d}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^6,x, algorithm="fracas")`

output `[1/60*(4*(2*b*c^6*x^6 - b*c^4*x^4 - 4*b*c^2*x^2 + 3*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + 4*(b*c^7*x^7 - b*c^5*x^5)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + (2*b*c^3*x^3 - (2*b*c^3 - 3*b*c)*x^5 - 3*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 4*(2*a*c^6*x^6 - a*c^4*x^4 - 4*a*c^2*x^2 + 3*a)*sqrt(-c^2*d*x^2 + d))/(c^2*x^7 - x^5), -1/60*(8*(b*c^7*x^7 - b*c^5*x^5)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 4*(2*b*c^6*x^6 - b*c^4*x^4 - 4*b*c^2*x^2 + 3*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (2*b*c^3*x^3 - (2*b*c^3 - 3*b*c)*x^5 - 3*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 4*(2*a*c^6*x^6 - a*c^4*x^4 - 4*a*c^2*x^2 + 3*a)*sqrt(-c^2*d*x^2 + d))/(c^2*x^7 - x^5)]`

### 3.63.6 SymPy [F]

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^6} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)}(a + b \operatorname{acosh}(cx))}{x^6} dx$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2)/x**6,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))/x**6, x)`

---

3.63.  $\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^6} dx$



**3.63.7 Maxima [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.73

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^6} dx$$

$$= -\frac{1}{60} \left( 8c^4 \sqrt{-d} \log(x) - \frac{2c^2 \sqrt{-dx^2} - 3\sqrt{-d}}{x^4} \right) bc$$

$$- \frac{1}{15} b \left( \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}} c^2}{dx^3} + \frac{3(-c^2 dx^2 + d)^{\frac{3}{2}}}{dx^5} \right) \operatorname{arccosh}(cx)$$

$$- \frac{1}{15} a \left( \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}} c^2}{dx^3} + \frac{3(-c^2 dx^2 + d)^{\frac{3}{2}}}{dx^5} \right)$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^6,x, algorithm="maxima")`

output `-1/60*(8*c^4*sqrt(-d)*log(x) - (2*c^2*sqrt(-d)*x^2 - 3*sqrt(-d))/x^4)*b*c - 1/15*b*(2*(-c^2*d*x^2 + d)^(3/2)*c^2/(d*x^3) + 3*(-c^2*d*x^2 + d)^(3/2)/(d*x^5))*arccosh(c*x) - 1/15*a*(2*(-c^2*d*x^2 + d)^(3/2)*c^2/(d*x^3) + 3*(-c^2*d*x^2 + d)^(3/2)/(d*x^5))`

**3.63.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^6} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^6,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.63.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^6} dx = \int \frac{(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2}}{x^6} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^6, x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^6, x)`

### 3.64 $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^8} dx$

3.64.1	Optimal result	682
3.64.2	Mathematica [A] (verified)	683
3.64.3	Rubi [A] (verified)	683
3.64.4	Maple [B] (verified)	685
3.64.5	Fricas [A] (verification not implemented)	686
3.64.6	Sympy [F(-1)]	686
3.64.7	Maxima [A] (verification not implemented)	687
3.64.8	Giac [F(-2)]	687
3.64.9	Mupad [F(-1)]	688

#### 3.64.1 Optimal result

Integrand size = 27, antiderivative size = 279

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^8} dx = -\frac{bc\sqrt{d-c^2dx^2}}{42x^6\sqrt{-1+cx}\sqrt{1+cx}} + \frac{bc^3\sqrt{d-c^2dx^2}}{140x^4\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2bc^5\sqrt{d-c^2dx^2}}{105x^2\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{7dx^7}$$

$$- \frac{4c^2(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{35dx^5}$$

$$- \frac{8c^4(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{105dx^3}$$

$$- \frac{8bc^7\sqrt{d-c^2dx^2}\log(x)}{105\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/7*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/d/x^7-4/35*c^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/d/x^5-8/105*c^4*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/d/x^3-1/42*b*c*(-c^2*d*x^2+d)^(1/2)/x^6/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/140*b*c^3*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/105*b*c^5*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-8/105*b*c^7*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.64.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.52

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^8} dx$$

$$= \frac{\sqrt{d - c^2 dx^2} (60(-1 + cx)^{3/2} (1 + cx)^{3/2} (a + \operatorname{barccosh}(cx)) + 16c^2 x^2 (-1 + cx)^{3/2} (1 + cx)^{3/2} (3 + 2c^2 x^2) (a + \operatorname{barccosh}(cx)) - bc^2 x^2 (10 - 3c^2 x^2 - 8c^4 x^4 + 32c^6 x^6 \operatorname{Log}[x]))}{420x^7 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^8,x]`

output `(Sqrt[d - c^2*d*x^2]*(60*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(a + b*ArcCosh[c*x]) + 16*c^2*x^2*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(3 + 2*c^2*x^2)*(a + b*ArcCosh[c*x]) - b*c*x*(10 - 3*c^2*x^2 - 8*c^4*x^4 + 32*c^6*x^6*Log[x]))/(420*x^7*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.64.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.64, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 2010, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^8} dx$$

$$\downarrow \text{6337}$$

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{-8c^6 x^6 - 4c^4 x^4 - 3c^2 x^2 + 15}{105x^7} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{7dx^7} - \frac{4c^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{35dx^5} - \frac{8c^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{105dx^3}$$

$$\downarrow \text{27}$$

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{-8c^6 x^6 - 4c^4 x^4 - 3c^2 x^2 + 15}{x^7} dx}{105\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{7dx^7} - \frac{4c^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{35dx^5} - \frac{8c^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{105dx^3}$$

$$\downarrow \text{2010}$$

---

3.64.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^8} dx$

$$\frac{bc\sqrt{d-c^2dx^2} \int \left(-\frac{8c^6}{x} - \frac{4c^4}{x^3} - \frac{3c^2}{x^5} + \frac{15}{x^7}\right) dx}{105\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{7dx^7} - \frac{4c^2(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{35dx^5} - \frac{8c^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{105dx^3}$$

↓ 2009

$$-\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{7dx^7} - \frac{4c^2(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{35dx^5} - \frac{8c^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{105dx^3} + \frac{bc\sqrt{d-c^2dx^2}\left(-8c^6\log(x) + \frac{2c^4}{x^2} + \frac{3c^2}{4x^4} - \frac{5}{2x^6}\right)}{105\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^8,x]`

output `-1/7*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(d*x^7) - (4*c^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(35*d*x^5) - (8*c^4*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(105*d*x^3) + (b*c*Sqrt[d - c^2*d*x^2]*(-5/(2*x^6) + (3*c^2)/(4*x^4) + (2*c^4)/x^2 - 8*c^6*Log[x]))/(105*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.64.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2010 `Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)] /; FreeQ[{a, b}, x] && InverseFunctionQ[v]`

rule 6337 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.64.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2536 vs.  $2(235) = 470$ .

Time = 1.17 (sec) , antiderivative size = 2537, normalized size of antiderivative = 9.09

method	result	size
default	Expression too large to display	2537
parts	Expression too large to display	2537

```
input int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^8,x,method=_RETURNVERBOSE)
```

```
output a*(-1/7/d/x^7*(-c^2*d*x^2+d)^(3/2)+4/7*c^2*(-1/5/d/x^5*(-c^2*d*x^2+d)^(3/2)
)-2/15*c^2/d/x^3*(-c^2*d*x^2+d)^(3/2))+8*b*(-d*(c^2*x^2-1))^(1/2)/(280*c^
8*x^8-105*c^6*x^6-21*c^4*x^4-315*c^2*x^2+225)*x^6/(c*x-1)^(1/2)/(c*x+1)^(1
/2)*arccosh(c*x)*c^13+8/5*b*(-d*(c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^
6-21*c^4*x^4-315*c^2*x^2+225)*x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)
*c^11+24*b*(-d*(c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^6-21*c^4*x^4-315*
c^2*x^2+225)*x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*c^9-64/3*b*(-d*(
c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^6-21*c^4*x^4-315*c^2*x^2+225)*x^8
/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*c^15-56/3*b*(-d*(c^2*x^2-1))^(1/
2)/(280*c^8*x^8-105*c^6*x^6-21*c^4*x^4-315*c^2*x^2+225)*x^7/(c*x-1)/(c*x+1
)*arccosh(c*x)*c^14-4/15*b*(-d*(c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^6
-21*c^4*x^4-315*c^2*x^2+225)*x^5/(c*x-1)/(c*x+1)*arccosh(c*x)*c^12-351/5*b
*(-d*(c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^6-21*c^4*x^4-315*c^2*x^2+22
5)*x^3/(c*x-1)/(c*x+1)*arccosh(c*x)*c^10+3057/35*b*(-d*(c^2*x^2-1))^(1/2)/
(280*c^8*x^8-105*c^6*x^6-21*c^4*x^4-315*c^2*x^2+225)*x/(c*x-1)/(c*x+1)*arc
cosh(c*x)*c^8-594/35*b*(-d*(c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^6-21*
c^4*x^4-315*c^2*x^2+225)/x/(c*x-1)/(c*x+1)*arccosh(c*x)*c^6+128/105*b*(-d*
(c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^6-21*c^4*x^4-315*c^2*x^2+225)*x^
11*c^18+16/15*b*(-d*(c^2*x^2-1))^(1/2)/(280*c^8*x^8-105*c^6*x^6-21*c^4*x^4
-315*c^2*x^2+225)*x^9*c^16-88/105*b*(-d*(c^2*x^2-1))^(1/2)/(280*c^8*x^8...
```

### 3.64.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 615, normalized size of antiderivative = 2.20

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^8} dx$$

$$= \frac{4(8bc^8x^8 - 4bc^6x^6 - bc^4x^4 - 18bc^2x^2 + 15b)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) + 16(bc^9x^9 - bc^7x^7)\sqrt{-c^2dx^2 + d}}{32(bc^9x^9 - bc^7x^7)\sqrt{d} \arctan\left(\frac{\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}(x^2 + 1)\sqrt{d}}{c^2dx^4 - (c^2 + 1)dx^2 + d}\right) - 4(8bc^8x^8 - 4bc^6x^6 - bc^4x^4 - 18bc^2x^2 + 15b)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) + 16(bc^9x^9 - bc^7x^7)\sqrt{-c^2dx^2 + d}}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^8,x, algorithm="fracas")`

output `[1/420*(4*(8*b*c^8*x^8 - 4*b*c^6*x^6 - b*c^4*x^4 - 18*b*c^2*x^2 + 15*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + 16*(b*c^9*x^9 - b*c^7*x^7)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + (8*b*c^5*x^5 - (8*b*c^5 + 3*b*c^3 - 10*b*c)*x^7 + 3*b*c^3*x^3 - 10*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 4*(8*a*c^8*x^8 - 4*a*c^6*x^6 - a*c^4*x^4 - 18*a*c^2*x^2 + 15*a)*sqrt(-c^2*d*x^2 + d))/(c^2*x^9 - x^7), -1/420*(32*(b*c^9*x^9 - b*c^7*x^7)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 4*(8*b*c^8*x^8 - 4*b*c^6*x^6 - b*c^4*x^4 - 18*b*c^2*x^2 + 15*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (8*b*c^5*x^5 - (8*b*c^5 + 3*b*c^3 - 10*b*c)*x^7 + 3*b*c^3*x^3 - 10*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 4*(8*a*c^8*x^8 - 4*a*c^6*x^6 - a*c^4*x^4 - 18*a*c^2*x^2 + 15*a)*sqrt(-c^2*d*x^2 + d))/(c^2*x^9 - x^7)]`

### 3.64.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2)/x**8,x)`

output `Timed out`

---

3.64.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^8} dx$

**3.64.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 207, normalized size of antiderivative = 0.74

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^8} dx$$

$$= -\frac{1}{420} \left( 32 c^6 \sqrt{-d} \log(x) - \frac{8 c^4 \sqrt{-d} x^4 + 3 c^2 \sqrt{-d} x^2 - 10 \sqrt{-d}}{x^6} \right) bc$$

$$- \frac{1}{105} \left( \frac{8 (-c^2 dx^2 + d)^{\frac{3}{2}} c^4}{dx^3} + \frac{12 (-c^2 dx^2 + d)^{\frac{3}{2}} c^2}{dx^5} + \frac{15 (-c^2 dx^2 + d)^{\frac{3}{2}}}{dx^7} \right) b \operatorname{arccosh}(cx)$$

$$- \frac{1}{105} \left( \frac{8 (-c^2 dx^2 + d)^{\frac{3}{2}} c^4}{dx^3} + \frac{12 (-c^2 dx^2 + d)^{\frac{3}{2}} c^2}{dx^5} + \frac{15 (-c^2 dx^2 + d)^{\frac{3}{2}}}{dx^7} \right) a$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^8,x, algorithm="maxima")`

output `-1/420*(32*c^6*sqrt(-d)*log(x) - (8*c^4*sqrt(-d)*x^4 + 3*c^2*sqrt(-d)*x^2 - 10*sqrt(-d))/x^6)*b*c - 1/105*(8*(-c^2*d*x^2 + d)^(3/2)*c^4/(d*x^3) + 12*(-c^2*d*x^2 + d)^(3/2)*c^2/(d*x^5) + 15*(-c^2*d*x^2 + d)^(3/2)/(d*x^7))*b*arccosh(c*x) - 1/105*(8*(-c^2*d*x^2 + d)^(3/2)*c^4/(d*x^3) + 12*(-c^2*d*x^2 + d)^(3/2)*c^2/(d*x^5) + 15*(-c^2*d*x^2 + d)^(3/2)/(d*x^7))*a`

**3.64.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^8} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^8,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.64.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^8} dx = \int \frac{(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2}}{x^8} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^8, x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^8, x)`

### 3.65 $\int x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

3.65.1	Optimal result	689
3.65.2	Mathematica [A] (verified)	690
3.65.3	Rubi [A] (verified)	690
3.65.4	Maple [B] (verified)	691
3.65.5	Fricas [A] (verification not implemented)	692
3.65.6	Sympy [F]	693
3.65.7	Maxima [A] (verification not implemented)	693
3.65.8	Giac [F(-2)]	694
3.65.9	Mupad [F(-1)]	694

#### 3.65.1 Optimal result

Integrand size = 27, antiderivative size = 272

$$\int x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \frac{8bx\sqrt{d - c^2 dx^2}}{105c^5\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{4bx^3\sqrt{d - c^2 dx^2}}{315c^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{bx^5\sqrt{d - c^2 dx^2}}{175c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcx^7\sqrt{d - c^2 dx^2}}{49\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6 d}$$

$$+ \frac{2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^6 d^2}$$

$$- \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d^3}$$

output

```
-1/3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/c^6/d+2/5*(-c^2*d*x^2+d)^(5/2)
)*(a+b*arccosh(c*x))/c^6/d^2-1/7*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/c
^6/d^3+8/105*b*x*(-c^2*d*x^2+d)^(1/2)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+4/31
5*b*x^3*(-c^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/175*b*x^5*(
-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/49*b*c*x^7*(-c^2*d*x^2
+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.65.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.62

$$\int x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{\sqrt{d - c^2 dx^2} (bcx(840 + 140c^2 x^2 + 63c^4 x^4 - 225c^6 x^6) + 105a\sqrt{-1 + cx}\sqrt{1 + cx}(-8 - 4c^2 x^2 - 3c^4 x^4 + 15c^6 x^6))}{11025c^6 \sqrt{-1 + cx}\sqrt{1 + cx}}$$

input `Integrate[x^5*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[d - c^2*d*x^2]*(b*c*x*(840 + 140*c^2*x^2 + 63*c^4*x^4 - 225*c^6*x^6) + 105*a*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-8 - 4*c^2*x^2 - 3*c^4*x^4 + 15*c^6*x^6) + 105*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-8 - 4*c^2*x^2 - 3*c^4*x^4 + 15*c^6*x^6)*ArcCosh[c*x]))/(11025*c^6*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.65.3 Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.64, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6337, 27, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6337$$

$$-\frac{bc\sqrt{d - c^2 dx^2} \int \frac{-15c^6 x^6 + 3c^4 x^4 + 4c^2 x^2 + 8}{105c^6} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d^3} +$$

$$\frac{2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^6 d^2} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6 d}$$

$$\downarrow 27$$

$$\frac{b\sqrt{d - c^2 dx^2} \int (-15c^6 x^6 + 3c^4 x^4 + 4c^2 x^2 + 8) dx}{105c^5 \sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d^3} +$$

$$\frac{2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^6 d^2} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6 d}$$

$$\downarrow 2009$$

---

3.65.  $\int x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

$$-\frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d^3} + \frac{2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^6 d^2} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6 d} + \frac{b \left( -\frac{15}{7} c^6 x^7 + \frac{3c^4 x^5}{5} + \frac{4c^2 x^3}{3} + 8x \right) \sqrt{d - c^2 dx^2}}{105c^5 \sqrt{cx - 1} \sqrt{cx + 1}}$$

input `Int[x^5*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `(b*Sqrt[d - c^2*d*x^2]*(8*x + (4*c^2*x^3)/3 + (3*c^4*x^5)/5 - (15*c^6*x^7)/7))/(105*c^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(3*c^6*d) + (2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(5*c^6*d^2) - ((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(7*c^6*d^3)`

### 3.65.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.65.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 987 vs.  $2(228) = 456$ .

Time = 0.69 (sec) , antiderivative size = 988, normalized size of antiderivative = 3.63

method	result
default	$a \left( -\frac{x^4(-c^2dx^2+d)^{\frac{3}{2}}}{7c^2d} + \frac{4x^2(-c^2dx^2+d)^{\frac{3}{2}}}{35c^2d} - \frac{8(-c^2dx^2+d)^{\frac{3}{2}}}{105dc^4} \right) + b \left( \frac{\sqrt{-d(c^2x^2-1)}(64c^8x^8-144c^6x^6+64\sqrt{cx+1}\sqrt{cx-1}}{\dots} \right)$
parts	$a \left( -\frac{x^4(-c^2dx^2+d)^{\frac{3}{2}}}{7c^2d} + \frac{4x^2(-c^2dx^2+d)^{\frac{3}{2}}}{35c^2d} - \frac{8(-c^2dx^2+d)^{\frac{3}{2}}}{105dc^4} \right) + b \left( \frac{\sqrt{-d(c^2x^2-1)}(64c^8x^8-144c^6x^6+64\sqrt{cx+1}\sqrt{cx-1}}{\dots} \right)$

input `int(x^5*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output

```

a*(-1/7*x^4*(-c^2*d*x^2+d)^(3/2)/c^2/d+4/7/c^2*(-1/5*x^2*(-c^2*d*x^2+d)^(3/2)/c^2/d-2/15/d/c^4*(-c^2*d*x^2+d)^(3/2)))+b*(1/6272*(-d*(c^2*x^2-1))^(1/2)*(64*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*x^4-112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-25*c^2*x^2+56*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+7*arccosh(c*x)))/(c*x+1)/c^6/(c*x-1)+3/3200*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c*x)))/(c*x+1)/c^6/(c*x-1)+1/1152*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x)))/(c*x+1)/c^6/(c*x-1)-5/128*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+arccosh(c*x)))/(c*x+1)/c^6/(c*x-1)-5/128*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x)))/(c*x+1)/c^6/(c*x-1)+1/1152*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x)))/(c*x+1)/c^6/(c*x-1)+3/3200*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-28*c^4*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(1+5*arccosh(c*x)))/(c*x+1)/c^6/(c*x-1)+1/6272*(-d*(c^2*x^2-1))^(1/2)*(-64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+64*c^8*x^8+112*(c*x+1)...
```

### 3.65.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.75

$$\int x^5 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx$$

$$= \frac{105(15bc^8x^8 - 18bc^6x^6 - bc^4x^4 - 4bc^2x^2 + 8b)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) - (225bc^7x^7 - 63bc^5x^5 - 21bc^3x^3 + 3bcx)}{c^6}$$

input `integrate(x^5*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `1/11025*(105*(15*b*c^8*x^8 - 18*b*c^6*x^6 - b*c^4*x^4 - 4*b*c^2*x^2 + 8*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (225*b*c^7*x^7 - 63*b*c^5*x^5 - 140*b*c^3*x^3 - 840*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 105*(15*a*c^8*x^8 - 18*a*c^6*x^6 - a*c^4*x^4 - 4*a*c^2*x^2 + 8*a)*sqrt(-c^2*d*x^2 + d))/(c^8*x^2 - c^6)`

### 3.65.6 Sympy [F]

$$\int x^5 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int x^5 \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate(x**5*(a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**5*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x)), x)`

### 3.65.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.75

$$\begin{aligned} & \int x^5 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx \\ &= -\frac{1}{105} \left( \frac{15(-c^2 dx^2 + d)^{\frac{3}{2}} x^4}{c^2 d} + \frac{12(-c^2 dx^2 + d)^{\frac{3}{2}} x^2}{c^4 d} + \frac{8(-c^2 dx^2 + d)^{\frac{3}{2}}}{c^6 d} \right) b \operatorname{arccosh}(cx) \\ & \quad - \frac{1}{105} \left( \frac{15(-c^2 dx^2 + d)^{\frac{3}{2}} x^4}{c^2 d} + \frac{12(-c^2 dx^2 + d)^{\frac{3}{2}} x^2}{c^4 d} + \frac{8(-c^2 dx^2 + d)^{\frac{3}{2}}}{c^6 d} \right) a \\ & \quad - \frac{(225 c^6 \sqrt{-dx^7} - 63 c^4 \sqrt{-dx^5} - 140 c^2 \sqrt{-dx^3} - 840 \sqrt{-dx}) b}{11025 c^5} \end{aligned}$$

input `integrate(x^5*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output 
$$-1/105*(15*(-c^2*d*x^2 + d)^{(3/2)}*x^4/(c^2*d) + 12*(-c^2*d*x^2 + d)^{(3/2)}*x^2/(c^4*d) + 8*(-c^2*d*x^2 + d)^{(3/2)}/(c^6*d))*b*\operatorname{arccosh}(c*x) - 1/105*(15*(-c^2*d*x^2 + d)^{(3/2)}*x^4/(c^2*d) + 12*(-c^2*d*x^2 + d)^{(3/2)}*x^2/(c^4*d) + 8*(-c^2*d*x^2 + d)^{(3/2)}/(c^6*d))*a - 1/11025*(225*c^6*\sqrt{-d}*x^7 - 63*c^4*\sqrt{-d}*x^5 - 140*c^2*\sqrt{-d}*x^3 - 840*\sqrt{-d}*x)*b/c^5$$

### 3.65.8 Giac [F(-2)]

Exception generated.

$$\int x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^5*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.65.9 Mupad [F(-1)]

Timed out.

$$\int x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int x^5 (a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2} dx$$

input `int(x^5*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2),x)`

output `int(x^5*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2), x)`

### 3.66 $\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

3.66.1	Optimal result . . . . .	695
3.66.2	Mathematica [A] (verified) . . . . .	696
3.66.3	Rubi [A] (verified) . . . . .	696
3.66.4	Maple [B] (verified) . . . . .	697
3.66.5	Fricas [A] (verification not implemented) . . . . .	698
3.66.6	Sympy [F] . . . . .	699
3.66.7	Maxima [A] (verification not implemented) . . . . .	699
3.66.8	Giac [F(-2)] . . . . .	700
3.66.9	Mupad [F(-1)] . . . . .	700

#### 3.66.1 Optimal result

Integrand size = 27, antiderivative size = 195

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \frac{2bx\sqrt{d - c^2 dx^2}}{15c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bx^3\sqrt{d - c^2 dx^2}}{45c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcx^5\sqrt{d - c^2 dx^2}}{25\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^4 d} + \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d^2}$$

```
output -1/3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/c^4/d+1/5*(-c^2*d*x^2+d)^(5/2)
)*(a+b*arccosh(c*x))/c^4/d^2+2/15*b*x*(-c^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(1/2)
)/(c*x+1)^(1/2)+1/45*b*x^3*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)
)-1/25*b*c*x^5*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```



**3.66.2 Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.65

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{\sqrt{d - c^2 dx^2} (bc(30x + 5c^2 x^3 - 9c^4 x^5) + 30(-1 + cx)^{3/2} (1 + cx)^{3/2} (a + \operatorname{barccosh}(cx)) + 45c^2 x^2 (-1 + cx)^{3/2})}{225c^4 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

input `Integrate[x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`output `(Sqrt[d - c^2*d*x^2]*(b*c*(30*x + 5*c^2*x^3 - 9*c^4*x^5) + 30*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(a + b*ArcCosh[c*x]) + 45*c^2*x^2*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(a + b*ArcCosh[c*x]))/(225*c^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`**3.66.3 Rubi [A] (verified)**Time = 0.40 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.68, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6337, 27, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6337}$$

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{-3c^4 x^4 + c^2 x^2 + 2}{15c^4} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d^2} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^4 d}$$

$$\downarrow \text{27}$$

$$\frac{b\sqrt{d - c^2 dx^2} \int (-3c^4 x^4 + c^2 x^2 + 2) dx}{15c^3 \sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d^2} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^4 d}$$

$$\downarrow \text{2009}$$

---

 3.66.  $\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d^2} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^4 d} + \frac{b \left( -\frac{3}{5}c^4 x^5 + \frac{c^2 x^3}{3} + 2x \right) \sqrt{d - c^2 dx^2}}{15c^3 \sqrt{cx - 1} \sqrt{cx + 1}}$$

input `Int[x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `(b*Sqrt[d - c^2*d*x^2]*(2*x + (c^2*x^3)/3 - (3*c^4*x^5)/5))/(15*c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(3*c^4*d) + ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(5*c^4*d^2)`

### 3.66.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.66.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 639 vs. 2(163) = 326.

Time = 1.32 (sec) , antiderivative size = 640, normalized size of antiderivative = 3.28

method	result
default	$a \left( -\frac{x^2(-c^2 dx^2 + d)^{\frac{3}{2}}}{5c^2 d} - \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}}}{15dc^4} \right) + b \left( \frac{\sqrt{-d(c^2 x^2 - 1)} (16c^6 x^6 - 28c^4 x^4 + 16\sqrt{cx+1} \sqrt{cx-1} c^5 x^5 + 13c^2 x^2 - 20\sqrt{cx-1})}{800(cx+1)c^4(cx-1)} \right)$
parts	$a \left( -\frac{x^2(-c^2 dx^2 + d)^{\frac{3}{2}}}{5c^2 d} - \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}}}{15dc^4} \right) + b \left( \frac{\sqrt{-d(c^2 x^2 - 1)} (16c^6 x^6 - 28c^4 x^4 + 16\sqrt{cx+1} \sqrt{cx-1} c^5 x^5 + 13c^2 x^2 - 20\sqrt{cx-1})}{800(cx+1)c^4(cx-1)} \right)$

3.66.  $\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

input `int(x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & a*(-1/5*x^2*(-c^2*d*x^2+d)^(3/2)/c^2/d-2/15/d/c^4*(-c^2*d*x^2+d)^(3/2))+b* \\ & (1/800*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x \\ & -1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x \\ & -1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c*x))/(c*x+1)/c^4/(c*x-1)+1/2 \\ & 88*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2) \\ & *c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))/(c*x+1 \\ & )/c^4/(c*x-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x \\ & +c^2*x^2-1)*(-1+arccosh(c*x))/(c*x+1)/c^4/(c*x-1)-1/16*(-d*(c^2*x^2-1))^(1 \\ & /2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))/(c*x+1)/ \\ & c^4/(c*x-1)+1/288*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c \\ & ^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arcco \\ & sh(c*x))/(c*x+1)/c^4/(c*x-1)+1/800*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/ \\ & 2)*(c*x-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3 \\ & -28*c^4*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(1+5*arccosh(c \\ & *x))/(c*x+1)/c^4/(c*x-1) \end{aligned}$$

### 3.66.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.90

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx)) dx = \frac{15(3bc^6x^6 - 4bc^4x^4 - bc^2x^2 + 2b)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) - (9bc^5x^5 - 5bc^3x^3 - 30bcx)\sqrt{-c^2dx^2 + d}}{225(c^6x^2 - c^4)}$$

input `integrate(x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output 
$$\frac{1}{225}*(15*(3*b*c^6*x^6 - 4*b*c^4*x^4 - b*c^2*x^2 + 2*b)*\operatorname{sqrt}(-c^2*d*x^2 + d)*\log(c*x + \operatorname{sqrt}(c^2*x^2 - 1)) - (9*b*c^5*x^5 - 5*b*c^3*x^3 - 30*b*c*x)*\operatorname{sqrt}(-c^2*d*x^2 + d)*\operatorname{sqrt}(c^2*x^2 - 1) + 15*(3*a*c^6*x^6 - 4*a*c^4*x^4 - a*c^2*x^2 + 2*a)*\operatorname{sqrt}(-c^2*d*x^2 + d))/(c^6*x^2 - c^4)$$

**3.66.6 Sympy [F]**

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int x^3 \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate(x**3*(a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**3*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x)), x)`

**3.66.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.74

$$\begin{aligned} & \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx \\ &= -\frac{1}{15} b \left( \frac{3(-c^2 dx^2 + d)^{\frac{3}{2}} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}}}{c^4 d} \right) \operatorname{arcosh}(cx) \\ & \quad - \frac{1}{15} a \left( \frac{3(-c^2 dx^2 + d)^{\frac{3}{2}} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}}}{c^4 d} \right) \\ & \quad - \frac{(9c^4 \sqrt{-dx^5} - 5c^2 \sqrt{-dx^3} - 30\sqrt{-dx})b}{225c^3} \end{aligned}$$

input `integrate(x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/15*b*(3*(-c^2*d*x^2 + d)^(3/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(3/2)/(c^4*d))*arccosh(c*x) - 1/15*a*(3*(-c^2*d*x^2 + d)^(3/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(3/2)/(c^4*d)) - 1/225*(9*c^4*sqrt(-d)*x^5 - 5*c^2*sqrt(-d)*x^3 - 30*sqrt(-d)*x)*b/c^3`

**3.66.8 Giac [F(-2)]**

Exception generated.

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.66.9 Mupad [F(-1)]**

Timed out.

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int x^3 (a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2} dx$$

input `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2),x)`

output `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2), x)`

### 3.67 $\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) dx$

3.67.1	Optimal result	701
3.67.2	Mathematica [A] (verified)	701
3.67.3	Rubi [A] (verified)	702
3.67.4	Maple [B] (verified)	703
3.67.5	Fricas [A] (verification not implemented)	704
3.67.6	Sympy [F]	704
3.67.7	Maxima [A] (verification not implemented)	705
3.67.8	Giac [F(-2)]	705
3.67.9	Mupad [F(-1)]	705

#### 3.67.1 Optimal result

Integrand size = 25, antiderivative size = 118

$$\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) dx = \frac{bx\sqrt{d - c^2dx^2}}{3c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcx^3\sqrt{d - c^2dx^2}}{9\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))}{3c^2d}$$

output 
$$-1/3*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))/c^2/d+1/3*b*x*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/9*b*c*x^3*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

#### 3.67.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.83

$$\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) dx = \frac{\sqrt{d - c^2dx^2}(bcx\sqrt{-1 + cx}\sqrt{1 + cx}(3 - c^2x^2) + 3a(-1 + c^2x^2)^2 + 3b(-1 + c^2x^2)^2 \operatorname{arccosh}(cx))}{9c^2(-1 + c^2x^2)}$$

input `Integrate[x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output  $(\text{Sqrt}[d - c^2*d*x^2]*(b*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(3 - c^2*x^2) + 3*a*(-1 + c^2*x^2)^2 + 3*b*(-1 + c^2*x^2)^2*\text{ArcCosh}[c*x]))/(9*c^2*(-1 + c^2*x^2))$

### 3.67.3 Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.74, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {6329, 25, 39, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x\sqrt{d - c^2dx^2}(a + \text{barccosh}(cx)) dx \\
 & \quad \downarrow 6329 \\
 & -\frac{b\sqrt{d - c^2dx^2} \int -((1 - cx)(cx + 1))dx}{3c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2dx^2)^{3/2} (a + \text{barccosh}(cx))}{3c^2d} \\
 & \quad \downarrow 25 \\
 & \frac{b\sqrt{d - c^2dx^2} \int (1 - cx)(cx + 1)dx}{3c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2dx^2)^{3/2} (a + \text{barccosh}(cx))}{3c^2d} \\
 & \quad \downarrow 39 \\
 & \frac{b\sqrt{d - c^2dx^2} \int (1 - c^2x^2) dx}{3c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2dx^2)^{3/2} (a + \text{barccosh}(cx))}{3c^2d} \\
 & \quad \downarrow 2009 \\
 & \frac{b\left(x - \frac{c^2x^3}{3}\right)\sqrt{d - c^2dx^2}}{3c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2dx^2)^{3/2} (a + \text{barccosh}(cx))}{3c^2d}
 \end{aligned}$$

input  $\text{Int}[x*\text{Sqrt}[d - c^2*d*x^2]*(a + b*\text{ArcCosh}[c*x]), x]$

output  $(b*\text{Sqrt}[d - c^2*d*x^2]*(x - (c^2*x^3)/3))/(3*c*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) - ((d - c^2*d*x^2)^(3/2)*(a + b*\text{ArcCosh}[c*x]))/(3*c^2*d)$

## 3.67.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 39 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(m_)), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6329 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

## 3.67.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 355 vs. 2(98) = 196.

Time = 0.51 (sec) , antiderivative size = 356, normalized size of antiderivative = 3.02

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{3}{2}}}{3c^2d} + b \left( \frac{\sqrt{-d(c^2x^2-1)}(4c^4x^4-5c^2x^2+4\sqrt{cx-1}\sqrt{cx+1}c^3x^3-3\sqrt{cx-1}\sqrt{cx+1}cx+1)(-1+3\operatorname{arccosh}(cx))}{72(cx+1)c^2(cx-1)} \right)$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{3}{2}}}{3c^2d} + b \left( \frac{\sqrt{-d(c^2x^2-1)}(4c^4x^4-5c^2x^2+4\sqrt{cx-1}\sqrt{cx+1}c^3x^3-3\sqrt{cx-1}\sqrt{cx+1}cx+1)(-1+3\operatorname{arccosh}(cx))}{72(cx+1)c^2(cx-1)} \right)$

input `int(x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`



```
output -1/3*a*(-c^2*d*x^2+d)^(3/2)/c^2/d+b*(1/72*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))/(c*x+1)/c^2/(c*x-1)-1/8*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+arccosh(c*x))/(c*x+1)/c^2/(c*x-1)-1/8*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))/(c*x+1)/c^2/(c*x-1)+1/72*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x))/(c*x+1)/c^2/(c*x-1)
```

### 3.67.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.20

$$\int x\sqrt{d-c^2x^2}(a+\operatorname{barccosh}(cx))dx$$

$$= \frac{3(bc^4x^4 - 2bc^2x^2 + b)\sqrt{-c^2dx^2 + d}\log(cx + \sqrt{c^2x^2 - 1}) - (bc^3x^3 - 3bcx)\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1} + 3}{9(c^4x^2 - c^2)}$$

```
input integrate(x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")
```

```
output 1/9*(3*(b*c^4*x^4 - 2*b*c^2*x^2 + b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (b*c^3*x^3 - 3*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 3*(a*c^4*x^4 - 2*a*c^2*x^2 + a)*sqrt(-c^2*d*x^2 + d))/(c^4*x^2 - c^2)
```

### 3.67.6 Sympy [F]

$$\int x\sqrt{d-c^2x^2}(a+\operatorname{barccosh}(cx))dx = \int x\sqrt{-d(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))dx$$

```
input integrate(x*(a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2),x)
```

```
output Integral(x*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x)), x)
```

**3.67.7 Maxima [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.69

$$\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) dx = -\frac{(-c^2 dx^2 + d)^{\frac{3}{2}} b \operatorname{arcosh}(cx)}{3 c^2 d} - \frac{(c^2 \sqrt{-d} dx^3 - 3 \sqrt{-d} dx) b}{9 cd} - \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} a}{3 c^2 d}$$

input `integrate(x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/3*(-c^2*d*x^2 + d)^(3/2)*b*arccosh(c*x)/(c^2*d) - 1/9*(c^2*sqrt(-d)*d*x^3 - 3*sqrt(-d)*d*x)*b/(c*d) - 1/3*(-c^2*d*x^2 + d)^(3/2)*a/(c^2*d)`

**3.67.8 Giac [F(-2)]**

Exception generated.

$$\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.67.9 Mupad [F(-1)]**

Timed out.

$$\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2} dx$$

input `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2),x)`

output `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2), x)`

### 3.68 $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x} dx$

3.68.1	Optimal result	706
3.68.2	Mathematica [A] (warning: unable to verify)	707
3.68.3	Rubi [A] (verified)	707
3.68.4	Maple [A] (verified)	710
3.68.5	Fricas [F]	710
3.68.6	Sympy [F]	711
3.68.7	Maxima [F]	711
3.68.8	Giac [F(-2)]	711
3.68.9	Mupad [F(-1)]	712

#### 3.68.1 Optimal result

Integrand size = 27, antiderivative size = 213

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x} dx = -\frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\arctan(e^{\operatorname{arccosh}(cx)})} - \frac{\sqrt{-1+cx}\sqrt{1+cx}}{ib\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})} + \frac{\sqrt{-1+cx}\sqrt{1+cx}}{ib\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(cx)})} - \frac{\sqrt{-1+cx}\sqrt{1+cx}}{\sqrt{-1+cx}\sqrt{1+cx}}$$

output  $(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}-b*c*x*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2*(a+b*\operatorname{arccosh}(c*x))*\arctan(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+I*b*\operatorname{polylog}(2,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-I*b*\operatorname{polylog}(2,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.68.2 Mathematica [A] (warning: unable to verify)

Time = 0.65 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x} dx$$

$$= a\sqrt{d - c^2 dx^2} + a\sqrt{d} \log(x) - a\sqrt{d} \log\left(d + \sqrt{d}\sqrt{d - c^2 dx^2}\right)$$

$$+ \frac{b\sqrt{d - c^2 dx^2}\left(-cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx\sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + i \operatorname{arccosh}(cx) \log(1 - ie^{-\operatorname{arccosh}(cx)})\right)}{\sqrt{\frac{-1+cx}{1+cx}}}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x,x]`

output `a*Sqrt[d - c^2*d*x^2] + a*Sqrt[d]*Log[x] - a*Sqrt[d]*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (b*Sqrt[d - c^2*d*x^2]*(-(c*x) + Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + I*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*PolyLog[2, I/E^ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`

### 3.68.3 Rubi [A] (verified)

Time = 0.82 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.70, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$ , Rules used = {6341, 24, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x} dx$$

$$\downarrow 6341$$

$$-\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d - c^2 dx^2} \int 1 dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))$$

$$\downarrow 24$$

$$-\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d - c^2 dx^2}}{\sqrt{cx-1}\sqrt{cx+1}}$$

---

3.68.  $\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x} dx$

$$\begin{aligned}
& \downarrow 6362 \\
& -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 3042 \\
& -\frac{\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a + \\
& \quad \operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 4668 \\
& -\frac{\sqrt{d-c^2dx^2}(-ib \int \log(1-ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1+ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)}))}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 2715 \\
& -\frac{\sqrt{d-c^2dx^2}(-ib \int e^{-\operatorname{arccosh}(cx)} \log(1-ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1+ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)}))}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 2838 \\
& -\frac{\sqrt{d-c^2dx^2}(2 \arctan(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}))}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x,x]`

output `-((b*c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.68.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x)) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`
- rule 6341 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x)) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`
- rule 6362 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)^(m_)/(Sqrt[(d1_) + (e1_)*(x_)]*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.68.4 Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 394, normalized size of antiderivative = 1.85

method	result
default	$-\sqrt{d} \ln \left( \frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x} \right) a + a\sqrt{-c^2dx^2+d} + \frac{b\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)x^2c^2}{(cx-1)(cx+1)} - \frac{b\sqrt{-d(c^2x^2-1)} cx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{b}{\sqrt{cx-1}\sqrt{cx+1}}$
parts	$-\sqrt{d} \ln \left( \frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x} \right) a + a\sqrt{-c^2dx^2+d} + \frac{b\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)x^2c^2}{(cx-1)(cx+1)} - \frac{b\sqrt{-d(c^2x^2-1)} cx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{b}{\sqrt{cx-1}\sqrt{cx+1}}$

input `int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x,x,method=_RETURNVERBOSE)`

output `-d^(1/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)*a+a*(-c^2*d*x^2+d)^(1/2)+b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)/(c*x+1)*arccosh(c*x)*x^2*c^2-b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c*x-b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)/(c*x+1)*arccosh(c*x)+I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))`

### 3.68.5 Fracas [F]

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x} dx = \int \frac{\sqrt{-c^2dx^2+d}(b\operatorname{arccosh}(cx)+a)}{x} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2+d)*(b*arccosh(c*x)+a)/x,x)`

**3.68.6 Sympy [F]**

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)}(a + b \operatorname{acosh}(cx))}{x} dx$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2)/x,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))/x, x)`

**3.68.7 Maxima [F]**

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{\sqrt{-c^2 dx^2 + d}(b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="maxima")`

output `-(sqrt(d)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - sqrt(-c^2*d*x^2 + d)*a + b*integrate(sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x, x)`

**3.68.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.68.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2}}{x} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x, x)`

**3.69** 
$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^3} dx$$

3.69.1	Optimal result	713
3.69.2	Mathematica [A] (warning: unable to verify)	714
3.69.3	Rubi [A] (verified)	714
3.69.4	Maple [A] (verified)	717
3.69.5	Fricas [F]	717
3.69.6	Sympy [F]	718
3.69.7	Maxima [F]	718
3.69.8	Giac [F(-2)]	718
3.69.9	Mupad [F(-1)]	719

**3.69.1 Optimal result**

Integrand size = 27, antiderivative size = 235

$$\begin{aligned} & \int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^3} dx \\ &= -\frac{bc\sqrt{d-c^2dx^2}}{2x\sqrt{-1+cx}\sqrt{1+cx}} - \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{2x^2} \\ & \quad + \frac{c^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & \quad - \frac{ibc^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{ibc^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output

```
-1/2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^2-1/2*b*c*(-c^2*d*x^2+d)^(1/2)/x/(c*x-1)^(1/2)/(c*x+1)^(1/2)+c^2*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/2*I*b*c^2*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/2*I*b*c^2*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.69.2 Mathematica [A] (warning: unable to verify)**

Time = 0.89 (sec) , antiderivative size = 307, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^3} dx$$

$$= \frac{1}{2} \left( -\frac{a\sqrt{d - c^2 dx^2}}{x^2} - ac^2\sqrt{d}\log(x) + ac^2\sqrt{d}\log\left(d + \sqrt{d}\sqrt{d - c^2 dx^2}\right) \right.$$

$$\left. + \frac{bd(1 + cx)\left(cx\sqrt{\frac{-1+cx}{1+cx}} - \operatorname{arccosh}(cx) + cx\operatorname{arccosh}(cx) + ic^2x^2\sqrt{\frac{-1+cx}{1+cx}}\operatorname{arccosh}(cx)\log\left(1 - ie^{-\operatorname{arccosh}(cx)}\right)\right)}{2} \right)$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^3,x]`output `(-((a*Sqrt[d - c^2*d*x^2])/x^2) - a*c^2*Sqrt[d]*Log[x] + a*c^2*Sqrt[d]*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (b*d*(1 + c*x)*(c*x*Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x] + c*x*ArcCosh[c*x] + I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*PolyLog[2, I/E^ArcCosh[c*x]]))/(x^2*Sqrt[d - c^2*d*x^2]))/2`**3.69.3 Rubi [A] (verified)**Time = 0.82 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.70, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$ , Rules used = {6339, 15, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^3} dx$$

$$\downarrow \text{6339}$$

$$\frac{c^2\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{bc\sqrt{d - c^2 dx^2} \int \frac{1}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{2x^2}$$

$$\downarrow \text{15}$$

---

3.69.  $\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^3} dx$

$$\begin{aligned}
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x \sqrt{cx - 1} \sqrt{cx + 1}} dx}{2 \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{bc \sqrt{d - c^2 dx^2}}{2x \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{6362} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{2 \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{bc \sqrt{d - c^2 dx^2}}{2x \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{3042} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx)) \operatorname{csc} \left( i \operatorname{arccosh}(cx) + \frac{\pi}{2} \right) \operatorname{darccosh}(cx)}{2 \sqrt{cx - 1} \sqrt{cx + 1}} - \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{bc \sqrt{d - c^2 dx^2}}{2x \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{4668} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} (-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)}))}{2 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{bc \sqrt{d - c^2 dx^2}}{2x \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{2715} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} (-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{2 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{bc \sqrt{d - c^2 dx^2}}{2x \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{2838} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}))}{2 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2x^2} - \frac{bc \sqrt{d - c^2 dx^2}}{2x \sqrt{cx - 1} \sqrt{cx + 1}}
\end{aligned}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^3,x]`

output `-1/2*(b*c*Sqrt[d - c^2*d*x^2])/(x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(2*x^2) + (c^2*Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

---

3.69.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^3} dx$

## 3.69.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`
- rule 6339 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`
- rule 6362 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.69.4 Maple [A] (verified)

Time = 1.22 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.86

method	result
default	$a \left( -\frac{(-c^2 d x^2 + d)^{\frac{3}{2}}}{2 d x^2} - \frac{c^2 \left( \sqrt{-c^2 d x^2 + d} - \sqrt{d} \ln \left( \frac{2 d + 2 \sqrt{d} \sqrt{-c^2 d x^2 + d}}{x} \right) \right)}{2} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(c x) c^2}{2(c x - 1)(c x + 1)} - \frac{b \sqrt{-d(c^2 x^2 - 1)}}{2 x \sqrt{c x - 1} \sqrt{c x + 1}}$
parts	$a \left( -\frac{(-c^2 d x^2 + d)^{\frac{3}{2}}}{2 d x^2} - \frac{c^2 \left( \sqrt{-c^2 d x^2 + d} - \sqrt{d} \ln \left( \frac{2 d + 2 \sqrt{d} \sqrt{-c^2 d x^2 + d}}{x} \right) \right)}{2} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(c x) c^2}{2(c x - 1)(c x + 1)} - \frac{b \sqrt{-d(c^2 x^2 - 1)}}{2 x \sqrt{c x - 1} \sqrt{c x + 1}}$

input `int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^3,x,method=_RETURNVERBOSE)`

output `a*(-1/2/d/x^2*(-c^2*d*x^2+d)^(3/2)-1/2*c^2*((-c^2*d*x^2+d)^(1/2)-d^(1/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x))-1/2*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)/(c*x+1)*arccosh(c*x)*c^2-1/2*b*(-d*(c^2*x^2-1))^(1/2)/x/(c*x-1)^(1/2)/(c*x+1)^(1/2)*c+1/2*b*(-d*(c^2*x^2-1))^(1/2)*arccosh(c*x)/x^2/(c*x-1)/(c*x+1)-1/2*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2+1/2*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2-1/2*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2+1/2*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2`

### 3.69.5 Fracas [F]

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{\sqrt{-c^2 dx^2 + d}(b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^3,x,algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/x^3, x)`

**3.69.6 Sympy [F]**

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)}(a + b \operatorname{acosh}(cx))}{x^3} dx$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2)/x**3,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))/x**3, x)`

**3.69.7 Maxima [F]**

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{\sqrt{-c^2 dx^2 + d}(b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^3,x, algorithm="maxima")`

output `1/2*(c^2*sqrt(d)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - sqrt(-c^2*d*x^2 + d)*c^2 - (-c^2*d*x^2 + d)^(3/2)/(d*x^2))*a + b*integrate(sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^3, x)`

**3.69.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.69.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2}}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^3, x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^3, x)`



### 3.70 $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$

3.70.1	Optimal result	720
3.70.2	Mathematica [A] (warning: unable to verify)	721
3.70.3	Rubi [A] (verified)	721
3.70.4	Maple [A] (verified)	725
3.70.5	Fricas [F]	725
3.70.6	Sympy [F]	726
3.70.7	Maxima [F]	726
3.70.8	Giac [F(-2)]	726
3.70.9	Mupad [F(-1)]	727

#### 3.70.1 Optimal result

Integrand size = 27, antiderivative size = 315

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$$

$$= -\frac{bc\sqrt{d-c^2dx^2}}{12x^3\sqrt{-1+cx}\sqrt{1+cx}} + \frac{bc^3\sqrt{d-c^2dx^2}}{8x\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{4x^4} + \frac{c^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{8x^2}$$

$$+ \frac{c^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{4\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{ibc^4\sqrt{d-c^2dx^2} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{8\sqrt{-1+cx}\sqrt{1+cx}} + \frac{ibc^4\sqrt{d-c^2dx^2} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{8\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/4*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^4+1/8*c^2*(a+b*arccosh(c*x))
)*(-c^2*d*x^2+d)^(1/2)/x^2-1/12*b*c*(-c^2*d*x^2+d)^(1/2)/x^3/(c*x-1)^(1/2)
/(c*x+1)^(1/2)+1/8*b*c^3*(-c^2*d*x^2+d)^(1/2)/x/(c*x-1)^(1/2)/(c*x+1)^(1/2)
)+1/4*c^4*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2
*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/8*I*b*c^4*polylog(2,-I*(c*x+
(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(
1/2)+1/8*I*b*c^4*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^
2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.70.2 Mathematica [A] (warning: unable to verify)

Time = 0.85 (sec) , antiderivative size = 290, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^5} dx$$

$$= \frac{1}{24} \left( \frac{3a(-2 + c^2 x^2) \sqrt{d - c^2 dx^2}}{x^4} - 3ac^4 \sqrt{d} \log(x) + 3ac^4 \sqrt{d} \log(d + \sqrt{d} \sqrt{d - c^2 dx^2}) \right.$$

$$\left. + \frac{b\sqrt{d - c^2 dx^2}(-2cx + 3c^3 x^3 - 6\sqrt{\frac{-1+cx}{1+cx}}(1+cx)\operatorname{arccosh}(cx) + 3c^2 x^2 \sqrt{\frac{-1+cx}{1+cx}}(1+cx)\operatorname{arccosh}(cx) - 3i \dots}{x^4} \right)$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^5, x]`

output `((3*a*(-2 + c^2*x^2)*Sqrt[d - c^2*d*x^2])/x^4 - 3*a*c^4*Sqrt[d]*Log[x] + 3*a*c^4*Sqrt[d]*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (b*Sqrt[d - c^2*d*x^2]*(-2*c*x + 3*c^3*x^3 - 6*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x] + 3*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x] - (3*I)*c^4*x^4*ArcCosh[c*x]*(Log[1 - I/E^ArcCosh[c*x]] - Log[1 + I/E^ArcCosh[c*x]]) - (3*I)*c^4*x^4*(PolyLog[2, (-I)/E^ArcCosh[c*x]] - PolyLog[2, I/E^ArcCosh[c*x]])))/x^4*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))/24`

### 3.70.3 Rubi [A] (verified)

Time = 1.18 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.68, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6339, 15, 6348, 15, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^5} dx$$

$$\downarrow \text{6339}$$

$$\frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x^3 \sqrt{cx - 1} \sqrt{cx + 1}} dx}{4\sqrt{cx - 1} \sqrt{cx + 1}} + \frac{bc \sqrt{d - c^2 dx^2} \int \frac{1}{x^4} dx}{4\sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{4x^4}$$

$$\downarrow \text{15}$$

---

3.70.  $\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^5} dx$

$$\begin{aligned}
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x^3 \sqrt{cx - 1} \sqrt{cx + 1}} dx}{4 \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4x^4} - \frac{bc \sqrt{d - c^2 dx^2}}{12x^3 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{6348} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x \sqrt{cx - 1} \sqrt{cx + 1}} dx - \frac{1}{2} bc \int \frac{1}{x^2} dx + \frac{\sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))}{2x^2} \right)}{4 \sqrt{cx - 1} \sqrt{cx + 1}} - \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4x^4} - \frac{bc \sqrt{d - c^2 dx^2}}{12x^3 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{15} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x \sqrt{cx - 1} \sqrt{cx + 1}} dx + \frac{\sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))}{2x^2} + \frac{bc}{2x} \right)}{4 \sqrt{cx - 1} \sqrt{cx + 1}} - \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4x^4} - \frac{bc \sqrt{d - c^2 dx^2}}{12x^3 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{6362} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx) + \frac{\sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))}{2x^2} + \frac{bc}{2x} \right)}{4 \sqrt{cx - 1} \sqrt{cx + 1}} - \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4x^4} - \frac{bc \sqrt{d - c^2 dx^2}}{12x^3 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{3042} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{2} c^2 \int (a + \operatorname{barccosh}(cx)) \csc(i \operatorname{arccosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx) + \frac{\sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))}{2x^2} + \frac{bc}{2x} \right)}{4 \sqrt{cx - 1} \sqrt{cx + 1}} - \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4x^4} - \frac{bc \sqrt{d - c^2 dx^2}}{12x^3 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{4668} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{2} c^2 (-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan(\frac{\sqrt{cx - 1} \sqrt{cx + 1}}{cx})) \right)}{4 \sqrt{cx - 1} \sqrt{cx + 1}} - \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4x^4} - \frac{bc \sqrt{d - c^2 dx^2}}{12x^3 \sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{2715} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{2} c^2 (-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} \right)}{4 \sqrt{cx - 1} \sqrt{cx + 1}} - \\
& \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4x^4} - \frac{bc \sqrt{d - c^2 dx^2}}{12x^3 \sqrt{cx - 1} \sqrt{cx + 1}}
\end{aligned}$$


---

3.70.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^5} dx$

↓ 2838

$$\frac{c^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{2} c^2 (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})) \right)}{4x^4} - \frac{4\sqrt{cx-1}\sqrt{cx+1} bc\sqrt{d-c^2 dx^2}}{12x^3\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x^5,x]`

output `-1/12*(b*c*Sqrt[d - c^2*d*x^2])/(x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(4*x^4) + (c^2*Sqrt[d - c^2*d*x^2]*((b*c)/(2*x) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(2*x^2) + (c^2*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/2))/(4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.70.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6339 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`

rule 6348 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d1*d2*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && ILtQ[m, -1]`

rule 6362 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.70.4 Maple [A] (verified)

Time = 1.11 (sec) , antiderivative size = 541, normalized size of antiderivative = 1.72

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{3}{2}}}{4dx^4} - \frac{ac^2(-c^2dx^2+d)^{\frac{3}{2}}}{8dx^2} + \frac{ac^4\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{8} - \frac{ac^4\sqrt{-c^2dx^2+d}}{8} + \frac{b\sqrt{-d(c^2x^2-1)}\operatorname{arccosh}(cx)}{8(cx+1)(cx-1)}$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{3}{2}}}{4dx^4} - \frac{ac^2(-c^2dx^2+d)^{\frac{3}{2}}}{8dx^2} + \frac{ac^4\sqrt{d}\ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{8} - \frac{ac^4\sqrt{-c^2dx^2+d}}{8} + \frac{b\sqrt{-d(c^2x^2-1)}\operatorname{arccosh}(cx)}{8(cx+1)(cx-1)}$

input `int((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^5,x,method=_RETURNVERBOSE)`

output

```
-1/4*a/d/x^4*(-c^2*d*x^2+d)^(3/2)-1/8*a*c^2/d/x^2*(-c^2*d*x^2+d)^(3/2)+1/8
*a*c^4*d^(1/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)-1/8*a*c^4*(-c^2*
d*x^2+d)^(1/2)+1/8*b*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)/(c*x-1)*arccosh(c*x)*c
^4+1/8*b*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)^(1/2)/x/(c*x-1)^(1/2)*c^3-3/8*b*(-
d*(c^2*x^2-1))^(1/2)/(c*x+1)/x^2/(c*x-1)*arccosh(c*x)*c^2-1/12*b*(-d*(c^2*
x^2-1))^(1/2)/(c*x+1)^(1/2)/x^3/(c*x-1)^(1/2)*c+1/4*b*(-d*(c^2*x^2-1))^(1/
2)/(c*x+1)/x^4/(c*x-1)*arccosh(c*x)-1/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)
^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))
)*c^4+1/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c
*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^4-1/8*I*b*(-d*(c^2*x^2-1))
^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1
/2)))*c^4+1/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog
(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^4
```

### 3.70.5 Fricas [F]

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^5} dx = \int \frac{\sqrt{-c^2dx^2+d}(b\operatorname{arccosh}(cx)+a)}{x^5} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^5,x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/x^5, x)`

---

3.70.  $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$

## 3.70.6 Sympy [F]

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)}(a + b \operatorname{acosh}(cx))}{x^5} dx$$

input `integrate((a+b*acosh(c*x))*(-c**2*d*x**2+d)**(1/2)/x**5,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))/x**5, x)`

## 3.70.7 Maxima [F]

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{\sqrt{-c^2 dx^2 + d}(b \operatorname{arcosh}(cx) + a)}{x^5} dx$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^5,x, algorithm="maxima")`

output `1/8*(c^4*sqrt(d)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - sqrt(-c^2*d*x^2 + d)*c^4 - (-c^2*d*x^2 + d)^(3/2)*c^2/(d*x^2) - 2*(-c^2*d*x^2 + d)^(3/2)/(d*x^4))*a + b*integrate(sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^5, x)`

## 3.70.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^5} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^5,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.70.  $\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x^5} dx$

**3.70.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{x^5} dx = \int \frac{(a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2}}{x^5} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^5, x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2))/x^5, x)`



### 3.71 $\int x^4(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.71.1	Optimal result	728
3.71.2	Mathematica [A] (warning: unable to verify)	729
3.71.3	Rubi [A] (verified)	729
3.71.4	Maple [B] (verified)	734
3.71.5	Fricas [F]	735
3.71.6	Sympy [F(-1)]	736
3.71.7	Maxima [F]	736
3.71.8	Giac [F]	736
3.71.9	Mupad [F(-1)]	737

#### 3.71.1 Optimal result

Integrand size = 27, antiderivative size = 360

$$\begin{aligned} \int x^4(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx &= \frac{3bdx^2\sqrt{d - c^2dx^2}}{256c^3\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &+ \frac{bdx^4\sqrt{d - c^2dx^2}}{256c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcdx^6\sqrt{d - c^2dx^2}}{32\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &+ \frac{bc^3dx^8\sqrt{d - c^2dx^2}}{64\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{3dx\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{128c^4} \\ &- \frac{dx^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{64c^2} + \frac{1}{16}dx^5\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) \\ &+ \frac{1}{8}x^5(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{3d\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{256bc^5\sqrt{-1 + cx}\sqrt{1 + cx}} \end{aligned}$$

output  $\frac{1}{8}x^5(-c^2dx^2+d)^{3/2}(a+b\operatorname{arccosh}(cx))-3/128dx^3(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}/c^4-1/64dx^5(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}/c^2+1/16dx^7(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}+3/256b^2dx^2(-c^2dx^2+d)^{1/2}/c^3/(cx-1)^{1/2}/(cx+1)^{1/2}+1/256b^2dx^4(-c^2dx^2+d)^{1/2}/c/(cx-1)^{1/2}/(cx+1)^{1/2}-1/32b^2cdx^6(-c^2dx^2+d)^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}+1/64b^2c^3dx^8(-c^2dx^2+d)^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}-3/256d^2(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{1/2}/b/c^5/(cx-1)^{1/2}/(cx+1)^{1/2}$

### 3.71.2 Mathematica [A] (warning: unable to verify)

Time = 3.47 (sec) , antiderivative size = 337, normalized size of antiderivative = 0.94

$$\int x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d \left( -576acx\sqrt{d - c^2 dx^2}(3 + 2c^2 x^2 - 24c^4 x^4 + 16c^6 x^6) - 1728a\sqrt{d} \arctan \left( \frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)} \right) + \operatorname{barccosh}(cx) \right)}{73728c^5}$$

input `Integrate[x^4*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]), x]`

output `(d*(-576*a*c*x*Sqrt[d - c^2*d*x^2]*(3 + 2*c^2*x^2 - 24*c^4*x^4 + 16*c^6*x^6) - 1728*a*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + (32*b*Sqrt[d - c^2*d*x^2]*(-72*ArcCosh[c*x]^2 + 18*Cosh[2*ArcCosh[c*x]] - 9*Cosh[4*ArcCosh[c*x]] - 2*Cosh[6*ArcCosh[c*x]] + 12*ArcCosh[c*x]*(-3*Sinh[2*ArcCosh[c*x]] + 3*Sinh[4*ArcCosh[c*x]] + Sinh[6*ArcCosh[c*x]]))) / (Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + (b*Sqrt[d - c^2*d*x^2]*(1440*ArcCosh[c*x]^2 - 576*Cosh[2*ArcCosh[c*x]] + 144*Cosh[4*ArcCosh[c*x]] + 64*Cosh[6*ArcCosh[c*x]] + 9*Cosh[8*ArcCosh[c*x]] - 24*ArcCosh[c*x]*(-48*Sinh[2*ArcCosh[c*x]] + 24*Sinh[4*ArcCosh[c*x]] + 16*Sinh[6*ArcCosh[c*x]] + 3*Sinh[8*ArcCosh[c*x]]))) / (Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))) / (73728*c^5)`

### 3.71.3 Rubi [A] (verified)

Time = 1.51 (sec) , antiderivative size = 325, normalized size of antiderivative = 0.90, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {6345, 25, 82, 244, 2009, 6341, 15, 6354, 15, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6345}$$

$$\frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd\sqrt{d - c^2 dx^2} \int -x^5(1 - cx)(cx + 1) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} +$$

$$\frac{1}{8}x^5(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{25}$$

---

3.71.  $\int x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int x^5 (1 - cx)(cx + 1) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{82} \\
& \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int x^5 (1 - c^2 x^2) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{244} \\
& \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int (x^5 - c^2 x^7) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2009} \\
& \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \\
& \quad \frac{bcd \left( \frac{x^6}{6} - \frac{c^2 x^8}{8} \right) \sqrt{d - c^2 dx^2}}{8\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6341} \\
& \frac{3}{8}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^4 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc \sqrt{d - c^2 dx^2} \int x^5 dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \right) + \\
& \quad \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^6}{6} - \frac{c^2 x^8}{8} \right) \sqrt{d - c^2 dx^2}}{8\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{15} \\
& \frac{3}{8}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^4 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^6 \sqrt{d - c^2 dx^2}}{36\sqrt{cx - 1}\sqrt{cx + 1}} \right) + \\
& \quad \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^6}{6} - \frac{c^2 x^8}{8} \right) \sqrt{d - c^2 dx^2}}{8\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6354}
\end{aligned}$$

$$\frac{3}{8}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{3 \int \frac{x^2(a + b \operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{b \int x^3 dx}{4c} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))}{4c^2} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{6}x^5 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))$$

$$- \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx)) - \frac{bcd \left( \frac{x^6}{6} - \frac{c^2 x^8}{8} \right) \sqrt{d - c^2 dx^2}}{8\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 15

$$\frac{3}{8}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{3 \int \frac{x^2(a + b \operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))}{4c^2} - \frac{bx^4}{16c} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{6}x^5 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))$$

$$- \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx)) - \frac{bcd \left( \frac{x^6}{6} - \frac{c^2 x^8}{8} \right) \sqrt{d - c^2 dx^2}}{8\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6354

$$\frac{3}{8}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{3 \left( \frac{\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{b \int x dx}{2c} + \frac{x \sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))}{2c^2} \right)}{4c^2} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))}{4c^2} - \frac{bx^4}{16c}$$

$$- \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx)) - \frac{bcd \left( \frac{x^6}{6} - \frac{c^2 x^8}{8} \right) \sqrt{d - c^2 dx^2}}{8\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 15

$$\begin{aligned}
& \frac{3}{8}d \left( \frac{\sqrt{d-c^2dx^2} \left( 3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx)) - \frac{bx^2}{4c}}{2c^2} \right) - \frac{bx^2}{4c} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx)) - \frac{bx^4}{16c}}{4c^2} - \frac{bx^4}{16c} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} \\
& \frac{1}{8}x^5(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd\left(\frac{x^6}{6} - \frac{c^2x^8}{8}\right)\sqrt{d-c^2dx^2}}{8\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow 6308 \\
& \frac{1}{8}x^5(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) + \\
& \frac{3}{8}d \left( \frac{\frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{\sqrt{d-c^2dx^2} \left( \frac{x^3\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} + \frac{3\left(\frac{(a+b\operatorname{arccosh}(cx))^2}{4bc^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx)) - \frac{bx^2}{4c}}{2c^2} \right)}{4c^2} \right)}{6\sqrt{cx-1}\sqrt{cx+1}}}{\frac{bcd\left(\frac{x^6}{6} - \frac{c^2x^8}{8}\right)\sqrt{d-c^2dx^2}}{8\sqrt{cx-1}\sqrt{cx+1}}} \right)
\end{aligned}$$

input `Int[x^4*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `-1/8*(b*c*d*Sqrt[d - c^2*d*x^2]*(x^6/6 - (c^2*x^8)/8))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^5*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/8 + (3*d*(-1/36*(b*c*x^6*Sqrt[d - c^2*d*x^2]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^5*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/6 - (Sqrt[d - c^2*d*x^2]*(-1/16*(b*x^4)/c + (x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(4*c^2) + (3*(-1/4*(b*x^2)/c + (x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(2*c^2) + (a + b*ArcCosh[c*x])^2/(4*b*c^3)))/(4*c^2)))/(6*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/8`

## 3.71.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 244 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

```
rule 6345 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*
x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m
+ 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m
+ 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1)
, x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n,
0] && GtQ[p, 0] && !LtQ[m, -1]
```

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_)^2)^(p_)*((d2_) + (e2_.)*(x_)^2)^(q_), x_Symbol] := Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(q + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^q*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^q/(
-1 + c*x)^q] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(q + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.71.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 782 vs. 2(304) = 608.

Time = 0.75 (sec) , antiderivative size = 783, normalized size of antiderivative = 2.18

method	result
default	$-\frac{ax^3(-c^2dx^2+d)^{\frac{5}{2}}}{8c^2d} - \frac{ax(-c^2dx^2+d)^{\frac{5}{2}}}{16c^4d} + \frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{64c^4} + \frac{3adx\sqrt{-c^2dx^2+d}}{128c^4} + \frac{3ad^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{128c^4\sqrt{c^2d}} + b\left(-\right)$
parts	$-\frac{ax^3(-c^2dx^2+d)^{\frac{5}{2}}}{8c^2d} - \frac{ax(-c^2dx^2+d)^{\frac{5}{2}}}{16c^4d} + \frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{64c^4} + \frac{3adx\sqrt{-c^2dx^2+d}}{128c^4} + \frac{3ad^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{128c^4\sqrt{c^2d}} + b\left(-\right)$

```
input int(x^4*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

---

3.71.  $\int x^4(d - c^2dx^2)^{3/2} (a + b\operatorname{arccosh}(cx)) dx$

output

```

-1/8*a*x^3*(-c^2*d*x^2+d)^(5/2)/c^2/d-1/16*a/c^4*x*(-c^2*d*x^2+d)^(5/2)/d+
1/64*a/c^4*x*(-c^2*d*x^2+d)^(3/2)+3/128*a/c^4*d*x*(-c^2*d*x^2+d)^(1/2)+3/1
28*a/c^4*d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b*
(-3/256*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/c^5*arccosh(c*x
)^2*d-1/16384*(-d*(c^2*x^2-1))^(1/2)*(128*c^9*x^9-320*c^7*x^7+128*(c*x-1)^(
1/2)*(c*x+1)^(1/2)*c^8*x^8+272*c^5*x^5-256*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^
6*x^6-88*c^3*x^3+160*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c*x-32*(c*x-1)^(
1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+8*arccosh(c*x
))*d/(c*x+1)/c^5/(c*x-1)+1/1024*(-d*(c^2*x^2-1))^(1/2)*(8*c^5*x^5-12*c^3*x
^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c*x-1)^(1/2)*(c*x+1)^(1/
2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+4*arccosh(c*x))*d/(c*x+1)/c^5/
(c*x-1)+1/1024*(-d*(c^2*x^2-1))^(1/2)*(-8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*
x^4+8*c^5*x^5+8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-12*c^3*x^3-(c*x-1)^(1/
2)*(c*x+1)^(1/2)+4*c*x)*(1+4*arccosh(c*x))*d/(c*x+1)/c^5/(c*x-1)-1/16384*(
-d*(c^2*x^2-1))^(1/2)*(-128*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^8*x^8+128*c^9*x^
9+256*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6-320*c^7*x^7-160*(c*x+1)^(1/2)*(c
*x-1)^(1/2)*c^4*x^4+272*c^5*x^5+32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-88*
c^3*x^3-(c*x-1)^(1/2)*(c*x+1)^(1/2)+8*c*x)*(1+8*arccosh(c*x))*d/(c*x+1)/c^
5/(c*x-1)

```

### 3.71.5 Fracas [F]

$$\int x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a) x^4 dx$$

input

```

integrate(x^4*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fracas
")

```

output

```

integral(-(a*c^2*d*x^6 - a*d*x^4 + (b*c^2*d*x^6 - b*d*x^4)*arccosh(c*x))*s
qrt(-c^2*d*x^2 + d), x)

```



**3.71.6 Sympy [F(-1)]**

Timed out.

$$\int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**4*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)`

output `Timed out`

**3.71.7 Maxima [F]**

$$\int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/128*(16*(-c^2*d*x^2 + d)^(5/2)*x^3/(c^2*d) - 2*(-c^2*d*x^2 + d)^(3/2)*x/c^4 + 8*(-c^2*d*x^2 + d)^(5/2)*x/(c^4*d) - 3*sqrt(-c^2*d*x^2 + d)*d*x/c^4 - 3*d^(3/2)*arcsin(c*x)/c^5)*a + b*integrate((-c^2*d*x^2 + d)^(3/2)*x^4*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)), x)`

**3.71.8 Giac [F]**

$$\int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)*x^4, x)`

**3.71.9 Mupad [F(-1)]**

Timed out.

$$\int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2} dx$$

input `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2),x)`output `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2), x)`

### 3.72 $\int x^2(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.72.1	Optimal result	738
3.72.2	Mathematica [A] (warning: unable to verify)	739
3.72.3	Rubi [A] (verified)	739
3.72.4	Maple [B] (verified)	743
3.72.5	Fricas [F]	744
3.72.6	Sympy [F(-1)]	745
3.72.7	Maxima [F]	745
3.72.8	Giac [F]	745
3.72.9	Mupad [F(-1)]	746

#### 3.72.1 Optimal result

Integrand size = 27, antiderivative size = 281

$$\begin{aligned} \int x^2(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx &= \frac{bdx^2\sqrt{d - c^2dx^2}}{32c\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &- \frac{7bcdx^4\sqrt{d - c^2dx^2}}{96\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx^6\sqrt{d - c^2dx^2}}{36\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &- \frac{dx\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{16c^2} + \frac{1}{8}dx^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) \\ &+ \frac{1}{6}x^3(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{d\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{32bc^3\sqrt{-1 + cx}\sqrt{1 + cx}} \end{aligned}$$

output  $\frac{1}{6}x^3(-c^2dx^2+d)^{3/2}(a+b\operatorname{arccosh}(cx))-\frac{1}{16}dx(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}/c^2+1/8dx^3(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}+1/32b^2dx^2(-c^2dx^2+d)^{1/2}/c/(cx-1)^{1/2}/(cx+1)^{1/2}-7/96b^2cdx^4(-c^2dx^2+d)^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}+1/36b^3c^3dx^6(-c^2dx^2+d)^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}-1/32d(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{1/2}/b/c^3/(cx-1)^{1/2}/(cx+1)^{1/2}$

### 3.72.2 Mathematica [A] (warning: unable to verify)

Time = 1.47 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.96

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d \left( -48acx\sqrt{d - c^2 dx^2}(3 - 14c^2 x^2 + 8c^4 x^4) - 144a\sqrt{d} \arctan \left( \frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)} \right) - \frac{18b\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)} \right) + \operatorname{barccosh}(cx)}{\dots}$$

input `Integrate[x^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `(d*(-48*a*c*x*Sqrt[d - c^2*d*x^2]*(3 - 14*c^2*x^2 + 8*c^4*x^4) - 144*a*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] - (18*b*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + (b*Sqrt[d - c^2*d*x^2]*(72*ArcCosh[c*x]^2 - 18*Cosh[2*ArcCosh[c*x]] + 9*Cosh[4*ArcCosh[c*x]] + 2*Cosh[6*ArcCosh[c*x]] - 12*ArcCosh[c*x]*(-3*Sinh[2*ArcCosh[c*x]] + 3*Sinh[4*ArcCosh[c*x]] + Sinh[6*ArcCosh[c*x]])))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))/(2304*c^3)`

### 3.72.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.96, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.370$ , Rules used = {6345, 25, 82, 244, 2009, 6341, 15, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6345} \\ & \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd\sqrt{d - c^2 dx^2} \int -x^3(1 - cx)(cx + 1) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \\ & \quad \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{25} \end{aligned}$$

$$\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int x^3 (1 - cx)(cx + 1) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))$$

↓ 82

$$\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int x^3 (1 - c^2 x^2) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))$$

↓ 244

$$\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int (x^3 - c^2 x^5) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))$$

↓ 2009

$$\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^4}{4} - \frac{c^2 x^6}{6} \right) \sqrt{d - c^2 dx^2}}{6\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6341

$$\frac{1}{2}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc \sqrt{d - c^2 dx^2} \int x^3 dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^4}{4} - \frac{c^2 x^6}{6} \right) \sqrt{d - c^2 dx^2}}{6\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 15

$$\frac{1}{2}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^4 \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}} \right) + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^4}{4} - \frac{c^2 x^6}{6} \right) \sqrt{d - c^2 dx^2}}{6\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6354

$$\frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \int x dx}{2c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \right. \\ \left. - \frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd\left(\frac{x^4}{4} - \frac{c^2x^6}{6}\right)\sqrt{d-c^2dx^2}}{6\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \downarrow 15 \\ \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} - \frac{bx^2}{4c} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \right. \\ \left. - \frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd\left(\frac{x^4}{4} - \frac{c^2x^6}{6}\right)\sqrt{d-c^2dx^2}}{6\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \downarrow 6308 \\ \frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) + \\ \frac{1}{2}d \left( \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{\sqrt{d-c^2dx^2} \left( \frac{(a+b\operatorname{arccosh}(cx))^2}{4bc^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} - \frac{bx^2}{4c} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{bcd\left(\frac{x^4}{4} - \frac{c^2x^6}{6}\right)\sqrt{d-c^2dx^2}}{6\sqrt{cx-1}\sqrt{cx+1}} \right)$$

input `Int[x^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]), x]`

output `-1/6*(b*c*d*Sqrt[d - c^2*d*x^2]*(x^4/4 - (c^2*x^6)/6))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^3*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/6 + (d*(-1/16*(b*c*x^4*Sqrt[d - c^2*d*x^2]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/4 - (Sqrt[d - c^2*d*x^2]*(-1/4*(b*x^2)/c + (x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(2*c^2) + (a + b*ArcCosh[c*x])^2/(4*b*c^3)))/(4*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/2`

## 3.72.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 244 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

```
rule 6345 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^(m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]
```

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NEqQ[m + 2*p + 1, 0]
```

### 3.72.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 882 vs.  $2(237) = 474$ .

Time = 0.71 (sec) , antiderivative size = 883, normalized size of antiderivative = 3.14

method	result
default	$-\frac{ax(-c^2dx^2+d)^{\frac{5}{2}}}{6c^2d} + \frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{24c^2} + \frac{adx\sqrt{-c^2dx^2+d}}{16c^2} + \frac{a d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{16c^2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)}{32\sqrt{cx-1}\sqrt{cx+1}c^3}\right)$
parts	$-\frac{ax(-c^2dx^2+d)^{\frac{5}{2}}}{6c^2d} + \frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{24c^2} + \frac{adx\sqrt{-c^2dx^2+d}}{16c^2} + \frac{a d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{16c^2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)}{32\sqrt{cx-1}\sqrt{cx+1}c^3}\right)$

```
input int(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```



output `-1/6*a*x*(-c^2*d*x^2+d)^(5/2)/c^2/d+1/24*a/c^2*x*(-c^2*d*x^2+d)^(3/2)+1/16*a/c^2*d*x*(-c^2*d*x^2+d)^(1/2)+1/16*a/c^2*d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b*(-1/32*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/c^3*arccosh(c*x)^2*d-1/2304*(-d*(c^2*x^2-1))^(1/2)*(32*c^7*x^7-64*c^5*x^5+32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6+38*c^3*x^3-48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-6*c*x+18*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+6*arccosh(c*x))*d/(c*x+1)/c^3/(c*x-1)+1/512*(-d*(c^2*x^2-1))^(1/2)*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+4*arccosh(c*x))*d/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+2*arccosh(c*x))*d/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)-2*c*x)*(1+2*arccosh(c*x))*d/(c*x+1)/c^3/(c*x-1)+1/512*(-d*(c^2*x^2-1))^(1/2)*(-8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c^5*x^5+8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-12*c^3*x^3-(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c*x)*(1+4*arccosh(c*x))*d/(c*x+1)/c^3/(c*x-1)-1/2304*(-d*(c^2*x^2-1))^(1/2)*(-32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6+32*c^7*x^7+48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-64*c^5*x^5-18*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+38*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)-6*c*x)*(1+6*arccosh(c*x))*d/(c...`

### 3.72.5 Fricas [F]

$$\int x^2(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a) x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^4 - a*d*x^2 + (b*c^2*d*x^4 - b*d*x^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

**3.72.6 Sympy [F(-1)]**

Timed out.

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**2*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)`output `Timed out`**3.72.7 Maxima [F]**

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a) x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`output `1/48*a*(2*(-c^2*d*x^2 + d)^(3/2)*x/c^2 - 8*(-c^2*d*x^2 + d)^(5/2)*x/(c^2*d) + 3*sqrt(-c^2*d*x^2 + d)*d*x/c^2 + 3*d^(3/2)*arcsin(c*x)/c^3) + b*integrate((-c^2*d*x^2 + d)^(3/2)*x^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)), x)`**3.72.8 Giac [F]**

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a) x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)*x^2, x)`

**3.72.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2} dx$$

input `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2),x)`output `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2), x)`

### 3.73 $\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.73.1	Optimal result	747
3.73.2	Mathematica [A] (warning: unable to verify)	748
3.73.3	Rubi [A] (verified)	748
3.73.4	Maple [B] (verified)	751
3.73.5	Fricas [F]	752
3.73.6	Sympy [F]	752
3.73.7	Maxima [F]	753
3.73.8	Giac [F(-2)]	753
3.73.9	Mupad [F(-1)]	753

#### 3.73.1 Optimal result

Integrand size = 24, antiderivative size = 200

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = -\frac{5bcdx^2\sqrt{d - c^2dx^2}}{16\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx^4\sqrt{d - c^2dx^2}}{16\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{3}{8}dx\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) + \frac{1}{4}x(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx)) - \frac{3d\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{16bc\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output `1/4*x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))+3/8*d*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)-5/16*b*c*d*x^2*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/16*b*c^3*d*x^4*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-3/16*d*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)`

### 3.73.2 Mathematica [A] (warning: unable to verify)

Time = 0.96 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.18

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{1}{8} adx(-5 + 2c^2 x^2) \sqrt{d - c^2 dx^2} - \frac{3ad^{3/2} \arctan\left(\frac{cx\sqrt{d-c^2 dx^2}}{\sqrt{d(-1+c^2 x^2)}}\right)}{8c}$$

$$- \frac{bd\sqrt{d - c^2 dx^2}(2\operatorname{arccosh}(cx)^2 + \cosh(2\operatorname{arccosh}(cx)) - 2\operatorname{arccosh}(cx) \sinh(2\operatorname{arccosh}(cx)))}{8c\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}$$

$$+ \frac{bd\sqrt{d - c^2 dx^2}(8\operatorname{arccosh}(cx)^2 + \cosh(4\operatorname{arccosh}(cx)) - 4\operatorname{arccosh}(cx) \sinh(4\operatorname{arccosh}(cx)))}{128c\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}$$

input `Integrate[(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]), x]`

output `-1/8*(a*d*x*(-5 + 2*c^2*x^2)*Sqrt[d - c^2*d*x^2]) - (3*a*d^(3/2)*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))])/(8*c) - (b*d*Sqrt[d - c^2*d*x^2]*(2*ArcCosh[c*x]^2 + Cosh[2*ArcCosh[c*x]] - 2*ArcCosh[c*x]*Sinh[2*ArcCosh[c*x]]))/(8*c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + (b*d*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]))/(128*c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`

### 3.73.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.08, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6312, 25, 82, 244, 2009, 6310, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6312}$$

$$\frac{3}{4} d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd\sqrt{d - c^2 dx^2} \int -x(1 - cx)(cx + 1) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} +$$

$$\frac{1}{4} x (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))$$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x(1 - cx)(cx + 1) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \\
& \downarrow 82 \\
& \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x(1 - c^2 x^2) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \\
& \quad \operatorname{barccosh}(cx)) \\
& \downarrow 244 \\
& \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int (x - c^2 x^3) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \\
& \quad \operatorname{barccosh}(cx)) \\
& \downarrow 2009 \\
& \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \\
& \quad \frac{bcd\left(\frac{x^2}{2} - \frac{c^2 x^4}{4}\right) \sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \downarrow 6310 \\
& \frac{3}{4}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \right) + \\
& \quad \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd\left(\frac{x^2}{2} - \frac{c^2 x^4}{4}\right) \sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \downarrow 15 \\
& \frac{3}{4}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^2\sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right) + \\
& \quad \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd\left(\frac{x^2}{2} - \frac{c^2 x^4}{4}\right) \sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \downarrow 6308
\end{aligned}$$

$$\frac{1}{4}x(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx)) + \frac{3}{4}d \left( \frac{1}{2}x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) - \frac{\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{4bc\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bcx^2\sqrt{d - c^2dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right) - \frac{bcd \left( \frac{x^2}{2} - \frac{c^2x^4}{4} \right) \sqrt{d - c^2dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}}$$

input `Int[(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `-1/4*(b*c*d*Sqrt[d - c^2*d*x^2]*(x^2/2 - (c^2*x^4)/4))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/4 + (3*d*(-1/4*(b*c*x^2*Sqrt[d - c^2*d*x^2]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(4*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/4`

### 3.73.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 244 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 6308 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]
```

```
rule 6310 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]
```

```
rule 6312 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]
```

### 3.73.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 545 vs. 2(168) = 336.

Time = 0.92 (sec) , antiderivative size = 546, normalized size of antiderivative = 2.73

method	result
default	$\frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{4} + \frac{3adx\sqrt{-c^2dx^2+d}}{8} + \frac{3ad^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{8\sqrt{c^2d}} + b\left(-\frac{3\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2d}{16\sqrt{cx-1}\sqrt{cx+1}c} - \frac{\sqrt{-d(c^2x^2-1)}}{16\sqrt{cx-1}\sqrt{cx+1}c}\right)$
parts	$\frac{ax(-c^2dx^2+d)^{\frac{3}{2}}}{4} + \frac{3adx\sqrt{-c^2dx^2+d}}{8} + \frac{3ad^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{8\sqrt{c^2d}} + b\left(-\frac{3\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2d}{16\sqrt{cx-1}\sqrt{cx+1}c} - \frac{\sqrt{-d(c^2x^2-1)}}{16\sqrt{cx-1}\sqrt{cx+1}c}\right)$

```
input int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

---

3.73.  $\int (d - c^2dx^2)^{3/2} (a + b\operatorname{arccosh}(cx)) dx$



output `1/4*a*x*(-c^2*d*x^2+d)^(3/2)+3/8*a*d*x*(-c^2*d*x^2+d)^(1/2)+3/8*a*d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b*(-3/16*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/c*arccosh(c*x)^2*d-1/256*(-d*(c^2*x^2-1))^(1/2)*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+4*arccosh(c*x))*d/(c*x-1)/(c*x+1)/c+1/16*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+2*arccosh(c*x))*d/(c*x-1)/(c*x+1)/c+1/16*(-d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2))-2*c*x*(1+2*arccosh(c*x))*d/(c*x-1)/(c*x+1)/c-1/256*(-d*(c^2*x^2-1))^(1/2)*(-8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c^5*x^5+8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-12*c^3*x^3-(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c*x)*(1+4*arccosh(c*x))*d/(c*x-1)/(c*x+1)/c)`

### 3.73.5 Fricas [F]

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a) dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

### 3.73.6 Sympy [F]

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x)), x)`

**3.73.7 Maxima [F]**

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a) dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/8*(2*(-c^2*d*x^2 + d)^(3/2)*x + 3*sqrt(-c^2*d*x^2 + d)*d*x + 3*d^(3/2)*arcsin(c*x)/c)*a + b*integrate((-c^2*d*x^2 + d)^(3/2)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1), x)`

**3.73.8 Giac [F(-2)]**

Exception generated.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.73.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2} dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2),x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2), x)`

### 3.74 $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.74.1	Optimal result	754
3.74.2	Mathematica [A] (warning: unable to verify)	754
3.74.3	Rubi [A] (verified)	755
3.74.4	Maple [A] (verified)	758
3.74.5	Fricas [F]	759
3.74.6	Sympy [F]	759
3.74.7	Maxima [F]	759
3.74.8	Giac [F(-2)]	760
3.74.9	Mupad [F(-1)]	760

#### 3.74.1 Optimal result

Integrand size = 27, antiderivative size = 197

$$\int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{x^2} dx = \frac{bc^3dx^2\sqrt{d - c^2dx^2}}{4\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{3}{2}c^2dx\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx)) - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{x} + \frac{3cd\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{4b\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bcd\sqrt{d - c^2dx^2} \log(x)}{\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output 
$$-(c^2d^2x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))/x-3/2*c^2*d*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}+1/4*b*c^3*d*x^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3/4*c*d*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/b/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+b*c*d*\ln(x)*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

#### 3.74.2 Mathematica [A] (warning: unable to verify)

Time = 0.97 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.13

$$\int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{x^2} dx = \frac{1}{8} \left( -\frac{4ad(2 + c^2x^2)\sqrt{d - c^2dx^2}}{x} + 12acd^{3/2} \arctan \left( \frac{cx\sqrt{d - c^2dx^2}}{\sqrt{d}(-1 + c^2x^2)} \right) + 4bcd\sqrt{d - c^2dx^2} \left( -\frac{2\operatorname{arccosh}(cx)}{cx} + \frac{\operatorname{arccosh}(cx)^2 + 2\log(cx)}{\sqrt{\frac{-1+cx}{1+cx}}(1 + cx)} \right) \right) + bcd$$

---

3.74.  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^2} dx$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^2,x]`

output `((-4*a*d*(2 + c^2*x^2)*Sqrt[d - c^2*d*x^2])/x + 12*a*c*d^(3/2)*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + 4*b*c*d*Sqrt[d - c^2*d*x^2]*((-2*ArcCosh[c*x])/(c*x) + (ArcCosh[c*x]^2 + 2*Log[c*x])/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))) + (b*c*d*Sqrt[d - c^2*d*x^2]*(2*ArcCosh[c*x]^2 + Cosh[2*ArcCosh[c*x]] - 2*ArcCosh[c*x]*Sinh[2*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))/8`

### 3.74.3 Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.06, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$ , Rules used = {6343, 25, 82, 244, 2009, 6310, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx \\
 & \quad \downarrow \text{6343} \\
 & -3c^2 d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)}{x} dx}{\sqrt{cx-1} \sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{25} \\
 & -3c^2 d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd \sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)}{x} dx}{\sqrt{cx-1} \sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{82} \\
 & -3c^2 d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd \sqrt{d - c^2 dx^2} \int \frac{1-c^2 x^2}{x} dx}{\sqrt{cx-1} \sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{244}
 \end{aligned}$$

---

3.74.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx$

$$\begin{aligned}
& -3c^2d \int \sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))dx + \frac{bcd\sqrt{d - c^2dx^2} \int \left(\frac{1}{x} - c^2x\right) dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \\
& \quad \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} \\
& \quad \downarrow \text{2009} \\
& -3c^2d \int \sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))dx - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} + \\
& \quad \frac{bcd\sqrt{d - c^2dx^2} \left(\log(x) - \frac{c^2x^2}{2}\right)}{\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6310} \\
& -3c^2d \left( -\frac{\sqrt{d - c^2dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2dx^2} \int x dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{2}x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) \right) - \\
& \quad \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} + \frac{bcd\sqrt{d - c^2dx^2} \left(\log(x) - \frac{c^2x^2}{2}\right)}{\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{15} \\
& -3c^2d \left( -\frac{\sqrt{d - c^2dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{2}x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) - \frac{bcx^2\sqrt{d - c^2dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right) - \\
& \quad \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} + \frac{bcd\sqrt{d - c^2dx^2} \left(\log(x) - \frac{c^2x^2}{2}\right)}{\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6308} \\
& -3c^2d \left( \frac{1}{2}x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) - \frac{\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{4bc\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bcx^2\sqrt{d - c^2dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right) - \\
& \quad \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} + \frac{bcd\sqrt{d - c^2dx^2} \left(\log(x) - \frac{c^2x^2}{2}\right)}{\sqrt{cx - 1}\sqrt{cx + 1}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^2,x]`

```
output -(((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x) - 3*c^2*d*(-1/4*(b*c*x^2
*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*Sqrt[d - c^2*d*x
^2]*(a + b*ArcCosh[c*x]))/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)
/(4*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (b*c*d*Sqrt[d - c^2*d*x^2]*(-1/2*
(c^2*x^2) + Log[x]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])
```

### 3.74.3.1 Defintions of rubi rules used

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 82 Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)
)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d,
e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]
```

```
rule 244 Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand
Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p
, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6308 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sq
rt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 +
c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[
c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1
] && EqQ[e2, (-c)*d2] && NeQ[n, -1]
```

```
rule 6310 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_
Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(
1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcC
osh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sq
rt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^
(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n
, 0]
```

```
rule 6343 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m
+ 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(
m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x],
x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && G
tQ[p, 0] && LtQ[m, -1]
```

### 3.74.4 Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.24

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{dx} - ac^2x(-c^2dx^2+d)^{\frac{3}{2}} - \frac{3ac^2dx\sqrt{-c^2dx^2+d}}{2} - \frac{3ac^2d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2\sqrt{c^2d}} + \frac{b\sqrt{-d(c^2x^2-1)}}{2\sqrt{c^2d}}$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{dx} - ac^2x(-c^2dx^2+d)^{\frac{3}{2}} - \frac{3ac^2dx\sqrt{-c^2dx^2+d}}{2} - \frac{3ac^2d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2\sqrt{c^2d}} + \frac{b\sqrt{-d(c^2x^2-1)}}{2\sqrt{c^2d}}$

```
input int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)
```

```
output -a/d/x*(-c^2*d*x^2+d)^(5/2)-a*c^2*x*(-c^2*d*x^2+d)^(3/2)-3/2*a*c^2*d*x*(-c
^2*d*x^2+d)^(1/2)-3/2*a*c^2*d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2
*d*x^2+d)^(1/2))+1/8*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/
x*(-4*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2+2*c^3*x^3+6*arccosh
(c*x)^2*x*c-8*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-8*c*x*arccosh(c*x)+
8*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-c*x)*d
```

$$3.74. \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^2} dx$$

**3.74.5 Fracas [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^2,x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^2, x)`

**3.74.6 Sympy [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(-d(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))}{x^2} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**2,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))/x**2, x)`

**3.74.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output `-1/2*(3*sqrt(-c^2*d*x^2 + d)*c^2*d*x + 3*c*d^(3/2)*arcsin(c*x) + 2*(-c^2*d*x^2 + d)^(3/2)/x)*a + b*integrate((-c^2*d*x^2 + d)^(3/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^2, x)`



**3.74.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.74.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^2,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^2, x)`

**3.75** 
$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^4} dx$$

3.75.1	Optimal result	761
3.75.2	Mathematica [A] (warning: unable to verify)	761
3.75.3	Rubi [A] (verified)	762
3.75.4	Maple [A] (verified)	765
3.75.5	Fricas [F]	765
3.75.6	Sympy [F]	766
3.75.7	Maxima [F]	766
3.75.8	Giac [F(-2)]	766
3.75.9	Mupad [F(-1)]	767

**3.75.1 Optimal result**

Integrand size = 27, antiderivative size = 203

$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^4} dx = -\frac{bcd\sqrt{d-c^2dx^2}}{6x^2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{c^2d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x} - \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{3x^3} - \frac{c^3d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2b\sqrt{-1+cx}\sqrt{1+cx}} - \frac{4bc^3d\sqrt{d-c^2dx^2}\log(x)}{3\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^3+c^2*d*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x-1/6*b*c*d*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/2*c^3*d*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/(c*x-1)^(1/2)/(c*x+1)^(1/2)-4/3*b*c^3*d*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.75.2 Mathematica [A] (warning: unable to verify)**

Time = 0.83 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.28

$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^4} dx = \frac{-2bd^2\sqrt{\frac{-1+cx}{1+cx}}(1-5c^2x^2+4c^4x^4)\operatorname{arccosh}(cx)+3bc^3d^2x^3(-1+cx)}{x^4}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^4,x]`

output `(-2*b*d^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 - 5*c^2*x^2 + 4*c^4*x^4)*ArcCosh[c*x] + 3*b*c^3*d^2*x^3*(-1 + c*x)*ArcCosh[c*x]^2 - 6*a*c^3*d^(3/2)*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + d^2*(b*c*x*(-1 + c*x) - 2*a*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 - 5*c^2*x^2 + 4*c^4*x^4) + 8*b*c^3*x^3*(-1 + c*x)*Log[c*x])/ (6*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2])`

### 3.75.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$ , Rules used = {6343, 25, 82, 244, 2009, 6339, 14, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx \\
 & \quad \downarrow \text{6343} \\
 & c^2(-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^2} dx - \frac{bcd\sqrt{d - c^2 dx^2} \int -\frac{(1-cx)(cx+1)}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{25} \\
 & c^2(-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{82} \\
 & c^2(-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{1-c^2 x^2}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3x^3} \\
 & \quad \downarrow \text{244}
 \end{aligned}$$

---

3.75.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx$

$$\begin{aligned}
& c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \left(\frac{1}{x^3} - \frac{c^2}{x}\right) dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{2009} \\
& c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^2} dx - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \\
& \quad \frac{bcd\sqrt{d-c^2dx^2}\left(c^2(-\log(x)) - \frac{1}{2x^2}\right)}{3\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{6339} \\
& c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{bc\sqrt{d-c^2dx^2} \int \frac{1}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd\sqrt{d-c^2dx^2}\left(c^2(-\log(x)) - \frac{1}{2x^2}\right)}{3\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{14} \\
& c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bc \log(x)\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd\sqrt{d-c^2dx^2}\left(c^2(-\log(x)) - \frac{1}{2x^2}\right)}{3\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{6308} \\
& c^2(-d) \left( \frac{c\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2b\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bc \log(x)\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd\sqrt{d-c^2dx^2}\left(c^2(-\log(x)) - \frac{1}{2x^2}\right)}{3\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^3 + (b*c*d*Sqrt[d - c^2*d*x^2]*(-1/2*1/x^2 - c^2*Log[x]))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - c^2*d*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x) + (c*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (b*c*Sqrt[d - c^2*d*x^2]*Log[x])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]))`

---

3.75.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x^4} dx$

## 3.75.3.1 Defintions of rubi rules used

- rule 14 `Int[(a_)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 82 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(Sqrt[(d1_) + (e1_)*(x_)]*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6339 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`

```
rule 6343 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1]
```

### 3.75.4 Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.33

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{3dx^3} + \frac{2ac^2(-c^2dx^2+d)^{\frac{5}{2}}}{3dx} + \frac{2ac^4x(-c^2dx^2+d)^{\frac{3}{2}}}{3} + ac^4dx\sqrt{-c^2dx^2+d} + \frac{ac^4d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{\sqrt{c^2d}}$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{3dx^3} + \frac{2ac^2(-c^2dx^2+d)^{\frac{5}{2}}}{3dx} + \frac{2ac^4x(-c^2dx^2+d)^{\frac{3}{2}}}{3} + ac^4dx\sqrt{-c^2dx^2+d} + \frac{ac^4d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{\sqrt{c^2d}}$

```
input int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)
```

```
output -1/3*a/d/x^3*(-c^2*d*x^2+d)^(5/2)+2/3*a*c^2/d/x*(-c^2*d*x^2+d)^(5/2)+2/3*a*c^4*x*(-c^2*d*x^2+d)^(3/2)+a*c^4*d*x*(-c^2*d*x^2+d)^(1/2)+a*c^4*d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))-1/6*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^3*(3*arccosh(c*x)^2*x^3*c^3-8*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-8*c^3*x^3*arccosh(c*x)+8*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3+2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*d
```

### 3.75.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a)}{x^4} dx$$

```
input integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")
```

---

3.75. 
$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^4} dx$$

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^4, x)`

### 3.75.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx))}{x^4} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**4,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))/x**4, x)`

### 3.75.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a)}{x^4} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")`

output `1/3*(3*sqrt(-c^2*d*x^2 + d)*c^4*d*x + 3*c^3*d^(3/2)*arcsin(c*x) + 2*(-c^2*d*x^2 + d)^(3/2)*c^2/x - (-c^2*d*x^2 + d)^(5/2)/(d*x^3))*a + b*integrate((-c^2*d*x^2 + d)^(3/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^4, x)`

### 3.75.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")`

---

3.75.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx$

output Exception raised: TypeError >> an error occurred running a Giac command:IN  
 PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const  
 index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.75.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^4,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^4, x)`



**3.76**  $\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^6} dx$

3.76.1	Optimal result	768
3.76.2	Mathematica [A] (verified)	768
3.76.3	Rubi [A] (verified)	769
3.76.4	Maple [B] (verified)	771
3.76.5	Fricas [A] (verification not implemented)	772
3.76.6	Sympy [F(-1)]	772
3.76.7	Maxima [C] (verification not implemented)	773
3.76.8	Giac [F(-2)]	773
3.76.9	Mupad [F(-1)]	774

**3.76.1 Optimal result**

Integrand size = 27, antiderivative size = 166

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^6} dx = -\frac{bcd\sqrt{d - c^2 dx^2}}{20x^4\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3 d\sqrt{d - c^2 dx^2}}{5x^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{5dx^5} + \frac{bc^5 d\sqrt{d - c^2 dx^2} \log(x)}{5\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
-1/5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^5-1/20*b*c*d*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/5*b*c^3*d*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/5*b*c^5*d*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.76.2 Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.57

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^6} dx = \frac{d\sqrt{d - c^2 dx^2} \left( \frac{(-1+cx)^{5/2}(1+cx)^{5/2}(a+b \operatorname{arccosh}(cx))}{x^5} - bc \left( -\frac{1}{4x^4} + \frac{c^2}{x^2} + c^4 \log(x) \right) \right)}{5\sqrt{-1 + cx}\sqrt{1 + cx}}$$

input

```
Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^6,x]
```

3.76.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^6} dx$

output  $-1/5*(d*\text{Sqrt}[d - c^2*d*x^2]*((-1 + c*x)^{(5/2)}*(1 + c*x)^{(5/2)}*(a + b*\text{ArcCosh}[c*x]))/x^5 - b*c*(-1/4*1/x^4 + c^2/x^2 + c^4*\text{Log}[x]))/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])$

### 3.76.3 Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.59, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6332, 82, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))}{x^6} dx \\
 & \quad \downarrow \text{6332} \\
 & \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2(cx+1)^2}{x^5} dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5dx^5} \\
 & \quad \downarrow \text{82} \\
 & \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2}{x^5} dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5dx^5} \\
 & \quad \downarrow \text{243} \\
 & \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2}{x^6} dx^2}{10\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5dx^5} \\
 & \quad \downarrow \text{49} \\
 & \frac{bcd\sqrt{d - c^2 dx^2} \int \left( \frac{c^4}{x^2} - \frac{2c^2}{x^4} + \frac{1}{x^6} \right) dx^2}{10\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5dx^5} \\
 & \quad \downarrow \text{2009} \\
 & \frac{bcd\sqrt{d - c^2 dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5dx^5}
 \end{aligned}$$

input  $\text{Int}[(d - c^2*d*x^2)^{(3/2)}*(a + b*\text{ArcCosh}[c*x])/x^6, x]$

---

3.76.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))}{x^6} dx$

output 
$$-1/5*((d - c^2*d*x^2)^{(5/2)}*(a + b*\text{ArcCosh}[c*x]))/(d*x^5) + (b*c*d*\text{Sqrt}[d - c^2*d*x^2]*(-1/2*1/x^4 + (2*c^2)/x^2 + c^4*\text{Log}[x^2]))/(10*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])$$

### 3.76.3.1 Defintions of rubi rules used

rule 49 
$$\text{Int}[(a + b*x)^m*(c + d*x)^n, x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, x\} \&\& \text{IGtQ}[m, 0] \&\& \text{IGtQ}[m + n + 2, 0]$$

rule 82 
$$\text{Int}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x] \rightarrow \text{Int}[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p, x\} \&\& \text{EqQ}[b*c + a*d, 0] \&\& \text{EqQ}[n, m] \&\& \text{IntegerQ}[m]$$

rule 243 
$$\text{Int}[(x + a + b*x^2)^p, x\_Symbol] \rightarrow \text{Simp}[1/2 \text{Subst}[\text{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, m, p, x\} \&\& \text{IntegerQ}[(m-1)/2]$$

rule 2009 
$$\text{Int}[u, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 6332 
$$\text{Int}[(a + \text{ArcCosh}[c*x]*b)^n*(f*x)^m*(d + e*x^2)^p, x\_Symbol] \rightarrow \text{Simp}[(f*x)^{m+1}*(d + e*x^2)^{p+1}*(a + b*\text{ArcCosh}[c*x])^n/(d*f*(m+1)), x] + \text{Simp}[b*c*(n/(f*(m+1)))*\text{Simp}[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] \text{Int}[(f*x)^{m+1}*(1 + c*x)^{p+1/2}*(-1 + c*x)^{p+1/2}*(a + b*\text{ArcCosh}[c*x])^{n-1}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, p, x\} \&\& \text{EqQ}[c^2*d + e, 0] \&\& \text{GtQ}[n, 0] \&\& \text{EqQ}[m + 2*p + 3, 0] \&\& \text{NeQ}[m, -1]$$

### 3.76.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2170 vs.  $2(138) = 276$ .

Time = 1.22 (sec) , antiderivative size = 2171, normalized size of antiderivative = 13.08

method	result	size
default	Expression too large to display	2171
parts	Expression too large to display	2171

```
input int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^6,x,method=_RETURNVERBOSE)
```

```
output 1/5*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6*x^6+10*c^4*x^4-5*c^2*x^2+
1)*x^7*c^12-9/20*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6*x^6+10*c^4*x
^4-5*c^2*x^2+1)*x^5*c^10+3/10*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6
*x^6+10*c^4*x^4-5*c^2*x^2+1)*x^3*c^8-1/20*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^
8*x^8-10*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)*x*c^6+1/5*b*(-d*(c^2*x^2-1))^(1/2
)*d/(5*c^8*x^8-10*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/(c*x+1)^(1/2)/(c*x-1)^(1
/2)*arccosh(c*x)*c^5+5/2*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6*x^6+
10*c^4*x^4-5*c^2*x^2+1)*x^2/(c*x+1)^(1/2)/(c*x-1)^(1/2)*c^7+9/20*b*(-d*(c^
2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/x^2/(c*x+1
)^(1/2)/(c*x-1)^(1/2)*c^3-1/20*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^
6*x^6+10*c^4*x^4-5*c^2*x^2+1)/x^4/(c*x+1)^(1/2)/(c*x-1)^(1/2)*c-56/5*b*(-d
*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)*x/(c*x
+1)/(c*x-1)*arccosh(c*x)*c^6+28/5*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10
*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/x/(c*x+1)/(c*x-1)*arccosh(c*x)*c^4-8/5*b*
(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/x^3
/(c*x+1)/(c*x-1)*arccosh(c*x)*c^2-b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10
*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)*x^9/(c*x+1)/(c*x-1)*arccosh(c*x)*c^14-1/5
*a/d/x^5*(-c^2*d*x^2+d)^(5/2)+5*b*(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c
^6*x^6+10*c^4*x^4-5*c^2*x^2+1)*x^7/(c*x+1)/(c*x-1)*arccosh(c*x)*c^12-11*b*
(-d*(c^2*x^2-1))^(1/2)*d/(5*c^8*x^8-10*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)*...
```

---

3.76.  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^6} dx$

**3.76.5 Fracas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 572, normalized size of antiderivative = 3.45

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \left[ -\frac{4(bc^6 dx^6 - 3bc^4 dx^4 + 3bc^2 dx^2 - bd)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{-c^2 dx^2 + d})}{x^6} \right]$$

```
input integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^6,x, algorithm="fracas")
```

```
output [-1/20*(4*(b*c^6*d*x^6 - 3*b*c^4*d*x^4 + 3*b*c^2*d*x^2 - b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - 2*(b*c^7*d*x^7 - b*c^5*d*x^5)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 - sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1))*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) - (4*b*c^3*d*x^3 - (4*b*c^3 - b*c)*d*x^5 - b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 4*(a*c^6*d*x^6 - 3*a*c^4*d*x^4 + 3*a*c^2*d*x^2 - a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^7 - x^5), 1/20*(4*(b*c^7*d*x^7 - b*c^5*d*x^5)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 4*(b*c^6*d*x^6 - 3*b*c^4*d*x^4 + 3*b*c^2*d*x^2 - b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + (4*b*c^3*d*x^3 - (4*b*c^3 - b*c)*d*x^5 - b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 4*(a*c^6*d*x^6 - 3*a*c^4*d*x^4 + 3*a*c^2*d*x^2 - a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^7 - x^5)]
```

**3.76.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \text{Timed out}$$

```
input integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**6,x)
```

```
output Timed out
```

**3.76.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.14

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^6} dx =$$

$$\frac{\left(2 c^6 d^3 \sqrt{-\frac{1}{c^4 d}} \log\left(x^2 - \frac{1}{c^2}\right) + 2i (-1)^{-2 c^2 dx^2 + 2d} c^4 d^{\frac{5}{2}} \log\left(-2 c^2 d + \frac{2d}{x^2}\right) + \frac{3\sqrt{-c^4 dx^4 + 2c^2 dx^2 - dc^2 d^2}}{x^2} - \frac{\sqrt{-c^4 dx^4}}{x^2}\right)}{20 d}$$

$$- \frac{(-c^2 dx^2 + d)^{\frac{5}{2}} b \operatorname{arccosh}(cx)}{5 dx^5} - \frac{(-c^2 dx^2 + d)^{\frac{5}{2}} a}{5 dx^5}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^6,x, algorithm="maxima")`

output `-1/20*(2*c^6*d^3*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2) + 2*I*(-1)^(-2*c^2*d*x^2 + 2*d)*c^4*d^(5/2)*log(-2*c^2*d + 2*d/x^2) + 3*sqrt(-c^4*d*x^4 + 2*c^2*d*x^2 - d)*c^2*d^2/x^2 - sqrt(-c^4*d*x^4 + 2*c^2*d*x^2 - d)*d^2/x^4)*b*c/d - 1/5*(-c^2*d*x^2 + d)^(5/2)*b*arccosh(c*x)/(d*x^5) - 1/5*(-c^2*d*x^2 + d)^(5/2)*a/(d*x^5)`

**3.76.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^6,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.76.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^6} dx$

**3.76.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^6} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^6,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^6, x)`

**3.77** 
$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^8} dx$$

3.77.1	Optimal result	775
3.77.2	Mathematica [A] (verified)	775
3.77.3	Rubi [A] (verified)	776
3.77.4	Maple [B] (verified)	778
3.77.5	Fricas [A] (verification not implemented)	779
3.77.6	Sympy [F(-1)]	779
3.77.7	Maxima [A] (verification not implemented)	780
3.77.8	Giac [F(-2)]	780
3.77.9	Mupad [F(-1)]	781

**3.77.1 Optimal result**

Integrand size = 27, antiderivative size = 247

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^8} dx = -\frac{bcd\sqrt{d - c^2 dx^2}}{42x^6\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{2bc^3d\sqrt{d - c^2 dx^2}}{35x^4\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bc^5d\sqrt{d - c^2 dx^2}}{70x^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{7dx^7} - \frac{2c^2(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{35dx^5} + \frac{2bc^7d\sqrt{d - c^2 dx^2} \log(x)}{35\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
-1/7*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^7-2/35*c^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^5-1/42*b*c*d*(-c^2*d*x^2+d)^(1/2)/x^6/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/35*b*c^3*d*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/70*b*c^5*d*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/35*b*c^7*d*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.77.2 Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.55

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^8} dx = \frac{d\sqrt{d - c^2 dx^2} (30(-1 + cx)^{5/2} (1 + cx)^{5/2} (a + b \operatorname{arccosh}(cx)) + 12c^2 x^2 (-1 + cx)^{5/2} (1 + cx)^{5/2} (a + b \operatorname{arccosh}(cx)))}{210x^7\sqrt{-1 + cx}\sqrt{1 + cx}}$$

---

3.77. 
$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^8} dx$$



input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^8,x]`

output `-1/210*(d*Sqrt[d - c^2*d*x^2]*(30*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(a + b*ArcCosh[c*x]) + 12*c^2*x^2*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(a + b*ArcCosh[c*x]) + b*c*x*(5 - 12*c^2*x^2 + 3*c^4*x^4 - 12*c^6*x^6*Log[x])))/(x^7*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.77.3 Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.58, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6337, 27, 354, 85, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx \\
 & \quad \downarrow \text{6337} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \frac{d(1 - c^2 x^2)^2 (2c^2 x^2 + 5)}{35x^7} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{7dx^7} \\
 & \quad \frac{2c^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{35dx^5} \\
 & \quad \downarrow \text{27} \\
 & \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1 - c^2 x^2)^2 (2c^2 x^2 + 5)}{x^7} dx}{35\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{7dx^7} \\
 & \quad \frac{2c^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{35dx^5} \\
 & \quad \downarrow \text{354} \\
 & \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1 - c^2 x^2)^2 (2c^2 x^2 + 5)}{x^8} dx^2}{70\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{7dx^7} \\
 & \quad \frac{2c^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{35dx^5} \\
 & \quad \downarrow \text{85}
 \end{aligned}$$

---

3.77.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx$

$$\frac{bcd\sqrt{d-c^2dx^2} \int \left( \frac{2c^6}{x^2} + \frac{c^4}{x^4} - \frac{8c^2}{x^6} + \frac{5}{x^8} \right) dx^2}{70\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{7dx^7} - \frac{2c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{35dx^5}$$

↓ 2009

$$-\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{7dx^7} - \frac{2c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{35dx^5} + \frac{bcd\sqrt{d-c^2dx^2} \left( 2c^6 \log(x^2) - \frac{c^4}{x^2} + \frac{4c^2}{x^4} - \frac{5}{3x^6} \right)}{70\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^8,x]`

output `-1/7*((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(d*x^7) - (2*c^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(35*d*x^5) + (b*c*d*Sqrt[d - c^2*d*x^2]*(-5/(3*x^6) + (4*c^2)/x^4 - c^4/x^2 + 2*c^6*Log[x^2]))/(70*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.77.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 85 `Int[((d_)*(x_))^(n_)*((a_) + (b_)*(x_))*((e_) + (f_)*(x_))^(p_), x_] :> Int[ExpandIntegrand[(a + b*x)*(d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, d, e, f, n}, x] && IGtQ[p, 0] && (NeQ[n, -1] || EqQ[p, 1]) && NeQ[b*e + a*f, 0] && (!IntegerQ[n] || LtQ[9*p + 5*n, 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, d, e, f])) && (NeQ[n + p + 3, 0] || EqQ[p, 1])`

rule 354 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

---

3.77.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x^8} dx$

```
rule 6337 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_
), x_Symbol] :> With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCo
sh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c
*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b
, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)]
&& (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])
```

### 3.77.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3144 vs.  $2(207) = 414$ .

Time = 1.17 (sec) , antiderivative size = 3145, normalized size of antiderivative = 12.73

method	result	size
default	Expression too large to display	3145
parts	Expression too large to display	3145

```
input int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^8,x,method=_RETURNVERBOSE)
```

```
output 142/105*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6+154
*c^4*x^4-105*c^2*x^2+25)*x^7/(c*x+1)/(c*x-1)*c^14-2/35*b*(-d*(c^2*x^2-1))^(
1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6+154*c^4*x^4-105*c^2*x^2+25)*x^
11*c^18+1/5*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6
+154*c^4*x^4-105*c^2*x^2+25)*x^9*c^16+26/105*b*(-d*(c^2*x^2-1))^(1/2)*d/(3
5*c^10*x^10-35*c^8*x^8-70*c^6*x^6+154*c^4*x^4-105*c^2*x^2+25)*x^7*c^14-116
/105*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6+154*c^
4*x^4-105*c^2*x^2+25)*x^5*c^12+20/21*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x
^10-35*c^8*x^8-70*c^6*x^6+154*c^4*x^4-105*c^2*x^2+25)*x^3*c^10-5/21*b*(-d*
(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6+154*c^4*x^4-105*c
^2*x^2+25)*x*c^8-164/5*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8*x^8
-70*c^6*x^6+154*c^4*x^4-105*c^2*x^2+25)*x^5/(c*x+1)/(c*x-1)*arccosh(c*x)*c
^12+52/5*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6+15
4*c^4*x^4-105*c^2*x^2+25)*x^3/(c*x+1)/(c*x-1)*arccosh(c*x)*c^10+1966/35*b*
(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6+154*c^4*x^4-1
05*c^2*x^2+25)*x/(c*x+1)/(c*x-1)*arccosh(c*x)*c^8-3272/35*b*(-d*(c^2*x^2-1
))^^(1/2)*d/(35*c^10*x^10-35*c^8*x^8-70*c^6*x^6+154*c^4*x^4-105*c^2*x^2+25)
/x/(c*x+1)/(c*x-1)*arccosh(c*x)*c^6+472/7*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c
^10*x^10-35*c^8*x^8-70*c^6*x^6+154*c^4*x^4-105*c^2*x^2+25)/x^3/(c*x+1)/(c*
x-1)*arccosh(c*x)*c^4-2*b*(-d*(c^2*x^2-1))^(1/2)*d/(35*c^10*x^10-35*c^8...
```

$$3.77. \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^8} dx$$

**3.77.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 648, normalized size of antiderivative = 2.62

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \left[ -\frac{6(2bc^8 dx^8 - bc^6 dx^6 - 9bc^4 dx^4 + 13bc^2 dx^2 - 5bd)\sqrt{-c^2 dx^2 - d}}{x^8} \right]$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^8,x, algorithm="fracas")`

output `[-1/210*(6*(2*b*c^8*d*x^8 - b*c^6*d*x^6 - 9*b*c^4*d*x^4 + 13*b*c^2*d*x^2 - 5*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - 6*(b*c^9*d*x^9 - b*c^7*d*x^7)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 - sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + (3*b*c^5*d*x^5 - (3*b*c^5 - 12*b*c^3 + 5*b*c)*d*x^7 - 12*b*c^3*d*x^3 + 5*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 6*(2*a*c^8*d*x^8 - a*c^6*d*x^6 - 9*a*c^4*d*x^4 + 13*a*c^2*d*x^2 - 5*a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^9 - x^7), 1/210*(12*(b*c^9*d*x^9 - b*c^7*d*x^7)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 6*(2*b*c^8*d*x^8 - b*c^6*d*x^6 - 9*b*c^4*d*x^4 + 13*b*c^2*d*x^2 - 5*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (3*b*c^5*d*x^5 - (3*b*c^5 - 12*b*c^3 + 5*b*c)*d*x^7 - 12*b*c^3*d*x^3 + 5*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 6*(2*a*c^8*d*x^8 - a*c^6*d*x^6 - 9*a*c^4*d*x^4 + 13*a*c^2*d*x^2 - 5*a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^9 - x^7)]`

**3.77.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**8,x)`

output `Timed out`

---

3.77.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx$

**3.77.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.66

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \frac{1}{210} \left( 12 c^6 \sqrt{-d} d \log(x) - \frac{3 c^4 \sqrt{-d} d x^4 - 12 c^2 \sqrt{-d} d x^2 + 5 \sqrt{-d} d}{x^6} \right. \\ \left. - \frac{1}{35} b \left( \frac{2 (-c^2 dx^2 + d)^{5/2} c^2}{dx^5} + \frac{5 (-c^2 dx^2 + d)^{5/2}}{dx^7} \right) \operatorname{arcosh}(cx) \right. \\ \left. - \frac{1}{35} a \left( \frac{2 (-c^2 dx^2 + d)^{5/2} c^2}{dx^5} + \frac{5 (-c^2 dx^2 + d)^{5/2}}{dx^7} \right) \right)$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^8,x, algorithm="maxima")`

output `1/210*(12*c^6*sqrt(-d)*d*log(x) - (3*c^4*sqrt(-d)*d*x^4 - 12*c^2*sqrt(-d)*d*x^2 + 5*sqrt(-d)*d)/x^6)*b*c - 1/35*b*(2*(-c^2*d*x^2 + d)^(5/2)*c^2/(d*x^5) + 5*(-c^2*d*x^2 + d)^(5/2)/(d*x^7))*arccosh(c*x) - 1/35*a*(2*(-c^2*d*x^2 + d)^(5/2)*c^2/(d*x^5) + 5*(-c^2*d*x^2 + d)^(5/2)/(d*x^7))`

**3.77.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^8,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.77.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^8} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^8,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^8, x)`

**3.78**  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^{10}} dx$

3.78.1	Optimal result . . . . .	782
3.78.2	Mathematica [A] (verified) . . . . .	783
3.78.3	Rubi [A] (verified) . . . . .	783
3.78.4	Maple [B] (verified) . . . . .	785
3.78.5	Fricas [A] (verification not implemented) . . . . .	786
3.78.6	Sympy [F(-1)] . . . . .	787
3.78.7	Maxima [A] (verification not implemented) . . . . .	787
3.78.8	Giac [F(-2)] . . . . .	788
3.78.9	Mupad [F(-1)] . . . . .	788

**3.78.1 Optimal result**

Integrand size = 27, antiderivative size = 328

$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^{10}} dx = -\frac{bcd\sqrt{d-c^2dx^2}}{72x^8\sqrt{-1+cx}\sqrt{1+cx}} + \frac{5bc^3d\sqrt{d-c^2dx^2}}{189x^6\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^5d\sqrt{d-c^2dx^2}}{420x^4\sqrt{-1+cx}\sqrt{1+cx}} - \frac{2bc^7d\sqrt{d-c^2dx^2}}{315x^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{9dx^9} - \frac{4c^2(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{63dx^7} - \frac{8c^4(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{315dx^5} + \frac{8bc^9d\sqrt{d-c^2dx^2}\log(x)}{315\sqrt{-1+cx}\sqrt{1+cx}}$$

```
output -1/9*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^9-4/63*c^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^7-8/315*c^4*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^5-1/72*b*c*d*(-c^2*d*x^2+d)^(1/2)/x^8/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/189*b*c^3*d*(-c^2*d*x^2+d)^(1/2)/x^6/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/420*b*c^5*d*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/315*b*c^7*d*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+8/315*b*c^9*d*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.78.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.47

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \frac{d\sqrt{d - c^2 dx^2} (840(-1 + cx)^{5/2}(1 + cx)^{5/2}(a + \operatorname{barccosh}(cx)) + 96c^2 x^2(-1 + cx)^{5/2}(1 + cx)^{5/2}(5 + 2c^2 x^2))}{7560x^9 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^10,x]`

output `-1/7560*(d*Sqrt[d - c^2*d*x^2]*(840*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(a + b*ArcCosh[c*x]) + 96*c^2*x^2*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(5 + 2*c^2*x^2)*(a + b*ArcCosh[c*x]) + b*c*x*(105 - 200*c^2*x^2 + 18*c^4*x^4 + 48*c^6*x^6 - 192*c^8*x^8*Log[x])))/(x^9*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.78.3 Rubi [A] (verified)

Time = 0.53 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.58, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6337, 27, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx \\ & \quad \downarrow \text{6337} \\ & \frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d(1 - c^2 x^2)^2 (8c^4 x^4 + 20c^2 x^2 + 35)}{315x^9} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{9dx^9} \\ & \quad \frac{4c^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{63dx^7} - \frac{8c^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{315dx^5} \\ & \quad \downarrow \text{27} \\ & \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1 - c^2 x^2)^2 (8c^4 x^4 + 20c^2 x^2 + 35)}{x^9} dx}{315\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{9dx^9} \\ & \quad \frac{4c^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{63dx^7} - \frac{8c^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{315dx^5} \end{aligned}$$

---

3.78.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx$



$$\begin{aligned}
& \downarrow 1578 \\
& \frac{bcd\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(8c^4x^4+20c^2x^2+35)}{x^{10}} dx^2}{630\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{9dx^9} - \\
& \frac{4c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{63dx^7} - \frac{8c^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{315dx^5} \\
& \downarrow 1195 \\
& \frac{bcd\sqrt{d-c^2dx^2} \int \left(\frac{8c^8}{x^2} + \frac{4c^6}{x^4} + \frac{3c^4}{x^6} - \frac{50c^2}{x^8} + \frac{35}{x^{10}}\right) dx^2}{630\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{9dx^9} - \\
& \frac{4c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{63dx^7} - \frac{8c^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{315dx^5} \\
& \downarrow 2009 \\
& \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{9dx^9} - \frac{4c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{63dx^7} - \\
& \frac{8c^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{315dx^5} + \frac{bcd\sqrt{d-c^2dx^2} \left(8c^8 \log(x^2) - \frac{4c^6}{x^2} - \frac{3c^4}{2x^4} + \frac{50c^2}{3x^6} - \frac{35}{4x^8}\right)}{630\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^10,x]`

output `-1/9*((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(d*x^9) - (4*c^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(63*d*x^7) - (8*c^4*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(315*d*x^5) + (b*c*d*Sqrt[d - c^2*d*x^2]*(-35/(4*x^8) + (50*c^2)/(3*x^6) - (3*c^4)/(2*x^4) - (4*c^6)/x^2 + 8*c^8*Log[x^2]))/(630*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.78.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 1195 `Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

---

3.78.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x^{10}} dx$

rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.78.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 4261 vs.  $2(276) = 552$ .

Time = 1.29 (sec) , antiderivative size = 4262, normalized size of antiderivative = 12.99

method	result	size
default	Expression too large to display	4262
parts	Expression too large to display	4262

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^10,x,method=_RETURNVERBOSE)`

output

```

-35/9*b*(-d*(c^2*x^2-1))^(1/2)*d/(840*c^12*x^12-945*c^10*x^10+189*c^8*x^8-
2730*c^6*x^6+6210*c^4*x^4-4725*c^2*x^2+1225)*x/(c*x+1)/(c*x-1)*c^10+1225/9
*b*(-d*(c^2*x^2-1))^(1/2)*d/(840*c^12*x^12-945*c^10*x^10+189*c^8*x^8-2730*
c^6*x^6+6210*c^4*x^4-4725*c^2*x^2+1225)/x^9/(c*x+1)/(c*x-1)*arccosh(c*x)+2
5915/126*b*(-d*(c^2*x^2-1))^(1/2)*d/(840*c^12*x^12-945*c^10*x^10+189*c^8*x
^8-2730*c^6*x^6+6210*c^4*x^4-4725*c^2*x^2+1225)/x^2/(c*x+1)^(1/2)/(c*x-1)^
(1/2)*c^7-1285/6*b*(-d*(c^2*x^2-1))^(1/2)*d/(840*c^12*x^12-945*c^10*x^10+1
89*c^8*x^8-2730*c^6*x^6+6210*c^4*x^4-4725*c^2*x^2+1225)/x^4/(c*x+1)^(1/2)/
(c*x-1)^(1/2)*c^5+21175/216*b*(-d*(c^2*x^2-1))^(1/2)*d/(840*c^12*x^12-945*
c^10*x^10+189*c^8*x^8-2730*c^6*x^6+6210*c^4*x^4-4725*c^2*x^2+1225)/x^6/(c*
x+1)^(1/2)/(c*x-1)^(1/2)*c^3-1225/72*b*(-d*(c^2*x^2-1))^(1/2)*d/(840*c^12*
x^12-945*c^10*x^10+189*c^8*x^8-2730*c^6*x^6+6210*c^4*x^4-4725*c^2*x^2+1225
)/x^8/(c*x+1)^(1/2)/(c*x-1)^(1/2)*c-1187/60*b*(-d*(c^2*x^2-1))^(1/2)*d/(84
0*c^12*x^12-945*c^10*x^10+189*c^8*x^8-2730*c^6*x^6+6210*c^4*x^4-4725*c^2*x
^2+1225)*x^4/(c*x+1)^(1/2)/(c*x-1)^(1/2)*c^13+4189/180*b*(-d*(c^2*x^2-1))^(
1/2)*d/(840*c^12*x^12-945*c^10*x^10+189*c^8*x^8-2730*c^6*x^6+6210*c^4*x^4
-4725*c^2*x^2+1225)*x^6/(c*x+1)^(1/2)/(c*x-1)^(1/2)*c^15+280/9*b*(-d*(c^2*
x^2-1))^(1/2)*d/(840*c^12*x^12-945*c^10*x^10+189*c^8*x^8-2730*c^6*x^6+6210
*c^4*x^4-4725*c^2*x^2+1225)/(c*x+1)^(1/2)/(c*x-1)^(1/2)*arccosh(c*x)*c^9-8
29/56*b*(-d*(c^2*x^2-1))^(1/2)*d/(840*c^12*x^12-945*c^10*x^10+189*c^8*x...

```

### 3.78.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 720, normalized size of antiderivative = 2.20

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^{10}} dx = \left[ -\frac{24(8bc^{10}dx^{10} - 4bc^8dx^8 - bc^6dx^6 - 53bc^4dx^4 + 85bc^2dx^2 - \dots}{\dots} \right]$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^10,x, algorithm="fracas")`

---

3.78.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^{10}} dx$

output `[-1/7560*(24*(8*b*c^10*d*x^10 - 4*b*c^8*d*x^8 - b*c^6*d*x^6 - 53*b*c^4*d*x^4 + 85*b*c^2*d*x^2 - 35*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - 96*(b*c^11*d*x^11 - b*c^9*d*x^9)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 - sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + (48*b*c^7*d*x^7 + 18*b*c^5*d*x^5 - (48*b*c^7 + 18*b*c^5 - 200*b*c^3 + 105*b*c)*d*x^9 - 200*b*c^3*d*x^3 + 105*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 24*(8*a*c^10*d*x^10 - 4*a*c^8*d*x^8 - a*c^6*d*x^6 - 53*a*c^4*d*x^4 + 85*a*c^2*d*x^2 - 35*a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^11 - x^9), 1/7560*(192*(b*c^11*d*x^11 - b*c^9*d*x^9)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 24*(8*b*c^10*d*x^10 - 4*b*c^8*d*x^8 - b*c^6*d*x^6 - 53*b*c^4*d*x^4 + 85*b*c^2*d*x^2 - 35*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (48*b*c^7*d*x^7 + 18*b*c^5*d*x^5 - (48*b*c^7 + 18*b*c^5 - 200*b*c^3 + 105*b*c)*d*x^9 - 200*b*c^3*d*x^3 + 105*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 24*(8*a*c^10*d*x^10 - 4*a*c^8*d*x^8 - a*c^6*d*x^6 - 53*a*c^4*d*x^4 + 85*a*c^2*d*x^2 - 35*a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^11 - x^9)]`

### 3.78.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**10,x)`

output `Timed out`

### 3.78.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.69

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \frac{1}{7560} \left( 192 c^8 \sqrt{-dd} \log(x) - \frac{48 c^6 \sqrt{-dd} dx^6 + 18 c^4 \sqrt{-dd} dx^4 - 200 c^2 \sqrt{-dd} dx^2 + d}{x^8} \right) - \frac{1}{315} b \left( \frac{8 (-c^2 dx^2 + d)^{5/2} c^4}{dx^5} + \frac{20 (-c^2 dx^2 + d)^{5/2} c^2}{dx^7} + \frac{35 (-c^2 dx^2 + d)^{5/2}}{dx^9} \right) \operatorname{arcosh}(cx) - \frac{1}{315} a \left( \frac{8 (-c^2 dx^2 + d)^{5/2} c^4}{dx^5} + \frac{20 (-c^2 dx^2 + d)^{5/2} c^2}{dx^7} + \frac{35 (-c^2 dx^2 + d)^{5/2}}{dx^9} \right)$$

---

3.78.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^10,x, algorithm="maxima")`

output `1/7560*(192*c^8*sqrt(-d)*d*log(x) - (48*c^6*sqrt(-d)*d*x^6 + 18*c^4*sqrt(-d)*d*x^4 - 200*c^2*sqrt(-d)*d*x^2 + 105*sqrt(-d)*d)/x^8)*b*c - 1/315*b*(8*(-c^2*d*x^2 + d)^(5/2)*c^4/(d*x^5) + 20*(-c^2*d*x^2 + d)^(5/2)*c^2/(d*x^7) + 35*(-c^2*d*x^2 + d)^(5/2)/(d*x^9))*arccosh(c*x) - 1/315*a*(8*(-c^2*d*x^2 + d)^(5/2)*c^4/(d*x^5) + 20*(-c^2*d*x^2 + d)^(5/2)*c^2/(d*x^7) + 35*(-c^2*d*x^2 + d)^(5/2)/(d*x^9))`

### 3.78.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^10,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command: INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.78.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^{10}} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^10,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^10, x)`

**3.79**  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^{12}} dx$

3.79.1	Optimal result	789
3.79.2	Mathematica [A] (verified)	790
3.79.3	Rubi [A] (verified)	790
3.79.4	Maple [B] (verified)	792
3.79.5	Fricas [A] (verification not implemented)	793
3.79.6	Sympy [F(-1)]	793
3.79.7	Maxima [A] (verification not implemented)	794
3.79.8	Giac [F(-2)]	794
3.79.9	Mupad [F(-1)]	795

**3.79.1 Optimal result**

Integrand size = 27, antiderivative size = 409

$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^{12}} dx = -\frac{bcd\sqrt{d-c^2dx^2}}{110x^{10}\sqrt{-1+cx}\sqrt{1+cx}} + \frac{bc^3d\sqrt{d-c^2dx^2}}{66x^8\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^5d\sqrt{d-c^2dx^2}}{1386x^6\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^7d\sqrt{d-c^2dx^2}}{770x^4\sqrt{-1+cx}\sqrt{1+cx}} - \frac{4bc^9d\sqrt{d-c^2dx^2}}{1155x^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{11dx^{11}} - \frac{2c^2(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{33dx^9} - \frac{8c^4(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{231dx^7} - \frac{16c^6(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{1155dx^5} + \frac{16bc^{11}d\sqrt{d-c^2dx^2}\log(x)}{1155\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/11*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^11-2/33*c^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^9-8/231*c^4*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^7-16/1155*c^6*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/d/x^5-1/110*b*c*d*(-c^2*d*x^2+d)^(1/2)/x^10/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/66*b*c^3*d*(-c^2*d*x^2+d)^(1/2)/x^8/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/1386*b*c^5*d*(-c^2*d*x^2+d)^(1/2)/x^6/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/770*b*c^7*d*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)-4/1155*b*c^9*d*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+16/1155*b*c^11*d*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.79.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.42

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \frac{d\sqrt{d - c^2 dx^2} (630(-1 + cx)^{5/2}(1 + cx)^{5/2}(a + \operatorname{barccosh}(cx)) + 12c^2 x^2(-1 + cx)^{5/2}(1 + cx)^{5/2} (35 + 20c^2 x^2 + 8c^4 x^4)(a + \operatorname{barccosh}(cx)) + bc^2 x^2(63 - 105c^2 x^2 + 5c^4 x^4 + 9c^6 x^6 + 24c^8 x^8 - 96c^{10} x^{10} \operatorname{Log}[x]))}{6930x^{11}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^12,x]`

output `-1/6930*(d*Sqrt[d - c^2*d*x^2]*(630*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(a + b*ArcCosh[c*x]) + 12*c^2*x^2*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(35 + 20*c^2*x^2 + 8*c^4*x^4)*(a + b*ArcCosh[c*x]) + b*c*x*(63 - 105*c^2*x^2 + 5*c^4*x^4 + 9*c^6*x^6 + 24*c^8*x^8 - 96*c^10*x^10*Log[x])))/(x^11*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.79.3 Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.56, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6337, 27, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx$$

↓ 6337

$$\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d(1 - c^2 x^2)^2 (16c^6 x^6 + 40c^4 x^4 + 70c^2 x^2 + 105)}{1155x^{11}} dx - (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{\frac{\sqrt{cx - 1}\sqrt{cx + 1}}{11dx^{11}}}$$

$$\frac{2c^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{33dx^9} - \frac{16c^6(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{1155dx^5} - \frac{8c^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{231dx^7}$$

↓ 27

---

3.79.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx$

$$\begin{aligned}
& \frac{bcd\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(16c^6x^6+40c^4x^4+70c^2x^2+105)}{x^{11}} dx}{1155\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{11dx^{11}} \\
& \frac{2c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{33dx^9} - \frac{16c^6(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{1155dx^5} - \\
& \frac{8c^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{231dx^7} \\
& \quad \downarrow \text{2331} \\
& \frac{bcd\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(16c^6x^6+40c^4x^4+70c^2x^2+105)}{x^{12}} dx^2}{2310\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{11dx^{11}} \\
& \frac{2c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{33dx^9} - \frac{16c^6(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{1155dx^5} - \\
& \frac{8c^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{231dx^7} \\
& \quad \downarrow \text{2123} \\
& \frac{bcd\sqrt{d-c^2dx^2} \int \left( \frac{16c^{10}}{x^2} + \frac{8c^8}{x^4} + \frac{6c^6}{x^6} + \frac{5c^4}{x^8} - \frac{140c^2}{x^{10}} + \frac{105}{x^{12}} \right) dx^2}{2310\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{11dx^{11}} - \frac{2c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{33dx^9} - \\
& \frac{16c^6(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{1155dx^5} - \frac{8c^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{231dx^7} \\
& \quad \downarrow \text{2009} \\
& \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{11dx^{11}} - \frac{2c^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{33dx^9} - \\
& \frac{16c^6(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{1155dx^5} - \frac{8c^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{231dx^7} + \\
& \frac{bcd\sqrt{d-c^2dx^2} \left( 16c^{10} \log(x^2) - \frac{8c^8}{x^2} - \frac{3c^6}{x^4} - \frac{5c^4}{3x^6} + \frac{35c^2}{x^8} - \frac{21}{x^{10}} \right)}{2310\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^12,x]`

output `-1/11*((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(d*x^11) - (2*c^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(33*d*x^9) - (8*c^4*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(231*d*x^7) - (16*c^6*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(1155*d*x^5) + (b*c*d*Sqrt[d - c^2*d*x^2]*(-21/x^10 + (35*c^2)/x^8 - (5*c^4)/(3*x^6) - (3*c^6)/x^4 - (8*c^8)/x^2 + 16*c^10*Log[x^2]))/(2310*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

$$3.79. \int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x^{12}} dx$$



## 3.79.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2123 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`
- rule 2331 `Int[(Pq_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`
- rule 6337 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

## 3.79.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5522 vs.  $2(345) = 690$ .

Time = 1.39 (sec) , antiderivative size = 5523, normalized size of antiderivative = 13.50

method	result	size
default	Expression too large to display	5523
parts	Expression too large to display	5523

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^12,x,method=_RETURNVERBOSE)`

output `result too large to display`

---

3.79.  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^{12}} dx$

**3.79.5 Fracas [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 792, normalized size of antiderivative = 1.94

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \left[ -\frac{6(16bc^{12}dx^{12} - 8bc^{10}dx^{10} - 2bc^8dx^8 - bc^6dx^6 - 145bc^4dx^4 - \dots}{x^{12}} \right]$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^12,x, algorithm="fracas")`

output `[-1/6930*(6*(16*b*c^12*d*x^12 - 8*b*c^10*d*x^10 - 2*b*c^8*d*x^8 - b*c^6*d*x^6 - 145*b*c^4*d*x^4 + 245*b*c^2*d*x^2 - 105*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - 48*(b*c^13*d*x^13 - b*c^11*d*x^11)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 - sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1))*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + (24*b*c^9*d*x^9 + 9*b*c^7*d*x^7 - (24*b*c^9 + 9*b*c^7 + 5*b*c^5 - 105*b*c^3 + 63*b*c)*d*x^11 + 5*b*c^5*d*x^5 - 105*b*c^3*d*x^3 + 63*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 6*(16*a*c^12*d*x^12 - 8*a*c^10*d*x^10 - 2*a*c^8*d*x^8 - a*c^6*d*x^6 - 145*a*c^4*d*x^4 + 245*a*c^2*d*x^2 - 105*a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^13 - x^11), 1/6930*(96*(b*c^13*d*x^13 - b*c^11*d*x^11)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 6*(16*b*c^12*d*x^12 - 8*b*c^10*d*x^10 - 2*b*c^8*d*x^8 - b*c^6*d*x^6 - 145*b*c^4*d*x^4 + 245*b*c^2*d*x^2 - 105*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (24*b*c^9*d*x^9 + 9*b*c^7*d*x^7 - (24*b*c^9 + 9*b*c^7 + 5*b*c^5 - 105*b*c^3 + 63*b*c)*d*x^11 + 5*b*c^5*d*x^5 - 105*b*c^3*d*x^3 + 63*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 6*(16*a*c^12*d*x^12 - 8*a*c^10*d*x^10 - 2*a*c^8*d*x^8 - a*c^6*d*x^6 - 145*a*c^4*d*x^4 + 245*a*c^2*d*x^2 - 105*a*d)*sqrt(-c^2*d*x^2 + d))/(c^2*x^13 - x^11)]`

**3.79.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**12,x)`

output `Timed out`

---

3.79.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx$

**3.79.7 Maxima [A] (verification not implemented)**

Time = 0.45 (sec) , antiderivative size = 287, normalized size of antiderivative = 0.70

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \frac{1}{6930} \left( 96 c^{10} \sqrt{-d} d \log(x) - \frac{24 c^8 \sqrt{-d} dx^8 + 9 c^6 \sqrt{-d} dx^6 + 5 c^4 \sqrt{-d} dx^4}{x^{10}} \right) b \operatorname{arccosh}(cx) - \frac{1}{1155} \left( \frac{16 (-c^2 dx^2 + d)^{5/2} c^6}{dx^5} + \frac{40 (-c^2 dx^2 + d)^{5/2} c^4}{dx^7} + \frac{70 (-c^2 dx^2 + d)^{5/2} c^2}{dx^9} + \frac{105 (-c^2 dx^2 + d)^{5/2}}{dx^{11}} \right) a$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^12,x, algorithm="maxima")`

output `1/6930*(96*c^10*sqrt(-d)*d*log(x) - (24*c^8*sqrt(-d)*d*x^8 + 9*c^6*sqrt(-d)*d*x^6 + 5*c^4*sqrt(-d)*d*x^4 - 105*c^2*sqrt(-d)*d*x^2 + 63*sqrt(-d)*d)/x^10)*b*c - 1/1155*(16*(-c^2*d*x^2 + d)^(5/2)*c^6/(d*x^5) + 40*(-c^2*d*x^2 + d)^(5/2)*c^4/(d*x^7) + 70*(-c^2*d*x^2 + d)^(5/2)*c^2/(d*x^9) + 105*(-c^2*d*x^2 + d)^(5/2)/(d*x^11))*b*arccosh(c*x) - 1/1155*(16*(-c^2*d*x^2 + d)^(5/2)*c^6/(d*x^5) + 40*(-c^2*d*x^2 + d)^(5/2)*c^4/(d*x^7) + 70*(-c^2*d*x^2 + d)^(5/2)*c^2/(d*x^9) + 105*(-c^2*d*x^2 + d)^(5/2)/(d*x^11))*a`

**3.79.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^12,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.79.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^{12}} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^12,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^12, x)`

### 3.80 $\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.80.1	Optimal result	796
3.80.2	Mathematica [A] (verified)	797
3.80.3	Rubi [A] (verified)	797
3.80.4	Maple [B] (verified)	799
3.80.5	Fricas [A] (verification not implemented)	800
3.80.6	Sympy [F(-1)]	801
3.80.7	Maxima [A] (verification not implemented)	801
3.80.8	Giac [F(-2)]	802
3.80.9	Mupad [F(-1)]	802

#### 3.80.1 Optimal result

Integrand size = 27, antiderivative size = 399

$$\begin{aligned} \int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = & \frac{16bdx\sqrt{d - c^2 dx^2}}{1155c^7\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & + \frac{8bdx^3\sqrt{d - c^2 dx^2}}{3465c^5\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{2bdx^5\sqrt{d - c^2 dx^2}}{1925c^3\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & + \frac{bdx^7\sqrt{d - c^2 dx^2}}{1617c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{4bcdx^9\sqrt{d - c^2 dx^2}}{297\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx^{11}\sqrt{d - c^2 dx^2}}{121\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^8d} + \frac{3(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^8d^2} \\ & - \frac{(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{3c^8d^3} + \frac{(d - c^2 dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{11c^8d^4} \end{aligned}$$

output

```
-1/5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/c^8/d+3/7*(-c^2*d*x^2+d)^(7/2)
)*(a+b*arccosh(c*x))/c^8/d^2-1/3*(-c^2*d*x^2+d)^(9/2)*(a+b*arccosh(c*x))/c
^8/d^3+1/11*(-c^2*d*x^2+d)^(11/2)*(a+b*arccosh(c*x))/c^8/d^4+16/1155*b*d*x
*(-c^2*d*x^2+d)^(1/2)/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)+8/3465*b*d*x^3*(-c^2
*d*x^2+d)^(1/2)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/1925*b*d*x^5*(-c^2*d*x^2
+d)^(1/2)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/1617*b*d*x^7*(-c^2*d*x^2+d)^(1
/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-4/297*b*c*d*x^9*(-c^2*d*x^2+d)^(1/2)/(c*
x-1)^(1/2)/(c*x+1)^(1/2)+1/121*b*c^3*d*x^11*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(
1/2)/(c*x+1)^(1/2)
```

### 3.80.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.52

$$\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$d\sqrt{d - c^2 dx^2} \left( 3465a\sqrt{-1 + cx}\sqrt{1 + cx}(-1 + c^2 x^2)^2 (16 + 40c^2 x^2 + 70c^4 x^4 + 105c^6 x^6) - bcx(55440 + 9240c^2 x^2 + 4158c^4 x^4 + 2475c^6 x^6 - 53900c^8 x^8 + 33075c^{10} x^{10}) + 3465b\sqrt{-1 + cx}\sqrt{1 + cx}(-1 + c^2 x^2)^2 (16 + 40c^2 x^2 + 70c^4 x^4 + 105c^6 x^6) \operatorname{ArcCosh}[cx] \right) / (c^8 \sqrt{-1 + cx} \sqrt{1 + cx})$$

input `Integrate[x^7*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `-1/4002075*(d*sqrt[d - c^2*d*x^2]*(3465*a*sqrt[-1 + c*x]*sqrt[1 + c*x]*(-1 + c^2*x^2)^2*(16 + 40*c^2*x^2 + 70*c^4*x^4 + 105*c^6*x^6) - b*c*x*(55440 + 9240*c^2*x^2 + 4158*c^4*x^4 + 2475*c^6*x^6 - 53900*c^8*x^8 + 33075*c^10*x^10) + 3465*b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(-1 + c^2*x^2)^2*(16 + 40*c^2*x^2 + 70*c^4*x^4 + 105*c^6*x^6)*ArcCosh[c*x]))/(c^8*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.80.3 Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.57, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6337}$$

$$-\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d(1 - c^2 x^2)^2 (105c^6 x^6 + 70c^4 x^4 + 40c^2 x^2 + 16)}{1155c^8} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{11c^8 d^4}$$

$$-\frac{(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{3c^8 d^3} + \frac{3(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^8 d^2}$$

$$-\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^8 d}$$

$$\downarrow \text{27}$$

---

3.80.  $\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& \frac{bd\sqrt{d-c^2dx^2} \int (1-c^2x^2)^2 (105c^6x^6 + 70c^4x^4 + 40c^2x^2 + 16) dx}{1155c^7\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{(d-c^2dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{3(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))} - \frac{(d-c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{7c^8d^2} + \\
& \frac{11c^8d^4}{7c^8d^2} - \frac{3c^8d^3}{5c^8d} \\
& \quad \downarrow \text{2341} \\
& \frac{bd\sqrt{d-c^2dx^2} \int (105c^{10}x^{10} - 140c^8x^8 + 5c^6x^6 + 6c^4x^4 + 8c^2x^2 + 16) dx}{1155c^7\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{(d-c^2dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{3(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))} - \frac{(d-c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{7c^8d^2} + \\
& \frac{11c^8d^4}{7c^8d^2} - \frac{3c^8d^3}{5c^8d} \\
& \quad \downarrow \text{2009} \\
& \frac{(d-c^2dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{3(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))} - \frac{(d-c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{7c^8d^2} + \\
& \frac{11c^8d^4}{7c^8d^2} - \frac{3c^8d^3}{5c^8d} + \\
& \frac{bd\left(\frac{105c^{10}x^{11}}{11} - \frac{140c^8x^9}{9} + \frac{5c^6x^7}{7} + \frac{6c^4x^5}{5} + \frac{8c^2x^3}{3} + 16x\right) \sqrt{d-c^2dx^2}}{1155c^7\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^7*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `(b*d*sqrt[d - c^2*d*x^2]*(16*x + (8*c^2*x^3)/3 + (6*c^4*x^5)/5 + (5*c^6*x^7)/7 - (140*c^8*x^9)/9 + (105*c^10*x^11)/11))/(1155*c^7*sqrt[-1 + c*x]*sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(5*c^8*d) + (3*(d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(7*c^8*d^2) - ((d - c^2*d*x^2)^(9/2)*(a + b*ArcCosh[c*x]))/(3*c^8*d^3) + ((d - c^2*d*x^2)^(11/2)*(a + b*ArcCosh[c*x]))/(11*c^8*d^4)`

## 3.80.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2341 `Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

## 3.80.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1845 vs.  $2(335) = 670$ .

Time = 0.89 (sec) , antiderivative size = 1846, normalized size of antiderivative = 4.63

method	result	size
default	Expression too large to display	1846
parts	Expression too large to display	1846

input `int(x^7*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`



output

```

a*(-1/11*x^6*(-c^2*d*x^2+d)^(5/2)/c^2/d+6/11/c^2*(-1/9*x^4*(-c^2*d*x^2+d)^(5/2)/c^2/d+4/9/c^2*(-1/7*x^2*(-c^2*d*x^2+d)^(5/2)/c^2/d-2/35/d/c^4*(-c^2*d*x^2+d)^(5/2))))+b*(-1/247808*(-d*(c^2*x^2-1))^(1/2)*(1+4096*c^8*x^8+220*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+2816*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+620*c^4*x^4-61*c^2*x^2-11*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1232*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+1024*c^12*x^12-2352*c^6*x^6-3328*c^10*x^10+1024*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^11*c^11-2816*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^9*c^9)*(-1+11*arccosh(c*x))*d/(c*x+1)/c^8/(c*x-1)-1/55296*(-d*(c^2*x^2-1))^(1/2)*(256*c^10*x^10-704*c^8*x^8+256*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^9*c^9+688*c^6*x^6-576*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7-280*c^4*x^4+432*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+41*c^2*x^2-120*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+9*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+9*arccosh(c*x))*d/(c*x+1)/c^8/(c*x-1)+1/100352*(-d*(c^2*x^2-1))^(1/2)*(64*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*x^4-112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-25*c^2*x^2+56*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+7*arccosh(c*x))*d/(c*x+1)/c^8/(c*x-1)+11/51200*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c*x))*d/(c*x+1)/c^8/(c*x-1)+1/3072*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*

```

### 3.80.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 275, normalized size of antiderivative = 0.69

$$\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{3465 (105 bc^{12} dx^{12} - 245 bc^{10} dx^{10} + 145 bc^8 dx^8 + bc^6 dx^6 + 2 bc^4 dx^4 + 8 bc^2 dx^2 - 16 bd) \sqrt{-c^2 dx^2 + d} \log ($$

input `integrate(x^7*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output

```

-1/4002075*(3465*(105*b*c^12*d*x^12 - 245*b*c^10*d*x^10 + 145*b*c^8*d*x^8 + b*c^6*d*x^6 + 2*b*c^4*d*x^4 + 8*b*c^2*d*x^2 - 16*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (33075*b*c^11*d*x^11 - 53900*b*c^9*d*x^9 + 2475*b*c^7*d*x^7 + 4158*b*c^5*d*x^5 + 9240*b*c^3*d*x^3 + 55440*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 3465*(105*a*c^12*d*x^12 - 245*a*c^10*d*x^10 + 145*a*c^8*d*x^8 + a*c^6*d*x^6 + 2*a*c^4*d*x^4 + 8*a*c^2*d*x^2 - 16*a*d)*sqrt(-c^2*d*x^2 + d))/(c^10*x^2 - c^8)

```

$$3.80. \quad \int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$$

### 3.80.6 Sympy [F(-1)]

Timed out.

$$\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**7*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)`

output `Timed out`

### 3.80.7 Maxima [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 285, normalized size of antiderivative = 0.71

$$\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{1}{1155} \left( \frac{105 (-c^2 dx^2 + d)^{5/2} x^6}{c^2 d} + \frac{70 (-c^2 dx^2 + d)^{5/2} x^4}{c^4 d} + \frac{40 (-c^2 dx^2 + d)^{5/2} x^2}{c^6 d} + \frac{16 (-c^2 dx^2 + d)^{5/2}}{c^8 d} \right) b \operatorname{arccosh}(cx)$$

$$-\frac{1}{1155} \left( \frac{105 (-c^2 dx^2 + d)^{5/2} x^6}{c^2 d} + \frac{70 (-c^2 dx^2 + d)^{5/2} x^4}{c^4 d} + \frac{40 (-c^2 dx^2 + d)^{5/2} x^2}{c^6 d} + \frac{16 (-c^2 dx^2 + d)^{5/2}}{c^8 d} \right) a$$

$$+ \frac{(33075 c^{10} \sqrt{-d} dx^{11} - 53900 c^8 \sqrt{-d} dx^9 + 2475 c^6 \sqrt{-d} dx^7 + 4158 c^4 \sqrt{-d} dx^5 + 9240 c^2 \sqrt{-d} dx^3 + 55440 \sqrt{-d} dx) b}{4002075 c^7}$$

input `integrate(x^7*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/1155*(105*(-c^2*d*x^2 + d)^(5/2)*x^6/(c^2*d) + 70*(-c^2*d*x^2 + d)^(5/2)*x^4/(c^4*d) + 40*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^6*d) + 16*(-c^2*d*x^2 + d)^(5/2)/(c^8*d))*b*arccosh(c*x) - 1/1155*(105*(-c^2*d*x^2 + d)^(5/2)*x^6/(c^2*d) + 70*(-c^2*d*x^2 + d)^(5/2)*x^4/(c^4*d) + 40*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^6*d) + 16*(-c^2*d*x^2 + d)^(5/2)/(c^8*d))*a + 1/4002075*(33075*c^10*sqrt(-d)*d*x^11 - 53900*c^8*sqrt(-d)*d*x^9 + 2475*c^6*sqrt(-d)*d*x^7 + 4158*c^4*sqrt(-d)*d*x^5 + 9240*c^2*sqrt(-d)*d*x^3 + 55440*sqrt(-d)*d*x)*b/c^7`

**3.80.8 Giac [F(-2)]**

Exception generated.

$$\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^7*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.80.9 Mupad [F(-1)]**

Timed out.

$$\int x^7 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int x^7 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2} dx$$

input `int(x^7*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2),x)`

output `int(x^7*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2), x)`

### 3.81 $\int x^5(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.81.1	Optimal result	803
3.81.2	Mathematica [A] (verified)	804
3.81.3	Rubi [A] (verified)	804
3.81.4	Maple [B] (verified)	806
3.81.5	Fricas [A] (verification not implemented)	807
3.81.6	Sympy [F(-1)]	807
3.81.7	Maxima [A] (verification not implemented)	808
3.81.8	Giac [F(-2)]	808
3.81.9	Mupad [F(-1)]	809

#### 3.81.1 Optimal result

Integrand size = 27, antiderivative size = 321

$$\int x^5(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{8bdx\sqrt{d - c^2dx^2}}{315c^5\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{4bdx^3\sqrt{d - c^2dx^2}}{945c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bdx^5\sqrt{d - c^2dx^2}}{525c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{10bcdx^7\sqrt{d - c^2dx^2}}{441\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx^9\sqrt{d - c^2dx^2}}{81\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^6d} + \frac{2(d - c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6d^2} - \frac{(d - c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^6d^3}$$

output

```
-1/5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/c^6/d+2/7*(-c^2*d*x^2+d)^(7/2)
)*(a+b*arccosh(c*x))/c^6/d-1/9*(-c^2*d*x^2+d)^(9/2)*(a+b*arccosh(c*x))/c
^6/d-3+8/315*b*d*x*(-c^2*d*x^2+d)^(1/2)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+4/
945*b*d*x^3*(-c^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/525*b*d
*x^5*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-10/441*b*c*d*x^7*(
-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/81*b*c^3*d*x^9*(-c^2*d*x
^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.81.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.48

$$\int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d\sqrt{d - c^2 dx^2} (-bcx(2520 + 420c^2 x^2 + 189c^4 x^4 - 2250c^6 x^6 + 1225c^8 x^8) + 11025c^4 x^4 (-1 + cx)^{5/2} (1 + cx))}{99225c^6 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

input `Integrate[x^5*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `-1/99225*(d*sqrt[d - c^2*d*x^2]*(-(b*c*x*(2520 + 420*c^2*x^2 + 189*c^4*x^4 - 2250*c^6*x^6 + 1225*c^8*x^8)) + 11025*c^4*x^4*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(a + b*ArcCosh[c*x]) + 1260*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(2 + 5*c^2*x^2)*(a + b*ArcCosh[c*x]))) / (c^6*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.81.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.58, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6337} \\ & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d(1 - c^2 x^2)^2 (35c^4 x^4 + 20c^2 x^2 + 8)}{315c^6} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^6 d^3} + \\ & \quad \frac{2(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d^2} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^6 d} \\ & \quad \downarrow \text{27} \\ & \frac{bd\sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^2 (35c^4 x^4 + 20c^2 x^2 + 8) dx}{315c^5 \sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^6 d^3} + \\ & \quad \frac{2(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d^2} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^6 d} \\ & \quad \downarrow \text{1467} \end{aligned}$$

---

3.81.  $\int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& \frac{bd\sqrt{d-c^2dx^2} \int (35c^8x^8 - 50c^6x^6 + 3c^4x^4 + 4c^2x^2 + 8) dx}{315c^5\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{9/2}(a+\operatorname{barccosh}(cx))}{9c^6d^3} + \\
& \frac{2(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{7c^6d^2} - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{5c^6d} \\
& \quad \downarrow \text{2009} \\
& - \frac{(d-c^2dx^2)^{9/2}(a+\operatorname{barccosh}(cx))}{9c^6d^3} + \frac{2(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{7c^6d^2} - \\
& \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{5c^6d} + \frac{bd\left(\frac{35c^8x^9}{9} - \frac{50c^6x^7}{7} + \frac{3c^4x^5}{5} + \frac{4c^2x^3}{3} + 8x\right)\sqrt{d-c^2dx^2}}{315c^5\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^5*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `(b*d*Sqrt[d - c^2*d*x^2]*(8*x + (4*c^2*x^3)/3 + (3*c^4*x^5)/5 - (50*c^6*x^7)/7 + (35*c^8*x^9)/9))/(315*c^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(5*c^6*d) + (2*(d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(7*c^6*d^2) - ((d - c^2*d*x^2)^(9/2)*(a + b*ArcCosh[c*x]))/(9*c^6*d^3)`

### 3.81.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 6337 Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_
), x_Symbol] :> With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCo
sh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c
*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b
, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)]
&& (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])
```

### 3.81.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1375 vs.  $2(269) = 538$ .

Time = 0.94 (sec) , antiderivative size = 1376, normalized size of antiderivative = 4.29

method	result	size
default	Expression too large to display	1376
parts	Expression too large to display	1376

```
input int(x^5*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(-1/9*x^4*(-c^2*d*x^2+d)^(5/2)/c^2/d+4/9/c^2*(-1/7*x^2*(-c^2*d*x^2+d)^(5
/2)/c^2/d-2/35/d/c^4*(-c^2*d*x^2+d)^(5/2)))+b*(-1/41472*(-d*(c^2*x^2-1))^(
1/2)*(256*c^10*x^10-704*c^8*x^8+256*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^9*c^9+68
8*c^6*x^6-576*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7-280*c^4*x^4+432*(c*x+1)^(
1/2)*(c*x-1)^(1/2)*c^5*x^5+41*c^2*x^2-120*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3
*x^3+9*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+9*arccosh(c*x))*d/(c*x+1)/c^
6/(c*x-1)-1/25088*(-d*(c^2*x^2-1))^(1/2)*(64*c^8*x^8-144*c^6*x^6+64*(c*x+1
)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*x^4-112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*
c^5*x^5-25*c^2*x^2+56*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*
(c*x+1)^(1/2)*c*x+1)*(-1+7*arccosh(c*x))*d/(c*x+1)/c^6/(c*x-1)+1/3200*(-d*
(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c
^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*
(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c*x))*d/(c*x+1)/c^6/(c*x-1)+1/1152*(-d*
(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x
^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))*d/(c*x+1)/c^6/
(c*x-1)-3/256*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*
x^2-1)*(-1+arccosh(c*x))*d/(c*x+1)/c^6/(c*x-1)-3/256*(-d*(c^2*x^2-1))^(1/2
)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))*d/(c*x+1)/
c^6/(c*x-1)+1/1152*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*
c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*a...
```

$$3.81. \quad \int x^5(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$$

**3.81.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 245, normalized size of antiderivative = 0.76

$$\int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{315 (35 bc^{10} dx^{10} - 85 bc^8 dx^8 + 53 bc^6 dx^6 + bc^4 dx^4 + 4 bc^2 dx^2 - 8 bd) \sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1}) -$$

input `integrate(x^5*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output `-1/99225*(315*(35*b*c^10*d*x^10 - 85*b*c^8*d*x^8 + 53*b*c^6*d*x^6 + b*c^4*d*x^4 + 4*b*c^2*d*x^2 - 8*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (1225*b*c^9*d*x^9 - 2250*b*c^7*d*x^7 + 189*b*c^5*d*x^5 + 420*b*c^3*d*x^3 + 2520*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 315*(35*a*c^10*d*x^10 - 85*a*c^8*d*x^8 + 53*a*c^6*d*x^6 + a*c^4*d*x^4 + 4*a*c^2*d*x^2 - 8*a*d)*sqrt(-c^2*d*x^2 + d))/(c^8*x^2 - c^6)`

**3.81.6 Sympy [F(-1)]**

Timed out.

$$\int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**5*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)`

output `Timed out`



### 3.81.7 Maxima [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.69

$$\int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{1}{315} \left( \frac{35(-c^2 dx^2 + d)^{5/2} x^4}{c^2 d} + \frac{20(-c^2 dx^2 + d)^{5/2} x^2}{c^4 d} + \frac{8(-c^2 dx^2 + d)^{5/2}}{c^6 d} \right) b \operatorname{arcosh}(cx)$$

$$-\frac{1}{315} \left( \frac{35(-c^2 dx^2 + d)^{5/2} x^4}{c^2 d} + \frac{20(-c^2 dx^2 + d)^{5/2} x^2}{c^4 d} + \frac{8(-c^2 dx^2 + d)^{5/2}}{c^6 d} \right) a$$

$$+ \frac{(1225 c^8 \sqrt{-d} dx^9 - 2250 c^6 \sqrt{-d} dx^7 + 189 c^4 \sqrt{-d} dx^5 + 420 c^2 \sqrt{-d} dx^3 + 2520 \sqrt{-d} dx) b}{99225 c^5}$$

input `integrate(x^5*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/315*(35*(-c^2*d*x^2 + d)^(5/2)*x^4/(c^2*d) + 20*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^4*d) + 8*(-c^2*d*x^2 + d)^(5/2)/(c^6*d))*b*arccosh(c*x) - 1/315*(35*(-c^2*d*x^2 + d)^(5/2)*x^4/(c^2*d) + 20*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^4*d) + 8*(-c^2*d*x^2 + d)^(5/2)/(c^6*d))*a + 1/99225*(1225*c^8*sqrt(-d)*d*x^9 - 2250*c^6*sqrt(-d)*d*x^7 + 189*c^4*sqrt(-d)*d*x^5 + 420*c^2*sqrt(-d)*d*x^3 + 2520*sqrt(-d)*d*x)*b/c^5`

### 3.81.8 Giac [F(-2)]

Exception generated.

$$\int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^5*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.81.9 Mupad [F(-1)]**

Timed out.

$$\int x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int x^5 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2} dx$$

input `int(x^5*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2),x)`output `int(x^5*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2), x)`

### 3.82 $\int x^3(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.82.1	Optimal result . . . . .	810
3.82.2	Mathematica [A] (verified) . . . . .	810
3.82.3	Rubi [A] (verified) . . . . .	811
3.82.4	Maple [B] (verified) . . . . .	813
3.82.5	Fricas [A] (verification not implemented) . . . . .	814
3.82.6	Sympy [F(-1)] . . . . .	814
3.82.7	Maxima [A] (verification not implemented) . . . . .	814
3.82.8	Giac [F(-2)] . . . . .	815
3.82.9	Mupad [F(-1)] . . . . .	815

#### 3.82.1 Optimal result

Integrand size = 27, antiderivative size = 243

$$\int x^3(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{2bdx\sqrt{d - c^2dx^2}}{35c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bdx^3\sqrt{d - c^2dx^2}}{105c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{8bcdx^5\sqrt{d - c^2dx^2}}{175\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx^7\sqrt{d - c^2dx^2}}{49\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4d} + \frac{(d - c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4d^2}$$

output

```
-1/5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/c^4/d+1/7*(-c^2*d*x^2+d)^(7/2)
)*(a+b*arccosh(c*x))/c^4/d^2+2/35*b*d*x*(-c^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(
1/2)/(c*x+1)^(1/2)+1/105*b*d*x^3*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x
+1)^(1/2)-8/175*b*c*d*x^5*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
+1/49*b*c^3*d*x^7*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

#### 3.82.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.56

$$\int x^3(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d\sqrt{d - c^2dx^2}(-bcx(210 + 35c^2x^2 - 168c^4x^4 + 75c^6x^6) + 210(-1 + cx)^{5/2}(1 + cx)^{5/2}(a + \operatorname{barccosh}(cx)) + 3675c^4\sqrt{-1 + cx}\sqrt{1 + cx}}{3675c^4\sqrt{-1 + cx}\sqrt{1 + cx}}$$

input `Integrate[x^3*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `-1/3675*(d*Sqrt[d - c^2*d*x^2]*(-(b*c*x*(210 + 35*c^2*x^2 - 168*c^4*x^4 + 75*c^6*x^6)) + 210*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(a + b*ArcCosh[c*x]) + 525*c^2*x^2*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2)*(a + b*ArcCosh[c*x]))) / (c^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.82.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.59, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 290, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6337} \\
 & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d(1 - c^2 x^2)^2 (5c^2 x^2 + 2)}{35c^4} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4 d^2} - \\
 & \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d} \\
 & \quad \downarrow \text{27} \\
 & \frac{bd\sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^2 (5c^2 x^2 + 2) dx}{35c^3 \sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4 d^2} - \\
 & \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d} \\
 & \quad \downarrow \text{290} \\
 & \frac{bd\sqrt{d - c^2 dx^2} \int (5c^6 x^6 - 8c^4 x^4 + c^2 x^2 + 2) dx}{35c^3 \sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4 d^2} - \\
 & \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4 d^2} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^4 d} + \frac{bd \left( \frac{5c^6 x^7}{7} - \frac{8c^4 x^5}{5} + \frac{c^2 x^3}{3} + 2x \right) \sqrt{d - c^2 dx^2}}{35c^3 \sqrt{cx - 1} \sqrt{cx + 1}}$$

input `Int[x^3*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `(b*d*sqrt[d - c^2*d*x^2]*(2*x + (c^2*x^3)/3 - (8*c^4*x^5)/5 + (5*c^6*x^7)/7))/(35*c^3*sqrt[-1 + c*x]*sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(5*c^4*d) + ((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(7*c^4*d^2)`

### 3.82.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 290 `Int[((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.82.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 965 vs.  $2(203) = 406$ .

Time = 0.53 (sec) , antiderivative size = 966, normalized size of antiderivative = 3.98

method	result
default	$a \left( -\frac{x^2(-c^2dx^2+d)^{\frac{5}{2}}}{7c^2d} - \frac{2(-c^2dx^2+d)^{\frac{5}{2}}}{35dc^4} \right) + b \left( -\frac{\sqrt{-d(c^2x^2-1)}(64c^8x^8-144c^6x^6+64\sqrt{cx+1}\sqrt{cx-1}x^7c^7+104c^4x^4-112c^2x^2+56(c^2x^2+d)^{\frac{5}{2}})}{c^4(c^2x^2-1)} \right)$
parts	$a \left( -\frac{x^2(-c^2dx^2+d)^{\frac{5}{2}}}{7c^2d} - \frac{2(-c^2dx^2+d)^{\frac{5}{2}}}{35dc^4} \right) + b \left( -\frac{\sqrt{-d(c^2x^2-1)}(64c^8x^8-144c^6x^6+64\sqrt{cx+1}\sqrt{cx-1}x^7c^7+104c^4x^4-112c^2x^2+56(c^2x^2+d)^{\frac{5}{2}})}{c^4(c^2x^2-1)} \right)$

input `int(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output

```
a*(-1/7*x^2*(-c^2*d*x^2+d)^(5/2)/c^2/d-2/35/d/c^4*(-c^2*d*x^2+d)^(5/2))+b*
(-1/6272*(-d*(c^2*x^2-1))^(1/2)*(64*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(
c*x-1)^(1/2)*x^7*c^7+104*c^4*x^4-112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-2
5*c^2*x^2+56*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x+1)^(
1/2)*c*x+1)*(-1+7*arccosh(c*x))*d/(c*x+1)/c^4/(c*x-1)+1/3200*(-d*(c^2*x^2-
1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13
*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1
/2)*c*x-1)*(-1+5*arccosh(c*x))*d/(c*x+1)/c^4/(c*x-1)+1/384*(-d*(c^2*x^2-1)
)^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-
1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))*d/(c*x+1)/c^4/(c*x-1)-3/
128*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1
+arccosh(c*x))*d/(c*x+1)/c^4/(c*x-1)-3/128*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)
)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))*d/(c*x+1)/c^4/(c*x-1
)+1/384*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c
^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x))*d
/(c*x+1)/c^4/(c*x-1)+1/3200*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)*(c*x
-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-28*c^4
*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(1+5*arccosh(c*x))*d/
(c*x+1)/c^4/(c*x-1)-1/6272*(-d*(c^2*x^2-1))^(1/2)*(-64*(c*x+1)^(1/2)*(c*x-
1)^(1/2)*x^7*c^7+64*c^8*x^8+112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-144...
```

**3.82.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.88

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$\frac{105 (5 bc^8 dx^8 - 13 bc^6 dx^6 + 9 bc^4 dx^4 + bc^2 dx^2 - 2 bd) \sqrt{-c^2 dx^2 + d} \log (cx + \sqrt{c^2 x^2 - 1}) - (75 bc^7 dx^7 - 168 b^2 c^6 dx^6 + 168 b^2 c^5 dx^5 - 35 b^2 c^4 dx^4 + 210 b^2 c^3 dx^3 + 210 b^2 c^2 dx^2 - 210 b^2 dx + 210 b^2) \sqrt{-c^2 dx^2 + d}}{3675 c^3}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fracas")
```

```
output -1/3675*(105*(5*b*c^8*d*x^8 - 13*b*c^6*d*x^6 + 9*b*c^4*d*x^4 + b*c^2*d*x^2 - 2*b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (75*b*c^7*d*x^7 - 168*b*c^5*d*x^5 + 35*b*c^3*d*x^3 + 210*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 105*(5*a*c^8*d*x^8 - 13*a*c^6*d*x^6 + 9*a*c^4*d*x^4 + a*c^2*d*x^2 - 2*a*d)*sqrt(-c^2*d*x^2 + d))/(c^6*x^2 - c^4)
```

**3.82.6 Sympy [F(-1)]**

Timed out.

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

```
input integrate(x**3*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)
```

```
output Timed out
```

**3.82.7 Maxima [A] (verification not implemented)**

Time = 0.38 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.66

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{1}{35} \left( \frac{5(-c^2 dx^2 + d)^{5/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{5/2}}{c^4 d} \right) b \operatorname{arcosh}(cx)$$

$$-\frac{1}{35} \left( \frac{5(-c^2 dx^2 + d)^{5/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{5/2}}{c^4 d} \right) a$$

$$+ \frac{(75 c^6 \sqrt{-d} dx^7 - 168 c^4 \sqrt{-d} dx^5 + 35 c^2 \sqrt{-d} dx^3 + 210 \sqrt{-d} dx) b}{3675 c^3}$$

3.82.  $\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

input `integrate(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/35*(5*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(5/2)/(c^4*d))*b*arccosh(c*x) - 1/35*(5*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(5/2)/(c^4*d))*a + 1/3675*(75*c^6*sqrt(-d)*d*x^7 - 168*c^4*sqrt(-d)*d*x^5 + 35*c^2*sqrt(-d)*d*x^3 + 210*sqrt(-d)*d*x)*b/c^3`

### 3.82.8 Giac [F(-2)]

Exception generated.

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.82.9 Mupad [F(-1)]

Timed out.

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx)) dx = \int x^3 (a + b \text{acosh}(cx)) (d - c^2 dx^2)^{3/2} dx$$

input `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2),x)`

output `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2), x)`



### 3.83 $\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.83.1	Optimal result	816
3.83.2	Mathematica [A] (verified)	816
3.83.3	Rubi [A] (verified)	817
3.83.4	Maple [B] (verified)	818
3.83.5	Fricas [A] (verification not implemented)	819
3.83.6	Sympy [F]	820
3.83.7	Maxima [A] (verification not implemented)	820
3.83.8	Giac [F(-2)]	820
3.83.9	Mupad [F(-1)]	821

#### 3.83.1 Optimal result

Integrand size = 25, antiderivative size = 165

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{bdx\sqrt{d - c^2 dx^2}}{5c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{2bcdx^3\sqrt{d - c^2 dx^2}}{15\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3 dx^5\sqrt{d - c^2 dx^2}}{25\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5c^2 d}$$

output 
$$\frac{-1/5*(-c^2*d*x^2+d)^{(5/2)*(a+b*\operatorname{arccosh}(c*x))}/c^2/d+1/5*b*d*x*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}-2/15*b*c*d*x^3*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}+1/25*b*c^3*d*x^5*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}}}{75c^2(-1+c^2x^2)}$$

#### 3.83.2 Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.65

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d\sqrt{d - c^2 dx^2} \left( 15a(-1 + c^2 x^2)^3 + bcx\sqrt{-1 + cx}\sqrt{1 + cx}(-15 + 10c^2 x^2 - 3c^4 x^4) + 15b(-1 + c^2 x^2)^3 \operatorname{arccosh}(cx) \right)}{75c^2(-1 + c^2 x^2)}$$

input `Integrate[x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output 
$$\frac{-1/75*(d*\text{Sqrt}[d - c^2*d*x^2]*(15*a*(-1 + c^2*x^2)^3 + b*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(-15 + 10*c^2*x^2 - 3*c^4*x^4) + 15*b*(-1 + c^2*x^2)^3*\text{ArcCosh}[c*x]))/(c^2*(-1 + c^2*x^2))$$

### 3.83.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.59, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {6329, 39, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx)) dx \\ & \quad \downarrow \text{6329} \\ & \frac{bd\sqrt{d - c^2 dx^2} \int (1 - cx)^2 (cx + 1)^2 dx}{5c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5c^2 d} \\ & \quad \downarrow \text{39} \\ & \frac{bd\sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^2 dx}{5c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5c^2 d} \\ & \quad \downarrow \text{210} \\ & \frac{bd\sqrt{d - c^2 dx^2} \int (c^4 x^4 - 2c^2 x^2 + 1) dx}{5c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5c^2 d} \\ & \quad \downarrow \text{2009} \\ & \frac{bd\left(\frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x\right) \sqrt{d - c^2 dx^2}}{5c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{5c^2 d} \end{aligned}$$

input 
$$\text{Int}[x*(d - c^2*d*x^2)^(3/2)*(a + b*\text{ArcCosh}[c*x]),x]$$

output 
$$(b*d*\text{Sqrt}[d - c^2*d*x^2]*(x - (2*c^2*x^3)/3 + (c^4*x^5)/5))/(5*c*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) - ((d - c^2*d*x^2)^(5/2)*(a + b*\text{ArcCosh}[c*x]))/(5*c^2*d)$$

## 3.83.3.1 Defintions of rubi rules used

rule 39 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

rule 210 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

## 3.83.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 619 vs.  $2(137) = 274$ .

Time = 1.01 (sec) , antiderivative size = 620, normalized size of antiderivative = 3.76

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{5c^2d} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}(16c^6x^6-28c^4x^4+16\sqrt{cx+1}\sqrt{cx-1}c^5x^5+13c^2x^2-20\sqrt{cx-1}\sqrt{cx+1}c^3x^3+5\sqrt{cx-1}\sqrt{cx+1}c^2x^2-20\sqrt{cx-1}\sqrt{cx+1}c^2x^2-20\sqrt{cx-1}\sqrt{cx+1}c^2x^2+5\sqrt{cx-1}\sqrt{cx+1}c^2x^2)}{800(cx+1)c^2(cx-1)}\right)$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{5c^2d} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}(16c^6x^6-28c^4x^4+16\sqrt{cx+1}\sqrt{cx-1}c^5x^5+13c^2x^2-20\sqrt{cx-1}\sqrt{cx+1}c^3x^3+5\sqrt{cx-1}\sqrt{cx+1}c^2x^2-20\sqrt{cx-1}\sqrt{cx+1}c^2x^2+5\sqrt{cx-1}\sqrt{cx+1}c^2x^2)}{800(cx+1)c^2(cx-1)}\right)$

input `int(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output

```
-1/5*a*(-c^2*d*x^2+d)^(5/2)/c^2/d+b*(-1/800*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c*x))*d/(c*x+1)/c^2/(c*x-1)+1/96*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))*d/(c*x+1)/c^2/(c*x-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+arccosh(c*x))*d/(c*x+1)/c^2/(c*x-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))*d/(c*x+1)/c^2/(c*x-1)+1/96*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x))*d/(c*x+1)/c^2/(c*x-1)-1/800*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-28*c^4*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(1+5*arccosh(c*x))*d/(c*x+1)/c^2/(c*x-1)
```

### 3.83.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.12

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \frac{15(bc^6 dx^6 - 3bc^4 dx^4 + 3bc^2 dx^2 - bd)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1}) - (3bc^5 dx^5 - 10bc^3 dx^3 + 15bc^2 dx^2 - 15bdx)\sqrt{-c^2 dx^2 + d}}{75(c^4 x^2 - c^2)}$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output

```
-1/75*(15*(b*c^6*d*x^6 - 3*b*c^4*d*x^4 + 3*b*c^2*d*x^2 - b*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (3*b*c^5*d*x^5 - 10*b*c^3*d*x^3 + 15*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 15*(a*c^6*d*x^6 - 3*a*c^4*d*x^4 + 3*a*c^2*d*x^2 - a*d)*sqrt(-c^2*d*x^2 + d))/(c^4*x^2 - c^2)
```

**3.83.6 Sympy [F]**

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int x(-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate(x*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)`

output `Integral(x*(-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x)), x)`

**3.83.7 Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.62

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = -\frac{(-c^2 dx^2 + d)^{5/2} b \operatorname{arcosh}(cx)}{5 c^2 d} - \frac{(-c^2 dx^2 + d)^{5/2} a}{5 c^2 d} + \frac{(3 c^4 \sqrt{-d} d^2 x^5 - 10 c^2 \sqrt{-d} d^2 x^3 + 15 \sqrt{-d} d^2 x) b}{75 c d}$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-1/5*(-c^2*d*x^2 + d)^(5/2)*b*arccosh(c*x)/(c^2*d) - 1/5*(-c^2*d*x^2 + d)^(5/2)*a/(c^2*d) + 1/75*(3*c^4*sqrt(-d)*d^2*x^5 - 10*c^2*sqrt(-d)*d^2*x^3 + 15*sqrt(-d)*d^2*x)*b/(c*d)`

**3.83.8 Giac [F(-2)]**

Exception generated.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.83.9 Mupad [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2} dx$$

input `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2),x)`output `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2), x)`

**3.84** 
$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x} dx$$

3.84.1	Optimal result	822
3.84.2	Mathematica [A] (warning: unable to verify)	823
3.84.3	Rubi [A] (verified)	823
3.84.4	Maple [A] (verified)	827
3.84.5	Fricas [F]	828
3.84.6	Sympy [F]	828
3.84.7	Maxima [F]	829
3.84.8	Giac [F(-2)]	829
3.84.9	Mupad [F(-1)]	829

**3.84.1 Optimal result**

Integrand size = 27, antiderivative size = 292

$$\int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{x} dx = -\frac{4bcdx\sqrt{d - c^2dx^2}}{3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx^3\sqrt{d - c^2dx^2}}{9\sqrt{-1 + cx}\sqrt{1 + cx}} + d\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx)) + \frac{1}{3}(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx)) - \frac{2d\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{ibd\sqrt{d - c^2dx^2} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{ibd\sqrt{d - c^2dx^2} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
1/3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))+d*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)-4/3*b*c*d*x*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/9*b*c^3*d*x^3*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*d*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+I*b*d*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-I*b*d*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.84.2 Mathematica [A] (warning: unable to verify)

Time = 0.96 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.15

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx = -\frac{1}{3} ad(-4 + c^2 x^2) \sqrt{d - c^2 dx^2} - \frac{bd\sqrt{d - c^2 dx^2} \left( 9cx + 12 \left( \frac{-1+cx}{1+cx} \right)^{3/2} (1+cx)^3 \operatorname{arccosh}(cx) - \cosh(3 \operatorname{arccosh}(cx)) \right)}{36 \sqrt{\frac{-1+cx}{1+cx}} (1+cx)} + ad^{3/2} \log(x) - ad^{3/2} \log \left( d + \sqrt{d} \sqrt{d - c^2 dx^2} \right) + \frac{bd\sqrt{d - c^2 dx^2} \left( -cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) \right)}{36 \sqrt{\frac{-1+cx}{1+cx}} (1+cx)}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x,x]`

output `-1/3*(a*d*(-4 + c^2*x^2)*Sqrt[d - c^2*d*x^2]) - (b*d*Sqrt[d - c^2*d*x^2]*(9*c*x + 12*((-1 + c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3*ArcCosh[c*x] - Cosh[3*ArcCosh[c*x]]))/(36*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + a*d^(3/2)*Log[x] - a*d^(3/2)*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (b*d*Sqrt[d - c^2*d*x^2]*(-(c*x) + Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + I*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*PolyLog[2, I/E^ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`

### 3.84.3 Rubi [A] (verified)

Time = 1.17 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.79, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.407$ , Rules used = {6345, 25, 39, 2009, 6341, 24, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx$$

↓ 6345

---

3.84.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx$



$$\begin{aligned}
& d \int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x} dx + \frac{bcd\sqrt{d - c^2 dx^2} \int -((1 - cx)(cx + 1)) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{25} \\
& d \int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x} dx - \frac{bcd\sqrt{d - c^2 dx^2} \int (1 - cx)(cx + 1) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{39} \\
& d \int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x} dx - \frac{bcd\sqrt{d - c^2 dx^2} \int (1 - c^2 x^2) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \\
& \quad \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2009} \\
& d \int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \\
& \quad \frac{bcd\left(x - \frac{c^2 x^3}{3}\right) \sqrt{d - c^2 dx^2}}{3\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6341} \\
& d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2 dx^2} \int 1 dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) \right) + \\
& \quad \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd\left(x - \frac{c^2 x^3}{3}\right) \sqrt{d - c^2 dx^2}}{3\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{24} \\
& d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d - c^2 dx^2}}{\sqrt{cx - 1}\sqrt{cx + 1}} \right) + \\
& \quad \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd\left(x - \frac{c^2 x^3}{3}\right) \sqrt{d - c^2 dx^2}}{3\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6362} \\
& d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx - 1}\sqrt{cx + 1}} + \sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d - c^2 dx^2}}{\sqrt{cx - 1}\sqrt{cx + 1}} \right) + \\
& \quad \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd\left(x - \frac{c^2 x^3}{3}\right) \sqrt{d - c^2 dx^2}}{3\sqrt{cx - 1}\sqrt{cx + 1}}
\end{aligned}$$

---

3.84.  $\int \frac{(d - c^2 dx^2)^{3/2}(a + \operatorname{barccosh}(cx))}{x} dx$

↓ 3042

$$d \left( -\frac{\sqrt{d-c^2dx^2} \int (a + \operatorname{barccosh}(cx)) \csc \left( i \operatorname{arccosh}(cx) + \frac{\pi}{2} \right) d \operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcd}{\sqrt{d-c^2dx^2}} \right) \\ \frac{1}{3} (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( x - \frac{c^2x^3}{3} \right) \sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 4668

$$d \left( -\frac{\sqrt{d-c^2dx^2} (-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)}))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \frac{1}{3} (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( x - \frac{c^2x^3}{3} \right) \sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2715

$$d \left( -\frac{\sqrt{d-c^2dx^2} (-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \frac{1}{3} (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( x - \frac{c^2x^3}{3} \right) \sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2838

$$d \left( -\frac{\sqrt{d-c^2dx^2} (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \frac{1}{3} (d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( x - \frac{c^2x^3}{3} \right) \sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x,x]`

output `-1/3*(b*c*d*Sqrt[d - c^2*d*x^2]*(x - (c^2*x^3)/3))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/3 + d*(-((b*c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]])))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]))`

## 3.84.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 39 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6362 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]]] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.84.4 Maple [A] (verified)

Time = 2.03 (sec) , antiderivative size = 499, normalized size of antiderivative = 1.71

method	result
default	$\frac{(-c^2 dx^2 + d)^{\frac{3}{2}} a}{3} - a \ln \left( \frac{2d + 2\sqrt{d}\sqrt{-c^2 dx^2 + d}}{x} \right) d^{\frac{3}{2}} + ad\sqrt{-c^2 dx^2 + d} + \frac{ib\sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(cx) \ln(1+i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{\sqrt{cx-1}\sqrt{cx+1}}$
parts	$\frac{(-c^2 dx^2 + d)^{\frac{3}{2}} a}{3} - a \ln \left( \frac{2d + 2\sqrt{d}\sqrt{-c^2 dx^2 + d}}{x} \right) d^{\frac{3}{2}} + ad\sqrt{-c^2 dx^2 + d} + \frac{ib\sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(cx) \ln(1+i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{\sqrt{cx-1}\sqrt{cx+1}}$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`

---

3.84. 
$$\int \frac{(d-c^2 dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x} dx$$

output  $1/3*(-c^2*d*x^2+d)^{(3/2)}*a-a*\ln((2*d+2*d^{(1/2)}*(-c^2*d*x^2+d)^{(1/2)})/x)*d^{(3/2)}+a*d*(-c^2*d*x^2+d)^{(1/2)}+I*b*(-d*(c^2*x^2-1))^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}*arccosh(c*x)*\ln(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*d-I*b*(-d*(c^2*x^2-1))^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}*arccosh(c*x)*\ln(1-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*d+1/9*b*(-d*(c^2*x^2-1))^{(1/2)*d/(c*x+1)^{(1/2)/(c*x-1)^{(1/2)}*c^3*x^3-4/3*b*(-d*(c^2*x^2-1))^{(1/2)*d/(c*x+1)^{(1/2)/(c*x-1)^{(1/2)}*c*x-4/3*b*(-d*(c^2*x^2-1))^{(1/2)*d/(c*x+1)/(c*x-1)*arccosh(c*x)-I*b*(-d*(c^2*x^2-1))^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)*dilog(1-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*d-1/3*b*(-d*(c^2*x^2-1))^{(1/2)*d/(c*x+1)/(c*x-1)*arccosh(c*x)*x^4*c^4+5/3*b*(-d*(c^2*x^2-1))^{(1/2)*d/(c*x+1)/(c*x-1)*arccosh(c*x)*x^2*c^2+I*b*(-d*(c^2*x^2-1))^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)*dilog(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*d}$

### 3.84.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x,x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x, x)`

### 3.84.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx))}{x} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))/x, x)`

**3.84.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `-1/3*(3*d^(3/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - (-c^2*d*x^2 + d)^(3/2) - 3*sqrt(-c^2*d*x^2 + d)*d)*a + b*integrate((-c^2*d*x^2 + d)^(3/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x, x)`

**3.84.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.84.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x, x)`

**3.85**  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^3} dx$

3.85.1	Optimal result	830
3.85.2	Mathematica [A] (warning: unable to verify)	831
3.85.3	Rubi [A] (verified)	831
3.85.4	Maple [A] (verified)	836
3.85.5	Fricas [F]	836
3.85.6	Sympy [F]	837
3.85.7	Maxima [F]	837
3.85.8	Giac [F(-2)]	837
3.85.9	Mupad [F(-1)]	838

**3.85.1 Optimal result**

Integrand size = 27, antiderivative size = 311

$$\int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{x^3} dx =$$

$$-\frac{bcd\sqrt{d - c^2dx^2}}{2x\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx\sqrt{d - c^2dx^2}}{\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$-\frac{3}{2}c^2d\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx)) - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{2x^2}$$

$$+ \frac{3c^2d\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{3ibc^2d\sqrt{d - c^2dx^2} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{3ibc^2d\sqrt{d - c^2dx^2} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
-1/2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^2-3/2*c^2*d*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)-1/2*b*c*d*(-c^2*d*x^2+d)^(1/2)/x/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*c^3*d*x*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+3*c^2*d*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-3/2*I*b*c^2*d*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+3/2*I*b*c^2*d*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

3.85.  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^3} dx$

### 3.85.2 Mathematica [A] (warning: unable to verify)

Time = 1.47 (sec) , antiderivative size = 500, normalized size of antiderivative = 1.61

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx = \frac{1}{2} \left( -\frac{ad(1 + 2c^2 x^2) \sqrt{d - c^2 dx^2}}{x^2} \right. \\ \left. - 3ac^2 d^{3/2} \log(x) + 3ac^2 d^{3/2} \log\left(d + \sqrt{d} \sqrt{d - c^2 dx^2}\right) - \frac{2bc^2 d \sqrt{d - c^2 dx^2} \left(-cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx\right)}{\sqrt{d - c^2 dx^2}} \right)$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^3,x]`

output `((-(a*d*(1 + 2*c^2*x^2)*Sqrt[d - c^2*d*x^2])/x^2) - 3*a*c^2*d^(3/2)*Log[x] + 3*a*c^2*d^(3/2)*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] - (2*b*c^2*d*Sqrt[d - c^2*d*x^2]*(-c*x) + Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + I*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*PolyLog[2, I/E^ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + (b*d^2*(1 + c*x)*(c*x*Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x] + c*x*ArcCosh[c*x] + I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*PolyLog[2, I/E^ArcCosh[c*x]]))/(x^2*Sqrt[d - c^2*d*x^2]))/2`

### 3.85.3 Rubi [A] (verified)

Time = 1.21 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.77, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {6343, 25, 82, 244, 2009, 6341, 24, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx$$

↓ 6343

---

3.85.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx$



$$\begin{aligned}
& -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx - \frac{bcd\sqrt{d-c^2dx^2} \int -\frac{(1-cx)(cx+1)}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow \text{25} \\
& -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \frac{(1-cx)(cx+1)}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow \text{82} \\
& -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \frac{1-c^2x^2}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow \text{244} \\
& -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx + \frac{bcd\sqrt{d-c^2dx^2} \int (\frac{1}{x^2} - c^2) dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \quad \downarrow \text{2009} \\
& -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \\
& \quad \frac{bcd(c^2(-x) - \frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{6341} \\
& -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \int 1 dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd(c^2(-x) - \frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{24}
\end{aligned}$$

$$\begin{aligned}
& -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd(c^2(-x)-\frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{6362} \\
& -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd(c^2(-x)-\frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{3042} \\
& -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd(c^2(-x)-\frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{4668} \\
& -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2}(-ib \int \log(1-ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1+ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan)}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd(c^2(-x)-\frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{2715} \\
& -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2}(-ib \int e^{-\operatorname{arccosh}(cx)} \log(1-ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1+ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd(c^2(-x)-\frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{2838} \\
& -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2}(2 \arctan(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}))}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \\
& \quad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd(c^2(-x)-\frac{1}{x})\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

---

3.85.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x^3} dx$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^3,x]`

output `(b*c*d*(-x^(-1) - c^2*x)*Sqrt[d - c^2*d*x^2])/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(2*x^2) - (3*c^2*d*(-(b*c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/2`

### 3.85.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

---

3.85.  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^3} dx$

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1]`

rule 6362 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.85.4 Maple [A] (verified)

Time = 0.97 (sec) , antiderivative size = 538, normalized size of antiderivative = 1.73

method	result
default	$a \left( -\frac{(-c^2 d x^2 + d)^{\frac{5}{2}}}{2 d x^2} - \frac{3 c^2 \left( \frac{(-c^2 d x^2 + d)^{\frac{3}{2}}}{3} + d \left( \sqrt{-c^2 d x^2 + d} - \sqrt{d} \ln \left( \frac{2 d + 2 \sqrt{d} \sqrt{-c^2 d x^2 + d}}{x} \right) \right) \right)}{2} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} c^4 d \operatorname{arccosh}(c x)}{(c x + 1)(c x - 1)}$
parts	$a \left( -\frac{(-c^2 d x^2 + d)^{\frac{5}{2}}}{2 d x^2} - \frac{3 c^2 \left( \frac{(-c^2 d x^2 + d)^{\frac{3}{2}}}{3} + d \left( \sqrt{-c^2 d x^2 + d} - \sqrt{d} \ln \left( \frac{2 d + 2 \sqrt{d} \sqrt{-c^2 d x^2 + d}}{x} \right) \right) \right)}{2} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} c^4 d \operatorname{arccosh}(c x)}{(c x + 1)(c x - 1)}$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)`

output

```

a*(-1/2/d/x^2*(-c^2*d*x^2+d)^(5/2)-3/2*c^2*(1/3*(-c^2*d*x^2+d)^(3/2)+d*((-
c^2*d*x^2+d)^(1/2)-d^(1/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x))))-b
*(-d*(c^2*x^2-1))^(1/2)*c^4*d/(c*x+1)/(c*x-1)*arccosh(c*x)*x^2+b*(-d*(c^2*
x^2-1))^(1/2)*c^3*d/(c*x+1)^(1/2)/(c*x-1)^(1/2)*x+1/2*b*(-d*(c^2*x^2-1))^(
1/2)*c^2*d/(c*x+1)/(c*x-1)*arccosh(c*x)-1/2*b*(-d*(c^2*x^2-1))^(1/2)*d/x/(
c*x+1)^(1/2)/(c*x-1)^(1/2)*c+1/2*b*(-d*(c^2*x^2-1))^(1/2)*d/x^2/(c*x+1)/(c
*x-1)*arccosh(c*x)-3/2*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1
/2)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2*d+3/2*I*b*(
-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1-I*(c*x
+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2*d-3/2*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1
)^(1/2)/(c*x+1)^(1/2)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2*d+3
/2*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1-I*(c*x+(
c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2*d

```

### 3.85.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^3,x, algorithm="fricas")`

3.85.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^3} dx$

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^3, x)`

### 3.85.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(-d(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))}{x^3} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**3,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))/x**3, x)`

### 3.85.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `1/2*(3*c^2*d^(3/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - (-c^2*d*x^2 + d)^(3/2)*c^2 - 3*sqrt(-c^2*d*x^2 + d)*c^2*d - (-c^2*d*x^2 + d)^(5/2)/(d*x^2))*a + b*integrate((-c^2*d*x^2 + d)^(3/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^3, x)`

### 3.85.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.85.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^3,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^3, x)`

**3.86**  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$

3.86.1	Optimal result	839
3.86.2	Mathematica [A] (warning: unable to verify)	840
3.86.3	Rubi [A] (verified)	840
3.86.4	Maple [A] (verified)	845
3.86.5	Fricas [F]	846
3.86.6	Sympy [F]	846
3.86.7	Maxima [F]	846
3.86.8	Giac [F(-2)]	847
3.86.9	Mupad [F(-1)]	847

**3.86.1 Optimal result**

Integrand size = 27, antiderivative size = 321

$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^5} dx =$$

$$-\frac{bcd\sqrt{d-c^2dx^2}}{12x^3\sqrt{-1+cx}\sqrt{1+cx}} + \frac{5bc^3d\sqrt{d-c^2dx^2}}{8x\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{3c^2d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{8x^2} - \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{4x^4}$$

$$- \frac{3c^4d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\arctan(e^{\operatorname{arccosh}(cx)})}{4\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{3ibc^4d\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})}{8\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{3ibc^4d\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(cx)})}{8\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/4*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^4+3/8*c^2*d*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x^2-1/12*b*c*d*(-c^2*d*x^2+d)^(1/2)/x^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/8*b*c^3*d*(-c^2*d*x^2+d)^(1/2)/x/(c*x-1)^(1/2)/(c*x+1)^(1/2)-3/4*c^4*d*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+3/8*I*b*c^4*d*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-3/8*I*b*c^4*d*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

3.86.  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$



### 3.86.2 Mathematica [A] (warning: unable to verify)

Time = 1.17 (sec) , antiderivative size = 574, normalized size of antiderivative = 1.79

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \frac{-2bcd^2x + 2bc^2d^2x^2 + 15bc^3d^2x^3 - 15bc^4d^2x^4 - 6ad^2 \sqrt{\frac{-1+cx}{1+cx}} + 2}{x^5}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^5,x]`

output `(-2*b*c*d^2*x + 2*b*c^2*d^2*x^2 + 15*b*c^3*d^2*x^3 - 15*b*c^4*d^2*x^4 - 6*a*d^2*Sqrt[(-1 + c*x)/(1 + c*x)] + 21*a*c^2*d^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)] - 15*a*c^4*d^2*x^4*Sqrt[(-1 + c*x)/(1 + c*x)] - 6*b*d^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + 21*b*c^2*d^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] - 15*b*c^4*d^2*x^4*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + (9*I)*b*c^4*d^2*x^4*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - (9*I)*b*c^5*d^2*x^5*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - (9*I)*b*c^4*d^2*x^4*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + (9*I)*b*c^5*d^2*x^5*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + 9*a*c^4*d^(3/2)*x^4*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2]*Log[x] - 9*a*c^4*d^(3/2)*x^4*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2]*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] - (9*I)*b*c^4*d^2*x^4*(-1 + c*x)*PolyLog[2, (-I)/E^ArcCosh[c*x]] + (9*I)*b*c^4*d^2*x^4*(-1 + c*x)*PolyLog[2, I/E^ArcCosh[c*x]])/(24*x^4*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2])`

### 3.86.3 Rubi [A] (verified)

Time = 1.20 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.81, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$ , Rules used = {6343, 25, 82, 244, 2009, 6339, 15, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx$$

↓ 6343

---

3.86.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx$

$$\begin{aligned}
& -\frac{3}{4}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^3} dx - \frac{bcd\sqrt{d-c^2dx^2} \int -\frac{(1-cx)(cx+1)}{x^4} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \qquad \qquad \qquad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} \\
& \qquad \qquad \qquad \downarrow \text{25} \\
& -\frac{3}{4}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^3} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \frac{(1-cx)(cx+1)}{x^4} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \qquad \qquad \qquad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} \\
& \qquad \qquad \qquad \downarrow \text{82} \\
& -\frac{3}{4}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^3} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \frac{1-c^2x^2}{x^4} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \qquad \qquad \qquad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} \\
& \qquad \qquad \qquad \downarrow \text{244} \\
& -\frac{3}{4}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^3} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \left(\frac{1}{x^4} - \frac{c^2}{x^2}\right) dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \qquad \qquad \qquad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} \\
& \qquad \qquad \qquad \downarrow \text{2009} \\
& -\frac{3}{4}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^3} dx - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \\
& \qquad \qquad \qquad \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \\
& \qquad \qquad \qquad \downarrow \text{6339} \\
& -\frac{3}{4}c^2d \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{bc\sqrt{d-c^2dx^2} \int \frac{1}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2x^2} \right) - \\
& \qquad \qquad \qquad \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \\
& \qquad \qquad \qquad \downarrow \text{15}
\end{aligned}$$

$$-\frac{3}{4}c^2d \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{bc\sqrt{d-c^2dx^2}}{2x\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6362

$$-\frac{3}{4}c^2d \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{bc\sqrt{d-c^2dx^2}}{2x\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 3042

$$-\frac{3}{4}c^2d \left( \frac{c^2\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2x^2} \right) - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 4668

$$-\frac{3}{4}c^2d \left( \frac{c^2\sqrt{d-c^2dx^2} (-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)}))}{2\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2715

$$-\frac{3}{4}c^2d \left( \frac{c^2\sqrt{d-c^2dx^2} (-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{2\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2838

$$-\frac{3}{4}c^2d \left( \frac{c^2\sqrt{d-c^2dx^2}(2\arctan(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) - ib\operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib\operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}))}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. + \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd\left(\frac{c^2}{x} - \frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right)$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^5,x]`

output `(b*c*d*(-1/3*1/x^3 + c^2/x)*Sqrt[d - c^2*d*x^2])/(4*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(4*x^4) - (3*c^2*d*(-1/2*(b*c*Sqrt[d - c^2*d*x^2])/(x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(2*x^2) + (c^2*Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))`/4

### 3.86.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 244 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))]^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6339 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) +
(e_.)*(x_)^2], x_Symbol] :> Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e
*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^
2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`

rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m
+ 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(
m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x],
x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && G
tQ[p, 0] && LtQ[m, -1]`

rule 6362 `Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst [Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.86.4 Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 570, normalized size of antiderivative = 1.78

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{4dx^4} + \frac{ac^2(-c^2dx^2+d)^{\frac{5}{2}}}{8dx^2} + \frac{ac^4(-c^2dx^2+d)^{\frac{3}{2}}}{8} - \frac{3ac^4d^{\frac{3}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{8} + \frac{3ac^4d\sqrt{-c^2dx^2+d}}{8} +$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{5}{2}}}{4dx^4} + \frac{ac^2(-c^2dx^2+d)^{\frac{5}{2}}}{8dx^2} + \frac{ac^4(-c^2dx^2+d)^{\frac{3}{2}}}{8} - \frac{3ac^4d^{\frac{3}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{8} + \frac{3ac^4d\sqrt{-c^2dx^2+d}}{8} +$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^5,x,method=_RETURNVERBOSE)`

output `-1/4*a/d/x^4*(-c^2*d*x^2+d)^(5/2)+1/8*a*c^2/d/x^2*(-c^2*d*x^2+d)^(5/2)+1/8*a*c^4*(-c^2*d*x^2+d)^(3/2)-3/8*a*c^4*d^(3/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)+3/8*a*c^4*d*(-c^2*d*x^2+d)^(1/2)+5/8*b*d*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)/(c*x-1)*arccosh(c*x)*c^4+5/8*b*d*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)^(1/2)/x/(c*x-1)^(1/2)*c^3-7/8*b*d*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)/x^2/(c*x-1)*arccosh(c*x)*c^2-1/12*b*d*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)^(1/2)/x^3/(c*x-1)^(1/2)*c+1/4*b*d*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)/x^4/(c*x-1)*arccosh(c*x)+3/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*d*c^4-3/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*d*c^4+3/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*d*c^4-3/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*d*c^4`

3.86. 
$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$$

**3.86.5 Fricas [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a)}{x^5} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^5,x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^5, x)`

**3.86.6 Sympy [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{(-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx))}{x^5} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))/x**5,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))/x**5, x)`

**3.86.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a)}{x^5} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^5,x, algorithm="maxima")`

output `-1/8*(3*c^4*d^(3/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - (-c^2*d*x^2 + d)^(3/2)*c^4 - 3*sqrt(-c^2*d*x^2 + d)*c^4*d - (-c^2*d*x^2 + d)^(5/2)*c^2/(d*x^2) + 2*(-c^2*d*x^2 + d)^(5/2)/(d*x^4))*a + b*integrate((-c^2*d*x^2 + d)^(3/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^5, x)`

**3.86.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^5,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.86.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2}}{x^5} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^5,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2))/x^5, x)`



### 3.87 $\int x^4(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

3.87.1	Optimal result	848
3.87.2	Mathematica [A] (warning: unable to verify)	849
3.87.3	Rubi [A] (verified)	849
3.87.4	Maple [B] (verified)	855
3.87.5	Fricas [F]	856
3.87.6	Sympy [F(-1)]	857
3.87.7	Maxima [F]	857
3.87.8	Giac [F]	857
3.87.9	Mupad [F(-1)]	858

#### 3.87.1 Optimal result

Integrand size = 27, antiderivative size = 454

$$\begin{aligned} \int x^4(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx &= \frac{3bd^2x^2\sqrt{d - c^2dx^2}}{512c^3\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &+ \frac{bd^2x^4\sqrt{d - c^2dx^2}}{512c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{31bcd^2x^6\sqrt{d - c^2dx^2}}{960\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &+ \frac{21bc^3d^2x^8\sqrt{d - c^2dx^2}}{640\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bc^5d^2x^{10}\sqrt{d - c^2dx^2}}{100\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &- \frac{3d^2x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{256c^4} - \frac{d^2x^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{128c^2} \\ &+ \frac{1}{32}d^2x^5\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) + \frac{1}{16}dx^5(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx)) \\ &+ \frac{1}{10}x^5(d - c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx)) - \frac{3d^2\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{512bc^5\sqrt{-1 + cx}\sqrt{1 + cx}} \end{aligned}$$

output  $1/16*d*x^5*(-c^2*d*x^2+d)^(3/2)*(a+b*\operatorname{arccosh}(c*x))+1/10*x^5*(-c^2*d*x^2+d)^(5/2)*(a+b*\operatorname{arccosh}(c*x))-3/256*d^2*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^(1/2)/c^4-1/128*d^2*x^3*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^(1/2)/c^2+1/32*d^2*x^5*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^(1/2)+3/512*b*d^2*x^2*(-c^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/512*b*d^2*x^4*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-31/960*b*c*d^2*x^6*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+21/640*b*c^3*d^2*x^8*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/100*b*c^5*d^2*x^10*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-3/512*d^2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)$

3.87.  $\int x^4(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

### 3.87.2 Mathematica [A] (warning: unable to verify)

Time = 6.04 (sec) , antiderivative size = 500, normalized size of antiderivative = 1.10

$$\int x^4(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{2880acd^2x\sqrt{\frac{-1+cx}{1+cx}}(1+cx)\sqrt{d-c^2dx^2}(-15-10c^2x^2+248c^4x^4-336c^6x^6+128c^8x^8) + \operatorname{barccosh}(cx)}{}$$

input `Integrate[x^4*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `(2880*a*c*d^2*x*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*sqrt[d - c^2*d*x^2]*(-15 - 10*c^2*x^2 + 248*c^4*x^4 - 336*c^6*x^6 + 128*c^8*x^8) - 43200*a*d^(5/2)*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcTan[(c*x*sqrt[d - c^2*d*x^2])/(sqrt[d]*(-1 + c^2*x^2))] + 1600*b*d^2*sqrt[d - c^2*d*x^2]*(-72*ArcCosh[c*x]^2 + 18*Cosh[2*ArcCosh[c*x]] - 9*Cosh[4*ArcCosh[c*x]] - 2*Cosh[6*ArcCosh[c*x]] + 12*ArcCosh[c*x]*(-3*Sinh[2*ArcCosh[c*x]] + 3*Sinh[4*ArcCosh[c*x]] + Sinh[6*ArcCosh[c*x]])) + 100*b*d^2*sqrt[d - c^2*d*x^2]*(1440*ArcCosh[c*x]^2 - 576*Cosh[2*ArcCosh[c*x]] + 144*Cosh[4*ArcCosh[c*x]] + 64*Cosh[6*ArcCosh[c*x]] + 9*Cosh[8*ArcCosh[c*x]] - 24*ArcCosh[c*x]*(-48*Sinh[2*ArcCosh[c*x]] + 24*Sinh[4*ArcCosh[c*x]] + 16*Sinh[6*ArcCosh[c*x]] + 3*Sinh[8*ArcCosh[c*x]])) + b*d^2*sqrt[d - c^2*d*x^2]*(-50400*ArcCosh[c*x]^2 + 25200*Cosh[2*ArcCosh[c*x]] - 3600*Cosh[4*ArcCosh[c*x]] - 2600*Cosh[6*ArcCosh[c*x]] - 675*Cosh[8*ArcCosh[c*x]] - 72*Cosh[10*ArcCosh[c*x]] + 120*ArcCosh[c*x]*(-420*Sinh[2*ArcCosh[c*x]] + 120*Sinh[4*ArcCosh[c*x]] + 130*Sinh[6*ArcCosh[c*x]] + 45*Sinh[8*ArcCosh[c*x]] + 6*Sinh[10*ArcCosh[c*x]])))/(3686400*c^5*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`

### 3.87.3 Rubi [A] (verified)

Time = 2.02 (sec) , antiderivative size = 431, normalized size of antiderivative = 0.95, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.630$ , Rules used = {6345, 82, 243, 49, 2009, 6345, 25, 82, 244, 2009, 6341, 15, 6354, 15, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$$

---

3.87.  $\int x^4(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& \downarrow \text{6345} \\
& \frac{1}{2}d \int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x^5 (1 - cx)^2 (cx + 1)^2 dx}{10\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{82} \\
& \frac{1}{2}d \int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x^5 (1 - c^2 x^2)^2 dx}{10\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{243} \\
& \frac{1}{2}d \int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x^4 (1 - c^2 x^2)^2 dx^2}{20\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{49} \\
& \frac{1}{2}d \int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (c^4 x^8 - 2c^2 x^6 + x^4) dx^2}{20\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \quad \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{2009} \\
& \frac{1}{2}d \int x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \\
& \quad \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \downarrow \text{6345} \\
& \frac{1}{2}d \left( \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd \sqrt{d - c^2 dx^2} \int -x^5 (1 - cx)(cx + 1) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) \\
& \quad - \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \downarrow \text{25} \\
& \frac{1}{2}d \left( \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int x^5 (1 - cx)(cx + 1) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) \\
& \quad - \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}}
\end{aligned}$$

---

3.87.  $\int x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

↓ 82

$$\frac{1}{2}d \left( \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x^5 (1 - c^2 x^2) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 244

$$\frac{1}{2}d \left( \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int (x^5 - c^2 x^7) dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 2009

$$\frac{1}{2}d \left( \frac{3}{8}d \int x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^6}{6} - \frac{c^2 x^8}{8} \right) \sqrt{d - c^2 dx^2}}{8\sqrt{cx - 1}\sqrt{cx + 1}} \right) - \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6341

$$\frac{1}{2}d \left( \frac{3}{8}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^4 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x^5 dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \right) \right) - \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 15

$$\frac{1}{2}d \left( \frac{3}{8}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^4 (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^6 \sqrt{d - c^2 dx^2}}{36\sqrt{cx - 1}\sqrt{cx + 1}} \right) \right) + \frac{1}{8}x^5 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{1}{10}x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6354

$$\frac{1}{2}d \left( \frac{3}{8}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} - \frac{b \int x^3 dx}{4c} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{1}{10}x^5(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^{10}}{5} - \frac{c^2x^8}{2} + \frac{x^6}{3} \right) \sqrt{d-c^2dx^2}}{20\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 15

$$\frac{1}{2}d \left( \frac{3}{8}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} - \frac{bx^4}{16c} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{1}{10}x^5(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^{10}}{5} - \frac{c^2x^8}{2} + \frac{x^6}{3} \right) \sqrt{d-c^2dx^2}}{20\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6354

$$\frac{1}{2}d \left( \frac{3}{8}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \int x dx}{2c} + \frac{x \sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} \right)}{4c^2} \right) + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{4c^2} \right)}{6\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{6}x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{1}{10}x^5(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^{10}}{5} - \frac{c^2x^8}{2} + \frac{x^6}{3} \right) \sqrt{d-c^2dx^2}}{20\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 15

$$\frac{1}{2}d \left( \frac{3}{8}d - \frac{\sqrt{d - c^2 dx^2} \left( 3 \left( \frac{\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{2c^2} + \frac{x \sqrt{cx - 1} \sqrt{cx + 1} (a + b \operatorname{arccosh}(cx)) - \frac{bx^2}{4c}}{2c^2} \right)}{4c^2} + \frac{x^3 \sqrt{cx - 1} \sqrt{cx + 1} (a + b \operatorname{arccosh}(cx))}{4c^2} \right)}{6\sqrt{cx - 1} \sqrt{cx + 1}} \right)$$

$$\frac{1}{10}x^5(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^{10}}{5} - \frac{c^2 x^8}{2} + \frac{x^6}{3} \right) \sqrt{d - c^2 dx^2}}{20\sqrt{cx - 1} \sqrt{cx + 1}}$$

↓ 6308

$$\frac{1}{10}x^5(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx)) + \frac{1}{2}d \left( \frac{1}{8}x^5(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx)) + \frac{3}{8}d \left( \frac{1}{6}x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx)) - \frac{\sqrt{d - c^2 dx^2} \left( \frac{x^3 \sqrt{cx - 1} \sqrt{cx + 1} (a + b \operatorname{arccosh}(cx))}{4c^2} \right)}{6\sqrt{cx - 1} \sqrt{cx + 1}} \right) \right)$$

input `Int[x^4*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `-1/20*(b*c*d^2*sqrt[d - c^2*d*x^2]*(x^6/3 - (c^2*x^8)/2 + (c^4*x^10)/5))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (x^5*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/10 + (d*(-1/8*(b*c*d*sqrt[d - c^2*d*x^2]*(x^6/6 - (c^2*x^8)/8)))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (x^5*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/8 + (3*d*(-1/36*(b*c*x^6*sqrt[d - c^2*d*x^2]))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (x^5*sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/6 - (sqrt[d - c^2*d*x^2]*(-1/16*(b*x^4)/c + (x^3*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(4*c^2) + (3*(-1/4*(b*x^2)/c + (x*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(2*c^2) + (a + b*ArcCosh[c*x])^2/(4*b*c^3)))/(4*c^2)))/(6*sqrt[-1 + c*x]*sqrt[1 + c*x]))/8)/2`

## 3.87.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`
- rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)])*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.87.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1742 vs.  $2(386) = 772$ .

Time = 0.90 (sec) , antiderivative size = 1743, normalized size of antiderivative = 3.84

method	result	size
default	Expression too large to display	1743
parts	Expression too large to display	1743

---

3.87.  $\int x^4(d - c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx)) dx$



input `int(x^4*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-1/10*a*x^3*(-c^2*d*x^2+d)^(7/2)/c^2/d-3/80*a/c^4*x*(-c^2*d*x^2+d)^(7/2)/d  
+1/160*a/c^4*x*(-c^2*d*x^2+d)^(5/2)+1/128*a/c^4*d*x*(-c^2*d*x^2+d)^(3/2)+3  
/256*a/c^4*d^2*x*(-c^2*d*x^2+d)^(1/2)+3/256*a/c^4*d^3/(c^2*d)^(1/2)*arctan  
((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b*(-3/512*(-d*(c^2*x^2-1))^(1/2)/(c  
*x-1)^(1/2)/(c*x+1)^(1/2)/c^5*arccosh(c*x)^2*d^2+1/102400*(-d*(c^2*x^2-1))  
^(1/2)*(-1280*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^8*x^8+512*(c*x+1)^(1/2)*(c*x-1  
)^(1/2)*c^10*x^10+50*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c  
x+1)^(1/2)-832*c^5*x^5+170*c^3*x^3+1696*c^7*x^7+1120*(c*x+1)^(1/2)*(c*x-1  
^(1/2)*c^6*x^6-400*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-1536*c^9*x^9-10*c*x  
+512*c^11*x^11)*(-1+10*arccosh(c*x))*d^2/(c*x+1)/c^5/(c*x-1)-1/32768*(-d*(  
c^2*x^2-1))^(1/2)*(128*c^9*x^9-320*c^7*x^7+128*(c*x-1)^(1/2)*(c*x+1)^(1/2)  
*c^8*x^8+272*c^5*x^5-256*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6-88*c^3*x^3+16  
0*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c*x-32*(c*x-1)^(1/2)*(c*x+1)^(1/2)  
*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+8*arccosh(c*x))*d^2/(c*x+1)/c^5/  
(c*x-1)-1/12288*(-d*(c^2*x^2-1))^(1/2)*(32*c^7*x^7-64*c^5*x^5+32*(c*x+1)^(  
1/2)*(c*x-1)^(1/2)*c^6*x^6+38*c^3*x^3-48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x  
^4-6*c*x+18*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2  
2)*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c  
x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+4*arc...`

### 3.87.5 Fricas [F]

$$\int x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fricas  
")`

output `integral((a*c^4*d^2*x^8 - 2*a*c^2*d^2*x^6 + a*d^2*x^4 + (b*c^4*d^2*x^8 - 2  
*b*c^2*d^2*x^6 + b*d^2*x^4)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

**3.87.6 Sympy [F(-1)]**

Timed out.

$$\int x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**4*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)`output `Timed out`**3.87.7 Maxima [F]**

$$\int x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`output `-1/1280*(128*(-c^2*d*x^2 + d)^(7/2)*x^3/(c^2*d) - 8*(-c^2*d*x^2 + d)^(5/2)*x/c^4 + 48*(-c^2*d*x^2 + d)^(7/2)*x/(c^4*d) - 10*(-c^2*d*x^2 + d)^(3/2)*d*x/c^4 - 15*sqrt(-c^2*d*x^2 + d)*d^2*x/c^4 - 15*d^(5/2)*arcsin(c*x)/c^5)*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)), x)`**3.87.8 Giac [F]**

$$\int x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a) x^4 dx$$

input `integrate(x^4*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)*x^4, x)`

**3.87.9 Mupad [F(-1)]**

Timed out.

$$\int x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} dx$$

input `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2),x)`output `int(x^4*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2), x)`

### 3.88 $\int x^2(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

3.88.1	Optimal result	859
3.88.2	Mathematica [A] (warning: unable to verify)	860
3.88.3	Rubi [A] (verified)	860
3.88.4	Maple [B] (verified)	865
3.88.5	Fricas [F]	866
3.88.6	Sympy [F(-1)]	867
3.88.7	Maxima [F]	867
3.88.8	Giac [F]	867
3.88.9	Mupad [F(-1)]	868

#### 3.88.1 Optimal result

Integrand size = 27, antiderivative size = 371

$$\begin{aligned} \int x^2(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx &= \frac{5bd^2x^2\sqrt{d - c^2dx^2}}{256c\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &- \frac{59bcd^2x^4\sqrt{d - c^2dx^2}}{768\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{17bc^3d^2x^6\sqrt{d - c^2dx^2}}{288\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &- \frac{bc^5d^2x^8\sqrt{d - c^2dx^2}}{64\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{5d^2x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{128c^2} \\ &+ \frac{5}{64}d^2x^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) + \frac{5}{48}dx^3(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx)) \\ &+ \frac{1}{8}x^3(d - c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx)) - \frac{5d^2\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{256bc^3\sqrt{-1 + cx}\sqrt{1 + cx}} \end{aligned}$$

output  $5/48*d*x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*\operatorname{arccosh}(c*x))+1/8*x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*\operatorname{arccosh}(c*x))-5/128*d^2*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^(1/2)/c^2+5/64*d^2*x^3*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^(1/2)+5/256*b*d^2*x^2*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-59/768*b*c*d^2*x^4*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+17/288*b*c^3*d^2*x^6*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/64*b*c^5*d^2*x^8*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-5/256*d^2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)$

**3.88.2 Mathematica [A] (warning: unable to verify)**

Time = 3.70 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.12

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{192acd^2 x \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \sqrt{d - c^2 dx^2} (-15 + 118c^2 x^2 - 136c^4 x^4 + 48c^6 x^6) - 2880ad^{5/2}}{\dots}$$

input `Integrate[x^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output

```
(192*a*c*d^2*x*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Sqrt[d - c^2*d*x^2]*(-15 + 118*c^2*x^2 - 136*c^4*x^4 + 48*c^6*x^6) - 2880*a*d^(5/2)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] - 576*b*d^2*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]) - 64*b*d^2*Sqrt[d - c^2*d*x^2]*(-72*ArcCosh[c*x]^2 + 18*Cosh[2*ArcCosh[c*x]] - 9*Cosh[4*ArcCosh[c*x]] - 2*Cosh[6*ArcCosh[c*x]] + 12*ArcCosh[c*x]*(-3*Sinh[2*ArcCosh[c*x]] + 3*Sinh[4*ArcCosh[c*x]] + Sinh[6*ArcCosh[c*x]])) + b*d^2*Sqrt[d - c^2*d*x^2]*(-1440*ArcCosh[c*x]^2 + 576*Cosh[2*ArcCosh[c*x]] - 144*Cosh[4*ArcCosh[c*x]] - 64*Cosh[6*ArcCosh[c*x]] - 9*Cosh[8*ArcCosh[c*x]] + 24*ArcCosh[c*x]*(-4*8*Sinh[2*ArcCosh[c*x]] + 24*Sinh[4*ArcCosh[c*x]] + 16*Sinh[6*ArcCosh[c*x]] + 3*Sinh[8*ArcCosh[c*x]])))/(73728*c^3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))
```

**3.88.3 Rubi [A] (verified)**Time = 1.57 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.01, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$ , Rules used = {6345, 82, 243, 49, 2009, 6345, 25, 82, 244, 2009, 6341, 15, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$$

↓ 6345

---

3.88.  $\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

$$\frac{5}{8}d \int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x^3(1 - cx)^2(cx + 1)^2 dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))$$

↓ 82

$$\frac{5}{8}d \int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x^3(1 - c^2 x^2)^2 dx}{8\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))$$

↓ 243

$$\frac{5}{8}d \int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x^2(1 - c^2 x^2)^2 dx^2}{16\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))$$

↓ 49

$$\frac{5}{8}d \int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (c^4 x^6 - 2c^2 x^4 + x^2) dx^2}{16\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))$$

↓ 2009

$$\frac{5}{8}d \int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6345

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd \sqrt{d - c^2 dx^2} \int -x^3(1 - cx)(cx + 1) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 25

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd \sqrt{d - c^2 dx^2} \int x^3(1 - cx)(cx + 1) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 82

---

3.88.  $\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x^3(1 - c^2 x^2) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 244

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int (x^3 - c^2 x^5) dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 2009

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^4}{4} - \frac{c^2 x^6}{6} \right) \sqrt{d - c^2 dx^2}}{6\sqrt{cx - 1}\sqrt{cx + 1}} \right) - \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6341

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x^3 dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \right) \right) - \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 15

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^4 \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}} \right) \right) + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^8}{4} - \frac{2c^2 x^6}{3} + \frac{x^4}{2} \right) \sqrt{d - c^2 dx^2}}{16\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6354

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \int x dx}{2c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \right. \\ \left. - \frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^8}{4} - \frac{2c^2x^6}{3} + \frac{x^4}{2} \right) \sqrt{d-c^2dx^2}}{16\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 15

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))}{2c^2} - \frac{bx^2}{4c} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \right. \\ \left. - \frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^8}{4} - \frac{2c^2x^6}{3} + \frac{x^4}{2} \right) \sqrt{d-c^2dx^2}}{16\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6308

$$\frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx)) + \\ \frac{5}{8}d \left( \frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) + \frac{1}{2}d \left( \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{\sqrt{d-c^2dx^2} \left( \frac{a+b\operatorname{arccosh}(cx)}{4bc^3} \right)}{16\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$

input `Int[x^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `-1/16*(b*c*d^2*Sqrt[d - c^2*d*x^2]*(x^4/2 - (2*c^2*x^6)/3 + (c^4*x^8)/4))/`  
`(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^3*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh`  
`[c*x]))/8 + (5*d*(-1/6*(b*c*d*Sqrt[d - c^2*d*x^2]*(x^4/4 - (c^2*x^6)/6)))/(`  
`Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x^3*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[`  
`c*x]))/6 + (d*(-1/16*(b*c*x^4*Sqrt[d - c^2*d*x^2]))/(Sqrt[-1 + c*x]*Sqrt[1`  
`+ c*x]) + (x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/4 - (Sqrt[d - c^2`  
`*d*x^2]*(-1/4*(b*x^2)/c + (x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c`  
`*x]))/(2*c^2) + (a + b*ArcCosh[c*x])^2/(4*b*c^3))/(4*Sqrt[-1 + c*x]*Sqrt[`  
`1 + c*x]))/2)/8`



## 3.88.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`
- rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)])*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.88.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1288 vs.  $2(315) = 630$ .

Time = 0.74 (sec) , antiderivative size = 1289, normalized size of antiderivative = 3.47

method	result	size
default	Expression too large to display	1289
parts	Expression too large to display	1289

---

3.88.  $\int x^2(d - c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx)) dx$

input `int(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-1/8*a*x*(-c^2*d*x^2+d)^(7/2)/c^2/d+1/48*a/c^2*x*(-c^2*d*x^2+d)^(5/2)+5/19  
2*a/c^2*d*x*(-c^2*d*x^2+d)^(3/2)+5/128*a/c^2*d^2*x*(-c^2*d*x^2+d)^(1/2)+5/  
128*a/c^2*d^3/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b  
*(-5/256*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/c^3*arccosh(c*  
x)^2*d^2+1/16384*(-d*(c^2*x^2-1))^(1/2)*(128*c^9*x^9-320*c^7*x^7+128*(c*x-  
1)^(1/2)*(c*x+1)^(1/2)*c^8*x^8+272*c^5*x^5-256*(c*x+1)^(1/2)*(c*x-1)^(1/2)  
*c^6*x^6-88*c^3*x^3+160*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c*x-32*(c*x-  
1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+8*arccosh(  
c*x))*d^2/(c*x+1)/c^3/(c*x-1)-1/2304*(-d*(c^2*x^2-1))^(1/2)*(32*c^7*x^7-64  
*c^5*x^5+32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6+38*c^3*x^3-48*(c*x+1)^(1/2  
)*(c*x-1)^(1/2)*c^4*x^4-6*c*x+18*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-  
1)^(1/2)*(c*x+1)^(1/2))*(-1+6*arccosh(c*x))*d^2/(c*x+1)/c^3/(c*x-1)+1/1024  
*(-d*(c^2*x^2-1))^(1/2)*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2  
)*c^4*x^4+4*c*x-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1  
)^^(1/2))*(-1+4*arccosh(c*x))*d^2/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c^2*x^2-1)  
)^^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/  
2)*(c*x+1)^(1/2))*(-1+2*arccosh(c*x))*d^2/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c  
^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)  
^(1/2)*(c*x+1)^(1/2)-2*c*x)*(1+2*arccosh(c*x))*d^2/(c*x+1)/c^3/(c*x-1)+1/1  
024*(-d*(c^2*x^2-1))^(1/2)*(-8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c^...`

### 3.88.5 Fricas [F]

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((a*c^4*d^2*x^6 - 2*a*c^2*d^2*x^4 + a*d^2*x^2 + (b*c^4*d^2*x^6 - 2  
*b*c^2*d^2*x^4 + b*d^2*x^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

**3.88.6 Sympy [F(-1)]**

Timed out.

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**2*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)`

output `Timed out`

**3.88.7 Maxima [F]**

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/384*(8*(-c^2*d*x^2 + d)^(5/2)*x/c^2 - 48*(-c^2*d*x^2 + d)^(7/2)*x/(c^2*d) + 10*(-c^2*d*x^2 + d)^(3/2)*d*x/c^2 + 15*sqrt(-c^2*d*x^2 + d)*d^2*x/c^2 + 15*d^(5/2)*arcsin(c*x)/c^3)*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*x^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)), x)`

**3.88.8 Giac [F]**

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)*x^2, x)`

**3.88.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} dx$$

input `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2),x)`output `int(x^2*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2), x)`

### 3.89 $\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

3.89.1	Optimal result	869
3.89.2	Mathematica [A] (warning: unable to verify)	870
3.89.3	Rubi [A] (verified)	870
3.89.4	Maple [B] (verified)	874
3.89.5	Fricas [F]	875
3.89.6	Sympy [F(-1)]	876
3.89.7	Maxima [F]	876
3.89.8	Giac [F(-2)]	876
3.89.9	Mupad [F(-1)]	877

#### 3.89.1 Optimal result

Integrand size = 24, antiderivative size = 293

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = -\frac{25bcd^2x^2\sqrt{d - c^2dx^2}}{96\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{5bc^3d^2x^4\sqrt{d - c^2dx^2}}{96\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bd^2(1 - c^2x^2)^3\sqrt{d - c^2dx^2}}{36c\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{5}{16}d^2x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx)) + \frac{5}{24}dx(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx)) + \frac{1}{6}x(d - c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx)) - \frac{5d^2\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{32bc\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output  $5/24*d*x*(-c^2*d*x^2+d)^(3/2)*(a+b*\operatorname{arccosh}(c*x))+1/6*x*(-c^2*d*x^2+d)^(5/2)*(a+b*\operatorname{arccosh}(c*x))+5/16*d^2*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^(1/2)-25/96*b*c*d^2*x^2*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/96*b*c^3*d^2*x^4*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/36*b*d^2*(-c^2*x^2+1)^3*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-5/32*d^2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)$

### 3.89.2 Mathematica [A] (warning: unable to verify)

Time = 1.93 (sec) , antiderivative size = 347, normalized size of antiderivative = 1.18

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{48acd^2 x \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \sqrt{d - c^2 dx^2} (33 - 26c^2 x^2 + 8c^4 x^4) - 720ad^{5/2} \sqrt{\frac{-1+cx}{1+cx}} (1+cx)}{\dots}$$

input `Integrate[(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `(48*a*c*d^2*x*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Sqrt[d - c^2*d*x^2]*(33 - 26*c^2*x^2 + 8*c^4*x^4) - 720*a*d^(5/2)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] - 288*b*d^2*Sqrt[d - c^2*d*x^2]*(Cosh[2*ArcCosh[c*x]] + 2*ArcCosh[c*x]*(ArcCosh[c*x] - Sinh[2*ArcCosh[c*x]])) + 36*b*d^2*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]) + b*d^2*Sqrt[d - c^2*d*x^2]*(-72*ArcCosh[c*x]^2 + 18*Cosh[2*ArcCosh[c*x]] - 9*Cosh[4*ArcCosh[c*x]] - 2*Cosh[6*ArcCosh[c*x]] + 12*ArcCosh[c*x]*(-3*Sinh[2*ArcCosh[c*x]] + 3*Sinh[4*ArcCosh[c*x]] + Sinh[6*ArcCosh[c*x]])))/(2304*c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`

### 3.89.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.04, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.458$ , Rules used = {6312, 82, 241, 6312, 25, 82, 244, 2009, 6310, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6312} \\ & \frac{5}{6}d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x(1 - cx)^2 (cx + 1)^2 dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \\ & \quad \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{82} \end{aligned}$$

---

3.89.  $\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

$$\frac{5}{6}d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int x(1 - c^2 x^2)^2 dx}{6\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))$$

↓ 241

$$\frac{5}{6}d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 6312

$$\frac{5}{6}d \left( \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd\sqrt{d - c^2 dx^2} \int -x(1 - cx)(cx + 1) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 25

$$\frac{5}{6}d \left( \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x(1 - cx)(cx + 1) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 82

$$\frac{5}{6}d \left( \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x(1 - c^2 x^2) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 244

$$\frac{5}{6}d \left( \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2 dx^2} \int (x - c^2 x^3) dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 2009



$$\frac{5}{6}d \left( \frac{3}{4}d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^2}{2} - \frac{c^2 x^4}{4} \right) \sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right. \\ \left. + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 6310

$$\frac{5}{6}d \left( \frac{3}{4}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \right) + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right. \\ \left. + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 15

$$\frac{5}{6}d \left( \frac{3}{4}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{bcx^2 \sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right) + \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right. \\ \left. + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 6308

$$\frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) + \\ \frac{5}{6}d \left( \frac{1}{4}x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) + \frac{3}{4}d \left( \frac{1}{2}x\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{4bc\sqrt{cx - 1}\sqrt{cx + 1}} \right. \right. \\ \left. \left. + \frac{bd^2(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{36c\sqrt{cx - 1}\sqrt{cx + 1}} \right) \right)$$

input `Int[(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `(b*d^2*(1 - c^2*x^2)^3*Sqrt[d - c^2*d*x^2])/(36*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/6 + (5*d*(-1/4*(b*c*d*Sqrt[d - c^2*d*x^2]*(x^2/2 - (c^2*x^4)/4)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/4 + (3*d*(-1/4*(b*c*x^2*Sqrt[d - c^2*d*x^2]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(4*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/4)/6`

## 3.89.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`
- rule 244 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^(n/2)), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

```
rule 6312 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),
x_Symbol] :> Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] +
(Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x
], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p
)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n
- 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n,
0] && GtQ[p, 0]
```

### 3.89.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 884 vs.  $2(249) = 498$ .

Time = 0.99 (sec) , antiderivative size = 885, normalized size of antiderivative = 3.02

method	result
default	$\frac{ax(-c^2dx^2+d)^{\frac{5}{2}}}{6} + \frac{5adx(-c^2dx^2+d)^{\frac{3}{2}}}{24} + \frac{5ad^2x\sqrt{-c^2dx^2+d}}{16} + \frac{5ad^3 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{16\sqrt{c^2d}} + b\left(-\frac{5\sqrt{-d(c^2x^2-1)} \arccos\left(\frac{cx-1}{\sqrt{cx-1}\sqrt{cx+1}}\right)}{32\sqrt{cx-1}\sqrt{cx+1}}\right)$
parts	$\frac{ax(-c^2dx^2+d)^{\frac{5}{2}}}{6} + \frac{5adx(-c^2dx^2+d)^{\frac{3}{2}}}{24} + \frac{5ad^2x\sqrt{-c^2dx^2+d}}{16} + \frac{5ad^3 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{16\sqrt{c^2d}} + b\left(-\frac{5\sqrt{-d(c^2x^2-1)} \arccos\left(\frac{cx-1}{\sqrt{cx-1}\sqrt{cx+1}}\right)}{32\sqrt{cx-1}\sqrt{cx+1}}\right)$

```
input int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

output  $\frac{1}{6}ax(-c^2dx^2+d)^{5/2} + \frac{5}{24}ad^2x(-c^2dx^2+d)^{3/2} + \frac{5}{16}ad^2x(-c^2dx^2+d)^{1/2} + \frac{5}{16}ad^3/(c^2d)^{1/2} \arctan((c^2d)^{1/2}x/(-c^2dx^2+d)^{1/2}) + b(-5/32(-d(c^2x^2-1))^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}/c \operatorname{arccosh}(cx))^2 d^2 + 1/2304(-d(c^2x^2-1))^{1/2} * (32c^7x^7 - 64c^5x^5 + 32(c^2x^2+d)^{1/2} * (cx-1)^{1/2} * c^6x^6 + 38c^3x^3 - 48(c^2x^2+d)^{1/2} * (cx+1)^{1/2} * (cx-1)^{1/2} * c^4x^4 - 6cx^2 + 18(c^2x^2-1)^{1/2} * (cx+1)^{1/2} * c^2x^2 - (cx-1)^{1/2} * (cx+1)^{1/2} * (-1+6 \operatorname{arccosh}(cx))) * d^2 / (cx-1) / (cx+1) / c - 3/512(-d(c^2x^2-1))^{1/2} * (8c^5x^5 - 12c^3x^3 + 8(c^2x^2+d)^{1/2} * (cx-1)^{1/2} * c^4x^4 + 4cx - 8(c^2x^2-1)^{1/2} * (cx+1)^{1/2} * c^2x^2 + (cx-1)^{1/2} * (cx+1)^{1/2} * (-1+4 \operatorname{arccosh}(cx))) * d^2 / (cx-1) / (cx+1) / c + 15/256(-d(c^2x^2-1))^{1/2} * (2c^3x^3 - 2cx^2 + (cx-1)^{1/2} * (cx+1)^{1/2} * c^2x^2 - (cx-1)^{1/2} * (cx+1)^{1/2}) * (-1+2 \operatorname{arccosh}(cx))) * d^2 / (cx-1) / (cx+1) / c + 15/256(-d(c^2x^2-1))^{1/2} * (-2(c^2x^2+d)^{1/2} * (cx+1)^{1/2} * c^2x^2 + 2c^3x^3 + (cx-1)^{1/2} * (cx+1)^{1/2} - 2cx) * (1+2 \operatorname{arccosh}(cx))) * d^2 / (cx-1) / (cx+1) / c - 3/512(-d(c^2x^2-1))^{1/2} * (-8(c^2x^2+d)^{1/2} * (cx+1)^{1/2} * (cx-1)^{1/2} * c^4x^4 + 8c^5x^5 + 8(c^2x^2+d)^{1/2} * (cx+1)^{1/2} * c^2x^2 - 12c^3x^3 - (cx-1)^{1/2} * (cx+1)^{1/2} + 4cx) * (1+4 \operatorname{arccosh}(cx))) * d^2 / (cx-1) / (cx+1) / c + 1/2304(-d(c^2x^2-1))^{1/2} * (-32(c^2x^2+d)^{1/2} * (cx+1)^{1/2} * (cx-1)^{1/2} * c^6x^6 + 32c^7x^7 + 48(c^2x^2+d)^{1/2} * (cx-1)^{1/2} * c^4x^4 - 64c^5x^5 - 18(c^2x^2+d)^{1/2} * (cx+1)^{1/2} * c^2x^2 + 38c^3x^3 + (cx-1)^{1/2} * (cx+1)^{1/2} - 6cx) * (1+6 \operatorname{arccosh}(cx))) * d^2 / (cx-1) / (cx+1) / c + \dots$

### 3.89.5 Fracas [F]

$$\int (d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a) dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

**3.89.6 Sympy [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)`output `Timed out`**3.89.7 Maxima [F]**

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a) dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`output `1/48*(8*(-c^2*d*x^2 + d)^(5/2)*x + 10*(-c^2*d*x^2 + d)^(3/2)*d*x + 15*sqrt(-c^2*d*x^2 + d)*d^2*x + 15*d^(5/2)*arcsin(c*x)/c)*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)), x)`**3.89.8 Giac [F(-2)]**

Exception generated.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.89.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2),x)`output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2), x)`

### 3.90 $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.90.1	Optimal result	878
3.90.2	Mathematica [A] (warning: unable to verify)	879
3.90.3	Rubi [A] (verified)	879
3.90.4	Maple [A] (verified)	884
3.90.5	Fricas [F]	885
3.90.6	Sympy [F(-1)]	885
3.90.7	Maxima [F]	885
3.90.8	Giac [F(-2)]	886
3.90.9	Mupad [F(-1)]	886

#### 3.90.1 Optimal result

Integrand size = 27, antiderivative size = 284

$$\int \frac{(d - c^2dx^2)^{5/2} (a + b\operatorname{arccosh}(cx))}{x^2} dx = \frac{9bc^3d^2x^2\sqrt{d - c^2dx^2}}{16\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bc^5d^2x^4\sqrt{d - c^2dx^2}}{16\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{15}{8}c^2d^2x\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx)) - \frac{5}{4}c^2dx(d - c^2dx^2)^{3/2}(a + b\operatorname{arccosh}(cx)) - \frac{(d - c^2dx^2)^{5/2}(a + b\operatorname{arccosh}(cx))}{x} + \frac{15cd^2\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^2}{16b\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bcd^2\sqrt{d - c^2dx^2}\log(x)}{\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
-5/4*c^2*d*x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))-(-c^2*d*x^2+d)^(5/2)*
(a+b*arccosh(c*x))/x-15/8*c^2*d^2*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2
)+9/16*b*c^3*d^2*x^2*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/16
*b*c^5*d^2*x^4*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+15/16*c*d^
2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/(c*x-1)^(1/2)/(c*x+1)^(1/2)+
b*c*d^2*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.90.2 Mathematica [A] (warning: unable to verify)

Time = 1.57 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.07

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \frac{1}{128} d^2 \left( \frac{16a\sqrt{d - c^2 dx^2}(-8 - 9c^2 x^2 + 2c^4 x^4)}{x} \right. \\ + 240ac\sqrt{d} \arctan \left( \frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)} \right) \\ + 64bc\sqrt{d - c^2 dx^2} \left( -\frac{2\operatorname{arccosh}(cx)}{cx} + \frac{\operatorname{arccosh}(cx)^2 + 2\log(cx)}{\sqrt{\frac{-1+cx}{1+cx}}(1+cx)} \right) \\ + \frac{32bc\sqrt{d - c^2 dx^2}(2\operatorname{arccosh}(cx)^2 + \cosh(2\operatorname{arccosh}(cx)) - 2\operatorname{arccosh}(cx) \sinh(2\operatorname{arccosh}(cx)))}{\sqrt{\frac{-1+cx}{1+cx}}(1+cx)} \\ \left. - \frac{bc\sqrt{d - c^2 dx^2}(8\operatorname{arccosh}(cx)^2 + \cosh(4\operatorname{arccosh}(cx)) - 4\operatorname{arccosh}(cx) \sinh(4\operatorname{arccosh}(cx)))}{\sqrt{\frac{-1+cx}{1+cx}}(1+cx)} \right)$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^2,x]`

output `(d^2*((16*a*Sqrt[d - c^2*d*x^2]*(-8 - 9*c^2*x^2 + 2*c^4*x^4))/x + 240*a*c*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + 64*b*c*Sqrt[d - c^2*d*x^2]*((-2*ArcCosh[c*x])/(c*x) + (ArcCosh[c*x]^2 + 2*Log[c*x])/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))) + (32*b*c*Sqrt[d - c^2*d*x^2]*(2*ArcCosh[c*x]^2 + Cosh[2*ArcCosh[c*x]] - 2*ArcCosh[c*x]*Sinh[2*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) - (b*c*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))/128`

### 3.90.3 Rubi [A] (verified)

Time = 1.03 (sec) , antiderivative size = 316, normalized size of antiderivative = 1.11, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.481$ , Rules used = {6343, 82, 243, 49, 2009, 6312, 25, 82, 244, 2009, 6310, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.90.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx$



$$\begin{aligned}
& \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx \\
& \quad \downarrow \text{6343} \\
& -5c^2 d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} \\
& \quad \downarrow \text{82} \\
& -5c^2 d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} \\
& \quad \downarrow \text{243} \\
& -5c^2 d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2}{x^2} dx^2}{2\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} \\
& \quad \downarrow \text{49} \\
& -5c^2 d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (x^2 c^4 - 2c^2 + \frac{1}{x^2}) dx^2}{2\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} \\
& \quad \downarrow \text{2009} \\
& -5c^2 d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} + \\
& \quad \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 x^4}{2} - 2c^2 x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{6312} \\
& -5c^2 d \left( \frac{3}{4} d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd \sqrt{d - c^2 dx^2} \int -x(1-cx)(cx+1) dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4} x (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} + \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 x^4}{2} - 2c^2 x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{25}
\end{aligned}$$

---

3.90.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx$

$$-5c^2d \left( \frac{3}{4}d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))dx - \frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right) \\ + \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4x^4}{2} - 2c^2x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 82

$$-5c^2d \left( \frac{3}{4}d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))dx - \frac{bcd\sqrt{d-c^2dx^2} \int x(1-c^2x^2)dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right) \\ + \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4x^4}{2} - 2c^2x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 244

$$-5c^2d \left( \frac{3}{4}d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))dx - \frac{bcd\sqrt{d-c^2dx^2} \int (x-c^2x^3)dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right) \\ + \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4x^4}{2} - 2c^2x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2009

$$-5c^2d \left( \frac{3}{4}d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))dx + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd \left( \frac{x^2}{2} - \frac{c^2x^4}{4} \right) \sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \\ + \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4x^4}{2} - 2c^2x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6310

$$-5c^2d \left( \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}}dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \int xdx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) \right) + \frac{1}{4} \\ + \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4x^4}{2} - 2c^2x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 15

$$\begin{aligned}
& -5c^2d \left( \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx^2\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{4}x \right. \\
& \quad \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4x^4}{2} - 2c^2x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \\
& \quad \downarrow \text{6308} \\
& \quad -\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x} - \\
& 5c^2d \left( \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) + \frac{3}{4}d \left( \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{4bc\sqrt{cx-1}\sqrt{cx+1}} \right) \right. \\
& \quad \left. \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4x^4}{2} - 2c^2x^2 + \log(x^2) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right)
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^2,x]`

output `-(((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x) - 5*c^2*d*(-1/4*(b*c*d*Sqrt[d - c^2*d*x^2]*(x^2/2 - (c^2*x^4)/4))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/4 + (3*d*(-1/4*(b*c*x^2*Sqrt[d - c^2*d*x^2]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(4*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/4 + (b*c*d^2*Sqrt[d - c^2*d*x^2]*(-2*c^2*x^2 + (c^4*x^4)/2 + Log[x^2]))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.90.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

---

3.90.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x^2} dx$

- rule 82 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`
- rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(Sqrt[(d1_) + (e1_)*(x_)]*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^(n/2)), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^(n/2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`
- rule 6312 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n/(2*p + 1))), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

```
rule 6343 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^(2*(m + 1)))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1]
```

### 3.90.4 Maple [A] (verified)

Time = 1.09 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.07

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{dx} - ac^2x(-c^2dx^2+d)^{\frac{5}{2}} - \frac{5ac^2dx(-c^2dx^2+d)^{\frac{3}{2}}}{4} - \frac{15ac^2d^2x\sqrt{-c^2dx^2+d}}{8} - \frac{15ac^2d^3 \arctan\left(\frac{\sqrt{-c^2dx^2+d}}{\sqrt{-c^2d}}\right)}{8\sqrt{c^2d}}$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{dx} - ac^2x(-c^2dx^2+d)^{\frac{5}{2}} - \frac{5ac^2dx(-c^2dx^2+d)^{\frac{3}{2}}}{4} - \frac{15ac^2d^2x\sqrt{-c^2dx^2+d}}{8} - \frac{15ac^2d^3 \arctan\left(\frac{\sqrt{-c^2dx^2+d}}{\sqrt{-c^2d}}\right)}{8\sqrt{c^2d}}$

```
input int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)
```

```
output -a/d/x*(-c^2*d*x^2+d)^(7/2)-a*c^2*x*(-c^2*d*x^2+d)^(5/2)-5/4*a*c^2*d*x*(-c^2*d*x^2+d)^(3/2)-15/8*a*c^2*d^2*x*(-c^2*d*x^2+d)^(1/2)-15/8*a*c^2*d^3/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+1/128*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x*(32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4-8*c^5*x^5-144*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2+72*c^3*x^3+120*arccosh(c*x)^2*x*c-128*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-128*c*x*arccosh(c*x)+128*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-33*c*x)*d^2
```

---

3.90. 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^2} dx$$

**3.90.5 Fricas [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^2,x, algorithm="fricas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^2, x)`

**3.90.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**2,x)`

output `Timed out`

**3.90.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output `-1/8*(10*(-c^2*d*x^2 + d)^(3/2)*c^2*d*x + 15*sqrt(-c^2*d*x^2 + d)*c^2*d^2*x + 15*c*d^(5/2)*arcsin(c*x) + 8*(-c^2*d*x^2 + d)^(5/2)/x)*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/x^2, x)`

---

3.90.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx$

**3.90.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.90.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^2,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^2, x)`

**3.91** 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^4} dx$$

3.91.1	Optimal result	887
3.91.2	Mathematica [A] (warning: unable to verify)	888
3.91.3	Rubi [A] (verified)	888
3.91.4	Maple [A] (verified)	893
3.91.5	Fricas [F]	893
3.91.6	Sympy [F(-1)]	894
3.91.7	Maxima [F]	894
3.91.8	Giac [F(-2)]	894
3.91.9	Mupad [F(-1)]	895

**3.91.1 Optimal result**

Integrand size = 27, antiderivative size = 293

$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^4} dx = -\frac{bcd^2\sqrt{d-c^2dx^2}}{6x^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^5d^2x^2\sqrt{d-c^2dx^2}}{4\sqrt{-1+cx}\sqrt{1+cx}} + \frac{5}{2}c^4d^2x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) + \frac{5c^2d(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{3x} - \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{3x^3} - \frac{5c^3d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{4b\sqrt{-1+cx}\sqrt{1+cx}} - \frac{7bc^3d^2\sqrt{d-c^2dx^2}\log(x)}{3\sqrt{-1+cx}\sqrt{1+cx}}$$

output 
$$\frac{5}{3}c^2d(-c^2dx^2+d)^{(3/2)}(a+b\operatorname{arccosh}(cx))/x-1/3(-c^2dx^2+d)^{(5/2)}(a+b\operatorname{arccosh}(cx))/x^3+5/2c^4d^2x(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{(1/2)}-1/6b*c*d^2(-c^2dx^2+d)^{(1/2)}/x^2/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-1/4*b*c^5d^2x^2(-c^2dx^2+d)^{(1/2)}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-5/4c^3d^2(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{(1/2)}/b/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-7/3b*c^3d^2\ln(x)(-c^2dx^2+d)^{(1/2)}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}$$



### 3.91.2 Mathematica [A] (warning: unable to verify)

Time = 1.31 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.09

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^4} dx = \frac{30bc^3 d^3 x^3 (-1 + cx) \operatorname{arccosh}(cx)^2 - 60ac^3 d^{5/2} x^3 \sqrt{\frac{-1+cx}{1+cx}} \sqrt{d - c^2 d}}{x^4}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^4,x]`

output  $(30*b*c^3*d^3*x^3*(-1 + c*x)*\operatorname{ArcCosh}[c*x]^2 - 60*a*c^3*d^{5/2}*x^3*\operatorname{Sqrt}[(1 + c*x)/(1 + c*x)]*\operatorname{Sqrt}[d - c^2*d*x^2]*\operatorname{ArcTan}[(c*x*\operatorname{Sqrt}[d - c^2*d*x^2])/(\operatorname{Sqrt}[d]*(-1 + c^2*x^2))] + 3*b*c^3*d^3*x^3*(-1 + c*x)*\operatorname{Cosh}[2*\operatorname{ArcCosh}[c*x]] - 4*d^3*(b*c*x*(1 - c*x) + a*\operatorname{Sqrt}[(1 + c*x)/(1 + c*x)]*(2 - 16*c^2*x^2 + 11*c^4*x^4 + 3*c^6*x^6) - 14*b*c^3*x^3*(-1 + c*x)*\operatorname{Log}[c*x]) - 2*b*d^3*(-1 + c*x)*\operatorname{ArcCosh}[c*x]*(4*\operatorname{Sqrt}[(1 + c*x)/(1 + c*x)]*(-1 - c*x + 7*c^2*x^2 + 7*c^3*x^3) + 3*c^3*x^3*\operatorname{Sinh}[2*\operatorname{ArcCosh}[c*x]]))/(24*x^3*\operatorname{Sqrt}[(1 + c*x)/(1 + c*x)]*\operatorname{Sqrt}[d - c^2*d*x^2])$

### 3.91.3 Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 312, normalized size of antiderivative = 1.06, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.481$ , Rules used = {6343, 82, 243, 49, 2009, 6343, 25, 82, 244, 2009, 6310, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^4} dx$$

↓ 6343

$$-\frac{5}{3}c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{x^2} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2 dx}{x^3}}{3\sqrt{cx-1}\sqrt{cx+1}} -$$

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{3x^3}$$

↓ 82

---

3.91.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^4} dx$

$$\begin{aligned}
& -\frac{5}{3}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd^2\sqrt{d - c^2dx^2} \int \frac{(1-c^2x^2)^2}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \\
& \quad \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{243} \\
& -\frac{5}{3}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd^2\sqrt{d - c^2dx^2} \int \frac{(1-c^2x^2)^2}{x^4} dx^2}{6\sqrt{cx - 1}\sqrt{cx + 1}} - \\
& \quad \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{49} \\
& -\frac{5}{3}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd^2\sqrt{d - c^2dx^2} \int \left(c^4 - \frac{2c^2}{x^2} + \frac{1}{x^4}\right) dx^2}{6\sqrt{cx - 1}\sqrt{cx + 1}} - \\
& \quad \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{3x^3} \\
& \quad \downarrow \text{2009} \\
& -\frac{5}{3}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^2} dx - \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{3x^3} + \\
& \quad \frac{bcd^2\sqrt{d - c^2dx^2} \left(c^4x^2 - 2c^2 \log(x^2) - \frac{1}{x^2}\right)}{6\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6343} \\
& -\frac{5}{3}c^2d \left( -3c^2d \int \sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx)) dx - \frac{bcd\sqrt{d - c^2dx^2} \int -\frac{(1-cx)(cx+1)}{x} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} \right. \\
& \quad \left. \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d - c^2dx^2} \left(c^4x^2 - 2c^2 \log(x^2) - \frac{1}{x^2}\right)}{6\sqrt{cx - 1}\sqrt{cx + 1}} \right) \\
& \quad \downarrow \text{25} \\
& -\frac{5}{3}c^2d \left( -3c^2d \int \sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx)) dx + \frac{bcd\sqrt{d - c^2dx^2} \int \frac{(1-cx)(cx+1)}{x} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} \right. \\
& \quad \left. \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d - c^2dx^2} \left(c^4x^2 - 2c^2 \log(x^2) - \frac{1}{x^2}\right)}{6\sqrt{cx - 1}\sqrt{cx + 1}} \right) \\
& \quad \downarrow \text{82}
\end{aligned}$$

---

3.91.  $\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^4} dx$

$$-\frac{5}{3}c^2d \left( -3c^2d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))dx + \frac{bcd\sqrt{d-c^2dx^2} \int \frac{1-c^2x^2}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d-c^2dx^2}(c^4x^2-2c^2\log(x^2)-\frac{1}{x^2})}{6\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 244

$$-\frac{5}{3}c^2d \left( -3c^2d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))dx + \frac{bcd\sqrt{d-c^2dx^2} \int (\frac{1}{x}-c^2x) dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d-c^2dx^2}(c^4x^2-2c^2\log(x^2)-\frac{1}{x^2})}{6\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 2009

$$-\frac{5}{3}c^2d \left( -3c^2d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))dx - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bcd\sqrt{d-c^2dx^2}(\log(x)-\frac{1}{x})}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d-c^2dx^2}(c^4x^2-2c^2\log(x^2)-\frac{1}{x^2})}{6\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6310

$$-\frac{5}{3}c^2d \left( -3c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \int x dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d-c^2dx^2}(c^4x^2-2c^2\log(x^2)-\frac{1}{x^2})}{6\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 15

$$-\frac{5}{3}c^2d \left( -3c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx^2\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d-c^2dx^2}(c^4x^2-2c^2\log(x^2)-\frac{1}{x^2})}{6\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6308

$$-\frac{5}{3}c^2d \left( -3c^2d \left( \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4bc\sqrt{cx-1}\sqrt{cx+1}} - \frac{bcx^2\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd^2\sqrt{d-c^2dx^2}(c^4x^2-2c^2\log(x^2)-\frac{1}{x^2})}{6\sqrt{cx-1}\sqrt{cx+1}} \right)$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^3 - (5*c^2*d*(-(((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x) - 3*c^2*d*(-1/4*(b*c*x^2*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(4*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (b*c*d*Sqrt[d - c^2*d*x^2]*(-1/2*(c^2*x^2) + Log[x]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/3 + (b*c*d^2*Sqrt[d - c^2*d*x^2]*(-x^(-2) + c^4*x^2 - 2*c^2*Log[x^2]))/(6*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.91.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

---

3.91.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x^4} dx$

- rule 244 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^(n/2)), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^(n/2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`
- rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1]`

### 3.91.4 Maple [A] (verified)

Time = 1.14 (sec) , antiderivative size = 340, normalized size of antiderivative = 1.16

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{3dx^3} + \frac{4ac^2(-c^2dx^2+d)^{\frac{7}{2}}}{3dx} + \frac{4ac^4x(-c^2dx^2+d)^{\frac{5}{2}}}{3} + \frac{5ac^4dx(-c^2dx^2+d)^{\frac{3}{2}}}{3} + \frac{5ac^4d^2x\sqrt{-c^2dx^2+d}}{2} + \dots$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{3dx^3} + \frac{4ac^2(-c^2dx^2+d)^{\frac{7}{2}}}{3dx} + \frac{4ac^4x(-c^2dx^2+d)^{\frac{5}{2}}}{3} + \frac{5ac^4dx(-c^2dx^2+d)^{\frac{3}{2}}}{3} + \frac{5ac^4d^2x\sqrt{-c^2dx^2+d}}{2} + \dots$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)`

output

$$-1/3*a/d/x^3*(-c^2*d*x^2+d)^(7/2)+4/3*a*c^2/d/x*(-c^2*d*x^2+d)^(7/2)+4/3*a*c^4*x*(-c^2*d*x^2+d)^(5/2)+5/3*a*c^4*d*x*(-c^2*d*x^2+d)^(3/2)+5/2*a*c^4*d^2*x*(-c^2*d*x^2+d)^(1/2)+5/2*a*c^4*d^3/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))-1/24*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^3*(-12*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4+6*c^5*x^5+30*arccosh(c*x)^2*x^3*c^3+56*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3-56*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-56*c^3*x^3*arccosh(c*x)-3*c^3*x^3+8*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c*x)*d^2$$

### 3.91.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)}{x^4} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^4, x)`

---

3.91.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^4} dx$

**3.91.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^4} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**4,x)`

output `Timed out`

**3.91.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)}{x^4} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")`

output `1/6*(10*(-c^2*d*x^2 + d)^(3/2)*c^4*d*x + 15*sqrt(-c^2*d*x^2 + d)*c^4*d^2*x + 15*c^3*d^(5/2)*arcsin(c*x) + 8*(-c^2*d*x^2 + d)^(5/2)*c^2/x - 2*(-c^2*d*x^2 + d)^(7/2)/(d*x^3))*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^4, x)`

**3.91.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.91.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^4} dx$

**3.91.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^4,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^4, x)`



**3.92**  $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^6} dx$

3.92.1	Optimal result	896
3.92.2	Mathematica [A] (warning: unable to verify)	897
3.92.3	Rubi [A] (verified)	897
3.92.4	Maple [B] (verified)	902
3.92.5	Fricas [F]	903
3.92.6	Sympy [F(-1)]	903
3.92.7	Maxima [F]	903
3.92.8	Giac [F(-2)]	904
3.92.9	Mupad [F(-1)]	904

**3.92.1 Optimal result**

Integrand size = 27, antiderivative size = 293

$$\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^6} dx = -\frac{bcd^2\sqrt{d - c^2dx^2}}{20x^4\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{11bc^3d^2\sqrt{d - c^2dx^2}}{30x^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{c^4d^2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{x} + \frac{c^2d(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))}{3x^3} - \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{5x^5} + \frac{c^5d^2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{2b\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{23bc^5d^2\sqrt{d - c^2dx^2} \log(x)}{15\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output 1/3*c^2*d*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/x^3-1/5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^5-c^4*d^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x-1/20*b*c*d^2*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)+11/30*b*c^3*d^2*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/2*c^5*d^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/b/(c*x-1)^(1/2)/(c*x+1)^(1/2)+23/15*b*c^5*d^2*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.92.2 Mathematica [A] (warning: unable to verify)

Time = 2.94 (sec) , antiderivative size = 400, normalized size of antiderivative = 1.37

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \frac{d^2 \left( 8ad \sqrt{\frac{-1+cx}{1+cx}} (-1 + c^2 x^2) (3 - 11c^2 x^2 + 23c^4 x^4) + 120ac^5 \sqrt{dx^5} \right)}{x^6}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^6,x]`

output `(d^2*(8*a*d*Sqrt[(-1 + c*x)/(1 + c*x)]*(-1 + c^2*x^2)*(3 - 11*c^2*x^2 + 23*c^4*x^4) + 120*a*c^5*Sqrt[d]*x^5*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + 40*b*c^2*d*x^2*(1 - c*x)*(c*x - 2*((-1 + c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3*ArcCosh[c*x] + 2*c^3*x^3*Log[c*x]) - 60*b*c^4*d*x^4*(1 - c*x)*(2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x] - c*x*(ArcCosh[c*x]^2 + 2*Log[c*x])) - b*d*(1 - c*x)*(20*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x] + Cosh[5*ArcCosh[c*x]]*Log[c*x] + Cosh[3*ArcCosh[c*x]]*(-1 + 5*Log[c*x]) + c*x*(3 + 10*Log[c*x]) - 5*ArcCosh[c*x]*Sinh[3*ArcCosh[c*x]] - ArcCosh[c*x]*Sinh[5*ArcCosh[c*x]])))/(120*x^5*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2])`

### 3.92.3 Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.07, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.481$ , Rules used = {6343, 82, 243, 49, 2009, 6343, 25, 82, 244, 2009, 6339, 14, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx$$

$$\downarrow \text{6343}$$

$$-c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2}{x^5} dx}{5\sqrt{cx - 1}\sqrt{cx + 1}}$$

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5}$$

$$\downarrow \text{82}$$

---

3.92.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx$

$$\begin{aligned}
& -c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1 - c^2 x^2)^2}{x^5} dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} - \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5} \\
& \quad \downarrow \text{243} \\
& -c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1 - c^2 x^2)^2}{x^6} dx^2}{10\sqrt{cx - 1}\sqrt{cx + 1}} - \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5} \\
& \quad \downarrow \text{49} \\
& -c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \left( \frac{c^4}{x^2} - \frac{2c^2}{x^4} + \frac{1}{x^6} \right) dx^2}{10\sqrt{cx - 1}\sqrt{cx + 1}} - \\
& \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5} \\
& \quad \downarrow \text{2009} \\
& -c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^4} dx - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5} + \\
& \quad \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx - 1}\sqrt{cx + 1}} \\
& \quad \downarrow \text{6343} \\
& -c^2 d \left( c^2 (-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^2} dx - \frac{bcd \sqrt{d - c^2 dx^2} \int \frac{(1 - cx)(cx + 1)}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3x^3} \right. \\
& \quad \left. \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5} + \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx - 1}\sqrt{cx + 1}} \right) \\
& \quad \downarrow \text{25} \\
& -c^2 d \left( c^2 (-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd \sqrt{d - c^2 dx^2} \int \frac{(1 - cx)(cx + 1)}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3x^3} \right. \\
& \quad \left. \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5} + \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx - 1}\sqrt{cx + 1}} \right) \\
& \quad \downarrow \text{82}
\end{aligned}$$

$$-c^2d \left( c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \frac{1-c^2x^2}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{5x^5} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 244

$$-c^2d \left( c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^2} dx + \frac{bcd\sqrt{d-c^2dx^2} \int \left( \frac{1}{x^3} - \frac{c^2}{x} \right) dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{5x^5} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 2009

$$-c^2d \left( c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x^2} dx - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} + \frac{bcd\sqrt{d-c^2dx^2}(c^2(-d))}{3\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{5x^5} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6339

$$-c^2d \left( c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{bc\sqrt{d-c^2dx^2} \int \frac{1}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} \right) - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{5x^5} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 14

$$-c^2d \left( c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} + \frac{bc \log(x)\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{3x^3} \right. \\ \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{5x^5} + \frac{bcd^2\sqrt{d-c^2dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6308

$$c^2 d \left( c^2 (-d) \left( \frac{c \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{2b \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x} + \frac{bc \log(x) \sqrt{d - c^2 dx^2}}{\sqrt{cx - 1} \sqrt{cx + 1}} \right) - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{5x^5} \right) - \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( c^4 \log(x^2) + \frac{2c^2}{x^2} - \frac{1}{2x^4} \right)}{10 \sqrt{cx - 1} \sqrt{cx + 1}}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^6,x]`

output `-1/5*((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^5 - c^2*d*(-1/3*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/x^3 + (b*c*d*Sqrt[d - c^2*d*x^2]*(-1/2*1/x^2 - c^2*Log[x]))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - c^2*d*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/x) + (c*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (b*c*Sqrt[d - c^2*d*x^2]*Log[x])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (b*c*d^2*Sqrt[d - c^2*d*x^2]*(-1/2*1/x^4 + (2*c^2)/x^2 + c^4*Log[x^2]))/(10*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.92.3.1 Defintions of rubi rules used

rule 14 `Int[(a_)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 82 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

- rule 244 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand  
Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p  
, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt  
[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 +  
c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[  
c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1  
&& EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6339 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_)*Sqrt[(d_) +  
(e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc  
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e  
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x  
)^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1  
+ c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1  
+ c*x]*Sqrt[-1 + c*x])), x], x) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`
- rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_)*((d_) + (e_  
.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc  
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m  
+ 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(  
m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)  
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x],  
x) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && G  
tQ[p, 0] && LtQ[m, -1]`

### 3.92.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2428 vs.  $2(251) = 502$ .

Time = 1.22 (sec) , antiderivative size = 2429, normalized size of antiderivative = 8.29

method	result	size
default	Expression too large to display	2429
parts	Expression too large to display	2429

```
input int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^6,x,method=_RETURNVERBOSE)
```

```
output 1/2*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)^2*d^
2*c^5+23/15*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*ln(1+(c*x
+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*d^2*c^5-46/15*b*(-d*(c^2*x^2-1))^(1/2)/(c
*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*d^2*c^5-175/4*b*(-d*(c^2*x^2-1))^(1
/2)*d^2/(1035*c^8*x^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x^2+9)/(c*x+1)^(1/2)/
(c*x-1)^(1/2)*c^5+2/15*a*c^2/d/x^3*(-c^2*d*x^2+d)^(7/2)-1329/4*b*(-d*(c^2*
x^2-1))^(1/2)*d^2/(1035*c^8*x^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x^2+9)*x^4/
(c*x+1)^(1/2)/(c*x-1)^(1/2)*c^9+1889/12*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(1035
*c^8*x^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x^2+9)*x^2/(c*x+1)^(1/2)/(c*x-1)^(
1/2)*c^7+141/20*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(1035*c^8*x^8-765*c^6*x^6+325
*c^4*x^4-75*c^2*x^2+9)/x^2/(c*x+1)^(1/2)/(c*x-1)^(1/2)*c^3-5819/30*b*(-d*(
c^2*x^2-1))^(1/2)*d^2/(1035*c^8*x^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x^2+9)*
x^9/(c*x+1)/(c*x-1)*c^14+18791/60*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(1035*c^8*x
^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x^2+9)*x^7/(c*x+1)/(c*x-1)*c^12-943/6*b*
(-d*(c^2*x^2-1))^(1/2)*d^2/(1035*c^8*x^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x^
2+9)*x^5/(c*x+1)/(c*x-1)*c^10+207/5*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(1035*c^8
*x^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x^2+9)*x^3/(c*x+1)/(c*x-1)*c^8-69/20*b
*(-d*(c^2*x^2-1))^(1/2)*d^2/(1035*c^8*x^8-765*c^6*x^6+325*c^4*x^4-75*c^2*x
^2+9)*x/(c*x+1)/(c*x-1)*c^6+9/5*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(1035*c^8*x^8
-765*c^6*x^6+325*c^4*x^4-75*c^2*x^2+9)/x^5/(c*x+1)/(c*x-1)*arccosh(c*x)...
```

### 3.92.5 Fracas [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x^6} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^6,x, algorithm="fracas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^6, x)`

### 3.92.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**6,x)`

output `Timed out`

### 3.92.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x^6} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^6,x, algorithm="maxima")`

output `-1/15*(10*(-c^2*d*x^2 + d)^(3/2)*c^6*d*x + 15*sqrt(-c^2*d*x^2 + d)*c^6*d^2*x + 15*c^5*d^(5/2)*arcsin(c*x) + 8*(-c^2*d*x^2 + d)^(5/2)*c^4/x - 2*(-c^2*d*x^2 + d)^(7/2)*c^2/(d*x^3) + 3*(-c^2*d*x^2 + d)^(7/2)/(d*x^5))*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/x^6, x)`

---

3.92.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx$



**3.92.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^6,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.92.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^6} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^6} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^6,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^6, x)`

### 3.93 $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^8} dx$

3.93.1	Optimal result	905
3.93.2	Mathematica [A] (verified)	905
3.93.3	Rubi [A] (verified)	906
3.93.4	Maple [B] (verified)	908
3.93.5	Fricas [A] (verification not implemented)	909
3.93.6	Sympy [F(-1)]	909
3.93.7	Maxima [C] (verification not implemented)	910
3.93.8	Giac [F(-2)]	910
3.93.9	Mupad [F(-1)]	911

#### 3.93.1 Optimal result

Integrand size = 27, antiderivative size = 219

$$\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^8} dx = -\frac{bcd^2\sqrt{d - c^2dx^2}}{42x^6\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{3bc^3d^2\sqrt{d - c^2dx^2}}{28x^4\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{3bc^5d^2\sqrt{d - c^2dx^2}}{14x^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{7/2} (a + \operatorname{arccosh}(cx))}{7dx^7} - \frac{bc^7d^2\sqrt{d - c^2dx^2} \log(x)}{7\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output 
$$-1/7*(-c^2*d*x^2+d)^{(7/2)}*(a+b*\operatorname{arccosh}(c*x))/d/x^7-1/42*b*c*d^2*(-c^2*d*x^2+d)^{(1/2)}/x^6/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3/28*b*c^3*d^2*(-c^2*d*x^2+d)^{(1/2)}/x^4/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3/14*b*c^5*d^2*(-c^2*d*x^2+d)^{(1/2)}/x^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/7*b*c^7*d^2*\ln(x)*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

#### 3.93.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.48

$$\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^8} dx = \frac{d^2\sqrt{d - c^2dx^2}(12(-1 + cx)^{7/2}(1 + cx)^{7/2}(a + \operatorname{arccosh}(cx)) - bc^7d^2\sqrt{d - c^2dx^2} \log(x))}{84x^7\sqrt{-1 + cx}\sqrt{1 + cx}}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^8,x]`

output `(d^2*Sqrt[d - c^2*d*x^2]*(12*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(a + b*ArcCosh[c*x]) - b*c*x*(2 - 9*c^2*x^2 + 18*c^4*x^4 + 12*c^6*x^6*Log[x])))/(84*x^7*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.93.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.51, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6332, 25, 82, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{x^8} dx \\
 & \quad \downarrow \text{6332} \\
 & - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int - \frac{(1-cx)^3 (cx+1)^3}{x^7} dx}{7\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \text{barccosh}(cx))}{7dx^7} \\
 & \quad \downarrow \text{25} \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^3 (cx+1)^3}{x^7} dx}{7\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \text{barccosh}(cx))}{7dx^7} \\
 & \quad \downarrow \text{82} \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^3}{x^7} dx}{7\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \text{barccosh}(cx))}{7dx^7} \\
 & \quad \downarrow \text{243} \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^3}{x^8} dx^2}{14\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \text{barccosh}(cx))}{7dx^7} \\
 & \quad \downarrow \text{49} \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \left( -\frac{c^6}{x^2} + \frac{3c^4}{x^4} - \frac{3c^2}{x^6} + \frac{1}{x^8} \right) dx^2}{14\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \text{barccosh}(cx))}{7dx^7} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.93.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{x^8} dx$

$$\frac{bcd^2\sqrt{d-c^2dx^2}\left(c^6(-\log(x^2))-\frac{3c^4}{x^2}+\frac{3c^2}{2x^4}-\frac{1}{3x^6}\right)}{14\sqrt{cx-1}\sqrt{cx+1}}-\frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{7dx^7}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^8,x]`

output `-1/7*((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(d*x^7) + (b*c*d^2*sqrt[d - c^2*d*x^2]*(-1/3*1/x^6 + (3*c^2)/(2*x^4) - (3*c^4)/x^2 - c^6*Log[x^2]))/(14*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.93.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

---

3.93.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x^8} dx$

### 3.93.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3774 vs.  $2(183) = 366$ .

Time = 1.33 (sec) , antiderivative size = 3775, normalized size of antiderivative = 17.24

method	result	size
default	Expression too large to display	3775
parts	Expression too large to display	3775

```
input int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^8,x,method=_RETURNVERBOSE)
```

```
output 2/7*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*d^2*
c^7-1/7*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*ln(1+(c*x+(c*
x-1)^(1/2)*(c*x+1)^(1/2))^2)*d^2*c^7+55/12*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(7
*c^12*x^12-21*c^10*x^10+35*c^8*x^8-35*c^6*x^6+21*c^4*x^4-7*c^2*x^2+1)/(c*x
+1)^(1/2)/(c*x-1)^(1/2)*c^7-17/84*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(7*c^12*x^1
2-21*c^10*x^10+35*c^8*x^8-35*c^6*x^6+21*c^4*x^4-7*c^2*x^2+1)*x^3/(c*x+1)/(
c*x-1)*c^10+1/42*b*(-d*(c^2*x^2-1))^(1/2)*d^2/(7*c^12*x^12-21*c^10*x^10+35
*c^8*x^8-35*c^6*x^6+21*c^4*x^4-7*c^2*x^2+1)*x/(c*x+1)/(c*x-1)*c^8+1/7*b*(-
d*(c^2*x^2-1))^(1/2)*d^2/(7*c^12*x^12-21*c^10*x^10+35*c^8*x^8-35*c^6*x^6+2
1*c^4*x^4-7*c^2*x^2+1)/x^7/(c*x+1)/(c*x-1)*arccosh(c*x)-3*b*(-d*(c^2*x^2-1
))^(1/2)*d^2/(7*c^12*x^12-21*c^10*x^10+35*c^8*x^8-35*c^6*x^6+21*c^4*x^4-7*
c^2*x^2+1)*x^4/(c*x+1)^(1/2)/(c*x-1)^(1/2)*arccosh(c*x)*c^11+b*(-d*(c^2*x^
2-1))^(1/2)*d^2/(7*c^12*x^12-21*c^10*x^10+35*c^8*x^8-35*c^6*x^6+21*c^4*x^4
-7*c^2*x^2+1)*x^2/(c*x+1)^(1/2)/(c*x-1)^(1/2)*arccosh(c*x)*c^9+3*b*(-d*(c^
2*x^2-1))^(1/2)*d^2/(7*c^12*x^12-21*c^10*x^10+35*c^8*x^8-35*c^6*x^6+21*c^4
*x^4-7*c^2*x^2+1)*x^10/(c*x+1)^(1/2)/(c*x-1)^(1/2)*arccosh(c*x)*c^17-5*b*(
-d*(c^2*x^2-1))^(1/2)*d^2/(7*c^12*x^12-21*c^10*x^10+35*c^8*x^8-35*c^6*x^6+
21*c^4*x^4-7*c^2*x^2+1)*x^8/(c*x+1)^(1/2)/(c*x-1)^(1/2)*arccosh(c*x)*c^15-
b*(-d*(c^2*x^2-1))^(1/2)*d^2/(7*c^12*x^12-21*c^10*x^10+35*c^8*x^8-35*c^6*x
^6+21*c^4*x^4-7*c^2*x^2+1)*x^12/(c*x+1)^(1/2)/(c*x-1)^(1/2)*arccosh(c*x...
```

**3.93.5 Fracas [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 703, normalized size of antiderivative = 3.21

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \frac{12 (bc^8 d^2 x^8 - 4bc^6 d^2 x^6 + 6bc^4 d^2 x^4 - 4bc^2 d^2 x^2 + bd^2) \sqrt{-c^2 dx^2 + d}}{c^2 dx^4 - (c^2 + 1) dx^2 + d} - 12 (bc^8 d^2 x^8 - 4bc^6 d^2 x^6 + 6bc^4 d^2 x^4 - 4bc^2 d^2 x^2 + bd^2) \sqrt{d} \arctan \left( \frac{\sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} (x^2 + 1) \sqrt{d}}{c^2 dx^4 - (c^2 + 1) dx^2 + d} \right)$$

```
input integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^8,x, algorithm="fracas")
```

```
output [1/84*(12*(b*c^8*d^2*x^8 - 4*b*c^6*d^2*x^6 + 6*b*c^4*d^2*x^4 - 4*b*c^2*d^2*x^2 + b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + 6*(b*c^9*d^2*x^9 - b*c^7*d^2*x^7)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) - (18*b*c^5*d^2*x^5 - (18*b*c^5 - 9*b*c^3 + 2*b*c)*d^2*x^7 - 9*b*c^3*d^2*x^3 + 2*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 12*(a*c^8*d^2*x^8 - 4*a*c^6*d^2*x^6 + 6*a*c^4*d^2*x^4 - 4*a*c^2*d^2*x^2 + a*d^2)*sqrt(-c^2*d*x^2 + d)/(c^2*x^9 - x^7), -1/84*(12*(b*c^9*d^2*x^9 - b*c^7*d^2*x^7)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 12*(b*c^8*d^2*x^8 - 4*b*c^6*d^2*x^6 + 6*b*c^4*d^2*x^4 - 4*b*c^2*d^2*x^2 + b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + (18*b*c^5*d^2*x^5 - (18*b*c^5 - 9*b*c^3 + 2*b*c)*d^2*x^7 - 9*b*c^3*d^2*x^3 + 2*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 12*(a*c^8*d^2*x^8 - 4*a*c^6*d^2*x^6 + 6*a*c^4*d^2*x^4 - 4*a*c^2*d^2*x^2 + a*d^2)*sqrt(-c^2*d*x^2 + d))/(c^2*x^9 - x^7)]
```

**3.93.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \text{Timed out}$$

```
input integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**8,x)
```

```
output Timed out
```

---

3.93.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^8} dx$

**3.93.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.02

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \frac{\left(6 c^8 d^4 \sqrt{-\frac{1}{c^4 d}} \log\left(x^2 - \frac{1}{c^2}\right) + 6i (-1)^{-2c^2 dx^2 + 2d} c^6 d^{7/2} \log(-2c^2 d)\right)}{7 dx^7} - \frac{(-c^2 dx^2 + d)^{7/2} b \operatorname{arccosh}(cx)}{7 dx^7} - \frac{(-c^2 dx^2 + d)^{7/2} a}{7 dx^7}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^8,x, algorithm="maxima")`

output `1/84*(6*c^8*d^4*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2) + 6*I*(-1)^(-2*c^2*d*x^2 + 2*d)*c^6*d^(7/2)*log(-2*c^2*d + 2*d/x^2) + 11*sqrt(-c^4*d*x^4 + 2*c^2*d*x^2 - d)*c^4*d^3/x^2 - 7*sqrt(-c^4*d*x^4 + 2*c^2*d*x^2 - d)*c^2*d^3/x^4 + 2*sqrt(-c^4*d*x^4 + 2*c^2*d*x^2 - d)*d^3/x^6)*b*c/d - 1/7*(-c^2*d*x^2 + d)^(7/2)*b*arccosh(c*x)/(d*x^7) - 1/7*(-c^2*d*x^2 + d)^(7/2)*a/(d*x^7)`

**3.93.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^8,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.93.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^8} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^8} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^8,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^8, x)`



**3.94**  $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^{10}} dx$

3.94.1	Optimal result . . . . .	912
3.94.2	Mathematica [A] (verified) . . . . .	913
3.94.3	Rubi [A] (verified) . . . . .	913
3.94.4	Maple [B] (verified) . . . . .	916
3.94.5	Fricas [A] (verification not implemented) . . . . .	916
3.94.6	Sympy [F(-1)] . . . . .	917
3.94.7	Maxima [A] (verification not implemented) . . . . .	918
3.94.8	Giac [F(-2)] . . . . .	918
3.94.9	Mupad [F(-1)] . . . . .	919

**3.94.1 Optimal result**

Integrand size = 27, antiderivative size = 314

$$\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^{10}} dx = -\frac{bc^3d^2\sqrt{d - c^2dx^2}}{189x^6\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^5d^2\sqrt{d - c^2dx^2}}{42x^4\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bc^7d^2\sqrt{d - c^2dx^2}}{21x^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcd^2(1 - c^2x^2)^4\sqrt{d - c^2dx^2}}{72x^8\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{7/2} (a + \operatorname{arccosh}(cx))}{9dx^9} - \frac{2c^2(d - c^2dx^2)^{7/2} (a + \operatorname{arccosh}(cx))}{63dx^7} - \frac{2bc^9d^2\sqrt{d - c^2dx^2} \log(x)}{63\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output -1/9*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/d/x^9-2/63*c^2*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/d/x^7-1/189*b*c^3*d^2*(-c^2*d*x^2+d)^(1/2)/x^6/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/42*b*c^5*d^2*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/21*b*c^7*d^2*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/72*b*c*d^2*(-c^2*x^2+1)^4*(-c^2*d*x^2+d)^(1/2)/x^8/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/63*b*c^9*d^2*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.94.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.47

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \frac{d^2 \sqrt{d - c^2 dx^2} (168(-1 + cx)^{7/2} (1 + cx)^{7/2} (a + \operatorname{barccosh}(cx)) + 48c^2 x^2 (-1 + cx)^{7/2} (1 + cx)^{7/2} (a + \operatorname{barccosh}(cx)) - b c x (21 - 76c^2 x^2 + 90c^4 x^4 - 12c^6 x^6 + 48c^8 x^8 \operatorname{Log}[x]))}{1512 x^9 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^10,x]`

output `(d^2*sqrt[d - c^2*d*x^2]*(168*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(a + b*ArcCosh[c*x]) + 48*c^2*x^2*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(a + b*ArcCosh[c*x]) - b*c*x*(21 - 76*c^2*x^2 + 90*c^4*x^4 - 12*c^6*x^6 + 48*c^8*x^8*Log[x]))/(1512*x^9*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.94.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.55, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6337, 27, 354, 87, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx \\ & \quad \downarrow \text{6337} \\ & \frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d^2(1-c^2x^2)^3(2c^2x^2+7)}{63x^9} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{9dx^9} - \\ & \quad \frac{2c^2(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{63dx^7} \\ & \quad \downarrow \text{27} \\ & \frac{bcd^2\sqrt{d - c^2 dx^2} \int \frac{(1-c^2x^2)^3(2c^2x^2+7)}{x^9} dx}{63\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{9dx^9} - \\ & \quad \frac{2c^2(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{63dx^7} \\ & \quad \downarrow \text{354} \end{aligned}$$

---

3.94.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx$

$$\begin{aligned}
& \frac{bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^3(2c^2x^2+7)}{x^{10}} dx^2}{126\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{9dx^9} - \\
& \quad \frac{2c^2(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{63dx^7} \\
& \quad \downarrow 87 \\
& \frac{bcd^2\sqrt{d-c^2dx^2} \left( 2c^2 \int \frac{(1-c^2x^2)^3}{x^8} dx^2 - \frac{7(1-c^2x^2)^4}{4x^8} \right)}{126\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{9dx^9} - \\
& \quad \frac{2c^2(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{63dx^7} \\
& \quad \downarrow 49 \\
& \frac{bcd^2\sqrt{d-c^2dx^2} \left( 2c^2 \int \left( -\frac{c^6}{x^2} + \frac{3c^4}{x^4} - \frac{3c^2}{x^6} + \frac{1}{x^8} \right) dx^2 - \frac{7(1-c^2x^2)^4}{4x^8} \right)}{126\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{9dx^9} - \frac{2c^2(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{63dx^7} \\
& \quad \downarrow 2009 \\
& - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{9dx^9} - \frac{2c^2(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{63dx^7} + \\
& \frac{bcd^2\sqrt{d-c^2dx^2} \left( 2c^2 \left( c^6(-\log(x^2)) - \frac{3c^4}{x^2} + \frac{3c^2}{2x^4} - \frac{1}{3x^6} \right) - \frac{7(1-c^2x^2)^4}{4x^8} \right)}{126\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^10,x]`

output `-1/9*((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(d*x^9) - (2*c^2*(d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(63*d*x^7) + (b*c*d^2*sqrt[d - c^2*d*x^2]*((-7*(1 - c^2*x^2)^4)/(4*x^8) + 2*c^2*(-1/3*1/x^6 + (3*c^2)/(2*x^4) - (3*c^4)/x^2 - c^6*Log[x^2])))/(126*sqrt[-1 + c*x]*sqrt[1 + c*x])`

## 3.94.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`
- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !EqQ[e, 0] || !EqQ[c, 0] || LtQ[p, n]))`
- rule 354 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[x^(m - 1)/2*(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[(m - 1)/2]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.94.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 5006 vs.  $2(266) = 532$ .

Time = 1.32 (sec) , antiderivative size = 5007, normalized size of antiderivative = 15.95

method	result	size
default	Expression too large to display	5007
parts	Expression too large to display	5007

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^10,x,method=_RETURNVERBOSE)`

output `result too large to display`

### 3.94.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 795, normalized size of antiderivative = 2.53

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^{10}} dx = \left[ \frac{24 (2 bc^{10} d^2 x^{10} - bc^8 d^2 x^8 - 16 bc^6 d^2 x^6 + 34 bc^4 d^2 x^4 - 26 bc^2 d^2 x^2 + 48 (bc^{11} d^2 x^{11} - bc^9 d^2 x^9) \sqrt{d} \arctan \left( \frac{\sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} (x^2 + 1) \sqrt{d}}{c^2 dx^4 - (c^2 + 1) dx^2 + d} \right) - 24 (2 bc^{10} d^2 x^{10} - bc^8 d^2 x^8 - 16 bc^6 d^2 x^6 + \dots}{\dots} \right]$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^10,x, algorithm="fracas")`

output `[1/1512*(24*(2*b*c^10*d^2*x^10 - b*c^8*d^2*x^8 - 16*b*c^6*d^2*x^6 + 34*b*c^4*d^2*x^4 - 26*b*c^2*d^2*x^2 + 7*b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + 24*(b*c^11*d^2*x^11 - b*c^9*d^2*x^9)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1))*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + (12*b*c^7*d^2*x^7 - 90*b*c^5*d^2*x^5 - (12*b*c^7 - 90*b*c^5 + 76*b*c^3 - 21*b*c)*d^2*x^9 + 76*b*c^3*d^2*x^3 - 21*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 24*(2*a*c^10*d^2*x^10 - a*c^8*d^2*x^8 - 16*a*c^6*d^2*x^6 + 34*a*c^4*d^2*x^4 - 26*a*c^2*d^2*x^2 + 7*a*d^2)*sqrt(-c^2*d*x^2 + d))/(c^2*x^11 - x^9), -1/1512*(48*(b*c^11*d^2*x^11 - b*c^9*d^2*x^9)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 24*(2*b*c^10*d^2*x^10 - b*c^8*d^2*x^8 - 16*b*c^6*d^2*x^6 + 34*b*c^4*d^2*x^4 - 26*b*c^2*d^2*x^2 + 7*b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (12*b*c^7*d^2*x^7 - 90*b*c^5*d^2*x^5 - (12*b*c^7 - 90*b*c^5 + 76*b*c^3 - 21*b*c)*d^2*x^9 + 76*b*c^3*d^2*x^3 - 21*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 24*(2*a*c^10*d^2*x^10 - a*c^8*d^2*x^8 - 16*a*c^6*d^2*x^6 + 34*a*c^4*d^2*x^4 - 26*a*c^2*d^2*x^2 + 7*a*d^2)*sqrt(-c^2*d*x^2 + d))/(c^2*x^11 - x^9)]`

### 3.94.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**10,x)`

output `Timed out`

**3.94.7 Maxima [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 187, normalized size of antiderivative = 0.60

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx =$$

$$-\frac{1}{1512} \left( 48 c^8 \sqrt{-d} d^2 \log(x) - \frac{12 c^6 \sqrt{-d} d^2 x^6 - 90 c^4 \sqrt{-d} d^2 x^4 + 76 c^2 \sqrt{-d} d^2 x^2 - 21 \sqrt{-d} d^2}{x^8} \right) bc$$

$$-\frac{1}{63} b \left( \frac{2 (-c^2 dx^2 + d)^{7/2} c^2}{dx^7} + \frac{7 (-c^2 dx^2 + d)^{7/2}}{dx^9} \right) \operatorname{arcosh}(cx)$$

$$-\frac{1}{63} a \left( \frac{2 (-c^2 dx^2 + d)^{7/2} c^2}{dx^7} + \frac{7 (-c^2 dx^2 + d)^{7/2}}{dx^9} \right)$$

```
input integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^10,x, algorithm="maxima")
```

```
output -1/1512*(48*c^8*sqrt(-d)*d^2*log(x) - (12*c^6*sqrt(-d)*d^2*x^6 - 90*c^4*sqrt(-d)*d^2*x^4 + 76*c^2*sqrt(-d)*d^2*x^2 - 21*sqrt(-d)*d^2)/x^8)*b*c - 1/63*b*(2*(-c^2*d*x^2 + d)^(7/2)*c^2/(d*x^7) + 7*(-c^2*d*x^2 + d)^(7/2)/(d*x^9))*arccosh(c*x) - 1/63*a*(2*(-c^2*d*x^2 + d)^(7/2)*c^2/(d*x^7) + 7*(-c^2*d*x^2 + d)^(7/2)/(d*x^9))
```

**3.94.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^10,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.94.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{10}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^{10}} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^10,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^10, x)`



### 3.95 $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^{12}} dx$

3.95.1	Optimal result	920
3.95.2	Mathematica [A] (verified)	921
3.95.3	Rubi [A] (verified)	921
3.95.4	Maple [B] (verified)	923
3.95.5	Fricas [A] (verification not implemented)	923
3.95.6	Sympy [F(-1)]	924
3.95.7	Maxima [A] (verification not implemented)	925
3.95.8	Giac [F(-2)]	925
3.95.9	Mupad [F(-1)]	926

#### 3.95.1 Optimal result

Integrand size = 27, antiderivative size = 385

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))}{x^{12}} dx = -\frac{bcd^2 \sqrt{d - c^2 dx^2}}{110x^{10} \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{23bc^3 d^2 \sqrt{d - c^2 dx^2}}{792x^8 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{113bc^5 d^2 \sqrt{d - c^2 dx^2}}{4158x^6 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{bc^7 d^2 \sqrt{d - c^2 dx^2}}{924x^4 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{2bc^9 d^2 \sqrt{d - c^2 dx^2}}{693x^2 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{arccosh}(cx))}{11dx^{11}} - \frac{4c^2 (d - c^2 dx^2)^{7/2} (a + \operatorname{arccosh}(cx))}{99dx^9} - \frac{8c^4 (d - c^2 dx^2)^{7/2} (a + \operatorname{arccosh}(cx))}{693dx^7} - \frac{8bc^{11} d^2 \sqrt{d - c^2 dx^2} \log(x)}{693 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

```
output -1/11*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/d/x^11-4/99*c^2*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/d/x^9-8/693*c^4*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/d/x^7-1/110*b*c*d^2*(-c^2*d*x^2+d)^(1/2)/x^10/(c*x-1)^(1/2)/(c*x+1)^(1/2)+23/792*b*c^3*d^2*(-c^2*d*x^2+d)^(1/2)/x^8/(c*x-1)^(1/2)/(c*x+1)^(1/2)-113/4158*b*c^5*d^2*(-c^2*d*x^2+d)^(1/2)/x^6/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/924*b*c^7*d^2*(-c^2*d*x^2+d)^(1/2)/x^4/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/693*b*c^9*d^2*(-c^2*d*x^2+d)^(1/2)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-8/693*b*c^11*d^2*ln(x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.95.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.43

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \frac{d^2 \sqrt{d - c^2 dx^2} (7560(-1 + cx)^{7/2} (1 + cx)^{7/2} (a + \operatorname{barccosh}(cx)) +$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^12,x]`

output `(d^2*Sqrt[d - c^2*d*x^2]*(7560*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(a + b*ArcCosh[c*x]) + 480*c^2*x^2*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(7 + 2*c^2*x^2)*(a + b*ArcCosh[c*x]) - b*c*x*(756 - 2415*c^2*x^2 + 2260*c^4*x^4 - 90*c^6*x^6 - 240*c^8*x^8 + 960*c^10*x^10*Log[x]))/(83160*x^11*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.95.3 Rubi [A] (verified)

Time = 0.52 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.53, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6337, 27, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx \\ & \quad \downarrow \text{6337} \\ & \frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d^2(1-c^2x^2)^3(8c^4x^4+28c^2x^2+63)}{693x^{11}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{11dx^{11}} \\ & \quad \frac{4c^2(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{99dx^9} - \frac{8c^4(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{693dx^7} \\ & \quad \downarrow \text{27} \\ & \frac{bcd^2\sqrt{d - c^2 dx^2} \int \frac{(1-c^2x^2)^3(8c^4x^4+28c^2x^2+63)}{x^{11}} dx}{693\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{11dx^{11}} \\ & \quad \frac{4c^2(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{99dx^9} - \frac{8c^4(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{693dx^7} \\ & \quad \downarrow \text{1578} \end{aligned}$$

---

3.95.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx$

$$\begin{aligned}
& \frac{bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^3(8c^4x^4+28c^2x^2+63)}{x^{12}} dx^2}{1386\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{11dx^{11}} - \\
& \frac{4c^2(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{99dx^9} - \frac{8c^4(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{693dx^7} \\
& \quad \downarrow \text{1195} \\
& \frac{bcd^2\sqrt{d-c^2dx^2} \int \left(-\frac{8c^{10}}{x^2} - \frac{4c^8}{x^4} - \frac{3c^6}{x^6} + \frac{113c^4}{x^8} - \frac{161c^2}{x^{10}} + \frac{63}{x^{12}}\right) dx^2}{1386\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{11dx^{11}} - \frac{4c^2(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{99dx^9} - \\
& \frac{8c^4(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{693dx^7} \\
& \quad \downarrow \text{2009} \\
& \frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{11dx^{11}} - \frac{4c^2(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{99dx^9} - \\
& \frac{8c^4(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{693dx^7} + \\
& \frac{bcd^2\sqrt{d-c^2dx^2} \left(-8c^{10} \log(x^2) + \frac{4c^8}{x^2} + \frac{3c^6}{2x^4} - \frac{113c^4}{3x^6} + \frac{161c^2}{4x^8} - \frac{63}{5x^{10}}\right)}{1386\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^12,x]`

output `-1/11*((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(d*x^11) - (4*c^2*(d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(99*d*x^9) - (8*c^4*(d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(693*d*x^7) + (b*c*d^2*sqrt[d - c^2*d*x^2]*(-63/(5*x^10) + (161*c^2)/(4*x^8) - (113*c^4)/(3*x^6) + (3*c^6)/(2*x^4) + (4*c^8)/x^2 - 8*c^10*Log[x^2]))/(1386*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.95.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1195 `Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))^(n_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

---

3.95.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x^{12}} dx$

rule 1578 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])]`

### 3.95.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 6381 vs.  $2(325) = 650$ .

Time = 1.47 (sec) , antiderivative size = 6382, normalized size of antiderivative = 16.58

method	result	size
default	Expression too large to display	6382
parts	Expression too large to display	6382

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^12,x,method=_RETURNVERBOSE)`

output `result too large to display`

### 3.95.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 879, normalized size of antiderivative = 2.28

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^{12}} dx = \frac{120 (8 bc^{12} d^2 x^{12} - 4 bc^{10} d^2 x^{10} - bc^8 d^2 x^8 - 116 bc^6 d^2 x^6 + 274 bc^4 d^2 x^4 - 960 (bc^{13} d^2 x^{13} - bc^{11} d^2 x^{11}) \sqrt{d} \arctan \left( \frac{\sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} (x^2 + 1) \sqrt{d}}{c^2 dx^4 - (c^2 + 1) dx^2 + d} \right) - 120 (8 bc^{12} d^2 x^{12} - 4 bc^{10} d^2 x^{10} - bc^8 d^2 x^8)}{c^2 dx^4 - (c^2 + 1) dx^2 + d}$$

---

3.95.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^{12}} dx$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^12,x, algorithm="fricas")`

output `[1/83160*(120*(8*b*c^12*d^2*x^12 - 4*b*c^10*d^2*x^10 - b*c^8*d^2*x^8 - 116*b*c^6*d^2*x^6 + 274*b*c^4*d^2*x^4 - 224*b*c^2*d^2*x^2 + 63*b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + 480*(b*c^13*d^2*x^13 - b*c^11*d^2*x^11)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + sqrt(-c^2*d*x^2 + d))*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) + (240*b*c^9*d^2*x^9 + 90*b*c^7*d^2*x^7 - (240*b*c^9 + 90*b*c^7 - 2260*b*c^5 + 2415*b*c^3 - 756*b*c)*d^2*x^11 - 2260*b*c^5*d^2*x^5 + 2415*b*c^3*d^2*x^3 - 756*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 120*(8*a*c^12*d^2*x^12 - 4*a*c^10*d^2*x^10 - a*c^8*d^2*x^8 - 116*a*c^6*d^2*x^6 + 274*a*c^4*d^2*x^4 - 224*a*c^2*d^2*x^2 + 63*a*d^2)*sqrt(-c^2*d*x^2 + d))/(c^2*x^13 - x^11), -1/83160*(960*(b*c^13*d^2*x^13 - b*c^11*d^2*x^11)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 120*(8*b*c^12*d^2*x^12 - 4*b*c^10*d^2*x^10 - b*c^8*d^2*x^8 - 116*b*c^6*d^2*x^6 + 274*b*c^4*d^2*x^4 - 224*b*c^2*d^2*x^2 + 63*b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (240*b*c^9*d^2*x^9 + 90*b*c^7*d^2*x^7 - (240*b*c^9 + 90*b*c^7 - 2260*b*c^5 + 2415*b*c^3 - 756*b*c)*d^2*x^11 - 2260*b*c^5*d^2*x^5 + 2415*b*c^3*d^2*x^3 - 756*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 120*(8*a*c^12*d^2*x^12 - 4*a*c^10*d^2*x^10 - a*c^8*d^2*x^8 - 116*a*c^6*d^2*x^6 + 274*a*c^4*d^2*x^4 - 224*a*c^2*d^2*x^2 + 63*a*d^2)*sqrt(-c^2*d*x^2 + d))/(c^2*x^13 - x^11)]`

### 3.95.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**12,x)`

output `Timed out`

---

3.95.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx$

**3.95.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.65

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx =$$

$$-\frac{1}{83160} \left( 960 c^{10} \sqrt{-dd^2} \log(x) - \frac{240 c^8 \sqrt{-dd^2} x^8 + 90 c^6 \sqrt{-dd^2} x^6 - 2260 c^4 \sqrt{-dd^2} x^4 + 2415 c^2 \sqrt{-dd^2} x^2}{x^{10}} \right.$$

$$-\frac{1}{693} b \left( \frac{8(-c^2 dx^2 + d)^{7/2} c^4}{dx^7} + \frac{28(-c^2 dx^2 + d)^{7/2} c^2}{dx^9} + \frac{63(-c^2 dx^2 + d)^{7/2}}{dx^{11}} \right) \operatorname{arcosh}(cx)$$

$$-\frac{1}{693} a \left( \frac{8(-c^2 dx^2 + d)^{7/2} c^4}{dx^7} + \frac{28(-c^2 dx^2 + d)^{7/2} c^2}{dx^9} + \frac{63(-c^2 dx^2 + d)^{7/2}}{dx^{11}} \right)$$

```
input integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^12,x, algorithm="maxima")
```

```
output -1/83160*(960*c^10*sqrt(-d)*d^2*log(x) - (240*c^8*sqrt(-d)*d^2*x^8 + 90*c^6*sqrt(-d)*d^2*x^6 - 2260*c^4*sqrt(-d)*d^2*x^4 + 2415*c^2*sqrt(-d)*d^2*x^2 - 756*sqrt(-d)*d^2)/x^10)*b*c - 1/693*b*(8*(-c^2*d*x^2 + d)^(7/2)*c^4/(d*x^7) + 28*(-c^2*d*x^2 + d)^(7/2)*c^2/(d*x^9) + 63*(-c^2*d*x^2 + d)^(7/2)/(d*x^11))*arccosh(c*x) - 1/693*a*(8*(-c^2*d*x^2 + d)^(7/2)*c^4/(d*x^7) + 28*(-c^2*d*x^2 + d)^(7/2)*c^2/(d*x^9) + 63*(-c^2*d*x^2 + d)^(7/2)/(d*x^11))
```

**3.95.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^12,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value
```

---

3.95.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx$

**3.95.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^{12}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^{12}} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^12,x)`output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^12, x)`

### 3.96 $\int x^7 (d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx)) dx$

3.96.1	Optimal result	927
3.96.2	Mathematica [A] (verified)	928
3.96.3	Rubi [A] (verified)	928
3.96.4	Maple [B] (verified)	930
3.96.5	Fricas [A] (verification not implemented)	931
3.96.6	Sympy [F(-1)]	932
3.96.7	Maxima [A] (verification not implemented)	932
3.96.8	Giac [F(-2)]	933
3.96.9	Mupad [F(-1)]	933

#### 3.96.1 Optimal result

Integrand size = 27, antiderivative size = 458

$$\begin{aligned} \int x^7 (d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx)) dx &= \frac{16bd^2 x \sqrt{d - c^2 dx^2}}{3003c^7 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &+ \frac{8bd^2 x^3 \sqrt{d - c^2 dx^2}}{9009c^5 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{2bd^2 x^5 \sqrt{d - c^2 dx^2}}{5005c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &+ \frac{5bd^2 x^7 \sqrt{d - c^2 dx^2}}{21021c \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{53bcd^2 x^9 \sqrt{d - c^2 dx^2}}{3861 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &+ \frac{27bc^3 d^2 x^{11} \sqrt{d - c^2 dx^2}}{1573 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{bc^5 d^2 x^{13} \sqrt{d - c^2 dx^2}}{169 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &- \frac{(d - c^2 dx^2)^{7/2} (a + b \operatorname{arccosh}(cx))}{7c^8 d} + \frac{(d - c^2 dx^2)^{9/2} (a + b \operatorname{arccosh}(cx))}{3c^8 d^2} \\ &- \frac{3(d - c^2 dx^2)^{11/2} (a + b \operatorname{arccosh}(cx))}{11c^8 d^3} + \frac{(d - c^2 dx^2)^{13/2} (a + b \operatorname{arccosh}(cx))}{13c^8 d^4} \end{aligned}$$

output 
$$\begin{aligned} &-1/7*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/c^8/d+1/3*(-c^2*d*x^2+d)^(9/2) \\ &)*(a+b*arccosh(c*x))/c^8/d^2-3/11*(-c^2*d*x^2+d)^(11/2)*(a+b*arccosh(c*x)) \\ &/c^8/d^3+1/13*(-c^2*d*x^2+d)^(13/2)*(a+b*arccosh(c*x))/c^8/d^4+16/3003*b*d \\ &^2*x*(-c^2*d*x^2+d)^(1/2)/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)+8/9009*b*d^2*x^3 \\ &*(-c^2*d*x^2+d)^(1/2)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/5005*b*d^2*x^5*(-c \\ &^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/21021*b*d^2*x^7*(-c^2* \\ &d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-53/3861*b*c*d^2*x^9*(-c^2*d*x \\ &^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+27/1573*b*c^3*d^2*x^11*(-c^2*d*x^2 \\ &+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/169*b*c^5*d^2*x^13*(-c^2*d*x^2+d)^( \\ &(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2) \end{aligned}$$



### 3.96.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.39

$$\int x^7 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d^2 \sqrt{d - c^2 dx^2} (bc(720720x + 120120c^2x^3 + 54054c^4x^5 + 32175c^6x^7 - 1856855c^8x^9 + 2321865c^{10}x^{11} - 800415c^{12}x^{13}) + 10405395c^6x^6(-1 + cx)^{7/2}(1 + cx)^{7/2}(a + b\operatorname{ArcCosh}[cx]) + 90090(-1 + cx)^{7/2}(1 + cx)^{7/2}(8 + 28c^2x^2 + 63c^4x^4)(a + b\operatorname{ArcCosh}[cx]))}{(135270135c^8\sqrt{-1 + cx}\sqrt{1 + cx})}$$

input `Integrate[x^7*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]), x]`

output  $(d^2\sqrt{d - c^2dx^2}*(b*c*(720720*x + 120120*c^2*x^3 + 54054*c^4*x^5 + 32175*c^6*x^7 - 1856855*c^8*x^9 + 2321865*c^{10}*x^{11} - 800415*c^{12}*x^{13}) + 10405395*c^6*x^6*(-1 + c*x)^{(7/2)}*(1 + c*x)^{(7/2)}*(a + b*\operatorname{ArcCosh}[c*x]) + 90090*(-1 + c*x)^{(7/2)}*(1 + c*x)^{(7/2)}*(8 + 28*c^2*x^2 + 63*c^4*x^4)*(a + b*\operatorname{ArcCosh}[c*x])))/(135270135*c^8*\sqrt{-1 + c*x}*\sqrt{1 + c*x})$

### 3.96.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.53, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 2341, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^7 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow 6337 \\ & - \frac{bc\sqrt{d - c^2 dx^2} \int - \frac{d^2(1 - c^2 x^2)^3 (231c^6 x^6 + 126c^4 x^4 + 56c^2 x^2 + 16)}{3003c^8} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \\ & \frac{(d - c^2 dx^2)^{13/2} (a + \operatorname{barccosh}(cx))}{3c^8 d^4} - \frac{3(d - c^2 dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{7c^8 d} + \\ & \frac{(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{3c^8 d^2} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^8 d} \\ & \quad \downarrow 27 \end{aligned}$$

$$\begin{aligned}
& \frac{bd^2\sqrt{d-c^2dx^2} \int (1-c^2x^2)^3 (231c^6x^6 + 126c^4x^4 + 56c^2x^2 + 16) dx}{3003c^7\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{(d-c^2dx^2)^{13/2} (a + \operatorname{barccosh}(cx))}{3c^8d^2} - \frac{3(d-c^2dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{7c^8d} + \\
& \frac{13c^8d^4}{3c^8d^2} \frac{(d-c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{3c^8d^2} - \frac{11c^8d^3}{7c^8d} \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^8d} \\
& \quad \downarrow \text{2341} \\
& \frac{bd^2\sqrt{d-c^2dx^2} \int (-231c^{12}x^{12} + 567c^{10}x^{10} - 371c^8x^8 + 5c^6x^6 + 6c^4x^4 + 8c^2x^2 + 16) dx}{3003c^7\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{(d-c^2dx^2)^{13/2} (a + \operatorname{barccosh}(cx))}{3c^8d^2} - \frac{3(d-c^2dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{7c^8d} + \\
& \frac{13c^8d^4}{3c^8d^2} \frac{(d-c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{3c^8d^2} - \frac{11c^8d^3}{7c^8d} \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^8d} \\
& \quad \downarrow \text{2009} \\
& \frac{(d-c^2dx^2)^{13/2} (a + \operatorname{barccosh}(cx))}{3c^8d^2} - \frac{3(d-c^2dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{7c^8d} + \\
& \frac{13c^8d^4}{3c^8d^2} \frac{(d-c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{3c^8d^2} - \frac{11c^8d^3}{7c^8d} \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^8d} + \\
& \frac{bd^2 \left( -\frac{231}{13}c^{12}x^{13} + \frac{567c^{10}x^{11}}{11} - \frac{371c^8x^9}{9} + \frac{5c^6x^7}{7} + \frac{6c^4x^5}{5} + \frac{8c^2x^3}{3} + 16x \right) \sqrt{d-c^2dx^2}}{3003c^7\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^7*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `(b*d^2*sqrt[d - c^2*d*x^2]*(16*x + (8*c^2*x^3)/3 + (6*c^4*x^5)/5 + (5*c^6*x^7)/7 - (371*c^8*x^9)/9 + (567*c^10*x^11)/11 - (231*c^12*x^13)/13))/(3003*c^7*sqrt[-1 + c*x]*sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(7*c^8*d) + ((d - c^2*d*x^2)^(9/2)*(a + b*ArcCosh[c*x]))/(3*c^8*d^2) - (3*(d - c^2*d*x^2)^(11/2)*(a + b*ArcCosh[c*x]))/(11*c^8*d^3) + ((d - c^2*d*x^2)^(13/2)*(a + b*ArcCosh[c*x]))/(13*c^8*d^4)`

## 3.96.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2341 Int[(Pq_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[p, -2]
```

```
rule 6337 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])
```

## 3.96.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2373 vs. 2(386) = 772.

Time = 0.97 (sec) , antiderivative size = 2374, normalized size of antiderivative = 5.18

method	result	size
default	Expression too large to display	2374
parts	Expression too large to display	2374

```
input int(x^7*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

output

```

a*(-1/13*x^6*(-c^2*d*x^2+d)^(7/2)/c^2/d+6/13/c^2*(-1/11*x^4*(-c^2*d*x^2+d)
^(7/2)/c^2/d+4/11/c^2*(-1/9*x^2*(-c^2*d*x^2+d)^(7/2)/c^2/d-2/63/d/c^4*(-c^
2*d*x^2+d)^(7/2))))+b*(1/1384448*(-d*(c^2*x^2-1))^(1/2)*(-1-16896*c^8*x^8-
364*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4096*c^14*x^14-9984*(c*x+1)^(1/2)*
(c*x-1)^(1/2)*x^7*c^7-1204*c^4*x^4+85*c^2*x^2+13*(c*x-1)^(1/2)*(c*x+1)^(1/
2)*c*x+2912*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-15360*c^12*x^12+6496*c^6*x
^6+22784*c^10*x^10+4096*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^13*c^13-13312*(c*x+1
)^(1/2)*(c*x-1)^(1/2)*x^11*c^11+16640*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^9*c^9)
*(-1+13*arccosh(c*x))*d^2/(c*x+1)/c^8/(c*x-1)+1/991232*(-d*(c^2*x^2-1))^(1
/2)*(1+4096*c^8*x^8+220*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+2816*(c*x+1)^(
1/2)*(c*x-1)^(1/2)*x^7*c^7+620*c^4*x^4-61*c^2*x^2-11*(c*x-1)^(1/2)*(c*x+1)
^(1/2)*c*x-1232*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+1024*c^12*x^12-2352*c^
6*x^6-3328*c^10*x^10+1024*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^11*c^11-2816*(c*x+
1)^(1/2)*(c*x-1)^(1/2)*x^9*c^9)*(-1+11*arccosh(c*x))*d^2/(c*x+1)/c^8/(c*x-
1)-1/110592*(-d*(c^2*x^2-1))^(1/2)*(256*c^10*x^10-704*c^8*x^8+256*(c*x+1)^(
1/2)*(c*x-1)^(1/2)*x^9*c^9+688*c^6*x^6-576*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^
7*c^7-280*c^4*x^4+432*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+41*c^2*x^2-120*(
c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+9*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-
1+9*arccosh(c*x))*d^2/(c*x+1)/c^8/(c*x-1)-3/200704*(-d*(c^2*x^2-1))^(1/2)*
(64*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*...

```

### 3.96.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 353, normalized size of antiderivative = 0.77

$$\int x^7 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{45045 (231 bc^{14} d^2 x^{14} - 798 bc^{12} d^2 x^{12} + 938 bc^{10} d^2 x^{10} - 376 bc^8 d^2 x^8 - bc^6 d^2 x^6 - 2 bc^4 d^2 x^4 + bc^2 d^2 x^2 + d^2)}{c^8 (c^2 x^2 - 1)^{5/2}}$$

input `integrate(x^7*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $1/135270135*(45045*(231*b*c^{14}*d^2*x^{14} - 798*b*c^{12}*d^2*x^{12} + 938*b*c^{10}*d^2*x^{10} - 376*b*c^8*d^2*x^8 - b*c^6*d^2*x^6 - 2*b*c^4*d^2*x^4 - 8*b*c^2*d^2*x^2 + 16*b*d^2)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) - (800415*b*c^{13}*d^2*x^{13} - 2321865*b*c^{11}*d^2*x^{11} + 1856855*b*c^9*d^2*x^9 - 32175*b*c^7*d^2*x^7 - 54054*b*c^5*d^2*x^5 - 120120*b*c^3*d^2*x^3 - 720720*b*c*d^2*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} + 45045*(231*a*c^{14}*d^2*x^{14} - 798*a*c^{12}*d^2*x^{12} + 938*a*c^{10}*d^2*x^{10} - 376*a*c^8*d^2*x^8 - a*c^6*d^2*x^6 - 2*a*c^4*d^2*x^4 - 8*a*c^2*d^2*x^2 + 16*a*d^2)*\sqrt{-c^2*d*x^2 + d})/(c^{10}*x^2 - c^8)$

### 3.96.6 Sympy [F(-1)]

Timed out.

$$\int x^7 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**7*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)`

output `Timed out`

### 3.96.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 313, normalized size of antiderivative = 0.68

$$\int x^7 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{1}{3003} \left( \frac{231 (-c^2 dx^2 + d)^{7/2} x^6}{c^2 d} + \frac{126 (-c^2 dx^2 + d)^{7/2} x^4}{c^4 d} + \frac{56 (-c^2 dx^2 + d)^{7/2} x^2}{c^6 d} + \frac{16 (-c^2 dx^2 + d)^{7/2}}{c^8 d} \right) b \operatorname{arccos}$$

$$-\frac{1}{3003} \left( \frac{231 (-c^2 dx^2 + d)^{7/2} x^6}{c^2 d} + \frac{126 (-c^2 dx^2 + d)^{7/2} x^4}{c^4 d} + \frac{56 (-c^2 dx^2 + d)^{7/2} x^2}{c^6 d} + \frac{16 (-c^2 dx^2 + d)^{7/2}}{c^8 d} \right) a$$

$$\frac{(800415 c^{12} \sqrt{-dd^2} x^{13} - 2321865 c^{10} \sqrt{-dd^2} x^{11} + 1856855 c^8 \sqrt{-dd^2} x^9 - 32175 c^6 \sqrt{-dd^2} x^7 - 54054 c^4 \sqrt{-dd^2})}{135270135 c^7}$$

input `integrate(x^7*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output 
$$\begin{aligned} & -1/3003*(231*(-c^2*d*x^2 + d)^{(7/2)}*x^6/(c^2*d) + 126*(-c^2*d*x^2 + d)^{(7/2)}*x^4/(c^4*d) + 56*(-c^2*d*x^2 + d)^{(7/2)}*x^2/(c^6*d) + 16*(-c^2*d*x^2 + d)^{(7/2)}/(c^8*d))*b*\operatorname{arccosh}(c*x) \\ & - 1/3003*(231*(-c^2*d*x^2 + d)^{(7/2)}*x^6/(c^2*d) + 126*(-c^2*d*x^2 + d)^{(7/2)}*x^4/(c^4*d) + 56*(-c^2*d*x^2 + d)^{(7/2)}*x^2/(c^6*d) + 16*(-c^2*d*x^2 + d)^{(7/2)}/(c^8*d))*a \\ & - 1/135270135*(800415*c^{12}*\sqrt{-d}*d^2*x^{13} - 2321865*c^{10}*\sqrt{-d}*d^2*x^{11} + 1856855*c^8*\sqrt{-d}*d^2*x^9 - 32175*c^6*\sqrt{-d}*d^2*x^7 - 54054*c^4*\sqrt{-d}*d^2*x^5 - 120120*c^2*\sqrt{-d}*d^2*x^3 - 720720*\sqrt{-d}*d^2*x)*b/c^7 \end{aligned}$$

### 3.96.8 Giac [F(-2)]

Exception generated.

$$\int x^7(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^7*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.96.9 Mupad [F(-1)]

Timed out.

$$\int x^7(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int x^7 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} dx$$

input `int(x^7*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2),x)`

output `int(x^7*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2), x)`

### 3.97 $\int x^5(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

3.97.1	Optimal result	934
3.97.2	Mathematica [A] (verified)	935
3.97.3	Rubi [A] (verified)	935
3.97.4	Maple [B] (verified)	937
3.97.5	Fricas [A] (verification not implemented)	938
3.97.6	Sympy [F(-1)]	938
3.97.7	Maxima [A] (verification not implemented)	939
3.97.8	Giac [F(-2)]	939
3.97.9	Mupad [F(-1)]	940

#### 3.97.1 Optimal result

Integrand size = 27, antiderivative size = 378

$$\int x^5(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{8bd^2x\sqrt{d - c^2dx^2}}{693c^5\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{4bd^2x^3\sqrt{d - c^2dx^2}}{2079c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bd^2x^5\sqrt{d - c^2dx^2}}{1155c\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{113bcd^2x^7\sqrt{d - c^2dx^2}}{4851\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{23bc^3d^2x^9\sqrt{d - c^2dx^2}}{891\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{bc^5d^2x^{11}\sqrt{d - c^2dx^2}}{121\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6d}$$

$$+ \frac{2(d - c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^6d^2} - \frac{(d - c^2dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{11c^6d^3}$$

output

```
-1/7*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/c^6/d+2/9*(-c^2*d*x^2+d)^(9/2)
*(a+b*arccosh(c*x))/c^6/d^2-1/11*(-c^2*d*x^2+d)^(11/2)*(a+b*arccosh(c*x))
/c^6/d^3+8/693*b*d^2*x*(-c^2*d*x^2+d)^(1/2)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)
)+4/2079*b*d^2*x^3*(-c^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/
1155*b*d^2*x^5*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-113/4851
*b*c*d^2*x^7*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+23/891*b*c^3
*d^2*x^9*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/121*b*c^5*d^2*
x^11*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.97.2 Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.43

$$\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d^2 \sqrt{d - c^2 dx^2} (bc(27720x + 4620c^2 x^3 + 2079c^4 x^5 - 55935c^6 x^7 + 61985c^8 x^9 - 19845c^{10} + \operatorname{barccosh}(cx))}{dx}$$

input `Integrate[x^5*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`output `(d^2*sqrt[d - c^2*d*x^2]*(b*c*(27720*x + 4620*c^2*x^3 + 2079*c^4*x^5 - 55935*c^6*x^7 + 61985*c^8*x^9 - 19845*c^10*x^11) + 218295*c^4*x^4*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(a + b*ArcCosh[c*x]) + 13860*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(2 + 7*c^2*x^2)*(a + b*ArcCosh[c*x]))/(2401245*c^6*sqrt[-1 + c*x]*sqrt[1 + c*x])`**3.97.3 Rubi [A] (verified)**Time = 0.54 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.52, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6337$$

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{d^2(1 - c^2 x^2)^3 (63c^4 x^4 + 28c^2 x^2 + 8)}{693c^6} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{11c^6 d^3} +$$

$$\frac{2(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^6 d^2} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d}$$

$$\downarrow 27$$

$$\frac{bd^2\sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^3 (63c^4 x^4 + 28c^2 x^2 + 8) dx}{693c^5\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{11/2} (a + \operatorname{barccosh}(cx))}{11c^6 d^3} +$$

$$\frac{2(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^6 d^2} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^6 d}$$

---

3.97.  $\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$



$$\begin{aligned}
 & \downarrow 1467 \\
 & \frac{bd^2\sqrt{d-c^2dx^2} \int (-63c^{10}x^{10} + 161c^8x^8 - 113c^6x^6 + 3c^4x^4 + 4c^2x^2 + 8) dx}{693c^5\sqrt{cx-1}\sqrt{cx+1}} \\
 & \frac{(d-c^2dx^2)^{11/2}(a+\operatorname{barccosh}(cx))}{11c^6d^3} + \frac{2(d-c^2dx^2)^{9/2}(a+\operatorname{barccosh}(cx))}{9c^6d^2} - \\
 & \frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{7c^6d} \\
 & \downarrow 2009 \\
 & -\frac{(d-c^2dx^2)^{11/2}(a+\operatorname{barccosh}(cx))}{11c^6d^3} + \frac{2(d-c^2dx^2)^{9/2}(a+\operatorname{barccosh}(cx))}{9c^6d^2} - \\
 & \frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{7c^6d} + \\
 & \frac{bd^2\left(-\frac{63}{11}c^{10}x^{11} + \frac{161c^8x^9}{9} - \frac{113c^6x^7}{7} + \frac{3c^4x^5}{5} + \frac{4c^2x^3}{3} + 8x\right)\sqrt{d-c^2dx^2}}{693c^5\sqrt{cx-1}\sqrt{cx+1}}
 \end{aligned}$$

input `Int[x^5*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `(b*d^2*sqrt[d - c^2*d*x^2]*(8*x + (4*c^2*x^3)/3 + (3*c^4*x^5)/5 - (113*c^6*x^7)/7 + (161*c^8*x^9)/9 - (63*c^10*x^11)/11))/(693*c^5*sqrt[-1 + c*x]*sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(7*c^6*d) + (2*(d - c^2*d*x^2)^(9/2)*(a + b*ArcCosh[c*x]))/(9*c^6*d^2) - ((d - c^2*d*x^2)^(11/2)*(a + b*ArcCosh[c*x]))/(11*c^6*d^3)`

### 3.97.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 6337 Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_
), x_Symbol] :> With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCo
sh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c
*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b
, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)]
&& (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])
```

### 3.97.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1839 vs.  $2(318) = 636$ .

Time = 0.76 (sec) , antiderivative size = 1840, normalized size of antiderivative = 4.87

method	result	size
default	Expression too large to display	1840
parts	Expression too large to display	1840

```
input int(x^5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(-1/11*x^4*(-c^2*d*x^2+d)^(7/2)/c^2/d+4/11/c^2*(-1/9*x^2*(-c^2*d*x^2+d)^(
7/2)/c^2/d-2/63/d/c^4*(-c^2*d*x^2+d)^(7/2)))+b*(1/247808*(-d*(c^2*x^2-1))
^(1/2)*(1+4096*c^8*x^8+220*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+2816*(c*x+1
)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+620*c^4*x^4-61*c^2*x^2-11*(c*x-1)^(1/2)*(c*x
+1)^(1/2)*c*x-1232*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+1024*c^12*x^12-2352
*c^6*x^6-3328*c^10*x^10+1024*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^11*c^11-2816*(c
*x+1)^(1/2)*(c*x-1)^(1/2)*x^9*c^9)*(-1+11*arccosh(c*x))*d^2/(c*x+1)/c^6/(c
*x-1)-1/165888*(-d*(c^2*x^2-1))^(1/2)*(256*c^10*x^10-704*c^8*x^8+256*(c*x+
1)^(1/2)*(c*x-1)^(1/2)*x^9*c^9+688*c^6*x^6-576*(c*x+1)^(1/2)*(c*x-1)^(1/2)
*x^7*c^7-280*c^4*x^4+432*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+41*c^2*x^2-12
0*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+9*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)
*(-1+9*arccosh(c*x))*d^2/(c*x+1)/c^6/(c*x-1)-5/100352*(-d*(c^2*x^2-1))^(1/
2)*(64*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*
x^4-112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-25*c^2*x^2+56*(c*x-1)^(1/2)*(c
*x+1)^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+7*arccosh(c*x
))*d^2/(c*x+1)/c^6/(c*x-1)+1/10240*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c
^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*
(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c
*x))*d^2/(c*x+1)/c^6/(c*x-1)+5/9216*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^
2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/...
```

**3.97.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 317, normalized size of antiderivative = 0.84

$$\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{3465 (63 bc^{12} d^2 x^{12} - 224 bc^{10} d^2 x^{10} + 274 bc^8 d^2 x^8 - 116 bc^6 d^2 x^6 - bc^4 d^2 x^4 - 4 bc^2 d^2 x^2 + \operatorname{barccosh}(cx))}{d^2}$$

```
input integrate(x^5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")
```

```
output 1/2401245*(3465*(63*b*c^12*d^2*x^12 - 224*b*c^10*d^2*x^10 + 274*b*c^8*d^2*x^8 - 116*b*c^6*d^2*x^6 - b*c^4*d^2*x^4 - 4*b*c^2*d^2*x^2 + 8*b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (19845*b*c^11*d^2*x^11 - 61985*b*c^9*d^2*x^9 + 55935*b*c^7*d^2*x^7 - 2079*b*c^5*d^2*x^5 - 4620*b*c^3*d^2*x^3 - 27720*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 3465*(63*a*c^12*d^2*x^12 - 224*a*c^10*d^2*x^10 + 274*a*c^8*d^2*x^8 - 116*a*c^6*d^2*x^6 - a*c^4*d^2*x^4 - 4*a*c^2*d^2*x^2 + 8*a*d^2)*sqrt(-c^2*d*x^2 + d))/(c^8*x^2 - c^6)
```

**3.97.6 Sympy [F(-1)]**

Timed out.

$$\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

```
input integrate(x**5*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)
```

```
output Timed out
```

**3.97.7 Maxima [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.66

$$\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{1}{693} \left( \frac{63(-c^2 dx^2 + d)^{7/2} x^4}{c^2 d} + \frac{28(-c^2 dx^2 + d)^{7/2} x^2}{c^4 d} + \frac{8(-c^2 dx^2 + d)^{7/2}}{c^6 d} \right) b \operatorname{arccosh}(cx)$$

$$-\frac{1}{693} \left( \frac{63(-c^2 dx^2 + d)^{7/2} x^4}{c^2 d} + \frac{28(-c^2 dx^2 + d)^{7/2} x^2}{c^4 d} + \frac{8(-c^2 dx^2 + d)^{7/2}}{c^6 d} \right) a$$

$$\frac{(19845 c^{10} \sqrt{-d} d^2 x^{11} - 61985 c^8 \sqrt{-d} d^2 x^9 + 55935 c^6 \sqrt{-d} d^2 x^7 - 2079 c^4 \sqrt{-d} d^2 x^5 - 4620 c^2 \sqrt{-d} d^2 x^3 - 27720 \sqrt{-d} d^2 x) b}{2401245 c^5}$$

```
input integrate(x^5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")
```

```
output -1/693*(63*(-c^2*d*x^2 + d)^(7/2)*x^4/(c^2*d) + 28*(-c^2*d*x^2 + d)^(7/2)*x^2/(c^4*d) + 8*(-c^2*d*x^2 + d)^(7/2)/(c^6*d))*b*arccosh(c*x) - 1/693*(63*(-c^2*d*x^2 + d)^(7/2)*x^4/(c^2*d) + 28*(-c^2*d*x^2 + d)^(7/2)*x^2/(c^4*d) + 8*(-c^2*d*x^2 + d)^(7/2)/(c^6*d))*a - 1/2401245*(19845*c^10*sqrt(-d)*d^2*x^11 - 61985*c^8*sqrt(-d)*d^2*x^9 + 55935*c^6*sqrt(-d)*d^2*x^7 - 2079*c^4*sqrt(-d)*d^2*x^5 - 4620*c^2*sqrt(-d)*d^2*x^3 - 27720*sqrt(-d)*d^2*x)*b/c^5
```

**3.97.8 Giac [F(-2)]**

Exception generated.

$$\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

```
input integrate(x^5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.97.9 Mupad [F(-1)]**

Timed out.

$$\int x^5 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int x^5 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} dx$$

input `int(x^5*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2),x)`output `int(x^5*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2), x)`

### 3.98 $\int x^3(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

3.98.1	Optimal result	941
3.98.2	Mathematica [A] (verified)	942
3.98.3	Rubi [A] (verified)	942
3.98.4	Maple [B] (verified)	944
3.98.5	Fricas [A] (verification not implemented)	945
3.98.6	Sympy [F(-1)]	945
3.98.7	Maxima [A] (verification not implemented)	946
3.98.8	Giac [F(-2)]	946
3.98.9	Mupad [F(-1)]	947

#### 3.98.1 Optimal result

Integrand size = 27, antiderivative size = 298

$$\begin{aligned} \int x^3(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = & \frac{2bd^2x\sqrt{d - c^2dx^2}}{63c^3\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & + \frac{bd^2x^3\sqrt{d - c^2dx^2}}{189c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcd^2x^5\sqrt{d - c^2dx^2}}{21\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & + \frac{19bc^3d^2x^7\sqrt{d - c^2dx^2}}{441\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bc^5d^2x^9\sqrt{d - c^2dx^2}}{81\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & - \frac{(d - c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4d} + \frac{(d - c^2dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^4d^2} \end{aligned}$$

output 
$$\begin{aligned} & -1/7*(-c^2*d*x^2+d)^{(7/2)}*(a+b*\operatorname{arccosh}(c*x))/c^4/d+1/9*(-c^2*d*x^2+d)^{(9/2)} \\ & *(a+b*\operatorname{arccosh}(c*x))/c^4/d^2+2/63*b*d^2*x*(-c^2*d*x^2+d)^{(1/2)}/c^3/(c*x-1) \\ & ^{(1/2)}/(c*x+1)^{(1/2)}+1/189*b*d^2*x^3*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/ \\ & (c*x+1)^{(1/2)}-1/21*b*c*d^2*x^5*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \\ & +19/441*b*c^3*d^2*x^7*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \\ & -1/81*b*c^5*d^2*x^9*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \end{aligned}$$

**3.98.2 Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.49

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d^2 \sqrt{d - c^2 dx^2} (bc(126x + 21c^2 x^3 - 189c^4 x^5 + 171c^6 x^7 - 49c^8 x^9) + 126(-1 + cx)^{7/2}(1 + cx)^{7/2})}{3969c^4 \sqrt{-1 + cx}}$$

input `Integrate[x^3*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`output `(d^2*sqrt[d - c^2*d*x^2]*(b*c*(126*x + 21*c^2*x^3 - 189*c^4*x^5 + 171*c^6*x^7 - 49*c^8*x^9) + 126*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(a + b*ArcCosh[c*x]) + 441*c^2*x^2*(-1 + c*x)^(7/2)*(1 + c*x)^(7/2)*(a + b*ArcCosh[c*x]))) / (3969*c^4*sqrt[-1 + c*x]*sqrt[1 + c*x])`**3.98.3 Rubi [A] (verified)**Time = 0.46 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.51, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 290, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6337} \\ & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{d^2(1 - c^2 x^2)^3 (7c^2 x^2 + 2)}{63c^4} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^4 d^2} - \\ & \quad \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4 d} \\ & \quad \downarrow \text{27} \\ & \frac{bd^2 \sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^3 (7c^2 x^2 + 2) dx}{63c^3 \sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{9/2} (a + \operatorname{barccosh}(cx))}{9c^4 d^2} - \\ & \quad \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^4 d} \\ & \quad \downarrow \text{290} \end{aligned}$$

---

3.98.  $\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

$$\frac{bd^2\sqrt{d-c^2dx^2} \int (-7c^8x^8 + 19c^6x^6 - 15c^4x^4 + c^2x^2 + 2) dx}{63c^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{(d-c^2dx^2)^{9/2}(a+\operatorname{barccosh}(cx))}{9c^4d^2} - \frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{7c^4d}$$

↓ 2009

$$\frac{(d-c^2dx^2)^{9/2}(a+\operatorname{barccosh}(cx))}{9c^4d^2} - \frac{(d-c^2dx^2)^{7/2}(a+\operatorname{barccosh}(cx))}{7c^4d} + \frac{bd^2\left(-\frac{7}{9}c^8x^9 + \frac{19c^6x^7}{7} - 3c^4x^5 + \frac{c^2x^3}{3} + 2x\right)\sqrt{d-c^2dx^2}}{63c^3\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[x^3*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `(b*d^2*sqrt[d - c^2*d*x^2]*(2*x + (c^2*x^3)/3 - 3*c^4*x^5 + (19*c^6*x^7)/7 - (7*c^8*x^9)/9))/(63*c^3*sqrt[-1 + c*x]*sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x]))/(7*c^4*d) + ((d - c^2*d*x^2)^(9/2)*(a + b*ArcCosh[c*x]))/(9*c^4*d^2)`

### 3.98.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 290 `Int[((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[p, 0] && IGtQ[q, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`



### 3.98.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1101 vs.  $2(250) = 500$ .

Time = 0.60 (sec) , antiderivative size = 1102, normalized size of antiderivative = 3.70

method	result	size
default	Expression too large to display	1102
parts	Expression too large to display	1102

```
input int(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(-1/9*x^2*(-c^2*d*x^2+d)^(7/2)/c^2/d-2/63/d/c^4*(-c^2*d*x^2+d)^(7/2))+b*
(1/41472*(-d*(c^2*x^2-1))^(1/2)*(256*c^10*x^10-704*c^8*x^8+256*(c*x+1)^(1/
2)*(c*x-1)^(1/2)*x^9*c^9+688*c^6*x^6-576*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c
^7-280*c^4*x^4+432*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+41*c^2*x^2-120*(c*x
-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+9*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+9
*arccosh(c*x))*d^2/(c*x+1)/c^4/(c*x-1)-3/25088*(-d*(c^2*x^2-1))^(1/2)*(64*
c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*x^4-112
*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-25*c^2*x^2+56*(c*x-1)^(1/2)*(c*x+1)^(
1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+7*arccosh(c*x))*d^2/
(c*x+1)/c^4/(c*x-1)+1/576*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c
*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1
+3*arccosh(c*x))*d^2/(c*x+1)/c^4/(c*x-1)-3/256*(-d*(c^2*x^2-1))^(1/2)*((c*
x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+arccosh(c*x))*d^2/(c*x+1)/c^4/
(c*x-1)-3/256*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2
*x^2-1)*(1+arccosh(c*x))*d^2/(c*x+1)/c^4/(c*x-1)+1/576*(-d*(c^2*x^2-1))^(1
/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x
+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x))*d^2/(c*x+1)/c^4/(c*x-1)-3/25
088*(-d*(c^2*x^2-1))^(1/2)*(-64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+64*c^8
*x^8+112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-144*c^6*x^6-56*(c*x-1)^(1/2)*
(c*x+1)^(1/2)*c^3*x^3+104*c^4*x^4+7*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-25*...
```

**3.98.5 Fracas [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 281, normalized size of antiderivative = 0.94

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{63 (7 bc^{10} d^2 x^{10} - 26 bc^8 d^2 x^8 + 34 bc^6 d^2 x^6 - 16 bc^4 d^2 x^4 - bc^2 d^2 x^2 + 2 bd^2) \sqrt{-c^2 dx^2 + d}}{c^6 x^2 - c^4}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fracas")
```

```
output 1/3969*(63*(7*b*c^10*d^2*x^10 - 26*b*c^8*d^2*x^8 + 34*b*c^6*d^2*x^6 - 16*b*c^4*d^2*x^4 - b*c^2*d^2*x^2 + 2*b*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (49*b*c^9*d^2*x^9 - 171*b*c^7*d^2*x^7 + 189*b*c^5*d^2*x^5 - 21*b*c^3*d^2*x^3 - 126*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 63*(7*a*c^10*d^2*x^10 - 26*a*c^8*d^2*x^8 + 34*a*c^6*d^2*x^6 - 16*a*c^4*d^2*x^4 - a*c^2*d^2*x^2 + 2*a*d^2)*sqrt(-c^2*d*x^2 + d))/(c^6*x^2 - c^4)
```

**3.98.6 Sympy [F(-1)]**

Timed out.

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

```
input integrate(x**3*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)
```

```
output Timed out
```

**3.98.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 185, normalized size of antiderivative = 0.62

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$-\frac{1}{63} \left( \frac{7(-c^2 dx^2 + d)^{7/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{7/2}}{c^4 d} \right) b \operatorname{arcosh}(cx)$$

$$-\frac{1}{63} \left( \frac{7(-c^2 dx^2 + d)^{7/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{7/2}}{c^4 d} \right) a$$

$$-\frac{(49 c^8 \sqrt{-dd^2} x^9 - 171 c^6 \sqrt{-dd^2} x^7 + 189 c^4 \sqrt{-dd^2} x^5 - 21 c^2 \sqrt{-dd^2} x^3 - 126 \sqrt{-dd^2} x) b}{3969 c^3}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")
```

```
output -1/63*(7*(-c^2*d*x^2 + d)^(7/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(7/2)/(c^4*d))*b*arccosh(c*x) - 1/63*(7*(-c^2*d*x^2 + d)^(7/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(7/2)/(c^4*d))*a - 1/3969*(49*c^8*sqrt(-d)*d^2*x^9 - 171*c^6*sqrt(-d)*d^2*x^7 + 189*c^4*sqrt(-d)*d^2*x^5 - 21*c^2*sqrt(-d)*d^2*x^3 - 126*sqrt(-d)*d^2*x)*b/c^3
```

**3.98.8 Giac [F(-2)]**

Exception generated.

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.98.9 Mupad [F(-1)]**

Timed out.

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int x^3 (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} dx$$

input `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2),x)`output `int(x^3*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2), x)`

### 3.99 $\int x(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx)) dx$

3.99.1	Optimal result . . . . .	948
3.99.2	Mathematica [A] (verified) . . . . .	948
3.99.3	Rubi [A] (verified) . . . . .	949
3.99.4	Maple [B] (verified) . . . . .	950
3.99.5	Fricas [A] (verification not implemented) . . . . .	951
3.99.6	Sympy [F(-1)] . . . . .	952
3.99.7	Maxima [A] (verification not implemented) . . . . .	952
3.99.8	Giac [F(-2)] . . . . .	952
3.99.9	Mupad [F(-1)] . . . . .	953

#### 3.99.1 Optimal result

Integrand size = 25, antiderivative size = 218

$$\int x(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx)) dx = \frac{bd^2 x \sqrt{d - c^2 dx^2}}{7c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcd^2 x^3 \sqrt{d - c^2 dx^2}}{7\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{3bc^3 d^2 x^5 \sqrt{d - c^2 dx^2}}{35\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bc^5 d^2 x^7 \sqrt{d - c^2 dx^2}}{49\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{7/2} (a + \text{barccosh}(cx))}{7c^2 d}$$

output

```
-1/7*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))/c^2/d+1/7*b*d^2*x*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/7*b*c*d^2*x^3*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+3/35*b*c^3*d^2*x^5*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/49*b*c^5*d^2*x^7*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

#### 3.99.2 Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.54

$$\int x(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx)) dx = \frac{d^2 \sqrt{d - c^2 dx^2} (35a(-1 + c^2 x^2)^4 + bcx\sqrt{-1 + cx}\sqrt{1 + cx}(35 - 35c^2 x^2 + 21c^4 x^4 - 5c^6 x^6))}{245c^2 (-1 + c^2 x^2)}$$

input

```
Integrate[x*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]
```

output  $(d^2 \sqrt{d - c^2 dx^2} (35a(-1 + c^2 x^2)^4 + b c x \sqrt{-1 + c x} \sqrt{1 + c x} (35 - 35c^2 x^2 + 21c^4 x^4 - 5c^6 x^6) + 35b(-1 + c^2 x^2)^4 \operatorname{ArcCosh}[c x])) / (245c^2(-1 + c^2 x^2))$

### 3.99.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.50, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6329, 25, 39, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6329$$

$$-\frac{bd^2 \sqrt{d - c^2 dx^2} \int -(1 - cx)^3 (cx + 1)^3 dx}{7c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^2 d}$$

$$\downarrow 25$$

$$\frac{bd^2 \sqrt{d - c^2 dx^2} \int (1 - cx)^3 (cx + 1)^3 dx}{7c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^2 d}$$

$$\downarrow 39$$

$$\frac{bd^2 \sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^3 dx}{7c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^2 d}$$

$$\downarrow 210$$

$$\frac{bd^2 \sqrt{d - c^2 dx^2} \int (-c^6 x^6 + 3c^4 x^4 - 3c^2 x^2 + 1) dx}{7c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^2 d}$$

$$\downarrow 2009$$

$$\frac{bd^2 \left( -\frac{1}{7}c^6 x^7 + \frac{3c^4 x^5}{5} - c^2 x^3 + x \right) \sqrt{d - c^2 dx^2}}{7c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))}{7c^2 d}$$

input  $\operatorname{Int}[x*(d - c^2*d*x^2)^(5/2)*(a + b*\operatorname{ArcCosh}[c*x]),x]$

output  $(b*d^2*\text{Sqrt}[d - c^2*d*x^2]*(x - c^2*x^3 + (3*c^4*x^5)/5 - (c^6*x^7)/7))/(7*c*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) - ((d - c^2*d*x^2)^{(7/2)}*(a + b*\text{ArcCosh}[c*x]))/(7*c^2*d)$

### 3.99.3.1 Defintions of rubi rules used

rule 25  $\text{Int}[-(F_x), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[F_x, x], x]$

rule 39  $\text{Int}[(a + (b \cdot x)^m) \cdot (c + (d \cdot x)^m), x\_Symbol] \rightarrow \text{Int}[(a \cdot c + b \cdot d \cdot x^2)^m, x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \ \&\& \ \text{EqQ}[b \cdot c + a \cdot d, 0] \ \&\& \ (\text{IntegerQ}[m] \ || \ (\text{GtQ}[a, 0] \ \&\& \ \text{GtQ}[c, 0]))$

rule 210  $\text{Int}[(a + (b \cdot x)^2)^p, x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b \cdot x^2)^p, x], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

rule 2009  $\text{Int}[u, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

rule 6329  $\text{Int}[(a + \text{ArcCosh}[c \cdot x]) \cdot (b \cdot x)^n \cdot (d + (e \cdot x)^2)^p, x\_Symbol] \rightarrow \text{Simp}[(d + e \cdot x^2)^{p+1} \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^n / (2 \cdot e \cdot (p+1)), x] - \text{Simp}[b \cdot (n / (2 \cdot c \cdot (p+1))) \cdot \text{Simp}[(d + e \cdot x^2)^p / ((1 + c \cdot x)^{p+1} - c \cdot x^p)], x] \ \text{Int}[(1 + c \cdot x)^{p+1/2} \cdot (-1 + c \cdot x)^{p+1/2} \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^{n-1}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, p\}, x] \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[p, -1]$

### 3.99.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 955 vs. 2(182) = 364.

Time = 0.56 (sec) , antiderivative size = 956, normalized size of antiderivative = 4.39

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{7c^2d} + b \left( \frac{\sqrt{-d(c^2x^2-1)}(64c^8x^8-144c^6x^6+64\sqrt{cx+1}\sqrt{cx-1}x^7c^7+104c^4x^4-112\sqrt{cx+1}\sqrt{cx-1}c^5x^5-25c^2x^2+5)}{6272(cx+1)c^2(cx-1)} \right)$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{7c^2d} + b \left( \frac{\sqrt{-d(c^2x^2-1)}(64c^8x^8-144c^6x^6+64\sqrt{cx+1}\sqrt{cx-1}x^7c^7+104c^4x^4-112\sqrt{cx+1}\sqrt{cx-1}c^5x^5-25c^2x^2+5)}{6272(cx+1)c^2(cx-1)} \right)$

---

3.99.  $\int x(d - c^2dx^2)^{5/2} (a + \text{barccosh}(cx)) dx$

input `int(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/7*a*(-c^2*d*x^2+d)^{(7/2)}/c^2/d+b*(1/6272*(-d*(c^2*x^2-1))^{(1/2)}*(64*c^8 \\ & *x^8-144*c^6*x^6+64*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*x^7*c^7+104*c^4*x^4-112*(c \\ & *x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^5*x^5-25*c^2*x^2+56*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)} \\ & )*c^3*x^3-7*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+1)*(-1+7*arccosh(c*x))*d^2/(c* \\ & x+1)/c^2/(c*x-1)-1/640*(-d*(c^2*x^2-1))^{(1/2)}*(16*c^6*x^6-28*c^4*x^4+16*(c \\ & *x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^5*x^5+13*c^2*x^2-20*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)} \\ & )*c^3*x^3+5*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x-1)*(-1+5*arccosh(c*x))*d^2/(c* \\ & x+1)/c^2/(c*x-1)+1/128*(-d*(c^2*x^2-1))^{(1/2)}*(4*c^4*x^4-5*c^2*x^2+4*(c*x- \\ & 1)^{(1/2)}*(c*x+1)^{(1/2)}*c^3*x^3-3*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+1)*(-1+3* \\ & arccosh(c*x))*d^2/(c*x+1)/c^2/(c*x-1)-5/128*(-d*(c^2*x^2-1))^{(1/2)}*((c*x-1) \\ & )^{(1/2)}*(c*x+1)^{(1/2)}*c*x+c^2*x^2-1)*(-1+arccosh(c*x))*d^2/(c*x+1)/c^2/(c* \\ & x-1)-5/128*(-d*(c^2*x^2-1))^{(1/2)}*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+c^2*x^ \\ & 2-1)*(1+arccosh(c*x))*d^2/(c*x+1)/c^2/(c*x-1)+1/128*(-d*(c^2*x^2-1))^{(1/2)} \\ & *(-4*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^3*x^3+4*c^4*x^4+3*(c*x-1)^{(1/2)}*(c*x+1) \\ & )^{(1/2)}*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x))*d^2/(c*x+1)/c^2/(c*x-1)-1/640*( \\ & -d*(c^2*x^2-1))^{(1/2)}*(-16*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^5*x^5+16*c^6*x^6+ \\ & 20*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^3*x^3-28*c^4*x^4-5*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)} \\ & )*c*x+13*c^2*x^2-1)*(1+5*arccosh(c*x))*d^2/(c*x+1)/c^2/(c*x-1)+1/6272* \\ & (-d*(c^2*x^2-1))^{(1/2)}*(-64*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*x^7*c^7+64*c^8*x^8 \\ & +112*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^5*x^5-144*c^6*x^6-56*(c*x-1)^{(1/2)}*(... \end{aligned}$$

### 3.99.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.11

$$\int x(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx)) dx = \frac{35(bc^8 d^2 x^8 - 4bc^6 d^2 x^6 + 6bc^4 d^2 x^4 - 4bc^2 d^2 x^2 + bd^2)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - d})}{c^4 x^2 - c^2}$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output 
$$\begin{aligned} & 1/245*(35*(b*c^8*d^2*x^8 - 4*b*c^6*d^2*x^6 + 6*b*c^4*d^2*x^4 - 4*b*c^2*d^2 \\ & *x^2 + b*d^2)*\text{sqrt}(-c^2*d*x^2 + d)*\log(c*x + \text{sqrt}(c^2*x^2 - 1)) - (5*b*c^7 \\ & *d^2*x^7 - 21*b*c^5*d^2*x^5 + 35*b*c^3*d^2*x^3 - 35*b*c*d^2*x)*\text{sqrt}(-c^2*d \\ & *x^2 + d)*\text{sqrt}(c^2*x^2 - 1) + 35*(a*c^8*d^2*x^8 - 4*a*c^6*d^2*x^6 + 6*a*c^4 \\ & *d^2*x^4 - 4*a*c^2*d^2*x^2 + a*d^2)*\text{sqrt}(-c^2*d*x^2 + d))/(c^4*x^2 - c^2) \end{aligned}$$

---

3.99.  $\int x(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx)) dx$



**3.99.6 Sympy [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)`output `Timed out`**3.99.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.54

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = -\frac{(-c^2 dx^2 + d)^{7/2} b \operatorname{arccosh}(cx)}{7 c^2 d} - \frac{(-c^2 dx^2 + d)^{7/2} a}{7 c^2 d} - \frac{(5 c^6 \sqrt{-d} d^3 x^7 - 21 c^4 \sqrt{-d} d^3 x^5 + 35 c^2 \sqrt{-d} d^3 x^3 - 35 \sqrt{-d} d^3 x) b}{245 cd}$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`output `-1/7*(-c^2*d*x^2 + d)^(7/2)*b*arccosh(c*x)/(c^2*d) - 1/7*(-c^2*d*x^2 + d)^(7/2)*a/(c^2*d) - 1/245*(5*c^6*sqrt(-d)*d^3*x^7 - 21*c^4*sqrt(-d)*d^3*x^5 + 35*c^2*sqrt(-d)*d^3*x^3 - 35*sqrt(-d)*d^3*x)*b/(c*d)`**3.99.8 Giac [F(-2)]**

Exception generated.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.99.  $\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

**3.99.9 Mupad [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} dx$$

input `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2),x)`output `int(x*(a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2), x)`

**3.100** 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x} dx$$

3.100.1 Optimal result . . . . . 954  
 3.100.2 Mathematica [A] (warning: unable to verify) . . . . . 955  
 3.100.3 Rubi [A] (verified) . . . . . 955  
 3.100.4 Maple [A] (verified) . . . . . 960  
 3.100.5 Fracas [F] . . . . . 961  
 3.100.6 Sympy [F(-1)] . . . . . 961  
 3.100.7 Maxima [F] . . . . . 962  
 3.100.8 Giac [F(-2)] . . . . . 962  
 3.100.9 Mupad [F(-1)] . . . . . 962

**3.100.1 Optimal result**

Integrand size = 27, antiderivative size = 379

$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x} dx = -\frac{23bcd^2x\sqrt{d-c^2dx^2}}{15\sqrt{-1+cx}\sqrt{1+cx}} + \frac{11bc^3d^2x^3\sqrt{d-c^2dx^2}}{45\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^5d^2x^5\sqrt{d-c^2dx^2}}{25\sqrt{-1+cx}\sqrt{1+cx}} + d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) + \frac{1}{3}d(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) + \frac{1}{5}(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx)) - \frac{2d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{ibd^2\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
1/3*d*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))+1/5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))+d^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)-23/15*b*c*d^2*x*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+11/45*b*c^3*d^2*x^3*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/25*b*c^5*d^2*x^5*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*d^2*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+I*b*d^2*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-I*b*d^2*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.100.2 Mathematica [A] (warning: unable to verify)**

Time = 2.68 (sec) , antiderivative size = 471, normalized size of antiderivative = 1.24

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx = \frac{1}{15} ad^2 \sqrt{d - c^2 dx^2} (23 - 11c^2 x^2 + 3c^4 x^4) - \frac{bd^2 \sqrt{d - c^2 dx^2} \left( 9cx + 12 \left( \frac{-1+cx}{1+cx} \right)^{3/2} (1+cx)^3 \operatorname{arccosh}(cx) - \cosh(3 \operatorname{arccosh}(cx)) \right)}{18 \sqrt{\frac{-1+cx}{1+cx}} (1+cx)} + ad^{5/2} \log(x) - ad^{5/2} \log \left( d + \sqrt{d} \sqrt{d - c^2 dx^2} \right) + \frac{bd^2 \sqrt{d - c^2 dx^2} \left( -cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) \right)}{18 \sqrt{\frac{-1+cx}{1+cx}} (1+cx)}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x,x]`

output

```
(a*d^2*Sqrt[d - c^2*d*x^2]*(23 - 11*c^2*x^2 + 3*c^4*x^4))/15 - (b*d^2*Sqrt[d - c^2*d*x^2]*(9*c*x + 12*((-1 + c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3*ArcCosh[c*x] - Cosh[3*ArcCosh[c*x]]))/(18*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + a*d^(5/2)*Log[x] - a*d^(5/2)*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (b*d^2*Sqrt[d - c^2*d*x^2]*(-(c*x) + Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + I*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*PolyLog[2, I/E^ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) - (b*d^2*Sqrt[d - c^2*d*x^2]*(25*Cosh[3*ArcCosh[c*x]] + 9*(-50*c*x + Cosh[5*ArcCosh[c*x]]) + 15*ArcCosh[c*x]*(30*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) - 5*Sinh[3*ArcCosh[c*x]] - 3*Sinh[5*ArcCosh[c*x]])))/(3600*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))
```

**3.100.3 Rubi [A] (verified)**Time = 1.56 (sec) , antiderivative size = 326, normalized size of antiderivative = 0.86, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$ , Rules used = {6345, 39, 210, 2009, 6345, 25, 39, 2009, 6341, 24, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx$$

---

3.100.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx$

$$\begin{aligned} & \downarrow 6345 \\ d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (1 - cx)^2 (cx + 1)^2 dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} + \\ & \quad \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \end{aligned}$$

$$\begin{aligned} & \downarrow 39 \\ d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^2 dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} + \\ & \quad \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \end{aligned}$$

$$\begin{aligned} & \downarrow 210 \\ d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (c^4 x^4 - 2c^2 x^2 + 1) dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} + \\ & \quad \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) \end{aligned}$$

$$\begin{aligned} & \downarrow 2009 \\ d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x} dx + \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \\ & \quad \frac{bcd^2 \left( \frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x \right) \sqrt{d - c^2 dx^2}}{5\sqrt{cx - 1}\sqrt{cx + 1}} \end{aligned}$$

$$\begin{aligned} & \downarrow 6345 \\ d \left( d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x} dx + \frac{bcd \sqrt{d - c^2 dx^2} \int -((1 - cx)(cx + 1)) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) \\ & \quad \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x \right) \sqrt{d - c^2 dx^2}}{5\sqrt{cx - 1}\sqrt{cx + 1}} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ d \left( d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x} dx - \frac{bcd \sqrt{d - c^2 dx^2} \int (1 - cx)(cx + 1) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) \right) \\ & \quad \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x \right) \sqrt{d - c^2 dx^2}}{5\sqrt{cx - 1}\sqrt{cx + 1}} \end{aligned}$$

$$\downarrow 39$$

---


$$3.100. \quad \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx$$

$$d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx - \frac{bcd\sqrt{d-c^2dx^2} \int (1-c^2x^2) dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right. \\ \left. - \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^5}{5} - \frac{2c^2x^3}{3} + x \right) \sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 2009

$$d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd \left( x - \frac{c^2x^3}{3} \right) \sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}} \right) + \\ \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^5}{5} - \frac{2c^2x^3}{3} + x \right) \sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6341

$$d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \int 1 dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right) \\ - \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^5}{5} - \frac{2c^2x^3}{3} + x \right) \sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 24

$$d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right) \\ - \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^5}{5} - \frac{2c^2x^3}{3} + x \right) \sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6362

$$d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{cx} d\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right) \\ - \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4x^5}{5} - \frac{2c^2x^3}{3} + x \right) \sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 3042

$$d \left( d \left( - \frac{\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx)) \csc(\operatorname{iarccosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx)}{\sqrt{cx - 1} \sqrt{cx + 1}} + \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) - \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x \right) \sqrt{d - c^2 dx^2}}{5 \sqrt{cx - 1} \sqrt{cx + 1}} \right) \right) \downarrow 4668$$

$$d \left( d \left( - \frac{\sqrt{d - c^2 dx^2} (-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan)}{\sqrt{cx - 1} \sqrt{cx + 1}} + \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x \right) \sqrt{d - c^2 dx^2}}{5 \sqrt{cx - 1} \sqrt{cx + 1}} \right) \right) \downarrow 2715$$

$$d \left( d \left( - \frac{\sqrt{d - c^2 dx^2} (-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) d)}{\sqrt{cx - 1} \sqrt{cx + 1}} + \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x \right) \sqrt{d - c^2 dx^2}}{5 \sqrt{cx - 1} \sqrt{cx + 1}} \right) \right) \downarrow 2838$$

$$d \left( d \left( - \frac{\sqrt{d - c^2 dx^2} (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}}{\sqrt{cx - 1} \sqrt{cx + 1}} + \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) - \frac{bcd^2 \left( \frac{c^4 x^5}{5} - \frac{2c^2 x^3}{3} + x \right) \sqrt{d - c^2 dx^2}}{5 \sqrt{cx - 1} \sqrt{cx + 1}} \right) \right)$$

input `Int(((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x,x]`

output `-1/5*(b*c*d^2*sqrt[d - c^2*d*x^2]*(x - (2*c^2*x^3)/3 + (c^4*x^5)/5))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/5 + d*(-1/3*(b*c*d*sqrt[d - c^2*d*x^2]*(x - (c^2*x^3)/3))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/3 + d*(-((b*c*x*sqrt[d - c^2*d*x^2])/(sqrt[-1 + c*x]*sqrt[1 + c*x])) + sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]) - (sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(sqrt[-1 + c*x]*sqrt[1 + c*x]))))`

---

3.100.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx$

## 3.100.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 39 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 210 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`



rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6362 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.100.4 Maple [A] (verified)

Time = 1.18 (sec) , antiderivative size = 620, normalized size of antiderivative = 1.64

method	result
default	$\frac{(-c^2dx^2+d)^{\frac{5}{2}}a}{5} + \frac{ad(-c^2dx^2+d)^{\frac{3}{2}}}{3} - ad^{\frac{5}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right) + ad^2\sqrt{-c^2dx^2+d} + \frac{ib\sqrt{-d(c^2x^2-1)}}{ar}$
parts	$\frac{(-c^2dx^2+d)^{\frac{5}{2}}a}{5} + \frac{ad(-c^2dx^2+d)^{\frac{3}{2}}}{3} - ad^{\frac{5}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right) + ad^2\sqrt{-c^2dx^2+d} + \frac{ib\sqrt{-d(c^2x^2-1)}}{ar}$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`

$$3.100. \int \frac{(d-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x} dx$$

output  $1/5*(-c^2*d*x^2+d)^{(5/2)}*a+1/3*a*d*(-c^2*d*x^2+d)^{(3/2)}-a*d^{(5/2)}*\ln((2*d+2*d^{(1/2)}*(-c^2*d*x^2+d)^{(1/2)})/x)+a*d^2*(-c^2*d*x^2+d)^{(1/2)}+I*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*\operatorname{arccosh}(c*x)*\ln(1+I*(c*x+(c*x-1))^{(1/2)}*(c*x+1)^{(1/2)}))*d^2+1/5*b*(-d*(c^2*x^2-1))^{(1/2)}*d^2/(c*x+1)/(c*x-1)*\operatorname{arccosh}(c*x)*x^6*c^6-14/15*b*(-d*(c^2*x^2-1))^{(1/2)}*d^2/(c*x+1)/(c*x-1)*\operatorname{arccosh}(c*x)*x^4*c^4+34/15*b*(-d*(c^2*x^2-1))^{(1/2)}*d^2/(c*x+1)/(c*x-1)*\operatorname{arccosh}(c*x)*x^2*c^2-I*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*\operatorname{arccosh}(c*x)*\ln(1-I*(c*x+(c*x-1))^{(1/2)}*(c*x+1)^{(1/2)}))*d^2-23/15*b*(-d*(c^2*x^2-1))^{(1/2)}*d^2/(c*x+1)/(c*x-1)^{(1/2)}/(c*x-1)^{(1/2)}*c^5*x^5+11/45*b*(-d*(c^2*x^2-1))^{(1/2)}*d^2/(c*x+1)^{(1/2)}/(c*x-1)^{(1/2)}*c^3*x^3-23/15*b*(-d*(c^2*x^2-1))^{(1/2)}*d^2/(c*x+1)^{(1/2)}/(c*x-1)^{(1/2)}*c*x+I*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*\operatorname{dilog}(1+I*(c*x+(c*x-1))^{(1/2)}*(c*x+1)^{(1/2)}))*d^2-I*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*\operatorname{dilog}(1-I*(c*x+(c*x-1))^{(1/2)}*(c*x+1)^{(1/2)}))*d^2$

### 3.100.5 Fracas [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x,x, algorithm="fricas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x, x)`

### 3.100.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x,x)`

output `Timed out`

---

3.100.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx$

**3.100.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `-1/15*(15*d^(5/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - 3*(-c^2*d*x^2 + d)^(5/2) - 5*(-c^2*d*x^2 + d)^(3/2)*d - 15*sqrt(-c^2*d*x^2 + d)*d^2)*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

**3.100.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.100.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x, x)`

---

3.100.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x} dx$

**3.101** 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^3} dx$$

3.101.1 Optimal result . . . . .	963
3.101.2 Mathematica [A] (warning: unable to verify) . . . . .	964
3.101.3 Rubi [A] (verified) . . . . .	965
3.101.4 Maple [A] (verified) . . . . .	970
3.101.5 Fricas [F] . . . . .	971
3.101.6 Sympy [F(-1)] . . . . .	971
3.101.7 Maxima [F] . . . . .	972
3.101.8 Giac [F(-2)] . . . . .	972
3.101.9 Mupad [F(-1)] . . . . .	972

**3.101.1 Optimal result**

Integrand size = 27, antiderivative size = 404

$$\begin{aligned} \int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^3} dx = & -\frac{bcd^2\sqrt{d-c^2dx^2}}{2x\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{7bc^3d^2x\sqrt{d-c^2dx^2}}{3\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^5d^2x^3\sqrt{d-c^2dx^2}}{9\sqrt{-1+cx}\sqrt{1+cx}} - \frac{5}{2}c^2d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \\ & - \frac{5}{6}c^2d(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx)) - \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{2x^2} \\ & + \frac{5c^2d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & - \frac{5ibc^2d^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{5ibc^2d^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output

```

-5/6*c^2*d*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))-1/2*(-c^2*d*x^2+d)^(5/2)
)*(a+b*arccosh(c*x))/x^2-5/2*c^2*d^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)
)-1/2*b*c*d^2*(-c^2*d*x^2+d)^(1/2)/x/(c*x-1)^(1/2)/(c*x+1)^(1/2)+7/3*b*c^
3*d^2*x*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/9*b*c^5*d^2*x^3
*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5*c^2*d^2*(a+b*arccosh(c
*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(
1/2)/(c*x+1)^(1/2)-5/2*I*b*c^2*d^2*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)
)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/2*I*b*c^2*d^2
*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-
1)^(1/2)/(c*x+1)^(1/2)

```

### 3.101.2 Mathematica [A] (warning: unable to verify)

Time = 3.71 (sec) , antiderivative size = 596, normalized size of antiderivative = 1.48

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^3} dx = \frac{1}{36} d^2 \left( \frac{6a \sqrt{d - c^2 dx^2} (-3 - 14c^2 x^2 + 2c^4 x^4)}{x^2} \right.$$

$$+ \frac{bc^2 \sqrt{d - c^2 dx^2} \left( 9cx + 12 \left( \frac{-1+cx}{1+cx} \right)^{3/2} (1+cx)^3 \operatorname{arccosh}(cx) - \cosh(3 \operatorname{arccosh}(cx)) \right)}{\sqrt{\frac{-1+cx}{1+cx}} (1+cx)}$$

$$- 90ac^2 \sqrt{d} \log(x) + 90ac^2 \sqrt{d} \log \left( d + \sqrt{d} \sqrt{d - c^2 dx^2} \right) - \frac{72bc^2 \sqrt{d - c^2 dx^2} \left( -cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx \right)}{x^3}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^3,x]`

output  $(d^2*((6*a*\sqrt{d - c^2*d*x^2})*(-3 - 14*c^2*x^2 + 2*c^4*x^4))/x^2 + (b*c^2*\sqrt{d - c^2*d*x^2}*(9*c*x + 12*((-1 + c*x)/(1 + c*x))^{3/2}*(1 + c*x)^3*\text{ArcCosh}[c*x] - \text{Cosh}[3*\text{ArcCosh}[c*x]]))/(\sqrt{(-1 + c*x)/(1 + c*x)}*(1 + c*x)) - 90*a*c^2*\sqrt{d}*\text{Log}[x] + 90*a*c^2*\sqrt{d}*\text{Log}[d + \sqrt{d}*\sqrt{d - c^2*d*x^2}] - (72*b*c^2*\sqrt{d - c^2*d*x^2}*(-(c*x) + \sqrt{(-1 + c*x)/(1 + c*x)})*\text{ArcCosh}[c*x] + c*x*\sqrt{(-1 + c*x)/(1 + c*x)}*\text{ArcCosh}[c*x] + I*\text{ArcCosh}[c*x]*\text{Log}[1 - I/E^{\text{ArcCosh}[c*x]}] - I*\text{ArcCosh}[c*x]*\text{Log}[1 + I/E^{\text{ArcCosh}[c*x]}]) + I*\text{PolyLog}[2, (-I)/E^{\text{ArcCosh}[c*x]}] - I*\text{PolyLog}[2, I/E^{\text{ArcCosh}[c*x]}])))/(\sqrt{(-1 + c*x)/(1 + c*x)}*(1 + c*x)) + (18*b*d*(1 + c*x)*(c*x*\sqrt{(-1 + c*x)/(1 + c*x)} - \text{ArcCosh}[c*x] + c*x*\text{ArcCosh}[c*x] + I*c^2*x^2*\sqrt{(-1 + c*x)/(1 + c*x)}*\text{ArcCosh}[c*x]*\text{Log}[1 - I/E^{\text{ArcCosh}[c*x]}] - I*c^2*x^2*\sqrt{(-1 + c*x)/(1 + c*x)}*\text{ArcCosh}[c*x]*\text{Log}[1 + I/E^{\text{ArcCosh}[c*x]}] + I*c^2*x^2*\sqrt{(-1 + c*x)/(1 + c*x)}*\text{PolyLog}[2, (-I)/E^{\text{ArcCosh}[c*x]}] - I*c^2*x^2*\sqrt{(-1 + c*x)/(1 + c*x)}*\text{PolyLog}[2, I/E^{\text{ArcCosh}[c*x]}])))/(x^2*\sqrt{d - c^2*d*x^2}))/36$

### 3.101.3 Rubi [A] (verified)

Time = 1.63 (sec) , antiderivative size = 335, normalized size of antiderivative = 0.83, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.556$ , Rules used = {6343, 82, 244, 2009, 6345, 25, 39, 2009, 6341, 24, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{x^3} dx$$

↓ 6343

$$-\frac{5}{2}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))}{x} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{2x^2}$$

↓ 82

$$-\frac{5}{2}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))}{x} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{2x^2}$$

↓ 244

---

3.101.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{x^3} dx$

$$\begin{aligned}
& -\frac{5}{2}c^2d \int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x} dx + \frac{bcd^2\sqrt{d-c^2dx^2} \int (x^2c^4-2c^2+\frac{1}{x^2}) dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \\
& \qquad \qquad \qquad \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{2009} \\
& -\frac{5}{2}c^2d \int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{x} dx - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \\
& \qquad \qquad \qquad \frac{bcd^2\left(\frac{c^4x^3}{3}-2c^2x-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \qquad \qquad \qquad \downarrow \text{6345} \\
& -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx + \frac{bcd\sqrt{d-c^2dx^2} \int -((1-cx)(cx+1))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right. \\
& \qquad \qquad \qquad \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2\left(\frac{c^4x^3}{3}-2c^2x-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \\
& \qquad \qquad \qquad \downarrow \text{25} \\
& -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx - \frac{bcd\sqrt{d-c^2dx^2} \int (1-cx)(cx+1)dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right. \\
& \qquad \qquad \qquad \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2\left(\frac{c^4x^3}{3}-2c^2x-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \\
& \qquad \qquad \qquad \downarrow \text{39} \\
& -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx - \frac{bcd\sqrt{d-c^2dx^2} \int (1-c^2x^2) dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) \right. \\
& \qquad \qquad \qquad \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2\left(\frac{c^4x^3}{3}-2c^2x-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \\
& \qquad \qquad \qquad \downarrow \text{2009} \\
& -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x} dx + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx)) - \frac{bcd\left(x-\frac{c^2x^3}{3}\right)\sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}} \right. \\
& \qquad \qquad \qquad \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2\left(\frac{c^4x^3}{3}-2c^2x-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \\
& \qquad \qquad \qquad \downarrow \text{6341}
\end{aligned}$$

---

3.101.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{x^3} dx$

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \int 1 dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) + \frac{1}{3}(d - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2 \left( \frac{c^4x^3}{3} - 2c^2x - \frac{1}{x} \right) \sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 24

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{3}(d - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2 \left( \frac{c^4x^3}{3} - 2c^2x - \frac{1}{x} \right) \sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6362

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{3}(d - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2 \left( \frac{c^4x^3}{3} - 2c^2x - \frac{1}{x} \right) \sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 3042

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \csc \left( i\operatorname{arccosh}(cx) + \frac{\pi}{2} \right) \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) + \frac{1}{3}(d - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2 \left( \frac{c^4x^3}{3} - 2c^2x - \frac{1}{x} \right) \sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 4668

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} (-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2a)}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{3}(d - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2 \left( \frac{c^4x^3}{3} - 2c^2x - \frac{1}{x} \right) \sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 2715



$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2}(-ib \int e^{-\operatorname{arccosh}(cx)} \log(1-ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1+ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\ \left. \left. + \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2\left(\frac{c^4x^3}{3}-2c^2x-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \right. \\ \left. \downarrow 2838 \right. \\ \left. -\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2}(2\arctan(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \right. \\ \left. \left. + \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{2x^2} + \frac{bcd^2\left(\frac{c^4x^3}{3}-2c^2x-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^3,x]`

output `(b*c*d^2*Sqrt[d - c^2*d*x^2]*(-x^(-1) - 2*c^2*x + (c^4*x^3)/3))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(2*x^2) - (5*c^2*d*(-1/3*(b*c*d*Sqrt[d - c^2*d*x^2]*(x - (c^2*x^3)/3)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/3 + d*(-((b*c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/2`

### 3.101.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 39 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

- rule 82 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 244 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_)^(m_)), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`
- rule 6341 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_)^(m_)*Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1]`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^(m + 1)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6362 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.101.4 Maple [A] (verified)

Time = 1.99 (sec) , antiderivative size = 390, normalized size of antiderivative = 0.97

method	result
default	$a \left( -\frac{(-c^2 d x^2 + d)^{\frac{7}{2}}}{2 d x^2} - \frac{5 c^2 \left( \frac{(-c^2 d x^2 + d)^{\frac{5}{2}}}{5} + d \left( \frac{(-c^2 d x^2 + d)^{\frac{3}{2}}}{3} + d \left( \sqrt{-c^2 d x^2 + d} - \sqrt{d} \ln \left( \frac{2 d + 2 \sqrt{d} \sqrt{-c^2 d x^2 + d}}{x} \right) \right) \right) \right)}{2} \right) + ib$
parts	$a \left( -\frac{(-c^2 d x^2 + d)^{\frac{7}{2}}}{2 d x^2} - \frac{5 c^2 \left( \frac{(-c^2 d x^2 + d)^{\frac{5}{2}}}{5} + d \left( \frac{(-c^2 d x^2 + d)^{\frac{3}{2}}}{3} + d \left( \sqrt{-c^2 d x^2 + d} - \sqrt{d} \ln \left( \frac{2 d + 2 \sqrt{d} \sqrt{-c^2 d x^2 + d}}{x} \right) \right) \right) \right)}{2} \right) + ib$

3.101.  $\int \frac{(d - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^3} dx$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)`

output `a*(-1/2/d/x^2*(-c^2*d*x^2+d)^(7/2)-5/2*c^2*(1/5*(-c^2*d*x^2+d)^(5/2)+d*(1/3*(-c^2*d*x^2+d)^(3/2)+d*((-c^2*d*x^2+d)^(1/2)-d^(1/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)))))+1/18*I*b*(-d*(c^2*x^2-1))^(1/2)*(-6*I*arccosh(c*x)*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^4*c^4+2*I*x^5*c^5+42*I*arccosh(c*x)*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^2*c^2-42*I*x^3*c^3+45*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*arccosh(c*x)*x^2*c^2-45*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*arccosh(c*x)*x^2*c^2-45*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*x^2*c^2+45*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*x^2*c^2+9*I*arccosh(c*x)*(c*x+1)^(1/2)*(c*x-1)^(1/2)+9*I*c*x)*d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^2`

### 3.101.5 Fracas [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{x^3} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \text{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^3,x, algorithm="fracas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^3, x)`

### 3.101.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{x^3} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**3,x)`

output `Timed out`

---

3.101.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))}{x^3} dx$

**3.101.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `1/6*(15*c^2*d^(5/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - 3*(-c^2*d*x^2 + d)^(5/2)*c^2 - 5*(-c^2*d*x^2 + d)^(3/2)*c^2*d - 15*sqrt(-c^2*d*x^2 + d)*c^2*d^2 - 3*(-c^2*d*x^2 + d)^(7/2)/(d*x^2))*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^3, x)`

**3.101.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.101.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^3,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^3, x)`

---

3.101.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^3} dx$

**3.102**  $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$

3.102.1 Optimal result . . . . . 973  
 3.102.2 Mathematica [A] (warning: unable to verify) . . . . . 974  
 3.102.3 Rubi [A] (verified) . . . . . 975  
 3.102.4 Maple [A] (verified) . . . . . 980  
 3.102.5 Fricas [F] . . . . . 981  
 3.102.6 Sympy [F(-1)] . . . . . 981  
 3.102.7 Maxima [F] . . . . . 981  
 3.102.8 Giac [F(-2)] . . . . . 982  
 3.102.9 Mupad [F(-1)] . . . . . 982

**3.102.1 Optimal result**

Integrand size = 27, antiderivative size = 407

$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^5} dx = -\frac{bcd^2\sqrt{d-c^2dx^2}}{12x^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{9bc^3d^2\sqrt{d-c^2dx^2}}{8x\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^5d^2x\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{15}{8}c^4d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))$$

$$+ \frac{5c^2d(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))}{8x^2} - \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{4x^4}$$

$$- \frac{15c^4d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\arctan(e^{\operatorname{arccosh}(cx)})}{4\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{15ibc^4d^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})}{8\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{15ibc^4d^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(cx)})}{8\sqrt{-1+cx}\sqrt{1+cx}}$$

output 
$$\frac{5}{8}c^2d(-c^2dx^2+d)^{3/2}(a+b\operatorname{arccosh}(cx))/x^2-1/4(-c^2dx^2+d)^{5/2}(a+b\operatorname{arccosh}(cx))/x^4+15/8c^4d^2(a+b\operatorname{arccosh}(cx))*(-c^2dx^2+d)^{1/2}-1/12b*c*d^2*(-c^2dx^2+d)^{1/2}/x^3/(c*x-1)^{1/2}/(c*x+1)^{1/2}+9/8b*c^3*d^2*(-c^2dx^2+d)^{1/2}/x/(c*x-1)^{1/2}/(c*x+1)^{1/2}-b*c^5*d^2*x*(-c^2dx^2+d)^{1/2}/(c*x-1)^{1/2}/(c*x+1)^{1/2}-15/4c^4d^2(a+b\operatorname{arccosh}(cx))*\arctan(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2})*(-c^2dx^2+d)^{1/2}/(c*x-1)^{1/2}/(c*x+1)^{1/2}+15/8I*b*c^4d^2\operatorname{polylog}(2,-I*(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*(-c^2dx^2+d)^{1/2}/(c*x-1)^{1/2}/(c*x+1)^{1/2}-15/8I*b*c^4d^2\operatorname{polylog}(2,I*(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*(-c^2dx^2+d)^{1/2}/(c*x-1)^{1/2}/(c*x+1)^{1/2}$$

### 3.102.2 Mathematica [A] (warning: unable to verify)

Time = 1.47 (sec) , antiderivative size = 660, normalized size of antiderivative = 1.62

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))}{x^5} dx = \frac{-2bcd^3x + 2bc^2d^3x^2 + 27bc^3d^3x^3 - 27bc^4d^3x^4 - 24bc^5d^3x^5 + 24b^2cd^3x^6}{x^5}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^5,x]`

output 
$$\begin{aligned} & (-2*b*c*d^3*x + 2*b*c^2*d^3*x^2 + 27*b*c^3*d^3*x^3 - 27*b*c^4*d^3*x^4 - 24 \\ & *b*c^5*d^3*x^5 + 24*b*c^6*d^3*x^6 - 6*a*d^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] + 3 \\ & 3*a*c^2*d^3*x^2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] - 3*a*c^4*d^3*x^4*\operatorname{Sqrt}[(-1 + c* \\ & x)/(1 + c*x)] - 24*a*c^6*d^3*x^6*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] - 6*b*d^3*\operatorname{Sqrt} \\ & [(-1 + c*x)/(1 + c*x)]*\operatorname{ArcCosh}[c*x] + 33*b*c^2*d^3*x^2*\operatorname{Sqrt}[(-1 + c*x)/(1 \\ & + c*x)]*\operatorname{ArcCosh}[c*x] - 3*b*c^4*d^3*x^4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*\operatorname{ArcCosh}[ \\ & c*x] - 24*b*c^6*d^3*x^6*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*\operatorname{ArcCosh}[c*x] + (45*I)*b \\ & *c^4*d^3*x^4*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - I/E^{\operatorname{ArcCosh}[c*x]}] - (45*I)*b*c^5*d^3*x^5 \\ & *\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - I/E^{\operatorname{ArcCosh}[c*x]}] - (45*I)*b*c^4*d^3*x^4*\operatorname{ArcCosh}[c*x \\ & ]*\operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[c*x]}] + (45*I)*b*c^5*d^3*x^5*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + I/ \\ & E^{\operatorname{ArcCosh}[c*x]}] + 45*a*c^4*d^{5/2}*x^4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*\operatorname{Sqrt}[d - \\ & c^2*d*x^2]*\operatorname{Log}[x] - 45*a*c^4*d^{5/2}*x^4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*\operatorname{Sqrt}[ \\ & d - c^2*d*x^2]*\operatorname{Log}[d + \operatorname{Sqrt}[d]*\operatorname{Sqrt}[d - c^2*d*x^2]] - (45*I)*b*c^4*d^3*x^4 \\ & *(-1 + c*x)*\operatorname{PolyLog}[2, (-I)/E^{\operatorname{ArcCosh}[c*x]}] + (45*I)*b*c^4*d^3*x^4*(-1 + c \\ & *x)*\operatorname{PolyLog}[2, I/E^{\operatorname{ArcCosh}[c*x]}] / (24*x^4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*\operatorname{Sqrt}[ \\ & d - c^2*d*x^2]) \end{aligned}$$

### 3.102.3 Rubi [A] (verified)

Time = 1.72 (sec) , antiderivative size = 343, normalized size of antiderivative = 0.84, number of steps used = 17, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.593$ , Rules used = {6343, 82, 244, 2009, 6343, 25, 82, 244, 2009, 6341, 24, 6362, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx \\
 & \quad \downarrow \text{6343} \\
 & -\frac{5}{4}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2}{x^4} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{4x^4} \\
 & \quad \downarrow \text{82} \\
 & -\frac{5}{4}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2}{x^4} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{4x^4} \\
 & \quad \downarrow \text{244} \\
 & -\frac{5}{4}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx + \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \left( c^4 - \frac{2c^2}{x^2} + \frac{1}{x^4} \right) dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{4x^4} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{5}{4}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{x^3} dx - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{4x^4} + \\
 & \quad \frac{bcd^2 \left( c^4 x + \frac{2c^2}{x} - \frac{1}{3x^3} \right) \sqrt{d - c^2 dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \\
 & \quad \downarrow \text{6343} \\
 & -\frac{5}{4}c^2d \left( -\frac{3}{2}c^2d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x} dx - \frac{bcd \sqrt{d - c^2 dx^2} \int -\frac{(1-cx)(cx+1)}{x^2} dx}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{2x^2} \right. \\
 & \quad \left. \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{4x^4} + \frac{bcd^2 \left( c^4 x + \frac{2c^2}{x} - \frac{1}{3x^3} \right) \sqrt{d - c^2 dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right)
 \end{aligned}$$

---

3.102.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx$



↓ 25

$$-\frac{5}{4}c^2d\left(-\frac{3}{2}c^2d\int\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x}dx+\frac{bcd\sqrt{d-c^2dx^2}\int\frac{(1-cx)(cx+1)}{x^2}dx}{2\sqrt{cx-1}\sqrt{cx+1}}-\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2}\right) \\ +\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4}+\frac{bcd^2\left(c^4x+\frac{2c^2}{x}-\frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 82

$$-\frac{5}{4}c^2d\left(-\frac{3}{2}c^2d\int\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x}dx+\frac{bcd\sqrt{d-c^2dx^2}\int\frac{1-c^2x^2}{x^2}dx}{2\sqrt{cx-1}\sqrt{cx+1}}-\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2}\right) \\ +\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4}+\frac{bcd^2\left(c^4x+\frac{2c^2}{x}-\frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 244

$$-\frac{5}{4}c^2d\left(-\frac{3}{2}c^2d\int\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x}dx+\frac{bcd\sqrt{d-c^2dx^2}\int\left(\frac{1}{x^2}-c^2\right)dx}{2\sqrt{cx-1}\sqrt{cx+1}}-\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2}\right) \\ +\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4}+\frac{bcd^2\left(c^4x+\frac{2c^2}{x}-\frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2009

$$-\frac{5}{4}c^2d\left(-\frac{3}{2}c^2d\int\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{x}dx-\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))}{2x^2}+\frac{bcd\left(c^2(-x)-\frac{1}{x}\right)\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}}\right) \\ +\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4}+\frac{bcd^2\left(c^4x+\frac{2c^2}{x}-\frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6341

$$-\frac{5}{4}c^2d\left(-\frac{3}{2}c^2d\left(-\frac{\sqrt{d-c^2dx^2}\int\frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}}dx}{\sqrt{cx-1}\sqrt{cx+1}}-\frac{bc\sqrt{d-c^2dx^2}\int 1dx}{\sqrt{cx-1}\sqrt{cx+1}}+\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))\right)\right) \\ +\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4}+\frac{bcd^2\left(c^4x+\frac{2c^2}{x}-\frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 24

$$-\frac{5}{4}c^2d \left( -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd^2 \left( c^4x + \frac{2c^2}{x} - \frac{1}{3x^3} \right) \sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \quad \downarrow \quad \mathbf{6362}$$

$$-\frac{5}{4}c^2d \left( -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd^2 \left( c^4x + \frac{2c^2}{x} - \frac{1}{3x^3} \right) \sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \quad \downarrow \quad \mathbf{3042}$$

$$-\frac{5}{4}c^2d \left( -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \csc \left( i\operatorname{arccosh}(cx) + \frac{\pi}{2} \right) \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd^2 \left( c^4x + \frac{2c^2}{x} - \frac{1}{3x^3} \right) \sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \quad \downarrow \quad \mathbf{4668}$$

$$-\frac{5}{4}c^2d \left( -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \left( -ib \int \log(1-ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1+ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) \right)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd^2 \left( c^4x + \frac{2c^2}{x} - \frac{1}{3x^3} \right) \sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \quad \downarrow \quad \mathbf{2715}$$

$$-\frac{5}{4}c^2d \left( -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \left( -ib \int e^{-\operatorname{arccosh}(cx)} \log(1-ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1+ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} \right)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) - \frac{bcx\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4} + \frac{bcd^2 \left( c^4x + \frac{2c^2}{x} - \frac{1}{3x^3} \right) \sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \quad \downarrow \quad \mathbf{2838}$$

$$-\frac{5}{4}c^2d\left(-\frac{3}{2}c^2d\left(-\frac{\sqrt{d-c^2dx^2}(2\arctan(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))-ib\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})+ib\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(cx)})\right)}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right. \\ \left.+\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))}{4x^4}+\frac{bcd^2\left(c^4x+\frac{2c^2}{x}-\frac{1}{3x^3}\right)\sqrt{d-c^2dx^2}}{4\sqrt{cx-1}\sqrt{cx+1}}\right)$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/x^5,x]`

output `(b*c*d^2*(-1/3*1/x^3 + (2*c^2)/x + c^4*x)*Sqrt[d - c^2*d*x^2]/(4*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(4*x^4) - (5*c^2*d*((b*c*d*(-x^(-1) - c^2*x)*Sqrt[d - c^2*d*x^2])/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(2*x^2) - (3*c^2*d*(-((b*c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]])))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/2)/4`

### 3.102.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) +
(e_.)*(x_)^2], x_Symbol] :> Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sq
rt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0]
&& IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m
+ 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(
m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x],
x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && G
tQ[p, 0] && LtQ[m, -1]`

rule 6362 `Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

### 3.102.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 691, normalized size of antiderivative = 1.70

method	result
default	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{4dx^4} + \frac{3ac^2(-c^2dx^2+d)^{\frac{7}{2}}}{8dx^2} + \frac{3ac^4(-c^2dx^2+d)^{\frac{5}{2}}}{8} + \frac{5ac^4d(-c^2dx^2+d)^{\frac{3}{2}}}{8} - \frac{15ac^4d^{\frac{5}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{8}$
parts	$-\frac{a(-c^2dx^2+d)^{\frac{7}{2}}}{4dx^4} + \frac{3ac^2(-c^2dx^2+d)^{\frac{7}{2}}}{8dx^2} + \frac{3ac^4(-c^2dx^2+d)^{\frac{5}{2}}}{8} + \frac{5ac^4d(-c^2dx^2+d)^{\frac{3}{2}}}{8} - \frac{15ac^4d^{\frac{5}{2}} \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{8}$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^5,x,method=_RETURNVERBOSE)`

output `-1/4*a/d/x^4*(-c^2*d*x^2+d)^(7/2)+3/8*a*c^2/d/x^2*(-c^2*d*x^2+d)^(7/2)+3/8*a*c^4*(-c^2*d*x^2+d)^(5/2)+5/8*a*c^4*d*(-c^2*d*x^2+d)^(3/2)-15/8*a*c^4*d^(5/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)+15/8*a*c^4*d^2*(-c^2*d*x^2+d)^(1/2)+b*(-d*(c^2*x^2-1))^(1/2)*c^6*d^2/(c*x+1)/(c*x-1)*arccosh(c*x)*x^2-b*(-d*(c^2*x^2-1))^(1/2)*c^5*d^2/(c*x+1)^(1/2)/(c*x-1)^(1/2)*x+1/8*b*(-d*(c^2*x^2-1))^(1/2)*c^4*d^2/(c*x+1)/(c*x-1)*arccosh(c*x)+9/8*b*d^2*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)^(1/2)/x/(c*x-1)^(1/2)*c^3-11/8*b*d^2*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)/x^2/(c*x-1)*arccosh(c*x)*c^2-1/12*b*d^2*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)/x^3/(c*x-1)^(1/2)*c+1/4*b*d^2*(-d*(c^2*x^2-1))^(1/2)/(c*x+1)/x^4/(c*x-1)*arccosh(c*x)+15/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^4*d^2-15/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^4*d^2-15/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^4*d^2+15/8*I*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^4*d^2`

3.102. 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))}{x^5} dx$$

**3.102.5 Fracas [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x^5} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^5,x, algorithm="fracas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^5, x)`

**3.102.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))/x**5,x)`

output `Timed out`

**3.102.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)}{x^5} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^5,x, algorithm="maxima")`

output `-1/8*(15*c^4*d^(5/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - 3*(-c^2*d*x^2 + d)^(5/2)*c^4 - 5*(-c^2*d*x^2 + d)^(3/2)*c^4*d - 15*sqrt(-c^2*d*x^2 + d)*c^4*d^2 - 3*(-c^2*d*x^2 + d)^(7/2)*c^2/(d*x^2) + 2*(-c^2*d*x^2 + d)^(7/2)/(d*x^4))*a + b*integrate((-c^2*d*x^2 + d)^(5/2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^5, x)`

---

3.102.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx$

**3.102.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))/x^5,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.102.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{x^5} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2}}{x^5} dx$$

input `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^5,x)`

output `int(((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2))/x^5, x)`

### 3.103 $\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx$

3.103.1 Optimal result . . . . .	983
3.103.2 Mathematica [A] (verified) . . . . .	983
3.103.3 Rubi [A] (verified) . . . . .	984
3.103.4 Maple [B] (verified) . . . . .	985
3.103.5 Fricas [F] . . . . .	985
3.103.6 Sympy [F] . . . . .	986
3.103.7 Maxima [F(-2)] . . . . .	986
3.103.8 Giac [F] . . . . .	986
3.103.9 Mupad [F(-1)] . . . . .	987

#### 3.103.1 Optimal result

Integrand size = 14, antiderivative size = 66

$$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx = -\frac{\sqrt{1-x^2}}{4\sqrt{-1+x}} + \frac{1}{2}x\sqrt{1-x^2} \operatorname{arccosh}(x) - \frac{\sqrt{1-x} \operatorname{arccosh}(x)^2}{4\sqrt{-1+x}}$$

output `-1/4*x^2*(1-x)^(1/2)/(-1+x)^(1/2)-1/4*arccosh(x)^2*(1-x)^(1/2)/(-1+x)^(1/2)+1/2*x*arccosh(x)*(-x^2+1)^(1/2)`

#### 3.103.2 Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.82

$$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx = \frac{-\sqrt{-((-1+x)(1+x))}(\cosh(2\operatorname{arccosh}(x)) + 2\operatorname{arccosh}(x)(\operatorname{arccosh}(x) - \sinh(2\operatorname{arccosh}(x))))}{8\sqrt{\frac{-1+x}{1+x}}(1+x)}$$

input `Integrate[Sqrt[1 - x^2]*ArcCosh[x], x]`

output `-1/8*(Sqrt[-((-1 + x)*(1 + x))]*(Cosh[2*ArcCosh[x]] + 2*ArcCosh[x]*(ArcCos h[x] - Sinh[2*ArcCosh[x]])))/(Sqrt[(-1 + x)/(1 + x)]*(1 + x))`



### 3.103.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {6310, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{1-x^2} \operatorname{arccosh}(x) dx \\
 & \quad \downarrow \text{6310} \\
 & -\frac{\sqrt{1-x} \int \frac{\operatorname{arccosh}(x)}{\sqrt{x-1}\sqrt{x+1}} dx}{2\sqrt{x-1}} - \frac{\sqrt{1-x} \int x dx}{2\sqrt{x-1}} + \frac{1}{2} x \sqrt{1-x^2} \operatorname{arccosh}(x) \\
 & \quad \downarrow \text{15} \\
 & -\frac{\sqrt{1-x} \int \frac{\operatorname{arccosh}(x)}{\sqrt{x-1}\sqrt{x+1}} dx}{2\sqrt{x-1}} + \frac{1}{2} \sqrt{1-x^2} x \operatorname{arccosh}(x) - \frac{\sqrt{1-xx^2}}{4\sqrt{x-1}} \\
 & \quad \downarrow \text{6308} \\
 & \frac{1}{2} \sqrt{1-x^2} x \operatorname{arccosh}(x) - \frac{\sqrt{1-x} \operatorname{arccosh}(x)^2}{4\sqrt{x-1}} - \frac{\sqrt{1-xx^2}}{4\sqrt{x-1}}
 \end{aligned}$$

input `Int[Sqrt[1 - x^2]*ArcCosh[x], x]`

output `-1/4*(Sqrt[1 - x]*x^2)/Sqrt[-1 + x] + (x*Sqrt[1 - x^2]*ArcCosh[x])/2 - (Sqrt[1 - x]*ArcCosh[x]^2)/(4*Sqrt[-1 + x])`

#### 3.103.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

```
rule 6310 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_
Symbol] :> Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(
1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcC
osh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sq
rt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^
(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n
, 0]
```

### 3.103.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 151 vs. 2(50) = 100.

Time = 0.94 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.30

method	result
default	$-\frac{\sqrt{-x^2+1} \operatorname{arccosh}(x)^2}{4\sqrt{x-1}\sqrt{1+x}} + \frac{\sqrt{-x^2+1} (2x^3-2x+2\sqrt{1+x}\sqrt{x-1}x^2-\sqrt{x-1}\sqrt{1+x})(-1+2 \operatorname{arccosh}(x))}{16(1+x)(x-1)} + \frac{\sqrt{-x^2+1} (-2\sqrt{1+x}\sqrt{x-1})}{16(1+x)(x-1)}$

```
input int(arccosh(x)*(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/4*(-x^2+1)^(1/2)/(x-1)^(1/2)/(1+x)^(1/2)*arccosh(x)^2+1/16*(-x^2+1)^(1/
2)*(2*x^3-2*x+2*(1+x)^(1/2)*(x-1)^(1/2)*x^2-(x-1)^(1/2)*(1+x)^(1/2))*(-1+2
*arccosh(x))/(1+x)/(x-1)+1/16*(-x^2+1)^(1/2)*(-2*(1+x)^(1/2)*(x-1)^(1/2)*x
^2+2*x^3+(x-1)^(1/2)*(1+x)^(1/2)-2*x)*(1+2*arccosh(x))/(1+x)/(x-1)
```

### 3.103.5 Fracas [F]

$$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx = \int \sqrt{-x^2+1} \operatorname{arccosh}(x) dx$$

```
input integrate(arccosh(x)*(-x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(sqrt(-x^2 + 1)*arccosh(x), x)
```

**3.103.6 Sympy [F]**

$$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx = \int \sqrt{-(x-1)(x+1)} \operatorname{acosh}(x) dx$$

input `integrate(acosh(x)*(-x**2+1)**(1/2),x)`

output `Integral(sqrt(-(x - 1)*(x + 1))*acosh(x), x)`

**3.103.7 Maxima [F(-2)]**

Exception generated.

$$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx = \text{Exception raised: RuntimeError}$$

input `integrate(arccosh(x)*(-x^2+1)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.103.8 Giac [F]**

$$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx = \int \sqrt{-x^2+1} \operatorname{arcosh}(x) dx$$

input `integrate(arccosh(x)*(-x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-x^2 + 1)*arccosh(x), x)`

**3.103.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{1-x^2} \operatorname{arccosh}(x) dx = \int \operatorname{acosh}(x) \sqrt{1-x^2} dx$$

input `int(acosh(x)*(1 - x^2)^(1/2),x)`output `int(acosh(x)*(1 - x^2)^(1/2), x)`

### 3.104 $\int \frac{x^5(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$

3.104.1 Optimal result . . . . .	988
3.104.2 Mathematica [A] (verified) . . . . .	989
3.104.3 Rubi [A] (verified) . . . . .	989
3.104.4 Maple [B] (verified) . . . . .	991
3.104.5 Fracas [A] (verification not implemented) . . . . .	992
3.104.6 Sympy [F] . . . . .	993
3.104.7 Maxima [A] (verification not implemented) . . . . .	993
3.104.8 Giac [F(-2)] . . . . .	994
3.104.9 Mupad [F(-1)] . . . . .	994

#### 3.104.1 Optimal result

Integrand size = 27, antiderivative size = 236

$$\int \frac{x^5(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = -\frac{8bx\sqrt{-1 + cx}\sqrt{1 + cx}}{15c^5\sqrt{d - c^2dx^2}} - \frac{4bx^3\sqrt{-1 + cx}\sqrt{1 + cx}}{45c^3\sqrt{d - c^2dx^2}} - \frac{bx^5\sqrt{-1 + cx}\sqrt{1 + cx}}{25c\sqrt{d - c^2dx^2}} - \frac{8\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{15c^6d} - \frac{4x^2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{15c^4d} - \frac{x^4\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{5c^2d}$$

output

```
-8/15*b*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5/(-c^2*d*x^2+d)^(1/2)-4/45*b*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/(-c^2*d*x^2+d)^(1/2)-1/25*b*x^5*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)-8/15*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^6/d-4/15*x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^4/d-1/5*x^4*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^2/d
```

**3.104.2 Mathematica [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.59

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{\sqrt{d - c^2 dx^2}(bcx\sqrt{-1 + cx}\sqrt{1 + cx}(120 + 20c^2x^2 + 9c^4x^4) - 15a(-8 + 4c^2x^2 + c^4x^4 + 3c^6x^6) - 15b(-8 + 4c^2x^2 + c^4x^4 + 3c^6x^6)*\operatorname{ArcCosh}[c*x])}{225c^6d(-1 + cx)(1 + cx)}$$

input `Integrate[(x^5*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2], x]`output `(Sqrt[d - c^2*d*x^2]*(b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(120 + 20*c^2*x^2 + 9*c^4*x^4) - 15*a*(-8 + 4*c^2*x^2 + c^4*x^4 + 3*c^6*x^6) - 15*b*(-8 + 4*c^2*x^2 + c^4*x^4 + 3*c^6*x^6)*ArcCosh[c*x]))/(225*c^6*d*(-1 + c*x)*(1 + c*x))`**3.104.3 Rubi [A] (verified)**Time = 0.72 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6353, 15, 6353, 15, 6329, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6353}$$

$$\frac{4 \int \frac{x^3(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{5c^2} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int x^4 dx}{5c\sqrt{d - c^2 dx^2}} - \frac{x^4\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{5c^2 d}$$

$$\downarrow \text{15}$$

$$\frac{4 \int \frac{x^3(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{5c^2} - \frac{x^4\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{5c^2 d} - \frac{bx^5\sqrt{cx - 1}\sqrt{cx + 1}}{25c\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{6353}$$

3.104.  $\int \frac{x^5(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$

$$\begin{aligned}
& 4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx)) dx}{\sqrt{d-c^2dx^2}}}{3c^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \int x^2 dx}{3c\sqrt{d-c^2dx^2}} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3c^2d} \right) \\
& \frac{5c^2}{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{bx^5\sqrt{cx-1}\sqrt{cx+1}}{25c\sqrt{d-c^2dx^2}}} \\
& \quad \downarrow 15 \\
& 4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx)) dx}{\sqrt{d-c^2dx^2}}}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3c^2d} - \frac{bx^3\sqrt{cx-1}\sqrt{cx+1}}{9c\sqrt{d-c^2dx^2}} \right) \\
& \frac{5c^2}{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{bx^5\sqrt{cx-1}\sqrt{cx+1}}{25c\sqrt{d-c^2dx^2}}} \\
& \quad \downarrow 6329 \\
& 4 \left( \frac{2 \left( -\frac{b\sqrt{cx-1}\sqrt{cx+1} \int 1 dx}{c\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{c^2d} \right)}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3c^2d} - \frac{bx^3\sqrt{cx-1}\sqrt{cx+1}}{9c\sqrt{d-c^2dx^2}} \right) \\
& \frac{5c^2}{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) - \frac{bx^5\sqrt{cx-1}\sqrt{cx+1}}{25c\sqrt{d-c^2dx^2}}} \\
& \quad \downarrow 24 \\
& \frac{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{5c^2d} + \\
& 4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3c^2d} + \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{c^2d} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c\sqrt{d-c^2dx^2}} \right)}{3c^2} - \frac{bx^3\sqrt{cx-1}\sqrt{cx+1}}{9c\sqrt{d-c^2dx^2}} \right) \\
& \frac{5c^2}{bx^5\sqrt{cx-1}\sqrt{cx+1}} \\
& \frac{5c^2}{25c\sqrt{d-c^2dx^2}}
\end{aligned}$$

input `Int[(x^5*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2], x]`

output `-1/25*(b*x^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*Sqrt[d - c^2*d*x^2]) - (x^4*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(5*c^2*d) + (4*(-1/9*(b*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*Sqrt[d - c^2*d*x^2]) - (x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(3*c^2*d) + (2*(-((b*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[d - c^2*d*x^2])) - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(c^2*d)))/(3*c^2)))/(5*c^2)`

## 3.104.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`
- rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

## 3.104.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 669 vs.  $2(200) = 400$ .

Time = 0.79 (sec) , antiderivative size = 670, normalized size of antiderivative = 2.84

method	result
default	$a \left( -\frac{x^4 \sqrt{-c^2 d x^2 + d}}{5c^2 d} + \frac{-4x^2 \sqrt{-c^2 d x^2 + d} - 8 \sqrt{-c^2 d x^2 + d}}{15c^2 d c^2} \right) + b \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (16c^6 x^6 - 28c^4 x^4 + 16\sqrt{cx+1} \sqrt{cx-1} c^5 x^5 + \dots}{\dots} \right)$
parts	$a \left( -\frac{x^4 \sqrt{-c^2 d x^2 + d}}{5c^2 d} + \frac{-4x^2 \sqrt{-c^2 d x^2 + d} - 8 \sqrt{-c^2 d x^2 + d}}{15c^2 d c^2} \right) + b \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (16c^6 x^6 - 28c^4 x^4 + 16\sqrt{cx+1} \sqrt{cx-1} c^5 x^5 + \dots}{\dots} \right)$

input `int(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`



output

```

a*(-1/5*x^4/c^2/d*(-c^2*d*x^2+d)^(1/2)+4/5/c^2*(-1/3*x^2/c^2/d*(-c^2*d*x^2+d)^(1/2)-2/3/d/c^4*(-c^2*d*x^2+d)^(1/2)))+b*(-1/800*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c*x))/c^6/d/(c^2*x^2-1)-5/288*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))/c^6/d/(c^2*x^2-1)-5/16*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+arccosh(c*x))/c^6/d/(c^2*x^2-1)-5/16*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))/c^6/d/(c^2*x^2-1)-5/288*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x))/c^6/d/(c^2*x^2-1)-1/800*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-28*c^4*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(1+5*arccosh(c*x))/c^6/d/(c^2*x^2-1)

```

### 3.104.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.75

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \frac{15(3bc^6x^6 + bc^4x^4 + 4bc^2x^2 - 8b)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) - (9bc^5x^5 + 20bc^3x^3 + 120bcx)}{225(c^8dx^2 - c^6d)}$$

input `integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output

```

-1/225*(15*(3*b*c^6*x^6 + b*c^4*x^4 + 4*b*c^2*x^2 - 8*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (9*b*c^5*x^5 + 20*b*c^3*x^3 + 120*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 15*(3*a*c^6*x^6 + a*c^4*x^4 + 4*a*c^2*x^2 - 8*a)*sqrt(-c^2*d*x^2 + d))/(c^8*d*x^2 - c^6*d)

```

## 3.104.6 Sympy [F]

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate(x**5*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**5*(a + b*acosh(c*x))/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

## 3.104.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.83

$$\begin{aligned} & \int \frac{x^5(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx \\ &= -\frac{1}{15} \left( \frac{3\sqrt{-c^2 dx^2 + dx^4}}{c^2 d} + \frac{4\sqrt{-c^2 dx^2 + dx^2}}{c^4 d} + \frac{8\sqrt{-c^2 dx^2 + d}}{c^6 d} \right) b \operatorname{arcosh}(cx) \\ & \quad - \frac{1}{15} \left( \frac{3\sqrt{-c^2 dx^2 + dx^4}}{c^2 d} + \frac{4\sqrt{-c^2 dx^2 + dx^2}}{c^4 d} + \frac{8\sqrt{-c^2 dx^2 + d}}{c^6 d} \right) a \\ & \quad + \frac{(9c^4\sqrt{-dx^5} + 20c^2\sqrt{-dx^3} + 120\sqrt{-dx})b}{225c^5d} \end{aligned}$$

input `integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/15*(3*sqrt(-c^2*d*x^2 + d)*x^4/(c^2*d) + 4*sqrt(-c^2*d*x^2 + d)*x^2/(c^4*d) + 8*sqrt(-c^2*d*x^2 + d)/(c^6*d))*b*arccosh(c*x) - 1/15*(3*sqrt(-c^2*d*x^2 + d)*x^4/(c^2*d) + 4*sqrt(-c^2*d*x^2 + d)*x^2/(c^4*d) + 8*sqrt(-c^2*d*x^2 + d)/(c^6*d))*a + 1/225*(9*c^4*sqrt(-d)*x^5 + 20*c^2*sqrt(-d)*x^3 + 120*sqrt(-d)*x)*b/(c^5*d)`

**3.104.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.104.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^5*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^5*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2), x)`

### 3.105 $\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$

3.105.1 Optimal result . . . . .	995
3.105.2 Mathematica [A] (verified) . . . . .	996
3.105.3 Rubi [A] (verified) . . . . .	996
3.105.4 Maple [B] (verified) . . . . .	998
3.105.5 Fricas [F] . . . . .	999
3.105.6 Sympy [F] . . . . .	999
3.105.7 Maxima [F] . . . . .	999
3.105.8 Giac [F] . . . . .	1000
3.105.9 Mupad [F(-1)] . . . . .	1000

#### 3.105.1 Optimal result

Integrand size = 27, antiderivative size = 212

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = -\frac{3bx^2\sqrt{-1 + cx}\sqrt{1 + cx}}{16c^3\sqrt{d - c^2dx^2}} - \frac{bx^4\sqrt{-1 + cx}\sqrt{1 + cx}}{16c\sqrt{d - c^2dx^2}} - \frac{3x\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{8c^4d} - \frac{x^3\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{4c^2d} + \frac{3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{16bc^5\sqrt{d - c^2dx^2}}$$

output

```
-3/16*b*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/(-c^2*d*x^2+d)^(1/2)-1/16*b*x^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)+3/16*(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^5/(-c^2*d*x^2+d)^(1/2)-3/8*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^4/d-1/4*x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^2/d
```

### 3.105.2 Mathematica [A] (verified)

Time = 0.94 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.81

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{-\frac{16acx(3+2c^2x^2)\sqrt{d-c^2dx^2}}{d} - \frac{48a \arctan\left(\frac{cx\sqrt{d-c^2dx^2}}{\sqrt{d}(-1+c^2x^2)}\right)}{\sqrt{d}} + \frac{b\sqrt{\frac{-1+cx}{1+cx}}(1+cx)(-16 \cosh(2\operatorname{arccosh}(cx)) - \cosh(4\operatorname{arccosh}(cx)) + 4\operatorname{arccosh}(cx))}{\sqrt{d}}}{128c^5}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2], x]`

output `((-16*a*c*x*(3 + 2*c^2*x^2)*Sqrt[d - c^2*d*x^2])/d - (48*a*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))])/Sqrt[d] + (b*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(-16*Cosh[2*ArcCosh[c*x]] - Cosh[4*ArcCosh[c*x]] + 4*ArcCosh[c*x]*(6*ArcCosh[c*x] + 8*Sinh[2*ArcCosh[c*x]] + Sinh[4*ArcCosh[c*x]])))/Sqrt[d - c^2*d*x^2])/(128*c^5)`

### 3.105.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6353, 15, 6353, 15, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6353}$$

$$\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{4c^2} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int x^3 dx}{4c\sqrt{d - c^2 dx^2}} - \frac{x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{4c^2 d}$$

$$\downarrow \text{15}$$

$$\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{4c^2} - \frac{x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{4c^2 d} - \frac{bx^4\sqrt{cx - 1}\sqrt{cx + 1}}{16c\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{6353}$$

---

3.105.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$

$$\begin{aligned}
& 3 \left( \frac{\int \frac{a + \operatorname{arccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx}{2c^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \int x dx}{2c\sqrt{d - c^2 dx^2}} - \frac{x\sqrt{d - c^2 dx^2}(a + \operatorname{arccosh}(cx))}{2c^2 d} \right) \\
& \frac{4c^2}{x^3\sqrt{d - c^2 dx^2}(a + \operatorname{arccosh}(cx)) - \frac{bx^4\sqrt{cx-1}\sqrt{cx+1}}{16c\sqrt{d - c^2 dx^2}}} \\
& \quad \downarrow 15 \\
& 3 \left( \frac{\int \frac{a + \operatorname{arccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx}{2c^2} - \frac{x\sqrt{d - c^2 dx^2}(a + \operatorname{arccosh}(cx))}{2c^2 d} - \frac{bx^2\sqrt{cx-1}\sqrt{cx+1}}{4c\sqrt{d - c^2 dx^2}} \right) \\
& \frac{4c^2}{x^3\sqrt{d - c^2 dx^2}(a + \operatorname{arccosh}(cx)) - \frac{bx^4\sqrt{cx-1}\sqrt{cx+1}}{16c\sqrt{d - c^2 dx^2}}} \\
& \quad \downarrow 6307 \\
& \frac{x^3\sqrt{d - c^2 dx^2}(a + \operatorname{arccosh}(cx))}{4c^2 d} + \\
& 3 \left( -\frac{x\sqrt{d - c^2 dx^2}(a + \operatorname{arccosh}(cx))}{2c^2 d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{arccosh}(cx))^2}{4bc^3\sqrt{d - c^2 dx^2}} - \frac{bx^2\sqrt{cx-1}\sqrt{cx+1}}{4c\sqrt{d - c^2 dx^2}} \right) \\
& \frac{4c^2}{bx^4\sqrt{cx-1}\sqrt{cx+1}} \\
& \frac{4c^2}{16c\sqrt{d - c^2 dx^2}}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2], x]`

output `-1/16*(b*x^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*Sqrt[d - c^2*d*x^2]) - (x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(4*c^2*d) + (3*(-1/4*(b*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*Sqrt[d - c^2*d*x^2]) - (x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(2*c^2*d) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(4*b*c^3*Sqrt[d - c^2*d*x^2])))/(4*c^2)`

### 3.105.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

---

3.105.  $\int \frac{x^4(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^(p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.105.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 567 vs. 2(180) = 360.

Time = 0.94 (sec) , antiderivative size = 568, normalized size of antiderivative = 2.68

method	result
default	$-\frac{ax^3\sqrt{-c^2dx^2+d}}{4c^2d} - \frac{3ax\sqrt{-c^2dx^2+d}}{8c^4d} + \frac{3a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{8c^4\sqrt{c^2d}} + b\left(-\frac{3\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx)^2}{16dc^5(c^2x^2-1)}\right)$
parts	$-\frac{ax^3\sqrt{-c^2dx^2+d}}{4c^2d} - \frac{3ax\sqrt{-c^2dx^2+d}}{8c^4d} + \frac{3a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{8c^4\sqrt{c^2d}} + b\left(-\frac{3\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx)^2}{16dc^5(c^2x^2-1)}\right)$

```
input int(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/4*a*x^3/c^2/d*(-c^2*d*x^2+d)^(1/2)-3/8*a/c^4*x/d*(-c^2*d*x^2+d)^(1/2)+3/8*a/c^4/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b*(-3/16*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/c^5/(c^2*x^2-1)*arccosh(c*x)^2-1/256*(-d*(c^2*x^2-1))^(1/2)*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+4*arccosh(c*x))/d/c^5/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+2*arccosh(c*x))/d/c^5/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)-2*c*x)*(1+2*arccosh(c*x))/d/c^5/(c^2*x^2-1)-1/256*(-d*(c^2*x^2-1))^(1/2)*(-8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c^5*x^5+8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-12*c^3*x^3-(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c*x)*(1+4*arccosh(c*x))/d/c^5/(c^2*x^2-1)
```

3.105. 
$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$$

**3.105.5 Fracas [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-(b*x^4*arccosh(c*x) + a*x^4)*sqrt(-c^2*d*x^2 + d)/(c^2*d*x^2 - d), x)`

**3.105.6 Sympy [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**4*(a + b*acosh(c*x))/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.105.7 Maxima [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/8*a*(2*sqrt(-c^2*d*x^2 + d)*x^3/(c^2*d) + 3*sqrt(-c^2*d*x^2 + d)*x/(c^4*d) - 3*arcsin(c*x)/(c^5*sqrt(d))) + b*integrate(x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/sqrt(-c^2*d*x^2 + d), x)`



**3.105.8 Giac [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^4/sqrt(-c^2*d*x^2 + d), x)`

**3.105.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2), x)`

### 3.106 $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$

3.106.1 Optimal result . . . . .	1001
3.106.2 Mathematica [A] (verified) . . . . .	1001
3.106.3 Rubi [A] (verified) . . . . .	1002
3.106.4 Maple [B] (verified) . . . . .	1003
3.106.5 Fricas [A] (verification not implemented) . . . . .	1004
3.106.6 Sympy [F] . . . . .	1005
3.106.7 Maxima [A] (verification not implemented) . . . . .	1005
3.106.8 Giac [F(-2)] . . . . .	1005
3.106.9 Mupad [F(-1)] . . . . .	1006

#### 3.106.1 Optimal result

Integrand size = 27, antiderivative size = 156

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = -\frac{2bx\sqrt{-1 + cx}\sqrt{1 + cx}}{3c^3\sqrt{d - c^2dx^2}} - \frac{bx^3\sqrt{-1 + cx}\sqrt{1 + cx}}{9c\sqrt{d - c^2dx^2}} - \frac{2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{3c^4d} - \frac{x^2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{3c^2d}$$

output

```
-2/3*b*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/(-c^2*d*x^2+d)^(1/2)-1/9*b*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)-2/3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^4/d-1/3*x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^2/d
```

#### 3.106.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.72

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{d - c^2dx^2}(bcx\sqrt{-1 + cx}\sqrt{1 + cx}(6 + c^2x^2) - 3a(-2 + c^2x^2 + c^4x^4) - 3b(-2 + c^2x^2 + c^4x^4) \operatorname{arccosh}(cx))}{9c^4d(-1 + cx)(1 + cx)}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2], x]`

output `(Sqrt[d - c^2*d*x^2]*(b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(6 + c^2*x^2) - 3*a*(-2 + c^2*x^2 + c^4*x^4) - 3*b*(-2 + c^2*x^2 + c^4*x^4)*ArcCosh[c*x]))/(9*c^4*d*(-1 + c*x)*(1 + c*x))`

### 3.106.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6353, 15, 6329, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{2 \int \frac{x(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{3c^2} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int x^2 dx}{3c\sqrt{d - c^2 dx^2}} - \frac{x^2\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{3c^2 d} \\
 & \quad \downarrow \text{15} \\
 & \frac{2 \int \frac{x(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{3c^2 d} - \frac{bx^3\sqrt{cx - 1}\sqrt{cx + 1}}{9c\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{6329} \\
 & \frac{2 \left( -\frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int 1 dx}{c\sqrt{d - c^2 dx^2}} - \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{c^2 d} \right)}{3c^2} - \frac{x^2\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{3c^2 d} - \frac{bx^3\sqrt{cx - 1}\sqrt{cx + 1}}{9c\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{24} \\
 & -\frac{x^2\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{3c^2 d} + \frac{2 \left( -\frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{c^2 d} - \frac{bx\sqrt{cx - 1}\sqrt{cx + 1}}{c\sqrt{d - c^2 dx^2}} \right)}{3c^2} - \frac{bx^3\sqrt{cx - 1}\sqrt{cx + 1}}{9c\sqrt{d - c^2 dx^2}}
 \end{aligned}$$

input `Int[(x^3*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2], x]`

---

3.106.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$

```
output -1/9*(b*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*Sqrt[d - c^2*d*x^2]) - (x^2*S
qrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(3*c^2*d) + (2*(-((b*x*Sqrt[-1 +
c*x]*Sqrt[1 + c*x])/(c*Sqrt[d - c^2*d*x^2]))) - (Sqrt[d - c^2*d*x^2]*(a + b
*ArcCosh[c*x]))/(c^2*d)))/(3*c^2)
```

### 3.106.3.1 Defintions of rubi rules used

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e
_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2
*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x]
- Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)
^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*Ar
cCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*
d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.106.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 381 vs.  $2(132) = 264$ .

Time = 0.96 (sec) , antiderivative size = 382, normalized size of antiderivative = 2.45

method	result
default	$a \left( -\frac{x^2 \sqrt{-c^2 d x^2 + d}}{3c^2 d} - \frac{2\sqrt{-c^2 d x^2 + d}}{3d c^4} \right) + b \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (4c^4 x^4 - 5c^2 x^2 + 4\sqrt{cx-1} \sqrt{cx+1} c^3 x^3 - 3\sqrt{cx-1} \sqrt{cx+1} cx + 1)}{72c^4 d(c^2 x^2 - 1)} \right)$
parts	$a \left( -\frac{x^2 \sqrt{-c^2 d x^2 + d}}{3c^2 d} - \frac{2\sqrt{-c^2 d x^2 + d}}{3d c^4} \right) + b \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (4c^4 x^4 - 5c^2 x^2 + 4\sqrt{cx-1} \sqrt{cx+1} c^3 x^3 - 3\sqrt{cx-1} \sqrt{cx+1} cx + 1)}{72c^4 d(c^2 x^2 - 1)} \right)$

input `int(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output `a*(-1/3*x^2/c^2/d*(-c^2*d*x^2+d)^(1/2)-2/3/d/c^4*(-c^2*d*x^2+d)^(1/2))+b*(-1/72*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))/c^4/d/(c^2*x^2-1)-3/8*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+arccosh(c*x))/c^4/d/(c^2*x^2-1)-3/8*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))/c^4/d/(c^2*x^2-1)-1/72*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c*x))/c^4/d/(c^2*x^2-1))`

### 3.106.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.94

$$\int \frac{x^3(a + b\operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \frac{3(bc^4 x^4 + bc^2 x^2 - 2b)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1}) - (bc^3 x^3 + 6bcx)\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1} + 9(c^6 dx^2 - c^4 d)}{9(c^6 dx^2 - c^4 d)}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output `-1/9*(3*(b*c^4*x^4 + b*c^2*x^2 - 2*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - (b*c^3*x^3 + 6*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 3*(a*c^4*x^4 + a*c^2*x^2 - 2*a)*sqrt(-c^2*d*x^2 + d))/(c^6*d*x^2 - c^4*d)`

**3.106.6 Sympy [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**3*(a + b*acosh(c*x))/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.106.7 Maxima [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.84

$$\begin{aligned} \int \frac{x^3(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx &= -\frac{1}{3} b \left( \frac{\sqrt{-c^2 dx^2 + dx^2}}{c^2 d} + \frac{2\sqrt{-c^2 dx^2 + d}}{c^4 d} \right) \operatorname{arcosh}(cx) \\ &\quad - \frac{1}{3} a \left( \frac{\sqrt{-c^2 dx^2 + dx^2}}{c^2 d} + \frac{2\sqrt{-c^2 dx^2 + d}}{c^4 d} \right) \\ &\quad + \frac{(c^2 \sqrt{-d} x^3 + 6\sqrt{-d} x) b}{9 c^3 d} \end{aligned}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/3*b*(sqrt(-c^2*d*x^2 + d)*x^2/(c^2*d) + 2*sqrt(-c^2*d*x^2 + d)/(c^4*d)) *arccosh(c*x) - 1/3*a*(sqrt(-c^2*d*x^2 + d)*x^2/(c^2*d) + 2*sqrt(-c^2*d*x^2 + d)/(c^4*d)) + 1/9*(c^2*sqrt(-d)*x^3 + 6*sqrt(-d)*x)*b/(c^3*d)`

**3.106.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.106.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2), x)`

### 3.107 $\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$

3.107.1 Optimal result . . . . .	1007
3.107.2 Mathematica [A] (verified) . . . . .	1007
3.107.3 Rubi [A] (verified) . . . . .	1008
3.107.4 Maple [B] (verified) . . . . .	1009
3.107.5 Fricas [F] . . . . .	1010
3.107.6 Sympy [F] . . . . .	1010
3.107.7 Maxima [F] . . . . .	1011
3.107.8 Giac [F] . . . . .	1011
3.107.9 Mupad [F(-1)] . . . . .	1011

#### 3.107.1 Optimal result

Integrand size = 27, antiderivative size = 132

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = -\frac{bx^2\sqrt{-1+cx}\sqrt{1+cx}}{4c\sqrt{d-c^2dx^2}} - \frac{x\sqrt{d-c^2dx^2}(a + \operatorname{arccosh}(cx))}{2c^2d} + \frac{\sqrt{-1+cx}\sqrt{1+cx}(a + \operatorname{arccosh}(cx))^2}{4bc^3\sqrt{d-c^2dx^2}}$$

```
output -1/4*b*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)+1/4*(a+b*arc
cosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^3/(-c^2*d*x^2+d)^(1/2)-1/2*x*
(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^2/d
```

#### 3.107.2 Mathematica [A] (verified)

Time = 0.64 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.07

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = \frac{-\frac{4acx\sqrt{d-c^2dx^2}}{d} - \frac{4a \arctan\left(\frac{cx\sqrt{d-c^2dx^2}}{\sqrt{d}(-1+c^2x^2)}\right)}{\sqrt{d}} + \frac{b\sqrt{\frac{-1+cx}{1+cx}}(1+cx)(-\cosh(2\operatorname{arccosh}(cx))+2\operatorname{arccosh}(cx)(\operatorname{arccosh}(cx)+\sinh(2\operatorname{arccosh}(cx))))}{\sqrt{d-c^2dx^2}}}{8c^3}$$

```
input Integrate[(x^2*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2],x]
```



output  $((-4*a*c*x*\text{Sqrt}[d - c^2*d*x^2])/d - (4*a*\text{ArcTan}[(c*x*\text{Sqrt}[d - c^2*d*x^2])/(\text{Sqrt}[d]*(-1 + c^2*x^2))])/ \text{Sqrt}[d] + (b*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(-\text{Cosh}[2*\text{ArcCosh}[c*x]] + 2*\text{ArcCosh}[c*x]*(\text{ArcCosh}[c*x] + \text{Sinh}[2*\text{ArcCosh}[c*x]])))/\text{Sqrt}[d - c^2*d*x^2])/(8*c^3)$

### 3.107.3 Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6353, 15, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \text{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6353

$$\frac{\int \frac{a + \text{barccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx}{2c^2} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int x dx}{2c\sqrt{d - c^2 dx^2}} - \frac{x\sqrt{d - c^2 dx^2}(a + \text{barccosh}(cx))}{2c^2 d}$$

↓ 15

$$\frac{\int \frac{a + \text{barccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx}{2c^2} - \frac{x\sqrt{d - c^2 dx^2}(a + \text{barccosh}(cx))}{2c^2 d} - \frac{bx^2\sqrt{cx - 1}\sqrt{cx + 1}}{4c\sqrt{d - c^2 dx^2}}$$

↓ 6307

$$-\frac{x\sqrt{d - c^2 dx^2}(a + \text{barccosh}(cx))}{2c^2 d} + \frac{\sqrt{cx - 1}\sqrt{cx + 1}(a + \text{barccosh}(cx))^2}{4bc^3\sqrt{d - c^2 dx^2}} - \frac{bx^2\sqrt{cx - 1}\sqrt{cx + 1}}{4c\sqrt{d - c^2 dx^2}}$$

input  $\text{Int}[(x^2*(a + b*\text{ArcCosh}[c*x]))/\text{Sqrt}[d - c^2*d*x^2], x]$

output  $-1/4*(b*x^2*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/ (c*\text{Sqrt}[d - c^2*d*x^2]) - (x*\text{Sqrt}[d - c^2*d*x^2]*(a + b*\text{ArcCosh}[c*x]))/(2*c^2*d) + (\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(a + b*\text{ArcCosh}[c*x])^2)/(4*b*c^3*\text{Sqrt}[d - c^2*d*x^2])$

### 3.107.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`
- rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.107.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. 2(112) = 224.

Time = 0.87 (sec) , antiderivative size = 300, normalized size of antiderivative = 2.27

method	result
default	$-\frac{ax\sqrt{-c^2dx^2+d}}{2c^2d} + \frac{a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2c^2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx)^2}{4dc^3(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}(2c^3x^3-2c^2x^2-2cx+d)}{4dc^3(c^2x^2-1)}\right)$
parts	$-\frac{ax\sqrt{-c^2dx^2+d}}{2c^2d} + \frac{a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2c^2\sqrt{c^2d}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx)^2}{4dc^3(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}(2c^3x^3-2c^2x^2-2cx+d)}{4dc^3(c^2x^2-1)}\right)$

input `int(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/2*a*x/c^2/d*(-c^2*d*x^2+d)^{(1/2)}+1/2*a/c^2/(c^2*d)^{(1/2)}*\arctan((c^2*d) \\ & ^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)})+b*(-1/4*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)} \\ & )*(c*x+1)^{(1/2)}/d/c^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)^2-1/16*(-d*(c^2*x^2-1))^{(1/2)} \\ & *(2*c^3*x^3-2*c*x+2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-(c*x-1)^{(1/2)}*(c \\ & *x+1)^{(1/2)}*(-1+2*\operatorname{arccosh}(c*x)))/d/c^3/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^{(1/2)} \\ & *(-2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+2*c^3*x^3+(c*x-1)^{(1/2)}*(c*x+ \\ & 1)^{(1/2)}-2*c*x)*(1+2*\operatorname{arccosh}(c*x))/d/c^3/(c^2*x^2-1) \end{aligned}$$

### 3.107.5 Fracas [F]

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*x^2*arccosh(c*x) + a*x^2)/(c^2*d*x^2 - d), x)`

### 3.107.6 Sympy [F]

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**2*(a + b*acosh(c*x))/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.107.7 Maxima [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/2*a*(sqrt(-c^2*d*x^2 + d)*x/(c^2*d) - arcsin(c*x)/(c^3*sqrt(d))) + b*integrate(x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/sqrt(-c^2*d*x^2 + d), x)`

**3.107.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^2/sqrt(-c^2*d*x^2 + d), x)`

**3.107.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2), x)`

### 3.108 $\int \frac{x(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$

3.108.1 Optimal result . . . . .	1012
3.108.2 Mathematica [A] (verified) . . . . .	1012
3.108.3 Rubi [A] (verified) . . . . .	1013
3.108.4 Maple [B] (verified) . . . . .	1014
3.108.5 Fricas [A] (verification not implemented) . . . . .	1014
3.108.6 Sympy [F] . . . . .	1015
3.108.7 Maxima [A] (verification not implemented) . . . . .	1015
3.108.8 Giac [F] . . . . .	1015
3.108.9 Mupad [F(-1)] . . . . .	1016

#### 3.108.1 Optimal result

Integrand size = 25, antiderivative size = 72

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = -\frac{bx\sqrt{-1 + cx}\sqrt{1 + cx}}{c\sqrt{d - c^2dx^2}} - \frac{\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{c^2d}$$

output `-b*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)-(a+b*arccosh(c*x))  
*(-c^2*d*x^2+d)^(1/2)/c^2/d`

#### 3.108.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.18

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{d - c^2dx^2}(a - ac^2x^2 + bcx\sqrt{-1 + cx}\sqrt{1 + cx} + (b - bc^2x^2) \operatorname{arccosh}(cx))}{c^2d(-1 + cx)(1 + cx)}$$

input `Integrate[(x*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2],x]`

output `(Sqrt[d - c^2*d*x^2]*(a - a*c^2*x^2 + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] +  
(b - b*c^2*x^2)*ArcCosh[c*x]))/(c^2*d*(-1 + c*x)*(1 + c*x))`

### 3.108.3 Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$ , Rules used = {6329, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow 6329$$

$$-\frac{b\sqrt{cx-1}\sqrt{cx+1} \int 1 dx}{c\sqrt{d-c^2 dx^2}} - \frac{\sqrt{d-c^2 dx^2}(a + \operatorname{arccosh}(cx))}{c^2 d}$$

$$\downarrow 24$$

$$-\frac{\sqrt{d-c^2 dx^2}(a + \operatorname{arccosh}(cx))}{c^2 d} - \frac{bx\sqrt{cx-1}\sqrt{cx+1}}{c\sqrt{d-c^2 dx^2}}$$

input `Int[(x*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2],x]`

output `-((b*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*Sqrt[d - c^2*d*x^2])) - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(c^2*d)`

#### 3.108.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^{(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^{(p_.)}, x_Symbol] :> Simp[(d + e*x^2)^{(p + 1)}*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^{(p + 1/2)}*(-1 + c*x)^{(p + 1/2)}*(a + b*ArcCosh[c*x])^{(n - 1)}, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.108.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. 2(64) = 128.

Time = 0.62 (sec) , antiderivative size = 158, normalized size of antiderivative = 2.19

method	result
default	$-\frac{a\sqrt{-c^2dx^2+d}}{c^2d} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)(-1+\operatorname{arccosh}(cx))}{2c^2d(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}(-\sqrt{cx-1}\sqrt{cx+1}cx}{2c^2d(c^2x^2-1)}\right)$
parts	$-\frac{a\sqrt{-c^2dx^2+d}}{c^2d} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)(-1+\operatorname{arccosh}(cx))}{2c^2d(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}(-\sqrt{cx-1}\sqrt{cx+1}cx}{2c^2d(c^2x^2-1)}\right)$

input `int(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-a/c^2/d*(-c^2*d*x^2+d)^(1/2)+b*(-1/2*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+\operatorname{arccosh}(c*x))/c^2/d/(c^2*x^2-1)-1/2*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+\operatorname{arccosh}(c*x))/c^2/d/(c^2*x^2-1))$$

### 3.108.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.62

$$\int \frac{x(a + b\operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}bcx - (bc^2x^2 - b)\sqrt{-c^2dx^2 + d}\log(cx + \sqrt{c^2x^2 - 1}) - (ac^2x^2 - a)\sqrt{-c^2dx^2 + d}}{c^4dx^2 - c^2d}$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output 
$$(\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*b*c*x - (b*c^2*x^2 - b)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) - (a*c^2*x^2 - a)*\sqrt{-c^2*d*x^2 + d})/(c^4*d*x^2 - c^2*d)$$

**3.108.6 Sympy [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate(x*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x*(a + b*acosh(c*x))/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.108.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.88

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \frac{b\sqrt{-d}x}{cd} - \frac{\sqrt{-c^2 dx^2 + d} \operatorname{arccosh}(cx)}{c^2 d} - \frac{\sqrt{-c^2 dx^2 + d} a}{c^2 d}$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `b*sqrt(-d)*x/(c*d) - sqrt(-c^2*d*x^2 + d)*b*arccosh(c*x)/(c^2*d) - sqrt(-c^2*d*x^2 + d)*a/(c^2*d)`

**3.108.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x/sqrt(-c^2*d*x^2 + d), x)`



**3.108.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2),x)`output `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(1/2), x)`

### 3.109 $\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{d-c^2dx^2}} dx$

3.109.1 Optimal result	1017
3.109.2 Mathematica [A] (verified)	1017
3.109.3 Rubi [A] (verified)	1018
3.109.4 Maple [A] (verified)	1018
3.109.5 Fracas [F]	1019
3.109.6 Sympy [F]	1019
3.109.7 Maxima [F]	1019
3.109.8 Giac [F]	1020
3.109.9 Mupad [F(-1)]	1020

#### 3.109.1 Optimal result

Integrand size = 24, antiderivative size = 53

$$\int \frac{a + \operatorname{arccosh}(cx)}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{2bc\sqrt{d - c^2dx^2}}$$

output  $1/2*(a+b*\operatorname{arccosh}(c*x))^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c/(-c^2*d*x^2+d)^{(1/2)}$

#### 3.109.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \frac{a + \operatorname{arccosh}(cx)}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{2bc\sqrt{d - c^2dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/Sqrt[d - c^2*d*x^2], x]`

output  $(\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(a + b*\operatorname{ArcCosh}[c*x])^2)/(2*b*c*\operatorname{Sqrt}[d - c^2*d*x^2])$

### 3.109.3 Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6307

$$\frac{\sqrt{cx - 1} \sqrt{cx + 1} (a + b \operatorname{arccosh}(cx))^2}{2bc \sqrt{d - c^2 dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])/Sqrt[d - c^2*d*x^2], x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(2*b*c*Sqrt[d - c^2*d*x^2])`

#### 3.109.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.109.4 Maple [A] (verified)

Time = 0.56 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.68

method	result	size
default	$\frac{a \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{\sqrt{c^2 d}} - \frac{b \sqrt{-d(cx-1)(cx+1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^2}{2d(c^2 x^2 - 1)c}$	89
parts	$\frac{a \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{\sqrt{c^2 d}} - \frac{b \sqrt{-d(cx-1)(cx+1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^2}{2d(c^2 x^2 - 1)c}$	89

input `int((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2), x, method=_RETURNVERBOSE)`

output  $a/(c^2d)^{(1/2)}*\arctan((c^2d)^{(1/2)}*x/(-c^2d*x^2+d)^{(1/2)})-1/2*b*(-d*(c*x-1)*(c*x+1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)/c*\operatorname{arccosh}(c*x)^2$

### 3.109.5 Fricas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^2*d*x^2 - d), x)`

### 3.109.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

### 3.109.7 Maxima [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/sqrt(-c^2*d*x^2 + d), x) + a*arcsin(c*x)/(c*sqrt(d))`

**3.109.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/sqrt(-c^2*d*x^2 + d), x)`

**3.109.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{\sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^(1/2),x)`

output `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^(1/2), x)`

### 3.110 $\int \frac{a+b\operatorname{arccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx$

3.110.1 Optimal result . . . . .	1021
3.110.2 Mathematica [A] (verified) . . . . .	1021
3.110.3 Rubi [A] (verified) . . . . .	1022
3.110.4 Maple [A] (verified) . . . . .	1024
3.110.5 Fricas [F] . . . . .	1024
3.110.6 Sympy [F] . . . . .	1025
3.110.7 Maxima [F] . . . . .	1025
3.110.8 Giac [F] . . . . .	1025
3.110.9 Mupad [F(-1)] . . . . .	1026

#### 3.110.1 Optimal result

Integrand size = 27, antiderivative size = 151

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x\sqrt{d - c^2dx^2}} dx = \frac{2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} - \frac{ib\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} + \frac{ib\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

```
output 2*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)
*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)-I*b*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c
*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)+I*b*polylog
(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*
d*x^2+d)^(1/2)
```

#### 3.110.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.01

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x\sqrt{d - c^2dx^2}} dx = \frac{a \log(x)}{\sqrt{d}} - \frac{a \log(d + \sqrt{d}\sqrt{d - c^2dx^2})}{\sqrt{d}} - \frac{ib\sqrt{\frac{-1+cx}{1+cx}}(1 + cx) (\operatorname{arccosh}(cx) (\log(1 - ie^{-\operatorname{arccosh}(cx)}) - \log(1 + ie^{-\operatorname{arccosh}(cx)})) + \operatorname{PolyLog}(2, -ie^{-\operatorname{arccosh}(cx)}))}{\sqrt{d - c^2dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*Sqrt[d - c^2*d*x^2]),x]`

output `(a*Log[x])/Sqrt[d] - (a*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]])/Sqrt[d] - (I*b*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(ArcCosh[c*x]*(Log[1 - I/E^ArcCosh[c*x]] - Log[1 + I/E^ArcCosh[c*x]]) + PolyLog[2, (-I)/E^ArcCosh[c*x]] - PolyLog[2, I/E^ArcCosh[c*x]]))/Sqrt[d - c^2*d*x^2]`

### 3.110.3 Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.57, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6361}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int (a + \operatorname{barccosh}(cx)) \csc\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{4668}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \left(-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)})\right)}{\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{2715}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \left(-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)}\right)}{\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{2838}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \left(2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})\right)}{\sqrt{d - c^2 dx^2}}$$

---

3.110.  $\int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx$

input `Int[(a + b*ArcCosh[c*x])/(x*Sqrt[d - c^2*d*x^2]),x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/Sqrt[d - c^2*d*x^2]`

### 3.110.3.1 Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))]^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6361 `Int[(((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)^(m_))/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`



### 3.110.4 Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 326, normalized size of antiderivative = 2.16

method	result
default	$-\frac{a \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{\sqrt{d}} + b\left(\frac{i\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx) \ln(1+i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{d(c^2x^2-1)} - \frac{i\sqrt{-d(c^2x^2-1)}}{d(c^2x^2-1)}\right)$
parts	$-\frac{a \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{\sqrt{d}} + b\left(\frac{i\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx) \ln(1+i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{d(c^2x^2-1)} - \frac{i\sqrt{-d(c^2x^2-1)}}{d(c^2x^2-1)}\right)$

input `int((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output `-a/d^(1/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)+b*(I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))+I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))`

### 3.110.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^2*d*x^3 - d*x), x)`

**3.110.6 Sympy [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))/x/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))/(x*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.110.7 Maxima [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(sqrt(-c^2*d*x^2 + d)*x), x) - a*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/sqrt(d)`

**3.110.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(sqrt(-c^2*d*x^2 + d)*x), x)`

**3.110.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x \sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^(1/2)),x)`output `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^(1/2)), x)`

### 3.111 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2\sqrt{d-c^2dx^2}} dx$

3.111.1 Optimal result . . . . .	1027
3.111.2 Mathematica [A] (verified) . . . . .	1027
3.111.3 Rubi [A] (verified) . . . . .	1028
3.111.4 Maple [B] (verified) . . . . .	1029
3.111.5 Fricas [A] (verification not implemented) . . . . .	1029
3.111.6 Sympy [F] . . . . .	1030
3.111.7 Maxima [C] (verification not implemented) . . . . .	1030
3.111.8 Giac [F] . . . . .	1030
3.111.9 Mupad [F(-1)] . . . . .	1031

#### 3.111.1 Optimal result

Integrand size = 27, antiderivative size = 71

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2\sqrt{d - c^2dx^2}} dx = -\frac{\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{dx} - \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}\log(x)}{\sqrt{d - c^2dx^2}}$$

output `-b*c*ln(x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)-(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/d/x`

#### 3.111.2 Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}\left(\frac{\sqrt{-1+cx}\sqrt{1+cx}(a+\operatorname{arccosh}(cx))}{x} - bc\log(x)\right)}{\sqrt{d - c^2dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^2*Sqrt[d - c^2*d*x^2]),x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCos h[c*x]))/x - b*c*Log[x]))/Sqrt[d - c^2*d*x^2]`

### 3.111.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$ , Rules used = {6332, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx$$

↓ 6332

$$-\frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{x} dx}{\sqrt{d-c^2 dx^2}} - \frac{\sqrt{d-c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{dx}$$

↓ 14

$$-\frac{\sqrt{d-c^2 dx^2}(a + b \operatorname{arccosh}(cx))}{dx} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \log(x)}{\sqrt{d-c^2 dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*Sqrt[d - c^2*d*x^2]),x]`

output `-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(d*x)) - (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[x])/Sqrt[d - c^2*d*x^2]`

#### 3.111.3.1 Defintions of rubi rules used

rule 14 `Int[(a_)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 6332 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

### 3.111.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 207 vs. 2(63) = 126.

Time = 1.00 (sec) , antiderivative size = 208, normalized size of antiderivative = 2.93

method	result
default	$-\frac{a\sqrt{-c^2dx^2+d}}{dx} + b\left(-\frac{2\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}\operatorname{arccosh}(cx)c}{d(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\operatorname{arccosh}(cx)c}{x(c^2x^2-1)d}\right)$
parts	$-\frac{a\sqrt{-c^2dx^2+d}}{dx} + b\left(-\frac{2\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}\operatorname{arccosh}(cx)c}{d(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\operatorname{arccosh}(cx)c}{x(c^2x^2-1)d}\right)$

input `int((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-a/d/x*(-c^2*d*x^2+d)^{(1/2)}+b*(-2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)*\operatorname{arccosh}(c*x)*c-(-d*(c^2*x^2-1))^{(1/2)}*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+c^2*x^2-1)*\operatorname{arccosh}(c*x)/x/(c^2*x^2-1)/d+(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*c)$$

### 3.111.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 265, normalized size of antiderivative = 3.73

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^2\sqrt{d - c^2dx^2}} dx$$

$$= \left[ \frac{bc\sqrt{-d}x \log\left(\frac{c^2dx^6 + c^2dx^2 - dx^4 + \sqrt{-c^2dx^2 + d}\sqrt{c^2x^2 - 1}(x^4 - 1)\sqrt{-d - d}}{c^2x^4 - x^2}\right) + 2\sqrt{-c^2dx^2 + d}b \log(cx + \sqrt{c^2x^2 - 1})}{2dx} \right] +$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output 
$$[-1/2*(b*c*\sqrt{-d})*x*\log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + \sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*(x^4 - 1)*\sqrt{-d} - d)/(c^2*x^4 - x^2)) + 2*\sqrt{-c^2*d*x^2 + d}*b*\log(c*x + \sqrt{c^2*x^2 - 1}) + 2*\sqrt{-c^2*d*x^2 + d}*a)/(d*x), (b*c*\sqrt{d})*x*\arctan(\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*(x^2 + 1)*\sqrt{d}/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - \sqrt{-c^2*d*x^2 + d}*b*\log(c*x + \sqrt{c^2*x^2 - 1}) - \sqrt{-c^2*d*x^2 + d}*a)/(d*x)]$$

---

3.111. 
$$\int \frac{a+b\operatorname{arccosh}(cx)}{x^2\sqrt{d-c^2dx^2}} dx$$

**3.111.6 Sympy [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{arccosh}(cx)}{x^2 \sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))/x**2/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))/(x**2*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.111.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.63

$$\begin{aligned} & \int \frac{a + b \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx \\ &= - \frac{\left( c^2 d \sqrt{-\frac{1}{c^4 d}} \log\left(x^2 - \frac{1}{c^2}\right) + i (-1)^{-2c^2 dx^2 + 2d} \sqrt{d} \log\left(-2c^2 d + \frac{2d}{x^2}\right) \right) bc}{\sqrt{-c^2 dx^2 + d} b \operatorname{arccosh}(cx) - \frac{2d}{\sqrt{-c^2 dx^2 + d} a}} \end{aligned}$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/2*(c^2*d*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2) + I*(-1)^(-2*c^2*d*x^2 + 2*d)*sqrt(d)*log(-2*c^2*d + 2*d/x^2))*b*c/d - sqrt(-c^2*d*x^2 + d)*b*arccosh(c*x)/(d*x) - sqrt(-c^2*d*x^2 + d)*a/(d*x)`

**3.111.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{\sqrt{-c^2 dx^2 + dx^2}} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(sqrt(-c^2*d*x^2 + d)*x^2), x)`

---

3.111.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx$

**3.111.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^(1/2)),x)`output `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^(1/2)), x)`



### 3.112 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3\sqrt{d-c^2dx^2}} dx$

3.112.1 Optimal result . . . . .	1032
3.112.2 Mathematica [A] (warning: unable to verify) . . . . .	1033
3.112.3 Rubi [A] (verified) . . . . .	1033
3.112.4 Maple [A] (verified) . . . . .	1036
3.112.5 Fracas [F] . . . . .	1036
3.112.6 Sympy [F] . . . . .	1037
3.112.7 Maxima [F] . . . . .	1037
3.112.8 Giac [F] . . . . .	1037
3.112.9 Mupad [F(-1)] . . . . .	1038

#### 3.112.1 Optimal result

Integrand size = 27, antiderivative size = 238

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^3\sqrt{d - c^2dx^2}} dx = \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{2x\sqrt{d - c^2dx^2}} - \frac{\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{2dx^2} + \frac{c^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} - \frac{ibc^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{2\sqrt{d - c^2dx^2}} + \frac{ibc^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{2\sqrt{d - c^2dx^2}}$$

output

```
1/2*b*c*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x/(-c^2*d*x^2+d)^(1/2)+c^2*(a+b*arccos
h(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2
)/(-c^2*d*x^2+d)^(1/2)-1/2*I*b*c^2*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)
^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)+1/2*I*b*c^2*poly
log(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c
^2*d*x^2+d)^(1/2)-1/2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/d/x^2
```

**3.112.2 Mathematica [A] (warning: unable to verify)**

Time = 0.89 (sec) , antiderivative size = 309, normalized size of antiderivative = 1.30

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx = \frac{1}{2} \left( -\frac{a \sqrt{d - c^2 dx^2}}{dx^2} + \frac{ac^2 \log(x)}{\sqrt{d}} - \frac{ac^2 \log(d + \sqrt{d} \sqrt{d - c^2 dx^2})}{\sqrt{d}} \right) + \frac{b(1 + cx) \left( cx \sqrt{\frac{-1+cx}{1+cx}} - \operatorname{arccosh}(cx) + cx \operatorname{arccosh}(cx) - ic^2 x^2 \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) \log(1 - ie^{-\operatorname{arccosh}(cx)}) \right)}{2}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*Sqrt[d - c^2*d*x^2]),x]`output `((-(a*Sqrt[d - c^2*d*x^2])/(d*x^2)) + (a*c^2*Log[x])/Sqrt[d] - (a*c^2*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]])/Sqrt[d] + (b*(1 + c*x)*(c*x*Sqrt[(-1 + c*x)/(1 + c*x)] - ArcCosh[c*x] + c*x*ArcCosh[c*x] - I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] + I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] - I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*PolyLog[2, (-I)/E^ArcCosh[c*x]] + I*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*PolyLog[2, I/E^ArcCosh[c*x]]))/(x^2*Sqrt[d - c^2*d*x^2]))/2`**3.112.3 Rubi [A] (verified)**Time = 0.72 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.71, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$ , Rules used = {6347, 15, 6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx$$

↓ 6347

$$\frac{1}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x \sqrt{d - c^2 dx^2}} dx - \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} \int \frac{1}{x^2} dx}{2 \sqrt{d - c^2 dx^2}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2 dx^2}$$

↓ 15

$$\frac{1}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x \sqrt{d - c^2 dx^2}} dx - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2 dx^2} + \frac{bc \sqrt{cx - 1} \sqrt{cx + 1}}{2x \sqrt{d - c^2 dx^2}}$$

---

3.112.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx$

$$\begin{aligned}
& \downarrow 6361 \\
& \frac{c^2 \sqrt{cx-1} \sqrt{cx+1} \int \frac{a + \operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx) - \frac{\sqrt{d-c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2dx^2}}{2\sqrt{d-c^2 dx^2}} + \\
& \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x\sqrt{d-c^2 dx^2}} \\
& \downarrow 3042 \\
& \frac{c^2 \sqrt{cx-1} \sqrt{cx+1} \int (a + \operatorname{barccosh}(cx)) \operatorname{csc} \left( \operatorname{iarccosh}(cx) + \frac{\pi}{2} \right) \operatorname{darccosh}(cx)}{2\sqrt{d-c^2 dx^2}} - \\
& \quad \frac{\sqrt{d-c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x\sqrt{d-c^2 dx^2}} \\
& \downarrow 4668 \\
& \frac{c^2 \sqrt{cx-1} \sqrt{cx+1} (-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan)}{2\sqrt{d-c^2 dx^2}} \\
& \quad \frac{\sqrt{d-c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x\sqrt{d-c^2 dx^2}} \\
& \downarrow 2715 \\
& \frac{c^2 \sqrt{cx-1} \sqrt{cx+1} (-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{2\sqrt{d-c^2 dx^2}} \\
& \quad \frac{\sqrt{d-c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x\sqrt{d-c^2 dx^2}} \\
& \downarrow 2838 \\
& \frac{c^2 \sqrt{cx-1} \sqrt{cx+1} (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}))}{2\sqrt{d-c^2 dx^2}} \\
& \quad \frac{\sqrt{d-c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2dx^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x\sqrt{d-c^2 dx^2}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*Sqrt[d - c^2*d*x^2]),x]`

output `(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*x*Sqrt[d - c^2*d*x^2]) - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])/(2*d*x^2) + (c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(2*Sqrt[d - c^2*d*x^2])`

## 3.112.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`
- rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((f_.)*(x_)^m)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`
- rule 6361 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*(x_)^m_/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

### 3.112.4 Maple [A] (verified)

Time = 1.11 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.81

method	result
default	$-\frac{a\sqrt{-c^2dx^2+d}}{2dx^2} - \frac{ac^2 \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{2\sqrt{d}} + b\left(-\frac{(c^2x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1}\sqrt{cx+1}cx - \operatorname{arccosh}(cx))\sqrt{-d(c^2x^2-1)}}{2d(c^2x^2-1)x^2} + \dots\right)$
parts	$-\frac{a\sqrt{-c^2dx^2+d}}{2dx^2} - \frac{ac^2 \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{2\sqrt{d}} + b\left(-\frac{(c^2x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1}\sqrt{cx+1}cx - \operatorname{arccosh}(cx))\sqrt{-d(c^2x^2-1)}}{2d(c^2x^2-1)x^2} + \dots\right)$

input `int((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output

```
-1/2*a/d/x^2*(-c^2*d*x^2+d)^(1/2)-1/2*a*c^2/d^(1/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)+b*(-1/2*(c^2*x^2*arccosh(c*x)+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-arccosh(c*x))*(-d*(c^2*x^2-1))^(1/2)/d/(c^2*x^2-1)/x^2+1/2*I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2-1/2*I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2+1/2*I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2-1/2*I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2)
```

### 3.112.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{\sqrt{-c^2 dx^2 + dx^3}} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^2*d*x^5 - d*x^3), x)`

**3.112.6 Sympy [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 \sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate((a+b*acosh(c*x))/x**3/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))/(x**3*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.112.7 Maxima [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + dx^3}} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/2*(c^2*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/sqrt(d) + sqrt(-c^2*d*x^2 + d)/(d*x^2))*a + b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(sqrt(-c^2*d*x^2 + d)*x^3), x)`

**3.112.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + dx^3}} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(sqrt(-c^2*d*x^2 + d)*x^3), x)`

**3.112.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^(1/2)),x)`output `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^(1/2)), x)`

### 3.113 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^4\sqrt{d-c^2dx^2}} dx$

3.113.1 Optimal result . . . . .	1039
3.113.2 Mathematica [A] (verified) . . . . .	1039
3.113.3 Rubi [A] (verified) . . . . .	1040
3.113.4 Maple [A] (verified) . . . . .	1042
3.113.5 Fricas [A] (verification not implemented) . . . . .	1042
3.113.6 Sympy [F] . . . . .	1043
3.113.7 Maxima [A] (verification not implemented) . . . . .	1043
3.113.8 Giac [F] . . . . .	1044
3.113.9 Mupad [F(-1)] . . . . .	1044

#### 3.113.1 Optimal result

Integrand size = 27, antiderivative size = 155

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^4\sqrt{d - c^2dx^2}} dx = \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{6x^2\sqrt{d - c^2dx^2}} - \frac{\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))}{3dx^3} - \frac{2c^2\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))}{3dx} - \frac{2bc^3\sqrt{-1 + cx}\sqrt{1 + cx} \log(x)}{3\sqrt{d - c^2dx^2}}$$

```
output 1/6*b*c*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x^2/(-c^2*d*x^2+d)^(1/2)-2/3*b*c^3*ln(
x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)-1/3*(a+b*arccosh(c*x))
*(-c^2*d*x^2+d)^(1/2)/d/x^3-2/3*c^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)
)/d/x
```

#### 3.113.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.12

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^4\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{d - c^2dx^2}(bcx + 6bc^3x^3 + 2a\sqrt{-1 + cx}\sqrt{1 + cx} + 4ac^2x^2\sqrt{-1 + cx}\sqrt{1 + cx} + 2b\sqrt{-1 + cx}\sqrt{1 + cx})}{6dx^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$



input `Integrate[(a + b*ArcCosh[c*x])/(x^4*Sqrt[d - c^2*d*x^2]),x]`

output `-1/6*(Sqrt[d - c^2*d*x^2]*(b*c*x + 6*b*c^3*x^3 + 2*a*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 4*a*c^2*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(1 + 2*c^2*x^2)*ArcCosh[c*x] - 4*b*c^3*x^3*Log[-1 + c*x] - 4*b*c^3*x^3*Log[1 + (-1 + c*x)^(-1)]))/(d*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.113.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.99, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6347, 15, 6332, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{arccosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx \\
 & \quad \downarrow \text{6347} \\
 & \frac{2}{3} c^2 \int \frac{a + \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx - \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} \int \frac{1}{x^3} dx}{3 \sqrt{d - c^2 dx^2}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))}{3 dx^3} \\
 & \quad \downarrow \text{15} \\
 & \frac{2}{3} c^2 \int \frac{a + \operatorname{arccosh}(cx)}{x^2 \sqrt{d - c^2 dx^2}} dx - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))}{3 dx^3} + \frac{bc \sqrt{cx - 1} \sqrt{cx + 1}}{6 x^2 \sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{6332} \\
 & \frac{2}{3} c^2 \left( - \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} \int \frac{1}{x} dx}{\sqrt{d - c^2 dx^2}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))}{dx} \right) - \\
 & \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))}{3 dx^3} + \frac{bc \sqrt{cx - 1} \sqrt{cx + 1}}{6 x^2 \sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{14} \\
 & \frac{2}{3} c^2 \left( - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))}{dx} - \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} \log(x)}{\sqrt{d - c^2 dx^2}} \right) - \\
 & \quad \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))}{3 dx^3} + \frac{bc \sqrt{cx - 1} \sqrt{cx + 1}}{6 x^2 \sqrt{d - c^2 dx^2}}
 \end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^4*Sqrt[d - c^2*d*x^2]),x]`

3.113.  $\int \frac{a + \operatorname{arccosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx$

output  $(b*c*\sqrt{-1 + c*x}*\sqrt{1 + c*x})/(6*x^2*\sqrt{d - c^2*d*x^2}) - (\sqrt{d - c^2*d*x^2}*(a + b*\text{ArcCosh}[c*x]))/(3*d*x^3) + (2*c^2*(-((\sqrt{d - c^2*d*x^2})*(a + b*\text{ArcCosh}[c*x]))/(d*x)) - (b*c*\sqrt{-1 + c*x}*\sqrt{1 + c*x}*\text{Log}[x])/(\sqrt{d - c^2*d*x^2}))/3$

### 3.113.3.1 Defintions of rubi rules used

rule 14  $\text{Int}[(a\_)/(x\_), x\_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$

rule 15  $\text{Int}[(a\_)*(x_)^(m\_), x\_Symbol] \rightarrow \text{Simp}[a*(x^(m + 1)/(m + 1)), x] \text{ ; FreeQ}[\{a, m\}, x] \ \&\& \ \text{NeQ}[m, -1]$

rule 6332  $\text{Int}[(a\_ + \text{ArcCosh}[(c\_)*(x\_)]*(b\_))^(n\_)*((f\_)*(x\_))^(m_)*((d_ + (e_)*(x_)^2)^(p_)), x\_Symbol] \rightarrow \text{Simp}[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*\text{ArcCosh}[c*x])^n/(d*f*(m + 1))), x] + \text{Simp}[b*c*(n/(f*(m + 1)))*\text{Simp}[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] \text{ Int}[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*\text{ArcCosh}[c*x])^(n - 1), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, m, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{EqQ}[m + 2*p + 3, 0] \ \&\& \ \text{NeQ}[m, -1]$

rule 6347  $\text{Int}[(a\_ + \text{ArcCosh}[(c\_)*(x\_)]*(b\_))^(n\_)*((f\_)*(x\_))^(m_)*((d_ + (e_)*(x_)^2)^(p_)), x\_Symbol] \rightarrow \text{Simp}[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*\text{ArcCosh}[c*x])^n/(d*f*(m + 1))), x] + (\text{Simp}[c^2*((m + 2*p + 3)/(f^2*(m + 1)))] \text{ Int}[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*\text{ArcCosh}[c*x])^n, x], x] + \text{Simp}[b*c*(n/(f*(m + 1)))*\text{Simp}[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] \text{ Int}[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*\text{ArcCosh}[c*x])^(n - 1), x], x] \text{ ; FreeQ}[\{a, b, c, d, e, f, p\}, x] \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{ILtQ}[m, -1]$

### 3.113.4 Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.24

method	result
default	$a \left( -\frac{\sqrt{-c^2 d x^2 + d}}{3 d x^3} - \frac{2 c^2 \sqrt{-c^2 d x^2 + d}}{3 d x} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{c x - 1} \sqrt{c x + 1} \left( 4 \sqrt{c x + 1} \operatorname{arccosh}(c x) \sqrt{c x - 1} c^2 x^2 + 4 c^3 x^3 \operatorname{arccosh}(c x) \right)}{6 d x^3 (c^2 x^2 - 1)}$
parts	$a \left( -\frac{\sqrt{-c^2 d x^2 + d}}{3 d x^3} - \frac{2 c^2 \sqrt{-c^2 d x^2 + d}}{3 d x} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{c x - 1} \sqrt{c x + 1} \left( 4 \sqrt{c x + 1} \operatorname{arccosh}(c x) \sqrt{c x - 1} c^2 x^2 + 4 c^3 x^3 \operatorname{arccosh}(c x) \right)}{6 d x^3 (c^2 x^2 - 1)}$

input `int((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output `a*(-1/3/d/x^3*(-c^2*d*x^2+d)^(1/2)-2/3*c^2/d/x*(-c^2*d*x^2+d)^(1/2))-1/6*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(4*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2+4*c^3*x^3*arccosh(c*x)-4*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3+2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)/d/x^3/(c^2*x^2-1)`

### 3.113.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 479, normalized size of antiderivative = 3.09

$$\int \frac{a + b \operatorname{arccosh}(c x)}{x^4 \sqrt{d - c^2 x^2}} dx$$

$$= \left[ \frac{2(2 b c^4 x^4 - b c^2 x^2 - b) \sqrt{-c^2 d x^2 + d} \log(c x + \sqrt{c^2 x^2 - 1}) + 2(b c^5 x^5 - b c^3 x^3) \sqrt{-d} \log\left(\frac{c^2 d x^6 + c^2 d x^2 - d x^4}{6(c^2 x^2 - 1)}\right)}{6(c^2 x^2 - 1)^2} \right]$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output `[-1/6*(2*(2*b*c^4*x^4 - b*c^2*x^2 - b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + 2*(b*c^5*x^5 - b*c^3*x^3)*sqrt(-d)*log((c^2*d*x^6 + c^2*d*x^2 - d*x^4 + sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^4 - 1)*sqrt(-d) - d)/(c^2*x^4 - x^2)) - sqrt(-c^2*d*x^2 + d)*(b*c*x^3 - b*c*x)*sqrt(c^2*x^2 - 1) + 2*(2*a*c^4*x^4 - a*c^2*x^2 - a)*sqrt(-c^2*d*x^2 + d)/(c^2*d*x^5 - d*x^3), 1/6*(4*(b*c^5*x^5 - b*c^3*x^3)*sqrt(d)*arctan(sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*(x^2 + 1)*sqrt(d)/(c^2*d*x^4 - (c^2 + 1)*d*x^2 + d)) - 2*(2*b*c^4*x^4 - b*c^2*x^2 - b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + sqrt(-c^2*d*x^2 + d)*(b*c*x^3 - b*c*x)*sqrt(c^2*x^2 - 1) - 2*(2*a*c^4*x^4 - a*c^2*x^2 - a)*sqrt(-c^2*d*x^2 + d))/(c^2*d*x^5 - d*x^3)]`

### 3.113.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4 \sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate((a+b*acosh(c*x))/x**4/(-c**2*d*x**2+d)**(1/2), x)`

output `Integral((a + b*acosh(c*x))/(x**4*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

### 3.113.7 Maxima [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.86

$$\begin{aligned} \int \frac{a + b \operatorname{arccosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx &= \frac{1}{6} \left( \frac{4c^2 \sqrt{-d} \log(x)}{d} - \frac{\sqrt{-d}}{dx^2} \right) bc \\ &\quad - \frac{1}{3} b \left( \frac{2\sqrt{-c^2 dx^2 + dc^2}}{dx} + \frac{\sqrt{-c^2 dx^2 + d}}{dx^3} \right) \operatorname{arccosh}(cx) \\ &\quad - \frac{1}{3} a \left( \frac{2\sqrt{-c^2 dx^2 + dc^2}}{dx} + \frac{\sqrt{-c^2 dx^2 + d}}{dx^3} \right) \end{aligned}$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(1/2), x, algorithm="maxima")`

output `1/6*(4*c^2*sqrt(-d)*log(x)/d - sqrt(-d)/(d*x^2))*b*c - 1/3*b*(2*sqrt(-c^2*d*x^2 + d)*c^2/(d*x) + sqrt(-c^2*d*x^2 + d)/(d*x^3))*arccosh(c*x) - 1/3*a*(2*sqrt(-c^2*d*x^2 + d)*c^2/(d*x) + sqrt(-c^2*d*x^2 + d)/(d*x^3))`

---

3.113.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx$

**3.113.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{-c^2 dx^2 + dx^4}} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(sqrt(-c^2*d*x^2 + d)*x^4), x)`

**3.113.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4 \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^(1/2)),x)`

output `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^(1/2)), x)`

**3.114** 
$$\int \frac{x^5(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$$

3.114.1 Optimal result . . . . . 1045  
 3.114.2 Mathematica [A] (verified) . . . . . 1046  
 3.114.3 Rubi [A] (verified) . . . . . 1046  
 3.114.4 Maple [A] (verified) . . . . . 1048  
 3.114.5 Fricas [A] (verification not implemented) . . . . . 1048  
 3.114.6 Sympy [F] . . . . . 1049  
 3.114.7 Maxima [F] . . . . . 1049  
 3.114.8 Giac [F(-2)] . . . . . 1050  
 3.114.9 Mupad [F(-1)] . . . . . 1050

**3.114.1 Optimal result**

Integrand size = 27, antiderivative size = 233

$$\int \frac{x^5(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = -\frac{5bx\sqrt{d - c^2dx^2}}{3c^5d^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bx^3\sqrt{d - c^2dx^2}}{9c^3d^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{a + \operatorname{arccosh}(cx)}{c^6d\sqrt{d - c^2dx^2}} + \frac{2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{c^6d^2} - \frac{(d - c^2dx^2)^{3/2}(a + \operatorname{arccosh}(cx))}{3c^6d^3} - \frac{b\sqrt{d - c^2dx^2}\operatorname{arctanh}(cx)}{c^6d^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output -1/3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))/c^6/d^3+(a+b*arccosh(c*x))/c^
6/d/(-c^2*d*x^2+d)^(1/2)+2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^6/d^2
-5/3*b*x*(-c^2*d*x^2+d)^(1/2)/c^5/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/9*b*x^
3*(-c^2*d*x^2+d)^(1/2)/c^3/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*arctanh(c*x)*
(-c^2*d*x^2+d)^(1/2)/c^6/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.114.2 Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.62

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \frac{24a - 12ac^2 x^2 - 3ac^4 x^4 + 15bcx\sqrt{-1 + cx}\sqrt{1 + cx} + bc^3 x^3 \sqrt{-1 + cx}\sqrt{1 + cx}}{9c^6 d \sqrt{d - c^2 dx^2}}$$

input `Integrate[(x^5*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]`output `(24*a - 12*a*c^2*x^2 - 3*a*c^4*x^4 + 15*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 3*b*(-8 + 4*c^2*x^2 + c^4*x^4)*ArcCosh[c*x] + 9*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*ArcTanh[c*x])/(9*c^6*d*Sqrt[d - c^2*d*x^2])`**3.114.3 Rubi [A] (verified)**Time = 0.49 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.70, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx \\ & \quad \downarrow \text{6337} \\ & -\frac{bc\sqrt{d - c^2 dx^2} \int \frac{-c^4 x^4 - 4c^2 x^2 + 8}{3c^6 d^2 (1 - c^2 x^2)} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6 d^3} + \\ & \quad \frac{2\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{c^6 d^2} + \frac{a + \operatorname{barccosh}(cx)}{c^6 d \sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{27} \\ & -\frac{b\sqrt{d - c^2 dx^2} \int \frac{-c^4 x^4 - 4c^2 x^2 + 8}{1 - c^2 x^2} dx}{3c^5 d^2 \sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6 d^3} + \\ & \quad \frac{2\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{c^6 d^2} + \frac{a + \operatorname{barccosh}(cx)}{c^6 d \sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{1467} \end{aligned}$$

---

3.114.  $\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& -\frac{b\sqrt{d-c^2dx^2} \int \left( c^2x^2 + \frac{3}{1-c^2x^2} + 5 \right) dx}{3c^5d^2\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6d^3} + \\
& \quad \frac{2\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))}{c^6d^2} + \frac{a + \operatorname{barccosh}(cx)}{c^6d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{2009} \\
& -\frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{3c^6d^3} + \frac{2\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))}{c^6d^2} + \frac{a + \operatorname{barccosh}(cx)}{c^6d\sqrt{d-c^2dx^2}} - \\
& \quad \frac{b\left(\frac{3\operatorname{arctanh}(cx)}{c} + \frac{c^2x^3}{3} + 5x\right)\sqrt{d-c^2dx^2}}{3c^5d^2\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(x^5*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]`

output `(a + b*ArcCosh[c*x])/(c^6*d*Sqrt[d - c^2*d*x^2]) + (2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(c^6*d^2) - ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(3*c^6*d^3) - (b*Sqrt[d - c^2*d*x^2]*(5*x + (c^2*x^3)/3 + (3*ArcTanh[c*x])/c))/(3*c^5*d^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.114.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

---

3.114.  $\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx$



**3.114.4 Maple [A] (verified)**

Time = 1.34 (sec) , antiderivative size = 325, normalized size of antiderivative = 1.39

method	result
default	$a \left( -\frac{x^4}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{-\frac{4x^2}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{8}{3d c^4 \sqrt{-c^2 d x^2 + d}}}{c^2} \right) + \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1} (3\sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx))}{c^2}$
parts	$a \left( -\frac{x^4}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{-\frac{4x^2}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{8}{3d c^4 \sqrt{-c^2 d x^2 + d}}}{c^2} \right) + \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1} (3\sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx))}{c^2}$

```
input int(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output a*(-1/3*x^4/c^2/d/(-c^2*d*x^2+d)^(1/2)+4/3/c^2*(-x^2/c^2/d/(-c^2*d*x^2+d)^(1/2)+2/d/c^4/(-c^2*d*x^2+d)^(1/2))+1/9*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4-c^5*x^5+12*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-14*c^3*x^3-9*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+9*ln((c*x-1)^(1/2)*(c*x+1)^(1/2))+c*x-1)*x^2*c^2-24*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+15*c*x+9*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-9*ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1))/(c^2*x^2-1)^2/d^2/c^6
```

**3.114.5 Fricas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 489, normalized size of antiderivative = 2.10

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \frac{\left[ \frac{12(bc^4 x^4 + 4bc^2 x^2 - 8b)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1}) - 9(bc^2 x^2 - b)}{18(c^8 d^2 x^2 - c^6 d)} \right.}{\left. - \frac{9(bc^2 x^2 - b)\sqrt{d} \arctan\left(\frac{2\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1}c\sqrt{dx}}{c^4 dx^4 - d}\right) - 6(bc^4 x^4 + 4bc^2 x^2 - 8b)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1})}{18(c^8 d^2 x^2 - c^6 d)} \right]}$$

```
input integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")
```

---

3.114.  $\int \frac{x^5(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$

output `[1/36*(12*(b*c^4*x^4 + 4*b*c^2*x^2 - 8*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - 9*(b*c^2*x^2 - b)*sqrt(-d)*log(-(c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 - 4*(c^3*x^3 + c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1))*sqrt(-d) - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)) - 4*(b*c^3*x^3 + 15*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) + 12*(a*c^4*x^4 + 4*a*c^2*x^2 - 8*a)*sqrt(-c^2*d*x^2 + d))/(c^8*d^2*x^2 - c^6*d^2), -1/18*(9*(b*c^2*x^2 - b)*sqrt(d)*arctan(2*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*c*sqrt(d)*x/(c^4*d*x^4 - d)) - 6*(b*c^4*x^4 + 4*b*c^2*x^2 - 8*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + 2*(b*c^3*x^3 + 15*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 6*(a*c^4*x^4 + 4*a*c^2*x^2 - 8*a)*sqrt(-c^2*d*x^2 + d))/(c^8*d^2*x^2 - c^6*d^2)]`

### 3.114.6 Sympy [F]

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**5*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x**5*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

### 3.114.7 Maxima [F]

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^5}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

```
output -1/3*a*(x^4/(sqrt(-c^2*d*x^2 + d)*c^2*d) + 4*x^2/(sqrt(-c^2*d*x^2 + d)*c^4
*d) - 8/(sqrt(-c^2*d*x^2 + d)*c^6*d) + 1/9*b*(((c^4*sqrt(d)*x^4 + 16*c^2*
sqrt(d)*x^2 - 8*sqrt(d))*sqrt(c*x + 1)*sqrt(c*x - 1)/sqrt(-c*x + 1) - 3*(c
^5*sqrt(d)*x^5 + 4*c^3*sqrt(d)*x^3 - 8*c*sqrt(d)*x + (c^4*sqrt(d)*x^4 + 4*
c^2*sqrt(d)*x^2 - 8*sqrt(d))*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c
*x + 1)*sqrt(c*x - 1))/sqrt(-c*x + 1))/(sqrt(c*x + 1)*c^7*d^2*x + (c*x + 1
)*sqrt(c*x - 1)*c^6*d^2) + 9*integrate(1/9*(3*c^7*sqrt(d)*x^7 + 9*c^5*sqrt
(d)*x^5 - 36*c^3*sqrt(d)*x^3 + 24*c*sqrt(d)*x + (3*c^6*sqrt(d)*x^6 + 8*c^4
*sqrt(d)*x^4 - 52*c^2*sqrt(d)*x^2 + 32*sqrt(d))*e^(1/2*log(c*x + 1) + 1/2*
log(c*x - 1)))/(sqrt(-c*x + 1)*((c^7*d^2*x^2 - c^5*d^2)*e^(3/2*log(c*x + 1
) + log(c*x - 1)) + 2*(c^8*d^2*x^3 - c^6*d^2*x)*e^(log(c*x + 1) + 1/2*log(
c*x - 1)) + (c^9*d^2*x^4 - c^7*d^2*x^2)*sqrt(c*x + 1))), x))
```

### 3.114.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.114.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$$

```
input int((x^5*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2),x)
```

```
output int((x^5*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2), x)
```

$$3.115 \quad \int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$$

3.115.1 Optimal result . . . . .	1051
3.115.2 Mathematica [A] (warning: unable to verify) . . . . .	1052
3.115.3 Rubi [A] (verified) . . . . .	1052
3.115.4 Maple [A] (verified) . . . . .	1055
3.115.5 Fricas [F] . . . . .	1056
3.115.6 Sympy [F] . . . . .	1056
3.115.7 Maxima [F] . . . . .	1056
3.115.8 Giac [F(-2)] . . . . .	1057
3.115.9 Mupad [F(-1)] . . . . .	1057

### 3.115.1 Optimal result

Integrand size = 27, antiderivative size = 226

$$\begin{aligned} \int \frac{x^4(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx &= \frac{bx^2\sqrt{-1 + cx}\sqrt{1 + cx}}{4c^3d\sqrt{d - c^2dx^2}} \\ &+ \frac{x^3(a + \operatorname{arccosh}(cx))}{c^2d\sqrt{d - c^2dx^2}} + \frac{3x\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{2c^4d^2} \\ &- \frac{3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{4bc^5d\sqrt{d - c^2dx^2}} - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx} \log(1 - c^2x^2)}{2c^5d\sqrt{d - c^2dx^2}} \end{aligned}$$

output  $x^3*(a+b*\operatorname{arccosh}(c*x))/c^2/d/(-c^2*d*x^2+d)^{(1/2)}+1/4*b*x^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/d/(-c^2*d*x^2+d)^{(1/2)}-3/4*(a+b*\operatorname{arccosh}(c*x))^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c^5/d/(-c^2*d*x^2+d)^{(1/2)}-1/2*b*\ln(-c^2*x^2+1)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^5/d/(-c^2*d*x^2+d)^{(1/2)}+3/2*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/c^4/d^2$

**3.115.2 Mathematica [A] (warning: unable to verify)**

Time = 1.24 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.85

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \frac{-4acdx(-3 + c^2 x^2) + 12a\sqrt{d}\sqrt{d - c^2 dx^2} \arctan\left(\frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right) + bd(8cx\operatorname{arccosh}(cx) - 8\sqrt{d}\sqrt{d - c^2 dx^2})}{(d - c^2 dx^2)^{3/2}}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2), x]`output `(-4*a*c*d*x*(-3 + c^2*x^2) + 12*a*Sqrt[d]*Sqrt[d - c^2*d*x^2]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + b*d*(8*c*x*ArcCosh[c*x] - Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(6*ArcCosh[c*x]^2 - Cosh[2*ArcCosh[c*x]]) + 8*Log[Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)] + 2*ArcCosh[c*x]*Sinh[2*ArcCosh[c*x]]))/(8*c^5*d^2*Sqrt[d - c^2*d*x^2])`**3.115.3 Rubi [A] (verified)**Time = 0.83 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.08, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6349, 25, 82, 243, 49, 2009, 6353, 15, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx \\ & \quad \downarrow 6349 \\ & -\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{c^2 d} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int -\frac{x^3}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))}{c^2 d\sqrt{d - c^2 dx^2}} \\ & \quad \downarrow 25 \\ & -\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{c^2 d} + \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x^3}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))}{c^2 d\sqrt{d - c^2 dx^2}} \\ & \quad \downarrow 82 \\ & -\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{c^2 d} + \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x^3}{1 - c^2 x^2} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))}{c^2 d\sqrt{d - c^2 dx^2}} \end{aligned}$$

---

3.115.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& \downarrow 243 \\
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2}{1-c^2x^2} dx^2}{2cd\sqrt{d-c^2dx^2}} + \frac{x^3(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} \\
& \downarrow 49 \\
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \left(-\frac{1}{c^2} - \frac{1}{c^2(c^2x^2-1)}\right) dx^2}{2cd\sqrt{d-c^2dx^2}} + \frac{x^3(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} \\
& \downarrow 2009 \\
& -\frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^3(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{2cd\sqrt{d-c^2dx^2}} \\
& \downarrow 6353 \\
& -\frac{3 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{d-c^2dx^2}} dx}{2c^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \int x dx}{2c\sqrt{d-c^2dx^2}} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2c^2d} \right)}{c^2d} + \\
& \frac{x^3(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{2cd\sqrt{d-c^2dx^2}} \\
& \downarrow 15 \\
& -\frac{3 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{d-c^2dx^2}} dx}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2c^2d} - \frac{bx^2\sqrt{cx-1}\sqrt{cx+1}}{4c\sqrt{d-c^2dx^2}} \right)}{c^2d} + \frac{x^3(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} + \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1} \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{2cd\sqrt{d-c^2dx^2}} \\
& \downarrow 6307 \\
& \frac{x^3(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} - \\
& 3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^2}{4bc^3\sqrt{d-c^2dx^2}} - \frac{bx^2\sqrt{cx-1}\sqrt{cx+1}}{4c\sqrt{d-c^2dx^2}} \right) + \\
& \frac{c^2d}{b\sqrt{cx-1}\sqrt{cx+1} \left(-\frac{x^2}{c^2} - \frac{\log(1-c^2x^2)}{c^4}\right)}{2cd\sqrt{d-c^2dx^2}}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]`

$$3.115. \quad \int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$$

```
output (x^3*(a + b*ArcCosh[c*x]))/(c^2*d*Sqrt[d - c^2*d*x^2]) - (3*(-1/4*(b*x^2*S
qrt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[d - c^2*d*x^2]) - (x*Sqrt[d - c^2*d*x
^2]*(a + b*ArcCosh[c*x]))/(2*c^2*d) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b
*ArcCosh[c*x])^2)/(4*b*c^3*Sqrt[d - c^2*d*x^2]))/(c^2*d) + (b*Sqrt[-1 + c
*x]*Sqrt[1 + c*x]*(-(x^2/c^2) - Log[1 - c^2*x^2]/c^4))/(2*c*d*Sqrt[d - c^2
*d*x^2])
```

### 3.115.3.1 Defintions of rubi rules used

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 49 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

```
rule 82 Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_
))^p, x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d,
e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]
```

```
rule 243 Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int
t[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && I
negerQ[(m - 1)/2]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6307 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_
Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d
+ e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x
] && EqQ[c^2*d + e, 0] && NeQ[n, -1]
```

rule 6349 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]`

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.115.4 Maple [A] (verified)

Time = 1.34 (sec) , antiderivative size = 301, normalized size of antiderivative = 1.33

method	result
default	$-\frac{ax^3}{2c^2d\sqrt{-c^2dx^2+d}} + \frac{3ax}{2c^4d\sqrt{-c^2dx^2+d}} - \frac{3a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2c^4d\sqrt{c^2d}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\sqrt{-d(c^2x^2-1)}}{4\sqrt{cx+1}\operatorname{arccosh}(cx)}$
parts	$-\frac{ax^3}{2c^2d\sqrt{-c^2dx^2+d}} + \frac{3ax}{2c^4d\sqrt{-c^2dx^2+d}} - \frac{3a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2c^4d\sqrt{c^2d}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\sqrt{-d(c^2x^2-1)}}{4\sqrt{cx+1}\operatorname{arccosh}(cx)}$

input `int(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)`

output 
$$-1/2*a*x^3/c^2/d/(-c^2*d*x^2+d)^(1/2)+3/2*a/c^4*x/d/(-c^2*d*x^2+d)^(1/2)-3/2*a/c^4/d/(c^2*d)^(1/2)*\arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+1/8*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-d*(c^2*x^2-1))^(1/2)*(4*(c*x+1)^(1/2)*\operatorname{arccosh}(c*x)*(c*x-1)^(1/2)*c^3*x^3-2*c^4*x^4+6*\operatorname{arccosh}(c*x)^2*x^2*c^2-12*(c*x+1)^(1/2)*\operatorname{arccosh}(c*x)*(c*x-1)^(1/2)*c*x-8*c^2*x^2*\operatorname{arccosh}(c*x)+8*\ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^2*c^2+3*c^2*x^2-6*\operatorname{arccosh}(c*x)^2+8*\operatorname{arccosh}(c*x)-8*\ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)-1)/(c^2*x^2-1)^2/d^2/c^5$$

3.115. 
$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$$



**3.115.5 Fricas [F]**

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral((b*x^4*arccosh(c*x) + a*x^4)*sqrt(-c^2*d*x^2 + d)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.115.6 Sympy [F]**

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x**4*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.115.7 Maxima [F]**

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-1/2*a*(x^3/(sqrt(-c^2*d*x^2 + d)*c^2*d) - 3*x/(sqrt(-c^2*d*x^2 + d)*c^4*d) + 3*arcsin(c*x)/(c^5*d^(3/2))) + b*integrate(x^4*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(-c^2*d*x^2 + d)^(3/2), x)`

**3.115.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.115.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2),x)`

output `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2), x)`

**3.116**  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$

3.116.1 Optimal result . . . . . 1058  
 3.116.2 Mathematica [A] (verified) . . . . . 1058  
 3.116.3 Rubi [A] (verified) . . . . . 1059  
 3.116.4 Maple [A] (verified) . . . . . 1060  
 3.116.5 Fricas [A] (verification not implemented) . . . . . 1061  
 3.116.6 Sympy [F] . . . . . 1062  
 3.116.7 Maxima [A] (verification not implemented) . . . . . 1062  
 3.116.8 Giac [F(-2)] . . . . . 1063  
 3.116.9 Mupad [F(-1)] . . . . . 1063

**3.116.1 Optimal result**

Integrand size = 27, antiderivative size = 150

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = -\frac{bx\sqrt{d - c^2dx^2}}{c^3d^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{a + \operatorname{arccosh}(cx)}{c^4d\sqrt{d - c^2dx^2}} + \frac{\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{c^4d^2} - \frac{b\sqrt{d - c^2dx^2}\operatorname{arctanh}(cx)}{c^4d^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
(a+b*arccosh(c*x))/c^4/d/(-c^2*d*x^2+d)^(1/2)+(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^4/d^2-b*x*(-c^2*d*x^2+d)^(1/2)/c^3/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*arctanh(c*x)*(-c^2*d*x^2+d)^(1/2)/c^4/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.116.2 Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.65

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \frac{2a - ac^2x^2 + bcx\sqrt{-1 + cx}\sqrt{1 + cx} + b(2 - c^2x^2) \operatorname{arccosh}(cx) + b\sqrt{-1 + cx}}{c^4d\sqrt{d - c^2dx^2}}$$

input

```
Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]
```

output  $(2*a - a*c^2*x^2 + b*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x] + b*(2 - c^2*x^2)*\text{ArcCosh}[c*x] + b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*\text{ArcTanh}[c*x])/(c^4*d*\text{Sqrt}[d - c^2*d*x^2])$

### 3.116.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 113, normalized size of antiderivative = 0.75, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 299, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$$

$$\downarrow 6337$$

$$-\frac{bc\sqrt{d - c^2 dx^2} \int \frac{2 - c^2 x^2}{c^4 d^2 (1 - c^2 x^2)} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{\sqrt{d - c^2 dx^2}(a + \text{barccosh}(cx))}{c^4 d^2} + \frac{a + \text{barccosh}(cx)}{c^4 d \sqrt{d - c^2 dx^2}}$$

$$\downarrow 27$$

$$-\frac{b\sqrt{d - c^2 dx^2} \int \frac{2 - c^2 x^2}{1 - c^2 x^2} dx}{c^3 d^2 \sqrt{cx - 1}\sqrt{cx + 1}} + \frac{\sqrt{d - c^2 dx^2}(a + \text{barccosh}(cx))}{c^4 d^2} + \frac{a + \text{barccosh}(cx)}{c^4 d \sqrt{d - c^2 dx^2}}$$

$$\downarrow 299$$

$$-\frac{b\sqrt{d - c^2 dx^2} \left( \int \frac{1}{1 - c^2 x^2} dx + x \right)}{c^3 d^2 \sqrt{cx - 1}\sqrt{cx + 1}} + \frac{\sqrt{d - c^2 dx^2}(a + \text{barccosh}(cx))}{c^4 d^2} + \frac{a + \text{barccosh}(cx)}{c^4 d \sqrt{d - c^2 dx^2}}$$

$$\downarrow 219$$

$$\frac{\sqrt{d - c^2 dx^2}(a + \text{barccosh}(cx))}{c^4 d^2} + \frac{a + \text{barccosh}(cx)}{c^4 d \sqrt{d - c^2 dx^2}} - \frac{b \left( \frac{\text{arctanh}(cx)}{c} + x \right) \sqrt{d - c^2 dx^2}}{c^3 d^2 \sqrt{cx - 1}\sqrt{cx + 1}}$$

input  $\text{Int}[(x^3*(a + b*\text{ArcCosh}[c*x]))/(d - c^2*d*x^2)^(3/2), x]$

output  $(a + b*\text{ArcCosh}[c*x])/(c^4*d*\text{Sqrt}[d - c^2*d*x^2]) + (\text{Sqrt}[d - c^2*d*x^2]*(a + b*\text{ArcCosh}[c*x]))/(c^4*d^2) - (b*\text{Sqrt}[d - c^2*d*x^2]*(x + \text{ArcTanh}[c*x]/c))/(c^3*d^2*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])$

## 3.116.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`
- rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

## 3.116.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.70

method	result
default	$a \left( -\frac{x^2}{c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{2}{d c^4 \sqrt{-c^2 d x^2 + d}} \right) + \frac{b \sqrt{c x - 1} \sqrt{c x + 1} \sqrt{-d(c^2 x^2 - 1)} (\sqrt{c x + 1} \operatorname{arccosh}(c x) \sqrt{c x - 1} c^2 x^2 - c^3 x^3 + \ln(\sqrt{c x - 1} \sqrt{c x + 1}))}{d c^4 \sqrt{-c^2 d x^2 + d}}$
parts	$a \left( -\frac{x^2}{c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{2}{d c^4 \sqrt{-c^2 d x^2 + d}} \right) + \frac{b \sqrt{c x - 1} \sqrt{c x + 1} \sqrt{-d(c^2 x^2 - 1)} (\sqrt{c x + 1} \operatorname{arccosh}(c x) \sqrt{c x - 1} c^2 x^2 - c^3 x^3 + \ln(\sqrt{c x - 1} \sqrt{c x + 1}))}{d c^4 \sqrt{-c^2 d x^2 + d}}$

input `int(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)`

3.116. 
$$\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$$

output  $a*(-x^2/c^2/d/(-c^2*d*x^2+d)^{(1/2)}+2/d/c^4/(-c^2*d*x^2+d)^{(1/2)})+b*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(-d*(c^2*x^2-1))^{(1/2)}*((c*x+1)^{(1/2)}*\operatorname{arccosh}(c*x)*(c*x-1)^{(1/2)}*c^2*x^2-c^3*x^3+\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)*x^2*c^2-2*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*x^2*c^2-2*\operatorname{arccosh}(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)+\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))/((c^2*x^2-1)^2/d^2/c^4$

### 3.116.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 429, normalized size of antiderivative = 2.86

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \left[ -\frac{4\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1}bcx - 4(bc^2 x^2 - 2b)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1})}{2\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1}bcx + (bc^2 x^2 - b)\sqrt{d} \arctan\left(\frac{2\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1}c\sqrt{dx}}{c^4 dx^4 - d}\right) - 2(bc^2 x^2 - 2b)\sqrt{-c^2 dx^2 + d}}{2(c^6 d^2 x^2 - c^4 d^2)} \right]$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="fracas")`

output  $[-1/4*(4*\sqrt{-c^2*d*x^2 + d})*\sqrt{c^2*x^2 - 1}*b*c*x - 4*(b*c^2*x^2 - 2*b)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) + (b*c^2*x^2 - b)*\sqrt{-d}*\log(-(c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 - 4*(c^3*x^3 + c*x))*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*\sqrt{-d} - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)) - 4*(a*c^2*x^2 - 2*a)*\sqrt{-c^2*d*x^2 + d}]/(c^6*d^2*x^2 - c^4*d^2), -1/2*(2*\sqrt{-c^2*d*x^2 + d})*\sqrt{c^2*x^2 - 1}*b*c*x + (b*c^2*x^2 - b)*\sqrt{d}*\arctan(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*c*\sqrt{d})*x/(c^4*d*x^4 - d) - 2*(b*c^2*x^2 - 2*b)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) - 2*(a*c^2*x^2 - 2*a)*\sqrt{-c^2*d*x^2 + d}]/(c^6*d^2*x^2 - c^4*d^2)]$

**3.116.6 Sympy [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x**3*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.116.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.05

$$\begin{aligned} \int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = & \\ & -\frac{1}{2}bc \left( \frac{2\sqrt{-d}x}{c^4 d^2} + \frac{\sqrt{-d} \log(cx + 1)}{c^5 d^2} - \frac{\sqrt{-d} \log(cx - 1)}{c^5 d^2} \right) \\ & - b \left( \frac{x^2}{\sqrt{-c^2 dx^2 + dc^2 d}} - \frac{2}{\sqrt{-c^2 dx^2 + dc^4 d}} \right) \operatorname{arcosh}(cx) \\ & - a \left( \frac{x^2}{\sqrt{-c^2 dx^2 + dc^2 d}} - \frac{2}{\sqrt{-c^2 dx^2 + dc^4 d}} \right) \end{aligned}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-1/2*b*c*(2*sqrt(-d)*x/(c^4*d^2) + sqrt(-d)*log(c*x + 1)/(c^5*d^2) - sqrt(-d)*log(c*x - 1)/(c^5*d^2)) - b*(x^2/(sqrt(-c^2*d*x^2 + d)*c^2*d) - 2/(sqrt(-c^2*d*x^2 + d)*c^4*d))*arccosh(c*x) - a*(x^2/(sqrt(-c^2*d*x^2 + d)*c^2*d) - 2/(sqrt(-c^2*d*x^2 + d)*c^4*d))`

**3.116.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.116.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2),x)`

output `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2), x)`



**3.117**  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$

3.117.1 Optimal result . . . . . 1064  
 3.117.2 Mathematica [A] (warning: unable to verify) . . . . . 1064  
 3.117.3 Rubi [A] (verified) . . . . . 1065  
 3.117.4 Maple [B] (verified) . . . . . 1067  
 3.117.5 Fricas [F] . . . . . 1067  
 3.117.6 Sympy [F] . . . . . 1068  
 3.117.7 Maxima [F] . . . . . 1068  
 3.117.8 Giac [F] . . . . . 1068  
 3.117.9 Mupad [F(-1)] . . . . . 1069

**3.117.1 Optimal result**

Integrand size = 27, antiderivative size = 143

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \frac{x(a + \operatorname{arccosh}(cx))}{c^2d\sqrt{d - c^2dx^2}} - \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{2bc^3d\sqrt{d - c^2dx^2}} - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx} \log(1 - c^2x^2)}{2c^3d\sqrt{d - c^2dx^2}}$$

output `x*(a+b*arccosh(c*x))/c^2/d/(-c^2*d*x^2+d)^(1/2)-1/2*(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^3/d/(-c^2*d*x^2+d)^(1/2)-1/2*b*ln(-c^2*x^2+1)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d/(-c^2*d*x^2+d)^(1/2)`

**3.117.2 Mathematica [A] (warning: unable to verify)**

Time = 0.61 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.11

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \frac{2acdx + 2a\sqrt{d}\sqrt{d - c^2dx^2} \arctan\left(\frac{cx\sqrt{d-c^2dx^2}}{\sqrt{d(-1+c^2x^2)}}\right) + bd\left(2cx\operatorname{arccosh}(cx) - \sqrt{\frac{-1}{1+c^2x^2}}\right)}{2c^3d^2\sqrt{d - c^2dx^2}}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]`

output  $(2*a*c*d*x + 2*a*\sqrt{d}*\sqrt{d - c^2*d*x^2}*\text{ArcTan}[(c*x*\sqrt{d - c^2*d*x^2})/(\sqrt{d}*(-1 + c^2*x^2))] + b*d*(2*c*x*\text{ArcCosh}[c*x] - \sqrt{(-1 + c*x)/(1 + c*x)}*(1 + c*x)*( \text{ArcCosh}[c*x]^2 + 2*\text{Log}[\sqrt{(-1 + c*x)/(1 + c*x)}]*(1 + c*x))))/(2*c^3*d^2*\sqrt{d - c^2*d*x^2})$

### 3.117.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6349, 25, 82, 240, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + \text{barccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6349} \\
 & -\frac{\int \frac{a + \text{barccosh}(cx)}{\sqrt{d - c^2dx^2}} dx}{c^2d} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int -\frac{x}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2dx^2}} + \frac{x(a + \text{barccosh}(cx))}{c^2d\sqrt{d - c^2dx^2}} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\int \frac{a + \text{barccosh}(cx)}{\sqrt{d - c^2dx^2}} dx}{c^2d} + \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2dx^2}} + \frac{x(a + \text{barccosh}(cx))}{c^2d\sqrt{d - c^2dx^2}} \\
 & \quad \downarrow \text{82} \\
 & -\frac{\int \frac{a + \text{barccosh}(cx)}{\sqrt{d - c^2dx^2}} dx}{c^2d} + \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x}{1 - c^2x^2} dx}{cd\sqrt{d - c^2dx^2}} + \frac{x(a + \text{barccosh}(cx))}{c^2d\sqrt{d - c^2dx^2}} \\
 & \quad \downarrow \text{240} \\
 & -\frac{\int \frac{a + \text{barccosh}(cx)}{\sqrt{d - c^2dx^2}} dx}{c^2d} + \frac{x(a + \text{barccosh}(cx))}{c^2d\sqrt{d - c^2dx^2}} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \log(1 - c^2x^2)}{2c^3d\sqrt{d - c^2dx^2}} \\
 & \quad \downarrow \text{6307} \\
 & \frac{x(a + \text{barccosh}(cx))}{c^2d\sqrt{d - c^2dx^2}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}(a + \text{barccosh}(cx))^2}{2bc^3d\sqrt{d - c^2dx^2}} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \log(1 - c^2x^2)}{2c^3d\sqrt{d - c^2dx^2}}
 \end{aligned}$$

input  $\text{Int}[(x^2*(a + b*\text{ArcCosh}[c*x]))/(d - c^2*d*x^2)^(3/2), x]$

---

3.117.  $\int \frac{x^2(a + \text{barccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx$

```
output (x*(a + b*ArcCosh[c*x]))/(c^2*d*Sqrt[d - c^2*d*x^2]) - (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(2*b*c^3*d*Sqrt[d - c^2*d*x^2]) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2])/(2*c^3*d*Sqrt[d - c^2*d*x^2])
```

### 3.117.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 82 Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]
```

```
rule 240 Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]
```

```
rule 6307 Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]
```

```
rule 6349 Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*(m - 1)/(2*e*(p + 1)) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.117.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 278 vs.  $2(125) = 250$ .

Time = 1.07 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.95

method	result
default	$\frac{ax}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{a \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{c^2 d \sqrt{c^2 d}} + \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^2}{2d^2 c^3 (c^2 x^2 - 1)} - \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1}}{d^2 c^3 (c^2 x^2 - 1)}$
parts	$\frac{ax}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{a \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{c^2 d \sqrt{c^2 d}} + \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^2}{2d^2 c^3 (c^2 x^2 - 1)} - \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1}}{d^2 c^3 (c^2 x^2 - 1)}$

input `int(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)`

output `a*x/c^2/d/(-c^2*d*x^2+d)^(1/2)-a/c^2/d/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+1/2*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/c^3/(c^2*x^2-1)*arccosh(c*x)^2-b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/c^3/(c^2*x^2-1)*arccosh(c*x)-b*(-d*(c^2*x^2-1))^(1/2)*arccosh(c*x)/d^2/c^2/(c^2*x^2-1)*x+b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/c^3/(c^2*x^2-1)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)`

### 3.117.5 Fracas [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*x^2*arccosh(c*x) + a*x^2)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.117.6 Sympy [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x**2*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.117.7 Maxima [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `a*(x/(sqrt(-c^2*d*x^2 + d)*c^2*d) - arcsin(c*x)/(c^3*d^(3/2))) + b*integrate(x^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(-c^2*d*x^2 + d)^(3/2), x)`

**3.117.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^2/(-c^2*d*x^2 + d)^(3/2), x)`

**3.117.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2),x)`output `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2), x)`

**3.118**  $\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$

3.118.1 Optimal result . . . . . 1070  
 3.118.2 Mathematica [A] (verified) . . . . . 1070  
 3.118.3 Rubi [A] (verified) . . . . . 1071  
 3.118.4 Maple [A] (verified) . . . . . 1072  
 3.118.5 Fricas [A] (verification not implemented) . . . . . 1073  
 3.118.6 Sympy [F] . . . . . 1073  
 3.118.7 Maxima [F] . . . . . 1074  
 3.118.8 Giac [F] . . . . . 1074  
 3.118.9 Mupad [F(-1)] . . . . . 1074

**3.118.1 Optimal result**

Integrand size = 25, antiderivative size = 76

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \frac{a + \operatorname{arccosh}(cx)}{c^2d\sqrt{d - c^2dx^2}} + \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arctanh}(cx)}{c^2d\sqrt{d - c^2dx^2}}$$

```
output (a+b*arccosh(c*x))/c^2/d/(-c^2*d*x^2+d)^(1/2)+b*arctanh(c*x)*(c*x-1)^(1/2)
*(c*x+1)^(1/2)/c^2/d/(-c^2*d*x^2+d)^(1/2)
```

**3.118.2 Mathematica [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.79

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = -\frac{a\sqrt{-d}(-1 + c^2x^2)}{c^2d^2(-1 + c^2x^2)} - \frac{b\sqrt{-d}(-1 + c^2x^2)\operatorname{arccosh}(cx)}{c^2d^2(-1 + c^2x^2)} + \frac{b\sqrt{d - c^2dx^2}(\log(-1 + cx) - \log(1 + cx))}{2c^2d^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
input Integrate[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]
```

```
output -((a*Sqrt[-(d*(-1 + c^2*x^2))])/(c^2*d^2*(-1 + c^2*x^2))) - (b*Sqrt[-(d*(-1 + c^2*x^2))]*ArcCosh[c*x])/(c^2*d^2*(-1 + c^2*x^2)) + (b*Sqrt[d - c^2*d*x^2]*(Log[-1 + c*x] - Log[1 + c*x]))/(2*c^2*d^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])
```

---

3.118.  $\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$

**3.118.3 Rubi [A] (verified)**

Time = 0.29 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {6329, 25, 39, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6329} \\
 & \frac{a + \operatorname{barccosh}(cx)}{c^2 d \sqrt{d - c^2 dx^2}} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int -\frac{1}{(1-cx)(cx+1)} dx}{cd\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{(1-cx)(cx+1)} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{c^2 d \sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{39} \\
 & \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{1-c^2 x^2} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{c^2 d \sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{219} \\
 & \frac{a + \operatorname{barccosh}(cx)}{c^2 d \sqrt{d - c^2 dx^2}} + \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \operatorname{arctanh}(cx)}{c^2 d \sqrt{d - c^2 dx^2}}
 \end{aligned}$$

input `Int[(x*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]`

output `(a + b*ArcCosh[c*x])/(c^2*d*Sqrt[d - c^2*d*x^2]) + (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*ArcTanh[c*x])/(c^2*d*Sqrt[d - c^2*d*x^2])`



## 3.118.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 39 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(m_)), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 6329 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n/(2*e*(p + 1)), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

## 3.118.4 Maple [A] (verified)

Time = 0.87 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.78

method	result
default	$\frac{a}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{b \sqrt{-d(c^2 x^2 - 1)} (\sqrt{cx-1} \sqrt{cx+1} \ln(1+cx+\sqrt{cx-1}\sqrt{cx+1}) - \sqrt{cx-1}\sqrt{cx+1} \ln(\sqrt{cx-1}\sqrt{cx+1}+cx-1) + \arccos(\frac{cx-1}{c^2 x^2 - 1}))}{d^2 c^2 (c^2 x^2 - 1)}$
parts	$\frac{a}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{b \sqrt{-d(c^2 x^2 - 1)} (\sqrt{cx-1} \sqrt{cx+1} \ln(1+cx+\sqrt{cx-1}\sqrt{cx+1}) - \sqrt{cx-1}\sqrt{cx+1} \ln(\sqrt{cx-1}\sqrt{cx+1}+cx-1) + \arccos(\frac{cx-1}{c^2 x^2 - 1}))}{d^2 c^2 (c^2 x^2 - 1)}$

input `int(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2), x, method=_RETURNVERBOSE)`

output `a/c^2/d/(-c^2*d*x^2+d)^(1/2)-b*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1)+arccosh(c*x))/d^2/c^2/(c^2*x^2-1)`

---

3.118. 
$$\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$$

**3.118.5 Fracas [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 327, normalized size of antiderivative = 4.30

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \frac{\left[ \frac{4\sqrt{-c^2 dx^2 + db} \log(cx + \sqrt{c^2 x^2 - 1}) + (bc^2 x^2 - b)\sqrt{-d} \log\left(-\frac{c^6 dx^6 + 5c^4 dx^4 - 5c^2 d x^2 - 4(c^3 x^3 + c^2 x^2 - 1)}{c^4 d^2 x^2 - c^2 d^2}\right)}{4(c^4 d^2 x^2 - c^2 d^2)} \right.}{(bc^2 x^2 - b)\sqrt{d} \arctan\left(\frac{2\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1}c\sqrt{dx}}{c^4 dx^4 - d}\right) + 2\sqrt{-c^2 dx^2 + db} \log(cx + \sqrt{c^2 x^2 - 1}) + 2\sqrt{-c^2 dx^2 + db}}{2(c^4 d^2 x^2 - c^2 d^2)}$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`output `[-1/4*(4*sqrt(-c^2*d*x^2 + d)*b*log(c*x + sqrt(c^2*x^2 - 1)) + (b*c^2*x^2 - b)*sqrt(-d)*log(-(c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 - 4*(c^3*x^3 + c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*sqrt(-d) - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)) + 4*sqrt(-c^2*d*x^2 + d)*a)/(c^4*d^2*x^2 - c^2*d^2), -1/2*((b*c^2*x^2 - b)*sqrt(d)*arctan(2*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*c*sqrt(d)*x/(c^4*d*x^4 - d)) + 2*sqrt(-c^2*d*x^2 + d)*b*log(c*x + sqrt(c^2*x^2 - 1)) + 2*sqrt(-c^2*d*x^2 + d)*a)/(c^4*d^2*x^2 - c^2*d^2)]`**3.118.6 Sympy [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(3/2),x)`output `Integral(x*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.118.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `b*(((c*sqrt(d)*x + sqrt(c*x + 1)*sqrt(c*x - 1)*sqrt(d))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/sqrt(-c*x + 1) + sqrt(c*x + 1)*sqrt(c*x - 1)*sqrt(d)/sqrt(-c*x + 1))/(sqrt(c*x + 1)*c^3*d^2*x + (c*x + 1)*sqrt(c*x - 1)*c^2*d^2) - integrate((c^2*x^3 + c*x^2*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1)) - x)/(sqrt(-c*x + 1)*((c^2*d^(3/2)*x^2 - d^(3/2))*e^(3/2*log(c*x + 1) + log(c*x - 1)) + 2*(c^3*d^(3/2)*x^3 - c*d^(3/2)*x)*e^(log(c*x + 1) + 1/2*log(c*x - 1)) + (c^4*d^(3/2)*x^4 - c^2*d^(3/2)*x^2)*sqrt(c*x + 1))), x) + a/(sqrt(-c^2*d*x^2 + d)*c^2*d)`

**3.118.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x/(-c^2*d*x^2 + d)^(3/2), x)`

**3.118.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2),x)`

output `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(3/2), x)`

---

3.118.  $\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$

$$3.119 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{(d-c^2dx^2)^{3/2}} dx$$

3.119.1 Optimal result . . . . .	1075
3.119.2 Mathematica [A] (verified) . . . . .	1075
3.119.3 Rubi [A] (verified) . . . . .	1076
3.119.4 Maple [B] (verified) . . . . .	1077
3.119.5 Fracas [F] . . . . .	1077
3.119.6 Sympy [F] . . . . .	1077
3.119.7 Maxima [A] (verification not implemented) . . . . .	1078
3.119.8 Giac [F] . . . . .	1078
3.119.9 Mupad [F(-1)] . . . . .	1078

### 3.119.1 Optimal result

Integrand size = 24, antiderivative size = 84

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d - c^2dx^2)^{3/2}} dx = \frac{x(a + \operatorname{arccosh}(cx))}{d\sqrt{d - c^2dx^2}} - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx} \log(1 - c^2x^2)}{2cd\sqrt{d - c^2dx^2}}$$

output `x*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(1/2)-1/2*b*ln(-c^2*x^2+1)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d/(-c^2*d*x^2+d)^(1/2)`

### 3.119.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.86

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d - c^2dx^2)^{3/2}} dx = \frac{2acx + 2bcx\operatorname{arccosh}(cx) - b\sqrt{-1 + cx}\sqrt{1 + cx} \log(1 - c^2x^2)}{2cd\sqrt{d - c^2dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^(3/2), x]`

output `(2*a*c*x + 2*b*c*x*ArcCosh[c*x] - b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2])/(2*c*d*Sqrt[d - c^2*d*x^2])`

**3.119.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$ , Rules used = {6314, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx$$

↓ 6314

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}}$$

↓ 240

$$\frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \log(1-c^2x^2)}{2cd\sqrt{d-c^2dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^(3/2),x]`

output `(x*(a + b*ArcCosh[c*x]))/(d*Sqrt[d - c^2*d*x^2]) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2])/(2*c*d*Sqrt[d - c^2*d*x^2])`

**3.119.3.1 Defintions of rubi rules used**

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

### 3.119.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(74) = 148.

Time = 0.93 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.14

method	result
default	$\frac{ax}{d\sqrt{-c^2dx^2+d}} - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}\operatorname{arccosh}(cx)}{d^2c(c^2x^2-1)} - \frac{b\sqrt{-d(c^2x^2-1)}\operatorname{arccosh}(cx)x}{d^2(c^2x^2-1)} + \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}}{d^2c(c^2x^2-1)}$
parts	$\frac{ax}{d\sqrt{-c^2dx^2+d}} - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}\operatorname{arccosh}(cx)}{d^2c(c^2x^2-1)} - \frac{b\sqrt{-d(c^2x^2-1)}\operatorname{arccosh}(cx)x}{d^2(c^2x^2-1)} + \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}}{d^2c(c^2x^2-1)}$

input `int((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)`

output `a/d*x/(-c^2*d*x^2+d)^(1/2)-b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/c/(c^2*x^2-1)*arccosh(c*x)-b*(-d*(c^2*x^2-1))^(1/2)*arccosh(c*x)/d^2/(c^2*x^2-1)*x+b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/c/(c^2*x^2-1)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)`

### 3.119.5 Fracas [F]

$$\int \frac{a + b\operatorname{arccosh}(cx)}{(d - c^2dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2dx^2 + d)^{3/2}} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

### 3.119.6 Sympy [F]

$$\int \frac{a + b\operatorname{arccosh}(cx)}{(d - c^2dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(-d(cx - 1)(cx + 1))^{3/2}} dx$$

input `integrate((a+b*acosh(c*x))/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

---

3.119. 
$$\int \frac{a+b\operatorname{arccosh}(cx)}{(d-c^2dx^2)^{3/2}} dx$$

**3.119.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.83

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx = -\frac{bc \sqrt{-\frac{1}{c^4 d}} \log(x^2 - \frac{1}{c^2})}{2d} + \frac{bx \operatorname{arcosh}(cx)}{\sqrt{-c^2 dx^2 + dd}} + \frac{ax}{\sqrt{-c^2 dx^2 + dd}}$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`output `-1/2*b*c*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2)/d + b*x*arccosh(c*x)/(sqrt(-c^2*d*x^2 + d)*d) + a*x/(sqrt(-c^2*d*x^2 + d)*d)`**3.119.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)/(-c^2*d*x^2 + d)^(3/2), x)`**3.119.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^(3/2),x)`output `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^(3/2), x)`

**3.120**  $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)^{3/2}} dx$

3.120.1 Optimal result . . . . . 1079  
 3.120.2 Mathematica [A] (warning: unable to verify) . . . . . 1080  
 3.120.3 Rubi [A] (verified) . . . . . 1080  
 3.120.4 Maple [B] (verified) . . . . . 1083  
 3.120.5 Fracas [F] . . . . . 1084  
 3.120.6 Sympy [F] . . . . . 1084  
 3.120.7 Maxima [F] . . . . . 1085  
 3.120.8 Giac [F] . . . . . 1085  
 3.120.9 Mupad [F(-1)] . . . . . 1085

**3.120.1 Optimal result**

Integrand size = 27, antiderivative size = 229

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x(d - c^2dx^2)^{3/2}} dx = \frac{a + b\operatorname{arccosh}(cx)}{d\sqrt{d - c^2dx^2}} + \frac{2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} + \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arctanh}(cx)}{d\sqrt{d - c^2dx^2}} - \frac{ib\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} + \frac{ib\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}}$$

output

```
(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(1/2)+2*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)+b*arctanh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)-I*b*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)+I*b*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)
```



**3.120.2 Mathematica [A] (warning: unable to verify)**

Time = 2.14 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.47

$$\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx = \frac{a\sqrt{d-c^2 dx^2}}{-1+c^2 x^2} - a\sqrt{d}\log(x) + a\sqrt{d}\log\left(d + \sqrt{d}\sqrt{d - c^2 dx^2}\right) + \frac{ibd\left(i\operatorname{arccosh}(cx) + \sqrt{\frac{-1+cx}{1+cx}}(1+cx)\operatorname{arccosh}(cx)\log(1-ie\right)}{}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^(3/2)),x]`output `-(((a*Sqrt[d - c^2*d*x^2])/(-1 + c^2*x^2) - a*Sqrt[d]*Log[x] + a*Sqrt[d]*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (I*b*d*(I*ArcCosh[c*x] + Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Cosh[ArcCosh[c*x]/2]] - I*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Sinh[ArcCosh[c*x]/2]] + Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, (-I)/E^ArcCosh[c*x]] - Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, I/E^ArcCosh[c*x]]))/Sqrt[d - c^2*d*x^2])/d^2`**3.120.3 Rubi [A] (verified)**Time = 0.73 (sec) , antiderivative size = 159, normalized size of antiderivative = 0.69, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6351, 25, 39, 219, 6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx$$

$$\downarrow 6351$$

$$\frac{\int \frac{a + \operatorname{arccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx}{d} - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{(1 - cx)(cx + 1)} dx}{d\sqrt{d - c^2 dx^2}} + \frac{a + \operatorname{arccosh}(cx)}{d\sqrt{d - c^2 dx^2}}$$

$$\downarrow 25$$

$$\frac{\int \frac{a + \operatorname{arccosh}(cx)}{x\sqrt{d - c^2 dx^2}} dx}{d} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{(1 - cx)(cx + 1)} dx}{d\sqrt{d - c^2 dx^2}} + \frac{a + \operatorname{arccosh}(cx)}{d\sqrt{d - c^2 dx^2}}$$

---

3.120.  $\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& \downarrow 39 \\
& \frac{\int \frac{a + b \operatorname{arccosh}(cx)}{x \sqrt{d - c^2 dx^2}} dx}{d} + \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} \int \frac{1}{1 - c^2 x^2} dx}{d \sqrt{d - c^2 dx^2}} + \frac{a + b \operatorname{arccosh}(cx)}{d \sqrt{d - c^2 dx^2}} \\
& \downarrow 219 \\
& \frac{\int \frac{a + b \operatorname{arccosh}(cx)}{x \sqrt{d - c^2 dx^2}} dx}{d} + \frac{a + b \operatorname{arccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \frac{b \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arctanh}(cx)}{d \sqrt{d - c^2 dx^2}} \\
& \downarrow 6361 \\
& \frac{\sqrt{cx - 1} \sqrt{cx + 1} \int \frac{a + b \operatorname{arccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \frac{a + b \operatorname{arccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \frac{b \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arctanh}(cx)}{d \sqrt{d - c^2 dx^2}} \\
& \downarrow 3042 \\
& \frac{\sqrt{cx - 1} \sqrt{cx + 1} \int (a + b \operatorname{arccosh}(cx)) \csc \left( i \operatorname{arccosh}(cx) + \frac{\pi}{2} \right) \operatorname{darccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \frac{a + b \operatorname{arccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \\
& \quad \frac{b \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arctanh}(cx)}{d \sqrt{d - c^2 dx^2}} \\
& \downarrow 4668 \\
& \frac{\sqrt{cx - 1} \sqrt{cx + 1} \left( -ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)}) \right)}{d \sqrt{d - c^2 dx^2}} \\
& \quad + \frac{a + b \operatorname{arccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \frac{b \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arctanh}(cx)}{d \sqrt{d - c^2 dx^2}} \\
& \downarrow 2715 \\
& \frac{\sqrt{cx - 1} \sqrt{cx + 1} \left( -ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} \right)}{d \sqrt{d - c^2 dx^2}} \\
& \quad + \frac{a + b \operatorname{arccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \frac{b \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arctanh}(cx)}{d \sqrt{d - c^2 dx^2}} \\
& \downarrow 2838 \\
& \frac{\sqrt{cx - 1} \sqrt{cx + 1} \left( 2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx)) - ib \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}) \right)}{d \sqrt{d - c^2 dx^2}} \\
& \quad + \frac{a + b \operatorname{arccosh}(cx)}{d \sqrt{d - c^2 dx^2}} + \frac{b \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arctanh}(cx)}{d \sqrt{d - c^2 dx^2}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^(3/2)),x]`

3.120.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx$

```
output (a + b*ArcCosh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (b*Sqrt[-1 + c*x]*Sqrt[1 +
c*x]*ArcTanh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]
*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^Ar
cCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(d*Sqrt[d - c^2*d*x^2])
```

### 3.120.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 39 Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(m_.), x_Symbol] := Int[(
a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (
IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))
```

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 2715 Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2838 Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4668 Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m, x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x)) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

```
rule 6351 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])
```

```
rule 6361 Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]
```

### 3.120.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 482 vs. 2(238) = 476.

Time = 1.22 (sec) , antiderivative size = 483, normalized size of antiderivative = 2.11

method	result
default	$\frac{a}{d\sqrt{-c^2dx^2+d}} - \frac{a \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{d^{\frac{3}{2}}} - \frac{b\sqrt{-d(c^2x^2-1)}(i \operatorname{arccosh}(cx) \ln(1-i(cx+\sqrt{cx-1}\sqrt{cx+1}))x^2c^2 - i \operatorname{arccosh}(cx))}{d^{\frac{3}{2}}}$
parts	$\frac{a}{d\sqrt{-c^2dx^2+d}} - \frac{a \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{d^{\frac{3}{2}}} - \frac{b\sqrt{-d(c^2x^2-1)}(i \operatorname{arccosh}(cx) \ln(1-i(cx+\sqrt{cx-1}\sqrt{cx+1}))x^2c^2 - i \operatorname{arccosh}(cx))}{d^{\frac{3}{2}}}$

```
input int((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

output  $a/d/(-c^2dx^2+d)^{(1/2)}-a/d^{(3/2)}*\ln((2*d+2*d^{(1/2)}*(-c^2*d*x^2+d)^{(1/2)})/x)-b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}*(I*\operatorname{arccosh}(c*x)*\ln(1-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*x^2*c^2-I*\operatorname{arccosh}(c*x)*\ln(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*x^2*c^2-I*\operatorname{dilog}(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*x^2*c^2+I*\operatorname{dilog}(1-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*x^2*c^2+2*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*x^2*c^2-\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)*x^2*c^2-I*\operatorname{arccosh}(c*x)*\ln(1-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))+I*\operatorname{arccosh}(c*x)*\ln(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))+\operatorname{arccosh}(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+I*\operatorname{dilog}(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))-I*\operatorname{dilog}(1-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))-\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1))/d^2/(c^2*x^2-1)$

### 3.120.5 Fracas [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2dx^2 + d)^{\frac{3}{2}}x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^4*d^2*x^5 - 2*c^2*d^2*x^3 + d^2*x), x)`

### 3.120.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))/x/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))/(x*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.120.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-a*(log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(3/2) - 1/(sqrt(-c^2*d*x^2 + d)*d)) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/((-c^2*d*x^2 + d)^(3/2)*x), x)`

**3.120.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(3/2)*x), x)`

**3.120.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^(3/2)),x)`

output `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^(3/2)), x)`

### 3.121 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d-c^2dx^2)^{3/2}} dx$

3.121.1 Optimal result . . . . .	1086
3.121.2 Mathematica [A] (verified) . . . . .	1086
3.121.3 Rubi [A] (verified) . . . . .	1087
3.121.4 Maple [A] (verified) . . . . .	1089
3.121.5 Fricas [F] . . . . .	1089
3.121.6 Sympy [F] . . . . .	1090
3.121.7 Maxima [A] (verification not implemented) . . . . .	1090
3.121.8 Giac [F] . . . . .	1090
3.121.9 Mupad [F(-1)] . . . . .	1091

#### 3.121.1 Optimal result

Integrand size = 27, antiderivative size = 158

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2(d - c^2dx^2)^{3/2}} dx = -\frac{a + \operatorname{arccosh}(cx)}{dx\sqrt{d - c^2dx^2}} + \frac{2c^2x(a + \operatorname{arccosh}(cx))}{d\sqrt{d - c^2dx^2}} + \frac{bc\sqrt{d - c^2dx^2} \log(x)}{d^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc\sqrt{d - c^2dx^2} \log(1 - c^2x^2)}{2d^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output (-a-b*arccosh(c*x))/d/x/(-c^2*d*x^2+d)^(1/2)+2*c^2*x*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(1/2)+b*c*ln(x)*(-c^2*d*x^2+d)^(1/2)/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/2*b*c*ln(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

#### 3.121.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.72

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2(d - c^2dx^2)^{3/2}} dx = \frac{-2a + 4ac^2x^2 + 2b(-1 + 2c^2x^2) \operatorname{arccosh}(cx) - 2bcx\sqrt{-1 + cx}\sqrt{1 + cx} \log(x) - b}{2dx\sqrt{d - c^2dx^2}}$$

```
input Integrate[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^(3/2)), x]
```

```
output (-2*a + 4*a*c^2*x^2 + 2*b*(-1 + 2*c^2*x^2)*ArcCosh[c*x] - 2*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[x] - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2])/(2*d*x*Sqrt[d - c^2*d*x^2])
```

**3.121.3 Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.77, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6337, 25, 27, 354, 86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6337} \\
 & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{1-2c^2 x^2}{d^2 x(1-c^2 x^2)} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{2c^2 x(a + \operatorname{barccosh}(cx))}{d\sqrt{d - c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{dx\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \frac{1-2c^2 x^2}{d^2 x(1-c^2 x^2)} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{2c^2 x(a + \operatorname{barccosh}(cx))}{d\sqrt{d - c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{dx\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \frac{1-2c^2 x^2}{x(1-c^2 x^2)} dx}{d^2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{2c^2 x(a + \operatorname{barccosh}(cx))}{d\sqrt{d - c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{dx\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{354} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \frac{1-2c^2 x^2}{x^2(1-c^2 x^2)} dx^2}{2d^2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{2c^2 x(a + \operatorname{barccosh}(cx))}{d\sqrt{d - c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{dx\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{86} \\
 & \frac{bc\sqrt{d - c^2 dx^2} \int \left(\frac{c^2}{c^2 x^2 - 1} + \frac{1}{x^2}\right) dx^2}{2d^2\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{2c^2 x(a + \operatorname{barccosh}(cx))}{d\sqrt{d - c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{dx\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2c^2 x(a + \operatorname{barccosh}(cx))}{d\sqrt{d - c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{dx\sqrt{d - c^2 dx^2}} + \frac{bc\sqrt{d - c^2 dx^2}(\log(1 - c^2 x^2) + \log(x^2))}{2d^2\sqrt{cx - 1}\sqrt{cx + 1}}
 \end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^(3/2)),x]`



output  $-\frac{(a + b \operatorname{ArcCosh}[c x])}{(d x \sqrt{d - c^2 d x^2})} + \frac{(2 c^2 x (a + b \operatorname{ArcCos} h[c x]))}{(d \sqrt{d - c^2 d x^2})} + \frac{(b c \sqrt{d - c^2 d x^2} (\operatorname{Log}[x^2] + \operatorname{Log}[1 - c^2 x^2]))}{(2 d^2 \sqrt{-1 + c x} \sqrt{1 + c x})}$

### 3.121.3.1 Defintions of rubi rules used

rule 25  $\operatorname{Int}[-(F x_), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F x, x], x]$

rule 27  $\operatorname{Int}[(a_*)(F x_), x\_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F x, x], x] /; \operatorname{FreeQ}[a, x] \&\& \operatorname{!MatchQ}[F x, (b_*)(G x_)] /; \operatorname{FreeQ}[b, x]$

rule 86  $\operatorname{Int}[(a_.) + (b_.)(x_)] * ((c_) + (d_.)(x_))^{(n_.)} * ((e_.) + (f_.)(x_))^{(p_.)}, x_] \rightarrow \operatorname{Int}[\operatorname{ExpandIntegrand}[(a + b x) * (c + d x)^n * (e + f x)^p, x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& ((\operatorname{ILtQ}[n, 0] \&\& \operatorname{ILtQ}[p, 0]) \|\operator\| \operatorname{EqQ}[p, 1] \|\operator\| (\operatorname{IGtQ}[p, 0] \&\& (\operatorname{!IntegerQ}[n] \|\operator\| \operatorname{LeQ}[9 p + 5(n + 2), 0] \|\operator\| \operatorname{GeQ}[n + p + 1, 0] \|\operator\| (\operatorname{GeQ}[n + p + 2, 0] \&\& \operatorname{RationalQ}[a, b, c, d, e, f])))$

rule 354  $\operatorname{Int}[(x_)^{(m_.)} * ((a_) + (b_.)(x_)^2)^{(p_.)} * ((c_) + (d_.)(x_)^2)^{(q_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[1/2 \operatorname{Subst}[\operatorname{Int}[x^{(m-1)/2} * (a + b x)^p * (c + d x)^q, x], x, x^2], x] /; \operatorname{FreeQ}[\{a, b, c, d, p, q\}, x] \&\& \operatorname{NeQ}[b * c - a * d, 0] \&\& \operatorname{IntegerQ}[(m - 1)/2]$

rule 2009  $\operatorname{Int}[u_, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$

rule 6337  $\operatorname{Int}[(a_.) + \operatorname{ArcCosh}[(c_.)(x_)] * (b_.)] * (x_)^{(m_.)} * ((d_) + (e_.)(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{u = \operatorname{IntHide}[x^m * (d + e x^2)^p, x]\}, \operatorname{Simp}[(a + b \operatorname{ArcCosh}[c x]) u, x] - \operatorname{Simp}[b * c * \operatorname{Simp}[\operatorname{Sqrt}[d + e x^2] / (\operatorname{Sqrt}[1 + c x] * \operatorname{Sqrt}[-1 + c x]), x], x] \operatorname{Int}[\operatorname{SimplifyIntegrand}[u / \operatorname{Sqrt}[d + e x^2], x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e\}, x] \&\& \operatorname{EqQ}[c^2 d + e, 0] \&\& \operatorname{IntegerQ}[p - 1/2] \&\& \operatorname{NeQ}[p, -2^{(-1)}] \&\& (\operatorname{IGtQ}[(m + 1)/2, 0] \|\operator\| \operatorname{ILtQ}[(m + 2 * p + 3)/2, 0])$

**3.121.4 Maple [A] (verified)**

Time = 1.22 (sec) , antiderivative size = 266, normalized size of antiderivative = 1.68

method	result
default	$a \left( -\frac{1}{dx\sqrt{-c^2dx^2+d}} + \frac{2c^2x}{d\sqrt{-c^2dx^2+d}} \right) - \frac{b \left( -2\sqrt{cx-1}\sqrt{cx+1} \ln \left( (cx+\sqrt{cx-1}\sqrt{cx+1})^4 - 1 \right) x^3 c^3 + 2 \ln \left( (cx+\sqrt{cx-1}\sqrt{cx+1}) \right) \right)}{d^2 \sqrt{-c^2dx^2+d}}$
parts	$a \left( -\frac{1}{dx\sqrt{-c^2dx^2+d}} + \frac{2c^2x}{d\sqrt{-c^2dx^2+d}} \right) - \frac{b \left( -2\sqrt{cx-1}\sqrt{cx+1} \ln \left( (cx+\sqrt{cx-1}\sqrt{cx+1})^4 - 1 \right) x^3 c^3 + 2 \ln \left( (cx+\sqrt{cx-1}\sqrt{cx+1}) \right) \right)}{d^2 \sqrt{-c^2dx^2+d}}$

```
input int((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output a*(-1/d/x/(-c^2*d*x^2+d)^(1/2)+2*c^2/d*x/(-c^2*d*x^2+d)^(1/2))-b*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^4-1)*x^3*c^3+2*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^4-1)*x^4*c^4+(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^4-1)*x*c-2*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^4-1)*x^2*c^2+arccosh(c*x))*(2*c^2*x^2-1+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x)*(-d*(c^2*x^2-1))^(1/2)/d^2/(c^2*x^2-1)/x
```

**3.121.5 Fracas [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(-c^2 dx^2 + d)^{3/2} x^2} dx$$

```
input integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fracas")
```

```
output integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2), x)
```

**3.121.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{arcosh}(cx)}{x^2 (-d (cx - 1) (cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))/x**2/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))/(x**2*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.121.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.91

$$\begin{aligned} \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{3/2}} dx &= \frac{1}{2} bc \left( \frac{\sqrt{-d} \log(cx + 1)}{d^2} + \frac{\sqrt{-d} \log(cx - 1)}{d^2} + \frac{2\sqrt{-d} \log(x)}{d^2} \right) \\ &+ \left( \frac{2c^2 x}{\sqrt{-c^2 dx^2 + dd}} - \frac{1}{\sqrt{-c^2 dx^2 + ddx}} \right) b \operatorname{arcosh}(cx) \\ &+ \left( \frac{2c^2 x}{\sqrt{-c^2 dx^2 + dd}} - \frac{1}{\sqrt{-c^2 dx^2 + ddx}} \right) a \end{aligned}$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `1/2*b*c*(sqrt(-d)*log(c*x + 1)/d^2 + sqrt(-d)*log(c*x - 1)/d^2 + 2*sqrt(-d)*log(x)/d^2) + (2*c^2*x/(sqrt(-c^2*d*x^2 + d)*d) - 1/(sqrt(-c^2*d*x^2 + d)*d*x))*b*arccosh(c*x) + (2*c^2*x/(sqrt(-c^2*d*x^2 + d)*d) - 1/(sqrt(-c^2*d*x^2 + d)*d*x))*a`

**3.121.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(3/2)*x^2), x)`

**3.121.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2 (d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^(3/2)),x)`output `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^(3/2)), x)`

# 3.122 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^{3/2}} dx$

3.122.1 Optimal result . . . . .	1092
3.122.2 Mathematica [A] (warning: unable to verify) . . . . .	1093
3.122.3 Rubi [A] (verified) . . . . .	1093
3.122.4 Maple [A] (verified) . . . . .	1098
3.122.5 Fracas [F] . . . . .	1099
3.122.6 Sympy [F] . . . . .	1099
3.122.7 Maxima [F] . . . . .	1099
3.122.8 Giac [F] . . . . .	1100
3.122.9 Mupad [F(-1)] . . . . .	1100

## 3.122.1 Optimal result

Integrand size = 27, antiderivative size = 329

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^3(d - c^2dx^2)^{3/2}} dx = \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{2dx\sqrt{d - c^2dx^2}} + \frac{3c^2(a + \operatorname{arccosh}(cx))}{2d\sqrt{d - c^2dx^2}} - \frac{a + \operatorname{arccosh}(cx)}{2dx^2\sqrt{d - c^2dx^2}} + \frac{3c^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} + \frac{bc^2\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arctanh}(cx)}{d\sqrt{d - c^2dx^2}} - \frac{3ibc^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{2d\sqrt{d - c^2dx^2}} + \frac{3ibc^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{2d\sqrt{d - c^2dx^2}}$$

output

```
3/2*c^2*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(1/2)+1/2*(-a-b*arccosh(c*x))/d/x^2/(-c^2*d*x^2+d)^(1/2)+1/2*b*c*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/x/(-c^2*d*x^2+d)^(1/2)+3*c^2*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)+b*c^2*arctanh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)-3/2*I*b*c^2*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)+3/2*I*b*c^2*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)
```

**3.122.2 Mathematica [A] (warning: unable to verify)**

Time = 3.68 (sec) , antiderivative size = 439, normalized size of antiderivative = 1.33

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx = \frac{1}{2} \left( -\frac{a(-1 + 3c^2 x^2) \sqrt{d - c^2 dx^2}}{d^2 x^2 (-1 + c^2 x^2)} \right. \\ \left. + \frac{3ac^2 \log(x)}{d^{3/2}} - \frac{3ac^2 \log\left(d + \sqrt{d} \sqrt{d - c^2 dx^2}\right)}{d^{3/2}} \right) \\ - bc^2 \left( -\frac{\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}{cx} + \left(-1 + \frac{1}{c^2 x^2}\right) \operatorname{arccosh}(cx) - 2 \operatorname{arccosh}(cx) \cosh^2\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) + 3i \sqrt{\frac{-1+cx}{1+cx}}(1+cx) \right)$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^(3/2)),x]`

output `((-((a*(-1 + 3*c^2*x^2)*Sqrt[d - c^2*d*x^2])/(d^2*x^2*(-1 + c^2*x^2))) + (3*a*c^2*Log[x])/d^(3/2) - (3*a*c^2*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]])/d^(3/2) - (b*c^2*(-((Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))/(c*x)) + (-1 + 1/(c^2*x^2))*ArcCosh[c*x] - 2*ArcCosh[c*x]*Cosh[ArcCosh[c*x]/2]^2 + (3*I)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - (3*I)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] - 2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Cosh[ArcCosh[c*x]/2]] + 2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Sinh[ArcCosh[c*x]/2]] + (3*I)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, (-I)/E^ArcCosh[c*x]] - (3*I)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, I/E^ArcCosh[c*x]] + 2*ArcCosh[c*x]*Sinh[ArcCosh[c*x]/2]^2))/(d*Sqrt[d - c^2*d*x^2]))/2`

**3.122.3 Rubi [A] (verified)**Time = 1.11 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.77, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.519$ , Rules used = {6347, 25, 82, 264, 219, 6351, 25, 39, 219, 6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.122.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
& \int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx \\
& \quad \downarrow \text{6347} \\
& \frac{3}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x (d - c^2 dx^2)^{3/2}} dx + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{1}{x^2(1-cx)(cx+1)} dx}{2d\sqrt{d-c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2 dx^2}} \\
& \quad \downarrow \text{25} \\
& \frac{3}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x (d - c^2 dx^2)^{3/2}} dx - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{x^2(1-cx)(cx+1)} dx}{2d\sqrt{d-c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2 dx^2}} \\
& \quad \downarrow \text{82} \\
& \frac{3}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x (d - c^2 dx^2)^{3/2}} dx - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{x^2(1-c^2 x^2)} dx}{2d\sqrt{d-c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2 dx^2}} \\
& \quad \downarrow \text{264} \\
& \frac{3}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x (d - c^2 dx^2)^{3/2}} dx - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{1}{1-c^2 x^2} dx - \frac{1}{x} \right)}{2d\sqrt{d-c^2 dx^2}} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2 dx^2}} \\
& \quad \downarrow \text{219} \\
& \frac{3}{2} c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x (d - c^2 dx^2)^{3/2}} dx - \frac{a + \operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2 dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} (\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2 dx^2}} \\
& \quad \downarrow \text{6351} \\
& \frac{3}{2} c^2 \left( \frac{\int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{d-c^2 dx^2}} dx}{d} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{1}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{d\sqrt{d-c^2 dx^2}} \right) - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2 dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} (\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2 dx^2}} \\
& \quad \downarrow \text{25} \\
& \frac{3}{2} c^2 \left( \frac{\int \frac{a + \operatorname{barccosh}(cx)}{x\sqrt{d-c^2 dx^2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{d\sqrt{d-c^2 dx^2}} \right) - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2 dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} (\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2 dx^2}} \\
& \quad \downarrow \text{39}
\end{aligned}$$

$$\begin{aligned}
& \frac{3}{2}c^2 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} \right) - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}(\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{219} \\
& \frac{3}{2}c^2 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}} \right) - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}(\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6361} \\
& \frac{3}{2}c^2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{cx} \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}} \right) - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}(\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{3042} \\
& \frac{3}{2}c^2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \csc\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}} \right) - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}(\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{4668} \\
& \frac{3}{2}c^2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(-ib \int \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + ib \int \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2 \operatorname{arctanh}(cx))}{d\sqrt{d-c^2dx^2}} \right) - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}(\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{2715} \\
& \frac{3}{2}c^2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(-ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2 \operatorname{arctanh}(cx))}{d\sqrt{d-c^2dx^2}} \right) - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}(\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{2838}
\end{aligned}$$



$$\frac{3}{2}c^2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(2\arctan(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) - ib\operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) + ib\operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{a+\operatorname{barccosh}(cx)}{2dx^2\sqrt{d-c^2dx^2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}(\operatorname{carctanh}(cx) - \frac{1}{x})}{2d\sqrt{d-c^2dx^2}} \right)$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^(3/2)), x]`

output `-1/2*(a + b*ArcCosh[c*x])/(d*x^2*Sqrt[d - c^2*d*x^2]) - (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-x^(-1) + c*ArcTanh[c*x]))/(2*d*Sqrt[d - c^2*d*x^2]) + (3*c^2*((a + b*ArcCosh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*ArcTanh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]])))/(d*Sqrt[d - c^2*d*x^2]))) / 2`

### 3.122.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 39 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.)), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 264 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^2)^(p + 1)/(a*c*(m + 1))), x] - Simp[b*((m + 2*p + 3)/(a*c^2*(m + 1)) Int[(c*x)^(m + 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && LtQ[m, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((f_.)*(x_))^m_.)*((d_) + (e_
.)*(x_)^2)^p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1
))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp
[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(
f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^
(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && ILtQ[m, -1]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((f_.)*(x_))^m_.)*((d_) + (e_
.)*(x_)^2)^p_.), x_Symbol] :> Simp[(-f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1
)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[
b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[
(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])
^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] &
& GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] ||
EqQ[n, 1])`

```
rule 6361 Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_))/Sqrt[(d_) + (e_.)
*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]
]/Sqrt[d + e*x^2])] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]]
, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && Int
egerQ[m]
```

### 3.122.4 Maple [A] (verified)

Time = 1.29 (sec) , antiderivative size = 588, normalized size of antiderivative = 1.79

method	result
default	$a \left( -\frac{1}{2dx^2\sqrt{-c^2dx^2+d}} + \frac{3c^2 \left( \frac{1}{d\sqrt{-c^2dx^2+d}} - \frac{\ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{d^{\frac{3}{2}}}\right)}{2} \right) + b \left( -\frac{\sqrt{-d(c^2x^2-1)}(3c^2x^2 \operatorname{arccosh}(cx) + \sqrt{c^2x^2-1})}{2d^2(c^2x^2-1)} \right)$
parts	$a \left( -\frac{1}{2dx^2\sqrt{-c^2dx^2+d}} + \frac{3c^2 \left( \frac{1}{d\sqrt{-c^2dx^2+d}} - \frac{\ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{d^{\frac{3}{2}}}\right)}{2} \right) + b \left( -\frac{\sqrt{-d(c^2x^2-1)}(3c^2x^2 \operatorname{arccosh}(cx) + \sqrt{c^2x^2-1})}{2d^2(c^2x^2-1)} \right)$

```
input int((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output a*(-1/2/d/x^2/(-c^2*d*x^2+d)^(1/2)+3/2*c^2*(1/d/(-c^2*d*x^2+d)^(1/2)-1/d^(
3/2)*ln((2*d+2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x))+b*(-1/2*(-d*(c^2*x^2-1))
^(1/2)*(3*c^2*x^2*arccosh(c*x)+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-arccosh(c*x
))/d^2/(c^2*x^2-1)/x^2-(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/
(c^2*x^2-1)/d^2*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*c^2+(-d*(c^2*x^2-1))
^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(c^2*x^2-1)/d^2*ln((c*x-1)^(1/2)*(c*x+1
)^(1/2)+c*x-1)*c^2+3/2*I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2
)/(c^2*x^2-1)/d^2*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2+3/2*I*(
-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(c^2*x^2-1)/d^2*arccosh(
c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*c^2-3/2*I*(-d*(c^2*x^2-1))^(
1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(c^2*x^2-1)/d^2*dilog(1-I*(c*x+(c*x-1)^(
1/2)*(c*x+1)^(1/2)))*c^2-3/2*I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1
)^(1/2)/(c^2*x^2-1)/d^2*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/
2)))*c^2)
```

$$3.122. \int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^{3/2}} dx$$

**3.122.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^4*d^2*x^7 - 2*c^2*d^2*x^5 + d^2*x^3), x)`

**3.122.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))/x**3/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))/(x**3*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.122.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-1/2*(3*c^2*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(3/2) - 3*c^2/(sqrt(-c^2*d*x^2 + d)*d) + 1/(sqrt(-c^2*d*x^2 + d)*d*x^2))*a + b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/((-c^2*d*x^2 + d)^(3/2)*x^3), x)`

**3.122.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(3/2)*x^3), x)`

**3.122.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^(3/2)),x)`

output `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^(3/2)), x)`

### 3.123 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^4(d-c^2dx^2)^{3/2}} dx$

3.123.1 Optimal result . . . . .	1101
3.123.2 Mathematica [A] (verified) . . . . .	1102
3.123.3 Rubi [A] (verified) . . . . .	1102
3.123.4 Maple [A] (verified) . . . . .	1104
3.123.5 Fricas [F] . . . . .	1105
3.123.6 Sympy [F] . . . . .	1105
3.123.7 Maxima [F] . . . . .	1105
3.123.8 Giac [F] . . . . .	1106
3.123.9 Mupad [F(-1)] . . . . .	1106

#### 3.123.1 Optimal result

Integrand size = 27, antiderivative size = 250

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^4(d - c^2dx^2)^{3/2}} dx = -\frac{bc\sqrt{d - c^2dx^2}}{6d^2x^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{a + \operatorname{arccosh}(cx)}{3dx^3\sqrt{d - c^2dx^2}} - \frac{4c^2(a + \operatorname{arccosh}(cx))}{3dx\sqrt{d - c^2dx^2}} + \frac{8c^4x(a + \operatorname{arccosh}(cx))}{3d\sqrt{d - c^2dx^2}} + \frac{5bc^3\sqrt{d - c^2dx^2} \log(x)}{3d^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3\sqrt{d - c^2dx^2} \log(1 - c^2x^2)}{2d^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output 1/3*(-a-b*arccosh(c*x))/d/x^3/(-c^2*d*x^2+d)^(1/2)-4/3*c^2*(a+b*arccosh(c*x))/d/x/(-c^2*d*x^2+d)^(1/2)+8/3*c^4*x*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(1/2)-1/6*b*c*(-c^2*d*x^2+d)^(1/2)/d^2/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/3*b*c^3*ln(x)*(-c^2*d*x^2+d)^(1/2)/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/2*b*c^3*ln(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/d^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.123.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 161, normalized size of antiderivative = 0.64

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx = \frac{-2a - 8ac^2 x^2 + 16ac^4 x^4 + bcx\sqrt{-1 + cx}\sqrt{1 + cx} + 2b(-1 - 4c^2 x^2 + 8c^4 x^4) \operatorname{arccosh}(cx)}{6dx^3 \sqrt{d - c^2 dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^(3/2)),x]`

output `(-2*a - 8*a*c^2*x^2 + 16*a*c^4*x^4 + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*b*(-1 - 4*c^2*x^2 + 8*c^4*x^4)*ArcCosh[c*x] - 10*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[x] - 3*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2])/(6*d*x^3*Sqrt[d - c^2*d*x^2])`

### 3.123.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 177, normalized size of antiderivative = 0.71, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6337, 27, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx \\ & \quad \downarrow \text{6337} \\ & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{8c^4 x^4 + 4c^2 x^2 + 1}{3d^2 x^3 (1 - c^2 x^2)} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{4c^2(a + b \operatorname{arccosh}(cx))}{3dx\sqrt{d - c^2 dx^2}} - \frac{a + b \operatorname{arccosh}(cx)}{3dx^3\sqrt{d - c^2 dx^2}} + \\ & \quad \frac{8c^4 x(a + b \operatorname{arccosh}(cx))}{3d\sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{27} \\ & \frac{bc\sqrt{d - c^2 dx^2} \int -\frac{8c^4 x^4 + 4c^2 x^2 + 1}{x^3(1 - c^2 x^2)} dx}{3d^2\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{4c^2(a + b \operatorname{arccosh}(cx))}{3dx\sqrt{d - c^2 dx^2}} - \frac{a + b \operatorname{arccosh}(cx)}{3dx^3\sqrt{d - c^2 dx^2}} + \\ & \quad \frac{8c^4 x(a + b \operatorname{arccosh}(cx))}{3d\sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{1578} \end{aligned}$$

---

3.123.  $\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& \frac{bc\sqrt{d-c^2dx^2} \int \frac{-8c^4x^4+4c^2x^2+1}{x^4(1-c^2x^2)} dx^2}{6d^2\sqrt{cx-1}\sqrt{cx+1}} - \frac{4c^2(a+\operatorname{barccosh}(cx))}{3dx\sqrt{d-c^2dx^2}} - \frac{a+\operatorname{barccosh}(cx)}{3dx^3\sqrt{d-c^2dx^2}} + \\
& \quad \frac{8c^4x(a+\operatorname{barccosh}(cx))}{3d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{1195} \\
& \frac{bc\sqrt{d-c^2dx^2} \int \left(\frac{3c^4}{c^2x^2-1} + \frac{5c^2}{x^2} + \frac{1}{x^4}\right) dx^2}{6d^2\sqrt{cx-1}\sqrt{cx+1}} - \frac{4c^2(a+\operatorname{barccosh}(cx))}{3dx\sqrt{d-c^2dx^2}} - \frac{a+\operatorname{barccosh}(cx)}{3dx^3\sqrt{d-c^2dx^2}} + \\
& \quad \frac{8c^4x(a+\operatorname{barccosh}(cx))}{3d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{2009} \\
& -\frac{4c^2(a+\operatorname{barccosh}(cx))}{3dx\sqrt{d-c^2dx^2}} - \frac{a+\operatorname{barccosh}(cx)}{3dx^3\sqrt{d-c^2dx^2}} + \frac{8c^4x(a+\operatorname{barccosh}(cx))}{3d\sqrt{d-c^2dx^2}} + \\
& \quad \frac{bc\sqrt{d-c^2dx^2} \left(5c^2 \log(x^2) + 3c^2 \log(1-c^2x^2) - \frac{1}{x^2}\right)}{6d^2\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^(3/2)),x]`

output `-1/3*(a + b*ArcCosh[c*x])/(d*x^3*Sqrt[d - c^2*d*x^2]) - (4*c^2*(a + b*ArcCosh[c*x]))/(3*d*x*Sqrt[d - c^2*d*x^2]) + (8*c^4*x*(a + b*ArcCosh[c*x]))/(3*d*Sqrt[d - c^2*d*x^2]) + (b*c*Sqrt[d - c^2*d*x^2]*(-x^(-2) + 5*c^2*Log[x^2] + 3*c^2*Log[1 - c^2*x^2]))/(6*d^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.123.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1195 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`



rule 1578 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])]`

### 3.123.4 Maple [A] (verified)

Time = 1.30 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.47

method	result
default	$a \left( -\frac{1}{3dx^3\sqrt{-c^2dx^2+d}} + \frac{4c^2 \left( -\frac{1}{dx\sqrt{-c^2dx^2+d}} + \frac{2c^2x}{d\sqrt{-c^2dx^2+d}} \right)}{3} \right) - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \left( 16\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx) \right)}{3}$
parts	$a \left( -\frac{1}{3dx^3\sqrt{-c^2dx^2+d}} + \frac{4c^2 \left( -\frac{1}{dx\sqrt{-c^2dx^2+d}} + \frac{2c^2x}{d\sqrt{-c^2dx^2+d}} \right)}{3} \right) - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \left( 16\sqrt{cx-1}\sqrt{cx+1} \operatorname{arccosh}(cx) \right)}{3}$

input `int((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)`

output `a*(-1/3/d/x^3/(-c^2*d*x^2+d)^(1/2)+4/3*c^2*(-1/d/x/(-c^2*d*x^2+d)^(1/2)+2*c^2/d*x/(-c^2*d*x^2+d)^(1/2))-1/6*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(16*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4+16*arccosh(c*x)*c^5*x^5-6*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^5*c^5-10*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^5*c^5-8*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-16*c^3*x^3*arccosh(c*x)+6*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^3*c^3+10*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3+c^3*x^3-2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-c*x)/d^2/(c^4*x^4-2*c^2*x^2+1)/x^3`

**3.123.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^4*d^2*x^8 - 2*c^2*d^2*x^6 + d^2*x^4), x)`

**3.123.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4 (-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))/x**4/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))/(x**4*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.123.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `1/3*(8*c^4*x/(sqrt(-c^2*d*x^2 + d)*d) - 4*c^2/(sqrt(-c^2*d*x^2 + d)*d*x) - 1/(sqrt(-c^2*d*x^2 + d)*d*x^3))*a + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(3/2)*x^4), x)`

**3.123.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(3/2)*x^4), x)`

**3.123.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4 (d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^(3/2)),x)`

output `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^(3/2)), x)`

**3.124**  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

3.124.1 Optimal result . . . . . 1107  
 3.124.2 Mathematica [A] (verified) . . . . . 1108  
 3.124.3 Rubi [A] (verified) . . . . . 1108  
 3.124.4 Maple [A] (verified) . . . . . 1110  
 3.124.5 Fricas [A] (verification not implemented) . . . . . 1111  
 3.124.6 Sympy [F] . . . . . 1112  
 3.124.7 Maxima [F] . . . . . 1112  
 3.124.8 Giac [F(-2)] . . . . . 1113  
 3.124.9 Mupad [F(-1)] . . . . . 1113

**3.124.1 Optimal result**

Integrand size = 27, antiderivative size = 243

$$\int \frac{x^5(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{bx\sqrt{d - c^2dx^2}}{c^5d^3\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bx\sqrt{d - c^2dx^2}}{6c^5d^3\sqrt{-1 + cx}\sqrt{1 + cx}(1 - c^2x^2)} + \frac{a + \operatorname{arccosh}(cx)}{3c^6d(d - c^2dx^2)^{3/2}} - \frac{2(a + \operatorname{arccosh}(cx))}{c^6d^2\sqrt{d - c^2dx^2}} - \frac{\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))}{c^6d^3} + \frac{11b\sqrt{d - c^2dx^2}\operatorname{arctanh}(cx)}{6c^6d^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output 1/3*(a+b*arccosh(c*x))/c^6/d/(-c^2*d*x^2+d)^(3/2)-2*(a+b*arccosh(c*x))/c^6/d^2/(-c^2*d*x^2+d)^(1/2)-(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c^6/d^3+b*x*(-c^2*d*x^2+d)^(1/2)/c^5/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/6*b*x*(-c^2*d*x^2+d)^(1/2)/c^5/d^3/(-c^2*x^2+1)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+11/6*b*arctanh(c*x)*(-c^2*d*x^2+d)^(1/2)/c^6/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.124.2 Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 167, normalized size of antiderivative = 0.69

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \frac{16a - 24ac^2 x^2 + 6ac^4 x^4 + 5bcx\sqrt{-1 + cx}\sqrt{1 + cx} - 6bc^3 x^3 \sqrt{-1 + cx}\sqrt{1 + cx}}{6c^6 d^2 (-1 + cx)^{3/2} (1 + cx)^{3/2}}$$

input `Integrate[(x^5*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2), x]`output `(16*a - 24*a*c^2*x^2 + 6*a*c^4*x^4 + 5*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - 6*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*b*(8 - 12*c^2*x^2 + 3*c^4*x^4)*ArcCosh[c*x] - 11*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1 + c^2*x^2)*ArcTan[h[c*x]])/(6*c^6*d^2*(-1 + c^2*x^2)*Sqrt[d - c^2*d*x^2])`**3.124.3 Rubi [A] (verified)**Time = 0.49 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6337, 27, 1471, 25, 299, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx \\ & \quad \downarrow \text{6337} \\ & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{3c^4 x^4 - 12c^2 x^2 + 8}{3c^6 d^3 (1 - c^2 x^2)^2} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{c^6 d^3} - \frac{2(a + \operatorname{barccosh}(cx))}{c^6 d^2 \sqrt{d - c^2 dx^2}} + \\ & \quad \frac{a + \operatorname{barccosh}(cx)}{3c^6 d (d - c^2 dx^2)^{3/2}} \\ & \quad \downarrow \text{27} \\ & -\frac{b\sqrt{d - c^2 dx^2} \int \frac{3c^4 x^4 - 12c^2 x^2 + 8}{(1 - c^2 x^2)^2} dx}{3c^5 d^3 \sqrt{cx - 1}\sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{c^6 d^3} - \frac{2(a + \operatorname{barccosh}(cx))}{c^6 d^2 \sqrt{d - c^2 dx^2}} + \\ & \quad \frac{a + \operatorname{barccosh}(cx)}{3c^6 d (d - c^2 dx^2)^{3/2}} \\ & \quad \downarrow \text{1471} \end{aligned}$$

---

3.124.  $\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{b\sqrt{d-c^2dx^2}\left(-\frac{1}{2}\int-\frac{17-6c^2x^2}{1-c^2x^2}dx-\frac{x}{2(1-c^2x^2)}\right)-\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{c^6d^3}}{3c^5d^3\sqrt{cx-1}\sqrt{cx+1}+\frac{2(a+\operatorname{barccosh}(cx))}{c^6d^2\sqrt{d-c^2dx^2}}+\frac{a+\operatorname{barccosh}(cx)}{3c^6d(d-c^2dx^2)^{3/2}}} \\
& \quad \downarrow 25 \\
& \frac{b\sqrt{d-c^2dx^2}\left(\frac{1}{2}\int\frac{17-6c^2x^2}{1-c^2x^2}dx-\frac{x}{2(1-c^2x^2)}\right)-\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{c^6d^3}}{3c^5d^3\sqrt{cx-1}\sqrt{cx+1}+\frac{2(a+\operatorname{barccosh}(cx))}{c^6d^2\sqrt{d-c^2dx^2}}+\frac{a+\operatorname{barccosh}(cx)}{3c^6d(d-c^2dx^2)^{3/2}}} \\
& \quad \downarrow 299 \\
& \frac{b\sqrt{d-c^2dx^2}\left(\frac{1}{2}\left(11\int\frac{1}{1-c^2x^2}dx+6x\right)-\frac{x}{2(1-c^2x^2)}\right)-\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{c^6d^3}}{3c^5d^3\sqrt{cx-1}\sqrt{cx+1}+\frac{2(a+\operatorname{barccosh}(cx))}{c^6d^2\sqrt{d-c^2dx^2}}+\frac{a+\operatorname{barccosh}(cx)}{3c^6d(d-c^2dx^2)^{3/2}}} \\
& \quad \downarrow 219 \\
& -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{c^6d^3}-\frac{2(a+\operatorname{barccosh}(cx))}{c^6d^2\sqrt{d-c^2dx^2}}+\frac{a+\operatorname{barccosh}(cx)}{3c^6d(d-c^2dx^2)^{3/2}}+ \\
& \quad \frac{b\left(\frac{1}{2}\left(\frac{11\operatorname{arctanh}(cx)}{c}+6x\right)-\frac{x}{2(1-c^2x^2)}\right)\sqrt{d-c^2dx^2}}{3c^5d^3\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(x^5*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2),x]`

output `(a + b*ArcCosh[c*x])/(3*c^6*d*(d - c^2*d*x^2)^(3/2)) - (2*(a + b*ArcCosh[c*x]))/(c^6*d^2*sqrt[d - c^2*d*x^2]) - (sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(c^6*d^3) + (b*sqrt[d - c^2*d*x^2]*(-1/2*x/(1 - c^2*x^2) + (6*x + (11*ArcTanh[c*x])/c)/2))/(3*c^5*d^3*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.124.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

---

3.124.  $\int \frac{x^5(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

```
rule 219 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

```
rule 299 Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x
*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2
*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && NeQ[2*p + 3, 0]
```

```
rule 1471 Int[((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.),
x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2
, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], x
, 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Simp[1/(2*d*(q
+ 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3),
x], x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^
2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]
```

```
rule 6337 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_
), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCo
sh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c
*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x], x]] /; FreeQ[{a, b
, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)]
&& (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])
```

### 3.124.4 Maple [A] (verified)

Time = 1.30 (sec) , antiderivative size = 398, normalized size of antiderivative = 1.64

method	result
default	$a \left( -\frac{x^4}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} + \frac{4x^2}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{8}{3d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} (6\sqrt{cx - 1} \sqrt{cx + 1} \arccos(\frac{cx}{c}))}{c^2}$
parts	$a \left( -\frac{x^4}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} + \frac{4x^2}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{8}{3d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}} \right) - \frac{b \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} (6\sqrt{cx - 1} \sqrt{cx + 1} \arccos(\frac{cx}{c}))}{c^2}$

3.124.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

input `int(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output  $a*(-x^4/c^2/d/(-c^2*d*x^2+d)^{(3/2)}+4/c^2*(x^2/c^2/d/(-c^2*d*x^2+d)^{(3/2)}-2/3/d/c^4/(-c^2*d*x^2+d)^{(3/2))}-1/6*b*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(6*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*arccosh(c*x)*c^4*x^4-6*c^5*x^5-11*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*c^4*x^4+11*\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)*c^4*x^4-24*(c*x+1)^{(1/2)}*arccosh(c*x)*(c*x-1)^{(1/2)}*c^2*x^2+11*c^3*x^3+22*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*x^2*c^2-22*\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)*x^2*c^2+16*arccosh(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}-5*c*x-11*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+11*\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1))/c^6/(c^6*x^6-3*c^4*x^4+3*c^2*x^2-1)/d^3$

### 3.124.5 Fracas [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 529, normalized size of antiderivative = 2.18

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \left[ -\frac{8(3bc^4x^4 - 12bc^2x^2 + 8b)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) + 11(bc^4x^4}{(d - c^2 dx^2)^{5/2}} \right]$$

input `integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="fracas")`

output  $[-1/24*(8*(3*b*c^4*x^4 - 12*b*c^2*x^2 + 8*b)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) + 11*(b*c^4*x^4 - 2*b*c^2*x^2 + b)*\sqrt{-d}*\log(-(c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 + 4*(c^3*x^3 + c*x)*\sqrt{-c^2*d*x^2 + d})*\sqrt{c^2*x^2 - 1}*\sqrt{-d} - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)) - 4*(6*b*c^3*x^3 - 5*b*c*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} + 8*(3*a*c^4*x^4 - 12*a*c^2*x^2 + 8*a)*\sqrt{-c^2*d*x^2 + d}]/(c^{10}*d^3*x^4 - 2*c^8*d^3*x^2 + c^6*d^3), 1/12*(11*(b*c^4*x^4 - 2*b*c^2*x^2 + b)*\sqrt{d}*\arctan(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*c*\sqrt{d})*x/(c^4*d*x^4 - d)) - 4*(3*b*c^4*x^4 - 12*b*c^2*x^2 + 8*b)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) + 2*(6*b*c^3*x^3 - 5*b*c*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} - 4*(3*a*c^4*x^4 - 12*a*c^2*x^2 + 8*a)*\sqrt{-c^2*d*x^2 + d}]/(c^{10}*d^3*x^4 - 2*c^8*d^3*x^2 + c^6*d^3)]$



## 3.124.6 Sympy [F]

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{5}{2}}} dx$$

input `integrate(x**5*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral(x**5*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`

## 3.124.7 Maxima [F]

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^5}{(-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `-1/9*b*(((9*c^4*sqrt(d)*x^4 - 8*sqrt(d))*sqrt(c*x + 1)*sqrt(c*x - 1)/sqrt(-c*x + 1) - 3*(3*c^5*sqrt(d)*x^5 - 12*c^3*sqrt(d)*x^3 + 8*c*sqrt(d)*x + 3*c^4*sqrt(d)*x^4 - 12*c^2*sqrt(d)*x^2 + 8*sqrt(d))*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/sqrt(-c*x + 1))/((c^8*d^3*x^2 - c^6*d^3)*(c*x + 1)*sqrt(c*x - 1) + (c^9*d^3*x^3 - c^7*d^3*x)*sqrt(c*x + 1)) + 9*integrate(1/9*(9*c^7*sqrt(d)*x^7 - 45*c^5*sqrt(d)*x^5 + 60*c^3*sqrt(d)*x^3 - 24*c*sqrt(d)*x + (9*c^6*sqrt(d)*x^6 - 54*c^4*sqrt(d)*x^4 + 60*c^2*sqrt(d)*x^2 - 16*sqrt(d))*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1)))/sqrt(-c*x + 1)*((c^9*d^3*x^4 - 2*c^7*d^3*x^2 + c^5*d^3)*e^(3/2*log(c*x + 1) + log(c*x - 1)) + 2*(c^10*d^3*x^5 - 2*c^8*d^3*x^3 + c^6*d^3*x)*e^(log(c*x + 1) + 1/2*log(c*x - 1)) + (c^11*d^3*x^6 - 2*c^9*d^3*x^4 + c^7*d^3*x^2)*sqrt(c*x + 1)), x) - 1/3*a*(3*x^4/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 12*x^2/((-c^2*d*x^2 + d)^(3/2)*c^4*d) + 8/((-c^2*d*x^2 + d)^(3/2)*c^6*d))`

**3.124.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^5*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.124.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x^5*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2),x)`

output `int((x^5*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2), x)`

**3.125**  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

3.125.1 Optimal result . . . . . 1114  
 3.125.2 Mathematica [A] (warning: unable to verify) . . . . . 1115  
 3.125.3 Rubi [A] (verified) . . . . . 1115  
 3.125.4 Maple [A] (verified) . . . . . 1118  
 3.125.5 Fricas [F] . . . . . 1119  
 3.125.6 Sympy [F] . . . . . 1119  
 3.125.7 Maxima [F] . . . . . 1120  
 3.125.8 Giac [F] . . . . . 1120  
 3.125.9 Mupad [F(-1)] . . . . . 1120

**3.125.1 Optimal result**

Integrand size = 27, antiderivative size = 224

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}}{6c^5d(d - c^2dx^2)^{3/2}} + \frac{x^3(a + \operatorname{arccosh}(cx))}{3c^2d(d - c^2dx^2)^{3/2}} - \frac{x(a + \operatorname{arccosh}(cx))}{c^4d^2\sqrt{d - c^2dx^2}} + \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{2bc^5d^2\sqrt{d - c^2dx^2}} + \frac{2b\sqrt{-1 + cx}\sqrt{1 + cx} \log(1 - c^2x^2)}{3c^5d^2\sqrt{d - c^2dx^2}}$$

```
output 1/3*x^3*(a+b*arccosh(c*x))/c^2/d/(-c^2*d*x^2+d)^(3/2)+1/6*b*(c*x-1)^(1/2)*
(c*x+1)^(1/2)/c^5/d/(-c^2*d*x^2+d)^(3/2)-x*(a+b*arccosh(c*x))/c^4/d^2/(-c^
2*d*x^2+d)^(1/2)+1/2*(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^
5/d^2/(-c^2*d*x^2+d)^(1/2)+2/3*b*ln(-c^2*x^2+1)*(c*x-1)^(1/2)*(c*x+1)^(1/2
)/c^5/d^2/(-c^2*d*x^2+d)^(1/2)
```

**3.125.2 Mathematica [A] (warning: unable to verify)**

Time = 0.72 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.00

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \frac{2acx(-3+4c^2x^2)\sqrt{d-c^2dx^2}}{(-1+c^2x^2)^2} - 6a\sqrt{d} \arctan\left(\frac{cx\sqrt{d-c^2dx^2}}{\sqrt{d(-1+c^2x^2)}}\right) + \frac{bd\left(-8cx \operatorname{arccosh}(cx) - \sqrt{\frac{-1}{1+c^2x^2}}\right)}{6c^5d^3}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2), x]`

output  $((2*a*c*x*(-3 + 4*c^2*x^2)*\operatorname{Sqrt}[d - c^2*d*x^2])/(-1 + c^2*x^2)^2 - 6*a*\operatorname{Sqrt}[d]*\operatorname{ArcTan}[(c*x*\operatorname{Sqrt}[d - c^2*d*x^2])/(\operatorname{Sqrt}[d]*(-1 + c^2*x^2))] + (b*d*(-8*c*x*\operatorname{ArcCosh}[c*x] - (\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x) + 2*c*x*\operatorname{ArcCosh}[c*x]))/(-1 + c^2*x^2) + \operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(3*\operatorname{ArcCosh}[c*x]^2 + 8*\operatorname{Log}[\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)])))/\operatorname{Sqrt}[d - c^2*d*x^2])/(6*c^5*d^3)$

**3.125.3 Rubi [A] (verified)**Time = 0.88 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.18, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.370$ , Rules used = {6349, 82, 243, 49, 2009, 6349, 25, 82, 240, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx \\ & \quad \downarrow \text{6349} \\ & -\frac{\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^3}{(1-cx)^2(cx+1)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} + \frac{x^3(a + b \operatorname{arccosh}(cx))}{3c^2d(d - c^2 dx^2)^{3/2}} \\ & \quad \downarrow \text{82} \\ & -\frac{\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx}{c^2 d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^3}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} + \frac{x^3(a + b \operatorname{arccosh}(cx))}{3c^2d(d - c^2 dx^2)^{3/2}} \\ & \quad \downarrow \text{243} \end{aligned}$$

---

3.125.  $\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& -\frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx}{c^2d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2}{(1-c^2x^2)^2} dx^2}{6cd^2\sqrt{d-c^2dx^2}} + \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow 49 \\
& -\frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx}{c^2d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \left( \frac{1}{c^2(c^2x^2-1)} + \frac{1}{c^2(c^2x^2-1)^2} \right) dx^2}{6cd^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow 2009 \\
& -\frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx}{c^2d} + \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{c^4(1-c^2x^2)} + \frac{\log(1-c^2x^2)}{c^4} \right)}{6cd^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 6349 \\
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{d-c^2dx^2}} dx}{c^2d} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \int -\frac{x}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} + \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{c^4(1-c^2x^2)} + \frac{\log(1-c^2x^2)}{c^4} \right)}{6cd^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 25 \\
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} + \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{c^4(1-c^2x^2)} + \frac{\log(1-c^2x^2)}{c^4} \right)}{6cd^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 82 \\
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x}{1-c^2x^2} dx}{cd\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} + \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{c^4(1-c^2x^2)} + \frac{\log(1-c^2x^2)}{c^4} \right)}{6cd^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 240
\end{aligned}$$

---

3.125.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
& -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}\log(1-c^2x^2)}{2c^3d\sqrt{d-c^2dx^2}} + \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} + \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{c^4(1-c^2x^2)} + \frac{\log(1-c^2x^2)}{c^4}\right)}{6cd^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6307} \\
& \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d(d-c^2dx^2)^{3/2}} - \frac{x(a+\operatorname{barccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^2}{2bc^3d\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}\log(1-c^2x^2)}{2c^3d\sqrt{d-c^2dx^2}} + \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{c^4(1-c^2x^2)} + \frac{\log(1-c^2x^2)}{c^4}\right)}{6cd^2\sqrt{d-c^2dx^2}}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2),x]`

output `(x^3*(a + b*ArcCosh[c*x]))/(3*c^2*d*(d - c^2*d*x^2)^(3/2)) + (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(1/(c^4*(1 - c^2*x^2)) + Log[1 - c^2*x^2]/c^4))/(6*c*d^2*Sqrt[d - c^2*d*x^2]) - ((x*(a + b*ArcCosh[c*x]))/(c^2*d*Sqrt[d - c^2*d*x^2]) - (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(2*b*c^3*d*Sqrt[d - c^2*d*x^2]) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2])/(2*c^3*d*Sqrt[d - c^2*d*x^2]))/(c^2*d)`

### 3.125.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

---

3.125.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

rule 243 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6307 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

rule 6349 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*(m - 1)/(2*e*(p + 1)) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]`

### 3.125.4 Maple [A] (verified)

Time = 1.20 (sec) , antiderivative size = 366, normalized size of antiderivative = 1.63

method	result
default	$\frac{ax^3}{3c^2d(-c^2dx^2+d)^{\frac{3}{2}}} - \frac{ax}{c^4d^2\sqrt{-c^2dx^2+d}} + \frac{a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{c^4d^2\sqrt{c^2d}} - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}}{3 \operatorname{arccosh}(cx)^2x^4c^4-8vd}$
parts	$\frac{ax^3}{3c^2d(-c^2dx^2+d)^{\frac{3}{2}}} - \frac{ax}{c^4d^2\sqrt{-c^2dx^2+d}} + \frac{a \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{c^4d^2\sqrt{c^2d}} - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}}{3 \operatorname{arccosh}(cx)^2x^4c^4-8vd}$

input `int(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

3.125. 
$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$$

output  $\frac{1}{3}ax^3/c^2/d/(-c^2dx^2+d)^{3/2}-a/c^4/d^2x/(-c^2dx^2+d)^{1/2}+a/c^4/d^2/(c^2d)^{1/2}*\arctan((c^2d)^{1/2}*x/(-c^2dx^2+d)^{1/2})-1/6*b*(-d*(c^2x^2-1))^{1/2}*(c*x-1)^{1/2}*(c*x+1)^{1/2}/(c^6*x^6-3*c^4*x^4+3*c^2*x^2-1)/d^3/c^5*(3*\operatorname{arccosh}(c*x)^2*x^4*c^4-8*(c*x+1)^{1/2}*\operatorname{arccosh}(c*x)*(c*x-1)^{1/2}*c^3*x^3-8*c^4*x^4*\operatorname{arccosh}(c*x)+8*\ln((c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))^2-1)*x^4*c^4-6*\operatorname{arccosh}(c*x)^2*x^2*c^2+6*(c*x+1)^{1/2}*\operatorname{arccosh}(c*x)*(c*x-1)^{1/2}*c*x+16*c^2*x^2*\operatorname{arccosh}(c*x)-16*\ln((c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))^2-1)*x^2*c^2-c^2*x^2+3*\operatorname{arccosh}(c*x)^2-8*\operatorname{arccosh}(c*x)+8*\ln((c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))^2-1)+1)$

### 3.125.5 Fracas [F]

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^4}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-(b*x^4*arccosh(c*x) + a*x^4)*sqrt(-c^2*d*x^2 + d)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

### 3.125.6 Sympy [F]

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{5/2}} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral(x**4*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`



**3.125.7 Maxima [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(-c^2dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/3*(x*(3*x^2/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 2/((-c^2*d*x^2 + d)^(3/2)*c^4*d)) - x/(sqrt(-c^2*d*x^2 + d)*c^4*d^2) + 3*arcsin(c*x)/(c^5*d^(5/2)))*a + b*integrate(x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(-c^2*d*x^2 + d)^(5/2), x)`

**3.125.8 Giac [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(-c^2dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^4/(-c^2*d*x^2 + d)^(5/2), x)`

**3.125.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(d - c^2dx^2)^{5/2}} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2),x)`

output `int((x^4*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2), x)`

$$3.126 \quad \int \frac{x^3(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

3.126.1 Optimal result . . . . .	1121
3.126.2 Mathematica [A] (verified) . . . . .	1121
3.126.3 Rubi [A] (verified) . . . . .	1122
3.126.4 Maple [A] (verified) . . . . .	1123
3.126.5 Fracas [A] (verification not implemented) . . . . .	1124
3.126.6 Sympy [F] . . . . .	1124
3.126.7 Maxima [A] (verification not implemented) . . . . .	1125
3.126.8 Giac [F(-2)] . . . . .	1125
3.126.9 Mupad [F(-1)] . . . . .	1126

### 3.126.1 Optimal result

Integrand size = 27, antiderivative size = 158

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \frac{bx\sqrt{d - c^2 dx^2}}{6c^3 d^3 (-1 + cx)^{3/2} (1 + cx)^{3/2}} + \frac{a + \operatorname{arccosh}(cx)}{3c^4 d (d - c^2 dx^2)^{3/2}} - \frac{a + \operatorname{arccosh}(cx)}{c^4 d^2 \sqrt{d - c^2 dx^2}} + \frac{5b\sqrt{d - c^2 dx^2} \operatorname{arctanh}(cx)}{6c^4 d^3 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

output  $\frac{1}{3} \frac{(a + b \operatorname{arccosh}(c x))}{c^4 d} \frac{1}{(-c^2 d x^2 + d)^{3/2}} + \frac{(-a - b \operatorname{arccosh}(c x))}{c^4 d^2} \frac{1}{(-c^2 d x^2 + d)^{1/2}} + \frac{1}{6} \frac{b x}{c^3 d^3} \frac{1}{(c x - 1)^{3/2}} \frac{1}{(c x + 1)^{3/2}} + \frac{5}{6} \frac{b \operatorname{arctanh}(c x)}{c^4 d^3} \frac{1}{\sqrt{-1 + c x} \sqrt{1 + c x}}$

### 3.126.2 Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.77

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \frac{4a - 6ac^2 x^2 - bcx\sqrt{-1 + cx}\sqrt{1 + cx} + b(4 - 6c^2 x^2) \operatorname{arccosh}(cx) - 5b\sqrt{-1 + cx}}{6c^4 d^2 (-1 + c^2 x^2) \sqrt{d - c^2 dx^2}}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2),x]`

---

3.126.  $\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

output  $(4*a - 6*a*c^2*x^2 - b*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x] + b*(4 - 6*c^2*x^2) * \text{ArcCosh}[c*x] - 5*b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(-1 + c^2*x^2)*\text{ArcTanh}[c*x]) / (6*c^4*d^2*(-1 + c^2*x^2)*\text{Sqrt}[d - c^2*d*x^2])$

### 3.126.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.87, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6337, 27, 298, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

↓ 6337

$$-\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{2-3c^2 x^2}{3c^4 d^3 (1-c^2 x^2)^2} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{a + \text{barccosh}(cx)}{c^4 d^2 \sqrt{d - c^2 dx^2}} + \frac{a + \text{barccosh}(cx)}{3c^4 d (d - c^2 dx^2)^{3/2}}$$

↓ 27

$$\frac{b\sqrt{d - c^2 dx^2} \int \frac{2-3c^2 x^2}{(1-c^2 x^2)^2} dx}{3c^3 d^3 \sqrt{cx - 1}\sqrt{cx + 1}} - \frac{a + \text{barccosh}(cx)}{c^4 d^2 \sqrt{d - c^2 dx^2}} + \frac{a + \text{barccosh}(cx)}{3c^4 d (d - c^2 dx^2)^{3/2}}$$

↓ 298

$$\frac{b\sqrt{d - c^2 dx^2} \left( \frac{5}{2} \int \frac{1}{1-c^2 x^2} dx - \frac{x}{2(1-c^2 x^2)} \right)}{3c^3 d^3 \sqrt{cx - 1}\sqrt{cx + 1}} - \frac{a + \text{barccosh}(cx)}{c^4 d^2 \sqrt{d - c^2 dx^2}} + \frac{a + \text{barccosh}(cx)}{3c^4 d (d - c^2 dx^2)^{3/2}}$$

↓ 219

$$-\frac{a + \text{barccosh}(cx)}{c^4 d^2 \sqrt{d - c^2 dx^2}} + \frac{a + \text{barccosh}(cx)}{3c^4 d (d - c^2 dx^2)^{3/2}} + \frac{b \left( \frac{5 \text{arctanh}(cx)}{2c} - \frac{x}{2(1-c^2 x^2)} \right) \sqrt{d - c^2 dx^2}}{3c^3 d^3 \sqrt{cx - 1}\sqrt{cx + 1}}$$

input  $\text{Int}[(x^3*(a + b*\text{ArcCosh}[c*x]))/(d - c^2*d*x^2)^(5/2), x]$

output  $(a + b*\text{ArcCosh}[c*x]) / (3*c^4*d*(d - c^2*d*x^2)^(3/2)) - (a + b*\text{ArcCosh}[c*x]) / (c^4*d^2*\text{Sqrt}[d - c^2*d*x^2]) + (b*\text{Sqrt}[d - c^2*d*x^2]*(-1/2*x/(1 - c^2*x^2) + (5*\text{ArcTanh}[c*x])/(2*c))) / (3*c^3*d^3*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])$

---

3.126.  $\int \frac{x^3(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

## 3.126.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 298 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[(-(b*c - a*d))*x*((a + b*x^2)^(p + 1)/(2*a*b*(p + 1))), x] - Simp[(a*d - b*c*(2*p + 3))/(2*a*b*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/2 + p, 0])`
- rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

## 3.126.4 Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.63

method	result
default	$a \left( \frac{x^2}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{2}{3 d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}} \right) + b \left( \frac{\sqrt{-d(c^2 x^2 - 1)} (6c^2 x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - 4 \operatorname{arccosh}(cx))}{6(c^2 x^2 - 1)^2 d^3 c^4} \right)$
parts	$a \left( \frac{x^2}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{2}{3 d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}} \right) + b \left( \frac{\sqrt{-d(c^2 x^2 - 1)} (6c^2 x^2 \operatorname{arccosh}(cx) + \sqrt{cx-1} \sqrt{cx+1} cx - 4 \operatorname{arccosh}(cx))}{6(c^2 x^2 - 1)^2 d^3 c^4} \right)$

input `int(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

3.126. 
$$\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$$

output  $a*(x^2/c^2/d/(-c^2*d*x^2+d)^{(3/2)}-2/3/d/c^4/(-c^2*d*x^2+d)^{(3/2)})+b*(1/6*(-d*(c^2*x^2-1))^{(1/2)}*(6*c^2*x^2*\operatorname{arccosh}(c*x)+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x-4*\operatorname{arccosh}(c*x))/(c^2*x^2-1)^2/d^3/c^4-5/6*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/c^4/(c^2*x^2-1)*\ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)+5/6*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/c^4/(c^2*x^2-1)*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))$

### 3.126.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 469, normalized size of antiderivative = 2.97

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \left[ \frac{4\sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} b c x + 8(3bc^2 x^2 - 2b)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1})}{(d - c^2 dx^2)^{5/2}} \right]$$

input `integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="fracas")`

output  $[1/24*(4*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*b*c*x + 8*(3*b*c^2*x^2 - 2*b)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) - 5*(b*c^4*x^4 - 2*b*c^2*x^2 + b)*\sqrt{-d}*\log(-(c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 + 4*(c^3*x^3 + c*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*\sqrt{-d} - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)) + 8*(3*a*c^2*x^2 - 2*a)*\sqrt{-c^2*d*x^2 + d})/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3), 1/12*(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*b*c*x + 5*(b*c^4*x^4 - 2*b*c^2*x^2 + b)*\sqrt{d}*\arctan(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*c*\sqrt{d}*x/(c^4*d*x^4 - d)) + 4*(3*b*c^2*x^2 - 2*b)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1}) + 4*(3*a*c^2*x^2 - 2*a)*\sqrt{-c^2*d*x^2 + d})/(c^8*d^3*x^4 - 2*c^6*d^3*x^2 + c^4*d^3)]$

### 3.126.6 Sympy [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{5/2}} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral(x**3*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`

---

3.126.  $\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

**3.126.7 Maxima [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.11

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \frac{1}{12} bc \left( \frac{2\sqrt{-d}x}{c^6 d^3 x^2 - c^4 d^3} + \frac{5\sqrt{-d} \log(cx + 1)}{c^5 d^3} - \frac{5\sqrt{-d} \log(cx - 1)}{c^5 d^3} \right) + \frac{1}{3} b \left( \frac{3x^2}{(-c^2 dx^2 + d)^{3/2} c^2 d} - \frac{2}{(-c^2 dx^2 + d)^{3/2} c^4 d} \right) \operatorname{arccosh}(cx) + \frac{1}{3} a \left( \frac{3x^2}{(-c^2 dx^2 + d)^{3/2} c^2 d} - \frac{2}{(-c^2 dx^2 + d)^{3/2} c^4 d} \right)$$

```
input integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")
```

```
output 1/12*b*c*(2*sqrt(-d)*x/(c^6*d^3*x^2 - c^4*d^3) + 5*sqrt(-d)*log(c*x + 1)/(c^5*d^3) - 5*sqrt(-d)*log(c*x - 1)/(c^5*d^3)) + 1/3*b*(3*x^2/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 2/((-c^2*d*x^2 + d)^(3/2)*c^4*d))*arccosh(c*x) + 1/3*a*(3*x^2/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 2/((-c^2*d*x^2 + d)^(3/2)*c^4*d))
```

**3.126.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(x^3*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.126.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2),x)`output `int((x^3*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2), x)`

$$3.127 \quad \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$$

3.127.1 Optimal result . . . . . 1127  
 3.127.2 Mathematica [A] (verified) . . . . . 1127  
 3.127.3 Rubi [A] (verified) . . . . . 1128  
 3.127.4 Maple [B] (verified) . . . . . 1130  
 3.127.5 Fracas [F] . . . . . 1130  
 3.127.6 Sympy [F] . . . . . 1131  
 3.127.7 Maxima [A] (verification not implemented) . . . . . 1131  
 3.127.8 Giac [F] . . . . . 1132  
 3.127.9 Mupad [F(-1)] . . . . . 1132

### 3.127.1 Optimal result

Integrand size = 27, antiderivative size = 133

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}}{6c^3d(d - c^2dx^2)^{3/2}} + \frac{x^3(a + \operatorname{arccosh}(cx))}{3d(d - c^2dx^2)^{3/2}} + \frac{b\sqrt{-1 + cx}\sqrt{1 + cx} \log(1 - c^2x^2)}{6c^3d^2\sqrt{d - c^2dx^2}}$$

output `1/3*x^3*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(3/2)+1/6*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d/(-c^2*d*x^2+d)^(3/2)+1/6*b*ln(-c^2*x^2+1)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d^2/(-c^2*d*x^2+d)^(1/2)`

### 3.127.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.76

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx} \left( -\frac{2x^3(a + \operatorname{arccosh}(cx))}{(-1 + cx)^{3/2}(1 + cx)^{3/2}} + \frac{b \left( \frac{1}{1 - c^2x^2} + \log(1 - c^2x^2) \right)}{c^3} \right)}{6d^2\sqrt{d - c^2dx^2}}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2), x]`



output  $(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*((-2*x^3*(a + b*\text{ArcCosh}[c*x]))/((-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (b*((1 - c^2*x^2)^(-1) + \text{Log}[1 - c^2*x^2]))/c^3))/ (6*d^2*\text{Sqrt}[d - c^2*d*x^2])$

### 3.127.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.81, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6332, 82, 243, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

↓ 6332

$$\frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x^3}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \text{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}}$$

↓ 82

$$\frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x^3}{(1-c^2 x^2)^2} dx}{3d^2\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \text{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}}$$

↓ 243

$$\frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x^2}{(1-c^2 x^2)^2} dx^2}{6d^2\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \text{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}}$$

↓ 49

$$\frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \left( \frac{1}{c^2(c^2 x^2 - 1)} + \frac{1}{c^2(c^2 x^2 - 1)^2} \right) dx^2}{6d^2\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \text{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}}$$

↓ 2009

$$\frac{x^3(a + \text{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{1}{c^4(1-c^2 x^2)} + \frac{\log(1-c^2 x^2)}{c^4} \right)}{6d^2\sqrt{d - c^2 dx^2}}$$

input  $\text{Int}[(x^2*(a + b*\text{ArcCosh}[c*x]))/(d - c^2*d*x^2)^(5/2), x]$

---

3.127.  $\int \frac{x^2(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

```
output (x^3*(a + b*ArcCosh[c*x]))/(3*d*(d - c^2*d*x^2)^(3/2)) + (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(1/(c^4*(1 - c^2*x^2)) + Log[1 - c^2*x^2]/c^4))/(6*d^2*Sqrt[d - c^2*d*x^2])
```

### 3.127.3.1 Defintions of rubi rules used

```
rule 49 Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

```
rule 82 Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]
```

```
rule 243 Int[(x_)^m_*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6332 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]
```

### 3.127.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 457 vs.  $2(113) = 226$ .

Time = 1.19 (sec) , antiderivative size = 458, normalized size of antiderivative = 3.44

method	result
default	$a \left( \frac{x}{2c^2 d(-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{\frac{x}{3d(-c^2 d x^2 + d)^{\frac{3}{2}}} + \frac{2x}{3d^2 \sqrt{-c^2 d x^2 + d}}}{2c^2} \right) + \frac{b \sqrt{-d(c^2 x^2 - 1)} (c^3 x^3 + \sqrt{cx-1} \sqrt{cx+1} c^2 x^2 - \sqrt{cx-1} \sqrt{cx+1})}{2c^2}$
parts	$a \left( \frac{x}{2c^2 d(-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{\frac{x}{3d(-c^2 d x^2 + d)^{\frac{3}{2}}} + \frac{2x}{3d^2 \sqrt{-c^2 d x^2 + d}}}{2c^2} \right) + \frac{b \sqrt{-d(c^2 x^2 - 1)} (c^3 x^3 + \sqrt{cx-1} \sqrt{cx+1} c^2 x^2 - \sqrt{cx-1} \sqrt{cx+1})}{2c^2}$

input `int(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output 
$$a*(1/2*x/c^2/d/(-c^2*d*x^2+d)^(3/2)-1/2/c^2*(1/3/d*x/(-c^2*d*x^2+d)^(3/2)+2/3/d^2*x/(-c^2*d*x^2+d)^(1/2)))+1/6*b*(-d*(c^2*x^2-1))^(1/2)*(c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^5*c^5+2*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^6*c^6+6*c^4*x^4*arccosh(c*x)+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^3*c^3-6*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^4*c^4+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-c^4*x^4-6*c^2*x^2*arccosh(c*x)+6*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^2*c^2+2*c^2*x^2*arccosh(c*x)-2*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)-1)/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3/c^3$$

### 3.127.5 Fracas [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^2}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*x^2*arccosh(c*x) + a*x^2)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

---

3.127. 
$$\int \frac{x^2(a+b \operatorname{arccosh}(cx))}{(d-c^2 dx^2)^{5/2}} dx$$

**3.127.6 Sympy [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{5}{2}}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral(x**2*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`

**3.127.7 Maxima [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.27

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \frac{1}{6} bc \left( \frac{\sqrt{-d}}{c^6 d^3 x^2 - c^4 d^3} - \frac{\sqrt{-d} \log(cx + 1)}{c^4 d^3} - \frac{\sqrt{-d} \log(cx - 1)}{c^4 d^3} \right) - \frac{1}{3} b \left( \frac{x}{\sqrt{-c^2 dx^2 + dc^2 d^2}} - \frac{x}{(-c^2 dx^2 + d)^{\frac{3}{2}} c^2 d} \right) \operatorname{arcosh}(cx) - \frac{1}{3} a \left( \frac{x}{\sqrt{-c^2 dx^2 + dc^2 d^2}} - \frac{x}{(-c^2 dx^2 + d)^{\frac{3}{2}} c^2 d} \right)$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/6*b*c*(sqrt(-d)/(c^6*d^3*x^2 - c^4*d^3) - sqrt(-d)*log(c*x + 1)/(c^4*d^3) - sqrt(-d)*log(c*x - 1)/(c^4*d^3)) - 1/3*b*(x/(sqrt(-c^2*d*x^2 + d)*c^2*d^2) - x/((-c^2*d*x^2 + d)^(3/2)*c^2*d))*arccosh(c*x) - 1/3*a*(x/(sqrt(-c^2*d*x^2 + d)*c^2*d^2) - x/((-c^2*d*x^2 + d)^(3/2)*c^2*d))`

**3.127.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^2/(-c^2*d*x^2 + d)^(5/2), x)`

**3.127.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2),x)`

output `int((x^2*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2), x)`

**3.128** 
$$\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$$

3.128.1 Optimal result . . . . . 1133  
 3.128.2 Mathematica [A] (verified) . . . . . 1133  
 3.128.3 Rubi [A] (verified) . . . . . 1134  
 3.128.4 Maple [B] (verified) . . . . . 1135  
 3.128.5 Fricas [A] (verification not implemented) . . . . . 1136  
 3.128.6 Sympy [F] . . . . . 1136  
 3.128.7 Maxima [F] . . . . . 1137  
 3.128.8 Giac [F] . . . . . 1137  
 3.128.9 Mupad [F(-1)] . . . . . 1137

**3.128.1 Optimal result**

Integrand size = 25, antiderivative size = 127

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{bx\sqrt{-1 + cx}\sqrt{1 + cx}}{6cd(d - c^2dx^2)^{3/2}} + \frac{a + \operatorname{arccosh}(cx)}{3c^2d(d - c^2dx^2)^{3/2}} + \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arctanh}(cx)}{6c^2d^2\sqrt{d - c^2dx^2}}$$

output  $1/3*(a+b*\operatorname{arccosh}(c*x))/c^2/d/(-c^2*d*x^2+d)^{(3/2)}+1/6*b*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c/d/(-c^2*d*x^2+d)^{(3/2)}+1/6*b*\operatorname{arctanh}(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^2/d^2/(-c^2*d*x^2+d)^{(1/2)}$

**3.128.2 Mathematica [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.28

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{\sqrt{d - c^2dx^2}(-2bcx + 2bc^3x^3 + 4a\sqrt{-1 + cx}\sqrt{1 + cx} + 4b\sqrt{-1 + cx}\sqrt{1 + cx})}{12c^2d^2}$$

input  $\operatorname{Integrate}[(x*(a + b*\operatorname{ArcCosh}[c*x]))/(d - c^2*d*x^2)^{(5/2)}, x]$

output  $(\text{Sqrt}[d - c^2*d*x^2]*(-2*b*c*x + 2*b*c^3*x^3 + 4*a*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x] + 4*b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*\text{ArcCosh}[c*x] + b*(-1 + c^2*x^2)^2*\text{Log}[-1 + c*x] - b*\text{Log}[1 + c*x] + 2*b*c^2*x^2*\text{Log}[1 + c*x] - b*c^4*x^4*\text{Log}[1 + c*x]))/(12*c^2*d^3*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))$

### 3.128.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.84, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {6329, 39, 215, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

$$\downarrow \text{6329}$$

$$\frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{(1-cx)^2(cx+1)^2} dx}{3cd^2\sqrt{d - c^2 dx^2}} + \frac{a + \text{barccosh}(cx)}{3c^2 d (d - c^2 dx^2)^{3/2}}$$

$$\downarrow \text{39}$$

$$\frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{(1-c^2 x^2)^2} dx}{3cd^2\sqrt{d - c^2 dx^2}} + \frac{a + \text{barccosh}(cx)}{3c^2 d (d - c^2 dx^2)^{3/2}}$$

$$\downarrow \text{215}$$

$$\frac{b\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{1}{2} \int \frac{1}{1-c^2 x^2} dx + \frac{x}{2(1-c^2 x^2)} \right)}{3cd^2\sqrt{d - c^2 dx^2}} + \frac{a + \text{barccosh}(cx)}{3c^2 d (d - c^2 dx^2)^{3/2}}$$

$$\downarrow \text{219}$$

$$\frac{a + \text{barccosh}(cx)}{3c^2 d (d - c^2 dx^2)^{3/2}} + \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{\text{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2 x^2)} \right)}{3cd^2\sqrt{d - c^2 dx^2}}$$

input  $\text{Int}[(x*(a + b*\text{ArcCosh}[c*x]))/(d - c^2*d*x^2)^(5/2), x]$

output  $(a + b*\text{ArcCosh}[c*x])/(3*c^2*d*(d - c^2*d*x^2)^(3/2)) + (b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(x/(2*(1 - c^2*x^2)) + \text{ArcTanh}[c*x]/(2*c)))/(3*c*d^2*\text{Sqrt}[d - c^2*d*x^2])$

---

3.128.  $\int \frac{x(a + \text{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

## 3.128.3.1 Defintions of rubi rules used

- rule 39 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^(n)/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^(p*(-1 + c*x)^p)) Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

## 3.128.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs.  $2(107) = 214$ .

Time = 1.14 (sec) , antiderivative size = 220, normalized size of antiderivative = 1.73

method	result
default	$\frac{a}{3c^2d(-c^2dx^2+d)^{\frac{3}{2}}} + b \left( \frac{\sqrt{-d(c^2x^2-1)}(\sqrt{cx-1}\sqrt{cx+1}cx+2 \operatorname{arccosh}(cx))}{6(c^2x^2-1)^2d^3c^2} - \frac{\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \ln(1+cx+\sqrt{cx-1})}{6d^3c^2(c^2x^2-1)} \right)$
parts	$\frac{a}{3c^2d(-c^2dx^2+d)^{\frac{3}{2}}} + b \left( \frac{\sqrt{-d(c^2x^2-1)}(\sqrt{cx-1}\sqrt{cx+1}cx+2 \operatorname{arccosh}(cx))}{6(c^2x^2-1)^2d^3c^2} - \frac{\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1} \ln(1+cx+\sqrt{cx-1})}{6d^3c^2(c^2x^2-1)} \right)$

input `int(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

3.128. 
$$\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$$



output  $\frac{1}{3} \frac{a}{c^2 d} (-c^2 d x^2 + d)^{3/2} + b \left( \frac{1}{6} (-d (c^2 x^2 - 1))^{1/2} ((c x - 1)^{1/2} (c x + 1)^{1/2} c x + 2 \operatorname{arccosh}(c x)) / (c^2 x^2 - 1)^{2/3} / c^2 - \frac{1}{6} (-d (c^2 x^2 - 1))^{1/2} (c x - 1)^{1/2} (c x + 1)^{1/2} / d^3 / c^2 / (c^2 x^2 - 1) \ln(1 + c x + (c x - 1)^{1/2} (c x + 1)^{1/2}) + \frac{1}{6} (-d (c^2 x^2 - 1))^{1/2} (c x - 1)^{1/2} (c x + 1)^{1/2} / d^3 / c^2 / (c^2 x^2 - 1) \ln((c x - 1)^{1/2} (c x + 1)^{1/2} + c x - 1) \right)$

### 3.128.5 Fricas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 421, normalized size of antiderivative = 3.31

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \left[ \frac{4 \sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} b c x + 8 \sqrt{-c^2 dx^2 + d} b \log(cx + \sqrt{c^2 x^2 - 1}) - (bc^4 x^4 - 2bc^2 x^2 + b) \sqrt{-d} \log(-c^6 dx^6 + 5c^4 dx^4 - 5c^2 dx^2 - 4(c^3 x^3 + cx) \sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} \sqrt{-d} - d) / (c^6 x^6 - 3c^4 x^4 + 3c^2 x^2 - 1) + 8 \sqrt{-c^2 dx^2 + d} a / (c^6 d^3 x^4 - 2c^4 d^3 x^2 + c^2 d^3), 1/12 (2 \sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} b c x - (bc^4 x^4 - 2bc^2 x^2 + b) \sqrt{d} \arctan(2 \sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} c \sqrt{d} x / (c^4 dx^4 - d)) + 4 \sqrt{-c^2 dx^2 + d} b \log(cx + \sqrt{c^2 x^2 - 1}) + 4 \sqrt{-c^2 dx^2 + d} a / (c^6 d^3 x^4 - 2c^4 d^3 x^2 + c^2 d^3)) \right]$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output  $[1/24*(4*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*b*c*x + 8*\sqrt{-c^2*d*x^2 + d}*b*\log(cx + \sqrt{c^2*x^2 - 1}) - (b*c^4*x^4 - 2*b*c^2*x^2 + b)*\sqrt{-d}*\log(-c^6*d*x^6 + 5*c^4*d*x^4 - 5*c^2*d*x^2 - 4*(c^3*x^3 + c*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*\sqrt{-d} - d)/(c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1) + 8*\sqrt{-c^2*d*x^2 + d}*a/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3), 1/12*(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*b*c*x - (b*c^4*x^4 - 2*b*c^2*x^2 + b)*\sqrt{d}*\arctan(2*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*c*\sqrt{d}*x/(c^4*d*x^4 - d)) + 4*\sqrt{-c^2*d*x^2 + d}*b*\log(cx + \sqrt{c^2*x^2 - 1}) + 4*\sqrt{-c^2*d*x^2 + d}*a/(c^6*d^3*x^4 - 2*c^4*d^3*x^2 + c^2*d^3)]$

### 3.128.6 Sympy [F]

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{5/2}} dx$$

input `integrate(x*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral(x*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`

**3.128.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `b*integrate(x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(-c^2*d*x^2 + d)^(5/2), x) + 1/3*a/((-c^2*d*x^2 + d)^(3/2)*c^2*d)`

**3.128.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x/(-c^2*d*x^2 + d)^(5/2), x)`

**3.128.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2),x)`

output `int((x*(a + b*acosh(c*x)))/(d - c^2*d*x^2)^(5/2), x)`

**3.129**  $\int \frac{a+b\operatorname{arccosh}(cx)}{(d-c^2dx^2)^{5/2}} dx$

3.129.1 Optimal result . . . . . 1138  
 3.129.2 Mathematica [A] (verified) . . . . . 1138  
 3.129.3 Rubi [A] (verified) . . . . . 1139  
 3.129.4 Maple [B] (verified) . . . . . 1141  
 3.129.5 Fricas [F] . . . . . 1141  
 3.129.6 Sympy [F] . . . . . 1142  
 3.129.7 Maxima [A] (verification not implemented) . . . . . 1142  
 3.129.8 Giac [F] . . . . . 1142  
 3.129.9 Mupad [F(-1)] . . . . . 1143

**3.129.1 Optimal result**

Integrand size = 24, antiderivative size = 162

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d - c^2dx^2)^{5/2}} dx = \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}}{6cd(d - c^2dx^2)^{3/2}} + \frac{x(a + \operatorname{arccosh}(cx))}{3d(d - c^2dx^2)^{3/2}} + \frac{2x(a + \operatorname{arccosh}(cx))}{3d^2\sqrt{d - c^2dx^2}} - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx} \log(1 - c^2x^2)}{3cd^2\sqrt{d - c^2dx^2}}$$

output `1/3*x*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(3/2)+1/6*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d/(-c^2*d*x^2+d)^(3/2)+2/3*x*(a+b*arccosh(c*x))/d^2/(-c^2*d*x^2+d)^(1/2)-1/3*b*ln(-c^2*x^2+1)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d^2/(-c^2*d*x^2+d)^(1/2)`

**3.129.2 Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.81

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d - c^2dx^2)^{5/2}} dx = \frac{-6acx + 4ac^3x^3 - b\sqrt{-1 + cx}\sqrt{1 + cx} + 2bcx(-3 + 2c^2x^2) \operatorname{arccosh}(cx) - 2b\sqrt{-1 + cx}\sqrt{1 + cx} \log[1 - c^2x^2]}{6cd^2(-1 + c^2x^2)\sqrt{d - c^2dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^(5/2), x]`

output `(-6*a*c*x + 4*a*c^3*x^3 - b*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*b*c*x*(-3 + 2*c^2*x^2)*ArcCosh[c*x] - 2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1 + c^2*x^2)*Log[1 - c^2*x^2])/(6*c*d^2*(-1 + c^2*x^2)*Sqrt[d - c^2*d*x^2])`

---

3.129.  $\int \frac{a+b\operatorname{arccosh}(cx)}{(d-c^2dx^2)^{5/2}} dx$

**3.129.3 Rubi [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.10, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$ , Rules used = {6316, 82, 241, 6314, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6316} \\
 & \frac{2 \int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx}{3d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{82} \\
 & \frac{2 \int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx}{3d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{241} \\
 & \frac{2 \int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{3/2}} dx}{3d} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{6cd^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} \\
 & \quad \downarrow \text{6314} \\
 & \frac{2 \left( \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right)}{3d} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}} + \\
 & \quad \frac{b\sqrt{cx-1}\sqrt{cx+1}}{6cd^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} \\
 & \quad \downarrow \text{240} \\
 & \frac{x(a + \operatorname{barccosh}(cx))}{3d(d - c^2 dx^2)^{3/2}} + \frac{2 \left( \frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \log(1-c^2x^2)}{2cd\sqrt{d-c^2dx^2}} \right)}{3d} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{6cd^2(1 - c^2x^2)\sqrt{d - c^2dx^2}}
 \end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d - c^2*d*x^2)^(5/2), x]`

```
output (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(6*c*d^2*(1 - c^2*x^2)*Sqrt[d - c^2*d*x^2
]) + (x*(a + b*ArcCosh[c*x]))/(3*d*(d - c^2*d*x^2)^(3/2)) + (2*((x*(a + b*
ArcCosh[c*x]))/(d*Sqrt[d - c^2*d*x^2]) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*L
og[1 - c^2*x^2])/(2*c*d*Sqrt[d - c^2*d*x^2])))/(3*d)
```

### 3.129.3.1 Defintions of rubi rules used

```
rule 82 Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)
)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d,
e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]
```

```
rule 240 Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x
^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]
```

```
rule 241 Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/
(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]
```

```
rule 6314 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2),
x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp
[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a
+ b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e},
x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]
```

```
rule 6316 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x
_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p +
1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*
ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 +
c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a
+ b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*
d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]
```

**3.129.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 476 vs.  $2(138) = 276$ .

Time = 1.01 (sec) , antiderivative size = 477, normalized size of antiderivative = 2.94

method	result
default	$a \left( \frac{x}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{2x}{3d^2\sqrt{-c^2dx^2+d}} \right) - \frac{b\sqrt{-d(c^2x^2-1)}(2c^3x^3-3cx+2\sqrt{cx-1}\sqrt{cx+1}c^2x^2-2\sqrt{cx-1}\sqrt{cx+1})(8\sqrt{cx-1}\sqrt{cx+1})}{(2c^3x^3-3cx+2\sqrt{cx-1}\sqrt{cx+1}c^2x^2-2\sqrt{cx-1}\sqrt{cx+1})(8\sqrt{cx-1}\sqrt{cx+1})}$
parts	$a \left( \frac{x}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{2x}{3d^2\sqrt{-c^2dx^2+d}} \right) - \frac{b\sqrt{-d(c^2x^2-1)}(2c^3x^3-3cx+2\sqrt{cx-1}\sqrt{cx+1}c^2x^2-2\sqrt{cx-1}\sqrt{cx+1})(8\sqrt{cx-1}\sqrt{cx+1})}{(2c^3x^3-3cx+2\sqrt{cx-1}\sqrt{cx+1}c^2x^2-2\sqrt{cx-1}\sqrt{cx+1})(8\sqrt{cx-1}\sqrt{cx+1})}$

input `int((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output `a*(1/3/d*x/(-c^2*d*x^2+d)^(3/2)+2/3/d^2*x/(-c^2*d*x^2+d)^(1/2))-1/6*b*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-3*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-2*(c*x-1)^(1/2)*(c*x+1)^(1/2))*(8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*(c*x+1)^(1/2))^2-1)*x^5*c^5-8*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^6*c^6-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^3*c^3+24*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^4*c^4+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-2*c^4*x^4+6*c^2*x^2*arccosh(c*x)+12*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x*c-24*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^2*c^2-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+4*c^2*x^2-8*arccosh(c*x)+8*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)-2)/(3*c^6*x^6-10*c^4*x^4+11*c^2*x^2-4)/d^3/c`

**3.129.5 Fracas [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

**3.129.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(-d(cx - 1)(cx + 1))^{\frac{5}{2}}} dx$$

input `integrate((a+b*acosh(c*x))/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral((a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`

**3.129.7 Maxima [A] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.97

$$\begin{aligned} \int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx &= \frac{1}{6} bc \left( \frac{\sqrt{-d}}{c^4 d^3 x^2 - c^2 d^3} + \frac{2\sqrt{-d} \log(cx + 1)}{c^2 d^3} + \frac{2\sqrt{-d} \log(cx - 1)}{c^2 d^3} \right) \\ &+ \frac{1}{3} b \left( \frac{2x}{\sqrt{-c^2 dx^2 + dd^2}} + \frac{x}{(-c^2 dx^2 + d)^{\frac{3}{2}} d} \right) \operatorname{arcosh}(cx) \\ &+ \frac{1}{3} a \left( \frac{2x}{\sqrt{-c^2 dx^2 + dd^2}} + \frac{x}{(-c^2 dx^2 + d)^{\frac{3}{2}} d} \right) \end{aligned}$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/6*b*c*(sqrt(-d)/(c^4*d^3*x^2 - c^2*d^3) + 2*sqrt(-d)*log(c*x + 1)/(c^2*d^3) + 2*sqrt(-d)*log(c*x - 1)/(c^2*d^3)) + 1/3*b*(2*x/(sqrt(-c^2*d*x^2 + d)*d^2) + x/((-c^2*d*x^2 + d)^(3/2)*d))*arccosh(c*x) + 1/3*a*(2*x/(sqrt(-c^2*d*x^2 + d)*d^2) + x/((-c^2*d*x^2 + d)^(3/2)*d))`

**3.129.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate((a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(-c^2*d*x^2 + d)^(5/2), x)`

---

3.129.  $\int \frac{a + \operatorname{barccosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx$

**3.129.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^(5/2), x)`output `int((a + b*acosh(c*x))/(d - c^2*d*x^2)^(5/2), x)`



**3.130**  $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)^{5/2}} dx$

3.130.1 Optimal result . . . . . 1144  
 3.130.2 Mathematica [A] (warning: unable to verify) . . . . . 1145  
 3.130.3 Rubi [A] (verified) . . . . . 1145  
 3.130.4 Maple [A] (verified) . . . . . 1150  
 3.130.5 Fracas [F] . . . . . 1150  
 3.130.6 Sympy [F] . . . . . 1151  
 3.130.7 Maxima [F] . . . . . 1151  
 3.130.8 Giac [F] . . . . . 1151  
 3.130.9 Mupad [F(-1)] . . . . . 1152

**3.130.1 Optimal result**

Integrand size = 27, antiderivative size = 317

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x(d - c^2dx^2)^{5/2}} dx = \frac{bcx\sqrt{-1 + cx}\sqrt{1 + cx}}{6d^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} + \frac{a + b\operatorname{arccosh}(cx)}{3d(d - c^2dx^2)^{3/2}}$$

$$+ \frac{a + b\operatorname{arccosh}(cx)}{d^2\sqrt{d - c^2dx^2}} + \frac{2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d - c^2dx^2}}$$

$$+ \frac{7b\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arctanh}(cx)}{6d^2\sqrt{d - c^2dx^2}} - \frac{ib\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d - c^2dx^2}}$$

$$+ \frac{ib\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d - c^2dx^2}}$$

```
output 1/3*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(3/2)+(a+b*arccosh(c*x))/d^2/(-c^2
*d*x^2+d)^(1/2)+1/6*b*c*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*x^2+1)/(-c
^2*d*x^2+d)^(1/2)+2*(a+b*arccosh(c*x))*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1
/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2)+7/6*b*arctanh(c*
x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2)-I*b*polylog(2,-I*(
c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*
x^2+d)^(1/2)+I*b*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1
/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2)
```

**3.130.2 Mathematica [A] (warning: unable to verify)**

Time = 7.07 (sec) , antiderivative size = 389, normalized size of antiderivative = 1.23

$$\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx = \sqrt{-d(-1 + c^2 x^2)} \left( \frac{a}{3d^3(-1 + c^2 x^2)^2} - \frac{a}{d^3(-1 + c^2 x^2)} \right) + \frac{a \log(x)}{d^{5/2}} - \frac{a \log\left(d + \sqrt{d} \sqrt{-d(-1 + c^2 x^2)}\right)}{d^{5/2}} + b \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \left( 14 \operatorname{arccosh}(cx) \coth\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) - \operatorname{csch}^2\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) - \frac{1}{2} \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \operatorname{arccosh}(cx) \right)$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^(5/2)),x]`

output

```
Sqrt[-(d*(-1 + c^2*x^2))]*(a/(3*d^3*(-1 + c^2*x^2)^2) - a/(d^3*(-1 + c^2*x^2))) + (a*Log[x])/d^(5/2) - (a*Log[d + Sqrt[d]*Sqrt[-(d*(-1 + c^2*x^2))]])/d^(5/2) + (b*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(14*ArcCosh[c*x]*Coth[ArcCosh[c*x]/2] - Csch[ArcCosh[c*x]/2]^2 - (Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Csch[ArcCosh[c*x]/2]^4)/2 - (24*I)*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] + (24*I)*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + 28*Log[Cosh[ArcCosh[c*x]/2]] - 28*Log[Sinh[ArcCosh[c*x]/2]] - (24*I)*PolyLog[2, (-I)/E^ArcCosh[c*x]] + (24*I)*PolyLog[2, I/E^ArcCosh[c*x]] - Sech[ArcCosh[c*x]/2]^2 - (8*ArcCosh[c*x]*Sinh[ArcCosh[c*x]/2]^4)/(((1 + c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3) - 14*ArcCosh[c*x]*Tanh[ArcCosh[c*x]/2]))/(24*d^2*Sqrt[-(d*(-1 + c*x)*(1 + c*x))])
```

**3.130.3 Rubi [A] (verified)**Time = 1.14 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.84, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.481$ , Rules used = {6351, 39, 215, 219, 6351, 25, 39, 219, 6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx$$

↓ 6351

---

3.130.  $\int \frac{a + \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{39} \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{215} \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{1}{1-c^2x^2} dx + \frac{x}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{219} \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6351} \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{25} \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{39} \\
& \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{219}
\end{aligned}$$

---

3.130.  $\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^{5/2}} dx$

$$\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx + \frac{a+b\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}}}{d} + \frac{a + \operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)}\right)}{3d^2\sqrt{d-c^2dx^2}}$$

↓ 6361

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} d\operatorname{arccosh}(cx) + \frac{a+b\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}}}{d} + \frac{a + \operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)}\right)}{3d^2\sqrt{d-c^2dx^2}}$$

↓ 3042

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx)) \csc\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+b\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)}\right)}{3d^2\sqrt{d-c^2dx^2}}$$

↓ 4668

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \left(-ib \int \log(1-ie^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + ib \int \log(1+ie^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)})\right) (a+b\operatorname{arccosh}(cx))}{d\sqrt{d-c^2dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)}\right)}{3d^2\sqrt{d-c^2dx^2}}$$

↓ 2715

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \left(-ib \int e^{-\operatorname{arccosh}(cx)} \log(1-ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1+ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2 \arctan(e^{\operatorname{arccosh}(cx)})\right) (a+b\operatorname{arccosh}(cx))}{d\sqrt{d-c^2dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)}\right)}{3d^2\sqrt{d-c^2dx^2}}$$

↓ 2838

---

3.130.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d-c^2dx^2)^{5/2}} dx$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}\left(2\arctan\left(e^{\operatorname{arccosh}(cx)}\right)(a+\operatorname{barccosh}(cx))-ib\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(cx)}\right)+ib\operatorname{PolyLog}\left(2,ie^{\operatorname{arccosh}(cx)}\right)\right)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}}$$

$$\frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)}\right)}{3d^2\sqrt{d-c^2dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d - c^2*d*x^2)^(5/2)),x]`

output `(a + b*ArcCosh[c*x])/(3*d*(d - c^2*d*x^2)^(3/2)) + (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(x/(2*(1 - c^2*x^2)) + ArcTanh[c*x]/(2*c)))/(3*d^2*Sqrt[d - c^2*d*x^2]) + ((a + b*ArcCosh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*ArcTanh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(d*Sqrt[d - c^2*d*x^2]))/d`

### 3.130.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 39 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_)^m)*((d_) + (e_
.)*(x_)^2)^p), x_Symbol] :> Simp[(-f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1
)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[
b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[
(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x]
)^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] &
& GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] ||
EqQ[n, 1])`

rule 6361 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*(x_)^m)/Sqrt[(d_) + (e_.
)*(x_)^2], x_Symbol] :> Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x
]/Sqrt[d + e*x^2])] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]]
, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && Int
egerQ[m]`

### 3.130.4 Maple [A] (verified)

Time = 1.33 (sec) , antiderivative size = 560, normalized size of antiderivative = 1.77

method	result
default	$\frac{a}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{a}{d^2\sqrt{-c^2dx^2+d}} - \frac{a \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{d^{\frac{5}{2}}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}(6c^2x^2 \operatorname{arccosh}(cx) - \sqrt{cx-1}\sqrt{cx+1})}{6d^3(c^2x^2-1)^2}\right)$
parts	$\frac{a}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{a}{d^2\sqrt{-c^2dx^2+d}} - \frac{a \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{d^{\frac{5}{2}}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}(6c^2x^2 \operatorname{arccosh}(cx) - \sqrt{cx-1}\sqrt{cx+1})}{6d^3(c^2x^2-1)^2}\right)$

input `int((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/3*a/d/(-c^2*d*x^2+d)^(3/2)+a/d^2/(-c^2*d*x^2+d)^(1/2)-a/d^(5/2)*\ln((2*d+ \\ & 2*d^(1/2)*(-c^2*d*x^2+d)^(1/2))/x)+b*(-1/6*(-d*(c^2*x^2-1))^(1/2)*(6*c^2*x \\ & ^2*\operatorname{arccosh}(c*x)-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-8*\operatorname{arccosh}(c*x))/d^3/(c^2*x \\ & ^2-1)^2-7/6*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/(c^2*x \\ & ^2-1)*\ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+7/6*(-d*(c^2*x^2-1))^(1/2)*(c*x \\ & -1)^(1/2)*(c*x+1)^(1/2)/d^3/(c^2*x^2-1)*\ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x \\ & -1)-I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/(c^2*x^2-1)*a \\ & \operatorname{rccosh}(c*x)*\ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+I*(-d*(c^2*x^2-1))^( \\ & 1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/(c^2*x^2-1)*\operatorname{dilog}(1+I*(c*x+(c*x-1)^( \\ & 1/2)*(c*x+1)^(1/2)))+I*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d \\ & ^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)*\ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-I*(- \\ & d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/(c^2*x^2-1)*\operatorname{dilog}(1-I \\ & *(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))) \end{aligned}$$

### 3.130.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(-c^2 dx^2 + d)^{5/2} x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^6*d^3*x^7 - 3*c^4*d^3*x^5 + 3*c^2*d^3*x^3 - d^3*x), x)`

**3.130.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(-d(cx - 1)(cx + 1))^{\frac{5}{2}}} dx$$

input `integrate((a+b*acosh(c*x))/x/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral((a + b*acosh(c*x))/(x*(-d*(c*x - 1)*(c*x + 1))**(5/2)), x)`

**3.130.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{5}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `-1/3*a*(3*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(5/2) - 3/(sqrt(-c^2*d*x^2 + d)*d^2) - 1/((-c^2*d*x^2 + d)^(3/2)*d) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(5/2)*x), x)`

**3.130.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{5}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(5/2)*x), x)`



**3.130.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^(5/2)),x)`output `int((a + b*acosh(c*x))/(x*(d - c^2*d*x^2)^(5/2)), x)`

### 3.131 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d-c^2dx^2)^{5/2}} dx$

3.131.1 Optimal result . . . . .	1153
3.131.2 Mathematica [A] (verified) . . . . .	1154
3.131.3 Rubi [A] (verified) . . . . .	1154
3.131.4 Maple [A] (verified) . . . . .	1156
3.131.5 Fracas [F] . . . . .	1157
3.131.6 Sympy [F(-1)] . . . . .	1157
3.131.7 Maxima [F] . . . . .	1157
3.131.8 Giac [F] . . . . .	1158
3.131.9 Mupad [F(-1)] . . . . .	1158

#### 3.131.1 Optimal result

Integrand size = 27, antiderivative size = 248

$$\int \frac{a + \operatorname{arccosh}(cx)}{x^2(d - c^2dx^2)^{5/2}} dx = -\frac{bc\sqrt{d - c^2dx^2}}{6d^3\sqrt{-1 + cx}\sqrt{1 + cx}(1 - c^2x^2)} - \frac{a + \operatorname{arccosh}(cx)}{dx(d - c^2dx^2)^{3/2}} + \frac{4c^2x(a + \operatorname{arccosh}(cx))}{3d(d - c^2dx^2)^{3/2}} + \frac{8c^2x(a + \operatorname{arccosh}(cx))}{3d^2\sqrt{d - c^2dx^2}} + \frac{bc\sqrt{d - c^2dx^2}\log(x)}{d^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{5bc\sqrt{d - c^2dx^2}\log(1 - c^2x^2)}{6d^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output (-a-b*arccosh(c*x))/d/x/(-c^2*d*x^2+d)^(3/2)+4/3*c^2*x*(a+b*arccosh(c*x))/d/(-c^2*d*x^2+d)^(3/2)+8/3*c^2*x*(a+b*arccosh(c*x))/d^2/(-c^2*d*x^2+d)^(1/2)-1/6*b*c*(-c^2*d*x^2+d)^(1/2)/d^3/(-c^2*x^2+1)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*c*ln(x)*(-c^2*d*x^2+d)^(1/2)/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/6*b*c*ln(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/d^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.131.2 Mathematica [A] (verified)**

Time = 0.26 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.70

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx = \frac{\sqrt{-1 + cx} \sqrt{1 + cx} \left( \frac{a + \operatorname{barccosh}(cx)}{x(-1 + cx)^{3/2}(1 + cx)^{3/2}} - \frac{4c^2 x(a + \operatorname{barccosh}(cx))}{3(-1 + cx)^{3/2}(1 + cx)^{3/2}} + \frac{8c^2 x(a + \operatorname{barccosh}(cx))}{3\sqrt{-1 + cx}\sqrt{1 + cx}} \right)}{d^2 \sqrt{d - c^2 dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^(5/2)), x]`output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((a + b*ArcCosh[c*x])/(x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) - (4*c^2*x*(a + b*ArcCosh[c*x]))/(3*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2)) + (8*c^2*x*(a + b*ArcCosh[c*x]))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - b*c*(1/(6*(-1 + c^2*x^2)) + Log[x] + (5*Log[1 - c^2*x^2])/6)))/(d^2*Sqrt[d - c^2*d*x^2])`**3.131.3 Rubi [A] (verified)**Time = 0.51 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.70, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6337, 27, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx \\ & \quad \downarrow \text{6337} \\ & -\frac{bc\sqrt{d - c^2 dx^2} \int -\frac{8c^4 x^4 - 12c^2 x^2 + 3}{3d^3 x(1 - c^2 x^2)^2} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{8c^2 x(a + \operatorname{barccosh}(cx))}{3d^2 \sqrt{d - c^2 dx^2}} + \frac{4c^2 x(a + \operatorname{barccosh}(cx))}{3d (d - c^2 dx^2)^{3/2}} - \\ & \quad \frac{a + \operatorname{barccosh}(cx)}{dx (d - c^2 dx^2)^{3/2}} \\ & \quad \downarrow \text{27} \\ & \frac{bc\sqrt{d - c^2 dx^2} \int \frac{8c^4 x^4 - 12c^2 x^2 + 3}{x(1 - c^2 x^2)^2} dx}{3d^3 \sqrt{cx - 1}\sqrt{cx + 1}} + \frac{8c^2 x(a + \operatorname{barccosh}(cx))}{3d^2 \sqrt{d - c^2 dx^2}} + \frac{4c^2 x(a + \operatorname{barccosh}(cx))}{3d (d - c^2 dx^2)^{3/2}} - \\ & \quad \frac{a + \operatorname{barccosh}(cx)}{dx (d - c^2 dx^2)^{3/2}} \end{aligned}$$

---

3.131.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \downarrow 1578 \\
& \frac{bc\sqrt{d-c^2dx^2} \int \frac{8c^4x^4-12c^2x^2+3}{x^2(1-c^2x^2)^2} dx^2}{6d^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{8c^2x(a+\operatorname{barccosh}(cx))}{3d^2\sqrt{d-c^2dx^2}} + \frac{4c^2x(a+\operatorname{barccosh}(cx))}{3d(d-c^2dx^2)^{3/2}} - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{dx(d-c^2dx^2)^{3/2}} \\
& \downarrow 1195 \\
& \frac{bc\sqrt{d-c^2dx^2} \int \left( \frac{5c^2}{c^2x^2-1} - \frac{c^2}{(c^2x^2-1)^2} + \frac{3}{x^2} \right) dx^2}{6d^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{8c^2x(a+\operatorname{barccosh}(cx))}{3d^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{4c^2x(a+\operatorname{barccosh}(cx))}{3d(d-c^2dx^2)^{3/2}} - \frac{a+\operatorname{barccosh}(cx)}{dx(d-c^2dx^2)^{3/2}} \\
& \downarrow 2009 \\
& \frac{8c^2x(a+\operatorname{barccosh}(cx))}{3d^2\sqrt{d-c^2dx^2}} + \frac{4c^2x(a+\operatorname{barccosh}(cx))}{3d(d-c^2dx^2)^{3/2}} - \frac{a+\operatorname{barccosh}(cx)}{dx(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{c^2x^2-1} + 5\log(1-c^2x^2) + 3\log(x^2) \right)}{6d^3\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*(d - c^2*d*x^2)^(5/2)),x]`

output `-((a + b*ArcCosh[c*x])/(d*x*(d - c^2*d*x^2)^(3/2))) + (4*c^2*x*(a + b*ArcCosh[c*x]))/(3*d*(d - c^2*d*x^2)^(3/2)) + (8*c^2*x*(a + b*ArcCosh[c*x]))/(3*d^2*sqrt[d - c^2*d*x^2]) + (b*c*sqrt[d - c^2*d*x^2]*((-1 + c^2*x^2)^(-1) + 3*Log[x^2] + 5*Log[1 - c^2*x^2]))/(6*d^3*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.131.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1195 `Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))^(n_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`

rule 1578 `Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.131.4 Maple [A] (verified)

Time = 1.26 (sec) , antiderivative size = 434, normalized size of antiderivative = 1.75

method	result
default	$a \left( -\frac{1}{dx(-c^2dx^2+d)^{\frac{3}{2}}} + 4c^2 \left( \frac{x}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{2x}{3d^2\sqrt{-c^2dx^2+d}} \right) \right) - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}}{(16\sqrt{cx-1}\sqrt{cx+1})}$
parts	$a \left( -\frac{1}{dx(-c^2dx^2+d)^{\frac{3}{2}}} + 4c^2 \left( \frac{x}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{2x}{3d^2\sqrt{-c^2dx^2+d}} \right) \right) - \frac{b\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}}{(16\sqrt{cx-1}\sqrt{cx+1})}$

input `int((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output `a*(-1/d/x/(-c^2*d*x^2+d)^(3/2)+4*c^2*(1/3/d*x/(-c^2*d*x^2+d)^(3/2)+2/3/d^2*x/(-c^2*d*x^2+d)^(1/2))-1/6*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(16*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4+16*arccosh(c*x)*c^5*x^5-6*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^5*c^5-10*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^5*c^5-24*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-32*c^3*x^3*arccosh(c*x)+12*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3+20*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^3*c^3-c^3*x^3+6*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+16*c*x*arccosh(c*x)-6*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-10*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x*c+c*x)/d^3/(c^6*x^6-3*c^4*x^4+3*c^2*x^2-1)/x`

**3.131.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{5}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^6*d^3*x^8 - 3*c^4*d^3*x^6 + 3*c^2*d^3*x^4 - d^3*x^2), x)`

**3.131.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**2/(-c**2*d*x**2+d)**(5/2),x)`

output `Timed out`

**3.131.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{5}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/3*a*(8*c^2*x/(sqrt(-c^2*d*x^2 + d)*d^2) + 4*c^2*x/((-c^2*d*x^2 + d)^(3/2)*d) - 3/((-c^2*d*x^2 + d)^(3/2)*d*x)) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(5/2)*x^2), x)`

**3.131.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{5}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(5/2)*x^2), x)`

**3.131.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2 (d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^(5/2)),x)`

output `int((a + b*acosh(c*x))/(x^2*(d - c^2*d*x^2)^(5/2)), x)`

### 3.132 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^{5/2}} dx$

3.132.1 Optimal result . . . . .	1159
3.132.2 Mathematica [A] (warning: unable to verify) . . . . .	1160
3.132.3 Rubi [A] (verified) . . . . .	1161
3.132.4 Maple [A] (verified) . . . . .	1167
3.132.5 Fricas [F] . . . . .	1168
3.132.6 Sympy [F(-1)] . . . . .	1168
3.132.7 Maxima [F] . . . . .	1169
3.132.8 Giac [F] . . . . .	1169
3.132.9 Mupad [F(-1)] . . . . .	1169

#### 3.132.1 Optimal result

Integrand size = 27, antiderivative size = 479

$$\begin{aligned}
\int \frac{a + b\operatorname{arccosh}(cx)}{x^3(d - c^2dx^2)^{5/2}} dx &= \frac{3bc\sqrt{-1 + cx}\sqrt{1 + cx}}{4d^2x\sqrt{d - c^2dx^2}} \\
&- \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{4d^2x(1 - c^2x^2)\sqrt{d - c^2dx^2}} + \frac{5bc^3x\sqrt{-1 + cx}\sqrt{1 + cx}}{12d^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} \\
&+ \frac{5c^2(a + b\operatorname{arccosh}(cx))}{6d(d - c^2dx^2)^{3/2}} - \frac{a + b\operatorname{arccosh}(cx)}{2dx^2(d - c^2dx^2)^{3/2}} + \frac{5c^2(a + b\operatorname{arccosh}(cx))}{2d^2\sqrt{d - c^2dx^2}} \\
&+ \frac{5c^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \arctan(e^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d - c^2dx^2}} \\
&+ \frac{13bc^2\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arctanh}(cx)}{6d^2\sqrt{d - c^2dx^2}} \\
&- \frac{5ibc^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{2d^2\sqrt{d - c^2dx^2}} \\
&+ \frac{5ibc^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{2d^2\sqrt{d - c^2dx^2}}
\end{aligned}$$



output  $5/6*c^2*(a+b*\operatorname{arccosh}(c*x))/d/(-c^2*d*x^2+d)^{(3/2)}+1/2*(-a-b*\operatorname{arccosh}(c*x))/d/x^2/(-c^2*d*x^2+d)^{(3/2)}+5/2*c^2*(a+b*\operatorname{arccosh}(c*x))/d^2/(-c^2*d*x^2+d)^{(1/2)}+3/4*b*c*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/x/(-c^2*d*x^2+d)^{(1/2)}-1/4*b*c*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/x/(-c^2*x^2+1)/(-c^2*d*x^2+d)^{(1/2)}+5/12*b*c^3*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*x^2+1)/(-c^2*d*x^2+d)^{(1/2)}+5*c^2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctan}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}+13/6*b*c^2*\operatorname{arctanh}(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-5/2*I*b*c^2*\operatorname{polylog}(2,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}+5/2*I*b*c^2*\operatorname{polylog}(2,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}$

### 3.132.2 Mathematica [A] (warning: unable to verify)

Time = 7.21 (sec) , antiderivative size = 534, normalized size of antiderivative = 1.11

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx = \sqrt{-d(-1 + c^2 x^2)} \left( -\frac{a}{2d^3 x^2} + \frac{ac^2}{3d^3 (-1 + c^2 x^2)^2} - \frac{2ac^2}{d^3 (-1 + c^2 x^2)} \right) + \frac{5ac^2 \log(x)}{2d^{5/2}} - \frac{5ac^2 \log\left(d + \sqrt{d} \sqrt{-d(-1 + c^2 x^2)}\right)}{2d^{5/2}} + \frac{bc^2 \left( \frac{6\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}{cx} + \frac{6(-1+cx)(1+cx)\operatorname{arccosh}(cx)}{c^2 x^2} + 26\operatorname{arccosh}(cx) \cosh^2\left(\frac{1}{2}\operatorname{arccosh}(cx)\right) - \coth\left(\frac{1}{2}\operatorname{arccosh}(cx)\right) \right)}{d^3}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^(5/2)),x]`

output  $\text{Sqrt}[-(d*(-1 + c^2*x^2))]*(-1/2*a/(d^3*x^2) + (a*c^2)/(3*d^3*(-1 + c^2*x^2)^2) - (2*a*c^2)/(d^3*(-1 + c^2*x^2))) + (5*a*c^2*\text{Log}[x])/(2*d^(5/2)) - (5*a*c^2*\text{Log}[d + \text{Sqrt}[d]*\text{Sqrt}[-(d*(-1 + c^2*x^2))]])/(2*d^(5/2)) + (b*c^2*((6*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x))/(c*x) + (6*(-1 + c*x)*(1 + c*x)*\text{ArcCosh}[c*x])/(c^2*x^2) + 26*\text{ArcCosh}[c*x]*\text{Cosh}[\text{ArcCosh}[c*x]/2]^2 - \text{Coth}[\text{ArcCosh}[c*x]/2] - \text{ArcCosh}[c*x]*\text{Coth}[\text{ArcCosh}[c*x]/2]^2 - (30*I)*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{ArcCosh}[c*x]*\text{Log}[1 - I/E^{\text{ArcCosh}[c*x]}] + (30*I)*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{ArcCosh}[c*x]*\text{Log}[1 + I/E^{\text{ArcCosh}[c*x]}] + 26*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{Log}[\text{Cosh}[\text{ArcCosh}[c*x]/2]] - 26*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{Log}[\text{Sinh}[\text{ArcCosh}[c*x]/2]] - (30*I)*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{PolyLog}[2, (-I)/E^{\text{ArcCosh}[c*x]}] + (30*I)*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{PolyLog}[2, I/E^{\text{ArcCosh}[c*x]}] - 26*\text{ArcCosh}[c*x]*\text{Sinh}[\text{ArcCosh}[c*x]/2]^2 - \text{Tanh}[\text{ArcCosh}[c*x]/2] - \text{ArcCosh}[c*x]*\text{Tanh}[\text{ArcCosh}[c*x]/2]^2))/(12*d^2*\text{Sqrt}[-(d*(-1 + c*x)*(1 + c*x))])$

### 3.132.3 Rubi [A] (verified)

Time = 1.67 (sec) , antiderivative size = 384, normalized size of antiderivative = 0.80, number of steps used = 19, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6347, 82, 253, 264, 219, 6351, 39, 215, 219, 6351, 25, 39, 219, 6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \text{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx$$

$$\downarrow 6347$$

$$\frac{5}{2}c^2 \int \frac{a + \text{barccosh}(cx)}{x (d - c^2 dx^2)^{5/2}} dx - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{x^2(1-cx)^2(cx+1)^2} dx}{2d^2\sqrt{d - c^2 dx^2}} - \frac{a + \text{barccosh}(cx)}{2dx^2 (d - c^2 dx^2)^{3/2}}$$

$$\downarrow 82$$

$$\frac{5}{2}c^2 \int \frac{a + \text{barccosh}(cx)}{x (d - c^2 dx^2)^{5/2}} dx - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{x^2(1-c^2x^2)^2} dx}{2d^2\sqrt{d - c^2 dx^2}} - \frac{a + \text{barccosh}(cx)}{2dx^2 (d - c^2 dx^2)^{3/2}}$$

$$\downarrow 253$$

---

3.132.  $\int \frac{a + \text{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{5}{2}c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{3}{2} \int \frac{1}{x^2(1 - c^2 x^2)} dx + \frac{1}{2x(1 - c^2 x^2)} \right)}{2d^2\sqrt{d - c^2 dx^2}} - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d - c^2 dx^2)^{3/2}} \\
& \quad \downarrow 264 \\
& \frac{5}{2}c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{3}{2} \left( c^2 \int \frac{1}{1 - c^2 x^2} dx - \frac{1}{x} \right) + \frac{1}{2x(1 - c^2 x^2)} \right)}{2d^2\sqrt{d - c^2 dx^2}} - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d - c^2 dx^2)^{3/2}} \\
& \quad \downarrow 219 \\
& \frac{\frac{5}{2}c^2 \int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{5/2}} dx - \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d - c^2 dx^2)^{3/2}} -}{2d^2\sqrt{d - c^2 dx^2}} \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{3}{2} (\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1 - c^2 x^2)} \right)}{2d^2\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow 6351 \\
& \frac{5}{2}c^2 \left( \frac{\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx}{d} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{(1 - cx)^2(cx + 1)^2} dx}{3d^2\sqrt{d - c^2 dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{3d(d - c^2 dx^2)^{3/2}} \right) - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d - c^2 dx^2)^{3/2}} - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{3}{2} (\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1 - c^2 x^2)} \right)}{2d^2\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow 39 \\
& \frac{5}{2}c^2 \left( \frac{\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx}{d} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{1}{(1 - c^2 x^2)^2} dx}{3d^2\sqrt{d - c^2 dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{3d(d - c^2 dx^2)^{3/2}} \right) - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d - c^2 dx^2)^{3/2}} - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{3}{2} (\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1 - c^2 x^2)} \right)}{2d^2\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow 215 \\
& \frac{5}{2}c^2 \left( \frac{\int \frac{a + \operatorname{barccosh}(cx)}{x(d - c^2 dx^2)^{3/2}} dx}{d} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{1}{2} \int \frac{1}{1 - c^2 x^2} dx + \frac{x}{2(1 - c^2 x^2)} \right)}{3d^2\sqrt{d - c^2 dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{3d(d - c^2 dx^2)^{3/2}} \right) - \\
& \quad \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d - c^2 dx^2)^{3/2}} - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{3}{2} (\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1 - c^2 x^2)} \right)}{2d^2\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow 219
\end{aligned}$$

---

3.132.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{5}{2}c^2 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) - \\
& \frac{a+\operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)} \right)}{2d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6351} \\
& \frac{5}{2}c^2 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{3d^2\sqrt{d-c^2dx^2}} \right) - \\
& \frac{a+\operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)} \right)}{2d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{25} \\
& \frac{5}{2}c^2 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{3d^2\sqrt{d-c^2dx^2}} \right) - \\
& \frac{a+\operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)} \right)}{2d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{39} \\
& \frac{5}{2}c^2 \left( \frac{\int \frac{a+\operatorname{barccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{1}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+\operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\operatorname{arctanh}(cx)}{2c} + \frac{x}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) - \\
& \frac{a+\operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)} \right)}{2d^2\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{219}
\end{aligned}$$

$$\frac{5}{2}c^2 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{a+b\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a + \operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a}{3d^2\sqrt{d-c^2dx^2}}\right)}{3d^2\sqrt{d-c^2dx^2}} \right) - \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)}\right)}{2d^2\sqrt{d-c^2dx^2}}$$

↓ 6361

$$\frac{5}{2}c^2 \left( \frac{\frac{\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{cx\sqrt{d-c^2dx^2}} d\operatorname{arccosh}(cx)}{d} + \frac{a+b\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}}}{d} + \frac{a + \operatorname{barccosh}(cx)}{3d(d-c^2dx^2)^{3/2}} + \dots \right) - \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)}\right)}{2d^2\sqrt{d-c^2dx^2}}$$

↓ 3042

$$\frac{5}{2}c^2 \left( \frac{\frac{\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx)) \operatorname{csc}\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{a+b\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{b\sqrt{cx-1}\sqrt{cx+1}\operatorname{arctanh}(cx)}{d\sqrt{d-c^2dx^2}}}{d} + \dots \right) - \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)}\right)}{2d^2\sqrt{d-c^2dx^2}}$$

↓ 4668

$$\frac{5}{2}c^2 \left( \frac{\frac{\sqrt{cx-1}\sqrt{cx+1} \left(-ib \int \log(1-ie^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + ib \int \log(1+ie^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2 \arctan(e^{\operatorname{arccosh}(cx)})\right) (a+b\operatorname{arccosh}(cx))}{d\sqrt{d-c^2dx^2}}}{d} + \dots \right) - \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)}\right)}{2d^2\sqrt{d-c^2dx^2}}$$

↓ 2715

$$\frac{5}{2}c^2 \left( \frac{\frac{\sqrt{cx-1}\sqrt{cx+1} \left(-ib \int e^{-\operatorname{arccosh}(cx)} \log(1-ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + ib \int e^{-\operatorname{arccosh}(cx)} \log(1+ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2 \arctan(e^{\operatorname{arccosh}(cx)})\right)}{d\sqrt{d-c^2dx^2}}}{d} + \dots \right) - \frac{a + \operatorname{barccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{3}{2}(\operatorname{carctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)}\right)}{2d^2\sqrt{d-c^2dx^2}}$$

---

3.132.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^{5/2}} dx$

↓ 2838

$$\frac{5}{2}c^2 \left( \frac{\frac{\sqrt{cx-1}\sqrt{cx+1} \left( 2 \arctan \left( e^{\operatorname{arccosh}(cx)} \right) (a + b \operatorname{arccosh}(cx)) - ib \operatorname{PolyLog} \left( 2, -ie^{\operatorname{arccosh}(cx)} \right) + ib \operatorname{PolyLog} \left( 2, ie^{\operatorname{arccosh}(cx)} \right) \right)}{d\sqrt{d-c^2dx^2}}}{d} + \frac{a + b \operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} \right) - \frac{a + b \operatorname{arccosh}(cx)}{2dx^2(d-c^2dx^2)^{3/2}} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{3}{2} (\operatorname{arctanh}(cx) - \frac{1}{x}) + \frac{1}{2x(1-c^2x^2)} \right)}{2d^2\sqrt{d-c^2dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*(d - c^2*d*x^2)^(5/2)), x]`

output `-1/2*(a + b*ArcCosh[c*x])/(d*x^2*(d - c^2*d*x^2)^(3/2)) - (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(1/(2*x*(1 - c^2*x^2)) + (3*(-x^(-1) + c*ArcTanh[c*x]))/2))/(2*d^2*Sqrt[d - c^2*d*x^2]) + (5*c^2*((a + b*ArcCosh[c*x])/(3*d*(d - c^2*d*x^2)^(3/2)) + (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(x/(2*(1 - c^2*x^2)) + ArcTanh[c*x]/(2*c)))/(3*d^2*Sqrt[d - c^2*d*x^2]) + ((a + b*ArcCosh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*ArcTanh[c*x])/(d*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(a + b*ArcCosh[c*x])*ArcTan[E^ArcCosh[c*x]] - I*b*PolyLog[2, (-I)*E^ArcCosh[c*x]] + I*b*PolyLog[2, I*E^ArcCosh[c*x]]))/(d*Sqrt[d - c^2*d*x^2]))/d)/2`

### 3.132.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 39 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

- rule 215  $\text{Int}[(a + b x^2)^p, x] \rightarrow \text{Simp}[(-x)(a + b x^2)^{p+1}/(2a(p+1)), x] + \text{Simp}[(2p+3)/(2a(p+1)) \text{Int}[(a + b x^2)^{p+1}, x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (\text{IntegerQ}[4p] \ || \ \text{IntegerQ}[6p])$
- rule 219  $\text{Int}[(a + b x^2)^{-1}, x] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \ \text{Rt}[-b, 2])) \ \text{ArcTanh}[\text{Rt}[-b, 2] \ (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 253  $\text{Int}[(c x)^m (a + b x^2)^p, x] \rightarrow \text{Simp}[(-c x)^{m+1} (a + b x^2)^{p+1}/(2a c (p+1)), x] + \text{Simp}[(m + 2p + 3)/(2a(p+1)) \text{Int}[(c x)^m (a + b x^2)^{p+1}, x], x] /; \text{FreeQ}\{a, b, c, m, x\} \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 264  $\text{Int}[(c x)^m (a + b x^2)^p, x] \rightarrow \text{Simp}[(c x)^{m+1} (a + b x^2)^{p+1}/(a c (m+1)), x] - \text{Simp}[b (m + 2p + 3)/(a c^2 (m+1)) \text{Int}[(c x)^{m+2} (a + b x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$
- rule 2715  $\text{Int}[\text{Log}[a + b x^2]^n, x] \rightarrow \text{Simp}[1/(d e^n \text{Log}[F]) \ \text{Subst}[\text{Int}[\text{Log}[a + b x]/x, x], x, (F^{e(c + d x)})^n], x] /; \text{FreeQ}\{F, a, b, c, d, e, n, x\} \ \&\& \ \text{GtQ}[a, 0]$
- rule 2838  $\text{Int}[\text{Log}[(d + e x^n)/x], x] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c) e x^n/n, x] /; \text{FreeQ}\{c, d, e, n, x\} \ \&\& \ \text{EqQ}[c d, 1]$
- rule 3042  $\text{Int}[u, x] \rightarrow \text{Int}[\text{DeactivateTrig}[u, x], x] /; \text{FunctionOfTrigOfLinearQ}[u, x]$
- rule 4668  $\text{Int}[\csc[e + \text{Pi} k + \text{Complex}[0, fz]](f x)^m, x] \rightarrow \text{Simp}[-2(c + d x)^m (\text{ArcTanh}[E^{(-I)e + f fz x}]/E^{I k \text{Pi}})/(f fz I), x] + (-\text{Simp}[d(m/(f fz I)) \text{Int}[(c + d x)^{m-1} \text{Log}[1 - E^{(-I)e + f fz x}]/E^{I k \text{Pi}}], x], x] + \text{Simp}[d(m/(f fz I)) \text{Int}[(c + d x)^{m-1} \text{Log}[1 + E^{(-I)e + f fz x}]/E^{I k \text{Pi}}], x], x) /; \text{FreeQ}\{c, d, e, f, fz, x\} \ \&\& \ \text{IntegerQ}[2k] \ \&\& \ \text{IGtQ}[m, 0]$

```
rule 6347 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1)))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]
```

```
rule 6351 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1))] Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])
```

```
rule 6361 Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]
```

### 3.132.4 Maple [A] (verified)

Time = 1.22 (sec) , antiderivative size = 655, normalized size of antiderivative = 1.37

method	result
default	$-\frac{a}{2dx^2(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{5ac^2}{6d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{5ac^2}{2d^2\sqrt{-c^2dx^2+d}} - \frac{5ac^2 \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{2d^{\frac{5}{2}}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}}{(15c}\right.$
parts	$-\frac{a}{2dx^2(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{5ac^2}{6d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{5ac^2}{2d^2\sqrt{-c^2dx^2+d}} - \frac{5ac^2 \ln\left(\frac{2d+2\sqrt{d}\sqrt{-c^2dx^2+d}}{x}\right)}{2d^{\frac{5}{2}}} + b\left(-\frac{\sqrt{-d(c^2x^2-1)}}{(15c}\right.$

```
input int((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)
```

3.132.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^{5/2}} dx$



output 
$$-1/2*a/d/x^2/(-c^2*d*x^2+d)^{(3/2)}+5/6*a*c^2/d/(-c^2*d*x^2+d)^{(3/2)}+5/2*a*c^2/d^2/(-c^2*d*x^2+d)^{(1/2)}-5/2*a*c^2/d^{(5/2)}*\ln((2*d+2*d^{(1/2)}*(-c^2*d*x^2+d)^{(1/2)})/x)+b*(-1/6*(-d*(c^2*x^2-1))^{(1/2)}*(15*c^4*x^4*\operatorname{arccosh}(c*x)+2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^3*x^3-20*c^2*x^2*\operatorname{arccosh}(c*x)-3*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+3*\operatorname{arccosh}(c*x))/d^3/(c^4*x^4-2*c^2*x^2+1)/x^2-13/6*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/(c^2*x^2-1)*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*c^2+13/6*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/(c^2*x^2-1)*\ln(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*c^2+5/2*I*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)*\ln(1-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*c^2+5/2*I*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)*\ln(1+I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*c^2$$

### 3.132.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(-c^2 dx^2 + d)^{5/2} x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(5/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^6*d^3*x^9 - 3*c^4*d^3*x^7 + 3*c^2*d^3*x^5 - d^3*x^3), x)`

### 3.132.6 Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**3/(-c**2*d*x**2+d)**(5/2),x)`

output Timed out

---

3.132.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d-c^2dx^2)^{5/2}} dx$

**3.132.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{5/2} x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `-1/6*a*(15*c^2*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(5/2) - 15*c^2/(sqrt(-c^2*d*x^2 + d)*d^2) - 5*c^2/((-c^2*d*x^2 + d)^(3/2)*d) + 3/((-c^2*d*x^2 + d)^(3/2)*d*x^2)) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/((-c^2*d*x^2 + d)^(5/2)*x^3), x)`

**3.132.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{5/2} x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(5/2)*x^3), x)`

**3.132.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^(5/2)),x)`

output `int((a + b*acosh(c*x))/(x^3*(d - c^2*d*x^2)^(5/2)), x)`

### 3.133 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^4(d-c^2dx^2)^{5/2}} dx$

3.133.1 Optimal result . . . . .	1170
3.133.2 Mathematica [A] (verified) . . . . .	.1171
3.133.3 Rubi [A] (verified) . . . . .	.1171
3.133.4 Maple [A] (verified) . . . . .	1173
3.133.5 Fracas [F] . . . . .	.1174
3.133.6 Sympy [F(-1)] . . . . .	.1174
3.133.7 Maxima [A] (verification not implemented) . . . . .	.1174
3.133.8 Giac [F] . . . . .	1175
3.133.9 Mupad [F(-1)] . . . . .	1175

#### 3.133.1 Optimal result

Integrand size = 27, antiderivative size = 338

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x^4(d - c^2dx^2)^{5/2}} dx = -\frac{bc\sqrt{d - c^2dx^2}}{6d^3x^2\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bc^3\sqrt{d - c^2dx^2}}{6d^3\sqrt{-1 + cx}\sqrt{1 + cx}(1 - c^2x^2)} - \frac{a + b\operatorname{arccosh}(cx)}{3dx^3(d - c^2dx^2)^{3/2}} - \frac{2c^2(a + b\operatorname{arccosh}(cx))}{dx(d - c^2dx^2)^{3/2}} + \frac{8c^4x(a + b\operatorname{arccosh}(cx))}{3d(d - c^2dx^2)^{3/2}} + \frac{16c^4x(a + b\operatorname{arccosh}(cx))}{3d^2\sqrt{d - c^2dx^2}} + \frac{8bc^3\sqrt{d - c^2dx^2} \log(x)}{3d^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{4bc^3\sqrt{d - c^2dx^2} \log(1 - c^2x^2)}{3d^3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output  $\frac{1}{3}*(-a-b*\operatorname{arccosh}(c*x))/d/x^3/(-c^2*d*x^2+d)^{(3/2)}-2*c^2*(a+b*\operatorname{arccosh}(c*x))/d/x/(-c^2*d*x^2+d)^{(3/2)}+8/3*c^4*x*(a+b*\operatorname{arccosh}(c*x))/d/(-c^2*d*x^2+d)^{(3/2)}+16/3*c^4*x*(a+b*\operatorname{arccosh}(c*x))/d^2/(-c^2*d*x^2+d)^{(1/2)}-1/6*b*c*(-c^2*d*x^2+d)^{(1/2)}/d^3/x^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/6*b*c^3*(-c^2*d*x^2+d)^{(1/2)}/d^3/(-c^2*x^2+1)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+8/3*b*c^3*\ln(x)*(-c^2*d*x^2+d)^{(1/2)}/d^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+4/3*b*c^3*\ln(-c^2*x^2+1)*(-c^2*d*x^2+d)^{(1/2)}/d^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

**3.133.2 Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.70

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx = \frac{2a + 12ac^2x^2 - 48ac^4x^4 + 32ac^6x^6 - bcx\sqrt{-1 + cx}\sqrt{1 + cx} + 2b(1 + 6c^2x^2 - 24c^4x^4)}{x^4 (d - c^2 dx^2)^{5/2}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^(5/2)),x]`

output `(2*a + 12*a*c^2*x^2 - 48*a*c^4*x^4 + 32*a*c^6*x^6 - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + 2*b*(1 + 6*c^2*x^2 - 24*c^4*x^4 + 16*c^6*x^6)*ArcCosh[c*x] - 16*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1 + c^2*x^2)*Log[x] + 8*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2] - 8*b*c^5*x^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2])/(6*d^2*x^3*(-1 + c^2*x^2)*Sqrt[d - c^2*d*x^2])`

**3.133.3 Rubi [A] (verified)**Time = 0.67 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.67, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6337, 27, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx$$

↓ 6337

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{16c^6 x^6 - 24c^4 x^4 + 6c^2 x^2 + 1}{3d^3 x^3 (1 - c^2 x^2)^2} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{2c^2(a + \operatorname{barccosh}(cx))}{dx (d - c^2 dx^2)^{3/2}} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3 (d - c^2 dx^2)^{3/2}} + \frac{16c^4 x(a + \operatorname{barccosh}(cx))}{3d^2 \sqrt{d - c^2 dx^2}} + \frac{8c^4 x(a + \operatorname{barccosh}(cx))}{3d (d - c^2 dx^2)^{3/2}}$$

↓ 27

$$\frac{bc\sqrt{d - c^2 dx^2} \int \frac{16c^6 x^6 - 24c^4 x^4 + 6c^2 x^2 + 1}{x^3 (1 - c^2 x^2)^2} dx}{3d^3 \sqrt{cx - 1}\sqrt{cx + 1}} - \frac{2c^2(a + \operatorname{barccosh}(cx))}{dx (d - c^2 dx^2)^{3/2}} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3 (d - c^2 dx^2)^{3/2}} + \frac{16c^4 x(a + \operatorname{barccosh}(cx))}{3d^2 \sqrt{d - c^2 dx^2}} + \frac{8c^4 x(a + \operatorname{barccosh}(cx))}{3d (d - c^2 dx^2)^{3/2}}$$

---

3.133.  $\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \downarrow \text{2331} \\
& \frac{bc\sqrt{d-c^2dx^2} \int \frac{16c^6x^6-24c^4x^4+6c^2x^2+1}{x^4(1-c^2x^2)^2} dx^2}{6d^3\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2(a+\operatorname{barccosh}(cx))}{dx(d-c^2dx^2)^{3/2}} - \frac{a+\operatorname{barccosh}(cx)}{3dx^3(d-c^2dx^2)^{3/2}} + \\
& \quad \frac{16c^4x(a+\operatorname{barccosh}(cx))}{3d^2\sqrt{d-c^2dx^2}} + \frac{8c^4x(a+\operatorname{barccosh}(cx))}{3d(d-c^2dx^2)^{3/2}} \\
& \downarrow \text{2123} \\
& \frac{bc\sqrt{d-c^2dx^2} \int \left( \frac{8c^4}{c^2x^2-1} - \frac{c^4}{(c^2x^2-1)^2} + \frac{8c^2}{x^2} + \frac{1}{x^4} \right) dx^2}{6d^3\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2(a+\operatorname{barccosh}(cx))}{dx(d-c^2dx^2)^{3/2}} - \\
& \quad \frac{a+\operatorname{barccosh}(cx)}{3dx^3(d-c^2dx^2)^{3/2}} + \frac{16c^4x(a+\operatorname{barccosh}(cx))}{3d^2\sqrt{d-c^2dx^2}} + \frac{8c^4x(a+\operatorname{barccosh}(cx))}{3d(d-c^2dx^2)^{3/2}} \\
& \downarrow \text{2009} \\
& -\frac{2c^2(a+\operatorname{barccosh}(cx))}{dx(d-c^2dx^2)^{3/2}} - \frac{a+\operatorname{barccosh}(cx)}{3dx^3(d-c^2dx^2)^{3/2}} + \frac{16c^4x(a+\operatorname{barccosh}(cx))}{3d^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{8c^4x(a+\operatorname{barccosh}(cx))}{3d(d-c^2dx^2)^{3/2}} + \frac{bc\sqrt{d-c^2dx^2} \left( -\frac{c^2}{1-c^2x^2} + 8c^2 \log(x^2) + 8c^2 \log(1-c^2x^2) - \frac{1}{x^2} \right)}{6d^3\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^4*(d - c^2*d*x^2)^(5/2)),x]`

output `-1/3*(a + b*ArcCosh[c*x])/(d*x^3*(d - c^2*d*x^2)^(3/2)) - (2*c^2*(a + b*ArcCosh[c*x]))/(d*x*(d - c^2*d*x^2)^(3/2)) + (8*c^4*x*(a + b*ArcCosh[c*x]))/(3*d*(d - c^2*d*x^2)^(3/2)) + (16*c^4*x*(a + b*ArcCosh[c*x]))/(3*d^2*sqrt[d - c^2*d*x^2]) + (b*c*sqrt[d - c^2*d*x^2]*(-x^(-2) - c^2/(1 - c^2*x^2) + 8*c^2*Log[x^2] + 8*c^2*Log[1 - c^2*x^2]))/(6*d^3*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.133.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2123 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol]
:> Int[ExpandIntegrand[Px*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c,
d, m, n}, x] && PolyQ[Px, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

rule 2331 `Int[(Pq_)*(x_)^((m_.)*((a_.) + (b_.)*(x_)^2)^(p_.), x_Symbol]
:> Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /;
FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`

rule 6337 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*(x_)^((m_.)*((d_.) + (e_.)*(x_)^2)^(p_.),
x_Symbol]
:> With[{u = IntHide[x^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[SimplifyIntegrand[u/Sqrt[d + e*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IntegerQ[p - 1/2] && NeQ[p, -2^(-1)] && (IGtQ[(m + 1)/2, 0] || ILtQ[(m + 2*p + 3)/2, 0])`

### 3.133.4 Maple [A] (verified)

Time = 1.15 (sec) , antiderivative size = 403, normalized size of antiderivative = 1.19

method	result
default	$a \left( -\frac{1}{3dx^3(-c^2dx^2+d)^{\frac{3}{2}}} + 2c^2 \left( -\frac{1}{dx(-c^2dx^2+d)^{\frac{3}{2}}} + 4c^2 \left( \frac{x}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{2x}{3d^2\sqrt{-c^2dx^2+d}} \right) \right) \right) - \frac{b\sqrt{-d(c^2x^2+d)}}{d^2}$
parts	$a \left( -\frac{1}{3dx^3(-c^2dx^2+d)^{\frac{3}{2}}} + 2c^2 \left( -\frac{1}{dx(-c^2dx^2+d)^{\frac{3}{2}}} + 4c^2 \left( \frac{x}{3d(-c^2dx^2+d)^{\frac{3}{2}}} + \frac{2x}{3d^2\sqrt{-c^2dx^2+d}} \right) \right) \right) - \frac{b\sqrt{-d(c^2x^2+d)}}{d^2}$

input `int((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output `a*(-1/3/d/x^3/(-c^2*d*x^2+d)^(3/2)+2*c^2*(-1/d/x/(-c^2*d*x^2+d)^(3/2)+4*c^2*(1/3/d*x/(-c^2*d*x^2+d)^(3/2)+2/3/d^2*x/(-c^2*d*x^2+d)^(1/2)))-1/6*b*(-d*(c^2*x^2-1)^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(32*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^6*x^6+32*arccosh(c*x)*c^7*x^7-16*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^4-1)*x^7*c^7-48*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4-64*arccosh(c*x)*c^5*x^5+32*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^4-1)*x^5*c^5+12*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2+32*c^3*x^3*arccosh(c*x)-16*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^4-1)*x^3*c^3-c^3*x^3+2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)/d^3/(c^6*x^6-3*c^4*x^4+3*c^2*x^2-1)/x^3`

3.133. 
$$\int \frac{a+b\operatorname{arccosh}(cx)}{x^4(d-c^2dx^2)^{5/2}} dx$$

**3.133.5 Fracas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{\frac{5}{2}} x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(5/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)/(c^6*d^3*x^10 - 3*c^4*d^3*x^8 + 3*c^2*d^3*x^6 - d^3*x^4), x)`

**3.133.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**4/(-c**2*d*x**2+d)**(5/2),x)`

output `Timed out`

**3.133.7 Maxima [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.82

$$\begin{aligned} \int \frac{a + \operatorname{barccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx &= \frac{1}{6} bc \left( \frac{8 c^2 \sqrt{-d} \log(cx + 1)}{d^3} + \frac{8 c^2 \sqrt{-d} \log(cx - 1)}{d^3} + \frac{16 c^2 \sqrt{-d} \log(x)}{d^3} + \frac{\sqrt{-d}}{c^2 d^3 x^4} \right) \\ &+ \frac{1}{3} \left( \frac{16 c^4 x}{\sqrt{-c^2 dx^2 + dd^2}} + \frac{8 c^4 x}{(-c^2 dx^2 + d)^{\frac{3}{2}} d} - \frac{6 c^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} dx} - \frac{1}{(-c^2 dx^2 + d)^{\frac{3}{2}} dx^3} \right) b \operatorname{arcosh}(cx) \\ &+ \frac{1}{3} \left( \frac{16 c^4 x}{\sqrt{-c^2 dx^2 + dd^2}} + \frac{8 c^4 x}{(-c^2 dx^2 + d)^{\frac{3}{2}} d} - \frac{6 c^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} dx} - \frac{1}{(-c^2 dx^2 + d)^{\frac{3}{2}} dx^3} \right) a \end{aligned}$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output  $1/6*b*c*(8*c^2*\sqrt{-d}*\log(c*x + 1)/d^3 + 8*c^2*\sqrt{-d}*\log(c*x - 1)/d^3 + 16*c^2*\sqrt{-d}*\log(x)/d^3 + \sqrt{-d}/(c^2*d^3*x^4 - d^3*x^2)) + 1/3*(16*c^4*x/(\sqrt{-c^2*d*x^2 + d}*d^2) + 8*c^4*x/((-c^2*d*x^2 + d)^{(3/2)*d}) - 6*c^2/((-c^2*d*x^2 + d)^{(3/2)*d*x}) - 1/((-c^2*d*x^2 + d)^{(3/2)*d*x^3}))*b*a$   
 $\text{rccosh}(c*x) + 1/3*(16*c^4*x/(\sqrt{-c^2*d*x^2 + d}*d^2) + 8*c^4*x/((-c^2*d*x^2 + d)^{(3/2)*d}) - 6*c^2/((-c^2*d*x^2 + d)^{(3/2)*d*x}) - 1/((-c^2*d*x^2 + d)^{(3/2)*d*x^3}))*a$

### 3.133.8 Giac [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(-c^2 dx^2 + d)^{5/2} x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((-c^2*d*x^2 + d)^(5/2)*x^4), x)`

### 3.133.9 Mupad [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4 (d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^(5/2)),x)`

output `int((a + b*acosh(c*x))/(x^4*(d - c^2*d*x^2)^(5/2)), x)`



### 3.134 $\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{7/2}} dx$

3.134.1 Optimal result . . . . .	1176
3.134.2 Mathematica [A] (verified) . . . . .	1177
3.134.3 Rubi [A] (verified) . . . . .	1177
3.134.4 Maple [A] (verified) . . . . .	1180
3.134.5 Fracas [F] . . . . .	1180
3.134.6 Sympy [F(-1)] . . . . .	1181
3.134.7 Maxima [A] (verification not implemented) . . . . .	1181
3.134.8 Giac [A] (verification not implemented) . . . . .	1182
3.134.9 Mupad [F(-1)] . . . . .	1182

#### 3.134.1 Optimal result

Integrand size = 20, antiderivative size = 246

$$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{7/2}} dx = \frac{\sqrt{-1+ax}\sqrt{1+ax}}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} + \frac{2\sqrt{-1+ax}\sqrt{1+ax}}{15ac^3(1-a^2x^2)\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)}{5c(c-a^2cx^2)^{5/2}} + \frac{4x\operatorname{arccosh}(ax)}{15c^2(c-a^2cx^2)^{3/2}} + \frac{8x\operatorname{arccosh}(ax)}{15c^3\sqrt{c-a^2cx^2}} - \frac{4\sqrt{-1+ax}\sqrt{1+ax}\log(1-a^2x^2)}{15ac^3\sqrt{c-a^2cx^2}}$$

```
output 1/5*x*arccosh(a*x)/c/(-a^2*c*x^2+c)^(5/2)+4/15*x*arccosh(a*x)/c^2/(-a^2*c*x^2+c)^(3/2)+8/15*x*arccosh(a*x)/c^3/(-a^2*c*x^2+c)^(1/2)+1/20*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/c^3/(-a^2*x^2+1)^2/(-a^2*c*x^2+c)^(1/2)+2/15*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/c^3/(-a^2*x^2+1)/(-a^2*c*x^2+c)^(1/2)-4/15*ln(-a^2*x^2+1)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/c^3/(-a^2*c*x^2+c)^(1/2)
```

**3.134.2 Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.47

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{7/2}} dx = \frac{4ax(15 - 20a^2x^2 + 8a^4x^4) \operatorname{arccosh}(ax) + \sqrt{-1 + ax}\sqrt{1 + ax} \left(11 - 8a^2x^2 - 16(-1 + \dots)\right)}{60ac^3(-1 + a^2x^2)^2 \sqrt{c - a^2cx^2}}$$

input `Integrate[ArcCosh[a*x]/(c - a^2*c*x^2)^(7/2),x]`output `(4*a*x*(15 - 20*a^2*x^2 + 8*a^4*x^4)*ArcCosh[a*x] + Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(11 - 8*a^2*x^2 - 16*(-1 + a^2*x^2)^2*Log[1 - a^2*x^2]))/(60*a*c^3*(-1 + a^2*x^2)^2*Sqrt[c - a^2*c*x^2])`**3.134.3 Rubi [A] (verified)**Time = 0.64 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.05, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$ , Rules used = {6316, 25, 82, 241, 6316, 82, 241, 6314, 240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{7/2}} dx \\ & \quad \downarrow \text{6316} \\ & \frac{4 \int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{5/2}} dx}{5c} - \frac{a\sqrt{ax - 1}\sqrt{ax + 1} \int -\frac{x}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c - a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)}{5c(c - a^2cx^2)^{5/2}} \\ & \quad \downarrow \text{25} \\ & \frac{4 \int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{5/2}} dx}{5c} + \frac{a\sqrt{ax - 1}\sqrt{ax + 1} \int \frac{x}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c - a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)}{5c(c - a^2cx^2)^{5/2}} \\ & \quad \downarrow \text{82} \\ & \frac{4 \int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{5/2}} dx}{5c} + \frac{a\sqrt{ax - 1}\sqrt{ax + 1} \int \frac{x}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c - a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)}{5c(c - a^2cx^2)^{5/2}} \\ & \quad \downarrow \text{241} \end{aligned}$$

---

3.134.  $\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{7/2}} dx$

$$\begin{aligned}
& \frac{4 \int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{5/2}} dx}{5c} + \frac{x \operatorname{arccosh}(ax)}{5c(c-a^2cx^2)^{5/2}} + \frac{\sqrt{ax-1}\sqrt{ax+1}}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} \\
& \quad \downarrow \text{6316} \\
& \frac{4 \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{3/2}} dx}{3c} + \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-ax)^2(ax+1)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)}{3c(c-a^2cx^2)^{3/2}} \right)}{5c} + \frac{x \operatorname{arccosh}(ax)}{5c(c-a^2cx^2)^{5/2}} + \\
& \quad \frac{\sqrt{ax-1}\sqrt{ax+1}}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} \\
& \quad \downarrow \text{82} \\
& \frac{4 \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{3/2}} dx}{3c} + \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)}{3c(c-a^2cx^2)^{3/2}} \right)}{5c} + \frac{x \operatorname{arccosh}(ax)}{5c(c-a^2cx^2)^{5/2}} + \\
& \quad \frac{\sqrt{ax-1}\sqrt{ax+1}}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} \\
& \quad \downarrow \text{241} \\
& \frac{4 \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{3/2}} dx}{3c} + \frac{x \operatorname{arccosh}(ax)}{3c(c-a^2cx^2)^{3/2}} + \frac{\sqrt{ax-1}\sqrt{ax+1}}{6ac^2(1-a^2x^2)\sqrt{c-a^2cx^2}} \right)}{5c} + \frac{x \operatorname{arccosh}(ax)}{5c(c-a^2cx^2)^{5/2}} + \\
& \quad \frac{\sqrt{ax-1}\sqrt{ax+1}}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} \\
& \quad \downarrow \text{6314} \\
& \frac{4 \left( \frac{2 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x \operatorname{arccosh}(ax)}{3c(c-a^2cx^2)^{3/2}} + \frac{\sqrt{ax-1}\sqrt{ax+1}}{6ac^2(1-a^2x^2)\sqrt{c-a^2cx^2}} \right)}{5c} + \\
& \quad \frac{x \operatorname{arccosh}(ax)}{5c(c-a^2cx^2)^{5/2}} + \frac{\sqrt{ax-1}\sqrt{ax+1}}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} \\
& \quad \downarrow \text{240}
\end{aligned}$$

---

3.134.  $\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{7/2}} dx$

$$4 \left( \frac{\operatorname{arccosh}(ax)}{3c(c-a^2cx^2)^{3/2}} + \frac{2 \left( \frac{\operatorname{arccosh}(ax)}{c\sqrt{c-a^2cx^2}} - \frac{\sqrt{ax-1}\sqrt{ax+1} \log(1-a^2x^2)}{2ac\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{\sqrt{ax-1}\sqrt{ax+1}}{6ac^2(1-a^2x^2)\sqrt{c-a^2cx^2}} \right) + \frac{\operatorname{arccosh}(ax)}{5c(c-a^2cx^2)^{5/2}} + \frac{5c\sqrt{ax-1}\sqrt{ax+1}}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}}$$

input `Int[ArcCosh[a*x]/(c - a^2*c*x^2)^(7/2), x]`

output `(Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(20*a*c^3*(1 - a^2*x^2)^2*Sqrt[c - a^2*c*x^2]) + (x*ArcCosh[a*x])/(5*c*(c - a^2*c*x^2)^(5/2)) + (4*((Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(6*a*c^2*(1 - a^2*x^2)*Sqrt[c - a^2*c*x^2]) + (x*ArcCosh[a*x])/(3*c*(c - a^2*c*x^2)^(3/2)) + (2*((x*ArcCosh[a*x])/(c*Sqrt[c - a^2*c*x^2]) - (Sqrt[-1 + a*x]*Sqrt[1 + a*x]*Log[1 - a^2*x^2])/(2*a*c*Sqrt[c - a^2*c*x^2])))/(3*c)))/(5*c)`

### 3.134.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

### 3.134.4 Maple [A] (verified)

Time = 1.37 (sec) , antiderivative size = 419, normalized size of antiderivative = 1.70

method	result
default	$-\frac{16\sqrt{-c(a^2x^2-1)}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{15c^4a(a^2x^2-1)} - \frac{\sqrt{-c(a^2x^2-1)}(8a^5x^5-20a^3x^3-8\sqrt{ax-1}\sqrt{ax+1}a^4x^4+15ax+16a^2x^2\sqrt{ax-1})}{15c^4a(a^2x^2-1)}$

input `int(arccosh(a*x)/(-a^2*c*x^2+c)^(7/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -16/15*(-c*(a^2*x^2-1))^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/c^4/a/(a^2*x^2-1) \\ & )*\operatorname{arccosh}(a*x)-1/60*(-c*(a^2*x^2-1))^(1/2)*(8*a^5*x^5-20*a^3*x^3-8*(a*x-1) \\ & ^{(1/2)}*(a*x+1)^(1/2)*a^4*x^4+15*a*x+16*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2) \\ & -8*(a*x-1)^(1/2)*(a*x+1)^(1/2))*(-64*(a*x+1)^(1/2)*(a*x-1)^(1/2)*a^7*x^7-6 \\ & 4*a^8*x^8+248*x^5*a^5*(a*x-1)^(1/2)*(a*x+1)^(1/2)+280*a^6*x^6+160*a^4*x^4* \\ & \operatorname{arccosh}(a*x)-340*a^3*x^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)-456*a^4*x^4-380*a^2*x \\ & ^2*\operatorname{arccosh}(a*x)+165*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+328*a^2*x^2+256*\operatorname{arccos} \\ & h(a*x)-88)/(40*a^10*x^10-215*a^8*x^8+469*a^6*x^6-517*a^4*x^4+287*a^2*x^2-6 \\ & 4)/c^4/a+8/15*(-c*(a^2*x^2-1))^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/c^4/a/(a^ \\ & 2*x^2-1)*\ln((a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))^2-1) \end{aligned}$$

### 3.134.5 Fricas [F]

$$\int \frac{\operatorname{arccosh}(ax)}{(c-a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{arccosh}(ax)}{(-a^2cx^2+c)^{7/2}} dx$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^(7/2),x, algorithm="fricas")`

output `integral(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)/(a^8*c^4*x^8 - 4*a^6*c^4*x^6 + 6*a^4*c^4*x^4 - 4*a^2*c^4*x^2 + c^4), x)`

### 3.134.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{7/2}} dx = \text{Timed out}$$

input `integrate(acosh(a*x)/(-a**2*c*x**2+c)**(7/2), x)`

output Timed out

### 3.134.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.78

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{7/2}} dx =$$

$$-\frac{1}{60}a \left( \frac{16 \sqrt{-\frac{1}{a^4c}} \log(x^2 - \frac{1}{a^2})}{c^3} + \frac{3}{(a^6c^3x^4\sqrt{-\frac{1}{c}} - 2a^4c^3x^2\sqrt{-\frac{1}{c}} + a^2c^3\sqrt{-\frac{1}{c}})c} - \frac{8}{(a^4c^2x^2\sqrt{-\frac{1}{c}} - a^2c^2\sqrt{-\frac{1}{c}})} \right)$$

$$+ \frac{1}{15} \left( \frac{8x}{\sqrt{-a^2cx^2 + cc^3}} + \frac{4x}{(-a^2cx^2 + c)^{\frac{3}{2}}c^2} + \frac{3x}{(-a^2cx^2 + c)^{\frac{5}{2}}c} \right) \operatorname{arccosh}(ax)$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^(7/2), x, algorithm="maxima")`

output `-1/60*a*(16*sqrt(-1/(a^4*c))*log(x^2 - 1/a^2)/c^3 + 3/((a^6*c^3*x^4*sqrt(-1/c) - 2*a^4*c^3*x^2*sqrt(-1/c) + a^2*c^3*sqrt(-1/c))*c) - 8/((a^4*c^2*x^2*sqrt(-1/c) - a^2*c^2*sqrt(-1/c))*c^2)) + 1/15*(8*x/(sqrt(-a^2*c*x^2 + c)*c^3) + 4*x/((-a^2*c*x^2 + c)^(3/2)*c^2) + 3*x/((-a^2*c*x^2 + c)^(5/2)*c))*arccosh(a*x)`

**3.134.8 Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.58

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{7/2}} dx = \frac{1}{60} \sqrt{-c} \left( \frac{16 \log(|a^2x^2 - 1|)}{ac^4} - \frac{24a^4x^4 - 56a^2x^2 + 35}{(a^2x^2 - 1)^2ac^4} \right) - \frac{\sqrt{-a^2cx^2 + c} \left( 4 \left( \frac{2a^4x^2}{c} - \frac{5a^2}{c} \right) x^2 + \frac{15}{c} \right) x \log(ax + \sqrt{a^2x^2 - 1})}{15(a^2cx^2 - c)^3}$$

input `integrate(arccosh(a*x)/(-a^2*c*x^2+c)^(7/2),x, algorithm="giac")`output `1/60*sqrt(-c)*(16*log(abs(a^2*x^2 - 1))/(a*c^4) - (24*a^4*x^4 - 56*a^2*x^2 + 35)/((a^2*x^2 - 1)^2*a*c^4)) - 1/15*sqrt(-a^2*c*x^2 + c)*(4*(2*a^4*x^2/c - 5*a^2/c)*x^2 + 15/c)*x*log(a*x + sqrt(a^2*x^2 - 1))/(a^2*c*x^2 - c)^3`**3.134.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{(c - a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{acosh}(ax)}{(c - a^2cx^2)^{7/2}} dx$$

input `int(acosh(a*x)/(c - a^2*c*x^2)^(7/2),x)`output `int(acosh(a*x)/(c - a^2*c*x^2)^(7/2), x)`

### 3.135 $\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$

3.135.1 Optimal result . . . . .	1183
3.135.2 Mathematica [A] (verified) . . . . .	1183
3.135.3 Rubi [A] (verified) . . . . .	1184
3.135.4 Maple [B] (verified) . . . . .	1186
3.135.5 Fracas [F] . . . . .	1186
3.135.6 Sympy [F] . . . . .	1187
3.135.7 Maxima [F(-2)] . . . . .	1187
3.135.8 Giac [F] . . . . .	1187
3.135.9 Mupad [F(-1)] . . . . .	1188

#### 3.135.1 Optimal result

Integrand size = 22, antiderivative size = 145

$$\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = -\frac{3x^2\sqrt{-1+ax}}{16a^3\sqrt{1-ax}} - \frac{x^4\sqrt{-1+ax}}{16a\sqrt{1-ax}} - \frac{3x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{8a^4} - \frac{x^3\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{4a^2} + \frac{3\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{16a^5\sqrt{1-ax}}$$

```
output -3/16*x^2*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-1/16*x^4*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)+3/16*arccosh(a*x)^2*(a*x-1)^(1/2)/a^5/(-a*x+1)^(1/2)-3/8*x*arccosh(a*x)*(-a^2*x^2+1)^(1/2)/a^4-1/4*x^3*arccosh(a*x)*(-a^2*x^2+1)^(1/2)/a^2
```

#### 3.135.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.64

$$\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax)(-16 \cosh(2\operatorname{arccosh}(ax)) - \cosh(4\operatorname{arccosh}(ax)) + 4\operatorname{arccosh}(ax)(6\operatorname{arccosh}(ax) + 8 \sinh(2\operatorname{arccosh}(ax))))}{128a^5\sqrt{-((-1+ax)(1+ax))}}$$

```
input Integrate[(x^4*ArcCosh[a*x])/Sqrt[1 - a^2*x^2], x]
```



output  $(\text{Sqrt}[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(-16*\text{Cosh}[2*\text{ArcCosh}[a*x]] - \text{Cosh}[4*\text{ArcCosh}[a*x]] + 4*\text{ArcCosh}[a*x]*(6*\text{ArcCosh}[a*x] + 8*\text{Sinh}[2*\text{ArcCosh}[a*x]] + \text{Sinh}[4*\text{ArcCosh}[a*x]])))/(128*a^5*\text{Sqrt}[(-1 + a*x)*(1 + a*x)])$

### 3.135.3 Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.06, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$ , Rules used = {6353, 15, 6353, 15, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{\sqrt{ax-1} \int x^3 dx}{4a\sqrt{1-ax}} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{4a^2} \\
 & \quad \downarrow \text{15} \\
 & \frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{4a^2} - \frac{x^4 \sqrt{ax-1}}{16a\sqrt{1-ax}} \\
 & \quad \downarrow \text{6353} \\
 & \frac{3 \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{\sqrt{ax-1} \int x dx}{2a\sqrt{1-ax}} - \frac{x \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{2a^2} \right)}{4a^2} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{4a^2} \\
 & \quad \quad \quad \frac{x^4 \sqrt{ax-1}}{16a\sqrt{1-ax}} \\
 & \quad \quad \quad \downarrow \text{15} \\
 & \frac{3 \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{x \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{2a^2} - \frac{x^2 \sqrt{ax-1}}{4a\sqrt{1-ax}} \right)}{4a^2} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{4a^2} - \frac{x^4 \sqrt{ax-1}}{16a\sqrt{1-ax}} \\
 & \quad \quad \quad \downarrow \text{6307}
 \end{aligned}$$

---

3.135.  $\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$

$$-\frac{x^3\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{4a^2} + \frac{3\left(\frac{\sqrt{ax-1}\operatorname{arccosh}(ax)^2}{4a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2a^2} - \frac{x^2\sqrt{ax-1}}{4a\sqrt{1-ax}}\right)}{4a^2} - \frac{x^4\sqrt{ax-1}}{16a\sqrt{1-ax}}$$

input `Int[(x^4*ArcCosh[a*x])/Sqrt[1 - a^2*x^2], x]`

output `-1/16*(x^4*Sqrt[-1 + a*x])/(a*Sqrt[1 - a*x]) - (x^3*Sqrt[1 - a^2*x^2]*ArcCosh[a*x])/(4*a^2) + (3*(-1/4*(x^2*Sqrt[-1 + a*x])/(a*Sqrt[1 - a*x]) - (x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x])/(2*a^2) + (Sqrt[-1 + a*x]*ArcCosh[a*x]^2)/(4*a^3*Sqrt[1 - a*x]))) / (4*a^2)`

### 3.135.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.135.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 455 vs.  $2(119) = 238$ .

Time = 0.90 (sec) , antiderivative size = 456, normalized size of antiderivative = 3.14

method	result
default	$-\frac{3\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^2}{16a^5(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(8a^5x^5-12a^3x^3+8\sqrt{ax-1}\sqrt{ax+1}a^4x^4+4ax-8a^2x^2\sqrt{ax-1}\sqrt{ax+1})}{256a^5(a^2x^2-1)}$

input `int(x^4*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -3/16*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a^5/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^2 \\ & -1/256*(-a^2*x^2+1)^(1/2)*(8*a^5*x^5-12*a^3*x^3+8*(a*x-1)^(1/2)*(a*x+1)^(1/2)* \\ & a^4*x^4+4*a*x-8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)* \\ & (a*x+1)^(1/2))*(-1+4*\operatorname{arccosh}(a*x))/a^5/(a^2*x^2-1)-1/16*(-a^2*x^2+1)^(1/2)* \\ & (2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)* \\ & (a*x+1)^(1/2))*(-1+2*\operatorname{arccosh}(a*x))/a^5/(a^2*x^2-1)-1/16*(-a^2*x^2+1)^(1/2)* \\ & (2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)^(1/2))* \\ & (1+2*\operatorname{arccosh}(a*x))/a^5/(a^2*x^2-1)-1/256*(-a^2*x^2+1)^(1/2)*(8*a^5*x^5-12*a^3*x^3-8*(a*x-1)^(1/2)* \\ & (a*x+1)^(1/2)*a^4*x^4+4*a*x+8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))* \\ & (1+4*\operatorname{arccosh}(a*x))/a^5/(a^2*x^2-1) \end{aligned}$$

### 3.135.5 Fracas [F]

$$\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^4*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^4*arccosh(a*x)/(a^2*x^2 - 1), x)`

**3.135.6 Sympy [F]**

$$\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{acosh}(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**4*acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**4*acosh(a*x)/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.135.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.135.8 Giac [F]**

$$\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^4*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^4*arccosh(a*x)/sqrt(-a^2*x^2 + 1), x)`

**3.135.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{acosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

input `int((x^4*acosh(a*x))/(1 - a^2*x^2)^(1/2), x)`output `int((x^4*acosh(a*x))/(1 - a^2*x^2)^(1/2), x)`

### 3.136 $\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$

3.136.1 Optimal result . . . . .	1189
3.136.2 Mathematica [A] (verified) . . . . .	1189
3.136.3 Rubi [A] (verified) . . . . .	1190
3.136.4 Maple [B] (verified) . . . . .	1191
3.136.5 Fricas [A] (verification not implemented) . . . . .	1192
3.136.6 Sympy [F] . . . . .	1192
3.136.7 Maxima [C] (verification not implemented) . . . . .	1193
3.136.8 Giac [F(-2)] . . . . .	1193
3.136.9 Mupad [F(-1)] . . . . .	1193

#### 3.136.1 Optimal result

Integrand size = 22, antiderivative size = 110

$$\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = -\frac{2x\sqrt{-1+ax}}{3a^3\sqrt{1-ax}} - \frac{x^3\sqrt{-1+ax}}{9a\sqrt{1-ax}} - \frac{2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{3a^4} - \frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{3a^2}$$

output 
$$-2/3*x*(a*x-1)^{(1/2)}/a^3/(-a*x+1)^{(1/2)}-1/9*x^3*(a*x-1)^{(1/2)}/a/(-a*x+1)^{(1/2)}-2/3*\operatorname{arccosh}(a*x)*(-a^2*x^2+1)^{(1/2)}/a^4-1/3*x^2*\operatorname{arccosh}(a*x)*(-a^2*x^2+1)^{(1/2)}/a^2$$

#### 3.136.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.67

$$\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = -\frac{ax\sqrt{-1+ax}\sqrt{1+ax}(6+a^2x^2)-3(-2+a^2x^2+a^4x^4)\operatorname{arccosh}(ax)}{9a^4\sqrt{1-a^2x^2}}$$

input `Integrate[(x^3*ArcCosh[a*x])/Sqrt[1 - a^2*x^2],x]`

output 
$$-1/9*(a*x*\operatorname{Sqrt}[-1+a*x]*\operatorname{Sqrt}[1+a*x]*(6+a^2*x^2)-3*(-2+a^2*x^2+a^4*x^4)*\operatorname{ArcCosh}[a*x])/(a^4*\operatorname{Sqrt}[1-a^2*x^2])$$

**3.136.3 Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {6353, 15, 6329, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{2 \int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{\sqrt{ax-1} \int x^2 dx}{3a\sqrt{1-ax}} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{3a^2} \\
 & \quad \downarrow \text{15} \\
 & \frac{2 \int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{3a^2} - \frac{x^3 \sqrt{ax-1}}{9a\sqrt{1-ax}} \\
 & \quad \downarrow \text{6329} \\
 & \frac{2 \left( -\frac{\sqrt{ax-1} \int 1 dx}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{a^2} \right)}{3a^2} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{3a^2} - \frac{x^3 \sqrt{ax-1}}{9a\sqrt{1-ax}} \\
 & \quad \downarrow \text{24} \\
 & -\frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{3a^2} + \frac{2 \left( -\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{a^2} - \frac{x \sqrt{ax-1}}{a\sqrt{1-ax}} \right)}{3a^2} - \frac{x^3 \sqrt{ax-1}}{9a\sqrt{1-ax}}
 \end{aligned}$$

input `Int[(x^3*ArcCosh[a*x])/Sqrt[1 - a^2*x^2],x]`

output `-1/9*(x^3*Sqrt[-1 + a*x])/(a*Sqrt[1 - a*x]) - (x^2*Sqrt[1 - a^2*x^2]*ArcCosh[a*x])/(3*a^2) + (2*(-((x*Sqrt[-1 + a*x])/(a*Sqrt[1 - a*x])) - (Sqrt[1 - a^2*x^2]*ArcCosh[a*x])/a^2))/(3*a^2)`

## 3.136.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`
- rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

## 3.136.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 310 vs. 2(90) = 180.

Time = 1.18 (sec) , antiderivative size = 311, normalized size of antiderivative = 2.83

method	result
default	$\frac{\sqrt{-a^2x^2+1} (4a^4x^4-5a^2x^2+4a^3x^3\sqrt{ax-1}\sqrt{ax+1}-3\sqrt{ax-1}\sqrt{ax+1}ax+1)(-1+3\operatorname{arccosh}(ax))}{72a^4(a^2x^2-1)} - \frac{3\sqrt{-a^2x^2+1}(\sqrt{ax-1}\sqrt{ax+1})}{8a^4}$

input `int(x^3*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`



output 
$$\begin{aligned} & -1/72*(-a^2*x^2+1)^{(1/2)}*(4*a^4*x^4-5*a^2*x^2+4*a^3*x^3*(a*x-1)^{(1/2)}*(a*x \\ & +1)^{(1/2)}-3*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a*x+1)*(-1+3*\operatorname{arccosh}(a*x))/a^4/(a^ \\ & 2*x^2-1)-3/8*(-a^2*x^2+1)^{(1/2)}*((a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a*x+a^2*x^2-1 \\ & )*(-1+\operatorname{arccosh}(a*x))/a^4/(a^2*x^2-1)-3/8*(-a^2*x^2+1)^{(1/2)}*(a^2*x^2-(a*x-1 \\ & )^{(1/2)}*(a*x+1)^{(1/2)}*a*x-1)*(1+\operatorname{arccosh}(a*x))/a^4/(a^2*x^2-1)-1/72*(-a^2*x \\ & ^2+1)^{(1/2)}*(4*a^4*x^4-5*a^2*x^2-4*a^3*x^3*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+3*( \\ & a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a*x+1)*(1+3*\operatorname{arccosh}(a*x))/a^4/(a^2*x^2-1) \end{aligned}$$

### 3.136.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.92

$$\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{3(a^4x^4 + a^2x^2 - 2)\sqrt{-a^2x^2 + 1} \log(ax + \sqrt{a^2x^2 - 1}) - (a^3x^3 + 6ax)\sqrt{a^2x^2 - 1}\sqrt{-a^2x^2 + 1}}{9(a^6x^2 - a^4)}$$

input `integrate(x^3*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output 
$$\begin{aligned} & -1/9*(3*(a^4*x^4 + a^2*x^2 - 2)*\operatorname{sqrt}(-a^2*x^2 + 1)*\log(a*x + \operatorname{sqrt}(a^2*x^2 \\ & - 1)) - (a^3*x^3 + 6*a*x)*\operatorname{sqrt}(a^2*x^2 - 1)*\operatorname{sqrt}(-a^2*x^2 + 1))/(a^6*x^2 - \\ & a^4) \end{aligned}$$

### 3.136.6 Sympy [F]

$$\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^3 \operatorname{acosh}(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**3*acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**3*acosh(a*x)/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.136.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.93 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.56

$$\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{1}{9} a \left( \frac{ix^3}{a^2} + \frac{6ix}{a^4} \right) - \frac{1}{3} \left( \frac{\sqrt{-a^2x^2+1}x^2}{a^2} + \frac{2\sqrt{-a^2x^2+1}}{a^4} \right) \operatorname{arccosh}(ax)$$

input `integrate(x^3*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `1/9*a*(I*x^3/a^2 + 6*I*x/a^4) - 1/3*(sqrt(-a^2*x^2 + 1)*x^2/a^2 + 2*sqrt(-a^2*x^2 + 1)/a^4)*arccosh(a*x)`

**3.136.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.136.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^3 \operatorname{acosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

input `int((x^3*acosh(a*x))/(1 - a^2*x^2)^(1/2),x)`

output `int((x^3*acosh(a*x))/(1 - a^2*x^2)^(1/2), x)`

### 3.137 $\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$

3.137.1 Optimal result . . . . .	1194
3.137.2 Mathematica [A] (verified) . . . . .	1194
3.137.3 Rubi [A] (verified) . . . . .	1195
3.137.4 Maple [B] (verified) . . . . .	1196
3.137.5 Fricas [F] . . . . .	1196
3.137.6 Sympy [F] . . . . .	1197
3.137.7 Maxima [F(-2)] . . . . .	1197
3.137.8 Giac [F] . . . . .	1197
3.137.9 Mupad [F(-1)] . . . . .	1198

#### 3.137.1 Optimal result

Integrand size = 22, antiderivative size = 88

$$\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = -\frac{x^2 \sqrt{-1+ax}}{4a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{2a^2} + \frac{\sqrt{-1+ax} \operatorname{arccosh}(ax)^2}{4a^3\sqrt{1-ax}}$$

output  $-1/4*x^2*(a*x-1)^{(1/2)}/a/(-a*x+1)^{(1/2)}+1/4*\operatorname{arccosh}(a*x)^2*(a*x-1)^{(1/2)}/a$   
 $\wedge 3/(-a*x+1)^{(1/2)}-1/2*x*\operatorname{arccosh}(a*x)*(-a^2*x^2+1)^{(1/2)}/a^2$

#### 3.137.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.85

$$\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-((-1+ax)(1+ax))}(-\cosh(2\operatorname{arccosh}(ax)) + 2\operatorname{arccosh}(ax)(\operatorname{arccosh}(ax) + \sinh(2\operatorname{arccosh}(ax))))}{8a^3 \sqrt{\frac{-1+ax}{1+ax}}(1+ax)}$$

input `Integrate[(x^2*ArcCosh[a*x])/Sqrt[1 - a^2*x^2],x]`

output  $-1/8*(\operatorname{Sqrt}[ -((-1+a*x)*(1+a*x))] * (-\operatorname{Cosh}[2*\operatorname{ArcCosh}[a*x]] + 2*\operatorname{ArcCosh}[a*x]$   
 $]*(\operatorname{ArcCosh}[a*x] + \operatorname{Sinh}[2*\operatorname{ArcCosh}[a*x]])))/(a^3*\operatorname{Sqrt}[(-1+a*x)/(1+a*x)]*(1+a*x))$

**3.137.3 Rubi [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$ , Rules used = {6353, 15, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{\sqrt{ax-1} \int x dx}{2a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{2a^2} \\
 & \quad \downarrow \text{15} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{2a^2} - \frac{x^2\sqrt{ax-1}}{4a\sqrt{1-ax}} \\
 & \quad \downarrow \text{6307} \\
 & \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^2}{4a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{2a^2} - \frac{x^2\sqrt{ax-1}}{4a\sqrt{1-ax}}
 \end{aligned}$$

input `Int[(x^2*ArcCosh[a*x])/Sqrt[1 - a^2*x^2], x]`

output `-1/4*(x^2*Sqrt[-1 + a*x])/(a*Sqrt[1 - a*x]) - (x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x])/(2*a^2) + (Sqrt[-1 + a*x]*ArcCosh[a*x]^2)/(4*a^3*Sqrt[1 - a*x])`

**3.137.3.1 Defintions of rubi rules used**

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1))/(m + 1), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

---

3.137.  $\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.137.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 222 vs.  $2(72) = 144$ .

Time = 0.68 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.53

method	result
default	$-\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^2}{4a^3(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(2a^3x^3-2ax+2a^2x^2\sqrt{ax-1}\sqrt{ax+1}-\sqrt{ax-1}\sqrt{ax+1})(-1+2\operatorname{arccosh}(ax))}{16a^3(a^2x^2-1)}$

input `int(x^2*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/4*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a^3/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^2 \\ & -1/16*(-a^2*x^2+1)^(1/2)*(2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2) \\ & - (a*x-1)^(1/2)*(a*x+1)^(1/2))*(-1+2*\operatorname{arccosh}(a*x))/a^3/(a^2*x^2-1) \\ & -1/16*(-a^2*x^2+1)^(1/2)*(2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2) \\ & + (a*x-1)^(1/2)*(a*x+1)^(1/2))*(1+2*\operatorname{arccosh}(a*x))/a^3/(a^2*x^2-1) \end{aligned}$$

### 3.137.5 Fracas [F]

$$\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^2*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^2*arccosh(a*x)/(a^2*x^2 - 1), x)`

**3.137.6 Sympy [F]**

$$\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{acosh}(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**2*acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**2*acosh(a*x)/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.137.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^2*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.137.8 Giac [F]**

$$\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^2*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^2*arccosh(a*x)/sqrt(-a^2*x^2 + 1), x)`

**3.137.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{acosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

input `int((x^2*acosh(a*x))/(1 - a^2*x^2)^(1/2), x)`output `int((x^2*acosh(a*x))/(1 - a^2*x^2)^(1/2), x)`

$$3.138 \quad \int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

3.138.1 Optimal result	1199
3.138.2 Mathematica [A] (verified)	1199
3.138.3 Rubi [A] (verified)	1200
3.138.4 Maple [B] (verified)	1201
3.138.5 Fricas [A] (verification not implemented)	1201
3.138.6 Sympy [F]	1201
3.138.7 Maxima [C] (verification not implemented)	1202
3.138.8 Giac [C] (verification not implemented)	1202
3.138.9 Mupad [F(-1)]	1202

### 3.138.1 Optimal result

Integrand size = 20, antiderivative size = 49

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = -\frac{x\sqrt{-1+ax}}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{a^2}$$

output `-x*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)-arccosh(a*x)*(-a^2*x^2+1)^(1/2)/a^2`

### 3.138.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.12

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{-ax\sqrt{-1+ax}\sqrt{1+ax} + (-1+a^2x^2) \operatorname{arccosh}(ax)}{a^2\sqrt{1-a^2x^2}}$$

input `Integrate[(x*ArcCosh[a*x])/Sqrt[1 - a^2*x^2],x]`

output `(-(a*x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + (-1 + a^2*x^2)*ArcCosh[a*x])/(a^2*Sqrt[1 - a^2*x^2])`

---

3.138.  $\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$



**3.138.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6329, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

$$\downarrow 6329$$

$$-\frac{\sqrt{ax-1} \int 1 dx}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{a^2}$$

$$\downarrow 24$$

$$-\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)}{a^2} - \frac{x\sqrt{ax-1}}{a\sqrt{1-ax}}$$

input `Int[(x*ArcCosh[a*x])/Sqrt[1 - a^2*x^2],x]`

output `-((x*Sqrt[-1 + a*x])/(a*Sqrt[1 - a*x])) - (Sqrt[1 - a^2*x^2]*ArcCosh[a*x])/a^2`

**3.138.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

**3.138.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 122 vs.  $2(43) = 86$ .

Time = 0.85 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.51

method	result
default	$-\frac{\sqrt{-a^2x^2+1}(\sqrt{ax-1}\sqrt{ax+1}ax+a^2x^2-1)(-1+\operatorname{arccosh}(ax))}{2a^2(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(a^2x^2-\sqrt{ax-1}\sqrt{ax+1}ax-1)(1+\operatorname{arccosh}(ax))}{2a^2(a^2x^2-1)}$

input `int(x*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/2*(-a^2*x^2+1)^{(1/2)}*((a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a*x+a^2*x^2-1)*(-1+\operatorname{arccosh}(a*x))/a^2/(a^2*x^2-1)-1/2*(-a^2*x^2+1)^{(1/2)}*(a^2*x^2-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a*x-1)*(1+\operatorname{arccosh}(a*x))/a^2/(a^2*x^2-1)$$

**3.138.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.47

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{a^2x^2-1}\sqrt{-a^2x^2+1}ax + (-a^2x^2+1)^{\frac{3}{2}} \log(ax + \sqrt{a^2x^2-1})}{a^4x^2 - a^2}$$

input `integrate(x*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fracas")`

output 
$$(\operatorname{sqrt}(a^2*x^2 - 1)*\operatorname{sqrt}(-a^2*x^2 + 1)*a*x + (-a^2*x^2 + 1)^{(3/2)}*\log(a*x + \operatorname{sqrt}(a^2*x^2 - 1)))/(a^4*x^2 - a^2)$$

**3.138.6 Sympy [F]**

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x \operatorname{acosh}(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x*acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x*acosh(a*x)/sqrt(-(a*x - 1)*(a*x + 1)), x)`

---

3.138. 
$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

**3.138.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.76 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.57

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{ix}{a} - \frac{\sqrt{-a^2x^2+1} \operatorname{arccosh}(ax)}{a^2}$$

input `integrate(x*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `I*x/a - sqrt(-a^2*x^2 + 1)*arccosh(a*x)/a^2`

**3.138.8 Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = -\frac{ix}{a} - \frac{\sqrt{-a^2x^2+1} \log(ax + \sqrt{a^2x^2-1})}{a^2}$$

input `integrate(x*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `-I*x/a - sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1))/a^2`

**3.138.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{x \operatorname{acosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

input `int((x*acosh(a*x))/(1 - a^2*x^2)^(1/2),x)`

output `int((x*acosh(a*x))/(1 - a^2*x^2)^(1/2), x)`

### 3.139 $\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$

3.139.1 Optimal result . . . . .	1203
3.139.2 Mathematica [A] (verified) . . . . .	1203
3.139.3 Rubi [A] (verified) . . . . .	1204
3.139.4 Maple [A] (verified) . . . . .	1204
3.139.5 Fricas [F] . . . . .	1205
3.139.6 Sympy [F] . . . . .	1205
3.139.7 Maxima [F] . . . . .	1205
3.139.8 Giac [F] . . . . .	1206
3.139.9 Mupad [F(-1)] . . . . .	1206

#### 3.139.1 Optimal result

Integrand size = 19, antiderivative size = 32

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{2a\sqrt{1-ax}}$$

output  $1/2*\operatorname{arccosh}(a*x)^2*(a*x-1)^{(1/2)}/a/(-a*x+1)^{(1/2)}$

#### 3.139.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.41

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{2a\sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]/Sqrt[1 - a^2*x^2], x]`

output  $(\operatorname{Sqrt}[-1 + a*x]*\operatorname{Sqrt}[1 + a*x]*\operatorname{ArcCosh}[a*x]^2)/(2*a*\operatorname{Sqrt}[1 - a^2*x^2])$

### 3.139.3 Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

↓ 6307

$$\frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^2}{2a\sqrt{1-ax}}$$

input `Int[ArcCosh[a*x]/Sqrt[1 - a^2*x^2], x]`

output `(Sqrt[-1 + a*x]*ArcCosh[a*x]^2)/(2*a*Sqrt[1 - a*x])`

#### 3.139.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.139.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.59

method	result	size
default	$-\frac{\sqrt{-(ax-1)(ax+1)}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^2}{2(a^2x^2-1)a}$	51

input `int(arccosh(a*x)/(-a^2*x^2+1)^(1/2), x, method=_RETURNVERBOSE)`

output `-1/2*(-(a*x-1)*(a*x+1))^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2-1)/a*arccosh(a*x)^2`

**3.139.5 Fricas [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)/(a^2*x^2 - 1), x)`

**3.139.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.139.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)/sqrt(-a^2*x^2 + 1), x)`

**3.139.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)/sqrt(-a^2*x^2 + 1), x)`

**3.139.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)/(1 - a^2*x^2)^(1/2),x)`

output `int(acosh(a*x)/(1 - a^2*x^2)^(1/2), x)`

### 3.140 $\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx$

3.140.1 Optimal result . . . . .	1207
3.140.2 Mathematica [A] (verified) . . . . .	1207
3.140.3 Rubi [A] (verified) . . . . .	1208
3.140.4 Maple [B] (verified) . . . . .	1210
3.140.5 Fricas [F] . . . . .	1210
3.140.6 Sympy [F] . . . . .	1210
3.140.7 Maxima [F] . . . . .	1211
3.140.8 Giac [F] . . . . .	1211
3.140.9 Mupad [F(-1)] . . . . .	1211

#### 3.140.1 Optimal result

Integrand size = 22, antiderivative size = 103

$$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx = \frac{2\sqrt{-1+ax}\operatorname{arccosh}(ax)\arctan(e^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{i\sqrt{-1+ax}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} + \frac{i\sqrt{-1+ax}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}}$$

```
output 2*arccosh(a*x)*arctan(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-I*polylog(2,-I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)+I*polylog(2,I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)
```

#### 3.140.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx = \frac{i\sqrt{-((-1+ax)(1+ax))}(\operatorname{arccosh}(ax)(\log(1-ie^{-\operatorname{arccosh}(ax)})-\log(1+ie^{-\operatorname{arccosh}(ax)})))+\operatorname{PolyLog}(2,-\sqrt{\frac{-1+ax}{1+ax}}(1+ax))}{\sqrt{\frac{-1+ax}{1+ax}}(1+ax)}$$



input `Integrate[ArcCosh[a*x]/(x*Sqrt[1 - a^2*x^2]),x]`

output `(I*Sqrt[-((-1 + a*x)*(1 + a*x))]*(ArcCosh[a*x]*(Log[1 - I/E^ArcCosh[a*x]] - Log[1 + I/E^ArcCosh[a*x]]) + PolyLog[2, (-I)/E^ArcCosh[a*x]] - PolyLog[2, I/E^ArcCosh[a*x]])/(Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x))`

### 3.140.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.64, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$ , Rules used = {6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6361} \\
 & \frac{\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)}{ax} d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{ax-1} \int \operatorname{arccosh}(ax) \csc\left(\operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{4668} \\
 & \frac{\sqrt{ax-1} \left(-i \int \log(1 - ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + i \int \log(1 + ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2\operatorname{arccosh}(ax) \arctan\right)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{2715} \\
 & \frac{\sqrt{ax-1} \left(-i \int e^{-\operatorname{arccosh}(ax)} \log(1 - ie^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} + i \int e^{-\operatorname{arccosh}(ax)} \log(1 + ie^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} +\right)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{2838} \\
 & \frac{\sqrt{ax-1} \left(2\operatorname{arccosh}(ax) \arctan(e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)})\right)}{\sqrt{1-ax}}
 \end{aligned}$$

---

3.140.  $\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx$

input `Int[ArcCosh[a*x]/(x*Sqrt[1 - a^2*x^2]),x]`

output `(Sqrt[-1 + a*x]*(2*ArcCosh[a*x]*ArcTan[E^ArcCosh[a*x]] - I*PolyLog[2, (-I)*E^ArcCosh[a*x]] + I*PolyLog[2, I*E^ArcCosh[a*x]]))/Sqrt[1 - a*x]`

### 3.140.3.1 Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol] :> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6361 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*(x_)^(m_)]/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

### 3.140.4 Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 269 vs.  $2(126) = 252$ .

Time = 0.87 (sec) , antiderivative size = 270, normalized size of antiderivative = 2.62

method	result
default	$\frac{i\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)\ln(1+i(ax+\sqrt{ax-1}\sqrt{ax+1}))}{a^2x^2-1} - \frac{i\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)\ln(1-i(ax+\sqrt{ax-1}\sqrt{ax+1}))}{a^2x^2-1}$

```
input int(arccosh(a*x)/x/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2-1)*arccosh(a*x)*
ln(1+I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))-I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)
*(a*x+1)^(1/2)/(a^2*x^2-1)*arccosh(a*x)*ln(1-I*(a*x+(a*x-1)^(1/2)*(a*x+1)
^(1/2)))+I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2-1)*dil
og(1+I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))-I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)
*(a*x+1)^(1/2)/(a^2*x^2-1)*dilog(1-I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))
```

### 3.140.5 Fricas [F]

$$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}x} dx$$

```
input integrate(arccosh(a*x)/x/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)/(a^2*x^3 - x), x)
```

### 3.140.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{x\sqrt{-(ax-1)(ax+1)}} dx$$

```
input integrate(acosh(a*x)/x/(-a**2*x**2+1)**(1/2),x)
```

```
output Integral(acosh(a*x)/(x*sqrt(-(a*x - 1)*(a*x + 1))), x)
```

---

3.140.  $\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx$

**3.140.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}x} dx$$

input `integrate(arccosh(a*x)/x/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)/(sqrt(-a^2*x^2 + 1)*x), x)`

**3.140.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}x} dx$$

input `integrate(arccosh(a*x)/x/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)/(sqrt(-a^2*x^2 + 1)*x), x)`

**3.140.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{x\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)/(x*(1 - a^2*x^2)^(1/2)),x)`

output `int(acosh(a*x)/(x*(1 - a^2*x^2)^(1/2)), x)`

### 3.141 $\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx$

3.141.1 Optimal result . . . . .	1212
3.141.2 Mathematica [A] (verified) . . . . .	1212
3.141.3 Rubi [A] (verified) . . . . .	1213
3.141.4 Maple [B] (verified) . . . . .	1214
3.141.5 Fricas [B] (verification not implemented) . . . . .	1214
3.141.6 Sympy [F] . . . . .	1215
3.141.7 Maxima [C] (verification not implemented) . . . . .	1215
3.141.8 Giac [C] (verification not implemented) . . . . .	1215
3.141.9 Mupad [F(-1)] . . . . .	1216

#### 3.141.1 Optimal result

Integrand size = 22, antiderivative size = 48

$$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx = -\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{x} - \frac{a\sqrt{-1+ax}\log(x)}{\sqrt{1-ax}}$$

output `-a*ln(x)*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-arccosh(a*x)*(-a^2*x^2+1)^(1/2)/x`

#### 3.141.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.19

$$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx = \frac{(-1+a^2x^2)\operatorname{arccosh}(ax) - ax\sqrt{-1+ax}\sqrt{1+ax}\log(x)}{x\sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]/(x^2*Sqrt[1 - a^2*x^2]),x]`

output `((-1 + a^2*x^2)*ArcCosh[a*x] - a*x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*Log[x])/(x*Sqrt[1 - a^2*x^2])`

**3.141.3 Rubi [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {6332, 14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx$$

↓ 6332

$$-\frac{a\sqrt{ax-1} \int \frac{1}{x} dx}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{x}$$

↓ 14

$$-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{x} - \frac{a\sqrt{ax-1} \log(x)}{\sqrt{1-ax}}$$

input `Int[ArcCosh[a*x]/(x^2*Sqrt[1 - a^2*x^2]),x]`

output `-((Sqrt[1 - a^2*x^2]*ArcCosh[a*x])/x) - (a*Sqrt[-1 + a*x]*Log[x])/Sqrt[1 - a*x]`

**3.141.3.1 Defintions of rubi rules used**

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^{(n_.)*((f_.)*(x_)^{(m_)}*((d_) + (e_.)*(x_)^2)^{(p_.)}, x_Symbol] := Simp[(f*x)^{(m+1)}*(d + e*x^2)^{(p+1)}*((a + b*ArcCosh[c*x])^n/(d*f*(m+1))), x] + Simp[b*c*(n/(f*(m+1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^{(m+1)}*(1 + c*x)^{(p+1/2)}*(-1 + c*x)^{(p+1/2)}*(a + b*ArcCosh[c*x])^{(n-1)}, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

**3.141.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 167 vs.  $2(42) = 84$ .

Time = 0.92 (sec) , antiderivative size = 168, normalized size of antiderivative = 3.50

method	result
default	$-\frac{2\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)a}{a^2x^2-1} - \frac{\sqrt{-a^2x^2+1}(a^2x^2-\sqrt{ax-1}\sqrt{ax+1}ax-1)\operatorname{arccosh}(ax)}{(a^2x^2-1)x} + \frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}}{(a^2x^2-1)x}$

input `int(arccosh(a*x)/x^2/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -2*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2-1)*\operatorname{arccosh}(a*x) \\ & *a-(-a^2*x^2+1)^(1/2)*(a^2*x^2-(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x-1)*\operatorname{arccosh}( \\ & a*x)/(a^2*x^2-1)/x+(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2 \\ & -1)*\ln(1+(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))^2)*a \end{aligned}$$

**3.141.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 89 vs.  $2(42) = 84$ .

Time = 0.28 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.85

$$\begin{aligned} & \int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx \\ & = \frac{ax \arctan\left(\frac{\sqrt{a^2x^2-1}\sqrt{-a^2x^2+1}(x^2+1)}{a^2x^4-(a^2+1)x^2+1}\right) - \sqrt{-a^2x^2+1} \log(ax + \sqrt{a^2x^2-1})}{x} \end{aligned}$$

input `integrate(arccosh(a*x)/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output 
$$(a*x*\arctan(\sqrt{a^2*x^2-1}*\sqrt{-a^2*x^2+1}*(x^2+1)/(a^2*x^4-(a^2+1)*x^2+1)) - \sqrt{-a^2*x^2+1}*\log(a*x + \sqrt{a^2*x^2-1}))/x$$

**3.141.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{x^2\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)/x**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)/(x**2*sqrt(-(a*x - 1)*(a*x + 1))), x)`

**3.141.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.52

$$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx = -\frac{1}{2} \left( a^2 \sqrt{-\frac{1}{a^4}} \log \left( x^2 - \frac{1}{a^2} \right) + i (-1)^{-2a^2x^2+2} \log \left( -2a^2 + \frac{2}{x^2} \right) \right) a - \frac{\sqrt{-a^2x^2+1} \operatorname{arccosh}(ax)}{x}$$

input `integrate(arccosh(a*x)/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `-1/2*(a^2*sqrt(-1/a^4)*log(x^2 - 1/a^2) + I*(-1)^(-2*a^2*x^2 + 2)*log(-2*a^2 + 2/x^2))*a - sqrt(-a^2*x^2 + 1)*arccosh(a*x)/x`

**3.141.8 Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.67

$$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx = \frac{1}{2} \left( \frac{a^4x}{(\sqrt{-a^2x^2+1}|a|+a)|a|} - \frac{\sqrt{-a^2x^2+1}|a|+a}{x|a|} \right) \log \left( ax + \sqrt{a^2x^2-1} \right) - ia \log(|x|)$$



input `integrate(arccosh(a*x)/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `1/2*(a^4*x/((sqrt(-a^2*x^2 + 1)*abs(a) + a)*abs(a)) - (sqrt(-a^2*x^2 + 1)*abs(a) + a)/(x*abs(a)))*log(a*x + sqrt(a^2*x^2 - 1)) - I*a*log(abs(x))`

### 3.141.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{x^2\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)/(x^2*(1 - a^2*x^2)^(1/2)),x)`

output `int(acosh(a*x)/(x^2*(1 - a^2*x^2)^(1/2)), x)`

### 3.142 $\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx$

3.142.1 Optimal result . . . . .	1217
3.142.2 Mathematica [A] (warning: unable to verify) . . . . .	1218
3.142.3 Rubi [A] (verified) . . . . .	1218
3.142.4 Maple [A] (verified) . . . . .	1221
3.142.5 Fricas [F] . . . . .	1221
3.142.6 Sympy [F] . . . . .	1221
3.142.7 Maxima [F] . . . . .	1222
3.142.8 Giac [F] . . . . .	1222
3.142.9 Mupad [F(-1)] . . . . .	1222

#### 3.142.1 Optimal result

Integrand size = 22, antiderivative size = 167

$$\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx = \frac{a\sqrt{-1+ax}}{2x\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2x^2} + \frac{a^2\sqrt{-1+ax}\operatorname{arccosh}(ax)\arctan(e^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{ia^2\sqrt{-1+ax}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(ax)})}{2\sqrt{1-ax}} + \frac{ia^2\sqrt{-1+ax}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(ax)})}{2\sqrt{1-ax}}$$

output

```
1/2*a*(a*x-1)^(1/2)/x/(-a*x+1)^(1/2)+a^2*arccosh(a*x)*arctan(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-1/2*I*a^2*polylog(2,-I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)+1/2*I*a^2*polylog(2,I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-1/2*arccosh(a*x)*(-a^2*x^2+1)^(1/2)/x^2
```

**3.142.2 Mathematica [A] (warning: unable to verify)**

Time = 0.28 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.40

$$\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx$$

$$= \frac{(1+ax) \left( ax\sqrt{\frac{-1+ax}{1+ax}} - \operatorname{arccosh}(ax) + ax\operatorname{arccosh}(ax) - ia^2x^2\sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax) \log(1 - ie^{-\operatorname{arccosh}(ax)}) \right)}{2x^2\sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]/(x^3*Sqrt[1 - a^2*x^2]),x]`output `((1 + a*x)*(a*x*Sqrt[(-1 + a*x)/(1 + a*x)] - ArcCosh[a*x] + a*x*ArcCosh[a*x] - I*a^2*x^2*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]*Log[1 - I/E^ArcCosh[a*x]] + I*a^2*x^2*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]*Log[1 + I/E^ArcCosh[a*x]] - I*a^2*x^2*Sqrt[(-1 + a*x)/(1 + a*x)]*PolyLog[2, (-I)/E^ArcCosh[a*x]] + I*a^2*x^2*Sqrt[(-1 + a*x)/(1 + a*x)]*PolyLog[2, I/E^ArcCosh[a*x]])/(2*x^2*Sqrt[1 - a^2*x^2])`**3.142.3 Rubi [A] (verified)**Time = 0.58 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.75, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.318$ , Rules used = {6347, 15, 6361, 3042, 4668, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx$$

$$\downarrow \text{6347}$$

$$\frac{1}{2}a^2 \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx - \frac{a\sqrt{ax-1}}{2\sqrt{1-ax}} \int \frac{1}{x^2} dx - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2x^2}$$

$$\downarrow \text{15}$$

$$\frac{1}{2}a^2 \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{1-a^2x^2}} dx - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2x^2} + \frac{a\sqrt{ax-1}}{2x\sqrt{1-ax}}$$

$$\downarrow \text{6361}$$

$$\frac{a^2\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)}{ax} d\operatorname{arccosh}(ax)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2x^2} + \frac{a\sqrt{ax-1}}{2x\sqrt{1-ax}}$$

↓ 3042

$$\frac{a^2\sqrt{ax-1} \int \operatorname{arccosh}(ax) \csc\left(i\operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d\operatorname{arccosh}(ax)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2x^2} + \frac{a\sqrt{ax-1}}{2x\sqrt{1-ax}}$$

↓ 4668

$$\frac{a^2\sqrt{ax-1}(-i \int \log(1 - ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + i \int \log(1 + ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2\operatorname{arccosh}(ax) \arctan\left(\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2\sqrt{1-ax}}\right) + \frac{a\sqrt{ax-1}}{2x\sqrt{1-ax}})}{2\sqrt{1-ax}}$$

↓ 2715

$$\frac{a^2\sqrt{ax-1}(-i \int e^{-\operatorname{arccosh}(ax)} \log(1 - ie^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} + i \int e^{-\operatorname{arccosh}(ax)} \log(1 + ie^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} + 2\operatorname{arccosh}(ax) \arctan\left(\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2\sqrt{1-ax}}\right) + \frac{a\sqrt{ax-1}}{2x\sqrt{1-ax}})}{2\sqrt{1-ax}}$$

↓ 2838

$$\frac{a^2\sqrt{ax-1}(2\operatorname{arccosh}(ax) \arctan(e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)}))}{2\sqrt{1-ax}} + \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}{2x^2} + \frac{a\sqrt{ax-1}}{2x\sqrt{1-ax}}$$

input `Int[ArcCosh[a*x]/(x^3*sqrt[1 - a^2*x^2]),x]`

output `(a*sqrt[-1 + a*x])/(2*x*sqrt[1 - a*x]) - (sqrt[1 - a^2*x^2]*ArcCosh[a*x])/(2*x^2) + (a^2*sqrt[-1 + a*x]*(2*ArcCosh[a*x]*ArcTan[E^ArcCosh[a*x]] - I*PolyLog[2, (-I)*E^ArcCosh[a*x]] + I*PolyLog[2, I*E^ArcCosh[a*x]]))/(2*sqrt[1 - a*x])`

## 3.142.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`
- rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`
- rule 6361 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

**3.142.4 Maple [A] (verified)**

Time = 0.99 (sec) , antiderivative size = 349, normalized size of antiderivative = 2.09

method	result
default	$-\frac{(a^2x^2 \operatorname{arccosh}(ax) + \sqrt{ax-1} \sqrt{ax+1} ax - \operatorname{arccosh}(ax)) \sqrt{-a^2x^2+1}}{2(a^2x^2-1)x^2} + \frac{i \sqrt{-a^2x^2+1} \sqrt{ax-1} \sqrt{ax+1} \operatorname{arccosh}(ax) \ln(1+i(ax+\sqrt{ax-1}\sqrt{-a^2x^2+1}))}{2a^2x^2-2}$

```
input int(arccosh(a*x)/x^3/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/2*(a^2*x^2*arccosh(a*x)+(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x-arccosh(a*x))*(-a^2*x^2+1)^(1/2)/(a^2*x^2-1)/x^2+I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*arccosh(a*x)*ln(1+I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*a^2/(2*a^2*x^2-2)-I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*arccosh(a*x)*ln(1-I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*a^2/(2*a^2*x^2-2)+I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*dilog(1+I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*a^2/(2*a^2*x^2-2)-I*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*dilog(1-I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*a^2/(2*a^2*x^2-2)
```

**3.142.5 Fricas [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{x^3 \sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1} x^3} dx$$

```
input integrate(arccosh(a*x)/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)/(a^2*x^5 - x^3), x)
```

**3.142.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{x^3 \sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{x^3 \sqrt{-(ax-1)(ax+1)}} dx$$

```
input integrate(acosh(a*x)/x**3/(-a**2*x**2+1)**(1/2),x)
```

```
output Integral(acosh(a*x)/(x**3*sqrt(-(a*x - 1)*(a*x + 1))), x)
```

---

3.142.  $\int \frac{\operatorname{arccosh}(ax)}{x^3 \sqrt{1-a^2x^2}} dx$

**3.142.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}x^3} dx$$

input `integrate(arccosh(a*x)/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)/(sqrt(-a^2*x^2 + 1)*x^3), x)`

**3.142.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}x^3} dx$$

input `integrate(arccosh(a*x)/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)/(sqrt(-a^2*x^2 + 1)*x^3), x)`

**3.142.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)}{x^3\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)/(x^3*(1 - a^2*x^2)^(1/2)),x)`

output `int(acosh(a*x)/(x^3*(1 - a^2*x^2)^(1/2)), x)`

### 3.143 $\int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{1-c^2x^2}} dx$

3.143.1 Optimal result	1223
3.143.2 Mathematica [A] (verified)	1223
3.143.3 Rubi [A] (verified)	1224
3.143.4 Maple [F]	1225
3.143.5 Fracas [F]	1225
3.143.6 Sympy [F(-1)]	1225
3.143.7 Maxima [F]	1226
3.143.8 Giac [F]	1226
3.143.9 Mupad [F(-1)]	1226

#### 3.143.1 Optimal result

Integrand size = 30, antiderivative size = 98

$$\int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{1-c^2x^2}} dx = \frac{2(fx)^{5/2}(a+b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{9}{4}, c^2x^2\right)}{5f} + \frac{4bc(fx)^{7/2}\sqrt{-1+cx} {}_3F_2\left(1, \frac{7}{4}, \frac{7}{4}, \frac{9}{4}, \frac{11}{4}; c^2x^2\right)}{35f^2\sqrt{1-cx}}$$

output `2/5*(f*x)^(5/2)*(a+b*arccosh(c*x))*hypergeom([1/2, 5/4], [9/4], c^2*x^2)/f+4/35*b*c*(f*x)^(7/2)*hypergeom([1, 7/4, 7/4], [9/4, 11/4], c^2*x^2)*(c*x-1)^(1/2)/f^2/(-c*x+1)^(1/2)`

#### 3.143.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.02

$$\int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{1-c^2x^2}} dx = \frac{2}{35}x(fx)^{3/2} \left( 7(a+b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{9}{4}, c^2x^2\right) + \dots \right)$$

input `Integrate[((f*x)^(3/2)*(a + b*ArcCosh[c*x]))/Sqrt[1 - c^2*x^2], x]`

output `(2*x*(f*x)^(3/2)*(7*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, 5/4, 9/4, c^2*x^2] + (2*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 7/4, 7/4}, {9/4, 11/4}, c^2*x^2])/Sqrt[1 - c^2*x^2]))/35`

---

3.143.  $\int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{1-c^2x^2}} dx$



### 3.143.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.033$ , Rules used = {6363}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^{3/2}(a + \text{barccosh}(cx))}{\sqrt{1 - c^2x^2}} dx$$

↓ 6363

$$\frac{4bc\sqrt{cx - 1}(fx)^{7/2} {}_3F_2\left(1, \frac{7}{4}, \frac{7}{4}; \frac{9}{4}, \frac{11}{4}; c^2x^2\right)}{35f^2\sqrt{1 - cx}} + \frac{2(fx)^{5/2} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{9}{4}, c^2x^2\right) (a + \text{barccosh}(cx))}{5f}$$

input `Int[((f*x)^(3/2)*(a + b*ArcCosh[c*x]))/Sqrt[1 - c^2*x^2],x]`

output `(2*(f*x)^(5/2)*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, 5/4, 9/4, c^2*x^2])/(5*f) + (4*b*c*(f*x)^(7/2)*Sqrt[-1 + c*x]*HypergeometricPFQ[{1, 7/4, 7/4}, {9/4, 11/4}, c^2*x^2])/(35*f^2*Sqrt[1 - c*x])`

#### 3.143.3.1 Defintions of rubi rules used

rule 6363 `Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.))/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && !IntegerQ[m]`

**3.143.4 Maple [F]**

$$\int \frac{(fx)^{\frac{3}{2}}(a + b \operatorname{arccosh}(cx))}{\sqrt{-c^2x^2 + 1}} dx$$

input `int((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

output `int((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

**3.143.5 Fricas [F]**

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{1 - c^2x^2}} dx = \int \frac{(fx)^{\frac{3}{2}}(b \operatorname{arccosh}(cx) + a)}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*(b*f*x*arccosh(c*x) + a*f*x)*sqrt(f*x)/(c^2*x^2 - 1), x)`

**3.143.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{1 - c^2x^2}} dx = \text{Timed out}$$

input `integrate((f*x)**(3/2)*(a+b*acosh(c*x))/(-c**2*x**2+1)**(1/2),x)`

output `Timed out`

**3.143.7 Maxima [F]**

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{1 - c^2x^2}} dx = \int \frac{(fx)^{3/2}(b \operatorname{arcosh}(cx) + a)}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((f*x)^(3/2)*(b*arccosh(c*x) + a)/sqrt(-c^2*x^2 + 1), x)`

**3.143.8 Giac [F]**

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{1 - c^2x^2}} dx = \int \frac{(fx)^{3/2}(b \operatorname{arcosh}(cx) + a)}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((f*x)^(3/2)*(b*arccosh(c*x) + a)/sqrt(-c^2*x^2 + 1), x)`

**3.143.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{1 - c^2x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))(fx)^{3/2}}{\sqrt{1 - c^2x^2}} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^(3/2))/(1 - c^2*x^2)^(1/2),x)`

output `int(((a + b*acosh(c*x))*(f*x)^(3/2))/(1 - c^2*x^2)^(1/2), x)`

$$3.144 \quad \int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$$

3.144.1 Optimal result	1227
3.144.2 Mathematica [A] (verified)	1227
3.144.3 Rubi [A] (verified)	1228
3.144.4 Maple [F]	1229
3.144.5 Fracas [F]	1229
3.144.6 Sympy [F(-1)]	1229
3.144.7 Maxima [F]	1230
3.144.8 Giac [F]	1230
3.144.9 Mupad [F(-1)]	1230

### 3.144.1 Optimal result

Integrand size = 31, antiderivative size = 141

$$\int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx = \frac{2(fx)^{5/2}\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{9}{4}, c^2x^2\right)}{5f\sqrt{d-c^2dx^2}} + \frac{4bc(fx)^{7/2}\sqrt{-1+cx}\sqrt{1+cx} {}_3F_2\left(1, \frac{7}{4}, \frac{7}{4}; \frac{9}{4}, \frac{11}{4}; c^2x^2\right)}{35f^2\sqrt{d-c^2dx^2}}$$

output `4/35*b*c*(f*x)^(7/2)*hypergeom([1, 7/4, 7/4], [9/4, 11/4], c^2*x^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/f^2/(-c^2*d*x^2+d)^(1/2)+2/5*(f*x)^(5/2)*(a+b*arccosh(c*x))*hypergeom([1/2, 5/4], [9/4], c^2*x^2)*(-c^2*x^2+1)^(1/2)/f/(-c^2*d*x^2+d)^(1/2)`

### 3.144.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.82

$$\int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx = \frac{2x(fx)^{3/2} (7\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{9}{4}, c^2x^2\right) + 2b*c*x*\sqrt{-1+cx}*\sqrt{1+cx}*\operatorname{HypergeometricPFQ}\{1, 7/4, 7/4\}, \{9/4, 11/4\}, c^2*x^2))}{35\sqrt{d-c^2dx^2}}$$

input `Integrate[((f*x)^(3/2)*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2], x]`

output `(2*x*(f*x)^(3/2)*(7*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, 5/4, 9/4, c^2*x^2] + 2*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 7/4, 7/4}, {9/4, 11/4}, c^2*x^2]))/(35*Sqrt[d - c^2*d*x^2])`

---


$$3.144. \quad \int \frac{(fx)^{3/2}(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$$

### 3.144.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.032$ , Rules used = {6363}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^{3/2}(a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6363

$$\frac{4bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{7/2} {}_3F_2\left(1, \frac{7}{4}, \frac{7}{4}; \frac{9}{4}, \frac{11}{4}; c^2 x^2\right)}{35f^2\sqrt{d-c^2 dx^2}} + \frac{2\sqrt{1-c^2 x^2}(fx)^{5/2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{4}, \frac{9}{4}, c^2 x^2\right)(a + \operatorname{barccosh}(cx))}{5f\sqrt{d-c^2 dx^2}}$$

input `Int[((f*x)^(3/2)*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2],x]`

output `(2*(f*x)^(5/2)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, 5/4, 9/4, c^2*x^2])/(5*f*Sqrt[d - c^2*d*x^2]) + (4*b*c*(f*x)^(7/2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 7/4, 7/4}, {9/4, 11/4}, c^2*x^2])/(35*f^2*Sqrt[d - c^2*d*x^2])`

#### 3.144.3.1 Defintions of rubi rules used

rule 6363 `Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && !IntegerQ[m]`

**3.144.4 Maple [F]**

$$\int \frac{(fx)^{\frac{3}{2}}(a + b \operatorname{arccosh}(cx))}{\sqrt{-c^2dx^2 + d}} dx$$

input `int((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x)`

output `int((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x)`

**3.144.5 Fricas [F]**

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = \int \frac{(fx)^{\frac{3}{2}}(b \operatorname{arccosh}(cx) + a)}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*f*x*arccosh(c*x) + a*f*x)*sqrt(f*x)/(c^2*d*x^2 - d), x)`

**3.144.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx = \text{Timed out}$$

input `integrate((f*x)**(3/2)*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2),x)`

output `Timed out`

**3.144.7 Maxima [F]**

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^{\frac{3}{2}}(b \operatorname{arccosh}(cx) + a)}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((f*x)^(3/2)*(b*arccosh(c*x) + a)/sqrt(-c^2*d*x^2 + d), x)`

**3.144.8 Giac [F]**

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^{\frac{3}{2}}(b \operatorname{arccosh}(cx) + a)}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^(3/2)*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((f*x)^(3/2)*(b*arccosh(c*x) + a)/sqrt(-c^2*d*x^2 + d), x)`

**3.144.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^{3/2}(a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^{3/2}}{\sqrt{d - c^2 dx^2}} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^(3/2))/(d - c^2*d*x^2)^(1/2),x)`

output `int(((a + b*acosh(c*x))*(f*x)^(3/2))/(d - c^2*d*x^2)^(1/2), x)`

### 3.145 $\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.145.1 Optimal result . . . . .	1231
3.145.2 Mathematica [A] (verified) . . . . .	1232
3.145.3 Rubi [A] (verified) . . . . .	1232
3.145.4 Maple [F] . . . . .	1237
3.145.5 Fricas [F] . . . . .	1237
3.145.6 Sympy [F(-1)] . . . . .	1238
3.145.7 Maxima [F] . . . . .	1238
3.145.8 Giac [F(-2)] . . . . .	1239
3.145.9 Mupad [F(-1)] . . . . .	1239

#### 3.145.1 Optimal result

Integrand size = 27, antiderivative size = 429

$$\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{bcd^3(2271 + 1329m + 284m^2 + 27m^3 + m^4)(fx)^{2+m}(1 - c^2x^2)}{f^2(3+m)^2(5+m)^2(7+m)^2\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{bc^3d^3(9+m)(13+2m)(fx)^{4+m}(1 - c^2x^2)}{f^4(5+m)^2(7+m)^2\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^5d^3(fx)^{6+m}(1 - c^2x^2)}{f^6(7+m)^2\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{d^3(fx)^{1+m}(a + \operatorname{barccosh}(cx))}{f(1+m)} - \frac{3c^2d^3(fx)^{3+m}(a + \operatorname{barccosh}(cx))}{f^3(3+m)}$$

$$+ \frac{3c^4d^3(fx)^{5+m}(a + \operatorname{barccosh}(cx))}{f^5(5+m)} - \frac{c^6d^3(fx)^{7+m}(a + \operatorname{barccosh}(cx))}{f^7(7+m)}$$

$$- \frac{3bcd^3(2161 + 1813m + 455m^2 + 35m^3)(fx)^{2+m}\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{f^2(1+m)(2+m)(3+m)^2(5+m)^2(7+m)^2\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
d^3*(f*x)^(1+m)*(a+b*arccosh(c*x))/f/(1+m)-3*c^2*d^3*(f*x)^(3+m)*(a+b*arccosh(c*x))/f^3/(3+m)+3*c^4*d^3*(f*x)^(5+m)*(a+b*arccosh(c*x))/f^5/(5+m)-c^6*d^3*(f*x)^(7+m)*(a+b*arccosh(c*x))/f^7/(7+m)-b*c*d^3*(m^4+27*m^3+284*m^2+1329*m+2271)*(f*x)^(2+m)*(-c^2*x^2+1)/f^2/(7+m)^2/(m^2+8*m+15)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*c^3*d^3*(9+m)*(13+2*m)*(f*x)^(4+m)*(-c^2*x^2+1)/f^4/(5+m)^2/(7+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*c^5*d^3*(f*x)^(6+m)*(-c^2*x^2+1)/f^6/(7+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-3*b*c*d^3*(35*m^3+455*m^2+1813*m+2161)*(f*x)^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/f^2/(7+m)^2/(m^2+3*m+2)/(m^2+8*m+15)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```



### 3.145.2 Mathematica [A] (verified)

Time = 0.79 (sec) , antiderivative size = 387, normalized size of antiderivative = 0.90

$$\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= d^3 x (fx)^m \left( \frac{a + \operatorname{barccosh}(cx)}{1 + m} - \frac{3c^2 x^2 (a + \operatorname{barccosh}(cx))}{3 + m} + \frac{3c^4 x^4 (a + \operatorname{barccosh}(cx))}{5 + m} - \frac{c^6 x^6 (a + \operatorname{barccosh}(cx))}{7 + m} + \frac{bc^7 x^7 \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 4 + \frac{m}{2}, 5 + \frac{m}{2}, c^2 x^2\right)}{(7 + m)(8 + m)\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcx\sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2 x^2\right)}{(2 + 3m + m^2)\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{3bc^3 x^3 \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4+m}{2}, \frac{6+m}{2}, c^2 x^2\right)}{(12 + 7m + m^2)\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{3bc^5 x^5 \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{6+m}{2}, \frac{8+m}{2}, c^2 x^2\right)}{(5 + m)(6 + m)\sqrt{-1 + cx}\sqrt{1 + cx}} \right)$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `d^3*x*(f*x)^m*((a + b*ArcCosh[c*x])/(1 + m) - (3*c^2*x^2*(a + b*ArcCosh[c*x]))/(3 + m) + (3*c^4*x^4*(a + b*ArcCosh[c*x]))/(5 + m) - (c^6*x^6*(a + b*ArcCosh[c*x]))/(7 + m) + (b*c^7*x^7*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, 4 + m/2, 5 + m/2, c^2*x^2])/((7 + m)*(8 + m)*sqrt[-1 + c*x]*sqrt[1 + c*x]) - (b*c*x*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, c^2*x^2])/((2 + 3*m + m^2)*sqrt[-1 + c*x]*sqrt[1 + c*x]) + (3*b*c^3*x^3*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m)/2, c^2*x^2])/((12 + 7*m + m^2)*sqrt[-1 + c*x]*sqrt[1 + c*x]) - (3*b*c^5*x^5*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (6 + m)/2, (8 + m)/2, c^2*x^2])/((5 + m)*(6 + m)*sqrt[-1 + c*x]*sqrt[1 + c*x]))`

### 3.145.3 Rubi [A] (verified)

Time = 2.39 (sec) , antiderivative size = 423, normalized size of antiderivative = 0.99, number of steps used = 9, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6336, 27, 2113, 2340, 1590, 27, 363, 279, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.145.  $\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
 & \int (d - c^2 dx^2)^3 (fx)^m (a + \operatorname{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6336} \\
 & -bc \int \frac{d^3 (fx)^{m+1} \left( -\frac{c^6 x^6}{m+7} + \frac{3c^4 x^4}{m+5} - \frac{3c^2 x^2}{m+3} + \frac{1}{m+1} \right)}{f \sqrt{cx-1} \sqrt{cx+1}} dx - \frac{c^6 d^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7 (m+7)} + \\
 & \quad \frac{3c^4 d^3 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5 (m+5)} - \frac{3c^2 d^3 (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3 (m+3)} + \\
 & \quad \frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
 & \quad \downarrow \text{27} \\
 & - \frac{bcd^3 \int \frac{(fx)^{m+1} \left( -\frac{c^6 x^6}{m+7} + \frac{3c^4 x^4}{m+5} - \frac{3c^2 x^2}{m+3} + \frac{1}{m+1} \right)}{\sqrt{cx-1} \sqrt{cx+1}} dx}{f} - \frac{c^6 d^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7 (m+7)} + \\
 & \quad \frac{3c^4 d^3 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5 (m+5)} - \frac{3c^2 d^3 (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3 (m+3)} + \\
 & \quad \frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
 & \quad \downarrow \text{2113} \\
 & - \frac{bcd^3 \sqrt{c^2 x^2 - 1} \int \frac{(fx)^{m+1} \left( -\frac{c^6 x^6}{m+7} + \frac{3c^4 x^4}{m+5} - \frac{3c^2 x^2}{m+3} + \frac{1}{m+1} \right)}{\sqrt{c^2 x^2 - 1}} dx}{f \sqrt{cx-1} \sqrt{cx+1}} - \frac{c^6 d^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7 (m+7)} + \\
 & \quad \frac{3c^4 d^3 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5 (m+5)} - \frac{3c^2 d^3 (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3 (m+3)} + \\
 & \quad \frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
 & \quad \downarrow \text{2340} \\
 & - \frac{bcd^3 \sqrt{c^2 x^2 - 1} \left( \int \frac{(fx)^{m+1} \left( \frac{(m+9)(2m+13)x^4 c^6}{(m+5)(m+7)} - \frac{3(m+7)x^2 c^4}{m+3} + \frac{(m+7)c^2}{m+1} \right)}{\sqrt{c^2 x^2 - 1}} dx - \frac{c^4 \sqrt{c^2 x^2 - 1} (fx)^{m+6}}{f^5 (m+7)^2} \right)}{f \sqrt{cx-1} \sqrt{cx+1}} - \\
 & \quad \frac{c^6 d^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7 (m+7)} + \frac{3c^4 d^3 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5 (m+5)} - \\
 & \quad \frac{3c^2 d^3 (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3 (m+3)} + \frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
 & \quad \downarrow \text{1590}
 \end{aligned}$$

---

3.145.  $\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx$

$$bcd^3\sqrt{c^2x^2-1} \left( \frac{c^4(fx)^{m+1} \left( \frac{(m+5)(m+7)}{m+1} - \frac{c^2(m^4+27m^3+284m^2+1329m+2271)x^2}{(m+3)(m+5)(m+7)} \right)}{\frac{\sqrt{c^2x^2-1}}{c^2(m+5)}} dx + \frac{c^4(m+9)(2m+13)\sqrt{c^2x^2-1}(fx)^{m+4}}{f^3(m+5)^2(m+7)} - \frac{c^4\sqrt{c^2x^2-1}}{f^5(m)} \right)$$

$$\frac{c^6d^3(fx)^{m+7}(a + \operatorname{barccosh}(cx))}{f^7(m+7)} + \frac{3c^4d^3(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{3c^2d^3(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^3(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)}$$

↓ 27

$$bcd^3\sqrt{c^2x^2-1} \left( \frac{c^2 \int \frac{(fx)^{m+1} \left( \frac{(m+5)(m+7)}{m+1} - \frac{c^2(m^4+27m^3+284m^2+1329m+2271)x^2}{(m+3)(m+5)(m+7)} \right)}{\sqrt{c^2x^2-1}} dx}{m+5} + \frac{c^4(m+9)(2m+13)\sqrt{c^2x^2-1}(fx)^{m+4}}{f^3(m+5)^2(m+7)} - \frac{c^4\sqrt{c^2x^2-1}}{f^5(m)} \right)$$

$$\frac{c^6d^3(fx)^{m+7}(a + \operatorname{barccosh}(cx))}{f^7(m+7)} + \frac{3c^4d^3(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{3c^2d^3(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^3(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)}$$

↓ 363

$$bcd^3\sqrt{c^2x^2-1} \left( \frac{c^2 \left( \frac{3(35m^3+455m^2+1813m+2161) \int \frac{(fx)^{m+1}}{\sqrt{c^2x^2-1}} dx}{(m+1)(m+3)^2(m+5)(m+7)} - \frac{(m^4+27m^3+284m^2+1329m+2271)\sqrt{c^2x^2-1}(fx)^{m+2}}{f(m+3)^2(m+5)(m+7)} \right)}{m+5} + \frac{c^4(m+9)(2m+13)\sqrt{c^2x^2-1}(fx)^{m+4}}{f^3(m+5)^2(m+7)} - \frac{c^4\sqrt{c^2x^2-1}}{f^5(m)} \right)$$

$$\frac{c^6d^3(fx)^{m+7}(a + \operatorname{barccosh}(cx))}{f^7(m+7)} + \frac{3c^4d^3(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{3c^2d^3(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^3(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)}$$

↓ 279

$$bcd^3\sqrt{c^2x^2-1} \left( \frac{c^2 \left( \frac{3(35m^3+455m^2+1813m+2161)\sqrt{1-c^2x^2} \int \frac{(fx)^{m+1}}{\sqrt{1-c^2x^2}} dx}{(m+1)(m+3)^2(m+5)(m+7)\sqrt{c^2x^2-1}} - \frac{(m^4+27m^3+284m^2+1329m+2271)\sqrt{c^2x^2-1}(fx)^{m+2}}{f(m+3)^2(m+5)(m+7)} \right)}{m+5} + \frac{c^4(m+9)}{c^2(m+7)} \right)$$


---


$$\frac{c^6d^3(fx)^{m+7}(a + \operatorname{barccosh}(cx))}{f^7(m+7)} + \frac{3c^4d^3(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{3c^2d^3(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^3(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)}$$

↓ 278

$$bcd^3\sqrt{c^2x^2-1} \left( \frac{c^2 \left( \frac{3(35m^3+455m^2+1813m+2161)\sqrt{1-c^2x^2}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{f(m+1)(m+2)(m+3)^2(m+5)(m+7)\sqrt{c^2x^2-1}} - \frac{(m^4+27m^3+284m^2+1329m+2271)}{f(m+3)^2(m+5)(m+7)} \right)}{m+5} + \frac{c^4(m+9)}{c^2(m+7)} \right)$$


---


$$f\sqrt{cx-1}\sqrt{cx+1}$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `(d^3*(f*x)^(1+m)*(a + b*ArcCosh[c*x]))/(f*(1+m)) - (3*c^2*d^3*(f*x)^(3+m)*(a + b*ArcCosh[c*x]))/(f^3*(3+m)) + (3*c^4*d^3*(f*x)^(5+m)*(a + b*ArcCosh[c*x]))/(f^5*(5+m)) - (c^6*d^3*(f*x)^(7+m)*(a + b*ArcCosh[c*x]))/(f^7*(7+m)) - (b*c*d^3*Sqrt[-1 + c^2*x^2]*(-(c^4*(f*x)^(6+m)*Sqrt[-1 + c^2*x^2]))/(f^5*(7+m)^2)) + ((c^4*(9+m)*(13+2*m)*(f*x)^(4+m)*Sqrt[-1 + c^2*x^2]))/(f^3*(5+m)^2*(7+m)) + (c^2*(-(((2271 + 1329*m + 284*m^2 + 27*m^3 + m^4)*(f*x)^(2+m)*Sqrt[-1 + c^2*x^2]))/(f*(3+m)^2*(5+m)*(7+m))) + (3*(2161 + 1813*m + 455*m^2 + 35*m^3)*(f*x)^(2+m)*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2+m)/2, (4+m)/2, c^2*x^2]))/(f*(1+m)*(2+m)*(3+m)^2*(5+m)*(7+m)*Sqrt[-1 + c^2*x^2]))/(5+m)/(c^2*(7+m)))/(f*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.145.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 279 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^(m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`
- rule 363 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^(m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`
- rule 1590 `Int[((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]`
- rule 2113 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[P_x*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

```
rule 2340 Int[(Pq_)*((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[
{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(c*x)^(m + q - 1)
)*((a + b*x^2)^(p + 1)/(b*c^(q - 1)*(m + q + 2*p + 1))), x] + Simp[1/(b*(m
+ q + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^p*ExpandToSum[b*(m + q + 2*p + 1)
*Pq - b*f*(m + q + 2*p + 1)*x^q - a*f*(m + q - 1)*x^(q - 2), x], x] /;
GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0]] /; FreeQ[{a, b, c, m, p}, x] && PolyQ
[Pq, x] && (!IGtQ[m, 0] || IGtQ[p + 1/2, -1])
```

```
rule 6336 Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)
)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]
```

### 3.145.4 Maple [F]

$$\int (fx)^m (-c^2 dx^2 + d)^3 (a + b \operatorname{arccosh}(cx)) dx$$

```
input int((f*x)^m*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x)
```

```
output int((f*x)^m*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x)
```

### 3.145.5 Fracas [F]

$$\int (fx)^m (d - c^2 dx^2)^3 (a + b \operatorname{arccosh}(cx)) dx = \int -(c^2 dx^2 - d)^3 (b \operatorname{arccosh}(cx) + a)(fx)^m dx$$

```
input integrate((f*x)^m*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fracas")
```

```
output integral(-(a*c^6*d^3*x^6 - 3*a*c^4*d^3*x^4 + 3*a*c^2*d^3*x^2 - a*d^3 + (b*
c^6*d^3*x^6 - 3*b*c^4*d^3*x^4 + 3*b*c^2*d^3*x^2 - b*d^3)*arccosh(c*x))*(f*
x)^m, x)
```

## 3.145.6 Sympy [F(-1)]

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `Timed out`

## 3.145.7 Maxima [F]

$$\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int -(c^2 dx^2 - d)^3 (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-a*c^6*d^3*f^m*x^7*x^m/(m + 7) + 3*a*c^4*d^3*f^m*x^5*x^m/(m + 5) - 3*a*c^2*d^3*f^m*x^3*x^m/(m + 3) + (f*x)^(m + 1)*a*d^3/(f*(m + 1)) - ((m^3 + 9*m^2 + 23*m + 15)*b*c^6*d^3*f^m*x^7 - 3*(m^3 + 11*m^2 + 31*m + 21)*b*c^4*d^3*f^m*x^5 + 3*(m^3 + 13*m^2 + 47*m + 35)*b*c^2*d^3*f^m*x^3 - (m^3 + 15*m^2 + 71*m + 105)*b*d^3*f^m*x)*x^m*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(m^4 + 16*m^3 + 86*m^2 + 176*m + 105) - integrate(((m^3 + 9*m^2 + 23*m + 15)*b*c^7*d^3*f^m*x^7 - 3*(m^3 + 11*m^2 + 31*m + 21)*b*c^5*d^3*f^m*x^5 + 3*(m^3 + 13*m^2 + 47*m + 35)*b*c^3*d^3*f^m*x^3 - (m^3 + 15*m^2 + 71*m + 105)*b*c*d^3*f^m*x)*x^m/((m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c^3*x^3 - (m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c*x + ((m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c^2*x^2 - m^4 - 16*m^3 - 86*m^2 - 176*m - 105)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) + integrate(((m^3 + 9*m^2 + 23*m + 15)*b*c^8*d^3*f^m*x^8 - 3*(m^3 + 11*m^2 + 31*m + 21)*b*c^6*d^3*f^m*x^6 + 3*(m^3 + 13*m^2 + 47*m + 35)*b*c^4*d^3*f^m*x^4 - (m^3 + 15*m^2 + 71*m + 105)*b*c^2*d^3*f^m*x^2)*x^m/((m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c^2*x^2 - m^4 - 16*m^3 - 86*m^2 - 176*m - 105), x)`

**3.145.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.145.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^3 (fx)^m dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^3*(f*x)^m,x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^3*(f*x)^m, x)`



### 3.146 $\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.146.1 Optimal result . . . . .	1240
3.146.2 Mathematica [A] (verified) . . . . .	1241
3.146.3 Rubi [A] (verified) . . . . .	1241
3.146.4 Maple [F] . . . . .	1245
3.146.5 Fricas [F] . . . . .	1245
3.146.6 Sympy [F] . . . . .	1245
3.146.7 Maxima [F] . . . . .	1246
3.146.8 Giac [F(-2)] . . . . .	1246
3.146.9 Mupad [F(-1)] . . . . .	1247

#### 3.146.1 Optimal result

Integrand size = 27, antiderivative size = 307

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{bcd^2(38 + 13m + m^2) (fx)^{2+m} (1 - c^2 x^2)}{f^2(3 + m)^2(5 + m)^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{bc^3 d^2 (fx)^{4+m} (1 - c^2 x^2)}{f^4(5 + m)^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{d^2 (fx)^{1+m} (a + \operatorname{barccosh}(cx))}{f(1 + m)}$$

$$- \frac{2c^2 d^2 (fx)^{3+m} (a + \operatorname{barccosh}(cx))}{f^3(3 + m)} + \frac{c^4 d^2 (fx)^{5+m} (a + \operatorname{barccosh}(cx))}{f^5(5 + m)}$$

$$- \frac{bcd^2(149 + 100m + 15m^2) (fx)^{2+m} \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2 x^2\right)}{f^2(1 + m)(2 + m)(3 + m)^2(5 + m)^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
d^2*(f*x)^(1+m)*(a+b*arccosh(c*x))/f/(1+m)-2*c^2*d^2*(f*x)^(3+m)*(a+b*arccosh(c*x))/f^3/(3+m)+c^4*d^2*(f*x)^(5+m)*(a+b*arccosh(c*x))/f^5/(5+m)-b*c*d^2*(m^2+13*m+38)*(f*x)^(2+m)*(-c^2*x^2+1)/f^2/(3+m)^2/(5+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*c^3*d^2*(f*x)^(4+m)*(-c^2*x^2+1)/f^4/(5+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*c*d^2*(15*m^2+100*m+149)*(f*x)^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/f^2/(m^2+3*m+2)/(m^2+8*m+15)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.146.2 Mathematica [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 290, normalized size of antiderivative = 0.94

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= d^2 x (fx)^m \left( \frac{a + \operatorname{barccosh}(cx)}{1 + m} - \frac{2c^2 x^2 (a + \operatorname{barccosh}(cx))}{3 + m} + \frac{c^4 x^4 (a + \operatorname{barccosh}(cx))}{5 + m} \right.$$

$$- \frac{bcx \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2 x^2\right)}{(2 + 3m + m^2) \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{2bc^3 x^3 \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4+m}{2}, \frac{6+m}{2}, c^2 x^2\right)}{(12 + 7m + m^2) \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$\left. - \frac{bc^5 x^5 \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{6+m}{2}, \frac{8+m}{2}, c^2 x^2\right)}{(5 + m)(6 + m) \sqrt{-1 + cx} \sqrt{1 + cx}} \right)$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`output `d^2*x*(f*x)^m*((a + b*ArcCosh[c*x])/(1 + m) - (2*c^2*x^2*(a + b*ArcCosh[c*x]))/(3 + m) + (c^4*x^4*(a + b*ArcCosh[c*x]))/(5 + m) - (b*c*x*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, c^2*x^2])/((2 + 3*m + m^2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (2*b*c^3*x^3*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m)/2, c^2*x^2])/((12 + 7*m + m^2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (b*c^5*x^5*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (6 + m)/2, (8 + m)/2, c^2*x^2])/((5 + m)*(6 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))`**3.146.3 Rubi [A] (verified)**Time = 0.72 (sec) , antiderivative size = 306, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$ , Rules used = {6336, 27, 1905, 1590, 27, 363, 279, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^2 (fx)^m (a + \operatorname{barccosh}(cx)) dx$$

↓ 6336

$$\begin{aligned}
& -bc \int \frac{d^2(fx)^{m+1} \left( \frac{c^4 x^4}{m+5} - \frac{2c^2 x^2}{m+3} + \frac{1}{m+1} \right)}{f \sqrt{cx-1} \sqrt{cx+1}} dx + \frac{c^4 d^2(fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \\
& \quad \frac{2c^2 d^2(fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 27 \\
& - \frac{bcd^2 \int \frac{(fx)^{m+1} \left( \frac{c^4 x^4}{m+5} - \frac{2c^2 x^2}{m+3} + \frac{1}{m+1} \right)}{\sqrt{cx-1} \sqrt{cx+1}} dx}{f} + \frac{c^4 d^2(fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \\
& \quad \frac{2c^2 d^2(fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 1905 \\
& - \frac{bcd^2 \sqrt{c^2 x^2 - 1} \int \frac{(fx)^{m+1} \left( \frac{c^4 x^4}{m+5} - \frac{2c^2 x^2}{m+3} + \frac{1}{m+1} \right)}{\sqrt{c^2 x^2 - 1}} dx}{f \sqrt{cx-1} \sqrt{cx+1}} + \frac{c^4 d^2(fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \\
& \quad \frac{2c^2 d^2(fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 1590 \\
& - \frac{bcd^2 \sqrt{c^2 x^2 - 1} \left( \int \frac{c^2 (fx)^{m+1} \left( \frac{m+5}{m+1} - \frac{c^2 (m^2+13m+38)x^2}{(m+3)(m+5)} \right)}{\sqrt{c^2 x^2 - 1}} dx + \frac{c^2 \sqrt{c^2 x^2 - 1} (fx)^{m+4}}{f^3(m+5)^2} \right)}{f \sqrt{cx-1} \sqrt{cx+1}} + \\
& \quad \frac{c^4 d^2(fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{2c^2 d^2(fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 27 \\
& - \frac{bcd^2 \sqrt{c^2 x^2 - 1} \left( \int \frac{(fx)^{m+1} \left( \frac{m+5}{m+1} - \frac{c^2 (m^2+13m+38)x^2}{(m+3)(m+5)} \right)}{\sqrt{c^2 x^2 - 1}} dx + \frac{c^2 \sqrt{c^2 x^2 - 1} (fx)^{m+4}}{f^3(m+5)^2} \right)}{m+5} + \\
& \quad \frac{c^4 d^2(fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{2c^2 d^2(fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 363
\end{aligned}$$

$$\begin{aligned}
& \frac{bcd^2\sqrt{c^2x^2-1} \left( \frac{(15m^2+100m+149) \int \frac{(fx)^{m+1}}{\sqrt{c^2x^2-1}} dx}{(m+1)(m+3)^2(m+5)} - \frac{(m^2+13m+38)\sqrt{c^2x^2-1}(fx)^{m+2}}{f(m+3)^2(m+5)} + \frac{c^2\sqrt{c^2x^2-1}(fx)^{m+4}}{f^3(m+5)^2} \right)}{f^5(m+5)} \\
& + \frac{c^4d^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{2c^2d^2(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \qquad \qquad \qquad \downarrow 279 \\
& \frac{bcd^2\sqrt{c^2x^2-1} \left( \frac{(15m^2+100m+149)\sqrt{1-c^2x^2} \int \frac{(fx)^{m+1}}{\sqrt{1-c^2x^2}} dx}{(m+1)(m+3)^2(m+5)\sqrt{c^2x^2-1}} - \frac{(m^2+13m+38)\sqrt{c^2x^2-1}(fx)^{m+2}}{f(m+3)^2(m+5)} + \frac{c^2\sqrt{c^2x^2-1}(fx)^{m+4}}{f^3(m+5)^2} \right)}{f^5(m+5)} \\
& + \frac{c^4d^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{2c^2d^2(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \qquad \qquad \qquad \downarrow 278 \\
& \frac{c^4d^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} - \frac{2c^2d^2(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& - \frac{bcd^2\sqrt{c^2x^2-1} \left( \frac{c^2\sqrt{c^2x^2-1}(fx)^{m+4}}{f^3(m+5)^2} + \frac{(15m^2+100m+149)\sqrt{1-c^2x^2}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{f(m+1)(m+2)(m+3)^2(m+5)\sqrt{c^2x^2-1}} - \frac{(m^2+13m+38)\sqrt{c^2x^2-1}(fx)^{m+2}}{f(m+3)^2(m+5)} \right)}{f^5(m+5)} \\
& \qquad \qquad \qquad \frac{f\sqrt{cx-1}\sqrt{cx+1}}{f(m+1)}
\end{aligned}$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output 
$$\begin{aligned}
& (d^2(fx)^{(1+m)}(a + b\operatorname{ArcCosh}[cx]))/(f(1+m)) - (2*c^2*d^2(fx)^{(3+m)}(a + b\operatorname{ArcCosh}[cx]))/(f^3(3+m)) + (c^4*d^2(fx)^{(5+m)}(a + b\operatorname{ArcCosh}[cx]))/(f^5(5+m)) - (b*c*d^2\sqrt{-1 + c^2*x^2}*((c^2(fx)^{(4+m)}\sqrt{-1 + c^2*x^2})/(f^3(5+m)^2) + (-((38 + 13*m + m^2)*(fx)^{(2+m)}\sqrt{-1 + c^2*x^2})/(f*(3+m)^2*(5+m)))) + ((149 + 100*m + 15*m^2)*(fx)^{(2+m)}\sqrt{1 - c^2*x^2}*\operatorname{Hypergeometric2F1}[1/2, (2+m)/2, (4+m)/2, c^2*x^2])/(f*(1+m)*(2+m)*(3+m)^2*(5+m)*\sqrt{-1 + c^2*x^2}))/((5+m)/(f*\sqrt{-1 + c*x}*\sqrt{1 + c*x}))
\end{aligned}$$

## 3.146.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 279 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^(m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`
- rule 363 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^(m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`
- rule 1590 `Int[((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^(m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]`
- rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(p_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^(m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

### 3.146.4 Maple [F]

$$\int (fx)^m (-c^2 dx^2 + d)^2 (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x)`

### 3.146.5 Fracas [F]

$$\int (fx)^m (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \int (c^2 dx^2 - d)^2 (b \operatorname{arccosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*(f*x)^m, x)`

### 3.146.6 Sympy [F]

$$\begin{aligned} \int (fx)^m (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx &= d^2 \left( \int a (fx)^m dx + \int b (fx)^m \operatorname{acosh}(cx) dx \right. \\ &\quad + \int (-2ac^2 x^2 (fx)^m) dx + \int ac^4 x^4 (fx)^m dx \\ &\quad \left. + \int (-2bc^2 x^2 (fx)^m \operatorname{acosh}(cx)) dx \right. \\ &\quad \left. + \int bc^4 x^4 (fx)^m \operatorname{acosh}(cx) dx \right) \end{aligned}$$

3.146.  $\int (fx)^m (d - c^2 dx^2)^2 (a + b \operatorname{arccosh}(cx)) dx$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `d**2*(Integral(a*(f*x)**m, x) + Integral(b*(f*x)**m*acosh(c*x), x) + Integral(-2*a*c**2*x**2*(f*x)**m, x) + Integral(a*c**4*x**4*(f*x)**m, x) + Integral(-2*b*c**2*x**2*(f*x)**m*acosh(c*x), x) + Integral(b*c**4*x**4*(f*x)**m*acosh(c*x), x))`

### 3.146.7 Maxima [F]

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int (c^2 dx^2 - d)^2 (b \operatorname{arccosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `a*c^4*d^2*f^m*x^5*x^m/(m + 5) - 2*a*c^2*d^2*f^m*x^3*x^m/(m + 3) + (f*x)^(m + 1)*a*d^2/(f*(m + 1)) + ((m^2 + 4*m + 3)*b*c^4*d^2*f^m*x^5 - 2*(m^2 + 6*m + 5)*b*c^2*d^2*f^m*x^3 + (m^2 + 8*m + 15)*b*d^2*f^m*x)*x^m*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(m^3 + 9*m^2 + 23*m + 15) + integrate(((m^2 + 4*m + 3)*b*c^5*d^2*f^m*x^5 - 2*(m^2 + 6*m + 5)*b*c^3*d^2*f^m*x^3 + (m^2 + 8*m + 15)*b*c*d^2*f^m*x)*x^m/((m^3 + 9*m^2 + 23*m + 15)*c^3*x^3 - (m^3 + 9*m^2 + 23*m + 15)*c*x + ((m^3 + 9*m^2 + 23*m + 15)*c^2*x^2 - m^3 - 9*m^2 - 23*m - 15)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) - integrate(((m^2 + 4*m + 3)*b*c^6*d^2*f^m*x^6 - 2*(m^2 + 6*m + 5)*b*c^4*d^2*f^m*x^4 + (m^2 + 8*m + 15)*b*c^2*d^2*f^m*x^2)*x^m/((m^3 + 9*m^2 + 23*m + 15)*c^2*x^2 - m^3 - 9*m^2 - 23*m - 15), x)`

### 3.146.8 Giac [F(-2)]

Exception generated.

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN  
 PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const  
 index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.146.9 Mupad [F(-1)]

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^2 (fx)^m dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^2*(f*x)^m,x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^2*(f*x)^m, x)`



### 3.147 $\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$

3.147.1 Optimal result . . . . .	1248
3.147.2 Mathematica [A] (verified) . . . . .	1249
3.147.3 Rubi [A] (verified) . . . . .	1249
3.147.4 Maple [F] . . . . .	1252
3.147.5 Fracas [F] . . . . .	1252
3.147.6 Sympy [F] . . . . .	1252
3.147.7 Maxima [F] . . . . .	1253
3.147.8 Giac [F(-2)] . . . . .	1253
3.147.9 Mupad [F(-1)] . . . . .	1253

#### 3.147.1 Optimal result

Integrand size = 25, antiderivative size = 184

$$\begin{aligned} & \int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx \\ &= \frac{bcd(fx)^{2+m} \sqrt{-1 + cx} \sqrt{1 + cx}}{f^2(3 + m)^2} + \frac{d(fx)^{1+m} (a + \operatorname{barccosh}(cx))}{f(1 + m)} \\ & \quad - \frac{c^2 d(fx)^{3+m} (a + \operatorname{barccosh}(cx))}{f^3(3 + m)} \\ & \quad - \frac{bcd(7 + 3m)(fx)^{2+m} \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2 x^2\right)}{f^2(1 + m)(2 + m)(3 + m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \end{aligned}$$

```
output d*(f*x)^(1+m)*(a+b*arccosh(c*x))/f/(1+m)-c^2*d*(f*x)^(3+m)*(a+b*arccosh(c*
x))/f^3/(3+m)+b*c*d*(f*x)^(2+m)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/f^2/(3+m)^2-b*
c*d*(7+3*m)*(f*x)^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*
x^2+1)^(1/2)/f^2/(3+m)^2/(m^2+3*m+2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.147.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.04

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx$$

$$= dx (fx)^m \left( \frac{a + \operatorname{barccosh}(cx)}{1 + m} - \frac{c^2 x^2 (a + \operatorname{barccosh}(cx))}{3 + m} \right. \\ \left. - \frac{bcx \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2 x^2\right)}{(2 + 3m + m^2) \sqrt{-1 + cx} \sqrt{1 + cx}} \right. \\ \left. + \frac{bc^3 x^3 \sqrt{1 - c^2 x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4+m}{2}, \frac{6+m}{2}, c^2 x^2\right)}{(12 + 7m + m^2) \sqrt{-1 + cx} \sqrt{1 + cx}} \right)$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `d*x*(f*x)^m*((a + b*ArcCosh[c*x])/(1 + m) - (c^2*x^2*(a + b*ArcCosh[c*x]))/(3 + m) - (b*c*x*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, c^2*x^2])/((2 + 3*m + m^2)*sqrt[-1 + c*x]*sqrt[1 + c*x]) + (b*c^3*x^3*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m)/2, c^2*x^2])/((12 + 7*m + m^2)*sqrt[-1 + c*x]*sqrt[1 + c*x]))`

### 3.147.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6336, 27, 960, 136, 279, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2) (fx)^m (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6336}$$

$$-bc \int \frac{d(fx)^{m+1} (-c^2(m+1)x^2 + m+3)}{f(m^2 + 4m + 3) \sqrt{cx-1} \sqrt{cx+1}} dx - \frac{c^2 d (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)}$$

$$\downarrow \text{27}$$

$$\begin{aligned}
& -\frac{bcd \int \frac{(fx)^{m+1}(-c^2(m+1)x^2+m+3)}{\sqrt{cx-1}\sqrt{cx+1}} dx}{f(m^2+4m+3)} - \frac{c^2 d(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \\
& \quad \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 960 \\
& -\frac{bcd \left( \frac{(3m+7) \int \frac{(fx)^{m+1}}{\sqrt{cx-1}\sqrt{cx+1}} dx}{m+3} - \frac{(m+1)\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2}}{f(m+3)} \right)}{f(m^2+4m+3)} - \frac{c^2 d(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \\
& \quad \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 136 \\
& -\frac{bcd \left( \frac{(3m+7)\sqrt{c^2x^2-1} \int \frac{(fx)^{m+1}}{\sqrt{c^2x^2-1}} dx}{(m+3)\sqrt{cx-1}\sqrt{cx+1}} - \frac{(m+1)\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2}}{f(m+3)} \right)}{f(m^2+4m+3)} - \frac{c^2 d(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \\
& \quad \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 279 \\
& -\frac{bcd \left( \frac{(3m+7)\sqrt{1-c^2x^2} \int \frac{(fx)^{m+1}}{\sqrt{1-c^2x^2}} dx}{(m+3)\sqrt{cx-1}\sqrt{cx+1}} - \frac{(m+1)\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2}}{f(m+3)} \right)}{f(m^2+4m+3)} - \frac{c^2 d(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \\
& \quad \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} \\
& \quad \downarrow 278 \\
& -\frac{c^2 d(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} - \\
& \frac{bcd \left( \frac{(3m+7)\sqrt{1-c^2x^2}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{f(m+2)(m+3)\sqrt{cx-1}\sqrt{cx+1}} - \frac{(m+1)\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2}}{f(m+3)} \right)}{f(m^2+4m+3)}
\end{aligned}$$

input `Int[(f*x)^m*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x]),x]`

output `(d*(f*x)^(1+m)*(a + b*ArcCosh[c*x]))/(f*(1+m)) - (c^2*d*(f*x)^(3+m)*(a + b*ArcCosh[c*x]))/(f^3*(3+m)) - (b*c*d*(-(((1+m)*(f*x)^(2+m)*Sqrt[-1+c*x]*Sqrt[1+c*x])/(f*(3+m))) + ((7+3*m)*(f*x)^(2+m)*Sqrt[1-c^2*x^2]*Hypergeometric2F1[1/2, (2+m)/2, (4+m)/2, c^2*x^2])/(f*(2+m)*(3+m)*Sqrt[-1+c*x]*Sqrt[1+c*x]))/(f*(3+4*m+m^2))`

## 3.147.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 136 `Int[((f_)*(x_)^(p_))*((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[(a*c + b*d*x^2)^m*(f*x)^p, x], x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m]`
- rule 278 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 279 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`
- rule 960 `Int[((e_)*(x_)^(m_))*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 6336 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_))*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

**3.147.4 Maple [F]**

$$\int (fx)^m (-c^2 dx^2 + d) (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x)`

**3.147.5 Fricas [F]**

$$\int (fx)^m (d - c^2 dx^2) (a + b \operatorname{arccosh}(cx)) dx = \int -(c^2 dx^2 - d)(b \operatorname{arccosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*(f*x)^m, x)`

**3.147.6 Sympy [F]**

$$\begin{aligned} \int (fx)^m (d - c^2 dx^2) (a + b \operatorname{arccosh}(cx)) dx = & -d \left( \int (-a(fx)^m) dx \right. \\ & + \int (-b(fx)^m \operatorname{acosh}(cx)) dx \\ & + \int ac^2 x^2 (fx)^m dx \\ & \left. + \int bc^2 x^2 (fx)^m \operatorname{acosh}(cx) dx \right) \end{aligned}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)*(a+b*acosh(c*x)),x)`

output `-d*(Integral(-a*(f*x)**m, x) + Integral(-b*(f*x)**m*acosh(c*x), x) + Integral(a*c**2*x**2*(f*x)**m, x) + Integral(b*c**2*x**2*(f*x)**m*acosh(c*x), x))`

**3.147.7 Maxima [F]**

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \int -(c^2 dx^2 - d)(b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `-a*c^2*d*f^m*x^3*x^m/(m + 3) - (b*c^2*d*f^m*(m + 1)*x^3 - b*d*f^m*(m + 3)*x)*x^m*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(m^2 + 4*m + 3) + (f*x)^(m + 1)*a*d/(f*(m + 1)) - integrate((b*c^3*d*f^m*(m + 1)*x^3 - b*c*d*f^m*(m + 3)*x)*x^m/((m^2 + 4*m + 3)*c^3*x^3 - (m^2 + 4*m + 3)*c*x + ((m^2 + 4*m + 3)*c^2*x^2 - m^2 - 4*m - 3)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) + integrate((b*c^4*d*f^m*(m + 1)*x^4 - b*c^2*d*f^m*(m + 3)*x^2)*x^m/((m^2 + 4*m + 3)*c^2*x^2 - m^2 - 4*m - 3), x)`

**3.147.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.147.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2) (fx)^m dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)*(f*x)^m, x)`

**3.148**  $\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx$

3.148.1 Optimal result . . . . . 1254  
 3.148.2 Mathematica [N/A] . . . . . 1254  
 3.148.3 Rubi [N/A] . . . . . 1255  
 3.148.4 Maple [N/A] (verified) . . . . . 1255  
 3.148.5 Fricas [N/A] . . . . . 1256  
 3.148.6 Sympy [N/A] . . . . . 1256  
 3.148.7 Maxima [N/A] . . . . . 1256  
 3.148.8 Giac [N/A] . . . . . 1257  
 3.148.9 Mupad [N/A] . . . . . 1257

**3.148.1 Optimal result**

Integrand size = 27, antiderivative size = 27

$$\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx = \operatorname{Int}\left(\frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d), x)`

**3.148.2 Mathematica [N/A]**

Not integrable

Time = 3.76 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx = \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2), x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2), x]`

**3.148.3 Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))}{d - c^2 dx^2} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2),x]`

output `$Aborted`

**3.148.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^m_.*((d_) + (e_.)*(x_)^2)^p_. , x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.148.4 Maple [N/A] (verified)**

Not integrable

Time = 1.34 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{-c^2 dx^2 + d} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x)`



**3.148.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{c^2 dx^2 - d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="fricas")`output `integral(-(b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d), x)`**3.148.6 Sympy [N/A]**

Not integrable

Time = 4.55 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.52

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = -\frac{\int \frac{a(fx)^m}{c^2 x^2 - 1} dx + \int \frac{b(fx)^m \operatorname{acosh}(cx)}{c^2 x^2 - 1} dx}{d}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(-c**2*d*x**2+d),x)`output `-(Integral(a*(f*x)**m/(c**2*x**2 - 1), x) + Integral(b*(f*x)**m*acosh(c*x)/(c**2*x**2 - 1), x))/d`**3.148.7 Maxima [N/A]**

Not integrable

Time = 0.81 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.19

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{c^2 dx^2 - d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="maxima")`output `-integrate((b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d), x)`

---

3.148.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx$

**3.148.8 Giac [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{c^2 dx^2 - d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x, algorithm="giac")`output `integrate(-(b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d), x)`**3.148.9 Mupad [N/A]**

Not integrable

Time = 3.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d - c^2 dx^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{d - c^2 dx^2} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2),x)`output `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2), x)`

**3.149** 
$$\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx$$

3.149.1 Optimal result . . . . . 1258  
 3.149.2 Mathematica [N/A] . . . . . 1259  
 3.149.3 Rubi [N/A] . . . . . 1259  
 3.149.4 Maple [N/A] (verified) . . . . . 1261  
 3.149.5 Fricas [N/A] . . . . . 1262  
 3.149.6 Sympy [N/A] . . . . . 1262  
 3.149.7 Maxima [N/A] . . . . . 1262  
 3.149.8 Giac [N/A] . . . . . 1263  
 3.149.9 Mupad [N/A] . . . . . 1263

**3.149.1 Optimal result**

Integrand size = 27, antiderivative size = 27

$$\begin{aligned} & \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^2} dx \\ &= \frac{(fx)^{1+m}(a+b\operatorname{arccosh}(cx))}{2d^2f(1-c^2x^2)} \\ & \quad - \frac{bc(fx)^{2+m}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{2d^2f^2(2+m)\sqrt{-1+cx}\sqrt{1+cx}} \\ & \quad + \frac{(1-m)\operatorname{Int}\left(\frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2}, x\right)}{2d} \end{aligned}$$

```
output 1/2*(f*x)^(1+m)*(a+b*arccosh(c*x))/d^2/f/(-c^2*x^2+1)-1/2*b*c*(f*x)^(2+m)*
hypergeom([3/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/d^2/f^2/(2+
m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/2*(1-m)*Unintegrable((f*x)^m*(a+b*arccosh
(c*x))/(-c^2*d*x^2+d), x)/d
```

**3.149.2 Mathematica [N/A]**

Not integrable

Time = 9.49 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx = \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2, x]`**3.149.3 Rubi [N/A]**

Not integrable

Time = 0.56 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6351, 27, 136, 279, 278, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx \\ & \quad \downarrow \text{6351} \\ & \frac{(1 - m) \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{d(1 - c^2x^2)} dx}{2d} + \frac{bc \int \frac{(fx)^{m+1}}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2d^2 f} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{2d^2 f(1 - c^2x^2)} \\ & \quad \downarrow \text{27} \\ & \frac{(1 - m) \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{1 - c^2x^2} dx}{2d^2} + \frac{bc \int \frac{(fx)^{m+1}}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2d^2 f} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{2d^2 f(1 - c^2x^2)} \\ & \quad \downarrow \text{136} \\ & \frac{(1 - m) \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{1 - c^2x^2} dx}{2d^2} + \frac{bc\sqrt{c^2x^2 - 1} \int \frac{(fx)^{m+1}}{(c^2x^2 - 1)^{3/2}} dx}{2d^2 f\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{2d^2 f(1 - c^2x^2)} \\ & \quad \downarrow \text{279} \end{aligned}$$

---

3.149.  $\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx$

$$\begin{aligned}
& \frac{(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx}{2d^2} - \frac{bc\sqrt{1-c^2x^2} \int \frac{(fx)^{m+1}}{(1-c^2x^2)^{3/2}} dx}{2d^2 f \sqrt{cx-1} \sqrt{cx+1}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2d^2 f (1-c^2x^2)} \\
& \quad \downarrow 278 \\
& \frac{(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx}{2d^2} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2d^2 f (1-c^2x^2)} - \\
& \quad \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{2d^2 f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow 6375 \\
& \frac{(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx}{2d^2} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2d^2 f (1-c^2x^2)} - \\
& \quad \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{2d^2 f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}}
\end{aligned}$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^2,x]`

output `$Aborted`

### 3.149.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 136 `Int[((f_)*(x_))^(p_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[(a*c + b*d*x^2)^m*(f*x)^p, x], x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m+1)/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/2, (m+1)/2+1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 279 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

### 3.149.4 Maple [N/A] (verified)

Not integrable

Time = 1.35 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(-c^2 dx^2 + d)^2} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x)`

**3.149.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.59

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(c^2dx^2 - d)^2} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="fracas")
```

```
output integral((b*arccosh(c*x) + a)*(f*x)^m/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)
```

**3.149.6 Sympy [N/A]**

Not integrable

Time = 64.46 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.15

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx = \int \frac{a(fx)^m}{c^4x^4 - 2c^2x^2 + 1} dx + \int \frac{b(fx)^m \operatorname{acosh}(cx)}{c^4x^4 - 2c^2x^2 + 1} dx$$

```
input integrate((f*x)**m*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**2,x)
```

```
output (Integral(a*(f*x)**m/(c**4*x**4 - 2*c**2*x**2 + 1), x) + Integral(b*(f*x)**m*acosh(c*x)/(c**4*x**4 - 2*c**2*x**2 + 1), x))/d**2
```

**3.149.7 Maxima [N/A]**

Not integrable

Time = 0.76 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.11

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d - c^2dx^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(c^2dx^2 - d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d)^2, x)`

### 3.149.8 Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.11

$$\int \frac{(fx)^m(a + b\operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{(c^2dx^2 - d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d)^2, x)`

### 3.149.9 Mupad [N/A]

Not integrable

Time = 3.05 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m(a + b\operatorname{arccosh}(cx))}{(d - c^2dx^2)^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{(d - c^2dx^2)^2} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^2,x)`

output `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^2, x)`



**3.150** 
$$\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx$$

3.150.1 Optimal result . . . . . 1264  
 3.150.2 Mathematica [N/A] . . . . . 1265  
 3.150.3 Rubi [N/A] . . . . . 1265  
 3.150.4 Maple [N/A] (verified) . . . . . 1268  
 3.150.5 Fricas [N/A] . . . . . 1268  
 3.150.6 Sympy [F(-1)] . . . . . 1269  
 3.150.7 Maxima [N/A] . . . . . 1269  
 3.150.8 Giac [N/A] . . . . . 1269  
 3.150.9 Mupad [N/A] . . . . . 1270

**3.150.1 Optimal result**

Integrand size = 27, antiderivative size = 27

$$\begin{aligned} & \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^3} dx \\ &= \frac{(fx)^{1+m}(a+b\operatorname{arccosh}(cx))}{4d^3f(1-c^2x^2)^2} + \frac{(3-m)(fx)^{1+m}(a+b\operatorname{arccosh}(cx))}{8d^3f(1-c^2x^2)} \\ & \quad - \frac{bc(3-m)(fx)^{2+m}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{8d^3f^2(2+m)\sqrt{-1+cx}\sqrt{1+cx}} \\ & \quad - \frac{bc(fx)^{2+m}\sqrt{1-c^2x^2}\operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{4d^3f^2(2+m)\sqrt{-1+cx}\sqrt{1+cx}} \\ & \quad + \frac{(1-m)(3-m)\operatorname{Int}\left(\frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{d-c^2dx^2}, x\right)}{8d^2} \end{aligned}$$

```
output 1/4*(f*x)^(1+m)*(a+b*arccosh(c*x))/d^3/f/(-c^2*x^2+1)^2+1/8*(3-m)*(f*x)^(1+m)*(a+b*arccosh(c*x))/d^3/f/(-c^2*x^2+1)-1/8*b*c*(3-m)*(f*x)^(2+m)*hypergeom([3/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/d^3/f^2/(2+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/4*b*c*(f*x)^(2+m)*hypergeom([5/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/d^3/f^2/(2+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/8*(1-m)*(3-m)*Unintegrable((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d),x)/d^2
```

**3.150.2 Mathematica [N/A]**

Not integrable

Time = 12.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3, x]`**3.150.3 Rubi [N/A]**

Not integrable

Time = 0.84 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6351, 27, 136, 279, 278, 6351, 136, 279, 278, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx \\ & \quad \downarrow \text{6351} \\ & \frac{(3 - m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d^2 (1 - c^2 x^2)^2} dx}{4d} - \frac{bc \int \frac{(fx)^{m+1}}{(cx-1)^{5/2} (cx+1)^{5/2}} dx}{4d^3 f} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1 - c^2 x^2)^2} \\ & \quad \downarrow \text{27} \\ & \frac{(3 - m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(1 - c^2 x^2)^2} dx}{4d^3} - \frac{bc \int \frac{(fx)^{m+1}}{(cx-1)^{5/2} (cx+1)^{5/2}} dx}{4d^3 f} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1 - c^2 x^2)^2} \\ & \quad \downarrow \text{136} \\ & \frac{(3 - m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(1 - c^2 x^2)^2} dx}{4d^3} - \frac{bc \sqrt{c^2 x^2 - 1} \int \frac{(fx)^{m+1}}{(c^2 x^2 - 1)^{5/2}} dx}{4d^3 f \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1 - c^2 x^2)^2} \\ & \quad \downarrow \text{279} \end{aligned}$$

---

3.150.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$

$$\begin{aligned}
& \frac{(3-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{4d^3} - \frac{bc\sqrt{1-c^2x^2} \int \frac{(fx)^{m+1}}{(1-c^2x^2)^{5/2}} dx}{4d^3 f \sqrt{cx-1} \sqrt{cx+1}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1-c^2x^2)^2} \\
& \quad \downarrow 278 \\
& \frac{(3-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{4d^3} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1-c^2x^2)^2} - \\
& \quad \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{4d^3 f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow 6351 \\
& \frac{(3-m) \left( \frac{1}{2}(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx + \frac{bc \int \frac{(fx)^{m+1}}{(cx-1)^{3/2} (cx+1)^{3/2}} dx}{2f} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2f(1-c^2x^2)} \right)}{4d^3} + \\
& \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1-c^2x^2)^2} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{4d^3 f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow 136 \\
& \frac{(3-m) \left( \frac{1}{2}(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx + \frac{bc\sqrt{c^2x^2-1} \int \frac{(fx)^{m+1}}{(c^2x^2-1)^{3/2}} dx}{2f\sqrt{cx-1}\sqrt{cx+1}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2f(1-c^2x^2)} \right)}{4d^3} + \\
& \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1-c^2x^2)^2} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{4d^3 f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow 279 \\
& \frac{(3-m) \left( \frac{1}{2}(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx - \frac{bc\sqrt{1-c^2x^2} \int \frac{(fx)^{m+1}}{(1-c^2x^2)^{3/2}} dx}{2f\sqrt{cx-1}\sqrt{cx+1}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2f(1-c^2x^2)} \right)}{4d^3} + \\
& \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1-c^2x^2)^2} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{4d^3 f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow 278 \\
& \frac{(3-m) \left( \frac{1}{2}(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2f(1-c^2x^2)} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{m+2}{2}\right)}{2f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}} \right)}{4d^3} + \\
& \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1-c^2x^2)^2} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{4d^3 f^2 (m+2) \sqrt{cx-1} \sqrt{cx+1}}
\end{aligned}$$

---

3.150.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d-c^2dx^2)^3} dx$

↓ 6375

$$(3-m) \left( \frac{1}{2}(1-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{1-c^2x^2} dx + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{2f(1-c^2x^2)} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{3}{2}, \frac{m+2}{2}, \frac{m+2}{2}, c^2x^2\right)}{2f^2(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)$$


---


$$\frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{4d^3 f (1-c^2x^2)^2} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(\frac{5}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{4d^3 f^2 (m+2)\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^3,x]`

output `$Aborted`

### 3.150.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 136 `Int[((f_)*(x_))^(p_)*((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[(a*c + b*d*x^2)^m*(f*x)^p, x], x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m+1)/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/2, (m+1)/2+1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 279 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

```
rule 6351 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])
```

```
rule 6375 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]
```

### 3.150.4 Maple [N/A] (verified)

Not integrable

Time = 1.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(-c^2 dx^2 + d)^3} dx$$

```
input int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x)
```

```
output int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x)
```

### 3.150.5 Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.11

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{(c^2 dx^2 - d)^3} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="fricas")
```

---

3.150.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx$

output `integral(-(b*arccosh(c*x) + a)*(f*x)^m/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

### 3.150.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**3,x)`

output Timed out

### 3.150.7 Maxima [N/A]

Not integrable

Time = 0.72 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.19

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(c^2 dx^2 - d)^3} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="maxima")`

output `-integrate((b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d)^3, x)`

### 3.150.8 Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int -\frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(c^2 dx^2 - d)^3} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^3,x, algorithm="giac")`

output `integrate(-(b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d)^3, x)`

---

3.150.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^3} dx$

**3.150.9 Mupad [N/A]**

Not integrable

Time = 3.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{(d - c^2 dx^2)^3} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^3,x)`output `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^3, x)`

### 3.151 $\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

3.151.1 Optimal result . . . . .	1271
3.151.2 Mathematica [A] (verified) . . . . .	1272
3.151.3 Rubi [A] (verified) . . . . .	1273
3.151.4 Maple [F] . . . . .	1277
3.151.5 Fricas [F] . . . . .	1278
3.151.6 Sympy [F(-1)] . . . . .	1278
3.151.7 Maxima [F] . . . . .	1278
3.151.8 Giac [F(-2)] . . . . .	1279
3.151.9 Mupad [F(-1)] . . . . .	1279

#### 3.151.1 Optimal result

Integrand size = 29, antiderivative size = 723

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = -\frac{bcd^2 (fx)^{2+m} \sqrt{d - c^2 dx^2}}{f^2(2+m)(6+m)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{15bcd^2 (fx)^{2+m} \sqrt{d - c^2 dx^2}}{f^2(2+m)^2(4+m)(6+m)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{5bcd^2 (fx)^{2+m} \sqrt{d - c^2 dx^2}}{f^2(2+m)(4+m)(6+m)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{5bc^3 d^2 (fx)^{4+m} \sqrt{d - c^2 dx^2}}{f^4(4+m)^2(6+m)\sqrt{-1+cx}\sqrt{1+cx}} + \frac{2bc^3 d^2 (fx)^{4+m} \sqrt{d - c^2 dx^2}}{f^4(4+m)(6+m)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{bc^5 d^2 (fx)^{6+m} \sqrt{d - c^2 dx^2}}{f^6(6+m)^2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{15d^2 (fx)^{1+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f(6+m)(8+6m+m^2)}$$

$$+ \frac{5d (fx)^{1+m} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))}{f(4+m)(6+m)} + \frac{(fx)^{1+m} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))}{f(6+m)}$$

$$+ \frac{15d^2 (fx)^{1+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2 x^2\right)}{f(4+m)(6+m)(2+3m+m^2)\sqrt{1-cx}\sqrt{1+cx}}$$

$$- \frac{15bcd^2 (fx)^{2+m} \sqrt{d - c^2 dx^2} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2 x^2\right)}{f^2(1+m)(2+m)^2(4+m)(6+m)\sqrt{-1+cx}\sqrt{1+cx}}$$



output  $5*d*(f*x)^{(1+m)}*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))/f/(4+m)/(6+m)+(f*x)^{(1+m)}*(-c^2*d*x^2+d)^{(5/2)}*(a+b*\operatorname{arccosh}(c*x))/f/(6+m)+15*d^2*(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/f/(6+m)/(m^2+6*m+8)+15*d^2*(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))*\operatorname{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], c^2*x^2)*(-c^2*d*x^2+d)^{(1/2)}/f/(6+m)/(m^3+7*m^2+14*m+8)/(-c*x+1)^{(1/2)}/(c*x+1)^{(1/2)}-b*c*d^2*(f*x)^{(2+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)/(6+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-15*b*c*d^2*(f*x)^{(2+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)^2/(4+m)/(6+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-5*b*c*d^2*(f*x)^{(2+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^2/(6+m)/(m^2+6*m+8)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+5*b*c^3*d^2*(f*x)^{(4+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^4/(4+m)^2/(6+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+2*b*c^3*d^2*(f*x)^{(4+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^4/(4+m)/(6+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b*c^5*d^2*(f*x)^{(6+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^6/(6+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-15*b*c*d^2*(f*x)^{(2+m)}*\operatorname{hypergeom}([1, 1+1/2*m, 1+1/2*m], [2+1/2*m, 3/2+1/2*m], c^2*x^2)*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)^2/(6+m)/(m^2+5*m+4)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.151.2 Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 350, normalized size of antiderivative = 0.48

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \frac{d^2 x (fx)^m \sqrt{d - c^2 dx^2}}{\left( \frac{5bcx \left( -\frac{1}{2+m} + \frac{c^2 x^2}{4+m} \right)}{4+m} - bcx \left( \frac{1}{2+m} - \frac{2c^2 x^2}{4+m} + \frac{c^4 x^4}{6+m} \right) - \frac{5(-1+cx)^{3/2}(1+cx)^3}{4+m} \right)}$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output  $(d^2*x*(f*x)^m*\text{Sqrt}[d - c^2*d*x^2]*((5*b*c*x*(-(2 + m)^{-1} + (c^2*x^2)/(4 + m)))/(4 + m) - b*c*x*((2 + m)^{-1} - (2*c^2*x^2)/(4 + m) + (c^4*x^4)/(6 + m)) - (5*(-1 + c*x)^{(3/2)}*(1 + c*x)^{(3/2)}*(a + b*\text{ArcCosh}[c*x]))/(4 + m) + (-1 + c*x)^{(5/2)}*(1 + c*x)^{(5/2)}*(a + b*\text{ArcCosh}[c*x]) + (15*(-(b*c*x) + (2 + m)*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(a + b*\text{ArcCosh}[c*x]) - ((2 + m)*(\text{Sqrt}[1 - c^2*x^2]*(a + b*\text{ArcCosh}[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])) + (b*c*x*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(2 + m)))/(1 + m)))/((2 + m)^2*(4 + m)))/((6 + m)*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])$

### 3.151.3 Rubi [A] (verified)

Time = 1.70 (sec) , antiderivative size = 548, normalized size of antiderivative = 0.76, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$ , Rules used = {6345, 82, 244, 2009, 6345, 25, 82, 244, 2009, 6341, 17, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{5/2} (fx)^m (a + \text{barccosh}(cx)) dx$$

$$\downarrow \text{6345}$$

$$\frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx)) dx}{m + 6} - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1 - cx)^2 (cx + 1)^2 dx}{f(m + 6) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 6)}$$

$$\downarrow \text{82}$$

$$\frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx)) dx}{m + 6} - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1 - c^2 x^2)^2 dx}{f(m + 6) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 6)}$$

$$\downarrow \text{244}$$

$$\frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx)) dx}{m + 6} - \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \left( (fx)^{m+1} - \frac{2c^2 (fx)^{m+3}}{f^2} + \frac{c^4 (fx)^{m+5}}{f^4} \right) dx}{f(m + 6) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 6)}$$

---

3.151.  $\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx)) dx$

$$\begin{aligned}
& \downarrow 2009 \\
& \frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx}{m+6} + \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \\
& \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} \\
& \downarrow 6345 \\
& 5d \left( \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} + \frac{bcd \sqrt{d - c^2 dx^2} \int -(fx)^{m+1} (1-cx)(cx+1) dx}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} + \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} \right) \\
& \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} \\
& \downarrow 25 \\
& 5d \left( \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} - \frac{bcd \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1-cx)(cx+1) dx}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} + \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} \right) \\
& \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} \\
& \downarrow 82 \\
& 5d \left( \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} - \frac{bcd \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1 - c^2 x^2) dx}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} + \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} \right) + \\
& \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} \\
& \downarrow 244 \\
& 5d \left( \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} - \frac{bcd \sqrt{d - c^2 dx^2} \int \left( (fx)^{m+1} - \frac{c^2 (fx)^{m+3}}{f^2} \right) dx}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} + \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} \right) \\
& \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} \\
& \downarrow 2009 \\
& 3.151. \quad \int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx
\end{aligned}$$

$$5d \left( \frac{3d \int (fx)^m \sqrt{d-c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} + \frac{(d-c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} - \frac{bcd \sqrt{d-c^2 dx^2} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} \right)$$

$$\frac{(d-c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d-c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}}$$

↓ 6341

$$5d \left( \frac{3d \left( -\frac{\sqrt{d-c^2 dx^2} \int (fx)^m (a + \operatorname{barccosh}(cx)) dx}{(m+2) \sqrt{cx-1} \sqrt{cx+1}} - \frac{bc \sqrt{d-c^2 dx^2} \int (fx)^{m+1} dx}{f(m+2) \sqrt{cx-1} \sqrt{cx+1}} + \frac{\sqrt{d-c^2 dx^2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+2)} \right)}{m+4} + \frac{(d-c^2 dx^2)^{3/2} (fx)^{m+1}}{f(m+4)} \right)$$

$$\frac{(d-c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d-c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}}$$

↓ 17

$$5d \left( \frac{3d \left( -\frac{\sqrt{d-c^2 dx^2} \int (fx)^m (a + \operatorname{barccosh}(cx)) dx}{(m+2) \sqrt{cx-1} \sqrt{cx+1}} + \frac{\sqrt{d-c^2 dx^2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \sqrt{d-c^2 dx^2} (fx)^{m+2}}{f^2(m+2)^2 \sqrt{cx-1} \sqrt{cx+1}} \right)}{m+4} + \frac{(d-c^2 dx^2)^{3/2} (fx)^{m+1}}{f(m+4)} \right)$$

$$\frac{(d-c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d-c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}}$$

↓ 6364

$$5d \left( \frac{3d \left( -\frac{\sqrt{d-c^2 dx^2} \left( \frac{bc (fx)^{m+2} {}_3F_2 \left( 1, \frac{m}{2} + 1, \frac{m}{2} + 1; \frac{m}{2} + \frac{3}{2}, \frac{m}{2} + 2; c^2 x^2 \right)}{f^2(m+1)(m+2)} + \frac{\sqrt{1-cx} (fx)^{m+1} \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2 x^2 \right) (a + \operatorname{barccosh}(cx))}{f(m+1) \sqrt{cx-1}} \right)}{(m+2) \sqrt{cx-1} \sqrt{cx+1}} \right)}{m+4}$$

$$\frac{(d-c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \frac{bcd^2 \sqrt{d-c^2 dx^2} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}}$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]),x]`

output `-((b*c*d^2*Sqrt[d - c^2*d*x^2]*((f*x)^(2 + m)/(f*(2 + m)) - (2*c^2*(f*x)^(4 + m))/(f^3*(4 + m)) + (c^4*(f*x)^(6 + m))/(f^5*(6 + m))))/(f*(6 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))/(f*(6 + m)) + (5*d*(-((b*c*d*Sqrt[d - c^2*d*x^2]*((f*x)^(2 + m)/(f*(2 + m)) - (c^2*(f*x)^(4 + m))/(f^3*(4 + m)))))/(f*(4 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(f*(4 + m)) + (3*d*(-((b*c*(f*x)^(2 + m)*Sqrt[d - c^2*d*x^2]))/(f^(2*(2 + m)^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(f*(2 + m)) - (Sqrt[d - c^2*d*x^2]*((f*x)^(1 + m)*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2]))/(f*(1 + m)*Sqrt[-1 + c*x]) + (b*c*(f*x)^(2 + m)*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2]))/(f^2*(1 + m)*(2 + m)))/((2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(4 + m))/(6 + m)`

### 3.151.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6364 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]`

### 3.151.4 Maple [F]

$$\int (fx)^m (-c^2 dx^2 + d)^{5/2} (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x)`

**3.151.5 Fricas [F]**

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((a*c^4*d^2*x^4 - 2*a*c^2*d^2*x^2 + a*d^2 + (b*c^4*d^2*x^4 - 2*b*c^2*d^2*x^2 + b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)*(f*x)^m, x)`

**3.151.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x)),x)`

output `Timed out`

**3.151.7 Maxima [F]**

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.151.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.151.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{5/2} (fx)^m dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(5/2)*(f*x)^m, x)`



### 3.152 $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.152.1 Optimal result	1280
3.152.2 Mathematica [A] (verified)	1281
3.152.3 Rubi [A] (verified)	1281
3.152.4 Maple [F]	1284
3.152.5 Fracas [F]	1285
3.152.6 Sympy [F(-1)]	1285
3.152.7 Maxima [F]	1285
3.152.8 Giac [F(-2)]	1286
3.152.9 Mupad [F(-1)]	1286

#### 3.152.1 Optimal result

Integrand size = 29, antiderivative size = 455

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$\begin{aligned} & -\frac{3bcd(fx)^{2+m}\sqrt{d - c^2 dx^2}}{f^2(2+m)^2(4+m)\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bcd(fx)^{2+m}\sqrt{d - c^2 dx^2}}{f^2(2+m)(4+m)\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{bc^3d(fx)^{4+m}\sqrt{d - c^2 dx^2}}{f^4(4+m)^2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{3d(fx)^{1+m}\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{f(8+6m+m^2)} \\ & + \frac{(fx)^{1+m}(d - c^2 dx^2)^{3/2}(a + \operatorname{barccosh}(cx))}{f(4+m)} \\ & + \frac{3d(fx)^{1+m}\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2 x^2\right)}{f(4+m)(2+3m+m^2)\sqrt{1-cx}\sqrt{1+cx}} \\ & - \frac{3bcd(fx)^{2+m}\sqrt{d - c^2 dx^2} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2 x^2\right)}{f^2(1+m)(2+m)^2(4+m)\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output  $(f*x)^{(1+m)}*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))/f/(4+m)+3*d*(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/f/(m^2+6*m+8)+3*d*(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))*\operatorname{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], c^2*x^2)*(-c^2*d*x^2+d)^{(1/2)}/f/(m^3+7*m^2+14*m+8)/(-c*x+1)^{(1/2)}/(c*x+1)^{(1/2)}-3*b*c*d*(f*x)^{(2+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)^2/(4+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b*c*d*(f*x)^{(2+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)/(4+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+b*c^3*d*(f*x)^{(4+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^4/(4+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3*b*c*d*(f*x)^{(2+m)}*\operatorname{hypergeom}([1, 1+1/2*m, 1+1/2*m], [2+1/2*m, 3/2+1/2*m], c^2*x^2)*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)^2/(m^2+5*m+4)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.152.2 Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 274, normalized size of antiderivative = 0.60

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$\frac{dx(fx)^m \sqrt{d - c^2 dx^2} \left( \frac{3bcx}{(2+m)^2} + bcx \left( \frac{1}{2+m} - \frac{c^2 x^2}{4+m} \right) - \frac{3\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{2+m} \right) + (-1 + cx)^{3/2}(1 + cx)^3}{(4 + m)\sqrt{\dots}}$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output 
$$-\left(\frac{d*x*(f*x)^m*\sqrt{d - c^2*d*x^2}*((3*b*c*x)/(2 + m)^2 + b*c*x*((2 + m)^(-1) - (c^2*x^2)/(4 + m)) - (3*\sqrt{-1 + c*x}*\sqrt{1 + c*x}*(a + b*ArcCosh[c*x])))/(2 + m) + (-1 + c*x)^(3/2)*(1 + c*x)^(3/2)*(a + b*ArcCosh[c*x]) + (3*\sqrt{1 - c^2*x^2}*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/((1 + m)*(2 + m)*\sqrt{-1 + c*x}*\sqrt{1 + c*x}) + (3*b*c*x*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/((1 + m)*(2 + m)^2)))/(4 + m)*\sqrt{-1 + c*x}*\sqrt{1 + c*x})$$

### 3.152.3 Rubi [A] (verified)

Time = 1.10 (sec) , antiderivative size = 396, normalized size of antiderivative = 0.87, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$ , Rules used = {6345, 25, 82, 244, 2009, 6341, 17, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{3/2} (fx)^m (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6345}$$

$$\frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m + 4} + \frac{bcd \sqrt{d - c^2 dx^2} \int -(fx)^{m+1} (1 - cx)(cx + 1) dx}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} +$$

$$\frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m + 4)}$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} - \frac{bcd \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1 - cx)(cx + 1) dx}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} + \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} \\
& \quad \downarrow \text{82} \\
& \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} - \frac{bcd \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1 - c^2 x^2) dx}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} + \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} \\
& \quad \downarrow \text{244} \\
& \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} - \frac{bcd \sqrt{d - c^2 dx^2} \int \left( (fx)^{m+1} - \frac{c^2 (fx)^{m+3}}{f^2} \right) dx}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} + \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} \\
& \quad \downarrow \text{2009} \\
& \frac{3d \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx}{m+4} + \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} - \\
& \quad \frac{bcd \sqrt{d - c^2 dx^2} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow \text{6341} \\
& 3d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx-1} \sqrt{cx+1}}}{(m+2) \sqrt{cx-1} \sqrt{cx+1}} - \frac{bc \sqrt{d - c^2 dx^2} \int (fx)^{m+1} dx}{f(m+2) \sqrt{cx-1} \sqrt{cx+1}} + \frac{\sqrt{d - c^2 dx^2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+2)} \right) + \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} - \frac{bcd \sqrt{d - c^2 dx^2} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow \text{17} \\
& 3d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx-1} \sqrt{cx+1}}}{(m+2) \sqrt{cx-1} \sqrt{cx+1}} + \frac{\sqrt{d - c^2 dx^2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \sqrt{d - c^2 dx^2} (fx)^{m+2}}{f^2(m+2)^2 \sqrt{cx-1} \sqrt{cx+1}} \right) + \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} - \frac{bcd \sqrt{d - c^2 dx^2} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m+4) \sqrt{cx-1} \sqrt{cx+1}} \\
& \quad \downarrow \text{6364}
\end{aligned}$$

---

3.152.  $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

$$3d \left( \frac{\sqrt{d-c^2dx^2} \left( \frac{bc(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2}+1, \frac{m}{2}+1; \frac{m}{2}+\frac{3}{2}, \frac{m}{2}+2; c^2x^2\right)}{f^{2(m+1)(m+2)}} + \frac{\sqrt{1-cx}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2x^2\right)}{f^{(m+1)\sqrt{cx-1}}} \right)}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+4)} - \frac{bcd\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}}{f^{(m+2)}} - \frac{c^2(fx)^{m+4}}{f^3(m+4)} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `-((b*c*d*Sqrt[d - c^2*d*x^2]*((f*x)^(2 + m)/(f*(2 + m)) - (c^2*(f*x)^(4 + m))/(f^3*(4 + m)))/(f*(4 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x]))/(f*(4 + m)) + (3*d*(-((b*c*(f*x)^(2 + m)*Sqrt[d - c^2*d*x^2])/(f^2*(2 + m)^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(f*(2 + m)) - (Sqrt[d - c^2*d*x^2]*(((f*x)^(1 + m)*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[-1 + c*x]) + (b*c*(f*x)^(2 + m)*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m)))/((2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/(4 + m)`

### 3.152.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_ + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_ + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6364 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_))/(Sqrt[(d1_ + (e1_.)*(x_)]*Sqrt[(d2_ + (e2_.)*(x_))], x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]`

### 3.152.4 Maple [F]

$$\int (fx)^m (-c^2 dx^2 + d)^{3/2} (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x)`

---

3.152.  $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx)) dx$

**3.152.5 Fricas [F]**

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-(a*c^2*d*x^2 - a*d + (b*c^2*d*x^2 - b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)*(f*x)^m, x)`

**3.152.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x)),x)`

output `Timed out`

**3.152.7 Maxima [F]**

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.152.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.152.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d - c^2 dx^2)^{3/2} (fx)^m dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(3/2)*(f*x)^m, x)`

### 3.153 $\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$

3.153.1 Optimal result . . . . .	1287
3.153.2 Mathematica [A] (verified) . . . . .	1288
3.153.3 Rubi [A] (verified) . . . . .	1288
3.153.4 Maple [F] . . . . .	1290
3.153.5 Fracas [F] . . . . .	1290
3.153.6 Sympy [F] . . . . .	1290
3.153.7 Maxima [F] . . . . .	1291
3.153.8 Giac [F(-2)] . . . . .	1291
3.153.9 Mupad [F(-1)] . . . . .	1292

#### 3.153.1 Optimal result

Integrand size = 29, antiderivative size = 278

$$\begin{aligned} & \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx \\ &= -\frac{bc(fx)^{2+m} \sqrt{d - c^2 dx^2}}{f^2(2+m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{(fx)^{1+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f(2+m)} \\ &+ \frac{(fx)^{1+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2 x^2\right)}{f(2+3m+m^2) \sqrt{1 - cx} \sqrt{1 + cx}} \\ &- \frac{bc(fx)^{2+m} \sqrt{d - c^2 dx^2} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2 x^2\right)}{f^2(1+m)(2+m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \end{aligned}$$

```
output (f*x)^(1+m)*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/f/(2+m)+(f*x)^(1+m)*(a
+b*arccosh(c*x))*hypergeom([1/2, 1/2+1/2*m],[3/2+1/2*m],c^2*x^2)*(-c^2*d*x
^2+d)^(1/2)/f/(m^2+3*m+2)/(-c*x+1)^(1/2)/(c*x+1)^(1/2)-b*c*(f*x)^(2+m)*(-c
^2*d*x^2+d)^(1/2)/f^2/(2+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*c*(f*x)^(2+m)*
hypergeom([1, 1+1/2*m, 1+1/2*m],[2+1/2*m, 3/2+1/2*m],c^2*x^2)*(-c^2*d*x^2+
d)^(1/2)/f^2/(1+m)/(2+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```



### 3.153.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.80

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{x(fx)^m \sqrt{d - c^2 dx^2} ((1 + m) (-bcx \sqrt{-1 + cx} \sqrt{1 + cx} + a(2 + m) (-1 + c^2 x^2) + b(2 + m) (-1 + c^2 x^2))}{(1 + m) (2 + m) (-1 + c^2 x^2)}$$

input `Integrate[(f*x)^m*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]),x]`

output `(x*(f*x)^m*Sqrt[d - c^2*d*x^2]*((1 + m)*(-b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + a*(2 + m)*(-1 + c^2*x^2) + b*(2 + m)*(-1 + c^2*x^2)*ArcCosh[c*x]) - (2 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2] - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/((1 + m)*(2 + m)^2*(-1 + c*x)*(1 + c*x))`

### 3.153.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.95, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$ , Rules used = {6341, 17, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d - c^2 dx^2} (fx)^m (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6341}$$

$$-\frac{\sqrt{d - c^2 dx^2} \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{(m + 2) \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{bc \sqrt{d - c^2 dx^2} \int (fx)^{m+1} dx}{f(m + 2) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{\sqrt{d - c^2 dx^2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m + 2)}$$

$$\downarrow \text{17}$$

$$\begin{aligned}
 & -\frac{\sqrt{d-c^2dx^2} \int \frac{(fx)^m(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{\sqrt{d-c^2dx^2}(fx)^{m+1}(a+\operatorname{barccosh}(cx))}{f(m+2)} \\
 & \quad - \frac{bc\sqrt{d-c^2dx^2}(fx)^{m+2}}{f^2(m+2)^2\sqrt{cx-1}\sqrt{cx+1}} \\
 & \quad \quad \quad \downarrow \text{6364} \\
 & -\frac{\sqrt{d-c^2dx^2} \left( \frac{bc(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2}+1, \frac{m}{2}+1; \frac{m}{2}+\frac{3}{2}, \frac{m}{2}+2; c^2x^2\right)}{f^2(m+1)(m+2)} + \frac{\sqrt{1-cx}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2x^2\right)(a+\operatorname{barccosh}(cx))}{f(m+1)\sqrt{cx-1}} \right)}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} \\
 & \quad - \frac{\sqrt{d-c^2dx^2}(fx)^{m+1}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{d-c^2dx^2}(fx)^{m+2}}{f^2(m+2)^2\sqrt{cx-1}\sqrt{cx+1}}
 \end{aligned}$$

input `Int[(f*x)^m*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]), x]`

output `-((b*c*(f*x)^(2 + m)*Sqrt[d - c^2*d*x^2])/(f^2*(2 + m)^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x]))/(f*(2 + m) - (Sqrt[d - c^2*d*x^2]*(((f*x)^(1 + m)*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[-1 + c*x]) + (b*c*(f*x)^(2 + m)*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m))))/( (2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.153.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6364 `Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_))/(Sqrt[(d1_) + (e1_.)*(x_.)]*Sqrt[(d2_) + (e2_.)*(x_.)]), x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]`

### 3.153.4 Maple [F]

$$\int (fx)^m \sqrt{-c^2 dx^2 + d} (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x)),x)`

### 3.153.5 Fracas [F]

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

### 3.153.6 Sympy [F]

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int (fx)^m \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx)) dx$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(1/2)*(a+b*acosh(c*x)),x)`

output `Integral((f*x)**m*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x)), x)`

### 3.153.7 Maxima [F]

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

### 3.153.8 Giac [F(-2)]

Exception generated.

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.153.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) \sqrt{d - c^2 dx^2} (fx)^m dx$$

input `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2)*(f*x)^m,x)`output `int((a + b*acosh(c*x))*(d - c^2*d*x^2)^(1/2)*(f*x)^m, x)`

$$3.154 \quad \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{\sqrt{d-c^2dx^2}} dx$$

3.154.1 Optimal result . . . . .	1293
3.154.2 Mathematica [A] (verified) . . . . .	1293
3.154.3 Rubi [A] (verified) . . . . .	1294
3.154.4 Maple [F] . . . . .	1295
3.154.5 Fricas [F] . . . . .	1295
3.154.6 Sympy [F] . . . . .	1296
3.154.7 Maxima [F] . . . . .	1296
3.154.8 Giac [F] . . . . .	1296
3.154.9 Mupad [F(-1)] . . . . .	1297

### 3.154.1 Optimal result

Integrand size = 29, antiderivative size = 176

$$\begin{aligned} & \int \frac{(fx)^m(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx \\ &= \frac{(fx)^{1+m}\sqrt{1 - c^2x^2}(a + \operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{f(1+m)\sqrt{d - c^2dx^2}} \\ &+ \frac{bc(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2x^2\right)}{f^2(1+m)(2+m)\sqrt{d - c^2dx^2}} \end{aligned}$$

```
output b*c*(f*x)^(2+m)*hypergeom([1, 1+1/2*m, 1+1/2*m],[2+1/2*m, 3/2+1/2*m],c^2*x
^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/f^2/(1+m)/(2+m)/(-c^2*d*x^2+d)^(1/2)+(f*x)
^(1+m)*(a+b*arccosh(c*x))*hypergeom([1/2, 1/2+1/2*m],[3/2+1/2*m],c^2*x^2)*
(-c^2*x^2+1)^(1/2)/f/(1+m)/(-c^2*d*x^2+d)^(1/2)
```

### 3.154.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.84

$$\begin{aligned} & \int \frac{(fx)^m(a + \operatorname{arccosh}(cx))}{\sqrt{d - c^2dx^2}} dx \\ &= \frac{x(fx)^m \left( (2+m)\sqrt{1 - c^2x^2}(a + \operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right) + bcx\sqrt{-1 + cx}\sqrt{1 + cx} \right)}{(1+m)(2+m)\sqrt{d - c^2dx^2}} \end{aligned}$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2],x]`

output `(x*(f*x)^m*((2 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2] + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2)]/((1 + m)*(2 + m)*Sqrt[d - c^2*d*x^2])`

### 3.154.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.034$ , Rules used = {6363}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \text{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6363

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2} + 1, \frac{m}{2} + 1; \frac{m}{2} + \frac{3}{2}, \frac{m}{2} + 2; c^2 x^2\right) + \sqrt{1 - c^2 x^2} (fx)^{m+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2 x^2\right) (a + \text{barccosh}(cx))}{f(m+1)\sqrt{d - c^2 dx^2}}$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/Sqrt[d - c^2*d*x^2],x]`

output `((f*x)^(1 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[d - c^2*d*x^2]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2)]/(f^2*(1 + m)*(2 + m)*Sqrt[d - c^2*d*x^2])`

## 3.154.3.1 Defintions of rubi rules used

```
rule 6363 Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && !IntegerQ[m]
```

## 3.154.4 Maple [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{-c^2 d x^2 + d}} dx$$

```
input int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x)
```

```
output int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x)
```

## 3.154.5 Fricas [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{\sqrt{-c^2 dx^2 + d}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")
```

```
output integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)*(f*x)^m/(c^2*d*x^2 - d), x)
```



**3.154.6 Sympy [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(1/2), x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.154.7 Maxima [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2), x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/sqrt(-c^2*d*x^2 + d), x)`

**3.154.8 Giac [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(1/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/sqrt(-c^2*d*x^2 + d), x)`

**3.154.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{\sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^(1/2), x)`

output `int((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^(1/2), x)`

**3.155** 
$$\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d-c^2dx^2)^{3/2}} dx$$

3.155.1 Optimal result . . . . . 1298  
 3.155.2 Mathematica [A] (verified) . . . . . 1299  
 3.155.3 Rubi [A] (verified) . . . . . 1299  
 3.155.4 Maple [F] . . . . . 1301  
 3.155.5 Fracas [F] . . . . . 1302  
 3.155.6 Sympy [F] . . . . . 1302  
 3.155.7 Maxima [F] . . . . . 1302  
 3.155.8 Giac [F] . . . . . 1303  
 3.155.9 Mupad [F(-1)] . . . . . 1303

**3.155.1 Optimal result**

Integrand size = 29, antiderivative size = 300

$$\int \frac{(fx)^m(a + \operatorname{arccosh}(cx))}{(d - c^2dx^2)^{3/2}} dx = \frac{(fx)^{1+m}(a + \operatorname{arccosh}(cx))}{df\sqrt{d - c^2dx^2}} - \frac{m(fx)^{1+m}\sqrt{1 - c^2x^2}(a + \operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{df(1+m)\sqrt{d - c^2dx^2}} + \frac{bc(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{df^2(2+m)\sqrt{d - c^2dx^2}} - \frac{bcm(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2x^2\right)}{df^2(1+m)(2+m)\sqrt{d - c^2dx^2}}$$

output

```
(f*x)^(1+m)*(a+b*arccosh(c*x))/d/f/(-c^2*d*x^2+d)^(1/2)+b*c*(f*x)^(2+m)*hy
pergeom([1, 1+1/2*m], [2+1/2*m], c^2*x^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/f^2/
(2+m)/(-c^2*d*x^2+d)^(1/2)-b*c*m*(f*x)^(2+m)*hypergeom([1, 1+1/2*m, 1+1/2*
m], [2+1/2*m, 3/2+1/2*m], c^2*x^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/f^2/(1+m)/(
2+m)/(-c^2*d*x^2+d)^(1/2)-m*(f*x)^(1+m)*(a+b*arccosh(c*x))*hypergeom([1/2,
1/2+1/2*m], [3/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/d/f/(1+m)/(-c^2*d*x^2+
d)^(1/2)
```

### 3.155.2 Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.72

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \frac{x(fx)^m (-m(2+m)\sqrt{1-c^2x^2}(a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}(\frac{1}{2}, (1+m)/2, (3+m)/2, c^2x^2)) + (1+m)((2+m)(a + \operatorname{barccosh}(cx)) + b*c*x*\sqrt{-1+cx}*\sqrt{1+cx}*\operatorname{Hypergeometric2F1}(1, 1+m/2, 2+m/2, c^2x^2)) - b*c*m*x*\sqrt{-1+cx}*\sqrt{1+cx}*\operatorname{HypergeometricPFQ}(\{1, 1+m/2, 1+m/2\}, \{3/2+m/2, 2+m/2\}, c^2x^2))}{d*(1+m)*(2+m)*\sqrt{d-c^2dx^2}}$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2),x]`

output `(x*(f*x)^m*(-(m*(2+m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1+m)/2, (3+m)/2, c^2*x^2]) + (1+m)*((2+m)*(a + b*ArcCosh[c*x]) + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[1, 1+m/2, 2+m/2, c^2*x^2]) - b*c*m*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1+m/2, 1+m/2}, {3/2+m/2, 2+m/2}, c^2*x^2]))/(d*(1+m)*(2+m)*Sqrt[d - c^2*d*x^2])`

### 3.155.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.172$ , Rules used = {6351, 25, 82, 278, 6363}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx \\ & \quad \downarrow \text{6351} \\ & -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{d} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{df\sqrt{d-c^2dx^2}} + \\ & \quad \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{df\sqrt{d-c^2dx^2}} \\ & \quad \downarrow \text{25} \\ & -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{df\sqrt{d-c^2dx^2}} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{df\sqrt{d-c^2dx^2}} \\ & \quad \downarrow \text{82} \end{aligned}$$

---

3.155.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{d} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{(fx)^{m+1}}{1 - c^2 x^2} dx}{df\sqrt{d - c^2 dx^2}} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{df\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow \text{278} \\
& -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{d} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{df\sqrt{d - c^2 dx^2}} + \\
& \frac{bc\sqrt{cx - 1}\sqrt{cx + 1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{df^2(m+2)\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow \text{6363} \\
& \frac{m \left( \frac{bc\sqrt{cx - 1}\sqrt{cx + 1}(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2} + 1, \frac{m}{2} + 1; \frac{m}{2} + \frac{3}{2}, \frac{m}{2} + 2; c^2 x^2\right)}{f^2(m+1)(m+2)\sqrt{d - c^2 dx^2}} + \frac{\sqrt{1 - c^2 x^2}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2 x^2\right)(a + \operatorname{barccosh}(cx))}{f(m+1)\sqrt{d - c^2 dx^2}} \right)}{d} \\
& \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{df\sqrt{d - c^2 dx^2}} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{df^2(m+2)\sqrt{d - c^2 dx^2}}
\end{aligned}$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(3/2), x]`

output `((f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(d*f*Sqrt[d - c^2*d*x^2]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, c^2*x^2])/(d*f^2*(2 + m)*Sqrt[d - c^2*d*x^2]) - (m*(((f*x)^(1 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[d - c^2*d*x^2]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m)*Sqrt[d - c^2*d*x^2]))/d`

### 3.155.3.1 Defintions of rubi rules used

rule 25 `Int[-(F*x_), x_Symbol] := Simp[Identity[-1] Int[F*x, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

rule 6363 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)]/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && !IntegerQ[m]`

### 3.155.4 Maple [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x)`

**3.155.5 Fricas [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)*(f*x)^m/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.155.6 Sympy [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.155.7 Maxima [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(-c^2*d*x^2 + d)^(3/2), x)`

**3.155.8 Giac [F]**

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(-c^2*d*x^2 + d)^(3/2), x)`

**3.155.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{(d - c^2 dx^2)^{3/2}} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^(3/2),x)`

output `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^(3/2), x)`



**3.156**  $\int \frac{(fx)^m(a+b\text{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

3.156.1 Optimal result . . . . . 1304  
 3.156.2 Mathematica [A] (verified) . . . . . 1305  
 3.156.3 Rubi [A] (verified) . . . . . 1306  
 3.156.4 Maple [F] . . . . . 1309  
 3.156.5 Fricas [F] . . . . . 1309  
 3.156.6 Sympy [F(-1)] . . . . . 1310  
 3.156.7 Maxima [F] . . . . . 1310  
 3.156.8 Giac [F] . . . . . 1310  
 3.156.9 Mupad [F(-1)] . . . . . 1311

**3.156.1 Optimal result**

Integrand size = 29, antiderivative size = 450

$$\int \frac{(fx)^m(a + \text{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{(fx)^{1+m}(a + \text{arccosh}(cx))}{3df(d - c^2dx^2)^{3/2}} + \frac{(2 - m)(fx)^{1+m}(a + \text{arccosh}(cx))}{3d^2f\sqrt{d - c^2dx^2}} - \frac{(2 - m)m(fx)^{1+m}\sqrt{1 - c^2x^2}(a + \text{arccosh}(cx)) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{3d^2f(1 + m)\sqrt{d - c^2dx^2}} + \frac{bc(2 - m)(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} \text{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{3d^2f^2(2 + m)\sqrt{d - c^2dx^2}} + \frac{bc(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} \text{Hypergeometric2F1}\left(2, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{3d^2f^2(2 + m)\sqrt{d - c^2dx^2}} - \frac{bc(2 - m)m(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2x^2\right)}{3d^2f^2(1 + m)(2 + m)\sqrt{d - c^2dx^2}}$$

---

3.156.  $\int \frac{(fx)^m(a+b\text{arccosh}(cx))}{(d-c^2dx^2)^{5/2}} dx$

output  $\frac{1}{3}(fx)^{(1+m)}(a+b\operatorname{arccosh}(cx))/d/f/(-c^2dx^2+d)^{(3/2)}+1/3(2-m)(fx)^{(1+m)}(a+b\operatorname{arccosh}(cx))/d^2/f/(-c^2dx^2+d)^{(1/2)}+1/3b*c*(2-m)(fx)^{(2+m)}\operatorname{hypergeom}([1, 1+1/2*m], [2+1/2*m], c^2*x^2)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/f^2/(2+m)/(-c^2dx^2+d)^{(1/2)}+1/3b*c*(fx)^{(2+m)}\operatorname{hypergeom}([2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/f^2/(2+m)/(-c^2dx^2+d)^{(1/2)}-1/3b*c*(2-m)*m*(fx)^{(2+m)}\operatorname{hypergeom}([1, 1+1/2*m, 1+1/2*m], [2+1/2*m, 3/2+1/2*m], c^2*x^2)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/f^2/(1+m)/(2+m)/(-c^2dx^2+d)^{(1/2)}-1/3(2-m)*m*(fx)^{(1+m)}(a+b\operatorname{arccosh}(cx))*\operatorname{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^{(1/2)}/d^2/f/(1+m)/(-c^2dx^2+d)^{(1/2)}$

### 3.156.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 319, normalized size of antiderivative = 0.71

$$\int \frac{(fx)^m(a + b\operatorname{arccosh}(cx))}{(d - c^2dx^2)^{5/2}} dx = \frac{x(fx)^m\sqrt{-1 + cx}\sqrt{1 + cx}\left(-\frac{a+b\operatorname{arccosh}(cx)}{(-1+cx)^{3/2}(1+cx)^{3/2}} + \frac{bcx \operatorname{Hypergeometric2F1}(2,1+\frac{m}{2}}{2+m}\right)}{d - c^2dx^2}$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2),x]`

output  $(x*(f*x)^m*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(-((a + b*\operatorname{ArcCosh}[c*x])/((-1 + c*x)^{(3/2)}*(1 + c*x)^{(3/2)}))) + (b*c*x*\operatorname{Hypergeometric2F1}[2, 1 + m/2, 2 + m/2, c^2*x^2])/(2 + m) + ((-2 + m)*(m*(2 + m)*\operatorname{Sqrt}[1 - c^2*x^2]*(a + b*\operatorname{ArcCosh}[c*x])*\operatorname{Hypergeometric2F1}[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2] - (1 + m)*((2 + m)*(a + b*\operatorname{ArcCosh}[c*x]) + b*c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*\operatorname{Hypergeometric2F1}[1, 1 + m/2, 2 + m/2, c^2*x^2]) + b*c*m*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*\operatorname{HypergeometricPFQ}[\{1, 1 + m/2, 1 + m/2\}, \{3/2 + m/2, 2 + m/2\}, c^2*x^2]))/((1 + m)*(2 + m)*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]))/(3*d^2*\operatorname{Sqrt}[d - c^2*d*x^2])$

**3.156.3 Rubi [A] (verified)**

Time = 0.97 (sec) , antiderivative size = 430, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$ , Rules used = {6351, 82, 278, 6351, 25, 82, 278, 6363}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6351} \\
 & \frac{(2-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx}{3d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{(1-cx)^2(cx+1)^2} dx}{3d^2 f \sqrt{d - c^2 dx^2}} + \\
 & \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3df (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{82} \\
 & \frac{(2-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx}{3d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{(1-c^2 x^2)^2} dx}{3d^2 f \sqrt{d - c^2 dx^2}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3df (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{278} \\
 & \frac{(2-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{3/2}} dx}{3d} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3df (d - c^2 dx^2)^{3/2}} + \\
 & \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{3d^2 f^2 (m+2) \sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{6351} \\
 & \frac{(2-m) \left( -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{d - c^2 dx^2}} dx}{d} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{df \sqrt{d - c^2 dx^2}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{df \sqrt{d - c^2 dx^2}} \right)}{3d} + \\
 & \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3df (d - c^2 dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{3d^2 f^2 (m+2) \sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

---

3.156.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx$

$$\frac{(2-m) \left( -\frac{m \int \frac{(fx)^m (a+b \operatorname{arccosh}(cx)) dx}{\sqrt{d-c^2 dx^2}}}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{df\sqrt{d-c^2 dx^2}} + \frac{(fx)^{m+1}(a+b \operatorname{arccosh}(cx))}{df\sqrt{d-c^2 dx^2}} \right)}{(fx)^{m+1}(a+b \operatorname{arccosh}(cx)) + \frac{3d}{3df(d-c^2 dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{3d^2 f^2 (m+2)\sqrt{d-c^2 dx^2}}}$$

↓ 82

$$\frac{(2-m) \left( -\frac{m \int \frac{(fx)^m (a+b \operatorname{arccosh}(cx)) dx}{\sqrt{d-c^2 dx^2}}}{d} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{1-c^2 x^2} dx}{df\sqrt{d-c^2 dx^2}} + \frac{(fx)^{m+1}(a+b \operatorname{arccosh}(cx))}{df\sqrt{d-c^2 dx^2}} \right)}{(fx)^{m+1}(a+b \operatorname{arccosh}(cx)) + \frac{3d}{3df(d-c^2 dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{3d^2 f^2 (m+2)\sqrt{d-c^2 dx^2}}}$$

↓ 278

$$\frac{(2-m) \left( -\frac{m \int \frac{(fx)^m (a+b \operatorname{arccosh}(cx)) dx}{\sqrt{d-c^2 dx^2}}}{d} + \frac{(fx)^{m+1}(a+b \operatorname{arccosh}(cx))}{df\sqrt{d-c^2 dx^2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{df^2 (m+2)\sqrt{d-c^2 dx^2}} \right)}{(fx)^{m+1}(a+b \operatorname{arccosh}(cx)) + \frac{3d}{3df(d-c^2 dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{3d^2 f^2 (m+2)\sqrt{d-c^2 dx^2}}}$$

↓ 6363

$$\frac{(2-m) \left( -\frac{m \left( \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2}+1, \frac{m}{2}+1; \frac{m}{2}+\frac{3}{2}, \frac{m}{2}+2; c^2 x^2\right)}{f^2 (m+1)(m+2)\sqrt{d-c^2 dx^2}} + \frac{\sqrt{1-c^2 x^2}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2 x^2\right)(a+b \operatorname{arccosh}(cx))}{f(m+1)\sqrt{d-c^2 dx^2}} \right)}{d}}{(fx)^{m+1}(a+b \operatorname{arccosh}(cx)) + \frac{3d}{3df(d-c^2 dx^2)^{3/2}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{3d^2 f^2 (m+2)\sqrt{d-c^2 dx^2}}}$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d - c^2*d*x^2)^(5/2), x]`

```
output ((f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(3*d*f*(d - c^2*d*x^2)^(3/2)) + (b*c*
(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[2, (2 + m)/2,
(4 + m)/2, c^2*x^2])/(3*d^2*f^2*(2 + m)*Sqrt[d - c^2*d*x^2]) + ((2 - m)*
((f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(d*f*Sqrt[d - c^2*d*x^2]) + (b*c*(f*x)
)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[1, (2 + m)/2, (4
+ m)/2, c^2*x^2])/(d*f^2*(2 + m)*Sqrt[d - c^2*d*x^2]) - (m*(((f*x)^(1 + m)
)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2,
(3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[d - c^2*d*x^2]) + (b*c*(f*x)^(2 + m)*
Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2
+ m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m)*Sqrt[d - c^2*d*x^2]))/d
)/(3*d)
```

### 3.156.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 82 Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))
)^(p_), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d,
e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]
```

```
rule 278 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

```
rule 6351 Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d_) + (e_
) *(x_)^2)^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1
)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[
b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[
(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])
^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] &
& GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] ||
EqQ[n, 1])
```

rule 6363 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && !IntegerQ[m]`

### 3.156.4 Maple [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x)`

### 3.156.5 Fracas [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d - c^2 dx^2)^{\frac{5}{2}}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{(-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)*(f*x)^m/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

**3.156.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(-c**2*d*x**2+d)**(5/2), x)`

output `Timed out`

**3.156.7 Maxima [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2), x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(-c^2*d*x^2 + d)^(5/2), x)`

**3.156.8 Giac [F]**

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(-c^2*d*x^2+d)^(5/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(-c^2*d*x^2 + d)^(5/2), x)`

**3.156.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{(d - c^2 dx^2)^{5/2}} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^(5/2), x)`

output `int(((a + b*acosh(c*x))*(f*x)^m)/(d - c^2*d*x^2)^(5/2), x)`



### 3.157 $\int (fx)^m (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

3.157.1 Optimal result . . . . .	1312
3.157.2 Mathematica [A] (verified) . . . . .	1313
3.157.3 Rubi [A] (verified) . . . . .	1314
3.157.4 Maple [F] . . . . .	1319
3.157.5 Fricas [F] . . . . .	1320
3.157.6 Sympy [F(-1)] . . . . .	1320
3.157.7 Maxima [F] . . . . .	1320
3.157.8 Giac [F] . . . . .	1321
3.157.9 Mupad [F(-1)] . . . . .	1321

#### 3.157.1 Optimal result

Integrand size = 35, antiderivative size = 817

$$\int (fx)^m (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + \operatorname{barccosh}(cx)) dx =$$

$$\frac{bcd1^2 d2^2 (fx)^{2+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x}}{f^2 (2+m) (6+m) \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{15bcd1^2 d2^2 (fx)^{2+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x}}{f^2 (2+m)^2 (4+m) (6+m) \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$- \frac{5bcd1^2 d2^2 (fx)^{2+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x}}{f^2 (2+m) (4+m) (6+m) \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{5bc^3 d1^2 d2^2 (fx)^{4+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x}}{f^4 (4+m)^2 (6+m) \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{2bc^3 d1^2 d2^2 (fx)^{4+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x}}{f^4 (4+m) (6+m) \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{bc^5 d1^2 d2^2 (fx)^{6+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x}}{f^6 (6+m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{15d1^2 d2^2 (fx)^{1+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx))}{f (6+m) (8 + 6m + m^2)}$$

$$+ \frac{5d1d2 (fx)^{1+m} (d1 + cd1x)^{3/2} (d2 - cd2x)^{3/2} (a + \operatorname{barccosh}(cx))}{f (4+m) (6+m)}$$

$$+ \frac{(fx)^{1+m} (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + \operatorname{barccosh}(cx))}{f (6+m)}$$

$$+ \frac{15d1^2 d2^2 (fx)^{1+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2 x^2\right)}{f (4+m) (6+m) (2 + 3m + m^2) \sqrt{1 - cx} \sqrt{1 + cx}}$$

$$- \frac{15bcd1^2 d2^2 (fx)^{2+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2 x^2\right)}{f^2 (1+m) (2+m)^2 (4+m) (6+m) \sqrt{-1 + cx} \sqrt{1 + cx}}$$

---

3.157.  $\int (fx)^m (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + \operatorname{barccosh}(cx)) dx$

```
output 5*d1*d2*(f*x)^(1+m)*(c*d1*x+d1)^(3/2)*(-c*d2*x+d2)^(3/2)*(a+b*arccosh(c*x)
)/f/(4+m)/(6+m)+(f*x)^(1+m)*(c*d1*x+d1)^(5/2)*(-c*d2*x+d2)^(5/2)*(a+b*arcc
osh(c*x))/f/(6+m)+15*d1^2*d2^2*(f*x)^(1+m)*(a+b*arccosh(c*x))*(c*d1*x+d1)^(
1/2)*(-c*d2*x+d2)^(1/2)/f/(6+m)/(m^2+6*m+8)+15*d1^2*d2^2*(f*x)^(1+m)*(a+b
*arccosh(c*x))*hypergeom([1/2, 1/2+1/2*m],[3/2+1/2*m],c^2*x^2)*(c*d1*x+d1)
^(1/2)*(-c*d2*x+d2)^(1/2)/f/(6+m)/(m^3+7*m^2+14*m+8)/(-c*x+1)^(1/2)/(c*x+1
)^(1/2)-b*c*d1^2*d2^2*(f*x)^(2+m)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f^2
/(2+m)/(6+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-15*b*c*d1^2*d2^2*(f*x)^(2+m)*(c*d
1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f^2/(2+m)^2/(4+m)/(6+m)/(c*x-1)^(1/2)/(c*
x+1)^(1/2)-5*b*c*d1^2*d2^2*(f*x)^(2+m)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2
)/f^2/(6+m)/(m^2+6*m+8)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5*b*c^3*d1^2*d2^2*(f*x
)^(4+m)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f^4/(4+m)^2/(6+m)/(c*x-1)^(1/
2)/(c*x+1)^(1/2)+2*b*c^3*d1^2*d2^2*(f*x)^(4+m)*(c*d1*x+d1)^(1/2)*(-c*d2*x+
d2)^(1/2)/f^4/(4+m)/(6+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*c^5*d1^2*d2^2*(f*x
)^(6+m)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f^6/(6+m)^2/(c*x-1)^(1/2)/(c*
x+1)^(1/2)-15*b*c*d1^2*d2^2*(f*x)^(2+m)*hypergeom([1, 1+1/2*m, 1+1/2*m],[2
+1/2*m, 3/2+1/2*m],c^2*x^2)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f^2/(2+m)
^2/(6+m)/(m^2+5*m+4)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.157.2 Mathematica [A] (verified)

Time = 1.65 (sec) , antiderivative size = 387, normalized size of antiderivative = 0.47

$$\int (fx)^m (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + barccosh(cx)) dx = \frac{d1^2 d2^2 x (fx)^m \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} \left( -\frac{bcx \left( \frac{1}{2+m} - \frac{2c^2 x^2}{4+m} + \frac{c^4 x^4}{6+m} \right)}{\sqrt{-1+cx} \sqrt{1+cx}} + (-1 + c \right.}{-cd2x)^{5/2} (a + barccosh(cx)) dx =$$

```
input Integrate[(f*x)^m*(d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2)*(a + b*ArcCosh[c
*x]),x]
```

output  $(d1^2*d2^2*x*(f*x)^m*\text{Sqrt}[d1 + c*d1*x]*\text{Sqrt}[d2 - c*d2*x]*(-((b*c*x*((2 + m)^{-1} - (2*c^2*x^2)/(4 + m) + (c^4*x^4)/(6 + m)))/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])) + (-1 + c^2*x^2)^2*(a + b*\text{ArcCosh}[c*x]) + (5*((b*c*x*(-(2 + m)^{-1} + (c^2*x^2)/(4 + m)))/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) - (-1 + c*x)*(1 + c*x)*(a + b*\text{ArcCosh}[c*x]) + (3*((1 + m)*(-b*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) + a*(2 + m)*(-1 + c^2*x^2) + b*(2 + m)*(-1 + c^2*x^2)*\text{ArcCosh}[c*x]) - (2 + m)*\text{Sqrt}[1 - c^2*x^2]*(a + b*\text{ArcCosh}[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2] - b*c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])))/((1 + m)*(2 + m)^2*(-1 + c*x)*(1 + c*x)))/((4 + m)))/(6 + m)$

### 3.157.3 Rubi [A] (verified)

Time = 2.60 (sec) , antiderivative size = 596, normalized size of antiderivative = 0.73, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.343$ , Rules used = {6346, 82, 244, 2009, 6346, 25, 82, 244, 2009, 6342, 17, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (cd1x + d1)^{5/2}(d2 - cd2x)^{5/2}(fx)^m(a + \text{barccosh}(cx)) dx$$

$$\downarrow \text{6346}$$

$$\frac{5d1d2 \int (fx)^m (cx d1 + d1)^{3/2} (d2 - cd2x)^{3/2} (a + \text{barccosh}(cx)) dx}{m + 6} - \frac{bcd1^2 d2^2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int (fx)^{m+1} (1 - cx)^2 (cx + 1)^2 dx}{f(m + 6) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{(cd1x + d1)^{5/2} (d2 - cd2x)^{5/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 6)}$$

$$\downarrow \text{82}$$

$$\frac{5d1d2 \int (fx)^m (cx d1 + d1)^{3/2} (d2 - cd2x)^{3/2} (a + \text{barccosh}(cx)) dx}{m + 6} - \frac{bcd1^2 d2^2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int (fx)^{m+1} (1 - c^2 x^2)^2 dx}{f(m + 6) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{(cd1x + d1)^{5/2} (d2 - cd2x)^{5/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 6)}$$

$$\downarrow \text{244}$$

---

3.157.  $\int (fx)^m (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + \text{barccosh}(cx)) dx$

$$\begin{aligned}
& \frac{5d_1d_2 \int (fx)^m (cxd_1 + d_1)^{3/2} (d_2 - cd_2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx}{m+6} \\
& \frac{bcd_1^2 d_2^2 \sqrt{cd_1x + d_1} \sqrt{d_2 - cd_2x} \int \left( (fx)^{m+1} - \frac{2c^2(fx)^{m+3}}{f^2} + \frac{c^4(fx)^{m+5}}{f^4} \right) dx}{\frac{f(m+6)\sqrt{cx-1}\sqrt{cx+1}}{(cd_1x + d_1)^{5/2} (d_2 - cd_2x)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}} + \\
& \qquad \qquad \qquad \downarrow \text{2009} \\
& \frac{5d_1d_2 \int (fx)^m (cxd_1 + d_1)^{3/2} (d_2 - cd_2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx}{m+6} + \\
& \frac{(cd_1x + d_1)^{5/2} (d_2 - cd_2x)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \\
& \frac{bcd_1^2 d_2^2 \sqrt{cd_1x + d_1} \sqrt{d_2 - cd_2x} \left( \frac{c^4(fx)^{m+6}}{f^5(m+6)} - \frac{2c^2(fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}} \\
& \qquad \qquad \qquad \downarrow \text{6346} \\
& \frac{5d_1d_2 \left( \frac{3d_1d_2 \int (fx)^m \sqrt{cxd_1+d_1} \sqrt{d_2-cd_2x} (a+\operatorname{barccosh}(cx)) dx}{m+4} + \frac{bcd_1d_2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \int -(fx)^{m+1}(1-cx)(cx+1) dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} \right)}{m+6} \\
& \frac{(cd_1x + d_1)^{5/2} (d_2 - cd_2x)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \\
& \frac{bcd_1^2 d_2^2 \sqrt{cd_1x + d_1} \sqrt{d_2 - cd_2x} \left( \frac{c^4(fx)^{m+6}}{f^5(m+6)} - \frac{2c^2(fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}} \\
& \qquad \qquad \qquad \downarrow \text{25} \\
& \frac{5d_1d_2 \left( \frac{3d_1d_2 \int (fx)^m \sqrt{cxd_1+d_1} \sqrt{d_2-cd_2x} (a+\operatorname{barccosh}(cx)) dx}{m+4} - \frac{bcd_1d_2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \int (fx)^{m+1}(1-cx)(cx+1) dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} \right)}{m+6} \\
& \frac{(cd_1x + d_1)^{5/2} (d_2 - cd_2x)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+6)} - \\
& \frac{bcd_1^2 d_2^2 \sqrt{cd_1x + d_1} \sqrt{d_2 - cd_2x} \left( \frac{c^4(fx)^{m+6}}{f^5(m+6)} - \frac{2c^2(fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}} \\
& \qquad \qquad \qquad \downarrow \text{82}
\end{aligned}$$

$$5d_1d_2 \left( \frac{3d_1d_2 \int (fx)^m \sqrt{cx}d_1+d_1 \sqrt{d_2-cd_2x} (a+\text{barccosh}(cx)) dx}{m+4} - \frac{bcd_1d_2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \int (fx)^{m+1} (1-c^2x^2) dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \frac{(cd_1x+d_1)^{5/2} (d_2-cd_2x)^{5/2} (fx)^{m+1} (a+\text{barccosh}(cx))}{f(m+6)} - \frac{bcd_1^2 d_2^2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 244

$$5d_1d_2 \left( \frac{3d_1d_2 \int (fx)^m \sqrt{cx}d_1+d_1 \sqrt{d_2-cd_2x} (a+\text{barccosh}(cx)) dx}{m+4} - \frac{bcd_1d_2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \int \left( (fx)^{m+1} - \frac{c^2 (fx)^{m+3}}{f^2} \right) dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \frac{(cd_1x+d_1)^{5/2} (d_2-cd_2x)^{5/2} (fx)^{m+1} (a+\text{barccosh}(cx))}{f(m+6)} - \frac{bcd_1^2 d_2^2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 2009

$$5d_1d_2 \left( \frac{3d_1d_2 \int (fx)^m \sqrt{cx}d_1+d_1 \sqrt{d_2-cd_2x} (a+\text{barccosh}(cx)) dx}{m+4} + \frac{(cd_1x+d_1)^{3/2} (d_2-cd_2x)^{3/2} (fx)^{m+1} (a+\text{barccosh}(cx))}{f(m+4)} - \frac{bcd_1^2 d_2^2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}} + \frac{(cd_1x+d_1)^{5/2} (d_2-cd_2x)^{5/2} (fx)^{m+1} (a+\text{barccosh}(cx))}{f(m+6)} \right)$$

↓ 6342

$$5d_1d_2 \left( \frac{3d_1d_2 \left( -\frac{\sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \int (fx)^m (a+\text{barccosh}(cx)) dx}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \int (fx)^{m+1} dx}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{\sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \int (fx)^{m+2} dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} \right)}{m+4} + \frac{(cd_1x+d_1)^{5/2} (d_2-cd_2x)^{5/2} (fx)^{m+1} (a+\text{barccosh}(cx))}{f(m+6)} - \frac{bcd_1^2 d_2^2 \sqrt{cd_1x+d_1} \sqrt{d_2-cd_2x} \left( \frac{c^4 (fx)^{m+6}}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}} \right)$$

3.157.  $\int (fx)^m (d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2} (a + \text{barccosh}(cx)) dx$

↓ 17

$$5d_1d_2 \left( \frac{3d_1d_2 \left( -\frac{\sqrt{cd_1x+d_1}\sqrt{d_2-cd_2x} \int \frac{(fx)^m (a+b\operatorname{arccosh}(cx)) dx}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{\sqrt{cd_1x+d_1}\sqrt{d_2-cd_2x} (fx)^{m+1} (a+b\operatorname{arccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{cd_1x+d_1}}{f^2(m+2)} \right)}{m+4} \right)$$

$$\frac{(cd_1x + d_1)^{5/2}(d_2 - cd_2x)^{5/2}(fx)^{m+1}(a + \operatorname{arccosh}(cx))}{f(m+6)} - \frac{bcd_1^2d_2^2\sqrt{cd_1x + d_1}\sqrt{d_2 - cd_2x} \left( \frac{c^4(fx)^{m+6}}{f^5(m+6)} - \frac{2c^2(fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6364

$$5d_1d_2 \left( \frac{3d_1d_2 \left( -\frac{\sqrt{cd_1x+d_1}\sqrt{d_2-cd_2x} \left( \frac{bc(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2}+1, \frac{m}{2}+1; \frac{m}{2}+\frac{3}{2}, \frac{m}{2}+2; c^2x^2\right)}{f^2(m+1)(m+2)} + \frac{\sqrt{1-cx}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2\right)}{f(m+1)\sqrt{cx-1}} \right)}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)}{m+4}$$

$$\frac{(cd_1x + d_1)^{5/2}(d_2 - cd_2x)^{5/2}(fx)^{m+1}(a + \operatorname{arccosh}(cx))}{f(m+6)} - \frac{bcd_1^2d_2^2\sqrt{cd_1x + d_1}\sqrt{d_2 - cd_2x} \left( \frac{c^4(fx)^{m+6}}{f^5(m+6)} - \frac{2c^2(fx)^{m+4}}{f^3(m+4)} + \frac{(fx)^{m+2}}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}\sqrt{cx+1}}$$

```
input Int[(f*x)^m*(d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2)*(a + b*ArcCosh[c*x]),x]
```

output 
$$-\left(\frac{(b*c*d1^2*d2^2*\sqrt{d1 + c*d1*x}*\sqrt{d2 - c*d2*x}*((f*x)^{(2 + m)})/(f*(2 + m)) - (2*c^2*(f*x)^{(4 + m)})/(f^3*(4 + m)) + (c^4*(f*x)^{(6 + m)})/(f^5*(6 + m)))}{(f*(6 + m)*\sqrt{-1 + c*x}*\sqrt{1 + c*x})} + \frac{((f*x)^{(1 + m)}*(d1 + c*d1*x)^{(5/2)}*(d2 - c*d2*x)^{(5/2)}*(a + b*\text{ArcCosh}[c*x]))}{(f*(6 + m))} + (5*d1*d2*(-((b*c*d1*d2*\sqrt{d1 + c*d1*x}*\sqrt{d2 - c*d2*x}*((f*x)^{(2 + m)})/(f*(2 + m)) - (c^2*(f*x)^{(4 + m)})/(f^3*(4 + m))))}{(f*(4 + m)*\sqrt{-1 + c*x}*\sqrt{1 + c*x})} + \frac{((f*x)^{(1 + m)}*(d1 + c*d1*x)^{(3/2)}*(d2 - c*d2*x)^{(3/2)}*(a + b*\text{ArcCosh}[c*x]))}{(f*(4 + m))} + \frac{(3*d1*d2*(-((b*c*(f*x)^{(2 + m)}*\sqrt{d1 + c*d1*x}*\sqrt{d2 - c*d2*x}))/f^{2*(2 + m)}*\sqrt{-1 + c*x}*\sqrt{1 + c*x})}{(f*(2 + m))} - \frac{(\sqrt{d1 + c*d1*x}*\sqrt{d2 - c*d2*x}*((f*x)^{(1 + m)}*\sqrt{1 - c*x}*(a + b*\text{ArcCosh}[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])}{(f*(1 + m)*\sqrt{-1 + c*x})} + \frac{(b*c*(f*x)^{(2 + m)}*HypergeometricPFQ[\{1, 1 + m/2, 1 + m/2\}, \{3/2 + m/2, 2 + m/2\}, c^2*x^2])}{(f^2*(1 + m)*(2 + m))})}{((2 + m)*\sqrt{-1 + c*x}*\sqrt{1 + c*x})})/(4 + m))/(6 + m)$$

### 3.157.3.1 Defintions of rubi rules used

rule 17  $\text{Int}[(c\_.)*((a\_.) + (b\_.)*(x\_))^m, x\_Symbol] \rightarrow \text{Simp}[c*((a + b*x)^{m+1})/(b*(m+1)), x] /; \text{FreeQ}\{a, b, c, m, x\} \ \&\& \ \text{NeQ}[m, -1]$

rule 25  $\text{Int}[-(F_x), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \ \text{Int}[F_x, x], x]$

rule 82  $\text{Int}[(a\_.) + (b\_.)*(x\_))^m*((c\_.) + (d\_.)*(x\_))^n*((e\_.) + (f\_.)*(x\_))^p, x_] \rightarrow \text{Int}[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[n, m] \ \&\& \ \text{IntegerQ}[m]$

rule 244  $\text{Int}[(c\_.)*(x\_))^m*((a\_.) + (b\_.)*(x\_)^2)^p, x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, m\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

rule 2009  $\text{Int}[u, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$

```
rule 6342 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d1_
+ (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] :> Simp[(f*x)^(m + 1)*S
qrt[d1 + e1*x]*Sqrt[d2 + e2*x]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (
-Simp[(1/(m + 2))*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/
Sqrt[-1 + c*x]] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-
1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d1 + e1*x]/Sqrt[1 +
c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[(f*x)^(m + 1)*(a + b*ArcC
osh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] &&
EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1
])
```

```
rule 6346 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_ + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[(f*x)^(m + 1)
*(d1 + e1*x)^p*(d2 + e2*x)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x]
+ (Simp[2*d1*d2*(p/(m + 2*p + 1)) Int[(f*x)^m*(d1 + e1*x)^(p - 1)*(d2 +
e2*x)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)
))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(
f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^
(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*
d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]
```

```
rule 6364 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)/(Sqrt[(d1_ + (
e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] :> Simp[((f*x)^(m + 1)/(f
*(m + 1))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b
*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] +
Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 +
e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2,
1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2
, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]
```

### 3.157.4 Maple [F]

$$\int (fx)^m (cd1x + d1)^{5/2} (-cd2x + d2)^{5/2} (a + b \operatorname{arccosh}(cx)) dx$$

```
input int((f*x)^m*(c*d1*x+d1)^(5/2)*(-c*d2*x+d2)^(5/2)*(a+b*arccosh(c*x)),x)
```

```
output int((f*x)^m*(c*d1*x+d1)^(5/2)*(-c*d2*x+d2)^(5/2)*(a+b*arccosh(c*x)),x)
```

---


$$3.157. \quad \int (fx)^m (d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2} (a + b \operatorname{arccosh}(cx)) dx$$



**3.157.5 Fracas [F]**

$$\int (fx)^m (d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (cd_1x + d_1)^{5/2} (-cd_2x + d_2)^{5/2} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(c*d1*x+d1)^(5/2)*(-c*d2*x+d2)^(5/2)*(a+b*arccosh(c*x)), x, algorithm="fricas")`

output `integral((a*c^4*d1^2*d2^2*x^4 - 2*a*c^2*d1^2*d2^2*x^2 + a*d1^2*d2^2 + (b*c^4*d1^2*d2^2*x^4 - 2*b*c^2*d1^2*d2^2*x^2 + b*d1^2*d2^2)*arccosh(c*x))*sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(f*x)^m, x)`

**3.157.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(c*d1*x+d1)**(5/2)*(-c*d2*x+d2)**(5/2)*(a+b*acosh(c*x)), x)`

output `Timed out`

**3.157.7 Maxima [F]**

$$\int (fx)^m (d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2} (a + \operatorname{barccosh}(cx)) dx = \int (cd_1x + d_1)^{5/2} (-cd_2x + d_2)^{5/2} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(c*d1*x+d1)^(5/2)*(-c*d2*x+d2)^(5/2)*(a+b*arccosh(c*x)), x, algorithm="maxima")`

output `integrate((c*d1*x + d1)^(5/2)*(-c*d2*x + d2)^(5/2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.157.8 Giac [F]**

$$\int (fx)^m (d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2} (a + b \operatorname{arccosh}(cx)) dx = \int (cd_1x + d_1)^{5/2} (-cd_2x + d_2)^{5/2} (b \operatorname{arccosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(c*d1*x+d1)^(5/2)*(-c*d2*x+d2)^(5/2)*(a+b*arccosh(c*x)), x, algorithm="giac")`

output `integrate((c*d1*x + d1)^(5/2)*(-c*d2*x + d2)^(5/2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.157.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2} (a + b \operatorname{arccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (fx)^m (d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2} dx$$

input `int((a + b*acosh(c*x))*(f*x)^m*(d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2), x)`

output `int((a + b*acosh(c*x))*(f*x)^m*(d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2), x)`

### 3.158 $\int (fx)^m (d1 + cd1x)^{3/2} (d2 - cd2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

3.158.1 Optimal result . . . . .	1322
3.158.2 Mathematica [A] (verified) . . . . .	1323
3.158.3 Rubi [A] (verified) . . . . .	1324
3.158.4 Maple [F] . . . . .	1327
3.158.5 Fricas [F] . . . . .	1328
3.158.6 Sympy [F(-1)] . . . . .	1328
3.158.7 Maxima [F] . . . . .	1328
3.158.8 Giac [F] . . . . .	1329
3.158.9 Mupad [F(-1)] . . . . .	1329

#### 3.158.1 Optimal result

Integrand size = 35, antiderivative size = 503

$$\int (fx)^m (d1 + cd1x)^{3/2} (d2 - cd2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx = -\frac{3bcd1d2(fx)^{2+m}\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}}{f^2(2+m)^2(4+m)\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{bcd1d2(fx)^{2+m}\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}}{f^2(2+m)(4+m)\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3d1d2(fx)^{4+m}\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}}{f^4(4+m)^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{3d1d2(fx)^{1+m}\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}(a + \operatorname{barccosh}(cx))}{f(8 + 6m + m^2)}$$

$$+ \frac{(fx)^{1+m}(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}(a + \operatorname{barccosh}(cx))}{f(4 + m)}$$

$$+ \frac{3d1d2(fx)^{1+m}\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}(a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{f(4+m)(2+3m+m^2)\sqrt{1 - cx}\sqrt{1 + cx}}$$

$$- \frac{3bcd1d2(fx)^{2+m}\sqrt{d1 + cd1x}\sqrt{d2 - cd2x} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2x^2\right)}{f^2(1+m)(2+m)^2(4+m)\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output  $(f*x)^{(1+m)}*(c*d1*x+d1)^{(3/2)}*(-c*d2*x+d2)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))/f/(4+m)+3*d1*d2*(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))*(c*d1*x+d1)^{(1/2)}*(-c*d2*x+d2)^{(1/2)}/f/(m^2+6*m+8)+3*d1*d2*(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))*\operatorname{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], c^2*x^2)*(c*d1*x+d1)^{(1/2)}*(-c*d2*x+d2)^{(1/2)}/f/(m^3+7*m^2+14*m+8)/(-c*x+1)^{(1/2)}/(c*x+1)^{(1/2)}-3*b*c*d1*d2*(f*x)^{(2+m)}*(c*d1*x+d1)^{(1/2)}*(-c*d2*x+d2)^{(1/2)}/f^2/(2+m)^2/(4+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b*c*d1*d2*(f*x)^{(2+m)}*(c*d1*x+d1)^{(1/2)}*(-c*d2*x+d2)^{(1/2)}/f^2/(2+m)/(4+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+b*c^3*d1*d2*(f*x)^{(4+m)}*(c*d1*x+d1)^{(1/2)}*(-c*d2*x+d2)^{(1/2)}/f^4/(4+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3*b*c*d1*d2*(f*x)^{(2+m)}*\operatorname{hypergeom}([1, 1+1/2*m, 1+1/2*m], [2+1/2*m, 3/2+1/2*m], c^2*x^2)*(c*d1*x+d1)^{(1/2)}*(-c*d2*x+d2)^{(1/2)}/f^2/(2+m)^2/(m^2+5*m+4)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.158.2 Mathematica [A] (verified)

Time = 0.76 (sec) , antiderivative size = 288, normalized size of antiderivative = 0.57

$$\int (fx)^m (d1 + cd1x)^{3/2} (d2 - cd2x)^{3/2} (a + b \operatorname{arccosh}(cx)) dx = \frac{d1d2x(fx)^m \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} \left( -\frac{3bcx}{(2+m)^2 \sqrt{-1+cx} \sqrt{1+cx}} + \frac{bcx \left( -\frac{1}{2+m} + \frac{c^2}{4} \right)}{\sqrt{-1+cx} \sqrt{1+cx}} \right)}{(2+m)^2 \sqrt{-1+cx} \sqrt{1+cx}}$$

input `Integrate[(f*x)^m*(d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output  $(d1*d2*x*(f*x)^m*\operatorname{Sqrt}[d1 + c*d1*x]*\operatorname{Sqrt}[d2 - c*d2*x]*((-3*b*c*x)/((2 + m)^2*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]) + (b*c*x*(-(2 + m)^{-1} + (c^2*x^2)/(4 + m)))/(\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]) + (3*(a + b*\operatorname{ArcCosh}[c*x]))/(2 + m) - (-1 + c*x)*(1 + c*x)*(a + b*\operatorname{ArcCosh}[c*x]) - (3*\operatorname{Sqrt}[1 - c^2*x^2]*(a + b*\operatorname{ArcCosh}[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/((1 + m)*(2 + m)*(-1 + c*x)*(1 + c*x)) - (3*b*c*x*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/((1 + m)*(2 + m)^2*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]))/(4 + m)$

**3.158.3 Rubi [A] (verified)**

Time = 1.65 (sec) , antiderivative size = 428, normalized size of antiderivative = 0.85, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$ , Rules used = {6346, 25, 82, 244, 2009, 6342, 17, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (cd1x + d1)^{3/2} (d2 - cd2x)^{3/2} (fx)^m (a + \text{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6346} \\
 & \frac{3d1d2 \int (fx)^m \sqrt{cxd1 + d1} \sqrt{d2 - cd2x} (a + \text{barccosh}(cx)) dx}{m + 4} + \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int -(fx)^{m+1} (1 - cx)(cx + 1) dx}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} + \\
 & \frac{(cd1x + d1)^{3/2} (d2 - cd2x)^{3/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 4)} \\
 & \quad \downarrow \text{25} \\
 & \frac{3d1d2 \int (fx)^m \sqrt{cxd1 + d1} \sqrt{d2 - cd2x} (a + \text{barccosh}(cx)) dx}{m + 4} - \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int (fx)^{m+1} (1 - cx)(cx + 1) dx}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} + \\
 & \frac{(cd1x + d1)^{3/2} (d2 - cd2x)^{3/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 4)} \\
 & \quad \downarrow \text{82} \\
 & \frac{3d1d2 \int (fx)^m \sqrt{cxd1 + d1} \sqrt{d2 - cd2x} (a + \text{barccosh}(cx)) dx}{m + 4} - \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int (fx)^{m+1} (1 - c^2 x^2) dx}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} + \\
 & \frac{(cd1x + d1)^{3/2} (d2 - cd2x)^{3/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 4)} \\
 & \quad \downarrow \text{244} \\
 & \frac{3d1d2 \int (fx)^m \sqrt{cxd1 + d1} \sqrt{d2 - cd2x} (a + \text{barccosh}(cx)) dx}{m + 4} - \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int \left( (fx)^{m+1} - \frac{c^2 (fx)^{m+3}}{f^2} \right) dx}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} + \\
 & \frac{(cd1x + d1)^{3/2} (d2 - cd2x)^{3/2} (fx)^{m+1} (a + \text{barccosh}(cx))}{f(m + 4)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 2009 \\
 & \frac{3d1d2 \int (fx)^m \sqrt{cd1x + d1} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx)) dx}{(cd1x + d1)^{3/2} (d2 - cd2x)^{3/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))} + \\
 & \frac{m + 4}{f(m + 4)} - \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} \\
 & \downarrow 6342 \\
 & 3d1d2 \left( - \frac{\sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx - 1} \sqrt{cx + 1}}}{(m+2) \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{bc \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int (fx)^{m+1} dx}{f(m+2) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{\sqrt{cd1x + d1} \sqrt{d2 - cd2x}}{f(m+2)} \right) \\
 & \frac{m + 4}{f(m + 4)} - \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} \\
 & \downarrow 17 \\
 & 3d1d2 \left( - \frac{\sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx - 1} \sqrt{cx + 1}}}{(m+2) \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{\sqrt{cd1x + d1} \sqrt{d2 - cd2x} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \sqrt{cd1x + d1} \sqrt{d2 - cd2x}}{f^2(m+2)} \right) \\
 & \frac{m + 4}{f(m + 4)} - \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}} \\
 & \downarrow 6364 \\
 & 3d1d2 \left( - \frac{\sqrt{cd1x + d1} \sqrt{d2 - cd2x} \left( \frac{bc (fx)^{m+2} {}_3F_2 \left( 1, \frac{m}{2} + 1, \frac{m}{2} + 1; \frac{m}{2} + \frac{3}{2}, \frac{m}{2} + 2; c^2 x^2 \right) + \sqrt{1 - cx} (fx)^{m+1} \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2 x^2 \right) (a + \operatorname{barccosh}(cx))}{f^2(m+1)(m+2)} \right)}{(m+2) \sqrt{cx - 1} \sqrt{cx + 1}} \right) \\
 & \frac{m + 4}{f(m + 4)} - \\
 & \frac{bcd1d2 \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \left( \frac{(fx)^{m+2}}{f(m+2)} - \frac{c^2 (fx)^{m+4}}{f^3(m+4)} \right)}{f(m + 4) \sqrt{cx - 1} \sqrt{cx + 1}}
 \end{aligned}$$

input `Int[(f*x)^m*(d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)*(a + b*ArcCosh[c*x]),x]`

output `-((b*c*d1*d2*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*((f*x)^(2 + m)/(f*(2 + m) - (c^2*(f*x)^(4 + m))/(f^3*(4 + m)))))/(f*(4 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*(d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)*(a + b*ArcCosh[c*x]))/(f*(4 + m)) + (3*d1*d2*(-((b*c*(f*x)^(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]))/(f^2*(2 + m)^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*(a + b*ArcCosh[c*x]))/(f*(2 + m)) - (Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*(((f*x)^(1 + m)*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2]))/(f*(1 + m)*Sqrt[-1 + c*x]) + (b*c*(f*x)^(2 + m)*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m)))/((2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(4 + m)`

### 3.158.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

```
rule 6342 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d1_
+ (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] :> Simp[(f*x)^(m + 1)*S
qrt[d1 + e1*x]*Sqrt[d2 + e2*x]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (
-Simp[(1/(m + 2))*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/
Sqrt[-1 + c*x]] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-
1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d1 + e1*x]/Sqrt[1 +
c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[(f*x)^(m + 1)*(a + b*ArcC
osh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] &&
EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1
])
```

```
rule 6346 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_ + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[(f*x)^(m + 1)
*(d1 + e1*x)^p*(d2 + e2*x)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x]
+ (Simp[2*d1*d2*(p/(m + 2*p + 1)) Int[(f*x)^m*(d1 + e1*x)^(p - 1)*(d2 +
e2*x)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)
))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(
f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^
(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*
d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]
```

```
rule 6364 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)/(Sqrt[(d1_) + (
e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] :> Simp[((f*x)^(m + 1)/(f
*(m + 1))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b
*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] +
Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 +
e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2,
1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2
, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]
```

### 3.158.4 Maple [F]

$$\int (fx)^m (cd1x + d1)^{\frac{3}{2}} (-cd2x + d2)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx)) dx$$

```
input int((f*x)^m*(c*d1*x+d1)^(3/2)*(-c*d2*x+d2)^(3/2)*(a+b*arccosh(c*x)),x)
```

```
output int((f*x)^m*(c*d1*x+d1)^(3/2)*(-c*d2*x+d2)^(3/2)*(a+b*arccosh(c*x)),x)
```



**3.158.5 Fracas [F]**

$$\int (fx)^m (d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (cd_1x + d_1)^{\frac{3}{2}} (-cd_2x + d_2)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(c*d1*x+d1)^(3/2)*(-c*d2*x+d2)^(3/2)*(a+b*arccosh(c*x)),  
x, algorithm="fricas")`

output `integral(-(a*c^2*d1*d2*x^2 - a*d1*d2 + (b*c^2*d1*d2*x^2 - b*d1*d2)*arccosh  
(c*x))*sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(f*x)^m, x)`

**3.158.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(c*d1*x+d1)**(3/2)*(-c*d2*x+d2)**(3/2)*(a+b*acosh(c*x)),  
x)`

output `Timed out`

**3.158.7 Maxima [F]**

$$\int (fx)^m (d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx = \int (cd_1x + d_1)^{\frac{3}{2}} (-cd_2x + d_2)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(c*d1*x+d1)^(3/2)*(-c*d2*x+d2)^(3/2)*(a+b*arccosh(c*x)),  
x, algorithm="maxima")`

output `integrate((c*d1*x + d1)^(3/2)*(-c*d2*x + d2)^(3/2)*(b*arccosh(c*x) + a)*(f  
*x)^m, x)`

---

3.158.  $\int (fx)^m (d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2} (a + \operatorname{barccosh}(cx)) dx$

**3.158.8 Giac [F]**

$$\int (fx)^m (d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2} (a + b \operatorname{arccosh}(cx)) dx = \int (cd_1x + d_1)^{\frac{3}{2}} (-cd_2x + d_2)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(c*d1*x+d1)^(3/2)*(-c*d2*x+d2)^(3/2)*(a+b*arccosh(c*x)), x, algorithm="giac")`

output `integrate((c*d1*x + d1)^(3/2)*(-c*d2*x + d2)^(3/2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.158.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2} (a + b \operatorname{arccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (fx)^m (d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2} dx$$

input `int((a + b*acosh(c*x))*(f*x)^m*(d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2), x)`

output `int((a + b*acosh(c*x))*(f*x)^m*(d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2), x)`

### 3.159 $\int (fx)^m \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx)) dx$

3.159.1 Optimal result . . . . .	1330
3.159.2 Mathematica [A] (verified) . . . . .	1331
3.159.3 Rubi [A] (verified) . . . . .	1331
3.159.4 Maple [F] . . . . .	1333
3.159.5 Fracas [F] . . . . .	1333
3.159.6 Sympy [F(-1)] . . . . .	1334
3.159.7 Maxima [F] . . . . .	1334
3.159.8 Giac [F] . . . . .	1334
3.159.9 Mupad [F(-1)] . . . . .	1335

#### 3.159.1 Optimal result

Integrand size = 35, antiderivative size = 302

$$\int (fx)^m \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{bc(fx)^{2+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x}}{f^2(2+m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{(fx)^{1+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx))}{f(2+m)}$$

$$+ \frac{(fx)^{1+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2 x^2\right)}{f(2+3m+m^2) \sqrt{1 - cx} \sqrt{1 + cx}}$$

$$- \frac{bc(fx)^{2+m} \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2 x^2\right)}{f^2(1+m)(2+m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

```
output (f*x)^(1+m)*(a+b*arccosh(c*x))*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f/(2+m)
)+(f*x)^(1+m)*(a+b*arccosh(c*x))*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], c^
2*x^2)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f/(m^2+3*m+2)/(-c*x+1)^(1/2)/(
c*x+1)^(1/2)-b*c*(f*x)^(2+m)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f^2/(2+m)
)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*c*(f*x)^(2+m)*hypergeom([1, 1+1/2*m, 1+1
/2*m], [2+1/2*m, 3/2+1/2*m], c^2*x^2)*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)/f
^2/(1+m)/(2+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.159.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.76

$$\int (fx)^m \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{x(fx)^m \sqrt{d1 + cd1x} \sqrt{d2 - cd2x} ((1 + m) (-bcx \sqrt{-1 + cx} \sqrt{1 + cx} + a(2 + m) (-1 + c^2x^2) + b(2 + m) ($$

input `Integrate[(f*x)^m*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*(a + b*ArcCosh[c*x]),x]`

output `(x*(f*x)^m*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*((1 + m)*(-b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + a*(2 + m)*(-1 + c^2*x^2) + b*(2 + m)*(-1 + c^2*x^2)*ArcCosh[c*x]) - (2 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2] - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2))/((1 + m)*(2 + m)^2*(-1 + c*x)*(1 + c*x))`

### 3.159.3 Rubi [A] (verified)

Time = 0.91 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.94, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.086$ , Rules used = {6342, 17, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{cd1x + d1} \sqrt{d2 - cd2x} (fx)^m (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6342}$$

$$\frac{\sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int \frac{(fx)^{m(a + \operatorname{barccosh}(cx))} dx}{\sqrt{cx - 1} \sqrt{cx + 1}}}{(m + 2) \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{bc \sqrt{cd1x + d1} \sqrt{d2 - cd2x} \int (fx)^{m+1} dx}{f(m + 2) \sqrt{cx - 1} \sqrt{cx + 1}} +$$

$$\frac{\sqrt{cd1x + d1} \sqrt{d2 - cd2x} (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m + 2)}$$

$$\downarrow \text{17}$$

$$-\frac{\sqrt{cd1x + d1}\sqrt{d2 - cd2x} \int \frac{(fx)^m(a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx-1}\sqrt{cx+1}}}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{\sqrt{cd1x + d1}\sqrt{d2 - cd2x}(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{cd1x + d1}\sqrt{d2 - cd2x}(fx)^{m+2}}{f^2(m+2)^2\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6364

$$-\frac{\sqrt{cd1x + d1}\sqrt{d2 - cd2x} \left( \frac{bc(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2}+1, \frac{m}{2}+1; \frac{m}{2}+\frac{3}{2}, \frac{m}{2}+2; c^2x^2\right)}{f^2(m+1)(m+2)} + \frac{\sqrt{1-cx}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2x^2\right)}{f(m+1)\sqrt{cx-1}} \right)}{(m+2)\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{cd1x + d1}\sqrt{d2 - cd2x}(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{cd1x + d1}\sqrt{d2 - cd2x}(fx)^{m+2}}{f^2(m+2)^2\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(f*x)^m*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*(a + b*ArcCosh[c*x]),x]`

output `-((b*c*(f*x)^(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x])/(f^2*(2 + m)^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + ((f*x)^(1 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*(a + b*ArcCosh[c*x]))/(f*(2 + m)) - (Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]*(((f*x)^(1 + m)*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[-1 + c*x]) + (b*c*(f*x)^(2 + m)*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m))))/((2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.159.3.1 Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1))], x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

rule 6342 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d1_ + (e1_.)*(x_)]*Sqrt[(d2_ + (e2_.)*(x_)]), x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6364 `Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]`

### 3.159.4 Maple [F]

$$\int (fx)^m \sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2} (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)*(a+b*arccosh(c*x)),x)`

### 3.159.5 Fracas [F]

$$\begin{aligned} & \int (fx)^m \sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x} (a + b \operatorname{arccosh}(cx)) dx \\ &= \int \sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2} (b \operatorname{arccosh}(cx) + a) (fx)^m dx \end{aligned}$$

input `integrate((f*x)^m*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)*(a+b*arccosh(c*x)), x, algorithm="fracas")`

output `integral(sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.159.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m \sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x} (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(c*d1*x+d1)**(1/2)*(-c*d2*x+d2)**(1/2)*(a+b*acosh(c*x)),x)`

output `Timed out`

**3.159.7 Maxima [F]**

$$\begin{aligned} & \int (fx)^m \sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x} (a + \operatorname{barccosh}(cx)) dx \\ &= \int \sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2} (b \operatorname{arcosh}(cx) + a) (fx)^m dx \end{aligned}$$

input `integrate((f*x)^m*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.159.8 Giac [F]**

$$\begin{aligned} & \int (fx)^m \sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x} (a + \operatorname{barccosh}(cx)) dx \\ &= \int \sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2} (b \operatorname{arcosh}(cx) + a) (fx)^m dx \end{aligned}$$

input `integrate((f*x)^m*(c*d1*x+d1)^(1/2)*(-c*d2*x+d2)^(1/2)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.159.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m \sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x} (a + b \operatorname{arccosh}(cx)) dx$$

$$= \int (a + b \operatorname{acosh}(cx)) (fx)^m \sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x} dx$$

input `int((a + b*acosh(c*x))*(f*x)^m*(d1 + c*d1*x)^(1/2)*(d2 - c*d2*x)^(1/2),x)`output `int((a + b*acosh(c*x))*(f*x)^m*(d1 + c*d1*x)^(1/2)*(d2 - c*d2*x)^(1/2), x)`



$$3.160 \quad \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{\sqrt{d1+cd1x}\sqrt{d2-cd2x}} dx$$

3.160.1 Optimal result . . . . .	1336
3.160.2 Mathematica [A] (verified) . . . . .	1336
3.160.3 Rubi [A] (verified) . . . . .	1337
3.160.4 Maple [F] . . . . .	1338
3.160.5 Fricas [F] . . . . .	1338
3.160.6 Sympy [F] . . . . .	1339
3.160.7 Maxima [F] . . . . .	1339
3.160.8 Giac [F] . . . . .	1339
3.160.9 Mupad [F(-1)] . . . . .	1340

### 3.160.1 Optimal result

Integrand size = 35, antiderivative size = 188

$$\begin{aligned} & \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{\sqrt{d1+cd1x}\sqrt{d2-cd2x}} dx \\ &= \frac{(fx)^{1+m}\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{f(1+m)\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \\ & \quad + \frac{bc(fx)^{2+m}\sqrt{-1+cx}\sqrt{1+cx} {}_3F_2\left(1, 1+\frac{m}{2}, 1+\frac{m}{2}; \frac{3}{2}+\frac{m}{2}, 2+\frac{m}{2}; c^2x^2\right)}{f^2(1+m)(2+m)\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \end{aligned}$$

```
output b*c*(f*x)^(2+m)*hypergeom([1, 1+1/2*m, 1+1/2*m], [2+1/2*m, 3/2+1/2*m], c^2*x
^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/f^(2/(1+m))/(2+m)/(c*d1*x+d1)^(1/2)/(-c*d2*x
+d2)^(1/2)+(f*x)^(1+m)*(a+b*arccosh(c*x))*hypergeom([1/2, 1/2+1/2*m], [3/2+
1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/f/(1+m)/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(
1/2)
```

### 3.160.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.81

$$\begin{aligned} & \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{\sqrt{d1+cd1x}\sqrt{d2-cd2x}} dx \\ &= \frac{x(fx)^m((2+m)\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right) + bcx\sqrt{-1+cx}\sqrt{1+cx}}{(1+m)(2+m)\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \end{aligned}$$

---

3.160.  $\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{\sqrt{d1+cd1x}\sqrt{d2-cd2x}} dx$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]),x]`

output `(x*(f*x)^m*((2 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2] + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2)]/((1 + m)*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x])`

### 3.160.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.029$ , Rules used = {6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{cd_1x + d_1}\sqrt{d_2 - cd_2x}} dx$$

↓ 6364

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2} + 1, \frac{m}{2} + 1; \frac{m}{2} + \frac{3}{2}, \frac{m}{2} + 2; c^2x^2\right)}{f^2(m+1)(m+2)\sqrt{cd_1x + d_1}\sqrt{d_2 - cd_2x}} + \frac{\sqrt{1 - c^2x^2}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2x^2\right) (a + \operatorname{barccosh}(cx))}{f(m+1)\sqrt{cd_1x + d_1}\sqrt{d_2 - cd_2x}}$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]),x]`

output `((f*x)^(1 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2)]/(f^2*(1 + m)*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x])`

## 3.160.3.1 Defintions of rubi rules used

```
rule 6364 Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.))/(Sqrt[(d1_) + (
e1_.)*(x_.)]*Sqrt[(d2_) + (e2_.)*(x_.)]), x_Symbol] := Simp[((f*x)^(m + 1)/(f
*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b
*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] +
Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 +
e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2,
1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2
, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]
```

## 3.160.4 Maple [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2}} dx$$

```
input int((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(1/2),x)
```

```
output int((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(1/2),x)
```

## 3.160.5 Fracas [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{\sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(1/2),
x, algorithm="fracas")
```

```
output integral(-sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(b*arccosh(c*x) + a)*(f*x)^
m/(c^2*d1*d2*x^2 - d1*d2), x)
```

**3.160.6 Sympy [F]**

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))}{\sqrt{d_1}(cx+1) \sqrt{-d_2}(cx-1)} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(c*d1*x+d1)**(1/2)/(-c*d2*x+d2)**(1/2),x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))/(sqrt(d1*(c*x + 1))*sqrt(-d2*(c*x - 1))), x)`

**3.160.7 Maxima [F]**

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{\sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(1/2), x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)), x)`

**3.160.8 Giac [F]**

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{d_1 + cd_1x} \sqrt{d_2 - cd_2x}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{\sqrt{cd_1x + d_1} \sqrt{-cd_2x + d_2}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(1/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)), x)`

**3.160.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{\sqrt{d_1 + cd_1 x} \sqrt{d_2 - cd_2 x}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{\sqrt{d_1 + cd_1 x} \sqrt{d_2 - cd_2 x}} dx$$

input `int((a + b*acosh(c*x))*(f*x)^m)/((d1 + c*d1*x)^(1/2)*(d2 - c*d2*x)^(1/2)),x)`

output `int((a + b*acosh(c*x))*(f*x)^m)/((d1 + c*d1*x)^(1/2)*(d2 - c*d2*x)^(1/2)), x)`

**3.161** 
$$\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d1+cd1x)^{3/2}(d2-cd2x)^{3/2}} dx$$

3.161.1 Optimal result . . . . . 1341  
 3.161.2 Mathematica [A] (verified) . . . . . 1342  
 3.161.3 Rubi [A] (verified) . . . . . 1342  
 3.161.4 Maple [F] . . . . . 1345  
 3.161.5 Fricas [F] . . . . . 1345  
 3.161.6 Sympy [F(-1)] . . . . . 1345  
 3.161.7 Maxima [F] . . . . . 1346  
 3.161.8 Giac [F] . . . . . 1346  
 3.161.9 Mupad [F(-1)] . . . . . 1346

**3.161.1 Optimal result**

Integrand size = 35, antiderivative size = 336

$$\int \frac{(fx)^m(a + b\operatorname{arccosh}(cx))}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx = \frac{(fx)^{1+m}(a + b\operatorname{arccosh}(cx))}{d1d2f\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}}$$

$$- \frac{m(fx)^{1+m}\sqrt{1 - c^2x^2}(a + b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{d1d2f(1 + m)\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}}$$

$$+ \frac{bc(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{d1d2f^2(2 + m)\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}}$$

$$- \frac{bcm(fx)^{2+m}\sqrt{-1 + cx}\sqrt{1 + cx} {}_3F_2\left(1, 1 + \frac{m}{2}, 1 + \frac{m}{2}; \frac{3}{2} + \frac{m}{2}, 2 + \frac{m}{2}; c^2x^2\right)}{d1d2f^2(1 + m)(2 + m)\sqrt{d1 + cd1x}\sqrt{d2 - cd2x}}$$

```
output (f*x)^(1+m)*(a+b*arccosh(c*x))/d1/d2/f/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(1/2)
)+b*c*(f*x)^(2+m)*hypergeom([1, 1+1/2*m], [2+1/2*m], c^2*x^2)*(c*x-1)^(1/2)*
(c*x+1)^(1/2)/d1/d2/f^2/(2+m)/(c*d1*x+d1)^(1/2)/(-c*d2*x+d2)^(1/2)-b*c*m*(
f*x)^(2+m)*hypergeom([1, 1+1/2*m, 1+1/2*m], [2+1/2*m, 3/2+1/2*m], c^2*x^2)*(
c*x-1)^(1/2)*(c*x+1)^(1/2)/d1/d2/f^2/(1+m)/(2+m)/(c*d1*x+d1)^(1/2)/(-c*d2*
x+d2)^(1/2)-m*(f*x)^(1+m)*(a+b*arccosh(c*x))*hypergeom([1/2, 1/2+1/2*m], [3
/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/d1/d2/f/(1+m)/(c*d1*x+d1)^(1/2)/(-c*
d2*x+d2)^(1/2)
```

**3.161.2 Mathematica [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 225, normalized size of antiderivative = 0.67

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx = \frac{x(fx)^m(-m(2+m)\sqrt{1-c^2x^2}(a + \operatorname{barccosh}(cx)) \operatorname{Hypergeometric2F1}[\dots])}{\dots}$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/((d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)),x]`

output `(x*(f*x)^m*(-(m*(2 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2]) + (1 + m)*((2 + m)*(a + b*ArcCosh[c*x]) + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[1, 1 + m/2, 2 + m/2, c^2*x^2]) - b*c*m*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2]))/(d1*d2*(1 + m)*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x])`

**3.161.3 Rubi [A] (verified)**Time = 1.19 (sec) , antiderivative size = 330, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6352, 25, 82, 278, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(cd1x + d1)^{3/2}(d2 - cd2x)^{3/2}} dx \\ & \quad \downarrow \text{6352} \\ & -\frac{m \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{\sqrt{cx}d1 + d1\sqrt{d2 - cd2x}} dx}{d1d2} - \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{d1d2f\sqrt{cd1x + d1}\sqrt{d2 - cd2x}} + \\ & \quad \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{d1d2f\sqrt{cd1x + d1}\sqrt{d2 - cd2x}} \\ & \quad \downarrow \text{25} \\ & -\frac{m \int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{\sqrt{cx}d1 + d1\sqrt{d2 - cd2x}} dx}{d1d2} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{d1d2f\sqrt{cd1x + d1}\sqrt{d2 - cd2x}} + \\ & \quad \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{d1d2f\sqrt{cd1x + d1}\sqrt{d2 - cd2x}} \end{aligned}$$

---

3.161.  $\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx$

$$\begin{aligned}
 & \downarrow 82 \\
 & -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx} d1 + d1 \sqrt{d2 - cd2x}}}{d1 d2} + \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} \int \frac{(fx)^{m+1} dx}{1 - c^2 x^2}}{d1 d2 f \sqrt{cd1x + d1} \sqrt{d2 - cd2x}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{d1 d2 f \sqrt{cd1x + d1} \sqrt{d2 - cd2x}} \\
 & \downarrow 278 \\
 & -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx} d1 + d1 \sqrt{d2 - cd2x}}}{d1 d2} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{d1 d2 f \sqrt{cd1x + d1} \sqrt{d2 - cd2x}} + \\
 & \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{d1 d2 f^2 (m+2) \sqrt{cd1x + d1} \sqrt{d2 - cd2x}} \\
 & \downarrow 6364 \\
 & m \left( \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} (fx)^{m+2} {}_3F_2\left(1, \frac{m}{2} + 1, \frac{m}{2} + 1; \frac{m}{2} + \frac{3}{2}, \frac{m}{2} + 2; c^2 x^2\right)}{f^2 (m+1) (m+2) \sqrt{cd1x + d1} \sqrt{d2 - cd2x}} + \frac{\sqrt{1 - c^2 x^2} (fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2 x^2\right) (a + \operatorname{barccosh}(cx))}{f (m+1) \sqrt{cd1x + d1} \sqrt{d2 - cd2x}} \right) \\
 & \frac{d1 d2}{(fx)^{m+1} (a + \operatorname{barccosh}(cx))} + \\
 & \frac{bc \sqrt{cx - 1} \sqrt{cx + 1} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, c^2 x^2\right)}{d1 d2 f^2 (m+2) \sqrt{cd1x + d1} \sqrt{d2 - cd2x}}
 \end{aligned}$$

input `Int(((f*x)^m*(a + b*ArcCosh[c*x]))/((d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)),x]`

output `((f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(d1*d2*f*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, c^2*x^2])/(d1*d2*f^2*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) - (m*(((f*x)^(1 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]))/(d1*d2)`



## 3.161.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 82 `Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 278 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`
- rule 6352 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*((f_)*(x_)^(m_))*((d1_) + (e1_)*(x_)^(p_))*((d2_) + (e2_)*(x_)^(p_)), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d1*d2*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d1*d2*(p + 1)) Int[(f*x)^m*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || EqQ[n, 1])`
- rule 6364 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_)^(m_))/(Sqrt[(d1_) + (e1_)*(x_)])*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] := Simp[((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]`

**3.161.4 Maple [F]**

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(cd_1x + d_1)^{\frac{3}{2}} (-cd_2x + d_2)^{\frac{3}{2}}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(3/2)/(-c*d2*x+d2)^(3/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(3/2)/(-c*d2*x+d2)^(3/2),x)`

**3.161.5 Fricas [F]**

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{(cd_1x + d_1)^{\frac{3}{2}} (-cd_2x + d_2)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(3/2)/(-c*d2*x+d2)^(3/2),  
x, algorithm="fricas")`

output `integral(sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(b*arccosh(c*x) + a)*(f*x)^m  
/(c^4*d1^2*d2^2*x^4 - 2*c^2*d1^2*d2^2*x^2 + d1^2*d2^2), x)`

**3.161.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d_1 + cd_1x)^{3/2} (d_2 - cd_2x)^{3/2}} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(c*d1*x+d1)**(3/2)/(-c*d2*x+d2)**(3/2),x)`

output `Timed out`

**3.161.7 Maxima [F]**

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(cd1x + d1)^{\frac{3}{2}}(-cd2x + d2)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(3/2)/(-c*d2*x+d2)^(3/2), x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/((c*d1*x + d1)^(3/2)*(-c*d2*x + d2)^(3/2)), x)`

**3.161.8 Giac [F]**

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(cd1x + d1)^{\frac{3}{2}}(-cd2x + d2)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(3/2)/(-c*d2*x+d2)^(3/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/((c*d1*x + d1)^(3/2)*(-c*d2*x + d2)^(3/2)), x)`

**3.161.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))(fx)^m}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/((d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)), x)`

output `int(((a + b*acosh(c*x))*(f*x)^m)/((d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)), x)`

---

3.161.  $\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{3/2}(d2 - cd2x)^{3/2}} dx$

$$3.162 \quad \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d1+cd1x)^{5/2}(d2-cd2x)^{5/2}} dx$$

3.162.1 Optimal result	1347
3.162.2 Mathematica [A] (verified)	1348
3.162.3 Rubi [A] (verified)	1349
3.162.4 Maple [F]	1352
3.162.5 Fricas [F]	1352
3.162.6 Sympy [F(-1)]	1353
3.162.7 Maxima [F]	1353
3.162.8 Giac [F]	1353
3.162.9 Mupad [F(-1)]	1354

### 3.162.1 Optimal result

Integrand size = 35, antiderivative size = 504

$$\begin{aligned} \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d1+cd1x)^{5/2}(d2-cd2x)^{5/2}} dx &= \frac{(fx)^{1+m}(a+b\operatorname{arccosh}(cx))}{3d1d2f(d1+cd1x)^{3/2}(d2-cd2x)^{3/2}} \\ &+ \frac{(2-m)(fx)^{1+m}(a+b\operatorname{arccosh}(cx))}{3d1^2d2^2f\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \\ &- \frac{(2-m)m(fx)^{1+m}\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx)) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, c^2x^2\right)}{3d1^2d2^2f(1+m)\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \\ &+ \frac{bc(2-m)(fx)^{2+m}\sqrt{-1+cx}\sqrt{1+cx} \operatorname{Hypergeometric2F1}\left(1, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{3d1^2d2^2f^2(2+m)\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \\ &+ \frac{bc(fx)^{2+m}\sqrt{-1+cx}\sqrt{1+cx} \operatorname{Hypergeometric2F1}\left(2, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{3d1^2d2^2f^2(2+m)\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \\ &- \frac{bc(2-m)m(fx)^{2+m}\sqrt{-1+cx}\sqrt{1+cx} {}_3F_2\left(1, 1+\frac{m}{2}, 1+\frac{m}{2}; \frac{3}{2}+\frac{m}{2}, 2+\frac{m}{2}; c^2x^2\right)}{3d1^2d2^2f^2(1+m)(2+m)\sqrt{d1+cd1x}\sqrt{d2-cd2x}} \end{aligned}$$

---


$$3.162. \quad \int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d1+cd1x)^{5/2}(d2-cd2x)^{5/2}} dx$$

output  $\frac{1}{3}(fx)^{(1+m)}(a+b\operatorname{arccosh}(cx))/d_1/d_2/f/(c*d_1*x+d_1)^{(3/2)/(-c*d_2*x+d_2)^{(3/2)+1/3*(2-m)}(fx)^{(1+m)}(a+b\operatorname{arccosh}(cx))/d_1^2/d_2^2/f/(c*d_1*x+d_1)^{(1/2)/(-c*d_2*x+d_2)^{(1/2)+1/3*b*c*(2-m)}(fx)^{(2+m)}\operatorname{hypergeom}([1, 1+1/2*m], [2+1/2*m], c^2*x^2)*(c*x-1)^{(1/2)}(c*x+1)^{(1/2)}/d_1^2/d_2^2/f^2/(2+m)/(c*d_1*x+d_1)^{(1/2)/(-c*d_2*x+d_2)^{(1/2)+1/3*b*c*(fx)^{(2+m)}\operatorname{hypergeom}([2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(c*x-1)^{(1/2)}(c*x+1)^{(1/2)}/d_1^2/d_2^2/f^2/(2+m)/(c*d_1*x+d_1)^{(1/2)/(-c*d_2*x+d_2)^{(1/2)-1/3*b*c*(2-m)*m*(fx)^{(2+m)}\operatorname{hypergeom}([1, 1+1/2*m], [2+1/2*m], 3/2+1/2*m], c^2*x^2)*(c*x-1)^{(1/2)}(c*x+1)^{(1/2)}/d_1^2/d_2^2/f^2/(1+m)/(2+m)/(c*d_1*x+d_1)^{(1/2)/(-c*d_2*x+d_2)^{(1/2)-1/3*(2-m)*m*(fx)^{(1+m)}(a+b\operatorname{arccosh}(cx))*\operatorname{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^{(1/2)}/d_1^2/d_2^2/f/(1+m)/(c*d_1*x+d_1)^{(1/2)/(-c*d_2*x+d_2)^{(1/2)}}$

### 3.162.2 Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 413, normalized size of antiderivative = 0.82

$$\int \frac{(fx)^m(a + b\operatorname{arccosh}(cx))}{(d_1 + cd_1x)^{5/2}(d_2 - cd_2x)^{5/2}} dx = \frac{x(fx)^m((1+m)(2+m)(2+3m+m^2)(a + b\operatorname{arccosh}(cx)) - bc(1 +$$

input `Integrate[(f*x)^m*(a + b*ArcCosh[c*x])/((d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2)),x]`

output  $(x*(fx)^m*((1+m)*(2+m)*(2+3m+m^2)*(a + b\operatorname{ArcCosh}[c*x]) - b*c*(1+m)*(2+3m+m^2)*x*(-1+c*x)^{(3/2)}(1+c*x)^{(3/2)}\operatorname{Hypergeometric2F1}[2, 1+m/2, 2+m/2, c^2*x^2] + (2-m)*(2+m)*(1-c*x)*(1+c*x)*((1+m)^2*(2+m)*(a + b\operatorname{ArcCosh}[c*x]) + b*c*(1+m)^2*x*\operatorname{Sqrt}[-1+c*x]*\operatorname{Sqrt}[1+c*x]*\operatorname{Hypergeometric2F1}[1, 1+m/2, 2+m/2, c^2*x^2] - m*(a*(2+3m+m^2)*\operatorname{Sqrt}[1-c^2*x^2]*\operatorname{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, c^2*x^2] + b*(2+3m+m^2)*\operatorname{Sqrt}[1-c^2*x^2]*\operatorname{ArcCosh}[c*x]*\operatorname{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, c^2*x^2] + b*c*(1+m)*x*\operatorname{Sqrt}[-1+c*x]*\operatorname{Sqrt}[1+c*x]*\operatorname{HypergeometricPFQ}[\{1, 1+m/2, 1+m/2\}, \{3/2+m/2, 2+m/2\}, c^2*x^2])))/(3*d_1^2*d_2^2*(1+m)*(2+m)*(2+3m+m^2)*(1-c*x)*(1+c*x)*\operatorname{Sqrt}[d_1 + c*d_1*x]*\operatorname{Sqrt}[d_2 - c*d_2*x])$

---

3.162.  $\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))}{(d_1+cd_1x)^{5/2}(d_2-cd_2x)^{5/2}} dx$

### 3.162.3 Rubi [A] (verified)

Time = 1.84 (sec) , antiderivative size = 484, normalized size of antiderivative = 0.96, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$ , Rules used = {6352, 82, 278, 6352, 25, 82, 278, 6364}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(cd1x + d1)^{5/2} (d2 - cd2x)^{5/2}} dx \\
 & \quad \downarrow \text{6352} \\
 & \frac{(2-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(cx d1 + d1)^{3/2} (d2 - cd2x)^{3/2}} dx}{3d1d2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{(1-cx)^2 (cx+1)^2} dx}{3d1^2 d2^2 f \sqrt{cd1x+d1} \sqrt{d2-cd2x}} + \\
 & \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3d1d2 f (cd1x + d1)^{3/2} (d2 - cd2x)^{3/2}} \\
 & \quad \downarrow \text{82} \\
 & \frac{(2-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(cx d1 + d1)^{3/2} (d2 - cd2x)^{3/2}} dx}{3d1d2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{(1-c^2x^2)^2} dx}{3d1^2 d2^2 f \sqrt{cd1x+d1} \sqrt{d2-cd2x}} + \\
 & \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3d1d2 f (cd1x + d1)^{3/2} (d2 - cd2x)^{3/2}} \\
 & \quad \downarrow \text{278} \\
 & \frac{(2-m) \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(cx d1 + d1)^{3/2} (d2 - cd2x)^{3/2}} dx}{3d1d2} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3d1d2 f (cd1x + d1)^{3/2} (d2 - cd2x)^{3/2}} + \\
 & \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{3d1^2 d2^2 f^2 (m+2) \sqrt{cd1x+d1} \sqrt{d2-cd2x}} \\
 & \quad \downarrow \text{6352} \\
 & (2-m) \left( -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{\sqrt{cx d1 + d1} \sqrt{d2 - cd2x}} dx}{d1d2} - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{d1d2 f \sqrt{cd1x+d1} \sqrt{d2-cd2x}} + \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{d1d2 f \sqrt{cd1x+d1} \sqrt{d2-cd2x}} \right) + \\
 & \quad \frac{(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{3d1d2} + \\
 & \quad \frac{bc\sqrt{cx-1}\sqrt{cx+1} (fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{3d1^2 d2^2 f^2 (m+2) \sqrt{cd1x+d1} \sqrt{d2-cd2x}} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

---

3.162.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{5/2} (d2 - cd2x)^{5/2}} dx$

$$(2-m) \left( -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx} d1 + d1 \sqrt{d2 - cd2x}}}{d1d2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{(1-cx)(cx+1)} dx}{d1d2f\sqrt{cd1x+d1}\sqrt{d2-cd2x}} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{d1d2f\sqrt{cd1x+d1}\sqrt{d2-cd2x}} \right) +$$

$$\frac{3d1d2}{(fx)^{m+1}(a + \operatorname{barccosh}(cx))} + \frac{3d1d2f(cd1x + d1)^{3/2}(d2 - cd2x)^{3/2} + bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{3d1^2d2^2f^2(m+2)\sqrt{cd1x+d1}\sqrt{d2-cd2x}}$$

↓ 82

$$(2-m) \left( -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx} d1 + d1 \sqrt{d2 - cd2x}}}{d1d2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{(fx)^{m+1}}{1-c^2x^2} dx}{d1d2f\sqrt{cd1x+d1}\sqrt{d2-cd2x}} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{d1d2f\sqrt{cd1x+d1}\sqrt{d2-cd2x}} \right) +$$

$$\frac{3d1d2}{(fx)^{m+1}(a + \operatorname{barccosh}(cx))} + \frac{3d1d2f(cd1x + d1)^{3/2}(d2 - cd2x)^{3/2} + bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{3d1^2d2^2f^2(m+2)\sqrt{cd1x+d1}\sqrt{d2-cd2x}}$$

↓ 278

$$(2-m) \left( -\frac{m \int \frac{(fx)^m (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx} d1 + d1 \sqrt{d2 - cd2x}}}{d1d2} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{d1d2f\sqrt{cd1x+d1}\sqrt{d2-cd2x}} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(1, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{d1d2f^2(m+2)\sqrt{cd1x+d1}\sqrt{d2-cd2x}} \right) +$$

$$\frac{3d1d2}{(fx)^{m+1}(a + \operatorname{barccosh}(cx))} + \frac{3d1d2f(cd1x + d1)^{3/2}(d2 - cd2x)^{3/2} + bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{3d1^2d2^2f^2(m+2)\sqrt{cd1x+d1}\sqrt{d2-cd2x}}$$

↓ 6364

$$(2-m) \left( -\frac{m \left( \frac{bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2}+1, \frac{m}{2}+1; \frac{m}{2}+\frac{3}{2}, \frac{m}{2}+2; c^2x^2\right)}{f^2(m+1)(m+2)\sqrt{cd1x+d1}\sqrt{d2-cd2x}} + \frac{\sqrt{1-c^2x^2}(fx)^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, c^2x^2\right)(a + \operatorname{barccosh}(cx))}{f(m+1)\sqrt{cd1x+d1}\sqrt{d2-cd2x}} \right)}{d1d2} + \frac{(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{d1d2f\sqrt{cd1x+d1}\sqrt{d2-cd2x}} \right) +$$

$$\frac{3d1d2}{(fx)^{m+1}(a + \operatorname{barccosh}(cx))} + \frac{3d1d2f(cd1x + d1)^{3/2}(d2 - cd2x)^{3/2} + bc\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+2} \operatorname{Hypergeometric2F1}\left(2, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{3d1^2d2^2f^2(m+2)\sqrt{cd1x+d1}\sqrt{d2-cd2x}}$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/((d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2)),x]`

output `((f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(3*d1*d2*f*(d1 + c*d1*x)^(3/2)*(d2 - c*d2*x)^(3/2)) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[2, (2 + m)/2, (4 + m)/2, c^2*x^2])/(3*d1^2*d2^2*f^2*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) + ((2 - m)*((f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(d1*d2*f*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Hypergeometric2F1[1, (2 + m)/2, (4 + m)/2, c^2*x^2])/(d1*d2*f^2*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) - (m*((f*x)^(1 + m)*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2])/(f*(1 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]) + (b*c*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2])/(f^2*(1 + m)*(2 + m)*Sqrt[d1 + c*d1*x]*Sqrt[d2 - c*d2*x]))/(d1*d2)))/(3*d1*d2)`

### 3.162.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`



rule 6352 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d1*d2*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d1*d2*(p + 1)) Int[(f*x)^m*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || EqQ[n, 1])`

rule 6364 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(((f*x)^(m + 1)/(f*(m + 1)))*Simp[Sqrt[1 - c^2*x^2]/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])]*(a + b*ArcCosh[c*x])*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, c^2*x^2], x] + Simp[b*c*((f*x)^(m + 2)/(f^2*(m + 1)*(m + 2)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*HypergeometricPFQ[{1, 1 + m/2, 1 + m/2}, {3/2 + m/2, 2 + m/2}, c^2*x^2], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && !IntegerQ[m]`

### 3.162.4 Maple [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(cd_1x + d_1)^{5/2} (-cd_2x + d_2)^{5/2}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(5/2)/(-c*d2*x+d2)^(5/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(5/2)/(-c*d2*x+d2)^(5/2),x)`

### 3.162.5 Fracas [F]

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{(cd_1x + d_1)^{5/2} (-cd_2x + d_2)^{5/2}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(5/2)/(-c*d2*x+d2)^(5/2), x, algorithm="fricas")`

---

3.162.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2}} dx$

output `integral(-sqrt(c*d1*x + d1)*sqrt(-c*d2*x + d2)*(b*arccosh(c*x) + a)*(f*x)^m/(c^6*d1^3*d2^3*x^6 - 3*c^4*d1^3*d2^3*x^4 + 3*c^2*d1^3*d2^3*x^2 - d1^3*d2^3), x)`

### 3.162.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{5/2}(d2 - cd2x)^{5/2}} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(c*d1*x+d1)**(5/2)/(-c*d2*x+d2)**(5/2),x)`

output `Timed out`

### 3.162.7 Maxima [F]

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{5/2}(d2 - cd2x)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{(cd1x + d1)^{\frac{5}{2}}(-cd2x + d2)^{\frac{5}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(5/2)/(-c*d2*x+d2)^(5/2), x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/((c*d1*x + d1)^(5/2)*(-c*d2*x + d2)^(5/2)), x)`

### 3.162.8 Giac [F]

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{5/2}(d2 - cd2x)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)(fx)^m}{(cd1x + d1)^{\frac{5}{2}}(-cd2x + d2)^{\frac{5}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(c*d1*x+d1)^(5/2)/(-c*d2*x+d2)^(5/2), x, algorithm="giac")`

---

3.162.  $\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d1 + cd1x)^{5/2}(d2 - cd2x)^{5/2}} dx$

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/((c*d1*x + d1)^(5/2)*(-c*d2*x + d2)^(5/2)), x)`

### 3.162.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{(d_1 + cd_1x)^{5/2} (d_2 - cd_2x)^{5/2}} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/((d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2)), x)`

output `int(((a + b*acosh(c*x))*(f*x)^m)/((d1 + c*d1*x)^(5/2)*(d2 - c*d2*x)^(5/2)), x)`

### 3.163 $\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$

3.163.1 Optimal result . . . . .	1355
3.163.2 Mathematica [A] (verified) . . . . .	1355
3.163.3 Rubi [A] (verified) . . . . .	1356
3.163.4 Maple [F] . . . . .	1357
3.163.5 Fricas [F] . . . . .	1357
3.163.6 Sympy [F] . . . . .	1357
3.163.7 Maxima [F] . . . . .	1358
3.163.8 Giac [F] . . . . .	1358
3.163.9 Mupad [F(-1)] . . . . .	1358

#### 3.163.1 Optimal result

Integrand size = 24, antiderivative size = 128

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{(fx)^{1+m} \operatorname{arccosh}(ax) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, a^2x^2\right)}{f(1+m)} + \frac{a(fx)^{2+m} \sqrt{-1+ax} {}_3F_2\left(1, 1+\frac{m}{2}, 1+\frac{m}{2}; \frac{3}{2}+\frac{m}{2}, 2+\frac{m}{2}; a^2x^2\right)}{f^2(1+m)(2+m)\sqrt{1-ax}}$$

output `(f*x)^(1+m)*arccosh(a*x)*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], a^2*x^2)/f/(1+m)+a*(f*x)^(2+m)*hypergeom([1, 1+1/2*m, 1+1/2*m], [2+1/2*m, 3/2+1/2*m], a^2*x^2)*(a*x-1)^(1/2)/f^2/(1+m)/(2+m)/(-a*x+1)^(1/2)`

#### 3.163.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.97

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \frac{x(fx)^m \left( \operatorname{arccosh}(ax) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+m}{2}, \frac{3+m}{2}, a^2x^2\right) + \frac{ax\sqrt{-1+ax}\sqrt{1+ax} {}_3F_2\left(1, 1+\frac{m}{2}, 1+\frac{m}{2}; \frac{3}{2}+\frac{m}{2}, 2+\frac{m}{2}; a^2x^2\right)}{(2+m)\sqrt{1-a^2x^2}} \right)}{1+m}$$

input `Integrate[((f*x)^m*ArcCosh[a*x])/Sqrt[1 - a^2*x^2], x]`

output  $(x*(f*x)^m*(\text{ArcCosh}[a*x]*\text{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, a^2*x^2] + (a*x*\text{Sqrt}[-1+a*x]*\text{Sqrt}[1+a*x]*\text{HypergeometricPFQ}[\{1, 1+m/2, 1+m/2\}, \{3/2+m/2, 2+m/2\}, a^2*x^2]))/((2+m)*\text{Sqrt}[1-a^2*x^2]))/(1+m)$

### 3.163.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6363}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\text{arccosh}(ax)(fx)^m}{\sqrt{1-a^2x^2}} dx$$

↓ 6363

$$\frac{a\sqrt{ax-1}(fx)^{m+2} {}_3F_2\left(1, \frac{m}{2}+1, \frac{m}{2}+1; \frac{m}{2}+\frac{3}{2}, \frac{m}{2}+2; a^2x^2\right)}{f^2(m+1)(m+2)\sqrt{1-ax}} + \frac{\text{arccosh}(ax)(fx)^{m+1} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+1}{2}, \frac{m+3}{2}, a^2x^2\right)}{f(m+1)}$$

input  $\text{Int}[(f*x)^m*\text{ArcCosh}[a*x]/\text{Sqrt}[1-a^2*x^2], x]$

output  $((f*x)^{(1+m)*\text{ArcCosh}[a*x]*\text{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, a^2*x^2])/(f*(1+m)) + (a*(f*x)^{(2+m)*\text{Sqrt}[-1+a*x]*\text{HypergeometricPFQ}[\{1, 1+m/2, 1+m/2\}, \{3/2+m/2, 2+m/2\}, a^2*x^2])/(f^2*(1+m)*(2+m)*\text{Sqrt}[1-a*x])$

#### 3.163.3.1 Defintions of rubi rules used

rule 6363  $\text{Int}[(a + \text{ArcCosh}[c*x])*(f*x)^m/\text{Sqrt}[d + e*x^2], x\_Symbol] := \text{Simp}[(f*x)^{(m+1)}/(f*(m+1))]*\text{Simp}[\text{Sqrt}[1-c^2*x^2]/\text{Sqrt}[d + e*x^2]]*(a + b*\text{ArcCosh}[c*x])* \text{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, c^2*x^2], x] + \text{Simp}[b*c*(f*x)^{(m+2)}/(f^2*(m+1)*(m+2))]*\text{Simp}[\text{Sqrt}[1+c*x]*(\text{Sqrt}[-1+c*x]/\text{Sqrt}[d + e*x^2])]*\text{HypergeometricPFQ}[\{1, 1+m/2, 1+m/2\}, \{3/2+m/2, 2+m/2\}, c^2*x^2], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{EqQ}[c^2*d + e, 0] \&\& !\text{IntegerQ}[m]$

$$3.163. \int \frac{(fx)^m \text{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx$$

**3.163.4 Maple [F]**

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{-a^2x^2 + 1}} dx$$

input `int((f*x)^m*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

output `int((f*x)^m*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

**3.163.5 Fracas [F]**

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1 - a^2x^2}} dx = \int \frac{(fx)^m \operatorname{arcosh}(ax)}{\sqrt{-a^2x^2 + 1}} dx$$

input `integrate((f*x)^m*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-a^2*x^2 + 1)*(f*x)^m*arccosh(a*x)/(a^2*x^2 - 1), x)`

**3.163.6 Sympy [F]**

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1 - a^2x^2}} dx = \int \frac{(fx)^m \operatorname{acosh}(ax)}{\sqrt{-(ax - 1)(ax + 1)}} dx$$

input `integrate((f*x)**m*acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral((f*x)**m*acosh(a*x)/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.163.7 Maxima [F]**

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{(fx)^m \operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate((f*x)^m*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((f*x)^m*arccosh(a*x)/sqrt(-a^2*x^2 + 1), x)`

**3.163.8 Giac [F]**

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{(fx)^m \operatorname{arcosh}(ax)}{\sqrt{-a^2x^2+1}} dx$$

input `integrate((f*x)^m*arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((f*x)^m*arccosh(a*x)/sqrt(-a^2*x^2 + 1), x)`

**3.163.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(fx)^m \operatorname{arccosh}(ax)}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax) (fx)^m}{\sqrt{1-a^2x^2}} dx$$

input `int((acosh(a*x)*(f*x)^m)/(1 - a^2*x^2)^(1/2),x)`

output `int((acosh(a*x)*(f*x)^m)/(1 - a^2*x^2)^(1/2), x)`

### 3.164 $\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^2 dx$

3.164.1 Optimal result . . . . .	1359
3.164.2 Mathematica [A] (verified) . . . . .	1360
3.164.3 Rubi [A] (verified) . . . . .	1360
3.164.4 Maple [A] (verified) . . . . .	1364
3.164.5 Fricas [A] (verification not implemented) . . . . .	1365
3.164.6 Sympy [F] . . . . .	1365
3.164.7 Maxima [A] (verification not implemented) . . . . .	1366
3.164.8 Giac [F(-2)] . . . . .	1366
3.164.9 Mupad [F(-1)] . . . . .	1367

#### 3.164.1 Optimal result

Integrand size = 20, antiderivative size = 266

$$\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^2 dx = \frac{4322c^3x}{3675} - \frac{1514a^2c^3x^3}{11025} + \frac{234a^4c^3x^5}{6125} - \frac{2}{343}a^6c^3x^7$$

$$- \frac{32c^3\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{35a} + \frac{16c^3(-1+ax)^{3/2}(1+ax)^{3/2}\operatorname{arccosh}(ax)}{105a}$$

$$- \frac{12c^3(-1+ax)^{5/2}(1+ax)^{5/2}\operatorname{arccosh}(ax)}{175a} + \frac{2c^3(-1+ax)^{7/2}(1+ax)^{7/2}\operatorname{arccosh}(ax)}{49a}$$

$$+ \frac{16}{35}c^3x\operatorname{arccosh}(ax)^2 + \frac{8}{35}c^3x(1-a^2x^2)\operatorname{arccosh}(ax)^2 + \frac{6}{35}c^3x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2 + \frac{1}{7}c^3x(1-a^2x^2)^3\operatorname{arccosh}(ax)^2$$

```
output 4322/3675*c^3*x-1514/11025*a^2*c^3*x^3+234/6125*a^4*c^3*x^5-2/343*a^6*c^3*x^7+16/105*c^3*(a*x-1)^(3/2)*(a*x+1)^(3/2)*arccosh(a*x)/a-12/175*c^3*(a*x-1)^(5/2)*(a*x+1)^(5/2)*arccosh(a*x)/a+2/49*c^3*(a*x-1)^(7/2)*(a*x+1)^(7/2)*arccosh(a*x)/a+16/35*c^3*x*arccosh(a*x)^2+8/35*c^3*x*(-a^2*x^2+1)*arccosh(a*x)^2+6/35*c^3*x*(-a^2*x^2+1)^2*arccosh(a*x)^2+1/7*c^3*x*(-a^2*x^2+1)^3*arccosh(a*x)^2-32/35*c^3*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a
```



**3.164.2 Mathematica [A] (verified)**

Time = 0.19 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.47

$$\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^2 dx$$

$$= \frac{c^3(453810ax - 52990a^3x^3 + 14742a^5x^5 - 2250a^7x^7 + 210\sqrt{-1+ax}\sqrt{1+ax}(-2161 + 757a^2x^2 - 351a^4x^4 + 75a^6x^6) \operatorname{ArcCosh}[ax] - 11025a^3x^3(-35 + 35a^2x^2 - 21a^4x^4 + 5a^6x^6) \operatorname{ArcCosh}[ax]^2)}{385875a}$$

input `Integrate[(c - a^2*c*x^2)^3*ArcCosh[a*x]^2,x]`output `(c^3*(453810*a*x - 52990*a^3*x^3 + 14742*a^5*x^5 - 2250*a^7*x^7 + 210*sqrt[-1 + a*x]*sqrt[1 + a*x]*(-2161 + 757*a^2*x^2 - 351*a^4*x^4 + 75*a^6*x^6)*ArcCosh[a*x] - 11025*a*x*(-35 + 35*a^2*x^2 - 21*a^4*x^4 + 5*a^6*x^6)*ArcCosh[a*x]^2))/(385875*a)`**3.164.3 Rubi [A] (verified)**Time = 2.14 (sec) , antiderivative size = 326, normalized size of antiderivative = 1.23, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$ , Rules used = {6312, 27, 6312, 6312, 6294, 6330, 24, 25, 39, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(ax)^2 (c - a^2cx^2)^3 dx$$

$$\downarrow \text{6312}$$

$$\frac{6}{7}c \int c^2(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^2 dx + \frac{2}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax) dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^2$$

$$\downarrow \text{27}$$

$$\frac{6}{7}c^3 \int (1 - a^2x^2)^2 \operatorname{arccosh}(ax)^2 dx + \frac{2}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax) dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^2$$

$$\downarrow \text{6312}$$

$$\frac{6}{7}c^3 \left( \frac{4}{5} \int (1 - a^2x^2) \operatorname{arccosh}(ax)^2 dx - \frac{2}{5}a \int x(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax) dx + \frac{1}{5}x(1 - a^2x^2)^2 \operatorname{arccosh}(ax) \right. \\ \left. + \frac{2}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax) dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right)$$

↓ 6312

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3}a \int x\sqrt{ax - 1}\sqrt{ax + 1} \operatorname{arccosh}(ax) dx + \frac{2}{3} \int \operatorname{arccosh}(ax)^2 dx + \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^2 \right) - \frac{2}{5}a \int x \right. \\ \left. + \frac{2}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax) dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right)$$

↓ 6294

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^2 - 2a \int \frac{x \operatorname{arccosh}(ax)}{\sqrt{ax - 1}\sqrt{ax + 1}} dx \right) + \frac{2}{3}a \int x\sqrt{ax - 1}\sqrt{ax + 1} \operatorname{arccosh}(ax) dx + \frac{1}{3}x(1 - a^2x^2) \right. \right. \\ \left. \left. + \frac{2}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax) dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right) \right)$$

↓ 6330

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax - 1}\sqrt{ax + 1} \operatorname{arccosh}(ax)}{a^2} - \frac{\int 1 dx}{a} \right) \right) + \frac{2}{3}a \left( \frac{(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right. \right. \\ \left. \left. + \frac{2}{7}ac^3 \left( \frac{(ax - 1)^{7/2}(ax + 1)^{7/2} \operatorname{arccosh}(ax)}{7a^2} - \frac{\int -(1 - ax)^3(ax + 1)^3 dx}{7a} \right) + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right) \right)$$

↓ 24

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3}a \left( \frac{(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} - \frac{\int -((1 - ax)(ax + 1)) dx}{3a} \right) + \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3} \right. \right. \\ \left. \left. + \frac{2}{7}ac^3 \left( \frac{(ax - 1)^{7/2}(ax + 1)^{7/2} \operatorname{arccosh}(ax)}{7a^2} - \frac{\int -(1 - ax)^3(ax + 1)^3 dx}{7a} \right) + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right) \right)$$

↓ 25

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3}a \left( \frac{\int (1-ax)(ax+1)dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( \frac{2}{7}ac^3 \left( \frac{\int (1-ax)^3(ax+1)^3 dx}{7a} + \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)}{7a^2} \right) + \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right) \right)$$

↓ 39

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3}a \left( \frac{\int (1-a^2x^2) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( \frac{2}{7}ac^3 \left( \frac{\int (1-a^2x^2)^3 dx}{7a} + \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)}{7a^2} \right) + \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right) \right)$$

↓ 210

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3}a \left( \frac{\int (1-a^2x^2) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( \frac{2}{7}ac^3 \left( \frac{\int (-a^6x^6 + 3a^4x^4 - 3a^2x^2 + 1) dx}{7a} + \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)}{7a^2} \right) + \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2 \right) \right)$$

↓ 2009

$$\frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2 + \frac{6}{7}c^3 \left( \frac{1}{5}x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^2 + \frac{4}{5} \left( \frac{2}{3}a \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} + \frac{x - \frac{a^2x^3}{3}}{3a} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{7}ac^3 \left( \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)}{7a^2} + \frac{-\frac{1}{7}a^6x^7 + \frac{3a^4x^5}{5} - a^2x^3 + x}{7a} \right) \right)$$

input `Int[(c - a^2*c*x^2)^3*ArcCosh[a*x]^2,x]`

```
output (c^3*x*(1 - a^2*x^2)^3*ArcCosh[a*x]^2)/7 + (2*a*c^3*((x - a^2*x^3 + (3*a^4
*x^5)/5 - (a^6*x^7)/7)/(7*a) + ((-1 + a*x)^(7/2)*(1 + a*x)^(7/2)*ArcCosh[a
*x])/(7*a^2)))/7 + (6*c^3*((x*(1 - a^2*x^2)^2*ArcCosh[a*x]^2)/5 - (2*a*(-1
/5*(x - (2*a^2*x^3)/3 + (a^4*x^5)/5)/a + ((-1 + a*x)^(5/2)*(1 + a*x)^(5/2)
*ArcCosh[a*x])/(5*a^2)))/5 + (4*((x*(1 - a^2*x^2)*ArcCosh[a*x]^2)/3 + (2*a
*((x - (a^2*x^3)/3)/(3*a) + ((-1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x]
)/(3*a^2)))/3 + (2*(x*ArcCosh[a*x]^2 - 2*a*(-(x/a) + (Sqrt[-1 + a*x]*Sqrt[
1 + a*x]*ArcCosh[a*x])/a^2)))/3))/5))/7
```

### 3.164.3.1 Defintions of rubi rules used

```
rule 24 Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 39 Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(m_.), x_Symbol] := Int[(
a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (
IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))
```

```
rule 210 Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2
)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6294 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*A
rcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt
[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]
```

```
rule 6312 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),
x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] +
(Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x
], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p
)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n
- 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n,
0] && GtQ[p, 0]
```

```
rule 6330 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p
_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 +
e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2
*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^
p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n -
1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && E
qQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]
```

### 3.164.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.71

method	result
derivativedivides	$-\frac{c^3 \left( 55125 \operatorname{arccosh}(ax)^2 a^7 x^7 - 15750 \operatorname{arccosh}(ax) \sqrt{ax-1} \sqrt{ax+1} a^6 x^6 - 231525 a^5 x^5 \operatorname{arccosh}(ax)^2 + 73710 a^4 x^4 \operatorname{arccosh}(ax) \right)}{\dots}$
default	$-\frac{c^3 \left( 55125 \operatorname{arccosh}(ax)^2 a^7 x^7 - 15750 \operatorname{arccosh}(ax) \sqrt{ax-1} \sqrt{ax+1} a^6 x^6 - 231525 a^5 x^5 \operatorname{arccosh}(ax)^2 + 73710 a^4 x^4 \operatorname{arccosh}(ax) \right)}{\dots}$

```
input int((-a^2*c*x^2+c)^3*arccosh(a*x)^2,x,method=_RETURNVERBOSE)
```

```
output -1/385875/a*c^3*(55125*arccosh(a*x)^2*a^7*x^7-15750*arccosh(a*x)*(a*x-1)^(
1/2)*(a*x+1)^(1/2)*a^6*x^6-231525*a^5*x^5*arccosh(a*x)^2+73710*a^4*x^4*arc
cosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)+2250*a^7*x^7+385875*a^3*x^3*arccosh(
a*x)^2-158970*a^2*x^2*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)-14742*a^5*x
^5-385875*a*x*arccosh(a*x)^2+453810*(a*x-1)^(1/2)*(a*x+1)^(1/2)*arccosh(a*
x)+52990*a^3*x^3-453810*a*x)
```

**3.164.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.66

$$\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^2 dx = \frac{2250 a^7 c^3 x^7 - 14742 a^5 c^3 x^5 + 52990 a^3 c^3 x^3 - 453810 a c^3 x + 11025 (5 a^7 c^3 x^7 - 21 a^5 c^3 x^5 + 35 a^3 c^3 x^3 - 35 a c^3 x) \log(ax + \sqrt{a^2 x^2 - 1})^2 - 210 (75 a^6 c^3 x^6 - 351 a^4 c^3 x^4 + 757 a^2 c^3 x^2 - 2161 c^3) \sqrt{a^2 x^2 - 1} \log(ax + \sqrt{a^2 x^2 - 1})}{a}$$

input `integrate((-a^2*c*x^2+c)^3*arccosh(a*x)^2,x, algorithm="fricas")`output `-1/385875*(2250*a^7*c^3*x^7 - 14742*a^5*c^3*x^5 + 52990*a^3*c^3*x^3 - 453810*a*c^3*x + 11025*(5*a^7*c^3*x^7 - 21*a^5*c^3*x^5 + 35*a^3*c^3*x^3 - 35*a*c^3*x)*log(a*x + sqrt(a^2*x^2 - 1))^2 - 210*(75*a^6*c^3*x^6 - 351*a^4*c^3*x^4 + 757*a^2*c^3*x^2 - 2161*c^3)*sqrt(a^2*x^2 - 1)*log(a*x + sqrt(a^2*x^2 - 1)))/a`**3.164.6 Sympy [F]**

$$\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^2 dx = -c^3 \left( \int 3a^2 x^2 \operatorname{acosh}^2(ax) dx + \int (-3a^4 x^4 \operatorname{acosh}^2(ax)) dx + \int a^6 x^6 \operatorname{acosh}^2(ax) dx + \int (-\operatorname{acosh}^2(ax)) dx \right)$$

input `integrate((-a**2*c*x**2+c)**3*acosh(a*x)**2,x)`output `-c**3*(Integral(3*a**2*x**2*acosh(a*x)**2, x) + Integral(-3*a**4*x**4*acosh(a*x)**2, x) + Integral(a**6*x**6*acosh(a*x)**2, x) + Integral(-acosh(a*x)**2, x))`

**3.164.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.67

$$\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^2 dx = -\frac{2}{343} a^6 c^3 x^7 + \frac{234}{6125} a^4 c^3 x^5 - \frac{1514}{11025} a^2 c^3 x^3 + \frac{4322}{3675} c^3 x$$

$$+ \frac{2}{3675} \left( 75 \sqrt{a^2 x^2 - 1} a^4 c^3 x^6 - 351 \sqrt{a^2 x^2 - 1} a^2 c^3 x^4 + 757 \sqrt{a^2 x^2 - 1} c^3 x^2 - \frac{2161 \sqrt{a^2 x^2 - 1} c^3}{a^2} \right) a \operatorname{arccosh}(ax)$$

$$- \frac{1}{35} (5 a^6 c^3 x^7 - 21 a^4 c^3 x^5 + 35 a^2 c^3 x^3 - 35 c^3 x) \operatorname{arccosh}(ax)^2$$

```
input integrate((-a^2*c*x^2+c)^3*arccosh(a*x)^2,x, algorithm="maxima")
```

```
output -2/343*a^6*c^3*x^7 + 234/6125*a^4*c^3*x^5 - 1514/11025*a^2*c^3*x^3 + 4322/
3675*c^3*x + 2/3675*(75*sqrt(a^2*x^2 - 1)*a^4*c^3*x^6 - 351*sqrt(a^2*x^2 -
1)*a^2*c^3*x^4 + 757*sqrt(a^2*x^2 - 1)*c^3*x^2 - 2161*sqrt(a^2*x^2 - 1)*c
^3/a^2)*a*arccosh(a*x) - 1/35*(5*a^6*c^3*x^7 - 21*a^4*c^3*x^5 + 35*a^2*c^3
*x^3 - 35*c^3*x)*arccosh(a*x)^2
```

**3.164.8 Giac [F(-2)]**

Exception generated.

$$\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^2 dx = \text{Exception raised: TypeError}$$

```
input integrate((-a^2*c*x^2+c)^3*arccosh(a*x)^2,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.164.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^2 dx = \int \operatorname{acosh}(ax)^2 (c - a^2 cx^2)^3 dx$$

input `int(acosh(a*x)^2*(c - a^2*c*x^2)^3,x)`output `int(acosh(a*x)^2*(c - a^2*c*x^2)^3, x)`



### 3.165 $\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2 dx$

3.165.1 Optimal result . . . . .	1368
3.165.2 Mathematica [A] (verified) . . . . .	1369
3.165.3 Rubi [A] (verified) . . . . .	1369
3.165.4 Maple [A] (verified) . . . . .	1373
3.165.5 Fricas [A] (verification not implemented) . . . . .	1373
3.165.6 Sympy [F] . . . . .	1374
3.165.7 Maxima [A] (verification not implemented) . . . . .	1374
3.165.8 Giac [F(-2)] . . . . .	1375
3.165.9 Mupad [F(-1)] . . . . .	1375

#### 3.165.1 Optimal result

Integrand size = 20, antiderivative size = 195

$$\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2 dx$$

$$= \frac{298c^2x}{225} - \frac{76}{675}a^2c^2x^3 + \frac{2}{125}a^4c^2x^5 - \frac{16c^2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{15a}$$

$$+ \frac{8c^2(-1+ax)^{3/2}(1+ax)^{3/2}\operatorname{arccosh}(ax)}{45a} - \frac{2c^2(-1+ax)^{5/2}(1+ax)^{5/2}\operatorname{arccosh}(ax)}{25a}$$

$$+ \frac{8}{15}c^2x\operatorname{arccosh}(ax)^2 + \frac{4}{15}c^2x(1-a^2x^2)\operatorname{arccosh}(ax)^2 + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2$$

```
output 298/225*c^2*x-76/675*a^2*c^2*x^3+2/125*a^4*c^2*x^5+8/45*c^2*(a*x-1)^(3/2)*
(a*x+1)^(3/2)*arccosh(a*x)/a-2/25*c^2*(a*x-1)^(5/2)*(a*x+1)^(5/2)*arccosh(
a*x)/a+8/15*c^2*x*arccosh(a*x)^2+4/15*c^2*x*(-a^2*x^2+1)*arccosh(a*x)^2+1/
5*c^2*x*(-a^2*x^2+1)^2*arccosh(a*x)^2-16/15*c^2*arccosh(a*x)*(a*x-1)^(1/2)
*(a*x+1)^(1/2)/a
```

**3.165.2 Mathematica [A] (verified)**

Time = 0.25 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.52

$$\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2 dx$$

$$= \frac{c^2(4470ax - 380a^3x^3 + 54a^5x^5 - 30\sqrt{-1 + ax}\sqrt{1 + ax}(149 - 38a^2x^2 + 9a^4x^4) \operatorname{arccosh}(ax) + 225ax(15 - 10a^2x^2 + 3a^4x^4) \operatorname{arccosh}(ax)^2)}{3375a}$$

input `Integrate[(c - a^2*c*x^2)^2*ArcCosh[a*x]^2,x]`

output `(c^2*(4470*a*x - 380*a^3*x^3 + 54*a^5*x^5 - 30*Sqrt[-1 + a*x]*Sqrt[1 + a*x])*(149 - 38*a^2*x^2 + 9*a^4*x^4)*ArcCosh[a*x] + 225*a*x*(15 - 10*a^2*x^2 + 3*a^4*x^4)*ArcCosh[a*x]^2)/(3375*a)`

**3.165.3 Rubi [A] (verified)**

Time = 1.32 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.16, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6312, 27, 6312, 6294, 6330, 24, 25, 39, 210, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(ax)^2 (c - a^2cx^2)^2 dx$$

$$\downarrow \text{6312}$$

$$\frac{4}{5}c \int c(1 - a^2x^2) \operatorname{arccosh}(ax)^2 dx - \frac{2}{5}ac^2 \int x(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax) dx + \frac{1}{5}c^2x(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^2$$

$$\downarrow \text{27}$$

$$\frac{4}{5}c^2 \int (1 - a^2x^2) \operatorname{arccosh}(ax)^2 dx - \frac{2}{5}ac^2 \int x(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax) dx + \frac{1}{5}c^2x(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^2$$

$$\downarrow \text{6312}$$

$$\frac{4}{5}c^2 \left( \frac{2}{3}a \int x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)dx + \frac{2}{3} \int \operatorname{arccosh}(ax)^2 dx + \frac{1}{3}x(1-a^2x^2)\operatorname{arccosh}(ax)^2 \right) - \frac{2}{5}ac^2 \int x(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)dx + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2$$

↓ 6294

$$\frac{4}{5}c^2 \left( \frac{2}{3} \left( x\operatorname{arccosh}(ax)^2 - 2a \int \frac{x\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right) + \frac{2}{3}a \int x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)dx + \frac{1}{3}x(1-a^2x^2)\operatorname{arccosh}(ax)^2 \right) - \frac{2}{5}ac^2 \int x(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)dx + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2$$

↓ 6330

$$\frac{4}{5}c^2 \left( \frac{2}{3} \left( x\operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{a^2} - \frac{\int 1dx}{a} \right) \right) + \frac{2}{3}a \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{3a^2} \right) \right) - \frac{2}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)}{5a^2} - \frac{\int (1-ax)^2(ax+1)^2 dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2$$

↓ 24

$$\frac{4}{5}c^2 \left( \frac{2}{3}a \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{3a^2} - \frac{\int -((1-ax)(ax+1))dx}{3a} \right) + \frac{1}{3}x(1-a^2x^2)\operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( x\operatorname{arccosh}(ax)^2 - 2a \int \frac{x\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right) \right) - \frac{2}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)}{5a^2} - \frac{\int (1-ax)^2(ax+1)^2 dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2$$

↓ 25

$$\frac{4}{5}c^2 \left( \frac{2}{3}a \left( \frac{\int (1-ax)(ax+1)dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2)\operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( x\operatorname{arccosh}(ax)^2 - 2a \int \frac{x\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right) \right) - \frac{2}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)}{5a^2} - \frac{\int (1-ax)^2(ax+1)^2 dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2$$

↓ 39

$$\frac{4}{5}c^2 \left( \frac{2}{3}a \left( \frac{\int (1 - a^2x^2) dx}{3a} + \frac{(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) + \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( x \operatorname{arccosh}(ax) \right) \right. \\ \left. + \frac{2}{5}ac^2 \left( \frac{(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)}{5a^2} - \frac{\int (1 - a^2x^2)^2 dx}{5a} \right) + \frac{1}{5}c^2x(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^2 \right)$$

↓ 210

$$\frac{4}{5}c^2 \left( \frac{2}{3}a \left( \frac{\int (1 - a^2x^2) dx}{3a} + \frac{(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) + \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( x \operatorname{arccosh}(ax) \right) \right. \\ \left. + \frac{2}{5}ac^2 \left( \frac{(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)}{5a^2} - \frac{\int (a^4x^4 - 2a^2x^2 + 1) dx}{5a} \right) + \frac{1}{5}c^2x(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^2 \right)$$

↓ 2009

$$\frac{1}{5}c^2x(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^2 + \\ \frac{4}{5}c^2 \left( \frac{2}{3}a \left( \frac{(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} + \frac{x - \frac{a^2x^3}{3}}{3a} \right) + \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3} \left( x \operatorname{arccosh}(ax) \right) \right. \\ \left. + \frac{2}{5}ac^2 \left( \frac{(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)}{5a^2} - \frac{\frac{a^4x^5}{5} - \frac{2a^2x^3}{3} + x}{5a} \right) \right)$$

input `Int[(c - a^2*c*x^2)^2*ArcCosh[a*x]^2,x]`

output `(c^2*x*(1 - a^2*x^2)^2*ArcCosh[a*x]^2)/5 - (2*a*c^2*(-1/5*(x - (2*a^2*x^3)/3 + (a^4*x^5)/5)/a + ((-1 + a*x)^(5/2)*(1 + a*x)^(5/2)*ArcCosh[a*x])/(5*a^2)))/5 + (4*c^2*((x*(1 - a^2*x^2)*ArcCosh[a*x]^2)/3 + (2*a*((x - (a^2*x^3)/3)/(3*a) + ((-1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x])/(3*a^2))))/3 + (2*(x*ArcCosh[a*x]^2 - 2*a*(-(x/a) + (Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/a^2)))/3)/5`

## 3.165.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 39 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 210 `Int[((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*x^2)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6294 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`
- rule 6312 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

```
rule 6330 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p
_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 +
e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2
*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^
p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n -
1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && E
qQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]
```

### 3.165.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.72

method	result
derivativedivides	$\frac{c^2 \left( 675a^5 x^5 \operatorname{arccosh}(ax)^2 - 270a^4 x^4 \operatorname{arccosh}(ax) \sqrt{ax-1} \sqrt{ax+1} - 2250a^3 x^3 \operatorname{arccosh}(ax)^2 + 1140a^2 x^2 \operatorname{arccosh}(ax) \sqrt{ax-1} \sqrt{ax+1} - 380a^3 x^3 + 4470a^2 x \right)}{3375a}$
default	$\frac{c^2 \left( 675a^5 x^5 \operatorname{arccosh}(ax)^2 - 270a^4 x^4 \operatorname{arccosh}(ax) \sqrt{ax-1} \sqrt{ax+1} - 2250a^3 x^3 \operatorname{arccosh}(ax)^2 + 1140a^2 x^2 \operatorname{arccosh}(ax) \sqrt{ax-1} \sqrt{ax+1} - 380a^3 x^3 + 4470a^2 x \right)}{3375a}$

```
input int((-a^2*c*x^2+c)^2*arccosh(a*x)^2,x,method=_RETURNVERBOSE)
```

```
output 1/3375/a*c^2*(675*a^5*x^5*arccosh(a*x)^2-270*a^4*x^4*arccosh(a*x)*(a*x-1)^(
(1/2)*(a*x+1)^(1/2)-2250*a^3*x^3*arccosh(a*x)^2+1140*a^2*x^2*arccosh(a*x)*
(a*x-1)^(1/2)*(a*x+1)^(1/2)+54*a^5*x^5+3375*a*x*arccosh(a*x)^2-4470*(a*x-1
)^(1/2)*(a*x+1)^(1/2)*arccosh(a*x)-380*a^3*x^3+4470*a*x)
```

### 3.165.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.73

$$\int (c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^2 dx$$

$$= \frac{54 a^5 c^2 x^5 - 380 a^3 c^2 x^3 + 4470 a c^2 x + 225 (3 a^5 c^2 x^5 - 10 a^3 c^2 x^3 + 15 a c^2 x) \log(ax + \sqrt{a^2 x^2 - 1})^2 - 30 (9 a^5 c^2 x^5 - 380 a^3 c^2 x^3 + 4470 a c^2 x)}{3375 a}$$

```
input integrate((-a^2*c*x^2+c)^2*arccosh(a*x)^2,x, algorithm="fricas")
```

output  $1/3375*(54*a^5*c^2*x^5 - 380*a^3*c^2*x^3 + 4470*a*c^2*x + 225*(3*a^5*c^2*x^5 - 10*a^3*c^2*x^3 + 15*a*c^2*x)*\log(a*x + \sqrt{a^2*x^2 - 1})^2 - 30*(9*a^4*c^2*x^4 - 38*a^2*c^2*x^2 + 149*c^2)*\sqrt{a^2*x^2 - 1}*\log(a*x + \sqrt{a^2*x^2 - 1}))/a$

### 3.165.6 Sympy [F]

$$\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2 dx = c^2 \left( \int (-2a^2x^2 \operatorname{acosh}^2(ax)) dx + \int a^4x^4 \operatorname{acosh}^2(ax) dx + \int \operatorname{acosh}^2(ax) dx \right)$$

input `integrate((-a**2*c*x**2+c)**2*acosh(a*x)**2,x)`

output `c**2*(Integral(-2*a**2*x**2*acosh(a*x)**2, x) + Integral(a**4*x**4*acosh(a*x)**2, x) + Integral(acosh(a*x)**2, x))`

### 3.165.7 Maxima [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.69

$$\begin{aligned} & \int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2 dx \\ &= \frac{2}{125} a^4c^2x^5 - \frac{76}{675} a^2c^2x^3 + \frac{298}{225} c^2x \\ & \quad - \frac{2}{225} \left( 9\sqrt{a^2x^2 - 1}a^2c^2x^4 - 38\sqrt{a^2x^2 - 1}c^2x^2 + \frac{149\sqrt{a^2x^2 - 1}c^2}{a^2} \right) a \operatorname{arccosh}(ax) \\ & \quad + \frac{1}{15} (3a^4c^2x^5 - 10a^2c^2x^3 + 15c^2x) \operatorname{arccosh}(ax)^2 \end{aligned}$$

input `integrate((-a^2*c*x^2+c)^2*arccosh(a*x)^2,x, algorithm="maxima")`

output  $2/125*a^4*c^2*x^5 - 76/675*a^2*c^2*x^3 + 298/225*c^2*x - 2/225*(9*\sqrt{a^2*x^2 - 1}*a^2*c^2*x^4 - 38*\sqrt{a^2*x^2 - 1}*c^2*x^2 + 149*\sqrt{a^2*x^2 - 1}*c^2/a^2)*a*\operatorname{arccosh}(a*x) + 1/15*(3*a^4*c^2*x^5 - 10*a^2*c^2*x^3 + 15*c^2*x)*\operatorname{arccosh}(a*x)^2$

**3.165.8 Giac [F(-2)]**

Exception generated.

$$\int (c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^2 dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^2*arccosh(a*x)^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.165.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^2 dx = \int \operatorname{acosh}(ax)^2 (c - a^2 cx^2)^2 dx$$

input `int(acosh(a*x)^2*(c - a^2*c*x^2)^2,x)`

output `int(acosh(a*x)^2*(c - a^2*c*x^2)^2, x)`



### 3.166 $\int (c - a^2cx^2) \operatorname{arccosh}(ax)^2 dx$

3.166.1 Optimal result . . . . .	1376
3.166.2 Mathematica [A] (verified) . . . . .	1376
3.166.3 Rubi [A] (verified) . . . . .	1377
3.166.4 Maple [A] (verified) . . . . .	1379
3.166.5 Fricas [A] (verification not implemented) . . . . .	1380
3.166.6 Sympy [F] . . . . .	1380
3.166.7 Maxima [A] (verification not implemented) . . . . .	1380
3.166.8 Giac [F(-2)] . . . . .	1381
3.166.9 Mupad [F(-1)] . . . . .	1381

#### 3.166.1 Optimal result

Integrand size = 18, antiderivative size = 112

$$\int (c - a^2cx^2) \operatorname{arccosh}(ax)^2 dx = \frac{14cx}{9} - \frac{2}{27}a^2cx^3 - \frac{4c\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{3a} \\ + \frac{2c(-1+ax)^{3/2}(1+ax)^{3/2}\operatorname{arccosh}(ax)}{9a} \\ + \frac{2}{3}cx\operatorname{arccosh}(ax)^2 + \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^2$$

output  $14/9*c*x-2/27*a^2*c*x^3+2/9*c*(a*x-1)^{(3/2)}*(a*x+1)^{(3/2)}*\operatorname{arccosh}(a*x)/a+2/3*c*x*\operatorname{arccosh}(a*x)^2+1/3*c*x*(-a^2*x^2+1)*\operatorname{arccosh}(a*x)^2-4/3*c*\operatorname{arccosh}(a*x)*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/a$

#### 3.166.2 Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.65

$$\int (c - a^2cx^2) \operatorname{arccosh}(ax)^2 dx \\ = \frac{c(42ax - 2a^3x^3 + 6\sqrt{-1+ax}\sqrt{1+ax}(-7 + a^2x^2) \operatorname{arccosh}(ax) - 9ax(-3 + a^2x^2) \operatorname{arccosh}(ax)^2)}{27a}$$

input `Integrate[(c - a^2*c*x^2)*ArcCosh[a*x]^2,x]`

output  $(c*(42*a*x - 2*a^3*x^3 + 6*\sqrt{-1 + a*x}*\sqrt{1 + a*x}*(-7 + a^2*x^2)*\text{ArcCosh}[a*x] - 9*a*x*(-3 + a^2*x^2)*\text{ArcCosh}[a*x]^2))/(27*a)$

### 3.166.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.14, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$ , Rules used = {6312, 6294, 6330, 24, 25, 39, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{arccosh}(ax)^2 (c - a^2 cx^2) dx \\
 & \quad \downarrow \text{6312} \\
 & \frac{2}{3}ac \int x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)dx + \frac{2}{3}c \int \operatorname{arccosh}(ax)^2 dx + \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^2 \\
 & \quad \downarrow \text{6294} \\
 & \frac{2}{3}c \left( x\operatorname{arccosh}(ax)^2 - 2a \int \frac{x\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right) + \frac{2}{3}ac \int x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)dx + \\
 & \quad \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^2 \\
 & \quad \downarrow \text{6330} \\
 & \frac{2}{3}c \left( x\operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{a^2} - \frac{\int 1 dx}{a} \right) \right) + \\
 & \frac{2}{3}ac \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{3a^2} - \frac{\int -((1-ax)(ax+1))dx}{3a} \right) + \\
 & \quad \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^2 \\
 & \quad \downarrow \text{24} \\
 & \frac{2}{3}ac \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{3a^2} - \frac{\int -((1-ax)(ax+1))dx}{3a} \right) + \\
 & \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^2 + \frac{2}{3}c \left( x\operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right) \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

$$\begin{aligned}
& \frac{2}{3}ac \left( \frac{\int (1-ax)(ax+1)dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) + \\
& \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^2 + \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right) \\
& \quad \downarrow \text{39} \\
& \frac{2}{3}ac \left( \frac{\int (1-a^2x^2) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) + \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^2 + \\
& \quad \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right) \\
& \quad \downarrow \text{2009} \\
& \frac{2}{3}ac \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} + \frac{x - \frac{a^2x^3}{3}}{3a} \right) + \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^2 + \\
& \quad \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right)
\end{aligned}$$

input `Int[(c - a^2*c*x^2)*ArcCosh[a*x]^2,x]`

output `(c*x*(1 - a^2*x^2)*ArcCosh[a*x]^2)/3 + (2*a*c*((x - (a^2*x^3)/3)/(3*a) + (-1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x])/(3*a^2))/3 + (2*c*(x*ArcCosh[a*x]^2 - 2*a*(-(x/a) + (Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/a^2)))/3`

### 3.166.3.1 Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 39 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(m_.), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

### 3.166.4 Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.80

method	result
derivativedivides	$-\frac{c(9a^3x^3 \operatorname{arccosh}(ax)^2 - 6a^2x^2 \operatorname{arccosh}(ax)\sqrt{ax-1}\sqrt{ax+1} - 27ax \operatorname{arccosh}(ax)^2 + 42\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) + 2a^3)}{27a}$
default	$-\frac{c(9a^3x^3 \operatorname{arccosh}(ax)^2 - 6a^2x^2 \operatorname{arccosh}(ax)\sqrt{ax-1}\sqrt{ax+1} - 27ax \operatorname{arccosh}(ax)^2 + 42\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) + 2a^3)}{27a}$

input `int((-a^2*c*x^2+c)*arccosh(a*x)^2,x,method=_RETURNVERBOSE)`

output `-1/27/a*c*(9*a^3*x^3*arccosh(a*x)^2-6*a^2*x^2*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)-27*a*x*arccosh(a*x)^2+42*(a*x-1)^(1/2)*(a*x+1)^(1/2)*arccosh(a*x)+2*a^3*x^3-42*a*x)`

**3.166.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.85

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^2 dx = \frac{2a^3 cx^3 - 42acx + 9(a^3 cx^3 - 3acx) \log(ax + \sqrt{a^2 x^2 - 1})^2 - 6(a^2 cx^2 - 7c)\sqrt{a^2 x^2 - 1} \log(ax + \sqrt{a^2 x^2 - 1})}{27a}$$

input `integrate((-a^2*c*x^2+c)*arccosh(a*x)^2,x, algorithm="fricas")`output `-1/27*(2*a^3*c*x^3 - 42*a*c*x + 9*(a^3*c*x^3 - 3*a*c*x)*log(a*x + sqrt(a^2*x^2 - 1))^2 - 6*(a^2*c*x^2 - 7*c)*sqrt(a^2*x^2 - 1)*log(a*x + sqrt(a^2*x^2 - 1)))/a`**3.166.6 Sympy [F]**

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^2 dx = -c \left( \int a^2 x^2 \operatorname{acosh}^2(ax) dx + \int (-\operatorname{acosh}^2(ax)) dx \right)$$

input `integrate((-a**2*c*x**2+c)*acosh(a*x)**2,x)`output `-c*(Integral(a**2*x**2*acosh(a*x)**2, x) + Integral(-acosh(a*x)**2, x))`**3.166.7 Maxima [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.68

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^2 dx = -\frac{2}{27} a^2 cx^3 + \frac{2}{9} \left( \sqrt{a^2 x^2 - 1} cx^2 - \frac{7\sqrt{a^2 x^2 - 1}c}{a^2} \right) a \operatorname{arccosh}(ax) - \frac{1}{3} (a^2 cx^3 - 3cx) \operatorname{arccosh}(ax)^2 + \frac{14}{9} cx$$

input `integrate((-a^2*c*x^2+c)*arccosh(a*x)^2,x, algorithm="maxima")`

output `-2/27*a^2*c*x^3 + 2/9*(sqrt(a^2*x^2 - 1)*c*x^2 - 7*sqrt(a^2*x^2 - 1)*c/a^2)*a*arccosh(a*x) - 1/3*(a^2*c*x^3 - 3*c*x)*arccosh(a*x)^2 + 14/9*c*x`

### 3.166.8 Giac [F(-2)]

Exception generated.

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^2 dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)*arccosh(a*x)^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.166.9 Mupad [F(-1)]

Timed out.

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^2 dx = \int \operatorname{acosh}(ax)^2 (c - a^2 cx^2) dx$$

input `int(acosh(a*x)^2*(c - a^2*c*x^2),x)`

output `int(acosh(a*x)^2*(c - a^2*c*x^2), x)`

### 3.167 $\int \frac{\operatorname{arccosh}(ax)^2}{c-a^2cx^2} dx$

3.167.1 Optimal result	1382
3.167.2 Mathematica [A] (verified)	1383
3.167.3 Rubi [C] (verified)	1383
3.167.4 Maple [A] (verified)	1385
3.167.5 Fricas [F]	1386
3.167.6 Sympy [F]	1386
3.167.7 Maxima [F]	1386
3.167.8 Giac [F]	1387
3.167.9 Mupad [F(-1)]	1387

#### 3.167.1 Optimal result

Integrand size = 20, antiderivative size = 98

$$\int \frac{\operatorname{arccosh}(ax)^2}{c-a^2cx^2} dx = \frac{2\operatorname{arccosh}(ax)^2 \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac} + \frac{2\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{ac} - \frac{2\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{ac} - \frac{2 \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)})}{ac} + \frac{2 \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)})}{ac}$$

```
output 2*arccosh(a*x)^2*arctanh(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c+2*arccosh(a*x)*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c-2*arccosh(a*x)*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c-2*polylog(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c+2*polylog(3,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c
```

**3.167.2 Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.97

$$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx$$

$$= \frac{-\operatorname{arccosh}(ax)^2 \log(1 - e^{\operatorname{arccosh}(ax)}) + \operatorname{arccosh}(ax)^2 \log(1 + e^{\operatorname{arccosh}(ax)}) + 2\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - 2\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) - 2\operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) + 2\operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)})}{a^2c}$$

input `Integrate[ArcCosh[a*x]^2/(c - a^2*c*x^2), x]`output `(-(ArcCosh[a*x]^2*Log[1 - E^ArcCosh[a*x]]) + ArcCosh[a*x]^2*Log[1 + E^ArcCosh[a*x]]) + 2*ArcCosh[a*x]*PolyLog[2, -E^ArcCosh[a*x]] - 2*ArcCosh[a*x]*PolyLog[2, E^ArcCosh[a*x]] - 2*PolyLog[3, -E^ArcCosh[a*x]] + 2*PolyLog[3, E^ArcCosh[a*x]])/(a*c)`**3.167.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.88, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$ , Rules used = {6318, 3042, 26, 4670, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx$$

$$\downarrow \text{6318}$$

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)$$

$$\frac{ac}{ac}$$

$$\downarrow \text{3042}$$

$$\int i \operatorname{arccosh}(ax)^2 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)$$

$$\frac{ac}{ac}$$

$$\downarrow \text{26}$$

$$\frac{i \int \operatorname{arccosh}(ax)^2 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{ac}$$

---

3.167.  $\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx$



↓ 4670

$$\frac{i(2i \int \operatorname{arccosh}(ax) \log(1 - e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - 2i \int \operatorname{arccosh}(ax) \log(1 + e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) + 2i \int \operatorname{arccosh}(ax) \log(1 - e^{-\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - 2i \int \operatorname{arccosh}(ax) \log(1 + e^{-\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax))}{ac}$$

↓ 3011

$$\frac{i(-2i(\int \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 2i(\int \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})) - 2i(\int \operatorname{PolyLog}(2, -e^{-\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{-\operatorname{arccosh}(ax)})) + 2i(\int \operatorname{PolyLog}(2, e^{-\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{-\operatorname{arccosh}(ax)}))}{ac}$$

↓ 2720

$$\frac{i(-2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 2i(\int e^{\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})) - 2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{-\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{-\operatorname{arccosh}(ax)})) + 2i(\int e^{\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{-\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{-\operatorname{arccosh}(ax)}))}{ac}$$

↓ 7143

$$\frac{i(2i \operatorname{arccosh}(ax)^2 \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) - 2i(\operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) - 2i(\operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})) - 2i(\operatorname{PolyLog}(3, -e^{-\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{-\operatorname{arccosh}(ax)})) + 2i(\operatorname{PolyLog}(3, e^{-\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{-\operatorname{arccosh}(ax)}))}{ac}$$

input `Int[ArcCosh[a*x]^2/(c - a^2*c*x^2), x]`

output `((-I)*((2*I)*ArcCosh[a*x]^2*ArcTanh[E^ArcCosh[a*x]] - (2*I)*(-(ArcCosh[a*x]*PolyLog[2, -E^ArcCosh[a*x]]) + PolyLog[3, -E^ArcCosh[a*x]]) + (2*I)*(-(ArcCosh[a*x]*PolyLog[2, E^ArcCosh[a*x]]) + PolyLog[3, E^ArcCosh[a*x]])))/(a*c)`

### 3.167.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

---

3.167.  $\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2 cx^2} dx$

```
rule 3011 Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4670 Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

```
rule 6318 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_.) + (e_.)*(x_)^2), x_Symb
ol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x
]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

### 3.167.4 Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.91

method	result
derivativedivides	$\frac{\operatorname{arccosh}(ax)^2 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{c} + \frac{2 \operatorname{arccosh}(ax) \operatorname{polylog}(2, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c} - \frac{2 \operatorname{polylog}(3, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c} - \frac{\operatorname{arccosh}(ax)}{c}$
default	$\frac{\operatorname{arccosh}(ax)^2 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{c} + \frac{2 \operatorname{arccosh}(ax) \operatorname{polylog}(2, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c} - \frac{2 \operatorname{polylog}(3, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c} - \frac{\operatorname{arccosh}(ax)}{c}$

```
input int(arccosh(a*x)^2/(-a^2*c*x^2+c), x, method=_RETURNVERBOSE)
```

$$3.167. \int \frac{\operatorname{arccosh}(ax)^2}{c-a^2cx^2} dx$$

output  $1/a*(1/c*\operatorname{arccosh}(a*x)^2*\ln(1+a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})+2/c*\operatorname{arccosh}(a*x)*\operatorname{polylog}(2,-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})-2/c*\operatorname{polylog}(3,-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})-1/c*\operatorname{arccosh}(a*x)^2*\ln(1-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})-2/c*\operatorname{arccosh}(a*x)*\operatorname{polylog}(2,a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})+2/c*\operatorname{polylog}(3,a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2))}$

### 3.167.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx = \int -\frac{\operatorname{arcosh}(ax)^2}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c),x, algorithm="fricas")`

output `integral(-arccosh(a*x)^2/(a^2*c*x^2 - c), x)`

### 3.167.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx = -\int \frac{\operatorname{acosh}^2(ax)}{a^2x^2-1} dx$$

input `integrate(acosh(a*x)**2/(-a**2*c*x**2+c),x)`

output `-Integral(acosh(a*x)**2/(a**2*x**2 - 1), x)/c`

### 3.167.7 Maxima [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx = \int -\frac{\operatorname{arcosh}(ax)^2}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c),x, algorithm="maxima")`

output `1/2*(log(a*x + 1) - log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^2 / (a*c) - integrate(((a*x*log(a*x + 1) - a*x*log(a*x - 1))*sqrt(a*x + 1)*sqrt(a*x - 1) + (a^2*x^2 - 1)*log(a*x + 1) - (a^2*x^2 - 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))/(a^3*c*x^3 - a*c*x + (a^2*c*x^2 - c)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

### 3.167.8 Giac [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx = \int -\frac{\operatorname{acosh}(ax)^2}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c),x, algorithm="giac")`

output `integrate(-arccosh(a*x)^2/(a^2*c*x^2 - c), x)`

### 3.167.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{c - a^2cx^2} dx = \int \frac{\operatorname{acosh}(ax)^2}{c - a^2cx^2} dx$$

input `int(acosh(a*x)^2/(c - a^2*c*x^2),x)`

output `int(acosh(a*x)^2/(c - a^2*c*x^2), x)`

**3.168**       $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^2} dx$

3.168.1 Optimal result . . . . . 1388  
 3.168.2 Mathematica [A] (verified) . . . . . 1389  
 3.168.3 Rubi [C] (verified) . . . . . 1389  
 3.168.4 Maple [A] (verified) . . . . . 1394  
 3.168.5 Fricas [F] . . . . . 1394  
 3.168.6 Sympy [F] . . . . . 1395  
 3.168.7 Maxima [F] . . . . . 1395  
 3.168.8 Giac [F] . . . . . 1395  
 3.168.9 Mupad [F(-1)] . . . . . 1396

**3.168.1 Optimal result**

Integrand size = 20, antiderivative size = 163

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^2} dx = -\frac{\operatorname{arccosh}(ax)}{ac^2\sqrt{-1+ax}\sqrt{1+ax}} + \frac{x\operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}$$

$$+ \frac{\operatorname{arccosh}(ax)^2\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac^2} - \frac{\operatorname{arctanh}(ax)}{ac^2}$$

$$+ \frac{\operatorname{arccosh}(ax)\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(ax)})}{ac^2}$$

$$- \frac{\operatorname{arccosh}(ax)\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(ax)})}{ac^2}$$

$$- \frac{\operatorname{PolyLog}(3,-e^{\operatorname{arccosh}(ax)})}{ac^2} + \frac{\operatorname{PolyLog}(3,e^{\operatorname{arccosh}(ax)})}{ac^2}$$

output  $1/2*x*\operatorname{arccosh}(a*x)^2/c^2/(-a^2*x^2+1)+\operatorname{arccosh}(a*x)^2*\operatorname{arctanh}(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^2-\operatorname{arctanh}(a*x)/a/c^2+\operatorname{arccosh}(a*x)*\operatorname{polylog}(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^2-\operatorname{arccosh}(a*x)*\operatorname{polylog}(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^2-\operatorname{polylog}(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^2+\operatorname{polylog}(3,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^2-\operatorname{arccosh}(a*x)/a/c^2/(a*x-1)^(1/2)/(a*x+1)^(1/2)$

### 3.168.2 Mathematica [A] (verified)

Time = 0.89 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.17

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^2} dx$$

$$= \frac{-4\operatorname{arccosh}(ax) \coth\left(\frac{1}{2}\operatorname{arccosh}(ax)\right) - \operatorname{arccosh}(ax)^2 \operatorname{csch}^2\left(\frac{1}{2}\operatorname{arccosh}(ax)\right) - 4\operatorname{arccosh}(ax)^2 \log\left(1 - e^{-\operatorname{arccosh}(ax)}\right)}{8a^2c^2}$$

input `Integrate[ArcCosh[a*x]^2/(c - a^2*c*x^2)^2,x]`

output `(-4*ArcCosh[a*x]*Coth[ArcCosh[a*x]/2] - ArcCosh[a*x]^2*Csch[ArcCosh[a*x]/2]^2 - 4*ArcCosh[a*x]^2*Log[1 - E^(-ArcCosh[a*x])] + 4*ArcCosh[a*x]^2*Log[1 + E^(-ArcCosh[a*x])] + 8*Log[Tanh[ArcCosh[a*x]/2]] - 8*ArcCosh[a*x]*PolyLog[2, -E^(-ArcCosh[a*x])] + 8*ArcCosh[a*x]*PolyLog[2, E^(-ArcCosh[a*x])] - 8*PolyLog[3, -E^(-ArcCosh[a*x])] + 8*PolyLog[3, E^(-ArcCosh[a*x])] - ArcCosh[a*x]^2*Sech[ArcCosh[a*x]/2]^2 + 4*ArcCosh[a*x]*Tanh[ArcCosh[a*x]/2])/(8*a*c^2)`

### 3.168.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.63 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.96, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.650$ , Rules used = {6316, 27, 6318, 3042, 26, 4670, 3011, 2720, 6330, 25, 39, 219, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^2} dx$$

$$\downarrow \text{6316}$$

$$\frac{\int \frac{\operatorname{arccosh}(ax)^2}{c(1-a^2x^2)} dx}{2c} + \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}$$

$$\downarrow \text{27}$$

$$\frac{\int \frac{\operatorname{arccosh}(ax)^2}{1-a^2x^2} dx}{2c^2} + \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}$$

$$\begin{aligned}
& \downarrow 6318 \\
& \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} - \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)} \\
& \downarrow 3042 \\
& \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} - \frac{\int i \operatorname{arccosh}(ax)^2 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)} \\
& \downarrow 26 \\
& \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} - \frac{i \int \operatorname{arccosh}(ax)^2 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)} \\
& \downarrow 4670 \\
& \frac{i(2i \int \operatorname{arccosh}(ax) \log(1 - e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - 2i \int \operatorname{arccosh}(ax) \log(1 + e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2ia \int \operatorname{arccosh}(ax) dx)}{2ac^2} \\
& \quad + \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)} \\
& \downarrow 3011 \\
& \frac{i(-2i(\int \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 2i(\int \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})))}{2ac^2} \\
& \quad + \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)} \\
& \downarrow 2720 \\
& \frac{i(-2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 2i(\int e^{\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})))}{2ac^2} \\
& \quad + \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{c^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)} \\
& \downarrow 6330 \\
& \frac{a \left( \frac{\int -\frac{1}{(1-ax)(ax+1)} dx}{a} - \frac{\operatorname{arccosh}(ax)}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right)}{c^2} - \\
& \frac{i(-2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 2i(\int e^{\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})))}{2ac^2} \\
& \quad + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}
\end{aligned}$$

---

3.168.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^2} dx$

$$\frac{a \left( -\frac{\int \frac{1}{(1-ax)(ax+1)} dx}{a} - \frac{\operatorname{arccosh}(ax)}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right)}{c^2} - \frac{i(-2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) dx - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}))}{c^2} - \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}$$

25

$$\frac{a \left( -\frac{\int \frac{1}{1-a^2x^2} dx}{a} - \frac{\operatorname{arccosh}(ax)}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right)}{c^2} - \frac{i(-2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) dx - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}))}{c^2} - \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}$$

39

$$\frac{a \left( -\frac{\operatorname{arccosh}(ax)}{a^2 \sqrt{ax-1} \sqrt{ax+1}} - \frac{\operatorname{arctanh}(ax)}{a^2} \right)}{c^2} + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}$$

219

$$\frac{i(2i \operatorname{arccosh}(ax)^2 \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) - 2i(\operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + \frac{x \operatorname{arccosh}(ax)^2}{2c^2(1-a^2x^2)}))}{2ac^2}$$

input `Int[ArcCosh[a*x]^2/(c - a^2*c*x^2)^2,x]`

output `(x*ArcCosh[a*x]^2)/(2*c^2*(1 - a^2*x^2)) + (a*(-(ArcCosh[a*x]/(a^2*sqrt[-1 + a*x]*sqrt[1 + a*x])) - ArcTanh[a*x]/a^2))/c^2 - ((I/2)*((2*I)*ArcCosh[a*x]^2*ArcTanh[E^ArcCosh[a*x]] - (2*I)*(-(ArcCosh[a*x]*PolyLog[2, -E^ArcCosh[a*x]]) + PolyLog[3, -E^ArcCosh[a*x]]) + (2*I)*(-(ArcCosh[a*x]*PolyLog[2, E^ArcCosh[a*x]]) + PolyLog[3, E^ArcCosh[a*x]])))/(a*c^2)`

---

3.168.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^2} dx$



## 3.168.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 39 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(m_.)), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_)^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_.) + (e1_.)*(x_))^(p_) * ((d2_.) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.168.4 Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.56

method	result
derivativedivides	$-\frac{\operatorname{arccosh}(ax)(ax \operatorname{arccosh}(ax)+2\sqrt{ax-1}\sqrt{ax+1})}{2(a^2x^2-1)c^2} + \frac{\operatorname{arccosh}(ax)^2 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{2c^2} + \frac{\operatorname{arccosh}(ax) \operatorname{polylog}(2,-ax-\sqrt{ax-1}\sqrt{ax+1})}{c^2}$
default	$-\frac{\operatorname{arccosh}(ax)(ax \operatorname{arccosh}(ax)+2\sqrt{ax-1}\sqrt{ax+1})}{2(a^2x^2-1)c^2} + \frac{\operatorname{arccosh}(ax)^2 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{2c^2} + \frac{\operatorname{arccosh}(ax) \operatorname{polylog}(2,-ax-\sqrt{ax-1}\sqrt{ax+1})}{c^2}$

input `int(arccosh(a*x)^2/(-a^2*c*x^2+c)^2,x,method=_RETURNVERBOSE)`

output `1/a*(-1/2/(a^2*x^2-1)*arccosh(a*x)*(a*x*arccosh(a*x)+2*(a*x-1)^(1/2)*(a*x+1)^(1/2))/c^2+1/2/c^2*arccosh(a*x)^2*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+1/c^2*arccosh(a*x)*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-1/c^2*polylog(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-1/2/c^2*arccosh(a*x)^2*ln(1-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-1/c^2*arccosh(a*x)*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+1/c^2*polylog(3,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))-2/c^2*arctanh(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))`

### 3.168.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^2} dx = \int \frac{\operatorname{arccosh}(ax)^2}{(a^2cx^2-c)^2} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^2,x, algorithm="fricas")`

output `integral(arccosh(a*x)^2/(a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2), x)`

**3.168.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{acosh}^2(ax)}{a^4x^4 - 2a^2x^2 + 1} dx$$

input `integrate(acosh(a*x)**2/(-a**2*c*x**2+c)**2,x)`

output `Integral(acosh(a*x)**2/(a**4*x**4 - 2*a**2*x**2 + 1), x)/c**2`

**3.168.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arcosh}(ax)^2}{(a^2cx^2 - c)^2} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^2,x, algorithm="maxima")`

output `-1/4*(2*a*x - (a^2*x^2 - 1)*log(a*x + 1) + (a^2*x^2 - 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^2/(a^3*c^2*x^2 - a*c^2) - integrate(-1/2*(2*a^3*x^3 + (2*a^2*x^2 - (a^3*x^3 - a*x))*log(a*x + 1) + (a^3*x^3 - a*x)*log(a*x - 1))*sqrt(a*x + 1)*sqrt(a*x - 1) - 2*a*x - (a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1) + (a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))/(a^5*c^2*x^5 - 2*a^3*c^2*x^3 + a*c^2*x + (a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

**3.168.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arcosh}(ax)^2}{(a^2cx^2 - c)^2} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^2,x, algorithm="giac")`

output `integrate(arccosh(a*x)^2/(a^2*c*x^2 - c)^2, x)`

**3.168.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2 cx^2)^2} dx = \int \frac{\operatorname{acosh}(ax)^2}{(c - a^2 cx^2)^2} dx$$

input `int(acosh(a*x)^2/(c - a^2*c*x^2)^2,x)`output `int(acosh(a*x)^2/(c - a^2*c*x^2)^2, x)`

**3.169**       $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^3} dx$

3.169.1 Optimal result . . . . . 1397  
 3.169.2 Mathematica [A] (warning: unable to verify) . . . . . 1398  
 3.169.3 Rubi [C] (verified) . . . . . 1398  
 3.169.4 Maple [A] (verified) . . . . . 1404  
 3.169.5 Fricas [F] . . . . . 1404  
 3.169.6 Sympy [F] . . . . . 1405  
 3.169.7 Maxima [F] . . . . . 1405  
 3.169.8 Giac [F] . . . . . 1406  
 3.169.9 Mupad [F(-1)] . . . . . 1406

**3.169.1 Optimal result**

Integrand size = 20, antiderivative size = 258

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^3} dx = -\frac{x}{12c^3(1-a^2x^2)} + \frac{\operatorname{arccosh}(ax)}{6ac^3(-1+ax)^{3/2}(1+ax)^{3/2}}$$

$$-\frac{3\operatorname{arccosh}(ax)}{4ac^3\sqrt{-1+ax}\sqrt{1+ax}} + \frac{x\operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

$$+\frac{3x\operatorname{arccosh}(ax)^2}{8c^3(1-a^2x^2)} + \frac{3\operatorname{arccosh}(ax)^2\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{4ac^3}$$

$$-\frac{5\operatorname{arctanh}(ax)}{6ac^3} + \frac{3\operatorname{arccosh}(ax)\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{4ac^3}$$

$$-\frac{3\operatorname{arccosh}(ax)\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{4ac^3}$$

$$-\frac{3\operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)})}{4ac^3} + \frac{3\operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)})}{4ac^3}$$

```
output -1/12*x/c^3/(-a^2*x^2+1)+1/6*arccosh(a*x)/a/c^3/(a*x-1)^(3/2)/(a*x+1)^(3/2)
)+1/4*x*arccosh(a*x)^2/c^3/(-a^2*x^2+1)^2+3/8*x*arccosh(a*x)^2/c^3/(-a^2*x
^2+1)+3/4*arccosh(a*x)^2*arctanh(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^3-5/
6*arctanh(a*x)/a/c^3+3/4*arccosh(a*x)*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)
^(1/2))/a/c^3-3/4*arccosh(a*x)*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/
a/c^3-3/4*polylog(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^3+3/4*polylog(3,
a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c^3-3/4*arccosh(a*x)/a/c^3/(a*x-1)^(1/2)
)/(a*x+1)^(1/2)
```

**3.169.2 Mathematica [A] (warning: unable to verify)**

Time = 3.30 (sec) , antiderivative size = 319, normalized size of antiderivative = 1.24

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^3} dx =$$

$$80\operatorname{arccosh}(ax) \operatorname{coth}\left(\frac{1}{2}\operatorname{arccosh}(ax)\right) + 2(-2 + 9\operatorname{arccosh}(ax)^2) \operatorname{csch}^2\left(\frac{1}{2}\operatorname{arccosh}(ax)\right) - 2\sqrt{\frac{-1+ax}{1+ax}}(1 + ax)$$

input `Integrate[ArcCosh[a*x]^2/(c - a^2*c*x^2)^3,x]`

output

```
-1/192*(80*ArcCosh[a*x]*Coth[ArcCosh[a*x]/2] + 2*(-2 + 9*ArcCosh[a*x]^2)*Csch[ArcCosh[a*x]/2]^2 - 2*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*ArcCosh[a*x]*Csch[ArcCosh[a*x]/2]^4 - 3*ArcCosh[a*x]^2*Csch[ArcCosh[a*x]/2]^4 - 160*Log[Tanh[ArcCosh[a*x]/2]] + 72*(ArcCosh[a*x]^2*Log[1 - E^(-ArcCosh[a*x])] - ArcCosh[a*x]^2*Log[1 + E^(-ArcCosh[a*x])]) + 2*ArcCosh[a*x]*PolyLog[2, -E^(-ArcCosh[a*x])] - 2*ArcCosh[a*x]*PolyLog[2, E^(-ArcCosh[a*x])] + 2*PolyLog[3, -E^(-ArcCosh[a*x])] - 2*PolyLog[3, E^(-ArcCosh[a*x])]) + 2*(-2 + 9*ArcCosh[a*x]^2)*Sech[ArcCosh[a*x]/2]^2 + 3*ArcCosh[a*x]^2*Sech[ArcCosh[a*x]/2]^4 - (32*ArcCosh[a*x]*Sinh[ArcCosh[a*x]/2]^4)/(((1 + a*x)/(1 + a*x))^3/2)*(1 + a*x)^3 - 80*ArcCosh[a*x]*Tanh[ArcCosh[a*x]/2])/(a*c^3)
```

**3.169.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.99 (sec) , antiderivative size = 256, normalized size of antiderivative = 0.99, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {6316, 27, 6316, 6318, 3042, 26, 4670, 3011, 2720, 6330, 25, 39, 215, 219, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^3} dx$$

↓ 6316

$$\frac{3 \int \frac{\operatorname{arccosh}(ax)^2}{c^2(1-a^2x^2)^2} dx}{4c} - \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

---

3.169.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^3} dx$

$$\begin{aligned}
& \downarrow 27 \\
& \frac{3 \int \frac{\operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{4c^3} - \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2} \\
& \downarrow 6316 \\
& \frac{3 \left( \frac{1}{2} \int \frac{\operatorname{arccosh}(ax)^2}{1-a^2x^2} dx + a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx + \frac{x \operatorname{arccosh}(ax)^2}{2(1-a^2x^2)} \right)}{4c^3} - \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \\
& \quad \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2} \\
& \downarrow 6318 \\
& \frac{3 \left( a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{\frac{ax-1}{ax+1}}} d\operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)^2}{2(1-a^2x^2)} \right)}{4c^3} - \\
& \quad \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2} \\
& \downarrow 3042 \\
& \frac{3 \left( a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{\int i \operatorname{arccosh}(ax)^2 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)^2}{2(1-a^2x^2)} \right)}{4c^3} - \\
& \quad \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2} \\
& \downarrow 26 \\
& \frac{3 \left( a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{i \int \operatorname{arccosh}(ax)^2 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)^2}{2(1-a^2x^2)} \right)}{4c^3} - \\
& \quad \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2} \\
& \downarrow 4670 \\
& \frac{3 \left( -\frac{i(2i \int \operatorname{arccosh}(ax) \log(1-e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - 2i \int \operatorname{arccosh}(ax) \log(1+e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2i \operatorname{arccosh}(ax)^2 \operatorname{arctanh}(ax))}{2a}}{4c^3} \right)}{4c^3} \\
& \quad \frac{a \int \frac{x \operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2} \\
& \downarrow 3011
\end{aligned}$$

---

3.169.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^3} dx$



$$3 \left( -\frac{i(-2i(\int \text{PolyLog}(2, -e^{\text{arccosh}(ax)}) d\text{arccosh}(ax) - \text{arccosh}(ax) \text{PolyLog}(2, -e^{\text{arccosh}(ax)})) + 2i(\int \text{PolyLog}(2, e^{\text{arccosh}(ax)}) d\text{arccosh}(ax) - \text{arccosh}(ax) \text{PolyLog}(2, e^{\text{arccosh}(ax)}))}{2a} \right)$$

$$\frac{a \int \frac{x \text{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \text{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

↓ 2720

$$3 \left( -\frac{i(-2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, -e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, -e^{\text{arccosh}(ax)})) + 2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, e^{\text{arccosh}(ax)}))}{2a} \right)$$

$$\frac{a \int \frac{x \text{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{2c^3} + \frac{x \text{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

↓ 6330

$$3 \left( a \left( \frac{\int -\frac{1}{(1-ax)(ax+1)} dx}{a} - \frac{\text{arccosh}(ax)}{a^2\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{i(-2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, -e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, -e^{\text{arccosh}(ax)})) + 2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, e^{\text{arccosh}(ax)}))}{2a} \right)$$

$$\frac{a \left( \frac{\int \frac{1}{(1-ax)^2(ax+1)^2} dx}{3a} - \frac{\text{arccosh}(ax)}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{2c^3} + \frac{x \text{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

↓ 25

$$3 \left( a \left( -\frac{\int \frac{1}{(1-ax)(ax+1)} dx}{a} - \frac{\text{arccosh}(ax)}{a^2\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{i(-2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, -e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, -e^{\text{arccosh}(ax)})) + 2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, e^{\text{arccosh}(ax)}))}{2a} \right)$$

$$\frac{a \left( \frac{\int \frac{1}{(1-ax)^2(ax+1)^2} dx}{3a} - \frac{\text{arccosh}(ax)}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{2c^3} + \frac{x \text{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

↓ 39

$$3 \left( a \left( -\frac{\int \frac{1}{1-a^2x^2} dx}{a} - \frac{\text{arccosh}(ax)}{a^2\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{i(-2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, -e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, -e^{\text{arccosh}(ax)})) + 2i(\int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, e^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, e^{\text{arccosh}(ax)}))}{2a} \right)$$

$$\frac{a \left( \frac{\int \frac{1}{(1-a^2x^2)^2} dx}{3a} - \frac{\text{arccosh}(ax)}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{2c^3} + \frac{x \text{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

---

3.169.  $\int \frac{\text{arccosh}(ax)^2}{(c-a^2cx^2)^3} dx$

↓ 215

$$3 \left( a \left( -\frac{\int \frac{1}{1-a^2x^2} dx}{a} - \frac{\operatorname{arccosh}(ax)}{a^2\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{i(-2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) dx)}{2c^3} \right) + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

↓ 219

$$3 \left( -\frac{i(-2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) dx) + 2i(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) dx)}{2a} \right) + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

↓ 7143

$$3 \left( a \left( -\frac{\operatorname{arccosh}(ax)}{a^2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arctanh}(ax)}{a^2} \right) + \frac{x \operatorname{arccosh}(ax)^2}{2(1-a^2x^2)} - \frac{i(2i \operatorname{arccosh}(ax)^2 \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) - 2i(\operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) dx)}{4c^3} \right) + \frac{x \operatorname{arccosh}(ax)^2}{4c^3(1-a^2x^2)^2}$$

input `Int[ArcCosh[a*x]^2/(c - a^2*c*x^2)^3,x]`

output `(x*ArcCosh[a*x]^2)/(4*c^3*(1 - a^2*x^2)^2) - (a*(-1/3*ArcCosh[a*x]/(a^2*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2)) + (x/(2*(1 - a^2*x^2)) + ArcTanh[a*x]/(2*a))/ (3*a)))/(2*c^3) + (3*((x*ArcCosh[a*x]^2)/(2*(1 - a^2*x^2)) + a*(-(ArcCosh[a*x]/(a^2*sqrt[-1 + a*x]*sqrt[1 + a*x])) - ArcTanh[a*x]/a^2) - ((I/2)*((2*I)*ArcCosh[a*x]^2*ArcTanh[E^ArcCosh[a*x]] - (2*I)*(-(ArcCosh[a*x]*PolyLog[2, -E^ArcCosh[a*x]]) + PolyLog[3, -E^ArcCosh[a*x]]) + (2*I)*(-(ArcCosh[a*x]*PolyLog[2, E^ArcCosh[a*x]]) + PolyLog[3, E^ArcCosh[a*x])))/a))/(4*c^3)`

---

3.169.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^3} dx$

## 3.169.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 39 `Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(m_.)), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`
- rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1)/(2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.169.4 Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 320, normalized size of antiderivative = 1.24

method	result
derivativedivides	$\frac{-9a^3x^3 \operatorname{arccosh}(ax)^2 + 18a^2x^2 \operatorname{arccosh}(ax)\sqrt{ax-1}\sqrt{ax+1} - 15ax \operatorname{arccosh}(ax)^2 - 22\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - 2a^3x^3 + 2ax}{24(a^4x^4 - 2a^2x^2 + 1)c^3} - \frac{5 \operatorname{arctanh}(ax)}{c^3}$
default	$\frac{-9a^3x^3 \operatorname{arccosh}(ax)^2 + 18a^2x^2 \operatorname{arccosh}(ax)\sqrt{ax-1}\sqrt{ax+1} - 15ax \operatorname{arccosh}(ax)^2 - 22\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - 2a^3x^3 + 2ax}{24(a^4x^4 - 2a^2x^2 + 1)c^3} - \frac{5 \operatorname{arctanh}(ax)}{c^3}$

input `int(arccosh(a*x)^2/(-a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{a} \left( -\frac{1}{24} (9a^3x^3 \operatorname{arccosh}(ax)^2 + 18a^2x^2 \operatorname{arccosh}(ax) \sqrt{ax-1} \sqrt{ax+1} - 15ax \operatorname{arccosh}(ax)^2 - 22\sqrt{ax-1} \sqrt{ax+1} \operatorname{arccosh}(ax) - 2a^3x^3 + 2ax) / (a^4x^4 - 2a^2x^2 + 1) / c^3 - \frac{5}{3} \operatorname{arctanh}(ax) \right) + \frac{3}{8} \operatorname{arccosh}(ax)^2 \ln(1 + ax + (ax-1)^{1/2} \sqrt{ax+1}) + \frac{3}{4} \operatorname{arccosh}(ax) \operatorname{polylog}(2, -ax - (ax-1)^{1/2} \sqrt{ax+1}) - \frac{3}{4} \operatorname{polylog}(3, -ax - (ax-1)^{1/2} \sqrt{ax+1}) - \frac{3}{8} \operatorname{arccosh}(ax)^2 \ln(1 - ax - (ax-1)^{1/2} \sqrt{ax+1}) - \frac{3}{4} \operatorname{arccosh}(ax) \operatorname{polylog}(2, ax + (ax-1)^{1/2} \sqrt{ax+1}) + \frac{3}{4} \operatorname{polylog}(3, ax + (ax-1)^{1/2} \sqrt{ax+1}) \right)$$

### 3.169.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arccosh}(ax)^2}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^3,x, algorithm="fricas")`

output `integral(-arccosh(a*x)^2/(a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3), x)`

## 3.169.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^3} dx = -\int \frac{\operatorname{acosh}^2(ax)}{a^6x^6 - 3a^4x^4 + 3a^2x^2 - 1} \frac{dx}{c^3}$$

input `integrate(acosh(a*x)**2/(-a**2*c*x**2+c)**3,x)`

output `-Integral(acosh(a*x)**2/(a**6*x**6 - 3*a**4*x**4 + 3*a**2*x**2 - 1), x)/c**3`

## 3.169.7 Maxima [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arcosh}(ax)^2}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^3,x, algorithm="maxima")`

output `-1/16*(6*a^3*x^3 - 10*a*x - 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1) + 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1))*sqrt(a*x - 1))^2/(a^5*c^3*x^4 - 2*a^3*c^3*x^2 + a*c^3) - integrate(-1/8*(6*a^5*x^5 - 16*a^3*x^3 + (6*a^4*x^4 - 10*a^2*x^2 - 3*(a^5*x^5 - 2*a^3*x^3 + a*x))*log(a*x + 1) + 3*(a^5*x^5 - 2*a^3*x^3 + a*x)*log(a*x - 1))*sqrt(a*x + 1)*sqrt(a*x - 1) + 10*a*x - 3*(a^6*x^6 - 3*a^4*x^4 + 3*a^2*x^2 - 1)*log(a*x + 1) + 3*(a^6*x^6 - 3*a^4*x^4 + 3*a^2*x^2 - 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1))*sqrt(a*x - 1))/(a^7*c^3*x^7 - 3*a^5*c^3*x^5 + 3*a^3*c^3*x^3 - a*c^3*x + (a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

**3.169.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arcosh}(ax)^2}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^3,x, algorithm="giac")`

output `integrate(-arccosh(a*x)^2/(a^2*c*x^2 - c)^3, x)`

**3.169.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^3} dx = \int \frac{\operatorname{acosh}(ax)^2}{(c - a^2cx^2)^3} dx$$

input `int(acosh(a*x)^2/(c - a^2*c*x^2)^3,x)`

output `int(acosh(a*x)^2/(c - a^2*c*x^2)^3, x)`

### 3.170 $\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$

3.170.1 Optimal result . . . . .	1407
3.170.2 Mathematica [A] (verified) . . . . .	1408
3.170.3 Rubi [A] (verified) . . . . .	1408
3.170.4 Maple [B] (verified) . . . . .	1414
3.170.5 Fricas [A] (verification not implemented) . . . . .	1415
3.170.6 Sympy [F] . . . . .	1416
3.170.7 Maxima [A] (verification not implemented) . . . . .	1416
3.170.8 Giac [F(-2)] . . . . .	1417
3.170.9 Mupad [F(-1)] . . . . .	1417

#### 3.170.1 Optimal result

Integrand size = 29, antiderivative size = 371

$$\begin{aligned} \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = & -\frac{856b^2 \sqrt{d - c^2 dx^2}}{3375c^4} + \frac{22b^2 x^2 \sqrt{d - c^2 dx^2}}{3375c^2} \\ & + \frac{2}{125} b^2 x^4 \sqrt{d - c^2 dx^2} + \frac{4abx \sqrt{d - c^2 dx^2}}{15c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ & + \frac{4b^2 x \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{15c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ & + \frac{2bx^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{45c \sqrt{-1 + cx} \sqrt{1 + cx}} \\ & - \frac{2bcx^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{25 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ & - \frac{2\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{15c^4} \\ & - \frac{x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{15c^2} \\ & + \frac{1}{5} x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \end{aligned}$$



output 
$$\begin{aligned} & -856/3375*b^2*(-c^2*d*x^2+d)^{(1/2)}/c^4+22/3375*b^2*x^2*(-c^2*d*x^2+d)^{(1/2)} \\ & )/c^2+2/125*b^2*x^4*(-c^2*d*x^2+d)^{(1/2)}-2/15*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d \\ & *x^2+d)^{(1/2)}/c^4-1/15*x^2*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d*x^2+d)^{(1/2)}/c^2+1 \\ & /5*x^4*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d*x^2+d)^{(1/2)}+4/15*a*b*x*(-c^2*d*x^2+d) \\ & ^{(1/2)}/c^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+4/15*b^2*x*\operatorname{arccosh}(c*x)*(-c^2*d*x^2 \\ & +d)^{(1/2)}/c^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+2/45*b*x^3*(a+b*\operatorname{arccosh}(c*x))*(- \\ & c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2/25*b*c*x^5*(a+b*\operatorname{arccosh} \\ & (c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \end{aligned}$$

### 3.170.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 237, normalized size of antiderivative = 0.64

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$= \frac{\sqrt{d - c^2 dx^2} \left( 225a^2(-1 + c^2 x^2)^2 (2 + 3c^2 x^2) - 30abcx \sqrt{-1 + cx} \sqrt{1 + cx} (-30 - 5c^2 x^2 + 9c^4 x^4) + 2b^2(428 \right)}{\dots}$$

input `Integrate[x^3*sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output 
$$\begin{aligned} & (\operatorname{sqrt}[d - c^2*d*x^2]*(225*a^2*(-1 + c^2*x^2)^2*(2 + 3*c^2*x^2) - 30*a*b*c* \\ & x*\operatorname{sqrt}[-1 + c*x]*\operatorname{sqrt}[1 + c*x]*(-30 - 5*c^2*x^2 + 9*c^4*x^4) + 2*b^2*(428 \\ & - 439*c^2*x^2 - 16*c^4*x^4 + 27*c^6*x^6) + 30*b*(15*a*(-1 + c^2*x^2)^2*(2 \\ & + 3*c^2*x^2) + b*c*x*\operatorname{sqrt}[-1 + c*x]*\operatorname{sqrt}[1 + c*x]*(30 + 5*c^2*x^2 - 9*c^4* \\ & x^4))*\operatorname{ArcCosh}[c*x] + 225*b^2*(-1 + c^2*x^2)^2*(2 + 3*c^2*x^2)*\operatorname{ArcCosh}[c*x] \\ & ^2)/(3375*c^4*(-1 + c^2*x^2)) \end{aligned}$$

### 3.170.3 Rubi [A] (verified)

Time = 2.17 (sec) , antiderivative size = 425, normalized size of antiderivative = 1.15, number of steps used = 14, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$ , Rules used = {6341, 6298, 111, 27, 111, 27, 83, 6354, 6298, 111, 27, 83, 6330, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$\begin{aligned}
& \downarrow 6341 \\
& \frac{2bc\sqrt{d-c^2dx^2} \int x^4(a+\operatorname{barccosh}(cx))dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \\
& \quad \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 6298 \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \int \frac{x^5}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{5\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 111 \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{\int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \\
& \quad \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 27 \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \\
& \quad \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 111 \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \\
& \quad \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 27
\end{aligned}$$

$$\frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{5\sqrt{cx-1}\sqrt{cx+1}} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} +$$

83

$$\frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{5\sqrt{cx-1}\sqrt{cx+1}}$$

6354

$$\frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \left( 2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{2b \int x^2(a+\operatorname{barccosh}(cx))dx}{3c} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^2}{3c^2} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{5\sqrt{cx-1}\sqrt{cx+1}}$$

6298

$$\frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \left( 2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{2b \left( \frac{1}{3}x^3(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{3c} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^2}{3c^2} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{5\sqrt{cx-1}\sqrt{cx+1}}$$

111

$$\begin{aligned}
& \sqrt{d - c^2 dx^2} \left( -\frac{2b \left( \frac{1}{3} x^3 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} bc \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c} + \frac{2 \int \frac{x(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \\
& \frac{5\sqrt{cx-1}\sqrt{cx+1}}{\frac{1}{5} x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 -} \\
& \frac{2bc\sqrt{d - c^2 dx^2} \left( \frac{1}{5} x^5 (a + \operatorname{barccosh}(cx)) - \frac{1}{5} bc \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \quad 27 \\
& \sqrt{d - c^2 dx^2} \left( -\frac{2b \left( \frac{1}{3} x^3 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} bc \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c} + \frac{2 \int \frac{x(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \\
& \frac{5\sqrt{cx-1}\sqrt{cx+1}}{\frac{1}{5} x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 -} \\
& \frac{2bc\sqrt{d - c^2 dx^2} \left( \frac{1}{5} x^5 (a + \operatorname{barccosh}(cx)) - \frac{1}{5} bc \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \quad 83 \\
& \sqrt{d - c^2 dx^2} \left( \frac{2 \int \frac{x(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1} (a + \operatorname{barccosh}(cx))^2}{3c^2} - \frac{2b \left( \frac{1}{3} x^3 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \right) \right)}{3c} \right) \\
& \frac{5\sqrt{cx-1}\sqrt{cx+1}}{\frac{1}{5} x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 -} \\
& \frac{2bc\sqrt{d - c^2 dx^2} \left( \frac{1}{5} x^5 (a + \operatorname{barccosh}(cx)) - \frac{1}{5} bc \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \quad 6330
\end{aligned}$$

$$\frac{\sqrt{d - c^2 dx^2} \left( \frac{2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{c^2} - \frac{2b \int (a+b\operatorname{arccosh}(cx)) dx}{c} \right)}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{3c^2} - \frac{2b \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) \right)}{3c^2} \right)}{5\sqrt{cx-1}\sqrt{cx+1}}$$


---


$$\frac{\frac{1}{5}x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2 - 2bc\sqrt{d - c^2 dx^2} \left( \frac{1}{5}x^5(a + \operatorname{arccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}}$$


---

↓ 2009

$$\frac{\sqrt{d - c^2 dx^2} \left( \frac{1}{5}x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2 - \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{3c^2} + \frac{2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{c^2} - \frac{2b \left( \frac{ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c} \right)}{3c^2} \right)}{5\sqrt{cx-1}\sqrt{cx+1}}$$


---


$$\frac{2bc\sqrt{d - c^2 dx^2} \left( \frac{1}{5}x^5(a + \operatorname{arccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `(x^4*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/5 - (2*b*c*Sqrt[d - c^2*d*x^2]*(-1/5*(b*c*((x^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(5*c^2) + (4*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2))))/(5*c^2))) + (x^5*(a + b*ArcCosh[c*x]))/5)/(5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*((x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(3*c^2) - (2*b*(-1/3*(b*c*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2))) + (x^3*(a + b*ArcCosh[c*x]))/3))/(3*c) + (2*((Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/c^2 - (2*b*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/c))/(3*c^2)))/(5*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.170.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 111 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*(x_)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

```
rule 6341 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) +
(e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sq
rt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0]
&& IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])
```

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.170.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1283 vs.  $2(315) = 630$ .

Time = 1.49 (sec) , antiderivative size = 1284, normalized size of antiderivative = 3.46

method	result	size
default	Expression too large to display	1284
parts	Expression too large to display	1284

```
input int(x^3*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output

```

a^2*(-1/5*x^2*(-c^2*d*x^2+d)^(3/2)/c^2/d-2/15/d/c^4*(-c^2*d*x^2+d)^(3/2))+
b^2*(1/4000*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)
*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(25*arccosh(c*x)^2-10*arccosh(c*x)+2)/
(c*x+1)/c^4/(c*x-1)+1/864*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c
*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(9*
arccosh(c*x)^2-6*arccosh(c*x)+2)/(c*x+1)/c^4/(c*x-1)-1/16*(-d*(c^2*x^2-1))
^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*arcco
sh(c*x)+2)/(c*x+1)/c^4/(c*x-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)
*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2+2*arccosh(c*x)+2)/(c*x+1)/c^
4/(c*x-1)+1/864*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3
*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(9*arccosh(c
*x)^2+6*arccosh(c*x)+2)/(c*x+1)/c^4/(c*x-1)+1/4000*(-d*(c^2*x^2-1))^(1/2)*
(-16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+
1)^(1/2)*c^3*x^3-28*c^4*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1
)*(25*arccosh(c*x)^2+10*arccosh(c*x)+2)/(c*x+1)/c^4/(c*x-1))+2*a*b*(1/800*
(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/
2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/
2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c*x)))/(c*x+1)/c^4/(c*x-1)+1/288*(-d*
(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c...

```

### 3.170.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 349, normalized size of antiderivative = 0.94

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$= \frac{225 (3 b^2 c^6 x^6 - 4 b^2 c^4 x^4 - b^2 c^2 x^2 + 2 b^2) \sqrt{-c^2 dx^2 + d} \log (cx + \sqrt{c^2 x^2 - 1})^2 - 30 (9 abc^5 x^5 - 5 abc^3 x^3 - \dots}{\dots}$$

input `integrate(x^3*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`



```
output 1/3375*(225*(3*b^2*c^6*x^6 - 4*b^2*c^4*x^4 - b^2*c^2*x^2 + 2*b^2)*sqrt(-c^
2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1))^2 - 30*(9*a*b*c^5*x^5 - 5*a*b*c^
3*x^3 - 30*a*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 30*((9*b^2*c^
5*x^5 - 5*b^2*c^3*x^3 - 30*b^2*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)
- 15*(3*a*b*c^6*x^6 - 4*a*b*c^4*x^4 - a*b*c^2*x^2 + 2*a*b)*sqrt(-c^2*d*x^
2 + d))*log(c*x + sqrt(c^2*x^2 - 1)) + (27*(25*a^2 + 2*b^2)*c^6*x^6 - 4*(2
25*a^2 + 8*b^2)*c^4*x^4 - (225*a^2 + 878*b^2)*c^2*x^2 + 450*a^2 + 856*b^2)
*sqrt(-c^2*d*x^2 + d))/(c^6*x^2 - c^4)
```

### 3.170.6 Sympy [F]

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int x^3 \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2 dx$$

```
input integrate(x**3*(a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2), x)
```

```
output Integral(x**3*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2, x)
```

### 3.170.7 Maxima [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 326, normalized size of antiderivative = 0.88

$$\begin{aligned} & \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx \\ &= -\frac{1}{15} b^2 \left( \frac{3(-c^2 dx^2 + d)^{\frac{3}{2}} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}}}{c^4 d} \right) \operatorname{arcosh}(cx)^2 \\ & \quad - \frac{2}{15} ab \left( \frac{3(-c^2 dx^2 + d)^{\frac{3}{2}} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}}}{c^4 d} \right) \operatorname{arcosh}(cx) \\ & \quad - \frac{1}{15} a^2 \left( \frac{3(-c^2 dx^2 + d)^{\frac{3}{2}} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}}}{c^4 d} \right) \\ & \quad + \frac{2}{3375} b^2 \left( \frac{27 \sqrt{c^2 x^2 - 1} c^2 \sqrt{-dx^4} + 11 \sqrt{c^2 x^2 - 1} \sqrt{-dx^2} - \frac{428 \sqrt{c^2 x^2 - 1} \sqrt{-d}}{c^2}}{c^2} - \frac{15(9c^4 \sqrt{-dx^5} - 5c^2 \sqrt{-dx^3} - 30 \sqrt{-dx})}{c^3} \right) \\ & \quad - \frac{2(9c^4 \sqrt{-dx^5} - 5c^2 \sqrt{-dx^3} - 30 \sqrt{-dx}) ab}{225 c^3} \end{aligned}$$

input `integrate(x^3*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/15*b^2*(3*(-c^2*d*x^2 + d)^(3/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(3/2)/(c^4*d))*arccosh(c*x)^2 - 2/15*a*b*(3*(-c^2*d*x^2 + d)^(3/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(3/2)/(c^4*d))*arccosh(c*x) - 1/15*a^2*(3*(-c^2*d*x^2 + d)^(3/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(3/2)/(c^4*d)) + 2/3375*b^2*(27*sqrt(c^2*x^2 - 1)*c^2*sqrt(-d)*x^4 + 11*sqrt(c^2*x^2 - 1)*sqrt(-d)*x^2 - 428*sqrt(c^2*x^2 - 1)*sqrt(-d)/c^2)/c^2 - 15*(9*c^4*sqrt(-d)*x^5 - 5*c^2*sqrt(-d)*x^3 - 30*sqrt(-d)*x)*arccosh(c*x)/c^3 - 2/225*(9*c^4*sqrt(-d)*x^5 - 5*c^2*sqrt(-d)*x^3 - 30*sqrt(-d)*x)*a*b/c^3`

### 3.170.8 Giac [F(-2)]

Exception generated.

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.170.9 Mupad [F(-1)]

Timed out.

$$\int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int x^3 (a + b \operatorname{acosh}(cx))^2 \sqrt{d - c^2 dx^2} dx$$

input `int(x^3*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2),x)`

output `int(x^3*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2), x)`

### 3.171 $\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$

3.171.1 Optimal result . . . . .	1418
3.171.2 Mathematica [A] (warning: unable to verify) . . . . .	1419
3.171.3 Rubi [A] (verified) . . . . .	1419
3.171.4 Maple [B] (verified) . . . . .	1424
3.171.5 Fricas [F] . . . . .	1425
3.171.6 Sympy [F] . . . . .	1425
3.171.7 Maxima [F] . . . . .	1426
3.171.8 Giac [F] . . . . .	1426
3.171.9 Mupad [F(-1)] . . . . .	1426

#### 3.171.1 Optimal result

Integrand size = 29, antiderivative size = 319

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = -\frac{b^2 x \sqrt{d - c^2 dx^2}}{64c^2} + \frac{1}{32} b^2 x^3 \sqrt{d - c^2 dx^2} - \frac{b^2 \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{64c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{b x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{8c \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{bcx^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{8 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{8c^2} + \frac{1}{4} x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^3}{24bc^3 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

output

```
-1/64*b^2*x*(-c^2*d*x^2+d)^(1/2)/c^2+1/32*b^2*x^3*(-c^2*d*x^2+d)^(1/2)-1/8
*x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/c^2+1/4*x^3*(a+b*arccosh(c*x)
)^2*(-c^2*d*x^2+d)^(1/2)-1/64*b^2*arccosh(c*x)*(-c^2*d*x^2+d)^(1/2)/c^3/(c
*x-1)^(1/2)/(c*x+1)^(1/2)+1/8*b*x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2
)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/8*b*c*x^4*(a+b*arccosh(c*x))*(-c^2*d*x^2
+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/24*(a+b*arccosh(c*x))^3*(-c^2*d*x^
2+d)^(1/2)/b/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.171.2 Mathematica [A] (warning: unable to verify)

Time = 2.13 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.76

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx =$$

$$-96a^2 cx(-1 + 2c^2 x^2) \sqrt{d - c^2 dx^2} + 96a^2 \sqrt{d} \arctan\left(\frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d(-1 + c^2 x^2)}}\right) + \frac{12ab\sqrt{d - c^2 dx^2} (\operatorname{barccosh}(cx)^2 + \cosh(4\operatorname{arccosh}(cx)))}{\sqrt{d(-1 + c^2 x^2)}} + \dots$$

input `Integrate[x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `-1/768*(-96*a^2*c*x*(-1 + 2*c^2*x^2)*Sqrt[d - c^2*d*x^2] + 96*a^2*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + (12*a*b*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + (b^2*Sqrt[d - c^2*d*x^2]*(32*ArcCosh[c*x]^3 + 12*ArcCosh[c*x]*Cosh[4*ArcCosh[c*x]] - 3*(1 + 8*ArcCosh[c*x]^2)*Sinh[4*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))/c^3`

### 3.171.3 Rubi [A] (verified)

Time = 1.87 (sec) , antiderivative size = 328, normalized size of antiderivative = 1.03, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$ , Rules used = {6341, 6298, 111, 27, 101, 43, 6354, 6298, 101, 43, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$\downarrow \text{6341}$$

$$-\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2 (a + \operatorname{barccosh}(cx))^2}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{4\sqrt{cx - 1} \sqrt{cx + 1}} - \frac{bc\sqrt{d - c^2 dx^2} \int x^3 (a + \operatorname{barccosh}(cx)) dx}{2\sqrt{cx - 1} \sqrt{cx + 1}} +$$

$$\frac{1}{4} x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2$$

$$\downarrow \text{6298}$$

$$\begin{aligned}
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \int \frac{x^4}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \quad \downarrow \text{111} \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{\int \frac{3x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \quad \downarrow \text{27} \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \quad \downarrow \text{101} \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \quad \downarrow \text{43} \\
& \frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \\
& \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{6354}
\end{aligned}$$

---

3.171.  $\int x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{\sqrt{d-c^2}dx^2 \left( \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \int x(a+b\operatorname{arccosh}(cx)) dx}{c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} +$$

$$\frac{\frac{1}{4}x^3\sqrt{d-c^2}dx^2(a+b\operatorname{arccosh}(cx))^2 - bc\sqrt{d-c^2}dx^2 \left( \frac{1}{4}x^4(a+b\operatorname{arccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}}{6298}$$

$$\frac{\sqrt{d-c^2}dx^2 \left( \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \left( \frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} +$$

$$\frac{\frac{1}{4}x^3\sqrt{d-c^2}dx^2(a+b\operatorname{arccosh}(cx))^2 - bc\sqrt{d-c^2}dx^2 \left( \frac{1}{4}x^4(a+b\operatorname{arccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}}{101}$$

$$\frac{\sqrt{d-c^2}dx^2 \left( - \frac{b \left( \frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} +$$

$$\frac{\frac{1}{4}x^3\sqrt{d-c^2}dx^2(a+b\operatorname{arccosh}(cx))^2 - bc\sqrt{d-c^2}dx^2 \left( \frac{1}{4}x^4(a+b\operatorname{arccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}}{43}$$

$$\begin{aligned}
& \sqrt{d - c^2 dx^2} \left( \frac{\int \frac{(a + b \operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{2c^2} - \frac{b \left( \frac{1}{2} x^2 (a + b \operatorname{arccosh}(cx)) - \frac{1}{2} bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c} \right) \\
& \frac{4\sqrt{cx-1}\sqrt{cx+1}}{\frac{1}{4} x^3 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - bc \sqrt{d - c^2 dx^2} \left( \frac{1}{4} x^4 (a + b \operatorname{arccosh}(cx)) - \frac{1}{4} bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}} \\
& \quad \downarrow \text{6308} \\
& \frac{1}{4} x^3 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - \sqrt{d - c^2 dx^2} \left( \frac{(a + b \operatorname{arccosh}(cx))^3}{6bc^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{2c^2} - \frac{b \left( \frac{1}{2} x^2 (a + b \operatorname{arccosh}(cx)) - \frac{1}{2} bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c} \right) \\
& \frac{4\sqrt{cx-1}\sqrt{cx+1}}{bc \sqrt{d - c^2 dx^2} \left( \frac{1}{4} x^4 (a + b \operatorname{arccosh}(cx)) - \frac{1}{4} bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `(x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/4 - (b*c*Sqrt[d - c^2*d*x^2]*((x^4*(a + b*ArcCosh[c*x]))/4 - (b*c*((x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/4 - (b*c*((x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/(4*c^2))/4)/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(2*c^2) + (a + b*ArcCosh[c*x])^3/(6*b*c^3) - (b*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/2)/c))/(4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.171.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 111 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 6298 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6308 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(Sqrt[(d1_) + (e1_)*(x_)]*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`



```
rule 6341 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) +
(e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqr
rt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0]
&& IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])
```

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.171.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 677 vs. 2(271) = 542.

Time = 0.65 (sec) , antiderivative size = 678, normalized size of antiderivative = 2.13

method	result
default	$-\frac{a^2x(-c^2dx^2+d)^{\frac{3}{2}}}{4c^2d} + \frac{a^2x\sqrt{-c^2dx^2+d}}{8c^2} + \frac{a^2d \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{8c^2\sqrt{c^2d}} + b^2 \left( -\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^3}{24\sqrt{cx-1}\sqrt{cx+1}c^3} + \frac{\sqrt{-d(c^2x^2-1)}}{24\sqrt{cx-1}\sqrt{cx+1}c^3} \right)$
parts	$-\frac{a^2x(-c^2dx^2+d)^{\frac{3}{2}}}{4c^2d} + \frac{a^2x\sqrt{-c^2dx^2+d}}{8c^2} + \frac{a^2d \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{8c^2\sqrt{c^2d}} + b^2 \left( -\frac{\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^3}{24\sqrt{cx-1}\sqrt{cx+1}c^3} + \frac{\sqrt{-d(c^2x^2-1)}}{24\sqrt{cx-1}\sqrt{cx+1}c^3} \right)$

```
input int(x^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output 
$$\begin{aligned} & -1/4*a^2*x*(-c^2*d*x^2+d)^{(3/2)}/c^2/d+1/8*a^2/c^2*x*(-c^2*d*x^2+d)^{(1/2)}+1 \\ & /8*a^2/c^2*d/(c^2*d)^{(1/2)}*\arctan((c^2*d)^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)})+b^ \\ & 2*(-1/24*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}/c^3*\operatorname{arccosh}(c* \\ & x)^3+1/512*(-d*(c^2*x^2-1))^{(1/2)}*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^{(1/2)}*(c \\ & *x-1)^{(1/2)}*c^4*x^4+4*c*x-8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+(c*x-1)^{(1 \\ & /2)}*(c*x+1)^{(1/2)})*(8*\operatorname{arccosh}(c*x)^2-4*\operatorname{arccosh}(c*x)+1)/(c*x+1)/c^3/(c*x-1) \\ & +1/512*(-d*(c^2*x^2-1))^{(1/2)}*(-8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^4*x^4+8*c^ \\ & 5*x^5+8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-12*c^3*x^3-(c*x-1)^{(1/2)}*(c*x+ \\ & 1)^{(1/2)}+4*c*x)*(8*\operatorname{arccosh}(c*x)^2+4*\operatorname{arccosh}(c*x)+1)/(c*x+1)/c^3/(c*x-1))+2 \\ & *a*b*(-1/16*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}/c^3*\operatorname{arccosh} \\ & (c*x)^2+1/256*(-d*(c^2*x^2-1))^{(1/2)}*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^{(1/2)} \\ & *(c*x-1)^{(1/2)}*c^4*x^4+4*c*x-8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+(c*x-1) \\ & ^{(1/2)}*(c*x+1)^{(1/2)})*(-1+4*\operatorname{arccosh}(c*x))/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c \\ & ^2*x^2-1))^{(1/2)}*(-8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^4*x^4+8*c^5*x^5+8*(c*x- \\ & 1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-12*c^3*x^3-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+4*c* \\ & x)*(1+4*\operatorname{arccosh}(c*x))/(c*x+1)/c^3/(c*x-1) \end{aligned}$$

### 3.171.5 Fricas [F]

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral((b^2*x^2*arccosh(c*x))^2 + 2*a*b*x^2*arccosh(c*x) + a^2*x^2)*sqrt(-c^2*d*x^2 + d), x)`

### 3.171.6 Sympy [F]

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int x^2 \sqrt{-d (cx - 1) (cx + 1)} (a + b \operatorname{acosh}(cx))^2 dx$$

input `integrate(x**2*(a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**2*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2, x)`

---

3.171.  $\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$

**3.171.7 Maxima [F]**

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2 dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `1/8*a^2*(sqrt(-c^2*d*x^2 + d)*x/c^2 - 2*(-c^2*d*x^2 + d)^(3/2)*x/(c^2*d) + sqrt(d)*arcsin(c*x)/c^3) + integrate(sqrt(-c^2*d*x^2 + d)*b^2*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2 + 2*sqrt(-c^2*d*x^2 + d)*a*b*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1), x)`

**3.171.8 Giac [F]**

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2 dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^2*x^2, x)`

**3.171.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2 dx = \int x^2 (a + b \operatorname{acosh}(cx))^2 \sqrt{d - c^2 dx^2} dx$$

input `int(x^2*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2),x)`

output `int(x^2*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2), x)`

### 3.172 $\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 dx$

3.172.1 Optimal result . . . . .	1427
3.172.2 Mathematica [A] (verified) . . . . .	1428
3.172.3 Rubi [A] (verified) . . . . .	1428
3.172.4 Maple [B] (verified) . . . . .	1431
3.172.5 Fricas [A] (verification not implemented) . . . . .	1431
3.172.6 Sympy [F] . . . . .	1432
3.172.7 Maxima [A] (verification not implemented) . . . . .	1432
3.172.8 Giac [F(-2)] . . . . .	1433
3.172.9 Mupad [F(-1)] . . . . .	1433

#### 3.172.1 Optimal result

Integrand size = 27, antiderivative size = 186

$$\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 dx = -\frac{14b^2\sqrt{d - c^2dx^2}}{27c^2} + \frac{2}{27}b^2x^2\sqrt{d - c^2dx^2} + \frac{2bx\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{3c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{2bcx^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{9\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2}{3c^2d}$$

output

```
-1/3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/c^2/d-14/27*b^2*(-c^2*d*x^2+d)^(1/2)/c^2+2/27*b^2*x^2*(-c^2*d*x^2+d)^(1/2)+2/3*b*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/9*b*c*x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.172.2 Mathematica [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.97

$$\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx$$

$$= \frac{\sqrt{d - c^2 dx^2} \left( -6abcx\sqrt{-1 + cx}\sqrt{1 + cx}(-3 + c^2 x^2) + 9a^2(-1 + c^2 x^2)^2 + 2b^2(7 - 8c^2 x^2 + c^4 x^4) + 6b(bc x \operatorname{barccosh}(cx) + 9b^2(-1 + c^2 x^2)^2 \operatorname{ArcCosh}[cx]^2) \right)}{27c^2(-1 + c^2 x^2)}$$

input `Integrate[x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`output `(Sqrt[d - c^2*d*x^2]*(-6*a*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-3 + c^2*x^2) + 9*a^2*(-1 + c^2*x^2)^2 + 2*b^2*(7 - 8*c^2*x^2 + c^4*x^4) + 6*b*(b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(3 - c^2*x^2) + 3*a*(-1 + c^2*x^2)^2)*ArcCosh[c*x] + 9*b^2*(-1 + c^2*x^2)^2*ArcCosh[c*x]^2))/(27*c^2*(-1 + c^2*x^2))`**3.172.3 Rubi [A] (verified)**Time = 0.57 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.88, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.259$ , Rules used = {6329, 25, 6304, 6309, 27, 960, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx$$

$$\downarrow 6329$$

$$-\frac{2b\sqrt{d - c^2 dx^2} \int -((1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))) dx}{3c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3c^2 d}$$

$$\downarrow 25$$

$$\frac{2b\sqrt{d - c^2 dx^2} \int (1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx)) dx}{3c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3c^2 d}$$

$$\downarrow 6304$$

$$\frac{2b\sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)(a + \operatorname{barccosh}(cx)) dx}{3c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3c^2 d}$$

---

 3.172.  $\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{array}{c} \downarrow 6309 \\ \frac{2b\sqrt{d-c^2dx^2}\left(-bc\int\frac{x(3-c^2x^2)}{3\sqrt{cx-1}\sqrt{cx+1}}dx-\frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}} \\ \frac{3c^2d}{3c^2d} \end{array}$$

$$\begin{array}{c} \downarrow 27 \\ \frac{2b\sqrt{d-c^2dx^2}\left(-\frac{1}{3}bc\int\frac{x(3-c^2x^2)}{\sqrt{cx-1}\sqrt{cx+1}}dx-\frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}} \\ \frac{3c^2d}{3c^2d} \end{array}$$

$$\begin{array}{c} \downarrow 960 \\ \frac{2b\sqrt{d-c^2dx^2}\left(-\frac{1}{3}bc\left(\frac{7}{3}\int\frac{x}{\sqrt{cx-1}\sqrt{cx+1}}dx-\frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1}\right)-\frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}} \\ \frac{3c^2d}{3c^2d} \end{array}$$

$$\begin{array}{c} \downarrow 83 \\ \frac{2b\sqrt{d-c^2dx^2}\left(-\frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))-\frac{1}{3}bc\left(\frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2}-\frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1}\right)\right)}{\frac{3c\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}} \\ \frac{3c^2d}{3c^2d} \end{array}$$

input `Int[x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `-1/3*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/(c^2*d) + (2*b*Sqrt[d - c^2*d*x^2]*(-1/3*(b*c*((7*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2) - (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/3)) + x*(a + b*ArcCosh[c*x]) - (c^2*x^3*(a + b*ArcCosh[c*x]))/3)/(3*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.172.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 960 `Int[((e_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x])^n, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`
- rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

**3.172.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 725 vs.  $2(158) = 316$ .

Time = 0.56 (sec) , antiderivative size = 726, normalized size of antiderivative = 3.90

method	result
default	$-\frac{a^2(-c^2dx^2+d)^{\frac{3}{2}}}{3c^2d} + b^2 \left( \frac{\sqrt{-d(c^2x^2-1)}(4c^4x^4-5c^2x^2+4\sqrt{cx-1}\sqrt{cx+1}c^3x^3-3\sqrt{cx-1}\sqrt{cx+1}cx+1)(9\operatorname{arccosh}(cx)^2-6\operatorname{arccosh}(cx))}{216(cx+1)c^2(cx-1)} \right)$
parts	$-\frac{a^2(-c^2dx^2+d)^{\frac{3}{2}}}{3c^2d} + b^2 \left( \frac{\sqrt{-d(c^2x^2-1)}(4c^4x^4-5c^2x^2+4\sqrt{cx-1}\sqrt{cx+1}c^3x^3-3\sqrt{cx-1}\sqrt{cx+1}cx+1)(9\operatorname{arccosh}(cx)^2-6\operatorname{arccosh}(cx))}{216(cx+1)c^2(cx-1)} \right)$

input `int(x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -1/3*a^2*(-c^2*d*x^2+d)^(3/2)/c^2/d+b^2*(1/216*(-d*(c^2*x^2-1))^(1/2)*(4*c \\ & ^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c* \\ & x+1)^(1/2)*c*x+1)*(9*arccosh(c*x)^2-6*arccosh(c*x)+2)/(c*x+1)/c^2/(c*x-1)- \\ & 1/8*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(ar \\ & ccosh(c*x)^2-2*arccosh(c*x)+2)/(c*x+1)/c^2/(c*x-1)-1/8*(-d*(c^2*x^2-1))^(1/2) \\ & *(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2+2*arccosh \\ & (c*x)+2)/(c*x+1)/c^2/(c*x-1)+1/216*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2) \\ &)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2* \\ & x^2+1)*(9*arccosh(c*x)^2+6*arccosh(c*x)+2)/(c*x+1)/c^2/(c*x-1))+2*a*b*(1/7 \\ & 2*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2) \\ &)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(-1+3*arccosh(c*x))/(c*x+1) \\ & /c^2/(c*x-1)-1/8*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c \\ & ^2*x^2-1)*(-1+arccosh(c*x))/(c*x+1)/c^2/(c*x-1)-1/8*(-d*(c^2*x^2-1))^(1/2) \\ & *(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))/(c*x+1)/c^2 \\ & /(c*x-1)+1/72*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x \\ & ^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(1+3*arccosh(c \\ & *x))/(c*x+1)/c^2/(c*x-1) \end{aligned}$$
**3.172.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.51

$$\int x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 dx$$

$$= \frac{9(b^2c^4x^4-2b^2c^2x^2+b^2)\sqrt{-c^2dx^2+d}\log(cx+\sqrt{c^2x^2-1})^2-6(abc^3x^3-3abcx)\sqrt{-c^2dx^2+d}\sqrt{c^2x^2-1}}{216}$$



input `integrate(x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `1/27*(9*(b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1))^2 - 6*(a*b*c^3*x^3 - 3*a*b*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 6*((b^2*c^3*x^3 - 3*b^2*c*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 3*(a*b*c^4*x^4 - 2*a*b*c^2*x^2 + a*b)*sqrt(-c^2*d*x^2 + d))*log(c*x + sqrt(c^2*x^2 - 1)) + ((9*a^2 + 2*b^2)*c^4*x^4 - 2*(9*a^2 + 8*b^2)*c^2*x^2 + 9*a^2 + 14*b^2)*sqrt(-c^2*d*x^2 + d))/(c^4*x^2 - c^2)`

### 3.172.6 Sympy [F]

$$\int x\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))^2 dx = \int x\sqrt{-d(cx - 1)(cx + 1)}(a + b \operatorname{acosh}(cx))^2 dx$$

input `integrate(x*(a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2, x)`

### 3.172.7 Maxima [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.10

$$\begin{aligned} & \int x\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))^2 dx \\ &= \frac{2}{27} b^2 \left( \frac{\sqrt{c^2 x^2 - 1} \sqrt{-d} dx^2 - \frac{7\sqrt{c^2 x^2 - 1} \sqrt{-d}}{c^2}}{d} - \frac{3(c^2 \sqrt{-d} dx^3 - 3\sqrt{-d} dx) \operatorname{arccosh}(cx)}{cd} \right) \\ & \quad - \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} b^2 \operatorname{arccosh}(cx)^2}{3c^2 d} - \frac{2(-c^2 dx^2 + d)^{\frac{3}{2}} ab \operatorname{arccosh}(cx)}{3c^2 d} \\ & \quad - \frac{2(c^2 \sqrt{-d} dx^3 - 3\sqrt{-d} dx) ab}{9cd} - \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} a^2}{3c^2 d} \end{aligned}$$

input `integrate(x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output  $\frac{2}{27}b^2\left(\sqrt{c^2x^2 - 1}\sqrt{-d}dx^2 - 7\sqrt{c^2x^2 - 1}\sqrt{-d}\frac{d}{c^2}\right) - 3\left(c^2\sqrt{-d}dx^3 - 3\sqrt{-d}dx\right)\operatorname{arccosh}(cx)/(cd) - \frac{1}{3}(-c^2dx^2 + d)^{3/2}b^2\operatorname{arccosh}(cx)^2/(c^2d) - \frac{2}{3}(-c^2dx^2 + d)^{3/2}ab\operatorname{arccosh}(cx)/(c^2d) - \frac{2}{9}(c^2\sqrt{-d}dx^3 - 3\sqrt{-d}dx)ab/(cd) - \frac{1}{3}(-c^2dx^2 + d)^{3/2}a^2/(c^2d)$

### 3.172.8 Giac [F(-2)]

Exception generated.

$$\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate(x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.172.9 Mupad [F(-1)]

Timed out.

$$\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 dx = \int x(a + b\operatorname{acosh}(cx))^2\sqrt{d - c^2dx^2} dx$$

input `int(x*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2),x)`

output `int(x*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2), x)`

### 3.173 $\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$

3.173.1 Optimal result	1434
3.173.2 Mathematica [A] (warning: unable to verify)	1435
3.173.3 Rubi [A] (verified)	1435
3.173.4 Maple [B] (verified)	1438
3.173.5 Fricas [F]	1438
3.173.6 Sympy [F]	1439
3.173.7 Maxima [F]	1439
3.173.8 Giac [F(-2)]	1439
3.173.9 Mupad [F(-1)]	1440

#### 3.173.1 Optimal result

Integrand size = 26, antiderivative size = 204

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{1}{4} b^2 x \sqrt{d - c^2 dx^2} + \frac{b^2 \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{4c \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{bcx^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{2 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{1}{2} x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^3}{6bc \sqrt{-1 + cx} \sqrt{1 + cx}}$$

output  $\frac{1}{4} b^2 x (-c^2 d x^2 + d)^{1/2} + \frac{1}{2} x (a + b \operatorname{arccosh}(c x))^2 (-c^2 d x^2 + d)^{1/2} + \frac{1}{4} b^2 \operatorname{arccosh}(c x) (-c^2 d x^2 + d)^{1/2} / c / (c x - 1)^{1/2} / (c x + 1)^{1/2} - \frac{1}{2} b c x^2 (a + b \operatorname{arccosh}(c x)) (-c^2 d x^2 + d)^{1/2} / (c x - 1)^{1/2} / (c x + 1)^{1/2} - \frac{1}{6} (a + b \operatorname{arccosh}(c x))^3 (-c^2 d x^2 + d)^{1/2} / b / c / (c x - 1)^{1/2} / (c x + 1)^{1/2}$

**3.173.2 Mathematica [A] (warning: unable to verify)**

Time = 1.16 (sec) , antiderivative size = 235, normalized size of antiderivative = 1.15

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{1}{24} \left( 12a^2 x \sqrt{d - c^2 dx^2} - \frac{12a^2 \sqrt{d} \arctan\left(\frac{cx \sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right)}{c} \right. \\ \left. - \frac{6ab \sqrt{d - c^2 dx^2} (2 \operatorname{arccosh}(cx))^2 + \cosh(2 \operatorname{arccosh}(cx)) - 2 \operatorname{arccosh}(cx) \sinh(2 \operatorname{arccosh}(cx))}{c \sqrt{\frac{-1 + cx}{1 + cx}} (1 + cx)} \right. \\ \left. + \frac{b^2 \sqrt{d - c^2 dx^2} (-4 \operatorname{arccosh}(cx))^3 - 6 \operatorname{arccosh}(cx) \cosh(2 \operatorname{arccosh}(cx)) + (3 + 6 \operatorname{arccosh}(cx))^2 \sinh(2 \operatorname{arccosh}(cx))}{c \sqrt{\frac{-1 + cx}{1 + cx}} (1 + cx)} \right)$$

input `Integrate[Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`output `(12*a^2*x*Sqrt[d - c^2*d*x^2] - (12*a^2*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))])/c - (6*a*b*Sqrt[d - c^2*d*x^2]*(2*ArcCosh[c*x]^2 + Cosh[2*ArcCosh[c*x]] - 2*ArcCosh[c*x]*Sinh[2*ArcCosh[c*x]]))/(c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + (b^2*Sqrt[d - c^2*d*x^2]*(-4*ArcCosh[c*x]^3 - 6*ArcCosh[c*x]*Cosh[2*ArcCosh[c*x]] + (3 + 6*ArcCosh[c*x]^2)*Sinh[2*ArcCosh[c*x]]))/(c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))/24`**3.173.3 Rubi [A] (verified)**Time = 0.75 (sec) , antiderivative size = 181, normalized size of antiderivative = 0.89, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$ , Rules used = {6310, 6298, 101, 43, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx \\ \downarrow \text{6310} \\ - \frac{bc \sqrt{d - c^2 dx^2} \int x(a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx - 1} \sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{2 \sqrt{cx - 1} \sqrt{cx + 1}} + \\ \frac{1}{2} x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2$$

$$\begin{aligned}
& \downarrow 6298 \\
& \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))-\frac{1}{2}bc\int\frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{\sqrt{d-c^2dx^2}\int\frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}}dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 101 \\
& \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))-\frac{1}{2}bc\left(\frac{\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx}{2c^2}+\frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)\right)}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{\sqrt{d-c^2dx^2}\int\frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}}dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 43 \\
& \frac{\sqrt{d-c^2dx^2}\int\frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}}dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \\
& \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))-\frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}+\frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)\right)}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 6308 \\
& \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \\
& \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))-\frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}+\frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)\right)}{\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `(x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^3)/(6*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (b*c*Sqrt[d - c^2*d*x^2]*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/2)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.173.3.1 Defintions of rubi rules used

- rule 43  $\text{Int}[1/(\text{Sqrt}[(a\_)+(b\_)(x\_)]*\text{Sqrt}[(c\_)+(d\_)(x\_)]), x\_Symbol] \rightarrow \text{Simp}[\text{ArcCosh}[b*(x/a)/(b*\text{Sqrt}[d/b]), x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{EqQ}[b*c + a*d, 0] \&\& \text{GtQ}[a, 0] \&\& \text{GtQ}[d/b, 0]$
- rule 101  $\text{Int}[(a\_)+(b\_)(x\_)]^{n\_}*((c\_)+(d\_)(x\_)]^{m\_}*((e\_)+(f\_)(x\_)]^{p\_}, x_] \rightarrow \text{Simp}[b*(a + b*x)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 3))), x] + \text{Simp}[1/(d*f*(n + p + 3)) \text{Int}[(c + d*x)^n*(e + f*x)^p*\text{Simp}[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \text{NeQ}[n + p + 3, 0]$
- rule 6298  $\text{Int}[(a\_)+(b\_)(x\_)]^{n\_}*((c\_)+(d\_)(x\_)]^{m\_}, x\_Symbol] \rightarrow \text{Simp}[(d*x)^{(m + 1)}*((a + b*\text{ArcCosh}[c*x])^n/(d*(m + 1))), x] - \text{Simp}[b*c*(n/(d*(m + 1))) \text{Int}[(d*x)^{(m + 1)}*((a + b*\text{ArcCosh}[c*x])^{(n - 1)}/(\text{Sqrt}[1 + c*x]*\text{Sqrt}[-1 + c*x])), x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x] \&\& \text{IGtQ}[n, 0] \& \& \text{NeQ}[m, -1]$
- rule 6308  $\text{Int}[(a\_)+(b\_)(x\_)]^{n\_}/(\text{Sqrt}[(d1\_)+(e1\_)(x\_)]*\text{Sqrt}[(d2\_)+(e2\_)(x\_)]), x\_Symbol] \rightarrow \text{Simp}[(1/(b*c*(n + 1)))*\text{Simp}[\text{Sqrt}[1 + c*x]/\text{Sqrt}[d1 + e1*x]]*\text{Simp}[\text{Sqrt}[-1 + c*x]/\text{Sqrt}[d2 + e2*x]]*(a + b*\text{ArcCosh}[c*x])^{(n + 1)}, x] /; \text{FreeQ}\{a, b, c, d1, e1, d2, e2, n\}, x] \&\& \text{EqQ}[e1, c*d1] \&\& \text{EqQ}[e2, (-c)*d2] \&\& \text{NeQ}[n, -1]$
- rule 6310  $\text{Int}[(a\_)+(b\_)(x\_)]^{n\_}*\text{Sqrt}[(d\_)+(e\_)(x_)^2], x\_Symbol] \rightarrow \text{Simp}[x*\text{Sqrt}[d + e*x^2]*((a + b*\text{ArcCosh}[c*x])^{n/2}), x] + (-\text{Simp}[(1/2)*\text{Simp}[\text{Sqrt}[d + e*x^2]/(\text{Sqrt}[1 + c*x]*\text{Sqrt}[-1 + c*x])] \text{Int}[(a + b*\text{ArcCosh}[c*x])^n/(\text{Sqrt}[1 + c*x]*\text{Sqrt}[-1 + c*x]), x], x] - \text{Simp}[b*c*(n/2)*\text{Simp}[\text{Sqrt}[d + e*x^2]/(\text{Sqrt}[1 + c*x]*\text{Sqrt}[-1 + c*x])] \text{Int}[x*(a + b*\text{ArcCosh}[c*x])^{(n - 1)}, x], x]) /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[c^2*d + e, 0] \&\& \text{GtQ}[n, 0]$

### 3.173.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 526 vs.  $2(172) = 344$ .

Time = 0.77 (sec) , antiderivative size = 527, normalized size of antiderivative = 2.58

method	result
default	$\frac{a^2 x \sqrt{-c^2 d x^2 + d}}{2} + \frac{a^2 d \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{2\sqrt{c^2 d}} + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(cx)^3}{6\sqrt{cx-1}\sqrt{cx+1}c} + \frac{\sqrt{-d(c^2 x^2 - 1)} (2c^3 x^3 - 2cx + 2\sqrt{cx-1})}{6\sqrt{cx-1}\sqrt{cx+1}c} \right)$
parts	$\frac{a^2 x \sqrt{-c^2 d x^2 + d}}{2} + \frac{a^2 d \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{2\sqrt{c^2 d}} + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(cx)^3}{6\sqrt{cx-1}\sqrt{cx+1}c} + \frac{\sqrt{-d(c^2 x^2 - 1)} (2c^3 x^3 - 2cx + 2\sqrt{cx-1})}{6\sqrt{cx-1}\sqrt{cx+1}c} \right)$

input `int((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & 1/2*a^2*x*(-c^2*d*x^2+d)^(1/2)+1/2*a^2*d/(c^2*d)^(1/2)*\arctan((c^2*d)^(1/2) \\ & )*x/(-c^2*d*x^2+d)^(1/2))+b^2*(-1/6*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/( \\ & c*x+1)^(1/2)/c*\operatorname{arccosh}(c*x)^3+1/16*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x \\ & +2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(2*\operatorname{arc} \\ & \operatorname{cosh}(c*x)^2-2*\operatorname{arccosh}(c*x)+1)/(c*x-1)/(c*x+1)/c+1/16*(-d*(c^2*x^2-1))^(1/2) \\ & )*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^( \\ & (1/2)-2*c*x)*(2*\operatorname{arccosh}(c*x)^2+2*\operatorname{arccosh}(c*x)+1)/(c*x-1)/(c*x+1)/c)+2*a*b* \\ & (-1/4*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/c*\operatorname{arccosh}(c*x)^2+ \\ & 1/16*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2) \\ & *c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+2*\operatorname{arccosh}(c*x))/(c*x-1)/(c*x+1)/ \\ & c+1/16*(-d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^ \\ & 3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)-2*c*x)*(1+2*\operatorname{arccosh}(c*x))/(c*x-1)/(c*x+1) \\ & )/c \end{aligned}$$

### 3.173.5 Fricas [F]

$$\int \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^2 dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x,algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

**3.173.6 Sympy [F]**

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2 dx$$

input `integrate((a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2, x)`

**3.173.7 Maxima [F]**

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^2 dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `1/2*(sqrt(-c^2*d*x^2 + d)*x + sqrt(d)*arcsin(c*x)/c)*a^2 + integrate(sqrt(-c^2*d*x^2 + d)*b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2 + 2*sqrt(-c^2*d*x^2 + d)*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1), x)`

**3.173.8 Giac [F(-2)]**

Exception generated.

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.173.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 \sqrt{d - c^2 dx^2} dx$$

input `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2),x)`output `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2), x)`

**3.174**  $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x} dx$

3.174.1 Optimal result . . . . . 1441  
 3.174.2 Mathematica [A] (warning: unable to verify) . . . . . 1442  
 3.174.3 Rubi [A] (verified) . . . . . 1443  
 3.174.4 Maple [F] . . . . . 1446  
 3.174.5 Fracas [F] . . . . . 1446  
 3.174.6 Sympy [F] . . . . . 1447  
 3.174.7 Maxima [F] . . . . . 1447  
 3.174.8 Giac [F(-2)] . . . . . 1447  
 3.174.9 Mupad [F(-1)] . . . . . 1448

**3.174.1 Optimal result**

Integrand size = 29, antiderivative size = 402

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x} dx$$

$$= 2b^2\sqrt{d-c^2dx^2} - \frac{2abcx\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}} - \frac{2b^2cx\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 - \frac{2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2ib\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2ib\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2ib^2\sqrt{d-c^2dx^2} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{2ib^2\sqrt{d-c^2dx^2} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```

2*b^2*(-c^2*d*x^2+d)^(1/2)+(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)-2*a*b
*c*x*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*b^2*c*x*arccosh(c*
x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*(a+b*arccosh(c*x))^2
*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2
)/(c*x+1)^(1/2)+2*I*b*(a+b*arccosh(c*x))*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(
c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*I*b*(a+b
*arccosh(c*x))*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+
d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*I*b^2*polylog(3,-I*(c*x+(c*x-1)^(1/
2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2*I*b^
2*polylog(3,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x
-1)^(1/2)/(c*x+1)^(1/2)

```

### 3.174.2 Mathematica [A] (warning: unable to verify)

Time = 1.23 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.12

$$\begin{aligned}
 & \int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x} dx \\
 &= a^2 \sqrt{d - c^2 dx^2} + a^2 \sqrt{d} \log(cx) - a^2 \sqrt{d} \log\left(d + \sqrt{d} \sqrt{d - c^2 dx^2}\right) \\
 &+ \frac{2ab \sqrt{d - c^2 dx^2} \left(-cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + i \operatorname{arccosh}(cx) \log\left(1 - ie^{-\operatorname{arccosh}(cx)}\right)\right)}{\sqrt{\frac{-1+cx}{1+cx}}} \\
 &+ b^2 \sqrt{d - c^2 dx^2} \left(2 + \frac{2cx \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx)}{1 - cx} + \operatorname{arccosh}(cx)^2\right) \\
 &+ \frac{i(\operatorname{arccosh}(cx))^2 \log\left(1 - ie^{-\operatorname{arccosh}(cx)}\right) - \operatorname{arccosh}(cx)^2 \log\left(1 + ie^{-\operatorname{arccosh}(cx)}\right) + 2 \operatorname{arccosh}(cx) \operatorname{PolyLog}\left(2, \frac{-1+cx}{1+cx}\right)}{\sqrt{\frac{-1+cx}{1+cx}}}
 \end{aligned}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x,x]`

output

```

a^2*Sqrt[d - c^2*d*x^2] + a^2*Sqrt[d]*Log[c*x] - a^2*Sqrt[d]*Log[d + Sqrt[
d]*Sqrt[d - c^2*d*x^2]] + (2*a*b*Sqrt[d - c^2*d*x^2]*(-(c*x) + Sqrt[(-1 +
c*x)/(1 + c*x)]*ArcCosh[c*x] + c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]
+ I*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh[c*x]*Log[1 + I/E^A
rcCosh[c*x]] + I*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*PolyLog[2, I/E^ArcCos
h[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + b^2*Sqrt[d - c^2*d*x^2]
*(2 + (2*c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x])/(1 - c*x) + ArcCosh[
c*x]^2 + (I*(ArcCosh[c*x]^2*Log[1 - I/E^ArcCosh[c*x]] - ArcCosh[c*x]^2*Log
[1 + I/E^ArcCosh[c*x]] + 2*ArcCosh[c*x]*PolyLog[2, (-I)/E^ArcCosh[c*x]] -
2*ArcCosh[c*x]*PolyLog[2, I/E^ArcCosh[c*x]] + 2*PolyLog[3, (-I)/E^ArcCosh[
c*x]] - 2*PolyLog[3, I/E^ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 +
c*x)))

```

### 3.174.3 Rubi [A] (verified)

Time = 1.33 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.59, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$ , Rules used = {6341, 2009, 6362, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx \\
 & \quad \downarrow \text{6341} \\
 & -\frac{2bc\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{\sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 - \\
 & \quad \frac{2bc\sqrt{d - c^2 dx^2} \left( ax + b\operatorname{barccosh}(cx) - \frac{b\sqrt{cx - 1}\sqrt{cx + 1}}{c} \right)}{\sqrt{cx - 1}\sqrt{cx + 1}} \\
 & \quad \downarrow \text{6362}
 \end{aligned}$$

---

3.174.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx$

$$\begin{aligned}
& - \frac{\sqrt{d - c^2 dx^2} \int \frac{(a + b \operatorname{arccosh}(cx))^2}{cx} d \operatorname{arccosh}(cx)}{\sqrt{cx - 1} \sqrt{cx + 1}} + \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - \\
& \frac{2bc \sqrt{d - c^2 dx^2} \left( ax + b \operatorname{arccosh}(cx) - \frac{b \sqrt{cx - 1} \sqrt{cx + 1}}{c} \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{3042} \\
& - \frac{\sqrt{d - c^2 dx^2} \int (a + b \operatorname{arccosh}(cx))^2 \csc \left( i \operatorname{arccosh}(cx) + \frac{\pi}{2} \right) d \operatorname{arccosh}(cx)}{\sqrt{cx - 1} \sqrt{cx + 1}} + \sqrt{d - c^2 dx^2} (a + \\
& \operatorname{arccosh}(cx))^2 - \frac{2bc \sqrt{d - c^2 dx^2} \left( ax + b \operatorname{arccosh}(cx) - \frac{b \sqrt{cx - 1} \sqrt{cx + 1}}{c} \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{4668} \\
& - \frac{\sqrt{d - c^2 dx^2} \left( -2ib \int (a + b \operatorname{arccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) + 2ib \int (a + b \operatorname{arccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - \frac{2bc \sqrt{d - c^2 dx^2} \left( ax + b \operatorname{arccosh}(cx) - \frac{b \sqrt{cx - 1} \sqrt{cx + 1}}{c} \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{3011} \\
& - \frac{\sqrt{d - c^2 dx^2} \left( 2ib (b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) d \operatorname{arccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx))) \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - \frac{2bc \sqrt{d - c^2 dx^2} \left( ax + b \operatorname{arccosh}(cx) - \frac{b \sqrt{cx - 1} \sqrt{cx + 1}}{c} \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{2720} \\
& - \frac{\sqrt{d - c^2 dx^2} \left( 2ib (b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx))) \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - \frac{2bc \sqrt{d - c^2 dx^2} \left( ax + b \operatorname{arccosh}(cx) - \frac{b \sqrt{cx - 1} \sqrt{cx + 1}}{c} \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \quad \downarrow \text{7143} \\
& - \frac{\sqrt{d - c^2 dx^2} \left( 2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + b \operatorname{arccosh}(cx))^2 + 2ib (b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \sqrt{cx - 1} \sqrt{cx + 1}) \right)}{\sqrt{cx - 1} \sqrt{cx + 1}} \\
& \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - \frac{2bc \sqrt{d - c^2 dx^2} \left( ax + b \operatorname{arccosh}(cx) - \frac{b \sqrt{cx - 1} \sqrt{cx + 1}}{c} \right)}{\sqrt{cx - 1} \sqrt{cx + 1}}
\end{aligned}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x,x]`

$$3.174. \quad \int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x} dx$$

```
output Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2 - (2*b*c*Sqrt[d - c^2*d*x^2]*(a
*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(Sqrt[-1 + c*
x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[
E^ArcCosh[c*x]] + (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCos
h[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-((a + b*ArcCosh[
c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]])))/(S
qrt[-1 + c*x]*Sqrt[1 + c*x])
```

### 3.174.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 3011 Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4668 Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_
))^m_, x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6362 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.174.4 Maple [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2 \sqrt{-c^2 dx^2 + d}}{x} dx$$

input `int((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x,x)`

output `int((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x,x)`

### 3.174.5 Fracas [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^2}{x} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/x, x)`

### 3.174.6 Sympy [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2}{x} dx$$

input `integrate((a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2)/x,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2/x, x)`

### 3.174.7 Maxima [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^2}{x} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="maxima")`

output `-(sqrt(d)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - sqrt(-c^2*d*x^2 + d))*a^2 + integrate(sqrt(-c^2*d*x^2 + d)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x + 2*sqrt(-c^2*d*x^2 + d)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x, x)`

### 3.174.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x} dx = \text{Exception raised: TypeError}$$



input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.174.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{x} dx = \int \frac{(a+b\operatorname{acosh}(cx))^2\sqrt{d-c^2dx^2}}{x} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x, x)`

**3.175**  $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx$

3.175.1 Optimal result . . . . . 1449  
 3.175.2 Mathematica [A] (warning: unable to verify) . . . . . 1450  
 3.175.3 Rubi [C] (warning: unable to verify) . . . . . 1451  
 3.175.4 Maple [B] (verified) . . . . . 1455  
 3.175.5 Fricas [F] . . . . . 1455  
 3.175.6 Sympy [F] . . . . . 1456  
 3.175.7 Maxima [F] . . . . . 1456  
 3.175.8 Giac [F(-2)] . . . . . 1456  
 3.175.9 Mupad [F(-1)] . . . . . 1457

**3.175.1 Optimal result**

Integrand size = 29, antiderivative size = 234

$$\begin{aligned} & \int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx \\ &= -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x} + \frac{c\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{c\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^3}{3b\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{2bc\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \log(1+e^{-2\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{b^2c\sqrt{d-c^2dx^2} \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output

```
-(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x+c*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/3*c*(a+b*arccosh(c*x))^3*(-c^2*d*x^2+d)^(1/2)/b/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2*b*c*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b^2*c*polylog(2,-1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.175.2 Mathematica [A] (warning: unable to verify)**

Time = 1.69 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.15

$$\begin{aligned}
& \int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^2} dx \\
&= -\frac{a^2 \sqrt{d - c^2 dx^2}}{x} + a^2 c \sqrt{d} \arctan \left( \frac{cx \sqrt{d - c^2 dx^2}}{\sqrt{d} (-1 + c^2 x^2)} \right) \\
&+ abc \sqrt{d - c^2 dx^2} \left( -\frac{2 \operatorname{arccosh}(cx)}{cx} + \frac{\operatorname{arccosh}(cx)^2 + 2 \log(cx)}{\sqrt{\frac{-1+cx}{1+cx}} (1+cx)} \right) \\
&+ \frac{1}{3} b^2 c \sqrt{d - c^2 dx^2} \left( \operatorname{arccosh}(cx) \left( -\frac{3 \operatorname{arccosh}(cx)}{cx} \right. \right. \\
&\quad \left. \left. + \frac{\operatorname{arccosh}(cx)(3 + \operatorname{arccosh}(cx)) + 6 \log(1 + e^{-2 \operatorname{arccosh}(cx)})}{\sqrt{\frac{-1+cx}{1+cx}} (1+cx)} \right) \right. \\
&\quad \left. + \frac{3 \sqrt{\frac{-1+cx}{1+cx}} \operatorname{PolyLog}(2, -e^{-2 \operatorname{arccosh}(cx)})}{1 - cx} \right)
\end{aligned}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x^2,x]`output `-((a^2*Sqrt[d - c^2*d*x^2])/x) + a^2*c*Sqrt[d]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + a*b*c*Sqrt[d - c^2*d*x^2]*((-2*ArcCosh[c*x])/(c*x) + (ArcCosh[c*x]^2 + 2*Log[c*x])/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))) + (b^2*c*Sqrt[d - c^2*d*x^2]*(ArcCosh[c*x]*((-3*ArcCosh[c*x])/(c*x) + (ArcCosh[c*x]*(3 + ArcCosh[c*x]) + 6*Log[1 + E^(-2*ArcCosh[c*x])]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))) + (3*Sqrt[(-1 + c*x)/(1 + c*x)]*PolyLog[2, -E^(-2*ArcCosh[c*x])])/(1 - c*x))/3`

**3.175.3 Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 1.85 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.79, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$ , Rules used = {6339, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx \\
 & \quad \downarrow \text{6339} \\
 & \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{2bc\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{6297} \\
 & \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \frac{2c\sqrt{d-c^2dx^2} \int -\left((a+\operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right)\right) d(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{25} \\
 & \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \frac{2c\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) d(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.175.  $\int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx$

$$\begin{aligned}
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{2ic\sqrt{d - c^2 dx^2} \int -i(a + \operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \frac{\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2} \\
& \quad \downarrow \text{26} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{2ic\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \frac{\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2} \\
& \quad \downarrow \text{4201} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{2ic\sqrt{d - c^2 dx^2} \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 + e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2 \right)}{\sqrt{cx-1}\sqrt{cx+1}} \\
& \frac{\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2} \\
& \quad \downarrow \text{2620} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{2ic\sqrt{d - c^2 dx^2} \left( 2i\left(\frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2} \\
& \quad \downarrow \text{2715} \\
& \frac{2ic\sqrt{d - c^2 dx^2} \left( 2i\left(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2} \\
& \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{x} \\
& \quad \downarrow \text{2838}
\end{aligned}$$

---

3.175.  $\int \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{x^2} dx$

$$\frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{2ic\sqrt{d - c^2 dx^2} \left( 2i \left( \frac{1}{4} b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) \right) - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{2} i (a + b)}{\sqrt{cx-1}\sqrt{cx+1}}}{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}$$

x  
↓ 6308

$$\frac{2ic\sqrt{d - c^2 dx^2} \left( 2i \left( \frac{1}{4} b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) \right) - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{2} i (a + b)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{c\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^3}{3b\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x^2,x]`

output `-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x) + (c*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^3)/(3*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((2*I)*c*Sqrt[d - c^2*d*x^2]*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]]/4)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.175.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(x_), x_Symbol] :> Simp[1/b
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a
, b, c}, x] && IGtQ[n, 0]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sq
rt[(d2_) + (e2_.)*(x_)]), x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 +
c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[
c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1
] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6339 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*Sqrt[(d_) +
(e_.)*(x_)^2], x_Symbol] :> Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^
2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`

### 3.175.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 581 vs.  $2(232) = 464$ .

Time = 0.86 (sec) , antiderivative size = 582, normalized size of antiderivative = 2.49

method	result
default	$-\frac{a^2(-c^2dx^2+d)^{\frac{3}{2}}}{dx} - a^2c^2x\sqrt{-c^2dx^2+d} - \frac{a^2c^2d \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{\sqrt{c^2d}} + \frac{b^2\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^3c}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{b^2\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2c}{3\sqrt{cx-1}\sqrt{cx+1}}$
parts	$-\frac{a^2(-c^2dx^2+d)^{\frac{3}{2}}}{dx} - a^2c^2x\sqrt{-c^2dx^2+d} - \frac{a^2c^2d \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{\sqrt{c^2d}} + \frac{b^2\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^3c}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{b^2\sqrt{-d(c^2x^2-1)} \operatorname{arccosh}(cx)^2c}{3\sqrt{cx-1}\sqrt{cx+1}}$

input `int((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & -a^2/d/x*(-c^2*d*x^2+d)^{(3/2)} - a^2*c^2*x*(-c^2*d*x^2+d)^{(1/2)} - a^2*c^2*d/(c^2*d)^{(1/2)} * \arctan((c^2*d)^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)}) + 1/3*b^2*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} * \operatorname{arccosh}(c*x)^3*c - b^2*(-d*(c^2*x^2-1))^{(1/2)} * \operatorname{arccosh}(c*x)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} * c - b^2*(-d*(c^2*x^2-1))^{(1/2)} * \operatorname{arccosh}(c*x)^2*x/(c*x-1)/(c*x+1) * c^2 + b^2*(-d*(c^2*x^2-1))^{(1/2)} * \operatorname{arccosh}(c*x)^2/x/(c*x-1)/(c*x+1) + 2*b^2*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} * \operatorname{arccosh}(c*x) * \ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2) * c + b^2*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} * \operatorname{polylog}(2, -(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2) * c + a*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} * \operatorname{arccosh}(c*x)^2 * c - 2*a*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} * \operatorname{arccosh}(c*x) * c - 2*a*b*(-d*(c^2*x^2-1))^{(1/2)} * \operatorname{arccosh}(c*x) * x/(c*x-1)/(c*x+1) * c^2 + 2*a*b*(-d*(c^2*x^2-1))^{(1/2)} * \operatorname{arccosh}(c*x) / x / (c*x-1)/(c*x+1) + 2*a*b*(-d*(c^2*x^2-1))^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} * \ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))^2) * c \end{aligned}$$

### 3.175.5 Fracas [F]

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx = \int \frac{\sqrt{-c^2dx^2+d}(b\operatorname{arccosh}(cx)+a)^2}{x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2+d)*(b^2*arccosh(c*x)^2+2*a*b*arccosh(c*x)+a^2)/x^2,x)`

---

3.175. 
$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx$$



**3.175.6 Sympy [F]**

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2}{x^2} dx$$

input `integrate((a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2)/x**2,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2/x**2, x)`

**3.175.7 Maxima [F]**

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^2}{x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="maxima")`

output `-(c*sqrt(d)*arcsin(c*x) + sqrt(-c^2*d*x^2 + d)/x)*a^2 + integrate(sqrt(-c^2*d*x^2 + d)*b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/x^2 + 2*sqrt(-c^2*d*x^2 + d)*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/x^2, x)`

**3.175.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.175.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 \sqrt{d - c^2 dx^2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x^2,x)`output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x^2, x)`

**3.176**  $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx$

3.176.1 Optimal result . . . . . 1458  
 3.176.2 Mathematica [B] (warning: unable to verify) . . . . . 1459  
 3.176.3 Rubi [A] (verified) . . . . . 1459  
 3.176.4 Maple [F] . . . . . 1463  
 3.176.5 Fracas [F] . . . . . 1463  
 3.176.6 Sympy [F] . . . . . 1464  
 3.176.7 Maxima [F] . . . . . 1464  
 3.176.8 Giac [F(-2)] . . . . . 1464  
 3.176.9 Mupad [F(-1)] . . . . . 1465

**3.176.1 Optimal result**

Integrand size = 29, antiderivative size = 427

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx$$

$$= -\frac{bc\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x\sqrt{-1+cx}\sqrt{1+cx}} - \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2x^2}$$

$$+ \frac{c^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{b^2c^2\sqrt{d-c^2dx^2} \arctan(\sqrt{-1+cx}\sqrt{1+cx})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{ibc^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{ibc^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{ib^2c^2\sqrt{d-c^2dx^2} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} - \frac{ib^2c^2\sqrt{d-c^2dx^2} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

output 
$$\begin{aligned} & -1/2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/x^2-b*c*(a+b*\operatorname{arccosh}(c*x))* \\ & (-c^2*d*x^2+d)^{(1/2)}/x/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+c^2*(a+b*\operatorname{arccosh}(c*x))^2 \\ & *2*\arctan(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)} \\ & /((c*x+1)^{(1/2)}+b^2*c^2*\arctan((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(-c^2*d*x^2+d)^{(1/2)} \\ & /((c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-I*b*c^2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,- \\ & I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c \\ & *x+1)^{(1/2)}+I*b*c^2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,I*(c*x+(c*x-1)^{(1/2)}*(c*x \\ & +1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+I*b^2*c^2*\operatorname{pol} \\ & \operatorname{ylog}(3,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)} \\ & /((c*x+1)^{(1/2)}-I*b^2*c^2*\operatorname{polylog}(3,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})) \\ & )*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \end{aligned}$$

### 3.176.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 5035 vs.  $2(427) = 854$ .

Time = 66.46 (sec) , antiderivative size = 5035, normalized size of antiderivative = 11.79

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx = \text{Result too large to show}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x^3,x]`

output Result too large to show

### 3.176.3 Rubi [A] (verified)

Time = 1.86 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.59, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$ , Rules used = {6339, 6298, 103, 218, 6362, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx$$

↓ 6339

---

3.176.  $\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx$

$$\begin{aligned}
& \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{bc\sqrt{d-c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x^2} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{6298} \\
& \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{bc\sqrt{d-c^2dx^2} \left( bc \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{103} \\
& \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{bc\sqrt{d-c^2dx^2} \left( bc^2 \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{218} \\
& \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{bc\sqrt{d-c^2dx^2} \left( bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{6362} \\
& \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} \operatorname{darccosh}(cx)}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{bc\sqrt{d-c^2dx^2} \left( bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{3042} \\
& \frac{c^2\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx))^2 \csc\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{bc\sqrt{d-c^2dx^2} \left( bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{4668}
\end{aligned}$$

---

3.176.  $\int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^3} dx$

$$\frac{c^2\sqrt{d-c^2dx^2}(-2ib \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2x^2}$$

↓ 3011

$$\frac{c^2\sqrt{d-c^2dx^2}(2ib(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))) - bc \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + bc \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2x^2}$$

↓ 2720

$$\frac{c^2\sqrt{d-c^2dx^2}(2ib(b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))) - bc \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + bc \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2x^2}$$

↓ 7143

$$\frac{c^2\sqrt{d-c^2dx^2}(2 \arctan(e^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))^2 + 2ib(b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))) - bc \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + bc \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2x^2}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x^3,x]`

output `-1/2*(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x^2 + (b*c*Sqrt[d - c^2*d*x^2]*(-(a + b*ArcCosh[c*x])/x) + b*c*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (c^2*Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-(a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-(a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]]))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.176.3.1 Defintions of rubi rules used

rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6339 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`

rule 6362 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.176.4 Maple [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2 \sqrt{-c^2 dx^2 + d}}{x^3} dx$$

input `int((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^3,x)`

output `int((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^3,x)`

### 3.176.5 Fracas [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^2}{x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^3,x, algorithm="fricas")`



output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/x^3, x)`

### 3.176.6 Sympy [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2}{x^3} dx$$

input `integrate((a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2)/x**3,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2/x**3, x)`

### 3.176.7 Maxima [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^2}{x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^3,x, algorithm="maxima")`

output `1/2*(c^2*sqrt(d)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - sqrt(-c^2*d*x^2 + d)*c^2 - (-c^2*d*x^2 + d)^(3/2)/(d*x^2))*a^2 + integrate(sqrt(-c^2*d*x^2 + d)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x^3 + 2*sqrt(-c^2*d*x^2 + d)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^3, x)`

### 3.176.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^3,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.176.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx = \int \frac{(a+b\operatorname{acosh}(cx))^2\sqrt{d-c^2dx^2}}{x^3} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x^3,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x^3, x)`

**3.177**  $\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx$

3.177.1 Optimal result . . . . . 1466  
 3.177.2 Mathematica [A] (warning: unable to verify) . . . . . 1467  
 3.177.3 Rubi [C] (warning: unable to verify) . . . . . 1467  
 3.177.4 Maple [B] (verified) . . . . . 1472  
 3.177.5 Fricas [F] . . . . . 1473  
 3.177.6 Sympy [F] . . . . . 1474  
 3.177.7 Maxima [F] . . . . . 1474  
 3.177.8 Giac [F(-2)] . . . . . 1474  
 3.177.9 Mupad [F(-1)] . . . . . 1475

**3.177.1 Optimal result**

Integrand size = 29, antiderivative size = 336

$$\begin{aligned} & \int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx \\ &= \frac{b^2c^2\sqrt{d-c^2dx^2}}{3x} - \frac{b^2c^3\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{3\sqrt{-1+cx}\sqrt{1+cx}} \\ & \quad - \frac{bc(1-c^2x^2)\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3x^2\sqrt{-1+cx}\sqrt{1+cx}} \\ & \quad - \frac{c^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3\sqrt{-1+cx}\sqrt{1+cx}} - \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{3dx^3} \\ & \quad - \frac{2bc^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\log(1+e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{-1+cx}\sqrt{1+cx}} \\ & \quad + \frac{b^2c^3\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output  $-1/3*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^2/d/x^3+1/3*b^2*c^2*(-c^2*d*x^2+d)^{(1/2)}/x-1/3*b^2*c^3*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/3*b*c*(-c^2*x^2+1)*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/x^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/3*c^3*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2/3*b*c^3*(a+b*\operatorname{arccosh}(c*x))*\ln(1+1/(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/3*b^2*c^3*\operatorname{polylog}(2,-1/(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

**3.177.2 Mathematica [A] (warning: unable to verify)**

Time = 0.90 (sec) , antiderivative size = 304, normalized size of antiderivative = 0.90

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx =$$


---


$$\frac{d(1 + cx) \left( a^2 - a^2 cx - a^2 c^2 x^2 - b^2 c^2 x^2 + a^2 c^3 x^3 + b^2 c^3 x^3 - abcx \sqrt{\frac{-1+cx}{1+cx}} - b^2 \left( -1 + cx + c^2 x^2 + c^3 x^3 \right) \right)}{x^3 \sqrt{d - c^2 dx^2}}$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x^4,x]`output

```

-1/3*(d*(1 + c*x)*(a^2 - a^2*c*x - a^2*c^2*x^2 - b^2*c^2*x^2 + a^2*c^3*x^3
+ b^2*c^3*x^3 - a*b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)] - b^2*(-1 + c*x + c^2*
x^2 + c^3*x^3*(-1 + Sqrt[(-1 + c*x)/(1 + c*x)]))*ArcCosh[c*x]^2 - b*ArcCos
h[c*x]*(b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)] - 2*a*(-1 + c*x)^2*(1 + c*x) + 2*
b*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*Log[1 + E^(-2*ArcCosh[c*x])]) - 2*a*b
*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*Log[c*x] + b^2*c^3*x^3*Sqrt[(-1 + c*x)
/(1 + c*x)]*PolyLog[2, -E^(-2*ArcCosh[c*x])]))/(x^3*Sqrt[d - c^2*d*x^2])

```

**3.177.3 Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 1.18 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.63, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.517$ , Rules used = {6332, 25, 6327, 6335, 108, 27, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$$

↓ 6332

$$\frac{2bc\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3}$$

↓ 25

$$\frac{2bc\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x^3} dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3}$$

---

3.177.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$

$$\begin{aligned}
& \downarrow 6327 \\
& \frac{2bc\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3dx^3} \\
& \downarrow 6335 \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( c^2 \left( -\int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{1}{2}bc \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2} dx - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3dx^3} \\
& \downarrow 108 \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( c^2 \left( -\int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{1}{2}bc \left( \int \frac{c^2}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3dx^3} \\
& \downarrow 27 \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( c^2 \left( -\int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{1}{2}bc \left( c^2 \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3dx^3} \\
& \downarrow 43 \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( c^2 \left( -\int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3dx^3} \\
& \downarrow 6297 \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( -\frac{c^2 \int -\left( (a+\operatorname{barccosh}(cx)) \tanh\left( \frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b} \right) \right) d(a+\operatorname{barccosh}(cx))}{b} - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3dx^3}
\end{aligned}$$

---

3.177.  $\int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^4} dx$

$$\downarrow \text{25}$$

$$2bc\sqrt{d - c^2 dx^2} \left( \frac{c^2 \int (a + \operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{b} - \frac{(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}\left(\frac{d - c^2 dx^2}{3\sqrt{cx - 1}\sqrt{cx + 1}}\right) \right) \right)$$

$$\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3}$$

$$\downarrow \text{3042}$$

$$2bc\sqrt{d - c^2 dx^2} \left( -\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3} + \frac{c^2 \int -i(a + \operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b}\right) d(a + \operatorname{barccosh}(cx))}{b} - \frac{(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}\left(\frac{d - c^2 dx^2}{3\sqrt{cx - 1}\sqrt{cx + 1}}\right) \right) \right)$$

$$\downarrow \text{26}$$

$$2bc\sqrt{d - c^2 dx^2} \left( -\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3} + \frac{ic^2 \int (a + \operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b}\right) d(a + \operatorname{barccosh}(cx))}{b} - \frac{(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}\left(\frac{d - c^2 dx^2}{3\sqrt{cx - 1}\sqrt{cx + 1}}\right) \right) \right)$$

$$\downarrow \text{4201}$$

$$2bc\sqrt{d - c^2 dx^2} \left( -\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3} + \frac{ic^2 \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)} (a + \operatorname{barccosh}(cx)) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2}{1 + e^{-2\operatorname{arccosh}(cx)}}}{b} \right) - \frac{(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}\left(\frac{d - c^2 dx^2}{3\sqrt{cx - 1}\sqrt{cx + 1}}\right) \right) \right)$$

$$\downarrow \text{2620}$$

$$2bc\sqrt{d - c^2 dx^2} \left( -\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3} + \frac{ic^2 \left( 2i \left( \frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2 \right)}{b} - \frac{(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}\left(\frac{d - c^2 dx^2}{3\sqrt{cx - 1}\sqrt{cx + 1}}\right) \right) \right)$$

$$\downarrow \text{2715}$$

---

3.177.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$

$$\begin{aligned}
 & -\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3} + \\
 & 2bc\sqrt{d - c^2 dx^2} \left( -\frac{ic^2 \left( 2i \left( -\frac{1}{4} b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) dx - 2\operatorname{arccosh}(cx) - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) \right)}{b} \right) \\
 & \hspace{20em} 3\sqrt{cx - 1}\sqrt{cx + 1} \\
 & \hspace{10em} \downarrow \text{2838} \\
 & -\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3dx^3} + \\
 & 2bc\sqrt{d - c^2 dx^2} \left( -\frac{ic^2 \left( 2i \left( \frac{1}{4} b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) - \frac{1}{2} i (a + \operatorname{barccosh}(cx))^2 \right) \right)}{b} \right) \\
 & \hspace{20em} 3\sqrt{cx - 1}\sqrt{cx + 1}
 \end{aligned}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x^4,x]`

output `-1/3*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/(d*x^3) + (2*b*c*Sqrt[d - c^2*d*x^2]*(-1/2*((1 - c^2*x^2)*(a + b*ArcCosh[c*x]))/x^2 - (b*c*(-((Sqrt[-1 + c*x]*Sqrt[1 + c*x])/x) + c*ArcCosh[c*x]))/2 - (I*c^2*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x]))] + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4))/b))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

3.177.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

---

3.177.  $\int \frac{\sqrt{d-c^2 dx^2}(a+\operatorname{barccosh}(cx))^2}{x^4} dx$

rule 108 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(m + 1))), x] - Simp[1/(b*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

rule 2620 `Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`



rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 6335 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (-Simp[b*c*((-d)^p/(f*(m + 1))) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x] - Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0] && ILtQ[(m + 1)/2, 0]`

### 3.177.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1750 vs.  $2(314) = 628$ .

Time = 1.22 (sec) , antiderivative size = 1751, normalized size of antiderivative = 5.21

method	result	size
default	Expression too large to display	1751
parts	Expression too large to display	1751

input `int((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^4,x,method=_RETURNVERBOSE)`

output  $\frac{1}{3}b^2(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)*x*\operatorname{arccosh}(cx)*c^4-$   
 $\frac{1}{3}b^2(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)/(cx+1)^{1/2}/(cx-$   
 $1)^{1/2}*c^3-1/3*b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)*x^3*\operatorname{ar}$   
 $\operatorname{ccosh}(cx)*c^6+1/3*b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)*x^3*$   
 $c^6-5/3*b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)*x^3/(cx+1)/(c*$   
 $x-1)*c^6+4/3*b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)*x/(cx+1)/$   
 $(cx-1)*c^4-1/3*b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)/x/(cx+$   
 $1)/(cx-1)*c^2+1/3*b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)/x^3/$   
 $(cx+1)/(cx-1)*\operatorname{arccosh}(cx)^2-2/3*b^2*(-d(c^2x^2-1))^{1/2}/(cx-1)^{1/2}$   
 $)/(cx+1)^{1/2}*\operatorname{arccosh}(cx)*\ln(1+(cx+(cx-1)^{1/2})*(cx+1)^{1/2})^2)*c^3$   
 $+2/3*b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)*x^5/(cx+1)/(cx-1)$   
 $)*c^8-b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)*x^4/(cx+1)^{1/2}$   
 $/(cx-1)^{1/2}*c^7+b^2*(-d(c^2x^2-1))^{1/2}/(3c^4x^4-3c^2x^2+1)/(cx$   
 $+1)^{1/2}/(cx-1)^{1/2}*\operatorname{arccosh}(cx)*c^3+b^2*(-d(c^2x^2-1))^{1/2}/(3c^4$   
 $*x^4-3c^2x^2+1)*x^2/(cx+1)^{1/2}/(cx-1)^{1/2}*c^5-1/3*b^2*(-d(c^2x^2$   
 $-1))^{1/2}/(3c^4x^4-3c^2x^2+1)/(cx+1)^{1/2}/(cx-1)^{1/2}*\operatorname{arccosh}(cx$   
 $)^2*c^3+1/3*a*b*(-d(c^2x^2-1))^{1/2}*(2*(cx+1)^{1/2}*\operatorname{arccosh}(cx)*(cx-$   
 $1)^{1/2}*c^2x^2+2*c^3*x^3*\operatorname{arccosh}(cx)-2*\ln(1+(cx+(cx-1)^{1/2})*(cx+1)^$   
 $(1/2))^2)*x^3*c^3-2*\operatorname{arccosh}(cx)*(cx-1)^{1/2}*(cx+1)^{1/2}-cx)/(cx-1)^$   
 $(1/2)/(cx+1)^{1/2}/x^3-1/3*a^2/d/x^3*(-c^2*d*x^2+d)^{3/2}-3*b^2*(-d(c...$

### 3.177.5 Fricas [F]

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx = \int \frac{\sqrt{-c^2dx^2+d}(b\operatorname{arccosh}(cx)+a)^2}{x^4} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^4,x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/x^4, x)`

## 3.177.6 Sympy [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2}{x^4} dx$$

input `integrate((a+b*acosh(c*x))**2*(-c**2*d*x**2+d)**(1/2)/x**4,x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2/x**4, x)`

## 3.177.7 Maxima [F]

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^2}{x^4} dx$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^4,x, algorithm="maxima")`

output `1/3*(c^4*d^2*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2) + I*(-1)^(-2*c^2*d*x^2 + 2*d)*c^2*d^(3/2)*log(-2*c^2*d + 2*d/x^2) + sqrt(-c^4*d*x^4 + 2*c^2*d*x^2 - d)*d/x^2)*a*b*c/d + 1/3*b^2*((c^2*sqrt(d)*x^2 - sqrt(d))*sqrt(c*x + 1)*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x^3 - 3*integrate(2/3*((c*x + 1)*sqrt(c*x - 1)*c^2*sqrt(d)*x + (c^3*sqrt(d)*x^2 - c*sqrt(d))*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(c*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*x^3), x) - 2/3*(-c^2*d*x^2 + d)^(3/2)*a*b*arccosh(c*x)/(d*x^3) - 1/3*(-c^2*d*x^2 + d)^(3/2)*a^2/(d*x^3)`

## 3.177.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/x^4,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.177.9 Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{x^4} dx = \int \frac{(a+b\operatorname{acosh}(cx))^2\sqrt{d-c^2dx^2}}{x^4} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x^4,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2))/x^4, x)`

### 3.178 $\int x^3(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.178.1 Optimal result . . . . .	1476
3.178.2 Mathematica [A] (verified) . . . . .	1477
3.178.3 Rubi [A] (verified) . . . . .	1478
3.178.4 Maple [B] (verified) . . . . .	1488
3.178.5 Fricas [A] (verification not implemented) . . . . .	1489
3.178.6 Sympy [F(-1)] . . . . .	1489
3.178.7 Maxima [A] (verification not implemented) . . . . .	1490
3.178.8 Giac [F(-2)] . . . . .	1491
3.178.9 Mupad [F(-1)] . . . . .	1491

#### 3.178.1 Optimal result

Integrand size = 29, antiderivative size = 495

$$\begin{aligned} \int x^3(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = & -\frac{37384b^2d\sqrt{d - c^2dx^2}}{385875c^4} \\ & + \frac{3358b^2dx^2\sqrt{d - c^2dx^2}}{385875c^2} + \frac{484b^2dx^4\sqrt{d - c^2dx^2}}{42875} - \frac{2}{343}b^2c^2dx^6\sqrt{d - c^2dx^2} \\ & + \frac{4abdx\sqrt{d - c^2dx^2}}{35c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{4b^2dx\sqrt{d - c^2dx^2}\operatorname{arccosh}(cx)}{35c^3\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & + \frac{2bdx^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{105c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{16bcdx^5\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{175\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & + \frac{2bc^3dx^7\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{49\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{2d\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{35c^4} \\ & - \frac{dx^2\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{35c^2} + \frac{3}{35}dx^4\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 \\ & + \frac{1}{7}x^4(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \end{aligned}$$

output  $\frac{1}{7}x^4(-c^2dx^2+d)^{3/2}(a+b\operatorname{arccosh}(cx))^2-37384/385875b^2d(-c^2dx^2+d)^{1/2}/c^4+3358/385875b^2d^2(-c^2dx^2+d)^{1/2}/c^2+484/42875b^2d^2(-c^2dx^2+d)^{1/2}-2/343b^2c^2d^2x^6(-c^2dx^2+d)^{1/2}-2/35d(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{1/2}/c^4-1/35d^2x^2(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{1/2}/c^2+3/35d^2x^4(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{1/2}+4/35abdx^2(-c^2dx^2+d)^{1/2}/c^3/(cx-1)^{1/2}/(cx+1)^{1/2}+4/35b^2dx\operatorname{arccosh}(cx)(-c^2dx^2+d)^{1/2}/c^3/(cx-1)^{1/2}/(cx+1)^{1/2}+2/105b^2d^2x^3(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}/c/(cx-1)^{1/2}/(cx+1)^{1/2}-16/175b^2cd^2x^5(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}+2/49b^2c^3d^2x^7(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}$

### 3.178.2 Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 262, normalized size of antiderivative = 0.53

$$\int x^3(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 dx = \frac{d\sqrt{d - c^2dx^2} \left( 11025a^2(-1 + c^2x^2)^3(2 + 5c^2x^2) - 210abcx\sqrt{-1 + cx}\sqrt{1 + cx}(210 + 35c^2x^2 - 168c^4x^4 + 75c^6x^6) + 2b^2(-18692 + 20371c^2x^2 + 499c^4x^4 - 3303c^6x^6 + 1125c^8x^8) - 210b(-105a(-1 + c^2x^2)^3(2 + 5c^2x^2) + b^2cx\sqrt{-1 + cx}\sqrt{1 + cx}(210 + 35c^2x^2 - 168c^4x^4 + 75c^6x^6)) \operatorname{ArcCosh}[cx] + 11025b^2(-1 + c^2x^2)^3(2 + 5c^2x^2)\operatorname{ArcCosh}[cx]^2 \right)}{c^4(-1 + c^2x^2)}$$

input `Integrate[x^3*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output  $-1/385875*(d\sqrt{d - c^2dx^2}*(11025a^2*(-1 + c^2x^2)^3*(2 + 5c^2x^2) - 210a^2bcx\sqrt{-1 + cx}\sqrt{1 + cx}*(210 + 35c^2x^2 - 168c^4x^4 + 75c^6x^6) + 2b^2*(-18692 + 20371c^2x^2 + 499c^4x^4 - 3303c^6x^6 + 1125c^8x^8) - 210b(-105a(-1 + c^2x^2)^3*(2 + 5c^2x^2) + b^2cx\sqrt{-1 + cx}\sqrt{1 + cx}*(210 + 35c^2x^2 - 168c^4x^4 + 75c^6x^6))\operatorname{ArcCosh}[cx] + 11025b^2(-1 + c^2x^2)^3*(2 + 5c^2x^2)\operatorname{ArcCosh}[cx]^2)/(c^4*(-1 + c^2x^2))$

**3.178.3 Rubi [A] (verified)**

Time = 3.76 (sec) , antiderivative size = 663, normalized size of antiderivative = 1.34, number of steps used = 25, number of rules used = 25,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.862$ , Rules used = {6345, 25, 6327, 6336, 27, 960, 111, 27, 111, 27, 83, 6341, 6298, 111, 27, 111, 27, 83, 6354, 6298, 111, 27, 83, 6330, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$\downarrow \text{6345}$$

$$\frac{2bcd\sqrt{d - c^2 dx^2} \int -x^4(1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))dx}{7\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{3}{7}d \int x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

$$\downarrow \text{25}$$

$$-\frac{2bcd\sqrt{d - c^2 dx^2} \int x^4(1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))dx}{7\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{3}{7}d \int x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

$$\downarrow \text{6327}$$

$$-\frac{2bcd\sqrt{d - c^2 dx^2} \int x^4(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))dx}{7\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{3}{7}d \int x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

$$\downarrow \text{6336}$$

$$\frac{\frac{3}{7}d \int x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx - 2bcd\sqrt{d - c^2 dx^2} \left( -bc \int \frac{x^5(7-5c^2 x^2)}{35\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^2 x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) \right)}{7\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

$$\downarrow \text{27}$$

$$\frac{\frac{3}{7}d \int x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2 dx - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \int \frac{x^5(7-5c^2 x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^2 x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) \right)}{7\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

---

3.178.  $\int x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned} & \downarrow 960 \\ & \frac{\frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \int \frac{x^5}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{5}{7}x^6 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \\ & \frac{\frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}} \end{aligned}$$

$$\begin{aligned} & \downarrow 111 \\ & \frac{\frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \frac{\int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \\ & \frac{\frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}} \end{aligned}$$

$$\begin{aligned} & \downarrow 27 \\ & \frac{\frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \\ & \frac{\frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}} \end{aligned}$$

$$\begin{aligned} & \downarrow 111 \\ & \frac{\frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \frac{4 \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \\ & \frac{\frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}} \end{aligned}$$

$$\downarrow 27$$



$$\frac{\frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) - \frac{5}{7}x^6 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{7}c \right)}{7\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 83

$$\frac{\frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{3c^4} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6341

$$\frac{\frac{3}{7}d \left( -\frac{2bc\sqrt{d - c^2 dx^2} \int x^4 (a + \operatorname{barccosh}(cx)) dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2} \int \frac{x^3 (a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \right) - \frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{3c^4} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6298

$$\frac{\frac{3}{7}d \left( -\frac{2bc\sqrt{d - c^2 dx^2} \left( \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{1}{5}bc \int \frac{x^5}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2} \int \frac{x^3 (a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \right) - \frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{3c^4} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}x^4 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{7\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 111

---

3.178.  $\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{\int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + x^4 \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \frac{\frac{1}{7}x^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{5c^2} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}{7\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 27

$$\frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + x^4 \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \frac{\frac{1}{7}x^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{5c^2} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}{7\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 111

$$\frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + x^2 \right)}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \frac{\frac{1}{7}x^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{5c^2} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}{7\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 27

---

3.178.  $\int x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{\frac{3}{7}d \left( \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx + x \right)}{3c^2} \right)}{5c^2} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right)}{\frac{1}{7}x^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{5c^2} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}$$

↓ 83

$$\frac{\frac{3}{7}d \left( \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}{\frac{1}{7}x^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{5c^2} \right) \right) \right)}$$

↓ 6354

$$\frac{\frac{3}{7}d \left( \frac{\sqrt{d-c^2dx^2} \left( \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} - \frac{2b \int x^2(a+\operatorname{barccosh}(cx)) dx}{3c} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^2}{3c^2} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \right)}{\frac{1}{7}x^4(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{5c^2} \right) \right) \right)}$$

↓ 6298

---

3.178.  $\int x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{3}{7}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{3c^2} - \frac{2b \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{3c} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{3c^2} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{\frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + \operatorname{arccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{arccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{3c^2} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right.$$

↓ 111

$$\frac{3}{7}d \left( \frac{\sqrt{d - c^2 dx^2} \left( -\frac{2b \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx + x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{3c^2} \right) \\ \left. - \frac{\frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + \operatorname{arccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{arccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{3c^2} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right.$$

↓ 27

$$\frac{3}{7}d \left( \frac{\sqrt{d - c^2 dx^2} \left( -\frac{2b \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{3c^2} \right) \\ \left. - \frac{\frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + \operatorname{arccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{arccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{3c^2} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right.$$

↓ 83

---

3.178.  $\int x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^2 dx$

$$\frac{3}{7}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{2 \int \frac{x(a + b \operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{3c^2} - \frac{2b \left( \frac{1}{3} x^3 (a + b \operatorname{arccosh}(cx)) - \frac{1}{3} bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right)}{\frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + b \operatorname{arccosh}(cx)) + \frac{1}{5}x^5 (a + b \operatorname{arccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{\right)} \right.$$

↓ 6330

$$\frac{3}{7}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{c^2} - \frac{2b \int (a + b \operatorname{arccosh}(cx)) dx}{c} \right)}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{3c^2} - \frac{2b \left( \frac{1}{3} x^3 (a + b \operatorname{arccosh}(cx)) - \frac{1}{3} bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right)}{\frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 - 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + b \operatorname{arccosh}(cx)) + \frac{1}{5}x^5 (a + b \operatorname{arccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right) \right) \right)}{\right)} \right.$$

↓ 2009

$$\frac{3}{7}d \left( \frac{\frac{1}{5}x^4 \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 - \sqrt{d - c^2 dx^2} \left( \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{3c^2} + \frac{2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{c^2} \right)}{\right)}{\frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 + 2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2 x^7 (a + b \operatorname{arccosh}(cx)) + \frac{1}{5}x^5 (a + b \operatorname{arccosh}(cx)) - \frac{1}{35}bc \left( \frac{19}{7} \left( \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right) \right) \right)}{\right)} \right.$$

input `Int[x^3*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output `(x^4*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/7 - (2*b*c*d*Sqrt[d - c^2*d*x^2]*(-1/35*(b*c*((-5*x^6*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/7 + (19*((x^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(5*c^2) + (4*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2)))/(5*c^2)))/7)) + (x^5*(a + b*ArcCosh[c*x]))/5 - (c^2*x^7*(a + b*ArcCosh[c*x]))/7)/(7*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (3*d*((x^4*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/5 - (2*b*c*Sqrt[d - c^2*d*x^2]*(-1/5*(b*c*((x^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(5*c^2) + (4*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2)))/(5*c^2)) + (x^5*(a + b*ArcCosh[c*x]))/5))/(5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*((x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(3*c^2) - (2*b*(-1/3*(b*c*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2)) + (x^3*(a + b*ArcCosh[c*x]))/3))/3*c) + (2*((Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/c^2 - (2*b*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/c))/(3*c^2)))/(5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/7`

### 3.178.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 111 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`

rule 960 `Int[((e_.)*(x_))^(m_.)*((a1_.) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_.) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_)^p)*((d2_) + (e2_.)*(x_)^p), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`



### 3.178.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1951 vs.  $2(423) = 846$ .

Time = 0.83 (sec) , antiderivative size = 1952, normalized size of antiderivative = 3.94

method	result	size
default	Expression too large to display	1952
parts	Expression too large to display	1952

input `int(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output

```

a^2*(-1/7*x^2*(-c^2*d*x^2+d)^(5/2)/c^2/d-2/35/d/c^4*(-c^2*d*x^2+d)^(5/2))+
b^2*(-1/43904*(-d*(c^2*x^2-1))^(1/2)*(64*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1
/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*x^4-112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*
x^5-25*c^2*x^2+56*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x
+1)^(1/2)*c*x+1)*(49*arccosh(c*x)^2-14*arccosh(c*x)+2)*d/(c*x+1)/c^4/(c*x-
1)+1/16000*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*
(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*
(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(25*arccosh(c*x)^2-10*arccosh(c*x)+2)*d
/(c*x+1)/c^4/(c*x-1)+1/1152*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*
(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(
9*arccosh(c*x)^2-6*arccosh(c*x)+2)*d/(c*x+1)/c^4/(c*x-1)-3/128*(-d*(c^2*x^
2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*
arccosh(c*x)+2)*d/(c*x+1)/c^4/(c*x-1)-3/128*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-
1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2+2*arccosh(c*x)+2)*d/
(c*x+1)/c^4/(c*x-1)+1/1152*(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1
)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(
9*arccosh(c*x)^2+6*arccosh(c*x)+2)*d/(c*x+1)/c^4/(c*x-1)+1/16000*(-d*(c^2*
x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-
1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-28*c^4*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*
x+13*c^2*x^2-1)*(25*arccosh(c*x)^2+10*arccosh(c*x)+2)*d/(c*x+1)/c^4/(c*...

```

**3.178.5 Fracas [A] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 432, normalized size of antiderivative = 0.87

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{11025 (5 b^2 c^8 dx^8 - 13 b^2 c^6 dx^6 + 9 b^2 c^4 dx^4 + b^2 c^2 dx^2 - 2 b^2 d) \sqrt{-c^2 dx^2 + d} \log (cx + \sqrt{c^2 x^2 - 1})^2 - 210 (7$$

```
input integrate(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")
```

```
output -1/385875*(11025*(5*b^2*c^8*d*x^8 - 13*b^2*c^6*d*x^6 + 9*b^2*c^4*d*x^4 + b^2*c^2*d*x^2 - 2*b^2*d)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1))^2 - 210*(75*a*b*c^7*d*x^7 - 168*a*b*c^5*d*x^5 + 35*a*b*c^3*d*x^3 + 210*a*b*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 210*((75*b^2*c^7*d*x^7 - 168*b^2*c^5*d*x^5 + 35*b^2*c^3*d*x^3 + 210*b^2*c*d*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 105*(5*a*b*c^8*d*x^8 - 13*a*b*c^6*d*x^6 + 9*a*b*c^4*d*x^4 + a*b*c^2*d*x^2 - 2*a*b*d)*sqrt(-c^2*d*x^2 + d))*log(c*x + sqrt(c^2*x^2 - 1)) + (1125*(49*a^2 + 2*b^2)*c^8*d*x^8 - 9*(15925*a^2 + 734*b^2)*c^6*d*x^6 + (99225*a^2 + 998*b^2)*c^4*d*x^4 + (11025*a^2 + 40742*b^2)*c^2*d*x^2 - 2*(11025*a^2 + 18692*b^2)*d)*sqrt(-c^2*d*x^2 + d))/(c^6*x^2 - c^4)
```

**3.178.6 Sympy [F(-1)]**

Timed out.

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Timed out}$$

```
input integrate(x**3*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2,x)
```

```
output Timed out
```

**3.178.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 388, normalized size of antiderivative = 0.78

$$\begin{aligned}
& \int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^2 dx = \\
& -\frac{1}{35} \left( \frac{5(-c^2 dx^2 + d)^{5/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{5/2}}{c^4 d} \right) b^2 \operatorname{arccosh}(cx)^2 \\
& -\frac{2}{35} \left( \frac{5(-c^2 dx^2 + d)^{5/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{5/2}}{c^4 d} \right) ab \operatorname{arccosh}(cx) \\
& -\frac{1}{35} \left( \frac{5(-c^2 dx^2 + d)^{5/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{5/2}}{c^4 d} \right) a^2 \\
& -\frac{2}{385875} b^2 \left( \frac{1125 \sqrt{c^2 x^2 - 1} c^4 \sqrt{-d} dx^6 - 2178 \sqrt{c^2 x^2 - 1} c^2 \sqrt{-d} dx^4 - 1679 \sqrt{c^2 x^2 - 1} \sqrt{-d} dx^2 + \frac{18692 \sqrt{c^2 x^2 - 1}}{c}}{c^2} \right) \\
& + \frac{2(75 c^6 \sqrt{-d} dx^7 - 168 c^4 \sqrt{-d} dx^5 + 35 c^2 \sqrt{-d} dx^3 + 210 \sqrt{-d} dx) ab}{3675 c^3}
\end{aligned}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")
```

```
output -1/35*(5*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(5/2)/(c^4*d))*b^2*arccosh(c*x)^2 - 2/35*(5*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(5/2)/(c^4*d))*a*b*arccosh(c*x) - 1/35*(5*(-c^2*d*x^2 + d)^(5/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(5/2)/(c^4*d))*a^2 - 2/385875*b^2*((1125*sqrt(c^2*x^2 - 1)*c^4*sqrt(-d)*d*x^6 - 2178*sqrt(c^2*x^2 - 1)*c^2*sqrt(-d)*d*x^4 - 1679*sqrt(c^2*x^2 - 1)*sqrt(-d)*d*x^2 + 18692*sqrt(c^2*x^2 - 1)*sqrt(-d)*d/c^2)/c^2 - 105*(75*c^6*sqrt(-d)*d*x^7 - 168*c^4*sqrt(-d)*d*x^5 + 35*c^2*sqrt(-d)*d*x^3 + 210*sqrt(-d)*d*x)*arccosh(c*x)/c^3 + 2/3675*(75*c^6*sqrt(-d)*d*x^7 - 168*c^4*sqrt(-d)*d*x^5 + 35*c^2*sqrt(-d)*d*x^3 + 210*sqrt(-d)*d*x)*a*b/c^3
```

**3.178.8 Giac [F(-2)]**

Exception generated.

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.178.9 Mupad [F(-1)]**

Timed out.

$$\int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int x^3 (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2} dx$$

input `int(x^3*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2),x)`

output `int(x^3*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2), x)`

### 3.179 $\int x^2(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.179.1 Optimal result	1492
3.179.2 Mathematica [A] (warning: unable to verify)	1493
3.179.3 Rubi [A] (verified)	1493
3.179.4 Maple [B] (verified)	1502
3.179.5 Fricas [F]	1503
3.179.6 Sympy [F(-1)]	1504
3.179.7 Maxima [F]	1504
3.179.8 Giac [F]	1504
3.179.9 Mupad [F(-1)]	1505

#### 3.179.1 Optimal result

Integrand size = 29, antiderivative size = 441

$$\begin{aligned} \int x^2(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx &= \frac{7b^2dx\sqrt{d - c^2dx^2}}{1152c^2} \\ &+ \frac{43b^2dx^3\sqrt{d - c^2dx^2}}{1728} - \frac{1}{108}b^2c^2dx^5\sqrt{d - c^2dx^2} \\ &+ \frac{7b^2d\sqrt{d - c^2dx^2}\operatorname{arccosh}(cx)}{1152c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bdx^2\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{16c\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &- \frac{7bcdx^4\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{48\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc^3dx^6\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))}{18\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &- \frac{dx\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{16c^2} + \frac{1}{8}dx^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 \\ &+ \frac{1}{6}x^3(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{d\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^3}{48bc^3\sqrt{-1 + cx}\sqrt{1 + cx}} \end{aligned}$$

output  $\frac{1}{6}x^3(-c^2dx^2+d)^{(3/2)}(a+b\operatorname{arccosh}(cx))^2+\frac{7}{1152}b^2d^2x^3(-c^2dx^2+d)^{(1/2)}/c^2+\frac{43}{1728}b^2d^2x^3(-c^2dx^2+d)^{(1/2)}-\frac{1}{108}b^2c^2d^2x^5(-c^2dx^2+d)^{(1/2)}-\frac{1}{16}d^2x^3(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{(1/2)}/c^2+\frac{1}{8}d^2x^3(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{(1/2)}+\frac{7}{1152}b^2d^2\operatorname{arccosh}(cx)(-c^2dx^2+d)^{(1/2)}/c^3/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}+\frac{1}{16}b^2d^2x^2(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{(1/2)}/c/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-\frac{7}{48}b^2c^3d^2x^4(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{(1/2)}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}+\frac{1}{18}b^2c^3d^2x^6(a+b\operatorname{arccosh}(cx))(-c^2dx^2+d)^{(1/2)}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-\frac{1}{48}d^2(a+b\operatorname{arccosh}(cx))^3(-c^2dx^2+d)^{(1/2)}/b/c^3/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}$

**3.179.2 Mathematica [A] (warning: unable to verify)**

Time = 4.83 (sec) , antiderivative size = 485, normalized size of antiderivative = 1.10

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{-288a^2 c dx \sqrt{\frac{-1+cx}{1+cx}}(1+cx) \sqrt{d - c^2 dx^2} (3 - 14c^2 x^2 + 8c^4 x^4) - 864a^2 d^{3/2} \sqrt{\frac{-1+cx}{1+cx}}(1 + \dots}{\dots}$$

input `Integrate[x^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output

```
(-288*a^2*c*d*x*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*sqrt[d - c^2*d*x^2]*(3 - 14*c^2*x^2 + 8*c^4*x^4) - 864*a^2*d^(3/2)*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcTan[(c*x*sqrt[d - c^2*d*x^2])/(sqrt[d]*(-1 + c^2*x^2))] - 216*a*b*d*sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*sinh[4*ArcCosh[c*x]]) - 18*b^2*d*sqrt[d - c^2*d*x^2]*(32*ArcCosh[c*x]^3 + 12*ArcCosh[c*x]*Cosh[4*ArcCosh[c*x]] - 3*(1 + 8*ArcCosh[c*x]^2)*sinh[4*ArcCosh[c*x]]) - 12*a*b*d*sqrt[d - c^2*d*x^2]*(-72*ArcCosh[c*x]^2 + 18*Cosh[2*ArcCosh[c*x]] - 9*Cosh[4*ArcCosh[c*x]] - 2*Cosh[6*ArcCosh[c*x]] + 12*ArcCosh[c*x]*(-3*sinh[2*ArcCosh[c*x]] + 3*sinh[4*ArcCosh[c*x]] + sinh[6*ArcCosh[c*x]])) + b^2*d*sqrt[d - c^2*d*x^2]*(288*ArcCosh[c*x]^3 + 12*ArcCosh[c*x]*(-18*Cosh[2*ArcCosh[c*x]] + 9*Cosh[4*ArcCosh[c*x]] + 2*Cosh[6*ArcCosh[c*x]]) + 108*sinh[2*ArcCosh[c*x]] - 27*sinh[4*ArcCosh[c*x]] - 4*sinh[6*ArcCosh[c*x]] - 72*ArcCosh[c*x]^2*(-3*sinh[2*ArcCosh[c*x]] + 3*sinh[4*ArcCosh[c*x]] + sinh[6*ArcCosh[c*x]])))/(13824*c^3*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))
```

**3.179.3 Rubi [A] (verified)**Time = 3.18 (sec) , antiderivative size = 550, normalized size of antiderivative = 1.25, number of steps used = 21, number of rules used = 21,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.724$ , Rules used = {6345, 25, 6327, 6336, 27, 960, 111, 27, 101, 43, 6341, 6298, 111, 27, 101, 43, 6354, 6298, 101, 43, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$$

---

3.179.  $\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
& \downarrow 6345 \\
& \frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + bcd\sqrt{d - c^2 dx^2} \int -x^3(1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))dx}{3\sqrt{cx - 1}\sqrt{cx + 1} \operatorname{barccosh}(cx)^2} + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \\
& \downarrow 25 \\
& \frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \int x^3(1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\
& \downarrow 6327 \\
& \frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \int x^3(1 - c^2 x^2)(a + \operatorname{barccosh}(cx))dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\
& \downarrow 6336 \\
& \frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \left( -bc \int \frac{x^4(3 - 2c^2 x^2)}{12\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\
& \downarrow 27 \\
& \frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \int \frac{x^4(3 - 2c^2 x^2)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\
& \downarrow 960 \\
& \frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \int \frac{x^4}{\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{3}x^5 \sqrt{cx - 1}\sqrt{cx + 1} \right) - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \\
& \frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\
& \downarrow 111
\end{aligned}$$

---

3.179.  $\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \left( \frac{\int \frac{3x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 27

$$\frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 101

$$\frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 43

$$\frac{\frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right) + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 6341

$$\frac{\frac{1}{2}d \left( -\frac{\sqrt{d - c^2 dx^2} \int \frac{x^2 (a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bcd\sqrt{d - c^2 dx^2} \int x^3 (a + \operatorname{barccosh}(cx)) dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 - \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right) + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) - \frac{1}{3}x^5 \sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2$$

---

3.179.  $\int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$



↓ 6298

$$\frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \int \frac{x^4}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) + \frac{1}{4}x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{6}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + x \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 111

$$\frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{\int \frac{3x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}} \right) \right) + \frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{6}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + x \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 27

$$\frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{d-c^2dx^2}}{3\sqrt{cx-1}\sqrt{cx+1}} \right) \right) + \frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{6}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + x \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 101

---

3.179.  $\int x^2(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx + x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$\frac{\frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{6}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x\sqrt{cx-1}\sqrt{cx+1} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 43

$$\frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x\sqrt{cx-1}\sqrt{cx+1} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$\frac{\frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{6}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x\sqrt{cx-1}\sqrt{cx+1} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6354

$$\frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \left( \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} - \frac{b \int x(a+\operatorname{barccosh}(cx)) dx}{c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^2}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x\sqrt{cx-1}\sqrt{cx+1} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$\frac{\frac{1}{6}x^3(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 - bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{6}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x\sqrt{cx-1}\sqrt{cx+1} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 6298

---

3.179.  $\int x^2(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{1}{2}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{\int \frac{(a + b \operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2c^2} - \frac{b \left( \frac{1}{2}x^2(a + b \operatorname{arccosh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{c} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{\frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6}c^2 x^6(a + b \operatorname{arccosh}(cx)) + \frac{1}{4}x^4(a + b \operatorname{arccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right.$$

↓ 101

$$\frac{1}{2}d \left( \frac{\sqrt{d - c^2 dx^2} \left( -\frac{b \left( \frac{1}{2}x^2(a + b \operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{\int \frac{(a + b \operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))}{2c^2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{\frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6}c^2 x^6(a + b \operatorname{arccosh}(cx)) + \frac{1}{4}x^4(a + b \operatorname{arccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right.$$

↓ 43

$$\frac{1}{2}d \left( \frac{\sqrt{d - c^2 dx^2} \left( \frac{\int \frac{(a + b \operatorname{arccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a + b \operatorname{arccosh}(cx))^2}{2c^2} - \frac{b \left( \frac{1}{2}x^2(a + b \operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} \right) \right)}{c} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{\frac{1}{6}x^3(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 - bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6}c^2 x^6(a + b \operatorname{arccosh}(cx)) + \frac{1}{4}x^4(a + b \operatorname{arccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} \right) + x \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right.$$

↓ 6308

---

3.179.  $\int x^2(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 dx$

$$\frac{1}{6}x^3(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 +$$

$$\frac{1}{2}d \left( \frac{1}{4}x^3\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{\sqrt{d - c^2dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))^3}{6bc^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))^2}{2c^2} - \frac{b}{2} \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$\frac{bcd\sqrt{d - c^2dx^2} \left( -\frac{1}{6}c^2x^6(a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a + \operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + x \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)}{3\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[x^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output `(x^3*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/6 - (b*c*d*Sqrt[d - c^2*d*x^2]*((x^4*(a + b*ArcCosh[c*x]))/4 - (c^2*x^6*(a + b*ArcCosh[c*x]))/6 - (b*c*(-1/3*(x^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (4*((x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(4*c^2) + (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/(4*c^2)))/3))/12)/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (d*((x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/4 - (b*c*Sqrt[d - c^2*d*x^2]*((x^4*(a + b*ArcCosh[c*x]))/4 - (b*c*((x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(4*c^2) + (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/(4*c^2)))/4))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^2)/(2*c^2) + (a + b*ArcCosh[c*x])^3/(6*b*c^3) - (b*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/2))/4)/(4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/2`

### 3.179.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

$$3.179. \quad \int x^2(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 dx$$

rule 101 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)*(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)n(e + f*x)pSimp[a2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 111 `Int[((a_.) + (b_.)*(x_))(m_)((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)(m - 1)(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)(m - 2)(c + d*x)n(e + f*x)pSimp[a2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`

rule 960 `Int[((e_.)*(x_))(m_)((a1_ + (b1_.)*(x_)(non2_))(p_)((a2_ + (b2_.)*(x_)(non2_))(p_)((c_) + (d_.)*(x_)(n_)), x_Symbol] := Simp[d*(e*x)(m + 1)(a1 + b1*x(n/2))(p + 1)((a2 + b2*x(n/2))(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)m(a1 + b1*x(n/2))p(a2 + b2*x(n/2))p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_)((d_.)*(x_))(m_), x_Symbol] := Simp[(d*x)(m + 1)((a + b*ArcCosh[c*x])n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)(m + 1)((a + b*ArcCosh[c*x])(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_)/(Sqrt[(d1_ + (e1_.)*(x_)]*Sqrt[(d2_ + (e2_.)*(x_))], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x])^n * u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x] - Simp[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1)), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.179.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1720 vs.  $2(377) = 754$ .

Time = 0.85 (sec) , antiderivative size = 1721, normalized size of antiderivative = 3.90

method	result	size
default	Expression too large to display	1721
parts	Expression too large to display	1721

```
input int(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

output

```

-1/6*a^2*x*(-c^2*d*x^2+d)^(5/2)/c^2/d+1/24*a^2/c^2*x*(-c^2*d*x^2+d)^(3/2)+
1/16*a^2/c^2*d*x*(-c^2*d*x^2+d)^(1/2)+1/16*a^2/c^2*d^2/(c^2*d)^(1/2)*arcta
n((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b^2*(-1/48*(-d*(c^2*x^2-1))^(1/2)/
(c*x-1)^(1/2)/(c*x+1)^(1/2)/c^3*arccosh(c*x)^3*d-1/6912*(-d*(c^2*x^2-1))^(
1/2)*(32*c^7*x^7-64*c^5*x^5+32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6+38*c^3*
x^3-48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-6*c*x+18*(c*x-1)^(1/2)*(c*x+1)^(
1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(18*arccosh(c*x)^2-6*arccosh(c*
x)+1)*d/(c*x+1)/c^3/(c*x-1)+1/1024*(-d*(c^2*x^2-1))^(1/2)*(8*c^5*x^5-12*c^
3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c*x-1)^(1/2)*(c*x+1)^(
1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(8*arccosh(c*x)^2-4*arccosh(c*x
)+1)*d/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(2*arcco
sh(c*x)^2-2*arccosh(c*x)+1)*d/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c^2*x^2-1))^(
1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+
1)^(1/2)-2*c*x)*(2*arccosh(c*x)^2+2*arccosh(c*x)+1)*d/(c*x+1)/c^3/(c*x-1)+
1/1024*(-d*(c^2*x^2-1))^(1/2)*(-8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c^
5*x^5+8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-12*c^3*x^3-(c*x-1)^(1/2)*(c*x+
1)^(1/2)+4*c*x)*(8*arccosh(c*x)^2+4*arccosh(c*x)+1)*d/(c*x+1)/c^3/(c*x-1)-
1/6912*(-d*(c^2*x^2-1))^(1/2)*(-32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6+32*
c^7*x^7+48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-64*c^5*x^5-18*(c*x-1)^(1...

```

### 3.179.5 Fracas [F]

$$\int x^2(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-(a^2*c^2*d*x^4 - a^2*d*x^2 + (b^2*c^2*d*x^4 - b^2*d*x^2)*arccosh(c*x)^2 + 2*(a*b*c^2*d*x^4 - a*b*d*x^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`



**3.179.6 Sympy [F(-1)]**

Timed out.

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Timed out}$$

input `integrate(x**2*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.179.7 Maxima [F]**

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `1/48*a^2*(2*(-c^2*d*x^2 + d)^(3/2)*x/c^2 - 8*(-c^2*d*x^2 + d)^(5/2)*x/(c^2*d) + 3*sqrt(-c^2*d*x^2 + d)*d*x/c^2 + 3*d^(3/2)*arcsin(c*x)/c^3) + integrate((-c^2*d*x^2 + d)^(3/2)*b^2*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2 + 2*(-c^2*d*x^2 + d)^(3/2)*a*b*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1), x)`

**3.179.8 Giac [F]**

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^2*x^2, x)`

**3.179.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int x^2 (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2} dx$$

input `int(x^2*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2),x)`output `int(x^2*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2), x)`

### 3.180 $\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.180.1 Optimal result . . . . .	1506
3.180.2 Mathematica [A] (verified) . . . . .	1507
3.180.3 Rubi [A] (verified) . . . . .	1507
3.180.4 Maple [B] (verified) . . . . .	1510
3.180.5 Fricas [A] (verification not implemented) . . . . .	1511
3.180.6 Sympy [F] . . . . .	1512
3.180.7 Maxima [A] (verification not implemented) . . . . .	1512
3.180.8 Giac [F(-2)] . . . . .	1513
3.180.9 Mupad [F(-1)] . . . . .	1513

#### 3.180.1 Optimal result

Integrand size = 27, antiderivative size = 348

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = -\frac{16b^2 d(1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{75c^2(1 - cx)(1 + cx)} - \frac{8b^2 d(1 - c^2 x^2)^2 \sqrt{d - c^2 dx^2}}{225c^2(1 - cx)(1 + cx)} - \frac{2b^2 d(1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{125c^2(1 - cx)(1 + cx)} + \frac{2bdx\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{5c\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{4bcdx^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{15\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{2bc^3 dx^5 \sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))}{25\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{5c^2 d}$$

```
output -1/5*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/c^2/d-16/75*b^2*d*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)-8/225*b^2*d*(-c^2*x^2+1)^2*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)-2/125*b^2*d*(-c^2*x^2+1)^3*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)+2/5*b*d*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-4/15*b*c*d*x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/25*b*c^3*d*x^5*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.180.2 Mathematica [A] (verified)**

Time = 1.48 (sec) , antiderivative size = 208, normalized size of antiderivative = 0.60

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx =$$

$$d\sqrt{d - c^2 dx^2} \left( 225a^2(-1 + c^2 x^2)^3 - 30abcx\sqrt{-1 + cx}\sqrt{1 + cx}(15 - 10c^2 x^2 + 3c^4 x^4) + 2b^2(-149 + 187c^2 x^2) \right)$$

input `Integrate[x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`output `-1/1125*(d*sqrt[d - c^2*d*x^2]*(225*a^2*(-1 + c^2*x^2)^3 - 30*a*b*c*x*sqrt[-1 + c*x]*sqrt[1 + c*x]*(15 - 10*c^2*x^2 + 3*c^4*x^4) + 2*b^2*(-149 + 187*c^2*x^2 - 47*c^4*x^4 + 9*c^6*x^6) - 30*b*(-15*a*(-1 + c^2*x^2)^3 + b*c*x*sqrt[-1 + c*x]*sqrt[1 + c*x]*(15 - 10*c^2*x^2 + 3*c^4*x^4))*ArcCosh[c*x] + 225*b^2*(-1 + c^2*x^2)^3*ArcCosh[c*x]^2))/(c^2*(-1 + c^2*x^2))`**3.180.3 Rubi [A] (verified)**Time = 0.81 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.64, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$ , Rules used = {6329, 6304, 6309, 27, 1905, 1576, 1140, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$\downarrow \text{6329}$$

$$\frac{2bd\sqrt{d - c^2 dx^2} \int (1 - cx)^2 (cx + 1)^2 (a + \operatorname{barccosh}(cx)) dx}{5c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{5c^2 d}$$

$$\downarrow \text{6304}$$

$$\frac{2bd\sqrt{d - c^2 dx^2} \int (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) dx}{5c\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{5c^2 d}$$

$$\downarrow \text{6309}$$

$$\frac{2bd\sqrt{d-c^2dx^2}\left(-bc\int\frac{x(3c^4x^4-10c^2x^2+15)}{15\sqrt{cx-1}\sqrt{cx+1}}dx+\frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx))-\frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{5c\sqrt{cx-1}\sqrt{cx+1}} \\ \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\ \downarrow 27$$

$$\frac{2bd\sqrt{d-c^2dx^2}\left(-\frac{1}{15}bc\int\frac{x(3c^4x^4-10c^2x^2+15)}{\sqrt{cx-1}\sqrt{cx+1}}dx+\frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx))-\frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{5c\sqrt{cx-1}\sqrt{cx+1}} \\ \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\ \downarrow 1905$$

$$\frac{2bd\sqrt{d-c^2dx^2}\left(-\frac{bc\sqrt{c^2x^2-1}\int\frac{x(3c^4x^4-10c^2x^2+15)}{\sqrt{c^2x^2-1}}dx}{15\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx))-\frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{5c\sqrt{cx-1}\sqrt{cx+1}} \\ \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\ \downarrow 1576$$

$$\frac{2bd\sqrt{d-c^2dx^2}\left(-\frac{bc\sqrt{c^2x^2-1}\int\frac{3c^4x^4-10c^2x^2+15}{\sqrt{c^2x^2-1}}dx^2}{30\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx))-\frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{5c\sqrt{cx-1}\sqrt{cx+1}} \\ \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\ \downarrow 1140$$

$$\frac{2bd\sqrt{d-c^2dx^2}\left(-\frac{bc\sqrt{c^2x^2-1}\int\left(3(c^2x^2-1)^{3/2}-4\sqrt{c^2x^2-1}+\frac{8}{\sqrt{c^2x^2-1}}\right)dx^2}{30\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx))-\frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx))+x(a+\operatorname{barccosh}(cx))\right)}{5c\sqrt{cx-1}\sqrt{cx+1}} \\ \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\ \downarrow 2009$$

---

3.180.  $\int x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$2bd\sqrt{d - c^2 dx^2} \left( \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)}{5c^2} \right)}{30} \right) - \frac{5c\sqrt{cx - 1}\sqrt{cx + 1}}{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} \frac{1}{5c^2 d}$$

input `Int[x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output `-1/5*((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/(c^2*d) + (2*b*d*Sqrt[d - c^2*d*x^2]*(-1/30*(b*c*Sqrt[-1 + c^2*x^2]*((16*Sqrt[-1 + c^2*x^2])/c^2 - (8*(-1 + c^2*x^2)^(3/2))/(3*c^2) + (6*(-1 + c^2*x^2)^(5/2))/(5*c^2)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + x*(a + b*ArcCosh[c*x]) - (2*c^2*x^3*(a + b*ArcCosh[c*x]))/3 + (c^4*x^5*(a + b*ArcCosh[c*x]))/5)/(5*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.180.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1140 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 0]`

rule 1576 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`

rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`

$$3.180. \quad \int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$$

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x])^u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.180.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1269 vs.  $2(308) = 616$ .

Time = 1.18 (sec) , antiderivative size = 1270, normalized size of antiderivative = 3.65

method	result	size
default	Expression too large to display	1270
parts	Expression too large to display	1270

input `int(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output

```

-1/5*a^2*(-c^2*d*x^2+d)^(5/2)/c^2/d+b^2*(-1/4000*(-d*(c^2*x^2-1))^(1/2)*(1
6*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*
(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(
25*arccosh(c*x)^2-10*arccosh(c*x)+2)*d/(c*x+1)/c^2/(c*x-1)+1/288*(-d*(c^2*
x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(9*arccosh(c*x)^2-6*arccosh(c*x)+2)*d/
(c*x+1)/c^2/(c*x-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/
2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*arccosh(c*x)+2)*d/(c*x+1)/c^2/(c*x-1)-
1/16*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(
arccosh(c*x)^2+2*arccosh(c*x)+2)*d/(c*x+1)/c^2/(c*x-1)+1/288*(-d*(c^2*x^2-
1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2
)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(9*arccosh(c*x)^2+6*arccosh(c*x)+2)*d/(c*
x+1)/c^2/(c*x-1)-1/4000*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)*(c*x-1)^(
1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-28*c^4*x^4
-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(25*arccosh(c*x)^2+10*arc
cosh(c*x)+2)*d/(c*x+1)/c^2/(c*x-1)+2*a*b*(-1/800*(-d*(c^2*x^2-1))^(1/2)*(
16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*
(-1+5*arccosh(c*x))*d/(c*x+1)/c^2/(c*x-1)+1/96*(-d*(c^2*x^2-1))^(1/2)*(4*c
^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3*(c*x-1)^(1/2)*...

```

### 3.180.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.05

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{225(b^2 c^6 dx^6 - 3b^2 c^4 dx^4 + 3b^2 c^2 dx^2 - b^2 d)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1})^2 - 30(3abc^5 dx^5 - 10abc^3 dx^3 + 5abc dx)}{...}$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="fracas")`



output 
$$\begin{aligned} & -1/1125*(225*(b^2*c^6*d*x^6 - 3*b^2*c^4*d*x^4 + 3*b^2*c^2*d*x^2 - b^2*d)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1})^2 - 30*(3*a*b*c^5*d*x^5 - \\ & 10*a*b*c^3*d*x^3 + 15*a*b*c*d*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} - \\ & 30*((3*b^2*c^5*d*x^5 - 10*b^2*c^3*d*x^3 + 15*b^2*c*d*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} - \\ & 15*(a*b*c^6*d*x^6 - 3*a*b*c^4*d*x^4 + 3*a*b*c^2*d*x^2 - a*b*d)*\sqrt{-c^2*d*x^2 + d})*\log(c*x + \sqrt{c^2*x^2 - 1}) + (9*(25*a^2 + 2*b^2)*c^6*d*x^6 - \\ & (675*a^2 + 94*b^2)*c^4*d*x^4 + (675*a^2 + 374*b^2)*c^2*d*x^2 - (225*a^2 + 298*b^2)*d)*\sqrt{-c^2*d*x^2 + d})/(c^4*x^2 - c^2) \end{aligned}$$

### 3.180.6 Sympy [F]

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int x(-d(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))^2 dx$$

input `integrate(x*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2,x)`

output `Integral(x*(-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2, x)`

### 3.180.7 Maxima [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 278, normalized size of antiderivative = 0.80

$$\begin{aligned} \int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx &= -\frac{(-c^2 dx^2 + d)^{\frac{5}{2}} b^2 \operatorname{arccosh}(cx)^2}{5 c^2 d} \\ &- \frac{2}{1125} b^2 \left( \frac{9 \sqrt{c^2 x^2 - 1} c^2 \sqrt{-d d^2} x^4 - 38 \sqrt{c^2 x^2 - 1} \sqrt{-d d^2} x^2 + \frac{149 \sqrt{c^2 x^2 - 1} \sqrt{-d d^2}}{c^2}}{d} - \frac{15 (3 c^4 \sqrt{-d d^2} x^5 - 10 c^2 \sqrt{-d d^2} x^3 + 15 \sqrt{-d d^2} x) a b}{75 c d} \right) \\ &- \frac{2(-c^2 dx^2 + d)^{\frac{5}{2}} a b \operatorname{arccosh}(cx)}{5 c^2 d} - \frac{(-c^2 dx^2 + d)^{\frac{5}{2}} a^2}{5 c^2 d} \\ &+ \frac{2(3 c^4 \sqrt{-d d^2} x^5 - 10 c^2 \sqrt{-d d^2} x^3 + 15 \sqrt{-d d^2} x) a b}{75 c d} \end{aligned}$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

---

3.180.  $\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

```
output -1/5*(-c^2*d*x^2 + d)^(5/2)*b^2*arccosh(c*x)^2/(c^2*d) - 2/1125*b^2*((9*sqrt(c^2*x^2 - 1)*c^2*sqrt(-d)*d^2*x^4 - 38*sqrt(c^2*x^2 - 1)*sqrt(-d)*d^2*x^2 + 149*sqrt(c^2*x^2 - 1)*sqrt(-d)*d^2/c^2)/d - 15*(3*c^4*sqrt(-d)*d^2*x^5 - 10*c^2*sqrt(-d)*d^2*x^3 + 15*sqrt(-d)*d^2*x)*arccosh(c*x)/(c*d)) - 2/5*(-c^2*d*x^2 + d)^(5/2)*a*b*arccosh(c*x)/(c^2*d) - 1/5*(-c^2*d*x^2 + d)^(5/2)*a^2/(c^2*d) + 2/75*(3*c^4*sqrt(-d)*d^2*x^5 - 10*c^2*sqrt(-d)*d^2*x^3 + 15*sqrt(-d)*d^2*x)*a*b/(c*d)
```

### 3.180.8 Giac [F(-2)]

Exception generated.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

```
input integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.180.9 Mupad [F(-1)]

Timed out.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int x(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2} dx$$

```
input int(x*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2),x)
```

```
output int(x*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2), x)
```

### 3.181 $\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.181.1 Optimal result	1514
3.181.2 Mathematica [A] (warning: unable to verify)	1515
3.181.3 Rubi [A] (verified)	1515
3.181.4 Maple [B] (verified)	1520
3.181.5 Fricas [F]	1521
3.181.6 Sympy [F]	1522
3.181.7 Maxima [F]	1522
3.181.8 Giac [F(-2)]	1522
3.181.9 Mupad [F(-1)]	1523

#### 3.181.1 Optimal result

Integrand size = 26, antiderivative size = 336

$$\begin{aligned} \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx &= \frac{15}{64} b^2 dx \sqrt{d - c^2 dx^2} \\ &+ \frac{1}{32} b^2 dx (1 - cx)(1 + cx) \sqrt{d - c^2 dx^2} + \frac{9b^2 d \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{64c \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &- \frac{3bcdx^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{8 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &+ \frac{bd(1 - c^2 x^2)^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{8c \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{3}{8} dx \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \\ &+ \frac{1}{4} x (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{d \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^3}{8bc \sqrt{-1 + cx} \sqrt{1 + cx}} \end{aligned}$$

output

```
1/4*x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2+15/64*b^2*d*x*(-c^2*d*x^2+d)^(1/2)+1/32*b^2*d*x*(-c*x+1)*(c*x+1)*(-c^2*d*x^2+d)^(1/2)+3/8*d*x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)+9/64*b^2*d*arccosh(c*x)*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-3/8*b*c*d*x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/8*b*d*(-c^2*x^2+1)^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/8*d*(a+b*arccosh(c*x))^3*(-c^2*d*x^2+d)^(1/2)/b/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.181.2 Mathematica [A] (warning: unable to verify)**

Time = 3.35 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.11

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{-96a^2 c dx \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (-5+2c^2 x^2) \sqrt{d-c^2 dx^2} - 288a^2 d^{3/2} \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \arctan\left(\frac{cx \sqrt{d-c^2 dx^2}}{\sqrt{d}(-1+c^2 x^2)}\right) - 192a^2 b d \sqrt{d-c^2 dx^2} (\cosh[2 \operatorname{ArcCosh}[cx]] + 2 \operatorname{ArcCosh}[cx] (\operatorname{ArcCosh}[cx] - \sinh[2 \operatorname{ArcCosh}[cx]])) - 32b^2 d \sqrt{d-c^2 dx^2} (4 \operatorname{ArcCosh}[cx]^3 + 6 \operatorname{ArcCosh}[cx] \cosh[2 \operatorname{ArcCosh}[cx]] - 3(1 + 2 \operatorname{ArcCosh}[cx]^2) \sinh[2 \operatorname{ArcCosh}[cx]]) + 12a^2 b d \sqrt{d-c^2 dx^2} (8 \operatorname{ArcCosh}[cx]^2 + \cosh[4 \operatorname{ArcCosh}[cx]] - 4 \operatorname{ArcCosh}[cx] \sinh[4 \operatorname{ArcCosh}[cx]]) + b^2 d \sqrt{d-c^2 dx^2} (32 \operatorname{ArcCosh}[cx]^3 + 12 \operatorname{ArcCosh}[cx] \cosh[4 \operatorname{ArcCosh}[cx]] - 3(1 + 8 \operatorname{ArcCosh}[cx]^2) \sinh[4 \operatorname{ArcCosh}[cx]])}{(768c \sqrt{\frac{-1+cx}{1+cx}} (1+cx))}$$

input `Integrate[(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output

```
(-96*a^2*c*d*x*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(-5 + 2*c^2*x^2)*Sqrt[d - c^2*d*x^2] - 288*a^2*d^(3/2)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] - 192*a*b*d*Sqrt[d - c^2*d*x^2]*(Cosh[2*ArcCosh[c*x]] + 2*ArcCosh[c*x]*(ArcCosh[c*x] - Sinh[2*ArcCosh[c*x]])) - 32*b^2*d*Sqrt[d - c^2*d*x^2]*(4*ArcCosh[c*x]^3 + 6*ArcCosh[c*x]*Cosh[2*ArcCosh[c*x]] - 3*(1 + 2*ArcCosh[c*x]^2)*Sinh[2*ArcCosh[c*x]]) + 12*a*b*d*Sqrt[d - c^2*d*x^2]*(8*ArcCosh[c*x]^2 + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]]) + b^2*d*Sqrt[d - c^2*d*x^2]*(32*ArcCosh[c*x]^3 + 12*ArcCosh[c*x]*Cosh[4*ArcCosh[c*x]] - 3*(1 + 8*ArcCosh[c*x]^2)*Sinh[4*ArcCosh[c*x]])/(768*c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))
```

**3.181.3 Rubi [A] (verified)**Time = 1.54 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.06, number of steps used = 12, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used = {6312, 25, 6310, 6298, 101, 43, 6308, 6327, 6329, 40, 40, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$$

↓ 6312

$$\frac{bcd\sqrt{d-c^2 dx^2} \int \frac{-x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \int \sqrt{d-c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{4}x(d-c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{}$$

↓ 25

---

3.181.  $\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
& -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \int \sqrt{d-c^2dx^2}(a + \\
& \quad \operatorname{barccosh}(cx))^2 dx + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \\
& \quad \downarrow \text{6310} \\
& -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \int x(a+\operatorname{barccosh}(cx))dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 \right. \\
& \quad \left. + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \right) \\
& \quad \downarrow \text{6298} \\
& -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 \right. \\
& \quad \left. + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \right) \\
& \quad \downarrow \text{101} \\
& -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 \right. \\
& \quad \left. + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \right) \\
& \quad \downarrow \text{43} \\
& -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \\
& \quad \left. + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \right) \\
& \quad \downarrow \text{6308}
\end{aligned}$$

$$\begin{aligned}
 & -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 + \\
 & \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{6327} \\
 & -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 + \\
 & \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{6329} \\
 & -\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b \int (cx-1)^{3/2}(cx+1)^{3/2}dx}{4c} - \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 + \\
 & \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{40} \\
 & -\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \int \sqrt{cx-1}\sqrt{cx+1}dx\right)}{4c} - \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \qquad \qquad \qquad \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 + \\
 & \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{40} \\
 & -\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)\right)}{4c} - \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \qquad \qquad \qquad \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 + \\
 & \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{43}
 \end{aligned}$$

---

3.181.  $\int (d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{\frac{1}{4}x(d - c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 - bcd\sqrt{d - c^2dx^2} \left( \frac{b \left( \frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c} \right) \right)}{4c} - \frac{(1-c^2x^2)^2(a + \operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d - c^2dx^2} \left( \frac{1}{2}x^2(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx}} \right)}{\sqrt{cx}}$$

input `Int[(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output `(x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/4 + (3*d*((x*Sqrt[d - c^2*d*x^2])*(a + b*ArcCosh[c*x])^2)/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^3)/(6*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (b*c*Sqrt[d - c^2*d*x^2]*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/2))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/4 - (b*c*d*Sqrt[d - c^2*d*x^2]*(-1/4*((1 - c^2*x^2)^2*(a + b*ArcCosh[c*x]))/c^2 + (b*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/(4*c)))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.181.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 101 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)*(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)n(e + f*x)pSimp[a2d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)((d_.)*(x_))(m_.), x_Symbol] := Simp[(d*x)(m + 1)((a + b*ArcCosh[c*x])n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)(m + 1)((a + b*ArcCosh[c*x])(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)/(Sqrt[(d1_)+(e1_.)*(x_)]*Sqrt[(d2_)+(e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)Sqrt[(d_)+(e_.)*(x_)2], x_Symbol] := Simp[x*Sqrt[d + e*x2]*((a + b*ArcCosh[c*x])(n/2)), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c2*d + e, 0] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)((d_)+(e_.)*(x_)2)(p_.), x_Symbol] := Simp[x*(d + e*x2)p((a + b*ArcCosh[c*x])n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x2)(p - 1)(a + b*ArcCosh[c*x])n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x2)p/((1 + c*x)p(-1 + c*x)p)] Int[x*(1 + c*x)(p - 1/2)(-1 + c*x)(p - 1/2)(a + b*ArcCosh[c*x])(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`



rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.181.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1060 vs.  $2(288) = 576$ .

Time = 1.05 (sec) , antiderivative size = 1061, normalized size of antiderivative = 3.16

method	result	size
default	Expression too large to display	1061
parts	Expression too large to display	1061

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output  $\frac{1}{4}x(-c^2dx^2+d)^{3/2}a^2+3/8a^2dxx(-c^2dx^2+d)^{1/2}+3/8a^2d^2/(c^2d)^{1/2}\arctan((c^2d)^{1/2}x/(-c^2dx^2+d)^{1/2})+b^2(-1/8(-d(c^2x^2-1))^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}/c\operatorname{arccosh}(cx))^3d-1/512(-d(c^2x^2-1))^{1/2}(8c^5x^5-12c^3x^3+8(cx+1)^{1/2}(cx-1)^{1/2}c^4x^4+4cx-8(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2+(cx-1)^{1/2}(cx+1)^{1/2})(8\operatorname{arccosh}(cx)^2-4\operatorname{arccosh}(cx)+1)d/(cx-1)/(cx+1)/c+1/16(-d(c^2x^2-1))^{1/2}(2c^3x^3-2cx+2(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2-(cx-1)^{1/2}(cx+1)^{1/2})(2\operatorname{arccosh}(cx)^2-2\operatorname{arccosh}(cx)+1)d/(cx-1)/(cx+1)/c+1/16(-d(c^2x^2-1))^{1/2}(-2(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2+2c^3x^3+(cx-1)^{1/2}(cx+1)^{1/2}-2cx)(2\operatorname{arccosh}(cx)^2+2\operatorname{arccosh}(cx)+1)d/(cx-1)/(cx+1)/c-1/512(-d(c^2x^2-1))^{1/2}(-8(cx+1)^{1/2}(cx-1)^{1/2}c^4x^4+8c^5x^5+8(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2-12c^3x^3-(cx-1)^{1/2}(cx+1)^{1/2}+4cx)(8\operatorname{arccosh}(cx)^2+4\operatorname{arccosh}(cx)+1)d/(cx-1)/(cx+1)/c)+2ab(-3/16(-d(c^2x^2-1))^{1/2}/(cx-1)^{1/2}/(cx+1)^{1/2}/c\operatorname{arccosh}(cx))^2d-1/256(-d(c^2x^2-1))^{1/2}(8c^5x^5-12c^3x^3+8(cx+1)^{1/2}(cx-1)^{1/2}c^4x^4+4cx-8(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2+(cx-1)^{1/2}(cx+1)^{1/2})(-1+4\operatorname{arccosh}(cx))d/(cx-1)/(cx+1)/c+1/16(-d(c^2x^2-1))^{1/2}(2c^3x^3-2cx+2(cx-1)^{1/2}(cx+1)^{1/2}c^2x^2-(cx-1)^{1/2}(cx+1)^{1/2})(-1+2\operatorname{arccosh}(cx))d/(cx-1)/(cx+1)/c+1/16(-d(c^2x^2-1))^{1/2}(-2(cx-1)^{1/2}(cx...$

### 3.181.5 Fracas [F]

$$\int (d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a)^2 dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-(a^2*c^2*d*x^2 - a^2*d + (b^2*c^2*d*x^2 - b^2*d)*arccosh(c*x))^2 + 2*(a*b*c^2*d*x^2 - a*b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

**3.181.6 Sympy [F]**

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-d(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))^2 dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2, x)`

**3.181.7 Maxima [F]**

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2 dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `1/8*(2*(-c^2*d*x^2 + d)^(3/2)*x + 3*sqrt(-c^2*d*x^2 + d)*d*x + 3*d^(3/2)*arcsin(c*x)/c)*a^2 + integrate((-c^2*d*x^2 + d)^(3/2)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2 + 2*(-c^2*d*x^2 + d)^(3/2)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)), x)`

**3.181.8 Giac [F(-2)]**

Exception generated.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.181.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2} dx$$

input `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2),x)`output `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2), x)`

**3.182** 
$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x} dx$$

3.182.1 Optimal result . . . . .	1524
3.182.2 Mathematica [A] (warning: unable to verify) . . . . .	1525
3.182.3 Rubi [A] (verified) . . . . .	1526
3.182.4 Maple [F] . . . . .	1532
3.182.5 Fricas [F] . . . . .	1533
3.182.6 Sympy [F] . . . . .	1533
3.182.7 Maxima [F] . . . . .	1533
3.182.8 Giac [F(-2)] . . . . .	1534
3.182.9 Mupad [F(-1)] . . . . .	1534

**3.182.1 Optimal result**

Integrand size = 29, antiderivative size = 573

$$\begin{aligned} \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x} dx &= \frac{68}{27}b^2d\sqrt{d-c^2dx^2} \\ &- \frac{2}{27}b^2c^2dx^2\sqrt{d-c^2dx^2} - \frac{2abcdx\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}} - \frac{2b^2cdx\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{2bcdx\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3\sqrt{-1+cx}\sqrt{1+cx}} + \frac{2bc^3dx^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{9\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2 \\ &- \frac{2d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{2ibd\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{2ibd\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{2ib^2d\sqrt{d-c^2dx^2} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{2ib^2d\sqrt{d-c^2dx^2} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output  $1/3*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^{2+68/27*b^2*d*(-c^2*d*x^2+d)^{(1/2)}-2/27*b^2*c^2*d*x^2*(-c^2*d*x^2+d)^{(1/2)}+d*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d*x^2+d)^{(1/2)}-2*a*b*c*d*x*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}-2*b^2*c*d*x*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}-2/3*b*c*d*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}+2/9*b*c^3*d*x^3*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}-2*d*(a+b*\operatorname{arccosh}(c*x))^{2*\arctan(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}+2*I*b*d*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}-2*I*b*d*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}-2*I*b^2*d*\operatorname{polylog}(3,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}+2*I*b^2*d*\operatorname{polylog}(3,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*(-c^2*d*x^2+d)^{(1/2)/(c*x-1)^{(1/2)/(c*x+1)^{(1/2)}}}$

### 3.182.2 Mathematica [A] (warning: unable to verify)

Time = 2.18 (sec) , antiderivative size = 650, normalized size of antiderivative = 1.13

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x} dx = -\frac{1}{3} a^2 d (-4 + c^2 x^2) \sqrt{d - c^2 dx^2}$$

$$- \frac{1}{54} b^2 d \sqrt{d - c^2 dx^2} \left( 2(-13 + \cosh(2 \operatorname{arccosh}(cx))) + 9 \operatorname{arccosh}(cx)^2 (-1 + \cosh(2 \operatorname{arccosh}(cx))) \right.$$

$$\left. + \frac{3 \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) (9cx - \cosh(3 \operatorname{arccosh}(cx)))}{-1 + cx} \right)$$

$$- \frac{abd \sqrt{d - c^2 dx^2} \left( 9cx + 12 \left( \frac{-1+cx}{1+cx} \right)^{3/2} (1 + cx)^3 \operatorname{arccosh}(cx) - \cosh(3 \operatorname{arccosh}(cx)) \right)}{18 \sqrt{\frac{-1+cx}{1+cx}} (1 + cx)}$$

$$+ a^2 d^{3/2} \log(cx) - a^2 d^{3/2} \log \left( d + \sqrt{d} \sqrt{d - c^2 dx^2} \right) + \frac{2abd \sqrt{d - c^2 dx^2} \left( -cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx \sqrt{\frac{-1+cx}{1+cx}} \right)}{18 \sqrt{\frac{-1+cx}{1+cx}} (1 + cx)}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x,x]`

output

```

-1/3*(a^2*d*(-4 + c^2*x^2)*Sqrt[d - c^2*d*x^2]) - (b^2*d*Sqrt[d - c^2*d*x^
2]*(2*(-13 + Cosh[2*ArcCosh[c*x]]) + 9*ArcCosh[c*x]^2*(-1 + Cosh[2*ArcCosh
[c*x]])) + (3*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*(9*c*x - Cosh[3*ArcCo
sh[c*x]])))/(-1 + c*x))/54 - (a*b*d*Sqrt[d - c^2*d*x^2]*(9*c*x + 12*((-1 +
c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3*ArcCosh[c*x] - Cosh[3*ArcCosh[c*x]]))/
(18*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + a^2*d^(3/2)*Log[c*x] - a^2*d^(3
/2)*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (2*a*b*d*Sqrt[d - c^2*d*x^2]*(-
(c*x) + Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + c*x*Sqrt[(-1 + c*x)/(1 +
c*x)]*ArcCosh[c*x] + I*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh
[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*Po
lyLog[2, I/E^ArcCosh[c*x]]))/((Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + b^2*
d*Sqrt[d - c^2*d*x^2]*(2 + (2*c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x])
/(1 - c*x) + ArcCosh[c*x]^2 + (I*(ArcCosh[c*x]^2*Log[1 - I/E^ArcCosh[c*x]]
- ArcCosh[c*x]^2*Log[1 + I/E^ArcCosh[c*x]] + 2*ArcCosh[c*x]*PolyLog[2, (-
I)/E^ArcCosh[c*x]] - 2*ArcCosh[c*x]*PolyLog[2, I/E^ArcCosh[c*x]] + 2*PolyL
og[3, (-I)/E^ArcCosh[c*x]] - 2*PolyLog[3, I/E^ArcCosh[c*x]]))/((Sqrt[(-1 +
c*x)/(1 + c*x)]*(1 + c*x)))

```

### 3.182.3 Rubi [A] (verified)

Time = 2.50 (sec) , antiderivative size = 396, normalized size of antiderivative = 0.69, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.517$ , Rules used = {6345, 25, 6304, 6309, 27, 960, 83, 6341, 2009, 6362, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx \\
 & \quad \downarrow \text{6345} \\
 & \frac{2bcd\sqrt{d - c^2 dx^2} \int -((1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \\
 & d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{25} \\
 & - \frac{2bcd\sqrt{d - c^2 dx^2} \int (1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx)) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \\
 & d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2
 \end{aligned}$$

---

3.182.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$

$$\begin{aligned}
& \downarrow 6304 \\
& \frac{2bcd\sqrt{d-c^2dx^2} \int (1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx + \\
& \quad \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 6309 \\
& \frac{d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx -}{2bcd\sqrt{d-c^2dx^2} \left( -bc \int \frac{x(3-c^2x^2)}{3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
& \quad \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 27 \\
& \frac{d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx -}{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}bc \int \frac{x(3-c^2x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
& \quad \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 960 \\
& \frac{d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx -}{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}bc \left( \frac{7}{3} \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) - \frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \\
& \downarrow 83 \\
& \frac{d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 -}{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow 6341
\end{aligned}$$



$$d \left( -\frac{2bc\sqrt{d-c^2dx^2} \int (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx)) \right. \\ \left. - \frac{\frac{1}{3}(d-c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}c^2x^3(a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 2009

$$d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax + b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{\frac{1}{3}(d-c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}c^2x^3(a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6362

$$d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax + b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{\frac{1}{3}(d-c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}c^2x^3(a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 3042

$$d \left( -\frac{\sqrt{d-c^2dx^2} \int (a + \operatorname{barccosh}(cx))^2 \csc \left( \operatorname{iarccosh}(cx) + \frac{\pi}{2} \right) \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax + b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{\frac{1}{3}(d-c^2dx^2)^{3/2}(a + \operatorname{barccosh}(cx))^2 - 2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}c^2x^3(a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 4668

---

3.182.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x} dx$

$$d \left( - \frac{\sqrt{d - c^2 dx^2} (-2ib \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) dx)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right. \\ \left. - \frac{\frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd \sqrt{d - c^2 dx^2} \left( -\frac{1}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3} bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3} x^2 \sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \downarrow \text{3011}$$

$$d \left( - \frac{\sqrt{d - c^2 dx^2} (2ib (b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) dx)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right. \\ \left. - \frac{\frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd \sqrt{d - c^2 dx^2} \left( -\frac{1}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3} bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3} x^2 \sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \downarrow \text{2720}$$

$$d \left( - \frac{\sqrt{d - c^2 dx^2} (2ib (b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) dx)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right. \\ \left. - \frac{\frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd \sqrt{d - c^2 dx^2} \left( -\frac{1}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3} bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3} x^2 \sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right) \\ \downarrow \text{7143}$$

$$d \left( - \frac{\sqrt{d - c^2 dx^2} (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx))^2 + 2ib (b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) dx)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right. \\ \left. - \frac{\frac{1}{3} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd \sqrt{d - c^2 dx^2} \left( -\frac{1}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{1}{3} bc \left( \frac{7\sqrt{cx-1}\sqrt{cx+1}}{3c^2} - \frac{1}{3} x^2 \sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x,x]`

```
output ((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/3 - (2*b*c*d*Sqrt[d - c^2*d
*x^2]*(-1/3*(b*c*((7*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2) - (x^2*Sqrt[-1
+ c*x]*Sqrt[1 + c*x])/3)) + x*(a + b*ArcCosh[c*x]) - (c^2*x^3*(a + b*ArcCo
sh[c*x]))/3))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + d*(Sqrt[d - c^2*d*x^2]*(a
+ b*ArcCosh[c*x])^2 - (2*b*c*Sqrt[d - c^2*d*x^2]*(a*x - (b*Sqrt[-1 + c*x]
*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (S
qrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I
)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3
, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, I*E^A
rcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]])))/(Sqrt[-1 + c*x]*Sqrt[1 +
c*x]))
```

### 3.182.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 83 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f
*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]
```

```
rule 960 Int[((e_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)
*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(
m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n
*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/
(b1*b2*(m + n*(p + 1) + 1) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n
/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2,
n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*x)))]^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6304 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(p_), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6309 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6362 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.182.4 Maple [F]

$$\int \frac{(-c^2 d x^2 + d)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x} dx$$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x,x)`

output `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x,x)`

---

3.182.  $\int \frac{(d - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x} dx$

**3.182.5 Fricas [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x,x, algorithm="fricas")`

output `integral(-(a^2*c^2*d*x^2 - a^2*d + (b^2*c^2*d*x^2 - b^2*d)*arccosh(c*x)^2 + 2*(a*b*c^2*d*x^2 - a*b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x, x)`

**3.182.6 Sympy [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \int \frac{(-d(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))^2}{x} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2/x,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2/x, x)`

**3.182.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x,x, algorithm="maxima")`

output `-1/3*(3*d^(3/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - (-c^2*d*x^2 + d)^(3/2) - 3*sqrt(-c^2*d*x^2 + d)*d*a^2 + integrate((-c^2*d*x^2 + d)^(3/2)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x + 2*(-c^2*d*x^2 + d)^(3/2)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x, x)`

**3.182.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.182.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2}}{x} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x, x)`

**3.183**  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx$

3.183.1 Optimal result . . . . .	1535
3.183.2 Mathematica [A] (warning: unable to verify) . . . . .	1536
3.183.3 Rubi [C] (warning: unable to verify) . . . . .	1537
3.183.4 Maple [A] (verified) . . . . .	1544
3.183.5 Fricas [F] . . . . .	1545
3.183.6 Sympy [F] . . . . .	1545
3.183.7 Maxima [F] . . . . .	1546
3.183.8 Giac [F(-2)] . . . . .	1546
3.183.9 Mupad [F(-1)] . . . . .	1546

**3.183.1 Optimal result**

Integrand size = 29, antiderivative size = 453

$$\begin{aligned} \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx &= -\frac{1}{4}b^2c^2dx\sqrt{d-c^2dx^2} \\ &- \frac{5b^2cd\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{4\sqrt{-1+cx}\sqrt{1+cx}} + \frac{3bc^3dx^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{2\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{bcd(1-c^2x^2)\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{3}{2}c^2dx\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 + \frac{cd\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x} + \frac{cd\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^3}{2b\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{2bcd\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\log(1+e^{-2\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{b^2cd\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-e^{-2\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$



output 
$$\begin{aligned} & -(-c^2 d x^2 + d)^{3/2} (a + b \operatorname{arccosh}(c x))^2 / x - 1/4 b^2 c^2 d x (-c^2 d x^2 + d)^{1/2} - 3/2 c^2 d x (a + b \operatorname{arccosh}(c x))^2 (-c^2 d x^2 + d)^{1/2} - 5/4 b^2 c^2 d \operatorname{arccosh}(c x) (-c^2 d x^2 + d)^{1/2} / (c x - 1)^{1/2} / (c x + 1)^{1/2} + 3/2 b^2 c^3 d x^2 (a + b \operatorname{arccosh}(c x)) (-c^2 d x^2 + d)^{1/2} / (c x - 1)^{1/2} / (c x + 1)^{1/2} + b^2 c^2 (-c^2 x^2 + 1) (a + b \operatorname{arccosh}(c x)) (-c^2 d x^2 + d)^{1/2} / (c x - 1)^{1/2} / (c x + 1)^{1/2} + c^2 d (a + b \operatorname{arccosh}(c x))^2 (-c^2 d x^2 + d)^{1/2} / (c x - 1)^{1/2} / (c x + 1)^{1/2} + 1/2 c^2 d (a + b \operatorname{arccosh}(c x))^3 (-c^2 d x^2 + d)^{1/2} / b / (c x - 1)^{1/2} / (c x + 1)^{1/2} + 2 b^2 c^2 d (a + b \operatorname{arccosh}(c x)) \ln(1 + 1 / (c x + (c x - 1)^{1/2} (c x + 1)^{1/2}))^2 (-c^2 d x^2 + d)^{1/2} / (c x - 1)^{1/2} / (c x + 1)^{1/2} - b^2 c^2 d \operatorname{polylog}(2, -1 / (c x + (c x - 1)^{1/2} (c x + 1)^{1/2}))^2 (-c^2 d x^2 + d)^{1/2} / (c x - 1)^{1/2} / (c x + 1)^{1/2} \end{aligned}$$

### 3.183.2 Mathematica [A] (warning: unable to verify)

Time = 4.19 (sec) , antiderivative size = 433, normalized size of antiderivative = 0.96

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^2} dx = \frac{-12a^2 d \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (2+c^2x^2) \sqrt{d-c^2dx^2} + 36a^2 cd^{3/2} x \sqrt{\frac{-1+cx}{1+cx}}}{x^2}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^2,x]`

output 
$$\begin{aligned} & (-12 a^2 d \operatorname{Sqrt}[(-1 + c x) / (1 + c x)] (1 + c x) (2 + c^2 x^2) \operatorname{Sqrt}[d - c^2 d x^2] + 36 a^2 c d^{3/2} x \operatorname{Sqrt}[(-1 + c x) / (1 + c x)] (1 + c x) \operatorname{ArcTan}[(c x \operatorname{Sqrt}[d - c^2 d x^2]) / (\operatorname{Sqrt}[d] (-1 + c^2 x^2))] - 24 a b d \operatorname{Sqrt}[d - c^2 d x^2] (2 \operatorname{Sqrt}[(-1 + c x) / (1 + c x)] (1 + c x) \operatorname{ArcCosh}[c x] - c x (\operatorname{ArcCos h}[c x]^2 + 2 \operatorname{Log}[c x])) - 8 b^2 d \operatorname{Sqrt}[d - c^2 d x^2] (\operatorname{ArcCosh}[c x] (3 \operatorname{Sqrt}[(-1 + c x) / (1 + c x)] (1 + c x) \operatorname{ArcCosh}[c x] - c x (\operatorname{ArcCosh}[c x] (3 + \operatorname{ArcCosh}[c x]) + 6 \operatorname{Log}[1 + E^{(-2 \operatorname{ArcCosh}[c x])}])) + 3 c x \operatorname{PolyLog}[2, -E^{(-2 \operatorname{ArcCosh}[c x])}]) + 6 a b c d x \operatorname{Sqrt}[d - c^2 d x^2] (\operatorname{Cosh}[2 \operatorname{ArcCosh}[c x]] + 2 \operatorname{ArcCosh}[c x] (\operatorname{ArcCosh}[c x] - \operatorname{Sinh}[2 \operatorname{ArcCosh}[c x]]) + b^2 c d x \operatorname{Sqrt}[d - c^2 d x^2] (4 \operatorname{ArcCosh}[c x]^3 + 6 \operatorname{ArcCosh}[c x] \operatorname{Cosh}[2 \operatorname{ArcCosh}[c x]] - 3 (1 + 2 \operatorname{ArcCosh}[c x]^2) \operatorname{Sinh}[2 \operatorname{ArcCosh}[c x])]) / (24 x \operatorname{Sqrt}[(-1 + c x) / (1 + c x)] (1 + c x)) \end{aligned}$$

**3.183.3 Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 2.36 (sec) , antiderivative size = 393, normalized size of antiderivative = 0.87, number of steps used = 20, number of rules used = 19,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.655$ , Rules used = {6343, 25, 6310, 6298, 101, 43, 6308, 6327, 6334, 40, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx \\
 & \quad \downarrow \text{6343} \\
 & -3c^2 d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2 dx^2} \int \frac{-(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{25} \\
 & -3c^2 d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{2bcd\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{6310} \\
 & -3c^2 d \left( -\frac{bc\sqrt{d - c^2 dx^2} \int x(a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2} x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \right) \\
 & \quad \frac{2bcd\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{6298} \\
 & -3c^2 d \left( -\frac{bc\sqrt{d - c^2 dx^2} \left( \frac{1}{2} x^2 (a + \operatorname{barccosh}(cx)) - \frac{1}{2} bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2} x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \right) \\
 & \quad \frac{2bcd\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{101}
 \end{aligned}$$

---

3.183.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx$

$$\begin{aligned}
 & -3c^2d \left( \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a + \operatorname{barccosh}(cx)) - \frac{1}{2}bc \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. \frac{2bcd\sqrt{d-c^2dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \right) \\
 & \qquad \qquad \qquad \downarrow 43 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & 3c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \right) \\
 & \qquad \qquad \qquad \downarrow 6308 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} - \\
 & 3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \right) \\
 & \qquad \qquad \qquad \downarrow 6327 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} - \\
 & 3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \right) \\
 & \qquad \qquad \qquad \downarrow 6334 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}bc \int \sqrt{cx-1}\sqrt{cx+1} dx + \frac{1}{2}(1-c^2x^2)(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} - \\
 & 3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a + \operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. \frac{(d-c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} \right) \\
 & \qquad \qquad \qquad \downarrow 40
 \end{aligned}$$

---

3.183.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx$

$$\frac{2bcd\sqrt{d-c^2dx^2}\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{2}bc\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d\left(-\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2-\frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}}\right)$$

↓ 43

$$\frac{2bcd\sqrt{d-c^2dx^2}\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))+\frac{1}{2}bc\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d\left(-\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2-\frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}}\right)$$

↓ 6297

$$\frac{2bcd\sqrt{d-c^2dx^2}\left(\frac{f-\left((a+\operatorname{barccosh}(cx))\tanh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)\right)d(a+\operatorname{barccosh}(cx))}{b}+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d\left(-\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2-\frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}}\right)$$

↓ 25

$$\frac{2bcd\sqrt{d-c^2dx^2}\left(-\frac{f(a+\operatorname{barccosh}(cx))\tanh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)d(a+\operatorname{barccosh}(cx))}{b}+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))+\frac{1}{2}bc\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d\left(-\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2-\frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}}\right)$$

↓ 3042

---

3.183.  $\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx$

$$\frac{2bcd\sqrt{d-c^2dx^2} \left( -\frac{\int -i(a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a+\operatorname{barccosh}(cx))}{b} + \frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 26

$$\frac{2bcd\sqrt{d-c^2dx^2} \left( \frac{i \int (a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a+\operatorname{barccosh}(cx))}{b} + \frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 4201

$$\frac{2bcd\sqrt{d-c^2dx^2} \left( \frac{i \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx)) d(a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2}{1+e^{-2\operatorname{arccosh}(cx)}} \right)}{b} + \frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 2620

$$\frac{2bcd\sqrt{d-c^2dx^2} \left( \frac{i \left( 2i \left( \frac{1}{2}b \int \log(1+e^{-2\operatorname{arccosh}(cx)}) d(a+\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1)(a+\operatorname{barccosh}(cx)) \right) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2 \right)}{b} \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2}\left(\frac{1}{2}x^2(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

---

3.183.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx$

↓ 2715

$$2bcd\sqrt{d-c^2dx^2} \left( \frac{i \left( 2i \left( -\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1+e^{-2\operatorname{arccosh}(cx)}) dx - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{arccosh}(cx)) - \frac{1}{2}i \right)}{b} \right)$$

---


$$\frac{(d-c^2dx^2)^{3/2} (a+\operatorname{arccosh}(cx))^2}{x} - \frac{3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{arccosh}(cx))^2 \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2838

$$2bcd\sqrt{d-c^2dx^2} \left( \frac{i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{arccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{arccosh}(cx)) - \frac{1}{2}i(a+\operatorname{arccosh}(cx))^2 \right)}{b} + \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{arccosh}(cx))^2 \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

---


$$\frac{(d-c^2dx^2)^{3/2} (a+\operatorname{arccosh}(cx))^2}{x} - \frac{3c^2d \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^3}{6bc\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{arccosh}(cx))^2 \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^2,x]`

output `-(((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x) - 3*c^2*d*((x*sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/2 - (sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^3)/(6*b*c*sqrt[-1 + c*x]*sqrt[1 + c*x]) - (b*c*sqrt[d - c^2*d*x^2]*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/2))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + (2*b*c*d*sqrt[d - c^2*d*x^2]*(((1 - c^2*x^2)*(a + b*ArcCosh[c*x]))/2 + (b*c*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*c) - ArcCosh[c*x]/(2*c))))/2 + (I*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])]) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4)))/b))/(sqrt[-1 + c*x]*sqrt[1 + c*x])`

## 3.183.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 40 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^(m/(2*m + 1))), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_) + (b_)*(x_))^(n_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]), x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`



```
rule 6334 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.)/(x_),
  x_Symbol] := Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d
  Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d
  )^p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x]) /; FreeQ
  [{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

```
rule 6343 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_
  .)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
  Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m
  + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(
  m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)
  *(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x],
  x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && G
  tQ[p, 0] && LtQ[m, -1]
```

### 3.183.4 Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.02

method	result
default	$-\frac{a^2(-c^2dx^2+d)^{\frac{5}{2}}}{dx} - a^2c^2x(-c^2dx^2+d)^{\frac{3}{2}} - \frac{3a^2c^2dx\sqrt{-c^2dx^2+d}}{2} - \frac{3a^2c^2d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2\sqrt{c^2d}} + \frac{b^2\sqrt{-d(c^2x^2+d)}}{\sqrt{c^2d}}$
parts	$-\frac{a^2(-c^2dx^2+d)^{\frac{5}{2}}}{dx} - a^2c^2x(-c^2dx^2+d)^{\frac{3}{2}} - \frac{3a^2c^2dx\sqrt{-c^2dx^2+d}}{2} - \frac{3a^2c^2d^2 \arctan\left(\frac{\sqrt{c^2dx}}{\sqrt{-c^2dx^2+d}}\right)}{2\sqrt{c^2d}} + \frac{b^2\sqrt{-d(c^2x^2+d)}}{\sqrt{c^2d}}$

```
input int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^2,x,method=_RETURNVERBOSE)
```

output `-a^2/d/x*(-c^2*d*x^2+d)^(5/2)-a^2*c^2*x*(-c^2*d*x^2+d)^(3/2)-3/2*a^2*c^2*d*x*(-c^2*d*x^2+d)^(1/2)-3/2*a^2*c^2*d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+1/4*b^2*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x*(-2*arccosh(c*x)^2*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^2*c^2+2*c^3*x^3*arccosh(c*x)-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*arccosh(c*x)^3*x*c-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2-4*arccosh(c*x)^2*x*c+8*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-c*x*arccosh(c*x)+4*polylg(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c)*d+1/4*a*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x*(-4*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2+2*c^3*x^3+6*arccosh(c*x)^2*x*c-8*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-8*c*x*arccosh(c*x)+8*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-c*x)*d`

### 3.183.5 Fracas [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a)^2}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^2,x, algorithm="fricas")`

output `integral(-(a^2*c^2*d*x^2 - a^2*d + (b^2*c^2*d*x^2 - b^2*d)*arccosh(c*x))^2 + 2*(a*b*c^2*d*x^2 - a*b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^2, x)`

### 3.183.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^2} dx = \int \frac{(-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx))^2}{x^2} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2/x**2,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2/x**2, x)`

**3.183.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a)^2}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^2,x, algorithm="maxima")`

output `-1/2*(3*sqrt(-c^2*d*x^2 + d)*c^2*d*x + 3*c*d^(3/2)*arcsin(c*x) + 2*(-c^2*d*x^2 + d)^(3/2)/x)*a^2 + integrate((-c^2*d*x^2 + d)^(3/2)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x^2 + 2*(-c^2*d*x^2 + d)^(3/2)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^2, x)`

**3.183.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.183.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x^2,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x^2, x)`

---

3.183.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx$

$$3.184 \quad \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx$$

3.184.1 Optimal result	1547
3.184.2 Mathematica [B] (warning: unable to verify)	1548
3.184.3 Rubi [A] (verified)	1548
3.184.4 Maple [F]	1555
3.184.5 Fricas [F]	1555
3.184.6 Sympy [F]	1556
3.184.7 Maxima [F]	1556
3.184.8 Giac [F(-2)]	1556
3.184.9 Mupad [F(-1)]	1557

### 3.184.1 Optimal result

Integrand size = 29, antiderivative size = 630

$$\begin{aligned} & \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx = -2b^2c^2d\sqrt{d-c^2dx^2} \\ & + \frac{3abc^3dx\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{3b^2c^3dx\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & - \frac{bcd\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{x\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bc^3dx\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & - \frac{3}{2}c^2d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 - \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{2x^2} \\ & + \frac{3c^2d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2\arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{b^2c^2d\sqrt{d-c^2dx^2}\arctan(\sqrt{-1+cx}\sqrt{1+cx})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & - \frac{3ibc^2d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{3ibc^2d\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{3ib^2c^2d\sqrt{d-c^2dx^2}\operatorname{PolyLog}(3,-ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & - \frac{3ib^2c^2d\sqrt{d-c^2dx^2}\operatorname{PolyLog}(3,ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

---


$$3.184. \quad \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx$$

output 
$$-1/2*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^2/x^2-2*b^2*c^2*d*(-c^2*d*x^2+d)^{(1/2)}-3/2*c^2*d*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}+3*a*b*c^3*d*x*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3*b^2*c^3*d*x*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b*c*d*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/x/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b*c^3*d*x*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/x/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3*c^2*d*(a+b*\operatorname{arccosh}(c*x))^2*\arctan(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+b^2*c^2*d*\arctan((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3*I*b*c^2*d*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3*I*b*c^2*d*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3*I*b^2*c^2*d*\operatorname{polylog}(3,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3*I*b^2*c^2*d*\operatorname{polylog}(3,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

### 3.184.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 5444 vs.  $2(630) = 1260$ .

Time = 63.65 (sec) , antiderivative size = 5444, normalized size of antiderivative = 8.64

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx = \text{Result too large to show}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^3,x]`

output `Result too large to show`

### 3.184.3 Rubi [A] (verified)

Time = 2.65 (sec) , antiderivative size = 387, normalized size of antiderivative = 0.61, number of steps used = 17, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.552$ , Rules used = {6343, 25, 6327, 6336, 25, 960, 103, 218, 6341, 2009, 6362, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.184. 
$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx$$

$$\begin{aligned}
& \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx \\
& \quad \downarrow \text{6343} \\
& -\frac{3}{2}c^2 d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx - \frac{bcd\sqrt{d - c^2 dx^2} \int -\frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x^2} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{25} \\
& -\frac{3}{2}c^2 d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x^2} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{6327} \\
& -\frac{3}{2}c^2 d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{bcd\sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)(a+\operatorname{barccosh}(cx))}{x^2} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{6336} \\
& -\frac{3}{2}c^2 d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
& \frac{bcd\sqrt{d - c^2 dx^2} \left( -bc \int -\frac{c^2 x^2 + 1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + c^2(-x)(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{25} \\
& -\frac{3}{2}c^2 d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
& \frac{bcd\sqrt{d - c^2 dx^2} \left( bc \int \frac{c^2 x^2 + 1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + c^2(-x)(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \\
& \quad \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
& \quad \downarrow \text{960}
\end{aligned}$$

---

3.184.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$

$$\begin{aligned}
 & -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx + \\
 & \frac{bcd\sqrt{d-c^2dx^2}\left(bc\left(\int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \sqrt{cx-1}\sqrt{cx+1}\right) + c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} 2x^2} \\
 & \qquad \qquad \qquad \downarrow \text{103} \\
 & -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx + \\
 & \frac{bcd\sqrt{d-c^2dx^2}\left(bc\left(c \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1}\right) + c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} 2x^2} \\
 & \qquad \qquad \qquad \downarrow \text{218} \\
 & -\frac{3}{2}c^2d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx + \\
 & \frac{bcd\sqrt{d-c^2dx^2}\left(c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1})\right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} 2x^2} \\
 & \qquad \qquad \qquad \downarrow \text{6341} \\
 & -\frac{3}{2}c^2d \left( -\frac{2bc\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) \\
 & \frac{bcd\sqrt{d-c^2dx^2}\left(c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1})\right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} 2x^2} \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & -\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\
 & \frac{bcd\sqrt{d-c^2dx^2}\left(c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1})\right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} 2x^2}
 \end{aligned}$$

---

3.184.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^3} dx$

↓ 6362

$$\frac{-\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+\sqrt{d-c^2dx^2})}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{bcd\sqrt{d-c^2dx^2} \left( c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1}) \right)}$$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{1}{2x^2}$$

↓ 3042

$$\frac{-\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx))^2 \csc\left(\operatorname{iarcosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+\sqrt{d-c^2dx^2})}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{bcd\sqrt{d-c^2dx^2} \left( c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1}) \right)}$$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{1}{2x^2}$$

↓ 4668

$$\frac{-\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} (-2ib \int (a+\operatorname{barccosh}(cx)) \log(1-ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a+\operatorname{barccosh}(cx)) \log(1+ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+\sqrt{d-c^2dx^2})}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{bcd\sqrt{d-c^2dx^2} \left( c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1}) \right)}$$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{1}{2x^2}$$

↓ 3011

$$\frac{-\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2} (2ib(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)))}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+\sqrt{d-c^2dx^2})}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{bcd\sqrt{d-c^2dx^2} \left( c^2(-x)(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1}) \right)}$$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{1}{2x^2}$$

↓ 2720

---

3.184.  $\int \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x^3} dx$



$$-\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2}(2ib(b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1})))}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} \right. \\ \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right) \quad \downarrow \quad 7143$$

$$-\frac{3}{2}c^2d \left( -\frac{\sqrt{d-c^2dx^2}(2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx))^2 + 2ib(b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + bc(\arctan(\sqrt{cx-1}\sqrt{cx+1}) + \sqrt{cx-1}\sqrt{cx+1})))}{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} \right. \\ \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^3, x]`

output `-1/2*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^2 + (b*c*d*Sqrt[d - c^2*d*x^2]*(-(a + b*ArcCosh[c*x])/x) - c^2*x*(a + b*ArcCosh[c*x]) + b*c*(Sqrt[-1 + c*x]*Sqrt[1 + c*x] + ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]]))/((Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (3*c^2*d*(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2 - (2*b*c*Sqrt[d - c^2*d*x^2]*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/c + b*x*ArcCosh[c*x]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-(a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-(a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]])))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/2`

## 3.184.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 103 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]*((e_) + (f_)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`
- rule 218 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 960 `Int[((e_)*(x_)^(m_))*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)^v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_)^(m_)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1]`

rule 6362 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_)]/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.184.4 Maple [F]

$$\int \frac{(-c^2 d x^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(c x))^2}{x^3} dx$$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^3,x)`

output `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^3,x)`

### 3.184.5 Fracas [F]

$$\int \frac{(d - c^2 d x^2)^{3/2} (a + b \operatorname{arccosh}(c x))^2}{x^3} dx = \int \frac{(-c^2 d x^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(c x) + a)^2}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^3,x, algorithm="fracas")`

output `integral(-(a^2*c^2*d*x^2 - a^2*d + (b^2*c^2*d*x^2 - b^2*d)*arccosh(c*x)^2 + 2*(a*b*c^2*d*x^2 - a*b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^3, x)`

**3.184.6 Sympy [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \int \frac{(-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx))^2}{x^3} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2/x**3,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2/x**3, x)`

**3.184.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a)^2}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^3,x, algorithm="maxima")`

output `1/2*(3*c^2*d^(3/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - (-c^2*d*x^2 + d)^(3/2)*c^2 - 3*sqrt(-c^2*d*x^2 + d)*c^2*d - (-c^2*d*x^2 + d)^(5/2)/(d*x^2))*a^2 + integrate((-c^2*d*x^2 + d)^(3/2)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x^3 + 2*(-c^2*d*x^2 + d)^(3/2)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^3, x)`

**3.184.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.184.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2}}{x^3} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x^3,x)`output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x^3, x)`

**3.185**  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx$

3.185.1 Optimal result . . . . . 1558  
 3.185.2 Mathematica [A] (warning: unable to verify) . . . . . 1559  
 3.185.3 Rubi [C] (warning: unable to verify) . . . . . 1560  
 3.185.4 Maple [A] (verified) . . . . . 1568  
 3.185.5 Fricas [F] . . . . . 1569  
 3.185.6 Sympy [F] . . . . . 1569  
 3.185.7 Maxima [F] . . . . . 1570  
 3.185.8 Giac [F(-2)] . . . . . 1570  
 3.185.9 Mupad [F(-1)] . . . . . 1570

**3.185.1 Optimal result**

Integrand size = 29, antiderivative size = 426

$$\int \frac{(d - c^2dx^2)^{3/2} (a + b\operatorname{arccosh}(cx))^2}{x^4} dx = \frac{b^2c^2d\sqrt{d - c^2dx^2}}{3x} - \frac{b^2c^3d\sqrt{d - c^2dx^2}\operatorname{arccosh}(cx)}{3\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcd(1 - c^2x^2)\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))}{3x^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{c^2d\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^2}{x} - \frac{4c^3d\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^2}{3\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2dx^2)^{3/2} (a + b\operatorname{arccosh}(cx))^2}{3x^3} - \frac{c^3d\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^3}{3b\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{8bc^3d\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{4b^2c^3d\sqrt{d - c^2dx^2} \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output 
$$-1/3*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^2/x^3+1/3*b^2*c^2*d*(-c^2*d*x^2+d)^{(1/2)}/x+c^2*d*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/x-1/3*b^2*c^3*d*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/3*b*c*d*(-c^2*x^2+1)*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/x^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-4/3*c^3*d*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/3*c^3*d*(a+b*\operatorname{arccosh}(c*x))^3*(-c^2*d*x^2+d)^{(1/2)}/b/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-8/3*b*c^3*d*(a+b*\operatorname{arccosh}(c*x))*\ln(1+1/(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+4/3*b^2*c^3*d*\operatorname{polylog}(2,-1/(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

### 3.185.2 Mathematica [A] (warning: unable to verify)

Time = 1.87 (sec) , antiderivative size = 583, normalized size of antiderivative = 1.37

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^4} dx = \frac{-abcd^2 x + abc^2 d^2 x^2 - a^2 d^2 \sqrt{\frac{-1+cx}{1+cx}} + 5a^2 c^2 d^2 x^2 \sqrt{\frac{-1+cx}{1+cx}} + b^2 c^2 d^2 x^2}{x^4}$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^4,x]`

output 
$$\begin{aligned} & (-a*b*c*d^2*x) + a*b*c^2*d^2*x^2 - a^2*d^2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] + 5 \\ & *a^2*c^2*d^2*x^2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] + b^2*c^2*d^2*x^2*\operatorname{Sqrt}[(-1 + c \\ & *x)/(1 + c*x)] - 4*a^2*c^4*d^2*x^4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] - b^2*c^4*d^2 \\ & *x^4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] - b*d^2*(-1 + c*x)*(-3*a*c^3*x^3 + b*(-\operatorname{S} \\ & \operatorname{rt}[(-1 + c*x)/(1 + c*x)] - c*x*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] + 4*c^2*x^2*\operatorname{S} \\ & \operatorname{qrt}[(-1 + c*x)/(1 + c*x)] + 4*c^3*x^3*(-1 + \operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)])))*\operatorname{Arc} \\ & \operatorname{Cosh}[c*x]^2 + b^2*c^3*d^2*x^3*(-1 + c*x)*\operatorname{ArcCosh}[c*x]^3 - 3*a^2*c^3*d^(3/2) \\ & )*x^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*\operatorname{Sqrt}[d - c^2*d*x^2]*\operatorname{ArcTan}[(c*x*\operatorname{S} \\ & \operatorname{qrt}[d - c^2*d*x^2])]/(\operatorname{Sqrt}[d]*(-1 + c^2*x^2))] + b*d^2*(-1 + c*x)*\operatorname{ArcCosh}[c*x]*(b*c \\ & *x + 2*a*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x - 4*c^2*x^2 - 4*c^3*x^3) + 8* \\ & b*c^3*x^3*\operatorname{Log}[1 + E^(-2*\operatorname{ArcCosh}[c*x])]) - 8*a*b*c^3*d^2*x^3*\operatorname{Log}[c*x] + 8*a \\ & *b*c^4*d^2*x^4*\operatorname{Log}[c*x] - 4*b^2*c^3*d^2*x^3*(-1 + c*x)*\operatorname{PolyLog}[2, -E^(-2*A \\ & \operatorname{rcCosh}[c*x])])]/(3*x^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*\operatorname{Sqrt}[d - c^2*d*x^2]) \end{aligned}$$



**3.185.3 Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 4.24 (sec) , antiderivative size = 403, normalized size of antiderivative = 0.95, number of steps used = 26, number of rules used = 25,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.862$ , Rules used = {6343, 25, 6327, 6335, 108, 27, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838, 6339, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx \\
 & \quad \downarrow \text{6343} \\
 & c^2(-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx - \\
 & \frac{2bcd\sqrt{d - c^2 dx^2} \int -\frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3} \\
 & \quad \downarrow \text{25} \\
 & c^2(-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx + \frac{2bcd\sqrt{d - c^2 dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3} \\
 & \quad \downarrow \text{6327} \\
 & c^2(-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx + \frac{2bcd\sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)(a+\operatorname{barccosh}(cx))}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3} \\
 & \quad \downarrow \text{6335} \\
 & c^2(-d) \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx + \\
 & \frac{2bcd\sqrt{d - c^2 dx^2} \left( c^2 \left( - \int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{1}{2} bc \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2} dx - \frac{(1-c^2 x^2)(a+\operatorname{barccosh}(cx))}{2x^2} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
 & \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3} \\
 & \quad \downarrow \text{108}
 \end{aligned}$$

---

3.185.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$

$$\begin{aligned}
 & \frac{c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + 2bcd\sqrt{d-c^2dx^2} \left( c^2 \left( -\int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{1}{2}bc \left( \int \frac{c^2}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} \right)}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \\
 & \qquad \qquad \qquad \frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & \frac{c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + 2bcd\sqrt{d-c^2dx^2} \left( c^2 \left( -\int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{1}{2}bc \left( c^2 \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} \right)}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \\
 & \qquad \qquad \qquad \frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3} \\
 & \qquad \qquad \qquad \downarrow 43 \\
 & \frac{c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + 2bcd\sqrt{d-c^2dx^2} \left( c^2 \left( -\int \frac{a+\operatorname{barccosh}(cx)}{x} dx \right) - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) \right)}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \\
 & \qquad \qquad \qquad \frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3} \\
 & \qquad \qquad \qquad \downarrow 6297 \\
 & \frac{c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + 2bcd\sqrt{d-c^2dx^2} \left( -\frac{c^2 \int (a+\operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) d(a+\operatorname{barccosh}(cx))}{b} - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) \right)}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \\
 & \qquad \qquad \qquad \frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3} \\
 & \qquad \qquad \qquad \downarrow 25 \\
 & \frac{c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + 2bcd\sqrt{d-c^2dx^2} \left( \frac{c^2 \int (a+\operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) d(a+\operatorname{barccosh}(cx))}{b} - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \left( \operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}\sqrt{cx+1}}{x} \right) \right)}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \\
 & \qquad \qquad \qquad \frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3}
 \end{aligned}$$

3.185.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^4} dx$

$$\begin{aligned}
 & \downarrow \text{3042} \\
 & c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + \\
 & 2bcd\sqrt{d-c^2dx^2} \left( \frac{c^2 \int -i(a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a+\operatorname{barccosh}(cx))}{b} - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \right)
 \end{aligned}$$

$$\frac{3\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \frac{1}{3x^3}$$

$$\begin{aligned}
 & \downarrow \text{26} \\
 & c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + \\
 & 2bcd\sqrt{d-c^2dx^2} \left( -\frac{ic^2 \int (a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a+\operatorname{barccosh}(cx))}{b} - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} - \frac{1}{2}bc \right)
 \end{aligned}$$

$$\frac{3\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \frac{1}{3x^3}$$

$$\begin{aligned}
 & \downarrow \text{4201} \\
 & c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + \\
 & 2bcd\sqrt{d-c^2dx^2} \left( -\frac{ic^2 \left( 2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1+e^{-2\operatorname{arccosh}(cx)}} d(a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2 \right)}{b} - \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{2x^2} \right)
 \end{aligned}$$

$$\frac{3\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \frac{1}{3x^3}$$

$$\begin{aligned}
 & \downarrow \text{2620} \\
 & c^2(-d) \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx + \\
 & 2bcd\sqrt{d-c^2dx^2} \left( -\frac{ic^2 \left( 2i \left( \frac{1}{2}b \int \log(1+e^{-2\operatorname{arccosh}(cx)}) d(a+\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1)(a+\operatorname{barccosh}(cx)) \right) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2 \right)}{b} \right)
 \end{aligned}$$

$$\frac{3\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2} \frac{1}{3x^3}$$

$$\downarrow \text{2715}$$

---

3.185.  $\int \frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^4} dx$

$$2bcd\sqrt{d - c^2dx^2} \left( -\frac{ic^2 \left( 2i \left( -\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1+e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) \right)}{b} \right)$$

---


$$c^2(-d) \int \frac{\sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 2838

$$c^2(-d) \int \frac{\sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx +$$

$$2bcd\sqrt{d - c^2dx^2} \left( -\frac{ic^2 \left( 2i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2 \right)}{b} \right)$$

$3\sqrt{cx - 1}\sqrt{cx + 1}$

$$\frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6339

$$c^2(-d) \left( \frac{c^2\sqrt{d - c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{2bc\sqrt{d - c^2dx^2} \int \frac{a+\operatorname{barccosh}(cx)}{x} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{\sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))}{x} \right)$$

$$2bcd\sqrt{d - c^2dx^2} \left( -\frac{ic^2 \left( 2i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2 \right)}{b} \right)$$

$3\sqrt{cx - 1}\sqrt{cx + 1}$

$$\frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6297

$$c^2(-d) \left( \frac{c^2\sqrt{d - c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{2c\sqrt{d - c^2dx^2} \int -\left( (a + \operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) \right)}{\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

$$2bcd\sqrt{d - c^2dx^2} \left( -\frac{ic^2 \left( 2i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2 \right)}{b} \right)$$

$3\sqrt{cx - 1}\sqrt{cx + 1}$

$$\frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 25

---

3.185.  $\int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$

$$c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) d(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\ 2bcd\sqrt{d-c^2dx^2} \left( -\frac{ic^2\left(2i\left(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1)\right)(a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2\right)}{b} \right)$$

$3\sqrt{cx-1}\sqrt{cx+1}$

$$\frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 3042

$$c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c\sqrt{d-c^2dx^2} \int -i(a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\ 2bcd\sqrt{d-c^2dx^2} \left( -\frac{ic^2\left(2i\left(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1)\right)(a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2\right)}{b} \right)$$

$3\sqrt{cx-1}\sqrt{cx+1}$

$$\frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 26

$$c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{2ic\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\ 2bcd\sqrt{d-c^2dx^2} \left( -\frac{ic^2\left(2i\left(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1)\right)(a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2\right)}{b} \right)$$

$3\sqrt{cx-1}\sqrt{cx+1}$

$$\frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 4201

$$c^2(-d) \left( \frac{c^2\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{2ic\sqrt{d-c^2dx^2} \left(2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1+e^{-2\operatorname{arccosh}(cx)}} d(a+\operatorname{barccosh}(cx))\right)}{\sqrt{cx-1}\sqrt{cx+1}} \right) \\ 2bcd\sqrt{d-c^2dx^2} \left( -\frac{ic^2\left(2i\left(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1)\right)(a+\operatorname{barccosh}(cx)) - \frac{1}{2}i(a+\operatorname{barccosh}(cx))^2\right)}{b} \right)$$

$3\sqrt{cx-1}\sqrt{cx+1}$

$$\frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

---

3.185.  $\int \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x^4} dx$

↓ 2620

$$c^2(-d) \left( \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{2ic\sqrt{d - c^2 dx^2} (2i(\frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)))}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$2bcd\sqrt{d - c^2 dx^2} \left( -\frac{ic^2 (2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2}{b}} \right)$$


---

$$\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 2715

$$c^2(-d) \left( \frac{2ic\sqrt{d - c^2 dx^2} (2i(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)))}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$2bcd\sqrt{d - c^2 dx^2} \left( -\frac{ic^2 (2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2}{b}} \right)$$


---

$$\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 2838

$$c^2(-d) \left( \frac{c^2 \sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{2ic\sqrt{d - c^2 dx^2} (2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)))}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$2bcd\sqrt{d - c^2 dx^2} \left( -\frac{ic^2 (2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2}{b}} \right)$$


---

$$\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6308

$$c^2(-d) \left( \frac{2ic\sqrt{d - c^2 dx^2} (2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx)))}{\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$2bcd\sqrt{d - c^2 dx^2} \left( -\frac{ic^2 (2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2}{b}} \right)$$


---

$$\frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

---

3.185.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^4,x]`

output `-1/3*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/x^3 + (2*b*c*d*Sqrt[d - c^2*d*x^2]*(-1/2*((1 - c^2*x^2)*(a + b*ArcCosh[c*x]))/x^2 - (b*c*(-((Sqrt[-1 + c*x]*Sqrt[1 + c*x])/x) + c*ArcCosh[c*x]))/2 - (I*c^2*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4)))/b))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - c^2*d*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/x) + (c*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^3)/(3*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((2*I)*c*Sqrt[d - c^2*d*x^2]*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]))`

### 3.185.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 108 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(m + 1))), x] - Simp[1/(b*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

---

3.185.  $\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx$

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6308 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/(Sqrt[(d1_)+(e1_)*(x_)]*Sqrt[(d2_)+(e2_)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6327 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*((f_)*(x_))^(m_)*((d1_)+(e1_)*(x_))^(p_)*((d2_)+(e2_)*(x_))^(p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

---

3.185. 
$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx$$



```
rule 6335 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_
)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*(a + b*ArcCosh[c
*x])/(f*(m + 1)), x] + (-Simp[b*c*((-d)^p/(f*(m + 1))) Int[(f*x)^(m + 1)
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x] - Simp[2*e*(p/(f^2*(m + 1
))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x]), x], x]) /
; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0] && ILtQ[(
m + 1)/2, 0]
```

```
rule 6339 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) +
(e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e*
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] - Simp[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 2)*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^
2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]
```

```
rule 6343 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*(a + b*Arc
Cosh[c*x])^n/(f*(m + 1)), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m
+ 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(
m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x],
x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && G
tQ[p, 0] && LtQ[m, -1]
```

### 3.185.4 Maple [A] (verified)

Time = 1.23 (sec) , antiderivative size = 500, normalized size of antiderivative = 1.17

method	result
default	$-\frac{a^2(-c^2dx^2+d)^{\frac{5}{2}}}{3dx^3} + \frac{2a^2c^2(-c^2dx^2+d)^{\frac{5}{2}}}{3dx} + \frac{2a^2c^4x(-c^2dx^2+d)^{\frac{3}{2}}}{3} + a^2c^4dx\sqrt{-c^2dx^2+d} + \frac{a^2c^4d^2\arctan\left(\frac{\sqrt{c^2d}}{\sqrt{-c^2d}}\right)}{\sqrt{c^2d}}$
parts	$-\frac{a^2(-c^2dx^2+d)^{\frac{5}{2}}}{3dx^3} + \frac{2a^2c^2(-c^2dx^2+d)^{\frac{5}{2}}}{3dx} + \frac{2a^2c^4x(-c^2dx^2+d)^{\frac{3}{2}}}{3} + a^2c^4dx\sqrt{-c^2dx^2+d} + \frac{a^2c^4d^2\arctan\left(\frac{\sqrt{c^2d}}{\sqrt{-c^2d}}\right)}{\sqrt{c^2d}}$

```
input int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^4,x,method=_RETURNVERBOSE)
```

$$3.185. \int \frac{(d-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx$$

output `-1/3*a^2/d/x^3*(-c^2*d*x^2+d)^(5/2)+2/3*a^2*c^2/d/x*(-c^2*d*x^2+d)^(5/2)+2/3*a^2*c^4*x*(-c^2*d*x^2+d)^(3/2)+a^2*c^4*d*x*(-c^2*d*x^2+d)^(1/2)+a^2*c^4*d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))-1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^3*(arccosh(c*x)^3*x^3*c^3-4*arccosh(c*x)^2*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^2*c^2-4*arccosh(c*x)^2*x^3*c^3+8*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3+4*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2+c*x*arccosh(c*x))*d-1/3*a*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^3*(3*arccosh(c*x)^2*x^3*c^3-8*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-8*c^3*x^3*arccosh(c*x)+8*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3+2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*d`

### 3.185.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arccosh}(cx) + a)^2}{x^4} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^4,x, algorithm="fricas")`

output `integral(-(a^2*c^2*d*x^2 - a^2*d + (b^2*c^2*d*x^2 - b^2*d)*arccosh(c*x))^2 + 2*(a*b*c^2*d*x^2 - a*b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^4, x)`

### 3.185.6 Sympy [F]

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2}{x^4} dx = \int \frac{(-d(cx - 1)(cx + 1))^{3/2} (a + b \operatorname{acosh}(cx))^2}{x^4} dx$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2/x**4,x)`

output `Integral((-d*(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2/x**4, x)`

**3.185.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{3/2} (b \operatorname{arcosh}(cx) + a)^2}{x^4} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^4,x, algorithm="maxima")`

output `1/3*(3*sqrt(-c^2*d*x^2 + d)*c^4*d*x + 3*c^3*d^(3/2)*arcsin(c*x) + 2*(-c^2*d*x^2 + d)^(3/2)*c^2/x - (-c^2*d*x^2 + d)^(5/2)/(d*x^3))*a^2 + integrate((-c^2*d*x^2 + d)^(3/2)*b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/x^4 + 2*(-c^2*d*x^2 + d)^(3/2)*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/x^4, x)`

**3.185.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/x^4,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command: INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.185.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2}}{x^4} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x^4,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2))/x^4, x)`

---

3.185.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$

### 3.186 $\int x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.186.1 Optimal result	.1571
3.186.2 Mathematica [A] (verified)	1572
3.186.3 Rubi [F]	1573
3.186.4 Maple [B] (verified)	1586
3.186.5 Fricas [A] (verification not implemented)	1587
3.186.6 Sympy [F(-1)]	1588
3.186.7 Maxima [A] (verification not implemented)	1589
3.186.8 Giac [F(-2)]	1590
3.186.9 Mupad [F(-1)]	1590

#### 3.186.1 Optimal result

Integrand size = 29, antiderivative size = 880

$$\begin{aligned}
 \int x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = & -\frac{37384b^2 d^2 \sqrt{d - c^2 dx^2}}{694575c^4} \\
 & + \frac{3358b^2 d^2 x^2 \sqrt{d - c^2 dx^2}}{694575c^2} + \frac{484b^2 d^2 x^4 \sqrt{d - c^2 dx^2}}{77175} - \frac{10b^2 c^2 d^2 x^6 \sqrt{d - c^2 dx^2}}{3087} \\
 & + \frac{4abd^2 x \sqrt{d - c^2 dx^2}}{63c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{16b^2 d^2 (1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{2835c^4 (1 - cx)(1 + cx)} \\
 & + \frac{8b^2 d^2 (1 - c^2 x^2)^2 \sqrt{d - c^2 dx^2}}{8505c^4 (1 - cx)(1 + cx)} + \frac{2b^2 d^2 (1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{4725c^4 (1 - cx)(1 + cx)} \\
 & - \frac{20b^2 d^2 (1 - c^2 x^2)^4 \sqrt{d - c^2 dx^2}}{3969c^4 (1 - cx)(1 + cx)} + \frac{2b^2 d^2 (1 - c^2 x^2)^5 \sqrt{d - c^2 dx^2}}{729c^4 (1 - cx)(1 + cx)} \\
 & + \frac{4b^2 d^2 x \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{63c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{2bd^2 x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{189c \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{2bcd^2 x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{21 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & + \frac{38bc^3 d^2 x^7 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{441 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{2bc^5 d^2 x^9 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{81 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{2d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{63c^4} - \frac{d^2 x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{63c^2} \\
 & + \frac{1}{21} d^2 x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 + \frac{5}{63} dx^4 (d \\
 & - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 + \frac{1}{9} x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2
 \end{aligned}$$

output  $5/63*d*x^4*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^{2+1/9*x^4*(-c^2*d*x^2+d)^{(5/2)}*(a+b*\operatorname{arccosh}(c*x))^{2-37384/694575*b^2*d^2*(-c^2*d*x^2+d)^{(1/2)}/c^4+3358/694575*b^2*d^2*x^2*(-c^2*d*x^2+d)^{(1/2)}/c^2+484/77175*b^2*d^2*x^4*(-c^2*d*x^2+d)^{(1/2)}-10/3087*b^2*c^2*d^2*x^6*(-c^2*d*x^2+d)^{(1/2)}+16/2835*b^2*d^2*(-c^2*x^2+1)*(-c^2*d*x^2+d)^{(1/2)}/c^4/(-c*x+1)/(c*x+1)+8/8505*b^2*d^2*(-c^2*x^2+1)^2*(-c^2*d*x^2+d)^{(1/2)}/c^4/(-c*x+1)/(c*x+1)+2/4725*b^2*d^2*(-c^2*x^2+1)^3*(-c^2*d*x^2+d)^{(1/2)}/c^4/(-c*x+1)/(c*x+1)-20/3969*b^2*d^2*(-c^2*x^2+1)^4*(-c^2*d*x^2+d)^{(1/2)}/c^4/(-c*x+1)/(c*x+1)+2/729*b^2*d^2*(-c^2*x^2+1)^5*(-c^2*d*x^2+d)^{(1/2)}/c^4/(-c*x+1)/(c*x+1)-2/63*d^2*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d*x^2+d)^{(1/2)}/c^2+1/21*d^2*x^4*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d*x^2+d)^{(1/2)}+4/63*a*b*d^2*x*(-c^2*d*x^2+d)^{(1/2)}/c^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+4/63*b^2*d^2*x*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d)^{(1/2)}/c^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+2/189*b*d^2*x^3*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2/21*b*c*d^2*x^5*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+38/441*b*c^3*d^2*x^7*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2/81*b*c^5*d^2*x^9*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.186.2 Mathematica [A] (verified)

Time = 0.60 (sec) , antiderivative size = 288, normalized size of antiderivative = 0.33

$$\int x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{d^2 \sqrt{d - c^2 dx^2} \left( 3969a^2(-1 + c^2 x^2)^4 (2 + 7c^2 x^2) - 126abcx \sqrt{-1 + cx} \sqrt{1 + cx} (-126 - \dots \right)}{\dots}$$

input `Integrate[x^3*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output  $(d^2*\operatorname{Sqrt}[d - c^2*d*x^2]*(3969*a^2*(-1 + c^2*x^2)^4*(2 + 7*c^2*x^2) - 126*a*b*c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(-126 - 21*c^2*x^2 + 189*c^4*x^4 - 171*c^6*x^6 + 49*c^8*x^8) + 2*b^2*(6140 - 7039*c^2*x^2 - 106*c^4*x^4 + 2152*c^6*x^6 - 1490*c^8*x^8 + 343*c^{10}*x^{10}) + 126*b*(63*a*(-1 + c^2*x^2)^4*(2 + 7*c^2*x^2) + b*c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(126 + 21*c^2*x^2 - 189*c^4*x^4 + 171*c^6*x^6 - 49*c^8*x^8))*\operatorname{ArcCosh}[c*x] + 3969*b^2*(-1 + c^2*x^2)^4*(2 + 7*c^2*x^2)*\operatorname{ArcCosh}[c*x]^2)/(250047*c^4*(-1 + c^2*x^2))$

**3.186.3 Rubi [F]**

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx \\
 & \quad \downarrow \text{6345} \\
 & \frac{2bcd^2\sqrt{d - c^2 dx^2} \int x^4(1 - cx)^2(cx + 1)^2(a + \operatorname{barccosh}(cx))dx}{9\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{5}{9}d \int x^3(d - c^2 dx^2)^{3/2} (a + \\
 & \quad \operatorname{barccosh}(cx))^2 dx + \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{6327} \\
 & \frac{2bcd^2\sqrt{d - c^2 dx^2} \int x^4(1 - c^2 x^2)^2(a + \operatorname{barccosh}(cx))dx}{9\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{5}{9}d \int x^3(d - c^2 dx^2)^{3/2} (a + \\
 & \quad \operatorname{barccosh}(cx))^2 dx + \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{6336} \\
 & \frac{\frac{5}{9}d \int x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -}{2bcd^2\sqrt{d - c^2 dx^2} \left( -bc \int \frac{x^5(35c^4 x^4 - 90c^2 x^2 + 63)}{315\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{9}c^4 x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \right.} \\
 & \quad \left. \operatorname{barccosh}(cx))^2 dx - \right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \\
 & \quad \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{27} \\
 & \frac{\frac{5}{9}d \int x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -}{2bcd^2\sqrt{d - c^2 dx^2} \left( -\frac{1}{315}bc \int \frac{x^5(35c^4 x^4 - 90c^2 x^2 + 63)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{9}c^4 x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \right.} \\
 & \quad \left. \operatorname{barccosh}(cx))^2 dx - \right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \\
 & \quad \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{1905} \\
 & \frac{\frac{5}{9}d \int x^3(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -}{2bcd^2\sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \int \frac{x^5(35c^4 x^4 - 90c^2 x^2 + 63)}{315\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{9}c^4 x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \right.} \\
 & \quad \left. \operatorname{barccosh}(cx))^2 dx - \right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \\
 & \quad \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2
 \end{aligned}$$

---

3.186.  $\int x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned} & \downarrow 1578 \\ & \frac{5}{9}d \int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx - \\ & 2bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \int \frac{x^4 (35c^4 x^4 - 90c^2 x^2 + 63)}{\sqrt{c^2 x^2 - 1}} dx^2}{630\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{9}c^4 x^9 (a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x \right) \end{aligned}$$

$$\begin{aligned} & \frac{9\sqrt{cx-1}\sqrt{cx+1}}{\frac{1}{9}x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} \\ & \downarrow 1195 \\ & \frac{5}{9}d \int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx - \\ & 2bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \int \left( \frac{35(c^2 x^2 - 1)^{7/2}}{c^4} + \frac{50(c^2 x^2 - 1)^{5/2}}{c^4} + \frac{3(c^2 x^2 - 1)^{3/2}}{c^4} - \frac{4\sqrt{c^2 x^2 - 1}}{c^4} + \frac{8}{c^4 \sqrt{c^2 x^2 - 1}} \right) dx^2}{630\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{9}c^4 x^9 (a + \operatorname{barccosh}(cx)) \right) \end{aligned}$$

$$\begin{aligned} & \frac{9\sqrt{cx-1}\sqrt{cx+1}}{\frac{1}{9}x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} \\ & \downarrow 2009 \end{aligned}$$

$$\begin{aligned} & \frac{5}{9}d \int x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{9}x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \\ & 2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4 x^9 (a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{70}{c^4} \right)}{\dots} \right) \end{aligned}$$

$$\begin{aligned} & \frac{9\sqrt{cx-1}\sqrt{cx+1}}{\dots} \\ & \downarrow 6345 \end{aligned}$$

$$\begin{aligned} & \frac{5}{9}d \left( \frac{2bcd\sqrt{d - c^2 dx^2} \int -x^4 (1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx)) dx}{7\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \right. \\ & \left. \frac{1}{9}x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \right. \\ & \left. 2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4 x^9 (a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{70}{c^4} \right)}{\dots} \right) \right) \end{aligned}$$

$$\begin{aligned} & \frac{9\sqrt{cx-1}\sqrt{cx+1}}{\dots} \\ & \downarrow 25 \end{aligned}$$

$$\frac{5}{9}d \left( -\frac{2bcd\sqrt{d-c^2dx^2} \int x^4(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{7\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{7}d \int x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 dx + \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \right.$$

$$\left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{70} \right) \right)$$


---


$$9\sqrt{cx-1}\sqrt{cx+1}$$

↓ 6327

$$\frac{5}{9}d \left( -\frac{2bcd\sqrt{d-c^2dx^2} \int x^4(1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{7\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{7}d \int x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 dx + \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \right.$$

$$\left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{70} \right) \right)$$


---


$$9\sqrt{cx-1}\sqrt{cx+1}$$

↓ 6336

$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d-c^2dx^2} \left( -bc \int \frac{x^5(7-5c^2x^2)}{35\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^2x^7(a+\operatorname{barccosh}(cx)) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \right.$$

$$\left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{70} \right) \right)$$


---


$$9\sqrt{cx-1}\sqrt{cx+1}$$

↓ 27



$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \int \frac{x^5(7-5c^2x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^2x^7(a + \operatorname{barccosh}(cx)) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) \right) - \frac{bc\sqrt{c^2x^2-1} \left( \frac{70}{9} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)}{9\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 960

$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \int \frac{x^5}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{5}{7}x^6\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) \right) - \frac{bc\sqrt{c^2x^2-1} \left( \frac{70}{9} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)}{9\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 111

$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right) \right)}{7\sqrt{cx-1}\sqrt{cx+1}} \right. \\ \left. - \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) \right) - \frac{bc\sqrt{c^2x^2-1} \left( \frac{70}{9} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)}{9\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 27

$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4 \sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$

$$\left. \frac{1}{9}x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4 x^9 (a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{70}{9} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 111

$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \frac{4 \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{5c^2} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$

$$\left. \frac{1}{9}x^4 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4 x^9 (a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7 (a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{70}{9} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 27

$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{35}bc \left( \frac{19}{7} \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx + x^2 \sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{\right)}{\right)} \right.$$

$$\left. - \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4 x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{70}{9} \right)}{\right)}{\right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 83

$$\frac{5}{9}d \left( \frac{3}{7}d \int x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{7}x^4(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7} \right)}{\right)} \right.$$

$$\left. - \frac{1}{9}x^4(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4 x^9(a + \operatorname{barccosh}(cx)) - \frac{2}{7}c^2 x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{70}{9} \right)}{\right)}{\right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 6341

$$\frac{5}{9}d \left( \frac{3}{7}d \left( -\frac{2bc\sqrt{d-c^2dx^2} \int x^4(a+\operatorname{barccosh}(cx))dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}}dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \right. \right. \\ \left. \left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{9\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{70}{9} \right) \right) \right) \right)$$

↓ 6298

$$\frac{5}{9}d \left( \frac{3}{7}d \left( -\frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \int \frac{x^5}{\sqrt{cx-1}\sqrt{cx+1}}dx \right)}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}}dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \right. \right. \\ \left. \left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{9\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{70}{9} \right) \right) \right) \right)$$

↓ 111

$$\frac{5}{9}d \left( \frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}}dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{\int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}}dx}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \right. \right. \\ \left. \left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{9\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{70}{9} \right) \right) \right) \right)$$

↓ 27

$$\frac{5}{9}d \left( \frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\ \left. \left. + \frac{1}{9}x^4(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{70} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$

↓ 111

$$\frac{5}{9}d \left( \frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} \frac{dx}{3c^2} \right) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right) \right. \right. \\ \left. \left. + \frac{1}{9}x^4(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{70} \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$

↓ 27

$$\frac{5}{9}d \left( \frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}}}{3c^2} \right)}{bc\sqrt{c^2x^2-1}} \left( \frac{70}{9} \right) \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \right. \\ \left. \left. \left. \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{9\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{70}{9} \right) \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right) \right) \right.$$

↓ 83

$$\frac{5}{9}d \left( \frac{3}{7}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}}}{3c^2} \right)}{bc\sqrt{c^2x^2-1}} \left( \frac{70}{9} \right) \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \right. \\ \left. \left. \left. \frac{1}{9}x^4(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{9}c^4x^9(a+\operatorname{barccosh}(cx)) - \frac{2}{7}c^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{9\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{70}{9} \right) \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right) \right) \right.$$

↓ 6354

$$\frac{5}{9}d \left( \frac{1}{7}(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2x^4 - \frac{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^2(a+\operatorname{barccosh}(cx))x^7 + \frac{1}{5}(a+\operatorname{barccosh}(cx))x^5 - \frac{bc\sqrt{c^2x^2-1}}{9\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{70}{9} \right) \right)}{9\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 6298

$$\frac{1}{9}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 x^4 -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{9} c^4 (a + \operatorname{barccosh}(cx)) x^9 - \frac{2}{7} c^2 (a + \operatorname{barccosh}(cx)) x^7 + \frac{1}{5} (a + \operatorname{barccosh}(cx)) x^5 - \frac{bc\sqrt{c^2 x^2 - 1}}{\left(\frac{70}{9}\right)} \right)$$


---

$$\frac{5}{9} d \left( \frac{1}{7} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 x^4 - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7} c^2 (a + \operatorname{barccosh}(cx)) x^7 + \frac{1}{5} (a + \operatorname{barccosh}(cx)) x^5 \right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 111

$$\frac{1}{9}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 x^4 -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{9} c^4 (a + \operatorname{barccosh}(cx)) x^9 - \frac{2}{7} c^2 (a + \operatorname{barccosh}(cx)) x^7 + \frac{1}{5} (a + \operatorname{barccosh}(cx)) x^5 - \frac{bc\sqrt{c^2 x^2 - 1}}{\left(\frac{70}{9}\right)} \right)$$


---

$$\frac{5}{9} d \left( \frac{1}{7} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 x^4 - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7} c^2 (a + \operatorname{barccosh}(cx)) x^7 + \frac{1}{5} (a + \operatorname{barccosh}(cx)) x^5 \right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

↓ 27

$$\frac{\frac{1}{9}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 x^4 - 2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{9}c^4(a + \operatorname{barccosh}(cx))x^9 - \frac{2}{7}c^2(a + \operatorname{barccosh}(cx))x^7 + \frac{1}{5}(a + \operatorname{barccosh}(cx))x^5 - \frac{bc\sqrt{c^2 x^2 - 1}}{70} \right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{5d}{9} \left( \frac{1}{7}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 x^4 - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{7}c^2(a + \operatorname{barccosh}(cx))x^7 + \frac{1}{5}(a + \operatorname{barccosh}(cx))x^5 \right)}{9\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

input `Int[x^3*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

### 3.186.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 111 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`



- rule 960  $\text{Int}[(e_*)(x_)^{(m_*)}((a1_)+(b1_*)(x_)^{(non2_)})^{(p_*)}((a2_)+(b2_*)\cdot(x_)^{(non2_)})^{(p_*)}((c_)+(d_*)(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}*(a1+b1*x^{(n/2)})^{(p+1)}*((a2+b2*x^{(n/2)})^{(p+1)})/(b1*b2*e*(m+n*(p+1)+1)), x] - \text{Simp}[(a1*a2*d*(m+1)-b1*b2*c*(m+n*(p+1)+1))/(b1*b2*(m+n*(p+1)+1)) \text{Int}[(e*x)^m*(a1+b1*x^{(n/2)})^p*(a2+b2*x^{(n/2)})^p, x], x] /; \text{FreeQ}[\{a1, b1, a2, b2, c, d, e, m, n, p\}, x] \&\& \text{EqQ}[\text{non2}, n/2] \&\& \text{EqQ}[a2*b1+a1*b2, 0] \&\& \text{NeQ}[m+n*(p+1)+1, 0]$
- rule 1195  $\text{Int}[(d_)+(e_*)(x_)^{(m_*)}((f_)+(g_*)(x_)^{(n_*)}((a_)+(b_*)(x_)+(c_)*(x_)^2)^{(p_)}), x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(d+e*x)^m*(f+g*x)^n*(a+b*x+c*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, m, n\}, x] \&\& \text{IGtQ}[p, 0]$
- rule 1578  $\text{Int}[(x_)^{(m_*)}((d_)+(e_*)(x_)^2)^{(q_*)}((a_)+(b_*)(x_)^2+(c_)*(x_)^4)^{(p_)}), x\_Symbol] \rightarrow \text{Simp}[1/2 \text{Subst}[\text{Int}[x^{(m-1)/2}*(d+e*x)^q*(a+b*x+c*x^2)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, c, d, e, p, q\}, x] \&\& \text{IntegerQ}[(m-1)/2]$
- rule 1905  $\text{Int}[(f_*)(x_)^{(m_*)}((d1_)+(e1_*)(x_)^{(non2_)})^{(q_*)}((d2_)+(e2_*)\cdot(x_)^{(non2_)})^{(q_*)}((a_)+(b_*)(x_)^{(n_)}+(c_)*(x_)^{(n2_)}), x\_Symbol] \rightarrow \text{Simp}[(d1+e1*x^{(n/2)})^{\text{FracPart}[q]}*((d2+e2*x^{(n/2)})^{\text{FracPart}[q]})/((d1*d2+e1*e2*x^n)^{\text{FracPart}[q]}) \text{Int}[(f*x)^m*(d1*d2+e1*e2*x^n)^q*(a+b*x^n+c*x^{(2*n)})^p, x], x] /; \text{FreeQ}[\{a, b, c, d1, e1, d2, e2, f, n, p, q\}, x] \&\& \text{EqQ}[n2, 2*n] \&\& \text{EqQ}[\text{non2}, n/2] \&\& \text{EqQ}[d2*e1+d1*e2, 0]$
- rule 2009  $\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$
- rule 6298  $\text{Int}[(a_)+(b_*)\cdot \text{ArcCosh}[(c_*)(x_)]*(d_*)(x_)^{(m_*)}((a_)+(b_*)(x_)^{(n_)}), x\_Symbol] \rightarrow \text{Simp}[(d*x)^{(m+1)}*((a+b*\text{ArcCosh}[c*x])^n/(d*(m+1))), x] - \text{Simp}[b*c*(n/(d*(m+1))) \text{Int}[(d*x)^{(m+1)}*((a+b*\text{ArcCosh}[c*x])^{(n-1)})/(\text{Sqrt}[1+c*x]*\text{Sqrt}[-1+c*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{NeQ}[m, -1]$

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x])^n * u, x] - Simp[b*c * Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x] - Simp[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) * Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)], x] - Simp[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^n, x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.186.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2223 vs.  $2(776) = 1552$ .

Time = 0.78 (sec) , antiderivative size = 2224, normalized size of antiderivative = 2.53

method	result	size
default	Expression too large to display	2224
parts	Expression too large to display	2224

```
input int(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

output

```

a^2*(-1/9*x^2*(-c^2*d*x^2+d)^(7/2)/c^2/d-2/63/d/c^4*(-c^2*d*x^2+d)^(7/2))+
b^2*(1/373248*(-d*(c^2*x^2-1))^(1/2)*(256*c^10*x^10-704*c^8*x^8+256*(c*x+1)
)^(1/2)*(c*x-1)^(1/2)*x^9*c^9+688*c^6*x^6-576*(c*x+1)^(1/2)*(c*x-1)^(1/2)*
x^7*c^7-280*c^4*x^4+432*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+41*c^2*x^2-120
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+9*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*
(81*arccosh(c*x)^2-18*arccosh(c*x)+2)*d^2/(c*x+1)/c^4/(c*x-1)-3/175616*(-d
*(c^2*x^2-1))^(1/2)*(64*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(c*x-1)^(1/2)
*x^7*c^7+104*c^4*x^4-112*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-25*c^2*x^2+56
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*
(49*arccosh(c*x)^2-14*arccosh(c*x)+2)*d^2/(c*x+1)/c^4/(c*x-1)+1/1728*(-d*(
c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x
^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(9*arccosh(c*x)^2-6*arccosh(c*x)+2
)*d^2/(c*x+1)/c^4/(c*x-1)-3/256*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x
+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*arccosh(c*x)+2)*d^2/(c*x+1)/c^4
/(c*x-1)-3/256*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^
2*x^2-1)*(arccosh(c*x)^2+2*arccosh(c*x)+2)*d^2/(c*x+1)/c^4/(c*x-1)+1/1728*
(-d*(c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(9*arccosh(c*x)^2+6*arccosh(
c*x)+2)*d^2/(c*x+1)/c^4/(c*x-1)-3/175616*(-d*(c^2*x^2-1))^(1/2)*(-64*(c*x+
1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+64*c^8*x^8+112*(c*x+1)^(1/2)*(c*x-1)^(1/2)

```

### 3.186.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 558, normalized size of antiderivative = 0.63

$$\int x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{3969(7b^2c^{10}d^2x^{10} - 26b^2c^8d^2x^8 + 34b^2c^6d^2x^6 - 16b^2c^4d^2x^4 - b^2c^2d^2x^2 + 2b^2d^2)\sqrt{-c}}{\dots}$$

input `integrate(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

```
output 1/250047*(3969*(7*b^2*c^10*d^2*x^10 - 26*b^2*c^8*d^2*x^8 + 34*b^2*c^6*d^2*x^6 - 16*b^2*c^4*d^2*x^4 - b^2*c^2*d^2*x^2 + 2*b^2*d^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1))^2 - 126*(49*a*b*c^9*d^2*x^9 - 171*a*b*c^7*d^2*x^7 + 189*a*b*c^5*d^2*x^5 - 21*a*b*c^3*d^2*x^3 - 126*a*b*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 126*((49*b^2*c^9*d^2*x^9 - 171*b^2*c^7*d^2*x^7 + 189*b^2*c^5*d^2*x^5 - 21*b^2*c^3*d^2*x^3 - 126*b^2*c*d^2*x)*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1) - 63*(7*a*b*c^10*d^2*x^10 - 26*a*b*c^8*d^2*x^8 + 34*a*b*c^6*d^2*x^6 - 16*a*b*c^4*d^2*x^4 - a*b*c^2*d^2*x^2 + 2*a*b*d^2)*sqrt(-c^2*d*x^2 + d))*log(c*x + sqrt(c^2*x^2 - 1)) + (343*(81*a^2 + 2*b^2)*c^10*d^2*x^10 - 2*(51597*a^2 + 1490*b^2)*c^8*d^2*x^8 + 2*(67473*a^2 + 2152*b^2)*c^6*d^2*x^6 - 4*(15876*a^2 + 53*b^2)*c^4*d^2*x^4 - (3969*a^2 + 14078*b^2)*c^2*d^2*x^2 + 2*(3969*a^2 + 6140*b^2)*d^2)*sqrt(-c^2*d*x^2 + d))/(c^6*x^2 - c^4)
```

### 3.186.6 Sympy [F(-1)]

Timed out.

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Timed out}$$

```
input integrate(x**3*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2,x)
```

```
output Timed out
```

**3.186.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 471, normalized size of antiderivative = 0.54

$$\begin{aligned}
& \int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \\
& -\frac{1}{63} \left( \frac{7(-c^2 dx^2 + d)^{7/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{7/2}}{c^4 d} \right) b^2 \operatorname{arcosh}(cx)^2 \\
& -\frac{2}{63} \left( \frac{7(-c^2 dx^2 + d)^{7/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{7/2}}{c^4 d} \right) ab \operatorname{arcosh}(cx) \\
& -\frac{1}{63} \left( \frac{7(-c^2 dx^2 + d)^{7/2} x^2}{c^2 d} + \frac{2(-c^2 dx^2 + d)^{7/2}}{c^4 d} \right) a^2 \\
& + \frac{2}{250047} b^2 \left( \frac{343 \sqrt{c^2 x^2 - 1} c^6 \sqrt{-dd^2} x^8 - 1147 \sqrt{c^2 x^2 - 1} c^4 \sqrt{-dd^2} x^6 + 1005 \sqrt{c^2 x^2 - 1} c^2 \sqrt{-dd^2} x^4 + 899 \sqrt{c^2 x^2 - 1} \sqrt{-dd^2}}{c^2} \right. \\
& \left. - \frac{2(49 c^8 \sqrt{-dd^2} x^9 - 171 c^6 \sqrt{-dd^2} x^7 + 189 c^4 \sqrt{-dd^2} x^5 - 21 c^2 \sqrt{-dd^2} x^3 - 126 \sqrt{-dd^2} x) ab}{3969 c^3} \right)
\end{aligned}$$

```
input integrate(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")
```

```
output -1/63*(7*(-c^2*d*x^2 + d)^(7/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(7/2)/(c^4*d))*b^2*arccosh(c*x)^2 - 2/63*(7*(-c^2*d*x^2 + d)^(7/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(7/2)/(c^4*d))*a*b*arccosh(c*x) - 1/63*(7*(-c^2*d*x^2 + d)^(7/2)*x^2/(c^2*d) + 2*(-c^2*d*x^2 + d)^(7/2)/(c^4*d))*a^2 + 2/250047*b^2*((343*sqrt(c^2*x^2 - 1)*c^6*sqrt(-d)*d^2*x^8 - 1147*sqrt(c^2*x^2 - 1)*c^4*sqrt(-d)*d^2*x^6 + 1005*sqrt(c^2*x^2 - 1)*c^2*sqrt(-d)*d^2*x^4 + 899*sqrt(c^2*x^2 - 1)*sqrt(-d)*d^2*x^2 - 6140*sqrt(c^2*x^2 - 1)*sqrt(-d)*d^2/c^2)/c^2 - 63*(49*c^8*sqrt(-d)*d^2*x^9 - 171*c^6*sqrt(-d)*d^2*x^7 + 189*c^4*sqrt(-d)*d^2*x^5 - 21*c^2*sqrt(-d)*d^2*x^3 - 126*sqrt(-d)*d^2*x)*arccosh(c*x)/c^3 - 2/3969*(49*c^8*sqrt(-d)*d^2*x^9 - 171*c^6*sqrt(-d)*d^2*x^7 + 189*c^4*sqrt(-d)*d^2*x^5 - 21*c^2*sqrt(-d)*d^2*x^3 - 126*sqrt(-d)*d^2*x)*a*b/c^3
```

**3.186.8 Giac [F(-2)]**

Exception generated.

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.186.9 Mupad [F(-1)]**

Timed out.

$$\int x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int x^3 (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2} dx$$

input `int(x^3*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2),x)`

output `int(x^3*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2), x)`

### 3.187 $\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.187.1 Optimal result . . . . .	1591
3.187.2 Mathematica [A] (warning: unable to verify) . . . . .	1592
3.187.3 Rubi [F] . . . . .	1593
3.187.4 Maple [B] (verified) . . . . .	1609
3.187.5 Fricas [F] . . . . .	1610
3.187.6 Sympy [F(-1)] . . . . .	1611
3.187.7 Maxima [F] . . . . .	1611
3.187.8 Giac [F] . . . . .	1611
3.187.9 Mupad [F(-1)] . . . . .	1612

#### 3.187.1 Optimal result

Integrand size = 29, antiderivative size = 841

$$\begin{aligned}
 \int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = & \frac{35b^2 d^2 x \sqrt{d - c^2 dx^2}}{9216c^2} + \frac{215b^2 d^2 x^3 \sqrt{d - c^2 dx^2}}{13824} \\
 & - \frac{5}{864} b^2 c^2 d^2 x^5 \sqrt{d - c^2 dx^2} + \frac{73b^2 d^2 x(1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{12288c^2(1 - cx)(1 + cx)} + \frac{73b^2 d^2 x^3(1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{18432(1 - cx)(1 + cx)} \\
 & - \frac{43b^2 c^2 d^2 x^5(1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{4608(1 - cx)(1 + cx)} + \frac{b^2 c^4 d^2 x^7(1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{256(1 - cx)(1 + cx)} \\
 & + \frac{35b^2 d^2 \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{9216c^3 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{5bd^2 x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{128c \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{59bcd^2 x^4 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{384 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{17bc^3 d^2 x^6 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{144 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{bc^5 d^2 x^8 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{32 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{5d^2 x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{128c^2} \\
 & + \frac{5}{64} d^2 x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 + \frac{5}{48} dx^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\
 & + \frac{1}{8} x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{5d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^3}{384bc^3 \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{73b^2 d^2 \sqrt{-1 + c^2 x^2} \sqrt{d - c^2 dx^2}}{12288c^3(1 - cx)}
 \end{aligned}$$



output

```

5/48*d*x^3*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2+1/8*x^3*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2+35/9216*b^2*d^2*x*(-c^2*d*x^2+d)^(1/2)/c^2+215/13824*b^2*d^2*x^3*(-c^2*d*x^2+d)^(1/2)-5/864*b^2*c^2*d^2*x^5*(-c^2*d*x^2+d)^(1/2)+73/12288*b^2*d^2*x*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)+73/18432*b^2*d^2*x^3*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/(-c*x+1)/(c*x+1)-43/4608*b^2*c^2*d^2*x^5*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/(-c*x+1)/(c*x+1)+1/256*b^2*c^4*d^2*x^7*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/(-c*x+1)/(c*x+1)-5/128*d^2*x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/c^2+5/64*d^2*x^3*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)+35/9216*b^2*d^2*arccosh(c*x)*(-c^2*d*x^2+d)^(1/2)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/128*b*d^2*x^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-59/384*b*c*d^2*x^4*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+17/144*b*c^3*d^2*x^6*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/32*b*c^5*d^2*x^8*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-5/384*d^2*(a+b*arccosh(c*x))^3*(-c^2*d*x^2+d)^(1/2)/b/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)-73/12288*b^2*d^2*arctanh(c*x)/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)*(-c^2*d*x^2+d)^(1/2)/c^3/(-c*x+1)/(c*x+1)

```

### 3.187.2 Mathematica [A] (warning: unable to verify)

Time = 5.92 (sec) , antiderivative size = 910, normalized size of antiderivative = 1.08

$$\int x^2 (d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2 dx =$$

$$\frac{d^2 \left( 34560 a^2 cx \sqrt{\frac{-1+cx}{1+cx}} \sqrt{d - c^2 dx^2} + 34560 a^2 c^2 x^2 \sqrt{\frac{-1+cx}{1+cx}} \sqrt{d - c^2 dx^2} - 271872 a^2 c^3 x^3 \sqrt{\frac{-1+cx}{1+cx}} \sqrt{d - c^2 dx^2} \right)}{c^3}$$

input `Integrate[x^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output

```

-1/884736*(d^2*(34560*a^2*c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^
2] + 34560*a^2*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] - 27
1872*a^2*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] - 271872*a
^2*c^4*x^4*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] + 313344*a^2*c^5
*x^5*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] + 313344*a^2*c^6*x^6*S
qrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] - 110592*a^2*c^7*x^7*Sqrt[(-
1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] - 110592*a^2*c^8*x^8*Sqrt[(-1 + c*
x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] + 11520*b^2*Sqrt[d - c^2*d*x^2]*ArcCosh[
c*x]^3 + 34560*a^2*Sqrt[d]*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcTan[(c*x*Sqrt[d -
c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + 34560*a^2*c*Sqrt[d]*x*Sqrt[(-1 +
c*x)/(1 + c*x)]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))]
+ 13824*a*b*Sqrt[d - c^2*d*x^2]*Cosh[2*ArcCosh[c*x]] + 3456*a*b*Sqrt[d -
c^2*d*x^2]*Cosh[4*ArcCosh[c*x]] - 1536*a*b*Sqrt[d - c^2*d*x^2]*Cosh[6*ArcC
osh[c*x]] + 216*a*b*Sqrt[d - c^2*d*x^2]*Cosh[8*ArcCosh[c*x]] - 6912*b^2*Sq
rt[d - c^2*d*x^2]*Sinh[2*ArcCosh[c*x]] - 864*b^2*Sqrt[d - c^2*d*x^2]*Sinh[
4*ArcCosh[c*x]] + 256*b^2*Sqrt[d - c^2*d*x^2]*Sinh[6*ArcCosh[c*x]] - 27*b^
2*Sqrt[d - c^2*d*x^2]*Sinh[8*ArcCosh[c*x]] + 24*b*Sqrt[d - c^2*d*x^2]*ArcC
osh[c*x]*(576*b*Cosh[2*ArcCosh[c*x]] + 144*b*Cosh[4*ArcCosh[c*x]] - 64*b*C
osh[6*ArcCosh[c*x]] + 9*b*Cosh[8*ArcCosh[c*x]] - 1152*a*Sinh[2*ArcCosh[c*x
]] - 576*a*Sinh[4*ArcCosh[c*x]] + 384*a*Sinh[6*ArcCosh[c*x]] - 72*a*Sin...

```

### 3.187.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx \\
 & \quad \downarrow \text{6345} \\
 & -\frac{bcd^2\sqrt{d - c^2 dx^2} \int x^3(1 - cx)^2(cx + 1)^2(a + \operatorname{barccosh}(cx))dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{5}{8}d \int x^2(d - c^2 dx^2)^{3/2} (a + \\
 & \quad \operatorname{barccosh}(cx))^2 dx + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{6327} \\
 & -\frac{bcd^2\sqrt{d - c^2 dx^2} \int x^3(1 - c^2 x^2)^2(a + \operatorname{barccosh}(cx))dx}{4\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{5}{8}d \int x^2(d - c^2 dx^2)^{3/2} (a + \\
 & \quad \operatorname{barccosh}(cx))^2 dx + \frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \quad \downarrow \text{6336}
 \end{aligned}$$

---

3.187.  $\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
 & \frac{\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx - bcd^2 \sqrt{d - c^2 dx^2} \left( -bc \int \frac{x^4 (3c^4 x^4 - 8c^2 x^2 + 6)}{24\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \\
 & \qquad \frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \qquad \qquad \qquad \downarrow \text{27} \\
 & \frac{\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx - bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{1}{24}bc \int \frac{x^4 (3c^4 x^4 - 8c^2 x^2 + 6)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \\
 & \qquad \frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \qquad \qquad \qquad \downarrow \text{1905} \\
 & \frac{\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx - bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \int \frac{x^4 (3c^4 x^4 - 8c^2 x^2 + 6)}{24\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{c^2 x^2 - 1}} + \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \\
 & \qquad \frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \qquad \qquad \qquad \downarrow \text{1590} \\
 & \frac{\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx - bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \left( \int \frac{c^2 x^4 (48 - 43c^2 x^2)}{8c^2} dx + \frac{3}{8}c^2 x^7 \sqrt{c^2 x^2 - 1} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \\
 & \qquad \frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \\
 & \qquad \qquad \qquad \downarrow \text{27}
 \end{aligned}$$

---

3.187.  $\int x^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \int \frac{x^4 (48 - 43c^2 x^2)}{\sqrt{c^2 x^2 - 1}} dx + \frac{3}{8} c^2 x^7 \sqrt{c^2 x^2 - 1} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{8} c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} c^2 x^6 (a + \operatorname{barccosh}(cx))$$

---


$$4\sqrt{cx-1}\sqrt{cx+1}$$

$$\frac{1}{8} x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 363

$$\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \left( \frac{73}{6} \int \frac{x^4}{\sqrt{c^2 x^2 - 1}} dx - \frac{43}{6} x^5 \sqrt{c^2 x^2 - 1} \right) + \frac{3}{8} c^2 x^7 \sqrt{c^2 x^2 - 1} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{8} c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} c^2 x^6 (a + \operatorname{barccosh}(cx))$$

---


$$4\sqrt{cx-1}\sqrt{cx+1}$$

$$\frac{1}{8} x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 262

$$\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \left( \frac{73}{6} \left( \frac{3 \int \frac{x^2}{\sqrt{c^2 x^2 - 1}} dx + x^3 \sqrt{c^2 x^2 - 1}}{4c^2} \right) - \frac{43}{6} x^5 \sqrt{c^2 x^2 - 1} \right) + \frac{3}{8} c^2 x^7 \sqrt{c^2 x^2 - 1} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{8} c^4 x^8 (a + \operatorname{barccosh}(cx))$$

---


$$4\sqrt{cx-1}\sqrt{cx+1}$$

$$\frac{1}{8} x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 262

$$\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \left( \frac{73}{6} \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{c^2 x^2 - 1}} dx + x \sqrt{c^2 x^2 - 1}}{2c^2} \right) + x^3 \sqrt{c^2 x^2 - 1}}{4c^2} \right) - \frac{43}{6} x^5 \sqrt{c^2 x^2 - 1} \right) + \frac{3}{8} c^2 x^7 \sqrt{c^2 x^2 - 1} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{1}{8} c^4 x^8 (a + \operatorname{barccosh}(cx))$$

---


$$4\sqrt{cx-1}\sqrt{cx+1}$$

$$\frac{1}{8} x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 224

---

3.187.  $\int x^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{bc\sqrt{c^2 x^2 - 1}}{24\sqrt{cx - 1}\sqrt{cx + 1}} \left( \frac{1}{8} \left( \frac{73}{6} \left( \frac{3 \left( \frac{\int \frac{1 - \frac{1}{c^2 x^2} d - \frac{x}{\sqrt{c^2 x^2 - 1}}}{1 - \frac{c^2 x^2 - 1}{2c^2}} + \frac{x\sqrt{c^2 x^2 - 1}}{2c^2} \right)}{4c^2} + \frac{x^3 \sqrt{c^2 x^2 - 1}}{4c^2} \right) - \frac{43}{6} x^5 \sqrt{c^2 x^2 - 1} + \frac{3}{8} c^2 x^7 \sqrt{c^2 x^2 - 1} \right) \right) \right) +$$

$4\sqrt{cx - 1}\sqrt{cx + 1}$

$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 219

$$\frac{5}{8}d \int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx -$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{8} c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3} c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4} x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1}}{24\sqrt{cx - 1}\sqrt{cx + 1}} \left( \frac{1}{8} \left( \frac{73}{6} \right) \right) \right) +$$

$4\sqrt{cx - 1}\sqrt{cx + 1}$

$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 6345

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{bcd\sqrt{d - c^2 dx^2} \int -x^3(1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \right.$$

$$\left. bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \left( \frac{1}{8} \frac{7}{6} \right) \right) \right.$$


---


$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 25

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x^3(1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6} \right.$$

$$\left. bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \left( \frac{1}{8} \frac{7}{6} \right) \right) \right.$$


---


$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 6327

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \int x^3 (1 - c^2 x^2) (a + \operatorname{barccosh}(cx)) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \right) + \frac{bcd^2\sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1}}{\frac{1}{8} \left( \frac{7}{6} \right)} \right)$$


---


$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 6336

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \left( -bc \int \frac{x^4 (3 - 2c^2 x^2)}{12\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{1}{6}c^2 x^6 (a + \operatorname{barccosh}(cx))^2 \right)}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{6}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \right) + \frac{bcd^2\sqrt{d - c^2 dx^2}}{4\sqrt{cx - 1}\sqrt{cx + 1}} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1}}{\frac{1}{8} \left( \frac{7}{6} \right)} \right)$$


---


$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 27

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \int \frac{x^4(3-2c^2x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{6}c^2x^6(a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4x^8(a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{\frac{1}{8}} \left( \frac{7}{6} \right) \right)$$

---


$$\frac{\frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 960

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \int \frac{x^4}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}x^5\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4x^8(a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{\frac{1}{8}} \left( \frac{7}{6} \right) \right)$$

---


$$\frac{\frac{1}{8}x^3(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{4\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 111



$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \left( \frac{\int \frac{3x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right) \right)}{3} \right)$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \left( \frac{7}{6} \right) \right)}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$


---


$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 27

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^3 \sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right) \right)}{3} \right)$$

$$bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \left( \frac{7}{6} \right) \right)}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$


---


$$\frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 101

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{12}bc \left( \frac{4}{3} \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx + x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \right) \right)}{4c^2} \right)}{bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \left( \frac{7}{6} \right) \right)}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right)} \right. \\ \left. \frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \right)$$

↓ 43

$$\frac{5}{8}d \left( \frac{1}{2}d \int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{6}x^3 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6}c \right)}{bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{8}c^4 x^8 (a + \operatorname{barccosh}(cx)) - \frac{1}{3}c^2 x^6 (a + \operatorname{barccosh}(cx)) + \frac{1}{4}x^4 (a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{1}{8} \left( \frac{7}{6} \right) \right)}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right)} \right. \\ \left. \frac{1}{8}x^3 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 \right)$$

↓ 6341

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \int x^3(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) \right.$$

$$\left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{8}c^4x^8(a+\operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{8} \left( \frac{7}{6} \right) \right) \right.$$


---


$$\frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2$$

↓ 6298

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \int \frac{x^4}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \right.$$

$$\left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{8}c^4x^8(a+\operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{8} \left( \frac{7}{6} \right) \right) \right.$$


---


$$\frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2$$

↓ 111

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{\int \frac{3x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \dots \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$

$$\left. \left. \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{8}c^4x^8(a+\operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{\frac{1}{8}} \left( \frac{7}{6} \dots \right) \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$

$$\left. \left. \frac{\frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$

↓ 27

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \dots \right) \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$

$$\left. \left. \frac{bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{8}c^4x^8(a+\operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{\frac{1}{8}} \left( \frac{7}{6} \dots \right) \right)}{4\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$

$$\left. \left. \frac{\frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{4\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$

↓ 101

---

3.187.  $\int x^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{2c^2} \right)}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$


---


$$\left. \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{8}c^4x^8(a+\operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{\frac{1}{8}} \frac{7}{6} \right) \right) \right.$$


---


$$\left. \frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 \right) \quad 4\sqrt{cx-1}\sqrt{cx+1}$$

↓ 43

$$\frac{5}{8}d \left( \frac{1}{2}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{2c^2} \right)}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right.$$


---


$$\left. \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{8}c^4x^8(a+\operatorname{barccosh}(cx)) - \frac{1}{3}c^2x^6(a+\operatorname{barccosh}(cx)) + \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{\frac{1}{8}} \frac{7}{6} \right) \right) \right.$$


---


$$\left. \frac{1}{8}x^3(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 \right) \quad 4\sqrt{cx-1}\sqrt{cx+1}$$

↓ 6354

---

3.187.  $\int x^2(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
 & \frac{1}{8}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 x^3 - \\
 & bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{8} c^4 (a + \operatorname{barccosh}(cx)) x^8 - \frac{1}{3} c^2 (a + \operatorname{barccosh}(cx)) x^6 + \frac{1}{4} (a + \operatorname{barccosh}(cx)) x^4 - \frac{bc\sqrt{c^2 x^2 - 1}}{\frac{3}{8} c^2 \sqrt{d - c^2 dx^2}} \right) \\
 \hline
 & \frac{5}{8} d \left( \frac{1}{6} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 x^3 - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6} c^2 (a + \operatorname{barccosh}(cx)) x^6 + \frac{1}{4} (a + \operatorname{barccosh}(cx)) x^4 - \frac{bc\sqrt{c^2 x^2 - 1}}{\frac{3}{8} c^2 \sqrt{d - c^2 dx^2}} \right)}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right) \\
 & \quad \downarrow \text{6298} \\
 & \frac{1}{8}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 x^3 - \\
 & bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{8} c^4 (a + \operatorname{barccosh}(cx)) x^8 - \frac{1}{3} c^2 (a + \operatorname{barccosh}(cx)) x^6 + \frac{1}{4} (a + \operatorname{barccosh}(cx)) x^4 - \frac{bc\sqrt{c^2 x^2 - 1}}{\frac{3}{8} c^2 \sqrt{d - c^2 dx^2}} \right) \\
 \hline
 & \frac{5}{8} d \left( \frac{1}{6} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 x^3 - \frac{bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{6} c^2 (a + \operatorname{barccosh}(cx)) x^6 + \frac{1}{4} (a + \operatorname{barccosh}(cx)) x^4 - \frac{bc\sqrt{c^2 x^2 - 1}}{\frac{3}{8} c^2 \sqrt{d - c^2 dx^2}} \right)}{4\sqrt{cx - 1}\sqrt{cx + 1}} \right)
 \end{aligned}$$

input `Int[x^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

## 3.187.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 111 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 262 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*(m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 363 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^(m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`

rule 960 `Int[((e_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^(m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 1590 `Int[((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^(m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]`

rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^(m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`



rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +
c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& NeQ[m, -1]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_ + (
e1_.)*(x_))^(p_.)*((d2_ + (e2_.)*(x_))^(p_.), x_Symbol] :> Int[(f*x)^m*(d1
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_ + (e_.)*(x_
)^2)^(p_.), x_Symbol] :> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*Sqrt[(d_ +
(e_.)*(x_)^2], x_Symbol] :> Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sq
rt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e
x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x
])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0]
&& IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_ + (e_
.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*
x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m
+ 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m
+ 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1)
, x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n,
0] && GtQ[p, 0] && !LtQ[m, -1]`

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.187.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2527 vs.  $2(741) = 1482$ .

Time = 0.88 (sec) , antiderivative size = 2528, normalized size of antiderivative = 3.01

method	result	size
default	Expression too large to display	2528
parts	Expression too large to display	2528

```
input int(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

output

```

-1/8*a^2*x*(-c^2*d*x^2+d)^(7/2)/c^2/d+1/48*a^2/c^2*x*(-c^2*d*x^2+d)^(5/2)+
5/192*a^2/c^2*d*x*(-c^2*d*x^2+d)^(3/2)+5/128*a^2/c^2*d^2*x*(-c^2*d*x^2+d)^(
1/2)+5/128*a^2/c^2*d^3/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d
)^(1/2))+b^2*(-5/384*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/c^
3*arccosh(c*x)^3*d^2+1/65536*(-d*(c^2*x^2-1))^(1/2)*(128*c^9*x^9-320*c^7*x
^7+128*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^8*x^8+272*c^5*x^5-256*(c*x+1)^(1/2)*(
c*x-1)^(1/2)*c^6*x^6-88*c^3*x^3+160*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*
c*x-32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(3
2*arccosh(c*x)^2-8*arccosh(c*x)+1)*d^2/(c*x+1)/c^3/(c*x-1)-1/6912*(-d*(c^2
*x^2-1))^(1/2)*(32*c^7*x^7-64*c^5*x^5+32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x
^6+38*c^3*x^3-48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-6*c*x+18*(c*x-1)^(1/2
)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(18*arccosh(c*x)^2-6*
arccosh(c*x)+1)*d^2/(c*x+1)/c^3/(c*x-1)+1/2048*(-d*(c^2*x^2-1))^(1/2)*(8*c
^5*x^5-12*c^3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c*x-1)^(1
/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(8*arccosh(c*x)^2-4
*arccosh(c*x)+1)*d^2/(c*x+1)/c^3/(c*x-1)+1/256*(-d*(c^2*x^2-1))^(1/2)*(2*c
^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(
1/2))*(2*arccosh(c*x)^2-2*arccosh(c*x)+1)*d^2/(c*x+1)/c^3/(c*x-1)+1/256*(-
d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*
x-1)^(1/2)*(c*x+1)^(1/2)-2*c*x)*(2*arccosh(c*x)^2+2*arccosh(c*x)+1)*d^2...

```

### 3.187.5 Fracas [F]

$$\int x^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((a^2*c^4*d^2*x^6 - 2*a^2*c^2*d^2*x^4 + a^2*d^2*x^2 + (b^2*c^4*d^2*x^6 - 2*b^2*c^2*d^2*x^4 + b^2*d^2*x^2)*arccosh(c*x)^2 + 2*(a*b*c^4*d^2*x^6 - 2*a*b*c^2*d^2*x^4 + a*b*d^2*x^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

**3.187.6 Sympy [F(-1)]**

Timed out.

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Timed out}$$

input `integrate(x**2*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.187.7 Maxima [F]**

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `1/384*(8*(-c^2*d*x^2 + d)^(5/2)*x/c^2 - 48*(-c^2*d*x^2 + d)^(7/2)*x/(c^2*d) + 10*(-c^2*d*x^2 + d)^(3/2)*d*x/c^2 + 15*sqrt(-c^2*d*x^2 + d)*d^2*x/c^2 + 15*d^(5/2)*arcsin(c*x)/c^3)*a^2 + integrate((-c^2*d*x^2 + d)^(5/2)*b^2*x^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2 + 2*(-c^2*d*x^2 + d)^(5/2)*a*b*x^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)), x)`

**3.187.8 Giac [F]**

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)^2 x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^2*x^2, x)`

**3.187.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int x^2 (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2} dx$$

input `int(x^2*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2),x)`output `int(x^2*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2), x)`

### 3.188 $\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.188.1 Optimal result . . . . .	1613
3.188.2 Mathematica [A] (verified) . . . . .	1614
3.188.3 Rubi [A] (verified) . . . . .	1614
3.188.4 Maple [B] (verified) . . . . .	1617
3.188.5 Fricas [A] (verification not implemented) . . . . .	1618
3.188.6 Sympy [F(-1)] . . . . .	1619
3.188.7 Maxima [A] (verification not implemented) . . . . .	1619
3.188.8 Giac [F(-2)] . . . . .	1620
3.188.9 Mupad [F(-1)] . . . . .	1620

#### 3.188.1 Optimal result

Integrand size = 27, antiderivative size = 470

$$\begin{aligned} \int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = & -\frac{32b^2 d^2 (1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{245c^2 (1 - cx)(1 + cx)} \\ & - \frac{16b^2 d^2 (1 - c^2 x^2)^2 \sqrt{d - c^2 dx^2}}{735c^2 (1 - cx)(1 + cx)} - \frac{12b^2 d^2 (1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2}}{1225c^2 (1 - cx)(1 + cx)} \\ & - \frac{2b^2 d^2 (1 - c^2 x^2)^4 \sqrt{d - c^2 dx^2}}{343c^2 (1 - cx)(1 + cx)} + \frac{2bd^2 x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{7c\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & - \frac{2bcd^2 x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{7\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{6bc^3 d^2 x^5 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{35\sqrt{-1 + cx}\sqrt{1 + cx}} \\ & - \frac{2bc^5 d^2 x^7 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{49\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2 d} \end{aligned}$$

output

```
-1/7*(-c^2*d*x^2+d)^(7/2)*(a+b*arccosh(c*x))^2/c^2/d-32/245*b^2*d^2*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)-16/735*b^2*d^2*(-c^2*x^2+1)^2*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)-12/1225*b^2*d^2*(-c^2*x^2+1)^3*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)-2/343*b^2*d^2*(-c^2*x^2+1)^4*(-c^2*d*x^2+d)^(1/2)/c^2/(-c*x+1)/(c*x+1)+2/7*b*d^2*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/7*b*c*d^2*x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+6/35*b*c^3*d^2*x^5*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/49*b*c^5*d^2*x^7*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.188.2 Mathematica [A] (verified)

Time = 1.51 (sec) , antiderivative size = 234, normalized size of antiderivative = 0.50

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{d^2 \sqrt{d - c^2 dx^2} \left( 3675a^2(-1 + c^2 x^2)^4 - 210abcx \sqrt{-1 + cx} \sqrt{1 + cx} (-35 + 35c^2 x^2 - 21c^4 x^4 + 5c^6 x^6) + 2b^2(2161 - 2918c^2 x^2 + 1108c^4 x^4 - 426c^6 x^6 + 75c^8 x^8) + 210b(35a(-1 + c^2 x^2)^4 + bcx \sqrt{-1 + cx} \sqrt{1 + cx} (35 - 35c^2 x^2 + 21c^4 x^4 - 5c^6 x^6)) \operatorname{ArcCosh}[cx] + 3675b^2(-1 + c^2 x^2)^4 \operatorname{ArcCosh}[cx]^2 \right)}{25725c^2(-1 + c^2 x^2)}$$

input `Integrate[x*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output `(d^2*sqrt[d - c^2*d*x^2]*(3675*a^2*(-1 + c^2*x^2)^4 - 210*a*b*c*x*sqrt[-1 + c*x]*sqrt[1 + c*x]*(-35 + 35*c^2*x^2 - 21*c^4*x^4 + 5*c^6*x^6) + 2*b^2*(2161 - 2918*c^2*x^2 + 1108*c^4*x^4 - 426*c^6*x^6 + 75*c^8*x^8) + 210*b*(35*a*(-1 + c^2*x^2)^4 + b*c*x*sqrt[-1 + c*x]*sqrt[1 + c*x]*(35 - 35*c^2*x^2 + 21*c^4*x^4 - 5*c^6*x^6))*ArcCosh[c*x] + 3675*b^2*(-1 + c^2*x^2)^4*ArcCosh[c*x]^2)/(25725*c^2*(-1 + c^2*x^2))`

### 3.188.3 Rubi [A] (verified)

Time = 1.02 (sec) , antiderivative size = 259, normalized size of antiderivative = 0.55, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6329, 25, 6304, 6309, 27, 2113, 2331, 2389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx \\ & \quad \downarrow \text{6329} \\ & -\frac{2bd^2 \sqrt{d - c^2 dx^2} \int -(1 - cx)^3 (cx + 1)^3 (a + \operatorname{barccosh}(cx)) dx}{7c \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2 d} \\ & \quad \downarrow \text{25} \\ & \frac{2bd^2 \sqrt{d - c^2 dx^2} \int (1 - cx)^3 (cx + 1)^3 (a + \operatorname{barccosh}(cx)) dx}{7c \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{(d - c^2 dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2 d} \\ & \quad \downarrow \text{6304} \end{aligned}$$

---

3.188.  $\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\frac{2bd^2\sqrt{d-c^2dx^2} \int (1-c^2x^2)^3 (a + \operatorname{barccosh}(cx)) dx}{7c\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2d}$$

↓ 6309

$$\frac{2bd^2\sqrt{d-c^2dx^2} \left( -bc \int \frac{x(-5c^6x^6+21c^4x^4-35c^2x^2+35)}{35\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^6x^7(a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4x^5(a + \operatorname{barccosh}(cx)) - c^2x \right)}{7c\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2d}$$

↓ 27

$$\frac{2bd^2\sqrt{d-c^2dx^2} \left( -\frac{1}{35}bc \int \frac{x(-5c^6x^6+21c^4x^4-35c^2x^2+35)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{7}c^6x^7(a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4x^5(a + \operatorname{barccosh}(cx)) - c^2x \right)}{7c\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2d}$$

↓ 2113

$$\frac{2bd^2\sqrt{d-c^2dx^2} \left( -\frac{bc\sqrt{c^2x^2-1} \int \frac{x(-5c^6x^6+21c^4x^4-35c^2x^2+35)}{\sqrt{c^2x^2-1}} dx}{35\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}c^6x^7(a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4x^5(a + \operatorname{barccosh}(cx)) - c^2x \right)}{7c\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2d}$$

↓ 2331

$$\frac{2bd^2\sqrt{d-c^2dx^2} \left( -\frac{bc\sqrt{c^2x^2-1} \int \frac{-5c^6x^6+21c^4x^4-35c^2x^2+35}{\sqrt{c^2x^2-1}} dx^2}{70\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}c^6x^7(a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4x^5(a + \operatorname{barccosh}(cx)) - c^2x \right)}{7c\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2d}$$

↓ 2389

$$\frac{2bd^2\sqrt{d-c^2dx^2} \left( -\frac{bc\sqrt{c^2x^2-1} \int \left( -5(c^2x^2-1)^{5/2} + 6(c^2x^2-1)^{3/2} - 8\sqrt{c^2x^2-1} + \frac{16}{\sqrt{c^2x^2-1}} \right) dx^2}{70\sqrt{cx-1}\sqrt{cx+1}} - \frac{1}{7}c^6x^7(a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4x^5(a + \operatorname{barccosh}(cx)) - c^2x \right)}{7c\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{7/2} (a + \operatorname{barccosh}(cx))^2}{7c^2d}$$

↓ 2009

---

3.188.  $\int x(d-c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$



$$2bd^2\sqrt{d-c^2dx^2} \left( -\frac{1}{7}c^6x^7(a + \operatorname{barccosh}(cx)) + \frac{3}{5}c^4x^5(a + \operatorname{barccosh}(cx)) - c^2x^3(a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) \right)$$


---


$$\frac{(d-c^2dx^2)^{7/2}(a + \operatorname{barccosh}(cx))^2}{7c^2d} \quad 7c\sqrt{cx-1}\sqrt{cx+1}$$

input `Int[x*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output `-1/7*((d - c^2*d*x^2)^(7/2)*(a + b*ArcCosh[c*x])^2)/(c^2*d) + (2*b*d^2*sqrt[d - c^2*d*x^2]*(-1/70*(b*c*sqrt[-1 + c^2*x^2]*((32*sqrt[-1 + c^2*x^2])/c^2 - (16*(-1 + c^2*x^2)^(3/2))/(3*c^2) + (12*(-1 + c^2*x^2)^(5/2))/(5*c^2) - (10*(-1 + c^2*x^2)^(7/2))/(7*c^2)))/(sqrt[-1 + c*x]*sqrt[1 + c*x]) + x*(a + b*ArcCosh[c*x]) - c^2*x^3*(a + b*ArcCosh[c*x]) + (3*c^4*x^5*(a + b*ArcCosh[c*x]))/5 - (c^6*x^7*(a + b*ArcCosh[c*x]))/7)/(7*c*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.188.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2113 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

```
rule 2331 Int[(Pq_)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Simp[1/2 S
ubst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /;
FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]
```

```
rule 2389 Int[(Pq_)*((a_) + (b_.)*(x_)^(n_.))^p, x_Symbol] := Int[ExpandIntegrand
[Pq*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, n}, x] && PolyQ[Pq, x] && (IGtQ[p
, 0] || EqQ[n, 1])
```

```
rule 6304 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n)*((d1_) + (e1_.)*(x_)^2)^(p_.)*
(d2_) + (e2_.)*(x_)^p, x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*A
rcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 +
d1*e2, 0] && IntegerQ[p]
```

```
rule 6309 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symb
ol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u,
x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x]
, x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

### 3.188.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1957 vs.  $2(418) = 836$ .

Time = 0.74 (sec) , antiderivative size = 1958, normalized size of antiderivative = 4.17

method	result	size
default	Expression too large to display	1958
parts	Expression too large to display	1958

```
input int(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

$$3.188. \quad \int x(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))^2 dx$$

output

```

-1/7*a^2*(-c^2*d*x^2+d)^(7/2)/c^2/d+b^2*(1/43904*(-d*(c^2*x^2-1))^(1/2)*(6
4*c^8*x^8-144*c^6*x^6+64*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^7*c^7+104*c^4*x^4-1
12*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5-25*c^2*x^2+56*(c*x-1)^(1/2)*(c*x+1)
^(1/2)*c^3*x^3-7*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(49*arccosh(c*x)^2-14*
arccosh(c*x)+2)*d^2/(c*x+1)/c^2/(c*x-1)-1/3200*(-d*(c^2*x^2-1))^(1/2)*(16*
c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c
*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(25
*arccosh(c*x)^2-10*arccosh(c*x)+2)*d^2/(c*x+1)/c^2/(c*x-1)+1/384*(-d*(c^2*
x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-3
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(9*arccosh(c*x)^2-6*arccosh(c*x)+2)*d^
2/(c*x+1)/c^2/(c*x-1)-5/128*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(
1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*arccosh(c*x)+2)*d^2/(c*x+1)/c^2/(c*
x-1)-5/128*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^
2-1)*(arccosh(c*x)^2+2*arccosh(c*x)+2)*d^2/(c*x+1)/c^2/(c*x-1)+1/384*(-d*(
c^2*x^2-1))^(1/2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x
-1)^(1/2)*(c*x+1)^(1/2)*c*x-5*c^2*x^2+1)*(9*arccosh(c*x)^2+6*arccosh(c*x)+
2)*d^2/(c*x+1)/c^2/(c*x-1)-1/3200*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)
*(c*x-1)^(1/2)*c^5*x^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-
28*c^4*x^4-5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(25*arccosh(c*x)
)^2+10*arccosh(c*x)+2)*d^2/(c*x+1)/c^2/(c*x-1)+1/43904*(-d*(c^2*x^2-1))...

```

### 3.188.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 477, normalized size of antiderivative = 1.01

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{3675 (b^2 c^8 d^2 x^8 - 4 b^2 c^6 d^2 x^6 + 6 b^2 c^4 d^2 x^4 - 4 b^2 c^2 d^2 x^2 + b^2 d^2) \sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{-c^2 dx^2 + d})}{\dots}$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output  $1/25725*(3675*(b^2*c^8*d^2*x^8 - 4*b^2*c^6*d^2*x^6 + 6*b^2*c^4*d^2*x^4 - 4*b^2*c^2*d^2*x^2 + b^2*d^2)*\sqrt{-c^2*d*x^2 + d}*\log(c*x + \sqrt{c^2*x^2 - 1})^2 - 210*(5*a*b*c^7*d^2*x^7 - 21*a*b*c^5*d^2*x^5 + 35*a*b*c^3*d^2*x^3 - 35*a*b*c*d^2*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} - 210*((5*b^2*c^7*d^2*x^7 - 21*b^2*c^5*d^2*x^5 + 35*b^2*c^3*d^2*x^3 - 35*b^2*c*d^2*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} - 35*(a*b*c^8*d^2*x^8 - 4*a*b*c^6*d^2*x^6 + 6*a*b*c^4*d^2*x^4 - 4*a*b*c^2*d^2*x^2 + a*b*d^2)*\sqrt{-c^2*d*x^2 + d})*\log(c*x + \sqrt{c^2*x^2 - 1}) + (75*(49*a^2 + 2*b^2)*c^8*d^2*x^8 - 12*(1225*a^2 + 71*b^2)*c^6*d^2*x^6 + 2*(11025*a^2 + 1108*b^2)*c^4*d^2*x^4 - 4*(3675*a^2 + 1459*b^2)*c^2*d^2*x^2 + (3675*a^2 + 4322*b^2)*d^2)*\sqrt{-c^2*d*x^2 + d})/(c^4*x^2 - c^2)$

### 3.188.6 Sympy [F(-1)]

Timed out.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Timed out}$$

input `integrate(x*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2,x)`

output Timed out

### 3.188.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 337, normalized size of antiderivative = 0.72

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx =$$

$$\frac{(-c^2 dx^2 + d)^{7/2} b^2 \operatorname{arccosh}(cx)^2}{7 c^2 d} - \frac{2(-c^2 dx^2 + d)^{7/2} ab \operatorname{arccosh}(cx)}{7 c^2 d}$$

$$+ \frac{2}{25725} b^2 \left( \frac{75 \sqrt{c^2 x^2 - 1} c^4 \sqrt{-d} d^3 x^6 - 351 \sqrt{c^2 x^2 - 1} c^2 \sqrt{-d} d^3 x^4 + 757 \sqrt{c^2 x^2 - 1} \sqrt{-d} d^3 x^2 - \frac{2161 \sqrt{c^2 x^2 - 1}}{c^2}}{d} \right)$$

$$- \frac{(-c^2 dx^2 + d)^{7/2} a^2}{7 c^2 d} - \frac{2(5 c^6 \sqrt{-d} d^3 x^7 - 21 c^4 \sqrt{-d} d^3 x^5 + 35 c^2 \sqrt{-d} d^3 x^3 - 35 \sqrt{-d} d^3 x) ab}{245 cd}$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

---

3.188.  $\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

```
output -1/7*(-c^2*d*x^2 + d)^(7/2)*b^2*arccosh(c*x)^2/(c^2*d) - 2/7*(-c^2*d*x^2 +
d)^(7/2)*a*b*arccosh(c*x)/(c^2*d) + 2/25725*b^2*((75*sqrt(c^2*x^2 - 1)*c^
4*sqrt(-d)*d^3*x^6 - 351*sqrt(c^2*x^2 - 1)*c^2*sqrt(-d)*d^3*x^4 + 757*sqrt
(c^2*x^2 - 1)*sqrt(-d)*d^3*x^2 - 2161*sqrt(c^2*x^2 - 1)*sqrt(-d)*d^3/c^2)/
d - 105*(5*c^6*sqrt(-d)*d^3*x^7 - 21*c^4*sqrt(-d)*d^3*x^5 + 35*c^2*sqrt(-d
)*d^3*x^3 - 35*sqrt(-d)*d^3*x)*arccosh(c*x)/(c*d)) - 1/7*(-c^2*d*x^2 + d)^(
7/2)*a^2/(c^2*d) - 2/245*(5*c^6*sqrt(-d)*d^3*x^7 - 21*c^4*sqrt(-d)*d^3*x^
5 + 35*c^2*sqrt(-d)*d^3*x^3 - 35*sqrt(-d)*d^3*x)*a*b/(c*d)
```

### 3.188.8 Giac [F(-2)]

Exception generated.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

```
input integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.188.9 Mupad [F(-1)]

Timed out.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int x(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2} dx$$

```
input int(x*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2),x)
```

```
output int(x*(a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2), x)
```

### 3.189 $\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

3.189.1 Optimal result	.1621
3.189.2 Mathematica [A] (warning: unable to verify)	.1622
3.189.3 Rubi [A] (verified)	.1623
3.189.4 Maple [B] (verified)	.1629
3.189.5 Fricas [F]	.1630
3.189.6 Sympy [F(-1)]	.1631
3.189.7 Maxima [F]	.1631
3.189.8 Giac [F(-2)]	.1631
3.189.9 Mupad [F(-1)]	.1632

#### 3.189.1 Optimal result

Integrand size = 26, antiderivative size = 486

$$\begin{aligned} \int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx &= \frac{245b^2 d^2 x \sqrt{d - c^2 dx^2}}{1152} \\ &+ \frac{65b^2 d^2 x (1 - cx)(1 + cx) \sqrt{d - c^2 dx^2}}{1728} + \frac{1}{108} b^2 d^2 x (1 - cx)^2 (1 + cx)^2 \sqrt{d - c^2 dx^2} \\ &+ \frac{115b^2 d^2 \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{1152c \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{5bcd^2 x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{16 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &+ \frac{5bd^2 (1 - c^2 x^2)^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{48c \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &+ \frac{bd^2 (1 - c^2 x^2)^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{18c \sqrt{-1 + cx} \sqrt{1 + cx}} \\ &+ \frac{5}{16} d^2 x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 + \frac{5}{24} dx (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 \\ &+ \frac{1}{6} x (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{5d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^3}{48bc \sqrt{-1 + cx} \sqrt{1 + cx}} \end{aligned}$$

output 
$$\begin{aligned} & 5/24*d*x*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^{2+1/6*x*(-c^2*d*x^2+d)^{(5/2)}*(a+b*\operatorname{arccosh}(c*x))^{2+245/1152*b^2*d^2*x*(-c^2*d*x^2+d)^{(1/2)}+65/1728*b^2*d^2*x*(-c*x+1)*(c*x+1)*(-c^2*d*x^2+d)^{(1/2)}+1/108*b^2*d^2*x*(-c*x+1)^2*(c*x+1)^2*(-c^2*d*x^2+d)^{(1/2)}+5/16*d^2*x*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d*x^2+d)^{(1/2)}+115/1152*b^2*d^2*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-5/16*b*c*d^2*x^2*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+5/48*b*d^2*(-c^2*x^2+1)^2*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/18*b*d^2*(-c^2*x^2+1)^3*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-5/48*d^2*(a+b*\operatorname{arccosh}(c*x))^{3*(-c^2*d*x^2+d)^{(1/2)}/b/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \end{aligned}$$

### 3.189.2 Mathematica [A] (warning: unable to verify)

Time = 3.65 (sec) , antiderivative size = 740, normalized size of antiderivative = 1.52

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \frac{d^2 \left( 9504a^2 cx \sqrt{\frac{-1+cx}{1+cx}} \sqrt{d - c^2 dx^2} + 9504a^2 c^2 x^2 \sqrt{\frac{-1+cx}{1+cx}} \sqrt{d - c^2 dx^2} - 7488a^2 c^3 x^3 \sqrt{\frac{-1+cx}{1+cx}} \right)}{c^3}$$

input `Integrate[(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output

```
(d^2*(9504*a^2*c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] + 9504*a^2*c^2*x^2*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] - 7488*a^2*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] - 7488*a^2*c^4*x^4*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] + 2304*a^2*c^5*x^5*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] + 2304*a^2*c^6*x^6*Sqrt[(-1 + c*x)/(1 + c*x)]*Sqrt[d - c^2*d*x^2] - 1440*b^2*Sqrt[d - c^2*d*x^2]*ArcCosh[c*x]^3 - 4320*a^2*Sqrt[d]*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] - 4320*a^2*c*Sqrt[d]*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] - 3240*a*b*Sqrt[d - c^2*d*x^2]*Cosh[2*ArcCosh[c*x]] + 324*a*b*Sqrt[d - c^2*d*x^2]*Cosh[4*ArcCosh[c*x]] - 24*a*b*Sqrt[d - c^2*d*x^2]*Cosh[6*ArcCosh[c*x]] + 1620*b^2*Sqrt[d - c^2*d*x^2]*Sinh[2*ArcCosh[c*x]] - 81*b^2*Sqrt[d - c^2*d*x^2]*Sinh[4*ArcCosh[c*x]] + 4*b^2*Sqrt[d - c^2*d*x^2]*Sinh[6*ArcCosh[c*x]] - 12*b*Sqrt[d - c^2*d*x^2]*ArcCosh[c*x]*(270*b*Cosh[2*ArcCosh[c*x]] - 27*b*Cosh[4*ArcCosh[c*x]] + 2*b*Cosh[6*ArcCosh[c*x]] - 540*a*Sinh[2*ArcCosh[c*x]] + 108*a*Sinh[4*ArcCosh[c*x]] - 12*a*Sinh[6*ArcCosh[c*x]]) + 72*b*Sqrt[d - c^2*d*x^2]*ArcCosh[c*x]^2*(-60*a + 45*b*Sinh[2*ArcCosh[c*x]] - 9*b*Sinh[4*ArcCosh[c*x]] + b*Sinh[6*ArcCosh[c*x]])))/(13824*c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))
```

### 3.189.3 Rubi [A] (verified)

Time = 2.37 (sec) , antiderivative size = 561, normalized size of antiderivative = 1.15, number of steps used = 14, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {6312, 6312, 25, 6310, 6298, 101, 43, 6308, 6327, 6329, 40, 40, 40, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$\downarrow \text{6312}$$

$$-\frac{bcd^2 \sqrt{d - c^2 dx^2} \int x(1 - cx)^2 (cx + 1)^2 (a + \operatorname{barccosh}(cx)) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{5}{6}d \int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{6}x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

$$\downarrow \text{6312}$$

---

3.189.  $\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$



$$\begin{aligned}
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( \frac{bcd\sqrt{d-c^2dx^2} \int -x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2dx + \frac{1}{4}x \right. \\
& \left. \frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 \right) \\
& \quad \downarrow \text{25} \\
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \int \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2dx + \frac{1}{4}x \right. \\
& \left. \frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 \right) \\
& \quad \downarrow \text{6310} \\
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \int x(a+\operatorname{barccosh}(cx))dx}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
& \left. \left. \frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 \right) \right) \\
& \quad \downarrow \text{6298} \\
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \right. \right.}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
& \left. \left. \frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 \right) \right) \\
& \quad \downarrow \text{101} \\
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \right. \right.}{\sqrt{cx-1}} \right. \right. \\
& \left. \left. \frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 \right) \right) \\
& \quad \downarrow \text{43}
\end{aligned}$$

$$\begin{aligned}
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x \right. \right. \\
& \left. \left. \frac{1}{6}x(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \right. \right. \\
& \quad \downarrow \text{6308} \\
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 + \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x \right. \right. \\
& \left. \left. \frac{1}{6}x(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \right. \right. \\
& \quad \downarrow \text{6327} \\
& -\frac{bcd^2\sqrt{d-c^2dx^2} \int x(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 + \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x \right. \right. \\
& \left. \left. \frac{1}{6}x(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \right. \right. \\
& \quad \downarrow \text{6329} \\
& -\frac{bcd^2\sqrt{d-c^2dx^2} \left( -\frac{b \int (cx-1)^{5/2}(cx+1)^{5/2} dx}{6c} - \frac{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))}{6c^2} \right)}{3\sqrt{cx-1}\sqrt{cx+1}} + \\
\frac{5}{6}d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b \int (cx-1)^{3/2}(cx+1)^{3/2} dx}{4c} - \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 \right. \\
& \left. \left. \frac{1}{6}x(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \right. \right. \\
& \quad \downarrow \text{40}
\end{aligned}$$

$$\frac{bcd^2\sqrt{d-c^2dx^2}\left(-\frac{b(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2}-\frac{5}{6}\int(cx-1)^{3/2}(cx+1)^{3/2}dx)}{6c}-\frac{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))}{6c^2}\right)}{3\sqrt{cx-1}\sqrt{cx+1}} +$$

$$\frac{5}{6}d\left(-\frac{bcd\sqrt{d-c^2dx^2}\left(\frac{b(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\int\sqrt{cx-1}\sqrt{cx+1}dx)}{4c}-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2}\right)}{2\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{4}x(d-c^2dx^2)^{3/2}\right)$$

$$\frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2$$

↓ 40

$$\frac{bcd^2\sqrt{d-c^2dx^2}\left(-\frac{b(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2}-\frac{5}{6}(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\int\sqrt{cx-1}\sqrt{cx+1}dx))}{6c}-\frac{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))}{6c^2}\right)}{3\sqrt{cx-1}\sqrt{cx+1}} +$$

$$\frac{5}{6}d\left(-\frac{bcd\sqrt{d-c^2dx^2}\left(\frac{b(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx))}{4c}-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2}\right)}{2\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{4}x(d-c^2dx^2)^{3/2}\right)$$

$$\frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2$$

↓ 40

$$\frac{bcd^2\sqrt{d-c^2dx^2}\left(-\frac{b(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2}-\frac{5}{6}(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx)))}{6c}-\frac{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))}{6c^2}\right)}{3\sqrt{cx-1}\sqrt{cx+1}} +$$

$$\frac{5}{6}d\left(-\frac{bcd\sqrt{d-c^2dx^2}\left(\frac{b(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx))}{4c}-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2}\right)}{2\sqrt{cx-1}\sqrt{cx+1}}+\frac{1}{4}x(d-c^2dx^2)^{3/2}\right)$$

$$\frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2$$

↓ 43

$$\frac{bcd^2\sqrt{d-c^2dx^2}\left(-\frac{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))}{6c^2}-\frac{b\left(\frac{1}{6}x(cx-1)^{5/2}(cx+1)^{5/2}-\frac{5}{6}\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{a}{2c}\right)\right)\right)}{6c}\right)}{\frac{1}{6}x(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2+\frac{3\sqrt{cx-1}\sqrt{cx+1}}{4c}}-\frac{\frac{5}{6}d\left(\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2-\frac{bcd\sqrt{d-c^2dx^2}\left(b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)\right)}{4c}\right)}{2\sqrt{cx-1}\sqrt{cx+1}}}$$

input `Int[(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output `(x*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/6 - (b*c*d^2*sqrt[d - c^2*d*x^2]*(-1/6*((1 - c^2*x^2)^3*(a + b*ArcCosh[c*x]))/c^2 - (b*((x*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/6 - (5*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c))))/4))/6))/(6*c)))/(3*sqrt[-1 + c*x]*sqrt[1 + c*x]) + (5*d*((x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/4 + (3*d*((x*sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/2 - (sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^3)/(6*b*c*sqrt[-1 + c*x]*sqrt[1 + c*x]) - (b*c*sqrt[d - c^2*d*x^2]*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/2)))/(sqrt[-1 + c*x]*sqrt[1 + c*x])))/4 - (b*c*d*sqrt[d - c^2*d*x^2]*(-1/4*((1 - c^2*x^2)^2*(a + b*ArcCosh[c*x]))/c^2 + (b*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/(4*c)))/(2*sqrt[-1 + c*x]*sqrt[1 + c*x])))/6`

### 3.189.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_) + (b_)*(x_))2*((c_) + (d_)*(x_))(n_)*((e_) + (f_)*(x_))(p_), x_] := Simp[b*(a + b*x)*(c + d*x)(n + 1)*((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)n*(e + f*x)p*Simp[a2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 6298 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))(n_)*((d_)*(x_))(m_), x_Symbol] := Simp[(d*x)(m + 1)*((a + b*ArcCosh[c*x])n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)(m + 1)*((a + b*ArcCosh[c*x])(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6308 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))(n_)/(Sqrt[(d1_) + (e1_)*(x_)]*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))(n_)*Sqrt[(d_) + (e_)*(x_)2], x_Symbol] := Simp[x*Sqrt[d + e*x2]*((a + b*ArcCosh[c*x])(n/2)), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c2*d + e, 0] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] +  
(Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x  
, x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p  
)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n  
- 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n,  
0] && GtQ[p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.)), x_Symbol] := Int[(f*x)^m*(d1  
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2  
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p  
_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p  
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +  
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x  
)^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&  
GtQ[n, 0] && NeQ[p, -1]`

### 3.189.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1736 vs.  $2(422) = 844$ .

Time = 1.05 (sec) , antiderivative size = 1737, normalized size of antiderivative = 3.57

method	result	size
default	Expression too large to display	1737
parts	Expression too large to display	1737

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output `1/6*x*(-c^2*d*x^2+d)^(5/2)*a^2+5/24*a^2*d*x*(-c^2*d*x^2+d)^(3/2)+5/16*a^2*d^2*x*(-c^2*d*x^2+d)^(1/2)+5/16*a^2*d^3/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b^2*(-5/48*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/c*arccosh(c*x)^3*d^2+1/6912*(-d*(c^2*x^2-1))^(1/2)*(32*c^7*x^7-64*c^5*x^5+32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6+38*c^3*x^3-48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-6*c*x+18*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(18*arccosh(c*x)^2-6*arccosh(c*x)+1)*d^2/(c*x-1)/(c*x+1)/c-3/1024*(-d*(c^2*x^2-1))^(1/2)*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+4*c*x-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(8*arccosh(c*x)^2-4*arccosh(c*x)+1)*d^2/(c*x-1)/(c*x+1)/c+15/256*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(2*arccosh(c*x)^2-2*arccosh(c*x)+1)*d^2/(c*x-1)/(c*x+1)/c+15/256*(-d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)-2*c*x)*(2*arccosh(c*x)^2+2*arccosh(c*x)+1)*d^2/(c*x-1)/(c*x+1)/c-3/1024*(-d*(c^2*x^2-1))^(1/2)*(-8*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+8*c^5*x^5+8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-12*c^3*x^3-(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c*x)*(8*arccosh(c*x)^2+4*arccosh(c*x)+1)*d^2/(c*x-1)/(c*x+1)/c+1/6912*(-d*(c^2*x^2-1))^(1/2)*(-32*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^6*x^6+32*c^7*x^7+48*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4-64*c^5*x^5-18*(c*x-1)^(1/2)*(c*x+1)^(...`

### 3.189.5 Fracas [F]

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2 dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output `integral((a^2*c^4*d^2*x^4 - 2*a^2*c^2*d^2*x^2 + a^2*d^2 + (b^2*c^4*d^2*x^4 - 2*b^2*c^2*d^2*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*c^4*d^2*x^4 - 2*a*b*c^2*d^2*x^2 + a*b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d), x)`

**3.189.6 Sympy [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.189.7 Maxima [F]**

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)^2 dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `1/48*(8*(-c^2*d*x^2 + d)^(5/2)*x + 10*(-c^2*d*x^2 + d)^(3/2)*d*x + 15*sqrt(-c^2*d*x^2 + d)*d^2*x + 15*d^(5/2)*arcsin(c*x)/c)*a^2 + integrate((-c^2*d*x^2 + d)^(5/2)*b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2 + 2*(-c^2*d*x^2 + d)^(5/2)*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1), x)`

**3.189.8 Giac [F(-2)]**

Exception generated.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.189.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2} dx$$

input `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2),x)`output `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2), x)`

**3.190**  $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x} dx$

3.190.1 Optimal result . . . . . 1633  
 3.190.2 Mathematica [A] (warning: unable to verify) . . . . . 1634  
 3.190.3 Rubi [A] (verified) . . . . . 1635  
 3.190.4 Maple [F] . . . . . 1645  
 3.190.5 Fricas [F] . . . . . 1646  
 3.190.6 Sympy [F(-1)] . . . . . 1646  
 3.190.7 Maxima [F] . . . . . 1646  
 3.190.8 Giac [F(-2)] . . . . . 1647  
 3.190.9 Mupad [F(-1)] . . . . . 1647

**3.190.1 Optimal result**

Integrand size = 29, antiderivative size = 836

$$\begin{aligned} \int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x} dx &= \frac{68}{27}b^2d^2\sqrt{d-c^2dx^2} \\ &- \frac{2}{27}b^2c^2d^2x^2\sqrt{d-c^2dx^2} - \frac{2abcd^2x\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{16b^2d^2(1-c^2x^2)\sqrt{d-c^2dx^2}}{75(1-cx)(1+cx)} \\ &+ \frac{8b^2d^2(1-c^2x^2)^2\sqrt{d-c^2dx^2}}{225(1-cx)(1+cx)} + \frac{2b^2d^2(1-c^2x^2)^3\sqrt{d-c^2dx^2}}{125(1-cx)(1+cx)} \\ &- \frac{2b^2cd^2x\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{\sqrt{-1+cx}\sqrt{1+cx}} - \frac{16bcd^2x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{15\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{22bc^3d^2x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{45\sqrt{-1+cx}\sqrt{1+cx}} - \frac{2bc^5d^2x^5\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{25\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 + \frac{1}{3}d(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2 \\ &+ \frac{1}{5}(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2 - \frac{2d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} + \frac{2ibd^2\sqrt{d-c^2dx^2}}{\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output

```

1/3*d*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2+1/5*(-c^2*d*x^2+d)^(5/2)*(
a+b*arccosh(c*x))^2+68/27*b^2*d^2*(-c^2*d*x^2+d)^(1/2)-2/27*b^2*c^2*d^2*x^
2*(-c^2*d*x^2+d)^(1/2)+16/75*b^2*d^2*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/(-c
*x+1)/(c*x+1)+8/225*b^2*d^2*(-c^2*x^2+1)^2*(-c^2*d*x^2+d)^(1/2)/(-c*x+1)/(
c*x+1)+2/125*b^2*d^2*(-c^2*x^2+1)^3*(-c^2*d*x^2+d)^(1/2)/(-c*x+1)/(c*x+1)+
d^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)-2*a*b*c*d^2*x*(-c^2*d*x^2+d)
^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*b^2*c*d^2*x*arccosh(c*x)*(-c^2*d*x^2+
d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-16/15*b*c*d^2*x*(a+b*arccosh(c*x))*(-
c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+22/45*b*c^3*d^2*x^3*(a+b*ar
ccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/25*b*c^5*d^
2*x^5*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-
2*d^2*(a+b*arccosh(c*x))^2*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-c^2*d
*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2*I*b^2*d^2*polylog(3,I*(c*x+(c*
x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2
)+2*I*b*d^2*(a+b*arccosh(c*x))*polylog(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/
2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*I*b^2*d^2*polylog(
3,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2
)/(c*x+1)^(1/2)-2*I*b*d^2*(a+b*arccosh(c*x))*polylog(2,I*(c*x+(c*x-1)^(1/2)
*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)

```

### 3.190.2 Mathematica [A] (warning: unable to verify)

Time = 5.84 (sec) , antiderivative size = 963, normalized size of antiderivative = 1.15

$$\begin{aligned}
& \int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2}{x} dx = \frac{1}{15} a^2 d^2 \sqrt{d - c^2 dx^2} (23 - 11c^2 x^2 + 3c^4 x^4) \\
& - \frac{1}{27} b^2 d^2 \sqrt{d - c^2 dx^2} \left( 2(-13 + \cosh(2 \operatorname{arccosh}(cx))) \right. \\
& + 9 \operatorname{arccosh}(cx)^2 (-1 + \cosh(2 \operatorname{arccosh}(cx))) \\
& \left. + \frac{3 \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) (9cx - \cosh(3 \operatorname{arccosh}(cx)))}{-1 + cx} \right) \\
& - \frac{abd^2 \sqrt{d - c^2 dx^2} \left( 9cx + 12 \left( \frac{-1+cx}{1+cx} \right)^{3/2} (1 + cx)^3 \operatorname{arccosh}(cx) - \cosh(3 \operatorname{arccosh}(cx)) \right)}{9 \sqrt{\frac{-1+cx}{1+cx}} (1 + cx)} \\
& + a^2 d^{5/2} \log(cx) - a^2 d^{5/2} \log \left( d + \sqrt{d} \sqrt{d - c^2 dx^2} \right) + \frac{2abd^2 \sqrt{d - c^2 dx^2} \left( -cx + \sqrt{\frac{-1+cx}{1+cx}} \operatorname{arccosh}(cx) + cx \sqrt{\frac{-1+cx}{1+cx}} \right)}{9 \sqrt{\frac{-1+cx}{1+cx}} (1 + cx)}
\end{aligned}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x,x]`

output `(a^2*d^2*Sqrt[d - c^2*d*x^2]*(23 - 11*c^2*x^2 + 3*c^4*x^4))/15 - (b^2*d^2*Sqrt[d - c^2*d*x^2]*(2*(-13 + Cosh[2*ArcCosh[c*x]]) + 9*ArcCosh[c*x]^2*(-1 + Cosh[2*ArcCosh[c*x]]) + (3*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]*(9*c*x - Cosh[3*ArcCosh[c*x]]))/(-1 + c*x))/27 - (a*b*d^2*Sqrt[d - c^2*d*x^2]*(9*c*x + 12*((-1 + c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3*ArcCosh[c*x] - Cosh[3*ArcCosh[c*x]]))/(9*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + a^2*d^(5/2)*Log[c*x] - a^2*d^(5/2)*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + (2*a*b*d^2*Sqrt[d - c^2*d*x^2]*(-(c*x) + Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x] + I*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh[c*x]*Log[1 + I/E^ArcCosh[c*x]] + I*PolyLog[2, (-I)/E^ArcCosh[c*x]] - I*PolyLog[2, I/E^ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + b^2*d^2*Sqrt[d - c^2*d*x^2]*(2 + (2*c*x*Sqrt[(-1 + c*x)/(1 + c*x)]*ArcCosh[c*x]))/(1 - c*x) + ArcCosh[c*x]^2 + (I*(ArcCosh[c*x]^2*Log[1 - I/E^ArcCosh[c*x]] - ArcCosh[c*x]^2*Log[1 + I/E^ArcCosh[c*x]] + 2*ArcCosh[c*x]*PolyLog[2, (-I)/E^ArcCosh[c*x]] - 2*ArcCosh[c*x]*PolyLog[2, I/E^ArcCosh[c*x]] + 2*PolyLog[3, (-I)/E^ArcCosh[c*x]] - 2*PolyLog[3, I/E^ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) - (a*b*d^2*Sqrt[d - c^2*d*x^2]*(25*Cosh[3*ArcCosh[c*x]] + 9*(-50*c*x + Cosh[5*ArcCosh[c*x]]) + 15*ArcCosh[c*x]*(30*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) - 5*Sinh[3*ArcCosh[c*x]] - 3*Sinh[5*ArcCosh[c*x]])))/(1800*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + ...`

### 3.190.3 Rubi [A] (verified)

Time = 4.46 (sec) , antiderivative size = 613, normalized size of antiderivative = 0.73, number of steps used = 24, number of rules used = 23,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.793$ , Rules used = {6345, 6304, 6309, 27, 1905, 1576, 1140, 2009, 6345, 25, 6304, 6309, 27, 960, 83, 6341, 2009, 6362, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2}{x} dx$$

↓ 6345

$$\frac{2bcd^2 \sqrt{d - c^2 dx^2} \int (1 - cx)^2 (cx + 1)^2 (a + \text{barccosh}(cx)) dx}{5\sqrt{cx - 1}\sqrt{cx + 1}} +$$

$$d \int \frac{(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^2}{x} dx + \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2$$

---

3.190.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2}{x} dx$

$$\begin{aligned}
 & \downarrow \text{6304} \\
 & -\frac{2bcd^2\sqrt{d-c^2dx^2} \int (1-c^2x^2)^2 (a+\operatorname{barccosh}(cx))dx}{5\sqrt{cx-1}\sqrt{cx+1}} + \\
 & d \int \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{5}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \\
 & \downarrow \text{6309} \\
 & \frac{d \int \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x} dx -}{2bcd^2\sqrt{d-c^2dx^2} \left( -bc \int \frac{x(3c^4x^4-10c^2x^2+15)}{15\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \\
 & \frac{1}{5}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \\
 & \downarrow \text{27} \\
 & \frac{d \int \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x} dx -}{2bcd^2\sqrt{d-c^2dx^2} \left( -\frac{1}{15}bc \int \frac{x(3c^4x^4-10c^2x^2+15)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \\
 & \frac{1}{5}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \\
 & \downarrow \text{1905} \\
 & \frac{d \int \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x} dx -}{2bcd^2\sqrt{d-c^2dx^2} \left( -\frac{bc\sqrt{c^2x^2-1} \int \frac{x(3c^4x^4-10c^2x^2+15)}{\sqrt{c^2x^2-1}} dx}{15\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \\
 & \frac{1}{5}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \\
 & \downarrow \text{1576} \\
 & \frac{d \int \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x} dx -}{2bcd^2\sqrt{d-c^2dx^2} \left( -\frac{bc\sqrt{c^2x^2-1} \int \frac{3c^4x^4-10c^2x^2+15}{\sqrt{c^2x^2-1}} dx}{30\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \\
 & \frac{1}{5}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 \\
 & \downarrow \text{1140}
 \end{aligned}$$

---

3.190.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} dx$

$$d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx - 2bcd^2 \sqrt{d - c^2 dx^2} \left( -\frac{bc\sqrt{c^2 x^2 - 1} \int (3(c^2 x^2 - 1)^{3/2} - 4\sqrt{c^2 x^2 - 1} + \frac{8}{\sqrt{c^2 x^2 - 1}}) dx^2}{30\sqrt{cx - 1}\sqrt{cx + 1}} + \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) \right)$$

---


$$\frac{1}{5}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2$$

↓ 2009

$$d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{5}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - 2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)}{5} \right)}{5\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

---


$$5\sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 6345

$$d \left( \frac{2bcd\sqrt{d - c^2 dx^2} \int -((1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{1}{5}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \right.$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)}{5} \right)}{5\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

---


$$5\sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 25

$$d \left( -\frac{2bcd\sqrt{d - c^2 dx^2} \int (1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx)) dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{1}{5}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \right.$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)}{5} \right)}{5\sqrt{cx - 1}\sqrt{cx + 1}} \right)$$

---


$$5\sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 6304

---

3.190.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$

$$d \left( -\frac{2bcd\sqrt{d-c^2dx^2} \int (1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{3\sqrt{cx-1}\sqrt{cx+1}} + d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 - \right.$$

$$\left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{5} \left( \frac{6(c^2x^2)}{5} \right) \right) \right.$$


---


$$5\sqrt{cx-1}\sqrt{cx+1}$$

↓ 6309

$$d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx - \frac{2bcd\sqrt{d-c^2dx^2} \left( -bc \int \frac{x(3-c^2x^2)}{3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + \right. \right.$$

$$\left. \frac{1}{5}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 - \right.$$

$$\left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{5} \left( \frac{6(c^2x^2)}{5} \right) \right) \right.$$


---


$$5\sqrt{cx-1}\sqrt{cx+1}$$

↓ 27

$$d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx - \frac{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}bc \int \frac{x(3-c^2x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) - \right. \right.$$

$$\left. \frac{1}{5}(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 - \right.$$

$$\left. 2bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{5} \left( \frac{6(c^2x^2)}{5} \right) \right) \right.$$


---


$$5\sqrt{cx-1}\sqrt{cx+1}$$

↓ 960

---

3.190.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} dx$

$$d \left( d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{3}bc \left( \frac{7}{3} \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \right.$$

$$\left. \frac{1}{5}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \right.$$

$$2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)}{5} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right)$$


---

↓ 83

$$d \left( d \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3}(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd\sqrt{d - c^2 dx^2} \left( -\frac{1}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} - \right.$$

$$\left. \frac{1}{5}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \right.$$

$$2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)}{5} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right)$$


---

↓ 6341

$$d \left( d \left( -\frac{2bc\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2 dx^2} \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \right) - \right.$$

$$\left. \frac{1}{5}(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \right.$$

$$2bcd^2\sqrt{d - c^2 dx^2} \left( \frac{1}{5}c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3}c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)}{5} \right)}{5\sqrt{cx-1}\sqrt{cx+1}} \right)$$


---

↓ 2009

---

3.190.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$



$$d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\ \left. \left. - \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{5} \left( \frac{6(c^2x^2-1)^{5/2}}{5} \right) \right) \right) \right)$$

↓ 6362

$$d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} d\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\ \left. \left. - \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{5} \left( \frac{6(c^2x^2-1)^{5/2}}{5} \right) \right) \right) \right)$$

↓ 3042

$$d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx))^2 \csc(i\operatorname{arccosh}(cx) + \frac{\pi}{2}) d\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \right. \right. \\ \left. \left. - \frac{1}{5}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bcd^2\sqrt{d-c^2dx^2}}{5\sqrt{cx-1}\sqrt{cx+1}} \left( \frac{1}{5}c^4x^5(a+\operatorname{barccosh}(cx)) - \frac{2}{3}c^2x^3(a+\operatorname{barccosh}(cx)) + x(a+\operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1}}{5} \left( \frac{6(c^2x^2-1)^{5/2}}{5} \right) \right) \right) \right)$$

↓ 4668

---

3.190.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} dx$

$$d \left( d \left( - \frac{\sqrt{d - c^2 dx^2} (-2ib \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) dx)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right. \right. \\ \left. \left. - \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2 \sqrt{d - c^2 dx^2}}{5\sqrt{cx - 1} \sqrt{cx + 1}} \left( \frac{1}{5} c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) \right) - \frac{bc\sqrt{c^2 x^2 - 1}}{5} \left( \frac{6(c^2 x^2 - 1)^{5/2}}{5} \right) \right) \right)$$

↓ 3011

$$d \left( d \left( - \frac{\sqrt{d - c^2 dx^2} (2ib(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) dx)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right. \right. \\ \left. \left. - \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2 \sqrt{d - c^2 dx^2}}{5\sqrt{cx - 1} \sqrt{cx + 1}} \left( \frac{1}{5} c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) \right) - \frac{bc\sqrt{c^2 x^2 - 1}}{5} \left( \frac{6(c^2 x^2 - 1)^{5/2}}{5} \right) \right) \right)$$

↓ 2720

$$d \left( d \left( - \frac{\sqrt{d - c^2 dx^2} (2ib(b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) dx)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right. \right. \\ \left. \left. - \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \frac{2bcd^2 \sqrt{d - c^2 dx^2}}{5\sqrt{cx - 1} \sqrt{cx + 1}} \left( \frac{1}{5} c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) \right) - \frac{bc\sqrt{c^2 x^2 - 1}}{5} \left( \frac{6(c^2 x^2 - 1)^{5/2}}{5} \right) \right) \right)$$

↓ 7143

---

3.190.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$

$$d \left( d \left( -\frac{\sqrt{d - c^2 dx^2} (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx))^2 + 2ib(b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(2, -\sqrt{ca})))}{\sqrt{ca}} \right. \right. \\ \left. \left. + \frac{1}{5} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 - \right. \right. \\ \left. \left. 2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{5} c^4 x^5 (a + \operatorname{barccosh}(cx)) - \frac{2}{3} c^2 x^3 (a + \operatorname{barccosh}(cx)) + x(a + \operatorname{barccosh}(cx)) \right) - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{6(c^2 x^2 - 1)^{5/2}}{5} \right)}{5\sqrt{cx - 1}\sqrt{cx + 1}} \right) \right)$$

```
input Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x,x]
```

```
output ((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/5 - (2*b*c*d^2*sqrt[d - c^2
*d*x^2]*(-1/30*(b*c*sqrt[-1 + c^2*x^2]*((16*sqrt[-1 + c^2*x^2])/c^2 - (8*(
-1 + c^2*x^2)^(3/2))/(3*c^2) + (6*(-1 + c^2*x^2)^(5/2))/(5*c^2)))/(sqrt[-1
+ c*x]*sqrt[1 + c*x]) + x*(a + b*ArcCosh[c*x]) - (2*c^2*x^3*(a + b*ArcCos
h[c*x]))/3 + (c^4*x^5*(a + b*ArcCosh[c*x]))/5)/(5*sqrt[-1 + c*x]*sqrt[1 +
c*x]) + d*(((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/3 - (2*b*c*d*sq
rt[d - c^2*d*x^2]*(-1/3*(b*c*((7*sqrt[-1 + c*x]*sqrt[1 + c*x])/(3*c^2) - (
x^2*sqrt[-1 + c*x]*sqrt[1 + c*x])/3)) + x*(a + b*ArcCosh[c*x]) - (c^2*x^3*
(a + b*ArcCosh[c*x]))/3))/(3*sqrt[-1 + c*x]*sqrt[1 + c*x]) + d*(sqrt[d - c
^2*d*x^2]*(a + b*ArcCosh[c*x])^2 - (2*b*c*sqrt[d - c^2*d*x^2]*(a*x - (b*sq
rt[-1 + c*x]*sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(sqrt[-1 + c*x]*sqrt[1
+ c*x]) - (sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[
c*x]] + (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) +
b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-((a + b*ArcCosh[c*x])*Poly
Log[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]])))/(sqrt[-1 + c
*x]*sqrt[1 + c*x]))))
```

3.190.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

---

3.190.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$

- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 960 `Int[((e_.)*(x_)^(m_.))*((a1_.) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_.) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*(a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1)), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 1140 `Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 0]`
- rule 1576 `Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`
- rule 1905 `Int[((f_.)*(x_)^(m_.))*((d1_.) + (e1_.)*(x_)^(non2_.))^(q_.)*((d2_.) + (e2_.)*(x_)^(non2_.))^(q_.)*((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.))^(p_.), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.) * (x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_) ^ (m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_)^(p_.)*(d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^(m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6362 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.190.4 Maple [F]

$$\int \frac{(-c^2 d x^2 + d)^{\frac{5}{2}} (a + b \operatorname{arccosh}(cx))^2}{x} dx$$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x,x)`

output `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x,x)`

**3.190.5 Fracas [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x,x, algorithm="fracas")`

output `integral((a^2*c^4*d^2*x^4 - 2*a^2*c^2*d^2*x^2 + a^2*d^2 + (b^2*c^4*d^2*x^4 - 2*b^2*c^2*d^2*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*c^4*d^2*x^4 - 2*a*b*c^2*d^2*x^2 + a*b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x, x)`

**3.190.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2/x,x)`

output `Timed out`

**3.190.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x,x, algorithm="maxima")`

output `-1/15*(15*d^(5/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - 3*(-c^2*d*x^2 + d)^(5/2) - 5*(-c^2*d*x^2 + d)^(3/2)*d - 15*sqrt(-c^2*d*x^2 + d)*d^2)*a^2 + integrate((-c^2*d*x^2 + d)^(5/2)*b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/x + 2*(-c^2*d*x^2 + d)^(5/2)*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

---

3.190.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx$

**3.190.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.190.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2}}{x} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x, x)`



**3.191** 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx$$

3.191.1 Optimal result . . . . .	1648
3.191.2 Mathematica [A] (warning: unable to verify) . . . . .	1649
3.191.3 Rubi [C] (warning: unable to verify) . . . . .	1650
3.191.4 Maple [A] (verified) . . . . .	1662
3.191.5 Fricas [F] . . . . .	1663
3.191.6 Sympy [F(-1)] . . . . .	1663
3.191.7 Maxima [F] . . . . .	1664
3.191.8 Giac [F(-2)] . . . . .	1664
3.191.9 Mupad [F(-1)] . . . . .	1664

**3.191.1 Optimal result**

Integrand size = 29, antiderivative size = 607

$$\begin{aligned} &\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx = \\ &-\frac{31}{64}b^2c^2d^2x\sqrt{d-c^2dx^2} - \frac{1}{32}b^2c^2d^2x(1-cx)(1+cx)\sqrt{d-c^2dx^2} \\ &-\frac{89b^2cd^2\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{64\sqrt{-1+cx}\sqrt{1+cx}} + \frac{15bc^3d^2x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{8\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{bcd^2(1-c^2x^2)\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &-\frac{bcd^2(1-c^2x^2)^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{8\sqrt{-1+cx}\sqrt{1+cx}} \\ &-\frac{15}{8}c^2d^2x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 + \frac{cd^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &-\frac{5}{4}c^2dx(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2 \\ &-\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x} + \frac{5cd^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^3}{8b\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{2bcd^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\log(1+e^{-2\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ &-\frac{b^2cd^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-e^{-2\operatorname{arccosh}(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output

$$\begin{aligned}
& -5/4*c^2*d*x*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^{-2}-(-c^2*d*x^2+d)^{(5/2)} \\
& *(a+b*\operatorname{arccosh}(c*x))^{-2}/x-31/64*b^2*c^2*d^2*x*(-c^2*d*x^2+d)^{(1/2)}-1/32*b^2 \\
& *c^2*d^2*x*(-c*x+1)*(c*x+1)*(-c^2*d*x^2+d)^{(1/2)}-15/8*c^2*d^2*x*(a+b*\operatorname{arcco} \\
& \operatorname{sh}(c*x))^{-2}*(-c^2*d*x^2+d)^{(1/2)}-89/64*b^2*c*d^2*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d) \\
& ^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+15/8*b*c^3*d^2*x^2*(a+b*\operatorname{arccosh}(c*x))* \\
& (-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+b*c*d^2*(-c^2*x^2+1)*(a+b \\
& *\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/8*b*c*d^ \\
& 2*(-c^2*x^2+1)^2*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c* \\
& x+1)^{(1/2)}+c*d^2*(a+b*\operatorname{arccosh}(c*x))^{-2}*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/( \\
& c*x+1)^{(1/2)}+5/8*c*d^2*(a+b*\operatorname{arccosh}(c*x))^{-3}*(-c^2*d*x^2+d)^{(1/2)}/b/(c*x-1) \\
& ^{(1/2)}/(c*x+1)^{(1/2)}+2*b*c*d^2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+1/(c*x+(c*x-1)^{(1/2)} \\
& )*(c*x+1)^{(1/2)})^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b^2*c \\
& *d^2*\operatorname{polylog}(2,-1/(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*(-c^2*d*x^2+d)^{(1/2)} \\
& /((c*x-1)^{(1/2)}/(c*x+1)^{(1/2)})
\end{aligned}$$

### 3.191.2 Mathematica [A] (warning: unable to verify)

Time = 5.26 (sec) , antiderivative size = 554, normalized size of antiderivative = 0.91

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2}{x^2} dx = \frac{d^2 \left( 96a^2 \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \sqrt{d - c^2 dx^2} (-8 - 9c^2 x^2 + 2c^4 x^4) + 144 \right)}{x^2}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x^2,x]`

output

$$\begin{aligned}
& (d^2*(96*a^2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{Sqrt}[d - c^2*d*x^2]*(-8 \\
& - 9*c^2*x^2 + 2*c^4*x^4) + 1440*a^2*c*\operatorname{Sqrt}[d]*x*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] \\
& *(1 + c*x)*\operatorname{ArcTan}[c*x*\operatorname{Sqrt}[d - c^2*d*x^2]]/(\operatorname{Sqrt}[d]*(-1 + c^2*x^2))) - 76 \\
& 8*a*b*\operatorname{Sqrt}[d - c^2*d*x^2]*(2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[ \\
& c*x] - c*x*(\operatorname{ArcCosh}[c*x]^2 + 2*\operatorname{Log}[c*x])) - 256*b^2*\operatorname{Sqrt}[d - c^2*d*x^2]*(A \\
& rcCosh[c*x]*(3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x] - c*x*(Ar \\
& cCosh[c*x]*(3 + \operatorname{ArcCosh}[c*x]) + 6*\operatorname{Log}[1 + E^(-2*\operatorname{ArcCosh}[c*x])])) + 3*c*x*P \\
& olyLog[2, -E^(-2*\operatorname{ArcCosh}[c*x])] + 384*a*b*c*x*\operatorname{Sqrt}[d - c^2*d*x^2]*(\operatorname{Cosh}[2 \\
& *\operatorname{ArcCosh}[c*x]] + 2*\operatorname{ArcCosh}[c*x]*(\operatorname{ArcCosh}[c*x] - \operatorname{Sinh}[2*\operatorname{ArcCosh}[c*x]])) + 6 \\
& 4*b^2*c*x*\operatorname{Sqrt}[d - c^2*d*x^2]*(4*\operatorname{ArcCosh}[c*x]^3 + 6*\operatorname{ArcCosh}[c*x]*\operatorname{Cosh}[2*Ar \\
& cCosh[c*x]] - 3*(1 + 2*\operatorname{ArcCosh}[c*x]^2)*\operatorname{Sinh}[2*\operatorname{ArcCosh}[c*x]]) - 12*a*b*c*x* \\
& \operatorname{Sqrt}[d - c^2*d*x^2]*(8*\operatorname{ArcCosh}[c*x]^2 + \operatorname{Cosh}[4*\operatorname{ArcCosh}[c*x]] - 4*\operatorname{ArcCosh}[c \\
& *x]*\operatorname{Sinh}[4*\operatorname{ArcCosh}[c*x]]) - b^2*c*x*\operatorname{Sqrt}[d - c^2*d*x^2]*(32*\operatorname{ArcCosh}[c*x]^3 \\
& + 12*\operatorname{ArcCosh}[c*x]*\operatorname{Cosh}[4*\operatorname{ArcCosh}[c*x]] - 3*(1 + 8*\operatorname{ArcCosh}[c*x]^2)*\operatorname{Sinh}[4* \\
& \operatorname{ArcCosh}[c*x]])))/(768*x*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x))
\end{aligned}$$

---

3.191.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2}{x^2} dx$

**3.191.3 Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 4.33 (sec) , antiderivative size = 663, normalized size of antiderivative = 1.09, number of steps used = 29, number of rules used = 28,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.966$ , Rules used = {6343, 6312, 25, 6310, 6298, 101, 43, 6308, 6327, 6329, 40, 40, 43, 6334, 40, 40, 43, 6334, 40, 43, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx \\
 & \quad \downarrow \text{6343} \\
 & \frac{2bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2 (a + \operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1} \sqrt{cx+1}} - 5c^2 d \int (d - c^2 dx^2)^{3/2} (a + \\
 & \operatorname{barccosh}(cx))^2 dx - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} \\
 & \quad \downarrow \text{6312} \\
 & \frac{2bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2 (a + \operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1} \sqrt{cx+1}} - \\
 & 5c^2 d \left( \frac{bcd \sqrt{d - c^2 dx^2} \int -x(1-cx)(cx+1)(a + \operatorname{barccosh}(cx)) dx}{2\sqrt{cx-1} \sqrt{cx+1}} + \frac{3}{4} d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{4} \int \right. \\
 & \left. \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{2bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2 (a + \operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1} \sqrt{cx+1}} - \\
 & 5c^2 d \left( - \frac{bcd \sqrt{d - c^2 dx^2} \int x(1-cx)(cx+1)(a + \operatorname{barccosh}(cx)) dx}{2\sqrt{cx-1} \sqrt{cx+1}} + \frac{3}{4} d \int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{1}{4} \int \right. \\
 & \left. \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x} \right) \\
 & \quad \downarrow \text{6310}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} dx - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \int x(a+\operatorname{barccosh}(cx))dx}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
 & \left. \left. \frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{x} \right) \right) \downarrow \mathbf{6298} \\
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} dx - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
 & \left. \left. \frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{x} \right) \right) \downarrow \mathbf{101} \\
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} dx - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
 & \left. \left. \frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{x} \right) \right) \downarrow \mathbf{43} \\
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} dx - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2} \right. \right. \\
 & \left. \left. \frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{x} \right) \right) \downarrow \mathbf{6308}
 \end{aligned}$$

---

3.191.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx$

$$\begin{aligned}
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 + \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int x(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
 & \quad \left. \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} \right) \right. \\
 & \quad \quad \quad \downarrow \text{6327} \\
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \int x(1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 + \frac{3}{4}d \left( -\frac{\sqrt{d-c^2dx^2} \int x(1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
 & \quad \left. \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} \right) \right) \\
 & \quad \quad \quad \downarrow \text{6329} \\
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b \int (cx-1)^{3/2}(cx+1)^{3/2} dx}{4c} - \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \right. \\
 & \quad \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} \right) \\
 & \quad \quad \quad \downarrow \text{40} \\
 & \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \\
 5c^2d & \left( -\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4} \int \sqrt{cx-1}\sqrt{cx+1} dx)}{4c} - \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2 \right. \\
 & \quad \left. \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} \right) \\
 & \quad \quad \quad \downarrow \text{40}
 \end{aligned}$$

---

3.191.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx$

$$5c^2d \left( \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx\right)\right)}{4c} - \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{4c^2} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

↓ 43

$$5c^2d \left( \frac{2bcd^2\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x} - \frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{4} \right)$$

↓ 6334

$$\frac{2bcd^2\sqrt{d-c^2dx^2} \left( \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx - \frac{1}{4}bc \int (cx-1)^{3/2}(cx+1)^{3/2} dx + \frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}}$$

$$\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x}$$

$$5c^2d \left( \frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{4} \right)$$

↓ 40

---

3.191.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^2} dx$

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx-\frac{1}{4}bc\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\int\sqrt{cx-1}\sqrt{cx+1}dx\right)+\frac{1}{4}(1-\right.}{\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x}}{\frac{bcd\sqrt{d-c^2dx^2}\left(\frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c}}{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2-\frac{2\sqrt{cx-1}\sqrt{cx+1}}{2\sqrt{cx-1}\sqrt{cx+1}}}}{5c^2d}\right)$$

↓ 40

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx-\frac{1}{4}bc\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\sqrt{cx-1}\sqrt{cx+1}dx\right)\right)+\frac{1}{4}(1-\right.}{\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x}}{\frac{bcd\sqrt{d-c^2dx^2}\left(\frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c}}{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2-\frac{2\sqrt{cx-1}\sqrt{cx+1}}{2\sqrt{cx-1}\sqrt{cx+1}}}}{5c^2d}\right)$$

↓ 43

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx+\frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))-\frac{1}{4}bc\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\int\sqrt{cx-1}\sqrt{cx+1}dx\right)\right)+\frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x}}{\frac{bcd\sqrt{d-c^2dx^2}\left(\frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c}}{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2-\frac{2\sqrt{cx-1}\sqrt{cx+1}}{2\sqrt{cx-1}\sqrt{cx+1}}}}{5c^2d}\right)$$

↓ 6334

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{2}bc\int\sqrt{cx-1}\sqrt{cx+1}dx+\frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))+\frac{1}{2}(1-c^2x^2)\sqrt{cx-1}\sqrt{cx+1}\right)}{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2-\frac{x}{bcd\sqrt{d-c^2dx^2}}\left(\frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c}\right)}-\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2-\frac{2\sqrt{cx-1}\sqrt{cx+1}}{2\sqrt{cx-1}\sqrt{cx+1}}}$$

↓ 40

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{2}bc\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)+\frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))+\frac{1}{2}(1-c^2x^2)\sqrt{cx-1}\sqrt{cx+1}\right)}{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2-\frac{x}{bcd\sqrt{d-c^2dx^2}}\left(\frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c}\right)}-\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2-\frac{2\sqrt{cx-1}\sqrt{cx+1}}{2\sqrt{cx-1}\sqrt{cx+1}}}$$

↓ 43

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{4}(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))+\frac{1}{2}bc\sqrt{cx-1}\sqrt{cx+1}\right)}{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2-\frac{x}{bcd\sqrt{d-c^2dx^2}}\left(\frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2}-\frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c}\right)}-\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2-\frac{2\sqrt{cx-1}\sqrt{cx+1}}{2\sqrt{cx-1}\sqrt{cx+1}}}$$

↓ 6297

---

3.191.  $\int\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx$



$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{\int -\left((a+b\operatorname{arccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)\right) d(a+b\operatorname{arccosh}(cx))}{b} + \frac{1}{4}(1-c^2x^2)^2(a+b\operatorname{arccosh}(cx)) \right)$$

$$5c^2d \left( \frac{\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x}}{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)} - \frac{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 25

$$2bcd^2\sqrt{d-c^2dx^2} \left( -\frac{\int (a+b\operatorname{arccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) d(a+b\operatorname{arccosh}(cx))}{b} + \frac{1}{4}(1-c^2x^2)^2(a+b\operatorname{arccosh}(cx)) \right)$$

$$5c^2d \left( \frac{\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x}}{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)} - \frac{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 3042

$$2bcd^2\sqrt{d-c^2dx^2} \left( -\frac{\int -i(a+b\operatorname{arccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right) d(a+b\operatorname{arccosh}(cx))}{b} + \frac{1}{4}(1-c^2x^2)^2(a+b\operatorname{arccosh}(cx)) \right)$$

$$5c^2d \left( \frac{\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x}}{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)} - \frac{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{2\sqrt{cx-1}\sqrt{cx+1}} \right)$$

↓ 26

---

3.191.  $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx$

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{i \int (a+b\operatorname{arccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right) d(a+b\operatorname{arccosh}(cx))}{b} + \frac{1}{4}(1-c^2x^2)^2(a+b\operatorname{arccosh}(cx)) \right)$$

$$5c^2d \left( \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x} - \frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2 \right)$$

↓ 4201

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{i\left(2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))}{1+e^{-2\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}i(a+b\operatorname{arccosh}(cx))^2\right)}{b} + \frac{1}{4}(1-c^2x^2)^2(a+b\operatorname{arccosh}(cx)) \right)$$

$$5c^2d \left( \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x} - \frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2 \right)$$

↓ 2620

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{i\left(2i\left(\frac{1}{2}b \int \log(1+e^{-2\operatorname{arccosh}(cx)}) d(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1)(a+b\operatorname{arccosh}(cx))\right) - \frac{1}{2}i(a+b\operatorname{arccosh}(cx))^2\right)}{b} + \frac{1}{4}(1-c^2x^2)^2(a+b\operatorname{arccosh}(cx)) \right)$$

$$5c^2d \left( \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x} - \frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)\right)}{4c} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}x(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2 \right)$$

↓ 2715

---

3.191.  $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx$

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{i\left(2i\left(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log\left(1+e^{-2\operatorname{arccosh}(cx)}\right) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log\left(e^{-2\operatorname{arccosh}(cx)}+1\right)\right)(a+\operatorname{arccosh}(cx))\right) - \frac{1}{2}}{b} \right)$$

$$5c^2d \left( \frac{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{arccosh}(cx))^2 - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{arccosh}(cx))^2}{x}}{\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)}{4c} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right)} \right)$$

↓ 2838

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{i\left(2i\left(\frac{1}{4}b^2 \operatorname{PolyLog}(2,-a-\operatorname{arccosh}(cx)) - \frac{1}{2}b \log\left(e^{-2\operatorname{arccosh}(cx)}+1\right)\right)(a+\operatorname{arccosh}(cx))\right) - \frac{1}{2}i(a+\operatorname{arccosh}(cx))^2}{b} \right) +$$

$$5c^2d \left( \frac{\frac{1}{4}x(d-c^2dx^2)^{3/2}(a+\operatorname{arccosh}(cx))^2 - \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{arccosh}(cx))^2}{x}}{\frac{bcd\sqrt{d-c^2dx^2} \left( \frac{b\left(\frac{1}{4}x(cx-1)^{3/2}(cx+1)^{3/2} - \frac{3}{4}\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{\operatorname{arccosh}(cx)}{2c}\right)}{4c} \right)}{2\sqrt{cx-1}\sqrt{cx+1}} \right)} \right)$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x^2,x]`

```

output -(((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x - 5*c^2*d*((x*(d - c^2
*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/4 + (3*d*((x*Sqrt[d - c^2*d*x^2]*(a
+ b*ArcCosh[c*x])^2)/2 - (Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^3)/(6*b
*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (b*c*Sqrt[d - c^2*d*x^2]*((x^2*(a + b*A
rcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh
[c*x]/(2*c^3)))/2))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/4 - (b*c*d*Sqrt[d - c
^2*d*x^2]*(-1/4*((1 - c^2*x^2)^2*(a + b*ArcCosh[c*x]))/c^2 + (b*((x*(-1 +
c*x)^(3/2)*(1 + c*x)^(3/2))/4 - (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - A
rcCosh[c*x]/(2*c)))/4))/(4*c)))/(2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (2*b*c
*d^2*Sqrt[d - c^2*d*x^2]*(((1 - c^2*x^2)*(a + b*ArcCosh[c*x]))/2 + ((1 - c
^2*x^2)^2*(a + b*ArcCosh[c*x]))/4 + (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])
/2 - ArcCosh[c*x]/(2*c)))/2 - (b*c*((x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/4
- (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/2 - ArcCosh[c*x]/(2*c)))/4))/4 + (
I*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*L
og[1 + E^(-2*ArcCosh[c*x])]) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4)))/
b))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])

```

### 3.191.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]

```

```

rule 40 Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*
(a + b*x)^m*((c + d*x)^m/(2*m + 1)), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a
+ b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[
b*c + a*d, 0] && IGtQ[m + 1/2, 0]

```

```

rule 43 Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[
ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a
*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]

```

rule 101 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)*(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)n(e + f*x)pSimp[a2d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 2620 `Int[((F_)(g_.)*(e_.) + (f_.)*(x_))(n_.)((c_.) + (d_.)*(x_))(m_.))/((a_.) + (b_.)*(F_)(g_.)*(e_.) + (f_.)*(x_))(n_.)), x_Symbol] := Simp[((c + d*x)m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)(m - 1)*Log[1 + b*((F^(g*(e + f*x)))n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)(e_.)*((c_.) + (d_.)*(x_))(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*xn/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_))(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)m(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[xn*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +
c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol]
:> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol]
:> Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^(n/2)), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol]
:> Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol]
:> Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 6334 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.)/(x_), x_Symbol] := Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d)^(p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]`

### 3.191.4 Maple [A] (verified)

Time = 1.24 (sec) , antiderivative size = 589, normalized size of antiderivative = 0.97

method	result
default	$-\frac{a^2(-c^2dx^2+d)^{\frac{7}{2}}}{dx} - a^2c^2x(-c^2dx^2+d)^{\frac{5}{2}} - \frac{5a^2c^2dx(-c^2dx^2+d)^{\frac{3}{2}}}{4} - \frac{15a^2c^2d^2x\sqrt{-c^2dx^2+d}}{8} - \frac{15a^2c^2d^3\arctan\left(\frac{cx}{\sqrt{-c^2dx^2+d}}\right)}{8\sqrt{c^2}}$
parts	$-\frac{a^2(-c^2dx^2+d)^{\frac{7}{2}}}{dx} - a^2c^2x(-c^2dx^2+d)^{\frac{5}{2}} - \frac{5a^2c^2dx(-c^2dx^2+d)^{\frac{3}{2}}}{4} - \frac{15a^2c^2d^2x\sqrt{-c^2dx^2+d}}{8} - \frac{15a^2c^2d^3\arctan\left(\frac{cx}{\sqrt{-c^2dx^2+d}}\right)}{8\sqrt{c^2}}$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^2,x,method=_RETURNVERBOSE)`

---

3.191. 
$$\int \frac{(d-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^2} dx$$

output

```
-a^2/d/x*(-c^2*d*x^2+d)^(7/2)-a^2*c^2*x*(-c^2*d*x^2+d)^(5/2)-5/4*a^2*c^2*d
*x*(-c^2*d*x^2+d)^(3/2)-15/8*a^2*c^2*d^2*x*(-c^2*d*x^2+d)^(1/2)-15/8*a^2*c
^2*d^3/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+1/64*b^2
*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x*(16*(c*x+1)^(1/2)*(c
*x-1)^(1/2)*arccosh(c*x)^2*x^4*c^4-8*arccosh(c*x)*c^5*x^5+2*(c*x+1)^(1/2)*
(c*x-1)^(1/2)*c^4*x^4-72*arccosh(c*x)^2*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^2*c^
2+72*c^3*x^3*arccosh(c*x)-33*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+40*arccos
h(c*x)^3*x*c-64*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2-64*arccosh(c*x)
^2*x*c+128*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-33*c
*x*arccosh(c*x)+64*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c)*d^
2+1/64*a*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x*(32*(c*x-1)
)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4-8*c^5*x^5-144*(c*x+1)^(1/2)*arc
cosh(c*x)*(c*x-1)^(1/2)*c^2*x^2+72*c^3*x^3+120*arccosh(c*x)^2*x*c-128*arcc
osh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-128*c*x*arccosh(c*x)+128*ln(1+(c*x+(c
*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-33*c*x)*d^2
```

### 3.191.5 Fracas [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2}{x^2} dx$$

input

```
integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^2,x, algorithm="fric
as")
```

output

```
integral((a^2*c^4*d^2*x^4 - 2*a^2*c^2*d^2*x^2 + a^2*d^2 + (b^2*c^4*d^2*x^4
- 2*b^2*c^2*d^2*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*c^4*d^2*x^4 - 2*a*
b*c^2*d^2*x^2 + a*b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^2, x)
```

### 3.191.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \text{Timed out}$$

input

```
integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2/x**2,x)
```

output

```
Timed out
```

---

3.191.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx$



**3.191.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^2,x, algorithm="maxima")`

output `-1/8*(10*(-c^2*d*x^2 + d)^(3/2)*c^2*d*x + 15*sqrt(-c^2*d*x^2 + d)*c^2*d^2*x + 15*c*d^(5/2)*arcsin(c*x) + 8*(-c^2*d*x^2 + d)^(5/2)/x)*a^2 + integrate((-c^2*d*x^2 + d)^(5/2)*b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/x^2 + 2*(-c^2*d*x^2 + d)^(5/2)*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x^2, x)`

**3.191.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.191.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x^2,x)`

output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x^2, x)`

---

3.191.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx$

$$3.192 \quad \int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx$$

3.192.1 Optimal result . . . . .	1666
3.192.2 Mathematica [B] (warning: unable to verify) . . . . .	1667
3.192.3 Rubi [A] (warning: unable to verify) . . . . .	1668
3.192.4 Maple [F] . . . . .	1678
3.192.5 Fricas [F] . . . . .	1678
3.192.6 Sympy [F(-1)] . . . . .	1679
3.192.7 Maxima [F] . . . . .	1679
3.192.8 Giac [F(-2)] . . . . .	1679
3.192.9 Mupad [F(-1)] . . . . .	1680

### 3.192.1 Optimal result

Integrand size = 29, antiderivative size = 890

$$\begin{aligned}
& \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = -\frac{170}{27} b^2 c^2 d^2 \sqrt{d - c^2 dx^2} \\
& + \frac{5}{27} b^2 c^4 d^2 x^2 \sqrt{d - c^2 dx^2} + \frac{5abc^3 d^2 x \sqrt{d - c^2 dx^2}}{\sqrt{-1 + cx} \sqrt{1 + cx}} \\
& + \frac{5b^2 c^2 d^2 (1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{3(1 - cx)(1 + cx)} + \frac{b^2 c^2 d^2 (1 - c^2 x^2)^2 \sqrt{d - c^2 dx^2}}{9(1 - cx)(1 + cx)} \\
& + \frac{5b^2 c^3 d^2 x \sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{\sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{bcd^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{x \sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{bc^3 d^2 x \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{3\sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{2bc^5 d^2 x^3 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{9\sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{5}{2} c^2 d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 - \frac{5}{6} c^2 d (d \\
& - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
& + \frac{5c^2 d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{b^2 c^2 d^2 \sqrt{-1 + c^2 x^2} \sqrt{d - c^2 dx^2} \arctan(\sqrt{-1 + c^2 x^2})}{(1 - cx)(1 + cx)} \\
& - \frac{5ibc^2 d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx} \sqrt{1 + cx}} \\
& + \frac{5ibc^2 d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx} \sqrt{1 + cx}} \\
& + \frac{5ib^2 c^2 d^2 \sqrt{d - c^2 dx^2} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{5ib^2 c^2 d^2 \sqrt{d - c^2 dx^2} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{\sqrt{-1 + cx} \sqrt{1 + cx}}
\end{aligned}$$

output

```

-5/6*c^2*d*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2-1/2*(-c^2*d*x^2+d)^(5
/2)*(a+b*arccosh(c*x))^2/x^2-170/27*b^2*c^2*d^2*(-c^2*d*x^2+d)^(1/2)+5/27*
b^2*c^4*d^2*x^2*(-c^2*d*x^2+d)^(1/2)+5/3*b^2*c^2*d^2*(-c^2*x^2+1)*(-c^2*d*
x^2+d)^(1/2)/(-c*x+1)/(c*x+1)+1/9*b^2*c^2*d^2*(-c^2*x^2+1)^2*(-c^2*d*x^2+d
)^(1/2)/(-c*x+1)/(c*x+1)-5/2*c^2*d^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(
1/2)+5*a*b*c^3*d^2*x*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5*b^
2*c^3*d^2*x*arccosh(c*x)*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-
b*c*d^2*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/x/(c*x-1)^(1/2)/(c*x+1)^(1
/2)-1/3*b*c^3*d^2*x*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/
(c*x+1)^(1/2)-2/9*b*c^5*d^2*x^3*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/(c
*x-1)^(1/2)/(c*x+1)^(1/2)+5*c^2*d^2*(a+b*arccosh(c*x))^2*arctan(c*x+(c*x-1
)^(1/2)*(c*x+1)^(1/2))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5*
I*b^2*c^2*d^2*polylog(3,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+
d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-5*I*b*c^2*d^2*(a+b*arccosh(c*x))*poly
log(2,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(
1/2)/(c*x+1)^(1/2)+5*I*b*c^2*d^2*(a+b*arccosh(c*x))*polylog(2,I*(c*x+(c*x-
1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-
5*I*b^2*c^2*d^2*polylog(3,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(-c^2*d*x^2
+d)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b^2*c^2*d^2*arctan((c^2*x^2-1)^(1/2)
)*(-c^2*x^2-1)^(1/2)*(-c^2*d*x^2+d)^(1/2)/(-c*x+1)/(c*x+1)

```

### 3.192.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 5694 vs.  $2(890) = 1780$ .

Time = 65.18 (sec) , antiderivative size = 5694, normalized size of antiderivative = 6.40

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2}{x^3} dx = \text{Result too large to show}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x^3,x]`

output `Result too large to show`

**3.192.3 Rubi [A] (warning: unable to verify)**

Time = 4.56 (sec) , antiderivative size = 609, normalized size of antiderivative = 0.68, number of steps used = 25, number of rules used = 24,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.828$ , Rules used = {6343, 6327, 6336, 27, 1905, 1578, 1192, 1467, 2009, 6345, 25, 6304, 6309, 27, 960, 83, 6341, 2009, 6362, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx \\
 & \quad \downarrow \text{6343} \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx}{\sqrt{cx-1} \sqrt{cx+1}} - \\
 & \frac{5}{2} c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx - \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{6327} \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx}{\sqrt{cx-1} \sqrt{cx+1}} - \frac{5}{2} c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx - \\
 & \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{6336} \\
 & -\frac{5}{2} c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( -bc \int -\frac{-c^4 x^4 + 6c^2 x^2 + 3}{3x \sqrt{cx-1} \sqrt{cx+1}} dx + \frac{1}{3} c^4 x^3 (a + \operatorname{barccosh}(cx)) - 2c^2 x (a + \operatorname{barccosh}(cx)) - \frac{a + \operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1} \sqrt{cx+1}} \\
 & \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{27} \\
 & -\frac{5}{2} c^2 d \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
 & \frac{bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{3} bc \int \frac{-c^4 x^4 + 6c^2 x^2 + 3}{x \sqrt{cx-1} \sqrt{cx+1}} dx + \frac{1}{3} c^4 x^3 (a + \operatorname{barccosh}(cx)) - 2c^2 x (a + \operatorname{barccosh}(cx)) - \frac{a + \operatorname{barccosh}(cx)}{x} \right)}{\sqrt{cx-1} \sqrt{cx+1}} \\
 & \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \\
 & \quad \downarrow \text{1905}
 \end{aligned}$$

---

3.192.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$

$$\begin{aligned}
 & -\frac{5}{2}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
 bcd^2\sqrt{d - c^2dx^2} & \left( \frac{bc\sqrt{c^2x^2-1} \int \frac{-c^4x^4+6c^2x^2+3}{x\sqrt{c^2x^2-1}} dx}{3\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} \right) \\
 & \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} \\
 & \frac{1}{2x^2} \\
 & \downarrow \text{1578} \\
 & -\frac{5}{2}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
 bcd^2\sqrt{d - c^2dx^2} & \left( \frac{bc\sqrt{c^2x^2-1} \int \frac{-c^4x^4+6c^2x^2+3}{x^2\sqrt{c^2x^2-1}} dx^2}{6\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} \right) \\
 & \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} \\
 & \frac{1}{2x^2} \\
 & \downarrow \text{1192} \\
 & -\frac{5}{2}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
 bcd^2\sqrt{d - c^2dx^2} & \left( \frac{b\sqrt{c^2x^2-1} \int \frac{-c^4x^8+4c^4x^4+8e^4d\sqrt{c^2x^2-1}}{x^4+1} dx}{3c^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} \right) \\
 & \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} \\
 & \frac{1}{2x^2} \\
 & \downarrow \text{1467} \\
 & -\frac{5}{2}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \\
 bcd^2\sqrt{d - c^2dx^2} & \left( \frac{b\sqrt{c^2x^2-1} \int \left(-x^4c^4 + \frac{3c^4}{x^4+1} + 5c^4\right) d\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} \right) \\
 & \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} \\
 & \frac{1}{2x^2} \\
 & \downarrow \text{2009}
 \end{aligned}$$

---

3.192.  $\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$

$$-\frac{5}{2}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x} dx +$$

$$bcd^2\sqrt{d - c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a + \operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2 - 1} (3c^4 \arctan(\sqrt{c^2x^2 - 1}))}{3c^3\sqrt{cx - 1}} \right)$$


---


$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 6345

$$-\frac{5}{2}c^2d \left( \frac{2bcd\sqrt{d - c^2dx^2} \int -((1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx)))dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + d \int \frac{\sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \right.$$

$$bcd^2\sqrt{d - c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a + \operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2 - 1} (3c^4 \arctan(\sqrt{c^2x^2 - 1}))}{3c^3\sqrt{cx - 1}} \right)$$


---


$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 25

$$-\frac{5}{2}c^2d \left( -\frac{2bcd\sqrt{d - c^2dx^2} \int (1 - cx)(cx + 1)(a + \operatorname{barccosh}(cx))dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + d \int \frac{\sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3} \right.$$

$$bcd^2\sqrt{d - c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a + \operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2 - 1} (3c^4 \arctan(\sqrt{c^2x^2 - 1}))}{3c^3\sqrt{cx - 1}} \right)$$


---


$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 6304

$$-\frac{5}{2}c^2d \left( -\frac{2bcd\sqrt{d - c^2dx^2} \int (1 - c^2x^2) (a + \operatorname{barccosh}(cx))dx}{3\sqrt{cx - 1}\sqrt{cx + 1}} + d \int \frac{\sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3} (d - \right.$$

$$bcd^2\sqrt{d - c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a + \operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2 - 1} (3c^4 \arctan(\sqrt{c^2x^2 - 1}))}{3c^3\sqrt{cx - 1}} \right)$$


---


$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{2x^2} \sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 6309

---

3.192.  $\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$

$$\begin{aligned}
 & -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx - \frac{2bcd\sqrt{d-c^2dx^2} \left( -bc \int \frac{x(3-c^2x^2)}{3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1} \left( 3c^4 \arctan\left(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}\right) \right)}{3c^3\sqrt{cx-1}} \right) \right) \right. \\
 & \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{27}
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx - \frac{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}bc \int \frac{x(3-c^2x^2)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}c^2x^3(a+\operatorname{barccosh}(cx)) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1} \left( 3c^4 \arctan\left(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}\right) \right)}{3c^3\sqrt{cx-1}} \right) \right) \right. \\
 & \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{960}
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx - \frac{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}bc \left( \frac{7}{3} \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{3}x^2\sqrt{cx-1}\sqrt{cx+1} \right) \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1} \left( 3c^4 \arctan\left(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}\right) \right)}{3c^3\sqrt{cx-1}} \right) \right) \right. \\
 & \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{83}
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{5}{2}c^2d \left( d \int \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{x} dx + \frac{1}{3}(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 - \frac{2bcd\sqrt{d-c^2dx^2} \left( -\frac{1}{3}c^2x \right)}{3\sqrt{cx-1}\sqrt{cx+1}} \right. \\
 & \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1} \left( 3c^4 \arctan\left(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}\right) \right)}{3c^3\sqrt{cx-1}} \right) \right) \right. \\
 & \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2} \right) \\
 & \qquad \qquad \qquad \downarrow \text{6341}
 \end{aligned}$$

---

3.192.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^3} dx$



$$-\frac{5}{2}c^2d \left( d \left( -\frac{2bc\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx))dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx)) \right) \right. \\ \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4\arctan(\sqrt{c^2x^2-1}))}{3c^3\sqrt{cx-1}} \right) \right) \\ \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} \\ 2x^2$$

↓ 2009

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right. \\ \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4\arctan(\sqrt{c^2x^2-1}))}{3c^3\sqrt{cx-1}} \right) \right) \\ \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} \\ 2x^2$$

↓ 6362

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 - \frac{2bc\sqrt{d-c^2dx^2}(ax+b\operatorname{arccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right. \\ \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4\arctan(\sqrt{c^2x^2-1}))}{3c^3\sqrt{cx-1}} \right) \right) \\ \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} \\ 2x^2$$

↓ 3042

$$-\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2} \int (a+\operatorname{barccosh}(cx))^2 \csc(i\operatorname{arccosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}} + \sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2 \right) \right. \\ \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a+\operatorname{barccosh}(cx)) - 2c^2x(a+\operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4\arctan(\sqrt{c^2x^2-1}))}{3c^3\sqrt{cx-1}} \right) \right) \\ \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} \\ 2x^2$$

↓ 4668

---

3.192.  $\int \frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^3} dx$

$$\begin{aligned}
& -\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2}(-2ib \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
& \left. \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4 \arctan(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}) \right. \right. \right. \\
& \left. \left. \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx))^2} \right) \right) \right) \\
& \qquad \qquad \qquad \downarrow \quad \mathbf{3011}
\end{aligned}$$

$$\begin{aligned}
& -\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2}(2ib(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
& \left. \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4 \arctan(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}) \right. \right. \right. \\
& \left. \left. \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx))^2} \right) \right) \right) \\
& \qquad \qquad \qquad \downarrow \quad \mathbf{2720}
\end{aligned}$$

$$\begin{aligned}
& -\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2}(2ib(b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
& \left. \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4 \arctan(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}) \right. \right. \right. \\
& \left. \left. \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx))^2} \right) \right) \right) \\
& \qquad \qquad \qquad \downarrow \quad \mathbf{7143}
\end{aligned}$$

$$\begin{aligned}
& -\frac{5}{2}c^2d \left( d \left( -\frac{\sqrt{d-c^2dx^2}(2 \arctan(e^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))^2 + 2ib(b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx)))}{\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\
& \left. \left. bcd^2\sqrt{d-c^2dx^2} \left( \frac{1}{3}c^4x^3(a + \operatorname{barccosh}(cx)) - 2c^2x(a + \operatorname{barccosh}(cx)) - \frac{a+\operatorname{barccosh}(cx)}{x} + \frac{b\sqrt{c^2x^2-1}(3c^4 \arctan(\frac{\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}}) \right. \right. \right. \\
& \left. \left. \left. \frac{\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{5/2}(a + \operatorname{barccosh}(cx))^2} \right) \right) \right)
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x^3, x]`

output `-1/2*((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x^2 + (b*c*d^2*Sqrt[d - c^2*d*x^2]*(-(a + b*ArcCosh[c*x])/x) - 2*c^2*x*(a + b*ArcCosh[c*x]) + (c^4*x^3*(a + b*ArcCosh[c*x]))/3 + (b*Sqrt[-1 + c^2*x^2]*(-1/3*(c^4*x^6) + 5*c^4*Sqrt[-1 + c^2*x^2] + 3*c^4*ArcTan[Sqrt[-1 + c^2*x^2]]))/(3*c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (5*c^2*d*((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2)/3 - (2*b*c*d*Sqrt[d - c^2*d*x^2]*(-1/3*(b*c*((7*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2) - (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/3)) + x*(a + b*ArcCosh[c*x]) - (c^2*x^3*(a + b*ArcCosh[c*x]))/3))/(3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + d*(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2 - (2*b*c*Sqrt[d - c^2*d*x^2]*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (Sqrt[d - c^2*d*x^2]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]])))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/2`

### 3.192.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 960 `Int[((e_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*  
*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(  
m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n  
*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/  
(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n  
/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2,  
n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 1192 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_)  
+ (c_)*(x_)^2)^(p_), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(  
2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 +  
c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] &&  
IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_),  
x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x],  
x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e  
+ a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)  
)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a  
+ b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && Int  
egerQ[(m - 1)/2]`

rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*  
*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x  
_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[  
q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a  
+ b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p,  
q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*x)))]^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6304 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(p_), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6309 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6327 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

---

3.192. 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^3} dx$$

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6341 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^(m + 1)*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/(f*(m + 2))), x] + (-Simp[(1/(m + 2))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]) Int[(f*x)^m*((a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] - Simp[b*c*(n/(f*(m + 2)))*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]) Int[(f*x)^(m + 1)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && (IGtQ[m, -2] || EqQ[n, 1])`

rule 6343 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1]`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6362 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_)]/(Sqrt[(d1_) + (e1_.)*(x_.)]*Sqrt[(d2_) + (e2_.)*(x_.)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.192.4 Maple [F]

$$\int \frac{(-c^2 d x^2 + d)^{\frac{5}{2}} (a + b \operatorname{arccosh}(c x))^2}{x^3} dx$$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^3,x)`

output `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^3,x)`

### 3.192.5 Fracas [F]

$$\int \frac{(d - c^2 d x^2)^{5/2} (a + b \operatorname{arccosh}(c x))^2}{x^3} dx = \int \frac{(-c^2 d x^2 + d)^{5/2} (b \operatorname{arccosh}(c x) + a)^2}{x^3} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^3,x, algorithm="fracas")`

output `integral((a^2*c^4*d^2*x^4 - 2*a^2*c^2*d^2*x^2 + a^2*d^2 + (b^2*c^4*d^2*x^4 - 2*b^2*c^2*d^2*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*c^4*d^2*x^4 - 2*a*b*c^2*d^2*x^2 + a*b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^3, x)`

**3.192.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \text{Timed out}$$

```
input integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2/x**3,x)
```

```
output Timed out
```

**3.192.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2}{x^3} dx$$

```
input integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^3,x, algorithm="maxima")
```

```
output 1/6*(15*c^2*d^(5/2)*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x)) - 3*(-c^2*d*x^2 + d)^(5/2)*c^2 - 5*(-c^2*d*x^2 + d)^(3/2)*c^2*d - 15*sqrt(-c^2*d*x^2 + d)*c^2*d^2 - 3*(-c^2*d*x^2 + d)^(7/2)/(d*x^2))*a^2 + integrate((-c^2*d*x^2 + d)^(5/2)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x^3 + 2*(-c^2*d*x^2 + d)^(5/2)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^3, x)
```

**3.192.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^3,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value
```

---

3.192.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx$



**3.192.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2}}{x^3} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x^3,x)`output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x^3, x)`

**3.193** 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx$$

3.193.1 Optimal result . . . . . 1681  
 3.193.2 Mathematica [A] (warning: unable to verify) . . . . . 1682  
 3.193.3 Rubi [F] . . . . . 1683  
 3.193.4 Maple [A] (verified) . . . . . 1693  
 3.193.5 Fracas [F] . . . . . 1694  
 3.193.6 Sympy [F(-1)] . . . . . 1694  
 3.193.7 Maxima [F] . . . . . 1695  
 3.193.8 Giac [F(-2)] . . . . . 1695  
 3.193.9 Mupad [F(-1)] . . . . . 1696

**3.193.1 Optimal result**

Integrand size = 29, antiderivative size = 638

$$\begin{aligned} \int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx &= \frac{7}{12}b^2c^4d^2x\sqrt{d-c^2dx^2} \\ &+ \frac{b^2c^2d^2(1-cx)(1+cx)\sqrt{d-c^2dx^2}}{3x} + \frac{23b^2c^3d^2\sqrt{d-c^2dx^2}\operatorname{arccosh}(cx)}{12\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{5bc^5d^2x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{2\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{7bc^3d^2(1-c^2x^2)\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{bcd^2(1-c^2x^2)^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{3x^2\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{5}{2}c^4d^2x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2 - \frac{7c^3d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{5c^2d(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{3x} \\ &- \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{3x^3} - \frac{5c^3d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^3}{6b\sqrt{-1+cx}\sqrt{1+cx}} \\ &- \frac{14bc^3d^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))\log(1+e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{-1+cx}\sqrt{1+cx}} \\ &+ \frac{7b^2c^3d^2\sqrt{d-c^2dx^2}\operatorname{PolyLog}(2,-e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output 
$$\begin{aligned} & 5/3*c^2*d*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^{2/x}-1/3*(-c^2*d*x^2+d)^{(5/2)}*(a+b*\operatorname{arccosh}(c*x))^{2/x^3}+7/12*b^2*c^4*d^2*x*(-c^2*d*x^2+d)^{(1/2)}+1/3*b^2*c^2*d^2*(-c*x+1)*(c*x+1)*(-c^2*d*x^2+d)^{(1/2)}/x+5/2*c^4*d^2*x*(a+b*\operatorname{arccosh}(c*x))^{2*(-c^2*d*x^2+d)^{(1/2)}+23/12*b^2*c^3*d^2*\operatorname{arccosh}(c*x)*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-5/2*b*c^5*d^2*x^2*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-7/3*b*c^3*d^2*(-c^2*x^2+1)*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/3*b*c*d^2*(-c^2*x^2+1)^2*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/x^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-7/3*c^3*d^2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-5/6*c^3*d^2*(a+b*\operatorname{arccosh}(c*x))^3*(-c^2*d*x^2+d)^{(1/2)}/b/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-14/3*b*c^3*d^2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+1/(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+7/3*b^2*c^3*d^2*\operatorname{polylog}(2,-1/(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))^2*(-c^2*d*x^2+d)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \end{aligned}$$

### 3.193.2 Mathematica [A] (warning: unable to verify)

Time = 2.73 (sec) , antiderivative size = 803, normalized size of antiderivative = 1.26

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2}{x^4} dx = \frac{-8abcd^3x + 8abc^2d^3x^2 - 8a^2d^3\sqrt{\frac{-1+cx}{1+cx}} + 64a^2c^2d^3x^2\sqrt{\frac{-1+cx}{1+cx}} + \dots}{x^4}$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x]))^2/x^4,x]`

output

$$\begin{aligned}
& (-8*a*b*c*d^3*x + 8*a*b*c^2*d^3*x^2 - 8*a^2*d^3*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] \\
& + 64*a^2*c^2*d^3*x^2*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] + 8*b^2*c^2*d^3*x^2*\text{Sqrt}[ \\
& (-1 + c*x)/(1 + c*x)] - 44*a^2*c^4*d^3*x^4*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] - 8* \\
& b^2*c^4*d^3*x^4*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] - 12*a^2*c^6*d^3*x^6*\text{Sqrt}[(-1 + \\
& c*x)/(1 + c*x)] + 20*b^2*c^3*d^3*x^3*(-1 + c*x)*\text{ArcCosh}[c*x]^3 - 60*a^2*c \\
& ^3*d^(5/2)*x^3*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*\text{Sqrt}[d - c^2*d*x^2]*\text{ArcTan}[(c*x* \\
& \text{Sqrt}[d - c^2*d*x^2])]/(\text{Sqrt}[d]*(-1 + c^2*x^2))] - 6*a*b*c^3*d^3*x^3*\text{Cosh}[2* \\
& \text{ArcCosh}[c*x]] + 6*a*b*c^4*d^3*x^4*\text{Cosh}[2*\text{ArcCosh}[c*x]] - 112*a*b*c^3*d^3*x \\
& ^3*\text{Log}[c*x] + 112*a*b*c^4*d^3*x^4*\text{Log}[c*x] - 56*b^2*c^3*d^3*x^3*(-1 + c*x) \\
& *\text{PolyLog}[2, -E^(-2*\text{ArcCosh}[c*x])] + 3*b^2*c^3*d^3*x^3*\text{Sinh}[2*\text{ArcCosh}[c*x]] \\
& - 3*b^2*c^4*d^3*x^4*\text{Sinh}[2*\text{ArcCosh}[c*x]] + 2*b*d^3*(-1 + c*x)*\text{ArcCosh}[c*x \\
& ]*(4*b*c*x + 8*a*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] + 8*a*c*x*\text{Sqrt}[(-1 + c*x)/(1 + \\
& c*x)] - 56*a*c^2*x^2*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] - 56*a*c^3*x^3*\text{Sqrt}[(-1 + \\
& c*x)/(1 + c*x)] + 3*b*c^3*x^3*\text{Cosh}[2*\text{ArcCosh}[c*x]] + 56*b*c^3*x^3*\text{Log}[1 + \\
& E^(-2*\text{ArcCosh}[c*x])] - 6*a*c^3*x^3*\text{Sinh}[2*\text{ArcCosh}[c*x]]) - 2*b*d^3*(-1 + \\
& c*x)*\text{ArcCosh}[c*x]^2*(-30*a*c^3*x^3 + 4*b*(-\text{Sqrt}[(-1 + c*x)/(1 + c*x)] - c* \\
& x*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] + 7*c^2*x^2*\text{Sqrt}[(-1 + c*x)/(1 + c*x)] + 7*c^ \\
& 3*x^3*(-1 + \text{Sqrt}[(-1 + c*x)/(1 + c*x)])) + 3*b*c^3*x^3*\text{Sinh}[2*\text{ArcCosh}[c*x] \\
& ])/((24*x^3*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*\text{Sqrt}[d - c^2*d*x^2])
\end{aligned}$$

### 3.193.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2}{x^4} dx \\
& \quad \downarrow \text{6343} \\
& \frac{2bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-cx)^2 (cx+1)^2 (a+\text{barccosh}(cx))}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \\
& \frac{5}{3}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^2}{x^2} dx - \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2}{3x^3} \\
& \quad \downarrow \text{6327} \\
& \frac{2bcd^2 \sqrt{d - c^2 dx^2} \int \frac{(1-c^2 x^2)^2 (a+\text{barccosh}(cx))}{x^3} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{5}{3}c^2d \int \frac{(d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^2}{x^2} dx - \\
& \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2}{3x^3} \\
& \quad \downarrow \text{6335}
\end{aligned}$$

---

3.193.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2}{x^4} dx$

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx+\frac{1}{2}bc\int\frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x^2}dx-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}dx}$$

↓ 108

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx+\frac{1}{2}bc\left(\int 3c^2\sqrt{cx-1}\sqrt{cx+1}dx-\frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x}\right)-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}dx}$$

↓ 27

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx+\frac{1}{2}bc\left(3c^2\int\sqrt{cx-1}\sqrt{cx+1}dx-\frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x}\right)-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}dx}$$

↓ 40

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx+\frac{1}{2}bc\left(3c^2\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2}\right)\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}dx}$$

↓ 43

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\int\frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x}dx-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2}+\frac{1}{2}bc\left(3c^2\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)-\frac{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))}{2x^2}\right)\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{3\sqrt{cx-1}\sqrt{cx+1}}{3x^3}(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}dx}$$

↓ 6334

---

3.193.  $\int\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^4}dx$

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{2}bc\int\sqrt{cx-1}\sqrt{cx+1}dx+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))\right)-\frac{(1-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{3\sqrt{cx-1}\sqrt{cx+1}}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{3x^3}}$$

↓ 40

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{2}bc\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{1}{2}\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))\right)-\frac{(1-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{3\sqrt{cx-1}\sqrt{cx+1}}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{3x^3}}$$

↓ 43

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\int\frac{a+\operatorname{barccosh}(cx)}{x}dx+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))+\frac{1}{2}bc\left(\frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1}-\frac{\operatorname{arccosh}(cx)}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)-\frac{(1-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{3\sqrt{cx-1}\sqrt{cx+1}}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{3x^3}}$$

↓ 6297

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\frac{\int-\left((a+\operatorname{barccosh}(cx))\tanh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)\right)d(a+\operatorname{barccosh}(cx))}{b}+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))\right)-\frac{(1-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{3\sqrt{cx-1}\sqrt{cx+1}}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{3x^3}}$$

↓ 25

$$\frac{2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(-\frac{\int(a+\operatorname{barccosh}(cx))\tanh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)d(a+\operatorname{barccosh}(cx))}{b}+\frac{1}{2}(1-c^2x^2)(a+\operatorname{barccosh}(cx))\right)-\frac{(1-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{3\sqrt{cx-1}\sqrt{cx+1}}\right)}{\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{3x^3}}$$

↓ 3042

---

3.193.  $\int\frac{(d-c^2dx^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}{x^4}dx$

$$2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(-\frac{\int -i(a+b\operatorname{arccosh}(cx))\tan\left(\frac{ia}{b}-\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)d(a+b\operatorname{arccosh}(cx))}{b}+\frac{1}{2}(1-c^2x^2)(a+b\operatorname{arccosh}(cx))\right)\right)$$

$$\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{3x^3}$$

↓ 26

$$2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\frac{i\int(a+b\operatorname{arccosh}(cx))\tan\left(\frac{ia}{b}-\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)d(a+b\operatorname{arccosh}(cx))}{b}+\frac{1}{2}(1-c^2x^2)(a+b\operatorname{arccosh}(cx))\right)\right)$$

$$\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{3x^3}$$

↓ 4201

$$2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\frac{i\left(2i\int\frac{e^{-2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))d(a+b\operatorname{arccosh}(cx))-\frac{1}{2}i(a+b\operatorname{arccosh}(cx))^2}{1+e^{-2\operatorname{arccosh}(cx)}}}{b}\right)+\frac{1}{2}(1-c^2x^2)(a+b\operatorname{arccosh}(cx))\right)\right)$$

$$\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{3x^3}$$

↓ 2620

$$2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\frac{i\left(2i\left(\frac{1}{2}b\int\log(1+e^{-2\operatorname{arccosh}(cx)})d(a+b\operatorname{arccosh}(cx))-\frac{1}{2}b\log(e^{-2\operatorname{arccosh}(cx)}+1)(a+b\operatorname{arccosh}(cx))\right)-\frac{1}{2}i(a+b\operatorname{arccosh}(cx))^2}{b}\right)+\frac{1}{2}(1-c^2x^2)(a+b\operatorname{arccosh}(cx))\right)\right)$$

$$\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{3x^3}$$

↓ 2715

$$2bcd^2\sqrt{d-c^2dx^2}\left(-2c^2\left(\frac{i\left(2i\left(-\frac{1}{4}b^2\int e^{2\operatorname{arccosh}(cx)}\log(1+e^{-2\operatorname{arccosh}(cx)})de^{-2\operatorname{arccosh}(cx)}-\frac{1}{2}b\log(e^{-2\operatorname{arccosh}(cx)}+1)(a+b\operatorname{arccosh}(cx))\right)-\frac{1}{2}i(a+b\operatorname{arccosh}(cx))^2}{b}\right)+\frac{1}{2}(1-c^2x^2)(a+b\operatorname{arccosh}(cx))\right)\right)$$

$$\frac{5}{3}c^2d\int\frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}{x^2}dx-\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{3x^3}$$

↓ 2838

---

3.193.  $\int\frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^4}dx$

$$-\frac{5}{3}c^2d \int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} dx +$$

$$2bcd^2\sqrt{d - c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1))(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6343

$$-\frac{5}{3}c^2d \left( -3c^2d \int \sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))^2 dx - \frac{2bcd\sqrt{d - c^2dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} \right)$$

$$2bcd^2\sqrt{d - c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1))(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 25

$$-\frac{5}{3}c^2d \left( -3c^2d \int \sqrt{d - c^2dx^2} (a + \operatorname{barccosh}(cx))^2 dx + \frac{2bcd\sqrt{d - c^2dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{x^2} \right)$$

$$2bcd^2\sqrt{d - c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1))(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6310

$$-\frac{5}{3}c^2d \left( -3c^2d \left( -\frac{bc\sqrt{d - c^2dx^2} \int x(a + \operatorname{barccosh}(cx)) dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d - c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d - c^2dx^2} \right) \right)$$

$$2bcd^2\sqrt{d - c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1))(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6298

---

3.193.  $\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx$



$$-\frac{5}{3}c^2d \left( -3c^2d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}}}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\ \left. \left. 2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i \left( 2i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) \right) - \frac{1}{2}i(a+\operatorname{barccosh}(cx)) \right)}{b} \right) \right) \right)$$

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 101

$$-\frac{5}{3}c^2d \left( -3c^2d \left( -\frac{bc\sqrt{d-c^2dx^2} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \left( \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right. \right. \\ \left. \left. 2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i \left( 2i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) \right) - \frac{1}{2}i(a+\operatorname{barccosh}(cx)) \right)}{b} \right) \right) \right)$$

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 43

$$-\frac{5}{3}c^2d \left( \frac{2bcd\sqrt{d-c^2dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - 3c^2d \left( -\frac{\sqrt{d-c^2dx^2} \int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}x\sqrt{d-c^2dx^2} \right) \right. \\ \left. 2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i \left( 2i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) \right) - \frac{1}{2}i(a+\operatorname{barccosh}(cx)) \right)}{b} \right) \right) \right)$$

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6308

$$-\frac{5}{3}c^2d \left( \frac{2bcd\sqrt{d-c^2dx^2} \int \frac{(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x} - 3c^2d \left( -\frac{\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \right. \\ \left. 2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i \left( 2i \left( \frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1) \right) (a+\operatorname{barccosh}(cx)) \right) - \frac{1}{2}i(a+\operatorname{barccosh}(cx)) \right)}{b} \right) \right) \right)$$

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

---

3.193.  $\int \frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{x^4} dx$

↓ 6327

$$-\frac{5}{3}c^2d \left( \frac{2bcd\sqrt{d-c^2dx^2} \int \frac{(1-c^2x^2)(a+\operatorname{barccosh}(cx))}{x} dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{(d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2}{x} - 3c^2d \left( -\frac{\sqrt{d-c^2dx^2}}{6bc} \right) \right)$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2,-a-\operatorname{barccosh}(cx))-\frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1))(a+\operatorname{barccosh}(cx)))-\frac{1}{2}i(a+\operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 6334

$$-\frac{5}{3}c^2d \left( \frac{2bcd\sqrt{d-c^2dx^2} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}bc \int \sqrt{cx-1}\sqrt{cx+1} dx + \frac{1}{2}(1-c^2x^2) (a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \left( \dots \right) \right)$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2,-a-\operatorname{barccosh}(cx))-\frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1))(a+\operatorname{barccosh}(cx)))-\frac{1}{2}i(a+\operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 40

$$-\frac{5}{3}c^2d \left( \frac{2bcd\sqrt{d-c^2dx^2} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}bc \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \frac{1}{2} \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx \right) + \frac{1}{2}(1-c^2x^2) (a+\operatorname{barccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{2}(1-c^2x^2) (a+\operatorname{barccosh}(cx)) \right)$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2,-a-\operatorname{barccosh}(cx))-\frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1))(a+\operatorname{barccosh}(cx)))-\frac{1}{2}i(a+\operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

↓ 43

$$-\frac{5}{3}c^2d \left( \frac{2bcd\sqrt{d-c^2dx^2} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x} dx + \frac{1}{2}(1-c^2x^2) (a+\operatorname{barccosh}(cx)) + \frac{1}{2}bc \left( \frac{1}{2}x\sqrt{cx-1}\sqrt{cx+1} - \operatorname{arcc} \right) \right)}{\sqrt{cx-1}\sqrt{cx+1}} - \left( \dots \right) \right)$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2,-a-\operatorname{barccosh}(cx))-\frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)}+1))(a+\operatorname{barccosh}(cx)))-\frac{1}{2}i(a+\operatorname{barccosh}(cx))}{b} \right) \right)$$


---

$$\frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{3x^3}$$

---

3.193.  $\int \frac{(d-c^2dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2}{x^4} dx$

↓ 6297

$$-\frac{5}{3}c^2d \left( \frac{2bcd\sqrt{d-c^2dx^2} \left( \frac{f - \left( (a+b\operatorname{arccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)\right) d(a+b\operatorname{arccosh}(cx))}{b} + \frac{1}{2}(1-c^2x^2)(a+b\operatorname{arccosh}(cx)) \right)}{\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{2bcd^2\sqrt{d-c^2dx^2} \left( -2c^2 \left( \frac{i(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-b\operatorname{arccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1))(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}i(a+b\operatorname{arccosh}(cx)))}{b} \right)}{\sqrt{d-c^2dx^2}} \right)}{\sqrt{d-c^2dx^2}} \right)$$

$$\frac{(d-c^2dx^2)^{5/2} (a+b\operatorname{arccosh}(cx))^2}{3x^3}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2)/x^4, x]`

output `$Aborted`

### 3.193.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 40 `Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(m_), x_Symbol] := Simp[x*(a + b*x)^m*((c + d*x)^(m/(2*m + 1))), x] + Simp[2*a*c*(m/(2*m + 1)) Int[(a + b*x)^(m - 1)*(c + d*x)^(m - 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && IGtQ[m + 1/2, 0]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

---

3.193.  $\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}{x^4} dx$

- rule 101 `Int[((a_.) + (b_.)*(x_))2((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[b*(a + b*x)*(c + d*x)(n + 1)((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)n(e + f*x)pSimp[a2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 108 `Int[((a_.) + (b_.)*(x_))(m_)((c_.) + (d_.)*(x_))(n_)((e_.) + (f_.)*(x_))(p_), x_] := Simp[(a + b*x)(m + 1)(c + d*x)n((e + f*x)p/(b*(m + 1))), x] - Simp[1/(b*(m + 1)) Int[(a + b*x)(m + 1)(c + d*x)(n - 1)(e + f*x)(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`
- rule 2620 `Int[((F_)(g_)((e_.) + (f_.)*(x_)))(n_)((c_.) + (d_.)*(x_))(m_) / ((a_) + (b_.)*(F_)(g_)((e_.) + (f_.)*(x_)))(n_), x_Symbol] := Simp[((c + d*x)m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))n/a), x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)(m - 1)*Log[1 + b*((F^(g*(e + f*x)))n/a), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)(e_)((c_.) + (d_.)*(x_)))(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*(d_) + (e_.)*(x_)(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*xn/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4201 `Int[((c_.) + (d_.)*(x_))(m_)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)m(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b  
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a  
, b, c}, x] && IGtQ[n, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]  
:= Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*  
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +  
c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &  
& NeQ[m, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sq  
rt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 +  
c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[  
c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1  
] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_  
Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(  
1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcC  
osh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sq  
rt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^  
(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n  
, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (  
e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1  
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2  
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6334 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.)/(x_),  
x_Symbol] := Simp[(d + e*x^2)^p*((a + b*ArcCosh[c*x])/(2*p)), x] + (Simp[d  
Int[(d + e*x^2)^(p - 1)*((a + b*ArcCosh[c*x])/x), x], x] - Simp[b*c*((-d  
)^p/(2*p)) Int[(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x] /; FreeQ  
[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

```
rule 6335 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_
)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c
*x])/(f*(m + 1))), x] + (-Simp[b*c*((-d)^p/(f*(m + 1))) Int[(f*x)^(m + 1)
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2), x], x] - Simp[2*e*(p/(f^2*(m + 1
))) Int[(f*x)^(m + 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x]), x], x] /
; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0] && ILtQ[(
m + 1)/2, 0]
```

```
rule 6343 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 1))), x] + (-Simp[2*e*(p/(f^2*(m + 1))) Int[(f*x)^(m
+ 2)*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(
m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)
*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x],
x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && G
tQ[p, 0] && LtQ[m, -1]
```

### 3.193.4 Maple [A] (verified)

Time = 1.29 (sec) , antiderivative size = 652, normalized size of antiderivative = 1.02

method	result
default	$-\frac{a^2(-c^2dx^2+d)^{\frac{7}{2}}}{3dx^3} + \frac{4a^2c^2(-c^2dx^2+d)^{\frac{7}{2}}}{3dx} + \frac{4a^2c^4x(-c^2dx^2+d)^{\frac{5}{2}}}{3} + \frac{5a^2c^4dx(-c^2dx^2+d)^{\frac{3}{2}}}{3} + \frac{5a^2c^4d^2x\sqrt{-c^2dx^2+d}}{2} +$
parts	$-\frac{a^2(-c^2dx^2+d)^{\frac{7}{2}}}{3dx^3} + \frac{4a^2c^2(-c^2dx^2+d)^{\frac{7}{2}}}{3dx} + \frac{4a^2c^4x(-c^2dx^2+d)^{\frac{5}{2}}}{3} + \frac{5a^2c^4dx(-c^2dx^2+d)^{\frac{3}{2}}}{3} + \frac{5a^2c^4d^2x\sqrt{-c^2dx^2+d}}{2} +$

```
input int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^4,x,method=_RETURNVERBOSE)
```

output `-1/3*a^2/d/x^3*(-c^2*d*x^2+d)^(7/2)+4/3*a^2*c^2/d/x*(-c^2*d*x^2+d)^(7/2)+4/3*a^2*c^4*x*(-c^2*d*x^2+d)^(5/2)+5/3*a^2*c^4*d*x*(-c^2*d*x^2+d)^(3/2)+5/2*a^2*c^4*d^2*x*(-c^2*d*x^2+d)^(1/2)+5/2*a^2*c^4*d^3/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))-1/12*b^2*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^3*(-6*(c*x+1)^(1/2)*(c*x-1)^(1/2)*arccosh(c*x)^2*x^4*c^4+6*arccosh(c*x)*c^5*x^5-3*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+10*arccosh(c*x)^3*x^3*c^3-28*arccosh(c*x)^2*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^2*c^2-28*arccosh(c*x)^2*x^3*c^3+56*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3-3*c^3*x^3*arccosh(c*x)+28*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+4*c^3*x^3+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2+4*c*x*arccosh(c*x))*d^2-1/12*a*b*(-d*(c^2*x^2-1))^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)/x^3*(-12*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4+6*c^5*x^5+30*arccosh(c*x)^2*x^3*c^3+56*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3-56*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-56*c^3*x^3*arccosh(c*x)-3*c^3*x^3+8*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c*x)*d^2`

### 3.193.5 Fricas [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)^2}{x^4} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^4,x, algorithm="fricas")`

output `integral((a^2*c^4*d^2*x^4 - 2*a^2*c^2*d^2*x^2 + a^2*d^2 + (b^2*c^4*d^2*x^4 - 2*b^2*c^2*d^2*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*c^4*d^2*x^4 - 2*a*b*c^2*d^2*x^2 + a*b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)/x^4, x)`

### 3.193.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2}{x^4} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2/x**4,x)`

output Timed out

### 3.193.7 Maxima [F]

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2}{x^4} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^4,x, algorithm="maxima")`

output `1/6*(10*(-c^2*d*x^2 + d)^(3/2)*c^4*d*x + 15*sqrt(-c^2*d*x^2 + d)*c^4*d^2*x + 15*c^3*d^(5/2)*arcsin(c*x) + 8*(-c^2*d*x^2 + d)^(5/2)*c^2/x - 2*(-c^2*d*x^2 + d)^(7/2)/(d*x^3))*a^2 + integrate((-c^2*d*x^2 + d)^(5/2)*b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/x^4 + 2*(-c^2*d*x^2 + d)^(5/2)*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x^4, x)`

### 3.193.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/x^4,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.193.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2}}{x^4} dx$$

input `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x^4,x)`output `int(((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2))/x^4, x)`

$$3.194 \quad \int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$$

3.194.1 Optimal result . . . . .	1697
3.194.2 Mathematica [A] (verified) . . . . .	1698
3.194.3 Rubi [A] (verified) . . . . .	1698
3.194.4 Maple [B] (verified) . . . . .	1704
3.194.5 Fricas [A] (verification not implemented) . . . . .	1705
3.194.6 Sympy [F] . . . . .	1706
3.194.7 Maxima [A] (verification not implemented) . . . . .	1706
3.194.8 Giac [F(-2)] . . . . .	1707
3.194.9 Mupad [F(-1)] . . . . .	1707

### 3.194.1 Optimal result

Integrand size = 29, antiderivative size = 421

$$\int \frac{x^5(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = -\frac{16abx\sqrt{-1 + cx}\sqrt{1 + cx}}{15c^5\sqrt{d - c^2dx^2}} - \frac{4144b^2(1 - cx)(1 + cx)}{3375c^6\sqrt{d - c^2dx^2}}$$

$$- \frac{272b^2x^2(1 - cx)(1 + cx)}{3375c^4\sqrt{d - c^2dx^2}} - \frac{2b^2x^4(1 - cx)(1 + cx)}{125c^2\sqrt{d - c^2dx^2}}$$

$$- \frac{16b^2x\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arccosh}(cx)}{15c^5\sqrt{d - c^2dx^2}}$$

$$- \frac{8bx^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{45c^3\sqrt{d - c^2dx^2}}$$

$$- \frac{2bx^5\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{25c\sqrt{d - c^2dx^2}}$$

$$- \frac{8\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{15c^6d}$$

$$- \frac{4x^2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{15c^4d}$$

$$- \frac{x^4\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{5c^2d}$$

output 
$$-4144/3375*b^2*(-c*x+1)*(c*x+1)/c^6/(-c^2*d*x^2+d)^{(1/2)}-272/3375*b^2*x^2*(-c*x+1)*(c*x+1)/c^4/(-c^2*d*x^2+d)^{(1/2)}-2/125*b^2*x^4*(-c*x+1)*(c*x+1)/c^2/(-c^2*d*x^2+d)^{(1/2)}-16/15*a*b*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^5/(-c^2*d*x^2+d)^{(1/2)}-16/15*b^2*x*arccosh(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^5/(-c^2*d*x^2+d)^{(1/2)}-8/45*b*x^3*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/(-c^2*d*x^2+d)^{(1/2)}-2/25*b*x^5*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c/(-c^2*d*x^2+d)^{(1/2)}-8/15*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/c^6/d-4/15*x^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/c^4/d-1/5*x^4*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/c^2/d$$

### 3.194.2 Mathematica [A] (verified)

Time = 0.48 (sec) , antiderivative size = 255, normalized size of antiderivative = 0.61

$$\int \frac{x^5(a + b\operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{d - c^2dx^2}(30abcx\sqrt{-1 + cx}\sqrt{1 + cx}(120 + 20c^2x^2 + 9c^4x^4) - 225a^2(-8 + 4c^2x^2 + c^4x^4 + 3c^6x^6) - 2b^2}$$

input `Integrate[(x^5*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2], x]`

output 
$$(\operatorname{Sqrt}[d - c^2*d*x^2]*(30*a*b*c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(120 + 20*c^2*x^2 + 9*c^4*x^4) - 225*a^2*(-8 + 4*c^2*x^2 + c^4*x^4 + 3*c^6*x^6) - 2*b^2*(-2072 + 1936*c^2*x^2 + 109*c^4*x^4 + 27*c^6*x^6) + 30*b*(b*c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(120 + 20*c^2*x^2 + 9*c^4*x^4) - 15*a*(-8 + 4*c^2*x^2 + c^4*x^4 + 3*c^6*x^6))*\operatorname{ArcCosh}[c*x] - 225*b^2*(-8 + 4*c^2*x^2 + c^4*x^4 + 3*c^6*x^6)*\operatorname{ArcCosh}[c*x]^2))/(3375*c^6*d*(-1 + c*x)*(1 + c*x))$$

### 3.194.3 Rubi [A] (verified)

Time = 1.66 (sec) , antiderivative size = 470, normalized size of antiderivative = 1.12, number of steps used = 14, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.483$ , Rules used = {6353, 6298, 111, 27, 111, 27, 83, 6353, 6298, 111, 27, 83, 6329, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5(a + b\operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx$$

3.194.  $\int \frac{x^5(a + b\operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx$

$$\begin{aligned}
& \downarrow \text{6353} \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int x^4(a+\operatorname{barccosh}(cx))dx}{5c\sqrt{d-c^2dx^2}} + \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{5c^2} \\
& \quad \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\
& \downarrow \text{6298} \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \int \frac{x^5}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{5c\sqrt{d-c^2dx^2}} + \\
& \quad \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{5c^2} - \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\
& \downarrow \text{111} \\
& \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{5c^2} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}} \\
& \quad \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\
& \downarrow \text{27} \\
& \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{5c^2} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}} \\
& \quad \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\
& \downarrow \text{111} \\
& \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{5c^2} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}} \\
& \quad \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} \\
& \downarrow \text{27}
\end{aligned}$$

---

3.194.  $\int \frac{x^5(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$

$$\begin{aligned}
& \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{5c^2} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}}}{5c^2d} \\
& \quad \downarrow 83 \\
& \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{5c^2} - \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}}}{5c^2d} \\
& \quad \downarrow 6353 \\
& 4 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int x^2(a+\operatorname{barccosh}(cx)) dx}{3c\sqrt{d-c^2dx^2}} + \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \right) - \\
& \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}}}{5c^2d} \\
& \quad \downarrow 6298 \\
& 4 \left( \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{3c\sqrt{d-c^2dx^2}} - \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \right) - \\
& \frac{x^4\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{5c^2d} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+\operatorname{barccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}}}{5c^2d} \\
& \quad \downarrow 111
\end{aligned}$$

$$\begin{aligned}
 & 4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{5c^2d} \right) \\
 & \frac{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{5c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+b\operatorname{arccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow 27 \\
 & 4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{5c^2d} \right) \\
 & \frac{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{5c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+b\operatorname{arccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow 83 \\
 & 4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} \right) \\
 & \frac{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{5c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+b\operatorname{arccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow 6329
 \end{aligned}$$

$$\begin{aligned}
 & 4 \left( \frac{2 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx)) dx - \sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c\sqrt{d-c^2dx^2}} \right)}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x \right)}{5c^2} \right) \\
 & \frac{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{5c^2d} - \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+b\operatorname{arccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{x^4\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{5c^2d} + \\
 & 4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( a+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2dx^2}} \right)}{3c^2} \right) - 2b\sqrt{cx-1}\sqrt{cx+1} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{5}x^5(a+b\operatorname{arccosh}(cx)) - \frac{1}{5}bc \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \right)}{5c\sqrt{d-c^2dx^2}}
 \end{aligned}$$

input `Int[(x^5*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `-1/5*(x^4*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/5*(b*c*((x^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(5*c^2) + (4*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2))))/(5*c^2))) + (x^5*(a + b*ArcCosh[c*x])/5)/(5*c*Sqrt[d - c^2*d*x^2]) + (4*(-1/3*(x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/3*(b*c*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2))) + (x^3*(a + b*ArcCosh[c*x])/3))/(3*c*Sqrt[d - c^2*d*x^2]) + (2*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d)) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(c*Sqrt[d - c^2*d*x^2])))/(3*c^2)))/(5*c^2)`

## 3.194.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 111 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6298 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6329 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`



```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.194.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1313 vs.  $2(365) = 730$ .

Time = 0.91 (sec) , antiderivative size = 1314, normalized size of antiderivative = 3.12

method	result	size
default	Expression too large to display	1314
parts	Expression too large to display	1314

```
input int(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output

```

a^2*(-1/5*x^4/c^2/d*(-c^2*d*x^2+d)^(1/2)+4/5/c^2*(-1/3*x^2/c^2/d*(-c^2*d*x
^2+d)^(1/2)-2/3/d/c^4*(-c^2*d*x^2+d)^(1/2)))+b^2*(-1/4000*(-d*(c^2*x^2-1))
^(1/2)*(16*c^6*x^6-28*c^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^
2*x^2-20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)
*c*x-1)*(25*arccosh(c*x)^2-10*arccosh(c*x)+2)/c^6/d/(c^2*x^2-1)-5/864*(-d*
(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^2+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*
x^3-3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+1)*(9*arccosh(c*x)^2-6*arccosh(c*x)+
2)/c^6/d/(c^2*x^2-1)-5/16*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1
/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*arccosh(c*x)+2)/c^6/d/(c^2*x^2-1)-5/1
6*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arc
cosh(c*x)^2+2*arccosh(c*x)+2)/c^6/d/(c^2*x^2-1)-5/864*(-d*(c^2*x^2-1))^(1/
2)*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3+4*c^4*x^4+3*(c*x-1)^(1/2)*(c*x+
1)^(1/2)*c*x-5*c^2*x^2+1)*(9*arccosh(c*x)^2+6*arccosh(c*x)+2)/c^6/d/(c^2*x
^2-1)-1/4000*(-d*(c^2*x^2-1))^(1/2)*(-16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x
^5+16*c^6*x^6+20*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*x^3-28*c^4*x^4-5*(c*x-1)^(
1/2)*(c*x+1)^(1/2)*c*x+13*c^2*x^2-1)*(25*arccosh(c*x)^2+10*arccosh(c*x)+2
)/c^6/d/(c^2*x^2-1))+2*a*b*(-1/800*(-d*(c^2*x^2-1))^(1/2)*(16*c^6*x^6-28*c
^4*x^4+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*x^5+13*c^2*x^2-20*(c*x-1)^(1/2)*
(c*x+1)^(1/2)*c^3*x^3+5*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-1)*(-1+5*arccosh(c
*x))/c^6/d/(c^2*x^2-1)-5/288*(-d*(c^2*x^2-1))^(1/2)*(4*c^4*x^4-5*c^2*x^...

```

### 3.194.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 348, normalized size of antiderivative = 0.83

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \frac{225(3b^2c^6x^6 + b^2c^4x^4 + 4b^2c^2x^2 - 8b^2)\sqrt{-c^2dx^2 + d} \log(cx + \sqrt{c^2x^2 - 1})^2 - 30(9abc^5x^5 + 20abc^3x^3 + 15ac^2x^2 + 5a^2x)}{2d^2}$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output 
$$\begin{aligned} & -1/3375*(225*(3*b^2*c^6*x^6 + b^2*c^4*x^4 + 4*b^2*c^2*x^2 - 8*b^2)*\sqrt{-c^2*d*x^2 + d})*\log(c*x + \sqrt{c^2*x^2 - 1})^2 - 30*(9*a*b*c^5*x^5 + 20*a*b*c^3*x^3 + 120*a*b*c*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} - 30*((9*b^2*c^5*x^5 + 20*b^2*c^3*x^3 + 120*b^2*c*x)*\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1} - 15*(3*a*b*c^6*x^6 + a*b*c^4*x^4 + 4*a*b*c^2*x^2 - 8*a*b)*\sqrt{-c^2*d*x^2 + d})*\log(c*x + \sqrt{c^2*x^2 - 1}) + (27*(25*a^2 + 2*b^2)*c^6*x^6 + (225*a^2 + 218*b^2)*c^4*x^4 + 4*(225*a^2 + 968*b^2)*c^2*x^2 - 1800*a^2 - 4144*b^2)*\sqrt{-c^2*d*x^2 + d})/(c^8*d*x^2 - c^6*d) \end{aligned}$$

### 3.194.6 Sympy [F]

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))^2}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate(x**5*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(1/2), x)`

output `Integral(x**5*(a + b*acosh(c*x))**2/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

### 3.194.7 Maxima [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 407, normalized size of antiderivative = 0.97

$$\begin{aligned} & \int \frac{x^5(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx \\ &= -\frac{1}{15} \left( \frac{3\sqrt{-c^2 dx^2 + dx^4}}{c^2 d} + \frac{4\sqrt{-c^2 dx^2 + dx^2}}{c^4 d} + \frac{8\sqrt{-c^2 dx^2 + d}}{c^6 d} \right) b^2 \operatorname{arccosh}(cx)^2 \\ & - \frac{2}{15} \left( \frac{3\sqrt{-c^2 dx^2 + dx^4}}{c^2 d} + \frac{4\sqrt{-c^2 dx^2 + dx^2}}{c^4 d} + \frac{8\sqrt{-c^2 dx^2 + d}}{c^6 d} \right) ab \operatorname{arccosh}(cx) \\ & - \frac{1}{15} \left( \frac{3\sqrt{-c^2 dx^2 + dx^4}}{c^2 d} + \frac{4\sqrt{-c^2 dx^2 + dx^2}}{c^4 d} + \frac{8\sqrt{-c^2 dx^2 + d}}{c^6 d} \right) a^2 \\ & - \frac{2}{3375} b^2 \left( \frac{27\sqrt{c^2 x^2 - 1} c^2 \sqrt{-dx^4} + 136\sqrt{c^2 x^2 - 1} \sqrt{-dx^2} + \frac{2072\sqrt{c^2 x^2 - 1} \sqrt{-d}}{c^2}}{c^4 d} - \frac{15(9c^4 \sqrt{-dx^5} + 20c^2 \sqrt{-dx^3} + 120\sqrt{-dx}) ab}{225c^5 d} \right) \end{aligned}$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/15*(3*sqrt(-c^2*d*x^2 + d)*x^4/(c^2*d) + 4*sqrt(-c^2*d*x^2 + d)*x^2/(c^4*d) + 8*sqrt(-c^2*d*x^2 + d)/(c^6*d))*b^2*arccosh(c*x)^2 - 2/15*(3*sqrt(-c^2*d*x^2 + d)*x^4/(c^2*d) + 4*sqrt(-c^2*d*x^2 + d)*x^2/(c^4*d) + 8*sqrt(-c^2*d*x^2 + d)/(c^6*d))*a*b*arccosh(c*x) - 1/15*(3*sqrt(-c^2*d*x^2 + d)*x^4/(c^2*d) + 4*sqrt(-c^2*d*x^2 + d)*x^2/(c^4*d) + 8*sqrt(-c^2*d*x^2 + d)/(c^6*d))*a^2 - 2/3375*b^2*((27*sqrt(c^2*x^2 - 1)*c^2*sqrt(-d)*x^4 + 136*sqrt(c^2*x^2 - 1)*sqrt(-d)*x^2 + 2072*sqrt(c^2*x^2 - 1)*sqrt(-d)/c^2)/(c^4*d) - 15*(9*c^4*sqrt(-d)*x^5 + 20*c^2*sqrt(-d)*x^3 + 120*sqrt(-d)*x)*arccosh(c*x)/(c^5*d) + 2/225*(9*c^4*sqrt(-d)*x^5 + 20*c^2*sqrt(-d)*x^3 + 120*sqrt(-d)*x)*a*b/(c^5*d)`

### 3.194.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.194.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^5*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^5*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2), x)`

---

3.194.  $\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$

### 3.195 $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$

3.195.1 Optimal result	1708
3.195.2 Mathematica [A] (verified)	1709
3.195.3 Rubi [A] (verified)	1709
3.195.4 Maple [B] (verified)	1714
3.195.5 Fracas [F]	1715
3.195.6 Sympy [F]	1716
3.195.7 Maxima [F]	1716
3.195.8 Giac [F]	1716
3.195.9 Mupad [F(-1)]	1717

#### 3.195.1 Optimal result

Integrand size = 29, antiderivative size = 355

$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx = -\frac{15b^2x(1-cx)(1+cx)}{64c^4\sqrt{d-c^2dx^2}} - \frac{b^2x^3(1-cx)(1+cx)}{32c^2\sqrt{d-c^2dx^2}}$$

$$+ \frac{15b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{arccosh}(cx)}{64c^5\sqrt{d-c^2dx^2}}$$

$$- \frac{3bx^2\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{8c^3\sqrt{d-c^2dx^2}}$$

$$- \frac{bx^4\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{8c\sqrt{d-c^2dx^2}}$$

$$- \frac{3x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{8c^4d}$$

$$- \frac{x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{4c^2d}$$

$$+ \frac{\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^3}{8bc^5\sqrt{d-c^2dx^2}}$$

output

```
-15/64*b^2*x*(-c*x+1)*(c*x+1)/c^4/(-c^2*d*x^2+d)^(1/2)-1/32*b^2*x^3*(-c*x+
1)*(c*x+1)/c^2/(-c^2*d*x^2+d)^(1/2)+15/64*b^2*arccosh(c*x)*(c*x-1)^(1/2)*(
c*x+1)^(1/2)/c^5/(-c^2*d*x^2+d)^(1/2)-3/8*b*x^2*(a+b*arccosh(c*x))*(c*x-1)
^(1/2)*(c*x+1)^(1/2)/c^3/(-c^2*d*x^2+d)^(1/2)-1/8*b*x^4*(a+b*arccosh(c*x))
*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)+1/8*(a+b*arccosh(c*x))
^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^5/(-c^2*d*x^2+d)^(1/2)-3/8*x*(a+b*arcco
sh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/c^4/d-1/4*x^3*(a+b*arccosh(c*x))^2*(-c^2*d
*x^2+d)^(1/2)/c^2/d
```

### 3.195.2 Mathematica [A] (verified)

Time = 1.54 (sec) , antiderivative size = 295, normalized size of antiderivative = 0.83

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{32a^2 c \sqrt{d} x (-1 + c^2 x^2) (3 + 2c^2 x^2) - 96a^2 \sqrt{d - c^2 dx^2} \arctan\left(\frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right) + b^2 \sqrt{d} \sqrt{\frac{-1 + cx}{1 + cx}} (1 + cx) (32a^2 c \sqrt{d} x (-1 + c^2 x^2) (3 + 2c^2 x^2) - 96a^2 \sqrt{d - c^2 dx^2} \arctan\left(\frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right) + b^2 \sqrt{d} \sqrt{\frac{-1 + cx}{1 + cx}} (1 + cx))}{(32a^2 c \sqrt{d} x (-1 + c^2 x^2) (3 + 2c^2 x^2) - 96a^2 \sqrt{d - c^2 dx^2} \arctan\left(\frac{cx\sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right) + b^2 \sqrt{d} \sqrt{\frac{-1 + cx}{1 + cx}} (1 + cx))}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `(32*a^2*c*Sqrt[d]*x*(-1 + c^2*x^2)*(3 + 2*c^2*x^2) - 96*a^2*Sqrt[d - c^2*d*x^2]*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + b^2*Sqrt[d]*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(32*ArcCosh[c*x]^3 - 4*ArcCosh[c*x]*(16*Cosh[2*ArcCosh[c*x]] + Cosh[4*ArcCosh[c*x]]) + 32*Sinh[2*ArcCosh[c*x]] + Sinh[4*ArcCosh[c*x]] + 8*ArcCosh[c*x]^2*(8*Sinh[2*ArcCosh[c*x]] + Sinh[4*ArcCosh[c*x]]) - 4*a*b*Sqrt[d]*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(16*Cosh[2*ArcCosh[c*x]] + Cosh[4*ArcCosh[c*x]] - 4*ArcCosh[c*x]*(6*ArcCosh[c*x] + 8*Sinh[2*ArcCosh[c*x]] + Sinh[4*ArcCosh[c*x]])))/(256*c^5*Sqrt[d]*Sqrt[d - c^2*d*x^2])`

### 3.195.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.05, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.379$ , Rules used = {6353, 6298, 111, 27, 101, 43, 6353, 6298, 101, 43, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6353}$$

$$\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{4c^2} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int x^3(a + \operatorname{barccosh}(cx)) dx}{2c\sqrt{d - c^2 dx^2}}$$

$$\frac{x^3\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{4c^2 d}$$

$$\downarrow \text{6298}$$

---

3.195.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$

$$\begin{aligned}
& \frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{4c^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \int \frac{x^4}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{2c\sqrt{d-c^2dx^2} \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d}} \\
& \quad \downarrow \text{111} \\
& \frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{4c^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{\int \frac{3x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2} \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d}} \\
& \quad \downarrow \text{27} \\
& \frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{4c^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2} \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d}} \\
& \quad \downarrow \text{101} \\
& \frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{4c^2} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2} \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d}} \\
& \quad \downarrow \text{43} \\
& \frac{3 \int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{4c^2} - \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6353}
\end{aligned}$$

---

3.195.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$

$$\begin{aligned}
 & \frac{3 \left( -\frac{b\sqrt{cx-1}\sqrt{cx+1} \int x(a+\operatorname{barccosh}(cx))dx}{c\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2c^2d} \right)}{4c^2} \\
 & \quad - \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d} \\
 & \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{6298} \\
 & \frac{3 \left( -\frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{c\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2c^2d} \right)}{4c^2} \\
 & \quad - \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d} \\
 & \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{101} \\
 & \frac{3 \left( -\frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2c^2d} \right)}{4c^2} \\
 & \quad - \frac{x^3\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{4c^2d} \\
 & \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+\operatorname{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{43}
 \end{aligned}$$

---

3.195.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$



$$\begin{aligned}
& 3 \left( \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c\sqrt{d-c^2dx^2}} \right) \\
& \frac{x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{4c^2} - \frac{4c^2}{4c^2d} \\
& b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+b\operatorname{arccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right) \\
& \frac{2c\sqrt{d-c^2dx^2}}{2c\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6307} \\
& -\frac{x^3\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{4c^2d} + \\
& 3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c\sqrt{d-c^2dx^2}} \right) \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}x^4(a+b\operatorname{arccosh}(cx)) - \frac{1}{4}bc \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \right)}{2c\sqrt{d-c^2dx^2}}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2], x]`

output `-1/4*(x^3*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((x^4*(a + b*ArcCosh[c*x]))/4 - (b*c*((x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(4*c^2) + (3*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/(4*c^2)))/4)/(2*c*Sqrt[d - c^2*d*x^2]) + (3*(-1/2*(x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^3)/(6*b*c^3*Sqrt[d - c^2*d*x^2]) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/2))/(c*Sqrt[d - c^2*d*x^2]))/(4*c^2)`

## 3.195.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 111 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 6298 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6307 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.195.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1091 vs.  $2(307) = 614$ .

Time = 0.93 (sec) , antiderivative size = 1092, normalized size of antiderivative = 3.08

method	result	size
default	Expression too large to display	1092
parts	Expression too large to display	1092

```
input int(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output 
$$\begin{aligned}
& -1/4*a^2*x^3/c^2/d*(-c^2*d*x^2+d)^{(1/2)}-3/8*a^2/c^4*x/d*(-c^2*d*x^2+d)^{(1/2)} \\
& +3/8*a^2/c^4/(c^2*d)^{(1/2)}*\arctan((c^2*d)^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)})+ \\
& b^2*(-1/8*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/c^5/(c^2*x^2-1) \\
& *\operatorname{arccosh}(c*x)^3-1/512*(-d*(c^2*x^2-1))^{(1/2)}*(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^{(1/2)} \\
& *(c*x-1)^{(1/2)}*c^4*x^4+4*c*x-8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+(c*x-1)^{(1/2)} \\
& *(c*x+1)^{(1/2)})*(8*\operatorname{arccosh}(c*x)^2-4*\operatorname{arccosh}(c*x)+1)/d/c^5/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^{(1/2)} \\
& *(2*c^3*x^3-2*c*x+2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)} \\
& *(2*\operatorname{arccosh}(c*x)^2-2*\operatorname{arccosh}(c*x)+1)/d/c^5/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^{(1/2)} \\
& *(-2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+2*c^3*x^3+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}-2*c*x)*(2* \\
& \operatorname{arccosh}(c*x)^2+2*\operatorname{arccosh}(c*x)+1)/d/c^5/(c^2*x^2-1)-1/512*(-d*(c^2*x^2-1))^{(1/2)} \\
& *(-8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^4*x^4+8*c^5*x^5+8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-12*c^3*x^3- \\
& (c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+4*c*x)*(8*\operatorname{arccosh}(c*x)^2+4*\operatorname{arccosh}(c*x)+1)/d/c^5/(c^2*x^2-1)+2*a*b*(-3/16*(-d*(c^2*x^2-1))^{(1/2)} \\
& *(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/c^5/(c^2*x^2-1)*\operatorname{arccosh}(c*x)^2-1/256*(-d*(c^2*x^2-1))^{(1/2)} \\
& *(8*c^5*x^5-12*c^3*x^3+8*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*c^4*x^4+4*c*x-8*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2+(c*x-1)^{(1/2)} \\
& *(c*x+1)^{(1/2)}*(-1+4*\operatorname{arccosh}(c*x)))/d/c^5/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^{(1/2)} \\
& *(2*c^3*x^3-2*c*x+2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c^2*x^2-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)} \\
& *(-1+2*\operatorname{arccosh}(c*x)))/d/c^5/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))\dots
\end{aligned}$$

### 3.195.5 Fracas [F]

$$\int \frac{x^4(a + b\operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{(b\operatorname{arccosh}(cx) + a)^2x^4}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-(b^2*x^4*arccosh(c*x)^2 + 2*a*b*x^4*arccosh(c*x) + a^2*x^4)*sqrt(-c^2*d*x^2 + d)/(c^2*d*x^2 - d), x)`

**3.195.6 Sympy [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))^2}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate(x**4*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**4*(a + b*acosh(c*x))**2/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.195.7 Maxima [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^4}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/8*a^2*(2*sqrt(-c^2*d*x^2 + d)*x^3/(c^2*d) + 3*sqrt(-c^2*d*x^2 + d)*x/(c^4*d) - 3*arcsin(c*x)/(c^5*sqrt(d))) + integrate(b^2*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/sqrt(-c^2*d*x^2 + d) + 2*a*b*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/sqrt(-c^2*d*x^2 + d), x)`

**3.195.8 Giac [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^4}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x^4/sqrt(-c^2*d*x^2 + d), x)`

**3.195.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4 (a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^4 (a + b \operatorname{acosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^4*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2), x)`output `int((x^4*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2), x)`

**3.196**  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$

3.196.1 Optimal result . . . . . 1718  
 3.196.2 Mathematica [A] (verified) . . . . . 1719  
 3.196.3 Rubi [A] (verified) . . . . . 1719  
 3.196.4 Maple [B] (verified) . . . . . 1722  
 3.196.5 Fricas [A] (verification not implemented) . . . . . 1723  
 3.196.6 Sympy [F] . . . . . 1724  
 3.196.7 Maxima [A] (verification not implemented) . . . . . 1724  
 3.196.8 Giac [F(-2)] . . . . . 1725  
 3.196.9 Mupad [F(-1)] . . . . . 1725

**3.196.1 Optimal result**

Integrand size = 29, antiderivative size = 292

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = -\frac{4abx\sqrt{-1 + cx}\sqrt{1 + cx}}{3c^3\sqrt{d - c^2dx^2}} - \frac{40b^2(1 - cx)(1 + cx)}{27c^4\sqrt{d - c^2dx^2}} - \frac{2b^2x^2(1 - cx)(1 + cx)}{27c^2\sqrt{d - c^2dx^2}} - \frac{4b^2x\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arccosh}(cx)}{3c^3\sqrt{d - c^2dx^2}} - \frac{2bx^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{9c\sqrt{d - c^2dx^2}} - \frac{2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{3c^4d} - \frac{x^2\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{3c^2d}$$

```
output -40/27*b^2*(-c*x+1)*(c*x+1)/c^4/(-c^2*d*x^2+d)^(1/2)-2/27*b^2*x^2*(-c*x+1)
*(c*x+1)/c^2/(-c^2*d*x^2+d)^(1/2)-4/3*a*b*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^
3/(-c^2*d*x^2+d)^(1/2)-4/3*b^2*x*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/
c^3/(-c^2*d*x^2+d)^(1/2)-2/9*b*x^3*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)
)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)-2/3*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/
2)/c^4/d-1/3*x^2*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/c^2/d
```

**3.196.2 Mathematica [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 201, normalized size of antiderivative = 0.69

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{\sqrt{d - c^2 dx^2}(6abcx\sqrt{-1 + cx}\sqrt{1 + cx}(6 + c^2 x^2) - 9a^2(-2 + c^2 x^2 + c^4 x^4) - 2b^2(-20 + 19c^2 x^2 + c^4 x^4) + 27c^4 d(-1 + cx)(1 + cx))}{27c^4 d(-1 + cx)(1 + cx)}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`output `(Sqrt[d - c^2*d*x^2]*(6*a*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(6 + c^2*x^2) - 9*a^2*(-2 + c^2*x^2 + c^4*x^4) - 2*b^2*(-20 + 19*c^2*x^2 + c^4*x^4) + 6*b*(b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(6 + c^2*x^2) - 3*a*(-2 + c^2*x^2 + c^4*x^4))*ArcCosh[c*x] - 9*b^2*(-2 + c^2*x^2 + c^4*x^4)*ArcCosh[c*x]^2))/ (27*c^4*d*(-1 + c*x)*(1 + c*x))`**3.196.3 Rubi [A] (verified)**Time = 0.88 (sec) , antiderivative size = 271, normalized size of antiderivative = 0.93, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.241$ , Rules used = {6353, 6298, 111, 27, 83, 6329, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6353}$$

$$-\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int x^2(a + \operatorname{barccosh}(cx))dx}{3c\sqrt{d - c^2 dx^2}} + \frac{2 \int \frac{x(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{3c^2} -$$

$$\frac{x^2\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{3c^2 d}$$

$$\downarrow \text{6298}$$

$$\frac{2 \int \frac{x(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{3c^2} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a + \operatorname{barccosh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{3c\sqrt{d - c^2 dx^2}} -$$

$$\frac{x^2\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{3c^2 d}$$

---

3.196.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$



$$\begin{aligned}
& \downarrow 111 \\
& \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} \\
& \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \\
& \downarrow 27 \\
& \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} \\
& \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \\
& \downarrow 83 \\
& \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} \\
& \downarrow 6329 \\
& \frac{2 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) dx}{c\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{c^2d} \right)}{3c^2} - \\
& \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} \\
& \downarrow 2009 \\
& -\frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} + \\
& \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} (ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c})}{c\sqrt{d-c^2dx^2}} \right)}{3c^2} \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+\operatorname{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}}
\end{aligned}$$

input `Int[(x^3*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `-1/3*(x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/3*(b*c*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2)))) + (x^3*(a + b*ArcCosh[c*x]))/3)/(3*c*Sqrt[d - c^2*d*x^2]) + (2*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d)) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/c + b*x*ArcCosh[c*x]))/(c*Sqrt[d - c^2*d*x^2])))/(3*c^2)`

### 3.196.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 111 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e
_.)*(x_)^2)^(p_), x_Symbol] :> Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2
*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x]
- Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)
^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*Ar
cCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*
d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.196.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 751 vs.  $2(252) = 504$ .

Time = 0.94 (sec) , antiderivative size = 752, normalized size of antiderivative = 2.58

method	result
default	$a^2 \left( -\frac{x^2 \sqrt{-c^2 d x^2 + d}}{3c^2 d} - \frac{2\sqrt{-c^2 d x^2 + d}}{3d c^4} \right) + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (4c^4 x^4 - 5c^2 x^2 + 4\sqrt{cx-1} \sqrt{cx+1} c^3 x^3 - 3\sqrt{cx-1} \sqrt{cx+1} cx + 216c^4 d(c^2 x^2 - 1))}{216c^4 d(c^2 x^2 - 1)} \right)$
parts	$a^2 \left( -\frac{x^2 \sqrt{-c^2 d x^2 + d}}{3c^2 d} - \frac{2\sqrt{-c^2 d x^2 + d}}{3d c^4} \right) + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (4c^4 x^4 - 5c^2 x^2 + 4\sqrt{cx-1} \sqrt{cx+1} c^3 x^3 - 3\sqrt{cx-1} \sqrt{cx+1} cx + 216c^4 d(c^2 x^2 - 1))}{216c^4 d(c^2 x^2 - 1)} \right)$

```
input int(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output

$$\begin{aligned}
& a^2 * (-1/3 * x^2 / c^2 / d * (-c^2 * d * x^2 + d)^{(1/2)} - 2/3 / d / c^4 * (-c^2 * d * x^2 + d)^{(1/2)}) + b \\
& ^2 * (-1/216 * (-d * (c^2 * x^2 - 1))^{(1/2)} * (4 * c^4 * x^4 - 5 * c^2 * x^2 + 4 * (c * x - 1)^{(1/2)} * (c * \\
& x + 1)^{(1/2)} * c^3 * x^3 - 3 * (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c * x + 1) * (9 * \operatorname{arccosh}(c * x)^2 - \\
& 6 * \operatorname{arccosh}(c * x) + 2) / c^4 / d / (c^2 * x^2 - 1) - 3/8 * (-d * (c^2 * x^2 - 1))^{(1/2)} * ((c * x - 1)^{(1/2)} * \\
& (c * x + 1)^{(1/2)} * c * x + c^2 * x^2 - 1) * (\operatorname{arccosh}(c * x)^2 - 2 * \operatorname{arccosh}(c * x) + 2) / c^4 / d / ( \\
& c^2 * x^2 - 1) - 3/8 * (-d * (c^2 * x^2 - 1))^{(1/2)} * (- (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c * x + c^2 * \\
& x^2 - 1) * (\operatorname{arccosh}(c * x)^2 + 2 * \operatorname{arccosh}(c * x) + 2) / c^4 / d / (c^2 * x^2 - 1) - 1/216 * (-d * (c^2 * \\
& x^2 - 1))^{(1/2)} * (-4 * (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c^3 * x^3 + 4 * c^4 * x^4 + 3 * (c * x - 1)^{(1/2)} * \\
& (c * x + 1)^{(1/2)} * c * x - 5 * c^2 * x^2 + 1) * (9 * \operatorname{arccosh}(c * x)^2 + 6 * \operatorname{arccosh}(c * x) + 2) \\
& / c^4 / d / (c^2 * x^2 - 1) + 2 * a * b * (-1/72 * (-d * (c^2 * x^2 - 1))^{(1/2)} * (4 * c^4 * x^4 - 5 * c^2 * x^2 + 4 * (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c^3 * x^3 - 3 * (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c * x + 1) * (-1 + 3 * \operatorname{arccosh}(c * x)) / c^4 / d / (c^2 * x^2 - 1) - 3/8 * (-d * (c^2 * x^2 - 1))^{(1/2)} * ((c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c * x + c^2 * x^2 - 1) * (-1 + \operatorname{arccosh}(c * x)) / c^4 / d / (c^2 * x^2 - 1) - 3/8 * (-d * (c^2 * x^2 - 1))^{(1/2)} * (- (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c * x + c^2 * x^2 - 1) * (1 + \operatorname{arccosh}(c * x)) / c^4 / d / (c^2 * x^2 - 1) - 1/72 * (-d * (c^2 * x^2 - 1))^{(1/2)} * (-4 * (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c^3 * x^3 + 4 * c^4 * x^4 + 3 * (c * x - 1)^{(1/2)} * (c * x + 1)^{(1/2)} * c * x - 5 * c^2 * x^2 + 1) * (1 + 3 * \operatorname{arccosh}(c * x)) / c^4 / d / (c^2 * x^2 - 1)
\end{aligned}$$

### 3.196.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 282, normalized size of antiderivative = 0.97

$$\int \frac{x^3 (a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \frac{9(b^2 c^4 x^4 + b^2 c^2 x^2 - 2b^2) \sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1})^2 - 6(abc^3 x^3 + 6abcx) \sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1}}{\dots}$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output

$$\begin{aligned}
& -1/27 * (9 * (b^2 * c^4 * x^4 + b^2 * c^2 * x^2 - 2 * b^2) * \operatorname{sqrt}(-c^2 * d * x^2 + d) * \log(c * x \\
& + \operatorname{sqrt}(c^2 * x^2 - 1))^{(2)} - 6 * (a * b * c^3 * x^3 + 6 * a * b * c * x) * \operatorname{sqrt}(-c^2 * d * x^2 + d) * \\
& \operatorname{sqrt}(c^2 * x^2 - 1) - 6 * ((b^2 * c^3 * x^3 + 6 * b^2 * c * x) * \operatorname{sqrt}(-c^2 * d * x^2 + d) * \operatorname{sqrt} \\
& (c^2 * x^2 - 1) - 3 * (a * b * c^4 * x^4 + a * b * c^2 * x^2 - 2 * a * b) * \operatorname{sqrt}(-c^2 * d * x^2 + d) \\
& ) * \log(c * x + \operatorname{sqrt}(c^2 * x^2 - 1)) + ((9 * a^2 + 2 * b^2) * c^4 * x^4 + (9 * a^2 + 38 * b^2) * c^2 * x^2 - 18 * a^2 - 40 * b^2) * \operatorname{sqrt}(-c^2 * d * x^2 + d)) / (c^6 * d * x^2 - c^4 * d)
\end{aligned}$$

**3.196.6 Sympy [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^2}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate(x**3*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(1/2), x)`

output `Integral(x**3*(a + b*acosh(c*x))**2/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.196.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.96

$$\begin{aligned} & \int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx \\ &= -\frac{1}{3}b^2 \left( \frac{\sqrt{-c^2dx^2 + dx^2}}{c^2d} + \frac{2\sqrt{-c^2dx^2 + d}}{c^4d} \right) \operatorname{arcosh}(cx)^2 \\ & \quad - \frac{2}{3}ab \left( \frac{\sqrt{-c^2dx^2 + dx^2}}{c^2d} + \frac{2\sqrt{-c^2dx^2 + d}}{c^4d} \right) \operatorname{arcosh}(cx) \\ & \quad - \frac{1}{3}a^2 \left( \frac{\sqrt{-c^2dx^2 + dx^2}}{c^2d} + \frac{2\sqrt{-c^2dx^2 + d}}{c^4d} \right) \\ & \quad - \frac{2}{27}b^2 \left( \frac{\sqrt{c^2x^2 - 1}\sqrt{-dx^2} + \frac{20\sqrt{c^2x^2 - 1}\sqrt{-d}}{c^2}}{c^2d} - \frac{3(c^2\sqrt{-dx^3} + 6\sqrt{-dx}) \operatorname{arcosh}(cx)}{c^3d} \right) \\ & \quad + \frac{2(c^2\sqrt{-dx^3} + 6\sqrt{-dx})ab}{9c^3d} \end{aligned}$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2), x, algorithm="maxima")`

output `-1/3*b^2*(sqrt(-c^2*d*x^2 + d)*x^2/(c^2*d) + 2*sqrt(-c^2*d*x^2 + d)/(c^4*d))*arccosh(c*x)^2 - 2/3*a*b*(sqrt(-c^2*d*x^2 + d)*x^2/(c^2*d) + 2*sqrt(-c^2*d*x^2 + d)/(c^4*d))*arccosh(c*x) - 1/3*a^2*(sqrt(-c^2*d*x^2 + d)*x^2/(c^2*d) + 2*sqrt(-c^2*d*x^2 + d)/(c^4*d)) - 2/27*b^2*((sqrt(c^2*x^2 - 1)*sqrt(-d)*x^2 + 20*sqrt(c^2*x^2 - 1)*sqrt(-d)/c^2)/(c^2*d) - 3*(c^2*sqrt(-d)*x^3 + 6*sqrt(-d)*x)*arccosh(c*x)/(c^3*d) + 2/9*(c^2*sqrt(-d)*x^3 + 6*sqrt(-d)*x)*a*b/(c^3*d)`

**3.196.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.196.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^3*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^3*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2), x)`

**3.197**  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$

3.197.1 Optimal result . . . . . 1726  
 3.197.2 Mathematica [A] (verified) . . . . . 1727  
 3.197.3 Rubi [A] (verified) . . . . . 1727  
 3.197.4 Maple [B] (verified) . . . . . 1730  
 3.197.5 Fricas [F] . . . . . 1730  
 3.197.6 Sympy [F] . . . . . 1731  
 3.197.7 Maxima [F] . . . . . 1731  
 3.197.8 Giac [F] . . . . . 1731  
 3.197.9 Mupad [F(-1)] . . . . . 1732

**3.197.1 Optimal result**

Integrand size = 29, antiderivative size = 226

$$\int \frac{x^2(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = -\frac{b^2x(1 - cx)(1 + cx)}{4c^2\sqrt{d - c^2dx^2}} + \frac{b^2\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arccosh}(cx)}{4c^3\sqrt{d - c^2dx^2}} - \frac{bx^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{2c\sqrt{d - c^2dx^2}} - \frac{x\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d - c^2dx^2}}$$

output

```
-1/4*b^2*x*(-c*x+1)*(c*x+1)/c^2/(-c^2*d*x^2+d)^(1/2)+1/4*b^2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/(-c^2*d*x^2+d)^(1/2)-1/2*b*x^2*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)+1/6*(a+b*arccosh(c*x))^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^3/(-c^2*d*x^2+d)^(1/2)-1/2*x*(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/c^2/d
```

### 3.197.2 Mathematica [A] (verified)

Time = 1.30 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.01

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{-\frac{12a^2 cx \sqrt{d - c^2 dx^2}}{d} - \frac{12a^2 \arctan\left(\frac{cx \sqrt{d - c^2 dx^2}}{\sqrt{d}(-1 + c^2 x^2)}\right)}{\sqrt{d}} + \frac{b^2 \sqrt{\frac{-1 + cx}{1 + cx}}(1 + cx)(4\operatorname{arccosh}(cx)^3 - 6\operatorname{arccosh}(cx) \cosh(2\operatorname{arccosh}(cx)) + (3 + 6\operatorname{arccosh}(cx)^2) \sinh(2\operatorname{arccosh}(cx))))}{\sqrt{d - c^2 dx^2}}}{24c^3}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2], x]`

output `((-12*a^2*c*x*Sqrt[d - c^2*d*x^2])/d - (12*a^2*ArcTan[(c*x*Sqrt[d - c^2*d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))])/Sqrt[d] + (b^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(4*ArcCosh[c*x]^3 - 6*ArcCosh[c*x]*Cosh[2*ArcCosh[c*x]] + (3 + 6*ArcCosh[c*x]^2)*Sinh[2*ArcCosh[c*x]]))/Sqrt[d - c^2*d*x^2] + (6*a*b*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(-Cosh[2*ArcCosh[c*x]] + 2*ArcCosh[c*x]*(ArcCosh[c*x] + Sinh[2*ArcCosh[c*x]])))/Sqrt[d - c^2*d*x^2])/(24*c^3)`

### 3.197.3 Rubi [A] (verified)

Time = 0.69 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.84, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.172$ , Rules used = {6353, 6298, 101, 43, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6353}$$

$$-\frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int x(a + \operatorname{barccosh}(cx)) dx}{c\sqrt{d - c^2 dx^2}} + \frac{\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{2c^2} - \frac{x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{2c^2 d}$$

$$\downarrow \text{6298}$$



$$\begin{aligned}
& \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+\operatorname{arccosh}(cx))-\frac{1}{2}bc\int\frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}}dx\right)}{c\sqrt{d-c^2dx^2}} + \frac{\int\frac{(a+\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}}dx}{2c^2} \\
& \quad \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} \\
& \quad \downarrow 101 \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+\operatorname{arccosh}(cx))-\frac{1}{2}bc\left(\frac{\int\frac{1}{\sqrt{cx-1}\sqrt{cx+1}}dx}{2c^2}+\frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)\right)}{c\sqrt{d-c^2dx^2}} + \\
& \quad \frac{\int\frac{(a+\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}}dx}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} \\
& \quad \downarrow 43 \\
& \frac{\int\frac{(a+\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}}dx}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} - \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+\operatorname{arccosh}(cx))-\frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}+\frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)\right)}{c\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 6307 \\
& \frac{-\frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}}}{c\sqrt{d-c^2dx^2}} - \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+\operatorname{arccosh}(cx))-\frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}+\frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)\right)}{c\sqrt{d-c^2dx^2}}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `-1/2*(x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^3)/(6*b*c^3*Sqrt[d - c^2*d*x^2]) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/2)/(c*Sqrt[d - c^2*d*x^2])`

## 3.197.3.1 Defintions of rubi rules used

- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_) + (b_)*(x_))^(n_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 6298 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6307 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`
- rule 6353 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

**3.197.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 562 vs.  $2(194) = 388$ .

Time = 0.84 (sec) , antiderivative size = 563, normalized size of antiderivative = 2.49

method	result
default	$-\frac{a^2 x \sqrt{-c^2 d x^2 + d}}{2c^2 d} + \frac{a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{2c^2 \sqrt{c^2 d}} + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^3}{6d c^3 (c^2 x^2 - 1)} - \frac{\sqrt{-d(c^2 x^2 - 1)} (2c^3 x}{6d c^3 (c^2 x^2 - 1)} \right)$
parts	$-\frac{a^2 x \sqrt{-c^2 d x^2 + d}}{2c^2 d} + \frac{a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{2c^2 \sqrt{c^2 d}} + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^3}{6d c^3 (c^2 x^2 - 1)} - \frac{\sqrt{-d(c^2 x^2 - 1)} (2c^3 x}{6d c^3 (c^2 x^2 - 1)} \right)$

input `int(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/2*a^2*x/c^2/d*(-c^2*d*x^2+d)^(1/2)+1/2*a^2/c^2/(c^2*d)^(1/2)*\arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))+b^2*(-1/6*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/c^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)^3-1/16*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(2*\operatorname{arccosh}(c*x)^2-2*\operatorname{arccosh}(c*x)+1)/d/c^3/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)-2*c*x)*(2*\operatorname{arccosh}(c*x)^2+2*\operatorname{arccosh}(c*x)+1)/d/c^3/(c^2*x^2-1)+2*a*b*(-1/4*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/c^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)^2-1/16*(-d*(c^2*x^2-1))^(1/2)*(2*c^3*x^3-2*c*x+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(-1+2*\operatorname{arccosh}(c*x))/d/c^3/(c^2*x^2-1)-1/16*(-d*(c^2*x^2-1))^(1/2)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+2*c^3*x^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)-2*c*x)*(1+2*\operatorname{arccosh}(c*x))/d/c^3/(c^2*x^2-1)) \end{aligned}$$

**3.197.5 Fracas [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-(b^2*x^2*arccosh(c*x)^2 + 2*a*b*x^2*arccosh(c*x) + a^2*x^2)*sqrt(-c^2*d*x^2 + d)/(c^2*d*x^2 - d), x)`

---

3.197. 
$$\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$$

**3.197.6 Sympy [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^2}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**2*(a + b*acosh(c*x))**2/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.197.7 Maxima [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^2}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/2*a^2*(sqrt(-c^2*d*x^2 + d)*x/(c^2*d) - arcsin(c*x)/(c^3*sqrt(d))) + integrate(b^2*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/sqrt(-c^2*d*x^2 + d) + 2*a*b*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/sqrt(-c^2*d*x^2 + d), x)`

**3.197.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^2}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x^2/sqrt(-c^2*d*x^2 + d), x)`

**3.197.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^2*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2), x)`output `int((x^2*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2), x)`

**3.198**  $\int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$

3.198.1 Optimal result . . . . . 1733  
 3.198.2 Mathematica [A] (verified) . . . . . 1733  
 3.198.3 Rubi [A] (verified) . . . . . 1734  
 3.198.4 Maple [B] (verified) . . . . . 1735  
 3.198.5 Fricas [A] (verification not implemented) . . . . . 1736  
 3.198.6 Sympy [F] . . . . . 1736  
 3.198.7 Maxima [A] (verification not implemented) . . . . . 1736  
 3.198.8 Giac [F] . . . . . 1737  
 3.198.9 Mupad [F(-1)] . . . . . 1737

**3.198.1 Optimal result**

Integrand size = 27, antiderivative size = 155

$$\int \frac{x(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = -\frac{2abx\sqrt{-1 + cx}\sqrt{1 + cx}}{c\sqrt{d - c^2dx^2}} - \frac{2b^2(1 - cx)(1 + cx)}{c^2\sqrt{d - c^2dx^2}} - \frac{2b^2x\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arccosh}(cx)}{c\sqrt{d - c^2dx^2}} - \frac{\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{c^2d}$$

output

```
-2*b^2*(-c*x+1)*(c*x+1)/c^2/(-c^2*d*x^2+d)^(1/2)-2*a*b*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)-2*b^2*x*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(-c^2*d*x^2+d)^(1/2)-(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/c^2/d
```

**3.198.2 Mathematica [A] (verified)**

Time = 0.74 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.96

$$\int \frac{x(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{d - c^2dx^2}(2abcx\sqrt{-1 + cx}\sqrt{1 + cx} + a^2(1 - c^2x^2) - 2b^2(-1 + c^2x^2) + 2b(a - ac^2x^2 + bcx\sqrt{-1 + cx}\sqrt{1 + cx}))}{c^2d(-1 + cx)(1 + cx)}$$

input `Integrate[(x*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `(Sqrt[d - c^2*d*x^2]*(2*a*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] + a^2*(1 - c^2*x^2) - 2*b^2*(-1 + c^2*x^2) + 2*b*(a - a*c^2*x^2 + b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])*ArcCosh[c*x] + b^2*(1 - c^2*x^2)*ArcCosh[c*x]^2))/(c^2*d*(-1 + c*x)*(1 + c*x))`

### 3.198.3 Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.70, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$ , Rules used = {6329, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6329}$$

$$-\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int (a + \operatorname{arccosh}(cx)) dx}{c\sqrt{d - c^2 dx^2}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2}{c^2 d}$$

$$\downarrow \text{2009}$$

$$-\frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2}{c^2 d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax + b \operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d - c^2 dx^2}}$$

input `Int[(x*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d)) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(c*Sqrt[d - c^2*d*x^2])`

### 3.198.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.198.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 313 vs. 2(139) = 278.

Time = 0.59 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.03

method	result
default	$-\frac{a^2\sqrt{-c^2dx^2+d}}{c^2d} + b^2 \left( -\frac{\sqrt{-d(c^2x^2-1)}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)(\operatorname{arccosh}(cx)^2-2\operatorname{arccosh}(cx)+2)}{2c^2d(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}}{c^2d} \right)$
parts	$-\frac{a^2\sqrt{-c^2dx^2+d}}{c^2d} + b^2 \left( -\frac{\sqrt{-d(c^2x^2-1)}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)(\operatorname{arccosh}(cx)^2-2\operatorname{arccosh}(cx)+2)}{2c^2d(c^2x^2-1)} - \frac{\sqrt{-d(c^2x^2-1)}}{c^2d} \right)$

input `int(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output `-a^2/c^2/d*(-c^2*d*x^2+d)^(1/2)+b^2*(-1/2*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*arccosh(c*x)+2)/c^2/d/(c^2*x^2-1)-1/2*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(arccosh(c*x)^2+2*arccosh(c*x)+2)/c^2/d/(c^2*x^2-1))+2*a*b*(-1/2*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-1+arccosh(c*x))/c^2/d/(c^2*x^2-1)-1/2*(-d*(c^2*x^2-1))^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(1+arccosh(c*x))/c^2/d/(c^2*x^2-1))`



**3.198.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.41

$$\int \frac{x(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{2\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1}abcx - (b^2 c^2 x^2 - b^2)\sqrt{-c^2 dx^2 + d} \log(cx + \sqrt{c^2 x^2 - 1})^2 + 2(\sqrt{-c^2 dx^2 + d}\sqrt{c^2 x^2 - 1})}{c^4 dx^2}$$

```
input integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")
```

```
output (2*sqrt(-c^2*d*x^2 + d)*sqrt(c^2*x^2 - 1)*a*b*c*x - (b^2*c^2*x^2 - b^2)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1))^2 + 2*(sqrt(-c^2*d*x^2 + d))*sqrt(c^2*x^2 - 1)*b^2*c*x - (a*b*c^2*x^2 - a*b)*sqrt(-c^2*d*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) - ((a^2 + 2*b^2)*c^2*x^2 - a^2 - 2*b^2)*sqrt(-c^2*d*x^2 + d))/(c^4*d*x^2 - c^2*d)
```

**3.198.6 Sympy [F]**

$$\int \frac{x(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^2}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

```
input integrate(x*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(1/2),x)
```

```
output Integral(x*(a + b*acosh(c*x))**2/sqrt(-d*(c*x - 1)*(c*x + 1)), x)
```

**3.198.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.94

$$\int \frac{x(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = 2b^2 \left( \frac{\sqrt{-d}x \operatorname{arccosh}(cx)}{cd} - \frac{\sqrt{c^2 x^2 - 1}\sqrt{-d}}{c^2 d} \right)$$

$$+ \frac{2ab\sqrt{-d}x}{cd} - \frac{\sqrt{-c^2 dx^2 + d}b^2 \operatorname{arccosh}(cx)}{c^2 d}$$

$$- \frac{2\sqrt{-c^2 dx^2 + d}ab \operatorname{arccosh}(cx)}{c^2 d} - \frac{\sqrt{-c^2 dx^2 + d}a^2}{c^2 d}$$

---

3.198.  $\int \frac{x(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$

input `integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `2*b^2*(sqrt(-d)*x*arccosh(c*x)/(c*d) - sqrt(c^2*x^2 - 1)*sqrt(-d)/(c^2*d) + 2*a*b*sqrt(-d)*x/(c*d) - sqrt(-c^2*d*x^2 + d)*b^2*arccosh(c*x)^2/(c^2*d) - 2*sqrt(-c^2*d*x^2 + d)*a*b*arccosh(c*x)/(c^2*d) - sqrt(-c^2*d*x^2 + d)*a^2/(c^2*d)`

### 3.198.8 Giac [F]

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x/sqrt(-c^2*d*x^2 + d), x)`

### 3.198.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2),x)`

output `int((x*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(1/2), x)`

**3.199**  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx$

3.199.1 Optimal result . . . . .	1738
3.199.2 Mathematica [A] (verified) . . . . .	1738
3.199.3 Rubi [A] (verified) . . . . .	1739
3.199.4 Maple [B] (verified) . . . . .	1739
3.199.5 Fricas [F] . . . . .	1740
3.199.6 Sympy [F] . . . . .	1740
3.199.7 Maxima [F] . . . . .	1741
3.199.8 Giac [F] . . . . .	1741
3.199.9 Mupad [F(-1)] . . . . .	1741

**3.199.1 Optimal result**

Integrand size = 26, antiderivative size = 53

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^3}{3bc\sqrt{d - c^2dx^2}}$$

output  $1/3*(a+b*\operatorname{arccosh}(c*x))^3*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c/(-c^2*d*x^2+d)^{(1/2)}$

**3.199.2 Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^3}{3bc\sqrt{d - c^2dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/Sqrt[d - c^2*d*x^2],x]`

output  $(\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(a + b*\operatorname{ArcCosh}[c*x])^3)/(3*b*c*\operatorname{Sqrt}[d - c^2*d*x^2])$

### 3.199.3 Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6307

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1}(a + \operatorname{arccosh}(cx))^3}{3bc\sqrt{d - c^2 dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/Sqrt[d - c^2*d*x^2],x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^3)/(3*b*c*Sqrt[d - c^2*d*x^2])`

#### 3.199.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^ (n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.199.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. 2(45) = 90.

Time = 0.48 (sec) , antiderivative size = 149, normalized size of antiderivative = 2.81

method	result
default	$\frac{a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{\sqrt{c^2 d}} - \frac{b^2 \sqrt{-d(cx-1)(cx+1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^3}{3d(c^2 x^2 - 1)c} - \frac{ab \sqrt{-d(cx-1)(cx+1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)}{d(c^2 x^2 - 1)c}$
parts	$\frac{a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{\sqrt{c^2 d}} - \frac{b^2 \sqrt{-d(cx-1)(cx+1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)^3}{3d(c^2 x^2 - 1)c} - \frac{ab \sqrt{-d(cx-1)(cx+1)} \sqrt{cx-1} \sqrt{cx+1} \operatorname{arccosh}(cx)}{d(c^2 x^2 - 1)c}$

3.199.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2 dx^2}} dx$

input `int((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output `a^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1/2))-1/3*b^2*(-d*(c*x-1)*(c*x+1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)/c*arccosh(c*x)^3-a*b*(-d*(c*x-1)*(c*x+1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)/c*arccosh(c*x)^2`

### 3.199.5 Fricas [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^2*d*x^2 - d), x)`

### 3.199.6 Sympy [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**2/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.199.7 Maxima [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `a^2*arcsin(c*x)/(c*sqrt(d)) + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/sqrt(-c^2*d*x^2 + d) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/sqrt(-c^2*d*x^2 + d), x)`

**3.199.8 Giac [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/sqrt(-c^2*d*x^2 + d), x)`

**3.199.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^2/(d - c^2*d*x^2)^(1/2),x)`

output `int((a + b*acosh(c*x))^2/(d - c^2*d*x^2)^(1/2), x)`

### 3.200 $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx$

3.200.1 Optimal result . . . . .	1742
3.200.2 Mathematica [A] (verified) . . . . .	1743
3.200.3 Rubi [A] (verified) . . . . .	1743
3.200.4 Maple [F] . . . . .	1745
3.200.5 Fracas [F] . . . . .	1746
3.200.6 Sympy [F] . . . . .	1746
3.200.7 Maxima [F] . . . . .	1746
3.200.8 Giac [F] . . . . .	1747
3.200.9 Mupad [F(-1)] . . . . .	1747

#### 3.200.1 Optimal result

Integrand size = 29, antiderivative size = 273

$$\int \frac{(a + b\operatorname{arccosh}(cx))^2}{x\sqrt{d - c^2dx^2}} dx$$

$$= \frac{2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

$$- \frac{2ib\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

$$+ \frac{2ib\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

$$+ \frac{2ib^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

$$- \frac{2ib^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

output

```
2*(a+b*arccosh(c*x))^2*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)
*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)-2*I*b*(a+b*arccosh(c*x))*polylog(2,-
I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x
^2+d)^(1/2)+2*I*b*(a+b*arccosh(c*x))*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)
)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)+2*I*b^2*polylog
(3,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2
*d*x^2+d)^(1/2)-2*I*b^2*polylog(3,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*(c*
x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)
```

### 3.200.2 Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.15

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x \sqrt{d - c^2 dx^2}} dx = \frac{a^2 \log(cx)}{\sqrt{d}} - \frac{a^2 \log(d + \sqrt{d} \sqrt{d - c^2 dx^2})}{\sqrt{d}} - \frac{2iab \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (\operatorname{arccosh}(cx) (\log(1 - ie^{-\operatorname{arccosh}(cx)}) - \log(1 + ie^{-\operatorname{arccosh}(cx)})) + \operatorname{PolyLog}(2, -ie^{-\operatorname{arccosh}(cx)}))}{\sqrt{d - c^2 dx^2}} + \frac{ib^2 \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (-\operatorname{arccosh}(cx))^2 (\log(1 - ie^{-\operatorname{arccosh}(cx)}) - \log(1 + ie^{-\operatorname{arccosh}(cx)})) - 2 \operatorname{arccosh}(cx) (\operatorname{PolyLog}(2, -ie^{-\operatorname{arccosh}(cx)}) + \operatorname{PolyLog}(3, -ie^{-\operatorname{arccosh}(cx)})))}{\sqrt{d - c^2 dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x*Sqrt[d - c^2*d*x^2]),x]`

output `(a^2*Log[c*x])/Sqrt[d] - (a^2*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]])/Sqrt[d] - ((2*I)*a*b*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(ArcCosh[c*x]*(Log[1 - I/E^ArcCosh[c*x]] - Log[1 + I/E^ArcCosh[c*x]]) + PolyLog[2, (-I)/E^ArcCosh[c*x]] - PolyLog[2, I/E^ArcCosh[c*x]]))/Sqrt[d - c^2*d*x^2] + (I*b^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(-ArcCosh[c*x]^2*(Log[1 - I/E^ArcCosh[c*x]] - Log[1 + I/E^ArcCosh[c*x]]) - 2*ArcCosh[c*x]*(PolyLog[2, (-I)/E^ArcCosh[c*x]] - PolyLog[2, I/E^ArcCosh[c*x]]) - 2*PolyLog[3, (-I)/E^ArcCosh[c*x]] + 2*PolyLog[3, I/E^ArcCosh[c*x]]))/Sqrt[d - c^2*d*x^2]`

### 3.200.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.51, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.207$ , Rules used = {6361, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x \sqrt{d - c^2 dx^2}} dx$$

↓ 6361

$$\frac{\sqrt{cx - 1} \sqrt{cx + 1}}{\sqrt{d - c^2 dx^2}} \int \frac{(a + b \operatorname{arccosh}(cx))^2}{cx} d \operatorname{arccosh}(cx)$$

↓ 3042

---

3.200.  $\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x \sqrt{d - c^2 dx^2}} dx$



$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int (a + \operatorname{barccosh}(cx))^2 \csc\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{\sqrt{d-c^2} dx^2}$$

↓ 4668

$$\frac{\sqrt{cx-1}\sqrt{cx+1} (-2ib \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{\sqrt{d-c^2} dx^2}$$

↓ 3011

$$\frac{\sqrt{cx-1}\sqrt{cx+1} (2ib(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)))}{\sqrt{d-c^2} dx^2}$$

↓ 2720

$$\frac{\sqrt{cx-1}\sqrt{cx+1} (2ib(b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)))}{\sqrt{d-c^2} dx^2}$$

↓ 7143

$$\frac{\sqrt{cx-1}\sqrt{cx+1} (2 \arctan(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx))^2 + 2ib(b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx))))}{\sqrt{d-c^2} dx^2}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x*Sqrt[d - c^2*d*x^2]),x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]])))/Sqrt[d - c^2*d*x^2]`

### 3.200.3.1 Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))]*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

$$3.200. \quad \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2} dx^2} dx$$

```
rule 3011 Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4668 Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

```
rule 6361 Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*(x_)^(m_.))/Sqrt[(d_.) + (e_.)
*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x
]/Sqrt[d + e*x^2])] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]]
, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && Int
egerQ[m]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

### 3.200.4 Maple [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x\sqrt{-c^2dx^2 + d}} dx$$

```
input int((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(1/2),x)
```

```
output int((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(1/2),x)
```

**3.200.5 Fricas [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^2*d*x^3 - d*x), x)`

**3.200.6 Sympy [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d - c^2dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate((a+b*acosh(c*x))**2/x/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.200.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-a^2*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/sqrt(d) + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/(sqrt(-c^2*d*x^2 + d)*x) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(sqrt(-c^2*d*x^2 + d)*x), x)`

**3.200.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/(sqrt(-c^2*d*x^2 + d)*x), x)`

**3.200.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x\sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^2/(x*(d - c^2*d*x^2)^(1/2)),x)`

output `int((a + b*acosh(c*x))^2/(x*(d - c^2*d*x^2)^(1/2)), x)`

**3.201**  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^2\sqrt{d-c^2dx^2}} dx$

3.201.1 Optimal result . . . . . 1748  
 3.201.2 Mathematica [A] (verified) . . . . . 1749  
 3.201.3 Rubi [C] (warning: unable to verify) . . . . . 1749  
 3.201.4 Maple [B] (verified) . . . . . 1752  
 3.201.5 Fricas [F] . . . . . 1753  
 3.201.6 Sympy [F] . . . . . 1753  
 3.201.7 Maxima [F] . . . . . 1754  
 3.201.8 Giac [F] . . . . . 1754  
 3.201.9 Mupad [F(-1)] . . . . . 1754

**3.201.1 Optimal result**

Integrand size = 29, antiderivative size = 186

$$\int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^2\sqrt{d - c^2dx^2}} dx$$

$$= -\frac{c\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))^2}{\sqrt{d - c^2dx^2}} - \frac{\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^2}{dx}$$

$$- \frac{2bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

$$+ \frac{b^2c\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}}$$

output

```
-c*(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)-2
*b*c*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)
)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)+b^2*c*polylog(2,-1/(c*x+(c*x-1)
)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(-c^2*d*x^2+d)^(1/2)-
(a+b*arccosh(c*x))^2*(-c^2*d*x^2+d)^(1/2)/d/x
```

**3.201.2 Mathematica [A] (verified)**

Time = 0.91 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.27

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx = -\frac{a^2 \sqrt{-d(-1 + c^2 x^2)}}{dx} - 2abc \left( \frac{\sqrt{d - c^2 dx^2} \operatorname{arccosh}(cx)}{cdx} + \frac{\sqrt{-1 + cx} \sqrt{1 + cx} (\log(-1 + \sqrt{1 + cx}) + \log(1 + \sqrt{1 + cx}))}{\sqrt{d - c^2 dx^2}} \right) + \frac{b^2 c \sqrt{\frac{-1 + cx}{1 + cx}} (1 + cx) \left( \operatorname{arccosh}(cx) \left( -\operatorname{arccosh}(cx) + \frac{\sqrt{\frac{-1 + cx}{1 + cx}} (1 + cx) \operatorname{arccosh}(cx)}{cx} - 2 \log(1 + e^{-2 \operatorname{arccosh}(cx)}) \right) \right)}{\sqrt{-d(-1 + cx)(1 + cx)}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^2*sqrt[d - c^2*d*x^2]),x]`output `-((a^2*sqrt[-(d*(-1 + c^2*x^2))])/(d*x)) - 2*a*b*c*((sqrt[d - c^2*d*x^2]*ArcCosh[c*x])/(c*d*x) + (sqrt[-1 + c*x]*sqrt[1 + c*x]*(Log[-1 + sqrt[1 + c*x]] + Log[1 + sqrt[1 + c*x]]))/sqrt[d - c^2*d*x^2]) + (b^2*c*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(ArcCosh[c*x]*(-ArcCosh[c*x] + (sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x])/(c*x) - 2*Log[1 + E^(-2*ArcCosh[c*x])])) + PolyLog[2, -E^(-2*ArcCosh[c*x])])/sqrt[-(d*(-1 + c*x)*(1 + c*x))]`**3.201.3 Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.74, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.310$ , Rules used = {6332, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx$$

↓ 6332

$$-\frac{2bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{arccosh}(cx)}{x} dx}{\sqrt{d - c^2 dx^2}} - \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{arccosh}(cx))^2}{dx}$$

↓ 6297

---

3.201.  $\int \frac{(a + \operatorname{arccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx$

$$\begin{aligned}
& \frac{2c\sqrt{cx-1}\sqrt{cx+1} \int -\left((a + \operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right)\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2} \frac{d}{dx} \frac{\sqrt{d-c^2dx^2}}{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}} \\
& \quad \downarrow 25 \\
& \frac{2c\sqrt{cx-1}\sqrt{cx+1} \int (a + \operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2} \frac{d}{dx} \frac{\sqrt{d-c^2dx^2}}{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}} \\
& \quad \downarrow 3042 \\
& \frac{2c\sqrt{cx-1}\sqrt{cx+1} \int -i(a + \operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2} \frac{d}{dx} \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2} +} \\
& \quad \downarrow 26 \\
& \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \int (a + \operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2} \frac{d}{dx} \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}} \\
& \quad \downarrow 4201 \\
& \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \left(2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1+e^{-2\operatorname{arccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i(a + \operatorname{barccosh}(cx))^2\right)}{\sqrt{d-c^2dx^2} \frac{d}{dx} \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}} \\
& \quad \downarrow 2620 \\
& \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \left(2i\left(\frac{1}{2}b \int \log(1 + e^{-2\operatorname{arccosh}(cx)}) d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx))\right)\right)}{\sqrt{d-c^2dx^2} \frac{d}{dx} \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}} \\
& \quad \downarrow 2715 \\
& \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \left(2i\left(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1 + e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)} - \frac{1}{2}b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx))\right)\right)}{\sqrt{d-c^2dx^2} \frac{d}{dx} \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}} \\
& \quad \downarrow 2838
\end{aligned}$$

---

3.201.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2\sqrt{d-c^2dx^2}} dx$

$$\frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{dx} - \frac{2ic\sqrt{cx - 1}\sqrt{cx + 1} \left( 2i \left( \frac{1}{4} b^2 \operatorname{PolyLog}(2, -a - \operatorname{barccosh}(cx)) \right) - \frac{1}{2} b \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx)) \right) - \frac{1}{2} i}{\sqrt{d - c^2 dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^2*Sqrt[d - c^2*d*x^2]),x]`

output `-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(d*x)) - ((2*I)*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4))/Sqrt[d - c^2*d*x^2]`

### 3.201.3.1 Defintions of rubi rules used

rule 25 `Int[-(F_x), x_Symbol] := Simp[Identity[-1] Int[F_x, x], x]`

rule 26 `Int[(Complex[0, a_]*(F_x), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`



rule 4201 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

### 3.201.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 475 vs. 2(192) = 384.

Time = 1.15 (sec) , antiderivative size = 476, normalized size of antiderivative = 2.56

method	result
default	$-\frac{a^2\sqrt{-c^2dx^2+d}}{dx} + b^2 \left( -\frac{\sqrt{-d(c^2x^2-1)}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\operatorname{arccosh}(cx)^2}{x(c^2x^2-1)d} - \frac{2\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}\operatorname{arccosh}(cx)}{d(c^2x^2-1)} \right)$
parts	$-\frac{a^2\sqrt{-c^2dx^2+d}}{dx} + b^2 \left( -\frac{\sqrt{-d(c^2x^2-1)}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\operatorname{arccosh}(cx)^2}{x(c^2x^2-1)d} - \frac{2\sqrt{-d(c^2x^2-1)}\sqrt{cx-1}\sqrt{cx+1}\operatorname{arccosh}(cx)}{d(c^2x^2-1)} \right)$

input `int((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-a^2/d/x*(-c^2*d*x^2+d)^{(1/2)}+b^2*(-(-d*(c^2*x^2-1))^{(1/2)}*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)*c*x+c^2*x^2-1}*arccosh(c*x)^2/x/(c^2*x^2-1)/d-2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)*arccosh(c*x)^2*c+2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)*arccosh(c*x)*ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*c+(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)*polylog(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*c)+2*a*b*(-2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)*arccosh(c*x)*c-(-d*(c^2*x^2-1))^{(1/2)}*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)*c*x+c^2*x^2-1}*arccosh(c*x)/x/(c^2*x^2-1)/d+(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d/(c^2*x^2-1)*ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*c)$$

### 3.201.5 Fracas [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^2*d*x^4 - d*x^2), x)`

### 3.201.6 Sympy [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^2 \sqrt{-d(cx-1)(cx+1)}} dx$$

input `integrate((a+b*acosh(c*x))**2/x**2/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x**2*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.201.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + dx^2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-(c^2*d*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2) + I*(-1)^(-2*c^2*d*x^2 + 2*d)*sqrt(d)*log(-2*c^2*d + 2*d/x^2))*a*b*c/d + b^2*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^(2/(sqrt(-c^2*d*x^2 + d)*x^2)), x) - 2*sqrt(-c^2*d*x^2 + d)*a*b*arccosh(c*x)/(d*x) - sqrt(-c^2*d*x^2 + d)*a^2/(d*x)`

**3.201.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + dx^2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/(sqrt(-c^2*d*x^2 + d)*x^2), x)`

**3.201.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^2*(d - c^2*d*x^2)^(1/2)),x)`

output `int((a + b*acosh(c*x))^2/(x^2*(d - c^2*d*x^2)^(1/2)), x)`

**3.202**      $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3\sqrt{d-c^2dx^2}} dx$

3.202.1 Optimal result . . . . . 1755  
 3.202.2 Mathematica [B] (warning: unable to verify) . . . . . 1756  
 3.202.3 Rubi [A] (verified) . . . . . 1756  
 3.202.4 Maple [F] . . . . . 1761  
 3.202.5 Fricas [F] . . . . . 1761  
 3.202.6 Sympy [F] . . . . . 1761  
 3.202.7 Maxima [F] . . . . . 1762  
 3.202.8 Giac [F] . . . . . 1762  
 3.202.9 Mupad [F(-1)] . . . . . 1762

**3.202.1 Optimal result**

Integrand size = 29, antiderivative size = 430

$$\begin{aligned} & \int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^3\sqrt{d - c^2dx^2}} dx \\ &= \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))}{x\sqrt{d - c^2dx^2}} - \frac{\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^2}{2dx^2} \\ &+ \frac{c^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} \\ &- \frac{b^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{\sqrt{d - c^2dx^2}} \\ &- \frac{ibc^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} \\ &+ \frac{ibc^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} \\ &+ \frac{ib^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} \\ &- \frac{ib^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{\sqrt{d - c^2dx^2}} \end{aligned}$$

output  $b*c*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/x/(-c^2*d*x^2+d)^{(1/2)}+c^2*(a+b*\operatorname{arccosh}(c*x))^2*\arctan(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(-c^2*d*x^2+d)^{(1/2)}-b^2*c^2*\arctan((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(-c^2*d*x^2+d)^{(1/2)}-I*b*c^2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(-c^2*d*x^2+d)^{(1/2)}+I*b*c^2*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(-c^2*d*x^2+d)^{(1/2)}+I*b^2*c^2*\operatorname{polylog}(3,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(-c^2*d*x^2+d)^{(1/2)}-I*b^2*c^2*\operatorname{polylog}(3,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(-c^2*d*x^2+d)^{(1/2)}-1/2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/d/x^2$

### 3.202.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 5036 vs.  $2(430) = 860$ .

Time = 63.66 (sec) , antiderivative size = 5036, normalized size of antiderivative = 11.71

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx = \text{Result too large to show}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^3*Sqrt[d - c^2*d*x^2]),x]`

output Result too large to show

### 3.202.3 Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.59, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.345$ , Rules used = {6347, 6298, 103, 218, 6361, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx$$

↓ 6347

---

3.202.  $\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx$

$$\begin{aligned}
& \frac{1}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^2} dx}{\sqrt{d-c^2dx^2}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2dx^2} \\
& \quad \downarrow \text{6298} \\
& \frac{1}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( bc \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{d-c^2dx^2}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2dx^2} \\
& \quad \downarrow \text{103} \\
& \frac{1}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx - \\
& \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( bc^2 \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{d-c^2dx^2}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2dx^2} \\
& \quad \downarrow \text{218} \\
& \frac{1}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx - \\
& \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{d-c^2dx^2}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2dx^2} \\
& \quad \downarrow \text{6361} \\
& \frac{c^2\sqrt{cx-1}\sqrt{cx+1} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} \operatorname{darccosh}(cx)}{2\sqrt{d-c^2dx^2}} - \\
& \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{d-c^2dx^2}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2dx^2} \\
& \quad \downarrow \text{3042} \\
& \frac{c^2\sqrt{cx-1}\sqrt{cx+1} \int (a + \operatorname{barccosh}(cx))^2 \csc\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{2\sqrt{d-c^2dx^2}} - \\
& \frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{\sqrt{d-c^2dx^2}} - \\
& \quad \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{2dx^2}
\end{aligned}$$

↓ 4668

$$\frac{c^2\sqrt{cx-1}\sqrt{cx+1}(-2ib \int (a + \operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a + \operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{2\sqrt{d-c^2dx^2}}$$

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\frac{\sqrt{d-c^2dx^2}}{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2} - \frac{2dx^2}}{2dx^2}}$$

↓ 3011

$$\frac{c^2\sqrt{cx-1}\sqrt{cx+1}(2ib(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx)))}{2\sqrt{d-c^2dx^2}}$$

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\frac{\sqrt{d-c^2dx^2}}{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2} - \frac{2dx^2}}{2dx^2}}$$

↓ 2720

$$\frac{c^2\sqrt{cx-1}\sqrt{cx+1}(2ib(b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + b)))}{2\sqrt{d-c^2dx^2}}$$

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\frac{\sqrt{d-c^2dx^2}}{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2} - \frac{2dx^2}}{2dx^2}}$$

↓ 7143

$$\frac{c^2\sqrt{cx-1}\sqrt{cx+1}(2 \arctan(e^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))^2 + 2ib(b \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)}) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a + \operatorname{barccosh}(cx))))}{2\sqrt{d-c^2dx^2}}$$

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(bc \arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x}\right)}{\frac{\sqrt{d-c^2dx^2}}{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2} - \frac{2dx^2}}{2dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^3*sqrt[d - c^2*d*x^2]),x]`

```
output -1/2*(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(d*x^2) - (b*c*Sqrt[-1 +
c*x]*Sqrt[1 + c*x]*(-(a + b*ArcCosh[c*x])/x) + b*c*ArcTan[Sqrt[-1 + c*x]
*Sqrt[1 + c*x]])/Sqrt[d - c^2*d*x^2] + (c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*
(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-(a + b*ArcCo
sh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*
x]]) - (2*I)*b*(-(a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*P
olyLog[3, I*E^ArcCosh[c*x]])))/(2*Sqrt[d - c^2*d*x^2])
```

### 3.202.3.1 Defintions of rubi rules used

```
rule 103 Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_
))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sq
rt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d
*e - f*(b*c + a*d), 0]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 3011 Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```



rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

rule 6361 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**3.202.4 Maple [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 \sqrt{-c^2 dx^2 + d}} dx$$

input `int((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(1/2),x)`

output `int((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(1/2),x)`

**3.202.5 Fricas [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + d} x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^2*d*x^5 - d*x^3), x)`

**3.202.6 Sympy [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^3 \sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))**2/x**3/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x**3*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.202.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + dx^3}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `-1/2*(c^2*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/sqrt(d) + sqrt(-c^2*d*x^2 + d)/(d*x^2))*a^2 + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^(2/(sqrt(-c^2*d*x^2 + d)*x^3) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(sqrt(-c^2*d*x^2 + d)*x^3), x)`

**3.202.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + dx^3}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/(sqrt(-c^2*d*x^2 + d)*x^3), x)`

**3.202.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^3 \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^3*(d - c^2*d*x^2)^(1/2)),x)`

output `int((a + b*acosh(c*x))^2/(x^3*(d - c^2*d*x^2)^(1/2)), x)`

### 3.203 $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^4\sqrt{d-c^2dx^2}} dx$

3.203.1 Optimal result . . . . .	1763
3.203.2 Mathematica [A] (warning: unable to verify) . . . . .	1764
3.203.3 Rubi [C] (warning: unable to verify) . . . . .	1764
3.203.4 Maple [B] (verified) . . . . .	1769
3.203.5 Fricas [F] . . . . .	1770
3.203.6 Sympy [F] . . . . .	1771
3.203.7 Maxima [F] . . . . .	1771
3.203.8 Giac [F] . . . . .	1771
3.203.9 Mupad [F(-1)] . . . . .	1772

#### 3.203.1 Optimal result

Integrand size = 29, antiderivative size = 328

$$\begin{aligned} & \int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^4\sqrt{d - c^2dx^2}} dx \\ &= \frac{b^2c^2(1 - cx)(1 + cx)}{3x\sqrt{d - c^2dx^2}} + \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))}{3x^2\sqrt{d - c^2dx^2}} \\ &\quad - \frac{2c^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))^2}{3\sqrt{d - c^2dx^2}} \\ &\quad - \frac{\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^2}{3dx^3} - \frac{2c^2\sqrt{d - c^2dx^2}(a + b\operatorname{arccosh}(cx))^2}{3dx} \\ &\quad - \frac{4bc^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{d - c^2dx^2}} \\ &\quad + \frac{2b^2c^3\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})}{3\sqrt{d - c^2dx^2}} \end{aligned}$$

output  $\frac{1}{3}b^2c^2(-cx+1)(cx+1)/x/(-c^2dx^2+d)^{(1/2)}+1/3b*c*(a+b*\operatorname{arccosh}(c*x))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/x^2/(-c^2dx^2+d)^{(1/2)}-2/3c^3*(a+b*\operatorname{arccosh}(c*x))^2*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/(-c^2dx^2+d)^{(1/2)}-4/3b*c^3*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)})^2*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/(-c^2dx^2+d)^{(1/2)}+2/3b^2*c^3*\operatorname{polylog}(2,-1/(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)})^2*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/(-c^2dx^2+d)^{(1/2)}-1/3*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2dx^2+d)^{(1/2)}/d/x^3-2/3c^2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2dx^2+d)^{(1/2)}/d/x$

**3.203.2 Mathematica [A] (warning: unable to verify)**

Time = 0.99 (sec) , antiderivative size = 370, normalized size of antiderivative = 1.13

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx$$

$$= \frac{-a^2 - a^2 c^2 x^2 + b^2 c^2 x^2 + 2a^2 c^4 x^4 - b^2 c^4 x^4 + abcx \sqrt{-1 + cx} \sqrt{1 + cx} - b^2(1 + cx) \left(1 - cx + 2c^2 x^2 + 2c^3 x^3\right)}{3x^3 \sqrt{d - c^2 dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^4*Sqrt[d - c^2*d*x^2]),x]`

output

```
(-a^2 - a^2*c^2*x^2 + b^2*c^2*x^2 + 2*a^2*c^4*x^4 - b^2*c^4*x^4 + a*b*c*x*
Sqrt[-1 + c*x]*Sqrt[1 + c*x] - b^2*(1 + c*x)*(1 - c*x + 2*c^2*x^2 + 2*c^3*
x^3*(-1 + Sqrt[(-1 + c*x)/(1 + c*x)]))*ArcCosh[c*x]^2 + b*(1 + c*x)*ArcCos
h[c*x]*(b*c*x*Sqrt[(-1 + c*x)/(1 + c*x)] + 2*a*(-1 + c*x - 2*c^2*x^2 + 2*c
^3*x^3) - 4*b*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*Log[1 + E^(-2*ArcCosh[c*x
])]) - 4*a*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[-1 + Sqrt[1 + c*x]]
- 4*a*b*c^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 + Sqrt[1 + c*x]] + 2*b^
2*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, -E^(-2*ArcCosh[c
*x])])/(3*x^3*Sqrt[d - c^2*d*x^2])
```

**3.203.3 Rubi [C] (warning: unable to verify)**

Result contains complex when optimal does not.

Time = 1.45 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.80, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$ , Rules used = {6347, 6298, 106, 6332, 6297, 25, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx$$

$$\downarrow 6347$$

$$\frac{2}{3}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 \sqrt{d - c^2 dx^2}} dx - \frac{2bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{x^3} dx}{3\sqrt{d - c^2 dx^2}} - \frac{\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{3dx^3}$$

---

3.203.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx$

$$\begin{aligned} & \downarrow 6298 \\ \frac{2}{3}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2\sqrt{d-c^2dx^2}} dx - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}bc \int \frac{1}{x^2\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} \right)}{3\sqrt{d-c^2dx^2} \sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2} - \frac{1}{3dx^3} \end{aligned}$$

$$\begin{aligned} & \downarrow 106 \\ \frac{2}{3}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2\sqrt{d-c^2dx^2}} dx - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2} \right)}{3\sqrt{d-c^2dx^2} \sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2} - \frac{1}{3dx^3} \end{aligned}$$

$$\begin{aligned} & \downarrow 6332 \\ \frac{2}{3}c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x} dx}{\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{dx} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2} \right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{3dx^3} \end{aligned}$$

$$\begin{aligned} & \downarrow 6297 \\ \frac{2}{3}c^2 \left( -\frac{2c\sqrt{cx-1}\sqrt{cx+1} \int -\left( (a + \operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) \right) d(a + \operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{3dx^3} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2} \right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{3dx^3} \end{aligned}$$

$$\begin{aligned} & \downarrow 25 \\ \frac{2}{3}c^2 \left( \frac{2c\sqrt{cx-1}\sqrt{cx+1} \int (a + \operatorname{barccosh}(cx)) \tanh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{3dx^3} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2} \right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a + \operatorname{barccosh}(cx))^2}{3dx^3} \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \end{aligned}$$

$$\frac{2}{3}c^2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{dx} + \frac{2c\sqrt{cx-1}\sqrt{cx+1} \int -i(a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) dx}{\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left(\frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2}\right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3dx^3} \right)$$

↓ 26

$$\frac{2}{3}c^2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{dx} - \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \tan\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right) dx}{\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left(\frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2}\right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3dx^3} \right)$$

↓ 4201

$$\frac{2}{3}c^2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{dx} - \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \left(2i \int \frac{e^{-2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1+e^{-2\operatorname{arccosh}(cx)}} d(a+\operatorname{barccosh}(cx))\right)}{\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left(\frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2}\right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3dx^3} \right)$$

↓ 2620

$$\frac{2}{3}c^2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{dx} - \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \left(2i \left(\frac{1}{2}b \int \log(1+e^{-2\operatorname{arccosh}(cx)}) d(a+\operatorname{barccosh}(cx))\right)\right)}{\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left(\frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2}\right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3dx^3} \right)$$

↓ 2715

$$\frac{2}{3}c^2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{dx} - \frac{2ic\sqrt{cx-1}\sqrt{cx+1} \left(2i \left(-\frac{1}{4}b^2 \int e^{2\operatorname{arccosh}(cx)} \log(1+e^{-2\operatorname{arccosh}(cx)}) de^{-2\operatorname{arccosh}(cx)}\right)\right)}{\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left(\frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2}\right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3dx^3} \right)$$

↓ 2838

---

3.203.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4\sqrt{d-c^2dx^2}} dx$

$$\frac{2}{3}c^2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{dx} - \frac{2ic\sqrt{cx-1}\sqrt{cx+1}(2i(\frac{1}{4}b^2 \operatorname{PolyLog}(2, -a-\operatorname{barccosh}(cx)) - \frac{1}{2}b \log(e^{\sqrt{d-c^2dx^2}}))}{\sqrt{d-c^2dx^2}} \right. \\ \left. - \frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} - \frac{a+\operatorname{barccosh}(cx)}{2x^2}\right)}{3\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3dx^3} \right)$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^4*sqrt[d - c^2*d*x^2]),x]`

output `-1/3*(sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(d*x^3) - (2*b*c*sqrt[-1 + c*x]*sqrt[1 + c*x]*((b*c*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*x) - (a + b*ArcCosh[c*x])/(2*x^2)))/(3*sqrt[d - c^2*d*x^2]) + (2*c^2*(-((sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(d*x)) - ((2*I)*c*sqrt[-1 + c*x]*sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^2 + (2*I)*(-1/2*(b*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x]])) + (b^2*PolyLog[2, -a - b*ArcCosh[c*x]])/4)))/sqrt[d - c^2*d*x^2]))/3`

### 3.203.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 106 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1), 0] && NeQ[m, -1]`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`



rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(x_), x_Symbol] :> Simp[1/b
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a
, b, c}, x] && IGtQ[n, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +
c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& NeQ[m, -1]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d +
e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2
)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b,
c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3
, 0] && NeQ[m, -1]`

```
rule 6347 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_
.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1
))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp
[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(
f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^
(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && ILtQ[m, -1]
```

### 3.203.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1520 vs.  $2(310) = 620$ .

Time = 1.21 (sec) , antiderivative size = 1521, normalized size of antiderivative = 4.64

method	result	size
default	Expression too large to display	1521
parts	Expression too large to display	1521

```
input int((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)
```

output `2/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*c^3+a^2*(-1/3/d/x^3*(-c^2*d*x^2+d)^(1/2)-2/3*c^2/d/x*(-c^2*d*x^2+d)^(1/2))-1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^3+2/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x^5*c^8+1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x^3*c^6-2/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x*c^4-1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)/x*c^2+1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)/x^3*arccosh(c*x)^2-4/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*arccosh(c*x)^2*c^3-4/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x^3*arccosh(c*x)*(c*x-1)*(c*x+1)*c^6+2*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x^2*arccosh(c*x)^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^5-2/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x*arccosh(c*x)*(c*x-1)*(c*x+1)*c^4-1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)/x^2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c+4/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*x^2-1)*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*c^3+4/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x^5*arccosh(c*x)*c^8-2*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x^3*arccosh(c*x)^2*c^6-2/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d/(3*c^4*x^4-2*c^2*x^2-1)*x^3*arccosh(c*x)*c^6+1/3*b^...`

### 3.203.5 Fricas [F]

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + dx^4}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^2*d*x^6 - d*x^4), x)`

**3.203.6 Sympy [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^4 \sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate((a+b*acosh(c*x))**2/x**4/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x**4*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.203.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + dx^4}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `1/3*(4*c^2*sqrt(-d)*log(x)/d - sqrt(-d)/(d*x^2))*a*b*c - 2/3*a*b*(2*sqrt(-c^2*d*x^2 + d)*c^2/(d*x) + sqrt(-c^2*d*x^2 + d)/(d*x^3))*arccosh(c*x) - 1/3*a^2*(2*sqrt(-c^2*d*x^2 + d)*c^2/(d*x) + sqrt(-c^2*d*x^2 + d)/(d*x^3)) + b^2*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/(sqrt(-c^2*d*x^2 + d)*x^4), x)`

**3.203.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{-c^2 dx^2 + dx^4}} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/(sqrt(-c^2*d*x^2 + d)*x^4), x)`

**3.203.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^4 \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^4*(d - c^2*d*x^2)^(1/2)),x)`output `int((a + b*acosh(c*x))^2/(x^4*(d - c^2*d*x^2)^(1/2)), x)`

$$3.204 \quad \int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$$

3.204.1 Optimal result	1773
3.204.2 Mathematica [A] (warning: unable to verify)	1774
3.204.3 Rubi [C] (verified)	1775
3.204.4 Maple [A] (verified)	1784
3.204.5 Fracas [F]	1785
3.204.6 Sympy [F]	1786
3.204.7 Maxima [F]	1786
3.204.8 Giac [F(-2)]	1786
3.204.9 Mupad [F(-1)]	1787

### 3.204.1 Optimal result

Integrand size = 29, antiderivative size = 556

$$\begin{aligned} \int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx &= \frac{16abx\sqrt{-1+cx}\sqrt{1+cx}}{3c^5d\sqrt{d-c^2dx^2}} + \frac{94b^2(1-cx)(1+cx)}{27c^6d\sqrt{d-c^2dx^2}} \\ &+ \frac{2b^2x^2(1-cx)(1+cx)}{27c^4d\sqrt{d-c^2dx^2}} + \frac{16b^2x\sqrt{-1+cx}\sqrt{1+cx}\operatorname{arccosh}(cx)}{3c^5d\sqrt{d-c^2dx^2}} \\ &- \frac{2bx\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{c^5d\sqrt{d-c^2dx^2}} \\ &+ \frac{2bx^3\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{9c^3d\sqrt{d-c^2dx^2}} + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\ &+ \frac{8\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^6d^2} + \frac{4x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^4d^2} \\ &+ \frac{4b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^6d\sqrt{d-c^2dx^2}} \\ &+ \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})}{c^6d\sqrt{d-c^2dx^2}} \\ &- \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{c^6d\sqrt{d-c^2dx^2}} \end{aligned}$$

---


$$3.204. \quad \int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$$

output  $94/27*b^2*(-c*x+1)*(c*x+1)/c^6/d/(-c^2*d*x^2+d)^{(1/2)}+2/27*b^2*x^2*(-c*x+1)*(c*x+1)/c^4/d/(-c^2*d*x^2+d)^{(1/2)}+x^4*(a+b*\operatorname{arccosh}(c*x))^2/c^2/d/(-c^2*d*x^2+d)^{(1/2)}+16/3*a*b*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^5/d/(-c^2*d*x^2+d)^{(1/2)}+16/3*b^2*x*\operatorname{arccosh}(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^5/d/(-c^2*d*x^2+d)^{(1/2)}-2*b*x*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^5/d/(-c^2*d*x^2+d)^{(1/2)}+2/9*b*x^3*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/d/(-c^2*d*x^2+d)^{(1/2)}+4*b*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^6/d/(-c^2*d*x^2+d)^{(1/2)}+2*b^2*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^6/d/(-c^2*d*x^2+d)^{(1/2)}-2*b^2*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^6/d/(-c^2*d*x^2+d)^{(1/2)}+8/3*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/c^6/d^2+4/3*x^2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/c^4/d^2$

### 3.204.2 Mathematica [A] (warning: unable to verify)

Time = 3.00 (sec) , antiderivative size = 392, normalized size of antiderivative = 0.71

$$\int \frac{x^5(a + b\operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \frac{-36a^2(-8 + 4c^2x^2 + c^4x^4) + 3ab(135\operatorname{arccosh}(cx) - 60\operatorname{arccosh}(cx)\cosh(2\operatorname{arccosh}(cx)))}{(d - c^2dx^2)^{3/2}}$$

input `Integrate[(x^5*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

output  $(-36*a^2*(-8 + 4*c^2*x^2 + c^4*x^4) + 3*a*b*(135*\operatorname{ArcCosh}[c*x] - 60*\operatorname{ArcCosh}[c*x]*\operatorname{Cosh}[2*\operatorname{ArcCosh}[c*x]] - 3*\operatorname{ArcCosh}[c*x]*\operatorname{Cosh}[4*\operatorname{ArcCosh}[c*x]] + 72*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{Log}[\operatorname{Cosh}[\operatorname{ArcCosh}[c*x]/2]] - 72*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{Log}[\operatorname{Sinh}[\operatorname{ArcCosh}[c*x]/2]] + 62*\operatorname{Sinh}[2*\operatorname{ArcCosh}[c*x]] + \operatorname{Sinh}[4*\operatorname{ArcCosh}[c*x]]) - b^2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(378*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x) - 378*c*x*\operatorname{ArcCosh}[c*x] + 189*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x]^2 - 6*\operatorname{ArcCosh}[c*x]*\operatorname{Cosh}[3*\operatorname{ArcCosh}[c*x]] - 54*\operatorname{ArcCosh}[c*x]^2*\operatorname{Coth}[\operatorname{ArcCosh}[c*x]/2] + 216*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - E^{(-\operatorname{ArcCosh}[c*x])}] - 216*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + E^{(-\operatorname{ArcCosh}[c*x])}] + 216*\operatorname{PolyLog}[2, -E^{(-\operatorname{ArcCosh}[c*x])}] - 216*\operatorname{PolyLog}[2, E^{(-\operatorname{ArcCosh}[c*x])}] + 2*\operatorname{Sinh}[3*\operatorname{ArcCosh}[c*x]] + 9*\operatorname{ArcCosh}[c*x]^2*\operatorname{Sinh}[3*\operatorname{ArcCosh}[c*x]] + 54*\operatorname{ArcCosh}[c*x]^2*\operatorname{Tanh}[\operatorname{ArcCosh}[c*x]/2]))/(108*c^6*d*\operatorname{Sqrt}[d - c^2*d*x^2])$

### 3.204.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 3.09 (sec) , antiderivative size = 535, normalized size of antiderivative = 0.96, number of steps used = 22, number of rules used = 21,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.724$ , Rules used = {6349, 25, 6327, 6353, 111, 27, 83, 6298, 111, 27, 83, 6329, 2009, 6353, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6349} \\
 & -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -\frac{x^4(a+\operatorname{barccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2 dx^2}} - \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2 dx^2}} dx}{c^2 d} + \frac{x^4(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^4(a+\operatorname{barccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2 dx^2}} - \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2 dx^2}} dx}{c^2 d} + \frac{x^4(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^4(a+\operatorname{barccosh}(cx))}{1-c^2 x^2} dx}{cd\sqrt{d-c^2 dx^2}} - \frac{4 \int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2 dx^2}} dx}{c^2 d} + \frac{x^4(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{6353} \\
 & -\frac{4 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int x^2(a+\operatorname{barccosh}(cx)) dx}{3c\sqrt{d-c^2 dx^2}} + \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2 dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2 dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2 d} \right)}{c^2 d} + \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2 x^2} dx}{c^2} + \frac{b \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2} \right)}{cd\sqrt{d-c^2 dx^2}} + \\
 & \frac{x^4(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{111}
 \end{aligned}$$

---

3.204.  $\int \frac{x^5(a+\operatorname{barccosh}(cx))^2}{(d-c^2 dx^2)^{3/2}} dx$



$$\begin{aligned}
& 4 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int x^2(a+\operatorname{barccosh}(cx))dx}{3c\sqrt{d-c^2dx^2}} + \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \right) \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} \right)}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 27
\end{aligned}$$

$$\begin{aligned}
& 4 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int x^2(a+\operatorname{barccosh}(cx))dx}{3c\sqrt{d-c^2dx^2}} + \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \right) \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} \right)}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 83
\end{aligned}$$

$$\begin{aligned}
& 4 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int x^2(a+\operatorname{barccosh}(cx))dx}{3c\sqrt{d-c^2dx^2}} + \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{3c^2d} \right) \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{x^3(a+\operatorname{barccosh}(cx))}{3c^2d} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} \right)}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow 6298
\end{aligned}$$

---

3.204.  $\int \frac{x^5(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{3c\sqrt{d-c^2dx^2}} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} \right)$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} \right)$$


---


$$\frac{cd\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} + \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}$$

↓ 111

$$4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} - \frac{x^2\sqrt{d-c^2dx^2}}{3c^2d} \right)$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} \right)$$


---


$$\frac{cd\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} + \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}$$

↓ 27

$$4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} - \frac{x^2\sqrt{d-c^2dx^2}}{3c^2d} \right)$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} \right)$$


---


$$\frac{cd\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} + \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}$$

↓ 83

---

3.204.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} \right)}{cd\sqrt{d-c^2dx^2}} \\
 & \frac{4 \left( \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + x^2\sqrt{d-c^2dx^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} \right)}{c^2d} \\
 & \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{6329} \\
 & \frac{4 \left( \frac{2 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx)) dx}{c\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} \right)}{3c^2} - \frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{3}x^3(a+b\operatorname{arccosh}(cx)) - \frac{1}{3}bc \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + x^2\sqrt{d-c^2dx^2} \right) \right)}{3c\sqrt{d-c^2dx^2}} \right)}{c^2d} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} \right)}{cd\sqrt{d-c^2dx^2}} + \\
 & \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x^2(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{3c} \right)}{cd\sqrt{d-c^2dx^2}} + \\
 & \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
 & \frac{4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2dx^2}} \right)}{3c^2} \right)}{c^2d} \\
 & \quad \downarrow \text{6353}
 \end{aligned}$$

3.204.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} + \frac{b \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c^2} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2}{3c} \right)}{3c} \right)$$

---


$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{\frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}}{3c^2} - 2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx)}{c\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{3c^2} \right) \right) - 2b\sqrt{cx-1}\sqrt{cx+1}$$


---

$c^2d$

↓ 83

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c} \right)}{3c} \right)$$

---


$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{\frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}}{3c^2} - 2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx)}{c\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{3c^2} \right) \right) - 2b\sqrt{cx-1}\sqrt{cx+1}$$


---

$c^2d$

↓ 6318

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}(cx+1)}} d\operatorname{arccosh}(cx)}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c} \right)}{3c} \right)$$

---


$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{\frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}}{3c^2} - 2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx)}{c\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{3c^2} \right) \right) - 2b\sqrt{cx-1}\sqrt{cx+1}$$


---

$c^2d$

↓ 3042

---

3.204.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{-\int i(a+b\operatorname{arccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} \right)$$


---


$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d} - 2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2dx^2}} \right) \right) - 2b\sqrt{cx-1}\sqrt{cx+1}$$


---

↓ 26

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{-\int i(a+b\operatorname{arccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} \right)$$


---


$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d} - 2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2dx^2}} \right) \right) - 2b\sqrt{cx-1}\sqrt{cx+1}$$


---

↓ 4670

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{-\left( ib \int \log(1-e\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - ib \int \log(1+e\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e\operatorname{arccosh}(cx)) \right) (a+b\operatorname{arccosh}(cx))}{c^3} + \frac{x(a+b\operatorname{arccosh}(cx))}{c^2}}{c^2} - \frac{x^3(a+b\operatorname{arccosh}(cx))}{3c^2} \right)$$


---


$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d} - 2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2dx^2}} \right) \right) - 2b\sqrt{cx-1}\sqrt{cx+1}$$


---

↓ 2715

---

3.204.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{-i \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) dx e^{\operatorname{arccosh}(cx)} - i b \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) dx e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^3} \right)$$

$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - 2 \left( \frac{-\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx)}{c\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{3c^2} \right) \right)$$

↓ 2838

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{-i \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right) (a+b\operatorname{arccosh}(cx)) + i b \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - i b \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} \right)$$

$$4 \left( -\frac{x^2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^2d} + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - 2 \left( \frac{-\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx)}{c\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{3c^2} \right) \right)$$

input `Int[(x^5*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

output `(x^4*(a + b*ArcCosh[c*x])^2)/(c^2*d*Sqrt[d - c^2*d*x^2]) - (4*(-1/3*(x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/3*(b*c*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2))) + (x^3*(a + b*ArcCosh[c*x]))/3))/3*c*Sqrt[d - c^2*d*x^2]) + (2*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d)) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(c*Sqrt[d - c^2*d*x^2])))/(3*c^2))/(c^2*d) + (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((b*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2)))/(3*c) - (x^3*(a + b*ArcCosh[c*x]))/(3*c^2) + ((b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^3 - (x*(a + b*ArcCosh[c*x]))/c^2 - (I*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c^3)/c^2))/(c*d*Sqrt[d - c^2*d*x^2])`

3.204.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

## 3.204.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 111 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/(f*fz*I)], x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 6349 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*(m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]`



```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.204.4 Maple [A] (verified)

Time = 1.46 (sec) , antiderivative size = 781, normalized size of antiderivative = 1.40

method	result
default	$a^2 \left( -\frac{x^4}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{-\frac{4x^2}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{8}{3d c^4 \sqrt{-c^2 d x^2 + d}}}{c^2} \right) + \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} (9\sqrt{cx + 1} \sqrt{cx - 1} \arccos(\frac{cx + 1}{\sqrt{c^2 x^2 - 1}}) - 9\sqrt{cx - 1} \sqrt{cx + 1} \arccos(\frac{cx - 1}{\sqrt{c^2 x^2 - 1}}))}{c^2 \sqrt{-d(c^2 x^2 - 1)}}$
parts	$a^2 \left( -\frac{x^4}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{-\frac{4x^2}{3c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{8}{3d c^4 \sqrt{-c^2 d x^2 + d}}}{c^2} \right) + \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} (9\sqrt{cx + 1} \sqrt{cx - 1} \arccos(\frac{cx + 1}{\sqrt{c^2 x^2 - 1}}) - 9\sqrt{cx - 1} \sqrt{cx + 1} \arccos(\frac{cx - 1}{\sqrt{c^2 x^2 - 1}}))}{c^2 \sqrt{-d(c^2 x^2 - 1)}}$

```
input int(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

output

```

a^2*(-1/3*x^4/c^2/d/(-c^2*d*x^2+d)^(1/2)+4/3/c^2*(-x^2/c^2/d/(-c^2*d*x^2+d)^(1/2)+2/d/c^4/(-c^2*d*x^2+d)^(1/2)))+1/27*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(9*(c*x+1)^(1/2)*(c*x-1)^(1/2)*arccosh(c*x)^2*x^4*c^4-6*arccosh(c*x)*c^5*x^5+2*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+36*arccosh(c*x)^2*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^2*c^2-84*c^3*x^3*arccosh(c*x)+92*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-54*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+54*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)))*x^2*c^2-54*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+54*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2-72*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2+90*c*x*arccosh(c*x)-94*(c*x-1)^(1/2)*(c*x+1)^(1/2)+54*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-54*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+54*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-54*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/(c^2*x^2-1)^2/d^2/c^6+2/9*a*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4-c^5*x^5+12*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-14*c^3*x^3-9*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+9*ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1))*x^2*c^2-24*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+15*c*x+9*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-9*ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1))/(c^2*x^2-1)^2/d^2/c^6

```

### 3.204.5 Fracas [F]

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^5}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral((b^2*x^5*arccosh(c*x)^2 + 2*a*b*x^5*arccosh(c*x) + a^2*x^5)*sqrt(-c^2*d*x^2 + d)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

## 3.204.6 Sympy [F]

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**5*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(3/2), x)`

output `Integral(x**5*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

## 3.204.7 Maxima [F]

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^5}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2), x, algorithm="maxima")`

output `-1/3*a^2*(x^4/(sqrt(-c^2*d*x^2 + d)*c^2*d) + 4*x^2/(sqrt(-c^2*d*x^2 + d)*c^4*d) - 8/(sqrt(-c^2*d*x^2 + d)*c^6*d) + 1/3*(b^2*c^4*sqrt(d)*x^4 + 4*b^2*c^2*sqrt(d)*x^2 - 8*b^2*sqrt(d))*sqrt(c*x + 1)*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/(c^8*d^2*x^2 - c^6*d^2) + integrate(-2/3*((4*b^2*c^3*x^3 - (3*a*b*c^5 - b^2*c^5)*x^5 - 8*b^2*c*x)*(c*x + 1)*sqrt(c*x - 1) + (3*b^2*c^4*x^4 - (3*a*b*c^6 - b^2*c^6)*x^6 - 12*b^2*c^2*x^2 + 8*b^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(c^10*d^(3/2)*x^5 - 2*c^8*d^(3/2)*x^3 + c^6*d^(3/2)*x + (c^9*d^(3/2)*x^4 - 2*c^7*d^(3/2)*x^2 + c^5*d^(3/2))*sqrt(c*x + 1)*sqrt(c*x - 1)), x)`

## 3.204.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.204.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^5(a + b\operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x^5(a + b\operatorname{acosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx$$

input `int((x^5*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2),x)`

output `int((x^5*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2), x)`

**3.205**  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

3.205.1 Optimal result . . . . .	1788
3.205.2 Mathematica [A] (warning: unable to verify) . . . . .	1789
3.205.3 Rubi [C] (verified) . . . . .	1790
3.205.4 Maple [A] (verified) . . . . .	1798
3.205.5 Fricas [F] . . . . .	1799
3.205.6 Sympy [F] . . . . .	1800
3.205.7 Maxima [F] . . . . .	1800
3.205.8 Giac [F(-2)] . . . . .	1800
3.205.9 Mupad [F(-1)] . . . . .	1801

**3.205.1 Optimal result**

Integrand size = 29, antiderivative size = 440

$$\int \frac{x^4(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \frac{b^2x(1 - cx)(1 + cx)}{4c^4d\sqrt{d - c^2dx^2}} - \frac{b^2\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arccosh}(cx)}{4c^5d\sqrt{d - c^2dx^2}} + \frac{bx^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{2c^3d\sqrt{d - c^2dx^2}} + \frac{x^3(a + \operatorname{arccosh}(cx))^2}{c^2d\sqrt{d - c^2dx^2}} + \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{c^5d\sqrt{d - c^2dx^2}} + \frac{3x\sqrt{d - c^2dx^2}(a + \operatorname{arccosh}(cx))^2}{2c^4d^2} - \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^3}{2bc^5d\sqrt{d - c^2dx^2}} - \frac{2b\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx)) \log(1 - e^{2\operatorname{arccosh}(cx)})}{c^5d\sqrt{d - c^2dx^2}} - \frac{b^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{c^5d\sqrt{d - c^2dx^2}}$$

output  $\frac{1}{4}b^2x(-cx+1)(cx+1)/c^4/d/(-c^2dx^2+d)^{1/2}+x^3(a+b\operatorname{arccosh}(cx))^2/c^2/d/(-c^2dx^2+d)^{1/2}-1/4b^2\operatorname{arccosh}(cx)(cx-1)^{1/2}(cx+1)^{1/2}/c^5/d/(-c^2dx^2+d)^{1/2}+1/2b^2x^2(a+b\operatorname{arccosh}(cx))(cx-1)^{1/2}(cx+1)^{1/2}/c^3/d/(-c^2dx^2+d)^{1/2}+(a+b\operatorname{arccosh}(cx))^2(cx-1)^{1/2}(cx+1)^{1/2}/c^5/d/(-c^2dx^2+d)^{1/2}-1/2(a+b\operatorname{arccosh}(cx))^3(cx-1)^{1/2}(cx+1)^{1/2}/b/c^5/d/(-c^2dx^2+d)^{1/2}-2b(a+b\operatorname{arccosh}(cx))\ln(1-(cx+(cx-1)^{1/2}(cx+1)^{1/2}))^2(cx-1)^{1/2}(cx+1)^{1/2}/c^5/d/(-c^2dx^2+d)^{1/2}-b^2\operatorname{polylog}(2,(cx+(cx-1)^{1/2}(cx+1)^{1/2}))^2(cx-1)^{1/2}(cx+1)^{1/2}/c^5/d/(-c^2dx^2+d)^{1/2}+3/2x(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{1/2}/c^4/d^2$

### 3.205.2 Mathematica [A] (warning: unable to verify)

Time = 1.69 (sec) , antiderivative size = 343, normalized size of antiderivative = 0.78

$$\int \frac{x^4(a + b\operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \frac{-4a^2c\sqrt{d}x(-3 + c^2x^2) + 12a^2\sqrt{d - c^2dx^2} \arctan\left(\frac{cx\sqrt{d - c^2dx^2}}{\sqrt{d}(-1 + c^2x^2)}\right) + 2ab\sqrt{d}\left(8c^2x^2\sqrt{d - c^2dx^2} - 4cx\sqrt{d} + 4d\right)}{(d - c^2dx^2)^{3/2}}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2),x]`

output  $(-4a^2c\sqrt{d}x^2(-3 + c^2x^2) + 12a^2\sqrt{d - c^2dx^2}\operatorname{ArcTan}[(cx\sqrt{d - c^2dx^2})/(\sqrt{d}(-1 + c^2x^2))] + 2ab\sqrt{d}(8c^2x^2\sqrt{d - c^2dx^2} - 4cx\sqrt{d} + 4d) + 8\operatorname{Log}[\sqrt{(-1 + cx)/(1 + cx)}(1 + cx)] + 2\operatorname{ArcCosh}[cx]\operatorname{Sinh}[2\operatorname{ArcCosh}[cx]]) + b^2\sqrt{d}(8cx\operatorname{ArcCosh}[cx]^2 + 8\sqrt{(-1 + cx)/(1 + cx)}(1 + cx)\operatorname{PolyLog}[2, E^{-2\operatorname{ArcCosh}[cx]}] - \sqrt{(-1 + cx)/(1 + cx)}(1 + cx)(4\operatorname{ArcCosh}[cx]^3 - 2\operatorname{ArcCosh}[cx](\operatorname{Cosh}[2\operatorname{ArcCosh}[cx]] - 8\operatorname{Log}[1 - E^{-2\operatorname{ArcCosh}[cx]}]) + \operatorname{Sinh}[2\operatorname{ArcCosh}[cx]] + 2\operatorname{ArcCosh}[cx]^2(4 + \operatorname{Sinh}[2\operatorname{ArcCosh}[cx]])))))/(8c^5d^{3/2}\sqrt{d - c^2dx^2})$

**3.205.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.47 (sec) , antiderivative size = 412, normalized size of antiderivative = 0.94, number of steps used = 19, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.621$ , Rules used = {6349, 25, 6327, 6353, 101, 43, 6298, 101, 43, 6307, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6349} \\
 & -\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{c^2 d} - \frac{2b\sqrt{cx - 1}\sqrt{cx + 1} \int -\frac{x^3(a + \operatorname{barccosh}(cx))}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{25} \\
 & -\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{c^2 d} + \frac{2b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x^3(a + \operatorname{barccosh}(cx))}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{6327} \\
 & -\frac{3 \int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{c^2 d} + \frac{2b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x^3(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{cd\sqrt{d - c^2 dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{6353} \\
 & \frac{2b\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{\int \frac{x(a + \operatorname{barccosh}(cx))}{1 - c^2 x^2} dx}{c^2} + \frac{b \int \frac{x^2}{\sqrt{cx - 1}\sqrt{cx + 1}} dx}{2c} - \frac{x^2(a + \operatorname{barccosh}(cx))}{2c^2} \right)}{cd\sqrt{d - c^2 dx^2}} \\
 & \frac{3 \left( -\frac{b\sqrt{cx - 1}\sqrt{cx + 1} \int x(a + \operatorname{barccosh}(cx)) dx}{c\sqrt{d - c^2 dx^2}} + \frac{\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx}{2c^2} - \frac{x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^2}{2c^2 d} \right)}{c^2 d} + \\
 & \frac{x^3(a + \operatorname{barccosh}(cx))^2}{c^2 d\sqrt{d - c^2 dx^2}} \\
 & \quad \downarrow \text{101}
 \end{aligned}$$

$$\begin{aligned}
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x(a+\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{c^2} + \frac{b \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2c} - \frac{x^2(a+\operatorname{arccosh}(cx))}{2c^2} \right) \\
& \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& 3 \left( -\frac{b\sqrt{cx-1}\sqrt{cx+1} \int x(a+\operatorname{arccosh}(cx)) dx}{c\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{arccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} \right) + \\
& \frac{c^2d}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& \downarrow 43 \\
& 3 \left( -\frac{b\sqrt{cx-1}\sqrt{cx+1} \int x(a+\operatorname{arccosh}(cx)) dx}{c\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{arccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} \right) + \\
& \frac{c^2d}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x(a+\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^2(a+\operatorname{arccosh}(cx))}{2c^2} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2c} \right) + \\
& \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& \downarrow 6298 \\
& 3 \left( -\frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+\operatorname{arccosh}(cx)) - \frac{1}{2}bc \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx \right)}{c\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{arccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} \right) + \\
& \frac{c^2d}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x(a+\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^2(a+\operatorname{arccosh}(cx))}{2c^2} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2c} \right) + \\
& \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{arccosh}(cx))^2} \\
& \downarrow 101
\end{aligned}$$

---

3.205.  $\int \frac{x^4(a+\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$



$$\begin{aligned}
 & 3 \left( \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))}{2c^2} \right) \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x(a+\operatorname{barccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2c} \right)}{c^2d} \\
 & \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{barccosh}(cx))^2} + \\
 & \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}} \\
 & \downarrow 43 \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x(a+\operatorname{barccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2c} \right) \\
 & \frac{cd\sqrt{d-c^2dx^2}}{c^2d} \\
 & 3 \left( \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{2c^2} - \frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2c^2d} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c\sqrt{d-c^2dx^2}} \right) \\
 & \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{barccosh}(cx))^2} + \\
 & \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}} \\
 & \downarrow 6307 \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{x(a+\operatorname{barccosh}(cx)) dx}{1-c^2x^2}}{c^2} - \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{2c} \right) \\
 & \frac{cd\sqrt{d-c^2dx^2}}{x^3(a+\operatorname{barccosh}(cx))^2} + \\
 & \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}} \\
 & 3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+\operatorname{barccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \right)}{c\sqrt{d-c^2dx^2}} \right) \\
 & \frac{cd\sqrt{d-c^2dx^2}}{c^2d} \\
 & \downarrow 6328
 \end{aligned}$$

3.205.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{cx(a+b\operatorname{arccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{c^4} - \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}{2c} \right)$$


---


$$\frac{cd\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}$$


---


$$3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}\right)\right)}{c\sqrt{d-c^2dx^2}} \right)$$


---

$c^2d$

↓ 3042

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int -i(a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{c^4} - \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}{2c} \right)$$


---


$$\frac{cd\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}$$


---


$$3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}\right)\right)}{c\sqrt{d-c^2dx^2}} \right)$$


---

$c^2d$

↓ 26

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{i \int (a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{c^4} - \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2}\right)}{2c} \right)$$


---


$$\frac{cd\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}$$


---


$$3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}\right)\right)}{c\sqrt{d-c^2dx^2}} \right)$$


---

$c^2d$

↓ 4199

---

3.205.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{i \left( 2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a+b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} - \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2} + \frac{b \left( \operatorname{arccosh}(cx) \right)}{2c^3} \right)$$


---


$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d}$$


---


$$3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} \right) \right)}{c\sqrt{d-c^2dx^2}} \right)$$


---

$c^2d$

↓ 25

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{i \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a+b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} - \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2} + \frac{b \left( \operatorname{arccosh}(cx) \right)}{2c^3} \right)$$


---


$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d}$$


---


$$3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} \right) \right)}{c\sqrt{d-c^2dx^2}} \right)$$


---

$c^2d$

↓ 2620

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{i \left( -2i \left( \frac{1}{2}b \int \log(1-e^{2\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - \frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx)) \right) - \frac{i(a+b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} - \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2} + \frac{b \left( \operatorname{arccosh}(cx) \right)}{2c^3} \right)$$


---


$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d}$$


---


$$3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}x^2(a+b\operatorname{arccosh}(cx)) - \frac{1}{2}bc \left( \frac{\operatorname{arccosh}(cx)}{2c^3} \right) \right)}{c\sqrt{d-c^2dx^2}} \right)$$


---

$c^2d$

↓ 2715

---

3.205.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{i\left(-2i\left(\frac{1}{4}b \int e^{-2\operatorname{arccosh}(cx)} \log\left(1-e^{2\operatorname{arccosh}(cx)}\right) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log\left(1-e^{2\operatorname{arccosh}(cx)}\right)\right)(a+\operatorname{arccosh}(cx))\right) - i(a+\operatorname{arccosh}(cx))}{c^4} \right) \\
 & \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}} - \frac{x^3(a+\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \\
 & 3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+\operatorname{arccosh}(cx)) - \frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}\right)\right)}{c\sqrt{d-c^2dx^2}} \right) \\
 & \frac{c^2d}{c^2d} \\
 & \downarrow 2838 \\
 & \frac{x^3(a+\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \\
 & 3 \left( -\frac{x\sqrt{d-c^2dx^2}(a+\operatorname{arccosh}(cx))^2}{2c^2d} + \frac{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{arccosh}(cx))^3}{6bc^3\sqrt{d-c^2dx^2}} - \frac{b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}x^2(a+\operatorname{arccosh}(cx)) - \frac{1}{2}bc\left(\frac{\operatorname{arccosh}(cx)}{2c^3}\right)\right)}{c\sqrt{d-c^2dx^2}} \right) \\
 & \frac{c^2d}{c^2d} \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{i\left(-2i\left(-\frac{1}{2} \log\left(1-e^{2\operatorname{arccosh}(cx)}\right)\right)(a+\operatorname{arccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog}\left(2, e^{2\operatorname{arccosh}(cx)}\right)\right) - \frac{i(a+\operatorname{arccosh}(cx))^2}{2b}}{c^4} \right) - \frac{x^2}{c^4} \\
 & \frac{cd\sqrt{d-c^2dx^2}}{cd\sqrt{d-c^2dx^2}}
 \end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

output `(x^3*(a + b*ArcCosh[c*x])^2)/(c^2*d*Sqrt[d - c^2*d*x^2]) - (3*(-1/2*(x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^3)/(6*b*c^3*Sqrt[d - c^2*d*x^2]) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((x^2*(a + b*ArcCosh[c*x]))/2 - (b*c*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/2))/(c*Sqrt[d - c^2*d*x^2]))/(c^2*d) + (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/2*(x^2*(a + b*ArcCosh[c*x]))/c^2 + (b*((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/(2*c) + (I*((( -1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/c^4))/(c*d*Sqrt[d - c^2*d*x^2])`

3.205.  $\int \frac{x^4(a+\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

## 3.205.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_) + (b_)*(x_))^(n_)*((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_))^(n_)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6349 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]`

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.205.4 Maple [A] (verified)

Time = 1.46 (sec) , antiderivative size = 756, normalized size of antiderivative = 1.72

method	result
default	$-\frac{a^2 x^3}{2c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{3a^2 x}{2c^4 d \sqrt{-c^2 d x^2 + d}} - \frac{3a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{2c^4 d \sqrt{c^2 d}} + \frac{b^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{-d(c^2 x^2 - 1)} (2 \operatorname{arccosh}(cx))^2 \sqrt{cx}}{2c^4 d \sqrt{c^2 d}}$
parts	$-\frac{a^2 x^3}{2c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{3a^2 x}{2c^4 d \sqrt{-c^2 d x^2 + d}} - \frac{3a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{2c^4 d \sqrt{c^2 d}} + \frac{b^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{-d(c^2 x^2 - 1)} (2 \operatorname{arccosh}(cx))^2 \sqrt{cx}}{2c^4 d \sqrt{c^2 d}}$

```
input int(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

output

$$\begin{aligned}
& -1/2*a^2*x^3/c^2/d/(-c^2*d*x^2+d)^{(1/2)}+3/2*a^2/c^4*x/d/(-c^2*d*x^2+d)^{(1/2)} \\
& -3/2*a^2/c^4/d/(c^2*d)^{(1/2)}*\arctan((c^2*d)^{(1/2)}*x/(-c^2*d*x^2+d)^{(1/2)}) \\
& +1/4*b^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(-d*(c^2*x^2-1))^{(1/2)}*(2*\operatorname{arccosh}(c*x) \\
& ^2*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*x^3*c^3-2*c^4*x^4*\operatorname{arccosh}(c*x)+(c*x-1)^{(1/2)} \\
& *(c*x+1)^{(1/2)}*c^3*x^3+2*\operatorname{arccosh}(c*x)^3*x^2*c^2-6*\operatorname{arccosh}(c*x)^2*(c*x+1) \\
& )^{(1/2)}*(c*x-1)^{(1/2)}*x*c-4*\operatorname{arccosh}(c*x)^2*x^2*c^2+8*\operatorname{arccosh}(c*x)*\ln(1+c*x \\
& +(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*x^2*c^2+8*\operatorname{arccosh}(c*x)*\ln(1-c*x-(c*x-1)^{(1/2)} \\
& )*(c*x+1)^{(1/2)})*x^2*c^2+3*c^2*x^2*\operatorname{arccosh}(c*x)+8*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)} \\
& )*(c*x+1)^{(1/2)})*x^2*c^2+8*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*x \\
& ^2*c^2-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x-2*\operatorname{arccosh}(c*x)^3+4*\operatorname{arccosh}(c*x)^2-8 \\
& *\operatorname{arccosh}(c*x)*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})-8*\operatorname{arccosh}(c*x)*\ln(1-c* \\
& x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})-\operatorname{arccosh}(c*x)-8*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)} \\
& )*(c*x+1)^{(1/2)}-8*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})))/(c^2*x^2-1)^ \\
& 2/d^2/c^5+1/4*a*b*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(-d*(c^2*x^2-1))^{(1/2)}*(4*(c \\
& *x+1)^{(1/2)}*\operatorname{arccosh}(c*x)*(c*x-1)^{(1/2)}*c^3*x^3-2*c^4*x^4+6*\operatorname{arccosh}(c*x)^2* \\
& x^2*c^2-12*(c*x+1)^{(1/2)}*\operatorname{arccosh}(c*x)*(c*x-1)^{(1/2)}*c*x-8*c^2*x^2*\operatorname{arccosh}( \\
& c*x)+8*\ln((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2-1)*x^2*c^2+3*c^2*x^2-6*\operatorname{arcco} \\
& sh(c*x)^2+8*\operatorname{arccosh}(c*x)-8*\ln((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2-1)-1)/(c \\
& ^2*x^2-1)^2/d^2/c^5
\end{aligned}$$

### 3.205.5 Fracas [F]

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^4}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral((b^2*x^4*arccosh(c*x)^2 + 2*a*b*x^4*arccosh(c*x) + a^2*x^4)*sqrt(-c^2*d*x^2 + d)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`



## 3.205.6 Sympy [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**4*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x**4*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

## 3.205.7 Maxima [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^4}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-1/2*a^2*(x^3/(sqrt(-c^2*d*x^2 + d)*c^2*d) - 3*x/(sqrt(-c^2*d*x^2 + d)*c^4*d) + 3*arcsin(c*x)/(c^5*d^(3/2))) + integrate(b^2*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/(-c^2*d*x^2 + d)^(3/2) + 2*a*b*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(-c^2*d*x^2 + d)^(3/2), x)`

## 3.205.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.205.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx$

**3.205.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4 (a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^4 (a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x^4*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2),x)`output `int((x^4*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2), x)`

**3.206**  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

3.206.1 Optimal result . . . . . 1802  
 3.206.2 Mathematica [A] (warning: unable to verify) . . . . . 1803  
 3.206.3 Rubi [C] (verified) . . . . . 1804  
 3.206.4 Maple [A] (verified) . . . . . 1809  
 3.206.5 Fracas [F] . . . . . 1810  
 3.206.6 Sympy [F] . . . . . 1810  
 3.206.7 Maxima [F] . . . . . 1810  
 3.206.8 Giac [F(-2)] . . . . . 1811  
 3.206.9 Mupad [F(-1)] . . . . . 1811

**3.206.1 Optimal result**

Integrand size = 29, antiderivative size = 413

$$\begin{aligned} \int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx &= \frac{4abx\sqrt{-1+cx}\sqrt{1+cx}}{c^3d\sqrt{d-c^2dx^2}} + \frac{2b^2(1-cx)(1+cx)}{c^4d\sqrt{d-c^2dx^2}} \\ &+ \frac{4b^2x\sqrt{-1+cx}\sqrt{1+cx}\operatorname{arccosh}(cx)}{c^3d\sqrt{d-c^2dx^2}} - \frac{2bx\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{c^3d\sqrt{d-c^2dx^2}} \\ &+ \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \frac{2\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^4d^2} \\ &+ \frac{4b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^4d\sqrt{d-c^2dx^2}} \\ &+ \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})}{c^4d\sqrt{d-c^2dx^2}} \\ &- \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{c^4d\sqrt{d-c^2dx^2}} \end{aligned}$$

output  $2*b^2*(-c*x+1)*(c*x+1)/c^4/d/(-c^2*d*x^2+d)^{(1/2)}+x^2*(a+b*\operatorname{arccosh}(c*x))^2/c^2/d/(-c^2*d*x^2+d)^{(1/2)}+4*a*b*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/d/(-c^2*d*x^2+d)^{(1/2)}+4*b^2*x*\operatorname{arccosh}(c*x)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/d/(-c^2*d*x^2+d)^{(1/2)}-2*b*x*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/d/(-c^2*d*x^2+d)^{(1/2)}+4*b*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^4/d/(-c^2*d*x^2+d)^{(1/2)}+2*b^2*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^4/d/(-c^2*d*x^2+d)^{(1/2)}-2*b^2*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^4/d/(-c^2*d*x^2+d)^{(1/2)}+2*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/c^4/d^2$

### 3.206.2 Mathematica [A] (warning: unable to verify)

Time = 1.42 (sec) , antiderivative size = 319, normalized size of antiderivative = 0.77

$$\int \frac{x^3(a + b\operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \frac{-2a^2(-2 + c^2x^2) + 2ab(3\operatorname{arccosh}(cx) - \operatorname{arccosh}(cx)\cosh(2\operatorname{arccosh}(cx)) + 2\cosh(2\operatorname{arccosh}(cx)))}{(d - c^2dx^2)^{3/2}}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2),x]`

output  $(-2*a^2*(-2 + c^2*x^2) + 2*a*b*(3*\operatorname{ArcCosh}[c*x] - \operatorname{ArcCosh}[c*x]*\operatorname{Cosh}[2*\operatorname{ArcCosh}[c*x]] + 2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(\operatorname{Log}[\operatorname{Cosh}[\operatorname{ArcCosh}[c*x]/2]] - \operatorname{Log}[\operatorname{Sinh}[\operatorname{ArcCosh}[c*x]/2]]) + \operatorname{Sinh}[2*\operatorname{ArcCosh}[c*x]]) + b^2*(2 + 3*\operatorname{ArcCosh}[c*x]^2 - 2*\operatorname{Cosh}[2*\operatorname{ArcCosh}[c*x]] - \operatorname{ArcCosh}[c*x]^2*\operatorname{Cosh}[2*\operatorname{ArcCosh}[c*x]] - 4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - E^{(-\operatorname{ArcCosh}[c*x])}] + 4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + E^{(-\operatorname{ArcCosh}[c*x])}] - 4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{PolyLog}[2, -E^{(-\operatorname{ArcCosh}[c*x])}] + 4*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{PolyLog}[2, E^{(-\operatorname{ArcCosh}[c*x])}] + 2*\operatorname{ArcCosh}[c*x]*\operatorname{Sinh}[2*\operatorname{ArcCosh}[c*x]]))/(2*c^4*d*\operatorname{Sqrt}[d - c^2*d*x^2])$

### 3.206.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.74 (sec) , antiderivative size = 287, normalized size of antiderivative = 0.69, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.448$ , Rules used = {6349, 25, 6327, 6329, 2009, 6353, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6349} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a + \operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a + \operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a + \operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{6329} \\
 & \frac{2 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) dx}{c\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{c^2d} \right)}{c^2d} + \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{cd\sqrt{d-c^2dx^2}} + \frac{x^2(a + \operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{cd\sqrt{d-c^2dx^2}} + \frac{x^2(a + \operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \\
 & 2 \left( -\frac{\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} (ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c})}{c\sqrt{d-c^2dx^2}} \right) \\
 & \quad \downarrow \text{6353}
 \end{aligned}$$

---

3.206.  $\int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} + \frac{b \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} \right)}{+} \\
& \frac{\frac{cd\sqrt{d-c^2dx^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c}}{c\sqrt{d-c^2dx^2}} \right)}{c^2d} \\
& \quad \downarrow 83 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{+} \\
& \frac{\frac{cd\sqrt{d-c^2dx^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c}}{c\sqrt{d-c^2dx^2}} \right)}{c^2d} \\
& \quad \downarrow 6318 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{+} \\
& \frac{\frac{cd\sqrt{d-c^2dx^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c}}{c\sqrt{d-c^2dx^2}} \right)}{c^2d} \\
& \quad \downarrow 3042 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int i(a+b\operatorname{arccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{+} \\
& \frac{\frac{cd\sqrt{d-c^2dx^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} - \frac{cd\sqrt{d-c^2dx^2}}{c^2d\sqrt{d-c^2dx^2}}}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c}}{c\sqrt{d-c^2dx^2}} \right)}{c^2d} \\
& \quad \downarrow 26
\end{aligned}$$

---

3.206.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \int (a+b\operatorname{arccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right) + \\
& \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} - \\
& 2 \left( -\frac{\sqrt{d-c^2 dx^2} (a+b\operatorname{arccosh}(cx))^2}{c^2 d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2 dx^2}} \right) \\
& \frac{cd\sqrt{d-c^2 dx^2}}{c^2 d} \downarrow \mathbf{4670}
\end{aligned}$$

$$\begin{aligned}
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^3} \right) + \\
& \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} - \\
& 2 \left( -\frac{\sqrt{d-c^2 dx^2} (a+b\operatorname{arccosh}(cx))^2}{c^2 d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2 dx^2}} \right) \\
& \frac{cd\sqrt{d-c^2 dx^2}}{c^2 d} \downarrow \mathbf{2715}
\end{aligned}$$

$$\begin{aligned}
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^3} \right) + \\
& \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} - \\
& 2 \left( -\frac{\sqrt{d-c^2 dx^2} (a+b\operatorname{arccosh}(cx))^2}{c^2 d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2 dx^2}} \right) \\
& \frac{cd\sqrt{d-c^2 dx^2}}{c^2 d} \downarrow \mathbf{2838}
\end{aligned}$$

$$\begin{aligned}
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{c^3} \right) + \\
& \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} - \\
& 2 \left( -\frac{\sqrt{d-c^2 dx^2} (a+b\operatorname{arccosh}(cx))^2}{c^2 d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( ax+b\operatorname{arccosh}(cx) - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c} \right)}{c\sqrt{d-c^2 dx^2}} \right) \\
& \frac{cd\sqrt{d-c^2 dx^2}}{c^2 d}
\end{aligned}$$

---

3.206.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2 dx^2)^{3/2}} dx$

input `Int[(x^3*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

output `(x^2*(a + b*ArcCosh[c*x])^2)/(c^2*d*Sqrt[d - c^2*d*x^2]) - (2*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2)/(c^2*d)) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a*x - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x])))/(c*Sqrt[d - c^2*d*x^2])))/(c^2*d) + (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^3 - (x*(a + b*ArcCosh[c*x]))/c^2 - (I*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c^3))/(c*d*Sqrt[d - c^2*d*x^2])`

### 3.206.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 83 `Int[((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

---

3.206.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$



rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 6349 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*(m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]`

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.206.4 Maple [A] (verified)

Time = 1.20 (sec) , antiderivative size = 648, normalized size of antiderivative = 1.57

method	result
default	$a^2 \left( -\frac{x^2}{c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{2}{d c^4 \sqrt{-c^2 d x^2 + d}} \right) + \frac{b^2 \sqrt{c x - 1} \sqrt{c x + 1} \sqrt{-d(c^2 x^2 - 1)} \left( \operatorname{arccosh}(c x)^2 \sqrt{c x + 1} \sqrt{c x - 1} x^2 c^2 - 2 c^3 x^3 \operatorname{arccosh}(c x) \right)}{d^2 \sqrt{-c^2 d x^2 + d}}$
parts	$a^2 \left( -\frac{x^2}{c^2 d \sqrt{-c^2 d x^2 + d}} + \frac{2}{d c^4 \sqrt{-c^2 d x^2 + d}} \right) + \frac{b^2 \sqrt{c x - 1} \sqrt{c x + 1} \sqrt{-d(c^2 x^2 - 1)} \left( \operatorname{arccosh}(c x)^2 \sqrt{c x + 1} \sqrt{c x - 1} x^2 c^2 - 2 c^3 x^3 \operatorname{arccosh}(c x) \right)}{d^2 \sqrt{-c^2 d x^2 + d}}$

```
input int(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output a^2*(-x^2/c^2/d/(-c^2*d*x^2+d)^(1/2)+2/d/c^4/(-c^2*d*x^2+d)^(1/2))+b^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-d*(c^2*x^2-1))^(1/2)*(arccosh(c*x)^2*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^2*c^2-2*c^3*x^3*arccosh(c*x)-2*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))*x^2*c^2+2*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-2*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+2*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2+2*c*x*arccosh(c*x)+2*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-2*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)+2*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-2*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/(c^2*x^2-1)^2/d^2/c^4+2*a*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-d*(c^2*x^2-1))^(1/2)*((c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-c^3*x^3+ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1)*x^2*c^2-ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2-2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1)+ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/(c^2*x^2-1)^2/d^2/c^4
```

**3.206.5 Fricas [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^3}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral((b^2*x^3*arccosh(c*x)^2 + 2*a*b*x^3*arccosh(c*x) + a^2*x^3)*sqrt(-c^2*d*x^2 + d)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.206.6 Sympy [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**3*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x**3*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.206.7 Maxima [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^3}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

```
output -a*b*c*(2*sqrt(-d)*x/(c^4*d^2) + sqrt(-d)*log(c*x + 1)/(c^5*d^2) - sqrt(-d)
)*log(c*x - 1)/(c^5*d^2)) - 2*a*b*(x^2/(sqrt(-c^2*d*x^2 + d)*c^2*d) - 2/(s
qrt(-c^2*d*x^2 + d)*c^4*d))*arccosh(c*x) - a^2*(x^2/(sqrt(-c^2*d*x^2 + d)*
c^2*d) - 2/(sqrt(-c^2*d*x^2 + d)*c^4*d)) - b^2*((c^2*x^2 - 2)*log(c*x + sq
rt(c*x + 1)*sqrt(c*x - 1))^2/(sqrt(c*x + 1)*sqrt(-c*x + 1)*c^4*d^(3/2)) -
integrate(2*(c^4*x^4 - 3*c^2*x^2 + (c^3*x^3 - 2*c*x)*sqrt(c*x + 1)*sqrt(c*
x - 1) + 2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(((c^5*d^(3/2)*x^2 - c^
3*d^(3/2))*(c*x + 1)*sqrt(c*x - 1) + (c^6*d^(3/2)*x^3 - c^4*d^(3/2)*x)*sq
rt(c*x + 1))*sqrt(-c*x + 1)), x))
```

### 3.206.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac
")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.206.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$$

```
input int((x^3*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2),x)
```

```
output int((x^3*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2), x)
```

**3.207**  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

3.207.1 Optimal result . . . . . 1812  
 3.207.2 Mathematica [A] (warning: unable to verify) . . . . . 1813  
 3.207.3 Rubi [C] (verified) . . . . . 1813  
 3.207.4 Maple [B] (verified) . . . . . 1817  
 3.207.5 Fricas [F] . . . . . 1818  
 3.207.6 Sympy [F] . . . . . 1818  
 3.207.7 Maxima [F] . . . . . 1819  
 3.207.8 Giac [F] . . . . . 1819  
 3.207.9 Mupad [F(-1)] . . . . . 1819

**3.207.1 Optimal result**

Integrand size = 29, antiderivative size = 257

$$\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx = \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \frac{\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^2}{c^3d\sqrt{d-c^2dx^2}} - \frac{\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^3}{3bc^3d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\log(1-e^{2\operatorname{arccosh}(cx)})}{c^3d\sqrt{d-c^2dx^2}} - \frac{b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,e^{2\operatorname{arccosh}(cx)})}{c^3d\sqrt{d-c^2dx^2}}$$

output

```
x*(a+b*arccosh(c*x))^2/c^2/d/(-c^2*d*x^2+d)^(1/2)+(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d/(-c^2*d*x^2+d)^(1/2)-1/3*(a+b*arccosh(c*x))^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^3/d/(-c^2*d*x^2+d)^(1/2)-2*b*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d/(-c^2*d*x^2+d)^(1/2)-b^2*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d/(-c^2*d*x^2+d)^(1/2)
```

**3.207.2 Mathematica [A] (warning: unable to verify)**

Time = 2.12 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.05

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \frac{3a^2cdx + 3a^2\sqrt{d}\sqrt{d - c^2dx^2} \arctan\left(\frac{cx\sqrt{d - c^2dx^2}}{\sqrt{d}(-1 + c^2x^2)}\right) + 3abd(2cx\operatorname{arccosh}(cx) -$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

```
output (3*a^2*c*d*x + 3*a^2*Sqrt[d]*Sqrt[d - c^2*d*x^2]*ArcTan[(c*x*Sqrt[d - c^2*
d*x^2])/(Sqrt[d]*(-1 + c^2*x^2))] + 3*a*b*d*(2*c*x*ArcCosh[c*x] - Sqrt[(-1
+ c*x)/(1 + c*x)]*(1 + c*x)*(ArcCosh[c*x]^2 + 2*Log[Sqrt[(-1 + c*x)/(1 +
c*x)]*(1 + c*x)]) - b^2*d*(ArcCosh[c*x]*(-3*c*x*ArcCosh[c*x] + Sqrt[(-1 +
c*x)/(1 + c*x)]*(1 + c*x)*(ArcCosh[c*x]*(3 + ArcCosh[c*x]) + 6*Log[1 - E^
(-2*ArcCosh[c*x])])) - 3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, E
^(-2*ArcCosh[c*x])]))/(3*c^3*d^2*Sqrt[d - c^2*d*x^2])
```

**3.207.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 1.27 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.77, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$ , Rules used = {6349, 25, 6307, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx$$

$$\downarrow 6349$$

$$\frac{2b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x(a + \operatorname{barccosh}(cx))}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2dx^2}} - \frac{\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx}{c^2d} + \frac{x(a + \operatorname{barccosh}(cx))^2}{c^2d\sqrt{d - c^2dx^2}}$$

$$\downarrow 25$$

$$\frac{2b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{x(a + \operatorname{barccosh}(cx))}{(1 - cx)(cx + 1)} dx}{cd\sqrt{d - c^2dx^2}} - \frac{\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2dx^2}} dx}{c^2d} + \frac{x(a + \operatorname{barccosh}(cx))^2}{c^2d\sqrt{d - c^2dx^2}}$$

$$\downarrow 6307$$

---

3.207.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)(cx+1)} dx}{\frac{cd\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \\
& \qquad \qquad \qquad \frac{3bc^3d\sqrt{d-c^2dx^2}}{3bc^3d\sqrt{d-c^2dx^2}} \\
& \qquad \qquad \qquad \downarrow \text{6327} \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{\frac{cd\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \\
& \qquad \qquad \qquad \frac{3bc^3d\sqrt{d-c^2dx^2}}{3bc^3d\sqrt{d-c^2dx^2}} \\
& \qquad \qquad \qquad \downarrow \text{6328} \\
& - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+\operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \\
& \qquad \qquad \qquad \frac{3bc^3d\sqrt{d-c^2dx^2}}{3bc^3d\sqrt{d-c^2dx^2}} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -i(a+\operatorname{barccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}} + \\
& \qquad \qquad \qquad \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{3bc^3d\sqrt{d-c^2dx^2}}{3bc^3d\sqrt{d-c^2dx^2}} \\
& \qquad \qquad \qquad \downarrow \text{26} \\
& \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}} + \\
& \qquad \qquad \qquad \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{3bc^3d\sqrt{d-c^2dx^2}}{3bc^3d\sqrt{d-c^2dx^2}} \\
& \qquad \qquad \qquad \downarrow \text{4199} \\
& \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( 2i \int -\frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b} \right)}{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}} + \\
& \qquad \qquad \qquad \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{3bc^3d\sqrt{d-c^2dx^2}}{3bc^3d\sqrt{d-c^2dx^2}} \\
& \qquad \qquad \qquad \downarrow \text{25} \\
& \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b} \right)}{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}(a+\operatorname{barccosh}(cx))^3}} + \\
& \qquad \qquad \qquad \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{3bc^3d\sqrt{d-c^2dx^2}}{3bc^3d\sqrt{d-c^2dx^2}}
\end{aligned}$$

---

3.207.  $\int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

↓ 2620

$$\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(\frac{1}{2}b\int\log(1-e^{2\operatorname{arccosh}(cx)})\operatorname{darccosh}(cx)-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))\right)\right)}{c^2d\sqrt{d-c^2dx^2}} - \frac{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}}(a+\operatorname{barccosh}(cx))^3}{3bc^3d\sqrt{d-c^2dx^2}}$$

↓ 2715

$$\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(\frac{1}{4}b\int e^{-2\operatorname{arccosh}(cx)}\log(1-e^{2\operatorname{arccosh}(cx)})de^{2\operatorname{arccosh}(cx)}-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))\right)\right)}{c^2d\sqrt{d-c^2dx^2}} - \frac{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}}(a+\operatorname{barccosh}(cx))^3}{3bc^3d\sqrt{d-c^2dx^2}}$$

↓ 2838

$$\frac{\frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + 2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))-\frac{1}{4}b\operatorname{PolyLog}(2,e^{2\operatorname{arccosh}(cx)})\right)\right) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b}}{\frac{\frac{c^3d\sqrt{d-c^2dx^2}}{\sqrt{cx-1}\sqrt{cx+1}}(a+\operatorname{barccosh}(cx))^3}{3bc^3d\sqrt{d-c^2dx^2}}}$$

input `Int[(x^2*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

output `(x*(a + b*ArcCosh[c*x])^2)/(c^2*d*Sqrt[d - c^2*d*x^2]) - (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^3)/(3*b*c^3*d*Sqrt[d - c^2*d*x^2]) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/(c^3*d*Sqrt[d - c^2*d*x^2])`

### 3.207.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

---

3.207.  $\int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$



rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6307 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

rule 6327 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.207.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 737 vs. 2(255) = 510.

Time = 1.09 (sec) , antiderivative size = 738, normalized size of antiderivative = 2.87

method	result
default	$\frac{a^2 x}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{c^2 d \sqrt{c^2 d}} + \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arccosh}(cx)^3}{3d^2 c^3 (c^2 x^2 - 1)} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx}}{d^2 c^3 (c^2 x^2 - 1)}$
parts	$\frac{a^2 x}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{a^2 \arctan\left(\frac{\sqrt{c^2 d x}}{\sqrt{-c^2 d x^2 + d}}\right)}{c^2 d \sqrt{c^2 d}} + \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arccosh}(cx)^3}{3d^2 c^3 (c^2 x^2 - 1)} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx}}{d^2 c^3 (c^2 x^2 - 1)}$

```
input int(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

---

3.207. 
$$\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$$

output  $a^2x/c^2/d/(-c^2dx^2+d)^{(1/2)}-a^2/c^2/d/(c^2d)^{(1/2)}*\arctan((c^2d)^{(1/2)}x/(-c^2dx^2+d)^{(1/2)})+1/3*b^2*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{arccosh}(c*x)^3-b^2*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{arccosh}(c*x)^2-b^2*(-d*(c^2x^2-1))^{(1/2)}*\operatorname{arccosh}(c*x)^2/d^2/c^2/(c^2x^2-1)*x+2*b^2*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{arccosh}(c*x)*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+2*b^2*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+2*b^2*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{arccosh}(c*x)*\ln(1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+2*b^2*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+a*b*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{arccosh}(c*x)^2-2*a*b*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\operatorname{arccosh}(c*x)-2*a*b*(-d*(c^2x^2-1))^{(1/2)}*\operatorname{arccosh}(c*x)/d^2/c^2/(c^2x^2-1)*x+2*a*b*(-d*(c^2x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c^3/(c^2x^2-1)*\ln((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2-1)$

### 3.207.5 Fricas [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral((b^2*x^2*arccosh(c*x))^2 + 2*a*b*x^2*arccosh(c*x) + a^2*x^2)*sqrt(-c^2*d*x^2 + d)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

### 3.207.6 Sympy [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{3/2}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(3/2),x)`

---

3.207.  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

output `Integral(x**2*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

### 3.207.7 Maxima [F]

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `a^2*(x/(sqrt(-c^2*d*x^2 + d)*c^2*d) - arcsin(c*x)/(c^3*d^(3/2))) + integrate(b^2*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/(-c^2*d*x^2 + d)^(3/2) + 2*a*b*x^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(-c^2*d*x^2 + d)^(3/2), x)`

### 3.207.8 Giac [F]

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x^2/(-c^2*d*x^2 + d)^(3/2), x)`

### 3.207.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x^2*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2),x)`

output `int((x^2*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2), x)`

---

3.207.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$

**3.208** 
$$\int \frac{x(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$$

3.208.1 Optimal result . . . . . 1820  
 3.208.2 Mathematica [A] (verified) . . . . . 1821  
 3.208.3 Rubi [C] (verified) . . . . . 1821  
 3.208.4 Maple [A] (verified) . . . . . 1824  
 3.208.5 Fracas [F] . . . . . 1824  
 3.208.6 Sympy [F] . . . . . 1825  
 3.208.7 Maxima [F] . . . . . 1825  
 3.208.8 Giac [F] . . . . . 1825  
 3.208.9 Mupad [F(-1)] . . . . . 1826

**3.208.1 Optimal result**

Integrand size = 27, antiderivative size = 196

$$\begin{aligned} \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx &= \frac{(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\ &+ \frac{4b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{c^2d\sqrt{d-c^2dx^2}} \\ &+ \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})}{c^2d\sqrt{d-c^2dx^2}} \\ &- \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{c^2d\sqrt{d-c^2dx^2}} \end{aligned}$$

output

```
(a+b*arccosh(c*x))^2/c^2/d/(-c^2*d*x^2+d)^(1/2)+4*b*(a+b*arccosh(c*x))*arc
tanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^2/d/(-
c^2*d*x^2+d)^(1/2)+2*b^2*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-
1)^(1/2)*(c*x+1)^(1/2)/c^2/d/(-c^2*d*x^2+d)^(1/2)-2*b^2*polylog(2,c*x+(c*x
-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^2/d/(-c^2*d*x^2+d)^(
1/2)
```

### 3.208.2 Mathematica [A] (verified)

Time = 1.43 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.12

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \frac{a^2 + 2ab \left( \operatorname{arccosh}(cx) + \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \left( \log \left( \cosh \left( \frac{1}{2} \operatorname{arccosh}(cx) \right) \right) \right) - \log \left( \sinh \left( \frac{1}{2} \operatorname{arccosh}(cx) \right) \right) \right)}{(d - c^2 dx^2)^{3/2}}$$

input `Integrate[(x*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

output `(a^2 + 2*a*b*(ArcCosh[c*x] + Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(Log[Cos h[ArcCosh[c*x]/2]] - Log[Sinh[ArcCosh[c*x]/2]])) - b^2*(-(ArcCosh[c*x]*(ArcCosh[c*x] - 2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(Log[1 - E^(-ArcCosh[c*x])]) - Log[1 + E^(-ArcCosh[c*x])])) + 2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, -E^(-ArcCosh[c*x])] - 2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, E^(-ArcCosh[c*x])])/(c^2*d*Sqrt[d - c^2*d*x^2])`

### 3.208.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.64, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6329, 25, 6304, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx \\ & \quad \downarrow \text{6329} \\ & \frac{(a + \operatorname{barccosh}(cx))^2}{c^2 d \sqrt{d - c^2 dx^2}} - \frac{2b \sqrt{cx - 1} \sqrt{cx + 1} \int -\frac{a + \operatorname{barccosh}(cx)}{(1 - cx)(cx + 1)} dx}{cd \sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{25} \\ & \frac{2b \sqrt{cx - 1} \sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{(1 - cx)(cx + 1)} dx}{cd \sqrt{d - c^2 dx^2}} + \frac{(a + \operatorname{barccosh}(cx))^2}{c^2 d \sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{6304} \end{aligned}$$

---

3.208.  $\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx}{cd\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6318} \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{3042} \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{26} \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{4670} \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} (ib \int \log(1-e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}))}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{2715} \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} (ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{c^2d\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{2838} \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} (2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{c^2d\sqrt{d-c^2dx^2}}
\end{aligned}$$

input `Int[(x*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2),x]`

output `(a + b*ArcCosh[c*x])^2/(c^2*d*Sqrt[d - c^2*d*x^2]) - ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/(c^2*d*Sqrt[d - c^2*d*x^2])`

---

3.208.  $\int \frac{x(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

## 3.208.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6304 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(p_), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6318 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`



```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

### 3.208.4 Maple [A] (verified)

Time = 0.99 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.74

method	result
default	$\frac{a^2}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \left( 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} \ln(1+cx+\sqrt{cx-1} \sqrt{cx+1}) - 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} \ln(1-cx+\sqrt{cx-1} \sqrt{cx+1}) \right)}{c^2 d \sqrt{-c^2 d x^2 + d}}$
parts	$\frac{a^2}{c^2 d \sqrt{-c^2 d x^2 + d}} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \left( 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} \ln(1+cx+\sqrt{cx-1} \sqrt{cx+1}) - 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} \ln(1-cx+\sqrt{cx-1} \sqrt{cx+1}) \right)}{c^2 d \sqrt{-c^2 d x^2 + d}}$

```
input int(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```

```
output a^2/c^2/d/(-c^2*d*x^2+d)^(1/2)-b^2*(-d*(c^2*x^2-1))^(1/2)*(2*arccosh(c*x)*
(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-2*arccos
h(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+2
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-2
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+ar
ccosh(c*x)^2/d^2/c^2/(c^2*x^2-1)-2*a*b*(-d*(c^2*x^2-1))^(1/2)*((c*x-1)^(1
/2)*(c*x+1)^(1/2)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-(c*x-1)^(1/2)*(c*x
+1)^(1/2)*ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1)+arccosh(c*x))/d^2/c^2/(c^2
*x^2-1)
```

### 3.208.5 Fracas [F]

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x}{(-c^2 dx^2 + d)^{3/2}} dx$$

```
input integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fracas
")
```

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*x*arccosh(c*x)^2 + 2*a*b*x*arccosh(c*x) + a^2*x)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

### 3.208.6 Sympy [F]

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(3/2), x)`

output `Integral(x*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

### 3.208.7 Maxima [F]

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2), x, algorithm="maxima")`

output `a^2/(sqrt(-c^2*d*x^2 + d)*c^2*d) + integrate(b^2*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/(-c^2*d*x^2 + d)^(3/2) + 2*a*b*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(-c^2*d*x^2 + d)^(3/2), x)`

### 3.208.8 Giac [F]

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x/(-c^2*d*x^2 + d)^(3/2), x)`

---

3.208.  $\int \frac{x(a+b \operatorname{arccosh}(cx))^2}{(d-c^2 dx^2)^{3/2}} dx$

**3.208.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2),x)`output `int((x*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(3/2), x)`

**3.209**  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

3.209.1 Optimal result . . . . . 1827  
 3.209.2 Mathematica [A] (verified) . . . . . 1828  
 3.209.3 Rubi [C] (verified) . . . . . 1828  
 3.209.4 Maple [B] (verified) . . . . . 1831  
 3.209.5 Fracas [F] . . . . . 1832  
 3.209.6 Sympy [F] . . . . . 1832  
 3.209.7 Maxima [F] . . . . . 1833  
 3.209.8 Giac [F] . . . . . 1833  
 3.209.9 Mupad [F(-1)] . . . . . 1833

**3.209.1 Optimal result**

Integrand size = 26, antiderivative size = 198

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \frac{x(a + \operatorname{arccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{cd\sqrt{d - c^2dx^2}} - \frac{2b\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx)) \log(1 - e^{2\operatorname{arccosh}(cx)})}{cd\sqrt{d - c^2dx^2}} - \frac{b^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{cd\sqrt{d - c^2dx^2}}$$

```
output x*(a+b*arccosh(c*x))^2/d/(-c^2*d*x^2+d)^(1/2)+(a+b*arccosh(c*x))^2*(c*x-1)
^(1/2)*(c*x+1)^(1/2)/c/d/(-c^2*d*x^2+d)^(1/2)-2*b*(a+b*arccosh(c*x))*ln(1-
(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d/(-c^2
*d*x^2+d)^(1/2)-b^2*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)
^(1/2)*(c*x+1)^(1/2)/c/d/(-c^2*d*x^2+d)^(1/2)
```

### 3.209.2 Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.64

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \frac{x(a + \operatorname{barccosh}(cx))^2 + \frac{\sqrt{-1+cx}\sqrt{1+cx}((a+\operatorname{barccosh}(cx))(a+\operatorname{barccosh}(cx))-2b \log(1-e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2 dx^2}}}{d\sqrt{d-c^2 dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(d - c^2*d*x^2)^(3/2), x]`

output `(x*(a + b*ArcCosh[c*x])^2 + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((a + b*ArcCosh[c*x])*(a + b*ArcCosh[c*x] - 2*b*Log[1 - E^ArcCosh[c*x]] - 2*b*Log[1 + E^ArcCosh[c*x]]) - 2*b^2*PolyLog[2, -E^ArcCosh[c*x]] - 2*b^2*PolyLog[2, E^ArcCosh[c*x]]))/c)/(d*Sqrt[d - c^2*d*x^2])`

### 3.209.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.66 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.71, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.346$ , Rules used = {6314, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx \\ & \quad \downarrow \text{6314} \\ & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\ & \quad \downarrow \text{6328} \\ & \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+\operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{cd\sqrt{d-c^2dx^2}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.209.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
& \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} - \\
& \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \int -i(a + \operatorname{barccosh}(cx)) \tan\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{cd\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{26} \\
& \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \\
& \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \int (a + \operatorname{barccosh}(cx)) \tan\left(\operatorname{iarccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{cd\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{4199} \\
& \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \\
& \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( 2i \int -\frac{e^{2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 - e^{2\operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{cd\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{25} \\
& \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \\
& \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 - e^{2\operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{cd\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{2620} \\
& \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \\
& \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \left( \frac{1}{2} b \int \log(1 - e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) \right) \right)}{cd\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{2715} \\
& \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \\
& \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \left( \frac{1}{4} b \int e^{-2\operatorname{arccosh}(cx)} \log(1 - e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) \right) \right)}{cd\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{2838} \\
& \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \\
& \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \left( -\frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right) - \frac{i(a + \operatorname{barccosh}(cx))^2}{2b} \right)}{cd\sqrt{d - c^2dx^2}}
\end{aligned}$$

---

3.209.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx$

input `Int[(a + b*ArcCosh[c*x])^2/(d - c^2*d*x^2)^(3/2),x]`

output `(x*(a + b*ArcCosh[c*x])^2)/(d*Sqrt[d - c^2*d*x^2]) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])]))/4))/c*d*Sqrt[d - c^2*d*x^2]`

### 3.209.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4199 Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)
*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp
[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x
)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && In
tegerQ[4*k] && IGtQ[m, 0]
```

```
rule 6314 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2),
x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp
[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a
+ b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e},
x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]
```

```
rule 6328 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2),
x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]
], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]
```

### 3.209.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 577 vs. 2(204) = 408.

Time = 0.98 (sec) , antiderivative size = 578, normalized size of antiderivative = 2.92

method	result
default	$\frac{a^2 x}{d\sqrt{-c^2 d x^2 + d}} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arccosh}(cx)^2}{d^2 c(c^2 x^2 - 1)} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(cx)^2 x}{d^2 (c^2 x^2 - 1)} + \frac{2b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1}}{d^2 (c^2 x^2 - 1)}$
parts	$\frac{a^2 x}{d\sqrt{-c^2 d x^2 + d}} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1} \sqrt{cx + 1} \operatorname{arccosh}(cx)^2}{d^2 c(c^2 x^2 - 1)} - \frac{b^2 \sqrt{-d(c^2 x^2 - 1)} \operatorname{arccosh}(cx)^2 x}{d^2 (c^2 x^2 - 1)} + \frac{2b^2 \sqrt{-d(c^2 x^2 - 1)} \sqrt{cx - 1}}{d^2 (c^2 x^2 - 1)}$

```
input int((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)
```



output  $a^2/d*x/(-c^2*d*x^2+d)^{(1/2)}-b^2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c/(c^2*x^2-1)*\operatorname{arccosh}(c*x)^2-b^2*(-d*(c^2*x^2-1))^{(1/2)}*\operatorname{arccosh}(c*x)^2/d^2/(c^2*x^2-1)*x+2*b^2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c/(c^2*x^2-1)*\operatorname{arccosh}(c*x)*\ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+2*b^2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c/(c^2*x^2-1)*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+2*b^2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c/(c^2*x^2-1)*\operatorname{arccosh}(c*x)*\ln(1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})+2*b^2*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c/(c^2*x^2-1)*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})-2*a*b*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c/(c^2*x^2-1)*\operatorname{arccosh}(c*x)-2*a*b*(-d*(c^2*x^2-1))^{(1/2)}*\operatorname{arccosh}(c*x)/d^2/(c^2*x^2-1)*x+2*a*b*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/c/(c^2*x^2-1)*\ln((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2-1)$

### 3.209.5 Fracas [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

### 3.209.6 Sympy [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{3/2}} dx$$

input `integrate((a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**3/2, x)`

**3.209.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-a*b*c*sqrt(-1/(c^4*d))*log(x^2 - 1/c^2)/d + b^2*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/(-c^2*d*x^2 + d)^(3/2), x) + 2*a*b*x*arccosh(c*x)/(sqrt(-c^2*d*x^2 + d)*d) + a^2*x/(sqrt(-c^2*d*x^2 + d)*d)`

**3.209.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/(-c^2*d*x^2 + d)^(3/2), x)`

**3.209.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^2/(d - c^2*d*x^2)^(3/2),x)`

output `int((a + b*acosh(c*x))^2/(d - c^2*d*x^2)^(3/2), x)`

$$3.210 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx$$

3.210.1 Optimal result . . . . .	1834
3.210.2 Mathematica [A] (warning: unable to verify) . . . . .	1835
3.210.3 Rubi [A] (verified) . . . . .	1836
3.210.4 Maple [F] . . . . .	1841
3.210.5 Fracas [F] . . . . .	1842
3.210.6 Sympy [F] . . . . .	1842
3.210.7 Maxima [F] . . . . .	1842
3.210.8 Giac [F] . . . . .	1843
3.210.9 Mupad [F(-1)] . . . . .	1843

### 3.210.1 Optimal result

Integrand size = 29, antiderivative size = 471

$$\begin{aligned} \int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx &= \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\ &+ \frac{2\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\ &+ \frac{4b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\ &+ \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\ &- \frac{2ib\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\ &+ \frac{2ib\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\ &- \frac{2b^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\ &+ \frac{2ib^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\ &- \frac{2ib^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \end{aligned}$$

output  $(a+b\operatorname{arccosh}(cx))^2/d/(-c^2dx^2+d)^{(1/2)}+2*(a+b\operatorname{arccosh}(cx))^2*\arctan(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)})*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}+4*b*(a+b\operatorname{arccosh}(cx))*\operatorname{arctanh}(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)})*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}+2*b^2*\operatorname{polylog}(2,-cx-(cx-1)^{(1/2)}*(cx+1)^{(1/2)})*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}+2*I*b*(a+b\operatorname{arccosh}(cx))*\operatorname{polylog}(2,-I*(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)}))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}+2*I*b*(a+b\operatorname{arccosh}(cx))*\operatorname{polylog}(2,I*(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)}))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}-2*b^2*\operatorname{polylog}(2,cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)})*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}+2*I*b^2*\operatorname{polylog}(3,-I*(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)}))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}-2*I*b^2*\operatorname{polylog}(3,I*(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)}))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}$

### 3.210.2 Mathematica [A] (warning: unable to verify)

Time = 3.08 (sec) , antiderivative size = 613, normalized size of antiderivative = 1.30

$$\int \frac{(a + b\operatorname{arccosh}(cx))^2}{x(d - c^2dx^2)^{3/2}} dx =$$

$$\frac{a^2\sqrt{d-c^2dx^2}}{-1+c^2x^2} - a^2\sqrt{d}\log(cx) + a^2\sqrt{d}\log\left(d + \sqrt{d}\sqrt{d - c^2dx^2}\right) + \frac{2iabd\left(i\operatorname{arccosh}(cx) + \sqrt{\frac{-1+cx}{1+cx}}(1+cx)\operatorname{arccosh}(cx)\right)\log\left(\frac{d + \sqrt{d}\sqrt{d - c^2dx^2}}{d - \sqrt{d}\sqrt{d - c^2dx^2}}\right)}{2d}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x*(d - c^2*d*x^2)^(3/2)), x]`

---

3.210.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx$

output

```

-(((a^2*Sqrt[d - c^2*d*x^2])/(-1 + c^2*x^2) - a^2*Sqrt[d]*Log[c*x] + a^2*S
qrt[d]*Log[d + Sqrt[d]*Sqrt[d - c^2*d*x^2]] + ((2*I)*a*b*d*(I*ArcCosh[c*x]
+ Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 - I/E^ArcCosh[c
*x]] - Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 + I/E^ArcCo
sh[c*x]] + I*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Cosh[ArcCosh[c*x]/2]
] - I*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Sinh[ArcCosh[c*x]/2]] + Sqr
t[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, (-I)/E^ArcCosh[c*x]] - Sqrt[(
-1 + c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, I/E^ArcCosh[c*x]]))/Sqrt[d - c^2
*d*x^2] + (b^2*d*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*((Sqrt[(-1 + c*x)/(1
+ c*x)]*ArcCosh[c*x]^2)/(1 - c*x) + 2*ArcCosh[c*x]*Log[1 - E^(-ArcCosh[c*
x])] + I*ArcCosh[c*x]^2*Log[1 - I/E^ArcCosh[c*x]] - I*ArcCosh[c*x]^2*Log[1
+ I/E^ArcCosh[c*x]] - 2*ArcCosh[c*x]*Log[1 + E^(-ArcCosh[c*x])] + 2*PolyL
og[2, -E^(-ArcCosh[c*x])] + (2*I)*ArcCosh[c*x]*PolyLog[2, (-I)/E^ArcCosh[c
*x]] - (2*I)*ArcCosh[c*x]*PolyLog[2, I/E^ArcCosh[c*x]] - 2*PolyLog[2, E^(-
ArcCosh[c*x])] + (2*I)*PolyLog[3, (-I)/E^ArcCosh[c*x]] - (2*I)*PolyLog[3,
I/E^ArcCosh[c*x]]))/Sqrt[d - c^2*d*x^2])/d^2)

```

### 3.210.3 Rubi [A] (verified)

Time = 2.33 (sec) , antiderivative size = 260, normalized size of antiderivative = 0.55, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.517$ , Rules used = {6351, 25, 6304, 6318, 3042, 26, 4670, 2715, 2838, 6361, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(a + b \operatorname{arccosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx \\
& \quad \downarrow \text{6351} \\
& -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{a+b\operatorname{arccosh}(cx)}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2 dx^2}} dx}{d} + \frac{(a + b \operatorname{arccosh}(cx))^2}{d\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow \text{25} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2 dx^2}} dx}{d} + \frac{(a + b \operatorname{arccosh}(cx))^2}{d\sqrt{d - c^2 dx^2}} \\
& \quad \downarrow \text{6304} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2 x^2} dx}{d\sqrt{d-c^2 dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2 dx^2}} dx}{d} + \frac{(a + b \operatorname{arccosh}(cx))^2}{d\sqrt{d - c^2 dx^2}}
\end{aligned}$$

---

3.210.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& \downarrow 6318 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\
& \downarrow 3042 \\
& \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \\
& \quad \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\
& \downarrow 26 \\
& \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \\
& \quad \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\
& \downarrow 4670 \\
& \frac{2ib\sqrt{cx-1}\sqrt{cx+1} (ib \int \log(1-e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2i\operatorname{arctanh})}{d\sqrt{d-c^2dx^2}} \\
& \quad \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\
& \downarrow 2715 \\
& \frac{2ib\sqrt{cx-1}\sqrt{cx+1} (ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \\
& \quad \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\
& \downarrow 2838 \\
& \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \\
& \frac{2ib\sqrt{cx-1}\sqrt{cx+1} (2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \\
& \quad \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \\
& \downarrow 6361
\end{aligned}$$

---

3.210.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx$

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}}$$

↓ 3042

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx))^2 \csc(\operatorname{iarcosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}}$$

↓ 4668

$$\frac{\sqrt{cx-1}\sqrt{cx+1}(-2ib \int (a+\operatorname{barccosh}(cx)) \log(1 - ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + 2ib \int (a+\operatorname{barccosh}(cx)) \log(1 + ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx))}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}}$$

↓ 3011

$$\frac{\sqrt{cx-1}\sqrt{cx+1}(2ib(b \int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)))}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}}$$

↓ 2720

$$\frac{\sqrt{cx-1}\sqrt{cx+1}(2ib(b \int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)))}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}}$$

↓ 7143

---

3.210.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}(2\arctan(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))^2+2ib(b\operatorname{PolyLog}(3,-ie^{\operatorname{arccosh}(cx)})-\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(cx)})\frac{d\sqrt{d-c^2dx^2}}{d\sqrt{d-c^2dx^2}})}{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})\frac{d\sqrt{d-c^2dx^2}}{d\sqrt{d-c^2dx^2}})}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x*(d - c^2*d*x^2)^(3/2)),x]`

output `(a + b*ArcCosh[c*x])^2/(d*Sqrt[d - c^2*d*x^2]) - ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/(d*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]])))/(d*Sqrt[d - c^2*d*x^2])`

### 3.210.3.1 Defintions of rubi rules used

rule 25 `Int[-(F_x_), x_Symbol] := Simp[Identity[-1] Int[F_x, x], x]`

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

---

3.210.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx$



rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.))*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

rule 6361 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.210.4 Maple [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `int((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(3/2),x)`

output `int((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(3/2),x)`

**3.210.5 Fricas [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^4*d^2*x^5 - 2*c^2*d^2*x^3 + d^2*x), x)`

**3.210.6 Sympy [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))**2/x/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.210.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-a^2*(log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(3/2) - 1/(sqrt(-c^2*d*x^2 + d)*d)) + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/((-c^2*d*x^2 + d)^(3/2)*x) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(3/2)*x), x)`

---

3.210.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx$

**3.210.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(3/2)*x), x)`

**3.210.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x(d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x*(d - c^2*d*x^2)^(3/2)),x)`

output `int((a + b*acosh(c*x))^2/(x*(d - c^2*d*x^2)^(3/2)), x)`

**3.211**  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^2(d-c^2dx^2)^{3/2}} dx$

3.211.1 Optimal result . . . . . 1844  
 3.211.2 Mathematica [A] (warning: unable to verify) . . . . . 1845  
 3.211.3 Rubi [C] (verified) . . . . . 1845  
 3.211.4 Maple [B] (verified) . . . . . 1852  
 3.211.5 Fricas [F] . . . . . 1853  
 3.211.6 Sympy [F] . . . . . 1853  
 3.211.7 Maxima [F] . . . . . 1853  
 3.211.8 Giac [F] . . . . . 1854  
 3.211.9 Mupad [F(-1)] . . . . . 1854

**3.211.1 Optimal result**

Integrand size = 29, antiderivative size = 341

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{x^2(d - c^2dx^2)^{3/2}} dx = -\frac{(a + \operatorname{arccosh}(cx))^2}{dx\sqrt{d - c^2dx^2}} + \frac{2c^2x(a + \operatorname{arccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \frac{2c\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} - \frac{4bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} - \frac{4bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))\log(1 - e^{2\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} - \frac{b^2c\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} - \frac{b^2c\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}}$$

output

```

-(a+b*arccosh(c*x))^2/d/x/(-c^2*d*x^2+d)^(1/2)+2*c^2*x*(a+b*arccosh(c*x))^2/d/(-c^2*d*x^2+d)^(1/2)+2*c*(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)-4*b*c*(a+b*arccosh(c*x))*arctanh((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)-4*b*c*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)-b^2*c*polylog(2, -(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2)-b^2*c*polylog(2, (c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(-c^2*d*x^2+d)^(1/2))
    
```

3.211.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^2(d-c^2dx^2)^{3/2}} dx$

**3.211.2 Mathematica [A] (warning: unable to verify)**

Time = 0.52 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.37

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx = \frac{2ab(-1 + 2c^2 x^2) \operatorname{arccosh}(cx) + b^2 \operatorname{arccosh}(cx)^2 + a(-a + 2ac^2 x^2 - 2bcx\sqrt{-1 + c^2 x^2})}{dx\sqrt{d - c^2 dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^2*(d - c^2*d*x^2)^(3/2)),x]`output `(2*a*b*(-1 + 2*c^2*x^2)*ArcCosh[c*x] + b^2*ArcCosh[c*x]^2 + a*(-a + 2*a*c^2*x^2 - 2*b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[c*x] - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Log[1 - c^2*x^2]))/(d*x*Sqrt[d - c^2*d*x^2])`**3.211.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.47 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.82, number of steps used = 20, number of rules used = 19,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.655$ , Rules used = {6347, 25, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx \\ & \quad \downarrow \text{6347} \\ & 2c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a + \operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d - c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{25} \\ & 2c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a + \operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d - c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d - c^2 dx^2}} \\ & \quad \downarrow \text{6314} \end{aligned}$$

---

3.211.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx$

$$\begin{aligned}
& 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right) - \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6327} \\
& 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right) - \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{6328} \\
& 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+\operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{cd\sqrt{d-c^2dx^2}} \right) - \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{3042} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} + \\
& 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -i(a+\operatorname{barccosh}(cx)) \tan(i\operatorname{arccosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx)}{cd\sqrt{d-c^2dx^2}} \right) - \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{26} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} + \\
& 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \tan(i\operatorname{arccosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx)}{cd\sqrt{d-c^2dx^2}} \right) - \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{4199}
\end{aligned}$$

---

3.211.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
 & 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( 2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 - e^{2\operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))}{2b} \right)}{cd\sqrt{d - c^2dx^2}} \right. \\
 & \qquad \left. \frac{2bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2x^2)} dx}{d\sqrt{d - c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d - c^2dx^2}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{25} \\
 & 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))}{1 - e^{2\operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a + \operatorname{barccosh}(cx))}{2b} \right)}{cd\sqrt{d - c^2dx^2}} \right. \\
 & \qquad \left. \frac{2bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2x^2)} dx}{d\sqrt{d - c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d - c^2dx^2}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{2620} \\
 & - \frac{2bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2x^2)} dx}{d\sqrt{d - c^2dx^2}} + \\
 & 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \left( \frac{1}{2} b \int \log(1 - e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) \right) \right)}{cd\sqrt{d - c^2dx^2}} \right. \\
 & \qquad \left. \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d - c^2dx^2}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{2715} \\
 & - \frac{2bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2x^2)} dx}{d\sqrt{d - c^2dx^2}} + \\
 & 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \left( \frac{1}{4} b \int e^{-2\operatorname{arccosh}(cx)} \log(1 - e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) \right) \right)}{cd\sqrt{d - c^2dx^2}} \right. \\
 & \qquad \left. \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d - c^2dx^2}} \right) \\
 & \qquad \qquad \qquad \downarrow \text{2838} \\
 & - \frac{2bc\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{x(1 - c^2x^2)} dx}{d\sqrt{d - c^2dx^2}} + \\
 & 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2dx^2}} + \frac{2ib\sqrt{cx - 1}\sqrt{cx + 1} \left( -2i \left( -\frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog} \right) \right)}{cd\sqrt{d - c^2dx^2}} \right. \\
 & \qquad \left. \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d - c^2dx^2}} \right)
 \end{aligned}$$

---

3.211.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2(d - c^2dx^2)^{3/2}} dx$



$$\begin{aligned}
 & \downarrow \text{6331} \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \\
 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog} \right. \right.}{cd\sqrt{d-c^2dx^2}} \\
 & \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right) \right. \\
 & \downarrow \text{5984} \\
 & \frac{4bc\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \\
 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog} \right. \right.}{cd\sqrt{d-c^2dx^2}} \\
 & \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right) \right. \\
 & \downarrow \text{3042} \\
 & \frac{4bc\sqrt{cx-1}\sqrt{cx+1} \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \\
 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog} \right. \right.}{cd\sqrt{d-c^2dx^2}} \\
 & \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right) \right. \\
 & \downarrow \text{26} \\
 & \frac{4ibc\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \\
 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog} \right. \right.}{cd\sqrt{d-c^2dx^2}} \\
 & \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right) \right. \\
 & \downarrow \text{4670}
 \end{aligned}$$

---

3.211.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{3/2}} dx$

$$\frac{4ibc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}ib \int \log(1-e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2}ib \int \log(1+e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + i\operatorname{arctan}\right)}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog}\right)}{cd\sqrt{d-c^2dx^2}}$$

$$2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right) \downarrow 2715$$

$$\frac{4ibc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1-e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1+e^{2\operatorname{arccosh}(cx)})\right)}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog}\right)}{cd\sqrt{d-c^2dx^2}}$$

$$2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right) \downarrow 2838$$

$$\frac{4ibc\sqrt{cx-1}\sqrt{cx+1}\left(i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + \frac{1}{4}ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4}ib \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})\right)}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx)) - \frac{1}{4}b \operatorname{PolyLog}\right)}{cd\sqrt{d-c^2dx^2}}$$

$$2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right)$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^2*(d - c^2*d*x^2)^(3/2)),x]`

output `-((a + b*ArcCosh[c*x])^2/(d*x*Sqrt[d - c^2*d*x^2])) + ((4*I)*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])]) + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])])/(d*Sqrt[d - c^2*d*x^2]) + 2*c^2*((x*(a + b*ArcCosh[c*x])^2)/(d*Sqrt[d - c^2*d*x^2]) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/(c*d*Sqrt[d - c^2*d*x^2]))`

## 3.211.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2620 `Int[((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)] / ((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp [((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4199 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`
- rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_) + (e_.)*(x_)^2)), x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1)))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

### 3.211.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1270 vs.  $2(359) = 718$ .

Time = 1.48 (sec) , antiderivative size = 1271, normalized size of antiderivative = 3.73

method	result	size
default	Expression too large to display	1271
parts	Expression too large to display	1271

input `int((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```
a^2*(-1/d/x/(-c^2*d*x^2+d)^(1/2)+2*c^2/d*x/(-c^2*d*x^2+d)^(1/2))-b^2*(-4*a
rccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2-4*arccosh(c*x)*l
n(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+4*arccosh(c*x)*ln(1+(c*x+(c*x
-1)^(1/2)*(c*x+1)^(1/2))^2)*x^4*c^4+4*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*
(c*x+1)^(1/2))*x^4*c^4+4*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)
)*x^4*c^4-4*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^2*c^2
+2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^4*c^4+4*polylog(2,-c*
x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^4*c^4+4*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+
1)^(1/2))*x^4*c^4-2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^2*c^
2-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2)
)*x^3*c^3+arccosh(c*x)^2-4*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2
*c^2-4*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^2*c^2+2*arccosh(c*x)*(
c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x*c-4
*(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^
3*c^3+(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1
/2))^2)*x*c+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,-c*x-(c*x-1)^(1/2)*(c*
x+1)^(1/2))*x*c+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,c*x+(c*x-1)^(1/2)*
(c*x+1)^(1/2))*x*c-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*polylog(2,-(c*x+(c*x-1)^(
1/2)*(c*x+1)^(1/2))^2)*x^3*c^3-4*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*
ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*x^3*c^3-4*arccosh(c*x)*(c*x-1)^(1...
```

**3.211.5 Fricas [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2), x)`

**3.211.6 Sympy [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^2 (-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))**2/x**2/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x**2*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.211.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `a*b*c*(sqrt(-d)*log(c*x + 1)/d^2 + sqrt(-d)*log(c*x - 1)/d^2 + 2*sqrt(-d)*log(x)/d^2) + 2*(2*c^2*x/(sqrt(-c^2*d*x^2 + d)*d) - 1/(sqrt(-c^2*d*x^2 + d)*d*x))*a*b*arccosh(c*x) + (2*c^2*x/(sqrt(-c^2*d*x^2 + d)*d) - 1/(sqrt(-c^2*d*x^2 + d)*d*x))*a^2 + b^2*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/((-c^2*d*x^2 + d)^(3/2)*x^2), x)`

---

3.211.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx$

**3.211.8 Giac [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(3/2)*x^2), x)`

**3.211.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^2*(d - c^2*d*x^2)^(3/2)),x)`

output `int((a + b*acosh(c*x))^2/(x^2*(d - c^2*d*x^2)^(3/2)), x)`

$$3.212 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{3/2}} dx$$

3.212.1 Optimal result . . . . .	1855
3.212.2 Mathematica [B] (warning: unable to verify) . . . . .	1856
3.212.3 Rubi [A] (verified) . . . . .	1857
3.212.4 Maple [F] . . . . .	1866
3.212.5 Fracas [F] . . . . .	1866
3.212.6 Sympy [F] . . . . .	1866
3.212.7 Maxima [F] . . . . .	1867
3.212.8 Giac [F] . . . . .	1867
3.212.9 Mupad [F(-1)] . . . . .	1867

### 3.212.1 Optimal result

Integrand size = 29, antiderivative size = 650

$$\begin{aligned} \int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^3(d - c^2dx^2)^{3/2}} dx = & \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))}{dx\sqrt{d - c^2dx^2}} \\ & + \frac{3c^2(a + b\operatorname{arccosh}(cx))^2}{2d\sqrt{d - c^2dx^2}} - \frac{(a + b\operatorname{arccosh}(cx))^2}{2dx^2\sqrt{d - c^2dx^2}} \\ & + \frac{3c^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \\ & - \frac{b^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d\sqrt{d - c^2dx^2}} \\ & + \frac{4bc^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \\ & + \frac{2b^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \\ & - \frac{3ibc^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \\ & + \frac{3ibc^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \\ & - \frac{2b^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \\ & + \frac{3ib^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \\ & - \frac{3ib^2c^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \end{aligned}$$

---

3.212.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{3/2}} dx$



output  $\frac{3}{2}c^2(a+b\operatorname{arccosh}(cx))^2/d/(-c^2dx^2+d)^{(1/2)}-1/2(a+b\operatorname{arccosh}(cx))$   
 $\wedge 2/d/x^2/(-c^2dx^2+d)^{(1/2)}+b*c*(a+b\operatorname{arccosh}(cx))*(cx-1)^{(1/2)}*(cx+1)$   
 $\wedge(1/2)/d/x/(-c^2dx^2+d)^{(1/2)}+3*c^2*(a+b\operatorname{arccosh}(cx))^2*\operatorname{arctan}(cx+(cx$   
 $-1)^{(1/2)}*(cx+1)^{(1/2))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}$   
 $-b^2*c^2*\operatorname{arctan}((cx-1)^{(1/2)}*(cx+1)^{(1/2))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/$   
 $d/(-c^2dx^2+d)^{(1/2)}+4*b*c^2*(a+b\operatorname{arccosh}(cx))*\operatorname{arctanh}(cx+(cx-1)^{(1/2)}$   
 $)*(cx+1)^{(1/2))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}+2*b^2*$   
 $c^2*\operatorname{polylog}(2,-cx-(cx-1)^{(1/2)}*(cx+1)^{(1/2))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}$   
 $)/d/(-c^2dx^2+d)^{(1/2)}-3*I*b*c^2*(a+b\operatorname{arccosh}(cx))*\operatorname{polylog}(2,-I*(cx+(c$   
 $*x-1)^{(1/2)}*(cx+1)^{(1/2)))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{($   
 $1/2)}+3*I*b*c^2*(a+b\operatorname{arccosh}(cx))*\operatorname{polylog}(2,I*(cx+(cx-1)^{(1/2)}*(cx+1)^{($   
 $1/2)))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}-2*b^2*c^2*\operatorname{polylo}$   
 $g(2,cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2d$   
 $*x^2+d)^{(1/2)}+3*I*b^2*c^2*\operatorname{polylog}(3,-I*(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)))*$   
 $(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2dx^2+d)^{(1/2)}-3*I*b^2*c^2*\operatorname{polylog}(3,I$   
 $*(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/d/(-c^2d*$   
 $x^2+d)^{(1/2)}$

### 3.212.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 5362 vs.  $2(650) = 1300$ .

Time = 63.67 (sec) , antiderivative size = 5362, normalized size of antiderivative = 8.25

$$\int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^3(d - c^2dx^2)^{3/2}} dx = \text{Result too large to show}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^3*(d - c^2*d*x^2)^(3/2)),x]`

output `Result too large to show`

**3.212.3 Rubi [A] (verified)**

Time = 4.98 (sec) , antiderivative size = 433, normalized size of antiderivative = 0.67, number of steps used = 28, number of rules used = 27,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.931$ , Rules used = {6347, 25, 6327, 6347, 103, 218, 6318, 3042, 26, 4670, 2715, 2838, 6351, 25, 6304, 6318, 3042, 26, 4670, 2715, 2838, 6361, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6347} \\
 & \frac{3}{2} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{3/2}} dx + \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{a+\operatorname{barccosh}(cx)}{x^2(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{3}{2} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{3/2}} dx - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{3}{2} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{3/2}} dx - \frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2 x^2)} dx}{d\sqrt{d-c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{6347} \\
 & \frac{\frac{3}{2} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{3/2}} dx - bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2 x^2} dx + bc \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{d\sqrt{d-c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2 dx^2}} \\
 & \quad \downarrow \text{103} \\
 & \frac{\frac{3}{2} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{3/2}} dx - bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2 x^2} dx + bc^2 \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{a+\operatorname{barccosh}(cx)}{x} \right)}{d\sqrt{d-c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2 dx^2}}
 \end{aligned}$$

---

3.212.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2 dx^2)^{3/2}} dx$

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(c^2\int\frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2}dx-\frac{a+\operatorname{barccosh}(cx)}{x}+bc\arctan(\sqrt{cx-1}\sqrt{cx+1})\right)}{d\sqrt{d-c^2dx^2}}+\frac{3}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 218

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-c\int\frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)}d\operatorname{arccosh}(cx)-\frac{a+\operatorname{barccosh}(cx)}{x}+bc\arctan(\sqrt{cx-1}\sqrt{cx+1})\right)}{d\sqrt{d-c^2dx^2}}+\frac{3}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 6318

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-c\int i(a+\operatorname{barccosh}(cx))\csc(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)-\frac{a+\operatorname{barccosh}(cx)}{x}+bc\arctan(\sqrt{cx-1}\sqrt{cx+1})\right)}{d\sqrt{d-c^2dx^2}}+\frac{3}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 3042

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-c\int i(a+\operatorname{barccosh}(cx))\csc(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)-\frac{a+\operatorname{barccosh}(cx)}{x}+bc\arctan(\sqrt{cx-1}\sqrt{cx+1})\right)}{d\sqrt{d-c^2dx^2}}+\frac{3}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 26

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-ic\int(a+\operatorname{barccosh}(cx))\csc(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)-\frac{a+\operatorname{barccosh}(cx)}{x}+bc\arctan(\sqrt{cx-1}\sqrt{cx+1})\right)}{d\sqrt{d-c^2dx^2}}+\frac{3}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 4670

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-ic(ib\int\log(1-e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)-ib\int\log(1+e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)+2i\arctan(\sqrt{cx-1}\sqrt{cx+1}))\right)}{d\sqrt{d-c^2dx^2}}+\frac{3}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 2715

---

3.212.  $\int\frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{3/2}}dx$

$$\begin{aligned}
 & \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-ic(ib\int e^{-\operatorname{arccosh}(cx)}\log(1-e^{\operatorname{arccosh}(cx)})de^{\operatorname{arccosh}(cx)}-ib\int e^{-\operatorname{arccosh}(cx)}\log(1+e^{\operatorname{arccosh}(cx)})\right)}{d\sqrt{d-c^2}} \\
 & \frac{\frac{3}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}}{2838} \\
 & \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,\right)}{d\sqrt{d-c^2dx^2}} \\
 & \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}} \\
 & \downarrow 6351 \\
 & \frac{\frac{3}{2}c^2\left(-\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{a+\operatorname{barccosh}(cx)}{(1-cx)(cx+1)}dx}{d\sqrt{d-c^2dx^2}}+\frac{\int\frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}}dx}{d}+\frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}\right)-}{d\sqrt{d-c^2dx^2}} \\
 & \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,\right)}{d\sqrt{d-c^2dx^2}} \\
 & \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}} \\
 & \downarrow 25 \\
 & \frac{\frac{3}{2}c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{a+\operatorname{barccosh}(cx)}{(1-cx)(cx+1)}dx}{d\sqrt{d-c^2dx^2}}+\frac{\int\frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}}dx}{d}+\frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}\right)-}{d\sqrt{d-c^2dx^2}} \\
 & \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,\right)}{d\sqrt{d-c^2dx^2}} \\
 & \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}} \\
 & \downarrow 6304 \\
 & \frac{\frac{3}{2}c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2}dx}{d\sqrt{d-c^2dx^2}}+\frac{\int\frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}}dx}{d}+\frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}\right)-}{d\sqrt{d-c^2dx^2}} \\
 & \frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,\right)}{d\sqrt{d-c^2dx^2}} \\
 & \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}
 \end{aligned}$$

3.212.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{3/2}} dx$

↓ 6318

$$\frac{\frac{3}{2}c^2 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right) - bc\sqrt{cx-1}\sqrt{cx+1} \left( -ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})) \right)}{d\sqrt{d-c^2dx^2}}$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 3042

$$\frac{\frac{3}{2}c^2 \left( \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right) - bc\sqrt{cx-1}\sqrt{cx+1} \left( -ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})) \right)}{d\sqrt{d-c^2dx^2}}$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 26

$$\frac{\frac{3}{2}c^2 \left( \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right) - bc\sqrt{cx-1}\sqrt{cx+1} \left( -ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})) \right)}{d\sqrt{d-c^2dx^2}}$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 4670

$$\frac{\frac{3}{2}c^2 \left( -\frac{2ib\sqrt{cx-1}\sqrt{cx+1} (ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)))}{d\sqrt{d-c^2dx^2}} \right) - bc\sqrt{cx-1}\sqrt{cx+1} \left( -ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})) \right)}{d\sqrt{d-c^2dx^2}}$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}}$$

↓ 2715

---

3.212.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{3/2}} dx$

$$\frac{3}{2}c^2 \left( -\frac{2ib\sqrt{cx-1}\sqrt{cx+1}(ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)})}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{bc\sqrt{cx-1}\sqrt{cx+1}(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}} \right. \\ \left. \downarrow 2838 \right.$$

$$\frac{3}{2}c^2 \left( \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{bc\sqrt{cx-1}\sqrt{cx+1}(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}} \right. \\ \left. \downarrow 6361 \right.$$

$$\frac{3}{2}c^2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1} \int \frac{(a+\operatorname{barccosh}(cx))^2}{cx} \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{bc\sqrt{cx-1}\sqrt{cx+1}(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}} \right. \\ \left. \downarrow 3042 \right.$$

$$\frac{3}{2}c^2 \left( \frac{\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx))^2 \csc(\operatorname{iarccosh}(cx) + \frac{\pi}{2}) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{bc\sqrt{cx-1}\sqrt{cx+1}(-ic(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2\sqrt{d-c^2dx^2}} \right. \\ \left. \downarrow 4668 \right.$$

---

3.212.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{3/2}} dx$



```

output -1/2*(a + b*ArcCosh[c*x])^2/(d*x^2*Sqrt[d - c^2*d*x^2]) - (b*c*Sqrt[-1 + c
*x]*Sqrt[1 + c*x]*(-(a + b*ArcCosh[c*x])/x) + b*c*ArcTan[Sqrt[-1 + c*x]*S
qrt[1 + c*x]] - I*c*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] +
I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/ (d*Sqr
t[d - c^2*d*x^2]) + (3*c^2*((a + b*ArcCosh[c*x])^2/(d*Sqrt[d - c^2*d*x^2])
- ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((2*I)*(a + b*ArcCosh[c*x])*ArcTa
nh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^Ar
cCosh[c*x]]))/ (d*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(
a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-((a + b*ArcCosh[c
*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]])
- (2*I)*b*(-((a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyL
og[3, I*E^ArcCosh[c*x]])))/ (d*Sqrt[d - c^2*d*x^2]))/2

```

### 3.212.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]

```

```

rule 103 Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x
_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sq
rt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d
*e - f*(b*c + a*d), 0]

```

```

rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

```

```

rule 2715 Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```



rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n/(b*c*n*Log[F])]), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(p_), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6318  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)]^{(n_.)}/((d_.) + (e_.)*(x_)^2), x\_Symbol] \rightarrow \text{Simp}[-(c*d)^{-1} \text{Subst}[\text{Int}[(a + b*x)^n * \text{Csch}[x], x], x, \text{ArcCosh}[c*x]], x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[n, 0]$

rule 6327  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)]^{(n_.)}*((f_.)*(x_))^{(m_.)}*((d_.) + (e_.)*(x_))^{(p_.)}*((d_2_.) + (e_2_.)*(x_))^{(p_2_.)}, x\_Symbol] \rightarrow \text{Int}[(f*x)^m*(d_1*d_2 + e_1*e_2*x^2)^p*(a + b*\text{ArcCosh}[c*x])^n, x] /;$   $\text{FreeQ}\{a, b, c, d_1, e_1, d_2, e_2, f, m, n\}, x\} \ \&\& \ \text{EqQ}[d_2*e_1 + d_1*e_2, 0] \ \&\& \ \text{IntegerQ}[p]$

rule 6347  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)]^{(n_.)}*((f_.)*(x_))^{(m_.)}*((d_.) + (e_.)*(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[(f*x)^{(m+1)}*(d + e*x^2)^{(p+1)}*((a + b*\text{ArcCosh}[c*x])^n/(d*f*(m+1))), x] + (\text{Simp}[c^2*((m+2*p+3)/(f^2*(m+1)))] \text{Int}[(f*x)^{(m+2)}*(d + e*x^2)^p*(a + b*\text{ArcCosh}[c*x])^n, x], x] + \text{Simp}[b*c*(n/(f*(m+1)))*\text{Simp}[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] \text{Int}[(f*x)^{(m+1)}*(1 + c*x)^{(p+1/2)}*(-1 + c*x)^{(p+1/2)}*(a + b*\text{ArcCosh}[c*x])^{(n-1)}, x], x]) /;$   $\text{FreeQ}\{a, b, c, d, e, f, p\}, x\} \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{ILtQ}[m, -1]$

rule 6351  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)]^{(n_.)}*((f_.)*(x_))^{(m_.)}*((d_.) + (e_.)*(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[(-(f*x)^{(m+1)}*(d + e*x^2)^{(p+1)}*((a + b*\text{ArcCosh}[c*x])^n/(2*d*f*(p+1))), x] + (\text{Simp}[(m+2*p+3)/(2*d*(p+1))] \text{Int}[(f*x)^m*(d + e*x^2)^{(p+1)}*(a + b*\text{ArcCosh}[c*x])^n, x], x] - \text{Simp}[b*c*(n/(2*f*(p+1)))*\text{Simp}[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] \text{Int}[(f*x)^{(m+1)}*(1 + c*x)^{(p+1/2)}*(-1 + c*x)^{(p+1/2)}*(a + b*\text{ArcCosh}[c*x])^{(n-1)}, x], x]) /;$   $\text{FreeQ}\{a, b, c, d, e, f, m\}, x\} \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{!GtQ}[m, 1] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{IntegerQ}[p] \ || \ \text{EqQ}[n, 1])$

rule 6361  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)]^{(n_.)}*(x_)^{(m_.)}/\text{Sqrt}[(d_.) + (e_.)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[(1/c^{(m+1)})*\text{Simp}[\text{Sqrt}[1 + c*x]*(\text{Sqrt}[-1 + c*x]/\text{Sqrt}[d + e*x^2])] \text{Subst}[\text{Int}[(a + b*x)^n * \text{Cosh}[x]^m, x], x, \text{ArcCosh}[c*x]], x] /;$   $\text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{EqQ}[c^2*d + e, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IntegerQ}[m]$

rule 7143  $\text{Int}[\text{PolyLog}[n_., (c_.)*((a_.) + (b_.)*(x_))^{(p_.)}]/((d_.) + (e_.)*(x_)), x\_Symbol] \rightarrow \text{Simp}[\text{PolyLog}[n + 1, c*(a + b*x)^p]/(e*p), x] /;$   $\text{FreeQ}\{a, b, c, d, e, n, p\}, x\} \ \&\& \ \text{EqQ}[b*d, a*e]$

---

3.212. 
$$\int \frac{(a+b\text{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{3/2}} dx$$

**3.212.4 Maple [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 (-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `int((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(3/2),x)`

output `int((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(3/2),x)`

**3.212.5 Fricas [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^4*d^2*x^7 - 2*c^2*d^2*x^5 + d^2*x^3), x)`

**3.212.6 Sympy [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^3 (-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))**2/x**3/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x**3*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.212.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `-1/2*(3*c^2*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(3/2) - 3*c^2/(sqrt(-c^2*d*x^2 + d)*d) + 1/(sqrt(-c^2*d*x^2 + d)*d*x^2))*a^2 + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/((-c^2*d*x^2 + d)^(3/2)*x^3) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(3/2)*x^3), x)`

**3.212.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(3/2)*x^3), x)`

**3.212.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^3*(d - c^2*d*x^2)^(3/2)),x)`

output `int((a + b*acosh(c*x))^2/(x^3*(d - c^2*d*x^2)^(3/2)), x)`

---

3.212.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{3/2}} dx$

**3.213**  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^4(d-c^2dx^2)^{3/2}} dx$

3.213.1 Optimal result . . . . . 1868  
 3.213.2 Mathematica [A] (warning: unable to verify) . . . . . 1869  
 3.213.3 Rubi [C] (verified) . . . . . 1870  
 3.213.4 Maple [B] (verified) . . . . . 1878  
 3.213.5 Fricas [F] . . . . . 1879  
 3.213.6 Sympy [F] . . . . . 1880  
 3.213.7 Maxima [F] . . . . . 1880  
 3.213.8 Giac [F] . . . . . 1880  
 3.213.9 Mupad [F(-1)] . . . . . 1881

**3.213.1 Optimal result**

Integrand size = 29, antiderivative size = 496

$$\begin{aligned} \int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^4(d - c^2dx^2)^{3/2}} dx &= \frac{b^2c^2(1 - cx)(1 + cx)}{3dx\sqrt{d - c^2dx^2}} \\ &+ \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))}{3dx^2\sqrt{d - c^2dx^2}} - \frac{(a + b\operatorname{arccosh}(cx))^2}{3dx^3\sqrt{d - c^2dx^2}} \\ &- \frac{4c^2(a + b\operatorname{arccosh}(cx))^2}{3dx\sqrt{d - c^2dx^2}} + \frac{8c^4x(a + b\operatorname{arccosh}(cx))^2}{3d\sqrt{d - c^2dx^2}} \\ &+ \frac{8c^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))^2}{3d\sqrt{d - c^2dx^2}} \\ &- \frac{20bc^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{3d\sqrt{d - c^2dx^2}} \\ &- \frac{16bc^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))\log(1 - e^{2\operatorname{arccosh}(cx)})}{3d\sqrt{d - c^2dx^2}} \\ &- \frac{5b^2c^3\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{3d\sqrt{d - c^2dx^2}} \\ &- \frac{b^2c^3\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{d\sqrt{d - c^2dx^2}} \end{aligned}$$

output  $\frac{1}{3}b^2c^2(-cx+1)(cx+1)/d/x/(-c^2dx^2+d)^{1/2}-1/3(a+b\operatorname{arccosh}(cx))^2/d/x^3/(-c^2dx^2+d)^{1/2}-4/3c^2(a+b\operatorname{arccosh}(cx))^2/d/x/(-c^2dx^2+d)^{1/2}+8/3c^4x(a+b\operatorname{arccosh}(cx))^2/d/(-c^2dx^2+d)^{1/2}+1/3b^2c(a+b\operatorname{arccosh}(cx))(cx-1)^{1/2}(cx+1)^{1/2}/d/x^2/(-c^2dx^2+d)^{1/2}+8/3c^3(a+b\operatorname{arccosh}(cx))^2(cx-1)^{1/2}(cx+1)^{1/2}/d/(-c^2dx^2+d)^{1/2}-20/3b^2c^3(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}((cx+(cx-1)^{1/2})(cx+1)^{1/2}))^2(cx-1)^{1/2}(cx+1)^{1/2}/d/(-c^2dx^2+d)^{1/2}-16/3b^2c^3(a+b\operatorname{arccosh}(cx))\ln(1-(cx+(cx-1)^{1/2})(cx+1)^{1/2}))^2(cx-1)^{1/2}(cx+1)^{1/2}/d/(-c^2dx^2+d)^{1/2}-5/3b^2c^3\operatorname{polylog}(2,-(cx+(cx-1)^{1/2})(cx+1)^{1/2}))^2(cx-1)^{1/2}(cx+1)^{1/2}/d/(-c^2dx^2+d)^{1/2}-b^2c^3\operatorname{polylog}(2,(cx+(cx-1)^{1/2})(cx+1)^{1/2}))^2(cx-1)^{1/2}(cx+1)^{1/2}/d/(-c^2dx^2+d)^{1/2}$

### 3.213.2 Mathematica [A] (warning: unable to verify)

Time = 1.97 (sec) , antiderivative size = 529, normalized size of antiderivative = 1.07

$$\int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^4(d - c^2dx^2)^{3/2}} dx = \frac{a^2(-1 - 4c^2x^2 + 8c^4x^4) + ab\left(6c^4x^4\operatorname{arccosh}(cx) + \sqrt{\frac{-1+cx}{1+cx}}(1+cx)\right)(cx + 2\sqrt{-1+cx})}{x^4(d - c^2dx^2)^{3/2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^4*(d - c^2*d*x^2)^(3/2)),x]`

output  $(a^2(-1 - 4c^2x^2 + 8c^4x^4) + a*b*(6c^4x^4\operatorname{ArcCosh}[c*x] + \operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(c*x + 2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x]) + 2*c^2*x^2*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(5*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x] - c*x*(5*\operatorname{Log}[c*x] + 3*\operatorname{Log}[\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)]))) + b^2*(c^2*x^2 - c^4*x^4 + c*x*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x] + 3*c^4*x^4*\operatorname{ArcCosh}[c*x]^2 + (-1 + c*x)*(1 + c*x)*\operatorname{ArcCosh}[c*x]^2 + 5*c^2*x^2*(-1 + c*x)*(1 + c*x)*\operatorname{ArcCosh}[c*x]^2 - 8*c^3*x^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x]^2 - 6*c^3*x^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - E^(-2*\operatorname{ArcCosh}[c*x])] - 10*c^3*x^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + E^(-2*\operatorname{ArcCosh}[c*x])] + 5*c^3*x^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{PolyLog}[2, -E^(-2*\operatorname{ArcCosh}[c*x])] + 3*c^3*x^3*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\operatorname{PolyLog}[2, E^(-2*\operatorname{ArcCosh}[c*x])])))/(3*d*x^3*\operatorname{Sqrt}[d - c^2*d*x^2])$

**3.213.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 4.18 (sec) , antiderivative size = 472, normalized size of antiderivative = 0.95, number of steps used = 24, number of rules used = 23,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.793$ , Rules used = {6347, 25, 6327, 6347, 25, 106, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6347} \\
 & \frac{4}{3} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{a+\operatorname{barccosh}(cx)}{x^3(1-cx)(cx+1)} dx}{3d\sqrt{d-c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{4}{3} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^3(1-cx)(cx+1)} dx}{3d\sqrt{d-c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{4}{3} c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{3/2}} dx - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^3(1-c^2x^2)} dx}{3d\sqrt{d-c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{6347} \\
 & \frac{4}{3} c^2 \left( 2c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int -\frac{a+\operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2dx^2}} \right) - \\
 & \quad \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2} bc \int \frac{1}{x^2\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} \right)}{3d\sqrt{d-c^2dx^2}} - \\
 & \quad \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

---

3.213.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{3/2}} dx$

$$\frac{4}{3}c^2 \left( 2c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2 dx^2}} \right) -$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2 x^2)} dx + \frac{1}{2}bc \int \frac{1}{x^2\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} \right)}{\frac{3d\sqrt{d-c^2 dx^2}}{(a + \operatorname{barccosh}(cx))^2} - \frac{3dx^3\sqrt{d-c^2 dx^2}}{3dx^3\sqrt{d-c^2 dx^2}}}$$

↓ 106

$$\frac{4}{3}c^2 \left( 2c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx\sqrt{d-c^2 dx^2}} \right) -$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2 x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{\frac{3d\sqrt{d-c^2 dx^2}}{(a + \operatorname{barccosh}(cx))^2} - \frac{3dx^3\sqrt{d-c^2 dx^2}}{3dx^3\sqrt{d-c^2 dx^2}}}$$

↓ 6314

$$\frac{4}{3}c^2 \left( 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2 x^2} dx}{d\sqrt{d-c^2 dx^2}} + \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2 dx^2}} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} \right) -$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2 x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{\frac{3d\sqrt{d-c^2 dx^2}}{(a + \operatorname{barccosh}(cx))^2} - \frac{3dx^3\sqrt{d-c^2 dx^2}}{3dx^3\sqrt{d-c^2 dx^2}}}$$

↓ 6327

$$\frac{4}{3}c^2 \left( 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2 x^2} dx}{d\sqrt{d-c^2 dx^2}} + \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2 dx^2}} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)(cx+1)} dx}{d\sqrt{d-c^2 dx^2}} \right) -$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2 x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{\frac{3d\sqrt{d-c^2 dx^2}}{(a + \operatorname{barccosh}(cx))^2} - \frac{3dx^3\sqrt{d-c^2 dx^2}}{3dx^3\sqrt{d-c^2 dx^2}}}$$

↓ 6328



$$\begin{aligned}
& \frac{4}{3}c^2 \left( 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+\operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{cd\sqrt{d-c^2dx^2}} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1}}{d\sqrt{d-c^2dx^2}} \right) \\
& \quad \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))^2} \\
& \quad \frac{3dx^3\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{3042} \\
& \quad \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2}} + \\
& \frac{4}{3}c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -i(a + \operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right) \right) \\
& \quad \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{26} \\
& \quad \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2}} + \\
& \frac{4}{3}c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a + \operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right) \right) \\
& \quad \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{4199} \\
& \frac{4}{3}c^2 \left( 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( 2i \int -\frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} \operatorname{darccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right)}{cd\sqrt{d-c^2dx^2}} \right) \right) \\
& \quad \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))^2} \\
& \quad \frac{3dx^3\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}} \\
& \quad \downarrow \text{25}
\end{aligned}$$

---

3.213.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{3/2}} dx$

$$\begin{aligned}
& \frac{4}{3}c^2 \left( 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))}{2x} \right)}{cd\sqrt{d-c^2dx^2}} \right. \right. \\
& \left. \left. - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))^2} \right) \right. \\
& \left. - \frac{3dx^3\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}} \right) \downarrow \text{2620} \\
& - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2}} + \\
& \frac{4}{3}c^2 \left( - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{2} b \int \log \right) \right)}{(a + \operatorname{barccosh}(cx))^2} \right. \right. \\
& \left. \left. - \frac{3dx^3\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}} \right) \right) \downarrow \text{2715} \\
& - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2}} + \\
& \frac{4}{3}c^2 \left( - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{4} b \int e^{-\operatorname{arccosh}(cx)} \right) \right)}{(a + \operatorname{barccosh}(cx))^2} \right. \right. \\
& \left. \left. - \frac{3dx^3\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}} \right) \right) \downarrow \text{2838} \\
& - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2}} + \\
& \frac{4}{3}c^2 \left( - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log \right) \right)}{(a + \operatorname{barccosh}(cx))^2} \right. \right. \\
& \left. \left. - \frac{3dx^3\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}} \right) \right) \downarrow \text{6331} \\
& - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a + \operatorname{barccosh}(cx))^2}
\end{aligned}$$

$$\frac{4}{3}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(-2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right) \right)$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( c^2 \left( - \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx) \right) - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{3d\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}}$$

↓ 5984

$$\frac{4}{3}c^2 \left( \frac{4bc\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(-2c^2 \int (a+\operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right) \right)$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2c^2 \int (a+\operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{3d\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}}$$

↓ 3042

$$\frac{4}{3}c^2 \left( \frac{4bc\sqrt{cx-1}\sqrt{cx+1} \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(-2c^2 \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right) \right)$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2c^2 \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{3d\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}}$$

↓ 26

$$\frac{4}{3}c^2 \left( \frac{4ibc\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{d\sqrt{d-c^2dx^2}} + 2c^2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}(-2ic^2 \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx))}{d\sqrt{d-c^2dx^2}} \right) \right)$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2ic^2 \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) - \frac{a+\operatorname{barccosh}(cx)}{2x^2} + \frac{bc\sqrt{cx-1}\sqrt{cx+1}}{2x} \right)}{3d\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2}$$

$$\frac{3d\sqrt{d-c^2dx^2}}{3dx^3\sqrt{d-c^2dx^2}}$$

↓ 4670

---

3.213.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{3/2}} dx$

$$\frac{\frac{4}{3}c^2 \left( \frac{4ibc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2}ib \int \log(1-e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2}ib \int \log(1+e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + \dots \right)}{d\sqrt{d-c^2dx^2}} \right)}{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2ic^2 \left( \frac{1}{2}ib \int \log(1-e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2}ib \int \log(1+e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) + \dots \right) \right)}{3d\sqrt{d-c^2dx^2}}$$

$$\frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}}$$

↓ 2715

$$\frac{\frac{4}{3}c^2 \left( \frac{4ibc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1-e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1+e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} \right)}{d\sqrt{d-c^2dx^2}} \right)}{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2ic^2 \left( \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1-e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} \log(1+e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} \right) \right)}{3d\sqrt{d-c^2dx^2}}$$

$$\frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}}$$

↓ 2838

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2ic^2 (i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})) (a + \operatorname{barccosh}(cx)) + \frac{1}{4}ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4}ib \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right)}{3d\sqrt{d-c^2dx^2}}$$

$$\frac{\frac{4}{3}c^2 \left( \frac{4ibc\sqrt{cx-1}\sqrt{cx+1} \left( i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4}ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4}ib \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right)}{d\sqrt{d-c^2dx^2}} \right)}{(a + \operatorname{barccosh}(cx))^2}{3dx^3\sqrt{d-c^2dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^4*(d - c^2*d*x^2)^(3/2)), x]`

```
output -1/3*(a + b*ArcCosh[c*x])^2/(d*x^3*Sqrt[d - c^2*d*x^2]) - (2*b*c*Sqrt[-1 +
c*x]*Sqrt[1 + c*x]*((b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*x) - (a + b*Arc
Cosh[c*x])/(2*x^2) - (2*I)*c^2*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCos
h[c*x])]) + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^
(2*ArcCosh[c*x])]))/(3*d*Sqrt[d - c^2*d*x^2]) + (4*c^2*(-((a + b*ArcCosh[
c*x])^2/(d*x*Sqrt[d - c^2*d*x^2])) + ((4*I)*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*
x]*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])]) + (I/4)*b*PolyLog[2
, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])]))/(d*Sqrt[
d - c^2*d*x^2]) + 2*c^2*((x*(a + b*ArcCosh[c*x])^2)/(d*Sqrt[d - c^2*d*x^2]
) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^
2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b
*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/(c*d*Sqrt[d - c^2*d*x^2])))/3
```

### 3.213.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 106 Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_
)^(p_)), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1
))/(m + 1)*(b*c - a*d)*(b*e - a*f)), x] /; FreeQ[{a, b, c, d, e, f, m, n,
p}, x] && EqQ[Simplify[m + n + p + 3], 0] && EqQ[a*d*f*(m + 1) + b*c*f*(n +
1) + b*d*e*(p + 1), 0] && NeQ[m, -1]
```

```
rule 2620 Int[(((F_)^(g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_)^(m_.)))/
((a_) + (b_.)*((F_)^(g_.)*((e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2715 Int[Log[(a_) + (b_.)*((F_)^(e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

---


$$3.213. \int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^4(d-c^2dx^2)^{3/2}} dx$$

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_)^(m_.))*Sech[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_.)/((d_.) + (e_.)*(x_.)^2),  
x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]  
], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)/((x_.)*((d_.) + (e_.)*(x_.)^2)),  
x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x  
, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IG  
tQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_.) + (e_  
.)*(x_.)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +  
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1  
))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp  
[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(  
f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^  
(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] &&  
GtQ[n, 0] && ILtQ[m, -1]`

### 3.213.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2162 vs.  $2(486) = 972$ .

Time = 1.40 (sec) , antiderivative size = 2163, normalized size of antiderivative = 4.36

method	result	size
default	Expression too large to display	2163
parts	Expression too large to display	2163

input `int((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(3/2),x,method=_RETURNVERBOSE)`

output

```

-1/3*a*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(16*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4+16*arccosh(c*x)*c^5*x^5-6*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^5*c^5-10*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^5*c^5-8*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-16*c^3*x^3*arccosh(c*x)+6*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^3*c^3+10*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^3*c^3+c^3*x^3-2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)-c*x)/d^2/(c^4*x^4-2*c^2*x^2+1)/x^3-32/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)*x^7*c^10+40/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)*x^5*c^8-7/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)*x*c^4-1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)/x*c^2+1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)/x^3*arccosh(c*x)^2-1/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)/x^2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c+10/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(c^2*x^2-1)/d^2*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*c^3+2*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(c^2*x^2-1)/d^2*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*c^3-32/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)*x^3*arccosh(c*x)*(c*x-1)*(c*x+1)*c^6+64/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7*c^2*x^2-1)*x^2*arccosh(c*x)^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^5-8/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^2/(8*c^4*x^4-7...

```

### 3.213.5 Fracas [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{3/2} x^4} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^4*d^2*x^8 - 2*c^2*d^2*x^6 + d^2*x^4), x)`



**3.213.6 Sympy [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^4 (-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))**2/x**4/(-c**2*d*x**2+d)**(3/2), x)`

output `Integral((a + b*acosh(c*x))**2/(x**4*(-d*(c*x - 1)*(c*x + 1))**(3/2)), x)`

**3.213.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^4} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(3/2), x, algorithm="maxima")`

output `1/3*(8*c^4*x/(sqrt(-c^2*d*x^2 + d)*d) - 4*c^2/(sqrt(-c^2*d*x^2 + d)*d*x) - 1/(sqrt(-c^2*d*x^2 + d)*d*x^3))*a^2 + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/((-c^2*d*x^2 + d)^(3/2)*x^4) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(3/2)*x^4), x)`

**3.213.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^4} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(3/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(3/2)*x^4), x)`

**3.213.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^4 (d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^4*(d - c^2*d*x^2)^(3/2)),x)`output `int((a + b*acosh(c*x))^2/(x^4*(d - c^2*d*x^2)^(3/2)), x)`

$$3.214 \quad \int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

3.214.1 Optimal result . . . . .	1882
3.214.2 Mathematica [A] (warning: unable to verify) . . . . .	1883
3.214.3 Rubi [C] (verified) . . . . .	1884
3.214.4 Maple [A] (verified) . . . . .	1893
3.214.5 Fracas [F] . . . . .	1894
3.214.6 Sympy [F(-1)] . . . . .	1895
3.214.7 Maxima [F] . . . . .	1895
3.214.8 Giac [F(-2)] . . . . .	1896
3.214.9 Mupad [F(-1)] . . . . .	1896

### 3.214.1 Optimal result

Integrand size = 29, antiderivative size = 568

$$\begin{aligned} \int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx = & -\frac{b^2x^2}{3c^4d^2\sqrt{d-c^2dx^2}} - \frac{16abx\sqrt{-1+cx}\sqrt{1+cx}}{3c^5d^2\sqrt{d-c^2dx^2}} \\ & - \frac{7b^2(1-cx)(1+cx)}{3c^6d^2\sqrt{d-c^2dx^2}} - \frac{16b^2x\sqrt{-1+cx}\sqrt{1+cx}\operatorname{arccosh}(cx)}{3c^5d^2\sqrt{d-c^2dx^2}} \\ & + \frac{11bx\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{3c^5d^2\sqrt{d-c^2dx^2}} \\ & + \frac{bx^3\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{3c^3d^2(1-c^2x^2)\sqrt{d-c^2dx^2}} + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\ & - \frac{4x^2(a+b\operatorname{arccosh}(cx))^2}{3c^4d^2\sqrt{d-c^2dx^2}} - \frac{8\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{3c^6d^3} \\ & - \frac{22b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{3c^6d^2\sqrt{d-c^2dx^2}} \\ & - \frac{11b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})}{3c^6d^2\sqrt{d-c^2dx^2}} \\ & + \frac{11b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{3c^6d^2\sqrt{d-c^2dx^2}} \end{aligned}$$

---


$$3.214. \quad \int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

output  $\frac{1}{3}x^4(a+b\operatorname{arccosh}(cx))^2/c^2/d/(-c^2dx^2+d)^{3/2}-1/3b^2x^2/c^4/d^2/(-c^2dx^2+d)^{1/2}-7/3b^2(-cx+1)(cx+1)/c^6/d^2/(-c^2dx^2+d)^{1/2}-4/3x^2(a+b\operatorname{arccosh}(cx))^2/c^4/d^2/(-c^2dx^2+d)^{1/2}-16/3abx(cx-1)^{1/2}(cx+1)^{1/2}/c^5/d^2/(-c^2dx^2+d)^{1/2}-16/3b^2x\operatorname{arccosh}(cx)(cx-1)^{1/2}(cx+1)^{1/2}/c^5/d^2/(-c^2dx^2+d)^{1/2}+11/3bxx(a+b\operatorname{arccosh}(cx))(cx-1)^{1/2}(cx+1)^{1/2}/c^5/d^2/(-c^2dx^2+d)^{1/2}+1/3bx^3(a+b\operatorname{arccosh}(cx))(cx-1)^{1/2}(cx+1)^{1/2}/c^3/d^2/(-c^2x^2+1)/(-c^2dx^2+d)^{1/2}-22/3b(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}(cx+(cx-1)^{1/2})(cx+1)^{1/2})(cx-1)^{1/2}(cx+1)^{1/2}/c^6/d^2/(-c^2dx^2+d)^{1/2}-11/3b^2\operatorname{polylog}(2,-cx-(cx-1)^{1/2})(cx+1)^{1/2})(cx-1)^{1/2}(cx+1)^{1/2}/c^6/d^2/(-c^2dx^2+d)^{1/2}+11/3b^2\operatorname{polylog}(2,cx+(cx-1)^{1/2})(cx+1)^{1/2})(cx-1)^{1/2}(cx+1)^{1/2}/c^6/d^2/(-c^2dx^2+d)^{1/2}-8/3(a+b\operatorname{arccosh}(cx))^2(-c^2dx^2+d)^{1/2}/c^6/d^3$

### 3.214.2 Mathematica [A] (warning: unable to verify)

Time = 4.39 (sec) , antiderivative size = 490, normalized size of antiderivative = 0.86

$$\int \frac{x^5(a + b\operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx =$$

$$\frac{8a^2(8 - 12c^2x^2 + 3c^4x^4) + 2ab(25\operatorname{arccosh}(cx) - 36\operatorname{arccosh}(cx)\cosh(2\operatorname{arccosh}(cx)) + 3\operatorname{arccosh}(cx)\cosh(4\operatorname{arccosh}(cx)))}{(d - c^2dx^2)^{3/2}}$$

input `Integrate[(x^5*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]`

output

```

-1/24*(8*a^2*(8 - 12*c^2*x^2 + 3*c^4*x^4) + 2*a*b*(25*ArcCosh[c*x] - 36*Ar
cCosh[c*x]*Cosh[2*ArcCosh[c*x]] + 3*ArcCosh[c*x]*Cosh[4*ArcCosh[c*x]] + 33
*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Cosh[ArcCosh[c*x]/2]] - 33*sqrt[
(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[Sinh[ArcCosh[c*x]/2]] + 4*Sinh[2*ArcCo
sh[c*x]] - 11*Log[Cosh[ArcCosh[c*x]/2]]*Sinh[3*ArcCosh[c*x]] + 11*Log[Sinh
[ArcCosh[c*x]/2]]*Sinh[3*ArcCosh[c*x]] - 3*Sinh[4*ArcCosh[c*x]]) + b^2*(22
+ 25*ArcCosh[c*x]^2 - 4*(7 + 9*ArcCosh[c*x]^2)*Cosh[2*ArcCosh[c*x]] + 3*(
2 + ArcCosh[c*x]^2)*Cosh[4*ArcCosh[c*x]] - 66*sqrt[(-1 + c*x)/(1 + c*x)]*(
1 + c*x)*ArcCosh[c*x]*Log[1 - E^(-ArcCosh[c*x])] + 66*sqrt[(-1 + c*x)/(1 +
c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 + E^(-ArcCosh[c*x])] + 88*((-1 + c*x)/
(1 + c*x))^(3/2)*(1 + c*x)^3*PolyLog[2, -E^(-ArcCosh[c*x])] - 88*((-1 + c*
x)/(1 + c*x))^(3/2)*(1 + c*x)^3*PolyLog[2, E^(-ArcCosh[c*x])] + 8*ArcCosh[
c*x]*Sinh[2*ArcCosh[c*x]] + 22*ArcCosh[c*x]*Log[1 - E^(-ArcCosh[c*x])] *Sin
h[3*ArcCosh[c*x]] - 22*ArcCosh[c*x]*Log[1 + E^(-ArcCosh[c*x])] *Sinh[3*ArcC
osh[c*x]] - 6*ArcCosh[c*x]*Sinh[4*ArcCosh[c*x]])))/(c^6*d*(d - c^2*d*x^2)^(
3/2))

```

### 3.214.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 3.47 (sec) , antiderivative size = 569, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.621$ , Rules used = {6349, 6327, 6349, 25, 109, 27, 83, 6327, 6329, 2009, 6353, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6349} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^4(a + \operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \frac{4 \int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{3c^2d} + \frac{x^4(a + \operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^4(a + \operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \frac{4 \int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{3c^2d} + \frac{x^4(a + \operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{6349}
 \end{aligned}$$

---

3.214.  $\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{2c^2} + \frac{b \int \frac{x^3}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3cd^2\sqrt{d-c^2dx^2}} \\
 & - \frac{4 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -\frac{x^2(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right)}{3c^2d} \\
 & + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{2c^2} + \frac{b \int \frac{x^3}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3cd^2\sqrt{d-c^2dx^2}} \\
 & - \frac{4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right)}{3c^2d} \\
 & + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{109} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{2c^2} + \frac{b \left( -\frac{\int -\frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c^2} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3cd^2\sqrt{d-c^2dx^2}} \\
 & - \frac{4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right)}{3c^2d} \\
 & + \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

---

3.214.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{2c^2} + \frac{b \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c^2} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right) \\
& \frac{3cd^2\sqrt{d-c^2dx^2}}{4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right)} + \\
& \frac{3c^2d}{x^4(a+b\operatorname{arccosh}(cx))^2} \\
& \frac{3c^2d}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 83 \\
& 4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right) + \\
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{2c^2} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right) + \\
& \frac{3cd^2\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} \\
& \frac{3c^2d}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 6327 \\
& 4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{cd\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right) + \\
& 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{2c^2} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right) + \\
& \frac{3cd^2\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} \\
& \frac{3c^2d}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 6329
\end{aligned}$$

---

3.214.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$4 \left( -\frac{2 \left( -\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx)) dx}{c\sqrt{d-c^2dx^2}} - \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} \right)}{c^2d} + \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{cd\sqrt{d-c^2dx^2}} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d} \right)$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{2c^2} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right)$$


---


$$\frac{3cd^2\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} - \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}}$$

↓ 2009

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{cd\sqrt{d-c^2dx^2}} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{c\sqrt{d-c^2dx^2}} \right)}{c^2d} \right)$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \int \frac{x^2(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{2c^2} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right)$$


---


$$\frac{3cd^2\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} - \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}}$$

↓ 6353

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx) dx}{1-c^2x^2}}{c^2} + \frac{b \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} \right)}{cd\sqrt{d-c^2dx^2}} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^2}{c^2d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^2}{c\sqrt{d-c^2dx^2}} \right)}{c^2d} \right)$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx) dx}{1-c^2x^2}}{c^2} + \frac{b \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} \right)}{2c^2} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{x^2}{c^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right)$$


---


$$\frac{3cd^2\sqrt{d-c^2dx^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} - \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}}$$

↓ 83

---

3.214.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$



$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{cd\sqrt{d-c^2dx^2}} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{c^2d} \right)}{c^2d} \right)$$

---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{c^2} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{2c^2} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{1}{2c} \right)}{2c} \right)$$

---


$$\frac{3cd^2\sqrt{d-c^2dx^2}}{3c^2d} \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}}$$

↓ 6318

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{cd\sqrt{d-c^2dx^2}} + \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2 \left( -\frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))}{c^2d} \right)}{c^2d} \right)$$

---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( \frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{c^3} - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3} \right)}{2c^2} + \frac{x^3(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{c^4} - \frac{1}{2c} \right)}{2c} \right)$$

---


$$\frac{3cd^2\sqrt{d-c^2dx^2}}{3c^2d} \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}}$$

↓ 3042

---

3.214.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int i(a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{cd\sqrt{d-c^2dx^2}} \right) + \frac{x^2(a+b\operatorname{arccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}}}{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( -\frac{\int i(a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{2c^2} \right) + \frac{3c^2d}{2c^2(1-}} \right. \right.$$

$$\left. \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \right) \frac{3cd^2\sqrt{d-c^2dx^2}}$$

↓ 26

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \int (a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{cd\sqrt{d-c^2dx^2}} \right) + \frac{x^2(a+b\operatorname{arccosh}(cx))}{c^2d\sqrt{d-c^2dx^2}}}{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( -\frac{i \int (a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx) - \frac{x(a+b\operatorname{arccosh}(cx))}{c^2} + \frac{b\sqrt{cx-1}\sqrt{cx+1}}{c^3}}{2c^2} \right) + \frac{3c^2d}{2c^2(1-}} \right. \right.$$

$$\left. \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \right) \frac{3cd^2\sqrt{d-c^2dx^2}}$$

↓ 4670

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right) (a+b\operatorname{arccosh}(cx))}{c^3}}{cd\sqrt{d-c^2dx^2}} \right)}{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( -\frac{i \left( ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right) (a+b\operatorname{arccosh}(cx))}{c^3}}{2c^2} \right) + \frac{3c^2d}{2c^2(1-}} \right. \right.$$

$$\left. \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \right) \frac{3cd^2\sqrt{d-c^2dx^2}}$$

↓ 2715

3.214.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) dx e^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) dx e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{c^3} \right)}{cd\sqrt{d-c^2dx^2}} \right)$$

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( -\frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) dx e^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) dx e^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{c^3} \right)}{2c^2} \right)$$

$$\frac{x^4(a + b\operatorname{arccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}}$$

↓ 2838

$$4 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{c^3} \right)}{cd\sqrt{d-c^2dx^2}} \right)$$

$$2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{3 \left( -\frac{i \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{c^3} \right)}{2c^2} \right)$$

$$\frac{x^4(a + b\operatorname{arccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}}$$

$3cd^2\sqrt{d - c^2dx^2}$

input `Int[(x^5*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]`

```

output (x^4*(a + b*ArcCosh[c*x])^2)/(3*c^2*d*(d - c^2*d*x^2)^(3/2)) + (2*b*Sqrt[-
1 + c*x]*Sqrt[1 + c*x]*((b*(-x^2/(c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (2
*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^4))/(2*c) + (x^3*(a + b*ArcCosh[c*x]))/(2
*c^2*(1 - c^2*x^2)) - (3*((b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^3 - (x*(a + b
*ArcCosh[c*x]))/c^2 - (I*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x
]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c^
3))/((2*c^2)))/(3*c*d^2*Sqrt[d - c^2*d*x^2]) - (4*((x^2*(a + b*ArcCosh[c*x]
)^2)/(c^2*d*Sqrt[d - c^2*d*x^2]) - (2*(-((Sqrt[d - c^2*d*x^2]*(a + b*ArcCo
sh[c*x])^2)/(c^2*d)) - (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a*x - (b*Sqrt[-1
+ c*x]*Sqrt[1 + c*x])/c + b*x*ArcCosh[c*x]))/(c*Sqrt[d - c^2*d*x^2])))/(c
^2*d) + (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]
)/c^3 - (x*(a + b*ArcCosh[c*x]))/c^2 - (I*((2*I)*(a + b*ArcCosh[c*x])*ArcT
anh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^A
rcCosh[c*x])))/c^3))/(c*d*Sqrt[d - c^2*d*x^2])))/(3*c^2*d)

```

### 3.214.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]

```

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]

```

```

rule 83 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f
*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]

```

rule 109 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Simp[1/(b*(b*e - a*f)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

---

3.214. 
$$\int \frac{x^{5(a+b\operatorname{arccosh}(cx))^2}}{(d-c^2dx^2)^{5/2}} dx$$

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_
.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1)))
Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - S
imp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)]
Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c
*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0]
&& GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_
.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a
+ b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2
*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x]
- Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)
^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*Ar
cCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*
d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.214.4 Maple [A] (verified)

Time = 1.46 (sec) , antiderivative size = 922, normalized size of antiderivative = 1.62

method	result
default	$a^2 \left( -\frac{x^4}{c^2 d(-c^2 d x^2 + d)^{\frac{3}{2}}} + \frac{\frac{4x^2}{c^2 d(-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{8}{3d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}}}{c^2} \right) + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (\sqrt{cx-1} \sqrt{cx+1} cx + c^2 x^2 - 1)}{2d^3 c^6 (c^2 x^2 - 1)} \right)$
parts	$a^2 \left( -\frac{x^4}{c^2 d(-c^2 d x^2 + d)^{\frac{3}{2}}} + \frac{\frac{4x^2}{c^2 d(-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{8}{3d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}}}{c^2} \right) + b^2 \left( -\frac{\sqrt{-d(c^2 x^2 - 1)} (\sqrt{cx-1} \sqrt{cx+1} cx + c^2 x^2 - 1)}{2d^3 c^6 (c^2 x^2 - 1)} \right)$

3.214.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

input `int(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & a^2*(-x^4/c^2/d/(-c^2*d*x^2+d)^{(3/2)}+4/c^2*(x^2/c^2/d/(-c^2*d*x^2+d)^{(3/2)} \\ & -2/3/d/c^4/(-c^2*d*x^2+d)^{(3/2)}))+b^2*(-1/2*(-d*(c^2*x^2-1))^{(1/2)}*((c*x-1) \\ & )^{(1/2)}*(c*x+1)^{(1/2)}*c*x+c^2*x^2-1)*(arccosh(c*x)^2-2*arccosh(c*x)+2)/d^3 \\ & /c^6/(c^2*x^2-1)-1/2*(-d*(c^2*x^2-1))^{(1/2)}*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}* \\ & c*x+c^2*x^2-1)*(arccosh(c*x)^2+2*arccosh(c*x)+2)/d^3/c^6/(c^2*x^2-1)+1/3*( \\ & -d*(c^2*x^2-1))^{(1/2)}*(6*arccosh(c*x)^2*x^2*c^2+(c*x+1)^{(1/2)}*arccosh(c*x) \\ & *(c*x-1)^{(1/2)}*c*x+c^2*x^2-5*arccosh(c*x)^2-1)/(c^2*x^2-1)^2/d^3/c^6+11/3* \\ & (-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/c^6/(c^2*x^2-1)*arc \\ & cosh(c*x)*ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))+11/3*(-d*(c^2*x^2-1))^{(1/2)} \\ & *(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/c^6/(c^2*x^2-1)*polylog(2,-c*x-(c*x-1)^{(1/2)} \\ & *(c*x+1)^{(1/2)})-11/3*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)} \\ & )/d^3/c^6/(c^2*x^2-1)*arccosh(c*x)*ln(1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})-1 \\ & 1/3*(-d*(c^2*x^2-1))^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^3/c^6/(c^2*x^2-1) \\ & *polylog(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))-1/3*a*b*(-d*(c^2*x^2-1))^{(1/2)} \\ & *(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(6*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*arccosh(c*x)* \\ & c^4*x^4-6*c^5*x^5-11*ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*c^4*x^4+11*ln(( \\ & c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)*c^4*x^4-24*(c*x+1)^{(1/2)}*arccosh(c*x)*(c \\ & *x-1)^{(1/2)}*c^2*x^2+11*c^3*x^3+22*ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*x^ \\ & 2*c^2-22*ln((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}+c*x-1)*x^2*c^2+16*arccosh(c*x)*(c* \\ & x-1)^{(1/2)}*(c*x+1)^{(1/2)}-5*c*x-11*ln(1+c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})...$$

### 3.214.5 Fracas [F]

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^5}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-(b^2*x^5*arccosh(c*x)^2 + 2*a*b*x^5*arccosh(c*x) + a^2*x^5)*sqrt(-c^2*d*x^2 + d)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

## 3.214.6 Sympy [F(-1)]

Timed out.

$$\int \frac{x^5(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \text{Timed out}$$

```
input integrate(x**5*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(5/2), x)
```

```
output Timed out
```

## 3.214.7 Maxima [F]

$$\int \frac{x^5(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^5}{(-c^2dx^2 + d)^{5/2}} dx$$

```
input integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2), x, algorithm="maxima")
```

```
output -1/3*a^2*(3*x^4/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 12*x^2/((-c^2*d*x^2 + d)^(3/2)*c^4*d) + 8/((-c^2*d*x^2 + d)^(3/2)*c^6*d)) - 1/3*(3*b^2*c^4*sqrt(d)*x^4 - 12*b^2*c^2*sqrt(d)*x^2 + 8*b^2*sqrt(d))*sqrt(c*x + 1)*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/(c^10*d^3*x^4 - 2*c^8*d^3*x^2 + c^6*d^3) - integrate(2/3*((12*b^2*c^3*x^3 + 3*(a*b*c^5 - b^2*c^5)*x^5 - 8*b^2*c*x)*(c*x + 1)*sqrt(c*x - 1) + (15*b^2*c^4*x^4 + 3*(a*b*c^6 - b^2*c^6)*x^6 - 20*b^2*c^2*x^2 + 8*b^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(c^12*d^(5/2)*x^7 - 3*c^10*d^(5/2)*x^5 + 3*c^8*d^(5/2)*x^3 - c^6*d^(5/2)*x + (c^11*d^(5/2)*x^6 - 3*c^9*d^(5/2)*x^4 + 3*c^7*d^(5/2)*x^2 - c^5*d^(5/2))*sqrt(c*x + 1)*sqrt(c*x - 1)), x)
```



**3.214.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^5*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.214.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x^5*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2),x)`

output `int((x^5*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2), x)`

**3.215** 
$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

3.215.1 Optimal result . . . . . 1897  
 3.215.2 Mathematica [A] (warning: unable to verify) . . . . . 1898  
 3.215.3 Rubi [C] (verified) . . . . . 1899  
 3.215.4 Maple [B] (verified) . . . . . 1907  
 3.215.5 Fracas [F] . . . . . 1908  
 3.215.6 Sympy [F] . . . . . 1909  
 3.215.7 Maxima [F] . . . . . 1909  
 3.215.8 Giac [F] . . . . . 1909  
 3.215.9 Mupad [F(-1)] . . . . . 1910

**3.215.1 Optimal result**

Integrand size = 29, antiderivative size = 482

$$\begin{aligned} \int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx = & -\frac{b^2}{3c^5d^2\sqrt{d-c^2dx^2}} \\ & + \frac{b^2(1-cx)}{3c^5d^2\sqrt{d-c^2dx^2}} + \frac{b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{arccosh}(cx)}{3c^5d^2\sqrt{d-c^2dx^2}} \\ & + \frac{bx^2\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{3c^3d^2(1-c^2x^2)\sqrt{d-c^2dx^2}} + \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\ & - \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^4d^2\sqrt{d-c^2dx^2}} - \frac{4\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^2}{3c^5d^2\sqrt{d-c^2dx^2}} \\ & + \frac{\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^3}{3bc^5d^2\sqrt{d-c^2dx^2}} \\ & + \frac{8b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\log(1-e^{2\operatorname{arccosh}(cx)})}{3c^5d^2\sqrt{d-c^2dx^2}} \\ & + \frac{4b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{3c^5d^2\sqrt{d-c^2dx^2}} \end{aligned}$$

output  $\frac{1}{3}x^3(a+b\operatorname{arccosh}(cx))^2/c^2/d/(-c^2dx^2+d)^{(3/2)}-1/3b^2/c^5/d^2/(-c^2dx^2+d)^{(1/2)}+1/3b^2*(-cx+1)/c^5/d^2/(-c^2dx^2+d)^{(1/2)}-x*(a+b\operatorname{arccosh}(cx))^2/c^4/d^2/(-c^2dx^2+d)^{(1/2)}+1/3b^2*\operatorname{arccosh}(cx)*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/c^5/d^2/(-c^2dx^2+d)^{(1/2)}+1/3b*x^2*(a+b\operatorname{arccosh}(cx))*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/c^3/d^2/(-c^2dx^2+d)^{(1/2)}-4/3*(a+b\operatorname{arccosh}(cx))^2*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/c^5/d^2/(-c^2dx^2+d)^{(1/2)}+1/3*(a+b\operatorname{arccosh}(cx))^3*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/b/c^5/d^2/(-c^2dx^2+d)^{(1/2)}+8/3b*(a+b\operatorname{arccosh}(cx))*\ln(1-(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)}))^2*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/c^5/d^2/(-c^2dx^2+d)^{(1/2)}+4/3b^2*\operatorname{polylog}(2,(cx+(cx-1)^{(1/2)}*(cx+1)^{(1/2)}))^2*(cx-1)^{(1/2)}*(cx+1)^{(1/2)}/c^5/d^2/(-c^2dx^2+d)^{(1/2)}$

### 3.215.2 Mathematica [A] (warning: unable to verify)

Time = 1.96 (sec) , antiderivative size = 382, normalized size of antiderivative = 0.79

$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx = \frac{a^2cx(-3+4c^2x^2)\sqrt{d-c^2dx^2}}{(-1+c^2x^2)^2} - 3a^2\sqrt{d}\arctan\left(\frac{cx\sqrt{d-c^2dx^2}}{\sqrt{d}(-1+c^2x^2)}\right) + \frac{abd\left(-8cx\operatorname{arccosh}(cx)-\sqrt{d-c^2dx^2}\right)}{\sqrt{d}(-1+c^2x^2)}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]`

output  $((a^2cx*(-3 + 4c^2x^2)*\operatorname{Sqrt}[d - c^2dx^2])/(-1 + c^2x^2)^2 - 3a^2*\operatorname{Sqrt}[d]*\operatorname{ArcTan}[(cx*\operatorname{Sqrt}[d - c^2dx^2])/(\operatorname{Sqrt}[d]*(-1 + c^2x^2))] + (a*b*d*(-8cx*\operatorname{ArcCosh}[c*x] - (\operatorname{Sqrt}[(-1 + cx)/(1 + cx)]*(1 + cx) + 2*cx*\operatorname{ArcCosh}[c*x]))/(-1 + c^2x^2) + \operatorname{Sqrt}[(-1 + cx)/(1 + cx)]*(1 + cx)*(3*\operatorname{ArcCosh}[c*x]^2 + 8*\operatorname{Log}[\operatorname{Sqrt}[(-1 + cx)/(1 + cx)]*(1 + cx)])))/\operatorname{Sqrt}[d - c^2dx^2] + (b^2*d*\operatorname{Sqrt}[(-1 + cx)/(1 + cx)]*(1 + cx)*(-((cx*(-1 + c^2x^2 + (-3 + 4c^2x^2)*\operatorname{ArcCosh}[c*x]^2))/(((1 + cx)/(1 + cx))^(3/2)*(1 + cx)^3)) + \operatorname{ArcCosh}[c*x]*((1 - c^2x^2)^(-1) + \operatorname{ArcCosh}[c*x]*(4 + \operatorname{ArcCosh}[c*x]) + 8*\operatorname{Log}[1 - E^(-2*\operatorname{ArcCosh}[c*x])]) - 4*\operatorname{PolyLog}[2, E^(-2*\operatorname{ArcCosh}[c*x])])))/\operatorname{Sqrt}[d - c^2dx^2])/(3*c^5*d^3)$

### 3.215.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 2.71 (sec) , antiderivative size = 461, normalized size of antiderivative = 0.96, number of steps used = 19, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.621$ , Rules used = {6349, 6327, 6349, 25, 100, 27, 87, 43, 6307, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6349} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^3(a+\operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{c^2d} + \frac{x^3(a + \operatorname{barccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^3(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \frac{\int \frac{x^2(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{c^2d} + \frac{x^3(a + \operatorname{barccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{6349} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \int \frac{x^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3cd^2\sqrt{d-c^2dx^2}} \\
 & \quad - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \\
 & \quad \frac{c^2d}{3c^2d(d - c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \int \frac{x^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{x^2(a+\operatorname{barccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3cd^2\sqrt{d-c^2dx^2}} \\
 & \quad - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \\
 & \quad \frac{c^2d}{3c^2d(d - c^2dx^2)^{3/2}}
 \end{aligned}$$

---

3.215.  $\int \frac{x^4(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 100 \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{\int \frac{e^{2x}}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c^3} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right) + \\
 & \frac{3cd^2\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} \\
 & \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 27 \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{\int \frac{x}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right) + \\
 & \frac{3cd^2\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} \\
 & \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 87 \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{c^2d} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right) + \\
 & \frac{3cd^2\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} \\
 & \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 43
 \end{aligned}$$

3.215.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} - \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{c^2d} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} + \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right) \\
 & \frac{3cd^2\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} \\
 & \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 6307 \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right) \\
 & \frac{3cd^2\sqrt{d-c^2dx^2}}{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+b\operatorname{arccosh}(cx))}{(1-cx)(cx+1)} dx} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{3bc^3d\sqrt{d-c^2dx^2}} + \\
 & \frac{c^2d}{x^3(a+b\operatorname{arccosh}(cx))^2} \\
 & \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 6327 \\
 & 2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx)}{c^2} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right) \\
 & \frac{3cd^2\sqrt{d-c^2dx^2}}{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{3bc^3d\sqrt{d-c^2dx^2}} + \\
 & \frac{c^2d}{x^3(a+b\operatorname{arccosh}(cx))^2} \\
 & \frac{3c^2d(d-c^2dx^2)^{3/2}}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 6328
 \end{aligned}$$

3.215.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
 & - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+b\operatorname{arccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{c^3 d \sqrt{d-c^2 dx^2}} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{3bc^3 d \sqrt{d-c^2 dx^2}} \\
 & + \frac{2b\sqrt{cx-1}\sqrt{cx+1}}{c^2 d} \left( \frac{\int \frac{cx(a+b\operatorname{arccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{c^4} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}}}{c} - \frac{1}{c^3 \sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right)
 \end{aligned}$$

$$\frac{3cd^2 \sqrt{d-c^2 dx^2}}{3c^2 d (d-c^2 dx^2)^{3/2}} \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2 d (d-c^2 dx^2)^{3/2}}$$

↓ 3042

$$\begin{aligned}
 & - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -i(a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{c^3 d \sqrt{d-c^2 dx^2}} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{3bc^3 d \sqrt{d-c^2 dx^2}} \\
 & + \frac{2b\sqrt{cx-1}\sqrt{cx+1}}{c^2 d} \left( \frac{\int -i(a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{c^4} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}}}{c} - \frac{1}{c^3 \sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right)
 \end{aligned}$$

$$\frac{3cd^2 \sqrt{d-c^2 dx^2}}{3c^2 d (d-c^2 dx^2)^{3/2}} \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2 d (d-c^2 dx^2)^{3/2}}$$

↓ 26

$$\begin{aligned}
 & - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{c^3 d \sqrt{d-c^2 dx^2}} + \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^3}{3bc^3 d \sqrt{d-c^2 dx^2}} \\
 & + \frac{2b\sqrt{cx-1}\sqrt{cx+1}}{c^2 d} \left( - \frac{i \int (a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{c^4} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2 x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c^2 \sqrt{cx+1}}}{c} - \frac{1}{c^3 \sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} \right)
 \end{aligned}$$

$$\frac{3cd^2 \sqrt{d-c^2 dx^2}}{3c^2 d (d-c^2 dx^2)^{3/2}} \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2 d (d-c^2 dx^2)^{3/2}}$$

↓ 4199

---

3.215.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2 dx^2)^{5/2}} dx$

$$\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\int\frac{e^{2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}}d\operatorname{arccosh}(cx)-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}\right)}{c^3d\sqrt{d-c^2dx^2}}+\frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}}-\frac{\sqrt{cx-1}\sqrt{cx+1}}{3bc^3}$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1}\left(-\frac{i\left(2i\int\frac{e^{2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}}d\operatorname{arccosh}(cx)-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}\right)}{c^4}+\frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)}+\frac{b}{c}\right)$$


---


$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}\quad \frac{3cd^2\sqrt{d-c^2dx^2}}{c^2d}$$

↓ 25

$$\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\int\frac{e^{2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}}d\operatorname{arccosh}(cx)-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}\right)}{c^3d\sqrt{d-c^2dx^2}}+\frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}}-\frac{\sqrt{cx-1}\sqrt{cx+1}}{3bc^3}$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1}\left(-\frac{i\left(-2i\int\frac{e^{2\operatorname{arccosh}(cx)}(a+b\operatorname{arccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}}d\operatorname{arccosh}(cx)-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}\right)}{c^4}+\frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)}+\frac{b}{c}\right)$$


---


$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}\quad \frac{3cd^2\sqrt{d-c^2dx^2}}{c^2d}$$

↓ 2620

$$\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(\frac{1}{2}b\int\log(1-e^{2\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}\right)\right)}{c^3d\sqrt{d-c^2dx^2}}+\frac{b}{c}$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1}\left(-\frac{i\left(-2i\left(\frac{1}{2}b\int\log(1-e^{2\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}\right)\right)}{c^4}+\frac{b}{c}\right)$$


---


$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}\quad \frac{3cd^2\sqrt{d-c^2dx^2}}{c^2d}$$

↓ 2715

3.215.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$



$$\begin{aligned}
 & \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{4} b \int e^{-2\operatorname{arccosh}(cx)} \log(1-e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) \right) - \frac{i(a+b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^3 d \sqrt{d-c^2 dx^2}} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( - \frac{i \left( -2i \left( \frac{1}{4} b \int e^{-2\operatorname{arccosh}(cx)} \log(1-e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) \right) - \frac{i(a+b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} \right)}{c^2 d} \\
 & \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2 d (d-c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{2838} \\
 & \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3c^2 d (d-c^2 dx^2)^{3/2}} - \frac{3cd^2 \sqrt{d-c^2 dx^2}}{3cd^2 \sqrt{d-c^2 dx^2}} \\
 & \frac{x(a+b\operatorname{arccosh}(cx))^2}{c^2 d \sqrt{d-c^2 dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right) - \frac{i(a+b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^3 d \sqrt{d-c^2 dx^2}} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( - \frac{i \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right) - \frac{i(a+b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} \right)}{c^2 d} + \dots
 \end{aligned}$$

```
input Int[(x^4*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]
```

```
output (x^3*(a + b*ArcCosh[c*x])^2)/(3*c^2*d*(d - c^2*d*x^2)^(3/2)) + (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((x^2*(a + b*ArcCosh[c*x]))/(2*c^2*(1 - c^2*x^2)) + (b*(-1/(c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (-Sqrt[-1 + c*x]/(c^2*Sqrt[1 + c*x])) + ArcCosh[c*x]/c^2)/c)/(2*c) - (I*(((1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/c^4)/(3*c*d^2*Sqrt[d - c^2*d*x^2]) - ((x*(a + b*ArcCosh[c*x])^2)/(c^2*d*Sqrt[d - c^2*d*x^2]) - (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^3)/(3*b*c^3*d*Sqrt[d - c^2*d*x^2]) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(((1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/c^3*d*Sqrt[d - c^2*d*x^2])/(c^2*d)
```

3.215.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2 dx^2)^{5/2}} dx$

## 3.215.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))`
- rule 100 `Int[((a_) + (b_)*(x_))^(2*((c_) + (d_)*(x_))^(n_))*((e_) + (f_)*(x_))^(p_), x_] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d^2*(d*e - c*f)*(n + 1))), x] - Simp[1/(d^2*(d*e - c*f)*(n + 1)) Int[(c + d*x)^(n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n + 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))`
- rule 2620 `Int[((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_
.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp
[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x
)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && In
tegerQ[4*k] && IGtQ[m, 0]`

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_
Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d
+ e*x^2])*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x
] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (
e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] :> Int[(f*x)^m*(d1
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2),
x_Symbol] :> Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]
], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.215.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2900 vs.  $2(446) = 892$ .

Time = 1.35 (sec) , antiderivative size = 2901, normalized size of antiderivative = 6.02

method	result	size
default	Expression too large to display	2901
parts	Expression too large to display	2901

```
input int(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)
```

```

output 4/3*b^2*(-d*(c^2*x^2-1))^(1/2)/(24*c^8*x^8-87*c^6*x^6+118*c^4*x^4-71*c^2*x
^2+16)/d^3*(c*x-1)*(c*x+1)*x^5+32*b^2*(-d*(c^2*x^2-1))^(1/2)/(24*c^8*x^8-8
7*c^6*x^6+118*c^4*x^4-71*c^2*x^2+16)/d^3*c^2*arccosh(c*x)^2*x^7+16/3*b^2*(
-d*(c^2*x^2-1))^(1/2)/(24*c^8*x^8-87*c^6*x^6+118*c^4*x^4-71*c^2*x^2+16)/d^
3*c^2*arccosh(c*x)*x^7+181/3*b^2*(-d*(c^2*x^2-1))^(1/2)/(24*c^8*x^8-87*c^6
*x^6+118*c^4*x^4-71*c^2*x^2+16)/d^3/c^2*arccosh(c*x)^2*x^3+40/3*b^2*(-d*(c
^2*x^2-1))^(1/2)/(24*c^8*x^8-87*c^6*x^6+118*c^4*x^4-71*c^2*x^2+16)/d^3/c^2
*arccosh(c*x)*x^3-16*b^2*(-d*(c^2*x^2-1))^(1/2)/(24*c^8*x^8-87*c^6*x^6+118
*c^4*x^4-71*c^2*x^2+16)/d^3/c^4*arccosh(c*x)^2*x-4*b^2*(-d*(c^2*x^2-1))^(1
/2)/(24*c^8*x^8-87*c^6*x^6+118*c^4*x^4-71*c^2*x^2+16)/d^3/c^4*arccosh(c*x)
*x+16/3*b^2*(-d*(c^2*x^2-1))^(1/2)/(24*c^8*x^8-87*c^6*x^6+118*c^4*x^4-71*c
^2*x^2+16)/d^3/c^5*(c*x-1)^(1/2)*(c*x+1)^(1/2)-a^2/c^4/d^2*x/(-c^2*d*x^2+d
)^(1/2)+a^2/c^4/d^2/(c^2*d)^(1/2)*arctan((c^2*d)^(1/2)*x/(-c^2*d*x^2+d)^(1
/2))+8/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^5/(c
^2*x^2-1)*arccosh(c*x)^2-8/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x
+1)^(1/2)/d^3/c^5/(c^2*x^2-1)*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-8
/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^5/(c^2*x^2
-1)*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-16/3*b^2*(-d*(c^2*x^2-1))^(
1/2)/(24*c^8*x^8-87*c^6*x^6+118*c^4*x^4-71*c^2*x^2+16)/d^3*(c*x-1)*(c*x+1
)*arccosh(c*x)*x^5-55/3*b^2*(-d*(c^2*x^2-1))^(1/2)/(24*c^8*x^8-87*c^6*x...

```

### 3.215.5 Fracas [F]

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^4}{(-c^2 dx^2 + d)^{5/2}} dx$$

```

input integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fric
as")

```

```

output integral(-(b^2*x^4*arccosh(c*x)^2 + 2*a*b*x^4*arccosh(c*x) + a^2*x^4)*sqrt
(-c^2*d*x^2 + d)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)

```

---

3.215.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

## 3.215.6 Sympy [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{\frac{5}{2}}} dx$$

input `integrate(x**4*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(5/2), x)`

output `Integral(x**4*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**5/2, x)`

## 3.215.7 Maxima [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^4}{(-c^2dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2), x, algorithm="maxima")`

output `1/3*(x*(3*x^2/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 2/((-c^2*d*x^2 + d)^(3/2)*c^4*d)) - x/(sqrt(-c^2*d*x^2 + d)*c^4*d^2) + 3*arcsin(c*x)/(c^5*d^(5/2)))*a^2 + integrate(b^2*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/(-c^2*d*x^2 + d)^(5/2) + 2*a*b*x^4*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(-c^2*d*x^2 + d)^(5/2), x)`

## 3.215.8 Giac [F]

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^4}{(-c^2dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x^4/(-c^2*d*x^2 + d)^(5/2), x)`

---

3.215.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx$

**3.215.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4 (a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^4 (a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x^4*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2),x)`output `int((x^4*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2), x)`

**3.216** 
$$\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

3.216.1 Optimal result . . . . .1911  
 3.216.2 Mathematica [A] (warning: unable to verify) . . . . .1912  
 3.216.3 Rubi [C] (verified) . . . . .1912  
 3.216.4 Maple [A] (verified) . . . . .1919  
 3.216.5 Fricas [F] . . . . .1920  
 3.216.6 Sympy [F] . . . . .1920  
 3.216.7 Maxima [F] . . . . .1920  
 3.216.8 Giac [F(-2)] . . . . .1921  
 3.216.9 Mupad [F(-1)] . . . . .1921

**3.216.1 Optimal result**

Integrand size = 29, antiderivative size = 336

$$\begin{aligned} \int \frac{x^3(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx &= -\frac{b^2}{3c^4d^2\sqrt{d - c^2dx^2}} \\ &+ \frac{bx\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{3c^3d^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} \\ &+ \frac{x^2(a + \operatorname{arccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}} - \frac{2(a + \operatorname{arccosh}(cx))^2}{3c^4d^2\sqrt{d - c^2dx^2}} \\ &- \frac{10b\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{3c^4d^2\sqrt{d - c^2dx^2}} \\ &- \frac{5b^2\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{3c^4d^2\sqrt{d - c^2dx^2}} \\ &+ \frac{5b^2\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{3c^4d^2\sqrt{d - c^2dx^2}} \end{aligned}$$

output 
$$\begin{aligned} &1/3*x^2*(a+b*\operatorname{arccosh}(c*x))^2/c^2/d/(-c^2*d*x^2+d)^{(3/2)}-1/3*b^2/c^4/d^2/(- \\ &c^2*d*x^2+d)^{(1/2)}-2/3*(a+b*\operatorname{arccosh}(c*x))^2/c^4/d^2/(-c^2*d*x^2+d)^{(1/2)}+1 \\ &/3*b*x*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/d^2/(-c^2*x^2+1) \\ &/(-c^2*d*x^2+d)^{(1/2)}-10/3*b*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}* \\ &(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^4/d^2/(-c^2*d*x^2+d)^{(1/2)}-5/ \\ &3*b^2*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1 \\ &/2)}/c^4/d^2/(-c^2*d*x^2+d)^{(1/2)}+5/3*b^2*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+ \\ &1)^{(1/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^4/d^2/(-c^2*d*x^2+d)^{(1/2)} \end{aligned}$$

3.216. 
$$\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$



**3.216.2 Mathematica [A] (warning: unable to verify)**

Time = 3.68 (sec) , antiderivative size = 356, normalized size of antiderivative = 1.06

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \frac{4a^2(-2 + 3c^2x^2) - ab(\operatorname{arccosh}(cx)(4 - 12\cosh(2\operatorname{arccosh}(cx))) - 2\sinh(2\operatorname{arccosh}(cx)))}{(d - c^2dx^2)^{5/2}}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]`

output

```
(4*a^2*(-2 + 3*c^2*x^2) - a*b*(ArcCosh[c*x]*(4 - 12*Cosh[2*ArcCosh[c*x]])
- 2*Sinh[2*ArcCosh[c*x]] + 5*(Log[Cosh[ArcCosh[c*x]/2]] - Log[Sinh[ArcCosh
[c*x]/2]))*(3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) - Sinh[3*ArcCosh[c*x]])
) - b^2*(2 + 2*ArcCosh[c*x]^2 - 2*(1 + 3*ArcCosh[c*x]^2)*Cosh[2*ArcCosh[c*
x]] - 15*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*(Log[1 - E^(-Ar
cCosh[c*x])] - Log[1 + E^(-ArcCosh[c*x])]) + 20*((-1 + c*x)/(1 + c*x))^(3/
2)*(1 + c*x)^3*PolyLog[2, -E^(-ArcCosh[c*x])] - 20*((-1 + c*x)/(1 + c*x))^(
3/2)*(1 + c*x)^3*PolyLog[2, E^(-ArcCosh[c*x])] - 2*ArcCosh[c*x]*Sinh[2*Ar
cCosh[c*x]] + 5*ArcCosh[c*x]*Log[1 - E^(-ArcCosh[c*x])]*Sinh[3*ArcCosh[c*x
]] - 5*ArcCosh[c*x]*Log[1 + E^(-ArcCosh[c*x])]*Sinh[3*ArcCosh[c*x]])))/(12*
c^4*d*(d - c^2*d*x^2)^(3/2))
```

**3.216.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.67 (sec) , antiderivative size = 330, normalized size of antiderivative = 0.98, number of steps used = 20, number of rules used = 19,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.655$ , Rules used = {6349, 6327, 6329, 25, 6304, 6318, 3042, 26, 4670, 2715, 2838, 6349, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx$$

↓ 6349

$$\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a + \operatorname{arccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3cd^2\sqrt{d - c^2dx^2}} - \frac{2 \int \frac{x(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx}{3c^2d} + \frac{x^2(a + \operatorname{arccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}}$$

---

3.216.  $\int \frac{x^3(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
& \downarrow 6327 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \frac{2 \int \frac{x(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{3c^2d} + \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 6329 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \\
& \frac{2 \left( \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} \right)}{3c^2d} + \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 25 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \\
& \frac{2 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{(1-cx)(cx+1)} dx}{cd\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right)}{3c^2d} + \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 6304 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \\
& \frac{2 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx}{cd\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} \right)}{3c^2d} + \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 6318 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \\
& \frac{2 \left( \frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{c^2d\sqrt{d-c^2dx^2}} \right)}{3c^2d} + \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 3042
\end{aligned}$$

---

3.216.  $\int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \\
 & \frac{2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)}{c^2d\sqrt{d-c^2dx^2}}\right)}{3c^2d} + \\
 & \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{26} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \\
 & \frac{2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)}{c^2d\sqrt{d-c^2dx^2}}\right)}{3c^2d} + \\
 & \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{4670} \\
 & \frac{2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left(ib \int \log(1-e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx) + 2i\operatorname{arctan}\right)}{c^2d\sqrt{d-c^2dx^2}}\right)}{3c^2d} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} + \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{2715} \\
 & \frac{2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left(ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)})de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)})\right)}{c^2d\sqrt{d-c^2dx^2}}\right)}{3c^2d} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} + \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{2838} \\
 & \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^2(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} - \\
 & \frac{2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})\right)}{c^2d\sqrt{d-c^2dx^2}}\right)}{3c^2d} \\
 & \frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}
 \end{aligned}$$

---

3.216.  $\int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{array}{c}
\downarrow 6349 \\
\frac{2b\sqrt{cx-1}\sqrt{cx+1}\left(-\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{2c^2} + \frac{b\int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{x(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)}\right)}{3cd^2\sqrt{d-c^2dx^2}} \\
\frac{2\left(\frac{(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}\left(2,-e^{\operatorname{arccosh}(cx)}\right)-ib\operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)\right)}{c^2d\sqrt{d-c^2dx^2}}\right)}{c^2d\sqrt{d-c^2dx^2}} \\
\frac{x^2(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
\downarrow 83 \\
\frac{2b\sqrt{cx-1}\sqrt{cx+1}\left(-\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{2c^2} + \frac{x(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}}\right)}{3cd^2\sqrt{d-c^2dx^2}} \\
\frac{2\left(\frac{(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}\left(2,-e^{\operatorname{arccosh}(cx)}\right)-ib\operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)\right)}{c^2d\sqrt{d-c^2dx^2}}\right)}{c^2d\sqrt{d-c^2dx^2}} \\
\frac{x^2(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
\downarrow 6318 \\
\frac{2b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{arccosh}(cx)}{2c^3} + \frac{x(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}}\right)}{3cd^2\sqrt{d-c^2dx^2}} \\
\frac{2\left(\frac{(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}\left(2,-e^{\operatorname{arccosh}(cx)}\right)-ib\operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)\right)}{c^2d\sqrt{d-c^2dx^2}}\right)}{c^2d\sqrt{d-c^2dx^2}} \\
\frac{x^2(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
\downarrow 3042 \\
\frac{2b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{\int i(a+b\operatorname{arccosh}(cx))\operatorname{csc}(i\operatorname{arccosh}(cx))\operatorname{arccosh}(cx)}{2c^3} + \frac{x(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}}\right)}{3cd^2\sqrt{d-c^2dx^2}} \\
\frac{2\left(\frac{(a+b\operatorname{arccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}\left(2,-e^{\operatorname{arccosh}(cx)}\right)-ib\operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)\right)}{c^2d\sqrt{d-c^2dx^2}}\right)}{c^2d\sqrt{d-c^2dx^2}} \\
\frac{x^2(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}
\end{array}$$

---

3.216.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

↓ 26

$$\frac{2b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{i\int(a+\operatorname{barccosh}(cx))\operatorname{csc}(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)}{2c^3} + \frac{x(a+\operatorname{barccosh}(cx))}{2c^2(1-c^2x^2)} - \frac{b}{2c^3\sqrt{cx-1}\sqrt{cx+1}}\right)}{3cd^2\sqrt{d-c^2dx^2}}$$


---


$$2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{c^2d\sqrt{d-c^2dx^2}}\right)$$


---


$$\frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}$$

↓ 4670

$$2b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{i\left(ib\int\log(1-e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)-ib\int\log(1+e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)+2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx))}{2c^3}\right)$$


---


$$2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{c^2d\sqrt{d-c^2dx^2}}\right)$$


---


$$\frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}$$

↓ 2715

$$2b\sqrt{cx-1}\sqrt{cx+1}\left(\frac{i\left(ib\int e^{-\operatorname{arccosh}(cx)}\log(1-e^{\operatorname{arccosh}(cx)})de^{\operatorname{arccosh}(cx)}-ib\int e^{-\operatorname{arccosh}(cx)}\log(1+e^{\operatorname{arccosh}(cx)})de^{\operatorname{arccosh}(cx)}+2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx))}{2c^3}\right)$$


---


$$2\left(\frac{(a+\operatorname{barccosh}(cx))^2}{c^2d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})\right)(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{c^2d\sqrt{d-c^2dx^2}}\right)$$


---


$$\frac{x^2(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}}$$

↓ 2838

---

3.216.  $\int \frac{x^3(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$2 \left( \frac{(a + \operatorname{barccosh}(cx))^2}{c^2 d \sqrt{d - c^2 dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{c^2 d \sqrt{d - c^2 dx^2}} \right)$$


---


$$2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{i \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c^3} + \frac{3c^2 d}{c^2 d \sqrt{d - c^2 dx^2}} \right)$$


---


$$\frac{x^2 (a + \operatorname{barccosh}(cx))^2}{3c^2 d (d - c^2 dx^2)^{3/2}}$$

input `Int[(x^3*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]`

output `(x^2*(a + b*ArcCosh[c*x])^2)/(3*c^2*d*(d - c^2*d*x^2)^(3/2)) + (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/2*b/(c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*c^2*(1 - c^2*x^2)) + ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c^3))/(3*c*d^2*Sqrt[d - c^2*d*x^2]) - (2*((a + b*ArcCosh[c*x])^2/(c^2*d*Sqrt[d - c^2*d*x^2]) - ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/(c^2*d*Sqrt[d - c^2*d*x^2])))/(3*c^2*d)`

### 3.216.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

---

3.216.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

```
rule 6349 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*((m - 1)/(2*e*(p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]
```

### 3.216.4 Maple [A] (verified)

Time = 1.00 (sec) , antiderivative size = 635, normalized size of antiderivative = 1.89

method	result
default	$a^2 \left( \frac{x^2}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{2}{3 d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}} \right) + b^2 \left( \frac{\sqrt{-d(c^2 x^2 - 1)} (3 \operatorname{arccosh}(cx)^2 x^2 c^2 + \sqrt{cx+1} \operatorname{arccosh}(cx) \sqrt{cx-1} cx + c^2)}{3(c^2 x^2 - 1)^2 d^3 c^4} \right)$
parts	$a^2 \left( \frac{x^2}{c^2 d (-c^2 d x^2 + d)^{\frac{3}{2}}} - \frac{2}{3 d c^4 (-c^2 d x^2 + d)^{\frac{3}{2}}} \right) + b^2 \left( \frac{\sqrt{-d(c^2 x^2 - 1)} (3 \operatorname{arccosh}(cx)^2 x^2 c^2 + \sqrt{cx+1} \operatorname{arccosh}(cx) \sqrt{cx-1} cx + c^2)}{3(c^2 x^2 - 1)^2 d^3 c^4} \right)$

```
input int(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)
```

```
output a^2*(x^2/c^2/d/(-c^2*d*x^2+d)^(3/2)-2/3/d/c^4/(-c^2*d*x^2+d)^(3/2))+b^2*(1/3*(-d*(c^2*x^2-1))^(1/2)*(3*arccosh(c*x)^2*x^2*c^2+(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c*x+c^2*x^2-2*arccosh(c*x)^2-1)/(c^2*x^2-1)^2/d^3/c^4+5/3*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^4/(c^2*x^2-1)*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+5/3*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^4/(c^2*x^2-1)*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))-5/3*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^4/(c^2*x^2-1)*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+2*a*b*(1/6*(-d*(c^2*x^2-1))^(1/2)*(6*c^2*x^2*arccosh(c*x)+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x-4*arccosh(c*x)))/(c^2*x^2-1)^2/d^3/c^4-5/6*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^4/(c^2*x^2-1)*ln((c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x-1)+5/6*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^4/(c^2*x^2-1)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))
```



**3.216.5 Fricas [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^3}{(-c^2dx^2 + d)^{5/2}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-(b^2*x^3*arccosh(c*x)^2 + 2*a*b*x^3*arccosh(c*x) + a^2*x^3)*sqrt(-c^2*d*x^2 + d)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

**3.216.6 Sympy [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{5/2}} dx$$

input `integrate(x**3*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral(x**3*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**5/2, x)`

**3.216.7 Maxima [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^3}{(-c^2dx^2 + d)^{5/2}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/6*a*b*c*(2*sqrt(-d)*x/(c^6*d^3*x^2 - c^4*d^3) + 5*sqrt(-d)*log(c*x + 1)/(c^5*d^3) - 5*sqrt(-d)*log(c*x - 1)/(c^5*d^3)) + 2/3*a*b*(3*x^2/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 2/((-c^2*d*x^2 + d)^(3/2)*c^4*d))*arccosh(c*x) + 1/3*a^2*(3*x^2/((-c^2*d*x^2 + d)^(3/2)*c^2*d) - 2/((-c^2*d*x^2 + d)^(3/2)*c^4*d)) + b^2*integrate(x^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/(-c^2*d*x^2 + d)^(5/2), x)`

---

3.216.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx$

**3.216.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.216.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x^3*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2),x)`

output `int((x^3*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2), x)`

**3.217** 
$$\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

3.217.1 Optimal result . . . . . 1922  
 3.217.2 Mathematica [A] (warning: unable to verify) . . . . . 1923  
 3.217.3 Rubi [C] (verified) . . . . . 1923  
 3.217.4 Maple [B] (verified) . . . . . 1929  
 3.217.5 Fricas [F] . . . . . 1930  
 3.217.6 Sympy [F] . . . . . 1931  
 3.217.7 Maxima [F] . . . . . 1931  
 3.217.8 Giac [F] . . . . . 1931  
 3.217.9 Mupad [F(-1)] . . . . . 1932

**3.217.1 Optimal result**

Integrand size = 29, antiderivative size = 389

$$\begin{aligned} \int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx &= -\frac{b^2}{3c^3d^2\sqrt{d-c^2dx^2}} + \frac{b^2(1-cx)}{3c^3d^2\sqrt{d-c^2dx^2}} \\ &+ \frac{b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{arccosh}(cx)}{3c^3d^2\sqrt{d-c^2dx^2}} + \frac{bx^2\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{3cd^2(1-c^2x^2)\sqrt{d-c^2dx^2}} \\ &+ \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} - \frac{\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^2}{3c^3d^2\sqrt{d-c^2dx^2}} \\ &+ \frac{2b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\log(1-e^{2\operatorname{arccosh}(cx)})}{3c^3d^2\sqrt{d-c^2dx^2}} \\ &+ \frac{b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,e^{2\operatorname{arccosh}(cx)})}{3c^3d^2\sqrt{d-c^2dx^2}} \end{aligned}$$

```
output 1/3*x^3*(a+b*arccosh(c*x))^2/d/(-c^2*d*x^2+d)^(3/2)-1/3*b^2/c^3/d^2/(-c^2*d*x^2+d)^(1/2)+1/3*b^2*(-c*x+1)/c^3/d^2/(-c^2*d*x^2+d)^(1/2)+1/3*b^2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d^2/(-c^2*d*x^2+d)^(1/2)+1/3*b*x^2*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d^2/(-c^2*x^2+1)/(-c^2*d*x^2+d)^(1/2)-1/3*(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d^2/(-c^2*d*x^2+d)^(1/2)+2/3*b*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d^2/(-c^2*d*x^2+d)^(1/2)+1/3*b^2*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/d^2/(-c^2*d*x^2+d)^(1/2)
```

**3.217.2 Mathematica [A] (warning: unable to verify)**

Time = 1.84 (sec) , antiderivative size = 264, normalized size of antiderivative = 0.68

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \frac{\frac{a^2 c^3 x^3}{1 - c^2 x^2} + ab \left( \frac{2c^3 x^3 \operatorname{arccosh}(cx)}{1 - c^2 x^2} + \frac{\sqrt{\frac{-1+cx}{1+cx}} (-1+2(-1+c^2 x^2) \log(\sqrt{\frac{-1+cx}{1+cx}}(1+cx)))}{-1+cx} \right)}{d - c^2 dx^2} + b^2$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2),x]`

output

```
((a^2*c^3*x^3)/(1 - c^2*x^2) + a*b*((2*c^3*x^3*ArcCosh[c*x])/(1 - c^2*x^2)
+ (Sqrt[(-1 + c*x)/(1 + c*x)]*(-1 + 2*(-1 + c^2*x^2)*Log[Sqrt[(-1 + c*x)/(
1 + c*x)]*(1 + c*x)]))/(-1 + c*x)) + b^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 +
c*x)*(-((c*x*(-1 + c^2*x^2 + c^2*x^2*ArcCosh[c*x]^2))/(((1 + c*x)/(1 + c*
x))^(3/2)*(1 + c*x)^3)) + ArcCosh[c*x]*((1 - c^2*x^2)^(-1) + ArcCosh[c*x]
+ 2*Log[1 - E^(-2*ArcCosh[c*x])]) - PolyLog[2, E^(-2*ArcCosh[c*x])]))/(3*c
^3*d^2*Sqrt[d - c^2*d*x^2])
```

**3.217.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 1.29 (sec) , antiderivative size = 249, normalized size of antiderivative = 0.64, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.517$ , Rules used = {6332, 6327, 6349, 100, 27, 87, 43, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

↓ 6332

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^3(a + \operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 6327

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x^3(a + \operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{x^3(a + \operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

---

3.217.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \downarrow 6349 \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \int \frac{x^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2} \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}} + \\
& \downarrow 100 \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{\int \frac{c^2x}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c^3} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2} \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}} + \\
& \downarrow 27 \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{\int \frac{x}{\sqrt{cx-1}(cx+1)^{3/2}} dx}{c} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2} \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}} + \\
& \downarrow 87 \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx))}{1-c^2x^2} dx}{c^2} + \frac{b \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}} \right)}{2c} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2} \frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}} + \\
& \downarrow 43
\end{aligned}$$

---

3.217.  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{x(a+b\operatorname{arccosh}(cx)) dx}{1-c^2x^2}}{c^2} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}}}{c} \right)}{2c} \right)$$

---


$$\frac{3d^2\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} - \frac{3d(d-c^2dx^2)^{3/2}}{3d(d-c^2dx^2)^{3/2}}$$

↓ 6328

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int \frac{cx(a+b\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)}}{c^4} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}}}{c} \right)}{2c} \right)$$

---


$$\frac{3d^2\sqrt{d-c^2dx^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} - \frac{3d(d-c^2dx^2)^{3/2}}{3d(d-c^2dx^2)^{3/2}}$$

↓ 3042

$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{\int -i(a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{c^4} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}}}{c} \right)}{2c} \right)$$

---


$$3d^2\sqrt{d-c^2dx^2}$$

↓ 26

$$\frac{x^3(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \int (a+b\operatorname{arccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{c^4} + \frac{x^2(a+b\operatorname{arccosh}(cx))}{2c^2(1-c^2x^2)} + \frac{b \left( \frac{\operatorname{arccosh}(cx) - \frac{\sqrt{cx-1}}{c^2\sqrt{cx+1}} - \frac{1}{c^3\sqrt{cx-1}\sqrt{cx+1}}}{c} \right)}{2c} \right)$$

---


$$3d^2\sqrt{d-c^2dx^2}$$

↓ 4199

---

3.217.  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\frac{x^3(a + \operatorname{barccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} + 2bc\sqrt{cx - 1}\sqrt{cx + 1} \left( -\frac{i \left( 2i \int -\frac{e^{2\operatorname{arccosh}(cx)}(a + b\operatorname{arccosh}(cx))}{1 - e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a + b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} + \frac{x^2(a + b\operatorname{arccosh}(cx))}{2c^2(1 - c^2x^2)} + \frac{b}{c} \right)$$

---


$$3d^2\sqrt{d - c^2dx^2}$$

↓ 25

$$\frac{x^3(a + \operatorname{barccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} + 2bc\sqrt{cx - 1}\sqrt{cx + 1} \left( -\frac{i \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a + b\operatorname{arccosh}(cx))}{1 - e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a + b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} + \frac{x^2(a + b\operatorname{arccosh}(cx))}{2c^2(1 - c^2x^2)} + \frac{b}{c} \right)$$

---


$$3d^2\sqrt{d - c^2dx^2}$$

↓ 2620

$$\frac{x^3(a + \operatorname{barccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} + 2bc\sqrt{cx - 1}\sqrt{cx + 1} \left( -\frac{i \left( -2i \left( \frac{1}{2} b \int \log(1 - e^{2\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - \frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) (a + b\operatorname{arccosh}(cx)) \right) - \frac{i(a + b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} + \frac{x^2(a + b\operatorname{arccosh}(cx))}{2c^2(1 - c^2x^2)} + \frac{b}{c} \right)$$

---


$$3d^2\sqrt{d - c^2dx^2}$$

↓ 2715

$$\frac{x^3(a + \operatorname{barccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} + 2bc\sqrt{cx - 1}\sqrt{cx + 1} \left( -\frac{i \left( -2i \left( \frac{1}{4} b \int e^{-2\operatorname{arccosh}(cx)} \log(1 - e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log(1 - e^{2\operatorname{arccosh}(cx)}) (a + b\operatorname{arccosh}(cx)) \right) - \frac{i(a + b\operatorname{arccosh}(cx))^2}{2b} \right)}{c^4} + \frac{x^2(a + b\operatorname{arccosh}(cx))}{2c^2(1 - c^2x^2)} + \frac{b}{c} \right)$$

---


$$3d^2\sqrt{d - c^2dx^2}$$

↓ 2838

---

3.217.  $\int \frac{x^2(a + b\operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx$

$$\frac{x^3(a + \operatorname{arccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1}}{3d^2\sqrt{d-c^2dx^2}} \left( -\frac{i\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)(a+\operatorname{arccosh}(cx))-\frac{1}{4}b\operatorname{PolyLog}(2,e^{2\operatorname{arccosh}(cx)})\right)-\frac{i(a+b\operatorname{arccosh}(cx))^2}{2b}}{c^4} \right) +$$

input `Int[(x^2*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2),x]`

output `(x^3*(a + b*ArcCosh[c*x])^2)/(3*d*(d - c^2*d*x^2)^(3/2)) + (2*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((x^2*(a + b*ArcCosh[c*x]))/(2*c^2*(1 - c^2*x^2)) + (b*(-1/(c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (-Sqrt[-1 + c*x]/(c^2*Sqrt[1 + c*x])) + ArcCosh[c*x]/c^2)/c))/(2*c) - (I*((( -1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/c^4)/(3*d^2*Sqrt[d - c^2*d*x^2])`

### 3.217.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`



- rule 87 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(-*(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))`
- rule 100 `Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d^2*(d*e - c*f)*(n + 1))), x] - Simp[1/(d^2*(d*e - c*f)*(n + 1)) Int[(c + d*x)^(n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n + 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))`
- rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4199 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]`

rule 6349 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] + (-Simp[f^2*(m - 1)/(2*e*(p + 1)) Int[(f*x)^(m - 2)*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && IGtQ[m, 1]`

### 3.217.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2656 vs. 2(363) = 726.

Time = 1.30 (sec) , antiderivative size = 2657, normalized size of antiderivative = 6.83

method	result	size
default	Expression too large to display	2657
parts	Expression too large to display	2657

input `int(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

$$3.217. \quad \int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

output 
$$-2/3*b^2*(-d*(c^2*x^2-1))^{1/2}*(c*x-1)^{1/2}*(c*x+1)^{1/2}/d^3/c^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)*\ln(1+c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2})-2/3*b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3*c^2*\operatorname{arccosh}(c*x)*x^5+1/3*b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3/c^3*(c*x-1)^{1/2}*(c*x+1)^{1/2}+1/3*b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3*x^3-2/3*b^2*(-d*(c^2*x^2-1))^{1/2}*(c*x-1)^{1/2}*(c*x+1)^{1/2}/d^3/c^3/(c^2*x^2-1)*\operatorname{polylog}(2,-c*x-(c*x-1)^{1/2}*(c*x+1)^{1/2})+1/3*b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3*\operatorname{arccosh}(c*x)*x^3+1/3*b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3*\operatorname{arccosh}(c*x)*(c*x-1)*(c*x+1)*x^3-2/3*b^2*(-d*(c^2*x^2-1))^{1/2}*(c*x-1)^{1/2}*(c*x+1)^{1/2}/d^3/c^3/(c^2*x^2-1)*\operatorname{polylog}(2,c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2})+2/3*b^2*(-d*(c^2*x^2-1))^{1/2}*(c*x-1)^{1/2}*(c*x+1)^{1/2}/d^3/c^3/(c^2*x^2-1)*\operatorname{arccosh}(c*x)^2+a^2*(1/2*x/c^2/d/(-c^2*d*x^2+d)^{3/2}-1/2/c^2*(1/3/d*x/(-c^2*d*x^2+d)^{3/2}+2/3/d^2*x/(-c^2*d*x^2+d)^{1/2}))+2/3*b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3*c^4*x^7-b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3*c^2*x^5+1/3*b^2*(-d*(c^2*x^2-1))^{1/2}/(3*c^8*x^8-9*c^6*x^6+10*c^4*x^4-5*c^2*x^2+1)/d^3*\operatorname{arccosh}(c*x)^2*x^3+1/3*a*b*(-d*(c^2*x^2-1))^{1/2}*(c^3*x^3+(c*x-1)^{1/2}*(c*x+1)^{1/2})*c^2*x^2-(c*x-1)^{1/2}*(c*x+1)^{1/2}*(-2*(c*x-1)^{1/2}...$$

### 3.217.5 Fracas [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 x^2}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-(b^2*x^2*arccosh(c*x))^2 + 2*a*b*x^2*arccosh(c*x) + a^2*x^2)*sqrt(-c^2*d*x^2 + d)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

## 3.217.6 Sympy [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{\frac{5}{2}}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(5/2), x)`

output `Integral(x**2*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**5/2, x)`

## 3.217.7 Maxima [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^2}{(-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2), x, algorithm="maxima")`

output `1/3*a*b*c*(sqrt(-d)/(c^6*d^3*x^2 - c^4*d^3) - sqrt(-d)*log(c*x + 1)/(c^4*d^3) - sqrt(-d)*log(c*x - 1)/(c^4*d^3)) - 2/3*a*b*(x/(sqrt(-c^2*d*x^2 + d)*c^2*d^2) - x/((-c^2*d*x^2 + d)^(3/2)*c^2*d))*arccosh(c*x) - 1/3*a^2*(x/(sqrt(-c^2*d*x^2 + d)*c^2*d^2) - x/((-c^2*d*x^2 + d)^(3/2)*c^2*d)) + b^2*integrate(x^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/(-c^2*d*x^2 + d)^(5/2), x)`

## 3.217.8 Giac [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x^2}{(-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x^2/(-c^2*d*x^2 + d)^(5/2), x)`

---

3.217.  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

**3.217.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x^2*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2), x)`output `int((x^2*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2), x)`

**3.218** 
$$\int \frac{x(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

3.218.1 Optimal result . . . . . 1933  
 3.218.2 Mathematica [A] (warning: unable to verify) . . . . . 1934  
 3.218.3 Rubi [C] (verified) . . . . . 1934  
 3.218.4 Maple [A] (verified) . . . . . 1938  
 3.218.5 Fracas [F] . . . . . 1939  
 3.218.6 Sympy [F] . . . . . 1939  
 3.218.7 Maxima [F] . . . . . 1939  
 3.218.8 Giac [F] . . . . . 1940  
 3.218.9 Mupad [F(-1)] . . . . . 1940

**3.218.1 Optimal result**

Integrand size = 27, antiderivative size = 298

$$\begin{aligned} \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx &= -\frac{b^2}{3c^2d^2\sqrt{d-c^2dx^2}} \\ &+ \frac{bx\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{3cd^2(1-c^2x^2)\sqrt{d-c^2dx^2}} + \frac{(a+b\operatorname{arccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\ &+ \frac{2b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{3c^2d^2\sqrt{d-c^2dx^2}} \\ &+ \frac{b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})}{3c^2d^2\sqrt{d-c^2dx^2}} \\ &- \frac{b^2\sqrt{-1+cx}\sqrt{1+cx}\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{3c^2d^2\sqrt{d-c^2dx^2}} \end{aligned}$$

```
output 1/3*(a+b*arccosh(c*x))^2/c^2/d/(-c^2*d*x^2+d)^(3/2)-1/3*b^2/c^2/d^2/(-c^2*
d*x^2+d)^(1/2)+1/3*b*x*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d^
2/(-c^2*x^2+1)/(-c^2*d*x^2+d)^(1/2)+2/3*b*(a+b*arccosh(c*x))*arctanh(c*x+(
c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^2/d^2/(-c^2*d*x^
2+d)^(1/2)+1/3*b^2*polylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/
2)*(c*x+1)^(1/2)/c^2/d^2/(-c^2*d*x^2+d)^(1/2)-1/3*b^2*polylog(2,c*x+(c*x-1
)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^2/d^2/(-c^2*d*x^2+d)^(
1/2)
```

### 3.218.2 Mathematica [A] (warning: unable to verify)

Time = 2.50 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.30

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \frac{4a^2 + b^2 \left( -2 + 4\operatorname{arccosh}(cx)^2 + 2 \cosh(2\operatorname{arccosh}(cx)) - 3\sqrt{\frac{-1+cx}{1+cx}}(1+cx)\operatorname{arccosh}(cx) \right)}{(d - c^2 dx^2)^{5/2}}$$

input `Integrate[(x*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2),x]`

output `(4*a^2 + b^2*(-2 + 4*ArcCosh[c*x]^2 + 2*Cosh[2*ArcCosh[c*x]] - 3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 - E^(-ArcCosh[c*x])] + 3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]*Log[1 + E^(-ArcCosh[c*x])] + 4*((-1 + c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3*PolyLog[2, -E^(-ArcCosh[c*x])] - 4*((-1 + c*x)/(1 + c*x))^(3/2)*(1 + c*x)^3*PolyLog[2, E^(-ArcCosh[c*x])]) + 2*ArcCosh[c*x]*Sinh[2*ArcCosh[c*x]] + ArcCosh[c*x]*Log[1 - E^(-ArcCosh[c*x])]*Sinh[3*ArcCosh[c*x]] - ArcCosh[c*x]*Log[1 + E^(-ArcCosh[c*x])]*Sinh[3*ArcCosh[c*x]]) + a*b*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(4*c*x + 3*(Log[Cosh[ArcCosh[c*x]/2]] - Log[Sinh[ArcCosh[c*x]/2]]) + (8*ArcCosh[c*x] + (-Log[Cosh[ArcCosh[c*x]/2]] + Log[Sinh[ArcCosh[c*x]/2]])*Sinh[3*ArcCosh[c*x]]))/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)))/(12*c^2*d*(d - c^2*d*x^2)^(3/2))`

### 3.218.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.94 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.63, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.370$ , Rules used = {6329, 6304, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

↓ 6329

$$\frac{2b\sqrt{cx - 1}\sqrt{cx + 1} \int \frac{a + \operatorname{barccosh}(cx)}{(1-cx)^2(cx+1)^2} dx}{3cd^2\sqrt{d - c^2 dx^2}} + \frac{(a + \operatorname{barccosh}(cx))^2}{3c^2 d (d - c^2 dx^2)^{3/2}}$$

---

3.218.  $\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$

$$\begin{aligned}
& \downarrow 6304 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx}{3cd^2\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} \\
& \downarrow 6316 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} \right)}{\frac{3cd^2\sqrt{d-c^2dx^2}}{(a+\operatorname{barccosh}(cx))^2} + 3c^2d(d-c^2dx^2)^{3/2}} + \\
& \downarrow 83 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{\frac{3cd^2\sqrt{d-c^2dx^2}}{(a+\operatorname{barccosh}(cx))^2} + 3c^2d(d-c^2dx^2)^{3/2}} + \\
& \downarrow 6318 \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{\frac{3cd^2\sqrt{d-c^2dx^2}}{(a+\operatorname{barccosh}(cx))^2} + 3c^2d(d-c^2dx^2)^{3/2}} + \\
& \downarrow 3042 \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} + \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3cd^2\sqrt{d-c^2dx^2}} \\
& \downarrow 26 \\
& \frac{(a+\operatorname{barccosh}(cx))^2}{3c^2d(d-c^2dx^2)^{3/2}} + \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3cd^2\sqrt{d-c^2dx^2}} \\
& \downarrow 4670
\end{aligned}$$

---

3.218.  $\int \frac{x(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$



$$\begin{aligned}
& \frac{(a + \operatorname{barccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}} + \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int \log(1 - e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1 + e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right)}{3cd^2\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{2715} \\
& \frac{(a + \operatorname{barccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}} + \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1 - e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1 + e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right)}{3cd^2\sqrt{d - c^2dx^2}} \\
& \quad \downarrow \text{2838} \\
& \frac{(a + \operatorname{barccosh}(cx))^2}{3c^2d(d - c^2dx^2)^{3/2}} + \\
& \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right)}{3cd^2\sqrt{d - c^2dx^2}} + \dots
\end{aligned}$$

input `Int[(x*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]`

output `(a + b*ArcCosh[c*x])^2/(3*c^2*d*(d - c^2*d*x^2)^(3/2)) + (2*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/2*b/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c))/(3*c*d^2*Sqrt[d - c^2*d*x^2])`

### 3.218.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

$$3.218. \quad \int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx$$

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x
_Symbol] :> Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x]
+ (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x
)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e
+ f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((d1_) + (e1_.)*(x_))^(p_.)*(
(d2_) + (e2_.)*(x_))^(p_.), x_Symbol] :> Int[(d1*d2 + e1*e2*x^2)^p*(a + b*A
rcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 +
d1*e2, 0] && IntegerQ[p]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((d_) + (e_.)*(x_)^2)^(p_), x
_Symbol] :> Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p +
1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*
ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 +
c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a
+ b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*
d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_/((d_) + (e_.)*(x_)^2), x_Symb
ol] :> Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x
]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_.)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

### 3.218.4 Maple [A] (verified)

Time = 1.19 (sec) , antiderivative size = 582, normalized size of antiderivative = 1.95

method	result
default	$\frac{a^2}{3c^2d(-c^2dx^2+d)^{\frac{3}{2}}} + b^2 \left( \frac{\sqrt{-d(c^2x^2-1)} (\sqrt{cx+1} \operatorname{arccosh}(cx)\sqrt{cx-1} cx+c^2x^2+\operatorname{arccosh}(cx)^2-1)}{3(c^2x^2-1)^2 d^3 c^2} - \frac{\sqrt{-d(c^2x^2-1)} \sqrt{cx-1} \sqrt{cx+1}}{3(c^2x^2-1)^2 d^3 c^2} \right)$
parts	$\frac{a^2}{3c^2d(-c^2dx^2+d)^{\frac{3}{2}}} + b^2 \left( \frac{\sqrt{-d(c^2x^2-1)} (\sqrt{cx+1} \operatorname{arccosh}(cx)\sqrt{cx-1} cx+c^2x^2+\operatorname{arccosh}(cx)^2-1)}{3(c^2x^2-1)^2 d^3 c^2} - \frac{\sqrt{-d(c^2x^2-1)} \sqrt{cx-1} \sqrt{cx+1}}{3(c^2x^2-1)^2 d^3 c^2} \right)$

```
input int(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)
```

```
output 1/3*a^2/c^2/d/(-c^2*d*x^2+d)^(3/2)+b^2*(1/3*(-d*(c^2*x^2-1))^(1/2)*((c*x+1)
)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c*x+c^2*x^2+arccosh(c*x)^2-1)/(c^2*x^2-
1)^2/d^3/c^2-1/3*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^
2/(c^2*x^2-1)*arccosh(c*x)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-1/3*(-d*(
c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^2/(c^2*x^2-1)*polylog(
2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/3*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/
2)*(c*x+1)^(1/2)/d^3/c^2/(c^2*x^2-1)*arccosh(c*x)*ln(1-c*x-(c*x-1)^(1/2)*(
c*x+1)^(1/2))+1/3*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c
^2/(c^2*x^2-1)*polylog(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+2*a*b*(1/6*(-d*(
c^2*x^2-1))^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+2*arccosh(c*x))/(c^2*x
^2-1)^2/d^3/c^2-1/6*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3
/c^2/(c^2*x^2-1)*ln(1+c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/6*(-d*(c^2*x^2-1)
)^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/c^2/(c^2*x^2-1)*ln((c*x-1)^(1/2)*(
c*x+1)^(1/2)+c*x-1))
```

$$3.218. \int \frac{x(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

**3.218.5 Fracas [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*x*arccosh(c*x)^2 + 2*a*b*x*arccosh(c*x) + a^2*x)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

**3.218.6 Sympy [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{5/2}} dx$$

input `integrate(x*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral(x*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`

**3.218.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/3*a^2/((-c^2*d*x^2 + d)^(3/2)*c^2*d) + integrate(b^2*x*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/(-c^2*d*x^2 + d)^(5/2) + 2*a*b*x*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(-c^2*d*x^2 + d)^(5/2), x)`

**3.218.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 x}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*x/(-c^2*d*x^2 + d)^(5/2), x)`

**3.218.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((x*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2),x)`

output `int((x*(a + b*acosh(c*x))^2)/(d - c^2*d*x^2)^(5/2), x)`

**3.219** 
$$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$$

3.219.1 Optimal result . . . . . 1941  
 3.219.2 Mathematica [A] (warning: unable to verify) . . . . . 1942  
 3.219.3 Rubi [C] (verified) . . . . . 1942  
 3.219.4 Maple [B] (verified) . . . . . 1948  
 3.219.5 Fracas [F] . . . . . 1949  
 3.219.6 Sympy [F] . . . . . 1949  
 3.219.7 Maxima [F] . . . . . 1949  
 3.219.8 Giac [F] . . . . . 1950  
 3.219.9 Mupad [F(-1)] . . . . . 1950

**3.219.1 Optimal result**

Integrand size = 26, antiderivative size = 331

$$\begin{aligned} \int \frac{(a + \operatorname{arccosh}(cx))^2}{(d - c^2dx^2)^{5/2}} dx &= -\frac{b^2x}{3d^2\sqrt{d - c^2dx^2}} \\ &+ \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{3cd^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} + \frac{x(a + \operatorname{arccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} \\ &+ \frac{2x(a + \operatorname{arccosh}(cx))^2}{3d^2\sqrt{d - c^2dx^2}} + \frac{2\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{3cd^2\sqrt{d - c^2dx^2}} \\ &- \frac{4b\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx)) \log(1 - e^{2\operatorname{arccosh}(cx)})}{3cd^2\sqrt{d - c^2dx^2}} \\ &- \frac{2b^2\sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{3cd^2\sqrt{d - c^2dx^2}} \end{aligned}$$

```
output 1/3*x*(a+b*arccosh(c*x))^2/d/(-c^2*d*x^2+d)^(3/2)-1/3*b^2*x/d^2/(-c^2*d*x^2+d)^(1/2)+2/3*x*(a+b*arccosh(c*x))^2/d^2/(-c^2*d*x^2+d)^(1/2)+1/3*b*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d^2/(-c^2*x^2+1)/(-c^2*d*x^2+d)^(1/2)+2/3*(a+b*arccosh(c*x))^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d^2/(-c^2*d*x^2+d)^(1/2)-4/3*b*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d^2/(-c^2*d*x^2+d)^(1/2)-2/3*b^2*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/d^2/(-c^2*d*x^2+d)^(1/2)
```

**3.219.2 Mathematica [A] (warning: unable to verify)**

Time = 1.64 (sec) , antiderivative size = 289, normalized size of antiderivative = 0.87

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \frac{\frac{a^2 cx(-3+2c^2 x^2)}{-1+c^2 x^2} + ab \left( 2cx \left( 2 + \frac{1}{1-c^2 x^2} \right) \operatorname{arccosh}(cx) + \frac{\sqrt{\frac{-1+cx}{1+cx}} (-1+(4-4c^2 x^2)) \log\left(\sqrt{\frac{-1+cx}{1+cx}}\right)}{-1+cx} \right)}{(d - c^2 dx^2)^{5/2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(d - c^2*d*x^2)^(5/2), x]`

output

```
((a^2*c*x*(-3 + 2*c^2*x^2))/(-1 + c^2*x^2) + a*b*(2*c*x*(2 + (1 - c^2*x^2)
^(-1))*ArcCosh[c*x] + (Sqrt[(-1 + c*x)/(1 + c*x)]*(-1 + (4 - 4*c^2*x^2)*Lo
g[Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)])))/(-1 + c*x)) + b^2*(-((ArcCosh[c*
x]*(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) + c*x*ArcCosh[c*x])))/(-1 + c^2*x^
2)) + c*x*(-1 + 2*ArcCosh[c*x]^2) - 2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)
*ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 - E^(-2*ArcCosh[c*x])]) + 2*Sqrt[(-1
+ c*x)/(1 + c*x)]*(1 + c*x)*PolyLog[2, E^(-2*ArcCosh[c*x])]))/(3*c*d^2*Sq
rt[d - c^2*d*x^2])
```

**3.219.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 1.71 (sec) , antiderivative size = 278, normalized size of antiderivative = 0.84, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6316, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838, 6329, 41}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

$$\downarrow \text{6316}$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{3d} + \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

$$\downarrow \text{6314}$$

---

3.219.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
& \frac{2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}\right)}{3d} + \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{6327} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
& \frac{2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}\right)}{3d} + \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{6328} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
& \frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+\operatorname{barccosh}(cx)) \operatorname{darccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} dx}{cd\sqrt{d-c^2dx^2}}\right)}{3d} + \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
& \frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -i(a+\operatorname{barccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx)+\frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{cd\sqrt{d-c^2dx^2}}\right)}{3d} + \\
& \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{26} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
& \frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \tan\left(i\operatorname{arccosh}(cx)+\frac{\pi}{2}\right) \operatorname{darccosh}(cx)}{cd\sqrt{d-c^2dx^2}}\right)}{3d} + \\
& \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}
\end{aligned}$$

---

3.219.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$



$$\begin{aligned}
 & \downarrow 4199 \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
 & 2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( 2i \int -\frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b} \right)}{cd\sqrt{d-c^2dx^2}} \right) \\
 & \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 25 \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
 & 2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} d\operatorname{arccosh}(cx) - \frac{i(a+\operatorname{barccosh}(cx))^2}{2b} \right)}{cd\sqrt{d-c^2dx^2}} \right) \\
 & \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 2620 \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
 & 2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{2}b \int \log(1-e^{2\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - \frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) \right) \right)}{cd\sqrt{d-c^2dx^2}} \right) \\
 & \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 2715 \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
 & 2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{4}b \int e^{-2\operatorname{arccosh}(cx)} \log(1-e^{2\operatorname{arccosh}(cx)}) de^{2\operatorname{arccosh}(cx)} - \frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) (a+\operatorname{barccosh}(cx)) \right) \right)}{cd\sqrt{d-c^2dx^2}} \right) \\
 & \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
 & \downarrow 2838
 \end{aligned}$$

---

3.219.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \\
& 2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right) - \frac{i(a+b\operatorname{arccosh}(cx))}{cd\sqrt{d-c^2dx^2}} \right)}{cd\sqrt{d-c^2dx^2}} \right) \\
& \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad 3d \\
& \quad \downarrow \quad 6329 \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{b \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} + \\
& 2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right) - \frac{i(a+b\operatorname{arccosh}(cx))}{cd\sqrt{d-c^2dx^2}} \right)}{cd\sqrt{d-c^2dx^2}} \right) \\
& \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad 3d \\
& \quad \downarrow \quad 41 \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)} - \frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \\
& 2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) (a+\operatorname{barccosh}(cx)) - \frac{1}{4} b \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)}) \right) - \frac{i(a+b\operatorname{arccosh}(cx))}{cd\sqrt{d-c^2dx^2}} \right)}{cd\sqrt{d-c^2dx^2}} \right) \\
& \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad 3d
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])^2/(d - c^2*d*x^2)^(5/2), x]`

output `(x*(a + b*ArcCosh[c*x])^2)/(3*d*(d - c^2*d*x^2)^(3/2)) + (2*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/2*(b*x)/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*ArcCosh[c*x])/(2*c^2*(1 - c^2*x^2)))/(3*d^2*Sqrt[d - c^2*d*x^2]) + (2*((x*(a + b*ArcCosh[c*x])^2)/(d*Sqrt[d - c^2*d*x^2]) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])]))/4))/(c*d*Sqrt[d - c^2*d*x^2]))/(3*d)`

---

3.219.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx$

## 3.219.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 41 `Int[1/(((a_) + (b_)*(x_))^(3/2)*((c_) + (d_)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4199 `Int[(((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2),  
x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp  
[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a  
+ b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e},  
x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x  
_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p +  
1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*  
ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 +  
c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a  
+ b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*  
d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (  
e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.)), x_Symbol] := Int[(f*x)^m*(d1  
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2  
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2),  
x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]  
], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p  
_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p  
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +  
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x  
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&  
GtQ[n, 0] && NeQ[p, -1]`

**3.219.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 2434 vs.  $2(313) = 626$ .

Time = 1.16 (sec) , antiderivative size = 2435, normalized size of antiderivative = 7.36

method	result	size
default	Expression too large to display	2435
parts	Expression too large to display	2435

input `int((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output

$$\frac{2}{3}b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3(c*x-1)*(c*x+1)*x-4/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^6\operatorname{arccosh}(c*x)*x^7-2b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^4\operatorname{arccosh}(c*x)^2x^5+14/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^4\operatorname{arccosh}(c*x)*x^5+17/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^2\operatorname{arccosh}(c*x)^2x^3-16/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^2\operatorname{arccosh}(c*x)*x^3-4/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3/c*(c*x-1)^{1/2}*(c*x+1)^{1/2}+4/3b^2(-d(c^2x^2-1))^{1/2}*(c*x-1)^{1/2}*(c*x+1)^{1/2}/d^3/c/(c^2x^2-1)*\operatorname{polylog}(2,-c*x-(c*x-1)^{1/2}*(c*x+1)^{1/2})+2/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^4*(c*x-1)*(c*x+1)*x^5-4/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^2*(c*x-1)*(c*x+1)*x^3+2b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3(c*x-1)*(c*x+1)*\operatorname{arccosh}(c*x)*x-4/3b^2(-d(c^2x^2-1))^{1/2}*(c*x-1)^{1/2}*(c*x+1)^{1/2}/d^3/c/(c^2x^2-1)*\operatorname{arccosh}(c*x)^2-4b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3\operatorname{arccosh}(c*x)^2x+2b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3\operatorname{arccosh}(c*x)*x-2/3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c^6x^7+3b^2(-d(c^2x^2-1))^{1/2}/(3c^6x^6-10c^4x^4+11c^2x^2-4)/d^3c\dots$$

## 3.219.5 Fricas [F]

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)`

## 3.219.6 Sympy [F]

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{5/2}} dx$$

input `integrate((a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral((a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**(5/2), x)`

## 3.219.7 Maxima [F]

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/3*a*b*c*(sqrt(-d)/(c^4*d^3*x^2 - c^2*d^3) + 2*sqrt(-d)*log(c*x + 1)/(c^2*d^3) + 2*sqrt(-d)*log(c*x - 1)/(c^2*d^3)) + 2/3*a*b*(2*x/(sqrt(-c^2*d*x^2 + d)*d^2) + x/((-c^2*d*x^2 + d)^(3/2)*d))*arccosh(c*x) + 1/3*a^2*(2*x/(sqrt(-c^2*d*x^2 + d)*d^2) + x/((-c^2*d*x^2 + d)^(3/2)*d)) + b^2*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)^2/(-c^2*d*x^2 + d)^(5/2), x)`

**3.219.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/(-c^2*d*x^2 + d)^(5/2), x)`

**3.219.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))^2/(d - c^2*d*x^2)^(5/2),x)`

output `int((a + b*acosh(c*x))^2/(d - c^2*d*x^2)^(5/2), x)`

$$3.220 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$$

3.220.1 Optimal result	. . . . .	1951
3.220.2 Mathematica [A] (warning: unable to verify)	. . . . .	1952
3.220.3 Rubi [A] (verified)	. . . . .	1953
3.220.4 Maple [F]	. . . . .	1962
3.220.5 Fricas [F]	. . . . .	1963
3.220.6 Sympy [F]	. . . . .	1963
3.220.7 Maxima [F]	. . . . .	1963
3.220.8 Giac [F]	. . . . .	1964
3.220.9 Mupad [F(-1)]	. . . . .	1964

### 3.220.1 Optimal result

Integrand size = 29, antiderivative size = 597

$$\begin{aligned} \int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx = & -\frac{b^2}{3d^2\sqrt{d-c^2dx^2}} \\ & + \frac{bcx\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}{3d^2(1-c^2x^2)\sqrt{d-c^2dx^2}} \\ & + \frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d^2\sqrt{d-c^2dx^2}} \\ & + \frac{2\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d-c^2dx^2}} \\ & + \frac{14b\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx)) \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{3d^2\sqrt{d-c^2dx^2}} \\ & + \frac{7b^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{3d^2\sqrt{d-c^2dx^2}} \\ & - \frac{2ib\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d-c^2dx^2}} \\ & + \frac{2ib\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d-c^2dx^2}} \\ & - \frac{7b^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{3d^2\sqrt{d-c^2dx^2}} \\ & + \frac{2ib^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d-c^2dx^2}} \\ & - \frac{2ib^2\sqrt{-1+cx}\sqrt{1+cx} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{d^2\sqrt{d-c^2dx^2}} \end{aligned}$$

---

3.220.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$



output  $1/3*(a+b*\operatorname{arccosh}(c*x))^2/d/(-c^2*d*x^2+d)^{(3/2)}-1/3*b^2/d^2/(-c^2*d*x^2+d)^{(1/2)}+(a+b*\operatorname{arccosh}(c*x))^2/d^2/(-c^2*d*x^2+d)^{(1/2)}+1/3*b*c*x*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*x^2+1)/(-c^2*d*x^2+d)^{(1/2)}+2*(a+b*\operatorname{arccosh}(c*x))^2*\operatorname{arctan}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}+14/3*b*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}+7/3*b^2*\operatorname{polylog}(2,-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-2*I*b*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}+2*I*b*(a+b*\operatorname{arccosh}(c*x))*\operatorname{polylog}(2,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-7/3*b^2*\operatorname{polylog}(2,c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}+2*I*b^2*\operatorname{polylog}(3,-I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-2*I*b^2*\operatorname{polylog}(3,I*(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}$

### 3.220.2 Mathematica [A] (warning: unable to verify)

Time = 9.53 (sec) , antiderivative size = 818, normalized size of antiderivative = 1.37

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx = \sqrt{-d(-1 + c^2 x^2)} \left( \frac{a^2}{3d^3(-1 + c^2 x^2)^2} - \frac{a^2}{d^3(-1 + c^2 x^2)} \right) + \frac{a^2 \log(cx)}{d^{5/2}} - \frac{a^2 \log\left(d + \sqrt{d} \sqrt{-d(-1 + c^2 x^2)}\right)}{d^{5/2}} + \frac{ab \sqrt{\frac{-1+cx}{1+cx}}(1+cx)}{d^{5/2}} \left( 14 \operatorname{arccosh}(cx) \coth\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) - \operatorname{csch}^2\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) - \frac{1}{2} \sqrt{\frac{-1+cx}{1+cx}}(1+cx) \operatorname{arccosh}(cx) \right) + \frac{b^2 \sqrt{\frac{-1+cx}{1+cx}}(1+cx)}{d^{5/2}} \left( -4 \coth\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) + 14 \operatorname{arccosh}(cx)^2 \coth\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) - 2 \operatorname{arccosh}(cx) \operatorname{csch}^2\left(\frac{1}{2} \operatorname{arccosh}(cx)\right) \right)$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x*(d - c^2*d*x^2)^(5/2)), x]`

output  $\text{Sqrt}[-(d*(-1 + c^2*x^2))]*(a^2/(3*d^3*(-1 + c^2*x^2)^2) - a^2/(d^3*(-1 + c^2*x^2))) + (a^2*\text{Log}[c*x])/d^{(5/2)} - (a^2*\text{Log}[d + \text{Sqrt}[d]*\text{Sqrt}[-(d*(-1 + c^2*x^2))]])/d^{(5/2)} + (a*b*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(14*\text{ArcCosh}[c*x]*\text{Coth}[\text{ArcCosh}[c*x]/2] - \text{Csch}[\text{ArcCosh}[c*x]/2]^2 - (\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{ArcCosh}[c*x]*\text{Csch}[\text{ArcCosh}[c*x]/2]^4)/2 - (24*I)*\text{ArcCosh}[c*x]*\text{Log}[1 - I/E^{\text{ArcCosh}[c*x]}] + (24*I)*\text{ArcCosh}[c*x]*\text{Log}[1 + I/E^{\text{ArcCosh}[c*x]}] + 28*\text{Log}[\text{Cosh}[\text{ArcCosh}[c*x]/2]] - 28*\text{Log}[\text{Sinh}[\text{ArcCosh}[c*x]/2]] - (24*I)*\text{PolyLog}[2, (-I)/E^{\text{ArcCosh}[c*x]}] + (24*I)*\text{PolyLog}[2, I/E^{\text{ArcCosh}[c*x]}] - \text{Sech}[\text{ArcCosh}[c*x]/2]^2 - (8*\text{ArcCosh}[c*x]*\text{Sinh}[\text{ArcCosh}[c*x]/2]^4)/(((1 + c*x)/(1 + c*x))^{(3/2)}*(1 + c*x)^3) - 14*\text{ArcCosh}[c*x]*\text{Tanh}[\text{ArcCosh}[c*x]/2])/((12*d^2*\text{Sqrt}[-(d*(-1 + c*x)*(1 + c*x))]) + (b^2*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(-4*\text{Coth}[\text{ArcCosh}[c*x]/2] + 14*\text{ArcCosh}[c*x]^2*\text{Coth}[\text{ArcCosh}[c*x]/2] - 2*\text{ArcCosh}[c*x]*\text{Csch}[\text{ArcCosh}[c*x]/2]^2 - (\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*\text{ArcCosh}[c*x]^2*\text{Csch}[\text{ArcCosh}[c*x]/2]^4)/2 - 56*\text{ArcCosh}[c*x]*\text{Log}[1 - E^{(-\text{ArcCosh}[c*x])}] - (24*I)*\text{ArcCosh}[c*x]^2*\text{Log}[1 - I/E^{\text{ArcCosh}[c*x]}] + (24*I)*\text{ArcCosh}[c*x]^2*\text{Log}[1 + I/E^{\text{ArcCosh}[c*x]}] + 56*\text{ArcCosh}[c*x]*\text{Log}[1 + E^{(-\text{ArcCosh}[c*x])}] - 56*\text{PolyLog}[2, -E^{(-\text{ArcCosh}[c*x])}] - (48*I)*\text{ArcCosh}[c*x]*\text{PolyLog}[2, (-I)/E^{\text{ArcCosh}[c*x]}] + (48*I)*\text{ArcCosh}[c*x]*\text{PolyLog}[2, I/E^{\text{ArcCosh}[c*x]}] + 56*\text{PolyLog}[2, E^{(-\text{ArcCosh}[c*x])}] - (48*I)*\text{PolyLog}[3, (-I)/E^{\text{ArcCosh}[c*x]}] + (48*I)*\text{PolyLog}[3, I/E^{\text{ArcCosh}[c*x]}] - 2*\text{ArcCosh}[c*x]*S...$

### 3.220.3 Rubi [A] (verified)

Time = 4.99 (sec) , antiderivative size = 448, normalized size of antiderivative = 0.75, number of steps used = 26, number of rules used = 25,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.862$ , Rules used = {6351, 6304, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838, 6351, 25, 6304, 6318, 3042, 26, 4670, 2715, 2838, 6361, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \text{barccosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx$$

$$\downarrow \text{6351}$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\text{barccosh}(cx)}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\text{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a + \text{barccosh}(cx))^2}{3d(d - c^2 dx^2)^{3/2}}$$

$$\downarrow \text{6304}$$

---

3.220.  $\int \frac{(a+\text{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx + \int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{6316} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{83} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{6318} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int i(a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{26} \\
& \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \int (a+\operatorname{barccosh}(cx)) \csc(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \\
& \quad \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \\
& \quad \downarrow \text{4670}
\end{aligned}$$

---

3.220.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right)$$

$$3d^2\sqrt{d-c^2dx^2}$$

$$\frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 2715

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right)$$

$$3d^2\sqrt{d-c^2dx^2}$$

$$\frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 2838

$$\frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right) +$$

$$3d^2\sqrt{d-c^2dx^2}$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 6351

$$-\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right) +$$

$$3d^2\sqrt{d-c^2dx^2}$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 25

---

3.220.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) +$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad 3d^2\sqrt{d-c^2dx^2}$$

↓ 6304

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) +$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad 3d^2\sqrt{d-c^2dx^2}$$

↓ 6318

$$-\frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) +$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad 3d^2\sqrt{d-c^2dx^2}$$

↓ 3042

$$\frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int i(a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} +$$

$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) +$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad 3d^2\sqrt{d-c^2dx^2}$$

↓ 26

---

3.220.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$

$$\frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} +$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right)$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad \frac{3d^2\sqrt{d-c^2dx^2}}{3d(d-c^2dx^2)^{3/2}}$$

↓ 4670

$$-\frac{2ib\sqrt{cx-1}\sqrt{cx+1} (ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx)-ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx)+2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx)))}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right)$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad \frac{3d^2\sqrt{d-c^2dx^2}}{3d(d-c^2dx^2)^{3/2}}$$

↓ 2715

$$-\frac{2ib\sqrt{cx-1}\sqrt{cx+1} (ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)}-ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)}+2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx)))}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right)$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad \frac{3d^2\sqrt{d-c^2dx^2}}{3d(d-c^2dx^2)^{3/2}}$$

↓ 2838

$$\frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} (2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib \operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib \operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right)$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \quad \frac{3d^2\sqrt{d-c^2dx^2}}{3d(d-c^2dx^2)^{3/2}}$$

↓ 6361

---

3.220.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int \frac{(a+b\operatorname{arccosh}(cx))^2}{cx} d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( 2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}\left(2, -e^{\operatorname{arccosh}(cx)}\right) \right)}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}\left(2, -e^{\operatorname{arccosh}(cx)}\right) \right) - ib \operatorname{PolyLog}\left(2, e^{\operatorname{arccosh}(cx)}\right)}{2c} \right) + \frac{d}{3d^2\sqrt{d-c^2dx^2}}$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 3042

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int (a+b\operatorname{arccosh}(cx))^2 \csc\left(i\operatorname{arccosh}(cx) + \frac{\pi}{2}\right) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} - \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( 2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}\left(2, -e^{\operatorname{arccosh}(cx)}\right) \right)}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}\left(2, -e^{\operatorname{arccosh}(cx)}\right) \right) - ib \operatorname{PolyLog}\left(2, e^{\operatorname{arccosh}(cx)}\right)}{2c} \right) + \frac{d}{3d^2\sqrt{d-c^2dx^2}}$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 4668

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \left( -2ib \int (a+b\operatorname{arccosh}(cx)) \log\left(1 - ie^{\operatorname{arccosh}(cx)}\right) d\operatorname{arccosh}(cx) + 2ib \int (a+b\operatorname{arccosh}(cx)) \log\left(1 + ie^{\operatorname{arccosh}(cx)}\right) d\operatorname{arccosh}(cx) \right)}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}\left(2, -e^{\operatorname{arccosh}(cx)}\right) \right) - ib \operatorname{PolyLog}\left(2, e^{\operatorname{arccosh}(cx)}\right)}{2c} \right) + \frac{d}{3d^2\sqrt{d-c^2dx^2}}$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 3011

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \left( 2ib \left( b \int \operatorname{PolyLog}\left(2, -ie^{\operatorname{arccosh}(cx)}\right) d\operatorname{arccosh}(cx) - \operatorname{PolyLog}\left(2, -ie^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) \right) - 2ib \left( b \int \operatorname{PolyLog}\left(2, ie^{\operatorname{arccosh}(cx)}\right) d\operatorname{arccosh}(cx) - \operatorname{PolyLog}\left(2, ie^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) \right) \right)}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}\left(2, -e^{\operatorname{arccosh}(cx)}\right) \right) - ib \operatorname{PolyLog}\left(2, e^{\operatorname{arccosh}(cx)}\right)}{2c} \right) + \frac{d}{3d^2\sqrt{d-c^2dx^2}}$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

↓ 2720

---

3.220.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}} dx$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}\left(2ib\left(\int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(cx)}\right)de^{\operatorname{arccosh}(cx)}-\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))\right)-2ib\left(\int e^{-\operatorname{arccosh}(cx)} \operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)de^{\operatorname{arccosh}(cx)}-\operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))\right)\right)}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1}\left(-\frac{i\left(2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}\left(2,-e^{\operatorname{arccosh}(cx)}\right)-ib\operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)\right)}{2c}\right)+\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$


---

↓ 7143

$$\frac{\sqrt{cx-1}\sqrt{cx+1}\left(2\operatorname{arctan}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))^2+2ib\left(b\operatorname{PolyLog}\left(3,-ie^{\operatorname{arccosh}(cx)}\right)-\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))\right)\right)}{d\sqrt{d-c^2dx^2}}$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1}\left(-\frac{i\left(2i\operatorname{arctanh}\left(e^{\operatorname{arccosh}(cx)}\right)(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}\left(2,-e^{\operatorname{arccosh}(cx)}\right)-ib\operatorname{PolyLog}\left(2,e^{\operatorname{arccosh}(cx)}\right)\right)}{2c}\right)+\frac{(a+b\operatorname{arccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x*(d - c^2*d*x^2)^(5/2)),x]`

output `(a + b*ArcCosh[c*x])^2/(3*d*(d - c^2*d*x^2)^(3/2)) + (2*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/2*b/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (x*(a + b*ArcCosh[c*x]))/(2*(1 - c^2*x^2)) - ((I/2)*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/c))/(3*d^2*Sqrt[d - c^2*d*x^2]) + ((a + b*ArcCosh[c*x])^2/(d*Sqrt[d - c^2*d*x^2]) - ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((2*I)*(a + b*ArcCosh[c*x])*ArcTanh[E^ArcCosh[c*x]] + I*b*PolyLog[2, -E^ArcCosh[c*x]] - I*b*PolyLog[2, E^ArcCosh[c*x]]))/(d*Sqrt[d - c^2*d*x^2]) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*(a + b*ArcCosh[c*x])^2*ArcTan[E^ArcCosh[c*x]] + (2*I)*b*(-(a + b*ArcCosh[c*x])*PolyLog[2, (-I)*E^ArcCosh[c*x]]) + b*PolyLog[3, (-I)*E^ArcCosh[c*x]]) - (2*I)*b*(-(a + b*ArcCosh[c*x])*PolyLog[2, I*E^ArcCosh[c*x]]) + b*PolyLog[3, I*E^ArcCosh[c*x]]))/(d*Sqrt[d - c^2*d*x^2])/d`



## 3.220.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_)^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/(f*fz*I)], x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x]], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*(d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

rule 6361 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.220.4 Maple [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x(-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `int((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(5/2),x)`

output `int((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(5/2),x)`

**3.220.5 Fricas [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^6*d^3*x^7 - 3*c^4*d^3*x^5 + 3*c^2*d^3*x^3 - d^3*x), x)`

**3.220.6 Sympy [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x(-d(cx - 1)(cx + 1))^{5/2}} dx$$

input `integrate((a+b*acosh(c*x))**2/x/(-c**2*d*x**2+d)**(5/2),x)`

output `Integral((a + b*acosh(c*x))**2/(x*(-d*(c*x - 1)*(c*x + 1))**(5/2)), x)`

**3.220.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `-1/3*a^2*(3*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(5/2) - 3/(sqrt(-c^2*d*x^2 + d)*d^2) - 1/((-c^2*d*x^2 + d)^(3/2)*d)) + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/((-c^2*d*x^2 + d)^(5/2)*x) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(5/2)*x), x)`

**3.220.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x} dx$$

input `integrate((a+b*arccosh(c*x))^2/x/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(5/2)*x), x)`

**3.220.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x(d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x*(d - c^2*d*x^2)^(5/2)),x)`

output `int((a + b*acosh(c*x))^2/(x*(d - c^2*d*x^2)^(5/2)), x)`

**3.221**  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$

3.221.1 Optimal result . . . . . 1965  
 3.221.2 Mathematica [A] (warning: unable to verify) . . . . . 1966  
 3.221.3 Rubi [C] (verified) . . . . . 1967  
 3.221.4 Maple [B] (verified) . . . . . 1977  
 3.221.5 Fracas [F] . . . . . 1978  
 3.221.6 Sympy [F(-1)] . . . . . 1979  
 3.221.7 Maxima [F] . . . . . 1979  
 3.221.8 Giac [F] . . . . . 1979  
 3.221.9 Mupad [F(-1)] . . . . . 1980

**3.221.1 Optimal result**

Integrand size = 29, antiderivative size = 476

$$\begin{aligned} \int \frac{(a + b\operatorname{arccosh}(cx))^2}{x^2(d - c^2dx^2)^{5/2}} dx &= -\frac{b^2c^2x}{3d^2\sqrt{d - c^2dx^2}} \\ &+ \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))}{3d^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} - \frac{(a + b\operatorname{arccosh}(cx))^2}{dx(d - c^2dx^2)^{3/2}} \\ &+ \frac{4c^2x(a + b\operatorname{arccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} + \frac{8c^2x(a + b\operatorname{arccosh}(cx))^2}{3d^2\sqrt{d - c^2dx^2}} \\ &+ \frac{8c\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))^2}{3d^2\sqrt{d - c^2dx^2}} \\ &- \frac{4bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{d^2\sqrt{d - c^2dx^2}} \\ &- \frac{16bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + b\operatorname{arccosh}(cx))\log(1 - e^{2\operatorname{arccosh}(cx)})}{3d^2\sqrt{d - c^2dx^2}} \\ &- \frac{b^2c\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{d^2\sqrt{d - c^2dx^2}} \\ &- \frac{5b^2c\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{3d^2\sqrt{d - c^2dx^2}} \end{aligned}$$

output  $-(a+b*\operatorname{arccosh}(c*x))^2/d/x/(-c^2*d*x^2+d)^{(3/2)}+4/3*c^2*x*(a+b*\operatorname{arccosh}(c*x))^2/d/(-c^2*d*x^2+d)^{(3/2)}-1/3*b^2*c^2*x/d^2/(-c^2*d*x^2+d)^{(1/2)}+8/3*c^2*x*(a+b*\operatorname{arccosh}(c*x))^2/d^2/(-c^2*d*x^2+d)^{(1/2)}+1/3*b*c*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}+8/3*c*(a+b*\operatorname{arccosh}(c*x))^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-4*b*c*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-16/3*b*c*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-b^2*c*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-5/3*b^2*c*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}$

### 3.221.2 Mathematica [A] (warning: unable to verify)

Time = 2.85 (sec) , antiderivative size = 457, normalized size of antiderivative = 0.96

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx = \frac{c \left( \frac{a^2(3 - 12c^2x^2 + 8c^4x^4)}{cx(-1 + c^2x^2)} + ab \left( 10cx \operatorname{arccosh}(cx) - \frac{\sqrt{\frac{-1+cx}{1+cx}}(1+cx) + 2cx \operatorname{arccosh}(cx)}{-1 + c^2x^2} - 2 \sqrt{\frac{-1+cx}{1+cx}} \right) \right)}{d - c^2 dx^2}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^2*(d - c^2*d*x^2)^(5/2)),x]`

output  $(c*((a^2*(3 - 12*c^2*x^2 + 8*c^4*x^4))/(c*x*(-1 + c^2*x^2)) + a*b*(10*c*x*ArcCosh[c*x] - (Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) + 2*c*x*ArcCosh[c*x])/(-1 + c^2*x^2) - 2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*((-3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x])/(c*x) + 3*Log[c*x] + 5*Log[Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)])) + b^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*((c*x*Sqrt[(-1 + c*x)/(1 + c*x)])/(1 - c*x) + ArcCosh[c*x]/(1 - c^2*x^2) - 8*ArcCosh[c*x]^2 - (c*x*ArcCosh[c*x]^2)/((( -1 + c*x)/(1 + c*x))^(3/2))*(1 + c*x)^3) + (5*c*x*ArcCosh[c*x]^2)/(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)) + (3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*ArcCosh[c*x]^2)/(c*x) - 10*ArcCosh[c*x]*Log[1 - E^(-2*ArcCosh[c*x])] - 6*ArcCosh[c*x]*Log[1 + E^(-2*ArcCosh[c*x])] + 3*PolyLog[2, -E^(-2*ArcCosh[c*x])] + 5*PolyLog[2, E^(-2*ArcCosh[c*x])])))/(3*d^2*Sqrt[d - c^2*d*x^2])$

**3.221.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 4.14 (sec) , antiderivative size = 469, normalized size of antiderivative = 0.99, number of steps used = 24, number of rules used = 23,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.793$ , Rules used = {6347, 6316, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838, 6329, 41, 6351, 41, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx$$

$$\downarrow \text{6347}$$

$$-\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{dx (d - c^2 dx^2)^{3/2}}$$

$$\downarrow \text{6316}$$

$$4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{3d} + \frac{x(a + \operatorname{barccosh}(cx))^2}{3d (d - c^2 dx^2)^{3/2}} \right) -$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx (d - c^2 dx^2)^{3/2}}$$

$$\downarrow \text{6314}$$

$$4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right)}{3d} \right) +$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a + \operatorname{barccosh}(cx))^2}{dx (d - c^2 dx^2)^{3/2}}$$

$$\downarrow \text{6327}$$

---

3.221.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$



$$\begin{aligned}
 & 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right)}{3d} \right) + \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{6328} \\
 & 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+\operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}(cx+1)}} d\operatorname{arccosh}}{cd\sqrt{d-c^2dx^2}} \right)}{3d} \right) + \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{3042} \\
 & - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} + \\
 & 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int -i(a+\operatorname{barccosh}(cx)) \tan(i\operatorname{arcc}}{cd\sqrt{d-c^2dx^2}} \right)}{3d} \right) + \\
 & \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{26} \\
 & - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} + \\
 & 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \int (a+\operatorname{barccosh}(cx)) \tan(i\operatorname{arcc}}{cd\sqrt{d-c^2dx^2}} \right)}{3d} \right) + \\
 & \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}
 \end{aligned}$$

---

3.221.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
 & \downarrow 4199 \\
 & 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( 2i \int -\frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} \right)}{cd\sqrt{d-c^2dx^2}} \right)}{3d} \right. \\
 & \qquad \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \right) \\
 & \downarrow 25 \\
 & 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \int \frac{e^{2\operatorname{arccosh}(cx)}(a+\operatorname{barccosh}(cx))}{1-e^{2\operatorname{arccosh}(cx)}} \right)}{cd\sqrt{d-c^2dx^2}} \right)}{3d} \right. \\
 & \qquad \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \right) \\
 & \downarrow 2620 \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} + \\
 & 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{2} b \int \log(1-e^{2\operatorname{arccosh}(cx)}) \right) \right)}{cd\sqrt{d-c^2dx^2}} \right)}{3d} \right. \\
 & \qquad \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \right) \\
 & \downarrow 2715
 \end{aligned}$$

---

3.221.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
 & -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} + \\
 4c^2 & \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{4} b \int e^{-2\operatorname{arccosh}(cx)} \log \right) \right)}{d\sqrt{d-c^2dx^2}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) \\
 & \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{2838}
 \end{aligned}$$

$$\begin{aligned}
 4c^2 & \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( -\frac{1}{2} \log(1-e^{2\operatorname{arccosh}(cx)}) \right) \right)}{d\sqrt{d-c^2dx^2}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{6329}
 \end{aligned}$$

$$\begin{aligned}
 4c^2 & \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{b \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \dots \right) \right)}{d\sqrt{d-c^2dx^2}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) \\
 & \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \\
 & \quad \downarrow \text{41}
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)}dx}{d^2\sqrt{d-c^2dx^2}}+ \\
 4c^2 & \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-\right.\right.\right.} \right. \\
 & \left. \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}\right)}{d^2\sqrt{d-c^2dx^2}} \right) + \\
 & \left. \downarrow 6351 \right. \\
 & -\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)}dx+\frac{1}{2}bc\int\frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}}dx+\frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)}\right)}{d^2\sqrt{d-c^2dx^2}}+ \\
 4c^2 & \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-\right.\right.\right.} \right. \\
 & \left. \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}\right)}{d^2\sqrt{d-c^2dx^2}} \right) + \\
 & \left. \downarrow 41 \right. \\
 & -\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)}dx+\frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)}-\frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2\sqrt{d-c^2dx^2}}+ \\
 4c^2 & \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-\right.\right.\right.} \right. \\
 & \left. \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}\right)}{d^2\sqrt{d-c^2dx^2}} \right) + \\
 & \left. \downarrow 6331 \right.
 \end{aligned}$$

---

3.221.  $\int\frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}}dx$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(-\int\frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)}\operatorname{darccosh}(cx)+\frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)}-\frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2\sqrt{d-c^2dx^2}}+$$

$$4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-\frac{cx-1}{cx+1})\right)\right)}{d\sqrt{d-c^2dx^2}}\right)}{d\sqrt{d-c^2dx^2}}\right)$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}$$

↓ 5984

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(-2\int(a+\operatorname{barccosh}(cx))\operatorname{csch}(2\operatorname{arccosh}(cx))\operatorname{darccosh}(cx)+\frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)}-\frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2\sqrt{d-c^2dx^2}}$$

$$4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-\frac{cx-1}{cx+1})\right)\right)}{d\sqrt{d-c^2dx^2}}\right)}{d\sqrt{d-c^2dx^2}}\right)$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}$$

↓ 3042

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(-2\int i(a+\operatorname{barccosh}(cx))\operatorname{csc}(2i\operatorname{arccosh}(cx))\operatorname{darccosh}(cx)+\frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)}-\frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2\sqrt{d-c^2dx^2}}$$

$$4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-\frac{cx-1}{cx+1})\right)\right)}{d\sqrt{d-c^2dx^2}}\right)}{d\sqrt{d-c^2dx^2}}\right)$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}$$

↓ 26

---

3.221.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(-2i\int(a+\operatorname{barccosh}(cx))\csc(2i\operatorname{arccosh}(cx))\operatorname{darccosh}(cx)+\frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)}-\frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2\sqrt{d-c^2dx^2}}$$


---


$$4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bcx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)\right)}{d\sqrt{d-c^2dx^2}}\right)}{d^2\sqrt{d-c^2dx^2}}\right)$$


---


$$\frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}$$

↓ 4670

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(\frac{1}{2}ib\int\log(1-e^{2\operatorname{arccosh}(cx)})\operatorname{darccosh}(cx)-\frac{1}{2}ib\int\log(1+e^{2\operatorname{arccosh}(cx)})\operatorname{darccosh}(cx)+\frac{d^2\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)\right)\right)\right)}{d^2\sqrt{d-c^2dx^2}}$$


---


$$4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bcx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)\right)}{d\sqrt{d-c^2dx^2}}\right)}{d^2\sqrt{d-c^2dx^2}}\right)$$


---


$$\frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}$$

↓ 2715

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(\frac{1}{4}ib\int e^{-2\operatorname{arccosh}(cx)}\log(1-e^{2\operatorname{arccosh}(cx)})de^{2\operatorname{arccosh}(cx)}-\frac{1}{4}ib\int e^{-2\operatorname{arccosh}(cx)}\log(1+e^{2\operatorname{arccosh}(cx)})de^{2\operatorname{arccosh}(cx)}+\frac{d^2\sqrt{d-c^2dx^2}}{2\sqrt{cx-1}\sqrt{cx+1}}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)\right)\right)\right)}{d^2\sqrt{d-c^2dx^2}}$$


---


$$4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bcx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)\right)}{d\sqrt{d-c^2dx^2}}\right)}{d^2\sqrt{d-c^2dx^2}}\right)$$


---


$$\frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}$$

↓ 2838

---

3.221.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$

$$2bc\sqrt{cx-1}\sqrt{cx+1}\left(-2i(\operatorname{iarctanh}(e^{2\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+\frac{1}{4}ib\operatorname{PolyLog}(2,-e^{2\operatorname{arccosh}(cx)})-\frac{1}{4}ib\operatorname{PolyLog}(2,e^{2\operatorname{arccosh}(cx)}))\right)$$


---


$$4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(\frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)}-\frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)\right)}{d^2\sqrt{d-c^2dx^2}}\right)}{d^2\sqrt{d-c^2dx^2}}\right)$$


---


$$\frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^2*(d - c^2*d*x^2)^(5/2)),x]`

output `-(a + b*ArcCosh[c*x])^2/(d*x*(d - c^2*d*x^2)^(3/2)) - (2*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/2*(b*c*x)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*ArcCosh[c*x])/(2*(1 - c^2*x^2)) - (2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCosh[c*x])]) + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^(2*ArcCosh[c*x])]))/(d^2*Sqrt[d - c^2*d*x^2]) + 4*c^2*((x*(a + b*ArcCosh[c*x])^2)/(3*d*(d - c^2*d*x^2)^(3/2)) + (2*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-1/2*(b*x)/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*ArcCosh[c*x])/(2*c^2*(1 - c^2*x^2))))/(3*d^2*Sqrt[d - c^2*d*x^2]) + (2*((x*(a + b*ArcCosh[c*x])^2)/(d*Sqrt[d - c^2*d*x^2]) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((-1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1 - E^(2*ArcCosh[c*x])]) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/(c*d*Sqrt[d - c^2*d*x^2]))/(3*d)`

### 3.221.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 41 `Int[1/(((a_) + (b_.)*(x_))^(3/2)*((c_) + (d_.)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`

---

3.221.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^2(d-c^2dx^2)^{5/2}} dx$

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[(c + d*x)^m/(b*f*g*n*Log[F])*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_))*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 5984 `Int[Csch[(a_) + (b_)*(x_)]^(n_)*((c_) + (d_)*(x_))^(m_)*Sech[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`



rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2),  
x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp  
[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a  
+ b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e},  
x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x  
_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p +  
1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*  
ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 +  
c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a  
+ b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*  
d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (  
e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.)), x_Symbol] := Int[(f*x)^m*(d1  
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2  
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2),  
x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]  
], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p  
_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p  
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +  
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x  
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&  
GtQ[n, 0] && NeQ[p, -1]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_) + (e_.)*(x_)^2)),  
x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x  
, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IG  
tQ[n, 0]`

```
rule 6347 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]
```

```
rule 6351 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[-(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])
```

### 3.221.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2856 vs.  $2(470) = 940$ .

Time = 1.50 (sec) , antiderivative size = 2857, normalized size of antiderivative = 6.00

method	result	size
default	Expression too large to display	2857
parts	Expression too large to display	2857

```
input int((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)
```

output

```

-64/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(8*c^6*x^6-25*c^4*x^4+26*c^2*x^2-9)*x
^5*arccosh(c*x)^2*c^6-280/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(8*c^6*x^6-25*c
^4*x^4+26*c^2*x^2-9)*x^5*arccosh(c*x)*c^6+56*b^2*(-d*(c^2*x^2-1))^(1/2)/d
^3/(8*c^6*x^6-25*c^4*x^4+26*c^2*x^2-9)*x^3*arccosh(c*x)^2*c^4+48*b^2*(-d*(c
^2*x^2-1))^(1/2)/d^3/(8*c^6*x^6-25*c^4*x^4+26*c^2*x^2-9)*x^3*arccosh(c*x)*
c^4-44*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(8*c^6*x^6-25*c^4*x^4+26*c^2*x^2-9)*
x*arccosh(c*x)^2*c^2-8*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(8*c^6*x^6-25*c^4*x
^4+26*c^2*x^2-9)*x*arccosh(c*x)*c^2-3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(8*c^6
*x^6-25*c^4*x^4+26*c^2*x^2-9)*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c-64/3*b^2*(-d*(
c^2*x^2-1))^(1/2)/d^3/(8*c^6*x^6-25*c^4*x^4+26*c^2*x^2-9)*x^9*arccosh(c*x)
*c^10-1/3*a*b*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(16*(c*x-
1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)*c^4*x^4+16*arccosh(c*x)*c^5*x^5-6*ln(1
+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^5*c^5-10*ln((c*x+(c*x-1)^(1/2)*(c*
x+1)^(1/2))^2-1)*x^5*c^5-24*(c*x+1)^(1/2)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x
^2-32*c^3*x^3*arccosh(c*x)+12*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*x^
3*c^3+20*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)*x^3*c^3-c^3*x^3+6*arcco
sh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+16*c*x*arccosh(c*x)-6*ln(1+(c*x+(c*x-1
)^(1/2)*(c*x+1)^(1/2))^2)*x*c-10*ln((c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2-1)
*x*c+c*x)/d^3/(c^6*x^6-3*c^4*x^4+3*c^2*x^2-1)/x-88/3*b^2*(-d*(c^2*x^2-1))^(
1/2)/d^3/(8*c^6*x^6-25*c^4*x^4+26*c^2*x^2-9)*x^5*(c*x-1)*(c*x+1)*c^6+8...

```

### 3.221.5 Fracas [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^6*d^3*x^8 - 3*c^4*d^3*x^6 + 3*c^2*d^3*x^4 - d^3*x^2), x)`

**3.221.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))**2/x**2/(-c**2*d*x**2+d)**(5/2),x)`output `Timed out`**3.221.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{5}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`output `1/3*a^2*(8*c^2*x/(sqrt(-c^2*d*x^2 + d)*d^2) + 4*c^2*x/((-c^2*d*x^2 + d)^(3/2)*d) - 3/((-c^2*d*x^2 + d)^(3/2)*d*x)) + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/((-c^2*d*x^2 + d)^(5/2)*x^2) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/((-c^2*d*x^2 + d)^(5/2)*x^2), x)`**3.221.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{5}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(5/2)*x^2), x)`

**3.221.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^2*(d - c^2*d*x^2)^(5/2)),x)`output `int((a + b*acosh(c*x))^2/(x^2*(d - c^2*d*x^2)^(5/2)), x)`

$$3.222 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$$

3.222.1 Optimal result . . . . .	1982
3.222.2 Mathematica [B] (warning: unable to verify) . . . . .	1983
3.222.3 Rubi [F] . . . . .	1984
3.222.4 Maple [F] . . . . .	1994
3.222.5 Fracas [F] . . . . .	1994
3.222.6 Sympy [F(-1)] . . . . .	1995
3.222.7 Maxima [F] . . . . .	1995
3.222.8 Giac [F] . . . . .	1995
3.222.9 Mupad [F(-1)] . . . . .	1996

**3.222.1 Optimal result**

Integrand size = 29, antiderivative size = 796

$$\begin{aligned}
& \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx = -\frac{b^2 c^2}{3d^2 \sqrt{d - c^2 dx^2}} \\
& + \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx))}{d^2 x (1 - c^2 x^2) \sqrt{d - c^2 dx^2}} \\
& - \frac{2bc^3 x \sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx))}{3d^2 (1 - c^2 x^2) \sqrt{d - c^2 dx^2}} + \frac{5c^2 (a + \operatorname{barccosh}(cx))^2}{6d (d - c^2 dx^2)^{3/2}} \\
& - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2 (d - c^2 dx^2)^{3/2}} + \frac{5c^2 (a + \operatorname{barccosh}(cx))^2}{2d^2 \sqrt{d - c^2 dx^2}} \\
& + \frac{5c^2 \sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx))^2 \arctan(e^{\operatorname{arccosh}(cx)})}{d^2 \sqrt{d - c^2 dx^2}} \\
& - \frac{b^2 c^2 \sqrt{-1 + cx}\sqrt{1 + cx} \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d^2 \sqrt{d - c^2 dx^2}} \\
& + \frac{26bc^2 \sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx)) \operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})}{3d^2 \sqrt{d - c^2 dx^2}} \\
& + \frac{13b^2 c^2 \sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)})}{3d^2 \sqrt{d - c^2 dx^2}} \\
& - \frac{5ibc^2 \sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(cx)})}{d^2 \sqrt{d - c^2 dx^2}} \\
& + \frac{5ibc^2 \sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(cx)})}{d^2 \sqrt{d - c^2 dx^2}} \\
& - \frac{13b^2 c^2 \sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)})}{3d^2 \sqrt{d - c^2 dx^2}} \\
& + \frac{5ib^2 c^2 \sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(cx)})}{d^2 \sqrt{d - c^2 dx^2}} \\
& - \frac{5ib^2 c^2 \sqrt{-1 + cx}\sqrt{1 + cx} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(cx)})}{d^2 \sqrt{d - c^2 dx^2}}
\end{aligned}$$

output

```

5/6*c^2*(a+b*arccosh(c*x))^2/d/(-c^2*d*x^2+d)^(3/2)-1/2*(a+b*arccosh(c*x))
^2/d/x^2/(-c^2*d*x^2+d)^(3/2)-1/3*b^2*c^2/d^2/(-c^2*d*x^2+d)^(1/2)+5/2*c^2
*(a+b*arccosh(c*x))^2/d^2/(-c^2*d*x^2+d)^(1/2)+b*c*(a+b*arccosh(c*x))*(c*x
-1)^(1/2)*(c*x+1)^(1/2)/d^2/x/(-c^2*x^2+1)/(-c^2*d*x^2+d)^(1/2)-2/3*b*c^3*
x*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*x^2+1)/(-c^2*d*
x^2+d)^(1/2)+5*c^2*(a+b*arccosh(c*x))^2*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(
1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2)-b^2*c^2*arctan(
(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d
)^(1/2)+26/3*b*c^2*(a+b*arccosh(c*x))*arctanh(c*x+(c*x-1)^(1/2)*(c*x+1)^(1
/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2)+13/3*b^2*c^2*pol
ylog(2,-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(
-c^2*d*x^2+d)^(1/2)-5*I*b*c^2*(a+b*arccosh(c*x))*polylog(2,-I*(c*x+(c*x-1)
^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2
)+5*I*b*c^2*(a+b*arccosh(c*x))*polylog(2,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2
)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2)-13/3*b^2*c^2*poly
log(2,c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c
^2*d*x^2+d)^(1/2)+5*I*b^2*c^2*polylog(3,-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2
)))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/(-c^2*d*x^2+d)^(1/2)-5*I*b^2*c^2*polyl
og(3,I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^2/
(-c^2*d*x^2+d)^(1/2)

```

### 3.222.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 5532 vs.  $2(796) = 1592$ .

Time = 65.92 (sec) , antiderivative size = 5532, normalized size of antiderivative = 6.95

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx = \text{Result too large to show}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^3*(d - c^2*d*x^2)^(5/2)),x]`

output `Result too large to show`



## 3.222.3 Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6347} \\
 & -\frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} + \frac{5}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{6327} \\
 & -\frac{bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^2(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} + \frac{5}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{6347} \\
 & -\frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - bc \int \frac{1}{x(cx-1)^{3/2}(cx+1)^{3/2}} dx - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} \right)}{d^2\sqrt{d-c^2dx^2}} + \\
 & \quad \frac{5}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{115} \\
 & -\frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - bc \left( -\frac{\int \frac{c}{x\sqrt{cx-1}\sqrt{cx+1}} dx}{c} - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} \right)}{d^2\sqrt{d-c^2dx^2}} + \\
 & \quad \frac{5}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{27} \\
 & -\frac{bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2} dx - bc \left( -\int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} \right)}{d^2\sqrt{d-c^2dx^2}} + \\
 & \quad \frac{5}{2}c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x (d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{2dx^2 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{103}
 \end{aligned}$$

---

3.222.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\int\frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2}dx - bc\left(-c\int\frac{1}{(cx-1)(cx+1)c+c}d(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}}\right) - \frac{a+b}{x}\right)}{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx - \frac{d^2\sqrt{d-c^2dx^2}}{2dx^2(d-c^2dx^2)^{3/2}}}$$

↓ 218

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\int\frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)^2}dx - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc\left(-\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx - \frac{d^2\sqrt{d-c^2dx^2}}{2dx^2(d-c^2dx^2)^{3/2}}}$$

↓ 6316

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\left(\frac{1}{2}\int\frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2}dx + \frac{1}{2}bc\int\frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}}dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)}\right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)}\right)}{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx - \frac{d^2\sqrt{d-c^2dx^2}}{2dx^2(d-c^2dx^2)^{3/2}}}$$

↓ 83

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\left(\frac{1}{2}\int\frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2}dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} - bc\left(-\arctan(\sqrt{cx-1}\sqrt{cx+1}) - \frac{1}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx - \frac{d^2\sqrt{d-c^2dx^2}}{2dx^2(d-c^2dx^2)^{3/2}}}$$

↓ 6318

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\left(-\frac{\int\frac{a+\operatorname{barccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)}d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right) - \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)}\right)}{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx - \frac{d^2\sqrt{d-c^2dx^2}}{2dx^2(d-c^2dx^2)^{3/2}}}$$

↓ 3042

---

3.222.  $\int\frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}}dx$

$$\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\left(-\frac{\int i(a+\operatorname{barccosh}(cx))\csc(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)}{2c}+\frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)}-\frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2\sqrt{d-c^2dx^2}}-\frac{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}}{26}}{d^2\sqrt{d-c^2dx^2}}-\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\left(-\frac{\int i(a+\operatorname{barccosh}(cx))\csc(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)}{2c}+\frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)}-\frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}}\right)}{d^2\sqrt{d-c^2dx^2}}-\frac{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}}{4670}}{d^2\sqrt{d-c^2dx^2}}-\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\left(-\frac{i\left(\int\log(1-e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)-\int\log(1+e^{\operatorname{arccosh}(cx)})d\operatorname{arccosh}(cx)+2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})\right)}{2c}\right)}{2715}}{d^2\sqrt{d-c^2dx^2}}-\frac{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}}{2838}}{d^2\sqrt{d-c^2dx^2}}-\frac{bc\sqrt{cx-1}\sqrt{cx+1}\left(3c^2\left(-\frac{i\left(\int e^{-\operatorname{arccosh}(cx)}\log(1-e^{\operatorname{arccosh}(cx)})de^{\operatorname{arccosh}(cx)}-\int e^{-\operatorname{arccosh}(cx)}\log(1+e^{\operatorname{arccosh}(cx)})de^{\operatorname{arccosh}(cx)}\right)}{2c}\right)}{6351}}{d^2\sqrt{d-c^2dx^2}}-\frac{\frac{5}{2}c^2\int\frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{5/2}}dx-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}}{2c}\left(-\frac{i\left(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+i\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-i\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})\right)}{2c}\right)}{d^2\sqrt{d-c^2dx^2}}-\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}}{6351}}{d^2\sqrt{d-c^2dx^2}}$$

3.222.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \right) -$$

$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right. \right.$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 6304

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{(1-c^2x^2)} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \right) -$$

$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right. \right.$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 6316

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{1}{2}bc \int \frac{x}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} \right) -$$

$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right. \right.$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 83

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{2} \int \frac{a+\operatorname{barccosh}(cx)}{1-c^2x^2} dx + \frac{x(a+\operatorname{barccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+\operatorname{barccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} \right) -$$

$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+\operatorname{barccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right. \right.$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 6318

---

3.222.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}}(cx+1)} d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+b\operatorname{arccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^3} dx}{d} \right)$$

$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) \right)$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 3042

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{\int i(a+b\operatorname{arccosh}(cx))\operatorname{csc}(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+b\operatorname{arccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right)$$

$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) \right)$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 26

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i\int (a+b\operatorname{arccosh}(cx))\operatorname{csc}(i\operatorname{arccosh}(cx))d\operatorname{arccosh}(cx)}{2c} + \frac{x(a+b\operatorname{arccosh}(cx))}{2(1-c^2x^2)} - \frac{b}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right)$$

$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) \right)$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 4670

---

3.222.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int \log(1-e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) - ib \int \log(1+e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) + 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) d\operatorname{arccosh}(cx) \right)}{2c}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right)$$


---


$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right) \right)$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 2715

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( ib \int e^{-\operatorname{arccosh}(cx)} \log(1-e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} - ib \int e^{-\operatorname{arccosh}(cx)} \log(1+e^{\operatorname{arccosh}(cx)}) de^{\operatorname{arccosh}(cx)} \right)}{2c} \right)}{3d^2\sqrt{d-c^2dx^2}} \right)$$


---


$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right) \right)$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 2838

$$\frac{5}{2}c^2 \left( \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x(d-c^2dx^2)^{3/2}} dx}{d} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right)}{3d^2\sqrt{d-c^2dx^2}} \right)$$


---


$$bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i \left( 2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)}) (a+b\operatorname{arccosh}(cx)) + ib \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(cx)}) - ib \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(cx)}) \right)}{2c} \right) \right)$$


---


$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 6351

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{2c} \right)}{d^2\sqrt{d-c^2dx^2}} \right)$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 25

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{(1-cx)(cx+1)} dx}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{2c} \right)}{d^2\sqrt{d-c^2dx^2}} \right)$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 6304

$$\frac{5}{2}c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -\frac{i(2\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)})}{2c} \right)}{d^2\sqrt{d-c^2dx^2}} \right)$$

$$\frac{(a+b\operatorname{arccosh}(cx))^2}{2dx^2(d-c^2dx^2)^{3/2}}$$

↓ 6318

---

3.222.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$

$$\frac{5}{2}c^2 \left( \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{\frac{cx-1}{cx+1}(cx+1)}} d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1}}{d} \right) \\ bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) \right)$$

$$\frac{(a + b\operatorname{arccosh}(cx))^2}{2dx^2 (d - c^2dx^2)^{3/2}}$$

↓ 3042

$$\frac{5}{2}c^2 \left( \frac{\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x\sqrt{d-c^2dx^2}} dx}{d} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int i(a+b\operatorname{arccosh}(cx)) \operatorname{csc}(i\operatorname{arccosh}(cx)) d\operatorname{arccosh}(cx)}{d\sqrt{d-c^2dx^2}} + \frac{(a+b\operatorname{arccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1}}{d} \right) \\ bc\sqrt{cx-1}\sqrt{cx+1} \left( 3c^2 \left( -\frac{i(2i\operatorname{arctanh}(e^{\operatorname{arccosh}(cx)})(a+b\operatorname{arccosh}(cx))+ib\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(cx)})-ib\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(cx)}))}{2c} \right) \right)$$

$$\frac{(a + b\operatorname{arccosh}(cx))^2}{2dx^2 (d - c^2dx^2)^{3/2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^3*(d - c^2*d*x^2)^(5/2)), x]`

output `$Aborted`

### 3.222.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

---

3.222.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^3(d-c^2dx^2)^{5/2}} dx$



- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`
- rule 115 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]`
- rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`
- rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(f*x)^(m + 1))*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] || EqQ[n, 1])`

### 3.222.4 Maple [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 (-c^2 dx^2 + d)^{\frac{5}{2}}} dx$$

input `int((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(5/2),x)`

output `int((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(5/2),x)`

### 3.222.5 Fracas [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{\frac{5}{2}}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{\frac{5}{2}} x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^6*d^3*x^9 - 3*c^4*d^3*x^7 + 3*c^2*d^3*x^5 - d^3*x^3), x)`

**3.222.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))**2/x**3/(-c**2*d*x**2+d)**(5/2),x)`

output `Timed out`

**3.222.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")`

output `-1/6*a^2*(15*c^2*log(2*sqrt(-c^2*d*x^2 + d)*sqrt(d)/abs(x) + 2*d/abs(x))/d^(5/2) - 15*c^2/(sqrt(-c^2*d*x^2 + d)*d^2) - 5*c^2/((-c^2*d*x^2 + d)^(3/2)*d) + 3/((-c^2*d*x^2 + d)^(3/2)*d*x^2) + integrate(b^2*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))^2/((-c^2*d*x^2 + d)^(5/2)*x^3) + 2*a*b*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/((-c^2*d*x^2 + d)^(5/2)*x^3), x)`

**3.222.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x^3} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^3/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(5/2)*x^3), x)`

**3.222.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^3 (d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^3*(d - c^2*d*x^2)^(5/2)),x)`output `int((a + b*acosh(c*x))^2/(x^3*(d - c^2*d*x^2)^(5/2)), x)`

**3.223** 
$$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$$

3.223.1 Optimal result . . . . .	1997
3.223.2 Mathematica [A] (warning: unable to verify) . . . . .	1998
3.223.3 Rubi [C] (verified) . . . . .	1999
3.223.4 Maple [B] (verified) . . . . .	2011
3.223.5 Fricas [F] . . . . .	2012
3.223.6 Sympy [F(-1)] . . . . .	2013
3.223.7 Maxima [F] . . . . .	2013
3.223.8 Giac [F] . . . . .	2013
3.223.9 Mupad [F(-1)] . . . . .	2014

**3.223.1 Optimal result**

Integrand size = 29, antiderivative size = 562

$$\begin{aligned} \int \frac{(a + \operatorname{arccosh}(cx))^2}{x^4(d - c^2dx^2)^{5/2}} dx &= \frac{b^2c^2}{3d^2x\sqrt{d - c^2dx^2}} - \frac{2b^2c^4x}{3d^2\sqrt{d - c^2dx^2}} \\ &+ \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))}{3d^2x^2(1 - c^2x^2)\sqrt{d - c^2dx^2}} - \frac{(a + \operatorname{arccosh}(cx))^2}{3dx^3(d - c^2dx^2)^{3/2}} \\ &- \frac{2c^2(a + \operatorname{arccosh}(cx))^2}{dx(d - c^2dx^2)^{3/2}} + \frac{8c^4x(a + \operatorname{arccosh}(cx))^2}{3d(d - c^2dx^2)^{3/2}} \\ &+ \frac{16c^4x(a + \operatorname{arccosh}(cx))^2}{3d^2\sqrt{d - c^2dx^2}} + \frac{16c^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^2}{3d^2\sqrt{d - c^2dx^2}} \\ &- \frac{32bc^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)})}{3d^2\sqrt{d - c^2dx^2}} \\ &- \frac{32bc^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))\log(1 - e^{2\operatorname{arccosh}(cx)})}{3d^2\sqrt{d - c^2dx^2}} \\ &- \frac{8b^2c^3\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)})}{3d^2\sqrt{d - c^2dx^2}} \\ &- \frac{8b^2c^3\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(cx)})}{3d^2\sqrt{d - c^2dx^2}} \end{aligned}$$

output 
$$-1/3*(a+b*\operatorname{arccosh}(c*x))^2/d/x^3/(-c^2*d*x^2+d)^{(3/2)}-2*c^2*(a+b*\operatorname{arccosh}(c*x))^2/d/x/(-c^2*d*x^2+d)^{(3/2)}+8/3*c^4*x*(a+b*\operatorname{arccosh}(c*x))^2/d/(-c^2*d*x^2+d)^{(3/2)}+1/3*b^2*c^2/d^2/x/(-c^2*d*x^2+d)^{(1/2)}-2/3*b^2*c^4*x/d^2/(-c^2*d*x^2+d)^{(1/2)}+16/3*c^4*x*(a+b*\operatorname{arccosh}(c*x))^2/d^2/(-c^2*d*x^2+d)^{(1/2)}+1/3*b*c*(a+b*\operatorname{arccosh}(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/x^2/(-c^2*x^2+1)/(-c^2*d*x^2+d)^{(1/2)}+16/3*c^3*(a+b*\operatorname{arccosh}(c*x))^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-32/3*b*c^3*(a+b*\operatorname{arccosh}(c*x))*\operatorname{arctanh}((c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-32/3*b*c^3*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-8/3*b^2*c^3*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)}-8/3*b^2*c^3*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/d^2/(-c^2*d*x^2+d)^{(1/2)})$$

### 3.223.2 Mathematica [A] (warning: unable to verify)

Time = 2.80 (sec) , antiderivative size = 534, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx = \frac{a^2(1+6c^2x^2-24c^4x^4+16c^6x^6)}{x^3(-1+c^2x^2)} + abc^3 \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \left( \frac{1}{c^2x^2} + \frac{1}{1-c^2x^2} + \frac{2\left(\frac{-1+cx}{1+cx}\right)^{3/2}}{1+c^2x^2} \right)$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(x^4*(d - c^2*d*x^2)^(5/2)),x]`

output 
$$\left( \frac{a^2(1+6c^2x^2-24c^4x^4+16c^6x^6)}{x^3(-1+c^2x^2)} + a*b*c^3*\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]*(1+c*x)*\left(\frac{1}{c^2*x^2} + (1-c^2*x^2)^{-1} + (2*((-1+c*x)/(1+c*x))^{3/2}*(1+6*c^2*x^2-24*c^4*x^4+16*c^6*x^6)*\operatorname{ArcCosh}[c*x])\right) / (c^3*x^3*(-1+c*x)^3) - 16*\operatorname{Log}[c*x] - 16*\operatorname{Log}\left[\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]*(1+c*x)\right] + b^2*c^3*\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]*(1+c*x)*\left(\frac{c*x*\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]}{1-c*x} - \left(\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]\right)*(1+c*x)\right) / (c*x) + \operatorname{ArcCosh}[c*x] / (c^2*x^2) + \operatorname{ArcCosh}[c*x] / (1-c^2*x^2) - 16*\operatorname{ArcCosh}[c*x]^2 - (c*x*\operatorname{ArcCosh}[c*x]^2) / (((-1+c*x)/(1+c*x))^{3/2}*(1+c*x)^3) + (8*c*x*\operatorname{ArcCosh}[c*x]^2) / (\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]*(1+c*x)) + (\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]*(1+c*x)*\operatorname{ArcCosh}[c*x]^2) / (c^3*x^3) + (8*\operatorname{Sqrt}\left[\frac{-1+c*x}{1+c*x}\right]*(1+c*x)*\operatorname{ArcCosh}[c*x]^2) / (c*x) - 16*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1-E^{-2*\operatorname{ArcCosh}[c*x]}] - 16*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1+E^{-2*\operatorname{ArcCosh}[c*x]}] + 8*\operatorname{PolyLog}[2,-E^{-2*\operatorname{ArcCosh}[c*x]}] + 8*\operatorname{PolyLog}[2,E^{-2*\operatorname{ArcCosh}[c*x]}]) / (3*d^2*\operatorname{Sqrt}[d-c^2*d*x^2])$$

3.223. 
$$\int \frac{(a+b\operatorname{arccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$$

**3.223.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 6.30 (sec) , antiderivative size = 750, normalized size of antiderivative = 1.33, number of steps used = 29, number of rules used = 28,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.966$ , Rules used = {6347, 6327, 6347, 114, 27, 41, 6316, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838, 6329, 41, 6351, 41, 6331, 5984, 3042, 26, 4670, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6347} \\
 & -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^3(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + 2c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{6327} \\
 & -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x^3(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + 2c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{x^2 (d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{6347} \\
 & \frac{2c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{dx (d - c^2 dx^2)^{3/2}} \right) -}{3d^2\sqrt{d-c^2dx^2}} \\
 & \quad \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{1}{2}bc \int \frac{1}{x^2(cx-1)^{3/2}(cx+1)^{3/2}} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} \right)}{(a + \operatorname{barccosh}(cx))^2} \\
 & \quad \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3 (d - c^2 dx^2)^{3/2}} \\
 & \quad \downarrow \text{114} \\
 & \frac{2c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \int \frac{(a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx - \frac{(a + \operatorname{barccosh}(cx))^2}{dx (d - c^2 dx^2)^{3/2}} \right) -}{3d^2\sqrt{d-c^2dx^2}} \\
 & \quad \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{1}{2}bc \left( \int \frac{2c^2}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} \right)}{(a + \operatorname{barccosh}(cx))^2} \\
 & \quad \frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3 (d - c^2 dx^2)^{3/2}}
 \end{aligned}$$

---

3.223.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$



↓ 27

$$\frac{2c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \right) - 2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{1}{2}bc \left( 2c^2 \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 3dx^3 (d-c^2dx^2)^{3/2}}$$

↓ 41

$$\frac{2c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-cx)^2(cx+1)^2} dx}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{5/2}} dx - \frac{(a+\operatorname{barccosh}(cx))^2}{dx(d-c^2dx^2)^{3/2}} \right) - 2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 3dx^3 (d-c^2dx^2)^{3/2}}$$

↓ 6316

$$\frac{2c^2 \left( 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \int \frac{(a+\operatorname{barccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx}{3d} + \frac{x(a+\operatorname{barccosh}(cx))^2}{3d(d-c^2dx^2)^{3/2}} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 3dx^3 (d-c^2dx^2)^{3/2}} \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 3dx^3 (d-c^2dx^2)^{3/2}}$$

↓ 6314

$$\frac{2c^2 \left( 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-cx)^2(cx+1)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right)}{3d} \right) - \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 3dx^3 (d-c^2dx^2)^{3/2}} \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 3dx^3 (d-c^2dx^2)^{3/2}}$$

---

3.223.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$

↓ 6327

$$2c^2 \left( 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{1-c^2x^2} dx}{d\sqrt{d-c^2dx^2}} + \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right)}{3d} \right) \right. \\ \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}} \right)$$

↓ 6328

$$2c^2 \left( 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} - \frac{2b\sqrt{cx-1}\sqrt{cx+1} \int \frac{cx(a+\operatorname{barccosh}(cx))}{\sqrt{\frac{cx-1}{cx+1}(cx+1)}} da}{cd\sqrt{d-c^2dx^2}} \right)}{3d} \right) \right. \\ \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2}} \right. \\ \left. \frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}} \right)$$

↓ 3042

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2}} + \\ 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2} dx}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} \right)}{3d} \right) \right. \\ \left. \frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}} \right)$$

↓ 26

---

3.223.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(2c^2\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx-\frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)}-\frac{1}{2}bc\left(\frac{1}{x\sqrt{cx-1}\sqrt{cx+1}}-\frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{3d^2\sqrt{d-c^2dx^2}}+$$

$$2c^2\left(-\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx}{d^2\sqrt{d-c^2dx^2}}+4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2}dx}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}}\right)^2}{3dx^3(d-c^2dx^2)^{3/2}}\right)\right)$$

↓ 4199

$$2c^2\left(4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2}dx}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(2i\int-\frac{e^{2\operatorname{arccosh}(cx)}(a+i)}{1-e^{2\operatorname{arccosh}(cx)}}\right)}{3d}\right)^2}{3d}\right)\right)$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(2c^2\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx-\frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)}-\frac{1}{2}bc\left(\frac{1}{x\sqrt{cx-1}\sqrt{cx+1}}-\frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{3d^2\sqrt{d-c^2dx^2}}$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 25

$$2c^2\left(4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2}dx}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\int\frac{e^{2\operatorname{arccosh}(cx)}(a+i)}{1-e^{2\operatorname{arccosh}(cx)}}\right)}{3d}\right)^2}{3d}\right)\right)$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(2c^2\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx-\frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)}-\frac{1}{2}bc\left(\frac{1}{x\sqrt{cx-1}\sqrt{cx+1}}-\frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{3d^2\sqrt{d-c^2dx^2}}$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 2620

---

3.223.  $\int\frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}}dx$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(2c^2\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx-\frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)}-\frac{1}{2}bc\left(\frac{1}{x\sqrt{cx-1}\sqrt{cx+1}}-\frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{3d^2\sqrt{d-c^2dx^2}}+$$

$$2c^2\left(-\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx}{d^2\sqrt{d-c^2dx^2}}+4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2}dx}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}}\right)^2}{(a+\operatorname{barccosh}(cx))^2}\right)\right)$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 2715

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(2c^2\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx-\frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)}-\frac{1}{2}bc\left(\frac{1}{x\sqrt{cx-1}\sqrt{cx+1}}-\frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{3d^2\sqrt{d-c^2dx^2}}+$$

$$2c^2\left(-\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx}{d^2\sqrt{d-c^2dx^2}}+4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2}dx}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))}{d\sqrt{d-c^2dx^2}}\right)^2}{(a+\operatorname{barccosh}(cx))^2}\right)\right)$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 2838

$$2c^2\left(4c^2\left(\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\int\frac{x(a+\operatorname{barccosh}(cx))}{(1-c^2x^2)^2}dx}{3d^2\sqrt{d-c^2dx^2}}+\frac{2\left(\frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}}+\frac{2ib\sqrt{cx-1}\sqrt{cx+1}\left(-2i\left(-\frac{1}{2}\log(1-e^{2\operatorname{arccosh}(cx)})\right)\right)}{2}\right)^2}{(a+\operatorname{barccosh}(cx))^2}\right)\right)$$

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1}\left(2c^2\int\frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2}dx-\frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)}-\frac{1}{2}bc\left(\frac{1}{x\sqrt{cx-1}\sqrt{cx+1}}-\frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}}\right)\right)}{3d^2\sqrt{d-c^2dx^2}}$$

$$\frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 6329

---

3.223.  $\int\frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}}dx$

$$2c^2 \left( 4c^2 \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{b \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx}{2c} + \frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) \\ \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2} \\ \frac{3d^2\sqrt{d-c^2dx^2}}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 41

$$2c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)} - \frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) \\ \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)^2} dx - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2} \\ \frac{3d^2\sqrt{d-c^2dx^2}}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 6351

$$2c^2 \left( -\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2}bc \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} \right)}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{a+\operatorname{barccosh}(cx)}{2c^2(1-c^2x^2)} - \frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} + \frac{2 \left( \frac{x(a+\operatorname{barccosh}(cx))^2}{d\sqrt{d-c^2dx^2}} + \frac{2ib\sqrt{cx-1}\sqrt{cx+1}}{2c^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \right) \\ \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{1}{2}bc \int \frac{1}{(cx-1)^{3/2}(cx+1)^{3/2}} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} \right) - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} - \frac{2c^2x}{\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2} \\ \frac{3d^2\sqrt{d-c^2dx^2}}{3dx^3(d-c^2dx^2)^{3/2}}$$

↓ 41

---

3.223.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$

$$\begin{aligned}
 & 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2} \right. \right. \\
 & \left. \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \left( \int \frac{a+\operatorname{barccosh}(cx)}{x(1-c^2x^2)} dx + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} - \frac{1}{2}bc \left( \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} \right) \right)}{3d^2\sqrt{d-c^2dx^2}} \right. \right. \\
 & \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}} \right) \right.
 \end{aligned}$$

↓ 6331

$$\begin{aligned}
 & 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( - \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( - \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right. \right. \\
 & \left. \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \left( - \int \frac{a+\operatorname{barccosh}(cx)}{cx\sqrt{\frac{cx-1}{cx+1}}(cx+1)} \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \right. \right. \\
 & \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}} \right) \right.
 \end{aligned}$$

↓ 5984

$$\begin{aligned}
 & 2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2 \int (a+\operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2\sqrt{d-c^2dx^2}} + 4c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2 \int (a+\operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d-c^2dx^2}} \right. \right. \\
 & \left. \left. \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \left( -2 \int (a+\operatorname{barccosh}(cx)) \operatorname{csch}(2\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right) - \frac{a+\operatorname{barccosh}(cx)}{2x^2(1-c^2x^2)} \right)}{3d^2\sqrt{d-c^2dx^2}} \right. \right. \\
 & \left. \left. \frac{(a+\operatorname{barccosh}(cx))^2}{3dx^3(d-c^2dx^2)^{3/2}} \right) \right.
 \end{aligned}$$

↓ 3042

---

3.223.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$

$$2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2 \int i(a + \operatorname{barccosh}(cx)) \csc(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2\sqrt{d-c^2dx^2}} \right)$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \left( -2 \int i(a + \operatorname{barccosh}(cx)) \csc(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$


---


$$\frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3 (d - c^2dx^2)^{3/2}}$$

↓ 26

$$2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2i \int (a + \operatorname{barccosh}(cx)) \csc(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right)}{d^2\sqrt{d-c^2dx^2}} \right)$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \left( -2i \int (a + \operatorname{barccosh}(cx)) \csc(2i\operatorname{arccosh}(cx)) \operatorname{darccosh}(cx) + \frac{a+\operatorname{barccosh}(cx)}{2(1-c^2x^2)} - \frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} \right) \right)$$


---


$$\frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3 (d - c^2dx^2)^{3/2}}$$

↓ 4670

$$2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2i \left( \frac{1}{2} ib \int \log(1 - e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} ib \int \log(1 + e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) \right) \right)}{d^2\sqrt{d-c^2dx^2}} \right)$$


---


$$2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \left( -2i \left( \frac{1}{2} ib \int \log(1 - e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) - \frac{1}{2} ib \int \log(1 + e^{2\operatorname{arccosh}(cx)}) \operatorname{darccosh}(cx) \right) \right) \right)$$


---


$$\frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3 (d - c^2dx^2)^{3/2}}$$

↓ 2715

---

3.223.  $\int \frac{(a+\operatorname{barccosh}(cx))^2}{x^4(d-c^2dx^2)^{5/2}} dx$

$$2 \left( 4 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{3d(d - c^2 dx^2)^{3/2}} + \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( \frac{a + \operatorname{barccosh}(cx)}{2c^2(1-c^2 x^2)} - \frac{bx}{2c\sqrt{cx-1}\sqrt{cx+1}} \right)}{3d^2\sqrt{d - c^2 dx^2}} + \frac{2 \left( \frac{x(a + \operatorname{barccosh}(cx))^2}{d\sqrt{d - c^2 dx^2}} + \frac{2bx}{d\sqrt{d - c^2 dx^2}} \right)}{3d^2\sqrt{d - c^2 dx^2}} \right) \right. \\ \left. + \frac{2b\sqrt{cx-1}\sqrt{cx+1} \left( 2 \left( -\frac{bcx}{2\sqrt{cx-1}\sqrt{cx+1}} + \frac{a + \operatorname{barccosh}(cx)}{2(1-c^2 x^2)} \right) - 2i(i(a + \operatorname{barccosh}(cx))\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) + \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} dx) \right)}{3d^2\sqrt{d - c^2 dx^2}} \right)$$

$$\frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3(d - c^2 dx^2)^{3/2}}$$

↓ 2838

$$\frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( 2c^2 \left( -2i(i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4}ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} dx) \right) \right)}{3d^2\sqrt{d - c^2 dx^2}}$$

$$2c^2 \left( \frac{2bc\sqrt{cx-1}\sqrt{cx+1} \left( -2i(i\operatorname{arctanh}(e^{2\operatorname{arccosh}(cx)}) (a + \operatorname{barccosh}(cx)) + \frac{1}{4}ib \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(cx)}) - \frac{1}{4}ib \int e^{-2\operatorname{arccosh}(cx)} dx) \right)}{d^2\sqrt{d - c^2 dx^2}} \right)$$

$$\frac{(a + \operatorname{barccosh}(cx))^2}{3dx^3(d - c^2 dx^2)^{3/2}}$$

input `Int[(a + b*ArcCosh[c*x])^2/(x^4*(d - c^2*d*x^2)^(5/2)),x]`



```

output -1/3*(a + b*ArcCosh[c*x])^2/(d*x^3*(d - c^2*d*x^2)^(3/2)) - (2*b*c*Sqrt[-1
+ c*x]*Sqrt[1 + c*x]*(-1/2*(b*c*(1/(x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) - (2*
c^2*x)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]))) - (a + b*ArcCosh[c*x])/(2*x^2*(1 -
c^2*x^2)) + 2*c^2*(-1/2*(b*c*x)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*A
rcCosh[c*x])/(2*(1 - c^2*x^2)) - (2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(
2*ArcCosh[c*x])] + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyL
og[2, E^(2*ArcCosh[c*x])])))/(3*d^2*Sqrt[d - c^2*d*x^2]) + 2*c^2*(-((a +
b*ArcCosh[c*x])^2/(d*x*(d - c^2*d*x^2)^(3/2))) - (2*b*c*Sqrt[-1 + c*x]*Sqr
t[1 + c*x]*(-1/2*(b*c*x)/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*ArcCosh[c
*x])/(2*(1 - c^2*x^2)) - (2*I)*(I*(a + b*ArcCosh[c*x])*ArcTanh[E^(2*ArcCos
h[c*x])] + (I/4)*b*PolyLog[2, -E^(2*ArcCosh[c*x])] - (I/4)*b*PolyLog[2, E^
(2*ArcCosh[c*x])])))/(d^2*Sqrt[d - c^2*d*x^2]) + 4*c^2*((x*(a + b*ArcCosh[
c*x])^2)/(3*d*(d - c^2*d*x^2)^(3/2)) + (2*b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]
*(-1/2*(b*x)/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (a + b*ArcCosh[c*x])/(2*c^
2*(1 - c^2*x^2)))/(3*d^2*Sqrt[d - c^2*d*x^2]) + (2*((x*(a + b*ArcCosh[c*x
])^2)/(d*Sqrt[d - c^2*d*x^2]) + ((2*I)*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(((
-1/2*I)*(a + b*ArcCosh[c*x])^2)/b - (2*I)*(-1/2*((a + b*ArcCosh[c*x])*Log[1
- E^(2*ArcCosh[c*x])])) - (b*PolyLog[2, E^(2*ArcCosh[c*x])])/4)))/(c*d*Sqr
t[d - c^2*d*x^2])))/(3*d)))

```

### 3.223.3.1 Defintions of rubi rules used

```

rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]

```

```

rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]

```

```

rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]

```

```

rule 41 Int[1/(((a_) + (b_.)*(x_))^(3/2)*((c_) + (d_.)*(x_))^(3/2)), x_Symbol] := S
imp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && Eq
Q[b*c + a*d, 0]

```

rule 114 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Simp[1/((m + 1)*(b*c - a*d)*(b*e - a*f)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])`

rule 2620 `Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 5984 `Int[Csch[(a_.) + (b_.)*(x_)]^(n_.)*((c_.) + (d_.)*(x_))^(m_.)*Sech[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Simp[2^n Int[(c + d*x)^m*Csch[2*a + 2*b*x]^n, x], x] /; FreeQ[{a, b, c, d}, x] && RationalQ[m] && IntegerQ[n]`

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 6331 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((x_)*((d_) + (e_.)*(x_)^2)),  
x_Symbol] := Simp[-d^(-1) Subst[Int[(a + b*x)^n/(Cosh[x]*Sinh[x]), x], x  
, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IG  
tQ[n, 0]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_.  
)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +  
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1  
))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp  
[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(  
f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^  
(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] &&  
GtQ[n, 0] && ILtQ[m, -1]`

rule 6351 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_.  
)*(x_)^2)^(p_), x_Symbol] := Simp[-(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a  
+ b*ArcCosh[c*x])^n/(2*d*f*(p + 1))), x] + (Simp[(m + 2*p + 3)/(2*d*(p + 1  
)) Int[(f*x)^m*(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[  
b*c*(n/(2*f*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[  
(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])  
(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] &  
& GtQ[n, 0] && LtQ[p, -1] && !GtQ[m, 1] && (IntegerQ[m] || IntegerQ[p] ||  
EqQ[n, 1])`

### 3.223.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 3744 vs.  $2(544) = 1088$ .

Time = 1.39 (sec) , antiderivative size = 3745, normalized size of antiderivative = 6.66

method	result	size
default	Expression too large to display	3745
parts	Expression too large to display	3745

input `int((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(5/2),x,method=_RETURNVERBOSE)`

output `a^2*(-1/3/d/x^3/(-c^2*d*x^2+d)^(3/2)+2*c^2*(-1/d/x/(-c^2*d*x^2+d)^(3/2)+4*c^2*(1/3/d*x/(-c^2*d*x^2+d)^(3/2)+2/3/d^2*x/(-c^2*d*x^2+d)^(1/2))))+128/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^5*(c*x-1)*(c*x+1)*c^8-8*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^9+256/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^9*(c*x-1)*(c*x+1)*arccosh(c*x)*c^12-640/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^7*(c*x-1)*(c*x+1)*arccosh(c*x)*c^10+64*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2*c^9+160*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^5*(c*x-1)*(c*x+1)*arccosh(c*x)*c^8-128*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2*c^7-80/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^3*(c*x-1)*(c*x+1)*arccosh(c*x)*c^6+176/3*b^2*(-d*(c^2*x^2-1))^(1/2)/d^3/(12*c^8*x^8-36*c^6*x^6+35*c^4*x^4-10*c^2*x^2-1)*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*arccosh(c*x)^2*c^5+8/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/(c^2*x^2-1)*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)*c^3+16/3*b^2*(-d*(c^2*x^2-1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/(c^2*x^2-1)*polylog(2,c*x+(c...`

### 3.223.5 Fracas [F]

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x^4} dx$$

input `integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(c^6*d^3*x^10 - 3*c^4*d^3*x^8 + 3*c^2*d^3*x^6 - d^3*x^4), x)`

**3.223.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

```
input integrate((a+b*acosh(c*x))**2/x**4/(-c**2*d*x**2+d)**(5/2),x)
```

```
output Timed out
```

**3.223.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x^4} dx$$

```
input integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(5/2),x, algorithm="maxima")
```

```
output 1/3*a*b*c*(8*c^2*sqrt(-d)*log(c*x + 1)/d^3 + 8*c^2*sqrt(-d)*log(c*x - 1)/d^3 + 16*c^2*sqrt(-d)*log(x)/d^3 + sqrt(-d)/(c^2*d^3*x^4 - d^3*x^2)) + 2/3*(16*c^4*x/(sqrt(-c^2*d*x^2 + d)*d^2) + 8*c^4*x/((-c^2*d*x^2 + d)^(3/2)*d) - 6*c^2/((-c^2*d*x^2 + d)^(3/2)*d*x) - 1/((-c^2*d*x^2 + d)^(3/2)*d*x^3))*a*b*arccosh(c*x) + 1/3*(16*c^4*x/(sqrt(-c^2*d*x^2 + d)*d^2) + 8*c^4*x/((-c^2*d*x^2 + d)^(3/2)*d) - 6*c^2/((-c^2*d*x^2 + d)^(3/2)*d*x) - 1/((-c^2*d*x^2 + d)^(3/2)*d*x^3))*a^2 + b^2*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))^2/((-c^2*d*x^2 + d)^(5/2)*x^4), x)
```

**3.223.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(-c^2 dx^2 + d)^{5/2} x^4} dx$$

```
input integrate((a+b*arccosh(c*x))^2/x^4/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")
```

```
output integrate((b*arccosh(c*x) + a)^2/((-c^2*d*x^2 + d)^(5/2)*x^4), x)
```

---

3.223.  $\int \frac{(a + \operatorname{barccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx$

**3.223.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{x^4 (d - c^2 dx^2)^{5/2}} dx$$

input `int((a + b*acosh(c*x))^2/(x^4*(d - c^2*d*x^2)^(5/2)),x)`output `int((a + b*acosh(c*x))^2/(x^4*(d - c^2*d*x^2)^(5/2)), x)`

### 3.224 $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{7/2}} dx$

3.224.1 Optimal result	2015
3.224.2 Mathematica [A] (warning: unable to verify)	2016
3.224.3 Rubi [C] (verified)	2016
3.224.4 Maple [B] (verified)	2024
3.224.5 Fricas [F]	2025
3.224.6 Sympy [F(-1)]	2026
3.224.7 Maxima [F]	2026
3.224.8 Giac [F(-2)]	2026
3.224.9 Mupad [F(-1)]	2027

#### 3.224.1 Optimal result

Integrand size = 22, antiderivative size = 429

$$\begin{aligned} \int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{7/2}} dx = & -\frac{x}{3c^3\sqrt{c-a^2cx^2}} - \frac{x}{30c^3(1-ax)(1+ax)\sqrt{c-a^2cx^2}} \\ & + \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{10ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} + \frac{4\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{15ac^3(1-a^2x^2)\sqrt{c-a^2cx^2}} \\ & + \frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}} + \frac{4x\operatorname{arccosh}(ax)^2}{15c^2(c-a^2cx^2)^{3/2}} \\ & + \frac{8x\operatorname{arccosh}(ax)^2}{15c^3\sqrt{c-a^2cx^2}} + \frac{8\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{15ac^3\sqrt{c-a^2cx^2}} \\ & - \frac{16\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)\log(1-e^{2\operatorname{arccosh}(ax)})}{15ac^3\sqrt{c-a^2cx^2}} \\ & - \frac{8\sqrt{-1+ax}\sqrt{1+ax}\operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)})}{15ac^3\sqrt{c-a^2cx^2}} \end{aligned}$$



output  $\frac{1}{5}x \operatorname{arccosh}(ax)^2/c/(-a^2cx^2+c)^{(5/2)} + 4/15x \operatorname{arccosh}(ax)^2/c^2/(-a^2cx^2+c)^{(3/2)} - 1/3x/c^3/(-a^2cx^2+c)^{(1/2)} - 1/30x/c^3/(-ax+1)/(ax+1)/(-a^2cx^2+c)^{(1/2)} + 8/15x \operatorname{arccosh}(ax)^2/c^3/(-a^2cx^2+c)^{(1/2)} + 1/10 \operatorname{arccosh}(ax) \cdot (ax-1)^{(1/2)} \cdot (ax+1)^{(1/2)}/a/c^3/(-a^2cx^2+c)^{(1/2)} + 4/15 \operatorname{arccosh}(ax) \cdot (ax-1)^{(1/2)} \cdot (ax+1)^{(1/2)}/a/c^3/(-a^2cx^2+c)^{(1/2)} + 8/15 \operatorname{arccosh}(ax)^2 \cdot (ax-1)^{(1/2)} \cdot (ax+1)^{(1/2)}/a/c^3/(-a^2cx^2+c)^{(1/2)} - 16/15 \operatorname{arccosh}(ax) \cdot \ln(1 - (ax+(ax-1)^{(1/2)} \cdot (ax+1)^{(1/2)})^2) \cdot (ax-1)^{(1/2)} \cdot (ax+1)^{(1/2)}/a/c^3/(-a^2cx^2+c)^{(1/2)} - 8/15 \operatorname{polylog}(2, (ax+(ax-1)^{(1/2)} \cdot (ax+1)^{(1/2)})^2) \cdot (ax-1)^{(1/2)} \cdot (ax+1)^{(1/2)}/a/c^3/(-a^2cx^2+c)^{(1/2)}$

### 3.224.2 Mathematica [A] (warning: unable to verify)

Time = 1.07 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.51

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx =$$

$$\frac{ax \left( 10 + \frac{1}{1-a^2x^2} \right) + 2 \left( 8 \sqrt{\frac{-1+ax}{1+ax}} + ax \left( -8 + 8 \sqrt{\frac{-1+ax}{1+ax}} - \frac{3}{(-1+a^2x^2)^2} + \frac{4}{-1+a^2x^2} \right) \right) \operatorname{arccosh}(ax)^2 + \left( \frac{-1+ax}{1+ax} \right)^{3/2}}{30ac^3 \sqrt{c}}$$

input `Integrate[ArcCosh[a*x]^2/(c - a^2*c*x^2)^(7/2),x]`

output  $\frac{-1/30 \cdot (ax \cdot (10 + (1 - a^2x^2)^{-1}) + 2 \cdot (8 \operatorname{Sqrt}[(-1 + ax)/(1 + ax)] + ax \cdot (-8 + 8 \operatorname{Sqrt}[(-1 + ax)/(1 + ax)] - 3/(-1 + a^2x^2)^2 + 4/(-1 + a^2x^2))) \cdot \operatorname{ArcCosh}[a*x]^2 + (((-1 + ax)/(1 + ax))^{3/2} \cdot \operatorname{ArcCosh}[a*x] \cdot (-11 + 8 \cdot a^2x^2 + 32 \cdot (-1 + a^2x^2)^2 \cdot \operatorname{Log}[1 - E^{(-2 \cdot \operatorname{ArcCosh}[a*x])}]))/(-1 + ax)^3 - 16 \cdot \operatorname{Sqrt}[(-1 + ax)/(1 + ax)] \cdot (1 + ax) \cdot \operatorname{PolyLog}[2, E^{(-2 \cdot \operatorname{ArcCosh}[a*x])}])}{(a \cdot c^3 \operatorname{Sqrt}[c - a^2 \cdot c \cdot x^2])}$

### 3.224.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 2.37 (sec) , antiderivative size = 407, normalized size of antiderivative = 0.95, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.773$ , Rules used = {6316, 25, 6316, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 2715, 2838, 6329, 41, 42, 41}

---

3.224.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx$

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx \\
 & \quad \downarrow \text{6316} \\
 & -\frac{2a\sqrt{ax-1}\sqrt{ax+1} \int -\frac{x\operatorname{arccosh}(ax)}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \frac{4 \int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{5/2}} dx}{5c} + \frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \frac{4 \int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{5/2}} dx}{5c} + \frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{6316} \\
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\
 & 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)}{(1-ax)^2(ax+1)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{3/2}} dx}{3c} + \frac{x\operatorname{arccosh}(ax)^2}{3c(c-a^2cx^2)^{3/2}} \right) \\
 & \quad \downarrow \text{6314} \\
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\
 & 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)}{(1-ax)^2(ax+1)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x\operatorname{arccosh}(ax)^2}{3c(c-a^2cx^2)^{3/2}} \right) \\
 & \quad \downarrow \text{6327} \\
 & \frac{5c}{5c(c-a^2cx^2)^{5/2}} \frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}} +
 \end{aligned}$$

---

3.224.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{7/2}} dx$

$$\begin{aligned}
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\
 & 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x \operatorname{arccosh}(ax)^2}{3c(c-a^2cx^2)^{3/2}} \right) + \\
 & \frac{5c}{5c(c-a^2cx^2)^{5/2}} x \operatorname{arccosh}(ax)^2 \\
 & \quad \downarrow \text{6328} \\
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\
 & 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1} \int \frac{ax \operatorname{arccosh}(ax) d \operatorname{arccosh}(ax)}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)}}{ac\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x \operatorname{arccosh}(ax)^2}{3c(c-a^2cx^2)^{3/2}} \right) + \\
 & \frac{5c}{5c(c-a^2cx^2)^{5/2}} x \operatorname{arccosh}(ax)^2 \\
 & \quad \downarrow \text{3042} \\
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\
 & 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1} \int -i \operatorname{arccosh}(ax) \tan\left(i \operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d \operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right) + \\
 & \frac{5c}{5c(c-a^2cx^2)^{5/2}} x \operatorname{arccosh}(ax)^2 \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

3.224.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{7/2}} dx$

$$\frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1} \int \operatorname{arccosh}(ax) \tan\left(\operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d\operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right) + \frac{x}{3}$$

$$\frac{x \operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}} \quad 5c$$

↓ 4199

$$\frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1} \left( 2i \int -\frac{e^{2\operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)}{1-e^{2\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2} i \operatorname{arccosh}(ax) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}} \quad 5c$$

↓ 25

$$\frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + 4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \int \frac{e^{2\operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)}{1-e^{2\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2} i \operatorname{arccosh}(ax) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}} \quad 5c$$

↓ 2620

$$\frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{2} \int \log(1-e^{2\operatorname{arccosh}(ax)}) \right) \operatorname{arccosh}(ax) - \frac{1}{2} \operatorname{arccosh}(ax) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

5c

$$\frac{x \operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}}$$

↓ 2715

$$\frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \log(1-e^{2\operatorname{arccosh}(ax)}) \right) de^{2\operatorname{arccosh}(ax)} \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

5c

$$\frac{x \operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}}$$

↓ 2838

$$\frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4 \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( -\frac{1}{4} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) - \frac{1}{2} \operatorname{arccosh}(ax) \log(1-e^{2\operatorname{arccosh}(ax)}) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

5c

$$\frac{x \operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}}$$

↓ 6329

---

3.224.  $\int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{7/2}} dx$

$$\begin{aligned}
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\operatorname{arccosh}(ax)}{4a^2(1-a^2x^2)^2} - \frac{\int \frac{1}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4a}\right)}{5c^3\sqrt{c-a^2cx^2}} + \\
 4 & \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\int \frac{1}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2a} + \frac{\operatorname{arccosh}(ax)}{2a^2(1-a^2x^2)}\right)}{3c^2\sqrt{c-a^2cx^2}} + \frac{2\left(\frac{x\operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1}\left(-2i\left(-\frac{1}{4}\operatorname{PolyLog}\left(2,e^{2\operatorname{arccosh}(ax)}\right) - \frac{1}{2}\operatorname{arccosh}(ax)\right)\right)}{ac\sqrt{c-a^2cx^2}}\right)}{3c} \right)
 \end{aligned}$$

5c

$$\frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}}$$

↓ 41

$$\begin{aligned}
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\operatorname{arccosh}(ax)}{4a^2(1-a^2x^2)^2} - \frac{\int \frac{1}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4a}\right)}{5c^3\sqrt{c-a^2cx^2}} + \\
 4 & \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\operatorname{arccosh}(ax)}{2a^2(1-a^2x^2)} - \frac{x}{2a\sqrt{ax-1}\sqrt{ax+1}}\right)}{3c^2\sqrt{c-a^2cx^2}} + \frac{2\left(\frac{x\operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1}\left(-2i\left(-\frac{1}{4}\operatorname{PolyLog}\left(2,e^{2\operatorname{arccosh}(ax)}\right) - \frac{1}{2}\operatorname{arccosh}(ax)\right)\right)}{ac\sqrt{c-a^2cx^2}}\right)}{3c} \right)
 \end{aligned}$$

5c

$$\frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}}$$

↓ 42

$$\begin{aligned}
 & \frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\operatorname{arccosh}(ax)}{4a^2(1-a^2x^2)^2} - \frac{-\frac{2}{3}\int \frac{1}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{x}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{4a}\right)}{5c^3\sqrt{c-a^2cx^2}} + \\
 4 & \left( \frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\operatorname{arccosh}(ax)}{2a^2(1-a^2x^2)} - \frac{x}{2a\sqrt{ax-1}\sqrt{ax+1}}\right)}{3c^2\sqrt{c-a^2cx^2}} + \frac{2\left(\frac{x\operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1}\left(-2i\left(-\frac{1}{4}\operatorname{PolyLog}\left(2,e^{2\operatorname{arccosh}(ax)}\right) - \frac{1}{2}\operatorname{arccosh}(ax)\right)\right)}{ac\sqrt{c-a^2cx^2}}\right)}{3c} \right)
 \end{aligned}$$

5c

$$\frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}}$$

↓ 41

$$\frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\operatorname{arccosh}(ax)}{4a^2(1-a^2x^2)^2} - \frac{\frac{2x}{3\sqrt{ax-1}\sqrt{ax+1}} - \frac{x}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{4a}\right)}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4\left(\frac{2a\sqrt{ax-1}\sqrt{ax+1}\left(\frac{\operatorname{arccosh}(ax)}{2a^2(1-a^2x^2)} - \frac{x}{2a\sqrt{ax-1}\sqrt{ax+1}}\right)}{3c^2\sqrt{c-a^2cx^2}} + \frac{2\left(\frac{x\operatorname{arccosh}(ax)^2}{c\sqrt{c-a^2cx^2}} + \frac{2i\sqrt{ax-1}\sqrt{ax+1}\left(-2i\left(-\frac{1}{4}\operatorname{PolyLog}\left(2,e^{2\operatorname{arccosh}(ax)}\right) - \frac{1}{2}\operatorname{arccosh}(ax)\right)\right)}{ac\sqrt{c-a^2cx^2}}\right)}{3c}\right)$$


---


$$\frac{x\operatorname{arccosh}(ax)^2}{5c(c-a^2cx^2)^{5/2}}$$

input `Int[ArcCosh[a*x]^2/(c - a^2*c*x^2)^(7/2),x]`

output `(x*ArcCosh[a*x]^2)/(5*c*(c - a^2*c*x^2)^(5/2)) + (2*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(-1/4*(-1/3*x/((-1 + a*x)^(3/2)*(1 + a*x)^(3/2)) + (2*x)/(3*Sqrt[-1 + a*x]*Sqrt[1 + a*x])))/a + ArcCosh[a*x]/(4*a^2*(1 - a^2*x^2)^2))/(5*c^3*Sqrt[c - a^2*c*x^2]) + (4*((x*ArcCosh[a*x]^2)/(3*c*(c - a^2*c*x^2)^(3/2)) + (2*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(-1/2*x/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + ArcCosh[a*x]/(2*a^2*(1 - a^2*x^2)))))/(3*c^2*Sqrt[c - a^2*c*x^2]) + (2*((x*ArcCosh[a*x]^2)/(c*Sqrt[c - a^2*c*x^2]) + ((2*I)*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*((-1/2*I)*ArcCosh[a*x]^2 - (2*I)*(-1/2*(ArcCosh[a*x]*Log[1 - E^(2*ArcCosh[a*x])]) - PolyLog[2, E^(2*ArcCosh[a*x])]/4)))/(a*c*Sqrt[c - a^2*c*x^2])))/(3*c)))/(5*c)`

### 3.224.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 41 `Int[1/(((a_) + (b_.)*(x_))^(3/2)*((c_) + (d_.)*(x_))^(3/2)), x_Symbol] := Simp[x/(a*c*Sqrt[a + b*x]*Sqrt[c + d*x]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0]`

- rule 42 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[(-x)*(a + b*x)^(m + 1)*((c + d*x)^(m + 1)/(2*a*c*(m + 1))), x] + Simp[(2*m + 3)/(2*a*c*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^(m + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && ILtQ[m + 3/2, 0]`
- rule 2620 `Int((((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2715 `Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`
- rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4199 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`
- rule 6314 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/((d_) + (e_)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(sqrt[-1 + c*x]/sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`



rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.)), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

### 3.224.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 793 vs. 2(395) = 790.

Time = 1.23 (sec) , antiderivative size = 794, normalized size of antiderivative = 1.85

method	result
default	$-\frac{\sqrt{-c(a^2x^2-1)}(8a^5x^5-20a^3x^3-8\sqrt{ax-1}\sqrt{ax+1}a^4x^4+15ax+16a^2x^2\sqrt{ax-1}\sqrt{ax+1}-8\sqrt{ax-1}\sqrt{ax+1})}{(c-a^2cx^2)^{7/2}}(-64 \operatorname{arccosh}(ax)\sqrt{ax}$

input `int(arccosh(a*x)^2/(-a^2*c*x^2+c)^(7/2),x,method=_RETURNVERBOSE)`

$$3.224. \int \frac{\operatorname{arccosh}(ax)^2}{(c-a^2cx^2)^{7/2}} dx$$

output

```
-1/30*(-c*(a^2*x^2-1))^(1/2)*(8*a^5*x^5-20*a^3*x^3-8*(a*x-1)^(1/2)*(a*x+1)
^(1/2)*a^4*x^4+15*a*x+16*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-8*(a*x-1)^(1/
2)*(a*x+1)^(1/2))*(-64*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^7*x^7-64
*arccosh(a*x)*a^8*x^8-32*(a*x+1)^(1/2)*(a*x-1)^(1/2)*a^7*x^7-32*a^8*x^8+24
8*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^5*x^5+280*arccosh(a*x)*a^6*x^
6+126*x^5*a^5*(a*x-1)^(1/2)*(a*x+1)^(1/2)+142*a^6*x^6+80*a^4*x^4*arccosh(a
*x)^2-340*a^3*x^3*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)-456*a^4*x^4*arc
cosh(a*x)-156*a^3*x^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)-265*a^4*x^4-190*a^2*x^2*
arccosh(a*x)^2+165*a*x*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)+328*a^2*x^
2*arccosh(a*x)+62*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+235*a^2*x^2+128*arccosh(
a*x)^2-88*arccosh(a*x)-80)/(40*a^10*x^10-215*a^8*x^8+469*a^6*x^6-517*a^4*x
^4+287*a^2*x^2-64)/c^4/a-16/15*(a*x+1)^(1/2)*(a*x-1)^(1/2)*(-c*(a^2*x^2-1)
)^(1/2)/c^4/a/(a^2*x^2-1)*arccosh(a*x)^2+16/15*(a*x+1)^(1/2)*(a*x-1)^(1/2)
*(-c*(a^2*x^2-1))^(1/2)/c^4/a/(a^2*x^2-1)*arccosh(a*x)*ln(1+a*x+(a*x-1)^(1
/2)*(a*x+1)^(1/2))+16/15*(a*x+1)^(1/2)*(a*x-1)^(1/2)*(-c*(a^2*x^2-1))^(1/2
)/c^4/a/(a^2*x^2-1)*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+16/15*(a*x
+1)^(1/2)*(a*x-1)^(1/2)*(-c*(a^2*x^2-1))^(1/2)/c^4/a/(a^2*x^2-1)*arccosh(a
*x)*ln(1-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+16/15*(a*x+1)^(1/2)*(a*x-1)^(1/2
)*(-c*(a^2*x^2-1))^(1/2)/c^4/a/(a^2*x^2-1)*polylog(2,a*x+(a*x-1)^(1/2)*(a*
x+1)^(1/2))
```

### 3.224.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{(-a^2cx^2 + c)^{7/2}} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^(7/2),x, algorithm="fricas")`

output `integral(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^2/(a^8*c^4*x^8 - 4*a^6*c^4*x^6 + 6*a^4*c^4*x^4 - 4*a^2*c^4*x^2 + c^4), x)`

**3.224.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx = \text{Timed out}$$

input `integrate(acosh(a*x)**2/(-a**2*c*x**2+c)**(7/2),x)`

output `Timed out`

**3.224.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{(-a^2cx^2 + c)^{7/2}} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^(7/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^2/(-a^2*c*x^2 + c)^(7/2), x)`

**3.224.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^2/(-a^2*c*x^2+c)^(7/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.224.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{acosh}(ax)^2}{(c - a^2cx^2)^{7/2}} dx$$

input `int(acosh(a*x)^2/(c - a^2*c*x^2)^(7/2), x)`output `int(acosh(a*x)^2/(c - a^2*c*x^2)^(7/2), x)`

### 3.225 $\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

3.225.1 Optimal result . . . . .	2028
3.225.2 Mathematica [A] (verified) . . . . .	2029
3.225.3 Rubi [A] (verified) . . . . .	2029
3.225.4 Maple [B] (verified) . . . . .	2033
3.225.5 Fricas [F] . . . . .	2034
3.225.6 Sympy [F] . . . . .	2034
3.225.7 Maxima [F(-2)] . . . . .	2035
3.225.8 Giac [F] . . . . .	2035
3.225.9 Mupad [F(-1)] . . . . .	2035

#### 3.225.1 Optimal result

Integrand size = 24, antiderivative size = 243

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = -\frac{15x\sqrt{1-ax}\sqrt{1+ax}}{64a^4} - \frac{x^3\sqrt{1-ax}\sqrt{1+ax}}{32a^2} + \frac{15\sqrt{-1+ax}\operatorname{arccosh}(ax)}{64a^5\sqrt{1-ax}} - \frac{3x^2\sqrt{-1+ax}\operatorname{arccosh}(ax)}{8a^3\sqrt{1-ax}} - \frac{x^4\sqrt{-1+ax}\operatorname{arccosh}(ax)}{8a\sqrt{1-ax}} - \frac{3x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{8a^4} - \frac{x^3\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{4a^2} + \frac{\sqrt{-1+ax}\operatorname{arccosh}(ax)^3}{8a^5\sqrt{1-ax}}$$

output

```
15/64*arccosh(a*x)*(a*x-1)^(1/2)/a^5/(-a*x+1)^(1/2)-3/8*x^2*arccosh(a*x)*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-1/8*x^4*arccosh(a*x)*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)+1/8*arccosh(a*x)^3*(a*x-1)^(1/2)/a^5/(-a*x+1)^(1/2)-15/64*x*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^4-1/32*x^3*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^2-3/8*x*arccosh(a*x)^2*(-a^2*x^2+1)^(1/2)/a^4-1/4*x^3*arccosh(a*x)^2*(-a^2*x^2+1)^(1/2)/a^2
```

### 3.225.2 Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.48

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

$$= \frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax)(32\operatorname{arccosh}(ax)^3 - 4\operatorname{arccosh}(ax)(16\cosh(2\operatorname{arccosh}(ax)) + \cosh(4\operatorname{arccosh}(ax))) + 32\sinh(2\operatorname{arccosh}(ax)))}{256a^5\sqrt{1-a^2x^2}}$$

input `Integrate[(x^4*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2],x]`

output `(Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(32*ArcCosh[a*x]^3 - 4*ArcCosh[a*x]*(16*Cosh[2*ArcCosh[a*x]] + Cosh[4*ArcCosh[a*x]]) + 32*Sinh[2*ArcCosh[a*x]] + Sinh[4*ArcCosh[a*x]] + 8*ArcCosh[a*x]^2*(8*Sinh[2*ArcCosh[a*x]] + Sinh[4*ArcCosh[a*x]])))/(256*a^5*Sqrt[1 - a^2*x^2])`

### 3.225.3 Rubi [A] (verified)

Time = 1.13 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.21, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.458$ , Rules used = {6353, 6298, 111, 27, 101, 43, 6353, 6298, 101, 43, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

$$\downarrow \text{6353}$$

$$\frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{\sqrt{ax-1} \int x^3 \operatorname{arccosh}(ax) dx}{2a\sqrt{1-ax}} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{4a^2}$$

$$\downarrow \text{6298}$$

$$\frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{\sqrt{ax-1} \left( \frac{1}{4} x^4 \operatorname{arccosh}(ax) - \frac{1}{4} a \int \frac{x^4}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2a\sqrt{1-ax}} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{4a^2}$$

$$\downarrow \text{111}$$

---

3.225.  $\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
& \frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{\int \frac{3x^2}{\sqrt{ax-1}\sqrt{ax+1}} dx}{4a^2} + \frac{x^3 \sqrt{ax-1}\sqrt{ax+1}}{4a^2} \right) \right)}{2a\sqrt{1-ax} \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{4a^2}} \\
& \quad \downarrow 27 \\
& \frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{3 \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}} dx}{4a^2} + \frac{x^3 \sqrt{ax-1}\sqrt{ax+1}}{4a^2} \right) \right)}{2a\sqrt{1-ax} \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{4a^2}} \\
& \quad \downarrow 101 \\
& \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{3 \left( \frac{\int \frac{\sqrt{ax-1}\sqrt{ax+1}}{2a^2} dx + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right)}{4a^2} + \frac{x^3 \sqrt{ax-1}\sqrt{ax+1}}{4a^2} \right) \right)}{2a\sqrt{1-ax} \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{4a^2}} \\
& \quad \downarrow 43 \\
& \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{x^3 \sqrt{ax-1}\sqrt{ax+1}}{4a^2} + \frac{3 \left( \frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right)}{4a^2} \right) \right)}{2a\sqrt{1-ax} \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{4a^2}} \\
& \quad \downarrow 6353 \\
& \frac{3 \left( \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{\sqrt{ax-1} \int x \operatorname{arccosh}(ax) dx}{a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} \right)}{4a^2 \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{4a^2}} \\
& \quad \downarrow 6298 \\
& \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{x^3 \sqrt{ax-1}\sqrt{ax+1}}{4a^2} + \frac{3 \left( \frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right)}{4a^2} \right) \right)}{2a\sqrt{1-ax}}
\end{aligned}$$

---

3.225.  $\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
& 3 \left( \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax) - \frac{1}{2}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} \right) \\
& \frac{4a^2}{x^3\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2} - \\
& \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{x^3\sqrt{ax-1}\sqrt{ax+1}}{4a^2} + \frac{3 \left( \frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right)}{4a^2} \right) \right)}{2a\sqrt{1-ax}} \\
& \downarrow 101 \\
& 3 \left( -\frac{\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax) - \frac{1}{2}a \left( \frac{\int \frac{1}{\sqrt{ax-1}\sqrt{ax+1}} dx + x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right) \right)}{a\sqrt{1-ax}} + \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} \right) \\
& \frac{4a^2}{x^3\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2} - \\
& \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{x^3\sqrt{ax-1}\sqrt{ax+1}}{4a^2} + \frac{3 \left( \frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right)}{4a^2} \right) \right)}{2a\sqrt{1-ax}} \\
& \downarrow 43 \\
& 3 \left( \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} - \frac{\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax) - \frac{1}{2}a \left( \frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right) \right)}{a\sqrt{1-ax}} \right) \\
& \frac{4a^2}{x^3\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2} - \\
& \frac{\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax) - \frac{1}{4}a \left( \frac{x^3\sqrt{ax-1}\sqrt{ax+1}}{4a^2} + \frac{3 \left( \frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right)}{4a^2} \right) \right)}{2a\sqrt{1-ax}} \\
& \downarrow 6307
\end{aligned}$$

---

3.225.  $\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$



$$\frac{-\frac{x^3\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{4a^2} + 3\left(\frac{\sqrt{ax-1}\operatorname{arccosh}(ax)^3}{6a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2a^2} - \frac{\sqrt{ax-1}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax) - \frac{1}{2}a\left(\frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2}\right)\right)}{a\sqrt{1-ax}}\right)}{\sqrt{ax-1}\left(\frac{1}{4}x^4\operatorname{arccosh}(ax) - \frac{1}{4}a\left(\frac{x^3\sqrt{ax-1}\sqrt{ax+1}}{4a^2} + \frac{3\left(\frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2}\right)}{4a^2}\right)\right)}{2a\sqrt{1-ax}}$$

input `Int[(x^4*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2], x]`

output `-1/4*(x^3*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/a^2 - (Sqrt[-1 + a*x]*((x^4*ArcCosh[a*x])/4 - (a*((x^3*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(4*a^2) + (3*((x*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(2*a^2) + ArcCosh[a*x]/(2*a^3)))/(4*a^2))))/(2*a*Sqrt[1 - a*x]) + (3*(-1/2*(x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/a^2 + (Sqrt[-1 + a*x]*ArcCosh[a*x]^3)/(6*a^3*Sqrt[1 - a*x]) - (Sqrt[-1 + a*x]*(x^2*ArcCosh[a*x])/2 - (a*((x*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(2*a^2) + ArcCosh[a*x]/(2*a^3))))/2))/(a*Sqrt[1 - a*x]))/(4*a^2)`

### 3.225.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 101 `Int[((a_) + (b_)*(x_))^(2*((c_) + (d_)*(x_))^(n_))*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

```
rule 111 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_] :> Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(d*f*(m + n + p + 1)), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

```
rule 6298 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((d_.)*(x_))^(m_), x_Symbol] :> Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

```
rule 6307 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]
```

```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] :> Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.225.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 487 vs. 2(199) = 398.

Time = 0.83 (sec) , antiderivative size = 488, normalized size of antiderivative = 2.01

method	result
default	$-\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^3}{8a^5(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(8a^5x^5-12a^3x^3+8\sqrt{ax-1}\sqrt{ax+1}a^4x^4+4ax-8a^2x^2\sqrt{ax-1}\sqrt{ax+1}+512a^5(a^2x^2-1))}{512a^5(a^2x^2-1)}$

3.225. 
$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

input `int(x^4*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/8*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a^5/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^3 \\ & -1/512*(-a^2*x^2+1)^(1/2)*(8*a^5*x^5-12*a^3*x^3+8*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4 \\ & +4*a*x-8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)^(1/2))* \\ & (8*\operatorname{arccosh}(a*x)^2-4*\operatorname{arccosh}(a*x)+1)/a^5/(a^2*x^2-1)-1/16*(-a^2*x^2+1)^(1/2)* \\ & (2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))* \\ & (2*\operatorname{arccosh}(a*x)^2-2*\operatorname{arccosh}(a*x)+1)/a^5/(a^2*x^2-1)-1/16*(-a^2*x^2+1)^(1/2)* \\ & (2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)^(1/2))* \\ & (2*\operatorname{arccosh}(a*x)^2+2*\operatorname{arccosh}(a*x)+1)/a^5/(a^2*x^2-1)-1/512*(-a^2*x^2+1)^(1/2)* \\ & (8*a^5*x^5-12*a^3*x^3-8*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4+4*a*x+8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2) \\ & -(a*x-1)^(1/2)*(a*x+1)^(1/2))* \\ & (8*\operatorname{arccosh}(a*x)^2+4*\operatorname{arccosh}(a*x)+1)/a^5/(a^2*x^2-1) \end{aligned}$$

### 3.225.5 Fricas [F]

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^4*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^4*arccosh(a*x)^2/(a^2*x^2 - 1), x)`

### 3.225.6 Sympy [F]

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{acosh}^2(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**4*acosh(a*x)**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**4*acosh(a*x)**2/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.225.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.225.8 Giac [F]**

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^4*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^4*arccosh(a*x)^2/sqrt(-a^2*x^2+1), x)`

**3.225.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{acosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

input `int((x^4*acosh(a*x)^2)/(1-a^2*x^2)^(1/2),x)`

output `int((x^4*acosh(a*x)^2)/(1-a^2*x^2)^(1/2), x)`

### 3.226 $\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

3.226.1 Optimal result . . . . .	2036
3.226.2 Mathematica [A] (verified) . . . . .	2036
3.226.3 Rubi [A] (verified) . . . . .	2037
3.226.4 Maple [B] (verified) . . . . .	2040
3.226.5 Fricas [A] (verification not implemented) . . . . .	2040
3.226.6 Sympy [F] . . . . .	2041
3.226.7 Maxima [C] (verification not implemented) . . . . .	2041
3.226.8 Giac [F(-2)] . . . . .	2042
3.226.9 Mupad [F(-1)] . . . . .	2042

#### 3.226.1 Optimal result

Integrand size = 24, antiderivative size = 177

$$\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = -\frac{40\sqrt{1-ax}\sqrt{1+ax}}{27a^4} - \frac{2x^2\sqrt{1-ax}\sqrt{1+ax}}{27a^2}$$

$$- \frac{4x\sqrt{-1+ax}\operatorname{arccosh}(ax)}{3a^3\sqrt{1-ax}} - \frac{2x^3\sqrt{-1+ax}\operatorname{arccosh}(ax)}{9a\sqrt{1-ax}}$$

$$- \frac{2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{3a^4} - \frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{3a^2}$$

output

```
-4/3*x*arccosh(a*x)*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-2/9*x^3*arccosh(a*x)*
(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)-40/27*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^4-2/27
*x^2*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^2-2/3*arccosh(a*x)^2*(-a^2*x^2+1)^(1/2
)/a^4-1/3*x^2*arccosh(a*x)^2*(-a^2*x^2+1)^(1/2)/a^2
```

#### 3.226.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.69

$$\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \left( -\frac{40}{27a^4} - \frac{2x^2}{27a^2} \right) \sqrt{1-a^2x^2} + \frac{2x\sqrt{1-a^2x^2}(6+a^2x^2)\operatorname{arccosh}(ax)}{9a^3\sqrt{-1+ax}\sqrt{1+ax}}$$

$$- \frac{\sqrt{1-a^2x^2}(2+a^2x^2)\operatorname{arccosh}(ax)^2}{3a^4}$$

input `Integrate[(x^3*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2],x]`

output  $(-40/(27*a^4) - (2*x^2)/(27*a^2))*Sqrt[1 - a^2*x^2] + (2*x*Sqrt[1 - a^2*x^2]*(6 + a^2*x^2)*ArcCosh[a*x])/(9*a^3*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) - (Sqrt[1 - a^2*x^2]*(2 + a^2*x^2)*ArcCosh[a*x]^2)/(3*a^4)$

### 3.226.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.21, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6353, 6298, 111, 27, 83, 6329, 6294, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{2 \int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{2\sqrt{ax-1} \int x^2 \operatorname{arccosh}(ax) dx}{3a\sqrt{1-ax}} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2} \\
 & \quad \downarrow \text{6298} \\
 & \frac{2 \int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{2\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax) - \frac{1}{3} a \int \frac{x^3}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{3a\sqrt{1-ax}} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2} \\
 & \quad \downarrow \text{111} \\
 & \frac{2 \int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{2\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax) - \frac{1}{3} a \left( \frac{\int \frac{2x}{\sqrt{ax-1}\sqrt{ax+1}} dx}{3a^2} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{3a\sqrt{1-ax}} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{2 \int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{2\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax) - \frac{1}{3} a \left( \frac{2 \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx}{3a^2} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{3a\sqrt{1-ax}} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2}
 \end{aligned}$$

---

3.226.  $\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
& \downarrow 83 \\
& \frac{2 \int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2} - \\
& \frac{2\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax) - \frac{1}{3} a \left( \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a^4} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{3a\sqrt{1-ax}} \\
& \downarrow 6329 \\
& \frac{2 \left( -\frac{2\sqrt{ax-1} \int \operatorname{arccosh}(ax) dx}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{a^2} \right)}{3a^2} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2} - \\
& \frac{2\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax) - \frac{1}{3} a \left( \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a^4} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{3a\sqrt{1-ax}} \\
& \downarrow 6294 \\
& \frac{2 \left( -\frac{2\sqrt{ax-1} \left( x \operatorname{arccosh}(ax) - a \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{a^2} \right)}{3a^2} - \\
& \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2} - \frac{2\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax) - \frac{1}{3} a \left( \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a^4} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{3a\sqrt{1-ax}} \\
& \downarrow 83 \\
& -\frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{3a^2} + \frac{2 \left( -\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2\sqrt{ax-1} \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a\sqrt{1-ax}} \right)}{3a^2} - \\
& \frac{2\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax) - \frac{1}{3} a \left( \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a^4} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{3a\sqrt{1-ax}}
\end{aligned}$$

input `Int[(x^3*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2], x]`

output `-1/3*(x^2*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/a^2 - (2*Sqrt[-1 + a*x]*(-1/3*(a*((2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(3*a^4) + (x^2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(3*a^2))) + (x^3*ArcCosh[a*x])/3))/(3*a*Sqrt[1 - a*x]) + (2*(-((Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/a^2) - (2*Sqrt[-1 + a*x]*(-((Sqrt[-1 + a*x]*Sqrt[1 + a*x])/a) + x*ArcCosh[a*x]))/(a*Sqrt[1 - a*x])))/(3*a^2)`

## 3.226.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 111 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 6294 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`
- rule 6298 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6329 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`



```
rule 6353 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] :> Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]
```

### 3.226.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 342 vs. 2(145) = 290.

Time = 1.12 (sec) , antiderivative size = 343, normalized size of antiderivative = 1.94

method	result
default	$-\frac{\sqrt{-a^2x^2+1} (4a^4x^4-5a^2x^2+4a^3x^3\sqrt{ax-1}\sqrt{ax+1}-3\sqrt{ax-1}\sqrt{ax+1}ax+1) (9 \operatorname{arccosh}(ax)^2-6 \operatorname{arccosh}(ax)+2)}{216a^4(a^2x^2-1)} - \frac{3\sqrt{-a^2x^2+1}}{216a^4(a^2x^2-1)}$

```
input int(x^3*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/216*(-a^2*x^2+1)^(1/2)*(4*a^4*x^4-5*a^2*x^2+4*a^3*x^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)-3*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+1)*(9*arccosh(a*x)^2-6*arccosh(a*x)+2)/a^4/(a^2*x^2-1)-3/8*(-a^2*x^2+1)^(1/2)*((a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+a^2*x^2-1)*(arccosh(a*x)^2-2*arccosh(a*x)+2)/a^4/(a^2*x^2-1)-3/8*(-a^2*x^2+1)^(1/2)*(a^2*x^2-(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x-1)*(arccosh(a*x)^2+2*arccosh(a*x)+2)/a^4/(a^2*x^2-1)-1/216*(-a^2*x^2+1)^(1/2)*(4*a^4*x^4-5*a^2*x^2-4*a^3*x^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)+3*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+1)*(9*arccosh(a*x)^2+6*arccosh(a*x)+2)/a^4/(a^2*x^2-1)
```

### 3.226.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.85

$$\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \frac{9(a^4x^4 + a^2x^2 - 2)\sqrt{-a^2x^2 + 1} \log(ax + \sqrt{a^2x^2 - 1})^2 - 6(a^3x^3 + 6ax)\sqrt{a^2x^2 - 1}\sqrt{-a^2x^2 + 1} \log(ax + \sqrt{a^2x^2 - 1})}{27(a^6x^2 - a^4)}$$

input `integrate(x^3*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `-1/27*(9*(a^4*x^4 + a^2*x^2 - 2)*sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1))^2 - 6*(a^3*x^3 + 6*a*x)*sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1)) + 2*(a^4*x^4 + 19*a^2*x^2 - 20)*sqrt(-a^2*x^2 + 1))/(a^6*x^2 - a^4)`

### 3.226.6 Sympy [F]

$$\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^3 \operatorname{acosh}^2(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**3*acosh(a*x)**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**3*acosh(a*x)**2/sqrt(-(a*x - 1)*(a*x + 1)), x)`

### 3.226.7 Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.59

$$\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = -\frac{1}{3} \left( \frac{\sqrt{-a^2x^2+1}x^2}{a^2} + \frac{2\sqrt{-a^2x^2+1}}{a^4} \right) \operatorname{arccosh}(ax)^2 + \frac{2 \left( -i\sqrt{a^2x^2-1}x^2 - \frac{20i\sqrt{a^2x^2-1}}{a^2} \right)}{27a^2} + \frac{2(i a^2 x^3 + 6i x) \operatorname{arccosh}(ax)}{9a^3}$$

input `integrate(x^3*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `-1/3*(sqrt(-a^2*x^2 + 1)*x^2/a^2 + 2*sqrt(-a^2*x^2 + 1)/a^4)*arccosh(a*x)^2 + 2/27*(-I*sqrt(a^2*x^2 - 1)*x^2 - 20*I*sqrt(a^2*x^2 - 1)/a^2)/a^2 + 2/9*(I*a^2*x^3 + 6*I*x)*arccosh(a*x)/a^3`

**3.226.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.226.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^3 \operatorname{acosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

input `int((x^3*acosh(a*x)^2)/(1 - a^2*x^2)^(1/2),x)`

output `int((x^3*acosh(a*x)^2)/(1 - a^2*x^2)^(1/2), x)`

### 3.227 $\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

3.227.1 Optimal result . . . . .	2043
3.227.2 Mathematica [A] (verified) . . . . .	2043
3.227.3 Rubi [A] (verified) . . . . .	2044
3.227.4 Maple [A] (verified) . . . . .	2046
3.227.5 Fricas [F] . . . . .	2046
3.227.6 Sympy [F] . . . . .	2047
3.227.7 Maxima [F(-2)] . . . . .	2047
3.227.8 Giac [F] . . . . .	2047
3.227.9 Mupad [F(-1)] . . . . .	2048

#### 3.227.1 Optimal result

Integrand size = 24, antiderivative size = 151

$$\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = -\frac{x\sqrt{1-ax}\sqrt{1+ax}}{4a^2} + \frac{\sqrt{-1+ax}\operatorname{arccosh}(ax)}{4a^3\sqrt{1-ax}} - \frac{x^2\sqrt{-1+ax}\operatorname{arccosh}(ax)}{2a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2a^2} + \frac{\sqrt{-1+ax}\operatorname{arccosh}(ax)^3}{6a^3\sqrt{1-ax}}$$

output  $1/4*\operatorname{arccosh}(a*x)*(a*x-1)^{(1/2)}/a^3/(-a*x+1)^{(1/2)}-1/2*x^2*\operatorname{arccosh}(a*x)*(a*x-1)^{(1/2)}/a/(-a*x+1)^{(1/2)}+1/6*\operatorname{arccosh}(a*x)^3*(a*x-1)^{(1/2)}/a^3/(-a*x+1)^{(1/2)}-1/4*x*(-a*x+1)^{(1/2)*(a*x+1)^{(1/2)}/a^2-1/2*x*\operatorname{arccosh}(a*x)^2*(-a^2*x^2+1)^{(1/2)}/a^2$

#### 3.227.2 Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.58

$$\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-((-1+ax)(1+ax))}(4\operatorname{arccosh}(ax)^3 - 6\operatorname{arccosh}(ax) \cosh(2\operatorname{arccosh}(ax)) + (3 + 6\operatorname{arccosh}(ax)^2) \sin^{-1}(ax))}{24a^3\sqrt{\frac{-1+ax}{1+ax}}(1+ax)}$$

input `Integrate[(x^2*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2],x]`

output `-1/24*(Sqrt[-((-1 + a*x)*(1 + a*x))]*(4*ArcCosh[a*x]^3 - 6*ArcCosh[a*x]*Cosh[2*ArcCosh[a*x]] + (3 + 6*ArcCosh[a*x]^2)*Sinh[2*ArcCosh[a*x]]))/(a^3*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x))`

### 3.227.3 Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.93, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$ , Rules used = {6353, 6298, 101, 43, 6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6353} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{\sqrt{ax-1} \int x \operatorname{arccosh}(ax) dx}{a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} \\
 & \quad \downarrow \text{6298} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{\sqrt{ax-1} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax) - \frac{1}{2} a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} \\
 & \quad \downarrow \text{101} \\
 & - \frac{\sqrt{ax-1} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax) - \frac{1}{2} a \left( \frac{\int \frac{1}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right) \right)}{a\sqrt{1-ax}} + \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \\
 & \quad \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} \\
 & \quad \downarrow \text{43} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{2a^2} - \\
 & \frac{\sqrt{ax-1} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax) - \frac{1}{2} a \left( \frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2} \right) \right)}{a\sqrt{1-ax}} \\
 & \quad \downarrow \text{6307}
 \end{aligned}$$

---

3.227.  $\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

$$\frac{\frac{\sqrt{ax-1}\operatorname{arccosh}(ax)^3}{6a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2a^2}}{\frac{\sqrt{ax-1}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax) - \frac{1}{2}a\left(\frac{\operatorname{arccosh}(ax)}{2a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a^2}\right)\right)}{a\sqrt{1-ax}}}$$

input `Int[(x^2*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2],x]`

output `-1/2*(x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/a^2 + (Sqrt[-1 + a*x]*ArcCosh[a*x]^3)/(6*a^3*Sqrt[1 - a*x]) - (Sqrt[-1 + a*x]*((x^2*ArcCosh[a*x])/2 - (a*(x*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(2*a^2) + ArcCosh[a*x]/(2*a^3))))/(a*Sqrt[1 - a*x])`

### 3.227.3.1 Defintions of rubi rules used

rule 43 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 101 `Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.227.4 Maple [A] (verified)

Time = 0.62 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.58

method	result
default	$-\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^3}{6a^3(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(2a^3x^3-2ax+2a^2x^2\sqrt{ax-1}\sqrt{ax+1}-\sqrt{ax-1}\sqrt{ax+1})(2\operatorname{arccosh}(ax))^2}{16a^3(a^2x^2-1)}$

input `int(x^2*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-1/6*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a^3/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^3-1/16*(-a^2*x^2+1)^(1/2)*(2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))*(2*\operatorname{arccosh}(a*x)^2-2*\operatorname{arccosh}(a*x)+1)/a^3/(a^2*x^2-1)-1/16*(-a^2*x^2+1)^(1/2)*(2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)^(1/2))*(2*\operatorname{arccosh}(a*x)^2+2*\operatorname{arccosh}(a*x)+1)/a^3/(a^2*x^2-1)$$

### 3.227.5 Fracas [F]

$$\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^2*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^2*arccosh(a*x)^2/(a^2*x^2 - 1), x)`

**3.227.6 Sympy [F]**

$$\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{acosh}^2(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**2*acosh(a*x)**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**2*acosh(a*x)**2/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.227.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^2*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.227.8 Giac [F]**

$$\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^2*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^2*arccosh(a*x)^2/sqrt(-a^2*x^2 + 1), x)`



**3.227.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2 \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{acosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

input `int((x^2*acosh(a*x)^2)/(1 - a^2*x^2)^(1/2),x)`output `int((x^2*acosh(a*x)^2)/(1 - a^2*x^2)^(1/2), x)`

### 3.228 $\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

3.228.1 Optimal result . . . . .	2049
3.228.2 Mathematica [A] (verified) . . . . .	2049
3.228.3 Rubi [A] (verified) . . . . .	2050
3.228.4 Maple [A] (verified) . . . . .	2051
3.228.5 Fricas [A] (verification not implemented) . . . . .	2051
3.228.6 Sympy [F] . . . . .	2052
3.228.7 Maxima [C] (verification not implemented) . . . . .	2052
3.228.8 Giac [C] (verification not implemented) . . . . .	2052
3.228.9 Mupad [F(-1)] . . . . .	2053

#### 3.228.1 Optimal result

Integrand size = 22, antiderivative size = 79

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = -\frac{2\sqrt{1-ax}\sqrt{1+ax}}{a^2} - \frac{2x\sqrt{-1+ax}\operatorname{arccosh}(ax)}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{a^2}$$

output  $-2*x*\operatorname{arccosh}(a*x)*(a*x-1)^{(1/2)}/a/(-a*x+1)^{(1/2)}-2*(-a*x+1)^{(1/2)}*(a*x+1)^{(1/2)}/a^2-\operatorname{arccosh}(a*x)^2*(-a^2*x^2+1)^{(1/2)}/a^2$

#### 3.228.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.68

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{1-a^2x^2} \left( -2 + \frac{2ax \operatorname{arccosh}(ax)}{\sqrt{-1+ax}\sqrt{1+ax}} - \operatorname{arccosh}(ax)^2 \right)}{a^2}$$

input `Integrate[(x*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2],x]`

output `(Sqrt[1 - a^2*x^2]*(-2 + (2*a*x*ArcCosh[a*x]))/(Sqrt[-1 + a*x]*Sqrt[1 + a*x]) - ArcCosh[a*x]^2)/a^2`

**3.228.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$ , Rules used = {6329, 6294, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

$$\downarrow 6329$$

$$-\frac{2\sqrt{ax-1} \int \operatorname{arccosh}(ax) dx}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{a^2}$$

$$\downarrow 6294$$

$$-\frac{2\sqrt{ax-1} \left( x \operatorname{arccosh}(ax) - a \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{a^2}$$

$$\downarrow 83$$

$$-\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2\sqrt{ax-1} \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a\sqrt{1-ax}}$$

input `Int[(x*ArcCosh[a*x]^2)/Sqrt[1 - a^2*x^2], x]`

output `-((Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/a^2) - (2*Sqrt[-1 + a*x]*(-(Sqrt[-1 + a*x]*Sqrt[1 + a*x])/a) + x*ArcCosh[a*x]))/(a*Sqrt[1 - a*x])`

**3.228.3.1 Defintions of rubi rules used**

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

---

3.228.  $\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

### 3.228.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.76

method	result
default	$-\frac{\sqrt{-a^2x^2+1}(\sqrt{ax-1}\sqrt{ax+1}ax+a^2x^2-1)(\operatorname{arccosh}(ax)^2-2\operatorname{arccosh}(ax)+2)}{2a^2(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(a^2x^2-\sqrt{ax-1}\sqrt{ax+1}ax-1)(\operatorname{arccosh}(ax)^2-2\operatorname{arccosh}(ax)+2)}{2a^2(a^2x^2-1)}$

```
input int(x*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/2*(-a^2*x^2+1)^(1/2)*((a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+a^2*x^2-1)*(arcco
sh(a*x)^2-2*arccosh(a*x)+2)/a^2/(a^2*x^2-1)-1/2*(-a^2*x^2+1)^(1/2)*(a^2*x^
2-(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x-1)*(arccosh(a*x)^2+2*arccosh(a*x)+2)/a^2
/(a^2*x^2-1)
```

### 3.228.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.44

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

$$= \frac{2\sqrt{a^2x^2-1}\sqrt{-a^2x^2+1}ax \log(ax + \sqrt{a^2x^2-1}) + (-a^2x^2+1)^{\frac{3}{2}} \log(ax + \sqrt{a^2x^2-1})^2 - 2(a^2x^2-1)\sqrt{-a^2x^2+1}}{a^4x^2-a^2}$$

```
input integrate(x*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="fracas")
```

```
output (2*sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)*a*x*log(a*x + sqrt(a^2*x^2 - 1)) +
(-a^2*x^2 + 1)^(3/2)*log(a*x + sqrt(a^2*x^2 - 1))^2 - 2*(a^2*x^2 - 1)*sqr
t(-a^2*x^2 + 1))/(a^4*x^2 - a^2)
```

**3.228.6 Sympy [F]**

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x \operatorname{acosh}^2(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x*acosh(a*x)**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x*acosh(a*x)**2/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.228.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.23 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.63

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \frac{2i x \operatorname{arccosh}(ax)}{a} - \frac{\sqrt{-a^2x^2+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2i \sqrt{a^2x^2-1}}{a^2}$$

input `integrate(x*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `2*I*x*arccosh(a*x)/a - sqrt(-a^2*x^2 + 1)*arccosh(a*x)^2/a^2 - 2*I*sqrt(a^2*x^2 - 1)/a^2`

**3.228.8 Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.96

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = -\frac{\sqrt{-a^2x^2+1} \log(ax + \sqrt{a^2x^2-1})^2}{a^2} - \frac{2i \left( x \log(ax + \sqrt{a^2x^2-1}) - \frac{\sqrt{a^2x^2-1}}{a} \right)}{a}$$

input `integrate(x*arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `-sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1))^2/a^2 - 2*I*(x*log(a*x + sqrt(a^2*x^2 - 1)) - sqrt(a^2*x^2 - 1)/a)/a`

**3.228.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{x \operatorname{acosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

input `int((x*acosh(a*x)^2)/(1 - a^2*x^2)^(1/2), x)`output `int((x*acosh(a*x)^2)/(1 - a^2*x^2)^(1/2), x)`

$$3.229 \quad \int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

3.229.1 Optimal result . . . . .	2054
3.229.2 Mathematica [A] (verified) . . . . .	2054
3.229.3 Rubi [A] (verified) . . . . .	2055
3.229.4 Maple [A] (verified) . . . . .	2055
3.229.5 Fricas [F] . . . . .	2056
3.229.6 Sympy [F] . . . . .	2056
3.229.7 Maxima [F] . . . . .	2056
3.229.8 Giac [F] . . . . .	2057
3.229.9 Mupad [F(-1)] . . . . .	2057

### 3.229.1 Optimal result

Integrand size = 21, antiderivative size = 32

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-1+ax}\operatorname{arccosh}(ax)^3}{3a\sqrt{1-ax}}$$

output  $1/3*\operatorname{arccosh}(a*x)^3*(a*x-1)^{(1/2)}/a/(-a*x+1)^{(1/2)}$

### 3.229.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.41

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^3}{3a\sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]^2/Sqrt[1 - a^2*x^2],x]`

output  $(\operatorname{Sqrt}[-1+a*x]*\operatorname{Sqrt}[1+a*x]*\operatorname{ArcCosh}[a*x]^3)/(3*a*\operatorname{Sqrt}[1-a^2*x^2])$

### 3.229.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

↓ 6307

$$\frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^3}{3a\sqrt{1-ax}}$$

input `Int[ArcCosh[a*x]^2/Sqrt[1 - a^2*x^2], x]`

output `(Sqrt[-1 + a*x]*ArcCosh[a*x]^3)/(3*a*Sqrt[1 - a*x])`

#### 3.229.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.229.4 Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.59

method	result	size
default	$-\frac{\sqrt{-(ax-1)(ax+1)}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^3}{3(a^2x^2-1)a}$	51

input `int(arccosh(a*x)^2/(-a^2*x^2+1)^(1/2), x, method=_RETURNVERBOSE)`

output `-1/3*(-(a*x-1)*(a*x+1))^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2-1)/a*arccosh(a*x)^3`



**3.229.5 Fracas [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^2/(a^2*x^2 - 1), x)`

**3.229.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}^2(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**2/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.229.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^2/sqrt(-a^2*x^2 + 1), x)`

**3.229.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^2/sqrt(-a^2*x^2 + 1), x)`

**3.229.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^2}{\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^2/(1 - a^2*x^2)^(1/2),x)`

output `int(acosh(a*x)^2/(1 - a^2*x^2)^(1/2), x)`

### 3.230 $\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx$

3.230.1 Optimal result . . . . .	2058
3.230.2 Mathematica [A] (verified) . . . . .	2059
3.230.3 Rubi [A] (verified) . . . . .	2059
3.230.4 Maple [F] . . . . .	2061
3.230.5 Fricas [F] . . . . .	2061
3.230.6 Sympy [F] . . . . .	2062
3.230.7 Maxima [F] . . . . .	2062
3.230.8 Giac [F] . . . . .	2062
3.230.9 Mupad [F(-1)] . . . . .	2063

#### 3.230.1 Optimal result

Integrand size = 24, antiderivative size = 183

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx = \frac{2\sqrt{-1+ax}\operatorname{arccosh}(ax)^2 \arctan(e^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{2i\sqrt{-1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} + \frac{2i\sqrt{-1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} + \frac{2i\sqrt{-1+ax} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{2i\sqrt{-1+ax} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}}$$

output

```
2*arccosh(a*x)^2*arctan(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-2*I*arccosh(a*x)*polylog(2,-I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)+2*I*arccosh(a*x)*polylog(2,I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)+2*I*polylog(3,-I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-2*I*polylog(3,I*(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))*(a*x-1)^(1/2)/(-a*x+1)^(1/2)
```

**3.230.2 Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx$$

$$= \frac{i\sqrt{\frac{-1+ax}{1+ax}}(1+ax)(-\operatorname{arccosh}(ax)^2(\log(1-ie^{-\operatorname{arccosh}(ax)}) - \log(1+ie^{-\operatorname{arccosh}(ax)})) - 2\operatorname{arccosh}(ax)(\operatorname{PolyLog}[2, (-1)/E^{\operatorname{arccosh}(ax)}] - \operatorname{PolyLog}[2, 1/E^{\operatorname{arccosh}(ax)}]) - 2\operatorname{PolyLog}[3, (-1)/E^{\operatorname{arccosh}(ax)}] + 2\operatorname{PolyLog}[3, 1/E^{\operatorname{arccosh}(ax)}]))}{x\sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]^2/(x*Sqrt[1 - a^2*x^2]),x]`output `(I*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(-(ArcCosh[a*x]^2*(Log[1 - 1/E^ArcCosh[a*x]] - Log[1 + 1/E^ArcCosh[a*x]]) - 2*ArcCosh[a*x]*(PolyLog[2, (-1)/E^ArcCosh[a*x]] - PolyLog[2, 1/E^ArcCosh[a*x]]) - 2*PolyLog[3, (-1)/E^ArcCosh[a*x]] + 2*PolyLog[3, 1/E^ArcCosh[a*x]])))/Sqrt[1 - a^2*x^2]`**3.230.3 Rubi [A] (verified)**Time = 0.54 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.58, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6361, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx$$

$$\downarrow 6361$$

$$\frac{\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)^2}{ax} d\operatorname{arccosh}(ax)}{\sqrt{1-ax}}$$

$$\downarrow 3042$$

$$\frac{\sqrt{ax-1} \int \operatorname{arccosh}(ax)^2 \csc\left(i\operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d\operatorname{arccosh}(ax)}{\sqrt{1-ax}}$$

$$\downarrow 4668$$

$$\frac{\sqrt{ax-1}(-2i \int \operatorname{arccosh}(ax) \log(1 - ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2i \int \operatorname{arccosh}(ax) \log(1 + ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax))}{\sqrt{1-ax}}$$

3.230.  $\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx$

↓ 3011

$$\frac{\sqrt{ax-1} (2i \int \text{PolyLog}(2, -ie^{\text{arccosh}(ax)}) \text{darccosh}(ax) - \text{arccosh}(ax) \text{PolyLog}(2, -ie^{\text{arccosh}(ax)}) - 2i \int \text{PolyLog}(\dots))}{\sqrt{1-ax}}$$

↓ 2720

$$\frac{\sqrt{ax-1} (2i \int e^{-\text{arccosh}(ax)} \text{PolyLog}(2, -ie^{\text{arccosh}(ax)}) de^{\text{arccosh}(ax)} - \text{arccosh}(ax) \text{PolyLog}(2, -ie^{\text{arccosh}(ax)}) - 2i \int \dots)}{\sqrt{1-ax}}$$

↓ 7143

$$\frac{\sqrt{ax-1} (2\text{arccosh}(ax)^2 \arctan(e^{\text{arccosh}(ax)}) + 2i (\text{PolyLog}(3, -ie^{\text{arccosh}(ax)}) - \text{arccosh}(ax) \text{PolyLog}(2, -ie^{\text{arccosh}(ax)}))}{\sqrt{1-ax}}$$

input `Int[ArcCosh[a*x]^2/(x*sqrt[1 - a^2*x^2]),x]`

output `(sqrt[-1 + a*x]*(2*ArcCosh[a*x]^2*ArcTan[E^ArcCosh[a*x]] + (2*I)*(-(ArcCosh[a*x]*PolyLog[2, (-I)*E^ArcCosh[a*x]]) + PolyLog[3, (-I)*E^ArcCosh[a*x]]) - (2*I)*(-(ArcCosh[a*x]*PolyLog[2, I*E^ArcCosh[a*x]]) + PolyLog[3, I*E^ArcCosh[a*x]])))/sqrt[1 - a*x]`

### 3.230.3.1 Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)^v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

---

3.230.  $\int \frac{\text{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx$

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_.)]*((c_.) + (d_.)*(x_.))^m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6361 `Int[(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n_.)*(x_)^m_)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.230.4 Maple [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{-a^2x^2+1}} dx$$

input `int(arccosh(a*x)^2/x/(-a^2*x^2+1)^(1/2),x)`

output `int(arccosh(a*x)^2/x/(-a^2*x^2+1)^(1/2),x)`

### 3.230.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arccosh}(ax)^2}{\sqrt{-a^2x^2+1}x} dx$$

input `integrate(arccosh(a*x)^2/x/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^2/(a^2*x^3 - x), x)`

**3.230.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}^2(ax)}{x\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**2/x/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**2/(x*sqrt(-(a*x - 1)*(a*x + 1))), x)`

**3.230.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}x} dx$$

input `integrate(arccosh(a*x)^2/x/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^2/(sqrt(-a^2*x^2 + 1)*x), x)`

**3.230.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}x} dx$$

input `integrate(arccosh(a*x)^2/x/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^2/(sqrt(-a^2*x^2 + 1)*x), x)`

**3.230.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^2/(x*(1 - a^2*x^2)^(1/2)),x)`output `int(acosh(a*x)^2/(x*(1 - a^2*x^2)^(1/2)), x)`



### 3.231 $\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx$

3.231.1 Optimal result . . . . .	2064
3.231.2 Mathematica [A] (verified) . . . . .	2064
3.231.3 Rubi [C] (verified) . . . . .	2065
3.231.4 Maple [A] (verified) . . . . .	2067
3.231.5 Fricas [F] . . . . .	2068
3.231.6 Sympy [F] . . . . .	2068
3.231.7 Maxima [F] . . . . .	2069
3.231.8 Giac [F] . . . . .	2069
3.231.9 Mupad [F(-1)] . . . . .	2069

#### 3.231.1 Optimal result

Integrand size = 24, antiderivative size = 124

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx = \frac{a\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} - \frac{2a\sqrt{-1+ax}\operatorname{arccosh}(ax)\log(1+e^{2\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{a\sqrt{-1+ax}\operatorname{PolyLog}(2,-e^{2\operatorname{arccosh}(ax)})}{\sqrt{1-ax}}$$

output `a*arccosh(a*x)^2*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-2*a*arccosh(a*x)*ln(1+(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))^2*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-a*polylog(2,-(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2)))^2*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-arccosh(a*x)^2*(-a^2*x^2+1)^(1/2)/x`

#### 3.231.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx = \frac{a\sqrt{\frac{-1+ax}{1+ax}}(1+ax)\left(\operatorname{arccosh}(ax)\left(-\operatorname{arccosh}(ax)+\frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax)\operatorname{arccosh}(ax)}{ax}-2\log(1+e^{-2\operatorname{arccosh}(ax)})\right)\right)}{\sqrt{-((-1+ax)(1+ax))}}$$

input `Integrate[ArcCosh[a*x]^2/(x^2*sqrt[1 - a^2*x^2]),x]`

output `(a*sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(ArcCosh[a*x]*(-ArcCosh[a*x] + (sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*ArcCosh[a*x])/(a*x) - 2*Log[1 + E^(-2*ArcCosh[a*x])])) + PolyLog[2, -E^(-2*ArcCosh[a*x])])/sqrt[-((-1 + a*x)*(1 + a*x))]`

### 3.231.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.57 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.83, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6332, 6297, 3042, 26, 4201, 2620, 2715, 2838}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6332} \\
 & -\frac{2a\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)}{x} dx}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} \\
 & \quad \downarrow \text{6297} \\
 & -\frac{2a\sqrt{ax-1} \int \frac{\sqrt{\frac{ax-1}{ax+1}}(ax+1)\operatorname{arccosh}(ax)}{ax} d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} - \frac{2a\sqrt{ax-1} \int -i\operatorname{arccosh}(ax) \tan(i\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{26} \\
 & -\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} + \frac{2ia\sqrt{ax-1} \int \operatorname{arccosh}(ax) \tan(i\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{4201}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} + 2ia\sqrt{ax-1}\left(2i\int\frac{e^{2\operatorname{arccosh}(ax)}\operatorname{arccosh}(ax)}{1+e^{2\operatorname{arccosh}(ax)}}d\operatorname{arccosh}(ax) - \frac{1}{2}i\operatorname{arccosh}(ax)^2\right)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{2620} \\
 & \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} + 2ia\sqrt{ax-1}\left(2i\left(\frac{1}{2}\operatorname{arccosh}(ax)\log\left(e^{2\operatorname{arccosh}(ax)}+1\right) - \frac{1}{2}\int\log\left(1+e^{2\operatorname{arccosh}(ax)}\right)d\operatorname{arccosh}(ax)\right) - \frac{1}{2}i\operatorname{arccosh}(ax)^2\right)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{2715} \\
 & \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} + 2ia\sqrt{ax-1}\left(2i\left(\frac{1}{2}\operatorname{arccosh}(ax)\log\left(e^{2\operatorname{arccosh}(ax)}+1\right) - \frac{1}{4}\int e^{-2\operatorname{arccosh}(ax)}\log\left(1+e^{2\operatorname{arccosh}(ax)}\right)de^{2\operatorname{arccosh}(ax)}\right) - \frac{1}{2}i\operatorname{arccosh}(ax)^2\right)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{2838} \\
 & \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{x} + 2ia\sqrt{ax-1}\left(2i\left(\frac{1}{4}\operatorname{PolyLog}\left(2,-e^{2\operatorname{arccosh}(ax)}\right) + \frac{1}{2}\operatorname{arccosh}(ax)\log\left(e^{2\operatorname{arccosh}(ax)}+1\right)\right) - \frac{1}{2}i\operatorname{arccosh}(ax)^2\right)}{\sqrt{1-ax}}
 \end{aligned}$$

input `Int[ArcCosh[a*x]^2/(x^2*Sqrt[1 - a^2*x^2]),x]`

output `-((Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/x) + ((2*I)*a*Sqrt[-1 + a*x]*((-1/2*I)*ArcCosh[a*x]^2 + (2*I)*((ArcCosh[a*x]*Log[1 + E^(2*ArcCosh[a*x])]))/2 + PolyLog[2, -E^(2*ArcCosh[a*x])]/4))/Sqrt[1 - a*x]`

### 3.231.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

---

3.231.  $\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx$

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
-> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))]^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4201 `Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)/(x_), x_Symbol] :> Simp[1/b
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a
, b, c}, x] && IGtQ[n, 0]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((f_.)*(x_)^(m_.)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d +
e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2
)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b,
c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3
, 0] && NeQ[m, -1]`

### 3.231.4 Maple [A] (verified)

Time = 0.90 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.94

method	result
default	$-\frac{\sqrt{-a^2x^2+1}(a^2x^2-\sqrt{ax-1}\sqrt{ax+1}ax-1)\operatorname{arccosh}(ax)^2}{x(a^2x^2-1)} - \frac{2\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^2a}{a^2x^2-1} + \frac{2\sqrt{-a^2x^2+1}\sqrt{ax-1}}{a^2x^2-1}$

---

3.231. 
$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx$$

input `int(arccosh(a*x)^2/x^2/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$-(a^2x^2+1)^{1/2}*(a^2x^2-(a*x-1)^{1/2}*(a*x+1)^{1/2})*a*x-1*\operatorname{arccosh}(a*x)^2/x/(a^2x^2-1)-2*(-a^2x^2+1)^{1/2}*(a*x-1)^{1/2}*(a*x+1)^{1/2}/(a^2x^2-1)*\operatorname{arccosh}(a*x)^2*a+2*(-a^2x^2+1)^{1/2}*(a*x-1)^{1/2}*(a*x+1)^{1/2}/(a^2x^2-1)*\operatorname{arccosh}(a*x)*\ln(1+(a*x+(a*x-1)^{1/2}*(a*x+1)^{1/2})^2)*a+(-a^2x^2+1)^{1/2}*(a*x-1)^{1/2}*(a*x+1)^{1/2}/(a^2x^2-1)*\operatorname{polylog}(2,-(a*x+(a*x-1)^{1/2}*(a*x+1)^{1/2}))^2)*a$$

### 3.231.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}x^2} dx$$

input `integrate(arccosh(a*x)^2/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^2/(a^2*x^4 - x^2), x)`

### 3.231.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}^2(ax)}{x^2\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**2/x**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**2/(x**2*sqrt(-(a*x - 1)*(a*x + 1))), x)`

**3.231.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}x^2} dx$$

input `integrate(arccosh(a*x)^2/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `(a^2*x^2 - 1)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^2/(sqrt(a*x + 1)*sqrt(-a*x + 1)*x) - integrate(2*(a^3*x^2 + sqrt(a*x + 1)*sqrt(a*x - 1)*a^2*x - a)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))/((sqrt(a*x + 1)*a*x^2 + (a*x + 1)*sqrt(a*x - 1)*x)*sqrt(-a*x + 1)), x)`

**3.231.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}x^2} dx$$

input `integrate(arccosh(a*x)^2/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^2/(sqrt(-a^2*x^2 + 1)*x^2), x)`

**3.231.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^2}{x^2\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^2/(x^2*(1 - a^2*x^2)^(1/2)),x)`

output `int(acosh(a*x)^2/(x^2*(1 - a^2*x^2)^(1/2)), x)`

### 3.232 $\int \frac{\operatorname{arccosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx$

3.232.1 Optimal result . . . . .	2070
3.232.2 Mathematica [A] (warning: unable to verify) . . . . .	2071
3.232.3 Rubi [A] (verified) . . . . .	2072
3.232.4 Maple [F] . . . . .	2076
3.232.5 Fricas [F] . . . . .	2076
3.232.6 Sympy [F] . . . . .	2076
3.232.7 Maxima [F] . . . . .	2077
3.232.8 Giac [F] . . . . .	2077
3.232.9 Mupad [F(-1)] . . . . .	2077

#### 3.232.1 Optimal result

Integrand size = 24, antiderivative size = 296

$$\begin{aligned} \int \frac{\operatorname{arccosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx = & \frac{a\sqrt{-1+ax}\operatorname{arccosh}(ax)}{x\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} \\ & + \frac{a^2\sqrt{-1+ax}\operatorname{arccosh}(ax)^2 \arctan(e^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\ & - \frac{a^2\sqrt{-1+ax} \arctan(\sqrt{-1+ax}\sqrt{1+ax})}{\sqrt{1-ax}} \\ & - \frac{ia^2\sqrt{-1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\ & + \frac{ia^2\sqrt{-1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\ & + \frac{ia^2\sqrt{-1+ax} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\ & - \frac{ia^2\sqrt{-1+ax} \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \end{aligned}$$

output  $a^2 \operatorname{arccosh}(ax) \sqrt{1-a^2x^2} / (x^3 \sqrt{1-a^2x^2}) + a^2 \operatorname{arccosh}(ax)^2 \arctan(ax) / (x^3 \sqrt{1-a^2x^2}) - a^2 \operatorname{arccosh}(ax) \operatorname{arctan}(ax) / (x^3 \sqrt{1-a^2x^2}) - a^2 \operatorname{arccosh}(ax) \operatorname{arccosh}(ax) / (x^3 \sqrt{1-a^2x^2}) - a^2 \operatorname{arccosh}(ax) \operatorname{polylog}(2, -I \operatorname{arccosh}(ax) \sqrt{1-a^2x^2}) / (x^3 \sqrt{1-a^2x^2}) - a^2 \operatorname{arccosh}(ax) \operatorname{polylog}(2, I \operatorname{arccosh}(ax) \sqrt{1-a^2x^2}) / (x^3 \sqrt{1-a^2x^2}) - a^2 \operatorname{arccosh}(ax) \operatorname{polylog}(3, -I \operatorname{arccosh}(ax) \sqrt{1-a^2x^2}) / (x^3 \sqrt{1-a^2x^2}) - a^2 \operatorname{arccosh}(ax) \operatorname{polylog}(3, I \operatorname{arccosh}(ax) \sqrt{1-a^2x^2}) / (x^3 \sqrt{1-a^2x^2}) - 1/2 \operatorname{arccosh}(ax)^2 (-a^2 x^2 + 1)^{1/2} / x^2$

### 3.232.2 Mathematica [A] (warning: unable to verify)

Time = 0.78 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.79

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^3 \sqrt{1-a^2x^2}} dx$$

$$= \frac{ia^2 \sqrt{-((-1+ax)(1+ax))} \left( \frac{2i \operatorname{arccosh}(ax)}{ax} + \frac{i \sqrt{\frac{-1+ax}{1+ax}} (1+ax) \operatorname{arccosh}(ax)^2}{a^2 x^2} - 4i \arctan \left( \tanh \left( \frac{1}{2} \operatorname{arccosh}(ax) \right) \right) \right)}{1}$$

input `Integrate[ArcCosh[a*x]^2/(x^3*Sqrt[1 - a^2*x^2]),x]`

output  $((I/2) a^2 \sqrt{-((-1+ax)(1+ax))} * (((2*I) \operatorname{ArcCosh}[a*x]) / (a*x) + (I * \sqrt{(-1+ax)/(1+ax)} * (1+ax) * \operatorname{ArcCosh}[a*x]^2) / (a^2 x^2) - (4*I) \operatorname{ArcTan}[\operatorname{Tanh}[\operatorname{ArcCosh}[a*x]/2]] + \operatorname{ArcCosh}[a*x]^2 \operatorname{Log}[1 - I/E^{\operatorname{ArcCosh}[a*x]}] - \operatorname{ArcCosh}[a*x]^2 \operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[a*x]}] + 2 * \operatorname{ArcCosh}[a*x] * \operatorname{PolyLog}[2, (-I)/E^{\operatorname{ArcCosh}[a*x]}] - 2 * \operatorname{ArcCosh}[a*x] * \operatorname{PolyLog}[2, I/E^{\operatorname{ArcCosh}[a*x]}] + 2 * \operatorname{PolyLog}[3, (-I)/E^{\operatorname{ArcCosh}[a*x]}] - 2 * \operatorname{PolyLog}[3, I/E^{\operatorname{ArcCosh}[a*x]}])) / (\sqrt{(-1+ax)/(1+ax)} * (1+ax))$



**3.232.3 Rubi [A] (verified)**

Time = 1.02 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.66, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6347, 6298, 103, 218, 6361, 3042, 4668, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6347} \\
 & \frac{1}{2}a^2 \int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx - \frac{a\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)}{x^2} dx}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} \\
 & \quad \downarrow \text{6298} \\
 & \frac{1}{2}a^2 \int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx - \frac{a\sqrt{ax-1} \left( a \int \frac{1}{x\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\operatorname{arccosh}(ax)}{x} \right)}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} \\
 & \quad \downarrow \text{103} \\
 & \frac{1}{2}a^2 \int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx - \frac{a\sqrt{ax-1} \left( a^2 \int \frac{1}{(ax-1)(ax+1)a+a} d(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x} \right)}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} \\
 & \quad \downarrow \text{218} \\
 & \frac{1}{2}a^2 \int \frac{\operatorname{arccosh}(ax)^2}{x\sqrt{1-a^2x^2}} dx - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} - \frac{a\sqrt{ax-1} \left( a \arctan(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x} \right)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{6361} \\
 & \frac{a^2\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)^2}{ax} d\operatorname{arccosh}(ax)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} - \frac{a\sqrt{ax-1} \left( a \arctan(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x} \right)}{\sqrt{1-ax}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\frac{a^2\sqrt{ax-1} \int \operatorname{arccosh}(ax)^2 \csc\left(i\operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d\operatorname{arccosh}(ax)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} - \frac{a\sqrt{ax-1}\left(a \arctan(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x}\right)}{\sqrt{1-ax}}$$

↓ 4668

$$\frac{a^2\sqrt{ax-1}\left(-2i \int \operatorname{arccosh}(ax) \log(1 - ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2i \int \operatorname{arccosh}(ax) \log(1 + ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax)\right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} - \frac{a\sqrt{ax-1}\left(a \arctan(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x}\right)}{\sqrt{1-ax}}$$

↓ 3011

$$\frac{a^2\sqrt{ax-1}\left(2i\left(\int \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)})\right) - 2i\left(\int \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)})\right)\right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} - \frac{a\sqrt{ax-1}\left(a \arctan(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x}\right)}{\sqrt{1-ax}}$$

↓ 2720

$$\frac{a^2\sqrt{ax-1}\left(2i\left(\int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)})\right) - 2i\left(\int e^{\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)})\right)\right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} - \frac{a\sqrt{ax-1}\left(a \arctan(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x}\right)}{\sqrt{1-ax}}$$

↓ 7143

$$\frac{a^2\sqrt{ax-1}\left(2\operatorname{arccosh}(ax)^2 \arctan(e^{\operatorname{arccosh}(ax)}) + 2i\left(\operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)})\right) - 2i\left(\operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)})\right)\right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}{2x^2} - \frac{a\sqrt{ax-1}\left(a \arctan(\sqrt{ax-1}\sqrt{ax+1}) - \frac{\operatorname{arccosh}(ax)}{x}\right)}{\sqrt{1-ax}}$$

input `Int[ArcCosh[a*x]^2/(x^3*sqrt[1 - a^2*x^2]),x]`

```
output -1/2*(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)/x^2 - (a*Sqrt[-1 + a*x]*(-ArcCosh
[a*x]/x) + a*ArcTan[Sqrt[-1 + a*x]*Sqrt[1 + a*x]]))/Sqrt[1 - a*x] + (a^2*S
qrt[-1 + a*x]*(2*ArcCosh[a*x]^2*ArcTan[E^ArcCosh[a*x]] + (2*I)*(-ArcCosh[
a*x]*PolyLog[2, (-I)*E^ArcCosh[a*x]]) + PolyLog[3, (-I)*E^ArcCosh[a*x]]) -
(2*I)*(-ArcCosh[a*x]*PolyLog[2, I*E^ArcCosh[a*x]]) + PolyLog[3, I*E^ArcC
osh[a*x]])))/(2*Sqrt[1 - a*x])
```

### 3.232.3.1 Defintions of rubi rules used

```
rule 103 Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_
))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sq
rt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d
*e - f*(b*c + a*d), 0]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 3011 Int[Log[1 + (e_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

rule 6361 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

**3.232.4 Maple [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^3 \sqrt{-a^2 x^2 + 1}} dx$$

input `int(arccosh(a*x)^2/x^3/(-a^2*x^2+1)^(1/2),x)`

output `int(arccosh(a*x)^2/x^3/(-a^2*x^2+1)^(1/2),x)`

**3.232.5 Fracas [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^3 \sqrt{1 - a^2 x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2 x^2 + 1} x^3} dx$$

input `integrate(arccosh(a*x)^2/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^2/(a^2*x^5 - x^3), x)`

**3.232.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^3 \sqrt{1 - a^2 x^2}} dx = \int \frac{\operatorname{acosh}^2(ax)}{x^3 \sqrt{-(ax - 1)(ax + 1)}} dx$$

input `integrate(acosh(a*x)**2/x**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**2/(x**3*sqrt(-(a*x - 1)*(a*x + 1))), x)`

**3.232.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}x^3} dx$$

input `integrate(arccosh(a*x)^2/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^2/(sqrt(-a^2*x^2 + 1)*x^3), x)`

**3.232.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^2}{\sqrt{-a^2x^2+1}x^3} dx$$

input `integrate(arccosh(a*x)^2/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^2/(sqrt(-a^2*x^2 + 1)*x^3), x)`

**3.232.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^2}{x^3\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^2/(x^3*(1 - a^2*x^2)^(1/2)),x)`

output `int(acosh(a*x)^2/(x^3*(1 - a^2*x^2)^(1/2)), x)`

### 3.233 $\int (fx)^m (d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2 dx$

3.233.1 Optimal result	2079
3.233.2 Mathematica [N/A]	2080
3.233.3 Rubi [N/A]	2081
3.233.4 Maple [N/A] (verified)	2094
3.233.5 Fricas [N/A]	2094
3.233.6 Sympy [F(-1)]	2094
3.233.7 Maxima [N/A]	2095
3.233.8 Giac [F(-2)]	2095
3.233.9 Mupad [N/A]	2095

### 3.233.1 Optimal result

Integrand size = 31, antiderivative size = 31

$$\begin{aligned}
& \int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = -\frac{10b^2 c^2 d^2 (fx)^{3+m} \sqrt{d - c^2 dx^2}}{f^3 (4+m)^3 (6+m)} \\
& - \frac{2b^2 c^2 d^2 (52 + 15m + m^2) (fx)^{3+m} (1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{f^3 (4+m)^2 (6+m)^3 (1 - cx)(1 + cx)} \\
& + \frac{2b^2 c^4 d^2 (fx)^{5+m} (1 - c^2 x^2) \sqrt{d - c^2 dx^2}}{f^5 (6+m)^3 (1 - cx)(1 + cx)} - \frac{2bcd^2 (fx)^{2+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f^2 (2+m)(6+m) \sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{30bcd^2 (fx)^{2+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f^2 (2+m)^2 (4+m)(6+m) \sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{10bcd^2 (fx)^{2+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f^2 (2+m)(4+m)(6+m) \sqrt{-1 + cx} \sqrt{1 + cx}} \\
& + \frac{10bc^3 d^2 (fx)^{4+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f^4 (4+m)^2 (6+m) \sqrt{-1 + cx} \sqrt{1 + cx}} \\
& + \frac{4bc^3 d^2 (fx)^{4+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f^4 (4+m)(6+m) \sqrt{-1 + cx} \sqrt{1 + cx}} \\
& - \frac{2bc^5 d^2 (fx)^{6+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f^6 (6+m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
& + \frac{15d^2 (fx)^{1+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{f(6+m)(8 + 6m + m^2)} \\
& + \frac{5d (fx)^{1+m} (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2}{f(4+m)(6+m)} \\
& + \frac{(fx)^{1+m} (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2}{f(6+m)} \\
& - \frac{30b^2 c^2 d^2 (fx)^{3+m} \sqrt{1 - c^2 x^2} \sqrt{d - c^2 dx^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2 x^2\right)}{f^3 (2+m)^2 (3+m)(4+m)(6+m)(1 - cx)(1 + cx)} \\
& - \frac{10b^2 c^2 d^2 (10 + 3m) (fx)^{3+m} \sqrt{1 - c^2 x^2} \sqrt{d - c^2 dx^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2 x^2\right)}{f^3 (2+m)(3+m)(4+m)^3 (6+m)(1 - cx)(1 + cx)} \\
& - \frac{2b^2 c^2 d^2 (264 + 130m + 15m^2) (fx)^{3+m} \sqrt{1 - c^2 x^2} \sqrt{d - c^2 dx^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2 x^2\right)}{f^3 (2+m)(3+m)(4+m)^2 (6+m)^3 (1 - cx)(1 + cx)} \\
& + \frac{15d^3 \operatorname{Int}\left(\frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}}, x\right)}{(6+m)(8 + 6m + m^2)}
\end{aligned}$$



output

```

5*d*(f*x)^(1+m)*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2/f/(4+m)/(6+m)+(f
*x)^(1+m)*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2/f/(6+m)-10*b^2*c^2*d^2
*(f*x)^(3+m)*(-c^2*d*x^2+d)^(1/2)/f^3/(4+m)^3/(6+m)-2*b^2*c^2*d^2*(m^2+15*
m+52)*(f*x)^(3+m)*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)/f^3/(4+m)^2/(6+m)^3/(-
c*x+1)/(c*x+1)+2*b^2*c^4*d^2*(f*x)^(5+m)*(-c^2*x^2+1)*(-c^2*d*x^2+d)^(1/2)
/f^5/(6+m)^3/(-c*x+1)/(c*x+1)+15*d^2*(f*x)^(1+m)*(a+b*arccosh(c*x))^2*(-c^
2*d*x^2+d)^(1/2)/f/(6+m)/(m^2+6*m+8)-2*b*c*d^2*(f*x)^(2+m)*(a+b*arccosh(c*
x))*(-c^2*d*x^2+d)^(1/2)/f^2/(2+m)/(6+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-30*b*
c*d^2*(f*x)^(2+m)*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/f^2/(2+m)^2/(4+m)
)/(6+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-10*b*c*d^2*(f*x)^(2+m)*(a+b*arccosh(c*
x))*(-c^2*d*x^2+d)^(1/2)/f^2/(6+m)/(m^2+6*m+8)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
+10*b*c^3*d^2*(f*x)^(4+m)*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/f^4/(4+m)
)^2/(6+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+4*b*c^3*d^2*(f*x)^(4+m)*(a+b*arccosh
(c*x))*(-c^2*d*x^2+d)^(1/2)/f^4/(4+m)/(6+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2*
b*c^5*d^2*(f*x)^(6+m)*(a+b*arccosh(c*x))*(-c^2*d*x^2+d)^(1/2)/f^6/(6+m)^2/
(c*x-1)^(1/2)/(c*x+1)^(1/2)-30*b^2*c^2*d^2*(f*x)^(3+m)*hypergeom([1/2, 3/2
+1/2*m], [5/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)*(-c^2*d*x^2+d)^(1/2)/f^3/(
2+m)^2/(3+m)/(4+m)/(6+m)/(-c*x+1)/(c*x+1)-10*b^2*c^2*d^2*(10+3*m)*(f*x)^(3
+m)*hypergeom([1/2, 3/2+1/2*m], [5/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)*(-c
^2*d*x^2+d)^(1/2)/f^3/(4+m)^3/(6+m)/(m^2+5*m+6)/(-c*x+1)/(c*x+1)-2*b^2*...

```

### 3.233.2 Mathematica [N/A]

Not integrable

Time = 2.23 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2 dx = \int (fx)^m (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^2 dx$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output `Integrate[(f*x)^m*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2, x]`

**3.233.3 Rubi [N/A]**

Not integrable

Time = 5.21 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 25, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6345, 6327, 6336, 27, 1905, 1590, 27, 363, 279, 278, 6345, 25, 6327, 6336, 27, 960, 136, 279, 278, 6345, 6298, 136, 279, 278, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d - c^2 dx^2)^{5/2} (fx)^m (a + \operatorname{barccosh}(cx))^2 dx \\
 & \quad \downarrow \text{6345} \\
 & -\frac{2bcd^2 \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1 - cx)^2 (cx + 1)^2 (a + \operatorname{barccosh}(cx)) dx}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} + \\
 & \frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{m+6} + \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} \\
 & \quad \downarrow \text{6327} \\
 & -\frac{2bcd^2 \sqrt{d - c^2 dx^2} \int (fx)^{m+1} (1 - c^2 x^2)^2 (a + \operatorname{barccosh}(cx)) dx}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} + \\
 & \frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{m+6} + \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} \\
 & \quad \downarrow \text{6336} \\
 & \frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{m+6} - \\
 & \frac{2bcd^2 \sqrt{d - c^2 dx^2} \left( -bc \int \frac{(fx)^{m+2} \left( \frac{c^4 x^4}{m+6} - \frac{2c^2 x^2}{m+4} + \frac{1}{m+2} \right) dx}{f \sqrt{cx-1} \sqrt{cx+1}} + \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5 (m+6)} - \frac{2c^2 (fx)^{m+4} (a + \operatorname{barccosh}(cx))}{f^3 (m+4)} + \dots \right)}{f(m+6) \sqrt{cx-1} \sqrt{cx+1}} \\
 & \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{2bcd^2 \sqrt{d - c^2 dx^2}} - \\
 & \left( - \frac{bc \int \frac{(fx)^{m+2} \left( \frac{c^4 x^4}{m+6} - \frac{2c^2 x^2}{m+4} + \frac{1}{m+2} \right) dx}{\sqrt{cx-1}\sqrt{cx+1}}}{f} + \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5 (m+6)} - \frac{2c^2 (fx)^{m+4} (a + \operatorname{barccosh}(cx))}{f^3 (m+4)} + \frac{(fx)^{m+2} (a + \operatorname{barccosh}(cx))^2}{f} \right) \\
 & \frac{f(m+6)\sqrt{cx-1}\sqrt{cx+1}}{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2} \\
 & \frac{f(m+6)}{f(m+6)} \\
 & \downarrow \text{1905} \\
 & \frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{2bcd^2 \sqrt{d - c^2 dx^2}} - \\
 & \left( - \frac{bc\sqrt{c^2 x^2 - 1} \int \frac{(fx)^{m+2} \left( \frac{c^4 x^4}{m+6} - \frac{2c^2 x^2}{m+4} + \frac{1}{m+2} \right) dx}{f\sqrt{cx-1}\sqrt{cx+1}}}{f\sqrt{cx-1}\sqrt{cx+1}} + \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5 (m+6)} - \frac{2c^2 (fx)^{m+4} (a + \operatorname{barccosh}(cx))}{f^3 (m+4)} + \frac{(fx)^{m+2} (a + \operatorname{barccosh}(cx))^2}{f} \right) \\
 & \frac{f(m+6)\sqrt{cx-1}\sqrt{cx+1}}{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2} \\
 & \frac{f(m+6)}{f(m+6)} \\
 & \downarrow \text{1590} \\
 & \frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{2bcd^2 \sqrt{d - c^2 dx^2}} - \\
 & \left( - \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{c^2 (fx)^{m+2} \left( \frac{m+6}{m+2} - \frac{c^2 (m^2 + 15m + 52)x^2}{(m+4)(m+6)} \right) dx}{\frac{\sqrt{c^2 x^2 - 1}}{c^2 (m+6)}} + \frac{c^2 \sqrt{c^2 x^2 - 1} (fx)^{m+5}}{f^3 (m+6)^2} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} + \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5 (m+6)} \right) \\
 & \frac{f(m+6)\sqrt{cx-1}\sqrt{cx+1}}{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2} \\
 & \frac{f(m+6)}{f(m+6)} \\
 & \downarrow \text{27}
 \end{aligned}$$

$$\frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{m+6} - \frac{2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{bc\sqrt{c^2 x^2 - 1} \left( \int \frac{(fx)^{m+2} \left( \frac{m+6}{m+2} - \frac{c^2(m^2+15m+52)x^2}{(m+4)(m+6)} \right) dx}{\sqrt{c^2 x^2 - 1}} + \frac{c^2 \sqrt{c^2 x^2 - 1} (fx)^{m+5}}{f^3 (m+6)^2} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} + \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5 (m+6)}$$

$$\frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)}$$

↓ 363

$$\frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{m+6} - \frac{2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{(15m^2+130m+264) \int \frac{(fx)^{m+2}}{\sqrt{c^2 x^2 - 1}} dx}{(m+2)(m+4)^2(m+6)} - \frac{(m^2+15m+52) \sqrt{c^2 x^2 - 1} (fx)^{m+3}}{f(m+4)^2(m+6)} + \frac{c^2 \sqrt{c^2 x^2 - 1} (fx)^{m+5}}{f^3 (m+6)^2} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} + \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5 (m+6)}$$

$$\frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)}$$

↓ 279

$$\frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{m+6} - \frac{2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{bc\sqrt{c^2 x^2 - 1} \left( \frac{(15m^2+130m+264) \sqrt{1-c^2 x^2} \int \frac{(fx)^{m+2}}{\sqrt{1-c^2 x^2}} dx}{(m+2)(m+4)^2(m+6)\sqrt{c^2 x^2 - 1}} - \frac{(m^2+15m+52) \sqrt{c^2 x^2 - 1} (fx)^{m+3}}{f(m+4)^2(m+6)} + \frac{c^2 \sqrt{c^2 x^2 - 1} (fx)^{m+5}}{f^3 (m+6)^2} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} + \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5 (m+6)}$$

$$\frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)}$$

↓ 278

3.233.  $\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\frac{5d \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx}{m+6} + \frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4} (a + \operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2 x^2 - 1}}{f(m+6)\sqrt{cx-1}} \left( \frac{c^2}{f(m+6)\sqrt{cx-1}} \right) \right)$$


---

↓ 6345

$$5d \left( \frac{2bcd\sqrt{d-c^2 dx^2} \int -(fx)^{m+1}(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \frac{3d \int (fx)^m \sqrt{d-c^2 dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2 dx^2)^{3/2} (fx)^m}{f(m+6)} \right)$$

$$\frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4} (a + \operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2 x^2 - 1}}{f(m+6)\sqrt{cx-1}} \left( \frac{c^2}{f(m+6)\sqrt{cx-1}} \right) \right)$$


---

↓ 25

$$5d \left( -\frac{2bcd\sqrt{d-c^2 dx^2} \int (fx)^{m+1}(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \frac{3d \int (fx)^m \sqrt{d-c^2 dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2 dx^2)^{3/2} (fx)^m}{f(m+6)} \right)$$

$$\frac{(d - c^2 dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{c^4 (fx)^{m+6} (a + \operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4} (a + \operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2 x^2 - 1}}{f(m+6)\sqrt{cx-1}} \left( \frac{c^2}{f(m+6)\sqrt{cx-1}} \right) \right)$$


---

↓ 6327

---

3.233.  $\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$5d \left( -\frac{2bcd\sqrt{d-c^2dx^2} \int (fx)^{m+1}(1-c^2x^2)(a+\operatorname{barccosh}(cx))dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)} \right)$$


---


$$\frac{(d-c^2dx^2)^{5/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+6)} -$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4 (fx)^{m+6} (a+\operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1} \left( \frac{c^2}{f(m+2)} \right)}{f(m+2)} \right)$$


---


$$f(m+6)\sqrt{cx-1}$$

↓ 6336

$$5d \left( -\frac{2bcd\sqrt{d-c^2dx^2} \left( -bc \int \frac{(fx)^{m+2} \left( \frac{1}{m+2} - \frac{c^2x^2}{m+4} \right) dx}{f\sqrt{cx-1}\sqrt{cx+1}} - \frac{c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} \right)$$


---


$$\frac{(d-c^2dx^2)^{5/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+6)} -$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4 (fx)^{m+6} (a+\operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1} \left( \frac{c^2}{f(m+2)} \right)}{f(m+2)} \right)$$


---


$$f(m+6)\sqrt{cx-1}$$

↓ 27

$$5d \left( - \frac{2bcd\sqrt{d-c^2dx^2} \left( - \frac{bc \int \frac{(fx)^{m+2} \left( \frac{1}{m+2} - \frac{c^2x^2}{m+4} \right) dx}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{c^2(fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} \right) + \frac{3d \int (fx)^m \sqrt{d-c^2dx^2}}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} \right)$$

---


$$\frac{(d - c^2dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} - \frac{m+6}{f(m+6)}$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4(fx)^{m+6} (a+\operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2(fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1}}{f(m+2)} \left( \frac{c^2}{f(m+2)} \right) \right)$$


---

$f(m+6)\sqrt{cx-1}$

↓ 960

$$5d \left( - \frac{2bcd\sqrt{d-c^2dx^2} \left( - \frac{bc \left( \frac{(3m+10) \int \frac{(fx)^{m+2}}{\sqrt{cx-1}\sqrt{cx+1}} dx}{(m+2)(m+4)^2} - \frac{\sqrt{cx-1}\sqrt{cx+1} (fx)^{m+3}}{f(m+4)^2} \right)}{f} - \frac{c^2(fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} \right)$$

---


$$\frac{(d - c^2dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+6)} - \frac{m+6}{f(m+6)}$$

$$2bcd^2\sqrt{d-c^2dx^2} \left( \frac{c^4(fx)^{m+6} (a+\operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2(fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1}}{f(m+2)} \left( \frac{c^2}{f(m+2)} \right) \right)$$


---

$f(m+6)\sqrt{cx-1}$

↓ 136

---

3.233.  $\int (fx)^m (d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$5d \left( \frac{2bcd\sqrt{d-c^2dx^2} \left( bc \left( \frac{(3m+10)\sqrt{c^2x^2-1} \int \frac{(fx)^{m+2}}{\sqrt{c^2x^2-1}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+3}}{f(m+4)^2} \right)}{f} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} - \frac{c^2(fx)^{m+4}(a+b\operatorname{arccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+b\operatorname{arccosh}(cx))}{f(m+2)} \right)$$

$$\frac{(d - c^2dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{arccosh}(cx))^2}{f(m+6)} - \frac{2bcd^2\sqrt{d - c^2dx^2} \left( \frac{c^4(fx)^{m+6}(a+b\operatorname{arccosh}(cx))}{f^5(m+6)} - \frac{2c^2(fx)^{m+4}(a+b\operatorname{arccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+b\operatorname{arccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1} \left( \frac{c^2}{f^3(m+4)} + \frac{(fx)^{m+2}(a+b\operatorname{arccosh}(cx))}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}} \right)}{f(m+6)\sqrt{cx-1}}$$

↓ 279

$$5d \left( \frac{2bcd\sqrt{d-c^2dx^2} \left( bc \left( \frac{(3m+10)\sqrt{1-c^2x^2} \int \frac{(fx)^{m+2}}{\sqrt{1-c^2x^2}} dx - \frac{\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+3}}{f(m+4)^2} \right)}{f} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} - \frac{c^2(fx)^{m+4}(a+b\operatorname{arccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+b\operatorname{arccosh}(cx))}{f(m+2)} \right)$$

$$\frac{(d - c^2dx^2)^{5/2} (fx)^{m+1} (a + \operatorname{arccosh}(cx))^2}{f(m+6)} - \frac{2bcd^2\sqrt{d - c^2dx^2} \left( \frac{c^4(fx)^{m+6}(a+b\operatorname{arccosh}(cx))}{f^5(m+6)} - \frac{2c^2(fx)^{m+4}(a+b\operatorname{arccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+b\operatorname{arccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1} \left( \frac{c^2}{f^3(m+4)} + \frac{(fx)^{m+2}(a+b\operatorname{arccosh}(cx))}{f(m+2)} \right)}{f(m+6)\sqrt{cx-1}} \right)}{f(m+6)\sqrt{cx-1}}$$

↓ 278

3.233.  $\int (fx)^m (d - c^2dx^2)^{5/2} (a + \operatorname{arccosh}(cx))^2 dx$



$$5d \left( \frac{3d \int (fx)^m \sqrt{d-c^2 dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} - \frac{2bcd\sqrt{d-c^2 dx^2} \left( -\frac{c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+1)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)$$

$m+6$

$$\frac{(d-c^2 dx^2)^{5/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+6)} - 2bcd^2 \sqrt{d-c^2 dx^2} \left( \frac{c^4 (fx)^{m+6} (a+\operatorname{barccosh}(cx))}{f^5(m+6)} - \frac{2c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2 x^2-1} \left( \frac{c^2}{f(m+6)\sqrt{cx-1}} \right)}{f(m+6)\sqrt{cx-1}} \right)$$

$f(m+6)\sqrt{cx-1}$

↓ 6345

$$\frac{(d-c^2 dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+6)} - 2bcd^2 \sqrt{d-c^2 dx^2} \left( \frac{(a+\operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{2c^2 (a+\operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} + \frac{c^4 (a+\operatorname{barccosh}(cx))(fx)^{m+6}}{f^5(m+6)} - \frac{bc\sqrt{c^2 x^2-1} \left( \frac{c^2}{f(m+6)\sqrt{cx-1}} \right)}{f(m+6)\sqrt{cx-1}} \right)$$

$f(m+6)\sqrt{cx-1}$

$$5d \left( \frac{(d-c^2 dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+4)} - \frac{2bcd\sqrt{d-c^2 dx^2} \left( \frac{(a+\operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{c^2 (a+\operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} - \frac{bc \left( \frac{(3m+10)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)$$

↓ 6298

3.233.  $\int (fx)^m (d-c^2 dx^2)^{5/2} (a+\operatorname{barccosh}(cx))^2 dx$

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{2c^2(a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} + \frac{c^4(a + \operatorname{barccosh}(cx))(fx)^{m+6}}{f^5(m+6)} - \frac{bc\sqrt{c^2 x^2 - 1}}{c^2} \right)$$


---


$$5d \left( \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+4)} - \frac{2bcd\sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{c^2(a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} - \frac{bc \left( \frac{(3m+10)}{c^2} \right)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)$$

↓ 136

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{2c^2(a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} + \frac{c^4(a + \operatorname{barccosh}(cx))(fx)^{m+6}}{f^5(m+6)} - \frac{bc\sqrt{c^2 x^2 - 1}}{c^2} \right)$$


---


$$5d \left( \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+4)} - \frac{2bcd\sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{c^2(a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} - \frac{bc \left( \frac{(3m+10)}{c^2} \right)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)$$

↓ 279

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{2c^2 (a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} + \frac{c^4 (a + \operatorname{barccosh}(cx))(fx)^{m+6}}{f^5(m+6)} - \frac{bc\sqrt{c^2 x^2 - 1}}{f(m+6)\sqrt{cx-1}} \left( \frac{c^2}{f(m+4)\sqrt{cx-1}} \right) \right)$$


---


$$5d \left( \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+4)} - \frac{2bcd\sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{c^2 (a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} - \frac{bc \left( \frac{(3m+10)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)$$

↓ 278

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{2c^2 (a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} + \frac{c^4 (a + \operatorname{barccosh}(cx))(fx)^{m+6}}{f^5(m+6)} - \frac{bc\sqrt{c^2 x^2 - 1}}{f(m+6)\sqrt{cx-1}} \left( \frac{c^2}{f(m+4)\sqrt{cx-1}} \right) \right)$$


---


$$5d \left( \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+4)} - \frac{2bcd\sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{c^2 (a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} - \frac{bc \left( \frac{(3m+10)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)$$

↓ 6375

$$\frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+6)} -$$

$$2bcd^2 \sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{2c^2 (a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} + \frac{c^4 (a + \operatorname{barccosh}(cx))(fx)^{m+6}}{f^5(m+6)} - \frac{bc\sqrt{c^2 x^2 - 1}}{c^2} \left( \frac{c^2}{f(m+6)\sqrt{cx-1}} \right) \right)$$


---


$$5d \left( \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 (fx)^{m+1}}{f(m+4)} - \frac{2bcd\sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))(fx)^{m+2}}{f(m+2)} - \frac{c^2 (a + \operatorname{barccosh}(cx))(fx)^{m+4}}{f^3(m+4)} - \frac{bc \left( \frac{(3m+10)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)}{f(m+4)\sqrt{cx-1}} \right)$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

### 3.233.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 136 `Int[((f_)*(x_))^(p_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[(a*c + b*d*x^2)^m*(f*x)^p, x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a), x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

- rule 279  $\text{Int}[(c_*)(x_*)^{(m_*)}((a_*) + (b_*)(x_*)^2)^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[a^{\text{IntPart}[p]} * ((a + b*x^2)^{\text{FracPart}[p]} / (1 + b*(x^2/a))^{\text{FracPart}[p]}) \text{Int}[(c*x)^m (1 + b*(x^2/a))^p, x], x] /;$  FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])
- rule 363  $\text{Int}[(e_*)(x_*)^{(m_*)}((a_*) + (b_*)(x_*)^2)^{(p_*)}((c_*) + (d_*)(x_*)^2), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}((a + b*x^2)^{(p+1}) / (b*e*(m+2*p+3))), x] - \text{Simp}[(a*d*(m+1) - b*c*(m+2*p+3)) / (b*(m+2*p+3)) \text{Int}[(e*x)^m (a + b*x^2)^p, x], x] /;$  FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b\*c - a\*d, 0] && NeQ[m + 2\*p + 3, 0]
- rule 960  $\text{Int}[(e_*)(x_*)^{(m_*)}((a1_*) + (b1_*)(x_*)^{(non2_*)})^{(p_*)}((a2_*) + (b2_*)(x_*)^{(non2_*)})^{(p_*)}((c_*) + (d_*)(x_*)^{(n_*)}), x\_Symbol] \rightarrow \text{Simp}[d*(e*x)^{(m+1)}(a1 + b1*x^{(n/2)})^{(p+1)}((a2 + b2*x^{(n/2)})^{(p+1)} / (b1*b2*e*(m+n*(p+1)+1))), x] - \text{Simp}[(a1*a2*d*(m+1) - b1*b2*c*(m+n*(p+1)+1)) / (b1*b2*(m+n*(p+1)+1)) \text{Int}[(e*x)^m (a1 + b1*x^{(n/2)})^p (a2 + b2*x^{(n/2)})^p, x], x] /;$  FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2\*b1 + a1\*b2, 0] && NeQ[m + n\*(p+1) + 1, 0]
- rule 1590  $\text{Int}[(f_*)(x_*)^{(m_*)}((d_*) + (e_*)(x_*)^2)^{(q_*)}((a_*) + (b_*)(x_*)^2 + (c_*)(x_*)^4)^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[c^p * (f*x)^{(m+4*p-1)} * ((d + e*x^2)^{(q+1)} / (e*f^{(4*p-1)} * (m+4*p+2*q+1))), x] + \text{Simp}[1 / (e*(m+4*p+2*q+1)) \text{Int}[(f*x)^m * (d + e*x^2)^q * \text{ExpandToSum}[e*(m+4*p+2*q+1) * ((a + b*x^2 + c*x^4)^p - c^p * x^{(4*p)}) - d*c^p * (m+4*p-1) * x^{(4*p-2)}, x], x], x] /;$  FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4\*a\*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4\*p + 2\*q + 1, 0]
- rule 1905  $\text{Int}[(f_*)(x_*)^{(m_*)}((d1_*) + (e1_*)(x_*)^{(non2_*)})^{(q_*)}((d2_*) + (e2_*)(x_*)^{(non2_*)})^{(q_*)}((a_*) + (b_*)(x_*)^{(n_*)} + (c_*)(x_*)^{(n2_*)})^{(p_*)}, x\_Symbol] \rightarrow \text{Simp}[(d1 + e1*x^{(n/2)})^{\text{FracPart}[q]} * ((d2 + e2*x^{(n/2)})^{\text{FracPart}[q]} / (d1*d2 + e1*e2*x^n)^{\text{FracPart}[q]}) \text{Int}[(f*x)^m * (d1*d2 + e1*e2*x^n)^q * (a + b*x^n + c*x^{(2*n)})^p, x], x] /;$  FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p}, q}, x] && EqQ[n2, 2\*n] && EqQ[non2, n/2] && EqQ[d2\*e1 + d1\*e2, 0]

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +
c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& NeQ[m, -1]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_ + (
e1_.)*(x_))^(p_.)*((d2_ + (e2_.)*(x_))^(p_.), x_Symbol] :> Int[(f*x)^m*(d1
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_ + (e_.)*(x_
)^2)^(p_.), x_Symbol] :> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_ + (e_
.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*
x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m
+ 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m
+ 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1)
, x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n,
0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_ + (e
_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*A
rcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.233.4 Maple [N/A] (verified)**

Not integrable

Time = 2.79 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int (fx)^m (-c^2 dx^2 + d)^{5/2} (a + b \operatorname{arccosh}(cx))^2 dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x)`output `int((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x)`**3.233.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 136, normalized size of antiderivative = 4.39

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)^2 (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")`output `integral((a^2*c^4*d^2*x^4 - 2*a^2*c^2*d^2*x^2 + a^2*d^2 + (b^2*c^4*d^2*x^4 - 2*b^2*c^2*d^2*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*c^4*d^2*x^4 - 2*a*b*c^2*d^2*x^2 + a*b*d^2)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)*(f*x)^m, x)`**3.233.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2 dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**2,x)`output `Timed out`

**3.233.7 Maxima [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^2 (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^2*(f*x)^m, x)`

**3.233.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.233.9 Mupad [N/A]**

Not integrable

Time = 3.22 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{5/2} (fx)^m dx$$

input `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(5/2)*(f*x)^m, x)`

---

3.233.  $\int (fx)^m (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^2 dx$



### 3.234 $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^2 dx$

3.234.1 Optimal result . . . . .	2096
3.234.2 Mathematica [N/A] . . . . .	2097
3.234.3 Rubi [N/A] . . . . .	2097
3.234.4 Maple [N/A] (verified) . . . . .	2104
3.234.5 Fricas [N/A] . . . . .	2104
3.234.6 Sympy [F(-1)] . . . . .	2104
3.234.7 Maxima [N/A] . . . . .	2105
3.234.8 Giac [F(-2)] . . . . .	2105
3.234.9 Mupad [N/A] . . . . .	2105

#### 3.234.1 Optimal result

Integrand size = 31, antiderivative size = 31

$$\begin{aligned} \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^2 dx = & -\frac{2b^2 c^2 d (fx)^{3+m} \sqrt{d - c^2 dx^2}}{f^3 (4 + m)^3} \\ & - \frac{6bcd (fx)^{2+m} \sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx))}{f^2 (2 + m)^2 (4 + m) \sqrt{-1 + cx} \sqrt{1 + cx}} - \frac{2bcd (fx)^{2+m} \sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx))}{f^2 (2 + m) (4 + m) \sqrt{-1 + cx} \sqrt{1 + cx}} \\ & + \frac{2bc^3 d (fx)^{4+m} \sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx))}{f^4 (4 + m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\ & + \frac{3d (fx)^{1+m} \sqrt{d - c^2 dx^2} (a + \text{barccosh}(cx))^2}{f (8 + 6m + m^2)} + \frac{(fx)^{1+m} (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^2}{f (4 + m)} \\ & - \frac{6b^2 c^2 d (fx)^{3+m} \sqrt{1 - c^2 x^2} \sqrt{d - c^2 dx^2} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2 x^2\right)}{f^3 (2 + m)^2 (3 + m) (4 + m) (1 - cx) (1 + cx)} \\ & - \frac{2b^2 c^2 d (10 + 3m) (fx)^{3+m} \sqrt{1 - c^2 x^2} \sqrt{d - c^2 dx^2} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2 x^2\right)}{f^3 (2 + m) (3 + m) (4 + m)^3 (1 - cx) (1 + cx)} \\ & + \frac{3d^2 \text{Int}\left(\frac{(fx)^m (a + \text{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}}, x\right)}{8 + 6m + m^2} \end{aligned}$$

output  $(f*x)^{(1+m)}*(-c^2*d*x^2+d)^{(3/2)}*(a+b*\operatorname{arccosh}(c*x))^2/f/(4+m)-2*b^2*c^2*d*(f*x)^{(3+m)}*(-c^2*d*x^2+d)^{(1/2)}/f^3/(4+m)^3+3*d*(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/f/(m^2+6*m+8)-6*b*c*d*(f*x)^{(2+m)}*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)^2/(4+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2*b*c*d*(f*x)^{(2+m)}*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)/(4+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+2*b*c^3*d*(f*x)^{(4+m)}*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/f^4/(4+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-6*b^2*c^2*d*(f*x)^{(3+m)}*\operatorname{hypergeom}([1/2, 3/2+1/2*m], [5/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^{(1/2)}*(-c^2*d*x^2+d)^{(1/2)}/f^3/(2+m)^2/(3+m)/(4+m)/(-c*x+1)/(c*x+1)-2*b^2*c^2*d*(10+3*m)*(f*x)^{(3+m)}*\operatorname{hypergeom}([1/2, 3/2+1/2*m], [5/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^{(1/2)}*(-c^2*d*x^2+d)^{(1/2)}/f^3/(4+m)^3/(m^2+5*m+6)/(-c*x+1)/(c*x+1)+3*d^2*\operatorname{Unintegrable}((f*x)^m*(a+b*\operatorname{arccosh}(c*x))^2/(-c^2*d*x^2+d)^{(1/2)}, x)/(m^2+6*m+8)$

### 3.234.2 Mathematica [N/A]

Not integrable

Time = 1.10 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

output `Integrate[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2, x]`

### 3.234.3 Rubi [N/A]

Not integrable

Time = 2.41 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6345, 25, 6327, 6336, 27, 960, 136, 279, 278, 6345, 6298, 136, 279, 278, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{3/2} (fx)^m (a + \operatorname{barccosh}(cx))^2 dx$$

---

3.234.  $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
 & \downarrow 6345 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \int -(fx)^{m+1}(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \downarrow 25 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \int (fx)^{m+1}(1-cx)(cx+1)(a+\operatorname{barccosh}(cx))dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \downarrow 6327 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \int (fx)^{m+1} (1-c^2x^2) (a+\operatorname{barccosh}(cx))dx}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \downarrow 6336 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \left( -bc \int \frac{(fx)^{m+2} \left( \frac{1}{m+2} - \frac{c^2x^2}{m+4} \right) dx}{f\sqrt{cx-1}\sqrt{cx+1}} - \frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \downarrow 27 \\
 & \frac{2bcd\sqrt{d-c^2dx^2} \left( -\frac{bc \int \frac{(fx)^{m+2} \left( \frac{1}{m+2} - \frac{c^2x^2}{m+4} \right) dx}{\sqrt{cx-1}\sqrt{cx+1}}}{f} - \frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} \right)}{f(m+4)\sqrt{cx-1}\sqrt{cx+1}} + \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \downarrow 960
 \end{aligned}$$

---

3.234.  $\int (fx)^m (d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
 & 2bcd\sqrt{d-c^2dx^2} \left( -\frac{bc \left( \frac{(3m+10) \int \frac{(fx)^{m+2}}{\sqrt{cx-1}\sqrt{cx+1}} dx}{(m+2)(m+4)^2} - \frac{\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+3}}{f(m+4)^2} \right)}{f} - \frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} \right) \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}} \frac{(fx)^{m+1}(a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \quad \downarrow 136 \\
 & 2bcd\sqrt{d-c^2dx^2} \left( -\frac{bc \left( \frac{(3m+10)\sqrt{c^2x^2-1} \int \frac{(fx)^{m+2}}{\sqrt{c^2x^2-1}} dx}{(m+2)(m+4)^2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+3}}{f(m+4)^2} \right)}{f} - \frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} \right) \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}} \frac{(fx)^{m+1}(a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \quad \downarrow 279 \\
 & 2bcd\sqrt{d-c^2dx^2} \left( -\frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2} \int \frac{(fx)^{m+2}}{\sqrt{1-c^2x^2}} dx}{(m+2)(m+4)^2\sqrt{cx-1}\sqrt{cx+1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(fx)^{m+3}}{f(m+4)^2} \right)}{f} - \frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} \right) \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} + \frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}} \frac{(fx)^{m+1}(a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \quad \downarrow 278 \\
 & \frac{3d \int (fx)^m \sqrt{d-c^2dx^2} (a+\operatorname{barccosh}(cx))^2 dx}{m+4} - \frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2}(fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \dots\right)}{f(m+2)(m+3)(m+4)^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{f} \\
 & \frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2}} \frac{(fx)^{m+1}(a+\operatorname{barccosh}(cx))^2}{f(m+4)} \\
 & \quad \downarrow 6345
 \end{aligned}$$

3.234.  $\int (fx)^m (d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
 & 3d \left( -\frac{2bc\sqrt{d-c^2dx^2} \int (fx)^{m+1} (a+\operatorname{barccosh}(cx)) dx}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{m+2} + \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+2)} \right) \\
 & \frac{m+4}{2bcd\sqrt{d-c^2dx^2} \left( -\frac{c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2} (fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \dots\right)}{f(m+2)(m+3)(m+4)^2 \sqrt{cx-1}\sqrt{cx+1}} \right)}{f} \right)} \\
 & \frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2} \\
 & \frac{f(m+4)}{f(m+4)}
 \end{aligned}$$

↓ 6298

$$\begin{aligned}
 & 3d \left( -\frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \int \frac{(fx)^{m+2}}{\sqrt{cx-1}\sqrt{cx+1}} dx}{f(m+2)} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{m+2} + \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+2)} \right) \\
 & \frac{m+4}{2bcd\sqrt{d-c^2dx^2} \left( -\frac{c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2} (fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \dots\right)}{f(m+2)(m+3)(m+4)^2 \sqrt{cx-1}\sqrt{cx+1}} \right)}{f} \right)} \\
 & \frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2} \\
 & \frac{f(m+4)}{f(m+4)}
 \end{aligned}$$

↓ 136

$$\begin{aligned}
 & 3d \left( -\frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1} \int \frac{(fx)^{m+2}}{\sqrt{c^2x^2-1}} dx}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2 dx}{\sqrt{d-c^2dx^2}}}{m+2} + \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+2)} \right) \\
 & \frac{m+4}{2bcd\sqrt{d-c^2dx^2} \left( -\frac{c^2 (fx)^{m+4} (a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2} (a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2} (fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \dots\right)}{f(m+2)(m+3)(m+4)^2 \sqrt{cx-1}\sqrt{cx+1}} \right)}{f} \right)} \\
 & \frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1}}{(d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2} \\
 & \frac{f(m+4)}{f(m+4)}
 \end{aligned}$$

↓ 279

---

3.234.  $\int (fx)^m (d-c^2dx^2)^{3/2} (a+\operatorname{barccosh}(cx))^2 dx$

$$3d \left( \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f^{(m+2)}} - \frac{bc\sqrt{1-c^2x^2} \int \frac{(fx)^{m+2}}{\sqrt{1-c^2x^2}} dx}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{m+2} + \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))}{f(m+2)} \right)$$

$$2bcd\sqrt{d-c^2dx^2} \left( -\frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2}(fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+3}{2}, \frac{m+5}{2}, c^2x^2\right)}{f(m+2)(m+3)(m+4)^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{f} \right)$$

$$\frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1} (d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)}$$

↓ 278

$$3d \left( \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{m+2} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{1-c^2x^2}(fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+3}{2}, \frac{m+5}{2}, c^2x^2\right)}{f^2(m+2)(m+3)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$2bcd\sqrt{d-c^2dx^2} \left( -\frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2}(fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+3}{2}, \frac{m+5}{2}, c^2x^2\right)}{f(m+2)(m+3)(m+4)^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{f} \right)$$

$$\frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1} (d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)}$$

↓ 6375

$$3d \left( \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{m+2} - \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{1-c^2x^2}(fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+3}{2}, \frac{m+5}{2}, c^2x^2\right)}{f^2(m+2)(m+3)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)$$

$$2bcd\sqrt{d-c^2dx^2} \left( -\frac{c^2(fx)^{m+4}(a+\operatorname{barccosh}(cx))}{f^3(m+4)} + \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \left( \frac{(3m+10)\sqrt{1-c^2x^2}(fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+3}{2}, \frac{m+5}{2}, c^2x^2\right)}{f(m+2)(m+3)(m+4)^2\sqrt{cx-1}\sqrt{cx+1}} \right)}{f} \right)$$

$$\frac{f(m+4)\sqrt{cx-1}\sqrt{cx+1} (d-c^2dx^2)^{3/2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+4)}$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2,x]`

3.234.  $\int (fx)^m (d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

output \$Aborted

### 3.234.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 136 `Int[((f_)*(x_))^(p_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[(a*c + b*d*x^2)^m*(f*x)^p, x], x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m]`

rule 278 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 279 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

rule 960 `Int[((e_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +
c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& NeQ[m, -1]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (
e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] :> Int[(f*x)^m*(d1
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6336 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_
)^2)^(p_.), x_Symbol] :> With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp
[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && E
qQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*Arc
Cosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*
x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m
+ 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m
+ 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1)
, x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n,
0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*A
rcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`



**3.234.4 Maple [N/A] (verified)**

Not integrable

Time = 2.77 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int (fx)^m (-c^2 dx^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^2 dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x)`

**3.234.5 Fracas [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.81

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^2 (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output `integral(-(a^2*c^2*d*x^2 - a^2*d + (b^2*c^2*d*x^2 - b^2*d)*arccosh(c*x)^2 + 2*(a*b*c^2*d*x^2 - a*b*d)*arccosh(c*x))*sqrt(-c^2*d*x^2 + d)*(f*x)^m, x)`

**3.234.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2 dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.234.7 Maxima [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2 (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^2*(f*x)^m, x)`

**3.234.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.234.9 Mupad [N/A]**

Not integrable

Time = 3.44 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (d - c^2 dx^2)^{3/2} (fx)^m dx$$

input `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(3/2)*(f*x)^m, x)`

---

3.234.  $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^2 dx$

### 3.235 $\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$

3.235.1 Optimal result	2106
3.235.2 Mathematica [N/A]	2107
3.235.3 Rubi [N/A]	2107
3.235.4 Maple [N/A] (verified)	2110
3.235.5 Fricas [N/A]	2110
3.235.6 Sympy [N/A]	2110
3.235.7 Maxima [N/A]	2111
3.235.8 Giac [F(-2)]	2111
3.235.9 Mupad [N/A]	2111

#### 3.235.1 Optimal result

Integrand size = 31, antiderivative size = 31

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$$

$$= -\frac{2bc(fx)^{2+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))}{f^2(2+m)^2 \sqrt{-1 + cx} \sqrt{1 + cx}} + \frac{(fx)^{1+m} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2}{f(2+m)}$$

$$- \frac{2b^2 c^2 (fx)^{3+m} \sqrt{1 - c^2 x^2} \sqrt{d - c^2 dx^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+m}{2}, \frac{5+m}{2}, c^2 x^2\right)}{f^3(2+m)^2(3+m)(1 - cx)(1 + cx)}$$

$$+ \frac{d \operatorname{Int}\left(\frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}}, x\right)}{2+m}$$

output  $(f*x)^{(1+m)}*(a+b*\operatorname{arccosh}(c*x))^2*(-c^2*d*x^2+d)^{(1/2)}/f/(2+m)-2*b*c*(f*x)^{(2+m)}*(a+b*\operatorname{arccosh}(c*x))*(-c^2*d*x^2+d)^{(1/2)}/f^2/(2+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2*b^2*c^2*(f*x)^{(3+m)}*\operatorname{hypergeom}([1/2, 3/2+1/2*m], [5/2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^{(1/2)}*(-c^2*d*x^2+d)^{(1/2)}/f^3/(2+m)^2/(3+m)/(-c*x+1)/(c*x+1)+d*\operatorname{Unintegrable}((f*x)^m*(a+b*\operatorname{arccosh}(c*x))^2/(-c^2*d*x^2+d)^{(1/2)}, x)/(2+m)$

**3.235.2 Mathematica [N/A]**

Not integrable

Time = 0.76 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$$

input `Integrate[(f*x)^m*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`output `Integrate[(f*x)^m*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2, x]`**3.235.3 Rubi [N/A]**

Not integrable

Time = 1.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6345, 6298, 136, 279, 278, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{d - c^2 dx^2} (fx)^m (a + \operatorname{barccosh}(cx))^2 dx \\ & \quad \downarrow \text{6345} \\ & -\frac{2bc\sqrt{d - c^2 dx^2} \int (fx)^{m+1} (a + \operatorname{barccosh}(cx)) dx}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{d \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2 dx}{\sqrt{d - c^2 dx^2}}}{m+2} + \\ & \quad \frac{\sqrt{d - c^2 dx^2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+2)} \\ & \quad \downarrow \text{6298} \\ & -\frac{2bc\sqrt{d - c^2 dx^2} \left( \frac{(fx)^{m+2} (a + \operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc \int \frac{(fx)^{m+2}}{\sqrt{cx-1}\sqrt{cx+1}} dx}{f(m+2)} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{d \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2 dx}{\sqrt{d - c^2 dx^2}}}{m+2} + \\ & \quad \frac{\sqrt{d - c^2 dx^2} (fx)^{m+1} (a + \operatorname{barccosh}(cx))^2}{f(m+2)} \\ & \quad \downarrow \text{136} \end{aligned}$$

$$\begin{aligned}
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{c^2x^2-1} \int \frac{(fx)^{m+2}}{\sqrt{c^2x^2-1}} dx}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{m+2} + \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+2)} \\
& \quad \downarrow 279 \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{1-c^2x^2} \int \frac{(fx)^{m+2}}{\sqrt{1-c^2x^2}} dx}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{m+2} + \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+2)} \\
& \quad \downarrow 278 \\
& \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{m+2} - \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+3}{2}, \frac{m+5}{2}, c^2x^2\right)}{f^2(m+2)(m+3)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+2)} \\
& \quad \downarrow 6375 \\
& \frac{d \int \frac{(fx)^m (a+\operatorname{barccosh}(cx))^2}{\sqrt{d-c^2dx^2}} dx}{m+2} - \\
& \frac{2bc\sqrt{d-c^2dx^2} \left( \frac{(fx)^{m+2}(a+\operatorname{barccosh}(cx))}{f(m+2)} - \frac{bc\sqrt{1-c^2x^2} (fx)^{m+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+3}{2}, \frac{m+5}{2}, c^2x^2\right)}{f^2(m+2)(m+3)\sqrt{cx-1}\sqrt{cx+1}} \right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \\
& \frac{\sqrt{d-c^2dx^2} (fx)^{m+1} (a+\operatorname{barccosh}(cx))^2}{f(m+2)}
\end{aligned}$$

input `Int[(f*x)^m*sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

## 3.235.3.1 Defintions of rubi rules used

rule 136 `Int[((f_.)*(x_))^(p_)*((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[(a*c + b*d*x^2)^m*(f*x)^p, x], x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 279 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6345 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(f*(m + 2*p + 1))), x] + (Simp[2*d*(p/(m + 2*p + 1)) Int[(f*x)^m*(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(f*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0] && !LtQ[m, -1]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.235.4 Maple [N/A] (verified)**

Not integrable

Time = 3.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int (fx)^m \sqrt{-c^2 dx^2 + d} (a + b \operatorname{arccosh}(cx))^2 dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^2,x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^2,x)`

**3.235.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.39

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^2 (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)*(f*x)^m, x)`

**3.235.6 Sympy [N/A]**

Not integrable

Time = 72.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^2 dx = \int (fx)^m \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2 dx$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(1/2)*(a+b*acosh(c*x))**2,x)`

output `Integral((f*x)**m*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2, x)`

**3.235.7 Maxima [N/A]**

Not integrable

Time = 0.35 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^2 (fx)^m dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^2*(f*x)^m, x)`

**3.235.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.235.9 Mupad [N/A]**

Not integrable

Time = 3.05 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 \sqrt{d - c^2 dx^2} (fx)^m dx$$

input `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^2*(d - c^2*d*x^2)^(1/2)*(f*x)^m, x)`

---

3.235.  $\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^2 dx$



$$3.236 \quad \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

3.236.1 Optimal result	2112
3.236.2 Mathematica [N/A]	2112
3.236.3 Rubi [N/A]	2113
3.236.4 Maple [N/A] (verified)	2113
3.236.5 Fricas [N/A]	2114
3.236.6 Sympy [N/A]	2114
3.236.7 Maxima [N/A]	2114
3.236.8 Giac [N/A]	2115
3.236.9 Mupad [N/A]	2115

### 3.236.1 Optimal result

Integrand size = 31, antiderivative size = 31

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \operatorname{Int}\left(\frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x)`

### 3.236.2 Mathematica [N/A]

Not integrable

Time = 3.78 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

---


$$3.236. \quad \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

**3.236.3 Rubi [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^2)/Sqrt[d - c^2*d*x^2],x]`

output `$Aborted`

**3.236.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(x_.)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.236.4 Maple [N/A] (verified)**

Not integrable

Time = 1.65 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{\sqrt{-c^2 d x^2 + d}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x)`

**3.236.5 Fracas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.87

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)*(f*x)^m/(c^2*d*x^2 - d), x)`

**3.236.6 Sympy [N/A]**

Not integrable

Time = 22.64 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))^2}{\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))**2/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.236.7 Maxima [N/A]**

Not integrable

Time = 0.36 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^2*(f*x)^m/sqrt(-c^2*d*x^2 + d), x)`

---

3.236.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx$

**3.236.8 Giac [N/A]**

Not integrable

Time = 0.36 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*(f*x)^m/sqrt(-c^2*d*x^2 + d), x)`

**3.236.9 Mupad [N/A]**

Not integrable

Time = 3.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (fx)^m}{\sqrt{d - c^2 dx^2}} dx$$

input `int(((a + b*acosh(c*x))^2*(f*x)^m)/(d - c^2*d*x^2)^(1/2),x)`

output `int(((a + b*acosh(c*x))^2*(f*x)^m)/(d - c^2*d*x^2)^(1/2), x)`

**3.237**  $\int \frac{(fx)^m(a+b\text{arccosh}(cx))^2}{(d-c^2dx^2)^{3/2}} dx$

3.237.1 Optimal result . . . . . 2116  
 3.237.2 Mathematica [N/A] . . . . . 2116  
 3.237.3 Rubi [N/A] . . . . . 2117  
 3.237.4 Maple [N/A] (verified) . . . . . 2117  
 3.237.5 Fricas [N/A] . . . . . 2118  
 3.237.6 Sympy [N/A] . . . . . 2118  
 3.237.7 Maxima [N/A] . . . . . 2118  
 3.237.8 Giac [N/A] . . . . . 2119  
 3.237.9 Mupad [N/A] . . . . . 2119

**3.237.1 Optimal result**

Integrand size = 31, antiderivative size = 31

$$\int \frac{(fx)^m(a + \text{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \text{Int}\left(\frac{(fx)^m(a + \text{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x)`

**3.237.2 Mathematica [N/A]**

Not integrable

Time = 4.46 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{(fx)^m(a + \text{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(fx)^m(a + \text{arccosh}(cx))^2}{(d - c^2dx^2)^{3/2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2),x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2), x]`

**3.237.3 Rubi [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(3/2),x]`

output `$Aborted`

**3.237.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^m_.*((d_) + (e_.)*(x_)^2)^p_.], x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.237.4 Maple [N/A] (verified)**

Not integrable

Time = 3.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x)`

---

3.237.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx$

**3.237.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.26

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)*(f*x)^m/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.237.6 Sympy [N/A]**

Not integrable

Time = 26.56 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))^2}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))**2/(-d*(c*x - 1)*(c*x + 1))**3/2, x)`

**3.237.7 Maxima [N/A]**

Not integrable

Time = 0.43 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^2*(f*x)^m/(-c^2*d*x^2 + d)^(3/2), x)`

### 3.237.8 Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2 (fx)^m}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*(f*x)^m/(-c^2*d*x^2 + d)^(3/2), x)`

### 3.237.9 Mupad [N/A]

Not integrable

Time = 3.38 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (fx)^m}{(d - c^2 dx^2)^{3/2}} dx$$

input `int(((a + b*acosh(c*x))^2*(f*x)^m)/(d - c^2*d*x^2)^(3/2),x)`

output `int(((a + b*acosh(c*x))^2*(f*x)^m)/(d - c^2*d*x^2)^(3/2), x)`



**3.238** 
$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

3.238.1 Optimal result . . . . .	2120
3.238.2 Mathematica [N/A] . . . . .	2120
3.238.3 Rubi [N/A] . . . . .	2121
3.238.4 Maple [N/A] (verified) . . . . .	2121
3.238.5 Fricas [N/A] . . . . .	2122
3.238.6 Sympy [F(-1)] . . . . .	2122
3.238.7 Maxima [N/A] . . . . .	2122
3.238.8 Giac [N/A] . . . . .	2123
3.238.9 Mupad [N/A] . . . . .	2123

**3.238.1 Optimal result**

Integrand size = 31, antiderivative size = 31

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \operatorname{Int} \left( \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}}, x \right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x)`

**3.238.2 Mathematica [N/A]**

Not integrable

Time = 4.76 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2),x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2), x]`

**3.238.3 Rubi [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^2)/(d - c^2*d*x^2)^(5/2),x]`

output `$Aborted`

**3.238.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.238.4 Maple [N/A] (verified)**

Not integrable

Time = 3.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x)`

---

3.238.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$

**3.238.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 84, normalized size of antiderivative = 2.71

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{(-c^2 dx^2 + d)^{5/2}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="
fricas")
```

```
output integral(-sqrt(-c^2*d*x^2 + d)*(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) +
a^2)*(f*x)^m/(c^6*d^3*x^6 - 3*c^4*d^3*x^4 + 3*c^2*d^3*x^2 - d^3), x)
```

**3.238.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \text{Timed out}$$

```
input integrate((f*x)**m*(a+b*acosh(c*x))**2/(-c**2*d*x**2+d)**(5/2),x)
```

```
output Timed out
```

**3.238.7 Maxima [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{(-c^2 dx^2 + d)^{5/2}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="
maxima")
```

```
output integrate((b*arccosh(c*x) + a)^2*(f*x)^m/(-c^2*d*x^2 + d)^(5/2), x)
```

---

3.238.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx$

**3.238.8 Giac [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2 (fx)^m}{(-c^2 dx^2 + d)^{5/2}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^2/(-c^2*d*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2*(f*x)^m/(-c^2*d*x^2 + d)^(5/2), x)`

**3.238.9 Mupad [N/A]**

Not integrable

Time = 3.44 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^2}{(d - c^2 dx^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2 (fx)^m}{(d - c^2 dx^2)^{5/2}} dx$$

input `int(((a + b*acosh(c*x))^2*(f*x)^m)/(d - c^2*d*x^2)^(5/2),x)`

output `int(((a + b*acosh(c*x))^2*(f*x)^m)/(d - c^2*d*x^2)^(5/2), x)`

$$3.239 \quad \int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx$$

3.239.1 Optimal result	2124
3.239.2 Mathematica [N/A]	2124
3.239.3 Rubi [N/A]	2125
3.239.4 Maple [N/A] (verified)	2125
3.239.5 Fricas [N/A]	2126
3.239.6 Sympy [N/A]	2126
3.239.7 Maxima [N/A]	2126
3.239.8 Giac [N/A]	2127
3.239.9 Mupad [N/A]	2127

### 3.239.1 Optimal result

Integrand size = 26, antiderivative size = 26

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx = \operatorname{Int}\left(\frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}}, x\right)$$

output `Unintegrable((f*x)^m*arccosh(c*x)^2/(-c^2*x^2+1)^(1/2), x)`

### 3.239.2 Mathematica [N/A]

Not integrable

Time = 0.74 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx = \int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx$$

input `Integrate[((f*x)^m*ArcCosh[c*x]^2)/Sqrt[1 - c^2*x^2], x]`

output `Integrate[((f*x)^m*ArcCosh[c*x]^2)/Sqrt[1 - c^2*x^2], x]`

**3.239.3 Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(cx)^2 (fx)^m}{\sqrt{1-c^2x^2}} dx$$

↓ 6375

$$\int \frac{\operatorname{arccosh}(cx)^2 (fx)^m}{\sqrt{1-c^2x^2}} dx$$

input `Int[((f*x)^m*ArcCosh[c*x]^2)/Sqrt[1 - c^2*x^2], x]`

output `$Aborted`

**3.239.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.239.4 Maple [N/A] (verified)**

Not integrable

Time = 1.50 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{-c^2x^2+1}} dx$$

input `int((f*x)^m*arccosh(c*x)^2/(-c^2*x^2+1)^(1/2), x)`

output `int((f*x)^m*arccosh(c*x)^2/(-c^2*x^2+1)^(1/2), x)`

**3.239.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.46

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx = \int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{-c^2x^2+1}} dx$$

input `integrate((f*x)^m*arccosh(c*x)^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`output `integral(-sqrt(-c^2*x^2 + 1)*(f*x)^m*arccosh(c*x)^2/(c^2*x^2 - 1), x)`**3.239.6 Sympy [N/A]**

Not integrable

Time = 11.45 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.04

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx = \int \frac{(fx)^m \operatorname{acosh}^2(cx)}{\sqrt{-(cx-1)(cx+1)}} dx$$

input `integrate((f*x)**m*acosh(c*x)**2/(-c**2*x**2+1)**(1/2),x)`output `Integral((f*x)**m*acosh(c*x)**2/sqrt(-(c*x - 1)*(c*x + 1)), x)`**3.239.7 Maxima [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx = \int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{-c^2x^2+1}} dx$$

input `integrate((f*x)^m*arccosh(c*x)^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`output `integrate((f*x)^m*arccosh(c*x)^2/sqrt(-c^2*x^2 + 1), x)`

---

3.239.  $\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx$

**3.239.8 Giac [N/A]**

Not integrable

Time = 0.36 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx = \int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{-c^2x^2+1}} dx$$

input `integrate((f*x)^m*arccosh(c*x)^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`output `integrate((f*x)^m*arccosh(c*x)^2/sqrt(-c^2*x^2 + 1), x)`**3.239.9 Mupad [N/A]**

Not integrable

Time = 3.24 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m \operatorname{arccosh}(cx)^2}{\sqrt{1-c^2x^2}} dx = \int \frac{\operatorname{acosh}(cx)^2 (fx)^m}{\sqrt{1-c^2x^2}} dx$$

input `int((acosh(c*x)^2*(f*x)^m)/(1 - c^2*x^2)^(1/2),x)`output `int((acosh(c*x)^2*(f*x)^m)/(1 - c^2*x^2)^(1/2), x)`



## 3.240 $\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx$

3.240.1 Optimal result . . . . .	2128
3.240.2 Mathematica [A] (verified) . . . . .	2129
3.240.3 Rubi [A] (verified) . . . . .	2129
3.240.4 Maple [A] (verified) . . . . .	2139
3.240.5 Fricas [A] (verification not implemented) . . . . .	2139
3.240.6 Sympy [F] . . . . .	2140
3.240.7 Maxima [A] (verification not implemented) . . . . .	2140
3.240.8 Giac [F(-2)] . . . . .	2141
3.240.9 Mupad [F(-1)] . . . . .	2141

### 3.240.1 Optimal result

Integrand size = 20, antiderivative size = 505

$$\begin{aligned}
 \int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx = & -\frac{976c^3\sqrt{-1+ax}\sqrt{1+ax}}{315a} \\
 & + \frac{16}{315}ac^3x^2\sqrt{-1+ax}\sqrt{1+ax} + \frac{7104c^3(1-a^2x^2)}{42875a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{1184c^3(1-a^2x^2)^2}{42875a\sqrt{-1+ax}\sqrt{1+ax}} \\
 & + \frac{2664c^3(1-a^2x^2)^3}{214375a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{6c^3(1-a^2x^2)^4}{2401a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{4322c^3x\operatorname{arccosh}(ax)}{1225} \\
 & - \frac{1514a^2c^3x^3\operatorname{arccosh}(ax)}{3675} + \frac{702a^4c^3x^5\operatorname{arccosh}(ax)}{6125} - \frac{6}{343}a^6c^3x^7\operatorname{arccosh}(ax) \\
 & - \frac{48c^3\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{35a} + \frac{8c^3(-1+ax)^{3/2}(1+ax)^{3/2}\operatorname{arccosh}(ax)^2}{35a} \\
 & - \frac{18c^3(-1+ax)^{5/2}(1+ax)^{5/2}\operatorname{arccosh}(ax)^2}{175a} + \frac{3c^3(-1+ax)^{7/2}(1+ax)^{7/2}\operatorname{arccosh}(ax)^2}{49a} \\
 & + \frac{16}{35}c^3x\operatorname{arccosh}(ax)^3 + \frac{8}{35}c^3x(1-a^2x^2)\operatorname{arccosh}(ax)^3 + \frac{6}{35}c^3x(1-a^2x^2)^2\operatorname{arccosh}(ax)^3 + \frac{1}{7}c^3x(1-a^2x^2)^3\operatorname{arccosh}(ax)^3
 \end{aligned}$$

output  $4322/1225*c^3*x*arccosh(a*x)-1514/3675*a^2*c^3*x^3*arccosh(a*x)+702/6125*a^4*c^3*x^5*arccosh(a*x)-6/343*a^6*c^3*x^7*arccosh(a*x)+8/35*c^3*(a*x-1)^(3/2)*(a*x+1)^(3/2)*arccosh(a*x)^2/a-18/175*c^3*(a*x-1)^(5/2)*(a*x+1)^(5/2)*arccosh(a*x)^2/a+3/49*c^3*(a*x-1)^(7/2)*(a*x+1)^(7/2)*arccosh(a*x)^2/a+16/35*c^3*x*arccosh(a*x)^3+8/35*c^3*x*(-a^2*x^2+1)*arccosh(a*x)^3+6/35*c^3*x*(-a^2*x^2+1)^2*arccosh(a*x)^3+1/7*c^3*x*(-a^2*x^2+1)^3*arccosh(a*x)^3+7104/42875*c^3*(-a^2*x^2+1)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1184/42875*c^3*(-a^2*x^2+1)^2/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+2664/214375*c^3*(-a^2*x^2+1)^3/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+6/2401*c^3*(-a^2*x^2+1)^4/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-976/315*c^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a+16/315*a*c^3*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-48/35*c^3*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a$

### 3.240.2 Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.35

$$\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx$$

$$= \frac{c^3(2\sqrt{-1 + ax}\sqrt{1 + ax}(-22329151 + 747937a^2x^2 - 134541a^4x^4 + 16875a^6x^6) - 210ax(-226905 + 26495a^2x^2 - 7371a^4x^4 + 1125a^6x^6)*\operatorname{ArcCosh}[a*x] + 11025*\sqrt{-1 + ax}*\sqrt{1 + ax}*(-2161 + 757a^2x^2 - 351a^4x^4 + 75a^6x^6)*\operatorname{ArcCosh}[a*x]^2 - 385875*a*x*(-35 + 35a^2x^2 - 21a^4x^4 + 5a^6x^6)*\operatorname{ArcCosh}[a*x]^3))/(13505625*a)}$$

input `Integrate[(c - a^2*c*x^2)^3*ArcCosh[a*x]^3,x]`

output  $(c^3*(2*\sqrt{-1 + a*x})*\sqrt{1 + a*x}*(-22329151 + 747937*a^2*x^2 - 134541*a^4*x^4 + 16875*a^6*x^6) - 210*a*x*(-226905 + 26495*a^2*x^2 - 7371*a^4*x^4 + 1125*a^6*x^6)*\operatorname{ArcCosh}[a*x] + 11025*\sqrt{-1 + a*x}*\sqrt{1 + a*x}*(-2161 + 757*a^2*x^2 - 351*a^4*x^4 + 75*a^6*x^6)*\operatorname{ArcCosh}[a*x]^2 - 385875*a*x*(-35 + 35*a^2*x^2 - 21*a^4*x^4 + 5*a^6*x^6)*\operatorname{ArcCosh}[a*x]^3))/(13505625*a)$

### 3.240.3 Rubi [A] (verified)

Time = 4.80 (sec) , antiderivative size = 665, normalized size of antiderivative = 1.32, number of steps used = 23, number of rules used = 22,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.100$ , Rules used = {6312, 27, 6312, 6312, 6294, 6330, 25, 6294, 83, 6304, 6309, 27, 960, 83, 1905, 1576, 1140, 2009, 2113, 2331, 2389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.240.  $\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx$

$$\begin{aligned}
& \int \operatorname{arccosh}(ax)^3 (c - a^2cx^2)^3 dx \\
& \quad \downarrow \text{6312} \\
& \frac{6}{7}c \int c^2(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^3 dx + \frac{3}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)^2 dx + \\
& \quad \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^3 \\
& \quad \downarrow \text{27} \\
& \frac{6}{7}c^3 \int (1 - a^2x^2)^2 \operatorname{arccosh}(ax)^3 dx + \frac{3}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)^2 dx + \\
& \quad \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^3 \\
& \quad \downarrow \text{6312} \\
& \frac{6}{7}c^3 \left( \frac{4}{5} \int (1 - a^2x^2) \operatorname{arccosh}(ax)^3 dx - \frac{3}{5}a \int x(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{5}x(1 - a^2x^2)^2 \operatorname{arccosh}(ax) \right. \\
& \quad \left. + \frac{3}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right) \\
& \quad \downarrow \text{6312} \\
& \frac{6}{7}c^3 \left( \frac{4}{5} \left( a \int x\sqrt{ax - 1}\sqrt{ax + 1} \operatorname{arccosh}(ax)^2 dx + \frac{2}{3} \int \operatorname{arccosh}(ax)^3 dx + \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^3 \right) - \frac{3}{5}a \int x \right. \\
& \quad \left. + \frac{3}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right) \\
& \quad \downarrow \text{6294} \\
& \frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{ax - 1}\sqrt{ax + 1}} dx \right) + a \int x\sqrt{ax - 1}\sqrt{ax + 1} \operatorname{arccosh}(ax)^2 dx + \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^3 \right) \right. \\
& \quad \left. + \frac{3}{7}ac^3 \int x(ax - 1)^{5/2}(ax + 1)^{5/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right) \\
& \quad \downarrow \text{6330} \\
& \frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax - 1}\sqrt{ax + 1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \int \operatorname{arccosh}(ax) dx}{a} \right) \right) + a \left( \frac{(ax - 1)^{3/2}(ax + 1)^{3/2}}{3a} \right) \right. \right. \\
& \quad \left. \left. + \frac{3}{7}ac^3 \left( \frac{(ax - 1)^{7/2}(ax + 1)^{7/2} \operatorname{arccosh}(ax)^2}{7a^2} - \frac{2 \int -(1 - ax)^3(ax + 1)^3 \operatorname{arccosh}(ax) dx}{7a} \right) + \frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right) \right) \\
& \quad \downarrow \text{25}
\end{aligned}$$

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \int \operatorname{arccosh}(ax) dx}{a} \right) \right) + a \left( \frac{2 \int (1-ax)(ax+1) \operatorname{arccosh}(ax) dx}{3a} + \frac{3}{7}ac^3 \left( \frac{2 \int (1-ax)^3(ax+1)^3 \operatorname{arccosh}(ax) dx}{7a} + \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)^2}{7a^2} \right) + \frac{1}{7}c^3 x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right) \right)$$

↓ 6294

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - a \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a} \right) \right) + \frac{3}{7}ac^3 \left( \frac{2 \int (1-ax)^3(ax+1)^3 \operatorname{arccosh}(ax) dx}{7a} + \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)^2}{7a^2} \right) + \frac{1}{7}c^3 x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 83

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( a \left( \frac{2 \int (1-ax)(ax+1) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + \frac{3}{7}ac^3 \left( \frac{2 \int (1-ax)^3(ax+1)^3 \operatorname{arccosh}(ax) dx}{7a} + \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)^2}{7a^2} \right) + \frac{1}{7}c^3 x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 6304

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( a \left( \frac{2 \int (1-a^2x^2) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + \frac{3}{7}ac^3 \left( \frac{2 \int (1-a^2x^2)^3 \operatorname{arccosh}(ax) dx}{7a} + \frac{(ax-1)^{7/2}(ax+1)^{7/2} \operatorname{arccosh}(ax)^2}{7a^2} \right) + \frac{1}{7}c^3 x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 6309

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( a \left( \frac{2 \left( -a \int \frac{x(3-a^2x^2)}{3\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} \right) + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) \right. \\ \left. \frac{3}{7}ac^3 \left( \frac{2 \left( -a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{35\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{7a} \right) \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 27

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( a \left( \frac{2 \left( -\frac{1}{3}a \int \frac{x(3-a^2x^2)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} \right) + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) \right. \\ \left. \frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{1}{35}a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{7a} \right) \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 960

$$\frac{6}{7}c^3 \left( \frac{4}{5} \left( a \left( \frac{2 \left( -\frac{1}{3}a \left( \frac{7}{3} \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} \right) + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{3a^2} \right) \right. \\ \left. \frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{1}{35}a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{7a} \right) \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 83

$$\frac{6}{7}c^3 \left( -\frac{3}{5}a \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{1}{15}a \int \frac{x(3a^4x^4-10a^2x^2+15)}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{5a} \right) \right. \\ \left. \frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{1}{35}a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{7a} \right) \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 1905

$$\frac{6}{7}c^3 \left( -\frac{3}{5}a \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \frac{x(3a^4x^4-10a^2x^2+15)}{\sqrt{a^2x^2-1}} dx}{15\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) \right. \\ \left. \frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{1}{35}a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + xa \right)}{7a} \right) \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 1576

$$\frac{6}{7}c^3 \left( -\frac{3}{5}a \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \frac{3a^4x^4-10a^2x^2+15}{\sqrt{a^2x^2-1}} dx^2}{30\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) \right. \\ \left. \frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{1}{35}a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + xa \right)}{7a} \right) \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 1140

$$\frac{6}{7}c^3 \left( -\frac{3}{5}a \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \left( 3(a^2x^2-1)^{3/2} - 4\sqrt{a^2x^2-1} + \frac{8}{\sqrt{a^2x^2-1}} \right) dx^2}{30\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) \right. \\ \left. \frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{1}{35}a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + xa \right)}{7a} \right) \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 \right)$$

↓ 2009

$$\frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{1}{35}a \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + xa \right)}{7a} \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 + \right. \\ \left. \frac{6}{7}c^3 \left( \frac{1}{5}x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3 + \frac{4}{5} \left( \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{7a} \right) \right) \right)$$

↓ 2113

$$\frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{a\sqrt{a^2x^2-1}}{35\sqrt{ax-1}\sqrt{ax+1}} \int \frac{x(-5a^6x^6+21a^4x^4-35a^2x^2+35)}{\sqrt{a^2x^2-1}} dx - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + xa \right)}{7a} \right. \\ \left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 + \right. \\ \left. \frac{6}{7}c^3 \left( \frac{1}{5}x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3 + \frac{4}{5} \left( \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{7a} \right) \right) \right)$$

↓ 2331

$$\frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \frac{-5a^6x^6+21a^4x^4-35a^2x^2+35}{\sqrt{a^2x^2-1}} dx^2}{70\sqrt{ax-1}\sqrt{ax+1}} - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + x \right)}{7a} \right.$$

$$\left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 + \right.$$

$$\left. \frac{6}{7}c^3 \left( \frac{1}{5}x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3 + \frac{4}{5} \left( \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{7a} \right) \right) \right)$$

↓ 2389

$$\frac{3}{7}ac^3 \left( \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \left( -5(a^2x^2-1)^{5/2} + 6(a^2x^2-1)^{3/2} - 8\sqrt{a^2x^2-1} + \frac{16}{\sqrt{a^2x^2-1}} \right) dx^2}{70\sqrt{ax-1}\sqrt{ax+1}} - \frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) + x \right)}{7a} \right.$$

$$\left. \frac{1}{7}c^3x(1-a^2x^2)^3 \operatorname{arccosh}(ax)^3 + \right.$$

$$\left. \frac{6}{7}c^3 \left( \frac{1}{5}x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3 + \frac{4}{5} \left( \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{7a} \right) \right) \right)$$

↓ 2009



$$\frac{1}{7}c^3x(1 - a^2x^2)^3 \operatorname{arccosh}(ax)^3 + \left( \frac{6}{7}c^3 \left[ \frac{1}{5}x(1 - a^2x^2)^2 \operatorname{arccosh}(ax)^3 + \frac{4}{5} \left( \frac{1}{3}x(1 - a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} \right) \right)}{2} \right) \right] \right) \right. \\ \left. \frac{3}{7}ac^3 \left[ \frac{(ax - 1)^{7/2}(ax + 1)^{7/2} \operatorname{arccosh}(ax)^2}{7a^2} + \frac{2 \left( -\frac{1}{7}a^6x^7 \operatorname{arccosh}(ax) + \frac{3}{5}a^4x^5 \operatorname{arccosh}(ax) - a^2x^3 \operatorname{arccosh}(ax) - \dots \right)}{\dots} \right] \right)$$

```
input Int[(c - a^2*c*x^2)^3*ArcCosh[a*x]^3,x]
```

```
output (c^3*x*(1 - a^2*x^2)^3*ArcCosh[a*x]^3)/7 + (3*a*c^3*((( -1 + a*x)^(7/2)*(1 + a*x)^(7/2)*ArcCosh[a*x]^2)/(7*a^2) + (2*(-1/70*(a*Sqrt[-1 + a^2*x^2]*((3*2*Sqrt[-1 + a^2*x^2])/a^2 - (16*(-1 + a^2*x^2)^(3/2))/(3*a^2) + (12*(-1 + a^2*x^2)^(5/2))/(5*a^2) - (10*(-1 + a^2*x^2)^(7/2))/(7*a^2)))/(Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + x*ArcCosh[a*x] - a^2*x^3*ArcCosh[a*x] + (3*a^4*x^5*ArcCosh[a*x])/5 - (a^6*x^7*ArcCosh[a*x])/7))/(7*a))/7 + (6*c^3*((x*(1 - a^2*x^2)^2*ArcCosh[a*x]^3)/5 - (3*a*((( -1 + a*x)^(5/2)*(1 + a*x)^(5/2)*ArcCosh[a*x]^2)/(5*a^2) - (2*(-1/30*(a*Sqrt[-1 + a^2*x^2]*((16*Sqrt[-1 + a^2*x^2])/a^2 - (8*(-1 + a^2*x^2)^(3/2))/(3*a^2) + (6*(-1 + a^2*x^2)^(5/2))/(5*a^2)))/(Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + x*ArcCosh[a*x] - (2*a^2*x^3*ArcCosh[a*x])/3 + (a^4*x^5*ArcCosh[a*x])/5))/(5*a))/5 + (4*((x*(1 - a^2*x^2)*ArcCosh[a*x]^3)/3 + a*((( -1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x]^2)/(3*a^2) + (2*(-1/3*(a*((7*Sqrt[-1 + a*x]*Sqrt[1 + a*x]))/(3*a^2) - (x^2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/3)) + x*ArcCosh[a*x] - (a^2*x^3*ArcCosh[a*x])/3))/(3*a)) + (2*(x*ArcCosh[a*x]^3 - 3*a*((Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^2)/a^2 - (2*(-((Sqrt[-1 + a*x]*Sqrt[1 + a*x])/a) + x*ArcCosh[a*x]))/a))/3))/5))/7
```

## 3.240.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 960 `Int[((e_.)*(x_)^(m_.))*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 1140 `Int[((d_.) + (e_.)*(x_)^(m_.))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 0]`
- rule 1576 `Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`
- rule 1905 `Int[((f_.)*(x_)^(m_.))*((d1_) + (e1_.)*(x_)^(non2_.))^(q_.)*((d2_) + (e2_.)*(x_)^(non2_.))^(q_.)*((a_.) + (b_.)*(x_)^(n_) + (c_.)*(x_)^(n2_))^(p_.), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2113 `Int[(Px_)*((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m])/(a*c + b*d*x^2)^FracPart[m] Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

rule 2331 `Int[(Pq_)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`

rule 2389 `Int[(Pq_)*((a_) + (b_.)*(x_)^(n_.))^(p_.), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, n}, x] && PolyQ[Pq, x] && (IGtQ[p, 0] || EqQ[n, 1])`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n/(2*p + 1))), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

```
rule 6330 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p
_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 +
e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2
*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^
p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n -
1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && E
qQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]
```

### 3.240.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 294, normalized size of antiderivative = 0.58

method	result
derivativedivides	$-\frac{c^3 \left( 1929375 \operatorname{arccosh}(ax)^3 a^7 x^7 - 826875 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^6 x^6 - 8103375 \operatorname{arccosh}(ax)^3 a^5 x^5 + 3869775 \operatorname{arccosh}(ax)^2 a^4 x^4 + 236250 \operatorname{arccosh}(ax) a^7 x^7 - 33750 a^6 x^6 \right)}{c^3 \left( 1929375 \operatorname{arccosh}(ax)^3 a^7 x^7 - 826875 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^6 x^6 - 8103375 \operatorname{arccosh}(ax)^3 a^5 x^5 + 3869775 \operatorname{arccosh}(ax)^2 a^4 x^4 + 236250 \operatorname{arccosh}(ax) a^7 x^7 - 33750 a^6 x^6 \right)}$
default	$-\frac{c^3 \left( 1929375 \operatorname{arccosh}(ax)^3 a^7 x^7 - 826875 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^6 x^6 - 8103375 \operatorname{arccosh}(ax)^3 a^5 x^5 + 3869775 \operatorname{arccosh}(ax)^2 a^4 x^4 + 236250 \operatorname{arccosh}(ax) a^7 x^7 - 33750 a^6 x^6 \right)}{c^3 \left( 1929375 \operatorname{arccosh}(ax)^3 a^7 x^7 - 826875 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^6 x^6 - 8103375 \operatorname{arccosh}(ax)^3 a^5 x^5 + 3869775 \operatorname{arccosh}(ax)^2 a^4 x^4 + 236250 \operatorname{arccosh}(ax) a^7 x^7 - 33750 a^6 x^6 \right)}$

```
input int((-a^2*c*x^2+c)^3*arccosh(a*x)^3,x,method=_RETURNVERBOSE)
```

```
output -1/13505625/a*c^3*(1929375*arccosh(a*x)^3*a^7*x^7-826875*arccosh(a*x)^2*(a
*x-1)^(1/2)*(a*x+1)^(1/2)*a^6*x^6-8103375*arccosh(a*x)^3*a^5*x^5+3869775*a
rccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4+236250*arccosh(a*x)*a^7*
x^7-33750*x^6*a^6*(a*x-1)^(1/2)*(a*x+1)^(1/2)+13505625*a^3*x^3*arccosh(a*x
)^3-8345925*a^2*x^2*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-1547910*a^5
*x^5*arccosh(a*x)+269082*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4-13505625*a*x*
arccosh(a*x)^3+23825025*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+5563950
*a^3*x^3*arccosh(a*x)-1495874*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-47650050
*a*x*arccosh(a*x)+44658302*(a*x-1)^(1/2)*(a*x+1)^(1/2))
```

### 3.240.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 248, normalized size of antiderivative = 0.49

$$\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^3 dx = \frac{385875 (5 a^7 c^3 x^7 - 21 a^5 c^3 x^5 + 35 a^3 c^3 x^3 - 35 a c^3 x) \log(ax + \sqrt{a^2 x^2 - 1})^3 - 11025 (75 a^6 c^3 x^6 - 351 a^5 c^3 x^5 + 525 a^4 c^3 x^4 - 210 a^3 c^3 x^3 + 35 a^2 c^3 x^2 - 35 a c^3 x) \operatorname{arccosh}(ax)^3 + 11025 (75 a^6 c^3 x^6 - 351 a^5 c^3 x^5 + 525 a^4 c^3 x^4 - 210 a^3 c^3 x^3 + 35 a^2 c^3 x^2 - 35 a c^3 x) \operatorname{arccosh}(ax)^2 + 11025 (75 a^6 c^3 x^6 - 351 a^5 c^3 x^5 + 525 a^4 c^3 x^4 - 210 a^3 c^3 x^3 + 35 a^2 c^3 x^2 - 35 a c^3 x) \operatorname{arccosh}(ax) + 11025 (75 a^6 c^3 x^6 - 351 a^5 c^3 x^5 + 525 a^4 c^3 x^4 - 210 a^3 c^3 x^3 + 35 a^2 c^3 x^2 - 35 a c^3 x) \operatorname{arccosh}(ax)^3}{c^3 (5 a^7 c^3 x^7 - 21 a^5 c^3 x^5 + 35 a^3 c^3 x^3 - 35 a c^3 x)}$$

---

3.240.  $\int (c - a^2 cx^2)^3 \operatorname{arccosh}(ax)^3 dx$

input `integrate((-a^2*c*x^2+c)^3*arccosh(a*x)^3,x, algorithm="fricas")`

output `-1/13505625*(385875*(5*a^7*c^3*x^7 - 21*a^5*c^3*x^5 + 35*a^3*c^3*x^3 - 35*a*c^3*x)*log(a*x + sqrt(a^2*x^2 - 1))^3 - 11025*(75*a^6*c^3*x^6 - 351*a^4*c^3*x^4 + 757*a^2*c^3*x^2 - 2161*c^3)*sqrt(a^2*x^2 - 1)*log(a*x + sqrt(a^2*x^2 - 1))^2 + 210*(1125*a^7*c^3*x^7 - 7371*a^5*c^3*x^5 + 26495*a^3*c^3*x^3 - 226905*a*c^3*x)*log(a*x + sqrt(a^2*x^2 - 1)) - 2*(16875*a^6*c^3*x^6 - 134541*a^4*c^3*x^4 + 747937*a^2*c^3*x^2 - 22329151*c^3)*sqrt(a^2*x^2 - 1)) /a`

### 3.240.6 Sympy [F]

$$\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx = -c^3 \left( \int 3a^2x^2 \operatorname{acosh}^3(ax) dx + \int (-3a^4x^4 \operatorname{acosh}^3(ax)) dx + \int a^6x^6 \operatorname{acosh}^3(ax) dx + \int (-\operatorname{acosh}^3(ax)) dx \right)$$

input `integrate((-a**2*c*x**2+c)**3*acosh(a*x)**3,x)`

output `-c**3*(Integral(3*a**2*x**2*acosh(a*x)**3, x) + Integral(-3*a**4*x**4*acosh(a*x)**3, x) + Integral(a**6*x**6*acosh(a*x)**3, x) + Integral(-acosh(a*x)**3, x))`

### 3.240.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.55

$$\begin{aligned} & \int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx \\ &= \frac{1}{1225} \left( 75 \sqrt{a^2x^2 - 1} a^4 c^3 x^6 - 351 \sqrt{a^2x^2 - 1} a^2 c^3 x^4 + 757 \sqrt{a^2x^2 - 1} c^3 x^2 - \frac{2161 \sqrt{a^2x^2 - 1} c^3}{a^2} \right) a \operatorname{arccosh}(ax) \\ & \quad - \frac{1}{35} (5 a^6 c^3 x^7 - 21 a^4 c^3 x^5 + 35 a^2 c^3 x^3 - 35 c^3 x) \operatorname{arccosh}(ax)^3 \\ & \quad + \frac{2}{13505625} \left( 16875 \sqrt{a^2x^2 - 1} a^4 c^3 x^6 - 134541 \sqrt{a^2x^2 - 1} a^2 c^3 x^4 + 747937 \sqrt{a^2x^2 - 1} c^3 x^2 - \frac{22329151 \sqrt{a^2x^2 - 1} c^3}{a} \right) \end{aligned}$$

input `integrate((-a^2*c*x^2+c)^3*arccosh(a*x)^3,x, algorithm="maxima")`

output `1/1225*(75*sqrt(a^2*x^2 - 1)*a^4*c^3*x^6 - 351*sqrt(a^2*x^2 - 1)*a^2*c^3*x^4 + 757*sqrt(a^2*x^2 - 1)*c^3*x^2 - 2161*sqrt(a^2*x^2 - 1)*c^3/a^2)*a*arc  
cosh(a*x)^2 - 1/35*(5*a^6*c^3*x^7 - 21*a^4*c^3*x^5 + 35*a^2*c^3*x^3 - 35*c  
^3*x)*arccosh(a*x)^3 + 2/13505625*(16875*sqrt(a^2*x^2 - 1)*a^4*c^3*x^6 - 1  
34541*sqrt(a^2*x^2 - 1)*a^2*c^3*x^4 + 747937*sqrt(a^2*x^2 - 1)*c^3*x^2 - 2  
2329151*sqrt(a^2*x^2 - 1)*c^3/a^2 - 105*(1125*a^6*c^3*x^7 - 7371*a^4*c^3*x  
^5 + 26495*a^2*c^3*x^3 - 226905*c^3*x)*arccosh(a*x)/a)*a`

### 3.240.8 Giac [F(-2)]

Exception generated.

$$\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^3*arccosh(a*x)^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.240.9 Mupad [F(-1)]

Timed out.

$$\int (c - a^2cx^2)^3 \operatorname{arccosh}(ax)^3 dx = \int \operatorname{acosh}(ax)^3 (c - a^2cx^2)^3 dx$$

input `int(acosh(a*x)^3*(c - a^2*c*x^2)^3,x)`

output `int(acosh(a*x)^3*(c - a^2*c*x^2)^3, x)`

### 3.241 $\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^3 dx$

3.241.1 Optimal result . . . . .	2142
3.241.2 Mathematica [A] (verified) . . . . .	2143
3.241.3 Rubi [A] (verified) . . . . .	2143
3.241.4 Maple [A] (verified) . . . . .	2150
3.241.5 Fricas [A] (verification not implemented) . . . . .	2150
3.241.6 Sympy [F] . . . . .	2151
3.241.7 Maxima [A] (verification not implemented) . . . . .	2151
3.241.8 Giac [F(-2)] . . . . .	2152
3.241.9 Mupad [F(-1)] . . . . .	2152

#### 3.241.1 Optimal result

Integrand size = 20, antiderivative size = 388

$$\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^3 dx$$

$$= -\frac{488c^2\sqrt{-1+ax}\sqrt{1+ax}}{135a} + \frac{8}{135}ac^2x^2\sqrt{-1+ax}\sqrt{1+ax} + \frac{16c^2(1-a^2x^2)}{125a\sqrt{-1+ax}\sqrt{1+ax}}$$

$$+ \frac{8c^2(1-a^2x^2)^2}{375a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{6c^2(1-a^2x^2)^3}{625a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{298}{75}c^2x\operatorname{arccosh}(ax)$$

$$- \frac{76}{225}a^2c^2x^3\operatorname{arccosh}(ax) + \frac{6}{125}a^4c^2x^5\operatorname{arccosh}(ax) - \frac{8c^2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{5a}$$

$$+ \frac{4c^2(-1+ax)^{3/2}(1+ax)^{3/2}\operatorname{arccosh}(ax)^2}{15a} - \frac{3c^2(-1+ax)^{5/2}(1+ax)^{5/2}\operatorname{arccosh}(ax)^2}{25a}$$

$$+ \frac{8}{15}c^2x\operatorname{arccosh}(ax)^3 + \frac{4}{15}c^2x(1-a^2x^2)\operatorname{arccosh}(ax)^3 + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^3$$

```
output 298/75*c^2*x*arccosh(a*x)-76/225*a^2*c^2*x^3*arccosh(a*x)+6/125*a^4*c^2*x^
5*arccosh(a*x)+4/15*c^2*(a*x-1)^(3/2)*(a*x+1)^(3/2)*arccosh(a*x)^2/a-3/25*
c^2*(a*x-1)^(5/2)*(a*x+1)^(5/2)*arccosh(a*x)^2/a+8/15*c^2*x*arccosh(a*x)^3
+4/15*c^2*x*(-a^2*x^2+1)*arccosh(a*x)^3+1/5*c^2*x*(-a^2*x^2+1)^2*arccosh(a
*x)^3+16/125*c^2*(-a^2*x^2+1)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+8/375*c^2*(-a^
2*x^2+1)^2/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+6/625*c^2*(-a^2*x^2+1)^3/a/(a*x-1
)^(1/2)/(a*x+1)^(1/2)-488/135*c^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a+8/135*a*c^
2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-8/5*c^2*arccosh(a*x)^2*(a*x-1)^(1/2)*(a
*x+1)^(1/2)/a
```

**3.241.2 Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.38

$$\int (c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^3 dx$$

$$= \frac{c^2(-2\sqrt{-1+ax}\sqrt{1+ax}(31841 - 842a^2x^2 + 81a^4x^4) + 30ax(2235 - 190a^2x^2 + 27a^4x^4) \operatorname{arccosh}(ax) - 225\sqrt{-1+ax}\sqrt{1+ax}(149 - 38a^2x^2 + 9a^4x^4) \operatorname{arccosh}(ax)^2 + 1125a^2x(15 - 10a^2x^2 + 3a^4x^4) \operatorname{arccosh}(ax)^3)}{16875a}$$

input `Integrate[(c - a^2*c*x^2)^2*ArcCosh[a*x]^3,x]`output `(c^2*(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(31841 - 842*a^2*x^2 + 81*a^4*x^4) + 30*a*x*(2235 - 190*a^2*x^2 + 27*a^4*x^4)*ArcCosh[a*x] - 225*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(149 - 38*a^2*x^2 + 9*a^4*x^4)*ArcCosh[a*x]^2 + 1125*a*x*(15 - 10*a^2*x^2 + 3*a^4*x^4)*ArcCosh[a*x]^3))/(16875*a)`**3.241.3 Rubi [A] (verified)**Time = 2.58 (sec) , antiderivative size = 431, normalized size of antiderivative = 1.11, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.850$ , Rules used = {6312, 27, 6312, 6294, 6330, 25, 6294, 83, 6304, 6309, 27, 960, 83, 1905, 1576, 1140, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(ax)^3 (c - a^2 cx^2)^2 dx$$

$$\downarrow \text{6312}$$

$$\frac{4}{5}c \int c(1 - a^2 x^2) \operatorname{arccosh}(ax)^3 dx - \frac{3}{5}ac^2 \int x(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{5}c^2 x(1 - a^2 x^2)^2 \operatorname{arccosh}(ax)^3$$

$$\downarrow \text{27}$$

$$\frac{4}{5}c^2 \int (1 - a^2 x^2) \operatorname{arccosh}(ax)^3 dx - \frac{3}{5}ac^2 \int x(ax - 1)^{3/2}(ax + 1)^{3/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{5}c^2 x(1 - a^2 x^2)^2 \operatorname{arccosh}(ax)^3$$

$$\downarrow \text{6312}$$



$$\frac{4}{5}c^2 \left( a \int x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2 dx + \frac{2}{3} \int \operatorname{arccosh}(ax)^3 dx + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 \right) - \frac{3}{5}ac^2 \int x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 6294

$$\frac{4}{5}c^2 \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \int \frac{x \operatorname{arccosh}(ax)^2}{\sqrt{ax-1}\sqrt{ax+1}} dx \right) + a \int x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2 dx + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 \right) - \frac{3}{5}ac^2 \int x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2 dx + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 6330

$$\frac{4}{5}c^2 \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \int \operatorname{arccosh}(ax) dx}{a} \right) \right) + a \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2}}{3a^2} \right) \right) - \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \int (1-ax)^2(ax+1)^2 \operatorname{arccosh}(ax) dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 25

$$\frac{4}{5}c^2 \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \int \operatorname{arccosh}(ax) dx}{a} \right) \right) + a \left( \frac{2 \int (1-ax)(ax+1) dx}{3a} \right) \right) - \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \int (1-ax)^2(ax+1)^2 \operatorname{arccosh}(ax) dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 6294

$$\frac{4}{5}c^2 \left( \frac{2}{3} \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - a \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a} \right) \right) \right) + a \left( \frac{2 \int (1-ax)(ax+1) dx}{3a} \right) - \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \int (1-ax)^2(ax+1)^2 \operatorname{arccosh}(ax) dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 83

$$\frac{4}{5}c^2 \left( a \left( \frac{2 \int (1-ax)(ax+1) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax) \right) + \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \int (1-ax)^2(ax+1)^2 \operatorname{arccosh}(ax) dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 6304

$$\frac{4}{5}c^2 \left( a \left( \frac{2 \int (1-a^2x^2) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + \frac{2}{3} \right) + \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \int (1-a^2x^2)^2 \operatorname{arccosh}(ax) dx}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 6309

$$\frac{4}{5}c^2 \left( a \left( \frac{2 \left( -a \int \frac{x(3-a^2x^2)}{3\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) \right) + \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -a \int \frac{x(3a^4x^4-10a^2x^2+15)}{15\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 27

$$\frac{4}{5}c^2 \left( a \left( \frac{2 \left( -\frac{1}{3}a \int \frac{x(3-a^2x^2)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) \right) + \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{1}{15}a \int \frac{x(3a^4x^4-10a^2x^2+15)}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 960

$$\frac{4}{5}c^2 \left( a \left( \frac{2 \left( -\frac{1}{3}a \left( \frac{7}{3} \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} \right) + \frac{(ax-1)^3}{5a} \right) \\ + \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{1}{15}a \int \frac{x(3a^4x^4-10a^2x^2+15)}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) \\ + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3$$

↓ 83

$$-\frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{1}{15}a \int \frac{x(3a^4x^4-10a^2x^2+15)}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) \\ + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3 +$$

$$\frac{4}{5}c^2 \left( \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) + x \operatorname{arccosh}(ax) \right)}{3a} \right) \right)$$

↓ 1905

$$-\frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \frac{x(3a^4x^4-10a^2x^2+15)}{\sqrt{a^2x^2-1}} dx + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) \\ + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3 +$$

$$\frac{4}{5}c^2 \left( \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) + x \operatorname{arccosh}(ax) \right)}{3a} \right) \right)$$

↓ 1576

$$-\frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \frac{3a^4x^4-10a^2x^2+15}{30\sqrt{ax-1}\sqrt{ax+1}} dx^2 + \frac{1}{5}a^4x^5 \operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3 \operatorname{arccosh}(ax) \right)}{5a} \right) \\ + \frac{1}{5}c^2x(1-a^2x^2)^2 \operatorname{arccosh}(ax)^3 +$$

$$\frac{4}{5}c^2 \left( \frac{1}{3}x(1-a^2x^2) \operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) + x \operatorname{arccosh}(ax) \right)}{3a} \right) \right)$$

$$\begin{aligned}
 & \downarrow 1140 \\
 & -\frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( -\frac{a\sqrt{a^2x^2-1} \int \left( 3(a^2x^2-1)^{3/2} - 4\sqrt{a^2x^2-1} + \frac{8}{\sqrt{a^2x^2-1}} \right) dx^2}{30\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{5}a^4x^5\operatorname{arccosh}(ax) \right)}{5a} \right. \\
 & \qquad \qquad \qquad \left. + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^3 + \right. \\
 & \frac{4}{5}c^2 \left( \frac{1}{3}x(1-a^2x^2)\operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3\operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) + x\operatorname{arccosh}(ax) \right)}{3a} \right) \right. \\
 & \qquad \qquad \qquad \downarrow 2009 \\
 & \qquad \qquad \qquad \left. + \frac{1}{5}c^2x(1-a^2x^2)^2\operatorname{arccosh}(ax)^3 + \right. \\
 & \frac{4}{5}c^2 \left( \frac{1}{3}x(1-a^2x^2)\operatorname{arccosh}(ax)^3 + a \left( \frac{2 \left( -\frac{1}{3}a^2x^3\operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) + x\operatorname{arccosh}(ax) \right)}{3a} \right) \right. \\
 & \qquad \qquad \qquad \left. \left. \left( \frac{3}{5}ac^2 \left( \frac{(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)^2}{5a^2} - \frac{2 \left( \frac{1}{5}a^4x^5\operatorname{arccosh}(ax) - \frac{2}{3}a^2x^3\operatorname{arccosh}(ax) - \frac{a\sqrt{a^2x^2-1} \left( \frac{6(a^2x^2-1)^{5/2}}{5a^2} \right)}{30\sqrt{ax-1}\sqrt{ax+1}} \right)}{5a} \right) \right) \right. \right.
 \end{aligned}$$

input `Int[(c - a^2*c*x^2)^2*ArcCosh[a*x]^3,x]`

output `(c^2*x*(1 - a^2*x^2)^2*ArcCosh[a*x]^3)/5 - (3*a*c^2*((( -1 + a*x)^(5/2)*(1 + a*x)^(5/2)*ArcCosh[a*x]^2)/(5*a^2) - (2*(-1/30*(a*Sqrt[-1 + a^2*x^2]*((1 + 6*Sqrt[-1 + a^2*x^2])/a^2 - (8*(-1 + a^2*x^2)^(3/2))/(3*a^2) + (6*(-1 + a^2*x^2)^(5/2))/(5*a^2))))/(Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + x*ArcCosh[a*x] - (2*a^2*x^3*ArcCosh[a*x])/3 + (a^4*x^5*ArcCosh[a*x])/5))/(5*a))/5 + (4*c^2*((x*(1 - a^2*x^2)*ArcCosh[a*x]^3)/3 + a*((( -1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x]^2)/(3*a^2) + (2*(-1/3*(a*((7*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(3*a^2) - (x^2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/3)) + x*ArcCosh[a*x] - (a^2*x^3*ArcCosh[a*x])/3))/(3*a)) + (2*(x*ArcCosh[a*x]^3 - 3*a*((Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^2)/a^2 - (2*(-((Sqrt[-1 + a*x]*Sqrt[1 + a*x])/a) + x*ArcCosh[a*x]))/a))/3))/5`

3.241.  $\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^3 dx$

## 3.241.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_)^(n_))*((e_) + (f_)*(x_)^(p_)), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 960 `Int[((e_)*(x_)^(m_))*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 1140 `Int[((d_) + (e_)*(x_)^(m_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 0]`
- rule 1576 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`
- rule 1905 `Int[((f_)*(x_)^(m_))*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c^n Int[x*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

**3.241.4 Maple [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 218, normalized size of antiderivative = 0.56

method	result
derivativedivides	$\frac{c^2 \left( 3375 \operatorname{arccosh}(ax)^3 a^5 x^5 - 2025 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^4 x^4 - 11250 a^3 x^3 \operatorname{arccosh}(ax)^3 + 8550 a^2 x^2 \operatorname{arccosh}(ax)^2 \right)}{c^2 \left( 3375 \operatorname{arccosh}(ax)^3 a^5 x^5 - 2025 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^4 x^4 - 11250 a^3 x^3 \operatorname{arccosh}(ax)^3 + 8550 a^2 x^2 \operatorname{arccosh}(ax)^2 \right)}$
default	$\frac{c^2 \left( 3375 \operatorname{arccosh}(ax)^3 a^5 x^5 - 2025 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^4 x^4 - 11250 a^3 x^3 \operatorname{arccosh}(ax)^3 + 8550 a^2 x^2 \operatorname{arccosh}(ax)^2 \right)}{c^2 \left( 3375 \operatorname{arccosh}(ax)^3 a^5 x^5 - 2025 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} a^4 x^4 - 11250 a^3 x^3 \operatorname{arccosh}(ax)^3 + 8550 a^2 x^2 \operatorname{arccosh}(ax)^2 \right)}$

```
input int((-a^2*c*x^2+c)^2*arccosh(a*x)^3,x,method=_RETURNVERBOSE)
```

```
output 1/16875/a*c^2*(3375*arccosh(a*x)^3*a^5*x^5-2025*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4-11250*a^3*x^3*arccosh(a*x)^3+8550*a^2*x^2*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+810*a^5*x^5*arccosh(a*x)-162*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4+16875*a*x*arccosh(a*x)^3-33525*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-5700*a^3*x^3*arccosh(a*x)+1684*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+67050*a*x*arccosh(a*x)-63682*(a*x-1)^(1/2)*(a*x+1)^(1/2))
```

**3.241.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.53

$$\int (c - a^2 c x^2)^2 \operatorname{arccosh}(ax)^3 dx$$

$$= \frac{1125 (3 a^5 c^2 x^5 - 10 a^3 c^2 x^3 + 15 a c^2 x) \log(ax + \sqrt{a^2 x^2 - 1})^3 - 225 (9 a^4 c^2 x^4 - 38 a^2 c^2 x^2 + 149 c^2) \sqrt{a^2 x^2 - 1}}{a}$$

```
input integrate((-a^2*c*x^2+c)^2*arccosh(a*x)^3,x, algorithm="fracas")
```

```
output 1/16875*(1125*(3*a^5*c^2*x^5 - 10*a^3*c^2*x^3 + 15*a*c^2*x)*log(a*x + sqrt(a^2*x^2 - 1))^3 - 225*(9*a^4*c^2*x^4 - 38*a^2*c^2*x^2 + 149*c^2)*sqrt(a^2*x^2 - 1)*log(a*x + sqrt(a^2*x^2 - 1))^2 + 30*(27*a^5*c^2*x^5 - 190*a^3*c^2*x^3 + 2235*a*c^2*x)*log(a*x + sqrt(a^2*x^2 - 1)) - 2*(81*a^4*c^2*x^4 - 842*a^2*c^2*x^2 + 31841*c^2)*sqrt(a^2*x^2 - 1))/a
```

**3.241.6 Sympy [F]**

$$\int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^3 dx = c^2 \left( \int (-2a^2x^2 \operatorname{acosh}^3(ax)) dx + \int a^4x^4 \operatorname{acosh}^3(ax) dx + \int \operatorname{acosh}^3(ax) dx \right)$$

input `integrate((-a**2*c*x**2+c)**2*acosh(a*x)**3,x)`

output `c**2*(Integral(-2*a**2*x**2*acosh(a*x)**3, x) + Integral(a**4*x**4*acosh(a*x)**3, x) + Integral(acosh(a*x)**3, x))`

**3.241.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 210, normalized size of antiderivative = 0.54

$$\begin{aligned} & \int (c - a^2cx^2)^2 \operatorname{arccosh}(ax)^3 dx \\ &= -\frac{1}{75} \left( 9\sqrt{a^2x^2 - 1}a^2c^2x^4 - 38\sqrt{a^2x^2 - 1}c^2x^2 + \frac{149\sqrt{a^2x^2 - 1}c^2}{a^2} \right) a \operatorname{arccosh}(ax)^2 \\ & \quad + \frac{1}{15} (3a^4c^2x^5 - 10a^2c^2x^3 + 15c^2x) \operatorname{arccosh}(ax)^3 \\ & \quad - \frac{2}{16875} \left( 81\sqrt{a^2x^2 - 1}a^2c^2x^4 - 842\sqrt{a^2x^2 - 1}c^2x^2 - \frac{15(27a^4c^2x^5 - 190a^2c^2x^3 + 2235c^2x) \operatorname{arccosh}(ax)}{a} \right) \end{aligned}$$

input `integrate((-a^2*c*x^2+c)^2*arccosh(a*x)^3,x, algorithm="maxima")`

output `-1/75*(9*sqrt(a^2*x^2 - 1)*a^2*c^2*x^4 - 38*sqrt(a^2*x^2 - 1)*c^2*x^2 + 149*sqrt(a^2*x^2 - 1)*c^2/a^2)*a*arccosh(a*x)^2 + 1/15*(3*a^4*c^2*x^5 - 10*a^2*c^2*x^3 + 15*c^2*x)*arccosh(a*x)^3 - 2/16875*(81*sqrt(a^2*x^2 - 1)*a^2*c^2*x^4 - 842*sqrt(a^2*x^2 - 1)*c^2*x^2 - 15*(27*a^4*c^2*x^5 - 190*a^2*c^2*x^3 + 2235*c^2*x)*arccosh(a*x)/a + 31841*sqrt(a^2*x^2 - 1)*c^2/a^2)*a`



**3.241.8 Giac [F(-2)]**

Exception generated.

$$\int (c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^2*arccosh(a*x)^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.241.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^3 dx = \int \operatorname{acosh}(ax)^3 (c - a^2 cx^2)^2 dx$$

input `int(acosh(a*x)^3*(c - a^2*c*x^2)^2,x)`

output `int(acosh(a*x)^3*(c - a^2*c*x^2)^2, x)`

### 3.242 $\int (c - a^2cx^2) \operatorname{arccosh}(ax)^3 dx$

3.242.1 Optimal result . . . . .	2153
3.242.2 Mathematica [A] (verified) . . . . .	2154
3.242.3 Rubi [A] (verified) . . . . .	2154
3.242.4 Maple [A] (verified) . . . . .	2158
3.242.5 Fracas [A] (verification not implemented) . . . . .	2159
3.242.6 Sympy [F] . . . . .	2159
3.242.7 Maxima [A] (verification not implemented) . . . . .	2159
3.242.8 Giac [F(-2)] . . . . .	2160
3.242.9 Mupad [F(-1)] . . . . .	2160

#### 3.242.1 Optimal result

Integrand size = 18, antiderivative size = 175

$$\int (c - a^2cx^2) \operatorname{arccosh}(ax)^3 dx = -\frac{122c\sqrt{-1+ax}\sqrt{1+ax}}{27a} + \frac{2}{27}acx^2\sqrt{-1+ax}\sqrt{1+ax}$$

$$+ \frac{14}{3}cx\operatorname{arccosh}(ax) - \frac{2}{9}a^2cx^3\operatorname{arccosh}(ax)$$

$$- \frac{2c\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{a}$$

$$+ \frac{c(-1+ax)^{3/2}(1+ax)^{3/2}\operatorname{arccosh}(ax)^2}{3a}$$

$$+ \frac{2}{3}cx\operatorname{arccosh}(ax)^3 + \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^3$$

```
output 14/3*c*x*arccosh(a*x)-2/9*a^2*c*x^3*arccosh(a*x)+1/3*c*(a*x-1)^(3/2)*(a*x+
1)^(3/2)*arccosh(a*x)^2/a+2/3*c*x*arccosh(a*x)^3+1/3*c*x*(-a^2*x^2+1)*arcc
osh(a*x)^3-122/27*c*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a+2/27*a*c*x^2*(a*x-1)^(1/
2)*(a*x+1)^(1/2)-2*c*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a
```

**3.242.2 Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.62

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^3 dx$$

$$= \frac{c(2\sqrt{-1+ax}\sqrt{1+ax}(-61+a^2x^2) - 6ax(-21+a^2x^2)\operatorname{arccosh}(ax) + 9\sqrt{-1+ax}\sqrt{1+ax}(-7+a^2x^2))}{27a}$$

input `Integrate[(c - a^2*c*x^2)*ArcCosh[a*x]^3,x]`

output `(c*(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(-61 + a^2*x^2) - 6*a*x*(-21 + a^2*x^2)*ArcCosh[a*x] + 9*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(-7 + a^2*x^2)*ArcCosh[a*x]^2 - 9*a*x*(-3 + a^2*x^2)*ArcCosh[a*x]^3))/(27*a)`

**3.242.3 Rubi [A] (verified)**

Time = 1.26 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.27, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$ , Rules used = {6312, 6294, 6330, 25, 6294, 83, 6304, 6309, 27, 960, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(ax)^3 (c - a^2 cx^2) dx$$

$$\downarrow \text{6312}$$

$$ac \int x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^2 dx + \frac{2}{3}c \int \operatorname{arccosh}(ax)^3 dx + \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^3$$

$$\downarrow \text{6294}$$

$$\frac{2}{3}c \left( x\operatorname{arccosh}(ax)^3 - 3a \int \frac{x\operatorname{arccosh}(ax)^2}{\sqrt{ax-1}\sqrt{ax+1}} dx \right) + ac \int x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^2 dx + \frac{1}{3}cx(1-a^2x^2)\operatorname{arccosh}(ax)^3$$

$$\downarrow \text{6330}$$

$$\begin{aligned}
& \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \int \operatorname{arccosh}(ax) dx}{a} \right) \right) + \\
& ac \left( \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} - \frac{2 \int -((1-ax)(ax+1) \operatorname{arccosh}(ax)) dx}{3a} \right) + \\
& \qquad \qquad \qquad \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^3 \\
& \qquad \qquad \qquad \downarrow \text{25} \\
& \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \int \operatorname{arccosh}(ax) dx}{a} \right) \right) + \\
& ac \left( \frac{2 \int (1-ax)(ax+1) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \\
& \qquad \qquad \qquad \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^3 \\
& \qquad \qquad \qquad \downarrow \text{6294} \\
& \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - a \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a} \right) \right) + \\
& ac \left( \frac{2 \int (1-ax)(ax+1) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \\
& \qquad \qquad \qquad \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^3 \\
& \qquad \qquad \qquad \downarrow \text{83} \\
& ac \left( \frac{2 \int (1-ax)(ax+1) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \\
& \qquad \qquad \qquad \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^3 + \\
& \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a} \right) \right) \\
& \qquad \qquad \qquad \downarrow \text{6304} \\
& ac \left( \frac{2 \int (1-a^2x^2) \operatorname{arccosh}(ax) dx}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) + \\
& \qquad \qquad \qquad \frac{1}{3}cx(1-a^2x^2) \operatorname{arccosh}(ax)^3 + \\
& \frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a} \right) \right) \\
& \qquad \qquad \qquad \downarrow \text{6309}
\end{aligned}$$

$$ac \left( \frac{2 \left( -a \int \frac{x(3-a^2x^2)}{3\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) +$$

$$\frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a} \right) \right)$$

↓ 27

$$ac \left( \frac{2 \left( -\frac{1}{3}a \int \frac{x(3-a^2x^2)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) +$$

$$\frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a} \right) \right)$$

↓ 960

$$ac \left( \frac{2 \left( -\frac{1}{3}a \left( \frac{7}{3} \int \frac{x}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) - \frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) + x \operatorname{arccosh}(ax) \right)}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) +$$

$$\frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a} \right) \right)$$

↓ 83

$$ac \left( \frac{2 \left( -\frac{1}{3}a^2x^3 \operatorname{arccosh}(ax) - \frac{1}{3}a \left( \frac{7\sqrt{ax-1}\sqrt{ax+1}}{3a^2} - \frac{1}{3}x^2\sqrt{ax-1}\sqrt{ax+1} \right) + x \operatorname{arccosh}(ax) \right)}{3a} + \frac{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)^2}{3a^2} \right) +$$

$$\frac{2}{3}c \left( x \operatorname{arccosh}(ax)^3 - 3a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^2}{a^2} - \frac{2 \left( x \operatorname{arccosh}(ax) - \frac{\sqrt{ax-1}\sqrt{ax+1}}{a} \right)}{a} \right) \right)$$

input `Int[(c - a^2*c*x^2)*ArcCosh[a*x]^3,x]`

```
output (c*x*(1 - a^2*x^2)*ArcCosh[a*x]^3)/3 + a*c*((( -1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x]^2)/(3*a^2) + (2*(-1/3*(a*((7*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/3)))/(3*a^2) - (x^2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/3)) + x*ArcCosh[a*x] - (a^2*x^3*ArcCosh[a*x])/3)/(3*a) + (2*c*(x*ArcCosh[a*x]^3 - 3*a*((Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^2)/a^2 - (2*(-((Sqrt[-1 + a*x]*Sqrt[1 + a*x])/a) + x*ArcCosh[a*x]))/a))/3
```

### 3.242.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 83 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]
```

```
rule 960 Int[((e_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

```
rule 6294 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]
```

```
rule 6304 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]
```

rule 6309 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

### 3.242.4 Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.80

method	result
derivativedivides	$-\frac{c(9a^3x^3 \operatorname{arccosh}(ax)^3 - 9a^2x^2 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} - 27ax \operatorname{arccosh}(ax)^3 + 63 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} + 6a^3 \operatorname{arccosh}(ax) - 2a^2 \sqrt{ax-1} \sqrt{ax+1} - 126a^2 \operatorname{arccosh}(ax) + 122 \sqrt{ax-1} \sqrt{ax+1})}{27a}$
default	$-\frac{c(9a^3x^3 \operatorname{arccosh}(ax)^3 - 9a^2x^2 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} - 27ax \operatorname{arccosh}(ax)^3 + 63 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} + 6a^3 \operatorname{arccosh}(ax) - 2a^2 \sqrt{ax-1} \sqrt{ax+1} - 126a^2 \operatorname{arccosh}(ax) + 122 \sqrt{ax-1} \sqrt{ax+1})}{27a}$

input `int((-a^2*c*x^2+c)*arccosh(a*x)^3,x,method=_RETURNVERBOSE)`

output `-1/27/a*c*(9*a^3*x^3*arccosh(a*x)^3-9*a^2*x^2*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-27*a*x*arccosh(a*x)^3+63*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+6*a^3*x^3*arccosh(a*x)-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-126*a*x*arccosh(a*x)+122*(a*x-1)^(1/2)*(a*x+1)^(1/2))`

---

3.242.  $\int (c - a^2cx^2) \operatorname{arccosh}(ax)^3 dx$

**3.242.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.80

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^3 dx = \frac{9(a^3 cx^3 - 3acx) \log(ax + \sqrt{a^2 x^2 - 1})^3 - 9(a^2 cx^2 - 7c) \sqrt{a^2 x^2 - 1} \log(ax + \sqrt{a^2 x^2 - 1})^2 + 6(a^3 cx^3 - 21acx) \log(ax + \sqrt{a^2 x^2 - 1}) - 2(a^2 cx^2 - 61c) \sqrt{a^2 x^2 - 1}}{27a}$$

input `integrate((-a^2*c*x^2+c)*arccosh(a*x)^3,x, algorithm="fricas")`output `-1/27*(9*(a^3*c*x^3 - 3*a*c*x)*log(a*x + sqrt(a^2*x^2 - 1))^3 - 9*(a^2*c*x^2 - 7*c)*sqrt(a^2*x^2 - 1)*log(a*x + sqrt(a^2*x^2 - 1))^2 + 6*(a^3*c*x^3 - 21*a*c*x)*log(a*x + sqrt(a^2*x^2 - 1)) - 2*(a^2*c*x^2 - 61*c)*sqrt(a^2*x^2 - 1))/a`**3.242.6 Sympy [F]**

$$\int (c - a^2 cx^2) \operatorname{arccosh}(ax)^3 dx = -c \left( \int a^2 x^2 \operatorname{acosh}^3(ax) dx + \int (-\operatorname{acosh}^3(ax)) dx \right)$$

input `integrate((-a**2*c*x**2+c)*acosh(a*x)**3,x)`output `-c*(Integral(a**2*x**2*acosh(a*x)**3, x) + Integral(-acosh(a*x)**3, x))`**3.242.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.71

$$\begin{aligned} & \int (c - a^2 cx^2) \operatorname{arccosh}(ax)^3 dx \\ &= \frac{1}{3} \left( \sqrt{a^2 x^2 - 1} cx^2 - \frac{7 \sqrt{a^2 x^2 - 1} c}{a^2} \right) a \operatorname{arccosh}(ax)^2 - \frac{1}{3} (a^2 cx^3 - 3cx) \operatorname{arccosh}(ax)^3 \\ &+ \frac{2}{27} \left( \sqrt{a^2 x^2 - 1} cx^2 - \frac{3(a^2 cx^3 - 21cx) \operatorname{arccosh}(ax)}{a} - \frac{61 \sqrt{a^2 x^2 - 1} c}{a^2} \right) a \end{aligned}$$



input `integrate((-a^2*c*x^2+c)*arccosh(a*x)^3,x, algorithm="maxima")`

output `1/3*(sqrt(a^2*x^2 - 1)*c*x^2 - 7*sqrt(a^2*x^2 - 1)*c/a^2)*a*arccosh(a*x)^2  
- 1/3*(a^2*c*x^3 - 3*c*x)*arccosh(a*x)^3 + 2/27*(sqrt(a^2*x^2 - 1)*c*x^2  
- 3*(a^2*c*x^3 - 21*c*x)*arccosh(a*x)/a - 61*sqrt(a^2*x^2 - 1)*c/a^2)*a`

### 3.242.8 Giac [F(-2)]

Exception generated.

$$\int (c - a^2cx^2) \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)*arccosh(a*x)^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.242.9 Mupad [F(-1)]

Timed out.

$$\int (c - a^2cx^2) \operatorname{arccosh}(ax)^3 dx = \int \operatorname{acosh}(ax)^3 (c - a^2cx^2) dx$$

input `int(acosh(a*x)^3*(c - a^2*c*x^2),x)`

output `int(acosh(a*x)^3*(c - a^2*c*x^2), x)`

### 3.243 $\int \frac{\operatorname{arccosh}(ax)^3}{c-a^2cx^2} dx$

3.243.1 Optimal result . . . . .	2161
3.243.2 Mathematica [A] (verified) . . . . .	2162
3.243.3 Rubi [C] (verified) . . . . .	2162
3.243.4 Maple [A] (verified) . . . . .	2165
3.243.5 Fricas [F] . . . . .	2165
3.243.6 Sympy [F] . . . . .	2166
3.243.7 Maxima [F] . . . . .	2166
3.243.8 Giac [F] . . . . .	2166
3.243.9 Mupad [F(-1)] . . . . .	2167

#### 3.243.1 Optimal result

Integrand size = 20, antiderivative size = 144

$$\int \frac{\operatorname{arccosh}(ax)^3}{c-a^2cx^2} dx = \frac{2\operatorname{arccosh}(ax)^3 \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac} + \frac{3\operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{ac} - \frac{3\operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{ac} - \frac{6\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)})}{ac} + \frac{6\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)})}{ac} + \frac{6 \operatorname{PolyLog}(4, -e^{\operatorname{arccosh}(ax)})}{ac} - \frac{6 \operatorname{PolyLog}(4, e^{\operatorname{arccosh}(ax)})}{ac}$$

```
output 2*arccosh(a*x)^3*arctanh(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c+3*arccosh(a*x)^2*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c-3*arccosh(a*x)^2*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c-6*arccosh(a*x)*polylog(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c+6*arccosh(a*x)*polylog(3,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c+6*polylog(4,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c-6*polylog(4,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))/a/c
```

**3.243.2 Mathematica [A] (verified)**

Time = 0.06 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.90

$$\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2cx^2} dx$$

$$= -\operatorname{arccosh}(ax)^3 \log(1 - e^{\operatorname{arccosh}(ax)}) + \operatorname{arccosh}(ax)^3 \log(1 + e^{\operatorname{arccosh}(ax)}) + 3\operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})$$

input `Integrate[ArcCosh[a*x]^3/(c - a^2*c*x^2), x]`

output `(-(ArcCosh[a*x]^3*Log[1 - E^ArcCosh[a*x]]) + ArcCosh[a*x]^3*Log[1 + E^ArcCosh[a*x]]) + 3*ArcCosh[a*x]^2*PolyLog[2, -E^ArcCosh[a*x]] - 3*ArcCosh[a*x]^2*PolyLog[2, E^ArcCosh[a*x]] - 6*ArcCosh[a*x]*PolyLog[3, -E^ArcCosh[a*x]] + 6*ArcCosh[a*x]*PolyLog[3, E^ArcCosh[a*x]] + 6*PolyLog[4, -E^ArcCosh[a*x]] - 6*PolyLog[4, E^ArcCosh[a*x]])/(a*c)`

**3.243.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.60 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.89, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$ , Rules used = {6318, 3042, 26, 4670, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2cx^2} dx$$

$$\downarrow \text{6318}$$

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)$$

$$\frac{ac}{ac}$$

$$\downarrow \text{3042}$$

$$\int i\operatorname{arccosh}(ax)^3 \csc(i\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)$$

$$\frac{ac}{ac}$$

$$\downarrow \text{26}$$

---

3.243.  $\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2cx^2} dx$

$$\frac{i \int \operatorname{arccosh}(ax)^3 \csc(i \operatorname{arccosh}(ax)) d \operatorname{arccosh}(ax)}{ac}$$

↓ 4670

$$\frac{i(3i \int \operatorname{arccosh}(ax)^2 \log(1 - e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - 3i \int \operatorname{arccosh}(ax)^2 \log(1 + e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2 \int \operatorname{arccosh}(ax) \log(1 - e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - 2 \int \operatorname{arccosh}(ax) \log(1 + e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2 \int \log(1 - e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - 2 \int \log(1 + e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax))}{ac}$$

↓ 3011

$$\frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})) - 2 \int \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2 \int \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax))}{ac}$$

↓ 7163

$$\frac{i(-3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \int \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax)) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)})) + 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) - \int \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax)) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)})) - 2 \int \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2 \int \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax))}{ac}$$

↓ 2720

$$\frac{i(-3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d e^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) + 2 \int \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax)) + 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) - \int e^{\operatorname{arccosh}(ax)} \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d e^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) + 2 \int \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax)) - 2 \int \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2 \int \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax))}{ac}$$

↓ 7143

$$\frac{i(2i \operatorname{arccosh}(ax)^3 \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) - 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \operatorname{PolyLog}(4, -e^{\operatorname{arccosh}(ax)}) + 2 \int \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax)) + 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) - \operatorname{PolyLog}(4, e^{\operatorname{arccosh}(ax)}) + 2 \int \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax)) - 2 \int \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2 \int \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax))}{ac}$$

input `Int[ArcCosh[a*x]^3/(c - a^2*c*x^2), x]`

output `((-I)*((2*I)*ArcCosh[a*x]^3*ArcTanh[E^ArcCosh[a*x]] - (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, -E^ArcCosh[a*x]]) + 2*(ArcCosh[a*x]*PolyLog[3, -E^ArcCosh[a*x]] - PolyLog[4, -E^ArcCosh[a*x]])) + (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, E^ArcCosh[a*x]]) + 2*(ArcCosh[a*x]*PolyLog[3, E^ArcCosh[a*x]] - PolyLog[4, E^ArcCosh[a*x]]))))/(a*c)`

## 3.243.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`
- rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`
- rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

```
rule 7163 Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.
)*(x_)))]^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### 3.243.4 Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.76

method	result
derivativedivides	$\frac{\operatorname{arccosh}(ax)^3 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{c} + \frac{3 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c} - \frac{6 \operatorname{arccosh}(ax) \operatorname{polylog}(3, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c}$
default	$\frac{\operatorname{arccosh}(ax)^3 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{c} + \frac{3 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c} - \frac{6 \operatorname{arccosh}(ax) \operatorname{polylog}(3, -ax-\sqrt{ax-1}\sqrt{ax+1})}{c}$

```
input int(arccosh(a*x)^3/(-a^2*c*x^2+c),x,method=_RETURNVERBOSE)
```

```
output 1/a*(1/c*arccosh(a*x)^3*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/c*arccosh(
a*x)^2*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-6/c*arccosh(a*x)*polylo
g(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+6/c*polylog(4,-a*x-(a*x-1)^(1/2)*(a
*x+1)^(1/2))-1/c*arccosh(a*x)^3*ln(1-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-3/c*a
rccosh(a*x)^2*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+6/c*arccosh(a*x)*
polylog(3,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))-6/c*polylog(4,a*x+(a*x-1)^(1/2)
*(a*x+1)^(1/2)))
```

### 3.243.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2cx^2} dx = \int -\frac{\operatorname{arccosh}(ax)^3}{a^2cx^2 - c} dx$$

```
input integrate(arccosh(a*x)^3/(-a^2*c*x^2+c),x, algorithm="fracas")
```

```
output integral(-arccosh(a*x)^3/(a^2*c*x^2 - c), x)
```

## 3.243.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2cx^2} dx = -\int \frac{\operatorname{acosh}^3(ax)}{a^2x^2 - 1} dx$$

input `integrate(acosh(a*x)**3/(-a**2*c*x**2+c), x)`

output `-Integral(acosh(a*x)**3/(a**2*x**2 - 1), x)/c`

## 3.243.7 Maxima [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2cx^2} dx = \int -\frac{\operatorname{arcosh}(ax)^3}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c), x, algorithm="maxima")`

output `1/2*(log(a*x + 1) - log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^3 / (a*c) - integrate(3/2*((a*x*log(a*x + 1) - a*x*log(a*x - 1))*sqrt(a*x + 1)*sqrt(a*x - 1) + (a^2*x^2 - 1)*log(a*x + 1) - (a^2*x^2 - 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^2/(a^3*c*x^3 - a*c*x + (a^2*c*x^2 - c)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

## 3.243.8 Giac [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2cx^2} dx = \int -\frac{\operatorname{arcosh}(ax)^3}{a^2cx^2 - c} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c), x, algorithm="giac")`

output `integrate(-arccosh(a*x)^3/(a^2*c*x^2 - c), x)`

**3.243.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{c - a^2 cx^2} dx = \int \frac{\operatorname{acosh}(ax)^3}{c - a^2 cx^2} dx$$

input `int(acosh(a*x)^3/(c - a^2*c*x^2), x)`output `int(acosh(a*x)^3/(c - a^2*c*x^2), x)`



$$3.244 \quad \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^2} dx$$

3.244.1 Optimal result . . . . .	2168
3.244.2 Mathematica [A] (verified) . . . . .	2169
3.244.3 Rubi [C] (verified) . . . . .	2170
3.244.4 Maple [A] (verified) . . . . .	2176
3.244.5 Fracas [F] . . . . .	2177
3.244.6 Sympy [F] . . . . .	2177
3.244.7 Maxima [F] . . . . .	2178
3.244.8 Giac [F] . . . . .	2178
3.244.9 Mupad [F(-1)] . . . . .	2178

### 3.244.1 Optimal result

Integrand size = 20, antiderivative size = 260

$$\begin{aligned} \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^2} dx = & -\frac{3\operatorname{arccosh}(ax)^2}{2ac^2\sqrt{-1+ax}\sqrt{1+ax}} + \frac{x\operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} \\ & - \frac{6\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac^2} \\ & + \frac{\operatorname{arccosh}(ax)^3\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac^2} - \frac{3\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{ac^2} \\ & + \frac{3\operatorname{arccosh}(ax)^2\operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})}{2ac^2} \\ & + \frac{3\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{ac^2} - \frac{3\operatorname{arccosh}(ax)^2\operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{2ac^2} \\ & - \frac{3\operatorname{arccosh}(ax)\operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)})}{ac^2} \\ & + \frac{3\operatorname{arccosh}(ax)\operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)})}{ac^2} \\ & + \frac{3\operatorname{PolyLog}(4, -e^{\operatorname{arccosh}(ax)})}{ac^2} - \frac{3\operatorname{PolyLog}(4, e^{\operatorname{arccosh}(ax)})}{ac^2} \end{aligned}$$

output  $\frac{1}{2}x \operatorname{arccosh}(ax)^3/c^2/(-a^2x^2+1)-6 \operatorname{arccosh}(ax) \operatorname{arctanh}(ax+(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2+\operatorname{arccosh}(ax)^3 \operatorname{arctanh}(ax+(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2-3 \operatorname{polylog}(2,-ax-(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2+3/2 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2,-ax-(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2+3 \operatorname{polylog}(2,ax+(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2-3/2 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2,ax+(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2-3 \operatorname{arccosh}(ax) \operatorname{polylog}(3,-ax-(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2+3 \operatorname{arccosh}(ax) \operatorname{polylog}(3,ax+(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2+3 \operatorname{polylog}(4,-ax-(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2-3 \operatorname{polylog}(4,ax+(ax-1)^{(1/2)}(ax+1)^{(1/2)})/a/c^2-3/2 \operatorname{arccosh}(ax)^2/a/c^2/(ax-1)^{(1/2)}/(ax+1)^{(1/2)}$

### 3.244.2 Mathematica [A] (verified)

Time = 1.41 (sec) , antiderivative size = 276, normalized size of antiderivative = 1.06

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^2} dx$$

$$= \frac{-\pi^4 + 2 \operatorname{arccosh}(ax)^4 - 12 \operatorname{arccosh}(ax)^2 \coth\left(\frac{1}{2} \operatorname{arccosh}(ax)\right) - 2 \operatorname{arccosh}(ax)^3 \operatorname{csch}^2\left(\frac{1}{2} \operatorname{arccosh}(ax)\right) + 48 \operatorname{arccosh}(ax) \operatorname{Log}[1 - E^{-\operatorname{arccosh}(ax)}] - 48 \operatorname{arccosh}(ax) \operatorname{Log}[1 + E^{-\operatorname{arccosh}(ax)}] + 8 \operatorname{arccosh}(ax)^3 \operatorname{Log}[1 + E^{-\operatorname{arccosh}(ax)}] - 8 \operatorname{arccosh}(ax)^3 \operatorname{Log}[1 - E^{\operatorname{arccosh}(ax)}] - 24(-2 + \operatorname{arccosh}(ax)^2) \operatorname{PolyLog}[2, -E^{-\operatorname{arccosh}(ax)}] - 48 \operatorname{PolyLog}[2, E^{-\operatorname{arccosh}(ax)}] - 24 \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}[2, E^{\operatorname{arccosh}(ax)}] - 48 \operatorname{arccosh}(ax) \operatorname{PolyLog}[3, -E^{-\operatorname{arccosh}(ax)}] + 48 \operatorname{arccosh}(ax) \operatorname{PolyLog}[3, E^{\operatorname{arccosh}(ax)}] - 48 \operatorname{PolyLog}[4, -E^{-\operatorname{arccosh}(ax)}] - 48 \operatorname{PolyLog}[4, E^{\operatorname{arccosh}(ax)}] - 2 \operatorname{arccosh}(ax)^3 \operatorname{Sech}[\operatorname{arccosh}(ax)/2]^2 + 12 \operatorname{arccosh}(ax)^2 \operatorname{Tanh}[\operatorname{arccosh}(ax)/2]}{(16a^2c^2)}$$

input `Integrate[ArcCosh[a*x]^3/(c - a^2*c*x^2)^2,x]`

output  $(-\pi^4 + 2 \operatorname{ArcCosh}[a*x]^4 - 12 \operatorname{ArcCosh}[a*x]^2 \operatorname{Coth}[\operatorname{ArcCosh}[a*x]/2] - 2 \operatorname{ArcCosh}[a*x]^3 \operatorname{Csch}[\operatorname{ArcCosh}[a*x]/2]^2 + 48 \operatorname{ArcCosh}[a*x] \operatorname{Log}[1 - E^{-\operatorname{ArcCosh}[a*x]}] - 48 \operatorname{ArcCosh}[a*x] \operatorname{Log}[1 + E^{-\operatorname{ArcCosh}[a*x]}] + 8 \operatorname{ArcCosh}[a*x]^3 \operatorname{Log}[1 + E^{-\operatorname{ArcCosh}[a*x]}] - 8 \operatorname{ArcCosh}[a*x]^3 \operatorname{Log}[1 - E^{\operatorname{ArcCosh}[a*x]}] - 24(-2 + \operatorname{ArcCosh}[a*x]^2) \operatorname{PolyLog}[2, -E^{-\operatorname{ArcCosh}[a*x]}] - 48 \operatorname{PolyLog}[2, E^{-\operatorname{ArcCosh}[a*x]}] - 24 \operatorname{ArcCosh}[a*x]^2 \operatorname{PolyLog}[2, E^{\operatorname{ArcCosh}[a*x]}] - 48 \operatorname{ArcCosh}[a*x] \operatorname{PolyLog}[3, -E^{-\operatorname{ArcCosh}[a*x]}] + 48 \operatorname{ArcCosh}[a*x] \operatorname{PolyLog}[3, E^{\operatorname{ArcCosh}[a*x]}] - 48 \operatorname{PolyLog}[4, -E^{-\operatorname{ArcCosh}[a*x]}] - 48 \operatorname{PolyLog}[4, E^{\operatorname{ArcCosh}[a*x]}] - 2 \operatorname{ArcCosh}[a*x]^3 \operatorname{Sech}[\operatorname{ArcCosh}[a*x]/2]^2 + 12 \operatorname{ArcCosh}[a*x]^2 \operatorname{Tanh}[\operatorname{ArcCosh}[a*x]/2])/(16a^2c^2)$

### 3.244.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 2.56 (sec) , antiderivative size = 244, normalized size of antiderivative = 0.94, number of steps used = 20, number of rules used = 19,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.950$ , Rules used = {6316, 27, 6318, 3042, 26, 4670, 3011, 6330, 25, 6304, 6318, 3042, 26, 4670, 2715, 2838, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^2} dx \\
 & \quad \downarrow \text{6316} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)^3}{c(1-a^2x^2)} dx}{2c} + \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} + \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} \\
 & \quad \downarrow \text{27} \\
 & \frac{\int \frac{\operatorname{arccosh}(ax)^3}{1-a^2x^2} dx}{2c^2} + \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} + \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} \\
 & \quad \downarrow \text{6318} \\
 & \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} - \frac{\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} \\
 & \quad \downarrow \text{3042} \\
 & \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} - \frac{\int i \operatorname{arccosh}(ax)^3 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} \\
 & \quad \downarrow \text{26} \\
 & \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} - \frac{i \int \operatorname{arccosh}(ax)^3 \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2ac^2} + \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} \\
 & \quad \downarrow \text{4670} \\
 & \frac{i(3i \int \operatorname{arccosh}(ax)^2 \log(1 - e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - 3i \int \operatorname{arccosh}(ax)^2 \log(1 + e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2}{2ac^2} \\
 & \quad \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} + \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}
 \end{aligned}$$

---

3.244.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^2} dx$

↓ 3011

$$\frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int$$

$$\frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{2c^2} + \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

↓ 6330

$$3a \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{(1-ax)(ax+1)} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right)$$

$$\frac{2c^2}{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int$$

$$\frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

↓ 25

$$3a \left( -\frac{2 \int \frac{\operatorname{arccosh}(ax)}{(1-ax)(ax+1)} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right)$$

$$\frac{2c^2}{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int$$

$$\frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

↓ 6304

$$3a \left( -\frac{2 \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right)$$

$$\frac{2c^2}{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int$$

$$\frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

↓ 6318

$$3a \left( \frac{2 \int \frac{\operatorname{arccosh}(ax) \operatorname{darccosh}(ax)}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} dx}{a^2} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right)$$

$$\frac{2c^2}{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int$$

$$\frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

---

3.244.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^2} dx$

$$\frac{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2 \int i \operatorname{arccosh}(ax) \csc(i \operatorname{arccosh}(ax)) d \operatorname{arccosh}(ax)}{a^2} \right)}{2c^2} -$$

$$\frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} dx$$

↓ 3042

$$\frac{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i \int \operatorname{arccosh}(ax) \csc(i \operatorname{arccosh}(ax)) d \operatorname{arccosh}(ax)}{a^2} \right)}{2c^2} -$$

$$\frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} dx$$

↓ 26

$$\frac{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i \left( i \int \log(1-e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - i \int \log(1+e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) \right)}{a^2} \right)}{2c^2} -$$

$$\frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} dx$$

↓ 4670

$$\frac{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i \left( i \int e^{-\operatorname{arccosh}(ax)} \log(1-e^{\operatorname{arccosh}(ax)}) d e^{\operatorname{arccosh}(ax)} - i \int e^{-\operatorname{arccosh}(ax)} \log(1+e^{\operatorname{arccosh}(ax)}) d e^{\operatorname{arccosh}(ax)} + 2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) \right)}{a^2} \right)}{2c^2} -$$

$$\frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \frac{x \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)} dx$$

↓ 2715

↓ 2838

---

3.244.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^2} dx$

$$\frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{a^2} \right)} + \frac{2c^2 \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

↓ 7163

$$\frac{i(-3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \int \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax)) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)})) + 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) - \int \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax)) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}))}{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{a^2} \right)} + \frac{2c^2 \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

↓ 2720

$$\frac{i(-3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d e^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)})) + 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) - \int e^{\operatorname{arccosh}(ax)} \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) d e^{\operatorname{arccosh}(ax)} - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}))}{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{a^2} \right)} + \frac{2c^2 \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

↓ 7143

$$\frac{i(2i \operatorname{arccosh}(ax)^3 \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) - 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - \operatorname{PolyLog}(4, -e^{\operatorname{arccosh}(ax)})) + 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, e^{\operatorname{arccosh}(ax)}) - \operatorname{PolyLog}(4, e^{\operatorname{arccosh}(ax)}))}{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{a^2} \right)} + \frac{2c^2 \operatorname{arccosh}(ax)^3}{2c^2(1-a^2x^2)}$$

input `Int[ArcCosh[a*x]^3/(c - a^2*c*x^2)^2,x]`

```
output (x*ArcCosh[a*x]^3)/(2*c^2*(1 - a^2*x^2)) + (3*a*(-(ArcCosh[a*x]^2/(a^2*Sqr
t[-1 + a*x]*Sqrt[1 + a*x])) + ((2*I)*((2*I)*ArcCosh[a*x]*ArcTanh[E^ArcCosh
[a*x]] + I*PolyLog[2, -E^ArcCosh[a*x]] - I*PolyLog[2, E^ArcCosh[a*x]]))/a^
2))/(2*c^2) - ((I/2)*((2*I)*ArcCosh[a*x]^3*ArcTanh[E^ArcCosh[a*x]] - (3*I)
*(-(ArcCosh[a*x]^2*PolyLog[2, -E^ArcCosh[a*x]]) + 2*(ArcCosh[a*x]*PolyLog[
3, -E^ArcCosh[a*x]] - PolyLog[4, -E^ArcCosh[a*x]])) + (3*I)*(-(ArcCosh[a*x
]^2*PolyLog[2, E^ArcCosh[a*x]]) + 2*(ArcCosh[a*x]*PolyLog[3, E^ArcCosh[a*x
]] - PolyLog[4, E^ArcCosh[a*x]])))))/(a*c^2)
```

### 3.244.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 2715 Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 2838 Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.) * (x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_)^(p_.)*(d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`



rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### 3.244.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 416, normalized size of antiderivative = 1.60

method	result
derivativedivides	$-\frac{\operatorname{arccosh}(ax)^2(ax \operatorname{arccosh}(ax) + 3\sqrt{ax-1}\sqrt{ax+1})}{2(a^2x^2-1)c^2} + \frac{\operatorname{arccosh}(ax)^3 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{2c^2} + \frac{3 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2, -ax-\sqrt{ax-1}\sqrt{ax+1})}{2c^2}$
default	$-\frac{\operatorname{arccosh}(ax)^2(ax \operatorname{arccosh}(ax) + 3\sqrt{ax-1}\sqrt{ax+1})}{2(a^2x^2-1)c^2} + \frac{\operatorname{arccosh}(ax)^3 \ln(1+ax+\sqrt{ax-1}\sqrt{ax+1})}{2c^2} + \frac{3 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2, -ax-\sqrt{ax-1}\sqrt{ax+1})}{2c^2}$

input `int(arccosh(a*x)^3/(-a^2*c*x^2+c)^2,x,method=_RETURNVERBOSE)`

```
output 1/a*(-1/2/(a^2*x^2-1)*arccosh(a*x)^2*(a*x*arccosh(a*x)+3*(a*x-1)^(1/2)*(a*x+1)^(1/2))/c^2+1/2/c^2*arccosh(a*x)^3*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/2/c^2*arccosh(a*x)^2*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-3/c^2*arccosh(a*x)*polylog(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/c^2*polylog(4,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-1/2/c^2*arccosh(a*x)^3*ln(1-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-3/2/c^2*arccosh(a*x)^2*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/c^2*arccosh(a*x)*polylog(3,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))-3/c^2*polylog(4,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))-3/c^2*arccosh(a*x)*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))-3/c^2*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/c^2*arccosh(a*x)*ln(1-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/c^2*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))
```

### 3.244.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(a^2cx^2 - c)^2} dx$$

```
input integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^2,x, algorithm="fricas")
```

```
output integral(arccosh(a*x)^3/(a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2), x)
```

### 3.244.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{acosh}^3(ax)}{a^4x^4 - 2a^2x^2 + 1} dx$$

```
input integrate(acosh(a*x)**3/(-a**2*c*x**2+c)**2,x)
```

```
output Integral(acosh(a*x)**3/(a**4*x**4 - 2*a**2*x**2 + 1), x)/c**2
```

**3.244.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(a^2cx^2 - c)^2} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^2,x, algorithm="maxima")`

output `-1/4*(2*a*x - (a^2*x^2 - 1)*log(a*x + 1) + (a^2*x^2 - 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^3/(a^3*c^2*x^2 - a*c^2) - integrate(-3/4*(2*a^3*x^3 + (2*a^2*x^2 - (a^3*x^3 - a*x)*log(a*x + 1) + (a^3*x^3 - a*x)*log(a*x - 1))*sqrt(a*x + 1)*sqrt(a*x - 1) - 2*a*x - (a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1) + (a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^2/(a^5*c^2*x^5 - 2*a^3*c^2*x^3 + a*c^2*x + (a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

**3.244.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(a^2cx^2 - c)^2} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^2,x, algorithm="giac")`

output `integrate(arccosh(a*x)^3/(a^2*c*x^2 - c)^2, x)`

**3.244.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^2} dx = \int \frac{\operatorname{acosh}(ax)^3}{(c - a^2cx^2)^2} dx$$

input `int(acosh(a*x)^3/(c - a^2*c*x^2)^2,x)`

output `int(acosh(a*x)^3/(c - a^2*c*x^2)^2, x)`

**3.245**      $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$

3.245.1 Optimal result . . . . .	2179
3.245.2 Mathematica [A] (warning: unable to verify) . . . . .	2180
3.245.3 Rubi [C] (verified) . . . . .	2181
3.245.4 Maple [A] (verified) . . . . .	2189
3.245.5 Fricas [F] . . . . .	2190
3.245.6 Sympy [F] . . . . .	2190
3.245.7 Maxima [F] . . . . .	2191
3.245.8 Giac [F] . . . . .	2191
3.245.9 Mupad [F(-1)] . . . . .	2191

**3.245.1 Optimal result**

Integrand size = 20, antiderivative size = 387

$$\begin{aligned} \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx = & \frac{1}{4ac^3\sqrt{-1+ax}\sqrt{1+ax}} - \frac{x\operatorname{arccosh}(ax)}{4c^3(1-a^2x^2)} \\ & + \frac{\operatorname{arccosh}(ax)^2}{4ac^3(-1+ax)^{3/2}(1+ax)^{3/2}} \\ & - \frac{9\operatorname{arccosh}(ax)^2}{8ac^3\sqrt{-1+ax}\sqrt{1+ax}} + \frac{x\operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\ & + \frac{3x\operatorname{arccosh}(ax)^3}{8c^3(1-a^2x^2)} - \frac{5\operatorname{arccosh}(ax)\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{ac^3} \\ & + \frac{3\operatorname{arccosh}(ax)^3\operatorname{arctanh}(e^{\operatorname{arccosh}(ax)})}{4ac^3} - \frac{5\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(ax)})}{2ac^3} \\ & + \frac{9\operatorname{arccosh}(ax)^2\operatorname{PolyLog}(2,-e^{\operatorname{arccosh}(ax)})}{8ac^3} \\ & + \frac{5\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(ax)})}{2ac^3} - \frac{9\operatorname{arccosh}(ax)^2\operatorname{PolyLog}(2,e^{\operatorname{arccosh}(ax)})}{8ac^3} \\ & - \frac{9\operatorname{arccosh}(ax)\operatorname{PolyLog}(3,-e^{\operatorname{arccosh}(ax)})}{4ac^3} \\ & + \frac{9\operatorname{arccosh}(ax)\operatorname{PolyLog}(3,e^{\operatorname{arccosh}(ax)})}{4ac^3} \\ & + \frac{9\operatorname{PolyLog}(4,-e^{\operatorname{arccosh}(ax)})}{4ac^3} - \frac{9\operatorname{PolyLog}(4,e^{\operatorname{arccosh}(ax)})}{4ac^3} \end{aligned}$$

output

$$\begin{aligned}
& -1/4*x*\operatorname{arccosh}(a*x)/c^3/(-a^2*x^2+1)+1/4*\operatorname{arccosh}(a*x)^2/a/c^3/(a*x-1)^{(3/2)} \\
& )/(a*x+1)^{(3/2)}+1/4*x*\operatorname{arccosh}(a*x)^3/c^3/(-a^2*x^2+1)^2+3/8*x*\operatorname{arccosh}(a*x) \\
& ^3/c^3/(-a^2*x^2+1)-5*\operatorname{arccosh}(a*x)*\operatorname{arctanh}(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)} \\
& )/a/c^3+3/4*\operatorname{arccosh}(a*x)^3*\operatorname{arctanh}(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})/a/c^3- \\
& 5/2*\operatorname{polylog}(2,-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})/a/c^3+9/8*\operatorname{arccosh}(a*x)^2*p \\
& \operatorname{olylog}(2,-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})/a/c^3+5/2*\operatorname{polylog}(2,a*x+(a*x-1) \\
& ^{(1/2)}*(a*x+1)^{(1/2)})/a/c^3-9/8*\operatorname{arccosh}(a*x)^2*\operatorname{polylog}(2,a*x+(a*x-1)^{(1/2)} \\
& *(a*x+1)^{(1/2)})/a/c^3-9/4*\operatorname{arccosh}(a*x)*\operatorname{polylog}(3,-a*x-(a*x-1)^{(1/2)}*(a*x+1) \\
& )^{(1/2)})/a/c^3+9/4*\operatorname{arccosh}(a*x)*\operatorname{polylog}(3,a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}) \\
& /a/c^3+9/4*\operatorname{polylog}(4,-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})/a/c^3-9/4*\operatorname{polylog}(4 \\
& ,a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})/a/c^3+1/4/a/c^3/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)} \\
& )-9/8*\operatorname{arccosh}(a*x)^2/a/c^3/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}
\end{aligned}$$

### 3.245.2 Mathematica [A] (warning: unable to verify)

Time = 6.83 (sec) , antiderivative size = 455, normalized size of antiderivative = 1.18

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx =$$

$$3\pi^4 - 6\operatorname{arccosh}(ax)^4 - 8\operatorname{coth}\left(\frac{1}{2}\operatorname{arccosh}(ax)\right) + 40\operatorname{arccosh}(ax)^2\operatorname{coth}\left(\frac{1}{2}\operatorname{arccosh}(ax)\right) - 4\operatorname{arccosh}(ax)\operatorname{csch}\left(\frac{1}{2}\operatorname{arccosh}(ax)\right)$$

input `Integrate[ArcCosh[a*x]^3/(c - a^2*c*x^2)^3,x]`

output

$$\begin{aligned}
& -1/64*(3*\operatorname{Pi}^4 - 6*\operatorname{ArcCosh}[a*x]^4 - 8*\operatorname{Coth}[\operatorname{ArcCosh}[a*x]/2] + 40*\operatorname{ArcCosh}[a*x] \\
& ]^2*\operatorname{Coth}[\operatorname{ArcCosh}[a*x]/2] - 4*\operatorname{ArcCosh}[a*x]*\operatorname{Csch}[\operatorname{ArcCosh}[a*x]/2]^2 + 6*\operatorname{ArcCo} \\
& \operatorname{sh}[a*x]^3*\operatorname{Csch}[\operatorname{ArcCosh}[a*x]/2]^2 - \operatorname{Sqrt}[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*\operatorname{Ar} \\
& \operatorname{cCosh}[a*x]^2*\operatorname{Csch}[\operatorname{ArcCosh}[a*x]/2]^4 - \operatorname{ArcCosh}[a*x]^3*\operatorname{Csch}[\operatorname{ArcCosh}[a*x]/2]^4 \\
& - 160*\operatorname{ArcCosh}[a*x]*\operatorname{Log}[1 - \operatorname{E}^{(-\operatorname{ArcCosh}[a*x])}] + 160*\operatorname{ArcCosh}[a*x]*\operatorname{Log}[1 + \\
& \operatorname{E}^{(-\operatorname{ArcCosh}[a*x])}] - 24*\operatorname{ArcCosh}[a*x]^3*\operatorname{Log}[1 + \operatorname{E}^{(-\operatorname{ArcCosh}[a*x])}] + 24*\operatorname{Ar} \\
& \operatorname{cCosh}[a*x]^3*\operatorname{Log}[1 - \operatorname{E}^{\operatorname{ArcCosh}[a*x]}] + 8*(-20 + 9*\operatorname{ArcCosh}[a*x]^2)*\operatorname{PolyLog}[ \\
& 2, -\operatorname{E}^{(-\operatorname{ArcCosh}[a*x])}] + 160*\operatorname{PolyLog}[2, \operatorname{E}^{(-\operatorname{ArcCosh}[a*x])}] + 72*\operatorname{ArcCosh}[a* \\
& x]^2*\operatorname{PolyLog}[2, \operatorname{E}^{\operatorname{ArcCosh}[a*x]}] + 144*\operatorname{ArcCosh}[a*x]*\operatorname{PolyLog}[3, -\operatorname{E}^{(-\operatorname{ArcCosh} \\
& [a*x])}] - 144*\operatorname{ArcCosh}[a*x]*\operatorname{PolyLog}[3, \operatorname{E}^{\operatorname{ArcCosh}[a*x]}] + 144*\operatorname{PolyLog}[4, -\operatorname{E}^{ \\
& (-\operatorname{ArcCosh}[a*x])}] + 144*\operatorname{PolyLog}[4, \operatorname{E}^{\operatorname{ArcCosh}[a*x]}] - 4*\operatorname{ArcCosh}[a*x]*\operatorname{Sech}[\operatorname{Ar} \\
& \operatorname{cCosh}[a*x]/2]^2 + 6*\operatorname{ArcCosh}[a*x]^3*\operatorname{Sech}[\operatorname{ArcCosh}[a*x]/2]^2 + \operatorname{ArcCosh}[a*x]^3 \\
& *\operatorname{Sech}[\operatorname{ArcCosh}[a*x]/2]^4 - (16*\operatorname{ArcCosh}[a*x]^2*\operatorname{Sinh}[\operatorname{ArcCosh}[a*x]/2]^4)/((( -1 \\
& + a*x)/(1 + a*x))^{(3/2)}*(1 + a*x)^3) + 8*\operatorname{Tanh}[\operatorname{ArcCosh}[a*x]/2] - 40*\operatorname{ArcCos} \\
& \operatorname{h}[a*x]^2*\operatorname{Tanh}[\operatorname{ArcCosh}[a*x]/2)]/(a*c^3)
\end{aligned}$$

$$3.245. \quad \int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx$$

### 3.245.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 4.78 (sec) , antiderivative size = 414, normalized size of antiderivative = 1.07, number of steps used = 23, number of rules used = 22,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.100$ , Rules used = {6316, 27, 6316, 6318, 3042, 26, 4670, 3011, 6330, 25, 6304, 6316, 83, 6318, 3042, 26, 4670, 2715, 2838, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx \\
 & \quad \downarrow \text{6316} \\
 & \frac{3 \int \frac{\operatorname{arccosh}(ax)^3}{c^2(1-a^2x^2)^2} dx}{4c} - \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{3 \int \frac{\operatorname{arccosh}(ax)^3}{(1-a^2x^2)^2} dx}{4c^3} - \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
 & \quad \downarrow \text{6316} \\
 & \frac{3 \left( \frac{1}{2} \int \frac{\operatorname{arccosh}(ax)^3}{1-a^2x^2} dx + \frac{3}{2} a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx + \frac{x \operatorname{arccosh}(ax)^3}{2(1-a^2x^2)} \right)}{4c^3} - \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4c^3} + \\
 & \quad \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
 & \quad \downarrow \text{6318} \\
 & \frac{3 \left( \frac{3}{2} a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d \operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)^3}{2(1-a^2x^2)} \right)}{4c^3} - \\
 & \quad \frac{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& 3 \left( \frac{\frac{3}{2}a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{\int i \operatorname{arccosh}(ax)^3 \csc(i \operatorname{arccosh}(ax)) d \operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)^3}{2(1-a^2x^2)} \right) \\
& \quad \frac{4c^3}{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
& \quad \downarrow 26 \\
& 3 \left( \frac{\frac{3}{2}a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{\int i \operatorname{arccosh}(ax)^3 \csc(i \operatorname{arccosh}(ax)) d \operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)^3}{2(1-a^2x^2)} \right) \\
& \quad \frac{4c^3}{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
& \quad \downarrow 4670 \\
& 3 \left( - \frac{i(3i \int \operatorname{arccosh}(ax)^2 \log(1-e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - 3i \int \operatorname{arccosh}(ax)^2 \log(1+e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) + 2i \operatorname{arccosh}(ax)^3 \operatorname{arccosh}(ax))}{2a} \right) \\
& \quad \frac{4c^3}{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
& \quad \downarrow 3011 \\
& 3 \left( - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{2a} \right) \\
& \quad \frac{4c^3}{3a \int \frac{x \operatorname{arccosh}(ax)^2}{(ax-1)^{5/2}(ax+1)^{5/2}} dx} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
& \quad \downarrow 6330 \\
& 3 \left( \frac{\frac{3}{2}a \left( \frac{2 \int -\frac{\operatorname{arccosh}(ax)}{(1-ax)(ax+1)} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})) + 3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) d \operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{2a}}{4c^3} \right) \\
& \quad \frac{3a \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{(1-ax)^2(ax+1)^2} dx}{3a} - \frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2} \\
& \quad \downarrow 25
\end{aligned}$$

---

3.245.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$

$$3 \left( \frac{3}{2} a \left( -\frac{2 \int \frac{\operatorname{arccosh}(ax)}{(1-ax)(ax+1)} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax))}{4c^3} \right)$$

$$\frac{3a \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{(1-ax)^2(ax+1)^2} dx}{3a} - \frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 6304

$$3 \left( \frac{3}{2} a \left( -\frac{2 \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax))}{4c^3} \right)$$

$$\frac{3a \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{(1-a^2x^2)^2} dx}{3a} - \frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 6316

$$3 \left( \frac{3}{2} a \left( -\frac{2 \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax))}{4c^3} \right)$$

$$\frac{3a \left( \frac{2 \left( \frac{1}{2} \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx + \frac{1}{2} a \int \frac{x}{(ax-1)^{3/2}(ax+1)^{3/2}} dx + \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} \right)}{3a} - \frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 83

$$3 \left( \frac{3}{2} a \left( -\frac{2 \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx}{a} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax))}{4c^3} \right)$$

$$\frac{3a \left( \frac{2 \left( \frac{1}{2} \int \frac{\operatorname{arccosh}(ax)}{1-a^2x^2} dx + \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}} \right)}{3a} - \frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 6318

---

3.245.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$



$$3 \left( \frac{3}{2} a \left( \frac{2 \int \frac{\operatorname{arccosh}(ax)}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)}{a^2} - \frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax))}{a^2} \right)$$

$$\frac{3a \left( \frac{2 \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}} \right)}{3a} - \frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 3042

$$3 \left( \frac{3}{2} a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2 \int i \operatorname{arccosh}(ax) \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{a^2} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax))}{a^2} \right)$$

$$\frac{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} + \frac{2 \left( -\frac{\int i \operatorname{arccosh}(ax) \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}} \right)}{3a} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 26

$$3 \left( \frac{3}{2} a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i \int \operatorname{arccosh}(ax) \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{a^2} \right) - \frac{i(-3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax))}{a^2} \right)$$

$$\frac{3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} + \frac{2 \left( -\frac{\int i \operatorname{arccosh}(ax) \csc(i \operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2a} + \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{1}{2a\sqrt{ax-1}\sqrt{ax+1}} \right)}{3a} \right)}{4c^3} + \frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 4670

---

3.245.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$

$$3 \left( \frac{3}{2} a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i \left( i \int \log(1-e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - i \int \log(1+e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2i \operatorname{arccosh}(ax) \arctan \right)}{a^2} \right) \right)$$

$$3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2 (ax-1)^{3/2} (ax+1)^{3/2}} + \frac{2 \left( -\frac{i \left( i \int \log(1-e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - i \int \log(1+e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 2i \operatorname{arccosh}(ax) \arctan \right)}{2a} \right)}{3a} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{4c^3 (1-a^2x^2)^2}$$

↓ 2715

$$3 \left( \frac{3}{2} a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i \left( i \int e^{-\operatorname{arccosh}(ax)} \log(1-e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - i \int e^{-\operatorname{arccosh}(ax)} \log(1+e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} \right)}{a^2} \right) \right)$$

$$3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2 (ax-1)^{3/2} (ax+1)^{3/2}} + \frac{2 \left( -\frac{i \left( i \int e^{-\operatorname{arccosh}(ax)} \log(1-e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} - i \int e^{-\operatorname{arccosh}(ax)} \log(1+e^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)} \right)}{2a} \right)}{3a} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{4c^3 (1-a^2x^2)^2}$$

↓ 2838

$$3 \left( -\frac{i \left( -3i \left( 2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) \right) \right) + 3i \left( 2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) \right) \right)}{2a} \right)$$

$$3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2 (ax-1)^{3/2} (ax+1)^{3/2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{i \left( 2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}) \right)}{2a} \right)}{3a} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{4c^3 (1-a^2x^2)^2}$$

↓ 7163

---

3.245.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$

$$3 \left( -\frac{i(-3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - f \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) \operatorname{arccosh}(ax)) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})$$

$$3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{2a} \right)}{3a} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 2720

$$3 \left( -\frac{i(-3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) - f e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(3, -e^{\operatorname{arccosh}(ax)}) d e^{\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)})$$

$$3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{2a} \right)}{3a} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

↓ 7143

$$3a \left( -\frac{\operatorname{arccosh}(ax)^2}{3a^2(ax-1)^{3/2}(ax+1)^{3/2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)}{2(1-a^2x^2)} - \frac{i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)})}{2a} \right)}{3a} \right)$$

$$3 \left( \frac{3}{2} a \left( -\frac{\operatorname{arccosh}(ax)^2}{a^2 \sqrt{ax-1} \sqrt{ax+1}} + \frac{2i(2i \operatorname{arccosh}(ax) \operatorname{arctanh}(e^{\operatorname{arccosh}(ax)}) + i \operatorname{PolyLog}(2, -e^{\operatorname{arccosh}(ax)}) - i \operatorname{PolyLog}(2, e^{\operatorname{arccosh}(ax)}))}{a^2} \right) \right) +$$

$$\frac{x \operatorname{arccosh}(ax)^3}{4c^3(1-a^2x^2)^2}$$

input `Int[ArcCosh[a*x]^3/(c - a^2*c*x^2)^3,x]`

3.245.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$

```
output (x*ArcCosh[a*x]^3)/(4*c^3*(1 - a^2*x^2)^2) - (3*a*(-1/3*ArcCosh[a*x]^2/(a^
2*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2)) + (2*(-1/2*1/(a*Sqrt[-1 + a*x]*Sqrt[1
+ a*x]) + (x*ArcCosh[a*x]))/(2*(1 - a^2*x^2)) - ((I/2)*((2*I)*ArcCosh[a*x]*
ArcTanh[E^ArcCosh[a*x]] + I*PolyLog[2, -E^ArcCosh[a*x]] - I*PolyLog[2, E^A
rcCosh[a*x]]))/a)/(3*a)))/(4*c^3) + (3*((x*ArcCosh[a*x]^3)/(2*(1 - a^2*x^
2)) + (3*a*(-(ArcCosh[a*x]^2/(a^2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])) + ((2*I)*
((2*I)*ArcCosh[a*x]*ArcTanh[E^ArcCosh[a*x]] + I*PolyLog[2, -E^ArcCosh[a*x]
] - I*PolyLog[2, E^ArcCosh[a*x]]))/a^2))/2 - ((I/2)*((2*I)*ArcCosh[a*x]^3*
ArcTanh[E^ArcCosh[a*x]] - (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, -E^ArcCosh[a*
x]]) + 2*(ArcCosh[a*x]*PolyLog[3, -E^ArcCosh[a*x]] - PolyLog[4, -E^ArcCosh
[a*x]])) + (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, E^ArcCosh[a*x]]) + 2*(ArcCos
h[a*x]*PolyLog[3, E^ArcCosh[a*x]] - PolyLog[4, E^ArcCosh[a*x]]))))/a)/(4*
c^3)
```

### 3.245.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 83 Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))),
x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f
*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]
```

```
rule 2715 Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:= Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x)) *(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2838 `Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_) *(x_)^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4670 `Int[csc[(e_) + (Complex[0, fz_])*(f_)*(x_)]*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

rule 6304 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(q_), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6316 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6318 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[-(c*d)^(-1) Subst[Int[(a + b*x)^n*Csch[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`

### 3.245.4 Maple [A] (verified)

Time = 0.55 (sec) , antiderivative size = 527, normalized size of antiderivative = 1.36

method	result
derivativedivides	$\frac{-3a^3x^3 \operatorname{arccosh}(ax)^3 + 9a^2x^2 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} - 5ax \operatorname{arccosh}(ax)^3 - 11 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} - 2a^3x^3 \operatorname{arccosh}(ax) - 2}{8(a^4x^4 - 2a^2x^2 + 1)c^3}$
default	$\frac{-3a^3x^3 \operatorname{arccosh}(ax)^3 + 9a^2x^2 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} - 5ax \operatorname{arccosh}(ax)^3 - 11 \operatorname{arccosh}(ax)^2 \sqrt{ax-1} \sqrt{ax+1} - 2a^3x^3 \operatorname{arccosh}(ax) - 2}{8(a^4x^4 - 2a^2x^2 + 1)c^3}$

input `int(arccosh(a*x)^3/(-a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)`

$$3.245. \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^3} dx$$

```
output 1/a*(-1/8*(3*a^3*x^3*arccosh(a*x)^3+9*a^2*x^2*arccosh(a*x)^2*(a*x-1)^(1/2)
*(a*x+1)^(1/2)-5*a*x*arccosh(a*x)^3-11*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)
)^(1/2)-2*a^3*x^3*arccosh(a*x)-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+2*a*x
*arccosh(a*x)+2*(a*x-1)^(1/2)*(a*x+1)^(1/2))/(a^4*x^4-2*a^2*x^2+1)/c^3-5/2
/c^3*arccosh(a*x)*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))-5/2/c^3*polylog(2,
-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+5/2/c^3*arccosh(a*x)*ln(1-a*x-(a*x-1)^(1
/2)*(a*x+1)^(1/2))+5/2/c^3*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+3/8/
c^3*arccosh(a*x)^3*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+9/8/c^3*arccosh(a
*x)^2*polylog(2,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))-9/4/c^3*arccosh(a*x)*pol
ylog(3,-a*x-(a*x-1)^(1/2)*(a*x+1)^(1/2))+9/4/c^3*polylog(4,-a*x-(a*x-1)^(1
/2)*(a*x+1)^(1/2))-3/8/c^3*arccosh(a*x)^3*ln(1-a*x-(a*x-1)^(1/2)*(a*x+1)^(
1/2))-9/8/c^3*arccosh(a*x)^2*polylog(2,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+9/
4/c^3*arccosh(a*x)*polylog(3,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))-9/4/c^3*pol
ylog(4,a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))
```

### 3.245.5 Fricas [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arccosh}(ax)^3}{(a^2cx^2 - c)^3} dx$$

```
input integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^3,x, algorithm="fricas")
```

```
output integral(-arccosh(a*x)^3/(a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^
3), x)
```

### 3.245.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx = -\int \frac{\operatorname{acosh}^3(ax)}{a^6x^6 - 3a^4x^4 + 3a^2x^2 - 1} \frac{dx}{c^3}$$

```
input integrate(acosh(a*x)**3/(-a**2*c*x**2+c)**3,x)
```

```
output -Integral(acosh(a*x)**3/(a**6*x**6 - 3*a**4*x**4 + 3*a**2*x**2 - 1), x)/c*
*3
```

---

3.245.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx$

**3.245.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arcosh}(ax)^3}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^3,x, algorithm="maxima")`

output `-1/16*(6*a^3*x^3 - 10*a*x - 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x + 1) + 3*(a^4*x^4 - 2*a^2*x^2 + 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1))*sqrt(a*x - 1))^3/(a^5*c^3*x^4 - 2*a^3*c^3*x^2 + a*c^3) - integrate(-3/16*(6*a^5*x^5 - 16*a^3*x^3 + (6*a^4*x^4 - 10*a^2*x^2 - 3*(a^5*x^5 - 2*a^3*x^3 + a*x)*log(a*x + 1) + 3*(a^5*x^5 - 2*a^3*x^3 + a*x)*log(a*x - 1))*sqrt(a*x + 1)*sqrt(a*x - 1) + 10*a*x - 3*(a^6*x^6 - 3*a^4*x^4 + 3*a^2*x^2 - 1)*log(a*x + 1) + 3*(a^6*x^6 - 3*a^4*x^4 + 3*a^2*x^2 - 1)*log(a*x - 1))*log(a*x + sqrt(a*x + 1))*sqrt(a*x - 1))^2/(a^7*c^3*x^7 - 3*a^5*c^3*x^5 + 3*a^3*c^3*x^3 - a*c^3*x + (a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3)*sqrt(a*x + 1)*sqrt(a*x - 1)), x)`

**3.245.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx = \int -\frac{\operatorname{arcosh}(ax)^3}{(a^2cx^2 - c)^3} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^3,x, algorithm="giac")`

output `integrate(-arccosh(a*x)^3/(a^2*c*x^2 - c)^3, x)`

**3.245.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx = \int \frac{\operatorname{acosh}(ax)^3}{(c - a^2cx^2)^3} dx$$

input `int(acosh(a*x)^3/(c - a^2*c*x^2)^3,x)`

output `int(acosh(a*x)^3/(c - a^2*c*x^2)^3, x)`

---

3.245.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^3} dx$



### 3.246 $\int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

3.246.1 Optimal result	2192
3.246.2 Mathematica [A] (warning: unable to verify)	2193
3.246.3 Rubi [A] (verified)	2194
3.246.4 Maple [A] (verified)	2205
3.246.5 Fricas [F]	2206
3.246.6 Sympy [F(-1)]	2207
3.246.7 Maxima [F(-2)]	2207
3.246.8 Giac [F(-2)]	2207
3.246.9 Mupad [F(-1)]	2208

#### 3.246.1 Optimal result

Integrand size = 22, antiderivative size = 605

$$\begin{aligned}
 \int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx = & -\frac{865ac^2x^2\sqrt{c - a^2cx^2}}{2304\sqrt{-1 + ax}\sqrt{1 + ax}} \\
 & + \frac{65a^3c^2x^4\sqrt{c - a^2cx^2}}{2304\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{c^2(1 - a^2x^2)^3\sqrt{c - a^2cx^2}}{216a\sqrt{-1 + ax}\sqrt{1 + ax}} \\
 & + \frac{245}{384}c^2x\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax) + \frac{65}{576}c^2x(1 - ax)(1 + ax)\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax) \\
 & + \frac{1}{36}c^2x(1 - ax)^2(1 + ax)^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax) + \frac{115c^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^2}{768a\sqrt{-1 + ax}\sqrt{1 + ax}} \\
 & - \frac{15ac^2x^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^2}{32\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{5c^2(1 - a^2x^2)^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^2}{32a\sqrt{-1 + ax}\sqrt{1 + ax}} \\
 & + \frac{c^2(1 - a^2x^2)^3\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^2}{12a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{5}{16}c^2x\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^3 + \frac{5}{24}cx(c \\
 & - a^2cx^2)^{3/2}\operatorname{arccosh}(ax)^3 + \frac{1}{6}x(c - a^2cx^2)^{5/2}\operatorname{arccosh}(ax)^3 - \frac{5c^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^4}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}
 \end{aligned}$$

output  $5/24*c*x*(-a^2*c*x^2+c)^{(3/2)}*\operatorname{arccosh}(a*x)^3+1/6*x*(-a^2*c*x^2+c)^{(5/2)}*\operatorname{arccosh}(a*x)^3+245/384*c^2*x*\operatorname{arccosh}(a*x)*(-a^2*c*x^2+c)^{(1/2)}+65/576*c^2*x*(-a*x+1)*(a*x+1)*\operatorname{arccosh}(a*x)*(-a^2*c*x^2+c)^{(1/2)}+1/36*c^2*x*(-a*x+1)^2*(a*x+1)^2*\operatorname{arccosh}(a*x)*(-a^2*c*x^2+c)^{(1/2)}+5/16*c^2*x*\operatorname{arccosh}(a*x)^3*(-a^2*c*x^2+c)^{(1/2)}-865/2304*a*c^2*x^2*(-a^2*c*x^2+c)^{(1/2)}/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}+65/2304*a^3*c^2*x^4*(-a^2*c*x^2+c)^{(1/2)}/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}+1/216*c^2*(-a^2*x^2+1)^3*(-a^2*c*x^2+c)^{(1/2)}/a/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}+115/768*c^2*\operatorname{arccosh}(a*x)^2*(-a^2*c*x^2+c)^{(1/2)}/a/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}-15/32*a*c^2*x^2*\operatorname{arccosh}(a*x)^2*(-a^2*c*x^2+c)^{(1/2)}/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}+5/32*c^2*(-a^2*x^2+1)^2*\operatorname{arccosh}(a*x)^2*(-a^2*c*x^2+c)^{(1/2)}/a/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}+1/12*c^2*(-a^2*x^2+1)^3*\operatorname{arccosh}(a*x)^2*(-a^2*c*x^2+c)^{(1/2)}/a/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}-5/64*c^2*\operatorname{arccosh}(a*x)^4*(-a^2*c*x^2+c)^{(1/2)}/a/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}$

### 3.246.2 Mathematica [A] (warning: unable to verify)

Time = 0.92 (sec) , antiderivative size = 189, normalized size of antiderivative = 0.31

$$\int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx = \frac{c^2\sqrt{c - a^2cx^2}(-4320\operatorname{arccosh}(ax)^4 - 9720\cosh(2\operatorname{arccosh}(ax)) + 243\cosh(4\operatorname{arccosh}(ax)))}{(55296a\sqrt{(-1 + ax)/(1 + ax)})(1 + ax)}$$

input `Integrate[(c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^3,x]`

output  $(c^2*\operatorname{Sqrt}[c - a^2*c*x^2]*(-4320*\operatorname{ArcCosh}[a*x]^4 - 9720*\operatorname{Cosh}[2*\operatorname{ArcCosh}[a*x]] + 243*\operatorname{Cosh}[4*\operatorname{ArcCosh}[a*x]] - 8*\operatorname{Cosh}[6*\operatorname{ArcCosh}[a*x]] - 72*\operatorname{ArcCosh}[a*x]^2*(270*\operatorname{Cosh}[2*\operatorname{ArcCosh}[a*x]] - 27*\operatorname{Cosh}[4*\operatorname{ArcCosh}[a*x]] + 2*\operatorname{Cosh}[6*\operatorname{ArcCosh}[a*x]]) + 288*\operatorname{ArcCosh}[a*x]^3*(45*\operatorname{Sinh}[2*\operatorname{ArcCosh}[a*x]] - 9*\operatorname{Sinh}[4*\operatorname{ArcCosh}[a*x]] + \operatorname{Sinh}[6*\operatorname{ArcCosh}[a*x]]) + 12*\operatorname{ArcCosh}[a*x]*(1620*\operatorname{Sinh}[2*\operatorname{ArcCosh}[a*x]] - 81*\operatorname{Sinh}[4*\operatorname{ArcCosh}[a*x]] + 4*\operatorname{Sinh}[6*\operatorname{ArcCosh}[a*x]])))/(55296*a*\operatorname{Sqrt}[(-1 + a*x)/(1 + a*x)]*(1 + a*x))$

**3.246.3 Rubi [A] (verified)**

Time = 6.94 (sec) , antiderivative size = 651, normalized size of antiderivative = 1.08, number of steps used = 28, number of rules used = 28,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.273$ , Rules used = {6312, 6312, 25, 6310, 6298, 6308, 6327, 6329, 6313, 25, 82, 241, 244, 2009, 6311, 15, 6308, 6313, 25, 82, 244, 2009, 6311, 15, 6308, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{arccosh}(ax)^3 (c - a^2cx^2)^{5/2} dx \\
 & \quad \downarrow \text{6312} \\
 & -\frac{ac^2\sqrt{c-a^2cx^2} \int x(1-ax)^2(ax+1)^2\operatorname{arccosh}(ax)^2 dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{5}{6}c \int (c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx + \\
 & \quad \frac{1}{6}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \\
 & \quad \downarrow \text{6312} \\
 & -\frac{ac^2\sqrt{c-a^2cx^2} \int x(1-ax)^2(ax+1)^2\operatorname{arccosh}(ax)^2 dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
 & \frac{5}{6}c \left( \frac{3ac\sqrt{c-a^2cx^2} \int -x(1-ax)(ax+1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \int \sqrt{c-a^2cx^2} \operatorname{arccosh}(ax)^3 dx + \frac{1}{4}x\operatorname{arccosh}(ax)^3 \right. \\
 & \quad \left. \frac{1}{6}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \\
 & \quad \downarrow \text{25} \\
 & -\frac{ac^2\sqrt{c-a^2cx^2} \int x(1-ax)^2(ax+1)^2\operatorname{arccosh}(ax)^2 dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
 & \frac{5}{6}c \left( -\frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \int \sqrt{c-a^2cx^2} \operatorname{arccosh}(ax)^3 dx + \frac{1}{4}x\operatorname{arccosh}(ax)^3 \right. \\
 & \quad \left. \frac{1}{6}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \\
 & \quad \downarrow \text{6310} \\
 & -\frac{ac^2\sqrt{c-a^2cx^2} \int x(1-ax)^2(ax+1)^2\operatorname{arccosh}(ax)^2 dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
 & \frac{5}{6}c \left( -\frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \int x\operatorname{arccosh}(ax)^2 dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2}}{2\sqrt{ax-1}\sqrt{ax+1}} \right) \right. \\
 & \quad \left. \frac{1}{6}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \\
 & \quad \downarrow \text{6298}
 \end{aligned}$$

---

3.246.  $\int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

$$\begin{aligned}
& -\frac{ac^2\sqrt{c-a^2cx^2}\int x(1-ax)^2(ax+1)^2\operatorname{arccosh}(ax)^2dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{5}{6}c & \left( -\frac{3ac\sqrt{c-a^2cx^2}\int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^2dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a\int\frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}}dx\right)}{2\sqrt{ax-1}\sqrt{ax+1}} \right. \right. \\
& \left. \left. + \frac{1}{6}x\operatorname{arccosh}(ax)^3(c-a^2cx^2)^{5/2} \right) \right. \\
& \quad \downarrow \text{6308} \\
& -\frac{ac^2\sqrt{c-a^2cx^2}\int x(1-ax)^2(ax+1)^2\operatorname{arccosh}(ax)^2dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{5}{6}c & \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a\int\frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right. \right. \\
& \left. \left. + \frac{1}{6}x\operatorname{arccosh}(ax)^3(c-a^2cx^2)^{5/2} \right) \right. \\
& \quad \downarrow \text{6327} \\
& -\frac{ac^2\sqrt{c-a^2cx^2}\int x(1-a^2x^2)^2\operatorname{arccosh}(ax)^2dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{5}{6}c & \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a\int\frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right. \right. \\
& \left. \left. + \frac{1}{6}x\operatorname{arccosh}(ax)^3(c-a^2cx^2)^{5/2} \right) \right. \\
& \quad \downarrow \text{6329} \\
& -\frac{ac^2\sqrt{c-a^2cx^2}\left(-\frac{\int(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)dx}{3a} - \frac{(1-a^2x^2)^3\operatorname{arccosh}(ax)^2}{6a^2}\right)}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{5}{6}c & \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a\int\frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right. \right. \\
& \left. \left. + \frac{1}{6}x\operatorname{arccosh}(ax)^3(c-a^2cx^2)^{5/2} \right) \right. \\
& \quad \downarrow \text{6313}
\end{aligned}$$

$$\begin{aligned}
& \frac{ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6} \int (ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) dx - \frac{1}{6} a \int x(1-ax)^2(ax+1)^2 dx + \frac{1}{6} x(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)}{3a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{\frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2}} \right)}{\downarrow 25}
\end{aligned}$$

$$\begin{aligned}
& \frac{ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6} \int (ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) dx - \frac{1}{6} a \int x(1-ax)^2(ax+1)^2 dx + \frac{1}{6} x(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)}{3a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{\frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2}} \right)}{\downarrow 82}
\end{aligned}$$

$$\begin{aligned}
& \frac{ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{1}{6} a \int x(1-a^2x^2)^2 dx - \frac{5}{6} \int (ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) dx + \frac{1}{6} x(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)}{3a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{\frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2}} \right)}{\downarrow 241}
\end{aligned}$$

$$\begin{aligned}
& \frac{ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6} \int (ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) dx + \frac{(1-a^2x^2)^3}{36a} + \frac{1}{6} x(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)}{3a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{\frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2}} \right)}{6a^2}
\end{aligned}$$

---

3.246.  $\int (c-a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

↓ 244

$$ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6}\int(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)dx + \frac{(1-a^2x^2)^3}{36a} + \frac{1}{6}x(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)}{3a} - \frac{(1-a^2x^2)^3\operatorname{arccosh}(ax)}{6a^2} \right)$$


---


$$\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a \int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \right.$$

$$\left. \frac{1}{6}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right)$$

↓ 2009

$$ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6}\int(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)dx + \frac{(1-a^2x^2)^3}{36a} + \frac{1}{6}x(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)}{3a} - \frac{(1-a^2x^2)^3\operatorname{arccosh}(ax)}{6a^2} \right)$$


---


$$\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a \int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \right.$$

$$\left. \frac{1}{6}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right)$$

↓ 6311

$$ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6}\int(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)dx + \frac{(1-a^2x^2)^3}{36a} + \frac{1}{6}x(ax-1)^{5/2}(ax+1)^{5/2}\operatorname{arccosh}(ax)}{3a} - \frac{(1-a^2x^2)^3\operatorname{arccosh}(ax)}{6a^2} \right)$$


---


$$\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a \int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \right.$$

$$\left. \frac{1}{6}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right)$$

↓ 15

---

3.246.  $\int (c-a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

$$\begin{aligned}
& \frac{ac^2\sqrt{c-a^2cx^2} \left( -\frac{5}{6} \int (ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) dx + \frac{(1-a^2x^2)^3}{36a} + \frac{1}{6} x(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax) - \frac{(1-a^2x^2)^3 \operatorname{arccosh}(ax)}{6a^2} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{5}{6} c \left( \frac{3}{4} c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2} x \operatorname{arccosh}(ax) \right) \right. \\
& \quad \left. + \frac{1}{6} x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \downarrow \text{6308} \\
& \frac{ac^2\sqrt{c-a^2cx^2} \left( -\frac{5}{6} \int (ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) dx + \frac{(1-a^2x^2)^3}{36a} + \frac{1}{6} x(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax) - \frac{(1-a^2x^2)^3 \operatorname{arccosh}(ax)}{6a^2} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{5}{6} c \left( \frac{3}{4} c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2} x \operatorname{arccosh}(ax) \right) \right. \\
& \quad \left. + \frac{1}{6} x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \downarrow \text{6313} \\
& \frac{ac^2\sqrt{c-a^2cx^2} \left( -\frac{5}{6} \left( -\frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx - \frac{1}{4} a \int -x(1-ax)(ax+1) dx + \frac{1}{4} x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right) + \frac{(1-a^2x^2)^3 \operatorname{arccosh}(ax)}{36a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{5}{6} c \left( \frac{3}{4} c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2} x \operatorname{arccosh}(ax) \right) \right. \\
& \quad \left. + \frac{1}{6} x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \downarrow \text{25}
\end{aligned}$$

---

3.246.  $\int (c-a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

$$\begin{aligned}
& ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6} \left( -\frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx + \frac{1}{4} a \int x(1-ax)(ax+1) dx + \frac{1}{4} x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right) + \frac{(1-a^2x)^3}{36a}}{3a} \right) \\
& \hline
& \frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \right. \\
& \qquad \qquad \qquad \left. \frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \\
& \qquad \qquad \qquad \downarrow \text{82} \\
& ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6} \left( \frac{1}{4} a \int x(1-a^2x^2) dx - \frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx + \frac{1}{4} x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right) + \frac{(1-a^2x^2)^3}{36a}}{3a} \right) \\
& \hline
& \frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \right. \\
& \qquad \qquad \qquad \left. \frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \\
& \qquad \qquad \qquad \downarrow \text{244} \\
& ac^2\sqrt{c-a^2cx^2} \left( -\frac{\frac{5}{6} \left( \frac{1}{4} a \int (x-a^2x^3) dx - \frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx + \frac{1}{4} x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right) + \frac{(1-a^2x^2)^3}{36a}}{3a} \right) \\
& \hline
& \frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \right. \\
& \qquad \qquad \qquad \left. \frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right) \\
& \qquad \qquad \qquad \downarrow \text{2009}
\end{aligned}$$



$$\begin{aligned}
 & \frac{ac^2\sqrt{c-a^2cx^2}}{2\sqrt{ax-1}\sqrt{ax+1}} \left( -\frac{\frac{5}{6}\left(-\frac{3}{4}\int\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)dx+\frac{1}{4}a\left(\frac{x^2}{2}-\frac{a^2x^4}{4}\right)+\frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)\right)+\frac{(1-a^2x^2)^3}{36a}+\frac{1}{6}}{3a} \right) \\
 & \frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax)^2-a\int\frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \right. \\
 & \qquad \qquad \qquad \left. \frac{1}{6}x\operatorname{arccosh}(ax)^3(c-a^2cx^2)^{5/2} \right) \downarrow \text{6311} \\
 & \frac{ac^2\sqrt{c-a^2cx^2}}{2\sqrt{ax-1}\sqrt{ax+1}} \left( -\frac{\frac{5}{6}\left(-\frac{3}{4}\left(-\frac{1}{2}\int\frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx-\frac{a}{2}\int\frac{xdx}{\sqrt{ax-1}\sqrt{ax+1}}+\frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)\right)+\frac{1}{4}a\left(\frac{x^2}{2}-\frac{a^2x^4}{4}\right)+\frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2}}{3a} \right) \right) \\
 & \frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax)^2-a\int\frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \right. \\
 & \qquad \qquad \qquad \left. \frac{1}{6}x\operatorname{arccosh}(ax)^3(c-a^2cx^2)^{5/2} \right) \downarrow \text{15} \\
 & \frac{ac^2\sqrt{c-a^2cx^2}}{2\sqrt{ax-1}\sqrt{ax+1}} \left( -\frac{\frac{5}{6}\left(-\frac{3}{4}\left(-\frac{1}{2}\int\frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx+\frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)-\frac{ax^2}{4}\right)+\frac{1}{4}a\left(\frac{x^2}{2}-\frac{a^2x^4}{4}\right)+\frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2}}{3a} \right) \right) \\
 & \frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2}\left(\frac{1}{2}x^2\operatorname{arccosh}(ax)^2-a\int\frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \right. \\
 & \qquad \qquad \qquad \left. \frac{1}{6}x\operatorname{arccosh}(ax)^3(c-a^2cx^2)^{5/2} \right) \downarrow \text{6308}
 \end{aligned}$$

3.246.  $\int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

$$\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \right. \\ \left. - \frac{(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2}{6a^2} - \frac{-\frac{5}{6} \left( \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)}{3a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \right. \\ \left. - \frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right)$$

↓ 6354

$$\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} - \frac{\int x dx}{2a} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \right. \\ \left. - \frac{(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2}{6a^2} - \frac{-\frac{5}{6} \left( \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)}{3a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \right. \\ \left. - \frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right)$$

↓ 15

$$\frac{5}{6}c \left( \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} - \frac{x^2}{4a} \right) \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \right. \\ \left. - \frac{(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2}{6a^2} - \frac{-\frac{5}{6} \left( \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)}{3a} \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \right. \\ \left. - \frac{1}{6}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{5/2} \right)$$

↓ 6308

---

3.246.  $\int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

$$ac^2 \left( -\frac{(1-a^2x^2)^3 \operatorname{arccosh}(ax)^2}{6a^2} - \frac{-\frac{5}{6} \left( \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax-1) \right)}{3a} \right)$$


---


$$\frac{1}{6} x \operatorname{arccosh}(ax)^3 (c - a^2cx^2)^{5/2} + \frac{3ac \left( \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax-1) \right)}{2a}$$


---


$$\frac{5}{6}c \left( \frac{1}{4} x \operatorname{arccosh}(ax)^3 (c - a^2cx^2)^{3/2} - \frac{\dots}{4\sqrt{ax-1}\sqrt{ax+1}} \right)$$

```
input Int[(c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^3,x]
```

```
output (x*(c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^3)/6 - (a*c^2*Sqrt[c - a^2*c*x^2]*(-1/6*((1 - a^2*x^2)^3*ArcCosh[a*x]^2)/a^2 - ((1 - a^2*x^2)^3/(36*a) + (x*(-1 + a*x)^(5/2)*(1 + a*x)^(5/2)*ArcCosh[a*x])/6 - (5*((a*(x^2/2 - (a^2*x^4)/4))/4 + (x*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x])/4 - (3*(-1/4*(a*x^2) + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/2 - ArcCosh[a*x]^2/(4*a)))/4)/6)/(3*a)))/(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + (5*c*((x*(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^3)/4 + (3*c*((x*Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^3)/2 - (Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^4)/(8*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) - (3*a*Sqrt[c - a^2*c*x^2]*((x^2*ArcCosh[a*x]^2)/2 - a*(-1/4*x^2/a + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]))/(2*a^2) + ArcCosh[a*x]^2/(4*a^3)))))/(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])))/4 - (3*a*c*Sqrt[c - a^2*c*x^2]*(-1/4*((1 - a^2*x^2)^2*ArcCosh[a*x]^2)/a^2 + ((a*(x^2/2 - (a^2*x^4)/4))/4 + (x*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x])/4 - (3*(-1/4*(a*x^2) + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/2 - ArcCosh[a*x]^2/(4*a))/4)/(2*a)))/(4*Sqrt[-1 + a*x]*Sqrt[1 + a*x])))/6
```

3.246.3.1 Defintions of rubi rules used

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

---

3.246.  $\int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

- rule 82 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`
- rule 241 `Int[(x_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`
- rule 244 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6298 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_)*(x_))^(m_), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6308 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(Sqrt[(d1_) + (e1_)*(x_)]*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^(n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6311  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)^{(n_.)}*\text{Sqrt}[(d1_.) + (e1_.)*(x_)]*\text{Sqrt}[(d2_.) + (e2_.)*(x_)], x\_Symbol] \rightarrow \text{Simp}[x*\text{Sqrt}[d1 + e1*x]*\text{Sqrt}[d2 + e2*x]*((a + b*\text{ArcCosh}[c*x])^{n/2}), x] + (-\text{Simp}[(1/2)*\text{Simp}[\text{Sqrt}[d1 + e1*x]/\text{Sqrt}[1 + c*x]]*\text{Simp}[\text{Sqrt}[d2 + e2*x]/\text{Sqrt}[-1 + c*x]] \text{Int}[(a + b*\text{ArcCosh}[c*x])^{n/2}/(\text{Sqrt}[1 + c*x]*\text{Sqrt}[-1 + c*x]), x], x] - \text{Simp}[b*c*(n/2)*\text{Simp}[\text{Sqrt}[d1 + e1*x]/\text{Sqrt}[1 + c*x]]*\text{Simp}[\text{Sqrt}[d2 + e2*x]/\text{Sqrt}[-1 + c*x]] \text{Int}[x*(a + b*\text{ArcCosh}[c*x])^{(n - 1)}, x], x) /; \text{FreeQ}\{a, b, c, d1, e1, d2, e2\}, x\} \&\& \text{EqQ}\{e1, c*d1\} \&\& \text{EqQ}\{e2, (-c)*d2\} \&\& \text{GtQ}\{n, 0\}$

rule 6312  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)^{(n_.)}*((d_.) + (e_.)*(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[x*(d + e*x^2)^p*((a + b*\text{ArcCosh}[c*x])^{n/(2*p + 1)}), x] + (\text{Simp}[2*d*(p/(2*p + 1)) \text{Int}[(d + e*x^2)^{(p - 1)}*(a + b*\text{ArcCosh}[c*x])^n, x], x] - \text{Simp}[b*c*(n/(2*p + 1))*\text{Simp}[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] \text{Int}[x*(1 + c*x)^{(p - 1/2)}*(-1 + c*x)^{(p - 1/2)}*(a + b*\text{ArcCosh}[c*x])^{(n - 1)}, x], x) /; \text{FreeQ}\{a, b, c, d, e\}, x\} \&\& \text{EqQ}\{c^2*d + e, 0\} \&\& \text{GtQ}\{n, 0\} \&\& \text{GtQ}\{p, 0\}$

rule 6313  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)^{(n_.)}*((d1_.) + (e1_.)*(x_))^{(p_.)}*(d2_.) + (e2_.)*(x_))^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[x*(d1 + e1*x)^p*(d2 + e2*x)^p*((a + b*\text{ArcCosh}[c*x])^{n/(2*p + 1)}), x] + (\text{Simp}[2*d1*d2*(p/(2*p + 1)) \text{Int}[(d1 + e1*x)^{(p - 1)}*(d2 + e2*x)^{(p - 1)}*(a + b*\text{ArcCosh}[c*x])^n, x], x] - \text{Simp}[b*c*(n/(2*p + 1))*\text{Simp}[(d1 + e1*x)^p/(1 + c*x)^p]*\text{Simp}[(d2 + e2*x)^p/(-1 + c*x)^p] \text{Int}[x*(1 + c*x)^{(p - 1/2)}*(-1 + c*x)^{(p - 1/2)}*(a + b*\text{ArcCosh}[c*x])^{(n - 1)}, x], x) /; \text{FreeQ}\{a, b, c, d1, e1, d2, e2\}, x\} \&\& \text{EqQ}\{e1, c*d1\} \&\& \text{EqQ}\{e2, (-c)*d2\} \&\& \text{GtQ}\{n, 0\} \&\& \text{GtQ}\{p, 0\}$

rule 6327  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)^{(n_.)}*((f_.)*(x_))^{(m_.)}*((d1_.) + (e1_.)*(x_))^{(p_.)}*(d2_.) + (e2_.)*(x_))^{(p_.)}, x\_Symbol] \rightarrow \text{Int}[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*\text{ArcCosh}[c*x])^n, x] /; \text{FreeQ}\{a, b, c, d1, e1, d2, e2, f, m, n\}, x\} \&\& \text{EqQ}\{d2*e1 + d1*e2, 0\} \&\& \text{IntegerQ}\{p\}$

rule 6329  $\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_)]*(b_.)^{(n_.)}*(x_)*((d_.) + (e_.)*(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[(d + e*x^2)^{(p + 1)}*((a + b*\text{ArcCosh}[c*x])^{n/(2*e*(p + 1))}), x] - \text{Simp}[b*(n/(2*c*(p + 1)))*\text{Simp}[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] \text{Int}[(1 + c*x)^{(p + 1/2)}*(-1 + c*x)^{(p + 1/2)}*(a + b*\text{ArcCosh}[c*x])^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x\} \&\& \text{EqQ}\{c^2*d + e, 0\} \&\& \text{GtQ}\{n, 0\} \&\& \text{NeQ}\{p, -1\}$

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.246.4 Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 887, normalized size of antiderivative = 1.47

method	result
default	$-\frac{5\sqrt{-c(a^2x^2-1)} \operatorname{arccosh}(ax)^4 c^2}{64\sqrt{ax-1}\sqrt{ax+1}a} + \frac{\sqrt{-c(a^2x^2-1)} (32a^7x^7-64a^5x^5+32a^6a^6\sqrt{ax-1}\sqrt{ax+1}+38a^3x^3-48\sqrt{ax-1}\sqrt{ax+1}a^4x^4-1382}{1382}$

```
input int((-a^2*c*x^2+c)^(5/2)*arccosh(a*x)^3,x,method=_RETURNVERBOSE)
```

---

3.246.  $\int (c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx$

output `-5/64*(-c*(a^2*x^2-1))^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)/a*arccosh(a*x)^4*c^2+1/13824*(-c*(a^2*x^2-1))^(1/2)*(32*a^7*x^7-64*a^5*x^5+32*x^6*a^6*(a*x-1)^(1/2)*(a*x+1)^(1/2)+38*a^3*x^3-48*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4-6*a*x+18*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))*(36*arccosh(a*x)^3-18*arccosh(a*x)^2+6*arccosh(a*x)-1)*c^2/(a*x-1)/(a*x+1)/a-3/4096*(-c*(a^2*x^2-1))^(1/2)*(8*a^5*x^5-12*a^3*x^3+8*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4+4*a*x-8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)^(1/2))*(32*arccosh(a*x)^3-24*arccosh(a*x)^2+12*arccosh(a*x)-3)*c^2/(a*x-1)/(a*x+1)/a+15/512*(-c*(a^2*x^2-1))^(1/2)*(2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))*(4*arccosh(a*x)^3-6*arccosh(a*x)^2+6*arccosh(a*x)-3)*c^2/(a*x-1)/(a*x+1)/a+15/512*(-c*(a^2*x^2-1))^(1/2)*(2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)^(1/2))*(4*arccosh(a*x)^3+6*arccosh(a*x)^2+6*arccosh(a*x)+3)*c^2/(a*x-1)/(a*x+1)/a-3/4096*(-c*(a^2*x^2-1))^(1/2)*(8*a^5*x^5-12*a^3*x^3-8*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4+4*a*x+8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))*(32*arccosh(a*x)^3+24*arccosh(a*x)^2+12*arccosh(a*x)+3)*c^2/(a*x-1)/(a*x+1)/a+1/13824*(-c*(a^2*x^2-1))^(1/2)*(-32*x^6*a^6*(a*x-1)^(1/2)*(a*x+1)^(1/2)+32*a^7*x^7+48*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^4*x^4-64*a^5*x^5-18*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+38*a^3*x^3+(a*x-1)^(1/2)*(a*x+1)^(1/2)-6*a*x)*(36*arccosh(a*x)^3+18*arcco...`

### 3.246.5 Fracas [F]

$$\int (c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx = \int (-a^2 cx^2 + c)^{5/2} \operatorname{arccosh}(ax)^3 dx$$

input `integrate((-a^2*c*x^2+c)^(5/2)*arccosh(a*x)^3,x, algorithm="fracas")`

output `integral((a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)*sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^3, x)`

**3.246.6 Sympy [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx = \text{Timed out}$$

input `integrate((-a**2*c*x**2+c)**(5/2)*acosh(a*x)**3,x)`

output `Timed out`

**3.246.7 Maxima [F(-2)]**

Exception generated.

$$\int (c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: RuntimeError}$$

input `integrate((-a^2*c*x^2+c)^(5/2)*arccosh(a*x)^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.246.8 Giac [F(-2)]**

Exception generated.

$$\int (c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(5/2)*arccosh(a*x)^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.246.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^3 dx = \int \operatorname{acosh}(ax)^3 (c - a^2 cx^2)^{5/2} dx$$

input `int(acosh(a*x)^3*(c - a^2*c*x^2)^(5/2),x)`output `int(acosh(a*x)^3*(c - a^2*c*x^2)^(5/2), x)`

### 3.247 $\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx$

3.247.1 Optimal result . . . . .	2209
3.247.2 Mathematica [A] (warning: unable to verify) . . . . .	2210
3.247.3 Rubi [A] (verified) . . . . .	2210
3.247.4 Maple [A] (verified) . . . . .	2218
3.247.5 Fricas [F] . . . . .	2218
3.247.6 Sympy [F] . . . . .	2219
3.247.7 Maxima [F(-2)] . . . . .	2219
3.247.8 Giac [F(-2)] . . . . .	2219
3.247.9 Mupad [F(-1)] . . . . .	2220

#### 3.247.1 Optimal result

Integrand size = 22, antiderivative size = 402

$$\begin{aligned} \int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx = & -\frac{51acx^2\sqrt{c - a^2cx^2}}{128\sqrt{-1 + ax}\sqrt{1 + ax}} \\ & + \frac{3a^3cx^4\sqrt{c - a^2cx^2}}{128\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{45}{64}cx\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax) \\ & + \frac{3}{32}cx(1 - ax)(1 + ax)\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax) \\ & + \frac{27c\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^2}{128a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{9acx^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^2}{16\sqrt{-1 + ax}\sqrt{1 + ax}} \\ & + \frac{3c(1 - a^2x^2)^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^2}{16a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{3}{8}cx\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^3 \\ & + \frac{1}{4}x(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 - \frac{3c\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^4}{32a\sqrt{-1 + ax}\sqrt{1 + ax}} \end{aligned}$$

```
output 1/4*x*(-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^3+45/64*c*x*arccosh(a*x)*(-a^2*c*x
^2+c)^(1/2)+3/32*c*x*(-a*x+1)*(a*x+1)*arccosh(a*x)*(-a^2*c*x^2+c)^(1/2)+3/
8*c*x*arccosh(a*x)^3*(-a^2*c*x^2+c)^(1/2)-51/128*a*c*x^2*(-a^2*c*x^2+c)^(1
/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/128*a^3*c*x^4*(-a^2*c*x^2+c)^(1/2)/(a*x
-1)^(1/2)/(a*x+1)^(1/2)+27/128*c*arccosh(a*x)^2*(-a^2*c*x^2+c)^(1/2)/a/(a*x
-1)^(1/2)/(a*x+1)^(1/2)-9/16*a*c*x^2*arccosh(a*x)^2*(-a^2*c*x^2+c)^(1/2)/(
a*x-1)^(1/2)/(a*x+1)^(1/2)+3/16*c*(-a^2*x^2+1)^2*arccosh(a*x)^2*(-a^2*c*x^
2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-3/32*c*arccosh(a*x)^4*(-a^2*c*x^2
+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)
```

**3.247.2 Mathematica [A] (warning: unable to verify)**

Time = 0.41 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.37

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx =$$


---


$$\frac{c\sqrt{c - a^2 cx^2}(96\operatorname{arccosh}(ax)^4 - 3(-64 \cosh(2\operatorname{arccosh}(ax)) + \cosh(4\operatorname{arccosh}(ax))) - 24\operatorname{arccosh}(ax)^2(-16$$

input `Integrate[(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^3,x]`output `-1/1024*(c*Sqrt[c - a^2*c*x^2]*(96*ArcCosh[a*x]^4 - 3*(-64*Cosh[2*ArcCosh[a*x]] + Cosh[4*ArcCosh[a*x]]) - 24*ArcCosh[a*x]^2*(-16*Cosh[2*ArcCosh[a*x]] + Cosh[4*ArcCosh[a*x]]) + 12*ArcCosh[a*x]*(-32*Sinh[2*ArcCosh[a*x]] + Sinh[4*ArcCosh[a*x]]) + 32*ArcCosh[a*x]^3*(-8*Sinh[2*ArcCosh[a*x]] + Sinh[4*ArcCosh[a*x]])))/(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x))`**3.247.3 Rubi [A] (verified)**Time = 3.88 (sec) , antiderivative size = 390, normalized size of antiderivative = 0.97, number of steps used = 18, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$ , Rules used = {6312, 25, 6310, 6298, 6308, 6327, 6329, 6313, 25, 82, 244, 2009, 6311, 15, 6308, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(ax)^3 (c - a^2 cx^2)^{3/2} dx$$

$$\downarrow \text{6312}$$

$$\frac{3ac\sqrt{c - a^2 cx^2} \int -x(1 - ax)(ax + 1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{3}{4}c \int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^3 dx +$$

$$\frac{1}{4}x\operatorname{arccosh}(ax)^3 (c - a^2 cx^2)^{3/2}$$

$$\downarrow \text{25}$$

$$-\frac{3ac\sqrt{c - a^2 cx^2} \int x(1 - ax)(ax + 1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{3}{4}c \int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^3 dx +$$

$$\frac{1}{4}x\operatorname{arccosh}(ax)^3 (c - a^2 cx^2)^{3/2}$$

---

3.247.  $\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx$

$$\begin{aligned}
& \downarrow \text{6310} \\
& -\frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \int x\operatorname{arccosh}(ax)^2 dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^3\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{1}{4}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2} \\
& \downarrow \text{6298} \\
& -\frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a \int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^3\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{1}{4}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2} \\
& \downarrow \text{6308} \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a \int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^3\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2} \\
& \downarrow \text{6327} \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a \int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^3\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{3ac\sqrt{c-a^2cx^2} \int x(1-a^2x^2)\operatorname{arccosh}(ax)^2 dx}{4\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2} \\
& \downarrow \text{6329} \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^2 - a \int \frac{x^2\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4\sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^3\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{3ac\sqrt{c-a^2cx^2} \left( \frac{\int (ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax) dx}{2a} - \frac{(1-a^2x^2)^2\operatorname{arccosh}(ax)^2}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{4}x\operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}
\end{aligned}$$

↓ 6313

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right)$$


---


$$\frac{3ac\sqrt{c-a^2cx^2} \left( -\frac{\frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx - \frac{1}{4}a \int -x(1-ax)(ax+1) dx + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}$$

↓ 25

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right)$$


---


$$\frac{3ac\sqrt{c-a^2cx^2} \left( -\frac{\frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx + \frac{1}{4}a \int x(1-ax)(ax+1) dx + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}$$

↓ 82

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right)$$


---


$$\frac{3ac\sqrt{c-a^2cx^2} \left( \frac{\frac{1}{4}a \int x(1-a^2x^2) dx - \frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}$$

↓ 244

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right)$$


---


$$\frac{3ac\sqrt{c-a^2cx^2} \left( \frac{\frac{1}{4}a \int (x-a^2x^3) dx - \frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}$$

↓ 2009

---

3.247.  $\int (c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx$

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right) - \frac{3ac\sqrt{c-a^2cx^2} \left( -\frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) dx + \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}$$

↓ 6311

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right) - \frac{3ac\sqrt{c-a^2cx^2} \left( -\frac{3}{4} \left( -\frac{1}{2} \int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{a \int x dx}{2} + \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) \right) + \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}$$

↓ 15

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right) - \frac{3ac\sqrt{c-a^2cx^2} \left( -\frac{3}{4} \left( -\frac{1}{2} \int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{ax^2}{4} \right) + \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2}$$

↓ 6308

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \right) - \frac{3ac\sqrt{c-a^2cx^2} \left( \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}$$


---


$$\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c-a^2cx^2)^{3/2} - \frac{3ac\sqrt{c-a^2cx^2} \left( \frac{1}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax) \right)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}$$

↓ 6354

---

3.247.  $\int (c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx$

$$\frac{\frac{3}{4}c \left( \frac{3a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} - \frac{\int x dx}{2a} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}} \right) + \frac{\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c - a^2cx^2)^{3/2} - \frac{3}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}}{4\sqrt{ax-1}\sqrt{ax+1}}$$

↓ 15

$$\frac{\frac{3}{4}c \left( \frac{3a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} - \frac{x^2}{4a} \right) \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4}{8a\sqrt{ax-1}} \right) + \frac{\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c - a^2cx^2)^{3/2} - \frac{3}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}}{4\sqrt{ax-1}\sqrt{ax+1}}$$

↓ 6308

$$\frac{\frac{3}{4}c \left( -\frac{\operatorname{arccosh}(ax)^4 \sqrt{c - a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c - a^2cx^2} - \frac{3a \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2\sqrt{ax-1}\sqrt{ax+1}} \right) + \frac{\frac{1}{4}x \operatorname{arccosh}(ax)^3 (c - a^2cx^2)^{3/2} - \frac{3}{4}a \left( \frac{x^2}{2} - \frac{a^2x^4}{4} \right) - \frac{3}{4} \left( \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{4a} - \frac{ax^2}{4} \right) + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2}\operatorname{arccosh}(ax)}{2a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)}{4a^2}}{4\sqrt{ax-1}\sqrt{ax+1}}$$

input `Int[(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^3,x]`

```
output (x*(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^3)/4 + (3*c*((x*Sqrt[c - a^2*c*x^2]*
ArcCosh[a*x]^3)/2 - (Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^4)/(8*a*Sqrt[-1 + a*
x]*Sqrt[1 + a*x]) - (3*a*Sqrt[c - a^2*c*x^2]*((x^2*ArcCosh[a*x]^2)/2 - a*(
-1/4*x^2/a + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/(2*a^2) + ArcCo
sh[a*x]^2/(4*a^3))))/(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]))/4 - (3*a*c*Sqrt[c
- a^2*c*x^2]*(-1/4*((1 - a^2*x^2)^2*ArcCosh[a*x]^2)/a^2 + ((a*(x^2/2 - (a^
2*x^4)/4))/4 + (x*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2)*ArcCosh[a*x])/4 - (3*(-
1/4*(a*x^2) + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/2 - ArcCosh[a*
x]^2/(4*a)))/4)/(2*a)))/(4*Sqrt[-1 + a*x]*Sqrt[1 + a*x])
```

### 3.247.3.1 Defintions of rubi rules used

```
rule 15 Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[
{a, m}, x] && NeQ[m, -1]
```

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 82 Int[((a_) + (b_.)*(x_)^(m_.))*((c_) + (d_.)*(x_)^(n_.))*((e_) + (f_.)*(x_)
)^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d,
e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]
```

```
rule 244 Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand
Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p
, 0]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6298 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((d_.)*(x_)^(m_.), x_Symbol]
:= Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +
c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& NeQ[m, -1]
```



rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6311 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] := Simp[x*Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6313 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[x*(d1 + e1*x)^p*(d2 + e2*x)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d1*d2*(p/(2*p + 1)) Int[(d1 + e1*x)^(p - 1)*(d2 + e2*x)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x)) /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && GtQ[p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1)) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.247.4 Maple [A] (verified)

Time = 1.09 (sec) , antiderivative size = 536, normalized size of antiderivative = 1.33

method	result
default	$-\frac{3\sqrt{-c(a^2x^2-1)} \operatorname{arccosh}(ax)^4 c}{32\sqrt{ax-1}\sqrt{ax+1}a} - \frac{\sqrt{-c(a^2x^2-1)}(8a^5x^5-12a^3x^3+8\sqrt{ax-1}\sqrt{ax+1}a^4x^4+4ax-8a^2x^2\sqrt{ax-1}\sqrt{ax+1}+\sqrt{ax-1})}{2048(ax-1)(ax+1)a}$

input `int((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^3,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -3/32*(-c*(a^2*x^2-1))^{(1/2)}/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}/a*\operatorname{arccosh}(a*x)^4* \\
 & c-1/2048*(-c*(a^2*x^2-1))^{(1/2)}*(8*a^5*x^5-12*a^3*x^3+8*(a*x-1)^{(1/2)}*(a*x \\
 & +1)^{(1/2)}*a^4*x^4+4*a*x-8*a^2*x^2*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+(a*x-1)^{(1/2} \\
 & )*(a*x+1)^{(1/2)})*(32*\operatorname{arccosh}(a*x)^3-24*\operatorname{arccosh}(a*x)^2+12*\operatorname{arccosh}(a*x)-3)*c \\
 & /(a*x-1)/(a*x+1)/a+1/32*(-c*(a^2*x^2-1))^{(1/2)}*(2*a^3*x^3-2*a*x+2*a^2*x^2* \\
 & (a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})*(4*\operatorname{arccosh}(a*x)^3 \\
 & -6*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)-3)*c/(a*x-1)/(a*x+1)/a+1/32*(-c*(a^2*x^2- \\
 & 1))^{(1/2)}*(2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+(a*x-1)^{(1/2} \\
 & )*(a*x+1)^{(1/2)})*(4*\operatorname{arccosh}(a*x)^3+6*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)+3)*c \\
 & /(a*x-1)/(a*x+1)/a-1/2048*(-c*(a^2*x^2-1))^{(1/2)}*(8*a^5*x^5-12*a^3*x^3-8*( \\
 & a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a^4*x^4+4*a*x+8*a^2*x^2*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2} \\
 & )-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})*(32*\operatorname{arccosh}(a*x)^3+24*\operatorname{arccosh}(a*x)^2+12* \\
 & \operatorname{arccosh}(a*x)+3)*c/(a*x-1)/(a*x+1)/a
 \end{aligned}$$

### 3.247.5 Fracas [F]

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx = \int (-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^3 dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^3,x, algorithm="fricas")`

output `integral(-a^2*c*x^2 - c)*sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^3, x)`

**3.247.6 Sympy [F]**

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx = \int (-c(ax - 1)(ax + 1))^{3/2} \operatorname{acosh}^3(ax) dx$$

input `integrate((-a**2*c*x**2+c)**(3/2)*acosh(a*x)**3,x)`

output `Integral((-c*(a*x - 1)*(a*x + 1))**(3/2)*acosh(a*x)**3, x)`

**3.247.7 Maxima [F(-2)]**

Exception generated.

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: RuntimeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^3,x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.247.8 Giac [F(-2)]**

Exception generated.

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.247.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^3 dx = \int \operatorname{acosh}(ax)^3 (c - a^2 cx^2)^{3/2} dx$$

input `int(acosh(a*x)^3*(c - a^2*c*x^2)^(3/2),x)`output `int(acosh(a*x)^3*(c - a^2*c*x^2)^(3/2), x)`

### 3.248 $\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 dx$

3.248.1 Optimal result . . . . .	2221
3.248.2 Mathematica [A] (verified) . . . . .	2222
3.248.3 Rubi [A] (verified) . . . . .	2222
3.248.4 Maple [A] (verified) . . . . .	2225
3.248.5 Fracas [F] . . . . .	2225
3.248.6 Sympy [F] . . . . .	2225
3.248.7 Maxima [F(-2)] . . . . .	2226
3.248.8 Giac [F(-2)] . . . . .	2226
3.248.9 Mupad [F(-1)] . . . . .	2226

#### 3.248.1 Optimal result

Integrand size = 22, antiderivative size = 231

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 dx = -\frac{3ax^2\sqrt{c - a^2cx^2}}{8\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{3}{4}x\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax) + \frac{3\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^2}{8a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{3ax^2\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^2}{4\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{1}{2}x\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 - \frac{\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^4}{8a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

output

```
3/4*x*arccosh(a*x)*(-a^2*c*x^2+c)^(1/2)+1/2*x*arccosh(a*x)^3*(-a^2*c*x^2+c)^(1/2)-3/8*a*x^2*(-a^2*c*x^2+c)^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/8*arccosh(a*x)^2*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-3/4*a*x^2*arccosh(a*x)^2*(-a^2*c*x^2+c)^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/8*arccosh(a*x)^4*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)
```

**3.248.2 Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.42

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^3 dx = \frac{\sqrt{-c(-1+ax)(1+ax)}(2\operatorname{arccosh}(ax)^4 + (3 + 6\operatorname{arccosh}(ax)^2) \cosh(2\operatorname{arccosh}(ax)) - 2\operatorname{arccosh}(ax)(3 + 2\operatorname{arccosh}(ax)^2) \sinh(2\operatorname{arccosh}(ax)))}{16a\sqrt{\frac{-1+ax}{1+ax}}(1+ax)}$$

input `Integrate[Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^3,x]`output `-1/16*(Sqrt[-(c*(-1 + a*x)*(1 + a*x))]*(2*ArcCosh[a*x]^4 + (3 + 6*ArcCosh[a*x]^2)*Cosh[2*ArcCosh[a*x]] - 2*ArcCosh[a*x]*(3 + 2*ArcCosh[a*x]^2)*Sinh[2*ArcCosh[a*x]]))/(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x))`**3.248.3 Rubi [A] (verified)**Time = 1.28 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {6310, 6298, 6308, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{arccosh}(ax)^3 \sqrt{c - a^2 cx^2} dx \\ & \quad \downarrow \text{6310} \\ & -\frac{3a\sqrt{c - a^2 cx^2} \int x \operatorname{arccosh}(ax)^2 dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\sqrt{c - a^2 cx^2} \int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{ax - 1}\sqrt{ax + 1}} dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2} x \operatorname{arccosh}(ax)^3 \sqrt{c - a^2 cx^2} \\ & \quad \downarrow \text{6298} \\ & -\frac{3a\sqrt{c - a^2 cx^2} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax - 1}\sqrt{ax + 1}} dx \right)}{2\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\sqrt{c - a^2 cx^2} \int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{ax - 1}\sqrt{ax + 1}} dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} + \\ & \quad \frac{1}{2} x \operatorname{arccosh}(ax)^3 \sqrt{c - a^2 cx^2} \\ & \quad \downarrow \text{6308} \end{aligned}$$

$$\begin{aligned}
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{6354} \\
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} - \frac{\int x dx}{2a} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \\
& \quad \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{15} \\
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} - \frac{x^2}{4a} \right) \right)}{2\sqrt{ax-1}\sqrt{ax+1}} - \\
& \quad \frac{\operatorname{arccosh}(ax)^4 \sqrt{c-a^2cx^2}}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^3 \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{6308} \\
& \frac{3a \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} - \frac{x^2}{4a} \right) \right) \sqrt{c-a^2cx^2}}{2\sqrt{ax-1}\sqrt{ax+1}}
\end{aligned}$$

input `Int[Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^3,x]`

output `(x*Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^3)/2 - (Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^4)/(8*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) - (3*a*Sqrt[c - a^2*c*x^2]*((x^2*ArcCosh[a*x]^2)/2 - a*(-1/4*x^2/a + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/(2*a^2) + ArcCosh[a*x]^2/(4*a^3))))/(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`



## 3.248.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)])*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`
- rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

**3.248.4 Maple [A] (verified)**

Time = 0.73 (sec) , antiderivative size = 256, normalized size of antiderivative = 1.11

method	result
default	$-\frac{\sqrt{-c(a^2x^2-1)} \operatorname{arccosh}(ax)^4}{8\sqrt{ax-1}\sqrt{ax+1}a} + \frac{\sqrt{-c(a^2x^2-1)} (2a^3x^3-2ax+2a^2x^2\sqrt{ax-1}\sqrt{ax+1}-\sqrt{ax-1}\sqrt{ax+1}) (4 \operatorname{arccosh}(ax)^3-6 \operatorname{arccosh}(ax)^2+6 \operatorname{arccosh}(ax)-3)}{32(ax-1)(ax+1)a}$

```
input int(arccosh(a*x)^3*(-a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -1/8*(-c*(a^2*x^2-1))^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)/a*arccosh(a*x)^4+1/32*(-c*(a^2*x^2-1))^(1/2)*(2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))*(4*arccosh(a*x)^3-6*arccosh(a*x)^2+6*arccosh(a*x)-3)/(a*x-1)/(a*x+1)/a+1/32*(-c*(a^2*x^2-1))^(1/2)*(2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)^(1/2))*(4*arccosh(a*x)^3+6*arccosh(a*x)^2+6*arccosh(a*x)+3)/(a*x-1)/(a*x+1)/a
```

**3.248.5 Fricas [F]**

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 dx = \int \sqrt{-a^2cx^2 + c} \operatorname{arccosh}(ax)^3 dx$$

```
input integrate(arccosh(a*x)^3*(-a^2*c*x^2+c)^(1/2),x, algorithm="fricas")
```

```
output integral(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^3, x)
```

**3.248.6 Sympy [F]**

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 dx = \int \sqrt{-c(ax - 1)(ax + 1)} \operatorname{acosh}^3(ax) dx$$

```
input integrate(acosh(a*x)**3*(-a**2*c*x**2+c)**(1/2),x)
```

```
output Integral(sqrt(-c*(a*x - 1)*(a*x + 1))*acosh(a*x)**3, x)
```

**3.248.7 Maxima [F(-2)]**

Exception generated.

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: RuntimeError}$$

```
input integrate(arccosh(a*x)^3*(-a^2*c*x^2+c)^(1/2),x, algorithm="maxima")
```

```
output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.
```

**3.248.8 Giac [F(-2)]**

Exception generated.

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 dx = \text{Exception raised: TypeError}$$

```
input integrate(arccosh(a*x)^3*(-a^2*c*x^2+c)^(1/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.248.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^3 dx = \int \operatorname{acosh}(ax)^3 \sqrt{c - a^2cx^2} dx$$

```
input int(acosh(a*x)^3*(c - a^2*c*x^2)^(1/2),x)
```

```
output int(acosh(a*x)^3*(c - a^2*c*x^2)^(1/2), x)
```

$$3.249 \quad \int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c-a^2cx^2}} dx$$

3.249.1 Optimal result . . . . .	2227
3.249.2 Mathematica [A] (verified) . . . . .	2227
3.249.3 Rubi [A] (verified) . . . . .	2228
3.249.4 Maple [A] (verified) . . . . .	2228
3.249.5 Fricas [F] . . . . .	2229
3.249.6 Sympy [F] . . . . .	2229
3.249.7 Maxima [F] . . . . .	2229
3.249.8 Giac [F] . . . . .	2230
3.249.9 Mupad [F(-1)] . . . . .	2230

### 3.249.1 Optimal result

Integrand size = 22, antiderivative size = 46

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c-a^2cx^2}} dx = \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^4}{4a\sqrt{c-a^2cx^2}}$$

output `1/4*arccosh(a*x)^4*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(-a^2*c*x^2+c)^(1/2)`

### 3.249.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c-a^2cx^2}} dx = \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^4}{4a\sqrt{c-a^2cx^2}}$$

input `Integrate[ArcCosh[a*x]^3/Sqrt[c - a^2*c*x^2],x]`

output `(Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^4)/(4*a*Sqrt[c - a^2*c*x^2])`

**3.249.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx$$

↓ 6307

$$\frac{\sqrt{ax - 1}\sqrt{ax + 1}\operatorname{arccosh}(ax)^4}{4a\sqrt{c - a^2cx^2}}$$

input `Int[ArcCosh[a*x]^3/Sqrt[c - a^2*c*x^2],x]`

output `(Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^4)/(4*a*Sqrt[c - a^2*c*x^2])`

**3.249.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.249.4 Maple [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.20

method	result	size
default	$-\frac{\sqrt{-c(ax-1)(ax+1)}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^4}{4c(a^2x^2-1)a}$	55

input `int(arccosh(a*x)^3/(-a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4*(-c*(a*x-1)*(a*x+1))^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/c/(a^2*x^2-1)/a*arccosh(a*x)^4`

---

3.249.  $\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx$

**3.249.5 Fracas [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^3/(a^2*c*x^2 - c), x)`

**3.249.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{acosh}^3(ax)}{\sqrt{-c(ax - 1)(ax + 1)}} dx$$

input `integrate(acosh(a*x)**3/(-a**2*c*x**2+c)**(1/2),x)`

output `Integral(acosh(a*x)**3/sqrt(-c*(a*x - 1)*(a*x + 1)), x)`

**3.249.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^3/sqrt(-a^2*c*x^2 + c), x)`

**3.249.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^3/sqrt(-a^2*c*x^2 + c), x)`

**3.249.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{\sqrt{c - a^2cx^2}} dx$$

input `int(acosh(a*x)^3/(c - a^2*c*x^2)^(1/2),x)`

output `int(acosh(a*x)^3/(c - a^2*c*x^2)^(1/2), x)`

**3.250**      $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx$

3.250.1 Optimal result . . . . . 2231  
 3.250.2 Mathematica [A] (verified) . . . . . 2232  
 3.250.3 Rubi [C] (verified) . . . . . 2232  
 3.250.4 Maple [B] (verified) . . . . . 2235  
 3.250.5 Fricas [F] . . . . . 2236  
 3.250.6 Sympy [F] . . . . . 2236  
 3.250.7 Maxima [F] . . . . . 2237  
 3.250.8 Giac [F] . . . . . 2237  
 3.250.9 Mupad [F(-1)] . . . . . 2237

**3.250.1 Optimal result**

Integrand size = 22, antiderivative size = 241

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx = \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^3}{ac\sqrt{c-a^2cx^2}}$$

$$- \frac{3\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2 \log(1-e^{2\operatorname{arccosh}(ax)})}{ac\sqrt{c-a^2cx^2}}$$

$$- \frac{3\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)})}{ac\sqrt{c-a^2cx^2}}$$

$$+ \frac{3\sqrt{-1+ax}\sqrt{1+ax} \operatorname{PolyLog}(3, e^{2\operatorname{arccosh}(ax)})}{2ac\sqrt{c-a^2cx^2}}$$

```
output x*arccosh(a*x)^3/c/(-a^2*c*x^2+c)^(1/2)+arccosh(a*x)^3*(a*x-1)^(1/2)*(a*x+
1)^(1/2)/a/c/(-a^2*c*x^2+c)^(1/2)-3*arccosh(a*x)^2*ln(1-(a*x+(a*x-1)^(1/2)
*(a*x+1)^(1/2))^2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/c/(-a^2*c*x^2+c)^(1/2)-3*
arccosh(a*x)*polylog(2,(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))^2)*(a*x-1)^(1/2)*
(a*x+1)^(1/2)/a/c/(-a^2*c*x^2+c)^(1/2)+3/2*polylog(3,(a*x+(a*x-1)^(1/2)*(a
*x+1)^(1/2))^2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/c/(-a^2*c*x^2+c)^(1/2)
```



### 3.250.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 162, normalized size of antiderivative = 0.67

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx = \frac{\sqrt{-1+ax}\sqrt{1+ax} \left( \operatorname{arccosh}(ax)^3 + \frac{ax \operatorname{arccosh}(ax)^3}{\sqrt{-1+ax}\sqrt{1+ax}} - 3 \operatorname{arccosh}(ax)^2 \log(1 - e^{\operatorname{arccosh}(ax)}) \right)}{(c - a^2cx^2)^{3/2}}$$

input `Integrate[ArcCosh[a*x]^3/(c - a^2*c*x^2)^(3/2),x]`

output `(Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(ArcCosh[a*x]^3 + (a*x*ArcCosh[a*x]^3)/(Sqrt[-1 + a*x]*Sqrt[1 + a*x])) - 3*ArcCosh[a*x]^2*Log[1 - E^ArcCosh[a*x]] - 3*ArcCosh[a*x]^2*Log[1 + E^ArcCosh[a*x]] - 6*ArcCosh[a*x]*PolyLog[2, -E^ArcCosh[a*x]] - 6*ArcCosh[a*x]*PolyLog[2, E^ArcCosh[a*x]] + 6*PolyLog[3, -E^ArcCosh[a*x]] + 6*PolyLog[3, E^ArcCosh[a*x]])/(a*c*Sqrt[c - a^2*c*x^2])`

### 3.250.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.77 (sec) , antiderivative size = 143, normalized size of antiderivative = 0.59, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.455$ , Rules used = {6314, 6328, 3042, 26, 4199, 25, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx \\ & \quad \downarrow \text{6314} \\ & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} \\ & \quad \downarrow \text{6328} \\ & \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} - \frac{3\sqrt{ax-1}\sqrt{ax+1} \int \frac{ax \operatorname{arccosh}(ax)^2}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d \operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \\ & \quad \downarrow \text{3042} \\ & \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} - \frac{3\sqrt{ax-1}\sqrt{ax+1} \int -i \operatorname{arccosh}(ax)^2 \tan\left(i \operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d \operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \end{aligned}$$

---

3.250.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx$

$$\begin{aligned}
& \downarrow 26 \\
& \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \int \operatorname{arccosh}(ax)^2 \tan\left(i \operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d \operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \\
& \downarrow 4199 \\
& \frac{\frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + 3i\sqrt{ax-1}\sqrt{ax+1} \left(2i \int -\frac{e^{2 \operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)^2}{1-e^{2 \operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax) - \frac{1}{3} i \operatorname{arccosh}(ax)^3\right)}{ac\sqrt{c-a^2cx^2}} \\
& \downarrow 25 \\
& \frac{\frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + 3i\sqrt{ax-1}\sqrt{ax+1} \left(-2i \int \frac{e^{2 \operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)^2}{1-e^{2 \operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax) - \frac{1}{3} i \operatorname{arccosh}(ax)^3\right)}{ac\sqrt{c-a^2cx^2}} \\
& \downarrow 2620 \\
& \frac{\frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + 3i\sqrt{ax-1}\sqrt{ax+1} \left(-2i \left(\int \operatorname{arccosh}(ax) \log\left(1-e^{2 \operatorname{arccosh}(ax)}\right) d \operatorname{arccosh}(ax) - \frac{1}{2} \operatorname{arccosh}(ax)^2 \log\left(1-e^{2 \operatorname{arccosh}(ax)}\right)\right)\right)}{ac\sqrt{c-a^2cx^2}} \\
& \downarrow 3011 \\
& \frac{\frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + 3i\sqrt{ax-1}\sqrt{ax+1} \left(-2i \left(\frac{1}{2} \int \operatorname{PolyLog}\left(2, e^{2 \operatorname{arccosh}(ax)}\right) d \operatorname{arccosh}(ax) - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}\left(2, e^{2 \operatorname{arccosh}(ax)}\right) - \frac{1}{2} \operatorname{arccosh}(ax)^2 \log\left(1-e^{2 \operatorname{arccosh}(ax)}\right)\right)\right)}{ac\sqrt{c-a^2cx^2}} \\
& \downarrow 2720 \\
& \frac{\frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + 3i\sqrt{ax-1}\sqrt{ax+1} \left(-2i \left(\frac{1}{4} \int e^{-2 \operatorname{arccosh}(ax)} \operatorname{PolyLog}\left(2, e^{2 \operatorname{arccosh}(ax)}\right) d e^{2 \operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}\left(2, e^{2 \operatorname{arccosh}(ax)}\right) - \frac{1}{2} \operatorname{arccosh}(ax)^2 \log\left(1-e^{2 \operatorname{arccosh}(ax)}\right)\right)\right)}{ac\sqrt{c-a^2cx^2}} \\
& \downarrow 7143 \\
& \frac{\frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + 3i\sqrt{ax-1}\sqrt{ax+1} \left(-2i \left(-\frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}\left(2, e^{2 \operatorname{arccosh}(ax)}\right) + \frac{1}{4} \operatorname{PolyLog}\left(3, e^{2 \operatorname{arccosh}(ax)}\right) - \frac{1}{2} \operatorname{arccosh}(ax)^2 \log\left(1-e^{2 \operatorname{arccosh}(ax)}\right)\right)\right)}{ac\sqrt{c-a^2cx^2}}
\end{aligned}$$

---

3.250.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx$

input `Int[ArcCosh[a*x]^3/(c - a^2*c*x^2)^(3/2),x]`

output `(x*ArcCosh[a*x]^3)/(c*Sqrt[c - a^2*c*x^2]) + ((3*I)*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*((-1/3*I)*ArcCosh[a*x]^3 - (2*I)*(-1/2*(ArcCosh[a*x]^2*Log[1 - E^(2*ArcCosh[a*x])]) - (ArcCosh[a*x]*PolyLog[2, E^(2*ArcCosh[a*x])]))/2 + PolyLog[3, E^(2*ArcCosh[a*x])]/4))/(a*c*Sqrt[c - a^2*c*x^2])`

### 3.250.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

```
rule 4199 Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_
.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp
[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x
)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && In
tegerQ[4*k] && IGtQ[m, 0]
```

```
rule 6314 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_.) + (e_.)*(x_)^2)^(3/2),
x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp
[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a
+ b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e},
x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]
```

```
rule 6328 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_.) + (e_.)*(x_)^2),
x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]
], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

### 3.250.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 547 vs.  $2(252) = 504$ .

Time = 1.14 (sec) , antiderivative size = 548, normalized size of antiderivative = 2.27

method	result
default	$-\frac{\sqrt{-c(a^2x^2-1)}(-\sqrt{ax-1}\sqrt{ax+1}+ax)\operatorname{arccosh}(ax)^3}{c^2a(a^2x^2-1)} - \frac{2\sqrt{ax+1}\sqrt{ax-1}\sqrt{-c(a^2x^2-1)}\operatorname{arccosh}(ax)^3}{c^2a(a^2x^2-1)} + \frac{3\sqrt{ax+1}\sqrt{ax-1}\sqrt{-c(a^2x^2-1)}\operatorname{arccosh}(ax)^3}{c^2a(a^2x^2-1)}$

```
input int(arccosh(a*x)^3/(-a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)
```

---

3.250. 
$$\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx$$

output 
$$\begin{aligned} & -(-c(a^2x^2-1))^{1/2} * (-a^2x^2-1)^{1/2} * (ax+1)^{1/2} + ax * \operatorname{arccosh}(ax)^3 / \\ & c^2/a/(a^2x^2-1) - 2*(ax+1)^{1/2} * (ax-1)^{1/2} * (-c(a^2x^2-1))^{1/2} / c^2 \\ & /a/(a^2x^2-1) * \operatorname{arccosh}(ax)^3 + 3*(ax+1)^{1/2} * (ax-1)^{1/2} * (-c(a^2x^2-1))^{1/2} / c^2 \\ & /a/(a^2x^2-1) * \operatorname{arccosh}(ax)^2 * \ln(1+ax+(ax-1)^{1/2} * (ax+1)^{1/2}) \\ & + 6*(ax+1)^{1/2} * (ax-1)^{1/2} * (-c(a^2x^2-1))^{1/2} / c^2 / a / (a^2x^2-1) \\ & * \operatorname{arccosh}(ax) * \operatorname{polylog}(2, -ax - (ax-1)^{1/2} * (ax+1)^{1/2}) - 6*(ax+1)^{1/2} \\ & * (ax-1)^{1/2} * (-c(a^2x^2-1))^{1/2} / c^2 / a / (a^2x^2-1) * \operatorname{polylog}(3, -ax - (ax-1)^{1/2} \\ & * (ax+1)^{1/2}) + 3*(ax+1)^{1/2} * (ax-1)^{1/2} * (-c(a^2x^2-1))^{1/2} / c^2 / a / (a^2x^2-1) \\ & * \operatorname{arccosh}(ax)^2 * \ln(1-ax - (ax-1)^{1/2} * (ax+1)^{1/2}) \\ & + 6*(ax+1)^{1/2} * (ax-1)^{1/2} * (-c(a^2x^2-1))^{1/2} / c^2 / a / (a^2x^2-1) \\ & * \operatorname{arccosh}(ax) * \operatorname{polylog}(2, ax + (ax-1)^{1/2} * (ax+1)^{1/2}) - 6*(ax+1)^{1/2} * (ax-1)^{1/2} \\ & * (-c(a^2x^2-1))^{1/2} / c^2 / a / (a^2x^2-1) * \operatorname{polylog}(3, ax + (ax-1)^{1/2} * (ax+1)^{1/2}) \end{aligned}$$

### 3.250.5 Fricas [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(-a^2cx^2 + c)^{3/2}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^3/(a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2), x)`

### 3.250.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{acosh}^3(ax)}{(-c(ax-1)(ax+1))^{3/2}} dx$$

input `integrate(acosh(a*x)**3/(-a**2*c*x**2+c)**(3/2),x)`

output `Integral(acosh(a*x)**3/(-c*(a*x - 1)*(a*x + 1))**(3/2), x)`

**3.250.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(-a^2cx^2 + c)^{3/2}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^3/(-a^2*c*x^2 + c)^(3/2), x)`

**3.250.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(-a^2cx^2 + c)^{3/2}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(3/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^3/(-a^2*c*x^2 + c)^(3/2), x)`

**3.250.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{(c - a^2cx^2)^{3/2}} dx$$

input `int(acosh(a*x)^3/(c - a^2*c*x^2)^(3/2),x)`

output `int(acosh(a*x)^3/(c - a^2*c*x^2)^(3/2), x)`

### 3.251 $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx$

3.251.1 Optimal result . . . . .	2238
3.251.2 Mathematica [C] (warning: unable to verify) . . . . .	2239
3.251.3 Rubi [C] (verified) . . . . .	2240
3.251.4 Maple [B] (verified) . . . . .	2246
3.251.5 Fricas [F] . . . . .	2247
3.251.6 Sympy [F] . . . . .	2247
3.251.7 Maxima [F] . . . . .	2247
3.251.8 Giac [F(-2)] . . . . .	2248
3.251.9 Mupad [F(-1)] . . . . .	2248

#### 3.251.1 Optimal result

Integrand size = 22, antiderivative size = 413

$$\begin{aligned}
 \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx &= -\frac{x\operatorname{arccosh}(ax)}{c^2\sqrt{c-a^2cx^2}} + \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{2ac^2(1-a^2x^2)\sqrt{c-a^2cx^2}} \\
 &+ \frac{x\operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} + \frac{2x\operatorname{arccosh}(ax)^3}{3c^2\sqrt{c-a^2cx^2}} + \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^3}{3ac^2\sqrt{c-a^2cx^2}} \\
 &- \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2 \log(1-e^{2\operatorname{arccosh}(ax)})}{ac^2\sqrt{c-a^2cx^2}} \\
 &+ \frac{\sqrt{-1+ax}\sqrt{1+ax} \log(1-a^2x^2)}{2ac^2\sqrt{c-a^2cx^2}} \\
 &- \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)})}{ac^2\sqrt{c-a^2cx^2}} \\
 &+ \frac{\sqrt{-1+ax}\sqrt{1+ax} \operatorname{PolyLog}(3, e^{2\operatorname{arccosh}(ax)})}{ac^2\sqrt{c-a^2cx^2}}
 \end{aligned}$$

output  $\frac{1}{3}x \operatorname{arccosh}(ax)^3 / c / (-a^2cx^2+c)^{(3/2)} - x \operatorname{arccosh}(ax) / c^2 / (-a^2cx^2+c)^{(1/2)} + 2/3x \operatorname{arccosh}(ax)^3 / c^2 / (-a^2cx^2+c)^{(1/2)} + 1/2 \operatorname{arccosh}(ax)^2 * (ax-1)^{(1/2)} * (ax+1)^{(1/2)} / a / c^2 / (-a^2cx^2+c)^{(1/2)} + 2/3 \operatorname{arccosh}(ax)^3 * (ax-1)^{(1/2)} * (ax+1)^{(1/2)} / a / c^2 / (-a^2cx^2+c)^{(1/2)} - 2 \operatorname{arccosh}(ax)^2 * \ln(1 - (ax+(ax-1)^{(1/2)} * (ax+1)^{(1/2)})^2) * (ax-1)^{(1/2)} * (ax+1)^{(1/2)} / a / c^2 / (-a^2cx^2+c)^{(1/2)} + 1/2 \ln(-a^2cx^2+c) * (ax-1)^{(1/2)} * (ax+1)^{(1/2)} / a / c^2 / (-a^2cx^2+c)^{(1/2)} - 2 \operatorname{arccosh}(ax) * \operatorname{polylog}(2, (ax+(ax-1)^{(1/2)} * (ax+1)^{(1/2)})^2) * (ax-1)^{(1/2)} * (ax+1)^{(1/2)} / a / c^2 / (-a^2cx^2+c)^{(1/2)} + \operatorname{polylog}(3, (ax+(ax-1)^{(1/2)} * (ax+1)^{(1/2)})^2) * (ax-1)^{(1/2)} * (ax+1)^{(1/2)} / a / c^2 / (-a^2cx^2+c)^{(1/2)}$

### 3.251.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.90 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.65

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx = \frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax) \left( -i\pi^3 - \frac{12ax\sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)}{-1+ax} + \frac{6\operatorname{arccosh}(ax)^2}{1-a^2x^2} + 8\operatorname{arccosh}(ax)^3 + \dots \right)}{(c - a^2cx^2)^{5/2}}$$

input `Integrate[ArcCosh[a*x]^3/(c - a^2*c*x^2)^(5/2),x]`

output  $(\operatorname{Sqrt}[(-1 + ax)/(1 + ax)] * (1 + ax) * ((-1) * \pi^3 - (12 * ax * \operatorname{Sqrt}[(-1 + ax)/(1 + ax)] * \operatorname{ArcCosh}[a*x]) / (-1 + ax) + (6 * \operatorname{ArcCosh}[a*x]^2) / (1 - a^2 * x^2) + 8 * \operatorname{ArcCosh}[a*x]^3 + (8 * ax * \operatorname{Sqrt}[(-1 + ax)/(1 + ax)] * \operatorname{ArcCosh}[a*x]^3) / (-1 + ax) - (4 * ax * ((-1 + ax)/(1 + ax))^{(3/2)} * \operatorname{ArcCosh}[a*x]^3) / (-1 + ax)^3 - 24 * \operatorname{ArcCosh}[a*x]^2 * \operatorname{Log}[1 - E^{(2 * \operatorname{ArcCosh}[a*x])}] + 12 * \operatorname{Log}[a*x] + 12 * \operatorname{Log}[(\operatorname{Sqrt}[(-1 + ax)/(1 + ax)] * (1 + ax)) / (a*x)] - 24 * \operatorname{ArcCosh}[a*x] * \operatorname{PolyLog}[2, E^{(2 * \operatorname{ArcCosh}[a*x])}] + 12 * \operatorname{PolyLog}[3, E^{(2 * \operatorname{ArcCosh}[a*x])}])) / (12 * a * c^2 * \operatorname{Sqrt}[c - a^2 * c * x^2])$



**3.251.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.12 (sec) , antiderivative size = 292, normalized size of antiderivative = 0.71, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.682$ , Rules used = {6316, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 3011, 2720, 6329, 6315, 240, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6316} \\
 & \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^2(ax+1)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx}{3c} + \frac{x\operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \\
 & \quad \downarrow \text{6314} \\
 & \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^2(ax+1)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \\
 & \quad \frac{x\operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \\
 & \quad \frac{x\operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \\
 & \quad \downarrow \text{6328} \\
 & \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \\
 & \quad 2 \left( \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} - \frac{3\sqrt{ax-1}\sqrt{ax+1} \int \frac{ax\operatorname{arccosh}(ax)^2}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} d\operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \right) + \frac{x\operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.251.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \\
& 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} - \frac{3\sqrt{ax-1}\sqrt{ax+1} \int -i \operatorname{arccosh}(ax)^2 \tan\left(i \operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d \operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \right) + \\
& \frac{3c}{3c(c-a^2cx^2)^{3/2}} x \operatorname{arccosh}(ax)^3 \\
& \quad \downarrow \text{26} \\
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \\
& 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \int \operatorname{arccosh}(ax)^2 \tan\left(i \operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d \operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \right) + \\
& \frac{3c}{3c(c-a^2cx^2)^{3/2}} x \operatorname{arccosh}(ax)^3 \\
& \quad \downarrow \text{4199} \\
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \\
& 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( 2i \int -\frac{e^{2 \operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)^2}{1-e^{2 \operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax) - \frac{1}{3} i \operatorname{arccosh}(ax)^3 \right)}{ac\sqrt{c-a^2cx^2}} \right) + \\
& \frac{3c}{3c(c-a^2cx^2)^{3/2}} x \operatorname{arccosh}(ax)^3 \\
& \quad \downarrow \text{25} \\
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \\
& 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \int \frac{e^{2 \operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)^2}{1-e^{2 \operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax) - \frac{1}{3} i \operatorname{arccosh}(ax)^3 \right)}{ac\sqrt{c-a^2cx^2}} \right) + \\
& \frac{3c}{3c(c-a^2cx^2)^{3/2}} x \operatorname{arccosh}(ax)^3 \\
& \quad \downarrow \text{2620}
\end{aligned}$$

---

3.251.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx$

$$\frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \int \operatorname{arccosh}(ax) \log(1-e^{2\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \frac{1}{2} \operatorname{arccosh}(ax)^2 \log(1-e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)$$

3c

$$\frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}}$$

↓ 3011

$$\frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{2} \int \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \operatorname{darccosh}(ax) - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) - \frac{1}{2} \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)$$

3c

$$\frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}}$$

↓ 2720

$$\frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) de^{2\operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)$$

3c

$$\frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}}$$

↓ 6329

$$\frac{a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx}{a} + \frac{\operatorname{arccosh}(ax)^2}{2a^2(1-a^2x^2)} \right)}{c^2\sqrt{c-a^2cx^2}} + 2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) de^{2\operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)$$

3c

$$\frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}}$$

↓ 6315

---

3.251.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{-a \int \frac{x}{1-a^2x^2} dx - \frac{x \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}}{a} + \frac{\operatorname{arccosh}(ax)^2}{2a^2(1-a^2x^2)} \right)}{c^2\sqrt{c-a^2cx^2}} + \\
& \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) de^{2\operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \\
& \frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \\
& \quad \downarrow \quad 240 \\
& \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) de^{2\operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \\
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{2a^2(1-a^2x^2)} + \frac{\frac{\log(1-a^2x^2)}{2a} - \frac{x \operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}}}{a} \right)}{c^2\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \\
& \quad \downarrow \quad 7143 \\
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{2a^2(1-a^2x^2)} + \frac{\frac{\log(1-a^2x^2)}{2a} - \frac{x \operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}}}{a} \right)}{c^2\sqrt{c-a^2cx^2}} + \\
& \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( -\frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) + \frac{1}{4} \operatorname{PolyLog}(3, e^{2\operatorname{arccosh}(ax)}) - \frac{1}{2} \operatorname{arccosh}(ax)^2 \operatorname{Log} \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \\
& \frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}}
\end{aligned}$$

input `Int[ArcCosh[a*x]^3/(c - a^2*c*x^2)^(5/2), x]`

output `(x*ArcCosh[a*x]^3)/(3*c*(c - a^2*c*x^2)^(3/2)) + (a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(ArcCosh[a*x]^2/(2*a^2*(1 - a^2*x^2)) + (-((x*ArcCosh[a*x])/(Sqrt[-1 + a*x]*Sqrt[1 + a*x])) + Log[1 - a^2*x^2]/(2*a))/a)/(c^2*Sqrt[c - a^2*c*x^2]) + (2*((x*ArcCosh[a*x]^3)/(c*Sqrt[c - a^2*c*x^2]) + ((3*I)*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*((-1/3*I)*ArcCosh[a*x]^3 - (2*I)*(-1/2*(ArcCosh[a*x]^2*Log[1 - E^(2*ArcCosh[a*x])])) - (ArcCosh[a*x]*PolyLog[2, E^(2*ArcCosh[a*x])]))/2 + PolyLog[3, E^(2*ArcCosh[a*x])]/4))/(a*c*Sqrt[c - a^2*c*x^2]))/(3*c)`

3.251.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx$

## 3.251.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 240 `Int[(x_)/((a_) + (b_)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`
- rule 2620 `Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_)^(m_)))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`
- rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`
- rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_)^(m_)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 4199 `Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[((c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6315 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(((d1_) + (e1_.)*(x_))^(3/2)*((d2_) + (e2_.)*(x_))^(3/2)), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d1*d2*Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x])), x] + Simp[b*c*(n/(d1*d2))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_))/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.251.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 954 vs.  $2(400) = 800$ .

Time = 1.32 (sec) , antiderivative size = 955, normalized size of antiderivative = 2.31

method	result
default	$-\frac{\sqrt{-c(a^2x^2-1)}(2a^3x^3-3ax-2a^2x^2\sqrt{ax-1}\sqrt{ax+1}+2\sqrt{ax-1}\sqrt{ax+1})\operatorname{arccosh}(ax)\left(6a^3x^3\operatorname{arccosh}(ax)\sqrt{ax-1}\sqrt{ax+1}+6a^4x^4\operatorname{arccosh}(ax)\right)}{\dots}$

input `int(arccosh(a*x)^3/(-a^2*c*x^2+c)^(5/2),x,method=_RETURNVERBOSE)`

output

$$\begin{aligned}
 & -1/6*(-c*(a^2*x^2-1))^{(1/2)}*(2*a^3*x^3-3*a*x-2*a^2*x^2*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+2*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})*\operatorname{arccosh}(a*x)*(6*a^3*x^3*\operatorname{arccosh}(a*x)*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+6*a^4*x^4*\operatorname{arccosh}(a*x)+6*a^3*x^3*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+6*a^4*x^4+6*a^2*x^2*\operatorname{arccosh}(a*x)^2-9*a*x*\operatorname{arccosh}(a*x)*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}-12*a^2*x^2*\operatorname{arccosh}(a*x)-6*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a*x-18*a^2*x^2-8*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)+12)/(3*a^6*x^6-10*a^4*x^4+11*a^2*x^2-4)/c^3/a-(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\ln((a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+a*x-1)-(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\ln(1+a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})+2*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\ln(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})-4/3*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^3+2*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^2*\ln(1+a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})+4*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\operatorname{arccosh}(a*x)*\operatorname{polylog}(2,-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})-4*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\operatorname{polylog}(3,-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})+2*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^2*\ln(1-a*x-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})+4*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)}*(-c*(a^2*x^2-1))^{(1/2)}/c^3/a/(a^2*x^2-1)*\operatorname{arccosh}(a*x)*\operatorname{polylog}(2,a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})
 \end{aligned}$$

**3.251.5 Fracas [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(-a^2cx^2 + c)^{5/2}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(5/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^3/(a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3), x)`

**3.251.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\operatorname{acosh}^3(ax)}{(-c(ax - 1)(ax + 1))^{5/2}} dx$$

input `integrate(acosh(a*x)**3/(-a**2*c*x**2+c)**(5/2),x)`

output `Integral(acosh(a*x)**3/(-c*(a*x - 1)*(a*x + 1))**(5/2), x)`

**3.251.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(-a^2cx^2 + c)^{5/2}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(5/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^3/(-a^2*c*x^2 + c)^(5/2), x)`



**3.251.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.251.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{(c - a^2cx^2)^{5/2}} dx$$

input `int(acosh(a*x)^3/(c - a^2*c*x^2)^(5/2),x)`

output `int(acosh(a*x)^3/(c - a^2*c*x^2)^(5/2), x)`

### 3.252 $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

3.252.1 Optimal result . . . . .	2249
3.252.2 Mathematica [C] (warning: unable to verify) . . . . .	2250
3.252.3 Rubi [C] (verified) . . . . .	2251
3.252.4 Maple [B] (verified) . . . . .	2261
3.252.5 Fricas [F] . . . . .	2262
3.252.6 Sympy [F(-1)] . . . . .	2263
3.252.7 Maxima [F] . . . . .	2263
3.252.8 Giac [F(-2)] . . . . .	2263
3.252.9 Mupad [F(-1)] . . . . .	2264

#### 3.252.1 Optimal result

Integrand size = 22, antiderivative size = 607

$$\begin{aligned}
\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx = & -\frac{\sqrt{-1+ax}\sqrt{1+ax}}{20ac^3(1-a^2x^2)\sqrt{c-a^2cx^2}} - \frac{x\operatorname{arccosh}(ax)}{c^3\sqrt{c-a^2cx^2}} \\
& - \frac{x\operatorname{arccosh}(ax)}{10c^3(1-ax)(1+ax)\sqrt{c-a^2cx^2}} + \frac{3\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{20ac^3(1-a^2x^2)^2\sqrt{c-a^2cx^2}} \\
& + \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2}{5ac^3(1-a^2x^2)\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}} \\
& + \frac{4x\operatorname{arccosh}(ax)^3}{15c^2(c-a^2cx^2)^{3/2}} + \frac{8x\operatorname{arccosh}(ax)^3}{15c^3\sqrt{c-a^2cx^2}} + \frac{8\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^3}{15ac^3\sqrt{c-a^2cx^2}} \\
& - \frac{8\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^2 \log(1-e^{2\operatorname{arccosh}(ax)})}{5ac^3\sqrt{c-a^2cx^2}} \\
& + \frac{\sqrt{-1+ax}\sqrt{1+ax} \log(1-a^2x^2)}{2ac^3\sqrt{c-a^2cx^2}} \\
& - \frac{8\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)})}{5ac^3\sqrt{c-a^2cx^2}} \\
& + \frac{4\sqrt{-1+ax}\sqrt{1+ax} \operatorname{PolyLog}(3, e^{2\operatorname{arccosh}(ax)})}{5ac^3\sqrt{c-a^2cx^2}}
\end{aligned}$$

output  $\frac{1}{5}x \operatorname{arccosh}(ax)^3 / c / (-a^2cx^2+c)^{(5/2)} + \frac{4}{15}x \operatorname{arccosh}(ax)^3 / c^2 / (-a^2cx^2+c)^{(3/2)} - x \operatorname{arccosh}(ax) / c^3 / (-a^2cx^2+c)^{(1/2)} - \frac{1}{10}x \operatorname{arccosh}(ax) / c^3 / (-ax+1) / (ax+1) / (-a^2cx^2+c)^{(1/2)} + \frac{8}{15}x \operatorname{arccosh}(ax)^3 / c^3 / (-a^2cx^2+c)^{(1/2)} - \frac{1}{20}(ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)} + \frac{3}{20} \operatorname{arccosh}(ax)^2 (ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)} + \frac{2}{5} \operatorname{arccosh}(ax)^2 (ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)} + \frac{8}{15} \operatorname{arccosh}(ax)^3 (ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)} - \frac{8}{5} \operatorname{arccosh}(ax)^2 \ln(1-(ax+(ax-1)^{(1/2)}(ax+1)^{(1/2}))^2) (ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)} + \frac{1}{2} \ln(-a^2cx^2+c)^{(1/2)} (ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)} - \frac{8}{5} \operatorname{arccosh}(ax) \operatorname{polylog}(2, (ax+(ax-1)^{(1/2)}(ax+1)^{(1/2}))^2) (ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)} + \frac{4}{5} \operatorname{polylog}(3, (ax+(ax-1)^{(1/2)}(ax+1)^{(1/2}))^2) (ax-1)^{(1/2)}(ax+1)^{(1/2)} / a / c^3 / (-a^2cx^2+c)^{(1/2)}$

### 3.252.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.64 (sec) , antiderivative size = 375, normalized size of antiderivative = 0.62

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx = \sqrt{\frac{-1+ax}{1+ax}}(1+ax) \left( 4i\pi^3 + \frac{3}{1-a^2x^2} + \frac{60ax\sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)}{-1+ax} - \frac{6ax\left(\frac{-1+ax}{1+ax}\right)^{3/2} \operatorname{arccosh}(ax)}{(-1+ax)^3} - \frac{9\operatorname{arccosh}(ax)^2}{(-1+a^2x^2)^2} + \frac{24\operatorname{arccosh}(ax)}{(-1+a^2x^2)^2} \right)$$

input `Integrate[ArcCosh[a*x]^3/(c - a^2*c*x^2)^(7/2),x]`

output 
$$\frac{-1/60 \left( \sqrt{\frac{-1+ax}{1+ax}} (1+ax) \left( (4I)\pi^3 + \frac{3}{1-a^2x^2} + \frac{60ax\sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)}{-1+ax} - \frac{6ax\left(\frac{-1+ax}{1+ax}\right)^{3/2} \operatorname{arccosh}(ax)}{(-1+ax)^3} - \frac{9\operatorname{arccosh}(ax)^2}{(-1+a^2x^2)^2} + \frac{24\operatorname{arccosh}(ax)}{(-1+a^2x^2)^2} \right) - (6ax\sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)) / (-1+ax) - (6ax\sqrt{\frac{-1+ax}{1+ax}})^{3/2} \operatorname{arccosh}(ax) / (-1+ax)^3 - (9\operatorname{arccosh}(ax)^2) / (-1+a^2x^2)^2 + (24\operatorname{arccosh}(ax)^2) / (-1+a^2x^2)^2 - 32\operatorname{arccosh}(ax)^3 - (32ax\sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)^3) / (-1+ax) + (16ax\sqrt{\frac{-1+ax}{1+ax}})^{3/2} \operatorname{arccosh}(ax)^3 / (-1+ax)^3 - (12ax\sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)^3) / ((-1+ax)^3(1+ax)^2) + 96\operatorname{arccosh}(ax)^2 \operatorname{Log}[1 - E^{(2\operatorname{arccosh}(ax))}] - 60\operatorname{Log}[ax] - 60\operatorname{Log}[\sqrt{\frac{-1+ax}{1+ax}}(1+ax)] / (ax) + 96\operatorname{arccosh}(ax) \operatorname{PolyLog}[2, E^{(2\operatorname{arccosh}(ax))}] - 48\operatorname{PolyLog}[3, E^{(2\operatorname{arccosh}(ax))}] \right)}{a^3c\sqrt{c - a^2cx^2}}$$

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

## 3.252.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 4.00 (sec) , antiderivative size = 498, normalized size of antiderivative = 0.82, number of steps used = 23, number of rules used = 22,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6316, 25, 6316, 6314, 6327, 6328, 3042, 26, 4199, 25, 2620, 3011, 2720, 6329, 6315, 240, 6317, 82, 241, 6315, 240, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx \\
 & \quad \downarrow \text{6316} \\
 & -\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int -\frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \frac{4 \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx}{5c} + \frac{x\operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \frac{4 \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{5/2}} dx}{5c} + \frac{x\operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{6316} \\
 & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\
 & 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^2(ax+1)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{3/2}} dx}{3c} + \frac{x\operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \right) \\
 & \quad \downarrow \text{6314} \\
 & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^3(ax+1)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\
 & 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{(1-ax)^2(ax+1)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\operatorname{arccosh}(ax)^2}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x\operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \right) \\
 & \quad \downarrow \\
 & \frac{5c}{5c(c-a^2cx^2)^{5/2}} \frac{x\operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}} +
 \end{aligned}$$

---

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

$$\begin{aligned} & \downarrow 6327 \\ & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\ & 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{1-a^2x^2} dx}{c\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \right) \\ & \frac{5c}{5c(c-a^2cx^2)^{5/2}} x \operatorname{arccosh}(ax)^3 \end{aligned}$$

$$\begin{aligned} & \downarrow 6328 \\ & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\ & 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} - \frac{3\sqrt{ax-1}\sqrt{ax+1} \int \frac{ax \operatorname{arccosh}(ax)^2}{\sqrt{\frac{ax-1}{ax+1}}(ax+1)} dx}{ac\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x \operatorname{arccosh}(ax)^3}{3c(c-a^2cx^2)^{3/2}} \right) \\ & \frac{5c}{5c(c-a^2cx^2)^{5/2}} x \operatorname{arccosh}(ax)^3 \end{aligned}$$

$$\begin{aligned} & \downarrow 3042 \\ & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + \\ & 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} - \frac{3\sqrt{ax-1}\sqrt{ax+1} \int -i \operatorname{arccosh}(ax)^2 \tan\left(i \operatorname{arccosh}(ax) + \frac{\pi}{2}\right) dx}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right) \\ & \frac{5c}{5c(c-a^2cx^2)^{5/2}} x \operatorname{arccosh}(ax)^3 \end{aligned}$$

$$\downarrow 26$$

---

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \int \operatorname{arccosh}(ax)^2 \tan\left(i \operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d \operatorname{arccosh}(ax)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right) +$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}} \quad 5c$$

↓ 4199

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( 2i \int -\frac{e^{2 \operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)^2}{1-e^{2 \operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax) - \frac{1}{3} i \operatorname{arccos} \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right) +$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}} \quad 5c$$

↓ 25

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} + 4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \int \frac{e^{2 \operatorname{arccosh}(ax)} \operatorname{arccosh}(ax)^2}{1-e^{2 \operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax) - \frac{1}{3} i \operatorname{arccos} \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right) +$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}} \quad 5c$$

↓ 2620

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \int \operatorname{arccosh}(ax) \log(1-e^{2\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \frac{1}{2} \operatorname{arccosh}(ax) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

5c

↓ 3011

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{2} \int \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \frac{1}{2} \operatorname{arccosh}(ax) \right) \right)}{3c} \right)}{3c} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

5c

↓ 2720

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^3} dx}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^2}{(1-a^2x^2)^2} dx}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) de^{2\operatorname{arccosh}(ax)} \right) \right)}{3c} \right)}{3c} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

5c

↓ 6329

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

$$\begin{aligned}
 & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{\int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2} dx}}{2a} \right)}{5c^3\sqrt{c-a^2cx^2}} + \\
 4 & \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2} dx}}{a} + \frac{\operatorname{arccosh}(ax)^2}{2a^2(1-a^2x^2)} \right)}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) dx \right) \right)}{c\sqrt{c-a^2cx^2}} \right)}{3c} \right)
 \end{aligned}$$

$$\frac{x\operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 6315

$$\begin{aligned}
 & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{\int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2} dx}}{2a} \right)}{5c^3\sqrt{c-a^2cx^2}} + \\
 4 & \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{-a \int \frac{x}{1-a^2x^2} dx - \frac{x\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}}{a} + \frac{\operatorname{arccosh}(ax)^2}{2a^2(1-a^2x^2)} \right)}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) dx \right) \right)}{c\sqrt{c-a^2cx^2}} \right)}{3c} \right)
 \end{aligned}$$

$$\frac{x\operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 240

$$\begin{aligned}
 & \frac{3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{\int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{5/2}(ax+1)^{5/2} dx}}{2a} \right)}{5c^3\sqrt{c-a^2cx^2}} + \\
 4 & \left( \frac{2 \left( \frac{x\operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) dx \right) \right)}{c\sqrt{c-a^2cx^2}} \right)}{3c} \right)
 \end{aligned}$$

$$\frac{x\operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 6317

---

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$



$$3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{-\frac{2}{3} \int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx + \frac{1}{3} a \int \frac{x}{(1-ax)^2(ax+1)^2} dx - \frac{x \operatorname{arccosh}(ax)}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{2a} \right) +$$


---


$$4 \left( \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) de^{2\operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)}{5c^3\sqrt{c-a^2cx^2}}$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 82

$$3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{\frac{1}{3} a \int \frac{x}{(1-a^2x^2)^2} dx - \frac{2}{3} \int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx - \frac{x \operatorname{arccosh}(ax)}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{2a} \right) +$$


---


$$4 \left( \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) de^{2\operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)}{5c^3\sqrt{c-a^2cx^2}}$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 241

$$3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{-\frac{2}{3} \int \frac{\operatorname{arccosh}(ax)}{(ax-1)^{3/2}(ax+1)^{3/2}} dx + \frac{1}{6a(1-a^2x^2)} - \frac{x \operatorname{arccosh}(ax)}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{2a} \right) +$$


---


$$4 \left( \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) de^{2\operatorname{arccosh}(ax)} - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog} \left( 2, e^{2\operatorname{arccosh}(ax)} \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)}{5c^3\sqrt{c-a^2cx^2}}$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 6315

---

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{-\frac{2}{3} \left( -a \int \frac{x}{1-a^2x^2} dx - \frac{x \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} \right) + \frac{1}{6a(1-a^2x^2)} - \frac{x \operatorname{arccosh}(ax)}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{2a} \right)}{5c^3\sqrt{c-a^2cx^2}} +$$

$$4 \left( \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) dx - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 240

$$4 \left( \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( \frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) dx - \frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{-\frac{2}{3} \left( \frac{\log(1-a^2x^2)}{2a} - \frac{x \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} \right) + \frac{1}{6a(1-a^2x^2)} - \frac{x \operatorname{arccosh}(ax)}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{2a} \right)}{5c^3\sqrt{c-a^2cx^2}} +$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

↓ 7143

$$3a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{4a^2(1-a^2x^2)^2} - \frac{-\frac{2}{3} \left( \frac{\log(1-a^2x^2)}{2a} - \frac{x \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} \right) + \frac{1}{6a(1-a^2x^2)} - \frac{x \operatorname{arccosh}(ax)}{3(ax-1)^{3/2}(ax+1)^{3/2}}}{2a} \right) +$$

$$4 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \left( \frac{\operatorname{arccosh}(ax)^2}{2a^2(1-a^2x^2)} + \frac{\log(1-a^2x^2)}{2a} - \frac{x \operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \right)}{c^2\sqrt{c-a^2cx^2}} + \frac{2 \left( \frac{x \operatorname{arccosh}(ax)^3}{c\sqrt{c-a^2cx^2}} + \frac{3i\sqrt{ax-1}\sqrt{ax+1} \left( -2i \left( -\frac{1}{2} \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, e^{2\operatorname{arccosh}(ax)}) \right) \right)}{ac\sqrt{c-a^2cx^2}} \right)}{3c} \right)$$

$$\frac{x \operatorname{arccosh}(ax)^3}{5c(c-a^2cx^2)^{5/2}}$$

5c

3.252.  $\int \frac{\operatorname{arccosh}(ax)^3}{(c-a^2cx^2)^{7/2}} dx$

input `Int[ArcCosh[a*x]^3/(c - a^2*c*x^2)^(7/2),x]`

output `(x*ArcCosh[a*x]^3)/(5*c*(c - a^2*c*x^2)^(5/2)) + (3*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(ArcCosh[a*x]^2/(4*a^2*(1 - a^2*x^2)^2) - (1/(6*a*(1 - a^2*x^2)) - (x*ArcCosh[a*x])/(3*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2)) - (2*(-((x*ArcCosh[a*x])/(Sqrt[-1 + a*x]*Sqrt[1 + a*x])) + Log[1 - a^2*x^2]/(2*a))))/(5*c^3*Sqrt[c - a^2*c*x^2]) + (4*((x*ArcCosh[a*x]^3)/(3*c*(c - a^2*c*x^2)^(3/2)) + (a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(ArcCosh[a*x]^2/(2*a^2*(1 - a^2*x^2)) + (-((x*ArcCosh[a*x])/(Sqrt[-1 + a*x]*Sqrt[1 + a*x])) + Log[1 - a^2*x^2]/(2*a))/a)/(c^2*Sqrt[c - a^2*c*x^2]) + (2*((x*ArcCosh[a*x]^3)/(c*Sqrt[c - a^2*c*x^2]) + ((3*I)*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*((-1/3*I)*ArcCosh[a*x]^3 - (2*I)*(-1/2*(ArcCosh[a*x]^2*Log[1 - E^(2*ArcCosh[a*x]])) - (ArcCosh[a*x]*PolyLog[2, E^(2*ArcCosh[a*x]]))/2 + PolyLog[3, E^(2*ArcCosh[a*x]])/4)))/(a*c*Sqrt[c - a^2*c*x^2])))/(3*c))/(5*c)`

### 3.252.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 82 `Int[((a_) + (b_.)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Int[(a*c + b*d*x^2)^m*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m] && IntegerQ[m]`

rule 240 `Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]`

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 2620 `Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[(c + d*x)^m/(b*f*g*n*Log[F])*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Simp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4199 `Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x))/(1 + E^(2*((-I)*e + f*fz*x)))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]`

rule 6314 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)/((d_) + (e_)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6315 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(((d1_) + (e1_.)*(x_))^(3/2)*((d2_) + (e2_.)*(x_))^(3/2)), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d1*d2*Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x]), x] + Simp[b*c*(n/(d1*d2))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6317 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(-x)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d1*d2*(p + 1))), x] + (Simp[(2*p + 3)/(2*d1*d2*(p + 1)) Int[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6328 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)/((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[1/e Subst[Int[(a + b*x)^n*Coth[x], x], x, ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0]`

```
rule 6329 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x
])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&
GtQ[n, 0] && NeQ[p, -1]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

### 3.252.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1318 vs.  $2(564) = 1128$ .

Time = 1.34 (sec) , antiderivative size = 1319, normalized size of antiderivative = 2.17

method	result	size
default	Expression too large to display	1319

```
input int(arccosh(a*x)^3/(-a^2*c*x^2+c)^(7/2),x,method=_RETURNVERBOSE)
```

```
output -1/60*(-c*(a^2*x^2-1))^(1/2)*(8*a^5*x^5-20*a^3*x^3-8*(a*x-1)^(1/2)*(a*x+1)
^(1/2)*a^4*x^4+15*a*x+16*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-8*(a*x-1)^(1/2)
*(a*x+1)^(1/2)*(24-380*a^2*x^2*arccosh(a*x)^3+1410*a^2*x^2*arccosh(a*x)
-1590*a^4*x^4*arccosh(a*x)+984*a^2*x^2*arccosh(a*x)^2-1368*a^4*x^4*arccosh
(a*x)^2-936*a^3*x^3*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)+24*(a*x+1)^(1
/2)*(a*x-1)^(1/2)*a^7*x^7-96*a^2*x^2+144*a^4*x^4-1020*a^3*x^3*arccosh(a*x)
^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-96*a^6*x^6+256*arccosh(a*x)^3-264*arccosh(a
*x)^2-480*arccosh(a*x)+24*a^8*x^8+105*a^3*x^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)-
84*x^5*a^5*(a*x-1)^(1/2)*(a*x+1)^(1/2)-192*arccosh(a*x)*a^8*x^8+852*arccos
h(a*x)*a^6*x^6-192*arccosh(a*x)^2*a^8*x^8+840*arccosh(a*x)^2*a^6*x^6+160*a
rccosh(a*x)^3*a^4*x^4-192*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^7*x^7
+756*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^5*x^5-45*(a*x-1)^(1/2)*(a
*x+1)^(1/2)*a*x-192*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^7*x^7+744*
arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a^5*x^5+372*a*x*arccosh(a*x)*(a
*x-1)^(1/2)*(a*x+1)^(1/2)+495*a*x*arccosh(a*x)^2*(a*x-1)^(1/2)*(a*x+1)^(1
/2))/(40*a^10*x^10-215*a^8*x^8+469*a^6*x^6-517*a^4*x^4+287*a^2*x^2-64)/c^4/
a-(a*x+1)^(1/2)*(a*x-1)^(1/2)*(-c*(a^2*x^2-1))^(1/2)/c^4/a/(a^2*x^2-1)*ln(
(a*x-1)^(1/2)*(a*x+1)^(1/2)+a*x-1)-(a*x+1)^(1/2)*(a*x-1)^(1/2)*(-c*(a^2*x^
2-1))^(1/2)/c^4/a/(a^2*x^2-1)*ln(1+a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))+2*(a*x
+1)^(1/2)*(a*x-1)^(1/2)*(-c*(a^2*x^2-1))^(1/2)/c^4/a/(a^2*x^2-1)*ln(a*x...
```

### 3.252.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(-a^2cx^2 + c)^{7/2}} dx$$

```
input integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(7/2),x, algorithm="fricas")
```

```
output integral(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^3/(a^8*c^4*x^8 - 4*a^6*c^4*x^6
+ 6*a^4*c^4*x^4 - 4*a^2*c^4*x^2 + c^4), x)
```

**3.252.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx = \text{Timed out}$$

```
input integrate(acosh(a*x)**3/(-a**2*c*x**2+c)**(7/2),x)
```

```
output Timed out
```

**3.252.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{(-a^2cx^2 + c)^{7/2}} dx$$

```
input integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(7/2),x, algorithm="maxima")
```

```
output integrate(arccosh(a*x)^3/(-a^2*c*x^2 + c)^(7/2), x)
```

**3.252.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(arccosh(a*x)^3/(-a^2*c*x^2+c)^(7/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```



**3.252.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{(c - a^2cx^2)^{7/2}} dx$$

input `int(acosh(a*x)^3/(c - a^2*c*x^2)^(7/2), x)`output `int(acosh(a*x)^3/(c - a^2*c*x^2)^(7/2), x)`

### 3.253 $\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

3.253.1 Optimal result . . . . .	2265
3.253.2 Mathematica [A] (verified) . . . . .	2266
3.253.3 Rubi [A] (verified) . . . . .	2266
3.253.4 Maple [B] (verified) . . . . .	2271
3.253.5 Fricas [F] . . . . .	2272
3.253.6 Sympy [F] . . . . .	2272
3.253.7 Maxima [F(-2)] . . . . .	2272
3.253.8 Giac [F] . . . . .	2273
3.253.9 Mupad [F(-1)] . . . . .	2273

#### 3.253.1 Optimal result

Integrand size = 24, antiderivative size = 315

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = -\frac{45x^2\sqrt{-1+ax}}{128a^3\sqrt{1-ax}} - \frac{3x^4\sqrt{-1+ax}}{128a\sqrt{1-ax}} - \frac{45x\sqrt{1-ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{64a^4} - \frac{3x^3\sqrt{1-ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{32a^2} + \frac{45\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{128a^5\sqrt{1-ax}} - \frac{9x^2\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{16a^3\sqrt{1-ax}} - \frac{3x^4\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{16a\sqrt{1-ax}} - \frac{3x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{8a^4} - \frac{x^3\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{4a^2} + \frac{3\sqrt{-1+ax}\operatorname{arccosh}(ax)^4}{32a^5\sqrt{1-ax}}$$

output

```
-45/128*x^2*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-3/128*x^4*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)+45/128*arccosh(a*x)^2*(a*x-1)^(1/2)/a^5/(-a*x+1)^(1/2)-9/16*x^2*arccosh(a*x)^2*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-3/16*x^4*arccosh(a*x)^2*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)+3/32*arccosh(a*x)^4*(a*x-1)^(1/2)/a^5/(-a*x+1)^(1/2)-45/64*x*arccosh(a*x)*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^4-3/32*x^3*arccosh(a*x)*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^2-3/8*x*arccosh(a*x)^3*(-a^2*x^2+1)^(1/2)/a^4-1/4*x^3*arccosh(a*x)^3*(-a^2*x^2+1)^(1/2)/a^2
```

### 3.253.2 Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.43

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

$$= \frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax)(-192(1+2\operatorname{arccosh}(ax))^2 \cosh(2\operatorname{arccosh}(ax)) - 3(1+8\operatorname{arccosh}(ax)^2) \cosh(4\operatorname{arccosh}(ax)))}{1024a^5}$$

1024a

input `Integrate[(x^4*ArcCosh[a*x]^3)/Sqrt[1 - a^2*x^2], x]`

output `(Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(-192*(1 + 2*ArcCosh[a*x]^2)*Cosh[2*ArcCosh[a*x]] - 3*(1 + 8*ArcCosh[a*x]^2)*Cosh[4*ArcCosh[a*x]] + 4*ArcCosh[a*x]*(24*ArcCosh[a*x]^3 + 32*(3 + 2*ArcCosh[a*x]^2)*Sinh[2*ArcCosh[a*x]] + (3 + 8*ArcCosh[a*x]^2)*Sinh[4*ArcCosh[a*x]]))/((1024*a^5*Sqrt[-((-1 + a*x)*(1 + a*x))])`

### 3.253.3 Rubi [A] (verified)

Time = 3.17 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.10, number of steps used = 11, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.458$ , Rules used = {6353, 6298, 6353, 6298, 6307, 6354, 15, 6308, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

$$\downarrow \text{6353}$$

$$\frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{3\sqrt{ax-1} \int x^3 \operatorname{arccosh}(ax)^2 dx}{4a\sqrt{1-ax}} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2}$$

$$\downarrow \text{6298}$$

$$\frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{4a^2} - \frac{3\sqrt{ax-1} \left( \frac{1}{4} x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2} a \int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4a\sqrt{1-ax}} - \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2}$$

$$\downarrow \text{6353}$$

---

3.253.  $\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
& 3 \left( \frac{\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{3\sqrt{ax-1} \int x \operatorname{arccosh}(ax)^2 dx}{2a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \right) \\
& \frac{4a^2}{3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)} - \frac{x^3\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2} \\
& \quad \downarrow \text{6298} \\
& 3 \left( \frac{\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \right) \\
& \frac{4a^2}{3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)} - \frac{x^3\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2} \\
& \quad \downarrow \text{6307} \\
& 3 \left( -\frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2a\sqrt{1-ax}} + \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \right) \\
& \frac{4a^2}{3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \int \frac{x^4 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)} - \frac{x^3\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2} \\
& \quad \downarrow \text{6354} \\
& 3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \left( \frac{3 \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{4a^2} - \frac{\int x^3 dx}{4a} + \frac{x^3\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{4a^2} \right) \right) \\
& \frac{4a\sqrt{1-ax}}{3 \left( \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} - \frac{\int x dx}{2a} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2a\sqrt{1-ax}} + \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}}{4a^2} \right)} \\
& \frac{4a^2}{x^3\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3} \\
& \quad \downarrow \text{15}
\end{aligned}$$

$$\begin{aligned}
 & \frac{3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \left( \frac{3 \int \frac{x^2 \operatorname{arccosh}(ax) dx}{\sqrt{ax-1}\sqrt{ax+1}}}{4a^2} + \frac{x^3 \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{4a^2} - \frac{x^4}{16a} \right) \right)}{4a\sqrt{1-ax}} + \\
 & 3 \left( \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax) dx}{\sqrt{ax-1}\sqrt{ax+1}}}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} - \frac{x^2}{4a} \right) \right)}{2a\sqrt{1-ax}} + \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \right) \\
 & \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2} \\
 & \quad \downarrow \text{6308} \\
 & \frac{3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \left( \frac{3 \int \frac{x^2 \operatorname{arccosh}(ax) dx}{\sqrt{ax-1}\sqrt{ax+1}}}{4a^2} + \frac{x^3 \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{4a^2} - \frac{x^4}{16a} \right) \right)}{4a\sqrt{1-ax}} - \\
 & \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2} + \\
 & 3 \left( \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} - \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2a\sqrt{1-ax}} - \frac{x^4}{16a} \right) \\
 & \quad \downarrow \text{6354} \\
 & 3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \left( \frac{3 \left( \frac{\int \frac{\operatorname{arccosh}(ax) dx}{\sqrt{ax-1}\sqrt{ax+1}}}{2a^2} - \frac{\int x dx}{2a} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} \right)}{4a^2} + \frac{x^3 \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{4a^2} \right) \right) \\
 & \frac{x^3 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{4a^2} + \\
 & 3 \left( \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} - \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2a\sqrt{1-ax}} - \frac{x^4}{16a} \right) \\
 & \quad \downarrow \text{15}
 \end{aligned}$$

3.253.  $\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
 & \frac{3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax) - \frac{x^2}{4a}}{2a^2} \right)}{4a^2} \right) + \frac{x^3\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{4a^2}}{4a\sqrt{1-ax}} \\
 & \frac{x^3\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{4a^2} + \\
 & \frac{3 \left( \frac{\sqrt{ax-1}\operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2a^2} - \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} \right) - \frac{x^2}{4a} \right)}{2a\sqrt{1-ax}} \right)}{4a^2}}{4a\sqrt{1-ax}} \\
 & \quad \downarrow \text{6308} \\
 & \frac{x^3\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{4a^2} + \\
 & \frac{3 \left( \frac{\sqrt{ax-1}\operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2a^2} - \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} \right) - \frac{x^2}{4a} \right)}{2a\sqrt{1-ax}} \right)}{4a^2}}{4a\sqrt{1-ax}} \\
 & \frac{3\sqrt{ax-1} \left( \frac{1}{4}x^4 \operatorname{arccosh}(ax)^2 - \frac{1}{2}a \left( \frac{x^3\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{4a^2} + \frac{3 \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} \right) - \frac{x^2}{4a}}{4a^2} \right)}{4a\sqrt{1-ax}} \right)}{4a\sqrt{1-ax}}
 \end{aligned}$$

input `Int[(x^4*ArcCosh[a*x]^3)/Sqrt[1 - a^2*x^2],x]`

output `-1/4*(x^3*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3)/a^2 - (3*Sqrt[-1 + a*x]*((x^4*ArcCosh[a*x]^2)/4 - (a*(-1/16*x^4/a + (x^3*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]))/(4*a^2) + (3*(-1/4*x^2/a + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]))/(2*a^2) + ArcCosh[a*x]^2/(4*a^3)))/(4*a^2)))/2)/(4*a*Sqrt[1 - a*x]) + (3*(-1/2*(x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3)/a^2 + (Sqrt[-1 + a*x]*ArcCosh[a*x]^4)/(8*a^3*Sqrt[1 - a*x]) - (3*Sqrt[-1 + a*x]*((x^2*ArcCosh[a*x]^2)/2 - a*(-1/4*x^2/a + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]))/(2*a^2) + ArcCosh[a*x]^2/(4*a^3)))/(2*a*Sqrt[1 - a*x])))/(4*a^2)`

3.253.  $\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

## 3.253.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x)) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

```
rule 6354 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e
1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] :> Simp[f*(f*x)^(m -
1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(
m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m
- 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f
*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(
-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(
a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f,
p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && N
eQ[m + 2*p + 1, 0]
```

### 3.253.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 519 vs. 2(259) = 518.

Time = 0.89 (sec) , antiderivative size = 520, normalized size of antiderivative = 1.65

method	result
default	$-\frac{3\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^4}{32a^5(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(8a^5x^5-12a^3x^3+8\sqrt{ax-1}\sqrt{ax+1}a^4x^4+4ax-8a^2x^2\sqrt{ax-1}\sqrt{ax+1})}{2048a^5(a^2x^2-1)}$

```
input int(x^4*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output -3/32*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a^5/(a^2*x^2-1)*arcco
sh(a*x)^4-1/2048*(-a^2*x^2+1)^(1/2)*(8*a^5*x^5-12*a^3*x^3+8*(a*x-1)^(1/2)*
(a*x+1)^(1/2)*a^4*x^4+4*a*x-8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(
1/2)*(a*x+1)^(1/2))*(32*arccosh(a*x)^3-24*arccosh(a*x)^2+12*arccosh(a*x)-
3)/a^5/(a^2*x^2-1)-1/32*(-a^2*x^2+1)^(1/2)*(2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x
-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(a*x+1)^(1/2))*(4*arccosh(a*x)^3-6*a
rccosh(a*x)^2+6*arccosh(a*x)-3)/a^5/(a^2*x^2-1)-1/32*(-a^2*x^2+1)^(1/2)*(2
*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)+(a*x-1)^(1/2)*(a*x+1)
^(1/2))*(4*arccosh(a*x)^3+6*arccosh(a*x)^2+6*arccosh(a*x)+3)/a^5/(a^2*x^2-
1)-1/2048*(-a^2*x^2+1)^(1/2)*(8*a^5*x^5-12*a^3*x^3-8*(a*x-1)^(1/2)*(a*x+1)
^(1/2)*a^4*x^4+4*a*x+8*a^2*x^2*(a*x-1)^(1/2)*(a*x+1)^(1/2)-(a*x-1)^(1/2)*(
a*x+1)^(1/2))*(32*arccosh(a*x)^3+24*arccosh(a*x)^2+12*arccosh(a*x)+3)/a^5/
(a^2*x^2-1)
```

$$3.253. \int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$



**3.253.5 Fricas [F]**

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^4*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^4*arccosh(a*x)^3/(a^2*x^2 - 1), x)`

**3.253.6 Sympy [F]**

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{acosh}^3(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**4*acosh(a*x)**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**4*acosh(a*x)**3/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.253.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

**3.253.8 Giac [F]**

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^4*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^4*arccosh(a*x)^3/sqrt(-a^2*x^2 + 1), x)`

**3.253.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^4 \operatorname{acosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

input `int((x^4*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2),x)`

output `int((x^4*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2), x)`

### 3.254 $\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

3.254.1 Optimal result . . . . .	2274
3.254.2 Mathematica [A] (verified) . . . . .	2275
3.254.3 Rubi [A] (verified) . . . . .	2275
3.254.4 Maple [A] (verified) . . . . .	2279
3.254.5 Fricas [A] (verification not implemented) . . . . .	2280
3.254.6 Sympy [F] . . . . .	2280
3.254.7 Maxima [C] (verification not implemented) . . . . .	2280
3.254.8 Giac [F(-2)] . . . . .	2281
3.254.9 Mupad [F(-1)] . . . . .	2282

#### 3.254.1 Optimal result

Integrand size = 24, antiderivative size = 243

$$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = -\frac{40x\sqrt{-1+ax}}{9a^3\sqrt{1-ax}} - \frac{2x^3\sqrt{-1+ax}}{27a\sqrt{1-ax}} - \frac{40\sqrt{1-ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{9a^4} - \frac{2x^2\sqrt{1-ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{9a^2} - \frac{2x\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{a^3\sqrt{1-ax}} - \frac{x^3\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{3a\sqrt{1-ax}} - \frac{2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{3a^4} - \frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{3a^2}$$

```
output -40/9*x*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-2/27*x^3*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-2*x*arccosh(a*x)^2*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-1/3*x^3*arccosh(a*x)^2*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)-40/9*arccosh(a*x)*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^4-2/9*x^2*arccosh(a*x)*(-a*x+1)^(1/2)*(a*x+1)^(1/2)/a^2-2/3*arccosh(a*x)^3*(-a^2*x^2+1)^(1/2)/a^4-1/3*x^2*arccosh(a*x)^3*(-a^2*x^2+1)^(1/2)/a^2
```

**3.254.2 Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.58

$$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

$$= \frac{\sqrt{1-a^2x^2}(2ax(6+a^2x^2) - 6\sqrt{-1+ax}\sqrt{1+ax}(20+a^2x^2) \operatorname{arccosh}(ax) + 9ax(6+a^2x^2) \operatorname{arccosh}(ax)^2)}{27a^4\sqrt{-1+ax}\sqrt{1+ax}}$$

input `Integrate[(x^3*ArcCosh[a*x]^3)/Sqrt[1 - a^2*x^2],x]`output `(Sqrt[1 - a^2*x^2]*(2*a*x*(60 + a^2*x^2) - 6*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(20 + a^2*x^2)*ArcCosh[a*x] + 9*a*x*(6 + a^2*x^2)*ArcCosh[a*x]^2 - 9*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(2 + a^2*x^2)*ArcCosh[a*x]^3))/(27*a^4*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`**3.254.3 Rubi [A] (verified)**Time = 2.02 (sec) , antiderivative size = 259, normalized size of antiderivative = 1.07, number of steps used = 10, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6353, 6298, 6329, 6294, 6330, 24, 6354, 15, 6330, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

$$\downarrow \text{6353}$$

$$\frac{2 \int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{\sqrt{ax-1} \int x^2 \operatorname{arccosh}(ax)^2 dx}{a\sqrt{1-ax}} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{3a^2}$$

$$\downarrow \text{6298}$$

$$\frac{2 \int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{3a^2} - \frac{\sqrt{ax-1} \left( \frac{1}{3} x^3 \operatorname{arccosh}(ax)^2 - \frac{2}{3} a \int \frac{x^3 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a\sqrt{1-ax}} - \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{3a^2}$$

$$\downarrow \text{6329}$$

3.254.  $\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
 & \frac{2\left(-\frac{3\sqrt{ax-1}\int\operatorname{arccosh}(ax)^2dx}{a\sqrt{1-ax}}-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{a^2}\right)}{3a^2} \\
 & \frac{\sqrt{ax-1}\left(\frac{1}{3}x^3\operatorname{arccosh}(ax)^2-\frac{2}{3}a\int\frac{x^3\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{a\sqrt{1-ax}}-\frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{3a^2} \\
 & \quad \downarrow 6294 \\
 & \frac{2\left(-\frac{3\sqrt{ax-1}\left(x\operatorname{arccosh}(ax)^2-2a\int\frac{x\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{a\sqrt{1-ax}}-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{a^2}\right)}{3a^2} \\
 & \frac{\sqrt{ax-1}\left(\frac{1}{3}x^3\operatorname{arccosh}(ax)^2-\frac{2}{3}a\int\frac{x^3\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{a\sqrt{1-ax}}-\frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{3a^2} \\
 & \quad \downarrow 6330 \\
 & \frac{2\left(-\frac{3\sqrt{ax-1}\left(x\operatorname{arccosh}(ax)^2-2a\left(\frac{\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{a^2}-\frac{\int 1dx}{a}\right)\right)}{a\sqrt{1-ax}}-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{a^2}\right)}{3a^2} \\
 & \frac{\sqrt{ax-1}\left(\frac{1}{3}x^3\operatorname{arccosh}(ax)^2-\frac{2}{3}a\int\frac{x^3\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{a\sqrt{1-ax}}-\frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{3a^2} \\
 & \quad \downarrow 24 \\
 & \frac{\sqrt{ax-1}\left(\frac{1}{3}x^3\operatorname{arccosh}(ax)^2-\frac{2}{3}a\int\frac{x^3\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx\right)}{a\sqrt{1-ax}}-\frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{3a^2}+ \\
 & \frac{2\left(-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{a^2}-\frac{3\sqrt{ax-1}\left(x\operatorname{arccosh}(ax)^2-2a\left(\frac{\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{a^2}-\frac{x}{a}\right)\right)}{a\sqrt{1-ax}}\right)}{3a^2} \\
 & \quad \downarrow 6354 \\
 & \frac{\sqrt{ax-1}\left(\frac{1}{3}x^3\operatorname{arccosh}(ax)^2-\frac{2}{3}a\left(\frac{2\int\frac{x\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}}dx}{3a^2}-\frac{\int x^2dx}{3a}+\frac{x^2\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{3a^2}\right)\right)}{a\sqrt{1-ax}} \\
 & \frac{x^2\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{3a^2}+ \\
 & \frac{2\left(-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{a^2}-\frac{3\sqrt{ax-1}\left(x\operatorname{arccosh}(ax)^2-2a\left(\frac{\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{a^2}-\frac{x}{a}\right)\right)}{a\sqrt{1-ax}}\right)}{3a^2} \\
 & \quad \downarrow 15
 \end{aligned}$$

3.254.  $\int \frac{x^3\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
 & \frac{\sqrt{ax-1} \left( \frac{1}{3}x^3 \operatorname{arccosh}(ax)^2 - \frac{2}{3}a \left( \frac{2 \int \frac{x \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{3a^2} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{3a^2} - \frac{x^3}{9a} \right) \right)}{a\sqrt{1-ax}} \\
 & \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{3a^2} + \\
 & 2 \left( -\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{a^2} - \frac{3\sqrt{ax-1} \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right)}{a\sqrt{1-ax}} \right) \\
 & \frac{3a^2}{\phantom{a\sqrt{1-ax}}} \quad \downarrow \quad 6330 \\
 & \frac{\sqrt{ax-1} \left( \frac{1}{3}x^3 \operatorname{arccosh}(ax)^2 - \frac{2}{3}a \left( \frac{2 \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{\int 1 dx}{a} \right)}{3a^2} + \frac{x^2 \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{3a^2} - \frac{x^3}{9a} \right) \right)}{a\sqrt{1-ax}} \\
 & \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{3a^2} + \\
 & 2 \left( -\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{a^2} - \frac{3\sqrt{ax-1} \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right)}{a\sqrt{1-ax}} \right) \\
 & \frac{3a^2}{\phantom{a\sqrt{1-ax}}} \quad \downarrow \quad 24 \\
 & \frac{x^2 \sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{3a^2} + \\
 & 2 \left( -\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{a^2} - \frac{3\sqrt{ax-1} \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right)}{a\sqrt{1-ax}} \right) \\
 & \frac{3a^2}{\phantom{a\sqrt{1-ax}}} \\
 & \frac{\sqrt{ax-1} \left( \frac{1}{3}x^3 \operatorname{arccosh}(ax)^2 - \frac{2}{3}a \left( \frac{x^2 \sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{3a^2} + \frac{2 \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right)}{3a^2} - \frac{x^3}{9a} \right) \right)}{a\sqrt{1-ax}}
 \end{aligned}$$

input `Int[(x^3*ArcCosh[a*x]^3)/Sqrt[1 - a^2*x^2],x]`

output `-1/3*(x^2*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3)/a^2 - (Sqrt[-1 + a*x]*((x^3*ArcCosh[a*x]^2)/3 - (2*a*(-1/9*x^3/a + (x^2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/(3*a^2) + (2*(-(x/a) + (Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/a^2))/(3*a^2)))/3)/(a*Sqrt[1 - a*x]) + (2*(-((Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3)/a^2) - (3*Sqrt[-1 + a*x]*(x*ArcCosh[a*x]^2 - 2*a*(-(x/a) + (Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/a^2)))/(a*Sqrt[1 - a*x])))/(3*a^2)`

3.254.  $\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

## 3.254.3.1 Defintions of rubi rules used

- rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`
- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`
- rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`
- rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`
- rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_)^(p_) * ((d2_) + (e2_.)*(x_)^(p_)), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_)^2)^(p_)*((d2_) + (e2_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x^2)^(p + 1)*(d2 + e2*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d1 + e1*x^2)^p*(d2 + e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x^2)^p/(1 + c*x)^p]*Simp[(d2 + e2*x^2)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.254.4 Maple [A] (verified)

Time = 1.38 (sec) , antiderivative size = 375, normalized size of antiderivative = 1.54

method	result
default	$-\frac{\sqrt{-a^2x^2+1}(4a^4x^4-5a^2x^2+4a^3x^3\sqrt{ax-1}\sqrt{ax+1}-3\sqrt{ax-1}\sqrt{ax+1}ax+1)(9\operatorname{arccosh}(ax)^3-9\operatorname{arccosh}(ax)^2+6\operatorname{arccosh}(ax)-2)}{216a^4(a^2x^2-1)}$

input `int(x^3*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/216*(-a^2*x^2+1)^(1/2)*(4*a^4*x^4-5*a^2*x^2+4*a^3*x^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)-3*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+1)*(9*\operatorname{arccosh}(a*x)^3-9*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)-2)/a^4/(a^2*x^2-1)-3/8*(-a^2*x^2+1)^(1/2)*((a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+a^2*x^2-1)*(\operatorname{arccosh}(a*x)^3-3*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)-6)/a^4/(a^2*x^2-1)-3/8*(-a^2*x^2+1)^(1/2)*(a^2*x^2-(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x-1)*(\operatorname{arccosh}(a*x)^3+3*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)+6)/a^4/(a^2*x^2-1)-1/216*(-a^2*x^2+1)^(1/2)*(4*a^4*x^4-5*a^2*x^2-4*a^3*x^3*(a*x-1)^(1/2)*(a*x+1)^(1/2)+3*(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+1)*(9*\operatorname{arccosh}(a*x)^3+9*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)+2)/a^4/(a^2*x^2-1) \end{aligned}$$

$$3.254. \quad \int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$



**3.254.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.84

$$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \frac{9(a^4x^4 + a^2x^2 - 2)\sqrt{-a^2x^2 + 1} \log(ax + \sqrt{a^2x^2 - 1})^3 - 9(a^3x^3 + 6ax)\sqrt{a^2x^2 - 1}\sqrt{-a^2x^2 + 1} \log(ax + \sqrt{a^2x^2 - 1})^2 + 9(a^2x^2 - 2)\sqrt{-a^2x^2 + 1} \log(ax + \sqrt{a^2x^2 - 1}) - 9(a^3x^3 + 6ax)\sqrt{-a^2x^2 + 1} \log(ax + \sqrt{a^2x^2 - 1})}{(a^6x^2 - a^4)}$$

input `integrate(x^3*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `-1/27*(9*(a^4*x^4 + a^2*x^2 - 2)*sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1))^3 - 9*(a^3*x^3 + 6*a*x)*sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1))^2 + 6*(a^4*x^4 + 19*a^2*x^2 - 20)*sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1)) - 2*(a^3*x^3 + 60*a*x)*sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1))/(a^6*x^2 - a^4)`

**3.254.6 Sympy [F]**

$$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^3 \operatorname{acosh}^3(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**3*acosh(a*x)**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**3*acosh(a*x)**3/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.254.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.54

$$\begin{aligned} & \int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx \\ &= -\frac{1}{3} \left( \frac{\sqrt{-a^2x^2+1}x^2}{a^2} + \frac{2\sqrt{-a^2x^2+1}}{a^4} \right) \operatorname{arccosh}(ax)^3 \\ & \quad + \frac{2}{27} a \left( \frac{3 \left( -i\sqrt{a^2x^2-1}x^2 - \frac{20i\sqrt{a^2x^2-1}}{a^2} \right) \operatorname{arccosh}(ax)}{a^3} + \frac{ia^2x^3+60ix}{a^4} \right) \\ & \quad + \frac{(ia^2x^3+6ix) \operatorname{arccosh}(ax)^2}{3a^3} \end{aligned}$$

input `integrate(x^3*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `-1/3*(sqrt(-a^2*x^2 + 1)*x^2/a^2 + 2*sqrt(-a^2*x^2 + 1)/a^4)*arccosh(a*x)^3 + 2/27*a*(3*(-I*sqrt(a^2*x^2 - 1)*x^2 - 20*I*sqrt(a^2*x^2 - 1)/a^2)*arccosh(a*x)/a^3 + (I*a^2*x^3 + 60*I*x)/a^4) + 1/3*(I*a^2*x^3 + 6*I*x)*arccosh(a*x)^2/a^3`

### 3.254.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.254.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^3 \operatorname{acosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

input `int((x^3*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2),x)`output `int((x^3*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2), x)`

### 3.255 $\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

3.255.1 Optimal result . . . . .	2283
3.255.2 Mathematica [A] (verified) . . . . .	2283
3.255.3 Rubi [A] (verified) . . . . .	2284
3.255.4 Maple [A] (verified) . . . . .	2286
3.255.5 Fricas [F] . . . . .	2287
3.255.6 Sympy [F] . . . . .	2287
3.255.7 Maxima [F(-2)] . . . . .	2287
3.255.8 Giac [F] . . . . .	2288
3.255.9 Mupad [F(-1)] . . . . .	2288

#### 3.255.1 Optimal result

Integrand size = 24, antiderivative size = 188

$$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = -\frac{3x^2\sqrt{-1+ax}}{8a\sqrt{1-ax}} - \frac{3x\sqrt{1-ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{4a^2} + \frac{3\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{8a^3\sqrt{1-ax}} - \frac{3x^2\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{4a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2a^2} + \frac{\sqrt{-1+ax}\operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}}$$

output

```
-3/8*x^2*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)+3/8*arccosh(a*x)^2*(a*x-1)^(1/2)/a
^3/(-a*x+1)^(1/2)-3/4*x^2*arccosh(a*x)^2*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)+1/
8*arccosh(a*x)^4*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)-3/4*x*arccosh(a*x)*(-a*x
+1)^(1/2)*(a*x+1)^(1/2)/a^2-1/2*x*arccosh(a*x)^3*(-a^2*x^2+1)^(1/2)/a^2
```

#### 3.255.2 Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.52

$$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-((-1+ax)(1+ax))}(-3(1+2\operatorname{arccosh}(ax))^2 \cosh(2\operatorname{arccosh}(ax)) + 2\operatorname{arccosh}(ax) (\operatorname{arccosh}(ax))^3 + 16a^3 \sqrt{\frac{-1+ax}{1+ax}} (1+ax))}{16a^3 \sqrt{\frac{-1+ax}{1+ax}} (1+ax)}$$

input `Integrate[(x^2*ArcCosh[a*x]^3)/Sqrt[1 - a^2*x^2], x]`

output 
$$\frac{-1/16 * (\text{Sqrt}[-((-1 + a*x)*(1 + a*x))] * (-3*(1 + 2*ArcCosh[a*x]^2)*Cosh[2*ArcCosh[a*x]] + 2*ArcCosh[a*x]*(ArcCosh[a*x]^3 + (3 + 2*ArcCosh[a*x]^2)*Sinh[2*ArcCosh[a*x]]))}{a^3 * \text{Sqrt}[(-1 + a*x)/(1 + a*x)] * (1 + a*x)}$$

### 3.255.3 Rubi [A] (verified)

Time = 1.15 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.84, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6353, 6298, 6307, 6354, 15, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx \\ & \quad \downarrow \text{6353} \\ & \frac{\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{3\sqrt{ax-1} \int x \operatorname{arccosh}(ax)^2 dx}{2a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \\ & \quad \downarrow \text{6298} \\ & \frac{\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx}{2a^2} - \frac{3\sqrt{ax-1} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2a\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \\ & \quad \downarrow \text{6307} \\ & - \frac{3\sqrt{ax-1} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^2 - a \int \frac{x^2 \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{2a\sqrt{1-ax}} + \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \\ & \quad \downarrow \text{6354} \\ & - \frac{3\sqrt{ax-1} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} - \frac{\int x dx}{2a} + \frac{x\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{2a^2} \right) \right)}{2a\sqrt{1-ax}} + \\ & \quad \frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2a^2} \\ & \quad \downarrow \text{15} \end{aligned}$$

---

3.255.  $\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
& \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\int \frac{\operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} - \frac{x^2}{4a} \right) \right)}{2a\sqrt{1-ax}} + \\
& \frac{\sqrt{ax-1}\operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2a^2} \\
& \quad \downarrow \text{6308} \\
& \frac{\sqrt{ax-1}\operatorname{arccosh}(ax)^4}{8a^3\sqrt{1-ax}} - \frac{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2a^2} - \\
& \frac{3\sqrt{ax-1} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^2 - a \left( \frac{\operatorname{arccosh}(ax)^2}{4a^3} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)}{2a^2} - \frac{x^2}{4a} \right) \right)}{2a\sqrt{1-ax}}
\end{aligned}$$

input `Int[(x^2*ArcCosh[a*x]^3)/Sqrt[1 - a^2*x^2], x]`

output `-1/2*(x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3)/a^2 + (Sqrt[-1 + a*x]*ArcCosh[a*x]^4)/(8*a^3*Sqrt[1 - a*x]) - (3*Sqrt[-1 + a*x]*((x^2*ArcCosh[a*x]^2)/2 - a*(-1/4*x^2/a + (x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x])/(2*a^2) + ArcCosh[a*x]^2/(4*a^3))))/(2*a*Sqrt[1 - a*x])`

### 3.255.3.1 Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_)^(m_.), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]`

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6353 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_)^p)*((d2_) + (e2_.)*(x_)^p), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*((m - 1)/(c^2*(m + 2*p + 1))) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.255.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.36

method	result
default	$-\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^4}{8a^3(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(2a^3x^3-2ax+2a^2x^2\sqrt{ax-1}\sqrt{ax+1}-\sqrt{ax-1}\sqrt{ax+1})(4\operatorname{arccosh}(ax)^3)}{32a^3(a^2x^2-1)}$

input `int(x^2*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

$$3.255. \quad \int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

output 
$$\begin{aligned} & -1/8*(-a^2*x^2+1)^{(1/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/a^3/(a^2*x^2-1)*\operatorname{arccosh}(a*x)^4 \\ & -1/32*(-a^2*x^2+1)^{(1/2)}*(2*a^3*x^3-2*a*x+2*a^2*x^2*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)} \\ & -(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})*(4*\operatorname{arccosh}(a*x)^3-6*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)-3) \\ & /a^3/(a^2*x^2-1)-1/32*(-a^2*x^2+1)^{(1/2)}*(2*a^3*x^3-2*a*x-2*a^2*x^2*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)} \\ & +(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})*(4*\operatorname{arccosh}(a*x)^3+6*\operatorname{arccosh}(a*x)^2+6*\operatorname{arccosh}(a*x)+3) \\ & /a^3/(a^2*x^2-1) \end{aligned}$$

### 3.255.5 Fracas [F]

$$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^2*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^2*arccosh(a*x)^3/(a^2*x^2 - 1), x)`

### 3.255.6 Sympy [F]

$$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{acosh}^3(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x**2*acosh(a*x)**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**2*acosh(a*x)**3/sqrt(-(a*x - 1)*(a*x + 1)), x)`

### 3.255.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \text{Exception raised: RuntimeError}$$



input `integrate(x^2*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

### 3.255.8 Giac [F]

$$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{acosh}(ax)^3}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(x^2*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^2*arccosh(a*x)^3/sqrt(-a^2*x^2 + 1), x)`

### 3.255.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^2 \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x^2 \operatorname{acosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

input `int((x^2*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2),x)`

output `int((x^2*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2), x)`

### 3.256 $\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$

3.256.1 Optimal result . . . . .	2289
3.256.2 Mathematica [A] (verified) . . . . .	2289
3.256.3 Rubi [A] (verified) . . . . .	2290
3.256.4 Maple [A] (verified) . . . . .	2291
3.256.5 Fricas [A] (verification not implemented) . . . . .	2292
3.256.6 Sympy [F] . . . . .	2292
3.256.7 Maxima [C] (verification not implemented) . . . . .	2292
3.256.8 Giac [C] (verification not implemented) . . . . .	2293
3.256.9 Mupad [F(-1)] . . . . .	2293

#### 3.256.1 Optimal result

Integrand size = 22, antiderivative size = 110

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = -\frac{6x\sqrt{-1+ax}}{a\sqrt{1-ax}} - \frac{6\sqrt{1-ax}\sqrt{1+ax}\operatorname{arccosh}(ax)}{a^2} - \frac{3x\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{a^2}$$

```
output -6*x*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)-3*x*arccosh(a*x)^2*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)-6*arccosh(a*x)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a^2-arccosh(a*x)^3*(-a^2*x^2+1)^(1/2)/a^2
```

#### 3.256.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.92

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{1-a^2x^2}(6ax - 6\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax) + 3ax\operatorname{arccosh}(ax)^2 - \sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax))}{a^2\sqrt{-1+ax}\sqrt{1+ax}}$$

```
input Integrate[(x*ArcCosh[a*x]^3)/Sqrt[1 - a^2*x^2], x]
```

output  $(\text{Sqrt}[1 - a^2x^2]*(6ax - 6\text{Sqrt}[-1 + ax]*\text{Sqrt}[1 + ax]*\text{ArcCosh}[ax] + 3ax*\text{ArcCosh}[ax]^2 - \text{Sqrt}[-1 + ax]*\text{Sqrt}[1 + ax]*\text{ArcCosh}[ax]^3))/(a^2*\text{Sqrt}[-1 + ax]*\text{Sqrt}[1 + ax])$

### 3.256.3 Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.86, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {6329, 6294, 6330, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1 - a^2x^2}} dx$$

↓ 6329

$$-\frac{3\sqrt{ax-1} \int \operatorname{arccosh}(ax)^2 dx}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{a^2}$$

↓ 6294

$$-\frac{3\sqrt{ax-1} \left( x \operatorname{arccosh}(ax)^2 - 2a \int \frac{x \operatorname{arccosh}(ax)}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{a^2}$$

↓ 6330

$$-\frac{3\sqrt{ax-1} \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{\int 1 dx}{a} \right) \right)}{a\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{a^2}$$

↓ 24

$$-\frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{a^2} - \frac{3\sqrt{ax-1} \left( x \operatorname{arccosh}(ax)^2 - 2a \left( \frac{\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}{a^2} - \frac{x}{a} \right) \right)}{a\sqrt{1-ax}}$$

input  $\text{Int}[(x*\text{ArcCosh}[a*x]^3)/\text{Sqrt}[1 - a^2*x^2], x]$

output  $-((\text{Sqrt}[1 - a^2x^2]*\text{ArcCosh}[a*x]^3)/a^2) - (3\text{Sqrt}[-1 + ax]*(x*\text{ArcCosh}[a*x]^2 - 2a*(-(x/a) + (\text{Sqrt}[-1 + ax]*\text{Sqrt}[1 + ax]*\text{ArcCosh}[a*x])/a^2)))/(a*\text{Sqrt}[1 - a*x])$

### 3.256.3.1 Defintions of rubi rules used

- rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`
- rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`
- rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`
- rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

### 3.256.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.41

method	result
default	$-\frac{\sqrt{-a^2x^2+1}(\sqrt{ax-1}\sqrt{ax+1}ax+a^2x^2-1)(\operatorname{arccosh}(ax)^3-3\operatorname{arccosh}(ax)^2+6\operatorname{arccosh}(ax)-6)}{2a^2(a^2x^2-1)} - \frac{\sqrt{-a^2x^2+1}(a^2x^2-\sqrt{ax-1}\sqrt{ax+1})}{2a^2(a^2x^2-1)}$

input `int(x*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/2*(-a^2*x^2+1)^(1/2)*((a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x+a^2*x^2-1)*(arccosh(a*x)^3-3*arccosh(a*x)^2+6*arccosh(a*x)-6)/a^2/(a^2*x^2-1)-1/2*(-a^2*x^2+1)^(1/2)*(a^2*x^2-(a*x-1)^(1/2)*(a*x+1)^(1/2)*a*x-1)*(arccosh(a*x)^3+3*arccosh(a*x)^2+6*arccosh(a*x)+6)/a^2/(a^2*x^2-1)`

3.256. 
$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

**3.256.5 Fricas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.45

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \frac{3\sqrt{a^2x^2-1}\sqrt{-a^2x^2+1}ax \log(ax + \sqrt{a^2x^2-1})^2 + (-a^2x^2+1)^{\frac{3}{2}} \log(ax + \sqrt{a^2x^2-1})^3 + 6\sqrt{a^2x^2-1}}{a^4x^2 - a^2}$$

input `integrate(x*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `(3*sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)*a*x*log(a*x + sqrt(a^2*x^2 - 1))^2 + (-a^2*x^2 + 1)^(3/2)*log(a*x + sqrt(a^2*x^2 - 1))^3 + 6*sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)*a*x - 6*(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1)))/(a^4*x^2 - a^2)`

**3.256.6 Sympy [F]**

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x \operatorname{acosh}^3(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(x*acosh(a*x)**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x*acosh(a*x)**3/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.256.7 Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.59

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \frac{3ix \operatorname{arccosh}(ax)^2}{a} - \frac{\sqrt{-a^2x^2+1} \operatorname{arccosh}(ax)^3}{a^2} + \frac{6 \left( ix - \frac{i\sqrt{a^2x^2-1} \operatorname{arccosh}(ax)}{a} \right)}{a}$$

input `integrate(x*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `3*I*x*arccosh(a*x)^2/a - sqrt(-a^2*x^2 + 1)*arccosh(a*x)^3/a^2 + 6*(I*x - I*sqrt(a^2*x^2 - 1)*arccosh(a*x)/a)/a`

### 3.256.8 Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.94

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = -\frac{\sqrt{-a^2x^2+1} \log(ax + \sqrt{a^2x^2-1})^3}{a^2} - \frac{3i \left( x \log(ax + \sqrt{a^2x^2-1})^2 + 2a \left( \frac{x}{a} - \frac{\sqrt{a^2x^2-1} \log(ax + \sqrt{a^2x^2-1})}{a^2} \right) \right)}{a}$$

input `integrate(x*arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `-sqrt(-a^2*x^2 + 1)*log(a*x + sqrt(a^2*x^2 - 1))^3/a^2 - 3*I*(x*log(a*x + sqrt(a^2*x^2 - 1))^2 + 2*a*(x/a - sqrt(a^2*x^2 - 1)*log(a*x + sqrt(a^2*x^2 - 1))/a^2))/a`

### 3.256.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x \operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{x \operatorname{acosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

input `int((x*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2),x)`

output `int((x*acosh(a*x)^3)/(1 - a^2*x^2)^(1/2), x)`

$$3.257 \quad \int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

3.257.1 Optimal result	2294
3.257.2 Mathematica [A] (verified)	2294
3.257.3 Rubi [A] (verified)	2295
3.257.4 Maple [A] (verified)	2295
3.257.5 Fricas [F]	2296
3.257.6 Sympy [F]	2296
3.257.7 Maxima [F]	2296
3.257.8 Giac [F]	2297
3.257.9 Mupad [F(-1)]	2297

### 3.257.1 Optimal result

Integrand size = 21, antiderivative size = 32

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-1+ax}\operatorname{arccosh}(ax)^4}{4a\sqrt{1-ax}}$$

output  $1/4*\operatorname{arccosh}(a*x)^4*(a*x-1)^{(1/2)}/a/(-a*x+1)^{(1/2)}$

### 3.257.2 Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.41

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \frac{\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^4}{4a\sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]^3/Sqrt[1 - a^2*x^2],x]`

output  $(\operatorname{Sqrt}[-1+a*x]*\operatorname{Sqrt}[1+a*x]*\operatorname{ArcCosh}[a*x]^4)/(4*a*\operatorname{Sqrt}[1-a^2*x^2])$

### 3.257.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

↓ 6307

$$\frac{\sqrt{ax-1} \operatorname{arccosh}(ax)^4}{4a\sqrt{1-ax}}$$

input `Int[ArcCosh[a*x]^3/Sqrt[1 - a^2*x^2], x]`

output `(Sqrt[-1 + a*x]*ArcCosh[a*x]^4)/(4*a*Sqrt[1 - a*x])`

#### 3.257.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.257.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.59

method	result	size
default	$-\frac{\sqrt{-(ax-1)(ax+1)}\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^4}{4(a^2x^2-1)a}$	51

input `int(arccosh(a*x)^3/(-a^2*x^2+1)^(1/2), x, method=_RETURNVERBOSE)`

output `-1/4*(-(a*x-1)*(a*x+1))^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2-1)/a*arccosh(a*x)^4`



**3.257.5 Fracas [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^3/(a^2*x^2 - 1), x)`

**3.257.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}^3(ax)}{\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**3/sqrt(-(a*x - 1)*(a*x + 1)), x)`

**3.257.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^3/sqrt(-a^2*x^2 + 1), x)`

**3.257.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}} dx$$

input `integrate(arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^3/sqrt(-a^2*x^2 + 1), x)`

**3.257.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^3/(1 - a^2*x^2)^(1/2),x)`

output `int(acosh(a*x)^3/(1 - a^2*x^2)^(1/2), x)`

**3.258**       $\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx$

3.258.1 Optimal result . . . . .	2298
3.258.2 Mathematica [A] (verified) . . . . .	2299
3.258.3 Rubi [A] (verified) . . . . .	2300
3.258.4 Maple [F] . . . . .	2302
3.258.5 Fracas [F] . . . . .	2302
3.258.6 Sympy [F] . . . . .	2303
3.258.7 Maxima [F] . . . . .	2303
3.258.8 Giac [F] . . . . .	2303
3.258.9 Mupad [F(-1)] . . . . .	2304

**3.258.1 Optimal result**

Integrand size = 24, antiderivative size = 265

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx = \frac{2\sqrt{-1+ax}\operatorname{arccosh}(ax)^3 \arctan(e^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{3i\sqrt{-1+ax}\operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} + \frac{3i\sqrt{-1+ax}\operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} + \frac{6i\sqrt{-1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{6i\sqrt{-1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{6i\sqrt{-1+ax} \operatorname{PolyLog}(4, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} + \frac{6i\sqrt{-1+ax} \operatorname{PolyLog}(4, ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}}$$

output  $2*\operatorname{arccosh}(a*x)^3*\arctan(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*(a*x-1)^{(1/2)})/(-a*x+1)^{(1/2)}-3*I*\operatorname{arccosh}(a*x)^2*\operatorname{polylog}(2,-I*(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}))*(a*x-1)^{(1/2)})/(-a*x+1)^{(1/2)}+3*I*\operatorname{arccosh}(a*x)^2*\operatorname{polylog}(2,I*(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}))*(a*x-1)^{(1/2)})/(-a*x+1)^{(1/2)}+6*I*\operatorname{arccosh}(a*x)*\operatorname{polylog}(3,-I*(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}))*(a*x-1)^{(1/2)})/(-a*x+1)^{(1/2)}-6*I*\operatorname{arccosh}(a*x)*\operatorname{polylog}(3,I*(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}))*(a*x-1)^{(1/2)})/(-a*x+1)^{(1/2)}-6*I*\operatorname{polylog}(4,-I*(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}))*(a*x-1)^{(1/2)})/(-a*x+1)^{(1/2)}+6*I*\operatorname{polylog}(4,I*(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}))*(a*x-1)^{(1/2)})/(-a*x+1)^{(1/2)}$

### 3.258.2 Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 488, normalized size of antiderivative = 1.84

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx$$

$$= \frac{i\sqrt{-((-1+ax)(1+ax))}(7\pi^4 + 8i\pi^3\operatorname{arccosh}(ax) + 24\pi^2\operatorname{arccosh}(ax)^2 - 32i\pi\operatorname{arccosh}(ax)^3 - 16\operatorname{arccosh}(ax)^4)}{x\sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]^3/(x*Sqrt[1 - a^2*x^2]),x]`

output  $((I/64)*\operatorname{Sqrt}[(-(-1+a*x)*(1+a*x))]*(7*\operatorname{Pi}^4 + (8*I)*\operatorname{Pi}^3*\operatorname{ArcCosh}[a*x] + 24*\operatorname{Pi}^2*\operatorname{ArcCosh}[a*x]^2 - (32*I)*\operatorname{Pi}*\operatorname{ArcCosh}[a*x]^3 - 16*\operatorname{ArcCosh}[a*x]^4 + (8*I)*\operatorname{Pi}^3*\operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[a*x]}] + 48*\operatorname{Pi}^2*\operatorname{ArcCosh}[a*x]*\operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[a*x]}] - (96*I)*\operatorname{Pi}*\operatorname{ArcCosh}[a*x]^2*\operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[a*x]}] - 64*\operatorname{ArcCosh}[a*x]^3*\operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[a*x]}] - 48*\operatorname{Pi}^2*\operatorname{ArcCosh}[a*x]*\operatorname{Log}[1 - I/E^{\operatorname{ArcCosh}[a*x]}] + (96*I)*\operatorname{Pi}*\operatorname{ArcCosh}[a*x]^2*\operatorname{Log}[1 - I/E^{\operatorname{ArcCosh}[a*x]}] - (8*I)*\operatorname{Pi}^3*\operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[a*x]}] + 64*\operatorname{ArcCosh}[a*x]^3*\operatorname{Log}[1 + I/E^{\operatorname{ArcCosh}[a*x]}] + (8*I)*\operatorname{Pi}^3*\operatorname{Log}[\operatorname{Tan}[(\operatorname{Pi} + (2*I)*\operatorname{ArcCosh}[a*x])/4]] - 48*(\operatorname{Pi} - (2*I)*\operatorname{ArcCosh}[a*x])^2*\operatorname{PolyLog}[2, (-I)/E^{\operatorname{ArcCosh}[a*x]}] + 192*\operatorname{ArcCosh}[a*x]^2*\operatorname{PolyLog}[2, (-I)*E^{\operatorname{ArcCosh}[a*x]}] - 48*\operatorname{Pi}^2*\operatorname{PolyLog}[2, I/E^{\operatorname{ArcCosh}[a*x]}] + (192*I)*\operatorname{Pi}*\operatorname{ArcCosh}[a*x]*\operatorname{PolyLog}[2, I/E^{\operatorname{ArcCosh}[a*x]}] + (192*I)*\operatorname{Pi}*\operatorname{PolyLog}[3, (-I)/E^{\operatorname{ArcCosh}[a*x]}] + 384*\operatorname{ArcCosh}[a*x]*\operatorname{PolyLog}[3, (-I)/E^{\operatorname{ArcCosh}[a*x]}] - 384*\operatorname{ArcCosh}[a*x]*\operatorname{PolyLog}[3, (-I)*E^{\operatorname{ArcCosh}[a*x]}] - (192*I)*\operatorname{Pi}*\operatorname{PolyLog}[3, I/E^{\operatorname{ArcCosh}[a*x]}] + 384*\operatorname{PolyLog}[4, (-I)/E^{\operatorname{ArcCosh}[a*x]}] + 384*\operatorname{PolyLog}[4, (-I)*E^{\operatorname{ArcCosh}[a*x]}]))/(\operatorname{Sqrt}[(-1+a*x)/(1+a*x)]*(1+a*x))$

**3.258.3 Rubi [A] (verified)**

Time = 0.68 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.58, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$ , Rules used = {6361, 3042, 4668, 3011, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx$$

$$\downarrow \text{6361}$$

$$\frac{\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)^3}{ax} d\operatorname{arccosh}(ax)}{\sqrt{1-ax}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{ax-1} \int \operatorname{arccosh}(ax)^3 \csc\left(i\operatorname{arccosh}(ax) + \frac{\pi}{2}\right) d\operatorname{arccosh}(ax)}{\sqrt{1-ax}}$$

$$\downarrow \text{4668}$$

$$\frac{\sqrt{ax-1}(-3i \int \operatorname{arccosh}(ax)^2 \log(1-ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 3i \int \operatorname{arccosh}(ax)^2 \log(1+ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax))}{\sqrt{1-ax}}$$

$$\downarrow \text{3011}$$

$$\frac{\sqrt{ax-1}(3i(2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}))}{\sqrt{1-ax}}$$

$$\downarrow \text{7163}$$

$$\frac{\sqrt{ax-1}(3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)}) - \int \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax)) - \operatorname{arccosh}(ax)^3 \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}}$$

$$\downarrow \text{2720}$$

$$\frac{\sqrt{ax-1}(3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)}) - \int e^{-\operatorname{arccosh}(ax)} \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)}) de^{\operatorname{arccosh}(ax)}) - \operatorname{arccosh}(ax)^3 \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}}$$

$$\downarrow \text{7143}$$

$$\frac{\sqrt{ax-1}(2\operatorname{arccosh}(ax)^3 \arctan(e^{\operatorname{arccosh}(ax)}) + 3i(2(\operatorname{arccosh}(ax) \operatorname{PolyLog}(3, -ie^{\operatorname{arccosh}(ax)}) - \operatorname{PolyLog}(4, -ie^{\operatorname{arccosh}(ax)})$$

---

3.258.  $\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx$

input `Int[ArcCosh[a*x]^3/(x*Sqrt[1 - a^2*x^2]),x]`

output `(Sqrt[-1 + a*x]*(2*ArcCosh[a*x]^3*ArcTan[E^ArcCosh[a*x]] + (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, (-I)*E^ArcCosh[a*x]]) + 2*(ArcCosh[a*x]*PolyLog[3, (-I)*E^ArcCosh[a*x]] - PolyLog[4, (-I)*E^ArcCosh[a*x]])) - (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, I*E^ArcCosh[a*x]]) + 2*(ArcCosh[a*x]*PolyLog[3, I*E^ArcCosh[a*x]] - PolyLog[4, I*E^ArcCosh[a*x]]))))/Sqrt[1 - a*x]`

### 3.258.3.1 Defintions of rubi rules used

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_)^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

```
rule 6361 Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/Sqrt[(d_) + (e_.)
*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]
]/Sqrt[d + e*x^2])] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]]
, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && Int
egerQ[m]
```

```
rule 7143 Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

```
rule 7163 Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)
)*(x_)))]^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)
^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c
, d, e, f, n, p}, x] && GtQ[m, 0]
```

### 3.258.4 Maple [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{-a^2x^2+1}} dx$$

```
input int(arccosh(a*x)^3/x/(-a^2*x^2+1)^(1/2),x)
```

```
output int(arccosh(a*x)^3/x/(-a^2*x^2+1)^(1/2),x)
```

### 3.258.5 Fracas [F]

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arccosh}(ax)^3}{\sqrt{-a^2x^2+1}x} dx$$

```
input integrate(arccosh(a*x)^3/x/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^3/(a^2*x^3 - x), x)
```

**3.258.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}^3(ax)}{x\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**3/x/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**3/(x*sqrt(-(a*x - 1)*(a*x + 1))), x)`

**3.258.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}x} dx$$

input `integrate(arccosh(a*x)^3/x/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^3/(sqrt(-a^2*x^2 + 1)*x), x)`

**3.258.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}x} dx$$

input `integrate(arccosh(a*x)^3/x/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^3/(sqrt(-a^2*x^2 + 1)*x), x)`



**3.258.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^3/(x*(1 - a^2*x^2)^(1/2)),x)`output `int(acosh(a*x)^3/(x*(1 - a^2*x^2)^(1/2)), x)`

### 3.259 $\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx$

3.259.1 Optimal result . . . . .	2305
3.259.2 Mathematica [A] (verified) . . . . .	2306
3.259.3 Rubi [C] (verified) . . . . .	2306
3.259.4 Maple [A] (verified) . . . . .	2309
3.259.5 Fricas [F] . . . . .	2310
3.259.6 Sympy [F] . . . . .	2310
3.259.7 Maxima [F] . . . . .	2310
3.259.8 Giac [F] . . . . .	2311
3.259.9 Mupad [F(-1)] . . . . .	2311

#### 3.259.1 Optimal result

Integrand size = 24, antiderivative size = 166

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx = \frac{a\sqrt{-1+ax}\operatorname{arccosh}(ax)^3}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} - \frac{3a\sqrt{-1+ax}\operatorname{arccosh}(ax)^2 \log(1+e^{2\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} - \frac{3a\sqrt{-1+ax}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} + \frac{3a\sqrt{-1+ax} \operatorname{PolyLog}(3, -e^{2\operatorname{arccosh}(ax)})}{2\sqrt{1-ax}}$$

output

```
a*arccosh(a*x)^3*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-3*a*arccosh(a*x)^2*ln(1+(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))^2)*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-3*a*arccosh(a*x)*polylog(2,-(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))^2)*(a*x-1)^(1/2)/(-a*x+1)^(1/2)+3/2*a*polylog(3,-(a*x+(a*x-1)^(1/2)*(a*x+1)^(1/2))^2)*(a*x-1)^(1/2)/(-a*x+1)^(1/2)-arccosh(a*x)^3*(-a^2*x^2+1)^(1/2)/x
```

**3.259.2 Mathematica [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx$$

$$= \frac{a\sqrt{\frac{-1+ax}{1+ax}}(1+ax) \left( 2\operatorname{arccosh}(ax)^2 \left( -\operatorname{arccosh}(ax) + \frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax)\operatorname{arccosh}(ax)}{ax} - 3\log(1+e^{-2\operatorname{arccosh}(ax)}) \right) \right)}{2\sqrt{-((-1+ax)(1+ax))}}$$

input `Integrate[ArcCosh[a*x]^3/(x^2*Sqrt[1 - a^2*x^2]),x]`output `(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(2*ArcCosh[a*x]^2*(-ArcCosh[a*x] + (Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*ArcCosh[a*x])/(a*x) - 3*Log[1 + E^(-2*ArcCosh[a*x])])) + 6*ArcCosh[a*x]*PolyLog[2, -E^(-2*ArcCosh[a*x])] + 3*PolyLog[3, -E^(-2*ArcCosh[a*x])])/(2*Sqrt[-((-1 + a*x)*(1 + a*x))])`**3.259.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.70 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.75, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {6332, 6297, 3042, 26, 4201, 2620, 3011, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx$$

$$\downarrow \text{6332}$$

$$-\frac{3a\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)^2}{x} dx}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x}$$

$$\downarrow \text{6297}$$

$$-\frac{3a\sqrt{ax-1} \int \frac{\sqrt{\frac{ax-1}{ax+1}}(ax+1)\operatorname{arccosh}(ax)^2}{ax} d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x}$$

$$\downarrow \text{3042}$$

---

3.259.  $\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx$

$$\begin{aligned}
& -\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} - \frac{3a\sqrt{ax-1} \int -i\operatorname{arccosh}(ax)^2 \tan(i\operatorname{arccosh}(ax))d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} \\
& \quad \downarrow 26 \\
& -\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} + \frac{3ia\sqrt{ax-1} \int \operatorname{arccosh}(ax)^2 \tan(i\operatorname{arccosh}(ax))d\operatorname{arccosh}(ax)}{\sqrt{1-ax}} \\
& \quad \downarrow 4201 \\
& \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} + 3ia\sqrt{ax-1} \left( 2i \int \frac{e^{2\operatorname{arccosh}(ax)}\operatorname{arccosh}(ax)^2}{1+e^{2\operatorname{arccosh}(ax)}}d\operatorname{arccosh}(ax) - \frac{1}{3}i\operatorname{arccosh}(ax)^3 \right)}{\sqrt{1-ax}} \\
& \quad \downarrow 2620 \\
& \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} + 3ia\sqrt{ax-1} \left( 2i \left( \frac{1}{2}\operatorname{arccosh}(ax)^2 \log(e^{2\operatorname{arccosh}(ax)}+1) - \int \operatorname{arccosh}(ax) \log(1+e^{2\operatorname{arccosh}(ax)})d\operatorname{arccosh}(ax) \right) - \frac{1}{3}i\operatorname{arccosh}(ax)^3 \right)}{\sqrt{1-ax}} \\
& \quad \downarrow 3011 \\
& \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} + 3ia\sqrt{ax-1} \left( 2i \left( -\frac{1}{2} \int \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(ax)})d\operatorname{arccosh}(ax) + \frac{1}{2}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(ax)}) + \frac{1}{2}\operatorname{arccosh}(ax)^2 \log(e^{2\operatorname{arccosh}(ax)}+1) \right) \right)}{\sqrt{1-ax}} \\
& \quad \downarrow 2720 \\
& \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} + 3ia\sqrt{ax-1} \left( 2i \left( -\frac{1}{4} \int e^{-2\operatorname{arccosh}(ax)} \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(ax)})de^{2\operatorname{arccosh}(ax)} + \frac{1}{2}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(ax)}) + \frac{1}{2}\operatorname{arccosh}(ax)^2 \log(e^{2\operatorname{arccosh}(ax)}+1) \right) \right)}{\sqrt{1-ax}} \\
& \quad \downarrow 7143 \\
& \frac{-\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{x} + 3ia\sqrt{ax-1} \left( 2i \left( \frac{1}{2}\operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -e^{2\operatorname{arccosh}(ax)}) - \frac{1}{4}\operatorname{PolyLog}(3, -e^{2\operatorname{arccosh}(ax)}) + \frac{1}{2}\operatorname{arccosh}(ax)^2 \log(e^{2\operatorname{arccosh}(ax)}+1) \right) \right)}{\sqrt{1-ax}}
\end{aligned}$$

input `Int[ArcCosh[a*x]^3/(x^2*Sqrt[1 - a^2*x^2]),x]`

```
output -((Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3)/x) + ((3*I)*a*Sqrt[-1 + a*x]*((-1/3*I)
)*ArcCosh[a*x]^3 + (2*I)*((ArcCosh[a*x]^2*Log[1 + E^(2*ArcCosh[a*x])])/2 +
(ArcCosh[a*x]*PolyLog[2, -E^(2*ArcCosh[a*x])])/2 - PolyLog[3, -E^(2*ArcCo
sh[a*x])])/4))/Sqrt[1 - a*x]
```

### 3.259.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 2620 Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_)^(m_))/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Si
mp[d*(m/(b*f*g*n*Log[F])) Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x
)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

```
rule 2720 Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

```
rule 3011 Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_)^(m_)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 4201 Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + (Complex[0, fz_]*(f_)*(x_)]], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Simp[2*I Int[
(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

rule 6297 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(x_), x_Symbol] := Simp[1/b  
Subst[Int[x^n*Tanh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a  
, b, c}, x] && IGtQ[n, 0]`

rule 6332 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.  
)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a +  
b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d +  
e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2  
)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b,  
c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3  
, 0] && NeQ[m, -1]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S  
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d  
, e, n, p}, x] && EqQ[b*d, a*e]`

### 3.259.4 Maple [A] (verified)

Time = 1.05 (sec) , antiderivative size = 313, normalized size of antiderivative = 1.89

method	result
default	$-\frac{\sqrt{-a^2x^2+1} (a^2x^2-\sqrt{ax-1}\sqrt{ax+1}ax-1) \operatorname{arccosh}(ax)^3}{x(a^2x^2-1)} - \frac{2\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)^3a}{a^2x^2-1} + \frac{3\sqrt{-a^2x^2+1}\sqrt{ax-1}}{a^2x^2-1}$

input `int(arccosh(a*x)^3/x^2/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -(-a^2x^2+1)^{(1/2)}*(a^2x^2-(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*a*x-1)*\operatorname{arccosh}(a*x)^3/x/(a^2x^2-1)-2*(-a^2x^2+1)^{(1/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/(a^2x^2-1)*\operatorname{arccosh}(a*x)^3*a+3*(-a^2x^2+1)^{(1/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/(a^2x^2-1)*\operatorname{arccosh}(a*x)^2*\ln(1+(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})^2)*a+3*(-a^2x^2+1)^{(1/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/(a^2x^2-1)*\operatorname{arccosh}(a*x)*\operatorname{polylog}(2,-(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})^2)*a-3/2*(-a^2x^2+1)^{(1/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/(a^2x^2-1)*\operatorname{polylog}(3,-(a*x+(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)})^2)*a \end{aligned}$$

**3.259.5 Fracas [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}x^2} dx$$

input `integrate(arccosh(a*x)^3/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^3/(a^2*x^4 - x^2), x)`

**3.259.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}^3(ax)}{x^2\sqrt{-(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**3/x**2/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**3/(x**2*sqrt(-(a*x - 1)*(a*x + 1))), x)`

**3.259.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}x^2} dx$$

input `integrate(arccosh(a*x)^3/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `(a^2*x^2 - 1)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^3/(sqrt(a*x + 1)*sqrt(-a*x + 1)*x) - integrate(3*(a^3*x^2 + sqrt(a*x + 1)*sqrt(a*x - 1)*a^2*x - a)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))^2/((sqrt(a*x + 1)*a*x^2 + (a*x + 1)*sqrt(a*x - 1)*x)*sqrt(-a*x + 1)), x)`

**3.259.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}x^2} dx$$

input `integrate(arccosh(a*x)^3/x^2/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^3/(sqrt(-a^2*x^2 + 1)*x^2), x)`

**3.259.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{x^2\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^3/(x^2*(1 - a^2*x^2)^(1/2)),x)`

output `int(acosh(a*x)^3/(x^2*(1 - a^2*x^2)^(1/2)), x)`



### 3.260 $\int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx$

3.260.1 Optimal result	2312
3.260.2 Mathematica [B] (warning: unable to verify)	2313
3.260.3 Rubi [A] (verified)	2314
3.260.4 Maple [F]	2320
3.260.5 Fricas [F]	2320
3.260.6 Sympy [F]	2320
3.260.7 Maxima [F]	2321
3.260.8 Giac [F]	2321
3.260.9 Mupad [F(-1)]	2321

#### 3.260.1 Optimal result

Integrand size = 24, antiderivative size = 460

$$\begin{aligned}
 \int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx = & \frac{3a\sqrt{-1+ax}\operatorname{arccosh}(ax)^2}{2x\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2} \\
 & - \frac{6a^2\sqrt{-1+ax}\operatorname{arccosh}(ax)\arctan(e^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\
 & + \frac{a^2\sqrt{-1+ax}\operatorname{arccosh}(ax)^3\arctan(e^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\
 & + \frac{3ia^2\sqrt{-1+ax}\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\
 & - \frac{3ia^2\sqrt{-1+ax}\operatorname{arccosh}(ax)^2\operatorname{PolyLog}(2,-ie^{\operatorname{arccosh}(ax)})}{2\sqrt{1-ax}} \\
 & - \frac{3ia^2\sqrt{-1+ax}\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\
 & + \frac{3ia^2\sqrt{-1+ax}\operatorname{arccosh}(ax)^2\operatorname{PolyLog}(2,ie^{\operatorname{arccosh}(ax)})}{2\sqrt{1-ax}} \\
 & + \frac{3ia^2\sqrt{-1+ax}\operatorname{arccosh}(ax)\operatorname{PolyLog}(3,-ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\
 & - \frac{3ia^2\sqrt{-1+ax}\operatorname{arccosh}(ax)\operatorname{PolyLog}(3,ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\
 & - \frac{3ia^2\sqrt{-1+ax}\operatorname{PolyLog}(4,-ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}} \\
 & + \frac{3ia^2\sqrt{-1+ax}\operatorname{PolyLog}(4,ie^{\operatorname{arccosh}(ax)})}{\sqrt{1-ax}}
 \end{aligned}$$

output  $\frac{3}{2}a \operatorname{arccosh}(ax)^2 (ax-1)^{1/2} / x / (-ax+1)^{1/2} - 6a^2 \operatorname{arccosh}(ax) \arctan(ax+(ax-1)^{1/2} (ax+1)^{1/2}) (ax-1)^{1/2} / (-ax+1)^{1/2} + a^2 \operatorname{arccosh}(ax)^3 \arctan(ax+(ax-1)^{1/2} (ax+1)^{1/2}) (ax-1)^{1/2} / (-ax+1)^{1/2} + 3Ia^2 \operatorname{polylog}(2, -I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} - 3/2 Ia^2 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2, -I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} - 3Ia^2 \operatorname{polylog}(2, I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} + 3/2 Ia^2 \operatorname{arccosh}(ax)^2 \operatorname{polylog}(2, I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} + 3Ia^2 \operatorname{arccosh}(ax) \operatorname{polylog}(3, -I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} - 3Ia^2 \operatorname{arccosh}(ax) \operatorname{polylog}(3, I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} - 3Ia^2 \operatorname{polylog}(4, -I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} + 3Ia^2 \operatorname{polylog}(4, I(ax+(ax-1)^{1/2} (ax+1)^{1/2})) (ax-1)^{1/2} / (-ax+1)^{1/2} - 1/2 \operatorname{arccosh}(ax)^3 (-a^2x^2+1)^{1/2} / x^2$

### 3.260.2 Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1051 vs.  $2(460) = 920$ .

Time = 4.23 (sec) , antiderivative size = 1051, normalized size of antiderivative = 2.28

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^3 \sqrt{1-a^2x^2}} dx = \frac{ia^2(1+ax) \left( 7\pi^4 \sqrt{\frac{-1+ax}{1+ax}} + 8i\pi^3 \sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax) + 24\pi^2 \sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)^2 + \frac{192i \sqrt{\frac{-1+ax}{1+ax}} \operatorname{arccosh}(ax)}{ax} \right)}{x^3 \sqrt{1-a^2x^2}}$$

input `Integrate[ArcCosh[a*x]^3/(x^3*Sqrt[1 - a^2*x^2]),x]`

output

```

((-1/128*I)*a^2*(1 + a*x)*(7*Pi^4*Sqrt[(-1 + a*x)/(1 + a*x)] + (8*I)*Pi^3*
Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x] + 24*Pi^2*Sqrt[(-1 + a*x)/(1 + a*x
)]*ArcCosh[a*x]^2 + ((192*I)*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]^2)/(a
*x) + ((64*I)*(-1 + a*x)*ArcCosh[a*x]^3)/(a^2*x^2) - (32*I)*Pi*Sqrt[(-1 +
a*x)/(1 + a*x)]*ArcCosh[a*x]^3 - 16*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x
]^4 - 384*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]*Log[1 - I/E^ArcCosh[a*x]
] + (8*I)*Pi^3*Sqrt[(-1 + a*x)/(1 + a*x)]*Log[1 + I/E^ArcCosh[a*x]] + 384*
Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]*Log[1 + I/E^ArcCosh[a*x]] + 48*Pi^
2*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]*Log[1 + I/E^ArcCosh[a*x]] - (96*
I)*Pi*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]^2*Log[1 + I/E^ArcCosh[a*x]]
- 64*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]^3*Log[1 + I/E^ArcCosh[a*x]] -
48*Pi^2*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]*Log[1 - I*E^ArcCosh[a*x]]
+ (96*I)*Pi*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]^2*Log[1 - I*E^ArcCosh
[a*x]] - (8*I)*Pi^3*Sqrt[(-1 + a*x)/(1 + a*x)]*Log[1 + I*E^ArcCosh[a*x]] +
64*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]^3*Log[1 + I*E^ArcCosh[a*x]] +
(8*I)*Pi^3*Sqrt[(-1 + a*x)/(1 + a*x)]*Log[Tan[(Pi + (2*I)*ArcCosh[a*x])/4]
] - 48*Sqrt[(-1 + a*x)/(1 + a*x)]*(8 + Pi^2 - (4*I)*Pi*ArcCosh[a*x] - 4*Ar
cCosh[a*x]^2)*PolyLog[2, (-I)/E^ArcCosh[a*x]] + 384*Sqrt[(-1 + a*x)/(1 + a
*x)]*PolyLog[2, I/E^ArcCosh[a*x]] + 192*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh
[a*x]^2*PolyLog[2, (-I)*E^ArcCosh[a*x]] - 48*Pi^2*Sqrt[(-1 + a*x)/(1 + ...

```

### 3.260.3 Rubi [A] (verified)

Time = 2.50 (sec) , antiderivative size = 273, normalized size of antiderivative = 0.59, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {6347, 6298, 6361, 3042, 4668, 3011, 6362, 3042, 4668, 2715, 2838, 7163, 2720, 7143}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\operatorname{arccosh}(ax)^3}{x^3 \sqrt{1-a^2x^2}} dx \\
 & \quad \downarrow \text{6347} \\
 & \frac{1}{2} a^2 \int \frac{\operatorname{arccosh}(ax)^3}{x \sqrt{1-a^2x^2}} dx - \frac{3a\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)^2}{x^2} dx}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)^3}{2x^2} \\
 & \quad \downarrow \text{6298}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{2}a^2 \int \frac{\operatorname{arccosh}(ax)^3}{x\sqrt{1-a^2x^2}} dx - \frac{3a\sqrt{ax-1} \left( 2a \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\operatorname{arccosh}(ax)^2}{x} \right)}{2\sqrt{1-ax}} - \\
& \qquad \qquad \qquad \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{6361} \\
& \frac{a^2\sqrt{ax-1} \int \frac{\operatorname{arccosh}(ax)^3}{ax} d\operatorname{arccosh}(ax)}{2\sqrt{1-ax}} - \frac{3a\sqrt{ax-1} \left( 2a \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\operatorname{arccosh}(ax)^2}{x} \right)}{2\sqrt{1-ax}} - \\
& \qquad \qquad \qquad \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{3042} \\
& \frac{a^2\sqrt{ax-1} \int \operatorname{arccosh}(ax)^3 \csc \left( i\operatorname{arccosh}(ax) + \frac{\pi}{2} \right) d\operatorname{arccosh}(ax)}{2\sqrt{1-ax}} - \\
& \frac{3a\sqrt{ax-1} \left( 2a \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\operatorname{arccosh}(ax)^2}{x} \right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{4668} \\
& \frac{a^2\sqrt{ax-1} \left( -3i \int \operatorname{arccosh}(ax)^2 \log(1 - ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) + 3i \int \operatorname{arccosh}(ax)^2 \log(1 + ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) \right)}{2\sqrt{1-ax}} - \\
& \frac{3a\sqrt{ax-1} \left( 2a \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\operatorname{arccosh}(ax)^2}{x} \right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{3011} \\
& \frac{a^2\sqrt{ax-1} \left( 3i \left( 2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) \right) \right)}{2\sqrt{1-ax}} - \\
& \frac{3a\sqrt{ax-1} \left( 2a \int \frac{\operatorname{arccosh}(ax)}{x\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\operatorname{arccosh}(ax)^2}{x} \right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{6362} \\
& \frac{a^2\sqrt{ax-1} \left( 3i \left( 2 \int \operatorname{arccosh}(ax) \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) d\operatorname{arccosh}(ax) - \operatorname{arccosh}(ax)^2 \operatorname{PolyLog}(2, -ie^{\operatorname{arccosh}(ax)}) \right) \right)}{2\sqrt{1-ax}} - \\
& \frac{3a\sqrt{ax-1} \left( 2a \int \frac{\operatorname{arccosh}(ax)}{ax} d\operatorname{arccosh}(ax) - \frac{\operatorname{arccosh}(ax)^2}{x} \right)}{2\sqrt{1-ax}} - \frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2} \\
& \qquad \qquad \qquad \downarrow \text{3042}
\end{aligned}$$

---

3.260.  $\int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx$

$$\frac{a^2\sqrt{ax-1}\left(3i\left(2\int\operatorname{arccosh}(ax)\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\operatorname{darccosh}(ax)-\operatorname{arccosh}(ax)^2\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\right)\right.}{3a\sqrt{ax-1}\left(-\frac{\operatorname{arccosh}(ax)^2}{x}+2a\int\operatorname{arccosh}(ax)\csc\left(i\operatorname{arccosh}(ax)+\frac{\pi}{2}\right)\operatorname{darccosh}(ax)\right)} -$$

$$\frac{\frac{2\sqrt{1-ax}}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}}{2x^2} -$$

↓ 4668

$$\frac{a^2\sqrt{ax-1}\left(3i\left(2\int\operatorname{arccosh}(ax)\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\operatorname{darccosh}(ax)-\operatorname{arccosh}(ax)^2\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\right)\right.}{3a\sqrt{ax-1}\left(-\frac{\operatorname{arccosh}(ax)^2}{x}+2a\left(-i\int\log\left(1-ie^{\operatorname{arccosh}(ax)}\right)\operatorname{darccosh}(ax)+i\int\log\left(1+ie^{\operatorname{arccosh}(ax)}\right)\operatorname{darccosh}(ax)\right)\right)} -$$

$$\frac{\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2}}{2\sqrt{1-ax}} -$$

↓ 2715

$$\frac{a^2\sqrt{ax-1}\left(3i\left(2\int\operatorname{arccosh}(ax)\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\operatorname{darccosh}(ax)-\operatorname{arccosh}(ax)^2\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\right)\right.}{3a\sqrt{ax-1}\left(-\frac{\operatorname{arccosh}(ax)^2}{x}+2a\left(-i\int e^{-\operatorname{arccosh}(ax)}\log\left(1-ie^{\operatorname{arccosh}(ax)}\right)de^{\operatorname{arccosh}(ax)}+i\int e^{-\operatorname{arccosh}(ax)}\log\left(1+ie^{\operatorname{arccosh}(ax)}\right)de^{\operatorname{arccosh}(ax)}\right)\right)} -$$

$$\frac{\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2}}{2\sqrt{1-ax}} -$$

↓ 2838

$$\frac{a^2\sqrt{ax-1}\left(3i\left(2\int\operatorname{arccosh}(ax)\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\operatorname{darccosh}(ax)-\operatorname{arccosh}(ax)^2\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)\right)\right.}{3a\sqrt{ax-1}\left(-\frac{\operatorname{arccosh}(ax)^2}{x}+2a\left(2\operatorname{arccosh}(ax)\arctan\left(e^{\operatorname{arccosh}(ax)}\right)-i\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)+i\operatorname{PolyLog}\left(2,ie^{\operatorname{arccosh}(ax)}\right)\right)\right)} -$$

$$\frac{\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2}}{2\sqrt{1-ax}} -$$

↓ 7163

$$\frac{a^2\sqrt{ax-1}\left(3i\left(2\left(\operatorname{arccosh}(ax)\operatorname{PolyLog}\left(3,-ie^{\operatorname{arccosh}(ax)}\right)-\int\operatorname{PolyLog}\left(3,-ie^{\operatorname{arccosh}(ax)}\right)\operatorname{darccosh}(ax)\right)\right)-\operatorname{arccosh}(ax)^2\operatorname{PolyLog}\left(3,-ie^{\operatorname{arccosh}(ax)}\right)\right)}{3a\sqrt{ax-1}\left(-\frac{\operatorname{arccosh}(ax)^2}{x}+2a\left(2\operatorname{arccosh}(ax)\arctan\left(e^{\operatorname{arccosh}(ax)}\right)-i\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)+i\operatorname{PolyLog}\left(2,ie^{\operatorname{arccosh}(ax)}\right)\right)\right)} -$$

$$\frac{\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2}}{2\sqrt{1-ax}} -$$

---

3.260.  $\int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx$

↓ 2720

$$\frac{a^2\sqrt{ax-1}\left(3i\left(2\operatorname{arccosh}(ax)\operatorname{PolyLog}\left(3,-ie^{\operatorname{arccosh}(ax)}\right)-\int e^{-\operatorname{arccosh}(ax)}\operatorname{PolyLog}\left(3,-ie^{\operatorname{arccosh}(ax)}\right)de^{\operatorname{arccosh}(ax)}\right)\right)}{2\sqrt{1-ax}}$$

$$\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2}$$

$$3a\sqrt{ax-1}\left(-\frac{\operatorname{arccosh}(ax)^2}{x}+2a\left(2\operatorname{arccosh}(ax)\arctan\left(e^{\operatorname{arccosh}(ax)}\right)-i\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)+i\operatorname{PolyLog}\left(2,ie^{\operatorname{arccosh}(ax)}\right)\right)\right)$$

↓ 7143

$$\frac{a^2\sqrt{ax-1}\left(2\operatorname{arccosh}(ax)^3\arctan\left(e^{\operatorname{arccosh}(ax)}\right)+3i\left(2\operatorname{arccosh}(ax)\operatorname{PolyLog}\left(3,-ie^{\operatorname{arccosh}(ax)}\right)-\operatorname{PolyLog}\left(4,-ie^{\operatorname{arccosh}(ax)}\right)\right)\right)}{2\sqrt{1-ax}}$$

$$\frac{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3}{2x^2}$$

$$3a\sqrt{ax-1}\left(-\frac{\operatorname{arccosh}(ax)^2}{x}+2a\left(2\operatorname{arccosh}(ax)\arctan\left(e^{\operatorname{arccosh}(ax)}\right)-i\operatorname{PolyLog}\left(2,-ie^{\operatorname{arccosh}(ax)}\right)+i\operatorname{PolyLog}\left(2,ie^{\operatorname{arccosh}(ax)}\right)\right)\right)$$

input `Int[ArcCosh[a*x]^3/(x^3*sqrt[1 - a^2*x^2]),x]`

output `-1/2*(sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3)/x^2 - (3*a*sqrt[-1 + a*x]*(-(ArcCosh[a*x]^2/x) + 2*a*(2*ArcCosh[a*x]*ArcTan[E^ArcCosh[a*x]] - I*PolyLog[2, (-I)*E^ArcCosh[a*x]] + I*PolyLog[2, I*E^ArcCosh[a*x]])))/(2*sqrt[1 - a*x]) + (a^2*sqrt[-1 + a*x]*(2*ArcCosh[a*x]^3*ArcTan[E^ArcCosh[a*x]] + (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, (-I)*E^ArcCosh[a*x]] + 2*(ArcCosh[a*x]*PolyLog[3, (-I)*E^ArcCosh[a*x]] - PolyLog[4, (-I)*E^ArcCosh[a*x]])) - (3*I)*(-(ArcCosh[a*x]^2*PolyLog[2, I*E^ArcCosh[a*x]] + 2*(ArcCosh[a*x]*PolyLog[3, I*E^ArcCosh[a*x]] - PolyLog[4, I*E^ArcCosh[a*x]])))))/(2*sqrt[1 - a*x])`

## 3.260.3.1 Defintions of rubi rules used

rule 2715 `Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_)))]^(n_.)], x_Symbol]
:> Simp[1/(d*e*n*Log[F]) Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

rule 2720 `Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)
*(F_)^v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

rule 2838 `Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

rule 3011 `Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))]^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Simp[g*(m/(b*c*n*Log[F])) Int[(f + g*x)^(
m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e
, f, g, n}, x] && GtQ[m, 0]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]`

rule 4668 `Int[csc[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)]*((c_.) + (d_.)*(x_
))^m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^((-I)*e + f*fz*x)/E^(
I*k*Pi)]/(f*fz*I)), x] + (-Simp[d*(m/(f*fz*I)) Int[(c + d*x)^(m - 1)*Log[
1 - E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x] + Simp[d*(m/(f*fz*I)) Int[(c
+ d*x)^(m - 1)*Log[1 + E^((-I)*e + f*fz*x)/E^(I*k*Pi)], x], x]) /; FreeQ[{c
, d, e, f, fz}, x] && IntegerQ[2*k] && IGtQ[m, 0]`

rule 6298 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.)*((d_.)*(x_)^m_.), x_Symbol]
:= Simp[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^n/(d*(m + 1))), x] - Simp[b*c*
(n/(d*(m + 1))) Int[(d*x)^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 +
c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& NeQ[m, -1]`

rule 6347 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(d*f*(m + 1))), x] + (Simp[c^2*((m + 2*p + 3)/(f^2*(m + 1)))) Int[(f*x)^(m + 2)*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x], x] + Simp[b*c*(n/(f*(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && ILtQ[m, -1]`

rule 6361 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[m]`

rule 6362 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_))/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/c^(m + 1))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]] Subst[Int[(a + b*x)^n*Cosh[x]^m, x], x, ArcCosh[c*x]] /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[n, 0] && IntegerQ[m]`

rule 7143 `Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]`

rule 7163 `Int(((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^(c_.)*((a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Simp[f*(m/(b*c*p*Log[F])) Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]`



**3.260.4 Maple [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^3 \sqrt{-a^2 x^2 + 1}} dx$$

input `int(arccosh(a*x)^3/x^3/(-a^2*x^2+1)^(1/2),x)`

output `int(arccosh(a*x)^3/x^3/(-a^2*x^2+1)^(1/2),x)`

**3.260.5 Fricas [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^3 \sqrt{1 - a^2 x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2 x^2 + 1} x^3} dx$$

input `integrate(arccosh(a*x)^3/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*arccosh(a*x)^3/(a^2*x^5 - x^3), x)`

**3.260.6 Sympy [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^3 \sqrt{1 - a^2 x^2}} dx = \int \frac{\operatorname{acosh}^3(ax)}{x^3 \sqrt{-(ax - 1)(ax + 1)}} dx$$

input `integrate(acosh(a*x)**3/x**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(acosh(a*x)**3/(x**3*sqrt(-(a*x - 1)*(a*x + 1))), x)`

**3.260.7 Maxima [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}x^3} dx$$

input `integrate(arccosh(a*x)^3/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^3/(sqrt(-a^2*x^2 + 1)*x^3), x)`

**3.260.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{arcosh}(ax)^3}{\sqrt{-a^2x^2+1}x^3} dx$$

input `integrate(arccosh(a*x)^3/x^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^3/(sqrt(-a^2*x^2 + 1)*x^3), x)`

**3.260.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx = \int \frac{\operatorname{acosh}(ax)^3}{x^3\sqrt{1-a^2x^2}} dx$$

input `int(acosh(a*x)^3/(x^3*(1 - a^2*x^2)^(1/2)),x)`

output `int(acosh(a*x)^3/(x^3*(1 - a^2*x^2)^(1/2)), x)`

**3.261**  $\int \frac{(fx)^m(a+b\operatorname{arccosh}(cx))^3}{\sqrt{1-c^2x^2}} dx$

3.261.1 Optimal result . . . . .	2322
3.261.2 Mathematica [N/A] . . . . .	2322
3.261.3 Rubi [N/A] . . . . .	2323
3.261.4 Maple [N/A] (verified) . . . . .	2323
3.261.5 Fricas [N/A] . . . . .	2324
3.261.6 Sympy [N/A] . . . . .	2324
3.261.7 Maxima [N/A] . . . . .	2324
3.261.8 Giac [N/A] . . . . .	2325
3.261.9 Mupad [N/A] . . . . .	2325

**3.261.1 Optimal result**

Integrand size = 30, antiderivative size = 30

$$\int \frac{(fx)^m(a + \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2x^2}} dx = \operatorname{Int}\left(\frac{(fx)^m(a + \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2x^2}}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^3/(-c^2*x^2+1)^(1/2), x)`

**3.261.2 Mathematica [N/A]**

Not integrable

Time = 3.94 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m(a + \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2x^2}} dx = \int \frac{(fx)^m(a + \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2x^2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^3)/Sqrt[1 - c^2*x^2], x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^3)/Sqrt[1 - c^2*x^2], x]`

**3.261.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^3)/Sqrt[1 - c^2*x^2], x]`

output `$Aborted`

**3.261.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.261.4 Maple [N/A] (verified)**

Not integrable

Time = 1.14 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^3}{\sqrt{-c^2 x^2 + 1}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^3/(-c^2*x^2+1)^(1/2), x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^3/(-c^2*x^2+1)^(1/2), x)`

**3.261.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.27

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^3 (fx)^m}{\sqrt{-c^2 x^2 + 1}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))^3/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(-(b^3*arccosh(c*x)^3 + 3*a*b^2*arccosh(c*x)^2 + 3*a^2*b*arccosh(c*x) + a^3)*sqrt(-c^2*x^2 + 1)*(f*x)^m/(c^2*x^2 - 1), x)
```

**3.261.6 Sympy [N/A]**

Not integrable

Time = 70.00 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))^3}{\sqrt{-(cx - 1)(cx + 1)}} dx$$

```
input integrate((f*x)**m*(a+b*acosh(c*x))**3/(-c**2*x**2+1)**(1/2),x)
```

```
output Integral((f*x)**m*(a + b*acosh(c*x))**3/sqrt(-(c*x - 1)*(c*x + 1)), x)
```

**3.261.7 Maxima [N/A]**

Not integrable

Time = 0.37 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^3 (fx)^m}{\sqrt{-c^2 x^2 + 1}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))^3/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")
```

```
output integrate((b*arccosh(c*x) + a)^3*(f*x)^m/sqrt(-c^2*x^2 + 1), x)
```

---

3.261.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx$

**3.261.8 Giac [N/A]**

Not integrable

Time = 0.37 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^3 (fx)^m}{\sqrt{-c^2 x^2 + 1}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^3/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^3*(f*x)^m/sqrt(-c^2*x^2 + 1), x)`

**3.261.9 Mupad [N/A]**

Not integrable

Time = 3.17 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^3}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^3 (fx)^m}{\sqrt{1 - c^2 x^2}} dx$$

input `int(((a + b*acosh(c*x))^3*(f*x)^m)/(1 - c^2*x^2)^(1/2),x)`

output `int(((a + b*acosh(c*x))^3*(f*x)^m)/(1 - c^2*x^2)^(1/2), x)`

$$3.262 \quad \int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx$$

3.262.1 Optimal result	2326
3.262.2 Mathematica [A] (verified)	2326
3.262.3 Rubi [C] (verified)	2327
3.262.4 Maple [A] (verified)	2328
3.262.5 Fracas [F]	2329
3.262.6 Sympy [F]	2329
3.262.7 Maxima [F]	2329
3.262.8 Giac [F]	2330
3.262.9 Mupad [F(-1)]	2330

### 3.262.1 Optimal result

Integrand size = 20, antiderivative size = 67

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx = \frac{35c^3 \operatorname{Shi}(\operatorname{arccosh}(ax))}{64a} - \frac{21c^3 \operatorname{Shi}(3\operatorname{arccosh}(ax))}{64a} + \frac{7c^3 \operatorname{Shi}(5\operatorname{arccosh}(ax))}{64a} - \frac{c^3 \operatorname{Shi}(7\operatorname{arccosh}(ax))}{64a}$$

output  $35/64*c^3*\operatorname{Shi}(\operatorname{arccosh}(a*x))/a - 21/64*c^3*\operatorname{Shi}(3*\operatorname{arccosh}(a*x))/a + 7/64*c^3*\operatorname{Shi}(5*\operatorname{arccosh}(a*x))/a - 1/64*c^3*\operatorname{Shi}(7*\operatorname{arccosh}(a*x))/a$

### 3.262.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.67

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx = \frac{c^3(35\operatorname{Shi}(\operatorname{arccosh}(ax)) - 21\operatorname{Shi}(3\operatorname{arccosh}(ax)) + 7\operatorname{Shi}(5\operatorname{arccosh}(ax)) - \operatorname{Shi}(7\operatorname{arccosh}(ax)))}{64a}$$

input  $\operatorname{Integrate}[(c - a^2*c*x^2)^3/\operatorname{ArcCosh}[a*x], x]$

output  $(c^3*(35*\operatorname{SinhIntegral}[\operatorname{ArcCosh}[a*x]] - 21*\operatorname{SinhIntegral}[3*\operatorname{ArcCosh}[a*x]] + 7*\operatorname{SinhIntegral}[5*\operatorname{ArcCosh}[a*x]] - \operatorname{SinhIntegral}[7*\operatorname{ArcCosh}[a*x]]))/(64*a)$

---


$$3.262. \quad \int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx$$

**3.262.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.36 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6321, 3042, 26, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx \\
 & \quad \downarrow \text{6321} \\
 & \frac{c^3 \int \frac{\left(\frac{ax-1}{ax+1}\right)^{7/2} (ax+1)^7}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{3042} \\
 & \frac{c^3 \int \frac{i \sin(i \operatorname{arccosh}(ax))^7}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{26} \\
 & \frac{ic^3 \int \frac{\sin(i \operatorname{arccosh}(ax))^7}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{3793} \\
 & \frac{ic^3 \int \left( \frac{35i \sqrt{\frac{ax-1}{ax+1}} (ax+1)}{64 \operatorname{arccosh}(ax)} - \frac{21i \sinh(3 \operatorname{arccosh}(ax))}{64 \operatorname{arccosh}(ax)} + \frac{7i \sinh(5 \operatorname{arccosh}(ax))}{64 \operatorname{arccosh}(ax)} - \frac{i \sinh(7 \operatorname{arccosh}(ax))}{64 \operatorname{arccosh}(ax)} \right) d\operatorname{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ic^3 \left( \frac{35}{64} i \operatorname{Shi}(\operatorname{arccosh}(ax)) - \frac{21}{64} i \operatorname{Shi}(3 \operatorname{arccosh}(ax)) + \frac{7}{64} i \operatorname{Shi}(5 \operatorname{arccosh}(ax)) - \frac{1}{64} i \operatorname{Shi}(7 \operatorname{arccosh}(ax)) \right)}{a}
 \end{aligned}$$

input `Int[(c - a^2*c*x^2)^3/ArcCosh[a*x], x]`

output `((-I)*c^3*(((35*I)/64)*SinhIntegral[ArcCosh[a*x]] - ((21*I)/64)*SinhIntegral[3*ArcCosh[a*x]] + ((7*I)/64)*SinhIntegral[5*ArcCosh[a*x]] - (I/64)*SinhIntegral[7*ArcCosh[a*x]]))/a`

---

3.262.  $\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx$



## 3.262.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`
- rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

## 3.262.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.66

method	result	size
derivativedivides	$\frac{c^3(35 \operatorname{Shi}(\operatorname{arccosh}(ax)) - 21 \operatorname{Shi}(3 \operatorname{arccosh}(ax)) + 7 \operatorname{Shi}(5 \operatorname{arccosh}(ax)) - \operatorname{Shi}(7 \operatorname{arccosh}(ax)))}{64a}$	44
default	$\frac{c^3(35 \operatorname{Shi}(\operatorname{arccosh}(ax)) - 21 \operatorname{Shi}(3 \operatorname{arccosh}(ax)) + 7 \operatorname{Shi}(5 \operatorname{arccosh}(ax)) - \operatorname{Shi}(7 \operatorname{arccosh}(ax)))}{64a}$	44

input `int((-a^2*c*x^2+c)^3/arccosh(a*x),x,method=_RETURNVERBOSE)`

output `1/64/a*c^3*(35*Shi(arccosh(a*x))-21*Shi(3*arccosh(a*x))+7*Shi(5*arccosh(a*x))-Shi(7*arccosh(a*x)))`

---

3.262.  $\int \frac{(c-a^2cx^2)^3}{\operatorname{arccosh}(ax)} dx$

**3.262.5 Fracas [F]**

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx = \int -\frac{(a^2 cx^2 - c)^3}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^3/arccosh(a*x),x, algorithm="fricas")`

output `integral(-(a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3)/arccosh(a*x), x)`

**3.262.6 Sympy [F]**

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx = -c^3 \left( \int \frac{3a^2 x^2}{\operatorname{acosh}(ax)} dx + \int \left( -\frac{3a^4 x^4}{\operatorname{acosh}(ax)} \right) dx + \int \frac{a^6 x^6}{\operatorname{acosh}(ax)} dx + \int \left( -\frac{1}{\operatorname{acosh}(ax)} \right) dx \right)$$

input `integrate((-a**2*c*x**2+c)**3/acosh(a*x),x)`

output `-c**3*(Integral(3*a**2*x**2/acosh(a*x), x) + Integral(-3*a**4*x**4/acosh(a*x), x) + Integral(a**6*x**6/acosh(a*x), x) + Integral(-1/acosh(a*x), x))`

**3.262.7 Maxima [F]**

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx = \int -\frac{(a^2 cx^2 - c)^3}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^3/arccosh(a*x),x, algorithm="maxima")`

output `-integrate((a^2*c*x^2 - c)^3/arccosh(a*x), x)`

**3.262.8 Giac [F]**

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx = \int -\frac{(a^2 cx^2 - c)^3}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^3/arccosh(a*x),x, algorithm="giac")`

output `integrate(-(a^2*c*x^2 - c)^3/arccosh(a*x), x)`

**3.262.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)} dx = \int \frac{(c - a^2 c x^2)^3}{\operatorname{acosh}(ax)} dx$$

input `int((c - a^2*c*x^2)^3/acosh(a*x),x)`

output `int((c - a^2*c*x^2)^3/acosh(a*x), x)`

### 3.263 $\int \frac{(c-a^2cx^2)^2}{\operatorname{arccosh}(ax)} dx$

3.263.1 Optimal result . . . . .	2331
3.263.2 Mathematica [A] (verified) . . . . .	2331
3.263.3 Rubi [C] (verified) . . . . .	2332
3.263.4 Maple [A] (verified) . . . . .	2333
3.263.5 Fricas [F] . . . . .	2334
3.263.6 Sympy [F] . . . . .	2334
3.263.7 Maxima [F] . . . . .	2334
3.263.8 Giac [F] . . . . .	2335
3.263.9 Mupad [F(-1)] . . . . .	2335

#### 3.263.1 Optimal result

Integrand size = 20, antiderivative size = 50

$$\int \frac{(c - a^2cx^2)^2}{\operatorname{arccosh}(ax)} dx = \frac{5c^2\operatorname{Shi}(\operatorname{arccosh}(ax))}{8a} - \frac{5c^2\operatorname{Shi}(3\operatorname{arccosh}(ax))}{16a} + \frac{c^2\operatorname{Shi}(5\operatorname{arccosh}(ax))}{16a}$$

output  $5/8*c^2*Shi(\operatorname{arccosh}(a*x))/a-5/16*c^2*Shi(3*\operatorname{arccosh}(a*x))/a+1/16*c^2*Shi(5*\operatorname{arccosh}(a*x))/a$

#### 3.263.2 Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.68

$$\int \frac{(c - a^2cx^2)^2}{\operatorname{arccosh}(ax)} dx = \frac{c^2(10\operatorname{Shi}(\operatorname{arccosh}(ax)) - 5\operatorname{Shi}(3\operatorname{arccosh}(ax)) + \operatorname{Shi}(5\operatorname{arccosh}(ax)))}{16a}$$

input `Integrate[(c - a^2*c*x^2)^2/ArcCosh[a*x], x]`

output  $(c^2*(10*\operatorname{SinhIntegral}[\operatorname{ArcCosh}[a*x]] - 5*\operatorname{SinhIntegral}[3*\operatorname{ArcCosh}[a*x]] + \operatorname{SinhIntegral}[5*\operatorname{ArcCosh}[a*x]]))/(16*a)$

**3.263.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6321, 3042, 26, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)} dx \\
 \downarrow \text{6321} \\
 \frac{c^2 \int \frac{\left(\frac{ax-1}{ax+1}\right)^{5/2} (ax+1)^5}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{3042} \\
 \frac{c^2 \int -\frac{i \sin(i \operatorname{arccosh}(ax))^5}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{26} \\
 \frac{ic^2 \int \frac{\sin(i \operatorname{arccosh}(ax))^5}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{3793} \\
 \frac{ic^2 \int \left( \frac{5i \sqrt{\frac{ax-1}{ax+1}} (ax+1)}{8 \operatorname{arccosh}(ax)} - \frac{5i \sinh(3 \operatorname{arccosh}(ax))}{16 \operatorname{arccosh}(ax)} + \frac{i \sinh(5 \operatorname{arccosh}(ax))}{16 \operatorname{arccosh}(ax)} \right) d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{2009} \\
 \frac{ic^2 \left( \frac{5}{8} i \operatorname{Shi}(\operatorname{arccosh}(ax)) - \frac{5}{16} i \operatorname{Shi}(3 \operatorname{arccosh}(ax)) + \frac{1}{16} i \operatorname{Shi}(5 \operatorname{arccosh}(ax)) \right)}{a}
 \end{array}$$

input `Int[(c - a^2*c*x^2)^2/ArcCosh[a*x], x]`

output `((-I)*c^2*(((5*I)/8)*SinhIntegral[ArcCosh[a*x]] - ((5*I)/16)*SinhIntegral[3*ArcCosh[a*x]] + (I/16)*SinhIntegral[5*ArcCosh[a*x]]))/a`

---

3.263.  $\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)} dx$

## 3.263.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`
- rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

## 3.263.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.66

method	result	size
derivativedivides	$\frac{c^2(10 \operatorname{Shi}(\operatorname{arccosh}(ax)) - 5 \operatorname{Shi}(3 \operatorname{arccosh}(ax)) + \operatorname{Shi}(5 \operatorname{arccosh}(ax)))}{16a}$	33
default	$\frac{c^2(10 \operatorname{Shi}(\operatorname{arccosh}(ax)) - 5 \operatorname{Shi}(3 \operatorname{arccosh}(ax)) + \operatorname{Shi}(5 \operatorname{arccosh}(ax)))}{16a}$	33

input `int((-a^2*c*x^2+c)^2/arccosh(a*x),x,method=_RETURNVERBOSE)`

output `1/16/a*c^2*(10*Shi(arccosh(a*x))-5*Shi(3*arccosh(a*x))+Shi(5*arccosh(a*x)))`

**3.263.5 Fracas [F]**

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)} dx = \int \frac{(a^2 cx^2 - c)^2}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^2/arccosh(a*x),x, algorithm="fricas")`

output `integral((a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)/arccosh(a*x), x)`

**3.263.6 Sympy [F]**

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)} dx = c^2 \left( \int \left( -\frac{2a^2 x^2}{\operatorname{acosh}(ax)} \right) dx + \int \frac{a^4 x^4}{\operatorname{acosh}(ax)} dx + \int \frac{1}{\operatorname{acosh}(ax)} dx \right)$$

input `integrate((-a**2*c*x**2+c)**2/acosh(a*x),x)`

output `c**2*(Integral(-2*a**2*x**2/acosh(a*x), x) + Integral(a**4*x**4/acosh(a*x), x) + Integral(1/acosh(a*x), x))`

**3.263.7 Maxima [F]**

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)} dx = \int \frac{(a^2 cx^2 - c)^2}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^2/arccosh(a*x),x, algorithm="maxima")`

output `integrate((a^2*c*x^2 - c)^2/arccosh(a*x), x)`

**3.263.8 Giac [F]**

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)} dx = \int \frac{(a^2 cx^2 - c)^2}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^2/arccosh(a*x),x, algorithm="giac")`

output `integrate((a^2*c*x^2 - c)^2/arccosh(a*x), x)`

**3.263.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)} dx = \int \frac{(c - a^2 cx^2)^2}{\operatorname{acosh}(ax)} dx$$

input `int((c - a^2*c*x^2)^2/acosh(a*x),x)`

output `int((c - a^2*c*x^2)^2/acosh(a*x), x)`



### 3.264 $\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx$

3.264.1 Optimal result	2336
3.264.2 Mathematica [A] (verified)	2336
3.264.3 Rubi [C] (verified)	2337
3.264.4 Maple [A] (verified)	2338
3.264.5 Fricas [F]	2339
3.264.6 Sympy [F]	2339
3.264.7 Maxima [F]	2339
3.264.8 Giac [F]	2340
3.264.9 Mupad [F(-1)]	2340

#### 3.264.1 Optimal result

Integrand size = 18, antiderivative size = 29

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx = \frac{3c \operatorname{Shi}(\operatorname{arccosh}(ax))}{4a} - \frac{c \operatorname{Shi}(3 \operatorname{arccosh}(ax))}{4a}$$

output `3/4*c*Shi(arccosh(a*x))/a-1/4*c*Shi(3*arccosh(a*x))/a`

#### 3.264.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx = \frac{c(3 \operatorname{Shi}(\operatorname{arccosh}(ax)) - \operatorname{Shi}(3 \operatorname{arccosh}(ax)))}{4a}$$

input `Integrate[(c - a^2*c*x^2)/ArcCosh[a*x], x]`

output `(c*(3*SinhIntegral[ArcCosh[a*x]] - SinhIntegral[3*ArcCosh[a*x]]))/(4*a)`

### 3.264.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$ , Rules used = {6321, 3042, 26, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx \\
 \downarrow \text{6321} \\
 \frac{c \int \frac{\left(\frac{ax-1}{ax+1}\right)^{3/2} (ax+1)^3}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{3042} \\
 \frac{c \int \frac{i \sin(i \operatorname{arccosh}(ax))^3}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{26} \\
 \frac{ic \int \frac{\sin(i \operatorname{arccosh}(ax))^3}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{3793} \\
 \frac{ic \int \left( \frac{3i \sqrt{\frac{ax-1}{ax+1}} (ax+1)}{4 \operatorname{arccosh}(ax)} - \frac{i \sinh(3 \operatorname{arccosh}(ax))}{4 \operatorname{arccosh}(ax)} \right) d\operatorname{arccosh}(ax)}{a} \\
 \downarrow \text{2009} \\
 \frac{ic \left( \frac{3}{4} i \operatorname{Shi}(\operatorname{arccosh}(ax)) - \frac{1}{4} i \operatorname{Shi}(3 \operatorname{arccosh}(ax)) \right)}{a}
 \end{array}$$

input `Int[(c - a^2*c*x^2)/ArcCosh[a*x], x]`

output `((-I)*c*(((3*I)/4)*SinhIntegral[ArcCosh[a*x]] - (I/4)*SinhIntegral[3*ArcCosh[a*x]]))/a`

## 3.264.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`
- rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

## 3.264.4 Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{c(3 \operatorname{Shi}(\operatorname{arccosh}(ax)) - \operatorname{Shi}(3 \operatorname{arccosh}(ax)))}{4a}$	24
default	$\frac{c(3 \operatorname{Shi}(\operatorname{arccosh}(ax)) - \operatorname{Shi}(3 \operatorname{arccosh}(ax)))}{4a}$	24

input `int((-a^2*c*x^2+c)/arccosh(a*x),x,method=_RETURNVERBOSE)`

output `1/4/a*c*(3*Shi(arccosh(a*x))-Shi(3*arccosh(a*x)))`

**3.264.5 Fracas [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx = \int -\frac{a^2 cx^2 - c}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)/arccosh(a*x),x, algorithm="fricas")`

output `integral(-(a^2*c*x^2 - c)/arccosh(a*x), x)`

**3.264.6 Sympy [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx = -c \left( \int \frac{a^2 x^2}{\operatorname{acosh}(ax)} dx + \int \left( -\frac{1}{\operatorname{acosh}(ax)} \right) dx \right)$$

input `integrate((-a**2*c*x**2+c)/acosh(a*x),x)`

output `-c*(Integral(a**2*x**2/acosh(a*x), x) + Integral(-1/acosh(a*x), x))`

**3.264.7 Maxima [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx = \int -\frac{a^2 cx^2 - c}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)/arccosh(a*x),x, algorithm="maxima")`

output `-integrate((a^2*c*x^2 - c)/arccosh(a*x), x)`

**3.264.8 Giac [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx = \int -\frac{a^2 cx^2 - c}{\operatorname{arcosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)/arccosh(a*x),x, algorithm="giac")`

output `integrate(-(a^2*c*x^2 - c)/arccosh(a*x), x)`

**3.264.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)} dx = \int \frac{c - a^2 cx^2}{\operatorname{acosh}(ax)} dx$$

input `int((c - a^2*c*x^2)/acosh(a*x),x)`

output `int((c - a^2*c*x^2)/acosh(a*x), x)`

$$3.265 \quad \int \frac{1}{(c-a^2cx^2)\operatorname{arccosh}(ax)} dx$$

3.265.1 Optimal result	2341
3.265.2 Mathematica [N/A]	2341
3.265.3 Rubi [N/A]	2342
3.265.4 Maple [N/A] (verified)	2342
3.265.5 Fricas [N/A]	2343
3.265.6 Sympy [N/A]	2343
3.265.7 Maxima [N/A]	2343
3.265.8 Giac [N/A]	2344
3.265.9 Mupad [N/A]	2344

### 3.265.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c-a^2cx^2)\operatorname{arccosh}(ax)} dx = \operatorname{Int}\left(\frac{1}{(c-a^2cx^2)\operatorname{arccosh}(ax)}, x\right)$$

output `Unintegrable(1/(-a^2*c*x^2+c)/arccosh(a*x), x)`

### 3.265.2 Mathematica [N/A]

Not integrable

Time = 1.58 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c-a^2cx^2)\operatorname{arccosh}(ax)} dx = \int \frac{1}{(c-a^2cx^2)\operatorname{arccosh}(ax)} dx$$

input `Integrate[1/((c - a^2*c*x^2)*ArcCosh[a*x]), x]`

output `Integrate[1/((c - a^2*c*x^2)*ArcCosh[a*x]), x]`

**3.265.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arccosh}(ax) (c - a^2 cx^2)} dx$$

↓ 6325

$$\int \frac{1}{\operatorname{arccosh}(ax) (c - a^2 cx^2)} dx$$

input `Int[1/((c - a^2*c*x^2)*ArcCosh[a*x]),x]`

output `$Aborted`

**3.265.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^ (n_.)*((d_) + (e_.)*(x_)^2)^ (p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.265.4 Maple [N/A] (verified)**

Not integrable

Time = 0.57 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-a^2 cx^2 + c) \operatorname{arccosh}(ax)} dx$$

input `int(1/(-a^2*c*x^2+c)/arccosh(a*x),x)`

output `int(1/(-a^2*c*x^2+c)/arccosh(a*x),x)`

**3.265.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)} dx = \int -\frac{1}{(a^2 cx^2 - c) \operatorname{arccosh}(ax)} dx$$

input `integrate(1/(-a^2*c*x^2+c)/arccosh(a*x),x, algorithm="fricas")`output `integral(-1/((a^2*c*x^2 - c)*arccosh(a*x)), x)`**3.265.6 Sympy [N/A]**

Not integrable

Time = 1.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)} dx = -\frac{\int \frac{1}{a^2 x^2 \operatorname{acosh}(ax) - \operatorname{acosh}(ax)} dx}{c}$$

input `integrate(1/(-a**2*c*x**2+c)/acosh(a*x),x)`output `-Integral(1/(a**2*x**2*acosh(a*x) - acosh(a*x)), x)/c`**3.265.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.25

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)} dx = \int -\frac{1}{(a^2 cx^2 - c) \operatorname{arccosh}(ax)} dx$$

input `integrate(1/(-a^2*c*x^2+c)/arccosh(a*x),x, algorithm="maxima")`output `-integrate(1/((a^2*c*x^2 - c)*arccosh(a*x)), x)`

---

3.265.  $\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)} dx$



**3.265.8 Giac [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)} dx = \int -\frac{1}{(a^2 cx^2 - c) \operatorname{arccosh}(ax)} dx$$

input `integrate(1/(-a^2*c*x^2+c)/arccosh(a*x),x, algorithm="giac")`output `integrate(-1/((a^2*c*x^2 - c)*arccosh(a*x)), x)`**3.265.9 Mupad [N/A]**

Not integrable

Time = 2.77 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)} dx = \int \frac{1}{\operatorname{acosh}(ax) (c - a^2 cx^2)} dx$$

input `int(1/(acosh(a*x)*(c - a^2*c*x^2)),x)`output `int(1/(acosh(a*x)*(c - a^2*c*x^2)), x)`

$$3.266 \quad \int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx$$

3.266.1 Optimal result	2345
3.266.2 Mathematica [N/A]	2345
3.266.3 Rubi [N/A]	2346
3.266.4 Maple [N/A] (verified)	2346
3.266.5 Fricas [N/A]	2347
3.266.6 Sympy [N/A]	2347
3.266.7 Maxima [N/A]	2347
3.266.8 Giac [N/A]	2348
3.266.9 Mupad [N/A]	2348

### 3.266.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx = \operatorname{Int}\left(\frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)}, x\right)$$

output `Unintegrable(1/(-a^2*c*x^2+c)^2/arccosh(a*x),x)`

### 3.266.2 Mathematica [N/A]

Not integrable

Time = 5.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx = \int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx$$

input `Integrate[1/((c - a^2*c*x^2)^2*ArcCosh[a*x]),x]`

output `Integrate[1/((c - a^2*c*x^2)^2*ArcCosh[a*x]), x]`

**3.266.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arccosh}(ax) (c - a^2cx^2)^2} dx$$

↓ 6325

$$\int \frac{1}{\operatorname{arccosh}(ax) (c - a^2cx^2)^2} dx$$

input `Int[1/((c - a^2*c*x^2)^2*ArcCosh[a*x]), x]`

output `$Aborted`

**3.266.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.266.4 Maple [N/A] (verified)**

Not integrable

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-a^2cx^2 + c)^2 \operatorname{arccosh}(ax)} dx$$

input `int(1/(-a^2*c*x^2+c)^2/arccosh(a*x), x)`

output `int(1/(-a^2*c*x^2+c)^2/arccosh(a*x), x)`

**3.266.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx = \int \frac{1}{(a^2 cx^2 - c)^2 \operatorname{arcosh}(ax)} dx$$

input `integrate(1/(-a^2*c*x^2+c)^2/arccosh(a*x),x, algorithm="fricas")`output `integral(1/((a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)*arccosh(a*x)), x)`**3.266.6 Sympy [N/A]**

Not integrable

Time = 4.23 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx = \int \frac{\frac{1}{a^4 x^4 \operatorname{acosh}(ax) - 2a^2 x^2 \operatorname{acosh}(ax) + \operatorname{acosh}(ax)}}{c^2} dx$$

input `integrate(1/(-a**2*c*x**2+c)**2/acosh(a*x),x)`output `Integral(1/(a**4*x**4*acosh(a*x) - 2*a**2*x**2*acosh(a*x) + acosh(a*x)), x)/c**2`**3.266.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx = \int \frac{1}{(a^2 cx^2 - c)^2 \operatorname{arcosh}(ax)} dx$$

input `integrate(1/(-a^2*c*x^2+c)^2/arccosh(a*x),x, algorithm="maxima")`output `integrate(1/((a^2*c*x^2 - c)^2*arccosh(a*x)), x)`

**3.266.8 Giac [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx = \int \frac{1}{(a^2 cx^2 - c)^2 \operatorname{arcosh}(ax)} dx$$

input `integrate(1/(-a^2*c*x^2+c)^2/arccosh(a*x),x, algorithm="giac")`output `integrate(1/((a^2*c*x^2 - c)^2*arccosh(a*x)), x)`**3.266.9 Mupad [N/A]**

Not integrable

Time = 3.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)} dx = \int \frac{1}{\operatorname{acosh}(ax) (c - a^2 cx^2)^2} dx$$

input `int(1/(acosh(a*x)*(c - a^2*c*x^2)^2),x)`output `int(1/(acosh(a*x)*(c - a^2*c*x^2)^2), x)`

**3.267**  $\int \frac{x^4 \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$

3.267.1 Optimal result . . . . .	2349
3.267.2 Mathematica [A] (warning: unable to verify) . . . . .	2350
3.267.3 Rubi [A] (verified) . . . . .	2350
3.267.4 Maple [A] (verified) . . . . .	2352
3.267.5 Fricas [F] . . . . .	2352
3.267.6 Sympy [F] . . . . .	2353
3.267.7 Maxima [F] . . . . .	2353
3.267.8 Giac [F] . . . . .	2353
3.267.9 Mupad [F(-1)] . . . . .	2354

**3.267.1 Optimal result**

Integrand size = 28, antiderivative size = 339

$$\int \frac{x^4 \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = -\frac{\sqrt{1-cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^5\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^5\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^5\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \log(a+b\operatorname{arccosh}(cx))}{16bc^5\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^5\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^5\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{6a}{b}\right) \operatorname{Shi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^5\sqrt{-1+cx}}$$

output 
$$-1/32*\text{Chi}(2*(a+b*\text{arccosh}(c*x))/b)*\cosh(2*a/b)*(-c*x+1)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}+1/16*\text{Chi}(4*(a+b*\text{arccosh}(c*x))/b)*\cosh(4*a/b)*(-c*x+1)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}+1/32*\text{Chi}(6*(a+b*\text{arccosh}(c*x))/b)*\cosh(6*a/b)*(-c*x+1)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}-1/16*\ln(a+b*\text{arccosh}(c*x))*(-c*x+1)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}+1/32*\text{Shi}(2*(a+b*\text{arccosh}(c*x))/b)*\sinh(2*a/b)*(-c*x+1)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}-1/16*\text{Shi}(4*(a+b*\text{arccosh}(c*x))/b)*\sinh(4*a/b)*(-c*x+1)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}-1/32*\text{Shi}(6*(a+b*\text{arccosh}(c*x))/b)*\sinh(6*a/b)*(-c*x+1)^{(1/2)}/b/c^5/(c*x-1)^{(1/2)}$$

### 3.267.2 Mathematica [A] (warning: unable to verify)

Time = 0.38 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.55

$$\int \frac{x^4 \sqrt{1-c^2x^2}}{a+b\text{arccosh}(cx)} dx$$

$$= \frac{\sqrt{1-c^2x^2} \left( -\cosh\left(\frac{2a}{b}\right) \text{Chi}\left(2\left(\frac{a}{b} + \text{arccosh}(cx)\right)\right) + 2 \cosh\left(\frac{4a}{b}\right) \text{Chi}\left(4\left(\frac{a}{b} + \text{arccosh}(cx)\right)\right) + \cosh\left(\frac{6a}{b}\right) \text{Chi}\left(6\left(\frac{a}{b} + \text{arccosh}(cx)\right)\right) \right)}{c^5}$$

input `Integrate[(x^4*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output 
$$\frac{(\text{Sqrt}[1 - c^2*x^2]*(-(\text{Cosh}[(2*a)/b]*\text{CoshIntegral}[2*(a/b + \text{ArcCosh}[c*x])]) + 2*\text{Cosh}[(4*a)/b]*\text{CoshIntegral}[4*(a/b + \text{ArcCosh}[c*x])]) + \text{Cosh}[(6*a)/b]*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]) - 2*\text{Log}[a + b*\text{ArcCosh}[c*x]] + \text{Sinh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])]) - 2*\text{Sinh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])]) - \text{Sinh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])])}{(32*c^5*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(b + b*c*x))}$$

### 3.267.3 Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.56, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4 \sqrt{1-c^2x^2}}{a+b\text{arccosh}(cx)} dx$$

---

3.267.  $\int \frac{x^4 \sqrt{1-c^2x^2}}{a+b\text{arccosh}(cx)} dx$

$$\begin{array}{c}
 \downarrow \text{6367} \\
 \frac{\sqrt{1-cx} \int \frac{\cosh^4\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^5\sqrt{cx-1}} \\
 \downarrow \text{5971} \\
 \frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{6a}{b} - \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} + \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} - \frac{1}{16(a+b\operatorname{arccosh}(cx))} \right)}{bc^5\sqrt{cx-1}} \\
 \downarrow \text{2009} \\
 \frac{\sqrt{1-cx} \left( -\frac{1}{32} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{32} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc^5\sqrt{cx-1}}
 \end{array}$$

input `Int[(x^4*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output `(sqrt[1 - c*x]*(-1/32*(Cosh[(2*a)/b]*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]) + (Cosh[(4*a)/b]*CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b])/16 + (Cosh[(6*a)/b]*CoshIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32 - Log[a + b*ArcCosh[c*x])/16 + (Sinh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/32 - (Sinh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/16 - (Sinh[(6*a)/b]*SinhIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32)/(b*c^5*sqrt[-1 + c*x])`

### 3.267.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`



rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.267.4 Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.88

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(4\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))+4\ln(a+b\operatorname{arccosh}(cx))cx+\operatorname{Ei}_1(6\operatorname{arccosh}(cx)+\frac{6a}{b})\right)}{\dots}$

input `int(x^4*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{64}(-c^2x^2+1)^{1/2}(-(cx-1)^{1/2}(cx+1)^{1/2}cx+c^2x^2-1)(4*(cx-1)^{1/2}(cx+1)^{1/2}\ln(a+b\operatorname{arccosh}(cx))+4*\ln(a+b\operatorname{arccosh}(cx))*cx+\operatorname{Ei}(1,6*\operatorname{arccosh}(cx)+6*a/b)*\exp((b*\operatorname{arccosh}(cx)+6*a)/b)+\operatorname{Ei}(1,-6*\operatorname{arccosh}(cx)-6*a/b)*\exp(-(-b*\operatorname{arccosh}(cx)+6*a)/b)+2*\operatorname{Ei}(1,4*\operatorname{arccosh}(cx)+4*a/b)*\exp((b*\operatorname{arccosh}(cx)+4*a)/b)-\operatorname{Ei}(1,2*\operatorname{arccosh}(cx)+2*a/b)*\exp((b*\operatorname{arccosh}(cx)+2*a)/b)-\operatorname{Ei}(1,-2*\operatorname{arccosh}(cx)-2*a/b)*\exp(-(-b*\operatorname{arccosh}(cx)+2*a)/b)+2*\operatorname{Ei}(1,-4*\operatorname{arccosh}(cx)-4*a/b)*\exp(-(-b*\operatorname{arccosh}(cx)+4*a)/b)))/(cx+1)/c^5/(cx-1)/b$$

### 3.267.5 Fracas [F]

$$\int \frac{x^4\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}x^4}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^4*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^4/(b*arccosh(c*x) + a), x)`

**3.267.6 Sympy [F]**

$$\int \frac{x^4 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{x^4 \sqrt{-(cx - 1)(cx + 1)}}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate(x**4*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**4*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x)), x)`

**3.267.7 Maxima [F]**

$$\int \frac{x^4 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^4}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^4*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)*x^4/(b*arccosh(c*x) + a), x)`

**3.267.8 Giac [F]**

$$\int \frac{x^4 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^4}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^4*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)*x^4/(b*arccosh(c*x) + a), x)`

**3.267.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4 \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{x^4 \sqrt{1 - c^2 x^2}}{a + b \operatorname{acosh}(cx)} dx$$

input `int((x^4*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)),x)`output `int((x^4*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)), x)`

**3.268**       $\int \frac{x^3 \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$

3.268.1 Optimal result	2355
3.268.2 Mathematica [A] (warning: unable to verify)	2356
3.268.3 Rubi [A] (verified)	2356
3.268.4 Maple [A] (verified)	2358
3.268.5 Fricas [F]	2358
3.268.6 Sympy [F]	2359
3.268.7 Maxima [F]	2359
3.268.8 Giac [F(-2)]	2359
3.268.9 Mupad [F(-1)]	2360

**3.268.1 Optimal result**

Integrand size = 28, antiderivative size = 297

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = -\frac{\sqrt{1-cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8bc^4\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^4\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^4\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8bc^4\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^4\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^4\sqrt{-1+cx}}$$

output 
$$\begin{aligned} & -1/8*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\text{cosh}(a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +1/16*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\text{cosh}(3*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +1/16*\text{Chi}(5*(a+b*\text{arccosh}(c*x))/b)*\text{cosh}(5*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +1/8*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*\text{sinh}(a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -1/16*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*\text{sinh}(3*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -1/16*\text{Shi}(5*(a+b*\text{arccosh}(c*x))/b)*\text{sinh}(5*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \end{aligned}$$

### 3.268.2 Mathematica [A] (warning: unable to verify)

Time = 0.34 (sec) , antiderivative size = 171, normalized size of antiderivative = 0.58

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{a+b\text{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2}(-2\cosh(\frac{a}{b})\text{Chi}(\frac{a}{b}+\text{arccosh}(cx)) + \cosh(\frac{3a}{b})\text{Chi}(3(\frac{a}{b}+\text{arccosh}(cx))) + \cosh(\frac{5a}{b})\text{Chi}(5(\frac{a}{b}+\text{arccosh}(cx))))}{16c^4\sqrt{(-1+cx)/(1+cx)}} + (b+bcx)$$

input `Integrate[(x^3*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output 
$$\begin{aligned} & (\text{Sqrt}[1 - c^2*x^2]*(-2*\text{Cosh}[a/b]*\text{CoshIntegral}[a/b + \text{ArcCosh}[c*x]] + \text{Cosh}[(3*a)/b]*\text{CoshIntegral}[3*(a/b + \text{ArcCosh}[c*x])] \\ & + \text{Cosh}[(5*a)/b]*\text{CoshIntegral}[5*(a/b + \text{ArcCosh}[c*x])] + 2*\text{Sinh}[a/b]*\text{SinhIntegral}[a/b + \text{ArcCosh}[c*x]] - \text{Sinh}[(3*a)/b]*\text{SinhIntegral}[3*(a/b + \text{ArcCosh}[c*x])] \\ & - \text{Sinh}[(5*a)/b]*\text{SinhIntegral}[5*(a/b + \text{ArcCosh}[c*x])])/(16*c^4*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(b + b*c*x)) \end{aligned}$$

### 3.268.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.58, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{a+b\text{arccosh}(cx)} dx$$

---

3.268.  $\int \frac{x^3 \sqrt{1-c^2x^2}}{a+b\text{arccosh}(cx)} dx$

$$\begin{array}{c}
 \downarrow \text{6367} \\
 \frac{\sqrt{1-cx} \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^4\sqrt{cx-1}} \\
 \downarrow \text{5971} \\
 \frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} + \frac{\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{bc^4\sqrt{cx-1}} \\
 \downarrow \text{2009} \\
 \frac{\sqrt{1-cx} \left( -\frac{1}{8} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) + \frac{1}{16} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc^4}
 \end{array}$$

input `Int[(x^3*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output `(sqrt[1 - c*x]*(-1/8*(Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b]) + (Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/16 + (Cosh[(5*a)/b]*CoshIntegral[(5*(a + b*ArcCosh[c*x])/b])/16 + (Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/8 - (Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/16 - (Sinh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/16))/(b*c^4*sqrt[-1 + c*x])`

### 3.268.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.268.4 Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 250, normalized size of antiderivative = 0.84

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(\text{Ei}_1\left(5\operatorname{arccosh}(cx)+\frac{5a}{b}\right)e^{\frac{b\operatorname{arccosh}(cx)+5a}{b}}+\text{Ei}_1\left(3\operatorname{arccosh}(cx)+\frac{3a}{b}\right)e^{\frac{b\operatorname{arccosh}(cx)+3a}{b}}+\text{Ei}_1\left(-3\operatorname{arccosh}(cx)-\frac{3a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)+3a}{b}}+\text{Ei}_1\left(-5\operatorname{arccosh}(cx)-\frac{5a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)+5a}{b}}-2\text{Ei}_1\left(\operatorname{arccosh}(cx)+\frac{a}{b}\right)e^{\frac{a+b\operatorname{arccosh}(cx)}{b}}-2\text{Ei}_1\left(-\operatorname{arccosh}(cx)-\frac{a}{b}\right)e^{-\frac{a+b\operatorname{arccosh}(cx)}{b}}\right)}{(c*x+1)/c^4/(c*x-1)/b}$

input `int(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `1/32*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(Ei(1,5*arccosh(c*x)+5*a/b)*exp((b*arccosh(c*x)+5*a)/b)+Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)+Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(-b*arccosh(c*x)+3*a)/b)+Ei(1,-5*arccosh(c*x)-5*a/b)*exp(-(-b*arccosh(c*x)+5*a)/b)-2*Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)-2*Ei(1,-arccosh(c*x)-a/b)*exp(-(-b*arccosh(c*x)+a)/b))/(c*x+1)/c^4/(c*x-1)/b`

### 3.268.5 Fracas [F]

$$\int \frac{x^3\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}x^3}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^3/(b*arccosh(c*x) + a), x)`

**3.268.6 Sympy [F]**

$$\int \frac{x^3 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{x^3 \sqrt{-(cx - 1)(cx + 1)}}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate(x**3*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**3*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x)), x)`

**3.268.7 Maxima [F]**

$$\int \frac{x^3 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^3}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)*x^3/(b*arccosh(c*x) + a), x)`

**3.268.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.268.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3 \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{x^3 \sqrt{1 - c^2 x^2}}{a + b \operatorname{acosh}(cx)} dx$$

input `int((x^3*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)),x)`output `int((x^3*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)), x)`

**3.269**  $\int \frac{x^2 \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$

3.269.1 Optimal result . . . . . 2361  
 3.269.2 Mathematica [A] (verified) . . . . . 2361  
 3.269.3 Rubi [A] (verified) . . . . . 2362  
 3.269.4 Maple [A] (verified) . . . . . 2363  
 3.269.5 Fricas [F] . . . . . 2364  
 3.269.6 Sympy [F] . . . . . 2364  
 3.269.7 Maxima [F] . . . . . 2364  
 3.269.8 Giac [F] . . . . . 2365  
 3.269.9 Mupad [F(-1)] . . . . . 2365

**3.269.1 Optimal result**

Integrand size = 28, antiderivative size = 139

$$\int \frac{x^2 \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{1-cx} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8bc^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \log(a+b\operatorname{arccosh}(cx))}{8bc^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8bc^3\sqrt{-1+cx}}$$

```
output 1/8*Chi(4*(a+b*arccosh(c*x))/b)*cosh(4*a/b)*(-c*x+1)^(1/2)/b/c^3/(c*x-1)^(1/2)-1/8*ln(a+b*arccosh(c*x))*(-c*x+1)^(1/2)/b/c^3/(c*x-1)^(1/2)-1/8*Shi(4*(a+b*arccosh(c*x))/b)*sinh(4*a/b)*(-c*x+1)^(1/2)/b/c^3/(c*x-1)^(1/2)
```

**3.269.2 Mathematica [A] (verified)**

Time = 0.31 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.74

$$\int \frac{x^2 \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{-((-1+cx)(1+cx))} \left(-\cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(4\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) + \log(a+b\operatorname{arccosh}(cx)) + \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(4\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right)\right)}{8bc^3\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}$$

input `Integrate[(x^2*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output `-1/8*(Sqrt[-((-1 + c*x)*(1 + c*x))]*(-(Cosh[(4*a)/b]*CoshIntegral[4*(a/b + ArcCosh[c*x])]) + Log[a + b*ArcCosh[c*x]] + Sinh[(4*a)/b]*SinhIntegral[4*(a/b + ArcCosh[c*x])]))/(b*c^3*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`

### 3.269.3 Rubi [A] (verified)

Time = 0.49 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.65, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{1 - cx} \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a + b \operatorname{barccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a + b \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc^3 \sqrt{cx - 1}}$$

$$\downarrow \text{5971}$$

$$\frac{\sqrt{1 - cx} \int \left( \frac{\cosh\left(\frac{4a}{b} - \frac{4(a + b \operatorname{barccosh}(cx))}{b}\right)}{8(a + \operatorname{barccosh}(cx))} - \frac{1}{8(a + \operatorname{barccosh}(cx))} \right) d(a + \operatorname{barccosh}(cx))}{bc^3 \sqrt{cx - 1}}$$

$$\downarrow \text{2009}$$

$$\frac{\sqrt{1 - cx} \left( \frac{1}{8} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a + b \operatorname{barccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a + b \operatorname{barccosh}(cx))}{b}\right) - \frac{1}{8} \log(a + \operatorname{barccosh}(cx)) \right)}{bc^3 \sqrt{cx - 1}}$$

input `Int[(x^2*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[1 - c*x]*((Cosh[(4*a)/b]*CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8 - Log[a + b*ArcCosh[c*x]]/8 - (Sinh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8))/(b*c^3*Sqrt[-1 + c*x])`

---

3.269.  $\int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx$

## 3.269.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

## 3.269.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.19

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(2\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))+2\ln(a+b\operatorname{arccosh}(cx))cx+\operatorname{Ei}_1(4\operatorname{arccosh}(cx)+\frac{4}{b})\right)}{16(cx+1)c^3(cx-1)b}$

input `int(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `1/16*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(a+b*arccosh(c*x))+2*ln(a+b*arccosh(c*x))*c*x+ Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)+Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(b*arccosh(c*x)+4*a)/b))/(c*x+1)/c^3/(c*x-1)/b`

**3.269.5 Fricas [F]**

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^2/(b*arccosh(c*x) + a), x)`

**3.269.6 Sympy [F]**

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{x^2 \sqrt{-(cx - 1)(cx + 1)}}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate(x**2*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**2*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x)), x)`

**3.269.7 Maxima [F]**

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)*x^2/(b*arccosh(c*x) + a), x)`

**3.269.8 Giac [F]**

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)*x^2/(b*arccosh(c*x) + a), x)`

**3.269.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{x^2 \sqrt{1 - c^2 x^2}}{a + b \operatorname{acosh}(cx)} dx$$

input `int((x^2*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)),x)`

output `int((x^2*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)), x)`

**3.270**  $\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$

3.270.1 Optimal result . . . . . 2366  
 3.270.2 Mathematica [A] (warning: unable to verify) . . . . . 2367  
 3.270.3 Rubi [A] (verified) . . . . . 2367  
 3.270.4 Maple [A] (verified) . . . . . 2368  
 3.270.5 Fricas [F] . . . . . 2369  
 3.270.6 Sympy [F] . . . . . 2369  
 3.270.7 Maxima [F] . . . . . 2369  
 3.270.8 Giac [F] . . . . . 2370  
 3.270.9 Mupad [F(-1)] . . . . . 2370

**3.270.1 Optimal result**

Integrand size = 26, antiderivative size = 197

$$\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = -\frac{\sqrt{1-cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4bc^2\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4bc^2\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4bc^2\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4bc^2\sqrt{-1+cx}}$$

output

```
-1/4*Chi((a+b*arccosh(c*x))/b)*cosh(a/b)*(-c*x+1)^(1/2)/b/c^2/(c*x-1)^(1/2)
)+1/4*Chi(3*(a+b*arccosh(c*x))/b)*cosh(3*a/b)*(-c*x+1)^(1/2)/b/c^2/(c*x-1)
)^(1/2)+1/4*Shi((a+b*arccosh(c*x))/b)*sinh(a/b)*(-c*x+1)^(1/2)/b/c^2/(c*x-1)
)^(1/2)-1/4*Shi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)*(-c*x+1)^(1/2)/b/c^2/(
c*x-1)^(1/2)
```

**3.270.2 Mathematica [A] (warning: unable to verify)**

Time = 0.29 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.64

$$\int \frac{x\sqrt{1-c^2x^2}}{a + \operatorname{barccosh}(cx)} dx$$

$$= \frac{\sqrt{1-c^2x^2} \left( -\cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) + \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) + \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \right)}{4c^2 \sqrt{\frac{-1+cx}{1+cx}} (b + bcx)}$$

input `Integrate[(x*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`output `(Sqrt[1 - c^2*x^2]*(-(Cosh[a/b]*CoshIntegral[a/b + ArcCosh[c*x]]) + Cosh[(3*a)/b]*CoshIntegral[3*(a/b + ArcCosh[c*x])] + Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] - Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])]))/(4*c^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(b + b*c*x))`**3.270.3 Rubi [A] (verified)**Time = 0.50 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.62, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x\sqrt{1-c^2x^2}}{a + \operatorname{barccosh}(cx)} dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{1-cx} \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc^2\sqrt{cx-1}}$$

$$\downarrow \text{5971}$$

$$\frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a + \operatorname{barccosh}(cx))}{bc^2\sqrt{cx-1}}$$

$$\downarrow \text{2009}$$

---

3.270.  $\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$



$$\frac{\sqrt{1-cx} \left( -\frac{1}{4} \cosh\left(\frac{a}{b}\right) \text{Chi}\left(\frac{a+b\text{arccosh}(cx)}{b}\right) + \frac{1}{4} \cosh\left(\frac{3a}{b}\right) \text{Chi}\left(\frac{3(a+b\text{arccosh}(cx))}{b}\right) + \frac{1}{4} \sinh\left(\frac{a}{b}\right) \text{Shi}\left(\frac{a+b\text{arccosh}(cx)}{b}\right) \right)}{bc^2\sqrt{cx-1}}$$

input `Int[(x*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output `(sqrt[1 - c*x]*(-1/4*(Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b]) + (Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/4 + (Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/4 - (Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/4))/(b*c^2*sqrt[-1 + c*x])`

### 3.270.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.270.4 Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.94

method	result
default	$\frac{\sqrt{-c^2x^2+1} (-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1) \left( \text{Ei}_1\left(3\text{arccosh}(cx)+\frac{3a}{b}\right)e^{\frac{b\text{arccosh}(cx)+3a}{b}} + \text{Ei}_1\left(-3\text{arccosh}(cx)-\frac{3a}{b}\right)e^{-\frac{b\text{arccosh}(cx)+3a}{b}} \right)}{8(cx+1)c^2(cx-1)b}$

input `int(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

3.270.  $\int \frac{x\sqrt{1-c^2x^2}}{a+b\text{arccosh}(cx)} dx$

output  $1/8*(-c^2*x^2+1)^{(1/2)}*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+c^2*x^2-1)*(Ei(1, 3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)+Ei(1, -3*arccosh(c*x)-3*a/b)*exp(-(-b*arccosh(c*x)+3*a)/b)-Ei(1, arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)-Ei(1, -arccosh(c*x)-a/b)*exp(-(-b*arccosh(c*x)+a)/b))/(c*x+1)/c^2/(c*x-1)/b$

### 3.270.5 Fracas [F]

$$\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}x}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x/(b*arccosh(c*x) + a), x)`

### 3.270.6 Sympy [F]

$$\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x\sqrt{-(cx-1)(cx+1)}}{a+b\operatorname{acosh}(cx)} dx$$

input `integrate(x*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x)),x)`

output `Integral(x*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x)), x)`

### 3.270.7 Maxima [F]

$$\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}x}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)*x/(b*arccosh(c*x) + a), x)`

**3.270.8 Giac [F]**

$$\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}x}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)*x/(b*arccosh(c*x) + a), x)`

**3.270.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x\sqrt{1-c^2x^2}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((x*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)),x)`

output `int((x*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)), x)`

**3.271**  $\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$

3.271.1 Optimal result . . . . . 2371  
 3.271.2 Mathematica [A] (warning: unable to verify) . . . . . 2371  
 3.271.3 Rubi [A] (verified) . . . . . 2372  
 3.271.4 Maple [A] (verified) . . . . . 2374  
 3.271.5 Fricas [F] . . . . . 2374  
 3.271.6 Sympy [F] . . . . . 2374  
 3.271.7 Maxima [F] . . . . . 2375  
 3.271.8 Giac [F] . . . . . 2375  
 3.271.9 Mupad [F(-1)] . . . . . 2375

**3.271.1 Optimal result**

Integrand size = 25, antiderivative size = 139

$$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{1-cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2bc\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \log(a+b\operatorname{arccosh}(cx))}{2bc\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2bc\sqrt{-1+cx}}$$

output  $1/2*\operatorname{Chi}(2*(a+b*\operatorname{arccosh}(c*x))/b)*\cosh(2*a/b)*(-c*x+1)^{(1/2)}/b/c/(c*x-1)^{(1/2)}-1/2*\ln(a+b*\operatorname{arccosh}(c*x))*(-c*x+1)^{(1/2)}/b/c/(c*x-1)^{(1/2)}-1/2*\operatorname{Shi}(2*(a+b*\operatorname{arccosh}(c*x))/b)*\sinh(2*a/b)*(-c*x+1)^{(1/2)}/b/c/(c*x-1)^{(1/2)}$

**3.271.2 Mathematica [A] (warning: unable to verify)**

Time = 0.31 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.76

$$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{-((-1+cx)(1+cx))}(\cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) - \log(a+b\operatorname{arccosh}(cx)) - \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right)}{2bc\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}$$

---

3.271.  $\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$

input `Integrate[Sqrt[1 - c^2*x^2]/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[-((-1 + c*x)*(1 + c*x))]*(Cosh[(2*a)/b]*CoshIntegral[2*(a/b + ArcCosh[c*x]]) - Log[a + b*ArcCosh[c*x]] - Sinh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[c*x])])/(2*b*c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`

### 3.271.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.65, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6321, 3042, 25, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx \\
 & \quad \downarrow \text{6321} \\
 & \frac{\sqrt{1 - cx} \int \frac{\sinh^2\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{a + b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{bc\sqrt{cx - 1}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{1 - cx} \int -\frac{\sin\left(\frac{ia}{b} - \frac{i(a + b \operatorname{arccosh}(cx))}{b}\right)^2}{a + b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{bc\sqrt{cx - 1}} \\
 & \quad \downarrow \text{25} \\
 & \frac{\sqrt{1 - cx} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a + b \operatorname{arccosh}(cx))}{b}\right)^2}{a + b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{bc\sqrt{cx - 1}} \\
 & \quad \downarrow \text{3793} \\
 & \frac{\sqrt{1 - cx} \int \left( \frac{1}{2(a + b \operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a + b \operatorname{arccosh}(cx))}{b}\right)}{2(a + b \operatorname{arccosh}(cx))} \right) d(a + b \operatorname{arccosh}(cx))}{bc\sqrt{cx - 1}} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.271.  $\int \frac{\sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx$

$$\frac{\sqrt{1-cx} \left( \frac{1}{2} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{2} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{2} \log(a+b\operatorname{arccosh}(cx)) \right)}{bc\sqrt{cx-1}}$$

input `Int[Sqrt[1 - c^2*x^2]/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[1 - c*x]*((Cosh[(2*a)/b]*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b])/2 - Log[a + b*ArcCosh[c*x]]/2 - (Sinh[(2*a)/b]*ShiIntegral[(2*(a + b*ArcCosh[c*x])/b])/2))/(b*c*Sqrt[-1 + c*x])`

### 3.271.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

**3.271.4 Maple [A] (verified)**

Time = 0.66 (sec) , antiderivative size = 165, normalized size of antiderivative = 1.19

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(2\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))+2\ln(a+b\operatorname{arccosh}(cx))cx+\operatorname{Ei}_1(2\operatorname{arccosh}(cx)+\frac{2}{b})\right)}{4(cx-1)(cx+1)cb}$

```
input int((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output 1/4*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(a+b*arccosh(c*x))+2*ln(a+b*arccosh(c*x))*c*x+Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)+Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b))/(c*x-1)/(c*x+1)/c/b
```

**3.271.5 Fricas [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}}{b\operatorname{arccosh}(cx)+a} dx$$

```
input integrate((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")
```

```
output integral(sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)
```

**3.271.6 Sympy [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{a+b\operatorname{acosh}(cx)} dx$$

```
input integrate((-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x)),x)
```

```
output Integral(sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x)), x)
```

**3.271.7 Maxima [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

**3.271.8 Giac [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{-c^2x^2+1}}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

**3.271.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{1-c^2x^2}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((1 - c^2*x^2)^(1/2)/(a + b*acosh(c*x)),x)`

output `int((1 - c^2*x^2)^(1/2)/(a + b*acosh(c*x)), x)`



$$3.272 \quad \int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx$$

3.272.1 Optimal result	2376
3.272.2 Mathematica [N/A]	2376
3.272.3 Rubi [N/A]	2377
3.272.4 Maple [N/A] (verified)	2378
3.272.5 Fricas [N/A]	2378
3.272.6 Sympy [N/A]	2379
3.272.7 Maxima [N/A]	2379
3.272.8 Giac [F(-2)]	2379
3.272.9 Mupad [N/A]	2380

### 3.272.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx = -\frac{\sqrt{-1+cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b\sqrt{1-cx}} + \frac{\sqrt{-1+cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b\sqrt{1-cx}} + \operatorname{Int}\left(\frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `-Chi((a+b*arccosh(c*x))/b)*cosh(a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+Shi((a+b*arccosh(c*x))/b)*sinh(a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+Unintegrateable(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

### 3.272.2 Mathematica [N/A]

Not integrable

Time = 1.53 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x*(a + b*ArcCosh[c*x])), x]`

output `Integrate[Sqrt[1 - c^2*x^2]/(x*(a + b*ArcCosh[c*x])), x]`

### 3.272.3 Rubi [N/A]

Not integrable

Time = 0.63 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1 - c^2 x^2}}{x(a + \operatorname{barccosh}(cx))} dx$$

↓ 6369

$$\int \left( \frac{1}{x\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} - \frac{c^2 x}{\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} \right) dx$$

↓ 2009

$$\int \frac{1}{x\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} dx - \frac{\sqrt{cx - 1} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{b\sqrt{1 - cx}} + \frac{\sqrt{cx - 1} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{b\sqrt{1 - cx}}$$

input `Int[Sqrt[1 - c^2*x^2]/(x*(a + b*ArcCosh[c*x])), x]`

output `$Aborted`

**3.272.3.1 Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

**3.272.4 Maple [N/A] (verified)**

Not integrable

Time = 0.97 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2 + 1}}{x(a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x)),x)`

**3.272.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1 - c^2x^2}}{x(a + b \operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2 + 1}}{(b \operatorname{arccosh}(cx) + a)x} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(b*x*arccosh(c*x) + a*x), x)`

**3.272.6 Sympy [N/A]**

Not integrable

Time = 1.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{x(a+b\operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x/(a+b*acosh(c*x)),x)`output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x*(a + b*acosh(c*x))), x)`**3.272.7 Maxima [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)x} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)*x), x)`**3.272.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x)),x, algorithm="giac")`output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.272.  $\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx$

**3.272.9 Mupad [N/A]**

Not integrable

Time = 3.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x*(a + b*acosh(c*x))),x)`output `int((1 - c^2*x^2)^(1/2)/(x*(a + b*acosh(c*x))), x)`

$$3.273 \quad \int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$$

3.273.1 Optimal result . . . . .	2381
3.273.2 Mathematica [N/A] . . . . .	2381
3.273.3 Rubi [N/A] . . . . .	2382
3.273.4 Maple [N/A] (verified) . . . . .	2383
3.273.5 Fricas [N/A] . . . . .	2383
3.273.6 Sympy [N/A] . . . . .	2383
3.273.7 Maxima [N/A] . . . . .	2384
3.273.8 Giac [N/A] . . . . .	2384
3.273.9 Mupad [N/A] . . . . .	2384

### 3.273.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx = -\frac{c\sqrt{-1+cx} \log(a+b\operatorname{arccosh}(cx))}{b\sqrt{1-cx}} + \operatorname{Int}\left(\frac{1}{x^2\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `-c*ln(a+b*arccosh(c*x))*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+Unintegrable(1/x^2/(a+b*arccosh(c*x)))/(-c^2*x^2+1)^(1/2), x)`

### 3.273.2 Mathematica [N/A]

Not integrable

Time = 1.80 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x^2*(a + b*ArcCosh[c*x])), x]`

output `Integrate[Sqrt[1 - c^2*x^2]/(x^2*(a + b*ArcCosh[c*x])), x]`

---


$$3.273. \quad \int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$$

**3.273.3 Rubi [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+\text{barccosh}(cx))} dx$$

↓ 6369

$$\int \left( \frac{1}{x^2\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))} - \frac{c^2}{\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))} \right) dx$$

↓ 2009

$$\int \frac{1}{x^2\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))} dx - \frac{c\sqrt{cx-1}\log(a+\text{barccosh}(cx))}{b\sqrt{1-cx}}$$

input `Int[Sqrt[1 - c^2*x^2]/(x^2*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.273.3.1 Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_))^m_*((d_) + (e_.)*(x_)^2)^p_, x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

**3.273.4 Maple [N/A] (verified)**

Not integrable

Time = 0.92 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2 + 1}}{x^2(a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x)),x)`output `int((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x)),x)`**3.273.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{1 - c^2x^2}}{x^2(a + b \operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2 + 1}}{(b \operatorname{arccosh}(cx) + a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x)),x, algorithm="fricas")`output `integral(sqrt(-c^2*x^2 + 1)/(b*x^2*arccosh(c*x) + a*x^2), x)`**3.273.6 Sympy [N/A]**

Not integrable

Time = 1.66 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{1 - c^2x^2}}{x^2(a + b \operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-(cx - 1)(cx + 1)}}{x^2(a + b \operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x**2/(a+b*acosh(c*x)),x)`output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x**2*(a + b*acosh(c*x))), x)`



**3.273.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)*x^2), x)`

**3.273.8 Giac [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)*x^2), x)`

**3.273.9 Mupad [N/A]**

Not integrable

Time = 2.70 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x^2*(a + b*acosh(c*x))),x)`

output `int((1 - c^2*x^2)^(1/2)/(x^2*(a + b*acosh(c*x))), x)`

---

3.273.  $\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$

**3.274**  $\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\text{arccosh}(cx))} dx$

3.274.1 Optimal result . . . . .	2385
3.274.2 Mathematica [N/A] . . . . .	2385
3.274.3 Rubi [N/A] . . . . .	2386
3.274.4 Maple [N/A] (verified) . . . . .	2386
3.274.5 Fricas [N/A] . . . . .	2387
3.274.6 Sympy [N/A] . . . . .	2387
3.274.7 Maxima [N/A] . . . . .	2387
3.274.8 Giac [F(-2)] . . . . .	2388
3.274.9 Mupad [N/A] . . . . .	2388

**3.274.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\text{arccosh}(cx))} dx = \text{Int}\left(\frac{\sqrt{1-c^2x^2}}{x^3(a+b\text{arccosh}(cx))}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x)), x)`

**3.274.2 Mathematica [N/A]**

Not integrable

Time = 7.71 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\text{arccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\text{arccosh}(cx))} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x^3*(a + b*ArcCosh[c*x])), x]`

output `Integrate[Sqrt[1 - c^2*x^2]/(x^3*(a + b*ArcCosh[c*x])), x]`

**3.274.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{arccosh}(cx))} dx$$

input `Int[Sqrt[1 - c^2*x^2]/(x^3*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.274.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.274.4 Maple [N/A] (verified)**

Not integrable

Time = 1.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2+1}}{x^3(a+b\operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x)),x)`

**3.274.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)x^3} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(b*x^3*arccosh(c*x) + a*x^3), x)`

**3.274.6 Sympy [N/A]**

Not integrable

Time = 3.08 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{x^3(a+b \operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x**3/(a+b*acosh(c*x)),x)`

output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x**3*(a + b*acosh(c*x))), x)`

**3.274.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)x^3} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)*x^3), x)`

---

3.274.  $\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$

**3.274.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.274.9 Mupad [N/A]**

Not integrable

Time = 3.05 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x^3*(a + b*acosh(c*x))),x)`

output `int((1 - c^2*x^2)^(1/2)/(x^3*(a + b*acosh(c*x))), x)`

$$3.275 \quad \int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

3.275.1 Optimal result . . . . .	2389
3.275.2 Mathematica [N/A] . . . . .	2389
3.275.3 Rubi [N/A] . . . . .	2390
3.275.4 Maple [N/A] (verified) . . . . .	2390
3.275.5 Fricas [N/A] . . . . .	2391
3.275.6 Sympy [N/A] . . . . .	2391
3.275.7 Maxima [N/A] . . . . .	2391
3.275.8 Giac [N/A] . . . . .	2392
3.275.9 Mupad [N/A] . . . . .	2392

### 3.275.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x)), x)`

### 3.275.2 Mathematica [N/A]

Not integrable

Time = 1.25 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x^4*(a + b*ArcCosh[c*x])), x]`

output `Integrate[Sqrt[1 - c^2*x^2]/(x^4*(a + b*ArcCosh[c*x])), x]`

**3.275.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{arccosh}(cx))} dx$$

input `Int[Sqrt[1 - c^2*x^2]/(x^4*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.275.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.275.4 Maple [N/A] (verified)**

Not integrable

Time = 1.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2+1}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x)),x)`

**3.275.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)x^4} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(b*x^4*arccosh(c*x) + a*x^4), x)`

**3.275.6 Sympy [N/A]**

Not integrable

Time = 8.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{x^4(a+b\operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x**4/(a+b*acosh(c*x)),x)`

output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x**4*(a + b*acosh(c*x))), x)`

**3.275.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)x^4} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)*x^4), x)`

---

3.275.  $\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$



**3.275.8 Giac [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{barccosh}(cx))} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)x^4} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)*x^4), x)`

**3.275.9 Mupad [N/A]**

Not integrable

Time = 2.91 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{barccosh}(cx))} dx = \int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x^4*(a + b*acosh(c*x))),x)`

output `int((1 - c^2*x^2)^(1/2)/(x^4*(a + b*acosh(c*x))), x)`

**3.276**  $\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

3.276.1 Optimal result . . . . . 2393  
 3.276.2 Mathematica [A] (warning: unable to verify) . . . . . 2394  
 3.276.3 Rubi [A] (verified) . . . . . 2394  
 3.276.4 Maple [A] (verified) . . . . . 2396  
 3.276.5 Fricas [F] . . . . . 2396  
 3.276.6 Sympy [F] . . . . . 2397  
 3.276.7 Maxima [F] . . . . . 2397  
 3.276.8 Giac [F] . . . . . 2397  
 3.276.9 Mupad [F(-1)] . . . . . 2398

**3.276.1 Optimal result**

Integrand size = 28, antiderivative size = 397

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = -\frac{3\sqrt{1-cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

$$+ \frac{3\sqrt{1-cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

$$+ \frac{\sqrt{1-cx} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

$$- \frac{\sqrt{1-cx} \cosh\left(\frac{7a}{b}\right) \operatorname{Chi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

$$+ \frac{3\sqrt{1-cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

$$- \frac{3\sqrt{1-cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

$$- \frac{\sqrt{1-cx} \sinh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

$$+ \frac{\sqrt{1-cx} \sinh\left(\frac{7a}{b}\right) \operatorname{Shi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^4\sqrt{-1+cx}}$$

output 
$$\begin{aligned} & -3/64*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\cosh(a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +3/64*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\cosh(3*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +1/64*\text{Chi}(5*(a+b*\text{arccosh}(c*x))/b)*\cosh(5*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -1/64*\text{Chi}(7*(a+b*\text{arccosh}(c*x))/b)*\cosh(7*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +3/64*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*\sinh(a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -3/64*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*\sinh(3*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -1/64*\text{Shi}(5*(a+b*\text{arccosh}(c*x))/b)*\sinh(5*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +1/64*\text{Shi}(7*(a+b*\text{arccosh}(c*x))/b)*\sinh(7*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \end{aligned}$$

### 3.276.2 Mathematica [A] (warning: unable to verify)

Time = 0.71 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.54

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\text{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2}(-3\cosh(\frac{a}{b})\text{Chi}(\frac{a}{b}+\text{arccosh}(cx)) + 3\cosh(\frac{3a}{b})\text{Chi}(3(\frac{a}{b}+\text{arccosh}(cx))))}{64c^4\sqrt{(-1+cx)/(1+cx)}(b+b*c*x)}$$

input `Integrate[(x^3*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]),x]`

output 
$$\begin{aligned} & (\text{Sqrt}[1 - c^2*x^2]*(-3*\text{Cosh}[a/b]*\text{CoshIntegral}[a/b + \text{ArcCosh}[c*x]] + 3*\text{Cosh} \\ & [(3*a)/b]*\text{CoshIntegral}[3*(a/b + \text{ArcCosh}[c*x])] + \text{Cosh}[(5*a)/b]*\text{CoshIntegral} \\ & [5*(a/b + \text{ArcCosh}[c*x])] - \text{Cosh}[(7*a)/b]*\text{CoshIntegral}[7*(a/b + \text{ArcCosh}[c* \\ & x]]) + 3*\text{Sinh}[a/b]*\text{SinhIntegral}[a/b + \text{ArcCosh}[c*x]] - 3*\text{Sinh}[(3*a)/b]*\text{Sinh} \\ & \text{Integral}[3*(a/b + \text{ArcCosh}[c*x])] - \text{Sinh}[(5*a)/b]*\text{SinhIntegral}[5*(a/b + \text{Arc} \\ & \text{Cosh}[c*x])] + \text{Sinh}[(7*a)/b]*\text{SinhIntegral}[7*(a/b + \text{ArcCosh}[c*x])])/(64*c^4 \\ & * \text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(b + b*c*x) \end{aligned}$$

### 3.276.3 Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.56, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\text{arccosh}(cx)} dx$$

---

3.276.  $\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\text{arccosh}(cx)} dx$

$$\begin{aligned}
 & \downarrow 6367 \\
 & \frac{\sqrt{1-cx} \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^4\sqrt{cx-1}} \\
 & \downarrow 5971 \\
 & \frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{7a}{b} - \frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} - \frac{3 \cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} + \frac{3 \cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} \right)}{bc^4\sqrt{cx-1}} \\
 & \downarrow 2009 \\
 & \frac{\sqrt{1-cx} \left( \frac{3}{64} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{3}{64} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{64} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc^4\sqrt{cx-1}}
 \end{aligned}$$

input `Int[(x^3*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]),x]`

output `-((Sqrt[1 - c*x]*((3*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/64 - (3*Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/64 - (Cosh[(5*a)/b]*CoshIntegral[(5*(a + b*ArcCosh[c*x])/b])/64 + (Cosh[(7*a)/b]*CoshIntegral[(7*(a + b*ArcCosh[c*x])/b])/64 - (3*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/64 + (3*Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/64 + (Sinh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/64 - (Sinh[(7*a)/b]*SinhIntegral[(7*(a + b*ArcCosh[c*x])/b])/64))/(b*c^4*Sqrt[-1 + c*x]))`

### 3.276.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

---

3.276.  $\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.276.4 Maple [A] (verified)

Time = 0.53 (sec) , antiderivative size = 318, normalized size of antiderivative = 0.80

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(\text{Ei}_1(5\operatorname{arccosh}(cx)+\frac{5a}{b})e^{\frac{b\operatorname{arccosh}(cx)+5a}{b}}-\text{Ei}_1(7\operatorname{arccosh}(cx)+\frac{7a}{b})e^{\frac{b\operatorname{arccosh}(cx)+7a}{b}}-\dots\right)}{\dots}$

input `int(x^3*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `1/128*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(Ei(1,5*arccosh(c*x)+5*a/b)*exp((b*arccosh(c*x)+5*a)/b)-Ei(1,7*arccosh(c*x)+7*a/b)*exp((b*arccosh(c*x)+7*a)/b)-Ei(1,-7*arccosh(c*x)-7*a/b)*exp(-(-b*arccosh(c*x)+7*a)/b)+3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)-3*Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)-3*Ei(1,-arccosh(c*x)-a/b)*exp(-(-b*arccosh(c*x)+a)/b)+3*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(-b*arccosh(c*x)+3*a)/b)+Ei(1,-5*arccosh(c*x)-5*a/b)*exp(-(-b*arccosh(c*x)+5*a)/b))/(c*x+1)/c^4/(c*x-1)/b`

### 3.276.5 Fracas [F]

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x^3}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-c^2*x^5 - x^3)*sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x`

---

3.276.  $\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

**3.276.6 Sympy [F]**

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+\operatorname{barccosh}(cx)} dx = \int \frac{x^3(-(cx-1)(cx+1))^{3/2}}{a+b\operatorname{acosh}(cx)} dx$$

input `integrate(x**3*(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**3*(-(c*x - 1)*(c*x + 1))**(3/2)/(a + b*acosh(c*x)), x)`

**3.276.7 Maxima [F]**

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+\operatorname{barccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{3/2}x^3}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x^3/(b*arccosh(c*x) + a), x)`

**3.276.8 Giac [F]**

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+\operatorname{barccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{3/2}x^3}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x^3/(b*arccosh(c*x) + a), x)`

**3.276.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x^3(1-c^2x^2)^{3/2}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((x^3*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)),x)`output `int((x^3*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)), x)`

**3.277**  $\int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

3.277.1 Optimal result . . . . . 2399  
 3.277.2 Mathematica [A] (warning: unable to verify) . . . . . 2400  
 3.277.3 Rubi [A] (verified) . . . . . 2400  
 3.277.4 Maple [A] (verified) . . . . . 2402  
 3.277.5 Fricas [F] . . . . . 2402  
 3.277.6 Sympy [F] . . . . . 2403  
 3.277.7 Maxima [F] . . . . . 2403  
 3.277.8 Giac [F] . . . . . 2403  
 3.277.9 Mupad [F(-1)] . . . . . 2404

**3.277.1 Optimal result**

Integrand size = 28, antiderivative size = 339

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{1-cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \log(a+b\operatorname{arccosh}(cx))}{16bc^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^3\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \sinh\left(\frac{6a}{b}\right) \operatorname{Shi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}}$$



output  $\frac{1}{32}\text{Chi}\left(\frac{2(a+b\text{arccosh}(cx))}{b}\right)\cosh\left(\frac{2a}{b}\right)(-cx+1)^{1/2}/b/c^3/(cx-1)^{1/2} + \frac{1}{16}\text{Chi}\left(\frac{4(a+b\text{arccosh}(cx))}{b}\right)\cosh\left(\frac{4a}{b}\right)(-cx+1)^{1/2}/b/c^3/(cx-1)^{1/2} - \frac{1}{32}\text{Chi}\left(\frac{6(a+b\text{arccosh}(cx))}{b}\right)\cosh\left(\frac{6a}{b}\right)(-cx+1)^{1/2}/b/c^3/(cx-1)^{1/2} - \frac{1}{16}\ln(a+b\text{arccosh}(cx))(-cx+1)^{1/2}/b/c^3/(cx-1)^{1/2} - \frac{1}{32}\text{Shi}\left(\frac{2(a+b\text{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)(-cx+1)^{1/2}/b/c^3/(cx-1)^{1/2} - \frac{1}{16}\text{Shi}\left(\frac{4(a+b\text{arccosh}(cx))}{b}\right)\sinh\left(\frac{4a}{b}\right)(-cx+1)^{1/2}/b/c^3/(cx-1)^{1/2} + \frac{1}{32}\text{Shi}\left(\frac{6(a+b\text{arccosh}(cx))}{b}\right)\sinh\left(\frac{6a}{b}\right)(-cx+1)^{1/2}/b/c^3/(cx-1)^{1/2}$

### 3.277.2 Mathematica [A] (warning: unable to verify)

Time = 0.62 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.55

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\text{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2}\left(-\cosh\left(\frac{2a}{b}\right)\text{Chi}\left(2\left(\frac{a}{b}+\text{arccosh}(cx)\right)\right) - 2\cosh\left(\frac{4a}{b}\right)\text{Chi}\left(4\left(\frac{a}{b}+\text{arccosh}(cx)\right)\right) + \cosh\left(\frac{6a}{b}\right)\text{Chi}\left(6\left(\frac{a}{b}+\text{arccosh}(cx)\right)\right)\right)}{c^3}$$

input `Integrate[(x^2*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]), x]`

output  $\frac{-1/32*(\text{Sqrt}[1 - c^2*x^2]*(-(\text{Cosh}[(2*a)/b]*\text{CoshIntegral}[2*(a/b + \text{ArcCosh}[c*x])]) - 2*\text{Cosh}[(4*a)/b]*\text{CoshIntegral}[4*(a/b + \text{ArcCosh}[c*x])]) + \text{Cosh}[(6*a)/b]*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]) + 2*\text{Log}[a + b*\text{ArcCosh}[c*x]) + \text{Sinh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])]) + 2*\text{Sinh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])]) - \text{Sinh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])])]/(c^3*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(b + b*c*x))$

### 3.277.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.56, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.277.  $\int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\text{arccosh}(cx)} dx$

$$\begin{aligned}
 & \int \frac{x^2(1-c^2x^2)^{3/2}}{a+\operatorname{arccosh}(cx)} dx \\
 & \quad \downarrow \text{6367} \\
 & \frac{\sqrt{1-cx} \int \frac{\cosh^2\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^3\sqrt{cx-1}} \\
 & \quad \downarrow \text{5971} \\
 & \frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{6a}{b}-\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{4a}{b}-\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{2a}{b}-\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} \right) + \frac{1}{16(a+b\operatorname{arccosh}(cx))}}{bc^3\sqrt{cx-1}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{1-cx} \left( -\frac{1}{32} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{16} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{32} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc^3\sqrt{cx-1}}
 \end{aligned}$$

input `Int[(x^2*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]),x]`

output `-((Sqrt[1 - c*x]*(-1/32*(Cosh[(2*a)/b]*CoshIntegral[(2*(a + b*ArcCosh[c*x])/b]) - (Cosh[(4*a)/b]*CoshIntegral[(4*(a + b*ArcCosh[c*x])/b])/16 + (Cosh[(6*a)/b]*CoshIntegral[(6*(a + b*ArcCosh[c*x])/b])/32 + Log[a + b*ArcCosh[c*x])/16 + (Sinh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x])/b])/32 + (Sinh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x])/b])/16 - (Sinh[(6*a)/b]*SinhIntegral[(6*(a + b*ArcCosh[c*x])/b])/32))/(b*c^3*Sqrt[-1 + c*x])`

### 3.277.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

$$3.277. \quad \int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$$

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.277.4 Maple [A] (verified)

Time = 0.71 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.88

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-4\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))-4\ln(a+b\operatorname{arccosh}(cx))cx+Ei_1(6\operatorname{arccosh}(cx))\right)}{\dots}$

input `int(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\begin{aligned} & -1/64*(-c^2*x^2+1)^{(1/2)}*(-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*c*x+c^2*x^2-1)*(-4* \\ & (c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*\ln(a+b*\operatorname{arccosh}(c*x))-4*\ln(a+b*\operatorname{arccosh}(c*x))*c* \\ & x+Ei(1,6*\operatorname{arccosh}(c*x)+6*a/b)*\exp((b*\operatorname{arccosh}(c*x)+6*a)/b)+Ei(1,-6*\operatorname{arccosh}(c \\ & *x)-6*a/b)*\exp(-(-b*\operatorname{arccosh}(c*x)+6*a)/b)-2*Ei(1,4*\operatorname{arccosh}(c*x)+4*a/b)*\exp( \\ & (b*\operatorname{arccosh}(c*x)+4*a)/b)-Ei(1,2*\operatorname{arccosh}(c*x)+2*a/b)*\exp((b*\operatorname{arccosh}(c*x)+2*a \\ & )/b)-Ei(1,-2*\operatorname{arccosh}(c*x)-2*a/b)*\exp(-(-b*\operatorname{arccosh}(c*x)+2*a)/b)-2*Ei(1,-4*a \\ & rccosh(c*x)-4*a/b)*\exp(-(-b*\operatorname{arccosh}(c*x)+4*a)/b))/(c*x+1)/c^3/(c*x-1)/b \end{aligned}$$

### 3.277.5 Fracas [F]

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{3/2}x^2}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-c^2*x^4 - x^2)*sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

**3.277.6 Sympy [F]**

$$\int \frac{x^2(1 - c^2x^2)^{3/2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{x^2(-(cx - 1)(cx + 1))^{3/2}}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate(x**2*(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**2*(-(c*x - 1)*(c*x + 1))**(3/2)/(a + b*acosh(c*x)), x)`

**3.277.7 Maxima [F]**

$$\int \frac{x^2(1 - c^2x^2)^{3/2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{(-c^2x^2 + 1)^{3/2}x^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x^2/(b*arccosh(c*x) + a), x)`

**3.277.8 Giac [F]**

$$\int \frac{x^2(1 - c^2x^2)^{3/2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{(-c^2x^2 + 1)^{3/2}x^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x^2/(b*arccosh(c*x) + a), x)`

**3.277.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x^2(1-c^2x^2)^{3/2}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((x^2*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)),x)`output `int((x^2*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)), x)`

**3.278**  $\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

3.278.1 Optimal result . . . . . 2405  
 3.278.2 Mathematica [A] (warning: unable to verify) . . . . . 2406  
 3.278.3 Rubi [A] (verified) . . . . . 2406  
 3.278.4 Maple [A] (verified) . . . . . 2407  
 3.278.5 Fricas [F] . . . . . 2408  
 3.278.6 Sympy [F] . . . . . 2408  
 3.278.7 Maxima [F] . . . . . 2409  
 3.278.8 Giac [F] . . . . . 2409  
 3.278.9 Mupad [F(-1)] . . . . . 2409

**3.278.1 Optimal result**

Integrand size = 26, antiderivative size = 297

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = -\frac{\sqrt{1-cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8bc^2\sqrt{-1+cx}} + \frac{3\sqrt{1-cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^2\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^2\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8bc^2\sqrt{-1+cx}} - \frac{3\sqrt{1-cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^2\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \sinh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^2\sqrt{-1+cx}}$$

```
output -1/8*Chi((a+b*arccosh(c*x))/b)*cosh(a/b)*(-c*x+1)^(1/2)/b/c^2/(c*x-1)^(1/2)
)+3/16*Chi(3*(a+b*arccosh(c*x))/b)*cosh(3*a/b)*(-c*x+1)^(1/2)/b/c^2/(c*x-1)
)^(1/2)-1/16*Chi(5*(a+b*arccosh(c*x))/b)*cosh(5*a/b)*(-c*x+1)^(1/2)/b/c^2/
(c*x-1)^(1/2)+1/8*Shi((a+b*arccosh(c*x))/b)*sinh(a/b)*(-c*x+1)^(1/2)/b/c^2
/(c*x-1)^(1/2)-3/16*Shi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)*(-c*x+1)^(1/2)
/b/c^2/(c*x-1)^(1/2)+1/16*Shi(5*(a+b*arccosh(c*x))/b)*sinh(5*a/b)*(-c*x+1)
)^(1/2)/b/c^2/(c*x-1)^(1/2)
```

3.278.  $\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

**3.278.2 Mathematica [A] (warning: unable to verify)**

Time = 0.60 (sec) , antiderivative size = 172, normalized size of antiderivative = 0.58

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2}(-2\cosh\left(\frac{a}{b}\right)\operatorname{Chi}\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right)+3\cosh\left(\frac{3a}{b}\right)\operatorname{Chi}\left(3\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right)\right)}{a+b\operatorname{arccosh}(cx)}$$

input `Integrate[(x*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[1 - c^2*x^2]*(-2*Cosh[a/b]*CoshIntegral[a/b + ArcCosh[c*x]] + 3*Cosh[(3*a)/b]*CoshIntegral[3*(a/b + ArcCosh[c*x])] - Cosh[(5*a)/b]*CoshIntegral[5*(a/b + ArcCosh[c*x])] + 2*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] - 3*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] + Sinh[(5*a)/b]*SinhIntegral[5*(a/b + ArcCosh[c*x])])/(16*c^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(b + b*c*x))`

**3.278.3 Rubi [A] (verified)**Time = 0.58 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.59, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$$

↓ 6367

$$\frac{\sqrt{1-cx} \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^2\sqrt{cx-1}}$$

↓ 5971

$$\frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{3\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} + \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{bc^2\sqrt{cx-1}}$$

↓ 2009

---

3.278.  $\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

$$\frac{\sqrt{1-cx} \left( \frac{1}{8} \cosh\left(\frac{a}{b}\right) \text{Chi}\left(\frac{a+b\text{arccosh}(cx)}{b}\right) - \frac{3}{16} \cosh\left(\frac{3a}{b}\right) \text{Chi}\left(\frac{3(a+b\text{arccosh}(cx))}{b}\right) + \frac{1}{16} \cosh\left(\frac{5a}{b}\right) \text{Chi}\left(\frac{5(a+b\text{arccosh}(cx))}{b}\right) \right)}{bc}$$

input `Int[(x*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]),x]`

output `-((Sqrt[1 - c*x]*((Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/8 - (3*Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/16 + (Cosh[(5*a)/b]*CoshIntegral[(5*(a + b*ArcCosh[c*x])/b])/16 - (Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/8 + (3*Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/16 - (Sinh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/16))/(b*c^2*Sqrt[-1 + c*x]))`

### 3.278.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.278.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 252, normalized size of antiderivative = 0.85

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(\text{Ei}_1\left(5\text{arccosh}(cx)+\frac{5a}{b}\right)e^{\frac{b\text{arccosh}(cx)+5a}{b}}+\text{Ei}_1\left(-5\text{arccosh}(cx)-\frac{5a}{b}\right)e^{-\frac{b\text{arccosh}(cx)+5a}{b}}\right)}{bc}$

3.278.  $\int \frac{x(1-c^2x^2)^{3/2}}{a+b\text{arccosh}(cx)} dx$



input `int(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-1/32*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(Ei(1,5*arccosh(c*x)+5*a/b)*exp((b*arccosh(c*x)+5*a)/b)+Ei(1,-5*arccosh(c*x)-5*a/b)*exp(-(-b*arccosh(c*x)+5*a)/b)-3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)+2*Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)+2*Ei(1,-arccosh(c*x)-a/b)*exp(-(-b*arccosh(c*x)+a)/b)-3*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(-b*arccosh(c*x)+3*a)/b))/(c*x+1)/c^2/(c*x-1)/b`

### 3.278.5 Fracas [F]

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-(c^2*x^3 - x)*sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

### 3.278.6 Sympy [F]

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x(-(cx-1)(cx+1))^{\frac{3}{2}}}{a+b\operatorname{acosh}(cx)} dx$$

input `integrate(x*(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(x*(-(c*x - 1)*(c*x + 1))**(3/2)/(a + b*acosh(c*x)), x)`

**3.278.7 Maxima [F]**

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+\operatorname{barccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x}{b \operatorname{arcosh}(cx)+a} dx$$

input `integrate(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x/(b*arccosh(c*x) + a), x)`

**3.278.8 Giac [F]**

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+\operatorname{barccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x}{b \operatorname{arcosh}(cx)+a} dx$$

input `integrate(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x/(b*arccosh(c*x) + a), x)`

**3.278.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(1-c^2x^2)^{3/2}}{a+\operatorname{barccosh}(cx)} dx = \int \frac{x(1-c^2x^2)^{3/2}}{a+b \operatorname{acosh}(cx)} dx$$

input `int((x*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)),x)`

output `int((x*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)), x)`

**3.279**  $\int \frac{(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

3.279.1 Optimal result . . . . . 2410  
 3.279.2 Mathematica [A] (warning: unable to verify) . . . . . 2411  
 3.279.3 Rubi [A] (verified) . . . . . 2411  
 3.279.4 Maple [A] (verified) . . . . . 2413  
 3.279.5 Fricas [F] . . . . . 2413  
 3.279.6 Sympy [F] . . . . . 2413  
 3.279.7 Maxima [F] . . . . . 2414  
 3.279.8 Giac [F] . . . . . 2414  
 3.279.9 Mupad [F(-1)] . . . . . 2414

**3.279.1 Optimal result**

Integrand size = 25, antiderivative size = 239

$$\int \frac{(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{1-cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2bc\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8bc\sqrt{-1+cx}} - \frac{3\sqrt{1-cx} \log(a+b\operatorname{arccosh}(cx))}{8bc\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2bc\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8bc\sqrt{-1+cx}}$$

```
output 1/2*Chi(2*(a+b*arccosh(c*x))/b)*cosh(2*a/b)*(-c*x+1)^(1/2)/b/c/(c*x-1)^(1/2)-1/8*Chi(4*(a+b*arccosh(c*x))/b)*cosh(4*a/b)*(-c*x+1)^(1/2)/b/c/(c*x-1)^(1/2)-3/8*ln(a+b*arccosh(c*x))*(-c*x+1)^(1/2)/b/c/(c*x-1)^(1/2)-1/2*Shi(2*(a+b*arccosh(c*x))/b)*sinh(2*a/b)*(-c*x+1)^(1/2)/b/c/(c*x-1)^(1/2)+1/8*Shi(4*(a+b*arccosh(c*x))/b)*sinh(4*a/b)*(-c*x+1)^(1/2)/b/c/(c*x-1)^(1/2)
```

**3.279.2 Mathematica [A] (warning: unable to verify)**

Time = 0.55 (sec) , antiderivative size = 147, normalized size of antiderivative = 0.62

$$\int \frac{(1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx = \frac{\sqrt{1 - c^2 x^2} \left( -4 \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) + \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(4\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) + 3 \log(a + b \operatorname{arccosh}(cx)) \right)}{8bc \sqrt{\frac{-1+cx}{1+cx}} (1 + cx)}$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(a + b*ArcCosh[c*x]),x]`output `-1/8*(Sqrt[1 - c^2*x^2]*(-4*Cosh[(2*a)/b]*CoshIntegral[2*(a/b + ArcCosh[c*x])] + Cosh[(4*a)/b]*CoshIntegral[4*(a/b + ArcCosh[c*x])] + 3*Log[a + b*ArcCosh[c*x]] + 4*Sinh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[c*x])] - Sinh[(4*a)/b]*SinhIntegral[4*(a/b + ArcCosh[c*x])]))/(b*c*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x))`**3.279.3 Rubi [A] (verified)**Time = 0.47 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.59, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$ , Rules used = {6321, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx \\ & \quad \downarrow \text{6321} \\ & - \frac{\sqrt{1 - cx} \int \frac{\sinh^4\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{a + b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{bc \sqrt{cx - 1}} \\ & \quad \downarrow \text{3042} \\ & - \frac{\sqrt{1 - cx} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a + b \operatorname{arccosh}(cx))}{b}\right)^4}{a + b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{bc \sqrt{cx - 1}} \\ & \quad \downarrow \text{3793} \end{aligned}$$

---

3.279.  $\int \frac{(1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx$

$$\frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2(a+b\operatorname{arccosh}(cx))} + \frac{3}{8(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{bc\sqrt{cx-1}}$$

↓ 2009

$$\frac{\sqrt{1-cx} \left( -\frac{1}{2} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{8} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{2} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc\sqrt{cx-1}}$$

input `Int[(1 - c^2*x^2)^(3/2)/(a + b*ArcCosh[c*x]),x]`

output `-((Sqrt[1 - c*x]*(-1/2*(Cosh[(2*a)/b]*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]) + (Cosh[(4*a)/b]*CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8 + (3*Log[a + b*ArcCosh[c*x]])/8 + (Sinh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/2 - (Sinh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8))/(b*c*Sqrt[-1 + c*x])`

### 3.279.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

---

3.279.  $\int \frac{(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

**3.279.4 Maple [A] (verified)**

Time = 0.86 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.97

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-6\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))-6\ln(a+b\operatorname{arccosh}(cx))cx+Ei_1(4\operatorname{arccosh}(cx))\right)}{c^2x^2+1}$

```
input int((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output -1/16*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(a+b*arccosh(c*x))-6*ln(a+b*arccosh(c*x))*c*x+Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)+Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b)-4*Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)-4*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b))/(c*x-1)/(c*x+1)/c/b
```

**3.279.5 Fracas [F]**

$$\int \frac{(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{3/2}}{b\operatorname{arccosh}(cx)+a} dx$$

```
input integrate((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fracas")
```

```
output integral((-c^2*x^2 + 1)^(3/2)/(b*arccosh(c*x) + a), x)
```

**3.279.6 Sympy [F]**

$$\int \frac{(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-(cx-1)(cx+1))^{3/2}}{a+b\operatorname{acosh}(cx)} dx$$

```
input integrate((-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)
```

```
output Integral((-c*x - 1)*(c*x + 1)**(3/2)/(a + b*acosh(c*x)), x)
```

---

3.279.  $\int \frac{(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx$

**3.279.7 Maxima [F]**

$$\int \frac{(1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(3/2)/(b*arccosh(c*x) + a), x)`

**3.279.8 Giac [F]**

$$\int \frac{(1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)/(b*arccosh(c*x) + a), x)`

**3.279.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{a + b \operatorname{acosh}(cx)} dx$$

input `int((1 - c^2*x^2)^(3/2)/(a + b*acosh(c*x)),x)`

output `int((1 - c^2*x^2)^(3/2)/(a + b*acosh(c*x)), x)`

$$3.280 \quad \int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))} dx$$

3.280.1 Optimal result	2415
3.280.2 Mathematica [N/A]	2416
3.280.3 Rubi [N/A]	2416
3.280.4 Maple [N/A] (verified)	2417
3.280.5 Fricas [N/A]	2417
3.280.6 Sympy [N/A]	2418
3.280.7 Maxima [N/A]	2418
3.280.8 Giac [F(-2)]	2418
3.280.9 Mupad [N/A]	2419

### 3.280.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\begin{aligned} \int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))} dx = & -\frac{5\sqrt{-1+cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4b\sqrt{1-cx}} \\ & + \frac{\sqrt{-1+cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b\sqrt{1-cx}} \\ & + \frac{5\sqrt{-1+cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4b\sqrt{1-cx}} \\ & - \frac{\sqrt{-1+cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b\sqrt{1-cx}} \\ & + \operatorname{Int}\left(\frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}, x\right) \end{aligned}$$

output `-5/4*Chi((a+b*arccosh(c*x))/b)*cosh(a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+1/4*Chi(3*(a+b*arccosh(c*x))/b)*cosh(3*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+5/4*Shi((a+b*arccosh(c*x))/b)*sinh(a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)-1/4*Shi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+Unintegrable(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

---


$$3.280. \quad \int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))} dx$$



**3.280.2 Mathematica [N/A]**

Not integrable

Time = 1.63 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(x*(a + b*ArcCosh[c*x])), x]`output `Integrate[(1 - c^2*x^2)^(3/2)/(x*(a + b*ArcCosh[c*x])), x]`**3.280.3 Rubi [N/A]**

Not integrable

Time = 0.98 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))} dx$$

↓ 6369

$$\int \left( -\frac{2c^2 x}{\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} + \frac{1}{x\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} + \frac{c^4 x^3}{\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} \right) dx$$

↓ 2009

$$\frac{\int \frac{1}{x\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} dx - \frac{5\sqrt{cx - 1} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{4b\sqrt{1 - cx}} + \frac{\sqrt{cx - 1} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{4b\sqrt{1 - cx}} + \frac{5\sqrt{cx - 1} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{4b\sqrt{1 - cx}} - \frac{\sqrt{cx - 1} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{4b\sqrt{1 - cx}}}{}$$

---

3.280.  $\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))} dx$

input `Int[(1 - c^2*x^2)^(3/2)/(x*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

### 3.280.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.280.4 Maple [N/A] (verified)

Not integrable

Time = 1.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{x(a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x)),x)`

### 3.280.5 Fracas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arccosh}(cx) + a)x} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x)),x, algorithm="fricas")`

---

3.280.  $\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))} dx$

output `integral((-c^2*x^2 + 1)^(3/2)/(b*x*arccosh(c*x) + a*x), x)`

### 3.280.6 Sympy [N/A]

Not integrable

Time = 10.39 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-(cx - 1)(cx + 1))^{\frac{3}{2}}}{x(a + b \operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/x/(a+b*acosh(c*x)),x)`

output `Integral((-c*x - 1)*(c*x + 1)**(3/2)/(x*(a + b*acosh(c*x))), x)`

### 3.280.7 Maxima [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)x} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)*x), x)`

### 3.280.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.280.9 Mupad [N/A]

Not integrable

Time = 2.91 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x(a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x*(a + b*acosh(c*x))),x)`

output `int((1 - c^2*x^2)^(3/2)/(x*(a + b*acosh(c*x))), x)`

**3.281**  $\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$

3.281.1 Optimal result . . . . . 2420  
 3.281.2 Mathematica [N/A] . . . . . 2421  
 3.281.3 Rubi [N/A] . . . . . 2421  
 3.281.4 Maple [N/A] (verified) . . . . . 2422  
 3.281.5 Fricas [N/A] . . . . . 2422  
 3.281.6 Sympy [N/A] . . . . . 2423  
 3.281.7 Maxima [N/A] . . . . . 2423  
 3.281.8 Giac [N/A] . . . . . 2423  
 3.281.9 Mupad [N/A] . . . . . 2424

**3.281.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))} dx = \frac{c\sqrt{-1+cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2b\sqrt{1-cx}} - \frac{3c\sqrt{-1+cx} \log(a+b\operatorname{arccosh}(cx))}{2b\sqrt{1-cx}} - \frac{c\sqrt{-1+cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2b\sqrt{1-cx}} + \operatorname{Int}\left(\frac{1}{x^2\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

```
output 1/2*c*Chi(2*(a+b*arccosh(c*x))/b)*cosh(2*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)-3/2*c*ln(a+b*arccosh(c*x))*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)-1/2*c*Shi(2*(a+b*arccosh(c*x))/b)*sinh(2*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+Unintegrate(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)
```

**3.281.2 Mathematica [N/A]**

Not integrable

Time = 2.41 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2(a + \operatorname{barccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x^2(a + \operatorname{barccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(x^2*(a + b*ArcCosh[c*x])), x]`output `Integrate[(1 - c^2*x^2)^(3/2)/(x^2*(a + b*ArcCosh[c*x])), x]`**3.281.3 Rubi [N/A]**

Not integrable

Time = 0.81 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2(a + \operatorname{barccosh}(cx))} dx$$

↓ 6369

$$\int \left( -\frac{2c^2}{\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} + \frac{1}{x^2 \sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} + \frac{c^4 x^2}{\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} \right) dx$$

↓ 2009

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} dx + \frac{c\sqrt{cx - 1} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{2b\sqrt{1 - cx}} - \frac{c\sqrt{cx - 1} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{2b\sqrt{1 - cx}} - \frac{3c\sqrt{cx - 1} \log(a + \operatorname{barccosh}(cx))}{2b\sqrt{1 - cx}}$$

input `Int[(1 - c^2*x^2)^(3/2)/(x^2*(a + b*ArcCosh[c*x])), x]`output `$Aborted`

---

3.281.  $\int \frac{(1 - c^2 x^2)^{3/2}}{x^2(a + \operatorname{barccosh}(cx))} dx$

**3.281.3.1** Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

**3.281.4** Maple [N/A] (verified)

Not integrable

Time = 0.90 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{x^2(a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x)),x)`

**3.281.5** Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{(1 - c^2x^2)^{3/2}}{x^2(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arccosh}(cx) + a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((-c^2*x^2 + 1)^(3/2)/(b*x^2*arccosh(c*x) + a*x^2), x)`

**3.281.6 Sympy [N/A]**

Not integrable

Time = 11.35 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{-(cx - 1)(cx + 1)^{\frac{3}{2}}}{x^2 (a + b \operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/x**2/(a+b*acosh(c*x)),x)`output `Integral((-c*x - 1)*(c*x + 1)**(3/2)/(x**2*(a + b*acosh(c*x))), x)`**3.281.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)*x^2), x)`**3.281.8 Giac [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)*x^2), x)`

---

3.281.  $\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))} dx$



**3.281.9 Mupad [N/A]**

Not integrable

Time = 3.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x^2*(a + b*acosh(c*x))),x)`output `int((1 - c^2*x^2)^(3/2)/(x^2*(a + b*acosh(c*x))), x)`

**3.282**  $\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$

3.282.1 Optimal result . . . . . 2425  
 3.282.2 Mathematica [N/A] . . . . . 2425  
 3.282.3 Rubi [N/A] . . . . . 2426  
 3.282.4 Maple [N/A] (verified) . . . . . 2426  
 3.282.5 Fricas [N/A] . . . . . 2427  
 3.282.6 Sympy [N/A] . . . . . 2427  
 3.282.7 Maxima [N/A] . . . . . 2427  
 3.282.8 Giac [F(-2)] . . . . . 2428  
 3.282.9 Mupad [N/A] . . . . . 2428

**3.282.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x)), x)`

**3.282.2 Mathematica [N/A]**

Not integrable

Time = 8.16 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx = \int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(x^3*(a + b*ArcCosh[c*x])), x]`

output `Integrate[(1 - c^2*x^2)^(3/2)/(x^3*(a + b*ArcCosh[c*x])), x]`

---

3.282.  $\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$

**3.282.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + \operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + \operatorname{arccosh}(cx))} dx$$

input `Int[(1 - c^2*x^2)^(3/2)/(x^3*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.282.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.282.4 Maple [N/A] (verified)**

Not integrable

Time = 1.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 x^2 + 1)^{3/2}}{x^3 (a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x)),x)`

---

3.282.  $\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$

**3.282.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)x^3} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="fricas")`output `integral((-c^2*x^2 + 1)^(3/2)/(b*x^3*arccosh(c*x) + a*x^3), x)`**3.282.6 Sympy [N/A]**

Not integrable

Time = 27.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-(cx - 1)(cx + 1))^{\frac{3}{2}}}{x^3(a + b \operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/x**3/(a+b*acosh(c*x)),x)`output `Integral((-(c*x - 1)*(c*x + 1))**(3/2)/(x**3*(a + b*acosh(c*x))), x)`**3.282.7 Maxima [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)x^3} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)*x^3), x)`

---

3.282.  $\int \frac{(1 - c^2 x^2)^{3/2}}{x^3(a + b \operatorname{arccosh}(cx))} dx$

**3.282.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.282.9 Mupad [N/A]**

Not integrable

Time = 3.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x^3*(a + b*acosh(c*x))),x)`

output `int((1 - c^2*x^2)^(3/2)/(x^3*(a + b*acosh(c*x))), x)`

$$3.283 \quad \int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

3.283.1 Optimal result	2429
3.283.2 Mathematica [N/A]	2429
3.283.3 Rubi [N/A]	2430
3.283.4 Maple [N/A] (verified)	2430
3.283.5 Fricas [N/A]	2431
3.283.6 Sympy [N/A]	2431
3.283.7 Maxima [N/A]	2431
3.283.8 Giac [N/A]	2432
3.283.9 Mupad [N/A]	2432

### 3.283.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x)), x)`

### 3.283.2 Mathematica [N/A]

Not integrable

Time = 1.34 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(x^4*(a + b*ArcCosh[c*x])), x]`

output `Integrate[(1 - c^2*x^2)^(3/2)/(x^4*(a + b*ArcCosh[c*x])), x]`

---


$$3.283. \quad \int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

**3.283.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + \operatorname{barccosh}(cx))} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + \operatorname{barccosh}(cx))} dx$$

input `Int[(1 - c^2*x^2)^(3/2)/(x^4*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.283.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.283.4 Maple [N/A] (verified)**

Not integrable

Time = 1.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 x^2 + 1)^{3/2}}{x^4 (a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x)),x)`

---

3.283.  $\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$

**3.283.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.14

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)x^4} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="fricas")`output `integral((-c^2*x^2 + 1)^(3/2)/(b*x^4*arccosh(c*x) + a*x^4), x)`**3.283.6 Sympy [N/A]**

Not integrable

Time = 71.91 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-(cx - 1)(cx + 1))^{\frac{3}{2}}}{x^4(a + b \operatorname{acosh}(cx))} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/x**4/(a+b*acosh(c*x)),x)`output `Integral((-(c*x - 1)*(c*x + 1))**(3/2)/(x**4*(a + b*acosh(c*x))), x)`**3.283.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)x^4} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)*x^4), x)`

---

3.283.  $\int \frac{(1 - c^2 x^2)^{3/2}}{x^4(a + b \operatorname{arccosh}(cx))} dx$



**3.283.8 Giac [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arccosh}(cx) + a) x^4} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)*x^4), x)`**3.283.9 Mupad [N/A]**

Not integrable

Time = 2.96 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x^4*(a + b*acosh(c*x))),x)`output `int((1 - c^2*x^2)^(3/2)/(x^4*(a + b*acosh(c*x))), x)`

**3.284** 
$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$$

3.284.1 Optimal result . . . . . 2433  
 3.284.2 Mathematica [A] (warning: unable to verify) . . . . . 2434  
 3.284.3 Rubi [A] (verified) . . . . . 2434  
 3.284.4 Maple [A] (verified) . . . . . 2436  
 3.284.5 Fricas [F] . . . . . 2436  
 3.284.6 Sympy [F(-1)] . . . . . 2437  
 3.284.7 Maxima [F] . . . . . 2437  
 3.284.8 Giac [F] . . . . . 2437  
 3.284.9 Mupad [F(-1)] . . . . . 2438

**3.284.1 Optimal result**

Integrand size = 28, antiderivative size = 397

$$\begin{aligned} \int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = & -\frac{3\sqrt{1-cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{128bc^4\sqrt{-1+cx}} \\ & + \frac{\sqrt{1-cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^4\sqrt{-1+cx}} \\ & - \frac{3\sqrt{1-cx} \cosh\left(\frac{7a}{b}\right) \operatorname{Chi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{256bc^4\sqrt{-1+cx}} \\ & + \frac{\sqrt{1-cx} \cosh\left(\frac{9a}{b}\right) \operatorname{Chi}\left(\frac{9(a+b\operatorname{arccosh}(cx))}{b}\right)}{256bc^4\sqrt{-1+cx}} \\ & + \frac{3\sqrt{1-cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{128bc^4\sqrt{-1+cx}} \\ & - \frac{\sqrt{1-cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^4\sqrt{-1+cx}} \\ & + \frac{3\sqrt{1-cx} \sinh\left(\frac{7a}{b}\right) \operatorname{Shi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{256bc^4\sqrt{-1+cx}} \\ & - \frac{\sqrt{1-cx} \sinh\left(\frac{9a}{b}\right) \operatorname{Shi}\left(\frac{9(a+b\operatorname{arccosh}(cx))}{b}\right)}{256bc^4\sqrt{-1+cx}} \end{aligned}$$

---

3.284. 
$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$$

output 
$$\begin{aligned} & -3/128*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\cosh(a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +1/32*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\cosh(3*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -3/256*\text{Chi}(7*(a+b*\text{arccosh}(c*x))/b)*\cosh(7*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +1/256*\text{Chi}(9*(a+b*\text{arccosh}(c*x))/b)*\cosh(9*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +3/128*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*\sinh(a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -1/32*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*\sinh(3*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & +3/256*\text{Shi}(7*(a+b*\text{arccosh}(c*x))/b)*\sinh(7*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \\ & -1/256*\text{Shi}(9*(a+b*\text{arccosh}(c*x))/b)*\sinh(9*a/b)*(-c*x+1)^{(1/2)}/b/c^4/(c*x-1)^{(1/2)} \end{aligned}$$

### 3.284.2 Mathematica [A] (warning: unable to verify)

Time = 1.05 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.54

$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2}(-6\cosh(\frac{a}{b})\text{Chi}(\frac{a}{b}+\text{arccosh}(cx)) + 8\cosh(\frac{3a}{b})\text{Chi}(3(\frac{a}{b}+\text{arccosh}(cx))))}{256c^4\sqrt{(-1+cx)/(1+cx)}}(b+b*cx)$$

input `Integrate[(x^3*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x]),x]`

output 
$$\begin{aligned} & (\text{Sqrt}[1 - c^2*x^2]*(-6*\text{Cosh}[a/b]*\text{CoshIntegral}[a/b + \text{ArcCosh}[c*x]] + 8*\text{Cosh} \\ & [(3*a)/b]*\text{CoshIntegral}[3*(a/b + \text{ArcCosh}[c*x])] - 3*\text{Cosh}[(7*a)/b]*\text{CoshIntegral}[7*(a/b + \text{ArcCosh}[c*x])] \\ & + \text{Cosh}[(9*a)/b]*\text{CoshIntegral}[9*(a/b + \text{ArcCosh}[c*x])] + 6*\text{Sinh}[a/b]*\text{SinhIntegral}[a/b + \text{ArcCosh}[c*x]] \\ & - 8*\text{Sinh}[(3*a)/b]*\text{SinhIntegral}[3*(a/b + \text{ArcCosh}[c*x])] + 3*\text{Sinh}[(7*a)/b]*\text{SinhIntegral}[7*(a/b + \text{ArcCosh}[c*x])] \\ & - \text{Sinh}[(9*a)/b]*\text{SinhIntegral}[9*(a/b + \text{ArcCosh}[c*x])])/(256*c^4*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(b + b*c*x)) \end{aligned}$$

### 3.284.3 Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.56, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx$$

---

3.284.  $\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx$

$$\begin{aligned}
 & \downarrow 6367 \\
 & \frac{\sqrt{1-cx} \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^6\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^4\sqrt{cx-1}} \\
 & \downarrow 5971 \\
 & \frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{9a}{b} - \frac{9(a+b\operatorname{arccosh}(cx))}{b}\right)}{256(a+b\operatorname{arccosh}(cx))} - \frac{3 \cosh\left(\frac{7a}{b} - \frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{256(a+b\operatorname{arccosh}(cx))} + \frac{\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} - \frac{3 \cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{128(a+b\operatorname{arccosh}(cx))} \right)}{bc^4\sqrt{cx-1}} \\
 & \downarrow 2009 \\
 & \frac{\sqrt{1-cx} \left( -\frac{3}{128} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) + \frac{1}{32} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{256} \cosh\left(\frac{7a}{b}\right) \operatorname{Chi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc^4\sqrt{cx-1}}
 \end{aligned}$$

input `Int[(x^3*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[1 - c*x]*((-3*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/128 + (Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/32 - (3*Cosh[(7*a)/b]*CoshIntegral[(7*(a + b*ArcCosh[c*x])/b])/256 + (Cosh[(9*a)/b]*CoshIntegral[(9*(a + b*ArcCosh[c*x])/b])/256 + (3*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/128 - (Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/32 + (3*Sinh[(7*a)/b]*SinhIntegral[(7*(a + b*ArcCosh[c*x])/b])/256 - (Sinh[(9*a)/b]*SinhIntegral[(9*(a + b*ArcCosh[c*x])/b])/256))/(b*c^4*Sqrt[-1 + c*x])`

### 3.284.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.284.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 320, normalized size of antiderivative = 0.81

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(3\operatorname{Ei}_1\left(7\operatorname{arccosh}(cx)+\frac{7a}{b}\right)e^{\frac{b\operatorname{arccosh}(cx)+7a}{b}}-\operatorname{Ei}_1\left(9\operatorname{arccosh}(cx)+\frac{9a}{b}\right)e^{\frac{b\operatorname{arccosh}(cx)+9a}{b}}\right)}{\dots}$

input `int(x^3*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-1/512*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(3*Ei(1,7*arccosh(c*x)+7*a/b)*exp((b*arccosh(c*x)+7*a)/b)-Ei(1,9*arccosh(c*x)+9*a/b)*exp((b*arccosh(c*x)+9*a)/b)-Ei(1,-9*arccosh(c*x)-9*a/b)*exp(-(-b*arccosh(c*x)+9*a)/b)-8*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)+6*Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)+6*Ei(1,-arccosh(c*x)-a/b)*exp(-(-b*arccosh(c*x)+a)/b)-8*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(-b*arccosh(c*x)+3*a)/b)+3*Ei(1,-7*arccosh(c*x)-7*a/b)*exp(-(-b*arccosh(c*x)+7*a)/b))/(c*x+1)/c^4/(c*x-1)/b`

### 3.284.5 Fracas [F]

$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{5/2}x^3}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((c^4*x^7 - 2*c^2*x^5 + x^3)*sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

---

3.284.  $\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$

**3.284.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \text{Timed out}$$

input `integrate(x**3*(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x)),x)`output `Timed out`**3.284.7 Maxima [F]**

$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{5/2}x^3}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate((-c^2*x^2 + 1)^(5/2)*x^3/(b*arccosh(c*x) + a), x)`**3.284.8 Giac [F]**

$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{5/2}x^3}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*x^2 + 1)^(5/2)*x^3/(b*arccosh(c*x) + a), x)`

**3.284.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x^3(1-c^2x^2)^{5/2}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((x^3*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x)),x)`output `int((x^3*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x)), x)`

**3.285**  $\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$

3.285.1 Optimal result . . . . . 2439  
 3.285.2 Mathematica [A] (warning: unable to verify) . . . . . 2440  
 3.285.3 Rubi [A] (verified) . . . . . 2440  
 3.285.4 Maple [A] (verified) . . . . . 2442  
 3.285.5 Fricas [F] . . . . . 2442  
 3.285.6 Sympy [F(-1)] . . . . . 2443  
 3.285.7 Maxima [F] . . . . . 2443  
 3.285.8 Giac [F] . . . . . 2443  
 3.285.9 Mupad [F(-1)] . . . . . 2444

**3.285.1 Optimal result**

Integrand size = 28, antiderivative size = 439

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{\sqrt{1-cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}}$$

$$+ \frac{\sqrt{1-cx} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}}$$

$$- \frac{\sqrt{1-cx} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}}$$

$$+ \frac{\sqrt{1-cx} \cosh\left(\frac{8a}{b}\right) \operatorname{Chi}\left(\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right)}{128bc^3\sqrt{-1+cx}}$$

$$- \frac{5\sqrt{1-cx} \log(a+b\operatorname{arccosh}(cx))}{128bc^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}}$$

$$- \frac{\sqrt{1-cx} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}}$$

$$+ \frac{\sqrt{1-cx} \sinh\left(\frac{6a}{b}\right) \operatorname{Shi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc^3\sqrt{-1+cx}}$$

$$- \frac{\sqrt{1-cx} \sinh\left(\frac{8a}{b}\right) \operatorname{Shi}\left(\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right)}{128bc^3\sqrt{-1+cx}}$$

---

3.285.  $\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$



output  $1/32*\text{Chi}(2*(a+b*\text{arccosh}(c*x))/b)*\cosh(2*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}+1/32*\text{Chi}(4*(a+b*\text{arccosh}(c*x))/b)*\cosh(4*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}-1/32*\text{Chi}(6*(a+b*\text{arccosh}(c*x))/b)*\cosh(6*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}+1/128*\text{Chi}(8*(a+b*\text{arccosh}(c*x))/b)*\cosh(8*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}-5/128*\ln(a+b*\text{arccosh}(c*x))*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}-1/32*\text{Shi}(2*(a+b*\text{arccosh}(c*x))/b)*\sinh(2*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}-1/32*\text{Shi}(4*(a+b*\text{arccosh}(c*x))/b)*\sinh(4*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}+1/32*\text{Shi}(6*(a+b*\text{arccosh}(c*x))/b)*\sinh(6*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}-1/128*\text{Shi}(8*(a+b*\text{arccosh}(c*x))/b)*\sinh(8*a/b)*(-c*x+1)^{(1/2)}/b/c^3/(c*x-1)^{(1/2)}$

### 3.285.2 Mathematica [A] (warning: unable to verify)

Time = 1.04 (sec) , antiderivative size = 233, normalized size of antiderivative = 0.53

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2}(4\cosh(\frac{2a}{b})\text{Chi}(2(\frac{a}{b}+\text{arccosh}(cx))))+4\cosh(\frac{4a}{b})\text{Chi}(4(\frac{a}{b}+\text{arccosh}(cx)))}{a+b\text{arccosh}(cx)}$$

input `Integrate[(x^2*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x]), x]`

output  $(\text{Sqrt}[1 - c^2*x^2]*(4*\text{Cosh}[(2*a)/b]*\text{CoshIntegral}[2*(a/b + \text{ArcCosh}[c*x])] + 4*\text{Cosh}[(4*a)/b]*\text{CoshIntegral}[4*(a/b + \text{ArcCosh}[c*x])] - 4*\text{Cosh}[(6*a)/b]*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])] + \text{Cosh}[(8*a)/b]*\text{CoshIntegral}[8*(a/b + \text{ArcCosh}[c*x])] - 5*\text{Log}[a + b*\text{ArcCosh}[c*x]] - 4*\text{Sinh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])] - 4*\text{Sinh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])] + 4*\text{Sinh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])] - \text{Sinh}[(8*a)/b]*\text{SinhIntegral}[8*(a/b + \text{ArcCosh}[c*x])])/(128*c^3*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(b + b*c*x))$

### 3.285.3 Rubi [A] (verified)

Time = 0.75 (sec) , antiderivative size = 240, normalized size of antiderivative = 0.55, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.285.  $\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx$

$$\begin{aligned}
 & \int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx \\
 & \quad \downarrow \text{6367} \\
 & \frac{\sqrt{1-cx} \int \frac{\cosh^2\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^6\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^3\sqrt{cx-1}} \\
 & \quad \downarrow \text{5971} \\
 & \frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{8a}{b}-\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right)}{128(a+b\operatorname{arccosh}(cx))} - \frac{\cosh\left(\frac{6a}{b}-\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} + \frac{\cosh\left(\frac{4a}{b}-\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} + \frac{\cosh\left(\frac{2a}{b}-\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} \right)}{bc^3\sqrt{cx-1}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{1-cx} \left( \frac{1}{32} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{32} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{32} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc^3\sqrt{cx-1}}
 \end{aligned}$$

input `Int[(x^2*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[1 - c*x]*((Cosh[(2*a)/b]*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b])/32 + (Cosh[(4*a)/b]*CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b])/32 - (Cosh[(6*a)/b]*CoshIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32 + (Cosh[(8*a)/b]*CoshIntegral[(8*(a + b*ArcCosh[c*x]))/b])/128 - (5*Log[a + b*ArcCosh[c*x]])/128 - (Sinh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/32 - (Sinh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/32 + (Sinh[(6*a)/b]*SinhIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32 - (Sinh[(8*a)/b]*SinhIntegral[(8*(a + b*ArcCosh[c*x]))/b])/128)/(b*c^3*Sqrt[-1 + c*x])`

### 3.285.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

$$3.285. \quad \int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$$

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.285.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 365, normalized size of antiderivative = 0.83

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-10\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))-10\ln(a+b\operatorname{arccosh}(cx))cx+4Ei_1(-6\operatorname{arccosh}(cx)+8a/b)\exp((b\operatorname{arccosh}(cx)+8a/b))-4Ei_1(4\operatorname{arccosh}(cx)+4a/b)\exp((b\operatorname{arccosh}(cx)+4a/b))-4Ei_1(2\operatorname{arccosh}(cx)+2a/b)\exp((b\operatorname{arccosh}(cx)+2a/b))-4Ei_1(-2\operatorname{arccosh}(cx)-2a/b)\exp(-(-b\operatorname{arccosh}(cx)+8a/b))-4Ei_1(-4\operatorname{arccosh}(cx)-4a/b)\exp(-(-b\operatorname{arccosh}(cx)+4a/b))\right)}{(c*x+1)/c^3/(c*x-1)/b}$

input `int(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `-1/256*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-10*(c*x-1)^(1/2)*(c*x+1)^(1/2)*ln(a+b*arccosh(c*x))-10*ln(a+b*arccosh(c*x))*c*x+4*Ei(1,-6*arccosh(c*x)-6*a/b)*exp(-(-b*arccosh(c*x)+8*a)/b)-Ei(1,8*arccosh(c*x)+8*a/b)*exp((b*arccosh(c*x)+8*a)/b)-Ei(1,-8*arccosh(c*x)-8*a/b)*exp(-(-b*arccosh(c*x)+8*a)/b)+4*Ei(1,6*arccosh(c*x)+6*a/b)*exp((b*arccosh(c*x)+6*a)/b)-4*Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)-4*Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)-4*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-4*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b))/(c*x+1)/c^3/(c*x-1)/b`

### 3.285.5 Fracas [F]

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{5/2}x^2}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((c^4*x^6 - 2*c^2*x^4 + x^2)*sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

---

3.285.  $\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$

**3.285.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^2(1 - c^2x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \text{Timed out}$$

input `integrate(x**2*(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x)),x)`output `Timed out`**3.285.7 Maxima [F]**

$$\int \frac{x^2(1 - c^2x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2 + 1)^{5/2}x^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate((-c^2*x^2 + 1)^(5/2)*x^2/(b*arccosh(c*x) + a), x)`**3.285.8 Giac [F]**

$$\int \frac{x^2(1 - c^2x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2 + 1)^{5/2}x^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*x^2 + 1)^(5/2)*x^2/(b*arccosh(c*x) + a), x)`

**3.285.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x^2(1-c^2x^2)^{5/2}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((x^2*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x)),x)`output `int((x^2*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x)), x)`

**3.286** 
$$\int \frac{x(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$$

3.286.1 Optimal result . . . . . 2445  
 3.286.2 Mathematica [A] (warning: unable to verify) . . . . . 2446  
 3.286.3 Rubi [A] (verified) . . . . . 2446  
 3.286.4 Maple [A] (verified) . . . . . 2448  
 3.286.5 Fracas [F] . . . . . 2448  
 3.286.6 Sympy [F(-1)] . . . . . 2449  
 3.286.7 Maxima [F] . . . . . 2449  
 3.286.8 Giac [F] . . . . . 2449  
 3.286.9 Mupad [F(-1)] . . . . . 2450

**3.286.1 Optimal result**

Integrand size = 26, antiderivative size = 397

$$\begin{aligned} \int \frac{x(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = & -\frac{5\sqrt{1-cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64bc^2\sqrt{-1+cx}} \\ & + \frac{9\sqrt{1-cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^2\sqrt{-1+cx}} \\ & - \frac{5\sqrt{1-cx} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^2\sqrt{-1+cx}} \\ & + \frac{\sqrt{1-cx} \cosh\left(\frac{7a}{b}\right) \operatorname{Chi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^2\sqrt{-1+cx}} \\ & + \frac{5\sqrt{1-cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64bc^2\sqrt{-1+cx}} \\ & - \frac{9\sqrt{1-cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^2\sqrt{-1+cx}} \\ & + \frac{5\sqrt{1-cx} \sinh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^2\sqrt{-1+cx}} \\ & - \frac{\sqrt{1-cx} \sinh\left(\frac{7a}{b}\right) \operatorname{Shi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64bc^2\sqrt{-1+cx}} \end{aligned}$$

output 
$$-5/64*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\cosh(a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}+9/64*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\cosh(3*a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}-5/64*\text{Chi}(5*(a+b*\text{arccosh}(c*x))/b)*\cosh(5*a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}+1/64*\text{Chi}(7*(a+b*\text{arccosh}(c*x))/b)*\cosh(7*a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}+5/64*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*\sinh(a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}-9/64*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*\sinh(3*a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}+5/64*\text{Shi}(5*(a+b*\text{arccosh}(c*x))/b)*\sinh(5*a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}-1/64*\text{Shi}(7*(a+b*\text{arccosh}(c*x))/b)*\sinh(7*a/b)*(-c*x+1)^{(1/2)}/b/c^2/(c*x-1)^{(1/2)}$$

### 3.286.2 Mathematica [A] (warning: unable to verify)

Time = 0.98 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.54

$$\int \frac{x(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2}(-5\cosh(\frac{a}{b})\text{Chi}(\frac{a}{b}+\text{arccosh}(cx))+9\cosh(\frac{3a}{b})\text{Chi}(3(\frac{a}{b}+\text{arccosh}(cx))))}{64*c^2*\sqrt{(-1+c*x)/(1+c*x)}*(b+b*c*x)}$$

input `Integrate[(x*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x]),x]`

output 
$$(\text{Sqrt}[1 - c^2*x^2]*(-5*\text{Cosh}[a/b]*\text{CoshIntegral}[a/b + \text{ArcCosh}[c*x]] + 9*\text{Cosh}[(3*a)/b]*\text{CoshIntegral}[3*(a/b + \text{ArcCosh}[c*x])] - 5*\text{Cosh}[(5*a)/b]*\text{CoshIntegral}[5*(a/b + \text{ArcCosh}[c*x])] + \text{Cosh}[(7*a)/b]*\text{CoshIntegral}[7*(a/b + \text{ArcCosh}[c*x])] + 5*\text{Sinh}[a/b]*\text{SinhIntegral}[a/b + \text{ArcCosh}[c*x]] - 9*\text{Sinh}[(3*a)/b]*\text{SinhIntegral}[3*(a/b + \text{ArcCosh}[c*x])] + 5*\text{Sinh}[(5*a)/b]*\text{SinhIntegral}[5*(a/b + \text{ArcCosh}[c*x])] - \text{Sinh}[(7*a)/b]*\text{SinhIntegral}[7*(a/b + \text{ArcCosh}[c*x])]))/(64*c^2*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(b + b*c*x))$$

### 3.286.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.56, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx$$

---

3.286.  $\int \frac{x(1-c^2x^2)^{5/2}}{a+b\text{arccosh}(cx)} dx$

$$\begin{array}{c}
 \downarrow 6367 \\
 \frac{\sqrt{1-cx} \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^6\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^2\sqrt{cx-1}} \\
 \downarrow 5971 \\
 \frac{\sqrt{1-cx} \int \left( \frac{\cosh\left(\frac{7a}{b} - \frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} - \frac{5 \cosh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} + \frac{9 \cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} - \frac{5 \cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} \right)}{bc^2\sqrt{cx-1}} \\
 \downarrow 2009 \\
 \frac{\sqrt{1-cx} \left( -\frac{5}{64} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) + \frac{9}{64} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{5}{64} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc^2\sqrt{cx-1}}
 \end{array}$$

input `Int[(x*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[1 - c*x]*((-5*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/64 + (9*Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/64 - (5*Cosh[(5*a)/b]*CoshIntegral[(5*(a + b*ArcCosh[c*x])/b])/64 + (Cosh[(7*a)/b]*CoshIntegral[(7*(a + b*ArcCosh[c*x])/b])/64 + (5*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/64 - (9*Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/64 + (5*Sinh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/64 - (Sinh[(7*a)/b]*SinhIntegral[(7*(a + b*ArcCosh[c*x])/b])/64))/(b*c^2*Sqrt[-1 + c*x])`

### 3.286.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

---

3.286.  $\int \frac{x(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$



rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.286.4 Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 318, normalized size of antiderivative = 0.80

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(\text{Ei}_1\left(7\operatorname{arccosh}(cx)+\frac{7a}{b}\right)e^{\frac{b\operatorname{arccosh}(cx)+7a}{b}}+\text{Ei}_1\left(-7\operatorname{arccosh}(cx)-\frac{7a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)+7a}{b}}\right)}{c^2(c*x+1)}$

input `int(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{128}(-c^2x^2+1)^{1/2}(-c*x-1)^{1/2}(c*x+1)^{1/2}c*x+c^2x^2-1*\left(\text{Ei}\left(1,7\operatorname{arccosh}(c*x)+\frac{7a}{b}\right)\exp\left(\frac{b\operatorname{arccosh}(c*x)+7a}{b}\right)+\text{Ei}\left(1,-7\operatorname{arccosh}(c*x)-\frac{7a}{b}\right)\exp\left(-\frac{b\operatorname{arccosh}(c*x)+7a}{b}\right)-5*\text{Ei}\left(1,5\operatorname{arccosh}(c*x)+\frac{5a}{b}\right)\exp\left(\frac{b\operatorname{arccosh}(c*x)+5a}{b}\right)+9*\text{Ei}\left(1,3\operatorname{arccosh}(c*x)+\frac{3a}{b}\right)\exp\left(\frac{b\operatorname{arccosh}(c*x)+3a}{b}\right)-5*\text{Ei}\left(1,\operatorname{arccosh}(c*x)+\frac{a}{b}\right)\exp\left(\frac{a+b\operatorname{arccosh}(c*x)}{b}\right)-5*\text{Ei}\left(1,-\operatorname{arccosh}(c*x)-\frac{a}{b}\right)\exp\left(-\frac{b\operatorname{arccosh}(c*x)+a}{b}\right)+9*\text{Ei}\left(1,-3\operatorname{arccosh}(c*x)-\frac{3a}{b}\right)\exp\left(-\frac{b\operatorname{arccosh}(c*x)+3a}{b}\right)-5*\text{Ei}\left(1,-5\operatorname{arccosh}(c*x)-\frac{5a}{b}\right)\exp\left(-\frac{b\operatorname{arccosh}(c*x)+5a}{b}\right)\right)/(c*x+1)/c^2/(c*x-1)/b$$

### 3.286.5 Fracas [F]

$$\int \frac{x(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{5/2}x}{b\operatorname{arccosh}(cx)+a} dx$$

input `integrate(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((c^4*x^5 - 2*c^2*x^3 + x)*sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

---

3.286. 
$$\int \frac{x(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$$

**3.286.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x(1 - c^2x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \text{Timed out}$$

input `integrate(x*(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x)),x)`output `Timed out`**3.286.7 Maxima [F]**

$$\int \frac{x(1 - c^2x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2 + 1)^{5/2} x}{b \operatorname{arccosh}(cx) + a} dx$$

input `integrate(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate((-c^2*x^2 + 1)^(5/2)*x/(b*arccosh(c*x) + a), x)`**3.286.8 Giac [F]**

$$\int \frac{x(1 - c^2x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2 + 1)^{5/2} x}{b \operatorname{arccosh}(cx) + a} dx$$

input `integrate(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*x^2 + 1)^(5/2)*x/(b*arccosh(c*x) + a), x)`

**3.286.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x(1-c^2x^2)^{5/2}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((x*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x)),x)`output `int((x*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x)), x)`

**3.287**  $\int \frac{(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$

3.287.1 Optimal result . . . . . 2451  
 3.287.2 Mathematica [A] (warning: unable to verify) . . . . . 2452  
 3.287.3 Rubi [A] (verified) . . . . . 2452  
 3.287.4 Maple [A] (verified) . . . . . 2454  
 3.287.5 Fricas [F] . . . . . 2455  
 3.287.6 Sympy [F(-1)] . . . . . 2455  
 3.287.7 Maxima [F] . . . . . 2455  
 3.287.8 Giac [F] . . . . . 2456  
 3.287.9 Mupad [F(-1)] . . . . . 2456

**3.287.1 Optimal result**

Integrand size = 25, antiderivative size = 339

$$\int \frac{(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx = \frac{15\sqrt{1-cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc\sqrt{-1+cx}} - \frac{3\sqrt{1-cx} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc\sqrt{-1+cx}} + \frac{\sqrt{1-cx} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc\sqrt{-1+cx}} - \frac{5\sqrt{1-cx} \log(a+b\operatorname{arccosh}(cx))}{16bc\sqrt{-1+cx}} - \frac{15\sqrt{1-cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc\sqrt{-1+cx}} + \frac{3\sqrt{1-cx} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \sinh\left(\frac{6a}{b}\right) \operatorname{Shi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32bc\sqrt{-1+cx}}$$

output  $15/32 \operatorname{Chi}(2(a+b \operatorname{arccosh}(cx))/b) \cosh(2a/b) (-cx+1)^{1/2} / b/c / (cx-1)^{(1/2)} - 3/16 \operatorname{Chi}(4(a+b \operatorname{arccosh}(cx))/b) \cosh(4a/b) (-cx+1)^{1/2} / b/c / (cx-1)^{(1/2)} + 1/32 \operatorname{Chi}(6(a+b \operatorname{arccosh}(cx))/b) \cosh(6a/b) (-cx+1)^{1/2} / b/c / (cx-1)^{(1/2)} - 5/16 \ln(a+b \operatorname{arccosh}(cx)) (-cx+1)^{1/2} / b/c / (cx-1)^{(1/2)} - 15/32 \operatorname{Shi}(2(a+b \operatorname{arccosh}(cx))/b) \sinh(2a/b) (-cx+1)^{1/2} / b/c / (cx-1)^{(1/2)} + 3/16 \operatorname{Shi}(4(a+b \operatorname{arccosh}(cx))/b) \sinh(4a/b) (-cx+1)^{1/2} / b/c / (cx-1)^{(1/2)} - 1/32 \operatorname{Shi}(6(a+b \operatorname{arccosh}(cx))/b) \sinh(6a/b) (-cx+1)^{1/2} / b/c / (cx-1)^{(1/2)}$

### 3.287.2 Mathematica [A] (warning: unable to verify)

Time = 0.76 (sec) , antiderivative size = 191, normalized size of antiderivative = 0.56

$$\int \frac{(1-c^2x^2)^{5/2}}{a+b \operatorname{arccosh}(cx)} dx = \frac{\sqrt{1-c^2x^2} (15 \cosh(\frac{2a}{b}) \operatorname{Chi}(2(\frac{a}{b} + \operatorname{arccosh}(cx))) - 6 \cosh(\frac{4a}{b}) \operatorname{Chi}(4(\frac{a}{b} + \operatorname{arccosh}(cx))))}{a+b \operatorname{arccosh}(cx)}$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(a + b*ArcCosh[c*x]),x]`

output  $(\operatorname{Sqrt}[1 - c^2x^2] * (15 * \operatorname{Cosh}[(2a)/b] * \operatorname{CoshIntegral}[2*(a/b + \operatorname{ArcCosh}[c*x])] - 6 * \operatorname{Cosh}[(4a)/b] * \operatorname{CoshIntegral}[4*(a/b + \operatorname{ArcCosh}[c*x])] + \operatorname{Cosh}[(6a)/b] * \operatorname{CoshIntegral}[6*(a/b + \operatorname{ArcCosh}[c*x])] - 10 * \operatorname{Log}[a + b * \operatorname{ArcCosh}[c*x]] - 15 * \operatorname{Sinh}[(2a)/b] * \operatorname{SinhIntegral}[2*(a/b + \operatorname{ArcCosh}[c*x])] + 6 * \operatorname{Sinh}[(4a)/b] * \operatorname{SinhIntegral}[4*(a/b + \operatorname{ArcCosh}[c*x])] - \operatorname{Sinh}[(6a)/b] * \operatorname{SinhIntegral}[6*(a/b + \operatorname{ArcCosh}[c*x])]) / (32 * b * c * \operatorname{Sqrt}[(-1 + cx)/(1 + cx)] * (1 + cx))$

### 3.287.3 Rubi [A] (verified)

Time = 0.57 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.56, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6321, 3042, 25, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-c^2x^2)^{5/2}}{a+b \operatorname{arccosh}(cx)} dx$$

↓ 6321

---

3.287.  $\int \frac{(1-c^2x^2)^{5/2}}{a+b \operatorname{arccosh}(cx)} dx$

$$\begin{aligned}
& \frac{\sqrt{1-cx} \int \frac{\sinh^6\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc\sqrt{cx-1}} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{1-cx} \int -\frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^6}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc\sqrt{cx-1}} \\
& \quad \downarrow \text{25} \\
& \frac{\sqrt{1-cx} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^6}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc\sqrt{cx-1}} \\
& \quad \downarrow \text{3793} \\
& \frac{\sqrt{1-cx} \int \left( -\frac{\cosh\left(\frac{6a}{b} - \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} + \frac{3 \cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{15 \cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} + \frac{1}{16(a+b\operatorname{arccosh}(cx))} \right)}{bc\sqrt{cx-1}} \\
& \quad \downarrow \text{2009} \\
& \frac{\sqrt{1-cx} \left( \frac{15}{32} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{16} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{32} \cosh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{bc\sqrt{cx-1}}
\end{aligned}$$

input `Int[(1 - c^2*x^2)^(5/2)/(a + b*ArcCosh[c*x]),x]`

output `(Sqrt[1 - c*x]*((15*Cosh[(2*a)/b]*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b])/32 - (3*Cosh[(4*a)/b]*CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b])/16 + (Cosh[(6*a)/b]*CoshIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32 - (5*Log[a + b*ArcCosh[c*x]])/16 - (15*Sinh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/32 + (3*Sinh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/16 - (Sinh[(6*a)/b]*SinhIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32))/(b*c*Sqrt[-1 + c*x])`

---

3.287.  $\int \frac{(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$

## 3.287.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

## 3.287.4 Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.88

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(20\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))+20\ln(a+b\operatorname{arccosh}(cx))cx+\operatorname{Ei}_1(6\operatorname{arccosh}(cx))\right)}{b}$

input `int((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{64}(-c^2x^2+1)^{1/2}(-cx-1)^{1/2}(cx+1)^{1/2}cx+c^2x^2-1\left(20(cx-1)^{1/2}(cx+1)^{1/2}\ln(a+b\operatorname{arccosh}(cx))+20\ln(a+b\operatorname{arccosh}(cx))cx+\operatorname{Ei}_1(6\operatorname{arccosh}(cx))+6\frac{a}{b}\exp\left(\frac{b\operatorname{arccosh}(cx)+6a}{b}\right)+\operatorname{Ei}_1(-6\operatorname{arccosh}(cx)-6\frac{a}{b})\exp\left(-\frac{b\operatorname{arccosh}(cx)+6a}{b}\right)-6\operatorname{Ei}_1(4\operatorname{arccosh}(cx)+4\frac{a}{b})\exp\left(\frac{b\operatorname{arccosh}(cx)+4a}{b}\right)+15\operatorname{Ei}_1(2\operatorname{arccosh}(cx)+2\frac{a}{b})\exp\left(\frac{b\operatorname{arccosh}(cx)+2a}{b}\right)+15\operatorname{Ei}_1(-2\operatorname{arccosh}(cx)-2\frac{a}{b})\exp\left(-\frac{b\operatorname{arccosh}(cx)+2a}{b}\right)-6\operatorname{Ei}_1(-4\operatorname{arccosh}(cx)-4\frac{a}{b})\exp\left(-\frac{b\operatorname{arccosh}(cx)+4a}{b}\right)\right)/(cx-1)/(cx+1)/c/b$$

---

3.287. 
$$\int \frac{(1-c^2x^2)^{5/2}}{a+b\operatorname{arccosh}(cx)} dx$$

**3.287.5 Fracas [F]**

$$\int \frac{(1 - c^2 x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a), x)`

**3.287.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x)),x)`

output `Timed out`

**3.287.7 Maxima [F]**

$$\int \frac{(1 - c^2 x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(5/2)/(b*arccosh(c*x) + a), x)`



**3.287.8 Giac [F]**

$$\int \frac{(1 - c^2 x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(5/2)/(b*arccosh(c*x) + a), x)`

**3.287.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{a + b \operatorname{acosh}(cx)} dx$$

input `int((1 - c^2*x^2)^(5/2)/(a + b*acosh(c*x)),x)`

output `int((1 - c^2*x^2)^(5/2)/(a + b*acosh(c*x)), x)`

**3.288** 
$$\int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))} dx$$

3.288.1 Optimal result	2457
3.288.2 Mathematica [N/A]	2458
3.288.3 Rubi [N/A]	2458
3.288.4 Maple [N/A] (verified)	2459
3.288.5 Fricas [N/A]	2460
3.288.6 Sympy [F(-1)]	2460
3.288.7 Maxima [N/A]	2460
3.288.8 Giac [F(-2)]	2461
3.288.9 Mupad [N/A]	2461

**3.288.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\begin{aligned} \int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))} dx = & -\frac{11\sqrt{-1+cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b\sqrt{1-cx}} \\ & + \frac{7\sqrt{-1+cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}} \\ & - \frac{\sqrt{-1+cx} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}} \\ & + \frac{11\sqrt{-1+cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b\sqrt{1-cx}} \\ & - \frac{7\sqrt{-1+cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}} \\ & + \frac{\sqrt{-1+cx} \sinh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}} \\ & + \operatorname{Int}\left(\frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}, x\right) \end{aligned}$$

---

3.288. 
$$\int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))} dx$$

output 
$$\begin{aligned} & -11/8*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\cosh(a/b)*(c*x-1)^{(1/2)}/b/(-c*x+1)^{(1/2)}+7 \\ & /16*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\cosh(3*a/b)*(c*x-1)^{(1/2)}/b/(-c*x+1)^{(1/2)} \\ & -1/16*\text{Chi}(5*(a+b*\text{arccosh}(c*x))/b)*\cosh(5*a/b)*(c*x-1)^{(1/2)}/b/(-c*x+1)^{(1/2)} \\ & +11/8*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*\sinh(a/b)*(c*x-1)^{(1/2)}/b/(-c*x+1)^{(1/2)} \\ & -7/16*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*\sinh(3*a/b)*(c*x-1)^{(1/2)}/b/(-c*x+1)^{(1/2)} \\ & +1/16*\text{Shi}(5*(a+b*\text{arccosh}(c*x))/b)*\sinh(5*a/b)*(c*x-1)^{(1/2)}/b/(-c*x+1)^{(1/2)} \\ & +\text{Unintegrable}(1/x/(a+b*\text{arccosh}(c*x))/(-c^2*x^2+1)^{(1/2)},x) \end{aligned}$$

### 3.288.2 Mathematica [N/A]

Not integrable

Time = 1.54 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1-c^2x^2)^{5/2}}{x(a+\text{barccosh}(cx))} dx = \int \frac{(1-c^2x^2)^{5/2}}{x(a+\text{barccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x*(a + b*ArcCosh[c*x])), x]`

output `Integrate[(1 - c^2*x^2)^(5/2)/(x*(a + b*ArcCosh[c*x])), x]`

### 3.288.3 Rubi [N/A]

Not integrable

Time = 1.38 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-c^2x^2)^{5/2}}{x(a+\text{barccosh}(cx))} dx$$

↓ 6369

$$\int \left( -\frac{3c^2x}{\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))} + \frac{1}{x\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))} - \frac{c^6x^5}{\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))} + \frac{1}{\sqrt{1-c^2x^2}} \right) dx$$

---

3.288.  $\int \frac{(1-c^2x^2)^{5/2}}{x(a+\text{barccosh}(cx))} dx$

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{arccosh}(cx))} dx - \frac{11\sqrt{cx-1} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{arccosh}(cx)}{b}\right)}{8b\sqrt{1-cx}} +$$

$$\frac{7\sqrt{cx-1} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}} - \frac{\sqrt{cx-1} \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}} +$$

$$\frac{11\sqrt{cx-1} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{arccosh}(cx)}{b}\right)}{8b\sqrt{1-cx}} - \frac{7\sqrt{cx-1} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}} +$$

$$\frac{\sqrt{cx-1} \sinh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+\operatorname{arccosh}(cx))}{b}\right)}{16b\sqrt{1-cx}}$$

input `Int[(1 - c^2*x^2)^(5/2)/(x*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

### 3.288.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.288.4 Maple [N/A] (verified)

Not integrable

Time = 1.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{5/2}}{x(a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x)),x)`

---

3.288.  $\int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))} dx$

**3.288.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.61

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)x} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b*x*arccosh(c*x) + a*x), x)`

**3.288.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + \operatorname{barccosh}(cx))} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x/(a+b*acosh(c*x)),x)`

output `Timed out`

**3.288.7 Maxima [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)x} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)*x), x)`

---

3.288.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + \operatorname{barccosh}(cx))} dx$

**3.288.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.288.9 Mupad [N/A]**

Not integrable

Time = 2.72 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(5/2)/(x*(a + b*acosh(c*x))),x)`

output `int((1 - c^2*x^2)^(5/2)/(x*(a + b*acosh(c*x))), x)`

**3.289**  $\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$

3.289.1 Optimal result . . . . . 2462  
 3.289.2 Mathematica [N/A] . . . . . 2463  
 3.289.3 Rubi [N/A] . . . . . 2463  
 3.289.4 Maple [N/A] (verified) . . . . . 2464  
 3.289.5 Fricas [N/A] . . . . . 2464  
 3.289.6 Sympy [F(-1)] . . . . . 2465  
 3.289.7 Maxima [N/A] . . . . . 2465  
 3.289.8 Giac [N/A] . . . . . 2465  
 3.289.9 Mupad [N/A] . . . . . 2466

**3.289.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))} dx = \frac{c\sqrt{-1+cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{b\sqrt{1-cx}} - \frac{c\sqrt{-1+cx} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8b\sqrt{1-cx}} - \frac{15c\sqrt{-1+cx} \log(a+b\operatorname{arccosh}(cx))}{8b\sqrt{1-cx}} - \frac{c\sqrt{-1+cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{b\sqrt{1-cx}} + \frac{c\sqrt{-1+cx} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8b\sqrt{1-cx}} + \operatorname{Int}\left(\frac{1}{x^2\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output

```
c*Chi(2*(a+b*arccosh(c*x))/b)*cosh(2*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)-1/8*c*Chi(4*(a+b*arccosh(c*x))/b)*cosh(4*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)-15/8*c*ln(a+b*arccosh(c*x))*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)-c*Shi(2*(a+b*arccosh(c*x))/b)*sinh(2*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+1/8*c*Shi(4*(a+b*arccosh(c*x))/b)*sinh(4*a/b)*(c*x-1)^(1/2)/b/(-c*x+1)^(1/2)+Unintegrabl e(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)
```

3.289.  $\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))} dx$

**3.289.2 Mathematica [N/A]**

Not integrable

Time = 2.59 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + \operatorname{barccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + \operatorname{barccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x^2*(a + b*ArcCosh[c*x])), x]`output `Integrate[(1 - c^2*x^2)^(5/2)/(x^2*(a + b*ArcCosh[c*x])), x]`**3.289.3 Rubi [N/A]**

Not integrable

Time = 1.20 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + \operatorname{barccosh}(cx))} dx$$

↓ 6369

$$\int \left( -\frac{3c^2}{\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} + \frac{1}{x^2 \sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} - \frac{c^6 x^4}{\sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} + \frac{1}{\sqrt{1 - c^2 x^2}} \right) dx$$

↓ 2009

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2}(a + \operatorname{barccosh}(cx))} dx + \frac{c\sqrt{cx - 1} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{b\sqrt{1 - cx}} - \frac{c\sqrt{cx - 1} \cosh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{8b\sqrt{1 - cx}} - \frac{c\sqrt{cx - 1} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{b\sqrt{1 - cx}} + \frac{c\sqrt{cx - 1} \sinh\left(\frac{4a}{b}\right) \operatorname{Shi}\left(\frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{8b\sqrt{1 - cx}} - \frac{15c\sqrt{cx - 1} \log(a + \operatorname{barccosh}(cx))}{8b\sqrt{1 - cx}}$$

---

3.289.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + \operatorname{barccosh}(cx))} dx$



input `Int[(1 - c^2*x^2)^(5/2)/(x^2*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

### 3.289.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.289.4 Maple [N/A] (verified)

Not integrable

Time = 0.98 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{5/2}}{x^2(a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x)),x)`

### 3.289.5 Fracas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{(1 - c^2x^2)^{5/2}}{x^2(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2x^2 + 1)^{5/2}}{(b \operatorname{arccosh}(cx) + a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x)),x, algorithm="fricas")`

---

3.289.  $\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b \operatorname{arccosh}(cx))} dx$

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b*x^2*arccosh(c*x) + a*x^2), x)`

### 3.289.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + b \operatorname{arccosh}(cx))} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x**2/(a+b*acosh(c*x)), x)`

output `Timed out`

### 3.289.7 Maxima [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arccosh}(cx) + a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x)), x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)*x^2), x)`

### 3.289.8 Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arccosh}(cx) + a)x^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x)), x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)*x^2), x)`

---

3.289.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x^2(a + b \operatorname{arccosh}(cx))} dx$

**3.289.9 Mupad [N/A]**

Not integrable

Time = 2.76 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(5/2)/(x^2*(a + b*acosh(c*x))),x)`output `int((1 - c^2*x^2)^(5/2)/(x^2*(a + b*acosh(c*x))), x)`

**3.290**  $\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$

3.290.1 Optimal result . . . . . 2467  
 3.290.2 Mathematica [N/A] . . . . . 2467  
 3.290.3 Rubi [N/A] . . . . . 2468  
 3.290.4 Maple [N/A] (verified) . . . . . 2468  
 3.290.5 Fricas [N/A] . . . . . 2469  
 3.290.6 Sympy [F(-1)] . . . . . 2469  
 3.290.7 Maxima [N/A] . . . . . 2469  
 3.290.8 Giac [F(-2)] . . . . . 2470  
 3.290.9 Mupad [N/A] . . . . . 2470

**3.290.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1 - c^2x^2)^{5/2}}{x^3(a + b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{(1 - c^2x^2)^{5/2}}{x^3(a + b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x)), x)`

**3.290.2 Mathematica [N/A]**

Not integrable

Time = 7.87 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1 - c^2x^2)^{5/2}}{x^3(a + b\operatorname{arccosh}(cx))} dx = \int \frac{(1 - c^2x^2)^{5/2}}{x^3(a + b\operatorname{arccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x^3*(a + b*ArcCosh[c*x])), x]`

output `Integrate[(1 - c^2*x^2)^(5/2)/(x^3*(a + b*ArcCosh[c*x])), x]`

---

3.290.  $\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$

**3.290.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{arccosh}(cx))} dx$$

input `Int[(1 - c^2*x^2)^(5/2)/(x^3*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.290.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.290.4 Maple [N/A] (verified)**

Not integrable

Time = 1.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 x^2 + 1)^{5/2}}{x^3 (a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x)),x)`

---

3.290.  $\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+b\operatorname{arccosh}(cx))} dx$

**3.290.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)x^3} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b*x^3*arccosh(c*x) + a*x^3), x)`

**3.290.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x**3/(a+b*acosh(c*x)),x)`

output `Timed out`

**3.290.7 Maxima [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)x^3} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)*x^3), x)`

---

3.290.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x^3(a + b \operatorname{arccosh}(cx))} dx$

**3.290.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.290.9 Mupad [N/A]**

Not integrable

Time = 2.71 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3(a + \operatorname{barccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(5/2)/(x^3*(a + b*acosh(c*x))),x)`

output `int((1 - c^2*x^2)^(5/2)/(x^3*(a + b*acosh(c*x))), x)`

**3.291** 
$$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

3.291.1 Optimal result . . . . . 2471  
 3.291.2 Mathematica [N/A] . . . . . 2471  
 3.291.3 Rubi [N/A] . . . . . 2472  
 3.291.4 Maple [N/A] (verified) . . . . . 2472  
 3.291.5 Fricas [N/A] . . . . . 2473  
 3.291.6 Sympy [F(-1)] . . . . . 2473  
 3.291.7 Maxima [N/A] . . . . . 2473  
 3.291.8 Giac [N/A] . . . . . 2474  
 3.291.9 Mupad [N/A] . . . . . 2474

**3.291.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x)), x)`

**3.291.2 Mathematica [N/A]**

Not integrable

Time = 1.53 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx = \int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x^4*(a + b*ArcCosh[c*x])), x]`

output `Integrate[(1 - c^2*x^2)^(5/2)/(x^4*(a + b*ArcCosh[c*x])), x]`

---

3.291. 
$$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$$



**3.291.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{barccosh}(cx))} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{barccosh}(cx))} dx$$

input `Int[(1 - c^2*x^2)^(5/2)/(x^4*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.291.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.291.4 Maple [N/A] (verified)**

Not integrable

Time = 1.30 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 x^2 + 1)^{5/2}}{x^4 (a + b \operatorname{arccosh}(cx))} dx$$

input `int((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x)),x)`

output `int((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x)),x)`

---

3.291.  $\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))} dx$

**3.291.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)x^4} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b*x^4*arccosh(c*x) + a*x^4), x)`

**3.291.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4(a + \operatorname{barccosh}(cx))} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x**4/(a+b*acosh(c*x)),x)`

output `Timed out`

**3.291.7 Maxima [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4(a + \operatorname{barccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)x^4} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)*x^4), x)`

---

3.291.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x^4(a + b \operatorname{arccosh}(cx))} dx$

**3.291.8 Giac [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a) x^4} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)*x^4), x)`**3.291.9 Mupad [N/A]**

Not integrable

Time = 2.71 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + b \operatorname{arccosh}(cx))} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + b \operatorname{acosh}(cx))} dx$$

input `int((1 - c^2*x^2)^(5/2)/(x^4*(a + b*acosh(c*x))),x)`output `int((1 - c^2*x^2)^(5/2)/(x^4*(a + b*acosh(c*x))), x)`

$$3.292 \quad \int \frac{x^4}{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)} dx$$

3.292.1 Optimal result . . . . .	2475
3.292.2 Mathematica [A] (verified) . . . . .	2475
3.292.3 Rubi [A] (verified) . . . . .	2476
3.292.4 Maple [A] (verified) . . . . .	2477
3.292.5 Fricas [F] . . . . .	2478
3.292.6 Sympy [F] . . . . .	2478
3.292.7 Maxima [F] . . . . .	2478
3.292.8 Giac [F] . . . . .	2479
3.292.9 Mupad [F(-1)] . . . . .	2479

### 3.292.1 Optimal result

Integrand size = 24, antiderivative size = 98

$$\int \frac{x^4}{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)} dx = \frac{\sqrt{-1+ax} \operatorname{Chi}(2\operatorname{arccosh}(ax))}{2a^5\sqrt{1-ax}} + \frac{\sqrt{-1+ax} \operatorname{Chi}(4\operatorname{arccosh}(ax))}{8a^5\sqrt{1-ax}} + \frac{3\sqrt{-1+ax} \log(\operatorname{arccosh}(ax))}{8a^5\sqrt{1-ax}}$$

output  $\frac{1}{2} \operatorname{Chi}(2 \operatorname{arccosh}(a x)) (a x - 1)^{1/2} / a^5 (-a x + 1)^{1/2} + \frac{1}{8} \operatorname{Chi}(4 \operatorname{arccosh}(a x)) (a x - 1)^{1/2} / a^5 (-a x + 1)^{1/2} + \frac{3}{8} \ln(\operatorname{arccosh}(a x)) (a x - 1)^{1/2} / a^5 (-a x + 1)^{1/2}$

### 3.292.2 Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.70

$$\int \frac{x^4}{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)} dx = \frac{\sqrt{\frac{-1+ax}{1+ax}} (1+ax) (4 \operatorname{Chi}(2 \operatorname{arccosh}(ax)) + \operatorname{Chi}(4 \operatorname{arccosh}(ax)) + 3 \log(\operatorname{arccosh}(ax)))}{8a^5 \sqrt{-((-1+ax)(1+ax))}}$$

input `Integrate[x^4/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(4*CoshIntegral[2*ArcCosh[a*x]] + CoshIntegral[4*ArcCosh[a*x]] + 3*Log[ArcCosh[a*x]])/(8*a^5*Sqrt[-((-1 + a*x)*(1 + a*x))])`

### 3.292.3 Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.56, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4}{\sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx \\
 & \quad \downarrow \text{6367} \\
 & \frac{\sqrt{ax - 1} \int \frac{a^4 x^4}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^5 \sqrt{1 - ax}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{ax - 1} \int \frac{\sin\left(i\operatorname{arccosh}(ax) + \frac{\pi}{2}\right)^4}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^5 \sqrt{1 - ax}} \\
 & \quad \downarrow \text{3793} \\
 & \frac{\sqrt{ax - 1} \int \left( \frac{\cosh(2\operatorname{arccosh}(ax))}{2\operatorname{arccosh}(ax)} + \frac{\cosh(4\operatorname{arccosh}(ax))}{8\operatorname{arccosh}(ax)} + \frac{3}{8\operatorname{arccosh}(ax)} \right) d\operatorname{arccosh}(ax)}{a^5 \sqrt{1 - ax}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{ax - 1} \left( \frac{1}{2} \operatorname{Chi}(2\operatorname{arccosh}(ax)) + \frac{1}{8} \operatorname{Chi}(4\operatorname{arccosh}(ax)) + \frac{3}{8} \log(\operatorname{arccosh}(ax)) \right)}{a^5 \sqrt{1 - ax}}
 \end{aligned}$$

input `Int[x^4/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[-1 + a*x]*(CoshIntegral[2*ArcCosh[a*x]]/2 + CoshIntegral[4*ArcCosh[a*x]]/8 + (3*Log[ArcCosh[a*x]]/8))/(a^5*Sqrt[1 - a*x])`

---

3.292.  $\int \frac{x^4}{\sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx$

3.292.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3793 Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

```
rule 6367 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]
```

3.292.4 Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.93

method	result
default	$-\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}(-Ei_1(4 \operatorname{arccosh}(ax))-Ei_1(-4 \operatorname{arccosh}(ax))+6 \ln(\operatorname{arccosh}(ax))-4 Ei_1(2 \operatorname{arccosh}(ax))-4 Ei_1(-2 \operatorname{arccosh}(ax)))}{16a^5(a^2x^2-1)}$

```
input int(x^4/arccosh(a*x)/(-a^2*x^2+1)^(1/2), x, method=_RETURNVERBOSE)
```

```
output -1/16*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a^5/(a^2*x^2-1)*(-Ei(1,4*arccosh(a*x))-Ei(1,-4*arccosh(a*x))+6*ln(arccosh(a*x))-4*Ei(1,2*arccosh(a*x))-4*Ei(1,-2*arccosh(a*x)))
```

3.292.  $\int \frac{x^4}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$

**3.292.5 Fracas [F]**

$$\int \frac{x^4}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^4}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^4/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^4/((a^2*x^2 - 1)*arccosh(a*x)), x)`

**3.292.6 Sympy [F]**

$$\int \frac{x^4}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^4}{\sqrt{-(ax-1)(ax+1)}\operatorname{acosh}(ax)} dx$$

input `integrate(x**4/acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**4/(sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)), x)`

**3.292.7 Maxima [F]**

$$\int \frac{x^4}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^4}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^4/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(x^4/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.292.8 Giac [F]**

$$\int \frac{x^4}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^4}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^4/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^4/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.292.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^4}{\operatorname{acosh}(ax)\sqrt{1-a^2x^2}} dx$$

input `int(x^4/(acosh(a*x)*(1 - a^2*x^2)^(1/2)),x)`

output `int(x^4/(acosh(a*x)*(1 - a^2*x^2)^(1/2)), x)`



### 3.293 $\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$

3.293.1 Optimal result	2480
3.293.2 Mathematica [A] (verified)	2480
3.293.3 Rubi [A] (verified)	2481
3.293.4 Maple [A] (verified)	2482
3.293.5 Fricas [F]	2483
3.293.6 Sympy [F]	2483
3.293.7 Maxima [F]	2483
3.293.8 Giac [F(-2)]	2484
3.293.9 Mupad [F(-1)]	2484

#### 3.293.1 Optimal result

Integrand size = 24, antiderivative size = 65

$$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \frac{3\sqrt{-1+ax}\operatorname{Chi}(\operatorname{arccosh}(ax))}{4a^4\sqrt{1-ax}} + \frac{\sqrt{-1+ax}\operatorname{Chi}(3\operatorname{arccosh}(ax))}{4a^4\sqrt{1-ax}}$$

output `3/4*Chi(arccosh(a*x))*(a*x-1)^(1/2)/a^4/(-a*x+1)^(1/2)+1/4*Chi(3*arccosh(a*x))*(a*x-1)^(1/2)/a^4/(-a*x+1)^(1/2)`

#### 3.293.2 Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.92

$$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax)(3\operatorname{Chi}(\operatorname{arccosh}(ax)) + \operatorname{Chi}(3\operatorname{arccosh}(ax)))}{4a^4\sqrt{-((-1+ax)(1+ax))}}$$

input `Integrate[x^3/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*(3*CoshIntegral[ArcCosh[a*x]] + CoshIntegral[3*ArcCosh[a*x]]))/(4*a^4*Sqrt[-((-1 + a*x)*(1 + a*x))])`

**3.293.3 Rubi [A] (verified)**

Time = 0.40 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx \\
 & \quad \downarrow \text{6367} \\
 & \frac{\sqrt{ax-1} \int \frac{a^3x^3}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^4\sqrt{1-ax}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{ax-1} \int \frac{\sin\left(i\operatorname{arccosh}(ax)+\frac{\pi}{2}\right)^3}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^4\sqrt{1-ax}} \\
 & \quad \downarrow \text{3793} \\
 & \frac{\sqrt{ax-1} \int \left(\frac{3ax}{4\operatorname{arccosh}(ax)} + \frac{\cosh(3\operatorname{arccosh}(ax))}{4\operatorname{arccosh}(ax)}\right) d\operatorname{arccosh}(ax)}{a^4\sqrt{1-ax}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{ax-1} \left(\frac{3}{4}\operatorname{Chi}(\operatorname{arccosh}(ax)) + \frac{1}{4}\operatorname{Chi}(3\operatorname{arccosh}(ax))\right)}{a^4\sqrt{1-ax}}
 \end{aligned}$$

input `Int[x^3/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[-1 + a*x]*((3*CoshIntegral[ArcCosh[a*x]])/4 + CoshIntegral[3*ArcCosh[a*x]]/4))/(a^4*Sqrt[1 - a*x])`

## 3.293.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

## 3.293.4 Maple [A] (verified)

Time = 1.03 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.20

method	result	size
default	$\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}(Ei_1(3\operatorname{arccosh}(ax))+Ei_1(-3\operatorname{arccosh}(ax))+3Ei_1(\operatorname{arccosh}(ax))+3Ei_1(-\operatorname{arccosh}(ax)))}{8a^4(a^2x^2-1)}$	78

input `int(x^3/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{8}(-a^2x^2+1)^{1/2}(ax-1)^{1/2}(ax+1)^{1/2}(Ei_1(3\operatorname{arccosh}(ax))+Ei_1(-3\operatorname{arccosh}(ax))+3Ei_1(\operatorname{arccosh}(ax))+3Ei_1(-\operatorname{arccosh}(ax)))/a^4/(a^2x^2-1)$$

**3.293.5 Fricas [F]**

$$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^3}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^3/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^3/((a^2*x^2 - 1)*arccosh(a*x)), x)`

**3.293.6 Sympy [F]**

$$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^3}{\sqrt{-(ax-1)(ax+1)}\operatorname{acosh}(ax)} dx$$

input `integrate(x**3/acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**3/(sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)), x)`

**3.293.7 Maxima [F]**

$$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^3}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^3/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(x^3/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.293.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.293.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^3}{\operatorname{acosh}(ax)\sqrt{1-a^2x^2}} dx$$

input `int(x^3/(acosh(a*x)*(1 - a^2*x^2)^(1/2)),x)`

output `int(x^3/(acosh(a*x)*(1 - a^2*x^2)^(1/2)), x)`

$$3.294 \quad \int \frac{x^2}{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)} dx$$

3.294.1 Optimal result	2485
3.294.2 Mathematica [A] (verified)	2485
3.294.3 Rubi [A] (verified)	2486
3.294.4 Maple [A] (verified)	2487
3.294.5 Fracas [F]	2488
3.294.6 Sympy [F]	2488
3.294.7 Maxima [F]	2488
3.294.8 Giac [F]	2489
3.294.9 Mupad [F(-1)]	2489

### 3.294.1 Optimal result

Integrand size = 24, antiderivative size = 65

$$\int \frac{x^2}{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)} dx = \frac{\sqrt{-1+ax} \operatorname{Chi}(2 \operatorname{arccosh}(ax))}{2a^3 \sqrt{1-ax}} + \frac{\sqrt{-1+ax} \log(\operatorname{arccosh}(ax))}{2a^3 \sqrt{1-ax}}$$

output `1/2*Chi(2*arccosh(a*x))*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)+1/2*ln(arccosh(a*x))*(a*x-1)^(1/2)/a^3/(-a*x+1)^(1/2)`

### 3.294.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.92

$$\int \frac{x^2}{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)} dx = -\frac{\sqrt{-((-1+ax)(1+ax))}(\operatorname{Chi}(2 \operatorname{arccosh}(ax)) + \log(\operatorname{arccosh}(ax)))}{2a^3 \sqrt{\frac{-1+ax}{1+ax}}(1+ax)}$$

input `Integrate[x^2/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `-1/2*(Sqrt[-((-1 + a*x)*(1 + a*x))]*(CoshIntegral[2*ArcCosh[a*x]] + Log[ArcCosh[a*x]]))/(a^3*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x))`

---


$$3.294. \quad \int \frac{x^2}{\sqrt{1-a^2x^2} \operatorname{arccosh}(ax)} dx$$

**3.294.3 Rubi [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx \\
 & \quad \downarrow \text{6367} \\
 & \frac{\sqrt{ax-1} \int \frac{a^2x^2}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^3\sqrt{1-ax}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{ax-1} \int \frac{\sin\left(i\operatorname{arccosh}(ax)+\frac{\pi}{2}\right)^2}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^3\sqrt{1-ax}} \\
 & \quad \downarrow \text{3793} \\
 & \frac{\sqrt{ax-1} \int \left(\frac{\cosh(2\operatorname{arccosh}(ax))}{2\operatorname{arccosh}(ax)} + \frac{1}{2\operatorname{arccosh}(ax)}\right) d\operatorname{arccosh}(ax)}{a^3\sqrt{1-ax}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{ax-1}\left(\frac{1}{2}\operatorname{Chi}(2\operatorname{arccosh}(ax)) + \frac{1}{2}\log(\operatorname{arccosh}(ax))\right)}{a^3\sqrt{1-ax}}
 \end{aligned}$$

input `Int[x^2/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[-1 + a*x]*(CoshIntegral[2*ArcCosh[a*x]]/2 + Log[ArcCosh[a*x]]/2))/(a^3*Sqrt[1 - a*x])`

## 3.294.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

## 3.294.4 Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.03

method	result	size
default	$\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}(Ei_1(2\operatorname{arccosh}(ax))+Ei_1(-2\operatorname{arccosh}(ax))-2\ln(\operatorname{arccosh}(ax)))}{4a^3(a^2x^2-1)}$	67

input `int(x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/4*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*(Ei(1,2*arccosh(a*x))+Ei(1,-2*arccosh(a*x))-2*ln(arccosh(a*x)))/a^3/(a^2*x^2-1)`



**3.294.5 Fracas [F]**

$$\int \frac{x^2}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^2}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x^2/((a^2*x^2 - 1)*arccosh(a*x)), x)`

**3.294.6 Sympy [F]**

$$\int \frac{x^2}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^2}{\sqrt{-(ax-1)(ax+1)}\operatorname{acosh}(ax)} dx$$

input `integrate(x**2/acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x**2/(sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)), x)`

**3.294.7 Maxima [F]**

$$\int \frac{x^2}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^2}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(x^2/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.294.8 Giac [F]**

$$\int \frac{x^2}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^2}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^2/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.294.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x^2}{\operatorname{acosh}(ax)\sqrt{1-a^2x^2}} dx$$

input `int(x^2/(acosh(a*x)*(1 - a^2*x^2)^(1/2)),x)`

output `int(x^2/(acosh(a*x)*(1 - a^2*x^2)^(1/2)), x)`

### 3.295 $\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$

3.295.1 Optimal result . . . . .	2490
3.295.2 Mathematica [A] (verified) . . . . .	2490
3.295.3 Rubi [A] (verified) . . . . .	2491
3.295.4 Maple [B] (verified) . . . . .	2492
3.295.5 Fricas [F] . . . . .	2492
3.295.6 Sympy [F] . . . . .	2493
3.295.7 Maxima [F] . . . . .	2493
3.295.8 Giac [F] . . . . .	2493
3.295.9 Mupad [F(-1)] . . . . .	2494

#### 3.295.1 Optimal result

Integrand size = 22, antiderivative size = 28

$$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \frac{\sqrt{-1+ax}\operatorname{Chi}(\operatorname{arccosh}(ax))}{a^2\sqrt{1-ax}}$$

output `Chi(arccosh(a*x))*(a*x-1)^(1/2)/a^2/(-a*x+1)^(1/2)`

#### 3.295.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.79

$$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = -\frac{\sqrt{-((-1+ax)(1+ax))}\operatorname{Chi}(\operatorname{arccosh}(ax))}{a^2\sqrt{\frac{-1+ax}{1+ax}}(1+ax)}$$

input `Integrate[x/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `-((Sqrt[-((-1 + a*x)*(1 + a*x))]*CoshIntegral[ArcCosh[a*x]])/(a^2*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)))`

**3.295.3 Rubi [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$ , Rules used = {6367, 3042, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx \\
 & \quad \downarrow \text{6367} \\
 & \frac{\sqrt{ax-1} \int \frac{ax}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^2\sqrt{1-ax}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{ax-1} \int \frac{\sin(i\operatorname{arccosh}(ax)+\frac{\pi}{2})}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a^2\sqrt{1-ax}} \\
 & \quad \downarrow \text{3782} \\
 & \frac{\sqrt{ax-1}\operatorname{Chi}(\operatorname{arccosh}(ax))}{a^2\sqrt{1-ax}}
 \end{aligned}$$

input `Int[x/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[-1 + a*x]*CoshIntegral[ArcCosh[a*x]])/(a^2*Sqrt[1 - a*x])`

**3.295.3.1 Defintions of rubi rules used**

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.295.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 57 vs.  $2(24) = 48$ .

Time = 0.78 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.07

method	result	size
default	$\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}(Ei_1(\operatorname{arccosh}(ax))+Ei_1(-\operatorname{arccosh}(ax)))}{2a^2(a^2x^2-1)}$	58

input `int(x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)*(Ei(1,arccosh(a*x))+Ei(1,-arccosh(a*x)))/a^2/(a^2*x^2-1)`

### 3.295.5 Fricas [F]

$$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)} dx$$

input `integrate(x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-a^2*x^2 + 1)*x/((a^2*x^2 - 1)*arccosh(a*x)), x)`

**3.295.6 Sympy [F]**

$$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x}{\sqrt{-(ax-1)(ax+1)}\operatorname{acosh}(ax)} dx$$

input `integrate(x/acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(x/(sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)), x)`

**3.295.7 Maxima [F]**

$$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x}{\sqrt{-a^2x^2+1}\operatorname{arcosh}(ax)} dx$$

input `integrate(x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(x/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.295.8 Giac [F]**

$$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x}{\sqrt{-a^2x^2+1}\operatorname{arcosh}(ax)} dx$$

input `integrate(x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.295.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{x}{\operatorname{acosh}(ax)\sqrt{1-a^2x^2}} dx$$

input `int(x/(acosh(a*x)*(1 - a^2*x^2)^(1/2)),x)`output `int(x/(acosh(a*x)*(1 - a^2*x^2)^(1/2)), x)`

**3.296**  $\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$

3.296.1 Optimal result . . . . . 2495  
 3.296.2 Mathematica [A] (verified) . . . . . 2495  
 3.296.3 Rubi [A] (verified) . . . . . 2496  
 3.296.4 Maple [A] (verified) . . . . . 2496  
 3.296.5 Fricas [B] (verification not implemented) . . . . . 2497  
 3.296.6 Sympy [F] . . . . . 2497  
 3.296.7 Maxima [F] . . . . . 2497  
 3.296.8 Giac [F] . . . . . 2498  
 3.296.9 Mupad [F(-1)] . . . . . 2498

**3.296.1 Optimal result**

Integrand size = 21, antiderivative size = 28

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \frac{\sqrt{-1+ax} \log(\operatorname{arccosh}(ax))}{a\sqrt{1-ax}}$$

output `ln(arccosh(a*x))*(a*x-1)^(1/2)/a/(-a*x+1)^(1/2)`

**3.296.2 Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.68

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \frac{\sqrt{\frac{-1+ax}{1+ax}}(1+ax) \log(\operatorname{arccosh}(ax))}{a\sqrt{-((-1+ax)(1+ax))}}$$

input `Integrate[1/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Log[ArcCosh[a*x]])/(a*Sqrt[-((-1 + a*x)*(1 + a*x))])`



### 3.296.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$ , Rules used = {6305}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$$

↓ 6305

$$\frac{\sqrt{ax-1}\log(\operatorname{arccosh}(ax))}{a\sqrt{1-ax}}$$

input `Int[1/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `(Sqrt[-1 + a*x]*Log[ArcCosh[a*x]])/(a*Sqrt[1 - a*x])`

#### 3.296.3.1 Defintions of rubi rules used

rule 6305 `Int[1/(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*Sqrt[(d_) + (e_.)*(x_)^2]), x_Symbol] :> Simp[(1/(b*c))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*Log[a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0]`

### 3.296.4 Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.71

method	result	size
default	$-\frac{\sqrt{-a^2x^2+1}\sqrt{ax-1}\sqrt{ax+1}\ln(\operatorname{arccosh}(ax))}{a(a^2x^2-1)}$	48

input `int(1/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `-(-a^2*x^2+1)^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(a^2*x^2-1)*ln(arccosh(a*x))`

**3.296.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 55 vs.  $2(24) = 48$ .

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.96

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = -\frac{\sqrt{a^2x^2-1}\sqrt{-a^2x^2+1}\log(\log(ax+\sqrt{a^2x^2-1}))}{a^3x^2-a}$$

input `integrate(1/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `-sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)*log(log(a*x + sqrt(a^2*x^2 - 1)))/(a^3*x^2 - a)`

**3.296.6 Sympy [F]**

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-(ax-1)(ax+1)}\operatorname{acosh}(ax)} dx$$

input `integrate(1/acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`

output `Integral(1/(sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)), x)`

**3.296.7 Maxima [F]**

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2x^2+1}\operatorname{arcosh}(ax)} dx$$

input `integrate(1/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.296.8 Giac [F]**

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2x^2+1}\operatorname{arcosh}(ax)} dx$$

input `integrate(1/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)), x)`

**3.296.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{\operatorname{acosh}(ax)\sqrt{1-a^2x^2}} dx$$

input `int(1/(acosh(a*x)*(1 - a^2*x^2)^(1/2)),x)`

output `int(1/(acosh(a*x)*(1 - a^2*x^2)^(1/2)), x)`

**3.297**      $\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$

3.297.1 Optimal result	2499
3.297.2 Mathematica [N/A]	2499
3.297.3 Rubi [N/A]	2500
3.297.4 Maple [N/A] (verified)	2500
3.297.5 Fricas [N/A]	2501
3.297.6 Sympy [N/A]	2501
3.297.7 Maxima [N/A]	2501
3.297.8 Giac [N/A]	2502
3.297.9 Mupad [N/A]	2502

**3.297.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \operatorname{Int}\left(\frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)}, x\right)$$

output `Unintegrable(1/x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

**3.297.2 Mathematica [N/A]**

Not integrable

Time = 0.69 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$$

input `Integrate[1/(x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `Integrate[1/(x*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]), x]`

**3.297.3 Rubi [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$$

↓ 6375

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx$$

input `Int[1/(x*sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `$Aborted`

**3.297.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.297.4 Maple [N/A] (verified)**

Not integrable

Time = 1.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{x \operatorname{arccosh}(ax) \sqrt{-a^2x^2 + 1}} dx$$

input `int(1/x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

output `int(1/x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

**3.297.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.46

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2x^2+1}x \operatorname{arccosh}(ax)} dx$$

```
input integrate(1/x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(-sqrt(-a^2*x^2 + 1)/((a^2*x^3 - x)*arccosh(a*x)), x)
```

**3.297.6 Sympy [N/A]**

Not integrable

Time = 1.37 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{x\sqrt{-(ax-1)(ax+1)}\operatorname{acosh}(ax)} dx$$

```
input integrate(1/x/acosh(a*x)/(-a**2*x**2+1)**(1/2),x)
```

```
output Integral(1/(x*sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)), x)
```

**3.297.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2x^2+1}x \operatorname{arccosh}(ax)} dx$$

```
input integrate(1/x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")
```

```
output integrate(1/(sqrt(-a^2*x^2 + 1)*x*arccosh(a*x)), x)
```

**3.297.8 Giac [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2x^2+1}x \operatorname{arccosh}(ax)} dx$$

input `integrate(1/x/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-a^2*x^2 + 1)*x*arccosh(a*x)), x)`

**3.297.9 Mupad [N/A]**

Not integrable

Time = 2.74 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)} dx = \int \frac{1}{x \operatorname{acosh}(ax) \sqrt{1-a^2x^2}} dx$$

input `int(1/(x*acosh(a*x)*(1 - a^2*x^2)^(1/2)),x)`

output `int(1/(x*acosh(a*x)*(1 - a^2*x^2)^(1/2)), x)`

$$3.298 \quad \int \frac{1}{x^2 \sqrt{1-a^2 x^2} \operatorname{arccosh}(ax)} dx$$

3.298.1 Optimal result	2503
3.298.2 Mathematica [N/A]	2503
3.298.3 Rubi [N/A]	2504
3.298.4 Maple [N/A] (verified)	2504
3.298.5 Fricas [N/A]	2505
3.298.6 Sympy [N/A]	2505
3.298.7 Maxima [N/A]	2505
3.298.8 Giac [N/A]	2506
3.298.9 Mupad [N/A]	2506

### 3.298.1 Optimal result

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{x^2 \sqrt{1-a^2 x^2} \operatorname{arccosh}(ax)} dx = \operatorname{Int}\left(\frac{1}{x^2 \sqrt{1-a^2 x^2} \operatorname{arccosh}(ax)}, x\right)$$

output `Unintegrable(1/x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

### 3.298.2 Mathematica [N/A]

Not integrable

Time = 0.89 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^2 \sqrt{1-a^2 x^2} \operatorname{arccosh}(ax)} dx = \int \frac{1}{x^2 \sqrt{1-a^2 x^2} \operatorname{arccosh}(ax)} dx$$

input `Integrate[1/(x^2*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `Integrate[1/(x^2*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]), x]`



**3.298.3 Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx$$

↓ 6375

$$\int \frac{1}{x^2 \sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx$$

input `Int[1/(x^2*sqrt[1 - a^2*x^2]*ArcCosh[a*x]),x]`

output `$Aborted`

**3.298.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.298.4 Maple [N/A] (verified)**

Not integrable

Time = 2.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{x^2 \operatorname{arccosh}(ax) \sqrt{-a^2 x^2 + 1}} dx$$

input `int(1/x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

output `int(1/x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x)`

**3.298.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.54

$$\int \frac{1}{x^2 \sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2 x^2 + 1} x^2 \operatorname{arccosh}(ax)} dx$$

input `integrate(1/x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`output `integral(-sqrt(-a^2*x^2 + 1)/((a^2*x^4 - x^2)*arccosh(a*x)), x)`**3.298.6 Sympy [N/A]**

Not integrable

Time = 4.13 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^2 \sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx = \int \frac{1}{x^2 \sqrt{-(ax - 1)(ax + 1)} \operatorname{acosh}(ax)} dx$$

input `integrate(1/x**2/acosh(a*x)/(-a**2*x**2+1)**(1/2),x)`output `Integral(1/(x**2*sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)), x)`**3.298.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2 x^2 + 1} x^2 \operatorname{arccosh}(ax)} dx$$

input `integrate(1/x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`output `integrate(1/(sqrt(-a^2*x^2 + 1)*x^2*arccosh(a*x)), x)`

**3.298.8 Giac [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx = \int \frac{1}{\sqrt{-a^2 x^2 + 1} x^2 \operatorname{arccosh}(ax)} dx$$

input `integrate(1/x^2/arccosh(a*x)/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`output `integrate(1/(sqrt(-a^2*x^2 + 1)*x^2*arccosh(a*x)), x)`**3.298.9 Mupad [N/A]**

Not integrable

Time = 2.72 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - a^2 x^2} \operatorname{arccosh}(ax)} dx = \int \frac{1}{x^2 \operatorname{acosh}(ax) \sqrt{1 - a^2 x^2}} dx$$

input `int(1/(x^2*acosh(a*x)*(1 - a^2*x^2)^(1/2)),x)`output `int(1/(x^2*acosh(a*x)*(1 - a^2*x^2)^(1/2)), x)`

**3.299**  $\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

3.299.1 Optimal result . . . . . 2507  
 3.299.2 Mathematica [A] (verified) . . . . . 2508  
 3.299.3 Rubi [A] (verified) . . . . . 2508  
 3.299.4 Maple [A] (verified) . . . . . 2510  
 3.299.5 Fricas [F] . . . . . 2510  
 3.299.6 Sympy [F] . . . . . 2510  
 3.299.7 Maxima [F] . . . . . 2511  
 3.299.8 Giac [F(-2)] . . . . . 2511  
 3.299.9 Mupad [F(-1)] . . . . . 2511

**3.299.1 Optimal result**

Integrand size = 28, antiderivative size = 197

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \frac{3\sqrt{-1+cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4bc^4\sqrt{1-cx}} + \frac{\sqrt{-1+cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4bc^4\sqrt{1-cx}} - \frac{3\sqrt{-1+cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4bc^4\sqrt{1-cx}} - \frac{\sqrt{-1+cx} \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4bc^4\sqrt{1-cx}}$$

output

```
3/4*Chi((a+b*arccosh(c*x))/b)*cosh(a/b)*(c*x-1)^(1/2)/b/c^4/(-c*x+1)^(1/2)
+1/4*Chi(3*(a+b*arccosh(c*x))/b)*cosh(3*a/b)*(c*x-1)^(1/2)/b/c^4/(-c*x+1)^(1/2)
-3/4*Shi((a+b*arccosh(c*x))/b)*sinh(a/b)*(c*x-1)^(1/2)/b/c^4/(-c*x+1)^(1/2)
-1/4*Shi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)*(c*x-1)^(1/2)/b/c^4/(-c*x+1)^(1/2)
```

**3.299.2 Mathematica [A] (verified)**

Time = 0.47 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.66

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$$

$$= \frac{\sqrt{\frac{-1+cx}{1+cx}}(1+cx) \left( 3 \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) + \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) - 3 \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \right)}{4bc^4\sqrt{-((-1+cx)(1+cx))}}$$

input `Integrate[x^3/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`output `(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(3*Cosh[a/b]*CoshIntegral[a/b + ArcCosh[c*x]] + Cosh[(3*a)/b]*CoshIntegral[3*(a/b + ArcCosh[c*x])] - 3*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] - Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])])/(4*b*c^4*Sqrt[-((-1 + c*x)*(1 + c*x))])`**3.299.3 Rubi [A] (verified)**Time = 0.55 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.62, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$$

$$\downarrow 6367$$

$$\frac{\sqrt{cx-1} \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^4\sqrt{1-cx}}$$

$$\downarrow 3042$$

$$\frac{\sqrt{cx-1} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)^3}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{bc^4\sqrt{1-cx}}$$

$$\downarrow 3793$$

---

3.299.  $\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

$$\frac{\sqrt{cx-1} \int \left( \frac{\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} + \frac{3 \cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{bc^4\sqrt{1-cx}}$$

↓ 2009

$$\frac{\sqrt{cx-1} \left( \frac{3}{4} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) + \frac{1}{4} \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{4} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{bc^4\sqrt{1-cx}}$$

input `Int[x^3/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `(Sqrt[-1 + c*x]*((3*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/4 + (Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/4 - (3*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/4 - (Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/4))/(b*c^4*Sqrt[1 - c*x])`

### 3.299.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

**3.299.4 Maple [A] (verified)**

Time = 1.24 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.92

method	result
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(\text{Ei}_1\left(3\operatorname{arccosh}(cx)+\frac{3a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)+3a}{b}}+\text{Ei}_1\left(-3\operatorname{arccosh}(cx)-\frac{3a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)+3a}{b}}\right)}{8b(c^2x^2-1)c^4}$

```
input int(x^3/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/8*(-c^2*x^2+1)^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(Ei(1,3
*arccosh(c*x)+3*a/b)*exp((-b*arccosh(c*x)+3*a)/b)+Ei(1,-3*arccosh(c*x)-3*a
/b)*exp(-b*arccosh(c*x)+3*a)/b)+3*Ei(1,arccosh(c*x)+a/b)*exp((-b*arccosh(
c*x)+a)/b)+3*Ei(1,-arccosh(c*x)-a/b)*exp(-a+b*arccosh(c*x)/b))/b/(c^2*x^
2-1)/c^4
```

**3.299.5 Fracas [F]**

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x^3}{\sqrt{-c^2x^2+1}(b\operatorname{arccosh}(cx)+a)} dx$$

```
input integrate(x^3/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(-sqrt(-c^2*x^2 + 1)*x^3/(a*c^2*x^2 + (b*c^2*x^2 - b)*arccosh(c*x)
- a), x)
```

**3.299.6 Sympy [F]**

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x^3}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))} dx$$

```
input integrate(x**3/(a+b*acosh(c*x))/(-c**2*x**2+1)**(1/2),x)
```

```
output Integral(x**3/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))), x)
```

---

3.299.  $\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

**3.299.7 Maxima [F]**

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^3}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x^3/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(x^3/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.299.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index\_m & i,const vecteur & l) Error: Bad Argument Value

**3.299.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^3}{(a+b\operatorname{acosh}(cx))\sqrt{1-c^2x^2}} dx$$

input `int(x^3/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)),x)`

output `int(x^3/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)), x)`



### 3.300 $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

3.300.1 Optimal result . . . . .	2512
3.300.2 Mathematica [A] (verified) . . . . .	2512
3.300.3 Rubi [A] (verified) . . . . .	2513
3.300.4 Maple [A] (verified) . . . . .	2514
3.300.5 Fricas [F] . . . . .	2515
3.300.6 Sympy [F] . . . . .	2515
3.300.7 Maxima [F] . . . . .	2515
3.300.8 Giac [F] . . . . .	2516
3.300.9 Mupad [F(-1)] . . . . .	2516

#### 3.300.1 Optimal result

Integrand size = 28, antiderivative size = 139

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \frac{\sqrt{-1+cx} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2bc^3\sqrt{1-cx}} + \frac{\sqrt{-1+cx} \log(a+b\operatorname{arccosh}(cx))}{2bc^3\sqrt{1-cx}} - \frac{\sqrt{-1+cx} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2bc^3\sqrt{1-cx}}$$

output `1/2*Chi(2*(a+b*arccosh(c*x))/b)*cosh(2*a/b)*(c*x-1)^(1/2)/b/c^3/(-c*x+1)^(1/2)+1/2*ln(a+b*arccosh(c*x))*(c*x-1)^(1/2)/b/c^3/(-c*x+1)^(1/2)-1/2*Shi(2*(a+b*arccosh(c*x))/b)*sinh(2*a/b)*(c*x-1)^(1/2)/b/c^3/(-c*x+1)^(1/2)`

#### 3.300.2 Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.71

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \frac{\sqrt{1-c^2x^2} \left( \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) + \log(a+b\operatorname{arccosh}(cx)) - \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \right)}{2c^3\sqrt{\frac{-1+cx}{1+cx}}(b+bcx)}$$

input `Integrate[x^2/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `-1/2*(Sqrt[1 - c^2*x^2]*(Cosh[(2*a)/b]*CoshIntegral[2*(a/b + ArcCosh[c*x])  
] + Log[a + b*ArcCosh[c*x]] - Sinh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[  
c*x]])))/(c^3*Sqrt[(-1 + c*x)/(1 + c*x)]*(b + b*c*x))`

### 3.300.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.65, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx \\
 & \quad \downarrow \text{6367} \\
 & \frac{\sqrt{cx-1} \int \frac{\cosh^2\left(\frac{a}{b}-\frac{a+b\operatorname{barccosh}(cx)}{b}\right)}{a+b\operatorname{barccosh}(cx)} d(a+b\operatorname{barccosh}(cx))}{bc^3\sqrt{1-cx}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{cx-1} \int \frac{\sin\left(\frac{ia}{b}-\frac{i(a+b\operatorname{barccosh}(cx))}{b}+\frac{\pi}{2}\right)^2}{a+b\operatorname{barccosh}(cx)} d(a+b\operatorname{barccosh}(cx))}{bc^3\sqrt{1-cx}} \\
 & \quad \downarrow \text{3793} \\
 & \frac{\sqrt{cx-1} \int \left( \frac{\cosh\left(\frac{2a}{b}-\frac{2(a+b\operatorname{barccosh}(cx))}{b}\right)}{2(a+b\operatorname{barccosh}(cx))} + \frac{1}{2(a+b\operatorname{barccosh}(cx))} \right) d(a+b\operatorname{barccosh}(cx))}{bc^3\sqrt{1-cx}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{cx-1} \left( \frac{1}{2} \cosh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{barccosh}(cx))}{b}\right) - \frac{1}{2} \sinh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{barccosh}(cx))}{b}\right) + \frac{1}{2} \log(a+b\operatorname{barccosh}(cx)) \right)}{bc^3\sqrt{1-cx}}
 \end{aligned}$$

input `Int[x^2/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

---

3.300.  $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx$

output  $(\text{Sqrt}[-1 + c*x]*((\text{Cosh}[(2*a)/b]*\text{CoshIntegral}[(2*(a + b*\text{ArcCosh}[c*x]))/b])/2 + \text{Log}[a + b*\text{ArcCosh}[c*x]]/2 - (\text{Sinh}[(2*a)/b]*\text{SinhIntegral}[(2*(a + b*\text{ArcCosh}[c*x]))/b])/2))/b)/b)/(b*c^3*\text{Sqrt}[1 - c*x])$

### 3.300.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.300.4 Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.16

method	result
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(2\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))-2\ln(a+b\operatorname{arccosh}(cx))cx+\operatorname{Ei}_1\left(2\operatorname{arccosh}(cx)+\frac{2a}{b}\right)\right)}{4b(c^2x^2-1)c^3}$

input `int(x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output  $1/4*(-c^2*x^2+1)^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*\ln(a+b*\operatorname{arccosh}(c*x))-2*\ln(a+b*\operatorname{arccosh}(c*x))*c*x+\operatorname{Ei}(1,2*\operatorname{arccosh}(c*x)+2*a/b)*\exp((-b*\operatorname{arccosh}(c*x)+2*a)/b)+\operatorname{Ei}(1,-2*\operatorname{arccosh}(c*x)-2*a/b)*\exp(-b*\operatorname{arccosh}(c*x)+2*a/b))/b/(c^2*x^2-1)/c^3$

3.300.  $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

**3.300.5 Fricas [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^2}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^2/(a*c^2*x^2 + (b*c^2*x^2 - b)*arccosh(c*x) - a), x)`

**3.300.6 Sympy [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^2}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))} dx$$

input `integrate(x**2/(a+b*acosh(c*x))/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x**2/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))), x)`

**3.300.7 Maxima [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^2}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(x^2/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.300.8 Giac [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^2}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^2/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.300.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^2}{(a+b\operatorname{acosh}(cx))\sqrt{1-c^2x^2}} dx$$

input `int(x^2/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)),x)`

output `int(x^2/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)), x)`

### 3.301 $\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

3.301.1 Optimal result . . . . .	2517
3.301.2 Mathematica [A] (verified) . . . . .	2517
3.301.3 Rubi [A] (verified) . . . . .	2518
3.301.4 Maple [A] (verified) . . . . .	2520
3.301.5 Fricas [F] . . . . .	2521
3.301.6 Sympy [F] . . . . .	2521
3.301.7 Maxima [F] . . . . .	2521
3.301.8 Giac [F] . . . . .	2522
3.301.9 Mupad [F(-1)] . . . . .	2522

#### 3.301.1 Optimal result

Integrand size = 26, antiderivative size = 92

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \frac{\sqrt{-1+cx} \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{bc^2\sqrt{1-cx}} - \frac{\sqrt{-1+cx} \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{bc^2\sqrt{1-cx}}$$

```
output Chi((a+b*arccosh(c*x))/b)*cosh(a/b)*(c*x-1)^(1/2)/b/c^2/(-c*x+1)^(1/2)-Shi
((a+b*arccosh(c*x))/b)*sinh(a/b)*(c*x-1)^(1/2)/b/c^2/(-c*x+1)^(1/2)
```

#### 3.301.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.88

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \frac{\sqrt{1-c^2x^2}(-\cosh\left(\frac{a}{b}\right)\operatorname{Chi}\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right)+\sinh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right))}{c^2\sqrt{\frac{-1+cx}{1+cx}}(b+bcx)}$$

```
input Integrate[x/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]
```

output (Sqrt[1 - c^2\*x^2]\*(-(Cosh[a/b]\*CoshIntegral[a/b + ArcCosh[c\*x]]) + Sinh[a/b]\*SinhIntegral[a/b + ArcCosh[c\*x]]))/(c^2\*Sqrt[(-1 + c\*x)/(1 + c\*x)]\*(b + b\*c\*x))

### 3.301.3 Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.74, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {6367, 3042, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{1 - c^2 x^2} (a + \operatorname{barccosh}(cx))} dx$$

↓ 6367

$$\frac{\sqrt{cx - 1} \int \frac{\cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc^2 \sqrt{1 - cx}}$$

↓ 3042

$$\frac{\sqrt{cx - 1} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc^2 \sqrt{1 - cx}}$$

↓ 3784

$$\frac{\sqrt{cx - 1} \left( \cosh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx)) - i \sinh\left(\frac{a}{b}\right) \int \frac{i \sinh\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx)) \right)}{bc^2 \sqrt{1 - cx}}$$

↓ 26

$$\frac{\sqrt{cx - 1} \left( \cosh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx)) - \sinh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx)) \right)}{bc^2 \sqrt{1 - cx}}$$

↓ 3042

---

3.301.  $\int \frac{x}{\sqrt{1 - c^2 x^2} (a + \operatorname{barccosh}(cx))} dx$

$$\frac{\sqrt{cx-1} \left( \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - \sinh\left(\frac{a}{b}\right) \int \frac{i \sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{bc^2\sqrt{1-cx}}$$

↓ 26

$$\frac{\sqrt{cx-1} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) + \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{bc^2\sqrt{1-cx}}$$

↓ 3779

$$\frac{\sqrt{cx-1} \left( -\sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) + \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{bc^2\sqrt{1-cx}}$$

↓ 3782

$$\frac{\sqrt{cx-1} \left( \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{bc^2\sqrt{1-cx}}$$

input `Int[x/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `(Sqrt[-1 + c*x]*(Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b] - Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b]))/(b*c^2*Sqrt[1 - c*x])`

### 3.301.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`



rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.301.4 Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.26

method	result	S
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(\text{Ei}_1\left(\text{arccosh}(cx)+\frac{a}{b}\right)e^{-\frac{b\text{arccosh}(cx)+a}{b}}+\text{Ei}_1\left(-\text{arccosh}(cx)-\frac{a}{b}\right)e^{-\frac{a+b\text{arccosh}(cx)}{b}}\right)}{2b(c^2x^2-1)c^2}$	1

input `int(x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*(-c^2*x^2+1)^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(Ei(1,a rccosh(c*x)+a/b)*exp((-b*arccosh(c*x)+a)/b)+Ei(1,-arccosh(c*x)-a/b)*exp(-(a+b*arccosh(c*x))/b))/b/(c^2*x^2-1)/c^2`

---

3.301.  $\int \frac{x}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))} dx$

**3.301.5 Fricas [F]**

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x/(a*c^2*x^2 + (b*c^2*x^2 - b)*arccosh(c*x) - a), x)`

**3.301.6 Sympy [F]**

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))} dx$$

input `integrate(x/(a+b*acosh(c*x))/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))), x)`

**3.301.7 Maxima [F]**

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(x/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.301.8 Giac [F]**

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.301.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x}{(a+b\operatorname{acosh}(cx))\sqrt{1-c^2x^2}} dx$$

input `int(x/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)),x)`

output `int(x/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)), x)`

**3.302**  $\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

3.302.1 Optimal result	2523
3.302.2 Mathematica [A] (verified)	2523
3.302.3 Rubi [A] (verified)	2524
3.302.4 Maple [A] (verified)	2524
3.302.5 Fricas [B] (verification not implemented)	2525
3.302.6 Sympy [F]	2525
3.302.7 Maxima [F]	2525
3.302.8 Giac [F]	2526
3.302.9 Mupad [F(-1)]	2526

**3.302.1 Optimal result**

Integrand size = 25, antiderivative size = 35

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \frac{\sqrt{-1+cx} \log(a+b\operatorname{arccosh}(cx))}{bc\sqrt{1-cx}}$$

output `ln(a+b*arccosh(c*x))*(c*x-1)^(1/2)/b/c/(-c*x+1)^(1/2)`

**3.302.2 Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \frac{\sqrt{\frac{-1+cx}{1+cx}}(1+cx) \log(a+b\operatorname{arccosh}(cx))}{bc\sqrt{-((-1+cx)(1+cx))}}$$

input `Integrate[1/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Log[a + b*ArcCosh[c*x]])/(b*c*Sqrt[-((-1 + c*x)*(1 + c*x))])`

### 3.302.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$ , Rules used = {6305}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$$

↓ 6305

$$\frac{\sqrt{cx-1} \log(a+b\operatorname{arccosh}(cx))}{bc\sqrt{1-cx}}$$

input `Int[1/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `(Sqrt[-1 + c*x]*Log[a + b*ArcCosh[c*x]])/(b*c*Sqrt[1 - c*x])`

#### 3.302.3.1 Defintions of rubi rules used

rule 6305 `Int[1/(((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*Sqrt[(d_) + (e_.)*(x_)^2]), x_Symbol] :> Simp[(1/(b*c))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*Log[a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0]`

### 3.302.4 Maple [A] (verified)

Time = 0.69 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

method	result	size
default	$-\frac{\sqrt{-c^2x^2+1}\sqrt{cx-1}\sqrt{cx+1}\ln(a+b\operatorname{arccosh}(cx))}{c(c^2x^2-1)b}$	55

input `int(1/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `-(-c^2*x^2+1)^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/(c^2*x^2-1)*ln(a+b*arccosh(c*x))/b`

---

3.302.  $\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

**3.302.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 65 vs.  $2(31) = 62$ .

Time = 0.25 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.86

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = -\frac{\sqrt{c^2x^2-1}\sqrt{-c^2x^2+1}\log\left(\frac{b\log(cx+\sqrt{c^2x^2-1})+a}{b}\right)}{bc^3x^2-bc}$$

input `integrate(1/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `-sqrt(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)*log((b*log(c*x + sqrt(c^2*x^2 - 1)) + a)/b)/(b*c^3*x^2 - b*c)`

**3.302.6 Sympy [F]**

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))} dx$$

input `integrate(1/(a+b*acosh(c*x))/(-c**2*x**2+1)**(1/2),x)`

output `Integral(1/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))), x)`

**3.302.7 Maxima [F]**

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(1/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.302.8 Giac [F]**

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(1/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.302.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{(a+b\operatorname{acosh}(cx))\sqrt{1-c^2x^2}} dx$$

input `int(1/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)),x)`

output `int(1/((a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)), x)`

### 3.303 $\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$

3.303.1 Optimal result	2527
3.303.2 Mathematica [N/A]	2527
3.303.3 Rubi [N/A]	2528
3.303.4 Maple [N/A] (verified)	2528
3.303.5 Fricas [N/A]	2529
3.303.6 Sympy [N/A]	2529
3.303.7 Maxima [N/A]	2529
3.303.8 Giac [F(-2)]	2530
3.303.9 Mupad [N/A]	2530

#### 3.303.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2), x)`

#### 3.303.2 Mathematica [N/A]

Not integrable

Time = 1.66 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[1/(x*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])), x]`

output `Integrate[1/(x*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])), x]`



**3.303.3 Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{arccosh}(cx))} dx$$

input `Int[1/(x*sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.303.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.303.4 Maple [N/A] (verified)**

Not integrable

Time = 1.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x(a+b\operatorname{arccosh}(cx))\sqrt{-c^2x^2+1}} dx$$

input `int(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

output `int(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

**3.303.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)x} dx$$

input `integrate(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)/(a*c^2*x^3 - a*x + (b*c^2*x^3 - b*x)*arccosh(c*x)), x)`

**3.303.6 Sympy [N/A]**

Not integrable

Time = 2.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{x\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))} dx$$

input `integrate(1/x/(a+b*acosh(c*x))/(-c**2*x**2+1)**(1/2),x)`

output `Integral(1/(x*sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))), x)`

**3.303.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)x} dx$$

input `integrate(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)*x), x)`

**3.303.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate(1/x/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.303.9 Mupad [N/A]**

Not integrable

Time = 2.75 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{x(a+b\operatorname{acosh}(cx))\sqrt{1-c^2x^2}} dx$$

input `int(1/(x*(a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)),x)`

output `int(1/(x*(a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)), x)`

$$\mathbf{3.304} \quad \int \frac{1}{x^2 \sqrt{1-c^2 x^2} (a+b \operatorname{arccosh}(cx))} dx$$

3.304.1 Optimal result	2531
3.304.2 Mathematica [N/A]	2531
3.304.3 Rubi [N/A]	2532
3.304.4 Maple [N/A] (verified)	2532
3.304.5 Fricas [N/A]	2533
3.304.6 Sympy [N/A]	2533
3.304.7 Maxima [N/A]	2533
3.304.8 Giac [N/A]	2534
3.304.9 Mupad [N/A]	2534

### 3.304.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x^2 \sqrt{1-c^2 x^2} (a+b \operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{1}{x^2 \sqrt{1-c^2 x^2} (a+b \operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2), x)`

### 3.304.2 Mathematica [N/A]

Not integrable

Time = 2.37 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x^2 \sqrt{1-c^2 x^2} (a+b \operatorname{arccosh}(cx))} dx = \int \frac{1}{x^2 \sqrt{1-c^2 x^2} (a+b \operatorname{arccosh}(cx))} dx$$

input `Integrate[1/(x^2*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])), x]`

output `Integrate[1/(x^2*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])), x]`

**3.304.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + \operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + \operatorname{arccosh}(cx))} dx$$

input `Int[1/(x^2*sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.304.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.304.4 Maple [N/A] (verified)**

Not integrable

Time = 1.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^2 (a + b \operatorname{arccosh}(cx)) \sqrt{-c^2 x^2 + 1}} dx$$

input `int(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

output `int(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x)`

**3.304.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.89

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{-c^2 x^2 + 1} (b \operatorname{arccosh}(cx) + a) x^2} dx$$

input `integrate(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)/(a*c^2*x^4 - a*x^2 + (b*c^2*x^4 - b*x^2)*arccosh(c*x)), x)`

**3.304.6 Sympy [N/A]**

Not integrable

Time = 3.68 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{x^2 \sqrt{-(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))} dx$$

input `integrate(1/x**2/(a+b*acosh(c*x))/(-c**2*x**2+1)**(1/2),x)`

output `Integral(1/(x**2*sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))), x)`

**3.304.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{-c^2 x^2 + 1} (b \operatorname{arccosh}(cx) + a) x^2} dx$$

input `integrate(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)*x^2), x)`

---

3.304.  $\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))} dx$

**3.304.8 Giac [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{-c^2 x^2 + 1} (b \operatorname{arcosh}(cx) + a) x^2} dx$$

input `integrate(1/x^2/(a+b*arccosh(c*x))/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`output `integrate(1/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)*x^2), x)`**3.304.9 Mupad [N/A]**

Not integrable

Time = 2.86 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{x^2 (a + b \operatorname{acosh}(cx)) \sqrt{1 - c^2 x^2}} dx$$

input `int(1/(x^2*(a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)),x)`output `int(1/(x^2*(a + b*acosh(c*x))*(1 - c^2*x^2)^(1/2)), x)`

$$3.305 \quad \int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

3.305.1 Optimal result	2535
3.305.2 Mathematica [N/A]	2535
3.305.3 Rubi [N/A]	2536
3.305.4 Maple [N/A] (verified)	2536
3.305.5 Fricas [N/A]	2537
3.305.6 Sympy [N/A]	2537
3.305.7 Maxima [N/A]	2537
3.305.8 Giac [N/A]	2538
3.305.9 Mupad [N/A]	2538

### 3.305.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)), x)`

### 3.305.2 Mathematica [N/A]

Not integrable

Time = 4.84 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[x^2/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

output `Integrate[x^2/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

---


$$3.305. \quad \int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$



**3.305.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))} dx$$

input `Int[x^2/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.305.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.305.4 Maple [N/A] (verified)**

Not integrable

Time = 0.52 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{(-c^2 x^2 + 1)^{3/2} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

---

3.305.  $\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$

**3.305.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.25

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^2/(a*c^4*x^4 - 2*a*c^2*x^2 + (b*c^4*x^4 - 2*b*c^2*x^2 + b)*arccosh(c*x) + a), x)`

**3.305.6 Sympy [N/A]**

Not integrable

Time = 8.70 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^2}{(-(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))} dx$$

input `integrate(x**2/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**2/((-c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))), x)`

**3.305.7 Maxima [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(x^2/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)), x)`

---

3.305.  $\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$

**3.305.8 Giac [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(x^2/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)), x)`**3.305.9 Mupad [N/A]**

Not integrable

Time = 3.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^2}{(a + b \operatorname{acosh}(cx)) (1 - c^2 x^2)^{3/2}} dx$$

input `int(x^2/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)),x)`output `int(x^2/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)), x)`

$$3.306 \quad \int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

3.306.1 Optimal result	2539
3.306.2 Mathematica [N/A]	2539
3.306.3 Rubi [N/A]	2540
3.306.4 Maple [N/A] (verified)	2540
3.306.5 Fracas [N/A]	2541
3.306.6 Sympy [N/A]	2541
3.306.7 Maxima [N/A]	2541
3.306.8 Giac [F(-2)]	2542
3.306.9 Mupad [N/A]	2542

### 3.306.1 Optimal result

Integrand size = 26, antiderivative size = 26

$$\int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

### 3.306.2 Mathematica [N/A]

Not integrable

Time = 6.91 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[x/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `Integrate[x/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

---


$$3.306. \quad \int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

**3.306.3 Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$$

↓ 6375

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$$

input `Int[x/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.306.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.306.4 Maple [N/A] (verified)**

Not integrable

Time = 1.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{x}{(-c^2 x^2 + 1)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

**3.306.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.35

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x/(a*c^4*x^4 - 2*a*c^2*x^2 + (b*c^4*x^4 - 2*b*c^2*x^2 + b)*arccosh(c*x) + a), x)`

**3.306.6 Sympy [N/A]**

Not integrable

Time = 8.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x}{(-(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))} dx$$

input `integrate(x/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(x/((-c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))), x)`

**3.306.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(x/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)), x)`

**3.306.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.306.9 Mupad [N/A]**

Not integrable

Time = 2.81 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x}{(a + b \operatorname{acosh}(cx)) (1 - c^2 x^2)^{3/2}} dx$$

input `int(x/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)),x)`

output `int(x/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)), x)`

$$3.307 \quad \int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

3.307.1 Optimal result	2543
3.307.2 Mathematica [N/A]	2543
3.307.3 Rubi [N/A]	2544
3.307.4 Maple [N/A] (verified)	2544
3.307.5 Fricas [N/A]	2545
3.307.6 Sympy [N/A]	2545
3.307.7 Maxima [N/A]	2545
3.307.8 Giac [N/A]	2546
3.307.9 Mupad [N/A]	2546

### 3.307.1 Optimal result

Integrand size = 25, antiderivative size = 25

$$\int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

### 3.307.2 Mathematica [N/A]

Not integrable

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[1/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `Integrate[1/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

---


$$3.307. \quad \int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$



**3.307.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))} dx$$

↓ 6325

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))} dx$$

input `Int[1/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.307.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.307.4 Maple [N/A] (verified)**

Not integrable

Time = 0.92 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

**3.307.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.40

$$\int \frac{1}{(1 - c^2x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(a*c^4*x^4 - 2*a*c^2*x^2 + (b*c^4*x^4 - 2*b*c^2*x^2 + b)*arccosh(c*x) + a), x)`

**3.307.6 Sympy [N/A]**

Not integrable

Time = 6.76 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{1}{(1 - c^2x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(-(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))} dx$$

input `integrate(1/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(1/((-c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))), x)`

**3.307.7 Maxima [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 - c^2x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(1/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)), x)`

---

3.307.  $\int \frac{1}{(1 - c^2x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$

**3.307.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(1/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)), x)`**3.307.9 Mupad [N/A]**

Not integrable

Time = 2.72 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx)) (1 - c^2 x^2)^{3/2}} dx$$

input `int(1/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)),x)`output `int(1/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)), x)`

**3.308** 
$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

3.308.1 Optimal result	2547
3.308.2 Mathematica [N/A]	2547
3.308.3 Rubi [N/A]	2548
3.308.4 Maple [N/A] (verified)	2548
3.308.5 Fricas [N/A]	2549
3.308.6 Sympy [N/A]	2549
3.308.7 Maxima [N/A]	2549
3.308.8 Giac [F(-2)]	2550
3.308.9 Mupad [N/A]	2550

**3.308.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)), x)`

**3.308.2 Mathematica [N/A]**

Not integrable

Time = 8.68 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[1/(x*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

output `Integrate[1/(x*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

**3.308.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\text{barccosh}(cx))} dx$$

↓ 6375

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\text{barccosh}(cx))} dx$$

input `Int[1/(x*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.308.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.308.4 Maple [N/A] (verified)**

Not integrable

Time = 1.00 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x(-c^2x^2+1)^{\frac{3}{2}}(a+b \operatorname{arccosh}(cx))} dx$$

input `int(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

**3.308.5 Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.29

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2x^2+1)^{\frac{3}{2}}(b\operatorname{arcosh}(cx)+a)x} dx$$

input `integrate(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2+1)/(a*c^4*x^5-2*a*c^2*x^3+a*x+(b*c^4*x^5-2*b*c^2*x^3+b*x)*arccosh(c*x)),x)`

**3.308.6 Sympy [N/A]**

Not integrable

Time = 21.22 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{x(-(cx-1)(cx+1))^{\frac{3}{2}}(a+b\operatorname{acosh}(cx))} dx$$

input `integrate(1/x/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(1/(x*(-(c*x-1)*(c*x+1))**(3/2)*(a+b*acosh(c*x))),x)`

**3.308.7 Maxima [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2x^2+1)^{\frac{3}{2}}(b\operatorname{arcosh}(cx)+a)x} dx$$

input `integrate(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(1/((-c^2*x^2+1)^(3/2)*(b*arccosh(c*x)+a)*x),x)`

**3.308.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx = \text{Exception raised: TypeError}$$

input `integrate(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.308.9 Mupad [N/A]**

Not integrable

Time = 3.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{1}{x(a+b\operatorname{acosh}(cx))(1-c^2x^2)^{3/2}} dx$$

input `int(1/(x*(a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)),x)`

output `int(1/(x*(a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)), x)`

$$3.309 \quad \int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

3.309.1 Optimal result	. . . . .	2551
3.309.2 Mathematica [N/A]	. . . . .	2551
3.309.3 Rubi [N/A]	. . . . .	2552
3.309.4 Maple [N/A] (verified)	. . . . .	2552
3.309.5 Fricas [N/A]	. . . . .	2553
3.309.6 Sympy [N/A]	. . . . .	2553
3.309.7 Maxima [N/A]	. . . . .	2553
3.309.8 Giac [N/A]	. . . . .	2554
3.309.9 Mupad [N/A]	. . . . .	2554

### 3.309.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)), x)`

### 3.309.2 Mathematica [N/A]

Not integrable

Time = 9.98 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[1/(x^2*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

output `Integrate[1/(x^2*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

---


$$3.309. \quad \int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$



**3.309.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \text{barccosh}(cx))} dx$$

↓ 6375

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \text{barccosh}(cx))} dx$$

input `Int[1/(x^2*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.309.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.309.4 Maple [N/A] (verified)**

Not integrable

Time = 1.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^2 (-c^2 x^2 + 1)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

**3.309.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.43

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a) x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(a*c^4*x^6 - 2*a*c^2*x^4 + a*x^2 + (b*c^4*x^6 - 2*b*c^2*x^4 + b*x^2)*arccosh(c*x)), x)`

**3.309.6 Sympy [N/A]**

Not integrable

Time = 48.17 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{x^2 (-(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))} dx$$

input `integrate(1/x**2/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(1/(x**2*(-(c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))), x)`

**3.309.7 Maxima [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a) x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(1/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)*x^2), x)`

---

3.309.  $\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$

**3.309.8 Giac [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a) x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(1/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)*x^2), x)`**3.309.9 Mupad [N/A]**

Not integrable

Time = 2.86 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{x^2 (a + b \operatorname{acosh}(cx)) (1 - c^2 x^2)^{3/2}} dx$$

input `int(1/(x^2*(a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)),x)`output `int(1/(x^2*(a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)), x)`

$$3.310 \quad \int \frac{x^m (1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx$$

3.310.1 Optimal result . . . . .	2555
3.310.2 Mathematica [N/A] . . . . .	2555
3.310.3 Rubi [N/A] . . . . .	2556
3.310.4 Maple [N/A] (verified) . . . . .	2556
3.310.5 Fricas [N/A] . . . . .	2557
3.310.6 Sympy [F(-1)] . . . . .	2557
3.310.7 Maxima [N/A] . . . . .	2557
3.310.8 Giac [F(-2)] . . . . .	2558
3.310.9 Mupad [N/A] . . . . .	2558

### 3.310.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^m (1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx = \operatorname{Int} \left( \frac{x^m (1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)}, x \right)$$

output `Unintegrable(x^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

### 3.310.2 Mathematica [N/A]

Not integrable

Time = 1.45 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^m (1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{x^m (1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx$$

input `Integrate[(x^m*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]),x]`

output `Integrate[(x^m*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]), x]`

---


$$3.310. \quad \int \frac{x^m (1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx$$

**3.310.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{3/2} x^m}{a + \operatorname{arccosh}(cx)} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{3/2} x^m}{a + \operatorname{arccosh}(cx)} dx$$

input `Int[(x^m*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x]),x]`

output `$Aborted`

**3.310.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.310.4 Maple [N/A] (verified)**

Not integrable

Time = 1.65 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^m (-c^2 x^2 + 1)^{\frac{3}{2}}}{a + b \operatorname{arccosh}(cx)} dx$$

input `int(x^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(x^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

---

3.310.  $\int \frac{x^m (1 - c^2 x^2)^{3/2}}{a + b \operatorname{arccosh}(cx)} dx$

**3.310.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.36

$$\int \frac{x^m(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x^m}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)*x^m/(b*arccosh(c*x) + a), x)`

**3.310.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^m(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \text{Timed out}$$

input `integrate(x**m*(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Timed out`

**3.310.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m(1-c^2x^2)^{3/2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x^m}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate(x^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x^m/(b*arccosh(c*x) + a), x)`

**3.310.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^m(1 - c^2x^2)^{3/2}}{a + b\operatorname{arccosh}(cx)} dx = \text{Exception raised: TypeError}$$

input `integrate(x^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.310.9 Mupad [N/A]**

Not integrable

Time = 2.74 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m(1 - c^2x^2)^{3/2}}{a + b\operatorname{arccosh}(cx)} dx = \int \frac{x^m(1 - c^2x^2)^{3/2}}{a + b\operatorname{acosh}(cx)} dx$$

input `int((x^m*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)),x)`

output `int((x^m*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x)), x)`

**3.311**  $\int \frac{x^m \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$

3.311.1 Optimal result . . . . .	2559
3.311.2 Mathematica [N/A] . . . . .	2559
3.311.3 Rubi [N/A] . . . . .	2560
3.311.4 Maple [N/A] (verified) . . . . .	2560
3.311.5 Fricas [N/A] . . . . .	2561
3.311.6 Sympy [N/A] . . . . .	2561
3.311.7 Maxima [N/A] . . . . .	2561
3.311.8 Giac [F(-2)] . . . . .	2562
3.311.9 Mupad [N/A] . . . . .	2562

**3.311.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^m \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \operatorname{Int}\left(\frac{x^m \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)}, x\right)$$

output `Unintegrable(x^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)), x)`

**3.311.2 Mathematica [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^m \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{x^m \sqrt{1-c^2x^2}}{a+b\operatorname{arccosh}(cx)} dx$$

input `Integrate[(x^m*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]), x]`

output `Integrate[(x^m*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]), x]`



**3.311.3 Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1 - c^2 x^2} x^m}{a + b \operatorname{arccosh}(cx)} dx$$

↓ 6375

$$\int \frac{\sqrt{1 - c^2 x^2} x^m}{a + b \operatorname{arccosh}(cx)} dx$$

input `Int[(x^m*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x]),x]`

output `$Aborted`

**3.311.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.311.4 Maple [N/A] (verified)**

Not integrable

Time = 1.35 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^m \sqrt{-c^2 x^2 + 1}}{a + b \operatorname{arccosh}(cx)} dx$$

input `int(x^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x)`

output `int(x^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x)`

**3.311.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^m}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^m/(b*arccosh(c*x) + a), x)`

**3.311.6 Sympy [N/A]**

Not integrable

Time = 2.67 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{x^m \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{x^m \sqrt{-(cx - 1)(cx + 1)}}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate(x**m*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**m*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x)), x)`

**3.311.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^m}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(x^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*x^2 + 1)*x^m/(b*arccosh(c*x) + a), x)`

---

3.311.  $\int \frac{x^m \sqrt{1 - c^2 x^2}}{a + b \operatorname{arccosh}(cx)} dx$

**3.311.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^m \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \text{Exception raised: TypeError}$$

input `integrate(x^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.311.9 Mupad [N/A]**

Not integrable

Time = 2.71 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m \sqrt{1 - c^2 x^2}}{a + \operatorname{barccosh}(cx)} dx = \int \frac{x^m \sqrt{1 - c^2 x^2}}{a + b \operatorname{acosh}(cx)} dx$$

input `int((x^m*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)),x)`

output `int((x^m*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x)), x)`

**3.312**  $\int \frac{x^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))} dx$

3.312.1 Optimal result . . . . .	2563
3.312.2 Mathematica [N/A] . . . . .	2563
3.312.3 Rubi [N/A] . . . . .	2564
3.312.4 Maple [N/A] (verified) . . . . .	2564
3.312.5 Fricas [N/A] . . . . .	2565
3.312.6 Sympy [N/A] . . . . .	2565
3.312.7 Maxima [N/A] . . . . .	2565
3.312.8 Giac [N/A] . . . . .	2566
3.312.9 Mupad [N/A] . . . . .	2566

**3.312.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))} dx = \text{Int}\left(\frac{x^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))}, x\right)$$

output `Unintegrable(x^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)), x)`

**3.312.2 Mathematica [N/A]**

Not integrable

Time = 1.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))} dx = \int \frac{x^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))} dx$$

input `Integrate[x^m/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])), x]`

output `Integrate[x^m/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])), x]`

**3.312.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+\operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+\operatorname{arccosh}(cx))} dx$$

input `Int[x^m/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.312.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.312.4 Maple [N/A] (verified)**

Not integrable

Time = 1.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^m}{\sqrt{-c^2x^2+1}(a+b\operatorname{arccosh}(cx))} dx$$

input `int(x^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x)`

output `int(x^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x)`

**3.312.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.79

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^m}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^m/(a*c^2*x^2 + (b*c^2*x^2 - b)*arccosh(c*x) - a), x)`

**3.312.6 Sympy [N/A]**

Not integrable

Time = 2.98 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^m}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))} dx$$

input `integrate(x**m/(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**m/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))), x)`

**3.312.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^m}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(x^m/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)), x)`

**3.312.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^m}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(x^m/(sqrt(-c^2*x^2+1)*(b*arccosh(c*x)+a)),x)`**3.312.9 Mupad [N/A]**

Not integrable

Time = 3.35 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(a+b\operatorname{acosh}(cx))\sqrt{1-c^2x^2}} dx$$

input `int(x^m/((a+b*acosh(c*x))*(1-c^2*x^2)^(1/2)),x)`output `int(x^m/((a+b*acosh(c*x))*(1-c^2*x^2)^(1/2)),x)`

**3.313**  $\int \frac{x^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$

3.313.1 Optimal result	2567
3.313.2 Mathematica [N/A]	2567
3.313.3 Rubi [N/A]	2568
3.313.4 Maple [N/A] (verified)	2568
3.313.5 Fricas [N/A]	2569
3.313.6 Sympy [N/A]	2569
3.313.7 Maxima [N/A]	2569
3.313.8 Giac [N/A]	2570
3.313.9 Mupad [N/A]	2570

**3.313.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{x^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(x^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)), x)`

**3.313.2 Mathematica [N/A]**

Not integrable

Time = 1.49 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[x^m/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

output `Integrate[x^m/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`



**3.313.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{arccosh}(cx))} dx$$

input `Int[x^m/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.313.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.313.4 Maple [N/A] (verified)**

Not integrable

Time = 1.46 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^m}{(-c^2 x^2 + 1)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(x^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(x^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x)`

**3.313.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.25

$$\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^m/(a*c^4*x^4 - 2*a*c^2*x^2 + (b*c^4*x^4 - 2*b*c^2*x^2 + b)*arccosh(c*x) + a), x)`

**3.313.6 Sympy [N/A]**

Not integrable

Time = 83.05 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(-(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))} dx$$

input `integrate(x**m/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(x**m/((-c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))), x)`

**3.313.7 Maxima [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(x^m/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)), x)`

---

3.313.  $\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$

**3.313.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(x^m/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)), x)`**3.313.9 Mupad [N/A]**

Not integrable

Time = 3.47 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(a + b \operatorname{acosh}(cx)) (1 - c^2 x^2)^{3/2}} dx$$

input `int(x^m/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)),x)`output `int(x^m/((a + b*acosh(c*x))*(1 - c^2*x^2)^(3/2)), x)`

**3.314**  $\int \frac{x^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))} dx$

3.314.1 Optimal result	2571
3.314.2 Mathematica [N/A]	2571
3.314.3 Rubi [N/A]	2572
3.314.4 Maple [N/A] (verified)	2572
3.314.5 Fricas [N/A]	2573
3.314.6 Sympy [F(-1)]	2573
3.314.7 Maxima [N/A]	2573
3.314.8 Giac [N/A]	2574
3.314.9 Mupad [N/A]	2574

**3.314.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{x^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(x^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)), x)`

**3.314.2 Mathematica [N/A]**

Not integrable

Time = 1.91 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{x^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[x^m/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])), x]`

output `Integrate[x^m/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])), x]`

**3.314.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{arccosh}(cx))} dx$$

↓ 6375

$$\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{arccosh}(cx))} dx$$

input `Int[x^m/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.314.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.314.4 Maple [N/A] (verified)**

Not integrable

Time = 1.99 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^m}{(-c^2 x^2 + 1)^{5/2} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(x^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x)`

output `int(x^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x)`

**3.314.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 86, normalized size of antiderivative = 3.07

$$\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(-c^2 x^2 + 1)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^m/(a*c^6*x^6 - 3*a*c^4*x^4 + 3*a*c^2*x^2 + (b*c^6*x^6 - 3*b*c^4*x^4 + 3*b*c^2*x^2 - b)*arccosh(c*x) - a), x)`

**3.314.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx = \text{Timed out}$$

input `integrate(x**m/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x)),x)`

output `Timed out`

**3.314.7 Maxima [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(-c^2 x^2 + 1)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(x^m/((-c^2*x^2 + 1)^(5/2)*(b*arccosh(c*x) + a)), x)`

---

3.314.  $\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx$

**3.314.8 Giac [N/A]**

Not integrable

Time = 0.35 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(-c^2 x^2 + 1)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(x^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(x^m/((-c^2*x^2 + 1)^(5/2)*(b*arccosh(c*x) + a)), x)`**3.314.9 Mupad [N/A]**

Not integrable

Time = 3.83 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^m}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx = \int \frac{x^m}{(a + b \operatorname{acosh}(cx)) (1 - c^2 x^2)^{5/2}} dx$$

input `int(x^m/((a + b*acosh(c*x))*(1 - c^2*x^2)^(5/2)),x)`output `int(x^m/((a + b*acosh(c*x))*(1 - c^2*x^2)^(5/2)), x)`

**3.315**  $\int \frac{(c-a^2cx^2)^3}{\operatorname{arccosh}(ax)^2} dx$

3.315.1 Optimal result . . . . .	2575
3.315.2 Mathematica [B] (warning: unable to verify) . . . . .	2575
3.315.3 Rubi [A] (verified) . . . . .	2576
3.315.4 Maple [A] (verified) . . . . .	2578
3.315.5 Fricas [F] . . . . .	2578
3.315.6 Sympy [F] . . . . .	2578
3.315.7 Maxima [F] . . . . .	2579
3.315.8 Giac [F] . . . . .	2579
3.315.9 Mupad [F(-1)] . . . . .	2580

**3.315.1 Optimal result**

Integrand size = 20, antiderivative size = 98

$$\int \frac{(c - a^2cx^2)^3}{\operatorname{arccosh}(ax)^2} dx = \frac{c^3(-1 + ax)^{7/2}(1 + ax)^{7/2}}{a\operatorname{arccosh}(ax)} + \frac{35c^3\operatorname{Chi}(\operatorname{arccosh}(ax))}{64a} - \frac{63c^3\operatorname{Chi}(3\operatorname{arccosh}(ax))}{64a} + \frac{35c^3\operatorname{Chi}(5\operatorname{arccosh}(ax))}{64a} - \frac{7c^3\operatorname{Chi}(7\operatorname{arccosh}(ax))}{64a}$$

```
output c^3*(a*x-1)^(7/2)*(a*x+1)^(7/2)/a/arccosh(a*x)+35/64*c^3*Chi(arccosh(a*x))
/a-63/64*c^3*Chi(3*arccosh(a*x))/a+35/64*c^3*Chi(5*arccosh(a*x))/a-7/64*c^
3*Chi(7*arccosh(a*x))/a
```

**3.315.2 Mathematica [B] (warning: unable to verify)**

Leaf count is larger than twice the leaf count of optimal. 257 vs. 2(98) = 196.

Time = 0.53 (sec) , antiderivative size = 257, normalized size of antiderivative = 2.62

$$\int \frac{(c - a^2cx^2)^3}{\operatorname{arccosh}(ax)^2} dx = \frac{c^3 \left( -64\sqrt{\frac{-1+ax}{1+ax}} - 64ax\sqrt{\frac{-1+ax}{1+ax}} + 192a^2x^2\sqrt{\frac{-1+ax}{1+ax}} + 192a^3x^3\sqrt{\frac{-1+ax}{1+ax}} - 192a^4x^4\sqrt{\frac{-1+ax}{1+ax}} - 192a^5x^5\sqrt{\frac{-1+ax}{1+ax}} \right)}{\dots}$$



input `Integrate[(c - a^2*c*x^2)^3/ArcCosh[a*x]^2,x]`

output  $(c^3*(-64*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] - 64*a*x*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 192*a^2*x^2*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 192*a^3*x^3*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] - 192*a^4*x^4*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] - 192*a^5*x^5*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 64*a^6*x^6*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 64*a^7*x^7*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 35*\text{ArcCosh}[a*x]*\text{CoshIntegral}[\text{ArcCosh}[a*x]] - 63*\text{ArcCosh}[a*x]*\text{CoshIntegral}[3*\text{ArcCosh}[a*x]] + 35*\text{ArcCosh}[a*x]*\text{CoshIntegral}[5*\text{ArcCosh}[a*x]] - 7*\text{ArcCosh}[a*x]*\text{CoshIntegral}[7*\text{ArcCosh}[a*x]]))/ (64*a*\text{ArcCosh}[a*x])$

### 3.315.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.85, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6319, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - a^2 cx^2)^3}{\text{arccosh}(ax)^2} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{c^3(ax - 1)^{7/2}(ax + 1)^{7/2}}{a \text{arccosh}(ax)} - 7ac^3 \int \frac{x(ax - 1)^{5/2}(ax + 1)^{5/2}}{\text{arccosh}(ax)} dx \\
 & \quad \downarrow \text{6368} \\
 & \frac{c^3(ax - 1)^{7/2}(ax + 1)^{7/2}}{a \text{arccosh}(ax)} - \frac{7c^3 \int \frac{ax(ax-1)^3(ax+1)^3}{\text{arccosh}(ax)} d\text{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{5971} \\
 & \frac{c^3(ax - 1)^{7/2}(ax + 1)^{7/2}}{a \text{arccosh}(ax)} - \\
 & \frac{7c^3 \int \left( -\frac{5ax}{64 \text{arccosh}(ax)} + \frac{9 \cosh(3 \text{arccosh}(ax))}{64 \text{arccosh}(ax)} - \frac{5 \cosh(5 \text{arccosh}(ax))}{64 \text{arccosh}(ax)} + \frac{\cosh(7 \text{arccosh}(ax))}{64 \text{arccosh}(ax)} \right) d\text{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.315.  $\int \frac{(c - a^2 cx^2)^3}{\text{arccosh}(ax)^2} dx$

$$\frac{c^3(ax-1)^{7/2}(ax+1)^{7/2}}{a \operatorname{arccosh}(ax)} - \frac{7c^3\left(-\frac{5}{64}\operatorname{Chi}(\operatorname{arccosh}(ax)) + \frac{9}{64}\operatorname{Chi}(3\operatorname{arccosh}(ax)) - \frac{5}{64}\operatorname{Chi}(5\operatorname{arccosh}(ax)) + \frac{1}{64}\operatorname{Chi}(7\operatorname{arccosh}(ax))\right)}{a}$$

input `Int[(c - a^2*c*x^2)^3/ArcCosh[a*x]^2,x]`

output `(c^3*(-1 + a*x)^(7/2)*(1 + a*x)^(7/2))/(a*ArcCosh[a*x]) - (7*c^3*((-5*CoshIntegral[ArcCosh[a*x]])/64 + (9*CoshIntegral[3*ArcCosh[a*x]])/64 - (5*CoshIntegral[5*ArcCosh[a*x]])/64 + CoshIntegral[7*ArcCosh[a*x]]/64))/a`

### 3.315.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(q_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^q/(-1 + c*x)^q] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

**3.315.4 Maple [A] (verified)**

Time = 0.85 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{c^3(35 \operatorname{Chi}(\operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 63 \operatorname{Chi}(3 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) + 35 \operatorname{Chi}(5 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 7 \operatorname{Chi}(7 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 35(a^2cx^2 - c)^{3/2} \operatorname{arccosh}(ax) + 64a \operatorname{arccosh}(ax))}{64a \operatorname{arccosh}(ax)^2}$
default	$\frac{c^3(35 \operatorname{Chi}(\operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 63 \operatorname{Chi}(3 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) + 35 \operatorname{Chi}(5 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 7 \operatorname{Chi}(7 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 35(a^2cx^2 - c)^{3/2} \operatorname{arccosh}(ax) + 64a \operatorname{arccosh}(ax))}{64a \operatorname{arccosh}(ax)^2}$

input `int((-a^2*c*x^2+c)^3/arccosh(a*x)^2,x,method=_RETURNVERBOSE)`output  $\frac{1/64/a*c^3*(35*\operatorname{Chi}(\operatorname{arccosh}(a*x))*\operatorname{arccosh}(a*x)-63*\operatorname{Chi}(3*\operatorname{arccosh}(a*x))*\operatorname{arccosh}(a*x)+35*\operatorname{Chi}(5*\operatorname{arccosh}(a*x))*\operatorname{arccosh}(a*x)-7*\operatorname{Chi}(7*\operatorname{arccosh}(a*x))*\operatorname{arccosh}(a*x)-35*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}+21*\sinh(3*\operatorname{arccosh}(a*x))-7*\sinh(5*\operatorname{arccosh}(a*x))+\sinh(7*\operatorname{arccosh}(a*x)))}{\operatorname{arccosh}(a*x)^2}$ **3.315.5 Fracas [F]**

$$\int \frac{(c - a^2cx^2)^3}{\operatorname{arccosh}(ax)^2} dx = \int -\frac{(a^2cx^2 - c)^3}{\operatorname{arccosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)^3/arccosh(a*x)^2,x, algorithm="fracas")`output `integral(-a^6*c^3*x^6 - 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 - c^3)/arccosh(a*x)^2, x)`**3.315.6 Sympy [F]**

$$\int \frac{(c - a^2cx^2)^3}{\operatorname{arccosh}(ax)^2} dx = -c^3 \left( \int \frac{3a^2x^2}{\operatorname{acosh}^2(ax)} dx + \int \left( -\frac{3a^4x^4}{\operatorname{acosh}^2(ax)} \right) dx + \int \frac{a^6x^6}{\operatorname{acosh}^2(ax)} dx + \int \left( -\frac{1}{\operatorname{acosh}^2(ax)} \right) dx \right)$$

input `integrate((-a**2*c*x**2+c)**3/acosh(a*x)**2,x)`

---

3.315.  $\int \frac{(c - a^2cx^2)^3}{\operatorname{arccosh}(ax)^2} dx$

output `-c**3*(Integral(3*a**2*x**2/acosh(a*x)**2, x) + Integral(-3*a**4*x**4/acosh(a*x)**2, x) + Integral(a**6*x**6/acosh(a*x)**2, x) + Integral(-1/acosh(a*x)**2, x))`

### 3.315.7 Maxima [F]

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)^2} dx = \int -\frac{(a^2 cx^2 - c)^3}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)^3/arccosh(a*x)^2,x, algorithm="maxima")`

output `(a^9*c^3*x^9 - 4*a^7*c^3*x^7 + 6*a^5*c^3*x^5 - 4*a^3*c^3*x^3 + a*c^3*x + (a^8*c^3*x^8 - 4*a^6*c^3*x^6 + 6*a^4*c^3*x^4 - 4*a^2*c^3*x^2 + c^3)*sqrt(a*x + 1)*sqrt(a*x - 1))/((a^3*x^2 + sqrt(a*x + 1)*sqrt(a*x - 1)*a^2*x - a)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))) - integrate((7*a^10*c^3*x^10 - 29*a^8*c^3*x^8 + 46*a^6*c^3*x^6 - 34*a^4*c^3*x^4 + 11*a^2*c^3*x^2 + (7*a^8*c^3*x^8 - 20*a^6*c^3*x^6 + 18*a^4*c^3*x^4 - 4*a^2*c^3*x^2 - c^3)*(a*x + 1)*(a*x - 1) - c^3 + 7*(2*a^9*c^3*x^9 - 7*a^7*c^3*x^7 + 9*a^5*c^3*x^5 - 5*a^3*c^3*x^3 + a*c^3*x)*sqrt(a*x + 1)*sqrt(a*x - 1))/((a^4*x^4 + (a*x + 1)*(a*x - 1)*a^2*x^2 - 2*a^2*x^2 + 2*(a^3*x^3 - a*x)*sqrt(a*x + 1)*sqrt(a*x - 1) + 1)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))), x)`

### 3.315.8 Giac [F]

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)^2} dx = \int -\frac{(a^2 cx^2 - c)^3}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)^3/arccosh(a*x)^2,x, algorithm="giac")`

output `integrate(-(a^2*c*x^2 - c)^3/arccosh(a*x)^2, x)`

**3.315.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^3}{\operatorname{arccosh}(ax)^2} dx = \int \frac{(c - a^2 cx^2)^3}{\operatorname{acosh}(ax)^2} dx$$

input `int((c - a^2*c*x^2)^3/acosh(a*x)^2,x)`output `int((c - a^2*c*x^2)^3/acosh(a*x)^2, x)`

**3.316**  $\int \frac{(c-a^2cx^2)^2}{\operatorname{arccosh}(ax)^2} dx$

3.316.1 Optimal result . . . . . 2581  
 3.316.2 Mathematica [B] (warning: unable to verify) . . . . . 2581  
 3.316.3 Rubi [A] (verified) . . . . . 2582  
 3.316.4 Maple [A] (verified) . . . . . 2583  
 3.316.5 Fricas [F] . . . . . 2584  
 3.316.6 Sympy [F] . . . . . 2584  
 3.316.7 Maxima [F] . . . . . 2584  
 3.316.8 Giac [F] . . . . . 2585  
 3.316.9 Mupad [F(-1)] . . . . . 2585

**3.316.1 Optimal result**

Integrand size = 20, antiderivative size = 82

$$\int \frac{(c - a^2cx^2)^2}{\operatorname{arccosh}(ax)^2} dx = -\frac{c^2(-1 + ax)^{5/2}(1 + ax)^{5/2}}{a\operatorname{arccosh}(ax)} + \frac{5c^2\operatorname{Chi}(\operatorname{arccosh}(ax))}{8a} - \frac{15c^2\operatorname{Chi}(3\operatorname{arccosh}(ax))}{16a} + \frac{5c^2\operatorname{Chi}(5\operatorname{arccosh}(ax))}{16a}$$

output

```
-c^2*(a*x-1)^(5/2)*(a*x+1)^(5/2)/a/arccosh(a*x)+5/8*c^2*Chi(arccosh(a*x))/a-15/16*c^2*Chi(3*arccosh(a*x))/a+5/16*c^2*Chi(5*arccosh(a*x))/a
```

**3.316.2 Mathematica [B] (warning: unable to verify)**

Leaf count is larger than twice the leaf count of optimal. 194 vs. 2(82) = 164.

Time = 0.34 (sec) , antiderivative size = 194, normalized size of antiderivative = 2.37

$$\int \frac{(c - a^2cx^2)^2}{\operatorname{arccosh}(ax)^2} dx = \frac{c^2 \left( 16\sqrt{\frac{-1+ax}{1+ax}} + 16ax\sqrt{\frac{-1+ax}{1+ax}} - 32a^2x^2\sqrt{\frac{-1+ax}{1+ax}} - 32a^3x^3\sqrt{\frac{-1+ax}{1+ax}} + 16a^4x^4\sqrt{\frac{-1+ax}{1+ax}} + 16a^5x^5\sqrt{\frac{-1+ax}{1+ax}} \right)}{16a\operatorname{arccosh}(ax)^2}$$

input

```
Integrate[(c - a^2*c*x^2)^2/ArcCosh[a*x]^2,x]
```

3.316.  $\int \frac{(c-a^2cx^2)^2}{\operatorname{arccosh}(ax)^2} dx$

output 
$$\frac{-1/16*(c^2*(16*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 16*a*x*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] - 32*a^2*x^2*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] - 32*a^3*x^3*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 16*a^4*x^4*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] + 16*a^5*x^5*\text{Sqrt}[(-1 + a*x)/(1 + a*x)] - 10*\text{ArcCosh}[a*x]*\text{CoshIntegral}[\text{ArcCosh}[a*x]] + 15*\text{ArcCosh}[a*x]*\text{CoshIntegral}[3*\text{ArcCosh}[a*x]] - 5*\text{ArcCosh}[a*x]*\text{CoshIntegral}[5*\text{ArcCosh}[a*x]])}{a*\text{ArcCosh}[a*x]}$$

### 3.316.3 Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.89, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {6319, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(c - a^2 cx^2)^2}{\text{arccosh}(ax)^2} dx \\ & \quad \downarrow \text{6319} \\ & 5ac^2 \int \frac{x(ax-1)^{3/2}(ax+1)^{3/2}}{\text{arccosh}(ax)} dx - \frac{c^2(ax-1)^{5/2}(ax+1)^{5/2}}{a\text{arccosh}(ax)} \\ & \quad \downarrow \text{6368} \\ & \frac{5c^2 \int \frac{ax(ax-1)^2(ax+1)^2}{\text{arccosh}(ax)} d\text{arccosh}(ax)}{a} - \frac{c^2(ax-1)^{5/2}(ax+1)^{5/2}}{a\text{arccosh}(ax)} \\ & \quad \downarrow \text{5971} \\ & \frac{5c^2 \int \left( \frac{ax}{8\text{arccosh}(ax)} - \frac{3\cosh(3\text{arccosh}(ax))}{16\text{arccosh}(ax)} + \frac{\cosh(5\text{arccosh}(ax))}{16\text{arccosh}(ax)} \right) d\text{arccosh}(ax)}{a} - \frac{c^2(ax-1)^{5/2}(ax+1)^{5/2}}{a\text{arccosh}(ax)} \\ & \quad \downarrow \text{2009} \\ & \frac{5c^2 \left( \frac{1}{8}\text{Chi}(\text{arccosh}(ax)) - \frac{3}{16}\text{Chi}(3\text{arccosh}(ax)) + \frac{1}{16}\text{Chi}(5\text{arccosh}(ax)) \right)}{a} - \frac{c^2(ax-1)^{5/2}(ax+1)^{5/2}}{a\text{arccosh}(ax)} \end{aligned}$$

input  $\text{Int}[(c - a^2*c*x^2)^2/\text{ArcCosh}[a*x]^2, x]$

---

3.316.  $\int \frac{(c - a^2 cx^2)^2}{\text{arccosh}(ax)^2} dx$

output  $-\left(\frac{c^2(-1+ax)^{5/2}(1+ax)^{5/2}}{a \operatorname{ArcCosh}[ax]}\right) + \frac{5c^2(\operatorname{CoshIntegral}[\operatorname{ArcCosh}[ax]]/8 - (3\operatorname{CoshIntegral}[3\operatorname{ArcCosh}[ax]])/16 + \operatorname{CoshIntegral}[5\operatorname{ArcCosh}[ax]]/16)}{a}$

### 3.316.3.1 Defintions of rubi rules used

rule 2009  $\operatorname{Int}[u_, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$

rule 5971  $\operatorname{Int}[\operatorname{Cosh}[a_] + (b_)(x_)^{(p_)} * ((c_) + (d_)(x_))^{(m_)} * \operatorname{Sinh}[(a_) + (b_)(x_)]^{(n_)}, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandTrigReduce}[(c + dx)^m, \operatorname{Sinh}[a + bx]^{n*} \operatorname{Cosh}[a + bx]^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, m, x\} \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{IGtQ}[p, 0]$

rule 6319  $\operatorname{Int}[(a_) + \operatorname{ArcCosh}[(c_)(x_)] * (b_)]^{(n_)} * ((d_) + (e_)(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Simp}[\operatorname{Sqrt}[1 + cx] * \operatorname{Sqrt}[-1 + cx] * (d + ex^2)^p * ((a + b \operatorname{ArcCosh}[cx])^{(n+1)}) / (b * c * (n+1)), x] - \operatorname{Simp}[c * ((2p + 1) / (b * (n + 1))) * \operatorname{Simp}[(d + ex^2)^p / ((1 + cx)^p * (-1 + cx)^p)] \operatorname{Int}[x * (1 + cx)^{(p-1/2)} * (-1 + cx)^{(p-1/2)} * (a + b \operatorname{ArcCosh}[cx])^{(n+1)}, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, p, x\} \ \&\& \operatorname{EqQ}[c^2 * d + e, 0] \ \&\& \operatorname{LtQ}[n, -1] \ \&\& \operatorname{IntegerQ}[2 * p]$

rule 6368  $\operatorname{Int}[(a_) + \operatorname{ArcCosh}[(c_)(x_)] * (b_)]^{(n_)} * (x_)^{(m_)} * ((d1_) + (e1_)(x_))^{(p_)} * ((d2_) + (e2_)(x_))^{(p_)}, x\_Symbol] \rightarrow \operatorname{Simp}[(1 / (b * c^{(m+1)})) * \operatorname{Simp}[(d1 + e1 * x)^p / (1 + cx)^p] * \operatorname{Simp}[(d2 + e2 * x)^p / (-1 + cx)^p] \operatorname{Subst}[\operatorname{Int}[x^n * \operatorname{Cosh}[-a/b + x/b]^m * \operatorname{Sinh}[-a/b + x/b]^{(2 * p + 1)}, x], x, a + b \operatorname{ArcCosh}[cx]], x] /; \operatorname{FreeQ}\{a, b, c, d1, e1, d2, e2, n, x\} \ \&\& \operatorname{EqQ}[e1, c * d1] \ \&\& \operatorname{EqQ}[e2, (-c) * d2] \ \&\& \operatorname{IGtQ}[p + 3/2, 0] \ \&\& \operatorname{IGtQ}[m, 0]$

### 3.316.4 Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.06

method	result
derivativedivides	$\frac{c^2(10 \operatorname{Chi}(\operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 15 \operatorname{Chi}(3 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) + 5 \operatorname{Chi}(5 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 10\sqrt{ax} - 16a \operatorname{arccosh}(ax))}{16a \operatorname{arccosh}(ax)}$
default	$\frac{c^2(10 \operatorname{Chi}(\operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 15 \operatorname{Chi}(3 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) + 5 \operatorname{Chi}(5 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 10\sqrt{ax} - 16a \operatorname{arccosh}(ax))}{16a \operatorname{arccosh}(ax)}$

3.316.  $\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)^2} dx$



input `int((-a^2*c*x^2+c)^2/arccosh(a*x)^2,x,method=_RETURNVERBOSE)`

output `1/16/a*c^2*(10*Chi(arccosh(a*x))*arccosh(a*x)-15*Chi(3*arccosh(a*x))*arccosh(a*x)+5*Chi(5*arccosh(a*x))*arccosh(a*x)-10*(a*x-1)^(1/2)*(a*x+1)^(1/2)+5*sinh(3*arccosh(a*x))-sinh(5*arccosh(a*x)))/arccosh(a*x)`

### 3.316.5 Fracas [F]

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)^2} dx = \int \frac{(a^2 cx^2 - c)^2}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)^2/arccosh(a*x)^2,x, algorithm="fricas")`

output `integral((a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)/arccosh(a*x)^2, x)`

### 3.316.6 Sympy [F]

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)^2} dx = c^2 \left( \int \left( -\frac{2a^2 x^2}{\operatorname{acosh}^2(ax)} \right) dx + \int \frac{a^4 x^4}{\operatorname{acosh}^2(ax)} dx + \int \frac{1}{\operatorname{acosh}^2(ax)} dx \right)$$

input `integrate((-a**2*c*x**2+c)**2/acosh(a*x)**2,x)`

output `c**2*(Integral(-2*a**2*x**2/acosh(a*x)**2, x) + Integral(a**4*x**4/acosh(a*x)**2, x) + Integral(acosh(a*x)**(-2), x))`

### 3.316.7 Maxima [F]

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)^2} dx = \int \frac{(a^2 cx^2 - c)^2}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)^2/arccosh(a*x)^2,x, algorithm="maxima")`

output `-(a^7*c^2*x^7 - 3*a^5*c^2*x^5 + 3*a^3*c^2*x^3 - a*c^2*x + (a^6*c^2*x^6 - 3*a^4*c^2*x^4 + 3*a^2*c^2*x^2 - c^2)*sqrt(a*x + 1)*sqrt(a*x - 1))/((a^3*x^2 + sqrt(a*x + 1)*sqrt(a*x - 1)*a^2*x - a)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))) + integrate((5*a^8*c^2*x^8 - 16*a^6*c^2*x^6 + 18*a^4*c^2*x^4 - 8*a^2*c^2*x^2 + (5*a^6*c^2*x^6 - 9*a^4*c^2*x^4 + 3*a^2*c^2*x^2 + c^2)*(a*x + 1)*(a*x - 1) + 5*(2*a^7*c^2*x^7 - 5*a^5*c^2*x^5 + 4*a^3*c^2*x^3 - a*c^2*x)*sqrt(a*x + 1)*sqrt(a*x - 1) + c^2)/((a^4*x^4 + (a*x + 1)*(a*x - 1)*a^2*x^2 - 2*a^2*x^2 + 2*(a^3*x^3 - a*x)*sqrt(a*x + 1)*sqrt(a*x - 1) + 1)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))), x)`

### 3.316.8 Giac [F]

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)^2} dx = \int \frac{(a^2 cx^2 - c)^2}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)^2/arccosh(a*x)^2,x, algorithm="giac")`

output `integrate((a^2*c*x^2 - c)^2/arccosh(a*x)^2, x)`

### 3.316.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(c - a^2 cx^2)^2}{\operatorname{arccosh}(ax)^2} dx = \int \frac{(c - a^2 cx^2)^2}{\operatorname{acosh}(ax)^2} dx$$

input `int((c - a^2*c*x^2)^2/acosh(a*x)^2,x)`

output `int((c - a^2*c*x^2)^2/acosh(a*x)^2, x)`

### 3.317 $\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx$

3.317.1 Optimal result . . . . .	2586
3.317.2 Mathematica [A] (verified) . . . . .	2586
3.317.3 Rubi [A] (verified) . . . . .	2587
3.317.4 Maple [A] (verified) . . . . .	2588
3.317.5 Fracas [F] . . . . .	2589
3.317.6 Sympy [F] . . . . .	2589
3.317.7 Maxima [F] . . . . .	2589
3.317.8 Giac [F] . . . . .	2590
3.317.9 Mupad [F(-1)] . . . . .	2590

#### 3.317.1 Optimal result

Integrand size = 18, antiderivative size = 58

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx = \frac{c(-1 + ax)^{3/2}(1 + ax)^{3/2}}{a \operatorname{arccosh}(ax)} + \frac{3c \operatorname{Chi}(\operatorname{arccosh}(ax))}{4a} - \frac{3c \operatorname{Chi}(3 \operatorname{arccosh}(ax))}{4a}$$

```
output c*(a*x-1)^(3/2)*(a*x+1)^(3/2)/a/arccosh(a*x)+3/4*c*Chi(arccosh(a*x))/a-3/4
*c*Chi(3*arccosh(a*x))/a
```

#### 3.317.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.12

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx = \frac{c \left( 4 \left( \frac{-1+ax}{1+ax} \right)^{3/2} (1 + ax)^3 + 3 \operatorname{arccosh}(ax) \operatorname{Chi}(\operatorname{arccosh}(ax)) - 3 \operatorname{arccosh}(ax) \operatorname{Chi}(3 \operatorname{arccosh}(ax)) \right)}{4a \operatorname{arccosh}(ax)}$$

```
input Integrate[(c - a^2*c*x^2)/ArcCosh[a*x]^2,x]
```

```
output (c*(4*((-1 + a*x)/(1 + a*x))^(3/2)*(1 + a*x)^3 + 3*ArcCosh[a*x]*CoshIntegral[ArcCosh[a*x]] - 3*ArcCosh[a*x]*CoshIntegral[3*ArcCosh[a*x]]))/(4*a*ArcCosh[a*x])
```

**3.317.3 Rubi [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.98, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {6319, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{c(ax - 1)^{3/2}(ax + 1)^{3/2}}{a \operatorname{arccosh}(ax)} - 3ac \int \frac{x\sqrt{ax - 1}\sqrt{ax + 1}}{\operatorname{arccosh}(ax)} dx \\
 & \quad \downarrow \text{6368} \\
 & \frac{c(ax - 1)^{3/2}(ax + 1)^{3/2}}{a \operatorname{arccosh}(ax)} - \frac{3c \int \frac{ax(ax-1)(ax+1)}{\operatorname{arccosh}(ax)} d\operatorname{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{5971} \\
 & \frac{c(ax - 1)^{3/2}(ax + 1)^{3/2}}{a \operatorname{arccosh}(ax)} - \frac{3c \int \left( \frac{\cosh(3\operatorname{arccosh}(ax))}{4\operatorname{arccosh}(ax)} - \frac{ax}{4\operatorname{arccosh}(ax)} \right) d\operatorname{arccosh}(ax)}{a} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c(ax - 1)^{3/2}(ax + 1)^{3/2}}{a \operatorname{arccosh}(ax)} - \frac{3c \left( \frac{1}{4} \operatorname{Chi}(3\operatorname{arccosh}(ax)) - \frac{1}{4} \operatorname{Chi}(\operatorname{arccosh}(ax)) \right)}{a}
 \end{aligned}$$

input `Int[(c - a^2*c*x^2)/ArcCosh[a*x]^2,x]`

output `(c*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2))/(a*ArcCosh[a*x]) - (3*c*(-1/4*CoshIntegral[ArcCosh[a*x]] + CoshIntegral[3*ArcCosh[a*x]]/4))/a`

### 3.317.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^(n)*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_.) + (e1_.)*(x_)^(p_.))*((d2_.) + (e2_.)*(x_)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.317.4 Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.05

method	result	size
derivativedivides	$\frac{c(3 \operatorname{Chi}(\operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 3 \operatorname{Chi}(3 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 3\sqrt{ax-1}\sqrt{ax+1} + \sinh(3 \operatorname{arccosh}(ax)))}{4a \operatorname{arccosh}(ax)}$	61
default	$\frac{c(3 \operatorname{Chi}(\operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 3 \operatorname{Chi}(3 \operatorname{arccosh}(ax)) \operatorname{arccosh}(ax) - 3\sqrt{ax-1}\sqrt{ax+1} + \sinh(3 \operatorname{arccosh}(ax)))}{4a \operatorname{arccosh}(ax)}$	61

input `int((-a^2*c*x^2+c)/arccosh(a*x)^2,x,method=_RETURNVERBOSE)`

output `1/4/a*c*(3*Chi(arccosh(a*x))*arccosh(a*x)-3*Chi(3*arccosh(a*x))*arccosh(a*x)-3*(a*x-1)^(1/2)*(a*x+1)^(1/2)+sinh(3*arccosh(a*x)))/arccosh(a*x)`

3.317. 
$$\int \frac{c-a^2cx^2}{\operatorname{arccosh}(ax)^2} dx$$

**3.317.5 Fracas [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx = \int -\frac{a^2 cx^2 - c}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)/arccosh(a*x)^2,x, algorithm="fricas")`

output `integral(-(a^2*c*x^2 - c)/arccosh(a*x)^2, x)`

**3.317.6 Sympy [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx = -c \left( \int \frac{a^2 x^2}{\operatorname{acosh}^2(ax)} dx + \int \left( -\frac{1}{\operatorname{acosh}^2(ax)} \right) dx \right)$$

input `integrate((-a**2*c*x**2+c)/acosh(a*x)**2,x)`

output `-c*(Integral(a**2*x**2/acosh(a*x)**2, x) + Integral(-1/acosh(a*x)**2, x))`

**3.317.7 Maxima [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx = \int -\frac{a^2 cx^2 - c}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)/arccosh(a*x)^2,x, algorithm="maxima")`

output `(a^5*c*x^5 - 2*a^3*c*x^3 + a*c*x + (a^4*c*x^4 - 2*a^2*c*x^2 + c)*sqrt(a*x + 1)*sqrt(a*x - 1))/((a^3*x^2 + sqrt(a*x + 1)*sqrt(a*x - 1)*a^2*x - a)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))) - integrate((3*a^6*c*x^6 - 7*a^4*c*x^4 + 5*a^2*c*x^2 + (3*a^4*c*x^4 - 2*a^2*c*x^2 - c)*(a*x + 1)*(a*x - 1) + 3*(2*a^5*c*x^5 - 3*a^3*c*x^3 + a*c*x)*sqrt(a*x + 1)*sqrt(a*x - 1) - c)/((a^4*x^4 + (a*x + 1)*(a*x - 1)*a^2*x^2 - 2*a^2*x^2 + 2*(a^3*x^3 - a*x)*sqrt(a*x + 1)*sqrt(a*x - 1) + 1)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))), x)`

**3.317.8 Giac [F]**

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx = \int -\frac{a^2 cx^2 - c}{\operatorname{arcosh}(ax)^2} dx$$

input `integrate((-a^2*c*x^2+c)/arccosh(a*x)^2,x, algorithm="giac")`

output `integrate(-(a^2*c*x^2 - c)/arccosh(a*x)^2, x)`

**3.317.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{c - a^2 cx^2}{\operatorname{arccosh}(ax)^2} dx = \int \frac{c - a^2 cx^2}{\operatorname{acosh}(ax)^2} dx$$

input `int((c - a^2*c*x^2)/acosh(a*x)^2,x)`

output `int((c - a^2*c*x^2)/acosh(a*x)^2, x)`

**3.318**  $\int \frac{1}{(c - a^2cx^2) \operatorname{arccosh}(ax)^2} dx$

3.318.1 Optimal result . . . . . 2591  
 3.318.2 Mathematica [N/A] . . . . . 2591  
 3.318.3 Rubi [N/A] . . . . . 2592  
 3.318.4 Maple [N/A] (verified) . . . . . 2593  
 3.318.5 Fricas [N/A] . . . . . 2593  
 3.318.6 Sympy [N/A] . . . . . 2593  
 3.318.7 Maxima [N/A] . . . . . 2594  
 3.318.8 Giac [N/A] . . . . . 2594  
 3.318.9 Mupad [N/A] . . . . . 2594

**3.318.1 Optimal result**

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c - a^2cx^2) \operatorname{arccosh}(ax)^2} dx = \frac{1}{ac\sqrt{-1 + ax}\sqrt{1 + ax}\operatorname{arccosh}(ax)} + \frac{a \operatorname{Int}\left(\frac{x}{(-1+ax)^{3/2}(1+ax)^{3/2}\operatorname{arccosh}(ax)}, x\right)}{c}$$

output `1/a/c/arccosh(a*x)/(a*x-1)^(1/2)/(a*x+1)^(1/2)+a*Unintegrable(x/(a*x-1)^(3/2)/(a*x+1)^(3/2)/arccosh(a*x), x)/c`

**3.318.2 Mathematica [N/A]**

Not integrable

Time = 2.90 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2cx^2) \operatorname{arccosh}(ax)^2} dx = \int \frac{1}{(c - a^2cx^2) \operatorname{arccosh}(ax)^2} dx$$

input `Integrate[1/((c - a^2*c*x^2)*ArcCosh[a*x]^2), x]`

output `Integrate[1/((c - a^2*c*x^2)*ArcCosh[a*x]^2), x]`



**3.318.3 Rubi [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 6376}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arccosh}(ax)^2 (c - a^2 cx^2)} dx$$

↓ 6319

$$\frac{a \int \frac{x}{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)} dx}{c} + \frac{1}{ac\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}$$

↓ 6376

$$\frac{a \int \frac{x}{(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)} dx}{c} + \frac{1}{ac\sqrt{ax-1}\sqrt{ax+1} \operatorname{arccosh}(ax)}$$

input `Int[1/((c - a^2*c*x^2)*ArcCosh[a*x]^2), x]`

output `$Aborted`

**3.318.3.1 Defintions of rubi rules used**

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6376 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.)), x_Symbol] :> Unintegrable[(f*x)^m*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n, p}, x]`

---

3.318.  $\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)^2} dx$

**3.318.4 Maple [N/A] (verified)**

Not integrable

Time = 0.66 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-a^2 c x^2 + c) \operatorname{arccosh}(ax)^2} dx$$

input `int(1/(-a^2*c*x^2+c)/arccosh(a*x)^2,x)`output `int(1/(-a^2*c*x^2+c)/arccosh(a*x)^2,x)`**3.318.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{1}{(c - a^2 c x^2) \operatorname{arccosh}(ax)^2} dx = \int -\frac{1}{(a^2 c x^2 - c) \operatorname{arccosh}(ax)^2} dx$$

input `integrate(1/(-a^2*c*x^2+c)/arccosh(a*x)^2,x, algorithm="fricas")`output `integral(-1/((a^2*c*x^2 - c)*arccosh(a*x)^2), x)`**3.318.6 Sympy [N/A]**

Not integrable

Time = 2.63 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.30

$$\int \frac{1}{(c - a^2 c x^2) \operatorname{arccosh}(ax)^2} dx = -\frac{\int \frac{1}{a^2 x^2 \operatorname{acosh}^2(ax) - \operatorname{acosh}^2(ax)} dx}{c}$$

input `integrate(1/(-a**2*c*x**2+c)/acosh(a*x)**2,x)`output `-Integral(1/(a**2*x**2*acosh(a*x)**2 - acosh(a*x)**2), x)/c`

---

3.318.  $\int \frac{1}{(c - a^2 c x^2) \operatorname{arccosh}(ax)^2} dx$

**3.318.7 Maxima [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 259, normalized size of antiderivative = 12.95

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)^2} dx = \int -\frac{1}{(a^2 cx^2 - c) \operatorname{arcosh}(ax)^2} dx$$

```
input integrate(1/(-a^2*c*x^2+c)/arccosh(a*x)^2,x, algorithm="maxima")
```

```
output (a*x + sqrt(a*x + 1)*sqrt(a*x - 1))/((a^3*c*x^2 + sqrt(a*x + 1)*sqrt(a*x - 1)*a^2*c*x - a*c)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))) + integrate((a^4*x^4 + (a^2*x^2 - 1)*(a*x + 1)*(a*x - 1) + (2*a^3*x^3 - a*x)*sqrt(a*x + 1)*sqrt(a*x - 1) - 1)/((a^6*c*x^6 - 3*a^4*c*x^4 + 3*a^2*c*x^2 + (a^4*c*x^4 - a^2*c*x^2)*(a*x + 1)*(a*x - 1) + 2*(a^5*c*x^5 - 2*a^3*c*x^3 + a*c*x)*sqrt(a*x + 1)*sqrt(a*x - 1) - c)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))), x)
```

**3.318.8 Giac [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)^2} dx = \int -\frac{1}{(a^2 cx^2 - c) \operatorname{arcosh}(ax)^2} dx$$

```
input integrate(1/(-a^2*c*x^2+c)/arccosh(a*x)^2,x, algorithm="giac")
```

```
output integrate(-1/((a^2*c*x^2 - c)*arccosh(a*x)^2), x)
```

**3.318.9 Mupad [N/A]**

Not integrable

Time = 2.70 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2 cx^2) \operatorname{arccosh}(ax)^2} dx = \int \frac{1}{\operatorname{acosh}(ax)^2 (c - a^2 cx^2)} dx$$

input `int(1/(acosh(a*x)^2*(c - a^2*c*x^2)),x)`

output `int(1/(acosh(a*x)^2*(c - a^2*c*x^2)), x)`

**3.319**  $\int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx$

3.319.1 Optimal result . . . . .	2596
3.319.2 Mathematica [N/A] . . . . .	2596
3.319.3 Rubi [N/A] . . . . .	2597
3.319.4 Maple [N/A] (verified) . . . . .	2598
3.319.5 Fricas [N/A] . . . . .	2598
3.319.6 Sympy [N/A] . . . . .	2598
3.319.7 Maxima [N/A] . . . . .	2599
3.319.8 Giac [N/A] . . . . .	2599
3.319.9 Mupad [N/A] . . . . .	2600

**3.319.1 Optimal result**

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx = -\frac{1}{ac^2(-1 + ax)^{3/2}(1 + ax)^{3/2}\operatorname{arccosh}(ax)} - \frac{3a\operatorname{Int}\left(\frac{x}{(-1+ax)^{5/2}(1+ax)^{5/2}\operatorname{arccosh}(ax)}, x\right)}{c^2}$$

output `-1/a/c^2/(a*x-1)^(3/2)/(a*x+1)^(3/2)/arccosh(a*x)-3*a*Unintegrable(x/(a*x-1)^(5/2)/(a*x+1)^(5/2)/arccosh(a*x),x)/c^2`

**3.319.2 Mathematica [N/A]**

Not integrable

Time = 8.90 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx = \int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx$$

input `Integrate[1/((c - a^2*c*x^2)^2*ArcCosh[a*x]^2),x]`

output `Integrate[1/((c - a^2*c*x^2)^2*ArcCosh[a*x]^2), x]`

**3.319.3 Rubi [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 6376}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arccosh}(ax)^2 (c - a^2 cx^2)^2} dx$$

↓ 6319

$$-\frac{3a \int \frac{x}{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)} dx}{c^2} - \frac{1}{ac^2(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}$$

↓ 6376

$$-\frac{3a \int \frac{x}{(ax-1)^{5/2}(ax+1)^{5/2} \operatorname{arccosh}(ax)} dx}{c^2} - \frac{1}{ac^2(ax-1)^{3/2}(ax+1)^{3/2} \operatorname{arccosh}(ax)}$$

input `Int[1/((c - a^2*c*x^2)^2*ArcCosh[a*x]^2), x]`

output `$Aborted`

**3.319.3.1 Defintions of rubi rules used**

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6376 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^ (n_.)*((f_.)*(x_)^m)*((d1_) + (e1_.)*(x_)^p)*((d2_) + (e2_.)*(x_)^p)*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n, p}, x]`

---

3.319.  $\int \frac{1}{(c - a^2 cx^2)^2 \operatorname{arccosh}(ax)^2} dx$

**3.319.4 Maple [N/A] (verified)**

Not integrable

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(-a^2cx^2 + c)^2 \operatorname{arccosh}(ax)^2} dx$$

input `int(1/(-a^2*c*x^2+c)^2/arccosh(a*x)^2,x)`output `int(1/(-a^2*c*x^2+c)^2/arccosh(a*x)^2,x)`**3.319.5 Fracas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx = \int \frac{1}{(a^2cx^2 - c)^2 \operatorname{arccosh}(ax)^2} dx$$

input `integrate(1/(-a^2*c*x^2+c)^2/arccosh(a*x)^2,x, algorithm="fricas")`output `integral(1/((a^4*c^2*x^4 - 2*a^2*c^2*x^2 + c^2)*arccosh(a*x)^2), x)`**3.319.6 Sympy [N/A]**

Not integrable

Time = 12.41 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.05

$$\int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx = \int \frac{1}{\frac{a^4x^4 \operatorname{acosh}^2(ax) - 2a^2x^2 \operatorname{acosh}^2(ax) + \operatorname{acosh}^2(ax)}{c^2}} dx$$

input `integrate(1/(-a**2*c*x**2+c)**2/acosh(a*x)**2,x)`output `Integral(1/(a**4*x**4*acosh(a*x)**2 - 2*a**2*x**2*acosh(a*x)**2 + acosh(a*x)**2), x)/c**2`

**3.319.7 Maxima [N/A]**

Not integrable

Time = 0.43 (sec) , antiderivative size = 351, normalized size of antiderivative = 17.55

$$\int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx = \int \frac{1}{(a^2cx^2 - c)^2 \operatorname{arccosh}(ax)^2} dx$$

```
input integrate(1/(-a^2*c*x^2+c)^2/arccosh(a*x)^2,x, algorithm="maxima")
```

```
output -(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))/((a^5*c^2*x^4 - 2*a^3*c^2*x^2 + a*c^2
+ (a^4*c^2*x^3 - a^2*c^2*x)*sqrt(a*x + 1)*sqrt(a*x - 1))*log(a*x + sqrt(a
*x + 1)*sqrt(a*x - 1))) - integrate((3*a^4*x^4 - 2*a^2*x^2 + (3*a^2*x^2 -
1)*(a*x + 1)*(a*x - 1) + 3*(2*a^3*x^3 - a*x)*sqrt(a*x + 1)*sqrt(a*x - 1) -
1)/((a^8*c^2*x^8 - 4*a^6*c^2*x^6 + 6*a^4*c^2*x^4 - 4*a^2*c^2*x^2 + (a^6*c
^2*x^6 - 2*a^4*c^2*x^4 + a^2*c^2*x^2)*(a*x + 1)*(a*x - 1) + 2*(a^7*c^2*x^7
- 3*a^5*c^2*x^5 + 3*a^3*c^2*x^3 - a*c^2*x)*sqrt(a*x + 1)*sqrt(a*x - 1) +
c^2)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1))), x)
```

**3.319.8 Giac [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{1}{(c - a^2cx^2)^2 \operatorname{arccosh}(ax)^2} dx = \int \frac{1}{(a^2cx^2 - c)^2 \operatorname{arccosh}(ax)^2} dx$$

```
input integrate(1/(-a^2*c*x^2+c)^2/arccosh(a*x)^2,x, algorithm="giac")
```

```
output integrate(1/((a^2*c*x^2 - c)^2*arccosh(a*x)^2), x)
```



**3.319.9 Mupad [N/A]**

Not integrable

Time = 2.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c - a^2 c x^2)^2 \operatorname{arccosh}(a x)^2} dx = \int \frac{1}{\operatorname{acosh}(a x)^2 (c - a^2 c x^2)^2} dx$$

input `int(1/(acosh(a*x)^2*(c - a^2*c*x^2)^2),x)`output `int(1/(acosh(a*x)^2*(c - a^2*c*x^2)^2), x)`

**3.320**      $\int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.320.1 Optimal result . . . . . 2601  
 3.320.2 Mathematica [A] (verified) . . . . . 2602  
 3.320.3 Rubi [A] (verified) . . . . . 2603  
 3.320.4 Maple [A] (verified) . . . . . 2605  
 3.320.5 Fricas [F] . . . . . 2606  
 3.320.6 Sympy [F] . . . . . 2606  
 3.320.7 Maxima [F] . . . . . 2607  
 3.320.8 Giac [F(-2)] . . . . . 2607  
 3.320.9 Mupad [F(-1)] . . . . . 2608

**3.320.1 Optimal result**

Integrand size = 28, antiderivative size = 350

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^3 \sqrt{-1+cx} \sqrt{1+cx} \sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} + \frac{\sqrt{1-cx} \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right)}{8b^2c^4\sqrt{-1+cx}} - \frac{3\sqrt{1-cx} \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{3a}{b}\right)}{16b^2c^4\sqrt{-1+cx}} - \frac{5\sqrt{1-cx} \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{5a}{b}\right)}{16b^2c^4\sqrt{-1+cx}} - \frac{\sqrt{1-cx} \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b^2c^4\sqrt{-1+cx}} + \frac{3\sqrt{1-cx} \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^4\sqrt{-1+cx}} + \frac{5\sqrt{1-cx} \cosh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^4\sqrt{-1+cx}}$$

output 
$$\begin{aligned} & -1/8*\cosh(a/b)*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^4/(c*x-1)^{(1/2)} \\ & +3/16*\cosh(3*a/b)*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^4/(c*x-1)^{(1/2)} \\ & +5/16*\cosh(5*a/b)*\text{Shi}(5*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^4/(c*x-1)^{(1/2)} \\ & +1/8*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\sinh(a/b)*(-c*x+1)^{(1/2)}/b^2/c^4/(c*x-1)^{(1/2)} \\ & -3/16*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\sinh(3*a/b)*(-c*x+1)^{(1/2)}/b^2/c^4/(c*x-1)^{(1/2)} \\ & -5/16*\text{Chi}(5*(a+b*\text{arccosh}(c*x))/b)*\sinh(5*a/b)*(-c*x+1)^{(1/2)}/b^2/c^4/(c*x-1)^{(1/2)} \\ & -x^3*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(-c^2*x^2+1)^{(1/2)}/b/c/(a+b*\text{arccosh}(c*x)) \end{aligned}$$

### 3.320.2 Mathematica [A] (verified)

Time = 0.56 (sec) , antiderivative size = 322, normalized size of antiderivative = 0.92

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\text{arccosh}(cx))^2} dx = \frac{\sqrt{1-c^2x^2}(16bc^3x^3 - 16bc^5x^5 + 2(a+b\text{arccosh}(cx))\text{Chi}(\frac{a}{b} + \text{arccosh}(cx))\sinh(\frac{a}{b}) - 3(a+b\text{arccosh}(cx)))}{(a+b\text{arccosh}(cx))^2}$$

input `Integrate[(x^3*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

output 
$$\begin{aligned} & (\text{sqrt}[1 - c^2*x^2]*(16*b*c^3*x^3 - 16*b*c^5*x^5 + 2*(a + b*\text{ArcCosh}[c*x])* \\ & \text{CoshIntegral}[a/b + \text{ArcCosh}[c*x]]*\text{Sinh}[a/b] - 3*(a + b*\text{ArcCosh}[c*x])* \\ & \text{CoshIntegral}[3*(a/b + \text{ArcCosh}[c*x])* \\ & \text{Sinh}[(3*a)/b] - 5*a*\text{CoshIntegral}[5*(a/b + \text{ArcCosh}[c*x])* \\ & \text{Sinh}[(5*a)/b] - 5*b*\text{ArcCosh}[c*x]*\text{CoshIntegral}[5*(a/b + \text{ArcCosh}[c*x])* \\ & \text{Sinh}[(5*a)/b] - 2*a*\text{Cosh}[a/b]*\text{SinhIntegral}[a/b + \text{ArcCosh}[c*x]] - \\ & 2*b*\text{ArcCosh}[c*x]*\text{Cosh}[a/b]*\text{SinhIntegral}[a/b + \text{ArcCosh}[c*x]] + 3*a*\text{Cosh}[(3*a)/b] \\ & *\text{SinhIntegral}[3*(a/b + \text{ArcCosh}[c*x])] + 3*b*\text{ArcCosh}[c*x]*\text{Cosh}[(3*a)/b] \\ & *\text{SinhIntegral}[3*(a/b + \text{ArcCosh}[c*x])] + 5*a*\text{Cosh}[(5*a)/b]*\text{SinhIntegral}[5*(a/b + \text{ArcCosh}[c*x])] \\ & + 5*b*\text{ArcCosh}[c*x]*\text{Cosh}[(5*a)/b]*\text{SinhIntegral}[5*(a/b + \text{ArcCosh}[c*x])]) \\ & )/(16*b^2*c^4*\text{sqrt}[-1 + c*x]*\text{sqrt}[1 + c*x]*(a + b*\text{ArcCosh}[c*x])) \end{aligned}$$

**3.320.3 Rubi [A] (verified)**

Time = 1.05 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.01, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$ , Rules used = {6357, 6302, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3 \sqrt{1-c^2x^2}}{(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{5c\sqrt{1-cx} \int \frac{x^4}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{3\sqrt{1-cx} \int \frac{x^2}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} - \frac{x^3\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6302} \\
 & \frac{5\sqrt{1-cx} \int -\frac{\cosh^4\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^4\sqrt{cx-1}} - \\
 & \frac{3\sqrt{1-cx} \int -\frac{\cosh^2\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^4\sqrt{cx-1}} - \\
 & \frac{x^3\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & -\frac{5\sqrt{1-cx} \int \frac{\cosh^4\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^4\sqrt{cx-1}} + \\
 & \frac{3\sqrt{1-cx} \int \frac{\cosh^2\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^4\sqrt{cx-1}} - \\
 & \frac{x^3\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{5971}
 \end{aligned}$$

$$\begin{aligned}
 & \frac{5\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} + \frac{3\sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{3\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))} \\
 & \frac{b^2 c^4 \sqrt{cx-1}}{x^3 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}} \\
 & \frac{b^2 c^4 \sqrt{cx-1}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{2009} \\
 & \frac{3\sqrt{1-cx} \left( -\frac{1}{4} \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4} \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{5\sqrt{1-cx} \left( -\frac{1}{8} \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{3}{16} \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{16} \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)} \\
 & \frac{b^2 c^4 \sqrt{cx-1}}{x^3 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}} \\
 & \frac{b^2 c^4 \sqrt{cx-1}}{bc(a+b\operatorname{arccosh}(cx))}
 \end{aligned}$$

input `Int[(x^3*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

output `-(x^3*sqrt[-1 + c*x]*sqrt[1 + c*x]*sqrt[1 - c^2*x^2])/(b*c*(a + b*ArcCosh[c*x])) - (3*sqrt[1 - c*x]*(-1/4*(CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b]) - (CoshIntegral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*a)/b])/4 + (Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/4 + (Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/4))/(b^2*c^4*sqrt[-1 + c*x]) + (5*sqrt[1 - c*x]*(-1/8*(CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b]) - (3*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*a)/b])/16 - (CoshIntegral[(5*(a + b*ArcCosh[c*x])/b]*Sinh[(5*a)/b])/16 + (Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/8 + (3*Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/16 + (Cosh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/16))/(b^2*c^4*sqrt[-1 + c*x])`

3.320.  $\int \frac{x^3 \sqrt{1-c^2 x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

## 3.320.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

## 3.320.4 Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 555, normalized size of antiderivative = 1.59

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(32\sqrt{cx-1}\sqrt{cx+1}bc^5x^5+32bc^6x^6-32\sqrt{cx-1}\sqrt{cx+1}bc^3x^3-32bc^4x^4+5\operatorname{arccosh}(cx)\right)}{\dots}$

input `int(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

$$3.320. \quad \int \frac{x^3\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

output `1/32*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^5*x^5+32*b*c^6*x^6-32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^3*x^3-32*b*c^4*x^4+5*arccosh(c*x)*b*Ei(1,-5*arccosh(c*x)-5*a/b)*exp(-(-b*arccosh(c*x)+5*a)/b)+3*arccosh(c*x)*b*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(-b*arccosh(c*x)+3*a)/b)-2*arccosh(c*x)*b*Ei(1,-arccosh(c*x)-a/b)*exp(-(-b*arccosh(c*x)+a)/b)-5*Ei(1,5*arccosh(c*x)+5*a/b)*exp((b*arccosh(c*x)+5*a)/b)*b*arccosh(c*x)-3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)*b*arccosh(c*x)+2*Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)*b*arccosh(c*x)+5*a*Ei(1,-5*arccosh(c*x)-5*a/b)*exp(-(-b*arccosh(c*x)+5*a)/b)+3*a*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(-b*arccosh(c*x)+3*a)/b)-2*a*Ei(1,-arccosh(c*x)-a/b)*exp(-(-b*arccosh(c*x)+a)/b)-5*Ei(1,5*arccosh(c*x)+5*a/b)*exp((b*arccosh(c*x)+5*a)/b)*a-3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)*a+2*Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)*a)/(c*x+1)/c^4/(c*x-1)/b^2/(a+b*arccosh(c*x))`

### 3.320.5 Fricas [F]

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}x^3}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^3/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

### 3.320.6 Sympy [F]

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^3 \sqrt{-(cx-1)(cx+1)}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x**3*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(x**3*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x))**2, x)`

---

3.320.  $\int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

**3.320.7 Maxima [F]**

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2 + 1}x^3}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^2*x^5 - x^3)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^6 - c*x^4)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((5*c^3*x^5 - 2*c*x^3)*(c*x + 1)^(3/2)*(c*x - 1) + (10*c^4*x^6 - 11*c^2*x^4 + 3*x^2)*(c*x + 1)*sqrt(c*x - 1) + (5*c^5*x^7 - 9*c^3*x^5 + 4*c*x^3)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.320.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{(a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`



**3.320.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^3 \sqrt{1-c^2x^2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((x^3*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2,x)`output `int((x^3*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2, x)`

**3.321**  $\int \frac{x^2\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.321.1 Optimal result . . . . . 2609  
 3.321.2 Mathematica [A] (verified) . . . . . 2609  
 3.321.3 Rubi [C] (verified) . . . . . 2610  
 3.321.4 Maple [B] (verified) . . . . . 2615  
 3.321.5 Fricas [F] . . . . . 2616  
 3.321.6 Sympy [F] . . . . . 2616  
 3.321.7 Maxima [F] . . . . . 2617  
 3.321.8 Giac [F] . . . . . 2617  
 3.321.9 Mupad [F(-1)] . . . . . 2618

**3.321.1 Optimal result**

Integrand size = 28, antiderivative size = 154

$$\int \frac{x^2\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^2\sqrt{-1+cx}\sqrt{1+cx}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{4a}{b}\right)}{2b^2c^3\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\cosh\left(\frac{4a}{b}\right)\operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{2b^2c^3\sqrt{-1+cx}}$$

```
output 1/2*cosh(4*a/b)*Shi(4*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c^3/(c*x-1)
^(1/2)-1/2*Chi(4*(a+b*arccosh(c*x))/b)*sinh(4*a/b)*(-c*x+1)^(1/2)/b^2/c^3/
(c*x-1)^(1/2)-x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/b/c/(a+b*
arccosh(c*x))
```

**3.321.2 Mathematica [A] (verified)**

Time = 0.44 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.84

$$\int \frac{x^2\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \frac{\sqrt{1-c^2x^2}(-2bc^2x^2(-1+c^2x^2)-(a+b\operatorname{arccosh}(cx))\operatorname{Chi}\left(4\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right)\right)\sinh\left(\frac{4a}{b}\right)+(a+b\operatorname{arccosh}(cx))\operatorname{Shi}\left(4\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right)\right)\cosh\left(\frac{4a}{b}\right))}{2b^2c^3\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}$$

input `Integrate[(x^2*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

output `(Sqrt[1 - c^2*x^2]*(-2*b*c^2*x^2*(-1 + c^2*x^2) - (a + b*ArcCosh[c*x])*CoshIntegral[4*(a/b + ArcCosh[c*x])*Sinh[(4*a)/b] + (a + b*ArcCosh[c*x])*Cosh[(4*a)/b]*SinhIntegral[4*(a/b + ArcCosh[c*x])]))/(2*b^2*c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))`

### 3.321.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.27 (sec) , antiderivative size = 262, normalized size of antiderivative = 1.70, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6357, 6302, 25, 5971, 27, 2009, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2 \sqrt{1 - c^2 x^2}}{(a + \operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{4c\sqrt{1 - cx} \int \frac{x^3}{a + \operatorname{barccosh}(cx)} dx}{b\sqrt{cx - 1}} - \frac{2\sqrt{1 - cx} \int \frac{x}{a + \operatorname{barccosh}(cx)} dx}{bc\sqrt{cx - 1}} - \frac{x^2 \sqrt{cx - 1} \sqrt{cx + 1} \sqrt{1 - c^2 x^2}}{bc(a + \operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6302} \\
 & \frac{4\sqrt{1 - cx} \int -\frac{\cosh^3\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{b^2 c^3 \sqrt{cx - 1}} - \\
 & \frac{2\sqrt{1 - cx} \int -\frac{\cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{a + \operatorname{barccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{b^2 c^3 \sqrt{cx - 1}} - \\
 & \frac{x^2 \sqrt{cx - 1} \sqrt{cx + 1} \sqrt{1 - c^2 x^2}}{bc(a + \operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

---

3.321.  $\int \frac{x^2 \sqrt{1 - c^2 x^2}}{(a + \operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{4\sqrt{1-cx} \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} + \\
 & \frac{2\sqrt{1-cx} \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} - \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{5971} \\
 & \frac{2\sqrt{1-cx} \int \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2(a+b\operatorname{arccosh}(cx))} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} - \\
 & \frac{4\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} - \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{27} \\
 & \frac{\sqrt{1-cx} \int \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} - \\
 & \frac{4\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} - \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{1-cx} \int \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} + \\
 & \frac{4\sqrt{1-cx} \left( -\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.321.  $\int \frac{x^2 \sqrt{1-c^2 x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{\sqrt{1-cx} \int -\frac{i \sin\left(\frac{2ia}{b} - \frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} + \\
 & \frac{4\sqrt{1-cx} \left(-\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{26} \\
 & -\frac{i\sqrt{1-cx} \int \frac{\sin\left(\frac{2ia}{b} - \frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} + \\
 & \frac{4\sqrt{1-cx} \left(-\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{3784} \\
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) + \cosh\left(\frac{2a}{b}\right) \int -\frac{i \sinh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{4\sqrt{1-cx} \left(-\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{26} \\
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \int \frac{\sinh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{4\sqrt{1-cx} \left(-\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.321.  $\int \frac{x^2 \sqrt{1-c^2 x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \int \frac{i \sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & - \frac{4\sqrt{1-cx} \left( -\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \quad 26 \\
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & - \frac{4\sqrt{1-cx} \left( -\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \quad 3779 \\
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} + \\
 & - \frac{4\sqrt{1-cx} \left( -\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \quad 3782 \\
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} + \\
 & - \frac{4\sqrt{1-cx} \left( -\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{1-c^2 x^2}}{bc(a+b\operatorname{arccosh}(cx))}
 \end{aligned}$$

input `Int[(x^2*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

3.321.  $\int \frac{x^2 \sqrt{1-c^2 x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

```
output 
$$-\left(\frac{x^2 \sqrt{-1 + cx} \sqrt{1 + cx} \sqrt{1 - c^2 x^2}}{b^2 c^3 \sqrt{-1 + cx}} - \frac{I \sqrt{1 - cx} (I \operatorname{CoshIntegral}[(2(a + b \operatorname{ArcCosh}[cx]))/b] \operatorname{Sinh}[(2a)/b] - I \operatorname{Cosh}[(2a)/b] \operatorname{SinhIntegral}[(2(a + b \operatorname{ArcCosh}[cx]))/b])}{b^2 c^3 \sqrt{-1 + cx}} + \frac{4 \sqrt{1 - cx} (-1/4 (\operatorname{CoshIntegral}[(2(a + b \operatorname{ArcCosh}[cx]))/b] \operatorname{Sinh}[(2a)/b] - (\operatorname{CoshIntegral}[(4(a + b \operatorname{ArcCosh}[cx]))/b] \operatorname{Sinh}[(4a)/b])/8 + (\operatorname{Cosh}[(2a)/b] \operatorname{SinhIntegral}[(2(a + b \operatorname{ArcCosh}[cx]))/b])/4 + (\operatorname{Cosh}[(4a)/b] \operatorname{SinhIntegral}[(4(a + b \operatorname{ArcCosh}[cx]))/b])/8)}{b^2 c^3 \sqrt{-1 + cx}}\right)$$

```

### 3.321.3.1 Defintions of rubi rules used

```
rule 25 
$$\operatorname{Int}[-(F_x), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$$

```

```
rule 26 
$$\operatorname{Int}[(\operatorname{Complex}[0, a]) (F_x), x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Complex}[\operatorname{Identity}[0], a]) \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ \operatorname{EqQ}[a^2, 1]$$

```

```
rule 27 
$$\operatorname{Int}[(a) (F_x), x\_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[F_x, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ \operatorname{MatchQ}[F_x, (b) (G_x)] /; \operatorname{FreeQ}[b, x]$$

```

```
rule 2009 
$$\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] /; \operatorname{SumQ}[u]$$

```

```
rule 3042 
$$\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] /; \operatorname{FunctionOfTrigOfLinearQ}[u, x]$$

```

```
rule 3779 
$$\operatorname{Int}[\sin[(e) + (\operatorname{Complex}[0, fz]) (f) (x)] / ((c) + (d) (x)), x\_Symbol] \rightarrow \operatorname{Simp}[I (\operatorname{SinhIntegral}[c f (fz/d) + f fz x / d], x) /; \operatorname{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \operatorname{EqQ}[d e - c f fz I, 0]$$

```

```
rule 3782 
$$\operatorname{Int}[\sin[(e) + (\operatorname{Complex}[0, fz]) (f) (x)] / ((c) + (d) (x)), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{CoshIntegral}[c f (fz/d) + f fz x / d], x] /; \operatorname{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \operatorname{EqQ}[d (e - \pi/2) - c f fz I, 0]$$

```

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

### 3.321.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 278 vs.  $2(136) = 272$ .

Time = 0.58 (sec) , antiderivative size = 279, normalized size of antiderivative = 1.81

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(4x^4c^4b\sqrt{cx-1}\sqrt{cx+1}+4bc^5x^5-4\sqrt{cx-1}\sqrt{cx+1}bc^2x^2-4bc^3x^3+\operatorname{arccosh}(cx)b\operatorname{Ei}_1(-\sqrt{-c^2x^2+1})\right)}{(a+b\operatorname{arccosh}(cx))^2}$

input `int(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

$$3.321. \quad \int \frac{x^2\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$



output  $\frac{1}{4}(-c^2x^2+1)^{1/2}(-cx-1)^{1/2}(cx+1)^{1/2}cx+c^2x^2-1)(4x^4*c^4*b*(cx-1)^{1/2}(cx+1)^{1/2}+4*b*c^5*x^5-4*(cx-1)^{1/2}(cx+1)^{1/2})*b*c^2*x^2-4*b*c^3*x^3+\operatorname{arccosh}(cx)*b*\operatorname{Ei}(1,-4*\operatorname{arccosh}(cx)-4*a/b)*\exp(-(-b*\operatorname{arccosh}(cx)+4*a)/b)-\operatorname{Ei}(1,4*\operatorname{arccosh}(cx)+4*a/b)*\exp((b*\operatorname{arccosh}(cx)+4*a)/b)*b*\operatorname{arccosh}(cx)+a*\operatorname{Ei}(1,-4*\operatorname{arccosh}(cx)-4*a/b)*\exp(-(-b*\operatorname{arccosh}(cx)+4*a)/b)-\operatorname{Ei}(1,4*\operatorname{arccosh}(cx)+4*a/b)*\exp((b*\operatorname{arccosh}(cx)+4*a)/b)*a)/(cx+1)/c^3/(cx-1)/b^2/(a+b*\operatorname{arccosh}(cx))$

### 3.321.5 Fracas [F]

$$\int \frac{x^2\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}x^2}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^2/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

### 3.321.6 Sympy [F]

$$\int \frac{x^2\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2\sqrt{-(cx-1)(cx+1)}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x**2*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(x**2*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x))**2, x)`

**3.321.7 Maxima [F]**

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^2}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^2*x^4 - x^2)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^5 - c*x^3)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((4*c^3*x^4 - c*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(4*c^4*x^5 - 4*c^2*x^3 + x)*(c*x + 1)*sqrt(c*x - 1) + (4*c^5*x^6 - 7*c^3*x^4 + 3*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.321.8 Giac [F]**

$$\int \frac{x^2 \sqrt{1 - c^2 x^2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2 x^2 + 1} x^2}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)*x^2/(b*arccosh(c*x) + a)^2, x)`

**3.321.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2 \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2 \sqrt{1-c^2x^2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((x^2*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2,x)`output `int((x^2*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2, x)`

**3.322**      $\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.322.1 Optimal result . . . . .	2619
3.322.2 Mathematica [A] (verified) . . . . .	2620
3.322.3 Rubi [C] (verified) . . . . .	2620
3.322.4 Maple [A] (verified) . . . . .	2625
3.322.5 Fracas [F] . . . . .	2626
3.322.6 Sympy [F] . . . . .	2626
3.322.7 Maxima [F] . . . . .	2627
3.322.8 Giac [F] . . . . .	2627
3.322.9 Mupad [F(-1)] . . . . .	2627

**3.322.1 Optimal result**

Integrand size = 26, antiderivative size = 248

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x\sqrt{-1+cx}\sqrt{1+cx}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} + \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{4b^2c^2\sqrt{-1+cx}} - \frac{3\sqrt{1-cx}\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{4b^2c^2\sqrt{-1+cx}} - \frac{\sqrt{1-cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4b^2c^2\sqrt{-1+cx}} + \frac{3\sqrt{1-cx}\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b^2c^2\sqrt{-1+cx}}$$

```
output -1/4*cosh(a/b)*Shi((a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)+3/4*cosh(3*a/b)*Shi(3*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)+1/4*Chi((a+b*arccosh(c*x))/b)*sinh(a/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)-3/4*Chi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)-x*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/b/c/(a+b*arccosh(c*x))
```

**3.322.2 Mathematica [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.88

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

$$= \frac{\sqrt{1-c^2x^2}(4bcx - 4bc^3x^3 + (a+b\operatorname{arccosh}(cx))\operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \sinh\left(\frac{a}{b}\right) - 3(a+b\operatorname{arccosh}(cx))\operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \cosh\left(\frac{a}{b}\right))}{(a+b\operatorname{arccosh}(cx))^2}$$

input `Integrate[(x*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

output `(Sqrt[1 - c^2*x^2]*(4*b*c*x - 4*b*c^3*x^3 + (a + b*ArcCosh[c*x])*CoshIntegral[a/b + ArcCosh[c*x]]*Sinh[a/b] - 3*(a + b*ArcCosh[c*x])*CoshIntegral[3*(a/b + ArcCosh[c*x])*Sinh[(3*a)/b] - a*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] - b*ArcCosh[c*x]*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + 3*a*Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] + 3*b*ArcCosh[c*x]*Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])]))/(4*b^2*c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))`

**3.322.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 1.30 (sec) , antiderivative size = 252, normalized size of antiderivative = 1.02, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.577$ , Rules used = {6357, 6296, 25, 3042, 26, 3784, 26, 3042, 26, 3779, 3782, 6302, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

$$\downarrow \text{6357}$$

$$\frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{1-cx} \int \frac{1}{a+b\operatorname{arccosh}(cx)} dx}{bc\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}$$

$$\downarrow \text{6296}$$

$$\begin{aligned}
 & \frac{\sqrt{1-cx} \int -\frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 25 \\
 & \frac{\sqrt{1-cx} \int \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 3042 \\
 & \frac{\sqrt{1-cx} \int -\frac{i \sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 26 \\
 & \frac{i\sqrt{1-cx} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 3784 \\
 & i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) + \cosh\left(\frac{a}{b}\right) \int -\frac{i \sinh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right) \\
 & \quad \frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{b^2c^2\sqrt{cx-1}}{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}} \frac{1}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 26
 \end{aligned}$$

3.322.  $\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)$$

$$\frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{b^2c^2\sqrt{cx-1}}{bc(a+b\operatorname{arccosh}(cx))} \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}$$

↓ 3042

$$i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \int \frac{i \sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)$$

$$\frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{b^2c^2\sqrt{cx-1}}{bc(a+b\operatorname{arccosh}(cx))} \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}$$

↓ 26

$$i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)$$

$$\frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{b^2c^2\sqrt{cx-1}}{bc(a+b\operatorname{arccosh}(cx))} \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}$$

↓ 3779

$$i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right) +$$

$$\frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{b^2c^2\sqrt{cx-1}}{bc(a+b\operatorname{arccosh}(cx))} \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}$$

↓ 3782

$$\frac{3c\sqrt{1-cx} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2c^2\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}$$

---

3.322.  $\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
& \downarrow \text{6302} \\
& \frac{3\sqrt{1-cx} \int -\frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} \\
& \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
& \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{25} \\
& \frac{3\sqrt{1-cx} \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} \\
& \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
& \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{5971} \\
& \frac{3\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} \\
& \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
& \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{2009} \\
& \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} + \\
& \frac{3\sqrt{1-cx} \left( -\frac{1}{4} \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4} \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
& \frac{x\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}
\end{aligned}$$

input `Int[(x*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

$$3.322. \quad \int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$



```
output 
$$-\left(\frac{x\sqrt{-1+cx}\sqrt{1+cx}\sqrt{1-c^2x^2}}{b^2c^2} - \frac{I\sqrt{1-cx}\left(I\operatorname{CoshIntegral}\left[\frac{a+b\operatorname{ArcCosh}[cx]}{b}\right]\operatorname{Sinh}\left[\frac{a}{b}\right] - I\operatorname{Cosh}\left[\frac{a}{b}\right]\operatorname{SinhIntegral}\left[\frac{a+b\operatorname{ArcCosh}[cx]}{b}\right]\right)}{b^2c^2\sqrt{-1+cx}} + \frac{3\sqrt{1-cx}\left(-\frac{1}{4}\left(\operatorname{CoshIntegral}\left[\frac{a+b\operatorname{ArcCosh}[cx]}{b}\right]\operatorname{Sinh}\left[\frac{a}{b}\right] - \operatorname{CoshIntegral}\left[\frac{3(a+b\operatorname{ArcCosh}[cx])}{b}\right]\operatorname{Sinh}\left[\frac{3a}{b}\right]\right)}{4} + \frac{\operatorname{Cosh}\left[\frac{a}{b}\right]\operatorname{SinhIntegral}\left[\frac{a+b\operatorname{ArcCosh}[cx]}{b}\right]}{4} + \frac{\operatorname{Cosh}\left[\frac{3a}{b}\right]\operatorname{SinhIntegral}\left[\frac{3(a+b\operatorname{ArcCosh}[cx])}{b}\right]}{4}\right)}{b^2c^2\sqrt{-1+cx}}\right)$$

```

### 3.322.3.1 Defintions of rubi rules used

```
rule 25 
$$\operatorname{Int}[-(F_x), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[F_x, x], x]$$

```

```
rule 26 
$$\operatorname{Int}[(\operatorname{Complex}[0, a])(F_x), x\_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Complex}[\operatorname{Identity}[0], a]) \operatorname{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \operatorname{EqQ}[a^2, 1]$$

```

```
rule 2009 
$$\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{IntSum}[u, x], x] \text{ ; SumQ}[u]$$

```

```
rule 3042 
$$\operatorname{Int}[u, x\_Symbol] \rightarrow \operatorname{Int}[\operatorname{DeactivateTrig}[u, x], x] \text{ ; FunctionOfTrigOfLinearQ}[u, x]$$

```

```
rule 3779 
$$\operatorname{Int}[\sin[(e.) + (\operatorname{Complex}[0, fz])(f.)*(x.)]/((c.) + (d.)*(x.)), x\_Symbol] \rightarrow \operatorname{Simp}[I*(\operatorname{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] \text{ ; FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \operatorname{EqQ}[d*e - c*f*fz*I, 0]$$

```

```
rule 3782 
$$\operatorname{Int}[\sin[(e.) + (\operatorname{Complex}[0, fz])(f.)*(x.)]/((c.) + (d.)*(x.)), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] \text{ ; FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \operatorname{EqQ}[d*(e - \operatorname{Pi}/2) - c*f*fz*I, 0]$$

```

```
rule 3784 
$$\operatorname{Int}[\sin[(e.) + (f.)*(x.)]/((c.) + (d.)*(x.)), x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[(d*e - c*f)/d] \operatorname{Int}[\operatorname{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \operatorname{Simp}[\operatorname{Sin}[(d*e - c*f)/d] \operatorname{Int}[\operatorname{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] \text{ ; FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \operatorname{NeQ}[d*e - c*f, 0]$$

```

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6296 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[1/(b*c) Subst[Int[x^n*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n*(x_)^(m_.), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*(m + 2*p + 1)/(b*f*(n + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

### 3.322.4 Maple [A] (verified)

Time = 0.64 (sec) , antiderivative size = 405, normalized size of antiderivative = 1.63

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-8\sqrt{cx-1}\sqrt{cx+1}bc^3x^3-8bc^4x^4+8\sqrt{cx-1}\sqrt{cx+1}bcx+8bc^2x^2+\operatorname{arccosh}(cx)\right)}{b^2E_1}$

input `int(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

$$3.322. \quad \int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

output `-1/8*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^3*x^3-8*b*c^4*x^4+8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c*x+8*b*c^2*x^2+arccosh(c*x)*b*Ei(1,-arccosh(c*x)-a/b)*exp(-(b*arccosh(c*x)+a)/b)-3*arccosh(c*x)*b*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(b*arccosh(c*x)+3*a)/b)+3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)*b*arccosh(c*x)-Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)*b*arccosh(c*x)+a*Ei(1,-arccosh(c*x)-a/b)*exp(-(b*arccosh(c*x)+a)/b)-3*a*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(b*arccosh(c*x)+3*a)/b)+3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)*a-Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)*a)/(c*x+1)/c^2/(c*x-1)/b^2/(a+b*arccosh(c*x))`

### 3.322.5 Fricas [F]

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}x}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

### 3.322.6 Sympy [F]

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x\sqrt{-(cx-1)(cx+1)}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(x*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x))**2, x)`

**3.322.7 Maxima [F]**

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}x}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-(c^2*x^3 - x)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^4 - c*x^2)*sqrt(c*x + 1) *sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b *c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate((3*(c*x + 1)^(3/2)*(c*x - 1)* c^3*x^3 + (6*c^4*x^4 - 5*c^2*x^2 + 1)*(c*x + 1)*sqrt(c*x - 1) + (3*c^5*x^5 - 5*c^3*x^3 + 2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.322.8 Giac [F]**

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}x}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)*x/(b*arccosh(c*x) + a)^2, x)`

**3.322.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x\sqrt{1-c^2x^2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((x*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2,x)`

output `int((x*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2, x)`

---

3.322.  $\int \frac{x\sqrt{1-c^2x^2}}{(a+b\operatorname{barccosh}(cx))^2} dx$

### 3.323 $\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.323.1 Optimal result . . . . .	2628
3.323.2 Mathematica [A] (verified) . . . . .	2628
3.323.3 Rubi [C] (verified) . . . . .	2629
3.323.4 Maple [B] (verified) . . . . .	2633
3.323.5 Fricas [F] . . . . .	2633
3.323.6 Sympy [F] . . . . .	2634
3.323.7 Maxima [F] . . . . .	2634
3.323.8 Giac [F] . . . . .	2634
3.323.9 Mupad [F(-1)] . . . . .	2635

#### 3.323.1 Optimal result

Integrand size = 25, antiderivative size = 146

$$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)}{b^2c\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\cosh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{b^2c\sqrt{-1+cx}}$$

output `cosh(2*a/b)*Shi(2*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c/(c*x-1)^(1/2) -Chi(2*(a+b*arccosh(c*x))/b)*sinh(2*a/b)*(-c*x+1)^(1/2)/b^2/c/(c*x-1)^(1/2) -(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/b/c/(a+b*arccosh(c*x))`

#### 3.323.2 Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \frac{\sqrt{1-c^2x^2}(b(-1+c^2x^2)+(a+b\operatorname{arccosh}(cx))\operatorname{Chi}(2(\frac{a}{b}+\operatorname{arccosh}(cx))))\sinh(\frac{2a}{b})-(a+b\operatorname{arccosh}(cx))}{b^2c\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}$$

input `Integrate[Sqrt[1 - c^2*x^2]/(a + b*ArcCosh[c*x])^2,x]`

output `-((Sqrt[1 - c^2*x^2]*(b*(-1 + c^2*x^2) + (a + b*ArcCosh[c*x])*CoshIntegral[2*(a/b + ArcCosh[c*x]])*Sinh[(2*a)/b] - (a + b*ArcCosh[c*x])*Cosh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[c*x])]))/(b^2*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))`

### 3.323.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.75 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.90, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.520$ , Rules used = {6319, 6302, 25, 5971, 27, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{1 - c^2 x^2}}{(a + b \operatorname{arccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{2c\sqrt{1 - cx} \int \frac{x}{a + b \operatorname{arccosh}(cx)} dx}{b\sqrt{cx - 1}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{1 - c^2 x^2}}{bc(a + b \operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{6302} \\
 & \frac{2\sqrt{1 - cx} \int -\frac{\cosh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{a + b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{\frac{b^2 c \sqrt{cx - 1}}{\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{1 - c^2 x^2}} - \frac{bc(a + b \operatorname{arccosh}(cx))}{bc(a + b \operatorname{arccosh}(cx))}} \\
 & \quad \downarrow \text{25} \\
 & \frac{2\sqrt{1 - cx} \int \frac{\cosh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{a + b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{\frac{b^2 c \sqrt{cx - 1}}{\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{1 - c^2 x^2}} - \frac{bc(a + b \operatorname{arccosh}(cx))}{bc(a + b \operatorname{arccosh}(cx))}} \\
 & \quad \downarrow \text{5971}
 \end{aligned}$$

---

3.323.  $\int \frac{\sqrt{1 - c^2 x^2}}{(a + b \operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{2\sqrt{1-cx} \int \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2(a+b\operatorname{arccosh}(cx))} d(a+b\operatorname{arccosh}(cx))}{b^2c\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 27 \\
 & \frac{\sqrt{1-cx} \int \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 3042 \\
 & \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{1-cx} \int -\frac{i \sin\left(\frac{2ia}{b} - \frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c\sqrt{cx-1}} \\
 & \quad \downarrow 26 \\
 & -\frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} + \frac{i\sqrt{1-cx} \int \frac{\sin\left(\frac{2ia}{b} - \frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c\sqrt{cx-1}} \\
 & \quad \downarrow 3784 \\
 & \frac{-\frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))} + i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) + \cosh\left(\frac{2a}{b}\right) \int -\frac{i \sinh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{b^2c\sqrt{cx-1}} \\
 & \quad \downarrow 26 \\
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \int \frac{\sinh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right) - \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}}{b^2c\sqrt{cx-1}} \\
 & \quad \downarrow 3042 \\
 & \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx)) + \frac{\pi}{2}}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \int -\frac{i \sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right) - \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+b\operatorname{arccosh}(cx))}}{b^2c\sqrt{cx-1}}
 \end{aligned}$$

---

3.323.  $\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
& \downarrow 26 \\
& -\frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+\operatorname{barccosh}(cx))} + \\
& \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) - \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+\operatorname{barccosh}(cx))}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) \right)}{b^2c\sqrt{cx-1}} \\
& \downarrow 3779 \\
& -\frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+\operatorname{barccosh}(cx))} + \\
& \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) \right)}{b^2c\sqrt{cx-1}} \\
& \downarrow 3782 \\
& -\frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bc(a+\operatorname{barccosh}(cx))} + \\
& \frac{i\sqrt{1-cx} \left( i \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) - i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) \right)}{b^2c\sqrt{cx-1}}
\end{aligned}$$

input `Int[Sqrt[1 - c^2*x^2]/(a + b*ArcCosh[c*x])^2,x]`

output `-((Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Sqrt[1 - c^2*x^2])/(b*c*(a + b*ArcCosh[c*x]))) + (I*Sqrt[1 - c*x]*(I*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]*Sinh[(2*a)/b] - I*Cosh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b]))/(b^2*c*Sqrt[-1 + c*x])`

### 3.323.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`



- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3779 `Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`
- rule 3782 `Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`
- rule 3784 `Int[sin[(e_) + (f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`
- rule 5971 `Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`
- rule 6302 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)^^(m_), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`
- rule 6319 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

**3.323.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 268 vs.  $2(132) = 264$ .

Time = 0.76 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.84

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(2\sqrt{cx-1}\sqrt{cx+1}bc^2x^2+2bc^3x^3+\operatorname{arccosh}(cx)b\operatorname{Ei}_1\left(-2\operatorname{arccosh}(cx)-\frac{2a}{b}\right)e^{-\frac{-b\operatorname{arccosh}(cx)+2a}{b}}\right)}{\dots}$

input `int((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output `1/2*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^2*x^2+2*b*c^3*x^3+arccosh(c*x)*b*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*b*arccosh(c*x)-2*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)+a*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*a-2*b*c*x)/(c*x-1)/(c*x+1)/c/b^2/(a+b*arccosh(c*x))`

**3.323.5 Fracas [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2+1)/(b^2*arccosh(c*x)^2+2*a*b*arccosh(c*x)+a^2),x)`

**3.323.6 Sympy [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{(a+b\operatorname{arcosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x))**2, x)`

**3.323.7 Maxima [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((2*c^2*x^2 + 1)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(2*c^3*x^3 - c*x)*(c*x + 1)*sqrt(c*x - 1) + (2*c^4*x^4 - 3*c^2*x^2 + 1)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^4*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^2*x^2 - 2*a*b*c^2*x^2 + 2*(a*b*c^3*x^3 - a*b*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + a*b + (b^2*c^4*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^2*x^2 - 2*b^2*c^2*x^2 + 2*(b^2*c^3*x^3 - b^2*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + b^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.323.8 Giac [F]**

$$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)/(b*arccosh(c*x) + a)^2, x)`

**3.323.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(1/2)/(a + b*acosh(c*x))^2,x)`output `int((1 - c^2*x^2)^(1/2)/(a + b*acosh(c*x))^2, x)`

$$3.324 \quad \int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$$

3.324.1 Optimal result	2636
3.324.2 Mathematica [N/A]	2637
3.324.3 Rubi [N/A]	2637
3.324.4 Maple [N/A] (verified)	2641
3.324.5 Fricas [N/A]	2641
3.324.6 Sympy [N/A]	2642
3.324.7 Maxima [N/A]	2642
3.324.8 Giac [F(-2)]	2643
3.324.9 Mupad [N/A]	2643

### 3.324.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}\sqrt{1-c^2x^2}}{bcx(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{b^2\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\operatorname{Int}\left(\frac{1}{x^2(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{-1+cx}}$$

output

```
cosh(a/b)*Shi((a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/(c*x-1)^(1/2)-Chi((a+b*arccosh(c*x))/b)*sinh(a/b)*(-c*x+1)^(1/2)/b^2/(c*x-1)^(1/2)-(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/b/c/x/(a+b*arccosh(c*x))+(c*x+1)^(1/2)*Unintegrable(1/x^2/(a+b*arccosh(c*x)),x)/b/c/(c*x-1)^(1/2)
```

**3.324.2 Mathematica [N/A]**

Not integrable

Time = 9.98 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x(a+\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x*(a + b*ArcCosh[c*x])^2), x]`output `Integrate[Sqrt[1 - c^2*x^2]/(x*(a + b*ArcCosh[c*x])^2), x]`**3.324.3 Rubi [N/A]**

Not integrable

Time = 1.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6357, 6296, 25, 3042, 26, 3784, 26, 3042, 26, 3779, 3782, 6303}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{1-c^2x^2}}{x(a+\operatorname{arccosh}(cx))^2} dx \\ & \quad \downarrow \text{6357} \\ & \frac{\sqrt{1-cx} \int \frac{1}{x^2(a+\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{c\sqrt{1-cx} \int \frac{1}{a+\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bcx(a+\operatorname{arccosh}(cx))} \\ & \quad \downarrow \text{6296} \\ & \frac{\sqrt{1-cx} \int -\frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) d(a+\operatorname{arccosh}(cx))}{a+\operatorname{arccosh}(cx)}}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{1}{x^2(a+\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\ & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bcx(a+\operatorname{arccosh}(cx))} \\ & \quad \downarrow \text{25} \end{aligned}$$

---

3.324.  $\int \frac{\sqrt{1-c^2x^2}}{x(a+\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
& -\frac{\sqrt{1-cx} \int \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
& \quad \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \quad \downarrow \text{3042} \\
& -\frac{\sqrt{1-cx} \int -\frac{i \sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
& \quad \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \quad \downarrow \text{26} \\
& \frac{i\sqrt{1-cx} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
& \quad \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \quad \downarrow \text{3784} \\
& i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) + \cosh\left(\frac{a}{b}\right) \int -\frac{i \sinh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right) \\
& \quad \frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{b^2\sqrt{cx-1}}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}} \\
& \quad \downarrow \text{26} \\
& i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right) \\
& \quad \frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{b^2\sqrt{cx-1}}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}} \\
& \quad \downarrow \text{3042}
\end{aligned}$$

---

3.324.  $\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$

$$i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))+\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \int \frac{i \sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)$$

---


$$\frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{b^2\sqrt{cx-1}}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2} bcx(a+b\operatorname{arccosh}(cx))}$$

↓ 26

$$i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))+\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)$$

---


$$\frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{b^2\sqrt{cx-1}}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2} bcx(a+b\operatorname{arccosh}(cx))}$$

↓ 3779

$$i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))+\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right) +$$

---


$$\frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{b^2\sqrt{cx-1}}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2} bcx(a+b\operatorname{arccosh}(cx))}$$

↓ 3782

$$\frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} + i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right) -$$

$$\frac{b^2\sqrt{cx-1}}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2} bcx(a+b\operatorname{arccosh}(cx))}$$

↓ 6303

$$\frac{\sqrt{1-cx} \int \frac{1}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} + i\sqrt{1-cx} \left( i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right) -$$

$$\frac{b^2\sqrt{cx-1}}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2} bcx(a+b\operatorname{arccosh}(cx))}$$



input `Int[Sqrt[1 - c^2*x^2]/(x*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

### 3.324.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`

rule 6296 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n, x_Symbol] := Simp[1/(b*c) Subst[Int[x^n*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x]`

rule 6303 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n*((d_.)*(x_)^m), x_Symbol] := Unintegrable[(d*x)^m*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, m, n}, x]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

### 3.324.4 Maple [N/A] (verified)

Not integrable

Time = 1.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2 + 1}}{x(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x))^2,x)`

### 3.324.5 Fracas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.54

$$\int \frac{\sqrt{1 - c^2x^2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2 + 1}}{(b \operatorname{arccosh}(cx) + a)^2 x} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(b^2*x*arccosh(c*x)^2 + 2*a*b*x*arccosh(c*x) + a^2*x), x)`

**3.324.6 Sympy [N/A]**

Not integrable

Time = 2.86 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{x(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x/(a+b*acosh(c*x))**2,x)`output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x*(a + b*acosh(c*x))**2), x)`**3.324.7 Maxima [N/A]**

Not integrable

Time = 0.69 (sec) , antiderivative size = 442, normalized size of antiderivative = 15.79

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)^2x} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`output `-((c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^3 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^2 - a*b*c*x + (b^2*c^3*x^3 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^2 - b^2*c*x)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((c^3*x^3 + 2*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + (2*c^4*x^4 + c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^5*x^5 - c^3*x^3)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^6 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^4 - 2*a*b*c^3*x^4 + a*b*c*x^2 + 2*(a*b*c^4*x^5 - a*b*c^2*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^6 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^4 - 2*b^2*c^3*x^4 + b^2*c*x^2 + 2*(b^2*c^4*x^5 - b^2*c^2*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.324.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(1/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.324.9 Mupad [N/A]**

Not integrable

Time = 3.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x*(a + b*acosh(c*x))^2),x)`

output `int((1 - c^2*x^2)^(1/2)/(x*(a + b*acosh(c*x))^2), x)`

**3.325**  $\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$

3.325.1 Optimal result . . . . .	2644
3.325.2 Mathematica [N/A] . . . . .	2644
3.325.3 Rubi [N/A] . . . . .	2645
3.325.4 Maple [N/A] (verified) . . . . .	2646
3.325.5 Fricas [N/A] . . . . .	2646
3.325.6 Sympy [N/A] . . . . .	2646
3.325.7 Maxima [N/A] . . . . .	2647
3.325.8 Giac [N/A] . . . . .	2647
3.325.9 Mupad [N/A] . . . . .	2648

**3.325.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}\sqrt{1-c^2x^2}}{bcx^2(a+b\operatorname{arccosh}(cx))} + \frac{2\sqrt{1-cx}\operatorname{Int}\left(\frac{1}{x^3(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{-1+cx}}$$

output `-(c*x-1)^(1/2)*(c*x+1)^(1/2)*(-c^2*x^2+1)^(1/2)/b/c/x^2/(a+b*arccosh(c*x)) + 2*(-c*x+1)^(1/2)*Unintegrable(1/x^3/(a+b*arccosh(c*x)), x)/b/c/(c*x-1)^(1/2)`

**3.325.2 Mathematica [N/A]**

Not integrable

Time = 3.96 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x^2*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[Sqrt[1 - c^2*x^2]/(x^2*(a + b*ArcCosh[c*x])^2), x]`

---

3.325.  $\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$

**3.325.3 Rubi [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6355, 6303}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+\operatorname{barccosh}(cx))^2} dx$$

↓ 6355

$$\frac{2\sqrt{1-cx} \int \frac{1}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bcx^2(a+\operatorname{barccosh}(cx))}$$

↓ 6303

$$\frac{2\sqrt{1-cx} \int \frac{1}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{1-c^2x^2}}{bcx^2(a+\operatorname{barccosh}(cx))}$$

input `Int[Sqrt[1 - c^2*x^2]/(x^2*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.325.3.1 Defintions of rubi rules used**

rule 6303 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((d_.)*(x_))^(m_.), x_Symbol]
:> Unintegrable[(d*x)^m*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, m, n}, x]`

rule 6355 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol]
:> Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && EqQ[m + 2*p + 1, 0]`

---

3.325.  $\int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$

**3.325.4 Maple [N/A] (verified)**

Not integrable

Time = 0.97 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2 + 1}}{x^2 (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x))^2,x)`output `int((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x))^2,x)`**3.325.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{\sqrt{1 - c^2x^2}}{x^2(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2 + 1}}{(b \operatorname{arccosh}(cx) + a)^2 x^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="fracas")`output `integral(sqrt(-c^2*x^2 + 1)/(b^2*x^2*arccosh(c*x)^2 + 2*a*b*x^2*arccosh(c*x) + a^2*x^2), x)`**3.325.6 Sympy [N/A]**

Not integrable

Time = 7.05 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt{1 - c^2x^2}}{x^2(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-(cx - 1)(cx + 1)}}{x^2 (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x**2/(a+b*acosh(c*x))**2,x)`output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x**2*(a + b*acosh(c*x))**2), x)`

---

3.325.  $\int \frac{\sqrt{1 - c^2x^2}}{x^2(a + b \operatorname{arccosh}(cx))^2} dx$

**3.325.7 Maxima [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 427, normalized size of antiderivative = 15.25

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)^2x^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^3 - a*b*c*x^2 + (b^2*c^3*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^3 - b^2*c*x^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate((3*(c*x + 1)^(3/2)*(c*x - 1)*c*x + 2*(2*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^7 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^5 - 2*a*b*c^3*x^5 + a*b*c*x^3 + 2*(a*b*c^4*x^6 - a*b*c^2*x^4)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^7 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^5 - 2*b^2*c^3*x^5 + b^2*c*x^3 + 2*(b^2*c^4*x^6 - b^2*c^2*x^4)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.325.8 Giac [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)^2x^2} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)^2*x^2), x)`



**3.325.9 Mupad [N/A]**

Not integrable

Time = 2.86 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^2(a+\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x^2(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x^2*(a + b*acosh(c*x))^2), x)`output `int((1 - c^2*x^2)^(1/2)/(x^2*(a + b*acosh(c*x))^2), x)`

**3.326**  $\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$

3.326.1 Optimal result . . . . .	2649
3.326.2 Mathematica [N/A] . . . . .	2649
3.326.3 Rubi [N/A] . . . . .	2650
3.326.4 Maple [N/A] (verified) . . . . .	2650
3.326.5 Fricas [N/A] . . . . .	2651
3.326.6 Sympy [N/A] . . . . .	2651
3.326.7 Maxima [N/A] . . . . .	2651
3.326.8 Giac [F(-2)] . . . . .	2652
3.326.9 Mupad [N/A] . . . . .	2652

**3.326.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x))^2, x)`

**3.326.2 Mathematica [N/A]**

Not integrable

Time = 97.63 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x^3*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[Sqrt[1 - c^2*x^2]/(x^3*(a + b*ArcCosh[c*x])^2), x]`

**3.326.3 Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{arccosh}(cx))^2} dx$$

input `Int[Sqrt[1 - c^2*x^2]/(x^3*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.326.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.326.4 Maple [N/A] (verified)**

Not integrable

Time = 1.21 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2+1}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x))^2,x)`

**3.326.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)^2 x^3} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(b^2*x^3*arccosh(c*x)^2 + 2*a*b*x^3*arccosh(c*x) + a^2*x^3), x)`

**3.326.6 Sympy [N/A]**

Not integrable

Time = 21.38 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{x^3(a+b \operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x**3/(a+b*acosh(c*x))**2,x)`

output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x**3*(a + b*acosh(c*x))**2), x)`

**3.326.7 Maxima [N/A]**

Not integrable

Time = 0.77 (sec) , antiderivative size = 453, normalized size of antiderivative = 16.18

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)^2 x^3} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^5 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^4 - a*b*c*x^3 + (b^2*c^3*x^5 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^4 - b^2*c*x^3)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate(((c^3*x^3 - 4*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + (2*c^4*x^4 - 7*c^2*x^2 + 3)*(c*x + 1)*sqrt(c*x - 1) + (c^5*x^5 - 3*c^3*x^3 + 2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^8 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^6 - 2*a*b*c^3*x^6 + a*b*c*x^4 + 2*(a*b*c^4*x^7 - a*b*c^2*x^5)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^8 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^6 - 2*b^2*c^3*x^6 + b^2*c*x^4 + 2*(b^2*c^4*x^7 - b^2*c^2*x^5)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.326.8 Giac [F(-2)]

Exception generated.

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(1/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.326.9 Mupad [N/A]

Not integrable

Time = 3.45 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x^3*(a + b*acosh(c*x))^2),x)`

---

3.326.  $\int \frac{\sqrt{1-c^2x^2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$

output `int((1 - c^2*x^2)^(1/2)/(x^3*(a + b*acosh(c*x))^2), x)`

**3.327**  $\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\text{arccosh}(cx))^2} dx$

3.327.1 Optimal result . . . . .	2654
3.327.2 Mathematica [N/A] . . . . .	2654
3.327.3 Rubi [N/A] . . . . .	2655
3.327.4 Maple [N/A] (verified) . . . . .	2655
3.327.5 Fricas [N/A] . . . . .	2656
3.327.6 Sympy [N/A] . . . . .	2656
3.327.7 Maxima [N/A] . . . . .	2656
3.327.8 Giac [N/A] . . . . .	2657
3.327.9 Mupad [N/A] . . . . .	2657

**3.327.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\text{arccosh}(cx))^2} dx = \text{Int}\left(\frac{\sqrt{1-c^2x^2}}{x^4(a+b\text{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x))^2, x)`

**3.327.2 Mathematica [N/A]**

Not integrable

Time = 5.62 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\text{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\text{arccosh}(cx))^2} dx$$

input `Integrate[Sqrt[1 - c^2*x^2]/(x^4*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[Sqrt[1 - c^2*x^2]/(x^4*(a + b*ArcCosh[c*x])^2), x]`

**3.327.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{arccosh}(cx))^2} dx$$

input `Int[Sqrt[1 - c^2*x^2]/(x^4*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.327.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.327.4 Maple [N/A] (verified)**

Not integrable

Time = 1.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt{-c^2x^2+1}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x))^2,x)`



**3.327.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(b^2*x^4*arccosh(c*x)^2 + 2*a*b*x^4*arccosh(c*x) + a^2*x^4), x)`

**3.327.6 Sympy [N/A]**

Not integrable

Time = 48.47 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-(cx-1)(cx+1)}}{x^4(a+b \operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(1/2)/x**4/(a+b*acosh(c*x))**2,x)`

output `Integral(sqrt(-(c*x - 1)*(c*x + 1))/(x**4*(a + b*acosh(c*x))**2), x)`

**3.327.7 Maxima [N/A]**

Not integrable

Time = 0.76 (sec) , antiderivative size = 456, normalized size of antiderivative = 16.29

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b \operatorname{arcosh}(cx)+a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^6 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^5 - a*b*c*x^4 + (b^2*c^3*x^6 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^5 - b^2*c*x^4)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate(((2*c^3*x^3 - 5*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(2*c^4*x^4 - 5*c^2*x^2 + 2)*(c*x + 1)*sqrt(c*x - 1) + (2*c^5*x^5 - 5*c^3*x^3 + 3*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^9 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^7 - 2*a*b*c^3*x^7 + a*b*c*x^5 + 2*(a*b*c^4*x^8 - a*b*c^2*x^6)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^9 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^7 - 2*b^2*c^3*x^7 + b^2*c*x^5 + 2*(b^2*c^4*x^8 - b^2*c^2*x^6)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.327.8 Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}}{(b\operatorname{arcosh}(cx)+a)^2x^4} dx$$

input `integrate((-c^2*x^2+1)^(1/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(sqrt(-c^2*x^2 + 1)/((b*arccosh(c*x) + a)^2*x^4), x)`

### 3.327.9 Mupad [N/A]

Not integrable

Time = 3.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(1/2)/(x^4*(a + b*acosh(c*x))^2),x)`

output `int((1 - c^2*x^2)^(1/2)/(x^4*(a + b*acosh(c*x))^2), x)`

---

3.327.  $\int \frac{\sqrt{1-c^2x^2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$

**3.328**  $\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.328.1 Optimal result . . . . . 2658  
 3.328.2 Mathematica [A] (verified) . . . . . 2659  
 3.328.3 Rubi [A] (verified) . . . . . 2660  
 3.328.4 Maple [A] (verified) . . . . . 2663  
 3.328.5 Fricas [F] . . . . . 2664  
 3.328.6 Sympy [F] . . . . . 2664  
 3.328.7 Maxima [F] . . . . . 2664  
 3.328.8 Giac [F] . . . . . 2665  
 3.328.9 Mupad [F(-1)] . . . . . 2665

**3.328.1 Optimal result**

Integrand size = 28, antiderivative size = 354

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^2\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)}{16b^2c^3\sqrt{-1+cx}} - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{4a}{b}\right)}{4b^2c^3\sqrt{-1+cx}} + \frac{3\sqrt{1-cx}\operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{6a}{b}\right)}{16b^2c^3\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\cosh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^3\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\cosh\left(\frac{4a}{b}\right)\operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b^2c^3\sqrt{-1+cx}} - \frac{3\sqrt{1-cx}\cosh\left(\frac{6a}{b}\right)\operatorname{Shi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^3\sqrt{-1+cx}}$$

output  $1/16*\cosh(2*a/b)*\text{Shi}(2*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}+1/4*\cosh(4*a/b)*\text{Shi}(4*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-3/16*\cosh(6*a/b)*\text{Shi}(6*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-1/16*\text{Chi}(2*(a+b*\text{arccosh}(c*x))/b)*\sinh(2*a/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-1/4*\text{Chi}(4*(a+b*\text{arccosh}(c*x))/b)*\sinh(4*a/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}+3/16*\text{Chi}(6*(a+b*\text{arccosh}(c*x))/b)*\sinh(6*a/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-x^2*(-c^2*x^2+1)^{(3/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c/(a+b*\text{arccosh}(c*x))$

### 3.328.2 Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 338, normalized size of antiderivative = 0.95

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\text{arccosh}(cx))^2} dx = \frac{\sqrt{-1+cx}\sqrt{1+cx}(16bc^2x^2-32bc^4x^4+16bc^6x^6-(a+b\text{arccosh}(cx))\text{Chi}(2(\frac{a}{b}+\text{arccosh}(cx)))\sinh(\frac{2a}{b}))}{(a+b\text{arccosh}(cx))^2}$$

input `Integrate[(x^2*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x])^2,x]`

output  $-1/16*(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(16*b*c^2*x^2 - 32*b*c^4*x^4 + 16*b*c^6*x^6 - (a + b*\text{ArcCosh}[c*x])*\text{CoshIntegral}[2*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(2*a)/b] - 4*(a + b*\text{ArcCosh}[c*x])*\text{CoshIntegral}[4*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(4*a)/b] + 3*a*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(6*a)/b] + 3*b*\text{ArcCosh}[c*x]*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(6*a)/b] + a*\text{Cosh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])] + b*\text{ArcCosh}[c*x]*\text{Cosh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])] + 4*a*\text{Cosh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])] + 4*b*\text{ArcCosh}[c*x]*\text{Cosh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])] - 3*a*\text{Cosh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])] - 3*b*\text{ArcCosh}[c*x]*\text{Cosh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])]))/(b^2*c^3*\text{Sqrt}[1 - c^2*x^2]*(a + b*\text{ArcCosh}[c*x]))$

**3.328.3 Rubi [A] (verified)**

Time = 1.30 (sec) , antiderivative size = 310, normalized size of antiderivative = 0.88, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6357, 25, 6327, 6367, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(1-c^2x^2)^{3/2}}{(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6357} \\
 & -\frac{6c\sqrt{1-cx} \int -\frac{x^3(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} + \frac{2\sqrt{1-cx} \int -\frac{x(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} - \\
 & \quad \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{6c\sqrt{1-cx} \int \frac{x^3(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{2\sqrt{1-cx} \int \frac{x(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} - \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & -\frac{2\sqrt{1-cx} \int \frac{x(1-c^2x^2)}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} + \frac{6c\sqrt{1-cx} \int \frac{x^3(1-c^2x^2)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6367} \\
 & -\frac{6\sqrt{1-cx} \int -\frac{\cosh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^3\sqrt{cx-1}} + \\
 & \quad \frac{2\sqrt{1-cx} \int -\frac{\cosh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^3\sqrt{cx-1}} - \\
 & \quad \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25}
 \end{aligned}$$

---

3.328.  $\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
& \frac{6\sqrt{1-cx} \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} \\
& \frac{2\sqrt{1-cx} \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} \\
& \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} (1-c^2 x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
& \quad \downarrow \text{5971} \\
& \frac{2\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} - \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} + \\
& \frac{6\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{6a}{b} - \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} - \frac{3 \sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^3 \sqrt{cx-1}} \\
& \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} (1-c^2 x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
& \quad \downarrow \text{2009} \\
& \frac{2\sqrt{1-cx} \left( \frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} \\
& \frac{6\sqrt{1-cx} \left( \frac{3}{32} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{32} \sinh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{32} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^3 \sqrt{cx-1}} \\
& \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} (1-c^2 x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))}
\end{aligned}$$

input `Int[(x^2*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x])^2,x]`

---

3.328.  $\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

output  $-\left(\frac{x^2\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{3/2}}{b^2c^3\sqrt{-1+cx}} + \frac{2\sqrt{1-cx}\left(\frac{\text{CoshIntegral}[2(a+b\text{ArcCosh}[cx])]}{b}\right)\text{Sinh}\left[\frac{2a}{b}\right]}{4} - \frac{\text{CoshIntegral}\left[\frac{4(a+b\text{ArcCosh}[cx])}{b}\right]\text{Sinh}\left[\frac{4a}{b}\right]}{8} - \frac{\text{Cosh}\left[\frac{2a}{b}\right]\text{SinhIntegral}\left[\frac{2(a+b\text{ArcCosh}[cx])}{b}\right]}{4} + \frac{\text{Cosh}\left[\frac{4a}{b}\right]\text{SinhIntegral}\left[\frac{4(a+b\text{ArcCosh}[cx])}{b}\right]}{8}\right)}{b^2c^3\sqrt{-1+cx}} - \frac{6\sqrt{1-cx}\left(\frac{3\text{CoshIntegral}[2(a+b\text{ArcCosh}[cx])]}{b}\right)\text{Sinh}\left[\frac{2a}{b}\right]}{32} - \frac{\text{CoshIntegral}\left[\frac{6(a+b\text{ArcCosh}[cx])}{b}\right]\text{Sinh}\left[\frac{6a}{b}\right]}{32} - \frac{3\text{Cosh}\left[\frac{2a}{b}\right]\text{SinhIntegral}\left[\frac{2(a+b\text{ArcCosh}[cx])}{b}\right]}{32} + \frac{\text{Cosh}\left[\frac{6a}{b}\right]\text{SinhIntegral}\left[\frac{6(a+b\text{ArcCosh}[cx])}{b}\right]}{32}\right)}{b^2c^3\sqrt{-1+cx}}$

### 3.328.3.1 Defintions of rubi rules used

rule 25  $\text{Int}[-(F_x), x\_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$

rule 2009  $\text{Int}[u, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] \text{ ; SumQ}[u]$

rule 5971  $\text{Int}[\text{Cosh}[a] + (b)(x)^{(p)}((c) + (d)(x))^{(m)}\text{Sinh}[a] + (b)(x)^{(n)}, x\_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(c + dx)^m, \text{Sinh}[a + bx]^n \text{Cosh}[a + bx]^p, x], x] \text{ ; FreeQ}\{a, b, c, d, m\}, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[p, 0]$

rule 6327  $\text{Int}[(a) + \text{ArcCosh}[c(x)](b)]^{(n)}((f)(x))^{(m)}((d) + (e)(x))^{(p)}, x\_Symbol] \rightarrow \text{Int}[(f*x)^m(d_1*d_2 + e_1*e_2*x^2)^p(a + b\text{ArcCosh}[cx])^n, x] \text{ ; FreeQ}\{a, b, c, d_1, e_1, d_2, e_2, f, m, n\}, x\} \ \&\& \ \text{EqQ}[d_2*e_1 + d_1*e_2, 0] \ \&\& \ \text{IntegerQ}[p]$

rule 6357  $\text{Int}[(a) + \text{ArcCosh}[c(x)](b)]^{(n)}((f)(x))^{(m)}((d) + (e)(x)^2)^{(p)}, x\_Symbol] \rightarrow \text{Simp}[(f*x)^m\text{Simp}[\sqrt{1+cx}\sqrt{-1+cx}](d + ex^2)^p(a + b\text{ArcCosh}[cx])^{(n+1)}(b^2c^{(n+1)}), x] + (\text{Simp}[f(m/(b^2c^{(n+1)}))\text{Simp}[(d + ex^2)^p/((1+cx)^p(-1+cx)^p)] \quad \text{Int}[(f*x)^{(m-1)}(1+cx)^{(p-1/2)}(-1+cx)^{(p-1/2)}(a + b\text{ArcCosh}[cx])^{(n+1)}, x], x] - \text{Simp}[c((m+2p+1)/(b*f^{(n+1)}))\text{Simp}[(d + ex^2)^p/((1+cx)^p(-1+cx)^p)] \quad \text{Int}[(f*x)^{(m+1)}(1+cx)^{(p-1/2)}(-1+cx)^{(p-1/2)}(a + b\text{ArcCosh}[cx])^{(n+1)}, x], x)] \text{ ; FreeQ}\{a, b, c, d, e, f, m, p\}, x\} \ \&\& \ \text{EqQ}[c^2d + e, 0] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IGtQ}[2p, 0] \ \&\& \ \text{NeQ}[m + 2p + 1, 0] \ \&\& \ \text{IGtQ}[m, -3]$

---

3.328.  $\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\text{arccosh}(cx))^2} dx$

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.328.4 Maple [A] (verified)

Time = 0.79 (sec) , antiderivative size = 599, normalized size of antiderivative = 1.69

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(32\sqrt{cx-1}\sqrt{cx+1}bc^6x^6+32bc^7x^7-64x^4c^4b\sqrt{cx-1}\sqrt{cx+1}-64bc^5x^5+32\sqrt{cx-1}\sqrt{cx+1}\right)}{\dots}$

input `int(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output `-1/32*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^6*x^6+32*b*c^7*x^7-64*x^4*c^4*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)-64*b*c^5*x^5+32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^2*x^2+32*b*c^3*x^3+3*arccosh(c*x)*b*Ei(1,-6*arccosh(c*x)-6*a/b)*exp(-(-b*arccosh(c*x)+6*a)/b)-arccosh(c*x)*b*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-4*arccosh(c*x)*b*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b)-3*Ei(1,6*arccosh(c*x)+6*a/b)*exp((b*arccosh(c*x)+6*a)/b)*b*arccosh(c*x)+4*Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)*b*arccosh(c*x)+Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*b*arccosh(c*x)+3*a*Ei(1,-6*arccosh(c*x)-6*a/b)*exp(-(-b*arccosh(c*x)+6*a)/b)-a*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-4*a*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b)-3*Ei(1,6*arccosh(c*x)+6*a/b)*exp((b*arccosh(c*x)+6*a)/b)*a+4*Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)*a+Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*a)/(c*x+1)/c^3/(c*x-1)/b^2/(a+b*arccosh(c*x))`

$$3.328. \int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$



**3.328.5 Fricas [F]**

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x^2}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-(c^2*x^4 - x^2)*sqrt(-c^2*x^2 + 1)/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

**3.328.6 Sympy [F]**

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2(-(cx-1)(cx+1))^{\frac{3}{2}}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x**2*(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(x**2*(-(c*x - 1)*(c*x + 1))**(3/2)/(a + b*acosh(c*x))**2, x)`

**3.328.7 Maxima [F]**

$$\int \frac{x^2(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x^2}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $((c^4x^6 - 2c^2x^4 + x^2)(cx + 1)\sqrt{cx - 1} + (c^5x^7 - 2c^3x^5 + cx^3)\sqrt{cx + 1})\sqrt{-cx + 1}/(abc^3x^2 + \sqrt{cx + 1})\sqrt{cx - 1} + abc^2x - abc + (b^2c^3x^2 + \sqrt{cx + 1})\sqrt{cx - 1} + b^2c^2x - b^2c) \log(cx + \sqrt{cx + 1})\sqrt{cx - 1}) - \text{integrate}(((6c^5x^6 - 7c^3x^4 + cx^2)(cx + 1)^{3/2}(cx - 1) + 2(6c^6x^7 - 11c^4x^5 + 6c^2x^3 - x)(cx + 1)\sqrt{cx - 1} + 3(2c^7x^8 - 5c^5x^6 + 4c^3x^4 - cx^2)\sqrt{cx + 1})\sqrt{-cx + 1}/(abc^5x^4 + (cx + 1)(cx - 1)abc^3x^2 - 2abc^3x^2 + abc + 2(abc^4x^3 - abc^2x)\sqrt{cx + 1})\sqrt{cx - 1} + (b^2c^5x^4 + (cx + 1)(cx - 1)b^2c^3x^2 - 2b^2c^3x^2 + b^2c + 2(b^2c^4x^3 - b^2c^2x)\sqrt{cx + 1})\sqrt{cx - 1}) \log(cx + \sqrt{cx + 1})\sqrt{cx - 1})), x)$

### 3.328.8 Giac [F]

$$\int \frac{x^2(1 - c^2x^2)^{3/2}}{(a + \text{barccosh}(cx))^2} dx = \int \frac{(-c^2x^2 + 1)^{3/2}x^2}{(b \text{arcosh}(cx) + a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x^2/(b*arccosh(c*x) + a)^2, x)`

### 3.328.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^2(1 - c^2x^2)^{3/2}}{(a + \text{barccosh}(cx))^2} dx = \int \frac{x^2(1 - c^2x^2)^{3/2}}{(a + b \text{acosh}(cx))^2} dx$$

input `int((x^2*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x))^2,x)`

output `int((x^2*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x))^2, x)`

**3.329** 
$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

3.329.1 Optimal result . . . . . 2666  
 3.329.2 Mathematica [A] (verified) . . . . . 2667  
 3.329.3 Rubi [C] (verified) . . . . . 2667  
 3.329.4 Maple [A] (verified) . . . . . 2672  
 3.329.5 Fricas [F] . . . . . 2673  
 3.329.6 Sympy [F] . . . . . 2673  
 3.329.7 Maxima [F] . . . . . 2674  
 3.329.8 Giac [F] . . . . . 2674  
 3.329.9 Mupad [F(-1)] . . . . . 2675

**3.329.1 Optimal result**

Integrand size = 26, antiderivative size = 348

$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} + \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{8b^2c^2\sqrt{-1+cx}} - \frac{9\sqrt{1-cx}\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{16b^2c^2\sqrt{-1+cx}} + \frac{5\sqrt{1-cx}\operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{5a}{b}\right)}{16b^2c^2\sqrt{-1+cx}} - \frac{\sqrt{1-cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b^2c^2\sqrt{-1+cx}} + \frac{9\sqrt{1-cx}\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^2\sqrt{-1+cx}} - \frac{5\sqrt{1-cx}\cosh\left(\frac{5a}{b}\right)\operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^2\sqrt{-1+cx}}$$

output

```
-1/8*cosh(a/b)*Shi((a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)+9/16*cosh(3*a/b)*Shi(3*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)-5/16*cosh(5*a/b)*Shi(5*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)+1/8*Chi((a+b*arccosh(c*x))/b)*sinh(a/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)-9/16*Chi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)+5/16*Chi(5*(a+b*arccosh(c*x))/b)*sinh(5*a/b)*(-c*x+1)^(1/2)/b^2/c^2/(c*x-1)^(1/2)-x*(-c^2*x^2+1)^(3/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(a+b*arccosh(c*x))
```

3.329. 
$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.329.2 Mathematica [A] (verified)**

Time = 0.75 (sec) , antiderivative size = 327, normalized size of antiderivative = 0.94

$$\int \frac{x(1 - c^2x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(-16bcx + 32bc^3x^3 - 16bc^5x^5 - 2(a + \operatorname{barccosh}(cx))\operatorname{Chi}\left(\frac{a}{b}\right))}{(a + \operatorname{barccosh}(cx))^2}$$

input `Integrate[(x*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x])^2,x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-16*b*c*x + 32*b*c^3*x^3 - 16*b*c^5*x^5 - 2*(a + b*ArcCosh[c*x])*CoshIntegral[a/b + ArcCosh[c*x]]*Sinh[a/b] + 9*(a + b*ArcCosh[c*x])*CoshIntegral[3*(a/b + ArcCosh[c*x]]*Sinh[(3*a)/b] - 5*a*CoshIntegral[5*(a/b + ArcCosh[c*x]]*Sinh[(5*a)/b] - 5*b*ArcCosh[c*x]*CoshIntegral[5*(a/b + ArcCosh[c*x]]*Sinh[(5*a)/b] + 2*a*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + 2*b*ArcCosh[c*x]*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] - 9*a*Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] - 9*b*ArcCosh[c*x]*Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] + 5*a*Cosh[(5*a)/b]*SinhIntegral[5*(a/b + ArcCosh[c*x])] + 5*b*ArcCosh[c*x]*Cosh[(5*a)/b]*SinhIntegral[5*(a/b + ArcCosh[c*x])])/(16*b^2*c^2*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x]))`

**3.329.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.05 (sec) , antiderivative size = 360, normalized size of antiderivative = 1.03, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {6357, 25, 6304, 6321, 25, 3042, 26, 3793, 2009, 6327, 6367, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(1 - c^2x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx$$

↓ 6357

$$-\frac{5c\sqrt{1 - cx} \int -\frac{x^2(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx - 1}} + \frac{\sqrt{1 - cx} \int -\frac{(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx - 1}}$$

$$\frac{x\sqrt{cx - 1}\sqrt{cx + 1}(1 - c^2x^2)^{3/2}}{bc(a + \operatorname{barccosh}(cx))}$$

---

3.329.  $\int \frac{x(1 - c^2x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \downarrow 6304 \\
& -\frac{\sqrt{1-cx} \int \frac{1-c^2x^2}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} + \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \downarrow 6321 \\
& \frac{\sqrt{1-cx} \int -\frac{\sinh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
& \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \downarrow 25 \\
& -\frac{\sqrt{1-cx} \int \frac{\sinh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
& \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \downarrow 3042 \\
& -\frac{\sqrt{1-cx} \int \frac{i \sin\left(\frac{ia}{b}-\frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^3}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
& \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \downarrow 26 \\
& -\frac{i\sqrt{1-cx} \int \frac{\sin\left(\frac{ia}{b}-\frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^3}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
& \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \downarrow 3793
\end{aligned}$$

---

3.329.  $\int \frac{x(1-c^2x^2)^{3/2}}{(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{i\sqrt{1-cx} \int \left( \frac{3i \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} - \frac{i \sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} + \\
 & \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{2009} \\
 & \frac{5c\sqrt{1-cx} \int \frac{x^2(1-cx)(cx+1)}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \frac{i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & \frac{5c\sqrt{1-cx} \int \frac{x^2(1-c^2x^2)}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \frac{i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{6367} \\
 & \frac{5\sqrt{1-cx} \int -\frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} - \\
 & \frac{i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{5\sqrt{1-cx} \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} - \\
 & \frac{i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))}
 \end{aligned}$$

---

3.329.  $\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned} & \downarrow \text{5971} \\ & 5\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{\sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} \right) d(a + \operatorname{arccosh}(cx)) \\ & \frac{i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\ & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a + \operatorname{arccosh}(cx))} \end{aligned}$$

$$\begin{aligned} & \downarrow \text{2009} \\ & \frac{i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{b^2 c^2 \sqrt{cx-1}} \\ & \frac{5\sqrt{1-cx} \left( \frac{1}{8} \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) + \frac{1}{16} \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{16} \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2 c^2} \\ & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a + \operatorname{arccosh}(cx))} \end{aligned}$$

input `Int[(x*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x])^2,x]`

output `-((x*sqrt[-1 + c*x]*sqrt[1 + c*x]*(1 - c^2*x^2)^(3/2))/(b*c*(a + b*ArcCosh[c*x]))) - (I*sqrt[1 - c*x]*(((3*I)/4)*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b] - (I/4)*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*a)/b] - ((3*I)/4)*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b] + (I/4)*Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b]))/(b^2*c^2*sqrt[-1 + c*x]) - (5*sqrt[1 - c*x]*((CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b])/8 + (CoshIntegral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*a)/b])/16 - (CoshIntegral[(5*(a + b*ArcCosh[c*x])/b]*Sinh[(5*a)/b])/16 - (Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/8 - (Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/16 + (Cosh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/16)))/(b^2*c^2*sqrt[-1 + c*x])`

## 3.329.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`
- rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`
- rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

---

3.329. 
$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$



rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.329.4 Maple [A] (verified)

Time = 0.96 (sec) , antiderivative size = 583, normalized size of antiderivative = 1.68

method	result
default	$\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-32\sqrt{cx-1}\sqrt{cx+1}bc^5x^5-32bc^6x^6+64\sqrt{cx-1}\sqrt{cx+1}bc^3x^3+64bc^4x^4-32\sqrt{cx-1}\sqrt{cx+1}bc^2x^2\right)}{(a+b\operatorname{arccosh}(cx))^2}$

input `int(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

---

3.329. 
$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

output  $\frac{1}{32}(-c^2x^2+1)^{1/2}(-(cx-1)^{1/2}(cx+1)^{1/2}cx+c^2x^2-1)(-32(cx-1)^{1/2}(cx+1)^{1/2}bc^5x^5-32bc^6x^6+64(cx-1)^{1/2}(cx+1)^{1/2}b^2c^3x^3+64b^2c^4x^4-32(cx-1)^{1/2}(cx+1)^{1/2}b^2cx-32b^2c^2x^2+9\operatorname{arccosh}(cx)bEi(1,-3\operatorname{arccosh}(cx)-3a/b)\exp(-(-b\operatorname{arccosh}(cx)+3a)/b)-2\operatorname{arccosh}(cx)bEi(1,-\operatorname{arccosh}(cx)-a/b)\exp(-(-b\operatorname{arccosh}(cx)+a)/b)-5\operatorname{arccosh}(cx)bEi(1,-5\operatorname{arccosh}(cx)-5a/b)\exp(-(-b\operatorname{arccosh}(cx)+5a)/b)+5Ei(1,5\operatorname{arccosh}(cx)+5a/b)\exp((b\operatorname{arccosh}(cx)+5a)/b)b\operatorname{arccosh}(cx)-9Ei(1,3\operatorname{arccosh}(cx)+3a/b)\exp((b\operatorname{arccosh}(cx)+3a)/b)b\operatorname{arccosh}(cx)+2Ei(1,\operatorname{arccosh}(cx)+a/b)\exp((a+b\operatorname{arccosh}(cx))/b)b\operatorname{arccosh}(cx)+9aEi(1,-3\operatorname{arccosh}(cx)-3a/b)\exp(-(-b\operatorname{arccosh}(cx)+3a)/b)-2aEi(1,-\operatorname{arccosh}(cx)-a/b)\exp(-(-b\operatorname{arccosh}(cx)+a)/b)-5aEi(1,-5\operatorname{arccosh}(cx)-5a/b)\exp(-(-b\operatorname{arccosh}(cx)+5a)/b)+5Ei(1,5\operatorname{arccosh}(cx)+5a/b)\exp((b\operatorname{arccosh}(cx)+5a)/b)a-9Ei(1,3\operatorname{arccosh}(cx)+3a/b)\exp((b\operatorname{arccosh}(cx)+3a)/b)a+2Ei(1,\operatorname{arccosh}(cx)+a/b)\exp((a+b\operatorname{arccosh}(cx))/b)a)/(cx+1)/c^2/(cx-1)/b^2/(a+b\operatorname{arccosh}(cx))$

### 3.329.5 Fracas [F]

$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{3/2}x}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-(c^2*x^3 - x)*sqrt(-c^2*x^2 + 1)/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

### 3.329.6 Sympy [F]

$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x(-(cx-1)(cx+1))^{3/2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x*(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(x*(-(c*x - 1)*(c*x + 1))**(3/2)/(a + b*acosh(c*x))**2, x)`

---

3.329.  $\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

**3.329.7 Maxima [F]**

$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x}{(b \operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `((c^4*x^5 - 2*c^2*x^3 + x)*(c*x + 1)*sqrt(c*x - 1) + (c^5*x^6 - 2*c^3*x^4 + c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate((5*(c^5*x^5 - c^3*x^3)*(c*x + 1)^(3/2)*(c*x - 1) + (10*c^6*x^6 - 17*c^4*x^4 + 8*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (5*c^7*x^7 - 12*c^5*x^5 + 9*c^3*x^3 - 2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.329.8 Giac [F]**

$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{\frac{3}{2}}x}{(b \operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)*x/(b*arccosh(c*x) + a)^2, x)`

**3.329.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x(1-c^2x^2)^{3/2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((x*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x))^2,x)`output `int((x*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x))^2, x)`

**3.330**  $\int \frac{(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.330.1 Optimal result . . . . . 2676  
 3.330.2 Mathematica [A] (verified) . . . . . 2677  
 3.330.3 Rubi [A] (verified) . . . . . 2677  
 3.330.4 Maple [B] (verified) . . . . . 2680  
 3.330.5 Fricas [F] . . . . . 2680  
 3.330.6 Sympy [F] . . . . . 2681  
 3.330.7 Maxima [F] . . . . . 2681  
 3.330.8 Giac [F] . . . . . 2682  
 3.330.9 Mupad [F(-1)] . . . . . 2682

**3.330.1 Optimal result**

Integrand size = 25, antiderivative size = 246

$$\int \frac{(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{3/2}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)}{b^2c\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{4a}{b}\right)}{2b^2c\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\cosh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{b^2c\sqrt{-1+cx}} - \frac{\sqrt{1-cx}\cosh\left(\frac{4a}{b}\right)\operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{2b^2c\sqrt{-1+cx}}$$

output

```
cosh(2*a/b)*Shi(2*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c/(c*x-1)^(1/2)
-1/2*cosh(4*a/b)*Shi(4*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/c/(c*x-1)^(1/2)
-Chi(2*(a+b*arccosh(c*x))/b)*sinh(2*a/b)*(-c*x+1)^(1/2)/b^2/c/(c*x-1)^(1/2)
+1/2*Chi(4*(a+b*arccosh(c*x))/b)*sinh(4*a/b)*(-c*x+1)^(1/2)/b^2/c/(c*x-1)^(1/2)
-(c^2*x^2+1)^(3/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(a+b*arccosh(c*x))
```

3.330.  $\int \frac{(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

**3.330.2 Mathematica [A] (verified)**

Time = 0.49 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.94

$$\int \frac{(1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(-2b + 4bc^2x^2 - 2bc^4x^4 + 2(a + \operatorname{barccosh}(cx))\operatorname{Chi}(2(\frac{a}{b} + \operatorname{arccosh}(cx))))}{(a + \operatorname{barccosh}(cx))^2}$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(a + b*ArcCosh[c*x])^2,x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(-2*b + 4*b*c^2*x^2 - 2*b*c^4*x^4 + 2*(a + b*ArcCosh[c*x])*CoshIntegral[2*(a/b + ArcCosh[c*x]])*Sinh[(2*a)/b] - (a + b*ArcCosh[c*x])*CoshIntegral[4*(a/b + ArcCosh[c*x]])*Sinh[(4*a)/b] - 2*a*Cosh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[c*x]]) - 2*b*ArcCosh[c*x]*Cosh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[c*x]]) + a*Cosh[(4*a)/b]*SinhIntegral[4*(a/b + ArcCosh[c*x]]) + b*ArcCosh[c*x]*Cosh[(4*a)/b]*SinhIntegral[4*(a/b + ArcCosh[c*x])])/(2*b^2*c*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x]))`

**3.330.3 Rubi [A] (verified)**Time = 0.76 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.73, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$ , Rules used = {6319, 25, 6327, 6367, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx \\ & \quad \downarrow \text{6319} \\ & -\frac{4c\sqrt{1 - cx} \int -\frac{x(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx - 1}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}(1 - c^2 x^2)^{3/2}}{bc(a + \operatorname{barccosh}(cx))} \\ & \quad \downarrow \text{25} \\ & \frac{4c\sqrt{1 - cx} \int \frac{x(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx - 1}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}(1 - c^2 x^2)^{3/2}}{bc(a + \operatorname{barccosh}(cx))} \\ & \quad \downarrow \text{6327} \end{aligned}$$

---

3.330.  $\int \frac{(1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
& \frac{4c\sqrt{1-cx} \int \frac{x(1-c^2x^2)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \quad \downarrow \text{6367} \\
& \frac{4\sqrt{1-cx} \int -\frac{\cosh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \quad \downarrow \text{25} \\
& \frac{4\sqrt{1-cx} \int \frac{\cosh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \quad \downarrow \text{5971} \\
& \frac{4\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{4a}{b}-\frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{8(a+\operatorname{barccosh}(cx))} - \frac{\sinh\left(\frac{2a}{b}-\frac{2(a+\operatorname{barccosh}(cx))}{b}\right)}{4(a+\operatorname{barccosh}(cx))} \right) d(a+\operatorname{barccosh}(cx))}{b^2c\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))} \\
& \quad \downarrow \text{2009} \\
& \frac{4\sqrt{1-cx} \left( \frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+\operatorname{barccosh}(cx))}{b}\right) - \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) \right)}{b^2c\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bc(a+\operatorname{barccosh}(cx))}
\end{aligned}$$

input `Int[(1 - c^2*x^2)^(3/2)/(a + b*ArcCosh[c*x])^2,x]`

output `-((Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(1 - c^2*x^2)^(3/2))/(b*c*(a + b*ArcCosh[c*x]))) - (4*Sqrt[1 - c*x]*((CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]*Sinh[(2*a)/b])/4 - (CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b]*Sinh[(4*a)/b])/8 - (Cosh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/4 + (Cosh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8))/(b^2*c*Sqrt[-1 + c*x])`

---

3.330.  $\int \frac{(1-c^2x^2)^{3/2}}{(a+\operatorname{barccosh}(cx))^2} dx$

## 3.330.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`
- rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1))]*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`



**3.330.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 444 vs.  $2(220) = 440$ .

Time = 0.92 (sec) , antiderivative size = 445, normalized size of antiderivative = 1.81

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(4x^4c^4b\sqrt{cx-1}\sqrt{cx+1}+4bc^5x^5-8\sqrt{cx-1}\sqrt{cx+1}bc^2x^2-8bc^3x^3+\operatorname{arccosh}(cx)b\operatorname{Ei}_1\right)}{\dots}$

```
input int((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output -1/4*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(4*x^4*c^4*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*b*c^5*x^5-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^2*x^2-8*b*c^3*x^3+arccosh(c*x)*b*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b)-2*arccosh(c*x)*b*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)*b*arccosh(c*x)+2*Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*b*arccosh(c*x)+4*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)+a*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b)-2*a*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)*a+2*Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*a+4*b*c*x)/(c*x-1)/(c*x+1)/c/b^2/(a+b*arccosh(c*x))
```

**3.330.5 Fracas [F]**

$$\int \frac{(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{3/2}}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

```
input integrate((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")
```

```
output integral((-c^2*x^2+1)^(3/2)/(b^2*arccosh(c*x)^2+2*a*b*arccosh(c*x)+a^2), x)
```

**3.330.6 Sympy [F]**

$$\int \frac{(1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-(cx - 1)(cx + 1))^{\frac{3}{2}}}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Integral((-c*x - 1)*(c*x + 1)**(3/2)/(a + b*acosh(c*x))**2, x)`

**3.330.7 Maxima [F]**

$$\int \frac{(1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `((c^4*x^4 - 2*c^2*x^2 + 1)*(c*x + 1)*sqrt(c*x - 1) + (c^5*x^5 - 2*c^3*x^3 + c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate(((4*c^4*x^4 - 3*c^2*x^2 - 1)*(c*x + 1)^(3/2)*(c*x - 1) + 4*(2*c^5*x^5 - 3*c^3*x^3 + c*x)*(c*x + 1)*sqrt(c*x - 1) + (4*c^6*x^6 - 9*c^4*x^4 + 6*c^2*x^2 - 1)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^4*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^2*x^2 - 2*a*b*c^2*x^2 + 2*(a*b*c^3*x^3 - a*b*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + a*b + (b^2*c^4*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^2*x^2 - 2*b^2*c^2*x^2 + 2*(b^2*c^3*x^3 - b^2*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + b^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.330.8 Giac [F]**

$$\int \frac{(1 - c^2 x^2)^{3/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)/(b*arccosh(c*x) + a)^2, x)`

**3.330.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(3/2)/(a + b*acosh(c*x))^2,x)`

output `int((1 - c^2*x^2)^(3/2)/(a + b*acosh(c*x))^2, x)`

**3.331**  $\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$

3.331.1 Optimal result . . . . . 2683  
 3.331.2 Mathematica [N/A] . . . . . 2684  
 3.331.3 Rubi [N/A] . . . . . 2684  
 3.331.4 Maple [N/A] (verified) . . . . . 2688  
 3.331.5 Fricas [N/A] . . . . . 2688  
 3.331.6 Sympy [N/A] . . . . . 2688  
 3.331.7 Maxima [N/A] . . . . . 2689  
 3.331.8 Giac [F(-2)] . . . . . 2689  
 3.331.9 Mupad [N/A] . . . . . 2690

**3.331.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{3/2}}{bcx(a+b\operatorname{arccosh}(cx))} - \frac{9\sqrt{1-cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{4b^2\sqrt{-1+cx}} + \frac{3\sqrt{1-cx}\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{4b^2\sqrt{-1+cx}} + \frac{9\sqrt{1-cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4b^2\sqrt{-1+cx}} - \frac{3\sqrt{1-cx}\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b^2\sqrt{-1+cx}} - \frac{\sqrt{1-cx}\operatorname{Int}\left(\frac{-1+c^2x^2}{x^2(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{-1+cx}}$$

```
output 9/4*cosh(a/b)*Shi((a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/(c*x-1)^(1/2)-3
/4*cosh(3*a/b)*Shi(3*(a+b*arccosh(c*x))/b)*(-c*x+1)^(1/2)/b^2/(c*x-1)^(1/2)
)-9/4*Chi((a+b*arccosh(c*x))/b)*sinh(a/b)*(-c*x+1)^(1/2)/b^2/(c*x-1)^(1/2)
+3/4*Chi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)*(-c*x+1)^(1/2)/b^2/(c*x-1)^(1
/2)-(-c^2*x^2+1)^(3/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/x/(a+b*arccosh(c*x)
)-(-c*x+1)^(1/2)*Unintegrable((c^2*x^2-1)/x^2/(a+b*arccosh(c*x)),x)/b/c/(c
*x-1)^(1/2)
```

3.331.  $\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$

**3.331.2 Mathematica [N/A]**

Not integrable

Time = 18.83 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))^2} dx$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(x*(a + b*ArcCosh[c*x])^2), x]`output `Integrate[(1 - c^2*x^2)^(3/2)/(x*(a + b*ArcCosh[c*x])^2), x]`**3.331.3 Rubi [N/A]**

Not integrable

Time = 1.67 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6357, 25, 6304, 6321, 25, 3042, 26, 3793, 2009, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))^2} dx \\ & \quad \downarrow \text{6357} \\ & -\frac{\sqrt{1 - cx} \int -\frac{(1-cx)(cx+1)}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{3c\sqrt{1 - cx} \int -\frac{(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \\ & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+\operatorname{barccosh}(cx))} \\ & \quad \downarrow \text{25} \\ & \frac{\sqrt{1 - cx} \int \frac{(1-cx)(cx+1)}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{3c\sqrt{1 - cx} \int \frac{(1-cx)(cx+1)}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+\operatorname{barccosh}(cx))} \\ & \quad \downarrow \text{6304} \\ & \frac{3c\sqrt{1 - cx} \int \frac{1-c^2x^2}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} + \frac{\sqrt{1 - cx} \int \frac{(1-cx)(cx+1)}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+\operatorname{barccosh}(cx))} \end{aligned}$$

---

3.331.  $\int \frac{(1-c^2x^2)^{3/2}}{x(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
& \downarrow \text{6321} \\
& \frac{3\sqrt{1-cx} \int -\frac{\sinh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
& \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{25} \\
& \frac{3\sqrt{1-cx} \int \frac{\sinh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
& \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{3042} \\
& \frac{3\sqrt{1-cx} \int \frac{i \sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^3}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
& \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{26} \\
& \frac{3i\sqrt{1-cx} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^3}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
& \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{3793} \\
& \frac{3i\sqrt{1-cx} \int \left( \frac{3i \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} - \frac{i \sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2\sqrt{cx-1}} + \\
& \quad \frac{\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+b\operatorname{arccosh}(cx))} \\
& \downarrow \text{2009}
\end{aligned}$$

---

3.331.  $\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$

$$\frac{\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{3i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) \right)}{b^2\sqrt{cx-1}}$$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 6327

$$\frac{\sqrt{1-cx} \int \frac{1-c^2x^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{3i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) \right)}{b^2\sqrt{cx-1}}$$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 6375

$$\frac{\sqrt{1-cx} \int \frac{1-c^2x^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{3i\sqrt{1-cx} \left( \frac{3}{4}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) - \frac{1}{4}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right) - \frac{3}{4}i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) \right)}{b^2\sqrt{cx-1}}$$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

input `Int[(1 - c^2*x^2)^(3/2)/(x*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

### 3.331.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

---

3.331.  $\int \frac{(1-c^2x^2)^{3/2}}{x(a+\operatorname{barccosh}(cx))^2} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_)^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_)^(p_.))*(d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

---

3.331. 
$$\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$$



**3.331.4 Maple [N/A] (verified)**

Not integrable

Time = 1.64 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{x(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x))^2,x)`output `int((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x))^2,x)`**3.331.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.54

$$\int \frac{(1 - c^2x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arccosh}(cx) + a)^2 x} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`output `integral((-c^2*x^2 + 1)^(3/2)/(b^2*x*arccosh(c*x)^2 + 2*a*b*x*arccosh(c*x) + a^2*x), x)`**3.331.6 Sympy [N/A]**

Not integrable

Time = 20.42 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{(1 - c^2x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-(cx - 1)(cx + 1))^{\frac{3}{2}}}{x(a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/x/(a+b*acosh(c*x))**2,x)`output `Integral((-c*x - 1)*(c*x + 1)**(3/2)/(x*(a + b*acosh(c*x))**2), x)`

---

3.331.  $\int \frac{(1-c^2x^2)^{3/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$

**3.331.7 Maxima [N/A]**

Not integrable

Time = 0.85 (sec) , antiderivative size = 476, normalized size of antiderivative = 17.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)^2 x} dx$$

```
input integrate((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="maxima")
```

```
output ((c^4*x^4 - 2*c^2*x^2 + 1)*(c*x + 1)*sqrt(c*x - 1) + (c^5*x^5 - 2*c^3*x^3
+ c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^3 + sqrt(c*x + 1)*sqrt(c*x
- 1)*a*b*c^2*x^2 - a*b*c*x + (b^2*c^3*x^3 + sqrt(c*x + 1)*sqrt(c*x - 1)*b
^2*c^2*x^2 - b^2*c*x)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate(
((3*c^5*x^5 - c^3*x^3 - 2*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + (6*c^6*x^6 - 7*
c^4*x^4 + 1)*(c*x + 1)*sqrt(c*x - 1) + 3*(c^7*x^7 - 2*c^5*x^5 + c^3*x^3)*s
qrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^6 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^
4 - 2*a*b*c^3*x^4 + a*b*c*x^2 + 2*(a*b*c^4*x^5 - a*b*c^2*x^3)*sqrt(c*x + 1
)*sqrt(c*x - 1) + (b^2*c^5*x^6 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^4 - 2*b^2*c
^3*x^4 + b^2*c*x^2 + 2*(b^2*c^4*x^5 - b^2*c^2*x^3)*sqrt(c*x + 1)*sqrt(c*x
- 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)
```

**3.331.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c^2*x^2+1)^(3/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.331.9 Mupad [N/A]**

Not integrable

Time = 3.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x(a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x*(a + b*acosh(c*x))^2),x)`output `int((1 - c^2*x^2)^(3/2)/(x*(a + b*acosh(c*x))^2), x)`

**3.332**  $\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$

3.332.1 Optimal result . . . . . 2691  
 3.332.2 Mathematica [N/A] . . . . . 2691  
 3.332.3 Rubi [N/A] . . . . . 2692  
 3.332.4 Maple [N/A] (verified) . . . . . 2693  
 3.332.5 Fricas [N/A] . . . . . 2694  
 3.332.6 Sympy [N/A] . . . . . 2694  
 3.332.7 Maxima [N/A] . . . . . 2695  
 3.332.8 Giac [N/A] . . . . . 2695  
 3.332.9 Mupad [N/A] . . . . . 2696

**3.332.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{3/2}}{bcx^2(a+b\operatorname{arccosh}(cx))} - \frac{2\sqrt{1-cx}\operatorname{Int}\left(\frac{-1+c^2x^2}{x^3(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{-1+cx}} - \frac{2c\sqrt{1-cx}\operatorname{Int}\left(\frac{-1+c^2x^2}{x(a+b\operatorname{arccosh}(cx))}, x\right)}{b\sqrt{-1+cx}}$$

output `-(-c^2*x^2+1)^(3/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/x^2/(a+b*arccosh(c*x)) - 2*(-c*x+1)^(1/2)*Unintegrable((c^2*x^2-1)/x^3/(a+b*arccosh(c*x)),x)/b/c/(c*x-1)^(1/2) - 2*c*(-c*x+1)^(1/2)*Unintegrable((c^2*x^2-1)/x/(a+b*arccosh(c*x)),x)/b/(c*x-1)^(1/2)`

**3.332.2 Mathematica [N/A]**

Not integrable

Time = 20.94 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(x^2*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[(1 - c^2*x^2)^(3/2)/(x^2*(a + b*ArcCosh[c*x])^2), x]`

---

3.332.  $\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$

**3.332.3 Rubi [N/A]**

Not integrable

Time = 0.83 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6357, 25, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(1-c^2x^2)^{3/2}}{x^2(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{2\sqrt{1-cx} \int -\frac{(1-cx)(cx+1)}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{2c\sqrt{1-cx} \int -\frac{(1-cx)(cx+1)}{x(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{cx-1}} \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx^2(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{2\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{2c\sqrt{1-cx} \int \frac{(1-cx)(cx+1)}{x(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{cx-1}} \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx^2(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & \frac{2c\sqrt{1-cx} \int \frac{1-c^2x^2}{x(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{cx-1}} + \frac{2\sqrt{1-cx} \int \frac{1-c^2x^2}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx^2(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6375} \\
 & \frac{2c\sqrt{1-cx} \int \frac{1-c^2x^2}{x(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{cx-1}} + \frac{2\sqrt{1-cx} \int \frac{1-c^2x^2}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{3/2}}{bcx^2(a+\operatorname{barccosh}(cx))}
 \end{aligned}$$

input `Int[(1 - c^2*x^2)^(3/2)/(x^2*(a + b*ArcCosh[c*x])^2), x]`

---

3.332.  $\int \frac{(1-c^2x^2)^{3/2}}{x^2(a+\operatorname{barccosh}(cx))^2} dx$

output \$Aborted

### 3.332.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*(m + 2*p + 1)/(b*f*(n + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

### 3.332.4 Maple [N/A] (verified)

Not integrable

Time = 1.38 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{x^2(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x))^2,x)`

### 3.332.5 Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arccosh}(cx) + a)^2 x^2} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((-c^2*x^2 + 1)^(3/2)/(b^2*x^2*arccosh(c*x)^2 + 2*a*b*x^2*arccosh(c*x) + a^2*x^2), x)`

### 3.332.6 Sympy [N/A]

Not integrable

Time = 65.61 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-(cx - 1)(cx + 1))^{\frac{3}{2}}}{x^2 (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/x**2/(a+b*acosh(c*x))**2,x)`

output `Integral((-c*x - 1)*(c*x + 1)**(3/2)/(x**2*(a + b*acosh(c*x))**2), x)`

**3.332.7 Maxima [N/A]**

Not integrable

Time = 0.94 (sec) , antiderivative size = 484, normalized size of antiderivative = 17.29

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

```
input integrate((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="maxima")
```

```
output ((c^4*x^4 - 2*c^2*x^2 + 1)*(c*x + 1)*sqrt(c*x - 1) + (c^5*x^5 - 2*c^3*x^3 + c*x)*sqrt(c*x + 1)*sqrt(-c*x + 1)/(a*b*c^3*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^3 - a*b*c*x^2 + (b^2*c^3*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^3 - b^2*c*x^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate(((2*c^5*x^5 + c^3*x^3 - 3*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(2*c^6*x^6 - c^4*x^4 - 2*c^2*x^2 + 1)*(c*x + 1)*sqrt(c*x - 1) + (2*c^7*x^7 - 3*c^5*x^5 + c*x)*sqrt(c*x + 1)*sqrt(-c*x + 1)/(a*b*c^5*x^7 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^5 - 2*a*b*c^3*x^5 + a*b*c*x^3 + 2*(a*b*c^4*x^6 - a*b*c^2*x^4)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^7 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^5 - 2*b^2*c^3*x^5 + b^2*c*x^3 + 2*(b^2*c^4*x^6 - b^2*c^2*x^4)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)
```

**3.332.8 Giac [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

```
input integrate((-c^2*x^2+1)^(3/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)^2*x^2), x)
```



**3.332.9 Mupad [N/A]**

Not integrable

Time = 2.92 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x^2 (a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x^2*(a + b*acosh(c*x))^2),x)`output `int((1 - c^2*x^2)^(3/2)/(x^2*(a + b*acosh(c*x))^2), x)`

$$3.333 \quad \int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$$

3.333.1 Optimal result	2697
3.333.2 Mathematica [F(-1)]	2697
3.333.3 Rubi [N/A]	2698
3.333.4 Maple [N/A] (verified)	2698
3.333.5 Fricas [N/A]	2699
3.333.6 Sympy [N/A]	2699
3.333.7 Maxima [N/A]	2699
3.333.8 Giac [F(-2)]	2700
3.333.9 Mupad [N/A]	2700

### 3.333.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x))^2,x)`

### 3.333.2 Mathematica [F(-1)]

Timed out.

$$\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx = \$Aborted$$

input `Integrate[(1 - c^2*x^2)^(3/2)/(x^3*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

---


$$3.333. \quad \int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.333.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + \operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + \operatorname{arccosh}(cx))^2} dx$$

input `Int[(1 - c^2*x^2)^(3/2)/(x^3*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.333.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.333.4 Maple [N/A] (verified)**

Not integrable

Time = 1.14 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 x^2 + 1)^{3/2}}{x^3 (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x))^2,x)`

---

3.333.  $\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$

**3.333.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{3/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^3} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((-c^2*x^2 + 1)^(3/2)/(b^2*x^3*arccosh(c*x)^2 + 2*a*b*x^3*arccosh(c*x) + a^2*x^3), x)`

**3.333.6 Sympy [N/A]**

Not integrable

Time = 176.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-(cx - 1)(cx + 1))^{3/2}}{x^3 (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((-c**2*x**2+1)**(3/2)/x**3/(a+b*acosh(c*x))**2,x)`

output `Integral((-c*x - 1)*(c*x + 1)**(3/2)/(x**3*(a + b*acosh(c*x))**2), x)`

**3.333.7 Maxima [N/A]**

Not integrable

Time = 0.92 (sec) , antiderivative size = 483, normalized size of antiderivative = 17.25

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{3/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^3} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `((c^4*x^4 - 2*c^2*x^2 + 1)*(c*x + 1)*sqrt(c*x - 1) + (c^5*x^5 - 2*c^3*x^3 + c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^5 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^4 - a*b*c*x^3 + (b^2*c^3*x^5 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^4 - b^2*c*x^3)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate(((c^5*x^5 + 3*c^3*x^3 - 4*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + (2*c^6*x^6 + 3*c^4*x^4 - 8*c^2*x^2 + 3)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^7 - 3*c^3*x^3 + 2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^8 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^6 - 2*a*b*c^3*x^6 + a*b*c*x^4 + 2*(a*b*c^4*x^7 - a*b*c^2*x^5)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^8 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^6 - 2*b^2*c^3*x^6 + b^2*c*x^4 + 2*(b^2*c^4*x^7 - b^2*c^2*x^5)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.333.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + b \operatorname{arccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(3/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.333.9 Mupad [N/A]

Not integrable

Time = 3.39 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x^3*(a + b*acosh(c*x))^2),x)`

---

3.333.  $\int \frac{(1 - c^2 x^2)^{3/2}}{x^3 (a + b \operatorname{arccosh}(cx))^2} dx$

output `int((1 - c^2*x^2)^(3/2)/(x^3*(a + b*acosh(c*x))^2), x)`

---

3.333.  $\int \frac{(1-c^2x^2)^{3/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$

**3.334**  $\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$

3.334.1 Optimal result	2702
3.334.2 Mathematica [F(-1)]	2702
3.334.3 Rubi [N/A]	2703
3.334.4 Maple [N/A] (verified)	2704
3.334.5 Fricas [N/A]	2705
3.334.6 Sympy [F(-1)]	2705
3.334.7 Maxima [N/A]	2705
3.334.8 Giac [N/A]	2706
3.334.9 Mupad [N/A]	2706

**3.334.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{3/2}}{bcx^4(a+b\operatorname{arccosh}(cx))} - \frac{4\sqrt{1-cx}\operatorname{Int}\left(\frac{-1+c^2x^2}{x^5(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{-1+cx}}$$

```
output -(-c^2*x^2+1)^(3/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/x^4/(a+b*arccosh(c*x))
-4*(-c*x+1)^(1/2)*Unintegrable((c^2*x^2-1)/x^5/(a+b*arccosh(c*x)),x)/b/c/(
c*x-1)^(1/2)
```

**3.334.2 Mathematica [F(-1)]**

Timed out.

$$\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx = \$Aborted$$

```
input Integrate[(1 - c^2*x^2)^(3/2)/(x^4*(a + b*ArcCosh[c*x])^2), x]
```

```
output $Aborted
```

---

3.334.  $\int \frac{(1-c^2x^2)^{3/2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$

**3.334.3 Rubi [N/A]**

Not integrable

Time = 0.65 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6355, 25, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6355} \\
 & - \frac{4\sqrt{1 - cx} \int -\frac{(1-cx)(cx+1)}{x^5(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx - 1}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}(1 - c^2 x^2)^{3/2}}{bcx^4(a + \operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{4\sqrt{1 - cx} \int \frac{(1-cx)(cx+1)}{x^5(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx - 1}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}(1 - c^2 x^2)^{3/2}}{bcx^4(a + \operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & \frac{4\sqrt{1 - cx} \int \frac{1-c^2 x^2}{x^5(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx - 1}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}(1 - c^2 x^2)^{3/2}}{bcx^4(a + \operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6375} \\
 & \frac{4\sqrt{1 - cx} \int \frac{1-c^2 x^2}{x^5(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx - 1}} - \frac{\sqrt{cx - 1}\sqrt{cx + 1}(1 - c^2 x^2)^{3/2}}{bcx^4(a + \operatorname{barccosh}(cx))}
 \end{aligned}$$

input `Int[(1 - c^2*x^2)^(3/2)/(x^4*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`



## 3.334.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6355 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && EqQ[m + 2*p + 1, 0]`
- rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

## 3.334.4 Maple [N/A] (verified)

Not integrable

Time = 1.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{x^4 (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x))^2,x)`

**3.334.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.75

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((-c^2*x^2 + 1)^(3/2)/(b^2*x^4*arccosh(c*x)^2 + 2*a*b*x^4*arccosh(c*x) + a^2*x^4), x)`

**3.334.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(3/2)/x**4/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.334.7 Maxima [N/A]**

Not integrable

Time = 0.77 (sec) , antiderivative size = 469, normalized size of antiderivative = 16.75

$$\int \frac{(1 - c^2 x^2)^{3/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}}}{(b \operatorname{arcosh}(cx) + a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $((c^4x^4 - 2c^2x^2 + 1)(cx + 1)\sqrt{cx - 1} + (c^5x^5 - 2c^3x^3 + cx)\sqrt{cx + 1})\sqrt{-cx + 1}/(abc^3x^6 + \sqrt{cx + 1}\sqrt{cx - 1}) * b^2c^2x^5 - abcx^4 + (b^2c^3x^6 + \sqrt{cx + 1}\sqrt{cx - 1}) * b^2c^2x^5 - b^2cx^4) * \log(cx + \sqrt{cx + 1}\sqrt{cx - 1})) - \text{integrate}((5(c^3x^3 - cx)(cx + 1)^{3/2}(cx - 1) + 4(2c^4x^4 - 3c^2x^2 + 1)(cx + 1)\sqrt{cx - 1} + 3(c^5x^5 - 2c^3x^3 + cx)\sqrt{cx + 1})\sqrt{-cx + 1}/(abc^5x^9 + (cx + 1)(cx - 1)abc^3x^7 - 2ab * c^3x^7 + abcx^5 + 2(abcx^4x^8 - abcx^2x^6)\sqrt{cx + 1}\sqrt{cx - 1} + (b^2c^5x^9 + (cx + 1)(cx - 1)b^2c^3x^7 - 2b^2c^3x^7 + b^2cx^5 + 2(b^2c^4x^8 - b^2c^2x^6)\sqrt{cx + 1}\sqrt{cx - 1})) * \log(cx + \sqrt{cx + 1}\sqrt{cx - 1})), x)$

### 3.334.8 Giac [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2x^2)^{3/2}}{x^4(a + \text{barccosh}(cx))^2} dx = \int \frac{(-c^2x^2 + 1)^{\frac{3}{2}}}{(b \text{arcosh}(cx) + a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(3/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(3/2)/((b*arccosh(c*x) + a)^2*x^4), x)`

### 3.334.9 Mupad [N/A]

Not integrable

Time = 3.52 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2x^2)^{3/2}}{x^4(a + \text{barccosh}(cx))^2} dx = \int \frac{(1 - c^2x^2)^{3/2}}{x^4(a + b \text{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(3/2)/(x^4*(a + b*acosh(c*x))^2),x)`

output `int((1 - c^2*x^2)^(3/2)/(x^4*(a + b*acosh(c*x))^2), x)`

---

3.334.  $\int \frac{(1 - c^2x^2)^{3/2}}{x^4(a + b \text{arccosh}(cx))^2} dx$

$$3.335 \quad \int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

3.335.1 Optimal result . . . . .	2707
3.335.2 Mathematica [A] (verified) . . . . .	2708
3.335.3 Rubi [A] (verified) . . . . .	2709
3.335.4 Maple [A] (verified) . . . . .	2712
3.335.5 Fricas [F] . . . . .	2712
3.335.6 Sympy [F(-1)] . . . . .	2713
3.335.7 Maxima [F] . . . . .	2713
3.335.8 Giac [F] . . . . .	2714
3.335.9 Mupad [F(-1)] . . . . .	2714

### 3.335.1 Optimal result

Integrand size = 28, antiderivative size = 454

$$\begin{aligned} \int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = & -\frac{x^2\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\ & - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)}{16b^2c^3\sqrt{-1+cx}} \\ & - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{4a}{b}\right)}{8b^2c^3\sqrt{-1+cx}} \\ & + \frac{3\sqrt{1-cx}\operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{6a}{b}\right)}{16b^2c^3\sqrt{-1+cx}} \\ & - \frac{\sqrt{1-cx}\operatorname{Chi}\left(\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{8a}{b}\right)}{16b^2c^3\sqrt{-1+cx}} \\ & + \frac{\sqrt{1-cx}\cosh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^3\sqrt{-1+cx}} \\ & + \frac{\sqrt{1-cx}\cosh\left(\frac{4a}{b}\right)\operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8b^2c^3\sqrt{-1+cx}} \\ & - \frac{3\sqrt{1-cx}\cosh\left(\frac{6a}{b}\right)\operatorname{Shi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^3\sqrt{-1+cx}} \\ & + \frac{\sqrt{1-cx}\cosh\left(\frac{8a}{b}\right)\operatorname{Shi}\left(\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^3\sqrt{-1+cx}} \end{aligned}$$

---


$$3.335. \quad \int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

output  $1/16*\cosh(2*a/b)*\text{Shi}(2*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}+1/8*\cosh(4*a/b)*\text{Shi}(4*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-3/16*\cosh(6*a/b)*\text{Shi}(6*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}+1/16*\cosh(8*a/b)*\text{Shi}(8*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-1/16*\text{Chi}(2*(a+b*\text{arccosh}(c*x))/b)*\sinh(2*a/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-1/8*\text{Chi}(4*(a+b*\text{arccosh}(c*x))/b)*\sinh(4*a/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}+3/16*\text{Chi}(6*(a+b*\text{arccosh}(c*x))/b)*\sinh(6*a/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-1/16*\text{Chi}(8*(a+b*\text{arccosh}(c*x))/b)*\sinh(8*a/b)*(-c*x+1)^{(1/2)}/b^2/c^3/(c*x-1)^{(1/2)}-x^2*(-c^2*x^2+1)^{(5/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c/(a+b*\text{arccosh}(c*x))$

### 3.335.2 Mathematica [A] (verified)

Time = 1.33 (sec) , antiderivative size = 446, normalized size of antiderivative = 0.98

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\text{arccosh}(cx))^2} dx = \frac{\sqrt{-1+cx}\sqrt{1+cx}(-16bc^2x^2+48bc^4x^4-48bc^6x^6+16bc^8x^8+(a+b\text{arccosh}(cx))^2)}{(a+b\text{arccosh}(cx))^2}$$

input `Integrate[(x^2*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x])^2,x]`

output  $(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(-16*b*c^2*x^2 + 48*b*c^4*x^4 - 48*b*c^6*x^6 + 16*b*c^8*x^8 + (a + b*\text{ArcCosh}[c*x])*\text{CoshIntegral}[2*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(2*a)/b] + 2*(a + b*\text{ArcCosh}[c*x])*\text{CoshIntegral}[4*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(4*a)/b] - 3*a*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(6*a)/b] - 3*b*\text{ArcCosh}[c*x]*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(6*a)/b] + a*\text{CoshIntegral}[8*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(8*a)/b] + b*\text{ArcCosh}[c*x]*\text{CoshIntegral}[8*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(8*a)/b] - a*\text{Cosh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])] - b*\text{ArcCosh}[c*x]*\text{Cosh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])] - 2*a*\text{Cosh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])] - 2*b*\text{ArcCosh}[c*x]*\text{Cosh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])] + 3*a*\text{Cosh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])] + 3*b*\text{ArcCosh}[c*x]*\text{Cosh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])] - a*\text{Cosh}[(8*a)/b]*\text{SinhIntegral}[8*(a/b + \text{ArcCosh}[c*x])] - b*\text{ArcCosh}[c*x]*\text{Cosh}[(8*a)/b]*\text{SinhIntegral}[8*(a/b + \text{ArcCosh}[c*x])]))/(16*b^2*c^3*\text{Sqrt}[1 - c^2*x^2]*(a + b*\text{ArcCosh}[c*x]))$

---

3.335.  $\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\text{arccosh}(cx))^2} dx$

**3.335.3 Rubi [A] (verified)**

Time = 1.74 (sec) , antiderivative size = 460, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {6357, 6327, 6367, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(1-c^2x^2)^{5/2}}{(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{8c\sqrt{1-cx} \int \frac{x^3(1-cx)^2(cx+1)^2}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{2\sqrt{1-cx} \int \frac{x(1-cx)^2(cx+1)^2}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} - \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & -\frac{2\sqrt{1-cx} \int \frac{x(1-c^2x^2)^2}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{cx-1}} + \frac{8c\sqrt{1-cx} \int \frac{x^3(1-c^2x^2)^2}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6367} \\
 & \frac{8\sqrt{1-cx} \int -\frac{\cosh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^3\sqrt{cx-1}} - \\
 & \frac{2\sqrt{1-cx} \int -\frac{\cosh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^3\sqrt{cx-1}} - \\
 & \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & -\frac{8\sqrt{1-cx} \int \frac{\cosh^3\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^3\sqrt{cx-1}} + \\
 & \frac{2\sqrt{1-cx} \int \frac{\cosh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^3\sqrt{cx-1}} - \\
 & \frac{x^2\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{5971}
 \end{aligned}$$

---

3.335.  $\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{8\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{8a}{b} - \frac{8(a+b\operatorname{arccosh}(cx))}{b}\right)}{128(a+b\operatorname{arccosh}(cx))} - \frac{\sinh\left(\frac{6a}{b} - \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} - \frac{\sinh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} + \frac{3\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} \right)}{2\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{6a}{b} - \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} - \frac{\sinh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} + \frac{5\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32(a+b\operatorname{arccosh}(cx))} \right)} d(a+b\operatorname{arccosh}(cx))} \\
 & \frac{b^2 c^3 \sqrt{cx-1}}{x^2 \sqrt{cx-1} \sqrt{cx+1} (1-c^2 x^2)^{5/2}} \\
 & \frac{bc(a+b\operatorname{arccosh}(cx))}{\downarrow 2009} \\
 & \frac{2\sqrt{1-cx} \left( -\frac{5}{32} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{32} \sinh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{8\sqrt{1-cx} \left( -\frac{3}{64} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{64} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{64} \sinh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) \right)} \\
 & \frac{x^2 \sqrt{cx-1} \sqrt{cx+1} (1-c^2 x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))}
 \end{aligned}$$

input `Int[(x^2*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x])^2,x]`

output `-(x^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(1 - c^2*x^2)^(5/2))/(b*c*(a + b*ArcCosh[c*x])) - (2*sqrt[1 - c*x]*((-5*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]*Sinh[(2*a)/b])/32 + (CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b]*Sinh[(4*a)/b])/8 - (CoshIntegral[(6*(a + b*ArcCosh[c*x]))/b]*Sinh[(6*a)/b])/32 + (5*Cosh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/32 - (Cosh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8 + (Cosh[(6*a)/b]*SinhIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32)/(b^2*c^3*sqrt[-1 + c*x]) + (8*sqrt[1 - c*x]*((-3*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]*Sinh[(2*a)/b])/64 + (CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b]*Sinh[(4*a)/b])/64 + (CoshIntegral[(6*(a + b*ArcCosh[c*x]))/b]*Sinh[(6*a)/b])/64 - (CoshIntegral[(8*(a + b*ArcCosh[c*x]))/b]*Sinh[(8*a)/b])/128 + (3*Cosh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/64 - (Cosh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/64 - (Cosh[(6*a)/b]*SinhIntegral[(6*(a + b*ArcCosh[c*x]))/b])/64 + (Cosh[(8*a)/b]*SinhIntegral[(8*(a + b*ArcCosh[c*x]))/b])/128)/(b^2*c^3*sqrt[-1 + c*x])`

3.335.  $\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

## 3.335.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`
- rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`



### 3.335.4 Maple [A] (verified)

Time = 0.91 (sec) , antiderivative size = 773, normalized size of antiderivative = 1.70

method	result	size
default	Expression too large to display	773

input `int(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output

```

1/32*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(32*(
c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^8*x^8-Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arc
cosh(c*x)+2*a)/b)*b*arccosh(c*x)+arccosh(c*x)*b*Ei(1,-2*arccosh(c*x)-2*a/b
)*exp(-(-b*arccosh(c*x)+2*a)/b)+3*Ei(1,6*arccosh(c*x)+6*a/b)*exp((b*arccos
h(c*x)+6*a)/b)*b*arccosh(c*x)+arccosh(c*x)*b*Ei(1,-8*arccosh(c*x)-8*a/b)*e
xp(-(-b*arccosh(c*x)+8*a)/b)-Ei(1,8*arccosh(c*x)+8*a/b)*exp((b*arccosh(c*x
)+8*a)/b)*b*arccosh(c*x)+2*arccosh(c*x)*b*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(
-(-b*arccosh(c*x)+4*a)/b)-3*arccosh(c*x)*b*Ei(1,-6*arccosh(c*x)-6*a/b)*exp
(-(-b*arccosh(c*x)+6*a)/b)-2*Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x
)+4*a)/b)*b*arccosh(c*x)-96*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^6*x^6+96*x^4*c
^4*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)-32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^2*x^2+
32*b*c^9*x^9-96*b*c^7*x^7-3*a*Ei(1,-6*arccosh(c*x)-6*a/b)*exp(-(-b*arccosh
(c*x)+6*a)/b)+3*Ei(1,6*arccosh(c*x)+6*a/b)*exp((b*arccosh(c*x)+6*a)/b)*a+a
*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-Ei(1,2*arccosh(
c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*a+2*a*Ei(1,-4*arccosh(c*x)-4*a/b)*
exp(-(-b*arccosh(c*x)+4*a)/b)-2*Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(
c*x)+4*a)/b)*a-Ei(1,8*arccosh(c*x)+8*a/b)*exp((b*arccosh(c*x)+8*a)/b)*a+a*
Ei(1,-8*arccosh(c*x)-8*a/b)*exp(-(-b*arccosh(c*x)+8*a)/b)-32*b*c^3*x^3+96*
b*c^5*x^5)/(c*x+1)/c^3/(c*x-1)/b^2/(a+b*arccosh(c*x))

```

### 3.335.5 Fracas [F]

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{5/2}x^2}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output `integral((c^4*x^6 - 2*c^2*x^4 + x^2)*sqrt(-c^2*x^2 + 1)/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

---

3.335.  $\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

**3.335.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \text{Timed out}$$

```
input integrate(x**2*(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)
```

```
output Timed out
```

**3.335.7 Maxima [F]**

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{5/2}x^2}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

```
input integrate(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")
```

```
output -((c^6*x^8 - 3*c^4*x^6 + 3*c^2*x^4 - x^2)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^9 - 3*c^5*x^7 + 3*c^3*x^5 - c*x^3)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((8*c^7*x^8 - 17*c^5*x^6 + 10*c^3*x^4 - c*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(8*c^8*x^9 - 22*c^6*x^7 + 21*c^4*x^5 - 8*c^2*x^3 + x)*(c*x + 1)*sqrt(c*x - 1) + (8*c^9*x^10 - 27*c^7*x^8 + 33*c^5*x^6 - 17*c^3*x^4 + 3*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1))*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)
```

**3.335.8 Giac [F]**

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{5/2}x^2}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^2*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(5/2)*x^2/(b*arccosh(c*x) + a)^2, x)`

**3.335.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(1-c^2x^2)^{5/2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^2(1-c^2x^2)^{5/2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((x^2*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x))^2,x)`

output `int((x^2*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x))^2, x)`

**3.336**  $\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.336.1 Optimal result . . . . . 2715  
 3.336.2 Mathematica [A] (verified) . . . . . 2716  
 3.336.3 Rubi [C] (verified) . . . . . 2717  
 3.336.4 Maple [A] (verified) . . . . . 2722  
 3.336.5 Fricas [F] . . . . . 2722  
 3.336.6 Sympy [F(-1)] . . . . . 2723  
 3.336.7 Maxima [F] . . . . . 2723  
 3.336.8 Giac [F] . . . . . 2724  
 3.336.9 Mupad [F(-1)] . . . . . 2724

**3.336.1 Optimal result**

Integrand size = 26, antiderivative size = 448

$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} + \frac{5\sqrt{1-cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{64b^2c^2\sqrt{-1+cx}} - \frac{27\sqrt{1-cx}\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{64b^2c^2\sqrt{-1+cx}} + \frac{25\sqrt{1-cx}\operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{5a}{b}\right)}{64b^2c^2\sqrt{-1+cx}} - \frac{7\sqrt{1-cx}\operatorname{Chi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{7a}{b}\right)}{64b^2c^2\sqrt{-1+cx}} - \frac{5\sqrt{1-cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64b^2c^2\sqrt{-1+cx}} + \frac{27\sqrt{1-cx}\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64b^2c^2\sqrt{-1+cx}} - \frac{25\sqrt{1-cx}\cosh\left(\frac{5a}{b}\right)\operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64b^2c^2\sqrt{-1+cx}} + \frac{7\sqrt{1-cx}\cosh\left(\frac{7a}{b}\right)\operatorname{Shi}\left(\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64b^2c^2\sqrt{-1+cx}}$$

---

3.336.  $\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

output 
$$\begin{aligned} & -5/64*\cosh(a/b)*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^2/(c*x-1)^{(1/2)} \\ & +27/64*\cosh(3*a/b)*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c^2/ \\ & (c*x-1)^{(1/2)}-25/64*\cosh(5*a/b)*\text{Shi}(5*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)} \\ & /b^2/c^2/(c*x-1)^{(1/2)}+7/64*\cosh(7*a/b)*\text{Shi}(7*(a+b*\text{arccosh}(c*x))/b)*(-c*x+ \\ & 1)^{(1/2)}/b^2/c^2/(c*x-1)^{(1/2)}+5/64*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\sinh(a/b)*(- \\ & c*x+1)^{(1/2)}/b^2/c^2/(c*x-1)^{(1/2)}-27/64*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\sinh( \\ & 3*a/b)*(-c*x+1)^{(1/2)}/b^2/c^2/(c*x-1)^{(1/2)}+25/64*\text{Chi}(5*(a+b*\text{arccosh}(c*x)) \\ & /b)*\sinh(5*a/b)*(-c*x+1)^{(1/2)}/b^2/c^2/(c*x-1)^{(1/2)}-7/64*\text{Chi}(7*(a+b*\text{arcco} \\ & \text{sh}(c*x))/b)*\sinh(7*a/b)*(-c*x+1)^{(1/2)}/b^2/c^2/(c*x-1)^{(1/2)}-x*(-c^2*x^2+1 \\ & )^{(5/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c/(a+b*\text{arccosh}(c*x)) \end{aligned}$$

### 3.336.2 Mathematica [A] (verified)

Time = 1.14 (sec) , antiderivative size = 436, normalized size of antiderivative = 0.97

$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\text{arccosh}(cx))^2} dx = \frac{\sqrt{-1+cx}\sqrt{1+cx}(-64bcx+192bc^3x^3-192bc^5x^5+64bc^7x^7-5(a+b\text{arccosh}(cx)))}{(a+b\text{arccosh}(cx))^2}$$

input `Integrate[(x*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x])^2,x]`

output 
$$\begin{aligned} & (\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(-64*b*c*x + 192*b*c^3*x^3 - 192*b*c^5*x^5 + \\ & 64*b*c^7*x^7 - 5*(a + b*\text{ArcCosh}[c*x])* \text{CoshIntegral}[a/b + \text{ArcCosh}[c*x]]*\text{Si} \\ & \text{nh}[a/b] + 27*(a + b*\text{ArcCosh}[c*x])* \text{CoshIntegral}[3*(a/b + \text{ArcCosh}[c*x])] * \text{Sin} \\ & \text{h}[(3*a)/b] - 25*a*\text{CoshIntegral}[5*(a/b + \text{ArcCosh}[c*x])* \text{Sinh}[(5*a)/b] - 25* \\ & b*\text{ArcCosh}[c*x]* \text{CoshIntegral}[5*(a/b + \text{ArcCosh}[c*x])* \text{Sinh}[(5*a)/b] + 7*a*\text{Co} \\ & \text{shIntegral}[7*(a/b + \text{ArcCosh}[c*x])* \text{Sinh}[(7*a)/b] + 7*b*\text{ArcCosh}[c*x]* \text{CoshIn} \\ & \text{tegral}[7*(a/b + \text{ArcCosh}[c*x])* \text{Sinh}[(7*a)/b] + 5*a*\text{Cosh}[a/b]* \text{SinhIntegral}[ \\ & a/b + \text{ArcCosh}[c*x]] + 5*b*\text{ArcCosh}[c*x]* \text{Cosh}[a/b]* \text{SinhIntegral}[a/b + \text{ArcCos} \\ & \text{h}[c*x]] - 27*a*\text{Cosh}[(3*a)/b]* \text{SinhIntegral}[3*(a/b + \text{ArcCosh}[c*x])] - 27*b*A \\ & \text{rcCosh}[c*x]* \text{Cosh}[(3*a)/b]* \text{SinhIntegral}[3*(a/b + \text{ArcCosh}[c*x])] + 25*a*\text{Cosh} \\ & [(5*a)/b]* \text{SinhIntegral}[5*(a/b + \text{ArcCosh}[c*x])] + 25*b*\text{ArcCosh}[c*x]* \text{Cosh}[(5 \\ & *a)/b]* \text{SinhIntegral}[5*(a/b + \text{ArcCosh}[c*x])] - 7*a*\text{Cosh}[(7*a)/b]* \text{SinhIntegr} \\ & \text{al}[7*(a/b + \text{ArcCosh}[c*x])] - 7*b*\text{ArcCosh}[c*x]* \text{Cosh}[(7*a)/b]* \text{SinhIntegral}[7 \\ & *(a/b + \text{ArcCosh}[c*x])])]/(64*b^2*c^2*\text{Sqrt}[1 - c^2*x^2]*(a + b*\text{ArcCosh}[c*x] \\ & )) \end{aligned}$$

---

3.336. 
$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\text{arccosh}(cx))^2} dx$$

**3.336.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.85 (sec) , antiderivative size = 464, normalized size of antiderivative = 1.04, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6357, 6304, 6321, 25, 3042, 26, 3793, 2009, 6327, 6367, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{bc\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{6304} \\
 & -\frac{\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{a+b\operatorname{arccosh}(cx)} dx}{bc\sqrt{cx-1}} + \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{6321} \\
 & -\frac{\sqrt{1-cx} \int -\frac{\sinh^5\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{\sqrt{1-cx} \int \frac{\sinh^5\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{1-cx} \int -\frac{i \sin\left(\frac{ia}{b}-\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^5}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))}
 \end{aligned}$$

---

3.336.  $\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \downarrow 26 \\
 & - \frac{i\sqrt{1-cx} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^5}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} + \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \quad \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \downarrow 3793 \\
 & \frac{i\sqrt{1-cx} \int \left( \frac{i \sinh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{5i \sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} + \frac{5i \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{cx-1}} \\
 & \quad - \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \downarrow 2009 \\
 & \frac{7c\sqrt{1-cx} \int \frac{x^2(1-cx)^2(cx+1)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \frac{i\sqrt{1-cx} \left( \frac{5}{8}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16}i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2} \\
 & \quad - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \downarrow 6327 \\
 & \frac{7c\sqrt{1-cx} \int \frac{x^2(1-c^2x^2)^2}{a+b\operatorname{arccosh}(cx)} dx}{b\sqrt{cx-1}} - \\
 & \frac{i\sqrt{1-cx} \left( \frac{5}{8}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16}i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2} \\
 & \quad - \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \downarrow 6367
 \end{aligned}$$

---

3.336.  $\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{7\sqrt{1-cx} \int -\frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} \\
 & \frac{i\sqrt{1-cx} \left(\frac{5}{8} i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16} i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16} i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{7\sqrt{1-cx} \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2 \sqrt{cx-1}} \\
 & \frac{i\sqrt{1-cx} \left(\frac{5}{8} i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16} i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16} i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{5971} \\
 & \frac{7\sqrt{1-cx} \int \left(\frac{\sinh\left(\frac{7a}{b} - \frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} - \frac{3 \sinh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64(a+b\operatorname{arccosh}(cx))} + \frac{5 \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64(a+b\operatorname{arccosh}(cx))}\right)}{b^2 c^2 \sqrt{cx-1}} \\
 & \frac{i\sqrt{1-cx} \left(\frac{5}{8} i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16} i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16} i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{2009} \\
 & \frac{i\sqrt{1-cx} \left(\frac{5}{8} i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16} i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16} i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2} \\
 & \frac{7\sqrt{1-cx} \left(-\frac{5}{64} \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{64} \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{3}{64} \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\right)}{b^2} \\
 & \frac{x\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))}
 \end{aligned}$$

input `Int[(x*(1 - c^2*x^2)^(5/2))/(a + b*ArcCosh[c*x])^2,x]`

3.336.  $\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$



```
output -((x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(1 - c^2*x^2)^(5/2))/(b*c*(a + b*ArcCosh
[c*x]))) - (I*Sqrt[1 - c*x]*(((5*I)/8)*CoshIntegral[(a + b*ArcCosh[c*x])/b
]*Sinh[a/b] - ((5*I)/16)*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*
a)/b] + (I/16)*CoshIntegral[(5*(a + b*ArcCosh[c*x])/b]*Sinh[(5*a)/b] - ((
5*I)/8)*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b] + ((5*I)/16)*Cosh[(
3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b] - (I/16)*Cosh[(5*a)/b]*Si
nhIntegral[(5*(a + b*ArcCosh[c*x])/b]))/(b^2*c^2*Sqrt[-1 + c*x]) + (7*Sqr
t[1 - c*x]*((-5*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b])/64 - (Cosh
Integral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*a)/b])/64 + (3*CoshIntegral[(
5*(a + b*ArcCosh[c*x])/b]*Sinh[(5*a)/b])/64 - (CoshIntegral[(7*(a + b*Arc
Cosh[c*x])/b]*Sinh[(7*a)/b])/64 + (5*Cosh[a/b]*SinhIntegral[(a + b*ArcCos
h[c*x])/b])/64 + (Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/
64 - (3*Cosh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/64 + (Cosh
[(7*a)/b]*SinhIntegral[(7*(a + b*ArcCosh[c*x])/b])/64))/(b^2*c^2*Sqrt[-1
+ c*x])
```

### 3.336.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3793 Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

```
rule 5971 Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

---

3.336. 
$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

---

3.336. 
$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.336.4 Maple [A] (verified)**

Time = 0.84 (sec) , antiderivative size = 759, normalized size of antiderivative = 1.69

method	result	size
default	Expression too large to display	759

```
input int(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output -1/128*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(38
4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^5*x^5-384*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*
c^3*x^3+128*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c*x-128*(c*x-1)^(1/2)*(c*x+1)^(1
/2)*b*c^7*x^7+384*b*c^6*x^6+5*a*Ei(1,-arccosh(c*x)-a/b)*exp(-(b*arccosh(c
*x)+a)/b)+25*a*Ei(1,-5*arccosh(c*x)-5*a/b)*exp(-(b*arccosh(c*x)+5*a)/b)-2
7*a*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(b*arccosh(c*x)+3*a)/b)-7*a*Ei(1,-7*
arccosh(c*x)-7*a/b)*exp(-(b*arccosh(c*x)+7*a)/b)+7*Ei(1,7*arccosh(c*x)+7*
a/b)*exp((b*arccosh(c*x)+7*a)/b)*a-25*Ei(1,5*arccosh(c*x)+5*a/b)*exp((b*ar
ccosh(c*x)+5*a)/b)*a+27*Ei(1,3*arccosh(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a
)/b)*a-5*Ei(1,arccosh(c*x)+a/b)*exp((a+b*arccosh(c*x))/b)*a+5*arccosh(c*x)
*b*Ei(1,-arccosh(c*x)-a/b)*exp(-(b*arccosh(c*x)+a)/b)+25*arccosh(c*x)*b*E
i(1,-5*arccosh(c*x)-5*a/b)*exp(-(b*arccosh(c*x)+5*a)/b)+7*Ei(1,7*arccosh(
c*x)+7*a/b)*exp((b*arccosh(c*x)+7*a)/b)*b*arccosh(c*x)-27*arccosh(c*x)*b*E
i(1,-3*arccosh(c*x)-3*a/b)*exp(-(b*arccosh(c*x)+3*a)/b)-7*arccosh(c*x)*b*
Ei(1,-7*arccosh(c*x)-7*a/b)*exp(-(b*arccosh(c*x)+7*a)/b)-25*Ei(1,5*arccos
h(c*x)+5*a/b)*exp((b*arccosh(c*x)+5*a)/b)*b*arccosh(c*x)+27*Ei(1,3*arccosh
(c*x)+3*a/b)*exp((b*arccosh(c*x)+3*a)/b)*b*arccosh(c*x)-5*Ei(1,arccosh(c*x
)+a/b)*exp((a+b*arccosh(c*x))/b)*b*arccosh(c*x)-384*b*c^4*x^4+128*b*c^2*x^
2-128*b*c^8*x^8)/(c*x+1)/c^2/(c*x-1)/b^2/(a+b*arccosh(c*x))
```

**3.336.5 Fracas [F]**

$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{5/2}x}{(b\operatorname{arccosh}(cx)+a)^2} dx$$

```
input integrate(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")
```

```
output integral((c^4*x^5 - 2*c^2*x^3 + x)*sqrt(-c^2*x^2 + 1)/(b^2*arccosh(c*x)^2
+ 2*a*b*arccosh(c*x) + a^2), x)
```

---

3.336.  $\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

**3.336.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x(1 - c^2x^2)^{5/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(x*(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.336.7 Maxima [F]**

$$\int \frac{x(1 - c^2x^2)^{5/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2 + 1)^{5/2}x}{(b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^6*x^7 - 3*c^4*x^5 + 3*c^2*x^3 - x)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^8 - 3*c^5*x^6 + 3*c^3*x^4 - c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate((7*(c^7*x^7 - 2*c^5*x^5 + c^3*x^3)*(c*x + 1)^(3/2)*(c*x - 1) + (14*c^8*x^8 - 37*c^6*x^6 + 33*c^4*x^4 - 11*c^2*x^2 + 1)*(c*x + 1)*sqrt(c*x - 1) + (7*c^9*x^9 - 23*c^7*x^7 + 27*c^5*x^5 - 13*c^3*x^3 + 2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.336.8 Giac [F]**

$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2+1)^{5/2}x}{(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x*(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(5/2)*x/(b*arccosh(c*x) + a)^2, x)`

**3.336.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x(1-c^2x^2)^{5/2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

input `int((x*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x))^2,x)`

output `int((x*(1 - c^2*x^2)^(5/2))/(a + b*acosh(c*x))^2, x)`

**3.337** 
$$\int \frac{(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

3.337.1 Optimal result . . . . .	2725
3.337.2 Mathematica [A] (verified) . . . . .	2726
3.337.3 Rubi [A] (verified) . . . . .	2726
3.337.4 Maple [B] (verified) . . . . .	2729
3.337.5 Fricas [F] . . . . .	2730
3.337.6 Sympy [F(-1)] . . . . .	2730
3.337.7 Maxima [F] . . . . .	2730
3.337.8 Giac [F] . . . . .	2731
3.337.9 Mupad [F(-1)] . . . . .	2731

**3.337.1 Optimal result**

Integrand size = 25, antiderivative size = 351

$$\int \frac{(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{5/2}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{15\sqrt{1-cx}\operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)}{16b^2c\sqrt{-1+cx}} + \frac{3\sqrt{1-cx}\operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{4a}{b}\right)}{4b^2c\sqrt{-1+cx}} - \frac{3\sqrt{1-cx}\operatorname{Chi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{6a}{b}\right)}{16b^2c\sqrt{-1+cx}} + \frac{15\sqrt{1-cx}\cosh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c\sqrt{-1+cx}} - \frac{3\sqrt{1-cx}\cosh\left(\frac{4a}{b}\right)\operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b^2c\sqrt{-1+cx}} + \frac{3\sqrt{1-cx}\cosh\left(\frac{6a}{b}\right)\operatorname{Shi}\left(\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c\sqrt{-1+cx}}$$

---

3.337. 
$$\int \frac{(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

output  $15/16*\cosh(2*a/b)*\text{Shi}(2*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c/(c*x-1)^{(1/2)}-3/4*\cosh(4*a/b)*\text{Shi}(4*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c/(c*x-1)^{(1/2)}+3/16*\cosh(6*a/b)*\text{Shi}(6*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/c/(c*x-1)^{(1/2)}-15/16*\text{Chi}(2*(a+b*\text{arccosh}(c*x))/b)*\sinh(2*a/b)*(-c*x+1)^{(1/2)}/b^2/c/(c*x-1)^{(1/2)}+3/4*\text{Chi}(4*(a+b*\text{arccosh}(c*x))/b)*\sinh(4*a/b)*(-c*x+1)^{(1/2)}/b^2/c/(c*x-1)^{(1/2)}-3/16*\text{Chi}(6*(a+b*\text{arccosh}(c*x))/b)*\sinh(6*a/b)*(-c*x+1)^{(1/2)}/b^2/c/(c*x-1)^{(1/2)}-(-c^2*x^2+1)^{(5/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c/(a+b*\text{arccosh}(c*x))$

### 3.337.2 Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 343, normalized size of antiderivative = 0.98

$$\int \frac{(1 - c^2 x^2)^{5/2}}{(a + b \text{arccosh}(cx))^2} dx = \frac{\sqrt{-1 + cx} \sqrt{1 + cx} (-16b + 48bc^2 x^2 - 48bc^4 x^4 + 16bc^6 x^6 + 15(a + b \text{arccosh}(cx)))}{(a + b \text{arccosh}(cx))^2}$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(a + b*ArcCosh[c*x])^2,x]`

output  $(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(-16*b + 48*b*c^2*x^2 - 48*b*c^4*x^4 + 16*b*c^6*x^6 + 15*(a + b*\text{ArcCosh}[c*x]))*\text{CoshIntegral}[2*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(2*a)/b] - 12*(a + b*\text{ArcCosh}[c*x])* \text{CoshIntegral}[4*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(4*a)/b] + 3*a*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(6*a)/b] + 3*b*\text{ArcCosh}[c*x]*\text{CoshIntegral}[6*(a/b + \text{ArcCosh}[c*x])]*\text{Sinh}[(6*a)/b] - 15*a*\text{Cosh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])] - 15*b*\text{ArcCosh}[c*x]*\text{Cosh}[(2*a)/b]*\text{SinhIntegral}[2*(a/b + \text{ArcCosh}[c*x])] + 12*a*\text{Cosh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])] + 12*b*\text{ArcCosh}[c*x]*\text{Cosh}[(4*a)/b]*\text{SinhIntegral}[4*(a/b + \text{ArcCosh}[c*x])] - 3*a*\text{Cosh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])] - 3*b*\text{ArcCosh}[c*x]*\text{Cosh}[(6*a)/b]*\text{SinhIntegral}[6*(a/b + \text{ArcCosh}[c*x])])/(16*b^2*c*\text{Sqrt}[1 - c^2*x^2]*(a + b*\text{ArcCosh}[c*x]))$

### 3.337.3 Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.65, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.240$ , Rules used = {6319, 6327, 6367, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.337.  $\int \frac{(1-c^2x^2)^{5/2}}{(a+b\text{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \int \frac{(1-c^2x^2)^{5/2}}{(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{6c\sqrt{1-cx} \int \frac{x(1-cx)^2(cx+1)^2}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & \frac{6c\sqrt{1-cx} \int \frac{x(1-c^2x^2)^2}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6367} \\
 & \frac{6\sqrt{1-cx} \int -\frac{\cosh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c\sqrt{cx-1}} - \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{6\sqrt{1-cx} \int -\frac{\cosh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c\sqrt{cx-1}} - \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{5971} \\
 & \frac{6\sqrt{1-cx} \int \left( \frac{\sinh\left(\frac{6a}{b}-\frac{6(a+\operatorname{barccosh}(cx))}{b}\right)}{32(a+\operatorname{barccosh}(cx))} - \frac{\sinh\left(\frac{4a}{b}-\frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{8(a+\operatorname{barccosh}(cx))} + \frac{5 \sinh\left(\frac{2a}{b}-\frac{2(a+\operatorname{barccosh}(cx))}{b}\right)}{32(a+\operatorname{barccosh}(cx))} \right) d(a+\operatorname{barccosh}(cx))}{b^2c\sqrt{cx-1}} \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{2009} \\
 & \frac{6\sqrt{1-cx} \left( -\frac{5}{32} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) + \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+\operatorname{barccosh}(cx))}{b}\right) - \frac{1}{32} \sinh\left(\frac{6a}{b}\right) \operatorname{Chi}\left(\frac{6(a+\operatorname{barccosh}(cx))}{b}\right) \right)}{b^2c\sqrt{cx-1}} \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bc(a+\operatorname{barccosh}(cx))}
 \end{aligned}$$

---

3.337.  $\int \frac{(1-c^2x^2)^{5/2}}{(a+\operatorname{barccosh}(cx))^2} dx$



input `Int[(1 - c^2*x^2)^(5/2)/(a + b*ArcCosh[c*x])^2,x]`

output `-((Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(1 - c^2*x^2)^(5/2))/(b*c*(a + b*ArcCosh[c*x]))) + (6*Sqrt[1 - c*x]*((-5*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]*Sinh[(2*a)/b])/32 + (CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b]*Sinh[(4*a)/b])/8 - (CoshIntegral[(6*(a + b*ArcCosh[c*x]))/b]*Sinh[(6*a)/b])/32 + (5*Cosh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/32 - (Cosh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8 + (Cosh[(6*a)/b]*SinhIntegral[(6*(a + b*ArcCosh[c*x]))/b])/32))/(b^2*c*Sqrt[-1 + c*x])`

### 3.337.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*((d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

---

3.337.  $\int \frac{(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.337.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 622 vs. 2(309) = 618.

Time = 0.95 (sec) , antiderivative size = 623, normalized size of antiderivative = 1.77

method	result
default	$-\frac{\sqrt{-c^2x^2+1}(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-32\sqrt{cx-1}\sqrt{cx+1}bc^6x^6-32bc^7x^7+96x^4c^4b\sqrt{cx-1}\sqrt{cx+1}+96bc^5x^5-96\sqrt{cx-1}\sqrt{cx+1}\right)}{\dots}$

input `int((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output `-1/32*(-c^2*x^2+1)^(1/2)*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-32*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^6*x^6-32*b*c^7*x^7+96*x^4*c^4*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)+96*b*c^5*x^5-96*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^2*x^2-96*b*c^3*x^3+12*arccosh(c*x)*b*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b)-15*arccosh(c*x)*b*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-3*arccosh(c*x)*b*Ei(1,-6*arccosh(c*x)-6*a/b)*exp(-(-b*arccosh(c*x)+6*a)/b)+3*Ei(1,6*arccosh(c*x)+6*a/b)*exp((b*arccosh(c*x)+6*a)/b)*b*arccosh(c*x)-12*Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)*b*arccosh(c*x)+15*Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*b*arccosh(c*x)+32*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)+12*a*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(-b*arccosh(c*x)+4*a)/b)-15*a*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(-b*arccosh(c*x)+2*a)/b)-3*a*Ei(1,-6*arccosh(c*x)-6*a/b)*exp(-(-b*arccosh(c*x)+6*a)/b)+3*Ei(1,6*arccosh(c*x)+6*a/b)*exp((b*arccosh(c*x)+6*a)/b)*a-12*Ei(1,4*arccosh(c*x)+4*a/b)*exp((b*arccosh(c*x)+4*a)/b)*a+15*Ei(1,2*arccosh(c*x)+2*a/b)*exp((b*arccosh(c*x)+2*a)/b)*a+32*b*c*x/(c*x-1)/(c*x+1)/c/b^2/(a+b*arccosh(c*x))`

$$3.337. \int \frac{(1-c^2x^2)^{5/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.337.5 Fricas [F]**

$$\int \frac{(1 - c^2 x^2)^{5/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

**3.337.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.337.7 Maxima [F]**

$$\int \frac{(1 - c^2 x^2)^{5/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  `-((c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^7 - 3*c^5*x^5 + 3*c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((6*c^6*x^6 - 11*c^4*x^4 + 4*c^2*x^2 + 1)*(c*x + 1)^(3/2)*(c*x - 1) + 6*(2*c^7*x^7 - 5*c^5*x^5 + 4*c^3*x^3 - c*x)*(c*x + 1)*sqrt(c*x - 1) + (6*c^8*x^8 - 19*c^6*x^6 + 21*c^4*x^4 - 9*c^2*x^2 + 1)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^4*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^2*x^2 - 2*a*b*c^2*x^2 + 2*(a*b*c^3*x^3 - a*b*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + a*b + (b^2*c^4*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^2*x^2 - 2*b^2*c^2*x^2 + 2*(b^2*c^3*x^3 - b^2*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + b^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.337.8 Giac [F]

$$\int \frac{(1 - c^2 x^2)^{5/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(5/2)/(b*arccosh(c*x) + a)^2, x)`

### 3.337.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(5/2)/(a + b*acosh(c*x))^2,x)`

output `int((1 - c^2*x^2)^(5/2)/(a + b*acosh(c*x))^2, x)`

**3.338** 
$$\int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$$

3.338.1 Optimal result . . . . .	2732
3.338.2 Mathematica [N/A] . . . . .	2733
3.338.3 Rubi [N/A] . . . . .	2733
3.338.4 Maple [N/A] (verified) . . . . .	2737
3.338.5 Fricas [N/A] . . . . .	2737
3.338.6 Sympy [F(-1)] . . . . .	2738
3.338.7 Maxima [N/A] . . . . .	2738
3.338.8 Giac [F(-2)] . . . . .	2739
3.338.9 Mupad [N/A] . . . . .	2739

**3.338.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\begin{aligned} \int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx = & -\frac{\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{5/2}}{bcx(a+b\operatorname{arccosh}(cx))} \\ & -\frac{25\sqrt{1-cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{8b^2\sqrt{-1+cx}} \\ & +\frac{25\sqrt{1-cx}\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{16b^2\sqrt{-1+cx}} \\ & -\frac{5\sqrt{1-cx}\operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{5a}{b}\right)}{16b^2\sqrt{-1+cx}} \\ & +\frac{25\sqrt{1-cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b^2\sqrt{-1+cx}} \\ & -\frac{25\sqrt{1-cx}\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2\sqrt{-1+cx}} \\ & +\frac{5\sqrt{1-cx}\cosh\left(\frac{5a}{b}\right)\operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2\sqrt{-1+cx}} + \frac{\sqrt{1-cx}\operatorname{Int}\left(\frac{(-1+c^2x^2)^2}{x^2(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{-1+cx}} \end{aligned}$$

---

3.338. 
$$\int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$$

output  $25/8*\cosh(a/b)*\text{Shi}((a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/(c*x-1)^{(1/2)}-25/16*\cosh(3*a/b)*\text{Shi}(3*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/(c*x-1)^{(1/2)}+5/16*\cosh(5*a/b)*\text{Shi}(5*(a+b*\text{arccosh}(c*x))/b)*(-c*x+1)^{(1/2)}/b^2/(c*x-1)^{(1/2)}-25/8*\text{Chi}((a+b*\text{arccosh}(c*x))/b)*\sinh(a/b)*(-c*x+1)^{(1/2)}/b^2/(c*x-1)^{(1/2)}+25/16*\text{Chi}(3*(a+b*\text{arccosh}(c*x))/b)*\sinh(3*a/b)*(-c*x+1)^{(1/2)}/b^2/(c*x-1)^{(1/2)}-5/16*\text{Chi}(5*(a+b*\text{arccosh}(c*x))/b)*\sinh(5*a/b)*(-c*x+1)^{(1/2)}/b^2/(c*x-1)^{(1/2)}-(-c^2*x^2+1)^{(5/2)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/c/x/(a+b*\text{arccosh}(c*x))+(-c*x+1)^{(1/2)}*\text{Unintegrable}((c^2*x^2-1)^2/x^2/(a+b*\text{arccosh}(c*x)),x)/b/c/(c*x-1)^{(1/2)}$

### 3.338.2 Mathematica [N/A]

Not integrable

Time = 16.72 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \text{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \text{arccosh}(cx))^2} dx$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[(1 - c^2*x^2)^(5/2)/(x*(a + b*ArcCosh[c*x])^2), x]`

### 3.338.3 Rubi [N/A]

Not integrable

Time = 2.37 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6357, 6304, 6321, 25, 3042, 26, 3793, 2009, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \text{arccosh}(cx))^2} dx$$

↓ 6357

---

3.338.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \text{arccosh}(cx))^2} dx$

$$\frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{5c\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 6304

$$\frac{5c\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{a+\operatorname{barccosh}(cx)} dx}{b\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 6321

$$\frac{5\sqrt{1-cx} \int -\frac{\sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 25

$$-\frac{5\sqrt{1-cx} \int \frac{\sinh^5\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 3042

$$-\frac{5\sqrt{1-cx} \int -\frac{i \sin\left(\frac{ia}{b}-\frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^5}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 26

$$\frac{5i\sqrt{1-cx} \int \frac{\sin\left(\frac{ia}{b}-\frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^5}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2\sqrt{cx-1}} + \frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+\operatorname{barccosh}(cx))}$$

↓ 3793

---

3.338.  $\int \frac{(1-c^2x^2)^{5/2}}{x(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
 & \frac{5i\sqrt{1-cx} \int \left( \frac{i \sinh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} - \frac{5i \sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16(a+b\operatorname{arccosh}(cx))} + \frac{5i \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{\frac{\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} - \frac{b^2\sqrt{cx-1}}{bcx(a+b\operatorname{arccosh}(cx))} \frac{(1-c^2x^2)^{5/2}}{\sqrt{cx-1}\sqrt{cx+1}}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{5i\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} + \\
 & \frac{5i\sqrt{1-cx} \left( \frac{5}{8}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16}i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+b\operatorname{arccosh}(cx))}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{5i\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} + \\
 & \frac{5i\sqrt{1-cx} \left( \frac{5}{8}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16}i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+b\operatorname{arccosh}(cx))}} \\
 & \quad \downarrow \text{6375} \\
 & \frac{5i\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{x^2(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{cx-1}} + \\
 & \frac{5i\sqrt{1-cx} \left( \frac{5}{8}i \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{5}{16}i \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{16}i \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{\frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx(a+b\operatorname{arccosh}(cx))}}
 \end{aligned}$$

input `Int[(1 - c^2*x^2)^(5/2)/(x*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

---

3.338.  $\int \frac{(1-c^2x^2)^{5/2}}{x(a+b\operatorname{arccosh}(cx))^2} dx$



## 3.338.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`
- rule 6304 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

### 3.338.4 Maple [N/A] (verified)

Not integrable

Time = 0.96 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{5/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x))^2,x)`

### 3.338.5 Fracas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.14

$$\int \frac{(1 - c^2x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2x^2 + 1)^{5/2}}{(b \operatorname{arccosh}(cx) + a)^2 x} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

---

3.338.  $\int \frac{(1 - c^2x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx$

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b^2*x*arccosh(c*x)^2 + 2*a*b*x*arccosh(c*x) + a^2*x), x)`

### 3.338.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x/(a+b*acosh(c*x))**2,x)`

output `Timed out`

### 3.338.7 Maxima [N/A]

Not integrable

Time = 1.10 (sec) , antiderivative size = 524, normalized size of antiderivative = 18.71

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arccosh}(cx) + a)^2 x} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-((c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^7 - 3*c^5*x^5 + 3*c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^3 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^2 - a*b*c*x + (b^2*c^3*x^3 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^2 - b^2*c*x)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((5*c^7*x^7 - 8*c^5*x^5 + c^3*x^3 + 2*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + (10*c^8*x^8 - 23*c^6*x^6 + 15*c^4*x^4 - c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + 5*(c^9*x^9 - 3*c^7*x^7 + 3*c^5*x^5 - c^3*x^3)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^6 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^4 - 2*a*b*c^3*x^4 + a*b*c*x^2 + 2*(a*b*c^4*x^5 - a*b*c^2*x^3)*sqrt(c*x + 1))*sqrt(c*x - 1) + (b^2*c^5*x^6 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^4 - 2*b^2*c^3*x^4 + b^2*c*x^2 + 2*(b^2*c^4*x^5 - b^2*c^2*x^3)*sqrt(c*x + 1))*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.338.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*x^2+1)^(5/2)/x/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.338.9 Mupad [N/A]**

Not integrable

Time = 3.46 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x(a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(5/2)/(x*(a + b*acosh(c*x))^2),x)`

output `int((1 - c^2*x^2)^(5/2)/(x*(a + b*acosh(c*x))^2), x)`

**3.339**  $\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$

3.339.1 Optimal result	2740
3.339.2 Mathematica [N/A]	2740
3.339.3 Rubi [N/A]	2741
3.339.4 Maple [N/A] (verified)	2742
3.339.5 Fricas [N/A]	2743
3.339.6 Sympy [F(-1)]	2743
3.339.7 Maxima [N/A]	2743
3.339.8 Giac [N/A]	2744
3.339.9 Mupad [N/A]	2744

**3.339.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}(1-c^2x^2)^{5/2}}{bcx^2(a+b\operatorname{arccosh}(cx))} + \frac{2\sqrt{1-cx}\operatorname{Int}\left(\frac{(-1+c^2x^2)^2}{x^3(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{-1+cx}} + \frac{4c\sqrt{1-cx}\operatorname{Int}\left(\frac{(-1+c^2x^2)^2}{x(a+b\operatorname{arccosh}(cx))}, x\right)}{b\sqrt{-1+cx}}$$

output `-(-c^2*x^2+1)^(5/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/x^2/(a+b*arccosh(c*x)) + 2*(-c*x+1)^(1/2)*Unintegrable((c^2*x^2-1)^2/x^3/(a+b*arccosh(c*x)),x)/b/c / (c*x-1)^(1/2)+ 4*c*(-c*x+1)^(1/2)*Unintegrable((c^2*x^2-1)^2/x/(a+b*arccosh(c*x)),x)/b/(c*x-1)^(1/2)`

**3.339.2 Mathematica [N/A]**

Not integrable

Time = 23.02 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x^2*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[(1 - c^2*x^2)^(5/2)/(x^2*(a + b*ArcCosh[c*x])^2), x]`

---

3.339.  $\int \frac{(1-c^2x^2)^{5/2}}{x^2(a+b\operatorname{arccosh}(cx))^2} dx$

**3.339.3 Rubi [N/A]**

Not integrable

Time = 1.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6357, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(1-c^2x^2)^{5/2}}{x^2(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{2\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} + \frac{4c\sqrt{1-cx} \int \frac{(1-cx)^2(cx+1)^2}{x(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{cx-1}} - \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx^2(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & \frac{4c\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{x(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{cx-1}} + \frac{2\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx^2(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6375} \\
 & \frac{4c\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{x(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{cx-1}} + \frac{2\sqrt{1-cx} \int \frac{(1-c^2x^2)^2}{x^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{cx-1}} - \\
 & \quad \frac{\sqrt{cx-1}\sqrt{cx+1}(1-c^2x^2)^{5/2}}{bcx^2(a+\operatorname{barccosh}(cx))}
 \end{aligned}$$

input `Int[(1 - c^2*x^2)^(5/2)/(x^2*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

## 3.339.3.1 Defintions of rubi rules used

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

## 3.339.4 Maple [N/A] (verified)

Not integrable

Time = 0.85 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2x^2 + 1)^{5/2}}{x^2(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x))^2,x)`

**3.339.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.36

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b^2*x^2*arccosh(c*x))^2 + 2*a*b*x^2*arccosh(c*x) + a^2*x^2), x)`

**3.339.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x**2/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.339.7 Maxima [N/A]**

Not integrable

Time = 1.16 (sec) , antiderivative size = 534, normalized size of antiderivative = 19.07

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`



output  `-((c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^7 - 3*c^5*x^5 + 3*c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^3 - a*b*c*x^2 + (b^2*c^3*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^3 - b^2*c*x^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((4*c^7*x^7 - 5*c^5*x^5 - 2*c^3*x^3 + 3*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(4*c^8*x^8 - 8*c^6*x^6 + 3*c^4*x^4 + 2*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (4*c^9*x^9 - 11*c^7*x^7 + 9*c^5*x^5 - c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^7 + (c*x - 1)*a*b*c^3*x^5 - 2*a*b*c^3*x^5 + a*b*c*x^3 + 2*(a*b*c^4*x^6 - a*b*c^2*x^4)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^7 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^5 - 2*b^2*c^3*x^5 + b^2*c*x^3 + 2*(b^2*c^4*x^6 - b^2*c^2*x^4)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.339.8 Giac [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^2/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)^2*x^2), x)`

### 3.339.9 Mupad [N/A]

Not integrable

Time = 3.33 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(5/2)/(x^2*(a + b*acosh(c*x))^2),x)`

output `int((1 - c^2*x^2)^(5/2)/(x^2*(a + b*acosh(c*x))^2), x)`

---

3.339.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x^2 (a + b \operatorname{arccosh}(cx))^2} dx$

$$3.340 \quad \int \frac{(1-c^2x^2)^{5/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$$

3.340.1 Optimal result	2745
3.340.2 Mathematica [N/A]	2745
3.340.3 Rubi [N/A]	2746
3.340.4 Maple [N/A] (verified)	2746
3.340.5 Fricas [N/A]	2747
3.340.6 Sympy [F(-1)]	2747
3.340.7 Maxima [N/A]	2747
3.340.8 Giac [F(-2)]	2748
3.340.9 Mupad [N/A]	2748

### 3.340.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{(1-c^2x^2)^{5/2}}{x^3(a+\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x))^2,x)`

### 3.340.2 Mathematica [N/A]

Not integrable

Time = 165.75 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+\operatorname{arccosh}(cx))^2} dx = \int \frac{(1-c^2x^2)^{5/2}}{x^3(a+\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x^3*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[(1 - c^2*x^2)^(5/2)/(x^3*(a + b*ArcCosh[c*x])^2), x]`

---


$$3.340. \quad \int \frac{(1-c^2x^2)^{5/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.340.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{arccosh}(cx))^2} dx$$

input `Int[(1 - c^2*x^2)^(5/2)/(x^3*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.340.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.340.4 Maple [N/A] (verified)**

Not integrable

Time = 1.72 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 x^2 + 1)^{5/2}}{x^3 (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x))^2,x)`

---

3.340.  $\int \frac{(1-c^2x^2)^{5/2}}{x^3(a+b\operatorname{arccosh}(cx))^2} dx$

**3.340.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.36

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^3} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b^2*x^3*arccosh(c*x))^2 + 2*a*b*x^3*arccosh(c*x) + a^2*x^3), x)`

**3.340.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x**3/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.340.7 Maxima [N/A]**

Not integrable

Time = 1.15 (sec) , antiderivative size = 534, normalized size of antiderivative = 19.07

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^3} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

```
output -((c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^7
- 3*c^5*x^5 + 3*c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^5
+ sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^4 - a*b*c*x^3 + (b^2*c^3*x^5 + sq
rt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^4 - b^2*c*x^3)*log(c*x + sqrt(c*x + 1)
*sqrt(c*x - 1))) + integrate(((3*c^7*x^7 - 2*c^5*x^5 - 5*c^3*x^3 + 4*c*x)*
(c*x + 1)^(3/2)*(c*x - 1) + 3*(2*c^8*x^8 - 3*c^6*x^6 - c^4*x^4 + 3*c^2*x^2
- 1)*(c*x + 1)*sqrt(c*x - 1) + (3*c^9*x^9 - 7*c^7*x^7 + 3*c^5*x^5 + 3*c^3
*x^3 - 2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^8 + (c*x + 1)*(c*x
- 1)*a*b*c^3*x^6 - 2*a*b*c^3*x^6 + a*b*c*x^4 + 2*(a*b*c^4*x^7 - a*b*c^2*x^
5)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^8 + (c*x + 1)*(c*x - 1)*b^2*c^
3*x^6 - 2*b^2*c^3*x^6 + b^2*c*x^4 + 2*(b^2*c^4*x^7 - b^2*c^2*x^5)*sqrt(c*x
+ 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)
```

### 3.340.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c^2*x^2+1)^(5/2)/x^3/(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.340.9 Mupad [N/A]

Not integrable

Time = 3.37 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + b \operatorname{acosh}(cx))^2} dx$$

```
input int((1 - c^2*x^2)^(5/2)/(x^3*(a + b*acosh(c*x))^2),x)
```

```
output int((1 - c^2*x^2)^(5/2)/(x^3*(a + b*acosh(c*x))^2), x)
```

---

3.340.  $\int \frac{(1 - c^2 x^2)^{5/2}}{x^3 (a + b \operatorname{barccosh}(cx))^2} dx$

**3.341** 
$$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\mathbf{arccosh}(cx))^2} dx$$

3.341.1 Optimal result	2749
3.341.2 Mathematica [ <b>F(-1)</b> ]	2749
3.341.3 Rubi [N/A]	2750
3.341.4 Maple [N/A] (verified)	2750
3.341.5 Fricas [N/A]	2751
3.341.6 Sympy [ <b>F(-1)</b> ]	2751
3.341.7 Maxima [N/A]	2751
3.341.8 Giac [N/A]	2752
3.341.9 Mupad [N/A]	2752

**3.341.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+\mathbf{barccosh}(cx))^2} dx = \mathbf{Int}\left(\frac{(1-c^2x^2)^{5/2}}{x^4(a+\mathbf{barccosh}(cx))^2}, x\right)$$

output `Unintegrable((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x))^2,x)`

**3.341.2 Mathematica [**F(-1)**]**

Timed out.

$$\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+\mathbf{barccosh}(cx))^2} dx = \mathbf{\$Aborted}$$

input `Integrate[(1 - c^2*x^2)^(5/2)/(x^4*(a + b*ArcCosh[c*x])^2),x]`

output `\$Aborted`

**3.341.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{arccosh}(cx))^2} dx$$

input `Int[(1 - c^2*x^2)^(5/2)/(x^4*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.341.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.341.4 Maple [N/A] (verified)**

Not integrable

Time = 1.43 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 x^2 + 1)^{5/2}}{x^4 (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x))^2,x)`

output `int((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x))^2,x)`

---

3.341.  $\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$

**3.341.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.36

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((c^4*x^4 - 2*c^2*x^2 + 1)*sqrt(-c^2*x^2 + 1)/(b^2*x^4*arccosh(c*x))^2 + 2*a*b*x^4*arccosh(c*x) + a^2*x^4), x)`

**3.341.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((-c**2*x**2+1)**(5/2)/x**4/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.341.7 Maxima [N/A]**

Not integrable

Time = 1.15 (sec) , antiderivative size = 533, normalized size of antiderivative = 19.04

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`



output `--((c^6*x^6 - 3*c^4*x^4 + 3*c^2*x^2 - 1)*(c*x + 1)*sqrt(c*x - 1) + (c^7*x^7 - 3*c^5*x^5 + 3*c^3*x^3 - c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^3*x^6 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^5 - a*b*c*x^4 + (b^2*c^3*x^6 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^5 - b^2*c*x^4)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((2*c^7*x^7 + c^5*x^5 - 8*c^3*x^3 + 5*c*x)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(2*c^8*x^8 - c^6*x^6 - 6*c^4*x^4 + 7*c^2*x^2 - 2)*(c*x + 1)*sqrt(c*x - 1) + (2*c^9*x^9 - 3*c^7*x^7 - 3*c^5*x^5 + 7*c^3*x^3 - 3*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)/(a*b*c^5*x^9 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^7 - 2*a*b*c^3*x^7 + a*b*c*x^5 + 2*(a*b*c^4*x^8 - a*b*c^2*x^6)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^9 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^7 - 2*b^2*c^3*x^7 + b^2*c*x^5 + 2*(b^2*c^4*x^8 - b^2*c^2*x^6)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.341.8 Giac [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{5/2}}{(b \operatorname{arcosh}(cx) + a)^2 x^4} dx$$

input `integrate((-c^2*x^2+1)^(5/2)/x^4/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((-c^2*x^2 + 1)^(5/2)/((b*arccosh(c*x) + a)^2*x^4), x)`

### 3.341.9 Mupad [N/A]

Not integrable

Time = 3.22 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(1 - c^2 x^2)^{5/2}}{x^4 (a + b \operatorname{acosh}(cx))^2} dx$$

input `int((1 - c^2*x^2)^(5/2)/(x^4*(a + b*acosh(c*x))^2),x)`

output `int((1 - c^2*x^2)^(5/2)/(x^4*(a + b*acosh(c*x))^2), x)`

---

3.341.  $\int \frac{(1-c^2x^2)^{5/2}}{x^4(a+b\operatorname{arccosh}(cx))^2} dx$

$$3.342 \quad \int \frac{x^5}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.342.1 Optimal result . . . . .	2753
3.342.2 Mathematica [A] (verified) . . . . .	2754
3.342.3 Rubi [A] (verified) . . . . .	2754
3.342.4 Maple [A] (verified) . . . . .	2757
3.342.5 Fracas [F] . . . . .	2757
3.342.6 Sympy [F] . . . . .	2758
3.342.7 Maxima [F] . . . . .	2758
3.342.8 Giac [F(-2)] . . . . .	2759
3.342.9 Mupad [F(-1)] . . . . .	2759

### 3.342.1 Optimal result

Integrand size = 28, antiderivative size = 337

$$\begin{aligned} \int \frac{x^5}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = & -\frac{x^5\sqrt{-1+cx}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\ & -\frac{5\sqrt{-1+cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{8b^2c^6\sqrt{1-cx}} \\ & -\frac{15\sqrt{-1+cx}\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{16b^2c^6\sqrt{1-cx}} \\ & -\frac{5\sqrt{-1+cx}\operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{5a}{b}\right)}{16b^2c^6\sqrt{1-cx}} \\ & +\frac{5\sqrt{-1+cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b^2c^6\sqrt{1-cx}} \\ & +\frac{15\sqrt{-1+cx}\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^6\sqrt{1-cx}} \\ & +\frac{5\sqrt{-1+cx}\cosh\left(\frac{5a}{b}\right)\operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^6\sqrt{1-cx}} \end{aligned}$$

output 
$$-x^5(c*x-1)^{(1/2)}/b/c/(a+b*\operatorname{arccosh}(c*x))/(-c*x+1)^{(1/2)}+5/8*\cosh(a/b)*\operatorname{Shi}((a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}/b^2/c^6/(-c*x+1)^{(1/2)}+15/16*\cosh(3*a/b)*\operatorname{Shi}(3*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}/b^2/c^6/(-c*x+1)^{(1/2)}+5/16*\cosh(5*a/b)*\operatorname{Shi}(5*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}/b^2/c^6/(-c*x+1)^{(1/2)}-5/8*\operatorname{Chi}((a+b*\operatorname{arccosh}(c*x))/b)*\sinh(a/b)*(c*x-1)^{(1/2)}/b^2/c^6/(-c*x+1)^{(1/2)}-15/16*\operatorname{Chi}(3*(a+b*\operatorname{arccosh}(c*x))/b)*\sinh(3*a/b)*(c*x-1)^{(1/2)}/b^2/c^6/(-c*x+1)^{(1/2)}-5/16*\operatorname{Chi}(5*(a+b*\operatorname{arccosh}(c*x))/b)*\sinh(5*a/b)*(c*x-1)^{(1/2)}/b^2/c^6/(-c*x+1)^{(1/2)}$$

### 3.342.2 Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 190, normalized size of antiderivative = 0.56

$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

$$= \frac{\sqrt{1-c^2x^2} \left( \frac{16bc^5x^5}{a+b\operatorname{arccosh}(cx)} + 5 \left( 2\operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \sinh\left(\frac{a}{b}\right) + 3\operatorname{Chi}\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \sinh\left(\frac{3a}{b}\right) + \dots \right) \right)}{\dots}$$

input `Integrate[x^5/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output 
$$\frac{(\operatorname{Sqrt}[1 - c^2*x^2]*((16*b*c^5*x^5)/(a + b*\operatorname{ArcCosh}[c*x]) + 5*(2*\operatorname{CoshIntegral}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Sinh}[a/b] + 3*\operatorname{CoshIntegral}[3*(a/b + \operatorname{ArcCosh}[c*x])]*\operatorname{Sinh}[(3*a)/b] + \operatorname{CoshIntegral}[5*(a/b + \operatorname{ArcCosh}[c*x])]*\operatorname{Sinh}[(5*a)/b] - 2*\operatorname{Cosh}[a/b]*\operatorname{SinhIntegral}[a/b + \operatorname{ArcCosh}[c*x]] - 3*\operatorname{Cosh}[(3*a)/b]*\operatorname{SinhIntegral}[3*(a/b + \operatorname{ArcCosh}[c*x])] - \operatorname{Cosh}[(5*a)/b]*\operatorname{SinhIntegral}[5*(a/b + \operatorname{ArcCosh}[c*x])])))/((16*b^2*c^6*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]))$$

### 3.342.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 215, normalized size of antiderivative = 0.64, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$ , Rules used = {6365, 6302, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

---

3.342. 
$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

$$\begin{aligned}
& \downarrow \text{6365} \\
& \frac{5\sqrt{cx-1} \int \frac{x^4}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{1-cx}} - \frac{x^5\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} \\
& \downarrow \text{6302} \\
& \frac{5\sqrt{cx-1} \int -\frac{\cosh^4\left(\frac{a-a+\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a-a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{\frac{b^2c^6\sqrt{1-cx}}{x^5\sqrt{cx-1}} bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} \\
& \downarrow \text{25} \\
& \frac{5\sqrt{cx-1} \int \frac{\cosh^4\left(\frac{a-a+\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a-a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{\frac{b^2c^6\sqrt{1-cx}}{x^5\sqrt{cx-1}} bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} \\
& \downarrow \text{5971} \\
& \frac{5\sqrt{cx-1} \int \left( \frac{\sinh\left(\frac{5a}{b}-\frac{5(a+\operatorname{barccosh}(cx))}{b}\right)}{16(a+\operatorname{barccosh}(cx))} + \frac{3\sinh\left(\frac{3a}{b}-\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{16(a+\operatorname{barccosh}(cx))} + \frac{\sinh\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{8(a+\operatorname{barccosh}(cx))} \right) d(a+\operatorname{barccosh}(cx))}{\frac{b^2c^6\sqrt{1-cx}}{x^5\sqrt{cx-1}} bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} \\
& \downarrow \text{2009} \\
& \frac{5\sqrt{cx-1} \left( -\frac{1}{8} \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) - \frac{3}{16} \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right) - \frac{1}{16} \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+\operatorname{barccosh}(cx))}{b}\right) \right)}{\frac{x^5\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))}}
\end{aligned}$$

input `Int[x^5/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

```
output  $-\left(\frac{x^5 \sqrt{-1 + cx}}{bc \sqrt{1 - cx} (a + b \operatorname{ArcCosh}[cx])}\right) + (5 \sqrt{-1 + cx} \left(-\frac{1}{8} \frac{\operatorname{CoshIntegral}[(a + b \operatorname{ArcCosh}[cx])/b] \operatorname{Sinh}[a/b]}{\operatorname{CoshIntegral}[(3(a + b \operatorname{ArcCosh}[cx])/b] \operatorname{Sinh}[(3a)/b])} - \frac{3 \operatorname{CoshIntegral}[(3(a + b \operatorname{ArcCosh}[cx])/b] \operatorname{Sinh}[(5a)/b]}{16} - \frac{\operatorname{CoshIntegral}[(5(a + b \operatorname{ArcCosh}[cx])/b] \operatorname{Sinh}[(5a)/b]}{16} + \frac{\operatorname{Cosh}[a/b] \operatorname{SinhIntegral}[(a + b \operatorname{ArcCosh}[cx])/b]}{8} + \frac{3 \operatorname{Cosh}[(3a)/b] \operatorname{SinhIntegral}[(3(a + b \operatorname{ArcCosh}[cx])/b]}{16} + \frac{\operatorname{Cosh}[(5a)/b] \operatorname{SinhIntegral}[(5(a + b \operatorname{ArcCosh}[cx])/b]}{16}\right) \right) / (b^2 c^6 \sqrt{1 - cx})$ 
```

### 3.342.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 5971 Int[Cosh[(a_.) + (b_.)(x_)]^(p_.)((c_.) + (d_.)(x_))^(m_.)*Sinh[(a_.) + (b_.)(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + dx)^m, Sinh[a + bx]^n Cosh[a + bx]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]
```

```
rule 6302 Int[((a_.) + ArcCosh[(c_.)(x_)]*(b_.))^(n_)*(x_)^(m_), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n Cosh[-a/b + x/b]^m Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]
```

```
rule 6365 Int[(((a_.) + ArcCosh[(c_.)(x_)]*(b_.))^(n_)*((f_.)(x_))^(m_.))/Sqrt[(d_) + (e_.)(x_)^2], x_Symbol] := Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2]), x] - Simp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2]) Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]
```

**3.342.4 Maple [A] (verified)**

Time = 0.97 (sec) , antiderivative size = 519, normalized size of antiderivative = 1.54

method	result
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-32\sqrt{cx-1}\sqrt{cx+1}bc^5x^5+32bc^6x^6+15\operatorname{arccosh}(cx)b\operatorname{Ei}_1\left(-3\operatorname{arccosh}(cx)-\frac{3a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)+a}{b}}\right)}{\dots}$

input `int(x^5/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{32}(-c^2x^2+1)^{1/2}((cx-1)^{1/2}(cx+1)^{1/2}cx+c^2x^2-1)\left(-32((cx-1)^{1/2}(cx+1)^{1/2}b^2c^5x^5+32b^2c^6x^6+15\operatorname{arccosh}(cx)b\operatorname{Ei}_1(-3\operatorname{arccosh}(cx)-\frac{3a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+a}{b})+10\operatorname{arccosh}(cx)b\operatorname{Ei}_1(-\operatorname{arccosh}(cx)-\frac{a}{b})\exp(-\frac{a+b\operatorname{arccosh}(cx)}{b})+5\operatorname{arccosh}(cx)b\operatorname{Ei}_1(-5\operatorname{arccosh}(cx)-\frac{5a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+5a}{b})-5\operatorname{Ei}_1(1,5\operatorname{arccosh}(cx)+\frac{5a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+5a}{b})b\operatorname{arccosh}(cx)-15\operatorname{Ei}_1(1,3\operatorname{arccosh}(cx)+\frac{3a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+3a}{b})b\operatorname{arccosh}(cx)-10\operatorname{Ei}_1(1,\operatorname{arccosh}(cx)+\frac{a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+a}{b})b\operatorname{arccosh}(cx)+15a\operatorname{Ei}_1(-3\operatorname{arccosh}(cx)-\frac{3a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+3a}{b})+10a\operatorname{Ei}_1(-\operatorname{arccosh}(cx)-\frac{a}{b})\exp(-\frac{a+b\operatorname{arccosh}(cx)}{b})+5a\operatorname{Ei}_1(-5\operatorname{arccosh}(cx)-\frac{5a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+5a}{b})-5\operatorname{Ei}_1(1,5\operatorname{arccosh}(cx)+\frac{5a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+5a}{b})a-15\operatorname{Ei}_1(1,3\operatorname{arccosh}(cx)+\frac{3a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+3a}{b})a-10\operatorname{Ei}_1(1,\operatorname{arccosh}(cx)+\frac{a}{b})\exp(-\frac{b\operatorname{arccosh}(cx)+a}{b})a\right)/c^6/(-c^2x^2+1)^{1/2}/(a+b\operatorname{arccosh}(cx))$$

**3.342.5 Fracas [F]**

$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^5}{\sqrt{-c^2x^2+1}(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate(x^5/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*x^2+1)*x^5/(a^2*c^2*x^2+(b^2*c^2*x^2-b^2)*arccosh(c*x)^2-a^2+2*(a*b*c^2*x^2-a*b)*arccosh(c*x)),x)`

## 3.342.6 Sympy [F]

$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^5}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x**5/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x**5/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2), x)`

## 3.342.7 Maxima [F]

$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^5}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^5/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `-(c^3*x^8 - c*x^6 + (c^2*x^7 - x^5)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((5*c^5*x^9 - 11*c^3*x^7 + 6*c*x^5 + (5*c^3*x^7 - 4*c*x^5)*(c*x + 1)*(c*x - 1) + 5*(2*c^4*x^8 - 3*c^2*x^6 + x^4)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)^(3/2)*(c*x - 1)*b^2*c^3*x^2 + 2*(b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x - 1)*a*b*c^3*x^2 + 2*(a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

**3.342.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^5/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.342.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^5}{(a+b\operatorname{acosh}(cx))^2\sqrt{1-c^2x^2}} dx$$

input `int(x^5/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)`

output `int(x^5/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)`



### 3.343 $\int \frac{x^4}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.343.1 Optimal result . . . . .	2760
3.343.2 Mathematica [A] (verified) . . . . .	2761
3.343.3 Rubi [A] (verified) . . . . .	2761
3.343.4 Maple [A] (verified) . . . . .	2763
3.343.5 Fricas [F] . . . . .	2764
3.343.6 Sympy [F] . . . . .	2764
3.343.7 Maxima [F] . . . . .	2764
3.343.8 Giac [F] . . . . .	2765
3.343.9 Mupad [F(-1)] . . . . .	2765

#### 3.343.1 Optimal result

Integrand size = 28, antiderivative size = 236

$$\int \frac{x^4}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^4\sqrt{-1+cx}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{-1+cx}\operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)}{b^2c^5\sqrt{1-cx}} - \frac{\sqrt{-1+cx}\operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{4a}{b}\right)}{2b^2c^5\sqrt{1-cx}} + \frac{\sqrt{-1+cx}\cosh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{b^2c^5\sqrt{1-cx}} + \frac{\sqrt{-1+cx}\cosh\left(\frac{4a}{b}\right)\operatorname{Shi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{2b^2c^5\sqrt{1-cx}}$$

output

```
-x^4*(c*x-1)^(1/2)/b/c/(a+b*arccosh(c*x))/(-c*x+1)^(1/2)+cosh(2*a/b)*Shi(2
*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/b^2/c^5/(-c*x+1)^(1/2)+1/2*cosh(4*a/b
)*Shi(4*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/b^2/c^5/(-c*x+1)^(1/2)-Chi(2*(
a+b*arccosh(c*x))/b)*sinh(2*a/b)*(c*x-1)^(1/2)/b^2/c^5/(-c*x+1)^(1/2)-1/2*
Chi(4*(a+b*arccosh(c*x))/b)*sinh(4*a/b)*(c*x-1)^(1/2)/b^2/c^5/(-c*x+1)^(1/
2)
```

**3.343.2 Mathematica [A] (verified)**

Time = 0.37 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.63

$$\int \frac{x^4}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$$

$$= \frac{\sqrt{1-c^2x^2} \left( \frac{2bc^4x^4}{a+b\operatorname{arccosh}(cx)} + 2\operatorname{Chi}\left(2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \sinh\left(\frac{2a}{b}\right) + \operatorname{Chi}\left(4\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \sinh\left(\frac{4a}{b}\right) - \operatorname{CoshIntegral}\left[2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right] \operatorname{SinhIntegral}\left[2\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right] - \operatorname{CoshIntegral}\left[4\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right] \operatorname{SinhIntegral}\left[4\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right] \right)}{2b^2c^5\sqrt{-1+cx}\sqrt{1+cx}}$$

input `Integrate[x^4/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`output `(Sqrt[1 - c^2*x^2]*((2*b*c^4*x^4)/(a + b*ArcCosh[c*x]) + 2*CoshIntegral[2*(a/b + ArcCosh[c*x])]*Sinh[(2*a)/b] + CoshIntegral[4*(a/b + ArcCosh[c*x])]*Sinh[(4*a)/b] - 2*Cosh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[c*x])] - Cosh[(4*a)/b]*SinhIntegral[4*(a/b + ArcCosh[c*x])])/(2*b^2*c^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`**3.343.3 Rubi [A] (verified)**Time = 0.69 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.72, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$ , Rules used = {6365, 6302, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$$

$$\downarrow \text{6365}$$

$$\frac{4\sqrt{cx-1} \int \frac{x^3}{a+b\operatorname{arccosh}(cx)} dx}{bc\sqrt{1-cx}} - \frac{x^4\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))}$$

$$\downarrow \text{6302}$$

$$\frac{4\sqrt{cx-1} \int -\frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{\frac{b^2c^5\sqrt{1-cx}}{x^4\sqrt{cx-1}}}$$

$$\frac{\phantom{4\sqrt{cx-1} \int -\frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))}$$

---

3.343.  $\int \frac{x^4}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{4\sqrt{cx-1} \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{\frac{b^2c^5\sqrt{1-cx}}{x^4\sqrt{cx-1}} bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\
 & \downarrow 5971 \\
 & \frac{4\sqrt{cx-1} \int \left( \frac{\sinh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{\frac{b^2c^5\sqrt{1-cx}}{x^4\sqrt{cx-1}} bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\
 & \downarrow 2009 \\
 & \frac{4\sqrt{cx-1} \left( -\frac{1}{4} \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - \frac{1}{8} \sinh\left(\frac{4a}{b}\right) \operatorname{Chi}\left(\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{x^4\sqrt{cx-1} bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))}
 \end{aligned}$$

input `Int[x^4/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `-(x^4*Sqrt[-1 + c*x])/(b*c*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x])) + (4*Sqrt[-1 + c*x]*(-1/4*(CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]*Sinh[(2*a)/b]) - (CoshIntegral[(4*(a + b*ArcCosh[c*x]))/b]*Sinh[(4*a)/b])/8 + (Cosh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b])/4 + (Cosh[(4*a)/b]*SinhIntegral[(4*(a + b*ArcCosh[c*x]))/b])/8))/(b^2*c^5*Sqrt[1 - c*x])`

### 3.343.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6365 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.))/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])], x] - Simp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]`

### 3.343.4 Maple [A] (verified)

Time = 0.87 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.64

method	result
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-4x^4c^4b\sqrt{cx-1}\sqrt{cx+1}+4bc^5x^5+\operatorname{arccosh}(cx)b\operatorname{Ei}_1(-4\operatorname{arccosh}(cx)-\frac{4a}{b})e^{-\frac{b\operatorname{arccosh}(cx)}{b}}\right)}{\dots}$

input `int(x^4/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/4*(-c^2*x^2+1)^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-4*x^4*c^4*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*b*c^5*x^5+arccosh(c*x)*b*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(b*arccosh(c*x)+4*a)/b)+2*arccosh(c*x)*b*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(b*arccosh(c*x)+2*a)/b)-Ei(1,4*arccosh(c*x)+4*a/b)*exp((-b*arccosh(c*x)+4*a)/b)*b*arccosh(c*x)-2*Ei(1,2*arccosh(c*x)+2*a/b)*exp((-b*arccosh(c*x)+2*a)/b)*b*arccosh(c*x)+a*Ei(1,-4*arccosh(c*x)-4*a/b)*exp(-(b*arccosh(c*x)+4*a)/b)+2*a*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(b*arccosh(c*x)+2*a)/b)-Ei(1,4*arccosh(c*x)+4*a/b)*exp((-b*arccosh(c*x)+4*a)/b)*a-2*Ei(1,2*arccosh(c*x)+2*a/b)*exp((-b*arccosh(c*x)+2*a)/b)*a)/c^5/(c^2*x^2-1)/b^2/(a+b*arccosh(c*x))`

**3.343.5 Fricas [F]**

$$\int \frac{x^4}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^4}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^4/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^4/(a^2*c^2*x^2 + (b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.343.6 Sympy [F]**

$$\int \frac{x^4}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^4}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x**4/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x**4/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2), x)`

**3.343.7 Maxima [F]**

$$\int \frac{x^4}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^4}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^4/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output  $-(c^3x^7 - cx^5 + (c^2x^6 - x^4)\sqrt{cx + 1}\sqrt{cx - 1})/(((cx + 1)\sqrt{cx - 1}b^2c^2x + (b^2c^3x^2 - b^2c)\sqrt{cx + 1})\sqrt{-cx + 1})\log(cx + \sqrt{cx + 1}\sqrt{cx - 1}) + ((cx + 1)\sqrt{cx - 1})ab^2c^2x + (ab^2c^3x^2 - ab^2c)\sqrt{cx + 1})\sqrt{-cx + 1}) + \text{integrate}((4c^5x^8 - 9c^3x^6 + 5cx^4 + (4c^3x^6 - 3cx^4)(cx + 1)(cx - 1) + 4(2c^4x^7 - 3c^2x^5 + x^3)\sqrt{cx + 1}\sqrt{cx - 1})/(((cx + 1)^{3/2}(cx - 1)b^2c^3x^2 + 2(b^2c^4x^3 - b^2c^2x)(cx + 1)\sqrt{cx - 1} + (b^2c^5x^4 - 2b^2c^3x^2 + b^2c)\sqrt{cx + 1})\sqrt{-cx + 1})\log(cx + \sqrt{cx + 1}\sqrt{cx - 1}) + ((cx + 1)^{3/2}(cx - 1)ab^2c^3x^2 + 2(ab^2c^4x^3 - ab^2c^2x)(cx + 1)\sqrt{cx - 1} + (ab^2c^5x^4 - 2ab^2c^3x^2 + ab^2c)\sqrt{cx + 1})\sqrt{-cx + 1}), x)$

### 3.343.8 Giac [F]

$$\int \frac{x^4}{\sqrt{1 - c^2x^2}(a + \text{barccosh}(cx))^2} dx = \int \frac{x^4}{\sqrt{-c^2x^2 + 1}(b \text{arcosh}(cx) + a)^2} dx$$

input `integrate(x^4/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^4/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^2), x)`

### 3.343.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^4}{\sqrt{1 - c^2x^2}(a + \text{barccosh}(cx))^2} dx = \int \frac{x^4}{(a + b \text{acosh}(cx))^2 \sqrt{1 - c^2x^2}} dx$$

input `int(x^4/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)`

output `int(x^4/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)`

### 3.344 $\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.344.1 Optimal result . . . . .	2766
3.344.2 Mathematica [A] (verified) . . . . .	2767
3.344.3 Rubi [A] (verified) . . . . .	2767
3.344.4 Maple [A] (verified) . . . . .	2769
3.344.5 Fricas [F] . . . . .	2770
3.344.6 Sympy [F] . . . . .	2770
3.344.7 Maxima [F] . . . . .	2770
3.344.8 Giac [F(-2)] . . . . .	2771
3.344.9 Mupad [F(-1)] . . . . .	2771

#### 3.344.1 Optimal result

Integrand size = 28, antiderivative size = 237

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^3\sqrt{-1+cx}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{3\sqrt{-1+cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{4b^2c^4\sqrt{1-cx}} - \frac{3\sqrt{-1+cx}\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{4b^2c^4\sqrt{1-cx}} + \frac{3\sqrt{-1+cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4b^2c^4\sqrt{1-cx}} + \frac{3\sqrt{-1+cx}\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b^2c^4\sqrt{1-cx}}$$

output  $-x^3(c*x-1)^{(1/2)}/b/c/(a+b*\operatorname{arccosh}(c*x))/(-c*x+1)^{(1/2)}+3/4*\cosh(a/b)*\operatorname{Shi}((a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}/b^2/c^4/(-c*x+1)^{(1/2)}+3/4*\cosh(3*a/b)*\operatorname{Shi}(3*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}/b^2/c^4/(-c*x+1)^{(1/2)}-3/4*\operatorname{Chi}((a+b*\operatorname{arccosh}(c*x))/b)*\sinh(a/b)*(c*x-1)^{(1/2)}/b^2/c^4/(-c*x+1)^{(1/2)}-3/4*\operatorname{Chi}(3*(a+b*\operatorname{arccosh}(c*x))/b)*\sinh(3*a/b)*(c*x-1)^{(1/2)}/b^2/c^4/(-c*x+1)^{(1/2)}$

**3.344.2 Mathematica [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.61

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$$

$$= \frac{\sqrt{1-c^2x^2} \left( \frac{4bc^3x^3}{a+b\operatorname{arccosh}(cx)} + 3\operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \sinh\left(\frac{a}{b}\right) + 3\operatorname{Chi}\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \sinh\left(\frac{3a}{b}\right) - 3\operatorname{Chi}\left(\frac{a}{b}\right) \sinh\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \right)}{4b^2c^4\sqrt{-1+cx}\sqrt{1+cx}}$$

input `Integrate[x^3/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2),x]`output `(Sqrt[1 - c^2*x^2]*((4*b*c^3*x^3)/(a + b*ArcCosh[c*x]) + 3*CoshIntegral[a/b + ArcCosh[c*x]]*Sinh[a/b] + 3*CoshIntegral[3*(a/b + ArcCosh[c*x]])*Sinh[(3*a)/b] - 3*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] - 3*Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])]))/(4*b^2*c^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`**3.344.3 Rubi [A] (verified)**Time = 0.68 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.70, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$ , Rules used = {6365, 6302, 25, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$$

$$\downarrow \text{6365}$$

$$\frac{3\sqrt{cx-1} \int \frac{x^2}{a+b\operatorname{arccosh}(cx)} dx}{bc\sqrt{1-cx}} - \frac{x^3\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))}$$

$$\downarrow \text{6302}$$

$$\frac{3\sqrt{cx-1} \int -\frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{\frac{b^2c^4\sqrt{1-cx}}{x^3\sqrt{cx-1}}}$$

$$\frac{\phantom{3\sqrt{cx-1} \int -\frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+\operatorname{barccosh}(cx))}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))}$$

---

3.344.  $\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$



$$\begin{aligned}
 & \downarrow 25 \\
 & \frac{3\sqrt{cx-1} \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{\frac{b^2c^4\sqrt{1-cx}}{x^3\sqrt{cx-1}} \operatorname{arccosh}(cx)} \\
 & \downarrow 5971 \\
 & \frac{3\sqrt{cx-1} \int \left( \frac{\sinh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} + \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4(a+b\operatorname{arccosh}(cx))} \right) d(a+b\operatorname{arccosh}(cx))}{\frac{b^2c^4\sqrt{1-cx}}{x^3\sqrt{cx-1}} \operatorname{arccosh}(cx)} \\
 & \downarrow 2009 \\
 & \frac{3\sqrt{cx-1} \left( -\frac{1}{4} \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \frac{1}{4} \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) + \frac{1}{4} \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{\frac{b^2c^4\sqrt{1-cx}}{x^3\sqrt{cx-1}} \operatorname{arccosh}(cx)}
 \end{aligned}$$

input `Int[x^3/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2),x]`

output `-((x^3*Sqrt[-1 + c*x])/(b*c*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x]))) + (3*Sqrt[-1 + c*x]*(-1/4*(CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b]) - (CoshIntegral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*a)/b])/4 + (Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/4 + (Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/4))/(b^2*c^4*Sqrt[1 - c*x])`

**3.344.3.1 Defintions of rubi rules used**

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

---

3.344.  $\int \frac{x^3}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6365 `Int((((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.))/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])], x] - Simp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]`

### 3.344.4 Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 375, normalized size of antiderivative = 1.58

method	result
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-8\sqrt{cx-1}\sqrt{cx+1}bc^3x^3+8bc^4x^4+3\operatorname{arccosh}(cx)b\operatorname{Ei}_1\left(-3\operatorname{arccosh}(cx)-\frac{3a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)}{b}}\right)}{\dots}$

input `int(x^3/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/8*(-c^2*x^2+1)^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^3*x^3+8*b*c^4*x^4+3*arccosh(c*x)*b*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(b*arccosh(c*x)+3*a)/b)+3*arccosh(c*x)*b*Ei(1,-arccosh(c*x)-a/b)*exp(-(a+b*arccosh(c*x))/b)-3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((-b*arccosh(c*x)+3*a)/b)*b*arccosh(c*x)-3*Ei(1,arccosh(c*x)+a/b)*exp((-b*arccosh(c*x)+a)/b)*b*arccosh(c*x)+3*a*Ei(1,-3*arccosh(c*x)-3*a/b)*exp(-(b*arccosh(c*x)+3*a)/b)+3*a*Ei(1,-arccosh(c*x)-a/b)*exp(-(a+b*arccosh(c*x))/b)-3*Ei(1,3*arccosh(c*x)+3*a/b)*exp((-b*arccosh(c*x)+3*a)/b)*a-3*Ei(1,arccosh(c*x)+a/b)*exp((-b*arccosh(c*x)+a)/b)*a)/c^4/(c^2*x^2-1)/b^2/(a+b*arccosh(c*x))`

**3.344.5 Fricas [F]**

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^3/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^3/(a^2*c^2*x^2 + (b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.344.6 Sympy [F]**

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x**3/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x**3/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2), x)`

**3.344.7 Maxima [F]**

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^3/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

```
output -(c^3*x^6 - c*x^4 + (c^2*x^5 - x^3)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x +
1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*
x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a
*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrat
e((3*c^5*x^7 - 7*c^3*x^5 + 4*c*x^3 + (3*c^3*x^5 - 2*c*x^3)*(c*x + 1)*(c*x
- 1) + 3*(2*c^4*x^6 - 3*c^2*x^4 + x^2)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x
+ 1)^(3/2)*(c*x - 1)*b^2*c^3*x^2 + 2*(b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*
sqrt(c*x - 1) + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*sqrt(
-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x
- 1)*a*b*c^3*x^2 + 2*(a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a
*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)
```

### 3.344.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

```
input integrate(x^3/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.344.9 Mupad [F(-1)]

Timed out.

$$\int \frac{x^3}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(a+b\operatorname{acosh}(cx))^2\sqrt{1-c^2x^2}} dx$$

```
input int(x^3/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)
```

```
output int(x^3/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)
```

### 3.345 $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.345.1 Optimal result . . . . .	2772
3.345.2 Mathematica [A] (verified) . . . . .	2772
3.345.3 Rubi [C] (verified) . . . . .	2773
3.345.4 Maple [A] (verified) . . . . .	2777
3.345.5 Fricas [F] . . . . .	2777
3.345.6 Sympy [F] . . . . .	2777
3.345.7 Maxima [F] . . . . .	2778
3.345.8 Giac [F] . . . . .	2778
3.345.9 Mupad [F(-1)] . . . . .	2779

#### 3.345.1 Optimal result

Integrand size = 28, antiderivative size = 136

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^2\sqrt{-1+cx}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{-1+cx}\operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{2a}{b}\right)}{b^2c^3\sqrt{1-cx}} + \frac{\sqrt{-1+cx}\cosh\left(\frac{2a}{b}\right)\operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{b^2c^3\sqrt{1-cx}}$$

output  $-x^2*(c*x-1)^{(1/2)}/b/c/(a+b*\operatorname{arccosh}(c*x))/(-c*x+1)^{(1/2)}+\cosh(2*a/b)*\operatorname{Shi}(2*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}/b^2/c^3/(-c*x+1)^{(1/2)}-\operatorname{Chi}(2*(a+b*\operatorname{arccosh}(c*x))/b)*\sinh(2*a/b)*(c*x-1)^{(1/2)}/b^2/c^3/(-c*x+1)^{(1/2)}$

#### 3.345.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.86

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \frac{\sqrt{1-c^2x^2}(bc^2x^2+(a+b\operatorname{arccosh}(cx))\operatorname{Chi}(2(\frac{a}{b}+\operatorname{arccosh}(cx))))\sinh(\frac{2a}{b})-(a+b\operatorname{arccosh}(cx))\cosh(\frac{2a}{b})}{b^2c^3\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}$$

input `Integrate[x^2/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `(Sqrt[1 - c^2*x^2]*(b*c^2*x^2 + (a + b*ArcCosh[c*x])*CoshIntegral[2*(a/b + ArcCosh[c*x]])*Sinh[(2*a)/b] - (a + b*ArcCosh[c*x])*Cosh[(2*a)/b]*SinhIntegral[2*(a/b + ArcCosh[c*x])])/(b^2*c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))`

### 3.345.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.81 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.89, number of steps used = 14, number of rules used = 13,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.464$ , Rules used = {6365, 6302, 25, 5971, 27, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6365} \\
 & \frac{2\sqrt{cx-1} \int \frac{x}{a+\operatorname{barccosh}(cx)} dx}{bc\sqrt{1-cx}} - \frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6302} \\
 & \frac{2\sqrt{cx-1} \int -\frac{\cosh\left(\frac{a}{b}-\frac{a+b\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}-\frac{a+b\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{\frac{b^2c^3\sqrt{1-cx}}{x^2\sqrt{cx-1}} bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{2\sqrt{cx-1} \int \frac{\cosh\left(\frac{a}{b}-\frac{a+b\operatorname{barccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}-\frac{a+b\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{\frac{b^2c^3\sqrt{1-cx}}{x^2\sqrt{cx-1}} bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{5971}
 \end{aligned}$$

---

3.345.  $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
 & -\frac{2\sqrt{cx-1} \int \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2(a+b\operatorname{arccosh}(cx))} d(a+b\operatorname{arccosh}(cx))}{b^2c^3\sqrt{1-cx}} - \frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 27 \\
 & -\frac{\sqrt{cx-1} \int \frac{\sinh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^3\sqrt{1-cx}} - \frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow 3042 \\
 & -\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{cx-1} \int \frac{i \sin\left(\frac{2ia}{b} - \frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^3\sqrt{1-cx}} \\
 & \quad \downarrow 26 \\
 & -\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} + \frac{i\sqrt{cx-1} \int \frac{\sin\left(\frac{2ia}{b} - \frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^3\sqrt{1-cx}} \\
 & \quad \downarrow 3784 \\
 & -\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} + \\
 & i\sqrt{cx-1} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) + \cosh\left(\frac{2a}{b}\right) \int \frac{i \sinh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right) \\
 & \quad \downarrow 26 \\
 & -\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} + \\
 & i\sqrt{cx-1} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\cosh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \int \frac{\sinh\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right) \\
 & \quad \downarrow 3042 \\
 & -\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} + \\
 & i\sqrt{cx-1} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \int \frac{i \sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)
 \end{aligned}$$

3.345.  $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

$$\begin{aligned}
 & \downarrow 26 \\
 & \frac{-\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{arccosh}(cx))} + i\sqrt{cx-1} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - \cosh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) \right)}{b^2c^3\sqrt{1-cx}} \\
 & \downarrow 3779 \\
 & \frac{-\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{arccosh}(cx))} + i\sqrt{cx-1} \left( i \sinh\left(\frac{2a}{b}\right) \int \frac{\sin\left(\frac{2i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2c^3\sqrt{1-cx}} \\
 & \downarrow 3782 \\
 & \frac{-\frac{x^2\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{arccosh}(cx))} + i\sqrt{cx-1} \left( i \sinh\left(\frac{2a}{b}\right) \operatorname{Chi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - i \cosh\left(\frac{2a}{b}\right) \operatorname{Shi}\left(\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) \right)}{b^2c^3\sqrt{1-cx}}
 \end{aligned}$$

input `Int[x^2/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2),x]`

output `-((x^2*Sqrt[-1 + c*x])/(b*c*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x]))) + (I*Sqrt[-1 + c*x]*(I*CoshIntegral[(2*(a + b*ArcCosh[c*x]))/b]*Sinh[(2*a)/b] - I*Cosh[(2*a)/b]*SinhIntegral[(2*(a + b*ArcCosh[c*x]))/b]))/(b^2*c^3*Sqrt[1 - c*x])`

### 3.345.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

---

3.345.  $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$



- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3779 `Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`
- rule 3782 `Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`
- rule 3784 `Int[sin[(e_) + (f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`
- rule 5971 `Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`
- rule 6302 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)^{(m_)}, x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`
- rule 6365 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)/Sqrt[(d_) + (e_)*(x_)^2], x_Symbol] := Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2]), x] - Simp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2]) Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]`

**3.345.4 Maple [A] (verified)**

Time = 0.77 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.79

method	result
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-2\sqrt{cx-1}\sqrt{cx+1}bc^2x^2+2bc^3x^3+\operatorname{arccosh}(cx)b\operatorname{Ei}_1\left(-2\operatorname{arccosh}(cx)-\frac{2a}{b}\right)e^{-\frac{b\operatorname{arccosh}(cx)}{b}}\right)}{2c^3}$

```
input int(x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output 1/2*(-c^2*x^2+1)^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*b*c^2*x^2+2*b*c^3*x^3+arccosh(c*x)*b*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(b*arccosh(c*x)+2*a)/b)-Ei(1,2*arccosh(c*x)+2*a/b)*exp((-b*arccosh(c*x)+2*a)/b)*b*arccosh(c*x)+a*Ei(1,-2*arccosh(c*x)-2*a/b)*exp(-(b*arccosh(c*x)+2*a)/b)-Ei(1,2*arccosh(c*x)+2*a/b)*exp((-b*arccosh(c*x)+2*a)/b)*a)/c^3/(c^2*x^2-1)/b^2/(a+b*arccosh(c*x))
```

**3.345.5 Fracas [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{\sqrt{-c^2x^2+1}(b\operatorname{arccosh}(cx)+a)^2} dx$$

```
input integrate(x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fracas")
```

```
output integral(-sqrt(-c^2*x^2+1)*x^2/(a^2*c^2*x^2+(b^2*c^2*x^2-b^2)*arccosh(c*x)^2-a^2+2*(a*b*c^2*x^2-a*b)*arccosh(c*x)),x)
```

**3.345.6 Sympy [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))^2} dx$$

```
input integrate(x**2/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2),x)
```

```
output Integral(x**2/(sqrt(-(c*x-1)*(c*x+1))*(a+b*acosh(c*x))**2),x)
```

---

3.345.  $\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.345.7 Maxima [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^2}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `-(c^3*x^5 - c*x^3 + (c^2*x^4 - x^2)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((2*c^5*x^6 - 5*c^3*x^4 + (2*c^3*x^4 - c*x^2)*(c*x + 1)*(c*x - 1) + 3*c*x^2 + 2*(2*c^4*x^5 - 3*c^2*x^3 + x)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)^(3/2)*(c*x - 1)*b^2*c^3*x^2 + 2*(b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x - 1)*a*b*c^3*x^2 + 2*(a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

**3.345.8 Giac [F]**

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^2}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x^2/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^2), x)`

**3.345.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{(a+b\operatorname{acosh}(cx))^2\sqrt{1-c^2x^2}} dx$$

input `int(x^2/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)`output `int(x^2/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)`

### 3.346 $\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.346.1 Optimal result	2780
3.346.2 Mathematica [A] (verified)	2780
3.346.3 Rubi [C] (verified)	2781
3.346.4 Maple [A] (verified)	2784
3.346.5 Fricas [F]	2784
3.346.6 Sympy [F]	2785
3.346.7 Maxima [F]	2785
3.346.8 Giac [F]	2786
3.346.9 Mupad [F(-1)]	2786

#### 3.346.1 Optimal result

Integrand size = 26, antiderivative size = 130

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x\sqrt{-1+cx}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{-1+cx}\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{b^2c^2\sqrt{1-cx}} + \frac{\sqrt{-1+cx}\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c^2\sqrt{1-cx}}$$

```
output -x*(c*x-1)^(1/2)/b/c/(a+b*arccosh(c*x))/(-c*x+1)^(1/2)+cosh(a/b)*Shi((a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/b^2/c^2/(-c*x+1)^(1/2)-Chi((a+b*arccosh(c*x))/b)*sinh(a/b)*(c*x-1)^(1/2)/b^2/c^2/(-c*x+1)^(1/2)
```

#### 3.346.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.82

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \frac{\sqrt{1-c^2x^2}(bcx+(a+b\operatorname{arccosh}(cx))\operatorname{Chi}\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right)\sinh\left(\frac{a}{b}\right)-(a+b\operatorname{arccosh}(cx))\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a}{b}+\operatorname{arccosh}(cx)\right))}{b^2c^2\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))}$$

input `Integrate[x/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2),x]`

output `(Sqrt[1 - c^2*x^2]*(b*c*x + (a + b*ArcCosh[c*x])*CoshIntegral[a/b + ArcCosh[c*x]]*Sinh[a/b] - (a + b*ArcCosh[c*x])*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]]))/(b^2*c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))`

### 3.346.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.65 (sec) , antiderivative size = 115, normalized size of antiderivative = 0.88, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.423$ , Rules used = {6365, 6296, 25, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6365} \\
 & \frac{\sqrt{cx-1} \int \frac{1}{a+b\operatorname{arccosh}(cx)} dx}{bc\sqrt{1-cx}} - \frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{6296} \\
 & \frac{\sqrt{cx-1} \int -\frac{\sinh\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{1-cx}} - \frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\sqrt{cx-1} \int \frac{\sinh\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{1-cx}} - \frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{cx-1} \int -\frac{i \sin\left(\frac{ia}{b}-\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c^2\sqrt{1-cx}} \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} + \frac{i\sqrt{cx-1} \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx))}{b^2c^2\sqrt{1-cx}} \\
& \quad \downarrow \text{3784} \\
& -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} + \\
& i\sqrt{cx-1} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) + \cosh\left(\frac{a}{b}\right) \int \frac{i \sinh\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) \right) \\
& \quad \downarrow \text{26} \\
& -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} + \\
& i\sqrt{cx-1} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) \right) \\
& \quad \downarrow \text{3042} \\
& -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} + \\
& i\sqrt{cx-1} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \int \frac{i \sin\left(\frac{i(a+\operatorname{barccosh}(cx))}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) \right) \\
& \quad \downarrow \text{26} \\
& -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} + \\
& i\sqrt{cx-1} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) - \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+\operatorname{barccosh}(cx))}{b}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) \right) \\
& \quad \downarrow \text{3779} \\
& -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{barccosh}(cx))} + \\
& i\sqrt{cx-1} \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+\operatorname{barccosh}(cx)} d(a+\operatorname{barccosh}(cx)) - i \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) \right)
\end{aligned}$$

---

3.346.  $\int \frac{x}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{array}{c} \downarrow \text{3782} \\ -\frac{x\sqrt{cx-1}}{bc\sqrt{1-cx}(a+\operatorname{arccosh}(cx))} + \\ \frac{i\sqrt{cx-1}\left(i\sinh\left(\frac{a}{b}\right)\operatorname{Chi}\left(\frac{a+\operatorname{arccosh}(cx)}{b}\right) - i\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+\operatorname{arccosh}(cx)}{b}\right)\right)}{b^2c^2\sqrt{1-cx}} \end{array}$$

input `Int[x/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2),x]`

output `-((x*Sqrt[-1 + c*x])/(b*c*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x]))) + (I*Sqrt[-1 + c*x]*(I*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b] - I*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b]))/(b^2*c^2*Sqrt[1 - c*x])`

### 3.346.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`

rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`

rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`



rule 6296 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[1/(b*c) S  
ubst[Int[x^n*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a,  
b, c, n}, x]`

rule 6365 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)/Sqrt[(d_ +  
(e_.)*(x_)^2], x_Symbol] := Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(  
b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])], x] - Si  
mp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]  
Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c  
, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]`

### 3.346.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.73

method	result
default	$\frac{\sqrt{-c^2x^2+1}(\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2-1)\left(-2\sqrt{cx-1}\sqrt{cx+1}bcx+2b^2c^2x^2+\operatorname{arccosh}(cx)b\operatorname{Ei}_1\left(-\operatorname{arccosh}(cx)-\frac{a}{b}\right)e^{-\frac{a+b\operatorname{arccosh}(cx)}{b}}\right)}{2c^2(c^2x^2-1)}$

input `int(x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*(-c^2*x^2+1)^(1/2)*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2-1)*(-2*(c*  
x-1)^(1/2)*(c*x+1)^(1/2)*b*c*x+2*b*c^2*x^2+arccosh(c*x)*b*Ei(1,-arccosh(c*  
x)-a/b)*exp(-(a+b*arccosh(c*x))/b)-Ei(1,arccosh(c*x)+a/b)*exp((-b*arccosh(  
c*x)+a)/b)*b*arccosh(c*x)+a*Ei(1,-arccosh(c*x)-a/b)*exp(-(a+b*arccosh(c*x)  
)/b)-Ei(1,arccosh(c*x)+a/b)*exp((-b*arccosh(c*x)+a)/b)*a)/c^2/(c^2*x^2-1)/  
b^2/(a+b*arccosh(c*x))`

### 3.346.5 Fricas [F]

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x}{\sqrt{-c^2x^2+1}(b\operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate(x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x/(a^2*c^2*x^2 + (b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

### 3.346.6 Sympy [F]

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(x/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2), x)`

output `Integral(x/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2), x)`

### 3.346.7 Maxima [F]

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2), x, algorithm="maxima")`

output `-(c^3*x^4 - c*x^2 + (c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((c^5*x^5 + (c*x + 1)*(c*x - 1)*c^3*x^3 - 3*c^3*x^3 + (2*c^4*x^4 - 3*c^2*x^2 + 1)*sqrt(c*x + 1)*sqrt(c*x - 1) + 2*c*x)/(((c*x + 1)^(3/2)*(c*x - 1)*b^2*c^3*x^2 + 2*(b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x - 1)*a*b*c^3*x^2 + 2*(a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

**3.346.8 Giac [F]**

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(x/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^2), x)`

**3.346.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x}{(a+b\operatorname{acosh}(cx))^2\sqrt{1-c^2x^2}} dx$$

input `int(x/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)`

output `int(x/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)`

**3.347**  $\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.347.1 Optimal result . . . . . 2787  
 3.347.2 Mathematica [A] (verified) . . . . . 2787  
 3.347.3 Rubi [A] (verified) . . . . . 2788  
 3.347.4 Maple [A] (verified) . . . . . 2788  
 3.347.5 Fricas [B] (verification not implemented) . . . . . 2789  
 3.347.6 Sympy [F] . . . . . 2789  
 3.347.7 Maxima [F] . . . . . 2789  
 3.347.8 Giac [F] . . . . . 2790  
 3.347.9 Mupad [B] (verification not implemented) . . . . . 2790

**3.347.1 Optimal result**

Integrand size = 25, antiderivative size = 37

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))}$$

output `-(c*x-1)^(1/2)/b/c/(a+b*arccosh(c*x))/(-c*x+1)^(1/2)`

**3.347.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.35

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}}{bc\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))}$$

input `Integrate[1/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `-((Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(b*c*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])))`

### 3.347.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

↓ 6307

$$\frac{\sqrt{cx-1}}{bc\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))}$$

input `Int[1/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2),x]`

output `-(Sqrt[-1 + c*x]/(b*c*Sqrt[1 - c*x]*(a + b*ArcCosh[c*x])))`

#### 3.347.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x])/Sqrt[d + e*x^2]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.347.4 Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.54

method	result	size
default	$\frac{\sqrt{-(cx-1)(cx+1)}\sqrt{cx-1}\sqrt{cx+1}}{(c^2x^2-1)c(a+b\operatorname{arccosh}(cx))^b}$	57

input `int(1/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `((-c*x-1)*(c*x+1))^(1/2)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(c^2*x^2-1)/c/(a+b*arccosh(c*x))/b`

---

3.347.  $\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.347.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 75 vs.  $2(33) = 66$ .

Time = 0.26 (sec) , antiderivative size = 75, normalized size of antiderivative = 2.03

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \frac{\sqrt{c^2x^2-1}\sqrt{-c^2x^2+1}}{abc^3x^2-abc+(b^2c^3x^2-b^2c)\log(cx+\sqrt{c^2x^2-1})}$$

input `integrate(1/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `sqrt(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)/(a*b*c^3*x^2 - a*b*c + (b^2*c^3*x^2 - b^2*c)*log(c*x + sqrt(c^2*x^2 - 1)))`

**3.347.6 Sympy [F]**

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(1/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2),x)`

output `Integral(1/(sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2), x)`

**3.347.7 Maxima [F]**

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(1/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

```
output -(c^3*x^3 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*x)/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate(-(c^2*x^2 - (c*x + 1)*(c*x - 1) - 1)/(((c*x + 1)^(3/2)*(c*x - 1)*b^2*c^2*x^2 + 2*(b^2*c^3*x^3 - b^2*c*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x - 1)*a*b*c^2*x^2 + 2*(a*b*c^3*x^3 - a*b*c*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^4*x^4 - 2*a*b*c^2*x^2 + a*b)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)
```

### 3.347.8 Giac [F]

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

```
input integrate(1/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")
```

```
output integrate(1/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^2), x)
```

### 3.347.9 Mupad [B] (verification not implemented)

Time = 3.35 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.59

$$\int \frac{1}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{b\sqrt{1-c^2x^2}\sqrt{cx-1}\sqrt{cx+1}}{c(a+b\operatorname{acosh}(cx))(b^2-b^2c^2x^2)}$$

```
input int(1/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)
```

```
output -(b*(1 - c^2*x^2)^(1/2)*(c*x - 1)^(1/2)*(c*x + 1)^(1/2))/(c*(a + b*acosh(c*x))*(b^2 - b^2*c^2*x^2))
```

**3.348**  $\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.348.1 Optimal result . . . . . 2791  
 3.348.2 Mathematica [N/A] . . . . . 2791  
 3.348.3 Rubi [N/A] . . . . . 2792  
 3.348.4 Maple [N/A] (verified) . . . . . 2793  
 3.348.5 Fricas [N/A] . . . . . 2793  
 3.348.6 Sympy [N/A] . . . . . 2793  
 3.348.7 Maxima [N/A] . . . . . 2794  
 3.348.8 Giac [F(-2)] . . . . . 2794  
 3.348.9 Mupad [N/A] . . . . . 2795

**3.348.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}}{bcx\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{\sqrt{-1+cx}\operatorname{Int}\left(\frac{1}{x^2(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{1-cx}}$$

output `-(c*x-1)^(1/2)/b/c/x/(a+b*arccosh(c*x))/(-c*x+1)^(1/2)-(c*x-1)^(1/2)*Unintegrable(1/x^2/(a+b*arccosh(c*x)),x)/b/c/(-c*x+1)^(1/2)`

**3.348.2 Mathematica [N/A]**

Not integrable

Time = 4.23 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/(x*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/(x*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`



**3.348.3 Rubi [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6365, 6303}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx$$

↓ 6365

$$-\frac{\sqrt{cx-1} \int \frac{1}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{\sqrt{cx-1}}{bcx\sqrt{1-cx}(a+\operatorname{barccosh}(cx))}$$

↓ 6303

$$-\frac{\sqrt{cx-1} \int \frac{1}{x^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{\sqrt{cx-1}}{bcx\sqrt{1-cx}(a+\operatorname{barccosh}(cx))}$$

input `Int[1/(x*sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.348.3.1 Defintions of rubi rules used**

rule 6303 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Unintegrable[(d*x)^m*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, m, n}, x]`

rule 6365 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.))/sqrt[(d_) + (e_.)*(x_)^2], x_Symbol]
:> Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])], x] - Simp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]
Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]`

**3.348.4 Maple [N/A] (verified)**

Not integrable

Time = 1.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x(a + b \operatorname{arccosh}(cx))^2 \sqrt{-c^2x^2 + 1}} dx$$

input `int(1/x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x)`output `int(1/x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x)`**3.348.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.86

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arccosh}(cx)+a)^2x} dx$$

input `integrate(1/x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`output `integral(-sqrt(-c^2*x^2+1)/(a^2*c^2*x^3-a^2*x+(b^2*c^2*x^3-b^2*x)*arccosh(c*x)^2+2*(a*b*c^2*x^3-a*b*x)*arccosh(c*x)),x)`**3.348.6 Sympy [N/A]**

Not integrable

Time = 7.06 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x\sqrt{-(cx-1)(cx+1)}(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(1/x/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2),x)`output `Integral(1/(x*sqrt(-(c*x-1)*(c*x+1))*(a+b*acosh(c*x))**2),x)`

---

3.348.  $\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.348.7 Maxima [N/A]**

Not integrable

Time = 0.83 (sec) , antiderivative size = 481, normalized size of antiderivative = 17.18

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-c^2x^2+1}(b\operatorname{arccosh}(cx)+a)^2x} dx$$

input `integrate(1/x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `-(c^3*x^3 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*x)/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^2 + (b^2*c^3*x^3 - b^2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^2 + (a*b*c^3*x^3 - a*b*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)) - integrate((c^5*x^5 - c^3*x^3 + (c^3*x^3 - 2*c*x)*(c*x + 1)*(c*x - 1) + (2*c^4*x^4 - 3*c^2*x^2 + 1)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)^(3/2)*(c*x - 1)*b^2*c^3*x^4 + 2*(b^2*c^4*x^5 - b^2*c^2*x^3)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^6 - 2*b^2*c^3*x^4 + b^2*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x - 1)*a*b*c^3*x^4 + 2*(a*b*c^4*x^5 - a*b*c^2*x^3)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^6 - 2*a*b*c^3*x^4 + a*b*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

**3.348.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(1/x/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.348.9 Mupad [N/A]**

Not integrable

Time = 3.15 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x(a+b\operatorname{acosh}(cx))^2\sqrt{1-c^2x^2}} dx$$

input `int(1/(x*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)`output `int(1/(x*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)`

**3.349**  $\int \frac{1}{x^2\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.349.1 Optimal result . . . . . 2796  
 3.349.2 Mathematica [N/A] . . . . . 2796  
 3.349.3 Rubi [N/A] . . . . . 2797  
 3.349.4 Maple [N/A] (verified) . . . . . 2798  
 3.349.5 Fracas [N/A] . . . . . 2798  
 3.349.6 Sympy [N/A] . . . . . 2798  
 3.349.7 Maxima [N/A] . . . . . 2799  
 3.349.8 Giac [N/A] . . . . . 2799  
 3.349.9 Mupad [N/A] . . . . . 2800

**3.349.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x^2\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}}{bcx^2\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))} - \frac{2\sqrt{-1+cx}\operatorname{Int}\left(\frac{1}{x^3(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{1-cx}}$$

output `-(c*x-1)^(1/2)/b/c/x^2/(a+b*arccosh(c*x))/(-c*x+1)^(1/2)-2*(c*x-1)^(1/2)*U  
 nintegrable(1/x^3/(a+b*arccosh(c*x)),x)/b/c/(-c*x+1)^(1/2)`

**3.349.2 Mathematica [N/A]**

Not integrable

Time = 1.42 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x^2\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x^2\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/(x^2*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/(x^2*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

**3.349.3 Rubi [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6365, 6303}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt{1-c^2x^2} (a + b \operatorname{arccosh}(cx))^2} dx$$

↓ 6365

$$-\frac{2\sqrt{cx-1} \int \frac{1}{x^3(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{\sqrt{cx-1}}{bcx^2\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))}$$

↓ 6303

$$-\frac{2\sqrt{cx-1} \int \frac{1}{x^3(a+b\operatorname{arccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{\sqrt{cx-1}}{bcx^2\sqrt{1-cx}(a+b\operatorname{arccosh}(cx))}$$

input `Int[1/(x^2*Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.349.3.1 Defintions of rubi rules used**

rule 6303 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Unintegrable[(d*x)^m*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, m, n}, x]`

rule 6365 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.))/Sqrt[(d_.) + (e_.)*(x_)^2], x_Symbol]
:> Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])], x] - Simp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]
Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]`

**3.349.4 Maple [N/A] (verified)**

Not integrable

Time = 0.99 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^2 (a + b \operatorname{arccosh}(cx))^2 \sqrt{-c^2 x^2 + 1}} dx$$

input `int(1/x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x)`output `int(1/x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x)`**3.349.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 86, normalized size of antiderivative = 3.07

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-c^2 x^2 + 1} (b \operatorname{arccosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`output `integral(-sqrt(-c^2*x^2 + 1)/(a^2*c^2*x^4 - a^2*x^2 + (b^2*c^2*x^4 - b^2*x^2)*arccosh(c*x)^2 + 2*(a*b*c^2*x^4 - a*b*x^2)*arccosh(c*x)), x)`**3.349.6 Sympy [N/A]**

Not integrable

Time = 21.61 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.11

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x^2 \sqrt{-(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate(1/x**2/(a+b*acosh(c*x))**2/(-c**2*x**2+1)**(1/2),x)`output `Integral(1/(x**2*sqrt(-(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**2), x)`

---

3.349.  $\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))^2} dx$

**3.349.7 Maxima [N/A]**

Not integrable

Time = 0.83 (sec) , antiderivative size = 491, normalized size of antiderivative = 17.54

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-c^2 x^2 + 1} (b \operatorname{arccosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `-(c^3*x^3 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*x)/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^3 + (b^2*c^3*x^4 - b^2*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^3 + (a*b*c^3*x^4 - a*b*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)) - integrate((2*c^5*x^5 - 3*c^3*x^3 + (2*c^3*x^3 - 3*c*x)*(c*x + 1)*(c*x - 1) + 2*(2*c^4*x^4 - 3*c^2*x^2 + 1)*sqrt(c*x + 1)*sqrt(c*x - 1) + c*x)/(((c*x + 1)^(3/2)*(c*x - 1)*b^2*c^3*x^5 + 2*(b^2*c^4*x^6 - b^2*c^2*x^4)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^7 - 2*b^2*c^3*x^5 + b^2*c*x^3)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x - 1)*a*b*c^3*x^5 + 2*(a*b*c^4*x^6 - a*b*c^2*x^4)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^7 - 2*a*b*c^3*x^5 + a*b*c*x^3)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

**3.349.8 Giac [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{\sqrt{-c^2 x^2 + 1} (b \operatorname{arccosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(a+b*arccosh(c*x))^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^2*x^2), x)`



**3.349.9 Mupad [N/A]**

Not integrable

Time = 3.43 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 \sqrt{1 - c^2 x^2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x^2 (a + b \operatorname{acosh}(cx))^2 \sqrt{1 - c^2 x^2}} dx$$

input `int(1/(x^2*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)`output `int(1/(x^2*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)`

**3.350** 
$$\int \frac{x^3}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.350.1 Optimal result	2801
3.350.2 Mathematica [N/A]	2801
3.350.3 Rubi [N/A]	2802
3.350.4 Maple [N/A] (verified)	2802
3.350.5 Fricas [N/A]	2803
3.350.6 Sympy [N/A]	2803
3.350.7 Maxima [N/A]	2803
3.350.8 Giac [F(-2)]	2804
3.350.9 Mupad [N/A]	2804

**3.350.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^3}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{x^3}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2}, x\right)$$

output `Unintegrable(x^3/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

**3.350.2 Mathematica [N/A]**

Not integrable

Time = 24.45 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^3}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$$

input `Integrate[x^3/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2),x]`

output `Integrate[x^3/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.350.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$$

input `Int[x^3/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.350.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n_.*((f_.)*(x_.))^m_.*((d_.) + (e_.)*(x_.^2))^p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.350.4 Maple [N/A] (verified)**

Not integrable

Time = 1.57 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^3}{(-c^2 x^2 + 1)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(x^3/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int(x^3/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

---

3.350.  $\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$

**3.350.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 106, normalized size of antiderivative = 3.79

$$\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^3/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x^3/(a^2*c^4*x^4 - 2*a^2*c^2*x^2 + (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*arccosh(c*x)^2 + a^2 + 2*(a*b*c^4*x^4 - 2*a*b*c^2*x^2 + a*b)*arccosh(c*x)), x)`

**3.350.6 Sympy [N/A]**

Not integrable

Time = 64.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(-(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate(x**3/((-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2), x)`

output `Integral(x**3/((-c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2), x)`

**3.350.7 Maxima [N/A]**

Not integrable

Time = 1.28 (sec) , antiderivative size = 531, normalized size of antiderivative = 18.96

$$\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^3/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `(c*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*x^3)/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) - integrate((c^5*x^7 - 5*c^3*x^5 + 4*c*x^3 + (c^3*x^5 - 2*c*x^3)*(c*x + 1)*(c*x - 1) + (2*c^4*x^6 - 7*c^2*x^4 + 3*x^2)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((b^2*c^5*x^4 - b^2*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^6*x^5 - 2*b^2*c^4*x^3 + b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^7*x^6 - 3*b^2*c^5*x^4 + 3*b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((a*b*c^5*x^4 - a*b*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^6*x^5 - 2*a*b*c^4*x^3 + a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^7*x^6 - 3*a*b*c^5*x^4 + 3*a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

### 3.350.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.350.9 Mupad [N/A]

Not integrable

Time = 3.46 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{3/2}} dx$$

---

3.350.  $\int \frac{x^3}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$

input `int(x^3/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)),x)`

output `int(x^3/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)), x)`

---

3.350.  $\int \frac{x^3}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$

### 3.351 $\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.351.1 Optimal result . . . . .	2806
3.351.2 Mathematica [N/A] . . . . .	2806
3.351.3 Rubi [N/A] . . . . .	2807
3.351.4 Maple [N/A] (verified) . . . . .	2808
3.351.5 Fricas [N/A] . . . . .	2808
3.351.6 Sympy [N/A] . . . . .	2809
3.351.7 Maxima [N/A] . . . . .	2809
3.351.8 Giac [N/A] . . . . .	2810
3.351.9 Mupad [N/A] . . . . .	2810

#### 3.351.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^2\sqrt{-1+cx}\sqrt{1+cx}}{bc(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} + \frac{2\sqrt{-1+cx}\operatorname{Int}\left(\frac{x}{(-1+c^2x^2)^2(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{1-cx}}$$

```
output -x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))
+2*(c*x-1)^(1/2)*Unintegrable(x/(c^2*x^2-1)^2/(a+b*arccosh(c*x)),x)/b/c/(-
c*x+1)^(1/2)
```

#### 3.351.2 Mathematica [N/A]

Not integrable

Time = 5.62 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

```
input Integrate[x^2/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]
```

```
output Integrate[x^2/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]
```

**3.351.3 Rubi [N/A]**

Not integrable

Time = 0.71 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6355, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$$

↓ 6355

$$\frac{2\sqrt{cx-1} \int \frac{x}{(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))}$$

↓ 6327

$$\frac{2\sqrt{cx-1} \int \frac{x}{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))}$$

↓ 6375

$$\frac{2\sqrt{cx-1} \int \frac{x}{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))}$$

input `Int[x^2/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.351.3.1 Defintions of rubi rules used**

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] :> Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

---

3.351.  $\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$



```
rule 6355 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && EqQ[m + 2*p + 1, 0]
```

```
rule 6375 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]
```

### 3.351.4 Maple [N/A] (verified)

Not integrable

Time = 0.54 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{(-c^2x^2 + 1)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^2} dx$$

```
input int(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)
```

```
output int(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)
```

### 3.351.5 Fracas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 106, normalized size of antiderivative = 3.79

$$\int \frac{x^2}{(1 - c^2x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{(-c^2x^2 + 1)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^2} dx$$

```
input integrate(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fracas")
```

```
output integral(sqrt(-c^2*x^2 + 1)*x^2/(a^2*c^4*x^4 - 2*a^2*c^2*x^2 + (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*arccosh(c*x)^2 + a^2 + 2*(a*b*c^4*x^4 - 2*a*b*c^2*x^2 + a*b)*arccosh(c*x)), x)
```

---

3.351.  $\int \frac{x^2}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.351.6 Sympy [N/A]**

Not integrable

Time = 71.19 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^2}{(- (cx - 1) (cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate(x**2/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`output `Integral(x**2/((-c*x - 1)*(c*x + 1))**3/2*(a + b*acosh(c*x))**2, x)`**3.351.7 Maxima [N/A]**

Not integrable

Time = 0.95 (sec) , antiderivative size = 503, normalized size of antiderivative = 17.96

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`output `(c*x^3 + sqrt(c*x + 1)*sqrt(c*x - 1)*x^2)/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((3*c^3*x^4 + (c*x + 1)*(c*x - 1)*c*x^2 - 3*c*x^2 + 2*(2*c^2*x^3 - x)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((b^2*c^5*x^4 - b^2*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^6*x^5 - 2*b^2*c^4*x^3 + b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^7*x^6 - 3*b^2*c^5*x^4 + 3*b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((a*b*c^5*x^4 - a*b*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^6*x^5 - 2*a*b*c^4*x^3 + a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^7*x^6 - 3*a*b*c^5*x^4 + 3*a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

**3.351.8 Giac [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`output `integrate(x^2/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)^2), x)`**3.351.9 Mupad [N/A]**

Not integrable

Time = 3.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{(a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{3/2}} dx$$

input `int(x^2/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)),x)`output `int(x^2/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)), x)`

**3.352** 
$$\int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.352.1 Optimal result . . . . .	2811
3.352.2 Mathematica [N/A] . . . . .	2811
3.352.3 Rubi [N/A] . . . . .	2812
3.352.4 Maple [N/A] (verified) . . . . .	2812
3.352.5 Fricas [N/A] . . . . .	2813
3.352.6 Sympy [N/A] . . . . .	2813
3.352.7 Maxima [N/A] . . . . .	2813
3.352.8 Giac [F(-2)] . . . . .	2814
3.352.9 Mupad [N/A] . . . . .	2814

**3.352.1 Optimal result**

Integrand size = 26, antiderivative size = 26

$$\int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

**3.352.2 Mathematica [N/A]**

Not integrable

Time = 19.57 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[x/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[x/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.352.3 Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$$

input `Int[x/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.352.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.352.4 Maple [N/A] (verified)**

Not integrable

Time = 1.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{x}{(-c^2 x^2 + 1)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

**3.352.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 104, normalized size of antiderivative = 4.00

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*x/(a^2*c^4*x^4 - 2*a^2*c^2*x^2 + (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*arccosh(c*x)^2 + a^2 + 2*(a*b*c^4*x^4 - 2*a*b*c^2*x^2 + a*b)*arccosh(c*x)), x)`

**3.352.6 Sympy [N/A]**

Not integrable

Time = 59.91 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.04

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x}{(-(cx - 1)(cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate(x/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(x/((-c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2), x)`

**3.352.7 Maxima [N/A]**

Not integrable

Time = 0.93 (sec) , antiderivative size = 511, normalized size of antiderivative = 19.65

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `(c*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*x)/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((c^5*x^5 + (c*x + 1)*(c*x - 1)*c^3*x^3 + c^3*x^3 + (2*c^4*x^4 + c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - 2*c*x)/(((b^2*c^5*x^4 - b^2*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^6*x^5 - 2*b^2*c^4*x^3 + b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^7*x^6 - 3*b^2*c^5*x^4 + 3*b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((a*b*c^5*x^4 - a*b*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^6*x^5 - 2*a*b*c^4*x^3 + a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^7*x^6 - 3*a*b*c^5*x^4 + 3*a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

### 3.352.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.352.9 Mupad [N/A]

Not integrable

Time = 3.59 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x}{(a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{3/2}} dx$$

input `int(x/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)),x)`

---

3.352.  $\int \frac{x}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$

output `int(x/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)), x)`

---

3.352.  $\int \frac{x}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$



**3.353** 
$$\int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.353.1 Optimal result	2816
3.353.2 Mathematica [N/A]	2816
3.353.3 Rubi [N/A]	2817
3.353.4 Maple [N/A] (verified)	2818
3.353.5 Fricas [N/A]	2818
3.353.6 Sympy [N/A]	2819
3.353.7 Maxima [N/A]	2819
3.353.8 Giac [N/A]	2820
3.353.9 Mupad [N/A]	2820

**3.353.1 Optimal result**

Integrand size = 25, antiderivative size = 25

$$\int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}}{bc(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))} + \frac{2c\sqrt{-1+cx}\operatorname{Int}\left(\frac{x}{(-1+c^2x^2)^2(a+b\operatorname{arccosh}(cx))}, x\right)}{b\sqrt{1-cx}}$$

output `-(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))+2*c*(c*x-1)^(1/2)*Unintegrable(x/(c^2*x^2-1)^2/(a+b*arccosh(c*x)),x)/b/(-c*x+1)^(1/2)`

**3.353.2 Mathematica [N/A]**

Not integrable

Time = 3.88 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.353.3 Rubi [N/A]**

Not integrable

Time = 0.61 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx$$

↓ 6319

$$\frac{2c\sqrt{cx-1} \int \frac{x}{(1-cx)^2(cx+1)^2(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{1-cx}} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))}$$

↓ 6327

$$\frac{2c\sqrt{cx-1} \int \frac{x}{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{1-cx}} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))}$$

↓ 6375

$$\frac{2c\sqrt{cx-1} \int \frac{x}{(1-c^2x^2)^2(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{1-cx}} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))}$$

input `Int[1/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.353.3.1 Defintions of rubi rules used**

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n*((d_) + (e_.)*(x_)^2)^(p_.), x  
_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] :> Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

### 3.353.4 Maple [N/A] (verified)

Not integrable

Time = 1.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{1}{(-c^2x^2 + 1)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

### 3.353.5 Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 4.12

$$\int \frac{1}{(1 - c^2x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(-c^2x^2 + 1)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(a^2*c^4*x^4 - 2*a^2*c^2*x^2 + (b^2*c^4*x^4 - 2*b^2*c^2*x^2 + b^2)*arccosh(c*x)^2 + a^2 + 2*(a*b*c^4*x^4 - 2*a*b*c^2*x^2 + a*b)*arccosh(c*x)), x)`

**3.353.6 Sympy [N/A]**

Not integrable

Time = 47.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(- (cx - 1) (cx + 1))^{\frac{3}{2}} (a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate(1/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`output `Integral(1/((-c*x - 1)*(c*x + 1))**(3/2)*(a + b*acosh(c*x))**2), x)`**3.353.7 Maxima [N/A]**

Not integrable

Time = 0.93 (sec) , antiderivative size = 502, normalized size of antiderivative = 20.08

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`output `(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((2*c^4*x^4 - c^2*x^2 + (2*c^2*x^2 - 1)*(c*x + 1)*(c*x - 1) + 2*(2*c^3*x^3 - c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) - 1)/(((b^2*c^4*x^4 - b^2*c^2*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^5*x^5 - 2*b^2*c^3*x^3 + b^2*c*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^6*x^6 - 3*b^2*c^4*x^4 + 3*b^2*c^2*x^2 - b^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1) + ((a*b*c^4*x^4 - a*b*c^2*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^5*x^5 - 2*a*b*c^3*x^3 + a*b*c*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^6*x^6 - 3*a*b*c^4*x^4 + 3*a*b*c^2*x^2 - a*b)*sqrt(c*x + 1))*sqrt(-c*x + 1))), x)`

**3.353.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`output `integrate(1/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)^2), x)`**3.353.9 Mupad [N/A]**

Not integrable

Time = 3.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{3/2}} dx$$

input `int(1/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)),x)`output `int(1/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)), x)`

$$3.354 \quad \int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.354.1 Optimal result	. . . . .	2821
3.354.2 Mathematica [N/A]	. . . . .	2821
3.354.3 Rubi [N/A]	. . . . .	2822
3.354.4 Maple [N/A] (verified)	. . . . .	2822
3.354.5 Fricas [N/A]	. . . . .	2823
3.354.6 Sympy [N/A]	. . . . .	2823
3.354.7 Maxima [N/A]	. . . . .	2823
3.354.8 Giac [F(-2)]	. . . . .	2824
3.354.9 Mupad [N/A]	. . . . .	2824

### 3.354.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

### 3.354.2 Mathematica [N/A]

Not integrable

Time = 21.38 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/(x*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/(x*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

---


$$3.354. \quad \int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.354.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\text{barccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\text{barccosh}(cx))^2} dx$$

input `Int[1/(x*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.354.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(x_.)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.354.4 Maple [N/A] (verified)**

Not integrable

Time = 1.40 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x(-c^2x^2+1)^{\frac{3}{2}}(a+b\text{arccosh}(cx))^2} dx$$

input `int(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

**3.354.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 108, normalized size of antiderivative = 3.86

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2x^2+1)^{3/2}(b\operatorname{arcosh}(cx)+a)^2x} dx$$

input `integrate(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output `integral(sqrt(-c^2*x^2+1)/(a^2*c^4*x^5-2*a^2*c^2*x^3+a^2*x+(b^2*c^4*x^5-2*b^2*c^2*x^3+b^2*x)*arccosh(c*x)^2+2*(a*b*c^4*x^5-2*a*b*c^2*x^3+a*b*x)*arccosh(c*x)), x)`

**3.354.6 Sympy [N/A]**

Not integrable

Time = 123.72 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{x(-cx-1)(cx+1)^{3/2}(a+b\operatorname{acosh}(cx))^2} dx$$

input `integrate(1/x/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(1/(x*(-(c*x-1)*(c*x+1))**(3/2)*(a+b*acosh(c*x))**2), x)`

**3.354.7 Maxima [N/A]**

Not integrable

Time = 1.27 (sec) , antiderivative size = 530, normalized size of antiderivative = 18.93

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2x^2+1)^{3/2}(b\operatorname{arcosh}(cx)+a)^2x} dx$$



input `integrate(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x^2 + (b^2*c^3*x^3 - b^2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x^2 + (a*b*c^3*x^3 - a*b*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((3*c^5*x^5 - 3*c^3*x^3 + (3*c^3*x^3 - 2*c*x)*(c*x + 1)*(c*x - 1) + (6*c^4*x^4 - 5*c^2*x^2 + 1)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((b^2*c^5*x^6 - b^2*c^3*x^4)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^6*x^7 - 2*b^2*c^4*x^5 + b^2*c^2*x^3)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^7*x^8 - 3*b^2*c^5*x^6 + 3*b^2*c^3*x^4 - b^2*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((a*b*c^5*x^6 - a*b*c^3*x^4)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^6*x^7 - 2*a*b*c^4*x^5 + a*b*c^2*x^3)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^7*x^8 - 3*a*b*c^5*x^6 + 3*a*b*c^3*x^4 - a*b*c*x^2)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

### 3.354.8 Giac [F(-2)]

Exception generated.

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(1/x/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.354.9 Mupad [N/A]

Not integrable

Time = 3.57 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x(a+b\operatorname{acosh}(cx))^2(1-c^2x^2)^{3/2}} dx$$

---

3.354.  $\int \frac{1}{x(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$

input `int(1/(x*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)),x)`

output `int(1/(x*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)), x)`

$$3.355 \quad \int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.355.1 Optimal result	2826
3.355.2 Mathematica [N/A]	2826
3.355.3 Rubi [N/A]	2827
3.355.4 Maple [N/A] (verified)	2827
3.355.5 Fricas [N/A]	2828
3.355.6 Sympy [F(-1)]	2828
3.355.7 Maxima [N/A]	2828
3.355.8 Giac [N/A]	2829
3.355.9 Mupad [N/A]	2829

### 3.355.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

### 3.355.2 Mathematica [N/A]

Not integrable

Time = 17.59 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/(x^2*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/(x^2*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

---


$$3.355. \quad \int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.355.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `Int[1/(x^2*(1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.355.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(x_.)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.355.4 Maple [N/A] (verified)**

Not integrable

Time = 1.30 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^2 (-c^2 x^2 + 1)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

**3.355.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 114, normalized size of antiderivative = 4.07

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)/(a^2*c^4*x^6 - 2*a^2*c^2*x^4 + a^2*x^2 + (b^2*c^4*x^6 - 2*b^2*c^2*x^4 + b^2*x^2)*arccosh(c*x)^2 + 2*(a*b*c^4*x^6 - 2*a*b*c^2*x^4 + a*b*x^2)*arccosh(c*x)), x)`

**3.355.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(1/x**2/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.355.7 Maxima [N/A]**

Not integrable

Time = 1.29 (sec) , antiderivative size = 538, normalized size of antiderivative = 19.21

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1})/(((c*x + 1)*\sqrt{c*x - 1}*b^2*c^2*x^3 + (b^2*c^3*x^4 - b^2*c*x^2)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((c*x + 1)*\sqrt{c*x - 1}*a*b*c^2*x^3 + (a*b*c^3*x^4 - a*b*c*x^2)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}) + \text{integrate}((4*c^5*x^5 - 5*c^3*x^3 + (4*c^3*x^3 - 3*c*x)*(c*x + 1)*(c*x - 1) + 2*(4*c^4*x^4 - 4*c^2*x^2 + 1)*\sqrt{c*x + 1}*\sqrt{c*x - 1} + c*x)/(((b^2*c^5*x^7 - b^2*c^3*x^5)*(c*x + 1)^{(3/2)}*(c*x - 1) + 2*(b^2*c^6*x^8 - 2*b^2*c^4*x^6 + b^2*c^2*x^4)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^7*x^9 - 3*b^2*c^5*x^7 + 3*b^2*c^3*x^5 - b^2*c*x^3)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^5*x^7 - a*b*c^3*x^5)*(c*x + 1)^{(3/2)}*(c*x - 1) + 2*(a*b*c^6*x^8 - 2*a*b*c^4*x^6 + a*b*c^2*x^4)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^7*x^9 - 3*a*b*c^5*x^7 + 3*a*b*c^3*x^5 - a*b*c*x^3)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}), x)$

### 3.355.8 Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(-c^2x^2+1)^{\frac{3}{2}}(b\operatorname{arcosh}(cx)+a)^2x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(1/((-c^2*x^2 + 1)^(3/2)*(b*arccosh(c*x) + a)^2*x^2), x)`

### 3.355.9 Mupad [N/A]

Not integrable

Time = 3.57 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x^2(a+b\operatorname{acosh}(cx))^2(1-c^2x^2)^{3/2}} dx$$

input `int(1/(x^2*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)),x)`

output `int(1/(x^2*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)), x)`

---

3.355.  $\int \frac{1}{x^2(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.356** 
$$\int \frac{x^4}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.356.1 Optimal result . . . . .	2830
3.356.2 Mathematica [N/A] . . . . .	2830
3.356.3 Rubi [N/A] . . . . .	2831
3.356.4 Maple [N/A] (verified) . . . . .	2832
3.356.5 Fricas [N/A] . . . . .	2833
3.356.6 Sympy [F(-1)] . . . . .	2833
3.356.7 Maxima [N/A] . . . . .	2833
3.356.8 Giac [N/A] . . . . .	2834
3.356.9 Mupad [N/A] . . . . .	2834

**3.356.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^4}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{x^4\sqrt{-1+cx}\sqrt{1+cx}}{bc(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))} - \frac{4\sqrt{-1+cx}\operatorname{Int}\left(\frac{x^3}{(-1+c^2x^2)^3(a+b\operatorname{arccosh}(cx))}, x\right)}{bc\sqrt{1-cx}}$$

output `-x^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x)) - 4*(c*x-1)^(1/2)*Unintegrable(x^3/(c^2*x^2-1)^3/(a+b*arccosh(c*x)),x)/b/c/(-c*x+1)^(1/2)`

**3.356.2 Mathematica [N/A]**

Not integrable

Time = 5.07 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^4}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x^4}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[x^4/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[x^4/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

---

3.356. 
$$\int \frac{x^4}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.356.3 Rubi [N/A]**

Not integrable

Time = 0.83 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6355, 25, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6355} \\
 & -\frac{4\sqrt{cx-1} \int -\frac{x^3}{(1-cx)^3(cx+1)^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{4\sqrt{cx-1} \int \frac{x^3}{(1-cx)^3(cx+1)^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & \frac{4\sqrt{cx-1} \int \frac{x^3}{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6375} \\
 & \frac{4\sqrt{cx-1} \int \frac{x^3}{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))} dx}{bc\sqrt{1-cx}} - \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))}
 \end{aligned}$$

input `Int[x^4/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`



## 3.356.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6355 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && EqQ[m + 2*p + 1, 0]`
- rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

## 3.356.4 Maple [N/A] (verified)

Not integrable

Time = 1.46 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^4}{(-c^2x^2 + 1)^{\frac{5}{2}} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(x^4/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int(x^4/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.356.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 5.14

$$\int \frac{x^4}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^4}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^4/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^4/(a^2*c^6*x^6 - 3*a^2*c^4*x^4 + 3*a^2*c^2*x^2 + (b^2*c^6*x^6 - 3*b^2*c^4*x^4 + 3*b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^6*x^6 - 3*a*b*c^4*x^4 + 3*a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.356.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^4}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(x**4/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.356.7 Maxima [N/A]**

Not integrable

Time = 1.35 (sec) , antiderivative size = 614, normalized size of antiderivative = 21.93

$$\int \frac{x^4}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^4}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^4/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

---

3.356.  $\int \frac{x^4}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$

output  $-(c*x^5 + \sqrt{c*x + 1}*\sqrt{c*x - 1})*x^4/(((b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}) - \text{integrate}((5*c^3*x^6 + 3*(c*x + 1)*(c*x - 1)*c*x^4 - 5*c*x^4 + 4*(2*c^2*x^5 - x^3)*\sqrt{c*x + 1}*\sqrt{c*x - 1}))/(((b^2*c^7*x^6 - 2*b^2*c^5*x^4 + b^2*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^8*x^7 - 3*b^2*c^6*x^5 + 3*b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^9*x^8 - 4*b^2*c^7*x^6 + 6*b^2*c^5*x^4 - 4*b^2*c^3*x^2 + b^2*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^7*x^6 - 2*a*b*c^5*x^4 + a*b*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^8*x^7 - 3*a*b*c^6*x^5 + 3*a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^9*x^8 - 4*a*b*c^7*x^6 + 6*a*b*c^5*x^4 - 4*a*b*c^3*x^2 + a*b*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}), x)$

### 3.356.8 Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^4}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^4}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^4/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(x^4/((-c^2*x^2 + 1)^(5/2)*(b*arccosh(c*x) + a)^2), x)`

### 3.356.9 Mupad [N/A]

Not integrable

Time = 3.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^4}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^4}{(a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{5/2}} dx$$

input `int(x^4/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)`

output `int(x^4/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)`

---

3.356.  $\int \frac{x^4}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$

$$3.357 \quad \int \frac{x^3}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.357.1 Optimal result . . . . .	2835
3.357.2 Mathematica [N/A] . . . . .	2835
3.357.3 Rubi [N/A] . . . . .	2836
3.357.4 Maple [N/A] (verified) . . . . .	2836
3.357.5 Fricas [N/A] . . . . .	2837
3.357.6 Sympy [F(-1)] . . . . .	2837
3.357.7 Maxima [N/A] . . . . .	2837
3.357.8 Giac [F(-2)] . . . . .	2838
3.357.9 Mupad [N/A] . . . . .	2838

### 3.357.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^3}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{x^3}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}, x\right)$$

output `Unintegrable(x^3/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

### 3.357.2 Mathematica [N/A]

Not integrable

Time = 39.63 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^3}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$$

input `Integrate[x^3/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `Integrate[x^3/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.357.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{arccosh}(cx))^2} dx$$

input `Int[x^3/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.357.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.357.4 Maple [N/A] (verified)**

Not integrable

Time = 1.36 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^3}{(-c^2 x^2 + 1)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(x^3/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int(x^3/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

---

3.357.  $\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$

**3.357.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 5.14

$$\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^3/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^3/(a^2*c^6*x^6 - 3*a^2*c^4*x^4 + 3*a^2*c^2*x^2 + (b^2*c^6*x^6 - 3*b^2*c^4*x^4 + 3*b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^6*x^6 - 3*a*b*c^4*x^4 + 3*a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.357.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(x**3/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.357.7 Maxima [N/A]**

Not integrable

Time = 1.96 (sec) , antiderivative size = 637, normalized size of antiderivative = 22.75

$$\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^3/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

---

3.357.  $\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$

output `-(c*x^4 + sqrt(c*x + 1)*sqrt(c*x - 1)*x^3)/(((b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) - integrate((c^5*x^7 + 3*c^3*x^5 - 4*c*x^3 + (c^3*x^5 + 2*c*x^3)*(c*x + 1)*(c*x - 1) + (2*c^4*x^6 + 5*c^2*x^4 - 3*x^2)*sqrt(c*x + 1)*sqrt(c*x - 1))/(((b^2*c^7*x^6 - 2*b^2*c^5*x^4 + b^2*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^8*x^7 - 3*b^2*c^6*x^5 + 3*b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^9*x^8 - 4*b^2*c^7*x^6 + 6*b^2*c^5*x^4 - 4*b^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((a*b*c^7*x^6 - 2*a*b*c^5*x^4 + a*b*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^8*x^7 - 3*a*b*c^6*x^5 + 3*a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^9*x^8 - 4*a*b*c^7*x^6 + 6*a*b*c^5*x^4 - 4*a*b*c^3*x^2 + a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)`

### 3.357.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.357.9 Mupad [N/A]

Not integrable

Time = 3.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^3}{(a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{5/2}} dx$$

---

3.357.  $\int \frac{x^3}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$

input `int(x^3/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)`

output `int(x^3/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)`

---

3.357.  $\int \frac{x^3}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$



**3.358** 
$$\int \frac{x^2}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.358.1 Optimal result	2840
3.358.2 Mathematica [N/A]	2840
3.358.3 Rubi [N/A]	2841
3.358.4 Maple [N/A] (verified)	2841
3.358.5 Fricas [N/A]	2842
3.358.6 Sympy [F(-1)]	2842
3.358.7 Maxima [N/A]	2842
3.358.8 Giac [N/A]	2843
3.358.9 Mupad [N/A]	2843

**3.358.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{x^2}{(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{x^2}{(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.358.2 Mathematica [N/A]**

Not integrable

Time = 8.99 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{x^2}{(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2} dx = \int \frac{x^2}{(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[x^2/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `Integrate[x^2/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.358.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$$

input `Int[x^2/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.358.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.358.4 Maple [N/A] (verified)**

Not integrable

Time = 1.43 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{(-c^2 x^2 + 1)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int(x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

---

3.358.  $\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$

**3.358.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 5.14

$$\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x^2/(a^2*c^6*x^6 - 3*a^2*c^4*x^4 + 3*a^2*c^2*x^2 + (b^2*c^6*x^6 - 3*b^2*c^4*x^4 + 3*b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^6*x^6 - 3*a*b*c^4*x^4 + 3*a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.358.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(x**2/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.358.7 Maxima [N/A]**

Not integrable

Time = 1.40 (sec) , antiderivative size = 635, normalized size of antiderivative = 22.68

$$\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $-(c*x^3 + \sqrt{c*x + 1}*\sqrt{c*x - 1})*x^2)/(((b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}) - \text{integrate}((2*c^5*x^6 + c^3*x^4 + (2*c^3*x^4 + c*x^2)*(c*x + 1)*(c*x - 1) - 3*c*x^2 + 2*(2*c^4*x^5 + c^2*x^3 - x)*\sqrt{c*x + 1}*\sqrt{c*x - 1}))/(((b^2*c^7*x^6 - 2*b^2*c^5*x^4 + b^2*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^8*x^7 - 3*b^2*c^6*x^5 + 3*b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^9*x^8 - 4*b^2*c^7*x^6 + 6*b^2*c^5*x^4 - 4*b^2*c^3*x^2 + b^2*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^7*x^6 - 2*a*b*c^5*x^4 + a*b*c^3*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^8*x^7 - 3*a*b*c^6*x^5 + 3*a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^9*x^8 - 4*a*b*c^7*x^6 + 6*a*b*c^5*x^4 - 4*a*b*c^3*x^2 + a*b*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1})), x)$

### 3.358.8 Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx = \int \frac{x^2}{(-c^2 x^2 + 1)^{5/2} (b \text{ arcosh}(cx) + a)^2} dx$$

input `integrate(x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(x^2/((-c^2*x^2 + 1)^(5/2)*(b*arccosh(c*x) + a)^2), x)`

### 3.358.9 Mupad [N/A]

Not integrable

Time = 3.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx = \int \frac{x^2}{(a + b \text{ acosh}(cx))^2 (1 - c^2 x^2)^{5/2}} dx$$

input `int(x^2/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)`

output `int(x^2/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)`

---

3.358.  $\int \frac{x^2}{(1 - c^2 x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx$

**3.359** 
$$\int \frac{x}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.359.1 Optimal result	2844
3.359.2 Mathematica [N/A]	2844
3.359.3 Rubi [N/A]	2845
3.359.4 Maple [N/A] (verified)	2845
3.359.5 Fricas [N/A]	2846
3.359.6 Sympy [F(-1)]	2846
3.359.7 Maxima [N/A]	2846
3.359.8 Giac [F(-2)]	2847
3.359.9 Mupad [N/A]	2847

**3.359.1 Optimal result**

Integrand size = 26, antiderivative size = 26

$$\int \frac{x}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{x}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.359.2 Mathematica [N/A]**

Not integrable

Time = 35.32 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{x}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{x}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[x/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[x/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.359.3 Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{arccosh}(cx))^2} dx$$

input `Int[x/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.359.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.359.4 Maple [N/A] (verified)**

Not integrable

Time = 1.33 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{x}{(-c^2 x^2 + 1)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int(x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.359.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 142, normalized size of antiderivative = 5.46

$$\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*x/(a^2*c^6*x^6 - 3*a^2*c^4*x^4 + 3*a^2*c^2*x^2 + (b^2*c^6*x^6 - 3*b^2*c^4*x^4 + 3*b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^6*x^6 - 3*a*b*c^4*x^4 + 3*a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.359.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(x/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.359.7 Maxima [N/A]**

Not integrable

Time = 1.69 (sec) , antiderivative size = 623, normalized size of antiderivative = 23.96

$$\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

```
output -(c*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*x)/(((b^2*c^4*x^3 - b^2*c^2*x)*(c*x
+ 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*
sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((a*b*c^4*x^3 - a*
b*c^2*x)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*s
qrt(c*x + 1))*sqrt(-c*x + 1)) - integrate((3*c^5*x^5 + 3*(c*x + 1)*(c*x -
1)*c^3*x^3 - c^3*x^3 + (6*c^4*x^4 - c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x -
1) - 2*c*x)/(((b^2*c^7*x^6 - 2*b^2*c^5*x^4 + b^2*c^3*x^2)*(c*x + 1)^(3/2)*
(c*x - 1) + 2*(b^2*c^8*x^7 - 3*b^2*c^6*x^5 + 3*b^2*c^4*x^3 - b^2*c^2*x)*(c
*x + 1)*sqrt(c*x - 1) + (b^2*c^9*x^8 - 4*b^2*c^7*x^6 + 6*b^2*c^5*x^4 - 4*b
^2*c^3*x^2 + b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*
sqrt(c*x - 1)) + ((a*b*c^7*x^6 - 2*a*b*c^5*x^4 + a*b*c^3*x^2)*(c*x + 1)^(3
/2)*(c*x - 1) + 2*(a*b*c^8*x^7 - 3*a*b*c^6*x^5 + 3*a*b*c^4*x^3 - a*b*c^2*x
)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^9*x^8 - 4*a*b*c^7*x^6 + 6*a*b*c^5*x^4 -
4*a*b*c^3*x^2 + a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)
```

### 3.359.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

```
input integrate(x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.359.9 Mupad [N/A]

Not integrable

Time = 3.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{x}{(a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{5/2}} dx$$

```
input int(x/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)
```

```
output int(x/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)
```

---

3.359.  $\int \frac{x}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$



**3.360** 
$$\int \frac{1}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.360.1 Optimal result	2848
3.360.2 Mathematica [N/A]	2848
3.360.3 Rubi [N/A]	2849
3.360.4 Maple [N/A] (verified)	2850
3.360.5 Fricas [N/A]	2851
3.360.6 Sympy [F(-1)]	2851
3.360.7 Maxima [N/A]	2851
3.360.8 Giac [N/A]	2852
3.360.9 Mupad [N/A]	2852

**3.360.1 Optimal result**

Integrand size = 25, antiderivative size = 25

$$\int \frac{1}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}}{bc(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))} - \frac{4c\sqrt{-1+cx}\operatorname{Int}\left(\frac{x}{(-1+c^2x^2)^3(a+b\operatorname{arccosh}(cx))}, x\right)}{b\sqrt{1-cx}}$$

output `-(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))-4*c*(c*x-1)^(1/2)*Unintegrable(x/(c^2*x^2-1)^3/(a+b*arccosh(c*x)),x)/b/(-c*x+1)^(1/2)`

**3.360.2 Mathematica [N/A]**

Not integrable

Time = 4.65 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

$$\int \frac{1}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.360.3 Rubi [N/A]**

Not integrable

Time = 0.63 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 25, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{4c\sqrt{cx-1} \int -\frac{x}{(1-cx)^3(cx+1)^3(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{1-cx}} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{25} \\
 & \frac{4c\sqrt{cx-1} \int \frac{x}{(1-cx)^3(cx+1)^3(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{1-cx}} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6327} \\
 & \frac{4c\sqrt{cx-1} \int \frac{x}{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{1-cx}} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{6375} \\
 & \frac{4c\sqrt{cx-1} \int \frac{x}{(1-c^2x^2)^3(a+\operatorname{barccosh}(cx))} dx}{b\sqrt{1-cx}} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))}
 \end{aligned}$$

input `Int[1/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

## 3.360.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_)^m_)*((d1_) + (e1_.)*(x_)^p_)*((d2_) + (e2_.)*(x_)^p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_)^m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

## 3.360.4 Maple [N/A] (verified)

Not integrable

Time = 1.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{1}{(-c^2x^2 + 1)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`output `int(1/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.360.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 141, normalized size of antiderivative = 5.64

$$\int \frac{1}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)/(a^2*c^6*x^6 - 3*a^2*c^4*x^4 + 3*a^2*c^2*x^2 + (b^2*c^6*x^6 - 3*b^2*c^4*x^4 + 3*b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^6*x^6 - 3*a*b*c^4*x^4 + 3*a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.360.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(1/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.360.7 Maxima [N/A]**

Not integrable

Time = 1.33 (sec) , antiderivative size = 609, normalized size of antiderivative = 24.36

$$\int \frac{1}{(1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $-(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1})/(((b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}) - \text{integrate}((4*c^4*x^4 - 3*c^2*x^2 + (4*c^2*x^2 - 1)*(c*x + 1)*(c*x - 1) + 4*(2*c^3*x^3 - c*x)*\sqrt{c*x + 1}*\sqrt{c*x - 1})/(((b^2*c^6*x^6 - 2*b^2*c^4*x^4 + b^2*c^2*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^7*x^7 - 3*b^2*c^5*x^5 + 3*b^2*c^3*x^3 - b^2*c*x)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^8*x^8 - 4*b^2*c^6*x^6 + 6*b^2*c^4*x^4 - 4*b^2*c^2*x^2 + b^2)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^6*x^6 - 2*a*b*c^4*x^4 + a*b*c^2*x^2)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^7*x^7 - 3*a*b*c^5*x^5 + 3*a*b*c^3*x^3 - a*b*c*x)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^8*x^8 - 4*a*b*c^6*x^6 + 6*a*b*c^4*x^4 - 4*a*b*c^2*x^2 + a*b)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}), x)$

### 3.360.8 Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 - c^2 x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{5/2} (b \text{ arcosh}(cx) + a)^2} dx$$

input `integrate(1/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(1/((-c^2*x^2 + 1)^(5/2)*(b*arccosh(c*x) + a)^2), x)`

### 3.360.9 Mupad [N/A]

Not integrable

Time = 3.51 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 - c^2 x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \text{ acosh}(cx))^2 (1 - c^2 x^2)^{5/2}} dx$$

input `int(1/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)`

output `int(1/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)`

---

3.360.  $\int \frac{1}{(1 - c^2 x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx$

**3.361** 
$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.361.1 Optimal result . . . . .	2853
3.361.2 Mathematica [N/A] . . . . .	2853
3.361.3 Rubi [N/A] . . . . .	2854
3.361.4 Maple [N/A] (verified) . . . . .	2854
3.361.5 Fricas [N/A] . . . . .	2855
3.361.6 Sympy [F(-1)] . . . . .	2855
3.361.7 Maxima [N/A] . . . . .	2855
3.361.8 Giac [F(-2)] . . . . .	2856
3.361.9 Mupad [N/A] . . . . .	2856

**3.361.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{x(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.361.2 Mathematica [N/A]**

Not integrable

Time = 30.70 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/(x*(1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/(x*(1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.361.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2} dx$$

input `Int[1/(x*(1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.361.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.361.4 Maple [N/A] (verified)**

Not integrable

Time = 1.46 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x(-c^2x^2+1)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `int(1/x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int(1/x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.361.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 144, normalized size of antiderivative = 5.14

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2x^2+1)^{\frac{5}{2}}(b\operatorname{arcosh}(cx)+a)^2x} dx$$

input `integrate(1/x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2+1)/(a^2*c^6*x^7-3*a^2*c^4*x^5+3*a^2*c^2*x^3-a^2*x+(b^2*c^6*x^7-3*b^2*c^4*x^5+3*b^2*c^2*x^3-b^2*x)*arccosh(c*x)^2+2*(a*b*c^6*x^7-3*a*b*c^4*x^5+3*a*b*c^2*x^3-a*b*x)*arccosh(c*x)),x)`

**3.361.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(1/x/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.361.7 Maxima [N/A]**

Not integrable

Time = 1.96 (sec) , antiderivative size = 638, normalized size of antiderivative = 22.79

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2x^2+1)^{\frac{5}{2}}(b\operatorname{arcosh}(cx)+a)^2x} dx$$

input `integrate(1/x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`



output  $-(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1})/(((b^2*c^4*x^4 - b^2*c^2*x^2)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^5*x^5 - 2*b^2*c^3*x^3 + b^2*c*x)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^4*x^4 - a*b*c^2*x^2)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^5*x^5 - 2*a*b*c^3*x^3 + a*b*c*x)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}) - \text{integrate}((5*c^5*x^5 - 5*c^3*x^3 + (5*c^3*x^3 - 2*c*x)*(c*x + 1)*(c*x - 1) + (10*c^4*x^4 - 7*c^2*x^2 + 1)*\sqrt{c*x + 1})*\sqrt{c*x - 1})/(((b^2*c^7*x^8 - 2*b^2*c^5*x^6 + b^2*c^3*x^4)*(c*x + 1)^{(3/2)}*(c*x - 1) + 2*(b^2*c^8*x^9 - 3*b^2*c^6*x^7 + 3*b^2*c^4*x^5 - b^2*c^2*x^3)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^9*x^{10} - 4*b^2*c^7*x^8 + 6*b^2*c^5*x^6 - 4*b^2*c^3*x^4 + b^2*c*x^2)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^7*x^8 - 2*a*b*c^5*x^6 + a*b*c^3*x^4)*(c*x + 1)^{(3/2)}*(c*x - 1) + 2*(a*b*c^8*x^9 - 3*a*b*c^6*x^7 + 3*a*b*c^4*x^5 - a*b*c^2*x^3)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^9*x^{10} - 4*a*b*c^7*x^8 + 6*a*b*c^5*x^6 - 4*a*b*c^3*x^4 + a*b*c*x^2)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}), x)$

### 3.361.8 Giac [F(-2)]

Exception generated.

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\text{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate(1/x/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.361.9 Mupad [N/A]

Not integrable

Time = 3.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\text{barccosh}(cx))^2} dx = \int \frac{1}{x(a+b\text{acosh}(cx))^2(1-c^2x^2)^{5/2}} dx$$

input `int(1/(x*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)`

---

3.361.  $\int \frac{1}{x(1-c^2x^2)^{5/2}(a+\text{barccosh}(cx))^2} dx$

output `int(1/(x*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)`

---

3.361.  $\int \frac{1}{x(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.362** 
$$\int \frac{1}{x^2(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.362.1 Optimal result . . . . .	2858
3.362.2 Mathematica [N/A] . . . . .	2858
3.362.3 Rubi [N/A] . . . . .	2859
3.362.4 Maple [N/A] (verified) . . . . .	2859
3.362.5 Fricas [N/A] . . . . .	2860
3.362.6 Sympy [F(-1)] . . . . .	2860
3.362.7 Maxima [N/A] . . . . .	2860
3.362.8 Giac [N/A] . . . . .	2861
3.362.9 Mupad [N/A] . . . . .	2861

**3.362.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{1}{x^2(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{x^2(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.362.2 Mathematica [N/A]**

Not integrable

Time = 12.77 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{1}{x^2(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{x^2(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/(x^2*(1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/(x^2*(1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.362.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `Int[1/(x^2*(1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.362.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(x_.)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.362.4 Maple [N/A] (verified)**

Not integrable

Time = 1.60 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^2 (-c^2 x^2 + 1)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int(1/x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.362.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 150, normalized size of antiderivative = 5.36

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arccosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)/(a^2*c^6*x^8 - 3*a^2*c^4*x^6 + 3*a^2*c^2*x^4 - a^2*x^2 + (b^2*c^6*x^8 - 3*b^2*c^4*x^6 + 3*b^2*c^2*x^4 - b^2*x^2)*arccosh(c*x)^2 + 2*(a*b*c^6*x^8 - 3*a*b*c^4*x^6 + 3*a*b*c^2*x^4 - a*b*x^2)*arccosh(c*x)), x)`

**3.362.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(1/x**2/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.362.7 Maxima [N/A]**

Not integrable

Time = 1.85 (sec) , antiderivative size = 647, normalized size of antiderivative = 23.11

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arccosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $-(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1})/(((b^2*c^4*x^5 - b^2*c^2*x^3)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^5*x^6 - 2*b^2*c^3*x^4 + b^2*c*x^2)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^4*x^5 - a*b*c^2*x^3)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^5*x^6 - 2*a*b*c^3*x^4 + a*b*c*x^2)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}) - \text{integrate}((6*c^5*x^5 - 7*c^3*x^3 + 3*(2*c^3*x^3 - c*x)*(c*x + 1)*(c*x - 1) + 2*(6*c^4*x^4 - 5*c^2*x^2 + 1)*\sqrt{c*x + 1}*\sqrt{c*x - 1} + c*x)/(((b^2*c^7*x^9 - 2*b^2*c^5*x^7 + b^2*c^3*x^5)*(c*x + 1)^{(3/2)}*(c*x - 1) + 2*(b^2*c^8*x^{10} - 3*b^2*c^6*x^8 + 3*b^2*c^4*x^6 - b^2*c^2*x^4)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^9*x^{11} - 4*b^2*c^7*x^9 + 6*b^2*c^5*x^7 - 4*b^2*c^3*x^5 + b^2*c*x^3)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^7*x^9 - 2*a*b*c^5*x^7 + a*b*c^3*x^5)*(c*x + 1)^{(3/2)}*(c*x - 1) + 2*(a*b*c^8*x^{10} - 3*a*b*c^6*x^8 + 3*a*b*c^4*x^6 - a*b*c^2*x^4)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^9*x^{11} - 4*a*b*c^7*x^9 + 6*a*b*c^5*x^7 - 4*a*b*c^3*x^5 + a*b*c*x^3)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}), x)$

### 3.362.8 Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(-c^2 x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2 x^2} dx$$

input `integrate(1/x^2/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(1/((-c^2*x^2 + 1)^(5/2)*(b*arccosh(c*x) + a)^2*x^2), x)`

### 3.362.9 Mupad [N/A]

Not integrable

Time = 3.36 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^2 (1 - c^2 x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{x^2 (a + b \operatorname{acosh}(cx))^2 (1 - c^2 x^2)^{5/2}} dx$$

input `int(1/(x^2*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)`

output `int(1/(x^2*(a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)`

$$3.363 \quad \int \frac{(fx)^m (1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

3.363.1 Optimal result	2863
3.363.2 Mathematica [N/A]	2863
3.363.3 Rubi [N/A]	2864
3.363.4 Maple [N/A] (verified)	2864
3.363.5 Fricas [N/A]	2865
3.363.6 Sympy [F(-1)]	2865
3.363.7 Maxima [N/A]	2865
3.363.8 Giac [F(-2)]	2866
3.363.9 Mupad [N/A]	2866

### 3.363.1 Optimal result

Integrand size = 30, antiderivative size = 30

$$\int \frac{(fx)^m (1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{(fx)^m (1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((f*x)^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

### 3.363.2 Mathematica [N/A]

Not integrable

Time = 1.61 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(fx)^m (1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[((f*x)^m*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x])^2,x]`

output `Integrate[((f*x)^m*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x])^2, x]`

---


$$3.363. \quad \int \frac{(fx)^m (1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$



**3.363.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1 - c^2 x^2)^{3/2} (fx)^m}{(a + \operatorname{barccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{(1 - c^2 x^2)^{3/2} (fx)^m}{(a + \operatorname{barccosh}(cx))^2} dx$$

input `Int[((f*x)^m*(1 - c^2*x^2)^(3/2))/(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

**3.363.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.363.4 Maple [N/A] (verified)**

Not integrable

Time = 2.60 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m (-c^2 x^2 + 1)^{3/2}}{(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((f*x)^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int((f*x)^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

---

3.363.  $\int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + b \operatorname{arccosh}(cx))^2} dx$

**3.363.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.80

$$\int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}} (fx)^m}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((f*x)^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)*(f*x)^m/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

**3.363.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.363.7 Maxima [N/A]**

Not integrable

Time = 1.95 (sec) , antiderivative size = 577, normalized size of antiderivative = 19.23

$$\int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(-c^2 x^2 + 1)^{\frac{3}{2}} (fx)^m}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((f*x)^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

```
output ((c^4*f^m*x^4 - 2*c^2*f^m*x^2 + f^m)*(c*x + 1)*sqrt(c*x - 1)*x^m + (c^5*f^m*x^5 - 2*c^3*f^m*x^3 + c*f^m*x)*sqrt(c*x + 1)*x^m)*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) - integrate(((c^5*f^m*(m + 4)*x^5 - c^3*f^m*(2*m + 3)*x^3 + c*f^m*(m - 1)*x)*(c*x + 1)^(3/2)*(c*x - 1)*x^m + (2*c^6*f^m*(m + 4)*x^6 - c^4*f^m*(5*m + 12)*x^4 + 4*c^2*f^m*(m + 1)*x^2 - f^m*m)*(c*x + 1)*sqrt(c*x - 1)*x^m + (c^7*f^m*(m + 4)*x^7 - 3*c^5*f^m*(m + 3)*x^5 + 3*c^3*f^m*(m + 2)*x^3 - c*f^m*(m + 1)*x)*sqrt(c*x + 1)*x^m)*sqrt(-c*x + 1)/(a*b*c^5*x^5 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^3 - 2*a*b*c^3*x^3 + a*b*c*x + 2*(a*b*c^4*x^4 - a*b*c^2*x^2)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^5 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^3 - 2*b^2*c^3*x^3 + b^2*c*x + 2*(b^2*c^4*x^4 - b^2*c^2*x^2)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)
```

### 3.363.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

```
input integrate((f*x)^m*(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.363.9 Mupad [N/A]

Not integrable

Time = 3.18 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + b \operatorname{acosh}(cx))^2} dx$$

```
input int(((f*x)^m*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x))^2,x)
```

---

3.363.  $\int \frac{(fx)^m (1 - c^2 x^2)^{3/2}}{(a + \operatorname{barccosh}(cx))^2} dx$

output `int(((f*x)^m*(1 - c^2*x^2)^(3/2))/(a + b*acosh(c*x))^2, x)`

---

3.363.  $\int \frac{(fx)^m(1-c^2x^2)^{3/2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

**3.364**  $\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.364.1 Optimal result	2868
3.364.2 Mathematica [N/A]	2868
3.364.3 Rubi [N/A]	2869
3.364.4 Maple [N/A] (verified)	2869
3.364.5 Fricas [N/A]	2870
3.364.6 Sympy [N/A]	2870
3.364.7 Maxima [N/A]	2870
3.364.8 Giac [F(-2)]	2871
3.364.9 Mupad [N/A]	2871

**3.364.1 Optimal result**

Integrand size = 30, antiderivative size = 30

$$\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((f*x)^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x)`

**3.364.2 Mathematica [N/A]**

Not integrable

Time = 0.37 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[((f*x)^m*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

output `Integrate[((f*x)^m*Sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2, x]`

**3.364.3 Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-c^2x^2}(fx)^m}{(a+b\operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{\sqrt{1-c^2x^2}(fx)^m}{(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Int[((f*x)^m*sqrt[1 - c^2*x^2])/(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

**3.364.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n_.*((f_.)*(x_))^m_.*((d_.) + (e_.)*(x_)^2)^p_. , x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.364.4 Maple [N/A] (verified)**

Not integrable

Time = 2.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m \sqrt{-c^2x^2 + 1}}{(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((f*x)^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x)`

output `int((f*x)^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x)`

---

3.364.  $\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

**3.364.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.47

$$\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}(fx)^m}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((f*x)^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2 + 1)*(f*x)^m/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

**3.364.6 Sympy [N/A]**

Not integrable

Time = 6.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m \sqrt{-(cx-1)(cx+1)}}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((f*x)**m*(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x))**2,x)`

output `Integral((f*x)**m*sqrt(-(c*x - 1)*(c*x + 1))/(a + b*acosh(c*x))**2, x)`

**3.364.7 Maxima [N/A]**

Not integrable

Time = 1.43 (sec) , antiderivative size = 514, normalized size of antiderivative = 17.13

$$\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{-c^2x^2+1}(fx)^m}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((f*x)^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-(c^2*f^m*x^2 - f^m)*(c*x + 1)*sqrt(c*x - 1)*x^m + (c^3*f^m*x^3 - c*f^m*x)*sqrt(c*x + 1)*x^m*sqrt(-c*x + 1)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((c^3*f^m*(m + 2)*x^3 - c*f^m*(m - 1)*x)*(c*x + 1)^(3/2)*(c*x - 1)*x^m + (2*c^4*f^m*(m + 2)*x^4 - c^2*f^m*(3*m + 2)*x^2 + f^m*m)*(c*x + 1)*sqrt(c*x - 1)*x^m + (c^5*f^m*(m + 2)*x^5 - c^3*f^m*(2*m + 3)*x^3 + c*f^m*(m + 1)*x)*sqrt(c*x + 1)*x^m)*sqrt(-c*x + 1)/(a*b*c^5*x^5 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^3 - 2*a*b*c^3*x^3 + a*b*c*x + 2*(a*b*c^4*x^4 - a*b*c^2*x^2)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^5 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^3 - 2*b^2*c^3*x^3 + b^2*c*x + 2*(b^2*c^4*x^4 - b^2*c^2*x^2)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.364.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

### 3.364.9 Mupad [N/A]

Not integrable

Time = 3.37 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{acosh}(cx))^2} dx$$

---

3.364.  $\int \frac{(fx)^m \sqrt{1-c^2x^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$



input `int(((f*x)^m*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2,x)`

output `int(((f*x)^m*(1 - c^2*x^2)^(1/2))/(a + b*acosh(c*x))^2, x)`

**3.365**  $\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))^2} dx$

3.365.1 Optimal result . . . . . 2873  
 3.365.2 Mathematica [N/A] . . . . . 2873  
 3.365.3 Rubi [N/A] . . . . . 2874  
 3.365.4 Maple [N/A] (verified) . . . . . 2875  
 3.365.5 Fricas [N/A] . . . . . 2875  
 3.365.6 Sympy [N/A] . . . . . 2875  
 3.365.7 Maxima [N/A] . . . . . 2876  
 3.365.8 Giac [N/A] . . . . . 2876  
 3.365.9 Mupad [N/A] . . . . . 2877

**3.365.1 Optimal result**

Integrand size = 30, antiderivative size = 30

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))^2} dx = -\frac{(fx)^m\sqrt{-1+cx}}{bc\sqrt{1-cx}(a+b\text{arccosh}(cx))} + \frac{fm\sqrt{-1+cx}\text{Int}\left(\frac{(fx)^{-1+m}}{a+b\text{arccosh}(cx)}, x\right)}{bc\sqrt{1-cx}}$$

output `-(f*x)^m*(c*x-1)^(1/2)/b/c/(a+b*arccosh(c*x))/(-c*x+1)^(1/2)+f*m*(c*x-1)^(1/2)*Unintegrable((f*x)^(-1+m)/(a+b*arccosh(c*x)),x)/b/c/(-c*x+1)^(1/2)`

**3.365.2 Mathematica [N/A]**

Not integrable

Time = 1.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))^2} dx = \int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b\text{arccosh}(cx))^2} dx$$

input `Integrate[(f*x)^m/(Sqrt[1-c^2*x^2]*(a+b*ArcCosh[c*x])^2),x]`

output `Integrate[(f*x)^m/(Sqrt[1-c^2*x^2]*(a+b*ArcCosh[c*x])^2),x]`

**3.365.3 Rubi [N/A]**

Not integrable

Time = 0.44 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6365, 6303}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))^2} dx$$

↓ 6365

$$\frac{fm\sqrt{cx-1} \int \frac{(fx)^{m-1}}{a+\text{barccosh}(cx)} dx}{bc\sqrt{1-cx}} - \frac{\sqrt{cx-1}(fx)^m}{bc\sqrt{1-cx}(a+\text{barccosh}(cx))}$$

↓ 6303

$$\frac{fm\sqrt{cx-1} \int \frac{(fx)^{m-1}}{a+\text{barccosh}(cx)} dx}{bc\sqrt{1-cx}} - \frac{\sqrt{cx-1}(fx)^m}{bc\sqrt{1-cx}(a+\text{barccosh}(cx))}$$

input `Int[(f*x)^m/(Sqrt[1 - c^2*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.365.3.1 Defintions of rubi rules used**

rule 6303 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Unintegrable[(d*x)^m*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, m, n}, x]`

rule 6365 `Int[(((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.))/Sqrt[(d_ + (e_.)*(x_)^2)], x_Symbol]
:> Simp[(f*x)^m*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])], x] - Simp[f*(m/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]
Int[(f*x)^(m - 1)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1]`

---

3.365.  $\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+\text{barccosh}(cx))^2} dx$

**3.365.4 Maple [N/A] (verified)**

Not integrable

Time = 1.36 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m}{\sqrt{-c^2x^2+1} (a+b \operatorname{arccosh}(cx))^2} dx$$

input `int((f*x)^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x)`output `int((f*x)^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x)`**3.365.5 Fricas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.73

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b \operatorname{arccosh}(cx))^2} dx = \int \frac{(fx)^m}{\sqrt{-c^2x^2+1}(b \operatorname{arccosh}(cx)+a)^2} dx$$

input `integrate((f*x)^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`output `integral(-sqrt(-c^2*x^2+1)*(f*x)^m/(a^2*c^2*x^2+(b^2*c^2*x^2-b^2)*arccosh(c*x)^2-a^2+2*(a*b*c^2*x^2-a*b)*arccosh(c*x)),x)`**3.365.6 Sympy [N/A]**

Not integrable

Time = 24.71 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b \operatorname{arccosh}(cx))^2} dx = \int \frac{(fx)^m}{\sqrt{-(cx-1)(cx+1)}(a+b \operatorname{acosh}(cx))^2} dx$$

input `integrate((f*x)**m/(-c**2*x**2+1)**(1/2)/(a+b*acosh(c*x))**2,x)`output `Integral((f*x)**m/(sqrt(-(c*x-1)*(c*x+1))*(a+b*acosh(c*x))**2),x)`

---

3.365.  $\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b \operatorname{arccosh}(cx))^2} dx$

**3.365.7 Maxima [N/A]**

Not integrable

Time = 0.71 (sec) , antiderivative size = 544, normalized size of antiderivative = 18.13

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

```
input integrate((f*x)^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")
```

```
output -((c^2*f^m*x^2 - f^m)*sqrt(c*x + 1)*sqrt(c*x - 1)*x^m + (c^3*f^m*x^3 - c*f^m*x)*x^m)/(((c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x + (b^2*c^3*x^2 - b^2*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x + (a*b*c^3*x^2 - a*b*c)*sqrt(c*x + 1))*sqrt(-c*x + 1)) + integrate((((c^3*f^m*m*x^3 - c*f^m*(m - 1)*x)*(c*x + 1)*(c*x - 1)*x^m + (2*c^4*f^m*m*x^4 - 3*c^2*f^m*m*x^2 + f^m*m)*sqrt(c*x + 1)*sqrt(c*x - 1)*x^m + (c^5*f^m*m*x^5 - c^3*f^m*(2*m + 1)*x^3 + c*f^m*(m + 1)*x)*x^m)/(((c*x + 1)^(3/2)*(c*x - 1)*b^2*c^3*x^3 + 2*(b^2*c^4*x^4 - b^2*c^2*x^2)*(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^5 - 2*b^2*c^3*x^3 + b^2*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + ((c*x + 1)^(3/2)*(c*x - 1)*a*b*c^3*x^3 + 2*(a*b*c^4*x^4 - a*b*c^2*x^2)*(c*x + 1)*sqrt(c*x - 1) + (a*b*c^5*x^5 - 2*a*b*c^3*x^3 + a*b*c*x)*sqrt(c*x + 1))*sqrt(-c*x + 1)), x)
```

**3.365.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{\sqrt{-c^2x^2+1}(b\operatorname{arcosh}(cx)+a)^2} dx$$

```
input integrate((f*x)^m/(-c^2*x^2+1)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output integrate((f*x)^m/(sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^2), x)
```

**3.365.9 Mupad [N/A]**

Not integrable

Time = 3.71 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m}{\sqrt{1-c^2x^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(fx)^m}{(a+b\operatorname{acosh}(cx))^2\sqrt{1-c^2x^2}} dx$$

input `int((f*x)^m/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)),x)`output `int((f*x)^m/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(1/2)), x)`

$$3.366 \quad \int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.366.1 Optimal result	2878
3.366.2 Mathematica [N/A]	2878
3.366.3 Rubi [N/A]	2879
3.366.4 Maple [N/A] (verified)	2879
3.366.5 Fricas [N/A]	2880
3.366.6 Sympy [F(-1)]	2880
3.366.7 Maxima [N/A]	2880
3.366.8 Giac [N/A]	2881
3.366.9 Mupad [N/A]	2881

### 3.366.1 Optimal result

Integrand size = 30, antiderivative size = 30

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((f*x)^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

### 3.366.2 Mathematica [N/A]

Not integrable

Time = 1.67 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[(f*x)^m/((1-c^2*x^2)^(3/2)*(a+b*ArcCosh[c*x])^2),x]`

output `Integrate[(f*x)^m/((1-c^2*x^2)^(3/2)*(a+b*ArcCosh[c*x])^2),x]`

**3.366.3 Rubi [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx$$

input `Int[(f*x)^m/((1 - c^2*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.366.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.366.4 Maple [N/A] (verified)**

Not integrable

Time = 1.63 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m}{(-c^2x^2 + 1)^{\frac{3}{2}}(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((f*x)^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int((f*x)^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x)`

---

3.366.  $\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$



**3.366.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 108, normalized size of antiderivative = 3.60

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(-c^2x^2+1)^{\frac{3}{2}}(b \operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate((f*x)^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(-c^2*x^2+1)*(f*x)^m/(a^2*c^4*x^4-2*a^2*c^2*x^2+(b^2*c^4*x^4-2*b^2*c^2*x^2+b^2)*arccosh(c*x)^2+a^2+2*(a*b*c^4*x^4-2*a*b*c^2*x^2+a*b)*arccosh(c*x)), x)`

**3.366.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((f*x)**m/(-c**2*x**2+1)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.366.7 Maxima [N/A]**

Not integrable

Time = 1.26 (sec) , antiderivative size = 596, normalized size of antiderivative = 19.87

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(-c^2x^2+1)^{\frac{3}{2}}(b \operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate((f*x)^m/(-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $(c^m f^m x^m + \sqrt{cx+1}\sqrt{cx-1}f^m x^m) / (((cx+1)\sqrt{cx-1})b^2c^2x + (b^2c^3x^2 - b^2c)\sqrt{cx+1})\sqrt{-cx+1}\log(cx + \sqrt{cx+1}\sqrt{cx-1}) + ((cx+1)\sqrt{cx-1}abc^2x + (abc^3x^2 - abc)\sqrt{cx+1})\sqrt{-cx+1}) - \text{integrate}(((c^3f^m(m-2)x^3 - cf^m(m-1)x)(cx+1)(cx-1)x^m + (2c^4f^m(m-2)x^4 - c^2f^m(3m-2)x^2 + f^m m)\sqrt{cx+1}\sqrt{cx-1})x^m + (c^5f^m(m-2)x^5 - c^3f^m(2m-1)x^3 + cf^m(m+1)x)x^m) / (((b^2c^5x^5 - b^2c^3x^3)(cx+1)^{(3/2)}(cx-1) + 2(b^2c^6x^6 - 2b^2c^4x^4 + b^2c^2x^2)(cx+1)\sqrt{cx-1}) + (b^2c^7x^7 - 3b^2c^5x^5 + 3b^2c^3x^3 - b^2cx)\sqrt{cx+1})\sqrt{-cx+1}\log(cx + \sqrt{cx+1}\sqrt{cx-1}) + ((abc^5x^5 - abc^3x^3)(cx+1)^{(3/2)}(cx-1) + 2(abc^6x^6 - 2abc^4x^4 + abc^2x^2)(cx+1)\sqrt{cx-1}) + (abc^7x^7 - 3abc^5x^5 + 3abc^3x^3 - abcx)\sqrt{cx+1})\sqrt{-cx+1}), x)$

### 3.366.8 Giac [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\text{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(-c^2x^2+1)^{3/2}(b\text{arcosh}(cx)+a)^2} dx$$

input `integrate((f*x)^m/((-c^2*x^2+1)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((f*x)^m/((-c^2*x^2+1)^(3/2)*(b*arccosh(c*x)+a)^2), x)`

### 3.366.9 Mupad [N/A]

Not integrable

Time = 4.08 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\text{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(a+b\text{acosh}(cx))^2(1-c^2x^2)^{3/2}} dx$$

input `int((f*x)^m/((a+b*acosh(c*x))^2*(1-c^2*x^2)^(3/2)),x)`

---

3.366.  $\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+\text{barccosh}(cx))^2} dx$

output `int((f*x)^m/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(3/2)), x)`

---

3.366.  $\int \frac{(fx)^m}{(1-c^2x^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$

$$3.367 \quad \int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.367.1 Optimal result	2883
3.367.2 Mathematica [N/A]	2883
3.367.3 Rubi [N/A]	2884
3.367.4 Maple [N/A] (verified)	2884
3.367.5 Fricas [N/A]	2885
3.367.6 Sympy [F(-1)]	2885
3.367.7 Maxima [N/A]	2885
3.367.8 Giac [N/A]	2886
3.367.9 Mupad [N/A]	2886

### 3.367.1 Optimal result

Integrand size = 30, antiderivative size = 30

$$\int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2}, x\right)$$

output `Unintegrable((f*x)^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

### 3.367.2 Mathematica [N/A]

Not integrable

Time = 2.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+\operatorname{barccosh}(cx))^2} dx$$

input `Integrate[(f*x)^m/((1-c^2*x^2)^(5/2)*(a+b*ArcCosh[c*x])^2),x]`

output `Integrate[(f*x)^m/((1-c^2*x^2)^(5/2)*(a+b*ArcCosh[c*x])^2),x]`

---


$$3.367. \quad \int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

**3.367.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2} dx$$

↓ 6375

$$\int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+\operatorname{arccosh}(cx))^2} dx$$

input `Int[(f*x)^m/((1 - c^2*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.367.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.367.4 Maple [N/A] (verified)**

Not integrable

Time = 2.46 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m}{(-c^2x^2 + 1)^{5/2}(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int((f*x)^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int((f*x)^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x)`

---

3.367.  $\int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.367.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 146, normalized size of antiderivative = 4.87

$$\int \frac{(fx)^m}{(1 - c^2x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(-c^2x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((f*x)^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*(f*x)^m/(a^2*c^6*x^6 - 3*a^2*c^4*x^4 + 3*a^2*c^2*x^2 + (b^2*c^6*x^6 - 3*b^2*c^4*x^4 + 3*b^2*c^2*x^2 - b^2)*arccosh(c*x)^2 - a^2 + 2*(a*b*c^6*x^6 - 3*a*b*c^4*x^4 + 3*a*b*c^2*x^2 - a*b)*arccosh(c*x)), x)`

**3.367.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m}{(1 - c^2x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate((f*x)**m/(-c**2*x**2+1)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.367.7 Maxima [N/A]**

Not integrable

Time = 1.28 (sec) , antiderivative size = 700, normalized size of antiderivative = 23.33

$$\int \frac{(fx)^m}{(1 - c^2x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(-c^2x^2 + 1)^{5/2} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((f*x)^m/(-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

---

3.367.  $\int \frac{(fx)^m}{(1 - c^2x^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx$

output  $-(c*f^m*x*x^m + \sqrt{c*x + 1}*\sqrt{c*x - 1}*f^m*x^m)/(((b^2*c^4*x^3 - b^2*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^5*x^4 - 2*b^2*c^3*x^2 + b^2*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^4*x^3 - a*b*c^2*x)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^5*x^4 - 2*a*b*c^3*x^2 + a*b*c)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}) + \text{integrate}(((c^3*f^m*(m - 4)*x^3 - c*f^m*(m - 1)*x)*(c*x + 1)*(c*x - 1)*x^m + (2*c^4*f^m*(m - 4)*x^4 - c^2*f^m*(3*m - 4)*x^2 + f^m*m)*\sqrt{c*x + 1}*\sqrt{c*x - 1}*x^m + (c^5*f^m*(m - 4)*x^5 - c^3*f^m*(2*m - 3)*x^3 + c*f^m*(m + 1)*x)*x^m)/(((b^2*c^7*x^7 - 2*b^2*c^5*x^5 + b^2*c^3*x^3)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(b^2*c^8*x^8 - 3*b^2*c^6*x^6 + 3*b^2*c^4*x^4 - b^2*c^2*x^2)*(c*x + 1)*\sqrt{c*x - 1} + (b^2*c^9*x^9 - 4*b^2*c^7*x^7 + 6*b^2*c^5*x^5 - 4*b^2*c^3*x^3 + b^2*c*x)*\sqrt{c*x + 1})*\sqrt{-c*x + 1}*\log(c*x + \sqrt{c*x + 1}*\sqrt{c*x - 1}) + ((a*b*c^7*x^7 - 2*a*b*c^5*x^5 + a*b*c^3*x^3)*(c*x + 1)^(3/2)*(c*x - 1) + 2*(a*b*c^8*x^8 - 3*a*b*c^6*x^6 + 3*a*b*c^4*x^4 - a*b*c^2*x^2)*(c*x + 1)*\sqrt{c*x - 1} + (a*b*c^9*x^9 - 4*a*b*c^7*x^7 + 6*a*b*c^5*x^5 - 4*a*b*c^3*x^3 + a*b*c*x)*\sqrt{c*x + 1})*\sqrt{-c*x + 1})), x)$

### 3.367.8 Giac [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m}{(1 - c^2x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(-c^2x^2 + 1)^{5/2} (b \text{ arcosh}(cx) + a)^2} dx$$

input `integrate((f*x)^m/((-c^2*x^2+1)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((f*x)^m/((-c^2*x^2 + 1)^(5/2)*(b*arccosh(c*x) + a)^2), x)`

### 3.367.9 Mupad [N/A]

Not integrable

Time = 3.89 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m}{(1 - c^2x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx = \int \frac{(fx)^m}{(a + b \text{ acosh}(cx))^2 (1 - c^2x^2)^{5/2}} dx$$

---

3.367.  $\int \frac{(fx)^m}{(1 - c^2x^2)^{5/2} (a + \text{barccosh}(cx))^2} dx$

input `int((f*x)^m/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)),x)`

output `int((f*x)^m/((a + b*acosh(c*x))^2*(1 - c^2*x^2)^(5/2)), x)`

---

3.367.  $\int \frac{(fx)^m}{(1-c^2x^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$



**3.368**  $\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx$

3.368.1 Optimal result	2888
3.368.2 Mathematica [A] (verified)	2888
3.368.3 Rubi [A] (verified)	2889
3.368.4 Maple [A] (verified)	2889
3.368.5 Fricas [B] (verification not implemented)	2890
3.368.6 Sympy [F]	2890
3.368.7 Maxima [F]	2890
3.368.8 Giac [F]	2891
3.368.9 Mupad [B] (verification not implemented)	2891

**3.368.1 Optimal result**

Integrand size = 21, antiderivative size = 32

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx = -\frac{\sqrt{-1+ax}}{2a\sqrt{1-ax}\operatorname{arccosh}(ax)^2}$$

output `-1/2*(a*x-1)^(1/2)/a/arccosh(a*x)^2/(-a*x+1)^(1/2)`

**3.368.2 Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.41

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx = -\frac{\sqrt{-1+ax}\sqrt{1+ax}}{2a\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^2}$$

input `Integrate[1/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3),x]`

output `-1/2*(Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(a*Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^2)`

### 3.368.3 Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx$$

↓ 6307

$$-\frac{\sqrt{ax-1}}{2a\sqrt{1-ax}\operatorname{arccosh}(ax)^2}$$

input `Int[1/(Sqrt[1 - a^2*x^2]*ArcCosh[a*x]^3),x]`

output `-1/2*Sqrt[-1 + a*x]/(a*Sqrt[1 - a*x]*ArcCosh[a*x]^2)`

#### 3.368.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.368.4 Maple [A] (verified)

Time = 0.70 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.59

method	result	size
default	$\frac{\sqrt{-(ax-1)(ax+1)}\sqrt{ax-1}\sqrt{ax+1}}{2(a^2x^2-1)a\operatorname{arccosh}(ax)^2}$	51

input `int(1/arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/2*(-(a*x-1)*(a*x+1))^(1/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/(a^2*x^2-1)/a/arc  
cosh(a*x)^2`

**3.368.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 56 vs.  $2(26) = 52$ .

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx = \frac{\sqrt{a^2x^2-1}\sqrt{-a^2x^2+1}}{2(a^3x^2-a)\log(ax+\sqrt{a^2x^2-1})^2}$$

input `integrate(1/arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(a^2*x^2 - 1)*sqrt(-a^2*x^2 + 1)/((a^3*x^2 - a)*log(a*x + sqrt(a^2*x^2 - 1))^2)`

**3.368.6 Sympy [F]**

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx = \int \frac{1}{\sqrt{-(ax-1)(ax+1)}\operatorname{acosh}^3(ax)} dx$$

input `integrate(1/acosh(a*x)**3/(-a**2*x**2+1)**(1/2),x)`

output `Integral(1/(sqrt(-(a*x - 1)*(a*x + 1))*acosh(a*x)**3), x)`

**3.368.7 Maxima [F]**

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx = \int \frac{1}{\sqrt{-a^2x^2+1}\operatorname{arcosh}(ax)^3} dx$$

input `integrate(1/arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="maxima")`

output

```
-1/2*(a^7*x^7 - 3*a^5*x^5 + 3*a^3*x^3 + (a^4*x^4 - a^2*x^2)*(a*x + 1)^(3/2)
)*(a*x - 1)^(3/2) + (3*a^5*x^5 - 5*a^3*x^3 + 2*a*x)*(a*x + 1)*(a*x - 1) +
(3*a^6*x^6 - 7*a^4*x^4 + 5*a^2*x^2 - 1)*sqrt(a*x + 1)*sqrt(a*x - 1) - a*x
- (a^5*x^5 - 2*a^3*x^3 - (a^2*x^2 - 1)*(a*x + 1)^(3/2)*(a*x - 1)^(3/2) - (
a^3*x^3 - a*x)*(a*x + 1)*(a*x - 1) + (a^4*x^4 - 2*a^2*x^2 + 1)*sqrt(a*x +
1)*sqrt(a*x - 1) + a*x)*log(a*x + sqrt(a*x + 1)*sqrt(a*x - 1)))/(((a*x + 1)
)^2*(a*x - 1)^(3/2)*a^4*x^3 + 3*(a^5*x^4 - a^3*x^2)*(a*x + 1)^(3/2)*(a*x -
1) + 3*(a^6*x^5 - 2*a^4*x^3 + a^2*x)*(a*x + 1)*sqrt(a*x - 1) + (a^7*x^6 -
3*a^5*x^4 + 3*a^3*x^2 - a)*sqrt(a*x + 1))*sqrt(-a*x + 1)*log(a*x + sqrt(a
*x + 1)*sqrt(a*x - 1))^2) - integrate(-1/2*(2*a^6*x^6 - 3*a^4*x^4 - (2*a^2
*x^2 - 3)*(a*x + 1)^2*(a*x - 1)^2 - 4*(a^3*x^3 - a*x)*(a*x + 1)^(3/2)*(a*x
- 1)^(3/2) - 4*(a^2*x^2 - 1)*(a*x + 1)*(a*x - 1) + 4*(a^5*x^5 - 2*a^3*x^3
+ a*x)*sqrt(a*x + 1)*sqrt(a*x - 1) + 1)/(((a*x + 1)^(5/2)*(a*x - 1)^2*a^4
*x^4 + 4*(a^5*x^5 - a^3*x^3)*(a*x + 1)^2*(a*x - 1)^(3/2) + 6*(a^6*x^6 - 2*
a^4*x^4 + a^2*x^2)*(a*x + 1)^(3/2)*(a*x - 1) + 4*(a^7*x^7 - 3*a^5*x^5 + 3*
a^3*x^3 - a*x)*(a*x + 1)*sqrt(a*x - 1) + (a^8*x^8 - 4*a^6*x^6 + 6*a^4*x^4
- 4*a^2*x^2 + 1)*sqrt(a*x + 1))*sqrt(-a*x + 1)*log(a*x + sqrt(a*x + 1)*sq
r(a*x - 1))), x)
```

### 3.368.8 Giac [F]

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx = \int \frac{1}{\sqrt{-a^2x^2+1}\operatorname{arccosh}(ax)^3} dx$$

input `integrate(1/arccosh(a*x)^3/(-a^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-a^2*x^2 + 1)*arccosh(a*x)^3), x)`

### 3.368.9 Mupad [B] (verification not implemented)

Time = 3.34 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx = \frac{\sqrt{1-a^2x^2}\sqrt{ax-1}\sqrt{ax+1}}{a\operatorname{acosh}(ax)^2(2a^2x^2-2)}$$

input `int(1/(acosh(a*x)^3*(1 - a^2*x^2)^(1/2)),x)`

---

3.368.  $\int \frac{1}{\sqrt{1-a^2x^2}\operatorname{arccosh}(ax)^3} dx$

output  $((1 - a^2x^2)^{1/2}(ax - 1)^{1/2}(ax + 1)^{1/2})/(a \operatorname{acosh}(ax)^2(2a^2x^2 - 2))$

$$3.369 \quad \int \frac{x^3(d-c^2 dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

3.369.1 Optimal result . . . . .	2893
3.369.2 Mathematica [A] (warning: unable to verify) . . . . .	2894
3.369.3 Rubi [A] (verified) . . . . .	2894
3.369.4 Maple [F] . . . . .	2897
3.369.5 Fricas [F(-2)] . . . . .	2897
3.369.6 Sympy [F] . . . . .	2898
3.369.7 Maxima [F] . . . . .	2898
3.369.8 Giac [F(-2)] . . . . .	2898
3.369.9 Mupad [F(-1)] . . . . .	2899

### 3.369.1 Optimal result

Integrand size = 27, antiderivative size = 259

$$\int \frac{x^3(d-c^2 dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = \frac{2dx^3(-1+cx)^{3/2}(1+cx)^{3/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{3de^{\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^4} - \frac{de^{\frac{6a}{b}}\sqrt{\frac{3\pi}{2}}\operatorname{erf}\left(\frac{\sqrt{6}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^4} + \frac{3de^{-\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^4} - \frac{de^{-\frac{6a}{b}}\sqrt{\frac{3\pi}{2}}\operatorname{erfi}\left(\frac{\sqrt{6}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^4}$$

```
output 3/32*d*exp(2*a/b)*erf(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(1/2)*Pi
^(1/2)/b^(3/2)/c^4+3/32*d*erfi(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2
^(1/2)*Pi^(1/2)/b^(3/2)/c^4/exp(2*a/b)-1/32*d*exp(6*a/b)*erf(6^(1/2)*(a+b*
arccosh(c*x))^(1/2)/b^(1/2))*6^(1/2)*Pi^(1/2)/b^(3/2)/c^4-1/32*d*erfi(6^(1
/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*6^(1/2)*Pi^(1/2)/b^(3/2)/c^4/exp(6*a
/b)+2*d*x^3*(c*x-1)^(3/2)*(c*x+1)^(3/2)/b/c/(a+b*arccosh(c*x))^(1/2)
```

**3.369.2 Mathematica [A] (warning: unable to verify)**

Time = 1.88 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.16

$$\int \frac{x^3(d - c^2 dx^2)}{(a + \operatorname{barccosh}(cx))^{3/2}} dx = \frac{de^{-\frac{6a}{b}} \left( -\sqrt{6} \sqrt{-\frac{a + \operatorname{barccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{6(a + \operatorname{barccosh}(cx))}{b}\right) \right) + 3\sqrt{2} e^{\frac{4a}{b}} \sqrt{-\frac{a + \operatorname{barccosh}(cx)}{b}}}{1}$$

input `Integrate[(x^3*(d - c^2*d*x^2))/(a + b*ArcCosh[c*x])^(3/2), x]`

output `(d*(-(Sqrt[6]*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-6*(a + b*ArcCosh[c*x])/b]) + 3*Sqrt[2]*E^((4*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-2*(a + b*ArcCosh[c*x])/b)] + E^((6*a)/b)*(-64*c^3*x^3*Sqrt[(-1 + c*x)/(1 + c*x)] - 64*c^4*x^4*Sqrt[(-1 + c*x)/(1 + c*x)] - 3*Sqrt[2]*E^((2*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (2*(a + b*ArcCosh[c*x])/b)] + Sqrt[6]*E^((6*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (6*(a + b*ArcCosh[c*x])/b)] + 10*Sinh[2*ArcCosh[c*x]] + 8*Sinh[4*ArcCosh[c*x]] + 2*Sinh[6*ArcCosh[c*x]]))/(32*b*c^4*E^((6*a)/b)*Sqrt[a + b*ArcCosh[c*x]])`

**3.369.3 Rubi [A] (verified)**Time = 1.99 (sec) , antiderivative size = 463, normalized size of antiderivative = 1.79, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6357, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(d - c^2 dx^2)}{(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

$$\downarrow \text{6357}$$

$$-\frac{12cd \int \frac{x^4 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a + \operatorname{barccosh}(cx)}} dx}{b} + \frac{6d \int \frac{x^2 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a + \operatorname{barccosh}(cx)}} dx}{bc} + \frac{2dx^3(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a + \operatorname{barccosh}(cx)}}$$

$$\downarrow \text{6368}$$

---

3.369.  $\int \frac{x^3(d - c^2 dx^2)}{(a + \operatorname{barccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
& \frac{12d \int \frac{\cosh^4\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{\phantom{12d \int}} + \\
& \frac{6d \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{\phantom{6d \int}} + \\
& \frac{b^2 c^4}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \frac{2dx^3(cx-1)^{3/2}(cx+1)^{3/2}}{\phantom{bc\sqrt{a+b\operatorname{arccosh}(cx)}}} \\
& \quad \downarrow \text{5971} \\
& \frac{6d \int \left( \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{1}{8\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{\phantom{6d \int}} - \\
& \frac{12d \int \left( \frac{\cosh\left(\frac{6a}{b} - \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{1}{16\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{\phantom{12d \int}} - \\
& \frac{b^2 c^4}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \frac{2dx^3(cx-1)^{3/2}(cx+1)^{3/2}}{\phantom{bc\sqrt{a+b\operatorname{arccosh}(cx)}}} \\
& \quad \downarrow \text{2009} \\
& \frac{6d \left( \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{-\frac{4a}{b}} \operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{4} \sqrt{a+b\operatorname{arccosh}(cx)} \right)}{\phantom{6d \left(}} - \\
& \frac{12d \left( \frac{1}{64} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{64} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{64} \sqrt{\frac{\pi}{6}} \sqrt{b} e^{\frac{6a}{b}} \operatorname{erf}\left(\frac{\sqrt{6}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)}{\phantom{12d \left(}} - \\
& \frac{b^2 c^4}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \frac{2dx^3(cx-1)^{3/2}(cx+1)^{3/2}}{\phantom{bc\sqrt{a+b\operatorname{arccosh}(cx)}}}
\end{aligned}$$

input `Int[(x^3*(d - c^2*d*x^2))/(a + b*ArcCosh[c*x])^(3/2),x]`



```
output (2*d*x^3*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]])
+ (6*d*(-1/4*Sqrt[a + b*ArcCosh[c*x]] + (Sqrt[b]*E^((4*a)/b)*Sqrt[Pi]*Erf[
(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 + (Sqrt[b]*Sqrt[Pi]*Erfi[(2*Sqrt
[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((4*a)/b)))/(b^2*c^4) - (12*d*(-1/8
*Sqrt[a + b*ArcCosh[c*x]] + (Sqrt[b]*E^((4*a)/b)*Sqrt[Pi]*Erf[(2*Sqrt[a +
b*ArcCosh[c*x]])/Sqrt[b]])/64 - (Sqrt[b]*E^((2*a)/b)*Sqrt[Pi/2]*Erf[(Sqrt[
2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 + (Sqrt[b]*E^((6*a)/b)*Sqrt[Pi/6
]*Erf[(Sqrt[6]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 + (Sqrt[b]*Sqrt[Pi]*
Erfi[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((4*a)/b)) - (Sqrt[b]*Sq
rt[Pi/2]*Erfi[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((2*a)/b)
) + (Sqrt[b]*Sqrt[Pi/6]*Erfi[(Sqrt[6]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/
(64*E^((6*a)/b)))/(b^2*c^4)
```

### 3.369.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 5971 Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

```
rule 6357 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*
x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[
f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f
*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(
n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((
1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x
)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e,
f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m +
2*p + 1, 0] && IGtQ[m, -3]
```

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.369.4 Maple [F]

$$\int \frac{x^3(-c^2 d x^2 + d)}{(a + b \operatorname{arccosh}(cx))^{\frac{3}{2}}} dx$$

input `int(x^3*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

output `int(x^3*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

### 3.369.5 Fracas [F(-2)]

Exception generated.

$$\int \frac{x^3(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="fracas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.369.6 Sympy [F]**

$$\int \frac{x^3(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx =$$

$$-d \left( \int \left( -\frac{x^3}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right)$$

$$+ \int \frac{c^2 x^5}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx$$

input `integrate(x**3*(-c**2*d*x**2+d)/(a+b*acosh(c*x))**(3/2),x)`

output `-d*(Integral(-x**3/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**2*x**5/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

**3.369.7 Maxima [F]**

$$\int \frac{x^3(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int -\frac{(c^2 dx^2 - d)x^3}{(b \operatorname{arccosh}(cx) + a)^{\frac{3}{2}}} dx$$

input `integrate(x^3*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `-integrate((c^2*d*x^2 - d)*x^3/(b*arccosh(c*x) + a)^(3/2), x)`

**3.369.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const ve  
cteur & l) Error: Bad Argument Value

### 3.369.9 Mupad [**F(-1)**]

Timed out.

$$\int \frac{x^3(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{x^3(d - c^2 dx^2)}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((x^3*(d - c^2*d*x^2))/(a + b*acosh(c*x))^(3/2),x)`

output `int((x^3*(d - c^2*d*x^2))/(a + b*acosh(c*x))^(3/2), x)`

$$3.370 \quad \int \frac{x^2(d-c^2dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

3.370.1 Optimal result	2900
3.370.2 Mathematica [A] (warning: unable to verify)	2901
3.370.3 Rubi [A] (verified)	2901
3.370.4 Maple [F]	2904
3.370.5 Fricas [F(-2)]	2904
3.370.6 Sympy [F]	2905
3.370.7 Maxima [F]	2905
3.370.8 Giac [F]	2905
3.370.9 Mupad [F(-1)]	2906

### 3.370.1 Optimal result

Integrand size = 27, antiderivative size = 340

$$\int \frac{x^2(d-c^2dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = \frac{2dx^2(-1+cx)^{3/2}(1+cx)^{3/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{de^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8b^{3/2}c^3} + \frac{de^{\frac{3a}{b}}\sqrt{3\pi}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^3} - \frac{de^{\frac{5a}{b}}\sqrt{5\pi}\operatorname{erf}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^3} + \frac{de^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8b^{3/2}c^3} + \frac{de^{-\frac{3a}{b}}\sqrt{3\pi}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^3} - \frac{de^{-\frac{5a}{b}}\sqrt{5\pi}\operatorname{erfi}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^3}$$

```
output 1/8*d*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^3+
1/8*d*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^3/exp(a/b)
+1/16*d*exp(3*a/b)*erf(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*P
i^(1/2)/b^(3/2)/c^3+1/16*d*erfi(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*
3^(1/2)*Pi^(1/2)/b^(3/2)/c^3/exp(3*a/b)-1/16*d*exp(5*a/b)*erf(5^(1/2)*(a+b
*arccosh(c*x))^(1/2)/b^(1/2))*5^(1/2)*Pi^(1/2)/b^(3/2)/c^3-1/16*d*erfi(5^(
1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*5^(1/2)*Pi^(1/2)/b^(3/2)/c^3/exp(5*
a/b)+2*d*x^2*(c*x-1)^(3/2)*(c*x+1)^(3/2)/b/c/(a+b*arccosh(c*x))^(1/2)
```

$$3.370. \quad \int \frac{x^2(d-c^2dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

**3.370.2 Mathematica [A] (warning: unable to verify)**

Time = 1.29 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.13

$$\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \frac{de^{-\frac{5a}{b}} \left( -4e^{\frac{5a}{b}} \sqrt{\frac{-1+cx}{1+cx}} - 4ce^{\frac{5a}{b}} x \sqrt{\frac{-1+cx}{1+cx}} - 2e^{\frac{6a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \right) \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right)}{(a + b \operatorname{arccosh}(cx))^{3/2}}$$

input `Integrate[(x^2*(d - c^2*d*x^2))/(a + b*ArcCosh[c*x])^(3/2), x]`

output

```
(d*(-4*E^((5*a)/b)*Sqrt[(-1 + c*x)/(1 + c*x)] - 4*c*E^((5*a)/b)*x*Sqrt[(-1 + c*x)/(1 + c*x)] - 2*E^((6*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, a/b + ArcCosh[c*x]] - Sqrt[5]*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-5*(a + b*ArcCosh[c*x])/b) + Sqrt[3]*E^((2*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-3*(a + b*ArcCosh[c*x])/b) + 2*E^((4*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, -(a + b*ArcCosh[c*x])/b] - Sqrt[3]*E^((8*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (3*(a + b*ArcCosh[c*x])/b) + Sqrt[5]*E^((10*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (5*(a + b*ArcCosh[c*x])/b) - 2*E^((5*a)/b)*Sinh[3*ArcCosh[c*x]] + 2*E^((5*a)/b)*Sinh[5*ArcCosh[c*x]]))/(16*b*c^3*E^((5*a)/b)*Sqrt[a + b*ArcCosh[c*x]])
```

**3.370.3 Rubi [A] (verified)**Time = 1.94 (sec) , antiderivative size = 525, normalized size of antiderivative = 1.54, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.148$ , Rules used = {6357, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

$$\downarrow \text{6357}$$

$$-\frac{10cd \int \frac{x^3 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+b \operatorname{arccosh}(cx)}} dx}{b} + \frac{4d \int \frac{x \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+b \operatorname{arccosh}(cx)}} dx}{bc} + \frac{2dx^2(cx-1)^{3/2}(cx+1)^{3/2}}{bc \sqrt{a+b \operatorname{arccosh}(cx)}}$$

$$\downarrow \text{6368}$$

---

3.370.  $\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{10d \int \frac{\cosh^3\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a + \operatorname{arccosh}(cx))}{\phantom{10d \int}} + \\
 & \frac{4d \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a + \operatorname{arccosh}(cx))}{\phantom{4d \int}} + \\
 & \frac{b^2 c^3}{2dx^2(cx-1)^{3/2}(cx+1)^{3/2}} \\
 & \frac{bc\sqrt{a+b\operatorname{arccosh}(cx)}}{\phantom{2dx^2(cx-1)^{3/2}(cx+1)^{3/2}}} \\
 & \quad \downarrow \text{5971} \\
 & \frac{4d \int \left( \frac{\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a + \operatorname{arccosh}(cx))}{\phantom{4d \int}} - \\
 & \frac{10d \int \left( \frac{\cosh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a + \operatorname{arccosh}(cx))}{\phantom{10d \int}} + \\
 & \frac{b^2 c^3}{2dx^2(cx-1)^{3/2}(cx+1)^{3/2}} \\
 & \frac{bc\sqrt{a+b\operatorname{arccosh}(cx)}}{\phantom{2dx^2(cx-1)^{3/2}(cx+1)^{3/2}}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{4d \left( -\frac{1}{8}\sqrt{\pi}\sqrt{b}e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{8}\sqrt{\frac{\pi}{3}}\sqrt{b}e^{\frac{3a}{b}} \operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{8}\sqrt{\pi}\sqrt{b}e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{\phantom{4d \left(}} \right. \\
 & \frac{10d \left( -\frac{1}{16}\sqrt{\pi}\sqrt{b}e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32}\sqrt{\frac{\pi}{3}}\sqrt{b}e^{\frac{3a}{b}} \operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32}\sqrt{\frac{\pi}{5}}\sqrt{b}e^{\frac{5a}{b}} \operatorname{erf}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{\phantom{10d \left(}} \right. \\
 & \frac{b^2 c^3}{2dx^2(cx-1)^{3/2}(cx+1)^{3/2}} \\
 & \frac{bc\sqrt{a+b\operatorname{arccosh}(cx)}}{\phantom{2dx^2(cx-1)^{3/2}(cx+1)^{3/2}}}
 \end{aligned}$$

input `Int[(x^2*(d - c^2*d*x^2))/(a + b*ArcCosh[c*x])^(3/2), x]`

```
output (2*d*x^2*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]])
+ (4*d*(-1/8*(Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]
+ (Sqrt[b]*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]]
)/Sqrt[b]])/8 - (Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])
/(8*E^(a/b)) + (Sqrt[b]*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]]
)/Sqrt[b]])/(8*E^((3*a)/b)))/(b^2*c^3) - (10*d*(-1/16*(Sqrt[b]*E^(a/b)*Sqr
t[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]) + (Sqrt[b]*E^((3*a)/b)*Sqrt[P
i/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 + (Sqrt[b]*E^((5*
a)/b)*Sqrt[Pi/5]*Erf[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 - (Sq
rt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(16*E^(a/b)) + (Sqr
t[b]*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((
3*a)/b)) + (Sqrt[b]*Sqrt[Pi/5]*Erfi[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqr
t[b]])/(32*E^((5*a)/b)))/(b^2*c^3)
```

### 3.370.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 5971 Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

```
rule 6357 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_
.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*
x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[
f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f
*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(
n + 1), x], x] - Simp[c*(m + 2*p + 1)/(b*f*(n + 1))*Simp[(d + e*x^2)^p/((
1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x
)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e,
f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m +
2*p + 1, 0] && IGtQ[m, -3]
```



```
rule 6368 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]
```

### 3.370.4 Maple [F]

$$\int \frac{x^2(-c^2 d x^2 + d)}{(a + b \operatorname{arccosh}(c x))^{\frac{3}{2}}} dx$$

```
input int(x^2*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)
```

```
output int(x^2*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)
```

### 3.370.5 Fracas [F(-2)]

Exception generated.

$$\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(x^2*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="fracas")
```

```
output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)
```

**3.370.6 Sympy [F]**

$$\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx =$$

$$-d \left( \int \left( -\frac{x^2}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right)$$

$$+ \int \frac{c^2 x^4}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx$$

input `integrate(x**2*(-c**2*d*x**2+d)/(a+b*acosh(c*x))**(3/2),x)`

output `-d*(Integral(-x**2/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**2*x**4/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

**3.370.7 Maxima [F]**

$$\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int -\frac{(c^2 dx^2 - d)x^2}{(b \operatorname{arccosh}(cx) + a)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `-integrate((c^2*d*x^2 - d)*x^2/(b*arccosh(c*x) + a)^(3/2), x)`

**3.370.8 Giac [F]**

$$\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int -\frac{(c^2 dx^2 - d)x^2}{(b \operatorname{arccosh}(cx) + a)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `integrate(-(c^2*d*x^2 - d)*x^2/(b*arccosh(c*x) + a)^(3/2), x)`

---

3.370.  $\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$

**3.370.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{x^2(d - c^2 dx^2)}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((x^2*(d - c^2*d*x^2))/(a + b*acosh(c*x))^(3/2),x)`output `int((x^2*(d - c^2*d*x^2))/(a + b*acosh(c*x))^(3/2), x)`

**3.371**  $\int \frac{x(d-c^2 dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.371.1 Optimal result . . . . . 2907  
 3.371.2 Mathematica [F] . . . . . 2908  
 3.371.3 Rubi [A] (verified) . . . . . 2908  
 3.371.4 Maple [F] . . . . . 2912  
 3.371.5 Fricas [F(-2)] . . . . . 2912  
 3.371.6 Sympy [F] . . . . . 2912  
 3.371.7 Maxima [F] . . . . . 2913  
 3.371.8 Giac [F(-2)] . . . . . 2913  
 3.371.9 Mupad [F(-1)] . . . . . 2913

**3.371.1 Optimal result**

Integrand size = 25, antiderivative size = 241

$$\int \frac{x(d-c^2 dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = \frac{2dx(-1+cx)^{3/2}(1+cx)^{3/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{de^{\frac{4a}{b}}\sqrt{\pi}\operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^2} + \frac{de^{\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2b^{3/2}c^2} - \frac{de^{-\frac{4a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^2} + \frac{de^{-\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2b^{3/2}c^2}$$

```
output 1/4*d*exp(2*a/b)*erf(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(1/2)*Pi^(1/2)/b^(3/2)/c^2+1/4*d*erfi(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(1/2)*Pi^(1/2)/b^(3/2)/c^2/exp(2*a/b)-1/4*d*exp(4*a/b)*erf(2*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^2-1/4*d*erfi(2*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^2/exp(4*a/b)+2*d*x*(c*x-1)^(3/2)*(c*x+1)^(3/2)/b/c/(a+b*arccosh(c*x))^(1/2)
```

## 3.371.2 Mathematica [F]

$$\int \frac{x(d - c^2 dx^2)}{(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{x(d - c^2 dx^2)}{(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

input `Integrate[(x*(d - c^2*d*x^2))/(a + b*ArcCosh[c*x])^(3/2), x]`

output `Integrate[(x*(d - c^2*d*x^2))/(a + b*ArcCosh[c*x])^(3/2), x]`

## 3.371.3 Rubi [A] (verified)

Time = 2.93 (sec) , antiderivative size = 275, normalized size of antiderivative = 1.14, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$ , Rules used = {6357, 6322, 3042, 25, 3793, 2009, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x(d - c^2 dx^2)}{(a + \operatorname{barccosh}(cx))^{3/2}} dx \\ & \quad \downarrow \text{6357} \\ & -\frac{8cd \int \frac{x^2 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} + \frac{2d \int \frac{\sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{bc} + \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{6322} \\ & \frac{2d \int \frac{\sinh^2\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{b^2 c^2} - \frac{8cd \int \frac{x^2 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} + \\ & \quad \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{3042} \\ & \frac{2d \int -\frac{\sin\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^2}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{b^2 c^2} - \frac{8cd \int \frac{x^2 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} + \\ & \quad \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \end{aligned}$$

---

3.371.  $\int \frac{x(d - c^2 dx^2)}{(a + \operatorname{barccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow 25 \\
& \frac{2d \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2} - \frac{8cd \int \frac{x^2 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{b} + \\
& \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow 3793 \\
& \frac{2d \int \left( \frac{1}{2\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^2} - \\
& \frac{8cd \int \frac{x^2 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{b} + \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow 2009 \\
& \frac{8cd \int \frac{x^2 \sqrt{cx-1} \sqrt{cx+1}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{b} + \\
& \frac{2d \left( \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{-\frac{2a}{b}} \operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \sqrt{a+b\operatorname{arccosh}(cx)} \right)}{b^2 c^2} + \\
& \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow 6368 \\
& \frac{8d \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2} + \\
& \frac{2d \left( \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{-\frac{2a}{b}} \operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \sqrt{a+b\operatorname{arccosh}(cx)} \right)}{b^2 c^2} + \\
& \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow 5971
\end{aligned}$$

---

3.371.  $\int \frac{x(d-c^2 dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{8d \int \left( \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{1}{8\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a + \operatorname{arccosh}(cx))}{b^2 c^2} + \\
 & \frac{2d \left( \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{-\frac{2a}{b}} \operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \sqrt{a + \operatorname{arccosh}(cx)} \right)}{b^2 c^2} + \\
 & \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a + \operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{8d \left( \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{-\frac{4a}{b}} \operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{4} \sqrt{a + \operatorname{arccosh}(cx)} \right)}{b^2 c^2} + \\
 & \frac{2d \left( \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{-\frac{2a}{b}} \operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \sqrt{a + \operatorname{arccosh}(cx)} \right)}{b^2 c^2} + \\
 & \frac{2dx(cx-1)^{3/2}(cx+1)^{3/2}}{bc\sqrt{a + \operatorname{arccosh}(cx)}}
 \end{aligned}$$

input `Int[(x*(d - c^2*d*x^2))/(a + b*ArcCosh[c*x])^(3/2),x]`

output `(2*d*x*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]]) - (8*d*(-1/4*Sqrt[a + b*ArcCosh[c*x]] + (Sqrt[b]*E^((4*a)/b)*Sqrt[Pi]*Erf[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 + (Sqrt[b]*Sqrt[Pi]*Erfi[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((4*a)/b)))/(b^2*c^2) + (2*d*(-Sqrt[a + b*ArcCosh[c*x]] + (Sqrt[b]*E^((2*a)/b)*Sqrt[Pi/2]*Erf[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/4 + (Sqrt[b]*Sqrt[Pi/2]*Erfi[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(4*E^((2*a)/b)))/(b^2*c^2)`

### 3.371.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

---

3.371.  $\int \frac{x(d-c^2 dx^2)}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6322 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[2*p, 0]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`



**3.371.4 Maple [F]**

$$\int \frac{x(-c^2 d x^2 + d)}{(a + b \operatorname{arccosh}(cx))^{\frac{3}{2}}} dx$$

input `int(x*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

output `int(x*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

**3.371.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{x(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.371.6 Sympy [F]**

$$\int \frac{x(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx =$$

$$-d \left( \int \left( -\frac{x}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right)$$

$$+ \int \frac{c^2 x^3}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx$$

input `integrate(x*(-c**2*d*x**2+d)/(a+b*acosh(c*x))**(3/2),x)`

output `-d*(Integral(-x/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**2*x**3/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

---

3.371.  $\int \frac{x(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$

**3.371.7 Maxima [F]**

$$\int \frac{x(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int -\frac{(c^2 dx^2 - d)x}{(b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}}} dx$$

input `integrate(x*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `-integrate((c^2*d*x^2 - d)*x/(b*arccosh(c*x) + a)^(3/2), x)`

**3.371.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x*(-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

**3.371.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(d - c^2 dx^2)}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{x(d - c^2 dx^2)}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((x*(d - c^2*d*x^2))/(a + b*acosh(c*x))^(3/2),x)`

output `int((x*(d - c^2*d*x^2))/(a + b*acosh(c*x))^(3/2), x)`

**3.372**  $\int \frac{d-c^2 dx^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.372.1 Optimal result . . . . . 2914  
 3.372.2 Mathematica [A] (warning: unable to verify) . . . . . 2915  
 3.372.3 Rubi [A] (verified) . . . . . 2915  
 3.372.4 Maple [F] . . . . . 2917  
 3.372.5 Fricas [F(-2)] . . . . . 2917  
 3.372.6 Sympy [F] . . . . . 2917  
 3.372.7 Maxima [F] . . . . . 2918  
 3.372.8 Giac [F] . . . . . 2918  
 3.372.9 Mupad [F(-1)] . . . . . 2918

**3.372.1 Optimal result**

Integrand size = 24, antiderivative size = 233

$$\int \frac{d - c^2 dx^2}{(a + b\operatorname{arccosh}(cx))^{3/2}} dx = \frac{2d(-1 + cx)^{3/2}(1 + cx)^{3/2}}{bc\sqrt{a + b\operatorname{arccosh}(cx)}} + \frac{3de^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c} - \frac{de^{\frac{3a}{b}}\sqrt{3\pi}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c} + \frac{3de^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c} - \frac{de^{-\frac{3a}{b}}\sqrt{3\pi}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c}$$

output

```
3/4*d*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c+3/4*d*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c/exp(a/b)-1/4*d*exp(3*a/b)*erf(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/b^(3/2)/c-1/4*d*erfi(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/b^(3/2)/c/exp(3*a/b)+2*d*(c*x-1)^(3/2)*(c*x+1)^(3/2)/b/c/(a+b*arccosh(c*x))^(1/2)
```

**3.372.2 Mathematica [A] (warning: unable to verify)**

Time = 1.10 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.06

$$\int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \frac{e^{-\frac{3a}{b}} \left( -3de^{\frac{4a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) - \sqrt{3}d \sqrt{-\frac{a+b \operatorname{arccosh}(cx)}{b}} \right)}{e^{-\frac{3a}{b}} \left( -3de^{\frac{4a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) - \sqrt{3}d \sqrt{-\frac{a+b \operatorname{arccosh}(cx)}{b}} \right)}$$

input `Integrate[(d - c^2*d*x^2)/(a + b*ArcCosh[c*x])^(3/2),x]`

output `(-3*d*E^((4*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, a/b + ArcCosh[c*x]] - Sqrt[3]*d*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-3*(a + b*ArcCosh[c*x]))/b] + d*E^((2*a)/b)*(3*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, -(a + b*ArcCosh[c*x])/b] + E^(a/b)*(-6*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) + Sqrt[3]*E^((3*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (3*(a + b*ArcCosh[c*x]))/b] + 2*Sinh[3*ArcCosh[c*x]]))/(4*b*c*E^((3*a)/b)*Sqrt[a + b*ArcCosh[c*x]])`

**3.372.3 Rubi [A] (verified)**Time = 0.92 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6319, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx \\ & \quad \downarrow \text{6319} \\ & \frac{2d(cx - 1)^{3/2}(cx + 1)^{3/2}}{bc\sqrt{a + b \operatorname{arccosh}(cx)}} - \frac{6cd \int \frac{x\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{a+b \operatorname{arccosh}(cx)}} dx}{b} \\ & \quad \downarrow \text{6368} \\ & \frac{2d(cx - 1)^{3/2}(cx + 1)^{3/2}}{bc\sqrt{a + b \operatorname{arccosh}(cx)}} - \frac{6d \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b \operatorname{arccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a+b \operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b \operatorname{arccosh}(cx)}} d(a + b \operatorname{arccosh}(cx))}{b^2c} \\ & \quad \downarrow \text{5971} \end{aligned}$$

---


$$3.372. \quad \int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

$$\frac{2d(cx - 1)^{3/2}(cx + 1)^{3/2}}{bc\sqrt{a + \operatorname{barccosh}(cx)}} - \frac{6d \int \left( \frac{\cosh\left(\frac{3a}{b} - \frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{4\sqrt{a + \operatorname{barccosh}(cx)}} - \frac{\cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{4\sqrt{a + \operatorname{barccosh}(cx)}} \right) d(a + \operatorname{barccosh}(cx))}{b^2c}$$

↓ 2009

$$\frac{2d(cx - 1)^{3/2}(cx + 1)^{3/2}}{bc\sqrt{a + \operatorname{barccosh}(cx)}} - \frac{6d \left( -\frac{1}{8}\sqrt{\pi}\sqrt{b}e^{a/b}\operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{8}\sqrt{\frac{\pi}{3}}\sqrt{b}e^{\frac{3a}{b}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{8}\sqrt{\pi}\sqrt{b}e^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right) \right)}{b^2c}$$

input `Int[(d - c^2*d*x^2)/(a + b*ArcCosh[c*x])^(3/2), x]`

output `(2*d*(-1 + c*x)^(3/2)*(1 + c*x)^(3/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]]) - (6*d*(-1/8*(Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]) + (Sqrt[b]*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/8 - (Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(8*E^(a/b)) + (Sqrt[b]*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(8*E^((3*a)/b)))/(b^2*c)`

### 3.372.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

$$3.372. \int \frac{d - c^2 dx^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

```
rule 6368 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]
```

### 3.372.4 Maple [F]

$$\int \frac{-c^2 dx^2 + d}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

```
input int((-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)
```

```
output int((-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)
```

### 3.372.5 Fracas [F(-2)]

Exception generated.

$$\int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")
```

```
output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)
```

### 3.372.6 Sympy [F]

$$\int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = -d \left( \int \frac{c^2 x^2}{a \sqrt{a + b \operatorname{arccosh}(cx)} + b \sqrt{a + b \operatorname{arccosh}(cx)} \operatorname{arccosh}(cx)} dx + \int \left( -\frac{1}{a \sqrt{a + b \operatorname{arccosh}(cx)} + b \sqrt{a + b \operatorname{arccosh}(cx)} \operatorname{arccosh}(cx)} \right) dx \right)$$

---

3.372.  $\int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$

input `integrate((-c**2*d*x**2+d)/(a+b*acosh(c*x))**(3/2),x)`

output `-d*(Integral(c**2*x**2/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(-1/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

### 3.372.7 Maxima [F]

$$\int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int -\frac{c^2 dx^2 - d}{(b \operatorname{arccosh}(cx) + a)^{3/2}} dx$$

input `integrate((-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `-integrate((c^2*d*x^2 - d)/(b*arccosh(c*x) + a)^(3/2), x)`

### 3.372.8 Giac [F]

$$\int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int -\frac{c^2 dx^2 - d}{(b \operatorname{arccosh}(cx) + a)^{3/2}} dx$$

input `integrate((-c^2*d*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `integrate(-(c^2*d*x^2 - d)/(b*arccosh(c*x) + a)^(3/2), x)`

### 3.372.9 Mupad [F(-1)]

Timed out.

$$\int \frac{d - c^2 dx^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{d - c^2 dx^2}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((d - c^2*d*x^2)/(a + b*acosh(c*x))^(3/2),x)`

output `int((d - c^2*d*x^2)/(a + b*acosh(c*x))^(3/2), x)`

**3.373**  $\int \frac{d-c^2 dx^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.373.1 Optimal result . . . . . 2919  
 3.373.2 Mathematica [N/A] . . . . . 2920  
 3.373.3 Rubi [N/A] . . . . . 2920  
 3.373.4 Maple [N/A] (verified) . . . . . 2923  
 3.373.5 Fricas [F(-2)] . . . . . 2924  
 3.373.6 Sympy [N/A] . . . . . 2924  
 3.373.7 Maxima [N/A] . . . . . 2925  
 3.373.8 Giac [F(-2)] . . . . . 2925  
 3.373.9 Mupad [N/A] . . . . . 2925

**3.373.1 Optimal result**

Integrand size = 27, antiderivative size = 27

$$\int \frac{d-c^2 dx^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx = \frac{2d(-1+cx)^{3/2}(1+cx)^{3/2}}{bcx\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{de^{\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}} - \frac{de^{-\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}} + \frac{2d\operatorname{Int}\left(\frac{1}{x^2\sqrt{-1+cx}\sqrt{1+cx}\sqrt{a+b\operatorname{arccosh}(cx)}}, x\right)}{bc}$$

```
output -1/2*d*exp(2*a/b)*erf(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(1/2)*Pi
^(1/2)/b^(3/2)-1/2*d*erfi(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(1/2)
)*Pi^(1/2)/b^(3/2)/exp(2*a/b)+2*d*(c*x-1)^(3/2)*(c*x+1)^(3/2)/b/c/x/(a+b*a
rccosh(c*x))^(1/2)+2*d*Unintegrable(1/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)/(a+b
*arccosh(c*x))^(1/2),x)/b/c
```



**3.373.2 Mathematica [N/A]**

Not integrable

Time = 1.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

input `Integrate[(d - c^2*d*x^2)/(x*(a + b*ArcCosh[c*x])^(3/2)), x]`output `Integrate[(d - c^2*d*x^2)/(x*(a + b*ArcCosh[c*x])^(3/2)), x]`**3.373.3 Rubi [N/A]**

Not integrable

Time = 3.30 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6357, 6322, 3042, 25, 3793, 2009, 6370, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx \\ & \quad \downarrow \text{6357} \\ & -\frac{2d \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2 \sqrt{a+\operatorname{barccosh}(cx)}} dx}{bc} - \frac{4cd \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} + \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx \sqrt{a + \operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{6322} \\ & -\frac{4d \int \frac{\sinh^2\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{b^2} - \frac{2d \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2 \sqrt{a+\operatorname{barccosh}(cx)}} dx}{bc} + \\ & \quad \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx \sqrt{a + \operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.373.  $\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
 & \frac{4d \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2} - \frac{2d \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} + \\
 & \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{25} \\
 & \frac{4d \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2} - \frac{2d \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} + \\
 & \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{3793} \\
 & \frac{4d \int \left( \frac{1}{2\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{b^2} - \\
 & \frac{2d \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} + \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{2d \int \frac{\sqrt{cx-1}\sqrt{cx+1}}{x^2\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} - \\
 & \frac{4d \left( \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{-\frac{2a}{b}} \operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \sqrt{a+b\operatorname{arccosh}(cx)} \right)}{b^2} + \\
 & \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{6370} \\
 & \frac{2d \int \left( \frac{c^2}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{1}{x^2\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+b\operatorname{arccosh}(cx)}} \right) dx}{bc} - \\
 & \frac{4d \left( \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{-\frac{2a}{b}} \operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \sqrt{a+b\operatorname{arccosh}(cx)} \right)}{b^2} + \\
 & \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.373.  $\int \frac{d-c^2 dx^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

$$\frac{2d \left( \frac{2c\sqrt{a+\operatorname{barccosh}(cx)}}{b} - \int \frac{1}{x^2\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}} dx \right)}{bc} - \frac{4d \left( \frac{1}{4}\sqrt{\frac{\pi}{2}}\sqrt{b}e^{\frac{2a}{b}} \operatorname{erf} \left( \frac{\sqrt{2}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\sqrt{b}e^{-\frac{2a}{b}} \operatorname{erfi} \left( \frac{\sqrt{2}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) - \sqrt{a+\operatorname{barccosh}(cx)} \right)}{b^2} + \frac{2d(cx-1)^{3/2}(cx+1)^{3/2}}{bcx\sqrt{a+\operatorname{barccosh}(cx)}}$$

input `Int[(d - c^2*d*x^2)/(x*(a + b*ArcCosh[c*x])^(3/2)),x]`

output `$Aborted`

### 3.373.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6322 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(q_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^q/(-1 + c*x)^p] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[2*p, 0]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6370 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_)^2)^(p_)*((d2_) + (e2_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x]), (f*x)^m*(d1 + e1*x)^(p + 1/2)*(d2 + e2*x)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[d1, 0] && LtQ[d2, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.373.4 Maple [N/A] (verified)

Not integrable

Time = 1.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{-c^2 dx^2 + d}{x (a + b \operatorname{arccosh}(cx))^{\frac{3}{2}}} dx$$

input `int((-c^2*d*x^2+d)/x/(a+b*arccosh(c*x))^(3/2),x)`

output `int((-c^2*d*x^2+d)/x/(a+b*arccosh(c*x))^(3/2),x)`

**3.373.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-c**2*d*x**2+d)/x/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.373.6 Sympy [N/A]**

Not integrable

Time = 8.80 (sec) , antiderivative size = 87, normalized size of antiderivative = 3.22

$$\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx =$$

$$-d \left( \int \frac{c^2 x^2}{ax \sqrt{a + b \operatorname{acosh}(cx)} + bx \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right.$$

$$\left. + \int \left( -\frac{1}{ax \sqrt{a + b \operatorname{acosh}(cx)} + bx \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right)$$

input `integrate((-c**2*d*x**2+d)/x/(a+b*acosh(c*x))**(3/2),x)`

output `-d*(Integral(c**2*x**2/(a*x*sqrt(a + b*acosh(c*x)) + b*x*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(-1/(a*x*sqrt(a + b*acosh(c*x)) + b*x*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

**3.373.7 Maxima [N/A]**

Not integrable

Time = 1.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.11

$$\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int -\frac{c^2 dx^2 - d}{(b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}} x} dx$$

input `integrate((-c^2*d*x^2+d)/x/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `-integrate((c^2*d*x^2 - d)/((b*arccosh(c*x) + a)^(3/2)*x), x)`

**3.373.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)/x/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.373.9 Mupad [N/A]**

Not integrable

Time = 3.92 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{d - c^2 dx^2}{x(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((d - c^2*d*x^2)/(x*(a + b*acosh(c*x))^(3/2)),x)`

output `int((d - c^2*d*x^2)/(x*(a + b*acosh(c*x))^(3/2)), x)`

---

3.373.  $\int \frac{d - c^2 dx^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx$

**3.374** 
$$\int \frac{x^3(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

3.374.1 Optimal result . . . . . 2926  
 3.374.2 Mathematica [A] (warning: unable to verify) . . . . . 2927  
 3.374.3 Rubi [A] (verified) . . . . . 2928  
 3.374.4 Maple [F] . . . . . 2930  
 3.374.5 Fricas [F(-2)] . . . . . 2930  
 3.374.6 Sympy [F] . . . . . 2931  
 3.374.7 Maxima [F] . . . . . 2931  
 3.374.8 Giac [F(-2)] . . . . . 2932  
 3.374.9 Mupad [F(-1)] . . . . . 2932

**3.374.1 Optimal result**

Integrand size = 29, antiderivative size = 479

$$\int \frac{x^3(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = -\frac{2d^2x^3(-1+cx)^{5/2}(1+cx)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{d^2e^{\frac{4a}{b}}\sqrt{\pi}\operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4} + \frac{3d^2e^{\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4} + \frac{d^2e^{\frac{8a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\frac{2\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4} - \frac{d^2e^{\frac{6a}{b}}\sqrt{\frac{3\pi}{2}}\operatorname{erf}\left(\frac{\sqrt{6}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4} - \frac{d^2e^{-\frac{4a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4} + \frac{3d^2e^{-\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4} + \frac{d^2e^{-\frac{8a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\frac{2\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4} - \frac{d^2e^{-\frac{6a}{b}}\sqrt{\frac{3\pi}{2}}\operatorname{erfi}\left(\frac{\sqrt{6}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32b^{3/2}c^4}$$

output 
$$\begin{aligned} & \frac{3}{64}d^2 \exp(2a/b) \operatorname{erf}(2^{1/2}(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2})^2 \cdot 2^{1/2} \cdot \pi^{1/2}/b^{3/2}/c^4 + \frac{1}{64}d^2 \exp(8a/b) \operatorname{erf}(2 \cdot 2^{1/2}(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2})^2 \cdot 2^{1/2} \cdot \pi^{1/2}/b^{3/2}/c^4 + \frac{3}{64}d^2 \operatorname{erfi}(2^{1/2}(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2})^2 \cdot 2^{1/2} \cdot \pi^{1/2}/b^{3/2}/c^4 \\ & + \frac{1}{64}d^2 \exp(2a/b) \operatorname{erfi}(2 \cdot 2^{1/2}(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2})^2 \cdot 2^{1/2} \cdot \pi^{1/2}/b^{3/2}/c^4 + \frac{1}{32}d^2 \exp(4a/b) \operatorname{erf}(2(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2}) \cdot \pi^{1/2}/b^{3/2}/c^4 \\ & - \frac{1}{32}d^2 \operatorname{erfi}(2(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2}) \cdot \pi^{1/2}/b^{3/2}/c^4 + \frac{1}{64}d^2 \exp(6a/b) \operatorname{erf}(6^{1/2}(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2}) \cdot 6^{1/2} \cdot \pi^{1/2}/b^{3/2}/c^4 \\ & - \frac{1}{64}d^2 \operatorname{erfi}(6^{1/2}(a+b\operatorname{arccosh}(cx))^{1/2}/b^{1/2}) \cdot 6^{1/2} \cdot \pi^{1/2}/b^{3/2}/c^4 + \frac{1}{64}d^2 \exp(6a/b) \cdot 2 \cdot d^2 \cdot x^3 \cdot (cx-1)^{5/2} \cdot (cx+1)^{5/2} / bc \cdot (a+b\operatorname{arccosh}(cx))^{1/2} \end{aligned}$$

### 3.374.2 Mathematica [A] (warning: unable to verify)

Time = 2.77 (sec) , antiderivative size = 527, normalized size of antiderivative = 1.10

$$\int \frac{x^3(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = d^2 e^{-\frac{8a}{b}} \left( 128c^3 e^{\frac{8a}{b}} x^3 \sqrt{\frac{-1+cx}{1+cx}} + 128c^4 e^{\frac{8a}{b}} x^4 \sqrt{\frac{-1+cx}{1+cx}} - \sqrt{2} \sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right) \right) + \sqrt{6}$$

input `Integrate[(x^3*(d - c^2*d*x^2)^2)/(a + b*ArcCosh[c*x])^(3/2), x]`

output 
$$\begin{aligned} & -\frac{1}{64}d^2 \left( 128c^3 E^{\frac{8a}{b}} x^3 \sqrt{\frac{-1+cx}{1+cx}} + 128c^4 E^{\frac{8a}{b}} x^4 \sqrt{\frac{-1+cx}{1+cx}} - \sqrt{2} \sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right) \right) + \sqrt{6} \\ & - \sqrt{2} \sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{8(a+b\operatorname{arccosh}(cx))}{b}\right) + \sqrt{6} E^{\frac{2a}{b}} \sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) + 2 E^{\frac{4a}{b}} \sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) \\ & - 3 \sqrt{2} E^{\frac{6a}{b}} \sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) + 3 \sqrt{2} E^{\frac{10a}{b}} \sqrt{\frac{a}{b} + \operatorname{ArcCosh}[cx]} \Gamma\left(\frac{1}{2}, \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right) - 2 E^{\frac{12a}{b}} \sqrt{\frac{a}{b} + \operatorname{ArcCosh}[cx]} \Gamma\left(\frac{1}{2}, \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right) \\ & - \sqrt{6} E^{\frac{14a}{b}} \sqrt{\frac{a}{b} + \operatorname{ArcCosh}[cx]} \Gamma\left(\frac{1}{2}, \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right) + \sqrt{2} E^{\frac{16a}{b}} \sqrt{\frac{a}{b} + \operatorname{ArcCosh}[cx]} \Gamma\left(\frac{1}{2}, \frac{8(a+b\operatorname{arccosh}(cx))}{b}\right) - 26 E^{\frac{8a}{b}} \operatorname{Sinh}[2\operatorname{ArcCosh}[cx]] - 18 E^{\frac{8a}{b}} \operatorname{Sinh}[4\operatorname{ArcCosh}[cx]] \\ & - 2 E^{\frac{8a}{b}} \operatorname{Sinh}[6\operatorname{ArcCosh}[cx]] + E^{\frac{8a}{b}} \operatorname{Sinh}[8\operatorname{ArcCosh}[cx]] \end{aligned}$$

$$3.374. \int \frac{x^3(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$



**3.374.3 Rubi [A] (verified)**

Time = 2.27 (sec) , antiderivative size = 571, normalized size of antiderivative = 1.19, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$ , Rules used = {6357, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(d-c^2dx^2)^2}{(a+\operatorname{barccosh}(cx))^{3/2}} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{16cd^2 \int \frac{x^4(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} - \frac{6d^2 \int \frac{x^2(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{bc} - \frac{2d^2x^3(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\
 & \quad \downarrow \text{6368} \\
 & \frac{16d^2 \int \frac{\cosh^4\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a+\operatorname{barccosh}(cx))}{b^2c^4} - \\
 & \frac{6d^2 \int \frac{\cosh^2\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b}-\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a+\operatorname{barccosh}(cx))}{b^2c^4} - \\
 & \frac{2d^2x^3(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\
 & \quad \downarrow \text{5971} \\
 & \frac{16d^2 \int \left( \frac{\cosh\left(\frac{8a}{b}-\frac{8(a+\operatorname{barccosh}(cx))}{b}\right)}{128\sqrt{a+\operatorname{barccosh}(cx)}} - \frac{\cosh\left(\frac{4a}{b}-\frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{a+\operatorname{barccosh}(cx)}} + \frac{3}{128\sqrt{a+\operatorname{barccosh}(cx)}} \right) d(a+\operatorname{barccosh}(cx))}{b^2c^4} \\
 & \frac{6d^2 \int \left( \frac{\cosh\left(\frac{6a}{b}-\frac{6(a+\operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{a+\operatorname{barccosh}(cx)}} - \frac{\cosh\left(\frac{4a}{b}-\frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{16\sqrt{a+\operatorname{barccosh}(cx)}} - \frac{\cosh\left(\frac{2a}{b}-\frac{2(a+\operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{a+\operatorname{barccosh}(cx)}} + \frac{1}{16\sqrt{a+\operatorname{barccosh}(cx)}} \right) d(a+\operatorname{barccosh}(cx))}{b^2c^4} \\
 & \frac{2d^2x^3(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.374.  $\int \frac{x^3(d-c^2dx^2)^2}{(a+\operatorname{barccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
& 16d^2 \left( -\frac{1}{128} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf} \left( \frac{2\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{512} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{8a}{b}} \operatorname{erf} \left( \frac{2\sqrt{2}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{128} \sqrt{\pi} \sqrt{b} e^{-\frac{4a}{b}} \operatorname{erfi} \left( \frac{2\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) \right. \\
& \left. 6d^2 \left( -\frac{1}{64} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf} \left( \frac{2\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{64} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf} \left( \frac{\sqrt{2}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{64} \sqrt{\frac{\pi}{6}} \sqrt{b} e^{\frac{6a}{b}} \operatorname{erf} \left( \frac{\sqrt{6}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) \right) \right) \\
& \frac{2d^2 x^3 (cx-1)^{5/2} (cx+1)^{5/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}}
\end{aligned}$$

input `Int[(x^3*(d - c^2*d*x^2)^2)/(a + b*ArcCosh[c*x])^(3/2), x]`

output `(-2*d^2*x^3*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]] + (16*d^2*((3*Sqrt[a + b*ArcCosh[c*x]])/64 - (Sqrt[b]*E^((4*a)/b)*Sqrt[Pi]*Erf[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/128 + (Sqrt[b]*E^((8*a)/b)*Sqrt[Pi/2]*Erf[(2*Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/512 - (Sqrt[b]*Sqrt[Pi]*Erfi[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(128*E^((4*a)/b)) + (Sqrt[b]*Sqrt[Pi/2]*Erfi[(2*Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(512*E^((8*a)/b)))/(b^2*c^4) - (6*d^2*(Sqrt[a + b*ArcCosh[c*x]])/8 - (Sqrt[b]*E^((4*a)/b)*Sqrt[Pi]*Erf[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 - (Sqrt[b]*E^((2*a)/b)*Sqrt[Pi/2]*Erf[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 + (Sqrt[b]*E^((6*a)/b)*Sqrt[Pi/6]*Erf[(Sqrt[6]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 - (Sqrt[b]*Sqrt[Pi]*Erfi[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((4*a)/b)) - (Sqrt[b]*Sqrt[Pi/2]*Erfi[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((2*a)/b)) + (Sqrt[b]*Sqrt[Pi/6]*Erfi[(Sqrt[6]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((6*a)/b)))/(b^2*c^4)`

### 3.374.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

$$3.374. \quad \int \frac{x^3(d-c^2 dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_)*((d1_) + (e1_.)*(x_))^(p_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.374.4 Maple [F]

$$\int \frac{x^3(-c^2dx^2 + d)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

input `int(x^3*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

output `int(x^3*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

### 3.374.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{x^3(d - c^2dx^2)^2}{(a + b\operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

---

3.374.  $\int \frac{x^3(d - c^2dx^2)^2}{(a + b\operatorname{arccosh}(cx))^{3/2}} dx$

output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

### 3.374.6 Sympy [F]

$$\int \frac{x^3(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = d^2 \left( \int \frac{x^3}{a \sqrt{a + b \operatorname{acosh}(cx)} + b \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right. \\ \left. + \int \left( -\frac{2c^2 x^5}{a \sqrt{a + b \operatorname{acosh}(cx)} + b \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right. \\ \left. + \int \frac{c^4 x^7}{a \sqrt{a + b \operatorname{acosh}(cx)} + b \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right)$$

input `integrate(x**3*(-c**2*d*x**2+d)**2/(a+b*acosh(c*x))**(3/2),x)`

output `d**2*(Integral(x**3/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(-2*c**2*x**5/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**4*x**7/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

### 3.374.7 Maxima [F]

$$\int \frac{x^3(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{(c^2 dx^2 - d)^2 x^3}{(b \operatorname{arccosh}(cx) + a)^{3/2}} dx$$

input `integrate(x^3*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((c^2*d*x^2 - d)^2*x^3/(b*arccosh(c*x) + a)^(3/2), x)`

**3.374.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^3*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve cteur & l) Error: Bad Argument Value`

**3.374.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{x^3(d - c^2 dx^2)^2}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((x^3*(d - c^2*d*x^2)^2)/(a + b*acosh(c*x))^(3/2),x)`

output `int((x^3*(d - c^2*d*x^2)^2)/(a + b*acosh(c*x))^(3/2), x)`

**3.375** 
$$\int \frac{x^2(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

3.375.1 Optimal result . . . . . 2933  
 3.375.2 Mathematica [A] (warning: unable to verify) . . . . . 2934  
 3.375.3 Rubi [A] (verified) . . . . . 2935  
 3.375.4 Maple [F] . . . . . 2937  
 3.375.5 Fracas [F(-2)] . . . . . 2937  
 3.375.6 Sympy [F] . . . . . 2938  
 3.375.7 Maxima [F] . . . . . 2938  
 3.375.8 Giac [F] . . . . . 2939  
 3.375.9 Mupad [F(-1)] . . . . . 2939

**3.375.1 Optimal result**

Integrand size = 29, antiderivative size = 462

$$\int \frac{x^2(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = -\frac{2d^2x^2(-1+cx)^{5/2}(1+cx)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{5d^2e^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3} + \frac{d^2e^{\frac{3a}{b}}\sqrt{3\pi}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3} - \frac{3d^2e^{\frac{5a}{b}}\sqrt{5\pi}\operatorname{erf}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3} + \frac{d^2e^{\frac{7a}{b}}\sqrt{7\pi}\operatorname{erf}\left(\frac{\sqrt{7}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3} + \frac{5d^2e^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3} + \frac{d^2e^{-\frac{3a}{b}}\sqrt{3\pi}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3} - \frac{3d^2e^{-\frac{5a}{b}}\sqrt{5\pi}\operatorname{erfi}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3} + \frac{d^2e^{-\frac{7a}{b}}\sqrt{7\pi}\operatorname{erfi}\left(\frac{\sqrt{7}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{64b^{3/2}c^3}$$

output  $5/64*d^2*\exp(a/b)*\operatorname{erf}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*\pi^{1/2}/b^{3/2}/c^3+5/64*d^2*\operatorname{erfi}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*\pi^{1/2}/b^{3/2}/c^3/\exp(a/b)+1/64*d^2*\exp(3*a/b)*\operatorname{erf}(3^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*3^{1/2}*\pi^{1/2}/b^{3/2}/c^3+1/64*d^2*\operatorname{erfi}(3^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*3^{1/2}*\pi^{1/2}/b^{3/2}/c^3/\exp(3*a/b)-3/64*d^2*\exp(5*a/b)*\operatorname{erf}(5^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*5^{1/2}*\pi^{1/2}/b^{3/2}/c^3-3/64*d^2*\operatorname{erfi}(5^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*5^{1/2}*\pi^{1/2}/b^{3/2}/c^3/\exp(5*a/b)+1/64*d^2*\exp(7*a/b)*\operatorname{erf}(7^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*7^{1/2}*\pi^{1/2}/b^{3/2}/c^3+1/64*d^2*\operatorname{erfi}(7^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*7^{1/2}*\pi^{1/2}/b^{3/2}/c^3/\exp(7*a/b)-2*d^2*x^2*(c*x-1)^{5/2}*(c*x+1)^{5/2}/b/c/(a+b*\operatorname{arccosh}(c*x))^{1/2}$

### 3.375.2 Mathematica [A] (warning: unable to verify)

Time = 2.48 (sec) , antiderivative size = 498, normalized size of antiderivative = 1.08

$$\int \frac{x^2(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx =$$

$$d^2 e^{-\frac{7a}{b}} \left( 10 e^{\frac{7a}{b}} \sqrt{\frac{-1+cx}{1+cx}} + 10 c e^{\frac{7a}{b}} x \sqrt{\frac{-1+cx}{1+cx}} + 5 e^{\frac{8a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) - \sqrt{7} \sqrt{-\frac{a+b \operatorname{arccosh}(cx)}{b}} \right)$$

input `Integrate[(x^2*(d - c^2*d*x^2)^2)/(a + b*ArcCosh[c*x])^(3/2), x]`

output  $-1/64*(d^2*(10*E^{((7*a)/b)}*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] + 10*c*E^{((7*a)/b)}*x*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)] + 5*E^{((8*a)/b)}*\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[1/2, a/b + \operatorname{ArcCosh}[c*x]] - \operatorname{Sqrt}[7]*\operatorname{Sqrt}[-(a + b*\operatorname{ArcCosh}[c*x])/b])*\operatorname{Gamma}[1/2, (-7*(a + b*\operatorname{ArcCosh}[c*x]))/b] + 3*\operatorname{Sqrt}[5]*E^{((2*a)/b)}*\operatorname{Sqrt}[-(a + b*\operatorname{ArcCosh}[c*x])/b])*\operatorname{Gamma}[1/2, (-5*(a + b*\operatorname{ArcCosh}[c*x]))/b] - \operatorname{Sqrt}[3]*E^{((4*a)/b)}*\operatorname{Sqrt}[-(a + b*\operatorname{ArcCosh}[c*x])/b])*\operatorname{Gamma}[1/2, (-3*(a + b*\operatorname{ArcCosh}[c*x]))/b] - 5*E^{((6*a)/b)}*\operatorname{Sqrt}[-(a + b*\operatorname{ArcCosh}[c*x])/b])*\operatorname{Gamma}[1/2, -(a + b*\operatorname{ArcCosh}[c*x])/b] + \operatorname{Sqrt}[3]*E^{((10*a)/b)}*\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[1/2, (3*(a + b*\operatorname{ArcCosh}[c*x]))/b] - 3*\operatorname{Sqrt}[5]*E^{((12*a)/b)}*\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[1/2, (5*(a + b*\operatorname{ArcCosh}[c*x]))/b] + \operatorname{Sqrt}[7]*E^{((14*a)/b)}*\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[1/2, (7*(a + b*\operatorname{ArcCosh}[c*x]))/b] + 2*E^{((7*a)/b)}*\operatorname{Sinh}[3*\operatorname{ArcCosh}[c*x]] - 6*E^{((7*a)/b)}*\operatorname{Sinh}[5*\operatorname{ArcCosh}[c*x]] + 2*E^{((7*a)/b)}*\operatorname{Sinh}[7*\operatorname{ArcCosh}[c*x]]))/(b*c^3*E^{((7*a)/b)}*\operatorname{Sqrt}[a + b*\operatorname{ArcCosh}[c*x]])$

$$3.375. \int \frac{x^2(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

**3.375.3 Rubi [A] (verified)**

Time = 2.40 (sec) , antiderivative size = 723, normalized size of antiderivative = 1.56, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$ , Rules used = {6357, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx \\
 & \quad \downarrow \text{6357} \\
 & \frac{14cd^2 \int \frac{x^3(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{b} - \frac{4d^2 \int \frac{x(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} - \frac{2d^2x^2(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{6368} \\
 & \frac{14d^2 \int \frac{\cosh^3\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2c^3} - \\
 & \frac{4d^2 \int \frac{\cosh\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2c^3} - \\
 & \frac{2d^2x^2(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{5971} \\
 & \frac{14d^2 \int \left( \frac{\cosh\left(\frac{7a}{b}-\frac{7(a+b\operatorname{arccosh}(cx))}{b}\right)}{64\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{5a}{b}-\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{64\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{3 \cosh\left(\frac{3a}{b}-\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{64\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{3 \cosh\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{64\sqrt{a+b\operatorname{arccosh}(cx)}} \right)}{b^2c^3} \\
 & \frac{4d^2 \int \left( \frac{\cosh\left(\frac{5a}{b}-\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{3 \cosh\left(\frac{3a}{b}-\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{\cosh\left(\frac{a}{b}-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{b^2c^3} \\
 & \frac{2d^2x^2(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

---

3.375.  $\int \frac{x^2(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$



$$\frac{4d^2 \left( \frac{1}{16} \sqrt{\pi} \sqrt{b} e^{a/b} \operatorname{erf} \left( \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{32} \sqrt{3\pi} \sqrt{b} e^{\frac{3a}{b}} \operatorname{erf} \left( \frac{\sqrt{3} \sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{32} \sqrt{\frac{\pi}{5}} \sqrt{b} e^{\frac{5a}{b}} \operatorname{erf} \left( \frac{\sqrt{5} \sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)}{14d^2 \left( \frac{3}{128} \sqrt{\pi} \sqrt{b} e^{a/b} \operatorname{erf} \left( \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{128} \sqrt{3\pi} \sqrt{b} e^{\frac{3a}{b}} \operatorname{erf} \left( \frac{\sqrt{3} \sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{128} \sqrt{\frac{\pi}{5}} \sqrt{b} e^{\frac{5a}{b}} \operatorname{erf} \left( \frac{\sqrt{5} \sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)}$$

$$\frac{2d^2 x^2 (cx - 1)^{5/2} (cx + 1)^{5/2}}{bc \sqrt{a + b \operatorname{arccosh}(cx)}}$$

input `Int[(x^2*(d - c^2*d*x^2)^2)/(a + b*ArcCosh[c*x])^(3/2), x]`

output `(-2*d^2*x^2*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]]) - (4*d^2*((Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/16 - (Sqrt[b]*E^((3*a)/b)*Sqrt[3*Pi]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 + (Sqrt[b]*E^((5*a)/b)*Sqrt[Pi/5]*Erf[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 + (Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(16*E^(a/b)) - (Sqrt[b]*Sqrt[3*Pi]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((3*a)/b)) + (Sqrt[b]*Sqrt[Pi/5]*Erfi[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((5*a)/b)))/(b^2*c^3) + (14*d^2*((3*Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/128 - (Sqrt[b]*E^((3*a)/b)*Sqrt[3*Pi]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/128 - (Sqrt[b]*E^((5*a)/b)*Sqrt[Pi/5]*Erf[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/128 + (Sqrt[b]*E^((7*a)/b)*Sqrt[Pi/7]*Erf[(Sqrt[7]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/128 + (3*Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(128*E^(a/b)) - (Sqrt[b]*Sqrt[3*Pi]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(128*E^((3*a)/b)) - (Sqrt[b]*Sqrt[Pi/5]*Erfi[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(128*E^((5*a)/b)) + (Sqrt[b]*Sqrt[Pi/7]*Erfi[(Sqrt[7]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(128*E^((7*a)/b)))/(b^2*c^3)`

### 3.375.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

$$3.375. \quad \int \frac{x^2(d-c^2 dx^2)^2}{(a+b \operatorname{arccosh}(cx))^{3/2}} dx$$

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.375.4 Maple [F]

$$\int \frac{x^2(-c^2dx^2 + d)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

input `int(x^2*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

output `int(x^2*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

### 3.375.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{x^2(d - c^2dx^2)^2}{(a + b\operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^2*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

---

3.375.  $\int \frac{x^2(d - c^2dx^2)^2}{(a + b\operatorname{arccosh}(cx))^{3/2}} dx$

output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

### 3.375.6 Sympy [F]

$$\int \frac{x^2(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = d^2 \left( \int \frac{x^2}{a \sqrt{a + b \operatorname{acosh}(cx)} + b \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right. \\ \left. + \int \left( -\frac{2c^2 x^4}{a \sqrt{a + b \operatorname{acosh}(cx)} + b \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right. \\ \left. + \int \frac{c^4 x^6}{a \sqrt{a + b \operatorname{acosh}(cx)} + b \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right)$$

input `integrate(x**2*(-c**2*d*x**2+d)**2/(a+b*acosh(c*x))**(3/2),x)`

output `d**2*(Integral(x**2/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(-2*c**2*x**4/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**4*x**6/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

### 3.375.7 Maxima [F]

$$\int \frac{x^2(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{(c^2 dx^2 - d)^2 x^2}{(b \operatorname{arccosh}(cx) + a)^{3/2}} dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((c^2*d*x^2 - d)^2*x^2/(b*arccosh(c*x) + a)^(3/2), x)`

**3.375.8 Giac [F]**

$$\int \frac{x^2(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{(c^2 dx^2 - d)^2 x^2}{(b \operatorname{arcosh}(cx) + a)^{3/2}} dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `integrate((c^2*d*x^2 - d)^2*x^2/(b*arccosh(c*x) + a)^(3/2), x)`

**3.375.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{x^2(d - c^2 dx^2)^2}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((x^2*(d - c^2*d*x^2)^2)/(a + b*acosh(c*x))^(3/2),x)`

output `int((x^2*(d - c^2*d*x^2)^2)/(a + b*acosh(c*x))^(3/2), x)`

**3.376** 
$$\int \frac{x(d-c^2 dx^2)^2}{(a+b \operatorname{arccosh}(cx))^{3/2}} dx$$

3.376.1 Optimal result . . . . . 2940  
 3.376.2 Mathematica [F] . . . . . 2941  
 3.376.3 Rubi [A] (verified) . . . . . 2941  
 3.376.4 Maple [F] . . . . . 2944  
 3.376.5 Fricas [F(-2)] . . . . . 2945  
 3.376.6 Sympy [F] . . . . . 2945  
 3.376.7 Maxima [F] . . . . . 2946  
 3.376.8 Giac [F(-2)] . . . . . 2946  
 3.376.9 Mupad [F(-1)] . . . . . 2946

**3.376.1 Optimal result**

Integrand size = 27, antiderivative size = 363

$$\int \frac{x(d-c^2 dx^2)^2}{(a+b \operatorname{arccosh}(cx))^{3/2}} dx = -\frac{2d^2 x(-1+cx)^{5/2}(1+cx)^{5/2}}{bc\sqrt{a+b \operatorname{arccosh}(cx)}} - \frac{d^2 e^{\frac{4a}{b}} \sqrt{\pi} \operatorname{erf}\left(\frac{2\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^2} + \frac{5d^2 e^{\frac{2a}{b}} \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^2} + \frac{d^2 e^{\frac{6a}{b}} \sqrt{\frac{3\pi}{2}} \operatorname{erf}\left(\frac{\sqrt{6}\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^2} - \frac{d^2 e^{-\frac{4a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{2\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^2} + \frac{5d^2 e^{-\frac{2a}{b}} \sqrt{\frac{\pi}{2}} \operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^2} + \frac{d^2 e^{-\frac{6a}{b}} \sqrt{\frac{3\pi}{2}} \operatorname{erfi}\left(\frac{\sqrt{6}\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c^2}$$

```
output 5/32*d^2*exp(2*a/b)*erf(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(1/2)*
Pi^(1/2)/b^(3/2)/c^2+5/32*d^2*erfi(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2)
)*2^(1/2)*Pi^(1/2)/b^(3/2)/c^2/exp(2*a/b)-1/4*d^2*exp(4*a/b)*erf(2*(a+b*a
rccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^2-1/4*d^2*erfi(2*(a+b*arcco
sh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^2/exp(4*a/b)+1/32*d^2*exp(6*a/b
)*erf(6^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*6^(1/2)*Pi^(1/2)/b^(3/2)/c
^2+1/32*d^2*erfi(6^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*6^(1/2)*Pi^(1/2
)/b^(3/2)/c^2/exp(6*a/b)-2*d^2*x*(c*x-1)^(5/2)*(c*x+1)^(5/2)/b/c/(a+b*arcc
osh(c*x))^(1/2)
```

3.376. 
$$\int \frac{x(d-c^2 dx^2)^2}{(a+b \operatorname{arccosh}(cx))^{3/2}} dx$$

## 3.376.2 Mathematica [F]

$$\int \frac{x(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{x(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

input `Integrate[(x*(d - c^2*d*x^2)^2)/(a + b*ArcCosh[c*x])^(3/2), x]`

output `Integrate[(x*(d - c^2*d*x^2)^2)/(a + b*ArcCosh[c*x])^(3/2), x]`

## 3.376.3 Rubi [A] (verified)

Time = 3.68 (sec) , antiderivative size = 567, normalized size of antiderivative = 1.56, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.296$ , Rules used = {6357, 6322, 3042, 3793, 2009, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx \\ & \quad \downarrow \text{6357} \\ & \frac{12cd^2 \int \frac{x^2(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} - \frac{2d^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{bc} - \frac{2d^2 x(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{6322} \\ & - \frac{2d^2 \int \frac{\sinh^4\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{b^2 c^2} + \frac{12cd^2 \int \frac{x^2(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} - \\ & \quad \frac{2d^2 x(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{3042} \\ & - \frac{2d^2 \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^4}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{b^2 c^2} + \frac{12cd^2 \int \frac{x^2(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} - \\ & \quad \frac{2d^2 x(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \end{aligned}$$

---

3.376.  $\int \frac{x(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
& \downarrow \mathbf{3793} \\
& \frac{2d^2 \int \left( \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{3}{8\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^2} + \\
& \frac{12cd^2 \int \frac{x^2(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{b} - \frac{2d^2 x(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow \mathbf{2009} \\
& \frac{12cd^2 \int \frac{x^2(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{b} - \\
& \frac{2d^2 \left( \frac{1}{32} \sqrt{\pi} \sqrt{be}^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{be}^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\pi} \sqrt{be}^{-\frac{4a}{b}} \operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)}{b^2 c^2} \\
& \frac{2d^2 x(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow \mathbf{6368} \\
& \frac{12d^2 \int \frac{\cosh^2\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2 c^2} - \\
& \frac{2d^2 \left( \frac{1}{32} \sqrt{\pi} \sqrt{be}^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{be}^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\pi} \sqrt{be}^{-\frac{4a}{b}} \operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)}{b^2 c^2} \\
& \frac{2d^2 x(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow \mathbf{5971} \\
& \frac{12d^2 \int \left( \frac{\cosh\left(\frac{6a}{b} - \frac{6(a+b\operatorname{arccosh}(cx))}{b}\right)}{32\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{32\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{1}{16\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{b^2 c^2} \\
& \frac{2d^2 \left( \frac{1}{32} \sqrt{\pi} \sqrt{be}^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{be}^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\pi} \sqrt{be}^{-\frac{4a}{b}} \operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)}{b^2 c^2} \\
& \frac{2d^2 x(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \downarrow \mathbf{2009}
\end{aligned}$$

---

3.376.  $\int \frac{x(d-c^2 dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

$$\frac{2d^2 \left( \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf} \left( \frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf} \left( \frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{-\frac{4a}{b}} \operatorname{erfi} \left( \frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)}{12d^2 \left( -\frac{1}{64} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf} \left( \frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{64} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf} \left( \frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{64} \sqrt{\frac{\pi}{6}} \sqrt{b} e^{\frac{6a}{b}} \operatorname{erfi} \left( \frac{\sqrt{6}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)} = \frac{2d^2 x (cx - 1)^{5/2} (cx + 1)^{5/2}}{bc \sqrt{a + b\operatorname{arccosh}(cx)}}$$

input `Int[(x*(d - c^2*d*x^2)^2)/(a + b*ArcCosh[c*x])^(3/2),x]`

output `(-2*d^2*x*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]]) - (2*d^2*((3*Sqrt[a + b*ArcCosh[c*x]])/4 + (Sqrt[b]*E^((4*a)/b)*Sqrt[Pi]*Erf[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 - (Sqrt[b]*E^((2*a)/b)*Sqrt[Pi/2]*Erf[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/4 + (Sqrt[b]*Sqrt[Pi]*Erfi[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((4*a)/b)) - (Sqrt[b]*Sqrt[Pi/2]*Erfi[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(4*E^((2*a)/b)))/(b^2*c^2) + (12*d^2*(Sqrt[a + b*ArcCosh[c*x]]/8 - (Sqrt[b]*E^((4*a)/b)*Sqrt[Pi]*Erf[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 - (Sqrt[b]*E^((2*a)/b)*Sqrt[Pi/2]*Erf[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 + (Sqrt[b]*E^((6*a)/b)*Sqrt[Pi/6]*Erf[(Sqrt[6]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/64 - (Sqrt[b]*Sqrt[Pi]*Erfi[(2*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((4*a)/b)) - (Sqrt[b]*Sqrt[Pi/2]*Erfi[(Sqrt[2]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((2*a)/b)) + (Sqrt[b]*Sqrt[Pi/6]*Erfi[(Sqrt[6]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*E^((6*a)/b)))/(b^2*c^2)`

### 3.376.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

$$3.376. \quad \int \frac{x(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$



rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6322 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[2*p, 0]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.376.4 Maple [F]

$$\int \frac{x(-c^2 d x^2 + d)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

input `int(x*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

output `int(x*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

---

3.376.  $\int \frac{x(d-c^2 dx^2)^2}{(a+b \operatorname{arccosh}(cx))^{3/2}} dx$

**3.376.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{x(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.376.6 Sympy [F]**

$$\begin{aligned} \int \frac{x(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx &= d^2 \left( \int \frac{x}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right. \\ &+ \int \left( -\frac{2c^2 x^3}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \\ &+ \left. \int \frac{c^4 x^5}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right) \end{aligned}$$

input `integrate(x*(-c**2*d*x**2+d)**2/(a+b*acosh(c*x))**(3/2),x)`

output `d**2*(Integral(x/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(-2*c**2*x**3/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**4*x**5/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`

**3.376.7 Maxima [F]**

$$\int \frac{x(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{(c^2 dx^2 - d)^2 x}{(b \operatorname{arcosh}(cx) + a)^{3/2}} dx$$

input `integrate(x*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((c^2*d*x^2 - d)^2*x/(b*arccosh(c*x) + a)^(3/2), x)`

**3.376.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

input `integrate(x*(-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve cteur & l) Error: Bad Argument Value`

**3.376.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{x(d - c^2 dx^2)^2}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((x*(d - c^2*d*x^2)^2)/(a + b*acosh(c*x))^(3/2),x)`

output `int((x*(d - c^2*d*x^2)^2)/(a + b*acosh(c*x))^(3/2), x)`

**3.377**  $\int \frac{(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.377.1 Optimal result . . . . . 2947  
 3.377.2 Mathematica [A] (warning: unable to verify) . . . . . 2948  
 3.377.3 Rubi [A] (verified) . . . . . 2948  
 3.377.4 Maple [F] . . . . . 2950  
 3.377.5 Fricas [F(-2)] . . . . . 2951  
 3.377.6 Sympy [F] . . . . . 2951  
 3.377.7 Maxima [F] . . . . . 2952  
 3.377.8 Giac [F] . . . . . 2952  
 3.377.9 Mupad [F(-1)] . . . . . 2952

**3.377.1 Optimal result**

Integrand size = 26, antiderivative size = 351

$$\int \frac{(d - c^2dx^2)^2}{(a + b\operatorname{arccosh}(cx))^{3/2}} dx = -\frac{2d^2(-1 + cx)^{5/2}(1 + cx)^{5/2}}{bc\sqrt{a + b\operatorname{arccosh}(cx)}} + \frac{5d^2e^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8b^{3/2}c} - \frac{5d^2e^{\frac{3a}{b}}\sqrt{3}\pi\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c} + \frac{d^2e^{\frac{5a}{b}}\sqrt{5}\pi\operatorname{erf}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c} + \frac{5d^2e^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8b^{3/2}c} - \frac{5d^2e^{-\frac{3a}{b}}\sqrt{3}\pi\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c} + \frac{d^2e^{-\frac{5a}{b}}\sqrt{5}\pi\operatorname{erfi}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16b^{3/2}c}$$

```
output 5/8*d^2*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c+
5/8*d^2*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c/exp(a/b)
-5/16*d^2*exp(3*a/b)*erf(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)
*Pi^(1/2)/b^(3/2)/c-5/16*d^2*erfi(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2)
)*3^(1/2)*Pi^(1/2)/b^(3/2)/c/exp(3*a/b)+1/16*d^2*exp(5*a/b)*erf(5^(1/2)*(a
+b*arccosh(c*x))^(1/2)/b^(1/2))*5^(1/2)*Pi^(1/2)/b^(3/2)/c+1/16*d^2*erfi(5
^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*5^(1/2)*Pi^(1/2)/b^(3/2)/c/exp(5*
a/b)-2*d^2*(c*x-1)^(5/2)*(c*x+1)^(5/2)/b/c/(a+b*arccosh(c*x))^(1/2)
```

3.377.  $\int \frac{(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

**3.377.2 Mathematica [A] (warning: unable to verify)**

Time = 1.59 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.10

$$\int \frac{(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx =$$

$$d^2 e^{-\frac{5a}{b}} \left( 20e^{\frac{5a}{b}} \sqrt{\frac{-1+cx}{1+cx}} + 20ce^{\frac{5a}{b}} x \sqrt{\frac{-1+cx}{1+cx}} + 10e^{\frac{6a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) - \sqrt{5} \sqrt{-\frac{a+ba}{b}} \right)$$

input `Integrate[(d - c^2*d*x^2)^2/(a + b*ArcCosh[c*x])^(3/2), x]`

output

$$\frac{-1/16*(d^2*(20*E^{((5*a)/b)*Sqrt[(-1 + c*x)/(1 + c*x)] + 20*c*E^{((5*a)/b)*x}*Sqrt[(-1 + c*x)/(1 + c*x)] + 10*E^{((6*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, a/b + ArcCosh[c*x]] - Sqrt[5]*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-5*(a + b*ArcCosh[c*x])/b)] + 5*Sqrt[3]*E^{((2*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-3*(a + b*ArcCosh[c*x])/b)] - 10*E^{((4*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, -((a + b*ArcCosh[c*x])/b)] - 5*Sqrt[3]*E^{((8*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (3*(a + b*ArcCosh[c*x])/b)] + Sqrt[5]*E^{((10*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (5*(a + b*ArcCosh[c*x])/b)] - 10*E^{((5*a)/b)*Sinh[3*ArcCosh[c*x]] + 2*E^{((5*a)/b)*Sinh[5*ArcCosh[c*x]]})/(b*c*E^{((5*a)/b)*Sqrt[a + b*ArcCosh[c*x]])}$$
**3.377.3 Rubi [A] (verified)**Time = 1.12 (sec) , antiderivative size = 331, normalized size of antiderivative = 0.94, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {6319, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

$$\downarrow \text{6319}$$

$$\frac{10cd^2 \int \frac{x(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx}{b} - \frac{2d^2 (cx-1)^{5/2} (cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}}$$

$$\downarrow \text{6368}$$

---

3.377.  $\int \frac{(d-c^2 dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

$$10d^2 \int \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a + b\operatorname{arccosh}(cx))$$

$$\frac{b^2c}{2d^2(cx-1)^{5/2}(cx+1)^{5/2}} \frac{1}{bc\sqrt{a+b\operatorname{arccosh}(cx)}}$$

↓ 5971

$$10d^2 \int \left( \frac{\cosh\left(\frac{5a}{b} - \frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{3\cosh\left(\frac{3a}{b} - \frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{\cosh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a + b\operatorname{arccosh}(cx))$$

$$\frac{b^2c}{2d^2(cx-1)^{5/2}(cx+1)^{5/2}} \frac{1}{bc\sqrt{a+b\operatorname{arccosh}(cx)}}$$

↓ 2009

$$10d^2 \left( \frac{1}{16} \sqrt{\pi} \sqrt{b} e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{32} \sqrt{3\pi} \sqrt{b} e^{\frac{3a}{b}} \operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\frac{\pi}{5}} \sqrt{b} e^{\frac{5a}{b}} \operatorname{erf}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)$$

$$\frac{2d^2(cx-1)^{5/2}(cx+1)^{5/2}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}}$$

input `Int[(d - c^2*d*x^2)^2/(a + b*ArcCosh[c*x])^(3/2),x]`

output `(-2*d^2*(-1 + c*x)^(5/2)*(1 + c*x)^(5/2))/(b*c*Sqrt[a + b*ArcCosh[c*x]]) + (10*d^2*((Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/16 - (Sqrt[b]*E^((3*a)/b)*Sqrt[3*Pi]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 + (Sqrt[b]*E^((5*a)/b)*Sqrt[Pi/5]*Erf[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/32 + (Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(16*E^(a/b)) - (Sqrt[b]*Sqrt[3*Pi]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((3*a)/b)) + (Sqrt[b]*Sqrt[Pi/5]*Erfi[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(32*E^((5*a)/b)))/(b^2*c)`

---

3.377.  $\int \frac{(d-c^2dx^2)^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

## 3.377.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(q_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^q/(-1 + c*x)^q] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

## 3.377.4 Maple [F]

$$\int \frac{(-c^2 dx^2 + d)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

input `int((-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

output `int((-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

**3.377.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.377.6 Sympy [F]**

$$\int \frac{(d - c^2 dx^2)^2}{(a + \operatorname{barccosh}(cx))^{3/2}} dx = d^2 \left( \int \left( -\frac{2c^2 x^2}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right. \\ \left. + \int \frac{c^4 x^4}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right. \\ \left. + \int \frac{1}{a\sqrt{a + b \operatorname{acosh}(cx)} + b\sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right)$$

input `integrate((-c**2*d*x**2+d)**2/(a+b*acosh(c*x))**(3/2),x)`

output `d**2*(Integral(-2*c**2*x**2/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**4*x**4/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(1/(a*sqrt(a + b*acosh(c*x)) + b*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))`



**3.377.7 Maxima [F]**

$$\int \frac{(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{(c^2 dx^2 - d)^2}{(b \operatorname{arcosh}(cx) + a)^{3/2}} dx$$

input `integrate((-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((c^2*d*x^2 - d)^2/(b*arccosh(c*x) + a)^(3/2), x)`

**3.377.8 Giac [F]**

$$\int \frac{(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{(c^2 dx^2 - d)^2}{(b \operatorname{arcosh}(cx) + a)^{3/2}} dx$$

input `integrate((-c^2*d*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `integrate((c^2*d*x^2 - d)^2/(b*arccosh(c*x) + a)^(3/2), x)`

**3.377.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{(d - c^2 dx^2)^2}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((d - c^2*d*x^2)^2/(a + b*acosh(c*x))^(3/2),x)`

output `int((d - c^2*d*x^2)^2/(a + b*acosh(c*x))^(3/2), x)`

**3.378**  $\int \frac{(d-c^2dx^2)^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.378.1 Optimal result . . . . . 2953  
 3.378.2 Mathematica [N/A] . . . . . 2954  
 3.378.3 Rubi [N/A] . . . . . 2954  
 3.378.4 Maple [N/A] (verified) . . . . . 2957  
 3.378.5 Fricas [F(-2)] . . . . . 2958  
 3.378.6 Sympy [N/A] . . . . . 2958  
 3.378.7 Maxima [N/A] . . . . . 2959  
 3.378.8 Giac [F(-2)] . . . . . 2959  
 3.378.9 Mupad [N/A] . . . . . 2959

**3.378.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(d - c^2dx^2)^2}{x(a + b\operatorname{arccosh}(cx))^{3/2}} dx = -\frac{2d^2(-1 + cx)^{5/2}(1 + cx)^{5/2}}{bcx\sqrt{a + b\operatorname{arccosh}(cx)}} + \frac{d^2e^{\frac{4a}{b}}\sqrt{\pi}\operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}} - \frac{3d^2e^{\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2b^{3/2}} + \frac{d^2e^{-\frac{4a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}} - \frac{3d^2e^{-\frac{2a}{b}}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2b^{3/2}} + \frac{2d^2\operatorname{Int}\left(\frac{1}{x^2\sqrt{-1+cx}\sqrt{1+cx}\sqrt{a+b\operatorname{arccosh}(cx)}}, x\right)}{bc}$$

output

```
-3/4*d^2*exp(2*a/b)*erf(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(1/2)*
Pi^(1/2)/b^(3/2)-3/4*d^2*erfi(2^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*2^(
1/2)*Pi^(1/2)/b^(3/2)/exp(2*a/b)+1/4*d^2*exp(4*a/b)*erf(2*(a+b*arccosh(c*
x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)+1/4*d^2*erfi(2*(a+b*arccosh(c*x))^(1/2
)/b^(1/2))*Pi^(1/2)/b^(3/2)/exp(4*a/b)-2*d^2*(c*x-1)^(5/2)*(c*x+1)^(5/2)/b
/c/x/(a+b*arccosh(c*x))^(1/2)+2*d^2*Unintegrable(1/x^2/(c*x-1)^(1/2)/(c*x+
1)^(1/2)/(a+b*arccosh(c*x))^(1/2),x)/b/c
```

3.378.  $\int \frac{(d-c^2dx^2)^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

**3.378.2 Mathematica [N/A]**

Not integrable

Time = 1.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

input `Integrate[(d - c^2*d*x^2)^2/(x*(a + b*ArcCosh[c*x])^(3/2)), x]`output `Integrate[(d - c^2*d*x^2)^2/(x*(a + b*ArcCosh[c*x])^(3/2)), x]`**3.378.3 Rubi [N/A]**

Not integrable

Time = 4.18 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6357, 6322, 3042, 3793, 2009, 6370, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx \\ & \quad \downarrow \text{6357} \\ & \frac{2d^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x^2 \sqrt{a+\operatorname{barccosh}(cx)}} dx}{bc} + \frac{8cd^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} - \frac{2d^2 (cx-1)^{5/2}(cx+1)^{5/2}}{bcx \sqrt{a + \operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{6322} \\ & \frac{8d^2 \int \frac{\sinh^4\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{b^2} + \frac{2d^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x^2 \sqrt{a+\operatorname{barccosh}(cx)}} dx}{bc} - \\ & \quad \frac{2d^2 (cx-1)^{5/2}(cx+1)^{5/2}}{bcx \sqrt{a + \operatorname{barccosh}(cx)}} \\ & \quad \downarrow \text{3042} \end{aligned}$$

---

3.378.  $\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
& \frac{8d^2 \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)^4}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx))}{b^2} + \frac{2d^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x^2 \sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} - \\
& \frac{2d^2 (cx-1)^{5/2}(cx+1)^{5/2}}{bcx \sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \quad \downarrow \text{3793} \\
& \frac{8d^2 \int \left( \frac{\cosh\left(\frac{4a}{b} - \frac{4(a+b\operatorname{arccosh}(cx))}{b}\right)}{8\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{\cosh\left(\frac{2a}{b} - \frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{2\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{3}{8\sqrt{a+b\operatorname{arccosh}(cx)}} \right) d(a+b\operatorname{arccosh}(cx))}{b^2} + \\
& \frac{2d^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x^2 \sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} - \frac{2d^2 (cx-1)^{5/2}(cx+1)^{5/2}}{bcx \sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \quad \downarrow \text{2009} \\
& \frac{2d^2 \int \frac{(cx-1)^{3/2}(cx+1)^{3/2}}{x^2 \sqrt{a+b\operatorname{arccosh}(cx)}} dx}{bc} + \\
& \frac{8d^2 \left( \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{-\frac{4a}{b}} \operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)}{b^2} \\
& \frac{2d^2 (cx-1)^{5/2}(cx+1)^{5/2}}{bcx \sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \quad \downarrow \text{6370} \\
& \frac{2d^2 \int \left( \frac{x^2 c^4}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{2c^2}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{1}{x^2 \sqrt{cx-1}\sqrt{cx+1}\sqrt{a+b\operatorname{arccosh}(cx)}} \right) dx}{bc} + \\
& \frac{8d^2 \left( \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) - \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf}\left(\frac{\sqrt{2}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{-\frac{4a}{b}} \operatorname{erfi}\left(\frac{2\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)}{b^2} \\
& \frac{2d^2 (cx-1)^{5/2}(cx+1)^{5/2}}{bcx \sqrt{a+b\operatorname{arccosh}(cx)}} \\
& \quad \downarrow \text{2009}
\end{aligned}$$

---

3.378.  $\int \frac{(d-c^2 dx^2)^2}{x(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

$$2d^2 \left( \int \frac{1}{x^2 \sqrt{cx-1} \sqrt{cx+1} \sqrt{a+b \operatorname{arccosh}(cx)}} dx + \frac{\sqrt{\frac{\pi}{2}} c e^{\frac{2a}{b}} \operatorname{erf} \left( \frac{\sqrt{2} \sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right)}{4\sqrt{b}} + \frac{\sqrt{\frac{\pi}{2}} c e^{-\frac{2a}{b}} \operatorname{erfi} \left( \frac{\sqrt{2} \sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right)}{4\sqrt{b}} - 3c \right)$$


---


$$8d^2 \left( \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{\frac{4a}{b}} \operatorname{erf} \left( \frac{2\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{4} \sqrt{\frac{\pi}{2}} \sqrt{b} e^{\frac{2a}{b}} \operatorname{erf} \left( \frac{\sqrt{2} \sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{32} \sqrt{\pi} \sqrt{b} e^{-\frac{4a}{b}} \operatorname{erfi} \left( \frac{2\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)$$


---


$$\frac{2d^2 (cx-1)^{5/2} (cx+1)^{5/2}}{bcx \sqrt{a+b \operatorname{arccosh}(cx)}} \quad b^2$$

input `Int[(d - c^2*d*x^2)^2/(x*(a + b*ArcCosh[c*x])^(3/2)),x]`

output `$Aborted`

### 3.378.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6322 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^q/(-1 + c*x)^q] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[2*p, 0]`

---

3.378.  $\int \frac{(d-c^2 dx^2)^2}{x(a+b \operatorname{arccosh}(cx))^{3/2}} dx$

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6370 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((f_.)*(x_))^(m_)*((d1_) + (e1_.)*(x_)^2)^(p_)*((d2_) + (e2_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/(Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x]), (f*x)^m*(d1 + e1*x)^(p + 1/2)*(d2 + e2*x)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[d1, 0] && LtQ[d2, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.378.4 Maple [N/A] (verified)

Not integrable

Time = 1.23 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 dx^2 + d)^2}{x (a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

input `int((-c^2*d*x^2+d)^2/x/(a+b*arccosh(c*x))^(3/2),x)`

output `int((-c^2*d*x^2+d)^2/x/(a+b*arccosh(c*x))^(3/2),x)`

**3.378.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate((-c**2*d*x**2+d)**2/x/(a+b*arccosh(c*x))**(3/2),x, algorithm="fricas")
```

```
output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)
```

**3.378.6 Sympy [N/A]**

Not integrable

Time = 9.34 (sec) , antiderivative size = 133, normalized size of antiderivative = 4.59

$$\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = d^2 \left( \int \left( -\frac{2c^2 x^2}{ax \sqrt{a + b \operatorname{acosh}(cx)} + bx \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} \right) dx \right. \\ \left. + \int \frac{c^4 x^4}{ax \sqrt{a + b \operatorname{acosh}(cx)} + bx \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right. \\ \left. + \int \frac{1}{ax \sqrt{a + b \operatorname{acosh}(cx)} + bx \sqrt{a + b \operatorname{acosh}(cx)} \operatorname{acosh}(cx)} dx \right)$$

```
input integrate((-c**2*d*x**2+d)**2/x/(a+b*acosh(c*x))**(3/2),x)
```

```
output d**2*(Integral(-2*c**2*x**2/(a*x*sqrt(a + b*acosh(c*x)) + b*x*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(c**4*x**4/(a*x*sqrt(a + b*acosh(c*x)) + b*x*sqrt(a + b*acosh(c*x))*acosh(c*x)), x) + Integral(1/(a*x*sqrt(a + b*acosh(c*x)) + b*x*sqrt(a + b*acosh(c*x))*acosh(c*x)), x))
```

**3.378.7 Maxima [N/A]**

Not integrable

Time = 1.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.03

$$\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{(c^2 dx^2 - d)^2}{(b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}} x} dx$$

input `integrate((-c^2*d*x^2+d)^2/x/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((c^2*d*x^2 - d)^2/((b*arccosh(c*x) + a)^(3/2)*x), x)`

**3.378.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^2/x/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.378.9 Mupad [N/A]**

Not integrable

Time = 3.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{(d - c^2 dx^2)^2}{x(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((d - c^2*d*x^2)^2/(x*(a + b*acosh(c*x))^(3/2)),x)`

output `int((d - c^2*d*x^2)^2/(x*(a + b*acosh(c*x))^(3/2)), x)`

---

3.378.  $\int \frac{(d - c^2 dx^2)^2}{x(a + \operatorname{barccosh}(cx))^{3/2}} dx$



### 3.379 $\int (c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx$

3.379.1 Optimal result . . . . .	2960
3.379.2 Mathematica [A] (verified) . . . . .	2961
3.379.3 Rubi [C] (verified) . . . . .	2961
3.379.4 Maple [F] . . . . .	2968
3.379.5 Fricas [F(-2)] . . . . .	2968
3.379.6 Sympy [F] . . . . .	2969
3.379.7 Maxima [F] . . . . .	2969
3.379.8 Giac [F(-2)] . . . . .	2969
3.379.9 Mupad [F(-1)] . . . . .	2970

#### 3.379.1 Optimal result

Integrand size = 24, antiderivative size = 351

$$\begin{aligned} \int (c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx &= \frac{3}{8}cx\sqrt{c - a^2cx^2}\sqrt{\operatorname{arccosh}(ax)} \\ &+ \frac{1}{4}x(c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} - \frac{c\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^{3/2}}{4a\sqrt{-1 + ax}\sqrt{1 + ax}} \\ &- \frac{c\sqrt{\pi}\sqrt{c - a^2cx^2}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{256a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{c\sqrt{\frac{\pi}{2}}\sqrt{c - a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1 + ax}\sqrt{1 + ax}} \\ &+ \frac{c\sqrt{\pi}\sqrt{c - a^2cx^2}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{256a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{c\sqrt{\frac{\pi}{2}}\sqrt{c - a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1 + ax}\sqrt{1 + ax}} \end{aligned}$$

```
output -1/4*c*arccosh(a*x)^(3/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/32*c*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/32*c*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/256*c*erf(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/256*c*erfi(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/4*x*(-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(1/2)+3/8*c*x*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)
```

**3.379.2 Mathematica [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 154, normalized size of antiderivative = 0.44

$$\int (c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx =$$

$$\frac{c\sqrt{c - a^2 cx^2} \left( -\sqrt{-\operatorname{arccosh}(ax)} \Gamma\left(\frac{3}{2}, -4\operatorname{arccosh}(ax)\right) + 8\sqrt{2}\sqrt{-\operatorname{arccosh}(ax)} \Gamma\left(\frac{3}{2}, -2\operatorname{arccosh}(ax)\right) + \sqrt{\operatorname{arccosh}(ax)} \right)}{128a\sqrt{\frac{-1+ax}{1+ax}}(1+ax)\sqrt{\operatorname{arccosh}(ax)}}$$

input `Integrate[(c - a^2*c*x^2)^(3/2)*Sqrt[ArcCosh[a*x]], x]`output `-1/128*(c*Sqrt[c - a^2*c*x^2]*(-(Sqrt[-ArcCosh[a*x]]*Gamma[3/2, -4*ArcCosh[a*x]]) + 8*Sqrt[2]*Sqrt[-ArcCosh[a*x]]*Gamma[3/2, -2*ArcCosh[a*x]] + Sqrt[ArcCosh[a*x]]*(32*ArcCosh[a*x]^(3/2) + 8*Sqrt[2]*Gamma[3/2, 2*ArcCosh[a*x]] - Gamma[3/2, 4*ArcCosh[a*x]])))/(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])`**3.379.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 2.85 (sec) , antiderivative size = 352, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 17,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.708$ , Rules used = {6312, 25, 6310, 6302, 5971, 27, 3042, 26, 3789, 2611, 2633, 2634, 6308, 6327, 6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{\operatorname{arccosh}(ax)} (c - a^2 cx^2)^{3/2} dx$$

$$\downarrow \text{6312}$$

$$\frac{ac\sqrt{c - a^2 cx^2} \int -\frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \int \sqrt{c - a^2 cx^2} \sqrt{\operatorname{arccosh}(ax)} dx +$$

$$\frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c - a^2 cx^2)^{3/2}$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& -\frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \int \sqrt{c-a^2cx^2} \sqrt{\operatorname{arccosh}(ax)} dx + \\
& \quad \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{6310} \\
& -\frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{3}{4}c \left( -\frac{a\sqrt{c-a^2cx^2} \int \frac{x}{\sqrt{\operatorname{arccosh}(ax)}} dx}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{6302} \\
& -\frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{3}{4}c \left( -\frac{\sqrt{c-a^2cx^2} \int \frac{ax\sqrt{\frac{ax-1}{ax+1}}(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{5971} \\
& -\frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{3}{4}c \left( -\frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sinh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{27}
\end{aligned}$$

$$\begin{aligned}
& \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{3}{4}c & \left( -\frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sinh(2\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \right. \\
& \left. - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{3}{4}c & \left( -\frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int -\frac{i\sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \right. \\
& \left. - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \right) \\
& \quad \downarrow \text{26} \\
& \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{3}{4}c & \left( -\frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{i\sqrt{c-a^2cx^2} \int \frac{\sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \right. \\
& \left. - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \right) \\
& \quad \downarrow \text{3789} \\
& \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{3}{4}c & \left( \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i \int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2}i \int \frac{e^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} \right. \\
& \left. - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2} \right) \\
& \quad \downarrow \text{2611}
\end{aligned}$$

$$\begin{aligned}
& - \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{3}{4}c & \left( \frac{i\sqrt{c-a^2cx^2} \left( i \int e^{2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}}}{2\sqrt{ax-1}\sqrt{ax+1}} \right) \\
& \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}
\end{aligned}$$

↓ 2633

$$\begin{aligned}
\frac{3}{4}c & \left( \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}}}{2\sqrt{ax-1}\sqrt{ax+1}} \right) \\
& \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}
\end{aligned}$$

↓ 2634

$$\begin{aligned}
\frac{3}{4}c & \left( - \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} \right) \\
& \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}
\end{aligned}$$

↓ 6308

$$\begin{aligned}
& - \frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
\frac{3}{4}c & \left( \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2}}{3a\sqrt{ax-1}\sqrt{ax+1}} \right) \\
& \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}
\end{aligned}$$

↓ 6327

$$\frac{ac\sqrt{c-a^2cx^2} \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \left( \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2}}{3a\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}$$

↓ 6367

$$\frac{c\sqrt{c-a^2cx^2} \int \frac{ax \left( \frac{ax-1}{ax+1} \right)^{3/2} (ax+1)^3}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \left( \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2}}{3a\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}$$

↓ 5971

$$\frac{c\sqrt{c-a^2cx^2} \int \left( \frac{\sinh(4\operatorname{arccosh}(ax))}{8\sqrt{\operatorname{arccosh}(ax)}} - \frac{\sinh(2\operatorname{arccosh}(ax))}{4\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \left( \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2}}{3a\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}$$

↓ 2009

$$\frac{c\sqrt{c-a^2cx^2} \left( -\frac{1}{32}\sqrt{\pi} \operatorname{erf}(2\sqrt{\operatorname{arccosh}(ax)}) + \frac{1}{8}\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) + \frac{1}{32}\sqrt{\pi} \operatorname{erfi}(2\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{8}\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \left( \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2}}{3a\sqrt{ax-1}\sqrt{ax+1}} \right) - \frac{1}{4}x\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}$$

input `Int[(c - a^2*c*x^2)^(3/2)*Sqrt[ArcCosh[a*x]], x]`

```
output (x*(c - a^2*c*x^2)^(3/2)*Sqrt[ArcCosh[a*x]])/4 + (c*Sqrt[c - a^2*c*x^2]*(-
1/32*(Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[a*x]]]) + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[A
rcCosh[a*x]]])/8 + (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[a*x]]])/32 - (Sqrt[Pi/2]*
Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/8))/(8*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) +
(3*c*((x*Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]])/2 - (Sqrt[c - a^2*c*x^2]
*ArcCosh[a*x]^(3/2))/(3*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + ((I/8)*Sqrt[c -
a^2*c*x^2]*((-1/2*I)*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + (I/2)*Sq
rt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])))/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x
]))/4
```

### 3.379.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2611 Int[(F_)^((g_)*((e_) + (f_)*(x_)))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] :
> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d
*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

```
rule 2633 Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt
[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{
F, a, b, c, d}, x] && PosQ[b]
```

```
rule 2634 Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt
[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; Fr
eeQ[{F, a, b, c, d}, x] && NegQ[b]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3789 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^(n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`



rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.)), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.379.4 Maple [F]

$$\int (-a^2cx^2 + c)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx$$

input `int((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(1/2),x)`

output `int((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(1/2),x)`

### 3.379.5 Fricas [F(-2)]

Exception generated.

$$\int (c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(1/2),x, algorithm="fricas")`

---

3.379.  $\int (c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx$

output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

### 3.379.6 Sympy [F]

$$\int (c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx = \int (-c(ax - 1)(ax + 1))^{3/2} \sqrt{\operatorname{acosh}(ax)} dx$$

input `integrate((-a**2*c*x**2+c)**(3/2)*acosh(a*x)**(1/2), x)`

output `Integral((-c*(a*x - 1)*(a*x + 1))**(3/2)*sqrt(acosh(a*x)), x)`

### 3.379.7 Maxima [F]

$$\int (c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx = \int (-a^2 cx^2 + c)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(1/2), x, algorithm="maxima")`

output `integrate((-a^2*c*x^2 + c)^(3/2)*sqrt(arccosh(a*x)), x)`

### 3.379.8 Giac [F(-2)]

Exception generated.

$$\int (c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(1/2), x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command: INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

**3.379.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx = \int \sqrt{\operatorname{acosh}(ax)} (c - a^2 cx^2)^{3/2} dx$$

input `int(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(3/2),x)`output `int(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(3/2), x)`

### 3.380 $\int \sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)} dx$

3.380.1 Optimal result . . . . .	2971
3.380.2 Mathematica [A] (verified) . . . . .	2972
3.380.3 Rubi [C] (verified) . . . . .	2972
3.380.4 Maple [F] . . . . .	2976
3.380.5 Fracas [F(-2)] . . . . .	2976
3.380.6 Sympy [F] . . . . .	2976
3.380.7 Maxima [F] . . . . .	2977
3.380.8 Giac [F(-2)] . . . . .	2977
3.380.9 Mupad [F(-1)] . . . . .	2977

#### 3.380.1 Optimal result

Integrand size = 24, antiderivative size = 205

$$\int \sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)} dx = \frac{1}{2}x\sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)} - \frac{\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2}}{3a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$+ \frac{\sqrt{\frac{\pi}{2}} \sqrt{c - a^2cx^2} \operatorname{erf}\left(\sqrt{2} \sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$- \frac{\sqrt{\frac{\pi}{2}} \sqrt{c - a^2cx^2} \operatorname{erfi}\left(\sqrt{2} \sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

```
output -1/3*arccosh(a*x)^(3/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)
+1/32*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)
)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/32*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1
/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/2*x*(-a^
2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)
```

### 3.380.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.57

$$\int \sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)} dx = \frac{\sqrt{-c(-1 + ax)(1 + ax)} \left( 16\operatorname{arccosh}(ax)^2 + 3\sqrt{2} \sqrt{-\operatorname{arccosh}(ax)} \Gamma\left(\frac{3}{2}, -2\operatorname{arccosh}(ax)\right) + 3\sqrt{2} \sqrt{\operatorname{arccosh}(ax)} \right)}{48a \sqrt{\frac{-1+ax}{1+ax}} (1 + ax) \sqrt{\operatorname{arccosh}(ax)}}$$

input `Integrate[Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]], x]`

output `-1/48*(Sqrt[-(c*(-1 + a*x)*(1 + a*x))]*(16*ArcCosh[a*x]^2 + 3*Sqrt[2]*Sqrt[-ArcCosh[a*x]]*Gamma[3/2, -2*ArcCosh[a*x]] + 3*Sqrt[2]*Sqrt[ArcCosh[a*x]]*Gamma[3/2, 2*ArcCosh[a*x]]))/(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])`

### 3.380.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 1.64 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.88, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.458$ , Rules used = {6310, 6302, 5971, 27, 3042, 26, 3789, 2611, 2633, 2634, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{\operatorname{arccosh}(ax)} \sqrt{c - a^2cx^2} dx \\ & \quad \downarrow \text{6310} \\ & -\frac{a\sqrt{c - a^2cx^2} \int \frac{x}{\sqrt{\operatorname{arccosh}(ax)}} dx}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\sqrt{c - a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax - 1}\sqrt{ax + 1}} dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c - a^2cx^2} \\ & \quad \downarrow \text{6302} \\ & -\frac{\sqrt{c - a^2cx^2} \int \frac{ax\sqrt{\frac{ax-1}{ax+1}}(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\sqrt{c - a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax - 1}\sqrt{ax + 1}} dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} + \\ & \quad \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c - a^2cx^2} \end{aligned}$$

$$\begin{aligned}
& \downarrow 5971 \\
& \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sinh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \downarrow 27 \\
& \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\sinh(2\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \downarrow 3042 \\
& \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int -\frac{i \sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \downarrow 26 \\
& -\frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{i\sqrt{c-a^2cx^2} \int \frac{\sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \downarrow 3789 \\
& \frac{i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i \int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2}i \int \frac{e^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \\
& \quad \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \downarrow 2611 \\
& \frac{i\sqrt{c-a^2cx^2} \left( i \int e^{2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \\
& \quad \frac{\sqrt{c-a^2cx^2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \downarrow 2633
\end{aligned}$$

$$\begin{aligned}
& \frac{i\sqrt{c-a^2cx^2}\left(\frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - i\int e^{-2\operatorname{arccosh}(ax)}d\sqrt{\operatorname{arccosh}(ax)}\right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \\
& \frac{\sqrt{c-a^2cx^2}\int\frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}}dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{2634} \\
& -\frac{\sqrt{c-a^2cx^2}\int\frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}}dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{i\sqrt{c-a^2cx^2}\left(\frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)\right)}{8a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{6308} \\
& \frac{i\sqrt{c-a^2cx^2}\left(\frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)\right)}{8a\sqrt{ax-1}\sqrt{ax+1}} - \\
& \frac{\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2}}{3a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2}
\end{aligned}$$

input `Int[Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]], x]`

output `(x*Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]])/2 - (Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2))/(3*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + ((I/8)*Sqrt[c - a^2*c*x^2]*((-1/2*I)*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + (I/2)*Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]]))/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`

### 3.380.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2611 `Int[(F_)^((g_)*((e_) + (f_)*(x_)))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 2633 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_))^2), x_Symbol] :> Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`

rule 2634 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_))^2), x_Symbol] :> Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3789 `Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)], x_Symbol] :> Simp[I/2 Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]`

rule 5971 `Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_))^(m_)*Sinh[(a_) + (b_)*(x_)]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6302 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*(x_)^m, x_Symbol] :> Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6308 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/(Sqrt[(d1_) + (e1_)*(x_)]*Sqrt[(d2_) + (e2_)*(x_)]), x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`



rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

### 3.380.4 Maple [F]

$$\int \sqrt{-a^2 c x^2 + c} \sqrt{\operatorname{arccosh}(ax)} dx$$

input `int((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2),x)`

output `int((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2),x)`

### 3.380.5 Fracas [F(-2)]

Exception generated.

$$\int \sqrt{c - a^2 c x^2} \sqrt{\operatorname{arccosh}(ax)} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.380.6 Sympy [F]

$$\int \sqrt{c - a^2 c x^2} \sqrt{\operatorname{arccosh}(ax)} dx = \int \sqrt{-c(ax - 1)(ax + 1)} \sqrt{\operatorname{acosh}(ax)} dx$$

input `integrate((-a**2*c*x**2+c)**(1/2)*acosh(a*x)**(1/2),x)`

output `Integral(sqrt(-c*(a*x - 1)*(a*x + 1))*sqrt(acosh(a*x)), x)`

**3.380.7 Maxima [F]**

$$\int \sqrt{c - a^2 cx^2} \sqrt{\operatorname{arccosh}(ax)} dx = \int \sqrt{-a^2 cx^2 + c} \sqrt{\operatorname{arccosh}(ax)} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-a^2*c*x^2 + c)*sqrt(arccosh(a*x)), x)`

**3.380.8 Giac [F(-2)]**

Exception generated.

$$\int \sqrt{c - a^2 cx^2} \sqrt{\operatorname{arccosh}(ax)} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.380.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{c - a^2 cx^2} \sqrt{\operatorname{arccosh}(ax)} dx = \int \sqrt{\operatorname{acosh}(ax)} \sqrt{c - a^2 cx^2} dx$$

input `int(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(1/2),x)`

output `int(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(1/2), x)`

**3.381**  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c-a^2cx^2}} dx$

3.381.1 Optimal result . . . . . 2978  
 3.381.2 Mathematica [A] (verified) . . . . . 2978  
 3.381.3 Rubi [A] (verified) . . . . . 2979  
 3.381.4 Maple [A] (verified) . . . . . 2979  
 3.381.5 Fricas [F(-2)] . . . . . 2980  
 3.381.6 Sympy [F] . . . . . 2980  
 3.381.7 Maxima [F] . . . . . 2980  
 3.381.8 Giac [F] . . . . . 2981  
 3.381.9 Mupad [F(-1)] . . . . . 2981

**3.381.1 Optimal result**

Integrand size = 24, antiderivative size = 48

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c-a^2cx^2}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^{3/2}}{3a\sqrt{c-a^2cx^2}}$$

output `2/3*arccosh(a*x)^(3/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(-a^2*c*x^2+c)^(1/2)`

**3.381.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c-a^2cx^2}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^{3/2}}{3a\sqrt{c-a^2cx^2}}$$

input `Integrate[Sqrt[ArcCosh[a*x]]/Sqrt[c - a^2*c*x^2],x]`

output `(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^(3/2))/(3*a*Sqrt[c - a^2*c*x^2])`

**3.381.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c - a^2 cx^2}} dx$$

↓ 6307

$$\frac{2\sqrt{ax-1}\sqrt{ax+1}\operatorname{arccosh}(ax)^{3/2}}{3a\sqrt{c - a^2 cx^2}}$$

input `Int[Sqrt[ArcCosh[a*x]]/Sqrt[c - a^2*c*x^2],x]`

output `(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^(3/2))/(3*a*Sqrt[c - a^2*c*x^2])`

**3.381.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.381.4 Maple [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{2 \operatorname{arccosh}(ax)^{\frac{3}{2}} \sqrt{ax-1} \sqrt{ax+1}}{3\sqrt{-c(ax-1)(ax+1)} a}$	41

input `int(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)`

output  $2/3*\operatorname{arccosh}(a*x)^{(3/2)/(-c*(a*x-1)*(a*x+1))^{(1/2)*(a*x-1)^{(1/2)*(a*x+1)^{(1/2)/a}}$

### 3.381.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c - a^2cx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.381.6 Sympy [F]

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\sqrt{\operatorname{acosh}(ax)}}{\sqrt{-c(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**(1/2)/(-a**2*c*x**2+c)**(1/2), x)`

output `Integral(sqrt(acosh(a*x))/sqrt(-c*(a*x - 1)*(a*x + 1)), x)`

### 3.381.7 Maxima [F]

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\sqrt{\operatorname{arcosh}(ax)}}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(arccosh(a*x))/sqrt(-a^2*c*x^2 + c), x)`

---

3.381.  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c - a^2cx^2}} dx$

**3.381.8 Giac [F]**

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c - a^2 cx^2}} dx = \int \frac{\sqrt{\operatorname{arcosh}(ax)}}{\sqrt{-a^2 cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(arccosh(a*x))/sqrt(-a^2*c*x^2 + c), x)`

**3.381.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{c - a^2 cx^2}} dx = \int \frac{\sqrt{\operatorname{acosh}(ax)}}{\sqrt{c - a^2 cx^2}} dx$$

input `int(acosh(a*x)^(1/2)/(c - a^2*c*x^2)^(1/2),x)`

output `int(acosh(a*x)^(1/2)/(c - a^2*c*x^2)^(1/2), x)`

**3.382** 
$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{3/2}} dx$$

3.382.1 Optimal result	2982
3.382.2 Mathematica [N/A]	2982
3.382.3 Rubi [N/A]	2983
3.382.4 Maple [N/A] (verified)	2984
3.382.5 Fricas [F(-2)]	2984
3.382.6 Sympy [N/A]	2984
3.382.7 Maxima [N/A]	2985
3.382.8 Giac [N/A]	2985
3.382.9 Mupad [N/A]	2985

**3.382.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{3/2}} dx = \frac{x\sqrt{\operatorname{arccosh}(ax)}}{c\sqrt{c-a^2cx^2}} + \frac{a\sqrt{-1+ax}\sqrt{1+ax}\operatorname{Int}\left(\frac{x}{(1-a^2x^2)\sqrt{\operatorname{arccosh}(ax)}}, x\right)}{2c\sqrt{c-a^2cx^2}}$$

output `x*arccosh(a*x)^(1/2)/c/(-a^2*c*x^2+c)^(1/2)+1/2*a*(a*x-1)^(1/2)*(a*x+1)^(1/2)*Unintegrable(x/(-a^2*x^2+1)/arccosh(a*x)^(1/2),x)/c/(-a^2*c*x^2+c)^(1/2)`

**3.382.2 Mathematica [N/A]**

Not integrable

Time = 5.54 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{3/2}} dx$$

input `Integrate[Sqrt[ArcCosh[a*x]]/(c - a^2*c*x^2)^(3/2), x]`

output `Integrate[Sqrt[ArcCosh[a*x]]/(c - a^2*c*x^2)^(3/2), x]`

---

3.382. 
$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{3/2}} dx$$

### 3.382.3 Rubi [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6314, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx$$

↓ 6314

$$\frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)\sqrt{\operatorname{arccosh}(ax)}} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{c\sqrt{c-a^2cx^2}}$$

↓ 6375

$$\frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)\sqrt{\operatorname{arccosh}(ax)}} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{c\sqrt{c-a^2cx^2}}$$

input `Int[Sqrt[ArcCosh[a*x]]/(c - a^2*c*x^2)^(3/2), x]`

output `$Aborted`

#### 3.382.3.1 Defintions of rubi rules used

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

---

3.382.  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx$



**3.382.4 Maple [N/A] (verified)**

Not integrable

Time = 1.45 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(-a^2cx^2 + c)^{\frac{3}{2}}} dx$$

input `int(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(3/2),x)`output `int(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(3/2),x)`**3.382.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`**3.382.6 Sympy [N/A]**

Not integrable

Time = 4.85 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{acosh}(ax)}}{(-c(ax - 1)(ax + 1))^{\frac{3}{2}}} dx$$

input `integrate(acosh(a*x)**(1/2)/(-a**2*c*x**2+c)**(3/2),x)`output `Integral(sqrt(acosh(a*x))/(-c*(a*x - 1)*(a*x + 1))**(3/2), x)`

---

3.382.  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx$

**3.382.7 Maxima [N/A]**

Not integrable

Time = 0.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}(ax)}}{(-a^2cx^2 + c)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`output `integrate(sqrt(arccosh(a*x))/(-a^2*c*x^2 + c)^(3/2), x)`**3.382.8 Giac [N/A]**

Not integrable

Time = 1.96 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}(ax)}}{(-a^2cx^2 + c)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="giac")`output `integrate(sqrt(arccosh(a*x))/(-a^2*c*x^2 + c)^(3/2), x)`**3.382.9 Mupad [N/A]**

Not integrable

Time = 2.79 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{acosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx$$

input `int(acosh(a*x)^(1/2)/(c - a^2*c*x^2)^(3/2),x)`output `int(acosh(a*x)^(1/2)/(c - a^2*c*x^2)^(3/2), x)`

---

3.382.  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{3/2}} dx$

**3.383**  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{5/2}} dx$

3.383.1 Optimal result . . . . . 2986  
 3.383.2 Mathematica [N/A] . . . . . 2987  
 3.383.3 Rubi [N/A] . . . . . 2987  
 3.383.4 Maple [N/A] (verified) . . . . . 2989  
 3.383.5 Fricas [F(-2)] . . . . . 2989  
 3.383.6 Sympy [N/A] . . . . . 2990  
 3.383.7 Maxima [N/A] . . . . . 2990  
 3.383.8 Giac [N/A] . . . . . 2990  
 3.383.9 Mupad [N/A] . . . . . 2991

**3.383.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{5/2}} dx = \frac{x\sqrt{\operatorname{arccosh}(ax)}}{3c(c-a^2cx^2)^{3/2}} + \frac{2x\sqrt{\operatorname{arccosh}(ax)}}{3c^2\sqrt{c-a^2cx^2}}$$

$$+ \frac{a\sqrt{-1+ax}\sqrt{1+ax}\operatorname{Int}\left(\frac{x}{(1-a^2x^2)\sqrt{\operatorname{arccosh}(ax)}}, x\right)}{3c^2\sqrt{c-a^2cx^2}}$$

$$+ \frac{a\sqrt{-1+ax}\sqrt{1+ax}\operatorname{Int}\left(\frac{x}{(-1+a^2x^2)^2\sqrt{\operatorname{arccosh}(ax)}}, x\right)}{6c^2\sqrt{c-a^2cx^2}}$$

output  $1/3*x*\operatorname{arccosh}(a*x)^{(1/2)}/c/(-a^2*c*x^2+c)^{(3/2)}+2/3*x*\operatorname{arccosh}(a*x)^{(1/2)}/c$   
 $^2/(-a^2*c*x^2+c)^{(1/2)}+1/3*a*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}*\operatorname{Unintegrable}(x/($   
 $-a^2*x^2+1)/\operatorname{arccosh}(a*x)^{(1/2)},x)/c^2/(-a^2*c*x^2+c)^{(1/2)}+1/6*a*(a*x-1)^{($   
 $1/2)}*(a*x+1)^{(1/2)}*\operatorname{Unintegrable}(x/(a^2*x^2-1)^2/\operatorname{arccosh}(a*x)^{(1/2)},x)/c^2/$   
 $(-a^2*c*x^2+c)^{(1/2)}$

---

3.383.  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{5/2}} dx$

**3.383.2 Mathematica [N/A]**

Not integrable

Time = 6.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx$$

input `Integrate[Sqrt[ArcCosh[a*x]]/(c - a^2*c*x^2)^(5/2), x]`output `Integrate[Sqrt[ArcCosh[a*x]]/(c - a^2*c*x^2)^(5/2), x]`**3.383.3 Rubi [N/A]**

Not integrable

Time = 0.96 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6316, 6314, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx \\ & \quad \downarrow \text{6316} \\ & \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-ax)^2(ax+1)^2\sqrt{\operatorname{arccosh}(ax)}} dx}{6c^2\sqrt{c-a^2cx^2}} + \frac{2 \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{3/2}} dx}{3c} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{3c(c-a^2cx^2)^{3/2}} \\ & \quad \downarrow \text{6314} \\ & \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-ax)^2(ax+1)^2\sqrt{\operatorname{arccosh}(ax)}} dx}{6c^2\sqrt{c-a^2cx^2}} + \\ & \frac{2 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)\sqrt{\operatorname{arccosh}(ax)}} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{c\sqrt{c-a^2cx^2}} \right)}{3c} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{3c(c-a^2cx^2)^{3/2}} \\ & \quad \downarrow \text{6327} \end{aligned}$$

---

3.383.  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c-a^2cx^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^2 \sqrt{\operatorname{arccosh}(ax)}} dx}{6c^2\sqrt{c-a^2cx^2}} + \\
& 2 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2) \sqrt{\operatorname{arccosh}(ax)}} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{c\sqrt{c-a^2cx^2}} \right) \\
& \frac{\hspace{10em}}{3c} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{3c(c-a^2cx^2)^{3/2}} \\
& \quad \downarrow \text{6375} \\
& \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^2 \sqrt{\operatorname{arccosh}(ax)}} dx}{6c^2\sqrt{c-a^2cx^2}} + \\
& 2 \left( \frac{a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2) \sqrt{\operatorname{arccosh}(ax)}} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{c\sqrt{c-a^2cx^2}} \right) \\
& \frac{\hspace{10em}}{3c} + \frac{x\sqrt{\operatorname{arccosh}(ax)}}{3c(c-a^2cx^2)^{3/2}}
\end{aligned}$$

input `Int[Sqrt[ArcCosh[a*x]]/(c - a^2*c*x^2)^(5/2),x]`

output `$Aborted`

### 3.383.3.1 Defintions of rubi rules used

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6316 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

### 3.383.4 Maple [N/A] (verified)

Not integrable

Time = 1.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(-a^2cx^2 + c)^{\frac{5}{2}}} dx$$

input `int(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(5/2),x)`

output `int(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(5/2),x)`

### 3.383.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(5/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.383.6 Sympy [N/A]**

Not integrable

Time = 98.58 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{acosh}(ax)}}{(-c(ax - 1)(ax + 1))^{5/2}} dx$$

input `integrate(acosh(a*x)**(1/2)/(-a**2*c*x**2+c)**(5/2), x)`output `Integral(sqrt(acosh(a*x))/(-c*(a*x - 1)*(a*x + 1))**(5/2), x)`**3.383.7 Maxima [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}(ax)}}{(-a^2cx^2 + c)^{5/2}} dx$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(5/2), x, algorithm="maxima")`output `integrate(sqrt(arccosh(a*x))/(-a^2*c*x^2 + c)^(5/2), x)`**3.383.8 Giac [N/A]**

Not integrable

Time = 1.95 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}(ax)}}{(-a^2cx^2 + c)^{5/2}} dx$$

input `integrate(arccosh(a*x)^(1/2)/(-a^2*c*x^2+c)^(5/2), x, algorithm="giac")`output `integrate(sqrt(arccosh(a*x))/(-a^2*c*x^2 + c)^(5/2), x)`

---

3.383.  $\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2cx^2)^{5/2}} dx$

**3.383.9 Mupad [N/A]**

Not integrable

Time = 2.75 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{(c - a^2 cx^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{acosh}(ax)}}{(c - a^2 cx^2)^{5/2}} dx$$

input `int(acosh(a*x)^(1/2)/(c - a^2*c*x^2)^(5/2), x)`output `int(acosh(a*x)^(1/2)/(c - a^2*c*x^2)^(5/2), x)`



### 3.384 $\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx$

3.384.1 Optimal result . . . . .	2992
3.384.2 Mathematica [A] (verified) . . . . .	2993
3.384.3 Rubi [A] (verified) . . . . .	2993
3.384.4 Maple [F] . . . . .	3000
3.384.5 Fricas [F(-2)] . . . . .	3000
3.384.6 Sympy [F(-1)] . . . . .	3001
3.384.7 Maxima [F] . . . . .	3001
3.384.8 Giac [F(-2)] . . . . .	3001
3.384.9 Mupad [F(-1)] . . . . .	3002

#### 3.384.1 Optimal result

Integrand size = 24, antiderivative size = 511

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx = \frac{27c\sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)}}{256a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{9acx^2\sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)}}{32\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{3c(1 - a^2x^2)^2 \sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)}}{32a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{3}{8}cx\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2} + \frac{1}{4}x(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} - \frac{3c\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{5/2}}{20a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{3c\sqrt{\pi}\sqrt{c - a^2cx^2} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{2048a\sqrt{-1 + ax}\sqrt{1 + ax}} + \dots$$

output `1/4*x*(-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(3/2)+3/8*c*x*arccosh(a*x)^(3/2)*(-a^2*c*x^2+c)^(1/2)-3/20*c*arccosh(a*x)^(5/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/128*c*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/128*c*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-3/2048*c*erf(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-3/2048*c*erfi(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+27/256*c*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-9/32*a*c*x^2*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/32*c*(-a^2*x^2+1)^2*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)`

**3.384.2 Mathematica [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.39

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx = \frac{c\sqrt{c - a^2cx^2} \left( -384\operatorname{arccosh}(ax)^3 - 480\operatorname{arccosh}(ax) \cosh(2\operatorname{arccosh}(ax)) + 60 \right)}{2560}$$

input `Integrate[(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(3/2), x]`

output `(c*Sqrt[c - a^2*c*x^2]*(-384*ArcCosh[a*x]^3 - 480*ArcCosh[a*x]*Cosh[2*ArcCosh[a*x]] + 60*Sqrt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + 60*Sqrt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]] - 5*Sqrt[-ArcCosh[a*x]]*Gamma[5/2, -4*ArcCosh[a*x]] + 5*Sqrt[ArcCosh[a*x]]*Gamma[5/2, 4*ArcCosh[a*x]] + 640*ArcCosh[a*x]^2*Sinh[2*ArcCosh[a*x]]))/(2560*a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])`

**3.384.3 Rubi [A] (verified)**Time = 4.11 (sec) , antiderivative size = 420, normalized size of antiderivative = 0.82, number of steps used = 16, number of rules used = 15,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$ , Rules used = {6312, 25, 6310, 6299, 6308, 6327, 6329, 6322, 3042, 3793, 2009, 6368, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{3/2} dx$$

$$\downarrow \text{6312}$$

$$\frac{3ac\sqrt{c - a^2cx^2} \int -x(1 - ax)(ax + 1)\sqrt{\operatorname{arccosh}(ax)} dx}{8\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{3}{4}c \int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2} dx + \frac{1}{4}x \operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{3/2}$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& -\frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\sqrt{\operatorname{arccosh}(ax)}dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{3}{4}c \int \sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{3/2}dx + \\
& \quad \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{6310} \\
& -\frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\sqrt{\operatorname{arccosh}(ax)}dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \int x\sqrt{\operatorname{arccosh}(ax)}dx}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{ax-1}\sqrt{ax+1}}dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2} \right) + \\
& \quad \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{6299} \\
& -\frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\sqrt{\operatorname{arccosh}(ax)}dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{ax-1}\sqrt{ax+1}}dx}{2\sqrt{ax-1}\sqrt{ax+1}} \right) + \\
& \quad \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{6308} \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \right) + \\
& \quad \frac{3ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\sqrt{\operatorname{arccosh}(ax)}dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{6327} \\
& \frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \right) + \\
& \quad \frac{3ac\sqrt{c-a^2cx^2} \int x(1-a^2x^2)\sqrt{\operatorname{arccosh}(ax)}dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2}(c-a^2cx^2)^{3/2} \\
& \quad \downarrow \text{6329}
\end{aligned}$$

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \right)$$

$$\frac{3ac\sqrt{c-a^2cx^2} \left( \frac{\int \frac{(ax-1)^{3/2}(ax+1)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx}{8a} - \frac{(1-a^2x^2)^2 \sqrt{\operatorname{arccosh}(ax)}}{4a^2} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x \operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}$$

↓ 6322

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \right)$$

$$\frac{3ac\sqrt{c-a^2cx^2} \left( \frac{\int \frac{(ax-1)^2(ax+1)^2 d\operatorname{arccosh}(ax)}{\sqrt{\operatorname{arccosh}(ax)}}}{8a^2} - \frac{(1-a^2x^2)^2 \sqrt{\operatorname{arccosh}(ax)}}{4a^2} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x \operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}$$

↓ 3042

$$\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \right)$$

$$\frac{3ac\sqrt{c-a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \sqrt{\operatorname{arccosh}(ax)}}{4a^2} + \frac{\int \frac{\sin(i\operatorname{arccosh}(ax))^4 d\operatorname{arccosh}(ax)}{\sqrt{\operatorname{arccosh}(ax)}}}{8a^2} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x \operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}$$

↓ 3793

$$\frac{\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2} \right) + 3ac\sqrt{c-a^2cx^2} \left( \frac{\int \left( -\frac{\cosh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} + \frac{\cosh(4\operatorname{arccosh}(ax))}{8\sqrt{\operatorname{arccosh}(ax)}} + \frac{3}{8\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{8a^2} - \frac{(1-a^2x^2)^2 \sqrt{\operatorname{arccosh}(ax)}}{4a^2} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}}$$

↓ 2009

$$\frac{\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2} \right) + 3ac\sqrt{c-a^2cx^2} \left( \frac{\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{8a^2}}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}}$$

↓ 6368

$$\frac{\frac{3}{4}c \left( -\frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}(ax)} - \frac{\int \frac{a^2x^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2} \right) + 3ac\sqrt{c-a^2cx^2} \left( \frac{\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{8a^2}}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}}$$

↓ 3042

$$\frac{3}{4}c \left( \frac{3a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{\int \frac{\sin\left(i\operatorname{arccosh}(ax) + \frac{\pi}{2}\right)^2}{\sqrt{\operatorname{arccosh}(ax)}} dx}{4a^2} \operatorname{arccosh}(ax) \right)}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2}}{5a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2} \right)$$

$$3ac\sqrt{c - a^2cx^2} \left( \frac{\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{8a^2}} \right)$$

---


$$\frac{1}{4}x\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{3/2}$$

↓ 3793

$$\frac{3}{4}c \left( \frac{3a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{\int \left( \frac{\cosh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} + \frac{1}{2\sqrt{\operatorname{arccosh}(ax)}} \right) dx}{4a^2} \operatorname{arccosh}(ax) \right)}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2}}{5a\sqrt{ax - 1}\sqrt{ax + 1}} \right)$$

$$3ac\sqrt{c - a^2cx^2} \left( \frac{\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{8a^2}} \right)$$

---


$$\frac{1}{4}x\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{3/2}$$

↓ 2009

$$3ac\sqrt{c - a^2cx^2} \left( \frac{\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{8a^2}} \right)$$

$$\frac{3}{4}c \left( \frac{3a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{\frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \sqrt{\operatorname{arccosh}(ax)}}}{4a^2}} \right)}{4\sqrt{ax - 1}\sqrt{ax + 1}} \right)$$

$$\frac{1}{4}x\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{3/2}$$

input `Int[(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(3/2),x]`

output `(x*(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(3/2))/4 - (3*a*c*Sqrt[c - a^2*c*x^2]*(-1/4*((1 - a^2*x^2)^2*Sqrt[ArcCosh[a*x]])/a^2 + ((3*Sqrt[ArcCosh[a*x]])/4 + (Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[a*x]])]/32 - (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]])]/4 + (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[a*x]])]/32 - (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]])]/4)/(8*a^2)))/(8*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) + (3*c*((x*Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2))/2 - (Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(5/2))/(5*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) - (3*a*Sqrt[c - a^2*c*x^2]*((x^2*Sqrt[ArcCosh[a*x]])/2 - (Sqrt[ArcCosh[a*x]] + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]])]/4 + (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]])]/4)/(4*a^2)))/(4*Sqrt[-1 + a*x]*Sqrt[1 + a*x])))/4`

### 3.384.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6299 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcCosh[c*x])^n/(m + 1)), x] - Simp[b*c*(n/(m + 1)) Int[x^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[m, 0] && GtQ[n, 0]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6322 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[2*p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`



rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.384.4 Maple [F]

$$\int (-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{3}{2}} dx$$

input `int((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(3/2),x)`

output `int((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(3/2),x)`

### 3.384.5 Fracas [F(-2)]

Exception generated.

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.384.6 Sympy [F(-1)]**

Timed out.

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx = \text{Timed out}$$

input `integrate((-a**2*c*x**2+c)**(3/2)*acosh(a*x)**(3/2),x)`output `Timed out`**3.384.7 Maxima [F]**

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx = \int (-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{3}{2}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(3/2),x, algorithm="maxima")`output `integrate((-a^2*c*x^2 + c)^(3/2)*arccosh(a*x)^(3/2), x)`**3.384.8 Giac [F(-2)]**

Exception generated.

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(3/2),x, algorithm="giac")`output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.384.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2} dx = \int \operatorname{acosh}(ax)^{3/2} (c - a^2 cx^2)^{3/2} dx$$

input `int(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(3/2),x)`output `int(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(3/2), x)`

### 3.385 $\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2} dx$

3.385.1 Optimal result . . . . .	3003
3.385.2 Mathematica [A] (verified) . . . . .	3004
3.385.3 Rubi [A] (verified) . . . . .	3004
3.385.4 Maple [F] . . . . .	3007
3.385.5 Fracas [F(-2)] . . . . .	3007
3.385.6 Sympy [F] . . . . .	3007
3.385.7 Maxima [F] . . . . .	3008
3.385.8 Giac [F(-2)] . . . . .	3008
3.385.9 Mupad [F(-1)] . . . . .	3008

#### 3.385.1 Optimal result

Integrand size = 24, antiderivative size = 302

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2} dx = \frac{3\sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)}}{16a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{3ax^2\sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)}}{8\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{1}{2}x\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2} - \frac{\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{5/2}}{5a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{3\sqrt{\frac{\pi}{2}}\sqrt{c - a^2cx^2} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{3\sqrt{\frac{\pi}{2}}\sqrt{c - a^2cx^2} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

output

```
1/2*x*arccosh(a*x)^(3/2)*(-a^2*c*x^2+c)^(1/2)-1/5*arccosh(a*x)^(5/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/128*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/128*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+3/16*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-3/8*a*x^2*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)
```

**3.385.2 Mathematica [A] (verified)**

Time = 0.35 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.45

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2} dx = \frac{\sqrt{c - a^2 cx^2} \left( 15\sqrt{2\pi} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + 15\sqrt{2\pi} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) \right)}{640a\sqrt{(-1 + ax)/(1 + ax)}(1 + ax)}$$

input `Integrate[Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2),x]`output `(Sqrt[c - a^2*c*x^2]*(15*Sqrt[2*Pi]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + 15*Sqrt[2*Pi]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]] - 8*Sqrt[ArcCosh[a*x]]*(16*ArcCosh[a*x]^2 + 15*Cosh[2*ArcCosh[a*x]] - 20*ArcCosh[a*x]*Sinh[2*ArcCosh[a*x]])))/(640*a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x))`**3.385.3 Rubi [A] (verified)**Time = 1.45 (sec) , antiderivative size = 203, normalized size of antiderivative = 0.67, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$ , Rules used = {6310, 6299, 6308, 6368, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{arccosh}(ax)^{3/2} \sqrt{c - a^2 cx^2} dx \\ & \quad \downarrow \text{6310} \\ & -\frac{3a\sqrt{c - a^2 cx^2} \int x \sqrt{\operatorname{arccosh}(ax)} dx}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\sqrt{c - a^2 cx^2} \int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{ax - 1}\sqrt{ax + 1}} dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} + \\ & \quad \frac{1}{2} x \operatorname{arccosh}(ax)^{3/2} \sqrt{c - a^2 cx^2} \\ & \quad \downarrow \text{6299} \\ & -\frac{3a\sqrt{c - a^2 cx^2} \left( \frac{1}{2} x^2 \sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4} a \int \frac{x^2}{\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{\operatorname{arccosh}(ax)}} dx \right)}{4\sqrt{ax - 1}\sqrt{ax + 1}} \\ & \quad -\frac{\sqrt{c - a^2 cx^2} \int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{ax - 1}\sqrt{ax + 1}} dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2} x \operatorname{arccosh}(ax)^{3/2} \sqrt{c - a^2 cx^2} \\ & \quad \downarrow \text{6308} \end{aligned}$$

---

3.385.  $\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2} dx$

$$\begin{aligned}
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{1}{4}a \int \frac{x^2}{\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{6368} \\
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{\int \frac{a^2x^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{2}x\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{3042} \\
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{\int \frac{\sin\left(\operatorname{arccosh}(ax) + \frac{\pi}{2}\right)^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{3793} \\
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{\int \left( \frac{\cosh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} + \frac{1}{2\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{2009} \\
& \frac{3a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}(ax)} - \frac{\frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \sqrt{\operatorname{arccosh}(ax)}}{4a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2}}{5a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{3/2}\sqrt{c-a^2cx^2}
\end{aligned}$$

input `Int[Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2),x]`

```
output (x*Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2))/2 - (Sqrt[c - a^2*c*x^2]*ArcCos
h[a*x]^(5/2))/(5*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) - (3*a*Sqrt[c - a^2*c*x^2
]*(x^2*Sqrt[ArcCosh[a*x]])/2 - (Sqrt[ArcCosh[a*x]] + (Sqrt[Pi/2]*Erf[Sqrt
[2]*Sqrt[ArcCosh[a*x]]])/4 + (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])
/4)/(4*a^2))/(4*Sqrt[-1 + a*x]*Sqrt[1 + a*x])
```

### 3.385.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3793 Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

```
rule 6299 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[
x^(m + 1)*((a + b*ArcCosh[c*x])^n/(m + 1)), x] - Simp[b*c*(n/(m + 1)) Int
[x^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x
], x] /; FreeQ[{a, b, c}, x] && IGtQ[m, 0] && GtQ[n, 0]
```

```
rule 6308 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sq
rt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 +
c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[
c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1
] && EqQ[e2, (-c)*d2] && NeQ[n, -1]
```

```
rule 6310 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_
Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(
1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcC
osh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sq
rt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^
(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n
, 0]
```

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.385.4 Maple [F]

$$\int \sqrt{-a^2 c x^2 + c} \operatorname{arccosh}(ax)^{\frac{3}{2}} dx$$

input `int((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(3/2),x)`

output `int((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(3/2),x)`

### 3.385.5 Fracas [F(-2)]

Exception generated.

$$\int \sqrt{c - a^2 c x^2} \operatorname{arccosh}(ax)^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.385.6 Sympy [F]

$$\int \sqrt{c - a^2 c x^2} \operatorname{arccosh}(ax)^{3/2} dx = \int \sqrt{-c(ax - 1)(ax + 1)} \operatorname{acosh}^{\frac{3}{2}}(ax) dx$$

input `integrate((-a**2*c*x**2+c)**(1/2)*acosh(a*x)**(3/2),x)`

output `Integral(sqrt(-c*(a*x - 1)*(a*x + 1))*acosh(a*x)**(3/2), x)`



**3.385.7 Maxima [F]**

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2} dx = \int \sqrt{-a^2 cx^2 + c} \operatorname{arccosh}(ax)^{\frac{3}{2}} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(3/2),x, algorithm="maxima")`

output `integrate(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^(3/2), x)`

**3.385.8 Giac [F(-2)]**

Exception generated.

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(3/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.385.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2} dx = \int \operatorname{acosh}(ax)^{3/2} \sqrt{c - a^2 cx^2} dx$$

input `int(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(1/2),x)`

output `int(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(1/2), x)`

$$3.386 \quad \int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c-a^2cx^2}} dx$$

3.386.1 Optimal result . . . . .	3009
3.386.2 Mathematica [A] (verified) . . . . .	3009
3.386.3 Rubi [A] (verified) . . . . .	3010
3.386.4 Maple [A] (verified) . . . . .	3010
3.386.5 Fracas [F(-2)] . . . . .	3011
3.386.6 Sympy [F] . . . . .	3011
3.386.7 Maxima [F] . . . . .	3011
3.386.8 Giac [F] . . . . .	3012
3.386.9 Mupad [F(-1)] . . . . .	3012

### 3.386.1 Optimal result

Integrand size = 24, antiderivative size = 48

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c-a^2cx^2}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^{5/2}}{5a\sqrt{c-a^2cx^2}}$$

output  $2/5*\operatorname{arccosh}(a*x)^{(5/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/a/(-a^2*c*x^2+c)^{(1/2)}$

### 3.386.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c-a^2cx^2}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^{5/2}}{5a\sqrt{c-a^2cx^2}}$$

input `Integrate[ArcCosh[a*x]^(3/2)/Sqrt[c - a^2*c*x^2],x]`

output  $(2*\operatorname{Sqrt}[-1 + a*x]*\operatorname{Sqrt}[1 + a*x]*\operatorname{ArcCosh}[a*x]^{(5/2)})/(5*a*\operatorname{Sqrt}[c - a^2*c*x^2])$

**3.386.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c - a^2 cx^2}} dx$$

↓ 6307

$$\frac{2\sqrt{ax - 1}\sqrt{ax + 1}\operatorname{arccosh}(ax)^{5/2}}{5a\sqrt{c - a^2 cx^2}}$$

input `Int[ArcCosh[a*x]^(3/2)/Sqrt[c - a^2*c*x^2],x]`

output `(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^(5/2))/(5*a*Sqrt[c - a^2*c*x^2])`

**3.386.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.386.4 Maple [A] (verified)**

Time = 0.55 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{2 \operatorname{arccosh}(ax)^{\frac{5}{2}} \sqrt{ax-1} \sqrt{ax+1}}{5 \sqrt{-c(ax-1)(ax+1)} a}$	41

input `int(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)`

output  $2/5*\operatorname{arccosh}(a*x)^{(5/2)/(-c*(a*x-1)*(a*x+1))^{(1/2)*(a*x-1)^{(1/2)*(a*x+1)^{(1/2)/a}}$

### 3.386.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c - a^2cx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.386.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{acosh}^{\frac{3}{2}}(ax)}{\sqrt{-c(ax-1)(ax+1)}} dx$$

input `integrate(acosh(a*x)**(3/2)/(-a**2*c*x**2+c)**(1/2),x)`

output `Integral(acosh(a*x)**(3/2)/sqrt(-c*(a*x - 1)*(a*x + 1)), x)`

### 3.386.7 Maxima [F]

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{arcosh}(ax)^{\frac{3}{2}}}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^(3/2)/sqrt(-a^2*c*x^2 + c), x)`

---

3.386.  $\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c - a^2cx^2}} dx$

**3.386.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{arcosh}(ax)^{3/2}}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^(3/2)/sqrt(-a^2*c*x^2 + c), x)`

**3.386.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{acosh}(ax)^{3/2}}{\sqrt{c - a^2cx^2}} dx$$

input `int(acosh(a*x)^(3/2)/(c - a^2*c*x^2)^(1/2),x)`

output `int(acosh(a*x)^(3/2)/(c - a^2*c*x^2)^(1/2), x)`

$$3.387 \quad \int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx$$

3.387.1 Optimal result	3013
3.387.2 Mathematica [N/A]	3013
3.387.3 Rubi [N/A]	3014
3.387.4 Maple [N/A] (verified)	3015
3.387.5 Fricas [F(-2)]	3015
3.387.6 Sympy [N/A]	3015
3.387.7 Maxima [N/A]	3016
3.387.8 Giac [N/A]	3016
3.387.9 Mupad [N/A]	3016

### 3.387.1 Optimal result

Integrand size = 24, antiderivative size = 24

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx = \frac{x \operatorname{arccosh}(ax)^{3/2}}{c\sqrt{c-a^2cx^2}} + \frac{3a\sqrt{-1+ax}\sqrt{1+ax} \operatorname{Int}\left(\frac{x\sqrt{\operatorname{arccosh}(ax)}}{1-a^2x^2}, x\right)}{2c\sqrt{c-a^2cx^2}}$$

output `x*arccosh(a*x)^(3/2)/c/(-a^2*c*x^2+c)^(1/2)+3/2*a*(a*x-1)^(1/2)*(a*x+1)^(1/2)*Unintegrable(x*arccosh(a*x)^(1/2)/(-a^2*x^2+1),x)/c/(-a^2*c*x^2+c)^(1/2)`

### 3.387.2 Mathematica [N/A]

Not integrable

Time = 5.25 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx$$

input `Integrate[ArcCosh[a*x]^(3/2)/(c - a^2*c*x^2)^(3/2), x]`

output `Integrate[ArcCosh[a*x]^(3/2)/(c - a^2*c*x^2)^(3/2), x]`

---


$$3.387. \quad \int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx$$

### 3.387.3 Rubi [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6314, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c - a^2cx^2)^{3/2}} dx$$

↓ 6314

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\sqrt{\operatorname{arccosh}(ax)}}{1-a^2x^2} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)^{3/2}}{c\sqrt{c-a^2cx^2}}$$

↓ 6375

$$\frac{3a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x\sqrt{\operatorname{arccosh}(ax)}}{1-a^2x^2} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x\operatorname{arccosh}(ax)^{3/2}}{c\sqrt{c-a^2cx^2}}$$

input `Int[ArcCosh[a*x]^(3/2)/(c - a^2*c*x^2)^(3/2),x]`

output `$Aborted`

#### 3.387.3.1 Defintions of rubi rules used

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.387.4 Maple [N/A] (verified)**

Not integrable

Time = 1.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arccosh}(ax)^{\frac{3}{2}}}{(-a^2cx^2+c)^{\frac{3}{2}}} dx$$

input `int(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(3/2),x)`output `int(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(3/2),x)`**3.387.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`**3.387.6 Sympy [N/A]**

Not integrable

Time = 46.95 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{acosh}^{\frac{3}{2}}(ax)}{(-c(ax-1)(ax+1))^{\frac{3}{2}}} dx$$

input `integrate(acosh(a*x)**(3/2)/(-a**2*c*x**2+c)**(3/2),x)`output `Integral(acosh(a*x)**(3/2)/(-c*(a*x - 1)*(a*x + 1))**(3/2), x)`

---

3.387.  $\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c-a^2cx^2)^{3/2}} dx$



**3.387.7 Maxima [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}(ax)^{\frac{3}{2}}}{(-a^2cx^2 + c)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`output `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2 + c)^(3/2), x)`**3.387.8 Giac [N/A]**

Not integrable

Time = 3.62 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}(ax)^{\frac{3}{2}}}{(-a^2cx^2 + c)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="giac")`output `integrate(arccosh(a*x)^(3/2)/(-a^2*c*x^2 + c)^(3/2), x)`**3.387.9 Mupad [N/A]**

Not integrable

Time = 2.71 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{acosh}(ax)^{3/2}}{(c - a^2cx^2)^{3/2}} dx$$

input `int(acosh(a*x)^(3/2)/(c - a^2*c*x^2)^(3/2),x)`output `int(acosh(a*x)^(3/2)/(c - a^2*c*x^2)^(3/2), x)`

---

3.387.  $\int \frac{\operatorname{arccosh}(ax)^{3/2}}{(c - a^2cx^2)^{3/2}} dx$

### 3.388 $\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx$

3.388.1 Optimal result . . . . .	3017
3.388.2 Mathematica [A] (verified) . . . . .	3018
3.388.3 Rubi [F] . . . . .	3018
3.388.4 Maple [F] . . . . .	3033
3.388.5 Fricas [F(-2)] . . . . .	3033
3.388.6 Sympy [F(-1)] . . . . .	3034
3.388.7 Maxima [F] . . . . .	3034
3.388.8 Giac [F(-2)] . . . . .	3034
3.388.9 Mupad [F(-1)] . . . . .	3035

#### 3.388.1 Optimal result

Integrand size = 24, antiderivative size = 580

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx = \frac{225}{512}cx\sqrt{c - a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}$$

$$+ \frac{15}{256}cx(1 - ax)(1 + ax)\sqrt{c - a^2cx^2}\sqrt{\operatorname{arccosh}(ax)} + \frac{45c\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^{3/2}}{256a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$- \frac{15acx^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^{3/2}}{32\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{5c(1 - a^2x^2)^2\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^{3/2}}{32a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$+ \frac{3}{8}cx\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^{5/2} + \frac{1}{4}x(c - a^2cx^2)^{3/2}\operatorname{arccosh}(ax)^{5/2} - \frac{3c\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^{7/2}}{28a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{15c\sqrt{\pi}\sqrt{c - a^2cx^2}\operatorname{arccosh}(ax)^{5/2}}{16a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

output

```
1/4*x*(-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(5/2)+3/8*c*x*arccosh(a*x)^(5/2)*(-a^2*c*x^2+c)^(1/2)+45/256*c*arccosh(a*x)^(3/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-15/32*a*c*x^2*arccosh(a*x)^(3/2)*(-a^2*c*x^2+c)^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)+5/32*c*(-a^2*x^2+1)^2*arccosh(a*x)^(3/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-3/28*c*arccosh(a*x)^(7/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+15/512*c*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-15/512*c*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-15/16384*c*erf(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+15/16384*c*erfi(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+225/512*c*x*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)+15/256*c*x*(-a*x+1)*(a*x+1)*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)
```

**3.388.2 Mathematica [A] (verified)**

Time = 0.48 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.37

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx = \frac{c\sqrt{c - a^2 cx^2} \left( -1536 \operatorname{arccosh}(ax)^4 - 4480 \operatorname{arccosh}(ax)^2 \cosh(2 \operatorname{arccosh}(ax)) - 4480 \operatorname{arccosh}(ax)^2 \cosh(2 \operatorname{arccosh}(ax)) \right)}{1}$$

input `Integrate[(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(5/2), x]`

output `(c*Sqrt[c - a^2*c*x^2]*(-1536*ArcCosh[a*x]^4 - 4480*ArcCosh[a*x]^2*Cosh[2*ArcCosh[a*x]] + 420*Sqrt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] - 420*Sqrt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + 7*Sqrt[-ArcCosh[a*x]]*Gamma[7/2, -4*ArcCosh[a*x]] + 7*Sqrt[ArcCosh[a*x]]*Gamma[7/2, 4*ArcCosh[a*x]] + 3360*ArcCosh[a*x]*Sinh[2*ArcCosh[a*x]] + 3584*ArcCosh[a*x]^3*Sinh[2*ArcCosh[a*x]])/(14336*a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])`

**3.388.3 Rubi [F]**

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \operatorname{arccosh}(ax)^{5/2} (c - a^2 cx^2)^{3/2} dx \\ & \quad \downarrow \text{6312} \\ & \frac{5ac\sqrt{c - a^2 cx^2} \int -x(1 - ax)(ax + 1)\operatorname{arccosh}(ax)^{3/2} dx}{8\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{3}{4}c \int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2} dx + \\ & \quad \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2 cx^2)^{3/2} \\ & \quad \downarrow \text{25} \\ & -\frac{5ac\sqrt{c - a^2 cx^2} \int x(1 - ax)(ax + 1)\operatorname{arccosh}(ax)^{3/2} dx}{8\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{3}{4}c \int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2} dx + \\ & \quad \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2 cx^2)^{3/2} \\ & \quad \downarrow \text{6310} \end{aligned}$$

$$-\frac{5ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^{3/2} dx}{8\sqrt{ax-1}\sqrt{ax+1}} +$$

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \int x\operatorname{arccosh}(ax)^{3/2} dx}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2} \right) +$$

$$\frac{1}{4}x\operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}$$

↓ 6299

$$-\frac{5ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^{3/2} dx}{8\sqrt{ax-1}\sqrt{ax+1}} +$$

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\sqrt{c-a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2} \right) +$$

$$\frac{1}{4}x\operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}$$

↓ 6308

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2} \right) +$$

$$\frac{5ac\sqrt{c-a^2cx^2} \int x(1-ax)(ax+1)\operatorname{arccosh}(ax)^{3/2} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}$$

↓ 6327

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c-a^2cx^2} \right) +$$

$$\frac{5ac\sqrt{c-a^2cx^2} \int x(1-a^2x^2)\operatorname{arccosh}(ax)^{3/2} dx}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x\operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}$$

↓ 6329

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) - \frac{5ac\sqrt{c-a^2cx^2} \left( \frac{3 \int (ax-1)^{3/2} (ax+1)^{3/2} \sqrt{\operatorname{arccosh}(ax)} dx}{8a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}$$

↓ 6313

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) - \frac{5ac\sqrt{c-a^2cx^2} \left( \frac{3 \left( -\frac{1}{8}a \int -\frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \sqrt{\operatorname{arccosh}(ax)} dx + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \sqrt{\operatorname{arccosh}(ax)} \right)}{8a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}$$

↓ 25

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) - \frac{5ac\sqrt{c-a^2cx^2} \left( \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \int \sqrt{ax-1}\sqrt{ax+1} \sqrt{\operatorname{arccosh}(ax)} dx + \frac{1}{4}x(ax-1)^{3/2}(ax+1)^{3/2} \sqrt{\operatorname{arccosh}(ax)} \right)}{8a} - \frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}$$

↓ 6311

$$\begin{aligned}
 & 5ac\sqrt{c - a^2cx^2} \left( \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( -\frac{1}{4}a \int \frac{x}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{1}{2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)} \right) \right)}{8a} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \\
 & \qquad \qquad \qquad \frac{1}{4}x\operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2} \\
 & \qquad \qquad \qquad \downarrow \text{6302} \\
 & 5ac\sqrt{c - a^2cx^2} \left( \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( -\frac{\int \frac{ax\sqrt{\frac{ax-1}{ax+1}}(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} \operatorname{arccosh}(ax)}{4a} - \frac{1}{2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)} \right) \right)}{8a} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2\operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax) \right) \\
 & \qquad \qquad \qquad \frac{1}{4}x\operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2} \\
 & \qquad \qquad \qquad \downarrow \text{5971}
 \end{aligned}$$

$$\frac{\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{5ac\sqrt{c-a^2cx^2} \left( \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( -\frac{1}{2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\int \frac{\sinh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} a \operatorname{arccosh}(ax) dx + \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)} \right)}{8a} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} \right)}{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}}$$

↓ 27

$$\frac{\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{5ac\sqrt{c-a^2cx^2} \left( \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( -\frac{1}{2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\int \frac{\sinh(2\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} a \operatorname{arccosh}(ax) dx + \frac{1}{2}x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)} \right)}{8a} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} \right)}{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}}$$

↓ 3042

$$\frac{\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{5ac\sqrt{c-a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( -\frac{1}{2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx - \frac{\int -\frac{i \sin(2i \operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax)}{8a} \right) \right)}{4} \right)}$$

---


$$\frac{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}}{8\sqrt{ax-1}\sqrt{ax+1}}$$

↓ 26

$$\frac{\frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right)}{5ac\sqrt{c-a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( -\frac{1}{2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{i \int \frac{\sin(2i \operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d \operatorname{arccosh}(ax)}{8a} \right) \right)}{4} \right)}$$

---


$$\frac{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}}{8\sqrt{ax-1}\sqrt{ax+1}}$$

↓ 3789



$$5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2} \int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2} \int \frac{e^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) \right)}{8a} \right)}{8\sqrt{ax - 1}\sqrt{ax + 1}} \right) \right)$$

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) - \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}$$

↓ 2611

$$5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \int e^{2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{8a} \right)}{8\sqrt{ax - 1}\sqrt{ax + 1}} \right) \right)$$

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) - \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}$$

↓ 2633

$$5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( -\frac{3}{4} \left( \frac{i \left( \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right) - \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)}}{8a} \right) - \frac{1}{2} \int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) \right)}{8\sqrt{ax - 1}\sqrt{ax + 1}} \right) \right)$$

$$\frac{3}{4}c \left( -\frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) - \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}$$

3.388.  $\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx$

$$\begin{aligned}
 & \downarrow 2634 \\
 & \frac{5ac\sqrt{c-a^2cx^2}}{4a^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( -\frac{3}{4} \left( -\frac{1}{2} \int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx + \frac{i \left( \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{ax-1}\sqrt{ax+1}) \right)}{8a} \right)}{8a} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \\
 & \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 6308 \\
 & \frac{5ac\sqrt{c-a^2cx^2}}{4a^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{ax-1}\sqrt{ax+1}) \right)}{8a} \right)}{8a} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \\
 & \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 6327 \\
 & \frac{5ac\sqrt{c-a^2cx^2}}{4a^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{ax-1}\sqrt{ax+1}) \right)}{8a} \right)}{8a} \right)}{8\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( -\frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax) \right) \\
 & \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c-a^2cx^2)^{3/2}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow \text{6354} \\
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a} \right)}{8a} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} - \frac{\int \frac{x}{\sqrt{\operatorname{arccosh}(ax)}} dx}{4a} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}{2a^2} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right)}{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}} \right) \\
 & \downarrow \text{6302} \\
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a} \right)}{8a} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( -\frac{\int \frac{ax\sqrt{\frac{ax-1}{ax+1}}(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{4a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{2a} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right)}{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}} \right) \\
 & \downarrow \text{5971}
 \end{aligned}$$

$$\begin{aligned}
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{ax-1}\sqrt{ax+1}) \right)}{8a} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( -\frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( -\frac{\int \frac{\sinh(2\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{2\sqrt{\operatorname{arccosh}(ax)}} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{8\sqrt{ax-1}\sqrt{ax+1}} \right)}{4a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2} \\
 & \quad \downarrow \quad 27 \\
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{ax-1}\sqrt{ax+1}) \right)}{8a} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( -\frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( -\frac{\int \frac{\sinh(2\operatorname{arccosh}(ax)) d\operatorname{arccosh}(ax)}{\sqrt{\operatorname{arccosh}(ax)}} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{8\sqrt{ax-1}\sqrt{ax+1}} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2} \\
 & \quad \downarrow \quad 3042
 \end{aligned}$$

$$\begin{aligned}
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{\int -\frac{i \sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{8\sqrt{ax-1}\sqrt{ax+1}} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right)}{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}} \right) \\
 & \qquad \qquad \qquad \downarrow \quad 26 \\
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) \right)}{8a} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i \int \frac{\sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}}{8\sqrt{ax-1}\sqrt{ax+1}} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right)}{\frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}} \right) \\
 & \qquad \qquad \qquad \downarrow \quad 3789
 \end{aligned}$$

$$\begin{aligned}
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{ax-1}\sqrt{ax+1}) \right)}{8a} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i \left( \frac{1}{2}i \int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2}i \int \frac{e^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2} \\
 & \quad \downarrow \text{2611} \\
 & 5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8}a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2}\sqrt{ax-1}\sqrt{ax+1}) \right)}{8a} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{3}{4}c \left( \frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i \left( i \int e^{2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{8a^3} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) \\
 & \frac{1}{4}x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2} \\
 & \quad \downarrow \text{2633}
 \end{aligned}$$

$$5ac\sqrt{c - a^2cx^2} \left( -\frac{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}{4a^2} + \frac{3 \left( \frac{1}{8} a \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx - \frac{3}{4} \left( \frac{i \left( \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2} \sqrt{\operatorname{arccosh}(ax)}) - \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erf}(\sqrt{2} \sqrt{ax+1}) \right)}{8a} \right) \right)}{8a^3} \right) + \frac{8\sqrt{ax-1}\sqrt{ax+1}}{4\sqrt{ax-1}\sqrt{ax+1}} \right) + \frac{3}{4} c \left( -\frac{5a\sqrt{c - a^2cx^2} \left( \frac{1}{2} x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4} a \left( \frac{i \left( \frac{1}{2} i \sqrt{\frac{\pi}{2}} \operatorname{erfi}(\sqrt{2} \sqrt{\operatorname{arccosh}(ax)}) - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{8a^3} \right) \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \right) + \frac{1}{4} x \operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}$$

input `Int[(c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(5/2),x]`

output `$Aborted`

### 3.388.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2611 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 2633 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`

rule 2634 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))2), x_Symbol] := Simp[Fa*Sqrt  
[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; Fr  
eeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3789 `Int[((c_.) + (d_.)*(x_))(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I  
/2 Int[(c + d*x)m/EI*(e + f*x)], x] - Simp[I/2 Int[(c + d*x)m*E  
I*(e + f*x)], x] /; FreeQ[{c, d, e, f, m}, x]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)](p_.)*((c_.) + (d_.)*(x_))(m_.)*Sinh[(a_.) +  
(b_.)*(x_)](n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)m, Sinh[a +  
b*x]n*Cosh[a + b*x]p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &  
& IGtQ[p, 0]`

rule 6299 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)(x_)(m_.), x_Symbol] := Simp[  
x(m + 1)*((a + b*ArcCosh[c*x])n/(m + 1)), x] - Simp[b*c*(n/(m + 1)) Int  
[x(m + 1)*((a + b*ArcCosh[c*x])(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x  
, x] /; FreeQ[{a, b, c}, x] && IGtQ[m, 0] && GtQ[n, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)(x_)(m_.), x_Symbol] := Simp[  
1/(b*c(m + 1)) Subst[Int[xn*Cosh[-a/b + x/b]m*Sinh[-a/b + x/b], x], x,  
a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sq  
rt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 +  
c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[  
c*x])(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1  
&& EqQ[e2, (-c)*d2] && NeQ[n, -1]`



rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6311 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)], x_Symbol] := Simp[x*Sqrt[d1 + e1*x]*Sqrt[d2 + e2*x]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d1 + e1*x]/Sqrt[1 + c*x]]*Simp[Sqrt[d2 + e2*x]/Sqrt[-1 + c*x]] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6313 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[x*(d1 + e1*x)^p*(d2 + e2*x)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d1*d2*(p/(2*p + 1)) Int[(d1 + e1*x)^(p - 1)*(d2 + e2*x)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && GtQ[p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]`

rule 6354 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_))*((d1_) + (e1_.)*(x_)^(p_))*((d2_) + (e2_.)*(x_)^(p_)), x_Symbol] := Simp[f*(f*x)^(m - 1)*(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(e1*e2*(m + 2*p + 1))), x] + (Simp[f^2*(m - 1)/(c^2*(m + 2*p + 1)) Int[(f*x)^(m - 2)*(d1 + e1*x)^p*(d2 + e2*x)^p*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*f*(n/(c*(m + 2*p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(f*x)^(m - 1)*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d1, e1, d2, e2, f, p}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && GtQ[n, 0] && IGtQ[m, 1] && NeQ[m + 2*p + 1, 0]`

### 3.388.4 Maple [F]

$$\int (-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{5}{2}} dx$$

input `int((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(5/2),x)`

output `int((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(5/2),x)`

### 3.388.5 Fricas [F(-2)]

Exception generated.

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(5/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

---

3.388.  $\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx$

**3.388.6 Sympy [F(-1)]**

Timed out.

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx = \text{Timed out}$$

input `integrate((-a**2*c*x**2+c)**(3/2)*acosh(a*x)**(5/2),x)`

output `Timed out`

**3.388.7 Maxima [F]**

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx = \int (-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{5}{2}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(5/2),x, algorithm="maxima")`

output `integrate((-a^2*c*x^2 + c)^(3/2)*arccosh(a*x)^(5/2), x)`

**3.388.8 Giac [F(-2)]**

Exception generated.

$$\int (c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)*arccosh(a*x)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.388.9 Mupad [F(-1)]**

Timed out.

$$\int (c - a^2 cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2} dx = \int \operatorname{acosh}(ax)^{5/2} (c - a^2 cx^2)^{3/2} dx$$

input `int(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(3/2),x)`output `int(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(3/2), x)`

### 3.389 $\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{5/2} dx$

3.389.1 Optimal result	3036
3.389.2 Mathematica [A] (verified)	3037
3.389.3 Rubi [C] (verified)	3037
3.389.4 Maple [F]	3043
3.389.5 Fracas [F(-2)]	3043
3.389.6 Sympy [F(-1)]	3043
3.389.7 Maxima [F]	3044
3.389.8 Giac [F(-2)]	3044
3.389.9 Mupad [F(-1)]	3044

#### 3.389.1 Optimal result

Integrand size = 24, antiderivative size = 330

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{5/2} dx = \frac{15}{32} x \sqrt{c - a^2cx^2} \sqrt{\operatorname{arccosh}(ax)}$$

$$+ \frac{5\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2}}{16a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{5ax^2\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{3/2}}{8\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$+ \frac{1}{2} x \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{5/2} - \frac{\sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{7/2}}{7a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{15\sqrt{\frac{\pi}{2}} \sqrt{c - a^2cx^2} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{256a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{15}{256a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

output

```

1/2*x*arccosh(a*x)^(5/2)*(-a^2*c*x^2+c)^(1/2)+5/16*arccosh(a*x)^(3/2)*(-a^
2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-5/8*a*x^2*arccosh(a*x)^(3/2
)*(-a^2*c*x^2+c)^(1/2)/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/7*arccosh(a*x)^(7/2)*
(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+15/512*erf(2^(1/2)*arcc
osh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x
+1)^(1/2)-15/512*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c
*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+15/32*x*(-a^2*c*x^2+c)^(1/2)*a
rccosh(a*x)^(1/2)

```

### 3.389.2 Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.45

$$\int \sqrt{c - a^2cx^2} \operatorname{arccosh}(ax)^{5/2} dx = \frac{\sqrt{-c(-1 + ax)(1 + ax)} \left( -105\sqrt{2\pi} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + 105\sqrt{2\pi} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + 8\sqrt{\operatorname{arccosh}(ax)} \right)}{3584a\sqrt{-c(-1 + ax)(1 + ax)}}$$

input `Integrate[Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(5/2),x]`

output `-1/3584*(Sqrt[-(c*(-1 + a*x)*(1 + a*x))]*(-105*Sqrt[2*Pi]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + 105*Sqrt[2*Pi]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + 8*Sqrt[ArcCosh[a*x]]*(64*ArcCosh[a*x]^3 + 140*ArcCosh[a*x]*Cosh[2*ArcCosh[a*x]] - 7*(15 + 16*ArcCosh[a*x]^2)*Sinh[2*ArcCosh[a*x]])))/(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x))`

### 3.389.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 2.60 (sec) , antiderivative size = 256, normalized size of antiderivative = 0.78, number of steps used = 15, number of rules used = 14,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$ , Rules used = {6310, 6299, 6308, 6354, 6302, 5971, 27, 3042, 26, 3789, 2611, 2633, 2634, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \operatorname{arccosh}(ax)^{5/2} \sqrt{c - a^2cx^2} dx$$

$$\downarrow \text{6310}$$

$$-\frac{5a\sqrt{c - a^2cx^2} \int x \operatorname{arccosh}(ax)^{3/2} dx}{4\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{\sqrt{c - a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{ax - 1}\sqrt{ax + 1}} dx}{2\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2} x \operatorname{arccosh}(ax)^{5/2} \sqrt{c - a^2cx^2}$$

$$\downarrow \text{6299}$$

$$\begin{aligned}
& \frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\sqrt{c-a^2cx^2} \int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{6308} \\
& \frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \int \frac{x^2 \sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx \right)}{4\sqrt{ax-1}\sqrt{ax+1}} - \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \\
& \quad \frac{1}{2}x \operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{6354} \\
& \frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} - \frac{\int \frac{x}{\sqrt{\operatorname{arccosh}(ax)}} dx}{4a} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}{2a^2} \right) \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{6302} \\
& \frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( -\frac{\int \frac{ax\sqrt{\frac{ax-1}{ax+1}} \operatorname{arccosh}(ax)}{\sqrt{\operatorname{arccosh}(ax)}}}{4a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}{2a^2} \right) \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{5971} \\
& \frac{5a\sqrt{c-a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( -\frac{\int \frac{\sinh(2\operatorname{arccosh}(ax)) \operatorname{arccosh}(ax)}{2\sqrt{\operatorname{arccosh}(ax)}}}{4a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}{2a^2} \right) \right)}{4\sqrt{ax-1}\sqrt{ax+1}} \\
& \frac{\operatorname{arccosh}(ax)^{7/2} \sqrt{c-a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x \operatorname{arccosh}(ax)^{5/2} \sqrt{c-a^2cx^2} \\
& \quad \downarrow \text{27}
\end{aligned}$$

$$5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( -\frac{\int \frac{\sinh(2\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{a}}{2a^2} \right) \right)$$


---


$$\frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2}$$

↓ 3042

$$5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( -\frac{\int -\frac{i\sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{a}}{2a^2} \right) \right)$$


---


$$\frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2}$$

↓ 26

$$5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i\int \frac{\sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{8a^3} + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{a}}{2a^2} \right) \right)$$


---


$$\frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2}$$

↓ 3789

$$5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i\left(\frac{1}{2}i\int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2}i\int \frac{e^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)\right)}{8a^3} + \frac{\int \frac{\sqrt{a}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{a}}{2a^2} \right) \right)$$


---


$$\frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2}$$

↓ 2611

$$5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i\left(i\int e^{2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - i\int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)}\right)}{8a^3} + \frac{\int \frac{\sqrt{a}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{x\sqrt{ax-1}\sqrt{ax+1}\sqrt{a}}{2a^2} \right) \right)$$


---


$$\frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax - 1}\sqrt{ax + 1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2}$$

↓ 2633



$$\begin{aligned}
 & 5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{8a^3} \right) + \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax}}}{4\sqrt{ax-1}\sqrt{ax+1}} \right. \\
 & \left. \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2} \right. \\
 & \quad \downarrow \text{2634} \\
 & 5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{\int \frac{\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{ax-1}\sqrt{ax+1}} dx}{2a^2} + \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) \right)}{8a^3} \right) \right. \\
 & \left. \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2} \right. \\
 & \quad \downarrow \text{6308} \\
 & - \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2} - \\
 & 5a\sqrt{c - a^2cx^2} \left( \frac{1}{2}x^2 \operatorname{arccosh}(ax)^{3/2} - \frac{3}{4}a \left( \frac{i \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) \right)}{8a^3} \right) + \frac{\operatorname{arccosh}(ax)}{3a^3} \right) \\
 & \quad \downarrow \\
 & \frac{\operatorname{arccosh}(ax)^{7/2}\sqrt{c - a^2cx^2}}{7a\sqrt{ax-1}\sqrt{ax+1}} + \frac{1}{2}x\operatorname{arccosh}(ax)^{5/2}\sqrt{c - a^2cx^2} - \frac{\operatorname{arccosh}(ax)}{3a^3}
 \end{aligned}$$

input `Int[Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(5/2),x]`

output `(x*Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(5/2))/2 - (Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(7/2))/(7*a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]) - (5*a*Sqrt[c - a^2*c*x^2]*((x^2*ArcCosh[a*x]^(3/2))/2 - (3*a*((x*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*Sqrt[ArcCosh[a*x]])/(2*a^2) + ArcCosh[a*x]^(3/2)/(3*a^3) + ((I/8)*((-1/2*I)*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]]) + (I/2)*Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]]))/a^3))/4)/(4*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`

## 3.389.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2611 `Int[(F_)^((g_)*((e_) + (f_)*(x_)))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] := Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`
- rule 2633 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2)), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`
- rule 2634 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2)), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3789 `Int[((c_) + (d_)*(x_)^m)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]`
- rule 5971 `Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_)^m)*Sinh[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6299  $\text{Int}[(a + \text{ArcCosh}[c \cdot x] \cdot b)^n \cdot x^m, x\_Symbol] \rightarrow \text{Simp}[x^{m+1} \cdot ((a + b \cdot \text{ArcCosh}[c \cdot x])^n / (m+1)), x] - \text{Simp}[b \cdot c \cdot n / (m+1) \cdot \text{Int}[x^{m+1} \cdot ((a + b \cdot \text{ArcCosh}[c \cdot x])^{n-1} / (\text{Sqrt}[1 + c \cdot x] \cdot \text{Sqrt}[-1 + c \cdot x])), x], x] /;$   $\text{FreeQ}\{a, b, c\}, x\} \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{GtQ}[n, 0]$

rule 6302  $\text{Int}[(a + \text{ArcCosh}[c \cdot x] \cdot b)^n \cdot x^m, x\_Symbol] \rightarrow \text{Simp}[1 / (b \cdot c^{m+1}) \cdot \text{Subst}[\text{Int}[x^n \cdot \text{Cosh}[-a/b + x/b]^m \cdot \text{Sinh}[-a/b + x/b], x], x, a + b \cdot \text{ArcCosh}[c \cdot x]], x] /;$   $\text{FreeQ}\{a, b, c, n\}, x\} \ \&\& \ \text{IGtQ}[m, 0]$

rule 6308  $\text{Int}[(a + \text{ArcCosh}[c \cdot x] \cdot b)^n / (\text{Sqrt}[(d_1 + e_1 \cdot x)] \cdot \text{Sqrt}[(d_2 + e_2 \cdot x)]), x\_Symbol] \rightarrow \text{Simp}[(1 / (b \cdot c \cdot (n+1))) \cdot \text{Simp}[\text{Sqrt}[1 + c \cdot x] / \text{Sqrt}[d_1 + e_1 \cdot x]] \cdot \text{Simp}[\text{Sqrt}[-1 + c \cdot x] / \text{Sqrt}[d_2 + e_2 \cdot x]] \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^{n+1}, x] /;$   $\text{FreeQ}\{a, b, c, d_1, e_1, d_2, e_2, n\}, x\} \ \&\& \ \text{EqQ}[e_1, c \cdot d_1] \ \&\& \ \text{EqQ}[e_2, (-c) \cdot d_2] \ \&\& \ \text{NeQ}[n, -1]$

rule 6310  $\text{Int}[(a + \text{ArcCosh}[c \cdot x] \cdot b)^n \cdot \text{Sqrt}[d + e \cdot x^2], x\_Symbol] \rightarrow \text{Simp}[x \cdot \text{Sqrt}[d + e \cdot x^2] \cdot ((a + b \cdot \text{ArcCosh}[c \cdot x])^{n/2}), x] + (-\text{Simp}[(1/2) \cdot \text{Simp}[\text{Sqrt}[d + e \cdot x^2] / (\text{Sqrt}[1 + c \cdot x] \cdot \text{Sqrt}[-1 + c \cdot x])] \cdot \text{Int}[(a + b \cdot \text{ArcCosh}[c \cdot x])^n / (\text{Sqrt}[1 + c \cdot x] \cdot \text{Sqrt}[-1 + c \cdot x]), x], x] - \text{Simp}[b \cdot c \cdot (n/2) \cdot \text{Simp}[\text{Sqrt}[d + e \cdot x^2] / (\text{Sqrt}[1 + c \cdot x] \cdot \text{Sqrt}[-1 + c \cdot x])] \cdot \text{Int}[x \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^{n-1}, x], x]) /;$   $\text{FreeQ}\{a, b, c, d, e\}, x\} \ \&\& \ \text{EqQ}[c^2 \cdot d + e, 0] \ \&\& \ \text{GtQ}[n, 0]$

rule 6354  $\text{Int}[(a + \text{ArcCosh}[c \cdot x] \cdot b)^n \cdot (f \cdot x)^m \cdot ((d_1 + e_1 \cdot x)^p \cdot (d_2 + e_2 \cdot x)^p), x\_Symbol] \rightarrow \text{Simp}[f \cdot (f \cdot x)^{m-1} \cdot (d_1 + e_1 \cdot x)^{p+1} \cdot (d_2 + e_2 \cdot x)^{p+1} \cdot ((a + b \cdot \text{ArcCosh}[c \cdot x])^n / (e_1 \cdot e_2 \cdot (m + 2 \cdot p + 1))), x] + (\text{Simp}[f^2 \cdot (m-1) / (c^2 \cdot (m + 2 \cdot p + 1))] \cdot \text{Int}[(f \cdot x)^{m-2} \cdot (d_1 + e_1 \cdot x)^p \cdot (d_2 + e_2 \cdot x)^p \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^n, x], x] - \text{Simp}[b \cdot f \cdot (n / (c \cdot (m + 2 \cdot p + 1))) \cdot \text{Simp}[(d_1 + e_1 \cdot x)^p / (1 + c \cdot x)^p] \cdot \text{Simp}[(d_2 + e_2 \cdot x)^p / (-1 + c \cdot x)^p] \cdot \text{Int}[(f \cdot x)^{m-1} \cdot (1 + c \cdot x)^{p+1/2} \cdot (-1 + c \cdot x)^{p+1/2} \cdot (a + b \cdot \text{ArcCosh}[c \cdot x])^{n-1}, x], x]) /;$   $\text{FreeQ}\{a, b, c, d_1, e_1, d_2, e_2, f, p\}, x\} \ \&\& \ \text{EqQ}[e_1, c \cdot d_1] \ \&\& \ \text{EqQ}[e_2, (-c) \cdot d_2] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{IGtQ}[m, 1] \ \&\& \ \text{NeQ}[m + 2 \cdot p + 1, 0]$

**3.389.4 Maple [F]**

$$\int \sqrt{-a^2 c x^2 + c} \operatorname{arccosh}(ax)^{\frac{5}{2}} dx$$

input `int((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(5/2),x)`

output `int((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(5/2),x)`

**3.389.5 Fricas [F(-2)]**

Exception generated.

$$\int \sqrt{c - a^2 c x^2} \operatorname{arccosh}(ax)^{5/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(5/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.389.6 Sympy [F(-1)]**

Timed out.

$$\int \sqrt{c - a^2 c x^2} \operatorname{arccosh}(ax)^{5/2} dx = \text{Timed out}$$

input `integrate((-a**2*c*x**2+c)**(1/2)*acosh(a*x)**(5/2),x)`

output `Timed out`

**3.389.7 Maxima [F]**

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2} dx = \int \sqrt{-a^2 cx^2 + c} \operatorname{arccosh}(ax)^{5/2} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(5/2),x, algorithm="maxima")`

output `integrate(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^(5/2), x)`

**3.389.8 Giac [F(-2)]**

Exception generated.

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(5/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.389.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2} dx = \int \operatorname{acosh}(ax)^{5/2} \sqrt{c - a^2 cx^2} dx$$

input `int(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(1/2),x)`

output `int(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(1/2), x)`

$$3.390 \quad \int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c-a^2cx^2}} dx$$

3.390.1 Optimal result . . . . .	3045
3.390.2 Mathematica [A] (verified) . . . . .	3045
3.390.3 Rubi [A] (verified) . . . . .	3046
3.390.4 Maple [A] (verified) . . . . .	3046
3.390.5 Fracas [F(-2)] . . . . .	3047
3.390.6 Sympy [F(-1)] . . . . .	3047
3.390.7 Maxima [F] . . . . .	3047
3.390.8 Giac [F] . . . . .	3048
3.390.9 Mupad [F(-1)] . . . . .	3048

### 3.390.1 Optimal result

Integrand size = 24, antiderivative size = 48

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c-a^2cx^2}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^{7/2}}{7a\sqrt{c-a^2cx^2}}$$

output  $2/7*\operatorname{arccosh}(a*x)^{(7/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/a/(-a^2*c*x^2+c)^{(1/2)}$

### 3.390.2 Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c-a^2cx^2}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\operatorname{arccosh}(ax)^{7/2}}{7a\sqrt{c-a^2cx^2}}$$

input `Integrate[ArcCosh[a*x]^(5/2)/Sqrt[c - a^2*c*x^2],x]`

output  $(2*\operatorname{Sqrt}[-1 + a*x]*\operatorname{Sqrt}[1 + a*x]*\operatorname{ArcCosh}[a*x]^{(7/2)})/(7*a*\operatorname{Sqrt}[c - a^2*c*x^2])$

**3.390.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c - a^2 cx^2}} dx$$

↓ 6307

$$\frac{2\sqrt{ax - 1}\sqrt{ax + 1}\operatorname{arccosh}(ax)^{7/2}}{7a\sqrt{c - a^2 cx^2}}$$

input `Int[ArcCosh[a*x]^(5/2)/Sqrt[c - a^2*c*x^2],x]`

output `(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*ArcCosh[a*x]^(7/2))/(7*a*Sqrt[c - a^2*c*x^2])`

**3.390.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.390.4 Maple [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{2 \operatorname{arccosh}(ax)^{\frac{7}{2}} \sqrt{ax-1} \sqrt{ax+1}}{7 \sqrt{-c(ax-1)(ax+1)} a}$	41

input `int(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)`

output  $2/7*\operatorname{arccosh}(a*x)^{(7/2)/(-c*(a*x-1)*(a*x+1))^{(1/2)*(a*x-1)^{(1/2)*(a*x+1)^{(1/2)/a}}$

### 3.390.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c - a^2cx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.390.6 Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c - a^2cx^2}} dx = \text{Timed out}$$

input `integrate(acosh(a*x)**(5/2)/(-a**2*c*x**2+c)**(1/2),x)`

output `Timed out`

### 3.390.7 Maxima [F]

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{arcosh}(ax)^{5/2}}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(a*x)^(5/2)/sqrt(-a^2*c*x^2 + c), x)`

---

3.390.  $\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c - a^2cx^2}} dx$



**3.390.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{arcosh}(ax)^{5/2}}{\sqrt{-a^2cx^2 + c}} dx$$

input `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(a*x)^(5/2)/sqrt(-a^2*c*x^2 + c), x)`

**3.390.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{\sqrt{c - a^2cx^2}} dx = \int \frac{\operatorname{acosh}(ax)^{5/2}}{\sqrt{c - a^2cx^2}} dx$$

input `int(acosh(a*x)^(5/2)/(c - a^2*c*x^2)^(1/2),x)`

output `int(acosh(a*x)^(5/2)/(c - a^2*c*x^2)^(1/2), x)`

**3.391**  $\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c-a^2cx^2)^{3/2}} dx$

3.391.1 Optimal result . . . . .	3049
3.391.2 Mathematica [N/A] . . . . .	3049
3.391.3 Rubi [N/A] . . . . .	3050
3.391.4 Maple [N/A] (verified) . . . . .	3051
3.391.5 Fricas [F(-2)] . . . . .	3051
3.391.6 Sympy [F(-1)] . . . . .	3051
3.391.7 Maxima [N/A] . . . . .	3052
3.391.8 Giac [N/A] . . . . .	3052
3.391.9 Mupad [N/A] . . . . .	3052

**3.391.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c-a^2cx^2)^{3/2}} dx = \frac{x\operatorname{arccosh}(ax)^{5/2}}{c\sqrt{c-a^2cx^2}} + \frac{5a\sqrt{-1+ax}\sqrt{1+ax}\operatorname{Int}\left(\frac{x\operatorname{arccosh}(ax)^{3/2}}{1-a^2x^2}, x\right)}{2c\sqrt{c-a^2cx^2}}$$

output `x*arccosh(a*x)^(5/2)/c/(-a^2*c*x^2+c)^(1/2)+5/2*a*(a*x-1)^(1/2)*(a*x+1)^(1/2)*Unintegrable(x*arccosh(a*x)^(3/2)/(-a^2*x^2+1),x)/c/(-a^2*c*x^2+c)^(1/2)`

**3.391.2 Mathematica [N/A]**

Not integrable

Time = 4.78 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c-a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c-a^2cx^2)^{3/2}} dx$$

input `Integrate[ArcCosh[a*x]^(5/2)/(c - a^2*c*x^2)^(3/2), x]`

output `Integrate[ArcCosh[a*x]^(5/2)/(c - a^2*c*x^2)^(3/2), x]`

**3.391.3 Rubi [N/A]**

Not integrable

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6314, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c - a^2cx^2)^{3/2}} dx$$

↓ 6314

$$\frac{5a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^{3/2}}{1-a^2x^2} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)^{5/2}}{c\sqrt{c-a^2cx^2}}$$

↓ 6375

$$\frac{5a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x \operatorname{arccosh}(ax)^{3/2}}{1-a^2x^2} dx}{2c\sqrt{c-a^2cx^2}} + \frac{x \operatorname{arccosh}(ax)^{5/2}}{c\sqrt{c-a^2cx^2}}$$

input `Int[ArcCosh[a*x]^(5/2)/(c - a^2*c*x^2)^(3/2),x]`

output `$Aborted`

**3.391.3.1 Defintions of rubi rules used**

rule 6314 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.391.4 Maple [N/A] (verified)**

Not integrable

Time = 1.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arccosh}(ax)^{\frac{5}{2}}}{(-a^2cx^2+c)^{\frac{3}{2}}} dx$$

input `int(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(3/2),x)`output `int(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(3/2),x)`**3.391.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c-a^2cx^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`**3.391.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c-a^2cx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate(acosch(a*x)**(5/2)/(-a**2*c*x**2+c)**(3/2),x)`output `Timed out`

**3.391.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}(ax)^{5/2}}{(-a^2cx^2 + c)^{3/2}} dx$$

input `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`output `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2 + c)^(3/2), x)`**3.391.8 Giac [N/A]**

Not integrable

Time = 3.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}(ax)^{5/2}}{(-a^2cx^2 + c)^{3/2}} dx$$

input `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2+c)^(3/2),x, algorithm="giac")`output `integrate(arccosh(a*x)^(5/2)/(-a^2*c*x^2 + c)^(3/2), x)`**3.391.9 Mupad [N/A]**

Not integrable

Time = 2.71 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c - a^2cx^2)^{3/2}} dx = \int \frac{\operatorname{acosh}(ax)^{5/2}}{(c - a^2cx^2)^{3/2}} dx$$

input `int(acosh(a*x)^(5/2)/(c - a^2*c*x^2)^(3/2),x)`output `int(acosh(a*x)^(5/2)/(c - a^2*c*x^2)^(3/2), x)`

---

3.391.  $\int \frac{\operatorname{arccosh}(ax)^{5/2}}{(c - a^2cx^2)^{3/2}} dx$

### 3.392 $\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

3.392.1 Optimal result	3053
3.392.2 Mathematica [A] (warning: unable to verify)	3054
3.392.3 Rubi [C] (verified)	3054
3.392.4 Maple [F]	3061
3.392.5 Fricas [F(-2)]	3061
3.392.6 Sympy [F]	3062
3.392.7 Maxima [F]	3062
3.392.8 Giac [F]	3062
3.392.9 Mupad [F(-1)]	3063

#### 3.392.1 Optimal result

Integrand size = 24, antiderivative size = 368

$$\begin{aligned} \int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx &= \frac{3}{8} a^2 x \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\ &+ \frac{1}{4} x (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{a^3 \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{4 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} \\ &- \frac{a^3 \sqrt{\pi} \sqrt{a^2 - x^2} \operatorname{erf}\left(2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{256 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} + \frac{a^3 \sqrt{\frac{\pi}{2}} \sqrt{a^2 - x^2} \operatorname{erf}\left(\sqrt{2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{16 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} \\ &+ \frac{a^3 \sqrt{\pi} \sqrt{a^2 - x^2} \operatorname{erfi}\left(2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{256 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} - \frac{a^3 \sqrt{\frac{\pi}{2}} \sqrt{a^2 - x^2} \operatorname{erfi}\left(\sqrt{2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{16 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} \end{aligned}$$

output  $-1/4*a^3*\operatorname{arccosh}(x/a)^{(3/2)}*(a^2-x^2)^{(1/2)}/(-1+x/a)^{(1/2)}/(1+x/a)^{(1/2)}+1/32*a^3*\operatorname{erf}(2^{(1/2)}*\operatorname{arccosh}(x/a)^{(1/2)})*2^{(1/2)}*Pi^{(1/2)}*(a^2-x^2)^{(1/2)}/(-1+x/a)^{(1/2)}/(1+x/a)^{(1/2)}-1/32*a^3*\operatorname{erfi}(2^{(1/2)}*\operatorname{arccosh}(x/a)^{(1/2)})*2^{(1/2)}*Pi^{(1/2)}*(a^2-x^2)^{(1/2)}/(-1+x/a)^{(1/2)}/(1+x/a)^{(1/2)}-1/256*a^3*\operatorname{erf}(2*\operatorname{arccosh}(x/a)^{(1/2)})*Pi^{(1/2)}*(a^2-x^2)^{(1/2)}/(-1+x/a)^{(1/2)}/(1+x/a)^{(1/2)}+1/256*a^3*\operatorname{erfi}(2*\operatorname{arccosh}(x/a)^{(1/2)})*Pi^{(1/2)}*(a^2-x^2)^{(1/2)}/(-1+x/a)^{(1/2)}/(1+x/a)^{(1/2)}+1/4*x*(a^2-x^2)^{(3/2)}*\operatorname{arccosh}(x/a)^{(1/2)}+3/8*a^2*x*(a^2-x^2)^{(1/2)}*\operatorname{arccosh}(x/a)^{(1/2)}$

### 3.392.2 Mathematica [A] (warning: unable to verify)

Time = 0.35 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.45

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx =$$

$$\frac{a^4 \sqrt{a^2 - x^2} \left( -\sqrt{-\operatorname{arccosh}\left(\frac{x}{a}\right)} \Gamma\left(\frac{3}{2}, -4\operatorname{arccosh}\left(\frac{x}{a}\right)\right) + 8\sqrt{2} \sqrt{-\operatorname{arccosh}\left(\frac{x}{a}\right)} \Gamma\left(\frac{3}{2}, -2\operatorname{arccosh}\left(\frac{x}{a}\right)\right) + \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{128 \sqrt{\frac{-a+x}{a+x}} (a+x) \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}$$

input `Integrate[(a^2 - x^2)^(3/2)*Sqrt[ArcCosh[x/a]],x]`

output `-1/128*(a^4*Sqrt[a^2 - x^2]*(-(Sqrt[-ArcCosh[x/a]]*Gamma[3/2, -4*ArcCosh[x/a]]) + 8*Sqrt[2]*Sqrt[-ArcCosh[x/a]]*Gamma[3/2, -2*ArcCosh[x/a]] + Sqrt[ArcCosh[x/a]]*(32*ArcCosh[x/a]^(3/2) + 8*Sqrt[2]*Gamma[3/2, 2*ArcCosh[x/a]] - Gamma[3/2, 4*ArcCosh[x/a]])))/(Sqrt[(-a + x)/(a + x)]*(a + x)*Sqrt[ArcCosh[x/a]])`

### 3.392.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 3.28 (sec) , antiderivative size = 369, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 18,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$ , Rules used = {6312, 25, 27, 6310, 6302, 5971, 27, 3042, 26, 3789, 2611, 2633, 2634, 6308, 6327, 6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$$

$$\downarrow \text{6312}$$

$$\frac{3}{4}a^2 \int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx + \frac{a\sqrt{a^2 - x^2} \int -\frac{(a-x)x(a+x)}{a^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8\sqrt{\frac{x}{a} - 1} \sqrt{\frac{x}{a} + 1}} +$$

$$\frac{1}{4}x(a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}$$

$$\downarrow \text{25}$$

---

3.392.  $\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

$$\begin{aligned}
& \frac{3}{4}a^2 \int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx - \frac{a\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{a^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \\
& \qquad \qquad \qquad \frac{1}{4}x(a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{3}{4}a^2 \int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx - \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \frac{1}{4}x(a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \qquad \qquad \qquad \downarrow 6310 \\
& \frac{3}{4}a^2 \left( -\frac{\sqrt{a^2 - x^2} \int \frac{x}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{4a\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} dx}{2\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) - \\
& \qquad \qquad \qquad \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \frac{1}{4}x(a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \qquad \qquad \qquad \downarrow 6302 \\
& \frac{3}{4}a^2 \left( -\frac{a\sqrt{a^2 - x^2} \int \frac{x\sqrt{\frac{x-1}{x+1}}\left(\frac{x}{a}+1\right)}{a\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right)}{4\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} dx}{2\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) - \\
& \qquad \qquad \qquad \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \frac{1}{4}x(a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \qquad \qquad \qquad \downarrow 5971 \\
& \qquad \qquad \qquad \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \\
& \frac{3}{4}a^2 \left( -\frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} dx}{2\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} - \frac{a\sqrt{a^2 - x^2} \int \frac{\sinh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right)}{4\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) + \\
& \qquad \qquad \qquad \frac{1}{4}x(a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \qquad \qquad \qquad \downarrow 27
\end{aligned}$$

---

3.392.  $\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$



$$\begin{aligned}
& \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
\frac{3}{4}a^2 & \left( -\frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2} \int \frac{\sinh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} d\operatorname{arccosh}\left(\frac{x}{a}\right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) \\
& \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
\frac{3}{4}a^2 & \left( -\frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2} \int -\frac{i \sin\left(2i\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} d\operatorname{arccosh}\left(\frac{x}{a}\right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) \\
& \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \quad \downarrow \text{26} \\
& \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
\frac{3}{4}a^2 & \left( -\frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{ia\sqrt{a^2 - x^2} \int \frac{\sin\left(2i\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} d\operatorname{arccosh}\left(\frac{x}{a}\right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) \\
& \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \quad \downarrow \text{3789} \\
\frac{3}{4}a^2 & \left( \frac{ia\sqrt{a^2 - x^2} \left( \frac{1}{2}i \int \frac{e^{2\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} d\operatorname{arccosh}\left(\frac{x}{a}\right) - \frac{1}{2}i \int \frac{e^{-2\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} d\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \right) \\
& \frac{\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}
\end{aligned}$$

---

3.392.  $\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

↓ 2611

$$\frac{3}{4}a^2 \left( \frac{ia\sqrt{a^2-x^2} \left( i \int e^{2\operatorname{arccosh}\left(\frac{x}{a}\right)} d\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - i \int e^{-2\operatorname{arccosh}\left(\frac{x}{a}\right)} d\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \right) \\ - \frac{\sqrt{a^2-x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}$$

↓ 2633

$$\frac{3}{4}a^2 \left( \frac{ia\sqrt{a^2-x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - i \int e^{-2\operatorname{arccosh}\left(\frac{x}{a}\right)} d\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \right) \\ - \frac{\sqrt{a^2-x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}$$

↓ 2634

$$\frac{3}{4}a^2 \left( - \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{ia\sqrt{a^2-x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \right) \\ - \frac{\sqrt{a^2-x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}$$

↓ 6308

$$- \frac{\sqrt{a^2-x^2} \int \frac{(a-x)x(a+x)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\ \frac{3}{4}a^2 \left( \frac{ia\sqrt{a^2-x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \right) \\ + \frac{1}{4}x(a^2-x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}$$

↓ 6327

---

3.392.  $\int (a^2-x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

$$\begin{aligned}
& \frac{\sqrt{a^2 - x^2} \int \frac{x(a^2 - x^2)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{8a\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \\
\frac{3}{4}a^2 & \left( \frac{ia\sqrt{a^2 - x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \right. \\
& \left. \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) \\
& \quad \downarrow \text{6367} \\
& \frac{a^3\sqrt{a^2 - x^2} \int \frac{x\left(\frac{x-1}{\frac{x}{a}+1}\right)^{3/2}\left(\frac{x+1}{\frac{x}{a}}\right)^3 d\operatorname{arccosh}\left(\frac{x}{a}\right)}{a\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \\
\frac{3}{4}a^2 & \left( \frac{ia\sqrt{a^2 - x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \right. \\
& \left. \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) \\
& \quad \downarrow \text{5971} \\
& \frac{a^3\sqrt{a^2 - x^2} \int \left( \frac{\sinh\left(4\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{8\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} - \frac{\sinh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{4\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \right) d\operatorname{arccosh}\left(\frac{x}{a}\right)}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \\
\frac{3}{4}a^2 & \left( \frac{ia\sqrt{a^2 - x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \right. \\
& \left. \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) \\
& \quad \downarrow \text{2009} \\
\frac{3}{4}a^2 & \left( \frac{ia\sqrt{a^2 - x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} + \right. \\
& \left. \frac{1}{4}x(a^2 - x^2)^{3/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} + \right. \\
& \left. \frac{a^3\sqrt{a^2 - x^2} \left( -\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + \frac{1}{8}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{8}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a} - 1}\sqrt{\frac{x}{a} + 1}} \right)
\end{aligned}$$

input `Int[(a^2 - x^2)^(3/2)*Sqrt[ArcCosh[x/a]], x]`

3.392.  $\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

```
output (x*(a^2 - x^2)^(3/2)*Sqrt[ArcCosh[x/a]]/4 + (a^3*Sqrt[a^2 - x^2]*(-1/32*(
Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[x/a]]]) + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh
[x/a]]])/8 + (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[x/a]]])/32 - (Sqrt[Pi/2]*Erfi[S
qrt[2]*Sqrt[ArcCosh[x/a]]])/8))/(8*Sqrt[-1 + x/a]*Sqrt[1 + x/a]) + (3*a^2*
((x*Sqrt[a^2 - x^2]*Sqrt[ArcCosh[x/a]])/2 - (a*Sqrt[a^2 - x^2]*ArcCosh[x/a
]^(3/2)))/(3*Sqrt[-1 + x/a]*Sqrt[1 + x/a]) + ((I/8)*a*Sqrt[a^2 - x^2]*((-1/
2*I)*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[x/a]]] + (I/2)*Sqrt[Pi/2]*Erfi[Sq
rt[2]*Sqrt[ArcCosh[x/a]]]))/(Sqrt[-1 + x/a]*Sqrt[1 + x/a]))/4
```

### 3.392.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 26 Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 2611 Int[(F_)^((g_)*((e_) + (f_)*(x_)))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] :
> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d
*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

```
rule 2633 Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt
[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{
F, a, b, c, d}, x] && PosQ[b]
```

```
rule 2634 Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt
[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; Fr
eeQ[{F, a, b, c, d}, x] && NegQ[b]
```

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3789 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^m, x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`

rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^(n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] + (Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && GtQ[p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_)^(p_.))*((d2_) + (e2_.)*(x_)^(p_.)), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.392.4 Maple [F]

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$$

input `int((a^2-x^2)^(3/2)*arccosh(x/a)^(1/2),x)`

output `int((a^2-x^2)^(3/2)*arccosh(x/a)^(1/2),x)`

### 3.392.5 Fracas [F(-2)]

Exception generated.

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \text{Exception raised: TypeError}$$

input `integrate((a^2-x^2)^(3/2)*arccosh(x/a)^(1/2),x, algorithm="fricas")`

---

3.392.  $\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

### 3.392.6 Sympy [F]

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int (-(-a + x)(a + x))^{\frac{3}{2}} \sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)} dx$$

input `integrate((a**2-x**2)**(3/2)*acosh(x/a)**(1/2),x)`

output `Integral((-(-a + x)*(a + x))**(3/2)*sqrt(acosh(x/a)), x)`

### 3.392.7 Maxima [F]

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int (a^2 - x^2)^{\frac{3}{2}} \sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)} dx$$

input `integrate((a^2-x^2)^(3/2)*arccosh(x/a)^(1/2),x, algorithm="maxima")`

output `integrate((a^2 - x^2)^(3/2)*sqrt(arccosh(x/a)), x)`

### 3.392.8 Giac [F]

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int (a^2 - x^2)^{\frac{3}{2}} \sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)} dx$$

input `integrate((a^2-x^2)^(3/2)*arccosh(x/a)^(1/2),x, algorithm="giac")`

output `integrate((a^2 - x^2)^(3/2)*sqrt(arccosh(x/a)), x)`

**3.392.9 Mupad [F(-1)]**

Timed out.

$$\int (a^2 - x^2)^{3/2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int \sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)} (a^2 - x^2)^{3/2} dx$$

input `int(acosh(x/a)^(1/2)*(a^2 - x^2)^(3/2), x)`output `int(acosh(x/a)^(1/2)*(a^2 - x^2)^(3/2), x)`



### 3.393 $\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

3.393.1 Optimal result . . . . .	3064
3.393.2 Mathematica [A] (verified) . . . . .	3065
3.393.3 Rubi [C] (verified) . . . . .	3065
3.393.4 Maple [F] . . . . .	3069
3.393.5 Fricas [F(-2)] . . . . .	3069
3.393.6 Sympy [F] . . . . .	3070
3.393.7 Maxima [F] . . . . .	3070
3.393.8 Giac [F] . . . . .	3070
3.393.9 Mupad [F(-1)] . . . . .	3071

#### 3.393.1 Optimal result

Integrand size = 24, antiderivative size = 211

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \frac{1}{2} x \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{a \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} + \frac{a \sqrt{\frac{\pi}{2}} \sqrt{a^2 - x^2} \operatorname{erf}\left(\sqrt{2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{16 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} - \frac{a \sqrt{\frac{\pi}{2}} \sqrt{a^2 - x^2} \operatorname{erfi}\left(\sqrt{2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{16 \sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}}$$

output

```
-1/3*a*arccosh(x/a)^(3/2)*(a^2-x^2)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)+1/3
2*a*erf(2^(1/2)*arccosh(x/a)^(1/2))*2^(1/2)*Pi^(1/2)*(a^2-x^2)^(1/2)/(-1+x
/a)^(1/2)/(1+x/a)^(1/2)-1/32*a*erfi(2^(1/2)*arccosh(x/a)^(1/2))*2^(1/2)*Pi
^(1/2)*(a^2-x^2)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)+1/2*x*(a^2-x^2)^(1/2)*
arccosh(x/a)^(1/2)
```

**3.393.2 Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.57

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \frac{a^2 \sqrt{a^2 - x^2} \left( 16 \operatorname{arccosh}\left(\frac{x}{a}\right)^2 + 3\sqrt{2} \sqrt{-\operatorname{arccosh}\left(\frac{x}{a}\right)} \Gamma\left(\frac{3}{2}, -2 \operatorname{arccosh}\left(\frac{x}{a}\right)\right) + 3\sqrt{2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \Gamma\left(\frac{3}{2}, 2 \operatorname{arccosh}\left(\frac{x}{a}\right)\right) \right)}{48 \sqrt{\frac{-a+x}{a+x}} (a+x) \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}$$

input `Integrate[Sqrt[a^2 - x^2]*Sqrt[ArcCosh[x/a]], x]`output `-1/48*(a^2*Sqrt[a^2 - x^2]*(16*ArcCosh[x/a]^2 + 3*Sqrt[2]*Sqrt[-ArcCosh[x/a]]*Gamma[3/2, -2*ArcCosh[x/a]] + 3*Sqrt[2]*Sqrt[ArcCosh[x/a]]*Gamma[3/2, 2*ArcCosh[x/a]]))/(Sqrt[(-a + x)/(a + x)]*(a + x)*Sqrt[ArcCosh[x/a]])`**3.393.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 1.91 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.88, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.458$ , Rules used = {6310, 6302, 5971, 27, 3042, 26, 3789, 2611, 2633, 2634, 6308}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx \\ & \quad \downarrow \text{6310} \\ & -\frac{\sqrt{a^2 - x^2} \int \frac{x}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{4a \sqrt{\frac{x}{a} - 1} \sqrt{\frac{x}{a} + 1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a} - 1} \sqrt{\frac{x}{a} + 1}} dx}{2 \sqrt{\frac{x}{a} - 1} \sqrt{\frac{x}{a} + 1}} + \frac{1}{2} x \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\ & \quad \downarrow \text{6302} \end{aligned}$$

$$\begin{aligned}
& \frac{a\sqrt{a^2-x^2} \int \frac{x\sqrt{\frac{x-1}{x}+1} \left(\frac{x}{a}+1\right)}{a\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \operatorname{darccosh}\left(\frac{x}{a}\right) - \sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{4\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{\frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{2} \\
& \quad \downarrow \text{5971} \\
& \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx - a\sqrt{a^2-x^2} \int \frac{\sinh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \operatorname{darccosh}\left(\frac{x}{a}\right)}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{\frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{2} \\
& \quad \downarrow \text{27} \\
& \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx - a\sqrt{a^2-x^2} \int \frac{\sinh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \operatorname{darccosh}\left(\frac{x}{a}\right)}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{\frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{2} \\
& \quad \downarrow \text{3042} \\
& \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx - a\sqrt{a^2-x^2} \int -\frac{i\sin\left(2i\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \operatorname{darccosh}\left(\frac{x}{a}\right)}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{\frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{2} \\
& \quad \downarrow \text{26} \\
& \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx + ia\sqrt{a^2-x^2} \int \frac{\sin\left(2i\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \operatorname{darccosh}\left(\frac{x}{a}\right)}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{\frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{2} \\
& \quad \downarrow \text{3789} \\
& \frac{ia\sqrt{a^2-x^2} \left( \frac{1}{2}i \int \frac{e^{2\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \operatorname{darccosh}\left(\frac{x}{a}\right) - \frac{1}{2}i \int \frac{e^{-2\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \operatorname{darccosh}\left(\frac{x}{a}\right) \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \\
& \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}
\end{aligned}$$

---

3.393.  $\int \sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$

$$\begin{aligned}
& \downarrow 2611 \\
& \frac{ia\sqrt{a^2-x^2} \left( i \int e^{2\operatorname{arccosh}\left(\frac{x}{a}\right)} d\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - i \int e^{-2\operatorname{arccosh}\left(\frac{x}{a}\right)} d\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \\
& \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \downarrow 2633 \\
& \frac{ia\sqrt{a^2-x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - i \int e^{-2\operatorname{arccosh}\left(\frac{x}{a}\right)} d\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \\
& \frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \downarrow 2634 \\
& -\frac{\sqrt{a^2-x^2} \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{ia\sqrt{a^2-x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \\
& \downarrow 6308 \\
& \frac{ia\sqrt{a^2-x^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \\
& \frac{a\sqrt{a^2-x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}
\end{aligned}$$

input `Int[Sqrt[a^2 - x^2]*Sqrt[ArcCosh[x/a]], x]`

output `(x*Sqrt[a^2 - x^2]*Sqrt[ArcCosh[x/a]])/2 - (a*Sqrt[a^2 - x^2]*ArcCosh[x/a]^(3/2))/(3*Sqrt[-1 + x/a]*Sqrt[1 + x/a]) + ((I/8)*a*Sqrt[a^2 - x^2]*((-1/2)*I)*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[x/a]]] + (I/2)*Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[x/a]]])/(Sqrt[-1 + x/a]*Sqrt[1 + x/a])`

## 3.393.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2611 `Int[(F_)^((g_)*((e_) + (f_)*(x_)))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`
- rule 2633 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`
- rule 2634 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3789 `Int[((c_) + (d_)*(x_)^(m_))*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]`
- rule 5971 `Int[Cosh[(a_) + (b_)*(x_)]^(p_)*((c_) + (d_)*(x_)^(m_))*Sinh[(a_) + (b_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`
- rule 6302 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*(x_)^(m_), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

```
rule 6308 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]
```

```
rule 6310 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]
```

### 3.393.4 Maple [F]

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx$$

```
input int((a^2-x^2)^(1/2)*arccosh(x/a)^(1/2),x)
```

```
output int((a^2-x^2)^(1/2)*arccosh(x/a)^(1/2),x)
```

### 3.393.5 Fracas [F(-2)]

Exception generated.

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \text{Exception raised: TypeError}$$

```
input integrate((a^2-x^2)^(1/2)*arccosh(x/a)^(1/2),x, algorithm="fricas")
```

```
output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)
```

**3.393.6 Sympy [F]**

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int \sqrt{-(-a + x)(a + x)} \sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)} dx$$

input `integrate((a**2-x**2)**(1/2)*acosh(x/a)**(1/2),x)`

output `Integral(sqrt(-(-a + x)*(a + x))*sqrt(acosh(x/a)), x)`

**3.393.7 Maxima [F]**

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int \sqrt{a^2 - x^2} \sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)} dx$$

input `integrate((a^2-x^2)^(1/2)*arccosh(x/a)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(a^2 - x^2)*sqrt(arccosh(x/a)), x)`

**3.393.8 Giac [F]**

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int \sqrt{a^2 - x^2} \sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)} dx$$

input `integrate((a^2-x^2)^(1/2)*arccosh(x/a)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(a^2 - x^2)*sqrt(arccosh(x/a)), x)`

**3.393.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx = \int \sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)} \sqrt{a^2 - x^2} dx$$

input `int(acosh(x/a)^(1/2)*(a^2 - x^2)^(1/2), x)`output `int(acosh(x/a)^(1/2)*(a^2 - x^2)^(1/2), x)`



**3.394**  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2-x^2}} dx$

3.394.1 Optimal result	3072
3.394.2 Mathematica [A] (verified)	3072
3.394.3 Rubi [A] (verified)	3073
3.394.4 Maple [A] (verified)	3073
3.394.5 Fricas [F(-2)]	3074
3.394.6 Sympy [F]	3074
3.394.7 Maxima [F]	3074
3.394.8 Giac [F]	3075
3.394.9 Mupad [F(-1)]	3075

**3.394.1 Optimal result**

Integrand size = 24, antiderivative size = 50

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2-x^2}} dx = \frac{2a\sqrt{-1+\frac{x}{a}}\sqrt{1+\frac{x}{a}}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{a^2-x^2}}$$

output `2/3*a*arccosh(x/a)^(3/2)*(-1+x/a)^(1/2)*(1+x/a)^(1/2)/(a^2-x^2)^(1/2)`

**3.394.2 Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2-x^2}} dx = \frac{2a\sqrt{-1+\frac{x}{a}}\sqrt{1+\frac{x}{a}}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{a^2-x^2}}$$

input `Integrate[Sqrt[ArcCosh[x/a]]/Sqrt[a^2 - x^2],x]`

output `(2*a*Sqrt[-1 + x/a]*Sqrt[1 + x/a]*ArcCosh[x/a]^(3/2))/(3*Sqrt[a^2 - x^2])`

**3.394.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx$$

↓ 6307

$$\frac{2a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{3\sqrt{a^2-x^2}}$$

input `Int[Sqrt[ArcCosh[x/a]]/Sqrt[a^2 - x^2],x]`

output `(2*a*Sqrt[-1 + x/a]*Sqrt[1 + x/a]*ArcCosh[x/a]^(3/2))/(3*Sqrt[a^2 - x^2])`

**3.394.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.394.4 Maple [A] (verified)**

Time = 0.57 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{2 \operatorname{arccosh}\left(\frac{x}{a}\right)^{\frac{3}{2}} \sqrt{-\frac{a-x}{a}} \sqrt{\frac{a+x}{a}} a}{3\sqrt{(a-x)(a+x)}}$	44

input `int(arccosh(x/a)^(1/2)/(a^2-x^2)^(1/2),x,method=_RETURNVERBOSE)`

output  $2/3*\operatorname{arccosh}(x/a)^{(3/2)/((a-x)*(a+x))^{(1/2)*(-(a-x)/a)^{(1/2)*((a+x)/a)^{(1/2)}}*a$

### 3.394.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.394.6 Sympy [F]

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx = \int \frac{\sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)}}{\sqrt{-(-a+x)(a+x)}} dx$$

input `integrate(acosh(x/a)**(1/2)/(a**2-x**2)**(1/2),x)`

output `Integral(sqrt(acosh(x/a))/sqrt(-(-a+x)*(a+x)), x)`

### 3.394.7 Maxima [F]

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx = \int \frac{\sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx$$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(arccosh(x/a))/sqrt(a^2 - x^2), x)`

---

3.394.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx$

**3.394.8 Giac [F]**

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx = \int \frac{\sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx$$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(arccosh(x/a))/sqrt(a^2 - x^2), x)`

**3.394.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx = \int \frac{\sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)}}{\sqrt{a^2 - x^2}} dx$$

input `int(acosh(x/a)^(1/2)/(a^2 - x^2)^(1/2),x)`

output `int(acosh(x/a)^(1/2)/(a^2 - x^2)^(1/2), x)`

**3.395**  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{3/2}} dx$

3.395.1 Optimal result . . . . .	3076
3.395.2 Mathematica [N/A] . . . . .	3076
3.395.3 Rubi [N/A] . . . . .	3077
3.395.4 Maple [N/A] (verified) . . . . .	3078
3.395.5 Fricas [F(-2)] . . . . .	3078
3.395.6 Sympy [N/A] . . . . .	3079
3.395.7 Maxima [N/A] . . . . .	3079
3.395.8 Giac [N/A] . . . . .	3079
3.395.9 Mupad [N/A] . . . . .	3080

**3.395.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{3/2}} dx = \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}} + \frac{\sqrt{-1+\frac{x}{a}}\sqrt{1+\frac{x}{a}}\operatorname{Int}\left(\frac{x}{(1-\frac{x^2}{a^2})\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}, x\right)}{2a^3\sqrt{a^2-x^2}}$$

```
output x*arccosh(x/a)^(1/2)/a^2/(a^2-x^2)^(1/2)+1/2*(-1+x/a)^(1/2)*(1+x/a)^(1/2)*
Unintegrable(x/(1-x^2/a^2)/arccosh(x/a)^(1/2),x)/a^3/(a^2-x^2)^(1/2)
```

**3.395.2 Mathematica [N/A]**

Not integrable

Time = 7.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{3/2}} dx$$

```
input Integrate[Sqrt[ArcCosh[x/a]]/(a^2 - x^2)^(3/2), x]
```

```
output Integrate[Sqrt[ArcCosh[x/a]]/(a^2 - x^2)^(3/2), x]
```

3.395.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{3/2}} dx$

**3.395.3 Rubi [N/A]**

Not integrable

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6314, 27, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx \\
 & \quad \downarrow \text{6314} \\
 & \frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{a^2 x}{(a^2-x^2)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{2a^3\sqrt{a^2-x^2}} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}} \\
 & \quad \downarrow \text{27} \\
 & \frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a^2-x^2)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{2a\sqrt{a^2-x^2}} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}} \\
 & \quad \downarrow \text{6375} \\
 & \frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a^2-x^2)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{2a\sqrt{a^2-x^2}} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}}
 \end{aligned}$$

input `Int[Sqrt[ArcCosh[x/a]]/(a^2 - x^2)^(3/2),x]`

output `$Aborted`

**3.395.3.1 Defintions of rubi rules used**

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

---

3.395.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{3/2}} dx$

```
rule 6314 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2),
x_Symbol] :> Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp
[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a
+ b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e},
x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]
```

```
rule 6375 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_)^(m_.))*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*Ar
cCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]
```

### 3.395.4 Maple [N/A] (verified)

Not integrable

Time = 1.85 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{\frac{3}{2}}} dx$$

```
input int(arccosh(x/a)^(1/2)/(a^2-x^2)^(3/2),x)
```

```
output int(arccosh(x/a)^(1/2)/(a^2-x^2)^(3/2),x)
```

### 3.395.5 Fracas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(3/2),x, algorithm="fracas")
```

```
output Exception raised: TypeError >> Error detected within library code: inte
grate: implementation incomplete (constant residues)
```

---

3.395.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx$

**3.395.6 Sympy [N/A]**

Not integrable

Time = 4.75 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)}}{(-(-a + x)(a + x))^{\frac{3}{2}}} dx$$

input `integrate(acosh(x/a)**(1/2)/(a**2-x**2)**(3/2),x)`output `Integral(sqrt(acosh(x/a))/(-(-a + x)*(a + x))**(3/2), x)`**3.395.7 Maxima [N/A]**

Not integrable

Time = 0.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(3/2),x, algorithm="maxima")`output `integrate(sqrt(arccosh(x/a))/(a^2 - x^2)^(3/2), x)`**3.395.8 Giac [N/A]**

Not integrable

Time = 1.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(3/2),x, algorithm="giac")`output `integrate(sqrt(arccosh(x/a))/(a^2 - x^2)^(3/2), x)`

---

3.395.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx$



**3.395.9 Mupad [N/A]**

Not integrable

Time = 3.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx$$

input `int(acosh(x/a)^(1/2)/(a^2 - x^2)^(3/2), x)`output `int(acosh(x/a)^(1/2)/(a^2 - x^2)^(3/2), x)`

**3.396**  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{5/2}} dx$

3.396.1 Optimal result . . . . . 3081  
 3.396.2 Mathematica [N/A] . . . . . 3082  
 3.396.3 Rubi [N/A] . . . . . 3082  
 3.396.4 Maple [N/A] (verified) . . . . . 3084  
 3.396.5 Fricas [F(-2)] . . . . . 3085  
 3.396.6 Sympy [N/A] . . . . . 3085  
 3.396.7 Maxima [N/A] . . . . . 3085  
 3.396.8 Giac [N/A] . . . . . 3086  
 3.396.9 Mupad [N/A] . . . . . 3086

**3.396.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{5/2}} dx = \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^2(a^2-x^2)^{3/2}} + \frac{2x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^4\sqrt{a^2-x^2}}$$

$$+ \frac{\sqrt{-1+\frac{x}{a}}\sqrt{1+\frac{x}{a}}\operatorname{Int}\left(\frac{x}{\left(1-\frac{x^2}{a^2}\right)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}, x\right)}{3a^5\sqrt{a^2-x^2}}$$

$$+ \frac{\sqrt{-1+\frac{x}{a}}\sqrt{1+\frac{x}{a}}\operatorname{Int}\left(\frac{x}{\left(-1+\frac{x^2}{a^2}\right)^2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}, x\right)}{6a^5\sqrt{a^2-x^2}}$$

```
output 1/3*x*arccosh(x/a)^(1/2)/a^2/(a^2-x^2)^(3/2)+2/3*x*arccosh(x/a)^(1/2)/a^4/
(a^2-x^2)^(1/2)+1/3*(-1+x/a)^(1/2)*(1+x/a)^(1/2)*Unintegrable(x/(1-x^2/a^2
)/arccosh(x/a)^(1/2),x)/a^5/(a^2-x^2)^(1/2)+1/6*(-1+x/a)^(1/2)*(1+x/a)^(1/
2)*Unintegrable(x/(-1+x^2/a^2)^2/arccosh(x/a)^(1/2),x)/a^5/(a^2-x^2)^(1/2)
```

**3.396.2 Mathematica [N/A]**

Not integrable

Time = 3.54 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx$$

input `Integrate[Sqrt[ArcCosh[x/a]]/(a^2 - x^2)^(5/2), x]`output `Integrate[Sqrt[ArcCosh[x/a]]/(a^2 - x^2)^(5/2), x]`**3.396.3 Rubi [N/A]**

Not integrable

Time = 0.95 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6316, 27, 6314, 27, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx \\ & \quad \downarrow \text{6316} \\ & \frac{2 \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx}{3a^2} + \frac{\sqrt{\frac{x}{a} - 1} \sqrt{\frac{x}{a} + 1} \int \frac{a^4 x}{(a-x)^2 (a+x)^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{6a^5 \sqrt{a^2 - x^2}} + \frac{x \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^2 (a^2 - x^2)^{3/2}} \\ & \quad \downarrow \text{27} \\ & \frac{\sqrt{\frac{x}{a} - 1} \sqrt{\frac{x}{a} + 1} \int \frac{x}{(a-x)^2 (a+x)^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{6a \sqrt{a^2 - x^2}} + \frac{2 \int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{3/2}} dx}{3a^2} + \frac{x \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^2 (a^2 - x^2)^{3/2}} \\ & \quad \downarrow \text{6314} \end{aligned}$$

---

3.396.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx$

$$\begin{aligned}
& \frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a-x)^2(a+x)^2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{6a\sqrt{a^2-x^2}} + \\
& \frac{2\left(\frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{a^2x}{(a^2-x^2)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{2a^3\sqrt{a^2-x^2}} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}}\right)}{3a^2} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^2(a^2-x^2)^{3/2}} \\
& \quad \downarrow 27 \\
& \frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a-x)^2(a+x)^2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{6a\sqrt{a^2-x^2}} + \\
& \frac{2\left(\frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a^2-x^2)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{2a\sqrt{a^2-x^2}} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}}\right)}{3a^2} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^2(a^2-x^2)^{3/2}} \\
& \quad \downarrow 6327 \\
& \frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a^2-x^2)^2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{6a\sqrt{a^2-x^2}} + \\
& \frac{2\left(\frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a^2-x^2)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{2a\sqrt{a^2-x^2}} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}}\right)}{3a^2} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^2(a^2-x^2)^{3/2}} \\
& \quad \downarrow 6375 \\
& \frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a^2-x^2)^2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{6a\sqrt{a^2-x^2}} + \\
& \frac{2\left(\frac{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x}{(a^2-x^2)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{2a\sqrt{a^2-x^2}} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2\sqrt{a^2-x^2}}\right)}{3a^2} + \frac{x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{3a^2(a^2-x^2)^{3/2}}
\end{aligned}$$

input `Int [Sqrt [ArcCosh [x/a]]/(a^2 - x^2)^(5/2), x]`

output `$Aborted`

3.396.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2-x^2)^{5/2}} dx$

## 3.396.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 6314 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)/((d_) + (e_)*(x_)^2)^(3/2), x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`
- rule 6316 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*d*(p + 1))), x] + (Simp[(2*p + 3)/(2*d*(p + 1)) Int[(d + e*x^2)^(p + 1)*(a + b*ArcCosh[c*x])^n, x], x] - Simp[b*c*(n/(2*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[p, -1] && NeQ[p, -3/2]`
- rule 6327 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_))^(p_)*((d2_) + (e2_)*(x_))^(p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6375 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))^(n_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

## 3.396.4 Maple [N/A] (verified)

Not integrable

Time = 1.68 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{\frac{5}{2}}} dx$$

input `int(arccosh(x/a)^(1/2)/(a^2-x^2)^(5/2), x)`

---

3.396.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{\frac{5}{2}}} dx$

output `int(arccosh(x/a)^(1/2)/(a^2-x^2)^(5/2), x)`

### 3.396.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(5/2), x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.396.6 Sympy [N/A]

Not integrable

Time = 97.63 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)}}{(-(-a + x)(a + x))^{\frac{5}{2}}} dx$$

input `integrate(acosh(x/a)**(1/2)/(a**2-x**2)**(5/2), x)`

output `Integral(sqrt(acosh(x/a))/(-(-a + x)*(a + x))**(5/2), x)`

### 3.396.7 Maxima [N/A]

Not integrable

Time = 0.53 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{\frac{5}{2}}} dx$$

---

3.396.  $\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(5/2),x, algorithm="maxima")`

output `integrate(sqrt(arccosh(x/a))/(a^2 - x^2)^(5/2), x)`

### 3.396.8 Giac [N/A]

Not integrable

Time = 1.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{arcosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx$$

input `integrate(arccosh(x/a)^(1/2)/(a^2-x^2)^(5/2),x, algorithm="giac")`

output `integrate(sqrt(arccosh(x/a))/(a^2 - x^2)^(5/2), x)`

### 3.396.9 Mupad [N/A]

Not integrable

Time = 2.76 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx = \int \frac{\sqrt{\operatorname{acosh}\left(\frac{x}{a}\right)}}{(a^2 - x^2)^{5/2}} dx$$

input `int(acosh(x/a)^(1/2)/(a^2 - x^2)^(5/2),x)`

output `int(acosh(x/a)^(1/2)/(a^2 - x^2)^(5/2), x)`

### 3.397 $\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$

3.397.1 Optimal result . . . . .	3087
3.397.2 Mathematica [A] (warning: unable to verify) . . . . .	3088
3.397.3 Rubi [A] (verified) . . . . .	3088
3.397.4 Maple [F] . . . . .	3096
3.397.5 Fricas [F(-2)] . . . . .	3096
3.397.6 Sympy [F(-1)] . . . . .	3096
3.397.7 Maxima [F] . . . . .	3097
3.397.8 Giac [F] . . . . .	3097
3.397.9 Mupad [F(-1)] . . . . .	3097

#### 3.397.1 Optimal result

Integrand size = 24, antiderivative size = 525

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \frac{27a^3\sqrt{a^2 - x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{256\sqrt{-1 + \frac{x}{a}}\sqrt{1 + \frac{x}{a}}} - \frac{9ax^2\sqrt{a^2 - x^2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{32\sqrt{-1 + \frac{x}{a}}\sqrt{1 + \frac{x}{a}}} + \frac{3(a^2 - x^2)^{5/2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{32a\sqrt{-1 + \frac{x}{a}}\sqrt{1 + \frac{x}{a}}} + \frac{3}{8}a^2x\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} + \frac{1}{4}x(a^2 - x^2)^{3/2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} - \frac{3a^3\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{20\sqrt{-1 + \frac{x}{a}}\sqrt{1 + \frac{x}{a}}} - \frac{3a^3\sqrt{\pi}\sqrt{a^2 - x^2}}{2048}$$

output

```
1/4*x*(a^2-x^2)^(3/2)*arccosh(x/a)^(3/2)+3/8*a^2*x*arccosh(x/a)^(3/2)*(a^2-x^2)^(1/2)-3/20*a^3*arccosh(x/a)^(5/2)*(a^2-x^2)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)+3/128*a^3*erf(2^(1/2)*arccosh(x/a)^(1/2))*2^(1/2)*Pi^(1/2)*(a^2-x^2)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)+3/128*a^3*erfi(2^(1/2)*arccosh(x/a)^(1/2))*2^(1/2)*Pi^(1/2)*(a^2-x^2)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)-3/2048*a^3*erf(2*arccosh(x/a)^(1/2))*Pi^(1/2)*(a^2-x^2)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)-3/2048*a^3*erfi(2*arccosh(x/a)^(1/2))*Pi^(1/2)*(a^2-x^2)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)+3/32*(a^2-x^2)^(5/2)*arccosh(x/a)^(1/2)/a/(-1+x/a)^(1/2)/(1+x/a)^(1/2)+27/256*a^3*(a^2-x^2)^(1/2)*arccosh(x/a)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)-9/32*a*x^2*(a^2-x^2)^(1/2)*arccosh(x/a)^(1/2)/(-1+x/a)^(1/2)/(1+x/a)^(1/2)
```



**3.397.2 Mathematica [A] (warning: unable to verify)**

Time = 0.59 (sec) , antiderivative size = 219, normalized size of antiderivative = 0.42

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right) dx = \frac{a^4 \sqrt{a^2 - x^2} \left( -384 \operatorname{arccosh}\left(\frac{x}{a}\right)^3 - 480 \operatorname{arccosh}\left(\frac{x}{a}\right) \cosh\left(2 \operatorname{arccosh}\left(\frac{x}{a}\right)\right) + 60 \sqrt{2\pi} \right)}{2560 \sqrt{\frac{-a+x}{a+x}} (a+x) \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}$$

input `Integrate[(a^2 - x^2)^(3/2)*ArcCosh[x/a]^(3/2),x]`

output `(a^4*Sqrt[a^2 - x^2]*(-384*ArcCosh[x/a]^3 - 480*ArcCosh[x/a]*Cosh[2*ArcCosh[x/a]] + 60*Sqrt[2*Pi]*Sqrt[ArcCosh[x/a]]*Erf[Sqrt[2]*Sqrt[ArcCosh[x/a]]] + 60*Sqrt[2*Pi]*Sqrt[ArcCosh[x/a]]*Erfi[Sqrt[2]*Sqrt[ArcCosh[x/a]]] - 5*Sqrt[-ArcCosh[x/a]]*Gamma[5/2, -4*ArcCosh[x/a]] + 5*Sqrt[ArcCosh[x/a]]*Gamma[5/2, 4*ArcCosh[x/a]] + 640*ArcCosh[x/a]^2*Sinh[2*ArcCosh[x/a]]))/(2560*Sqrt[(-a + x)/(a + x)]*(a + x)*Sqrt[ArcCosh[x/a]])`

**3.397.3 Rubi [A] (verified)**

Time = 5.06 (sec) , antiderivative size = 447, normalized size of antiderivative = 0.85, number of steps used = 17, number of rules used = 16,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6312, 25, 27, 6310, 6299, 6308, 6327, 6329, 6322, 3042, 3793, 2009, 6368, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right) dx$$

$$\downarrow \text{6312}$$

$$\frac{3}{4}a^2 \int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right) dx + \frac{3a\sqrt{a^2 - x^2} \int -\frac{(a-x)x(a+x)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2} dx}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)$$

$$\downarrow \text{25}$$

$$\begin{aligned}
& \frac{3}{4}a^2 \int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx - \frac{3a\sqrt{a^2 - x^2} \int \frac{(a-x)x(a+x)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{a^2}}{8\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \qquad \qquad \qquad \frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{3}{4}a^2 \int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx - \frac{3\sqrt{a^2 - x^2} \int (a-x)x(a+x)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \qquad \qquad \qquad \frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\
& \qquad \qquad \qquad \downarrow 6310 \\
& \frac{3}{4}a^2 \left( -\frac{3\sqrt{a^2 - x^2} \int x\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \right) - \\
& \qquad \qquad \qquad \frac{3\sqrt{a^2 - x^2} \int (a-x)x(a+x)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\
& \qquad \qquad \qquad \downarrow 6299 \\
& \frac{3}{4}a^2 \left( -\frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \right) - \\
& \qquad \qquad \qquad \frac{3\sqrt{a^2 - x^2} \int (a-x)x(a+x)\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\
& \qquad \qquad \qquad \downarrow 6308
\end{aligned}$$

$$\begin{aligned}
& \frac{3}{4}a^2 \left( \frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \operatorname{arccosh}\left(\frac{x}{a}\right) dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2} \right) \\
& \frac{3\sqrt{a^2-x^2} \int (a-x)x(a+x) \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\
& \quad \downarrow \text{6327} \\
& \frac{3}{4}a^2 \left( \frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \operatorname{arccosh}\left(\frac{x}{a}\right) dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2} \right) \\
& \frac{3\sqrt{a^2-x^2} \int x(a^2-x^2) \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\
& \quad \downarrow \text{6329} \\
& \frac{3}{4}a^2 \left( \frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \operatorname{arccosh}\left(\frac{x}{a}\right) dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2} \right) \\
& \frac{3\sqrt{a^2-x^2} \left( \frac{1}{8}a^3 \int \frac{(\frac{x}{a}-1)^{3/2}(\frac{x}{a}+1)^{3/2}}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx - \frac{1}{4}(a^2-x^2)^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\
& \frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\
& \quad \downarrow \text{6322}
\end{aligned}$$

---

3.397.  $\int (a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$

$$\frac{3}{4}a^2 \left( \frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2} \right) \\ - \frac{3\sqrt{a^2-x^2} \left( \frac{1}{8}a^4 \int \frac{\left(\frac{x}{a}-1\right)^2\left(\frac{x}{a}+1\right)^2}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right) - \frac{1}{4}(a^2-x^2)^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

↓ 3042

$$\frac{3}{4}a^2 \left( \frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2} \right) \\ - \frac{3\sqrt{a^2-x^2} \left( -\frac{1}{4}(a^2-x^2)^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} + \frac{1}{8}a^4 \int \frac{\sin\left(i\operatorname{arccosh}\left(\frac{x}{a}\right)\right)^4}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right) \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

↓ 3793

$$\frac{3}{4}a^2 \left( \frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2-x^2} \right) \\ - \frac{3\sqrt{a^2-x^2} \left( \frac{1}{8}a^4 \int \left( -\frac{\cosh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} + \frac{\cosh\left(4\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{8\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} + \frac{3}{8\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \right) d\operatorname{arccosh}\left(\frac{x}{a}\right) - \frac{1}{4}(a^2-x^2)^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

↓ 2009

---

3.397.  $\int (a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$

$$\frac{3}{4}a^2 \left( \frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \operatorname{arccosh}\left(\frac{x}{a}\right) dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \right)$$

$$\frac{\frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} - 3\sqrt{a^2 - x^2} \left( \frac{1}{8}a^4 \left( \frac{1}{32}\sqrt{\pi} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right) + \frac{1}{32}\sqrt{\pi} \operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}}$$

↓ 6368

$$\frac{3}{4}a^2 \left( \frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \int \frac{x^2}{a^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right) \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \right)$$

$$\frac{\frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} - 3\sqrt{a^2 - x^2} \left( \frac{1}{8}a^4 \left( \frac{1}{32}\sqrt{\pi} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right) + \frac{1}{32}\sqrt{\pi} \operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}}$$

↓ 3042

$$\frac{3}{4}a^2 \left( \frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \int \frac{\sin\left(i\operatorname{arccosh}\left(\frac{x}{a}\right) + \frac{\pi}{2}\right)^2}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right) \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \right)$$

$$\frac{\frac{1}{4}x(a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} - 3\sqrt{a^2 - x^2} \left( \frac{1}{8}a^4 \left( \frac{1}{32}\sqrt{\pi} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right) + \frac{1}{32}\sqrt{\pi} \operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}}$$

↓ 3793

$$\frac{3}{4}a^2 \left( -\frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \int \left( \frac{\cosh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} + \frac{1}{2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \right) d\operatorname{arccosh}\left(\frac{x}{a}\right) \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2-x^2}}{5\sqrt{\frac{x}{a}}} \right) - \frac{\frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} - 3\sqrt{a^2-x^2} \left( \frac{1}{8}a^4 \left( \frac{1}{32}\sqrt{\pi} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + \frac{1}{32}\sqrt{\pi} \operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}}$$

↓ 2009

$$\frac{3}{4}a^2 \left( -\frac{3\sqrt{a^2-x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \left( \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right) \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\frac{1}{4}x(a^2-x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} - 3\sqrt{a^2-x^2} \left( \frac{1}{8}a^4 \left( \frac{1}{32}\sqrt{\pi} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + \frac{1}{32}\sqrt{\pi} \operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right)}{8a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}}$$

input `Int[(a^2 - x^2)^(3/2)*ArcCosh[x/a]^(3/2), x]`

output `(x*(a^2 - x^2)^(3/2)*ArcCosh[x/a]^(3/2))/4 - (3*Sqrt[a^2 - x^2]*(-1/4*((a^2 - x^2)^2*Sqrt[ArcCosh[x/a]]) + (a^4*((3*Sqrt[ArcCosh[x/a]]))/4 + (Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[x/a]]])/32 - (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[x/a]]])/4 + (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[x/a]]])/32 - (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[x/a]]])/4))/8)/(8*a*Sqrt[-1 + x/a]*Sqrt[1 + x/a]) + (3*a^2*((x*Sqrt[a^2 - x^2]*ArcCosh[x/a]^(3/2))/2 - (a*Sqrt[a^2 - x^2]*ArcCosh[x/a]^(5/2))/(5*Sqrt[-1 + x/a]*Sqrt[1 + x/a]) - (3*Sqrt[a^2 - x^2]*((x^2*Sqrt[ArcCosh[x/a]]))/2 - (a^2*(Sqrt[ArcCosh[x/a]] + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[x/a]]])/4 + (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[x/a]]])/4)))/4)/(4*a*Sqrt[-1 + x/a]*Sqrt[1 + x/a]))/4`

## 3.397.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`
- rule 6299 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcCosh[c*x])^n/(m + 1)), x] - Simp[b*c*(n/(m + 1)) Int[x^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[m, 0] && GtQ[n, 0]`
- rule 6308 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sqrt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && NeQ[n, -1]`
- rule 6310 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcCosh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]`

rule 6312 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] :> Simp[x*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^n/(2*p + 1)), x] +  
(Simp[2*d*(p/(2*p + 1)) Int[(d + e*x^2)^(p - 1)*(a + b*ArcCosh[c*x])^n, x  
, x] - Simp[b*c*(n/(2*p + 1))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p  
)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n  
- 1), x], x]) /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n,  
0] && GtQ[p, 0]`

rule 6322 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*  
(d2_) + (e2_.)*(x_))^(p_.), x_Symbol] :> Simp[(1/(b*c))*Simp[(d1 + e1*x)^p/  
(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Sinh[-a/b + x  
/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2  
, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[2*p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (  
e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] :> Int[(f*x)^m*(d1  
*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2  
, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6329 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p  
_.), x_Symbol] :> Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e*(p  
+ 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 +  
c*x)^p)] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x  
)^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] &&  
GtQ[n, 0] && NeQ[p, -1]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x  
_)^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*  
Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int  
t[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c  
*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[  
e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`



**3.397.4 Maple [F]**

$$\int (a^2 - x^2)^{\frac{3}{2}} \operatorname{arccosh}\left(\frac{x}{a}\right)^{\frac{3}{2}} dx$$

input `int((a^2-x^2)^(3/2)*arccosh(x/a)^(3/2),x)`

output `int((a^2-x^2)^(3/2)*arccosh(x/a)^(3/2),x)`

**3.397.5 Fricas [F(-2)]**

Exception generated.

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \text{Exception raised: TypeError}$$

input `integrate((a^2-x^2)^(3/2)*arccosh(x/a)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.397.6 Sympy [F(-1)]**

Timed out.

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \text{Timed out}$$

input `integrate((a**2-x**2)**(3/2)*acosh(x/a)**(3/2),x)`

output `Timed out`

**3.397.7 Maxima [F]**

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \int (a^2 - x^2)^{\frac{3}{2}} \operatorname{arcosh}\left(\frac{x}{a}\right)^{\frac{3}{2}} dx$$

input `integrate((a^2-x^2)^(3/2)*arccosh(x/a)^(3/2),x, algorithm="maxima")`

output `integrate((a^2 - x^2)^(3/2)*arccosh(x/a)^(3/2), x)`

**3.397.8 Giac [F]**

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \int (a^2 - x^2)^{\frac{3}{2}} \operatorname{arcosh}\left(\frac{x}{a}\right)^{\frac{3}{2}} dx$$

input `integrate((a^2-x^2)^(3/2)*arccosh(x/a)^(3/2),x, algorithm="giac")`

output `integrate((a^2 - x^2)^(3/2)*arccosh(x/a)^(3/2), x)`

**3.397.9 Mupad [F(-1)]**

Timed out.

$$\int (a^2 - x^2)^{3/2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \int \operatorname{acosh}\left(\frac{x}{a}\right)^{3/2} (a^2 - x^2)^{3/2} dx$$

input `int(acosh(x/a)^(3/2)*(a^2 - x^2)^(3/2),x)`

output `int(acosh(x/a)^(3/2)*(a^2 - x^2)^(3/2), x)`

### 3.398 $\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$

3.398.1 Optimal result	3098
3.398.2 Mathematica [A] (verified)	3099
3.398.3 Rubi [A] (verified)	3099
3.398.4 Maple [F]	3102
3.398.5 Fracas [F(-2)]	3102
3.398.6 Sympy [F]	3102
3.398.7 Maxima [F]	3103
3.398.8 Giac [F]	3103
3.398.9 Mupad [F(-1)]	3103

#### 3.398.1 Optimal result

Integrand size = 24, antiderivative size = 316

$$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \frac{3a\sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{16\sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} - \frac{3x^2\sqrt{a^2 - x^2} \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{8a\sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} + \frac{1}{2}x\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} - \frac{a\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} + \frac{3a\sqrt{\frac{\pi}{2}}\sqrt{a^2 - x^2} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{64\sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}} + \frac{3a\sqrt{\frac{\pi}{2}}\sqrt{a^2 - x^2} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)}{64\sqrt{-1 + \frac{x}{a}} \sqrt{1 + \frac{x}{a}}}$$

output  $1/2*x*\operatorname{arccosh}(x/a)^{(3/2)}*(a^2-x^2)^{(1/2)}-1/5*a*\operatorname{arccosh}(x/a)^{(5/2)}*(a^2-x^2)^{(1/2)} / (-1+x/a)^{(1/2)} / (1+x/a)^{(1/2)} + 3/128*a*\operatorname{erf}(2^{(1/2)}*\operatorname{arccosh}(x/a)^{(1/2)}) * 2^{(1/2)} * \operatorname{Pi}^{(1/2)} * (a^2-x^2)^{(1/2)} / (-1+x/a)^{(1/2)} / (1+x/a)^{(1/2)} + 3/128*a*\operatorname{erfi}(2^{(1/2)}*\operatorname{arccosh}(x/a)^{(1/2)}) * 2^{(1/2)} * \operatorname{Pi}^{(1/2)} * (a^2-x^2)^{(1/2)} / (-1+x/a)^{(1/2)} / (1+x/a)^{(1/2)} + 3/16*a*(a^2-x^2)^{(1/2)}*\operatorname{arccosh}(x/a)^{(1/2)} / (-1+x/a)^{(1/2)} / (1+x/a)^{(1/2)} - 3/8*x^2*(a^2-x^2)^{(1/2)}*\operatorname{arccosh}(x/a)^{(1/2)} / a / (-1+x/a)^{(1/2)} / (1+x/a)^{(1/2)}$

**3.398.2 Mathematica [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.46

$$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \frac{a^2 \sqrt{a^2 - x^2} \left(15\sqrt{2\pi} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + 15\sqrt{2\pi} \operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right)\right)}{640 \sqrt{(-a+x)/(a+x)} (a+x)}$$

input `Integrate[Sqrt[a^2 - x^2]*ArcCosh[x/a]^(3/2),x]`

output `(a^2*Sqrt[a^2 - x^2]*(15*Sqrt[2*Pi]*Erf[Sqrt[2]*Sqrt[ArcCosh[x/a]]] + 15*Sqrt[2*Pi]*Erfi[Sqrt[2]*Sqrt[ArcCosh[x/a]]] - 8*Sqrt[ArcCosh[x/a]]*(16*ArcCosh[x/a]^2 + 15*Cosh[2*ArcCosh[x/a]] - 20*ArcCosh[x/a]*Sinh[2*ArcCosh[x/a]])))/(640*Sqrt[(-a+x)/(a+x)]*(a+x))`

**3.398.3 Rubi [A] (verified)**Time = 1.75 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.69, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$ , Rules used = {6310, 6299, 6308, 6368, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx \\ & \quad \downarrow \text{6310} \\ & -\frac{3\sqrt{a^2 - x^2} \int x \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\ & \quad \downarrow \text{6299} \\ & -\frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} \operatorname{arccosh}\left(\frac{x}{a}\right) dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{\sqrt{a^2 - x^2} \int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} dx}{2\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \\ & \quad \frac{1}{2}x\sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} \\ & \quad \downarrow \text{6308} \end{aligned}$$

---

3.398.  $\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$

$$\frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{\int \frac{x^2}{\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} dx}{4a} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

↓ 6368

$$\frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \int \frac{x^2}{a^2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right) \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

↓ 3042

$$\frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \int \frac{\sin\left(i\operatorname{arccosh}\left(\frac{x}{a}\right) + \frac{\pi}{2}\right)^2}{\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} d\operatorname{arccosh}\left(\frac{x}{a}\right) \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

↓ 3793

$$\frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \int \left( \frac{\cosh\left(2\operatorname{arccosh}\left(\frac{x}{a}\right)\right)}{2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} + \frac{1}{2\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}} \right) d\operatorname{arccosh}\left(\frac{x}{a}\right) \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

↓ 2009

$$\frac{3\sqrt{a^2 - x^2} \left( \frac{1}{2}x^2 \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} - \frac{1}{4}a^2 \left( \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}\right) \right) + \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)} \right)}{4a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} - \frac{a\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}} + \frac{1}{2}x\sqrt{a^2 - x^2}\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}$$

input `Int[Sqrt[a^2 - x^2]*ArcCosh[x/a]^(3/2), x]`

```
output (x*Sqrt[a^2 - x^2]*ArcCosh[x/a]^(3/2))/2 - (a*Sqrt[a^2 - x^2]*ArcCosh[x/a]
^(5/2))/(5*Sqrt[-1 + x/a]*Sqrt[1 + x/a]) - (3*Sqrt[a^2 - x^2]*((x^2*Sqrt[A
rcCosh[x/a]]))/2 - (a^2*(Sqrt[ArcCosh[x/a]] + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[
ArcCosh[x/a]]))/4 + (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[x/a]]])/4))/4)/
(4*a*Sqrt[-1 + x/a]*Sqrt[1 + x/a])
```

### 3.398.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3793 Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

```
rule 6299 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*(x_)^(m_.), x_Symbol] := Simp[
x^(m + 1)*((a + b*ArcCosh[c*x])^n/(m + 1)), x] - Simp[b*c*(n/(m + 1)) Int
[x^(m + 1)*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x
], x] /; FreeQ[{a, b, c}, x] && IGtQ[m, 0] && GtQ[n, 0]
```

```
rule 6308 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/(Sqrt[(d1_) + (e1_.)*(x_)]*Sq
rt[(d2_) + (e2_.)*(x_)]), x_Symbol] := Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 +
c*x]/Sqrt[d1 + e1*x]]*Simp[Sqrt[-1 + c*x]/Sqrt[d2 + e2*x]]*(a + b*ArcCosh[
c*x])^(n + 1), x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1
] && EqQ[e2, (-c)*d2] && NeQ[n, -1]
```

```
rule 6310 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*Sqrt[(d_) + (e_.)*(x_)^2], x_
Symbol] := Simp[x*Sqrt[d + e*x^2]*((a + b*ArcCosh[c*x])^n/2), x] + (-Simp[(
1/2)*Simp[Sqrt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[(a + b*ArcC
osh[c*x])^n/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] - Simp[b*c*(n/2)*Simp[Sq
rt[d + e*x^2]/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])] Int[x*(a + b*ArcCosh[c*x])^
(n - 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && GtQ[n
, 0]
```

```
rule 6368 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]
```

### 3.398.4 Maple [F]

$$\int \operatorname{arccosh}\left(\frac{x}{a}\right)^{\frac{3}{2}} \sqrt{a^2 - x^2} dx$$

```
input int(arccosh(x/a)^(3/2)*(a^2-x^2)^(1/2),x)
```

```
output int(arccosh(x/a)^(3/2)*(a^2-x^2)^(1/2),x)
```

### 3.398.5 Fricas [F(-2)]

Exception generated.

$$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \text{Exception raised: TypeError}$$

```
input integrate(arccosh(x/a)^(3/2)*(a^2-x^2)^(1/2),x, algorithm="fricas")
```

```
output Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)
```

### 3.398.6 Sympy [F]

$$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \int \sqrt{-(-a + x)(a + x)} \operatorname{acosh}^{\frac{3}{2}}\left(\frac{x}{a}\right) dx$$

```
input integrate(acosh(x/a)**(3/2)*(a**2-x**2)**(1/2),x)
```

```
output Integral(sqrt(-(-a + x)*(a + x))*acosh(x/a)**(3/2), x)
```

---

3.398.  $\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx$

**3.398.7 Maxima [F]**

$$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{\frac{3}{2}} dx$$

input `integrate(arccosh(x/a)^(3/2)*(a^2-x^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(a^2 - x^2)*arccosh(x/a)^(3/2), x)`

**3.398.8 Giac [F]**

$$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{\frac{3}{2}} dx$$

input `integrate(arccosh(x/a)^(3/2)*(a^2-x^2)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(a^2 - x^2)*arccosh(x/a)^(3/2), x)`

**3.398.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a^2 - x^2} \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2} dx = \int \operatorname{acosh}\left(\frac{x}{a}\right)^{3/2} \sqrt{a^2 - x^2} dx$$

input `int(acosh(x/a)^(3/2)*(a^2 - x^2)^(1/2),x)`

output `int(acosh(x/a)^(3/2)*(a^2 - x^2)^(1/2), x)`



**3.399**  $\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2-x^2}} dx$

3.399.1 Optimal result . . . . .	3104
3.399.2 Mathematica [A] (verified) . . . . .	3104
3.399.3 Rubi [A] (verified) . . . . .	3105
3.399.4 Maple [A] (verified) . . . . .	3105
3.399.5 Fricas [F(-2)] . . . . .	3106
3.399.6 Sympy [F] . . . . .	3106
3.399.7 Maxima [F] . . . . .	3106
3.399.8 Giac [F] . . . . .	3107
3.399.9 Mupad [F(-1)] . . . . .	3107

**3.399.1 Optimal result**

Integrand size = 24, antiderivative size = 50

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2-x^2}} dx = \frac{2a\sqrt{-1+\frac{x}{a}}\sqrt{1+\frac{x}{a}}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{a^2-x^2}}$$

output `2/5*a*arccosh(x/a)^(5/2)*(-1+x/a)^(1/2)*(1+x/a)^(1/2)/(a^2-x^2)^(1/2)`

**3.399.2 Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2-x^2}} dx = \frac{2a\sqrt{-1+\frac{x}{a}}\sqrt{1+\frac{x}{a}}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{a^2-x^2}}$$

input `Integrate[ArcCosh[x/a]^(3/2)/Sqrt[a^2 - x^2], x]`

output `(2*a*Sqrt[-1 + x/a]*Sqrt[1 + x/a]*ArcCosh[x/a]^(5/2))/(5*Sqrt[a^2 - x^2])`

**3.399.3 Rubi [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx$$

↓ 6307

$$\frac{2a\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1}\operatorname{arccosh}\left(\frac{x}{a}\right)^{5/2}}{5\sqrt{a^2-x^2}}$$

input `Int[ArcCosh[x/a]^(3/2)/Sqrt[a^2 - x^2], x]`

output `(2*a*Sqrt[-1 + x/a]*Sqrt[1 + x/a]*ArcCosh[x/a]^(5/2))/(5*Sqrt[a^2 - x^2])`

**3.399.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.399.4 Maple [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{2 \operatorname{arccosh}\left(\frac{x}{a}\right)^{\frac{5}{2}} \sqrt{-\frac{a-x}{a}} \sqrt{\frac{a+x}{a}} a}{5\sqrt{(a-x)(a+x)}}$	44

input `int(arccosh(x/a)^(3/2)/(a^2-x^2)^(1/2), x, method=_RETURNVERBOSE)`

output  $2/5*\operatorname{arccosh}(x/a)^{(5/2)/((a-x)*(a+x))^{(1/2)*(-(a-x)/a)^{(1/2)*((a+x)/a)^{(1/2)}}*a$

### 3.399.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(arccosh(x/a)^(3/2)/(a^2-x^2)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.399.6 Sympy [F]

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx = \int \frac{\operatorname{acosh}^{\frac{3}{2}}\left(\frac{x}{a}\right)}{\sqrt{-(-a+x)(a+x)}} dx$$

input `integrate(acosh(x/a)**(3/2)/(a**2-x**2)**(1/2),x)`

output `Integral(acosh(x/a)**(3/2)/sqrt(-(-a+x)*(a+x)), x)`

### 3.399.7 Maxima [F]

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx = \int \frac{\operatorname{arcosh}\left(\frac{x}{a}\right)^{\frac{3}{2}}}{\sqrt{a^2 - x^2}} dx$$

input `integrate(arccosh(x/a)^(3/2)/(a^2-x^2)^(1/2),x, algorithm="maxima")`

output `integrate(arccosh(x/a)^(3/2)/sqrt(a^2 - x^2), x)`

---

3.399.  $\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx$

**3.399.8 Giac [F]**

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx = \int \frac{\operatorname{arcosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx$$

input `integrate(arccosh(x/a)^(3/2)/(a^2-x^2)^(1/2),x, algorithm="giac")`

output `integrate(arccosh(x/a)^(3/2)/sqrt(a^2 - x^2), x)`

**3.399.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx = \int \frac{\operatorname{acosh}\left(\frac{x}{a}\right)^{3/2}}{\sqrt{a^2 - x^2}} dx$$

input `int(acosh(x/a)^(3/2)/(a^2 - x^2)^(1/2), x)`

output `int(acosh(x/a)^(3/2)/(a^2 - x^2)^(1/2), x)`

**3.400**  $\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2-x^2)^{3/2}} dx$

3.400.1 Optimal result . . . . . 3108  
 3.400.2 Mathematica [N/A] . . . . . 3108  
 3.400.3 Rubi [N/A] . . . . . 3109  
 3.400.4 Maple [N/A] (verified) . . . . . 3110  
 3.400.5 Fricas [F(-2)] . . . . . 3110  
 3.400.6 Sympy [N/A] . . . . . 3111  
 3.400.7 Maxima [N/A] . . . . . 3111  
 3.400.8 Giac [N/A] . . . . . 3111  
 3.400.9 Mupad [N/A] . . . . . 3112

**3.400.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2-x^2)^{3/2}} dx = \frac{x \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{a^2 \sqrt{a^2-x^2}} + \frac{3 \sqrt{-1+\frac{x}{a}} \sqrt{1+\frac{x}{a}} \operatorname{Int}\left(\frac{x \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{1-\frac{x^2}{a^2}}, x\right)}{2a^3 \sqrt{a^2-x^2}}$$

output `x*arccosh(x/a)^(3/2)/a^2/(a^2-x^2)^(1/2)+3/2*(-1+x/a)^(1/2)*(1+x/a)^(1/2)*  
 Unintegrable(x*arccosh(x/a)^(1/2)/(1-x^2/a^2),x)/a^3/(a^2-x^2)^(1/2)`

**3.400.2 Mathematica [N/A]**

Not integrable

Time = 7.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2-x^2)^{3/2}} dx = \int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2-x^2)^{3/2}} dx$$

input `Integrate[ArcCosh[x/a]^(3/2)/(a^2-x^2)^(3/2),x]`

output `Integrate[ArcCosh[x/a]^(3/2)/(a^2-x^2)^(3/2),x]`

---

3.400.  $\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2-x^2)^{3/2}} dx$

**3.400.3 Rubi [N/A]**

Not integrable

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6314, 27, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx$$

↓ 6314

$$\frac{3\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{a^2 x \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2 - x^2} dx}{2a^3 \sqrt{a^2 - x^2}} + \frac{x \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{a^2 \sqrt{a^2 - x^2}}$$

↓ 27

$$\frac{3\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2 - x^2} dx}{2a \sqrt{a^2 - x^2}} + \frac{x \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{a^2 \sqrt{a^2 - x^2}}$$

↓ 6375

$$\frac{3\sqrt{\frac{x}{a}-1}\sqrt{\frac{x}{a}+1} \int \frac{x \sqrt{\operatorname{arccosh}\left(\frac{x}{a}\right)}}{a^2 - x^2} dx}{2a \sqrt{a^2 - x^2}} + \frac{x \operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{a^2 \sqrt{a^2 - x^2}}$$

input `Int[ArcCosh[x/a]^(3/2)/(a^2 - x^2)^(3/2), x]`

output `$Aborted`

**3.400.3.1 Defintions of rubi rules used**

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

```
rule 6314 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/((d_) + (e_.)*(x_)^2)^(3/2),
x_Symbol] := Simp[x*((a + b*ArcCosh[c*x])^n/(d*Sqrt[d + e*x^2])), x] + Simp
[b*c*(n/d)*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])] Int[x*((a
+ b*ArcCosh[c*x])^(n - 1)/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e},
x] && EqQ[c^2*d + e, 0] && GtQ[n, 0]
```

```
rule 6375 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*A
rcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]
```

### 3.400.4 Maple [N/A] (verified)

Not integrable

Time = 1.43 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{\frac{3}{2}}}{(a^2 - x^2)^{\frac{3}{2}}} dx$$

```
input int(arccosh(x/a)^(3/2)/(a^2-x^2)^(3/2), x)
```

```
output int(arccosh(x/a)^(3/2)/(a^2-x^2)^(3/2), x)
```

### 3.400.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx = \text{Exception raised: TypeError}$$

```
input integrate(arccosh(x/a)^(3/2)/(a^2-x^2)^(3/2), x, algorithm="fricas")
```

```
output Exception raised: TypeError >> Error detected within library code: inte
grate: implementation incomplete (constant residues)
```

**3.400.6 Sympy [N/A]**

Not integrable

Time = 48.71 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\operatorname{acosh}^{\frac{3}{2}}\left(\frac{x}{a}\right)}{\left(-(-a + x)(a + x)\right)^{\frac{3}{2}}} dx$$

input `integrate(acosh(x/a)**(3/2)/(a**2-x**2)**(3/2),x)`output `Integral(acosh(x/a)**(3/2)/(-(-a + x)*(a + x))**(3/2), x)`**3.400.7 Maxima [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}\left(\frac{x}{a}\right)^{\frac{3}{2}}}{(a^2 - x^2)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(x/a)^(3/2)/(a^2-x^2)^(3/2),x, algorithm="maxima")`output `integrate(arccosh(x/a)^(3/2)/(a^2 - x^2)^(3/2), x)`**3.400.8 Giac [N/A]**

Not integrable

Time = 4.40 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\operatorname{arcosh}\left(\frac{x}{a}\right)^{\frac{3}{2}}}{(a^2 - x^2)^{\frac{3}{2}}} dx$$

input `integrate(arccosh(x/a)^(3/2)/(a^2-x^2)^(3/2),x, algorithm="giac")`output `integrate(arccosh(x/a)^(3/2)/(a^2 - x^2)^(3/2), x)`

---

3.400.  $\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx$



**3.400.9 Mupad [N/A]**

Not integrable

Time = 2.87 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\operatorname{arccosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx = \int \frac{\operatorname{acosh}\left(\frac{x}{a}\right)^{3/2}}{(a^2 - x^2)^{3/2}} dx$$

input `int(acosh(x/a)^(3/2)/(a^2 - x^2)^(3/2), x)`output `int(acosh(x/a)^(3/2)/(a^2 - x^2)^(3/2), x)`

**3.401**  $\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx$

3.401.1 Optimal result . . . . . 3113  
 3.401.2 Mathematica [A] (verified) . . . . . 3113  
 3.401.3 Rubi [A] (verified) . . . . . 3114  
 3.401.4 Maple [F] . . . . . 3116  
 3.401.5 Fricas [F(-2)] . . . . . 3116  
 3.401.6 Sympy [F] . . . . . 3117  
 3.401.7 Maxima [F] . . . . . 3117  
 3.401.8 Giac [F] . . . . . 3117  
 3.401.9 Mupad [F(-1)] . . . . . 3118

**3.401.1 Optimal result**

Integrand size = 19, antiderivative size = 65

$$\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx = \frac{\sqrt{\pi}\sqrt{-1+x}\operatorname{erf}\left(\sqrt{\operatorname{arccosh}(x)}\right)}{2\sqrt{1-x}} + \frac{\sqrt{\pi}\sqrt{-1+x}\operatorname{erfi}\left(\sqrt{\operatorname{arccosh}(x)}\right)}{2\sqrt{1-x}}$$

output `1/2*erf(arccosh(x)^(1/2))*Pi^(1/2)*(-1+x)^(1/2)/(1-x)^(1/2)+1/2*erfi(arccosh(x)^(1/2))*Pi^(1/2)*(-1+x)^(1/2)/(1-x)^(1/2)`

**3.401.2 Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11

$$\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx = \frac{\sqrt{-((-1+x)(1+x))}\left(\sqrt{-\operatorname{arccosh}(x)}\Gamma\left(\frac{1}{2}, -\operatorname{arccosh}(x)\right) - \sqrt{\operatorname{arccosh}(x)}\Gamma\left(\frac{1}{2}, \operatorname{arccosh}(x)\right)\right)}{2\sqrt{\frac{-1+x}{1+x}}(1+x)\sqrt{\operatorname{arccosh}(x)}}$$

input `Integrate[x/(Sqrt[1 - x^2]*Sqrt[ArcCosh[x]]),x]`

---

3.401.  $\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx$

output 
$$\frac{-1/2*(\text{Sqrt}[ -((-1 + x)*(1 + x))] * (\text{Sqrt}[-\text{ArcCosh}[x]] * \text{Gamma}[1/2, -\text{ArcCosh}[x]] - \text{Sqrt}[\text{ArcCosh}[x]] * \text{Gamma}[1/2, \text{ArcCosh}[x]]))}{(\text{Sqrt}[(-1 + x)/(1 + x)] * (1 + x) * \text{Sqrt}[\text{ArcCosh}[x]])}$$

### 3.401.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.77, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {6367, 3042, 3788, 26, 2611, 2633, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{\sqrt{1-x^2} \sqrt{\text{arccosh}(x)}} dx \\ & \quad \downarrow \text{6367} \\ & \frac{\sqrt{x-1} \int \frac{x}{\sqrt{\text{arccosh}(x)}} d\text{arccosh}(x)}{\sqrt{1-x}} \\ & \quad \downarrow \text{3042} \\ & \frac{\sqrt{x-1} \int \frac{\sin\left(i\text{arccosh}(x) + \frac{\pi}{2}\right)}{\sqrt{\text{arccosh}(x)}} d\text{arccosh}(x)}{\sqrt{1-x}} \\ & \quad \downarrow \text{3788} \\ & \frac{\sqrt{x-1} \left( \frac{1}{2} i \int -\frac{ie^{\text{arccosh}(x)}}{\sqrt{\text{arccosh}(x)}} d\text{arccosh}(x) - \frac{1}{2} i \int \frac{ie^{-\text{arccosh}(x)}}{\sqrt{\text{arccosh}(x)}} d\text{arccosh}(x) \right)}{\sqrt{1-x}} \\ & \quad \downarrow \text{26} \\ & \frac{\sqrt{x-1} \left( \frac{1}{2} \int \frac{e^{-\text{arccosh}(x)}}{\sqrt{\text{arccosh}(x)}} d\text{arccosh}(x) + \frac{1}{2} \int \frac{e^{\text{arccosh}(x)}}{\sqrt{\text{arccosh}(x)}} d\text{arccosh}(x) \right)}{\sqrt{1-x}} \\ & \quad \downarrow \text{2611} \\ & \frac{\sqrt{x-1} \left( \int e^{-\text{arccosh}(x)} d\sqrt{\text{arccosh}(x)} + \int e^{\text{arccosh}(x)} d\sqrt{\text{arccosh}(x)} \right)}{\sqrt{1-x}} \\ & \quad \downarrow \text{2633} \end{aligned}$$

---

3.401.  $\int \frac{x}{\sqrt{1-x^2} \sqrt{\text{arccosh}(x)}} dx$

$$\frac{\sqrt{x-1} \left( \int e^{-\operatorname{arccosh}(x)} d\sqrt{\operatorname{arccosh}(x)} + \frac{1}{2} \sqrt{\pi} \operatorname{erfi} \left( \sqrt{\operatorname{arccosh}(x)} \right) \right)}{\sqrt{1-x}}$$

↓ 2634

$$\frac{\sqrt{x-1} \left( \frac{1}{2} \sqrt{\pi} \operatorname{erf} \left( \sqrt{\operatorname{arccosh}(x)} \right) + \frac{1}{2} \sqrt{\pi} \operatorname{erfi} \left( \sqrt{\operatorname{arccosh}(x)} \right) \right)}{\sqrt{1-x}}$$

input `Int[x/(Sqrt[1 - x^2]*Sqrt[ArcCosh[x]]), x]`

output `(Sqrt[-1 + x]*((Sqrt[Pi]*Erf[Sqrt[ArcCosh[x]]])/2 + (Sqrt[Pi]*Erfi[Sqrt[ArcCosh[x]]])/2))/Sqrt[1 - x]`

### 3.401.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2611 `Int[(F_)^((g_)*((e_) + (f_)*(x_)))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 2633 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`

rule 2634 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3788 `Int[((c_.) + (d_.)*(x_))^(m_.)*sin[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol]
-> Simp[I/2 Int[(c + d*x)^m/(E^(I*k*Pi)*E^(I*(e + f*x))), x], x] - Simp
[I/2 Int[(c + d*x)^m*E^(I*k*Pi)*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]
&& IntegerQ[2*k]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol]
-> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x]
&& EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.401.4 Maple [F]

$$\int \frac{x}{\sqrt{-x^2+1} \sqrt{\operatorname{arccosh}(x)}} dx$$

input `int(x/(-x^2+1)^(1/2)/arccosh(x)^(1/2),x)`

output `int(x/(-x^2+1)^(1/2)/arccosh(x)^(1/2),x)`

### 3.401.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{x}{\sqrt{1-x^2} \sqrt{\operatorname{arccosh}(x)}} dx = \text{Exception raised: TypeError}$$

input `integrate(x/(-x^2+1)^(1/2)/arccosh(x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.401.6 Sympy [F]**

$$\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx = \int \frac{x}{\sqrt{-(x-1)(x+1)}\sqrt{\operatorname{acosh}(x)}} dx$$

input `integrate(x/(-x**2+1)**(1/2)/acosh(x)**(1/2),x)`

output `Integral(x/(sqrt(-(x - 1)*(x + 1))*sqrt(acosh(x))), x)`

**3.401.7 Maxima [F]**

$$\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx = \int \frac{x}{\sqrt{-x^2+1}\sqrt{\operatorname{arcosh}(x)}} dx$$

input `integrate(x/(-x^2+1)^(1/2)/arccosh(x)^(1/2),x, algorithm="maxima")`

output `integrate(x/(sqrt(-x^2 + 1)*sqrt(arccosh(x))), x)`

**3.401.8 Giac [F]**

$$\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx = \int \frac{x}{\sqrt{-x^2+1}\sqrt{\operatorname{arcosh}(x)}} dx$$

input `integrate(x/(-x^2+1)^(1/2)/arccosh(x)^(1/2),x, algorithm="giac")`

output `integrate(x/(sqrt(-x^2 + 1)*sqrt(arccosh(x))), x)`

**3.401.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x}{\sqrt{1-x^2}\sqrt{\operatorname{arccosh}(x)}} dx = \int \frac{x}{\sqrt{\operatorname{acosh}(x)}\sqrt{1-x^2}} dx$$

input `int(x/(acosh(x)^(1/2)*(1 - x^2)^(1/2)), x)`output `int(x/(acosh(x)^(1/2)*(1 - x^2)^(1/2)), x)`

**3.402**      $\int \frac{(c-a^2cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$

3.402.1 Optimal result	3119
3.402.2 Mathematica [A] (verified)	3120
3.402.3 Rubi [A] (verified)	3121
3.402.4 Maple [F]	3122
3.402.5 Fracas [F(-2)]	3123
3.402.6 Sympy [F(-1)]	3123
3.402.7 Maxima [F]	3123
3.402.8 Giac [F]	3124
3.402.9 Mupad [F(-1)]	3124

**3.402.1 Optimal result**

Integrand size = 24, antiderivative size = 438

$$\int \frac{(c - a^2cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = -\frac{5c^2\sqrt{c - a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}}{8a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$- \frac{3c^2\sqrt{\pi}\sqrt{c - a^2cx^2}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$+ \frac{15c^2\sqrt{\frac{\pi}{2}}\sqrt{c - a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$+ \frac{c^2\sqrt{\frac{\pi}{6}}\sqrt{c - a^2cx^2}\operatorname{erf}\left(\sqrt{6}\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$- \frac{3c^2\sqrt{\pi}\sqrt{c - a^2cx^2}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$+ \frac{15c^2\sqrt{\frac{\pi}{2}}\sqrt{c - a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$+ \frac{c^2\sqrt{\frac{\pi}{6}}\sqrt{c - a^2cx^2}\operatorname{erfi}\left(\sqrt{6}\sqrt{\operatorname{arccosh}(ax)}\right)}{64a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

---

3.402.      $\int \frac{(c-a^2cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$



output  $\frac{1}{384}c^2 \operatorname{erf}(6^{1/2} \operatorname{arccosh}(ax)^{1/2}) 6^{1/2} \pi^{1/2} (-a^2cx^2+c)^{1/2} / a (ax-1)^{1/2} (ax+1)^{1/2} + \frac{1}{384}c^2 \operatorname{erfi}(6^{1/2} \operatorname{arccosh}(ax)^{1/2}) 6^{1/2} \pi^{1/2} (-a^2cx^2+c)^{1/2} / a (ax-1)^{1/2} (ax+1)^{1/2} + \frac{5}{128}c^2 \operatorname{erf}(2^{1/2} \operatorname{arccosh}(ax)^{1/2}) 2^{1/2} \pi^{1/2} (-a^2cx^2+c)^{1/2} / a (ax-1)^{1/2} (ax+1)^{1/2} + \frac{15}{128}c^2 \operatorname{erfi}(2^{1/2} \operatorname{arccosh}(ax)^{1/2}) 2^{1/2} \pi^{1/2} (-a^2cx^2+c)^{1/2} / a (ax-1)^{1/2} (ax+1)^{1/2} - \frac{3}{64}c^2 \operatorname{erf}(2 \operatorname{arccosh}(ax)^{1/2}) \pi^{1/2} (-a^2cx^2+c)^{1/2} / a (ax-1)^{1/2} (ax+1)^{1/2} - \frac{3}{64}c^2 \operatorname{erfi}(2 \operatorname{arccosh}(ax)^{1/2}) \pi^{1/2} (-a^2cx^2+c)^{1/2} / a (ax-1)^{1/2} (ax+1)^{1/2} - \frac{5}{8}c^2 (-a^2cx^2+c)^{1/2} \operatorname{arccosh}(ax)^{1/2} / a (ax-1)^{1/2} (ax+1)^{1/2}$

### 3.402.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 209, normalized size of antiderivative = 0.48

$$\int \frac{(c - a^2cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \frac{c^2 \sqrt{c - a^2cx^2} \left( 240 \operatorname{arccosh}(ax) - \sqrt{6} \sqrt{-\operatorname{arccosh}(ax)} \Gamma\left(\frac{1}{2}, -6 \operatorname{arccosh}(ax)\right) + 18 \sqrt{-\operatorname{arccosh}(ax)} \Gamma\left(\frac{1}{2}, -4 \operatorname{arccosh}(ax)\right) \right)}{\dots}$$

input `Integrate[(c - a^2*c*x^2)^(5/2)/Sqrt[ArcCosh[a*x]], x]`

output  $-1/384(c^2 \operatorname{Sqrt}[c - a^2cx^2] * (240 \operatorname{ArcCosh}[a*x] - \operatorname{Sqrt}[6] \operatorname{Sqrt}[-\operatorname{ArcCosh}[a*x]] * \operatorname{Gamma}[1/2, -6 \operatorname{ArcCosh}[a*x]] + 18 \operatorname{Sqrt}[-\operatorname{ArcCosh}[a*x]] * \operatorname{Gamma}[1/2, -4 \operatorname{ArcCosh}[a*x]] - 45 \operatorname{Sqrt}[2] \operatorname{Sqrt}[-\operatorname{ArcCosh}[a*x]] * \operatorname{Gamma}[1/2, -2 \operatorname{ArcCosh}[a*x]] + 45 \operatorname{Sqrt}[2] \operatorname{Sqrt}[\operatorname{ArcCosh}[a*x]] * \operatorname{Gamma}[1/2, 2 \operatorname{ArcCosh}[a*x]] - 18 \operatorname{Sqrt}[\operatorname{ArcCosh}[a*x]] * \operatorname{Gamma}[1/2, 4 \operatorname{ArcCosh}[a*x]] + \operatorname{Sqrt}[6] \operatorname{Sqrt}[\operatorname{ArcCosh}[a*x]] * \operatorname{Gamma}[1/2, 6 \operatorname{ArcCosh}[a*x]])) / (a \operatorname{Sqrt}[(-1 + a*x)/(1 + a*x)] * (1 + a*x) \operatorname{Sqrt}[\operatorname{ArcCosh}[a*x]]))$

**3.402.3 Rubi [A] (verified)**

Time = 0.54 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.47, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$ , Rules used = {6321, 3042, 25, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx \\
 & \quad \downarrow \text{6321} \\
 & \frac{c^2 \sqrt{c - a^2 cx^2} \int \frac{(ax-1)^3 (ax+1)^3}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{c^2 \sqrt{c - a^2 cx^2} \int -\frac{\sin(i\operatorname{arccosh}(ax))^6}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{25} \\
 & -\frac{c^2 \sqrt{c - a^2 cx^2} \int \frac{\sin(i\operatorname{arccosh}(ax))^6}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{3793} \\
 & -\frac{c^2 \sqrt{c - a^2 cx^2} \int \left( -\frac{15 \cosh(2\operatorname{arccosh}(ax))}{32\sqrt{\operatorname{arccosh}(ax)}} + \frac{3 \cosh(4\operatorname{arccosh}(ax))}{16\sqrt{\operatorname{arccosh}(ax)}} - \frac{\cosh(6\operatorname{arccosh}(ax))}{32\sqrt{\operatorname{arccosh}(ax)}} + \frac{5}{16\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{c^2 \sqrt{c - a^2 cx^2} \left( -\frac{3}{64} \sqrt{\pi} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{15}{64} \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{64} \sqrt{\frac{\pi}{6}} \operatorname{erf}\left(\sqrt{6}\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{3}{64} \right)}{a\sqrt{ax-1}\sqrt{ax+1}}
 \end{aligned}$$

input `Int[(c - a^2*c*x^2)^(5/2)/Sqrt[ArcCosh[a*x]], x]`

---

3.402.  $\int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$

```
output (c^2*Sqrt[c - a^2*c*x^2]*((-5*Sqrt[ArcCosh[a*x]])/8 - (3*Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[a*x]]])/64 + (15*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/64 + (Sqrt[Pi/6]*Erf[Sqrt[6]*Sqrt[ArcCosh[a*x]]])/64 - (3*Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[a*x]]])/64 + (15*Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/64 + (Sqrt[Pi/6]*Erfi[Sqrt[6]*Sqrt[ArcCosh[a*x]]])/64)/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x])
```

### 3.402.3.1 Defintions of rubi rules used

```
rule 25 Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]
```

```
rule 2009 Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 3042 Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]
```

```
rule 3793 Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

```
rule 6321 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]
```

### 3.402.4 Maple [F]

$$\int \frac{(-a^2cx^2 + c)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

```
input int((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x)
```

```
output int((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x)
```

**3.402.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.402.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \text{Timed out}$$

input `integrate((-a**2*c*x**2+c)**(5/2)/acosh(a*x)**(1/2),x)`

output `Timed out`

**3.402.7 Maxima [F]**

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{(-a^2 cx^2 + c)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

input `integrate((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x, algorithm="maxima")`

output `integrate((-a^2*c*x^2 + c)^(5/2)/sqrt(arccosh(a*x)), x)`

**3.402.8 Giac [F]**

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{(-a^2 cx^2 + c)^{5/2}}{\sqrt{\operatorname{arcosh}(ax)}} dx$$

input `integrate((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x, algorithm="giac")`

output `integrate((-a^2*c*x^2 + c)^(5/2)/sqrt(arccosh(a*x)), x)`

**3.402.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{(c - a^2 cx^2)^{5/2}}{\sqrt{\operatorname{acosh}(ax)}} dx$$

input `int((c - a^2*c*x^2)^(5/2)/acosh(a*x)^(1/2),x)`

output `int((c - a^2*c*x^2)^(5/2)/acosh(a*x)^(1/2), x)`

**3.403** 
$$\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

3.403.1 Optimal result . . . . . 3125  
 3.403.2 Mathematica [A] (verified) . . . . . 3126  
 3.403.3 Rubi [A] (verified) . . . . . 3126  
 3.403.4 Maple [F] . . . . . 3128  
 3.403.5 Fricas [F(-2)] . . . . . 3128  
 3.403.6 Sympy [F] . . . . . 3128  
 3.403.7 Maxima [F] . . . . . 3129  
 3.403.8 Giac [F] . . . . . 3129  
 3.403.9 Mupad [F(-1)] . . . . . 3129

**3.403.1 Optimal result**

Integrand size = 24, antiderivative size = 294

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = -\frac{3c\sqrt{c - a^2 cx^2}\sqrt{\operatorname{arccosh}(ax)}}{4a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$- \frac{c\sqrt{\pi}\sqrt{c - a^2 cx^2}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{32a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{c\sqrt{\frac{\pi}{2}}\sqrt{c - a^2 cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{4a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

$$- \frac{c\sqrt{\pi}\sqrt{c - a^2 cx^2}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{32a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{c\sqrt{\frac{\pi}{2}}\sqrt{c - a^2 cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{4a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

output

```
1/8*c*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)
)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/8*c*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(
1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/32*c*er
f(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x
+1)^(1/2)-1/32*c*erfi(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/
a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-3/4*c*(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2
)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)
```

**3.403.2 Mathematica [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.52

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx =$$

$$\frac{c\sqrt{c - a^2 cx^2} \left( \sqrt{-\operatorname{arccosh}(ax)} \Gamma\left(\frac{1}{2}, -4\operatorname{arccosh}(ax)\right) - 4\sqrt{2} \sqrt{-\operatorname{arccosh}(ax)} \Gamma\left(\frac{1}{2}, -2\operatorname{arccosh}(ax)\right) + \sqrt{\operatorname{arccosh}(ax)} \Gamma\left(\frac{1}{2}, -\operatorname{arccosh}(ax)\right) \right)}{32a \sqrt{\frac{-1+ax}{1+ax}} (1+ax) \sqrt{\operatorname{arccosh}(ax)}}$$

input `Integrate[(c - a^2*c*x^2)^(3/2)/Sqrt[ArcCosh[a*x]], x]`output `-1/32*(c*Sqrt[c - a^2*c*x^2]*(Sqrt[-ArcCosh[a*x]]*Gamma[1/2, -4*ArcCosh[a*x]] - 4*Sqrt[2]*Sqrt[-ArcCosh[a*x]]*Gamma[1/2, -2*ArcCosh[a*x]] + Sqrt[ArcCosh[a*x]]*(24*Sqrt[ArcCosh[a*x]] + 4*Sqrt[2]*Gamma[1/2, 2*ArcCosh[a*x]] - Gamma[1/2, 4*ArcCosh[a*x]])))/(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])`**3.403.3 Rubi [A] (verified)**Time = 0.47 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.50, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {6321, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

$$\downarrow \text{6321}$$

$$\frac{c\sqrt{c - a^2 cx^2} \int \frac{(ax-1)^2(ax+1)^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}}$$

$$\downarrow \text{3042}$$

$$\frac{c\sqrt{c - a^2 cx^2} \int \frac{\sin(i\operatorname{arccosh}(ax))^4}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}}$$

$$\downarrow \text{3793}$$

---

3.403.  $\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$

$$\frac{c\sqrt{c-a^2cx^2} \int \left( -\frac{\cosh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} + \frac{\cosh(4\operatorname{arccosh}(ax))}{8\sqrt{\operatorname{arccosh}(ax)}} + \frac{3}{8\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}}$$

↓ 2009

$$\frac{c\sqrt{c-a^2cx^2} \left( \frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) \right)}{a\sqrt{ax-1}\sqrt{ax+1}}$$

input `Int[(c - a^2*c*x^2)^(3/2)/Sqrt[ArcCosh[a*x]],x]`

output `-((c*Sqrt[c - a^2*c*x^2]*((3*Sqrt[ArcCosh[a*x]])/4 + (Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[a*x]]])/32 - (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/4 + (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[a*x]]])/32 - (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/4))/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x]))`

### 3.403.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`



**3.403.4 Maple [F]**

$$\int \frac{(-a^2cx^2 + c)^{\frac{3}{2}}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

input `int((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x)`

output `int((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x)`

**3.403.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{(c - a^2cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.403.6 Sympy [F]**

$$\int \frac{(c - a^2cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{(-c(ax - 1)(ax + 1))^{\frac{3}{2}}}{\sqrt{\operatorname{acosh}(ax)}} dx$$

input `integrate((-a**2*c*x**2+c)**(3/2)/acosh(a*x)**(1/2),x)`

output `Integral((-c*(a*x - 1)*(a*x + 1))**(3/2)/sqrt(acosh(a*x)), x)`

**3.403.7 Maxima [F]**

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{(-a^2 cx^2 + c)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x, algorithm="maxima")`

output `integrate((-a^2*c*x^2 + c)^(3/2)/sqrt(arccosh(a*x)), x)`

**3.403.8 Giac [F]**

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{(-a^2 cx^2 + c)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x, algorithm="giac")`

output `integrate((-a^2*c*x^2 + c)^(3/2)/sqrt(arccosh(a*x)), x)`

**3.403.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{(c - a^2 cx^2)^{3/2}}{\sqrt{\operatorname{acosh}(ax)}} dx$$

input `int((c - a^2*c*x^2)^(3/2)/acosh(a*x)^(1/2),x)`

output `int((c - a^2*c*x^2)^(3/2)/acosh(a*x)^(1/2), x)`

**3.404**  $\int \frac{\sqrt{c-a^2cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx$

3.404.1 Optimal result . . . . . 3130  
 3.404.2 Mathematica [A] (verified) . . . . . 3130  
 3.404.3 Rubi [A] (verified) . . . . . 3131  
 3.404.4 Maple [F] . . . . . 3133  
 3.404.5 Fracas [F(-2)] . . . . . 3133  
 3.404.6 Sympy [F] . . . . . 3133  
 3.404.7 Maxima [F] . . . . . 3134  
 3.404.8 Giac [F] . . . . . 3134  
 3.404.9 Mupad [F(-1)] . . . . . 3134

**3.404.1 Optimal result**

Integrand size = 24, antiderivative size = 175

$$\int \frac{\sqrt{c-a^2cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = -\frac{\sqrt{c-a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}}{a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{4a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{4a\sqrt{-1+ax}\sqrt{1+ax}}$$

output `1/8*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/8*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-(-a^2*c*x^2+c)^(1/2)*arccosh(a*x)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)`

**3.404.2 Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.65

$$\int \frac{\sqrt{c-a^2cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \frac{\sqrt{-c(-1+ax)(1+ax)}\left(8\operatorname{arccosh}(ax) - \sqrt{2}\sqrt{-\operatorname{arccosh}(ax)}\Gamma\left(\frac{1}{2}, -2\operatorname{arccosh}(ax)\right) + \sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{8a\sqrt{\frac{-1+ax}{1+ax}}(1+ax)\sqrt{\operatorname{arccosh}(ax)}}$$

input `Integrate[Sqrt[c - a^2*c*x^2]/Sqrt[ArcCosh[a*x]],x]`

output `-1/8*(Sqrt[-(c*(-1 + a*x)*(1 + a*x))]*(8*ArcCosh[a*x] - Sqrt[2]*Sqrt[-ArcCosh[a*x]]*Gamma[1/2, -2*ArcCosh[a*x]] + Sqrt[2]*Sqrt[ArcCosh[a*x]]*Gamma[1/2, 2*ArcCosh[a*x]]))/(a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])`

### 3.404.3 Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.59, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$ , Rules used = {6321, 3042, 25, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c - a^2 cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx \\
 & \quad \downarrow \text{6321} \\
 & \frac{\sqrt{c - a^2 cx^2} \int \frac{(ax-1)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\sqrt{c - a^2 cx^2} \int -\frac{\sin(i\operatorname{arccosh}(ax))^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{25} \\
 & -\frac{\sqrt{c - a^2 cx^2} \int \frac{\sin(i\operatorname{arccosh}(ax))^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{3793} \\
 & -\frac{\sqrt{c - a^2 cx^2} \int \left( \frac{1}{2\sqrt{\operatorname{arccosh}(ax)}} - \frac{\cosh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\frac{\sqrt{c - a^2 cx^2} \left( \frac{1}{4} \sqrt{\frac{\pi}{2}} \operatorname{erf} \left( \sqrt{2} \sqrt{\operatorname{arccosh}(ax)} \right) + \frac{1}{4} \sqrt{\frac{\pi}{2}} \operatorname{erfi} \left( \sqrt{2} \sqrt{\operatorname{arccosh}(ax)} \right) - \sqrt{\operatorname{arccosh}(ax)} \right)}{a \sqrt{ax - 1} \sqrt{ax + 1}}$$

input `Int[Sqrt[c - a^2*c*x^2]/Sqrt[ArcCosh[a*x]],x]`

output `(Sqrt[c - a^2*c*x^2]*(-Sqrt[ArcCosh[a*x]] + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/4 + (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/4))/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`

### 3.404.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

**3.404.4 Maple [F]**

$$\int \frac{\sqrt{-a^2cx^2 + c}}{\sqrt{\operatorname{arccosh}(ax)}} dx$$

input `int((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x)`

output `int((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x)`

**3.404.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{c - a^2cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.404.6 Sympy [F]**

$$\int \frac{\sqrt{c - a^2cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{\sqrt{-c(ax - 1)(ax + 1)}}{\sqrt{\operatorname{acosh}(ax)}} dx$$

input `integrate((-a**2*c*x**2+c)**(1/2)/acosh(a*x)**(1/2),x)`

output `Integral(sqrt(-c*(a*x - 1)*(a*x + 1))/sqrt(acosh(a*x)), x)`

**3.404.7 Maxima [F]**

$$\int \frac{\sqrt{c - a^2 cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{\sqrt{-a^2 cx^2 + c}}{\sqrt{\operatorname{arcosh}(ax)}} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-a^2*c*x^2 + c)/sqrt(arccosh(a*x)), x)`

**3.404.8 Giac [F]**

$$\int \frac{\sqrt{c - a^2 cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{\sqrt{-a^2 cx^2 + c}}{\sqrt{\operatorname{arcosh}(ax)}} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-a^2*c*x^2 + c)/sqrt(arccosh(a*x)), x)`

**3.404.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{c - a^2 cx^2}}{\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{\sqrt{c - a^2 cx^2}}{\sqrt{\operatorname{acosh}(ax)}} dx$$

input `int((c - a^2*c*x^2)^(1/2)/acosh(a*x)^(1/2),x)`

output `int((c - a^2*c*x^2)^(1/2)/acosh(a*x)^(1/2), x)`

**3.405**  $\int \frac{1}{\sqrt{c-a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}} dx$

3.405.1 Optimal result . . . . . 3135  
 3.405.2 Mathematica [A] (verified) . . . . . 3135  
 3.405.3 Rubi [A] (verified) . . . . . 3136  
 3.405.4 Maple [A] (verified) . . . . . 3136  
 3.405.5 Fricas [F(-2)] . . . . . 3137  
 3.405.6 Sympy [F] . . . . . 3137  
 3.405.7 Maxima [F] . . . . . 3137  
 3.405.8 Giac [F] . . . . . 3138  
 3.405.9 Mupad [F(-1)] . . . . . 3138

**3.405.1 Optimal result**

Integrand size = 24, antiderivative size = 46

$$\int \frac{1}{\sqrt{c-a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\sqrt{\operatorname{arccosh}(ax)}}{a\sqrt{c-a^2cx^2}}$$

output `2*(a*x-1)^(1/2)*(a*x+1)^(1/2)*arccosh(a*x)^(1/2)/a/(-a^2*c*x^2+c)^(1/2)`

**3.405.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{c-a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}} dx = \frac{2\sqrt{-1+ax}\sqrt{1+ax}\sqrt{\operatorname{arccosh}(ax)}}{a\sqrt{c-a^2cx^2}}$$

input `Integrate[1/(Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]]),x]`

output `(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*Sqrt[ArcCosh[a*x]])/(a*Sqrt[c - a^2*c*x^2])`



### 3.405.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{\operatorname{arccosh}(ax)}\sqrt{c-a^2cx^2}} dx$$

↓ 6307

$$\frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}{a\sqrt{c-a^2cx^2}}$$

input `Int[1/(Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]]),x]`

output `(2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*Sqrt[ArcCosh[a*x]])/(a*Sqrt[c - a^2*c*x^2])`

#### 3.405.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.405.4 Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{\operatorname{arccosh}(ax)}}{\sqrt{-c(ax-1)(ax+1)}a}$	41

input `int(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x,method=_RETURNVERBOSE)`

output  $2/(-c*(a*x-1)*(a*x+1))^{(1/2)*(a*x-1)^{(1/2)*(a*x+1)^{(1/2)*\operatorname{arccosh}(a*x)^{(1/2)}}$   
)/a

### 3.405.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{\sqrt{c - a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

### 3.405.6 Sympy [F]

$$\int \frac{1}{\sqrt{c - a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{\sqrt{-c(ax-1)(ax+1)}\sqrt{\operatorname{acosh}(ax)}} dx$$

input `integrate(1/(-a**2*c*x**2+c)**(1/2)/acosh(a*x)**(1/2),x)`

output `Integral(1/(sqrt(-c*(a*x - 1)*(a*x + 1))*sqrt(acosh(a*x))), x)`

### 3.405.7 Maxima [F]

$$\int \frac{1}{\sqrt{c - a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{\sqrt{-a^2cx^2 + c}\sqrt{\operatorname{arccosh}(ax)}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-a^2*c*x^2 + c)*sqrt(arccosh(a*x))), x)`

**3.405.8 Giac [F]**

$$\int \frac{1}{\sqrt{c - a^2 c x^2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{\sqrt{-a^2 c x^2 + c} \sqrt{\operatorname{arccosh}(ax)}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-a^2*c*x^2 + c)*sqrt(arccosh(a*x))), x)`

**3.405.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{c - a^2 c x^2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{\sqrt{\operatorname{acosh}(ax)} \sqrt{c - a^2 c x^2}} dx$$

input `int(1/(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(1/2)),x)`

output `int(1/(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(1/2)), x)`

$$3.406 \quad \int \frac{1}{(c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx$$

3.406.1 Optimal result	3139
3.406.2 Mathematica [N/A]	3139
3.406.3 Rubi [N/A]	3140
3.406.4 Maple [N/A] (verified)	3140
3.406.5 Fricas [F(-2)]	3141
3.406.6 Sympy [N/A]	3141
3.406.7 Maxima [N/A]	3141
3.406.8 Giac [N/A]	3142
3.406.9 Mupad [N/A]	3142

### 3.406.1 Optimal result

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{(c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \operatorname{Int}\left(\frac{1}{(c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}}, x\right)$$

output `Unintegrable(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2), x)`

### 3.406.2 Mathematica [N/A]

Not integrable

Time = 5.50 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{(c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{(c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx$$

input `Integrate[1/((c - a^2*c*x^2)^(3/2)*Sqrt[ArcCosh[a*x]]), x]`

output `Integrate[1/((c - a^2*c*x^2)^(3/2)*Sqrt[ArcCosh[a*x]]), x]`

**3.406.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{\operatorname{arccosh}(ax)} (c - a^2 cx^2)^{3/2}} dx$$

↓ 6325

$$\int \frac{1}{\sqrt{\operatorname{arccosh}(ax)} (c - a^2 cx^2)^{3/2}} dx$$

input `Int[1/((c - a^2*c*x^2)^(3/2)*Sqrt[ArcCosh[a*x]]),x]`

output `$Aborted`

**3.406.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.406.4 Maple [N/A] (verified)**

Not integrable

Time = 1.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-a^2 cx^2 + c)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx$$

input `int(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x)`

output `int(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x)`

**3.406.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.406.6 Sympy [N/A]**

Not integrable

Time = 15.14 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{(-c(ax - 1)(ax + 1))^{\frac{3}{2}} \sqrt{\operatorname{acosh}(ax)}} dx$$

input `integrate(1/(-a**2*c*x**2+c)**(3/2)/acosh(a*x)**(1/2),x)`

output `Integral(1/((-c*(a*x - 1)*(a*x + 1))**(3/2)*sqrt(acosh(a*x))), x)`

**3.406.7 Maxima [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{(-a^2cx^2 + c)^{\frac{3}{2}} \sqrt{\operatorname{arcosh}(ax)}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x, algorithm="maxima")`

output `integrate(1/((-a^2*c*x^2 + c)^(3/2)*sqrt(arccosh(a*x))), x)`

---

3.406.  $\int \frac{1}{(c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx$

**3.406.8 Giac [N/A]**

Not integrable

Time = 2.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{(-a^2 cx^2 + c)^{3/2} \sqrt{\operatorname{arcosh}(ax)}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2),x, algorithm="giac")`output `integrate(1/((-a^2*c*x^2 + c)^(3/2)*sqrt(arccosh(a*x))), x)`**3.406.9 Mupad [N/A]**

Not integrable

Time = 2.79 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{\sqrt{\operatorname{acosh}(ax)} (c - a^2 cx^2)^{3/2}} dx$$

input `int(1/(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(3/2)),x)`output `int(1/(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(3/2)), x)`

**3.407**  $\int \frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx$

3.407.1 Optimal result . . . . . 3143  
 3.407.2 Mathematica [N/A] . . . . . 3143  
 3.407.3 Rubi [N/A] . . . . . 3144  
 3.407.4 Maple [N/A] (verified) . . . . . 3144  
 3.407.5 Fricas [F(-2)] . . . . . 3145  
 3.407.6 Sympy [F(-1)] . . . . . 3145  
 3.407.7 Maxima [N/A] . . . . . 3145  
 3.407.8 Giac [N/A] . . . . . 3146  
 3.407.9 Mupad [N/A] . . . . . 3146

**3.407.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \operatorname{Int}\left(\frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}}, x\right)$$

output `Unintegrable(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2), x)`

**3.407.2 Mathematica [N/A]**

Not integrable

Time = 5.84 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx$$

input `Integrate[1/((c - a^2*c*x^2)^(5/2)*Sqrt[ArcCosh[a*x]]), x]`

output `Integrate[1/((c - a^2*c*x^2)^(5/2)*Sqrt[ArcCosh[a*x]]), x]`



**3.407.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{\operatorname{arccosh}(ax)} (c - a^2 cx^2)^{5/2}} dx$$

↓ 6325

$$\int \frac{1}{\sqrt{\operatorname{arccosh}(ax)} (c - a^2 cx^2)^{5/2}} dx$$

input `Int[1/((c - a^2*c*x^2)^(5/2)*Sqrt[ArcCosh[a*x]]),x]`

output `$Aborted`

**3.407.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.407.4 Maple [N/A] (verified)**

Not integrable

Time = 1.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-a^2 cx^2 + c)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx$$

input `int(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x)`

output `int(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x)`

**3.407.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.407.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \text{Timed out}$$

input `integrate(1/(-a**2*c*x**2+c)**(5/2)/acosh(a*x)**(1/2),x)`

output `Timed out`

**3.407.7 Maxima [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{(-a^2cx^2 + c)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x, algorithm="maxima")`

output `integrate(1/((-a^2*c*x^2 + c)^(5/2)*sqrt(arccosh(a*x))), x)`

**3.407.8 Giac [N/A]**

Not integrable

Time = 2.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{(-a^2 cx^2 + c)^{5/2} \sqrt{\operatorname{arcosh}(ax)}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2),x, algorithm="giac")`output `integrate(1/((-a^2*c*x^2 + c)^(5/2)*sqrt(arccosh(a*x))), x)`**3.407.9 Mupad [N/A]**

Not integrable

Time = 2.88 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} dx = \int \frac{1}{\sqrt{\operatorname{acosh}(ax)} (c - a^2 cx^2)^{5/2}} dx$$

input `int(1/(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(5/2)),x)`output `int(1/(acosh(a*x)^(1/2)*(c - a^2*c*x^2)^(5/2)), x)`

**3.408**       $\int \frac{(c-a^2cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$

3.408.1 Optimal result . . . . . 3147  
 3.408.2 Mathematica [A] (verified) . . . . . 3148  
 3.408.3 Rubi [A] (verified) . . . . . 3149  
 3.408.4 Maple [F] . . . . . 3151  
 3.408.5 Fricas [F(-2)] . . . . . 3151  
 3.408.6 Sympy [F(-1)] . . . . . 3151  
 3.408.7 Maxima [F] . . . . . 3152  
 3.408.8 Giac [F] . . . . . 3152  
 3.408.9 Mupad [F(-1)] . . . . . 3152

**3.408.1 Optimal result**

Integrand size = 24, antiderivative size = 433

$$\int \frac{(c-a^2cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}(c-a^2cx^2)^{5/2}}{a\sqrt{\operatorname{arccosh}(ax)}} + \frac{3c^2\sqrt{\pi}\sqrt{c-a^2cx^2}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{8a\sqrt{-1+ax}\sqrt{1+ax}} - \frac{15c^2\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1+ax}\sqrt{1+ax}} - \frac{c^2\sqrt{\frac{3\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erf}\left(\sqrt{6}\sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1+ax}\sqrt{1+ax}} - \frac{3c^2\sqrt{\pi}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{8a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{15c^2\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{c^2\sqrt{\frac{3\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(\sqrt{6}\sqrt{\operatorname{arccosh}(ax)}\right)}{16a\sqrt{-1+ax}\sqrt{1+ax}}$$

```
output -15/32*c^2*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)
^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+15/32*c^2*erfi(2^(1/2)*arccosh(a*x)^(
1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+
3/8*c^2*erf(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(
1/2)/(a*x+1)^(1/2)-3/8*c^2*erfi(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^
2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/32*c^2*erf(6^(1/2)*arccosh(a*x)
^(1/2))*6^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2
)+1/32*c^2*erfi(6^(1/2)*arccosh(a*x)^(1/2))*6^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c
)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-2*(-a^2*c*x^2+c)^(5/2)*(a*x-1)^(1/2)
*(a*x+1)^(1/2)/a/arccosh(a*x)^(1/2)
```

### 3.408.2 Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 411, normalized size of antiderivative = 0.95

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \frac{c^2 e^{-6 \operatorname{arccosh}(ax)} \sqrt{c - a^2 cx^2} \left( -1 + 6e^{2 \operatorname{arccosh}(ax)} + e^{4 \operatorname{arccosh}(ax)} + 52e^{6 \operatorname{arccosh}(ax)} + e^{8 \operatorname{arccosh}(ax)} \right)}{\dots}$$

```
input Integrate[(c - a^2*c*x^2)^(5/2)/ArcCosh[a*x]^(3/2), x]
```

```
output (c^2*Sqrt[c - a^2*c*x^2]*(-1 + 6*E^(2*ArcCosh[a*x]) + E^(4*ArcCosh[a*x]) +
52*E^(6*ArcCosh[a*x]) + E^(8*ArcCosh[a*x]) + 6*E^(10*ArcCosh[a*x]) - E^(1
2*ArcCosh[a*x]) - 64*a^2*E^(6*ArcCosh[a*x])*x^2 - 16*E^(6*ArcCosh[a*x])*Sq
rt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + 16*E^(6*ArcC
osh[a*x])*Sqrt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]] +
Sqrt[6]*E^(6*ArcCosh[a*x])*Sqrt[-ArcCosh[a*x]]*Gamma[1/2, -6*ArcCosh[a*x]
] - 12*E^(6*ArcCosh[a*x])*Sqrt[-ArcCosh[a*x]]*Gamma[1/2, -4*ArcCosh[a*x]]
- Sqrt[2]*E^(6*ArcCosh[a*x])*Sqrt[-ArcCosh[a*x]]*Gamma[1/2, -2*ArcCosh[a*x
]] - Sqrt[2]*E^(6*ArcCosh[a*x])*Sqrt[ArcCosh[a*x]]*Gamma[1/2, 2*ArcCosh[a*
x]] - 12*E^(6*ArcCosh[a*x])*Sqrt[ArcCosh[a*x]]*Gamma[1/2, 4*ArcCosh[a*x]]
+ Sqrt[6]*E^(6*ArcCosh[a*x])*Sqrt[ArcCosh[a*x]]*Gamma[1/2, 6*ArcCosh[a*x]]
)/(32*a*E^(6*ArcCosh[a*x])*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcC
osh[a*x]])
```

**3.408.3 Rubi [A] (verified)**

Time = 0.84 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.56, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$ , Rules used = {6319, 6327, 6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(c - a^2 cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{12ac^2 \sqrt{c - a^2 cx^2} \int \frac{x(1-ax)^2(ax+1)^2}{\sqrt{\operatorname{arccosh}(ax)}} dx}{\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c - a^2 cx^2)^{5/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{12ac^2 \sqrt{c - a^2 cx^2} \int \frac{x(1-a^2 x^2)^2}{\sqrt{\operatorname{arccosh}(ax)}} dx}{\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c - a^2 cx^2)^{5/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{6367} \\
 & \frac{12c^2 \sqrt{c - a^2 cx^2} \int \frac{ax \left(\frac{ax-1}{ax+1}\right)^{5/2} (ax+1)^5}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c - a^2 cx^2)^{5/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{5971} \\
 & \frac{12c^2 \sqrt{c - a^2 cx^2} \int \left( \frac{5 \sinh(2\operatorname{arccosh}(ax))}{32\sqrt{\operatorname{arccosh}(ax)}} - \frac{\sinh(4\operatorname{arccosh}(ax))}{8\sqrt{\operatorname{arccosh}(ax)}} + \frac{\sinh(6\operatorname{arccosh}(ax))}{32\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c - a^2 cx^2)^{5/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{12c^2 \sqrt{c - a^2 cx^2} \left( \frac{1}{32} \sqrt{\pi} \operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{5}{64} \sqrt{\frac{\pi}{2}} \operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{64} \sqrt{\frac{\pi}{6}} \operatorname{erf}\left(\sqrt{6}\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{32} \right)}{a\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c - a^2 cx^2)^{5/2}}{a\sqrt{\operatorname{arccosh}(ax)}}
 \end{aligned}$$

---

3.408.  $\int \frac{(c - a^2 cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$

input `Int[(c - a^2*c*x^2)^(5/2)/ArcCosh[a*x]^(3/2),x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(c - a^2*c*x^2)^(5/2))/(a*Sqrt[ArcCosh[a*x]]) + (12*c^2*Sqrt[c - a^2*c*x^2]*((Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[a*x]]])/32 - (5*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/64 - (Sqrt[Pi/6]*Erf[Sqrt[6]*Sqrt[ArcCosh[a*x]]])/64 - (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[a*x]]])/32 + (5*Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/64 + (Sqrt[Pi/6]*Erfi[Sqrt[6]*Sqrt[ArcCosh[a*x]]])/64))/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`

### 3.408.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_.) + (e1_.)*(x_))^(p_.)*((d2_.) + (e2_.)*(x_))^(q_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

---

3.408.  $\int \frac{(c-a^2cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$

**3.408.4 Maple [F]**

$$\int \frac{(-a^2cx^2 + c)^{\frac{5}{2}}}{\operatorname{arccosh}(ax)^{\frac{3}{2}}} dx$$

input `int((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x)`

output `int((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x)`

**3.408.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{(c - a^2cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.408.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(c - a^2cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \text{Timed out}$$

input `integrate((-a**2*c*x**2+c)**(5/2)/acosh(a*x)**(3/2),x)`

output `Timed out`



**3.408.7 Maxima [F]**

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{(-a^2 cx^2 + c)^{5/2}}{\operatorname{arcosh}(ax)^{3/2}} dx$$

input `integrate((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x, algorithm="maxima")`

output `integrate((-a^2*c*x^2 + c)^(5/2)/arccosh(a*x)^(3/2), x)`

**3.408.8 Giac [F]**

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{(-a^2 cx^2 + c)^{5/2}}{\operatorname{arcosh}(ax)^{3/2}} dx$$

input `integrate((-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x, algorithm="giac")`

output `integrate((-a^2*c*x^2 + c)^(5/2)/arccosh(a*x)^(3/2), x)`

**3.408.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^{5/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{(c - a^2 cx^2)^{5/2}}{\operatorname{acosh}(ax)^{3/2}} dx$$

input `int((c - a^2*c*x^2)^(5/2)/acosh(a*x)^(3/2),x)`

output `int((c - a^2*c*x^2)^(5/2)/acosh(a*x)^(3/2), x)`

**3.409**  $\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$

3.409.1 Optimal result . . . . . 3153  
 3.409.2 Mathematica [A] (verified) . . . . . 3154  
 3.409.3 Rubi [A] (verified) . . . . . 3154  
 3.409.4 Maple [F] . . . . . 3157  
 3.409.5 Fricas [F(-2)] . . . . . 3157  
 3.409.6 Sympy [F] . . . . . 3157  
 3.409.7 Maxima [F] . . . . . 3158  
 3.409.8 Giac [F] . . . . . 3158  
 3.409.9 Mupad [F(-1)] . . . . . 3158

**3.409.1 Optimal result**

Integrand size = 24, antiderivative size = 286

$$\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}(c-a^2cx^2)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} + \frac{c\sqrt{\pi}\sqrt{c-a^2cx^2}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{4a\sqrt{-1+ax}\sqrt{1+ax}} - \frac{c\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{a\sqrt{-1+ax}\sqrt{1+ax}} - \frac{c\sqrt{\pi}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{4a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{c\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{a\sqrt{-1+ax}\sqrt{1+ax}}$$

output

```
-1/2*c*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/2*c*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+1/4*c*erf(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-1/4*c*erfi(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-2*(-a^2*c*x^2+c)^(3/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/arccosh(a*x)^(1/2)
```

**3.409.2 Mathematica [A] (verified)**

Time = 0.45 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.84

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx =$$


---


$$ce^{-4\operatorname{arccosh}(ax)} \sqrt{c - a^2 cx^2} \left( -1 - 14e^{4\operatorname{arccosh}(ax)} - e^{8\operatorname{arccosh}(ax)} + 16a^2 e^{4\operatorname{arccosh}(ax)} x^2 + 4e^{4\operatorname{arccosh}(ax)} \sqrt{2\pi} \sqrt{\operatorname{arccosh}(ax)} \right)$$


---

input `Integrate[(c - a^2*c*x^2)^(3/2)/ArcCosh[a*x]^(3/2),x]`output

```

-1/8*(c*Sqrt[c - a^2*c*x^2]*(-1 - 14*E^(4*ArcCosh[a*x]) - E^(8*ArcCosh[a*x]
]) + 16*a^2*E^(4*ArcCosh[a*x])*x^2 + 4*E^(4*ArcCosh[a*x])*Sqrt[2*Pi]*Sqrt[
ArcCosh[a*x]]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] - 4*E^(4*ArcCosh[a*x])*Sqrt[
2*Pi]*Sqrt[ArcCosh[a*x]]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + 2*E^(4*ArcCosh
[a*x])*Sqrt[-ArcCosh[a*x]]*Gamma[1/2, -4*ArcCosh[a*x]] + 2*E^(4*ArcCosh[a*
x])*Sqrt[ArcCosh[a*x]]*Gamma[1/2, 4*ArcCosh[a*x]]))/(a*E^(4*ArcCosh[a*x])*
Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])

```

**3.409.3 Rubi [A] (verified)**Time = 0.70 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.64, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6319, 25, 6327, 6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$$

$$\downarrow \text{6319}$$

$$\frac{8ac\sqrt{c - a^2 cx^2} \int -\frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}(c - a^2 cx^2)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}}$$

$$\downarrow \text{25}$$

---

3.409.  $\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx$

$$\begin{aligned}
& \frac{8ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
& \quad \downarrow \text{6327} \\
& \frac{8ac\sqrt{c-a^2cx^2} \int \frac{x(1-a^2x^2)}{\sqrt{\operatorname{arccosh}(ax)}} dx}{\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
& \quad \downarrow \text{6367} \\
& \frac{8c\sqrt{c-a^2cx^2} \int \frac{ax\left(\frac{ax-1}{ax+1}\right)^{3/2} (ax+1)^3}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
& \quad \downarrow \text{5971} \\
& \frac{8c\sqrt{c-a^2cx^2} \int \left( \frac{\sinh(4\operatorname{arccosh}(ax))}{8\sqrt{\operatorname{arccosh}(ax)}} - \frac{\sinh(2\operatorname{arccosh}(ax))}{4\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
& \quad \downarrow \text{2009} \\
& \frac{8c\sqrt{c-a^2cx^2} \left( -\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{8}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{8}\sqrt{\pi}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) \right)}{a\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}}
\end{aligned}$$

input `Int[(c - a^2*c*x^2)^(3/2)/ArcCosh[a*x]^(3/2), x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(c - a^2*c*x^2)^(3/2))/(a*Sqrt[ArcCosh[a*x]]) - (8*c*Sqrt[c - a^2*c*x^2]*(-1/32*(Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[a*x]]]) + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/8 + (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[a*x]]])/32 - (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/8))/(a*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`

## 3.409.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`
- rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1))]*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

**3.409.4 Maple [F]**

$$\int \frac{(-a^2 c x^2 + c)^{\frac{3}{2}}}{\operatorname{arccosh}(ax)^{\frac{3}{2}}} dx$$

input `int((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x)`

output `int((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x)`

**3.409.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{(c - a^2 c x^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.409.6 Sympy [F]**

$$\int \frac{(c - a^2 c x^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{(-c(ax - 1)(ax + 1))^{\frac{3}{2}}}{\operatorname{acosh}^{\frac{3}{2}}(ax)} dx$$

input `integrate((-a**2*c*x**2+c)**(3/2)/acosh(a*x)**(3/2),x)`

output `Integral((-c*(a*x - 1)*(a*x + 1))**(3/2)/acosh(a*x)**(3/2), x)`

**3.409.7 Maxima [F]**

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{(-a^2 cx^2 + c)^{3/2}}{\operatorname{arcosh}(ax)^{3/2}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x, algorithm="maxima")`

output `integrate((-a^2*c*x^2 + c)^(3/2)/arccosh(a*x)^(3/2), x)`

**3.409.8 Giac [F]**

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{(-a^2 cx^2 + c)^{3/2}}{\operatorname{arcosh}(ax)^{3/2}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x, algorithm="giac")`

output `integrate((-a^2*c*x^2 + c)^(3/2)/arccosh(a*x)^(3/2), x)`

**3.409.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{acosh}(ax)^{3/2}} dx$$

input `int((c - a^2*c*x^2)^(3/2)/acosh(a*x)^(3/2),x)`

output `int((c - a^2*c*x^2)^(3/2)/acosh(a*x)^(3/2), x)`

### 3.410 $\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx$

3.410.1 Optimal result . . . . .	3159
3.410.2 Mathematica [A] (verified) . . . . .	3159
3.410.3 Rubi [C] (verified) . . . . .	3160
3.410.4 Maple [F] . . . . .	3163
3.410.5 Fracas [F(-2)] . . . . .	3163
3.410.6 Sympy [F] . . . . .	3164
3.410.7 Maxima [F] . . . . .	3164
3.410.8 Giac [F] . . . . .	3164
3.410.9 Mupad [F(-1)] . . . . .	3165

#### 3.410.1 Optimal result

Integrand size = 24, antiderivative size = 170

$$\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}\sqrt{c-a^2cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} - \frac{\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{\sqrt{\frac{\pi}{2}}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{a\sqrt{-1+ax}\sqrt{1+ax}}$$

output 
$$\begin{aligned} & -1/2*\operatorname{erf}\left(2^{(1/2)}*\operatorname{arccosh}(a*x)^{(1/2)}\right)*2^{(1/2)}*\operatorname{Pi}^{(1/2)}*(-a^2*c*x^2+c)^{(1/2)} \\ & /a/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}+1/2*\operatorname{erfi}\left(2^{(1/2)}*\operatorname{arccosh}(a*x)^{(1/2)}\right)*2^{(1/2)} \\ & )*\operatorname{Pi}^{(1/2)}*(-a^2*c*x^2+c)^{(1/2)}/a/(a*x-1)^{(1/2)}/(a*x+1)^{(1/2)}-2*(a*x-1)^{(1/2)} \\ & /2*(a*x+1)^{(1/2)}*(-a^2*c*x^2+c)^{(1/2)}/a/\operatorname{arccosh}(a*x)^{(1/2)} \end{aligned}$$

#### 3.410.2 Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.75

$$\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \frac{\sqrt{c-a^2cx^2}\left(4-4a^2x^2-\sqrt{2\pi}\sqrt{\operatorname{arccosh}(ax)}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)+\sqrt{2\pi}\sqrt{\operatorname{arccosh}(ax)}\right)}{2a\sqrt{\frac{-1+ax}{1+ax}}(1+ax)\sqrt{\operatorname{arccosh}(ax)}}$$

input `Integrate[Sqrt[c - a^2*c*x^2]/ArcCosh[a*x]^(3/2),x]`



```
output (Sqrt[c - a^2*c*x^2]*(4 - 4*a^2*x^2 - Sqrt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]] + Sqrt[2*Pi]*Sqrt[ArcCosh[a*x]]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]]))/(2*a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*Sqrt[ArcCosh[a*x]])
```

### 3.410.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.65 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.87, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$ , Rules used = {6319, 6302, 5971, 27, 3042, 26, 3789, 2611, 2633, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c - a^2 cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{4a\sqrt{c - a^2 cx^2} \int \frac{x}{\sqrt{\operatorname{arccosh}(ax)}} dx}{\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{c - a^2 cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{6302} \\
 & \frac{4\sqrt{c - a^2 cx^2} \int \frac{ax\sqrt{\frac{ax-1}{ax+1}}(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{c - a^2 cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{5971} \\
 & \frac{4\sqrt{c - a^2 cx^2} \int \frac{\sinh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{c - a^2 cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{27} \\
 & \frac{2\sqrt{c - a^2 cx^2} \int \frac{\sinh(2\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{c - a^2 cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

$$\begin{aligned}
& -\frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} + \frac{2\sqrt{c-a^2cx^2} \int -\frac{i \sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
& \quad \downarrow 26 \\
& -\frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} - \frac{2i\sqrt{c-a^2cx^2} \int \frac{\sin(2i\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
& \quad \downarrow 3789 \\
& \frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} - \frac{2i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i \int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2}i \int \frac{e^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) \right)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
& \quad \downarrow 2611 \\
& \frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} - \frac{2i\sqrt{c-a^2cx^2} \left( i \int e^{2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
& \quad \downarrow 2633 \\
& \frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} - \frac{2i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) - i \int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} \right)}{a\sqrt{ax-1}\sqrt{ax+1}} \\
& \quad \downarrow 2634 \\
& \frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{a\sqrt{\operatorname{arccosh}(ax)}} - \frac{2i\sqrt{c-a^2cx^2} \left( \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erfi} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) - \frac{1}{2}i\sqrt{\frac{\pi}{2}} \operatorname{erf} \left( \sqrt{2}\sqrt{\operatorname{arccosh}(ax)} \right) \right)}{a\sqrt{ax-1}\sqrt{ax+1}}
\end{aligned}$$

input `Int[Sqrt[c - a^2*c*x^2]/ArcCosh[a*x]^(3/2), x]`

```
output (-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*Sqrt[c - a^2*c*x^2])/(a*Sqrt[ArcCosh[a*x]]
) - ((2*I)*Sqrt[c - a^2*c*x^2]*((-1/2*I)*Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcC
osh[a*x]]] + (I/2)*Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]]))/(a*Sqrt[-
1 + a*x]*Sqrt[1 + a*x])
```

### 3.410.3.1 Defintions of rubi rules used

```
rule 26 Int[(Complex[0, a_]*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) I
nt[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]
```

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2611 Int[(F_)^((g_.)*(e_.) + (f_.)*(x_))/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] :
> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d
*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]
```

```
rule 2633 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt
[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{
F, a, b, c, d}, x] && PosQ[b]
```

```
rule 2634 Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt
[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; Fr
eeQ[{F, a, b, c, d}, x] && NegQ[b]
```

```
rule 3042 Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear
Q[u, x]
```

```
rule 3789 Int[((c_.) + (d_.)*(x_)^m)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I
/2 Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)^m*E
^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]
```

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6302 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[1/(b*c^(m + 1)) Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[m, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

### 3.410.4 Maple [F]

$$\int \frac{\sqrt{-a^2cx^2 + c}}{\operatorname{arccosh}(ax)^{\frac{3}{2}}} dx$$

input `int((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2),x)`

output `int((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2),x)`

### 3.410.5 Fracas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

---

3.410.  $\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx$

**3.410.6 Sympy [F]**

$$\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{\sqrt{-c(ax - 1)(ax + 1)}}{\operatorname{acosh}^{\frac{3}{2}}(ax)} dx$$

input `integrate((-a**2*c*x**2+c)**(1/2)/acosh(a*x)**(3/2), x)`

output `Integral(sqrt(-c*(a*x - 1)*(a*x + 1))/acosh(a*x)**(3/2), x)`

**3.410.7 Maxima [F]**

$$\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{\sqrt{-a^2cx^2 + c}}{\operatorname{arcosh}(ax)^{\frac{3}{2}}} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2), x, algorithm="maxima")`

output `integrate(sqrt(-a^2*c*x^2 + c)/arccosh(a*x)^(3/2), x)`

**3.410.8 Giac [F]**

$$\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{\sqrt{-a^2cx^2 + c}}{\operatorname{arcosh}(ax)^{\frac{3}{2}}} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2), x, algorithm="giac")`

output `integrate(sqrt(-a^2*c*x^2 + c)/arccosh(a*x)^(3/2), x)`

**3.410.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{c - a^2 c x^2}}{\operatorname{arccosh}(a x)^{3/2}} dx = \int \frac{\sqrt{c - a^2 c x^2}}{\operatorname{acosh}(a x)^{3/2}} dx$$

input `int((c - a^2*c*x^2)^(1/2)/acosh(a*x)^(3/2),x)`output `int((c - a^2*c*x^2)^(1/2)/acosh(a*x)^(3/2), x)`

**3.411**  $\int \frac{1}{\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{3/2}} dx$

3.411.1 Optimal result . . . . . 3166  
 3.411.2 Mathematica [A] (verified) . . . . . 3166  
 3.411.3 Rubi [A] (verified) . . . . . 3167  
 3.411.4 Maple [A] (verified) . . . . . 3167  
 3.411.5 Fricas [A] (verification not implemented) . . . . . 3168  
 3.411.6 Sympy [F] . . . . . 3168  
 3.411.7 Maxima [F] . . . . . 3168  
 3.411.8 Giac [F] . . . . . 3169  
 3.411.9 Mupad [F(-1)] . . . . . 3169

**3.411.1 Optimal result**

Integrand size = 24, antiderivative size = 46

$$\int \frac{1}{\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{3/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}}{a\sqrt{c-a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}}$$

output `-2*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(1/2)`

**3.411.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{3/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}}{a\sqrt{c-a^2cx^2}\sqrt{\operatorname{arccosh}(ax)}}$$

input `Integrate[1/(Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2)),x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(a*Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]])`

**3.411.3 Rubi [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arccosh}(ax)^{3/2} \sqrt{c - a^2 cx^2}} dx$$

↓ 6307

$$-\frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}\sqrt{c - a^2 cx^2}}$$

input `Int[1/(Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2)),x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(a*Sqrt[c - a^2*c*x^2]*Sqrt[ArcCosh[a*x]])`

**3.411.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.411.4 Maple [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.89

method	result	size
default	$-\frac{2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{-c(ax-1)(ax+1)}\sqrt{\operatorname{arccosh}(ax)}a}$	41

input `int(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2),x,method=_RETURNVERBOSE)`



output  $-2/(-c*(a*x-1)*(a*x+1))^{(1/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/\operatorname{arccosh}(a*x)^{(1/2)}/a$

### 3.411.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.28

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2}} dx = \frac{2\sqrt{-a^2 cx^2 + c} \sqrt{a^2 x^2 - 1}}{(a^3 cx^2 - ac) \sqrt{\log(ax + \sqrt{a^2 x^2 - 1})}}$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `2*sqrt(-a^2*c*x^2 + c)*sqrt(a^2*x^2 - 1)/((a^3*c*x^2 - a*c)*sqrt(log(a*x + sqrt(a^2*x^2 - 1))))`

### 3.411.6 Sympy [F]

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{\sqrt{-c(ax-1)(ax+1)} \operatorname{acosh}^{\frac{3}{2}}(ax)} dx$$

input `integrate(1/(-a**2*c*x**2+c)**(1/2)/acosh(a*x)**(3/2),x)`

output `Integral(1/(sqrt(-c*(a*x - 1)*(a*x + 1))*acosh(a*x)**(3/2)), x)`

### 3.411.7 Maxima [F]

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{\sqrt{-a^2 cx^2 + c} \operatorname{arcosh}(ax)^{\frac{3}{2}}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^(3/2)), x)`

**3.411.8 Giac [F]**

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{\sqrt{-a^2 cx^2 + c} \operatorname{arccosh}(ax)^{3/2}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(3/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^(3/2)), x)`

**3.411.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{\operatorname{acosh}(ax)^{3/2} \sqrt{c - a^2 cx^2}} dx$$

input `int(1/(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(1/2)),x)`

output `int(1/(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(1/2)), x)`

**3.412**  $\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx$

3.412.1 Optimal result . . . . . 3170  
 3.412.2 Mathematica [N/A] . . . . . 3170  
 3.412.3 Rubi [N/A] . . . . . 3171  
 3.412.4 Maple [N/A] (verified) . . . . . 3172  
 3.412.5 Fricas [F(-2)] . . . . . 3172  
 3.412.6 Sympy [N/A] . . . . . 3173  
 3.412.7 Maxima [N/A] . . . . . 3173  
 3.412.8 Giac [N/A] . . . . . 3173  
 3.412.9 Mupad [N/A] . . . . . 3174

**3.412.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx = -\frac{2\sqrt{-1 + ax}\sqrt{1 + ax}}{a(c - a^2cx^2)^{3/2} \sqrt{\operatorname{arccosh}(ax)}} + \frac{4a\sqrt{-1 + ax}\sqrt{1 + ax} \operatorname{Int}\left(\frac{x}{(-1 + a^2x^2)^2 \sqrt{\operatorname{arccosh}(ax)}}, x\right)}{c\sqrt{c - a^2cx^2}}$$

output `-2*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(1/2)+4*a*(a*x-1)^(1/2)*(a*x+1)^(1/2)*Unintegrateable(x/(a^2*x^2-1)^2/arccosh(a*x)^(1/2),x)/c/(-a^2*c*x^2+c)^(1/2)`

**3.412.2 Mathematica [N/A]**

Not integrable

Time = 4.76 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx$$

input `Integrate[1/((c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(3/2)), x]`

output `Integrate[1/((c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(3/2)), x]`

### 3.412.3 Rubi [N/A]

Not integrable

Time = 0.56 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{3/2}} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{4a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-ax)^2(ax+1)^2\sqrt{\operatorname{arccosh}(ax)}} dx}{c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{4a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^2\sqrt{\operatorname{arccosh}(ax)}} dx}{c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}} \\
 & \quad \downarrow \text{6375} \\
 & \frac{4a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^2\sqrt{\operatorname{arccosh}(ax)}} dx}{c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{3/2}}
 \end{aligned}$$

input `Int[1/((c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(3/2)),x]`

output `$Aborted`

#### 3.412.3.1 Defintions of rubi rules used

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

---

3.412.  $\int \frac{1}{(c-a^2cx^2)^{3/2}\operatorname{arccosh}(ax)^{3/2}} dx$

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

### 3.412.4 Maple [N/A] (verified)

Not integrable

Time = 1.40 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{3}{2}}} dx$$

input `int(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x)`

output `int(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x)`

### 3.412.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.412.6 Sympy [N/A]**

Not integrable

Time = 170.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{(-c(ax - 1)(ax + 1))^{\frac{3}{2}} \operatorname{acosh}^{\frac{3}{2}}(ax)} dx$$

input `integrate(1/(-a**2*c*x**2+c)**(3/2)/acosh(a*x)**(3/2), x)`output `Integral(1/((-c*(a*x - 1)*(a*x + 1))**(3/2)*acosh(a*x)**(3/2)), x)`**3.412.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{(-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arcosh}(ax)^{\frac{3}{2}}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2), x, algorithm="maxima")`output `integrate(1/((-a^2*c*x^2 + c)^(3/2)*arccosh(a*x)^(3/2)), x)`**3.412.8 Giac [N/A]**

Not integrable

Time = 0.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{(-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arcosh}(ax)^{\frac{3}{2}}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2), x, algorithm="giac")`output `integrate(1/((-a^2*c*x^2 + c)^(3/2)*arccosh(a*x)^(3/2)), x)`

**3.412.9 Mupad [N/A]**

Not integrable

Time = 3.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 c x^2)^{3/2} \operatorname{arccosh}(a x)^{3/2}} dx = \int \frac{1}{\operatorname{acosh}(a x)^{3/2} (c - a^2 c x^2)^{3/2}} dx$$

input `int(1/(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(3/2)),x)`output `int(1/(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(3/2)), x)`

**3.413**  $\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx$

3.413.1 Optimal result	3175
3.413.2 Mathematica [N/A]	3175
3.413.3 Rubi [N/A]	3176
3.413.4 Maple [N/A] (verified)	3177
3.413.5 Fricas [F(-2)]	3178
3.413.6 Sympy [F(-1)]	3178
3.413.7 Maxima [N/A]	3178
3.413.8 Giac [N/A]	3179
3.413.9 Mupad [N/A]	3179

**3.413.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx = -\frac{2\sqrt{-1 + ax}\sqrt{1 + ax}}{a(c - a^2cx^2)^{5/2} \sqrt{\operatorname{arccosh}(ax)}} - \frac{8a\sqrt{-1 + ax}\sqrt{1 + ax} \operatorname{Int}\left(\frac{x}{(-1 + a^2x^2)^3 \sqrt{\operatorname{arccosh}(ax)}}, x\right)}{c^2 \sqrt{c - a^2cx^2}}$$

output `-2*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(1/2)-8*a*(a*x-1)^(1/2)*(a*x+1)^(1/2)*Unintegrateable(x/(a^2*x^2-1)^3/arccosh(a*x)^(1/2),x)/c^2/(-a^2*c*x^2+c)^(1/2)`

**3.413.2 Mathematica [N/A]**

Not integrable

Time = 5.77 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx$$

input `Integrate[1/((c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^(3/2)), x]`

output `Integrate[1/((c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^(3/2)), x]`



**3.413.3 Rubi [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 25, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{8a\sqrt{ax-1}\sqrt{ax+1} \int -\frac{x}{(1-ax)^3(ax+1)^3\sqrt{\operatorname{arccosh}(ax)}} dx}{c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{8a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-ax)^3(ax+1)^3\sqrt{\operatorname{arccosh}(ax)}} dx}{c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{8a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^3\sqrt{\operatorname{arccosh}(ax)}} dx}{c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{6375} \\
 & \frac{8a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^3\sqrt{\operatorname{arccosh}(ax)}} dx}{c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}(c-a^2cx^2)^{5/2}}
 \end{aligned}$$

input `Int[1/((c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^(3/2)),x]`

output `$Aborted`

## 3.413.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_)^m_)*((d1_) + (e1_.)*(x_)^p_)*((d2_) + (e2_.)*(x_)^p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_)^m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

## 3.413.4 Maple [N/A] (verified)

Not integrable

Time = 1.58 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-a^2cx^2 + c)^{\frac{5}{2}} \operatorname{arccosh}(ax)^{\frac{3}{2}}} dx$$

input `int(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x)`

output `int(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x)`

**3.413.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.413.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx = \text{Timed out}$$

input `integrate(1/(-a**2*c*x**2+c)**(5/2)/acosh(a*x)**(3/2),x)`

output `Timed out`

**3.413.7 Maxima [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{(-a^2cx^2 + c)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x, algorithm="maxima")`

output `integrate(1/((-a^2*c*x^2 + c)^(5/2)*arccosh(a*x)^(3/2)), x)`

**3.413.8 Giac [N/A]**

Not integrable

Time = 0.39 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{(-a^2 cx^2 + c)^{5/2} \operatorname{arcosh}(ax)^{3/2}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2),x, algorithm="giac")`output `integrate(1/((-a^2*c*x^2 + c)^(5/2)*arccosh(a*x)^(3/2)), x)`**3.413.9 Mupad [N/A]**

Not integrable

Time = 2.86 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} dx = \int \frac{1}{\operatorname{acosh}(ax)^{3/2} (c - a^2 cx^2)^{5/2}} dx$$

input `int(1/(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(5/2)),x)`output `int(1/(acosh(a*x)^(3/2)*(c - a^2*c*x^2)^(5/2)), x)`

**3.414**  $\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

3.414.1 Optimal result . . . . . 3180  
 3.414.2 Mathematica [A] (warning: unable to verify) . . . . . 3181  
 3.414.3 Rubi [A] (verified) . . . . . 3181  
 3.414.4 Maple [F] . . . . . 3185  
 3.414.5 Fricas [F(-2)] . . . . . 3186  
 3.414.6 Sympy [F(-1)] . . . . . 3186  
 3.414.7 Maxima [F] . . . . . 3186  
 3.414.8 Giac [F] . . . . . 3187  
 3.414.9 Mupad [F(-1)] . . . . . 3187

**3.414.1 Optimal result**

Integrand size = 24, antiderivative size = 329

$$\int \frac{(c - a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx = -\frac{2\sqrt{-1 + ax}\sqrt{1 + ax}(c - a^2cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}} - \frac{16cx(1 - ax)(1 + ax)\sqrt{c - a^2cx^2}}{3\sqrt{\operatorname{arccosh}(ax)}} - \frac{2c\sqrt{\pi}\sqrt{c - a^2cx^2}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{3a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{2c\sqrt{2\pi}\sqrt{c - a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{3a\sqrt{-1 + ax}\sqrt{1 + ax}} - \frac{2c\sqrt{\pi}\sqrt{c - a^2cx^2}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right)}{3a\sqrt{-1 + ax}\sqrt{1 + ax}} + \frac{2c\sqrt{2\pi}\sqrt{c - a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{3a\sqrt{-1 + ax}\sqrt{1 + ax}}$$

output

```
-2/3*(-a^2*c*x^2+c)^(3/2)*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/arccosh(a*x)^(3/2)
-2/3*c*erf(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(
1/2)/(a*x+1)^(1/2)-2/3*c*erfi(2*arccosh(a*x)^(1/2))*Pi^(1/2)*(-a^2*c*x^2+c
)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+2/3*c*erf(2^(1/2)*arccosh(a*x)^(1/2)
)*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+2/3*
c*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a
/(a*x-1)^(1/2)/(a*x+1)^(1/2)-16/3*c*x*(-a*x+1)*(a*x+1)*(-a^2*c*x^2+c)^(1/2)
)/arccosh(a*x)^(1/2)
```

3.414.  $\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

**3.414.2 Mathematica [A] (warning: unable to verify)**

Time = 0.45 (sec) , antiderivative size = 317, normalized size of antiderivative = 0.96

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx =$$


---


$$ce^{-4\operatorname{arccosh}(ax)}\sqrt{c - a^2 cx^2}\left(-1 - 14e^{4\operatorname{arccosh}(ax)} - e^{8\operatorname{arccosh}(ax)} + 16a^2 e^{4\operatorname{arccosh}(ax)}x^2 + 8\operatorname{arccosh}(ax) - 8e^{8\operatorname{arccosh}(ax)}\right)$$

input `Integrate[(c - a^2*c*x^2)^(3/2)/ArcCosh[a*x]^(5/2),x]`

output

```
-1/24*(c*Sqrt[c - a^2*c*x^2]*(-1 - 14*E^(4*ArcCosh[a*x]) - E^(8*ArcCosh[a*x]) + 16*a^2*E^(4*ArcCosh[a*x])*x^2 + 8*ArcCosh[a*x] - 8*E^(8*ArcCosh[a*x])*ArcCosh[a*x] + 64*a*E^(4*ArcCosh[a*x])*x*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x] + 64*a^2*E^(4*ArcCosh[a*x])*x^2*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x] - 16*E^(4*ArcCosh[a*x])*(-ArcCosh[a*x])^(3/2)*Gamma[1/2, -4*ArcCosh[a*x]] + 16*Sqrt[2]*E^(4*ArcCosh[a*x])*(-ArcCosh[a*x])^(3/2)*Gamma[1/2, -2*ArcCosh[a*x]] + 16*Sqrt[2]*E^(4*ArcCosh[a*x])*ArcCosh[a*x]^(3/2)*Gamma[1/2, 2*ArcCosh[a*x]] - 16*E^(4*ArcCosh[a*x])*ArcCosh[a*x]^(3/2)*Gamma[1/2, 4*ArcCosh[a*x]]))/(a*E^(4*ArcCosh[a*x])*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*ArcCosh[a*x]^(3/2))
```

**3.414.3 Rubi [A] (verified)**Time = 2.20 (sec) , antiderivative size = 251, normalized size of antiderivative = 0.76, number of steps used = 13, number of rules used = 12,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$ , Rules used = {6319, 25, 6327, 6357, 6322, 3042, 25, 3793, 2009, 6368, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$$

↓ 6319

$$-\frac{8ac\sqrt{c - a^2 cx^2} \int -\frac{x(1-ax)(ax+1)}{\operatorname{arccosh}(ax)^{3/2}} dx}{3\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}(c - a^2 cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}$$

↓ 25

---

3.414.  $\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

$$\begin{aligned}
& \frac{8ac\sqrt{c-a^2cx^2} \int \frac{x(1-ax)(ax+1)}{\operatorname{arccosh}(ax)^{3/2}} dx}{3\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
& \quad \downarrow \text{6327} \\
& \frac{8ac\sqrt{c-a^2cx^2} \int \frac{x(1-a^2x^2)}{\operatorname{arccosh}(ax)^{3/2}} dx}{3\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
& \quad \downarrow \text{6357} \\
& \frac{8ac\sqrt{c-a^2cx^2} \left( -8a \int \frac{x^2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx + \frac{2 \int \frac{\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx}{a} + \frac{2x(ax-1)^{3/2}(ax+1)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{3\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
& \quad \downarrow \text{6322} \\
& \frac{8ac\sqrt{c-a^2cx^2} \left( \frac{2 \int \frac{(ax-1)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} - 8a \int \frac{x^2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx + \frac{2x(ax-1)^{3/2}(ax+1)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{3\sqrt{ax-1}\sqrt{ax+1}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
& \quad \downarrow \text{3042} \\
& \frac{8ac\sqrt{c-a^2cx^2} \left( \frac{2 \int -\frac{\sin(i\operatorname{arccosh}(ax))^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} - 8a \int \frac{x^2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx + \frac{2x(ax-1)^{3/2}(ax+1)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{3\sqrt{ax-1}\sqrt{ax+1}} + \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
& \quad \downarrow \text{25} \\
& \frac{8ac\sqrt{c-a^2cx^2} \left( \frac{2 \int \frac{\sin(i\operatorname{arccosh}(ax))^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} - 8a \int \frac{x^2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx + \frac{2x(ax-1)^{3/2}(ax+1)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{3\sqrt{ax-1}\sqrt{ax+1}} + \frac{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
& \quad \downarrow \text{3793} \\
& \frac{8ac\sqrt{c-a^2cx^2} \left( \frac{2 \int \frac{\sin(i\operatorname{arccosh}(ax))^2}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} - 8a \int \frac{x^2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx + \frac{2x(ax-1)^{3/2}(ax+1)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{3\sqrt{ax-1}\sqrt{ax+1}}
\end{aligned}$$

---

3.414.  $\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

$$8ac\sqrt{c - a^2cx^2} \left( -\frac{2 \int \left( \frac{1}{2\sqrt{\operatorname{arccosh}(ax)}} - \frac{\cosh(2\operatorname{arccosh}(ax))}{2\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{a^2} - 8a \int \frac{x^2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx + \frac{2x(ax-1)^{3/2}(ax+1)^{3/2}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)$$


---


$$\frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}} - \frac{3a\operatorname{arccosh}(ax)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}$$

↓ 2009

$$8ac\sqrt{c - a^2cx^2} \left( -8a \int \frac{x^2\sqrt{ax-1}\sqrt{ax+1}}{\sqrt{\operatorname{arccosh}(ax)}} dx + \frac{2\left(\frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \sqrt{\operatorname{arccosh}(ax)}\right)}{a^2} \right)$$


---


$$\frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}} - \frac{3a\operatorname{arccosh}(ax)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}$$

↓ 6368

$$8ac\sqrt{c - a^2cx^2} \left( -\frac{8 \int \frac{a^2x^2(ax-1)(ax+1)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} + \frac{2\left(\frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \sqrt{\operatorname{arccosh}(ax)}\right)}{a^2} \right)$$


---


$$\frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}} - \frac{3a\operatorname{arccosh}(ax)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}$$

↓ 5971

$$8ac\sqrt{c - a^2cx^2} \left( -\frac{8 \int \left( \frac{\cosh(4\operatorname{arccosh}(ax))}{8\sqrt{\operatorname{arccosh}(ax)}} - \frac{1}{8\sqrt{\operatorname{arccosh}(ax)}} \right) d\operatorname{arccosh}(ax)}{a^2} + \frac{2\left(\frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \sqrt{\operatorname{arccosh}(ax)}\right)}{a^2} \right)$$


---


$$\frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}} - \frac{3a\operatorname{arccosh}(ax)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}$$

↓ 2009

$$8ac\sqrt{c - a^2cx^2} \left( -\frac{8\left(\frac{1}{32}\sqrt{\pi}\operatorname{erf}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{32}\sqrt{\pi}\operatorname{erfi}\left(2\sqrt{\operatorname{arccosh}(ax)}\right) - \frac{1}{4}\sqrt{\operatorname{arccosh}(ax)}\right)}{a^2} + \frac{2\left(\frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) + \frac{1}{4}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right) - \sqrt{\operatorname{arccosh}(ax)}\right)}{a^2} \right)$$


---


$$\frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}(c-a^2cx^2)^{3/2}} - \frac{3a\operatorname{arccosh}(ax)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}$$

---

3.414.  $\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$



input `Int[(c - a^2*c*x^2)^(3/2)/ArcCosh[a*x]^(5/2),x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*(c - a^2*c*x^2)^(3/2))/(3*a*ArcCosh[a*x]^(3/2)) + (8*a*c*Sqrt[c - a^2*c*x^2]*((2*x*(-1 + a*x)^(3/2)*(1 + a*x)^(3/2))/(a*Sqrt[ArcCosh[a*x]]) - (8*(-1/4*Sqrt[ArcCosh[a*x]] + (Sqrt[Pi]*Erf[2*Sqrt[ArcCosh[a*x]]])/32 + (Sqrt[Pi]*Erfi[2*Sqrt[ArcCosh[a*x]]])/32))/a^2 + (2*(-Sqrt[ArcCosh[a*x]] + (Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/4 + (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/4))/a^2))/(3*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`

### 3.414.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

---

3.414. 
$$\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$$

rule 6322 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[2*p, 0]`

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6357 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(f*x)^m*Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + (Simp[f*(m/(b*c*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m - 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] - Simp[c*((m + 2*p + 1)/(b*f*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[(f*x)^(m + 1)*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x]) /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IGtQ[2*p, 0] && NeQ[m + 2*p + 1, 0] && IGtQ[m, -3]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.414.4 Maple [F]

$$\int \frac{(-a^2cx^2 + c)^{\frac{3}{2}}}{\operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `int((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x)`

output `int((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x)`

---

3.414.  $\int \frac{(c-a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

**3.414.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{(c - a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.414.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(c - a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \text{Timed out}$$

input `integrate((-a**2*c*x**2+c)**(3/2)/acosh(a*x)**(5/2), x)`

output `Timed out`

**3.414.7 Maxima [F]**

$$\int \frac{(c - a^2cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{(-a^2cx^2 + c)^{\frac{3}{2}}}{\operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x, algorithm="maxima")`

output `integrate((-a^2*c*x^2 + c)^(3/2)/arccosh(a*x)^(5/2), x)`

**3.414.8 Giac [F]**

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{(-a^2 cx^2 + c)^{\frac{3}{2}}}{\operatorname{arcosh}(ax)^{\frac{5}{2}}} dx$$

input `integrate((-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x, algorithm="giac")`

output `integrate((-a^2*c*x^2 + c)^(3/2)/arccosh(a*x)^(5/2), x)`

**3.414.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(c - a^2 cx^2)^{3/2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{(c - a^2 c x^2)^{3/2}}{\operatorname{acosh}(ax)^{5/2}} dx$$

input `int((c - a^2*c*x^2)^(3/2)/acosh(a*x)^(5/2),x)`

output `int((c - a^2*c*x^2)^(3/2)/acosh(a*x)^(5/2), x)`

### 3.415 $\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

3.415.1 Optimal result	3188
3.415.2 Mathematica [A] (warning: unable to verify)	3188
3.415.3 Rubi [A] (verified)	3189
3.415.4 Maple [F]	3192
3.415.5 Fricas [F(-2)]	3193
3.415.6 Sympy [F]	3193
3.415.7 Maxima [F]	3193
3.415.8 Giac [F]	3194
3.415.9 Mupad [F(-1)]	3194

#### 3.415.1 Optimal result

Integrand size = 24, antiderivative size = 201

$$\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}\sqrt{c-a^2cx^2}}{3a\operatorname{arccosh}(ax)^{3/2}} - \frac{8x\sqrt{c-a^2cx^2}}{3\sqrt{\operatorname{arccosh}(ax)}} + \frac{2\sqrt{2\pi}\sqrt{c-a^2cx^2}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{3a\sqrt{-1+ax}\sqrt{1+ax}} + \frac{2\sqrt{2\pi}\sqrt{c-a^2cx^2}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)}{3a\sqrt{-1+ax}\sqrt{1+ax}}$$

output

```
2/3*erf(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/
a/(a*x-1)^(1/2)/(a*x+1)^(1/2)+2/3*erfi(2^(1/2)*arccosh(a*x)^(1/2))*2^(1/2)
*Pi^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/(a*x-1)^(1/2)/(a*x+1)^(1/2)-2/3*(a*x-1)^(
1/2)*(a*x+1)^(1/2)*(-a^2*c*x^2+c)^(1/2)/a/arccosh(a*x)^(3/2)-8/3*x*(-a^2*c
*x^2+c)^(1/2)/arccosh(a*x)^(1/2)
```

#### 3.415.2 Mathematica [A] (warning: unable to verify)

Time = 0.26 (sec) , antiderivative size = 141, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \frac{2\sqrt{c-a^2cx^2}\left((1+ax)\left(-1+ax+4ax\sqrt{\frac{-1+ax}{1+ax}}\operatorname{arccosh}(ax)\right)+\sqrt{2}(-\operatorname{arccosh}(ax))^{3/2}\Gamma\left(\frac{1}{2},-2\operatorname{arccosh}(ax)\right)\right)}{3a\sqrt{\frac{-1+ax}{1+ax}}(1+ax)\operatorname{arccosh}(ax)^{3/2}}$$

input `Integrate[Sqrt[c - a^2*c*x^2]/ArcCosh[a*x]^(5/2),x]`

output `(-2*Sqrt[c - a^2*c*x^2]*((1 + a*x)*(-1 + a*x + 4*a*x*Sqrt[(-1 + a*x)/(1 + a*x)]*ArcCosh[a*x]) + Sqrt[2]*(-ArcCosh[a*x])^(3/2)*Gamma[1/2, -2*ArcCosh[a*x]] + Sqrt[2]*ArcCosh[a*x]^(3/2)*Gamma[1/2, 2*ArcCosh[a*x]])/(3*a*Sqrt[(-1 + a*x)/(1 + a*x)]*(1 + a*x)*ArcCosh[a*x]^(3/2))`

### 3.415.3 Rubi [A] (verified)

Time = 0.62 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.91, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$ , Rules used = {6319, 6300, 25, 3042, 3788, 26, 2611, 2633, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{c - a^2 cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx \\
 & \quad \downarrow \text{6319} \\
 & \frac{4a\sqrt{c - a^2 cx^2} \int \frac{x}{\operatorname{arccosh}(ax)^{3/2}} dx}{3\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{c - a^2 cx^2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
 & \quad \downarrow \text{6300} \\
 & \frac{4a\sqrt{c - a^2 cx^2} \left( -\frac{2 \int \frac{\cosh(2\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} - \frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{\frac{3\sqrt{ax - 1}\sqrt{ax + 1}}{2\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{c - a^2 cx^2}} - \frac{3a\operatorname{arccosh}(ax)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}} \\
 & \quad \downarrow \text{25} \\
 & \frac{4a\sqrt{c - a^2 cx^2} \left( \frac{2 \int \frac{\cosh(2\operatorname{arccosh}(ax))}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} - \frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{3\sqrt{ax - 1}\sqrt{ax + 1}} - \frac{2\sqrt{ax - 1}\sqrt{ax + 1}\sqrt{c - a^2 cx^2}}{3a\operatorname{arccosh}(ax)^{3/2}} \\
 & \quad \downarrow \text{3042}
 \end{aligned}$$

---

3.415.  $\int \frac{\sqrt{c - a^2 cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

$$\begin{aligned}
 & -\frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{3\operatorname{arccosh}(ax)^{3/2}} + \\
 & 4a\sqrt{c-a^2cx^2} \left( -\frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} + \frac{2\int \frac{\sin\left(2i\operatorname{arccosh}(ax)+\frac{\pi}{2}\right)}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)}{a^2} \right) \\
 & \hline
 & \frac{3\sqrt{ax-1}\sqrt{ax+1}}{\downarrow 3788} \\
 & -\frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{3\operatorname{arccosh}(ax)^{3/2}} + \\
 & 4a\sqrt{c-a^2cx^2} \left( -\frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} - \frac{2\left(\frac{1}{2}i\int \frac{ie^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2}i\int \frac{-ie^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)\right)}{a^2} \right) \\
 & \hline
 & \frac{3\sqrt{ax-1}\sqrt{ax+1}}{\downarrow 26} \\
 & 4a\sqrt{c-a^2cx^2} \left( -\frac{2\left(-\frac{1}{2}\int \frac{e^{-2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax) - \frac{1}{2}\int \frac{e^{2\operatorname{arccosh}(ax)}}{\sqrt{\operatorname{arccosh}(ax)}} d\operatorname{arccosh}(ax)\right)}{a^2} - \frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} \right) \\
 & \hline
 & \frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}} \\
 & \frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{3\operatorname{arccosh}(ax)^{3/2}} \\
 & \downarrow 2611 \\
 & 4a\sqrt{c-a^2cx^2} \left( -\frac{2\left(-\int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - \int e^{2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)}\right)}{a^2} - \frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} \right) \\
 & \hline
 & \frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}} \\
 & \frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{3\operatorname{arccosh}(ax)^{3/2}} \\
 & \downarrow 2633 \\
 & 4a\sqrt{c-a^2cx^2} \left( -\frac{2\left(-\int e^{-2\operatorname{arccosh}(ax)} d\sqrt{\operatorname{arccosh}(ax)} - \frac{1}{2}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)\right)}{a^2} - \frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} \right) \\
 & \hline
 & \frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}} \\
 & \frac{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}}{3\operatorname{arccosh}(ax)^{3/2}} \\
 & \downarrow 2634
 \end{aligned}$$

3.415.  $\int \frac{\sqrt{c-a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx$

$$\frac{4a\sqrt{c-a^2cx^2} \left( -\frac{2\left(-\frac{1}{2}\sqrt{\frac{\pi}{2}}\operatorname{erf}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)-\frac{1}{2}\sqrt{\frac{\pi}{2}}\operatorname{erfi}\left(\sqrt{2}\sqrt{\operatorname{arccosh}(ax)}\right)\right)}{a^2} - \frac{2x\sqrt{ax-1}\sqrt{ax+1}}{a\sqrt{\operatorname{arccosh}(ax)}} \right)}{\frac{3\sqrt{ax-1}\sqrt{ax+1}}{2\sqrt{ax-1}\sqrt{ax+1}\sqrt{c-a^2cx^2}} - \frac{3a\operatorname{arccosh}(ax)^{3/2}}{3a\operatorname{arccosh}(ax)^{3/2}}}$$

input `Int[Sqrt[c - a^2*c*x^2]/ArcCosh[a*x]^(5/2), x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x]*Sqrt[c - a^2*c*x^2])/(3*a*ArcCosh[a*x]^(3/2)) + (4*a*Sqrt[c - a^2*c*x^2]*((-2*x*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(a*Sqrt[ArcCosh[a*x]]) - (2*(-1/2*(Sqrt[Pi/2]*Erf[Sqrt[2]*Sqrt[ArcCosh[a*x]]]) - (Sqrt[Pi/2]*Erfi[Sqrt[2]*Sqrt[ArcCosh[a*x]]])/2))/a^2))/(3*Sqrt[-1 + a*x]*Sqrt[1 + a*x])`

### 3.415.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2611 `Int[(F_)^((g_)*((e_) + (f_)*(x_)))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] := Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 2633 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`

rule 2634 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`



rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3788 `Int[((c_.) + (d_.)*(x_)^(m_.)*sin[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/(E^(I*k*Pi)*E^(I*(e + f*x))), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*k*Pi)*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x] && IntegerQ[2*k]`

rule 6300 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*(x_)^(m_.), x_Symbol] := Simp[x^m*Sqrt[1 + c*x]*Sqrt[-1 + c*x]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] + Simp[1/(b^2*c^(m + 1)*(n + 1)) Subst[Int[ExpandTrigReduce[x^(n + 1), Cosh[-a/b + x/b]^(m - 1)*(m - (m + 1)*Cosh[-a/b + x/b]^2), x], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c}, x] && IGtQ[m, 0] && GeQ[n, -2] && LtQ[n, -1]`

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p) Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

### 3.415.4 Maple [F]

$$\int \frac{\sqrt{-a^2cx^2 + c}}{\operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `int((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x)`

output `int((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x)`

**3.415.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.415.6 Sympy [F]**

$$\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{\sqrt{-c(ax - 1)(ax + 1)}}{\operatorname{acosh}^{\frac{5}{2}}(ax)} dx$$

input `integrate((-a**2*c*x**2+c)**(1/2)/acosh(a*x)**(5/2),x)`

output `Integral(sqrt(-c*(a*x - 1)*(a*x + 1))/acosh(a*x)**(5/2), x)`

**3.415.7 Maxima [F]**

$$\int \frac{\sqrt{c - a^2cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{\sqrt{-a^2cx^2 + c}}{\operatorname{arcosh}(ax)^{\frac{5}{2}}} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x, algorithm="maxima")`

output `integrate(sqrt(-a^2*c*x^2 + c)/arccosh(a*x)^(5/2), x)`

**3.415.8 Giac [F]**

$$\int \frac{\sqrt{c - a^2 cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{\sqrt{-a^2 cx^2 + c}}{\operatorname{arcosh}(ax)^{5/2}} dx$$

input `integrate((-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x, algorithm="giac")`

output `integrate(sqrt(-a^2*c*x^2 + c)/arccosh(a*x)^(5/2), x)`

**3.415.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{\sqrt{c - a^2 cx^2}}{\operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{\sqrt{c - a^2 c x^2}}{\operatorname{acosh}(ax)^{5/2}} dx$$

input `int((c - a^2*c*x^2)^(1/2)/acosh(a*x)^(5/2),x)`

output `int((c - a^2*c*x^2)^(1/2)/acosh(a*x)^(5/2), x)`

**3.416**  $\int \frac{1}{\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{5/2}} dx$

3.416.1 Optimal result	3195
3.416.2 Mathematica [A] (verified)	3195
3.416.3 Rubi [A] (verified)	3196
3.416.4 Maple [A] (verified)	3196
3.416.5 Fricas [A] (verification not implemented)	3197
3.416.6 Sympy [F(-1)]	3197
3.416.7 Maxima [F]	3197
3.416.8 Giac [F]	3198
3.416.9 Mupad [F(-1)]	3198

**3.416.1 Optimal result**

Integrand size = 24, antiderivative size = 48

$$\int \frac{1}{\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{5/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}}{3a\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{3/2}}$$

output `-2/3*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/arccosh(a*x)^(3/2)/(-a^2*c*x^2+c)^(1/2)`

**3.416.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{5/2}} dx = -\frac{2\sqrt{-1+ax}\sqrt{1+ax}}{3a\sqrt{c-a^2cx^2}\operatorname{arccosh}(ax)^{3/2}}$$

input `Integrate[1/(Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(5/2)),x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(3*a*Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2))`

**3.416.3 Rubi [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arccosh}(ax)^{5/2} \sqrt{c - a^2 cx^2}} dx$$

↓ 6307

$$-\frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a\operatorname{arccosh}(ax)^{3/2}\sqrt{c - a^2 cx^2}}$$

input `Int[1/(Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(5/2)),x]`

output `(-2*Sqrt[-1 + a*x]*Sqrt[1 + a*x])/(3*a*Sqrt[c - a^2*c*x^2]*ArcCosh[a*x]^(3/2))`

**3.416.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.416.4 Maple [A] (verified)**

Time = 0.58 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.85

method	result	size
default	$-\frac{2\sqrt{ax-1}\sqrt{ax+1}}{3\sqrt{-c(ax-1)(ax+1)}\operatorname{arccosh}(ax)^{\frac{3}{2}}a}$	41

input `int(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x,method=_RETURNVERBOSE)`

output  $-2/3/(-c*(a*x-1)*(a*x+1))^{(1/2)}*(a*x-1)^{(1/2)}*(a*x+1)^{(1/2)}/\operatorname{arccosh}(a*x)^{(3/2)}/a$

### 3.416.5 Fricas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.23

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2}} dx = \frac{2\sqrt{-a^2 cx^2 + c} \sqrt{a^2 x^2 - 1}}{3(a^3 cx^2 - ac) \log(ax + \sqrt{a^2 x^2 - 1})^{3/2}}$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x, algorithm="fricas")`

output  $2/3*\sqrt{-a^2*c*x^2 + c}*\sqrt{a^2*x^2 - 1}/((a^3*c*x^2 - a*c)*\log(a*x + \sqrt{a^2*x^2 - 1}))^{(3/2)}$

### 3.416.6 Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2}} dx = \text{Timed out}$$

input `integrate(1/(-a**2*c*x**2+c)**(1/2)/acosh(a*x)**(5/2),x)`

output Timed out

### 3.416.7 Maxima [F]

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{\sqrt{-a^2 cx^2 + c} \operatorname{arccosh}(ax)^{5/2}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x, algorithm="maxima")`

output `integrate(1/(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^(5/2)), x)`

**3.416.8 Giac [F]**

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{\sqrt{-a^2 cx^2 + c} \operatorname{arccosh}(ax)^{5/2}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(1/2)/arccosh(a*x)^(5/2),x, algorithm="giac")`

output `integrate(1/(sqrt(-a^2*c*x^2 + c)*arccosh(a*x)^(5/2)), x)`

**3.416.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{c - a^2 cx^2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{\operatorname{acosh}(ax)^{5/2} \sqrt{c - a^2 cx^2}} dx$$

input `int(1/(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(1/2)),x)`

output `int(1/(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(1/2)), x)`

**3.417**  $\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx$

3.417.1 Optimal result	3199
3.417.2 Mathematica [N/A]	3199
3.417.3 Rubi [N/A]	3200
3.417.4 Maple [N/A] (verified)	3201
3.417.5 Fricas [F(-2)]	3201
3.417.6 Sympy [F(-1)]	3202
3.417.7 Maxima [N/A]	3202
3.417.8 Giac [N/A]	3202
3.417.9 Mupad [N/A]	3203

**3.417.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx = -\frac{2\sqrt{-1 + ax}\sqrt{1 + ax}}{3a(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{3/2}} + \frac{4a\sqrt{-1 + ax}\sqrt{1 + ax} \operatorname{Int}\left(\frac{x}{(-1 + a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}}, x\right)}{3c\sqrt{c - a^2cx^2}}$$

output `-2/3*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(3/2)+4/3*a*(a*x-1)^(1/2)*(a*x+1)^(1/2)*Unintegrable(x/(a^2*x^2-1)^2/arccosh(a*x)^(3/2),x)/c/(-a^2*c*x^2+c)^(1/2)`

**3.417.2 Mathematica [N/A]**

Not integrable

Time = 4.75 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx$$

input `Integrate[1/((c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(5/2)), x]`

output `Integrate[1/((c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(5/2)), x]`



### 3.417.3 Rubi [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{3/2}} dx$$

↓ 6319

$$\frac{4a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-ax)^2(ax+1)^2 \operatorname{arccosh}(ax)^{3/2}} dx}{3c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a \operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}}$$

↓ 6327

$$\frac{4a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}} dx}{3c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a \operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}}$$

↓ 6375

$$\frac{4a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^2 \operatorname{arccosh}(ax)^{3/2}} dx}{3c\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a \operatorname{arccosh}(ax)^{3/2} (c-a^2cx^2)^{3/2}}$$

input `Int[1/((c - a^2*c*x^2)^(3/2)*ArcCosh[a*x]^(5/2)),x]`

output `$Aborted`

#### 3.417.3.1 Defintions of rubi rules used

rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_)*((d_) + (e_.)*(x_)^2)^(p_.), x  
_Symbol] :> Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`

---

3.417.  $\int \frac{1}{(c-a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx$

rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_))^(p_.)*((d2_) + (e2_.)*(x_))^(p_.), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

### 3.417.4 Maple [N/A] (verified)

Not integrable

Time = 1.52 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `int(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x)`

output `int(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x)`

### 3.417.5 Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.417.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx = \text{Timed out}$$

input `integrate(1/(-a**2*c*x**2+c)**(3/2)/acosh(a*x)**(5/2), x)`output `Timed out`**3.417.7 Maxima [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{(-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2), x, algorithm="maxima")`output `integrate(1/((-a^2*c*x^2 + c)^(3/2)*arccosh(a*x)^(5/2)), x)`**3.417.8 Giac [N/A]**

Not integrable

Time = 0.38 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{3/2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{(-a^2cx^2 + c)^{\frac{3}{2}} \operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(3/2)/arccosh(a*x)^(5/2), x, algorithm="giac")`output `integrate(1/((-a^2*c*x^2 + c)^(3/2)*arccosh(a*x)^(5/2)), x)`

**3.417.9 Mupad [N/A]**

Not integrable

Time = 2.76 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 c x^2)^{3/2} \operatorname{arccosh}(a x)^{5/2}} dx = \int \frac{1}{\operatorname{acosh}(a x)^{5/2} (c - a^2 c x^2)^{3/2}} dx$$

input `int(1/(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(3/2)),x)`output `int(1/(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(3/2)), x)`

**3.418**  $\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx$

3.418.1 Optimal result	3204
3.418.2 Mathematica [N/A]	3204
3.418.3 Rubi [N/A]	3205
3.418.4 Maple [N/A] (verified)	3206
3.418.5 Fricas [F(-2)]	3207
3.418.6 Sympy [F(-1)]	3207
3.418.7 Maxima [N/A]	3207
3.418.8 Giac [N/A]	3208
3.418.9 Mupad [N/A]	3208

**3.418.1 Optimal result**

Integrand size = 24, antiderivative size = 24

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx = -\frac{2\sqrt{-1 + ax}\sqrt{1 + ax}}{3a(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{3/2}} - \frac{8a\sqrt{-1 + ax}\sqrt{1 + ax} \operatorname{Int}\left(\frac{x}{(-1 + a^2x^2)^3 \operatorname{arccosh}(ax)^{3/2}}, x\right)}{3c^2\sqrt{c - a^2cx^2}}$$

output `-2/3*(a*x-1)^(1/2)*(a*x+1)^(1/2)/a/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(3/2)  
-8/3*a*(a*x-1)^(1/2)*(a*x+1)^(1/2)*Unintegrable(x/(a^2*x^2-1)^3/arccosh(a*x)^(3/2),x)/c^2/(-a^2*c*x^2+c)^(1/2)`

**3.418.2 Mathematica [N/A]**

Not integrable

Time = 4.58 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx$$

input `Integrate[1/((c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^(5/2)), x]`

output `Integrate[1/((c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^(5/2)), x]`

### 3.418.3 Rubi [N/A]

Not integrable

Time = 0.56 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6319, 25, 6327, 6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\operatorname{arccosh}(ax)^{5/2} (c - a^2cx^2)^{5/2}} dx \\
 & \quad \downarrow \text{6319} \\
 & -\frac{8a\sqrt{ax-1}\sqrt{ax+1} \int -\frac{x}{(1-ax)^3(ax+1)^3 \operatorname{arccosh}(ax)^{3/2}} dx}{3c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{25} \\
 & \frac{8a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-ax)^3(ax+1)^3 \operatorname{arccosh}(ax)^{3/2}} dx}{3c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{6327} \\
 & \frac{8a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^3 \operatorname{arccosh}(ax)^{3/2}} dx}{3c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{5/2}} \\
 & \quad \downarrow \text{6375} \\
 & \frac{8a\sqrt{ax-1}\sqrt{ax+1} \int \frac{x}{(1-a^2x^2)^3 \operatorname{arccosh}(ax)^{3/2}} dx}{3c^2\sqrt{c-a^2cx^2}} - \frac{2\sqrt{ax-1}\sqrt{ax+1}}{3a\operatorname{arccosh}(ax)^{3/2} (c - a^2cx^2)^{5/2}}
 \end{aligned}$$

input `Int[1/((c - a^2*c*x^2)^(5/2)*ArcCosh[a*x]^(5/2)),x]`

output `$Aborted`

## 3.418.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 6319 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*(d + e*x^2)^p]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c*((2*p + 1)/(b*(n + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Int[x*(1 + c*x)^(p - 1/2)*(-1 + c*x)^(p - 1/2)*(a + b*ArcCosh[c*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && LtQ[n, -1] && IntegerQ[2*p]`
- rule 6327 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_)^m_)*((d1_) + (e1_.)*(x_)^p_)*((d2_) + (e2_.)*(x_)^p_), x_Symbol] := Int[(f*x)^m*(d1*d2 + e1*e2*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, m, n}, x] && EqQ[d2*e1 + d1*e2, 0] && IntegerQ[p]`
- rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_)^m_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

## 3.418.4 Maple [N/A] (verified)

Not integrable

Time = 1.59 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{1}{(-a^2cx^2 + c)^{\frac{5}{2}} \operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `int(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(5/2),x)`output `int(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(5/2),x)`

**3.418.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(5/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.418.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx = \text{Timed out}$$

input `integrate(1/(-a**2*c*x**2+c)**(5/2)/acosh(a*x)**(5/2),x)`

output `Timed out`

**3.418.7 Maxima [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{(-a^2cx^2 + c)^{\frac{5}{2}} \operatorname{arccosh}(ax)^{\frac{5}{2}}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(5/2),x, algorithm="maxima")`

output `integrate(1/((-a^2*c*x^2 + c)^(5/2)*arccosh(a*x)^(5/2)), x)`



**3.418.8 Giac [N/A]**

Not integrable

Time = 0.39 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{(-a^2 cx^2 + c)^{5/2} \operatorname{arcosh}(ax)^{5/2}} dx$$

input `integrate(1/(-a^2*c*x^2+c)^(5/2)/arccosh(a*x)^(5/2),x, algorithm="giac")`output `integrate(1/((-a^2*c*x^2 + c)^(5/2)*arccosh(a*x)^(5/2)), x)`**3.418.9 Mupad [N/A]**

Not integrable

Time = 2.75 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{1}{(c - a^2 cx^2)^{5/2} \operatorname{arccosh}(ax)^{5/2}} dx = \int \frac{1}{\operatorname{acosh}(ax)^{5/2} (c - a^2 cx^2)^{5/2}} dx$$

input `int(1/(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(5/2)),x)`output `int(1/(acosh(a*x)^(5/2)*(c - a^2*c*x^2)^(5/2)), x)`

### 3.419 $\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$

3.419.1 Optimal result . . . . .	3209
3.419.2 Mathematica [A] (verified) . . . . .	3210
3.419.3 Rubi [A] (verified) . . . . .	3210
3.419.4 Maple [F] . . . . .	3211
3.419.5 Fracas [F] . . . . .	3212
3.419.6 Sympy [F] . . . . .	3212
3.419.7 Maxima [F] . . . . .	3212
3.419.8 Giac [F(-2)] . . . . .	3213
3.419.9 Mupad [F(-1)] . . . . .	3213

#### 3.419.1 Optimal result

Integrand size = 29, antiderivative size = 253

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = -\frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^{1+n}}{8bc^3(1+n)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2^{-2(3+n)} e^{-\frac{4a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{c^3 \sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2^{-2(3+n)} e^{\frac{4a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{c^3 \sqrt{-1+cx}\sqrt{1+cx}}$$

```
output -1/8*(a+b*arccosh(c*x))^(1+n)*(-c^2*d*x^2+d)^(1/2)/b/c^3/(1+n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+(a+b*arccosh(c*x))^n*GAMMA(1+n,-4*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/(2^(6+2*n))/c^3/exp(4*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-exp(4*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,4*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/(2^(6+2*n))/c^3/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.419.2 Mathematica [A] (verified)**

Time = 0.75 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.72

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx =$$

$$\frac{d \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (a + \operatorname{barccosh}(cx))^n \left( -\frac{8(a+\operatorname{barccosh}(cx))}{b(1+n)} + 4^{-n} e^{-\frac{4a}{b}} \left( -\frac{(a+\operatorname{barccosh}(cx))^2}{b^2} \right)^{-n} \left( \frac{a}{b} + \operatorname{arccosh}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right) \right) \right)}{64c^3 \sqrt{-d(-1+cx)}}$$

input `Integrate[x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`output `-1/64*(d*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])^n*((-8*(a + b*ArcCosh[c*x]))/(b*(1 + n)) + ((a/b + ArcCosh[c*x])^n*Gamma[1 + n, (-4*(a + b*ArcCosh[c*x]))/b] - E^((8*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, (4*(a + b*ArcCosh[c*x])/b)]/(4^n*E^((4*a)/b)*(-(a + b*ArcCosh[c*x])^2/b^2))^n)/(c^3*Sqrt[-(d*(-1 + c*x)*(1 + c*x))])`**3.419.3 Rubi [A] (verified)**Time = 0.58 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.73, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx))^n \cosh^2\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^3 \sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow \text{5971}$$

$$\frac{\sqrt{d - c^2 dx^2} \int \left(\frac{1}{8}(a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{4a}{b} - \frac{4(a + \operatorname{barccosh}(cx))}{b}\right) - \frac{1}{8}(a + \operatorname{barccosh}(cx))^n\right) d(a + \operatorname{barccosh}(cx))}{bc^3 \sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow \text{2009}$$

---

3.419.  $\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$

$$\frac{\sqrt{d - c^2 dx^2} \left( -\frac{(a + b \operatorname{arccosh}(cx))^{n+1}}{8(n+1)} + b 2^{-2(n+3)} e^{-\frac{4a}{b}} (a + b \operatorname{arccosh}(cx))^n \left( -\frac{a + b \operatorname{arccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{4(a + b \operatorname{arccosh}(cx))}{b} \right) \right)}{bc^3 \sqrt{cx - 1} \sqrt{cx + 1}}$$

input `Int[x^2*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`

output `(Sqrt[d - c^2*d*x^2]*(-1/8*(a + b*ArcCosh[c*x])^(1 + n)/(1 + n) + (b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-4*(a + b*ArcCosh[c*x]))/b])/(2^(2*(3 + n)))*E^((4*a)/b)*(-((a + b*ArcCosh[c*x])/b))^n) - (b*E^((4*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (4*(a + b*ArcCosh[c*x]))/b])/(2^(2*(3 + n))*((a + b*ArcCosh[c*x])/b)^n))/(b*c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.419.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n*(x_)^m*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.419.4 Maple [F]

$$\int x^2(a + b \operatorname{arccosh}(cx))^n \sqrt{-c^2 dx^2 + d} dx$$

input `int(x^2*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x)`

output `int(x^2*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x)`

**3.419.5 Fracas [F]**

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^n x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x^2, x)`

**3.419.6 Sympy [F]**

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int x^2 \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^n dx$$

input `integrate(x**2*(a+b*acosh(c*x))**n*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**2*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**n, x)`

**3.419.7 Maxima [F]**

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^n x^2 dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x^2, x)`

**3.419.8 Giac [F(-2)]**

Exception generated.

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate(x^2*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

**3.419.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int x^2 (a + b \operatorname{acosh}(cx))^n \sqrt{d - c^2 dx^2} dx$$

input `int(x^2*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2),x)`

output `int(x^2*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2), x)`

### 3.420 $\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n dx$

3.420.1 Optimal result . . . . .	3214
3.420.2 Mathematica [A] (verified) . . . . .	3215
3.420.3 Rubi [A] (verified) . . . . .	3215
3.420.4 Maple [F] . . . . .	3217
3.420.5 Fracas [F] . . . . .	3217
3.420.6 Sympy [F] . . . . .	3217
3.420.7 Maxima [F] . . . . .	3218
3.420.8 Giac [F(-2)] . . . . .	3218
3.420.9 Mupad [F(-1)] . . . . .	3218

#### 3.420.1 Optimal result

Integrand size = 27, antiderivative size = 379

$$\int x\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n dx$$

$$= \frac{3^{-1-n}e^{-\frac{3a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{8c^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{e^{-\frac{a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{8c^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{e^{a/b}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{a+\operatorname{barccosh}(cx)}{b}\right)}{8c^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{3^{-1-n}e^{\frac{3a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{8c^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
1/8*3^(-1-n)*(a+b*arccosh(c*x))^n*GAMMA(1+n,-3*(a+b*arccosh(c*x))/b)*(-c^2
*d*x^2+d)^(1/2)/c^2/exp(3*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(
c*x+1)^(1/2)-1/8*(a+b*arccosh(c*x))^n*GAMMA(1+n,(-a-b*arccosh(c*x))/b)*(-c
^2*d*x^2+d)^(1/2)/c^2/exp(a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(
c*x+1)^(1/2)+1/8*exp(a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,(a+b*arccosh(c*x)
)/b)*(-c^2*d*x^2+d)^(1/2)/c^2/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*
x+1)^(1/2)-1/8*3^(-1-n)*exp(3*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,3*(a+b*a
rccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^2/(((a+b*arccosh(c*x))/b)^n)/(c*x-1
)^(1/2)/(c*x+1)^(1/2)
```

**3.420.2 Mathematica [A] (verified)**

Time = 0.97 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.64

$$\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^n dx =$$

$$de^{-\frac{3a}{b}} \sqrt{\frac{-1+cx}{1+cx}}(1 + cx)(a + \operatorname{barccosh}(cx))^n \left( 3e^{\frac{4a}{b}} \left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)^{-n} \Gamma\left(1 + n, \frac{a}{b} + \operatorname{arccosh}(cx)\right) + \left(-\right.$$

input `Integrate[x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`output `-1/24*(d*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])^n*((3*E^((4*a)/b)*Gamma[1 + n, a/b + ArcCosh[c*x]])/(a/b + ArcCosh[c*x])^n + (Gamma[1 + n, (-3*(a + b*ArcCosh[c*x]))/b]/3^n - 3*E^((2*a)/b)*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b]) - (E^((6*a)/b)*(-(a + b*ArcCosh[c*x])/b)^(2*n)*Gamma[1 + n, (3*(a + b*ArcCosh[c*x]))/b])/(3^n*(-(a + b*ArcCosh[c*x])^2/b^2))^n)/(-(a + b*ArcCosh[c*x])/b)^n)/(c^2*E^((3*a)/b)*Sqrt[-(d*(-1 + c*x)*(1 + c*x))])`**3.420.3 Rubi [A] (verified)**Time = 0.64 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.74, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^n dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) \sinh^2\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^2\sqrt{cx - 1}\sqrt{cx + 1}}$$

$$\downarrow \text{5971}$$

$$\frac{\sqrt{d - c^2 dx^2} \int \left(\frac{1}{4}(a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{3a}{b} - \frac{3(a + \operatorname{barccosh}(cx))}{b}\right) - \frac{1}{4}(a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right)\right)}{bc^2\sqrt{cx - 1}\sqrt{cx + 1}}$$

---


$$3.420. \quad \int x\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^n dx$$



↓ 2009

$$\sqrt{d - c^2 dx^2} \left( \frac{1}{8} b 3^{-n-1} e^{-\frac{3a}{b}} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{3(a + \operatorname{barccosh}(cx))}{b}\right) - \frac{1}{8} b e^{-\frac{a}{b}} (a + \right.$$

input `Int[x*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`

output `(Sqrt[d - c^2*d*x^2]*((3^(-1 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-3*(a + b*ArcCosh[c*x]))/b])/(8*E^((3*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n) - (b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b])/(8*E^(a/b)*(-(a + b*ArcCosh[c*x])/b))^n + (b*E^(a/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (a + b*ArcCosh[c*x])/b])/(8*((a + b*ArcCosh[c*x])/b)^n) - (3^(-1 - n)*b*E^((3*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (3*(a + b*ArcCosh[c*x]))/b])/(8*((a + b*ArcCosh[c*x])/b)^n))/(b*c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.420.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

**3.420.4 Maple [F]**

$$\int x(a + b \operatorname{arccosh}(cx))^n \sqrt{-c^2 dx^2 + d} dx$$

input `int(x*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x)`

output `int(x*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x)`

**3.420.5 Fricas [F]**

$$\int x\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))^n dx = \int \sqrt{-c^2 dx^2 + d}(b \operatorname{arccosh}(cx) + a)^n x dx$$

input `integrate(x*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x, x)`

**3.420.6 Sympy [F]**

$$\int x\sqrt{d - c^2 dx^2}(a + b \operatorname{arccosh}(cx))^n dx = \int x\sqrt{-d(cx - 1)(cx + 1)}(a + b \operatorname{acosh}(cx))^n dx$$

input `integrate(x*(a+b*acosh(c*x))**n*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x*sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**n, x)`

**3.420.7 Maxima [F]**

$$\int x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^n dx = \int \sqrt{-c^2dx^2+d}(b\operatorname{arcosh}(cx)+a)^n x dx$$

input `integrate(x*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x, x)`

**3.420.8 Giac [F(-2)]**

Exception generated.

$$\int x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate(x*(a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.420.9 Mupad [F(-1)]**

Timed out.

$$\int x\sqrt{d-c^2dx^2}(a+\operatorname{barccosh}(cx))^n dx = \int x(a+b\operatorname{acosh}(cx))^n \sqrt{d-c^2dx^2} dx$$

input `int(x*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2),x)`

output `int(x*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2), x)`

### 3.421 $\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$

3.421.1 Optimal result . . . . .	3219
3.421.2 Mathematica [A] (verified) . . . . .	3220
3.421.3 Rubi [A] (verified) . . . . .	3220
3.421.4 Maple [F] . . . . .	3222
3.421.5 Fracas [F] . . . . .	3222
3.421.6 Sympy [F] . . . . .	3222
3.421.7 Maxima [F] . . . . .	3223
3.421.8 Giac [F(-2)] . . . . .	3223
3.421.9 Mupad [F(-1)] . . . . .	3223

#### 3.421.1 Optimal result

Integrand size = 26, antiderivative size = 253

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = -\frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^{1+n}}{2bc(1+n)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2^{-3-n} e^{-\frac{2a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2^{-3-n} e^{\frac{2a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

```
output -1/2*(a+b*arccosh(c*x))^(1+n)*(-c^2*d*x^2+d)^(1/2)/b/c/(1+n)/(c*x-1)^(1/2)
/(c*x+1)^(1/2)+2^(-3-n)*(a+b*arccosh(c*x))^n*GAMMA(1+n,-2*(a+b*arccosh(c*x)
)/b)*(-c^2*d*x^2+d)^(1/2)/c/exp(2*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1
)^(1/2)/(c*x+1)^(1/2)-2^(-3-n)*exp(2*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,2
*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c/(((a+b*arccosh(c*x))/b)^n)/(
c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.421.2 Mathematica [A] (verified)**

Time = 0.53 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.85

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$$

$$= \frac{2^{-3-n} d e^{-\frac{2a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (a + \operatorname{barccosh}(cx))^n \left(-\frac{(a+\operatorname{barccosh}(cx))^2}{b^2}\right)^{-n} \left(2^{2+n} e^{\frac{2a}{b}} (a + \operatorname{barccosh}(cx)) \left(-\frac{(a+\operatorname{barccosh}(cx))^2}{b^2}\right)^{-n} \right)}{}$$

input `Integrate[Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`

output

$$\frac{(2^{-3-n} d \operatorname{Sqrt}[-(1+cx)/(1+cx)] (1+cx) (a + b \operatorname{ArcCosh}[c*x])^n (2^{2+n} E^{((2*a)/b)} (a + b \operatorname{ArcCosh}[c*x]) (-(a + b \operatorname{ArcCosh}[c*x])^2/b^2))^n - b(1+n)(a/b + \operatorname{ArcCosh}[c*x])^n \operatorname{Gamma}[1+n, (-2(a + b \operatorname{ArcCosh}[c*x])/b) + b E^{((4*a)/b)} (1+n) (-(a + b \operatorname{ArcCosh}[c*x])/b))^n \operatorname{Gamma}[1+n, (2(a + b \operatorname{ArcCosh}[c*x])/b)])/ (b*c E^{((2*a)/b)} (1+n) \operatorname{Sqrt}[d - c^2*d*x^2] (-(a + b \operatorname{ArcCosh}[c*x])^2/b^2))^n}{}$$
**3.421.3 Rubi [A] (verified)**Time = 0.51 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.73, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$ , Rules used = {6321, 3042, 25, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$$

$$\downarrow \text{6321}$$

$$\frac{\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx))^n \sinh^2 \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{bc\sqrt{cx - 1}\sqrt{cx + 1}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{d - c^2 dx^2} \int -(a + \operatorname{barccosh}(cx))^n \sin \left( \frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} \right)^2 d(a + \operatorname{barccosh}(cx))}{bc\sqrt{cx - 1}\sqrt{cx + 1}}$$

$$\downarrow \text{25}$$

$$\frac{\sqrt{d-c^2dx^2} \int (a + \operatorname{barccosh}(cx))^n \sin\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^2 d(a + \operatorname{barccosh}(cx))}{bc\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 3793

$$\frac{\sqrt{d-c^2dx^2} \int \left(\frac{1}{2}(a + \operatorname{barccosh}(cx))^n - \frac{1}{2}(a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{2a}{b} - \frac{2(a+\operatorname{barccosh}(cx))}{b}\right)\right) d(a + \operatorname{barccosh}(cx))}{bc\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2009

$$\frac{\sqrt{d-c^2dx^2} \left( -\frac{(a+\operatorname{barccosh}(cx))^{n+1}}{2(n+1)} + b2^{-n-3}e^{-\frac{2a}{b}}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n+1, -\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) \right)}{bc\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`

output `(Sqrt[d - c^2*d*x^2]*(-1/2*(a + b*ArcCosh[c*x])^(1 + n)/(1 + n) + (2^(-3 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x]))/b])/(E^((2*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n - (2^(-3 - n)*b*E^((2*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x])/b])/((a + b*ArcCosh[c*x])/b)^n))/(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.421.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] :> Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)]  
Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x]  
/; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

### 3.421.4 Maple [F]

$$\int (a + b \operatorname{arccosh}(cx))^n \sqrt{-c^2 dx^2 + d} dx$$

input `int((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x)`

output `int((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x)`

### 3.421.5 Fracas [F]

$$\int \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^n dx$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n, x)`

### 3.421.6 Sympy [F]

$$\int \sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n dx = \int \sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^n dx$$

input `integrate((a+b*acosh(c*x))**n*(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**n, x)`

**3.421.7 Maxima [F]**

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int \sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n, x)`

**3.421.8 Giac [F(-2)]**

Exception generated.

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.421.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n \sqrt{d - c^2 dx^2} dx$$

input `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2),x)`

output `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2), x)`



**3.422**  $\int \frac{\sqrt{d-c^2x^2}(a+b\operatorname{arccosh}(cx))^n}{x} dx$

3.422.1 Optimal result	3224
3.422.2 Mathematica [N/A]	3225
3.422.3 Rubi [N/A]	3225
3.422.4 Maple [N/A] (verified)	3226
3.422.5 Fricas [N/A]	3226
3.422.6 Sympy [N/A]	3227
3.422.7 Maxima [N/A]	3227
3.422.8 Giac [F(-2)]	3227
3.422.9 Mupad [N/A]	3228

**3.422.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{\sqrt{d-c^2x^2}(a+b\operatorname{arccosh}(cx))^n}{x} dx =$$

$$\frac{de^{-\frac{a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2\sqrt{d-c^2x^2}}$$

$$+ \frac{de^{a/b}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2\sqrt{d-c^2x^2}}$$

$$+ d\operatorname{Int}\left(\frac{(a+b\operatorname{arccosh}(cx))^n}{x\sqrt{d-c^2x^2}},x\right)$$

output

```
-1/2*d*(a+b*arccosh(c*x))^n*GAMMA(1+n,(-a-b*arccosh(c*x))/b)*(c*x-1)^(1/2)
*(c*x+1)^(1/2)/exp(a/b)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+1
/2*d*exp(a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,(a+b*arccosh(c*x))/b)*(c*x-1)
^(1/2)*(c*x+1)^(1/2)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+d*Uni
ntegrable((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2),x)
```

**3.422.2 Mathematica [N/A]**

Not integrable

Time = 0.34 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x} dx$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n)/x, x]`output `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n)/x, x]`**3.422.3 Rubi [N/A]**

Not integrable

Time = 0.78 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x} dx \\ & \quad \downarrow \text{6369} \\ & \int \left( \frac{d(a + \operatorname{barccosh}(cx))^n}{x\sqrt{d - c^2 dx^2}} - \frac{c^2 dx (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} \right) dx \\ & \quad \downarrow \text{2009} \\ & d \int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{d - c^2 dx^2}} dx - \\ & \frac{de^{-\frac{a}{b}} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{2\sqrt{d - c^2 dx^2}} + \\ & \frac{de^{a/b} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{2\sqrt{d - c^2 dx^2}} \end{aligned}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n)/x, x]`

---

3.422.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x} dx$

output `$Aborted`

### 3.422.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.422.4 Maple [N/A] (verified)

Not integrable

Time = 1.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n \sqrt{-c^2 d x^2 + d}}{x} dx$$

input `int((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x,x)`

output `int((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x,x)`

### 3.422.5 Fracas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^n}{x} dx$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/x, x)`

---

3.422.  $\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x} dx$

**3.422.6 Sympy [N/A]**

Not integrable

Time = 4.59 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^n}{x} dx$$

input `integrate((a+b*acosh(c*x))**n*(-c**2*d*x**2+d)**(1/2)/x,x)`output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**n/x, x)`**3.422.7 Maxima [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^n}{x} dx$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="maxima")`output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/x, x)`**3.422.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x,x, algorithm="giac")`output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.422.  $\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x} dx$

**3.422.9 Mupad [N/A]**

Not integrable

Time = 3.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n \sqrt{d - c^2 dx^2}}{x} dx$$

input `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2))/x,x)`output `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2))/x, x)`

**3.423** 
$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx$$

3.423.1 Optimal result	3229
3.423.2 Mathematica [N/A]	3229
3.423.3 Rubi [N/A]	3230
3.423.4 Maple [N/A] (verified)	3231
3.423.5 Fricas [N/A]	3231
3.423.6 Sympy [N/A]	3231
3.423.7 Maxima [N/A]	3232
3.423.8 Giac [F(-2)]	3232
3.423.9 Mupad [N/A]	3232

**3.423.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx = -\frac{cd\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^{1+n}}{b(1+n)\sqrt{d-c^2dx^2}} + d\operatorname{Int}\left(\frac{(a+b\operatorname{arccosh}(cx))^n}{x^2\sqrt{d-c^2dx^2}}, x\right)$$

output `-c*d*(a+b*arccosh(c*x))^(1+n)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/(1+n)/(-c^2*d*x^2+d)^(1/2)+d*Unintegrable((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(1/2), x)`

**3.423.2 Mathematica [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx = \int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx$$

input `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n)/x^2, x]`

output `Integrate[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n)/x^2, x]`

---

3.423. 
$$\int \frac{\sqrt{d-c^2dx^2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx$$

**3.423.3 Rubi [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx$$

↓ 6369

$$\int \left( \frac{d(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} - \frac{c^2 d(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} \right) dx$$

↓ 2009

$$d \int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx - \frac{cd \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^{n+1}}{b(n + 1) \sqrt{d - c^2 dx^2}}$$

input `Int[(Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n)/x^2,x]`

output `$Aborted`

**3.423.3.1 Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

**3.423.4 Maple [N/A] (verified)**

Not integrable

Time = 1.41 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n \sqrt{-c^2 dx^2 + d}}{x^2} dx$$

input `int((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x^2,x)`output `int((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x^2,x)`**3.423.5 Fricas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arccosh}(cx) + a)^n}{x^2} dx$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="fricas")`output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/x^2, x)`**3.423.6 Sympy [N/A]**

Not integrable

Time = 6.43 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt{d - c^2 dx^2} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx = \int \frac{\sqrt{-d(cx - 1)(cx + 1)} (a + b \operatorname{acosh}(cx))^n}{x^2} dx$$

input `integrate((a+b*acosh(c*x))**n*(-c**2*d*x**2+d)**(1/2)/x**2,x)`output `Integral(sqrt(-d*(c*x - 1)*(c*x + 1))*(a + b*acosh(c*x))**n/x**2, x)`



**3.423.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \int \frac{\sqrt{-c^2 dx^2 + d} (b \operatorname{arcosh}(cx) + a)^n}{x^2} dx$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="maxima")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/x^2, x)`

**3.423.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^n*(-c^2*d*x^2+d)^(1/2)/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.423.9 Mupad [N/A]**

Not integrable

Time = 3.23 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n \sqrt{d - c^2 dx^2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2))/x^2,x)`

output `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2))/x^2, x)`

---

3.423.  $\int \frac{\sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx$

### 3.424 $\int x^2(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$

3.424.1 Optimal result . . . . .	3233
3.424.2 Mathematica [A] (verified) . . . . .	3234
3.424.3 Rubi [A] (verified) . . . . .	3235
3.424.4 Maple [F] . . . . .	3236
3.424.5 Fricas [F] . . . . .	3236
3.424.6 Sympy [F(-1)] . . . . .	3237
3.424.7 Maxima [F] . . . . .	3237
3.424.8 Giac [F(-2)] . . . . .	3237
3.424.9 Mupad [F(-1)] . . . . .	3238

#### 3.424.1 Optimal result

Integrand size = 29, antiderivative size = 658

$$\int x^2(d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = -\frac{d\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^{1+n}}{16bc^3(1+n)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$-\frac{2^{-7-n}3^{-1-n}de^{-\frac{6a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n\left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n, -\frac{6(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+\frac{2^{-7-2n}de^{-\frac{4a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n\left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n, -\frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+\frac{2^{-7-n}de^{-\frac{2a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n\left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n, -\frac{2(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$-\frac{2^{-7-n}de^{\frac{2a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n, \frac{2(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$-\frac{2^{-7-2n}de^{\frac{4a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n, \frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+\frac{2^{-7-n}3^{-1-n}de^{\frac{6a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n, \frac{6(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/16*d*(a+b*arccosh(c*x))^(1+n)*(-c^2*d*x^2+d)^(1/2)/b/c^3/(1+n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2^(-7-n)*3^(-1-n)*d*(a+b*arccosh(c*x))^n*GAMMA(1+n,-6*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/exp(6*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2^(-7-2*n)*d*(a+b*arccosh(c*x))^n*GAMMA(1+n,-4*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/exp(4*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2^(-7-n)*d*(a+b*arccosh(c*x))^n*GAMMA(1+n,-2*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/exp(2*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2^(-7-2*n)*d*exp(2*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,2*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2^(-7-2*n)*d*exp(4*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,4*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2^(-7-n)*3^(-1-n)*d*exp(6*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,6*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.424.2 Mathematica [A] (verified)

Time = 2.31 (sec) , antiderivative size = 438, normalized size of antiderivative = 0.67

$$\int x^2(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n dx = \frac{2^{-7-2n} 3^{-1-n} d^2 e^{-\frac{6a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (a + b \operatorname{arccosh}(cx))^n \left( -\frac{(a+b \operatorname{arccosh}(cx))^2}{b^2} \right)^{-n} \left( 2^n b \operatorname{arccosh}(cx) \right)^n}{\dots}$$

input `Integrate[x^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output

```
(2^(-7 - 2*n)*3^(-1 - n)*d^2*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])^n*(2^n*b*(1 + n)*(a/b + ArcCosh[c*x])^n*Gamma[1 + n, (-6*(a + b*ArcCosh[c*x]))/b] - 3^(1 + n)*b*E^((2*a)/b)*(1 + n)*(a/b + ArcCosh[c*x])^n*Gamma[1 + n, (-4*(a + b*ArcCosh[c*x]))/b] - 2^n*3^(1 + n)*b*E^((4*a)/b)*(1 + n)*(a/b + ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x]))/b] + 2^n*3^(1 + n)*b*E^((8*a)/b)*(1 + n)*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x]))/b] + 3^(1 + n)*b*E^((10*a)/b)*(1 + n)*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, (4*(a + b*ArcCosh[c*x]))/b] + 2^n*E^((6*a)/b)*(2^(3 + n)*3^(1 + n)*(a + b*ArcCosh[c*x])*(-((a + b*ArcCosh[c*x])^2/b^2)))^n - b*E^((6*a)/b)*(1 + n)*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, (6*(a + b*ArcCosh[c*x]))/b]))/(b*c^3*E^((6*a)/b)*(1 + n)*Sqrt[d - c^2*d*x^2]*(-(a + b*ArcCosh[c*x])^2/b^2))^n
```

---

3.424.  $\int x^2(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n dx$

**3.424.3 Rubi [A] (verified)**

Time = 0.90 (sec) , antiderivative size = 444, normalized size of antiderivative = 0.67, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$$

$$\downarrow \text{6367}$$

$$\frac{d\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx))^n \cosh^2\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^3\sqrt{cx - 1}\sqrt{cx + 1}}$$

$$\downarrow \text{5971}$$

$$\frac{d\sqrt{d - c^2 dx^2} \int \left(\frac{1}{32} \cosh\left(\frac{6a}{b} - \frac{6(a + \operatorname{barccosh}(cx))}{b}\right)\right) (a + \operatorname{barccosh}(cx))^n - \frac{1}{16} \cosh\left(\frac{4a}{b} - \frac{4(a + \operatorname{barccosh}(cx))}{b}\right) (a + \operatorname{barccosh}(cx))}{bc^3\sqrt{cx - 1}\sqrt{cx + 1}}$$

$$\downarrow \text{2009}$$

$$\frac{d\sqrt{d - c^2 dx^2} \left( \frac{(a + \operatorname{barccosh}(cx))^{n+1}}{16(n+1)} + b2^{-n-7}3^{-n-1}e^{-\frac{6a}{b}} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n + 1, -\frac{6(a + \operatorname{barccosh}(cx))}{b}\right) \right)}{bc^3\sqrt{cx - 1}\sqrt{cx + 1}}$$

input `Int[x^2*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output

$$\begin{aligned} & -\left(\frac{d\sqrt{d - c^2 dx^2} \left( (a + b \operatorname{ArcCosh}[c x])^{n+1} \right)}{16(n+1)} + b 2^{-n-7} 3^{-n-1} e^{-\frac{6a}{b}} (a + b \operatorname{ArcCosh}[c x])^n \left(-\frac{a + b \operatorname{ArcCosh}[c x]}{b}\right)^{-n} \Gamma\left(n + 1, -\frac{6(a + b \operatorname{ArcCosh}[c x])}{b}\right) \right)}{bc^3\sqrt{cx - 1}\sqrt{cx + 1}} \end{aligned}$$

$$3.424. \quad \int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$$

## 3.424.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

## 3.424.4 Maple [F]

$$\int x^2(-c^2dx^2 + d)^{\frac{3}{2}}(a + b \operatorname{arccosh}(cx))^n dx$$

input `int(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

output `int(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

## 3.424.5 Fracas [F]

$$\int x^2(d - c^2dx^2)^{3/2}(a + b\operatorname{arccosh}(cx))^n dx = \int (-c^2dx^2 + d)^{\frac{3}{2}}(b \operatorname{arccosh}(cx) + a)^n x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="fricas")`

output `integral(-(c^2*d*x^4 - d*x^2)*sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n, x)`

**3.424.6 Sympy [F(-1)]**

Timed out.

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate(x**2*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.424.7 Maxima [F]**

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^n x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n*x^2, x)`

**3.424.8 Giac [F(-2)]**

Exception generated.

$$\int x^2(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate(x^2*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

**3.424.9 Mupad [F(-1)]**

Timed out.

$$\int x^2 (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int x^2 (a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{3/2} dx$$

input `int(x^2*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2),x)`output `int(x^2*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2), x)`

### 3.425 $\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$

3.425.1 Optimal result . . . . .	3239
3.425.2 Mathematica [A] (verified) . . . . .	3240
3.425.3 Rubi [A] (verified) . . . . .	3241
3.425.4 Maple [F] . . . . .	3242
3.425.5 Fricas [F] . . . . .	3242
3.425.6 Sympy [F(-1)] . . . . .	3243
3.425.7 Maxima [F] . . . . .	3243
3.425.8 Giac [F(-2)] . . . . .	3243
3.425.9 Mupad [F(-1)] . . . . .	3244

#### 3.425.1 Optimal result

Integrand size = 27, antiderivative size = 578

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx =$$

$$\frac{5^{-1-n} de^{-\frac{5a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, -\frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{32c^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{3^{-n} de^{-\frac{3a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, -\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{32c^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$- \frac{de^{-\frac{a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, -\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{16c^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{de^{a/b} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{16c^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$- \frac{3^{-n} de^{\frac{3a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, \frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{32c^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$

$$+ \frac{5^{-1-n} de^{\frac{5a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, \frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{32c^2 \sqrt{-1 + cx} \sqrt{1 + cx}}$$



output

$$\begin{aligned}
& -1/32*5^{(-1-n)}*d*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,-5*(a+b*\operatorname{arccosh}(c*x))/b)* \\
& (-c^2*d*x^2+d)^{(1/2)}/c^2/\exp(5*a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)} \\
& /((c*x+1)^{(1/2)}+1/32*d*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,-3*(a+b*\operatorname{arccosh}(c*x))/b) \\
& )/b*(-c^2*d*x^2+d)^{(1/2)}/(3^n)/c^2/\exp(3*a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n) \\
& /((c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/16*d*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,(-a-b* \\
& \operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c^2/\exp(a/b)/(((a+b*\operatorname{arccosh}(c*x))/b) \\
& )^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/16*d*\exp(a/b)*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMM} \\
& A(1+n,(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c^2/(((a+b*\operatorname{arccosh}(c*x))/ \\
& b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/32*d*\exp(3*a/b)*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{G} \\
& AMMA(1+n,3*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/(3^n)/c^2/(((a+b*\operatorname{arc} \\
& \operatorname{cosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/32*5^{(-1-n)}*d*\exp(5*a/b)*(a \\
& +b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,5*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/ \\
& c^2/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}
\end{aligned}$$

### 3.425.2 Mathematica [A] (verified)

Time = 1.65 (sec) , antiderivative size = 500, normalized size of antiderivative = 0.87

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx =$$

$$\frac{15^{-1-n} d^2 e^{-\frac{5a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (a + \operatorname{barccosh}(cx))^n \left( -\frac{(a + \operatorname{barccosh}(cx))^2}{b^2} \right)^{-3n} \left( 2 \cdot 15^{1+n} e^{\frac{6a}{b}} \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right) \right)}{1}$$

input `Integrate[x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output

$$\begin{aligned}
& -1/32*(15^{(-1-n)}*d^2*\operatorname{Sqrt}[(-1+c*x)/(1+c*x)]*(1+c*x)*(a+b*\operatorname{ArcCosh} \\
& [c*x])^n*(2*15^{(1+n)}*E^{((6*a)/b)}*(-((a+b*\operatorname{ArcCosh}[c*x])/b))^n*(-((a+b \\
& *\operatorname{ArcCosh}[c*x])^2/b^2))^{(2*n)}*\operatorname{Gamma}[1+n,a/b+\operatorname{ArcCosh}[c*x]]+(a/b+\operatorname{Arc} \\
& \operatorname{Cosh}[c*x])^n*(-(3^{(1+n)}*(-((a+b*\operatorname{ArcCosh}[c*x])^2/b^2))^{(2*n)}*\operatorname{Gamma}[1+ \\
& n,(-5*(a+b*\operatorname{ArcCosh}[c*x]))/b])+(3*5^{(1+n)}*E^{((2*a)/b)}*(-((a+b*\operatorname{ArcCo} \\
& sh[c*x])^2/b^2))^{(2*n)}*\operatorname{Gamma}[1+n,(-3*(a+b*\operatorname{ArcCosh}[c*x]))/b]-2*15^{(1 \\
& +n)}*E^{((4*a)/b)}*(-((a+b*\operatorname{ArcCosh}[c*x])^2/b^2))^{(2*n)}*\operatorname{Gamma}[1+n,-((a \\
& +b*\operatorname{ArcCosh}[c*x])/b])+(5^{(1+n)}*E^{((8*a)/b)}*(a/b+\operatorname{ArcCosh}[c*x])^n*(-((a \\
& +b*\operatorname{ArcCosh}[c*x])/b))^{(3*n)}*\operatorname{Gamma}[1+n,(3*(a+b*\operatorname{ArcCosh}[c*x]))/b]-4* \\
& 5^{(1+n)}*E^{((8*a)/b)}*(-((a+b*\operatorname{ArcCosh}[c*x])/b))^{(2*n)}*(-((a+b*\operatorname{ArcCosh}[ \\
& c*x])^2/b^2))^{(2*n)}*\operatorname{Gamma}[1+n,(3*(a+b*\operatorname{ArcCosh}[c*x]))/b]+3^{(1+n)}*E^{((1 \\
& 0*a)/b)}*(a/b+\operatorname{ArcCosh}[c*x])^n*(-((a+b*\operatorname{ArcCosh}[c*x])/b))^{(3*n)}*\operatorname{Gamma}[1+ \\
& n,(5*(a+b*\operatorname{ArcCosh}[c*x]))/b]))/(c^2*E^{((5*a)/b)}*\operatorname{Sqrt}[d-c^2*d*x^2]*(- \\
& ((a+b*\operatorname{ArcCosh}[c*x])^2/b^2))^{(3*n)})
\end{aligned}$$

$$3.425. \quad \int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$$

**3.425.3 Rubi [A] (verified)**

Time = 0.78 (sec) , antiderivative size = 404, normalized size of antiderivative = 0.70, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$$

↓ 6367

$$\frac{d\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) \sinh^4\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^2\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 5971

$$\frac{d\sqrt{d - c^2 dx^2} \int \left(\frac{1}{16} \cosh\left(\frac{5a}{b} - \frac{5(a + \operatorname{barccosh}(cx))}{b}\right)\right) (a + \operatorname{barccosh}(cx))^n - \frac{3}{16} \cosh\left(\frac{3a}{b} - \frac{3(a + \operatorname{barccosh}(cx))}{b}\right) (a + \operatorname{barccosh}(cx))^n}{bc^2\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 2009

$$\frac{d\sqrt{d - c^2 dx^2} \left(\frac{1}{32} b^5 n^{-1} e^{-\frac{5a}{b}} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n + 1, -\frac{5(a + \operatorname{barccosh}(cx))}{b}\right) - \frac{1}{32} b^3 n^{-1} e^{-\frac{3a}{b}} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n + 1, -\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)\right)}{bc^2\sqrt{cx - 1}\sqrt{cx + 1}}$$

input `Int[x*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output

```

-((d*Sqrt[d - c^2*d*x^2]*((5^(-1 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n
, (-5*(a + b*ArcCosh[c*x]))/b])/(32*E^((5*a)/b)*(-(a + b*ArcCosh[c*x])/b)
)^n) - (b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-3*(a + b*ArcCosh[c*x]))/b
])/(32*3^n*E^((3*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n) + (b*(a + b*ArcCosh[c
*x])^n*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b])/(16*E^(a/b)*(-(a + b*ArcC
osh[c*x])/b))^n) - (b*E^(a/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (a + b*A
rcCosh[c*x])/b])/(16*((a + b*ArcCosh[c*x])/b)^n) + (b*E^((3*a)/b)*(a + b*A
rcCosh[c*x])^n*Gamma[1 + n, (3*(a + b*ArcCosh[c*x]))/b])/(32*3^n*((a + b*A
rcCosh[c*x])/b)^n) - (5^(-1 - n)*b*E^((5*a)/b)*(a + b*ArcCosh[c*x])^n*Gamm
a[1 + n, (5*(a + b*ArcCosh[c*x]))/b])/(32*((a + b*ArcCosh[c*x])/b)^n))/(b
*c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])

```

## 3.425.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

## 3.425.4 Maple [F]

$$\int x(-c^2 dx^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^n dx$$

input `int(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

output `int(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

## 3.425.5 Fracas [F]

$$\int x(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^n x dx$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="fracas")`

output `integral(-(c^2*d*x^3 - d*x)*sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n, x)`

**3.425.6 Sympy [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate(x*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.425.7 Maxima [F]**

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^n x dx$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n*x, x)`

**3.425.8 Giac [F(-2)]**

Exception generated.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.425.9 Mupad [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int x(a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{3/2} dx$$

input `int(x*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2),x)`output `int(x*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2), x)`

### 3.426 $\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$

3.426.1 Optimal result	3245
3.426.2 Mathematica [A] (verified)	3246
3.426.3 Rubi [A] (verified)	3246
3.426.4 Maple [F]	3248
3.426.5 Fricas [F]	3248
3.426.6 Sympy [F(-1)]	3248
3.426.7 Maxima [F]	3249
3.426.8 Giac [F(-2)]	3249
3.426.9 Mupad [F(-1)]	3249

#### 3.426.1 Optimal result

Integrand size = 26, antiderivative size = 450

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = -\frac{3d\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^{1+n}}{8bc(1+n)\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$-\frac{2^{-2(3+n)}de^{-\frac{4a}{b}}\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+\frac{2^{-3-n}de^{-\frac{2a}{b}}\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$-\frac{2^{-3-n}de^{\frac{2a}{b}}\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+\frac{2^{-2(3+n)}de^{\frac{4a}{b}}\sqrt{d - c^2 dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
-3/8*d*(a+b*arccosh(c*x))^(1+n)*(-c^2*d*x^2+d)^(1/2)/b/c/(1+n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-d*(a+b*arccosh(c*x))^n*GAMMA(1+n,-4*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/(2^(6+2*n))/c/exp(4*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2^(-3-n)*d*(a+b*arccosh(c*x))^n*GAMMA(1+n,-2*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c/exp(2*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2^(-3-n)*d*exp(2*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,2*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+d*exp(4*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,4*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/(2^(6+2*n))/c/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

3.426.  $\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$

### 3.426.2 Mathematica [A] (verified)

Time = 1.60 (sec) , antiderivative size = 384, normalized size of antiderivative = 0.85

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \frac{4^{-3-n} d^2 e^{-\frac{4a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx) (a + \operatorname{barccosh}(cx))^n \left( -\frac{(a+\operatorname{barccosh}(cx))^2}{b^2} \right)^{-2n} \left( 3 \cdot 2^{3+2n} e^{\frac{4a}{b}} \right)}{3 \cdot 2^{3+2n} e^{\frac{4a}{b}}}$$

input `Integrate[(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output `(4^(-3 - n)*d^2*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])^n*(3*2^(3 + 2*n)*E^((4*a)/b)*(a + b*ArcCosh[c*x])*(-(a + b*ArcCosh[c*x])^2/b^2))^(2*n) + b*(1 + n)*(a/b + ArcCosh[c*x])^(2*n)*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, (-4*(a + b*ArcCosh[c*x]))/b] - 2^(3 + n)*b*E^((2*a)/b)*(1 + n)*(a/b + ArcCosh[c*x])^n*(-(a + b*ArcCosh[c*x])^2/b^2))^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x]))/b] + 2^(3 + n)*b*E^((6*a)/b)*(1 + n)*(-(a + b*ArcCosh[c*x])/b))^n*(-(a + b*ArcCosh[c*x])^2/b^2))^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x]))/b] - b*E^((8*a)/b)*(1 + n)*(a/b + ArcCosh[c*x])^n*(-(a + b*ArcCosh[c*x])/b)^(2*n)*Gamma[1 + n, (4*(a + b*ArcCosh[c*x]))/b])/b*c*E^((4*a)/b)*(1 + n)*sqrt[d - c^2*d*x^2]*(-(a + b*ArcCosh[c*x])^2/b^2))^(2*n))`

### 3.426.3 Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 308, normalized size of antiderivative = 0.68, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {6321, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$$

↓ 6321

$$\frac{d\sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx))^n \sinh^4 \left( \frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b} \right) d(a + \operatorname{barccosh}(cx))}{bc\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 3042

---

3.426.  $\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$

$$\frac{d\sqrt{d-c^2dx^2} \int (a + \operatorname{barccosh}(cx))^n \sin\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b}\right)^4 d(a + \operatorname{barccosh}(cx))}{bc\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 3793

$$\frac{d\sqrt{d-c^2dx^2} \int \left(\frac{1}{8} \cosh\left(\frac{4a}{b} - \frac{4(a+\operatorname{barccosh}(cx))}{b}\right) (a + \operatorname{barccosh}(cx))^n - \frac{1}{2} \cosh\left(\frac{2a}{b} - \frac{2(a+\operatorname{barccosh}(cx))}{b}\right) (a + \operatorname{barccosh}(cx))^n\right)}{bc\sqrt{cx-1}\sqrt{cx+1}}$$

↓ 2009

$$\frac{d\sqrt{d-c^2dx^2} \left(\frac{3(a+\operatorname{barccosh}(cx))^{n+1}}{8(n+1)} + b2^{-2(n+3)}e^{-\frac{4a}{b}}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n+1, -\frac{4(a+\operatorname{barccosh}(cx))}{b}\right)\right)}{bc\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output `-(d*Sqrt[d - c^2*d*x^2]*((3*(a + b*ArcCosh[c*x])^(1 + n))/(8*(1 + n)) + (b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-4*(a + b*ArcCosh[c*x])/b)]/(2^(2*(3 + n))*E^((4*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n - (2^(-3 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x])/b)]/(E^((2*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n + (2^(-3 - n)*b*E^((2*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x])/b)]/((a + b*ArcCosh[c*x])/b)^n - (b*E^((4*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (4*(a + b*ArcCosh[c*x])/b)]/(2^(2*(3 + n))*((a + b*ArcCosh[c*x])/b)^n)))/(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.426.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

---

3.426.  $\int (d - c^2dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$



rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] :> Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)]  
Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x]  
/; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

### 3.426.4 Maple [F]

$$\int (-c^2 dx^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^n dx$$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

output `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

### 3.426.5 Fricas [F]

$$\int (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^n dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="fricas")`

output `integral((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n, x)`

### 3.426.6 Sympy [F(-1)]

Timed out.

$$\int (d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.426.7 Maxima [F]**

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n, x)`

**3.426.8 Giac [F(-2)]**

Exception generated.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.426.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{3/2} dx$$

input `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2),x)`

output `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2), x)`

$$3.427 \quad \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^n}{x} dx$$

3.427.1 Optimal result . . . . .	3250
3.427.2 Mathematica [N/A] . . . . .	3251
3.427.3 Rubi [N/A] . . . . .	3251
3.427.4 Maple [N/A] (verified) . . . . .	3253
3.427.5 Fricas [N/A] . . . . .	3253
3.427.6 Sympy [F(-1)] . . . . .	3253
3.427.7 Maxima [N/A] . . . . .	3254
3.427.8 Giac [F(-2)] . . . . .	3254
3.427.9 Mupad [N/A] . . . . .	3254

### 3.427.1 Optimal result

Integrand size = 29, antiderivative size = 29

$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^n}{x} dx = \frac{3^{-1-n}d^2e^{-\frac{3a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8\sqrt{d-c^2dx^2}}$$

$$- \frac{5d^2e^{-\frac{a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8\sqrt{d-c^2dx^2}}$$

$$+ \frac{5d^2e^{a/b}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8\sqrt{d-c^2dx^2}}$$

$$- \frac{3^{-1-n}d^2e^{\frac{3a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{8\sqrt{d-c^2dx^2}}$$

$$+ d^2\operatorname{Int}\left(\frac{(a+b\operatorname{arccosh}(cx))^n}{x\sqrt{d-c^2dx^2}},x\right)$$

---


$$3.427. \quad \int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^n}{x} dx$$

output  $1/8*3^{(-1-n)}*d^2*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,-3*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/\exp(3*a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}-5/8*d^2*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,(-a-b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/\exp(a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}+5/8*d^2*\exp(a/b)*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}-1/8*3^{(-1-n)}*d^2*\exp(3*a/b)*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,3*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}+d^2*\operatorname{Unintegrable}((a+b*\operatorname{arccosh}(c*x))^n/x/(-c^2*d*x^2+d)^{(1/2)},x)$

### 3.427.2 Mathematica [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n)/x,x]`

output `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n)/x, x]`

### 3.427.3 Rubi [N/A]

Not integrable

Time = 1.35 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$$

↓ 6369

---

3.427.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$

$$\begin{aligned}
& \int \left( -\frac{2c^2 d^2 x (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} + \frac{d^2 (a + \operatorname{barccosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} + \frac{c^4 d^2 x^3 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} \right) dx \\
& \quad \downarrow \text{2009} \\
& \quad d^2 \int \frac{(a + \operatorname{barccosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} dx + \\
& \frac{d^2 3^{-n-1} e^{-\frac{3a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n+1, -\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{8\sqrt{d - c^2 dx^2}} \\
& \frac{5d^2 e^{-\frac{a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n+1, -\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{8\sqrt{d - c^2 dx^2}} + \\
& \frac{5d^2 e^{a/b} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n+1, \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{8\sqrt{d - c^2 dx^2}} - \\
& \frac{d^2 3^{-n-1} e^{\frac{3a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n+1, \frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{8\sqrt{d - c^2 dx^2}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n)/x,x]`

output `$Aborted`

### 3.427.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

**3.427.4 Maple [N/A] (verified)**

Not integrable

Time = 2.47 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 d x^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^n}{x} dx$$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x,x)`output `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x,x)`**3.427.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^n}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x,x, algorithm="fricas")`output `integral((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n/x, x)`**3.427.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**n/x,x)`output `Timed out`

**3.427.7 Maxima [N/A]**

Not integrable

Time = 0.62 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^n}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n/x, x)`

**3.427.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.427.9 Mupad [N/A]**

Not integrable

Time = 3.16 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{3/2}}{x} dx$$

input `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2))/x,x)`

output `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2))/x, x)`

---

3.427.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$

**3.428** 
$$\int \frac{(d-c^2dx^2)^{3/2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx$$

3.428.1 Optimal result . . . . .	3255
3.428.2 Mathematica [N/A] . . . . .	3256
3.428.3 Rubi [N/A] . . . . .	3256
3.428.4 Maple [N/A] (verified) . . . . .	3257
3.428.5 Fracas [N/A] . . . . .	3257
3.428.6 Sympy [F(-1)] . . . . .	3258
3.428.7 Maxima [N/A] . . . . .	3258
3.428.8 Giac [F(-2)] . . . . .	3258
3.428.9 Mupad [N/A] . . . . .	3259

**3.428.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(d - c^2dx^2)^{3/2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx = -\frac{3cd^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^{1+n}}{2b(1 + n)\sqrt{d - c^2dx^2}}$$

$$+ \frac{2^{-3-n}cd^2e^{-\frac{2a}{b}}\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^n \left(-\frac{a + \operatorname{arccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{2(a + \operatorname{arccosh}(cx))}{b}\right)}{\sqrt{d - c^2dx^2}}$$

$$- \frac{2^{-3-n}cd^2e^{\frac{2a}{b}}\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^n \left(\frac{a + \operatorname{arccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{2(a + \operatorname{arccosh}(cx))}{b}\right)}{\sqrt{d - c^2dx^2}}$$

$$+ d^2 \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{d - c^2dx^2}}, x\right)$$

output

```
-3/2*c*d^2*(a+b*arccosh(c*x))^(1+n)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/(1+n)/(-
c^2*d*x^2+d)^(1/2)+2^(-3-n)*c*d^2*(a+b*arccosh(c*x))^n*GAMMA(1+n,-2*(a+b*a
rccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/exp(2*a/b)/(((a-b*arccosh(c*x
))/b)^n)/(-c^2*d*x^2+d)^(1/2)-2^(-3-n)*c*d^2*exp(2*a/b)*(a+b*arccosh(c*x)
)^n*GAMMA(1+n,2*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(((a+b*ar
ccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+d^2*Unintegrate((a+b*arccosh(c*x)
)^n/x^2/(-c^2*d*x^2+d)^(1/2),x)
```



**3.428.2 Mathematica [N/A]**

Not integrable

Time = 0.83 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx$$

input `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n)/x^2,x]`output `Integrate[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n)/x^2, x]`**3.428.3 Rubi [N/A]**

Not integrable

Time = 1.05 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx \\ & \quad \downarrow \text{6369} \\ & \int \left( -\frac{2c^2 d^2 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} + \frac{d^2 (a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} + \frac{c^4 d^2 x^2 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} \right) dx \\ & \quad \downarrow \text{2009} \\ & d^2 \int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx - \frac{3cd^2 \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^{n+1}}{2b(n + 1) \sqrt{d - c^2 dx^2}} + \\ & \frac{cd^2 2^{-n-3} e^{-\frac{2a}{b}} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{d - c^2 dx^2}} \\ & \frac{cd^2 2^{-n-3} e^{\frac{2a}{b}} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, \frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{d - c^2 dx^2}} \end{aligned}$$

input `Int[((d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n)/x^2,x]`

---

3.428.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx$

output `$Aborted`

### 3.428.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_))^(m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.428.4 Maple [N/A] (verified)

Not integrable

Time = 1.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 d x^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx$$

input `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x^2,x)`

output `int((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x^2,x)`

### 3.428.5 Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^n}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x^2,x, algorithm="fricas")`

output `integral((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n/x^2, x)`

---

3.428.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx$

**3.428.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**n/x**2,x)`

output `Timed out`

**3.428.7 Maxima [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^n}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x^2,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^n/x^2, x)`

**3.428.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

---

3.428.  $\int \frac{(d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx$

**3.428.9 Mupad [N/A]**

Not integrable

Time = 3.42 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{3/2} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{3/2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2))/x^2,x)`output `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2))/x^2, x)`

### 3.429 $\int x^2(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$

3.429.1 Optimal result	3260
3.429.2 Mathematica [A] (verified)	3261
3.429.3 Rubi [A] (verified)	3262
3.429.4 Maple [F]	3264
3.429.5 Fricas [F]	3264
3.429.6 Sympy [F(-1)]	3264
3.429.7 Maxima [F]	3265
3.429.8 Giac [F(-2)]	3265
3.429.9 Mupad [F(-1)]	3265

#### 3.429.1 Optimal result

Integrand size = 29, antiderivative size = 870

$$\int x^2(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = -\frac{5d^2\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^{1+n}}{128bc^3(1+n)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2^{-11-3n}d^2e^{-\frac{8a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{8(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2^{-7-n}3^{-1-n}d^2e^{-\frac{6a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{6(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2^{-2(4+n)}d^2e^{-\frac{4a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2^{-7-n}d^2e^{-\frac{2a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{2(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2^{-7-n}d^2e^{\frac{2a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{2(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2^{-2(4+n)}d^2e^{\frac{4a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{4(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2^{-7-n}3^{-1-n}d^2e^{\frac{6a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{6(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2^{-11-3n}d^2e^{\frac{8a}{b}}\sqrt{d - c^2dx^2}(a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{8(a+\operatorname{barccosh}(cx))}{b}\right)}{c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

---

3.429.  $\int x^2(d - c^2dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$

output

```

-5/128*d^2*(a+b*arccosh(c*x))^(1+n)*(-c^2*d*x^2+d)^(1/2)/b/c^3/(1+n)/(c*x-
1)^(1/2)/(c*x+1)^(1/2)+2^(-11-3*n)*d^2*(a+b*arccosh(c*x))^n*GAMMA(1+n,-8*(
a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/exp(8*a/b)/(((a-b*arccosh(c
*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2^(-7-n)*3^(-1-n)*d^2*(a+b*arccosh(
c*x))^n*GAMMA(1+n,-6*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/exp(6*
a/b)/(((a-b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+d^2*(a+b*arcc
osh(c*x))^n*GAMMA(1+n,-4*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/(2^(8+
2*n))/c^3/exp(4*a/b)/(((a-b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/
2)+2^(-7-n)*d^2*(a+b*arccosh(c*x))^n*GAMMA(1+n,-2*(a+b*arccosh(c*x))/b)*(-
c^2*d*x^2+d)^(1/2)/c^3/exp(2*a/b)/(((a-b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2
)/(c*x+1)^(1/2)-2^(-7-n)*d^2*exp(2*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,2*(
a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^3/(((a+b*arccosh(c*x))/b)^n)/(
c*x-1)^(1/2)/(c*x+1)^(1/2)-d^2*exp(4*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,4
*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/(2^(8+2*n))/c^3/(((a+b*arccosh
(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2^(-7-n)*3^(-1-n)*d^2*exp(6*a/b)*
(a+b*arccosh(c*x))^n*GAMMA(1+n,6*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2
)/c^3/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2^(-11-3*n)*d
^2*exp(8*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,8*(a+b*arccosh(c*x))/b)*(-c^2
*d*x^2+d)^(1/2)/c^3/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)

```

### 3.429.2 Mathematica [A] (verified)

Time = 5.62 (sec) , antiderivative size = 677, normalized size of antiderivative = 0.78

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \frac{2^{-11-3n} 3^{-1-n} d^3 e^{-\frac{8a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx)(a + \operatorname{barccosh}(cx))^n \left(-\frac{(a+\operatorname{barccosh}(cx))^2}{b^2}\right)^{-n} \left(-3\right)}{\dots}$$

input `Integrate[x^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n,x]`

output  $(2^{(-11 - 3*n)} * 3^{(-1 - n)} * d^3 * \text{Sqrt}[(-1 + c*x)/(1 + c*x)] * (1 + c*x) * (a + b * \text{ArcCosh}[c*x])^n * (-3^{(1 + n)} * b * (1 + n) * (a/b + \text{ArcCosh}[c*x])^n * \text{Gamma}[1 + n, (-8 * (a + b * \text{ArcCosh}[c*x]))/b]) + 4^{(2 + n)} * b * E^{((2*a)/b)} * (1 + n) * (a/b + \text{ArcCosh}[c*x])^n * \text{Gamma}[1 + n, (-6 * (a + b * \text{ArcCosh}[c*x]))/b] - 2^{(3 + n)} * 3^{(1 + n)} * b * E^{((4*a)/b)} * (1 + n) * (a/b + \text{ArcCosh}[c*x])^n * \text{Gamma}[1 + n, (-4 * (a + b * \text{ArcCosh}[c*x]))/b] - 3^{(1 + n)} * 4^{(2 + n)} * b * E^{((6*a)/b)} * (1 + n) * (a/b + \text{ArcCosh}[c*x])^n * \text{Gamma}[1 + n, (-2 * (a + b * \text{ArcCosh}[c*x]))/b] + E^{((8*a)/b)} * (5 * 2^{(4 + 3*n)} * 3^{(1 + n)} * a * (-((a + b * \text{ArcCosh}[c*x])^2/b^2))^n + 5 * 2^{(4 + 3*n)} * 3^{(1 + n)} * b * \text{ArcCosh}[c*x] * (-((a + b * \text{ArcCosh}[c*x])^2/b^2))^n + 3^{(1 + n)} * 4^{(2 + n)} * b * E^{((2*a)/b)} * (1 + n) * (-((a + b * \text{ArcCosh}[c*x])/b))^n * \text{Gamma}[1 + n, (2 * (a + b * \text{ArcCosh}[c*x]))/b] + 2^{(3 + n)} * 3^{(1 + n)} * b * E^{((4*a)/b)} * (1 + n) * (-((a + b * \text{ArcCosh}[c*x])/b))^n * \text{Gamma}[1 + n, (4 * (a + b * \text{ArcCosh}[c*x]))/b] - 4^{(2 + n)} * b * E^{((6*a)/b)} * (-((a + b * \text{ArcCosh}[c*x])/b))^n * \text{Gamma}[1 + n, (6 * (a + b * \text{ArcCosh}[c*x]))/b] - 4^{(2 + n)} * b * E^{((6*a)/b)} * n * (-((a + b * \text{ArcCosh}[c*x])/b))^n * \text{Gamma}[1 + n, (6 * (a + b * \text{ArcCosh}[c*x]))/b] + 3^{(1 + n)} * b * E^{((8*a)/b)} * (-((a + b * \text{ArcCosh}[c*x])/b))^n * \text{Gamma}[1 + n, (8 * (a + b * \text{ArcCosh}[c*x]))/b] + 3^{(1 + n)} * b * E^{((8*a)/b)} * n * (-((a + b * \text{ArcCosh}[c*x])/b))^n * \text{Gamma}[1 + n, (8 * (a + b * \text{ArcCosh}[c*x]))/b])) / (b * c^3 * E^{((8*a)/b)} * (1 + n) * \text{Sqrt}[d - c^2 * d * x^2] * (-((a + b * \text{ArcCosh}[c*x])^2/b^2))^n)$

### 3.429.3 Rubi [A] (verified)

Time = 1.05 (sec) , antiderivative size = 567, normalized size of antiderivative = 0.65, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.103$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2 (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^n dx$$

$$\downarrow 6367$$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int (a + \text{barccosh}(cx))^n \cosh^2 \left( \frac{a}{b} - \frac{a + \text{barccosh}(cx)}{b} \right) \sinh^6 \left( \frac{a}{b} - \frac{a + \text{barccosh}(cx)}{b} \right) d(a + \text{barccosh}(cx))}{bc^3 \sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow 5971$$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int \left( \frac{1}{128} \cosh \left( \frac{8a}{b} - \frac{8(a + \text{barccosh}(cx))}{b} \right) (a + \text{barccosh}(cx))^n - \frac{1}{32} \cosh \left( \frac{6a}{b} - \frac{6(a + \text{barccosh}(cx))}{b} \right) (a + \text{barccosh}(cx))^n \right) dx}{bc^3 \sqrt{cx - 1} \sqrt{cx + 1}}$$

---

3.429.  $\int x^2 (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^n dx$

↓ 2009

$$d^2 \sqrt{d - c^2 dx^2} \left( -\frac{5(a + \operatorname{barccosh}(cx))^{n+1}}{128(n+1)} + b2^{-3n-11} e^{-\frac{8a}{b}} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{8(a+b)}{b}\right) \right)$$

input `Int[x^2*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n,x]`

output `(d^2*sqrt[d - c^2*d*x^2]*((-5*(a + b*ArcCosh[c*x])^(1 + n))/(128*(1 + n)) + (2^(-11 - 3*n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-8*(a + b*ArcCosh[c*x]))/b])/ (E^((8*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n - (2^(-7 - n)*3^(-1 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-6*(a + b*ArcCosh[c*x]))/b])/ (E^((6*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n + (b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-4*(a + b*ArcCosh[c*x]))/b])/ (2^(2*(4 + n))*E^((4*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n + (2^(-7 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x]))/b])/ (E^((2*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n - (2^(-7 - n)*b*E^((2*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x]))/b])/ ((a + b*ArcCosh[c*x])/b)^n - (b*E^((4*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (4*(a + b*ArcCosh[c*x]))/b])/ (2^(2*(4 + n))*((a + b*ArcCosh[c*x])/b)^n) + (2^(-7 - n)*3^(-1 - n)*b*E^((6*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (6*(a + b*ArcCosh[c*x]))/b])/ ((a + b*ArcCosh[c*x])/b)^n - (2^(-11 - 3*n)*b*E^((8*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (8*(a + b*ArcCosh[c*x]))/b])/ ((a + b*ArcCosh[c*x])/b)^n) / (b*c^3*sqrt[1 + c*x])`

### 3.429.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] & IGtQ[p, 0]`



rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.429.4 Maple [F]

$$\int x^2(-c^2 d x^2 + d)^{\frac{5}{2}}(a + b \operatorname{arccosh}(cx))^n dx$$

input `int(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x)`

output `int(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x)`

### 3.429.5 Fricas [F]

$$\int x^2(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)^n x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="fricas")`

output `integral((c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2)*sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n, x)`

### 3.429.6 Sympy [F(-1)]

Timed out.

$$\int x^2(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n dx = \text{Timed out}$$

input `integrate(x**2*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

---

3.429.  $\int x^2(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n dx$

**3.429.7 Maxima [F]**

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^n x^2 dx$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^n*x^2, x)`

**3.429.8 Giac [F(-2)]**

Exception generated.

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate(x^2*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Valuesym2poly/r2sym(const gen &`

**3.429.9 Mupad [F(-1)]**

Timed out.

$$\int x^2(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \int x^2 (a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{5/2} dx$$

input `int(x^2*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2),x)`

output `int(x^2*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2), x)`

### 3.430 $\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$

3.430.1 Optimal result	3266
3.430.2 Mathematica [A] (verified)	3267
3.430.3 Rubi [A] (verified)	3268
3.430.4 Maple [F]	3270
3.430.5 Fricas [F]	3270
3.430.6 Sympy [F(-1)]	3270
3.430.7 Maxima [F]	3271
3.430.8 Giac [F(-2)]	3271
3.430.9 Mupad [F(-1)]	3271

#### 3.430.1 Optimal result

Integrand size = 27, antiderivative size = 793

$$\begin{aligned}
 & \int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \\
 & \frac{7^{-1-n} d^2 e^{-\frac{7a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{7(a + \operatorname{barccosh}(cx))}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{5^{-n} d^2 e^{-\frac{5a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & + \frac{3^{1-n} d^2 e^{-\frac{3a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{5d^2 e^{-\frac{a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & + \frac{5d^2 e^{a/b} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{3^{1-n} d^2 e^{\frac{3a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & + \frac{5^{-n} d^2 e^{\frac{5a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}} \\
 & - \frac{7^{-1-n} d^2 e^{\frac{7a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{7(a + \operatorname{barccosh}(cx))}{b}\right)}{128c^2 \sqrt{-1 + cx} \sqrt{1 + cx}}
 \end{aligned}$$

---

3.430.  $\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$

output

```

1/128*7^(-1-n)*d^2*(a+b*arccosh(c*x))^n*GAMMA(1+n,-7*(a+b*arccosh(c*x))/b)
*(-c^2*d*x^2+d)^(1/2)/c^2/exp(7*a/b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(
1/2)/(c*x+1)^(1/2)-1/128*d^2*(a+b*arccosh(c*x))^n*GAMMA(1+n,-5*(a+b*arccos
h(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/(5^n)/c^2/exp(5*a/b)/(((a+b*arccosh(c*x))
/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/128*3^(1-n)*d^2*(a+b*arccosh(c*x))^n*
GAMMA(1+n,-3*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^2/exp(3*a/b)/(((
-a-b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-5/128*d^2*(a+b*arccos
h(c*x))^n*GAMMA(1+n,(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^2/exp(a/
b)/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+5/128*d^2*exp(a
/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1
/2)/c^2/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/128*3^(1-
n)*d^2*exp(3*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,3*(a+b*arccosh(c*x))/b)*(-
c^2*d*x^2+d)^(1/2)/c^2/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(
1/2)+1/128*d^2*exp(5*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,5*(a+b*arccosh(c*
x))/b)*(-c^2*d*x^2+d)^(1/2)/(5^n)/c^2/(((a+b*arccosh(c*x))/b)^n)/(c*x-1)^(
1/2)/(c*x+1)^(1/2)-1/128*7^(-1-n)*d^2*exp(7*a/b)*(a+b*arccosh(c*x))^n*GAMM
A(1+n,7*(a+b*arccosh(c*x))/b)*(-c^2*d*x^2+d)^(1/2)/c^2/(((a+b*arccosh(c*x)
)/b)^n)/(c*x-1)^(1/2)/(c*x+1)^(1/2)

```

### 3.430.2 Mathematica [A] (verified)

Time = 3.07 (sec) , antiderivative size = 633, normalized size of antiderivative = 0.80

$$\int x(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n dx$$

$$= \frac{5^{-n} 2^{1-n} d^3 e^{-\frac{7a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx)(a + b \operatorname{arccosh}(cx))^n \left( -\frac{(a+b \operatorname{arccosh}(cx))^2}{b^2} \right)^{-3n} \left( -10 \right)}{\dots}$$

input `Integrate[x*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n,x]`

output

```
(21^(-1 - n)*d^3*sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])
^n*(-(105^(1 + n)*E^((8*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n*(-((a + b*ArcC
osh[c*x])^2/b^2))^(2*n)*Gamma[1 + n, a/b + ArcCosh[c*x]]) + (a/b + ArcCosh
[c*x])^n*(-(3^(1 + n)*5^n*(-((a + b*ArcCosh[c*x])^2/b^2))^(2*n)*Gamma[1 +
n, (-7*(a + b*ArcCosh[c*x]))/b]) + E^((2*a)/b)*(21^(1 + n)*(-(a + b*ArcCo
sh[c*x])^2/b^2))^(2*n)*Gamma[1 + n, (-5*(a + b*ArcCosh[c*x]))/b] - 9*5^n*7
^(1 + n)*E^((2*a)/b)*(-(a + b*ArcCosh[c*x])^2/b^2))^(2*n)*Gamma[1 + n, (-
3*(a + b*ArcCosh[c*x]))/b] + 105^(1 + n)*E^((4*a)/b)*(-(a + b*ArcCosh[c*x
])^2/b^2))^(2*n)*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b] - 5^n*7^(2 + n)*E
^((8*a)/b)*(a/b + ArcCosh[c*x])^n*(-((a + b*ArcCosh[c*x])/b))^(3*n)*Gamma[
1 + n, (3*(a + b*ArcCosh[c*x]))/b] + 16*5^n*7^(1 + n)*E^((8*a)/b)*(-(a +
b*ArcCosh[c*x])/b))^(2*n)*(-(a + b*ArcCosh[c*x])^2/b^2))^n*Gamma[1 + n, (
3*(a + b*ArcCosh[c*x]))/b] - 21^(1 + n)*E^((10*a)/b)*(a/b + ArcCosh[c*x])^
n*(-((a + b*ArcCosh[c*x])/b))^(3*n)*Gamma[1 + n, (5*(a + b*ArcCosh[c*x]))/
b] + 3^(1 + n)*5^n*E^((12*a)/b)*(a/b + ArcCosh[c*x])^n*(-((a + b*ArcCosh[c
*x])/b))^(3*n)*Gamma[1 + n, (7*(a + b*ArcCosh[c*x]))/b]])))/(128*5^n*c^2*E
^((7*a)/b)*sqrt[d - c^2*d*x^2]*(-(a + b*ArcCosh[c*x])^2/b^2))^(3*n))
```

### 3.430.3 Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 532, normalized size of antiderivative = 0.67, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6367, 5971, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$$

$$\downarrow \text{6367}$$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int (a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) \sinh^6\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^2 \sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow \text{5971}$$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int \left(\frac{1}{64} \cosh\left(\frac{7a}{b} - \frac{7(a + \operatorname{barccosh}(cx))}{b}\right)\right) (a + \operatorname{barccosh}(cx))^n - \frac{5}{64} \cosh\left(\frac{5a}{b} - \frac{5(a + \operatorname{barccosh}(cx))}{b}\right) (a + \operatorname{barccosh}(cx))^n}{1}}$$

$$\downarrow \text{2009}$$

---

3.430.  $\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$

$$d^2 \sqrt{d - c^2 dx^2} \left( \frac{1}{128} b 7^{-n-1} e^{-\frac{7a}{b}} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{7(a + \operatorname{barccosh}(cx))}{b}\right) - \frac{1}{128} b 5^{-n} \right)$$

input `Int[x*(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n,x]`

output `(d^2*sqrt[d - c^2*d*x^2]*((7^(-1 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-7*(a + b*ArcCosh[c*x]))/b])/(128*E^((7*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n - (b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-5*(a + b*ArcCosh[c*x]))/b])/(128*5^n*E^((5*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n + (3^(1 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-3*(a + b*ArcCosh[c*x]))/b])/(128*E^((3*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n - (5*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b])/(128*E^(a/b)*(-(a + b*ArcCosh[c*x])/b))^n + (5*b*E^(a/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (a + b*ArcCosh[c*x])/b])/(128*((a + b*ArcCosh[c*x])/b)^n - (3^(1 - n)*b*E^((3*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (3*(a + b*ArcCosh[c*x])/b])/(128*((a + b*ArcCosh[c*x])/b)^n) + (b*E^((5*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (5*(a + b*ArcCosh[c*x])/b])/(128*5^n*((a + b*ArcCosh[c*x])/b)^n - (7^(-1 - n)*b*E^((7*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (7*(a + b*ArcCosh[c*x])/b])/(128*((a + b*ArcCosh[c*x])/b)^n)))/(b*c^2*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.430.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 5971 `Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

$$3.430. \quad \int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$$

**3.430.4 Maple [F]**

$$\int x(-c^2dx^2 + d)^{\frac{5}{2}}(a + b \operatorname{arccosh}(cx))^n dx$$

input `int(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x)`

output `int(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x)`

**3.430.5 Fracas [F]**

$$\int x(d - c^2dx^2)^{5/2}(a + b \operatorname{arccosh}(cx))^n dx = \int (-c^2dx^2 + d)^{\frac{5}{2}}(b \operatorname{arccosh}(cx) + a)^n x dx$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="fracas")`

output `integral((c^4*d^2*x^5 - 2*c^2*d^2*x^3 + d^2*x)*sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n, x)`

**3.430.6 Sympy [F(-1)]**

Timed out.

$$\int x(d - c^2dx^2)^{5/2}(a + b \operatorname{arccosh}(cx))^n dx = \text{Timed out}$$

input `integrate(x*(-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.430.7 Maxima [F]**

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^n x dx$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^n*x, x)`

**3.430.8 Giac [F(-2)]**

Exception generated.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate(x*(-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.430.9 Mupad [F(-1)]**

Timed out.

$$\int x(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \int x(a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{5/2} dx$$

input `int(x*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2),x)`

output `int(x*(a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2), x)`



### 3.431 $\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$

3.431.1 Optimal result . . . . .	3272
3.431.2 Mathematica [A] (verified) . . . . .	3273
3.431.3 Rubi [A] (verified) . . . . .	3274
3.431.4 Maple [F] . . . . .	3276
3.431.5 Fricas [F] . . . . .	3276
3.431.6 Sympy [F(-1)] . . . . .	3276
3.431.7 Maxima [F] . . . . .	3277
3.431.8 Giac [F(-2)] . . . . .	3277
3.431.9 Mupad [F(-1)] . . . . .	3277

#### 3.431.1 Optimal result

Integrand size = 26, antiderivative size = 674

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = -\frac{5d^2 \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^{1+n}}{16bc(1+n)\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2^{-7-n} 3^{-1-n} d^2 e^{-\frac{6a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{6(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{3 \cdot 2^{-7-2n} d^2 e^{-\frac{4a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{15 \cdot 2^{-7-n} d^2 e^{-\frac{2a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, -\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{15 \cdot 2^{-7-n} d^2 e^{\frac{2a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{3 \cdot 2^{-7-2n} d^2 e^{\frac{4a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{2^{-7-n} 3^{-1-n} d^2 e^{\frac{6a}{b}} \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n \left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(1+n, \frac{6(a + \operatorname{barccosh}(cx))}{b}\right)}{c\sqrt{-1+cx}\sqrt{1+cx}}$$

output 
$$\begin{aligned} & -5/16*d^2*(a+b*\operatorname{arccosh}(c*x))^{(1+n)}*(-c^2*d*x^2+d)^{(1/2)}/b/c/(1+n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+2^{(-7-n)}*3^{(-1-n)}*d^2*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,- \\ & 6*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c/\exp(6*a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3*2^{(-7-2*n)}*d^2*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,-4*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c/\exp(4*a/b)/ \\ & (((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+15*2^{(-7-n)}*d^2*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,-2*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c/\exp(2*a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-15*2^{(-7-n)}*d^2*\exp(2*a/b)*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,2*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3*2^{(-7-2*n)}*d^2*\exp(4*a/b)*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,4*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-2^{(-7-n)}*3^{(-1-n)}*d^2*\exp(6*a/b)*(a+b*\operatorname{arccosh}(c*x))^n*\operatorname{GAMMA}(1+n,6*(a+b*\operatorname{arccosh}(c*x))/b)*(-c^2*d*x^2+d)^{(1/2)}/c/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \end{aligned}$$

### 3.431.2 Mathematica [A] (verified)

Time = 3.95 (sec) , antiderivative size = 538, normalized size of antiderivative = 0.80

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \frac{2^{-7-2n} 3^{-1-n} d^3 e^{-\frac{6a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx)(a + \operatorname{barccosh}(cx))^n \left(-\frac{(a+b\operatorname{arccosh}(cx))^2}{b^2}\right)^{-2n} \left(-2\right)}{\dots}$$

input `Integrate[(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n,x]`

output  $(2^{(-7 - 2*n)}*3^{(-1 - n)}*d^3*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*\text{ArcCosh}[c*x])^n*(-(2^n*b*(1 + n)*(a/b + \text{ArcCosh}[c*x])^{(2*n)}*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)}*\text{Gamma}[1 + n, (-6*(a + b*\text{ArcCosh}[c*x])/b)] + 3^{(2 + n)}*b*\text{E}^{((2*a)/b)*(1 + n)*(a/b + \text{ArcCosh}[c*x])^{(2*n)}*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)}*\text{Gamma}[1 + n, (-4*(a + b*\text{ArcCosh}[c*x])/b]} - 5*2^n*3^{(2 + n)}*b*\text{E}^{((4*a)/b)*(1 + n)*(a/b + \text{ArcCosh}[c*x])^{(2*n)}*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)}*\text{Gamma}[1 + n, (-2*(a + b*\text{ArcCosh}[c*x])/b]} + 5*2^n*3^{(2 + n)}*b*\text{E}^{((8*a)/b)*(1 + n)*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)}*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)}*\text{Gamma}[1 + n, (2*(a + b*\text{ArcCosh}[c*x])/b]} - 3^{(2 + n)}*b*\text{E}^{((10*a)/b)*(1 + n)*(a/b + \text{ArcCosh}[c*x])^{(2*n)}*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)}*\text{Gamma}[1 + n, (4*(a + b*\text{ArcCosh}[c*x])/b]} + 2^n*\text{E}^{((6*a)/b)*(5*2^{(3 + n)}*3^{(1 + n)}*(a + b*\text{ArcCosh}[c*x])*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)} + b*\text{E}^{((6*a)/b)*(1 + n)*(a/b + \text{ArcCosh}[c*x])^{(2*n)}*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)}*\text{Gamma}[1 + n, (6*(a + b*\text{ArcCosh}[c*x])/b)])))/(b*c*\text{E}^{((6*a)/b)*(1 + n)}*\text{Sqrt}[d - c^2*d*x^2]*(-((a + b*\text{ArcCosh}[c*x])/b))^{(2*n)})$

### 3.431.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 447, normalized size of antiderivative = 0.66, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$ , Rules used = {6321, 3042, 25, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^n dx$$

$$\downarrow \text{6321}$$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int (a + \text{barccosh}(cx))^n \sinh^6 \left( \frac{a}{b} - \frac{a + \text{barccosh}(cx)}{b} \right) d(a + \text{barccosh}(cx))}{bc \sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow \text{3042}$$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int -(a + \text{barccosh}(cx))^n \sin \left( \frac{ia}{b} - \frac{i(a + \text{barccosh}(cx))}{b} \right)^6 d(a + \text{barccosh}(cx))}{bc \sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow \text{25}$$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int (a + \text{barccosh}(cx))^n \sin \left( \frac{ia}{b} - \frac{i(a + \text{barccosh}(cx))}{b} \right)^6 d(a + \text{barccosh}(cx))}{bc \sqrt{cx - 1} \sqrt{cx + 1}}$$

$$\downarrow \text{3793}$$

---

3.431.  $\int (d - c^2 dx^2)^{5/2} (a + \text{barccosh}(cx))^n dx$

$$\frac{d^2 \sqrt{d - c^2 dx^2} \int \left( -\frac{1}{32} \cosh \left( \frac{6a}{b} - \frac{6(a + \operatorname{barccosh}(cx))}{b} \right) (a + \operatorname{barccosh}(cx))^n + \frac{3}{16} \cosh \left( \frac{4a}{b} - \frac{4(a + \operatorname{barccosh}(cx))}{b} \right) (a + \operatorname{barccosh}(cx))^n \right)}{bc\sqrt{c}}$$

↓ 2009

$$\frac{d^2 \sqrt{d - c^2 dx^2} \left( -\frac{5(a + \operatorname{barccosh}(cx))^{n+1}}{16(n+1)} + b 2^{-n-7} 3^{-n-1} e^{-\frac{6a}{b}} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma(n + 1, -\frac{6(a + \operatorname{barccosh}(cx))}{b}) \right)}{bc\sqrt{c}}$$

input `Int[(d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n,x]`

output `(d^2*sqrt[d - c^2*d*x^2]*((-5*(a + b*ArcCosh[c*x])^(1 + n))/(16*(1 + n)) + (2^(-7 - n)*3^(-1 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-6*(a + b*ArcCosh[c*x]))/b])/E^((6*a)/b)*(-(a + b*ArcCosh[c*x])/b)^n - (3*2^(-7 - 2*n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-4*(a + b*ArcCosh[c*x]))/b])/E^((4*a)/b)*(-(a + b*ArcCosh[c*x])/b)^n + (15*2^(-7 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x]))/b])/E^((2*a)/b)*(-(a + b*ArcCosh[c*x])/b)^n - (15*2^(-7 - n)*b*E^((2*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x]))/b])/((a + b*ArcCosh[c*x])/b)^n + (3*2^(-7 - 2*n)*b*E^((4*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (4*(a + b*ArcCosh[c*x]))/b])/((a + b*ArcCosh[c*x])/b)^n - (2^(-7 - n)*3^(-1 - n)*b*E^((6*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (6*(a + b*ArcCosh[c*x]))/b])/((a + b*ArcCosh[c*x])/b)^n)/(b*c*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.431.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

---

3.431.  $\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx$

rule 6321 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((d_.) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] :> Simp[(1/(b*c))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)]  
Subst[Int[x^n*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x]  
/; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p, 0]`

### 3.431.4 Maple [F]

$$\int (-c^2 dx^2 + d)^{\frac{5}{2}} (a + b \operatorname{arccosh}(cx))^n dx$$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x)`

output `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x)`

### 3.431.5 Fracas [F]

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{\frac{5}{2}} (b \operatorname{arccosh}(cx) + a)^n dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="fracas")`

output `integral((c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2)*sqrt(-c^2*d*x^2 + d)*(b*arcco  
sh(c*x) + a)^n, x)`

### 3.431.6 Sympy [F(-1)]

Timed out.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.431.7 Maxima [F]**

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^n, x)`

**3.431.8 Giac [F(-2)]**

Exception generated.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.431.9 Mupad [F(-1)]**

Timed out.

$$\int (d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{5/2} dx$$

input `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2),x)`

output `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2), x)`

$$3.432 \quad \int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^n}{x} dx$$

3.432.1 Optimal result	3279
3.432.2 Mathematica [N/A]	3280
3.432.3 Rubi [N/A]	3281
3.432.4 Maple [N/A] (verified)	3282
3.432.5 Fricas [N/A]	3282
3.432.6 Sympy [F(-1)]	3283
3.432.7 Maxima [N/A]	3283
3.432.8 Giac [F(-2)]	3283
3.432.9 Mupad [N/A]	3284

---


$$3.432. \quad \int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^n}{x} dx$$

**3.432.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\begin{aligned}
& \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \\
& \frac{5^{-1-n} d^3 e^{-\frac{5a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, -\frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{d - c^2 dx^2}} \\
& - \frac{5 \cdot 3^{-1-n} d^3 e^{-\frac{3a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, -\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{d - c^2 dx^2}} \\
& + \frac{3^{-n} d^3 e^{-\frac{3a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, -\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{8\sqrt{d - c^2 dx^2}} \\
& - \frac{11 d^3 e^{-\frac{a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, -\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{16\sqrt{d - c^2 dx^2}} \\
& + \frac{11 d^3 e^{a/b} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{16\sqrt{d - c^2 dx^2}} \\
& + \frac{5 \cdot 3^{-1-n} d^3 e^{\frac{3a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, \frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{d - c^2 dx^2}} \\
& - \frac{3^{-n} d^3 e^{\frac{3a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, \frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{8\sqrt{d - c^2 dx^2}} \\
& + \frac{5^{-1-n} d^3 e^{\frac{5a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(1 + n, \frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{d - c^2 dx^2}} \\
& + d^3 \operatorname{Int}\left(\frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{d - c^2 dx^2}}, x\right)
\end{aligned}$$



output

```

-1/32*5^(-1-n)*d^3*(a+b*arccosh(c*x))^n*GAMMA(1+n,-5*(a+b*arccosh(c*x))/b)
*(c*x-1)^(1/2)*(c*x+1)^(1/2)/exp(5*a/b)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*
d*x^2+d)^(1/2)-5/32*3^(-1-n)*d^3*(a+b*arccosh(c*x))^n*GAMMA(1+n,-3*(a+b*ar
ccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/exp(3*a/b)/(((a+b*arccosh(c*x)
)/b)^n)/(-c^2*d*x^2+d)^(1/2)+1/8*d^3*(a+b*arccosh(c*x))^n*GAMMA(1+n,-3*(a+
b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(3^n)/exp(3*a/b)/(((a+b*ar
ccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)-11/16*d^3*(a+b*arccosh(c*x))^n*GAMM
A(1+n,(-a-b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/exp(a/b)/(((a+b*
arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+11/16*d^3*exp(a/b)*(a+b*arccosh(c
*x))^n*GAMMA(1+n,(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/(((a+b*
arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+5/32*3^(-1-n)*d^3*exp(3*a/b)*(a+b
*arccosh(c*x))^n*GAMMA(1+n,3*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(
1/2)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)-1/8*d^3*exp(3*a/b)*(a
+b*arccosh(c*x))^n*GAMMA(1+n,3*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)
^(1/2)/(3^n)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+1/32*5^(-1-n)
*d^3*exp(5*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,5*(a+b*arccosh(c*x))/b)*(c*
x-1)^(1/2)*(c*x+1)^(1/2)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+d
^3*Unintegrable((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2),x)

```

### 3.432.2 Mathematica [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n)/x,x]`

output `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n)/x, x]`

**3.432.3 Rubi [N/A]**

Not integrable

Time = 2.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$$

↓ 6369

$$\int \left( -\frac{3c^2 d^3 x (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} + \frac{d^3 (a + \operatorname{barccosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} - \frac{c^6 d^3 x^5 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} + \frac{3c^4 d^3 x^3 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} \right) dx$$

↓ 2009

$$\frac{5^{-n-1} d^3 e^{-\frac{5a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \Gamma\left(n+1, -\frac{5(a+\operatorname{barccosh}(cx))}{b}\right) \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}}{32\sqrt{d-c^2 dx^2}} -$$

$$\frac{5 \cdot 3^{-n-1} d^3 e^{-\frac{3a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \Gamma\left(n+1, -\frac{3(a+\operatorname{barccosh}(cx))}{b}\right) \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}}{32\sqrt{d-c^2 dx^2}} +$$

$$\frac{3^{-n} d^3 e^{-\frac{3a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \Gamma\left(n+1, -\frac{3(a+\operatorname{barccosh}(cx))}{b}\right) \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}}{8\sqrt{d-c^2 dx^2}} -$$

$$\frac{11 d^3 e^{-\frac{a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \Gamma\left(n+1, -\frac{a+\operatorname{barccosh}(cx)}{b}\right) \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n}}{16\sqrt{d-c^2 dx^2}} +$$

$$\frac{11 d^3 e^{a/b} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n+1, \frac{a+\operatorname{barccosh}(cx)}{b}\right)}{16\sqrt{d-c^2 dx^2}} +$$

$$\frac{5 \cdot 3^{-n-1} d^3 e^{\frac{3a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n+1, \frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{d-c^2 dx^2}} -$$

$$\frac{3^{-n} d^3 e^{\frac{3a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n+1, \frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{8\sqrt{d-c^2 dx^2}} +$$

$$\frac{5^{-n-1} d^3 e^{\frac{5a}{b}} \sqrt{cx-1} \sqrt{cx+1} (a + \operatorname{barccosh}(cx))^n \left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n+1, \frac{5(a+\operatorname{barccosh}(cx))}{b}\right)}{32\sqrt{d-c^2 dx^2}} +$$

$$d^3 \int \frac{(a + \operatorname{barccosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} dx$$

---

3.432.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n)/x,x]`

output `$Aborted`

### 3.432.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n_)*((f_.)*(x_.))^m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`

### 3.432.4 Maple [N/A] (verified)

Not integrable

Time = 1.66 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 d x^2 + d)^{\frac{5}{2}} (a + b \operatorname{arccosh}(cx))^n}{x} dx$$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x,x)`

output `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x,x)`

### 3.432.5 Fracas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.86

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)^n}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x,x, algorithm="fracas")`

---

3.432.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n}{x} dx$

output `integral((c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2)*sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/x, x)`

### 3.432.6 Sympy [F(-1)]

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**n/x,x)`

output Timed out

### 3.432.7 Maxima [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)^n}{x} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^n/x, x)`

### 3.432.8 Giac [F(-2)]

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x,x, algorithm="giac")`

output Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vecteur & l) Error: Bad Argument Value

### 3.432.9 Mupad [N/A]

Not integrable

Time = 3.20 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{5/2}}{x} dx$$

input `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2))/x,x)`

output `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2))/x, x)`

**3.433** 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx$$

3.433.1 Optimal result . . . . .	3285
3.433.2 Mathematica [N/A] . . . . .	3286
3.433.3 Rubi [N/A] . . . . .	3286
3.433.4 Maple [N/A] (verified) . . . . .	3288
3.433.5 Fracas [N/A] . . . . .	3288
3.433.6 Sympy [F(-1)] . . . . .	3288
3.433.7 Maxima [N/A] . . . . .	3289
3.433.8 Giac [F(-2)] . . . . .	3289
3.433.9 Mupad [N/A] . . . . .	3289

**3.433.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(d - c^2dx^2)^{5/2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx = -\frac{15cd^3\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^{1+n}}{8b(1 + n)\sqrt{d - c^2dx^2}}$$

$$-\frac{2^{-2(3+n)}cd^3e^{-\frac{4a}{b}}\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^n\left(-\frac{a+\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1 + n, -\frac{4(a+\operatorname{arccosh}(cx))}{b}\right)}{\sqrt{d - c^2dx^2}}$$

$$+\frac{2^{-2-n}cd^3e^{-\frac{2a}{b}}\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^n\left(-\frac{a+\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1 + n, -\frac{2(a+\operatorname{arccosh}(cx))}{b}\right)}{\sqrt{d - c^2dx^2}}$$

$$-\frac{2^{-2-n}cd^3e^{\frac{2a}{b}}\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^n\left(\frac{a+\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1 + n, \frac{2(a+\operatorname{arccosh}(cx))}{b}\right)}{\sqrt{d - c^2dx^2}}$$

$$+\frac{2^{-2(3+n)}cd^3e^{\frac{4a}{b}}\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^n\left(\frac{a+\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1 + n, \frac{4(a+\operatorname{arccosh}(cx))}{b}\right)}{\sqrt{d - c^2dx^2}}$$

$$+ d^3\operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{d - c^2dx^2}}, x\right)$$

---

3.433. 
$$\int \frac{(d-c^2dx^2)^{5/2}(a+b\operatorname{arccosh}(cx))^n}{x^2} dx$$

output 
$$-15/8*c*d^3*(a+b*\operatorname{arccosh}(c*x))^{(1+n)}*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/b/(1+n)/(-c^2*d*x^2+d)^{(1/2)}-c*d^3*(a+b*\operatorname{arccosh}(c*x))^{(1+n)}*\operatorname{GAMMA}(1+n,-4*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(2^{(6+2*n)})/\exp(4*a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}+2^{(-2-n)}*c*d^3*(a+b*\operatorname{arccosh}(c*x))^{(1+n)}*\operatorname{GAMMA}(1+n,-2*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/\exp(2*a/b)/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}-2^{(-2-n)}*c*d^3*\exp(2*a/b)*(a+b*\operatorname{arccosh}(c*x))^{(1+n)}*\operatorname{GAMMA}(1+n,2*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}+c*d^3*\exp(4*a/b)*(a+b*\operatorname{arccosh}(c*x))^{(1+n)}*\operatorname{GAMMA}(1+n,4*(a+b*\operatorname{arccosh}(c*x))/b)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/(2^{(6+2*n)})/(((a+b*\operatorname{arccosh}(c*x))/b)^n)/(-c^2*d*x^2+d)^{(1/2)}+d^3*\operatorname{Unintegrable}((a+b*\operatorname{arccosh}(c*x))^{(1+n)}/x^2/(-c^2*d*x^2+d)^{(1/2)},x)$$

### 3.433.2 Mathematica [N/A]

Not integrable

Time = 0.62 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx = \int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx$$

input `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n)/x^2,x]`

output `Integrate[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n)/x^2, x]`

### 3.433.3 Rubi [N/A]

Not integrable

Time = 1.60 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6369, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx$$

↓ 6369

---

3.433.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{arccosh}(cx))^n}{x^2} dx$

$$\begin{aligned}
& \int \left( -\frac{3c^2 d^3 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} + \frac{d^3 (a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} - \frac{c^6 d^3 x^4 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} + \frac{3c^4 d^3 x^2 (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} \right) \\
& \quad \downarrow \text{2009} \\
& d^3 \int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx - \frac{15cd^3 \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^{n+1}}{8b(n+1) \sqrt{d - c^2 dx^2}} - \\
& \frac{cd^3 2^{-2(n+3)} e^{-\frac{4a}{b}} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{d - c^2 dx^2}} + \\
& \frac{cd^3 2^{-n-2} e^{-\frac{2a}{b}} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, -\frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{d - c^2 dx^2}} - \\
& \frac{cd^3 2^{-n-2} e^{\frac{2a}{b}} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, \frac{2(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{d - c^2 dx^2}} + \\
& \frac{cd^3 2^{-2(n+3)} e^{\frac{4a}{b}} \sqrt{cx - 1} \sqrt{cx + 1} (a + \operatorname{barccosh}(cx))^n \left( \frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n + 1, \frac{4(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{d - c^2 dx^2}}
\end{aligned}$$

input `Int[((d - c^2*d*x^2)^(5/2)*(a + b*ArcCosh[c*x])^n)/x^2, x]`

output `$Aborted`

### 3.433.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6369 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n_)*((f_.)*(x_)^m_)*((d_) + (e_.)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n/Sqrt[d + e*x^2], (f*x)^m*(d + e*x^2)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && !IGtQ[(m + 1)/2, 0] && (EqQ[m, -1] || EqQ[m, -2])`



**3.433.4 Maple [N/A] (verified)**

Not integrable

Time = 1.40 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(-c^2 d x^2 + d)^{\frac{5}{2}} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx$$

input `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x^2,x)`output `int((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x^2,x)`**3.433.5 Fricas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.86

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{\frac{5}{2}} (b \operatorname{arccosh}(cx) + a)^n}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x^2,x, algorithm="fricas")`output `integral((c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2)*sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/x^2, x)`**3.433.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + b \operatorname{arccosh}(cx))^n}{x^2} dx = \text{Timed out}$$

input `integrate((-c**2*d*x**2+d)**(5/2)*(a+b*acosh(c*x))**n/x**2,x)`output `Timed out`

**3.433.7 Maxima [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \int \frac{(-c^2 dx^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)^n}{x^2} dx$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x^2,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^n/x^2, x)`

**3.433.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \text{Exception raised: TypeError}$$

input `integrate((-c^2*d*x^2+d)^(5/2)*(a+b*arccosh(c*x))^n/x^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.433.9 Mupad [N/A]**

Not integrable

Time = 3.49 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{5/2}}{x^2} dx$$

input `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2))/x^2,x)`

output `int(((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(5/2))/x^2, x)`

---

3.433.  $\int \frac{(d - c^2 dx^2)^{5/2} (a + \operatorname{barccosh}(cx))^n}{x^2} dx$

**3.434**  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$

3.434.1 Optimal result . . . . . 3290  
 3.434.2 Mathematica [A] (verified) . . . . . 3291  
 3.434.3 Rubi [A] (verified) . . . . . 3291  
 3.434.4 Maple [F] . . . . . 3293  
 3.434.5 Fricas [F] . . . . . 3293  
 3.434.6 Sympy [F] . . . . . 3293  
 3.434.7 Maxima [F] . . . . . 3294  
 3.434.8 Giac [F(-2)] . . . . . 3294  
 3.434.9 Mupad [F(-1)] . . . . . 3294

**3.434.1 Optimal result**

Integrand size = 28, antiderivative size = 323

$$\int \frac{x^3(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$$

$$= \frac{3^{-1-n}e^{-\frac{3a}{b}}\sqrt{-1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{8c^4\sqrt{1-cx}}$$

$$+ \frac{3e^{-\frac{a}{b}}\sqrt{-1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8c^4\sqrt{1-cx}}$$

$$- \frac{3e^{a/b}\sqrt{-1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8c^4\sqrt{1-cx}}$$

$$- \frac{3^{-1-n}e^{\frac{3a}{b}}\sqrt{-1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{8c^4\sqrt{1-cx}}$$

output

```
1/8*3^(-1-n)*(a+b*arccosh(c*x))^n*GAMMA(1+n,-3*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^4/exp(3*a/b)/(((a+b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)+3/8*(a+b*arccosh(c*x))^n*GAMMA(1+n,(-a-b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^4/exp(a/b)/(((a+b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)-3/8*exp(a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^4/(((a+b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)-1/8*3^(-1-n)*exp(3*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,3*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^4/(((a+b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)
```

3.434.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$

### 3.434.2 Mathematica [A] (verified)

Time = 0.97 (sec) , antiderivative size = 292, normalized size of antiderivative = 0.90

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

$$= \frac{3^{-1-n} e^{-\frac{3a}{b}} \sqrt{1 - c^2x^2} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b^2}\right)^{-2n} \left(3^{2+n} e^{\frac{4a}{b}} \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^n \left(-\frac{a + \operatorname{barccosh}(cx)}{b}\right)^n \right)}{\dots}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2], x]`

output  $(3^{-1-n} \sqrt{1 - c^2x^2} (a + b \operatorname{ArcCosh}[cx])^n (3^{2+n} E^{(4a/b)} (-(a + b \operatorname{ArcCosh}[cx])/b)^n (-(a + b \operatorname{ArcCosh}[cx])^2/b^2)^n \Gamma[1 + n, a/b + \operatorname{ArcCosh}[cx]] - (a/b + \operatorname{ArcCosh}[cx])^n (-(a + b \operatorname{ArcCosh}[cx])^2/b^2)^n \Gamma[1 + n, (-3(a + b \operatorname{ArcCosh}[cx]))/b] + 3^{2+n} E^{(2a/b)} (-(a + b \operatorname{ArcCosh}[cx])^2/b^2)^n \Gamma[1 + n, -(a + b \operatorname{ArcCosh}[cx])/b]) - E^{(6a/b)} (-(a + b \operatorname{ArcCosh}[cx])/b)^{2n} \Gamma[1 + n, (3(a + b \operatorname{ArcCosh}[cx])/b)]) / (8c^4 E^{(3a/b)} \sqrt{(-1 + cx)/(1 + cx)} (1 + cx) (-(a + b \operatorname{ArcCosh}[cx])^2/b^2)^{2n})$

### 3.434.3 Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.82, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{cx - 1} \int (a + \operatorname{barccosh}(cx))^n \cosh^3\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^4 \sqrt{1 - cx}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{cx-1} \int (a + \operatorname{barccosh}(cx))^n \sin\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)^3 d(a + \operatorname{barccosh}(cx))}{bc^4 \sqrt{1-cx}}$$

↓ 3793

$$\frac{\sqrt{cx-1} \int \left(\frac{1}{4} \cosh\left(\frac{3a}{b} - \frac{3(a+\operatorname{barccosh}(cx))}{b}\right) (a + \operatorname{barccosh}(cx))^n + \frac{3}{4} \cosh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) (a + \operatorname{barccosh}(cx))\right)}{bc^4 \sqrt{1-cx}}$$

↓ 2009

$$\frac{\sqrt{cx-1} \left(\frac{1}{8} b 3^{-n-1} e^{-\frac{3a}{b}} (a + \operatorname{barccosh}(cx))^n \left(-\frac{a+\operatorname{barccosh}(cx)}{b}\right)^{-n} \Gamma\left(n+1, -\frac{3(a+\operatorname{barccosh}(cx))}{b}\right) + \frac{3}{8} b e^{-\frac{a}{b}} (a + \operatorname{barccosh}(cx))\right)}{bc^4 \sqrt{1-cx}}$$

input `Int[(x^3*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2],x]`

output `(Sqrt[-1 + c*x]*((3^(-1 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-3*(a + b*ArcCosh[c*x])/b)]/(8*E^((3*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n) + (3*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b])/(8*E^(a/b)*(-(a + b*ArcCosh[c*x])/b))^n - (3*b*E^(a/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (a + b*ArcCosh[c*x])/b])/(8*((a + b*ArcCosh[c*x])/b)^n) - (3^(-1 - n)*b*E^((3*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (3*(a + b*ArcCosh[c*x])/b)]/(8*((a + b*ArcCosh[c*x])/b)^n)))/(b*c^4*Sqrt[1 - c*x])`

### 3.434.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.434.4 Maple [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2x^2 + 1}} dx$$

input `int(x^3*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

output `int(x^3*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

### 3.434.5 Fracas [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^n x^3}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^n*x^3/(c^2*x^2 - 1), x)`

### 3.434.6 Sympy [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^n}{\sqrt{-(cx - 1)(cx + 1)}} dx$$

input `integrate(x**3*(a+b*acosh(c*x))**n/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x**3*(a + b*acosh(c*x))**n/sqrt(-(c*x - 1)*(c*x + 1)), x)`

---

3.434.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$

**3.434.7 Maxima [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^3}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x^3/sqrt(-c^2*x^2 + 1), x)`

**3.434.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.434.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

input `int((x^3*(a + b*acosh(c*x))^n)/(1 - c^2*x^2)^(1/2),x)`

output `int((x^3*(a + b*acosh(c*x))^n)/(1 - c^2*x^2)^(1/2), x)`

**3.435**  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$

3.435.1 Optimal result . . . . . 3295  
 3.435.2 Mathematica [A] (verified) . . . . . 3296  
 3.435.3 Rubi [A] (verified) . . . . . 3296  
 3.435.4 Maple [F] . . . . . 3298  
 3.435.5 Fracas [F] . . . . . 3298  
 3.435.6 Sympy [F] . . . . . 3298  
 3.435.7 Maxima [F] . . . . . 3299  
 3.435.8 Giac [F] . . . . . 3299  
 3.435.9 Mupad [F(-1)] . . . . . 3299

**3.435.1 Optimal result**

Integrand size = 28, antiderivative size = 211

$$\int \frac{x^2(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx = \frac{\sqrt{-1+cx}(a+b\operatorname{arccosh}(cx))^{1+n}}{2bc^3(1+n)\sqrt{1-cx}} + \frac{2^{-3-n}e^{-\frac{2a}{b}}\sqrt{-1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{c^3\sqrt{1-cx}} - \frac{2^{-3-n}e^{\frac{2a}{b}}\sqrt{-1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{c^3\sqrt{1-cx}}$$

```
output 1/2*(a+b*arccosh(c*x))^(1+n)*(c*x-1)^(1/2)/b/c^3/(1+n)/(-c*x+1)^(1/2)+2^(-3-n)*(a+b*arccosh(c*x))^n*GAMMA(1+n,-2*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^3/exp(2*a/b)/(((a+b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)-2^(-3-n)*exp(2*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,2*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^3/(((a+b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)
```



**3.435.2 Mathematica [A] (verified)**

Time = 0.61 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

$$= \frac{2^{-3-n} e^{-\frac{2a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx)(a + \operatorname{barccosh}(cx))^n \left(-\frac{(a+\operatorname{barccosh}(cx))^2}{b^2}\right)^{-n} \left(2^{2+n} e^{\frac{2a}{b}} (a + \operatorname{barccosh}(cx)) \left(-\frac{(a+\operatorname{barccosh}(cx))^2}{b^2}\right)^{-n} \right)}{\dots}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2],x]`output `(2^(-3 - n)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])^n*(2^(2 + n)*E^((2*a)/b)*(a + b*ArcCosh[c*x])*(-(a + b*ArcCosh[c*x])^2/b^2))^n + b*(1 + n)*(a/b + ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x])/b] - b*E^((4*a)/b)*(1 + n)*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x])/b)])/b))/b*c^3*E^((2*a)/b)*(1 + n)*Sqrt[1 - c^2*x^2]*(-(a + b*ArcCosh[c*x])^2/b^2))^n`**3.435.3 Rubi [A] (verified)**Time = 0.53 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.81, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{cx - 1} \int (a + \operatorname{barccosh}(cx))^n \cosh^2\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^3\sqrt{1 - cx}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{cx - 1} \int (a + \operatorname{barccosh}(cx))^n \sin\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)^2 d(a + \operatorname{barccosh}(cx))}{bc^3\sqrt{1 - cx}}$$

$$\downarrow \text{3793}$$

---

3.435.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$

$$\frac{\sqrt{cx-1} \int \left( \frac{1}{2} \cosh \left( \frac{2a}{b} - \frac{2(a+\operatorname{barccosh}(cx))}{b} \right) (a + \operatorname{barccosh}(cx))^n + \frac{1}{2} (a + \operatorname{barccosh}(cx))^n \right) d(a + \operatorname{barccosh}(cx))}{bc^3 \sqrt{1-cx}}$$

↓ 2009

$$\frac{\sqrt{cx-1} \left( \frac{(a+\operatorname{barccosh}(cx))^{n+1}}{2(n+1)} + b2^{-n-3} e^{-\frac{2a}{b}} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a+\operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma \left( n+1, -\frac{2(a+\operatorname{barccosh}(cx))}{b} \right) \right)}{bc^3 \sqrt{1-cx}}$$

input `Int[(x^2*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2],x]`

output `(Sqrt[-1 + c*x]*((a + b*ArcCosh[c*x])^(1 + n)/(2*(1 + n)) + (2^(-3 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x])/b)]/(E^((2*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n) - (2^(-3 - n)*b*E^((2*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x])/b)]/((a + b*ArcCosh[c*x])/b)^n))/(b*c^3*Sqrt[1 - c*x])`

### 3.435.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

**3.435.4 Maple [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2x^2 + 1}} dx$$

input `int(x^2*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

output `int(x^2*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

**3.435.5 Fracas [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^n x^2}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^n*x^2/(c^2*x^2 - 1), x)`

**3.435.6 Sympy [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^n}{\sqrt{-(cx - 1)(cx + 1)}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))**n/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x**2*(a + b*acosh(c*x))**n/sqrt(-(c*x - 1)*(c*x + 1)), x)`

**3.435.7 Maxima [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^2}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x^2/sqrt(-c^2*x^2 + 1), x)`

**3.435.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^2}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n*x^2/sqrt(-c^2*x^2 + 1), x)`

**3.435.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

input `int((x^2*(a + b*acosh(c*x))^n)/(1 - c^2*x^2)^(1/2),x)`

output `int((x^2*(a + b*acosh(c*x))^n)/(1 - c^2*x^2)^(1/2), x)`

**3.436**  $\int \frac{x(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$

3.436.1 Optimal result . . . . . 3300  
 3.436.2 Mathematica [A] (verified) . . . . . 3300  
 3.436.3 Rubi [A] (verified) . . . . . 3301  
 3.436.4 Maple [F] . . . . . 3303  
 3.436.5 Fricas [F] . . . . . 3303  
 3.436.6 Sympy [F] . . . . . 3303  
 3.436.7 Maxima [F] . . . . . 3304  
 3.436.8 Giac [F] . . . . . 3304  
 3.436.9 Mupad [F(-1)] . . . . . 3304

**3.436.1 Optimal result**

Integrand size = 26, antiderivative size = 154

$$\int \frac{x(a + b\operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

$$= \frac{e^{-\frac{a}{b}}\sqrt{-1 + cx}(a + b\operatorname{arccosh}(cx))^n \left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2c^2\sqrt{1 - cx}} - \frac{e^{a/b}\sqrt{-1 + cx}(a + b\operatorname{arccosh}(cx))^n \left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2c^2\sqrt{1 - cx}}$$

```
output 1/2*(a+b*arccosh(c*x))^n*GAMMA(1+n, (-a-b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^2/exp(a/b)/(((a-b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)-1/2*exp(a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n, (a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)/c^2/(((a+b*arccosh(c*x))/b)^n)/(-c*x+1)^(1/2)
```

**3.436.2 Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.00

$$\int \frac{x(a + b\operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \frac{e^{-\frac{a}{b}}\sqrt{-((-1 + cx)(1 + cx))}(a + b\operatorname{arccosh}(cx))^n \left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n} \left(-e^{\frac{2a}{b}}\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\right)^n \Gamma\left(1 + n, -\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2c^2\sqrt{\frac{-1+cx}{1+cx}}(1 + cx)}$$

3.436.  $\int \frac{x(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$

input `Integrate[(x*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2], x]`

output 
$$-1/2*(\text{Sqrt}[ -((-1 + c*x)*(1 + c*x))] * (a + b*\text{ArcCosh}[c*x])^n * (-E^{((2*a)/b)} * (-((a + b*\text{ArcCosh}[c*x])/b))^n * \text{Gamma}[1 + n, a/b + \text{ArcCosh}[c*x]]) + (a/b + \text{ArcCosh}[c*x])^n * \text{Gamma}[1 + n, -((a + b*\text{ArcCosh}[c*x])/b)])) / (c^2 * E^{(a/b)} * \text{Sqrt}[(-1 + c*x)/(1 + c*x)] * (1 + c*x) * (-((a + b*\text{ArcCosh}[c*x])^2/b^2))^n)$$

### 3.436.3 Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.90, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$ , Rules used = {6367, 3042, 3788, 26, 2612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x(a + \text{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx \\ & \quad \downarrow \text{6367} \\ & \frac{\sqrt{cx - 1} \int (a + \text{barccosh}(cx))^n \cosh\left(\frac{a}{b} - \frac{a + \text{barccosh}(cx)}{b}\right) d(a + \text{barccosh}(cx))}{bc^2\sqrt{1 - cx}} \\ & \quad \downarrow \text{3042} \\ & \frac{\sqrt{cx - 1} \int (a + \text{barccosh}(cx))^n \sin\left(\frac{ia}{b} - \frac{i(a + \text{barccosh}(cx))}{b} + \frac{\pi}{2}\right) d(a + \text{barccosh}(cx))}{bc^2\sqrt{1 - cx}} \\ & \quad \downarrow \text{3788} \\ & \frac{\sqrt{cx - 1} \left(\frac{1}{2}i \int -ie^{-\text{arccosh}(cx)}(a + \text{barccosh}(cx))^n d(a + \text{barccosh}(cx)) - \frac{1}{2}i \int ie^{\text{arccosh}(cx)}(a + \text{barccosh}(cx))^n d(a + \text{barccosh}(cx))\right)}{bc^2\sqrt{1 - cx}} \\ & \quad \downarrow \text{26} \\ & \frac{\sqrt{cx - 1} \left(\frac{1}{2} \int e^{-\text{arccosh}(cx)}(a + \text{barccosh}(cx))^n d(a + \text{barccosh}(cx)) + \frac{1}{2} \int e^{\text{arccosh}(cx)}(a + \text{barccosh}(cx))^n d(a + \text{barccosh}(cx))\right)}{bc^2\sqrt{1 - cx}} \\ & \quad \downarrow \text{2612} \end{aligned}$$

$$\frac{\sqrt{cx-1} \left( \frac{1}{2} b e^{-\frac{a}{b}} (a + \operatorname{barccosh}(cx))^n \left( -\frac{a + \operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma \left( n + 1, -\frac{a + \operatorname{barccosh}(cx)}{b} \right) - \frac{1}{2} b e^{a/b} (a + \operatorname{barccosh}(cx)) \right)}{bc^2 \sqrt{1 - cx}}$$

input `Int[(x*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2],x]`

output `(Sqrt[-1 + c*x]*((b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, -((a + b*ArcCosh[c*x])/b)])/(2*E^(a/b)*(-(a + b*ArcCosh[c*x])/b))^n - (b*E^(a/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (a + b*ArcCosh[c*x])/b])/(2*((a + b*ArcCosh[c*x])/b)^n))/(b*c^2*Sqrt[1 - c*x])`

### 3.436.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2612 `Int[(F_)^((g_)*(e_) + (f_)*(x_))*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*(-f)*g*(Log[F]/d))^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m])*Gamma[m + 1, ((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] && !IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3788 `Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + Pi*(k_) + (f_)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/(E^(I*k*Pi)*E^(I*(e + f*x))), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*k*Pi)*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x] && IntegerQ[2*k]`

rule 6367 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

---

3.436.  $\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$

**3.436.4 Maple [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2x^2 + 1}} dx$$

input `int(x*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

output `int(x*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

**3.436.5 Fricas [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^n x}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^n*x/(c^2*x^2 - 1), x)`

**3.436.6 Sympy [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^n}{\sqrt{-(cx - 1)(cx + 1)}} dx$$

input `integrate(x*(a+b*acosh(c*x))**n/(-c**2*x**2+1)**(1/2),x)`

output `Integral(x*(a + b*acosh(c*x))**n/sqrt(-(c*x - 1)*(c*x + 1)), x)`



**3.436.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x/sqrt(-c^2*x^2 + 1), x)`

**3.436.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x}{\sqrt{-c^2x^2 + 1}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n*x/sqrt(-c^2*x^2 + 1), x)`

**3.436.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

input `int((x*(a + b*acosh(c*x))^n)/(1 - c^2*x^2)^(1/2),x)`

output `int((x*(a + b*acosh(c*x))^n)/(1 - c^2*x^2)^(1/2), x)`

**3.437**  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{\sqrt{1-c^2x^2}} dx$

3.437.1 Optimal result . . . . . 3305  
 3.437.2 Mathematica [A] (verified) . . . . . 3305  
 3.437.3 Rubi [A] (verified) . . . . . 3306  
 3.437.4 Maple [A] (verified) . . . . . 3306  
 3.437.5 Fricas [B] (verification not implemented) . . . . . 3307  
 3.437.6 Sympy [F] . . . . . 3307  
 3.437.7 Maxima [F] . . . . . 3307  
 3.437.8 Giac [F] . . . . . 3308  
 3.437.9 Mupad [F(-1)] . . . . . 3308

**3.437.1 Optimal result**

Integrand size = 25, antiderivative size = 43

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \frac{\sqrt{-1 + cx}(a + \operatorname{arccosh}(cx))^{1+n}}{bc(1 + n)\sqrt{1 - cx}}$$

output `(a+b*arccosh(c*x))^(1+n)*(c*x-1)^(1/2)/b/c/(1+n)/(-c*x+1)^(1/2)`

**3.437.2 Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.30

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^{1+n}}{bc(1 + n)\sqrt{1 - c^2x^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^n/Sqrt[1 - c^2*x^2], x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^(1 + n))/(b*c*(1 + n)*Sqrt[1 - c^2*x^2])`

### 3.437.3 Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2x^2}} dx$$

↓ 6307

$$\frac{\sqrt{cx - 1}(a + \operatorname{arccosh}(cx))^{n+1}}{bc(n + 1)\sqrt{1 - cx}}$$

input `Int[(a + b*ArcCosh[c*x])^n/Sqrt[1 - c^2*x^2], x]`

output `(Sqrt[-1 + c*x]*(a + b*ArcCosh[c*x])^(1 + n))/(b*c*(1 + n)*Sqrt[1 - c*x])`

#### 3.437.3.1 Defintions of rubi rules used

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_ Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

### 3.437.4 Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.23

method	result	size
default	$\frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^{1+n}}{bc(1+n)\sqrt{-(cx-1)(cx+1)}}$	53

input `int((a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2), x, method=_RETURNVERBOSE)`

output `1/b/c/(1+n)*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(a+b*arccosh(c*x))^(1+n)/(-(c*x-1)*(c*x+1))^(1/2)`

**3.437.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 213 vs. 2(39) = 78.

Time = 0.28 (sec) , antiderivative size = 213, normalized size of antiderivative = 4.95

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \frac{(\sqrt{c^2 x^2 - 1} \sqrt{-c^2 x^2 + 1} b \log(cx + \sqrt{c^2 x^2 - 1}) + \sqrt{c^2 x^2 - 1} \sqrt{-c^2 x^2 + 1} a) \cosh(n \log(b \log(cx + \sqrt{c^2 x^2 - 1})))}{(b^2 c^2 x^2 - 1) \sqrt{-c^2 x^2 + 1}}$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `((sqrt(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)*b*log(c*x + sqrt(c^2*x^2 - 1)) + sqrt(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)*a)*cosh(n*log(b*log(c*x + sqrt(c^2*x^2 - 1)) + a)) + (sqrt(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)*b*log(c*x + sqrt(c^2*x^2 - 1)) + sqrt(c^2*x^2 - 1)*sqrt(-c^2*x^2 + 1)*a)*sinh(n*log(b*log(c*x + sqrt(c^2*x^2 - 1)) + a)))/(b*c*n - (b*c^3*n + b*c^3)*x^2 + b*c)`

**3.437.6 Sympy [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{\sqrt{-(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))**n/(-c**2*x**2+1)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**n/sqrt(-(c*x - 1)*(c*x + 1)), x)`

**3.437.7 Maxima [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1}} dx$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/sqrt(-c^2*x^2 + 1), x)`

---

3.437.  $\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$

**3.437.8 Giac [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1}} dx$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n/sqrt(-c^2*x^2 + 1), x)`

**3.437.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$$

input `int((a + b*acosh(c*x))^n/(1 - c^2*x^2)^(1/2),x)`

output `int((a + b*acosh(c*x))^n/(1 - c^2*x^2)^(1/2), x)`

$$3.438 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^n}{x\sqrt{1-c^2x^2}} dx$$

3.438.1 Optimal result	3309
3.438.2 Mathematica [N/A]	3309
3.438.3 Rubi [N/A]	3310
3.438.4 Maple [N/A] (verified)	3310
3.438.5 Fricas [N/A]	3311
3.438.6 Sympy [N/A]	3311
3.438.7 Maxima [N/A]	3311
3.438.8 Giac [N/A]	3312
3.438.9 Mupad [N/A]	3312

### 3.438.1 Optimal result

Integrand size = 28, antiderivative size = 28

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{1 - c^2x^2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^n/x/(-c^2*x^2+1)^(1/2), x)`

### 3.438.2 Mathematica [N/A]

Not integrable

Time = 2.09 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^n/(x*Sqrt[1 - c^2*x^2]), x]`

output `Integrate[(a + b*ArcCosh[c*x])^n/(x*Sqrt[1 - c^2*x^2]), x]`

---


$$3.438. \quad \int \frac{(a+b\operatorname{arccosh}(cx))^n}{x\sqrt{1-c^2x^2}} dx$$

**3.438.3 Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx$$

↓ 6375

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^n/(x*Sqrt[1 - c^2*x^2]),x]`

output `$Aborted`

**3.438.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.438.4 Maple [N/A] (verified)**

Not integrable

Time = 1.43 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x\sqrt{-c^2x^2 + 1}} dx$$

input `int((a+b*arccosh(c*x))^n/x/(-c^2*x^2+1)^(1/2),x)`

output `int((a+b*arccosh(c*x))^n/x/(-c^2*x^2+1)^(1/2),x)`

**3.438.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.39

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2x^2 + 1}x} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^n/(c^2*x^3 - x), x)`

**3.438.6 Sympy [N/A]**

Not integrable

Time = 6.48 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x\sqrt{-(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))**n/x/(-c**2*x**2+1)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**n/(x*sqrt(-(c*x - 1)*(c*x + 1))), x)`

**3.438.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2x^2 + 1}x} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*x^2 + 1)*x), x)`

---

3.438.  $\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{1 - c^2x^2}} dx$



**3.438.8 Giac [N/A]**

Not integrable

Time = 12.44 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x \sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1} x} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*x^2 + 1)*x), x)`**3.438.9 Mupad [N/A]**

Not integrable

Time = 3.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x \sqrt{1 - c^2 x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x \sqrt{1 - c^2 x^2}} dx$$

input `int((a + b*acosh(c*x))^n/(x*(1 - c^2*x^2)^(1/2)),x)`output `int((a + b*acosh(c*x))^n/(x*(1 - c^2*x^2)^(1/2)), x)`

**3.439**  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{x^2\sqrt{1-c^2x^2}} dx$

3.439.1 Optimal result . . . . .	3313
3.439.2 Mathematica [N/A] . . . . .	3313
3.439.3 Rubi [N/A] . . . . .	3314
3.439.4 Maple [N/A] (verified) . . . . .	3314
3.439.5 Fricas [N/A] . . . . .	3315
3.439.6 Sympy [N/A] . . . . .	3315
3.439.7 Maxima [N/A] . . . . .	3315
3.439.8 Giac [N/A] . . . . .	3316
3.439.9 Mupad [N/A] . . . . .	3316

**3.439.1 Optimal result**

Integrand size = 28, antiderivative size = 28

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{1 - c^2x^2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{1 - c^2x^2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^n/x^2/(-c^2*x^2+1)^(1/2), x)`

**3.439.2 Mathematica [N/A]**

Not integrable

Time = 1.18 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.07

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{1 - c^2x^2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{1 - c^2x^2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^n/(x^2*Sqrt[1 - c^2*x^2]), x]`

output `Integrate[(a + b*ArcCosh[c*x])^n/(x^2*Sqrt[1 - c^2*x^2]), x]`

**3.439.3 Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx$$

↓ 6375

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^n/(x^2*Sqrt[1 - c^2*x^2]),x]`

output `$Aborted`

**3.439.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.439.4 Maple [N/A] (verified)**

Not integrable

Time = 1.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x^2 \sqrt{-c^2 x^2 + 1}} dx$$

input `int((a+b*arccosh(c*x))^n/x^2/(-c^2*x^2+1)^(1/2),x)`

output `int((a+b*arccosh(c*x))^n/x^2/(-c^2*x^2+1)^(1/2),x)`

**3.439.5 Fricas [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.46

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1} x^2} dx$$

```
input integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")
```

```
output integral(-sqrt(-c^2*x^2 + 1)*(b*arccosh(c*x) + a)^n/(c^2*x^4 - x^2), x)
```

**3.439.6 Sympy [N/A]**

Not integrable

Time = 31.45 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x^2 \sqrt{-(cx - 1)(cx + 1)}} dx$$

```
input integrate((a+b*acosh(c*x))**n/x**2/(-c**2*x**2+1)**(1/2),x)
```

```
output Integral((a + b*acosh(c*x))**n/(x**2*sqrt(-(c*x - 1)*(c*x + 1))), x)
```

**3.439.7 Maxima [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1} x^2} dx$$

```
input integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")
```

```
output integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*x^2 + 1)*x^2), x)
```

---

3.439.  $\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx$

**3.439.8 Giac [N/A]**

Not integrable

Time = 12.45 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*x^2 + 1)*x^2), x)`

**3.439.9 Mupad [N/A]**

Not integrable

Time = 3.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x^2 \sqrt{1 - c^2 x^2}} dx$$

input `int((a + b*acosh(c*x))^n/(x^2*(1 - c^2*x^2)^(1/2)),x)`

output `int((a + b*acosh(c*x))^n/(x^2*(1 - c^2*x^2)^(1/2)), x)`

**3.440**  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx$

3.440.1 Optimal result . . . . . 3317  
 3.440.2 Mathematica [A] (verified) . . . . . 3318  
 3.440.3 Rubi [A] (verified) . . . . . 3318  
 3.440.4 Maple [F] . . . . . 3320  
 3.440.5 Fricas [F] . . . . . 3320  
 3.440.6 Sympy [F] . . . . . 3320  
 3.440.7 Maxima [F] . . . . . 3321  
 3.440.8 Giac [F(-2)] . . . . . 3321  
 3.440.9 Mupad [F(-1)] . . . . . 3321

**3.440.1 Optimal result**

Integrand size = 29, antiderivative size = 379

$$\int \frac{x^3(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx$$

$$= \frac{3^{-1-n}e^{-\frac{3a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{8c^4\sqrt{d-c^2dx^2}}$$

$$+ \frac{3e^{-\frac{a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8c^4\sqrt{d-c^2dx^2}}$$

$$- \frac{3e^{a/b}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8c^4\sqrt{d-c^2dx^2}}$$

$$- \frac{3^{-1-n}e^{\frac{3a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{8c^4\sqrt{d-c^2dx^2}}$$

output

```
1/8*3^(-1-n)*(a+b*arccosh(c*x))^n*GAMMA(1+n,-3*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^4/exp(3*a/b)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)+3/8*(a+b*arccosh(c*x))^n*GAMMA(1+n,(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^4/exp(a/b)/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)-3/8*exp(a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^4/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)-1/8*3^(-1-n)*exp(3*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n,3*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^4/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)
```

3.440.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx$

**3.440.2 Mathematica [A] (verified)**

Time = 0.90 (sec) , antiderivative size = 291, normalized size of antiderivative = 0.77

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx =$$

$$3^{-1-n} e^{-\frac{3a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx)(a + b \operatorname{arccosh}(cx))^n \left( -\frac{(a+b \operatorname{arccosh}(cx))^2}{b^2} \right)^{-2n} \left( 3^{2+n} e^{\frac{4a}{b}} \left( -\frac{a+b \operatorname{arccosh}(cx)}{b} \right)^n \right)$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2],x]`

output

```
-1/8*(3^(-1 - n)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])
^n*(3^(2 + n)*E^((4*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n*(-((a + b*ArcCosh[
c*x])^2/b^2))^n*Gamma[1 + n, a/b + ArcCosh[c*x]] - (a/b + ArcCosh[c*x])^n*
((-(a + b*ArcCosh[c*x])^2/b^2))^n*Gamma[1 + n, (-3*(a + b*ArcCosh[c*x]))/
b] + 3^(2 + n)*E^((2*a)/b)*(-(a + b*ArcCosh[c*x])^2/b^2))^n*Gamma[1 + n,
-(a + b*ArcCosh[c*x])/b] - E^((6*a)/b)*(-(a + b*ArcCosh[c*x])/b))^(2*n)
*Gamma[1 + n, (3*(a + b*ArcCosh[c*x])/b)))/(c^4*E^((3*a)/b)*Sqrt[d - c^2
*d*x^2]*(-(a + b*ArcCosh[c*x])^2/b^2))^(2*n))
```

**3.440.3 Rubi [A] (verified)**Time = 0.66 (sec) , antiderivative size = 279, normalized size of antiderivative = 0.74, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int (a + b \operatorname{arccosh}(cx))^n \cosh^3 \left( \frac{a}{b} - \frac{a+b \operatorname{arccosh}(cx)}{b} \right) d(a + b \operatorname{arccosh}(cx))}{bc^4 \sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int (a + b \operatorname{arccosh}(cx))^n \sin \left( \frac{ia}{b} - \frac{i(a+b \operatorname{arccosh}(cx))}{b} + \frac{\pi}{2} \right)^3 d(a + b \operatorname{arccosh}(cx))}{bc^4 \sqrt{d - c^2 dx^2}}$$

---

3.440.  $\int \frac{x^3(a+b \operatorname{arccosh}(cx))^n}{\sqrt{d-c^2 dx^2}} dx$

↓ 3793

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int \left( \frac{1}{4} \cosh\left(\frac{3a}{b} - \frac{3(a+\operatorname{arccosh}(cx))}{b}\right) (a + \operatorname{arccosh}(cx))^n + \frac{3}{4} \cosh\left(\frac{a}{b} - \frac{a+\operatorname{arccosh}(cx)}{b}\right) (a + \operatorname{arccosh}(cx))^n \right)}{bc^4\sqrt{d-c^2x^2}}$$

↓ 2009

$$\sqrt{cx-1}\sqrt{cx+1} \left( \frac{1}{8} b 3^{-n-1} e^{-\frac{3a}{b}} (a + \operatorname{arccosh}(cx))^n \left( -\frac{a+\operatorname{arccosh}(cx)}{b} \right)^{-n} \Gamma\left(n+1, -\frac{3(a+\operatorname{arccosh}(cx))}{b}\right) + \frac{3}{8} b e^{-\frac{a}{b}} \right)$$

input `Int[(x^3*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2], x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((3^(-1 - n))*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-3*(a + b*ArcCosh[c*x]))/b])/(8*E^((3*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n + (3*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b])/(8*E^(a/b)*(-(a + b*ArcCosh[c*x])/b))^n - (3*b*E^(a/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (a + b*ArcCosh[c*x])/b])/(8*((a + b*ArcCosh[c*x])/b)^n) - (3^(-1 - n))*b*E^((3*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (3*(a + b*ArcCosh[c*x]))/b])/(8*((a + b*ArcCosh[c*x])/b)^n))/(b*c^4*Sqrt[d - c^2*d*x^2])`

### 3.440.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`



rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

### 3.440.4 Maple [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2dx^2 + d}} dx$$

input `int(x^3*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

output `int(x^3*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

### 3.440.5 Fracas [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^n x^3}{\sqrt{-c^2dx^2 + d}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x^3/(c^2*d*x^2 - d), x)`

### 3.440.6 Sympy [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2dx^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^n}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate(x**3*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**3*(a + b*acosh(c*x))**n/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

---

3.440.  $\int \frac{x^3(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2dx^2}} dx$

**3.440.7 Maxima [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^3}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x^3/sqrt(-c^2*d*x^2 + d), x)`

**3.440.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.440.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^3*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^3*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(1/2), x)`

**3.441**  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx$

3.441.1 Optimal result . . . . . 3322  
 3.441.2 Mathematica [A] (verified) . . . . . 3323  
 3.441.3 Rubi [A] (verified) . . . . . 3323  
 3.441.4 Maple [F] . . . . . 3325  
 3.441.5 Fracas [F] . . . . . 3325  
 3.441.6 Sympy [F] . . . . . 3325  
 3.441.7 Maxima [F] . . . . . 3326  
 3.441.8 Giac [F] . . . . . 3326  
 3.441.9 Mupad [F(-1)] . . . . . 3326

**3.441.1 Optimal result**

Integrand size = 29, antiderivative size = 253

$$\int \frac{x^2(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx = \frac{\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^{1+n}}{2bc^3(1+n)\sqrt{d-c^2dx^2}} + \frac{2^{-3-n}e^{-\frac{2a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(-\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,-\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{c^3\sqrt{d-c^2dx^2}} - \frac{2^{-3-n}e^{\frac{2a}{b}}\sqrt{-1+cx}\sqrt{1+cx}(a+b\operatorname{arccosh}(cx))^n\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(1+n,\frac{2(a+b\operatorname{arccosh}(cx))}{b}\right)}{c^3\sqrt{d-c^2dx^2}}$$

```
output 1/2*(a+b*arccosh(c*x))^(1+n)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c^3/(1+n)/(-c^2
*d*x^2+d)^(1/2)+2^(-3-n)*(a+b*arccosh(c*x))^n*GAMMA(1+n,-2*(a+b*arccosh(c*
x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/exp(2*a/b)/(((a+b*arccosh(c*x))/b)
^n)/(-c^2*d*x^2+d)^(1/2)-2^(-3-n)*exp(2*a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+
n,2*(a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/(((a+b*arccosh(c
*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)
```

**3.441.2 Mathematica [A] (verified)**

Time = 0.63 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.84

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{2^{-3-n} e^{-\frac{2a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1+cx)(a + \operatorname{barccosh}(cx))^n \left(-\frac{(a+\operatorname{barccosh}(cx))^2}{b^2}\right)^{-n} \left(2^{2+n} e^{\frac{2a}{b}} (a + \operatorname{barccosh}(cx)) \left(-\frac{(a+\operatorname{barccosh}(cx))^2}{b^2}\right)^{-n} \right)}{\dots}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2],x]`output `(2^(-3 - n)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])^n*(2^(2 + n)*E^((2*a)/b)*(a + b*ArcCosh[c*x])*(-(a + b*ArcCosh[c*x])^2/b^2))^n + b*(1 + n)*(a/b + ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x])/b] - b*E^((4*a)/b)*(1 + n)*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x])/b)])/ (b*c^3*E^((2*a)/b)*(1 + n)*Sqrt[d - c^2*d*x^2]*(-(a + b*ArcCosh[c*x])^2/b^2))^n`**3.441.3 Rubi [A] (verified)**Time = 0.57 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.73, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.138$ , Rules used = {6367, 3042, 3793, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int (a + \operatorname{barccosh}(cx))^n \cosh^2\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^3\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int (a + \operatorname{barccosh}(cx))^n \sin\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)^2 d(a + \operatorname{barccosh}(cx))}{bc^3\sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{3793}$$

---

3.441.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \int \left( \frac{1}{2} \cosh\left(\frac{2a}{b} - \frac{2(a+\operatorname{barccosh}(cx))}{b}\right) (a + \operatorname{barccosh}(cx))^n + \frac{1}{2}(a + \operatorname{barccosh}(cx))^n \right) d(a + \operatorname{barccosh}(cx))}{bc^3\sqrt{d-c^2dx^2}}$$

↓ 2009

$$\frac{\sqrt{cx-1}\sqrt{cx+1} \left( \frac{(a+\operatorname{barccosh}(cx))^{n+1}}{2(n+1)} + b2^{-n-3}e^{-\frac{2a}{b}}(a + \operatorname{barccosh}(cx))^n \left( -\frac{a+\operatorname{barccosh}(cx)}{b} \right)^{-n} \Gamma\left(n+1, -\frac{2(a+\operatorname{barccosh}(cx))}{b}\right) \right)}{bc^3\sqrt{d-c^2dx^2}}$$

input `Int[(x^2*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2], x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((a + b*ArcCosh[c*x])^(1 + n)/(2*(1 + n)) + (2^(-3 - n)*b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (-2*(a + b*ArcCosh[c*x])/b)]/b))/(E^((2*a)/b)*(-(a + b*ArcCosh[c*x])/b))^n - (2^(-3 - n)*b*E^((2*a)/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (2*(a + b*ArcCosh[c*x])/b)]/(a + b*ArcCosh[c*x])/b))^n)/(b*c^3*Sqrt[d - c^2*d*x^2])`

### 3.441.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3793 `Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))`

rule 6367 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

**3.441.4 Maple [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2 d x^2 + d}} dx$$

input `int(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

output `int(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

**3.441.5 Fracas [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^n x^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x^2/(c^2*d*x^2 - d), x)`

**3.441.6 Sympy [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^n}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x**2*(a + b*acosh(c*x))**n/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.441.7 Maxima [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x^2/sqrt(-c^2*d*x^2 + d), x)`

**3.441.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^2}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n*x^2/sqrt(-c^2*d*x^2 + d), x)`

**3.441.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x^2*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(1/2),x)`

output `int((x^2*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(1/2), x)`

$$3.442 \quad \int \frac{x(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx$$

3.442.1 Optimal result . . . . .	3327
3.442.2 Mathematica [A] (verified) . . . . .	3327
3.442.3 Rubi [A] (verified) . . . . .	3328
3.442.4 Maple [F] . . . . .	3330
3.442.5 Fricas [F] . . . . .	3330
3.442.6 Sympy [F] . . . . .	3330
3.442.7 Maxima [F] . . . . .	3331
3.442.8 Giac [F] . . . . .	3331
3.442.9 Mupad [F(-1)] . . . . .	3331

### 3.442.1 Optimal result

Integrand size = 27, antiderivative size = 182

$$\int \frac{x(a + \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2dx^2}} dx$$

$$= \frac{e^{-\frac{a}{b}} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{arccosh}(cx))^n \left(-\frac{a + \operatorname{arccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, -\frac{a + \operatorname{arccosh}(cx)}{b}\right)}{2c^2 \sqrt{d - c^2dx^2}} - \frac{e^{a/b} \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{arccosh}(cx))^n \left(\frac{a + \operatorname{arccosh}(cx)}{b}\right)^{-n} \Gamma\left(1 + n, \frac{a + \operatorname{arccosh}(cx)}{b}\right)}{2c^2 \sqrt{d - c^2dx^2}}$$

```
output 1/2*(a+b*arccosh(c*x))^n*GAMMA(1+n, (-a-b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^2/exp(a/b)/((((-a-b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)-1/2*exp(a/b)*(a+b*arccosh(c*x))^n*GAMMA(1+n, (a+b*arccosh(c*x))/b)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^2/(((a+b*arccosh(c*x))/b)^n)/(-c^2*d*x^2+d)^(1/2)
```

### 3.442.2 Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 153, normalized size of antiderivative = 0.84

$$\int \frac{x(a + \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2dx^2}} dx$$

$$= \frac{e^{-\frac{a}{b}} \sqrt{\frac{-1+cx}{1+cx}} (1 + cx) (a + \operatorname{arccosh}(cx))^n \left(-\frac{(a + \operatorname{arccosh}(cx))^2}{b^2}\right)^{-n} \left(-e^{\frac{2a}{b}} \left(-\frac{a + \operatorname{arccosh}(cx)}{b}\right)^n \Gamma\left(1 + n, \frac{a}{b} + \dots\right)\right)}{2c^2 \sqrt{-d(-1 + cx)(1 + cx)}}$$

---

3.442.  $\int \frac{x(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx$



input `Integrate[(x*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2],x]`

output `(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*(a + b*ArcCosh[c*x])^n*(-(E^((2*a)/b))*(-(a + b*ArcCosh[c*x])/b))^n*Gamma[1 + n, a/b + ArcCosh[c*x]]) + (a/b + ArcCosh[c*x])^n*Gamma[1 + n, -(a + b*ArcCosh[c*x])/b])/(2*c^2*E^(a/b)*Sqrt[-(d*(-1 + c*x)*(1 + c*x))]*(-(a + b*ArcCosh[c*x])^2/b^2))^n)`

### 3.442.3 Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.84, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$ , Rules used = {6367, 3042, 3788, 26, 2612}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

$$\downarrow \text{6367}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int (a + \operatorname{barccosh}(cx))^n \cosh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{bc^2 \sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{3042}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \int (a + \operatorname{barccosh}(cx))^n \sin\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right) d(a + \operatorname{barccosh}(cx))}{bc^2 \sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{3788}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \left(\frac{1}{2}i \int -ie^{-\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))^n d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i \int ie^{\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))^n d(a + \operatorname{barccosh}(cx))\right)}{bc^2 \sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{26}$$

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1} \left(\frac{1}{2} \int e^{-\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))^n d(a + \operatorname{barccosh}(cx)) + \frac{1}{2} \int e^{\operatorname{arccosh}(cx)}(a + \operatorname{barccosh}(cx))^n d(a + \operatorname{barccosh}(cx))\right)}{bc^2 \sqrt{d - c^2 dx^2}}$$

$$\downarrow \text{2612}$$

---

3.442.  $\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$

$$\frac{\sqrt{cx-1}\sqrt{cx+1}\left(\frac{1}{2}be^{-\frac{a}{b}}(a+\operatorname{arccosh}(cx))^n\left(-\frac{a+\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(n+1,-\frac{a+\operatorname{arccosh}(cx)}{b}\right)-\frac{1}{2}be^{a/b}(a+\operatorname{arccosh}(cx))^n\left(\frac{a+\operatorname{arccosh}(cx)}{b}\right)^{-n}\Gamma\left(n+1,\frac{a+\operatorname{arccosh}(cx)}{b}\right)\right)}{bc^2\sqrt{d-c^2dx^2}}$$

input `Int[(x*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2],x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*((b*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, -((a + b*ArcCosh[c*x])/b)])/(2*E^(a/b)*(-(a + b*ArcCosh[c*x])/b))^n - (b*E^(a/b)*(a + b*ArcCosh[c*x])^n*Gamma[1 + n, (a + b*ArcCosh[c*x])/b])/(2*((a + b*ArcCosh[c*x])/b)^n)))/(b*c^2*Sqrt[d - c^2*d*x^2])`

### 3.442.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2612 `Int[(F_)^((g_)*(e_) + (f_)*(x_))*((c_) + (d_)*(x_))^(m_), x_Symbol] := Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*(-f)*g*(Log[F]/d))^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m])*Gamma[m + 1, ((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] && !IntegerQ[m]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3788 `Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + Pi*(k_) + (f_)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/(E^(I*k*Pi)*E^(I*(e + f*x))), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*k*Pi)*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x] && IntegerQ[2*k]`

rule 6367 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d + e*x^2)^p/((1 + c*x)^p*(-1 + c*x)^p)] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && IGtQ[2*p + 2, 0] && IGtQ[m, 0]`

**3.442.4 Maple [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2 d x^2 + d}} dx$$

input `int(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

output `int(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

**3.442.5 Fricas [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^n x}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x/(c^2*d*x^2 - d), x)`

**3.442.6 Sympy [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^n}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate(x*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral(x*(a + b*acosh(c*x))**n/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.442.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x/sqrt(-c^2*d*x^2 + d), x)`

**3.442.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n*x/sqrt(-c^2*d*x^2 + d), x)`

**3.442.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

input `int((x*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(1/2),x)`

output `int((x*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(1/2), x)`

**3.443**  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{\sqrt{d-c^2dx^2}} dx$

3.443.1 Optimal result . . . . . 3332  
 3.443.2 Mathematica [A] (verified) . . . . . 3332  
 3.443.3 Rubi [A] (verified) . . . . . 3333  
 3.443.4 Maple [A] (verified) . . . . . 3333  
 3.443.5 Fricas [B] (verification not implemented) . . . . . 3334  
 3.443.6 Sympy [F] . . . . . 3334  
 3.443.7 Maxima [F] . . . . . 3335  
 3.443.8 Giac [F] . . . . . 3335  
 3.443.9 Mupad [F(-1)] . . . . . 3335

**3.443.1 Optimal result**

Integrand size = 26, antiderivative size = 57

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^{1+n}}{bc(1 + n)\sqrt{d - c^2dx^2}}$$

output `(a+b*arccosh(c*x))^(1+n)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(1+n)/(-c^2*d*x^2+d)^(1/2)`

**3.443.2 Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2dx^2}} dx = \frac{\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{arccosh}(cx))^{1+n}}{bc(1 + n)\sqrt{d - c^2dx^2}}$$

input `Integrate[(a + b*ArcCosh[c*x])^n/Sqrt[d - c^2*d*x^2],x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^(1 + n))/(b*c*(1 + n)*Sqrt[d - c^2*d*x^2])`

**3.443.3 Rubi [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$ , Rules used = {6307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6307

$$\frac{\sqrt{cx - 1}\sqrt{cx + 1}(a + \operatorname{arccosh}(cx))^{n+1}}{bc(n + 1)\sqrt{d - c^2 dx^2}}$$

input `Int[(a + b*ArcCosh[c*x])^n/Sqrt[d - c^2*d*x^2],x]`

output `(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x])^(1 + n))/(b*c*(1 + n)*Sqrt[d - c^2*d*x^2])`

**3.443.3.1 Defintions of rubi rules used**

rule 6307 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)/Sqrt[(d_) + (e_.)*(x_)^2], x_Symbol] :> Simp[(1/(b*c*(n + 1)))*Simp[Sqrt[1 + c*x]*(Sqrt[-1 + c*x]/Sqrt[d + e*x^2])]*(a + b*ArcCosh[c*x])^(n + 1), x] /; FreeQ[{a, b, c, d, e, n}, x] && EqQ[c^2*d + e, 0] && NeQ[n, -1]`

**3.443.4 Maple [A] (verified)**

Time = 0.56 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.95

method	result	size
default	$\frac{\sqrt{cx-1}\sqrt{cx+1}(a+b\operatorname{arccosh}(cx))^{1+n}}{bc(1+n)\sqrt{-d(cx-1)(cx+1)}}$	54

input `int((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x,method=_RETURNVERBOSE)`

output  $1/b/c/(1+n)*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(a+b*\operatorname{arccosh}(c*x))^{(1+n)/(-d*(c*x-1)*(c*x+1))^{(1/2)}$

### 3.443.5 Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 221 vs.  $2(51) = 102$ .

Time = 0.28 (sec) , antiderivative size = 221, normalized size of antiderivative = 3.88

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

$$= \frac{(\sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} b \log(cx + \sqrt{c^2 x^2 - 1}) + \sqrt{-c^2 dx^2 + d} \sqrt{c^2 x^2 - 1} a) \cosh(n \log(b \log(cx + \sqrt{c^2 x^2 - 1}) + a))}{(b^2 c^2 d^2 n^2 + b^2 c^2 d^2 - (b^2 c^3 d^2 n^2 + b^2 c^3 d^2) x^2)}$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output  $((\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*b*\log(c*x + \sqrt{c^2*x^2 - 1}) + \sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*a)*\cosh(n*\log(b*\log(c*x + \sqrt{c^2*x^2 - 1}) + a)) + (\sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*b*\log(c*x + \sqrt{c^2*x^2 - 1}) + \sqrt{-c^2*d*x^2 + d}*\sqrt{c^2*x^2 - 1}*a)*\sinh(n*\log(b*\log(c*x + \sqrt{c^2*x^2 - 1}) + a)))/(b*c*d*n^2 + b*c*d - (b*c^3*d*n^2 + b*c^3*d)*x^2)$

### 3.443.6 Sympy [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**n/sqrt(-d*(c*x - 1)*(c*x + 1)), x)`

**3.443.7 Maxima [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/sqrt(-c^2*d*x^2 + d), x)`

**3.443.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n/sqrt(-c^2*d*x^2 + d), x)`

**3.443.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^n/(d - c^2*d*x^2)^(1/2),x)`

output `int((a + b*acosh(c*x))^n/(d - c^2*d*x^2)^(1/2), x)`



**3.444**  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{x\sqrt{d-c^2dx^2}} dx$

3.444.1 Optimal result	3336
3.444.2 Mathematica [N/A]	3336
3.444.3 Rubi [N/A]	3337
3.444.4 Maple [N/A] (verified)	3337
3.444.5 Fricas [N/A]	3338
3.444.6 Sympy [N/A]	3338
3.444.7 Maxima [N/A]	3338
3.444.8 Giac [N/A]	3339
3.444.9 Mupad [N/A]	3339

**3.444.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{d - c^2dx^2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{d - c^2dx^2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2), x)`

**3.444.2 Mathematica [N/A]**

Not integrable

Time = 0.48 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{d - c^2dx^2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{d - c^2dx^2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^n/(x*Sqrt[d - c^2*d*x^2]), x]`

output `Integrate[(a + b*ArcCosh[c*x])^n/(x*Sqrt[d - c^2*d*x^2]), x]`

**3.444.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{d - c^2 dx^2}} dx$$

↓ 6375

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x\sqrt{d - c^2 dx^2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^n/(x*sqrt[d - c^2*d*x^2]),x]`

output `$Aborted`

**3.444.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.444.4 Maple [N/A] (verified)**

Not integrable

Time = 1.32 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x\sqrt{-c^2 dx^2 + d}} dx$$

input `int((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2),x)`

output `int((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2),x)`

**3.444.5 Fracas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.45

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2dx^2 + dx}} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="fracas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/(c^2*d*x^3 - d*x), x)`

**3.444.6 Sympy [N/A]**

Not integrable

Time = 6.59 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{d - c^2dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x\sqrt{-d}(cx - 1)(cx + 1)} dx$$

input `integrate((a+b*acosh(c*x))**n/x/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**n/(x*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.444.7 Maxima [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{d - c^2dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2dx^2 + dx}} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*d*x^2 + d)*x), x)`

---

3.444.  $\int \frac{(a + \operatorname{barccosh}(cx))^n}{x\sqrt{d - c^2dx^2}} dx$

**3.444.8 Giac [N/A]**

Not integrable

Time = 12.45 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*d*x^2 + d)*x), x)`**3.444.9 Mupad [N/A]**

Not integrable

Time = 3.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^n/(x*(d - c^2*d*x^2)^(1/2)),x)`output `int((a + b*acosh(c*x))^n/(x*(d - c^2*d*x^2)^(1/2)), x)`

**3.445**  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{x^2\sqrt{d-c^2dx^2}} dx$

3.445.1 Optimal result . . . . .	3340
3.445.2 Mathematica [N/A] . . . . .	3340
3.445.3 Rubi [N/A] . . . . .	3341
3.445.4 Maple [N/A] (verified) . . . . .	3341
3.445.5 Fricas [N/A] . . . . .	3342
3.445.6 Sympy [N/A] . . . . .	3342
3.445.7 Maxima [N/A] . . . . .	3342
3.445.8 Giac [N/A] . . . . .	3343
3.445.9 Mupad [N/A] . . . . .	3343

**3.445.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{d - c^2dx^2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{d - c^2dx^2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(1/2), x)`

**3.445.2 Mathematica [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{d - c^2dx^2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2\sqrt{d - c^2dx^2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^n/(x^2*Sqrt[d - c^2*d*x^2]), x]`

output `Integrate[(a + b*ArcCosh[c*x])^n/(x^2*Sqrt[d - c^2*d*x^2]), x]`

**3.445.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx$$

↓ 6375

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^n/(x^2*sqrt[d - c^2*d*x^2]),x]`

output `$Aborted`

**3.445.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.445.4 Maple [N/A] (verified)**

Not integrable

Time = 1.47 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x^2 \sqrt{-c^2 d x^2 + d}} dx$$

input `int((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(1/2),x)`

output `int((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(1/2),x)`

**3.445.5 Fracas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.52

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + dx^2}} dx$$

input `integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/(c^2*d*x^4 - d*x^2), x)`

**3.445.6 Sympy [N/A]**

Not integrable

Time = 31.69 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x^2 \sqrt{-d(cx - 1)(cx + 1)}} dx$$

input `integrate((a+b*acosh(c*x))**n/x**2/(-c**2*d*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**n/(x**2*sqrt(-d*(c*x - 1)*(c*x + 1))), x)`

**3.445.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + dx^2}} dx$$

input `integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*d*x^2 + d)*x^2), x)`

---

3.445.  $\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx$

**3.445.8 Giac [N/A]**

Not integrable

Time = 12.59 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + dx^2}} dx$$

input `integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n/(sqrt(-c^2*d*x^2 + d)*x^2), x)`

**3.445.9 Mupad [N/A]**

Not integrable

Time = 3.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x^2 \sqrt{d - c^2 dx^2}} dx$$

input `int((a + b*acosh(c*x))^n/(x^2*(d - c^2*d*x^2)^(1/2)),x)`

output `int((a + b*acosh(c*x))^n/(x^2*(d - c^2*d*x^2)^(1/2)), x)`



$$3.446 \quad \int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

3.446.1 Optimal result	3344
3.446.2 Mathematica [N/A]	3344
3.446.3 Rubi [N/A]	3345
3.446.4 Maple [N/A] (verified)	3345
3.446.5 Fricas [N/A]	3346
3.446.6 Sympy [N/A]	3346
3.446.7 Maxima [N/A]	3346
3.446.8 Giac [N/A]	3347
3.446.9 Mupad [N/A]	3347

### 3.446.1 Optimal result

Integrand size = 29, antiderivative size = 29

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \operatorname{Int}\left(\frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}}, x\right)$$

output `Unintegrable(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2), x)`

### 3.446.2 Mathematica [N/A]

Not integrable

Time = 0.94 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2), x]`

output `Integrate[(x^2*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2), x]`

---


$$3.446. \quad \int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

**3.446.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

↓ 6375

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `Int[(x^2*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2),x]`

output `$Aborted`

**3.446.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.446.4 Maple [N/A] (verified)**

Not integrable

Time = 1.45 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))^n}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `int(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

output `int(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

---

3.446.  $\int \frac{x^2(a+b \operatorname{arccosh}(cx))^n}{(d-c^2 dx^2)^{3/2}} dx$

**3.446.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.93

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^2}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x^2/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.446.6 Sympy [N/A]**

Not integrable

Time = 67.56 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^n}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x**2*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x**2*(a + b*acosh(c*x))**n/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.446.7 Maxima [N/A]**

Not integrable

Time = 0.58 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^2}{(-c^2dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x^2/(-c^2*d*x^2 + d)^(3/2), x)`

---

3.446.  $\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx$

**3.446.8 Giac [N/A]**

Not integrable

Time = 12.74 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x^2}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n*x^2/(-c^2*d*x^2 + d)^(3/2), x)`

**3.446.9 Mupad [N/A]**

Not integrable

Time = 3.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x^2*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(3/2),x)`

output `int((x^2*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(3/2), x)`

**3.447**  $\int \frac{x(a+b\operatorname{arccosh}(cx))^n}{(d-c^2dx^2)^{3/2}} dx$

3.447.1 Optimal result . . . . . 3348  
 3.447.2 Mathematica [N/A] . . . . . 3348  
 3.447.3 Rubi [N/A] . . . . . 3349  
 3.447.4 Maple [N/A] (verified) . . . . . 3349  
 3.447.5 Fricas [N/A] . . . . . 3350  
 3.447.6 Sympy [N/A] . . . . . 3350  
 3.447.7 Maxima [N/A] . . . . . 3350  
 3.447.8 Giac [N/A] . . . . . 3351  
 3.447.9 Mupad [N/A] . . . . . 3351

**3.447.1 Optimal result**

Integrand size = 27, antiderivative size = 27

$$\int \frac{x(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx = \operatorname{Int}\left(\frac{x(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}}, x\right)$$

output `Unintegrable(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2), x)`

**3.447.2 Mathematica [N/A]**

Not integrable

Time = 0.63 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{x(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx = \int \frac{x(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx$$

input `Integrate[(x*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2), x]`

output `Integrate[(x*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2), x]`

**3.447.3 Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

↓ 6375

$$\int \frac{x(a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `Int[(x*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2),x]`

output `$Aborted`

**3.447.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_)^m)*((d_) + (e_.)*(x_)^2)^p, x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.447.4 Maple [N/A] (verified)**

Not integrable

Time = 1.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `int(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

output `int(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

---

3.447.  $\int \frac{x(a+b\operatorname{arccosh}(cx))^n}{(d-c^2dx^2)^{3/2}} dx$

**3.447.5 Fracas [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.00

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n*x/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.447.6 Sympy [N/A]**

Not integrable

Time = 67.70 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^n}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral(x*(a + b*acosh(c*x))**n/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.447.7 Maxima [N/A]**

Not integrable

Time = 0.52 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{x(a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n*x/(-c^2*d*x^2 + d)^(3/2), x)`

---

3.447.  $\int \frac{x(a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$

**3.447.8 Giac [N/A]**

Not integrable

Time = 12.57 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n x}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate(x*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^n*x/(-c^2*d*x^2 + d)^(3/2), x)`**3.447.9 Mupad [N/A]**

Not integrable

Time = 3.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{x(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{x(a + b \operatorname{acosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((x*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(3/2),x)`output `int((x*(a + b*acosh(c*x))^n)/(d - c^2*d*x^2)^(3/2), x)`



**3.448**  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{(d-c^2dx^2)^{3/2}} dx$

3.448.1 Optimal result . . . . .	3352
3.448.2 Mathematica [N/A] . . . . .	3352
3.448.3 Rubi [N/A] . . . . .	3353
3.448.4 Maple [N/A] (verified) . . . . .	3353
3.448.5 Fricas [N/A] . . . . .	3354
3.448.6 Sympy [N/A] . . . . .	3354
3.448.7 Maxima [N/A] . . . . .	3354
3.448.8 Giac [N/A] . . . . .	3355
3.448.9 Mupad [N/A] . . . . .	3355

**3.448.1 Optimal result**

Integrand size = 26, antiderivative size = 26

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2), x)`

**3.448.2 Mathematica [N/A]**

Not integrable

Time = 0.13 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2dx^2)^{3/2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^n/(d - c^2*d*x^2)^(3/2), x]`

output `Integrate[(a + b*ArcCosh[c*x])^n/(d - c^2*d*x^2)^(3/2), x]`

**3.448.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

↓ 6325

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^n/(d - c^2*d*x^2)^(3/2),x]`

output `$Aborted`

**3.448.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.448.4 Maple [N/A] (verified)**

Not integrable

Time = 1.38 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `int((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

output `int((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

---

3.448.  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{(d-c^2 dx^2)^{3/2}} dx$

**3.448.5 Fracas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.04

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.448.6 Sympy [N/A]**

Not integrable

Time = 49.66 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.04

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{(-d(cx - 1)(cx + 1))^{\frac{3}{2}}} dx$$

input `integrate((a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(3/2),x)`

output `Integral((a + b*acosh(c*x))**n/(-d*(c*x - 1)*(c*x + 1))**(3/2), x)`

**3.448.7 Maxima [N/A]**

Not integrable

Time = 0.55 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/(-c^2*d*x^2 + d)^(3/2), x)`

---

3.448.  $\int \frac{(a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$

**3.448.8 Giac [N/A]**

Not integrable

Time = 12.62 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^n/(-c^2*d*x^2 + d)^(3/2), x)`**3.448.9 Mupad [N/A]**

Not integrable

Time = 3.19 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^n/(d - c^2*d*x^2)^(3/2),x)`output `int((a + b*acosh(c*x))^n/(d - c^2*d*x^2)^(3/2), x)`

**3.449**  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{x(d-c^2dx^2)^{3/2}} dx$

3.449.1 Optimal result . . . . .	3356
3.449.2 Mathematica [N/A] . . . . .	3356
3.449.3 Rubi [N/A] . . . . .	3357
3.449.4 Maple [N/A] (verified) . . . . .	3357
3.449.5 Fricas [N/A] . . . . .	3358
3.449.6 Sympy [F(-1)] . . . . .	3358
3.449.7 Maxima [N/A] . . . . .	3358
3.449.8 Giac [N/A] . . . . .	3359
3.449.9 Mupad [N/A] . . . . .	3359

**3.449.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2dx^2)^{3/2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2dx^2)^{3/2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(3/2), x)`

**3.449.2 Mathematica [N/A]**

Not integrable

Time = 0.67 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2dx^2)^{3/2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2dx^2)^{3/2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^n/(x*(d - c^2*d*x^2)^(3/2)), x]`

output `Integrate[(a + b*ArcCosh[c*x])^n/(x*(d - c^2*d*x^2)^(3/2)), x]`

**3.449.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx$$

↓ 6375

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^n/(x*(d - c^2*d*x^2)^(3/2)),x]`

output `$Aborted`

**3.449.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_)^m)*((d_) + (e_.)*(x_)^2)^p, x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.449.4 Maple [N/A] (verified)**

Not integrable

Time = 1.52 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x(-c^2 dx^2 + d)^{3/2}} dx$$

input `int((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(3/2),x)`

output `int((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(3/2),x)`

---

3.449.  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{x(d-c^2 dx^2)^{3/2}} dx$

**3.449.5 Fricas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.90

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/(c^4*d^2*x^5 - 2*c^2*d^2*x^3 + d^2*x), x)`

**3.449.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))**n/x/(-c**2*d*x**2+d)**(3/2), x)`

output `Timed out`

**3.449.7 Maxima [N/A]**

Not integrable

Time = 0.54 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/((-c^2*d*x^2 + d)^(3/2)*x), x)`

---

3.449.  $\int \frac{(a + \operatorname{barccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx$

**3.449.8 Giac [N/A]**

Not integrable

Time = 12.58 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}} x} dx$$

input `integrate((a+b*arccosh(c*x))^n/x/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^n/((-c^2*d*x^2 + d)^(3/2)*x), x)`**3.449.9 Mupad [N/A]**

Not integrable

Time = 3.70 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x(d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^n/(x*(d - c^2*d*x^2)^(3/2)),x)`output `int((a + b*acosh(c*x))^n/(x*(d - c^2*d*x^2)^(3/2)), x)`



$$3.450 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^n}{x^2(d-c^2dx^2)^{3/2}} dx$$

3.450.1 Optimal result	3360
3.450.2 Mathematica [N/A]	3360
3.450.3 Rubi [N/A]	3361
3.450.4 Maple [N/A] (verified)	3361
3.450.5 Fricas [N/A]	3362
3.450.6 Sympy [F(-1)]	3362
3.450.7 Maxima [N/A]	3362
3.450.8 Giac [N/A]	3363
3.450.9 Mupad [N/A]	3363

### 3.450.1 Optimal result

Integrand size = 29, antiderivative size = 29

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2(d - c^2dx^2)^{3/2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^n}{x^2(d - c^2dx^2)^{3/2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(3/2), x)`

### 3.450.2 Mathematica [N/A]

Not integrable

Time = 0.69 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2(d - c^2dx^2)^{3/2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2(d - c^2dx^2)^{3/2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^n/(x^2*(d - c^2*d*x^2)^(3/2)), x]`

output `Integrate[(a + b*ArcCosh[c*x])^n/(x^2*(d - c^2*d*x^2)^(3/2)), x]`

---


$$3.450. \quad \int \frac{(a+b\operatorname{arccosh}(cx))^n}{x^2(d-c^2dx^2)^{3/2}} dx$$

**3.450.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx$$

↓ 6375

$$\int \frac{(a + \operatorname{arccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^n/(x^2*(d - c^2*d*x^2)^(3/2)),x]`

output `$Aborted`

**3.450.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_)^m)*((d_) + (e_.)*(x_)^2)^p, x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.450.4 Maple [N/A] (verified)**

Not integrable

Time = 1.55 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x^2 (-c^2 dx^2 + d)^{3/2}} dx$$

input `int((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(3/2),x)`

output `int((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(3/2),x)`

---

3.450.  $\int \frac{(a+b\operatorname{arccosh}(cx))^n}{x^2(d-c^2dx^2)^{3/2}} dx$

**3.450.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.97

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="fricas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(b*arccosh(c*x) + a)^n/(c^4*d^2*x^6 - 2*c^2*d^2*x^4 + d^2*x^2), x)`

**3.450.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))**n/x**2/(-c**2*d*x**2+d)**(3/2),x)`

output `Timed out`

**3.450.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^n/((-c^2*d*x^2 + d)^(3/2)*x^2), x)`

---

3.450.  $\int \frac{(a + \operatorname{barccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx$

**3.450.8 Giac [N/A]**

Not integrable

Time = 12.52 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}} x^2} dx$$

input `integrate((a+b*arccosh(c*x))^n/x^2/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^n/((-c^2*d*x^2 + d)^(3/2)*x^2), x)`

**3.450.9 Mupad [N/A]**

Not integrable

Time = 3.38 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n}{x^2 (d - c^2 dx^2)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^n/(x^2*(d - c^2*d*x^2)^(3/2)),x)`

output `int((a + b*acosh(c*x))^n/(x^2*(d - c^2*d*x^2)^(3/2)), x)`

**3.451** 
$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$$

3.451.1 Optimal result	3364
3.451.2 Mathematica [N/A]	3364
3.451.3 Rubi [N/A]	3365
3.451.4 Maple [N/A] (verified)	3365
3.451.5 Fricas [N/A]	3366
3.451.6 Sympy [N/A]	3366
3.451.7 Maxima [N/A]	3366
3.451.8 Giac [N/A]	3367
3.451.9 Mupad [N/A]	3367

**3.451.1 Optimal result**

Integrand size = 30, antiderivative size = 30

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \operatorname{Int}\left(\frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2), x)`

**3.451.2 Mathematica [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2], x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2], x]`

**3.451.3 Rubi [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^n)/Sqrt[1 - c^2*x^2],x]`

output `$Aborted`

**3.451.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n_.*((f_.)*(x_))^(m_.)*((d_ + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.451.4 Maple [N/A] (verified)**

Not integrable

Time = 1.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2 x^2 + 1}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x)`

**3.451.5 Fricas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.40

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(fx)^m (b \operatorname{arccosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="fricas")`

output `integral(-sqrt(-c^2*x^2 + 1)*(f*x)^m*(b*arccosh(c*x) + a)^n/(c^2*x^2 - 1), x)`

**3.451.6 Sympy [N/A]**

Not integrable

Time = 89.72 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.03

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))^n}{\sqrt{-(cx - 1)(cx + 1)}} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))**n/(-c**2*x**2+1)**(1/2),x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))**n/sqrt(-(c*x - 1)*(c*x + 1)), x)`

**3.451.7 Maxima [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(fx)^m (b \operatorname{arccosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="maxima")`

output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n/sqrt(-c^2*x^2 + 1), x)`

---

3.451.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx$

**3.451.8 Giac [N/A]**

Not integrable

Time = 12.61 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 x^2 + 1}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*x^2+1)^(1/2),x, algorithm="giac")`

output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n/sqrt(-c^2*x^2 + 1), x)`

**3.451.9 Mupad [N/A]**

Not integrable

Time = 3.44 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{1 - c^2 x^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (fx)^m}{\sqrt{1 - c^2 x^2}} dx$$

input `int(((a + b*acosh(c*x))^n*(f*x)^m)/(1 - c^2*x^2)^(1/2),x)`

output `int(((a + b*acosh(c*x))^n*(f*x)^m)/(1 - c^2*x^2)^(1/2), x)`



### 3.452 $\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx$

3.452.1 Optimal result . . . . .	3368
3.452.2 Mathematica [N/A] . . . . .	3368
3.452.3 Rubi [N/A] . . . . .	3369
3.452.4 Maple [N/A] (verified) . . . . .	3369
3.452.5 Fricas [N/A] . . . . .	3370
3.452.6 Sympy [F(-1)] . . . . .	3370
3.452.7 Maxima [N/A] . . . . .	3370
3.452.8 Giac [F(-2)] . . . . .	3371
3.452.9 Mupad [N/A] . . . . .	3371

#### 3.452.1 Optimal result

Integrand size = 29, antiderivative size = 29

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx = \operatorname{Int}\left((fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n, x\right)$$

output `Unintegrable((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))^n,x)`

#### 3.452.2 Mathematica [N/A]

Not integrable

Time = 0.62 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx = \int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x])^n,x]`

output `Integrate[(f*x)^m*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x])^n, x]`

**3.452.3 Rubi [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^2 (fx)^m (a + \operatorname{barccosh}(cx))^n dx$$

↓ 6375

$$\int (d - c^2 dx^2)^2 (fx)^m (a + \operatorname{barccosh}(cx))^n dx$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^2*(a + b*ArcCosh[c*x])^n,x]`

output `$Aborted`

**3.452.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n]*((f_.)*(x_.))^m*((d_.) + (e_.)*(x_)^2)^p, x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.452.4 Maple [N/A] (verified)**

Not integrable

Time = 0.52 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int (fx)^m (-c^2 dx^2 + d)^2 (a + b \operatorname{arccosh}(cx))^n dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))^n,x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))^n,x)`

**3.452.5 Fricas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.48

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx = \int (c^2 dx^2 - d)^2 (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))^n,x, algorithm="fricas")`

output `integral((c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2)*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

**3.452.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**2*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.452.7 Maxima [N/A]**

Not integrable

Time = 1.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.10

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx = \int (c^2 dx^2 - d)^2 (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((c^2*d*x^2 - d)^2*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

---

3.452.  $\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx$

**3.452.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^2*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.452.9 Mupad [N/A]**

Not integrable

Time = 3.10 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int (fx)^m (d - c^2 dx^2)^2 (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^2 (fx)^m dx$$

input `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^2*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^2*(f*x)^m, x)`

### 3.453 $\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx$

3.453.1 Optimal result . . . . .	3372
3.453.2 Mathematica [N/A] . . . . .	3372
3.453.3 Rubi [N/A] . . . . .	3373
3.453.4 Maple [N/A] (verified) . . . . .	3373
3.453.5 Fricas [N/A] . . . . .	3374
3.453.6 Sympy [F(-1)] . . . . .	3374
3.453.7 Maxima [N/A] . . . . .	3374
3.453.8 Giac [F(-2)] . . . . .	3375
3.453.9 Mupad [N/A] . . . . .	3375

#### 3.453.1 Optimal result

Integrand size = 27, antiderivative size = 27

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx = \operatorname{Int}((fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n, x)$$

output `Unintegrable((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x))^n,x)`

#### 3.453.2 Mathematica [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx = \int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x])^n,x]`

output `Integrate[(f*x)^m*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x])^n, x]`

**3.453.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2) (fx)^m (a + \text{barccosh}(cx))^n dx$$

↓ 6375

$$\int (d - c^2 dx^2) (fx)^m (a + \text{barccosh}(cx))^n dx$$

input `Int[(f*x)^m*(d - c^2*d*x^2)*(a + b*ArcCosh[c*x])^n,x]`

output `$Aborted`

**3.453.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n]*((f_.)*(x_.))^m*((d_.) + (e_.)*(x_.)^2)^p, x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.453.4 Maple [N/A] (verified)**

Not integrable

Time = 0.31 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int (fx)^m (-c^2 dx^2 + d) (a + b \text{arccosh}(cx))^n dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x))^n,x)`

output `int((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x))^n,x)`

**3.453.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.15

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx = \int -(c^2 dx^2 - d)(fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x))^n,x, algorithm="fricas")`

output `integral(-(c^2*d*x^2 - d)*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

**3.453.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.453.7 Maxima [N/A]**

Not integrable

Time = 1.05 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.19

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx = \int -(c^2 dx^2 - d)(fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `-integrate((c^2*d*x^2 - d)*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

**3.453.8 Giac [F(-2)]**

Exception generated.

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx = \text{Exception raised: TypeError}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.453.9 Mupad [N/A]**

Not integrable

Time = 3.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int (fx)^m (d - c^2 dx^2) (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2) (fx)^m dx$$

input `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)*(f*x)^m, x)`



### 3.454 $\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx$

3.454.1 Optimal result	3376
3.454.2 Mathematica [N/A]	3376
3.454.3 Rubi [N/A]	3377
3.454.4 Maple [N/A] (verified)	3377
3.454.5 Fricas [N/A]	3378
3.454.6 Sympy [N/A]	3378
3.454.7 Maxima [N/A]	3378
3.454.8 Giac [F(-1)]	3379
3.454.9 Mupad [N/A]	3379

#### 3.454.1 Optimal result

Integrand size = 16, antiderivative size = 16

$$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx = \operatorname{Int}((fx)^m (a + \operatorname{barccosh}(cx))^n, x)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^n,x)`

#### 3.454.2 Mathematica [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx = \int (fx)^m (a + \operatorname{barccosh}(cx))^n dx$$

input `Integrate[(f*x)^m*(a + b*ArcCosh[c*x])^n,x]`

output `Integrate[(f*x)^m*(a + b*ArcCosh[c*x])^n, x]`

**3.454.3 Rubi [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6303}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (fx)^m (a + \operatorname{arccosh}(cx))^n dx$$

↓ 6303

$$\int (fx)^m (a + \operatorname{arccosh}(cx))^n dx$$

input `Int[(f*x)^m*(a + b*ArcCosh[c*x])^n,x]`

output `$Aborted`

**3.454.3.1 Defintions of rubi rules used**

rule 6303 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n]*((d_.)*(x_.))^m, x_Symbol]
:= Unintegrable[(d*x)^m*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, m, n}, x]`

**3.454.4 Maple [N/A] (verified)**

Not integrable

Time = 0.30 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int (fx)^m (a + b \operatorname{arccosh}(cx))^n dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^n,x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^n,x)`

**3.454.5 Fracas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx = \int (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n,x, algorithm="fricas")`output `integral((f*x)^m*(b*arccosh(c*x) + a)^n, x)`**3.454.6 Sympy [N/A]**

Not integrable

Time = 31.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx = \int (fx)^m (a + b \operatorname{acosh}(cx))^n dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))**n,x)`output `Integral((f*x)**m*(a + b*acosh(c*x))**n, x)`**3.454.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx = \int (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n, x)`

**3.454.8 Giac [F(-1)]**

Timed out.

$$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Timed out`

**3.454.9 Mupad [N/A]**

Not integrable

Time = 2.94 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int (fx)^m (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n (fx)^m dx$$

input `int((a + b*acosh(c*x))^n*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^n*(f*x)^m, x)`

**3.455**  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx$

3.455.1 Optimal result . . . . . 3380  
 3.455.2 Mathematica [N/A] . . . . . 3380  
 3.455.3 Rubi [N/A] . . . . . 3381  
 3.455.4 Maple [N/A] (verified) . . . . . 3381  
 3.455.5 Fricas [N/A] . . . . . 3382  
 3.455.6 Sympy [N/A] . . . . . 3382  
 3.455.7 Maxima [N/A] . . . . . 3382  
 3.455.8 Giac [N/A] . . . . . 3383  
 3.455.9 Mupad [N/A] . . . . . 3383

**3.455.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx = \operatorname{Int} \left( \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2}, x \right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d), x)`

**3.455.2 Mathematica [N/A]**

Not integrable

Time = 1.06 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx = \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2), x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2), x]`

**3.455.3 Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2),x]`

output `$Aborted`

**3.455.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((f_.)*(x_))^m_.*((d_) + (e_.)*(x_)^2)^p_., x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.455.4 Maple [N/A] (verified)**

Not integrable

Time = 0.55 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{-c^2 dx^2 + d} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d),x)`

**3.455.5 Fracas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx = \int -\frac{(fx)^m (b \operatorname{arccosh}(cx) + a)^n}{c^2 dx^2 - d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d),x, algorithm="fracas")`

output `integral(-(f*x)^m*(b*arccosh(c*x) + a)^n/(c^2*d*x^2 - d), x)`

**3.455.6 Sympy [N/A]**

Not integrable

Time = 79.71 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx = -\int \frac{(fx)^m (a + b \operatorname{acosh}(cx))^n}{c^2 x^2 - 1} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d),x)`

output `-Integral((f*x)**m*(a + b*acosh(c*x))**n/(c**2*x**2 - 1), x)/d`

**3.455.7 Maxima [N/A]**

Not integrable

Time = 0.55 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx = \int -\frac{(fx)^m (b \operatorname{arccosh}(cx) + a)^n}{c^2 dx^2 - d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d),x, algorithm="maxima")`

output `-integrate((f*x)^m*(b*arccosh(c*x) + a)^n/(c^2*d*x^2 - d), x)`

---

3.455.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx$

**3.455.8 Giac [N/A]**

Not integrable

Time = 13.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx = \int -\frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{c^2 dx^2 - d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d),x, algorithm="giac")`output `integrate(-(f*x)^m*(b*arccosh(c*x) + a)^n/(c^2*d*x^2 - d), x)`**3.455.9 Mupad [N/A]**

Not integrable

Time = 3.33 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{d - c^2 dx^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (fx)^m}{d - c^2 dx^2} dx$$

input `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2),x)`output `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2), x)`



**3.456** 
$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx$$

3.456.1 Optimal result . . . . .	3384
3.456.2 Mathematica [N/A] . . . . .	3384
3.456.3 Rubi [N/A] . . . . .	3385
3.456.4 Maple [N/A] (verified) . . . . .	3385
3.456.5 Fricas [N/A] . . . . .	3386
3.456.6 Sympy [F(-1)] . . . . .	3386
3.456.7 Maxima [N/A] . . . . .	3386
3.456.8 Giac [N/A] . . . . .	3387
3.456.9 Mupad [N/A] . . . . .	3387

**3.456.1 Optimal result**

Integrand size = 29, antiderivative size = 29

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx = \operatorname{Int}\left(\frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^2,x)`

**3.456.2 Mathematica [N/A]**

Not integrable

Time = 1.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx = \int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^2,x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^2, x]`

**3.456.3 Rubi [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^2,x]`

output `$Aborted`

**3.456.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_]*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.456.4 Maple [N/A] (verified)**

Not integrable

Time = 0.66 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(-c^2 dx^2 + d)^2} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^2,x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^2,x)`

**3.456.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.55

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{(c^2 dx^2 - d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^2,x, algorithm="fricas")`

output `integral((f*x)^m*(b*arccosh(c*x) + a)^n/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.456.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**2,x)`

output `Timed out`

**3.456.7 Maxima [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.10

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{(c^2 dx^2 - d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^2,x, algorithm="maxima")`

output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n/(c^2*d*x^2 - d)^2, x)`

---

3.456.  $\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^2} dx$

**3.456.8 Giac [N/A]**

Not integrable

Time = 12.33 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.10

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^2} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{(c^2 dx^2 - d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^2,x, algorithm="giac")`

output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n/(c^2*d*x^2 - d)^2, x)`

**3.456.9 Mupad [N/A]**

Not integrable

Time = 3.42 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.07

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (fx)^m}{(d - c^2 dx^2)^2} dx$$

input `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2)^2,x)`

output `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2)^2, x)`

### 3.457 $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^n dx$

3.457.1 Optimal result . . . . .	3388
3.457.2 Mathematica [N/A] . . . . .	3388
3.457.3 Rubi [N/A] . . . . .	3389
3.457.4 Maple [N/A] (verified) . . . . .	3389
3.457.5 Fricas [N/A] . . . . .	3390
3.457.6 Sympy [F(-1)] . . . . .	3390
3.457.7 Maxima [N/A] . . . . .	3390
3.457.8 Giac [F(-1)] . . . . .	3391
3.457.9 Mupad [N/A] . . . . .	3391

#### 3.457.1 Optimal result

Integrand size = 31, antiderivative size = 31

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^n dx = \text{Int}\left((fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^n, x\right)$$

output `Unintegrable((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

#### 3.457.2 Mathematica [N/A]

Not integrable

Time = 1.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^n dx = \int (fx)^m (d - c^2 dx^2)^{3/2} (a + \text{barccosh}(cx))^n dx$$

input `Integrate[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output `Integrate[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n, x]`

**3.457.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d - c^2 dx^2)^{3/2} (fx)^m (a + \operatorname{barccosh}(cx))^n dx$$

↓ 6375

$$\int (d - c^2 dx^2)^{3/2} (fx)^m (a + \operatorname{barccosh}(cx))^n dx$$

input `Int[(f*x)^m*(d - c^2*d*x^2)^(3/2)*(a + b*ArcCosh[c*x])^n,x]`

output `$Aborted`

**3.457.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.457.4 Maple [N/A] (verified)**

Not integrable

Time = 1.58 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int (fx)^m (-c^2 dx^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^n dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x)`

**3.457.5 Fracas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{3/2} (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="fricas")`

output `integral(-(c^2*d*x^2 - d)*sqrt(-c^2*d*x^2 + d)*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

**3.457.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(3/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.457.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int (-c^2 dx^2 + d)^{3/2} (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate((-c^2*d*x^2 + d)^(3/2)*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

---

3.457.  $\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx$

**3.457.8 Giac [F(-1)]**

Timed out.

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(3/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Timed out`

**3.457.9 Mupad [N/A]**

Not integrable

Time = 3.31 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m (d - c^2 dx^2)^{3/2} (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n (d - c^2 dx^2)^{3/2} (fx)^m dx$$

input `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(3/2)*(f*x)^m, x)`



### 3.458 $\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$

3.458.1 Optimal result . . . . .	3392
3.458.2 Mathematica [N/A] . . . . .	3392
3.458.3 Rubi [N/A] . . . . .	3393
3.458.4 Maple [N/A] (verified) . . . . .	3393
3.458.5 Fricas [N/A] . . . . .	3394
3.458.6 Sympy [F(-1)] . . . . .	3394
3.458.7 Maxima [N/A] . . . . .	3394
3.458.8 Giac [F(-1)] . . . . .	3395
3.458.9 Mupad [N/A] . . . . .	3395

#### 3.458.1 Optimal result

Integrand size = 31, antiderivative size = 31

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \operatorname{Int}\left((fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n, x\right)$$

output `Unintegrable((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^n,x)`

#### 3.458.2 Mathematica [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx$$

input `Integrate[(f*x)^m*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`

output `Integrate[(f*x)^m*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n, x]`

**3.458.3 Rubi [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d - c^2 dx^2} (fx)^m (a + \operatorname{barccosh}(cx))^n dx$$

↓ 6375

$$\int \sqrt{d - c^2 dx^2} (fx)^m (a + \operatorname{barccosh}(cx))^n dx$$

input `Int[(f*x)^m*Sqrt[d - c^2*d*x^2]*(a + b*ArcCosh[c*x])^n,x]`

output `$Aborted`

**3.458.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^n_.*((f_.)*(x_.))^m_.*((d_.) + (e_.)*(x_.)^2)^p_., x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.458.4 Maple [N/A] (verified)**

Not integrable

Time = 1.56 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int (fx)^m \sqrt{-c^2 dx^2 + d} (a + b \operatorname{arccosh}(cx))^n dx$$

input `int((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^n,x)`

output `int((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^n,x)`

**3.458.5 Fracas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int \sqrt{-c^2 dx^2 + d} (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^n,x, algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

**3.458.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((f*x)**m*(-c**2*d*x**2+d)**(1/2)*(a+b*acosh(c*x))**n,x)`

output `Timed out`

**3.458.7 Maxima [N/A]**

Not integrable

Time = 0.54 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int \sqrt{-c^2 dx^2 + d} (fx)^m (b \operatorname{arcosh}(cx) + a)^n dx$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^n,x, algorithm="maxima")`

output `integrate(sqrt(-c^2*d*x^2 + d)*(f*x)^m*(b*arccosh(c*x) + a)^n, x)`

**3.458.8 Giac [F(-1)]**

Timed out.

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \text{Timed out}$$

input `integrate((f*x)^m*(-c^2*d*x^2+d)^(1/2)*(a+b*arccosh(c*x))^n,x, algorithm="giac")`

output `Timed out`

**3.458.9 Mupad [N/A]**

Not integrable

Time = 3.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int (fx)^m \sqrt{d - c^2 dx^2} (a + \operatorname{barccosh}(cx))^n dx = \int (a + b \operatorname{acosh}(cx))^n \sqrt{d - c^2 dx^2} (fx)^m dx$$

input `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2)*(f*x)^m,x)`

output `int((a + b*acosh(c*x))^n*(d - c^2*d*x^2)^(1/2)*(f*x)^m, x)`

**3.459**  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$

3.459.1 Optimal result . . . . .	3396
3.459.2 Mathematica [N/A] . . . . .	3396
3.459.3 Rubi [N/A] . . . . .	3397
3.459.4 Maple [N/A] (verified) . . . . .	3397
3.459.5 Fricas [N/A] . . . . .	3398
3.459.6 Sympy [N/A] . . . . .	3398
3.459.7 Maxima [N/A] . . . . .	3398
3.459.8 Giac [N/A] . . . . .	3399
3.459.9 Mupad [N/A] . . . . .	3399

**3.459.1 Optimal result**

Integrand size = 31, antiderivative size = 31

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \operatorname{Int}\left(\frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2), x)`

**3.459.2 Mathematica [N/A]**

Not integrable

Time = 0.69 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2], x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2], x]`

**3.459.3 Rubi [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^n)/Sqrt[d - c^2*d*x^2],x]`

output `$Aborted`

**3.459.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n_.*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.459.4 Maple [N/A] (verified)**

Not integrable

Time = 1.60 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{\sqrt{-c^2 d x^2 + d}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x)`

**3.459.5 Fracas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.48

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + d}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="
fricas")
```

```
output integral(-sqrt(-c^2*d*x^2 + d)*(f*x)^m*(b*arccosh(c*x) + a)^n/(c^2*d*x^2 -
d), x)
```

**3.459.6 Sympy [N/A]**

Not integrable

Time = 92.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))^n}{\sqrt{-d(cx - 1)(cx + 1)}} dx$$

```
input integrate((f*x)**m*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(1/2),x)
```

```
output Integral((f*x)**m*(a + b*acosh(c*x))**n/sqrt(-d*(c*x - 1)*(c*x + 1)), x)
```

**3.459.7 Maxima [N/A]**

Not integrable

Time = 0.54 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + d}} dx$$

```
input integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="
maxima")
```

```
output integrate((f*x)^m*(b*arccosh(c*x) + a)^n/sqrt(-c^2*d*x^2 + d), x)
```

---

3.459.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx$

**3.459.8 Giac [N/A]**

Not integrable

Time = 13.10 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{\sqrt{-c^2 dx^2 + d}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(1/2),x, algorithm="giac")`

output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n/sqrt(-c^2*d*x^2 + d), x)`

**3.459.9 Mupad [N/A]**

Not integrable

Time = 3.50 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{\sqrt{d - c^2 dx^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (fx)^m}{\sqrt{d - c^2 dx^2}} dx$$

input `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2)^(1/2),x)`

output `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2)^(1/2), x)`



**3.460** 
$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

3.460.1 Optimal result . . . . .	3400
3.460.2 Mathematica [N/A] . . . . .	3400
3.460.3 Rubi [N/A] . . . . .	3401
3.460.4 Maple [N/A] (verified) . . . . .	3401
3.460.5 Fricas [N/A] . . . . .	3402
3.460.6 Sympy [F(-1)] . . . . .	3402
3.460.7 Maxima [N/A] . . . . .	3402
3.460.8 Giac [N/A] . . . . .	3403
3.460.9 Mupad [N/A] . . . . .	3403

**3.460.1 Optimal result**

Integrand size = 31, antiderivative size = 31

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \operatorname{Int} \left( \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}}, x \right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

**3.460.2 Mathematica [N/A]**

Not integrable

Time = 1.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.06

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2),x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2), x]`

**3.460.3 Rubi [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x])^n)/(d - c^2*d*x^2)^(3/2),x]`

output `$Aborted`

**3.460.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n]*((f_.)*(x_))^m*((d_) + (e_.)*(x_)^2)^p, x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.460.4 Maple [N/A] (verified)**

Not integrable

Time = 2.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(-c^2 dx^2 + d)^{3/2}} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x)`

---

3.460.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$

**3.460.5 Fracas [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.87

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="fracas")`

output `integral(sqrt(-c^2*d*x^2 + d)*(f*x)^m*(b*arccosh(c*x) + a)^n/(c^4*d^2*x^4 - 2*c^2*d^2*x^2 + d^2), x)`

**3.460.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))**n/(-c**2*d*x**2+d)**(3/2),x)`

output `Timed out`

**3.460.7 Maxima [N/A]**

Not integrable

Time = 0.56 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="maxima")`

output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n/(-c^2*d*x^2 + d)^(3/2), x)`

---

3.460.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx$

**3.460.8 Giac [N/A]**

Not integrable

Time = 12.81 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(fx)^m (b \operatorname{arcosh}(cx) + a)^n}{(-c^2 dx^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))^n/(-c^2*d*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((f*x)^m*(b*arccosh(c*x) + a)^n/(-c^2*d*x^2 + d)^(3/2), x)`

**3.460.9 Mupad [N/A]**

Not integrable

Time = 3.53 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))^n}{(d - c^2 dx^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^n (fx)^m}{(d - c^2 dx^2)^{3/2}} dx$$

input `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2)^(3/2),x)`

output `int(((a + b*acosh(c*x))^n*(f*x)^m)/(d - c^2*d*x^2)^(3/2), x)`

### 3.461 $\int x^4(d + ex^2) (a + \operatorname{barccosh}(cx)) dx$

3.461.1 Optimal result . . . . .	3404
3.461.2 Mathematica [A] (verified) . . . . .	3405
3.461.3 Rubi [A] (verified) . . . . .	3405
3.461.4 Maple [A] (verified) . . . . .	3408
3.461.5 Fricas [A] (verification not implemented) . . . . .	3408
3.461.6 Sympy [F] . . . . .	3409
3.461.7 Maxima [A] (verification not implemented) . . . . .	3409
3.461.8 Giac [F(-2)] . . . . .	3409
3.461.9 Mupad [F(-1)] . . . . .	3410

#### 3.461.1 Optimal result

Integrand size = 19, antiderivative size = 177

$$\int x^4(d + ex^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{8b(49c^2d + 30e) \sqrt{-1 + cx}\sqrt{1 + cx}}{3675c^7} - \frac{4b(49c^2d + 30e) x^2 \sqrt{-1 + cx}\sqrt{1 + cx}}{3675c^5} - \frac{b(49c^2d + 30e) x^4 \sqrt{-1 + cx}\sqrt{1 + cx}}{1225c^3} - \frac{be x^6 \sqrt{-1 + cx}\sqrt{1 + cx}}{49c} + \frac{1}{5} dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7} ex^7(a + \operatorname{barccosh}(cx))$$

```
output 1/5*d*x^5*(a+b*arccosh(c*x))+1/7*e*x^7*(a+b*arccosh(c*x))-8/3675*b*(49*c^2*d+30*e)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^7-4/3675*b*(49*c^2*d+30*e)*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5-1/1225*b*(49*c^2*d+30*e)*x^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-1/49*b*e*x^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c
```

**3.461.2 Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.69

$$\int x^4(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \frac{1}{35}ax^5(7d + 5ex^2) - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}(240e + 8c^2(49d + 15ex^2) + 2c^4(98dx^2 + 45ex^4) + 3c^6(49dx^4 + 25ex^6))}{3675c^7} + \frac{1}{35}bx^5(7d + 5ex^2) \operatorname{arccosh}(cx)$$

input `Integrate[x^4*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`output `(a*x^5*(7*d + 5*e*x^2))/35 - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(240*e + 8*c^2*(49*d + 15*e*x^2) + 2*c^4*(98*d*x^2 + 45*e*x^4) + 3*c^6*(49*d*x^4 + 25*e*x^6)))/(3675*c^7) + (b*x^5*(7*d + 5*e*x^2)*ArcCosh[c*x])/35`**3.461.3 Rubi [A] (verified)**Time = 0.36 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {6371, 960, 111, 27, 111, 27, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int x^4(d + ex^2)(a + \operatorname{barccosh}(cx)) dx \\ & \quad \downarrow \text{6371} \\ & -\frac{1}{35}bc \int \frac{x^5(5ex^2 + 7d)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}ex^7(a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{960} \\ & -\frac{1}{35}bc \left( \frac{1}{7} \left( \frac{30e}{c^2} + 49d \right) \int \frac{x^5}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{5ex^6\sqrt{cx - 1}\sqrt{cx + 1}}{7c^2} \right) + \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}ex^7(a + \operatorname{barccosh}(cx)) \\ & \quad \downarrow \text{111} \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{35}bc \left( \frac{1}{7} \left( \frac{30e}{c^2} + 49d \right) \left( \frac{\int \frac{4x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) + \frac{5ex^6\sqrt{cx-1}\sqrt{cx+1}}{7c^2} \right) + \\
& \quad \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}ex^7(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow 27 \\
& -\frac{1}{35}bc \left( \frac{1}{7} \left( \frac{30e}{c^2} + 49d \right) \left( \frac{4 \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) + \frac{5ex^6\sqrt{cx-1}\sqrt{cx+1}}{7c^2} \right) + \\
& \quad \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}ex^7(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow 111 \\
& -\frac{1}{35}bc \left( \frac{1}{7} \left( \frac{30e}{c^2} + 49d \right) \left( \frac{4 \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) + \frac{5ex^6\sqrt{cx-1}\sqrt{cx+1}}{7c^2} \right) + \\
& \quad \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}ex^7(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow 27 \\
& -\frac{1}{35}bc \left( \frac{1}{7} \left( \frac{30e}{c^2} + 49d \right) \left( \frac{4 \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} + \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) + \frac{5ex^6\sqrt{cx-1}\sqrt{cx+1}}{7c^2} \right) + \\
& \quad \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}ex^7(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow 83 \\
& \quad \frac{1}{5}dx^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}ex^7(a + \operatorname{barccosh}(cx)) - \\
& \quad \frac{1}{35}bc \left( \frac{5ex^6\sqrt{cx-1}\sqrt{cx+1}}{7c^2} + \frac{1}{7} \left( \frac{x^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{4 \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)}{5c^2} \right) \left( \frac{30e}{c^2} + 49d \right) \right)
\end{aligned}$$

input `Int[x^4*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`

output `-1/35*(b*c*((5*e*x^6*sqrt[-1 + c*x]*sqrt[1 + c*x])/(7*c^2) + ((49*d + (30*e)/c^2)*(x^4*sqrt[-1 + c*x]*sqrt[1 + c*x])/(5*c^2) + (4*((2*sqrt[-1 + c*x])*sqrt[1 + c*x])/(3*c^4) + (x^2*sqrt[-1 + c*x]*sqrt[1 + c*x])/(3*c^2))))/(5*c^2)))/7) + (d*x^5*(a + b*ArcCosh[c*x])/5 + (e*x^7*(a + b*ArcCosh[c*x]))/7`

## 3.461.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`
- rule 111 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 960 `Int[((e_.)*(x_))^(m_.)*((a1_.) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_.) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_.) + (d_.)*(x_)^(n_.)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`
- rule 6371 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[d*(f*x)^(m + 1)*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (Simp[e*(f*x)^(m + 3)*((a + b*ArcCosh[c*x])/(f^3*(m + 3))), x] - Simp[b*(c/(f*(m + 1)*(m + 3))) Int[(f*x)^(m + 1)*((d*(m + 3) + e*(m + 1)*x^2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x)) /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && NeQ[m, -1] && NeQ[m, -3]`



### 3.461.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.69

method	result
parts	$a\left(\frac{1}{7}e x^7 + \frac{1}{5}d x^5\right) + \frac{b\left(\frac{c^5 \operatorname{arccosh}(cx)e x^7}{7} + \frac{\operatorname{arccosh}(cx)c^5 x^5 d}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(75c^6 e x^6 + 147c^6 d x^4 + 90c^4 e x^4 + 196c^4 d x^2 + 120c^2 e x^2 + 392c^2 d + 240e)}{3675c^2}\right)}{c^5}$
derivativedivides	$\frac{a\left(\frac{1}{5}d c^7 x^5 + \frac{1}{7}e c^7 x^7\right)}{c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)d c^7 x^5}{5} + \frac{\operatorname{arccosh}(cx)e c^7 x^7}{7} - \frac{\sqrt{cx-1}\sqrt{cx+1}(75c^6 e x^6 + 147c^6 d x^4 + 90c^4 e x^4 + 196c^4 d x^2 + 120c^2 e x^2 + 392c^2 d + 240e)}{3675}\right)}{c^5}$
default	$\frac{a\left(\frac{1}{5}d c^7 x^5 + \frac{1}{7}e c^7 x^7\right)}{c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)d c^7 x^5}{5} + \frac{\operatorname{arccosh}(cx)e c^7 x^7}{7} - \frac{\sqrt{cx-1}\sqrt{cx+1}(75c^6 e x^6 + 147c^6 d x^4 + 90c^4 e x^4 + 196c^4 d x^2 + 120c^2 e x^2 + 392c^2 d + 240e)}{3675}\right)}{c^5}$

input `int(x^4*(e*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/7*e*x^7+1/5*d*x^5)+b/c^5*(1/7*c^5*arccosh(c*x)*e*x^7+1/5*arccosh(c*x)*c^5*x^5*d-1/3675/c^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(75*c^6*e*x^6+147*c^6*d*x^4+90*c^4*e*x^4+196*c^4*d*x^2+120*c^2*e*x^2+392*c^2*d+240*e))`

### 3.461.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.79

$$\int x^4(d + ex^2)(a + b\operatorname{arccosh}(cx)) dx = \frac{525 ac^7 ex^7 + 735 ac^7 dx^5 + 105(5 bc^7 ex^7 + 7 bc^7 dx^5) \log(cx + \sqrt{c^2 x^2 - 1}) - (75 bc^6 ex^6 + 3(49 bc^6 d + 30 b^2 c^4 e)x^4 + 392 b^2 c^2 d + 4(49 b^2 c^4 d + 30 b^2 c^2 e)x^2 + 240 b^2 e) \sqrt{c^2 x^2 - 1}}{3675 c^7}$$

input `integrate(x^4*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `1/3675*(525*a*c^7*e*x^7 + 735*a*c^7*d*x^5 + 105*(5*b*c^7*e*x^7 + 7*b*c^7*d*x^5)*log(c*x + sqrt(c^2*x^2 - 1)) - (75*b*c^6*e*x^6 + 3*(49*b*c^6*d + 30*b*c^4*e)*x^4 + 392*b*c^2*d + 4*(49*b*c^4*d + 30*b*c^2*e)*x^2 + 240*b*e)*sqrt(c^2*x^2 - 1)/c^7`

**3.461.6 Sympy [F]**

$$\int x^4(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \int x^4(a + b \operatorname{acosh}(cx))(d + ex^2) dx$$

input `integrate(x**4*(e*x**2+d)*(a+b*acosh(c*x)),x)`

output `Integral(x**4*(a + b*acosh(c*x))*(d + e*x**2), x)`

**3.461.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.01

$$\begin{aligned} \int x^4(d + ex^2)(a + \operatorname{barccosh}(cx)) dx &= \frac{1}{7} aex^7 + \frac{1}{5} adx^5 \\ &+ \frac{1}{75} \left( 15x^5 \operatorname{arcosh}(cx) - \left( \frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6} \right) c \right) bd \\ &+ \frac{1}{245} \left( 35x^7 \operatorname{arcosh}(cx) - \left( \frac{5\sqrt{c^2x^2 - 1}x^6}{c^2} + \frac{6\sqrt{c^2x^2 - 1}x^4}{c^4} + \frac{8\sqrt{c^2x^2 - 1}x^2}{c^6} + \frac{16\sqrt{c^2x^2 - 1}}{c^8} \right) c \right) be \end{aligned}$$

input `integrate(x^4*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/7*a*e*x^7 + 1/5*a*d*x^5 + 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d + 1/245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*e`

**3.461.8 Giac [F(-2)]**

Exception generated.

$$\int x^4(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

### 3.461.9 Mupad [F(-1)]

Timed out.

$$\int x^4(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \int x^4(a + b \operatorname{acosh}(cx))(ex^2 + d) dx$$

input `int(x^4*(a + b*acosh(c*x))*(d + e*x^2),x)`

output `int(x^4*(a + b*acosh(c*x))*(d + e*x^2), x)`

### 3.462 $\int x^3(d + ex^2)(a + \operatorname{barccosh}(cx)) dx$

3.462.1 Optimal result . . . . .	3411
3.462.2 Mathematica [A] (warning: unable to verify) . . . . .	3412
3.462.3 Rubi [A] (verified) . . . . .	3412
3.462.4 Maple [A] (verified) . . . . .	3415
3.462.5 Fricas [A] (verification not implemented) . . . . .	3415
3.462.6 Sympy [F] . . . . .	3416
3.462.7 Maxima [A] (verification not implemented) . . . . .	3416
3.462.8 Giac [F(-2)] . . . . .	3417
3.462.9 Mupad [F(-1)] . . . . .	3417

#### 3.462.1 Optimal result

Integrand size = 19, antiderivative size = 161

$$\int x^3(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = -\frac{b(9c^2d + 5e)x\sqrt{-1 + cx}\sqrt{1 + cx}}{96c^5} - \frac{b(9c^2d + 5e)x^3\sqrt{-1 + cx}\sqrt{1 + cx}}{144c^3} - \frac{be x^5\sqrt{-1 + cx}\sqrt{1 + cx}}{36c} - \frac{b(9c^2d + 5e)\operatorname{arccosh}(cx)}{96c^6} + \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx))$$

```
output -1/96*b*(9*c^2*d+5*e)*arccosh(c*x)/c^6+1/4*d*x^4*(a+b*arccosh(c*x))+1/6*e*x^6*(a+b*arccosh(c*x))-1/96*b*(9*c^2*d+5*e)*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5-1/144*b*(9*c^2*d+5*e)*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-1/36*b*e*x^5*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c
```

**3.462.2 Mathematica [A] (warning: unable to verify)**

Time = 0.13 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.87

$$\int x^3(d + ex^2)(a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{24ac^6x^4(3d + 2ex^2) - bcx\sqrt{-1 + cx}\sqrt{1 + cx}(15e + c^2(27d + 10ex^2) + 2c^4(9dx^2 + 4ex^4)) + 24bc^6x^4(3d + 2ex^2)\operatorname{ArcCosh}[cx] - 6b^2(9c^2d + 5e)\operatorname{ArcTanh}\left[\sqrt{\frac{-1 + cx}{1 + cx}}\right]}{288c^6}$$

input `Integrate[x^3*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`output `(24*a*c^6*x^4*(3*d + 2*e*x^2) - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(15*e + c^2*(27*d + 10*e*x^2) + 2*c^4*(9*d*x^2 + 4*e*x^4)) + 24*b*c^6*x^4*(3*d + 2*e*x^2)*ArcCosh[c*x] - 6*b*(9*c^2*d + 5*e)*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]])/(288*c^6)`**3.462.3 Rubi [A] (verified)**Time = 0.35 (sec) , antiderivative size = 157, normalized size of antiderivative = 0.98, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {6371, 27, 960, 111, 27, 101, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(d + ex^2)(a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6371$$

$$-\frac{1}{24}bc \int \frac{2x^4(2ex^2 + 3d)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx))$$

$$\downarrow 27$$

$$-\frac{1}{12}bc \int \frac{x^4(2ex^2 + 3d)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx))$$

$$\downarrow 960$$

$$-\frac{1}{12}bc \left( \frac{1}{3} \left( \frac{5e}{c^2} + 9d \right) \int \frac{x^4}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{ex^5\sqrt{cx - 1}\sqrt{cx + 1}}{3c^2} \right) + \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx))$$

$$\begin{aligned}
& \downarrow 111 \\
& -\frac{1}{12}bc \left( \frac{1}{3} \left( \frac{5e}{c^2} + 9d \right) \left( \frac{\int \frac{3x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) + \frac{ex^5\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) + \\
& \quad \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx)) \\
& \downarrow 27 \\
& -\frac{1}{12}bc \left( \frac{1}{3} \left( \frac{5e}{c^2} + 9d \right) \left( \frac{3 \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) + \frac{ex^5\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) + \\
& \quad \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx)) \\
& \downarrow 101 \\
& -\frac{1}{12}bc \left( \frac{1}{3} \left( \frac{5e}{c^2} + 9d \right) \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) + \frac{ex^5\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) + \\
& \quad \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx)) \\
& \downarrow 43 \\
& \quad \frac{1}{4}dx^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}ex^6(a + \operatorname{barccosh}(cx)) - \\
& \frac{1}{12}bc \left( \frac{1}{3} \left( \frac{3 \left( \frac{\operatorname{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) \left( \frac{5e}{c^2} + 9d \right) + \frac{ex^5\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right)
\end{aligned}$$

input `Int[x^3*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`

output `(d*x^4*(a + b*ArcCosh[c*x]))/4 + (e*x^6*(a + b*ArcCosh[c*x]))/6 - (b*c*((e*x^5*sqrt[-1 + c*x]*sqrt[1 + c*x])/(3*c^2) + ((9*d + (5*e)/c^2)*((x^3*sqrt[-1 + c*x]*sqrt[1 + c*x])/(4*c^2) + (3*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3)))/(4*c^2))/3))/12`

## 3.462.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 43 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*Sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`
- rule 101 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`
- rule 111 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`
- rule 960 `Int[((e_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

```
rule 6371 Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[d*(f*x)^(m + 1)*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (Simp[e*(f*x)^(m + 3)*((a + b*ArcCosh[c*x])/(f^3*(m + 3))), x] - Simp[b*(c/(f*(m + 1)*(m + 3))) Int[(f*x)^(m + 1)*((d*(m + 3) + e*(m + 1)*x^2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && NeQ[m, -1] && NeQ[m, -3]
```

### 3.462.4 Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 215, normalized size of antiderivative = 1.34

method	result
parts	$a\left(\frac{1}{6}ex^6 + \frac{1}{4}dx^4\right) + \frac{b\left(\frac{c^4 \operatorname{arccosh}(cx)ex^6}{6} + \frac{\operatorname{arccosh}(cx)c^4x^4d}{4} - \sqrt{cx-1}\sqrt{cx+1}\left(18c^5d\sqrt{c^2x^2-1}x^3 + 8e\sqrt{c^2x^2-1}c^5x^5 + 27d^3c^3\right)\right)}{c^2}$
derivativedivides	$\frac{a\left(\frac{1}{4}c^6dx^4 + \frac{1}{8}c^6ex^6\right)}{c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)d c^6x^4}{4} + \frac{\operatorname{arccosh}(cx)ec^6x^6}{6} - \sqrt{cx-1}\sqrt{cx+1}\left(18c^5d\sqrt{c^2x^2-1}x^3 + 8e\sqrt{c^2x^2-1}c^5x^5 + 27d^3c^3\right)\right)}{c^4}$
default	$\frac{a\left(\frac{1}{4}c^6dx^4 + \frac{1}{8}c^6ex^6\right)}{c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)d c^6x^4}{4} + \frac{\operatorname{arccosh}(cx)ec^6x^6}{6} - \sqrt{cx-1}\sqrt{cx+1}\left(18c^5d\sqrt{c^2x^2-1}x^3 + 8e\sqrt{c^2x^2-1}c^5x^5 + 27d^3c^3\right)\right)}{c^4}$

```
input int(x^3*(e*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(1/6*e*x^6+1/4*d*x^4)+b/c^4*(1/6*c^4*arccosh(c*x)*e*x^6+1/4*arccosh(c*x)*c^4*x^4*d-1/288/c^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(18*c^5*d*(c^2*x^2-1)^(1/2)*x^3+8*e*(c^2*x^2-1)^(1/2)*c^5*x^5+27*d*c^3*x*(c^2*x^2-1)^(1/2)+10*e*c^3*x^3*(c^2*x^2-1)^(1/2)+27*d*c^2*ln(c*x+(c^2*x^2-1)^(1/2))+15*e*c*x*(c^2*x^2-1)^(1/2)+15*e*ln(c*x+(c^2*x^2-1)^(1/2)))/(c^2*x^2-1)^(1/2)
```

### 3.462.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.84

$$\int x^3(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \frac{48ac^6ex^6 + 72ac^6dx^4 + 3(16bc^6ex^6 + 24bc^6dx^4 - 9bc^2d - 5be) \log(cx + \sqrt{c^2x^2 - 1}) - (8bc^5ex^5 + 2(9d + 3e))c^4}{288c^6}$$

```
input integrate(x^3*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")
```



output  $\frac{1}{288}(48ac^6e^6x^6 + 72ac^6d^2x^4 + 3(16b^2c^6e^6x^6 + 24b^2c^6d^2x^4 - 9b^2c^6d - 5b^2e))\log(cx + \sqrt{c^2x^2 - 1}) - (8b^2c^5e^6x^5 + 2(9b^2c^5d + 5b^2c^3e))x^3 + 3(9b^2c^3d + 5b^2c^3e)x)\sqrt{c^2x^2 - 1})/c^6$

### 3.462.6 Sympy [F]

$$\int x^3(d + ex^2)(a + b \operatorname{arccosh}(cx)) dx = \int x^3(a + b \operatorname{acosh}(cx))(d + ex^2) dx$$

input `integrate(x**3*(e*x**2+d)*(a+b*acosh(c*x)),x)`

output `Integral(x**3*(a + b*acosh(c*x))*(d + e*x**2), x)`

### 3.462.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.22

$$\begin{aligned} \int x^3(d + ex^2)(a + b \operatorname{arccosh}(cx)) dx &= \frac{1}{6} aex^6 + \frac{1}{4} adx^4 \\ &+ \frac{1}{32} \left( 8x^4 \operatorname{arccosh}(cx) - \left( \frac{2\sqrt{c^2x^2 - 1}x^3}{c^2} + \frac{3\sqrt{c^2x^2 - 1}x}{c^4} + \frac{3 \log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^5} \right) c \right) bd \\ &+ \frac{1}{288} \left( 48x^6 \operatorname{arccosh}(cx) - \left( \frac{8\sqrt{c^2x^2 - 1}x^5}{c^2} + \frac{10\sqrt{c^2x^2 - 1}x^3}{c^4} + \frac{15\sqrt{c^2x^2 - 1}x}{c^6} + \frac{15 \log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^7} \right) c \right) b^2e \end{aligned}$$

input `integrate(x^3*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output  $\frac{1}{6}a^2e^6x^6 + \frac{1}{4}a^2d^2x^4 + \frac{1}{32}(8x^4 \operatorname{arccosh}(cx) - (2\sqrt{c^2x^2 - 1})x^3/c^2 + 3\sqrt{c^2x^2 - 1}x/c^4 + 3\log(2c^2x + 2\sqrt{c^2x^2 - 1})c)/c^5)c)b^2d + \frac{1}{288}(48x^6 \operatorname{arccosh}(cx) - (8\sqrt{c^2x^2 - 1})x^5/c^2 + 10\sqrt{c^2x^2 - 1}x^3/c^4 + 15\sqrt{c^2x^2 - 1}x/c^6 + 15\log(2c^2x + 2\sqrt{c^2x^2 - 1})c)/c^7)c)b^2e$

**3.462.8 Giac [F(-2)]**

Exception generated.

$$\int x^3(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:IN  
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const  
index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.462.9 Mupad [F(-1)]**

Timed out.

$$\int x^3(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \int x^3(a + b \operatorname{acosh}(cx))(ex^2 + d) dx$$

input `int(x^3*(a + b*acosh(c*x))*(d + e*x^2),x)`

output `int(x^3*(a + b*acosh(c*x))*(d + e*x^2), x)`

### 3.463 $\int x^2(d + ex^2) (a + \operatorname{barccosh}(cx)) dx$

3.463.1 Optimal result . . . . .	3418
3.463.2 Mathematica [A] (verified) . . . . .	3418
3.463.3 Rubi [A] (verified) . . . . .	3419
3.463.4 Maple [A] (verified) . . . . .	3421
3.463.5 Fricas [A] (verification not implemented) . . . . .	3421
3.463.6 Sympy [F] . . . . .	3422
3.463.7 Maxima [A] (verification not implemented) . . . . .	3422
3.463.8 Giac [F(-2)] . . . . .	3423
3.463.9 Mupad [F(-1)] . . . . .	3423

#### 3.463.1 Optimal result

Integrand size = 19, antiderivative size = 138

$$\int x^2(d + ex^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{2b(25c^2d + 12e) \sqrt{-1 + cx}\sqrt{1 + cx}}{225c^5} - \frac{b(25c^2d + 12e) x^2 \sqrt{-1 + cx}\sqrt{1 + cx}}{225c^3} - \frac{bex^4 \sqrt{-1 + cx}\sqrt{1 + cx}}{25c} + \frac{1}{3}dx^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}ex^5(a + \operatorname{barccosh}(cx))$$

```
output 1/3*d*x^3*(a+b*arccosh(c*x))+1/5*e*x^5*(a+b*arccosh(c*x))-2/225*b*(25*c^2*d+12*e)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5-1/225*b*(25*c^2*d+12*e)*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-1/25*b*e*x^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c
```

#### 3.463.2 Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.73

$$\int x^2(d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \frac{1}{225} \left( 15ax^3(5d + 3ex^2) - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}(24e + 2c^2(25d + 6ex^2) + c^4(25dx^2 + 9ex^4))}{c^5} + 15bx^3(5d + 3ex^2) \operatorname{arccosh}(cx) \right)$$

input `Integrate[x^2*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`

output  $(15*a*x^3*(5*d + 3*e*x^2) - (b*\sqrt{-1 + c*x})*\sqrt{1 + c*x}*(24*e + 2*c^2*(25*d + 6*e*x^2) + c^4*(25*d*x^2 + 9*e*x^4)))/c^5 + 15*b*x^3*(5*d + 3*e*x^2)*\text{ArcCosh}[c*x])/225$

### 3.463.3 Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6371, 960, 111, 27, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^2 (d + ex^2) (a + \text{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6371} \\
 & -\frac{1}{15}bc \int \frac{x^3(3ex^2 + 5d)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{3}dx^3(a + \text{barccosh}(cx)) + \frac{1}{5}ex^5(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{960} \\
 & -\frac{1}{15}bc \left( \frac{1}{5} \left( \frac{12e}{c^2} + 25d \right) \int \frac{x^3}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{3ex^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) + \frac{1}{3}dx^3(a + \text{barccosh}(cx)) + \frac{1}{5}ex^5(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{111} \\
 & -\frac{1}{15}bc \left( \frac{1}{5} \left( \frac{12e}{c^2} + 25d \right) \left( \frac{\int \frac{2x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) + \frac{3ex^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) + \frac{1}{3}dx^3(a + \text{barccosh}(cx)) + \frac{1}{5}ex^5(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{15}bc \left( \frac{1}{5} \left( \frac{12e}{c^2} + 25d \right) \left( \frac{2 \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}} dx}{3c^2} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) + \frac{3ex^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} \right) + \frac{1}{3}dx^3(a + \text{barccosh}(cx)) + \frac{1}{5}ex^5(a + \text{barccosh}(cx)) \\
 & \quad \downarrow \text{83}
 \end{aligned}$$

$$\frac{1}{15}bc \left( \frac{3ex^4\sqrt{cx-1}\sqrt{cx+1}}{5c^2} + \frac{1}{5} \left( \frac{2\sqrt{cx-1}\sqrt{cx+1}}{3c^4} + \frac{x^2\sqrt{cx-1}\sqrt{cx+1}}{3c^2} \right) \left( \frac{12e}{c^2} + 25d \right) \right) - \frac{1}{3}dx^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}ex^5(a + \operatorname{barccosh}(cx)) -$$

input `Int[x^2*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`

output `-1/15*(b*c*((3*e*x^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(5*c^2) + ((25*d + (12*e)/c^2)*((2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^4) + (x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2))))/5) + (d*x^3*(a + b*ArcCosh[c*x]))/3 + (e*x^5*(a + b*ArcCosh[c*x]))/5`

### 3.463.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 83 `Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]`

rule 111 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Simp[1/(d*f*(m + n + p + 1)) Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`

rule 960 `Int[((e_.)*(x_))^(m_.)*((a1_.) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_.) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

```
rule 6371 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2), x_Symbol] :> Simp[d*(f*x)^(m + 1)*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (Simp[e*(f*x)^(m + 3)*((a + b*ArcCosh[c*x])/(f^3*(m + 3))), x] - Simp[b*(c/(f*(m + 1)*(m + 3))) Int[(f*x)^(m + 1)*((d*(m + 3) + e*(m + 1)*x^2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && NeQ[m, -1] && NeQ[m, -3]
```

### 3.463.4 Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.76

method	result
parts	$a\left(\frac{1}{5}ex^5 + \frac{1}{3}dx^3\right) + \frac{b\left(\frac{c^3 \operatorname{arccosh}(cx)ex^5}{5} + \frac{\operatorname{arccosh}(cx)c^3x^3d}{3} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4ex^4+25c^4dx^2+12c^2ex^2+50c^2d+24e)}{225c^2}\right)}{c^3}$
derivativedivides	$\frac{a\left(\frac{1}{3}dc^5x^3 + \frac{1}{5}ec^5x^5\right)}{c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)dc^5x^3}{3} + \frac{\operatorname{arccosh}(cx)ec^5x^5}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4ex^4+25c^4dx^2+12c^2ex^2+50c^2d+24e)}{225}\right)}{c^3}$
default	$\frac{a\left(\frac{1}{3}dc^5x^3 + \frac{1}{5}ec^5x^5\right)}{c^2} + \frac{b\left(\frac{\operatorname{arccosh}(cx)dc^5x^3}{3} + \frac{\operatorname{arccosh}(cx)ec^5x^5}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4ex^4+25c^4dx^2+12c^2ex^2+50c^2d+24e)}{225}\right)}{c^3}$

```
input int(x^2*(e*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

```
output a*(1/5*e*x^5+1/3*d*x^3)+b/c^3*(1/5*c^3*arccosh(c*x)*e*x^5+1/3*arccosh(c*x)*c^3*x^3*d-1/225/c^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(9*c^4*e*x^4+25*c^4*d*x^2+12*c^2*e*x^2+50*c^2*d+24*e))
```

### 3.463.5 Fracas [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.86

$$\int x^2(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \frac{45ac^5ex^5 + 75ac^5dx^3 + 15(3bc^5ex^5 + 5bc^5dx^3) \log(cx + \sqrt{c^2x^2 - 1}) - (9bc^4ex^4 + 50bc^2d + (25bc^4d + 225c^5))}{225c^5}$$

```
input integrate(x^2*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")
```

output  $1/225*(45*a*c^5*e*x^5 + 75*a*c^5*d*x^3 + 15*(3*b*c^5*e*x^5 + 5*b*c^5*d*x^3)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (9*b*c^4*e*x^4 + 50*b*c^2*d + (25*b*c^4*d + 12*b*c^2*e)*x^2 + 24*b*e)*\sqrt{c^2*x^2 - 1})/c^5$

### 3.463.6 Sympy [F]

$$\int x^2(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \int x^2(a + b \operatorname{acosh}(cx))(d + ex^2) dx$$

input `integrate(x**2*(e*x**2+d)*(a+b*acosh(c*x)),x)`

output `Integral(x**2*(a + b*acosh(c*x))*(d + e*x**2), x)`

### 3.463.7 Maxima [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.01

$$\begin{aligned} & \int x^2(d + ex^2)(a + \operatorname{barccosh}(cx)) dx \\ &= \frac{1}{5} aex^5 + \frac{1}{3} adx^3 + \frac{1}{9} \left( 3x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \right) bd \\ &+ \frac{1}{75} \left( 15x^5 \operatorname{arcosh}(cx) - \left( \frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6} \right) c \right) be \end{aligned}$$

input `integrate(x^2*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output  $1/5*a*e*x^5 + 1/3*a*d*x^3 + 1/9*(3*x^3*\operatorname{arccosh}(c*x) - c*(\sqrt{c^2*x^2 - 1}*x^2/c^2 + 2*\sqrt{c^2*x^2 - 1}/c^4))*b*d + 1/75*(15*x^5*\operatorname{arccosh}(c*x) - (3*\sqrt{c^2*x^2 - 1}*x^4/c^2 + 4*\sqrt{c^2*x^2 - 1}*x^2/c^4 + 8*\sqrt{c^2*x^2 - 1}/c^6)*c)*b*e$

**3.463.8 Giac [F(-2)]**

Exception generated.

$$\int x^2(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^2*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

**3.463.9 Mupad [F(-1)]**

Timed out.

$$\int x^2(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \int x^2(a + b \operatorname{acosh}(cx))(ex^2 + d) dx$$

input `int(x^2*(a + b*acosh(c*x))*(d + e*x^2),x)`

output `int(x^2*(a + b*acosh(c*x))*(d + e*x^2), x)`



### 3.464 $\int x(d + ex^2) (a + \operatorname{barccosh}(cx)) dx$

3.464.1 Optimal result . . . . .	3424
3.464.2 Mathematica [A] (warning: unable to verify) . . . . .	3424
3.464.3 Rubi [A] (verified) . . . . .	3425
3.464.4 Maple [B] (verified) . . . . .	3427
3.464.5 Fricas [A] (verification not implemented) . . . . .	3427
3.464.6 Sympy [F] . . . . .	3428
3.464.7 Maxima [A] (verification not implemented) . . . . .	3428
3.464.8 Giac [F(-2)] . . . . .	3428
3.464.9 Mupad [F(-1)] . . . . .	3429

#### 3.464.1 Optimal result

Integrand size = 17, antiderivative size = 122

$$\int x(d + ex^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{b(8c^2d + 3e) x\sqrt{-1 + cx}\sqrt{1 + cx}}{32c^3} - \frac{bex^3\sqrt{-1 + cx}\sqrt{1 + cx}}{16c} - \frac{b(8c^2d + 3e) \operatorname{arccosh}(cx)}{32c^4} + \frac{1}{2}dx^2(a + \operatorname{barccosh}(cx)) + \frac{1}{4}ex^4(a + \operatorname{barccosh}(cx))$$

```
output -1/32*b*(8*c^2*d+3*e)*arccosh(c*x)/c^4+1/2*d*x^2*(a+b*arccosh(c*x))+1/4*e*x^4*(a+b*arccosh(c*x))-1/32*b*(8*c^2*d+3*e)*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-1/16*b*e*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c
```

#### 3.464.2 Mathematica [A] (warning: unable to verify)

Time = 0.09 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.98

$$\int x(d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \frac{cx(8ac^3x(2d + ex^2) - b\sqrt{-1 + cx}\sqrt{1 + cx}(3e + 2c^2(4d + ex^2))) + 8bc^4x^2(2d + ex^2) \operatorname{arccosh}(cx) - 2b(8c^2d + 3e)x\sqrt{-1 + cx}\sqrt{1 + cx}}{32c^4}$$

```
input Integrate[x*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]
```

output  $(c*x*(8*a*c^3*x*(2*d + e*x^2) - b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(3*e + 2*c^2*(4*d + e*x^2))) + 8*b*c^4*x^2*(2*d + e*x^2)*\text{ArcCosh}[c*x] - 2*b*(8*c^2*d + 3*e)*\text{ArcTanh}[\text{Sqrt}[(-1 + c*x)/(1 + c*x)])]/(32*c^4)$

### 3.464.3 Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.99, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$ , Rules used = {6371, 27, 960, 101, 43}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d + ex^2)(a + \text{barccosh}(cx)) dx$$

$$\downarrow 6371$$

$$-\frac{1}{8}bc \int \frac{2x^2(ex^2 + 2d)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{2}dx^2(a + \text{barccosh}(cx)) + \frac{1}{4}ex^4(a + \text{barccosh}(cx))$$

$$\downarrow 27$$

$$-\frac{1}{4}bc \int \frac{x^2(ex^2 + 2d)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{1}{2}dx^2(a + \text{barccosh}(cx)) + \frac{1}{4}ex^4(a + \text{barccosh}(cx))$$

$$\downarrow 960$$

$$-\frac{1}{4}bc \left( \frac{1}{4} \left( \frac{3e}{c^2} + 8d \right) \int \frac{x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{ex^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) + \frac{1}{2}dx^2(a + \text{barccosh}(cx)) + \frac{1}{4}ex^4(a + \text{barccosh}(cx))$$

$$\downarrow 101$$

$$-\frac{1}{4}bc \left( \frac{1}{4} \left( \frac{3e}{c^2} + 8d \right) \left( \frac{\int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}} dx}{2c^2} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) + \frac{ex^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right) + \frac{1}{2}dx^2(a + \text{barccosh}(cx)) + \frac{1}{4}ex^4(a + \text{barccosh}(cx))$$

$$\downarrow 43$$

$$\frac{1}{2}dx^2(a + \text{barccosh}(cx)) + \frac{1}{4}ex^4(a + \text{barccosh}(cx)) - \frac{1}{4}bc \left( \frac{1}{4} \left( \frac{\text{arccosh}(cx)}{2c^3} + \frac{x\sqrt{cx-1}\sqrt{cx+1}}{2c^2} \right) \left( \frac{3e}{c^2} + 8d \right) + \frac{ex^3\sqrt{cx-1}\sqrt{cx+1}}{4c^2} \right)$$

input `Int[x*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`

output `(d*x^2*(a + b*ArcCosh[c*x]))/2 + (e*x^4*(a + b*ArcCosh[c*x]))/4 - (b*c*((e*x^3*sqrt[-1 + c*x]*sqrt[1 + c*x])/(4*c^2) + ((8*d + (3*e)/c^2)*((x*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*c^2) + ArcCosh[c*x]/(2*c^3))))/4)`

### 3.464.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 43 `Int[1/(sqrt[(a_) + (b_)*(x_)]*sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[ArcCosh[b*(x/a)]/(b*sqrt[d/b]), x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && GtQ[a, 0] && GtQ[d/b, 0]`

rule 101 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Simp[1/(d*f*(n + p + 3)) Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]`

rule 960 `Int[((e_)*(x_))^(m_)*((a1_) + (b1_)*(x_))^(non2_))^(p_)*((a2_) + (b2_)*(x_))^(non2_))^(p_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6371 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2), x_Symbol] := Simp[d*(f*x)^(m + 1)*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (Simp[e*(f*x)^(m + 3)*((a + b*ArcCosh[c*x])/(f^3*(m + 3))), x] - Simp[b*(c/(f*(m + 1)*(m + 3))) Int[(f*x)^(m + 1)*((d*(m + 3) + e*(m + 1)*x^2)/(sqrt[1 + c*x]*sqrt[-1 + c*x])], x], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && NeQ[m, -1] && NeQ[m, -3]`

### 3.464.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 224 vs. 2(104) = 208.

Time = 0.34 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.84

method	result
parts	$\frac{a(e^2x^2+d)^2}{4e} + \frac{b \left( \frac{c^2 e \operatorname{arccosh}(cx)x^4}{4} + \frac{\operatorname{arccosh}(cx)c^2x^2d}{2} + \frac{c^2 \operatorname{arccosh}(cx)d^2}{4e} - \frac{\sqrt{cx-1}\sqrt{cx+1} (8c^4d^2 \ln(cx+\sqrt{c^2x^2-1}) + 8dc^3ex^3 + 8c^2d^2x^2 + 8cd^3) \operatorname{arccosh}(cx)}{4} \right)}{c^2}$
derivativedivides	$\frac{a(c^2ex^2+c^2d)^2}{4c^2e} + \frac{b \left( \frac{\operatorname{arccosh}(cx)c^4d^2}{4e} + \frac{\operatorname{arccosh}(cx)c^4dx^2}{2} + \frac{e \operatorname{arccosh}(cx)c^4x^4}{4} - \frac{\sqrt{cx-1}\sqrt{cx+1} (8c^4d^2 \ln(cx+\sqrt{c^2x^2-1}) + 8dc^3ex^3 + 8c^2d^2x^2 + 8cd^3) \operatorname{arccosh}(cx)}{4} \right)}{c^2}$
default	$\frac{a(c^2ex^2+c^2d)^2}{4c^2e} + \frac{b \left( \frac{\operatorname{arccosh}(cx)c^4d^2}{4e} + \frac{\operatorname{arccosh}(cx)c^4dx^2}{2} + \frac{e \operatorname{arccosh}(cx)c^4x^4}{4} - \frac{\sqrt{cx-1}\sqrt{cx+1} (8c^4d^2 \ln(cx+\sqrt{c^2x^2-1}) + 8dc^3ex^3 + 8c^2d^2x^2 + 8cd^3) \operatorname{arccosh}(cx)}{4} \right)}{c^2}$

input `int(x*(e*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output  $\frac{1}{4}a*(e*x^2+d)^2/e+b/c^2*(1/4*c^2*e*arccosh(c*x)*x^4+1/2*arccosh(c*x)*c^2*x^2*d+1/4*c^2/e*arccosh(c*x)*d^2-1/32/c^2/e*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(8*c^4*d^2*\ln(c*x+(c^2*x^2-1)^{(1/2)})+8*d*c^3*e*x*(c^2*x^2-1)^{(1/2)}+2*e^2*(c^2*x^2-1)^{(1/2)}*c^3*x^3+8*d*c^2*e*\ln(c*x+(c^2*x^2-1)^{(1/2)})+3*e^2*c*x*(c^2*x^2-1)^{(1/2)}+3*e^2*\ln(c*x+(c^2*x^2-1)^{(1/2)}))/((c^2*x^2-1)^{(1/2)})$

### 3.464.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.93

$$\int x(d+ex^2)(a+b\operatorname{arccosh}(cx))dx = \frac{8ac^4ex^4 + 16ac^4dx^2 + (8bc^4ex^4 + 16bc^4dx^2 - 8bc^2d - 3be) \log(cx + \sqrt{c^2x^2 - 1}) - (2bc^3ex^3 + (8bc^3d + 3b^2c^2e)x)\sqrt{c^2x^2 - 1}}{32c^4}$$

input `integrate(x*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $\frac{1}{32}*(8*a*c^4*e*x^4 + 16*a*c^4*d*x^2 + (8*b*c^4*e*x^4 + 16*b*c^4*d*x^2 - 8*b*c^2*d - 3*b*e)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (2*b*c^3*e*x^3 + (8*b*c^3*d + 3*b*c^2*e)*x)*\sqrt{c^2*x^2 - 1})/c^4$

**3.464.6 Sympy [F]**

$$\int x(d + ex^2) (a + b \operatorname{arccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (d + ex^2) dx$$

input `integrate(x*(e*x**2+d)*(a+b*acosh(c*x)),x)`

output `Integral(x*(a + b*acosh(c*x))*(d + e*x**2), x)`

**3.464.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.28

$$\begin{aligned} \int x(d + ex^2) (a + b \operatorname{arccosh}(cx)) dx &= \frac{1}{4} aex^4 + \frac{1}{2} adx^2 \\ &+ \frac{1}{4} \left( 2x^2 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x}{c^2} + \frac{\log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^3} \right) \right) bd \\ &+ \frac{1}{32} \left( 8x^4 \operatorname{arccosh}(cx) - \left( \frac{2\sqrt{c^2x^2 - 1}x^3}{c^2} + \frac{3\sqrt{c^2x^2 - 1}x}{c^4} + \frac{3\log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^5} \right) c \right) be \end{aligned}$$

input `integrate(x*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/4*a*e*x^4 + 1/2*a*d*x^2 + 1/4*(2*x^2*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x/c^2 + log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^3))*b*d + 1/32*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*e`

**3.464.8 Giac [F(-2)]**

Exception generated.

$$\int x(d + ex^2) (a + b \operatorname{arccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const ve  
cteur & l) Error: Bad Argument Value

### 3.464.9 Mupad [F(-1)]

Timed out.

$$\int x(d + ex^2)(a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx))(ex^2 + d) dx$$

input `int(x*(a + b*acosh(c*x))*(d + e*x^2),x)`

output `int(x*(a + b*acosh(c*x))*(d + e*x^2), x)`

### 3.465 $\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx$

3.465.1 Optimal result . . . . .	3430
3.465.2 Mathematica [A] (verified) . . . . .	3430
3.465.3 Rubi [A] (verified) . . . . .	3431
3.465.4 Maple [A] (verified) . . . . .	3432
3.465.5 Fricas [A] (verification not implemented) . . . . .	3433
3.465.6 Sympy [F] . . . . .	3433
3.465.7 Maxima [A] (verification not implemented) . . . . .	3434
3.465.8 Giac [F(-2)] . . . . .	3434
3.465.9 Mupad [F(-1)] . . . . .	3434

#### 3.465.1 Optimal result

Integrand size = 16, antiderivative size = 94

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = -\frac{b(9c^2d + 2e) \sqrt{-1 + cx} \sqrt{1 + cx}}{9c^3} - \frac{bex^2 \sqrt{-1 + cx} \sqrt{1 + cx}}{9c} + dx(a + \operatorname{barccosh}(cx)) + \frac{1}{3}ex^3(a + \operatorname{barccosh}(cx))$$

output `d*x*(a+b*arccosh(c*x))+1/3*e*x^3*(a+b*arccosh(c*x))-1/9*b*(9*c^2*d+2*e)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-1/9*b*e*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c`

#### 3.465.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.81

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \frac{1}{9} \left( 3ax(3d + ex^2) - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}(2e + c^2(9d + ex^2))}{c^3} + 3bx(3d + ex^2) \operatorname{arccosh}(cx) \right)$$

input `Integrate[(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`

output  $(3*a*x*(3*d + e*x^2) - (b*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]*(2*e + c^2*(9*d + e*x^2)))/c^3 + 3*b*x*(3*d + e*x^2)*\text{ArcCosh}[c*x])/9$

### 3.465.3 Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.05, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {6323, 27, 960, 83}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)(a + \text{barccosh}(cx)) dx$$

$$\downarrow 6323$$

$$-bc \int \frac{x(ex^2 + 3d)}{3\sqrt{cx - 1}\sqrt{cx + 1}} dx + dx(a + \text{barccosh}(cx)) + \frac{1}{3}ex^3(a + \text{barccosh}(cx))$$

$$\downarrow 27$$

$$-\frac{1}{3}bc \int \frac{x(ex^2 + 3d)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + dx(a + \text{barccosh}(cx)) + \frac{1}{3}ex^3(a + \text{barccosh}(cx))$$

$$\downarrow 960$$

$$-\frac{1}{3}bc \left( \frac{1}{3} \left( \frac{2e}{c^2} + 9d \right) \int \frac{x}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{ex^2\sqrt{cx - 1}\sqrt{cx + 1}}{3c^2} \right) + dx(a + \text{barccosh}(cx)) + \frac{1}{3}ex^3(a + \text{barccosh}(cx))$$

$$\downarrow 83$$

$$dx(a + \text{barccosh}(cx)) + \frac{1}{3}ex^3(a + \text{barccosh}(cx)) - \frac{1}{3}bc \left( \frac{\sqrt{cx - 1}\sqrt{cx + 1} \left( \frac{2e}{c^2} + 9d \right)}{3c^2} + \frac{ex^2\sqrt{cx - 1}\sqrt{cx + 1}}{3c^2} \right)$$

input  $\text{Int}[(d + e*x^2)*(a + b*\text{ArcCosh}[c*x]), x]$

output  $-1/3*(b*c*((9*d + (2*e)/c^2)*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/(3*c^2) + (e*x^2*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/(3*c^2)) + d*x*(a + b*\text{ArcCosh}[c*x]) + (e*x^3*(a + b*\text{ArcCosh}[c*x]))/3$



3.465.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 83 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && EqQ[a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)), 0]
```

```
rule 960 Int[((e_)*(x_))^(m_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*(a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1)), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]
```

```
rule 6323 Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d + e, 0] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])
```

3.465.4 Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.81

method	result	size
parts	$a\left(\frac{1}{3}x^3e + dx\right) + \frac{b\left(\frac{c \operatorname{arccosh}(cx)x^3e + \operatorname{arccosh}(cx)dcx - \frac{\sqrt{cx-1}\sqrt{cx+1}(c^2ex^2+9c^2d+2e)}{9c^2}}{c}\right)}{c}$	76
derivativedivides	$\frac{a\left(dc^3x + \frac{1}{3}ec^3x^3\right)}{c^2} + \frac{b\left(\operatorname{arccosh}(cx)dc^3x + \frac{\operatorname{arccosh}(cx)ec^3x^3 - \frac{\sqrt{cx-1}\sqrt{cx+1}(c^2ex^2+9c^2d+2e)}{9}}{c}\right)}{c^2}$	90
default	$\frac{a\left(dc^3x + \frac{1}{3}ec^3x^3\right)}{c^2} + \frac{b\left(\operatorname{arccosh}(cx)dc^3x + \frac{\operatorname{arccosh}(cx)ec^3x^3 - \frac{\sqrt{cx-1}\sqrt{cx+1}(c^2ex^2+9c^2d+2e)}{9}}{c}\right)}{c^2}$	90

3.465.  $\int (d + ex^2)(a + \operatorname{arccosh}(cx)) dx$

input `int((e*x^2+d)*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/3*x^3*e+d*x)+b/c*(1/3*c*arccosh(c*x)*x^3*e+arccosh(c*x)*d*c*x-1/9/c^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(c^2*e*x^2+9*c^2*d+2*e))`

### 3.465.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{3ac^3ex^3 + 9ac^3dx + 3(bc^3ex^3 + 3bc^3dx) \log(cx + \sqrt{c^2x^2 - 1}) - (bc^2ex^2 + 9bc^2d + 2be)\sqrt{c^2x^2 - 1}}{9c^3}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `1/9*(3*a*c^3*e*x^3 + 9*a*c^3*d*x + 3*(b*c^3*e*x^3 + 3*b*c^3*d*x)*log(c*x + sqrt(c^2*x^2 - 1)) - (b*c^2*e*x^2 + 9*b*c^2*d + 2*b*e)*sqrt(c^2*x^2 - 1))/c^3`

### 3.465.6 Sympy [F]

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d + ex^2) dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x)),x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2), x)`

**3.465.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.97

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{1}{3} aex^3 + \frac{1}{9} \left( 3x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \right) be$$

$$+ adx + \frac{(cx \operatorname{arcosh}(cx) - \sqrt{c^2x^2 - 1})bd}{c}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`output `1/3*a*e*x^3 + 1/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*e + a*d*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*d/c`**3.465.8 Giac [F(-2)]**

Exception generated.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`output `Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`**3.465.9 Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (ex^2 + d) dx$$

input `int((a + b*acosh(c*x))*(d + e*x^2),x)`output `int((a + b*acosh(c*x))*(d + e*x^2), x)`

**3.466**  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x} dx$

3.466.1 Optimal result . . . . .	3435
3.466.2 Mathematica [A] (verified) . . . . .	3436
3.466.3 Rubi [A] (verified) . . . . .	3436
3.466.4 Maple [A] (verified) . . . . .	3438
3.466.5 Fricas [F] . . . . .	3438
3.466.6 Sympy [F] . . . . .	3438
3.466.7 Maxima [F] . . . . .	3439
3.466.8 Giac [F] . . . . .	3439
3.466.9 Mupad [F(-1)] . . . . .	3439

**3.466.1 Optimal result**

Integrand size = 19, antiderivative size = 264

$$\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x} dx = -\frac{bex\sqrt{-1+cx}\sqrt{1+cx}}{4c} - \frac{b\operatorname{arccosh}(cx)}{4c^2} + \frac{1}{2}ex^2(a+b\operatorname{arccosh}(cx)) - \frac{ibd\sqrt{1-c^2x^2}\arcsin(cx)^2}{2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{bd\sqrt{1-c^2x^2}\arcsin(cx)\log(1-e^{2i\arcsin(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} + d(a+b\operatorname{arccosh}(cx))\log(x) - \frac{bd\sqrt{1-c^2x^2}\arcsin(cx)\log(x)}{\sqrt{-1+cx}\sqrt{1+cx}} - \frac{ibd\sqrt{1-c^2x^2}\operatorname{PolyLog}(2, e^{2i\arcsin(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}}$$

```
output -1/4*b*e*arccosh(c*x)/c^2+1/2*e*x^2*(a+b*arccosh(c*x))+d*(a+b*arccosh(c*x)
)*ln(x)-1/4*b*e*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-1/2*I*b*d*arcsin(c*x)^2*(-
c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*d*arcsin(c*x)*ln(1-(I*c*x+(
-c^2*x^2+1)^(1/2))^2*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*d*a
rcsin(c*x)*ln(x)*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/2*I*b*d*
polylog(2,(I*c*x+(-c^2*x^2+1)^(1/2))^2*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(
c*x+1)^(1/2))
```

**3.466.2 Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.49

$$\int \frac{(d + ex^2)(a + \operatorname{arccosh}(cx))}{x} dx = \frac{1}{2} aex^2 - \frac{bex\sqrt{-1+cx}\sqrt{1+cx}}{4c} + \frac{1}{2} bex^2 \operatorname{arccosh}(cx) - \frac{bearctanh\left(\frac{\sqrt{-1+cx}}{\sqrt{1+cx}}\right)}{2c^2} + ad \log(x) + \frac{1}{2} bd(\operatorname{arccosh}(cx) (\operatorname{arccosh}(cx) + 2 \log(1 + e^{-2\operatorname{arccosh}(cx)})) - \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)}))$$

input `Integrate[((d + e*x^2)*(a + b*ArcCosh[c*x]))/x,x]`output `(a*e*x^2)/2 - (b*e*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(4*c) + (b*e*x^2*ArcCosh[c*x])/2 - (b*e*ArcTanh[Sqrt[-1 + c*x]/Sqrt[1 + c*x]])/(2*c^2) + a*d*Log[x] + (b*d*(ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLog[2, -E^(-2*ArcCosh[c*x])]))/2`**3.466.3 Rubi [A] (verified)**Time = 0.92 (sec) , antiderivative size = 274, normalized size of antiderivative = 1.04, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)(a + \operatorname{arccosh}(cx))}{x} dx$$

$$\downarrow 6373$$

$$-bc \int \frac{ex^2 + 2d \log(x)}{2\sqrt{cx} - 1\sqrt{cx} + 1} dx + d \log(x)(a + \operatorname{arccosh}(cx)) + \frac{1}{2} ex^2(a + \operatorname{arccosh}(cx))$$

$$\downarrow 27$$

$$-\frac{1}{2} bc \int \frac{ex^2 + 2d \log(x)}{\sqrt{cx} - 1\sqrt{cx} + 1} dx + d \log(x)(a + \operatorname{arccosh}(cx)) + \frac{1}{2} ex^2(a + \operatorname{arccosh}(cx))$$

$$\downarrow 7293$$

---

3.466.  $\int \frac{(d+ex^2)(a+\operatorname{arccosh}(cx))}{x} dx$

$$-\frac{1}{2}bc \int \left( \frac{ex^2}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{2d \log(x)}{\sqrt{cx-1}\sqrt{cx+1}} \right) dx + d \log(x)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}ex^2(a + \operatorname{barccosh}(cx))$$

↓ 2009

$$\frac{1}{2}bc \left( \frac{\operatorname{earccosh}(cx)}{2c^3} + \frac{id\sqrt{1-c^2x^2} \operatorname{PolyLog}(2, e^{2i \arcsin(cx)})}{c\sqrt{cx-1}\sqrt{cx+1}} + \frac{id\sqrt{1-c^2x^2} \arcsin(cx)^2}{c\sqrt{cx-1}\sqrt{cx+1}} - \frac{2d\sqrt{1-c^2x^2} \arcsin(cx)}{c\sqrt{cx-1}\sqrt{cx+1}} \right)$$

input `Int[((d + e*x^2)*(a + b*ArcCosh[c*x]))/x,x]`

output `(e*x^2*(a + b*ArcCosh[c*x])/2 + d*(a + b*ArcCosh[c*x])*Log[x] - (b*c*((e*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) + (e*ArcCosh[c*x])/(2*c^3) + (I*d*Sqrt[1 - c^2*x^2]*ArcSin[c*x]^2)/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (2*d*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[1 - E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (2*d*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[x])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (I*d*Sqrt[1 - c^2*x^2]*PolyLog[2, E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/2`

### 3.466.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

**3.466.4 Maple [A] (verified)**

Time = 1.24 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.48

method	result
parts	$\frac{ae x^2}{2} + ad \ln(x) - \frac{db \operatorname{arccosh}(cx)^2}{2} + \frac{b \operatorname{arccosh}(cx) e x^2}{2} - \frac{be x \sqrt{cx-1} \sqrt{cx+1}}{4c} - \frac{be \operatorname{arccosh}(cx)}{4c^2} + db \operatorname{arccosh}(cx)$
derivativedivides	$\frac{ae x^2}{2} + ad \ln(cx) - \frac{db \operatorname{arccosh}(cx)^2}{2} - \frac{be x \sqrt{cx-1} \sqrt{cx+1}}{4c} + \frac{b \operatorname{arccosh}(cx) e x^2}{2} - \frac{be \operatorname{arccosh}(cx)}{4c^2} + db \operatorname{arccosh}(cx)$
default	$\frac{ae x^2}{2} + ad \ln(cx) - \frac{db \operatorname{arccosh}(cx)^2}{2} - \frac{be x \sqrt{cx-1} \sqrt{cx+1}}{4c} + \frac{b \operatorname{arccosh}(cx) e x^2}{2} - \frac{be \operatorname{arccosh}(cx)}{4c^2} + db \operatorname{arccosh}(cx)$

input `int((e*x^2+d)*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`output  $\frac{1}{2} a e x^2 + a d \ln(x) - \frac{1}{2} d b \operatorname{arccosh}(c x)^2 + \frac{1}{2} b \operatorname{arccosh}(c x) e x^2 - \frac{1}{4} b e x (c x - 1)^{1/2} (c x + 1)^{1/2} / c - \frac{1}{4} b e \operatorname{arccosh}(c x) / c^2 + d b \operatorname{arccosh}(c x) \ln(1 + (c x + (c x - 1)^{1/2} (c x + 1)^{1/2})^2) + \frac{1}{2} d b \operatorname{polylog}(2, -(c x + (c x - 1)^{1/2} (c x + 1)^{1/2})^2)$ **3.466.5 Fracas [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)(b \operatorname{arccosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x,x, algorithm="fracas")`output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arccosh(c*x))/x, x)`**3.466.6 Sympy [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d + ex^2)}{x} dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x))/x,x)`output `Integral((a + b*acosh(c*x))*(d + e*x**2)/x, x)`

---

3.466.  $\int \frac{(d+ex^2)(a+b \operatorname{arccosh}(cx))}{x} dx$

**3.466.7 Maxima [F]**

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `1/2*a*e*x^2 + a*d*log(x) + integrate(b*e*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) + b*d*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/x, x)`

**3.466.8 Giac [F]**

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arccosh(c*x) + a)/x, x)`

**3.466.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx))(ex^2 + d)}{x} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2))/x,x)`

output `int(((a + b*acosh(c*x))*(d + e*x^2))/x, x)`



**3.467**  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.467.1 Optimal result . . . . .	3440
3.467.2 Mathematica [A] (verified) . . . . .	3440
3.467.3 Rubi [A] (verified) . . . . .	3441
3.467.4 Maple [A] (verified) . . . . .	3442
3.467.5 Fricas [A] (verification not implemented) . . . . .	3443
3.467.6 Sympy [F] . . . . .	3444
3.467.7 Maxima [A] (verification not implemented) . . . . .	3444
3.467.8 Giac [F] . . . . .	3444
3.467.9 Mupad [F(-1)] . . . . .	3445

**3.467.1 Optimal result**

Integrand size = 19, antiderivative size = 75

$$\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^2} dx = -\frac{be\sqrt{-1+cx}\sqrt{1+cx}}{c} - \frac{d(a+b\operatorname{arccosh}(cx))}{x} + ex(a+b\operatorname{arccosh}(cx)) + bcd \arctan\left(\sqrt{-1+cx}\sqrt{1+cx}\right)$$

output `-d*(a+b*arccosh(c*x))/x+e*x*(a+b*arccosh(c*x))+b*c*d*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))-b*e*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c`

**3.467.2 Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.40

$$\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^2} dx = -\frac{ad}{x} + aex - \frac{be\sqrt{-1+cx}\sqrt{1+cx}}{c} - \frac{bd\operatorname{arccosh}(cx)}{x} + bex\operatorname{arccosh}(cx) + \frac{bcd\sqrt{-1+c^2x^2} \arctan\left(\sqrt{-1+c^2x^2}\right)}{\sqrt{-1+cx}\sqrt{1+cx}}$$

input `Integrate[((d + e*x^2)*(a + b*ArcCosh[c*x]))/x^2,x]`

output  $-\left(\frac{a*d}{x} + a*e*x - \frac{b*e*\sqrt{-1 + c*x}*\sqrt{1 + c*x}}{c} - \frac{b*d*\text{ArcCosh}[c*x]}{x} + b*e*x*\text{ArcCosh}[c*x] + \frac{b*c*d*\sqrt{-1 + c^2*x^2}*\text{ArcTan}[\sqrt{-1 + c*x}*\sqrt{1 + c*x}]}{\sqrt{-1 + c*x}*\sqrt{1 + c*x}}\right)$

### 3.467.3 Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.01, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {6371, 960, 103, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)(a + \text{barccosh}(cx))}{x^2} dx$$

↓ 6371

$$bc \int \frac{d - ex^2}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{d(a + \text{barccosh}(cx))}{x} + ex(a + \text{barccosh}(cx))$$

↓ 960

$$bc \left( d \int \frac{1}{x\sqrt{cx - 1}\sqrt{cx + 1}} dx - \frac{e\sqrt{cx - 1}\sqrt{cx + 1}}{c^2} \right) - \frac{d(a + \text{barccosh}(cx))}{x} + ex(a + \text{barccosh}(cx))$$

↓ 103

$$bc \left( cd \int \frac{1}{(cx - 1)(cx + 1)c + c} d(\sqrt{cx - 1}\sqrt{cx + 1}) - \frac{e\sqrt{cx - 1}\sqrt{cx + 1}}{c^2} \right) - \frac{d(a + \text{barccosh}(cx))}{x} + ex(a + \text{barccosh}(cx))$$

↓ 218

$$-\frac{d(a + \text{barccosh}(cx))}{x} + ex(a + \text{barccosh}(cx)) + bc \left( d \arctan(\sqrt{cx - 1}\sqrt{cx + 1}) - \frac{e\sqrt{cx - 1}\sqrt{cx + 1}}{c^2} \right)$$

input  $\text{Int}[\left(\frac{d + e*x^2}{x^2}\right)*(a + b*\text{ArcCosh}[c*x]),x]$

output  $-\left(\frac{d*(a + b*\text{ArcCosh}[c*x])}{x} + e*x*(a + b*\text{ArcCosh}[c*x]) + b*c*\left(-\left(\frac{e*\sqrt{-1 + c*x}*\sqrt{1 + c*x}}{c^2}\right) + d*\text{ArcTan}[\sqrt{-1 + c*x}*\sqrt{1 + c*x}]\right)\right)$

---

3.467.  $\int \frac{(d+ex^2)(a+\text{barccosh}(cx))}{x^2} dx$

## 3.467.3.1 Defintions of rubi rules used

rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 960 `Int[((e_.)*(x_)^(m_.))*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*(a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1)), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6371 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[d*(f*x)^(m + 1)*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (Simp[e*(f*x)^(m + 3)*((a + b*ArcCosh[c*x])/(f^3*(m + 3))), x] - Simp[b*(c/(f*(m + 1)*(m + 3))) Int[(f*x)^(m + 1)*((d*(m + 3) + e*(m + 1)*x^2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && NeQ[m, -1] && NeQ[m, -3]`

## 3.467.4 Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.36

---

3.467.  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^2} dx$

method	result
parts	$a\left(ex - \frac{d}{x}\right) + bc\left(\frac{\operatorname{arccosh}(cx)ex}{c} - \frac{\operatorname{arccosh}(cx)d}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}\left(d c^2 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) + e\sqrt{c^2x^2-1}\right)}{c^2\sqrt{c^2x^2-1}}\right)$
derivativedivides	$c\left(\frac{a\left(ecx - \frac{dc}{x}\right)}{c^2} + \frac{b\left(\operatorname{arccosh}(cx)ecx - \frac{\operatorname{arccosh}(cx)dc}{x} + \frac{\left(-d c^2 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) - e\sqrt{c^2x^2-1}\right)\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{c^2x^2-1}}\right)}{c^2}\right)$
default	$c\left(\frac{a\left(ecx - \frac{dc}{x}\right)}{c^2} + \frac{b\left(\operatorname{arccosh}(cx)ecx - \frac{\operatorname{arccosh}(cx)dc}{x} + \frac{\left(-d c^2 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) - e\sqrt{c^2x^2-1}\right)\sqrt{cx-1}\sqrt{cx+1}}{\sqrt{c^2x^2-1}}\right)}{c^2}\right)$

input `int((e*x^2+d)*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)`

output `a*(e*x-d/x)+b*c*(1/c*arccosh(c*x)*e*x-arccosh(c*x)*d/c/x-1/c^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(d*c^2*arctan(1/(c^2*x^2-1)^(1/2))+e*(c^2*x^2-1)^(1/2))/(c^2*x^2-1)^(1/2))`

### 3.467.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.76

$$\int \frac{(d + ex^2)(a + b\operatorname{arccosh}(cx))}{x^2} dx$$

$$= \frac{2bc^2dx \arctan(-cx + \sqrt{c^2x^2 - 1}) + acex^2 - \sqrt{c^2x^2 - 1}bex - acd + (bcd - bce)x \log(-cx + \sqrt{c^2x^2 - 1})}{cx}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^2,x, algorithm="fracas")`

output `(2*b*c^2*d*x*arctan(-c*x + sqrt(c^2*x^2 - 1)) + a*c*e*x^2 - sqrt(c^2*x^2 - 1)*b*e*x - a*c*d + (b*c*d - b*c*e)*x*log(-c*x + sqrt(c^2*x^2 - 1)) + (b*c*e*x^2 - b*c*d + (b*c*d - b*c*e)*x)*log(c*x + sqrt(c^2*x^2 - 1)))/(c*x)`

**3.467.6 Sympy [F]**

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d + ex^2)}{x^2} dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x))/x**2,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)/x**2, x)`

**3.467.7 Maxima [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.84

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^2} dx = -\left(c \arcsin\left(\frac{1}{c|x|}\right) + \frac{\operatorname{arcosh}(cx)}{x}\right)bd + aex$$

$$+ \frac{(cx \operatorname{arcosh}(cx) - \sqrt{c^2x^2 - 1})be}{c} - \frac{ad}{x}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output `-(c*arcsin(1/(c*abs(x)))) + arccosh(c*x)/x)*b*d + a*e*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*e/c - a*d/x`

**3.467.8 Giac [F]**

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arccosh(c*x) + a)/x^2, x)`

**3.467.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))(ex^2 + d)}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2))/x^2,x)`output `int(((a + b*acosh(c*x))*(d + e*x^2))/x^2, x)`

**3.468**  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^3} dx$

3.468.1 Optimal result . . . . .	3446
3.468.2 Mathematica [A] (verified) . . . . .	3447
3.468.3 Rubi [A] (verified) . . . . .	3447
3.468.4 Maple [A] (verified) . . . . .	3449
3.468.5 Fricas [F] . . . . .	3449
3.468.6 Sympy [F] . . . . .	3450
3.468.7 Maxima [F] . . . . .	3450
3.468.8 Giac [F] . . . . .	3450
3.468.9 Mupad [F(-1)] . . . . .	3451

**3.468.1 Optimal result**

Integrand size = 19, antiderivative size = 251

$$\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^3} dx = \frac{bcd\sqrt{-1+cx}\sqrt{1+cx}}{2x} - \frac{d(a+b\operatorname{arccosh}(cx))}{2x^2} - \frac{ibe\sqrt{1-c^2x^2}\arcsin(cx)^2}{2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{be\sqrt{1-c^2x^2}\arcsin(cx)\log(1-e^{2i\arcsin(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} + e(a+b\operatorname{arccosh}(cx))\log(x) - \frac{be\sqrt{1-c^2x^2}\arcsin(cx)\log(x)}{\sqrt{-1+cx}\sqrt{1+cx}} - \frac{ibe\sqrt{1-c^2x^2}\operatorname{PolyLog}(2, e^{2i\arcsin(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}}$$

output  $-1/2*d*(a+b*\operatorname{arccosh}(c*x))/x^2+e*(a+b*\operatorname{arccosh}(c*x))*\ln(x)+1/2*b*c*d*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/x-1/2*I*b*e*\arcsin(c*x)^2*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+b*e*\arcsin(c*x)*\ln(1-(I*c*x+(-c^2*x^2+1)^{(1/2)})^2*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b*e*\arcsin(c*x)*\ln(x)*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/2*I*b*e*\operatorname{polylog}(2,(I*c*x+(-c^2*x^2+1)^{(1/2)})^2*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)})$

**3.468.2 Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.40

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^3} dx = -\frac{ad}{2x^2} + \frac{bcd\sqrt{-1+cx}\sqrt{1+cx}}{2x} - \frac{bd\operatorname{barccosh}(cx)}{2x^2} \\ + ae \log(x) + \frac{1}{2}be(\operatorname{arccosh}(cx)(\operatorname{arccosh}(cx)) \\ + 2\log(1 + e^{-2\operatorname{arccosh}(cx)})) \\ - \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})$$

input `Integrate[((d + e*x^2)*(a + b*ArcCosh[c*x]))/x^3,x]`output `-1/2*(a*d)/x^2 + (b*c*d*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*x) - (b*d*ArcCosh[c*x])/(2*x^2) + a*e*Log[x] + (b*e*(ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLog[2, -E^(-2*ArcCosh[c*x])]))/2`**3.468.3 Rubi [A] (verified)**Time = 0.88 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^3} dx \\ \downarrow \text{6373} \\ -bc \int -\frac{\frac{d}{x^2} - 2e \log(x)}{2\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d(a + \operatorname{barccosh}(cx))}{2x^2} + e \log(x)(a + \operatorname{barccosh}(cx)) \\ \downarrow \text{27} \\ \frac{1}{2}bc \int \frac{\frac{d}{x^2} - 2e \log(x)}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d(a + \operatorname{barccosh}(cx))}{2x^2} + e \log(x)(a + \operatorname{barccosh}(cx)) \\ \downarrow \text{7293} \\ \frac{1}{2}bc \int \left( \frac{d}{x^2\sqrt{cx-1}\sqrt{cx+1}} - \frac{2e \log(x)}{\sqrt{cx-1}\sqrt{cx+1}} \right) dx - \frac{d(a + \operatorname{barccosh}(cx))}{2x^2} + e \log(x)(a + \operatorname{barccosh}(cx))$$

---

3.468.  $\int \frac{(d+ex^2)(a+\operatorname{barccosh}(cx))}{x^3} dx$



$$\begin{aligned} & \downarrow 2009 \\ & -\frac{d(a + \operatorname{barccosh}(cx))}{2x^2} + e \log(x)(a + \operatorname{barccosh}(cx)) + \\ & \frac{1}{2}bc \left( -\frac{ie\sqrt{1-c^2x^2} \operatorname{PolyLog}(2, e^{2i \arcsin(cx)})}{c\sqrt{cx-1}\sqrt{cx+1}} - \frac{ie\sqrt{1-c^2x^2} \arcsin(cx)^2}{c\sqrt{cx-1}\sqrt{cx+1}} + \frac{2e\sqrt{1-c^2x^2} \arcsin(cx) \log(1 - e^{2i \arcsin(cx)})}{c\sqrt{cx-1}\sqrt{cx+1}} \right) \end{aligned}$$

input `Int[((d + e*x^2)*(a + b*ArcCosh[c*x]))/x^3,x]`

output `-1/2*(d*(a + b*ArcCosh[c*x]))/x^2 + e*(a + b*ArcCosh[c*x])*Log[x] + (b*c*(d*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/x - (I*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]^2)/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (2*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[1 - E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (2*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[x])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (I*e*Sqrt[1 - c^2*x^2]*PolyLog[2, E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/2`

### 3.468.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6373 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

### 3.468.4 Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 135, normalized size of antiderivative = 0.54

method	result
derivativedivides	$c^2 \left( \frac{ae \ln(cx)}{c^2} - \frac{ad}{2c^2x^2} + \frac{b \left( -\frac{e \operatorname{arccosh}(cx)^2}{2} - \frac{d(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx))}{2x^2} + \ln(1+(cx+\sqrt{cx-1}\sqrt{cx+1})^2) \right)}{c^2} \right)$
default	$c^2 \left( \frac{ae \ln(cx)}{c^2} - \frac{ad}{2c^2x^2} + \frac{b \left( -\frac{e \operatorname{arccosh}(cx)^2}{2} - \frac{d(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx))}{2x^2} + \ln(1+(cx+\sqrt{cx-1}\sqrt{cx+1})^2) \right)}{c^2} \right)$
parts	$-\frac{ad}{2x^2} + ae \ln(x) + b c^2 \left( -\frac{e \operatorname{arccosh}(cx)^2}{2c^2} - \frac{d(-\sqrt{cx-1}\sqrt{cx+1}cx+c^2x^2+\operatorname{arccosh}(cx))}{2c^2x^2} + \frac{e \operatorname{arccosh}(cx) \ln(1+(cx+\sqrt{cx-1}\sqrt{cx+1})^2)}{c^2} \right)$

input `int((e*x^2+d)*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)`

output `c^2*(a/c^2*e*ln(c*x)-1/2*a*d/c^2/x^2+b/c^2*(-1/2*e*arccosh(c*x)^2-1/2*d*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+c^2*x^2+arccosh(c*x))/x^2+ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*e*arccosh(c*x)+1/2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)*e))`

### 3.468.5 Fracas [F]

$$\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(ex^2+d)(b\operatorname{arccosh}(cx)+a)}{x^3} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^3,x, algorithm="fracas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arccosh(c*x))/x^3, x)`

**3.468.6 Sympy [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d + ex^2)}{x^3} dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x))/x**3,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)/x**3, x)`

**3.468.7 Maxima [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `1/2*b*d*(sqrt(c^2*x^2 - 1)*c/x - arccosh(c*x)/x^2) + b*e*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/x, x) + a*e*log(x) - 1/2*a*d/x^2`

**3.468.8 Giac [F]**

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arccosh(c*x) + a)/x^3, x)`

**3.468.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx))(ex^2 + d)}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2))/x^3,x)`output `int(((a + b*acosh(c*x))*(d + e*x^2))/x^3, x)`

**3.469**  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^4} dx$

3.469.1 Optimal result . . . . .	3452
3.469.2 Mathematica [A] (verified) . . . . .	3452
3.469.3 Rubi [A] (verified) . . . . .	3453
3.469.4 Maple [A] (verified) . . . . .	3455
3.469.5 Fricas [A] (verification not implemented) . . . . .	3455
3.469.6 Sympy [F] . . . . .	3456
3.469.7 Maxima [A] (verification not implemented) . . . . .	3456
3.469.8 Giac [F] . . . . .	3456
3.469.9 Mupad [F(-1)] . . . . .	3457

**3.469.1 Optimal result**

Integrand size = 19, antiderivative size = 94

$$\int \frac{(d + ex^2)(a + \operatorname{arccosh}(cx))}{x^4} dx = \frac{bcd\sqrt{-1 + cx}\sqrt{1 + cx}}{6x^2} - \frac{d(a + \operatorname{arccosh}(cx))}{3x^3} - \frac{e(a + \operatorname{arccosh}(cx))}{x} + \frac{1}{6}bc(c^2d + 6e) \arctan\left(\sqrt{-1 + cx}\sqrt{1 + cx}\right)$$

output `-1/3*d*(a+b*arccosh(c*x))/x^3-e*(a+b*arccosh(c*x))/x+1/6*b*c*(c^2*d+6*e)*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))+1/6*b*c*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x^2`

**3.469.2 Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.36

$$\int \frac{(d + ex^2)(a + \operatorname{arccosh}(cx))}{x^4} dx = \frac{-2b(d + 3ex^2) \operatorname{arccosh}(cx) + \frac{bcdx(-1+c^2x^2)-2a\sqrt{-1+cx}\sqrt{1+cx}(d+3ex^2)+bc(c^2d+6e)x^3\sqrt{-1+c^2x^2} \arctan(\sqrt{-1+c^2x^2})}{\sqrt{-1+cx}\sqrt{1+cx}}}{6x^3}$$

input `Integrate[((d + e*x^2)*(a + b*ArcCosh[c*x]))/x^4,x]`

3.469.  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^4} dx$

output  $(-2*b*(d + 3*e*x^2)*ArcCosh[c*x] + (b*c*d*x*(-1 + c^2*x^2) - 2*a*sqrt[-1 + c*x]*sqrt[1 + c*x]*(d + 3*e*x^2) + b*c*(c^2*d + 6*e)*x^3*sqrt[-1 + c^2*x^2]*ArcTan[sqrt[-1 + c^2*x^2]])/(sqrt[-1 + c*x]*sqrt[1 + c*x])/(6*x^3)$

### 3.469.3 Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6371, 25, 956, 103, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^4} dx \\
 & \quad \downarrow \text{6371} \\
 & -\frac{1}{3}bc \int -\frac{3ex^2 + d}{x^3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{e(a + \operatorname{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3}bc \int \frac{3ex^2 + d}{x^3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{e(a + \operatorname{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{956} \\
 & \frac{1}{3}bc \left( \frac{1}{2}(c^2d + 6e) \int \frac{1}{x\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{d\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right) - \frac{d(a + \operatorname{barccosh}(cx))}{3x^3} - \\
 & \quad \frac{x}{e(a + \operatorname{barccosh}(cx))} \\
 & \quad \downarrow \text{103} \\
 & \frac{1}{3}bc \left( \frac{1}{2}c(c^2d + 6e) \int \frac{1}{(cx-1)(cx+1)c+c} d(\sqrt{cx-1}\sqrt{cx+1}) + \frac{d\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right) - \\
 & \quad \frac{d(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{e(a + \operatorname{barccosh}(cx))}{x} \\
 & \quad \downarrow \text{218} \\
 & -\frac{d(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{e(a + \operatorname{barccosh}(cx))}{x} + \\
 & \frac{1}{3}bc \left( \frac{1}{2} \arctan(\sqrt{cx-1}\sqrt{cx+1}) (c^2d + 6e) + \frac{d\sqrt{cx-1}\sqrt{cx+1}}{2x^2} \right)
 \end{aligned}$$

input `Int[((d + e*x^2)*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*(d*(a + b*ArcCosh[c*x]))/x^3 - (e*(a + b*ArcCosh[c*x]))/x + (b*c*((d*  
Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*x^2) + ((c^2*d + 6*e)*ArcTan[Sqrt[-1 + c*  
x]*Sqrt[1 + c*x]]/2))/3`

### 3.469.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 103 `Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]*((e_.) + (f_.)*(x_))), x_] := Simp[b*f Subst[Int[1/(d*(b*e - a*f)^2 + b*f^2*x^2), x], x, Sqrt[a + b*x]*Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - f*(b*c + a*d), 0]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 956 `Int[((e_.)*(x_)^(m_.))*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[c*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*(a2 + b2*x^(n/2))^(p + 1)/(a1*a2*e*(m + 1))), x] + Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(a1*a2*e^n*(m + 1)) Int[(e*x)^(m + n)*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]`

rule 6371 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[d*(f*x)^(m + 1)*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (Simp[e*(f*x)^(m + 3)*((a + b*ArcCosh[c*x])/(f^3*(m + 3))), x] - Simp[b*(c/(f*(m + 1)*(m + 3)) Int[(f*x)^(m + 1)*((d*(m + 3) + e*(m + 1)*x^2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x)], x) /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && NeQ[m, -1] && NeQ[m, -3]`

### 3.469.4 Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.50

method	result
parts	$a\left(-\frac{e}{x} - \frac{d}{3x^3}\right) + b c^3 \left( -\frac{\operatorname{arccosh}(cx)e}{c^3 x} - \frac{\operatorname{arccosh}(cx)d}{3c^3 x^3} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{6c^4\sqrt{c^2x^2-1}} \left( \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) c^4 d x^2 + 6 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) \right) \right)$
derivativedivides	$c^3 \left( \frac{a\left(-\frac{d}{3c x^3} - \frac{e}{cx}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arccosh}(cx)d}{3c x^3} - \frac{\operatorname{arccosh}(cx)e}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{6\sqrt{c^2x^2-1}} \left( \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) c^4 d x^2 + 6 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) \right) \right)}{c^2} \right)$
default	$c^3 \left( \frac{a\left(-\frac{d}{3c x^3} - \frac{e}{cx}\right)}{c^2} + \frac{b\left(-\frac{\operatorname{arccosh}(cx)d}{3c x^3} - \frac{\operatorname{arccosh}(cx)e}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{6\sqrt{c^2x^2-1}} \left( \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) c^4 d x^2 + 6 \arctan\left(\frac{1}{\sqrt{c^2x^2-1}}\right) \right) \right)}{c^2} \right)$

input `int((e*x^2+d)*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)`

output `a*(-e/x-1/3*d/x^3)+b*c^3*(-1/c^3*arccosh(c*x)*e/x-1/3*arccosh(c*x)*d/c^3/x^3-1/6/c^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(arctan(1/(c^2*x^2-1)^(1/2))*c^4*d*x^2+6*arctan(1/(c^2*x^2-1)^(1/2))*e*c^2*x^2-(c^2*x^2-1)^(1/2)*c^2*d)/(c^2*x^2-1)^(1/2)/x^2)`

### 3.469.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.48

$$\int \frac{(d + ex^2)(a + b\operatorname{arccosh}(cx))}{x^4} dx = \frac{2(bc^3d + 6bce)x^3 \arctan(-cx + \sqrt{c^2x^2 - 1}) + 2(bd + 3be)x^3 \log(-cx + \sqrt{c^2x^2 - 1}) + \sqrt{c^2x^2 - 1}bcdx - 6x^3}{6x^3}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")`

output `1/6*(2*(b*c^3*d + 6*b*c*e)*x^3*arctan(-c*x + sqrt(c^2*x^2 - 1)) + 2*(b*d + 3*b*e)*x^3*log(-c*x + sqrt(c^2*x^2 - 1)) + sqrt(c^2*x^2 - 1)*b*c*d*x - 6*a*e*x^2 - 2*a*d - 2*(3*b*e*x^2 - (b*d + 3*b*e)*x^3 + b*d)*log(c*x + sqrt(c^2*x^2 - 1)))/x^3`

3.469.  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^4} dx$



**3.469.6 Sympy [F]**

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx))(d + ex^2)}{x^4} dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x))/x**4,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)/x**4, x)`

**3.469.7 Maxima [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.90

$$\begin{aligned} & \int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^4} dx \\ &= -\frac{1}{6} \left( \left( c^2 \arcsin\left(\frac{1}{c|x|}\right) - \frac{\sqrt{c^2x^2 - 1}}{x^2} \right) c + \frac{2 \operatorname{arcosh}(cx)}{x^3} \right) bd \\ & \quad - \left( c \arcsin\left(\frac{1}{c|x|}\right) + \frac{\operatorname{arcosh}(cx)}{x} \right) be - \frac{ae}{x} - \frac{ad}{3x^3} \end{aligned}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")`

output `-1/6*((c^2*arcsin(1/(c*abs(x))) - sqrt(c^2*x^2 - 1)/x^2)*c + 2*arccosh(c*x)/x^3)*b*d - (c*arcsin(1/(c*abs(x))) + arccosh(c*x)/x)*b*e - a*e/x - 1/3*a*d/x^3`

**3.469.8 Giac [F]**

$$\int \frac{(d + ex^2)(a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)}{x^4} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arccosh(c*x) + a)/x^4, x)`

---

3.469.  $\int \frac{(d+ex^2)(a+b\operatorname{arccosh}(cx))}{x^4} dx$

**3.469.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)(a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx))(ex^2 + d)}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2))/x^4,x)`output `int(((a + b*acosh(c*x))*(d + e*x^2))/x^4, x)`

### 3.470 $\int x^4(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.470.1 Optimal result . . . . .	3458
3.470.2 Mathematica [A] (verified) . . . . .	3459
3.470.3 Rubi [A] (verified) . . . . .	3459
3.470.4 Maple [A] (verified) . . . . .	3462
3.470.5 Fricas [A] (verification not implemented) . . . . .	3462
3.470.6 Sympy [F] . . . . .	3463
3.470.7 Maxima [A] (verification not implemented) . . . . .	3463
3.470.8 Giac [F(-2)] . . . . .	3464
3.470.9 Mupad [F(-1)] . . . . .	3464

#### 3.470.1 Optimal result

Integrand size = 21, antiderivative size = 319

$$\int x^4(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{b(63c^4d^2 + 90c^2de + 35e^2)(1 - c^2x^2)}{315c^9\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{2b(63c^4d^2 + 135c^2de + 70e^2)(1 - c^2x^2)^2}{945c^9\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{b(21c^4d^2 + 90c^2de + 70e^2)(1 - c^2x^2)^3}{525c^9\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{2be(9c^2d + 14e)(1 - c^2x^2)^4}{441c^9\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{be^2(1 - c^2x^2)^5}{81c^9\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{1}{5}d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^2x^9(a + \operatorname{barccosh}(cx))$$

output  $\frac{1}{5}d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^2x^9(a + \operatorname{barccosh}(cx)) + \frac{1}{315}b(63c^4d^2 + 90c^2de + 35e^2)(-c^2x^2 + 1)/c^9 / (cx - 1)^{1/2} / (cx + 1)^{1/2} - \frac{2}{945}b(63c^4d^2 + 135c^2de + 70e^2)(-c^2x^2 + 1)^2/c^9 / (cx - 1)^{1/2} / (cx + 1)^{1/2} + \frac{1}{525}b(21c^4d^2 + 90c^2de + 70e^2)(-c^2x^2 + 1)^3/c^9 / (cx - 1)^{1/2} / (cx + 1)^{1/2} - \frac{2}{441}be(9c^2d + 14e)(-c^2x^2 + 1)^4/c^9 / (cx - 1)^{1/2} / (cx + 1)^{1/2} + \frac{1}{81}be^2(-c^2x^2 + 1)^5/c^9 / (cx - 1)^{1/2} / (cx + 1)^{1/2}$

**3.470.2 Mathematica [A] (verified)**

Time = 0.18 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.60

$$\int x^4(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx$$

$$= \frac{315ax^5(63d^2+90dex^2+35e^2x^4) - \frac{b\sqrt{-1+cx}\sqrt{1+cx}(4480e^2+160c^2e(81d+14ex^2)+24c^4(441d^2+270dex^2+70e^2x^4)+4c^6(1323d^2+90d^2ex^2+35e^2x^4))}{c^9} + 315b^2x^5(63d^2+90dex^2+35e^2x^4)\operatorname{ArcCosh}[cx]}{99225}$$

99225

input `Integrate[x^4*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `(315*a*x^5*(63*d^2 + 90*d*e*x^2 + 35*e^2*x^4) - (b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(4480*e^2 + 160*c^2*e*(81*d + 14*e*x^2) + 24*c^4*(441*d^2 + 270*d*e*x^2 + 70*e^2*x^4) + 4*c^6*(1323*d^2*x^2 + 1215*d*e*x^4 + 350*e^2*x^6) + c^8*(3969*d^2*x^4 + 4050*d*e*x^6 + 1225*e^2*x^8)))/c^9 + 315*b*x^5*(63*d^2 + 90*d*e*x^2 + 35*e^2*x^4)*ArcCosh[c*x])/99225`

**3.470.3 Rubi [A] (verified)**Time = 0.60 (sec) , antiderivative size = 267, normalized size of antiderivative = 0.84, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6373, 27, 1905, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^4(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx$$

$$\downarrow 6373$$

$$-bc \int \frac{x^5(35e^2x^4+90dex^2+63d^2)}{315\sqrt{cx-1}\sqrt{cx+1}}dx + \frac{1}{5}d^2x^5(a+\operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{9}e^2x^9(a+\operatorname{barccosh}(cx))$$

$$\downarrow 27$$

$$-\frac{1}{315}bc \int \frac{x^5(35e^2x^4+90dex^2+63d^2)}{\sqrt{cx-1}\sqrt{cx+1}}dx + \frac{1}{5}d^2x^5(a+\operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{9}e^2x^9(a+\operatorname{barccosh}(cx))$$

$$\downarrow 1905$$

---

3.470.  $\int x^4(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx$

$$\begin{aligned}
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^5(35e^2x^4+90dex^2+63d^2)}{\sqrt{c^2x^2-1}} dx}{315\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a + \operatorname{barccosh}(cx)) + \\
 & \qquad \qquad \qquad \frac{1}{9}e^2x^9(a + \operatorname{barccosh}(cx)) \\
 & \qquad \qquad \qquad \downarrow \text{1578} \\
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^4(35e^2x^4+90dex^2+63d^2)}{\sqrt{c^2x^2-1}} dx^2}{630\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a + \operatorname{barccosh}(cx)) + \\
 & \qquad \qquad \qquad \frac{1}{9}e^2x^9(a + \operatorname{barccosh}(cx)) \\
 & \qquad \qquad \qquad \downarrow \text{1195} \\
 & \frac{bc\sqrt{c^2x^2-1} \int \left( \frac{35e^2(c^2x^2-1)^{7/2}}{c^8} + \frac{10e(9dc^2+14e)(c^2x^2-1)^{5/2}}{c^8} + \frac{3(21d^2c^4+90dec^2+70e^2)(c^2x^2-1)^{3/2}}{c^8} + \frac{2(63d^2c^4+135dec^2+70e^2)}{c^8} \right)}{630\sqrt{cx-1}\sqrt{cx+1}} \\
 & \qquad \qquad \qquad \frac{1}{5}d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^2x^9(a + \operatorname{barccosh}(cx)) \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & \frac{1}{5}d^2x^5(a + \operatorname{barccosh}(cx)) + \frac{2}{7}dex^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^2x^9(a + \operatorname{barccosh}(cx)) - \\
 & \frac{bc\sqrt{c^2x^2-1} \left( \frac{20e(c^2x^2-1)^{7/2}(9c^2d+14e)}{7c^{10}} + \frac{70e^2(c^2x^2-1)^{9/2}}{9c^{10}} + \frac{6(c^2x^2-1)^{5/2}(21c^4d^2+90c^2de+70e^2)}{5c^{10}} + \frac{4(c^2x^2-1)^{3/2}(63c^4d^2+135c^2de+70e^2)}{3c^{10}} \right)}{630\sqrt{cx-1}\sqrt{cx+1}}
 \end{aligned}$$

input `Int[x^4*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `-1/630*(b*c*Sqrt[-1 + c^2*x^2]*((2*(63*c^4*d^2 + 90*c^2*d*e + 35*e^2)*Sqrt[-1 + c^2*x^2])/c^10 + (4*(63*c^4*d^2 + 135*c^2*d*e + 70*e^2)*(-1 + c^2*x^2)^(3/2))/(3*c^10) + (6*(21*c^4*d^2 + 90*c^2*d*e + 70*e^2)*(-1 + c^2*x^2)^(5/2))/(5*c^10) + (20*e*(9*c^2*d + 14*e)*(-1 + c^2*x^2)^(7/2))/(7*c^10) + (70*e^2*(-1 + c^2*x^2)^(9/2))/(9*c^10)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (d^2*x^5*(a + b*ArcCosh[c*x]))/5 + (2*d*e*x^7*(a + b*ArcCosh[c*x]))/7 + (e^2*x^9*(a + b*ArcCosh[c*x]))/9`

## 3.470.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 1195 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`
- rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`
- rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

**3.470.4 Maple [A] (verified)**

Time = 0.84 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.67

method	result
parts	$a\left(\frac{1}{9}e^2x^9 + \frac{2}{7}dex^7 + \frac{1}{5}d^2x^5\right) + \frac{b\left(\frac{c^5 \operatorname{arccosh}(cx)e^2x^9}{9} + \frac{2c^5 \operatorname{arccosh}(cx)dex^7}{7} + \frac{\operatorname{arccosh}(cx)c^5x^5d^2}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{c^4}\right)}{\sqrt{cx-1}\sqrt{cx+1}}$
derivativedivides	$\frac{a\left(\frac{1}{5}c^9d^2x^5 + \frac{2}{7}dc^9ex^7 + \frac{1}{9}e^2c^9x^9\right)}{c^4} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^9d^2x^5}{5} + \frac{2 \operatorname{arccosh}(cx)dc^9ex^7}{7} + \frac{\operatorname{arccosh}(cx)e^2c^9x^9}{9} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{c^4}\right)}{\sqrt{cx-1}\sqrt{cx+1}}$
default	$\frac{a\left(\frac{1}{5}c^9d^2x^5 + \frac{2}{7}dc^9ex^7 + \frac{1}{9}e^2c^9x^9\right)}{c^4} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^9d^2x^5}{5} + \frac{2 \operatorname{arccosh}(cx)dc^9ex^7}{7} + \frac{\operatorname{arccosh}(cx)e^2c^9x^9}{9} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{c^4}\right)}{\sqrt{cx-1}\sqrt{cx+1}}$

input `int(x^4*(e*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`output `a*(1/9*e^2*x^9+2/7*d*e*x^7+1/5*d^2*x^5)+b/c^5*(1/9*c^5*arccosh(c*x)*e^2*x^9+2/7*c^5*arccosh(c*x)*d*e*x^7+1/5*arccosh(c*x)*c^5*x^5*d^2-1/99225/c^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(1225*c^8*e^2*x^8+4050*c^8*d*e*x^6+3969*c^8*d^2*x^4+1400*c^6*e^2*x^6+4860*c^6*d*e*x^4+5292*c^6*d^2*x^2+1680*c^4*e^2*x^4+6480*c^4*d*e*x^2+10584*c^4*d^2+2240*c^2*e^2*x^2+12960*c^2*d*e+4480*e^2))`**3.470.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 231, normalized size of antiderivative = 0.72

$$\int x^4(d+ex^2)^2(a+b\operatorname{arccosh}(cx))dx$$

$$= \frac{11025ac^9e^2x^9 + 28350ac^9dex^7 + 19845ac^9d^2x^5 + 315(35bc^9e^2x^9 + 90bc^9dex^7 + 63bc^9d^2x^5)\log(cx + \sqrt{c^2x^2 - 1}) - (1225b^2c^8e^2x^8 + 10584b^2c^4d^2 + 50(81b^2c^8d^2e + 28b^2c^6e^2)x^6 + 12960b^2c^2d^2e + 3(1323b^2c^8d^2 + 1620b^2c^6d^2e + 560b^2c^4e^2)x^4 + 4480b^2e^2 + 4(1323b^2c^6d^2 + 1620b^2c^4d^2e + 560b^2c^2e^2)x^2)\sqrt{c^2x^2 - 1}}{c^9}$$

input `integrate(x^4*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fracas")`output `1/99225*(11025*a*c^9*e^2*x^9 + 28350*a*c^9*d*e*x^7 + 19845*a*c^9*d^2*x^5 + 315*(35*b*c^9*e^2*x^9 + 90*b*c^9*d*e*x^7 + 63*b*c^9*d^2*x^5)*log(c*x + sqrt(c^2*x^2 - 1)) - (1225*b^2*c^8*e^2*x^8 + 10584*b^2*c^4*d^2 + 50*(81*b^2*c^8*d^2*e + 28*b^2*c^6*e^2)*x^6 + 12960*b^2*c^2*d^2*e + 3*(1323*b^2*c^8*d^2 + 1620*b^2*c^6*d^2*e + 560*b^2*c^4*e^2)*x^4 + 4480*b^2*e^2 + 4*(1323*b^2*c^6*d^2 + 1620*b^2*c^4*d^2*e + 560*b^2*c^2*e^2)*x^2)*sqrt(c^2*x^2 - 1)/c^9`

---

3.470.  $\int x^4(d+ex^2)^2(a+b\operatorname{arccosh}(cx))dx$

**3.470.6 Sympy [F]**

$$\int x^4 (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (d + ex^2)^2 dx$$

input `integrate(x**4*(e*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `Integral(x**4*(a + b*acosh(c*x))*(d + e*x**2)**2, x)`

**3.470.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 305, normalized size of antiderivative = 0.96

$$\begin{aligned} \int x^4 (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx &= \frac{1}{9} ae^2 x^9 + \frac{2}{7} adex^7 + \frac{1}{5} ad^2 x^5 \\ &+ \frac{1}{75} \left( 15x^5 \operatorname{arcosh}(cx) - \left( \frac{3\sqrt{c^2x^2-1}x^4}{c^2} + \frac{4\sqrt{c^2x^2-1}x^2}{c^4} + \frac{8\sqrt{c^2x^2-1}}{c^6} \right) c \right) bd^2 \\ &+ \frac{2}{245} \left( 35x^7 \operatorname{arcosh}(cx) - \left( \frac{5\sqrt{c^2x^2-1}x^6}{c^2} + \frac{6\sqrt{c^2x^2-1}x^4}{c^4} + \frac{8\sqrt{c^2x^2-1}x^2}{c^6} + \frac{16\sqrt{c^2x^2-1}}{c^8} \right) c \right) bde \\ &+ \frac{1}{2835} \left( 315x^9 \operatorname{arcosh}(cx) - \left( \frac{35\sqrt{c^2x^2-1}x^8}{c^2} + \frac{40\sqrt{c^2x^2-1}x^6}{c^4} + \frac{48\sqrt{c^2x^2-1}x^4}{c^6} + \frac{64\sqrt{c^2x^2-1}x^2}{c^8} \right) c \right) bde^2 \end{aligned}$$

input `integrate(x^4*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/9*a*e^2*x^9 + 2/7*a*d*e*x^7 + 1/5*a*d^2*x^5 + 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d^2 + 2/245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*d*e + 1/2835*(315*x^9*arccosh(c*x) - (35*sqrt(c^2*x^2 - 1)*x^8/c^2 + 40*sqrt(c^2*x^2 - 1)*x^6/c^4 + 48*sqrt(c^2*x^2 - 1)*x^4/c^6 + 64*sqrt(c^2*x^2 - 1)*x^2/c^8 + 128*sqrt(c^2*x^2 - 1)/c^10)*c)*b*e^2`



**3.470.8 Giac [F(-2)]**

Exception generated.

$$\int x^4(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

**3.470.9 Mupad [F(-1)]**

Timed out.

$$\int x^4(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (ex^2 + d)^2 dx$$

input `int(x^4*(a + b*acosh(c*x))*(d + e*x^2)^2,x)`

output `int(x^4*(a + b*acosh(c*x))*(d + e*x^2)^2, x)`

### 3.471 $\int x^3(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.471.1 Optimal result . . . . .	3465
3.471.2 Mathematica [A] (warning: unable to verify) . . . . .	3466
3.471.3 Rubi [A] (verified) . . . . .	3466
3.471.4 Maple [A] (verified) . . . . .	3470
3.471.5 Fricas [A] (verification not implemented) . . . . .	3471
3.471.6 Sympy [F] . . . . .	3471
3.471.7 Maxima [A] (verification not implemented) . . . . .	3472
3.471.8 Giac [F(-2)] . . . . .	3472
3.471.9 Mupad [F(-1)] . . . . .	3473

#### 3.471.1 Optimal result

Integrand size = 21, antiderivative size = 341

$$\int x^3(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{b(288c^4d^2 + 320c^2de + 105e^2) x(1 - c^2x^2)}{3072c^7\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{b(288c^4d^2 + 320c^2de + 105e^2) x^3(1 - c^2x^2)}{4608c^5\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{be(64c^2d + 21e) x^5(1 - c^2x^2)}{1152c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{be^2x^7(1 - c^2x^2)}{64c\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx))$$

$$- \frac{b(288c^4d^2 + 320c^2de + 105e^2) \sqrt{-1 + c^2x^2} \operatorname{arctanh}\left(\frac{cx}{\sqrt{-1 + c^2x^2}}\right)}{3072c^8\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output 1/4*d^2*x^4*(a+b*arccosh(c*x))+1/3*d*e*x^6*(a+b*arccosh(c*x))+1/8*e^2*x^8*
(a+b*arccosh(c*x))+1/3072*b*(288*c^4*d^2+320*c^2*d*e+105*e^2)*x*(-c^2*x^2+
1)/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/4608*b*(288*c^4*d^2+320*c^2*d*e+105*e
^2)*x^3*(-c^2*x^2+1)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/1152*b*e*(64*c^2*d+
21*e)*x^5*(-c^2*x^2+1)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/64*b*e^2*x^7*(-c^
2*x^2+1)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/3072*b*(288*c^4*d^2+320*c^2*d*e+1
05*e^2)*arctanh(c*x/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/c^8/(c*x-1)^(1/2)
/(c*x+1)^(1/2)
```

**3.471.2 Mathematica [A] (warning: unable to verify)**

Time = 0.22 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.63

$$\int x^3(d + ex^2)^2(a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{384ac^8x^4(6d^2 + 8dex^2 + 3e^2x^4) - bcx\sqrt{-1 + cx}\sqrt{1 + cx}(315e^2 + 30c^2e(32d + 7ex^2) + 8c^4(108d^2 + 80dex^2 + 21e^2x^4) + 16c^6(36d^2x^2 + 32d^2ex^4 + 9e^2x^6)) + 384b^2c^8x^4(6d^2 + 8dex^2 + 3e^2x^4)\operatorname{ArcCosh}[cx] - 6b^2(288c^4d^2 + 320c^2de + 105e^2)\operatorname{ArcTanh}[\sqrt{(-1 + cx)/(1 + cx)}}]}{9216c^8}$$

input `Integrate[x^3*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`output `(384*a*c^8*x^4*(6*d^2 + 8*d*e*x^2 + 3*e^2*x^4) - b*c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(315*e^2 + 30*c^2*e*(32*d + 7*e*x^2) + 8*c^4*(108*d^2 + 80*d*e*x^2 + 21*e^2*x^4) + 16*c^6*(36*d^2*x^2 + 32*d^2*e*x^4 + 9*e^2*x^6)) + 384*b*c^8*x^4*(6*d^2 + 8*d*e*x^2 + 3*e^2*x^4)*ArcCosh[c*x] - 6*b*(288*c^4*d^2 + 320*c^2*d*e + 105*e^2)*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]])/(9216*c^8)`**3.471.3 Rubi [A] (verified)**Time = 0.56 (sec) , antiderivative size = 266, normalized size of antiderivative = 0.78, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6373, 27, 1905, 1590, 363, 262, 262, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^3(d + ex^2)^2(a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6373}$$

$$-bc \int \frac{x^4(3e^2x^4 + 8dex^2 + 6d^2)}{24\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{27}$$

$$-\frac{1}{24}bc \int \frac{x^4(3e^2x^4 + 8dex^2 + 6d^2)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{1905}$$

---

 3.471.  $\int x^3(d + ex^2)^2(a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^4(3e^2x^4+8dex^2+6d^2)}{\sqrt{c^2x^2-1}} dx}{24\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \\
& \qquad \qquad \qquad \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx)) \\
& \qquad \qquad \qquad \downarrow \text{1590} \\
& -\frac{bc\sqrt{c^2x^2-1} \left( \frac{\int \frac{x^4(48c^2d^2+e(64dc^2+21e)x^2)}{\sqrt{c^2x^2-1}} dx}{8c^2} + \frac{3e^2x^7\sqrt{c^2x^2-1}}{8c^2} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \\
& \qquad \qquad \qquad \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx)) \\
& \qquad \qquad \qquad \downarrow \text{363} \\
& -\frac{bc\sqrt{c^2x^2-1} \left( \frac{(288c^4d^2+5e(64c^2d+21e)) \int \frac{x^4}{\sqrt{c^2x^2-1}} dx}{6c^2} + \frac{e^2x^5\sqrt{c^2x^2-1}(64c^2d+21e)}{6c^2} + \frac{3e^2x^7\sqrt{c^2x^2-1}}{8c^2} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{4}d^2x^4(a + \\
& \qquad \qquad \qquad \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx)) \\
& \qquad \qquad \qquad \downarrow \text{262} \\
& -\frac{bc\sqrt{c^2x^2-1} \left( \frac{(288c^4d^2+5e(64c^2d+21e)) \left( \frac{3 \int \frac{x^2}{\sqrt{c^2x^2-1}} dx}{4c^2} + \frac{x^3\sqrt{c^2x^2-1}}{4c^2} \right)}{6c^2} + \frac{e^2x^5\sqrt{c^2x^2-1}(64c^2d+21e)}{6c^2} + \frac{3e^2x^7\sqrt{c^2x^2-1}}{8c^2} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} + \\
& \qquad \qquad \qquad \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx)) \\
& \qquad \qquad \qquad \downarrow \text{262} \\
& -\frac{bc\sqrt{c^2x^2-1} \left( \frac{(288c^4d^2+5e(64c^2d+21e)) \left( \frac{3 \left( \frac{\int \frac{1}{\sqrt{c^2x^2-1}} dx}{2c^2} + \frac{x\sqrt{c^2x^2-1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{c^2x^2-1}}{4c^2} \right)}{6c^2} + \frac{e^2x^5\sqrt{c^2x^2-1}(64c^2d+21e)}{6c^2} + \frac{3e^2x^7\sqrt{c^2x^2-1}}{8c^2} \right)}{24\sqrt{cx-1}\sqrt{cx+1}} + \\
& \qquad \qquad \qquad \frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx))
\end{aligned}$$

---

3.471.  $\int x^3(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx$

↓ 224

$$bc\sqrt{c^2x^2 - 1} \left( \frac{(288c^4d^2 + 5e(64c^2d + 21e)) \left( \frac{3 \left( \frac{\int \frac{1}{1 - \frac{c^2x^2}{c^2x^2 - 1}} d \frac{x}{\sqrt{c^2x^2 - 1}} + \frac{x\sqrt{c^2x^2 - 1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{c^2x^2 - 1}}{4c^2} \right)}{6c^2} + \frac{e x^5 \sqrt{c^2x^2 - 1} (64c^2d + 21e)}{8c^2} + \frac{3e^2x^7\sqrt{c^2x^2 - 1}}{8c^2} \right)$$

$$\frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx))$$

↓ 219

$$\frac{1}{4}d^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{3}dex^6(a + \operatorname{barccosh}(cx)) + \frac{1}{8}e^2x^8(a + \operatorname{barccosh}(cx)) - bc\sqrt{c^2x^2 - 1} \left( \frac{\left( \frac{3 \left( \frac{\operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2 - 1}}\right)}{2c^3} + \frac{x\sqrt{c^2x^2 - 1}}{2c^2} \right)}{4c^2} + \frac{x^3\sqrt{c^2x^2 - 1}}{4c^2} \right) (288c^4d^2 + 5e(64c^2d + 21e))}{6c^2} + \frac{e x^5 \sqrt{c^2x^2 - 1} (64c^2d + 21e)}{8c^2} + \frac{3e^2x^7\sqrt{c^2x^2 - 1}}{8c^2} \right)$$

$$24\sqrt{cx - 1}\sqrt{cx + 1}$$

input `Int[x^3*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `(d^2*x^4*(a + b*ArcCosh[c*x]))/4 + (d*e*x^6*(a + b*ArcCosh[c*x]))/3 + (e^2*x^8*(a + b*ArcCosh[c*x]))/8 - (b*c*sqrt[-1 + c^2*x^2]*((3*e^2*x^7*sqrt[-1 + c^2*x^2])/(8*c^2) + ((e*(64*c^2*d + 21*e))*x^5*sqrt[-1 + c^2*x^2])/(6*c^2) + ((288*c^4*d^2 + 5*e*(64*c^2*d + 21*e))*((x^3*sqrt[-1 + c^2*x^2])/(4*c^2) + (3*((x*sqrt[-1 + c^2*x^2])/(2*c^2) + ArcTanh[(c*x)/sqrt[-1 + c^2*x^2]])/(2*c^3)))/(4*c^2)))/(6*c^2))/(8*c^2))/(24*sqrt[-1 + c*x]*sqrt[1 + c*x])`

3.471.  $\int x^3(d + ex^2)^2(a + \operatorname{barccosh}(cx)) dx$

## 3.471.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 224 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 262 `Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[c*(c*x)^(m - 1)*((a + b*x^2)^(p + 1)/(b*(m + 2*p + 1))), x] - Simp[a*c^2*((m - 1)/(b*(m + 2*p + 1)) Int[(c*x)^(m - 2)*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[m, 2 - 1] && NeQ[m + 2*p + 1, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`
- rule 363 `Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^(m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`
- rule 1590 `Int[((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^(m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]`

```
rule 1905 Int[((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_)^(non2_.))^(q_.)*((d2_) + (e2_.)
*(x_)^(non2_.))^(q_.)*((a_.) + (b_.)*(x_)^(n_) + (c_.)*(x_)^(n2_))^(p_.), x
_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[
q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a
+ b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p,
q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]
```

```
rule 6373 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x
_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Sim
p[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] &&
NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && Le
Q[m + p, 0]))
```

### 3.471.4 Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 364, normalized size of antiderivative = 1.07

method	result
parts	$a\left(\frac{1}{8}e^2x^8 + \frac{1}{3}dex^6 + \frac{1}{4}d^2x^4\right) + \frac{b\left(\frac{c^4 \operatorname{arccosh}(cx)e^2x^8}{8} + \frac{c^4 \operatorname{arccosh}(cx)dex^6}{3} + \frac{\operatorname{arccosh}(cx)c^4x^4d^2}{4} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{5}\right)}{e^4}$
derivativedivides	$\frac{a\left(\frac{1}{4}c^8d^2x^4 + \frac{1}{3}c^8dex^6 + \frac{1}{8}c^8e^2x^8\right)}{e^4} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^8d^2x^4}{4} + \frac{\operatorname{arccosh}(cx)d c^8e x^6}{3} + \frac{\operatorname{arccosh}(cx)e^2c^8x^8}{8} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{576c^7d^2}\right)}{e^4}$
default	$\frac{a\left(\frac{1}{4}c^8d^2x^4 + \frac{1}{3}c^8dex^6 + \frac{1}{8}c^8e^2x^8\right)}{e^4} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^8d^2x^4}{4} + \frac{\operatorname{arccosh}(cx)d c^8e x^6}{3} + \frac{\operatorname{arccosh}(cx)e^2c^8x^8}{8} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{576c^7d^2}\right)}{e^4}$

```
input int(x^3*(e*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)
```

output  $a*(1/8*e^{2*x^8}+1/3*d*e*x^6+1/4*d^2*x^4)+b/c^4*(1/8*c^4*\operatorname{arccosh}(c*x)*e^{2*x^8}+1/3*c^4*\operatorname{arccosh}(c*x)*d*e*x^6+1/4*\operatorname{arccosh}(c*x)*c^4*x^4*d^2-1/9216/c^4*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(576*c^7*d^2*(c^2*x^2-1)^{(1/2)}*x^3+512*c^7*d*e*(c^2*x^2-1)^{(1/2)}*x^5+144*e^2*(c^2*x^2-1)^{(1/2)}*c^7*x^7+864*c^5*d^2*x*(c^2*x^2-1)^{(1/2)}+640*(c^2*x^2-1)^{(1/2)}*c^5*d*e*x^3+168*e^2*c^5*x^5*(c^2*x^2-1)^{(1/2)}+864*c^4*d^2*\ln(c*x+(c^2*x^2-1)^{(1/2)})+960*d*c^3*e*x*(c^2*x^2-1)^{(1/2)}+210*e^2*(c^2*x^2-1)^{(1/2)}*c^3*x^3+960*d*c^2*e*\ln(c*x+(c^2*x^2-1)^{(1/2)})+315*e^2*c*x*(c^2*x^2-1)^{(1/2)}+315*e^2*\ln(c*x+(c^2*x^2-1)^{(1/2)}))/(c^2*x^2-1)^{(1/2)}$

### 3.471.5 Fricas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.67

$$\int x^3(d+ex^2)^2(a+b\operatorname{arccosh}(cx))dx$$

$$= \frac{1152ac^8e^2x^8 + 3072ac^8dex^6 + 2304ac^8d^2x^4 + 3(384bc^8e^2x^8 + 1024bc^8dex^6 + 768bc^8d^2x^4 - 288bc^4d^2 - 320bc^2de - 105b^2e^2)\log(cx + \sqrt{c^2x^2 - 1}) - (144bc^7e^{2x^7} + 8(64bc^7d^2e + 21bc^5e^2)x^5 + 2(288bc^7d^2 + 320bc^5de + 105bc^3e^2)x^3 + 3(288bc^5d^2 + 320bc^3de + 105b^2e^2)x)\sqrt{c^2x^2 - 1}}{c^8}$$

input `integrate(x^3*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $1/9216*(1152*a*c^8*e^{2*x^8} + 3072*a*c^8*d*e*x^6 + 2304*a*c^8*d^2*x^4 + 3*(384*b*c^8*e^{2*x^8} + 1024*b*c^8*d*e*x^6 + 768*b*c^8*d^2*x^4 - 288*b*c^4*d^2 - 320*b*c^2*d*e - 105*b^2*e^2)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (144*b*c^7*e^{2*x^7} + 8*(64*b*c^7*d^2*e + 21*b*c^5*e^2)*x^5 + 2*(288*b*c^7*d^2 + 320*b*c^5*d*e + 105*b*c^3*e^2)*x^3 + 3*(288*b*c^5*d^2 + 320*b*c^3*d*e + 105*b*c^2*e^2)*x)*\sqrt{c^2*x^2 - 1})/c^8$

### 3.471.6 Sympy [F]

$$\int x^3(d+ex^2)^2(a+b\operatorname{arccosh}(cx))dx = \int x^3(a+b\operatorname{acosh}(cx))(d+ex^2)^2dx$$

input `integrate(x**3*(e*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `Integral(x**3*(a + b*acosh(c*x))*(d + e*x**2)**2, x)`

---

3.471.  $\int x^3(d+ex^2)^2(a+b\operatorname{arccosh}(cx))dx$



**3.471.7 Maxima [A] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 332, normalized size of antiderivative = 0.97

$$\int x^3(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx = \frac{1}{8}ae^2x^8 + \frac{1}{3}adex^6 + \frac{1}{4}ad^2x^4 + \frac{1}{32}\left(8x^4\operatorname{arcosh}(cx) - \left(\frac{2\sqrt{c^2x^2-1}x^3}{c^2} + \frac{3\sqrt{c^2x^2-1}x}{c^4} + \frac{3\log(2c^2x+2\sqrt{c^2x^2-1}c)}{c^5}\right)c\right)bd^2 + \frac{1}{144}\left(48x^6\operatorname{arcosh}(cx) - \left(\frac{8\sqrt{c^2x^2-1}x^5}{c^2} + \frac{10\sqrt{c^2x^2-1}x^3}{c^4} + \frac{15\sqrt{c^2x^2-1}x}{c^6} + \frac{15\log(2c^2x+2\sqrt{c^2x^2-1}c)}{c^7}\right)c\right)bd^2 + \frac{1}{3072}\left(384x^8\operatorname{arcosh}(cx) - \left(\frac{48\sqrt{c^2x^2-1}x^7}{c^2} + \frac{56\sqrt{c^2x^2-1}x^5}{c^4} + \frac{70\sqrt{c^2x^2-1}x^3}{c^6} + \frac{105\sqrt{c^2x^2-1}x}{c^8}\right)c\right)bd^2$$

input `integrate(x^3*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`output `1/8*a*e^2*x^8 + 1/3*a*d*e*x^6 + 1/4*a*d^2*x^4 + 1/32*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*d^2 + 1/144*(48*x^6*arccosh(c*x) - (8*sqrt(c^2*x^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c^4 + 15*sqrt(c^2*x^2 - 1)*x/c^6 + 15*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^7)*c)*b*d*e + 1/3072*(384*x^8*arccosh(c*x) - (48*sqrt(c^2*x^2 - 1)*x^7/c^2 + 56*sqrt(c^2*x^2 - 1)*x^5/c^4 + 70*sqrt(c^2*x^2 - 1)*x^3/c^6 + 105*sqrt(c^2*x^2 - 1)*x/c^8 + 105*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^9)*c)*b*e^2`**3.471.8 Giac [F(-2)]**

Exception generated.

$$\int x^3(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`output `Exception raised: TypeError >> an error occurred running a Giac command:IN PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.471.9 Mupad [F(-1)]**

Timed out.

$$\int x^3 (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x^3 (a + b \operatorname{acosh}(cx)) (ex^2 + d)^2 dx$$

input `int(x^3*(a + b*acosh(c*x))*(d + e*x^2)^2,x)`output `int(x^3*(a + b*acosh(c*x))*(d + e*x^2)^2, x)`

### 3.472 $\int x^2(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.472.1 Optimal result . . . . .	3474
3.472.2 Mathematica [A] (verified) . . . . .	3475
3.472.3 Rubi [A] (verified) . . . . .	3475
3.472.4 Maple [A] (verified) . . . . .	3478
3.472.5 Fricas [A] (verification not implemented) . . . . .	3478
3.472.6 Sympy [F] . . . . .	3479
3.472.7 Maxima [A] (verification not implemented) . . . . .	3479
3.472.8 Giac [F(-2)] . . . . .	3480
3.472.9 Mupad [F(-1)] . . . . .	3480

#### 3.472.1 Optimal result

Integrand size = 21, antiderivative size = 260

$$\int x^2(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{b(35c^4d^2 + 42c^2de + 15e^2)(1 - c^2x^2)}{105c^7\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{b(35c^4d^2 + 84c^2de + 45e^2)(1 - c^2x^2)^2}{315c^7\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{be(14c^2d + 15e)(1 - c^2x^2)^3}{175c^7\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{be^2(1 - c^2x^2)^4}{49c^7\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) + \frac{2}{5}dex^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^2x^7(a + \operatorname{barccosh}(cx))$$

```
output 1/3*d^2*x^3*(a+b*arccosh(c*x))+2/5*d*e*x^5*(a+b*arccosh(c*x))+1/7*e^2*x^7*
(a+b*arccosh(c*x))+1/105*b*(35*c^4*d^2+42*c^2*d*e+15*e^2)*(-c^2*x^2+1)/c^7
/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/315*b*(35*c^4*d^2+84*c^2*d*e+45*e^2)*(-c^2*
x^2+1)^2/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/175*b*e*(14*c^2*d+15*e)*(-c^2*x
^2+1)^3/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/49*b*e^2*(-c^2*x^2+1)^4/c^7/(c*x
-1)^(1/2)/(c*x+1)^(1/2)
```

**3.472.2 Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.63

$$\int x^2(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx$$

$$= \frac{105ax^3(35d^2+42dex^2+15e^2x^4) - \frac{b\sqrt{-1+cx}\sqrt{1+cx}(720e^2+24c^2e(98d+15ex^2)+2c^4(1225d^2+588dex^2+135e^2x^4)+c^6(1225d^2x^2+135e^2x^4)+c^6(1225d^2x^2+882d^2ex^2+225e^2x^6))}{c^7} + 105b^2x^3(35d^2+42dex^2+15e^2x^4)\operatorname{ArcCosh}[cx]}{11025}$$

input `Integrate[x^2*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`output `(105*a*x^3*(35*d^2 + 42*d*e*x^2 + 15*e^2*x^4) - (b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(720*e^2 + 24*c^2*e*(98*d + 15*e*x^2) + 2*c^4*(1225*d^2 + 588*d*e*x^2 + 135*e^2*x^4) + c^6*(1225*d^2*x^2 + 882*d^2*e*x^2 + 225*e^2*x^6)))/c^7 + 105*b*x^3*(35*d^2 + 42*d*e*x^2 + 15*e^2*x^4)*ArcCosh[c*x])/11025`**3.472.3 Rubi [A] (verified)**Time = 0.56 (sec) , antiderivative size = 226, normalized size of antiderivative = 0.87, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6373, 27, 1905, 1578, 1195, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx$$

$$\downarrow \text{6373}$$

$$-bc \int \frac{x^3(15e^2x^4+42dex^2+35d^2)}{105\sqrt{cx-1}\sqrt{cx+1}}dx + \frac{1}{3}d^2x^3(a+\operatorname{barccosh}(cx)) + \frac{2}{5}dex^5(a+\operatorname{barccosh}(cx)) + \frac{1}{7}e^2x^7(a+\operatorname{barccosh}(cx))$$

$$\downarrow \text{27}$$

$$-\frac{1}{105}bc \int \frac{x^3(15e^2x^4+42dex^2+35d^2)}{\sqrt{cx-1}\sqrt{cx+1}}dx + \frac{1}{3}d^2x^3(a+\operatorname{barccosh}(cx)) + \frac{2}{5}dex^5(a+\operatorname{barccosh}(cx)) + \frac{1}{7}e^2x^7(a+\operatorname{barccosh}(cx))$$

$$\downarrow \text{1905}$$

$$\begin{aligned}
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^3(15e^2x^4+42dex^2+35d^2)}{\sqrt{c^2x^2-1}} dx}{105\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) + \frac{2}{5}dex^5(a + \operatorname{barccosh}(cx)) + \\
 & \qquad \qquad \qquad \frac{1}{7}e^2x^7(a + \operatorname{barccosh}(cx)) \\
 & \qquad \qquad \qquad \downarrow \text{1578} \\
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^2(15e^2x^4+42dex^2+35d^2)}{\sqrt{c^2x^2-1}} dx^2}{210\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) + \frac{2}{5}dex^5(a + \operatorname{barccosh}(cx)) + \\
 & \qquad \qquad \qquad \frac{1}{7}e^2x^7(a + \operatorname{barccosh}(cx)) \\
 & \qquad \qquad \qquad \downarrow \text{1195} \\
 & \frac{bc\sqrt{c^2x^2-1} \int \left( \frac{15e^2(c^2x^2-1)^{5/2}}{c^6} + \frac{3e(14dc^2+15e)(c^2x^2-1)^{3/2}}{c^6} + \frac{(35d^2c^4+84dec^2+45e^2)\sqrt{c^2x^2-1}}{c^6} + \frac{35d^2c^4+42dec^2+15e^2}{c^6\sqrt{c^2x^2-1}} \right) dx^2}{210\sqrt{cx-1}\sqrt{cx+1}} \\
 & \qquad \qquad \qquad \frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) + \frac{2}{5}dex^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^2x^7(a + \operatorname{barccosh}(cx)) \\
 & \qquad \qquad \qquad \downarrow \text{2009} \\
 & \frac{\frac{1}{3}d^2x^3(a + \operatorname{barccosh}(cx)) + \frac{2}{5}dex^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^2x^7(a + \operatorname{barccosh}(cx)) -}{bc\sqrt{c^2x^2-1} \left( \frac{6e(c^2x^2-1)^{5/2}(14c^2d+15e)}{5c^8} + \frac{30e^2(c^2x^2-1)^{7/2}}{7c^8} + \frac{2(c^2x^2-1)^{3/2}(35c^4d^2+84c^2de+45e^2)}{3c^8} + \frac{2\sqrt{c^2x^2-1}(35c^4d^2+42c^2de+15e^2)}{c^8} \right)}{210\sqrt{cx-1}\sqrt{cx+1}}
 \end{aligned}$$

input `Int[x^2*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `-1/210*(b*c*Sqrt[-1 + c^2*x^2]*((2*(35*c^4*d^2 + 42*c^2*d*e + 15*e^2)*Sqrt[-1 + c^2*x^2])/c^8 + (2*(35*c^4*d^2 + 84*c^2*d*e + 45*e^2)*(-1 + c^2*x^2)^(3/2))/(3*c^8) + (6*e*(14*c^2*d + 15*e)*(-1 + c^2*x^2)^(5/2))/(5*c^8) + (30*e^2*(-1 + c^2*x^2)^(7/2))/(7*c^8))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (d^2*x^3*(a + b*ArcCosh[c*x]))/3 + (2*d*e*x^5*(a + b*ArcCosh[c*x]))/5 + (e^2*x^7*(a + b*ArcCosh[c*x]))/7`

## 3.472.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 1195 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n}, x] && IGtQ[p, 0]`
- rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`
- rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(q_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.472.4 Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.70

method	result
parts	$a\left(\frac{1}{7}e^2x^7 + \frac{2}{5}dex^5 + \frac{1}{3}d^2x^3\right) + \frac{b\left(\frac{c^3 \operatorname{arccosh}(cx)e^2x^7}{7} + \frac{2c^3 \operatorname{arccosh}(cx)dex^5}{5} + \frac{\operatorname{arccosh}(cx)c^3x^3d^2}{3} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{c^3}\right)}{c^3}$
derivativedivides	$\frac{a\left(\frac{1}{3}c^7d^2x^3 + \frac{2}{5}dc^7ex^5 + \frac{1}{7}e^2c^7x^7\right)}{c^4} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^7d^2x^3}{3} + \frac{2 \operatorname{arccosh}(cx)dc^7ex^5}{5} + \frac{\operatorname{arccosh}(cx)e^2c^7x^7}{7} - \frac{\sqrt{cx-1}\sqrt{cx+1}(225c^6e^2)}{c^3}\right)}{c^3}$
default	$\frac{a\left(\frac{1}{3}c^7d^2x^3 + \frac{2}{5}dc^7ex^5 + \frac{1}{7}e^2c^7x^7\right)}{c^4} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^7d^2x^3}{3} + \frac{2 \operatorname{arccosh}(cx)dc^7ex^5}{5} + \frac{\operatorname{arccosh}(cx)e^2c^7x^7}{7} - \frac{\sqrt{cx-1}\sqrt{cx+1}(225c^6e^2)}{c^3}\right)}{c^3}$

input `int(x^2*(e*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/7*e^2*x^7+2/5*d*e*x^5+1/3*d^2*x^3)+b/c^3*(1/7*c^3*arccosh(c*x)*e^2*x^7+2/5*c^3*arccosh(c*x)*d*e*x^5+1/3*arccosh(c*x)*c^3*x^3*d^2-1/11025/c^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(225*c^6*e^2*x^6+882*c^6*d*e*x^4+1225*c^6*d^2*x^2+270*c^4*e^2*x^4+1176*c^4*d*e*x^2+2450*c^4*d^2+360*c^2*e^2*x^2+2352*c^2*d*e+720*e^2))`

### 3.472.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.76

$$\int x^2(d + ex^2)^2(a + b\operatorname{arccosh}(cx)) dx$$

$$= \frac{1575 ac^7e^2x^7 + 4410 ac^7dex^5 + 3675 ac^7d^2x^3 + 105(15 bc^7e^2x^7 + 42 bc^7dex^5 + 35 bc^7d^2x^3) \log(cx + \sqrt{c^2x^2 - 1})}{c^7}$$

input `integrate(x^2*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `1/11025*(1575*a*c^7*e^2*x^7 + 4410*a*c^7*d*e*x^5 + 3675*a*c^7*d^2*x^3 + 105*(15*b*c^7*e^2*x^7 + 42*b*c^7*d*e*x^5 + 35*b*c^7*d^2*x^3)*log(c*x + sqrt(c^2*x^2 - 1)) - (225*b*c^6*e^2*x^6 + 2450*b*c^4*d^2 + 2352*b*c^2*d*e + 18*(49*b*c^6*d*e + 15*b*c^4*e^2)*x^4 + 720*b*e^2 + (1225*b*c^6*d^2 + 1176*b*c^4*d*e + 360*b*c^2*e^2)*x^2)*sqrt(c^2*x^2 - 1)/c^7`

**3.472.6 Sympy [F]**

$$\int x^2(d + ex^2)^2(a + \operatorname{barccosh}(cx)) dx = \int x^2(a + b \operatorname{acosh}(cx))(d + ex^2)^2 dx$$

input `integrate(x**2*(e*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `Integral(x**2*(a + b*acosh(c*x))*(d + e*x**2)**2, x)`

**3.472.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 247, normalized size of antiderivative = 0.95

$$\begin{aligned} & \int x^2(d + ex^2)^2(a + \operatorname{barccosh}(cx)) dx \\ &= \frac{1}{7} ae^2x^7 + \frac{2}{5} adex^5 + \frac{1}{3} ad^2x^3 + \frac{1}{9} \left( 3x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \right) bd^2 \\ &+ \frac{2}{75} \left( 15x^5 \operatorname{arcosh}(cx) - \left( \frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6} \right) c \right) bde \\ &+ \frac{1}{245} \left( 35x^7 \operatorname{arcosh}(cx) - \left( \frac{5\sqrt{c^2x^2 - 1}x^6}{c^2} + \frac{6\sqrt{c^2x^2 - 1}x^4}{c^4} + \frac{8\sqrt{c^2x^2 - 1}x^2}{c^6} + \frac{16\sqrt{c^2x^2 - 1}}{c^8} \right) c \right) be^2 \end{aligned}$$

input `integrate(x^2*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/7*a*e^2*x^7 + 2/5*a*d*e*x^5 + 1/3*a*d^2*x^3 + 1/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*d^2 + 2/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d*e + 1/245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*e^2`



**3.472.8 Giac [F(-2)]**

Exception generated.

$$\int x^2(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^2*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

**3.472.9 Mupad [F(-1)]**

Timed out.

$$\int x^2(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) (ex^2 + d)^2 dx$$

input `int(x^2*(a + b*acosh(c*x))*(d + e*x^2)^2,x)`

output `int(x^2*(a + b*acosh(c*x))*(d + e*x^2)^2, x)`

### 3.473 $\int x(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.473.1 Optimal result . . . . .	3481
3.473.2 Mathematica [A] (warning: unable to verify) . . . . .	3482
3.473.3 Rubi [A] (verified) . . . . .	3482
3.473.4 Maple [A] (verified) . . . . .	3485
3.473.5 Fricas [A] (verification not implemented) . . . . .	3486
3.473.6 Sympy [F] . . . . .	3486
3.473.7 Maxima [A] (verification not implemented) . . . . .	3487
3.473.8 Giac [F(-2)] . . . . .	3487
3.473.9 Mupad [F(-1)] . . . . .	3488

#### 3.473.1 Optimal result

Integrand size = 19, antiderivative size = 269

$$\begin{aligned} & \int x(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx \\ &= \frac{b(44c^4d^2 + 44c^2de + 15e^2)x(1 - c^2x^2)}{288c^5\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{5b(2c^2d + e)x(1 - c^2x^2)(d + ex^2)}{144c^3\sqrt{-1 + cx}\sqrt{1 + cx}} \\ &+ \frac{bx(1 - c^2x^2)(d + ex^2)^2}{36c\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{(d + ex^2)^3(a + \operatorname{barccosh}(cx))}{6e} \\ &- \frac{b(2c^2d + e)(8c^4d^2 + 8c^2de + 5e^2)\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{cx}{\sqrt{-1 + c^2x^2}}\right)}{96c^6e\sqrt{-1 + cx}\sqrt{1 + cx}} \end{aligned}$$

output  $\frac{1}{6}(ex^2+d)^3(a+b\operatorname{arccosh}(cx))/e+1/288*b*(44*c^4*d^2+44*c^2*d*e+15*e^2)*x*(-c^2*x^2+1)/c^5/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+5/144*b*(2*c^2*d+e)*x*(-c^2*x^2+1)*(e*x^2+d)/c^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/36*b*x*(-c^2*x^2+1)*(e*x^2+d)^2/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/96*b*(2*c^2*d+e)*(8*c^4*d^2+8*c^2*d*e+5*e^2)*\operatorname{arctanh}(c*x/(c^2*x^2-1)^{(1/2)})*(c^2*x^2-1)^{(1/2)}/c^6/e/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

**3.473.2 Mathematica [A] (warning: unable to verify)**

Time = 0.19 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.68

$$\int x(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{cx(48ac^5x(3d^2 + 3dex^2 + e^2x^4) - b\sqrt{-1 + cx}\sqrt{1 + cx}(15e^2 + 2c^2e(27d + 5ex^2) + 4c^4(18d^2 + 9dex^2 + 2e^2x^4)) - 48b^2c^6x^2(3d^2 + 3dex^2 + e^2x^4)\operatorname{ArcCosh}[cx] - 6b^2(24c^4d^2 + 18c^2de + 5e^2)\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1 + cx)/(1 + cx)])]}{288c^6}$$

input `Integrate[x*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`output `(c*x*(48*a*c^5*x*(3*d^2 + 3*d*e*x^2 + e^2*x^4) - b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(15*e^2 + 2*c^2*e*(27*d + 5*e*x^2) + 4*c^4*(18*d^2 + 9*d*e*x^2 + 2*e^2*x^4))) + 48*b*c^6*x^2*(3*d^2 + 3*d*e*x^2 + e^2*x^4)*ArcCosh[c*x] - 6*b*(24*c^4*d^2 + 18*c^2*d*e + 5*e^2)*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]])/(288*c^6)`**3.473.3 Rubi [A] (verified)**Time = 0.45 (sec) , antiderivative size = 246, normalized size of antiderivative = 0.91, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.368$ , Rules used = {6372, 648, 318, 403, 299, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6372$$

$$\frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{6e} - \frac{bc \int \frac{(ex^2+d)^3}{\sqrt{cx-1}\sqrt{cx+1}} dx}{6e}$$

$$\downarrow 648$$

$$\frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{6e} - \frac{bc\sqrt{c^2x^2 - 1} \int \frac{(ex^2+d)^3}{\sqrt{c^2x^2-1}} dx}{6e\sqrt{cx-1}\sqrt{cx+1}}$$

$$\downarrow 318$$

$$\begin{aligned}
 & \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\int \frac{(ex^2 + d)(5e(2dc^2 + e)x^2 + d(6dc^2 + e))}{\sqrt{c^2x^2 - 1}} dx}{6c^2} + \frac{ex\sqrt{c^2x^2 - 1}(d + ex^2)^2}{6c^2} \right)} \\
 & \qquad \qquad \qquad \frac{6e\sqrt{cx - 1}\sqrt{cx + 1}}{\downarrow 403} \\
 & \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\int \frac{e(44d^2c^4 + 44dec^2 + 15e^2)x^2 + d(24d^2c^4 + 14dec^2 + 5e^2)}{\sqrt{c^2x^2 - 1}} dx}{4c^2} + \frac{5ex\sqrt{c^2x^2 - 1}(2c^2d + e)(d + ex^2)}{4c^2} + \frac{ex\sqrt{c^2x^2 - 1}(d + ex^2)^2}{6c^2} \right)} \\
 & \qquad \qquad \qquad \frac{6e\sqrt{cx - 1}\sqrt{cx + 1}}{\downarrow 299} \\
 & \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\frac{3(2c^2d + e)(8c^4d^2 + 8c^2de + 5e^2) \int \frac{1}{\sqrt{c^2x^2 - 1}} dx}{2c^2} + \frac{ex\sqrt{c^2x^2 - 1}(44c^4d^2 + 44c^2de + 15e^2)}{4c^2}}{6c^2} + \frac{5ex\sqrt{c^2x^2 - 1}(2c^2d + e)(d + ex^2)}{4c^2} + \frac{ex\sqrt{c^2x^2 - 1}}{6c} \right)} \\
 & \qquad \qquad \qquad \frac{6e\sqrt{cx - 1}\sqrt{cx + 1}}{\downarrow 224} \\
 & \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\frac{3(2c^2d + e)(8c^4d^2 + 8c^2de + 5e^2) \int \frac{1}{1 - \frac{cx}{c^2x^2 - 1}} d \frac{x}{\sqrt{c^2x^2 - 1}}}{2c^2} + \frac{ex\sqrt{c^2x^2 - 1}(44c^4d^2 + 44c^2de + 15e^2)}{4c^2}}{6c^2} + \frac{5ex\sqrt{c^2x^2 - 1}(2c^2d + e)(d + ex^2)}{4c^2} + \frac{ex\sqrt{c^2x^2 - 1}}{6c} \right)} \\
 & \qquad \qquad \qquad \frac{6e\sqrt{cx - 1}\sqrt{cx + 1}}{\downarrow 219} \\
 & \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\frac{3 \operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2 - 1}}\right)(2c^2d + e)(8c^4d^2 + 8c^2de + 5e^2)}{2c^3} + \frac{ex\sqrt{c^2x^2 - 1}(44c^4d^2 + 44c^2de + 15e^2)}{4c^2}}{6c^2} + \frac{5ex\sqrt{c^2x^2 - 1}(2c^2d + e)(d + ex^2)}{4c^2} + \frac{ex\sqrt{c^2x^2 - 1}}{6c} \right)} \\
 & \qquad \qquad \qquad \frac{6e\sqrt{cx - 1}\sqrt{cx + 1}}{\downarrow 219}
 \end{aligned}$$

3.473.  $\int x(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

input `Int[x*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/(6*e) - (b*c*Sqrt[-1 + c^2*x^2]*((e*x*Sqrt[-1 + c^2*x^2]*(d + e*x^2)^2)/(6*c^2) + ((5*e*(2*c^2*d + e)*x*Sqrt[-1 + c^2*x^2]*(d + e*x^2))/(4*c^2) + ((e*(44*c^4*d^2 + 44*c^2*d*e + 15*e^2)*x*Sqrt[-1 + c^2*x^2])/(2*c^2) + (3*(2*c^2*d + e)*(8*c^4*d^2 + 8*c^2*d*e + 5*e^2)*ArcTanh[(c*x)/Sqrt[-1 + c^2*x^2]])/(2*c^3))/(4*c^2))/(6*e*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.473.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 318 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[d*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1)/(b*(2*(p + q) + 1))), x] + Simp[1/(b*(2*(p + q) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*(b*c*(2*(p + q) + 1) - a*d) + d*(b*c*(2*(p + 2*q - 1) + 1) - a*d*(2*(q - 1) + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[2*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, 2, p, q, x]`

rule 403 `Int[((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2), x_Symbol] := Simp[f*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(b*(2*(p + q + 1) + 1))), x] + Simp[1/(b*(2*(p + q + 1) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[c*(b*e - a*f + b*e*2*(p + q + 1)) + (d*(b*e - a*f) + f*2*q*(b*c - a*d) + b*d*e*2*(p + q + 1))*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && GtQ[q, 0] && NeQ[2*(p + q + 1) + 1, 0]`

rule 648 `Int[((c_) + (d_)*(x_)^(m_))*((e_) + (f_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c + d*x)^FracPart[m]*((e + f*x)^FracPart[m]/(c*e + d*f*x^2)^FracPart[m]) Int[(c*e + d*f*x^2)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m, n] && EqQ[d*e + c*f, 0] && !(EqQ[p, 2] && LtQ[m, -1])`

rule 6372 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])/(2*e*(p + 1))), x] - Simp[b*(c/(2*e*(p + 1))) Int[(d + e*x^2)^(p + 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[c^2*d + e, 0] && NeQ[p, -1]`

### 3.473.4 Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.26

method	result
parts	$\frac{a(e x^2+d)^3}{6e} + \frac{b \left( \frac{c^2 e^2 \operatorname{arccosh}(c x) x^6}{6} + \frac{c^2 e \operatorname{arccosh}(c x) x^4 d}{2} + \frac{\operatorname{arccosh}(c x) c^2 x^2 d^2}{2} + \frac{c^2 \operatorname{arccosh}(c x) d^3}{6e} - \frac{\sqrt{c x-1} \sqrt{c x+1} (48 c^6 d^3 \ln)}{\dots} \right)}{\dots}$
derivativedivides	$\frac{a(c^2 e x^2+c^2 d)^3}{6e^4 e} + \frac{b \left( \frac{\operatorname{arccosh}(c x) c^6 d^3}{6e} + \frac{\operatorname{arccosh}(c x) c^6 d^2 x^2}{2} + \frac{e \operatorname{arccosh}(c x) c^6 d x^4}{2} + \frac{e^2 \operatorname{arccosh}(c x) c^6 x^6}{6} - \frac{\sqrt{c x-1} \sqrt{c x+1} (48 c^6 d^3 \ln)}{\dots} \right)}{\dots}$
default	$\frac{a(c^2 e x^2+c^2 d)^3}{6e^4 e} + \frac{b \left( \frac{\operatorname{arccosh}(c x) c^6 d^3}{6e} + \frac{\operatorname{arccosh}(c x) c^6 d^2 x^2}{2} + \frac{e \operatorname{arccosh}(c x) c^6 d x^4}{2} + \frac{e^2 \operatorname{arccosh}(c x) c^6 x^6}{6} - \frac{\sqrt{c x-1} \sqrt{c x+1} (48 c^6 d^3 \ln)}{\dots} \right)}{\dots}$

input `int(x*(e*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output  $\frac{1}{6}a(e^2x+d)^3/e+b/c^2(1/6c^2e^2\operatorname{arccosh}(cx)x^6+1/2c^2e\operatorname{arccosh}(cx)x^4d+1/2\operatorname{arccosh}(cx)c^2x^2d^2+1/6c^2/e\operatorname{arccosh}(cx)d^3-1/288/c^4/e(c^2x-1)^{1/2}(c^2x+1)^{1/2}(48c^6d^3\ln(c^2x+(c^2x^2-1)^{1/2}))+72c^5d^2e^2x(c^2x^2-1)^{1/2}+36c^5de^2(c^2x^2-1)^{1/2}x^3+8e^3(c^2x^2-1)^{1/2}c^5x^5+72c^4d^2e\ln(c^2x+(c^2x^2-1)^{1/2}))+54c^3de^2x(c^2x^2-1)^{1/2}+10e^3c^3x^3(c^2x^2-1)^{1/2}+54c^2de^2\ln(c^2x+(c^2x^2-1)^{1/2}))+15e^3cx(c^2x^2-1)^{1/2}+15e^3\ln(c^2x+(c^2x^2-1)^{1/2}))/c^2x^2-1)^{1/2})$

### 3.473.5 Fricas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.72

$$\int x(d+ex^2)^2(a+b\operatorname{arccosh}(cx))dx$$

$$= \frac{48ac^6e^2x^6 + 144ac^6dex^4 + 144ac^6d^2x^2 + 3(16bc^6e^2x^6 + 48bc^6dex^4 + 48bc^6d^2x^2 - 24bc^4d^2 - 18bc^2de - 18bc^2de - 5b^2c^4d^2 - 18b^2c^2de - 5b^2e^2)\log(cx + \sqrt{c^2x^2 - 1}) - (8b^2c^5e^2x^5 + 2(18b^2c^5de + 5b^2c^3e^2)x^3 + 3(24b^2c^5d^2 + 18b^2c^3de + 5b^2ce^2)x)\sqrt{c^2x^2 - 1}}{c^6}$$

input `integrate(x*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $\frac{1}{288}(48ac^6e^2x^6 + 144ac^6dex^4 + 144ac^6d^2x^2 + 3(16bc^6e^2x^6 + 48bc^6dex^4 + 48bc^6d^2x^2 - 24bc^4d^2 - 18bc^2de - 5b^2c^4d^2 - 18b^2c^2de - 5b^2e^2)\log(cx + \sqrt{c^2x^2 - 1}) - (8b^2c^5e^2x^5 + 2(18b^2c^5de + 5b^2c^3e^2)x^3 + 3(24b^2c^5d^2 + 18b^2c^3de + 5b^2ce^2)x)\sqrt{c^2x^2 - 1})/c^6$

### 3.473.6 Sympy [F]

$$\int x(d+ex^2)^2(a+b\operatorname{arccosh}(cx))dx = \int x(a+b\operatorname{acosh}(cx))(d+ex^2)^2dx$$

input `integrate(x*(e*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `Integral(x*(a + b*acosh(c*x))*(d + e*x**2)**2, x)`

**3.473.7 Maxima [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.01

$$\int x(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx = \frac{1}{6}ae^2x^6 + \frac{1}{2}adex^4 + \frac{1}{2}ad^2x^2 + \frac{1}{4}\left(2x^2\operatorname{arcosh}(cx) - c\left(\frac{\sqrt{c^2x^2-1}x}{c^2} + \frac{\log(2c^2x+2\sqrt{c^2x^2-1}c)}{c^3}\right)\right)bd^2 + \frac{1}{16}\left(8x^4\operatorname{arcosh}(cx) - \left(\frac{2\sqrt{c^2x^2-1}x^3}{c^2} + \frac{3\sqrt{c^2x^2-1}x}{c^4} + \frac{3\log(2c^2x+2\sqrt{c^2x^2-1}c)}{c^5}\right)c\right)bde + \frac{1}{288}\left(48x^6\operatorname{arcosh}(cx) - \left(\frac{8\sqrt{c^2x^2-1}x^5}{c^2} + \frac{10\sqrt{c^2x^2-1}x^3}{c^4} + \frac{15\sqrt{c^2x^2-1}x}{c^6} + \frac{15\log(2c^2x+2\sqrt{c^2x^2-1}c)}{c^7}\right)c\right)bde$$

```
input integrate(x*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")
```

```
output 1/6*a*e^2*x^6 + 1/2*a*d*e*x^4 + 1/2*a*d^2*x^2 + 1/4*(2*x^2*arccosh(c*x) -
c*(sqrt(c^2*x^2 - 1)*x/c^2 + log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^3)*b*
d^2 + 1/16*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2
*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*d*e + 1
/288*(48*x^6*arccosh(c*x) - (8*sqrt(c^2*x^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2
- 1)*x^3/c^4 + 15*sqrt(c^2*x^2 - 1)*x/c^6 + 15*log(2*c^2*x + 2*sqrt(c^2*x
^2 - 1)*c)/c^7)*c)*b*e^2
```

**3.473.8 Giac [F(-2)]**

Exception generated.

$$\int x(d+ex^2)^2(a+\operatorname{barccosh}(cx))dx = \text{Exception raised: RuntimeError}$$

```
input integrate(x*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve
cteur & l) Error: Bad Argument Value
```



**3.473.9 Mupad [F(-1)]**

Timed out.

$$\int x(d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (ex^2 + d)^2 dx$$

input `int(x*(a + b*acosh(c*x))*(d + e*x^2)^2,x)`output `int(x*(a + b*acosh(c*x))*(d + e*x^2)^2, x)`

### 3.474 $\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.474.1 Optimal result . . . . .	3489
3.474.2 Mathematica [A] (verified) . . . . .	3490
3.474.3 Rubi [A] (verified) . . . . .	3490
3.474.4 Maple [A] (verified) . . . . .	3493
3.474.5 Fricas [A] (verification not implemented) . . . . .	3493
3.474.6 Sympy [F] . . . . .	3494
3.474.7 Maxima [A] (verification not implemented) . . . . .	3494
3.474.8 Giac [F(-2)] . . . . .	3495
3.474.9 Mupad [F(-1)] . . . . .	3495

#### 3.474.1 Optimal result

Integrand size = 18, antiderivative size = 196

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \frac{b(15c^4d^2 + 10c^2de + 3e^2)(1 - c^2x^2)}{15c^5\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{2be(5c^2d + 3e)(1 - c^2x^2)^2}{45c^5\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{be^2(1 - c^2x^2)^3}{25c^5\sqrt{-1 + cx}\sqrt{1 + cx}} + d^2x(a + \operatorname{barccosh}(cx)) + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx))$$

```
output d^2*x*(a+b*arccosh(c*x))+2/3*d*e*x^3*(a+b*arccosh(c*x))+1/5*e^2*x^5*(a+b*arccosh(c*x))+1/15*b*(15*c^4*d^2+10*c^2*d*e+3*e^2)*(-c^2*x^2+1)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)-2/45*b*e*(5*c^2*d+3*e)*(-c^2*x^2+1)^2/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/25*b*e^2*(-c^2*x^2+1)^3/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.474.2 Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.66

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{1}{225} \left( \frac{15ax(15d^2 + 10dex^2 + 3e^2x^4) + b\sqrt{-1 + cx}\sqrt{1 + cx}(24e^2 + 4c^2e(25d + 3ex^2) + c^4(225d^2 + 50dex^2 + 9e^2x^4))}{c^5} + 15bx(15d^2 + 10dex^2 + 3e^2x^4) \operatorname{arccosh}(cx) \right)$$

input `Integrate[(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`output `(15*a*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4) - (b*sqrt[-1 + c*x]*sqrt[1 + c*x])*(24*e^2 + 4*c^2*e*(25*d + 3*e*x^2) + c^4*(225*d^2 + 50*d*e*x^2 + 9*e^2*x^4)))/c^5 + 15*b*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4)*ArcCosh[c*x])/225`**3.474.3 Rubi [A] (verified)**Time = 0.47 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.92, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6323, 27, 1905, 1576, 1140, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6323}$$

$$-bc \int \frac{x(3e^2x^4 + 10dex^2 + 15d^2)}{15\sqrt{cx - 1}\sqrt{cx + 1}} dx + d^2x(a + \operatorname{barccosh}(cx)) + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{27}$$

$$-\frac{1}{15}bc \int \frac{x(3e^2x^4 + 10dex^2 + 15d^2)}{\sqrt{cx - 1}\sqrt{cx + 1}} dx + d^2x(a + \operatorname{barccosh}(cx)) + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx))$$

$$\begin{aligned}
 & \downarrow 1905 \\
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{x(3e^2x^4+10dex^2+15d^2)}{\sqrt{c^2x^2-1}} dx}{15\sqrt{cx-1}\sqrt{cx+1}} + d^2x(a + \operatorname{barccosh}(cx)) + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx)) + \\
 & \quad \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx)) \\
 & \downarrow 1576 \\
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{3e^2x^4+10dex^2+15d^2}{\sqrt{c^2x^2-1}} dx^2}{30\sqrt{cx-1}\sqrt{cx+1}} + d^2x(a + \operatorname{barccosh}(cx)) + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx)) + \\
 & \quad \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx)) \\
 & \downarrow 1140 \\
 & -\frac{bc\sqrt{c^2x^2-1} \int \left( \frac{3(c^2x^2-1)^{3/2}e^2}{c^4} + \frac{2(5dc^2+3e)\sqrt{c^2x^2-1}e}{c^4} + \frac{15d^2c^4+10dec^2+3e^2}{c^4\sqrt{c^2x^2-1}} \right) dx^2}{30\sqrt{cx-1}\sqrt{cx+1}} + d^2x(a + \\
 & \quad \operatorname{barccosh}(cx)) + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx)) \\
 & \downarrow 2009 \\
 & \frac{d^2x(a + \operatorname{barccosh}(cx)) + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx)) -}{30\sqrt{cx-1}\sqrt{cx+1}} \\
 & \quad bc\sqrt{c^2x^2-1} \left( \frac{4e(c^2x^2-1)^{3/2}(5c^2d+3e)}{3c^6} + \frac{6e^2(c^2x^2-1)^{5/2}}{5c^6} + \frac{2\sqrt{c^2x^2-1}(15c^4d^2+10c^2de+3e^2)}{c^6} \right)
 \end{aligned}$$

input `Int[(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `-1/30*(b*c*Sqrt[-1 + c^2*x^2]*((2*(15*c^4*d^2 + 10*c^2*d*e + 3*e^2)*Sqrt[-1 + c^2*x^2])/c^6 + (4*e*(5*c^2*d + 3*e)*(-1 + c^2*x^2)^(3/2))/(3*c^6) + (6*e^2*(-1 + c^2*x^2)^(5/2))/(5*c^6)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + d^2*x*(a + b*ArcCosh[c*x]) + (2*d*e*x^3*(a + b*ArcCosh[c*x]))/3 + (e^2*x^5*(a + b*ArcCosh[c*x]))/5`

## 3.474.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 1140 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[p, 0]`
- rule 1576 `Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]`
- rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6323 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d + e, 0] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])`

### 3.474.4 Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.70

method	result
parts	$a\left(\frac{1}{5}e^2x^5 + \frac{2}{3}dex^3 + d^2x\right) + \frac{b\left(\frac{c \operatorname{arccosh}(cx)e^2x^5}{5} + \frac{2c \operatorname{arccosh}(cx)dex^3}{3} + \operatorname{arccosh}(cx)cx d^2 - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4e^2x^4 + 50c^4e^2x^2 + 10c^4e^2)}{c}\right)}{c}$
derivativedivides	$\frac{a\left(\frac{d^2c^5x + \frac{2}{3}dc^5ex^3 + \frac{1}{5}e^2c^5x^5}{c^4}\right) + \frac{b\left(\operatorname{arccosh}(cx)d^2c^5x + \frac{2 \operatorname{arccosh}(cx)dc^5ex^3}{3} + \frac{\operatorname{arccosh}(cx)e^2c^5x^5}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4e^2x^4 + 50c^4e^2x^2 + 10c^4e^2)}{c}\right)}{c^4}}{c}$
default	$\frac{a\left(\frac{d^2c^5x + \frac{2}{3}dc^5ex^3 + \frac{1}{5}e^2c^5x^5}{c^4}\right) + \frac{b\left(\operatorname{arccosh}(cx)d^2c^5x + \frac{2 \operatorname{arccosh}(cx)dc^5ex^3}{3} + \frac{\operatorname{arccosh}(cx)e^2c^5x^5}{5} - \frac{\sqrt{cx-1}\sqrt{cx+1}(9c^4e^2x^4 + 50c^4e^2x^2 + 10c^4e^2)}{c}\right)}{c^4}}{c}$

input `int((e*x^2+d)^2*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/5*e^2*x^5+2/3*d*e*x^3+d^2*x)+b/c*(1/5*c*arccosh(c*x)*e^2*x^5+2/3*c*arccosh(c*x)*d*e*x^3+arccosh(c*x)*c*x*d^2-1/225/c^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(9*c^4*e^2*x^4+50*c^4*d*e*x^2+225*c^4*d^2+12*c^2*e^2*x^2+100*c^2*d*e+24*e^2))`

### 3.474.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.83

$$\int (d + ex^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \frac{45 ac^5 e^2 x^5 + 150 ac^5 dex^3 + 225 ac^5 d^2 x + 15 (3 bc^5 e^2 x^5 + 10 bc^5 dex^3 + 15 bc^5 d^2 x) \log (cx + \sqrt{c^2 x^2 - 1}) - 225 c^5}{225 c^5}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `1/225*(45*a*c^5*e^2*x^5 + 150*a*c^5*d*e*x^3 + 225*a*c^5*d^2*x + 15*(3*b*c^5*e^2*x^5 + 10*b*c^5*d*e*x^3 + 15*b*c^5*d^2*x)*log(c*x + sqrt(c^2*x^2 - 1)) - (9*b*c^4*e^2*x^4 + 225*b*c^4*d^2 + 100*b*c^2*d*e + 24*b*e^2 + 2*(25*b*c^4*d*e + 6*b*c^2*e^2)*x^2)*sqrt(c^2*x^2 - 1))/c^5`

**3.474.6 Sympy [F]**

$$\int (d + ex^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d + ex^2)^2 dx$$

input `integrate((e*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**2, x)`

**3.474.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 180, normalized size of antiderivative = 0.92

$$\begin{aligned} & \int (d + ex^2)^2 (a + b \operatorname{arccosh}(cx)) dx \\ &= \frac{1}{5} ae^2 x^5 + \frac{2}{3} adex^3 + \frac{2}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bde \\ &+ \frac{1}{75} \left( 15x^5 \operatorname{arccosh}(cx) - \left( \frac{3\sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4\sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8\sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) be^2 \\ &+ ad^2 x + \frac{(cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) bd^2}{c} \end{aligned}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/5*a*e^2*x^5 + 2/3*a*d*e*x^3 + 2/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*d*e + 1/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*e^2 + a*d^2*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*d^2/c`

**3.474.8 Giac [F(-2)]**

Exception generated.

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

**3.474.9 Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (ex^2 + d)^2 dx$$

input `int((a + b*acosh(c*x))*(d + e*x^2)^2,x)`

output `int((a + b*acosh(c*x))*(d + e*x^2)^2, x)`



**3.475**  $\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x} dx$

3.475.1 Optimal result . . . . . 3496  
 3.475.2 Mathematica [A] (warning: unable to verify) . . . . . 3497  
 3.475.3 Rubi [A] (verified) . . . . . 3498  
 3.475.4 Maple [A] (verified) . . . . . 3500  
 3.475.5 Fricas [F] . . . . . 3500  
 3.475.6 Sympy [F] . . . . . 3501  
 3.475.7 Maxima [F] . . . . . 3501  
 3.475.8 Giac [F] . . . . . 3501  
 3.475.9 Mupad [F(-1)] . . . . . 3502

**3.475.1 Optimal result**

Integrand size = 21, antiderivative size = 342

$$\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x} dx = -\frac{be(16c^2d+3e)x\sqrt{-1+cx}\sqrt{1+cx}}{32c^3} - \frac{be^2x^3\sqrt{-1+cx}\sqrt{1+cx}}{16c} - \frac{be(16c^2d+3e)\operatorname{arccosh}(cx)}{32c^4} + dex^2(a+b\operatorname{arccosh}(cx)) + \frac{1}{4}e^2x^4(a+b\operatorname{arccosh}(cx)) - \frac{ibd^2\sqrt{1-c^2x^2}\arcsin(cx)^2}{2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{bd^2\sqrt{1-c^2x^2}\arcsin(cx)\log(1-e^{2i\arcsin(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} + d^2(a+b\operatorname{arccosh}(cx))\log(x) - \frac{bd^2\sqrt{1-c^2x^2}\arcsin(cx)\log(x)}{\sqrt{-1+cx}\sqrt{1+cx}} - \frac{ibd^2\sqrt{1-c^2x^2}\operatorname{PolyLog}(2,e^{2i\arcsin(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}}$$

output 
$$\begin{aligned} & -1/32*b*e*(16*c^2*d+3*e)*\operatorname{arccosh}(c*x)/c^4+d*e*x^2*(a+b*\operatorname{arccosh}(c*x))+1/4*e \\ & ^2*x^4*(a+b*\operatorname{arccosh}(c*x))+d^2*(a+b*\operatorname{arccosh}(c*x))*\ln(x)-1/32*b*e*(16*c^2*d+ \\ & 3*e)*x*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3-1/16*b*e^2*x^3*(c*x-1)^{(1/2)}*(c*x+1) \\ & )^{(1/2)}/c-1/2*I*b*d^2*\operatorname{arcsin}(c*x)^2*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+ \\ & 1)^{(1/2)}+b*d^2*\operatorname{arcsin}(c*x)*\ln(1-(I*c*x+(-c^2*x^2+1)^{(1/2)})^2*(-c^2*x^2+1) \\ & )^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-b*d^2*\operatorname{arcsin}(c*x)*\ln(x)*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \\ & -1/2*I*b*d^2*\operatorname{polylog}(2,(I*c*x+(-c^2*x^2+1)^{(1/2)})^2*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}) \end{aligned}$$

### 3.475.2 Mathematica [A] (warning: unable to verify)

Time = 0.39 (sec) , antiderivative size = 228, normalized size of antiderivative = 0.67

$$\begin{aligned} & \int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x} dx \\ & = adex^2 + \frac{1}{4}ae^2x^4 + bdex^2\operatorname{arccosh}(cx) + \frac{1}{4}be^2x^4\operatorname{arccosh}(cx) \\ & \quad - \frac{bde\left(cx\sqrt{-1+cx}\sqrt{1+cx} + 2\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right)}{2c^2} \\ & \quad - \frac{be^2\left(cx\sqrt{\frac{-1+cx}{1+cx}}(3+3cx+2c^2x^2+2c^3x^3) + 6\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right)}{32c^4} \\ & \quad + \frac{1}{2}bd^2\operatorname{arccosh}(cx)\left(\operatorname{arccosh}(cx) + 2\log\left(1+e^{-2\operatorname{arccosh}(cx)}\right)\right) \\ & \quad + ad^2\log(x) - \frac{1}{2}bd^2\operatorname{PolyLog}\left(2,-e^{-2\operatorname{arccosh}(cx)}\right) \end{aligned}$$

input `Integrate[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x,x]`

output 
$$\begin{aligned} & a*d*e*x^2 + (a*e^2*x^4)/4 + b*d*e*x^2*\operatorname{ArcCosh}[c*x] + (b*e^2*x^4*\operatorname{ArcCosh}[c* \\ & x])/4 - (b*d*e*(c*x*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x] + 2*\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1 + c* \\ & x)/(1 + c*x)]])/((2*c^2) - (b*e^2*(c*x*\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(3 + 3*c \\ & *x + 2*c^2*x^2 + 2*c^3*x^3) + 6*\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]]))/(32* \\ & c^4) + (b*d^2*\operatorname{ArcCosh}[c*x]*(\operatorname{ArcCosh}[c*x] + 2*\operatorname{Log}[1 + E^(-2*\operatorname{ArcCosh}[c*x])])) \\ & )/2 + a*d^2*\operatorname{Log}[x] - (b*d^2*\operatorname{PolyLog}[2, -E^(-2*\operatorname{ArcCosh}[c*x])])/2 \end{aligned}$$

**3.475.3 Rubi [A] (verified)**

Time = 1.08 (sec) , antiderivative size = 372, normalized size of antiderivative = 1.09, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx$$

↓ 6373

$$-bc \int \frac{e^2 x^4 + 4dex^2 + 4d^2 \log(x)}{4\sqrt{cx-1}\sqrt{cx+1}} dx + d^2 \log(x)(a + \operatorname{barccosh}(cx)) + dex^2(a + \operatorname{barccosh}(cx)) + \frac{1}{4}e^2 x^4(a + \operatorname{barccosh}(cx))$$

↓ 27

$$-\frac{1}{4}bc \int \frac{e^2 x^4 + 4dex^2 + 4d^2 \log(x)}{\sqrt{cx-1}\sqrt{cx+1}} dx + d^2 \log(x)(a + \operatorname{barccosh}(cx)) + dex^2(a + \operatorname{barccosh}(cx)) + \frac{1}{4}e^2 x^4(a + \operatorname{barccosh}(cx))$$

↓ 7293

$$-\frac{1}{4}bc \int \left( \frac{e^2 x^4}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{4dex^2}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{4d^2 \log(x)}{\sqrt{cx-1}\sqrt{cx+1}} \right) dx + d^2 \log(x)(a + \operatorname{barccosh}(cx)) + dex^2(a + \operatorname{barccosh}(cx)) + \frac{1}{4}e^2 x^4(a + \operatorname{barccosh}(cx))$$

↓ 2009

$$\frac{1}{4}bc \left( \frac{3e^2 \operatorname{arccosh}(cx)}{8c^5} + \frac{2d \operatorname{arccosh}(cx)}{c^3} + \frac{2id^2 \sqrt{1-c^2x^2} \operatorname{PolyLog}(2, e^{2i \operatorname{arcsin}(cx)})}{c\sqrt{cx-1}\sqrt{cx+1}} + \frac{2id^2 \sqrt{1-c^2x^2} \operatorname{arcsin}(cx)^2}{c\sqrt{cx-1}\sqrt{cx+1}} \right) + d^2 \log(x)(a + \operatorname{barccosh}(cx)) + dex^2(a + \operatorname{barccosh}(cx)) + \frac{1}{4}e^2 x^4(a + \operatorname{barccosh}(cx)) -$$

input `Int[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x,x]`

```
output d*e*x^2*(a + b*ArcCosh[c*x]) + (e^2*x^4*(a + b*ArcCosh[c*x]))/4 + d^2*(a +
  b*ArcCosh[c*x])*Log[x] - (b*c*((2*d*e*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^2
  + (3*e^2*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(8*c^4) + (e^2*x^3*Sqrt[-1 + c*x]
  ]*Sqrt[1 + c*x])/(4*c^2) + (2*d*e*ArcCosh[c*x])/c^3 + (3*e^2*ArcCosh[c*x])
  /(8*c^5) + ((2*I)*d^2*Sqrt[1 - c^2*x^2]*ArcSin[c*x]^2)/(c*Sqrt[-1 + c*x]*S
  qrt[1 + c*x]) - (4*d^2*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[1 - E^((2*I)*ArcS
  in[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (4*d^2*Sqrt[1 - c^2*x^2]*Arc
  Sin[c*x]*Log[x])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((2*I)*d^2*Sqrt[1 - c^
  2*x^2]*PolyLog[2, E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])
  ))/4
```

### 3.475.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
  tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6373 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x
  _)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Sim
  p[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
  + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] &&
  NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && Le
  Q[m + p, 0]))
```

```
rule 7293 Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
  ]
```

**3.475.4 Maple [A] (verified)**

Time = 0.99 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.65

method	result
parts	$a \left( \frac{e^2 x^4}{4} + d e x^2 + d^2 \ln(x) \right) + \frac{b \operatorname{arccosh}(cx) e^2 x^4}{4} - \frac{b d e \operatorname{arccosh}(cx)}{2c^2} - \frac{b \sqrt{cx-1} \sqrt{cx+1} d e x}{2c} + b d^2 \operatorname{arccosh}(cx)$
derivativedivides	$a d e x^2 + \frac{a e^2 x^4}{4} + a d^2 \ln(cx) + \frac{b \operatorname{arccosh}(cx) e^2 x^4}{4} - \frac{3 b e^2 \operatorname{arccosh}(cx)}{32 c^4} - \frac{b \sqrt{cx-1} \sqrt{cx+1} d e x}{2c} + b \operatorname{arccosh}(cx)$
default	$a d e x^2 + \frac{a e^2 x^4}{4} + a d^2 \ln(cx) + \frac{b \operatorname{arccosh}(cx) e^2 x^4}{4} - \frac{3 b e^2 \operatorname{arccosh}(cx)}{32 c^4} - \frac{b \sqrt{cx-1} \sqrt{cx+1} d e x}{2c} + b \operatorname{arccosh}(cx)$

input `int((e*x^2+d)^2*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`output `a*(1/4*e^2*x^4+d*e*x^2+d^2*ln(x))+1/4*b*arccosh(c*x)*e^2*x^4-1/2*b/c^2*d*e*arccosh(c*x)-1/2*b/c*(c*x-1)^(1/2)*(c*x+1)^(1/2)*d*e*x+b*d^2*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)-3/32*b/c^4*e^2*arccosh(c*x)-1/16*b*e^2*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-3/32*b/c^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)*e^2*x+b*arccosh(c*x)*d*e*x^2-1/2*b*d^2*arccosh(c*x)^2+1/2*b*d^2*polylg(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)`**3.475.5 Fracas [F]**

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arccosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x,x, algorithm="fracas")`output `integral((a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arccosh(c*x))/x, x)`

**3.475.6 Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^2}{x} dx$$

input `integrate((e*x**2+d)**2*(a+b*acosh(c*x))/x,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**2/x, x)`

**3.475.7 Maxima [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `1/4*a*e^2*x^4 + a*d*e*x^2 + a*d^2*log(x) + integrate(b*e^2*x^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 2*b*d*e*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + b*d^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

**3.475.8 Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arccosh(c*x) + a)/x, x)`

**3.475.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^2}{x} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x,x)`output `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x, x)`

**3.476**  $\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.476.1 Optimal result . . . . . 3503  
 3.476.2 Mathematica [A] (verified) . . . . . 3504  
 3.476.3 Rubi [A] (warning: unable to verify) . . . . . 3504  
 3.476.4 Maple [A] (verified) . . . . . 3507  
 3.476.5 Fricas [A] (verification not implemented) . . . . . 3507  
 3.476.6 Sympy [F] . . . . . 3508  
 3.476.7 Maxima [A] (verification not implemented) . . . . . 3508  
 3.476.8 Giac [F] . . . . . 3509  
 3.476.9 Mupad [F(-1)] . . . . . 3509

**3.476.1 Optimal result**

Integrand size = 21, antiderivative size = 160

$$\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^2} dx = \frac{be(6c^2d+e)(1-c^2x^2)}{3c^3\sqrt{-1+cx}\sqrt{1+cx}} - \frac{be^2(1-c^2x^2)^2}{9c^3\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{d^2(a+b\operatorname{arccosh}(cx))}{x} + 2dex(a+b\operatorname{arccosh}(cx))$$

$$+ \frac{1}{3}e^2x^3(a+b\operatorname{arccosh}(cx))$$

$$+ bcd^2 \arctan\left(\sqrt{-1+cx}\sqrt{1+cx}\right)$$

```
output -d^2*(a+b*arccosh(c*x))/x+2*d*e*x*(a+b*arccosh(c*x))+1/3*e^2*x^3*(a+b*arccosh(c*x))+b*c*d^2*arctan((c*x-1)^(1/2)*(c*x+1)^(1/2))+1/3*b*e*(6*c^2*d+e)*(-c^2*x^2+1)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/9*b*e^2*(-c^2*x^2+1)^2/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```



**3.476.2 Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.80

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx = \frac{1}{3} \left( -\frac{3ad^2}{x} + 6adex + ae^2x^3 - \frac{be\sqrt{-1+cx}\sqrt{1+cx}(2e + c^2(18d + ex^2))}{3c^3} + \frac{b(-3d^2 + 6dex^2 + e^2x^4) \operatorname{arccosh}(cx)}{x} - 3bcd^2 \arctan \left( \frac{1}{\sqrt{-1+cx}\sqrt{1+cx}} \right) \right)$$

input `Integrate[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x^2,x]`output `((-3*a*d^2)/x + 6*a*d*e*x + a*e^2*x^3 - (b*e*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*e + c^2*(18*d + e*x^2)))/(3*c^3) + (b*(-3*d^2 + 6*d*e*x^2 + e^2*x^4)*ArcCosh[c*x])/x - 3*b*c*d^2*ArcTan[1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])])/3`**3.476.3 Rubi [A] (warning: unable to verify)**Time = 0.52 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.90, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6373, 27, 1905, 1578, 1192, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx$$

$$\downarrow 6373$$

$$-bc \int -\frac{-e^2x^4 - 6dex^2 + 3d^2}{3x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^2(a + \operatorname{barccosh}(cx))}{x} + 2dex(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^2x^3(a + \operatorname{barccosh}(cx))$$

$$\downarrow 27$$

$$\frac{1}{3}bc \int \frac{-e^2x^4 - 6dex^2 + 3d^2}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^2(a + \operatorname{barccosh}(cx))}{x} + 2dex(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^2x^3(a + \operatorname{barccosh}(cx))$$

---

3.476.  $\int \frac{(d+ex^2)^2(a+\operatorname{barccosh}(cx))}{x^2} dx$

$$\begin{aligned}
& \downarrow 1905 \\
& \frac{bc\sqrt{c^2x^2-1} \int \frac{-e^2x^4-6dex^2+3d^2}{x\sqrt{c^2x^2-1}} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{x} + 2dex(a+\operatorname{barccosh}(cx)) + \frac{1}{3}e^2x^3(a+\operatorname{barccosh}(cx)) \\
& \downarrow 1578 \\
& \frac{bc\sqrt{c^2x^2-1} \int \frac{-e^2x^4-6dex^2+3d^2}{x^2\sqrt{c^2x^2-1}} dx^2}{6\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{x} + 2dex(a+\operatorname{barccosh}(cx)) + \frac{1}{3}e^2x^3(a+\operatorname{barccosh}(cx)) \\
& \downarrow 1192 \\
& \frac{b\sqrt{c^2x^2-1} \int \frac{-e^2x^8-2e(3dc^2+e)x^4+3c^4d^2-e^2-6c^2de}{x^4+1} d\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{x} + 2dex(a+\operatorname{barccosh}(cx)) + \frac{1}{3}e^2x^3(a+\operatorname{barccosh}(cx)) \\
& \downarrow 1467 \\
& \frac{b\sqrt{c^2x^2-1} \int \left(\frac{3d^2c^4}{x^4+1} - e^2x^4 - e(6dc^2+e)\right) d\sqrt{c^2x^2-1}}{3c^3\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{x} + 2dex(a+\operatorname{barccosh}(cx)) + \frac{1}{3}e^2x^3(a+\operatorname{barccosh}(cx)) \\
& \downarrow 2009 \\
& -\frac{d^2(a+\operatorname{barccosh}(cx))}{x} + 2dex(a+\operatorname{barccosh}(cx)) + \frac{1}{3}e^2x^3(a+\operatorname{barccosh}(cx)) + \\
& \frac{b\sqrt{c^2x^2-1} \left(3c^4d^2 \arctan(\sqrt{c^2x^2-1}) - e\sqrt{c^2x^2-1}(6c^2d+e) - \frac{1}{3}e^2x^6\right)}{3c^3\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x^2,x]`

output `-((d^2*(a + b*ArcCosh[c*x]))/x) + 2*d*e*x*(a + b*ArcCosh[c*x]) + (e^2*x^3*(a + b*ArcCosh[c*x]))/3 + (b*Sqrt[-1 + c^2*x^2]*(-1/3*(e^2*x^6) - e*(6*c^2*d + e)*Sqrt[-1 + c^2*x^2] + 3*c^4*d^2*ArcTan[Sqrt[-1 + c^2*x^2]]))/(3*c^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

## 3.476.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 1192 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`
- rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`
- rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`
- rule 1905 `Int[((f_)*(x_))^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.476.4 Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.11

method	result
parts	$a\left(\frac{e^2x^3}{3} + 2dex - \frac{d^2}{x}\right) + bc\left(\frac{\operatorname{arccosh}(cx)x^3e^2}{3c} + \frac{2\operatorname{arccosh}(cx)xde}{c} - \frac{\operatorname{arccosh}(cx)d^2}{cx} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{c^4}\right)$
derivativedivides	$c\left(\frac{a\left(2c^3xde + \frac{c^3x^3e^2}{3} - \frac{c^3d^2}{x}\right)}{c^4} + \frac{b\left(2\operatorname{arccosh}(cx)d c^3ex + \frac{\operatorname{arccosh}(cx)e^2c^3x^3}{3} - \frac{\operatorname{arccosh}(cx)c^3d^2}{x} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{c^4}\right)}{c^4}\right)$
default	$c\left(\frac{a\left(2c^3xde + \frac{c^3x^3e^2}{3} - \frac{c^3d^2}{x}\right)}{c^4} + \frac{b\left(2\operatorname{arccosh}(cx)d c^3ex + \frac{\operatorname{arccosh}(cx)e^2c^3x^3}{3} - \frac{\operatorname{arccosh}(cx)c^3d^2}{x} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{c^4}\right)}{c^4}\right)$

input `int((e*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)`

output `a*(1/3*e^2*x^3+2*d*e*x-d^2/x)+b*c*(1/3/c*arccosh(c*x)*x^3*e^2+2/c*arccosh(c*x)*x*d*e-arccosh(c*x)*d^2/c/x-1/9/c^4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(9*c^4*d^2*arctan(1/(c^2*x^2-1)^(1/2))+18*d*c^2*e*(c^2*x^2-1)^(1/2)+e^2*(c^2*x^2-1)^(1/2)*c^2*x^2+2*e^2*(c^2*x^2-1)^(1/2))/(c^2*x^2-1)^(1/2))`

### 3.476.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.48

$$\int \frac{(d + ex^2)^2 (a + b\operatorname{arccosh}(cx))}{x^2} dx$$

$$= \frac{3ac^3e^2x^4 + 18bc^4d^2x \arctan(-cx + \sqrt{c^2x^2 - 1}) + 18ac^3dex^2 - 9ac^3d^2 + 3(3bc^3d^2 - 6bc^3de - bc^3e^2)x}{c^4}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x, algorithm="fricas")`

output  $1/9*(3*a*c^3*e^2*x^4 + 18*b*c^4*d^2*x*\arctan(-c*x + \sqrt{c^2*x^2 - 1}) + 18*a*c^3*d*e*x^2 - 9*a*c^3*d^2 + 3*(3*b*c^3*d^2 - 6*b*c^3*d*e - b*c^3*e^2)*x*\log(-c*x + \sqrt{c^2*x^2 - 1}) + 3*(b*c^3*e^2*x^4 + 6*b*c^3*d*e*x^2 - 3*b*c^3*d^2 + (3*b*c^3*d^2 - 6*b*c^3*d*e - b*c^3*e^2)*x)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (b*c^2*e^2*x^3 + 2*(9*b*c^2*d*e + b*e^2)*x)*\sqrt{c^2*x^2 - 1})/(c^3*x)$

### 3.476.6 Sympy [F]

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^2}{x^2} dx$$

input `integrate((e*x**2+d)**2*(a+b*acosh(c*x))/x**2,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**2/x**2, x)`

### 3.476.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 134, normalized size of antiderivative = 0.84

$$\begin{aligned} & \int \frac{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))}{x^2} dx \\ &= \frac{1}{3} ae^2 x^3 - \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) bd^2 \\ &+ \frac{1}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) be^2 \\ &+ 2 adex + \frac{2 (cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) bde}{c} - \frac{ad^2}{x} \end{aligned}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output  $1/3*a*e^2*x^3 - (c*\arcsin(1/(c*\operatorname{abs}(x))) + \operatorname{arccosh}(c*x)/x)*b*d^2 + 1/9*(3*x^3*\operatorname{arccosh}(c*x) - c*(\sqrt{c^2*x^2 - 1}*x^2/c^2 + 2*\sqrt{c^2*x^2 - 1}/c^4))*b*e^2 + 2*a*d*e*x + 2*(c*x*\operatorname{arccosh}(c*x) - \sqrt{c^2*x^2 - 1})*b*d*e/c - a*d^2/x$

---

3.476.  $\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^2} dx$

**3.476.8 Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arccosh(c*x) + a)/x^2, x)`

**3.476.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^2}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x^2,x)`

output `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x^2, x)`

**3.477**  $\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^3} dx$

3.477.1 Optimal result . . . . . 3510  
 3.477.2 Mathematica [A] (verified) . . . . . 3511  
 3.477.3 Rubi [A] (verified) . . . . . 3512  
 3.477.4 Maple [A] (verified) . . . . . 3514  
 3.477.5 Fricas [F] . . . . . 3514  
 3.477.6 Sympy [F] . . . . . 3515  
 3.477.7 Maxima [F] . . . . . 3515  
 3.477.8 Giac [F] . . . . . 3515  
 3.477.9 Mupad [F(-1)] . . . . . 3516

**3.477.1 Optimal result**

Integrand size = 21, antiderivative size = 321

$$\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^3} dx = \frac{bcd^2\sqrt{-1+cx}\sqrt{1+cx}}{2x} - \frac{be^2x\sqrt{-1+cx}\sqrt{1+cx}}{4c}$$

$$- \frac{be^2\operatorname{arccosh}(cx)}{4c^2} - \frac{d^2(a+b\operatorname{arccosh}(cx))}{2x^2}$$

$$+ \frac{1}{2}e^2x^2(a+b\operatorname{arccosh}(cx))$$

$$- \frac{ibde\sqrt{1-c^2x^2}\arcsin(cx)^2}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{2bde\sqrt{1-c^2x^2}\arcsin(cx)\log(1-e^{2i\arcsin(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ 2de(a+b\operatorname{arccosh}(cx))\log(x)$$

$$- \frac{2bde\sqrt{1-c^2x^2}\arcsin(cx)\log(x)}{\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{ibde\sqrt{1-c^2x^2}\operatorname{PolyLog}(2, e^{2i\arcsin(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

output 
$$-1/4*b*e^2*arccosh(c*x)/c^2-1/2*d^2*(a+b*arccosh(c*x))/x^2+1/2*e^2*x^2*(a+b*arccosh(c*x))+2*d*e*(a+b*arccosh(c*x))*ln(x)+1/2*b*c*d^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x-1/4*b*e^2*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-I*b*d*e*arcsin(c*x)^2*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2*b*d*e*arcsin(c*x)*ln(1-(I*c*x+(-c^2*x^2+1)^(1/2))^2*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2))-2*b*d*e*arcsin(c*x)*ln(x)*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-I*b*d*e*polylog(2,(I*c*x+(-c^2*x^2+1)^(1/2))^2*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2))$$

### 3.477.2 Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.54

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))}{x^3} dx$$

$$= \frac{1}{4} \left( -\frac{2ad^2}{x^2} + 2ae^2x^2 + \frac{2bd^2 (cx\sqrt{-1+cx}\sqrt{1+cx} - \operatorname{arccosh}(cx))}{x^2} + \frac{be^2 (-cx\sqrt{-1+cx}\sqrt{1+cx} + 2c^2x^2 \operatorname{arccosh}(cx) - 2 \operatorname{arctanh}(\sqrt{\frac{-1+cx}{1+cx}}))}{c^2} + 8ade \log(x) + 4bde (\operatorname{arccosh}(cx) (\operatorname{arccosh}(cx) + 2 \log(1 + e^{-2 \operatorname{arccosh}(cx)})) - \operatorname{PolyLog}(2, -e^{-2 \operatorname{arccosh}(cx)})) \right)$$

input `Integrate[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x^3,x]`

output 
$$((-2*a*d^2)/x^2 + 2*a*e^2*x^2 + (2*b*d^2*(c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - \operatorname{ArcCosh}[c*x]))/x^2 + (b*e^2*(-(c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + 2*c^2*x^2*\operatorname{ArcCosh}[c*x] - 2*\operatorname{ArcTanh}[Sqrt[(-1 + c*x)/(1 + c*x)]]))/c^2 + 8*a*d*e*\operatorname{Log}[x] + 4*b*d*e*(\operatorname{ArcCosh}[c*x]*(\operatorname{ArcCosh}[c*x] + 2*\operatorname{Log}[1 + E^(-2*\operatorname{ArcCosh}[c*x])])) - \operatorname{PolyLog}[2, -E^(-2*\operatorname{ArcCosh}[c*x])]))/4$$



**3.477.3 Rubi [A] (verified)**

Time = 1.10 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.02, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx$$

↓ 6373

$$-bc \int -\frac{\frac{d^2}{x^2} - 4e \log(x)d - e^2 x^2}{2\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^2(a + \operatorname{barccosh}(cx))}{2x^2} + 2de \log(x)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}e^2 x^2(a + \operatorname{barccosh}(cx))$$

↓ 27

$$\frac{1}{2}bc \int \frac{\frac{d^2}{x^2} - 4e \log(x)d - e^2 x^2}{\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^2(a + \operatorname{barccosh}(cx))}{2x^2} + 2de \log(x)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}e^2 x^2(a + \operatorname{barccosh}(cx))$$

↓ 7293

$$\frac{1}{2}bc \int \left( \frac{d^2}{x^2 \sqrt{cx-1}\sqrt{cx+1}} - \frac{4e \log(x)d}{\sqrt{cx-1}\sqrt{cx+1}} - \frac{e^2 x^2}{\sqrt{cx-1}\sqrt{cx+1}} \right) dx - \frac{d^2(a + \operatorname{barccosh}(cx))}{2x^2} + 2de \log(x)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}e^2 x^2(a + \operatorname{barccosh}(cx))$$

↓ 2009

$$-\frac{d^2(a + \operatorname{barccosh}(cx))}{2x^2} + 2de \log(x)(a + \operatorname{barccosh}(cx)) + \frac{1}{2}e^2 x^2(a + \operatorname{barccosh}(cx)) + \frac{1}{2}bc \left( -\frac{e^2 \operatorname{arccosh}(cx)}{2c^3} - \frac{2ide\sqrt{1-c^2x^2} \operatorname{PolyLog}(2, e^{2i \operatorname{arcsin}(cx)})}{c\sqrt{cx-1}\sqrt{cx+1}} - \frac{2ide\sqrt{1-c^2x^2} \operatorname{arcsin}(cx)^2}{c\sqrt{cx-1}\sqrt{cx+1}} + \frac{4de\sqrt{1-c^2x^2} \operatorname{arcsin}(cx)}{c\sqrt{cx-1}\sqrt{cx+1}} \right)$$

input `Int[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x^3,x]`

```
output -1/2*(d^2*(a + b*ArcCosh[c*x])/x^2 + (e^2*x^2*(a + b*ArcCosh[c*x]))/2 + 2
*d*e*(a + b*ArcCosh[c*x])*Log[x] + (b*c*((d^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]
)/x - (e^2*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*c^2) - (e^2*ArcCosh[c*x])/(2
*c^3) - ((2*I)*d*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]^2)/(c*Sqrt[-1 + c*x]*Sqrt
[1 + c*x]) + (4*d*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[1 - E^((2*I)*ArcSin[
c*x]))]/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (4*d*e*Sqrt[1 - c^2*x^2]*ArcSin
[c*x]*Log[x])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((2*I)*d*e*Sqrt[1 - c^2*x
^2]*PolyLog[2, E^((2*I)*ArcSin[c*x]))]/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])))/
2
```

### 3.477.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6373 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x
_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Sim
p[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1
+ c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] &&
NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && Le
Q[m + p, 0]))
```

```
rule 7293 Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

## 3.477.4 Maple [A] (verified)

Time = 1.16 (sec) , antiderivative size = 196, normalized size of antiderivative = 0.61

method	result
parts	$a \left( \frac{e^2 x^2}{2} - \frac{d^2}{2x^2} + 2de \ln(x) \right) - b \operatorname{arccosh}(cx)^2 de + \frac{b e^2 \operatorname{arccosh}(cx)x^2}{2} - \frac{b e^2 x \sqrt{cx-1} \sqrt{cx+1}}{4c} - \frac{b e^2 \operatorname{arccosh}(cx)}{4c}$
derivativedivides	$c^2 \left( \frac{a x^2 e^2}{2c^2} + \frac{2ade \ln(cx)}{c^2} - \frac{a d^2}{2c^2 x^2} - \frac{bde \operatorname{arccosh}(cx)^2}{c^2} + \frac{b \operatorname{arccosh}(cx)x^2 e^2}{2c^2} - \frac{b \sqrt{cx-1} \sqrt{cx+1} e^2 x}{4c^3} - \frac{b e^2 \operatorname{arccosh}(cx)}{4c^3} \right)$
default	$c^2 \left( \frac{a x^2 e^2}{2c^2} + \frac{2ade \ln(cx)}{c^2} - \frac{a d^2}{2c^2 x^2} - \frac{bde \operatorname{arccosh}(cx)^2}{c^2} + \frac{b \operatorname{arccosh}(cx)x^2 e^2}{2c^2} - \frac{b \sqrt{cx-1} \sqrt{cx+1} e^2 x}{4c^3} - \frac{b e^2 \operatorname{arccosh}(cx)}{4c^3} \right)$

```
input int((e*x^2+d)^2*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)
```

```
output a*(1/2*e^2*x^2-1/2*d^2/x^2+2*d*e*ln(x))-b*arccosh(c*x)^2*d*e+1/2*b*e^2*arccosh(c*x)*x^2-1/4*b*e^2*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-1/4*b*e^2*arccosh(c*x)/c^2+1/2*b*c*d^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/x-1/2*b*c^2*d^2-1/2*b*d^2/x^2*arccosh(c*x)+2*b*e*d*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)+b*e*d*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)
```

## 3.477.5 Fracas [F]

$$\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(ex^2+d)^2(b\operatorname{arccosh}(cx)+a)}{x^3} dx$$

```
input integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^3,x, algorithm="fracas")
```

```
output integral((a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arccosh(c*x))/x^3, x)
```

**3.477.6 Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^2}{x^3} dx$$

input `integrate((e*x**2+d)**2*(a+b*acosh(c*x))/x**3,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**2/x**3, x)`

**3.477.7 Maxima [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `1/2*a*e^2*x^2 + 1/2*b*d^2*(sqrt(c^2*x^2 - 1)*c/x - arccosh(c*x)/x^2) + 2*a*d*e*log(x) - 1/2*a*d^2/x^2 + integrate(b*e^2*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 2*b*d*e*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

**3.477.8 Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arccosh(c*x) + a)/x^3, x)`

**3.477.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^2}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x^3,x)`output `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x^3, x)`

**3.478**  $\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^4} dx$

3.478.1 Optimal result . . . . . 3517  
 3.478.2 Mathematica [A] (verified) . . . . . 3518  
 3.478.3 Rubi [A] (warning: unable to verify) . . . . . 3518  
 3.478.4 Maple [A] (verified) . . . . . 3521  
 3.478.5 Fricas [A] (verification not implemented) . . . . . 3522  
 3.478.6 Sympy [F] . . . . . 3523  
 3.478.7 Maxima [A] (verification not implemented) . . . . . 3523  
 3.478.8 Giac [F] . . . . . 3524  
 3.478.9 Mupad [F(-1)] . . . . . 3524

**3.478.1 Optimal result**

Integrand size = 21, antiderivative size = 184

$$\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^4} dx = \frac{be^2(1-c^2x^2)}{c\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bcd^2(1-c^2x^2)}{6x^2\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{d^2(a+b\operatorname{arccosh}(cx))}{3x^3} - \frac{2de(a+b\operatorname{arccosh}(cx))}{x}$$

$$+ e^2x(a+b\operatorname{arccosh}(cx))$$

$$+ \frac{bcd(c^2d+12e)\sqrt{-1+c^2x^2}\arctan(\sqrt{-1+c^2x^2})}{6\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/3*d^2*(a+b*arccosh(c*x))/x^3-2*d*e*(a+b*arccosh(c*x))/x+e^2*x*(a+b*arccosh(c*x))+b*e^2*(-c^2*x^2+1)/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/6*b*c*d^2*(-c^2*x^2+1)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/6*b*c*d*(c^2*d+12*e)*arctan((c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.478.2 Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.72

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx = -\frac{ad^2}{3x^3} - \frac{2ade}{x} + ae^2x + b\left(-\frac{e^2}{c} + \frac{cd^2}{6x^2}\right) \sqrt{-1 + cx} \sqrt{1 + cx} - \frac{b(d^2 + 6dex^2 - 3e^2x^4) \operatorname{arccosh}(cx)}{3x^3} - \frac{1}{6}bcd(c^2d + 12e) \arctan\left(\frac{1}{\sqrt{-1 + cx} \sqrt{1 + cx}}\right)$$

input `Integrate[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x^4,x]`output `-1/3*(a*d^2)/x^3 - (2*a*d*e)/x + a*e^2*x + b*(-(e^2/c) + (c*d^2)/(6*x^2))*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - (b*(d^2 + 6*d*e*x^2 - 3*e^2*x^4)*ArcCosh[c*x])/(3*x^3) - (b*c*d*(c^2*d + 12*e)*ArcTan[1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])])/6`**3.478.3 Rubi [A] (warning: unable to verify)**Time = 0.52 (sec) , antiderivative size = 168, normalized size of antiderivative = 0.91, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6373, 27, 1905, 1578, 1192, 1471, 25, 299, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx$$

↓ 6373

$$-bc \int -\frac{-3e^2x^4 + 6dex^2 + d^2}{3x^3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^2(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a + \operatorname{barccosh}(cx))}{x} + e^2x(a + \operatorname{barccosh}(cx))$$

↓ 27

$$\frac{1}{3}bc \int \frac{-3e^2x^4 + 6dex^2 + d^2}{x^3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^2(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a + \operatorname{barccosh}(cx))}{x} + e^2x(a + \operatorname{barccosh}(cx))$$

---

3.478.  $\int \frac{(d+ex^2)^2(a+\operatorname{barccosh}(cx))}{x^4} dx$

$$\begin{aligned}
& \downarrow 1905 \\
& \frac{bc\sqrt{c^2x^2-1} \int \frac{-3e^2x^4+6dex^2+d^2}{x^3\sqrt{c^2x^2-1}} dx}{3\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a+\operatorname{barccosh}(cx))}{x} + e^2x(a+\operatorname{barccosh}(cx)) \\
& \downarrow 1578 \\
& \frac{bc\sqrt{c^2x^2-1} \int \frac{-3e^2x^4+6dex^2+d^2}{x^4\sqrt{c^2x^2-1}} dx^2}{6\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a+\operatorname{barccosh}(cx))}{x} + e^2x(a+\operatorname{barccosh}(cx)) \\
& \downarrow 1192 \\
& \frac{b\sqrt{c^2x^2-1} \int \frac{-3e^2x^8+6(c^2d-e)ex^4+c^4d^2-3e^2+6c^2de}{(x^4+1)^2} d\sqrt{c^2x^2-1}}{3c\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a+\operatorname{barccosh}(cx))}{x} + e^2x(a+\operatorname{barccosh}(cx)) \\
& \downarrow 1471 \\
& \frac{b\sqrt{c^2x^2-1} \left( \frac{c^4d^2\sqrt{c^2x^2-1}}{2(x^4+1)} - \frac{1}{2} \int \frac{d^2c^4+12dec^2-6e^2x^4-6e^2}{x^4+1} d\sqrt{c^2x^2-1} \right)}{3c\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a+\operatorname{barccosh}(cx))}{x} + e^2x(a+\operatorname{barccosh}(cx)) \\
& \downarrow 25 \\
& \frac{b\sqrt{c^2x^2-1} \left( \frac{1}{2} \int \frac{d^2c^4+12dec^2-6e^2x^4-6e^2}{x^4+1} d\sqrt{c^2x^2-1} + \frac{c^4d^2\sqrt{c^2x^2-1}}{2(x^4+1)} \right)}{3c\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a+\operatorname{barccosh}(cx))}{x} + e^2x(a+\operatorname{barccosh}(cx)) \\
& \downarrow 299 \\
& \frac{b\sqrt{c^2x^2-1} \left( \frac{1}{2} \left( c^2d(c^2d+12e) \int \frac{1}{x^4+1} d\sqrt{c^2x^2-1} - 6e^2\sqrt{c^2x^2-1} \right) + \frac{c^4d^2\sqrt{c^2x^2-1}}{2(x^4+1)} \right)}{3c\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a+\operatorname{barccosh}(cx))}{x} + e^2x(a+\operatorname{barccosh}(cx)) \\
& \downarrow 216 \\
& \frac{d^2(a+\operatorname{barccosh}(cx))}{3x^3} - \frac{2de(a+\operatorname{barccosh}(cx))}{x} + e^2x(a+\operatorname{barccosh}(cx)) + \\
& \frac{b\sqrt{c^2x^2-1} \left( \frac{1}{2} \left( c^2d \arctan \left( \sqrt{c^2x^2-1} \right) (c^2d+12e) - 6e^2\sqrt{c^2x^2-1} \right) + \frac{c^4d^2\sqrt{c^2x^2-1}}{2(x^4+1)} \right)}{3c\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

---

3.478.  $\int \frac{(d+ex^2)^2(a+\operatorname{barccosh}(cx))}{x^4} dx$



input `Int[((d + e*x^2)^2*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*(d^2*(a + b*ArcCosh[c*x]))/x^3 - (2*d*e*(a + b*ArcCosh[c*x]))/x + e^2*x*(a + b*ArcCosh[c*x]) + (b*Sqrt[-1 + c^2*x^2]*((c^4*d^2*Sqrt[-1 + c^2*x^2]))/(2*(1 + x^4)) + (-6*e^2*Sqrt[-1 + c^2*x^2] + c^2*d*(c^2*d + 12*e)*ArcTan[Sqrt[-1 + c^2*x^2]]/2))/(3*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.478.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`

rule 1192 `Int[((d_.) + (e_.)*(x_)^(m_))*((f_.) + (g_.)*(x_)^(n_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1471 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], R = Coeff[PolynomialRemainder[(a + b*x^2 + c*x^4)^p, d + e*x^2, x], 0]}, Simp[(-R)*x*((d + e*x^2)^(q + 1)/(2*d*(q + 1))), x] + Simp[1/(2*d*(q + 1)) Int[(d + e*x^2)^(q + 1)*ExpandToSum[2*d*(q + 1)*Qx + R*(2*q + 3), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && LtQ[q, -1]`

rule 1578 `Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]`

rule 1905 `Int[((f_)*(x_)^(m_)*((d1_) + (e1_)*(x_)^(non2_))^(q_)*((d2_) + (e2_)*(x_)^(non2_))^(p_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`

rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_)^(m_)*((d_) + (e_)*(x_)^2)^(p_)), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.478.4 Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.04

---


$$3.478. \quad \int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^4} dx$$

method	result
parts	$a \left( e^2 x - \frac{2de}{x} - \frac{d^2}{3x^3} \right) + b c^3 \left( \frac{\operatorname{arccosh}(cx) x e^2}{c^3} - \frac{2 \operatorname{arccosh}(cx) de}{c^3 x} - \frac{\operatorname{arccosh}(cx) d^2}{3c^3 x^3} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{c^3} \left( \operatorname{arctan} \left( \frac{1}{\sqrt{c^2 x^2 - 1}} \right) \right) \right)$
derivativedivides	$c^3 \left( \frac{a \left( cx e^2 - \frac{c d^2}{3x^3} - \frac{2dce}{x} \right)}{c^4} + \frac{b \left( \operatorname{arccosh}(cx) e^2 cx - \frac{\operatorname{arccosh}(cx) c d^2}{3x^3} - \frac{2 \operatorname{arccosh}(cx) dce}{x} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{c^3} \left( \operatorname{arctan} \left( \frac{1}{\sqrt{c^2 x^2 - 1}} \right) \right) \right)}{c^4} \right)$
default	$c^3 \left( \frac{a \left( cx e^2 - \frac{c d^2}{3x^3} - \frac{2dce}{x} \right)}{c^4} + \frac{b \left( \operatorname{arccosh}(cx) e^2 cx - \frac{\operatorname{arccosh}(cx) c d^2}{3x^3} - \frac{2 \operatorname{arccosh}(cx) dce}{x} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{c^3} \left( \operatorname{arctan} \left( \frac{1}{\sqrt{c^2 x^2 - 1}} \right) \right) \right)}{c^4} \right)$

input `int((e*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)`

output `a*(e^2*x-2*d*e/x-1/3*d^2/x^3)+b*c^3*(1/c^3*arccosh(c*x)*x*e^2-2/c^3*arccosh(c*x)*d*e/x-1/3*arccosh(c*x)*d^2/c^3/x^3-1/6/c^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(arctan(1/(c^2*x^2-1)^(1/2)))*c^6*d^2*x^2+12*arctan(1/(c^2*x^2-1)^(1/2))*c^4*d*e*x^2-(c^2*x^2-1)^(1/2)*c^4*d^2+6*e^2*(c^2*x^2-1)^(1/2)*c^2*x^2)/(c^2*x^2-1)^(1/2)/x^2)`

### 3.478.5 Fracas [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.17

$$\int \frac{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \frac{6ace^2x^4 - 12acdex^2 + 2(bc^4d^2 + 12bc^2de)x^3 \arctan(-cx + \sqrt{c^2x^2 - 1}) + 2(bcd^2 + 6bcde - 3bce^2)x^3 \log(-cx + \sqrt{c^2x^2 - 1})}{c^4}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")`

output `1/6*(6*a*c*e^2*x^4 - 12*a*c*d*e*x^2 + 2*(b*c^4*d^2 + 12*b*c^2*d*e)*x^3*arctan(-c*x + sqrt(c^2*x^2 - 1)) + 2*(b*c*d^2 + 6*b*c*d*e - 3*b*c*e^2)*x^3*log(-c*x + sqrt(c^2*x^2 - 1)) - 2*a*c*d^2 + 2*(3*b*c*e^2*x^4 - 6*b*c*d*e*x^2 - b*c*d^2 + (b*c*d^2 + 6*b*c*d*e - 3*b*c*e^2)*x^3)*log(c*x + sqrt(c^2*x^2 - 1)) + (b*c^2*d^2*x - 6*b*e^2*x^3)*sqrt(c^2*x^2 - 1)/(c*x^3)`

3.478.  $\int \frac{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}{x^4} dx$

**3.478.6 Sympy [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^2}{x^4} dx$$

input `integrate((e*x**2+d)**2*(a+b*acosh(c*x))/x**4,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**2/x**4, x)`

**3.478.7 Maxima [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.68

$$\begin{aligned} & \int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx \\ &= -\frac{1}{6} \left( \left( c^2 \arcsin \left( \frac{1}{c|x|} \right) - \frac{\sqrt{c^2 x^2 - 1}}{x^2} \right) c + \frac{2 \operatorname{arccosh}(cx)}{x^3} \right) b d^2 \\ & \quad - 2 \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) b d e + a e^2 x \\ & \quad + \frac{(cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) b e^2}{c} - \frac{2 a d e}{x} - \frac{a d^2}{3 x^3} \end{aligned}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")`

output `-1/6*((c^2*arcsin(1/(c*abs(x))) - sqrt(c^2*x^2 - 1)/x^2)*c + 2*arccosh(c*x)/x^3)*b*d^2 - 2*(c*arcsin(1/(c*abs(x))) + arccosh(c*x)/x)*b*d*e + a*e^2*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*e^2/c - 2*a*d*e/x - 1/3*a*d^2/x^3`

**3.478.8 Giac [F]**

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)}{x^4} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arccosh(c*x) + a)/x^4, x)`

**3.478.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^2}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x^4,x)`

output `int(((a + b*acosh(c*x))*(d + e*x^2)^2)/x^4, x)`

### 3.479 $\int x^4(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.479.1 Optimal result . . . . .	3525
3.479.2 Mathematica [A] (verified) . . . . .	3526
3.479.3 Rubi [A] (verified) . . . . .	3527
3.479.4 Maple [A] (verified) . . . . .	3529
3.479.5 Fricas [A] (verification not implemented) . . . . .	3530
3.479.6 Sympy [F(-1)] . . . . .	3530
3.479.7 Maxima [A] (verification not implemented) . . . . .	3531
3.479.8 Giac [F(-2)] . . . . .	3531
3.479.9 Mupad [F(-1)] . . . . .	3532

#### 3.479.1 Optimal result

Integrand size = 21, antiderivative size = 435

$$\int x^4(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{b(231c^6d^3 + 495c^4d^2e + 385c^2de^2 + 105e^3)(1 - c^2x^2)}{1155c^{11}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{b(462c^6d^3 + 1485c^4d^2e + 1540c^2de^2 + 525e^3)(1 - c^2x^2)^2}{3465c^{11}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{b(77c^6d^3 + 495c^4d^2e + 770c^2de^2 + 350e^3)(1 - c^2x^2)^3}{1925c^{11}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{be(99c^4d^2 + 308c^2de + 210e^2)(1 - c^2x^2)^4}{1617c^{11}\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{be^2(11c^2d + 15e)(1 - c^2x^2)^5}{297c^{11}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{be^3(1 - c^2x^2)^6}{121c^{11}\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{1}{5}d^3x^5(a + \operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a + \operatorname{barccosh}(cx))$$

$$+ \frac{1}{3}de^2x^9(a + \operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a + \operatorname{barccosh}(cx))$$

output  $\frac{1}{5}d^3x^5(a+b\operatorname{arccosh}(cx))+\frac{3}{7}d^2e^x(x^7(a+b\operatorname{arccosh}(cx))+\frac{1}{3}d^2e^{2x^9}(a+b\operatorname{arccosh}(cx))+\frac{1}{11}e^{3x^{11}}(a+b\operatorname{arccosh}(cx))+\frac{1}{1155}b(231c^6d^3+495c^4d^2e+385c^2de^2+105e^3)(-c^2x^2+1)/c^{11}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-1/3465b(462c^6d^3+1485c^4d^2e+1540c^2de^2+525e^3)(-c^2x^2+1)^2/c^{11}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}+1/1925b(77c^6d^3+495c^4d^2e+770c^2de^2+350e^3)(-c^2x^2+1)^3/c^{11}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-1/1617b(99c^4d^2+308c^2de+210e^2)(-c^2x^2+1)^4/c^{11}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}+1/297be^2(11c^2d+15e)(-c^2x^2+1)^5/c^{11}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-1/121be^3(-c^2x^2+1)^6/c^{11}/(cx-1)^{(1/2)}/(cx+1)^{(1/2)})$

### 3.479.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.63

$$\int x^4(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx$$

$$= \frac{3465ax^5(231d^3+495d^2ex^2+385de^2x^4+105e^3x^6) - \frac{b\sqrt{-1+cx}\sqrt{1+cx}(134400e^3+4480c^2e^2(121d+15ex^2)+80c^4e(9801d^2+3388de^2x^2+630e^2x^4)+24c^6(17787d^3+16335d^2e^2x^2+8470de^2x^4+1750e^3x^6)+c^{10}x^4(160083d^3+245025d^2e^2x^2+148225de^2x^4+33075e^3x^6)+2c^8(106722d^3x^2+147015d^2e^2x^4+84700de^2x^6+18375e^3x^8))}{c^{11}+3465bx^5(231d^3+495d^2e^2x^2+385de^2x^4+105e^3x^6)\operatorname{ArcCosh}[cx]}}{4002075}$$

input `Integrate[x^4*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output  $(3465ax^5(231d^3+495d^2e^2x^2+385de^2x^4+105e^3x^6) - (b\operatorname{Sqrt}[-1+cx]\operatorname{Sqrt}[1+cx](134400e^3+4480c^2e^2(121d+15e^2x^2)+80c^4e(9801d^2+3388de^2x^2+630e^2x^4)+24c^6(17787d^3+16335d^2e^2x^2+8470de^2x^4+1750e^3x^6)+c^{10}x^4(160083d^3+245025d^2e^2x^2+148225de^2x^4+33075e^3x^6)+2c^8(106722d^3x^2+147015d^2e^2x^4+84700de^2x^6+18375e^3x^8)))/c^{11}+3465bx^5(231d^3+495d^2e^2x^2+385de^2x^4+105e^3x^6)\operatorname{ArcCosh}[c*x])/4002075$

**3.479.3 Rubi [A] (verified)**

Time = 0.96 (sec) , antiderivative size = 365, normalized size of antiderivative = 0.84, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6373, 27, 2113, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^4(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx \\
 & \quad \downarrow \text{6373} \\
 & -bc \int \frac{x^5(105e^3x^6+385de^2x^4+495d^2ex^2+231d^3)}{1155\sqrt{cx-1}\sqrt{cx+1}}dx + \frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{3}de^2x^9(a+\operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a+\operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{27} \\
 & -\frac{bc \int \frac{x^5(105e^3x^6+385de^2x^4+495d^2ex^2+231d^3)}{\sqrt{cx-1}\sqrt{cx+1}}dx}{1155} + \frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{3}de^2x^9(a+\operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a+\operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{2113} \\
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^5(105e^3x^6+385de^2x^4+495d^2ex^2+231d^3)}{\sqrt{c^2x^2-1}}dx}{1155\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{3}de^2x^9(a+\operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a+\operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{2331} \\
 & -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^4(105e^3x^6+385de^2x^4+495d^2ex^2+231d^3)}{\sqrt{c^2x^2-1}}dx^2}{2310\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{3}de^2x^9(a+\operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a+\operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{2123} \\
 & \frac{bc\sqrt{c^2x^2-1} \int \left( \frac{105e^3(c^2x^2-1)^{9/2}}{c^{10}} + \frac{35e^2(11dc^2+15e)(c^2x^2-1)^{7/2}}{c^{10}} + \frac{5e(99d^2c^4+308dec^2+210e^2)(c^2x^2-1)^{5/2}}{c^{10}} + \frac{3(77d^3c^6+495d^2e^2)}{c^{10}} \right) dx}{2310\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{3}de^2x^9(a+\operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a+\operatorname{barccosh}(cx)) \\
 & \quad \downarrow \text{2009} \\
 & \frac{bc\sqrt{c^2x^2-1} \int \left( \frac{105e^3(c^2x^2-1)^{9/2}}{c^{10}} + \frac{35e^2(11dc^2+15e)(c^2x^2-1)^{7/2}}{c^{10}} + \frac{5e(99d^2c^4+308dec^2+210e^2)(c^2x^2-1)^{5/2}}{c^{10}} + \frac{3(77d^3c^6+495d^2e^2)}{c^{10}} \right) dx}{2310\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{5}d^3x^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a+\operatorname{barccosh}(cx)) + \frac{1}{3}de^2x^9(a+\operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a+\operatorname{barccosh}(cx))
 \end{aligned}$$



$$\frac{\frac{1}{5}d^3x^5(a + \operatorname{barccosh}(cx)) + \frac{3}{7}d^2ex^7(a + \operatorname{barccosh}(cx)) + \frac{1}{3}de^2x^9(a + \operatorname{barccosh}(cx)) + \frac{1}{11}e^3x^{11}(a + \operatorname{barccosh}(cx)) - bc\sqrt{c^2x^2 - 1} \left( \frac{70e^2(c^2x^2 - 1)^{9/2}(11c^2d + 15e)}{9c^{12}} + \frac{210e^3(c^2x^2 - 1)^{11/2}}{11c^{12}} + \frac{10e(c^2x^2 - 1)^{7/2}(99c^4d^2 + 308c^2de + 210e^2)}{7c^{12}} + \frac{6(c^2x^2 - 1)^{5/2}(77c^6d + 2310\sqrt{cx})}{7c^{12}} \right)}{2310\sqrt{cx}}$$

input `Int[x^4*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/2310*(b*c*Sqrt[-1 + c^2*x^2]*((2*(231*c^6*d^3 + 495*c^4*d^2*e + 385*c^2*d*e^2 + 105*e^3)*Sqrt[-1 + c^2*x^2])/c^12 + (2*(462*c^6*d^3 + 1485*c^4*d^2*e + 1540*c^2*d*e^2 + 525*e^3)*(-1 + c^2*x^2)^(3/2))/(3*c^12) + (6*(77*c^6*d^3 + 495*c^4*d^2*e + 770*c^2*d*e^2 + 350*e^3)*(-1 + c^2*x^2)^(5/2))/(5*c^12) + (10*e*(99*c^4*d^2 + 308*c^2*d*e + 210*e^2)*(-1 + c^2*x^2)^(7/2))/(7*c^12) + (70*e^2*(11*c^2*d + 15*e)*(-1 + c^2*x^2)^(9/2))/(9*c^12) + (210*e^3*(-1 + c^2*x^2)^(11/2))/(11*c^12))/Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (d^3*x^5*(a + b*ArcCosh[c*x]))/5 + (3*d^2*e*x^7*(a + b*ArcCosh[c*x]))/7 + (d*e^2*x^9*(a + b*ArcCosh[c*x]))/3 + (e^3*x^11*(a + b*ArcCosh[c*x]))/11`

### 3.479.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2113 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[P_x*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

rule 2123 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[P_x*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[P_x, x] && (IntegersQ[m, n] || IGtQ[m, -2])`

rule 2331 `Int[(Pq_)*(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`

rule 6373 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.479.4 Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 319, normalized size of antiderivative = 0.73

method	result
parts	$a\left(\frac{1}{11}e^3x^{11} + \frac{1}{3}de^2x^9 + \frac{3}{7}d^2ex^7 + \frac{1}{5}d^3x^5\right) + \frac{b\left(\frac{c^5 \operatorname{arccosh}(cx)e^3x^{11}}{11} + \frac{c^5 \operatorname{arccosh}(cx)de^2x^9}{3} + \frac{3e^5 \operatorname{arccosh}(cx)d^2ex^7}{7}\right)}{c^6}$
derivativedivides	$\frac{a\left(\frac{1}{5}c^{11}d^3x^5 + \frac{3}{7}c^{11}d^2ex^7 + \frac{1}{3}c^{11}de^2x^9 + \frac{1}{11}e^3c^{11}x^{11}\right)}{c^6} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^{11}d^3x^5}{5} + \frac{3 \operatorname{arccosh}(cx)c^{11}d^2ex^7}{7} + \frac{\operatorname{arccosh}(cx)c^{11}de^2x^9}{3}\right)}{c^6}$
default	$\frac{a\left(\frac{1}{5}c^{11}d^3x^5 + \frac{3}{7}c^{11}d^2ex^7 + \frac{1}{3}c^{11}de^2x^9 + \frac{1}{11}e^3c^{11}x^{11}\right)}{c^6} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^{11}d^3x^5}{5} + \frac{3 \operatorname{arccosh}(cx)c^{11}d^2ex^7}{7} + \frac{\operatorname{arccosh}(cx)c^{11}de^2x^9}{3}\right)}{c^6}$

input `int(x^4*(e*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/11*e^3*x^11+1/3*d*e^2*x^9+3/7*d^2*e*x^7+1/5*d^3*x^5)+b/c^5*(1/11*c^5*arccosh(c*x)*e^3*x^11+1/3*c^5*arccosh(c*x)*d*e^2*x^9+3/7*c^5*arccosh(c*x)*d^2*e*x^7+1/5*arccosh(c*x)*c^5*x^5*d^3-1/4002075/c^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(33075*c^10*e^3*x^10+148225*c^10*d*e^2*x^8+245025*c^10*d^2*e*x^6+36750*c^8*e^3*x^8+160083*c^10*d^3*x^4+169400*c^8*d*e^2*x^6+294030*c^8*d^2*e*x^4+42000*c^6*e^3*x^6+213444*c^8*d^3*x^2+203280*c^6*d*e^2*x^4+392040*c^6*d^2*e*x^2+50400*c^4*e^3*x^4+426888*c^6*d^3+271040*c^4*d*e^2*x^2+784080*c^4*d^2*e+67200*c^2*e^3*x^2+542080*c^2*d*e^2+134400*e^3))`

**3.479.5 Fracas [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 334, normalized size of antiderivative = 0.77

$$\int x^4 (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{363825 ac^{11} e^3 x^{11} + 1334025 ac^{11} d e^2 x^9 + 1715175 ac^{11} d^2 e x^7 + 800415 ac^{11} d^3 x^5 + 3465 (105 bc^{11} e^3 x^{11} + 385 b^2 c^{11} d e^2 x^9 + 495 b^2 c^{11} d^2 e x^7 + 231 b^2 c^{11} d^3 x^5) \log(cx + \sqrt{c^2 x^2 - 1}) - (33075 b^3 c^{10} e^3 x^{10} + 426888 b^3 c^6 d^3 + 1225 (121 b^3 c^{10} d e^2 + 30 b^3 c^8 e^3) x^8 + 784080 b^3 c^4 d^2 e + 25 (9801 b^3 c^{10} d^2 e + 6776 b^3 c^8 d e^2 + 1680 b^3 c^6 e^3) x^6 + 542080 b^3 c^2 d e^2 + 3 (53361 b^3 c^{10} d^3 + 98010 b^3 c^8 d^2 e + 67760 b^3 c^6 d e^2 + 16800 b^3 c^4 e^3) x^4 + 134400 b^3 e^3 + 4 (53361 b^3 c^8 d^3 + 98010 b^3 c^6 d^2 e + 67760 b^3 c^4 d e^2 + 16800 b^3 c^2 e^3) x^2) \sqrt{c^2 x^2 - 1}}{c^{11}}$$

input `integrate(x^4*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`output `1/4002075*(363825*a*c^11*e^3*x^11 + 1334025*a*c^11*d*e^2*x^9 + 1715175*a*c^11*d^2*e*x^7 + 800415*a*c^11*d^3*x^5 + 3465*(105*b*c^11*e^3*x^11 + 385*b*c^11*d*e^2*x^9 + 495*b*c^11*d^2*e*x^7 + 231*b*c^11*d^3*x^5)*log(c*x + sqrt(c^2*x^2 - 1)) - (33075*b*c^10*e^3*x^10 + 426888*b*c^6*d^3 + 1225*(121*b*c^10*d*e^2 + 30*b*c^8*e^3)*x^8 + 784080*b*c^4*d^2*e + 25*(9801*b*c^10*d^2*e + 6776*b*c^8*d*e^2 + 1680*b*c^6*e^3)*x^6 + 542080*b*c^2*d*e^2 + 3*(53361*b*c^10*d^3 + 98010*b*c^8*d^2*e + 67760*b*c^6*d*e^2 + 16800*b*c^4*e^3)*x^4 + 134400*b*e^3 + 4*(53361*b*c^8*d^3 + 98010*b*c^6*d^2*e + 67760*b*c^4*d*e^2 + 16800*b*c^2*e^3)*x^2)*sqrt(c^2*x^2 - 1)/c^11`**3.479.6 Sympy [F(-1)]**

Timed out.

$$\int x^4 (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate(x**4*(e*x**2+d)**3*(a+b*acosh(c*x)),x)`output `Timed out`

**3.479.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 451, normalized size of antiderivative = 1.04

$$\int x^4(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx = \frac{1}{11}ae^3x^{11} + \frac{1}{3}ade^2x^9 + \frac{3}{7}ad^2ex^7 + \frac{1}{5}ad^3x^5$$

$$+ \frac{1}{75}\left(15x^5\operatorname{arccosh}(cx) - \left(\frac{3\sqrt{c^2x^2-1}x^4}{c^2} + \frac{4\sqrt{c^2x^2-1}x^2}{c^4} + \frac{8\sqrt{c^2x^2-1}}{c^6}\right)c\right)bd^3$$

$$+ \frac{3}{245}\left(35x^7\operatorname{arccosh}(cx) - \left(\frac{5\sqrt{c^2x^2-1}x^6}{c^2} + \frac{6\sqrt{c^2x^2-1}x^4}{c^4} + \frac{8\sqrt{c^2x^2-1}x^2}{c^6} + \frac{16\sqrt{c^2x^2-1}}{c^8}\right)c\right)bd^2e$$

$$+ \frac{1}{945}\left(315x^9\operatorname{arccosh}(cx) - \left(\frac{35\sqrt{c^2x^2-1}x^8}{c^2} + \frac{40\sqrt{c^2x^2-1}x^6}{c^4} + \frac{48\sqrt{c^2x^2-1}x^4}{c^6} + \frac{64\sqrt{c^2x^2-1}x^2}{c^8} + \frac{128\sqrt{c^2x^2-1}}{c^{10}}\right)c\right)bd^2e^2$$

$$+ \frac{1}{7623}\left(693x^{11}\operatorname{arccosh}(cx) - \left(\frac{63\sqrt{c^2x^2-1}x^{10}}{c^2} + \frac{70\sqrt{c^2x^2-1}x^8}{c^4} + \frac{80\sqrt{c^2x^2-1}x^6}{c^6} + \frac{96\sqrt{c^2x^2-1}x^4}{c^8} + \frac{128\sqrt{c^2x^2-1}x^2}{c^{10}} + \frac{256\sqrt{c^2x^2-1}}{c^{12}}\right)c\right)bd^2e^3$$

input `integrate(x^4*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

```
output 1/11*a*e^3*x^11 + 1/3*a*d*e^2*x^9 + 3/7*a*d^2*e*x^7 + 1/5*a*d^3*x^5 + 1/75
*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)
*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d^3 + 3/245*(35*x^7*arccosh(c*x)
- (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*
x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*d^2*e + 1/945*(315*x^9*a
rccosh(c*x) - (35*sqrt(c^2*x^2 - 1)*x^8/c^2 + 40*sqrt(c^2*x^2 - 1)*x^6/c^4
+ 48*sqrt(c^2*x^2 - 1)*x^4/c^6 + 64*sqrt(c^2*x^2 - 1)*x^2/c^8 + 128*sqrt(
c^2*x^2 - 1)/c^10)*c)*b*d*e^2 + 1/7623*(693*x^11*arccosh(c*x) - (63*sqrt(c
^2*x^2 - 1)*x^10/c^2 + 70*sqrt(c^2*x^2 - 1)*x^8/c^4 + 80*sqrt(c^2*x^2 - 1)
*x^6/c^6 + 96*sqrt(c^2*x^2 - 1)*x^4/c^8 + 128*sqrt(c^2*x^2 - 1)*x^2/c^10 +
256*sqrt(c^2*x^2 - 1)/c^12)*c)*b*e^3
```

**3.479.8 Giac [F(-2)]**

Exception generated.

$$\int x^4(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx = \text{Exception raised: RuntimeError}$$

input `integrate(x^4*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")`

output Exception raised: RuntimeError >> an error occurred running a Giac command  
 :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const ve  
 cteur & l) Error: Bad Argument Value

### 3.479.9 Mupad [F(-1)]

Timed out.

$$\int x^4 (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x^4 (a + b \operatorname{acosh}(cx)) (ex^2 + d)^3 dx$$

input `int(x^4*(a + b*acosh(c*x))*(d + e*x^2)^3,x)`

output `int(x^4*(a + b*acosh(c*x))*(d + e*x^2)^3, x)`

### 3.480 $\int x^3(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.480.1 Optimal result . . . . .	3533
3.480.2 Mathematica [A] (warning: unable to verify) . . . . .	3534
3.480.3 Rubi [A] (verified) . . . . .	3535
3.480.4 Maple [A] (verified) . . . . .	3539
3.480.5 Fricas [A] (verification not implemented) . . . . .	3539
3.480.6 Sympy [F] . . . . .	3540
3.480.7 Maxima [A] (verification not implemented) . . . . .	3540
3.480.8 Giac [F(-2)] . . . . .	3541
3.480.9 Mupad [F(-1)] . . . . .	3541

#### 3.480.1 Optimal result

Integrand size = 21, antiderivative size = 494

$$\int x^3(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= -\frac{b(1232c^8d^4 - 2536c^6d^3e - 7758c^4d^2e^2 - 6615c^2de^3 - 1890e^4)x(1 - c^2x^2)}{76800c^9e\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{b(136c^6d^3 - 1096c^4d^2e - 1617c^2de^2 - 630e^3)x(1 - c^2x^2)(d + ex^2)}{38400c^7e\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{b(26c^4d^2 + 201c^2de + 126e^2)x(1 - c^2x^2)(d + ex^2)^2}{9600c^5e\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{b(11c^2d + 18e)x(1 - c^2x^2)(d + ex^2)^3}{1600c^3e\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bx(1 - c^2x^2)(d + ex^2)^4}{100ce\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{d(d + ex^2)^4(a + \operatorname{barccosh}(cx))}{8e^2} + \frac{(d + ex^2)^5(a + \operatorname{barccosh}(cx))}{10e^2}$$

$$+ \frac{b(128c^{10}d^5 - 480c^6d^3e^2 - 800c^4d^2e^3 - 525c^2de^4 - 126e^5)\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{cx}{\sqrt{-1 + c^2x^2}}\right)}{5120c^{10}e^2\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output 
$$\begin{aligned} & -1/8*d*(e*x^2+d)^4*(a+b*\operatorname{arccosh}(c*x))/e^2+1/10*(e*x^2+d)^5*(a+b*\operatorname{arccosh}(c*x))/e^2-1/76800*b*(1232*c^8*d^4-2536*c^6*d^3*e-7758*c^4*d^2*e^2-6615*c^2*d^3*e^3-1890*e^4)*x*(-c^2*x^2+1)/c^9/e/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-1/38400*b*(136*c^6*d^3-1096*c^4*d^2*e-1617*c^2*d*e^2-630*e^3)*x*(-c^2*x^2+1)*(e*x^2+d)/c^7/e/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/9600*b*(26*c^4*d^2+201*c^2*d*e+126*e^2)*x*(-c^2*x^2+1)*(e*x^2+d)^2/c^5/e/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/1600*b*(11*c^2*d+18*e)*x*(-c^2*x^2+1)*(e*x^2+d)^3/c^3/e/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/100*b*x*(-c^2*x^2+1)*(e*x^2+d)^4/c/e/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/5120*b*(128*c^10*d^5-480*c^6*d^3*e^2-800*c^4*d^2*e^3-525*c^2*d*e^4-126*e^5)*\operatorname{arctanh}(c*x/(c^2*x^2-1)^{(1/2)})*(c^2*x^2-1)^{(1/2)}/c^10/e^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)} \end{aligned}$$

### 3.480.2 Mathematica [A] (warning: unable to verify)

Time = 0.36 (sec) , antiderivative size = 294, normalized size of antiderivative = 0.60

$$\int x^3(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx = \frac{1920ax^4(10d^3+20d^2ex^2+15de^2x^4+4e^3x^6) - \frac{bx\sqrt{-1+cx}\sqrt{1+cx}(1890e^3+315c^2e^2(25d+4ex^2)+6c^4e(2000d^2+875dex^2+168e^2x^4))}{c^9} + 1920b*x^4*(10*d^3+20*d^2*e*x^2+15*d*e^2*x^4+4*e^3*x^6)*\operatorname{ArcCosh}[c*x] - (30*b*(480*c^6*d^3+800*c^4*d^2*e+525*c^2*d*e^2+126*e^3)*\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1+c*x)/(1+c*x)])]}{c^{10}}}{76800}$$

input `Integrate[x^3*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output 
$$\begin{aligned} & (1920*a*x^4*(10*d^3+20*d^2*e*x^2+15*d*e^2*x^4+4*e^3*x^6) - (b*x*\operatorname{Sqrt}[-1+c*x]*\operatorname{Sqrt}[1+c*x]*(1890*e^3+315*c^2*e^2*(25*d+4*e*x^2)+6*c^4*e*(2000*d^2+875*d*e*x^2+168*e^2*x^4)+8*c^6*(900*d^3+1000*d^2*e*x^2+525*d*e^2*x^4+108*e^3*x^6)+16*c^8*(300*d^3*x^2+400*d^2*e*x^4+225*d*e^2*x^6+48*e^3*x^8)))/c^9+1920*b*x^4*(10*d^3+20*d^2*e*x^2+15*d*e^2*x^4+4*e^3*x^6)*\operatorname{ArcCosh}[c*x] - (30*b*(480*c^6*d^3+800*c^4*d^2*e+525*c^2*d*e^2+126*e^3)*\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1+c*x)/(1+c*x)])]/c^{10})/76800 \end{aligned}$$

**3.480.3 Rubi [A] (verified)**

Time = 0.86 (sec) , antiderivative size = 442, normalized size of antiderivative = 0.89, number of steps used = 12, number of rules used = 11,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.524$ , Rules used = {6373, 27, 2041, 403, 27, 403, 403, 403, 299, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6373} \\
 & -bc \int -\frac{(d - 4ex^2)(ex^2 + d)^4}{40e^2 \sqrt{cx - 1} \sqrt{cx + 1}} dx + \frac{(d + ex^2)^5 (a + \operatorname{barccosh}(cx))}{10e^2} - \\
 & \quad \frac{d(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{bc \int \frac{(d - 4ex^2)(ex^2 + d)^4}{\sqrt{cx - 1} \sqrt{cx + 1}} dx}{40e^2} + \frac{(d + ex^2)^5 (a + \operatorname{barccosh}(cx))}{10e^2} - \frac{d(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e^2} \\
 & \quad \downarrow \text{2041} \\
 & \frac{bc \sqrt{c^2 x^2 - 1} \int \frac{(d - 4ex^2)(ex^2 + d)^4}{\sqrt{c^2 x^2 - 1}} dx}{40e^2 \sqrt{cx - 1} \sqrt{cx + 1}} + \frac{(d + ex^2)^5 (a + \operatorname{barccosh}(cx))}{10e^2} - \\
 & \quad \frac{d(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e^2} \\
 & \quad \downarrow \text{403} \\
 & \frac{bc \sqrt{c^2 x^2 - 1} \left( \int \frac{2(ex^2 + d)^3 (d(5c^2 d - 2e) - e(11dc^2 + 18e)x^2)}{\sqrt{c^2 x^2 - 1} 10c^2} dx - \frac{2ex \sqrt{c^2 x^2 - 1} (d + ex^2)^4}{5c^2} \right)}{40e^2 \sqrt{cx - 1} \sqrt{cx + 1}} + \\
 & \quad \frac{(d + ex^2)^5 (a + \operatorname{barccosh}(cx))}{10e^2} - \frac{d(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e^2} \\
 & \quad \downarrow \text{27} \\
 & \frac{bc \sqrt{c^2 x^2 - 1} \left( \int \frac{(ex^2 + d)^3 (d(5c^2 d - 2e) - e(11dc^2 + 18e)x^2)}{\sqrt{c^2 x^2 - 1} 5c^2} dx - \frac{2ex \sqrt{c^2 x^2 - 1} (d + ex^2)^4}{5c^2} \right)}{40e^2 \sqrt{cx - 1} \sqrt{cx + 1}} + \\
 & \quad \frac{(d + ex^2)^5 (a + \operatorname{barccosh}(cx))}{10e^2} - \frac{d(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e^2}
 \end{aligned}$$



$$\begin{aligned}
 & \downarrow 403 \\
 bc\sqrt{c^2x^2 - 1} & \left( \frac{\int \frac{(ex^2+d)^2(d(40d^2c^4-27dec^2-18e^2)-e(26d^2c^4+201dec^2+126e^2)x^2)}{\sqrt{c^2x^2-1}} dx}{8c^2} - \frac{ex\sqrt{c^2x^2-1}(11c^2d+18e)(d+ex^2)^3}{8c^2} - \frac{2ex\sqrt{c^2x^2-1}(d+ex^2)^3}{5c^2} \right) \\
 & \frac{(d+ex^2)^5(a+\operatorname{barccosh}(cx))}{10e^2} - \frac{40e^2\sqrt{cx-1}\sqrt{cx+1}d(d+ex^2)^4(a+\operatorname{barccosh}(cx))}{8e^2} \\
 & \downarrow 403 \\
 bc\sqrt{c^2x^2 - 1} & \left( \frac{\int \frac{(ex^2+d)(e(136d^3c^6-1096d^2ec^4-1617de^2c^2-630e^3)x^2+d(240d^3c^6-188d^2ec^4-309de^2c^2-126e^3))}{\sqrt{c^2x^2-1}} dx}{6c^2} - \frac{ex\sqrt{c^2x^2-1}(26c^4d^2+201c^2de+18e^2d^2+18e^2d+18e^2)}{6c^2} - \frac{2ex\sqrt{c^2x^2-1}(d+ex^2)^3}{5c^2} \right) \\
 & \frac{(d+ex^2)^5(a+\operatorname{barccosh}(cx))}{10e^2} - \frac{40e^2\sqrt{cx-1}\sqrt{cx+1}d(d+ex^2)^4(a+\operatorname{barccosh}(cx))}{8e^2} \\
 & \downarrow 403 \\
 bc\sqrt{c^2x^2 - 1} & \left( \frac{\int \frac{e(1232d^4c^8-2536d^3ec^6-7758d^2e^2c^4-6615de^3c^2-1890e^4)x^2+d(960d^4c^8-616d^3ec^6-2332d^2e^2c^4-2121de^3c^2-630e^4)}{\sqrt{c^2x^2-1}} dx}{4c^2} + \frac{ex\sqrt{c^2x^2-1}(1232c^8d^4-2536c^6d^3e-7758c^4d^2e^2-6615c^2de^3-1890e^4)}{4c^2} + \frac{2ex\sqrt{c^2x^2-1}(d+ex^2)^3}{5c^2} \right) \\
 & \frac{(d+ex^2)^5(a+\operatorname{barccosh}(cx))}{10e^2} - \frac{40e^2\sqrt{cx-1}\sqrt{cx+1}d(d+ex^2)^4(a+\operatorname{barccosh}(cx))}{8e^2} \\
 & \downarrow 299 \\
 bc\sqrt{c^2x^2 - 1} & \left( \frac{15(128c^{10}d^5-480c^6d^3e^2-800c^4d^2e^3-525c^2de^4-126e^5)}{2c^2} \int \frac{1}{\sqrt{c^2x^2-1}} dx + \frac{ex\sqrt{c^2x^2-1}(1232c^8d^4-2536c^6d^3e-7758c^4d^2e^2-6615c^2de^3-1890e^4)}{4c^2} + \frac{2ex\sqrt{c^2x^2-1}(d+ex^2)^3}{6c^2} + \frac{2ex\sqrt{c^2x^2-1}(d+ex^2)^3}{8c^2} \right) \\
 & \frac{(d+ex^2)^5(a+\operatorname{barccosh}(cx))}{10e^2} - \frac{40e^2\sqrt{cx-1}\sqrt{cx+1}d(d+ex^2)^4(a+\operatorname{barccosh}(cx))}{8e^2} \\
 & \downarrow 224
 \end{aligned}$$

3.480.  $\int x^3(d+ex^2)^3(a+\operatorname{barccosh}(cx)) dx$

$$bc\sqrt{c^2x^2 - 1} \left( \frac{15(128c^{10}d^5 - 480c^6d^3e^2 - 800c^4d^2e^3 - 525c^2de^4 - 126e^5)}{2c^2} \int \frac{1}{1 - \frac{c^2x^2}{c^2x^2 - 1}} d \frac{x}{\sqrt{c^2x^2 - 1}} + \frac{ex\sqrt{c^2x^2 - 1}(1232c^8d^4 - 2536c^6d^3e - 7758c^4d^2e^2 - 6615c^2de^3 - 1890e^4)}{2c^2} \right)$$

$$\frac{(d + ex^2)^5 (a + \operatorname{barccosh}(cx))}{10e^2} - \frac{d(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e^2}$$

219

$$bc\sqrt{c^2x^2 - 1} \left( \frac{(d + ex^2)^5 (a + \operatorname{barccosh}(cx))}{10e^2} - \frac{d(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e^2} + \frac{15\operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2 - 1}}\right)(128c^{10}d^5 - 480c^6d^3e^2 - 800c^4d^2e^3 - 525c^2de^4 - 126e^5)}{2c^3} + \frac{ex\sqrt{c^2x^2 - 1}(1232c^8d^4 - 2536c^6d^3e - 7758c^4d^2e^2 - 6615c^2de^3 - 1890e^4)}{2c^2} \right)$$

input `Int[x^3*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/8*(d*(d + e*x^2)^4*(a + b*ArcCosh[c*x]))/e^2 + ((d + e*x^2)^5*(a + b*ArcCosh[c*x]))/(10*e^2) + (b*c*sqrt[-1 + c^2*x^2]*((-2*e*x*sqrt[-1 + c^2*x^2])*(d + e*x^2)^4)/(5*c^2) + (-1/8*(e*(11*c^2*d + 18*e)*x*sqrt[-1 + c^2*x^2])*(d + e*x^2)^3)/c^2 + (-1/6*(e*(26*c^4*d^2 + 201*c^2*d*e + 126*e^2)*x*sqrt[-1 + c^2*x^2])*(d + e*x^2)^2)/c^2 + ((e*(136*c^6*d^3 - 1096*c^4*d^2*e - 1617*c^2*d*e^2 - 630*e^3)*x*sqrt[-1 + c^2*x^2])*(d + e*x^2))/(4*c^2) + ((e*(1232*c^8*d^4 - 2536*c^6*d^3*e - 7758*c^4*d^2*e^2 - 6615*c^2*d*e^3 - 1890*e^4)*x*sqrt[-1 + c^2*x^2])/(2*c^2) + (15*(128*c^10*d^5 - 480*c^6*d^3*e^2 - 800*c^4*d^2*e^3 - 525*c^2*d*e^4 - 126*e^5)*ArcTanh[(c*x)/sqrt[-1 + c^2*x^2]])/(2*c^3))/(4*c^2)/(6*c^2)/(8*c^2)/(5*c^2))/(40*e^2*sqrt[-1 + c*x]*sqrt[1 + c*x])`

## 3.480.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`
- rule 403 `Int[((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[f*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(b*(2*(p + q + 1) + 1))), x] + Simp[1/(b*(2*(p + q + 1) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[c*(b*e - a*f + b*e*2*(p + q + 1)) + (d*(b*e - a*f) + f*2*q*(b*c - a*d) + b*d*e*2*(p + q + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && GtQ[q, 0] && NeQ[2*(p + q + 1) + 1, 0]`
- rule 2041 `Int[((e1_) + (f1_.)*(x_)^(n2_.))^(r_.)*((e2_) + (f2_.)*(x_)^(n2_.))^(r_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[(e1 + f1*x^(n/2))^FracPart[r]*((e2 + f2*x^(n/2))^FracPart[r]/(e1*e2 + f1*f2*x^n)^FracPart[r]) Int[(a + b*x^n)^p*(c + d*x^n)^q*(e1*e2 + f1*f2*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e1, f1, e2, f2, n, p, q, r}, x] && EqQ[n2, n/2] && EqQ[e2*f1 + e1*f2, 0]`
- rule 6373 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^(m)*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LtQ[m + p, 0]))`

### 3.480.4 Maple [A] (verified)

Time = 0.77 (sec) , antiderivative size = 538, normalized size of antiderivative = 1.09

method	result
parts	$a\left(\frac{1}{10}e^3x^{10} + \frac{3}{8}de^2x^8 + \frac{1}{2}d^2ex^6 + \frac{1}{4}d^3x^4\right) + \frac{b\left(\frac{c^4 \operatorname{arccosh}(cx)e^3x^{10}}{10} + \frac{3c^4 \operatorname{arccosh}(cx)de^2x^8}{8} + \frac{c^4 \operatorname{arccosh}(cx)d^2ex^6}{2}\right)}{c^6}$
derivativedivides	$\frac{a\left(\frac{1}{4}c^{10}d^3x^4 + \frac{1}{2}c^{10}d^2ex^6 + \frac{3}{8}c^{10}de^2x^8 + \frac{1}{10}c^{10}e^3x^{10}\right)}{c^6} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^{10}d^3x^4}{4} + \frac{\operatorname{arccosh}(cx)c^{10}d^2ex^6}{2} + \frac{3 \operatorname{arccosh}(cx)c^{10}de^2x^8}{8}\right)}{c^6}$
default	$\frac{a\left(\frac{1}{4}c^{10}d^3x^4 + \frac{1}{2}c^{10}d^2ex^6 + \frac{3}{8}c^{10}de^2x^8 + \frac{1}{10}c^{10}e^3x^{10}\right)}{c^6} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^{10}d^3x^4}{4} + \frac{\operatorname{arccosh}(cx)c^{10}d^2ex^6}{2} + \frac{3 \operatorname{arccosh}(cx)c^{10}de^2x^8}{8}\right)}{c^6}$

input `int(x^3*(e*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/10*e^3*x^10+3/8*d*e^2*x^8+1/2*d^2*e*x^6+1/4*d^3*x^4)+b/c^4*(1/10*c^4*arccosh(c*x)*e^3*x^10+3/8*c^4*arccosh(c*x)*d*e^2*x^8+1/2*c^4*arccosh(c*x)*d^2*e*x^6+1/4*arccosh(c*x)*c^4*x^4*d^3-1/76800/c^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(4800*c^9*d^3*(c^2*x^2-1)^(1/2)*x^3+6400*c^9*d^2*e*(c^2*x^2-1)^(1/2)*x^5+3600*c^9*d*e^2*(c^2*x^2-1)^(1/2)*x^7+768*e^3*(c^2*x^2-1)^(1/2)*c^9*x^9+7200*c^7*d^3*x*(c^2*x^2-1)^(1/2)+8000*(c^2*x^2-1)^(1/2)*c^7*d^2*e*x^3+4200*(c^2*x^2-1)^(1/2)*c^7*d*e^2*x^5+864*e^3*c^7*x^7*(c^2*x^2-1)^(1/2)+7200*c^6*d^3*ln(c*x+(c^2*x^2-1)^(1/2))+12000*c^5*d^2*e*x*(c^2*x^2-1)^(1/2)+5250*c^5*d*e^2*(c^2*x^2-1)^(1/2)*x^3+1008*e^3*(c^2*x^2-1)^(1/2)*c^5*x^5+12000*c^4*d^2*e*ln(c*x+(c^2*x^2-1)^(1/2))+7875*c^3*d*e^2*x*(c^2*x^2-1)^(1/2)+1260*e^3*c^3*x^3*(c^2*x^2-1)^(1/2)+7875*c^2*d*e^2*ln(c*x+(c^2*x^2-1)^(1/2))+1890*e^3*c*x*(c^2*x^2-1)^(1/2)+1890*e^3*ln(c*x+(c^2*x^2-1)^(1/2)))/(c^2*x^2-1)^(1/2)`

### 3.480.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 330, normalized size of antiderivative = 0.67

$$\int x^3(d + ex^2)^3(a + b\operatorname{arccosh}(cx)) dx = \frac{7680 ac^{10}e^3x^{10} + 28800 ac^{10}de^2x^8 + 38400 ac^{10}d^2ex^6 + 19200 ac^{10}d^3x^4 + 15(512 bc^{10}e^3x^{10} + 1920 bc^{10}de^2x^8 + \dots)}{c^6}$$

input `integrate(x^3*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`

3.480.  $\int x^3(d + ex^2)^3(a + b\operatorname{arccosh}(cx)) dx$

```
output 1/76800*(7680*a*c^10*e^3*x^10 + 28800*a*c^10*d*e^2*x^8 + 38400*a*c^10*d^2*
e*x^6 + 19200*a*c^10*d^3*x^4 + 15*(512*b*c^10*e^3*x^10 + 1920*b*c^10*d*e^2
*x^8 + 2560*b*c^10*d^2*e*x^6 + 1280*b*c^10*d^3*x^4 - 480*b*c^6*d^3 - 800*b
*c^4*d^2*e - 525*b*c^2*d*e^2 - 126*b*e^3)*log(c*x + sqrt(c^2*x^2 - 1)) - (
768*b*c^9*e^3*x^9 + 144*(25*b*c^9*d*e^2 + 6*b*c^7*e^3)*x^7 + 8*(800*b*c^9*
d^2*e + 525*b*c^7*d*e^2 + 126*b*c^5*e^3)*x^5 + 10*(480*b*c^9*d^3 + 800*b*c
^7*d^2*e + 525*b*c^5*d*e^2 + 126*b*c^3*e^3)*x^3 + 15*(480*b*c^7*d^3 + 800*
b*c^5*d^2*e + 525*b*c^3*d*e^2 + 126*b*c*e^3)*x)*sqrt(c^2*x^2 - 1))/c^10
```

### 3.480.6 Sympy [F]

$$\int x^3(d + ex^2)^3(a + \operatorname{barccosh}(cx)) dx = \int x^3(a + b \operatorname{acosh}(cx))(d + ex^2)^3 dx$$

```
input integrate(x**3*(e*x**2+d)**3*(a+b*acosh(c*x)),x)
```

```
output Integral(x**3*(a + b*acosh(c*x))*(d + e*x**2)**3, x)
```

### 3.480.7 Maxima [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 487, normalized size of antiderivative = 0.99

$$\begin{aligned} \int x^3(d + ex^2)^3(a + \operatorname{barccosh}(cx)) dx &= \frac{1}{10} ae^3x^{10} + \frac{3}{8} ade^2x^8 + \frac{1}{2} ad^2ex^6 + \frac{1}{4} ad^3x^4 \\ &+ \frac{1}{32} \left( 8x^4 \operatorname{arcosh}(cx) - \left( \frac{2\sqrt{c^2x^2 - 1}x^3}{c^2} + \frac{3\sqrt{c^2x^2 - 1}x}{c^4} + \frac{3 \log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^5} \right) c \right) bd^3 \\ &+ \frac{1}{96} \left( 48x^6 \operatorname{arcosh}(cx) - \left( \frac{8\sqrt{c^2x^2 - 1}x^5}{c^2} + \frac{10\sqrt{c^2x^2 - 1}x^3}{c^4} + \frac{15\sqrt{c^2x^2 - 1}x}{c^6} + \frac{15 \log(2c^2x + 2\sqrt{c^2x^2 - 1}c)}{c^7} \right) c \right) bd^3 \\ &+ \frac{1}{1024} \left( 384x^8 \operatorname{arcosh}(cx) - \left( \frac{48\sqrt{c^2x^2 - 1}x^7}{c^2} + \frac{56\sqrt{c^2x^2 - 1}x^5}{c^4} + \frac{70\sqrt{c^2x^2 - 1}x^3}{c^6} + \frac{105\sqrt{c^2x^2 - 1}x}{c^8} \right) c \right) bd^3 \\ &+ \frac{1}{12800} \left( 1280x^{10} \operatorname{arcosh}(cx) - \left( \frac{128\sqrt{c^2x^2 - 1}x^9}{c^2} + \frac{144\sqrt{c^2x^2 - 1}x^7}{c^4} + \frac{168\sqrt{c^2x^2 - 1}x^5}{c^6} + \frac{210\sqrt{c^2x^2 - 1}x^3}{c^8} \right) c \right) bd^3 \end{aligned}$$

```
input integrate(x^3*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")
```

```
output 1/10*a*e^3*x^10 + 3/8*a*d*e^2*x^8 + 1/2*a*d^2*e*x^6 + 1/4*a*d^3*x^4 + 1/32
*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*
x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^5)*c)*b*d^3 + 1/96*(48*x^
6*arccosh(c*x) - (8*sqrt(c^2*x^2 - 1)*x^5/c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c
^4 + 15*sqrt(c^2*x^2 - 1)*x/c^6 + 15*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/
c^7)*c)*b*d^2*e + 1/1024*(384*x^8*arccosh(c*x) - (48*sqrt(c^2*x^2 - 1)*x^7
/c^2 + 56*sqrt(c^2*x^2 - 1)*x^5/c^4 + 70*sqrt(c^2*x^2 - 1)*x^3/c^6 + 105*s
qrt(c^2*x^2 - 1)*x/c^8 + 105*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^9)*c)*
b*d*e^2 + 1/12800*(1280*x^10*arccosh(c*x) - (128*sqrt(c^2*x^2 - 1)*x^9/c^2
+ 144*sqrt(c^2*x^2 - 1)*x^7/c^4 + 168*sqrt(c^2*x^2 - 1)*x^5/c^6 + 210*sqr
t(c^2*x^2 - 1)*x^3/c^8 + 315*sqrt(c^2*x^2 - 1)*x/c^10 + 315*log(2*c^2*x +
2*sqrt(c^2*x^2 - 1)*c)/c^11)*c)*b*e^3
```

### 3.480.8 Giac [F(-2)]

Exception generated.

$$\int x^3 (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: TypeError}$$

```
input integrate(x^3*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

### 3.480.9 Mupad [F(-1)]

Timed out.

$$\int x^3 (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x^3 (a + b \operatorname{acosh}(cx)) (ex^2 + d)^3 dx$$

```
input int(x^3*(a + b*acosh(c*x))*(d + e*x^2)^3,x)
```

```
output int(x^3*(a + b*acosh(c*x))*(d + e*x^2)^3, x)
```

### 3.481 $\int x^2(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.481.1 Optimal result . . . . .	3542
3.481.2 Mathematica [A] (verified) . . . . .	3543
3.481.3 Rubi [A] (verified) . . . . .	3543
3.481.4 Maple [A] (verified) . . . . .	3546
3.481.5 Fricas [A] (verification not implemented) . . . . .	3546
3.481.6 Sympy [F] . . . . .	3547
3.481.7 Maxima [A] (verification not implemented) . . . . .	3547
3.481.8 Giac [F(-2)] . . . . .	3548
3.481.9 Mupad [F(-1)] . . . . .	3548

#### 3.481.1 Optimal result

Integrand size = 21, antiderivative size = 365

$$\int x^2(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{b(105c^6d^3 + 189c^4d^2e + 135c^2de^2 + 35e^3)(1 - c^2x^2)}{315c^9\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{b(105c^6d^3 + 378c^4d^2e + 405c^2de^2 + 140e^3)(1 - c^2x^2)^2}{945c^9\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{be(63c^4d^2 + 135c^2de + 70e^2)(1 - c^2x^2)^3}{525c^9\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{be^2(27c^2d + 28e)(1 - c^2x^2)^4}{441c^9\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{be^3(1 - c^2x^2)^5}{81c^9\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{1}{3}d^3x^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}d^2ex^5(a + \operatorname{barccosh}(cx))$$

$$+ \frac{3}{7}de^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^3x^9(a + \operatorname{barccosh}(cx))$$

output  $\frac{1}{3}d^3x^3(a+b\operatorname{arccosh}(cx))+\frac{3}{5}d^2ex^5(a+b\operatorname{arccosh}(cx))+\frac{3}{7}d^2ex^7(a+b\operatorname{arccosh}(cx))+\frac{1}{9}e^3x^9(a+b\operatorname{arccosh}(cx))+\frac{1}{315}b(105c^6d^3+189c^4d^2e+135c^2de^2+35e^3)(-c^2x^2+1)/c^9/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-1/945b(105c^6d^3+378c^4d^2e+405c^2de^2+140e^3)(-c^2x^2+1)^2/c^9/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}+1/525b(63c^4d^2+135c^2de+70e^2)(-c^2x^2+1)^3/c^9/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}-1/441be^2(27c^2d+28e)(-c^2x^2+1)^4/c^9/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}+1/81be^3(-c^2x^2+1)^5/c^9/(cx-1)^{(1/2)}/(cx+1)^{(1/2)}$

**3.481.2 Mathematica [A] (verified)**

Time = 0.23 (sec) , antiderivative size = 236, normalized size of antiderivative = 0.65

$$\int x^2(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx$$

$$= \frac{315ax^3(105d^3+189d^2ex^2+135de^2x^4+35e^3x^6) - \frac{b\sqrt{-1+cx}\sqrt{1+cx}(4480e^3+80c^2e^2(243d+28ex^2)+24c^4e(1323d^2+405dex^2+70e^2x^4))+2c^6(11025d^3+7938d^2ex^2+3645de^2x^4+700e^3x^6)+c^8(11025d^3x^2+11907d^2ex^4+6075de^2x^6+1225e^3x^8))}{c^9} + 315bx^3(105d^3+189d^2ex^2+135de^2x^4+35e^3x^6)\operatorname{ArcCosh}[cx]}{99225}$$

input `Integrate[x^2*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `(315*a*x^3*(105*d^3 + 189*d^2*e*x^2 + 135*d*e^2*x^4 + 35*e^3*x^6) - (b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(4480*e^3 + 80*c^2*e^2*(243*d + 28*e*x^2) + 24*c^4*e*(1323*d^2 + 405*d*e*x^2 + 70*e^2*x^4) + 2*c^6*(11025*d^3 + 7938*d^2*e*x^2 + 3645*d*e^2*x^4 + 700*e^3*x^6) + c^8*(11025*d^3*x^2 + 11907*d^2*e*x^4 + 6075*d*e^2*x^6 + 1225*e^3*x^8)))/c^9 + 315*b*x^3*(105*d^3 + 189*d^2*e*x^2 + 135*d*e^2*x^4 + 35*e^3*x^6)*ArcCosh[c*x])/99225`

**3.481.3 Rubi [A] (verified)**Time = 0.88 (sec) , antiderivative size = 313, normalized size of antiderivative = 0.86, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6373, 27, 2113, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^2(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx$$

$$\downarrow \text{6373}$$

$$-bc \int \frac{x^3(35e^3x^6+135de^2x^4+189d^2ex^2+105d^3)}{315\sqrt{cx-1}\sqrt{cx+1}}dx + \frac{1}{3}d^3x^3(a+\operatorname{barccosh}(cx)) + \frac{3}{5}d^2ex^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}de^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{9}e^3x^9(a+\operatorname{barccosh}(cx))$$

$$\downarrow \text{27}$$

$$-\frac{1}{315}bc \int \frac{x^3(35e^3x^6+135de^2x^4+189d^2ex^2+105d^3)}{\sqrt{cx-1}\sqrt{cx+1}}dx + \frac{1}{3}d^3x^3(a+\operatorname{barccosh}(cx)) + \frac{3}{5}d^2ex^5(a+\operatorname{barccosh}(cx)) + \frac{3}{7}de^2x^7(a+\operatorname{barccosh}(cx)) + \frac{1}{9}e^3x^9(a+\operatorname{barccosh}(cx))$$

---

3.481.  $\int x^2(d+ex^2)^3(a+\operatorname{barccosh}(cx))dx$



$$\begin{aligned}
& \downarrow \text{2113} \\
& -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^3(35e^3x^6+135de^2x^4+189d^2ex^2+105d^3)}{\sqrt{c^2x^2-1}} dx}{315\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}d^3x^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}d^2ex^5(a + \\
& \operatorname{barccosh}(cx)) + \frac{3}{7}de^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^3x^9(a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{2331} \\
& -\frac{bc\sqrt{c^2x^2-1} \int \frac{x^2(35e^3x^6+135de^2x^4+189d^2ex^2+105d^3)}{\sqrt{c^2x^2-1}} dx^2}{630\sqrt{cx-1}\sqrt{cx+1}} + \frac{1}{3}d^3x^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}d^2ex^5(a + \\
& \operatorname{barccosh}(cx)) + \frac{3}{7}de^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^3x^9(a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{2123} \\
& \frac{bc\sqrt{c^2x^2-1} \int \left( \frac{35e^3(c^2x^2-1)^{7/2}}{c^8} + \frac{5e^2(27dc^2+28e)(c^2x^2-1)^{5/2}}{c^8} + \frac{3e(63d^2c^4+135dec^2+70e^2)(c^2x^2-1)^{3/2}}{c^8} + \frac{(105d^3c^6+378d^2ec^4+}{630\sqrt{cx-1}\sqrt{cx+1}} \right. \\
& \left. \frac{1}{3}d^3x^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}d^2ex^5(a + \operatorname{barccosh}(cx)) + \frac{3}{7}de^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^3x^9(a + \operatorname{barccosh}(cx)) \right)}{630\sqrt{cx-1}\sqrt{cx+1}} \\
& \downarrow \text{2009} \\
& \frac{1}{3}d^3x^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}d^2ex^5(a + \operatorname{barccosh}(cx)) + \frac{3}{7}de^2x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^3x^9(a + \\
& \operatorname{barccosh}(cx)) - \\
& \frac{bc\sqrt{c^2x^2-1} \left( \frac{10e^2(c^2x^2-1)^{7/2}(27c^2d+28e)}{7c^{10}} + \frac{70e^3(c^2x^2-1)^{9/2}}{9c^{10}} + \frac{6e(c^2x^2-1)^{5/2}(63c^4d^2+135c^2de+70e^2)}{5c^{10}} + \frac{2(c^2x^2-1)^{3/2}(105c^6d^3+}{630\sqrt{cx-1}\sqrt{cx+1}} \right)}{630\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[x^2*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `-1/630*(b*c*Sqrt[-1 + c^2*x^2]*((2*(105*c^6*d^3 + 189*c^4*d^2*e + 135*c^2*d*e^2 + 35*e^3)*Sqrt[-1 + c^2*x^2])/c^10 + (2*(105*c^6*d^3 + 378*c^4*d^2*e + 405*c^2*d*e^2 + 140*e^3)*(-1 + c^2*x^2)^(3/2))/(3*c^10) + (6*e*(63*c^4*d^2 + 135*c^2*d*e + 70*e^2)*(-1 + c^2*x^2)^(5/2))/(5*c^10) + (10*e^2*(27*c^2*d + 28*e)*(-1 + c^2*x^2)^(7/2))/(7*c^10) + (70*e^3*(-1 + c^2*x^2)^(9/2))/(9*c^10))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (d^3*x^3*(a + b*ArcCosh[c*x]))/3 + (3*d^2*e*x^5*(a + b*ArcCosh[c*x]))/5 + (3*d*e^2*x^7*(a + b*ArcCosh[c*x]))/7 + (e^3*x^9*(a + b*ArcCosh[c*x]))/9`

---

3.481.  $\int x^2(d + ex^2)^3(a + \operatorname{barccosh}(cx)) dx$

## 3.481.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2113 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[P_x*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`
- rule 2123 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[P_x*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[P_x, x] && (IntegersQ[m, n] || IGtQ[m, -2])`
- rule 2331 `Int[(P_q)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, P_q, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[P_q, x^2] && IntegerQ[(m - 1)/2]`
- rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.481.4 Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 273, normalized size of antiderivative = 0.75

method	result
parts	$a\left(\frac{1}{9}e^3x^9 + \frac{3}{7}de^2x^7 + \frac{3}{5}d^2ex^5 + \frac{1}{3}d^3x^3\right) + \frac{b\left(\frac{c^3\operatorname{arccosh}(cx)e^3x^9}{9} + \frac{3c^3\operatorname{arccosh}(cx)de^2x^7}{7} + \frac{3c^3\operatorname{arccosh}(cx)d^2e}{5}\right)}{c^6}$
derivativedivides	$\frac{a\left(\frac{1}{3}c^9d^3x^3 + \frac{3}{5}c^9d^2ex^5 + \frac{3}{7}c^9de^2x^7 + \frac{1}{9}e^3c^9x^9\right)}{c^6} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^9d^3x^3}{3} + \frac{3\operatorname{arccosh}(cx)c^9d^2ex^5}{5} + \frac{3\operatorname{arccosh}(cx)c^9de^2x^7}{7} + \frac{\operatorname{arccosh}(cx)c^9e^3}{9}\right)}{c^6}$
default	$\frac{a\left(\frac{1}{3}c^9d^3x^3 + \frac{3}{5}c^9d^2ex^5 + \frac{3}{7}c^9de^2x^7 + \frac{1}{9}e^3c^9x^9\right)}{c^6} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^9d^3x^3}{3} + \frac{3\operatorname{arccosh}(cx)c^9d^2ex^5}{5} + \frac{3\operatorname{arccosh}(cx)c^9de^2x^7}{7} + \frac{\operatorname{arccosh}(cx)c^9e^3}{9}\right)}{c^6}$

input `int(x^2*(e*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/9*e^3*x^9+3/7*d*e^2*x^7+3/5*d^2*e*x^5+1/3*d^3*x^3)+b/c^3*(1/9*c^3*arccosh(c*x)*e^3*x^9+3/7*c^3*arccosh(c*x)*d*e^2*x^7+3/5*c^3*arccosh(c*x)*d^2*e*x^5+1/3*arccosh(c*x)*c^3*x^3*d^3-1/99225/c^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(1225*c^8*e^3*x^8+6075*c^8*d*e^2*x^6+11907*c^8*d^2*e*x^4+1400*c^6*e^3*x^6+11025*c^8*d^3*x^2+7290*c^6*d*e^2*x^4+15876*c^6*d^2*e*x^2+1680*c^4*e^3*x^4+22050*c^6*d^3+9720*c^4*d*e^2*x^2+31752*c^4*d^2*e+2240*c^2*e^3*x^2+19440*c^2*d*e^2+4480*e^3))`

### 3.481.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 289, normalized size of antiderivative = 0.79

$$\int x^2(d + ex^2)^3(a + b\operatorname{arccosh}(cx)) dx$$

$$= \frac{11025 ac^9 e^3 x^9 + 42525 ac^9 d e^2 x^7 + 59535 ac^9 d^2 e x^5 + 33075 ac^9 d^3 x^3 + 315 (35 bc^9 e^3 x^9 + 135 bc^9 d e^2 x^7 + 18 bc^9 d^2 e x^5 + 18 bc^9 d^3 x^3)}{c^6}$$

input `integrate(x^2*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $\frac{1}{99225}(11025ac^9e^3x^9 + 42525ac^9d^2e^2x^7 + 59535ac^9d^2e^2x^5 + 33075ac^9d^3x^3 + 315(35b^2c^9e^3x^9 + 135b^2c^9d^2e^2x^7 + 189b^2c^9d^2e^2x^5 + 105b^2c^9d^3x^3))\log(cx + \sqrt{c^2x^2 - 1}) - (1225b^2c^8e^3x^8 + 22050b^2c^6d^3 + 31752b^2c^4d^2e + 25(243b^2c^8d^2e^2 + 56b^2c^6e^3))x^6 + 19440b^2c^2d^2e^2 + 3(3969b^2c^8d^2e + 2430b^2c^6d^2e^2 + 560b^2c^4e^3)x^4 + 4480b^2e^3 + (11025b^2c^8d^3 + 15876b^2c^6d^2e + 9720b^2c^4d^2e^2 + 2240b^2c^2e^3)x^2)\sqrt{c^2x^2 - 1}/c^9$

### 3.481.6 Sympy [F]

$$\int x^2(d + ex^2)^3(a + \operatorname{barccosh}(cx)) dx = \int x^2(a + b \operatorname{acosh}(cx))(d + ex^2)^3 dx$$

input `integrate(x**2*(e*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `Integral(x**2*(a + b*acosh(c*x))*(d + e*x**2)**3, x)`

### 3.481.7 Maxima [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.02

$$\begin{aligned} \int x^2(d + ex^2)^3(a + \operatorname{barccosh}(cx)) dx &= \frac{1}{9}ae^3x^9 + \frac{3}{7}ade^2x^7 + \frac{3}{5}ad^2ex^5 \\ &+ \frac{1}{3}ad^3x^3 + \frac{1}{9}\left(3x^3 \operatorname{arcosh}(cx) - c\left(\frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4}\right)\right)bd^3 \\ &+ \frac{1}{25}\left(15x^5 \operatorname{arcosh}(cx) - \left(\frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6}\right)c\right)bd^2e \\ &+ \frac{3}{245}\left(35x^7 \operatorname{arcosh}(cx) - \left(\frac{5\sqrt{c^2x^2 - 1}x^6}{c^2} + \frac{6\sqrt{c^2x^2 - 1}x^4}{c^4} + \frac{8\sqrt{c^2x^2 - 1}x^2}{c^6} + \frac{16\sqrt{c^2x^2 - 1}}{c^8}\right)c\right)bde \\ &+ \frac{1}{2835}\left(315x^9 \operatorname{arcosh}(cx) - \left(\frac{35\sqrt{c^2x^2 - 1}x^8}{c^2} + \frac{40\sqrt{c^2x^2 - 1}x^6}{c^4} + \frac{48\sqrt{c^2x^2 - 1}x^4}{c^6} + \frac{64\sqrt{c^2x^2 - 1}x^2}{c^8}\right)c\right)bd^2e \end{aligned}$$

input `integrate(x^2*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

```
output 1/9*a*e^3*x^9 + 3/7*a*d*e^2*x^7 + 3/5*a*d^2*e*x^5 + 1/3*a*d^3*x^3 + 1/9*(3
*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4
))*b*d^3 + 1/25*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sq
rt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d^2*e + 3/245*(35*
x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/
c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*d*e^2 +
1/2835*(315*x^9*arccosh(c*x) - (35*sqrt(c^2*x^2 - 1)*x^8/c^2 + 40*sqrt(c^
2*x^2 - 1)*x^6/c^4 + 48*sqrt(c^2*x^2 - 1)*x^4/c^6 + 64*sqrt(c^2*x^2 - 1)*x
^2/c^8 + 128*sqrt(c^2*x^2 - 1)/c^10)*c)*b*e^3
```

### 3.481.8 Giac [F(-2)]

Exception generated.

$$\int x^2(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

```
input integrate(x^2*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve
cteur & l) Error: Bad Argument Value
```

### 3.481.9 Mupad [F(-1)]

Timed out.

$$\int x^2(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x^2 (a + b \operatorname{acosh}(cx)) (ex^2 + d)^3 dx$$

```
input int(x^2*(a + b*acosh(c*x))*(d + e*x^2)^3,x)
```

```
output int(x^2*(a + b*acosh(c*x))*(d + e*x^2)^3, x)
```

### 3.482 $\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.482.1 Optimal result . . . . .	3549
3.482.2 Mathematica [A] (warning: unable to verify) . . . . .	3550
3.482.3 Rubi [A] (verified) . . . . .	3550
3.482.4 Maple [A] (verified) . . . . .	3554
3.482.5 Fricas [A] (verification not implemented) . . . . .	3554
3.482.6 Sympy [F] . . . . .	3555
3.482.7 Maxima [A] (verification not implemented) . . . . .	3555
3.482.8 Giac [F(-2)] . . . . .	3556
3.482.9 Mupad [F(-1)] . . . . .	3556

#### 3.482.1 Optimal result

Integrand size = 19, antiderivative size = 358

$$\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{5b(2c^2d + e)(40c^4d^2 + 40c^2de + 21e^2)x(1 - c^2x^2)}{3072c^7\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{b(104c^4d^2 + 104c^2de + 35e^2)x(1 - c^2x^2)(d + ex^2)}{1536c^5\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{7b(2c^2d + e)x(1 - c^2x^2)(d + ex^2)^2}{384c^3\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bx(1 - c^2x^2)(d + ex^2)^3}{64c\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{(d + ex^2)^4(a + \operatorname{barccosh}(cx))}{8e} - \frac{b(128c^8d^4 + 256c^6d^3e + 288c^4d^2e^2 + 160c^2de^3 + 35e^4)\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{cx}{\sqrt{-1 + c^2x^2}}\right)}{1024c^8e\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output 1/8*(e*x^2+d)^4*(a+b*arccosh(c*x))/e+5/3072*b*(2*c^2*d+e)*(40*c^4*d^2+40*c^2*d*e+21*e^2)*x*(-c^2*x^2+1)/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/1536*b*(104*c^4*d^2+104*c^2*d*e+35*e^2)*x*(-c^2*x^2+1)*(e*x^2+d)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+7/384*b*(2*c^2*d+e)*x*(-c^2*x^2+1)*(e*x^2+d)^2/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/64*b*x*(-c^2*x^2+1)*(e*x^2+d)^3/c/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/1024*b*(128*c^8*d^4+256*c^6*d^3*e+288*c^4*d^2*e^2+160*c^2*d*e^3+35*e^4)*arctanh(c*x/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/c^8/e/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.482.2 Mathematica [A] (warning: unable to verify)**

Time = 0.28 (sec) , antiderivative size = 256, normalized size of antiderivative = 0.72

$$\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{cx(384ac^7x(4d^3 + 6d^2ex^2 + 4de^2x^4 + e^3x^6) - b\sqrt{-1 + cx}\sqrt{1 + cx}(105e^3 + 10c^2e^2(48d + 7ex^2) + 8c^4e(108d^2 + 40d*ex^2 + 7e^2x^4) + 16c^6(48d^3 + 36d^2*ex^2 + 16d*e^2x^4 + 3e^3x^6))) + 384b*c^8*x^2*(4d^3 + 6d^2*ex^2 + 4d*e^2*x^4 + e^3*x^6)*\operatorname{ArcCosh}[c*x] - 6*b*(256*c^6*d^3 + 288*c^4*d^2*e + 160*c^2*d*e^2 + 35*e^3)*\operatorname{ArcTanh}[\operatorname{Sqrt}[(-1 + cx)/(1 + cx)]])}{(3072*c^8)}$$

input `Integrate[x*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`output `(c*x*(384*a*c^7*x*(4*d^3 + 6*d^2*e*x^2 + 4*d*e^2*x^4 + e^3*x^6) - b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(105*e^3 + 10*c^2*e^2*(48*d + 7*e*x^2) + 8*c^4*e*(108*d^2 + 40*d*e*x^2 + 7*e^2*x^4) + 16*c^6*(48*d^3 + 36*d^2*e*x^2 + 16*d*e^2*x^4 + 3*e^3*x^6))) + 384*b*c^8*x^2*(4*d^3 + 6*d^2*e*x^2 + 4*d*e^2*x^4 + e^3*x^6)*ArcCosh[c*x] - 6*b*(256*c^6*d^3 + 288*c^4*d^2*e + 160*c^2*d*e^2 + 35*e^3)*ArcTanh[Sqrt[(-1 + c*x)/(1 + c*x)]])/(3072*c^8)`**3.482.3 Rubi [A] (verified)**Time = 0.55 (sec) , antiderivative size = 328, normalized size of antiderivative = 0.92, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$ , Rules used = {6372, 648, 318, 403, 403, 299, 224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow 6372$$

$$\frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e} - \frac{bc \int \frac{(ex^2+d)^4}{\sqrt{cx-1}\sqrt{cx+1}} dx}{8e}$$

$$\downarrow 648$$

$$\frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e} - \frac{bc\sqrt{c^2x^2 - 1} \int \frac{(ex^2+d)^4}{\sqrt{c^2x^2 - 1}} dx}{8e\sqrt{cx - 1}\sqrt{cx + 1}}$$

$$\downarrow 318$$

$$\begin{aligned}
 & \frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\int \frac{(ex^2+d)^2 (7e(2dc^2+e)x^2+d(8dc^2+e))}{\sqrt{c^2x^2-1}} dx}{8c^2} + \frac{ex\sqrt{c^2x^2-1}(d+ex^2)^3}{8c^2} \right)} \\
 & \qquad \qquad \qquad \frac{8e\sqrt{cx-1}\sqrt{cx+1}}{\qquad \qquad \qquad \downarrow \text{403}} \\
 & \frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\int \frac{(ex^2+d)(e(104d^2c^4+104dec^2+35e^2)x^2+d(48d^2c^4+20dec^2+7e^2))}{\sqrt{c^2x^2-1}} dx}{6c^2} + \frac{7ex\sqrt{c^2x^2-1}(2c^2d+e)(d+ex^2)^2}{6e^2} + \frac{ex\sqrt{c^2x^2-1}(d+ex^2)^3}{8c^2} \right)} \\
 & \qquad \qquad \qquad \frac{8e\sqrt{cx-1}\sqrt{cx+1}}{\qquad \qquad \qquad \downarrow \text{403}} \\
 & \frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{\int \frac{5e(2dc^2+e)(40d^2c^4+40dec^2+21e^2)x^2+d(192d^3c^6+184d^2ec^4+132de^2c^2+35e^3)}{\sqrt{c^2x^2-1}} dx}{4c^2} + \frac{ex\sqrt{c^2x^2-1}(104c^4d^2+104c^2de+35e^2)(d+ex^2)}{4c^2} + \frac{ex\sqrt{c^2x^2-1}(d+ex^2)^3}{8c^2} \right)} \\
 & \qquad \qquad \qquad \frac{8e\sqrt{cx-1}\sqrt{cx+1}}{\qquad \qquad \qquad \downarrow \text{299}} \\
 & \frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{bc\sqrt{c^2x^2 - 1} \left( \frac{3(128c^8d^4+256c^6d^3e+288c^4d^2e^2+160c^2de^3+35e^4) \int \frac{1}{\sqrt{c^2x^2-1}} dx}{2c^2} + \frac{5ex\sqrt{c^2x^2-1}(2c^2d+e)(40c^4d^2+40c^2de+21e^2)}{2c^2} + \frac{ex\sqrt{c^2x^2-1}(104c^4d^2+104c^2de+35e^2)(d+ex^2)^3}{8c^2} \right)} \\
 & \qquad \qquad \qquad \frac{8e\sqrt{cx-1}\sqrt{cx+1}}{\qquad \qquad \qquad \downarrow \text{224}}
 \end{aligned}$$

---

3.482.  $\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$



$$bc\sqrt{c^2x^2 - 1} \left( \frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e} - \frac{3(128c^8d^4 + 256c^6d^3e + 288c^4d^2e^2 + 160c^2de^3 + 35e^4) \int \frac{1 - \frac{c^2x^2}{c^2x^2 - 1} d - \frac{x}{\sqrt{c^2x^2 - 1}}}{2c^2} + \frac{5ex\sqrt{c^2x^2 - 1}(2c^2d + e)(40c^4d^2 + 40c^2de + 21e^2)}{4c^2} + \frac{ex\sqrt{c^2x^2 - 1}}{6c^2} + \frac{ex\sqrt{c^2x^2 - 1}}{8c^2} \right)$$


---


$$8e\sqrt{cx - 1}\sqrt{cx + 1}$$

↓ 219

$$bc\sqrt{c^2x^2 - 1} \left( \frac{(d + ex^2)^4 (a + \operatorname{barccosh}(cx))}{8e} - \frac{3\operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2 - 1}}\right)(128c^8d^4 + 256c^6d^3e + 288c^4d^2e^2 + 160c^2de^3 + 35e^4)}{2c^3} + \frac{5ex\sqrt{c^2x^2 - 1}(2c^2d + e)(40c^4d^2 + 40c^2de + 21e^2)}{4c^2} + \frac{ex\sqrt{c^2x^2 - 1}}{6c^2} + \frac{ex\sqrt{c^2x^2 - 1}}{8c^2} \right)$$


---


$$8e\sqrt{cx - 1}\sqrt{cx + 1}$$

input `Int[x*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

output `((d + e*x^2)^4*(a + b*ArcCosh[c*x]))/(8*e) - (b*c*Sqrt[-1 + c^2*x^2]*((e*x*Sqrt[-1 + c^2*x^2]*(d + e*x^2)^3)/(8*c^2) + ((7*e*(2*c^2*d + e))*x*Sqrt[-1 + c^2*x^2]*(d + e*x^2)^2)/(6*c^2) + ((e*(104*c^4*d^2 + 104*c^2*d*e + 35*e^2))*x*Sqrt[-1 + c^2*x^2]*(d + e*x^2))/(4*c^2) + ((5*e*(2*c^2*d + e))*(40*c^4*d^2 + 40*c^2*d*e + 21*e^2))*x*Sqrt[-1 + c^2*x^2])/(2*c^2) + (3*(128*c^8*d^4 + 256*c^6*d^3*e + 288*c^4*d^2*e^2 + 160*c^2*d*e^3 + 35*e^4)*ArcTanh[(c*x)/Sqrt[-1 + c^2*x^2]])/(2*c^3)/(4*c^2)/(6*c^2)/(8*c^2))/(8*e*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.482.3.1 Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

---

3.482.  $\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

- rule 299 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*x*((a + b*x^2)^(p + 1)/(b*(2*p + 3))), x] - Simp[(a*d - b*c*(2*p + 3))/(b*(2*p + 3)) Int[(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NeQ[2*p + 3, 0]`
- rule 318 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[d*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q - 1)/(b*(2*(p + q) + 1))), x] + Simp[1/(b*(2*(p + q) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 2)*Simp[c*(b*c*(2*(p + q) + 1) - a*d) + d*(b*c*(2*(p + 2*q - 1) + 1) - a*d*(2*(q - 1) + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[2*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, 2, p, q, x]`
- rule 403 `Int[((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.)*((e_) + (f_.)*(x_)^2), x_Symbol] := Simp[f*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^q/(b*(2*(p + q + 1) + 1))), x] + Simp[1/(b*(2*(p + q + 1) + 1)) Int[(a + b*x^2)^p*(c + d*x^2)^(q - 1)*Simp[c*(b*e - a*f + b*e*2*(p + q + 1)) + (d*(b*e - a*f) + f*2*q*(b*c - a*d) + b*d*e*2*(p + q + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && GtQ[q, 0] && NeQ[2*(p + q + 1) + 1, 0]`
- rule 648 `Int[((c_) + (d_.)*(x_)^(m_.))*((e_) + (f_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c + d*x)^FracPart[m]*((e + f*x)^FracPart[m]/(c*e + d*f*x^2)^FracPart[m]) Int[(c*e + d*f*x^2)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m, n] && EqQ[d*e + c*f, 0] && !(EqQ[p, 2] && LtQ[m, -1])`
- rule 6372 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])/(2*e*(p + 1))), x] - Simp[b*(c/(2*e*(p + 1))) Int[(d + e*x^2)^(p + 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[c^2*d + e, 0] && NeQ[p, -1]`

### 3.482.4 Maple [A] (verified)

Time = 0.76 (sec) , antiderivative size = 476, normalized size of antiderivative = 1.33

method	result
parts	$\frac{a(e^2x^2+d)^4}{8e} + \frac{b\left(\frac{c^2e^3 \operatorname{arccosh}(cx)x^8}{8} + \frac{c^2e^2 \operatorname{arccosh}(cx)x^6d}{2} + \frac{3c^2e \operatorname{arccosh}(cx)x^4d^2}{4} + \frac{\operatorname{arccosh}(cx)c^2x^2d^3}{2} + \frac{c^2 \operatorname{arccosh}(cx)d^4}{8e}\right)}{8e^6}$
derivativedivides	$\frac{a(c^2ex^2+c^2d)^4}{8c^6e} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^8d^4}{8e} + \frac{\operatorname{arccosh}(cx)c^8d^3x^2}{2} + \frac{3e \operatorname{arccosh}(cx)c^8d^2x^4}{4} + \frac{e^2 \operatorname{arccosh}(cx)c^8dx^6}{2} + \frac{e^3 \operatorname{arccosh}(cx)c^8x^8}{8}\right)}{8c^6e}$
default	$\frac{a(c^2ex^2+c^2d)^4}{8c^6e} + \frac{b\left(\frac{\operatorname{arccosh}(cx)c^8d^4}{8e} + \frac{\operatorname{arccosh}(cx)c^8d^3x^2}{2} + \frac{3e \operatorname{arccosh}(cx)c^8d^2x^4}{4} + \frac{e^2 \operatorname{arccosh}(cx)c^8dx^6}{2} + \frac{e^3 \operatorname{arccosh}(cx)c^8x^8}{8}\right)}{8c^6e}$

input `int(x*(e*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output  $\frac{1}{8}ax^8(e^2x^2+d)^4/e + \frac{b}{c^2} \left( \frac{1}{8}c^2e^3 \operatorname{arccosh}(cx)x^8 + \frac{1}{2}c^2e^2 \operatorname{arccosh}(cx)x^6d + \frac{3}{4}c^2e \operatorname{arccosh}(cx)x^4d^2 + \frac{1}{2} \operatorname{arccosh}(cx)c^2x^2d^3 + \frac{1}{8}c^2 \operatorname{arccosh}(cx)d^4 \right) / e^6$   
 $- \frac{1}{3072} \frac{b}{c^6} \frac{e^3}{e} (cx-1)^{1/2} (cx+1)^{1/2} (384c^8d^4 \ln(cx+(c^2x^2-1)^{1/2}) + 768c^7d^3ex(c^2x^2-1)^{1/2} + 576c^7d^2e^2(c^2x^2-1)^{1/2}x^3 + 256c^7d^2e^3(c^2x^2-1)^{1/2}x^5 + 48e^4(c^2x^2-1)^{1/2}c^7x^7 + 768c^6d^3e \ln(cx+(c^2x^2-1)^{1/2}) + 864c^5d^2e^2x^3(c^2x^2-1)^{1/2} + 320(c^2x^2-1)^{1/2}c^5d^2e^3x^3 + 56e^4c^5x^5(c^2x^2-1)^{1/2} + 864c^4d^2e^2 \ln(cx+(c^2x^2-1)^{1/2}) + 480c^3d^2e^3x^3(c^2x^2-1)^{1/2} + 70e^4c^3x^3(c^2x^2-1)^{1/2} + 480c^2d^2e^3 \ln(cx+(c^2x^2-1)^{1/2}) + 105e^4c^2x^2(c^2x^2-1)^{1/2} + 105e^4 \ln(cx+(c^2x^2-1)^{1/2})) / (c^2x^2-1)^{1/2}$

### 3.482.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 286, normalized size of antiderivative = 0.80

$$\int x(d + ex^2)^3 (a + \operatorname{arccosh}(cx)) dx = \frac{384ac^8e^3x^8 + 1536ac^8de^2x^6 + 2304ac^8d^2ex^4 + 1536ac^8d^3x^2 + 3(128bc^8e^3x^8 + 512bc^8de^2x^6 + 768bc^8d^2)}{8e^6}$$

input `integrate(x*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fracas")`

```
output 1/3072*(384*a*c^8*e^3*x^8 + 1536*a*c^8*d*e^2*x^6 + 2304*a*c^8*d^2*e*x^4 +
1536*a*c^8*d^3*x^2 + 3*(128*b*c^8*e^3*x^8 + 512*b*c^8*d*e^2*x^6 + 768*b*c^
8*d^2*e*x^4 + 512*b*c^8*d^3*x^2 - 256*b*c^6*d^3 - 288*b*c^4*d^2*e - 160*b*
c^2*d*e^2 - 35*b*e^3)*log(c*x + sqrt(c^2*x^2 - 1)) - (48*b*c^7*e^3*x^7 + 8
*(32*b*c^7*d*e^2 + 7*b*c^5*e^3)*x^5 + 2*(288*b*c^7*d^2*e + 160*b*c^5*d*e^2
+ 35*b*c^3*e^3)*x^3 + 3*(256*b*c^7*d^3 + 288*b*c^5*d^2*e + 160*b*c^3*d*e^
2 + 35*b*c*e^3)*x)*sqrt(c^2*x^2 - 1)/c^8
```

### 3.482.6 Sympy [F]

$$\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (d + ex^2)^3 dx$$

```
input integrate(x*(e*x**2+d)**3*(a+b*acosh(c*x)),x)
```

```
output Integral(x*(a + b*acosh(c*x))*(d + e*x**2)**3, x)
```

### 3.482.7 Maxima [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 409, normalized size of antiderivative = 1.14

$$\begin{aligned} \int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx &= \frac{1}{8} ae^3 x^8 + \frac{1}{2} ade^2 x^6 + \frac{3}{4} ad^2 ex^4 + \frac{1}{2} ad^3 x^2 \\ &+ \frac{1}{4} \left( 2x^2 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x}{c^2} + \frac{\log(2c^2 x + 2\sqrt{c^2 x^2 - 1} c)}{c^3} \right) \right) bd^3 \\ &+ \frac{3}{32} \left( 8x^4 \operatorname{arcosh}(cx) - \left( \frac{2\sqrt{c^2 x^2 - 1} x^3}{c^2} + \frac{3\sqrt{c^2 x^2 - 1} x}{c^4} + \frac{3 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1} c)}{c^5} \right) c \right) bd^2 e \\ &+ \frac{1}{96} \left( 48x^6 \operatorname{arcosh}(cx) - \left( \frac{8\sqrt{c^2 x^2 - 1} x^5}{c^2} + \frac{10\sqrt{c^2 x^2 - 1} x^3}{c^4} + \frac{15\sqrt{c^2 x^2 - 1} x}{c^6} + \frac{15 \log(2c^2 x + 2\sqrt{c^2 x^2 - 1} c)}{c^7} \right) c \right) bde \\ &+ \frac{1}{3072} \left( 384x^8 \operatorname{arcosh}(cx) - \left( \frac{48\sqrt{c^2 x^2 - 1} x^7}{c^2} + \frac{56\sqrt{c^2 x^2 - 1} x^5}{c^4} + \frac{70\sqrt{c^2 x^2 - 1} x^3}{c^6} + \frac{105\sqrt{c^2 x^2 - 1} x}{c^8} \right) c \right) bde \end{aligned}$$

```
input integrate(x*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")
```

```
output 1/8*a*e^3*x^8 + 1/2*a*d*e^2*x^6 + 3/4*a*d^2*e*x^4 + 1/2*a*d^3*x^2 + 1/4*(2
*x^2*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x/c^2 + log(2*c^2*x + 2*sqrt(c^2*
x^2 - 1)*c)/c^3))*b*d^3 + 3/32*(8*x^4*arccosh(c*x) - (2*sqrt(c^2*x^2 - 1)*
x^3/c^2 + 3*sqrt(c^2*x^2 - 1)*x/c^4 + 3*log(2*c^2*x + 2*sqrt(c^2*x^2 - 1)*
c)/c^5)*c)*b*d^2*e + 1/96*(48*x^6*arccosh(c*x) - (8*sqrt(c^2*x^2 - 1)*x^5/
c^2 + 10*sqrt(c^2*x^2 - 1)*x^3/c^4 + 15*sqrt(c^2*x^2 - 1)*x/c^6 + 15*log(2
*c^2*x + 2*sqrt(c^2*x^2 - 1)*c)/c^7)*c)*b*d*e^2 + 1/3072*(384*x^8*arccosh(
c*x) - (48*sqrt(c^2*x^2 - 1)*x^7/c^2 + 56*sqrt(c^2*x^2 - 1)*x^5/c^4 + 70*s
qrt(c^2*x^2 - 1)*x^3/c^6 + 105*sqrt(c^2*x^2 - 1)*x/c^8 + 105*log(2*c^2*x +
2*sqrt(c^2*x^2 - 1)*c)/c^9)*c)*b*e^3
```

### 3.482.8 Giac [F(-2)]

Exception generated.

$$\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

```
input integrate(x*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")
```

```
output Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve
cteur & l) Error: Bad Argument Value
```

### 3.482.9 Mupad [F(-1)]

Timed out.

$$\int x(d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int x(a + b \operatorname{acosh}(cx)) (ex^2 + d)^3 dx$$

```
input int(x*(a + b*acosh(c*x))*(d + e*x^2)^3,x)
```

```
output int(x*(a + b*acosh(c*x))*(d + e*x^2)^3, x)
```

### 3.483 $\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.483.1 Optimal result . . . . .	3557
3.483.2 Mathematica [A] (verified) . . . . .	3558
3.483.3 Rubi [A] (verified) . . . . .	3558
3.483.4 Maple [A] (verified) . . . . .	3561
3.483.5 Fricas [A] (verification not implemented) . . . . .	3561
3.483.6 Sympy [F] . . . . .	3562
3.483.7 Maxima [A] (verification not implemented) . . . . .	3562
3.483.8 Giac [F(-2)] . . . . .	3563
3.483.9 Mupad [F(-1)] . . . . .	3563

#### 3.483.1 Optimal result

Integrand size = 18, antiderivative size = 287

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \frac{b(35c^6d^3 + 35c^4d^2e + 21c^2de^2 + 5e^3)(1 - c^2x^2)}{35c^7\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{be(35c^4d^2 + 42c^2de + 15e^2)(1 - c^2x^2)^2}{105c^7\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{3be^2(7c^2d + 5e)(1 - c^2x^2)^3}{175c^7\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{be^3(1 - c^2x^2)^4}{49c^7\sqrt{-1 + cx}\sqrt{1 + cx}} + d^3x(a + \operatorname{barccosh}(cx)) + d^2ex^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^3x^7(a + \operatorname{barccosh}(cx))$$

output

```
d^3*x*(a+b*arccosh(c*x))+d^2*e*x^3*(a+b*arccosh(c*x))+3/5*d*e^2*x^5*(a+b*arccosh(c*x))+1/7*e^3*x^7*(a+b*arccosh(c*x))+1/35*b*(35*c^6*d^3+35*c^4*d^2*e+21*c^2*d*e^2+5*e^3)*(-c^2*x^2+1)/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/105*b*e*(35*c^4*d^2+42*c^2*d*e+15*e^2)*(-c^2*x^2+1)^2/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)+3/175*b*e^2*(7*c^2*d+5*e)*(-c^2*x^2+1)^3/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/49*b*e^3*(-c^2*x^2+1)^4/c^7/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.483.2 Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.67

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = a \left( d^3x + d^2ex^3 + \frac{3}{5}de^2x^5 + \frac{e^3x^7}{7} \right) - \frac{b\sqrt{-1 + cx}\sqrt{1 + cx}(240e^3 + 24c^2e^2(49d + 5ex^2) + 2c^4e(1225d^2 + 294dex^2 + 45e^2x^4) + c^6(3675d^3 + 1225d^2ex^2 + 441de^2x^4 + 75e^3x^6))}{3675c^7} + \frac{1}{35}bx(35d^3 + 35d^2ex^2 + 21de^2x^4 + 5e^3x^6) \operatorname{arccosh}(cx)$$

input `Integrate[(d + e*x^2)^3*(a + b*ArcCosh[c*x]), x]`

```
output a*(d^3*x + d^2*e*x^3 + (3*d*e^2*x^5)/5 + (e^3*x^7)/7) - (b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(240*e^3 + 24*c^2*e^2*(49*d + 5*e*x^2) + 2*c^4*e*(1225*d^2 + 294*d*e*x^2 + 45*e^2*x^4) + c^6*(3675*d^3 + 1225*d^2*e*x^2 + 441*d*e^2*x^4 + 75*e^3*x^6)))/(3675*c^7) + (b*x*(35*d^3 + 35*d^2*e*x^2 + 21*d*e^2*x^4 + 5*e^3*x^6)*ArcCosh[c*x])/35
```

**3.483.3 Rubi [A] (verified)**Time = 0.72 (sec) , antiderivative size = 253, normalized size of antiderivative = 0.88, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6323, 27, 2113, 2331, 2389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

↓ 6323

$$-bc \int \frac{x(5e^3x^6 + 21de^2x^4 + 35d^2ex^2 + 35d^3)}{35\sqrt{cx-1}\sqrt{cx+1}} dx + d^3x(a + \operatorname{barccosh}(cx)) + d^2ex^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^3x^7(a + \operatorname{barccosh}(cx))$$

↓ 27

$$-\frac{1}{35}bc \int \frac{x(5e^3x^6 + 21de^2x^4 + 35d^2ex^2 + 35d^3)}{\sqrt{cx-1}\sqrt{cx+1}} dx + d^3x(a + \operatorname{barccosh}(cx)) + d^2ex^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^3x^7(a + \operatorname{barccosh}(cx))$$

---

3.483.  $\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
& \downarrow \text{2113} \\
& \frac{bc\sqrt{c^2x^2-1} \int \frac{x(5e^3x^6+21de^2x^4+35d^2ex^2+35d^3)}{\sqrt{c^2x^2-1}} dx}{35\sqrt{cx-1}\sqrt{cx+1}} + d^3x(a + \operatorname{barccosh}(cx)) + d^2ex^3(a + \\
& \operatorname{barccosh}(cx)) + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^3x^7(a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{2331} \\
& \frac{bc\sqrt{c^2x^2-1} \int \frac{5e^3x^6+21de^2x^4+35d^2ex^2+35d^3}{\sqrt{c^2x^2-1}} dx^2}{70\sqrt{cx-1}\sqrt{cx+1}} + d^3x(a + \operatorname{barccosh}(cx)) + d^2ex^3(a + \\
& \operatorname{barccosh}(cx)) + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^3x^7(a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{2389} \\
& \frac{bc\sqrt{c^2x^2-1} \int \left( \frac{5(c^2x^2-1)^{5/2}e^3}{c^6} + \frac{3(7dc^2+5e)(c^2x^2-1)^{3/2}e^2}{c^6} + \frac{(35d^2c^4+42dec^2+15e^2)\sqrt{c^2x^2-1}e}{c^6} + \frac{35d^3c^6+35d^2ec^4+21de^2c^2+5e^3}{c^6\sqrt{c^2x^2-1}} \right) dx}{70\sqrt{cx-1}\sqrt{cx+1}} \\
& d^3x(a + \operatorname{barccosh}(cx)) + d^2ex^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^3x^7(a + \operatorname{barccosh}(cx)) \\
& \downarrow \text{2009} \\
& d^3x(a + \operatorname{barccosh}(cx)) + d^2ex^3(a + \operatorname{barccosh}(cx)) + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx)) + \frac{1}{7}e^3x^7(a + \\
& \operatorname{barccosh}(cx)) - \\
& \frac{bc\sqrt{c^2x^2-1} \left( \frac{6e^2(c^2x^2-1)^{5/2}(7c^2d+5e)}{5c^8} + \frac{10e^3(c^2x^2-1)^{7/2}}{7c^8} + \frac{2e(c^2x^2-1)^{3/2}(35c^4d^2+42c^2de+15e^2)}{3c^8} + \frac{2\sqrt{c^2x^2-1}(35c^6d^3+35c^4d^2e+35c^2d^2e^2+5e^3)}{c^8\sqrt{c^2x^2-1}} \right)}{70\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(d + e*x^2)^3*(a + b*ArcCosh[c*x]), x]`

output `-1/70*(b*c*Sqrt[-1 + c^2*x^2]*((2*(35*c^6*d^3 + 35*c^4*d^2*e + 21*c^2*d*e^2 + 5*e^3)*Sqrt[-1 + c^2*x^2])/c^8 + (2*e*(35*c^4*d^2 + 42*c^2*d*e + 15*e^2)*(-1 + c^2*x^2)^(3/2))/(3*c^8) + (6*e^2*(7*c^2*d + 5*e)*(-1 + c^2*x^2)^(5/2))/(5*c^8) + (10*e^3*(-1 + c^2*x^2)^(7/2))/(7*c^8)))/(Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + d^3*x*(a + b*ArcCosh[c*x]) + d^2*e*x^3*(a + b*ArcCosh[c*x]) + (3*d*e^2*x^5*(a + b*ArcCosh[c*x]))/5 + (e^3*x^7*(a + b*ArcCosh[c*x]))/7`



## 3.483.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2113 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`
- rule 2331 `Int[(Pq_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`
- rule 2389 `Int[(Pq_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Int[ExpandIntegrand[Pq*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, n}, x] && PolyQ[Pq, x] && (IGtQ[p, 0] || EqQ[n, 1])`
- rule 6323 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d + e, 0] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])`

### 3.483.4 Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.74

method	result
parts	$a\left(\frac{1}{7}e^3x^7 + \frac{3}{5}de^2x^5 + d^2ex^3 + d^3x\right) + \frac{b\left(\frac{c \operatorname{arccosh}(cx)e^3x^7}{7} + \frac{3c \operatorname{arccosh}(cx)de^2x^5}{5} + c \operatorname{arccosh}(cx)d^2ex^3 + \operatorname{arccosh}(cx)d^3x\right)}{c^6}$
derivativedivides	$\frac{a\left(d^3c^7x+d^2c^7ex^3+\frac{3}{5}dc^7e^2x^5+\frac{1}{7}e^3c^7x^7\right)}{c^6} + \frac{b\left(\operatorname{arccosh}(cx)d^3c^7x+\operatorname{arccosh}(cx)d^2c^7ex^3+\frac{3 \operatorname{arccosh}(cx)dc^7e^2x^5}{5}+\frac{\operatorname{arccosh}(cx)e^3c^7x^7}{7}\right)}{c^6}$
default	$\frac{a\left(d^3c^7x+d^2c^7ex^3+\frac{3}{5}dc^7e^2x^5+\frac{1}{7}e^3c^7x^7\right)}{c^6} + \frac{b\left(\operatorname{arccosh}(cx)d^3c^7x+\operatorname{arccosh}(cx)d^2c^7ex^3+\frac{3 \operatorname{arccosh}(cx)dc^7e^2x^5}{5}+\frac{\operatorname{arccosh}(cx)e^3c^7x^7}{7}\right)}{c^6}$

input `int((e*x^2+d)^3*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/7*e^3*x^7+3/5*d*e^2*x^5+d^2*e*x^3+d^3*x)+b/c*(1/7*c*arccosh(c*x)*e^3*x^7+3/5*c*arccosh(c*x)*d*e^2*x^5+c*arccosh(c*x)*d^2*e*x^3+arccosh(c*x)*c*x*d^3-1/3675/c^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(75*c^6*e^3*x^6+441*c^6*d*e^2*x^4+1225*c^6*d^2*e*x^2+90*c^4*e^3*x^4+3675*c^6*d^3+588*c^4*d*e^2*x^2+2450*c^4*d^2*e+120*c^2*e^3*x^2+1176*c^2*d*e^2+240*e^3))`

### 3.483.5 Fracas [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 241, normalized size of antiderivative = 0.84

$$\int (d + ex^2)^3 (a + b \operatorname{arccosh}(cx)) dx$$

$$= \frac{525 ac^7 e^3 x^7 + 2205 ac^7 d e^2 x^5 + 3675 ac^7 d^2 e x^3 + 3675 ac^7 d^3 x + 105 (5 bc^7 e^3 x^7 + 21 bc^7 d e^2 x^5 + 35 bc^7 d^2 e x^3 + 35 bc^7 d^3 x) \log(cx + \sqrt{c^2 x^2 - 1}) - (75 b^2 c^6 e^3 x^6 + 3675 b^2 c^6 d^3 + 2450 b^2 c^4 d^2 e + 1176 b^2 c^2 d e^2 + 9 (49 b^2 c^6 d e^2 + 10 b^2 c^4 e^3) x^4 + 240 b^2 e^3 + (1225 b^2 c^6 d^2 e + 588 b^2 c^4 d e^2 + 120 b^2 c^2 e^3) x^2) \sqrt{c^2 x^2 - 1}}{c^7}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output `1/3675*(525*a*c^7*e^3*x^7 + 2205*a*c^7*d*e^2*x^5 + 3675*a*c^7*d^2*e*x^3 + 3675*a*c^7*d^3*x + 105*(5*b*c^7*e^3*x^7 + 21*b*c^7*d*e^2*x^5 + 35*b*c^7*d^2*e*x^3 + 35*b*c^7*d^3*x)*log(c*x + sqrt(c^2*x^2 - 1)) - (75*b*c^6*e^3*x^6 + 3675*b*c^6*d^3 + 2450*b*c^4*d^2*e + 1176*b*c^2*d*e^2 + 9*(49*b*c^6*d*e^2 + 10*b*c^4*e^3)*x^4 + 240*b*e^3 + (1225*b*c^6*d^2*e + 588*b*c^4*d*e^2 + 120*b*c^2*e^3)*x^2)*sqrt(c^2*x^2 - 1)/c^7`

**3.483.6 Sympy [F]**

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d + ex^2)^3 dx$$

input `integrate((e*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**3, x)`

**3.483.7 Maxima [A] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.00

$$\begin{aligned} \int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx &= \frac{1}{7} ae^3 x^7 + \frac{3}{5} ade^2 x^5 + ad^2 ex^3 \\ &+ \frac{1}{3} \left( 3x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bd^2 e \\ &+ \frac{1}{25} \left( 15x^5 \operatorname{arcosh}(cx) - \left( \frac{3\sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4\sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8\sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bde^2 \\ &+ \frac{1}{245} \left( 35x^7 \operatorname{arcosh}(cx) - \left( \frac{5\sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6\sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8\sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16\sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) be^3 \\ &+ ad^3 x + \frac{(cx \operatorname{arcosh}(cx) - \sqrt{c^2 x^2 - 1}) bd^3}{c} \end{aligned}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `1/7*a*e^3*x^7 + 3/5*a*d*e^2*x^5 + a*d^2*e*x^3 + 1/3*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*b*d^2*e + 1/25*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d*e^2 + 1/245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b*e^3 + a*d^3*x + (c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*b*d^3/c`

**3.483.8 Giac [F(-2)]**

Exception generated.

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

**3.483.9 Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (ex^2 + d)^3 dx$$

input `int((a + b*acosh(c*x))*(d + e*x^2)^3,x)`

output `int((a + b*acosh(c*x))*(d + e*x^2)^3, x)`

$$3.484 \quad \int \frac{(d+ex^2)^3 (a+b\operatorname{arccosh}(cx))}{x} dx$$

3.484.1 Optimal result . . . . .	3565
3.484.2 Mathematica [A] (warning: unable to verify) . . . . .	3566
3.484.3 Rubi [A] (verified) . . . . .	3567
3.484.4 Maple [A] (verified) . . . . .	3569
3.484.5 Fricas [F] . . . . .	3569
3.484.6 Sympy [F] . . . . .	3570
3.484.7 Maxima [F] . . . . .	3570
3.484.8 Giac [F] . . . . .	3570
3.484.9 Mupad [F(-1)] . . . . .	3571

**3.484.1 Optimal result**

Integrand size = 21, antiderivative size = 509

$$\begin{aligned}
\int \frac{(d+ex^2)^3 (a+\operatorname{barccosh}(cx))}{x} dx = & -\frac{3bd^2ex\sqrt{-1+cx}\sqrt{1+cx}}{4c} \\
& -\frac{9bde^2x\sqrt{-1+cx}\sqrt{1+cx}}{32c^3} \\
& -\frac{5be^3x\sqrt{-1+cx}\sqrt{1+cx}}{96c^5} \\
& -\frac{3bde^2x^3\sqrt{-1+cx}\sqrt{1+cx}}{16c} \\
& -\frac{5be^3x^3\sqrt{-1+cx}\sqrt{1+cx}}{144c^3} \\
& -\frac{be^3x^5\sqrt{-1+cx}\sqrt{1+cx}}{36c} - \frac{3bd^2e\operatorname{arccosh}(cx)}{4c^2} \\
& -\frac{9bde^2\operatorname{arccosh}(cx)}{32c^4} - \frac{5be^3\operatorname{arccosh}(cx)}{96c^6} \\
& + \frac{3}{2}d^2ex^2(a+\operatorname{barccosh}(cx)) \\
& + \frac{3}{4}de^2x^4(a+\operatorname{barccosh}(cx)) + \frac{1}{6}e^3x^6(a+\operatorname{barccosh}(cx)) \\
& - \frac{ibd^3\sqrt{1-c^2x^2}\arcsin(cx)^2}{2\sqrt{-1+cx}\sqrt{1+cx}} \\
& + \frac{bd^3\sqrt{1-c^2x^2}\arcsin(cx)\log(1-e^{2i\arcsin(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\
& + d^3(a+\operatorname{barccosh}(cx))\log(x) \\
& - \frac{bd^3\sqrt{1-c^2x^2}\arcsin(cx)\log(x)}{\sqrt{-1+cx}\sqrt{1+cx}} \\
& - \frac{ibd^3\sqrt{1-c^2x^2}\operatorname{PolyLog}(2, e^{2i\arcsin(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}}
\end{aligned}$$

output

$$\begin{aligned}
& -3/4*b*d^2*e*arccosh(c*x)/c^2-9/32*b*d*e^2*arccosh(c*x)/c^4-5/96*b*e^3*arc \\
& cosh(c*x)/c^6+3/2*d^2*e*x^2*(a+b*arccosh(c*x))+3/4*d*e^2*x^4*(a+b*arccosh( \\
& c*x))+1/6*e^3*x^6*(a+b*arccosh(c*x))+d^3*(a+b*arccosh(c*x))*ln(x)-3/4*b*d^ \\
& 2*e*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-9/32*b*d*e^2*x*(c*x-1)^(1/2)*(c*x+1)^( \\
& 1/2)/c^3-5/96*b*e^3*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5-3/16*b*d*e^2*x^3*(c* \\
& x-1)^(1/2)*(c*x+1)^(1/2)/c-5/144*b*e^3*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3 \\
& -1/36*b*e^3*x^5*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-1/2*I*b*d^3*arcsin(c*x)^2*(- \\
& c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*d^3*arcsin(c*x)*ln(1-(I*c*x \\
& +(-c^2*x^2+1)^(1/2))^2*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*d \\
& ^3*arcsin(c*x)*ln(x)*(-c^2*x^2+1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/2*I* \\
& b*d^3*polylog(2,(I*c*x+(-c^2*x^2+1)^(1/2))^2*(-c^2*x^2+1)^(1/2)/(c*x-1)^( \\
& 1/2)/(c*x+1)^(1/2)
\end{aligned}$$

### 3.484.2 Mathematica [A] (warning: unable to verify)

Time = 0.79 (sec) , antiderivative size = 344, normalized size of antiderivative = 0.68

$$\begin{aligned}
& \int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x} dx = \frac{3}{2}ad^2ex^2 + \frac{3}{4}ade^2x^4 + \frac{1}{6}ae^3x^6 \\
& - \frac{3bd^2e\left(cx\sqrt{-1+cx}\sqrt{1+cx} - 2c^2x^2\operatorname{arccosh}(cx) + 2\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right)}{4c^2} \\
& - \frac{3bde^2\left(cx\sqrt{\frac{-1+cx}{1+cx}}(3+3cx+2c^2x^2+2c^3x^3) - 8c^4x^4\operatorname{arccosh}(cx) + 6\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right)}{32c^4} \\
& - \frac{be^3\left(cx\sqrt{\frac{-1+cx}{1+cx}}(15+15cx+10c^2x^2+10c^3x^3+8c^4x^4+8c^5x^5) - 48c^6x^6\operatorname{arccosh}(cx) + 30\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right)}{288c^6} \\
& + ad^3\log(x) \\
& + \frac{1}{2}bd^3\left(\operatorname{arccosh}(cx)\left(\operatorname{arccosh}(cx) + 2\log\left(1+e^{-2\operatorname{arccosh}(cx)}\right)\right) - \operatorname{PolyLog}\left(2,-e^{-2\operatorname{arccosh}(cx)}\right)\right)
\end{aligned}$$

input `Integrate[((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x,x]`

output  $(3*a*d^2*e*x^2)/2 + (3*a*d*e^2*x^4)/4 + (a*e^3*x^6)/6 - (3*b*d^2*e*(c*x*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x] - 2*c^2*x^2*\text{ArcCosh}[c*x] + 2*\text{ArcTanh}[\text{Sqrt}[(-1 + c*x)/(1 + c*x)]])/(4*c^2) - (3*b*d*e^2*(c*x*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(3 + 3*c*x + 2*c^2*x^2 + 2*c^3*x^3) - 8*c^4*x^4*\text{ArcCosh}[c*x] + 6*\text{ArcTanh}[\text{Sqrt}[(-1 + c*x)/(1 + c*x)]])/(32*c^4) - (b*e^3*(c*x*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(15 + 15*c*x + 10*c^2*x^2 + 10*c^3*x^3 + 8*c^4*x^4 + 8*c^5*x^5) - 48*c^6*x^6*\text{ArcCosh}[c*x] + 30*\text{ArcTanh}[\text{Sqrt}[(-1 + c*x)/(1 + c*x)]])/(288*c^6) + a*d^3*\text{Log}[x] + (b*d^3*(\text{ArcCosh}[c*x]*(\text{ArcCosh}[c*x] + 2*\text{Log}[1 + E^(-2*\text{ArcCosh}[c*x])]) - \text{PolyLog}[2, -E^(-2*\text{ArcCosh}[c*x])])))/2$

### 3.484.3 Rubi [A] (verified)

Time = 1.33 (sec) , antiderivative size = 508, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^3 (a + \text{barccosh}(cx))}{x} dx$$

↓ 6373

$$-bc \int \frac{2e^3x^6 + 9de^2x^4 + 18d^2ex^2 + 12d^3 \log(x)}{12\sqrt{cx-1}\sqrt{cx+1}} dx + d^3 \log(x)(a + \text{barccosh}(cx)) + \frac{3}{2}d^2ex^2(a + \text{barccosh}(cx)) + \frac{3}{4}de^2x^4(a + \text{barccosh}(cx)) + \frac{1}{6}e^3x^6(a + \text{barccosh}(cx))$$

↓ 27

$$-\frac{1}{12}bc \int \frac{2e^3x^6 + 9de^2x^4 + 18d^2ex^2 + 12d^3 \log(x)}{\sqrt{cx-1}\sqrt{cx+1}} dx + d^3 \log(x)(a + \text{barccosh}(cx)) + \frac{3}{2}d^2ex^2(a + \text{barccosh}(cx)) + \frac{3}{4}de^2x^4(a + \text{barccosh}(cx)) + \frac{1}{6}e^3x^6(a + \text{barccosh}(cx))$$

↓ 7293

$$-\frac{1}{12}bc \int \left( \frac{2e^3x^6}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{9de^2x^4}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{18d^2ex^2}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{12d^3 \log(x)}{\sqrt{cx-1}\sqrt{cx+1}} \right) dx + d^3 \log(x)(a + \text{barccosh}(cx)) + \frac{3}{2}d^2ex^2(a + \text{barccosh}(cx)) + \frac{3}{4}de^2x^4(a + \text{barccosh}(cx)) + \frac{1}{6}e^3x^6(a + \text{barccosh}(cx))$$

↓ 2009

---

3.484.  $\int \frac{(d+ex^2)^3(a+\text{barccosh}(cx))}{x} dx$



$$d^3 \log(x)(a + \operatorname{barccosh}(cx)) + \frac{3}{2}d^2ex^2(a + \operatorname{barccosh}(cx)) + \frac{3}{4}de^2x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{6}e^3x^6(a + \operatorname{barccosh}(cx)) - \frac{1}{12}bc \left( \frac{5e^3 \operatorname{arccosh}(cx)}{8c^7} + \frac{27de^2 \operatorname{arccosh}(cx)}{8c^5} + \frac{9d^2e \operatorname{arccosh}(cx)}{c^3} + \frac{6id^3 \sqrt{1-c^2x^2} \operatorname{PolyLog}(2, e^{2i \operatorname{arcsin}(cx)})}{c\sqrt{cx-1}\sqrt{cx+1}} + \frac{6id^3}{c} \right)$$

input `Int[((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x,x]`

output `(3*d^2*e*x^2*(a + b*ArcCosh[c*x]))/2 + (3*d*e^2*x^4*(a + b*ArcCosh[c*x]))/4 + (e^3*x^6*(a + b*ArcCosh[c*x]))/6 + d^3*(a + b*ArcCosh[c*x])*Log[x] - (b*c*((9*d^2*e*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c^2 + (27*d*e^2*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(8*c^4) + (5*e^3*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(8*c^6) + (9*d*e^2*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(4*c^2) + (5*e^3*x^3*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(12*c^4) + (e^3*x^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(3*c^2) + (9*d^2*e*ArcCosh[c*x])/c^3 + (27*d*e^2*ArcCosh[c*x])/(8*c^5) + (5*e^3*ArcCosh[c*x])/(8*c^7) + ((6*I)*d^3*Sqrt[1 - c^2*x^2]*ArcSin[c*x]^2)/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (12*d^3*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[1 - E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (12*d^3*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[x])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((6*I)*d^3*Sqrt[1 - c^2*x^2]*PolyLog[2, E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/12`

### 3.484.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]`  
`]`

### 3.484.4 Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 348, normalized size of antiderivative = 0.68

method	result
parts	$a \left( \frac{e^3 x^6}{6} + \frac{3d e^2 x^4}{4} + \frac{3d^2 e x^2}{2} + d^3 \ln(x) \right) + \frac{b \operatorname{arccosh}(cx) e^3 x^6}{6} - \frac{9bd e^2 \operatorname{arccosh}(cx)}{32c^4} - \frac{3bd^2 e \operatorname{arccosh}(cx)}{4c^2}$
derivativedivides	$\frac{a \left( \frac{3c^6 d^2 e x^2}{2} + \frac{3c^6 d e^2 x^4}{4} + \frac{c^6 e^3 x^6}{6} + c^6 d^3 \ln(cx) \right)}{c^6} + \frac{b d^3 \operatorname{polylog} \left( 2, - (cx + \sqrt{cx-1} \sqrt{cx+1})^2 \right)}{2} + \frac{b \operatorname{arccosh}(cx) e^3 x^6}{6} +$
default	$\frac{a \left( \frac{3c^6 d^2 e x^2}{2} + \frac{3c^6 d e^2 x^4}{4} + \frac{c^6 e^3 x^6}{6} + c^6 d^3 \ln(cx) \right)}{c^6} + \frac{b d^3 \operatorname{polylog} \left( 2, - (cx + \sqrt{cx-1} \sqrt{cx+1})^2 \right)}{2} + \frac{b \operatorname{arccosh}(cx) e^3 x^6}{6} +$

input `int((e*x^2+d)^3*(a+b*arccosh(c*x))/x,x,method=_RETURNVERBOSE)`

output `a*(1/6*e^3*x^6+3/4*d*e^2*x^4+3/2*d^2*e*x^2+d^3*ln(x))+1/6*b*arccosh(c*x)*e^3*x^6-9/32*b*d*e^2*arccosh(c*x)/c^4-3/4*b*d^2*e*arccosh(c*x)/c^2-1/36*b*e^3*x^5*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-5/144*b*e^3*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-5/96*b*e^3*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5+3/4*b*arccosh(c*x)*d*e^2*x^4+3/2*b*arccosh(c*x)*d^2*e*x^2+b*d^3*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)-5/96*b*e^3*arccosh(c*x)/c^6-3/16*b*d*e^2*x^3*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-9/32*b*d*e^2*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-3/4*b*d^2*e*x*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-1/2*b*d^3*arccosh(c*x)^2+1/2*b*d^3*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)`

### 3.484.5 Fracas [F]

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)^3 (b \operatorname{arccosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x,x, algorithm="fracas")`

output `integral((a*e^3*x^6 + 3*a*d*e^2*x^4 + 3*a*d^2*e*x^2 + a*d^3 + (b*e^3*x^6 + 3*b*d*e^2*x^4 + 3*b*d^2*e*x^2 + b*d^3)*arccosh(c*x))/x, x)`

3.484.  $\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x} dx$

**3.484.6 Sympy [F]**

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^3}{x} dx$$

input `integrate((e*x**2+d)**3*(a+b*acosh(c*x))/x,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**3/x, x)`

**3.484.7 Maxima [F]**

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)^3 (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x,x, algorithm="maxima")`

output `1/6*a*e^3*x^6 + 3/4*a*d*e^2*x^4 + 3/2*a*d^2*e*x^2 + a*d^3*log(x) + integrate(b*e^3*x^5*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 3*b*d*e^2*x^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 3*b*d^2*e*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + b*d^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/x, x)`

**3.484.8 Giac [F]**

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x} dx = \int \frac{(ex^2 + d)^3 (b \operatorname{arcosh}(cx) + a)}{x} dx$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x,x, algorithm="giac")`

output `integrate((e*x^2 + d)^3*(b*arccosh(c*x) + a)/x, x)`

**3.484.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^3 (a + \operatorname{arccosh}(cx))}{x} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^3}{x} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x,x)`output `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x, x)`

**3.485**  $\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^2} dx$

3.485.1 Optimal result . . . . .	3572
3.485.2 Mathematica [A] (verified) . . . . .	3573
3.485.3 Rubi [A] (verified) . . . . .	3573
3.485.4 Maple [A] (verified) . . . . .	3576
3.485.5 Fricas [A] (verification not implemented) . . . . .	3576
3.485.6 Sympy [F] . . . . .	3577
3.485.7 Maxima [A] (verification not implemented) . . . . .	3577
3.485.8 Giac [F] . . . . .	3578
3.485.9 Mupad [F(-1)] . . . . .	3578

**3.485.1 Optimal result**

Integrand size = 21, antiderivative size = 265

$$\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^2} dx = \frac{be(15c^4d^2+5c^2de+e^2)(1-c^2x^2)}{5c^5\sqrt{-1+cx}\sqrt{1+cx}} - \frac{be^2(5c^2d+2e)(1-c^2x^2)^2}{15c^5\sqrt{-1+cx}\sqrt{1+cx}} + \frac{be^3(1-c^2x^2)^3}{25c^5\sqrt{-1+cx}\sqrt{1+cx}} - \frac{d^3(a+b\operatorname{arccosh}(cx))}{x} + 3d^2ex(a+b\operatorname{arccosh}(cx)) + de^2x^3(a+b\operatorname{arccosh}(cx)) + \frac{1}{5}e^3x^5(a+b\operatorname{arccosh}(cx)) + \frac{bcd^3\sqrt{-1+c^2x^2}\arctan(\sqrt{-1+c^2x^2})}{\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-d^3*(a+b*arccosh(c*x))/x+3*d^2*e*x*(a+b*arccosh(c*x))+d*e^2*x^3*(a+b*arccosh(c*x))+1/5*e^3*x^5*(a+b*arccosh(c*x))+1/5*b*e*(15*c^4*d^2+5*c^2*d*e+e^2)*(-c^2*x^2+1)/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/15*b*e^2*(5*c^2*d+2*e)*(-c^2*x^2+1)^2/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/25*b*e^3*(-c^2*x^2+1)^3/c^5/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*c*d^3*arctan((c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.485.2 Mathematica [A] (verified)**

Time = 0.22 (sec) , antiderivative size = 182, normalized size of antiderivative = 0.69

$$\int \frac{(d + ex^2)^3 (a + \operatorname{arccosh}(cx))}{x^2} dx$$

$$= -\frac{ad^3}{x} + 3ad^2ex + ade^2x^3 + \frac{1}{5}ae^3x^5$$

$$- \frac{be\sqrt{-1+cx}\sqrt{1+cx}(8e^2 + 2c^2e(25d + 2ex^2) + c^4(225d^2 + 25dex^2 + 3e^2x^4))}{75c^5}$$

$$+ \frac{b(-5d^3 + 15d^2ex^2 + 5de^2x^4 + e^3x^6) \operatorname{arccosh}(cx)}{5x} - bcd^3 \arctan\left(\frac{1}{\sqrt{-1+cx}\sqrt{1+cx}}\right)$$

input `Integrate[((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x^2,x]`output `-((a*d^3)/x) + 3*a*d^2*e*x + a*d*e^2*x^3 + (a*e^3*x^5)/5 - (b*e*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(8*e^2 + 2*c^2*e*(25*d + 2*e*x^2) + c^4*(225*d^2 + 25*d*e*x^2 + 3*e^2*x^4)))/(75*c^5) + (b*(-5*d^3 + 15*d^2*e*x^2 + 5*d*e^2*x^4 + e^3*x^6)*ArcCosh[c*x])/(5*x) - b*c*d^3*ArcTan[1/(Sqrt[-1 + c*x]*Sqrt[1 + c*x])]`**3.485.3 Rubi [A] (verified)**Time = 0.79 (sec) , antiderivative size = 217, normalized size of antiderivative = 0.82, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$ , Rules used = {6373, 27, 2113, 2331, 2123, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^3 (a + \operatorname{arccosh}(cx))}{x^2} dx$$

$$\downarrow \text{6373}$$

$$-bc \int -\frac{-e^3x^6 - 5de^2x^4 - 15d^2ex^2 + 5d^3}{5x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^3(a + \operatorname{arccosh}(cx))}{x} + 3d^2ex(a + \operatorname{arccosh}(cx)) +$$

$$de^2x^3(a + \operatorname{arccosh}(cx)) + \frac{1}{5}e^3x^5(a + \operatorname{arccosh}(cx))$$

$$\downarrow \text{27}$$

---

3.485.  $\int \frac{(d+ex^2)^3(a+\operatorname{arccosh}(cx))}{x^2} dx$

$$\frac{1}{5}bc \int \frac{-e^3x^6 - 5de^2x^4 - 15d^2ex^2 + 5d^3}{x\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^3(a + \operatorname{barccosh}(cx))}{x} + 3d^2ex(a + \operatorname{barccosh}(cx)) + de^2x^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^3x^5(a + \operatorname{barccosh}(cx))$$

↓ 2113

$$\frac{bc\sqrt{c^2x^2-1} \int \frac{-e^3x^6 - 5de^2x^4 - 15d^2ex^2 + 5d^3}{x\sqrt{c^2x^2-1}} dx}{5\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^3(a + \operatorname{barccosh}(cx))}{x} + 3d^2ex(a + \operatorname{barccosh}(cx)) + de^2x^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^3x^5(a + \operatorname{barccosh}(cx))$$

↓ 2331

$$\frac{bc\sqrt{c^2x^2-1} \int \frac{-e^3x^6 - 5de^2x^4 - 15d^2ex^2 + 5d^3}{x^2\sqrt{c^2x^2-1}} dx^2}{10\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^3(a + \operatorname{barccosh}(cx))}{x} + 3d^2ex(a + \operatorname{barccosh}(cx)) + de^2x^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^3x^5(a + \operatorname{barccosh}(cx))$$

↓ 2123

$$\frac{bc\sqrt{c^2x^2-1} \int \left( \frac{5d^3}{x^2\sqrt{c^2x^2-1}} - \frac{e^3(c^2x^2-1)^{3/2}}{c^4} - \frac{e^2(5dc^2+2e)\sqrt{c^2x^2-1}}{c^4} - \frac{e(15d^2c^4+5dec^2+e^2)}{c^4\sqrt{c^2x^2-1}} \right) dx^2}{10\sqrt{cx-1}\sqrt{cx+1}} - \frac{d^3(a + \operatorname{barccosh}(cx))}{x} + 3d^2ex(a + \operatorname{barccosh}(cx)) + de^2x^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^3x^5(a + \operatorname{barccosh}(cx))$$

↓ 2009

$$-\frac{d^3(a + \operatorname{barccosh}(cx))}{x} + 3d^2ex(a + \operatorname{barccosh}(cx)) + de^2x^3(a + \operatorname{barccosh}(cx)) + \frac{1}{5}e^3x^5(a + \operatorname{barccosh}(cx)) + \frac{bc\sqrt{c^2x^2-1} \left( 10d^3 \arctan(\sqrt{c^2x^2-1}) - \frac{2e^2(c^2x^2-1)^{3/2}(5c^2d+2e)}{3c^6} - \frac{2e^3(c^2x^2-1)^{5/2}}{5c^6} - \frac{2e\sqrt{c^2x^2-1}(15c^4d^2+5c^2de+e^2)}{c^6} \right)}{10\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int(((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x^2,x)`

output `-((d^3*(a + b*ArcCosh[c*x]))/x) + 3*d^2*e*x*(a + b*ArcCosh[c*x]) + d*e^2*x^3*(a + b*ArcCosh[c*x]) + (e^3*x^5*(a + b*ArcCosh[c*x]))/5 + (b*c*Sqrt[-1 + c^2*x^2])*((-2*e*(15*c^4*d^2 + 5*c^2*d*e + e^2)*Sqrt[-1 + c^2*x^2])/c^6 - (2*e^2*(5*c^2*d + 2*e)*(-1 + c^2*x^2)^(3/2))/(3*c^6) - (2*e^3*(-1 + c^2*x^2)^(5/2))/(5*c^6) + 10*d^3*ArcTan[Sqrt[-1 + c^2*x^2]])/(10*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

---

3.485.  $\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x^2} dx$

## 3.485.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`
- rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`
- rule 2113 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[P_x*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[P_x, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`
- rule 2123 `Int[(P_x)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[P_x*(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && PolyQ[P_x, x] && (IntegersQ[m, n] || IGtQ[m, -2])`
- rule 2331 `Int[(P_q)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, P_q, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[P_q, x^2] && IntegerQ[(m - 1)/2]`
- rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`



### 3.485.4 Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.03

method	result
parts	$a \left( \frac{e^3 x^5}{5} + d e^2 x^3 + 3 d^2 e x - \frac{d^3}{x} \right) + b c \left( \frac{\operatorname{arccosh}(c x) e^3 x^5}{5 c} + \frac{\operatorname{arccosh}(c x) d e^2 x^3}{c} + \frac{3 \operatorname{arccosh}(c x) d^2 e x}{c} - \frac{d^3}{x} \right)$
derivativedivides	$c \left( \frac{a \left( 3 c^5 x d^2 e + c^5 x^3 d e^2 + \frac{c^5 x^5 e^3}{5} - \frac{c^5 d^3}{x} \right)}{c^6} + \frac{b \left( 3 \operatorname{arccosh}(c x) c^5 d^2 e x + \operatorname{arccosh}(c x) c^5 d e^2 x^3 + \frac{\operatorname{arccosh}(c x) e^3 c^5 x^5}{5} - \operatorname{arccosh}(c x) d^3}{x} \right)}{c^6} \right)$
default	$c \left( \frac{a \left( 3 c^5 x d^2 e + c^5 x^3 d e^2 + \frac{c^5 x^5 e^3}{5} - \frac{c^5 d^3}{x} \right)}{c^6} + \frac{b \left( 3 \operatorname{arccosh}(c x) c^5 d^2 e x + \operatorname{arccosh}(c x) c^5 d e^2 x^3 + \frac{\operatorname{arccosh}(c x) e^3 c^5 x^5}{5} - \operatorname{arccosh}(c x) d^3}{x} \right)}{c^6} \right)$

input `int((e*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x,method=_RETURNVERBOSE)`

output `a*(1/5*e^3*x^5+d*e^2*x^3+3*d^2*e*x-d^3/x)+b*c*(1/5/c*arccosh(c*x)*e^3*x^5+1/c*arccosh(c*x)*d*e^2*x^3+3/c*arccosh(c*x)*d^2*e*x-arccosh(c*x)*d^3/c/x-1/75/c^6*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(75*c^6*d^3*arctan(1/(c^2*x^2-1)^(1/2))+225*c^4*d^2*e*(c^2*x^2-1)^(1/2)+25*c^4*d*e^2*(c^2*x^2-1)^(1/2)*x^2+3*e^3*(c^2*x^2-1)^(1/2)*c^4*x^4+50*c^2*d*e^2*(c^2*x^2-1)^(1/2)+4*e^3*c^2*x^2*(c^2*x^2-1)^(1/2)+8*e^3*(c^2*x^2-1)^(1/2))/(c^2*x^2-1)^(1/2))`

### 3.485.5 Fracas [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 327, normalized size of antiderivative = 1.23

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^2} dx$$

$$= \frac{15 ac^5 e^3 x^6 + 75 ac^5 d e^2 x^4 + 150 bc^6 d^3 x \arctan(-cx + \sqrt{c^2 x^2 - 1}) + 225 ac^5 d^2 e x^2 - 75 ac^5 d^3 + 15 (5 bc^5 d^3 \operatorname{arccosh}(cx) - 3 bc^5 d^3)}{c^6}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x, algorithm="fricas")`

output  $1/75*(15*a*c^5*e^3*x^6 + 75*a*c^5*d*e^2*x^4 + 150*b*c^6*d^3*x*\arctan(-c*x + \sqrt{c^2*x^2 - 1})) + 225*a*c^5*d^2*e*x^2 - 75*a*c^5*d^3 + 15*(5*b*c^5*d^3 - 15*b*c^5*d^2*e - 5*b*c^5*d*e^2 - b*c^5*e^3)*x*\log(-c*x + \sqrt{c^2*x^2 - 1}) + 15*(b*c^5*e^3*x^6 + 5*b*c^5*d*e^2*x^4 + 15*b*c^5*d^2*e*x^2 - 5*b*c^5*d^3 + (5*b*c^5*d^3 - 15*b*c^5*d^2*e - 5*b*c^5*d*e^2 - b*c^5*e^3)*x)*\log(c*x + \sqrt{c^2*x^2 - 1}) - (3*b*c^4*e^3*x^5 + (25*b*c^4*d*e^2 + 4*b*c^2*e^3)*x^3 + (225*b*c^4*d^2*e + 50*b*c^2*d*e^2 + 8*b*e^3)*x)*\sqrt{c^2*x^2 - 1})/(c^5*x)$

### 3.485.6 Sympy [F]

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^3}{x^2} dx$$

input `integrate((e*x**2+d)**3*(a+b*acosh(c*x))/x**2,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**3/x**2, x)`

### 3.485.7 Maxima [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 222, normalized size of antiderivative = 0.84

$$\begin{aligned} & \int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^2} dx \\ &= \frac{1}{5} a e^3 x^5 + a d e^2 x^3 - \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) b d^3 \\ &+ \frac{1}{3} \left( 3 x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2 \sqrt{c^2 x^2 - 1}}{c^4} \right) \right) b d e^2 \\ &+ \frac{1}{75} \left( 15 x^5 \operatorname{arccosh}(cx) - \left( \frac{3 \sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4 \sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8 \sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) b e^3 \\ &+ 3 a d^2 e x + \frac{3 (c x \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) b d^2 e}{c} - \frac{a d^3}{x} \end{aligned}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x, algorithm="maxima")`

output  $1/5*a*e^3*x^5 + a*d*e^2*x^3 - (c*\arcsin(1/(c*\text{abs}(x))) + \text{arccosh}(c*x)/x)*b*d^3 + 1/3*(3*x^3*\text{arccosh}(c*x) - c*(\text{sqrt}(c^2*x^2 - 1)*x^2/c^2 + 2*\text{sqrt}(c^2*x^2 - 1)/c^4))*b*d*e^2 + 1/75*(15*x^5*\text{arccosh}(c*x) - (3*\text{sqrt}(c^2*x^2 - 1)*x^4/c^2 + 4*\text{sqrt}(c^2*x^2 - 1)*x^2/c^4 + 8*\text{sqrt}(c^2*x^2 - 1)/c^6)*c)*b*e^3 + 3*a*d^2*e*x + 3*(c*x*\text{arccosh}(c*x) - \text{sqrt}(c^2*x^2 - 1))*b*d^2*e/c - a*d^3/x$

### 3.485.8 Giac [F]

$$\int \frac{(d + ex^2)^3 (a + b \text{arccosh}(cx))}{x^2} dx = \int \frac{(ex^2 + d)^3 (b \text{arcosh}(cx) + a)}{x^2} dx$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^2,x, algorithm="giac")`

output `integrate((e*x^2 + d)^3*(b*arccosh(c*x) + a)/x^2, x)`

### 3.485.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^2)^3 (a + b \text{arccosh}(cx))}{x^2} dx = \int \frac{(a + b \text{acosh}(cx)) (ex^2 + d)^3}{x^2} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x^2,x)`

output `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x^2, x)`

**3.486** 
$$\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^3} dx$$

3.486.1 Optimal result . . . . .	3579
3.486.2 Mathematica [A] (warning: unable to verify) . . . . .	3580
3.486.3 Rubi [A] (verified) . . . . .	3581
3.486.4 Maple [A] (verified) . . . . .	3583
3.486.5 Fricas [F] . . . . .	3583
3.486.6 Sympy [F] . . . . .	3584
3.486.7 Maxima [F] . . . . .	3584
3.486.8 Giac [F] . . . . .	3584
3.486.9 Mupad [F(-1)] . . . . .	3585

**3.486.1 Optimal result**

Integrand size = 21, antiderivative size = 476

$$\begin{aligned} \int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^3} dx = & -\frac{bcd^3(1-c^2x^2)}{2x\sqrt{-1+cx}\sqrt{1+cx}} + \frac{3be^2(8c^2d+e)x(1-c^2x^2)}{32c^3\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{be^3x^3(1-c^2x^2)}{16c\sqrt{-1+cx}\sqrt{1+cx}} - \frac{d^3(a+b\operatorname{arccosh}(cx))}{2x^2} \\ & + \frac{3}{2}de^2x^2(a+b\operatorname{arccosh}(cx)) + \frac{1}{4}e^3x^4(a+b\operatorname{arccosh}(cx)) \\ & - \frac{3ibd^2e\sqrt{1-c^2x^2}\arcsin(cx)^2}{2\sqrt{-1+cx}\sqrt{1+cx}} \\ & - \frac{3be^2(8c^2d+e)\sqrt{-1+c^2x^2}\operatorname{arctanh}\left(\frac{cx}{\sqrt{-1+c^2x^2}}\right)}{32c^4\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{3bd^2e\sqrt{1-c^2x^2}\arcsin(cx)\log(1-e^{2i\arcsin(cx)})}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & + 3d^2e(a+b\operatorname{arccosh}(cx))\log(x) \\ & - \frac{3bd^2e\sqrt{1-c^2x^2}\arcsin(cx)\log(x)}{\sqrt{-1+cx}\sqrt{1+cx}} \\ & - \frac{3ibd^2e\sqrt{1-c^2x^2}\operatorname{PolyLog}(2,e^{2i\arcsin(cx)})}{2\sqrt{-1+cx}\sqrt{1+cx}} \end{aligned}$$

output 
$$-1/2*d^3*(a+b*\operatorname{arccosh}(c*x))/x^2+3/2*d*e^2*x^2*(a+b*\operatorname{arccosh}(c*x))+1/4*e^3*x^4*(a+b*\operatorname{arccosh}(c*x))+3*d^2*e*(a+b*\operatorname{arccosh}(c*x))*\ln(x)-1/2*b*c*d^3*(-c^2*x^2+1)/x/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3/32*b*e^2*(8*c^2*d+e)*x*(-c^2*x^2+1)/c^3/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+1/16*b*e^3*x^3*(-c^2*x^2+1)/c/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3/2*I*b*d^2*e*\operatorname{arcsin}(c*x)^2*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}+3*b*d^2*e*\operatorname{arcsin}(c*x)*\ln(1-(I*c*x+(-c^2*x^2+1)^{(1/2}))^2)*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3*b*d^2*e*\operatorname{arcsin}(c*x)*\ln(x)*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3/2*I*b*d^2*e*\operatorname{polylog}(2,(I*c*x+(-c^2*x^2+1)^{(1/2}))^2)*(-c^2*x^2+1)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}-3/32*b*e^2*(8*c^2*d+e)*\operatorname{arctanh}(c*x/(c^2*x^2-1)^{(1/2}))*(-c^2*x^2-1)^{(1/2)}/c^4/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$$

### 3.486.2 Mathematica [A] (warning: unable to verify)

Time = 0.56 (sec) , antiderivative size = 278, normalized size of antiderivative = 0.58

$$\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^3} dx$$

$$= \frac{1}{4} \left( -\frac{2ad^3}{x^2} + 6ade^2x^2 + ae^3x^4 + \frac{2bd^3(cx\sqrt{-1+cx}\sqrt{1+cx} - \operatorname{arccosh}(cx))}{x^2} \right.$$

$$+ \frac{6bde^2x^2\operatorname{arccosh}(cx) + be^3x^4\operatorname{arccosh}(cx)}{c^2}$$

$$- \frac{3bde^2\left(cx\sqrt{-1+cx}\sqrt{1+cx} + 2\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right)}{c^2}$$

$$- \frac{be^3\left(cx\sqrt{\frac{-1+cx}{1+cx}}(3+3cx+2c^2x^2+2c^3x^3) + 6\operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)\right)}{8c^4}$$

$$+ 6bd^2e\operatorname{arccosh}(cx) \left( \operatorname{arccosh}(cx) + 2\log(1+e^{-2\operatorname{arccosh}(cx)}) \right) + 12ad^2e\log(x)$$

$$\left. - 6bd^2e\operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)}) \right)$$

input `Integrate[((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x^3,x]`

output  $((-2*a*d^3)/x^2 + 6*a*d*e^2*x^2 + a*e^3*x^4 + (2*b*d^3*(c*x*\sqrt{-1 + c*x})*\sqrt{1 + c*x} - \text{ArcCosh}[c*x]))/x^2 + 6*b*d*e^2*x^2*\text{ArcCosh}[c*x] + b*e^3*x^4*\text{ArcCosh}[c*x] - (3*b*d*e^2*(c*x*\sqrt{-1 + c*x})*\sqrt{1 + c*x} + 2*\text{ArcTanh}[\sqrt{(-1 + c*x)/(1 + c*x)}]))/c^2 - (b*e^3*(c*x*\sqrt{(-1 + c*x)/(1 + c*x)})*(3 + 3*c*x + 2*c^2*x^2 + 2*c^3*x^3) + 6*\text{ArcTanh}[\sqrt{(-1 + c*x)/(1 + c*x)}]))/(8*c^4) + 6*b*d^2*e*\text{ArcCosh}[c*x]*(\text{ArcCosh}[c*x] + 2*\text{Log}[1 + E^{(-2*\text{ArcCosh}[c*x])}]) + 12*a*d^2*e*\text{Log}[x] - 6*b*d^2*e*\text{PolyLog}[2, -E^{(-2*\text{ArcCosh}[c*x])}]))/4$

### 3.486.3 Rubi [A] (verified)

Time = 2.00 (sec) , antiderivative size = 480, normalized size of antiderivative = 1.01, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$ , Rules used = {6373, 27, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^3 (a + \text{barccosh}(cx))}{x^3} dx$$

↓ 6373

$$-bc \int -\frac{-e^3 x^6 - 6de^2 x^4 - 12d^2 e \log(x) x^2 + 2d^3}{4x^2 \sqrt{cx - 1} \sqrt{cx + 1}} dx - \frac{d^3 (a + \text{barccosh}(cx))}{2x^2} + 3d^2 e \log(x) (a + \text{barccosh}(cx)) + \frac{3}{2} de^2 x^2 (a + \text{barccosh}(cx)) + \frac{1}{4} e^3 x^4 (a + \text{barccosh}(cx))$$

↓ 27

$$\frac{1}{4} bc \int \frac{-e^3 x^6 - 6de^2 x^4 - 12d^2 e \log(x) x^2 + 2d^3}{x^2 \sqrt{cx - 1} \sqrt{cx + 1}} dx - \frac{d^3 (a + \text{barccosh}(cx))}{2x^2} + 3d^2 e \log(x) (a + \text{barccosh}(cx)) + \frac{3}{2} de^2 x^2 (a + \text{barccosh}(cx)) + \frac{1}{4} e^3 x^4 (a + \text{barccosh}(cx))$$

↓ 7293

$$\frac{1}{4} bc \int \left( \frac{-e^3 x^6 - 6de^2 x^4 + 2d^3}{x^2 \sqrt{cx - 1} \sqrt{cx + 1}} - \frac{12d^2 e \log(x)}{\sqrt{cx - 1} \sqrt{cx + 1}} \right) dx - \frac{d^3 (a + \text{barccosh}(cx))}{2x^2} + 3d^2 e \log(x) (a + \text{barccosh}(cx)) + \frac{3}{2} de^2 x^2 (a + \text{barccosh}(cx)) + \frac{1}{4} e^3 x^4 (a + \text{barccosh}(cx))$$

↓ 2009

$$-\frac{d^3(a + \operatorname{barccosh}(cx))}{2x^2} + 3d^2e \log(x)(a + \operatorname{barccosh}(cx)) + \frac{3}{2}de^2x^2(a + \operatorname{barccosh}(cx)) + \frac{1}{4}e^3x^4(a + \operatorname{barccosh}(cx)) + \frac{1}{4}bc \left( -\frac{6id^2e\sqrt{1-c^2x^2} \operatorname{PolyLog}(2, e^{2i \arcsin(cx)})}{c\sqrt{cx-1}\sqrt{cx+1}} - \frac{6id^2e\sqrt{1-c^2x^2} \arcsin(cx)^2}{c\sqrt{cx-1}\sqrt{cx+1}} + \frac{12d^2e\sqrt{1-c^2x^2} \arcsin(cx) \log(x)}{c\sqrt{cx-1}\sqrt{cx+1}} \right)$$

input `Int[((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x^3,x]`

output `-1/2*(d^3*(a + b*ArcCosh[c*x]))/x^2 + (3*d*e^2*x^2*(a + b*ArcCosh[c*x]))/2 + (e^3*x^4*(a + b*ArcCosh[c*x]))/4 + 3*d^2*e*(a + b*ArcCosh[c*x])*Log[x] + (b*c*((-2*d^3*(1 - c^2*x^2))/(x*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (3*e^2*(8*c^2*d + e)*x*(1 - c^2*x^2))/(8*c^4*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (e^3*x^3*(1 - c^2*x^2))/(4*c^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((6*I)*d^2*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]^2)/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (3*e^2*(8*c^2*d + e)*Sqrt[-1 + c^2*x^2]*ArcTanh[(c*x)/Sqrt[-1 + c^2*x^2]])/(8*c^5*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (12*d^2*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[1 - E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (12*d^2*e*Sqrt[1 - c^2*x^2]*ArcSin[c*x]*Log[x])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - ((6*I)*d^2*e*Sqrt[1 - c^2*x^2]*PolyLog[2, E^((2*I)*ArcSin[c*x])])/(c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/4`

### 3.486.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6373 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]`

---

3.486.  $\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x^3} dx$

## 3.486.4 Maple [A] (verified)

Time = 1.52 (sec) , antiderivative size = 293, normalized size of antiderivative = 0.62

method	result
parts	$a \left( \frac{e^3 x^4}{4} + \frac{3d e^2 x^2}{2} - \frac{d^3}{2x^2} + 3d^2 e \ln(x) \right) - \frac{bc^2 d^3}{2} - \frac{3be^2 \sqrt{cx-1} \sqrt{cx+1} x d}{4c} + \frac{be^3 \operatorname{arccosh}(cx) x^4}{4} - \frac{3bd}{4c^2}$
derivativeldivides	$c^2 \left( \frac{a \left( \frac{3c^4 d e^2 x^2}{2} + \frac{c^4 x^4 e^3}{4} + 3c^4 d^2 e \ln(cx) - \frac{c^4 d^3}{2x^2} \right)}{c^6} - \frac{3be^3 x \sqrt{cx-1} \sqrt{cx+1}}{32c^5} - \frac{3bd e^2 x \sqrt{cx-1} \sqrt{cx+1}}{4c^3} + \frac{b \operatorname{arccosh}(cx) x^4}{4c^2} \right)$
default	$c^2 \left( \frac{a \left( \frac{3c^4 d e^2 x^2}{2} + \frac{c^4 x^4 e^3}{4} + 3c^4 d^2 e \ln(cx) - \frac{c^4 d^3}{2x^2} \right)}{c^6} - \frac{3be^3 x \sqrt{cx-1} \sqrt{cx+1}}{32c^5} - \frac{3bd e^2 x \sqrt{cx-1} \sqrt{cx+1}}{4c^3} + \frac{b \operatorname{arccosh}(cx) x^4}{4c^2} \right)$

```
input int((e*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x,method=_RETURNVERBOSE)
```

```
output a*(1/4*e^3*x^4+3/2*d*e^2*x^2-1/2*d^3/x^2+3*d^2*e*ln(x))-1/2*b*c^2*d^3-3/4*b/c*e^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*x*d+1/4*b*e^3*arccosh(c*x)*x^4-3/2*b*d^2*e*arccosh(c*x)^2+3/2*b*e*d^2*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)-3/4*b/c^2*e^2*arccosh(c*x)*d+3*b*e*d^2*arccosh(c*x)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2)-1/16*b/c*e^3*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x^3-3/32*b/c^3*e^3*(c*x+1)^(1/2)*(c*x-1)^(1/2)*x+3/2*b*e^2*arccosh(c*x)*x^2*d+1/2*b*c*d^3/x*(c*x+1)^(1/2)*(c*x-1)^(1/2)-3/32*b/c^4*e^3*arccosh(c*x)-1/2*b*d^3/x^2*arccosh(c*x)
```

## 3.486.5 Fracas [F]

$$\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(ex^2+d)^3(b\operatorname{arccosh}(cx)+a)}{x^3} dx$$

```
input integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x, algorithm="fricas")
```

```
output integral((a*e^3*x^6 + 3*a*d*e^2*x^4 + 3*a*d^2*e*x^2 + a*d^3 + (b*e^3*x^6 + 3*b*d*e^2*x^4 + 3*b*d^2*e*x^2 + b*d^3)*arccosh(c*x))/x^3, x)
```



**3.486.6 Sympy [F]**

$$\int \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^3}{x^3} dx$$

input `integrate((e*x**2+d)**3*(a+b*acosh(c*x))/x**3,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**3/x**3, x)`

**3.486.7 Maxima [F]**

$$\int \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^3 (b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x, algorithm="maxima")`

output `1/4*a*e^3*x^4 + 3/2*a*d*e^2*x^2 + 1/2*b*d^3*(sqrt(c^2*x^2 - 1)*c/x - arccosh(c*x)/x^2) + 3*a*d^2*e*log(x) - 1/2*a*d^3/x^2 + integrate(b*e^3*x^3*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) + 3*b*d*e^2*x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1) + 3*b*d^2*e*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/x, x)`

**3.486.8 Giac [F]**

$$\int \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{x^3} dx = \int \frac{(ex^2 + d)^3 (b \operatorname{arcosh}(cx) + a)}{x^3} dx$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^3,x, algorithm="giac")`

output `integrate((e*x^2 + d)^3*(b*arccosh(c*x) + a)/x^3, x)`

**3.486.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^3 (a + \operatorname{arccosh}(cx))}{x^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^3}{x^3} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x^3,x)`output `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x^3, x)`

**3.487**  $\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^4} dx$

3.487.1 Optimal result . . . . . 3586  
 3.487.2 Mathematica [A] (verified) . . . . . 3587  
 3.487.3 Rubi [A] (warning: unable to verify) . . . . . 3587  
 3.487.4 Maple [A] (verified) . . . . . 3591  
 3.487.5 Fricas [A] (verification not implemented) . . . . . 3591  
 3.487.6 Sympy [F] . . . . . 3592  
 3.487.7 Maxima [A] (verification not implemented) . . . . . 3592  
 3.487.8 Giac [F] . . . . . 3593  
 3.487.9 Mupad [F(-1)] . . . . . 3593

**3.487.1 Optimal result**

Integrand size = 21, antiderivative size = 260

$$\int \frac{(d+ex^2)^3(a+b\operatorname{arccosh}(cx))}{x^4} dx = \frac{be^2(9c^2d+e)(1-c^2x^2)}{3c^3\sqrt{-1+cx}\sqrt{1+cx}} - \frac{bcd^3(1-c^2x^2)}{6x^2\sqrt{-1+cx}\sqrt{1+cx}}$$

$$- \frac{be^3(1-c^2x^2)^2}{9c^3\sqrt{-1+cx}\sqrt{1+cx}} - \frac{d^3(a+b\operatorname{arccosh}(cx))}{3x^3}$$

$$- \frac{3d^2e(a+b\operatorname{arccosh}(cx))}{x} + 3de^2x(a+b\operatorname{arccosh}(cx))$$

$$+ \frac{1}{3}e^3x^3(a+b\operatorname{arccosh}(cx))$$

$$+ \frac{bcd^2(c^2d+18e)\sqrt{-1+c^2x^2}\arctan(\sqrt{-1+c^2x^2})}{6\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
-1/3*d^3*(a+b*arccosh(c*x))/x^3-3*d^2*e*(a+b*arccosh(c*x))/x+3*d*e^2*x*(a+b*arccosh(c*x))+1/3*e^3*x^3*(a+b*arccosh(c*x))+1/3*b*e^2*(9*c^2*d+e)*(-c^2*x^2+1)/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/6*b*c*d^3*(-c^2*x^2+1)/x^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/9*b*e^3*(-c^2*x^2+1)^2/c^3/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/6*b*c*d^2*(c^2*d+18*e)*arctan((c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.487.2 Mathematica [A] (verified)**

Time = 0.24 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.71

$$\int \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx$$

$$= \frac{1}{6} \left( -\frac{2ad^3}{x^3} - \frac{18ad^2e}{x} + 18ade^2x + 2ae^3x^3 - \frac{b\sqrt{-1+cx}\sqrt{1+cx}(-3c^4d^3 + 4e^3x^2 + 2c^2e^2x^2(27d + ex^2))}{3c^3x^2} + \frac{2b(-d^3 - 9d^2ex^2 + 9de^2x^4 + e^3x^6) \operatorname{arccosh}(cx)}{x^3} - bcd^2(c^2d + 18e) \operatorname{arctan} \left( \frac{1}{\sqrt{-1+cx}\sqrt{1+cx}} \right) \right)$$

input `Integrate[((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x^4,x]`output `((-2*a*d^3)/x^3 - (18*a*d^2*e)/x + 18*a*d*e^2*x + 2*a*e^3*x^3 - (b*sqrt[-1 + c*x]*sqrt[1 + c*x]*(-3*c^4*d^3 + 4*e^3*x^2 + 2*c^2*e^2*x^2*(27*d + e*x^2)))/(3*c^3*x^2) + (2*b*(-d^3 - 9*d^2*e*x^2 + 9*d*e^2*x^4 + e^3*x^6)*ArcCosh[c*x])/x^3 - b*c*d^2*(c^2*d + 18*e)*ArcTan[1/(sqrt[-1 + c*x]*sqrt[1 + c*x])])/6`**3.487.3 Rubi [A] (warning: unable to verify)**Time = 0.85 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.76, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$ , Rules used = {6373, 27, 2113, 2331, 2124, 27, 1192, 1467, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^3 (a + \operatorname{barccosh}(cx))}{x^4} dx$$

$$\downarrow 6373$$

$$-bc \int -\frac{-e^3x^6 - 9de^2x^4 + 9d^2ex^2 + d^3}{3x^3\sqrt{cx-1}\sqrt{cx+1}} dx - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx))$$

---

3.487.  $\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x^4} dx$

$$\begin{aligned}
& \frac{1}{3}bc \int \frac{-e^3x^6 - 9de^2x^4 + 9d^2ex^2 + d^3}{x^3\sqrt{cx} - 1\sqrt{cx} + 1} dx - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + \\
& \quad 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{27} \\
& \frac{bc\sqrt{c^2x^2 - 1} \int \frac{-e^3x^6 - 9de^2x^4 + 9d^2ex^2 + d^3}{x^3\sqrt{c^2x^2 - 1}} dx}{3\sqrt{cx} - 1\sqrt{cx} + 1} - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + \\
& \quad 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2113} \\
& \frac{bc\sqrt{c^2x^2 - 1} \int \frac{-e^3x^6 - 9de^2x^4 + 9d^2ex^2 + d^3}{x^4\sqrt{c^2x^2 - 1}} dx^2}{6\sqrt{cx} - 1\sqrt{cx} + 1} - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + \\
& \quad 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2331} \\
& \frac{bc\sqrt{c^2x^2 - 1} \left( \int \frac{-2e^3x^4 - 18de^2x^2 + d^2(dc^2 + 18e)}{2x^2\sqrt{c^2x^2 - 1}} dx^2 + \frac{d^3\sqrt{c^2x^2 - 1}}{x^2} \right)}{6\sqrt{cx} - 1\sqrt{cx} + 1} - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \\
& \quad \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{2124} \\
& \frac{bc\sqrt{c^2x^2 - 1} \left( \frac{1}{2} \int \frac{-2e^3x^4 - 18de^2x^2 + d^2(dc^2 + 18e)}{x^2\sqrt{c^2x^2 - 1}} dx^2 + \frac{d^3\sqrt{c^2x^2 - 1}}{x^2} \right)}{6\sqrt{cx} - 1\sqrt{cx} + 1} - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \\
& \quad \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{27} \\
& \frac{bc\sqrt{c^2x^2 - 1} \left( \frac{1}{2} \int \frac{-2e^3x^4 - 18de^2x^2 + d^2(dc^2 + 18e)}{x^2\sqrt{c^2x^2 - 1}} dx^2 + \frac{d^3\sqrt{c^2x^2 - 1}}{x^2} \right)}{6\sqrt{cx} - 1\sqrt{cx} + 1} - \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \\
& \quad \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{1192} \\
& \frac{bc\sqrt{c^2x^2 - 1} \left( \frac{\int \frac{-2e^3x^8 - 2e^2(9dc^2 + 2e)x^4 + c^6d^3 - 2e^3 - 18c^2de^2 + 18c^4d^2e}{x^4 + 1} d\sqrt{c^2x^2 - 1}}{c^4} + \frac{d^3\sqrt{c^2x^2 - 1}}{x^2} \right)}{6\sqrt{cx} - 1\sqrt{cx} + 1} - \\
& \quad \frac{d^3(a + \operatorname{barccosh}(cx))}{3x^3} - \frac{3d^2e(a + \operatorname{barccosh}(cx))}{x} + 3de^2x(a + \operatorname{barccosh}(cx)) + \frac{1}{3}e^3x^3(a + \operatorname{barccosh}(cx)) \\
& \quad \downarrow \text{1467}
\end{aligned}$$

---

3.487.  $\int \frac{(d+ex^2)^3(a+\operatorname{barccosh}(cx))}{x^4} dx$

$$\frac{bc\sqrt{c^2x^2-1}\left(\frac{\int(-2e^3x^4-2e^2(9dc^2+e)+\frac{d^3c^6+18d^2ec^4}{x^4+1})d\sqrt{c^2x^2-1}}{c^4}+\frac{d^3\sqrt{c^2x^2-1}}{x^2}\right)}{6\sqrt{cx-1}\sqrt{cx+1}}-\frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3}-\frac{3d^2e(a+\operatorname{barccosh}(cx))}{x}+3de^2x(a+\operatorname{barccosh}(cx))+\frac{1}{3}e^3x^3(a+\operatorname{barccosh}(cx))}{2009}$$

$$-\frac{d^3(a+\operatorname{barccosh}(cx))}{3x^3}-\frac{3d^2e(a+\operatorname{barccosh}(cx))}{x}+3de^2x(a+\operatorname{barccosh}(cx))+\frac{1}{3}e^3x^3(a+\operatorname{barccosh}(cx))+\frac{bc\sqrt{c^2x^2-1}\left(\frac{c^4d^2\arctan(\sqrt{c^2x^2-1})(c^2d+18e)-2e^2\sqrt{c^2x^2-1}(9c^2d+e)-\frac{2}{3}e^3x^6}{c^4}+\frac{d^3\sqrt{c^2x^2-1}}{x^2}\right)}{6\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[((d + e*x^2)^3*(a + b*ArcCosh[c*x]))/x^4,x]`

output `-1/3*(d^3*(a + b*ArcCosh[c*x]))/x^3 - (3*d^2*e*(a + b*ArcCosh[c*x]))/x + 3*d*e^2*x*(a + b*ArcCosh[c*x]) + (e^3*x^3*(a + b*ArcCosh[c*x]))/3 + (b*c*sqrt[-1 + c^2*x^2]*((d^3*sqrt[-1 + c^2*x^2])/x^2 + ((-2*e^3*x^6)/3 - 2*e^2*(9*c^2*d + e)*sqrt[-1 + c^2*x^2] + c^4*d^2*(c^2*d + 18*e)*ArcTan[sqrt[-1 + c^2*x^2]])/c^4)/(6*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.487.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 1192 `Int[((d_) + (e_)*(x_)^(m_))*((f_) + (g_)*(x_)^(n_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[2/e^(n + 2*p + 1) Subst[Int[x^(2*m + 1)*(e*f - d*g + g*x^2)^n*(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4)^p, x], x, Sqrt[d + e*x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && ILtQ[n, 0] && IntegerQ[m + 1/2]`

rule 1467 `Int[((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IGtQ[p, 0] && IGtQ[q, -2]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2113 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

rule 2124 `Int[(Px_)*((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := With[{Qx = PolynomialQuotient[Px, a + b*x, x], R = PolynomialRemainder[Px, a + b*x, x]}, Simp[R*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((m + 1)*(b*c - a*d))), x] + Simp[1/((m + 1)*(b*c - a*d)) Int[(a + b*x)^(m + 1)*(c + d*x)^n*ExpandToSum[(m + 1)*(b*c - a*d)*Qx - d*R*(m + n + 2), x], x], x]] /; FreeQ[{a, b, c, d, n}, x] && PolyQ[Px, x] && LtQ[m, -1] && (IntegerQ[m] || !ILtQ[n, -1])`

rule 2331 `Int[(Pq_)*(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*SubstFor[x^2, Pq, x]*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x^2] && IntegerQ[(m - 1)/2]`

rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_))^(m_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.487.4 Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.04

method	result
parts	$a \left( \frac{e^3 x^3}{3} + 3d e^2 x - \frac{3d^2 e}{x} - \frac{d^3}{3x^3} \right) + b c^3 \left( \frac{\operatorname{arccosh}(cx) x^3 e^3}{3c^3} + \frac{3 \operatorname{arccosh}(cx) x d e^2}{c^3} - \frac{3 \operatorname{arccosh}(cx) d^2 e}{c^3 x} - \dots \right)$
derivativedivides	$c^3 \left( \frac{a \left( 3c^3 x d e^2 + \frac{c^3 x^3 e^3}{3} - \frac{c^3 d^3}{3x^3} - \frac{3c^3 d^2 e}{x} \right)}{c^6} + \dots \right) + \dots$
default	$c^3 \left( \frac{a \left( 3c^3 x d e^2 + \frac{c^3 x^3 e^3}{3} - \frac{c^3 d^3}{3x^3} - \frac{3c^3 d^2 e}{x} \right)}{c^6} + \dots \right) + \dots$

input `int((e*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x,method=_RETURNVERBOSE)`

output `a*(1/3*e^3*x^3+3*d*e^2*x-3*d^2*e/x-1/3*d^3/x^3)+b*c^3*(1/3/c^3*arccosh(c*x)*x^3*e^3+3/c^3*arccosh(c*x)*x*d*e^2-3/c^3*arccosh(c*x)*d^2*e/x-1/3*arccosh(c*x)*d^3/c^3/x^3-1/18/c^8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(3*arctan(1/(c^2*x^2-1))^(1/2))*c^8*d^3*x^2+54*arctan(1/(c^2*x^2-1))^(1/2))*c^6*d^2*e*x^2-3*(c^2*x^2-1)^(1/2)*c^6*d^3+54*c^4*d*e^2*(c^2*x^2-1)^(1/2)*x^2+2*e^3*(c^2*x^2-1)^(1/2)*c^4*x^4+4*e^3*c^2*x^2*(c^2*x^2-1)^(1/2))/(c^2*x^2-1)^(1/2)/x^2)`

### 3.487.5 Fracas [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.24

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx$$

$$= \frac{6ac^3e^3x^6 + 54ac^3de^2x^4 - 54ac^3d^2ex^2 - 6ac^3d^3 + 6(bc^6d^3 + 18bc^4d^2e)x^3 \arctan(-cx + \sqrt{c^2x^2 - 1}) + 6 \dots}{\dots}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x, algorithm="fricas")`



output 
$$\frac{1}{18}(6ac^3e^3x^6 + 54a^2c^3de^2x^4 - 54a^2c^3d^2e^2x^2 - 6a^2c^3d^3 + 6(b^2c^6d^3 + 18b^2c^4d^2e)x^3 \arctan(-cx + \sqrt{c^2x^2 - 1}) + 6(b^2c^3d^3 + 9b^2c^3d^2e - 9b^2c^3de^2 - b^2c^3e^3)x^3 \log(-cx + \sqrt{c^2x^2 - 1}) + 6(b^2c^3e^3x^6 + 9b^2c^3d^2e^2x^4 - 9b^2c^3d^2e^2x^2 - b^2c^3d^3 + (b^2c^3d^3 + 9b^2c^3d^2e - 9b^2c^3de^2 - b^2c^3e^3)x^3) \log(cx + \sqrt{c^2x^2 - 1}) - (2b^2c^2e^3x^5 - 3b^2c^4d^3x + 2(27b^2c^2de^2 + 2b^2e^3)x^3) \sqrt{c^2x^2 - 1}) / (c^3x^3)$$

### 3.487.6 Sympy [F]

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (d + ex^2)^3}{x^4} dx$$

input `integrate((e*x**2+d)**3*(a+b*acosh(c*x))/x**4,x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**3/x**4, x)`

### 3.487.7 Maxima [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.76

$$\begin{aligned} & \int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx \\ &= \frac{1}{3} ae^3 x^3 - \frac{1}{6} \left( \left( c^2 \arcsin \left( \frac{1}{c|x|} \right) - \frac{\sqrt{c^2 x^2 - 1}}{x^2} \right) c + \frac{2 \operatorname{arccosh}(cx)}{x^3} \right) bd^3 \\ & \quad - 3 \left( c \arcsin \left( \frac{1}{c|x|} \right) + \frac{\operatorname{arccosh}(cx)}{x} \right) bd^2 e \\ & \quad + \frac{1}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) be^3 \\ & \quad + 3ade^2 x + \frac{3(cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) bde^2}{c} - \frac{3ad^2 e}{x} - \frac{ad^3}{3x^3} \end{aligned}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x, algorithm="maxima")`

output  $1/3*a*e^3*x^3 - 1/6*((c^2*\arcsin(1/(c*abs(x)))) - \sqrt{c^2*x^2 - 1}/x^2)*c + 2*\operatorname{arccosh}(c*x)/x^3)*b*d^3 - 3*(c*\arcsin(1/(c*abs(x))) + \operatorname{arccosh}(c*x)/x)*b*d^2*e + 1/9*(3*x^3*\operatorname{arccosh}(c*x) - c*(\sqrt{c^2*x^2 - 1})*x^2/c^2 + 2*\sqrt{c^2*x^2 - 1}/c^4))*b*e^3 + 3*a*d*e^2*x + 3*(c*x*\operatorname{arccosh}(c*x) - \sqrt{c^2*x^2 - 1})*b*d*e^2/c - 3*a*d^2*e/x - 1/3*a*d^3/x^3$

### 3.487.8 Giac [F]

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(ex^2 + d)^3 (b \operatorname{arccosh}(cx) + a)}{x^4} dx$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))/x^4,x, algorithm="giac")`

output `integrate((e*x^2 + d)^3*(b*arccosh(c*x) + a)/x^4, x)`

### 3.487.9 Mupad [F(-1)]

Timed out.

$$\int \frac{(d + ex^2)^3 (a + b \operatorname{arccosh}(cx))}{x^4} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^3}{x^4} dx$$

input `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x^4,x)`

output `int(((a + b*acosh(c*x))*(d + e*x^2)^3)/x^4, x)`

### 3.488 $\int (d + ex^2)^4 (a + \operatorname{barccosh}(cx)) dx$

3.488.1 Optimal result . . . . .	3594
3.488.2 Mathematica [A] (verified) . . . . .	3595
3.488.3 Rubi [A] (verified) . . . . .	3595
3.488.4 Maple [A] (verified) . . . . .	3598
3.488.5 Fricas [A] (verification not implemented) . . . . .	3598
3.488.6 Sympy [F] . . . . .	3599
3.488.7 Maxima [A] (verification not implemented) . . . . .	3599
3.488.8 Giac [F(-2)] . . . . .	3600
3.488.9 Mupad [F(-1)] . . . . .	3600

#### 3.488.1 Optimal result

Integrand size = 18, antiderivative size = 395

$$\int (d + ex^2)^4 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{b(315c^8d^4 + 420c^6d^3e + 378c^4d^2e^2 + 180c^2de^3 + 35e^4)(1 - c^2x^2)}{315c^9\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$- \frac{4be(105c^6d^3 + 189c^4d^2e + 135c^2de^2 + 35e^3)(1 - c^2x^2)^2}{945c^9\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{2be^2(63c^4d^2 + 90c^2de + 35e^2)(1 - c^2x^2)^3}{525c^9\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{4be^3(9c^2d + 7e)(1 - c^2x^2)^4}{441c^9\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{be^4(1 - c^2x^2)^5}{81c^9\sqrt{-1 + cx}\sqrt{1 + cx}} + d^4x(a + \operatorname{barccosh}(cx)) + \frac{4}{3}d^3ex^3(a + \operatorname{barccosh}(cx))$$

$$+ \frac{6}{5}d^2e^2x^5(a + \operatorname{barccosh}(cx)) + \frac{4}{7}de^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^4x^9(a + \operatorname{barccosh}(cx))$$

```
output d^4*x*(a+b*arccosh(c*x))+4/3*d^3*e*x^3*(a+b*arccosh(c*x))+6/5*d^2*e^2*x^5*
(a+b*arccosh(c*x))+4/7*d*e^3*x^7*(a+b*arccosh(c*x))+1/9*e^4*x^9*(a+b*arcco
sh(c*x))+1/315*b*(315*c^8*d^4+420*c^6*d^3*e+378*c^4*d^2*e^2+180*c^2*d*e^3+
35*e^4)*(-c^2*x^2+1)/c^9/(c*x-1)^(1/2)/(c*x+1)^(1/2)-4/945*b*e*(105*c^6*d^
3+189*c^4*d^2*e+135*c^2*d*e^2+35*e^3)*(-c^2*x^2+1)^2/c^9/(c*x-1)^(1/2)/(c*
x+1)^(1/2)+2/525*b*e^2*(63*c^4*d^2+90*c^2*d*e+35*e^2)*(-c^2*x^2+1)^3/c^9/(
c*x-1)^(1/2)/(c*x+1)^(1/2)-4/441*b*e^3*(9*c^2*d+7*e)*(-c^2*x^2+1)^4/c^9/(c
*x-1)^(1/2)/(c*x+1)^(1/2)+1/81*b*e^4*(-c^2*x^2+1)^5/c^9/(c*x-1)^(1/2)/(c*x
+1)^(1/2)
```

### 3.488.2 Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 265, normalized size of antiderivative = 0.67

$$\int (d + ex^2)^4 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{315ax(315d^4 + 420d^3ex^2 + 378d^2e^2x^4 + 180de^3x^6 + 35e^4x^8) - \frac{b\sqrt{-1+cx}\sqrt{1+cx}(4480e^4+320c^2e^3(81d+7ex^2)+48c^4e^2}{99225}}$$

input `Integrate[(d + e*x^2)^4*(a + b*ArcCosh[c*x]),x]`

output `(315*a*x*(315*d^4 + 420*d^3*e*x^2 + 378*d^2*e^2*x^4 + 180*d*e^3*x^6 + 35*e^4*x^8) - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(4480*e^4 + 320*c^2*e^3*(81*d + 7*e*x^2) + 48*c^4*e^2*(1323*d^2 + 270*d*e*x^2 + 35*e^2*x^4) + 8*c^6*e*(110*25*d^3 + 3969*d^2*e*x^2 + 1215*d*e^2*x^4 + 175*e^3*x^6) + c^8*(99225*d^4 + 44100*d^3*e*x^2 + 23814*d^2*e^2*x^4 + 8100*d*e^3*x^6 + 1225*e^4*x^8)))/c^9 + 315*b*x*(315*d^4 + 420*d^3*e*x^2 + 378*d^2*e^2*x^4 + 180*d*e^3*x^6 + 35*e^4*x^8)*ArcCosh[c*x])/99225`

### 3.488.3 Rubi [A] (verified)

Time = 0.87 (sec) , antiderivative size = 343, normalized size of antiderivative = 0.87, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$ , Rules used = {6323, 27, 2113, 2331, 2389, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^4 (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6323}$$

$$-bc \int \frac{x(35e^4x^8 + 180de^3x^6 + 378d^2e^2x^4 + 420d^3ex^2 + 315d^4)}{315\sqrt{cx-1}\sqrt{cx+1}} dx + d^4x(a + \operatorname{barccosh}(cx)) + \frac{4}{3}d^3ex^3(a + \operatorname{barccosh}(cx)) + \frac{6}{5}d^2e^2x^5(a + \operatorname{barccosh}(cx)) + \frac{4}{7}de^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^4x^9(a + \operatorname{barccosh}(cx))$$

$$\downarrow \text{27}$$

$$-\frac{1}{315}bc \int \frac{x(35e^4x^8 + 180de^3x^6 + 378d^2e^2x^4 + 420d^3ex^2 + 315d^4)}{\sqrt{cx-1}\sqrt{cx+1}} dx + d^4x(a + \operatorname{barccosh}(cx)) + \frac{4}{3}d^3ex^3(a + \operatorname{barccosh}(cx)) + \frac{6}{5}d^2e^2x^5(a + \operatorname{barccosh}(cx)) + \frac{4}{7}de^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^4x^9(a + \operatorname{barccosh}(cx))$$

↓ 2113

$$-\frac{bc\sqrt{c^2x^2-1} \int \frac{x(35e^4x^8+180de^3x^6+378d^2e^2x^4+420d^3ex^2+315d^4)}{\sqrt{c^2x^2-1}} dx}{315\sqrt{cx-1}\sqrt{cx+1}} + d^4x(a + \operatorname{barccosh}(cx)) + \frac{4}{3}d^3ex^3(a + \operatorname{barccosh}(cx)) + \frac{6}{5}d^2e^2x^5(a + \operatorname{barccosh}(cx)) + \frac{4}{7}de^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^4x^9(a + \operatorname{barccosh}(cx))$$

↓ 2331

$$-\frac{bc\sqrt{c^2x^2-1} \int \frac{35e^4x^8+180de^3x^6+378d^2e^2x^4+420d^3ex^2+315d^4}{\sqrt{c^2x^2-1}} dx^2}{630\sqrt{cx-1}\sqrt{cx+1}} + d^4x(a + \operatorname{barccosh}(cx)) + \frac{4}{3}d^3ex^3(a + \operatorname{barccosh}(cx)) + \frac{6}{5}d^2e^2x^5(a + \operatorname{barccosh}(cx)) + \frac{4}{7}de^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^4x^9(a + \operatorname{barccosh}(cx))$$

↓ 2389

$$-\frac{bc\sqrt{c^2x^2-1} \int \left( \frac{35(c^2x^2-1)^{7/2}e^4}{c^8} + \frac{20(9dc^2+7e)(c^2x^2-1)^{5/2}e^3}{c^8} + \frac{6(63d^2c^4+90dec^2+35e^2)(c^2x^2-1)^{3/2}e^2}{c^8} + \frac{4(105d^3c^6+189d^2ec^4+126d^4e^2c^2+35d^5e^4)}{c^8} \right) dx}{630\sqrt{cx-1}\sqrt{cx+1}} + d^4x(a + \operatorname{barccosh}(cx)) + \frac{4}{3}d^3ex^3(a + \operatorname{barccosh}(cx)) + \frac{6}{5}d^2e^2x^5(a + \operatorname{barccosh}(cx)) + \frac{4}{7}de^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^4x^9(a + \operatorname{barccosh}(cx))$$

↓ 2009

$$d^4x(a + \operatorname{barccosh}(cx)) + \frac{4}{3}d^3ex^3(a + \operatorname{barccosh}(cx)) + \frac{6}{5}d^2e^2x^5(a + \operatorname{barccosh}(cx)) + \frac{4}{7}de^3x^7(a + \operatorname{barccosh}(cx)) + \frac{1}{9}e^4x^9(a + \operatorname{barccosh}(cx)) - \frac{bc\sqrt{c^2x^2-1} \left( \frac{40e^3(c^2x^2-1)^{7/2}(9c^2d+7e)}{7c^{10}} + \frac{70e^4(c^2x^2-1)^{9/2}}{9c^{10}} + \frac{12e^2(c^2x^2-1)^{5/2}(63c^4d^2+90c^2de+35e^2)}{5c^{10}} + \frac{8e(c^2x^2-1)^{3/2}(105c^6d^3+126c^5de^2+35c^4e^4)}{3c^{10}} \right)}{630\sqrt{cx-1}\sqrt{cx+1}}$$

input `Int[(d + e*x^2)^4*(a + b*ArcCosh[c*x]),x]`

output 
$$\begin{aligned} & -1/630*(b*c*\text{Sqrt}[-1 + c^2*x^2]*((2*(315*c^8*d^4 + 420*c^6*d^3*e + 378*c^4*d^2*e^2 + 180*c^2*d*e^3 + 35*e^4)*\text{Sqrt}[-1 + c^2*x^2])/c^{10} + (8*e*(105*c^6*d^3 + 189*c^4*d^2*e + 135*c^2*d*e^2 + 35*e^3)*(-1 + c^2*x^2)^{(3/2)})/(3*c^{10}) + (12*e^2*(63*c^4*d^2 + 90*c^2*d*e + 35*e^2)*(-1 + c^2*x^2)^{(5/2)})/(5*c^{10}) + (40*e^3*(9*c^2*d + 7*e)*(-1 + c^2*x^2)^{(7/2)})/(7*c^{10}) + (70*e^4*(-1 + c^2*x^2)^{(9/2)})/(9*c^{10}))/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) + d^4*x*(a + b*\text{ArcCosh}[c*x]) + (4*d^3*e*x^3*(a + b*\text{ArcCosh}[c*x]))/3 + (6*d^2*e^2*x^5*(a + b*\text{ArcCosh}[c*x]))/5 + (4*d*e^3*x^7*(a + b*\text{ArcCosh}[c*x]))/7 + (e^4*x^9*(a + b*\text{ArcCosh}[c*x]))/9 \end{aligned}$$

### 3.488.3.1 Defintions of rubi rules used

rule 27 
$$\text{Int}[(a_*)(F_x_), x\_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_*)(G_x_)] /; \text{FreeQ}[b, x]$$

rule 2009 
$$\text{Int}[u_, x\_Symbol] \rightarrow \text{Simp}[\text{IntSum}[u, x], x] /; \text{SumQ}[u]$$

rule 2113 
$$\text{Int}[(P_x)*((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[(a + b*x)^{\text{FracPart}[m]}*((c + d*x)^{\text{FracPart}[m]})/(\text{FracPart}[m]) \quad \text{Int}[P_x*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{PolyQ}[P_x, x] \ \&\& \ \text{EqQ}[b*c + a*d, 0] \ \&\& \ \text{EqQ}[m, n] \ \&\& \ !\text{IntegerQ}[m]$$

rule 2331 
$$\text{Int}[(P_q)*(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{Simp}[1/2 \quad \text{Subst}[\text{Int}[x^{((m-1)/2)*\text{SubstFor}[x^2, P_q, x]*(a + b*x)^p, x], x, x^2], x] /; \text{FreeQ}[\{a, b, p\}, x] \ \&\& \ \text{PolyQ}[P_q, x^2] \ \&\& \ \text{IntegerQ}[(m-1)/2]$$

rule 2389 
$$\text{Int}[(P_q)*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}, x\_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[P_q*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, n\}, x] \ \&\& \ \text{PolyQ}[P_q, x] \ \&\& \ (\text{IGtQ}[p, 0] \ || \ \text{EqQ}[n, 1])$$

rule 6323 
$$\text{Int}[(a_.) + \text{ArcCosh}[(c_.)*(x_.)]*(b_.)*((d_.) + (e_.)*(x_.)^2)^{(p_.)}, x\_Symbol] \rightarrow \text{With}[\{u = \text{IntHide}[(d + e*x^2)^p, x]\}, \text{Simp}[(a + b*\text{ArcCosh}[c*x]) \quad u, x] - \text{Simp}[b*c \quad \text{Int}[\text{SimplifyIntegrand}[u/(\text{Sqrt}[1 + c*x]*\text{Sqrt}[-1 + c*x]), x], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[c^2*d + e, 0] \ \&\& \ (\text{IGtQ}[p, 0] \ || \ \text{ILtQ}[p + 1/2, 0])$$

### 3.488.4 Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 302, normalized size of antiderivative = 0.76

method	result
parts	$a\left(\frac{1}{9}e^4x^9 + \frac{4}{7}de^3x^7 + \frac{6}{5}d^2e^2x^5 + \frac{4}{3}x^3d^3e + d^4x\right) + \frac{b\left(\frac{c \operatorname{arccosh}(cx)e^4x^9}{9} + \frac{4c \operatorname{arccosh}(cx)de^3x^7}{7} + \frac{6c \operatorname{arccosh}(cx)d^2e^2x^5}{5}\right)}{c^8}$
derivativedivides	$\frac{a\left(d^4c^9x + \frac{4}{3}d^3c^9ex^3 + \frac{6}{5}d^2c^9e^2x^5 + \frac{4}{7}dc^9e^3x^7 + \frac{1}{9}e^4c^9x^9\right)}{c^8} + \frac{b\left(\operatorname{arccosh}(cx)d^4c^9x + \frac{4 \operatorname{arccosh}(cx)d^3c^9ex^3}{3} + \frac{6 \operatorname{arccosh}(cx)d^2c^9e^2x^5}{5}\right)}{c^8}$
default	$\frac{a\left(d^4c^9x + \frac{4}{3}d^3c^9ex^3 + \frac{6}{5}d^2c^9e^2x^5 + \frac{4}{7}dc^9e^3x^7 + \frac{1}{9}e^4c^9x^9\right)}{c^8} + \frac{b\left(\operatorname{arccosh}(cx)d^4c^9x + \frac{4 \operatorname{arccosh}(cx)d^3c^9ex^3}{3} + \frac{6 \operatorname{arccosh}(cx)d^2c^9e^2x^5}{5}\right)}{c^8}$

input `int((e*x^2+d)^4*(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `a*(1/9*e^4*x^9+4/7*d*e^3*x^7+6/5*d^2*e^2*x^5+4/3*x^3*d^3*e+d^4*x)+b/c*(1/9*c*arccosh(c*x)*e^4*x^9+4/7*c*arccosh(c*x)*d*e^3*x^7+6/5*c*arccosh(c*x)*d^2*e^2*x^5+4/3*c*arccosh(c*x)*d^3*e*x^3+arccosh(c*x)*d^4*c*x-1/99225/c^8*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(1225*c^8*e^4*x^8+8100*c^8*d*e^3*x^6+23814*c^8*d^2*e^2*x^4+1400*c^6*e^4*x^6+44100*c^8*d^3*e*x^2+9720*c^6*d*e^3*x^4+99225*c^8*d^4+31752*c^6*d^2*e^2*x^2+1680*c^4*e^4*x^4+88200*c^6*d^3*e+12960*c^4*d*e^3*x^2+63504*c^4*d^2*e^2+2240*c^2*e^4*x^2+25920*c^2*d*e^3+4480*e^4))`

### 3.488.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 333, normalized size of antiderivative = 0.84

$$\int (d + ex^2)^4 (a + b \operatorname{arccosh}(cx)) dx$$


---


$$= \frac{11025 ac^9 e^4 x^9 + 56700 ac^9 d e^3 x^7 + 119070 ac^9 d^2 e^2 x^5 + 132300 ac^9 d^3 e x^3 + 99225 ac^9 d^4 x + 315 (35 bc^9 e^4 x^9$$

input `integrate((e*x^2+d)^4*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output  $\frac{1}{99225}(11025a^9c^4e^4x^9 + 56700a^9c^9d^3e^3x^7 + 119070a^9c^9d^2e^2x^5 + 132300a^9c^9d^3e^3x^3 + 99225a^9c^9d^4e^4x + 315(35b^3c^9e^4x^9 + 180b^3c^9d^3e^3x^7 + 378b^3c^9d^2e^2x^5 + 420b^3c^9d^3e^3x^3 + 315b^3c^9d^4e^4x)\log(cx + \sqrt{c^2x^2 - 1}) - (1225b^8c^8e^4x^8 + 99225b^8c^8d^4 + 88200b^6c^6d^3e + 63504b^4c^4d^2e^2 + 25920b^2c^2d^3e^3 + 100(81b^8c^8d^3e^3 + 14b^6c^6e^4)x^6 + 4480b^6e^4 + 6(3969b^8c^8d^2e^2 + 1620b^6c^6d^3e^3 + 280b^4c^4e^4)x^4 + 4(11025b^8c^8d^3e + 7938b^6c^6d^2e^2 + 3240b^4c^4d^3e^3 + 560b^2c^2e^4)x^2)\sqrt{c^2x^2 - 1})/c^9$

### 3.488.6 Sympy [F]

$$\int (d + ex^2)^4 (a + b \operatorname{arccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (d + ex^2)^4 dx$$

input `integrate((e*x**2+d)**4*(a+b*acosh(c*x)),x)`

output `Integral((a + b*acosh(c*x))*(d + e*x**2)**4, x)`

### 3.488.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 415, normalized size of antiderivative = 1.05

$$\begin{aligned} \int (d + ex^2)^4 (a + b \operatorname{arccosh}(cx)) dx &= \frac{1}{9} ae^4 x^9 + \frac{4}{7} ade^3 x^7 + \frac{6}{5} ad^2 e^2 x^5 \\ &+ \frac{4}{3} ad^3 ex^3 + \frac{4}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) bd^3 e \\ &+ \frac{2}{25} \left( 15x^5 \operatorname{arccosh}(cx) - \left( \frac{3\sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4\sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8\sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) bd^2 e^2 \\ &+ \frac{4}{245} \left( 35x^7 \operatorname{arccosh}(cx) - \left( \frac{5\sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6\sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8\sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16\sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) bde^3 \\ &+ \frac{1}{2835} \left( 315x^9 \operatorname{arccosh}(cx) - \left( \frac{35\sqrt{c^2 x^2 - 1} x^8}{c^2} + \frac{40\sqrt{c^2 x^2 - 1} x^6}{c^4} + \frac{48\sqrt{c^2 x^2 - 1} x^4}{c^6} + \frac{64\sqrt{c^2 x^2 - 1} x^2}{c^8} \right) c \right) bde^4 \\ &+ ad^4 x + \frac{(cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) bd^4}{c} \end{aligned}$$



input `integrate((e*x^2+d)^4*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output 
$$\begin{aligned} & 1/9*a*e^4*x^9 + 4/7*a*d*e^3*x^7 + 6/5*a*d^2*e^2*x^5 + 4/3*a*d^3*e*x^3 + 4/ \\ & 9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1) \\ & /c^4))*b*d^3*e + 2/25*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 \\ & + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c)*b*d^2*e^2 + 4/ \\ & 245*(35*x^7*arccosh(c*x) - (5*sqrt(c^2*x^2 - 1)*x^6/c^2 + 6*sqrt(c^2*x^2 - \\ & 1)*x^4/c^4 + 8*sqrt(c^2*x^2 - 1)*x^2/c^6 + 16*sqrt(c^2*x^2 - 1)/c^8)*c)*b \\ & *d*e^3 + 1/2835*(315*x^9*arccosh(c*x) - (35*sqrt(c^2*x^2 - 1)*x^8/c^2 + 40 \\ & *sqrt(c^2*x^2 - 1)*x^6/c^4 + 48*sqrt(c^2*x^2 - 1)*x^4/c^6 + 64*sqrt(c^2*x^ \\ & 2 - 1)*x^2/c^8 + 128*sqrt(c^2*x^2 - 1)/c^{10})*c)*b*e^4 + a*d^4*x + (c*x*arc \\ & cosh(c*x) - sqrt(c^2*x^2 - 1))*b*d^4/c \end{aligned}$$

### 3.488.8 Giac [F(-2)]

Exception generated.

$$\int (d + ex^2)^4 (a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: RuntimeError}$$

input `integrate((e*x^2+d)^4*(a+b*arccosh(c*x)),x, algorithm="giac")`

output Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const ve  
cteur & l) Error: Bad Argument Value

### 3.488.9 Mupad [F(-1)]

Timed out.

$$\int (d + ex^2)^4 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (ex^2 + d)^4 dx$$

input `int((a + b*acosh(c*x))*(d + e*x^2)^4,x)`

output `int((a + b*acosh(c*x))*(d + e*x^2)^4, x)`

$$3.489 \quad \int \frac{x^4(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx$$

3.489.1 Optimal result	3602
3.489.2 Mathematica [C] (warning: unable to verify)	3603
3.489.3 Rubi [A] (verified)	3604
3.489.4 Maple [C] (verified)	3606
3.489.5 Fricas [F]	3607
3.489.6 Sympy [F]	3607
3.489.7 Maxima [F(-2)]	3608
3.489.8 Giac [F]	3608
3.489.9 Mupad [F(-1)]	3608

**3.489.1 Optimal result**

Integrand size = 21, antiderivative size = 627

$$\begin{aligned}
\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d + ex^2} dx = & -\frac{adx}{e^2} + \frac{bd\sqrt{-1+cx}\sqrt{1+cx}}{ce^2} \\
& - \frac{2b\sqrt{-1+cx}\sqrt{1+cx}}{9c^3e} - \frac{bx^2\sqrt{-1+cx}\sqrt{1+cx}}{9ce} \\
& - \frac{bdx\operatorname{arccosh}(cx)}{e^2} + \frac{x^3(a + \operatorname{barccosh}(cx))}{3e} \\
& + \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2e^{5/2}} \\
& - \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2e^{5/2}} \\
& + \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2e^{5/2}} \\
& - \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2e^{5/2}} \\
& - \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2e^{5/2}} \\
& + \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2e^{5/2}} \\
& - \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2e^{5/2}} \\
& + \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2e^{5/2}}
\end{aligned}$$

output

```

-a*d*x/e^2-b*d*x*arccosh(c*x)/e^2+1/3*x^3*(a+b*arccosh(c*x))/e+1/2*(-d)^(3/2)*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(5/2)-1/2*(-d)^(3/2)*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(5/2)+1/2*(-d)^(3/2)*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)-1/2*(-d)^(3/2)*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)-1/2*b*(-d)^(3/2)*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(5/2)+1/2*b*(-d)^(3/2)*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(5/2)-1/2*b*(-d)^(3/2)*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)+1/2*b*(-d)^(3/2)*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)+b*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/e^2-2/9*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3/e-1/9*b*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/e

```

### 3.489.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 0.86 (sec) , antiderivative size = 524, normalized size of antiderivative = 0.84

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = -\frac{adx}{e^2} + \frac{ax^3}{3e} + \frac{ad^{3/2} \arctan\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{e^{5/2}}$$

$$+ \frac{b\left(\frac{4d\sqrt{e}\left(\sqrt{\frac{-1+cx}{1+cx}}(1+cx) - cx \operatorname{arccosh}(cx)\right)}{c} - \frac{4e^{3/2}\left(\sqrt{-1+cx}\sqrt{1+cx}(2+c^2x^2) - 3c^3x^3 \operatorname{arccosh}(cx)\right)}{9c^3}\right)}{e^{5/2}} - id^{3/2}\left(\operatorname{arccosh}(cx)\right)$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

output

```

-((a*d*x)/e^2) + (a*x^3)/(3*e) + (a*d^(3/2)*ArcTan[(Sqrt[e]*x)/Sqrt[d]])/e
^(5/2) + (b*((4*d*Sqrt[e]*(Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) - c*x*ArcC
osh[c*x])))/c - (4*e^(3/2)*(Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2 + c^2*x^2) - 3*
c^3*x^3*ArcCosh[c*x]))/(9*c^3) - I*d^(3/2)*(ArcCosh[c*x]*(-ArcCosh[c*x] +
2*(Log[1 + (I*Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[d] - Sqrt[c^2*d + e]))] + Log
[1 + (I*Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[d] + Sqrt[c^2*d + e]))]) + 2*PolyL
og[2, (I*Sqrt[e]*E^ArcCosh[c*x])/(-c*Sqrt[d] + Sqrt[c^2*d + e])] + 2*Pol
yLog[2, ((-I)*Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[d] + Sqrt[c^2*d + e])] + I*
d^(3/2)*(ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(Log[1 + (I*Sqrt[e]*E^ArcCosh[c*x
]))/(-c*Sqrt[d] + Sqrt[c^2*d + e])] + Log[1 - (I*Sqrt[e]*E^ArcCosh[c*x])/
(c*Sqrt[d] + Sqrt[c^2*d + e]))]) + 2*PolyLog[2, (I*Sqrt[e]*E^ArcCosh[c*x])
/(c*Sqrt[d] - Sqrt[c^2*d + e])] + 2*PolyLog[2, (I*Sqrt[e]*E^ArcCosh[c*x])/
(c*Sqrt[d] + Sqrt[c^2*d + e]))))/(4*e^(5/2))

```

### 3.489.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 627, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^4(a + \operatorname{barccosh}(cx))}{d + ex^2} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{d^2(a + \operatorname{barccosh}(cx))}{e^2(d + ex^2)} - \frac{d(a + \operatorname{barccosh}(cx))}{e^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{e} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\begin{aligned}
& \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2e^{5/2}} - \\
& \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2e^{5/2}} + \\
& \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2e^{5/2}} - \\
& \frac{(-d)^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2e^{5/2}} + \frac{x^3(a + \operatorname{barccosh}(cx))}{e^2} - \frac{adx}{e^2} - \\
& \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2e^{5/2}} + \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2e^{5/2}} - \\
& \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2e^{5/2}} + \frac{b(-d)^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2e^{5/2}} - \\
& \frac{bdx \operatorname{arccosh}(cx)}{e^2} - \frac{2b\sqrt{cx - 1}\sqrt{cx + 1}}{9c^3e} + \frac{bd\sqrt{cx - 1}\sqrt{cx + 1}}{ce^2} - \frac{bx^2\sqrt{cx - 1}\sqrt{cx + 1}}{9ce}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

output

```

-((a*d*x)/e^2) + (b*d*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*e^2) - (2*b*Sqrt[-1
+ c*x]*Sqrt[1 + c*x])/(9*c^3*e) - (b*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(9
*c*e) - (b*d*x*ArcCosh[c*x])/e^2 + (x^3*(a + b*ArcCosh[c*x]))/(3*e) + ((-d
)^(3/2)*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d]
- Sqrt[-(c^2*d) - e])])/(2*e^(5/2)) - ((-d)^(3/2)*(a + b*ArcCosh[c*x])*Log
[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^(5/
2)) + ((-d)^(3/2)*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c
*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^(5/2)) - ((-d)^(3/2)*(a + b*ArcCosh
[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])
)/(2*e^(5/2)) - (b*(-d)^(3/2)*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqr
t[-d] - Sqrt[-(c^2*d) - e])])]/(2*e^(5/2)) + (b*(-d)^(3/2)*PolyLog[2, (Sqr
t[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])]/(2*e^(5/2)) - (b*
(-d)^(3/2)*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*
d) - e])])]/(2*e^(5/2)) + (b*(-d)^(3/2)*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x]
)/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])]/(2*e^(5/2))

```

3.489.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

3.489.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 45.91 (sec) , antiderivative size = 364, normalized size of antiderivative = 0.58

method	result
parts	$\frac{ax^3}{3e} - \frac{adx}{e^2} + \frac{ad^2 \arctan\left(\frac{ex}{\sqrt{de}}\right)}{e^2\sqrt{de}} - \frac{bx^2\sqrt{cx-1}\sqrt{cx+1}}{9ce} - \frac{bdx \operatorname{arccosh}(cx)}{e^2} + \frac{bd\sqrt{cx-1}\sqrt{cx+1}}{ce^2} + \frac{bcd^2}{ce^2} \left( \frac{-R1}{-R1} \right)$
derivativedivides	$\frac{-\frac{ac^5dx}{e^2} + \frac{ac^5x^3}{3e} + \frac{ac^5d^2 \arctan\left(\frac{ex}{\sqrt{de}}\right)}{e^2\sqrt{de}} + \frac{be^4\sqrt{cx+1}\sqrt{cx-1}d}{e^2} - \frac{bc^5 \operatorname{arccosh}(cx)dx}{e^2} + \frac{bc^6d^2}{e^2} \left( \frac{-R1 = \operatorname{RootOf}(e_{-Z^4 + (4c^2d+2e)} - \dots)}{-R1 = \operatorname{RootOf}(e_{-Z^4 + (4c^2d+2e)} - \dots)} \right)$
default	$\frac{-\frac{ac^5dx}{e^2} + \frac{ac^5x^3}{3e} + \frac{ac^5d^2 \arctan\left(\frac{ex}{\sqrt{de}}\right)}{e^2\sqrt{de}} + \frac{be^4\sqrt{cx+1}\sqrt{cx-1}d}{e^2} - \frac{bc^5 \operatorname{arccosh}(cx)dx}{e^2} + \frac{bc^6d^2}{e^2} \left( \frac{-R1 = \operatorname{RootOf}(e_{-Z^4 + (4c^2d+2e)} - \dots)}{-R1 = \operatorname{RootOf}(e_{-Z^4 + (4c^2d+2e)} - \dots)} \right)$

```
input int(x^4*(a+b*arccosh(c*x))/(e*x^2+d), x, method=_RETURNVERBOSE)
```

3.489.  $\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx$

output  $\frac{1}{3}ax^3/e - a*d*x/e^2 + a*d^2/e^2/(d*e)^{(1/2)}*\arctan(e*x/(d*e)^{(1/2)}) - 1/9*b*x^2*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c/e - b*d*x*\operatorname{arccosh}(c*x)/e^2 + b*d*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c/e^2 + 1/2*b*c/e^2*d^2*\sum(_R1/(_R1^2*e+2*c^2*d+e))*(\operatorname{arccosh}(c*x)*\ln((\_R1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/_R1)+\operatorname{dilog}((\_R1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/_R1)),\_R1=\operatorname{RootOf}(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e)) - 1/2*b*c/e^2*d^2*\sum(1/_R1/(_R1^2*e+2*c^2*d+e))*(\operatorname{arccosh}(c*x)*\ln((\_R1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/_R1)+\operatorname{dilog}((\_R1-c*x-(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/_R1)),\_R1=\operatorname{RootOf}(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/3*b/e*\operatorname{arccosh}(c*x)*x^3 - 2/9*b*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3/e$

### 3.489.5 Fricas [F]

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^4}{ex^2 + d} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*x^4*arccosh(c*x) + a*x^4)/(e*x^2 + d), x)`

### 3.489.6 Sympy [F]

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{d + ex^2} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(e*x**2+d),x)`

output `Integral(x**4*(a + b*acosh(c*x))/(d + e*x**2), x)`



**3.489.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.489.8 Giac [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{ex^2 + d} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^4/(e*x^2 + d), x)`

**3.489.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{d + ex^2} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{ex^2 + d} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d + e*x^2),x)`

output `int((x^4*(a + b*acosh(c*x)))/(d + e*x^2), x)`

# 3.490 $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx$

3.490.1 Optimal result	3609
3.490.2 Mathematica [A] (warning: unable to verify)	3610
3.490.3 Rubi [A] (verified)	3611
3.490.4 Maple [C] (warning: unable to verify)	3612
3.490.5 Fricas [F]	3613
3.490.6 Sympy [F]	3614
3.490.7 Maxima [F]	3614
3.490.8 Giac [F(-2)]	3614
3.490.9 Mupad [F(-1)]	3615

## 3.490.1 Optimal result

Integrand size = 21, antiderivative size = 521

$$\int \frac{x^3(a + b\operatorname{arccosh}(cx))}{d + ex^2} dx = -\frac{bx\sqrt{-1 + cx}\sqrt{1 + cx}}{4ce} - \frac{b\operatorname{arccosh}(cx)}{4c^2e} + \frac{x^2(a + b\operatorname{arccosh}(cx))}{2e} + \frac{d(a + b\operatorname{arccosh}(cx))^2}{2be^2} - \frac{d(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2e^2} - \frac{d(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2e^2} - \frac{d(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2e^2} - \frac{d(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2e^2} - \frac{bd \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2e^2} - \frac{bd \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2e^2} - \frac{bd \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2e^2} - \frac{bd \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2e^2}$$

output

$$\begin{aligned}
& -1/4*b*arccosh(c*x)/c^2/e+1/2*x^2*(a+b*arccosh(c*x))/e+1/2*d*(a+b*arccosh(c*x))^2/b/e^2-1/2*d*(a+b*arccosh(c*x))*\ln(1-(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2-1/2*d*(a+b*arccosh(c*x))*\ln(1+(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2-1/2*d*(a+b*arccosh(c*x))*\ln(1-(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2-1/2*d*(a+b*arccosh(c*x))*\ln(1+(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2-1/2*b*d*polylog(2,-(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2-1/2*b*d*polylog(2,(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2-1/2*b*d*polylog(2,-(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2-1/2*b*d*polylog(2,(c*x+(c*x-1)^{1/2}*(c*x+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2-1/4*b*x*(c*x-1)^{1/2}*(c*x+1)^{1/2}/c/e
\end{aligned}$$

### 3.490.2 Mathematica [A] (warning: unable to verify)

Time = 0.42 (sec) , antiderivative size = 512, normalized size of antiderivative = 0.98

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \frac{-2ac^2ex^2 + bcex\sqrt{-1 + cx}\sqrt{1 + cx} - 2bc^2ex^2 \operatorname{arccosh}(cx) - 2bc^2d \operatorname{arccosh}(cx)^2 + 2b \operatorname{arctanh}\left(\sqrt{\frac{-1+cx}{1+cx}}\right)}{d + ex^2}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

output

$$\begin{aligned}
& -1/4*(-2*a*c^2*e*x^2 + b*c*e*x*\sqrt{-1 + c*x}*\sqrt{1 + c*x} - 2*b*c^2*e*x^2*\operatorname{ArcCosh}[c*x] - 2*b*c^2*d*\operatorname{ArcCosh}[c*x]^2 + 2*b*e*\operatorname{ArcTanh}[\sqrt{(-1 + c*x)/(1 + c*x)}]) + 2*b*c^2*d*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + (\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(c*\sqrt{-d} - \sqrt{-(c^2*d) - e})] + 2*b*c^2*d*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + (\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(-c*\sqrt{-d}) + \sqrt{-(c^2*d) - e})] + 2*b*c^2*d*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - (\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(c*\sqrt{-d} + \sqrt{-(c^2*d) - e})] + 2*b*c^2*d*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + (\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(c*\sqrt{-d} + \sqrt{-(c^2*d) - e})] + 2*a*c^2*d*\operatorname{Log}[d + e*x^2] + 2*b*c^2*d*\operatorname{PolyLog}[2, (\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(c*\sqrt{-d} - \sqrt{-(c^2*d) - e})] + 2*b*c^2*d*\operatorname{PolyLog}[2, (\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(-c*\sqrt{-d}) + \sqrt{-(c^2*d) - e})] + 2*b*c^2*d*\operatorname{PolyLog}[2, -((\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(c*\sqrt{-d} + \sqrt{-(c^2*d) - e}))] + 2*b*c^2*d*\operatorname{PolyLog}[2, (\sqrt{e}*E^{\operatorname{ArcCosh}[c*x]})/(c*\sqrt{-d} + \sqrt{-(c^2*d) - e})])]/(c^2*e^2)
\end{aligned}$$

**3.490.3 Rubi [A] (verified)**

Time = 1.28 (sec) , antiderivative size = 521, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^3(a + \operatorname{barccosh}(cx))}{d + ex^2} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{x(a + \operatorname{barccosh}(cx))}{e} - \frac{dx(a + \operatorname{barccosh}(cx))}{e(d + ex^2)} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{d(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} \right)}{2e^2} - \\
 & \frac{d(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1 \right)}{2e^2} - \\
 & \frac{d(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} \right)}{2e^2} - \frac{d(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1 \right)}{2e^2} + \\
 & \frac{d(a + \operatorname{barccosh}(cx))^2}{2be^2} + \frac{x^2(a + \operatorname{barccosh}(cx))}{2e} - \frac{bd \operatorname{PolyLog} \left( 2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2e^2} - \\
 & \frac{bd \operatorname{PolyLog} \left( 2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2e^2} - \frac{bd \operatorname{PolyLog} \left( 2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2e^2} - \\
 & \frac{bd \operatorname{PolyLog} \left( 2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2e^2} - \frac{\operatorname{barccosh}(cx)}{4c^2e} - \frac{bx\sqrt{cx - 1}\sqrt{cx + 1}}{4ce}
 \end{aligned}$$

input `Int[(x^3*(a + b*ArcCosh[c*x]))/(d + e*x^2), x]`

```
output -1/4*(b*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*e) - (b*ArcCosh[c*x])/(4*c^2*e)
+ (x^2*(a + b*ArcCosh[c*x]))/(2*e) + (d*(a + b*ArcCosh[c*x])^2)/(2*b*e^2)
- (d*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] -
Sqrt[-(c^2*d) - e])])/(2*e^2) - (d*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E
^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^2) - (d*(a + b*Arc
Cosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) -
e])])/(2*e^2) - (d*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(
c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^2) - (b*d*PolyLog[2, -((Sqrt[e]*E^
ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])]/(2*e^2) - (b*d*PolyLog[
2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^2) -
(b*d*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e
])])]/(2*e^2) - (b*d*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqr
t[-(c^2*d) - e])])/(2*e^2)
```

### 3.490.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n_.)*((f_.)*(x_)^m_.)*((d_) + (e
_.)*(x_)^2)^p_., x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

### 3.490.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.17 (sec) , antiderivative size = 2111, normalized size of antiderivative = 4.05

method	result	size
derivativedivides	Expression too large to display	2111
default	Expression too large to display	2111
parts	Expression too large to display	2118

```
input int(x^3*(a+b*arccosh(c*x))/(e*x^2+d),x,method=_RETURNVERBOSE)
```

output `1/c^4*(1/2*a*c^4/e*x^2-1/2*a*c^4*d/e^2*ln(c^2*e*x^2+c^2*d)+b*c^2*(1/2*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^2*d/e^3/(c^2*d+e)*polylog(2,e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))+(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^2*d/e^3/(c^2*d+e)*ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*arccosh(c*x)+1/2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d/e^2/(c^2*d+e)*arccosh(c*x)*ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e))+(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^4*d^2/e^4/(c^2*d+e)*ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*arccosh(c*x)+1/16*(-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x+2*c^2*x^2-1)*(1+2*arccosh(c*x))/e+1/16*(-1+2*arccosh(c*x))/e*(2*c^2*x^2-1+2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c*x)-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)/e^4*c^4*d^2*ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*arccosh(c*x)+1/8*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)/e^2/(c^2*d+e)*polylog(2,e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))-1/2*c^2*d/e^2*sum((_R1^2*e+4*c^2*d+2*e)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*...`

### 3.490.5 Fracas [F]

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^3}{ex^2 + d} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*x^3*arccosh(c*x) + a*x^3)/(e*x^2 + d), x)`

**3.490.6 Sympy [F]**

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{d + ex^2} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(e*x**2+d),x)`

output `Integral(x**3*(a + b*acosh(c*x))/(d + e*x**2), x)`

**3.490.7 Maxima [F]**

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^3}{ex^2 + d} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="maxima")`

output `1/2*a*(x^2/e - d*log(e*x^2 + d)/e^2) + b*integrate(x^3*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(e*x^2 + d), x)`

**3.490.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.490.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{ex^2 + d} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d + e*x^2),x)`output `int((x^3*(a + b*acosh(c*x)))/(d + e*x^2), x)`



$$3.491 \quad \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx$$

3.491.1 Optimal result	3616
3.491.2 Mathematica [C] (verified)	3617
3.491.3 Rubi [A] (verified)	3618
3.491.4 Maple [C] (verified)	3619
3.491.5 Fricas [F]	3620
3.491.6 Sympy [F]	3621
3.491.7 Maxima [F(-2)]	3621
3.491.8 Giac [F]	3621
3.491.9 Mupad [F(-1)]	3622

### 3.491.1 Optimal result

Integrand size = 21, antiderivative size = 544

$$\begin{aligned} \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx = & \frac{ax}{e} - \frac{b\sqrt{-1+cx}\sqrt{1+cx}}{ce} + \frac{b\operatorname{arccosh}(cx)}{e} \\ & + \frac{\sqrt{-d}(a+b\operatorname{arccosh}(cx))\log\left(1-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \\ & - \frac{\sqrt{-d}(a+b\operatorname{arccosh}(cx))\log\left(1+\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \\ & + \frac{\sqrt{-d}(a+b\operatorname{arccosh}(cx))\log\left(1-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \\ & - \frac{\sqrt{-d}(a+b\operatorname{arccosh}(cx))\log\left(1+\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \\ & - \frac{b\sqrt{-d}\operatorname{PolyLog}\left(2,-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \\ & + \frac{b\sqrt{-d}\operatorname{PolyLog}\left(2,\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \\ & - \frac{b\sqrt{-d}\operatorname{PolyLog}\left(2,-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \\ & + \frac{b\sqrt{-d}\operatorname{PolyLog}\left(2,\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^{3/2}} \end{aligned}$$

output  $a*x/e+b*x*\operatorname{arccosh}(c*x)/e+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}-1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}-1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}-1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}+1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}-1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}+1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2}))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2}))*(-d)^{(1/2)}/e^{(3/2)}-b*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c/e$

### 3.491.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 457, normalized size of antiderivative = 0.84

$$\int \frac{x^2(a + b\operatorname{arccosh}(cx))}{d + ex^2} dx$$

$$= \frac{4ac\sqrt{ex} - 4ac\sqrt{d}\arctan\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) + ib\left(4i\sqrt{e}\left(\sqrt{\frac{-1+cx}{1+cx}}(1+cx) - cx\operatorname{arccosh}(cx)\right) - c\sqrt{d}\left(\operatorname{arccosh}(cx)\left(\operatorname{arccosh}(cx) + \frac{cx}{\sqrt{d}}\right)\right)\right)}{d + ex^2}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

output  $(4*a*c*\operatorname{Sqrt}[e]*x - 4*a*c*\operatorname{Sqrt}[d]*\operatorname{ArcTan}[(\operatorname{Sqrt}[e]*x)/\operatorname{Sqrt}[d]] + I*b*((4*I)*\operatorname{Sqrt}[e]*(\operatorname{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x) - c*x*\operatorname{ArcCosh}[c*x]) - c*\operatorname{Sqrt}[d]*(\operatorname{ArcCosh}[c*x]*(\operatorname{ArcCosh}[c*x] - 2*(\operatorname{Log}[1 + (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] - \operatorname{Sqrt}[c^2*d + e])]) + \operatorname{Log}[1 + (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])])) - 2*\operatorname{PolyLog}[2, (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(- (c*\operatorname{Sqrt}[d]) + \operatorname{Sqrt}[c^2*d + e])]) - 2*\operatorname{PolyLog}[2, ((-I)*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])]) + c*\operatorname{Sqrt}[d]*(\operatorname{ArcCosh}[c*x]*(\operatorname{ArcCosh}[c*x] - 2*(\operatorname{Log}[1 + (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(- (c*\operatorname{Sqrt}[d]) + \operatorname{Sqrt}[c^2*d + e])]) + \operatorname{Log}[1 - (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])])) - 2*\operatorname{PolyLog}[2, (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] - \operatorname{Sqrt}[c^2*d + e])]) - 2*\operatorname{PolyLog}[2, (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])])))/(4*c*e^{(3/2)})$

---

3.491.  $\int \frac{x^2(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx$

**3.491.3 Rubi [A] (verified)**

Time = 1.23 (sec) , antiderivative size = 544, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + \operatorname{barccosh}(cx))}{d + ex^2} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{a + \operatorname{barccosh}(cx)}{e} - \frac{d(a + \operatorname{barccosh}(cx))}{e(d + ex^2)} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} \right)}{2e^{3/2}} - \\
 & \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1 \right)}{2e^{3/2}} + \\
 & \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} \right)}{2e^{3/2}} - \\
 & \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1 \right)}{2e^{3/2}} + \frac{ax}{e} - \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2e^{3/2}} + \\
 & \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2e^{3/2}} - \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2e^{3/2}} + \\
 & \frac{b\sqrt{-d} \operatorname{PolyLog} \left( 2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2e^{3/2}} + \frac{b \operatorname{arccosh}(cx)}{e} - \frac{b\sqrt{cx - 1}\sqrt{cx + 1}}{ce}
 \end{aligned}$$

input `Int[(x^2*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

```
output (a*x)/e - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*e) + (b*x*ArcCosh[c*x])/e +
(Sqrt[-d]*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d]
- Sqrt[-(c^2*d) - e])])/(2*e^(3/2)) - (Sqrt[-d]*(a + b*ArcCosh[c*x])*Log
[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^(3/
2)) + (Sqrt[-d]*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*S
qrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^(3/2)) - (Sqrt[-d]*(a + b*ArcCosh[c*x
])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2
*e^(3/2)) - (b*Sqrt[-d]*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d]
- Sqrt[-(c^2*d) - e])])/(2*e^(3/2)) + (b*Sqrt[-d]*PolyLog[2, (Sqrt[e]*E^A
rcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^(3/2)) - (b*Sqrt[-d]
*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])
)/(2*e^(3/2)) + (b*Sqrt[-d]*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d]
+ Sqrt[-(c^2*d) - e])])/(2*e^(3/2))
```

### 3.491.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

### 3.491.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 23.72 (sec) , antiderivative size = 284, normalized size of antiderivative = 0.52

method	result
parts	$\frac{ax}{e} - \frac{ad \arctan\left(\frac{ex}{\sqrt{de}}\right)}{e\sqrt{de}} + \frac{bx \operatorname{arccosh}(cx)}{e} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{ce} - \frac{bcd \left( \frac{\sum \dots}{\dots} \right)}{\dots}$
derivativedivides	$\frac{\frac{ac^3x}{e} - \frac{ac^3d \arctan\left(\frac{ex}{\sqrt{de}}\right)}{e\sqrt{de}} - \frac{bc^2\sqrt{cx-1}\sqrt{cx+1}}{e} + \frac{bc^3 \operatorname{arccosh}(cx)x}{e}}{\dots}$
default	$\frac{\frac{ac^3x}{e} - \frac{ac^3d \arctan\left(\frac{ex}{\sqrt{de}}\right)}{e\sqrt{de}} - \frac{bc^2\sqrt{cx-1}\sqrt{cx+1}}{e} + \frac{bc^3 \operatorname{arccosh}(cx)x}{e}}{\dots}$

```
input int(x^2*(a+b*arccosh(c*x))/(e*x^2+d),x,method=_RETURNVERBOSE)
```

```
output a*x/e-a*d/e/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2))+b*x*arccosh(c*x)/e-b*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/e-1/2*b*c/e*d*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/2*b*c/e*d*sum(1/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))
```

### 3.491.5 Fracas [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^2}{ex^2 + d} dx$$

```
input integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="fricas")
```

```
output integral((b*x^2*arccosh(c*x) + a*x^2)/(e*x^2 + d), x)
```

**3.491.6 Sympy [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{d + ex^2} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(e*x**2+d), x)`

output `Integral(x**2*(a + b*acosh(c*x))/(d + e*x**2), x)`

**3.491.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d), x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.491.8 Giac [F]**

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{ex^2 + d} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^2/(e*x^2 + d), x)`

**3.491.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{ex^2 + d} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d + e*x^2),x)`output `int((x^2*(a + b*acosh(c*x)))/(d + e*x^2), x)`

### 3.492 $\int \frac{x(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx$

3.492.1 Optimal result . . . . .	3623
3.492.2 Mathematica [A] (verified) . . . . .	3624
3.492.3 Rubi [A] (verified) . . . . .	3625
3.492.4 Maple [C] (warning: unable to verify) . . . . .	3626
3.492.5 Fricas [F] . . . . .	3627
3.492.6 Sympy [F] . . . . .	3628
3.492.7 Maxima [F] . . . . .	3628
3.492.8 Giac [F] . . . . .	3628
3.492.9 Mupad [F(-1)] . . . . .	3629

#### 3.492.1 Optimal result

Integrand size = 19, antiderivative size = 449

$$\int \frac{x(a + b\operatorname{arccosh}(cx))}{d + ex^2} dx = -\frac{(a + b\operatorname{arccosh}(cx))^2}{2be} + \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2e} + \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2e} + \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2e} + \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2e} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2e}$$



output

```

-1/2*(a+b*arccosh(c*x))^2/b/e+1/2*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e+1/2*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e+1/2*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e+1/2*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e+1/2*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e+1/2*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e+1/2*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e+1/2*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e

```

### 3.492.2 Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 447, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int \frac{x(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = & -\frac{b \operatorname{arccosh}(cx)^2}{2e} + \frac{b \operatorname{arccosh}(cx) \log\left(1 - \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{2e} \\
& + \frac{b \operatorname{arccosh}(cx) \log\left(1 + \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{2e} \\
& + \frac{b \operatorname{arccosh}(cx) \log\left(1 - \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{2e} \\
& + \frac{b \operatorname{arccosh}(cx) \log\left(1 + \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{2e} \\
& + \frac{a \log(d + ex^2)}{2e} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{2e} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{2e} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{2e} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{2e}
\end{aligned}$$

input `Integrate[(x*(a + b*ArcCosh[c*x]))/(d + e*x^2), x]`

output 
$$-1/2*(b*\text{ArcCosh}[c*x]^2)/e + (b*\text{ArcCosh}[c*x]*\text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})]/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e]))/(2*e) + (b*\text{ArcCosh}[c*x]*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})]/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e]))/(2*e) + (b*\text{ArcCosh}[c*x]*\text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})]/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e]))/(2*e) + (b*\text{ArcCosh}[c*x]*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})]/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e]))/(2*e) + (a*\text{Log}[d + e*x^2])/(2*e) + (b*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e]))]/(2*e) + (b*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e]))]/(2*e) + (b*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e]))]/(2*e) + (b*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e]))]/(2*e)$$

### 3.492.3 Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 449, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx$$

↓ 6374

$$\int \left( \frac{a + b \operatorname{arccosh}(cx)}{2\sqrt{e}(\sqrt{-d} + \sqrt{ex})} - \frac{a + b \operatorname{arccosh}(cx)}{2\sqrt{e}(\sqrt{-d} - \sqrt{ex})} \right) dx$$

↓ 2009

$$\frac{(a + b \operatorname{arccosh}(cx)) \log \left( 1 - \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} \right)}{2e} + \frac{(a + b \operatorname{arccosh}(cx)) \log \left( \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1 \right)}{2e} +$$

$$\frac{(a + b \operatorname{arccosh}(cx)) \log \left( 1 - \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} \right)}{2e} + \frac{(a + b \operatorname{arccosh}(cx)) \log \left( \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1 \right)}{2e} -$$

$$\frac{(a + b \operatorname{arccosh}(cx))^2}{2be} + \frac{b \operatorname{PolyLog} \left( 2, -\frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{dc^2 - e}} \right)}{2e} + \frac{b \operatorname{PolyLog} \left( 2, \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{dc^2 - e}} \right)}{2e} +$$

$$\frac{b \operatorname{PolyLog} \left( 2, -\frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2e} + \frac{b \operatorname{PolyLog} \left( 2, \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2e}$$

input `Int[(x*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

---

3.492.  $\int \frac{x(a+b\operatorname{arccosh}(cx))}{d+ex^2} dx$

```
output -1/2*(a + b*ArcCosh[c*x])^2/(b*e) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]
 *E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e) + ((a + b*ArcCo
 sh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e]
 )])/(2*e) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt
 [-d] + Sqrt[-(c^2*d) - e])])/(2*e) + ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]
 ]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e) + (b*PolyLog[2
 , -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e) +
 (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])
 / (2*e) + (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2
 *d) - e])])/(2*e) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] +
 Sqrt[-(c^2*d) - e])])/(2*e)
```

### 3.492.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((f_.)*(x_))^m_)*((d_) + (e
 _)*(x_)^2)^p_., x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
 (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
 + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

### 3.492.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.88 (sec) , antiderivative size = 1987, normalized size of antiderivative = 4.43

method	result	size
derivativedivides	Expression too large to display	1987
default	Expression too large to display	1987
parts	Expression too large to display	1988

```
input int(x*(a+b*arccosh(c*x))/(e*x^2+d), x, method=_RETURNVERBOSE)
```

output  $\frac{1}{c^2} \left( \frac{1}{2} a c^2 / e \ln(c^2 e x^2 + c^2 d) + b c^2 \left( -\frac{1}{2} / e \operatorname{arccosh}(c x)^2 + \frac{1}{2} (2 c^2 d - 2 (d c^2 (c^2 d + e))^{1/2} + e) / e^2 \ln(1 - e (c x + (c x - 1)^{1/2}) (c x + 1)^{1/2}) \right)^2 / (-2 c^2 d - 2 (d c^2 (c^2 d + e))^{1/2} - e) \operatorname{arccosh}(c x) + \frac{1}{4} (-2 (d c^2 (c^2 d + e))^{1/2} c^2 d + 2 c^4 d^2 + 2 c^2 d e - (d c^2 (c^2 d + e))^{1/2} e) / c^2 d e / (c^2 d + e) \operatorname{arccosh}(c x)^2 - \frac{1}{8} (-2 (d c^2 (c^2 d + e))^{1/2} c^2 d + 2 c^4 d^2 + 2 c^2 d e - (d c^2 (c^2 d + e))^{1/2} e) / c^2 d e / (c^2 d + e) \operatorname{arccosh}(c x)^2 - \frac{1}{4} (-2 (d c^2 (c^2 d + e))^{1/2} c^2 d + 2 c^4 d^2 + 2 c^2 d e - (d c^2 (c^2 d + e))^{1/2} e) / c^2 d e / (c^2 d + e) \operatorname{polylog}(2, e (c x + (c x - 1)^{1/2}) (c x + 1)^{1/2})^2 / (-2 c^2 d - 2 (d c^2 (c^2 d + e))^{1/2} - e) \right) - \frac{1}{4} (d c^2 (c^2 d + e))^{1/2} / d c^2 / (c^2 d + e) \operatorname{arccosh}(c x) * \ln(1 - e (c x + (c x - 1)^{1/2}) (c x + 1)^{1/2})^2 / (-2 c^2 d + 2 (d c^2 (c^2 d + e))^{1/2} - e) + (-2 (d c^2 (c^2 d + e))^{1/2} c^2 d + 2 c^4 d^2 + 2 c^2 d e - (d c^2 (c^2 d + e))^{1/2} e) / e^2 / (c^2 d + e) \operatorname{arccosh}(c x)^2 - \frac{1}{2} (-2 (d c^2 (c^2 d + e))^{1/2} c^2 d + 2 c^4 d^2 + 2 c^2 d e - (d c^2 (c^2 d + e))^{1/2} e) / e^2 / (c^2 d + e) \operatorname{polylog}(2, e (c x + (c x - 1)^{1/2}) (c x + 1)^{1/2})^2 / (-2 c^2 d - 2 (d c^2 (c^2 d + e))^{1/2} - e) - (2 c^2 d - 2 (d c^2 (c^2 d + e))^{1/2} + e) / e^3 \operatorname{polylog}(2, e (c x + (c x - 1)^{1/2}) (c x + 1)^{1/2})^2 / (-2 c^2 d - 2 (d c^2 (c^2 d + e))^{1/2} - e) * d c^2 + \frac{1}{2} (d c^2 (c^2 d + e))^{1/2} / e / (c^2 d + e) \operatorname{arccosh}(c x)^2 - \frac{1}{4} (d c^2 (c^2 d + e))^{1/2} / e / (c^2 d + e) \operatorname{polylog}(2, e (c x + (c x - 1)^{1/2}) (c x + 1)^{1/2})^2 / (-2 c^2 d + 2 (d c^2 (c^2 d + e))^{1/2} - e) - \frac{1}{4} (-2 (d c^2 (c^2 d + e))^{1/2} c^2 d + 2 c^4 d^2 + 2 c^2 d e - (d c^2 (c^2 d + e))^{1/2} e) / c^2 d e / (c^2 d + e) * \ln(1 - e (c x + (c x - 1)^{1/2}) (c x + 1) \dots$

### 3.492.5 Fracas [F]

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x}{ex^2 + d} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*x*arccosh(c*x) + a*x)/(e*x^2 + d), x)`

**3.492.6 Sympy [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{d + ex^2} dx$$

input `integrate(x*(a+b*acosh(c*x))/(e*x**2+d), x)`

output `Integral(x*(a + b*acosh(c*x))/(d + e*x**2), x)`

**3.492.7 Maxima [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{ex^2 + d} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d), x, algorithm="maxima")`

output `b*integrate(x*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(e*x^2 + d), x) + 1/2  
*a*log(e*x^2 + d)/e`

**3.492.8 Giac [F]**

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{ex^2 + d} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x/(e*x^2 + d), x)`

**3.492.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{ex^2 + d} dx$$

input `int((x*(a + b*acosh(c*x)))/(d + e*x^2), x)`output `int((x*(a + b*acosh(c*x)))/(d + e*x^2), x)`

### 3.493 $\int \frac{a+b\operatorname{arccosh}(cx)}{d+ex^2} dx$

3.493.1 Optimal result . . . . .	3630
3.493.2 Mathematica [A] (verified) . . . . .	3631
3.493.3 Rubi [A] (verified) . . . . .	3632
3.493.4 Maple [C] (verified) . . . . .	3633
3.493.5 Fracas [F] . . . . .	3634
3.493.6 Sympy [F] . . . . .	3635
3.493.7 Maxima [F(-2)] . . . . .	3635
3.493.8 Giac [F] . . . . .	3635
3.493.9 Mupad [F(-1)] . . . . .	3636

#### 3.493.1 Optimal result

Integrand size = 18, antiderivative size = 501

$$\int \frac{a + b\operatorname{arccosh}(cx)}{d + ex^2} dx = \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}} + \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2\sqrt{-d}\sqrt{e}}$$

output  $\frac{1}{2}(a+b\operatorname{arccosh}(cx))\ln(1-(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}-(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}-\frac{1}{2}(a+b\operatorname{arccosh}(cx))\ln(1+(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}-(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}+\frac{1}{2}(a+b\operatorname{arccosh}(cx))\ln(1-(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}+(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}-\frac{1}{2}(a+b\operatorname{arccosh}(cx))\ln(1+(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}+(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}-\frac{1}{2}b\operatorname{polylog}(2,-(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}-(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}+\frac{1}{2}b\operatorname{polylog}(2,(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}-(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}-\frac{1}{2}b\operatorname{polylog}(2,-(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}+(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}+\frac{1}{2}b\operatorname{polylog}(2,(cx+(cx-1)^{1/2})(cx+1)^{1/2})e^{1/2}/(c(-d)^{1/2}+(-c^2d-e)^{1/2}))/(-d)^{1/2}/e^{1/2}$

### 3.493.2 Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 397, normalized size of antiderivative = 0.79

$$\int \frac{a + b\operatorname{arccosh}(cx)}{d + ex^2} dx$$

$$= -\left((a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)\right) + (a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-c\sqrt{-d}+\sqrt{-c^2d-e}}\right) + (a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right) + (a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-c\sqrt{-d}+\sqrt{-c^2d-e}}\right)$$

input `Integrate[(a + b*ArcCosh[c*x])/(d + e*x^2), x]`

output  $(-((a + b\operatorname{ArcCosh}[c*x])\operatorname{Log}[1 + (\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c\operatorname{Sqrt}[-d] - \operatorname{Sqrt}[-(c^2*d) - e])]) + (a + b\operatorname{ArcCosh}[c*x])\operatorname{Log}[1 + (\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(-(c\operatorname{Sqrt}[-d]) + \operatorname{Sqrt}[-(c^2*d) - e])]) + (a + b\operatorname{ArcCosh}[c*x])\operatorname{Log}[1 - (\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c\operatorname{Sqrt}[-d] + \operatorname{Sqrt}[-(c^2*d) - e])]) - (a + b\operatorname{ArcCosh}[c*x])\operatorname{Log}[1 + (\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c\operatorname{Sqrt}[-d] + \operatorname{Sqrt}[-(c^2*d) - e])]) + b\operatorname{PolyLog}[2, (\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c\operatorname{Sqrt}[-d] - \operatorname{Sqrt}[-(c^2*d) - e])]) - b\operatorname{PolyLog}[2, (\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(-(c\operatorname{Sqrt}[-d]) + \operatorname{Sqrt}[-(c^2*d) - e])]) - b\operatorname{PolyLog}[2, -((\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c\operatorname{Sqrt}[-d] + \operatorname{Sqrt}[-(c^2*d) - e])]) + b\operatorname{PolyLog}[2, (\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c\operatorname{Sqrt}[-d] + \operatorname{Sqrt}[-(c^2*d) - e])])]/(2\operatorname{Sqrt}[-d]*\operatorname{Sqrt}[e])$



**3.493.3 Rubi [A] (verified)**

Time = 1.16 (sec) , antiderivative size = 501, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{d + ex^2} dx \\
 & \quad \downarrow \text{6324} \\
 & \int \left( \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx))}{2d(\sqrt{-d} - \sqrt{ex})} + \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx))}{2d(\sqrt{-d} + \sqrt{ex})} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} \right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1 \right)}{2\sqrt{-d}\sqrt{e}} + \\
 & \frac{(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} \right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1 \right)}{2\sqrt{-d}\sqrt{e}} - \\
 & \frac{b \operatorname{PolyLog} \left( 2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2\sqrt{-d}\sqrt{e}} + \frac{b \operatorname{PolyLog} \left( 2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2\sqrt{-d}\sqrt{e}} - \\
 & \frac{b \operatorname{PolyLog} \left( 2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2\sqrt{-d}\sqrt{e}} + \frac{b \operatorname{PolyLog} \left( 2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2\sqrt{-d}\sqrt{e}}
 \end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d + e*x^2), x]`

```
output ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) - ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) - ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) - (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e]))])/(2*Sqrt[-d]*Sqrt[e]) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) - (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e]))])/(2*Sqrt[-d]*Sqrt[e]) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e])
```

### 3.493.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6324 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])
```

### 3.493.4 Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.28 (sec) , antiderivative size = 232, normalized size of antiderivative = 0.46

method	result
parts	$\frac{a \arctan\left(\frac{ex}{\sqrt{de}}\right)}{\sqrt{de}} + \frac{bc \left( \frac{-R1 \left( \operatorname{arccosh}(cx) \ln\left(\frac{R1-cx-\sqrt{cx-1}\sqrt{cx+1}}{-R1}\right) + \operatorname{dilog}\left(\frac{-R1-cx-\sqrt{cx-1}\sqrt{cx+1}}{-R1}\right)\right)}{-R1^2 e+2c^2 d+e} \right)}{2}$
derivativedivides	$\frac{ac \arctan\left(\frac{ex}{\sqrt{de}}\right)}{\sqrt{de}} + bc^2 \left( \frac{-R1 \left( \operatorname{arccosh}(cx) \ln\left(\frac{R1-cx-\sqrt{cx-1}\sqrt{cx+1}}{-R1}\right) + \operatorname{dilog}\left(\frac{-R1-cx-\sqrt{cx-1}\sqrt{cx+1}}{-R1}\right)\right)}{-R1^2 e+2c^2 d+e} \right)$
default	$\frac{ac \arctan\left(\frac{ex}{\sqrt{de}}\right)}{\sqrt{de}} + bc^2 \left( \frac{-R1 \left( \operatorname{arccosh}(cx) \ln\left(\frac{R1-cx-\sqrt{cx-1}\sqrt{cx+1}}{-R1}\right) + \operatorname{dilog}\left(\frac{-R1-cx-\sqrt{cx-1}\sqrt{cx+1}}{-R1}\right)\right)}{-R1^2 e+2c^2 d+e} \right)$

input `int((a+b*arccosh(c*x))/(e*x^2+d),x,method=_RETURNVERBOSE)`

output `a/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2))+1/2*b*c*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))-1/2*b*c*sum(1/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))`

### 3.493.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{d + ex^2} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{ex^2 + d} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e*x^2 + d), x)`

**3.493.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{d + ex^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{d + ex^2} dx$$

input `integrate((a+b*acosh(c*x))/(e*x**2+d), x)`

output `Integral((a + b*acosh(c*x))/(d + e*x**2), x)`

**3.493.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + \operatorname{barccosh}(cx)}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d), x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.493.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{d + ex^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{ex^2 + d} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d), x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(e*x^2 + d), x)`

**3.493.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{d + ex^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{ex^2 + d} dx$$

input `int((a + b*acosh(c*x))/(d + e*x^2),x)`output `int((a + b*acosh(c*x))/(d + e*x^2), x)`

# 3.494 $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d+ex^2)} dx$

3.494.1 Optimal result . . . . .	3637
3.494.2 Mathematica [C] (verified) . . . . .	3638
3.494.3 Rubi [A] (verified) . . . . .	3639
3.494.4 Maple [C] (warning: unable to verify) . . . . .	3640
3.494.5 Fracas [F] . . . . .	3641
3.494.6 Sympy [F] . . . . .	3642
3.494.7 Maxima [F] . . . . .	3642
3.494.8 Giac [F] . . . . .	3642
3.494.9 Mupad [F(-1)] . . . . .	3643

## 3.494.1 Optimal result

Integrand size = 21, antiderivative size = 489

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x(d + ex^2)} dx = \frac{(a + b\operatorname{arccosh}(cx))^2}{bd} + \frac{(a + b\operatorname{arccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)})}{d}$$

$$- \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d}$$

$$- \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d}$$

$$- \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d}$$

$$- \frac{(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d}$$

$$- \frac{b \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d}$$

$$- \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d}$$

$$- \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d}$$

output  $(a+b\operatorname{arccosh}(cx))^2/b/d+(a+b\operatorname{arccosh}(cx))*\ln(1+(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))^2/d-1/2*(a+b\operatorname{arccosh}(cx))*\ln(1-(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/d-1/2*(a+b\operatorname{arccosh}(cx))*\ln(1+(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/d-1/2*(a+b\operatorname{arccosh}(cx))*\ln(1-(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/d-1/2*(a+b\operatorname{arccosh}(cx))*\ln(1+(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/d-1/2*b*\operatorname{polylog}(2,-1/(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))/d-1/2*b*\operatorname{polylog}(2,-(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/d-1/2*b*\operatorname{polylog}(2,(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/d-1/2*b*\operatorname{polylog}(2,-(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/d-1/2*b*\operatorname{polylog}(2,(cx+(cx-1)^{1/2})*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/d$

### 3.494.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.22 (sec) , antiderivative size = 410, normalized size of antiderivative = 0.84

$$\int \frac{a + b\operatorname{arccosh}(cx)}{x(d + ex^2)} dx =$$

$$\frac{-2b\operatorname{arccosh}(cx)^2 - 2b\operatorname{arccosh}(cx) \log\left(1 + e^{-2\operatorname{arccosh}(cx)}\right) + b\operatorname{arccosh}(cx) \log\left(1 + \frac{i\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{d - \sqrt{c^2d + e}}}\right) + b\operatorname{arccosh}(cx) \log\left(1 + \frac{i\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{d - \sqrt{c^2d + e}}}\right)}{d}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*(d + e*x^2)),x]`

output  $-1/2*(-2*b*\operatorname{ArcCosh}[c*x]^2 - 2*b*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + E^{(-2*\operatorname{ArcCosh}[c*x])}] + b*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] - \operatorname{Sqrt}[c^2*d + e])] + b*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(-c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])] + b*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 - (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])] + b*\operatorname{ArcCosh}[c*x]*\operatorname{Log}[1 + (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])] - 2*a*\operatorname{Log}[x] + a*\operatorname{Log}[d + e*x^2] + b*\operatorname{PolyLog}[2, -E^{(-2*\operatorname{ArcCosh}[c*x])}] + b*\operatorname{PolyLog}[2, (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] - \operatorname{Sqrt}[c^2*d + e])] + b*\operatorname{PolyLog}[2, (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(-c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])] + b*\operatorname{PolyLog}[2, ((-I)*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])] + b*\operatorname{PolyLog}[2, (I*\operatorname{Sqrt}[e]*E^{\operatorname{ArcCosh}[c*x]})/(c*\operatorname{Sqrt}[d] + \operatorname{Sqrt}[c^2*d + e])])/d$

---

3.494.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d+ex^2)} dx$

**3.494.3 Rubi [A] (verified)**

Time = 1.34 (sec) , antiderivative size = 489, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{a + \operatorname{barccosh}(cx)}{dx} - \frac{ex(a + \operatorname{barccosh}(cx))}{d(d + ex^2)} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} \right)}{2d} - \frac{(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1 \right)}{2d} \\
 & - \frac{(a + \operatorname{barccosh}(cx)) \log \left( 1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} \right)}{2d} - \frac{(a + \operatorname{barccosh}(cx)) \log \left( \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1 \right)}{2d} + \\
 & \frac{(a + \operatorname{barccosh}(cx))^2}{bd} + \frac{\log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))}{d} - \\
 & \frac{b \operatorname{PolyLog} \left( 2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2d} - \frac{b \operatorname{PolyLog} \left( 2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}} \right)}{2d} - \\
 & \frac{b \operatorname{PolyLog} \left( 2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2d} - \frac{b \operatorname{PolyLog} \left( 2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}} \right)}{2d} - \frac{b \operatorname{PolyLog} \left( 2, -e^{-2\operatorname{arccosh}(cx)} \right)}{2d}
 \end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d + e*x^2)),x]`



```
output (a + b*ArcCosh[c*x])^2/(b*d) + ((a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh
[c*x])])/d - ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqr
t[-d] - Sqrt[-(c^2*d) - e])])/(2*d) - ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[
e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*d) - ((a + b*Arc
Cosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) -
e])])/(2*d) - ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sq
rt[-d] + Sqrt[-(c^2*d) - e])])/(2*d) - (b*PolyLog[2, -E^(-2*ArcCosh[c*x])])
)/(2*d) - (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^
2*d) - e])]))/(2*d) - (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] -
Sqrt[-(c^2*d) - e])])/(2*d) - (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c
*Sqrt[-d] + Sqrt[-(c^2*d) - e])]))/(2*d) - (b*PolyLog[2, (Sqrt[e]*E^ArcCos
h[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*d)
```

### 3.494.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((f_.)*(x_)^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

### 3.494.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.60 (sec) , antiderivative size = 381, normalized size of antiderivative = 0.78

method	result
parts	$-\frac{a \ln(ex^2+d)}{2d} + \frac{a \ln(x)}{d} + b \left( -\frac{e \left( \frac{(-R1^2+1) \left( \operatorname{arccosh}(cx) \ln \left( \frac{R1-}{-R1=\operatorname{RootOf}(e-Z^4+(4c^2d+2e)-Z^2+e)} \right)}{4d} \right)}{\right)}{\right)$
derivativedivides	$\frac{a \ln(cx)}{d} - \frac{a \ln(c^2ex^2+c^2d)}{2d} + \frac{b \operatorname{arccosh}(cx) \ln(1+i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{d} + \frac{b \operatorname{arccosh}(cx) \ln(1-i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{d}$
default	$\frac{a \ln(cx)}{d} - \frac{a \ln(c^2ex^2+c^2d)}{2d} + \frac{b \operatorname{arccosh}(cx) \ln(1+i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{d} + \frac{b \operatorname{arccosh}(cx) \ln(1-i(cx+\sqrt{cx-1}\sqrt{cx+1}))}{d}$

input `int((a+b*arccosh(c*x))/x/(e*x^2+d),x,method=_RETURNVERBOSE)`

output

```
-1/2*a/d*ln(e*x^2+d)+a/d*ln(x)+b*(-1/4*e/d*sum((_R1^2+1)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/d*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/d*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/d*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/d*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-1/4/d*sum((_R1^2*e+4*c^2*d+e)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e)))
```

### 3.494.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(ex^2 + d)x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e*x^3 + d*x), x)`

**3.494.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(d + ex^2)} dx$$

input `integrate((a+b*acosh(c*x))/x/(e*x**2+d),x)`

output `Integral((a + b*acosh(c*x))/(x*(d + e*x**2)), x)`

**3.494.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d),x, algorithm="maxima")`

output `-1/2*a*(log(e*x^2 + d)/d - 2*log(x)/d) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(e*x^3 + d*x), x)`

**3.494.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)*x), x)`

**3.494.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(ex^2 + d)} dx$$

input `int((a + b*acosh(c*x))/(x*(d + e*x^2)),x)`output `int((a + b*acosh(c*x))/(x*(d + e*x^2)), x)`

### 3.495 $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d+ex^2)} dx$

3.495.1 Optimal result . . . . .	3644
3.495.2 Mathematica [A] (verified) . . . . .	3646
3.495.3 Rubi [A] (verified) . . . . .	3647
3.495.4 Maple [C] (warning: unable to verify) . . . . .	3649
3.495.5 Fracas [F] . . . . .	3650
3.495.6 Sympy [F] . . . . .	3650
3.495.7 Maxima [F(-2)] . . . . .	3651
3.495.8 Giac [F] . . . . .	3651
3.495.9 Mupad [F(-1)] . . . . .	3651

#### 3.495.1 Optimal result

Integrand size = 21, antiderivative size = 543

$$\begin{aligned} \int \frac{a + b\operatorname{arccosh}(cx)}{x^2(d + ex^2)} dx = & -\frac{a + b\operatorname{arccosh}(cx)}{dx} + \frac{bc \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d} \\ & + \frac{\sqrt{e}(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \\ & - \frac{\sqrt{e}(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \\ & + \frac{\sqrt{e}(a + b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \\ & - \frac{\sqrt{e}(a + b\operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \\ & - \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \\ & + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \\ & - \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \\ & + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2(-d)^{3/2}} \end{aligned}$$

output  $(-a-b*\operatorname{arccosh}(c*x))/d/x+b*c*\arctan((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}-1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}-1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}-1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}+1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}-1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}+1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(3/2)}$

**3.495.2 Mathematica [A] (verified)**

Time = 0.99 (sec) , antiderivative size = 549, normalized size of antiderivative = 1.01

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x^2(d + ex^2)} dx = & \frac{1}{2} \left( -\frac{2(a + \operatorname{barccosh}(cx))}{dx} + \frac{2bc\sqrt{-1 + c^2x^2} \arctan(\sqrt{-1 + c^2x^2})}{d\sqrt{-1 + cx}\sqrt{1 + cx}} \right. \\
& + \frac{d\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{(-d)^{5/2}} \\
& + \frac{\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{(-d)^{3/2}} \\
& + \frac{\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{(-d)^{3/2}} \\
& + \frac{d\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{(-d)^{5/2}} \\
& + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{(-d)^{3/2}} \\
& + \frac{bd\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{(-d)^{5/2}} \\
& + \frac{bd\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{(-d)^{5/2}} \\
& \left. + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{(-d)^{3/2}} \right)
\end{aligned}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^2*(d + e*x^2)),x]`

output  $((-2*(a + b*\text{ArcCosh}[c*x]))/(d*x) + (2*b*c*\text{Sqrt}[-1 + c^2*x^2]*\text{ArcTan}[\text{Sqrt}[-1 + c^2*x^2]])/(d*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) + (d*\text{Sqrt}[e]*(a + b*\text{ArcCosh}[c*x])*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (\text{Sqrt}[e]*(a + b*\text{ArcCosh}[c*x])*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (\text{Sqrt}[e]*(a + b*\text{ArcCosh}[c*x])*\text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(3/2)} + (\text{Sqrt}[e]*(a + b*\text{ArcCosh}[c*x])*\text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(3/2)} + (d*\text{Sqrt}[e]*(a + b*\text{ArcCosh}[c*x])*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (b*\text{Sqrt}[e]*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(3/2)} + (b*d*\text{Sqrt}[e]*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(-(c*\text{Sqrt}[-d]) + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (b*d*\text{Sqrt}[e]*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (b*\text{Sqrt}[e]*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(3/2}))/2$

### 3.495.3 Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 543, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d + ex^2)} dx$$

$$\downarrow \text{6374}$$

$$\int \left( \frac{a + b \operatorname{arccosh}(cx)}{dx^2} - \frac{e(a + b \operatorname{arccosh}(cx))}{d(d + ex^2)} \right) dx$$

$$\downarrow \text{2009}$$



$$\begin{aligned}
& \frac{\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2(-d)^{3/2}} - \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2(-d)^{3/2}} + \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2(-d)^{3/2}} - \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2(-d)^{3/2}} - \frac{a + \operatorname{barccosh}(cx)}{d} - \\
& \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2(-d)^{3/2}} + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2(-d)^{3/2}} - \\
& \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2(-d)^{3/2}} + \frac{b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2(-d)^{3/2}} + \\
& \frac{bc \arctan(\sqrt{cx - 1}\sqrt{cx + 1})}{d}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*(d + e*x^2)),x]`

output `-((a + b*ArcCosh[c*x])/(d*x)) + (b*c*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/d + (Sqrt[e]*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2)) - (Sqrt[e]*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2)) + (Sqrt[e]*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2)) - (Sqrt[e]*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2)) - (b*Sqrt[e]*PolyLog[2, -(Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2)) + (b*Sqrt[e]*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2)) - (b*Sqrt[e]*PolyLog[2, -(Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2)) + (b*Sqrt[e]*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*(-d)^(3/2))`

### 3.495.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

### 3.495.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 26.27 (sec) , antiderivative size = 331, normalized size of antiderivative = 0.61

method	result
parts	$-\frac{a}{dx} - \frac{ae \arctan\left(\frac{ex}{\sqrt{de}}\right)}{d\sqrt{de}} + bc \left( -\frac{\operatorname{arccosh}(cx)}{cxd} + \frac{2 \arctan(cx + \sqrt{cx-1} \sqrt{cx+1})}{d} + \frac{e \left( \sum_{-R1=\text{RootOf}(e-Z^4+(4c^2} \right)} \right)}{\dots} \right)$
derivativedivides	$c \left( -\frac{ae \arctan\left(\frac{ex}{\sqrt{de}}\right)}{cd\sqrt{de}} - \frac{a}{dcx} - \frac{b \operatorname{arccosh}(cx)}{cxd} + \frac{2b \arctan(cx + \sqrt{cx-1} \sqrt{cx+1})}{d} - \frac{be \left( \sum_{-R1=\text{RootOf}(e-Z^4+(4c^2} \right)} \right)}{\dots} \right)$
default	$c \left( -\frac{ae \arctan\left(\frac{ex}{\sqrt{de}}\right)}{cd\sqrt{de}} - \frac{a}{dcx} - \frac{b \operatorname{arccosh}(cx)}{cxd} + \frac{2b \arctan(cx + \sqrt{cx-1} \sqrt{cx+1})}{d} - \frac{be \left( \sum_{-R1=\text{RootOf}(e-Z^4+(4c^2} \right)} \right)}{\dots} \right)$

input `int((a+b*arccosh(c*x))/x^2/(e*x^2+d), x, method=_RETURNVERBOSE)`

3.495.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d+ex^2)} dx$

output `-a/d/x-a*e/d/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2))+b*c*(-arccosh(c*x)/c/x/d+2/d*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))+1/8/d^2*e/c^2*sum((_R1^2*e+4*c^2*d+e)/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))-1/8/d^2*e/c^2*sum((4*_R1^2*c^2*d+_R1^2*e+e)/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))`

### 3.495.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e*x^4 + d*x^2), x)`

### 3.495.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2(d + ex^2)} dx$$

input `integrate((a+b*acosh(c*x))/x**2/(e*x**2+d),x)`

output `Integral((a + b*acosh(c*x))/(x**2*(d + e*x**2)), x)`

**3.495.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2(d + ex^2)} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))/x^2/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.495.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)*x^2), x)`

**3.495.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2(e x^2 + d)} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d + e*x^2)),x)`

output `int((a + b*acosh(c*x))/(x^2*(d + e*x^2)), x)`

$$3.496 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d+ex^2)} dx$$

3.496.1 Optimal result . . . . .	3653
3.496.2 Mathematica [C] (verified) . . . . .	3654
3.496.3 Rubi [A] (verified) . . . . .	3655
3.496.4 Maple [C] (warning: unable to verify) . . . . .	3657
3.496.5 Fricas [F] . . . . .	3658
3.496.6 Sympy [F] . . . . .	3658
3.496.7 Maxima [F] . . . . .	3658
3.496.8 Giac [F] . . . . .	3659
3.496.9 Mupad [F(-1)] . . . . .	3659

**3.496.1 Optimal result**

Integrand size = 21, antiderivative size = 550

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d + ex^2)} dx = & \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{2dx} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} - \frac{e(a + \operatorname{barccosh}(cx))^2}{bd^2} \\
& - \frac{e(a + \operatorname{barccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)})}{d^2} \\
& + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^2} \\
& + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^2} \\
& + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^2} \\
& + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^2} \\
& + \frac{be \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d^2} \\
& + \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^2} \\
& + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^2} \\
& + \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^2} \\
& + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^2}
\end{aligned}$$

output

$$\begin{aligned} & \frac{1}{2}(-a-b\operatorname{arccosh}(cx))/d/x^2 - e*(a+b\operatorname{arccosh}(cx))^2/b/d^2 - e*(a+b\operatorname{arccosh}(cx)) \\ & * \ln(1+(cx+(cx-1)^{1/2})*(cx+1)^{1/2})^2/d^2 + 1/2*e*(a+b\operatorname{arccosh}(cx)) \\ & * \ln(1-(cx+(cx-1)^{1/2})*(cx+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}) \\ & )/d^2 + 1/2*e*(a+b\operatorname{arccosh}(cx))*\ln(1+(cx+(cx-1)^{1/2})*(cx+1)^{1/2})*e^{1/2} \\ & / (c*(-d)^{1/2}-(-c^2*d-e)^{1/2})/d^2 + 1/2*e*(a+b\operatorname{arccosh}(cx)) \\ & * \ln(1-(cx+(cx-1)^{1/2})*(cx+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}) \\ & )/d^2 + 1/2*e*(a+b\operatorname{arccosh}(cx))*\ln(1+(cx+(cx-1)^{1/2})*(cx+1)^{1/2})*e^{1/2} \\ & / (c*(-d)^{1/2}+(-c^2*d-e)^{1/2})/d^2 + 1/2*b*e*\operatorname{polylog}(2,-1/(cx+(cx-1)^{1/2})*(cx+1)^{1/2}) \\ & )^2/d^2 + 1/2*b*e*\operatorname{polylog}(2,-(cx+(cx-1)^{1/2})*(cx+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}) \\ & )/d^2 + 1/2*b*e*\operatorname{polylog}(2,(cx+(cx-1)^{1/2})*(cx+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}) \\ & )/d^2 + 1/2*b*e*\operatorname{polylog}(2,-(cx+(cx-1)^{1/2})*(cx+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}) \\ & )/d^2 + 1/2*b*c*(cx-1)^{1/2}*(cx+1)^{1/2}/d/x \end{aligned}$$

### 3.496.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 518, normalized size of antiderivative = 0.94

$$\begin{aligned} & \int \frac{a + b\operatorname{arccosh}(cx)}{x^3(d + ex^2)} dx \\ & = -\frac{a}{2dx^2} - \frac{ae \log(x)}{d^2} + \frac{ae \log(d + ex^2)}{2d^2} + b \left( \frac{cx\sqrt{-1 + cx}\sqrt{1 + cx} - \operatorname{arccosh}(cx)}{2dx^2} \right. \\ & \quad - \frac{e(\operatorname{arccosh}(cx)(\operatorname{arccosh}(cx) + 2\log(1 + e^{-2\operatorname{arccosh}(cx)})) - \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)}))}{2d^2} \\ & \quad + \frac{e(\operatorname{arccosh}(cx)(-\operatorname{arccosh}(cx) + 2(\log(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d-\sqrt{-c^2d-e}}}) + \log(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d+\sqrt{-c^2d-e}}}))}{4d^2} + 2\operatorname{PolyLog}(2, \frac{e^{-2\operatorname{arccosh}(cx)}}{1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d-\sqrt{-c^2d-e}}}}))}{4d^2} \\ & \quad \left. + \frac{e(\operatorname{arccosh}(cx)(-\operatorname{arccosh}(cx) + 2(\log(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-ic\sqrt{d+\sqrt{-c^2d-e}}}) + \log(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d+\sqrt{-c^2d-e}}}))}{4d^2} + 2\operatorname{PolyLog}(2, \frac{e^{-2\operatorname{arccosh}(cx)}}{1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d+\sqrt{-c^2d-e}}}}))}{4d^2} \right) \end{aligned}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d + e*x^2)),x]`

output

```

-1/2*a/(d*x^2) - (a*e*Log[x])/d^2 + (a*e*Log[d + e*x^2])/(2*d^2) + b*((c*x
*sqrt[-1 + c*x]*sqrt[1 + c*x] - ArcCosh[c*x])/(2*d*x^2) - (e*(ArcCosh[c*x]
*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLog[2, -E^(-2*ArcCo
sh[c*x])]))/(2*d^2) + (e*(ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(Log[1 + (sqrt[e
]*E^ArcCosh[c*x])/(I*c*sqrt[d] - sqrt[-(c^2*d) - e]))] + Log[1 + (sqrt[e]*E
^ArcCosh[c*x])/(I*c*sqrt[d] + sqrt[-(c^2*d) - e]))]) + 2*PolyLog[2, (sqrt[
e]*E^ArcCosh[c*x])/((-I)*c*sqrt[d] + sqrt[-(c^2*d) - e])] + 2*PolyLog[2, -
((sqrt[e]*E^ArcCosh[c*x])/(I*c*sqrt[d] + sqrt[-(c^2*d) - e]))])/(4*d^2) +
(e*(ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(Log[1 + (sqrt[e]*E^ArcCosh[c*x])/((-
I)*c*sqrt[d] + sqrt[-(c^2*d) - e]))] + Log[1 - (sqrt[e]*E^ArcCosh[c*x])/(I*
c*sqrt[d] + sqrt[-(c^2*d) - e]))]) + 2*PolyLog[2, -((sqrt[e]*E^ArcCosh[c*x
])/((-I)*c*sqrt[d] + sqrt[-(c^2*d) - e]))] + 2*PolyLog[2, (sqrt[e]*E^ArcCo
sh[c*x])/(I*c*sqrt[d] + sqrt[-(c^2*d) - e]))])/(4*d^2)

```

### 3.496.3 Rubi [A] (verified)

Time = 1.45 (sec) , antiderivative size = 550, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{arccosh}(cx)}{x^3(d + ex^2)} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{e^2x(a + \operatorname{arccosh}(cx))}{d^2(d + ex^2)} - \frac{e(a + \operatorname{arccosh}(cx))}{d^2x} + \frac{a + \operatorname{arccosh}(cx)}{dx^3} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$



$$\begin{aligned} & \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2d^2} + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2d^2} + \\ & \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2d^2} + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2d^2} - \\ & \frac{e(a + \operatorname{barccosh}(cx))^2}{bd^2} - \frac{e \log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx))}{d^2} - \frac{a + \operatorname{barccosh}(cx)}{2dx^2} + \\ & \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2d^2} + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2d^2} + \\ & \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2d^2} + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2d^2} + \\ & \frac{be \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d^2} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1}}{2dx} \end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*(d + e*x^2)),x]`

output `(b*c*sqrt[-1 + c*x]*sqrt[1 + c*x])/(2*d*x) - (a + b*ArcCosh[c*x])/(2*d*x^2) - (e*(a + b*ArcCosh[c*x])^2)/(b*d^2) - (e*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])])/d^2 + (e*(a + b*ArcCosh[c*x])*Log[1 - (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/d^2 + (e*(a + b*ArcCosh[c*x])*Log[1 + (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/d^2 + (e*(a + b*ArcCosh[c*x])*Log[1 - (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/d^2 + (e*(a + b*ArcCosh[c*x])*Log[1 + (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/d^2 + (b*e*PolyLog[2, -E^(-2*ArcCosh[c*x])])/d^2 + (b*e*PolyLog[2, -((sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e]))])/d^2 + (b*e*PolyLog[2, (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/d^2 + (b*e*PolyLog[2, -((sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e]))])/d^2 + (b*e*PolyLog[2, (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/d^2`

### 3.496.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

### 3.496.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.09 (sec) , antiderivative size = 466, normalized size of antiderivative = 0.85

method	result
parts	$a \left( \frac{e \ln(e x^2 + d)}{2d^2} - \frac{1}{2d x^2} - \frac{e \ln(x)}{d^2} \right) + b c^2 \left( -\frac{-\sqrt{cx-1} \sqrt{cx+1} cx + c^2 x^2 + \operatorname{arccosh}(cx)}{2c^2 x^2 d} + \frac{e^2}{-R1 = \operatorname{RootOf}(e^2 - \dots)} \right)$
derivativedivides	$c^2 \left( -\frac{a}{2d c^2 x^2} - \frac{ae \ln(cx)}{c^2 d^2} + \frac{ae \ln(c^2 e x^2 + c^2 d)}{2c^2 d^2} \right) + b c^2 \left( -\frac{-\sqrt{cx-1} \sqrt{cx+1} cx + c^2 x^2 + \operatorname{arccosh}(cx)}{2c^4 x^2 d} - \frac{e \operatorname{arccosh}(cx)}{c^2 d^2} \right)$
default	$c^2 \left( -\frac{a}{2d c^2 x^2} - \frac{ae \ln(cx)}{c^2 d^2} + \frac{ae \ln(c^2 e x^2 + c^2 d)}{2c^2 d^2} \right) + b c^2 \left( -\frac{-\sqrt{cx-1} \sqrt{cx+1} cx + c^2 x^2 + \operatorname{arccosh}(cx)}{2c^4 x^2 d} - \frac{e \operatorname{arccosh}(cx)}{c^2 d^2} \right)$

input `int((a+b*arccosh(c*x))/x^3/(e*x^2+d),x,method=_RETURNVERBOSE)`

output

```

a*(1/2*e/d^2*ln(e*x^2+d)-1/2/d/x^2-e/d^2*ln(x))+b*c^2*(-1/2*(-(c*x-1)^(1/2)
)*(c*x+1)^(1/2)*c*x+c^2*x^2+arccosh(c*x))/c^2/x^2/d+1/4*e^2/d^2/c^2*sum((_
R1^2+1)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)
)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf
(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))-e/d^2/c^2*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(
1/2)*(c*x+1)^(1/2)))-e/d^2/c^2*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*
x+1)^(1/2)))-e/d^2/c^2*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-e/d^2/
c^2*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/4*e/d^2/c^2*sum((_R1^2*
e+4*c^2*d+e)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(
c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=R
ootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))
    
```

**3.496.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e*x^5 + d*x^3), x)`

**3.496.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3(d + ex^2)} dx$$

input `integrate((a+b*acosh(c*x))/x**3/(e*x**2+d),x)`

output `Integral((a + b*acosh(c*x))/(x**3*(d + e*x**2)), x)`

**3.496.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d),x, algorithm="maxima")`

output `1/2*a*(e*log(e*x^2 + d)/d^2 - 2*e*log(x)/d^2 - 1/(d*x^2)) + b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(e*x^5 + d*x^3), x)`

**3.496.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)*x^3), x)`

**3.496.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3(ex^2 + d)} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d + e*x^2)),x)`

output `int((a + b*acosh(c*x))/(x^3*(d + e*x^2)), x)`

$$3.497 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{x^4(d+ex^2)} dx$$

3.497.1 Optimal result . . . . .	3661
3.497.2 Mathematica [A] (verified) . . . . .	3663
3.497.3 Rubi [A] (verified) . . . . .	3664
3.497.4 Maple [C] (warning: unable to verify) . . . . .	3666
3.497.5 Fricas [F] . . . . .	3667
3.497.6 Sympy [F] . . . . .	3667
3.497.7 Maxima [F(-2)] . . . . .	3668
3.497.8 Giac [F] . . . . .	3668
3.497.9 Mupad [F(-1)] . . . . .	3668

**3.497.1 Optimal result**

Integrand size = 21, antiderivative size = 624

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d + ex^2)} dx = & \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{6dx^2} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} \\
& + \frac{e(a + \operatorname{barccosh}(cx))}{d^2x} + \frac{bc^3 \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{6d} \\
& - \frac{bce \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d^2} \\
& + \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}} \\
& - \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}} \\
& + \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}} \\
& - \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}} \\
& - \frac{be^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}} \\
& + \frac{be^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}} \\
& - \frac{be^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}} \\
& + \frac{be^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{2(-d)^{5/2}}
\end{aligned}$$

output  $\frac{1}{3}(-a-b\operatorname{arccosh}(cx))/d/x^3+e(a+b\operatorname{arccosh}(cx))/d^2/x+1/6*b*c^3*\arctan((cx-1)^{1/2}*(cx+1)^{1/2})/d-b*c*e*\arctan((cx-1)^{1/2}*(cx+1)^{1/2})/d^2+1/2*e^{3/2}*(a+b\operatorname{arccosh}(cx))*\ln(1-(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/(-d)^{5/2}-1/2*e^{3/2}*(a+b\operatorname{arccosh}(cx))*\ln(1+(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/(-d)^{5/2}+1/2*e^{3/2}*(a+b\operatorname{arccosh}(cx))*\ln(1-(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/(-d)^{5/2}-1/2*e^{3/2}*(a+b\operatorname{arccosh}(cx))*\ln(1+(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/(-d)^{5/2}-1/2*b*e^{3/2}*polylog(2,-(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/(-d)^{5/2}+1/2*b*e^{3/2}*polylog(2,(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/(-d)^{5/2}+1/2*b*e^{3/2}*polylog(2,-(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/(-d)^{5/2}+1/2*b*e^{3/2}*polylog(2,(cx+(cx-1)^{1/2}*(cx+1)^{1/2}))*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/(-d)^{5/2}+1/6*b*c*(cx-1)^{1/2}*(cx+1)^{1/2}/d/x^2$

**3.497.2 Mathematica [A] (verified)**

Time = 1.20 (sec) , antiderivative size = 657, normalized size of antiderivative = 1.05

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d + ex^2)} dx = & \frac{1}{6} \left( -\frac{2a}{dx^3} + \frac{bc\sqrt{-1+cx}\sqrt{1+cx}}{dx^2} - \frac{2\operatorname{barccosh}(cx)}{dx^3} \right. \\
& + \frac{6e(a + \operatorname{barccosh}(cx))}{d^2x} + \frac{bc^3\sqrt{-1+c^2x^2} \arctan(\sqrt{-1+c^2x^2})}{d\sqrt{-1+cx}\sqrt{1+cx}} \\
& - \frac{6bce\sqrt{-1+c^2x^2} \arctan(\sqrt{-1+c^2x^2})}{d^2\sqrt{-1+cx}\sqrt{1+cx}} \\
& - \frac{3e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \\
& + \frac{3e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \\
& + \frac{3e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \\
& - \frac{3e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \\
& + \frac{3be^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \\
& - \frac{3be^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \\
& - \frac{3be^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \\
& \left. + \frac{3be^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{(-d)^{5/2}} \right)
\end{aligned}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^4*(d + e*x^2)),x]`



output  $((-2*a)/(d*x^3) + (b*c*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])/(d*x^2) - (2*b*\text{ArcCos}h[c*x])/(d*x^3) + (6*e*(a + b*\text{ArcCosh}[c*x]))/(d^2*x) + (b*c^3*\text{Sqrt}[-1 + c^2*x^2]*\text{ArcTan}[\text{Sqrt}[-1 + c^2*x^2]])/(d*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) - (6*b*c*e*\text{Sqrt}[-1 + c^2*x^2]*\text{ArcTan}[\text{Sqrt}[-1 + c^2*x^2]])/(d^2*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]) - (3*e^{(3/2)}*(a + b*\text{ArcCosh}[c*x])*Log[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (3*e^{(3/2)}*(a + b*\text{ArcCosh}[c*x])*Log[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(-(c*\text{Sqrt}[-d]) + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (3*e^{(3/2)}*(a + b*\text{ArcCosh}[c*x])*Log[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} - (3*e^{(3/2)}*(a + b*\text{ArcCosh}[c*x])*Log[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (3*b*e^{(3/2)}*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} - (3*b*e^{(3/2)}*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(-(c*\text{Sqrt}[-d]) + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} - (3*b*e^{(3/2)}*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2)} + (3*b*e^{(3/2)}*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(-d)^{(5/2}))/6$

### 3.497.3 Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 624, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \text{barccosh}(cx)}{x^4 (d + ex^2)} dx$$

$$\downarrow 6374$$

$$\int \left( \frac{e^2(a + \text{barccosh}(cx))}{d^2 (d + ex^2)} - \frac{e(a + \text{barccosh}(cx))}{d^2 x^2} + \frac{a + \text{barccosh}(cx)}{dx^4} \right) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2(-d)^{5/2}} - \\
& \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2(-d)^{5/2}} + \\
& \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2(-d)^{5/2}} - \\
& \frac{e^{3/2}(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2(-d)^{5/2}} + \frac{e(a + \operatorname{barccosh}(cx))}{d^2 x} - \frac{a + \operatorname{barccosh}(cx)}{3dx^3} - \\
& \frac{be^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2(-d)^{5/2}} + \frac{be^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2(-d)^{5/2}} - \\
& \frac{be^{3/2} \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2(-d)^{5/2}} + \frac{be^{3/2} \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2(-d)^{5/2}} + \\
& \frac{bc^3 \arctan(\sqrt{cx - 1}\sqrt{cx + 1})}{6d} - \frac{bce \arctan(\sqrt{cx - 1}\sqrt{cx + 1})}{d^2} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1}}{6dx^2}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^4*(d + e*x^2)),x]`

output `(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(6*d*x^2) - (a + b*ArcCosh[c*x])/(3*d*x^3) + (e*(a + b*ArcCosh[c*x]))/(d^2*x) + (b*c^3*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/(6*d) - (b*c*e*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/d^2 + (e^(3/2)*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*(-d)^(5/2)) - (e^(3/2)*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*(-d)^(5/2)) + (e^(3/2)*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*(-d)^(5/2)) - (e^(3/2)*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*(-d)^(5/2)) - (b*e^(3/2)*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])]/(2*(-d)^(5/2)) + (b*e^(3/2)*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*(-d)^(5/2)) - (b*e^(3/2)*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])]/(2*(-d)^(5/2)) + (b*e^(3/2)*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*(-d)^(5/2))`

3.497.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

3.497.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 32.84 (sec) , antiderivative size = 424, normalized size of antiderivative = 0.68

method	result
parts	$a \left( \frac{e^2 \arctan\left(\frac{ex}{\sqrt{de}}\right)}{d^2 \sqrt{de}} - \frac{1}{3d x^3} + \frac{e}{d^2 x} \right) + \frac{b \left( 8 \arctan(cx + \sqrt{cx-1} \sqrt{cx+1}) c^7 d^2 x^3 + 4 \sqrt{cx+1} \sqrt{cx-1} c^5 d^2 x - 48 \arctan\left(\frac{ex}{\sqrt{de}}\right) c^3 d^2 x^3 - 4 \sqrt{cx+1} \sqrt{cx-1} c^5 d^2 x \right)}{c^3 d^2 \sqrt{de}}$
derivativedivides	$c^3 \left( -\frac{a}{3d c^3 x^3} + \frac{ae}{c^3 d^2 x} + \frac{a e^2 \arctan\left(\frac{ex}{\sqrt{de}}\right)}{c^3 d^2 \sqrt{de}} - \frac{b \left( -8 \arctan(cx + \sqrt{cx-1} \sqrt{cx+1}) c^7 d^2 x^3 - 4 \sqrt{cx+1} \sqrt{cx-1} c^5 d^2 x \right)}{c^3 d^2 \sqrt{de}} \right)$
default	$c^3 \left( -\frac{a}{3d c^3 x^3} + \frac{ae}{c^3 d^2 x} + \frac{a e^2 \arctan\left(\frac{ex}{\sqrt{de}}\right)}{c^3 d^2 \sqrt{de}} - \frac{b \left( -8 \arctan(cx + \sqrt{cx-1} \sqrt{cx+1}) c^7 d^2 x^3 - 4 \sqrt{cx+1} \sqrt{cx-1} c^5 d^2 x \right)}{c^3 d^2 \sqrt{de}} \right)$

input `int((a+b*arccosh(c*x))/x^4/(e*x^2+d),x,method=_RETURNVERBOSE)`

output `a*(e^2/d^2/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2))-1/3/d/x^3+e/d^2/x)+1/24*b/c^4*(8*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*c^7*d^2*x^3+4*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^5*d^2*x-48*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*c^5*d*e*x^3+24*arccosh(c*x)*c^4*d*e*x^2-8*c^4*d^2*arccosh(c*x)+3*sum((4*_R1^2*c^2*d+_R1^2*e+e)/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))*e^2*c^3*x^3-3*sum((*_R1^2*e+4*c^2*d+e)/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))*e^2*c^3*x^3)/x^3/d^3`

### 3.497.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(e*x^2+d),x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e*x^6 + d*x^4), x)`

### 3.497.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^4(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4(d + ex^2)} dx$$

input `integrate((a+b*acosh(c*x))/x**4/(e*x**2+d),x)`

output `Integral((a + b*acosh(c*x))/(x**4*(d + e*x**2)), x)`

**3.497.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d + ex^2)} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))/x^4/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.497.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d + ex^2)} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)x^4} dx$$

input `integrate((a+b*arccosh(c*x))/x^4/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)*x^4), x)`

**3.497.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^4(d + ex^2)} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^4(e x^2 + d)} dx$$

input `int((a + b*acosh(c*x))/(x^4*(d + e*x^2)),x)`

output `int((a + b*acosh(c*x))/(x^4*(d + e*x^2)), x)`

$$3.498 \quad \int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx$$

3.498.1 Optimal result . . . . .	3669
3.498.2 Mathematica [C] (verified) . . . . .	3670
3.498.3 Rubi [A] (verified) . . . . .	3671
3.498.4 Maple [C] (warning: unable to verify) . . . . .	3673
3.498.5 Fricas [F] . . . . .	3674
3.498.6 Sympy [F] . . . . .	3674
3.498.7 Maxima [F] . . . . .	3674
3.498.8 Giac [F(-2)] . . . . .	3675
3.498.9 Mupad [F(-1)] . . . . .	3675

### 3.498.1 Optimal result

Integrand size = 21, antiderivative size = 562

$$\begin{aligned} \int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx = & \frac{d(a+b\operatorname{arccosh}(cx))}{2e^2(d+ex^2)} - \frac{(a+b\operatorname{arccosh}(cx))^2}{2be^2} \\ & - \frac{bc\sqrt{d}\sqrt{-1+c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{2e^2\sqrt{c^2d+e}\sqrt{-1+cx}\sqrt{1+cx}} \\ & + \frac{(a+b\operatorname{arccosh}(cx))\log\left(1-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^2} \\ & + \frac{(a+b\operatorname{arccosh}(cx))\log\left(1+\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^2} \\ & + \frac{(a+b\operatorname{arccosh}(cx))\log\left(1-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^2} \\ & + \frac{(a+b\operatorname{arccosh}(cx))\log\left(1+\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^2} \\ & + \frac{b\operatorname{PolyLog}\left(2,-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^2} \\ & + \frac{b\operatorname{PolyLog}\left(2,\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^2} \\ & + \frac{b\operatorname{PolyLog}\left(2,-\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^2} \\ & + \frac{b\operatorname{PolyLog}\left(2,\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^2} \end{aligned}$$

---

3.498.  $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx$

output  $\frac{1}{2}d*(a+b*\operatorname{arccosh}(c*x))/e^2/(e*x^2+d)-1/2*(a+b*\operatorname{arccosh}(c*x))^2/b/e^2+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2+1/2*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2+1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2+1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}-(-c^2*d-e)^{1/2}))/e^2+1/2*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2+1/2*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{1/2})*(c*x+1)^{1/2})*e^{1/2}/(c*(-d)^{1/2}+(-c^2*d-e)^{1/2}))/e^2-1/2*b*c*\operatorname{arctanh}(x*(c^2*d+e)^{1/2}/d^{1/2}/(c^2*x^2-1)^{1/2})*d^{1/2}*(c^2*x^2-1)^{1/2}/e^2/(c^2*d+e)^{1/2}/(c*x-1)^{1/2}/(c*x+1)^{1/2}$

### 3.498.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.38 (sec) , antiderivative size = 694, normalized size of antiderivative = 1.23

$$\int \frac{x^3(a + b\operatorname{arccosh}(cx))}{(d + ex^2)^2} dx$$

$$\frac{2ad}{d+ex^2} + 2a \log(d + ex^2) + b \left( -2\operatorname{arccosh}(cx)^2 + 2\operatorname{arccosh}(cx) \left( \log \left( 1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-ic\sqrt{d} + \sqrt{-c^2d-e}} \right) + \log \left( 1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} + \sqrt{-c^2d-e}} \right) \right) \right)$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output  $((2*a*d)/(d + e*x^2) + 2*a*Log[d + e*x^2] + b*(-2*ArcCosh[c*x]^2 + 2*ArcCosh[c*x]*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/((-I)*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) + Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) + 2*ArcCosh[c*x]*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] - Sqrt[-(c^2*d) - e])]) + Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) - I*Sqrt[d]*(ArcCosh[c*x])/((-I)*Sqrt[d] + Sqrt[e]*x) + (c*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])]/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*Sqrt[e]*x)))/Sqrt[-(c^2*d) - e] - I*Sqrt[d]*(-(ArcCosh[c*x])/(I*Sqrt[d] + Sqrt[e]*x)) - (c*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])]/(c*Sqrt[-(c^2*d) - e]*(I*Sqrt[d] + Sqrt[e]*x)))/Sqrt[-(c^2*d) - e]) + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] - Sqrt[-(c^2*d) - e])] + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/((-I)*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] + 2*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])])]/(4*e^2)$

### 3.498.3 Rubi [A] (verified)

Time = 1.39 (sec) , antiderivative size = 562, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \text{barccosh}(cx))}{(d + ex^2)^2} dx$$

↓ 6374

$$\int \left( \frac{x(a + \text{barccosh}(cx))}{e(d + ex^2)} - \frac{dx(a + \text{barccosh}(cx))}{e(d + ex^2)^2} \right) dx$$

↓ 2009



$$\begin{aligned}
& \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2e^2} + \frac{(a + \operatorname{arccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2e^2} + \\
& \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2e^2} + \frac{(a + \operatorname{arccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2e^2} + \\
& \frac{d(a + \operatorname{arccosh}(cx))}{2e^2(d + ex^2)} - \frac{(a + \operatorname{arccosh}(cx))^2}{2be^2} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2e^2} + \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2e^2} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2e^2} + \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2e^2} - \frac{bc\sqrt{d}\sqrt{c^2x^2 - 1} \operatorname{arctanh}\left(\frac{x\sqrt{c^2d + e}}{\sqrt{d}\sqrt{c^2x^2 - 1}}\right)}{2e^2\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{c^2d + e}}
\end{aligned}$$

input `Int[(x^3*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output `(d*(a + b*ArcCosh[c*x]))/(2*e^2*(d + e*x^2)) - (a + b*ArcCosh[c*x])^2/(2*b*e^2) - (b*c*Sqrt[d]*Sqrt[-1 + c^2*x^2]*ArcTanh[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2])])/(2*e^2*Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^2) + ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^2) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^2) + ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^2) + (b*PolyLog[2, -(Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^2) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^2) + (b*PolyLog[2, -(Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^2) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^2)`

### 3.498.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n_.*((f_.)*(x_)^m_.)*((d_.) + (e_.)*(x_)^2)^p_.], x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

**3.498.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 3.01 (sec) , antiderivative size = 2132, normalized size of antiderivative = 3.79

method	result	size
derivativdivides	Expression too large to display	2132
default	Expression too large to display	2132
parts	Expression too large to display	2144

input `int(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

$$\begin{aligned} & 1/c^4*(1/2*a*c^6/e^2*d/(c^2*e*x^2+c^2*d)+1/2*a*c^4/e^2*\ln(c^2*e*x^2+c^2*d) \\ & +b*c^4*(-1/8*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2* \\ & (c^2*d+e))^(1/2)*e)/c^2/d/e^2/(c^2*d+e)*polylog(2,e*(c*x+(c*x-1)^(1/2)*(c* \\ & x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))+(-2*(d*c^2*(c^2*d+e) \\ & )^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^2*d/e^4/(c^ \\ & 2*d+e)*arccosh(c*x)^2-1/2*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^ \\ & 2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^2*d/e^4/(c^2*d+e)*polylog(2,e*(c*x+(c*x \\ & -1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))+1/4*(d* \\ & c^2*(c^2*d+e))^(1/2)/e/d/c^2/(c^2*d+e)*arccosh(c*x)^2+1/4*(-2*(d*c^2*(c^2* \\ & d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)/c^2/d/e^2 \\ & / (c^2*d+e)*arccosh(c*x)^2+(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)/e^4*\ln(1-e \\ & *(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e \\ & ))*c^2*d*arccosh(c*x)+1/2*(d*c^2*(c^2*d+e))^(1/2)/e^2/(c^2*d+e)*arctanh(1/ \\ & 4*(4*c^2*d+2*e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2+2*e)/(c^4*d^2+c^2*d*e)^( \\ & 1/2))-1/4*(d*c^2*(c^2*d+e))^(1/2)/e^2/(c^2*d+e)*polylog(2,e*(c*x+(c*x-1)^( \\ & 1/2)*(c*x+1)^(1/2))^2/(-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e))-1/8*(d*c^2* \\ & (c^2*d+e))^(1/2)/e/d/c^2/(c^2*d+e)*polylog(2,e*(c*x+(c*x-1)^(1/2)*(c*x+1)^( \\ & 1/2))^2/(-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e))+(-2*(d*c^2*(c^2*d+e))^(1/ \\ & 2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)/e^3/(c^2*d+e)*arcc \\ & osh(c*x)^2+1/2*(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)/e^3*\ln(1-e*(c*x+(c... \end{aligned}$$

**3.498.5 Fracas [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^3}{(ex^2 + d)^2} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*x^3*arccosh(c*x) + a*x^3)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**3.498.6 Sympy [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(d + ex^2)^2} dx$$

input `integrate(x**3*(a+b*acosh(c*x))/(e*x**2+d)**2,x)`

output `Integral(x**3*(a + b*acosh(c*x))/(d + e*x**2)**2, x)`

**3.498.7 Maxima [F]**

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^3}{(ex^2 + d)^2} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `1/2*a*(d/(e^3*x^2 + d*e^2) + log(e*x^2 + d)/e^2) + b*integrate(x^3*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**3.498.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.498.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^2} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d + e*x^2)^2,x)`

output `int((x^3*(a + b*acosh(c*x)))/(d + e*x^2)^2, x)`

$$3.499 \quad \int \frac{x(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx$$

3.499.1 Optimal result . . . . .	3676
3.499.2 Mathematica [A] (verified) . . . . .	3676
3.499.3 Rubi [A] (verified) . . . . .	3677
3.499.4 Maple [B] (verified) . . . . .	3678
3.499.5 Fricas [B] (verification not implemented) . . . . .	3679
3.499.6 Sympy [F] . . . . .	3680
3.499.7 Maxima [F(-2)] . . . . .	3681
3.499.8 Giac [F] . . . . .	3681
3.499.9 Mupad [F(-1)] . . . . .	3681

### 3.499.1 Optimal result

Integrand size = 19, antiderivative size = 113

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx = -\frac{a + \operatorname{arccosh}(cx)}{2e(d + ex^2)} + \frac{bc\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{2\sqrt{de}\sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output  $1/2*(-a-b*\operatorname{arccosh}(c*x))/e/(e*x^2+d)+1/2*b*c*\operatorname{arctanh}(x*(c^2*d+e)^{(1/2)}/d^{(1/2)}/(c^2*x^2-1)^{(1/2)})*(c^2*x^2-1)^{(1/2)}/e/d^{(1/2)}/(c^2*d+e)^{(1/2)}/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.499.2 Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.09

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx = -\frac{a}{d+ex^2} + \frac{\operatorname{arccosh}(cx)}{d+ex^2} - \frac{bc\sqrt{-1+cx}\sqrt{1+cx}\operatorname{arctan}\left(\frac{\sqrt{-c^2d-ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{2e}$$

input `Integrate[(x*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output  $-1/2*(a/(d + e*x^2) + (b*\operatorname{ArcCosh}[c*x])/(d + e*x^2) - (b*c*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*\operatorname{ArcTan}[(\operatorname{Sqrt}[-(c^2*d) - e]*x)/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[-1 + c^2*x^2])])/( \operatorname{Sqrt}[d]*\operatorname{Sqrt}[-(c^2*d) - e]*\operatorname{Sqrt}[-1 + c^2*x^2]))/e$

---

3.499.  $\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx$

**3.499.3 Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$ , Rules used = {6372, 648, 291, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx$$

↓ 6372

$$\frac{bc \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}(ex^2+d)} dx}{2e} - \frac{a + \operatorname{barccosh}(cx)}{2e(d + ex^2)}$$

↓ 648

$$\frac{bc\sqrt{c^2x^2-1} \int \frac{1}{\sqrt{c^2x^2-1}(ex^2+d)} dx}{2e\sqrt{cx-1}\sqrt{cx+1}} - \frac{a + \operatorname{barccosh}(cx)}{2e(d + ex^2)}$$

↓ 291

$$\frac{bc\sqrt{c^2x^2-1} \int \frac{1}{d - \frac{(dc^2+e)x^2}{c^2x^2-1}} d \frac{x}{\sqrt{c^2x^2-1}}}{2e\sqrt{cx-1}\sqrt{cx+1}} - \frac{a + \operatorname{barccosh}(cx)}{2e(d + ex^2)}$$

↓ 221

$$\frac{bc\sqrt{c^2x^2-1} \operatorname{arctanh}\left(\frac{x\sqrt{c^2d+e}}{\sqrt{d}\sqrt{c^2x^2-1}}\right)}{2\sqrt{de}\sqrt{cx-1}\sqrt{cx+1}\sqrt{c^2d+e}} - \frac{a + \operatorname{barccosh}(cx)}{2e(d + ex^2)}$$

input `Int[(x*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output `-1/2*(a + b*ArcCosh[c*x])/(e*(d + e*x^2)) + (b*c*Sqrt[-1 + c^2*x^2]*ArcTan h[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2])])/(2*Sqrt[d]*e*Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

**3.499.3.1 Defintions of rubi rules used**

- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 291 `Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*((c_) + (d_)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 648 `Int[((c_) + (d_)*(x_))^(m_)*((e_) + (f_)*(x_))^(n_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c + d*x)^FracPart[m]*((e + f*x)^FracPart[m]/(c*e + d*f*x^2)^FracPart[m]) Int[(c*e + d*f*x^2)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m, n] && EqQ[d*e + c*f, 0] && !(EqQ[p, 2] && LtQ[m, -1])`
- rule 6372 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*(x_)*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])/(2*e*(p + 1))), x] - Simp[b*(c/(2*e*(p + 1))) Int[(d + e*x^2)^(p + 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[c^2*d + e, 0] && NeQ[p, -1]`

**3.499.4 Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 398 vs.  $2(96) = 192$ .

Time = 7.56 (sec) , antiderivative size = 399, normalized size of antiderivative = 3.53

method	result
parts	$-\frac{a}{2e(e x^2+d)} + b \left( -\frac{c^4 \operatorname{arccosh}(cx)}{2e(c^2 e x^2+c^2 d)} - \frac{c^4 \sqrt{cx-1} \sqrt{cx+1}}{e c x-\sqrt{-c^2 d e}} \ln \left( \frac{2\sqrt{-\frac{c^2 d+e}{e}} \sqrt{c^2 x^2-1} e+2\sqrt{-c^2 d e} c x-2 e}{e c x-\sqrt{-c^2 d e}} \right) c^2 d-\ln \left( \frac{2\sqrt{-\frac{c^2 d+e}{e}} \sqrt{c^2 x^2-1} e+2\sqrt{-c^2 d e} c x-2 e}{e c x-\sqrt{-c^2 d e}} \right) \right)$
derivativedivides	$-\frac{a c^4}{2e(c^2 e x^2+c^2 d)} + b c^4 \left( -\frac{\operatorname{arccosh}(cx)}{2e(c^2 e x^2+c^2 d)} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{e c x-\sqrt{-c^2 d e}} \ln \left( \frac{2\sqrt{-\frac{c^2 d+e}{e}} \sqrt{c^2 x^2-1} e+2\sqrt{-c^2 d e} c x-2 e}{e c x-\sqrt{-c^2 d e}} \right) c^2 d-\ln \left( \frac{2\sqrt{-\frac{c^2 d+e}{e}} \sqrt{c^2 x^2-1} e+2\sqrt{-c^2 d e} c x-2 e}{e c x-\sqrt{-c^2 d e}} \right) \right)$
default	$-\frac{a c^4}{2e(c^2 e x^2+c^2 d)} + b c^4 \left( -\frac{\operatorname{arccosh}(cx)}{2e(c^2 e x^2+c^2 d)} - \frac{\sqrt{cx-1} \sqrt{cx+1}}{e c x-\sqrt{-c^2 d e}} \ln \left( \frac{2\sqrt{-\frac{c^2 d+e}{e}} \sqrt{c^2 x^2-1} e+2\sqrt{-c^2 d e} c x-2 e}{e c x-\sqrt{-c^2 d e}} \right) c^2 d-\ln \left( \frac{2\sqrt{-\frac{c^2 d+e}{e}} \sqrt{c^2 x^2-1} e+2\sqrt{-c^2 d e} c x-2 e}{e c x-\sqrt{-c^2 d e}} \right) \right)$

input `int(x*(a+b*arccosh(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output 
$$-1/2*a/e/(e*x^2+d)+b/c^2*(-1/2*c^4/e/(c^2*e*x^2+c^2*d)*\operatorname{arccosh}(c*x)-1/4*c^4*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}*(\ln(2*((-c^2*d+e)/e)^{(1/2)}*(c^2*x^2-1)^{(1/2)})*e+(-c^2*d*e)^{(1/2)}*c*x-e)/(e*c*x-(-c^2*d*e)^{(1/2)}))*c^2*d-\ln(-2*((-c^2*d+e)/e)^{(1/2)}*(c^2*x^2-1)^{(1/2)}*e+(-c^2*d*e)^{(1/2)}*c*x+e)/(e*c*x+(-c^2*d*e)^{(1/2)}))*c^2*d+\ln(2*((-c^2*d+e)/e)^{(1/2)}*(c^2*x^2-1)^{(1/2)}*e+(-c^2*d*e)^{(1/2)}*c*x-e)/(e*c*x-(-c^2*d*e)^{(1/2)}))*e-\ln(-2*((-c^2*d+e)/e)^{(1/2)}*(c^2*x^2-1)^{(1/2)}*e+(-c^2*d*e)^{(1/2)}*c*x+e)/(e*c*x+(-c^2*d*e)^{(1/2)}))*e)/(c^2*x^2-1)^{(1/2)}/((-c^2*d*e)^{(1/2)}+e)/(-(-c^2*d*e)^{(1/2)}+e)/(-c^2*d*e)^{(1/2)}/(-c^2*d+e)/e)^{(1/2)}$$

### 3.499.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs.  $2(93) = 186$ .

3.499. 
$$\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx$$



Time = 0.28 (sec) , antiderivative size = 537, normalized size of antiderivative = 4.75

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx$$

$$= \left[ \frac{2ac^2d^2 - 2(bc^2de + be^2)x^2 \log(cx + \sqrt{c^2x^2 - 1}) + 2ade - (bcex^2 + bcd)\sqrt{c^2d^2 + de} \log\left(-\frac{2c^2d^2 - (4c^4d^2 + 4c^2d^2e + e^2)x^2 + d^2e - 2\sqrt{c^2d^2 + de}}{2(c^2d^3e + d^2e^2 + (c^2d^2e^2 + d^2e^3)x^2)}\right)}{2(c^2d^3e + d^2e^2 + (c^2d^2e^2 + d^2e^3)x^2)} \right]$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `[-1/4*(2*a*c^2*d^2 - 2*(b*c^2*d*e + b*e^2)*x^2*log(c*x + sqrt(c^2*x^2 - 1)) + 2*a*d*e - (b*c*e*x^2 + b*c*d)*sqrt(c^2*d^2 + d*e)*log(-(2*c^2*d^2 - (4*c^4*d^2 + 4*c^2*d^2*e + e^2)*x^2 + d^2e - 2*sqrt(c^2*d^2 + d*e))*((2*c^3*d + c*e)*x^2 - c*d) - 2*sqrt(c^2*x^2 - 1)*(sqrt(c^2*d^2 + d*e)*(2*c^2*d + e)*x + 2*(c^3*d^2 + c*d*e)*x))/(e*x^2 + d) - 2*(b*c^2*d^2 + b*d*e + (b*c^2*d*e + b*e^2)*x^2)*log(-c*x + sqrt(c^2*x^2 - 1))/(c^2*d^3*e + d^2*e^2 + (c^2*d^2*e^2 + d*e^3)*x^2), -1/2*(a*c^2*d^2 - (b*c^2*d*e + b*e^2)*x^2*log(c*x + sqrt(c^2*x^2 - 1)) + a*d*e - (b*c*e*x^2 + b*c*d)*sqrt(-c^2*d^2 - d*e)*arctan((sqrt(-c^2*d^2 - d*e)*sqrt(c^2*x^2 - 1)*e*x - sqrt(-c^2*d^2 - d*e)*(c*e*x^2 + c*d))/(c^2*d^2 + d*e)) - (b*c^2*d^2 + b*d*e + (b*c^2*d*e + b*e^2)*x^2)*log(-c*x + sqrt(c^2*x^2 - 1))/(c^2*d^3*e + d^2*e^2 + (c^2*d^2*e^2 + d*e^3)*x^2)]`

### 3.499.6 Sympy [F]

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(d + ex^2)^2} dx$$

input `integrate(x*(a+b*acosh(c*x))/(e*x**2+d)**2,x)`

output `Integral(x*(a + b*acosh(c*x))/(d + e*x**2)**2, x)`

---

3.499.  $\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx$

**3.499.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.499.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(ex^2 + d)^2} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x/(e*x^2 + d)^2, x)`

**3.499.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^2} dx$$

input `int((x*(a + b*acosh(c*x)))/(d + e*x^2)^2,x)`

output `int((x*(a + b*acosh(c*x)))/(d + e*x^2)^2, x)`

# 3.500 $\int \frac{a+b\operatorname{arccosh}(cx)}{x(d+ex^2)^2} dx$

3.500.1 Optimal result . . . . .	3682
3.500.2 Mathematica [C] (verified) . . . . .	3683
3.500.3 Rubi [A] (verified) . . . . .	3684
3.500.4 Maple [C] (warning: unable to verify) . . . . .	3686
3.500.5 Fricas [F] . . . . .	3687
3.500.6 Sympy [F] . . . . .	3687
3.500.7 Maxima [F] . . . . .	3688
3.500.8 Giac [F] . . . . .	3688
3.500.9 Mupad [F(-1)] . . . . .	3688

## 3.500.1 Optimal result

Integrand size = 21, antiderivative size = 598

$$\int \frac{a + \operatorname{arccosh}(cx)}{x(d + ex^2)^2} dx = \frac{a + \operatorname{arccosh}(cx)}{2d(d + ex^2)} + \frac{(a + \operatorname{arccosh}(cx))^2}{bd^2}$$

$$- \frac{bc\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{2d^{3/2}\sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

$$+ \frac{(a + \operatorname{arccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)})}{d^2}$$

$$- \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^2}$$

$$- \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^2}$$

$$- \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^2}$$

$$- \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 + \frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^2}$$

$$- \frac{b \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d^2} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^2}$$

$$- \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^2} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^2}$$

$$- \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}\operatorname{arccosh}(cx)}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^2}$$

output

```

1/2*(a+b*arccosh(c*x))/d/(e*x^2+d)+(a+b*arccosh(c*x))^2/b/d^2+(a+b*arccosh
(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d^2-1/2*(a+b*arccosh(c*
x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e
)^(1/2)))/d^2-1/2*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)
)*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^2-1/2*(a+b*arccosh(c*x))*ln(1
-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)
))/d^2-1/2*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2
))/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^2-1/2*b*polylog(2,-1/(c*x+(c*x-1)^(1/
2)*(c*x+1)^(1/2))^2)/d^2-1/2*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)
)*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^2-1/2*b*polylog(2,(c*x+(c*x-1
)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^2-1/2*b*
polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d
-e)^(1/2)))/d^2-1/2*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/
(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^2-1/2*b*c*arctanh(x*(c^2*d+e)^(1/2)/d^(
1/2)/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/d^(3/2)/(c^2*d+e)^(1/2)/(c*x-1)^(
1/2)/(c*x+1)^(1/2)

```

### 3.500.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.16 (sec) , antiderivative size = 754, normalized size of antiderivative = 1.26

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^2} dx = \frac{a}{2d^2 + 2dex^2} + \frac{a \log(x)}{d^2} - \frac{a \log(d + ex^2)}{2d^2} + b \left( -\frac{\sqrt{d} \operatorname{arccosh}(cx)}{\sqrt{d-i}\sqrt{ex}} - \frac{\sqrt{d} \operatorname{arccosh}(cx)}{\sqrt{d+i}\sqrt{ex}} - 4 \operatorname{arccosh}(cx)^2 - 4 \operatorname{arccosh}(cx) \log(1 + e^{-2 \operatorname{arccosh}(cx)}) + 2 \operatorname{arccosh}(cx) \right)$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*(d + e*x^2)^2), x]`

output

```

a/(2*d^2 + 2*d*e*x^2) + (a*Log[x])/d^2 - (a*Log[d + e*x^2])/(2*d^2) - (b*(
-((Sqrt[d]*ArcCosh[c*x])/(Sqrt[d] - I*Sqrt[e]*x)) - (Sqrt[d]*ArcCosh[c*x])
/(Sqrt[d] + I*Sqrt[e]*x) - 4*ArcCosh[c*x]^2 - 4*ArcCosh[c*x]*Log[1 + E^(-2
*ArcCosh[c*x])) + 2*ArcCosh[c*x]*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqr
t[d] - Sqrt[-(c^2*d) - e])] + 2*ArcCosh[c*x]*Log[1 + (Sqrt[e]*E^ArcCosh[c*
x])/((-I)*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] + 2*ArcCosh[c*x]*Log[1 - (Sqrt[
e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] + 2*ArcCosh[c*x]*Lo
g[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] + (I*c*
Sqrt[d]*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*d) - e]*Sqrt[-1
+ c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*Sqrt[e]*x))])/S
qrt[-(c^2*d) - e] - (I*c*Sqrt[d]*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x + Sq
rt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) - e]*(I*S
qrt[d] + Sqrt[e]*x))])/Sqrt[-(c^2*d) - e] + 2*PolyLog[2, -E^(-2*ArcCosh[c*
x])] + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] - Sqrt[-(c^2*d)
- e])] + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/((-I)*c*Sqrt[d] + Sqrt[-(c^
2*d) - e])] + 2*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[
-(c^2*d) - e]))] + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sq
rt[-(c^2*d) - e]))])/(4*d^2)

```

### 3.500.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 598, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)^2} dx$$

$$\downarrow \text{6374}$$

$$\int \left( -\frac{ex(a + \operatorname{barccosh}(cx))}{d^2(d + ex^2)} + \frac{a + \operatorname{barccosh}(cx)}{d^2x} - \frac{ex(a + \operatorname{barccosh}(cx))}{d(d + ex^2)^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2d^2} - \frac{(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2d^2} \\
& \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2d^2} - \frac{(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2d^2} + \\
& \frac{(a + \operatorname{barccosh}(cx))^2}{bd^2} + \frac{\log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx))}{d^2} + \frac{a + \operatorname{barccosh}(cx)}{2d(d + ex^2)} - \\
& \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2d^2} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2d^2} - \\
& \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2d^2} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2d^2} - \\
& \frac{b \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d^2} - \frac{bc\sqrt{c^2x^2 - 1} \operatorname{arctanh}\left(\frac{x\sqrt{c^2d + e}}{\sqrt{d}\sqrt{c^2x^2 - 1}}\right)}{2d^{3/2}\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{c^2d + e}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d + e*x^2)^2), x]`

output `(a + b*ArcCosh[c*x])/(2*d*(d + e*x^2)) + (a + b*ArcCosh[c*x])^2/(b*d^2) - (b*c*sqrt[-1 + c^2*x^2]*ArcTanh[(sqrt[c^2*d + e]*x)/(sqrt[d]*sqrt[-1 + c^2*x^2])])/(2*d^(3/2)*sqrt[c^2*d + e]*sqrt[-1 + c*x]*sqrt[1 + c*x]) + ((a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])])/d^2 - ((a + b*ArcCosh[c*x])*Log[1 - (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/(2*d^2) - ((a + b*ArcCosh[c*x])*Log[1 + (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/(2*d^2) - ((a + b*ArcCosh[c*x])*Log[1 - (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(2*d^2) - ((a + b*ArcCosh[c*x])*Log[1 + (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(2*d^2) - (b*PolyLog[2, -E^(-2*ArcCosh[c*x])])/(2*d^2) - (b*PolyLog[2, -((sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/(2*d^2) - (b*PolyLog[2, (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/(2*d^2) - (b*PolyLog[2, -((sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(2*d^2) - (b*PolyLog[2, (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(2*d^2) - (b*PolyLog[2, (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(2*d^2)`

### 3.500.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

### 3.500.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.49 (sec) , antiderivative size = 505, normalized size of antiderivative = 0.84

method	result
parts	$-\frac{a \ln(e x^2+d)}{2d^2} + \frac{a}{2d(e x^2+d)} + \frac{a \ln(x)}{d^2} + b \left( \frac{c^2 \operatorname{arccosh}(cx)}{2d(c^2 e x^2+c^2 d)} + \frac{\sqrt{d c^2(c^2 d+e)} \operatorname{arctanh}\left(\frac{4c^2 d+2e(cx+\sqrt{cx-1}\sqrt{cx-1})}{4\sqrt{c^4 d^2+c^2 d e}}\right)}{2d^2(c^2 d+e)} \right)$
derivativedivides	$\frac{a \ln(cx)}{d^2} + \frac{a c^2}{2d(c^2 e x^2+c^2 d)} - \frac{a \ln(c^2 e x^2+c^2 d)}{2d^2} + \frac{b c^2 \operatorname{arccosh}(cx)}{2d(c^2 e x^2+c^2 d)} + \frac{b \sqrt{d c^2(c^2 d+e)} \operatorname{arctanh}\left(\frac{4c^2 d+2e(cx+\sqrt{cx-1}\sqrt{cx-1})}{4\sqrt{c^4 d^2+c^2 d e}}\right)}{2d^2(c^2 d+e)}$
default	$\frac{a \ln(cx)}{d^2} + \frac{a c^2}{2d(c^2 e x^2+c^2 d)} - \frac{a \ln(c^2 e x^2+c^2 d)}{2d^2} + \frac{b c^2 \operatorname{arccosh}(cx)}{2d(c^2 e x^2+c^2 d)} + \frac{b \sqrt{d c^2(c^2 d+e)} \operatorname{arctanh}\left(\frac{4c^2 d+2e(cx+\sqrt{cx-1}\sqrt{cx-1})}{4\sqrt{c^4 d^2+c^2 d e}}\right)}{2d^2(c^2 d+e)}$

input `int((a+b*arccosh(c*x))/x/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output `-1/2*a/d^2*ln(e*x^2+d)+1/2*a/d/(e*x^2+d)+a/d^2*ln(x)+b*(1/2*c^2*arccosh(c*x)/d/(c^2*e*x^2+c^2*d)+1/2*(d*c^2*(c^2*d+e))^(1/2)/d^2/(c^2*d+e)*arctanh(1/4*(4*c^2*d+2*e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2+2*e)/(c^4*d^2+c^2*d*e)^(1/2))-1/4/d^2*sum((_R1^2*e+4*c^2*d+e)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/d^2*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/d^2*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/d^2*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/d^2*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-1/4/d^2*e*sum((_R1^2+1)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))`

### 3.500.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e^2*x^5 + 2*d*e*x^3 + d^2*x), x)`

### 3.500.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(d + ex^2)^2} dx$$

input `integrate((a+b*acosh(c*x))/x/(e*x**2+d)**2,x)`

output `Integral((a + b*acosh(c*x))/(x*(d + e*x**2)**2), x)`



**3.500.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d)^2,x, algorithm="maxima")`

output `1/2*a*(1/(d*e*x^2 + d^2) - log(e*x^2 + d)/d^2 + 2*log(x)/d^2) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(e^2*x^5 + 2*d*e*x^3 + d^2*x), x)`

**3.500.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)^2*x), x)`

**3.500.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(ex^2 + d)^2} dx$$

input `int((a + b*acosh(c*x))/(x*(d + e*x^2)^2), x)`

output `int((a + b*acosh(c*x))/(x*(d + e*x^2)^2), x)`

$$3.501 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d+ex^2)^2} dx$$

3.501.1 Optimal result . . . . .	3690
3.501.2 Mathematica [C] (verified) . . . . .	3691
3.501.3 Rubi [A] (verified) . . . . .	3692
3.501.4 Maple [C] (warning: unable to verify) . . . . .	3694
3.501.5 Fricas [F] . . . . .	3695
3.501.6 Sympy [F(-1)] . . . . .	3695
3.501.7 Maxima [F] . . . . .	3696
3.501.8 Giac [F] . . . . .	3696
3.501.9 Mupad [F(-1)] . . . . .	3696

**3.501.1 Optimal result**

Integrand size = 21, antiderivative size = 634

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^2} dx &= \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{2d^2x} - \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2} - \frac{e(a + \operatorname{barccosh}(cx))}{2d^2(d + ex^2)} \\
&- \frac{2e(a + \operatorname{barccosh}(cx))^2}{bd^3} + \frac{bce\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d\sqrt{-1+c^2x^2}}}\right)}{2d^{5/2}\sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx}} \\
&- \frac{2e(a + \operatorname{barccosh}(cx)) \log(1 + e^{-2\operatorname{arccosh}(cx)})}{d^3} \\
&+ \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{d^3} \\
&+ \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{d^3} \\
&+ \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{d^3} \\
&+ \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{d^3} \\
&+ \frac{be \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{d^3} \\
&+ \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{d^3} \\
&+ \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{d^3} \\
&+ \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{d^3} \\
&+ \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{d^3}
\end{aligned}$$

output

```

1/2*(-a-b*arccosh(c*x))/d^2/x^2-1/2*e*(a+b*arccosh(c*x))/d^2/(e*x^2+d)-2*
e*(a+b*arccosh(c*x))^2/b/d^3-2*e*(a+b*arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/
2)*(c*x+1)^(1/2))^2)/d^3+e*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x
+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^3+e*(a+b*arccosh(c*x
))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)
^(1/2)))/d^3+e*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e
^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^3+e*(a+b*arccosh(c*x))*ln(1+(c*x
+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^3
+b*e*polylog(2,-1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d^3+b*e*polylog(2,-
(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2))
)/d^3+b*e*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)
-(-c^2*d-e)^(1/2)))/d^3+b*e*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e
^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^3+b*e*polylog(2,(c*x+(c*x-1)^(1/
2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^3+1/2*b*c*(c*
x-1)^(1/2)*(c*x+1)^(1/2)/d^2/x+1/2*b*c*e*arctanh(x*(c^2*d+e)^(1/2)/d^(1/2)
/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/d^(5/2)/(c^2*d+e)^(1/2)/(c*x-1)^(1/2
)/(c*x+1)^(1/2)

```

### 3.501.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.55 (sec) , antiderivative size = 792, normalized size of antiderivative = 1.25

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d + ex^2)^2} dx$$

$$= \frac{-\frac{2ad}{x^2} - \frac{2ade}{d+ex^2} - 8ae \log(x) + 4ae \log(d + ex^2) + b \left( \frac{2d(cx\sqrt{-1+cx}\sqrt{1+cx} - \operatorname{arccosh}(cx))}{x^2} - 4e \operatorname{arccosh}(cx)^2 - 4e \operatorname{arccosh}(cx) \right)}{d^2}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d + e*x^2)^2), x]`

output  $((-2*a*d)/x^2 - (2*a*d*e)/(d + e*x^2) - 8*a*e*Log[x] + 4*a*e*Log[d + e*x^2] + b*((2*d*(c*x*Sqrt[-1 + c*x]*Sqrt[1 + c*x] - ArcCosh[c*x]))/x^2 - 4*e*ArcCosh[c*x]^2 - 4*e*ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x]))] + 4*e*ArcCosh[c*x]*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/((-I)*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) + Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) + 4*e*ArcCosh[c*x]*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] - Sqrt[-(c^2*d) - e])]) + Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) + I*Sqrt[d]*e*(ArcCosh[c*x])/((-I)*Sqrt[d] + Sqrt[e]*x) + (c*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*d) - e])*Sqrt[-1 + c*x]*Sqrt[1 + c*x])]/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*Sqrt[e]*x)))/Sqrt[-(c^2*d) - e] + I*Sqrt[d]*e*(-(ArcCosh[c*x])/(I*Sqrt[d] + Sqrt[e]*x)) - (c*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[-(c^2*d) - e])*Sqrt[-1 + c*x]*Sqrt[1 + c*x])]/(c*Sqrt[-(c^2*d) - e]*(I*Sqrt[d] + Sqrt[e]*x)))/Sqrt[-(c^2*d) - e] + 4*e*PolyLog[2, -E^(-2*ArcCosh[c*x])] + 4*e*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] - Sqrt[-(c^2*d) - e])] + 4*e*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/((-I)*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] + 4*e*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])]) + 4*e*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])])/(4*d^3)$

### 3.501.3 Rubi [A] (verified)

Time = 1.61 (sec) , antiderivative size = 634, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \text{barccosh}(cx)}{x^3 (d + ex^2)^2} dx$$

↓ 6374

$$\int \left( \frac{2e^2 x (a + \text{barccosh}(cx))}{d^3 (d + ex^2)} - \frac{2e (a + \text{barccosh}(cx))}{d^3 x} + \frac{e^2 x (a + \text{barccosh}(cx))}{d^2 (d + ex^2)^2} + \frac{a + \text{barccosh}(cx)}{d^2 x^3} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{d^3} + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{d^3} + \\
& \frac{e(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{d^3} + \frac{e(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{d^3} - \\
& \frac{2e(a + \operatorname{barccosh}(cx))^2}{bd^3} - \frac{2e \log(e^{-2\operatorname{arccosh}(cx)} + 1)(a + \operatorname{barccosh}(cx))}{d^3} - \frac{e(a + \operatorname{barccosh}(cx))}{2d^2(d + ex^2)} - \\
& \frac{a + \operatorname{barccosh}(cx)}{2d^2x^2} + \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{d^3} + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{d^3} + \\
& \frac{be \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{d^3} + \frac{be \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{d^3} + \\
& \frac{be \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{d^3} + \frac{bce\sqrt{c^2x^2 - 1} \operatorname{arctanh}\left(\frac{x\sqrt{c^2d + e}}{\sqrt{d}\sqrt{c^2x^2 - 1}}\right)}{2d^{5/2}\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{c^2d + e}} + \frac{bc\sqrt{cx - 1}\sqrt{cx + 1}}{2d^2x}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^3*(d + e*x^2)^2), x]`

output `(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(2*d^2*x) - (a + b*ArcCosh[c*x])/(2*d^2*x^2) - (e*(a + b*ArcCosh[c*x]))/(2*d^2*(d + e*x^2)) - (2*e*(a + b*ArcCosh[c*x])^2)/(b*d^3) + (b*c*e*Sqrt[-1 + c^2*x^2]*ArcTanh[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2])])/(2*d^(5/2)*Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (2*e*(a + b*ArcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])])/d^3 + (e*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/d^3 + (e*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/d^3 + (e*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/d^3 + (e*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/d^3 + (b*e*PolyLog[2, -E^(-2*ArcCosh[c*x])])/d^3 + (b*e*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e]))])/d^3 + (b*e*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/d^3 + (b*e*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e]))])/d^3 + (b*e*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/d^3`

3.501.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

3.501.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 1.61 (sec) , antiderivative size = 635, normalized size of antiderivative = 1.00

method	result
parts	$\frac{ae \ln(ex^2+d)}{d^3} - \frac{ae}{2d^2(ex^2+d)} - \frac{a}{2d^2x^2} - \frac{2ae \ln(x)}{d^3} + b c^2 \left( -\frac{-\sqrt{cx-1}\sqrt{cx+1}c^3 dx - \sqrt{cx-1}\sqrt{cx+1}c^3 e x^3 + c^3 d}{2c^2 x^2 (c^2 d + e)} \right)$
derivativedivides	$c^2 \left( -\frac{a}{2d^2 c^2 x^2} - \frac{2ae \ln(cx)}{c^2 d^3} + \frac{ae \ln(c^2 e x^2 + c^2 d)}{c^2 d^3} - \frac{ae}{2d^2 (c^2 e x^2 + c^2 d)} + b c^4 \left( -\frac{-\sqrt{cx-1}\sqrt{cx+1}c^3 dx - \sqrt{cx-1}\sqrt{cx+1}c^3 e x^3 + c^3 d}{2c^2 x^2 (c^2 d + e)} \right) \right)$
default	$c^2 \left( -\frac{a}{2d^2 c^2 x^2} - \frac{2ae \ln(cx)}{c^2 d^3} + \frac{ae \ln(c^2 e x^2 + c^2 d)}{c^2 d^3} - \frac{ae}{2d^2 (c^2 e x^2 + c^2 d)} + b c^4 \left( -\frac{-\sqrt{cx-1}\sqrt{cx+1}c^3 dx - \sqrt{cx-1}\sqrt{cx+1}c^3 e x^3 + c^3 d}{2c^2 x^2 (c^2 d + e)} \right) \right)$

```
input int((a+b*arccosh(c*x))/x^3/(e*x^2+d)^2,x,method=_RETURNVERBOSE)
```

3.501.  $\int \frac{a+b\text{arccosh}(cx)}{x^3(d+ex^2)^2} dx$

output `a*e/d^3*ln(e*x^2+d)-1/2*a*e/d^2/(e*x^2+d)-1/2*a/d^2/x^2-2*a/d^3*e*ln(x)+b*c^2*(-1/2*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^3*d*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))*c^3*e*x^3+c^4*d*x^2+c^4*e*x^4+arccosh(c*x)*c^2*d+2*arccosh(c*x)*c^2*e*x^2)/c^2/x^2/(c^2*e*x^2+c^2*d)/d^2-1/2*(d*c^2*(c^2*d+e))^(1/2)/d^3/c^2/(c^2*d+e)*e*arctanh(1/4*(4*c^2*d+2*e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2+2*e)/(c^4*d^2+c^2*d*e)^(1/2))+1/2*e/d^3/c^2*sum((_R1^2*e+4*c^2*d+e)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))-2*e/d^3/c^2*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))-2*e/d^3/c^2*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))))-2*e/d^3/c^2*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-2*e/d^3/c^2*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/2*e^2/d^3/c^2*sum((_R1^2+1)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))`

### 3.501.5 Fracas [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e^2*x^7 + 2*d*e*x^5 + d^2*x^3), x)`

### 3.501.6 Sympy [F(-1)]

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^2} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**3/(e*x**2+d)**2,x)`

output `Timed out`



**3.501.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d)^2,x, algorithm="maxima")`

output `-1/2*a*((2*e*x^2 + d)/(d^2*e*x^4 + d^3*x^2) - 2*e*log(e*x^2 + d)/d^3 + 4*e*log(x)/d^3) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(e^2*x^7 + 2*d*e*x^5 + d^2*x^3), x)`

**3.501.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)^2*x^3), x)`

**3.501.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (ex^2 + d)^2} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d + e*x^2)^2),x)`

output `int((a + b*acosh(c*x))/(x^3*(d + e*x^2)^2), x)`

$$3.502 \quad \int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx$$

3.502.1 Optimal result . . . . .	3698
3.502.2 Mathematica [C] (warning: unable to verify) . . . . .	3699
3.502.3 Rubi [A] (verified) . . . . .	3700
3.502.4 Maple [C] (warning: unable to verify) . . . . .	3703
3.502.5 Fricas [F] . . . . .	3704
3.502.6 Sympy [F] . . . . .	3704
3.502.7 Maxima [F(-2)] . . . . .	3704
3.502.8 Giac [F] . . . . .	3705
3.502.9 Mupad [F(-1)] . . . . .	3705

## 3.502.1 Optimal result

Integrand size = 21, antiderivative size = 839

$$\begin{aligned}
\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx &= \frac{ax}{e^2} - \frac{b\sqrt{-1+cx}\sqrt{1+cx}}{ce^2} + \frac{bx\operatorname{arccosh}(cx)}{e^2} \\
&- \frac{d(a + \operatorname{barccosh}(cx))}{4e^{5/2}(\sqrt{-d} - \sqrt{ex})} + \frac{d(a + \operatorname{barccosh}(cx))}{4e^{5/2}(\sqrt{-d} + \sqrt{ex})} \\
&+ \frac{bcd\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}\right)}{2\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{c\sqrt{-d}+\sqrt{e}e^{5/2}}}} \\
&- \frac{bcd\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}\right)}{2\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{c\sqrt{-d}+\sqrt{e}e^{5/2}}}} \\
&+ \frac{3\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{4e^{5/2}} \\
&- \frac{3\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{4e^{5/2}} \\
&+ \frac{3\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{4e^{5/2}} \\
&- \frac{3\sqrt{-d}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{4e^{5/2}} \\
&- \frac{3b\sqrt{-d}\operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{4e^{5/2}} \\
&+ \frac{3b\sqrt{-d}\operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{4e^{5/2}} \\
&- \frac{3b\sqrt{-d}\operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{4e^{5/2}} \\
&+ \frac{3b\sqrt{-d}\operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{4e^{5/2}}
\end{aligned}$$

output

```

a*x/e^2+b*x*arccosh(c*x)/e^2+3/4*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)
)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2))*(-d)^(1/2)/e^(5/
2)-3/4*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(
c*(-d)^(1/2)-(-c^2*d-e)^(1/2))*(-d)^(1/2)/e^(5/2)+3/4*(a+b*arccosh(c*x))*
ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1
/2))*(-d)^(1/2)/e^(5/2)-3/4*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c
*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2))*(-d)^(1/2)/e^(5/2)-3
/4*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-
c^2*d-e)^(1/2))*(-d)^(1/2)/e^(5/2)+3/4*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x
+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2))*(-d)^(1/2)/e^(5/2)-3/
4*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c
^2*d-e)^(1/2))*(-d)^(1/2)/e^(5/2)+3/4*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x
+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2))*(-d)^(1/2)/e^(5/2)-1/4
*d*(a+b*arccosh(c*x))/e^(5/2)/((-d)^(1/2)-x*e^(1/2))+1/4*d*(a+b*arccosh(c*
x))/e^(5/2)/((-d)^(1/2)+x*e^(1/2))-b*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/e^2+1/2
*b*c*d*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*x-1)^(1/2)/(c
*(-d)^(1/2)+e^(1/2))^(1/2))/e^(5/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*(-d)^(
1/2)+e^(1/2))^(1/2)-1/2*b*c*d*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)+e^(1/2))
^(1/2)/(c*x-1)^(1/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2))/e^(5/2)/(c*(-d)^(1/2)-e
^(1/2))^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(1/2)

```

### 3.502.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 1.32 (sec) , antiderivative size = 777, normalized size of antiderivative = 0.93

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx$$

$$= \frac{8a\sqrt{e}x + \frac{4ad\sqrt{e}x}{d+ex^2} - 12a\sqrt{d} \arctan\left(\frac{\sqrt{e}x}{\sqrt{d}}\right) + b \left( \frac{8\sqrt{e} \left( -\sqrt{\frac{-1+cx}{1+cx}}(1+cx) + cx \operatorname{arccosh}(cx) \right)}{c} + 2d \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d} + \sqrt{e}x} + \frac{c \log\left(\dots\right)}{\dots} \right)}{\dots} \right)}{\dots}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output  $(8*a*\sqrt{e}*x + (4*a*d*\sqrt{e}*x)/(d + e*x^2) - 12*a*\sqrt{d}*ArcTan[(\sqrt{e}*x)/\sqrt{d}] + b*((8*\sqrt{e}*(-(\sqrt{(-1 + c*x)/(1 + c*x)}*(1 + c*x)) + c*x*ArcCosh[c*x]))/c + 2*d*(ArcCosh[c*x]/((-I)*\sqrt{d} + \sqrt{e}*x) + (c*\text{Log}[(2*e*(I*\sqrt{e} + c^2*\sqrt{d}*x - I*\sqrt{-(c^2*d) - e})*\sqrt{-1 + c*x}*\sqrt{1 + c*x}))/(\sqrt{-(c^2*d) - e}*(\sqrt{d} + I*\sqrt{e}*x)))/\sqrt{-(c^2*d) - e}) + 2*d*(ArcCosh[c*x]/(I*\sqrt{d} + \sqrt{e}*x) + (c*\text{Log}[(2*e*(-\sqrt{e} - I*c^2*\sqrt{d}*x + \sqrt{-(c^2*d) - e})*\sqrt{-1 + c*x}*\sqrt{1 + c*x}))/(\sqrt{-(c^2*d) - e}*(I*\sqrt{d} + \sqrt{e}*x)))/\sqrt{-(c^2*d) - e}) - (3*I)*\sqrt{d}*(ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(\text{Log}[1 + (\sqrt{e})*E^{\text{ArcCosh}[c*x]})]/(I*c*\sqrt{d} - \sqrt{-(c^2*d) - e}])) + \text{Log}[1 + (\sqrt{e})*E^{\text{ArcCosh}[c*x]})]/(I*c*\sqrt{d} + \sqrt{-(c^2*d) - e}))) + 2*\text{PolyLog}[2, (\sqrt{e})*E^{\text{ArcCosh}[c*x]})/((-I)*c*\sqrt{d} + \sqrt{-(c^2*d) - e}]] + 2*\text{PolyLog}[2, -((\sqrt{e})*E^{\text{ArcCosh}[c*x]})/(I*c*\sqrt{d} + \sqrt{-(c^2*d) - e}))] + (3*I)*\sqrt{d}*(ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(\text{Log}[1 + (\sqrt{e})*E^{\text{ArcCosh}[c*x]})]/((-I)*c*\sqrt{d} + \sqrt{-(c^2*d) - e}])) + \text{Log}[1 - (\sqrt{e})*E^{\text{ArcCosh}[c*x]})/(I*c*\sqrt{d} + \sqrt{-(c^2*d) - e}]])) + 2*\text{PolyLog}[2, (\sqrt{e})*E^{\text{ArcCosh}[c*x]})/(I*c*\sqrt{d} - \sqrt{-(c^2*d) - e}]] + 2*\text{PolyLog}[2, (\sqrt{e})*E^{\text{ArcCosh}[c*x]})/(I*c*\sqrt{d} + \sqrt{-(c^2*d) - e}]])))/(8*e^{(5/2)})$

### 3.502.3 Rubi [A] (verified)

Time = 2.63 (sec) , antiderivative size = 839, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + \text{barccosh}(cx))}{(d + ex^2)^2} dx$$

$$\downarrow 6374$$

$$\int \left( \frac{d^2(a + \text{barccosh}(cx))}{e^2(d + ex^2)^2} - \frac{2d(a + \text{barccosh}(cx))}{e^2(d + ex^2)} + \frac{a + \text{barccosh}(cx)}{e^2} \right) dx$$

$$\downarrow 2009$$

---

3.502.  $\int \frac{x^4(a + \text{barccosh}(cx))}{(d + ex^2)^2} dx$

$$\begin{aligned}
& \frac{x \operatorname{arccosh}(cx) b}{e^2} + \frac{c d \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{-dc+\sqrt{e}\sqrt{cx-1}}}\right) b}{2\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{-dc+\sqrt{e}e^{5/2}}}} - \frac{c d \operatorname{arctanh}\left(\frac{\sqrt{\sqrt{-dc+\sqrt{e}\sqrt{cx+1}}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right) b}{2\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{-dc+\sqrt{e}e^{5/2}}}} \\
& \frac{3\sqrt{-d} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right) b}{4e^{5/2}} + \frac{3\sqrt{-d} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right) b}{4e^{5/2}} - \\
& \frac{3\sqrt{-d} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc+\sqrt{-dc^2-e}}}\right) b}{4e^{5/2}} + \frac{3\sqrt{-d} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc+\sqrt{-dc^2-e}}}\right) b}{4e^{5/2}} - \\
& \frac{\sqrt{cx-1}\sqrt{cx+1}b}{ce^2} + \frac{ax}{e^2} - \frac{d(a+b\operatorname{arccosh}(cx))}{4e^{5/2}(\sqrt{-d}-\sqrt{ex})} + \frac{d(a+b\operatorname{arccosh}(cx))}{4e^{5/2}(\sqrt{ex}+\sqrt{-d})} + \\
& \frac{3\sqrt{-d}(a+b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{4e^{5/2}} - \\
& \frac{3\sqrt{-d}(a+b\operatorname{arccosh}(cx)) \log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{c\sqrt{-d}-\sqrt{-dc^2-e}} + 1\right)}{4e^{5/2}} + \\
& \frac{3\sqrt{-d}(a+b\operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc+\sqrt{-dc^2-e}}}\right)}{4e^{5/2}} - \\
& \frac{3\sqrt{-d}(a+b\operatorname{arccosh}(cx)) \log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{\sqrt{-dc+\sqrt{-dc^2-e}} + 1}\right)}{4e^{5/2}}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

```

output (a*x)/e^2 - (b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c*e^2) + (b*x*ArcCosh[c*x])/
e^2 - (d*(a + b*ArcCosh[c*x]))/(4*e^(5/2)*(Sqrt[-d] - Sqrt[e]*x)) + (d*(a
+ b*ArcCosh[c*x]))/(4*e^(5/2)*(Sqrt[-d] + Sqrt[e]*x)) + (b*c*d*ArcTanh[(Sq
rt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[-
1 + c*x])])/(2*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*e^(5/
2)) - (b*c*d*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sq
rt[-d] - Sqrt[e]]*Sqrt[-1 + c*x])])/(2*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*S
qrt[-d] + Sqrt[e]]*e^(5/2)) + (3*Sqrt[-d]*(a + b*ArcCosh[c*x])*Log[1 - (Sq
rt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(4*e^(5/2)) - (3
*Sqrt[-d]*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d
] - Sqrt[-(c^2*d) - e])])/(4*e^(5/2)) + (3*Sqrt[-d]*(a + b*ArcCosh[c*x])*L
og[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(4*e^(
5/2)) - (3*Sqrt[-d]*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/
(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(4*e^(5/2)) - (3*b*Sqrt[-d]*PolyLog[2,
-((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e]))])/(4*e^(5/2
)) + (3*b*Sqrt[-d]*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[
-(c^2*d) - e])])/(4*e^(5/2)) - (3*b*Sqrt[-d]*PolyLog[2, -((Sqrt[e]*E^ArcCo
sh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e]))])/(4*e^(5/2)) + (3*b*Sqrt[-d]*
PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(4
*e^(5/2))

```

### 3.502.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

### 3.502.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 31.12 (sec) , antiderivative size = 897, normalized size of antiderivative = 1.07

method	result	size
parts	Expression too large to display	897
derivativedivides	Expression too large to display	913
default	Expression too large to display	913

```
input int(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)
```

```
output a*(1/e^2*x-1/e^2*d*(-1/2*x/(e*x^2+d)+3/2/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2)
))))+b/c^5*(1/2*c^4*(-1+arccosh(c*x))/e^2*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)
)+1/2*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*c^4*(1+arccosh(c*x))/e^2+1/2*d*ar
ccosh(c*x)*c^7*x/e^2/(c^2*e*x^2+c^2*d)+1/2*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(
1/2)+e)*e)^(1/2)*(2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e+(d*
c^2*(c^2*d+e))^(1/2)*e)*d*c^6*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/
((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))/e^5/(c^2*d+e)-1/2*(-(2*c
^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d+2*(d*c^2*(c^2*d+e))^(1
/2)+e)*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((-2*c^2*d+2*(d*c^2*(c
^2*d+e))^(1/2)-e)*e)^(1/2))*d*c^6/e^5+1/2*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/
2)+e)*e)^(1/2)*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c
^2*(c^2*d+e))^(1/2)*e)*d*c^6*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((2
*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2))/e^5/(c^2*d+e)-1/2*((2*c^2*d+
2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e
)*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((2*c^2*d+2*(d*c^2*(c^2*d+e))
^(1/2)+e)*e)^(1/2))*d*c^6/e^5+3/4*d/e^2*c^6*sum(1/_R1/(_R1^2*e+2*c^2*d+e)*
(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x
-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e
))-3/4*d/e^2*c^6*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*
x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/...
```



**3.502.5 Fracas [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(ex^2 + d)^2} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*x^4*arccosh(c*x) + a*x^4)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

**3.502.6 Sympy [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(d + ex^2)^2} dx$$

input `integrate(x**4*(a+b*acosh(c*x))/(e*x**2+d)**2,x)`

output `Integral(x**4*(a + b*acosh(c*x))/(d + e*x**2)**2, x)`

**3.502.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.502.8 Giac [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(ex^2 + d)^2} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^4/(e*x^2 + d)^2, x)`

**3.502.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^2} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d + e*x^2)^2,x)`

output `int((x^4*(a + b*acosh(c*x)))/(d + e*x^2)^2, x)`

$$3.503 \quad \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^2} dx$$

3.503.1 Optimal result . . . . .	3707
3.503.2 Mathematica [C] (verified) . . . . .	3708
3.503.3 Rubi [A] (verified) . . . . .	3709
3.503.4 Maple [C] (warning: unable to verify) . . . . .	3711
3.503.5 Fricas [F] . . . . .	3712
3.503.6 Sympy [F] . . . . .	3712
3.503.7 Maxima [F(-2)] . . . . .	3713
3.503.8 Giac [F] . . . . .	3713
3.503.9 Mupad [F(-1)] . . . . .	3713

## 3.503.1 Optimal result

Integrand size = 21, antiderivative size = 792

$$\begin{aligned}
\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = & \frac{a + \operatorname{barccosh}(cx)}{4e^{3/2}(\sqrt{-d} - \sqrt{ex})} - \frac{a + \operatorname{barccosh}(cx)}{4e^{3/2}(\sqrt{-d} + \sqrt{ex})} \\
& - \frac{b \operatorname{carctanh}\left(\frac{\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{1+cx}}}\right)}{2\sqrt{c\sqrt{-d} - \sqrt{e}}\sqrt{c\sqrt{-d} + \sqrt{e}}e^{3/2}} \\
& + \frac{b \operatorname{carctanh}\left(\frac{\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{1+cx}}}\right)}{2\sqrt{c\sqrt{-d} - \sqrt{e}}\sqrt{c\sqrt{-d} + \sqrt{e}}e^{3/2}} \\
& + \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}} \\
& - \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}} \\
& + \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}} \\
& - \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}} \\
& - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}} \\
& - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4\sqrt{-d}e^{3/2}}
\end{aligned}$$

output

```

1/4*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)-1/4*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)+1/4*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)-1/4*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)-1/4*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)+1/4*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)+1/4*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)+1/4*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(3/2)/(-d)^(1/2)+1/4*(a+b*arccosh(c*x))/e^(3/2)/(-d)^(1/2)-x*e^(1/2))+1/4*(-a-b*arccosh(c*x))/e^(3/2)/((-d)^(1/2)+x*e^(1/2))-1/2*b*c*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*x-1)^(1/2))/(c*(-d)^(1/2)+e^(1/2))^(1/2))/e^(3/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(1/2)+1/2*b*c*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)+e^(1/2))^(1/2)/(c*x-1)^(1/2))/(c*(-d)^(1/2)-e^(1/2))^(1/2))/e^(3/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(1/2)
    
```

### 3.503.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.08 (sec) , antiderivative size = 719, normalized size of antiderivative = 0.91

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx$$

$$= \frac{-\frac{4a\sqrt{ex}}{d+ex^2} + \frac{4a \arctan\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{\sqrt{d}}}{b} + b \left( -\frac{2\operatorname{arccosh}(cx)}{i\sqrt{d}+\sqrt{ex}} - 2 \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d}+\sqrt{ex}} + \frac{c \log\left(\frac{2e(i\sqrt{e}+c^2\sqrt{dx}-i\sqrt{-c^2d-e}\sqrt{-1+cx}\sqrt{1+cx})}{c\sqrt{-c^2d-e}(\sqrt{d}+i\sqrt{ex})}\right)}{\sqrt{-c^2d-e}} \right) \right)$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output  $((-4*a*\text{Sqrt}[e]*x)/(d + e*x^2) + (4*a*\text{ArcTan}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d]])/\text{Sqrt}[d] + b*((-2*\text{ArcCosh}[c*x])/(I*\text{Sqrt}[d] + \text{Sqrt}[e]*x) - 2*(\text{ArcCosh}[c*x]/((-I)*\text{Sqrt}[d] + \text{Sqrt}[e]*x) + (c*\text{Log}[(2*e*(I*\text{Sqrt}[e] + c^2*\text{Sqrt}[d]*x - I*\text{Sqrt}[-(c^2*d) - e]*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])]/(c*\text{Sqrt}[-(c^2*d) - e]*(\text{Sqrt}[d] + I*\text{Sqrt}[e]*x)))/\text{Sqrt}[-(c^2*d) - e]) - (2*c*\text{Log}[(2*e*(-\text{Sqrt}[e] - I*c^2*\text{Sqrt}[d]*x + \text{Sqrt}[-(c^2*d) - e]*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])]/(c*\text{Sqrt}[-(c^2*d) - e]*(I*\text{Sqrt}[d] + \text{Sqrt}[e]*x)))/\text{Sqrt}[-(c^2*d) - e] + (I*(\text{ArcCosh}[c*x]*(-\text{ArcCosh}[c*x] + 2*(\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] - \text{Sqrt}[-(c^2*d) - e])) + \text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e]))]) + 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/((-I)*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])] + 2*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])))/\text{Sqrt}[d] + (I*(\text{ArcCosh}[c*x]*(\text{ArcCosh}[c*x] - 2*(\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/((-I)*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])) + \text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e]))]) - 2*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/((-I)*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e]))] - 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])))/\text{Sqrt}[d]))/(8*e^{(3/2)})$

### 3.503.3 Rubi [A] (verified)

Time = 2.38 (sec) , antiderivative size = 792, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2(a + \text{barccosh}(cx))}{(d + ex^2)^2} dx$$

$$\downarrow \text{6374}$$

$$\int \left( \frac{a + \text{barccosh}(cx)}{e(d + ex^2)} - \frac{d(a + \text{barccosh}(cx))}{e(d + ex^2)^2} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{4\sqrt{-d}e^{3/2}} - \frac{(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{4\sqrt{-d}e^{3/2}} + \\
& \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{4\sqrt{-d}e^{3/2}} - \frac{(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{4\sqrt{-d}e^{3/2}} + \\
& \frac{a + \operatorname{barccosh}(cx)}{4e^{3/2}(\sqrt{-d} - \sqrt{ex})} - \frac{a + \operatorname{barccosh}(cx)}{4e^{3/2}(\sqrt{-d} + \sqrt{ex})} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{dc^2 - e}}\right)}{4\sqrt{-d}e^{3/2}} + \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{dc^2 - e}}\right)}{4\sqrt{-d}e^{3/2}} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{dc^2 - e}}\right)}{4\sqrt{-d}e^{3/2}} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{dc^2 - e}}\right)}{4\sqrt{-d}e^{3/2}} - \\
& \frac{b \operatorname{arctanh}\left(\frac{\sqrt{cx+1}\sqrt{c\sqrt{-d}-\sqrt{e}}}{\sqrt{cx-1}\sqrt{c\sqrt{-d}+\sqrt{e}}}\right)}{2e^{3/2}\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{cx+1}\sqrt{c\sqrt{-d}+\sqrt{e}}}{\sqrt{cx-1}\sqrt{c\sqrt{-d}-\sqrt{e}}}\right)}{2e^{3/2}\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output `(a + b*ArcCosh[c*x])/(4*e^(3/2)*(Sqrt[-d] - Sqrt[e]*x)) - (a + b*ArcCosh[c*x])/(4*e^(3/2)*(Sqrt[-d] + Sqrt[e]*x)) - (b*c*ArcTanh[(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[-1 + c*x])])/(2*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*e^(3/2)) + (b*c*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[-1 + c*x])])/(2*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*e^(3/2)) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(4*Sqrt[-d]*e^(3/2)) - ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(4*Sqrt[-d]*e^(3/2)) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(4*Sqrt[-d]*e^(3/2)) - ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(4*Sqrt[-d]*e^(3/2)) - (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])]/(4*Sqrt[-d]*e^(3/2)) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(4*Sqrt[-d]*e^(3/2)) - (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])]/(4*Sqrt[-d]*e^(3/2)) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(4*Sqrt[-d]*e^(3/2))`

3.503.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

3.503.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 24.16 (sec) , antiderivative size = 813, normalized size of antiderivative = 1.03

method	result
derivativedivides	$-\frac{a c^5 x}{2e(c^2 e x^2 + c^2 d)} + \frac{a c^3 \arctan\left(\frac{e x}{\sqrt{d e}}\right)}{2e \sqrt{d e}} + b c^4 \left( -\frac{\operatorname{arccosh}(c x) c x}{2e(c^2 e x^2 + c^2 d)} - \frac{\sqrt{(2c^2 d + 2\sqrt{d c^2(c^2 d + e)} + e)} e (-2\sqrt{d c^2(c^2 d + e)} c^2 d + 2c^4 d^2 + 2e^2 d^2)}{2e(c^2 e x^2 + c^2 d)} \right)$
default	$-\frac{a c^5 x}{2e(c^2 e x^2 + c^2 d)} + \frac{a c^3 \arctan\left(\frac{e x}{\sqrt{d e}}\right)}{2e \sqrt{d e}} + b c^4 \left( -\frac{\operatorname{arccosh}(c x) c x}{2e(c^2 e x^2 + c^2 d)} - \frac{\sqrt{(2c^2 d + 2\sqrt{d c^2(c^2 d + e)} + e)} e (-2\sqrt{d c^2(c^2 d + e)} c^2 d + 2c^4 d^2 + 2e^2 d^2)}{2e(c^2 e x^2 + c^2 d)} \right)$
parts	$-\frac{a x}{2e(e x^2 + d)} + \frac{a \arctan\left(\frac{e x}{\sqrt{d e}}\right)}{2e \sqrt{d e}} + b \left( -\frac{c^5 \operatorname{arccosh}(c x) x}{2e(c^2 e x^2 + c^2 d)} - \frac{\sqrt{-(2c^2 d - 2\sqrt{d c^2(c^2 d + e)} + e)} e (2\sqrt{d c^2(c^2 d + e)} c^2 d + 2c^4 d^2 + 2e^2 d^2)}{2e(c^2 e x^2 + c^2 d)} \right)$

input `int(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`



output `1/c^3*(-1/2*a*c^5/e*x/(c^2*e*x^2+c^2*d)+1/2*a*c^3/e/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2))+b*c^4*(-1/2*arccosh(c*x)/e*c*x/(c^2*e*x^2+c^2*d)-1/2*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2))/e^4/(c^2*d+e)+1/2*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2))/e^4-1/2*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e+(d*c^2*(c^2*d+e))^(1/2)*e)*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))/e^4/(c^2*d+e)+1/2*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))/e^4+1/4/e*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))-1/4/e*sum(1/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e)))`

### 3.503.5 Fracas [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^2}{(ex^2 + d)^2} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*x^2*arccosh(c*x) + a*x^2)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

### 3.503.6 Sympy [F]

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(d + ex^2)^2} dx$$

input `integrate(x**2*(a+b*acosh(c*x))/(e*x**2+d)**2,x)`

output `Integral(x**2*(a + b*acosh(c*x))/(d + e*x**2)**2, x)`

---

3.503.  $\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx$

**3.503.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.503.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(ex^2 + d)^2} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^2/(e*x^2 + d)^2, x)`

**3.503.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^2} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d + e*x^2)^2,x)`

output `int((x^2*(a + b*acosh(c*x)))/(d + e*x^2)^2, x)`

$$3.504 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^2} dx$$

3.504.1 Optimal result . . . . .	3715
3.504.2 Mathematica [C] (verified) . . . . .	3716
3.504.3 Rubi [A] (verified) . . . . .	3717
3.504.4 Maple [C] (warning: unable to verify) . . . . .	3719
3.504.5 Fricas [F] . . . . .	3720
3.504.6 Sympy [F] . . . . .	3720
3.504.7 Maxima [F(-2)] . . . . .	3721
3.504.8 Giac [F] . . . . .	3721
3.504.9 Mupad [F(-1)] . . . . .	3721

**3.504.1 Optimal result**

Integrand size = 18, antiderivative size = 804

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^2} dx = & -\frac{a + \operatorname{barccosh}(cx)}{4d\sqrt{e}(\sqrt{-d} - \sqrt{ex})} + \frac{a + \operatorname{barccosh}(cx)}{4d\sqrt{e}(\sqrt{-d} + \sqrt{ex})} \\
& + \frac{b \operatorname{carctanh}\left(\frac{\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{1+cx}}}\right)}{2d\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{e}}}} \\
& - \frac{b \operatorname{carctanh}\left(\frac{\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{1+cx}}}\right)}{2d\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{e}}}} \\
& - \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
& + \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
& - \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
& + \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2d - e}}\right)}{4(-d)^{3/2}\sqrt{e}}
\end{aligned}$$

output

```

-1/4*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*
(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(1/2)+1/4*(a+b*arccosh(c*x))*ln
(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2
)))/(-d)^(3/2)/e^(1/2)-1/4*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x
+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(1/2)+1/4
*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)
^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(1/2)+1/4*b*polylog(2,-(c*x+(c*x-1)
^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(3/2)/
e^(1/2)-1/4*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(
1/2)-(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(1/2)+1/4*b*polylog(2,-(c*x+(c*x-1)^(
1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e
^(1/2)-1/4*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(
1/2)+(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(1/2)+1/4*(-a-b*arccosh(c*x))/d/e^(1/
2)/((-d)^(1/2)-x*e^(1/2))+1/4*(a+b*arccosh(c*x))/d/e^(1/2)/((-d)^(1/2)+x*e
^(1/2))+1/2*b*c*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*x-1)
^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(1/2))/d/e^(1/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2
)/(c*(-d)^(1/2)+e^(1/2))^(1/2)-1/2*b*c*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)
+e^(1/2))^(1/2)/(c*x-1)^(1/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2))/d/e^(1/2)/(c*(-
d)^(1/2)-e^(1/2))^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(1/2)

```

### 3.504.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.39 (sec) , antiderivative size = 734, normalized size of antiderivative = 0.91

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx = \frac{1}{2} \left( \frac{ax}{d^2 + dex^2} + \frac{a \arctan\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{d^{3/2} \sqrt{e}} \right) + b \left( 2\sqrt{d} \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d} + \sqrt{ex}} + \frac{\operatorname{clog}\left(\frac{2e(i\sqrt{e} + c^2\sqrt{dx} - i\sqrt{-c^2d - e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c\sqrt{-c^2d - e}(\sqrt{d} + i\sqrt{ex})}\right)}{\sqrt{-c^2d - e}} \right) - 2\sqrt{d} \left( -\frac{\operatorname{arccosh}(cx)}{i\sqrt{d} + \sqrt{ex}} - \frac{\operatorname{clog}\left(\frac{2e(-\sqrt{e} - ic^2\sqrt{dx} - i\sqrt{-c^2d - e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c\sqrt{-c^2d - e}(\sqrt{d} + i\sqrt{ex})}\right)}{\sqrt{-c^2d - e}} \right) \right)$$

input `Integrate[(a + b*ArcCosh[c*x])/(d + e*x^2)^2,x]`

3.504.  $\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx$

output 
$$\begin{aligned} & ((a*x)/(d^2 + d*e*x^2) + (a*\text{ArcTan}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d]])/(d^{3/2}*\text{Sqrt}[e]) \\ & + (b*(2*\text{Sqrt}[d]*(\text{ArcCosh}[c*x]/((-I)*\text{Sqrt}[d] + \text{Sqrt}[e]*x) + (c*\text{Log}[(2*e*(I \\ & * \text{Sqrt}[e] + c^2*\text{Sqrt}[d]*x - I*\text{Sqrt}[-(c^2*d) - e]*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c* \\ & x]))/(c*\text{Sqrt}[-(c^2*d) - e]*(\text{Sqrt}[d] + I*\text{Sqrt}[e]*x)))/\text{Sqrt}[-(c^2*d) - e]) \\ & - 2*\text{Sqrt}[d]*(-\text{ArcCosh}[c*x]/(I*\text{Sqrt}[d] + \text{Sqrt}[e]*x)) - (c*\text{Log}[(2*e*(-\text{Sqrt}[ \\ & e] - I*c^2*\text{Sqrt}[d]*x + \text{Sqrt}[-(c^2*d) - e]*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]))/( \\ & c*\text{Sqrt}[-(c^2*d) - e]*(I*\text{Sqrt}[d] + \text{Sqrt}[e]*x)))/\text{Sqrt}[-(c^2*d) - e]) + I*(\text{A} \\ & \text{rcCosh}[c*x]*(-\text{ArcCosh}[c*x] + 2*(\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt} \\ & [d] - \text{Sqrt}[-(c^2*d) - e])]) + \text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] \\ & + \text{Sqrt}[-(c^2*d) - e])])) + 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/((-I)*c* \\ & \text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])] + 2*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/( \\ & I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])]) - I*(\text{ArcCosh}[c*x]*(-\text{ArcCosh}[c*x] + 2* \\ & (\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/((-I)*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])]) + \\ & \text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])])) + \\ & 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] - \text{Sqrt}[-(c^2*d) - e])] \\ & + 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e]) \\ & ])))/(4*d^{3/2}*\text{Sqrt}[e])/2 \end{aligned}$$

### 3.504.3 Rubi [A] (verified)

Time = 1.52 (sec) , antiderivative size = 804, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \text{barccosh}(cx)}{(d + ex^2)^2} dx \\ & \quad \downarrow \text{6324} \\ & \int \left( -\frac{e(a + \text{barccosh}(cx))}{2d(-de - e^2x^2)} - \frac{e(a + \text{barccosh}(cx))}{4d(\sqrt{-d}\sqrt{e} - ex)^2} - \frac{e(a + \text{barccosh}(cx))}{4d(\sqrt{-d}\sqrt{e} + ex)^2} \right) dx \\ & \quad \downarrow \text{2009} \end{aligned}$$

$$\begin{aligned}
& -\frac{\log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)(a + \operatorname{barccosh}(cx))}{4(-d)^{3/2}\sqrt{e}} + \frac{\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{c\sqrt{-d}-\sqrt{-dc^2-e}} + 1\right)(a + \operatorname{barccosh}(cx))}{4(-d)^{3/2}\sqrt{e}} - \\
& \frac{\log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)(a + \operatorname{barccosh}(cx))}{4(-d)^{3/2}\sqrt{e}} + \frac{\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{\sqrt{-dc}+\sqrt{-dc^2-e}} + 1\right)(a + \operatorname{barccosh}(cx))}{4(-d)^{3/2}\sqrt{e}} - \\
& \frac{a + \operatorname{barccosh}(cx)}{4d\sqrt{e}(\sqrt{-d} - \sqrt{ex})} + \frac{a + \operatorname{barccosh}(cx)}{4d\sqrt{e}(\sqrt{ex} + \sqrt{-d})} + \frac{b\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{-dc}+\sqrt{e}\sqrt{cx-1}}\right)}{2d\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{-dc} + \sqrt{e}\sqrt{e}}} - \\
& \frac{b\operatorname{arctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right)}{2d\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{-dc} + \sqrt{e}\sqrt{e}}} + \frac{b\operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{4(-d)^{3/2}\sqrt{e}} - \\
& \frac{b\operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{4(-d)^{3/2}\sqrt{e}} + \frac{b\operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{4(-d)^{3/2}\sqrt{e}} - \frac{b\operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{4(-d)^{3/2}\sqrt{e}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d + e*x^2)^2, x]`

output `-1/4*(a + b*ArcCosh[c*x])/(d*sqrt[e]*(sqrt[-d] - sqrt[e]*x)) + (a + b*ArcCosh[c*x])/(4*d*sqrt[e]*(sqrt[-d] + sqrt[e]*x)) + (b*c*ArcTanh[(sqrt[c*sqrt[-d] - sqrt[e]]*sqrt[1 + c*x])/(sqrt[c*sqrt[-d] + sqrt[e]]*sqrt[-1 + c*x])])/(2*d*sqrt[c*sqrt[-d] - sqrt[e]]*sqrt[c*sqrt[-d] + sqrt[e]]*sqrt[e]) - (b*c*ArcTanh[(sqrt[c*sqrt[-d] + sqrt[e]]*sqrt[1 + c*x])/(sqrt[c*sqrt[-d] - sqrt[e]]*sqrt[-1 + c*x])])/(2*d*sqrt[c*sqrt[-d] - sqrt[e]]*sqrt[c*sqrt[-d] + sqrt[e]]*sqrt[e]) - ((a + b*ArcCosh[c*x])*Log[1 - (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/(4*(-d)^(3/2)*sqrt[e]) + ((a + b*ArcCosh[c*x])*Log[1 + (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/(4*(-d)^(3/2)*sqrt[e]) - ((a + b*ArcCosh[c*x])*Log[1 - (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(4*(-d)^(3/2)*sqrt[e]) + ((a + b*ArcCosh[c*x])*Log[1 + (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(4*(-d)^(3/2)*sqrt[e]) + (b*PolyLog[2, -((sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e]))]/(4*(-d)^(3/2)*sqrt[e]) - (b*PolyLog[2, (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] - sqrt[-(c^2*d) - e])])/(4*(-d)^(3/2)*sqrt[e]) + (b*PolyLog[2, -((sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e]))]/(4*(-d)^(3/2)*sqrt[e]) - (b*PolyLog[2, (sqrt[e]*E^ArcCosh[c*x])/(c*sqrt[-d] + sqrt[-(c^2*d) - e])])/(4*(-d)^(3/2)*sqrt[e])`

3.504.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

3.504.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 20.61 (sec) , antiderivative size = 828, normalized size of antiderivative = 1.03

method	result
parts	$\frac{ax}{2d(ex^2+d)} + \frac{a \arctan\left(\frac{ex}{\sqrt{de}}\right)}{2d\sqrt{de}} + b \left( \frac{c^3 \operatorname{arccosh}(cx)x}{2d(c^2ex^2+c^2d)} + \frac{c^2 \left( \frac{-R1\left(\operatorname{arccosh}(cx)\right)}{-R1=\operatorname{RootOf}\left(e-Z^4+(4c^2d+2e)-Z^2+e\right)} \right)}{2d(c^2ex^2+c^2d)} \right)$
derivativedivides	$\frac{ac^3x}{2d(c^2ex^2+c^2d)} + \frac{ac \arctan\left(\frac{ex}{\sqrt{de}}\right)}{2d\sqrt{de}} + bc^4 \left( \frac{\operatorname{arccosh}(cx)x}{2cd(c^2ex^2+c^2d)} + \frac{-R1\left(\operatorname{arccosh}(cx)\right)}{-R1=\operatorname{RootOf}\left(e-Z^4+(4c^2d+2e)-Z^2+e\right)} \right)$
default	$\frac{ac^3x}{2d(c^2ex^2+c^2d)} + \frac{ac \arctan\left(\frac{ex}{\sqrt{de}}\right)}{2d\sqrt{de}} + bc^4 \left( \frac{\operatorname{arccosh}(cx)x}{2cd(c^2ex^2+c^2d)} + \frac{-R1\left(\operatorname{arccosh}(cx)\right)}{-R1=\operatorname{RootOf}\left(e-Z^4+(4c^2d+2e)-Z^2+e\right)} \right)$

input `int((a+b*arccosh(c*x))/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

3.504.  $\int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^2} dx$



output `1/2*a*x/d/(e*x^2+d)+1/2*a/d/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2))+b/c*(1/2*c^3*arccosh(c*x)*x/d/(c^2*e*x^2+c^2*d)+1/4/d*c^2*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/2*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e+(d*c^2*(c^2*d+e))^(1/2)*e)*c^2*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))/d/(c^2*d+e)/e^3-1/2*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))*c^2/d/e^3+1/2*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^2*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2))/d/(c^2*d+e)/e^3-1/2*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2))*c^2/d/e^3-1/4/d*c^2*sum(1/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))`

### 3.504.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`

### 3.504.6 Sympy [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(d + ex^2)^2} dx$$

input `integrate((a+b*acosh(c*x))/(e*x**2+d)**2,x)`

output `Integral((a + b*acosh(c*x))/(d + e*x**2)**2, x)`

---

3.504.  $\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx$

**3.504.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.504.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(e*x^2 + d)^2, x)`

**3.504.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(ex^2 + d)^2} dx$$

input `int((a + b*acosh(c*x))/(d + e*x^2)^2,x)`

output `int((a + b*acosh(c*x))/(d + e*x^2)^2, x)`

$$3.505 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d+ex^2)^2} dx$$

3.505.1 Optimal result . . . . .	3723
3.505.2 Mathematica [C] (verified) . . . . .	3724
3.505.3 Rubi [A] (verified) . . . . .	3725
3.505.4 Maple [C] (warning: unable to verify) . . . . .	3728
3.505.5 Fricas [F] . . . . .	3729
3.505.6 Sympy [F(-1)] . . . . .	3729
3.505.7 Maxima [F(-2)] . . . . .	3729
3.505.8 Giac [F] . . . . .	3730
3.505.9 Mupad [F(-1)] . . . . .	3730

## 3.505.1 Optimal result

Integrand size = 21, antiderivative size = 846

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d + ex^2)^2} dx = & -\frac{a + \operatorname{barccosh}(cx)}{d^2 x} + \frac{\sqrt{e}(a + \operatorname{barccosh}(cx))}{4d^2 (\sqrt{-d} - \sqrt{ex})} \\
& - \frac{\sqrt{e}(a + \operatorname{barccosh}(cx))}{4d^2 (\sqrt{-d} + \sqrt{ex})} + \frac{bc \arctan(\sqrt{-1 + cx}\sqrt{1 + cx})}{d^2} \\
& - \frac{bc\sqrt{e} \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{1+cx}}}\right)}{2d^2 \sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{c\sqrt{-d} + \sqrt{e}}}} \\
& + \frac{bc\sqrt{e} \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d} + \sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{1+cx}}}\right)}{2d^2 \sqrt{c\sqrt{-d} - \sqrt{e}\sqrt{c\sqrt{-d} + \sqrt{e}}}} \\
& - \frac{3\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}} \\
& + \frac{3\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}} \\
& - \frac{3\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}} \\
& + \frac{3\sqrt{e}(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}} \\
& + \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}} \\
& - \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}} \\
& + \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}} \\
& - \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{4(-d)^{5/2}}
\end{aligned}$$

output

$$\begin{aligned}
& (-a-b*\operatorname{arccosh}(c*x))/d^2/x+b*c*\arctan((c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})/d^2-3/4* \\
& (a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*e^{(1/2)}/(c*(-d)^{(1/2)} \\
& -(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(5/2)}+3/4*(a+b*\operatorname{arccosh}(c*x))*\ln(1+(c \\
& *x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2)}))*e \\
& ^{(1/2)}/(-d)^{(5/2)}-3/4*(a+b*\operatorname{arccosh}(c*x))*\ln(1-(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)} \\
& ))*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(5/2)}+3/4*(a+b \\
& *\operatorname{arccosh}(c*x))*\ln(1+(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*e^{(1/2)}/(c*(-d)^{(1/2)} \\
& )+(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(5/2)}+3/4*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)} \\
& )*(c*x+1)^{(1/2)})*e^{(1/2)}/(c*(-d)^{(1/2)}-(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(5/2)} \\
& -3/4*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*e^{(1/2)}/(c*(-d)^{(1/2)} \\
& -(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(5/2)}+3/4*b*\operatorname{polylog}(2,-(c*x+(c*x-1)^{(1/2)} \\
& )*(c*x+1)^{(1/2)})*e^{(1/2)}/(c*(-d)^{(1/2)}+(-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(5/2)} \\
& -3/4*b*\operatorname{polylog}(2,(c*x+(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)})*e^{(1/2)}/(c*(-d)^{(1/2)}+ \\
& (-c^2*d-e)^{(1/2)}))*e^{(1/2)}/(-d)^{(5/2)}+1/4*(a+b*\operatorname{arccosh}(c*x))*e^{(1/2)}/d^2/( \\
& (-d)^{(1/2)}-x*e^{(1/2)})-1/4*(a+b*\operatorname{arccosh}(c*x))*e^{(1/2)}/d^2/((-d)^{(1/2)}+x*e^{(1/2)}) \\
& -1/2*b*c*\operatorname{arctanh}((c*x+1)^{(1/2)}*(c*(-d)^{(1/2)}-e^{(1/2)})^{(1/2)}/(c*x-1)^{(1/2)} \\
& )/(c*(-d)^{(1/2)}+e^{(1/2)})^{(1/2)})*e^{(1/2)}/d^2/(c*(-d)^{(1/2)}-e^{(1/2)})^{(1/2)} \\
& )/(c*(-d)^{(1/2)}+e^{(1/2)})^{(1/2)}+1/2*b*c*\operatorname{arctanh}((c*x+1)^{(1/2)}*(c*(-d)^{(1/2)} \\
& +e^{(1/2)})^{(1/2)}/(c*x-1)^{(1/2)}/(c*(-d)^{(1/2)}-e^{(1/2)})^{(1/2)})*e^{(1/2)}/d^2/(c \\
& *(-d)^{(1/2)}-e^{(1/2)})^{(1/2)}/(c*(-d)^{(1/2)}+e^{(1/2)})^{(1/2)}
\end{aligned}$$

### 3.505.2 Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.72 (sec) , antiderivative size = 821, normalized size of antiderivative = 0.97

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^2 (d + ex^2)^2} dx$$

$$\begin{aligned}
& -\frac{8a\sqrt{d}}{x} - \frac{4a\sqrt{d}ex}{d+ex^2} - 12a\sqrt{e} \arctan\left(\frac{\sqrt{ex}}{\sqrt{d}}\right) + b \left( 8\sqrt{d} \left( -\frac{\operatorname{arccosh}(cx)}{x} + \frac{c\sqrt{-1+c^2x^2} \arctan\left(\frac{\sqrt{-1+c^2x^2}}{\sqrt{-1+cx}\sqrt{1+cx}}\right)}{\sqrt{-1+cx}\sqrt{1+cx}} \right) - 2\sqrt{d}\sqrt{e} \right) \\
& = \dots
\end{aligned}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^2*(d + e*x^2)^2), x]`

$$3.505. \quad \int \frac{a+b\operatorname{arccosh}(cx)}{x^2(d+ex^2)^2} dx$$

output 
$$\begin{aligned} & ((-8*a*\text{Sqrt}[d])/x - (4*a*\text{Sqrt}[d]*e*x)/(d + e*x^2) - 12*a*\text{Sqrt}[e]*\text{ArcTan}[(\text{Sqrt}[e]*x)/\text{Sqrt}[d]] + b*(8*\text{Sqrt}[d]*(-\text{ArcCosh}[c*x]/x) + (c*\text{Sqrt}[-1 + c^2*x^2]*\text{ArcTan}[\text{Sqrt}[-1 + c^2*x^2]])/(\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x])) - 2*\text{Sqrt}[d]*\text{Sqrt}[e]*(\text{ArcCosh}[c*x]/((-I)*\text{Sqrt}[d] + \text{Sqrt}[e]*x) + (c*\text{Log}[(2*e*(I*\text{Sqrt}[e] + c^2*\text{Sqrt}[d]*x - I*\text{Sqrt}[-(c^2*d) - e]*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]))/(c*\text{Sqrt}[-(c^2*d) - e]*(\text{Sqrt}[d] + I*\text{Sqrt}[e]*x)))]/\text{Sqrt}[-(c^2*d) - e]) + 2*\text{Sqrt}[d]*\text{Sqrt}[e]*(-\text{ArcCosh}[c*x]/(I*\text{Sqrt}[d] + \text{Sqrt}[e]*x) - (c*\text{Log}[(2*e*(-\text{Sqrt}[e] - I*c^2*\text{Sqrt}[d]*x + \text{Sqrt}[-(c^2*d) - e]*\text{Sqrt}[-1 + c*x]*\text{Sqrt}[1 + c*x]))/(c*\text{Sqrt}[-(c^2*d) - e]*(I*\text{Sqrt}[d] + \text{Sqrt}[e]*x)))]/\text{Sqrt}[-(c^2*d) - e]) - (3*I)*\text{Sqrt}[e]*(\text{ArcCosh}[c*x]*(-\text{ArcCosh}[c*x] + 2*(\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})]/(I*c*\text{Sqrt}[d] - \text{Sqrt}[-(c^2*d) - e])) + \text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e]))]) + 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/((-I)*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])] + 2*\text{PolyLog}[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e]))]) + (3*I)*\text{Sqrt}[e]*(\text{ArcCosh}[c*x]*(-\text{ArcCosh}[c*x] + 2*(\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/((-I)*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])) + \text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e]))]) + 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] - \text{Sqrt}[-(c^2*d) - e])] + 2*\text{PolyLog}[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(I*c*\text{Sqrt}[d] + \text{Sqrt}[-(c^2*d) - e])))]/(8*d^(5/2)) \end{aligned}$$

### 3.505.3 Rubi [A] (verified)

Time = 2.47 (sec) , antiderivative size = 846, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + \text{barccosh}(cx)}{x^2 (d + ex^2)^2} dx \\ & \quad \downarrow \text{6374} \\ & \int \left( -\frac{e(a + \text{barccosh}(cx))}{d^2 (d + ex^2)} + \frac{a + \text{barccosh}(cx)}{d^2 x^2} - \frac{e(a + \text{barccosh}(cx))}{d (d + ex^2)^2} \right) dx \\ & \quad \downarrow \text{2009} \end{aligned}$$

$$\begin{aligned}
& \frac{3\sqrt{e} \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right) (a + \operatorname{barccosh}(cx))}{4(-d)^{5/2}} + \\
& \frac{3\sqrt{e} \log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{c\sqrt{-d}-\sqrt{-dc^2-e}} + 1\right) (a + \operatorname{barccosh}(cx))}{4(-d)^{5/2}} - \\
& \frac{3\sqrt{e} \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right) (a + \operatorname{barccosh}(cx))}{4(-d)^{5/2}} + \\
& \frac{3\sqrt{e} \log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{\sqrt{-dc}+\sqrt{-dc^2-e}} + 1\right) (a + \operatorname{barccosh}(cx))}{4(-d)^{5/2}} - \frac{a + \operatorname{barccosh}(cx)}{d^2 x} + \frac{\sqrt{e}(a + \operatorname{barccosh}(cx))}{4d^2(\sqrt{-d} - \sqrt{ex})} - \\
& \frac{\sqrt{e}(a + \operatorname{barccosh}(cx))}{4d^2(\sqrt{ex} + \sqrt{-d})} + \frac{bc \arctan(\sqrt{cx-1}\sqrt{cx+1})}{d^2} - \frac{bc\sqrt{e} \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx-1}}}\right)}{2d^2\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{\sqrt{-dc}+\sqrt{e}}}} + \\
& \frac{bc\sqrt{e} \operatorname{arctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right)}{2d^2\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{\sqrt{-dc}+\sqrt{e}}}} + \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{4(-d)^{5/2}} - \\
& \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{4(-d)^{5/2}} + \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{4(-d)^{5/2}} - \\
& \frac{3b\sqrt{e} \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{4(-d)^{5/2}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x^2*(d + e*x^2)^2), x]`

```

output -((a + b*ArcCosh[c*x])/(d^2*x)) + (Sqrt[e]*(a + b*ArcCosh[c*x]))/(4*d^2*(S
qrt[-d] - Sqrt[e]*x)) - (Sqrt[e]*(a + b*ArcCosh[c*x]))/(4*d^2*(Sqrt[-d] +
Sqrt[e]*x)) + (b*c*ArcTan[Sqrt[-1 + c*x]*Sqrt[1 + c*x]])/d^2 - (b*c*Sqrt[e
]*ArcTanh[(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] + Sq
rt[e]]*Sqrt[-1 + c*x])])/(2*d^2*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d]
+ Sqrt[e]]) + (b*c*Sqrt[e]*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[1 + c
*x])/(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[-1 + c*x])])/(2*d^2*Sqrt[c*Sqrt[-d]
- Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]) - (3*Sqrt[e]*(a + b*ArcCosh[c*x])*L
og[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(4*(-d
)^(5/2)) + (3*Sqrt[e]*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x]
)/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(4*(-d)^(5/2)) - (3*Sqrt[e]*(a + b*A
rcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d)
- e])])/(4*(-d)^(5/2)) + (3*Sqrt[e]*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*
E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(4*(-d)^(5/2)) + (3*b
*Sqrt[e]*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d)
- e]))])/(4*(-d)^(5/2)) - (3*b*Sqrt[e]*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])
/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(4*(-d)^(5/2)) + (3*b*Sqrt[e]*PolyLog
[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e]))])/(4*(-d
)^(5/2)) - (3*b*Sqrt[e]*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] +
Sqrt[-(c^2*d) - e])])/(4*(-d)^(5/2))

```

### 3.505.3.1 Defintions of rubi rules used

```

rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

```

rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]

```



### 3.505.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 37.20 (sec) , antiderivative size = 900, normalized size of antiderivative = 1.06

method	result
parts	$a \left( -\frac{e \left( \frac{x}{2e x^2 + 2d} + \frac{3 \arctan\left(\frac{ex}{\sqrt{de}}\right)}{2\sqrt{de}} \right)}{d^2} - \frac{1}{d^2 x} \right) + bc \left( -\frac{\operatorname{arccosh}(cx)(3c^2 e x^2 + 2c^2 d)}{2cx d^2 (c^2 e x^2 + c^2 d)} + \frac{2 \arctan(cx + \sqrt{cx-1} \sqrt{c}}{d^2} \right)$
derivativedivides	Expression too large to display
default	Expression too large to display

input `int((a+b*arccosh(c*x))/x^2/(e*x^2+d)^2,x,method=_RETURNVERBOSE)`

output

```

a*(-e/d^2*(1/2*x/(e*x^2+d)+3/2/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2)))-1/d^2/
x)+b*c*(-1/2/c/x*arccosh(c*x)*(3*c^2*e*x^2+2*c^2*d)/d^2/(c^2*e*x^2+c^2*d)+
2/d^2*arctan(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))-3/16/d^3*e/c^2*sum((4*_R1^2*
c^2*d+_R1^2*e+e)/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)
^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1
)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/2*(-(2*c^2*d-2*(d*c^2*(c^2*d
+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctanh(e*(c*
x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(
1/2))/d^2/e^2+1/2*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*
d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/
((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2))/d^2/e^2-1/2*(-(2*c^2*d-2*
(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4
*d^2+2*c^2*d*e+(d*c^2*(c^2*d+e))^(1/2)*e)*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*
x+1)^(1/2))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))/d^2/(c^2*d+e
)/e^2-1/2*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(-2*(d*c^2*(c^2*
d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*arctan(e*
(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e
)^(1/2))/d^2/(c^2*d+e)/e^2+3/16/d^3*e/c^2*sum((_R1^2*e+4*c^2*d+e)/_R1/(_R1
^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1
)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(...

```

**3.505.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(e*x^2+d)^2,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e^2*x^6 + 2*d*e*x^4 + d^2*x^2), x)`

**3.505.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d + ex^2)^2} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x**2/(e*x**2+d)**2,x)`

output `Timed out`

**3.505.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d + ex^2)^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))/x^2/(e*x^2+d)^2,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.505.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d + ex^2)^2} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^2 x^2} dx$$

input `integrate((a+b*arccosh(c*x))/x^2/(e*x^2+d)^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)^2*x^2), x)`

**3.505.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^2 (d + ex^2)^2} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^2 (ex^2 + d)^2} dx$$

input `int((a + b*acosh(c*x))/(x^2*(d + e*x^2)^2),x)`

output `int((a + b*acosh(c*x))/(x^2*(d + e*x^2)^2), x)`

$$3.506 \quad \int \frac{x^5(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$$

3.506.1 Optimal result . . . . .	3732
3.506.2 Mathematica [C] (warning: unable to verify) . . . . .	3733
3.506.3 Rubi [A] (verified) . . . . .	3734
3.506.4 Maple [C] (warning: unable to verify) . . . . .	3736
3.506.5 Fricas [F] . . . . .	3737
3.506.6 Sympy [F(-1)] . . . . .	3738
3.506.7 Maxima [F] . . . . .	3738
3.506.8 Giac [F(-2)] . . . . .	3738
3.506.9 Mupad [F(-1)] . . . . .	3739

## 3.506.1 Optimal result

Integrand size = 21, antiderivative size = 737

$$\begin{aligned}
\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = & \frac{bcdx(1 - c^2x^2)}{8e^2(c^2d + e)\sqrt{-1 + cx}\sqrt{1 + cx}(d + ex^2)} \\
& - \frac{d^2(a + \operatorname{barccosh}(cx))}{4e^3(d + ex^2)^2} \\
& + \frac{d(a + \operatorname{barccosh}(cx))}{e^3(d + ex^2)} - \frac{(a + \operatorname{barccosh}(cx))^2}{2be^3} \\
& - \frac{bc\sqrt{d}\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{e^3\sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx}} \\
& + \frac{bc\sqrt{d}(2c^2d + e)\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{8e^3(c^2d + e)^{3/2}\sqrt{-1 + cx}\sqrt{1 + cx}} \\
& + \frac{(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^3} \\
& + \frac{(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^3} \\
& + \frac{(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^3} \\
& + \frac{(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^3} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^3} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{2e^3} \\
& + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^3} \\
& + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{2e^3}
\end{aligned}$$

output

```

-1/4*d^2*(a+b*arccosh(c*x))/e^3/(e*x^2+d)^2+d*(a+b*arccosh(c*x))/e^3/(e*x^
2+d)-1/2*(a+b*arccosh(c*x))^2/b/e^3+1/2*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-
1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^3+1/2*(
a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(
1/2)-(-c^2*d-e)^(1/2)))/e^3+1/2*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)
*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^3+1/2*(a+b*arcc
osh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c
^2*d-e)^(1/2)))/e^3+1/2*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(
1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^3+1/2*b*polylog(2,(c*x+(c*x-1)^(1/
2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^3+1/2*b*polyl
og(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(
1/2)))/e^3+1/2*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-
d)^(1/2)+(-c^2*d-e)^(1/2)))/e^3+1/8*b*c*d*x*(-c^2*x^2+1)/e^2/(c^2*d+e)/(e*
x^2+d)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/8*b*c*(2*c^2*d+e)*arctanh(x*(c^2*d+e)
^(1/2)/d^(1/2)/(c^2*x^2-1)^(1/2))*d^(1/2)*(c^2*x^2-1)^(1/2)/e^3/(c^2*d+e)^(
3/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*c*arctanh(x*(c^2*d+e)^(1/2)/d^(1/2)/(c
^2*x^2-1)^(1/2))*d^(1/2)*(c^2*x^2-1)^(1/2)/e^3/(c^2*d+e)^(1/2)/(c*x-1)^(1/
2)/(c*x+1)^(1/2)

```

### 3.506.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 6.63 (sec) , antiderivative size = 1097, normalized size of antiderivative = 1.49

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$$

$$= -\frac{4ad^2}{(d+ex^2)^2} + \frac{16ad}{d+ex^2} + 8a \log(d + ex^2) + b \left( -\frac{cd\sqrt{e}\sqrt{-1+cx}\sqrt{1+cx}}{(c^2d+e)(-i\sqrt{d}+\sqrt{ex})} - \frac{cd\sqrt{e}\sqrt{-1+cx}\sqrt{1+cx}}{(c^2d+e)(i\sqrt{d}+\sqrt{ex})} + \frac{7\sqrt{d}\operatorname{arccosh}(cx)}{\sqrt{d}-i\sqrt{ex}} - \frac{\operatorname{arccosh}(cx)}{\sqrt{d}} \right)$$

input `Integrate[(x^5*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

---

3.506.  $\int \frac{x^5(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$

output

```

((-4*a*d^2)/(d + e*x^2)^2 + (16*a*d)/(d + e*x^2) + 8*a*Log[d + e*x^2] + b*
(-(c*d*Sqrt[e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/((c^2*d + e)*((-I)*Sqrt[d] +
Sqrt[e]*x))) - (c*d*Sqrt[e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/((c^2*d + e)*(I
*Sqrt[d] + Sqrt[e]*x)) + (7*Sqrt[d]*ArcCosh[c*x])/(Sqrt[d] - I*Sqrt[e]*x)
- (d*ArcCosh[c*x])/(Sqrt[d] + I*Sqrt[e]*x)^2 + (7*Sqrt[d]*ArcCosh[c*x])/(S
qrt[d] + I*Sqrt[e]*x) + (d*ArcCosh[c*x])/(I*Sqrt[d] + Sqrt[e]*x)^2 - 8*Arc
Cosh[c*x]^2 + 8*ArcCosh[c*x]*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d]
- Sqrt[-(c^2*d) - e])] + 8*ArcCosh[c*x]*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/
((-I)*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] + 8*ArcCosh[c*x]*Log[1 - (Sqrt[e]*E
^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] + 8*ArcCosh[c*x]*Log[1
+ (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e])] - ((7*I)*c*
Sqrt[d]*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*d) - e]*Sqrt[-1
+ c*x]*Sqrt[1 + c*x])]/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*Sqrt[e]*x)))]/S
qrt[-(c^2*d) - e] + ((7*I)*c*Sqrt[d]*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x
+ Sqrt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])]/(c*Sqrt[-(c^2*d) - e]*
(I*Sqrt[d] + Sqrt[e]*x)))]/Sqrt[-(c^2*d) - e] - (c^3*d^(3/2)*Log[(e*Sqrt[c
^2*d + e]*((-I)*Sqrt[e] - c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*S
qrt[1 + c*x])]/(c^3*(d + I*Sqrt[d]*Sqrt[e]*x)))]/(c^2*d + e)^(3/2) + (c^3*
d^(3/2)*Log[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[e] + c^2*Sqrt[d]*x + Sqrt[c^2*d
+ e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])]/(c^3*(d - I*Sqrt[d]*Sqrt[e]*x)))]/(...

```

### 3.506.3 Rubi [A] (verified)

Time = 1.65 (sec) , antiderivative size = 737, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{d^2 x(a + \operatorname{barccosh}(cx))}{e^2 (d + ex^2)^3} - \frac{2dx(a + \operatorname{barccosh}(cx))}{e^2 (d + ex^2)^2} + \frac{x(a + \operatorname{barccosh}(cx))}{e^2 (d + ex^2)} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\begin{aligned}
& \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2e^3} + \frac{(a + \operatorname{arccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2e^3} + \\
& \frac{(a + \operatorname{arccosh}(cx)) \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2e^3} + \frac{(a + \operatorname{arccosh}(cx)) \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2e^3} - \\
& \frac{d^2(a + \operatorname{arccosh}(cx))}{4e^3(d + ex^2)^2} + \frac{d(a + \operatorname{arccosh}(cx))}{e^3(d + ex^2)} - \frac{(a + \operatorname{arccosh}(cx))^2}{2be^3} + \\
& \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2e^3} + \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2e^3} + \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2e^3} + \\
& \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2e^3} + \frac{bc\sqrt{d}\sqrt{c^2x^2 - 1}(2c^2d + e) \operatorname{arctanh}\left(\frac{x\sqrt{c^2d + e}}{\sqrt{d}\sqrt{c^2x^2 - 1}}\right)}{8e^3\sqrt{cx - 1}\sqrt{cx + 1}(c^2d + e)^{3/2}} - \\
& \frac{bc\sqrt{d}\sqrt{c^2x^2 - 1} \operatorname{arctanh}\left(\frac{x\sqrt{c^2d + e}}{\sqrt{d}\sqrt{c^2x^2 - 1}}\right)}{e^3\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{c^2d + e}} + \frac{bcdx(1 - c^2x^2)}{8e^2\sqrt{cx - 1}\sqrt{cx + 1}(c^2d + e)(d + ex^2)}
\end{aligned}$$

input `Int[(x^5*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

output `(b*c*d*x*(1 - c^2*x^2))/(8*e^2*(c^2*d + e)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(d + e*x^2) - (d^2*(a + b*ArcCosh[c*x]))/(4*e^3*(d + e*x^2)^2) + (d*(a + b*ArcCosh[c*x]))/(e^3*(d + e*x^2)) - (a + b*ArcCosh[c*x])^2/(2*b*e^3) - (b*c*Sqrt[d]*Sqrt[-1 + c^2*x^2]*ArcTanh[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2])])/(e^3*Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + (b*c*Sqrt[d]*(2*c^2*d + e)*Sqrt[-1 + c^2*x^2]*ArcTanh[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2])])/(8*e^3*(c^2*d + e)^(3/2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^3) + ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^3) + ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^3) + ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^3) + (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e]))]/(2*e^3) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*e^3) + (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e]))]/(2*e^3) + (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*e^3))`



**3.506.3.1 Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

**3.506.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 5.52 (sec) , antiderivative size = 3551, normalized size of antiderivative = 4.82

method	result	size
derivativedivides	Expression too large to display	3551
default	Expression too large to display	3551
parts	Expression too large to display	3556

input `int(x^5*(a+b*arccosh(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output  $1/c^6*(a*c^6*(-1/4*c^4*d^2/e^3/(c^2*e*x^2+c^2*d)^2+1/2/e^3*\ln(c^2*e*x^2+c^2*d)+1/e^3*d*c^2/(c^2*e*x^2+c^2*d))+b*c^6*((2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)/e^5/(c^2*d+e)*c^4*d^2*\ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^(1/2)-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*\operatorname{arccosh}(c*x)-1/4*(d*c^2*(c^2*d+e))^(1/2)/e/(c^2*d+e)^2/d/c^2*\operatorname{arccosh}(c*x)*\ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^(1/2)-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e))-1/2*(d*c^2*(c^2*d+e))^(1/2)/e^3/(c^2*d+e)^2*d*c^2*\operatorname{arccosh}(c*x)*\ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^(1/2)-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e))+3/2*(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)/e^4/(c^2*d+e)*c^2*d*\ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^(1/2)-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*\operatorname{arccosh}(c*x)-(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^4*d^2/e^5/(c^4*d^2+2*c^2*d*e+e^2)*\ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^(1/2)-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*\operatorname{arccosh}(c*x)-2*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^2*d/e^4/(c^4*d^2+2*c^2*d*e+e^2)*\ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^(1/2)-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*\operatorname{arccosh}(c*x)-1/4*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)/c^2/d/e^2/(c^4*d^2+2*c^2*d*e+e^2)*\ln(1-e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^(1/2)-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)-e))*\operatorname{arccosh}(c*x)+(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^4*d^2/e^5/(c^4*d^2+2*c^2*d*e+e^2)...$

### 3.506.5 Fracas [F]

$$\int \frac{x^5(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)x^5}{(ex^2 + d)^3} dx$$

input `integrate(x^5*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*x^5*arccosh(c*x) + a*x^5)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**3.506.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

```
input integrate(x**5*(a+b*acosh(c*x))/(e*x**2+d)**3,x)
```

```
output Timed out
```

**3.506.7 Maxima [F]**

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^5}{(ex^2 + d)^3} dx$$

```
input integrate(x^5*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="maxima")
```

```
output 1/4*a*((4*d*e*x^2 + 3*d^2)/(e^5*x^4 + 2*d*e^4*x^2 + d^2*e^3) + 2*log(e*x^2
+ d)/e^3) + b*integrate(x^5*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(e^3*x
^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)
```

**3.506.8 Giac [F(-2)]**

Exception generated.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: TypeError}$$

```
input integrate(x^5*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="giac")
```

```
output Exception raised: TypeError >> an error occurred running a Giac command:IN
PUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const
index_m & i,const vecteur & l) Error: Bad Argument Value
```

**3.506.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^5(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{x^5(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^3} dx$$

input `int((x^5*(a + b*acosh(c*x)))/(d + e*x^2)^3,x)`output `int((x^5*(a + b*acosh(c*x)))/(d + e*x^2)^3, x)`

### 3.507 $\int \frac{x^3(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$

3.507.1 Optimal result . . . . .	3740
3.507.2 Mathematica [A] (verified) . . . . .	3741
3.507.3 Rubi [A] (verified) . . . . .	3741
3.507.4 Maple [B] (verified) . . . . .	3745
3.507.5 Fricas [B] (verification not implemented) . . . . .	3746
3.507.6 Sympy [F(-1)] . . . . .	3746
3.507.7 Maxima [F] . . . . .	3747
3.507.8 Giac [F(-2)] . . . . .	3747
3.507.9 Mupad [F(-1)] . . . . .	3748

#### 3.507.1 Optimal result

Integrand size = 21, antiderivative size = 231

$$\int \frac{x^3(a + b\operatorname{arccosh}(cx))}{(d + ex^2)^3} dx = -\frac{bcx(1 - c^2x^2)}{8e(c^2d + e)\sqrt{-1 + cx}\sqrt{1 + cx}(d + ex^2)} + \frac{x^4(a + b\operatorname{arccosh}(cx))}{4d(d + ex^2)^2} - \frac{b\sqrt{1 - c^2x^2} \arcsin(cx)}{4de^2\sqrt{-1 + cx}\sqrt{1 + cx}} + \frac{bc(2c^2d + 3e)\sqrt{1 - c^2x^2} \arctan\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{1-c^2x^2}}\right)}{8\sqrt{de^2}(c^2d + e)^{3/2}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
1/4*x^4*(a+b*arccosh(c*x))/d/(e*x^2+d)^2-1/8*b*c*x*(-c^2*x^2+1)/e/(c^2*d+e)/(e*x^2+d)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-1/4*b*arcsin(c*x)*(-c^2*x^2+1)^(1/2)/d/e^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/8*b*c*(2*c^2*d+3*e)*arctan(x*(c^2*d+e)^(1/2)/d^(1/2)/(-c^2*x^2+1)^(1/2))*(-c^2*x^2+1)^(1/2)/e^2/(c^2*d+e)^(3/2)/d^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

### 3.507.2 Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 192, normalized size of antiderivative = 0.83

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$$

$$= \frac{\frac{bcex\sqrt{-1+cx}\sqrt{1+cx}(d+ex^2)}{c^2d+e} - 2a(d+2ex^2)}{(d+ex^2)^2} - \frac{2b(d+2ex^2)\operatorname{arccosh}(cx)}{(d+ex^2)^2} - \frac{bc(2c^2d+3e)\sqrt{-1+cx}\sqrt{1+cx}\arctan\left(\frac{\sqrt{-c^2d-ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{\sqrt{d}(-c^2d-e)^{3/2}\sqrt{-1+c^2x^2}}$$

$$= \frac{\hspace{15em}}{8e^2}$$

input `Integrate[(x^3*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

output `((b*c*e*x*sqrt[-1 + c*x]*sqrt[1 + c*x]*(d + e*x^2))/(c^2*d + e) - 2*a*(d + 2*e*x^2))/(d + e*x^2)^2 - (2*b*(d + 2*e*x^2)*ArcCosh[c*x])/(d + e*x^2)^2 - (b*c*(2*c^2*d + 3*e)*sqrt[-1 + c*x]*sqrt[1 + c*x]*ArcTan[(sqrt[-(c^2*d - e)]*x)/(sqrt[d]*sqrt[-1 + c^2*x^2])])/(sqrt[d]*(-(c^2*d) - e)^(3/2)*sqrt[-1 + c^2*x^2])/(8*e^2)`

### 3.507.3 Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 220, normalized size of antiderivative = 0.95, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.476$ , Rules used = {6373, 27, 2038, 372, 25, 398, 224, 219, 291, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$$

$$\downarrow \text{6373}$$

$$\frac{x^4(a + \operatorname{barccosh}(cx))}{4d(d + ex^2)^2} - bc \int \frac{x^4}{4d\sqrt{cx-1}\sqrt{cx+1}(ex^2+d)^2} dx$$

$$\downarrow \text{27}$$

$$\frac{x^4(a + \operatorname{barccosh}(cx))}{4d(d + ex^2)^2} - \frac{bc \int \frac{x^4}{\sqrt{cx-1}\sqrt{cx+1}(ex^2+d)^2} dx}{4d}$$

$$\downarrow \text{2038}$$

---

3.507.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$

$$\begin{aligned}
& \frac{x^4(a + \operatorname{arccosh}(cx))}{4d(d+ex^2)^2} - \frac{bc\sqrt{c^2x^2-1} \int \frac{x^4}{\sqrt{c^2x^2-1}(ex^2+d)^2} dx}{4d\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{372} \\
& \frac{x^4(a + \operatorname{arccosh}(cx))}{4d(d+ex^2)^2} - \frac{bc\sqrt{c^2x^2-1} \left( \frac{\int -\frac{d-2(dc^2+e)x^2}{\sqrt{c^2x^2-1}(ex^2+d)} dx}{2e(c^2d+e)} - \frac{dx\sqrt{c^2x^2-1}}{2e(c^2d+e)(d+ex^2)} \right)}{4d\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{25} \\
& \frac{x^4(a + \operatorname{arccosh}(cx))}{4d(d+ex^2)^2} - \frac{bc\sqrt{c^2x^2-1} \left( -\frac{\int \frac{d-2(dc^2+e)x^2}{\sqrt{c^2x^2-1}(ex^2+d)} dx}{2e(c^2d+e)} - \frac{dx\sqrt{c^2x^2-1}}{2e(c^2d+e)(d+ex^2)} \right)}{4d\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{398} \\
& \frac{x^4(a + \operatorname{arccosh}(cx))}{4d(d+ex^2)^2} - \frac{bc\sqrt{c^2x^2-1} \left( -\frac{d(2c^2d+3e) \int \frac{1}{\sqrt{c^2x^2-1}(ex^2+d)} dx}{e} - \frac{2(c^2d+e) \int \frac{1}{\sqrt{c^2x^2-1}} dx}{e} - \frac{dx\sqrt{c^2x^2-1}}{2e(c^2d+e)(d+ex^2)} \right)}{4d\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{224} \\
& \frac{x^4(a + \operatorname{arccosh}(cx))}{4d(d+ex^2)^2} - \frac{bc\sqrt{c^2x^2-1} \left( -\frac{d(2c^2d+3e) \int \frac{1}{\sqrt{c^2x^2-1}(ex^2+d)} dx}{e} - \frac{2(c^2d+e) \int \frac{1}{1-\frac{c^2x^2}{c^2x^2-1}} d \frac{x}{\sqrt{c^2x^2-1}}}{e} - \frac{dx\sqrt{c^2x^2-1}}{2e(c^2d+e)(d+ex^2)} \right)}{4d\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{219} \\
& \frac{x^4(a + \operatorname{arccosh}(cx))}{4d(d+ex^2)^2} - \frac{bc\sqrt{c^2x^2-1} \left( -\frac{d(2c^2d+3e) \int \frac{1}{\sqrt{c^2x^2-1}(ex^2+d)} dx}{e} - \frac{2\operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2-1}}\right)(c^2d+e)}{ce} - \frac{dx\sqrt{c^2x^2-1}}{2e(c^2d+e)(d+ex^2)} \right)}{4d\sqrt{cx-1}\sqrt{cx+1}} \\
& \quad \downarrow \text{291}
\end{aligned}$$

---

3.507.  $\int \frac{x^3(a+\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$

$$\begin{array}{c}
\frac{x^4(a + \operatorname{barccosh}(cx))}{4d(d + ex^2)^2} - \\
bc\sqrt{c^2x^2 - 1} \left( -\frac{\frac{d(2c^2d+3e) \int \frac{1}{d - \frac{(dc^2+e)x^2}{c^2x^2-1}} dx \frac{x}{\sqrt{c^2x^2-1}}}{e} \operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2-1}}\right)(c^2d+e)}{2e(c^2d+e)} - \frac{dx\sqrt{c^2x^2-1}}{2e(c^2d+e)(d+ex^2)} \right) \\
\hline
4d\sqrt{cx - 1}\sqrt{cx + 1} \\
\downarrow \text{221} \\
\frac{x^4(a + \operatorname{barccosh}(cx))}{4d(d + ex^2)^2} - \\
bc\sqrt{c^2x^2 - 1} \left( -\frac{\frac{\sqrt{d}(2c^2d+3e) \operatorname{arctanh}\left(\frac{x\sqrt{c^2d+e}}{\sqrt{d}\sqrt{c^2x^2-1}}\right) \operatorname{arctanh}\left(\frac{cx}{\sqrt{c^2x^2-1}}\right)(c^2d+e)}{e\sqrt{c^2d+e}}}{2e(c^2d+e)} - \frac{dx\sqrt{c^2x^2-1}}{2e(c^2d+e)(d+ex^2)} \right) \\
\hline
4d\sqrt{cx - 1}\sqrt{cx + 1}
\end{array}$$

input `Int[(x^3*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

output `(x^4*(a + b*ArcCosh[c*x]))/(4*d*(d + e*x^2)^2) - (b*c*sqrt[-1 + c^2*x^2]*( -1/2*(d*x*sqrt[-1 + c^2*x^2])/(e*(c^2*d + e)*(d + e*x^2)) - ((-2*(c^2*d + e)*ArcTanh[(c*x)/sqrt[-1 + c^2*x^2]])/(c*e) + (sqrt[d]*(2*c^2*d + 3*e)*ArcTanh[(sqrt[c^2*d + e]*x)/(sqrt[d]*sqrt[-1 + c^2*x^2])])/(e*sqrt[c^2*d + e]))/(2*e*(c^2*d + e)))/(4*d*sqrt[-1 + c*x]*sqrt[1 + c*x])`

### 3.507.3.1 Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

---

3.507.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$



- rule 224 `Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`
- rule 291 `Int[1/(Sqrt[(a_) + (b_)*(x_)^2]*((c_) + (d_)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 372 `Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_), x_Symbol] := Simp[(-a)*e^3*(e*x)^(m - 3)*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*b*(b*c - a*d)*(p + 1))), x] + Simp[e^4/(2*b*(b*c - a*d)*(p + 1)) Int[(e*x)^(m - 4)*(a + b*x^2)^(p + 1)*(c + d*x^2)^q*Simp[a*c*(m - 3) + (a*d*(m + 2*q - 1) + 2*b*c*(p + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && GtQ[m, 3] && IntBinomialQ[a, b, c, d, e, m, 2, p, q, x]`
- rule 398 `Int[((e_) + (f_)*(x_)^2)/(((a_) + (b_)*(x_)^2)*Sqrt[(c_) + (d_)*(x_)^2]), x_Symbol] := Simp[f/b Int[1/Sqrt[c + d*x^2], x], x] + Simp[(b*e - a*f)/b Int[1/((a + b*x^2)*Sqrt[c + d*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`
- rule 2038 `Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[(a1 + b1*x^(n/2))^FracPart[p]*((a2 + b2*x^(n/2))^FracPart[p]/(a1*a2 + b1*b2*x^n)^FracPart[p]) Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && !(EqQ[n, 2] && IGtQ[q, 0])`
- rule 6373 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x]] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.507.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1168 vs.  $2(197) = 394$ .

Time = 0.55 (sec) , antiderivative size = 1169, normalized size of antiderivative = 5.06

method	result	size
parts	Expression too large to display	1169
derivativedivides	Expression too large to display	1199
default	Expression too large to display	1199

input `int(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output

```
a*(1/4*d/e^2/(e*x^2+d)^2-1/2/e^2/(e*x^2+d))+b/c^4*(1/4*c^8*arccosh(c*x)/e^
2*d/(c^2*e*x^2+c^2*d)^2-1/2*c^6*arccosh(c*x)/e^2/(c^2*e*x^2+c^2*d)+1/16*c^
6*e*(2*ln(-2*(-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c
*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^6*x^2*d^2*e+2*ln(-2*(-(c^2*d+e)/e)^(1/
2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c
^6*d^3-2*ln(2*((-c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c
*x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*c^6*d^2*e*x^2-2*ln(2*((-c^2*d+e)/e)^(1/2)
*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*c^6
*d^3+5*ln(-2*(-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c
*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^4*x^2*d*e^2+5*ln(-2*(-(c^2*d+e)/e)^(1/
2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c
^4*d^2*e-5*ln(2*((-c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)
*c*x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*c^4*d*e^2*x^2-5*ln(2*((-c^2*d+e)/e)^(1/
2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*c
^4*d^2*e+2*c^3*d*e*(-c^2*d*e)^(1/2)*(c^2*x^2-1)^(1/2)*(-c^2*d+e)/e)^(1/2)
*x+3*ln(-2*(-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x
+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^2*x^2*e^3+3*ln(-2*(-(c^2*d+e)/e)^(1/2)*(
c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^2*d
*e^2-3*ln(2*((-c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x
-e)/(e*c*x-(-c^2*d*e)^(1/2)))*e^3*c^2*x^2-3*ln(2*((-c^2*d+e)/e)^(1/2)*...
```

**3.507.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 577 vs.  $2(196) = 392$ .

Time = 0.36 (sec) , antiderivative size = 1217, normalized size of antiderivative = 5.27

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `[-1/16*(2*(2*a - b)*c^4*d^4 + 2*(4*a - b)*c^2*d^3*e - 4*(b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4*log(c*x + sqrt(c^2*x^2 - 1)) + 4*a*d^2*e^2 - 2*(b*c^4*d^2*e^2 + b*c^2*d*e^3)*x^4 + 4*((2*a - b)*c^4*d^3*e + (4*a - b)*c^2*d^2*e^2 + 2*a*d*e^3)*x^2 - (2*b*c^3*d^3 + 3*b*c*d^2*e + (2*b*c^3*d*e^2 + 3*b*c*e^3)*x^4 + 2*(2*b*c^3*d^2*e + 3*b*c*d*e^2)*x^2)*sqrt(c^2*d^2 + d*e)*log(-(2*c^2*d^2 - (4*c^4*d^2 + 4*c^2*d*e + e^2)*x^2 + d*e - 2*sqrt(c^2*d^2 + d*e))*((2*c^3*d + c*e)*x^2 - c*d) - 2*sqrt(c^2*x^2 - 1)*(sqrt(c^2*d^2 + d*e))*(2*c^2*d + e)*x + 2*(c^3*d^2 + c*d*e)*x))/(e*x^2 + d) - 4*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2*e^2 + (b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^4*d^3*e + 2*b*c^2*d^2*e^2 + b*d*e^3)*x^2)*log(-c*x + sqrt(c^2*x^2 - 1)) - 2*sqrt(c^2*x^2 - 1)*((b*c^3*d^2*e^2 + b*c*d*e^3)*x^3 + (b*c^3*d^3*e + b*c*d^2*e^2)*x))/(c^4*d^5*e^2 + 2*c^2*d^4*e^3 + d^3*e^4 + (c^4*d^3*e^4 + 2*c^2*d^2*e^5 + d*e^6)*x^4 + 2*(c^4*d^4*e^3 + 2*c^2*d^3*e^4 + d^2*e^5)*x^2), -1/8*((2*a - b)*c^4*d^4 + (4*a - b)*c^2*d^3*e - 2*(b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4*log(c*x + sqrt(c^2*x^2 - 1)) + 2*a*d^2*e^2 - (b*c^4*d^2*e^2 + b*c^2*d*e^3)*x^4 + 2*((2*a - b)*c^4*d^3*e + (4*a - b)*c^2*d^2*e^2 + 2*a*d*e^3)*x^2 - (2*b*c^3*d^3 + 3*b*c*d^2*e + (2*b*c^3*d*e^2 + 3*b*c*e^3)*x^4 + 2*(2*b*c^3*d^2*e + 3*b*c*d*e^2)*x^2)*sqrt(-c^2*d^2 - d*e)*arctan((sqrt(-c^2*d^2 - d*e)*sqrt(c^2*x^2 - 1))*e*x - sqrt(-c^2*d^2 - d*e)*(c*e*x^2 + c*d))/(c^2*d^2 + d*e)) - 2*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2...`

**3.507.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x**3*(a+b*acosh(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

---

3.507.  $\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$

## 3.507.7 Maxima [F]

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^3}{(ex^2 + d)^3} dx$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `-1/8*b*((c^4*d + 2*c^2*e)*log(e*x^2 + d)/(c^4*d^2*e^2 + 2*c^2*d*e^3 + e^4) + (c^4*d^3 + c^2*d^2*e + (c^4*d^2*e + c^2*d*e^2)*x^2 + 2*(c^4*d^3 + 2*c^2*d^2*e + d^2*e + 2*(c^4*d^2*e + 2*c^2*d*e^2 + e^3)*x^2)*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)) - (c^4*d^3 + 2*c^2*d^2*e + (c^4*d*e^2 + 2*c^2*e^3)*x^4 + 2*(c^4*d^2*e + 2*c^2*d*e^2)*x^2)*log(c*x + 1) - (c^4*d^3 + 2*c^2*d^2*e + (c^4*d*e^2 + 2*c^2*e^3)*x^4 + 2*(c^4*d^2*e + 2*c^2*d*e^2)*x^2)*log(c*x - 1))/(c^4*d^4*e^2 + 2*c^2*d^3*e^3 + d^2*e^4 + (c^4*d^2*e^4 + 2*c^2*d*e^5 + e^6)*x^4 + 2*(c^4*d^3*e^3 + 2*c^2*d^2*e^4 + d*e^5)*x^2) + 8*integrate(1/4*(2*c*e*x^2 + c*d)/(c^3*e^4*x^7 - c*d^2*e^2*x + (2*c^3*d*e^3 - c*e^4)*x^5 + (c^3*d^2*e^2 - 2*c*d*e^3)*x^3 + (c^2*e^4*x^6 + (2*c^2*d*e^3 - e^4)*x^4 - d^2*e^2 + (c^2*d^2*e^2 - 2*d*e^3)*x^2)*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1))), x) - 1/4*(2*e*x^2 + d)*a/(e^4*x^4 + 2*d*e^3*x^2 + d^2*e^2)`

## 3.507.8 Giac [F(-2)]

Exception generated.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: TypeError}$$

input `integrate(x^3*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value`

**3.507.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^3(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{x^3(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^3} dx$$

input `int((x^3*(a + b*acosh(c*x)))/(d + e*x^2)^3,x)`output `int((x^3*(a + b*acosh(c*x)))/(d + e*x^2)^3, x)`

**3.508** 
$$\int \frac{x(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$$

3.508.1 Optimal result . . . . .	3749
3.508.2 Mathematica [A] (verified) . . . . .	3749
3.508.3 Rubi [A] (verified) . . . . .	3750
3.508.4 Maple [B] (verified) . . . . .	3752
3.508.5 Fricas [B] (verification not implemented) . . . . .	3753
3.508.6 Sympy [F(-1)] . . . . .	3753
3.508.7 Maxima [F] . . . . .	3754
3.508.8 Giac [F] . . . . .	3754
3.508.9 Mupad [F(-1)] . . . . .	3754

**3.508.1 Optimal result**

Integrand size = 19, antiderivative size = 177

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx = \frac{bcx(1 - c^2x^2)}{8d(c^2d + e)\sqrt{-1 + cx}\sqrt{1 + cx}(d + ex^2)} - \frac{a + \operatorname{arccosh}(cx)}{4e(d + ex^2)^2} + \frac{bc(2c^2d + e)\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{8d^{3/2}e(c^2d + e)^{3/2}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output `1/4*(-a-b*arccosh(c*x))/e/(e*x^2+d)^2+1/8*b*c*x*(-c^2*x^2+1)/d/(c^2*d+e)/(e*x^2+d)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/8*b*c*(2*c^2*d+e)*arctanh(x*(c^2*d+e)^(1/2)/d^(1/2)/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/d^(3/2)/e/(c^2*d+e)^(3/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)`

**3.508.2 Mathematica [A] (verified)**

Time = 0.83 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.03

$$\int \frac{x(a + \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx = \frac{1}{8} \left( -\frac{\frac{2a}{e} + \frac{bcx\sqrt{-1+cx}\sqrt{1+cx}(d+ex^2)}{d(c^2d+e)}}{(d + ex^2)^2} - \frac{2\operatorname{arccosh}(cx)}{e(d + ex^2)^2} - \frac{bc(2c^2d + e)\sqrt{-1 + cx}\sqrt{1 + cx}\operatorname{arctan}\left(\frac{\sqrt{-c^2d-ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{d^{3/2}(-c^2d - e)^{3/2}e\sqrt{-1 + c^2x^2}} \right)$$

input `Integrate[(x*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

output 
$$\frac{-(((2a)/e + (b*c*x*\sqrt{-1 + c*x}*\sqrt{1 + c*x}*(d + e*x^2))/(d*(c^2*d + e)))/(d + e*x^2)^2 - (2*b*ArcCosh[c*x])/(e*(d + e*x^2)^2 - (b*c*(2*c^2*d + e)*\sqrt{-1 + c*x}*\sqrt{1 + c*x}*\text{ArcTan}[(\sqrt{-c^2*d} - e)*x]/(\sqrt{d}*\sqrt{-1 + c^2*x^2}]))/(d^{(3/2)*(-c^2*d - e)^{(3/2)*e*\sqrt{-1 + c^2*x^2}})/8$$

### 3.508.3 Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 166, normalized size of antiderivative = 0.94, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$ , Rules used = {6372, 648, 296, 291, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x(a + \text{barccosh}(cx))}{(d + ex^2)^3} dx \\ & \quad \downarrow 6372 \\ & \frac{bc \int \frac{1}{\sqrt{cx-1}\sqrt{cx+1}(ex^2+d)^2} dx}{4e} - \frac{a + \text{barccosh}(cx)}{4e(d + ex^2)^2} \\ & \quad \downarrow 648 \\ & \frac{bc\sqrt{c^2x^2 - 1} \int \frac{1}{\sqrt{c^2x^2-1}(ex^2+d)^2} dx}{4e\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{a + \text{barccosh}(cx)}{4e(d + ex^2)^2} \\ & \quad \downarrow 296 \\ & \frac{bc\sqrt{c^2x^2 - 1} \left( \frac{(2c^2d+e) \int \frac{1}{\sqrt{c^2x^2-1}(ex^2+d)} dx}{2d(c^2d+e)} - \frac{ex\sqrt{c^2x^2-1}}{2d(c^2d+e)(d+ex^2)} \right)}{4e\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{a + \text{barccosh}(cx)}{4e(d + ex^2)^2} \\ & \quad \downarrow 291 \\ & \frac{bc\sqrt{c^2x^2 - 1} \left( \frac{(2c^2d+e) \int \frac{1}{d - \frac{(dc^2+e)x^2}{c^2x^2-1}} d \frac{x}{\sqrt{c^2x^2-1}}}{2d(c^2d+e)} - \frac{ex\sqrt{c^2x^2-1}}{2d(c^2d+e)(d+ex^2)} \right)}{4e\sqrt{cx - 1}\sqrt{cx + 1}} - \frac{a + \text{barccosh}(cx)}{4e(d + ex^2)^2} \\ & \quad \downarrow 221 \end{aligned}$$

---

3.508.  $\int \frac{x(a + \text{barccosh}(cx))}{(d + ex^2)^3} dx$

$$\frac{bc\sqrt{c^2x^2-1} \left( \frac{(2c^2d+e)\operatorname{arctanh}\left(\frac{x\sqrt{c^2d+e}}{\sqrt{d}\sqrt{c^2x^2-1}}\right)}{2d^{3/2}(c^2d+e)^{3/2}} - \frac{ex\sqrt{c^2x^2-1}}{2d(c^2d+e)(d+ex^2)} \right)}{4e\sqrt{cx-1}\sqrt{cx+1}} - \frac{a + \operatorname{barccosh}(cx)}{4e(d+ex^2)^2}$$

input `Int[(x*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

output `-1/4*(a + b*ArcCosh[c*x])/(e*(d + e*x^2)^2) + (b*c*Sqrt[-1 + c^2*x^2]*(-1/2*(e*x*Sqrt[-1 + c^2*x^2]))/(d*(c^2*d + e)*(d + e*x^2)) + ((2*c^2*d + e)*ArcTanh[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2])])/(2*d^(3/2)*(c^2*d + e)^(3/2)))/(4*e*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.508.3.1 Defintions of rubi rules used

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 296 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))], x] + Simp[(b*c + 2*(p + 1)*(b*c - a*d))/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && EqQ[2*(p + q + 2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]`

rule 648 `Int[((c_) + (d_.)*(x_)^(m_.))*((e_) + (f_.)*(x_)^(n_.))*((a_.) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(c + d*x)^FracPart[m]*((e + f*x)^FracPart[m]/(c*e + d*f*x^2)^FracPart[m]) Int[(c*e + d*f*x^2)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[m, n] && EqQ[d*e + c*f, 0] && !(EqQ[p, 2] && LtQ[m, -1])`



```
rule 6372 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x
_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcCosh[c*x])/(2*e*(p + 1))),
x] - Simp[b*(c/(2*e*(p + 1))) Int[(d + e*x^2)^(p + 1)/(Sqrt[1 + c*x]*Sqrt
[-1 + c*x]), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[c^2*d + e, 0] &&
NeQ[p, -1]
```

### 3.508.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1125 vs.  $2(154) = 308$ .

Time = 0.62 (sec) , antiderivative size = 1126, normalized size of antiderivative = 6.36

method	result	size
parts	Expression too large to display	1126
derivativedivides	Expression too large to display	1149
default	Expression too large to display	1149

```
input int(x*(a+b*arccosh(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)
```

```
output -1/4*a/e/(e*x^2+d)^2+b/c^2*(-1/4*c^6/e/(c^2*e*x^2+c^2*d)^2*arccosh(c*x)+1/
16*c^4*e^2*(2*ln(-2*(-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(
1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^6*x^2*d^2*e+2*ln(-2*(-(c^2*d+e)
/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1
/2)))*c^6*d^3-2*ln(2*((-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(
1/2)*c*x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*c^6*d^2*e*x^2-2*ln(2*((-(c^2*d+e)/e)
^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x-e)/(e*c*x-(-c^2*d*e)^(1/2
)))*c^6*d^3+3*ln(-2*(-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(
1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^4*x^2*d*e^2+3*ln(-2*(-(c^2*d+e)
/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1
/2)))*c^4*d^2*e-3*ln(2*((-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)
^(1/2)*c*x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*c^4*d*e^2*x^2-3*ln(2*((-(c^2*d+e)
/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x-e)/(e*c*x-(-c^2*d*e)^(1
/2)))*c^4*d^2*e-2*c^3*d*e*(-c^2*d*e)^(1/2)*(c^2*x^2-1)^(1/2)*(-c^2*d+e)/e)
^(1/2)*x+ln(-2*(-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)
)*c*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^2*x^2*e^3+ln(-2*(-(c^2*d+e)/e)^(1/2)
*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x+e)/(e*c*x+(-c^2*d*e)^(1/2)))*c^
2*d*e^2-ln(2*((-(c^2*d+e)/e)^(1/2)*(c^2*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*
x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*e^3*c^2*x^2-ln(2*((-(c^2*d+e)/e)^(1/2)*(c^2
*x^2-1)^(1/2)*e+(-c^2*d*e)^(1/2)*c*x-e)/(e*c*x-(-c^2*d*e)^(1/2)))*c^2*d...
```

**3.508.5 Fracas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 584 vs.  $2(150) = 300$ .

Time = 0.38 (sec) , antiderivative size = 1233, normalized size of antiderivative = 6.97

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Too large to display}$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `[-1/16*(2*(2*a + b)*c^4*d^4 + 2*(4*a + b)*c^2*d^3*e + 4*a*d^2*e^2 + 2*(b*c^4*d^2*e^2 + b*c^2*d*e^3)*x^4 + 4*(b*c^4*d^3*e + b*c^2*d^2*e^2)*x^2 - (2*b*c^3*d^3 + b*c*d^2*e + (2*b*c^3*d*e^2 + b*c*e^3)*x^4 + 2*(2*b*c^3*d^2*e + b*c*d*e^2)*x^2)*sqrt(c^2*d^2 + d*e)*log(-(2*c^2*d^2 - (4*c^4*d^2 + 4*c^2*d*e + e^2)*x^2 + d*e - 2*sqrt(c^2*d^2 + d*e))*((2*c^3*d + c*e)*x^2 - c*d) - 2*sqrt(c^2*x^2 - 1)*(sqrt(c^2*d^2 + d*e)*(2*c^2*d + e)*x + 2*(c^3*d^2 + c*d*e)*x))/(e*x^2 + d) - 4*((b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^4*d^3*e + 2*b*c^2*d^2*e^2 + b*d*e^3)*x^2)*log(c*x + sqrt(c^2*x^2 - 1)) - 4*(b*c^4*d^4 + 2*b*c^2*d^3*e + b*d^2*e^2 + (b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^4*d^3*e + 2*b*c^2*d^2*e^2 + b*d*e^3)*x^2)*log(-c*x + sqrt(c^2*x^2 - 1)) + 2*sqrt(c^2*x^2 - 1)*((b*c^3*d^2*e^2 + b*c*d*e^3)*x^3 + (b*c^3*d^3*e + b*c*d^2*e^2)*x))/(c^4*d^6*e + 2*c^2*d^5*e^2 + d^4*e^3 + (c^4*d^4*e^3 + 2*c^2*d^3*e^4 + d^2*e^5)*x^4 + 2*(c^4*d^5*e^2 + 2*c^2*d^4*e^3 + d^3*e^4)*x^2), -1/8*((2*a + b)*c^4*d^4 + (4*a + b)*c^2*d^3*e + 2*a*d^2*e^2 + (b*c^4*d^2*e^2 + b*c^2*d*e^3)*x^4 + 2*(b*c^4*d^3*e + b*c^2*d^2*e^2)*x^2 - (2*b*c^3*d^3 + b*c*d^2*e + (2*b*c^3*d*e^2 + b*c*e^3)*x^4 + 2*(2*b*c^3*d^2*e + b*c*d*e^2)*x^2)*sqrt(-c^2*d^2 - d*e)*arctan((sqrt(-c^2*d^2 - d*e)*sqrt(c^2*x^2 - 1)*e*x - sqrt(-c^2*d^2 - d*e)*(c*e*x^2 + c*d))/(c^2*d^2 + d*e)) - 2*((b*c^4*d^2*e^2 + 2*b*c^2*d*e^3 + b*e^4)*x^4 + 2*(b*c^4*d^3*e + 2*b*c^2*d^2*e^2 + b*d*e^3)*x^2)*log(c*x + sqrt(c^2*x^2 - 1)) - ...`

**3.508.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x*(a+b*acosh(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

---

3.508.  $\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$

**3.508.7 Maxima [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(ex^2 + d)^3} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `-1/8*(c^4*log(e*x^2 + d)/(c^4*d^2*e + 2*c^2*d*e^2 + e^3) + 8*c*integrate(1/4/(c^3*e^3*x^7 + (2*c^3*d*e^2 - c*e^3)*x^5 - c*d^2*e*x + (c^3*d^2*e - 2*c*d*e^2)*x^3 + (c^2*e^3*x^6 + (2*c^2*d*e^2 - e^3)*x^4 - d^2*e + (c^2*d^2*e - 2*d*e^2)*x^2)*e^(1/2*log(c*x + 1) + 1/2*log(c*x - 1))), x) - (c^4*d^2 + c^2*d*e + (c^4*d*e + c^2*e^2)*x^2 - 2*(c^4*d^2 + 2*c^2*d*e + e^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + (c^4*e^2*x^4 + 2*c^4*d*e*x^2 + c^4*d^2)*log(c*x + 1) + (c^4*e^2*x^4 + 2*c^4*d*e*x^2 + c^4*d^2)*log(c*x - 1))/(c^4*d^4*e + 2*c^2*d^3*e^2 + d^2*e^3 + (c^4*d^2*e^3 + 2*c^2*d*e^4 + e^5)*x^4 + 2*(c^4*d^3*e^2 + 2*c^2*d^2*e^3 + d*e^4)*x^2))*b - 1/4*a/(e^3*x^4 + 2*d*e^2*x^2 + d^2*e)`

**3.508.8 Giac [F]**

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x}{(ex^2 + d)^3} dx$$

input `integrate(x*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x/(e*x^2 + d)^3, x)`

**3.508.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{x(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^3} dx$$

input `int((x*(a + b*acosh(c*x)))/(d + e*x^2)^3,x)`

output `int((x*(a + b*acosh(c*x)))/(d + e*x^2)^3, x)`

---

3.508.  $\int \frac{x(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$

$$3.509 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{x(d+ex^2)^3} dx$$

3.509.1 Optimal result . . . . .	3756
3.509.2 Mathematica [C] (warning: unable to verify) . . . . .	3757
3.509.3 Rubi [A] (verified) . . . . .	3759
3.509.4 Maple [C] (warning: unable to verify) . . . . .	3761
3.509.5 Fricas [F] . . . . .	3762
3.509.6 Sympy [F(-1)] . . . . .	3763
3.509.7 Maxima [F] . . . . .	3763
3.509.8 Giac [F] . . . . .	3763
3.509.9 Mupad [F(-1)] . . . . .	3764

## 3.509.1 Optimal result

Integrand size = 21, antiderivative size = 772

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x(d+ex^2)^3} dx = & -\frac{bcex(1-c^2x^2)}{8d^2(c^2d+e)\sqrt{-1+cx}\sqrt{1+cx}(d+ex^2)} \\
& + \frac{a + \operatorname{barccosh}(cx)}{4d(d+ex^2)^2} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(d+ex^2)} \\
& + \frac{(a + \operatorname{barccosh}(cx))^2}{bd^3} - \frac{bc\sqrt{-1+c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{2d^{5/2}\sqrt{c^2d+e}\sqrt{-1+cx}\sqrt{1+cx}} \\
& - \frac{bc(2c^2d+e)\sqrt{-1+c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{8d^{5/2}(c^2d+e)^{3/2}\sqrt{-1+cx}\sqrt{1+cx}} \\
& + \frac{(a + \operatorname{barccosh}(cx))\log(1+e^{-2\operatorname{arccosh}(cx)})}{d^3} \\
& - \frac{(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^3} \\
& - \frac{(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^3} \\
& - \frac{(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^3} \\
& - \frac{(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^3} \\
& - \frac{b \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d^3} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^3} \\
& - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d-\sqrt{-c^2d-e}}}\right)}{2d^3} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^3} \\
& - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d+\sqrt{-c^2d-e}}}\right)}{2d^3}
\end{aligned}$$

output

```

1/4*(a+b*arccosh(c*x))/d/(e*x^2+d)^2+1/2*(a+b*arccosh(c*x))/d^2/(e*x^2+d)+
(a+b*arccosh(c*x))^2/b/d^3+(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c
*x+1)^(1/2))^2)/d^3-1/2*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1
)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^3-1/2*(a+b*arccosh(c*x)
)*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(
1/2)))/d^3-1/2*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*
e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^3-1/2*(a+b*arccosh(c*x))*ln(1+(
c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/
d^3-1/2*b*polylog(2,-1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d^3-1/2*b*poly
log(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(
1/2)))/d^3-1/2*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(
-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^3-1/2*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+
1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^3-1/2*b*polylog(2,(c*
x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^
3-1/8*b*c*e*x*(-c^2*x^2+1)/d^2/(c^2*d+e)/(e*x^2+d)/(c*x-1)^(1/2)/(c*x+1)^(
1/2)-1/8*b*c*(2*c^2*d+e)*arctanh(x*(c^2*d+e)^(1/2)/d^(1/2)/(c^2*x^2-1)^(1/
2))*(c^2*x^2-1)^(1/2)/d^(5/2)/(c^2*d+e)^(3/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)-
1/2*b*c*arctanh(x*(c^2*d+e)^(1/2)/d^(1/2)/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(
1/2)/d^(5/2)/(c^2*d+e)^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)

```

### 3.509.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 6.07 (sec) , antiderivative size = 1204, normalized size of antiderivative = 1.56

$$\begin{aligned}
 \int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^3} dx &= \frac{a}{4d(d + ex^2)^2} + \frac{a}{2d^2(d + ex^2)} + \frac{a \log(x)}{d^3} \\
 &- \frac{a \log(d + ex^2)}{2d^3} + b \left( - \frac{5i \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d} + \sqrt{ex}} + \frac{c \log \left( \frac{2e(i\sqrt{e} + c^2\sqrt{dx} - i\sqrt{-c^2d - e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c\sqrt{-c^2d - e}(\sqrt{d} + i\sqrt{ex})} \right)}{\sqrt{-c^2d - e}} \right)}{16d^{5/2}} \right. \\
 &\quad \left. - \frac{5i \left( - \frac{\operatorname{arccosh}(cx)}{i\sqrt{d} + \sqrt{ex}} - \frac{c \log \left( \frac{2e(-\sqrt{e} - ic^2\sqrt{dx} + \sqrt{-c^2d - e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c\sqrt{-c^2d - e}(i\sqrt{d} + \sqrt{ex})} \right)}{\sqrt{-c^2d - e}} \right)}{16d^{5/2}} \right) \\
 &+ \frac{\sqrt{e} \left( \frac{c\sqrt{-1 + cx}\sqrt{1 + cx}}{(c^2d + e)(-i\sqrt{d} + \sqrt{ex})} - \frac{\operatorname{arccosh}(cx)}{\sqrt{e}(-i\sqrt{d} + \sqrt{ex})^2} + \frac{c^3\sqrt{d} \left( \log(4) + \log \left( \frac{e\sqrt{c^2d + e}(-i\sqrt{e} - c^2\sqrt{dx} + \sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c^3(d + i\sqrt{d}\sqrt{ex})} \right) \right)}{\sqrt{e}(c^2d + e)^{3/2}} \right)}{16d^2} \\
 &+ \frac{\sqrt{e} \left( \frac{c\sqrt{-1 + cx}\sqrt{1 + cx}}{(c^2d + e)(i\sqrt{d} + \sqrt{ex})} - \frac{\operatorname{arccosh}(cx)}{\sqrt{e}(i\sqrt{d} + \sqrt{ex})^2} - \frac{c^3\sqrt{d} \left( \log(4) + \log \left( \frac{e\sqrt{c^2d + e}(-i\sqrt{e} + c^2\sqrt{dx} + \sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c^3(d - i\sqrt{d}\sqrt{ex})} \right) \right)}{\sqrt{e}(c^2d + e)^{3/2}} \right)}{16d^2} \\
 &+ \frac{\operatorname{arccosh}(cx) \left( \operatorname{arccosh}(cx) + 2 \log(1 + e^{-2\operatorname{arccosh}(cx)}) \right) - \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})}{2d^3} \\
 &- \frac{\operatorname{arccosh}(cx) \left( -\operatorname{arccosh}(cx) + 2 \left( \log \left( 1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} - \sqrt{-c^2d - e}} \right) + \log \left( 1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} + \sqrt{-c^2d - e}} \right) \right) \right) + 2 \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})}{4d^3} \\
 &- \frac{\operatorname{arccosh}(cx) \left( -\operatorname{arccosh}(cx) + 2 \left( \log \left( 1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-ic\sqrt{d} + \sqrt{-c^2d - e}} \right) + \log \left( 1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} + \sqrt{-c^2d - e}} \right) \right) \right) + 2 \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)})}{4d^3}
 \end{aligned}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x*(d + e*x^2)^3), x]`

output

```

a/(4*d*(d + e*x^2)^2) + a/(2*d^2*(d + e*x^2)) + (a*Log[x])/d^3 - (a*Log[d
+ e*x^2])/(2*d^3) + b*(((5*I)/16)*(ArcCosh[c*x]/((-I)*Sqrt[d] + Sqrt[e]*
x) + (c*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*d) - e]*Sqrt[-1
+ c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*Sqrt[e]*x)))/S
qrt[-(c^2*d) - e])/d^(5/2) - (((5*I)/16)*(-(ArcCosh[c*x]/(I*Sqrt[d] + Sqr
t[e]*x)) - (c*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[-(c^2*d) - e]*Sq
rt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) - e]*(I*Sqrt[d] + Sqrt[e]*x
)))/Sqrt[-(c^2*d) - e])/d^(5/2) + (Sqrt[e]*((c*Sqrt[-1 + c*x]*Sqrt[1 + c*
x]))/((c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*((-I)
*Sqrt[d] + Sqrt[e]*x)^2) + (c^3*Sqrt[d]*(Log[4] + Log[(e*Sqrt[c^2*d + e]*
(-I)*Sqrt[e] - c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x
]))/(c^3*(d + I*Sqrt[d]*Sqrt[e]*x)))/((Sqrt[e]*(c^2*d + e)^(3/2))))/(16*d
^2) + (Sqrt[e]*((c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/((c^2*d + e)*(I*Sqrt[d] +
Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*(I*Sqrt[d] + Sqrt[e]*x)^2) - (c^3*Sqr
t[d]*(Log[4] + Log[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[e] + c^2*Sqrt[d]*x + Sqrt
[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c^3*(d - I*Sqrt[d]*Sqrt[e]*x)
)))/((Sqrt[e]*(c^2*d + e)^(3/2))))/(16*d^2) + (ArcCosh[c*x]*(ArcCosh[c*x] +
2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLog[2, -E^(-2*ArcCosh[c*x])])/(2*d^
3) - (ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x])]/(I
*c*Sqrt[d] - Sqrt[-(c^2*d) - e])) + Log[1 + (Sqrt[e]*E^ArcCosh[c*x])]/(I...

```

### 3.509.3 Rubi [A] (verified)

Time = 1.71 (sec) , antiderivative size = 772, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x(d + ex^2)^3} dx$$

↓ 6374

$$\int \left( -\frac{ex(a + \operatorname{barccosh}(cx))}{d^3(d + ex^2)} + \frac{a + \operatorname{barccosh}(cx)}{d^3x} - \frac{ex(a + \operatorname{barccosh}(cx))}{d^2(d + ex^2)^2} - \frac{ex(a + \operatorname{barccosh}(cx))}{d(d + ex^2)^3} \right) dx$$

↓ 2009



$$\begin{aligned}
& \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}}\right)}{2d^3} - \frac{(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{c^2(-d) - e}} + 1\right)}{2d^3} \\
& \frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}}\right)}{2d^3} - \frac{(a + \operatorname{barccosh}(cx)) \log\left(\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{c^2(-d) - e} + c\sqrt{-d}} + 1\right)}{2d^3} + \\
& \frac{(a + \operatorname{barccosh}(cx))^2}{bd^3} + \frac{\log(e^{-2\operatorname{arccosh}(cx)} + 1) (a + \operatorname{barccosh}(cx))}{d^3} + \frac{a + \operatorname{barccosh}(cx)}{2d^2(d + ex^2)} + \\
& \frac{a + \operatorname{barccosh}(cx)}{4d(d + ex^2)^2} - \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2d^3} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-dc^2 - e}}\right)}{2d^3} - \\
& \frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2d^3} - \frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{\sqrt{-dc} + \sqrt{-dc^2 - e}}\right)}{2d^3} - \\
& \frac{b \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d^3} - \frac{bc\sqrt{c^2x^2 - 1}(2c^2d + e) \operatorname{arctanh}\left(\frac{x\sqrt{c^2d + e}}{\sqrt{d}\sqrt{c^2x^2 - 1}}\right)}{8d^{5/2}\sqrt{cx - 1}\sqrt{cx + 1}(c^2d + e)^{3/2}} - \\
& \frac{bc\sqrt{c^2x^2 - 1} \operatorname{arctanh}\left(\frac{x\sqrt{c^2d + e}}{\sqrt{d}\sqrt{c^2x^2 - 1}}\right)}{2d^{5/2}\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{c^2d + e}} - \frac{bcex(1 - c^2x^2)}{8d^2\sqrt{cx - 1}\sqrt{cx + 1}(c^2d + e)(d + ex^2)}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(x*(d + e*x^2)^3), x]`

output

```

-1/8*(b*c*e*x*(1 - c^2*x^2))/(d^2*(c^2*d + e)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]
*(d + e*x^2)) + (a + b*ArcCosh[c*x])/(4*d*(d + e*x^2)^2) + (a + b*ArcCosh[
c*x])/(2*d^2*(d + e*x^2)) + (a + b*ArcCosh[c*x])^2/(b*d^3) - (b*c*Sqrt[-1
+ c^2*x^2]*ArcTanh[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2])])/(2*d
^(5/2)*Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (b*c*(2*c^2*d + e)*
Sqrt[-1 + c^2*x^2]*ArcTanh[(Sqrt[c^2*d + e]*x)/(Sqrt[d]*Sqrt[-1 + c^2*x^2]
)])/((8*d^(5/2)*(c^2*d + e)^(3/2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) + ((a + b*A
rcCosh[c*x])*Log[1 + E^(-2*ArcCosh[c*x])])/d^3 - ((a + b*ArcCosh[c*x])*Log
[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*d^3)
- ((a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqr
t[-(c^2*d) - e])])/(2*d^3) - ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcC
osh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*d^3) - ((a + b*ArcCosh[c*
x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(
2*d^3) - (b*PolyLog[2, -E^(-2*ArcCosh[c*x])])/(2*d^3) - (b*PolyLog[2, -((S
qrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*d^3) - (b*P
olyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*
d^3) - (b*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d
) - e])])/(2*d^3) - (b*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] +
Sqrt[-(c^2*d) - e])])/(2*d^3)

```

**3.509.3.1 Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

**3.509.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.44 (sec) , antiderivative size = 1225, normalized size of antiderivative = 1.59

method	result	size
parts	Expression too large to display	1225
derivativedivides	Expression too large to display	1278
default	Expression too large to display	1278

input `int((a+b*arccosh(c*x))/x/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

```

output 1/4*a/d/(e*x^2+d)^2-1/2*a/d^3*ln(e*x^2+d)+1/2*a/d^2/(e*x^2+d)+a/d^3*ln(x)+
b*(1/8*c^2*(6*c^4*d^2*arccosh(c*x)+4*arccosh(c*x)*c^4*d*e*x^2+(c*x-1)^(1/2)
)*(c*x+1)^(1/2)*c^3*d*e*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)*e^2*c^3*x^3-c^4*d^2-
2*c^4*d*e*x^2-c^4*e^2*x^4+6*c^2*d*e*arccosh(c*x)+4*arccosh(c*x)*e^2*c^2*x^
2)/d^2/(c^2*d+e)/(c^2*e*x^2+c^2*d)^2+5/8*(d*c^2*(c^2*d+e))^(1/2)/(c^2*d+e)
^2/d^3*e*arctanh(1/4*(4*c^2*d+2*e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2+2*e)
/(c^4*d^2+c^2*d*e)^(1/2))+3/4*(d*c^2*(c^2*d+e))^(1/2)/(c^2*d+e)^2/d^2*c^2*
arctanh(1/4*(4*c^2*d+2*e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))^2+2*e)/(c^4*d^2
+c^2*d*e)^(1/2))-1/4/(c^2*d+e)/d^3*e*sum((_R1^2*e+4*c^2*d+e)/(_R1^2*e+2*c^
2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((
_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)
*_Z^2+e))+1/(c^2*d+e)/d^3*e*arccosh(c*x)*ln(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)
^(1/2)))+1/(c^2*d+e)/d^3*e*arccosh(c*x)*ln(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)
^(1/2)))+1/(c^2*d+e)/d^3*e*dilog(1+I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))+1/(
c^2*d+e)/d^3*e*dilog(1-I*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))-1/4/(c^2*d+e)/
d^3*e^2*sum((_R1^2+1)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)
)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R
1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))-1/4/(c^2*d+e)/d^2*c^2*sum((_R
1^2*e+4*c^2*d+e)/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/
2)*(c*x+1)^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)...

```

### 3.509.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^3} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(ex^2 + d)^3 x} dx$$

```
input integrate((a+b*arccosh(c*x))/x/(e*x^2+d)^3,x, algorithm="fricas")
```

```
output integral((b*arccosh(c*x) + a)/(e^3*x^7 + 3*d*e^2*x^5 + 3*d^2*e*x^3 + d^3*x
), x)
```

**3.509.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/x/(e*x**2+d)**3,x)`output `Timed out`**3.509.7 Maxima [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^3} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(ex^2 + d)^3 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d)^3,x, algorithm="maxima")`output `1/4*a*((2*e*x^2 + 3*d)/(d^2*e^2*x^4 + 2*d^3*e*x^2 + d^4) - 2*log(e*x^2 + d)/d^3 + 4*log(x)/d^3) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(e^3*x^7 + 3*d*e^2*x^5 + 3*d^2*e*x^3 + d^3*x), x)`**3.509.8 Giac [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^3} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(ex^2 + d)^3 x} dx$$

input `integrate((a+b*arccosh(c*x))/x/(e*x^2+d)^3,x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)/((e*x^2 + d)^3*x), x)`

**3.509.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x(d + ex^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x(ex^2 + d)^3} dx$$

input `int((a + b*acosh(c*x))/(x*(d + e*x^2)^3), x)`output `int((a + b*acosh(c*x))/(x*(d + e*x^2)^3), x)`

$$3.510 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d+ex^2)^3} dx$$

3.510.1 Optimal result . . . . .	3766
3.510.2 Mathematica [C] (warning: unable to verify) . . . . .	3767
3.510.3 Rubi [A] (verified) . . . . .	3769
3.510.4 Maple [C] (warning: unable to verify) . . . . .	3771
3.510.5 Fricas [F] . . . . .	3772
3.510.6 Sympy [F(-1)] . . . . .	3773
3.510.7 Maxima [F] . . . . .	3773
3.510.8 Giac [F] . . . . .	3773
3.510.9 Mupad [F(-1)] . . . . .	3774

## 3.510.1 Optimal result

Integrand size = 21, antiderivative size = 834

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^3} dx = & \frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{2d^3x} + \frac{bce^2x(1 - c^2x^2)}{8d^3(c^2d + e)\sqrt{-1 + cx}\sqrt{1 + cx}(d + ex^2)} \\
& - \frac{a + \operatorname{barccosh}(cx)}{2d^3x^2} - \frac{e(a + \operatorname{barccosh}(cx))}{4d^2(d + ex^2)^2} - \frac{e(a + \operatorname{barccosh}(cx))}{d^3(d + ex^2)} \\
& - \frac{3e(a + \operatorname{barccosh}(cx))^2}{bd^4} + \frac{bce\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{d^{7/2}\sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx}} \\
& + \frac{bce(2c^2d + e)\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{c^2d+ex}}{\sqrt{d}\sqrt{-1+c^2x^2}}\right)}{8d^{7/2}(c^2d + e)^{3/2}\sqrt{-1 + cx}\sqrt{1 + cx}} \\
& - \frac{3e(a + \operatorname{barccosh}(cx))\log(1 + e^{-2\operatorname{arccosh}(cx)})}{d^4} \\
& + \frac{3e(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^4} \\
& + \frac{3e(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^4} \\
& + \frac{3e(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^4} \\
& + \frac{3e(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^4} \\
& + \frac{3be \operatorname{PolyLog}\left(2, -e^{-2\operatorname{arccosh}(cx)}\right)}{2d^4} \\
& + \frac{3be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^4} \\
& + \frac{3be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d - \sqrt{-c^2d - e}}}\right)}{2d^4} \\
& + \frac{3be \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^4} \\
& + \frac{3be \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d + \sqrt{-c^2d - e}}}\right)}{2d^4}
\end{aligned}$$

output

```

1/2*(-a-b*arccosh(c*x))/d^3/x^2-1/4*e*(a+b*arccosh(c*x))/d^2/(e*x^2+d)^2-e
*(a+b*arccosh(c*x))/d^3/(e*x^2+d)-3*e*(a+b*arccosh(c*x))^2/b/d^4-3*e*(a+b*
arccosh(c*x))*ln(1+1/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d^4+3/2*e*(a+b*a
rccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-
(-c^2*d-e)^(1/2)))/d^4+3/2*e*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c
*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^4+3/2*e*(a+b*arcco
sh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^
2*d-e)^(1/2)))/d^4+3/2*e*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1
)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^4+3/2*b*e*polylog(2,-1
/(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))^2)/d^4+3/2*b*e*polylog(2,-(c*x+(c*x-1)^(
1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/d^4+3/2*b*e*
polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-
e)^(1/2)))/d^4+3/2*b*e*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2
)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^4+3/2*b*e*polylog(2,(c*x+(c*x-1)^(1/2
)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/d^4+1/8*b*c*e^2*
x*(-c^2*x^2+1)/d^3/(c^2*d+e)/(e*x^2+d)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+1/2*b*c
*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d^3/x+1/8*b*c*e*(2*c^2*d+e)*arctanh(x*(c^2*d+
e)^(1/2)/d^(1/2)/(c^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/d^(7/2)/(c^2*d+e)^(3
/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*c*e*arctanh(x*(c^2*d+e)^(1/2)/d^(1/2)/(c
^2*x^2-1)^(1/2))*(c^2*x^2-1)^(1/2)/d^(7/2)/(c^2*d+e)^(1/2)/(c*x-1)^(1/2)...

```

### 3.510.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.



Time = 6.09 (sec) , antiderivative size = 1261, normalized size of antiderivative = 1.51

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d + ex^2)^3} dx = -\frac{a}{2d^3 x^2} - \frac{ae}{4d^2 (d + ex^2)^2} - \frac{ae}{d^3 (d + ex^2)}$$

$$- \frac{3ae \log(x)}{d^4} + \frac{3ae \log(d + ex^2)}{2d^4} + b \left( \frac{cx\sqrt{-1+cx}\sqrt{1+cx} - \operatorname{arccosh}(cx)}{2d^3 x^2} \right.$$

$$+ \frac{9ie \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d} + \sqrt{ex}} + \frac{c \log\left(\frac{2e(i\sqrt{e} + c^2\sqrt{dx} - i\sqrt{-c^2d-e}\sqrt{-1+cx}\sqrt{1+cx})}{c\sqrt{-c^2d-e}(\sqrt{d} + i\sqrt{ex})}\right)}{\sqrt{-c^2d-e}} \right)}{16d^{7/2}}$$

$$+ \frac{9ie \left( -\frac{\operatorname{arccosh}(cx)}{i\sqrt{d} + \sqrt{ex}} - \frac{c \log\left(\frac{2e(-\sqrt{e} - ic^2\sqrt{dx} + \sqrt{-c^2d-e}\sqrt{-1+cx}\sqrt{1+cx})}{c\sqrt{-c^2d-e}(i\sqrt{d} + \sqrt{ex})}\right)}{\sqrt{-c^2d-e}} \right)}{16d^{7/2}}$$

$$+ \frac{e^{3/2} \left( \frac{c\sqrt{-1+cx}\sqrt{1+cx}}{(c^2d+e)(-i\sqrt{d} + \sqrt{ex})} - \frac{\operatorname{arccosh}(cx)}{\sqrt{e}(-i\sqrt{d} + \sqrt{ex})^2} + \frac{c^3\sqrt{d} \left( \log(4) + \log\left(\frac{e\sqrt{c^2d+e}(-i\sqrt{e} - c^2\sqrt{dx} + \sqrt{c^2d+e}\sqrt{-1+cx}\sqrt{1+cx})}{c^3(d+i\sqrt{d}\sqrt{ex})}\right) \right)}{\sqrt{e}(c^2d+e)^{3/2}} \right)}{16d^3}$$

$$- \frac{e^{3/2} \left( \frac{c\sqrt{-1+cx}\sqrt{1+cx}}{(c^2d+e)(i\sqrt{d} + \sqrt{ex})} - \frac{\operatorname{arccosh}(cx)}{\sqrt{e}(i\sqrt{d} + \sqrt{ex})^2} - \frac{c^3\sqrt{d} \left( \log(4) + \log\left(\frac{e\sqrt{c^2d+e}(-i\sqrt{e} + c^2\sqrt{dx} + \sqrt{c^2d+e}\sqrt{-1+cx}\sqrt{1+cx})}{c^3(d-i\sqrt{d}\sqrt{ex})}\right) \right)}{\sqrt{e}(c^2d+e)^{3/2}} \right)}{16d^3}$$

$$- \frac{3e(\operatorname{arccosh}(cx) (\operatorname{arccosh}(cx) + 2 \log(1 + e^{-2\operatorname{arccosh}(cx)})) - \operatorname{PolyLog}(2, -e^{-2\operatorname{arccosh}(cx)}))}{2d^4}$$

$$+ \frac{3e \left( \operatorname{arccosh}(cx) \left( -\operatorname{arccosh}(cx) + 2 \left( \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} - \sqrt{-c^2d-e}}\right) + \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} + \sqrt{-c^2d-e}}\right) \right) \right) + 2 \operatorname{PolyLog}}{4d^4}$$

$$+ \frac{3e \left( \operatorname{arccosh}(cx) \left( -\operatorname{arccosh}(cx) + 2 \left( \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-ic\sqrt{d} + \sqrt{-c^2d-e}}\right) + \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} + \sqrt{-c^2d-e}}\right) \right) \right) + 2 \operatorname{PolyLog}}{4d^4}$$

input `Integrate[(a + b*ArcCosh[c*x])/(x^3*(d + e*x^2)^3), x]`

3.510.  $\int \frac{a+b\operatorname{arccosh}(cx)}{x^3(d+ex^2)^3} dx$

output 
$$\begin{aligned}
 & -1/2*a/(d^3*x^2) - (a*e)/(4*d^2*(d + e*x^2)^2) - (a*e)/(d^3*(d + e*x^2)) - \\
 & (3*a*e*Log[x])/d^4 + (3*a*e*Log[d + e*x^2])/(2*d^4) + b*((c*x*Sqrt[-1 + c \\
 & *x]*Sqrt[1 + c*x] - ArcCosh[c*x])/(2*d^3*x^2) + (((9*I)/16)*e*(ArcCosh[c*x \\
 & ]/((-I)*Sqrt[d + Sqrt[e]*x) + (c*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I* \\
 & Sqrt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) - e]*(S \\
 & qrt[d + I*Sqrt[e]*x)))]/Sqrt[-(c^2*d) - e]))/d^(7/2) + (((9*I)/16)*e*(-(A \\
 & rcCosh[c*x]/(I*Sqrt[d + Sqrt[e]*x)) - (c*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[ \\
 & d]*x + Sqrt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) \\
 & - e]*(I*Sqrt[d + Sqrt[e]*x)))]/Sqrt[-(c^2*d) - e]))/d^(7/2) - (e^(3/2))*(( \\
 & c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/((c^2*d + e)*((-I)*Sqrt[d + Sqrt[e]*x)) - \\
 & ArcCosh[c*x]/(Sqrt[e]*((-I)*Sqrt[d + Sqrt[e]*x)^2) + (c^3*Sqrt[d]*(Log[4 \\
 & ] + Log[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[e] - c^2*Sqrt[d]*x + Sqrt[c^2*d + e] \\
 & *Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c^3*(d + I*Sqrt[d]*Sqrt[e]*x)))]/(Sqrt[e] \\
 & ]*(c^2*d + e)^(3/2)))/(16*d^3) - (e^(3/2))*((c*Sqrt[-1 + c*x]*Sqrt[1 + c*x] \\
 & )/((c^2*d + e)*(I*Sqrt[d + Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*(I*Sqrt[d \\
 & ] + Sqrt[e]*x)^2) - (c^3*Sqrt[d]*(Log[4 + Log[(e*Sqrt[c^2*d + e]*((-I)*Sq \\
 & rt[e] + c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c^ \\
 & 3*(d - I*Sqrt[d]*Sqrt[e]*x)))]/(Sqrt[e]*(c^2*d + e)^(3/2)))/(16*d^3) - ( \\
 & 3*e*(ArcCosh[c*x]*(ArcCosh[c*x] + 2*Log[1 + E^(-2*ArcCosh[c*x])]) - PolyLo \\
 & g[2, -E^(-2*ArcCosh[c*x])]))/(2*d^4) + (3*e*(ArcCosh[c*x]*(-ArcCosh[c*x]...
 \end{aligned}$$

### 3.510.3 Rubi [A] (verified)

Time = 1.83 (sec) , antiderivative size = 834, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \text{barccosh}(cx)}{x^3 (d + ex^2)^3} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{3e^2 x(a + \text{barccosh}(cx))}{d^4 (d + ex^2)} - \frac{3e(a + \text{barccosh}(cx))}{d^4 x} + \frac{2e^2 x(a + \text{barccosh}(cx))}{d^3 (d + ex^2)^2} + \frac{a + \text{barccosh}(cx)}{d^3 x^3} + \frac{e^2 x(a + \text{barccosh}(cx))}{d^2 (d + ex^2)} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$



**3.510.3.1 Defintions of rubi rules used**

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6374 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]`

**3.510.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 2.69 (sec) , antiderivative size = 1455, normalized size of antiderivative = 1.74

method	result	size
parts	Expression too large to display	1455
derivativedivides	Expression too large to display	1515
default	Expression too large to display	1515

input `int((a+b*arccosh(c*x))/x^3/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output  $\frac{3}{2}ae/d^4 \ln(ex^2+d) - \frac{1}{4}ae/d^2 (ex^2+d)^{-2} - ae/d^3 (ex^2+d)^{-1} - \frac{1}{2}a/d^3 x^2 - 3a/d^4 e \ln(x) + bc^2(-\frac{1}{8}(-4(c*x-1)^{1/2}(c*x+1)^{1/2})c^7d^3x - 8(c*x-1)^{1/2}(c*x+1)^{1/2})c^7d^2e^2x^3 - 4(c*x-1)^{1/2}(c*x+1)^{1/2})c^7d^2e^2x^5 + 4c^8d^3x^2 + 8c^8d^2e^2x^4 + 4c^8d^2e^2x^6 + 4c^6d^3 \operatorname{arccosh}(c*x) + 18 \operatorname{arccosh}(c*x)c^6d^2e^2x^2 + 12 \operatorname{arccosh}(c*x)c^6d^2e^2x^4 - 4(c*x-1)^{1/2}(c*x+1)^{1/2})c^5d^2e^2x^7 - (c*x-1)^{1/2}(c*x+1)^{1/2})c^5d^2e^2x^3 - 3(c*x-1)^{1/2}(c*x+1)^{1/2})e^3c^5x^5 + 3c^6d^2e^2x^2 + 6c^6d^2e^2x^4 + 3c^6e^3x^6 + 4c^4d^2e \operatorname{arccosh}(c*x) + 18 \operatorname{arccosh}(c*x)c^4d^2e^2x^2 + 12 \operatorname{arccosh}(c*x)e^3c^4x^4) / c^2/x^2/d^3 / (c^2d+e) / (c^2e^2x^2+c^2d)^2 - 9/8(d*c^2(c^2d+e))^{1/2} / (c^2d+e)^2/d^4/c^2e^2 \operatorname{arctanh}(1/4(4*c^2d+2*e*(c*x+(c*x-1)^{1/2}(c*x+1)^{1/2}))^2+2*e) / (c^4d^2+c^2d*e)^{1/2}) - 5/4(d*c^2(c^2d+e))^{1/2} / (c^2d+e)^2/d^3e \operatorname{arctanh}(1/4(4*c^2d+2*e*(c*x+(c*x-1)^{1/2}(c*x+1)^{1/2}))^2+2*e) / (c^4d^2+c^2d*e)^{1/2}) + 3/4/(c^2d+e)e^2/d^4/c^2 \operatorname{sum}((\_R1^2e+4c^2d+e) / (\_R1^2e+2c^2d+e) * (\operatorname{arccosh}(c*x) \ln((\_R1-c*x-(c*x-1)^{1/2}(c*x+1)^{1/2}) / \_R1) + \operatorname{dilog}((\_R1-c*x-(c*x-1)^{1/2}(c*x+1)^{1/2}) / \_R1)), \_R1=RootOf(e*_Z^4+(4*c^2d+2*e)*_Z^2+e)) - 3/(c^2d+e)e^2/d^4/c^2 \operatorname{arccosh}(c*x) \ln(1+I*(c*x+(c*x-1)^{1/2}(c*x+1)^{1/2})) - 3/(c^2d+e)e^2/d^4/c^2 \operatorname{arccosh}(c*x) \ln(1-I*(c*x+(c*x-1)^{1/2}(c*x+1)^{1/2})) - 3/(c^2d+e)e^2/d^4/c^2 \operatorname{dilog}(1+I*(c*x+(c*x-1)^{1/2}(c*x+1)^{1/2})) - 3/(c^2d+e)e^2/d^4/c^2 \operatorname{dilog}(1-I*(c*x+(c*x-1)^{1/2}(c*x+1)^{1/2})) + 3/4/(c^2d+e)...$

### 3.510.5 Fracas [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3(d + ex^2)^3} dx = \int \frac{b \operatorname{arccosh}(cx) + a}{(ex^2 + d)^3 x^3} dx$$

input `integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e^3*x^9 + 3*d*e^2*x^7 + 3*d^2*e*x^5 + d^3*x^3), x)`

**3.510.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^3} dx = \text{Timed out}$$

```
input integrate((a+b*acosh(c*x))/x**3/(e*x**2+d)**3,x)
```

```
output Timed out
```

**3.510.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^3} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^3 x^3} dx$$

```
input integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d)^3,x, algorithm="maxima")
```

```
output -1/4*a*((6*e^2*x^4 + 9*d*e*x^2 + 2*d^2)/(d^3*e^2*x^6 + 2*d^4*e*x^4 + d^5*x^2) - 6*e*log(e*x^2 + d)/d^4 + 12*e*log(x)/d^4) + b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(e^3*x^9 + 3*d*e^2*x^7 + 3*d^2*e*x^5 + d^3*x^3), x)
```

**3.510.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{x^3 (d + ex^2)^3} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^3 x^3} dx$$

```
input integrate((a+b*arccosh(c*x))/x^3/(e*x^2+d)^3,x, algorithm="giac")
```

```
output integrate((b*arccosh(c*x) + a)/((e*x^2 + d)^3*x^3), x)
```

**3.510.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{x^3 (d + ex^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{x^3 (ex^2 + d)^3} dx$$

input `int((a + b*acosh(c*x))/(x^3*(d + e*x^2)^3), x)`output `int((a + b*acosh(c*x))/(x^3*(d + e*x^2)^3), x)`

$$3.511 \quad \int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$$

3.511.1 Optimal result . . . . .	3776
3.511.2 Mathematica [C] (warning: unable to verify) . . . . .	3777
3.511.3 Rubi [A] (verified) . . . . .	3779
3.511.4 Maple [C] (warning: unable to verify) . . . . .	3782
3.511.5 Fricas [F] . . . . .	3783
3.511.6 Sympy [F(-1)] . . . . .	3783
3.511.7 Maxima [F(-2)] . . . . .	3783
3.511.8 Giac [F] . . . . .	3784
3.511.9 Mupad [F(-1)] . . . . .	3784



### 3.511.1 Optimal result

Integrand size = 21, antiderivative size = 1224

$$\begin{aligned}
 \int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = & -\frac{bc\sqrt{-d}\sqrt{-1+cx}\sqrt{1+cx}}{16e^2(c^2d+e)(\sqrt{-d}-\sqrt{ex})} - \frac{bc\sqrt{-d}\sqrt{-1+cx}\sqrt{1+cx}}{16e^2(c^2d+e)(\sqrt{-d}+\sqrt{ex})} \\
 & - \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx))}{16e^{5/2}(\sqrt{-d}-\sqrt{ex})^2} + \frac{5(a + \operatorname{barccosh}(cx))}{16e^{5/2}(\sqrt{-d}-\sqrt{ex})} \\
 & + \frac{\sqrt{-d}(a + \operatorname{barccosh}(cx))}{16e^{5/2}(\sqrt{-d}+\sqrt{ex})^2} - \frac{5(a + \operatorname{barccosh}(cx))}{16e^{5/2}(\sqrt{-d}+\sqrt{ex})} \\
 & - \frac{bc^3 \operatorname{darctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}\right)}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(c\sqrt{-d}+\sqrt{e})^{3/2}e^{5/2}} \\
 & - \frac{5b \operatorname{carctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}\right)}{8\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}e^{5/2}} \\
 & + \frac{bc^3 \operatorname{darctanh}\left(\frac{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}\right)}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(c\sqrt{-d}+\sqrt{e})^{3/2}e^{5/2}} \\
 & + \frac{5b \operatorname{carctanh}\left(\frac{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}\right)}{8\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}e^{5/2}} \\
 & + \frac{3(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & - \frac{3(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & + \frac{3(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & - \frac{3(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & - \frac{3b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & + \frac{3b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & - \frac{3b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & + \frac{3b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}}
 \end{aligned}$$


---

3.511.  $\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx$

$$\begin{aligned}
 & + \frac{3b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
 & + \frac{3b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee}^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16\sqrt{-d}e^{5/2}}
 \end{aligned}$$

output

```

3/16*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*
(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(5/2)/(-d)^(1/2)-3/16*(a+b*arccosh(c*x))*l
n(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/
2)))/e^(5/2)/(-d)^(1/2)+3/16*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c
*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)/(-d)^(1/2)-3
/16*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(
-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)/(-d)^(1/2)-3/16*b*polylog(2,-(c*x+(c*
x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(5/2)
/(-d)^(1/2)+3/16*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*
(-d)^(1/2)-(-c^2*d-e)^(1/2)))/e^(5/2)/(-d)^(1/2)-3/16*b*polylog(2,-(c*x+(c
*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)
)/(-d)^(1/2)+3/16*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c
*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/e^(5/2)/(-d)^(1/2)-1/8*b*c^3*d*arctanh((c*x
+1)^(1/2)*(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*x-1)^(1/2)/(c*(-d)^(1/2)+e^(1/2)
)^(1/2))/e^(5/2)/(c*(-d)^(1/2)-e^(1/2))^(3/2)/(c*(-d)^(1/2)+e^(1/2))^(3/2)
+1/8*b*c^3*d*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)+e^(1/2))^(1/2)/(c*x-1)^(1
/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2))/e^(5/2)/(c*(-d)^(1/2)-e^(1/2))^(3/2)/(c*
(-d)^(1/2)+e^(1/2))^(3/2)-1/16*(a+b*arccosh(c*x))*(-d)^(1/2)/e^(5/2)/((-d)
^(1/2)-x*e^(1/2))^2+5/16*(a+b*arccosh(c*x))/e^(5/2)/((-d)^(1/2)-x*e^(1/2))
+1/16*(a+b*arccosh(c*x))*(-d)^(1/2)/e^(5/2)/((-d)^(1/2)+x*e^(1/2))^2-5/...

```

### 3.511.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

---

3.511. 
$$\int \frac{x^4(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$$

Time = 11.03 (sec) , antiderivative size = 1185, normalized size of antiderivative = 0.97

$$\begin{aligned}
 \int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx &= \frac{adx}{4e^2(d + ex^2)^2} - \frac{5ax}{8e^2(d + ex^2)} + \frac{3a \arctan\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{8\sqrt{d}e^{5/2}} \\
 &+ b \left( -\frac{5 \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d} + \sqrt{ex}} + \frac{c \log\left(\frac{2e(i\sqrt{e} + c^2\sqrt{dx} - i\sqrt{-c^2d - e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c\sqrt{-c^2d - e}(\sqrt{d} + i\sqrt{ex})}\right)}{\sqrt{-c^2d - e}} \right)}{16e^{5/2}} \right. \\
 &\quad \left. + \frac{5 \left( -\frac{\operatorname{arccosh}(cx)}{i\sqrt{d} + \sqrt{ex}} - \frac{c \log\left(\frac{2e(-\sqrt{e} - ic^2\sqrt{dx} + \sqrt{-c^2d - e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c\sqrt{-c^2d - e}(i\sqrt{d} + \sqrt{ex})}\right)}{\sqrt{-c^2d - e}} \right)}{16e^{5/2}} \right) \\
 &+ \frac{i\sqrt{d} \left( \frac{c\sqrt{-1 + cx}\sqrt{1 + cx}}{(c^2d + e)(-i\sqrt{d} + \sqrt{ex})} - \frac{\operatorname{arccosh}(cx)}{\sqrt{e}(-i\sqrt{d} + \sqrt{ex})^2} + \frac{c^3\sqrt{d} \left( \log(4) + \log\left(\frac{e\sqrt{c^2d + e}(-i\sqrt{e} - c^2\sqrt{dx} + \sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c^3(d + i\sqrt{d}\sqrt{ex})}\right) \right)}{\sqrt{e}(c^2d + e)^{3/2}} \right)}{16e^2} \\
 &+ \frac{i\sqrt{d} \left( \frac{c\sqrt{-1 + cx}\sqrt{1 + cx}}{(c^2d + e)(i\sqrt{d} + \sqrt{ex})} - \frac{\operatorname{arccosh}(cx)}{\sqrt{e}(i\sqrt{d} + \sqrt{ex})^2} - \frac{c^3\sqrt{d} \left( \log(4) + \log\left(\frac{e\sqrt{c^2d + e}(-i\sqrt{e} + c^2\sqrt{dx} + \sqrt{c^2d + e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c^3(d - i\sqrt{d}\sqrt{ex})}\right) \right)}{\sqrt{e}(c^2d + e)^{3/2}} \right)}{16e^2} \\
 &+ \frac{3i \left( \operatorname{arccosh}(cx) \left( -\operatorname{arccosh}(cx) + 2 \left( \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} - \sqrt{-c^2d - e}}\right) + \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} + \sqrt{-c^2d - e}}\right) \right) \right) + 2 \operatorname{PolyLog}}{32\sqrt{d}e^{5/2}} \\
 &- \frac{3i \left( \operatorname{arccosh}(cx) \left( -\operatorname{arccosh}(cx) + 2 \left( \log\left(1 + \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{-ic\sqrt{d} + \sqrt{-c^2d - e}}\right) + \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{ic\sqrt{d} + \sqrt{-c^2d - e}}\right) \right) \right) + 2 \operatorname{PolyLog}}{32\sqrt{d}e^{5/2}}
 \end{aligned}$$

input `Integrate[(x^4*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

3.511.  $\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$

output

```
(a*d*x)/(4*e^2*(d + e*x^2)^2) - (5*a*x)/(8*e^2*(d + e*x^2)) + (3*a*ArcTan[
(Sqrt[e]*x)/Sqrt[d]])/(8*Sqrt[d]*e^(5/2)) + b*((-5*(ArcCosh[c*x]/((-I)*Sqr
t[d] + Sqrt[e]*x) + (c*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*
d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))]/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*
Sqrt[e]*x)))]/Sqrt[-(c^2*d) - e]))/(16*e^(5/2)) + (5*(-(ArcCosh[c*x]/(I*Sq
rt[d] + Sqrt[e]*x)) - (c*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[-(c^2
*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))]/(c*Sqrt[-(c^2*d) - e]*(I*Sqrt[d] +
Sqrt[e]*x)))]/Sqrt[-(c^2*d) - e]))/(16*e^(5/2)) + ((I/16)*Sqrt[d]*((c*Sqr
t[-1 + c*x]*Sqrt[1 + c*x])/((c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)) - ArcC
osh[c*x]/(Sqrt[e]*((-I)*Sqrt[d] + Sqrt[e]*x)^2) + (c^3*Sqrt[d]*(Log[4] + L
og[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[e] - c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt
[-1 + c*x]*Sqrt[1 + c*x]))]/(c^3*(d + I*Sqrt[d]*Sqrt[e]*x)))))/(Sqrt[e]*(c^
2*d + e)^(3/2)))/e^2 - ((I/16)*Sqrt[d]*((c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/
((c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*(I*Sqrt[d] +
Sqrt[e]*x)^2) - (c^3*Sqrt[d]*(Log[4] + Log[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[
e] + c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))]/(c^3*(
d - I*Sqrt[d]*Sqrt[e]*x)))))/(Sqrt[e]*(c^2*d + e)^(3/2)))/e^2 + (((3*I)/3
2)*(ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x])]/(I*c
*Sqrt[d] - Sqrt[-(c^2*d) - e])) + Log[1 + (Sqrt[e]*E^ArcCosh[c*x])]/(I*c*Sq
rt[d] + Sqrt[-(c^2*d) - e])))) + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/...
```

### 3.511.3 Rubi [A] (verified)

Time = 4.14 (sec) , antiderivative size = 1224, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$$

↓ 6374

$$\int \left( \frac{d^2(a + b \operatorname{arccosh}(cx))}{e^2(d + ex^2)^3} - \frac{2d(a + b \operatorname{arccosh}(cx))}{e^2(d + ex^2)^2} + \frac{a + b \operatorname{arccosh}(cx)}{e^2(d + ex^2)} \right) dx$$

↓ 2009

---

3.511.  $\int \frac{x^4(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$

$$\begin{aligned}
& - \frac{bd\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx-1}}}\right)c^3}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(\sqrt{-dc}+\sqrt{e})^{3/2}e^{5/2}} + \frac{bd\operatorname{arctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right)c^3}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(\sqrt{-dc}+\sqrt{e})^{3/2}e^{5/2}} \\
& - \frac{5b\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx-1}}}\right)c}{8\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{\sqrt{-dc}+\sqrt{e}}e^{5/2}} + \frac{5b\operatorname{arctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right)c}{8\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{\sqrt{-dc}+\sqrt{e}}e^{5/2}} \\
& - \frac{b\sqrt{-d}\sqrt{cx-1}\sqrt{cx+1}c}{16e^2(dc^2+e)(\sqrt{-d}-\sqrt{ex})} - \frac{b\sqrt{-d}\sqrt{cx-1}\sqrt{cx+1}c}{16e^2(dc^2+e)(\sqrt{ex}+\sqrt{-d})} + \frac{5(a+b\operatorname{arccosh}(cx))}{16e^{5/2}(\sqrt{-d}-\sqrt{ex})} \\
& - \frac{5(a+b\operatorname{arccosh}(cx))}{16e^{5/2}(\sqrt{ex}+\sqrt{-d})} - \frac{\sqrt{-d}(a+b\operatorname{arccosh}(cx))}{16e^{5/2}(\sqrt{-d}-\sqrt{ex})^2} + \frac{\sqrt{-d}(a+b\operatorname{arccosh}(cx))}{16e^{5/2}(\sqrt{ex}+\sqrt{-d})^2} + \\
& \frac{3(a+b\operatorname{arccosh}(cx))\log\left(1-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16\sqrt{-d}e^{5/2}} - \frac{3(a+b\operatorname{arccosh}(cx))\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{c\sqrt{-d}-\sqrt{-dc^2-e}}+1\right)}{16\sqrt{-d}e^{5/2}} + \\
& \frac{3(a+b\operatorname{arccosh}(cx))\log\left(1-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16\sqrt{-d}e^{5/2}} - \frac{3(a+b\operatorname{arccosh}(cx))\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{\sqrt{-dc}+\sqrt{-dc^2-e}}+1\right)}{16\sqrt{-d}e^{5/2}} \\
& - \frac{3b\operatorname{PolyLog}\left(2,-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16\sqrt{-d}e^{5/2}} + \frac{3b\operatorname{PolyLog}\left(2,\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16\sqrt{-d}e^{5/2}} \\
& - \frac{3b\operatorname{PolyLog}\left(2,-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16\sqrt{-d}e^{5/2}} + \frac{3b\operatorname{PolyLog}\left(2,\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16\sqrt{-d}e^{5/2}}
\end{aligned}$$

input `Int[(x^4*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

```

output -1/16*(b*c*Sqrt[-d]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(e^2*(c^2*d + e)*(Sqrt[-d] - Sqrt[e]*x)) - (b*c*Sqrt[-d]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(16*e^2*(c^2*d + e)*(Sqrt[-d] + Sqrt[e]*x)) - (Sqrt[-d]*(a + b*ArcCosh[c*x]))/(16*e^(5/2)*(Sqrt[-d] - Sqrt[e]*x)^2) + (5*(a + b*ArcCosh[c*x]))/(16*e^(5/2)*(Sqrt[-d] - Sqrt[e]*x)) + (Sqrt[-d]*(a + b*ArcCosh[c*x]))/(16*e^(5/2)*(Sqrt[-d] + Sqrt[e]*x)^2) - (5*(a + b*ArcCosh[c*x]))/(16*e^(5/2)*(Sqrt[-d] + Sqrt[e]*x)) - (b*c^3*d*ArcTanh[(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[-1 + c*x])])/(8*(c*Sqrt[-d] - Sqrt[e])^(3/2)*(c*Sqrt[-d] + Sqrt[e])^(3/2)*e^(5/2)) - (5*b*c*ArcTanh[(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[-1 + c*x])])/(8*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*e^(5/2)) + (b*c^3*d*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[-1 + c*x])])/(8*(c*Sqrt[-d] - Sqrt[e])^(3/2)*(c*Sqrt[-d] + Sqrt[e])^(3/2)*e^(5/2)) + (5*b*c*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[-1 + c*x])])/(8*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*e^(5/2)) + (3*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(16*Sqrt[-d]*e^(5/2)) - (3*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(16*Sqrt[-d]*e^(5/2)) + (3*(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])...

```

### 3.511.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d + e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]
```

**3.511.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 49.46 (sec) , antiderivative size = 1752, normalized size of antiderivative = 1.43

method	result	size
parts	Expression too large to display	1752
derivativedivides	Expression too large to display	1754
default	Expression too large to display	1754

input `int(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)`

output

```
a*((-5/8*x^3/e-3/8*d/e^2*x)/(e*x^2+d)^2+3/8/e^2/(d*e)^(1/2)*arctan(e*x/(d*
e)^(1/2)))+b/c^5*(-1/8*c^6*((c*x-1)^(1/2)*(c*x+1)^(1/2)*c^4*d^2+(c*x+1)^(1
/2)*(c*x-1)^(1/2)*c^4*d*e*x^2+3*arccosh(c*x)*d^2*c^5*x+5*arccosh(c*x)*d*c^
5*e*x^3+3*arccosh(c*x)*d*c^3*e*x+5*arccosh(c*x)*e^2*c^3*x^3)/e^2/(c^2*d+e)
/(c^2*e*x^2+c^2*d)^2-5/8*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*
(2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e+(d*c^2*(c^2*d+e))^(1/
2)*e)*c^6*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((-2*c^2*d+2*(d*c^2*
(c^2*d+e))^(1/2)-e)*e)^(1/2)/(c^2*d+e)^2/e^4+5/8*(-(2*c^2*d-2*(d*c^2*(c^2
*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctanh(e*(
c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e
)^(1/2)*c^6/e^4/(c^2*d+e)-5/8*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(
1/2)*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e
))^(1/2)*e)*c^6*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((2*c^2*d+2*(d*
c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)/(c^2*d+e)^2/e^4+5/8*((2*c^2*d+2*(d*c^2*(
c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctan(e
*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*
e)^(1/2)*c^6/e^4/(c^2*d+e)-3/16/(c^2*d+e)/e*c^6*sum(1/_R1/(_R1^2*e+2*c^2*
d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog((_R
1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_
Z^2+e))+3/16/(c^2*d+e)/e*c^6*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*...
```

**3.511.5 Fracas [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(ex^2 + d)^3} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*x^4*arccosh(c*x) + a*x^4)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**3.511.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x**4*(a+b*acosh(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**3.511.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: ValueError}$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`



**3.511.8 Giac [F]**

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^4}{(ex^2 + d)^3} dx$$

input `integrate(x^4*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^4/(e*x^2 + d)^3, x)`

**3.511.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^4(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{x^4(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^3} dx$$

input `int((x^4*(a + b*acosh(c*x)))/(d + e*x^2)^3,x)`

output `int((x^4*(a + b*acosh(c*x)))/(d + e*x^2)^3, x)`

$$3.512 \quad \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx$$

3.512.1 Optimal result . . . . .	3786
3.512.2 Mathematica [C] (warning: unable to verify) . . . . .	3787
3.512.3 Rubi [A] (verified) . . . . .	3788
3.512.4 Maple [C] (warning: unable to verify) . . . . .	3791
3.512.5 Fricas [F] . . . . .	3792
3.512.6 Sympy [F(-1)] . . . . .	3792
3.512.7 Maxima [F(-2)] . . . . .	3792
3.512.8 Giac [F] . . . . .	3793
3.512.9 Mupad [F(-1)] . . . . .	3793

## 3.512.1 Optimal result

Integrand size = 21, antiderivative size = 1234

$$\begin{aligned}
\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = & -\frac{bc\sqrt{-1+cx}\sqrt{1+cx}}{16\sqrt{-de}(c^2d+e)(\sqrt{-d}-\sqrt{ex})} \\
& -\frac{bc\sqrt{-1+cx}\sqrt{1+cx}}{16\sqrt{-de}(c^2d+e)(\sqrt{-d}+\sqrt{ex})} \\
& -\frac{a + \operatorname{barccosh}(cx)}{16\sqrt{-de}e^{3/2}(\sqrt{-d}-\sqrt{ex})^2} - \frac{a + \operatorname{barccosh}(cx)}{16de^{3/2}(\sqrt{-d}-\sqrt{ex})} \\
& +\frac{a + \operatorname{barccosh}(cx)}{16\sqrt{-de}e^{3/2}(\sqrt{-d}+\sqrt{ex})^2} + \frac{a + \operatorname{barccosh}(cx)}{16de^{3/2}(\sqrt{-d}+\sqrt{ex})} \\
& +\frac{bc^3 \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}\right)}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(c\sqrt{-d}+\sqrt{e})^{3/2}e^{3/2}} \\
& +\frac{b \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}\right)}{8d\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}e^{3/2}} \\
& -\frac{bc^3 \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}\right)}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(c\sqrt{-d}+\sqrt{e})^{3/2}e^{3/2}} \\
& -\frac{b \operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}\right)}{8d\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}e^{3/2}} \\
& -\frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{3/2}e^{3/2}} \\
& +\frac{(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{3/2}e^{3/2}} \\
& -\frac{(a + \operatorname{barccosh}(cx)) \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16(-d)^{3/2}e^{3/2}} \\
& +\frac{(a + \operatorname{barccosh}(cx)) \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16(-d)^{3/2}e^{3/2}} \\
& +\frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{3/2}e^{3/2}} \\
& -\frac{b \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{3/2}e^{3/2}} \\
3.512. \quad \int \frac{x^2(a+b\operatorname{arccosh}(cx))}{(d+ex^2)^3} dx & +\frac{b \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16(-d)^{3/2}e^{3/2}}
\end{aligned}$$

output

```

-1/16*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c
*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(3/2)+1/16*(a+b*arccosh(c*x))*
ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1
/2)))/(-d)^(3/2)/e^(3/2)-1/16*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(
c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(3/2)+
1/16*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*
(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(3/2)+1/16*b*polylog(2,-(c*x+(c
*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(
3/2)/e^(3/2)-1/16*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c
*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(3/2)+1/16*b*polylog(2,-(c*x+(
c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(
3/2)/e^(3/2)-1/16*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(
c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(3/2)/e^(3/2)+1/8*b*c^3*arctanh((c*x+
1)^(1/2)*(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*x-1)^(1/2)/(c*(-d)^(1/2)+e^(1/2))
^(1/2))/e^(3/2)/(c*(-d)^(1/2)-e^(1/2))^(3/2)/(c*(-d)^(1/2)+e^(1/2))^(3/2)-
1/8*b*c^3*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)+e^(1/2))^(1/2)/(c*x-1)^(1/2)
/(c*(-d)^(1/2)-e^(1/2))^(1/2))/e^(3/2)/(c*(-d)^(1/2)-e^(1/2))^(3/2)/(c*(-d)
)^(1/2)+e^(1/2))^(3/2)+1/16*(-a-b*arccosh(c*x))/d/e^(3/2)/((-d)^(1/2)-x*e^(1/2)
)+1/16*(a+b*arccosh(c*x))/e^(3/2)/(-d)^(1/2)/((-d)^(1/2)+x*e^(1/2))^2+1...

```

### 3.512.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 11.03 (sec) , antiderivative size = 1193, normalized size of antiderivative = 0.97

$$\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx = -\frac{ax}{4e(d + ex^2)^2} + \frac{ax}{8de(d + ex^2)} + \frac{a \arctan\left(\frac{\sqrt{ex}}{\sqrt{d}}\right)}{8d^{3/2}e^{3/2}}$$

$$+ b \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d} + \sqrt{ex}} + \frac{c \log\left(\frac{2e(i\sqrt{e} + c^2\sqrt{dx} - i\sqrt{-c^2d - e}\sqrt{-1 + cx}\sqrt{1 + cx})}{c\sqrt{-c^2d - e}(\sqrt{d} + i\sqrt{ex})}\right)}{\sqrt{-c^2d - e}} \right) - \frac{\operatorname{arccosh}(cx)}{i\sqrt{d} + \sqrt{ex}} - \frac{c \log\left(\frac{2e(-\sqrt{e} - ic^2\sqrt{dx} + \sqrt{-c^2d - e}\sqrt{-1 + cx})}{c\sqrt{-c^2d - e}(i\sqrt{d} + \sqrt{ex})}\right)}{\sqrt{-c^2d - e}}$$

input `Integrate[(x^2*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

---

3.512.  $\int \frac{x^2(a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$

output

```

-1/4*(a*x)/(e*(d + e*x^2)^2) + (a*x)/(8*d*e*(d + e*x^2)) + (a*ArcTan[(Sqrt
[e]*x)/Sqrt[d]])/(8*d^(3/2)*e^(3/2)) + b*((ArcCosh[c*x]/((-I)*Sqrt[d] + Sq
rt[e]*x) + (c*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*d) - e]*S
qrt[-1 + c*x]*Sqrt[1 + c*x]))]/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*Sqrt[e]*x
)))/Sqrt[-(c^2*d) - e])/(16*d*e^(3/2)) - ((ArcCosh[c*x]/(I*Sqrt[d] + Sqr
t[e]*x)) - (c*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[-(c^2*d) - e]*Sq
rt[-1 + c*x]*Sqrt[1 + c*x]))]/(c*Sqrt[-(c^2*d) - e]*(I*Sqrt[d] + Sqrt[e]*x
)))/Sqrt[-(c^2*d) - e])/(16*d*e^(3/2)) - ((I/16)*((c*Sqrt[-1 + c*x]*Sqrt[1
+ c*x])/((c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*
((-I)*Sqrt[d] + Sqrt[e]*x)^2) + (c^3*Sqrt[d]*(Log[4] + Log[(e*Sqrt[c^2*d +
e]*((-I)*Sqrt[e] - c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1
+ c*x]))]/(c^3*(d + I*Sqrt[d]*Sqrt[e]*x))))/(Sqrt[e]*(c^2*d + e)^(3/2)))/
(Sqrt[d]*e) + ((I/16)*((c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/((c^2*d + e)*(I*Sq
rt[d] + Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*(I*Sqrt[d] + Sqrt[e]*x)^2) - (
c^3*Sqrt[d]*(Log[4] + Log[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[e] + c^2*Sqrt[d]*x
+ Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))]/(c^3*(d - I*Sqrt[d]*Sqrt
[e]*x))))/(Sqrt[e]*(c^2*d + e)^(3/2)))/((Sqrt[d]*e) + ((I/32)*(ArcCosh[c*
x]*(-ArcCosh[c*x] + 2*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x]))/(I*c*Sqrt[d] - Sqr
t[-(c^2*d) - e]]) + Log[1 + (Sqrt[e]*E^ArcCosh[c*x]))/(I*c*Sqrt[d] + Sqrt[-
(c^2*d) - e]]))) + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/((-I)*c*Sqrt[d]...

```

### 3.512.3 Rubi [A] (verified)

Time = 3.11 (sec) , antiderivative size = 1234, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$ , Rules used = {6374, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx \\
 & \quad \downarrow \text{6374} \\
 & \int \left( \frac{a + \operatorname{barccosh}(cx)}{e(d + ex^2)^2} - \frac{d(a + \operatorname{barccosh}(cx))}{e(d + ex^2)^3} \right) dx \\
 & \quad \downarrow \text{2009}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\operatorname{barctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx-1}}}\right) c^3}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(\sqrt{-dc}+\sqrt{e})^{3/2}e^{3/2}} - \frac{\operatorname{barctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right) c^3}{8(c\sqrt{-d}-\sqrt{e})^{3/2}(\sqrt{-dc}+\sqrt{e})^{3/2}e^{3/2}} + \\
& \frac{\operatorname{barctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx-1}}}\right) c}{8d\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{\sqrt{-dc}+\sqrt{e}}e^{3/2}} - \frac{\operatorname{barctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right) c}{8d\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{\sqrt{-dc}+\sqrt{e}}e^{3/2}} - \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1}}{16\sqrt{-de}(dc^2+e)(\sqrt{-d}-\sqrt{ex})} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{16\sqrt{-de}(dc^2+e)(\sqrt{ex}+\sqrt{-d})} - \frac{a+\operatorname{barccosh}(cx)}{16de^{3/2}(\sqrt{-d}-\sqrt{ex})} + \\
& \frac{a+\operatorname{barccosh}(cx)}{16de^{3/2}(\sqrt{ex}+\sqrt{-d})} - \frac{16\sqrt{-de}^{3/2}(\sqrt{-d}-\sqrt{ex})^2}{16\sqrt{-de}^{3/2}(\sqrt{-d}-\sqrt{ex})^2} + \frac{16\sqrt{-de}^{3/2}(\sqrt{ex}+\sqrt{-d})^2}{16\sqrt{-de}^{3/2}(\sqrt{ex}+\sqrt{-d})^2} - \\
& \frac{(a+\operatorname{barccosh}(cx))\log\left(1-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16(-d)^{3/2}e^{3/2}} + \frac{(a+\operatorname{barccosh}(cx))\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{c\sqrt{-d}-\sqrt{-dc^2-e}}+1\right)}{16(-d)^{3/2}e^{3/2}} - \\
& \frac{(a+\operatorname{barccosh}(cx))\log\left(1-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16(-d)^{3/2}e^{3/2}} + \frac{(a+\operatorname{barccosh}(cx))\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{\sqrt{-dc}+\sqrt{-dc^2-e}}+1\right)}{16(-d)^{3/2}e^{3/2}} + \\
& \frac{b\operatorname{PolyLog}\left(2,-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16(-d)^{3/2}e^{3/2}} - \frac{b\operatorname{PolyLog}\left(2,\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16(-d)^{3/2}e^{3/2}} + \\
& \frac{b\operatorname{PolyLog}\left(2,-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16(-d)^{3/2}e^{3/2}} - \frac{b\operatorname{PolyLog}\left(2,\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16(-d)^{3/2}e^{3/2}}
\end{aligned}$$

input `Int[(x^2*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

```

output -1/16*(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(Sqrt[-d]*e*(c^2*d + e)*(Sqrt[-d]
- Sqrt[e]*x)) - (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(16*Sqrt[-d]*e*(c^2*d
+ e)*(Sqrt[-d] + Sqrt[e]*x)) - (a + b*ArcCosh[c*x])/(16*Sqrt[-d]*e^(3/2)*(
Sqrt[-d] - Sqrt[e]*x)^2) - (a + b*ArcCosh[c*x])/(16*d*e^(3/2)*(Sqrt[-d] -
Sqrt[e]*x)) + (a + b*ArcCosh[c*x])/(16*Sqrt[-d]*e^(3/2)*(Sqrt[-d] + Sqrt[e
]*x)^2) + (a + b*ArcCosh[c*x])/(16*d*e^(3/2)*(Sqrt[-d] + Sqrt[e]*x)) + (b*
c^3*ArcTanh[(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] +
Sqrt[e]]*Sqrt[-1 + c*x])])/(8*(c*Sqrt[-d] - Sqrt[e])^(3/2)*(c*Sqrt[-d] + S
qrt[e])^(3/2)*e^(3/2)) + (b*c*ArcTanh[(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 +
c*x])/(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[-1 + c*x])])/(8*d*Sqrt[c*Sqrt[-d]
- Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*e^(3/2)) - (b*c^3*ArcTanh[(Sqrt[c*Sq
rt[-d] + Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[-1 + c*x
])])/(8*(c*Sqrt[-d] - Sqrt[e])^(3/2)*(c*Sqrt[-d] + Sqrt[e])^(3/2)*e^(3/2))
- (b*c*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d]
- Sqrt[e]]*Sqrt[-1 + c*x])])/(8*d*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt
[-d] + Sqrt[e]]*e^(3/2)) - ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCos
h[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(16*(-d)^(3/2)*e^(3/2)) + ((a
+ b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^
2*d) - e])])/(16*(-d)^(3/2)*e^(3/2)) - ((a + b*ArcCosh[c*x])*Log[1 - (Sqrt
[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(16*(-d)^(3/2)*...

```

### 3.512.3.1 Defintions of rubi rules used

```

rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]

```

```

rule 6374 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e
_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n,
(f*x)^m*(d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[c^2*d
+ e, 0] && IGtQ[n, 0] && IntegerQ[p] && IntegerQ[m]

```

### 3.512.4 Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 23.20 (sec) , antiderivative size = 1222, normalized size of antiderivative = 0.99

method	result	size
parts	Expression too large to display	1222
derivativedivides	Expression too large to display	1237
default	Expression too large to display	1237

```
input int(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)
```

```
output a*((1/8/d*x^3-1/8/e*x)/(e*x^2+d)^2+1/8/e/d/(d*e)^(1/2)*arctan(e*x/(d*e)^(1/2)))+b/c^3*(1/8*c^4*((c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*d*e*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^4*d^2+arccosh(c*x)*d*c^5*e*x^3-arccosh(c*x)*d^2*c^5*x+arccosh(c*x)*e^2*c^3*x^3-arccosh(c*x)*d*c^3*e*x)/e/(c^2*e*x^2+c^2*d)^2/d/(c^2*d+e)+1/16/(c^2*d+e)/d*c^4*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/16/(c^2*d+e)/e*c^6*sum(_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)+dilog(( _R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+1/8*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e+(d*c^2*(c^2*d+e))^(1/2)*e)*c^4*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))/(c^2*d+e)^2/d/e^3-1/8*(-(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)-e)*e)^(1/2))*c^4/(c^2*d+e)/d/e^3+1/8*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^2*(c^2*d+e))^(1/2)*e)*c^4*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2))/(c^2*d+e)^2/d/e^3-1/8*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e)^(1/2)*(2*c^2*d-2*(d*c^2*(c^...
```



**3.512.5 Fracas [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(ex^2 + d)^3} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*x^2*arccosh(c*x) + a*x^2)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**3.512.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate(x**2*(a+b*acosh(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**3.512.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Exception raised: ValueError}$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.512.8 Giac [F]**

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)x^2}{(ex^2 + d)^3} dx$$

input `integrate(x^2*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)*x^2/(e*x^2 + d)^3, x)`

**3.512.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{x^2(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{x^2(a + b \operatorname{acosh}(cx))}{(ex^2 + d)^3} dx$$

input `int((x^2*(a + b*acosh(c*x)))/(d + e*x^2)^3,x)`

output `int((x^2*(a + b*acosh(c*x)))/(d + e*x^2)^3, x)`

$$3.513 \quad \int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^3} dx$$

3.513.1 Optimal result . . . . .	3795
3.513.2 Mathematica [C] (warning: unable to verify) . . . . .	3796
3.513.3 Rubi [A] (verified) . . . . .	3797
3.513.4 Maple [C] (warning: unable to verify) . . . . .	3800
3.513.5 Fricas [F] . . . . .	3801
3.513.6 Sympy [F(-1)] . . . . .	3801
3.513.7 Maxima [F(-2)] . . . . .	3801
3.513.8 Giac [F] . . . . .	3802
3.513.9 Mupad [F(-1)] . . . . .	3802

## 3.513.1 Optimal result

Integrand size = 18, antiderivative size = 1234

$$\begin{aligned}
\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^3} dx = & -\frac{bc\sqrt{-1+cx}\sqrt{1+cx}}{16(-d)^{3/2}(c^2d+e)(\sqrt{-d}-\sqrt{ex})} \\
& -\frac{bc\sqrt{-1+cx}\sqrt{1+cx}}{16(-d)^{3/2}(c^2d+e)(\sqrt{-d}+\sqrt{ex})} \\
& -\frac{a + \operatorname{barccosh}(cx)}{16(-d)^{3/2}\sqrt{e}(\sqrt{-d}-\sqrt{ex})^2} - \frac{3(a + \operatorname{barccosh}(cx))}{16d^2\sqrt{e}(\sqrt{-d}-\sqrt{ex})} \\
& + \frac{a + \operatorname{barccosh}(cx)}{16(-d)^{3/2}\sqrt{e}(\sqrt{-d}+\sqrt{ex})^2} + \frac{3(a + \operatorname{barccosh}(cx))}{16d^2\sqrt{e}(\sqrt{-d}+\sqrt{ex})} \\
& -\frac{bc^3\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{-1+cx}}}\right)}{8d(c\sqrt{-d}-\sqrt{e})^{3/2}(c\sqrt{-d}+\sqrt{e})^{3/2}\sqrt{e}} \\
& +\frac{3bc\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{-1+cx}}}\right)}{8d^2\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}\sqrt{e}} \\
& +\frac{bc^3\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{-1+cx}}}\right)}{8d(c\sqrt{-d}-\sqrt{e})^{3/2}(c\sqrt{-d}+\sqrt{e})^{3/2}\sqrt{e}} \\
& -\frac{3bc\operatorname{arctanh}\left(\frac{\sqrt{c\sqrt{-d}+\sqrt{e}\sqrt{1+cx}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{-1+cx}}}\right)}{8d^2\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{c\sqrt{-d}+\sqrt{e}}\sqrt{e}} \\
& +\frac{3(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{5/2}\sqrt{e}} \\
& -\frac{3(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{5/2}\sqrt{e}} \\
& +\frac{3(a + \operatorname{barccosh}(cx))\log\left(1 - \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16(-d)^{5/2}\sqrt{e}} \\
& -\frac{3(a + \operatorname{barccosh}(cx))\log\left(1 + \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16(-d)^{5/2}\sqrt{e}} \\
& -\frac{3b\operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{5/2}\sqrt{e}} \\
& +\frac{3b\operatorname{PolyLog}\left(2, \frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}-\sqrt{-c^2d-e}}\right)}{16(-d)^{5/2}\sqrt{e}} \\
\hline
3.513. \quad & \int \frac{a+\operatorname{barccosh}(cx)}{(d+ex^2)^3} dx - \frac{3b\operatorname{PolyLog}\left(2, -\frac{\sqrt{ee}\operatorname{arccosh}(cx)}{c\sqrt{-d}+\sqrt{-c^2d-e}}\right)}{16(-d)^{5/2}\sqrt{e}}
\end{aligned}$$

output `3/16*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)-3/16*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)+3/16*(a+b*arccosh(c*x))*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)-3/16*(a+b*arccosh(c*x))*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)-3/16*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)+3/16*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)+3/16*b*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)+3/16*b*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(5/2)/e^(1/2)-1/8*b*c^3*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)-e^(1/2))^(1/2)/(c*x-1)^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(1/2))/d/(c*(-d)^(1/2)-e^(1/2))^(3/2)/e^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(3/2)+1/8*b*c^3*arctanh((c*x+1)^(1/2)*(c*(-d)^(1/2)+e^(1/2))^(1/2)/(c*x-1)^(1/2)/(c*(-d)^(1/2)-e^(1/2))^(1/2))/d/(c*(-d)^(1/2)-e^(1/2))^(3/2)/e^(1/2)/(c*(-d)^(1/2)+e^(1/2))^(3/2)+1/16*(-a-b*arccosh(c*x))/(-d)^(3/2)/e^(1/2)/((-d)^(1/2)-x*e^(1/2))^2-3/16*(a+b*arccosh(c*x))/d^2/e^(1/2)/((-d)^(1/2)-x*e^(1/2))+1/16*(a+b*arccosh(c*x))/(-d)^(3/2)/e^(1/2)/((-d)^(1/2)+x*e^(1/2))...`

### 3.513.2 Mathematica [C] (warning: unable to verify)

Result contains complex when optimal does not.

Time = 10.35 (sec) , antiderivative size = 1161, normalized size of antiderivative = 0.94

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^3} dx$$

$$= \frac{8ad^{3/2}x}{(d+ex^2)^2} + \frac{12a\sqrt{d}x}{d+ex^2} + \frac{12a \arctan\left(\frac{\sqrt{e}x}{\sqrt{d}}\right)}{\sqrt{e}} + \frac{6b\sqrt{d} \left( \frac{\operatorname{arccosh}(cx)}{-i\sqrt{d}+\sqrt{ex}} + \frac{c \log\left(\frac{2e(i\sqrt{e}+c^2\sqrt{d}x-i\sqrt{-c^2d-e}\sqrt{-1+cx}\sqrt{1+cx})}{c\sqrt{-c^2d-e}(\sqrt{d}+i\sqrt{ex})}\right)}{\sqrt{-c^2d-e}} \right)}{\sqrt{e}} - \frac{6b\sqrt{d} \left( -\frac{\operatorname{arccosh}(cx)}{i\sqrt{d}+\sqrt{ex}} + \frac{c \log\left(\frac{2e(-i\sqrt{e}+c^2\sqrt{d}x+i\sqrt{-c^2d-e}\sqrt{-1+cx}\sqrt{1+cx})}{c\sqrt{-c^2d-e}(\sqrt{d}-i\sqrt{ex})}\right)}{\sqrt{-c^2d-e}} \right)}{\sqrt{e}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(d + e*x^2)^3,x]`

output  $((8*a*d^{(3/2)*x})/(d + e*x^2)^2 + (12*a*Sqrt[d]*x)/(d + e*x^2) + (12*a*ArcTan[(Sqrt[e]*x)/Sqrt[d]])/Sqrt[e] + (6*b*Sqrt[d]*(ArcCosh[c*x]/((-I)*Sqrt[d] + Sqrt[e]*x) + (c*Log[(2*e*(I*Sqrt[e] + c^2*Sqrt[d]*x - I*Sqrt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) - e]*(Sqrt[d] + I*Sqrt[e]*x))])/Sqrt[-(c^2*d) - e])/Sqrt[e] - (6*b*Sqrt[d]*(-ArcCosh[c*x]/(I*Sqrt[d] + Sqrt[e]*x)) - (c*Log[(2*e*(-Sqrt[e] - I*c^2*Sqrt[d]*x + Sqrt[-(c^2*d) - e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c*Sqrt[-(c^2*d) - e]*(I*Sqrt[d] + Sqrt[e]*x))])/Sqrt[-(c^2*d) - e])/Sqrt[e] + (2*I)*b*d*((c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/((c^2*d + e)*((-I)*Sqrt[d] + Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*((-I)*Sqrt[d] + Sqrt[e]*x)^2) + (c^3*Sqrt[d]*(Log[4] + Log[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[e] - c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c^3*(d + I*Sqrt[d]*Sqrt[e]*x))]))/(Sqrt[e]*(c^2*d + e)^(3/2)) - (2*I)*b*d*((c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/((c^2*d + e)*(I*Sqrt[d] + Sqrt[e]*x)) - ArcCosh[c*x]/(Sqrt[e]*(I*Sqrt[d] + Sqrt[e]*x)^2) - (c^3*Sqrt[d]*(Log[4] + Log[(e*Sqrt[c^2*d + e]*((-I)*Sqrt[e] + c^2*Sqrt[d]*x + Sqrt[c^2*d + e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(c^3*(d - I*Sqrt[d]*Sqrt[e]*x))]))/(Sqrt[e]*(c^2*d + e)^(3/2)) + ((3*I)*b*(ArcCosh[c*x]*(-ArcCosh[c*x] + 2*(Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] - Sqrt[-(c^2*d) - e]]) + Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(I*c*Sqrt[d] + Sqrt[-(c^2*d) - e]]) + 2*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/((-I)*c*Sqrt[d] + Sqrt[-(c^2...$

### 3.513.3 Rubi [A] (verified)

Time = 1.99 (sec) , antiderivative size = 1234, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \text{barccosh}(cx)}{(d + ex^2)^3} dx$$

↓ 6324

$$\int \left( -\frac{3e(a + \text{barccosh}(cx))}{8d^2(-de - e^2x^2)} - \frac{3e(a + \text{barccosh}(cx))}{16d^2(\sqrt{-d}\sqrt{e} - ex)^2} - \frac{3e(a + \text{barccosh}(cx))}{16d^2(\sqrt{-d}\sqrt{e} + ex)^2} - \frac{e^{3/2}(a + \text{barccosh}(cx))}{8(-d)^{3/2}(\sqrt{-d}\sqrt{e} - ex)^3} - \frac{e^3}{8(-d)^{3/2}(\sqrt{-d}\sqrt{e} + ex)^3} \right) dx$$

↓ 2009

$$\begin{aligned}
& - \frac{\operatorname{barctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{-dc+\sqrt{e}\sqrt{cx-1}}}\right) c^3}{8d(c\sqrt{-d}-\sqrt{e})^{3/2}(\sqrt{-dc}+\sqrt{e})^{3/2}\sqrt{e}} + \frac{\operatorname{barctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right) c^3}{8d(c\sqrt{-d}-\sqrt{e})^{3/2}(\sqrt{-dc}+\sqrt{e})^{3/2}\sqrt{e}} + \\
& \frac{3\operatorname{barctanh}\left(\frac{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx+1}}}{\sqrt{-dc+\sqrt{e}\sqrt{cx-1}}}\right) c}{8d^2\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{e}}} - \frac{3\operatorname{barctanh}\left(\frac{\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{cx+1}}}{\sqrt{c\sqrt{-d}-\sqrt{e}\sqrt{cx-1}}}\right) c}{8d^2\sqrt{c\sqrt{-d}-\sqrt{e}}\sqrt{\sqrt{-dc}+\sqrt{e}\sqrt{e}}} - \\
& \frac{b\sqrt{cx-1}\sqrt{cx+1}}{16(-d)^{3/2}(dc^2+e)(\sqrt{-d}-\sqrt{ex})} - \frac{b\sqrt{cx-1}\sqrt{cx+1}}{16(-d)^{3/2}(dc^2+e)(\sqrt{ex}+\sqrt{-d})} - \frac{3(a+\operatorname{barccosh}(cx))}{16d^2\sqrt{e}(\sqrt{-d}-\sqrt{ex})} + \\
& \frac{3(a+\operatorname{barccosh}(cx))}{16d^2\sqrt{e}(\sqrt{ex}+\sqrt{-d})} - \frac{3(a+\operatorname{barccosh}(cx))}{16(-d)^{3/2}\sqrt{e}(\sqrt{-d}-\sqrt{ex})^2} + \frac{3(a+\operatorname{barccosh}(cx))}{16(-d)^{3/2}\sqrt{e}(\sqrt{ex}+\sqrt{-d})^2} + \\
& \frac{3(a+\operatorname{barccosh}(cx))\log\left(1-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16(-d)^{5/2}\sqrt{e}} - \frac{3(a+\operatorname{barccosh}(cx))\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{c\sqrt{-d}-\sqrt{-dc^2-e}}+1\right)}{16(-d)^{5/2}\sqrt{e}} + \\
& \frac{3(a+\operatorname{barccosh}(cx))\log\left(1-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16(-d)^{5/2}\sqrt{e}} - \frac{3(a+\operatorname{barccosh}(cx))\log\left(\frac{e^{\operatorname{arccosh}(cx)}\sqrt{e}}{\sqrt{-dc}+\sqrt{-dc^2-e}}+1\right)}{16(-d)^{5/2}\sqrt{e}} - \\
& \frac{3b\operatorname{PolyLog}\left(2,-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16(-d)^{5/2}\sqrt{e}} + \frac{3b\operatorname{PolyLog}\left(2,\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{16(-d)^{5/2}\sqrt{e}} - \\
& \frac{3b\operatorname{PolyLog}\left(2,-\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16(-d)^{5/2}\sqrt{e}} + \frac{3b\operatorname{PolyLog}\left(2,\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{16(-d)^{5/2}\sqrt{e}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d + e*x^2)^3,x]`

output

```

-1/16*(b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/((-d)^(3/2)*(c^2*d + e)*(Sqrt[-d]
- Sqrt[e]*x)) - (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(16*(-d)^(3/2)*(c^2*d
+ e)*(Sqrt[-d] + Sqrt[e]*x)) - (a + b*ArcCosh[c*x])/(16*(-d)^(3/2)*Sqrt[e]
*(Sqrt[-d] - Sqrt[e]*x)^2) - (3*(a + b*ArcCosh[c*x]))/(16*d^2*Sqrt[e]*(Sqr
t[-d] - Sqrt[e]*x)) + (a + b*ArcCosh[c*x])/(16*(-d)^(3/2)*Sqrt[e]*(Sqrt[-d]
+ Sqrt[e]*x)^2) + (3*(a + b*ArcCosh[c*x]))/(16*d^2*Sqrt[e]*(Sqrt[-d] + S
qrt[e]*x)) - (b*c^3*ArcTanh[(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[1 + c*x])/(Sq
rt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[-1 + c*x])])/(8*d*(c*Sqrt[-d] - Sqrt[e])^(3/
2)*(c*Sqrt[-d] + Sqrt[e])^(3/2)*Sqrt[e]) + (3*b*c*ArcTanh[(Sqrt[c*Sqrt[-d]
- Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[-1 + c*x])])/(
8*d^2*Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[e]) + (b*
c^3*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[1 + c*x])/(Sqrt[c*Sqrt[-d] -
Sqrt[e]]*Sqrt[-1 + c*x])])/(8*d*(c*Sqrt[-d] - Sqrt[e])^(3/2)*(c*Sqrt[-d] +
Sqrt[e])^(3/2)*Sqrt[e]) - (3*b*c*ArcTanh[(Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt
[1 + c*x])/(Sqrt[c*Sqrt[-d] - Sqrt[e]]*Sqrt[-1 + c*x])])/(8*d^2*Sqrt[c*Sqr
t[-d] - Sqrt[e]]*Sqrt[c*Sqrt[-d] + Sqrt[e]]*Sqrt[e]) + (3*(a + b*ArcCosh[c
*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(
16*(-d)^(5/2)*Sqrt[e]) - (3*(a + b*ArcCosh[c*x])*Log[1 + (Sqrt[e]*E^ArcCo
sh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(16*(-d)^(5/2)*Sqrt[e]) + (3*
(a + b*ArcCosh[c*x])*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqr...

```

### 3.513.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`



**3.513.4 Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 29.03 (sec) , antiderivative size = 1778, normalized size of antiderivative = 1.44

method	result	size
parts	Expression too large to display	1778
derivativedivides	Expression too large to display	1803
default	Expression too large to display	1803

```
input int((a+b*arccosh(c*x))/(e*x^2+d)^3,x,method=_RETURNVERBOSE)
```

```
output 1/4*a*x/d/(e*x^2+d)^2+3/8*a/d^2*x/(e*x^2+d)+3/8*a/d^2/(d*e)^(1/2)*arctan(e
*x/(d*e)^(1/2))+b/c*(1/8*c^2*(5*arccosh(c*x)*d^2*c^5*x+3*arccosh(c*x)*d*c^
5*e*x^3-(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^4*d^2-(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^
4*d*e*x^2+5*arccosh(c*x)*d*c^3*e*x+3*arccosh(c*x)*e^2*c^3*x^3)/d^2/(c^2*e*
x^2+c^2*d)^2/(c^2*d+e)+3/8*(-2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/2)+e)*e^(1/2
)*(2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e+(d*c^2*(c^2*d+e))^(
1/2)*e)*c^2*arctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((-2*c^2*d+2*(d*c^
2*(c^2*d+e))^(1/2)-e)*e^(1/2))/(c^2*d+e)^2/d^2/e^2-3/8*(-2*c^2*d-2*(d*c^
2*(c^2*d+e))^(1/2)+e)*e^(1/2)*(2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*c^2*a
rctanh(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((-2*c^2*d+2*(d*c^2*(c^2*d+e))^(
1/2)-e)*e^(1/2))/(c^2*d+e)/d^2/e^2+3/8*((2*c^2*d+2*(d*c^2*(c^2*d+e))^(1/
2)+e)*e^(1/2)*(-2*(d*c^2*(c^2*d+e))^(1/2)*c^2*d+2*c^4*d^2+2*c^2*d*e-(d*c^
2*(c^2*d+e))^(1/2)*e)*c^2*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((2*c^
2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e^(1/2))/(c^2*d+e)^2/d^2/e^2-3/8*((2*c^
2*d+2*(d*c^2*(c^2*d+e))^(1/2)+e)*e^(1/2)*(2*c^2*d-2*(d*c^2*(c^2*d+e))^(1/
2)+e)*c^2*arctan(e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2)))/((2*c^2*d+2*(d*c^2*(c
^2*d+e))^(1/2)+e)*e^(1/2))/(c^2*d+e)/d^2/e^2-3/16/(c^2*d+e)/d^2*c^2*e*sum
(1/_R1/(_R1^2*e+2*c^2*d+e)*(arccosh(c*x)*ln((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)
^(1/2))/_R1)+dilog((_R1-c*x-(c*x-1)^(1/2)*(c*x+1)^(1/2))/_R1)),_R1=RootOf(
e*_Z^4+(4*c^2*d+2*e)*_Z^2+e))+3/16/(c^2*d+e)/d^2*c^2*e*sum(_R1/(_R1^2*e...
```

**3.513.5 Fricas [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^3} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^3} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**3.513.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**3.513.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^3} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.513.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^3} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^3} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(e*x^2 + d)^3, x)`

**3.513.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^3} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(ex^2 + d)^3} dx$$

input `int((a + b*acosh(c*x))/(d + e*x^2)^3,x)`

output `int((a + b*acosh(c*x))/(d + e*x^2)^3, x)`

### 3.514 $\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx$

3.514.1 Optimal result	3803
3.514.2 Mathematica [N/A]	3803
3.514.3 Rubi [N/A]	3804
3.514.4 Maple [N/A] (verified)	3804
3.514.5 Fricas [N/A]	3805
3.514.6 Sympy [N/A]	3805
3.514.7 Maxima [F(-2)]	3805
3.514.8 Giac [N/A]	3806
3.514.9 Mupad [N/A]	3806

#### 3.514.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx = \operatorname{Int}\left(\sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)), x\right)$$

output `Unintegrable((a+b*arccosh(c*x))*(e*x^2+d)^(1/2),x)`

#### 3.514.2 Mathematica [N/A]

Not integrable

Time = 6.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx = \int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx$$

input `Integrate[Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x]),x]`

output `Integrate[Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x]), x]`

**3.514.3 Rubi [N/A]**

Not integrable

Time = 0.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d + ex^2}(a + \operatorname{arccosh}(cx)) dx$$

↓ 6325

$$\int \sqrt{d + ex^2}(a + \operatorname{arccosh}(cx)) dx$$

input `Int[Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x]),x]`

output `$Aborted`

**3.514.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.514.4 Maple [N/A] (verified)**

Not integrable

Time = 1.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int (a + b \operatorname{arccosh}(cx)) \sqrt{ex^2 + d} dx$$

input `int((a+b*arccosh(c*x))*(e*x^2+d)^(1/2),x)`

output `int((a+b*arccosh(c*x))*(e*x^2+d)^(1/2),x)`

**3.514.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx = \int \sqrt{ex^2 + d}(b \operatorname{arcosh}(cx) + a) dx$$

input `integrate((a+b*arccosh(c*x))*(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)*(b*arccosh(c*x) + a), x)`

**3.514.6 Sympy [N/A]**

Not integrable

Time = 1.72 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) \sqrt{d + ex^2} dx$$

input `integrate((a+b*acosh(c*x))*(e*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))*sqrt(d + e*x**2), x)`

**3.514.7 Maxima [F(-2)]**

Exception generated.

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))*(e*x^2+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.514.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx = \int \sqrt{ex^2 + d}(b \operatorname{arcosh}(cx) + a) dx$$

input `integrate((a+b*arccosh(c*x))*(e*x^2+d)^(1/2),x, algorithm="giac")`output `integrate(sqrt(e*x^2 + d)*(b*arccosh(c*x) + a), x)`**3.514.9 Mupad [N/A]**

Not integrable

Time = 3.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) \sqrt{ex^2 + d} dx$$

input `int((a + b*acosh(c*x))*(d + e*x^2)^(1/2),x)`output `int((a + b*acosh(c*x))*(d + e*x^2)^(1/2), x)`

### 3.515 $\int \frac{a+b\operatorname{arccosh}(cx)}{\sqrt{d+ex^2}} dx$

3.515.1 Optimal result	3807
3.515.2 Mathematica [N/A]	3807
3.515.3 Rubi [N/A]	3808
3.515.4 Maple [N/A] (verified)	3808
3.515.5 Fricas [N/A]	3809
3.515.6 Sympy [N/A]	3809
3.515.7 Maxima [F(-2)]	3809
3.515.8 Giac [N/A]	3810
3.515.9 Mupad [N/A]	3810

#### 3.515.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{a + b\operatorname{arccosh}(cx)}{\sqrt{d + ex^2}} dx = \operatorname{Int}\left(\frac{a + b\operatorname{arccosh}(cx)}{\sqrt{d + ex^2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x)`

#### 3.515.2 Mathematica [N/A]

Not integrable

Time = 2.91 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{a + b\operatorname{arccosh}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{a + b\operatorname{arccosh}(cx)}{\sqrt{d + ex^2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])/Sqrt[d + e*x^2],x]`

output `Integrate[(a + b*ArcCosh[c*x])/Sqrt[d + e*x^2], x]`



**3.515.3 Rubi [N/A]**

Not integrable

Time = 0.20 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d + ex^2}} dx$$

↓ 6325

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d + ex^2}} dx$$

input `Int[(a + b*ArcCosh[c*x])/Sqrt[d + e*x^2], x]`

output `$Aborted`

**3.515.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.515.4 Maple [N/A] (verified)**

Not integrable

Time = 1.16 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{ex^2 + d}} dx$$

input `int((a+b*arccosh(c*x))/(e*x^2+d)^(1/2), x)`

output `int((a+b*arccosh(c*x))/(e*x^2+d)^(1/2), x)`

**3.515.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{ex^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")`output `integral((b*arccosh(c*x) + a)/sqrt(e*x^2 + d), x)`**3.515.6 Sympy [N/A]**

Not integrable

Time = 1.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{\sqrt{d + ex^2}} dx$$

input `integrate((a+b*acosh(c*x))/(e*x**2+d)**(1/2),x)`output `Integral((a + b*acosh(c*x))/sqrt(d + e*x**2), x)`**3.515.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{a + \operatorname{barccosh}(cx)}{\sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.515.8 Giac [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{\sqrt{ex^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)/sqrt(e*x^2 + d), x)`**3.515.9 Mupad [N/A]**

Not integrable

Time = 3.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{a + b \operatorname{arccosh}(cx)}{\sqrt{d + ex^2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{\sqrt{ex^2 + d}} dx$$

input `int((a + b*acosh(c*x))/(d + e*x^2)^(1/2),x)`output `int((a + b*acosh(c*x))/(d + e*x^2)^(1/2), x)`

# 3.516 $\int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^{3/2}} dx$

3.516.1 Optimal result . . . . .	3811
3.516.2 Mathematica [C] (verified) . . . . .	3811
3.516.3 Rubi [A] (verified) . . . . .	3812
3.516.4 Maple [F] . . . . .	3814
3.516.5 Fricas [A] (verification not implemented) . . . . .	3814
3.516.6 Sympy [F] . . . . .	3815
3.516.7 Maxima [F(-2)] . . . . .	3815
3.516.8 Giac [F] . . . . .	3816
3.516.9 Mupad [F(-1)] . . . . .	3816

## 3.516.1 Optimal result

Integrand size = 20, antiderivative size = 101

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d + ex^2)^{3/2}} dx = \frac{x(a + \operatorname{arccosh}(cx))}{d\sqrt{d + ex^2}} - \frac{b\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{-1+c^2x^2}}{c\sqrt{d+ex^2}}\right)}{d\sqrt{e}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output -b*arctanh(e^(1/2)*(c^2*x^2-1)^(1/2)/c/(e*x^2+d)^(1/2))*(c^2*x^2-1)^(1/2)/
d/e^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+x*(a+b*arccosh(c*x))/d/(e*x^2+d)^(1/2)
```

## 3.516.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 13.47 (sec) , antiderivative size = 556, normalized size of antiderivative = 5.50

$$\int \frac{a + \operatorname{arccosh}(cx)}{(d + ex^2)^{3/2}} dx = ax + b\operatorname{arccosh}(cx) + \frac{2b(-1+cx)^{3/2} \sqrt{\frac{(c\sqrt{d}-i\sqrt{e})(1+cx)}{(c\sqrt{d}+i\sqrt{e})(-1+cx)}}}{c(-ic\sqrt{d}+\sqrt{e})(i\sqrt{d}+\sqrt{ex}) \sqrt{\frac{1+\frac{ic\sqrt{d}}{\sqrt{e}}-cx+\frac{i\sqrt{e}}{\sqrt{e}}}{1-cx}}}$$

input `Integrate[(a + b*ArcCosh[c*x])/(d + e*x^2)^(3/2),x]`

output `(a*x + b*x*ArcCosh[c*x] + (2*b*(-1 + c*x)^(3/2)*Sqrt[((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))/((c*Sqrt[d] + I*Sqrt[e])*(-1 + c*x))]*((c*(-I)*c*Sqrt[d] + Sqrt[e])*(I*Sqrt[d] + Sqrt[e]*x)*Sqrt[(1 + (I*c*Sqrt[d])/Sqrt[e] - c*x + (I*Sqrt[e]*x)/Sqrt[d])/(1 - c*x)]*EllipticF[ArcSin[Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(2 - 2*c*x))]], ((4*I)*c*Sqrt[d]*Sqrt[e])/(c*Sqrt[d] + I*Sqrt[e])^2)/(-1 + c*x) + c*Sqrt[d]*(-(c*Sqrt[d]) + I*Sqrt[e])*Sqrt[((c^2*d + e)*(d + e*x^2))/(d*e*(-1 + c*x)^2)]*Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(1 - c*x))]*EllipticPi[(2*c*Sqrt[d])/(c*Sqrt[d] + I*Sqrt[e]), ArcSin[Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(2 - 2*c*x))]], ((4*I)*c*Sqrt[d]*Sqrt[e])/(c*Sqrt[d] + I*Sqrt[e])^2))/((c*(c^2*d + e)*Sqrt[1 + c*x]*Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(1 - c*x))])/((d*Sqrt[d + e*x^2]))`

### 3.516.3 Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {6323, 27, 2038, 353, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{3/2}} dx \\
 & \quad \downarrow \text{6323} \\
 & \frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d + ex^2}} - bc \int \frac{x}{d\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{ex^2 + d}} dx \\
 & \quad \downarrow \text{27} \\
 & \frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d + ex^2}} - \frac{bc \int \frac{x}{\sqrt{cx - 1}\sqrt{cx + 1}\sqrt{ex^2 + d}} dx}{d} \\
 & \quad \downarrow \text{2038} \\
 & \frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d + ex^2}} - \frac{bc\sqrt{c^2x^2 - 1} \int \frac{x}{\sqrt{c^2x^2 - 1}\sqrt{ex^2 + d}} dx}{d\sqrt{cx - 1}\sqrt{cx + 1}} \\
 & \quad \downarrow \text{353}
 \end{aligned}$$

---

3.516.  $\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{3/2}} dx$

$$\frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d + ex^2}} - \frac{bc\sqrt{c^2x^2 - 1} \int \frac{1}{\sqrt{c^2x^2 - 1}\sqrt{ex^2 + d}} dx^2}{2d\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 66

$$\frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d + ex^2}} - \frac{bc\sqrt{c^2x^2 - 1} \int \frac{1}{c^2 - ex^4} d\frac{\sqrt{c^2x^2 - 1}}{\sqrt{ex^2 + d}}}{d\sqrt{cx - 1}\sqrt{cx + 1}}$$

↓ 221

$$\frac{x(a + \operatorname{barccosh}(cx))}{d\sqrt{d + ex^2}} - \frac{b\sqrt{c^2x^2 - 1} \operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{c^2x^2 - 1}}{c\sqrt{d + ex^2}}\right)}{d\sqrt{e}\sqrt{cx - 1}\sqrt{cx + 1}}$$

input `Int[(a + b*ArcCosh[c*x])/(d + e*x^2)^(3/2),x]`

output `(x*(a + b*ArcCosh[c*x]))/(d*Sqrt[d + e*x^2]) - (b*Sqrt[-1 + c^2*x^2]*ArcTanh[(Sqrt[e]*Sqrt[-1 + c^2*x^2])/(c*Sqrt[d + e*x^2])])/(d*Sqrt[e]*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

### 3.516.3.1 Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]`

rule 66 `Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 353 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]`

```
rule 2038 Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[(a1 + b1*x^(n/2))^FracPart[p]*((a2 + b2*x^(n/2))^FracPart[p]/(a1*a2 + b1*b2*x^n)^FracPart[p]) Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && !(EqQ[n, 2] && IGtQ[q, 0])
```

```
rule 6323 Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d + e, 0] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])
```

### 3.516.4 Maple [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(ex^2 + d)^{\frac{3}{2}}} dx$$

```
input int((a+b*arccosh(c*x))/(e*x^2+d)^(3/2), x)
```

```
output int((a+b*arccosh(c*x))/(e*x^2+d)^(3/2), x)
```

### 3.516.5 Fracas [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.29

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^{3/2}} dx = \left[ \frac{4 \sqrt{ex^2 + d} b e x \log(cx + \sqrt{c^2 x^2 - 1}) + 4 \sqrt{ex^2 + d} a e x + (b e x^2 + b d) \sqrt{e} \log(8 c^4}{4} \right]$$

```
input integrate((a+b*arccosh(c*x))/(e*x^2+d)^(3/2), x, algorithm="fracas")
```

```
output [1/4*(4*sqrt(e*x^2 + d)*b*e*x*log(c*x + sqrt(c^2*x^2 - 1)) + 4*sqrt(e*x^2
+ d)*a*e*x + (b*e*x^2 + b*d)*sqrt(e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d
*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^3*e*x^2 + c^3*d - c*e)*sqrt(c^2*x^
2 - 1)*sqrt(e*x^2 + d)*sqrt(e + e^2))/(d*e^2*x^2 + d^2*e), 1/2*(2*sqrt(e*
x^2 + d)*b*e*x*log(c*x + sqrt(c^2*x^2 - 1)) + 2*sqrt(e*x^2 + d)*a*e*x + (b
*e*x^2 + b*d)*sqrt(-e)*arctan(1/2*(2*c^2*e*x^2 + c^2*d - e)*sqrt(c^2*x^2 -
1)*sqrt(e*x^2 + d)*sqrt(-e)/(c^3*e^2*x^4 - c*d*e + (c^3*d*e - c*e^2)*x^2
))/(d*e^2*x^2 + d^2*e)]
```

### 3.516.6 Sympy [F]

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(d + ex^2)^{3/2}} dx$$

```
input integrate((a+b*acosh(c*x))/(e*x**2+d)**(3/2),x)
```

```
output Integral((a + b*acosh(c*x))/(d + e*x**2)**(3/2), x)
```

### 3.516.7 Maxima [F(-2)]

Exception generated.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

```
input integrate((a+b*arccosh(c*x))/(e*x^2+d)^(3/2),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e+c^2*d>0)', see `assume?` for m
ore detail
```



**3.516.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(e*x^2 + d)^(3/2), x)`

**3.516.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{3/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(ex^2 + d)^{3/2}} dx$$

input `int((a + b*acosh(c*x))/(d + e*x^2)^(3/2),x)`

output `int((a + b*acosh(c*x))/(d + e*x^2)^(3/2), x)`

# 3.517 $\int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^{5/2}} dx$

3.517.1 Optimal result . . . . .	3817
3.517.2 Mathematica [C] (verified) . . . . .	3817
3.517.3 Rubi [A] (verified) . . . . .	3818
3.517.4 Maple [F] . . . . .	3821
3.517.5 Fricas [B] (verification not implemented) . . . . .	3821
3.517.6 Sympy [F] . . . . .	3822
3.517.7 Maxima [F] . . . . .	3822
3.517.8 Giac [F] . . . . .	3822
3.517.9 Mupad [F(-1)] . . . . .	3823

## 3.517.1 Optimal result

Integrand size = 20, antiderivative size = 182

$$\int \frac{a + b\operatorname{arccosh}(cx)}{(d + ex^2)^{5/2}} dx = -\frac{bc\sqrt{-1 + cx}\sqrt{1 + cx}}{3d(c^2d + e)\sqrt{d + ex^2}} + \frac{x(a + b\operatorname{arccosh}(cx))}{3d(d + ex^2)^{3/2}} + \frac{2x(a + b\operatorname{arccosh}(cx))}{3d^2\sqrt{d + ex^2}} + \frac{2b\sqrt{1 - c^2x^2} \arctan\left(\frac{\sqrt{e}\sqrt{1 - c^2x^2}}{c\sqrt{d + ex^2}}\right)}{3d^2\sqrt{e}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

```
output 1/3*x*(a+b*arccosh(c*x))/d/(e*x^2+d)^(3/2)+2/3*b*arctan(e^(1/2)*(-c^2*x^2+1)^(1/2)/c/(e*x^2+d)^(1/2))*(-c^2*x^2+1)^(1/2)/d^2/e^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+2/3*x*(a+b*arccosh(c*x))/d^2/(e*x^2+d)^(1/2)-1/3*b*c*(c*x-1)^(1/2)*(c*x+1)^(1/2)/d/(c^2*d+e)/(e*x^2+d)^(1/2)
```

## 3.517.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.  
Time = 1.87 (sec) , antiderivative size = 633, normalized size of antiderivative = 3.48

$$\int \frac{a + b\operatorname{arccosh}(cx)}{(d + ex^2)^{5/2}} dx = -\frac{bc\sqrt{-1+cx}\sqrt{1+cx}(d+ex^2)}{d(c^2d+e)} + \frac{ax(3d+2ex^2)}{d^2} + \frac{bx(3d+2ex^2)\operatorname{arccosh}(cx)}{d^2} + \frac{4b(-1+cx)^{3/2}\sqrt{\frac{c\sqrt{d-ix}\sqrt{d+ix}}{c\sqrt{d+ix}\sqrt{d-ix}}}}{d^2}$$

3.517.  $\int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^{5/2}} dx$

input `Integrate[(a + b*ArcCosh[c*x])/(d + e*x^2)^(5/2),x]`

output `(-((b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(d + e*x^2))/(d*(c^2*d + e))) + (a*x*(3*d + 2*e*x^2))/d^2 + (b*x*(3*d + 2*e*x^2)*ArcCosh[c*x])/d^2 + (4*b*(-1 + c*x)^(3/2)*Sqrt[((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))/((c*Sqrt[d] + I*Sqrt[e])*(-1 + c*x))]*(d + e*x^2)*((c*(-I)*c*Sqrt[d] + Sqrt[e])*(I*Sqrt[d] + Sqrt[e]*x)*Sqrt[(1 + (I*c*Sqrt[d])/Sqrt[e] - c*x + (I*Sqrt[e]*x)/Sqrt[d])/(1 - c*x)]*EllipticF[ArcSin[Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x)))/(2 - 2*c*x))], ((4*I)*c*Sqrt[d]*Sqrt[e])/(c*Sqrt[d] + I*Sqrt[e])^2))/(-1 + c*x) + c*Sqrt[d]*(-(c*Sqrt[d]) + I*Sqrt[e])*Sqrt[((c^2*d + e)*(d + e*x^2))/(d*e*(-1 + c*x)^2)]*Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(1 - c*x))]*EllipticPi[(2*c*Sqrt[d])/(c*Sqrt[d] + I*Sqrt[e]), ArcSin[Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x)))/(2 - 2*c*x))], ((4*I)*c*Sqrt[d]*Sqrt[e])/(c*Sqrt[d] + I*Sqrt[e])^2))/((c*d^2*(c^2*d + e)*Sqrt[1 + c*x]*Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(1 - c*x)))]/(3*(d + e*x^2)^(3/2))`

### 3.517.3 Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$ , Rules used = {6323, 27, 1076, 435, 87, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{5/2}} dx$$

$$\downarrow \text{6323}$$

$$-bc \int \frac{x(2ex^2 + 3d)}{3d^2 \sqrt{cx - 1} \sqrt{cx + 1} (ex^2 + d)^{3/2}} dx + \frac{2x(a + \operatorname{barccosh}(cx))}{3d^2 \sqrt{d + ex^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d (d + ex^2)^{3/2}}$$

$$\downarrow \text{27}$$

$$-\frac{bc \int \frac{x(2ex^2 + 3d)}{\sqrt{cx - 1} \sqrt{cx + 1} (ex^2 + d)^{3/2}} dx}{3d^2} + \frac{2x(a + \operatorname{barccosh}(cx))}{3d^2 \sqrt{d + ex^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d (d + ex^2)^{3/2}}$$

$$\downarrow \text{1076}$$

---

3.517.  $\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{5/2}} dx$

$$\begin{aligned}
& -\frac{bc\sqrt{c^2x^2-1} \int \frac{x(2ex^2+3d)}{\sqrt{c^2x^2-1}(ex^2+d)^{3/2}} dx}{3d^2\sqrt{cx-1}\sqrt{cx+1}} + \frac{2x(a + \operatorname{barccosh}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d+ex^2)^{3/2}} \\
& \quad \downarrow 435 \\
& -\frac{bc\sqrt{c^2x^2-1} \int \frac{2ex^2+3d}{\sqrt{c^2x^2-1}(ex^2+d)^{3/2}} dx^2}{6d^2\sqrt{cx-1}\sqrt{cx+1}} + \frac{2x(a + \operatorname{barccosh}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d+ex^2)^{3/2}} \\
& \quad \downarrow 87 \\
& -\frac{bc\sqrt{c^2x^2-1} \left( 2 \int \frac{1}{\sqrt{c^2x^2-1}\sqrt{ex^2+d}} dx^2 + \frac{2d\sqrt{c^2x^2-1}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{6d^2\sqrt{cx-1}\sqrt{cx+1}} + \frac{2x(a + \operatorname{barccosh}(cx))}{3d^2\sqrt{d+ex^2}} + \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{3d(d+ex^2)^{3/2}} \\
& \quad \downarrow 66 \\
& -\frac{bc\sqrt{c^2x^2-1} \left( 4 \int \frac{1}{c^2-ex^4} d \frac{\sqrt{c^2x^2-1}}{\sqrt{ex^2+d}} + \frac{2d\sqrt{c^2x^2-1}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{6d^2\sqrt{cx-1}\sqrt{cx+1}} + \frac{2x(a + \operatorname{barccosh}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d+ex^2)^{3/2}} \\
& \quad \downarrow 221 \\
& \frac{2x(a + \operatorname{barccosh}(cx))}{3d^2\sqrt{d+ex^2}} + \frac{x(a + \operatorname{barccosh}(cx))}{3d(d+ex^2)^{3/2}} - \\
& \frac{bc\sqrt{c^2x^2-1} \left( \frac{4\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{c^2x^2-1}}{c\sqrt{d+ex^2}}\right)}{c\sqrt{e}} + \frac{2d\sqrt{c^2x^2-1}}{(c^2d+e)\sqrt{d+ex^2}} \right)}{6d^2\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d + e*x^2)^(5/2),x]`

output `(x*(a + b*ArcCosh[c*x]))/(3*d*(d + e*x^2)^(3/2)) + (2*x*(a + b*ArcCosh[c*x]))/(3*d^2*sqrt[d + e*x^2]) - (b*c*sqrt[-1 + c^2*x^2]*((2*d*sqrt[-1 + c^2*x^2]))/((c^2*d + e)*sqrt[d + e*x^2]) + (4*ArcTanh[(sqrt[e]*sqrt[-1 + c^2*x^2])]/(c*sqrt[d + e*x^2]))/(c*sqrt[e]))/(6*d^2*sqrt[-1 + c*x]*sqrt[1 + c*x])`

## 3.517.3.1 Defintions of rubi rules used

- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`
- rule 66 `Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]`
- rule 87 `Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !EqQ[e, 0] || !EqQ[c, 0] || LtQ[p, n])))`
- rule 221 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`
- rule 435 `Int[(x_)^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2)^(q_)*((e_) + (f_)*(x_)^2)^(r_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p*(c + d*x)^q*(e + f*x)^r, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, f, p, q, r}, x] && IntegerQ[(m - 1)/2]`
- rule 1076 `Int[((g_)*(x_)^(m_)*((e1_) + (f1_)*(x_)^(n2_))^(r_)*((e2_) + (f2_)*(x_)^(n2_))^(r_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(e1 + f1*x^(n/2))^FracPart[r]*((e2 + f2*x^(n/2))^FracPart[r]/(e1*e2 + f1*f2*x^n)^FracPart[r]) Int[(g*x)^m*(a + b*x^n)^p*(c + d*x^n)^q*(e1*e2 + f1*f2*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e1, f1, e2, f2, g, m, n, p, q, r}, x] && EqQ[n2, n/2] && EqQ[e2*f1 + e1*f2, 0]`
- rule 6323 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d + e, 0] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])`

**3.517.4 Maple [F]**

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(ex^2 + d)^{\frac{5}{2}}} dx$$

input `int((a+b*arccosh(c*x))/(e*x^2+d)^(5/2),x)`

output `int((a+b*arccosh(c*x))/(e*x^2+d)^(5/2),x)`

**3.517.5 Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 355 vs. 2(150) = 300.

Time = 0.33 (sec) , antiderivative size = 724, normalized size of antiderivative = 3.98

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^{5/2}} dx = \left[ \frac{(bc^2d^3 + (bc^2de^2 + be^3)x^4 + bd^2e + 2(bc^2d^2e + bde^2)x^2)\sqrt{e} \log(8c^4e^2x^4 + c^4d^2 - 6c^2d^2e + 8(c^4d^2e - c^2e^2)x^2 - 4(2c^3e^2x^2 + c^3d - ce)\sqrt{c^2x^2 - 1})\sqrt{ex^2 + d} + e^2 + 2(2(b*c^2*d*e^2 + b*e^3)*x^3 + 3*(b*c^2*d^2*e + b*d*e^2)*x)\sqrt{ex^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) + 2(2(a*c^2*d*e^2 + a*e^3)*x^3 + 3*(a*c^2*d^2*e + a*d*e^2)*x - (b*c*d*e^2*x^2 + b*c*d^2*e)\sqrt{c^2x^2 - 1})\sqrt{ex^2 + d}}{(c^2*d^5*e + d^4*e^2 + (c^2*d^3*e^3 + d^2*e^4)*x^4 + 2*(c^2*d^4*e^2 + d^3*e^3)*x^2), 1/3*((b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)\sqrt{-e}) \operatorname{arctan}(1/2*(2*c^2*e*x^2 + c^2*d - e)\sqrt{c^2x^2 - 1})\sqrt{ex^2 + d} \sqrt{-e}/(c^3e^2x^4 - c*d*e + (c^3d*e - ce^2)x^2) + (2*(b*c^2*d*e^2 + b*e^3)*x^3 + 3*(b*c^2*d^2*e + b*d*e^2)*x)\sqrt{ex^2 + d} \log(cx + \sqrt{c^2x^2 - 1}) + (2*(a*c^2*d*e^2 + a*e^3)*x^3 + 3*(a*c^2*d^2*e + a*d*e^2)*x - (b*c*d*e^2*x^2 + b*c*d^2*e)\sqrt{c^2x^2 - 1})\sqrt{ex^2 + d}}{(c^2*d^5*e + d^4*e^2 + (c^2*d^3*e^3 + d^2*e^4)*x^4 + 2*(c^2*d^4*e^2 + d^3*e^3)*x^2)} \right]$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `[1/6*((b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d^2*e + 8*(c^4*d^2e - c^2*e^2)*x^2 - 4*(2*c^3*e^2*x^2 + c^3*d - c*e)*sqrt(c^2*x^2 - 1))*sqrt(ex^2 + d)*sqrt(e) + e^2 + 2*(2*(b*c^2*d*e^2 + b*e^3)*x^3 + 3*(b*c^2*d^2*e + b*d*e^2)*x)*sqrt(ex^2 + d)*log(cx + sqrt(c^2*x^2 - 1)) + 2*(2*(a*c^2*d*e^2 + a*e^3)*x^3 + 3*(a*c^2*d^2*e + a*d*e^2)*x - (b*c*d*e^2*x^2 + b*c*d^2*e)*sqrt(c^2*x^2 - 1))*sqrt(ex^2 + d))/(c^2*d^5*e + d^4*e^2 + (c^2*d^3*e^3 + d^2*e^4)*x^4 + 2*(c^2*d^4*e^2 + d^3*e^3)*x^2), 1/3*((b*c^2*d^3 + (b*c^2*d*e^2 + b*e^3)*x^4 + b*d^2*e + 2*(b*c^2*d^2*e + b*d*e^2)*x^2)*sqrt(-e)*arctan(1/2*(2*c^2*e*x^2 + c^2*d - e)*sqrt(c^2*x^2 - 1))*sqrt(ex^2 + d)*sqrt(-e)/(c^3*e^2*x^4 - c*d*e + (c^3*d*e - c*e^2)*x^2) + (2*(b*c^2*d*e^2 + b*e^3)*x^3 + 3*(b*c^2*d^2*e + b*d*e^2)*x)*sqrt(ex^2 + d)*log(cx + sqrt(c^2*x^2 - 1)) + (2*(a*c^2*d*e^2 + a*e^3)*x^3 + 3*(a*c^2*d^2*e + a*d*e^2)*x - (b*c*d*e^2*x^2 + b*c*d^2*e)*sqrt(c^2*x^2 - 1))*sqrt(ex^2 + d))/(c^2*d^5*e + d^4*e^2 + (c^2*d^3*e^3 + d^2*e^4)*x^4 + 2*(c^2*d^4*e^2 + d^3*e^3)*x^2)]`

**3.517.6 Sympy [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(d + ex^2)^{5/2}} dx$$

input `integrate((a+b*acosh(c*x))/(e*x**2+d)**(5/2),x)`

output `Integral((a + b*acosh(c*x))/(d + e*x**2)**(5/2), x)`

**3.517.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/3*a*(2*x/(sqrt(e*x^2 + d)*d^2) + x/((e*x^2 + d)^(3/2)*d)) + b*integrate(log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(e*x^2 + d)^(5/2), x)`

**3.517.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(5/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(e*x^2 + d)^(5/2), x)`

**3.517.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^{5/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(ex^2 + d)^{5/2}} dx$$

input `int((a + b*acosh(c*x))/(d + e*x^2)^(5/2), x)`output `int((a + b*acosh(c*x))/(d + e*x^2)^(5/2), x)`



# 3.518 $\int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^{7/2}} dx$

3.518.1 Optimal result . . . . .	3824
3.518.2 Mathematica [C] (verified) . . . . .	3825
3.518.3 Rubi [A] (verified) . . . . .	3825
3.518.4 Maple [F] . . . . .	3829
3.518.5 Fricas [B] (verification not implemented) . . . . .	3829
3.518.6 Sympy [F(-1)] . . . . .	3830
3.518.7 Maxima [F] . . . . .	3831
3.518.8 Giac [F] . . . . .	3831
3.518.9 Mupad [F(-1)] . . . . .	3831

## 3.518.1 Optimal result

Integrand size = 20, antiderivative size = 284

$$\int \frac{a + b\operatorname{arccosh}(cx)}{(d + ex^2)^{7/2}} dx = \frac{bc(1 - c^2x^2)}{15d(c^2d + e)\sqrt{-1 + cx}\sqrt{1 + cx}(d + ex^2)^{3/2}} + \frac{2bc(3c^2d + 2e)(1 - c^2x^2)}{15d^2(c^2d + e)^2\sqrt{-1 + cx}\sqrt{1 + cx}\sqrt{d + ex^2}} + \frac{x(a + b\operatorname{arccosh}(cx))}{5d(d + ex^2)^{5/2}} + \frac{4x(a + b\operatorname{arccosh}(cx))}{15d^2(d + ex^2)^{3/2}} + \frac{8x(a + b\operatorname{arccosh}(cx))}{15d^3\sqrt{d + ex^2}} - \frac{8b\sqrt{-1 + c^2x^2}\operatorname{arctanh}\left(\frac{\sqrt{e}\sqrt{-1 + c^2x^2}}{c\sqrt{d + ex^2}}\right)}{15d^3\sqrt{e}\sqrt{-1 + cx}\sqrt{1 + cx}}$$

output

```
1/5*x*(a+b*arccosh(c*x))/d/(e*x^2+d)^(5/2)+4/15*x*(a+b*arccosh(c*x))/d^2/(
e*x^2+d)^(3/2)+1/15*b*c*(-c^2*x^2+1)/d/(c^2*d+e)/(e*x^2+d)^(3/2)/(c*x-1)^(
1/2)/(c*x+1)^(1/2)-8/15*b*arctanh(e^(1/2)*(c^2*x^2-1)^(1/2)/c/(e*x^2+d)^(1
/2))*(c^2*x^2-1)^(1/2)/d^3/e^(1/2)/(c*x-1)^(1/2)/(c*x+1)^(1/2)+8/15*x*(a+b
*arccosh(c*x))/d^3/(e*x^2+d)^(1/2)+2/15*b*c*(3*c^2*d+2*e)*(-c^2*x^2+1)/d^2
/(c^2*d+e)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)/(e*x^2+d)^(1/2)
```

### 3.518.2 Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 3.00 (sec) , antiderivative size = 685, normalized size of antiderivative = 2.41

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^{7/2}} dx = \frac{ax(15d^2 + 20dex^2 + 8e^2x^4)}{d^3} - \frac{bc\sqrt{-1+cx}\sqrt{1+cx}(d+ex^2)(e(5d+4ex^2)+c^2d(7d+6ex^2))}{d^2(c^2d+e)^2} + \frac{bx(15d^2+20dex^2+8e^2x^4)}{d^3}$$

input `Integrate[(a + b*ArcCosh[c*x])/(d + e*x^2)^(7/2), x]`

output

```
((a*x*(15*d^2 + 20*d*e*x^2 + 8*e^2*x^4))/d^3 - (b*c*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(d + e*x^2)*(e*(5*d + 4*e*x^2) + c^2*d*(7*d + 6*e*x^2)))/(d^2*(c^2*d + e)^2) + (b*x*(15*d^2 + 20*d*e*x^2 + 8*e^2*x^4)*ArcCosh[c*x])/d^3 + (16*b*(-1 + c*x)^(3/2)*Sqrt[((c*Sqrt[d] - I*Sqrt[e])*(1 + c*x))/((c*Sqrt[d] + I*Sqrt[e])*(-1 + c*x))]*(d + e*x^2)^2*((c*((-I)*c*Sqrt[d] + Sqrt[e])*(I*Sqrt[d] + Sqrt[e]*x)*Sqrt[(1 + (I*c*Sqrt[d])/Sqrt[e] - c*x + (I*Sqrt[e]*x)/Sqrt[d])/(1 - c*x)]*EllipticF[ArcSin[Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(2 - 2*c*x))]], ((4*I)*c*Sqrt[d]*Sqrt[e])/(c*Sqrt[d] + I*Sqrt[e])^2))/(-1 + c*x) + c*Sqrt[d]*(-(c*Sqrt[d]) + I*Sqrt[e])*Sqrt[((c^2*d + e)*(d + e*x^2))/(d*e*(-1 + c*x)^2)]*Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(1 - c*x))]*EllipticPi[(2*c*Sqrt[d])/((c*Sqrt[d] + I*Sqrt[e])^2)], ArcSin[Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(2 - 2*c*x))]], ((4*I)*c*Sqrt[d]*Sqrt[e])/(c*Sqrt[d] + I*Sqrt[e])^2))/((c*d^3*(c^2*d + e)*Sqrt[1 + c*x]*Sqrt[-((-1 + (I*Sqrt[e]*x)/Sqrt[d] + c*((I*Sqrt[d])/Sqrt[e] + x))/(1 - c*x))]))/(15*(d + e*x^2)^(5/2))
```

### 3.518.3 Rubi [A] (verified)

Time = 1.16 (sec) , antiderivative size = 270, normalized size of antiderivative = 0.95, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$ , Rules used = {6323, 27, 2038, 7266, 1193, 27, 87, 66, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.518.  $\int \frac{a+b\operatorname{arccosh}(cx)}{(d+ex^2)^{7/2}} dx$

$$\begin{aligned}
& \int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{7/2}} dx \\
& \quad \downarrow \text{6323} \\
& -bc \int \frac{x(8e^2x^4 + 20dex^2 + 15d^2)}{15d^3\sqrt{cx-1}\sqrt{cx+1}(ex^2+d)^{5/2}} dx + \frac{8x(a + \operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}} + \frac{4x(a + \operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}} \\
& \quad \downarrow \text{27} \\
& -\frac{bc \int \frac{x(8e^2x^4+20dex^2+15d^2)}{\sqrt{cx-1}\sqrt{cx+1}(ex^2+d)^{5/2}} dx}{15d^3} + \frac{8x(a + \operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}} + \frac{4x(a + \operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}} \\
& \quad \downarrow \text{2038} \\
& -\frac{bc\sqrt{c^2x^2-1} \int \frac{x(8e^2x^4+20dex^2+15d^2)}{\sqrt{c^2x^2-1}(ex^2+d)^{5/2}} dx}{15d^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{8x(a + \operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}} + \frac{4x(a + \operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}} \\
& \quad \downarrow \text{7266} \\
& -\frac{bc\sqrt{c^2x^2-1} \int \frac{8e^2x^4+20dex^2+15d^2}{\sqrt{c^2x^2-1}(ex^2+d)^{5/2}} dx^2}{30d^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{8x(a + \operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}} + \frac{4x(a + \operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \\
& \quad \frac{x(a + \operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}} \\
& \quad \downarrow \text{1193} \\
& -\frac{bc\sqrt{c^2x^2-1} \left( \frac{2 \int \frac{3(4e(dc^2+e)x^2+d(7dc^2+6e))}{\sqrt{c^2x^2-1}(ex^2+d)^{3/2}} dx^2}{3(c^2d+e)} + \frac{2d^2\sqrt{c^2x^2-1}}{(c^2d+e)(d+ex^2)^{3/2}} \right)}{30d^3\sqrt{cx-1}\sqrt{cx+1}} + \frac{8x(a + \operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}} + \\
& \quad \frac{4x(a + \operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \frac{x(a + \operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}} \\
& \quad \downarrow \text{27}
\end{aligned}$$

---

3.518.  $\int \frac{a+\operatorname{barccosh}(cx)}{(d+ex^2)^{7/2}} dx$

$$\begin{aligned}
& \frac{bc\sqrt{c^2x^2-1} \left( \frac{2 \int \frac{4e(dc^2+e)x^2+d(7dc^2+6e)}{\sqrt{c^2x^2-1}(ex^2+d)^{3/2}} dx^2}{c^2d+e} + \frac{2d^2\sqrt{c^2x^2-1}}{(c^2d+e)(d+ex^2)^{3/2}} \right)}{\frac{30d^3\sqrt{cx-1}\sqrt{cx+1}}{4x(a+\operatorname{barccosh}(cx))} + \frac{x(a+\operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \frac{x(a+\operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}}} + \frac{8x(a+\operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}}} \\
& \quad \downarrow 87 \\
& \frac{bc\sqrt{c^2x^2-1} \left( \frac{2 \left( \frac{4(c^2d+e) \int \frac{1}{\sqrt{c^2x^2-1}\sqrt{ex^2+d}} dx^2 + \frac{2d\sqrt{c^2x^2-1}(3c^2d+2e)}{(c^2d+e)\sqrt{d+ex^2}} \right)}{c^2d+e} + \frac{2d^2\sqrt{c^2x^2-1}}{(c^2d+e)(d+ex^2)^{3/2}} \right)}{\frac{30d^3\sqrt{cx-1}\sqrt{cx+1}}{4x(a+\operatorname{barccosh}(cx))} + \frac{x(a+\operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \frac{x(a+\operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}}} + \frac{8x(a+\operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}}} \\
& \quad \downarrow 66 \\
& \frac{bc\sqrt{c^2x^2-1} \left( \frac{2 \left( \frac{8(c^2d+e) \int \frac{1}{c^2-ex^4} d \frac{\sqrt{c^2x^2-1}}{\sqrt{ex^2+d}} + \frac{2d\sqrt{c^2x^2-1}(3c^2d+2e)}{(c^2d+e)\sqrt{d+ex^2}} \right)}{c^2d+e} + \frac{2d^2\sqrt{c^2x^2-1}}{(c^2d+e)(d+ex^2)^{3/2}} \right)}{\frac{30d^3\sqrt{cx-1}\sqrt{cx+1}}{4x(a+\operatorname{barccosh}(cx))} + \frac{x(a+\operatorname{barccosh}(cx))}{15d^2(d+ex^2)^{3/2}} + \frac{x(a+\operatorname{barccosh}(cx))}{5d(d+ex^2)^{5/2}}} + \frac{8x(a+\operatorname{barccosh}(cx))}{15d^3\sqrt{d+ex^2}}} \\
& \quad \downarrow 221 \\
& \frac{bc\sqrt{c^2x^2-1} \left( \frac{2 \left( \frac{8(c^2d+e) \operatorname{arctanh} \left( \frac{\sqrt{e}\sqrt{c^2x^2-1}}{c\sqrt{d+ex^2}} \right) + \frac{2d\sqrt{c^2x^2-1}(3c^2d+2e)}{(c^2d+e)\sqrt{d+ex^2}} \right)}{c\sqrt{e}} + \frac{2d^2\sqrt{c^2x^2-1}}{(c^2d+e)(d+ex^2)^{3/2}} \right)}{30d^3\sqrt{cx-1}\sqrt{cx+1}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])/(d + e*x^2)^(7/2),x]`

```
output (x*(a + b*ArcCosh[c*x]))/(5*d*(d + e*x^2)^(5/2)) + (4*x*(a + b*ArcCosh[c*x
]))/(15*d^2*(d + e*x^2)^(3/2)) + (8*x*(a + b*ArcCosh[c*x]))/(15*d^3*Sqrt[d
+ e*x^2]) - (b*c*Sqrt[-1 + c^2*x^2]*((2*d^2*Sqrt[-1 + c^2*x^2])/((c^2*d +
e)*(d + e*x^2)^(3/2)) + (2*((2*d*(3*c^2*d + 2*e)*Sqrt[-1 + c^2*x^2])/((c^
2*d + e)*Sqrt[d + e*x^2]) + (8*(c^2*d + e)*ArcTanh[(Sqrt[e]*Sqrt[-1 + c^2*
x^2])/(c*Sqrt[d + e*x^2])])/(c*Sqrt[e])))/(c^2*d + e)))/(30*d^3*Sqrt[-1 +
c*x]*Sqrt[1 + c*x])
```

### 3.518.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(G_x_) /; FreeQ[b, x]]
```

```
rule 66 Int[1/(Sqrt[(a_) + (b_)*(x_)]*Sqrt[(c_) + (d_)*(x_)]), x_Symbol] := Simp[
2 Subst[Int[1/(b - d*x^2), x], x, Sqrt[a + b*x]/Sqrt[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && !GtQ[c - a*(d/b), 0]
```

```
rule 87 Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p
_), x_] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p
+ 1)*(c*f - d*e))), x] - Simp[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p
+ 1)))/(f*(p + 1)*(c*f - d*e)) Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x]
/; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || Intege
rQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n])))
```

```
rule 221 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

```
rule 1193 Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_)
+ (c_)*(x_)^2)^(p_), x_Symbol] := With[{Qx = PolynomialQuotient[(a + b*x
+ c*x^2)^p, d + e*x, x], R = PolynomialRemainder[(a + b*x + c*x^2)^p, d +
e*x, x]}, Simp[R*(d + e*x)^(m + 1)*((f + g*x)^(n + 1)/((m + 1)*(e*f - d*g))
), x] + Simp[1/((m + 1)*(e*f - d*g)) Int[(d + e*x)^(m + 1)*(f + g*x)^n*Ex
pandToSum[(m + 1)*(e*f - d*g)*Qx - g*R*(m + n + 2), x], x], x]] /; FreeQ[{a
, b, c, d, e, f, g, n}, x] && IGtQ[p, 0] && ILtQ[2*m, -2] && !IntegerQ[n]
&& !(EqQ[m, -2] && EqQ[p, 1] && EqQ[2*c*d - b*e, 0])
```

rule 2038 `Int[(u_)*((c_) + (d_)*(x_)^(n_))^(q_)*((a1_) + (b1_)*(x_)^(non2_))^(p_)*((a2_) + (b2_)*(x_)^(non2_))^(p_), x_Symbol] := Simp[(a1 + b1*x^(n/2))^FracPart[p]*((a2 + b2*x^(n/2))^FracPart[p]/(a1*a2 + b1*b2*x^n)^FracPart[p]) Int[u*(a1*a2 + b1*b2*x^n)^p*(c + d*x^n)^q, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, n, p, q}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && !(EqQ[n, 2] && IGtQ[q, 0])`

rule 6323 `Int[((a_) + ArcCosh[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_)^2)^(p_), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d + e, 0] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])`

rule 7266 `Int[(u_)*(x_)^(m_), x_Symbol] := Simp[1/(m + 1) Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]`

### 3.518.4 Maple [F]

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(ex^2 + d)^{7/2}} dx$$

input `int((a+b*arccosh(c*x))/(e*x^2+d)^(7/2),x)`

output `int((a+b*arccosh(c*x))/(e*x^2+d)^(7/2),x)`

### 3.518.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 674 vs.  $2(238) = 476$ .

Time = 0.39 (sec) , antiderivative size = 1360, normalized size of antiderivative = 4.79

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^{7/2}} dx = \text{Too large to display}$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(7/2),x, algorithm="fracas")`

output

```
[1/15*(2*(b*c^4*d^5 + 2*b*c^2*d^4*e + (b*c^4*d^2*e^3 + 2*b*c^2*d*e^4 + b*e^5)*x^6 + b*d^3*e^2 + 3*(b*c^4*d^3*e^2 + 2*b*c^2*d^2*e^3 + b*d*e^4)*x^4 + 3*(b*c^4*d^4*e + 2*b*c^2*d^3*e^2 + b*d^2*e^3)*x^2)*sqrt(e)*log(8*c^4*e^2*x^4 + c^4*d^2 - 6*c^2*d*e + 8*(c^4*d*e - c^2*e^2)*x^2 - 4*(2*c^3*e*x^2 + c^3*d - c*e)*sqrt(c^2*x^2 - 1)*sqrt(e*x^2 + d)*sqrt(e) + e^2) + (8*(b*c^4*d^2*e^3 + 2*b*c^2*d*e^4 + b*e^5)*x^5 + 20*(b*c^4*d^3*e^2 + 2*b*c^2*d^2*e^3 + b*d*e^4)*x^3 + 15*(b*c^4*d^4*e + 2*b*c^2*d^3*e^2 + b*d^2*e^3)*x)*sqrt(e*x^2 + d)*log(c*x + sqrt(c^2*x^2 - 1)) + (8*(a*c^4*d^2*e^3 + 2*a*c^2*d*e^4 + a*e^5)*x^5 + 20*(a*c^4*d^3*e^2 + 2*a*c^2*d^2*e^3 + a*d*e^4)*x^3 + 15*(a*c^4*d^4*e + 2*a*c^2*d^3*e^2 + a*d^2*e^3)*x - (7*b*c^3*d^4*e + 5*b*c*d^3*e^2 + 2*(3*b*c^3*d^2*e^3 + 2*b*c*d*e^4)*x^4 + (13*b*c^3*d^3*e^2 + 9*b*c*d^2*e^3)*x^2)*sqrt(c^2*x^2 - 1)*sqrt(e*x^2 + d))/(c^4*d^8*e + 2*c^2*d^7*e^2 + d^6*e^3 + (c^4*d^5*e^4 + 2*c^2*d^4*e^5 + d^3*e^6)*x^6 + 3*(c^4*d^6*e^3 + 2*c^2*d^5*e^4 + d^4*e^5)*x^4 + 3*(c^4*d^7*e^2 + 2*c^2*d^6*e^3 + d^5*e^4)*x^2), 1/15*(4*(b*c^4*d^5 + 2*b*c^2*d^4*e + (b*c^4*d^2*e^3 + 2*b*c^2*d*e^4 + b*e^5)*x^6 + b*d^3*e^2 + 3*(b*c^4*d^3*e^2 + 2*b*c^2*d^2*e^3 + b*d*e^4)*x^4 + 3*(b*c^4*d^4*e + 2*b*c^2*d^3*e^2 + b*d^2*e^3)*x^2)*sqrt(-e)*arctan(1/2*(2*c^2*e*x^2 + c^2*d - e)*sqrt(c^2*x^2 - 1)*sqrt(e*x^2 + d)*sqrt(-e)/(c^3*e^2*x^4 - c*d*e + (c^3*d*e - c*e^2)*x^2)) + (8*(b*c^4*d^2*e^3 + 2*b*c^2*d*e^4 + b*e^5)*x^5 + 20*(b*c^4*d^3*e^2 + 2*b*c^2*d^2*e^3 + b*d*e^4)*x^3 + ...
```

### 3.518.6 Sympy [F(-1)]

Timed out.

$$\int \frac{a + b \operatorname{arccosh}(cx)}{(d + ex^2)^{7/2}} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))/(e*x**2+d)**(7/2),x)`

output `Timed out`

**3.518.7 Maxima [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{7/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^{7/2}} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(7/2),x, algorithm="maxima")`

output `1/15*a*(8*x/(sqrt(e*x^2 + d)*d^3) + 4*x/((e*x^2 + d)^(3/2)*d^2) + 3*x/((e*x^2 + d)^(5/2)*d)) + b*integrate(log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1))/(e*x^2 + d)^(7/2), x)`

**3.518.8 Giac [F]**

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{7/2}} dx = \int \frac{b \operatorname{arcosh}(cx) + a}{(ex^2 + d)^{7/2}} dx$$

input `integrate((a+b*arccosh(c*x))/(e*x^2+d)^(7/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)/(e*x^2 + d)^(7/2), x)`

**3.518.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{a + \operatorname{barccosh}(cx)}{(d + ex^2)^{7/2}} dx = \int \frac{a + b \operatorname{acosh}(cx)}{(ex^2 + d)^{7/2}} dx$$

input `int((a + b*acosh(c*x))/(d + e*x^2)^(7/2),x)`

output `int((a + b*acosh(c*x))/(d + e*x^2)^(7/2), x)`



### 3.519 $\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

3.519.1 Optimal result . . . . .	3832
3.519.2 Mathematica [A] (verified) . . . . .	3833
3.519.3 Rubi [A] (verified) . . . . .	3834
3.519.4 Maple [F] . . . . .	3838
3.519.5 Fricas [F] . . . . .	3839
3.519.6 Sympy [F(-1)] . . . . .	3839
3.519.7 Maxima [F] . . . . .	3839
3.519.8 Giac [F] . . . . .	3840
3.519.9 Mupad [F(-1)] . . . . .	3840

#### 3.519.1 Optimal result

Integrand size = 23, antiderivative size = 558

$$\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{be \left( 3c^2 de(7+m)^2 (12+7m+m^2) + 3c^4 d^2 (35+12m+m^2)^2 + e^2 (360+342m+119m^2+18m^3+m^4) \right)}{c^5 f^2 (3+m)^2 (5+m)^2 (7+m)^2 \sqrt{-1+cx} \sqrt{1+cx}}$$

$$+ \frac{be^2 (3c^2 d(7+m)^2 + e(30+11m+m^2)) (fx)^{4+m} (1-c^2 x^2)}{c^3 f^4 (5+m)^2 (7+m)^2 \sqrt{-1+cx} \sqrt{1+cx}}$$

$$+ \frac{be^3 (fx)^{6+m} (1-c^2 x^2)}{cf^6 (7+m)^2 \sqrt{-1+cx} \sqrt{1+cx}} + \frac{d^3 (fx)^{1+m} (a + \operatorname{barccosh}(cx))}{f(1+m)}$$

$$+ \frac{3d^2 e (fx)^{3+m} (a + \operatorname{barccosh}(cx))}{f^3 (3+m)}$$

$$+ \frac{3de^2 (fx)^{5+m} (a + \operatorname{barccosh}(cx))}{f^5 (5+m)} + \frac{e^3 (fx)^{7+m} (a + \operatorname{barccosh}(cx))}{f^7 (7+m)}$$

$$+ \frac{b \left( \frac{c^6 d^3 (3+m)(5+m)(7+m)}{1+m} + \frac{e^{(2+m)} (3c^2 de(7+m)^2 (12+7m+m^2) + 3c^4 d^2 (35+12m+m^2)^2 + e^2 (360+342m+119m^2+18m^3+m^4))}{(3+m)(5+m)(7+m)} \right)}{c^5 f^2 (2+m)(3+m)(5+m)(7+m) \sqrt{-1+cx} \sqrt{1+cx}}$$

output  $d^3(fx)^{(1+m)}(a+b\operatorname{arccosh}(cx))/f/(1+m)+3d^2e^*(fx)^{(3+m)}(a+b\operatorname{arccosh}(cx))/f^3/(3+m)+3d^2e^{2*(fx)^{(5+m)}(a+b\operatorname{arccosh}(cx))/f^5/(5+m)+e^{3*(fx)^{(7+m)}(a+b\operatorname{arccosh}(cx))/f^7/(7+m)+b^*e^*(3*c^2*d^*e^*(7+m)^2*(m^2+7*m+12)+3*c^4*d^2*(m^2+12*m+35)^2+e^{2*(m^4+18*m^3+119*m^2+342*m+360)}*(fx)^{(2+m)}(-c^2*x^2+1)/c^5/f^2/(3+m)^2/(5+m)^2/(7+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)+b^*e^{2*(3*c^2*d^*(7+m)^2+e^{2*(m^2+11*m+30)}*(fx)^{(4+m)}(-c^2*x^2+1)/c^3/f^4/(5+m)^2/(7+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)+b^*e^{3*(fx)^{(6+m)}(-c^2*x^2+1)/c/f^6/(7+m)^2/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)-b^*(c^6*d^3*(3+m)*(5+m)*(7+m)/(1+m)+e^{(2+m)}(3*c^2*d^*e^*(7+m)^2*(m^2+7*m+12)+3*c^4*d^2*(m^2+12*m+35)^2+e^{2*(m^4+18*m^3+119*m^2+342*m+360)})/(m^3+15*m^2+71*m+105)}*(fx)^{(2+m)}\operatorname{hypergeom}([1/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^{(1/2)}/c^5/f^2/(2+m)/(3+m)/(5+m)/(7+m)/(c*x-1)^{(1/2)}/(c*x+1)^{(1/2)}$

### 3.519.2 Mathematica [A] (verified)

Time = 0.87 (sec) , antiderivative size = 397, normalized size of antiderivative = 0.71

$$\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$$

$$= x(fx)^m \left( \frac{d^3(a + \operatorname{barccosh}(cx))}{1 + m} + \frac{3d^2ex^2(a + \operatorname{barccosh}(cx))}{3 + m} + \frac{3de^2x^4(a + \operatorname{barccosh}(cx))}{5 + m} + \frac{e^3x^6(a + \operatorname{barccosh}(cx))}{7 + m} - \frac{bce^3x^7\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, 4 + \frac{m}{2}, 5 + \frac{m}{2}, c^2x^2\right)}{(7 + m)(8 + m)\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{bcd^3x\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{(2 + 3m + m^2)\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{3bcd^2ex^3\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4+m}{2}, \frac{6+m}{2}, c^2x^2\right)}{(12 + 7m + m^2)\sqrt{-1 + cx}\sqrt{1 + cx}} - \frac{3bcde^2x^5\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{6+m}{2}, \frac{8+m}{2}, c^2x^2\right)}{(5 + m)(6 + m)\sqrt{-1 + cx}\sqrt{1 + cx}} \right)$$

input `Integrate[(f*x)^m*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

```

output x*(f*x)^m*((d^3*(a + b*ArcCosh[c*x]))/(1 + m) + (3*d^2*e*x^2*(a + b*ArcCos
h[c*x]))/(3 + m) + (3*d*e^2*x^4*(a + b*ArcCosh[c*x]))/(5 + m) + (e^3*x^6*(
a + b*ArcCosh[c*x]))/(7 + m) - (b*c*e^3*x^7*sqrt[1 - c^2*x^2]*Hypergeometr
ic2F1[1/2, 4 + m/2, 5 + m/2, c^2*x^2])/((7 + m)*(8 + m)*sqrt[-1 + c*x]*Sqr
t[1 + c*x]) - (b*c*d^3*x*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/
2, (4 + m)/2, c^2*x^2])/((2 + 3*m + m^2)*sqrt[-1 + c*x]*sqrt[1 + c*x]) - (
3*b*c*d^2*e*x^3*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m
)/2, c^2*x^2])/((12 + 7*m + m^2)*sqrt[-1 + c*x]*sqrt[1 + c*x]) - (3*b*c*d*
e^2*x^5*sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (6 + m)/2, (8 + m)/2, c^2
*x^2])/((5 + m)*(6 + m)*sqrt[-1 + c*x]*sqrt[1 + c*x]))

```

### 3.519.3 Rubi [A] (verified)

Time = 2.47 (sec) , antiderivative size = 544, normalized size of antiderivative = 0.97, number of steps used = 8, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$ , Rules used = {6373, 27, 2113, 2340, 1590, 363, 279, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex^2)^3 (fx)^m (a + \operatorname{barccosh}(cx)) dx \\
 & \quad \downarrow \text{6373} \\
 & -bc \int \frac{(fx)^{m+1} \left( \frac{e^3 x^6}{m+7} + \frac{3de^2 x^4}{m+5} + \frac{3d^2 ex^2}{m+3} + \frac{d^3}{m+1} \right)}{f \sqrt{cx-1} \sqrt{cx+1}} dx + \frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} + \\
 & \frac{3d^2 e (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3 (m+3)} + \frac{3de^2 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5 (m+5)} + \frac{e^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7 (m+7)} \\
 & \quad \downarrow \text{27} \\
 & -bc \int \frac{(fx)^{m+1} \left( \frac{e^3 x^6}{m+7} + \frac{3de^2 x^4}{m+5} + \frac{3d^2 ex^2}{m+3} + \frac{d^3}{m+1} \right)}{\sqrt{cx-1} \sqrt{cx+1}} dx + \frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} + \\
 & \frac{3d^2 e (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3 (m+3)} + \frac{3de^2 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5 (m+5)} + \frac{e^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7 (m+7)} \\
 & \quad \downarrow \text{2113} \\
 & -\frac{bc \sqrt{c^2 x^2 - 1} \int \frac{(fx)^{m+1} \left( \frac{e^3 x^6}{m+7} + \frac{3de^2 x^4}{m+5} + \frac{3d^2 ex^2}{m+3} + \frac{d^3}{m+1} \right)}{\sqrt{c^2 x^2 - 1}} dx}{f \sqrt{cx-1} \sqrt{cx+1}} + \frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} + \\
 & \frac{3d^2 e (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3 (m+3)} + \frac{3de^2 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5 (m+5)} + \frac{e^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7 (m+7)}
 \end{aligned}$$

---

3.519.  $\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

↓ 2340

$$bc\sqrt{c^2x^2 - 1} \left( \int \frac{(fx)^{m+1} \left( \frac{e^2(3c^2d(m+7)^2 + e(m^2 + 11m + 30))x^4}{(m+5)(m+7)} + \frac{3c^2d^2e(m+7)x^2}{m+3} + \frac{c^2d^3(m+7)}{m+1} \right)}{\sqrt{c^2x^2 - 1} c^2(m+7)} dx + \frac{e^3\sqrt{c^2x^2 - 1}(fx)^{m+6}}{c^2f^5(m+7)^2} \right) +$$

$$\frac{d^3(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{3d^2e(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{3de^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))} + \frac{f^3(m+3)}{f^5(m+5)} + \frac{e^3(fx)^{m+7}(a + \operatorname{barccosh}(cx))}{f^7(m+7)}$$

↓ 1590

$$bc\sqrt{c^2x^2 - 1} \left( \int \frac{(fx)^{m+1} \left( \frac{d^3(m+5)(m+7)c^4}{m+1} + \frac{e(3d^2(m^2 + 12m + 35)^2c^4 + 3de(m+7)^2(m^2 + 7m + 12)c^2 + e^2(m^4 + 18m^3 + 119m^2 + 342m + 360))x^2}{(m+3)(m+5)(m+7)} \right)}{\sqrt{c^2x^2 - 1} c^2(m+5)} dx \right) +$$

$$\frac{d^3(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{3d^2e(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{3de^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))} + \frac{f^3(m+3)}{f^5(m+5)} + \frac{e^3(fx)^{m+7}(a + \operatorname{barccosh}(cx))}{f^7(m+7)}$$

↓ 363

$$bc\sqrt{c^2x^2 - 1} \left( \int \frac{\left( \frac{c^4d^3(m+5)(m+7)}{m+1} + \frac{e(m+2)(3c^4d^2(m^2 + 12m + 35)^2 + 3c^2de(m+7)^2(m^2 + 7m + 12) + e^2(m^4 + 18m^3 + 119m^2 + 342m + 360))}{c^2(m+3)^2(m+5)(m+7)} \right) \int \frac{(fx)^{m+1}}{\sqrt{c^2x^2 - 1}} dx}{c^2(m+5)} \right) +$$

$$\frac{d^3(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{3d^2e(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{3de^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))} + \frac{f^3(m+3)}{f^5(m+5)} + \frac{e^3(fx)^{m+7}(a + \operatorname{barccosh}(cx))}{f^7(m+7)}$$

↓ 279

3.519.  $\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

$$bc\sqrt{c^2x^2 - 1} \left( \frac{\sqrt{1-c^2x^2} \left( \frac{c^4 d^3 (m+5)(m+7)}{m+1} + \frac{e^{(m+2)} (3c^4 d^2 (m^2+12m+35)^2 + 3c^2 de(m+7)^2 (m^2+7m+12) + e^2 (m^4+18m^3+119m^2+342m+360))}{c^2 (m+3)^2 (m+5)(m+7)} \right)}{\sqrt{c^2x^2-1}} \right) \frac{1}{c^2(m+5)}$$

$$\frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{3d^2 e (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{3de^2 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5(m+5)} + \frac{e^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7(m+7)}$$

↓ 278

$$\frac{d^3 (fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{3d^2 e (fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{3de^2 (fx)^{m+5} (a + \operatorname{barccosh}(cx))}{f^5(m+5)} + \frac{e^3 (fx)^{m+7} (a + \operatorname{barccosh}(cx))}{f^7(m+7)}$$

$$bc\sqrt{c^2x^2 - 1} \left( \frac{e^3 \sqrt{c^2x^2-1} (fx)^{m+6}}{c^2 f^5 (m+7)^2} + \frac{e^2 \sqrt{c^2x^2-1} (fx)^{m+4} (3c^2 d(m+7)^2 + e(m^2+11m+30))}{c^2 f^3 (m+5)^2 (m+7)} + \frac{e^2 \sqrt{c^2x^2-1} (fx)^{m+2} (3c^4 d^2 (m^2+12m+35)^2 + 3c^2 de(m+7)^2 (m^2+7m+12) + e^2 (m^4+18m^3+119m^2+342m+360))}{c^2 f(m+3)^2 (m+5)(m+7)} \right)$$

input `Int[(f*x)^m*(d + e*x^2)^3*(a + b*ArcCosh[c*x]),x]`

```
output (d^3*(f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(f*(1 + m)) + (3*d^2*e*(f*x)^(3 +
m)*(a + b*ArcCosh[c*x]))/(f^3*(3 + m)) + (3*d*e^2*(f*x)^(5 + m)*(a + b*Ar
cCosh[c*x]))/(f^5*(5 + m)) + (e^3*(f*x)^(7 + m)*(a + b*ArcCosh[c*x]))/(f^7
*(7 + m)) - (b*c*Sqrt[-1 + c^2*x^2]*((e^3*(f*x)^(6 + m)*Sqrt[-1 + c^2*x^2]
)/(c^2*f^5*(7 + m)^2) + ((e^2*(3*c^2*d*(7 + m)^2 + e*(30 + 11*m + m^2))*(f
*x)^(4 + m)*Sqrt[-1 + c^2*x^2]))/(c^2*f^3*(5 + m)^2*(7 + m)) + ((e*(3*c^2*d
*e*(7 + m)^2*(12 + 7*m + m^2) + 3*c^4*d^2*(35 + 12*m + m^2)^2 + e^2*(360 +
342*m + 119*m^2 + 18*m^3 + m^4))*(f*x)^(2 + m)*Sqrt[-1 + c^2*x^2]))/(c^2*f
*(3 + m)^2*(5 + m)*(7 + m)) + (((c^4*d^3*(5 + m)*(7 + m))/(1 + m) + (e*(2
+ m)*(3*c^2*d*e*(7 + m)^2*(12 + 7*m + m^2) + 3*c^4*d^2*(35 + 12*m + m^2)^2
+ e^2*(360 + 342*m + 119*m^2 + 18*m^3 + m^4)))/(c^2*(3 + m)^2*(5 + m)*(7
+ m)))*(f*x)^(2 + m)*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (
4 + m)/2, c^2*x^2])/(f*(2 + m)*Sqrt[-1 + c^2*x^2]))/(c^2*(5 + m)))/(c^2*(7
+ m)))/(f*Sqrt[-1 + c*x]*Sqrt[1 + c*x])
```

### 3.519.3.1 Defintions of rubi rules used

```
rule 27 Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

```
rule 278 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((
c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (
-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0
] || GtQ[a, 0])
```

```
rule 279 Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntP
art[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(
1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] &&
!(ILtQ[p, 0] || GtQ[a, 0])
```

```
rule 363 Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x
_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))),
x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^
m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d
, 0] && NeQ[m + 2*p + 3, 0]
```

rule 1590 `Int[((f_.)*(x_.))^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]`

rule 2113 `Int[(Px_)*((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_Symbol] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[Px*(a*c + b*d*x^2)^m*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && PolyQ[Px, x] && EqQ[b*c + a*d, 0] && EqQ[m, n] && !IntegerQ[m]`

rule 2340 `Int[(Pq_)*((c_.)*(x_.))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(c*x)^(m + q - 1)*((a + b*x^2)^(p + 1)/(b*c^(q - 1)*(m + q + 2*p + 1))), x] + Simp[1/(b*(m + q + 2*p + 1)) Int[(c*x)^m*(a + b*x^2)^p*ExpandToSum[b*(m + q + 2*p + 1)*Pq - b*f*(m + q + 2*p + 1)*x^q - a*f*(m + q - 1)*x^(q - 2), x], x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, m, p}, x] && PolyQ[Pq, x] && (!IGtQ[m, 0] || IGtQ[p + 1/2, -1])`

rule 6373 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))*((f_.)*(x_.))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

### 3.519.4 Maple [F]

$$\int (fx)^m (ex^2 + d)^3 (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(e*x^2+d)^3*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(e*x^2+d)^3*(a+b*arccosh(c*x)),x)`

---

3.519.  $\int (fx)^m (d + ex^2)^3 (a + b \operatorname{arccosh}(cx)) dx$

**3.519.5 Fracas [F]**

$$\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (ex^2 + d)^3 (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((a*e^3*x^6 + 3*a*d*e^2*x^4 + 3*a*d^2*e*x^2 + a*d^3 + (b*e^3*x^6 + 3*b*d*e^2*x^4 + 3*b*d^2*e*x^2 + b*d^3)*arccosh(c*x))*(f*x)^m, x)`

**3.519.6 Sympy [F(-1)]**

Timed out.

$$\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \text{Timed out}$$

input `integrate((f*x)**m*(e*x**2+d)**3*(a+b*acosh(c*x)),x)`

output `Timed out`

**3.519.7 Maxima [F]**

$$\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (ex^2 + d)^3 (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="maxima")`



output `a*e^3*f^m*x^7*x^m/(m + 7) + 3*a*d*e^2*f^m*x^5*x^m/(m + 5) + 3*a*d^2*e*f^m*x^3*x^m/(m + 3) + (f*x)^(m + 1)*a*d^3/(f*(m + 1)) + ((m^3 + 9*m^2 + 23*m + 15)*b*e^3*f^m*x^7 + 3*(m^3 + 11*m^2 + 31*m + 21)*b*d*e^2*f^m*x^5 + 3*(m^3 + 13*m^2 + 47*m + 35)*b*d^2*e*f^m*x^3 + (m^3 + 15*m^2 + 71*m + 105)*b*d^3*f^m*x)*x^m*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(m^4 + 16*m^3 + 86*m^2 + 176*m + 105) + integrate(((m^3 + 9*m^2 + 23*m + 15)*b*c*e^3*f^m*x^7 + 3*(m^3 + 11*m^2 + 31*m + 21)*b*c*d*e^2*f^m*x^5 + 3*(m^3 + 13*m^2 + 47*m + 35)*b*c*d^2*e*f^m*x^3 + (m^3 + 15*m^2 + 71*m + 105)*b*c*d^3*f^m*x)*x^m/((m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c^3*x^3 - (m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c*x + ((m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c^2*x^2 - m^4 - 16*m^3 - 86*m^2 - 176*m - 105)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) - integrate(((m^3 + 9*m^2 + 23*m + 15)*b*c^2*e^3*f^m*x^8 + 3*(m^3 + 11*m^2 + 31*m + 21)*b*c^2*d*e^2*f^m*x^6 + 3*(m^3 + 13*m^2 + 47*m + 35)*b*c^2*d^2*e*f^m*x^4 + (m^3 + 15*m^2 + 71*m + 105)*b*c^2*d^3*f^m*x^2)*x^m/((m^4 + 16*m^3 + 86*m^2 + 176*m + 105)*c^2*x^2 - m^4 - 16*m^3 - 86*m^2 - 176*m - 105), x)`

### 3.519.8 Giac [F]

$$\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (ex^2 + d)^3 (b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^3*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^3*(b*arccosh(c*x) + a)*(f*x)^m, x)`

### 3.519.9 Mupad [F(-1)]

Timed out.

$$\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (fx)^m (ex^2 + d)^3 dx$$

input `int((a + b*acosh(c*x))*(f*x)^m*(d + e*x^2)^3,x)`

output `int((a + b*acosh(c*x))*(f*x)^m*(d + e*x^2)^3, x)`

---

3.519.  $\int (fx)^m (d + ex^2)^3 (a + \operatorname{barccosh}(cx)) dx$

### 3.520 $\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

3.520.1 Optimal result . . . . .	3841
3.520.2 Mathematica [A] (verified) . . . . .	3842
3.520.3 Rubi [A] (verified) . . . . .	3842
3.520.4 Maple [F] . . . . .	3846
3.520.5 Fricas [F] . . . . .	3846
3.520.6 Sympy [F] . . . . .	3846
3.520.7 Maxima [F] . . . . .	3847
3.520.8 Giac [F] . . . . .	3847
3.520.9 Mupad [F(-1)] . . . . .	3847

#### 3.520.1 Optimal result

Integrand size = 23, antiderivative size = 353

$$\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= \frac{be(2c^2d(5+m)^2 + e(12+7m+m^2))(fx)^{2+m}(1-c^2x^2)}{c^3f^2(3+m)^2(5+m)^2\sqrt{-1+cx}\sqrt{1+cx}}$$

$$+ \frac{be^2(fx)^{4+m}(1-c^2x^2)}{cf^4(5+m)^2\sqrt{-1+cx}\sqrt{1+cx}} + \frac{d^2(fx)^{1+m}(a + \operatorname{barccosh}(cx))}{f(1+m)}$$

$$+ \frac{2de(fx)^{3+m}(a + \operatorname{barccosh}(cx))}{f^3(3+m)} + \frac{e^2(fx)^{5+m}(a + \operatorname{barccosh}(cx))}{f^5(5+m)}$$

$$- \frac{b\left(\frac{c^4d^2(3+m)(5+m)}{1+m} + \frac{e(2+m)(2c^2d(5+m)^2+e(12+7m+m^2))}{(3+m)(5+m)}\right)(fx)^{2+m}\sqrt{1-c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}\right)}{c^3f^2(2+m)(3+m)(5+m)\sqrt{-1+cx}\sqrt{1+cx}}$$

```
output d^2*(f*x)^(1+m)*(a+b*arccosh(c*x))/f/(1+m)+2*d*e*(f*x)^(3+m)*(a+b*arccosh(c*x))/f^3/(3+m)+e^2*(f*x)^(5+m)*(a+b*arccosh(c*x))/f^5/(5+m)+b*e*(2*c^2*d*(5+m)^2+e*(m^2+7*m+12))*(f*x)^(2+m)*(-c^2*x^2+1)/c^3/f^2/(3+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)+b*e^2*(f*x)^(4+m)*(-c^2*x^2+1)/c/f^4/(5+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)-b*(c^4*d^2*(3+m)*(5+m)/(1+m)+e*(2+m)*(2*c^2*d*(5+m)^2+e*(m^2+7*m+12))/(3+m)/(5+m))*(f*x)^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/c^3/f^2/(2+m)/(3+m)/(5+m)/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.520.2 Mathematica [A] (verified)**

Time = 0.32 (sec) , antiderivative size = 293, normalized size of antiderivative = 0.83

$$\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$$

$$= x(fx)^m \left( \frac{d^2(a + \operatorname{barccosh}(cx))}{1 + m} + \frac{2dex^2(a + \operatorname{barccosh}(cx))}{3 + m} + \frac{e^2x^4(a + \operatorname{barccosh}(cx))}{5 + m} \right. \\ \left. - \frac{bcd^2x\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{(2 + 3m + m^2)\sqrt{-1 + cx}\sqrt{1 + cx}} \right. \\ \left. - \frac{2bcdex^3\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4+m}{2}, \frac{6+m}{2}, c^2x^2\right)}{(12 + 7m + m^2)\sqrt{-1 + cx}\sqrt{1 + cx}} \right. \\ \left. - \frac{bce^2x^5\sqrt{1 - c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{6+m}{2}, \frac{8+m}{2}, c^2x^2\right)}{(5 + m)(6 + m)\sqrt{-1 + cx}\sqrt{1 + cx}} \right)$$

input `Integrate[(f*x)^m*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`output `x*(f*x)^m*((d^2*(a + b*ArcCosh[c*x]))/(1 + m) + (2*d*e*x^2*(a + b*ArcCosh[c*x]))/(3 + m) + (e^2*x^4*(a + b*ArcCosh[c*x]))/(5 + m) - (b*c*d^2*x*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, c^2*x^2])/((2 + 3*m + m^2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (2*b*c*d*e*x^3*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m)/2, c^2*x^2])/((12 + 7*m + m^2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]) - (b*c*e^2*x^5*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (6 + m)/2, (8 + m)/2, c^2*x^2])/((5 + m)*(6 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))`**3.520.3 Rubi [A] (verified)**Time = 0.74 (sec) , antiderivative size = 350, normalized size of antiderivative = 0.99, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.304$ , Rules used = {6373, 27, 1905, 1590, 363, 279, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^2 (fx)^m (a + \operatorname{barccosh}(cx)) dx$$

↓ 6373

---

3.520.  $\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

$$\begin{aligned}
 & -bc \int \frac{(fx)^{m+1} \left( \frac{e^2 x^4}{m+5} + \frac{2dex^2}{m+3} + \frac{d^2}{m+1} \right)}{f\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \\
 & \quad \frac{2de(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} \\
 & \quad \downarrow 27 \\
 & -bc \int \frac{(fx)^{m+1} \left( \frac{e^2 x^4}{m+5} + \frac{2dex^2}{m+3} + \frac{d^2}{m+1} \right)}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \\
 & \quad \frac{2de(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} \\
 & \quad \downarrow 1905 \\
 & -bc\sqrt{c^2x^2-1} \int \frac{(fx)^{m+1} \left( \frac{e^2 x^4}{m+5} + \frac{2dex^2}{m+3} + \frac{d^2}{m+1} \right)}{\sqrt{c^2x^2-1}} dx + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \\
 & \quad \frac{2de(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} \\
 & \quad \downarrow 1590 \\
 & -bc\sqrt{c^2x^2-1} \left( \int \frac{(fx)^{m+1} \left( \frac{c^2(m+5)d^2}{m+1} + \frac{e(2c^2d(m+5)^2 + e(m^2+7m+12))x^2}{(m+3)(m+5)} \right)}{\sqrt{c^2x^2-1}} dx + \frac{e^2\sqrt{c^2x^2-1}(fx)^{m+4}}{c^2f^3(m+5)^2} \right) \\
 & \quad + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{2de(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} \\
 & \quad \downarrow 363 \\
 & bc\sqrt{c^2x^2-1} \left( \int \frac{\left( \frac{c^2d^2(m+5)}{m+1} + \frac{e(m+2)(2c^2d(m+5)^2 + e(m^2+7m+12))}{c^2(m+3)^2(m+5)} \right) (fx)^{m+1}}{\sqrt{c^2x^2-1}} dx + \frac{e\sqrt{c^2x^2-1}(fx)^{m+2}(2c^2d(m+5)^2 + e(m^2+7m+12))}{c^2f(m+3)^2(m+5)} \right) + e^2 \\
 & \quad + \frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{2de(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} \\
 & \quad \downarrow 279
 \end{aligned}$$

---

3.520.  $\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

$$bc\sqrt{c^2x^2 - 1} \left( \frac{\sqrt{1-c^2x^2} \left( \frac{c^2d^2(m+5)}{m+1} + \frac{e(m+2)(2c^2d(m+5)^2 + e(m^2+7m+12))}{c^2(m+3)^2(m+5)} \right)}{\sqrt{c^2x^2-1}} \int \frac{(fx)^{m+1}}{\sqrt{1-c^2x^2}} dx + \frac{e\sqrt{c^2x^2-1}(fx)^{m+2}(2c^2d(m+5)^2 + e(m^2+7m+12))}{c^2f(m+3)^2(m+5)} \right)$$


---


$$\frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{2de(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{f\sqrt{cx-1}\sqrt{cx+1}}{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))} + \frac{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)}$$

↓ 278

$$\frac{d^2(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{2de(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} + \frac{e^2(fx)^{m+5}(a + \operatorname{barccosh}(cx))}{f^5(m+5)} -$$

$$bc\sqrt{c^2x^2 - 1} \left( \frac{\sqrt{1-c^2x^2}(fx)^{m+2} \left( \frac{c^2d^2(m+5)}{m+1} + \frac{e(m+2)(2c^2d(m+5)^2 + e(m^2+7m+12))}{c^2(m+3)^2(m+5)} \right)}{f(m+2)\sqrt{c^2x^2-1}} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right) + \frac{e\sqrt{c^2x^2-1}(fx)^m}{c^2(m+5)} \right)$$


---


$$f\sqrt{cx-1}\sqrt{cx+1}$$

input `Int[(f*x)^m*(d + e*x^2)^2*(a + b*ArcCosh[c*x]),x]`

output `(d^2*(f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(f*(1 + m)) + (2*d*e*(f*x)^(3 + m)*(a + b*ArcCosh[c*x]))/(f^3*(3 + m)) + (e^2*(f*x)^(5 + m)*(a + b*ArcCosh[c*x]))/(f^5*(5 + m)) - (b*c*Sqrt[-1 + c^2*x^2]*((e^2*(f*x)^(4 + m)*Sqrt[-1 + c^2*x^2]))/(c^2*f^3*(5 + m)^2) + ((e*(2*c^2*d*(5 + m)^2 + e*(12 + 7*m + m^2))*(f*x)^(2 + m)*Sqrt[-1 + c^2*x^2]))/(c^2*f*(3 + m)^2*(5 + m)) + (((c^2*d^2*(5 + m))/(1 + m) + (e*(2 + m)*(2*c^2*d*(5 + m)^2 + e*(12 + 7*m + m^2)))/(c^2*(3 + m)^2*(5 + m)))*(f*x)^(2 + m)*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, c^2*x^2])/(f*(2 + m)*Sqrt[-1 + c^2*x^2]))/(c^2*(5 + m)))/(f*Sqrt[-1 + c*x]*Sqrt[1 + c*x])`

**3.520.3.1 Defintions of rubi rules used**

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

3.520.  $\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx$

rule 279 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

rule 363 `Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`

rule 1590 `Int[((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Simp[c^p*(f*x)^(m + 4*p - 1)*((d + e*x^2)^(q + 1)/(e*f^(4*p - 1)*(m + 4*p + 2*q + 1))), x] + Simp[1/(e*(m + 4*p + 2*q + 1)) Int[(f*x)^m*(d + e*x^2)^q*ExpandToSum[e*(m + 4*p + 2*q + 1)*((a + b*x^2 + c*x^4)^p - c^p*x^(4*p)) - d*c^p*(m + 4*p - 1)*x^(4*p - 2), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p, 0] && !IntegerQ[q] && NeQ[m + 4*p + 2*q + 1, 0]`

rule 1905 `Int[((f_.)*(x_))^(m_.)*((d1_) + (e1_.)*(x_)^(non2_.))^q_.)*((d2_) + (e2_.)*(x_)^(non2_.))^q_.)*((a_.) + (b_.)*(x_)^(n_) + (c_.)*(x_)^(n2_))^(p_.), x_Symbol] := Simp[(d1 + e1*x^(n/2))^FracPart[q]*((d2 + e2*x^(n/2))^FracPart[q]/(d1*d2 + e1*e2*x^n)^FracPart[q]) Int[(f*x)^m*(d1*d2 + e1*e2*x^n)^q*(a + b*x^n + c*x^(2*n))^p, x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, f, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[non2, n/2] && EqQ[d2*e1 + d1*e2, 0]`

rule 6373 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^p, x]}, Simp[(a + b*ArcCosh[c*x]) u, x] - Simp[b*c Int[SimplifyIntegrand[u/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (GtQ[p, 0] || (IGtQ[(m - 1)/2, 0] && LeQ[m + p, 0]))`

**3.520.4 Maple [F]**

$$\int (fx)^m (ex^2 + d)^2 (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(e*x^2+d)^2*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(e*x^2+d)^2*(a+b*arccosh(c*x)),x)`

**3.520.5 Fracas [F]**

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{arccosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output `integral((a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arccosh(c*x))*(f*x)^m, x)`

**3.520.6 Sympy [F]**

$$\int (fx)^m (d + ex^2)^2 (a + b \operatorname{arccosh}(cx)) dx = \int (fx)^m (a + b \operatorname{acosh}(cx)) (d + ex^2)^2 dx$$

input `integrate((f*x)**m*(e*x**2+d)**2*(a+b*acosh(c*x)),x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))*(d + e*x**2)**2, x)`

**3.520.7 Maxima [F]**

$$\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `a*e^2*f^m*x^5*x^m/(m + 5) + 2*a*d*e*f^m*x^3*x^m/(m + 3) + (f*x)^(m + 1)*a*d^2/(f*(m + 1)) + ((m^2 + 4*m + 3)*b*e^2*f^m*x^5 + 2*(m^2 + 6*m + 5)*b*d*e*f^m*x^3 + (m^2 + 8*m + 15)*b*d^2*f^m*x)*x^m*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(m^3 + 9*m^2 + 23*m + 15) + integrate(((m^2 + 4*m + 3)*b*c*e^2*f^m*x^5 + 2*(m^2 + 6*m + 5)*b*c*d*e*f^m*x^3 + (m^2 + 8*m + 15)*b*c*d^2*f^m*x)*x^m/((m^3 + 9*m^2 + 23*m + 15)*c^3*x^3 - (m^3 + 9*m^2 + 23*m + 15)*c*x + ((m^3 + 9*m^2 + 23*m + 15)*c^2*x^2 - m^3 - 9*m^2 - 23*m - 15)*sqrt(c*x + 1))*sqrt(c*x - 1), x) - integrate(((m^2 + 4*m + 3)*b*c^2*e^2*f^m*x^6 + 2*(m^2 + 6*m + 5)*b*c^2*d*e*f^m*x^4 + (m^2 + 8*m + 15)*b*c^2*d^2*f^m*x^2)*x^m/((m^3 + 9*m^2 + 23*m + 15)*c^2*x^2 - m^3 - 9*m^2 - 23*m - 15), x)`

**3.520.8 Giac [F]**

$$\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int (ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)^2*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.520.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d + ex^2)^2 (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (fx)^m (ex^2 + d)^2 dx$$

input `int((a + b*acosh(c*x))*(f*x)^m*(d + e*x^2)^2,x)`

output `int((a + b*acosh(c*x))*(f*x)^m*(d + e*x^2)^2, x)`



### 3.521 $\int (fx)^m (d + ex^2) (a + \operatorname{arccosh}(cx)) dx$

3.521.1 Optimal result . . . . .	3848
3.521.2 Mathematica [A] (verified) . . . . .	3849
3.521.3 Rubi [A] (verified) . . . . .	3849
3.521.4 Maple [F] . . . . .	3852
3.521.5 Fricas [F] . . . . .	3852
3.521.6 Sympy [F] . . . . .	3852
3.521.7 Maxima [F] . . . . .	3853
3.521.8 Giac [F] . . . . .	3853
3.521.9 Mupad [F(-1)] . . . . .	3853

#### 3.521.1 Optimal result

Integrand size = 21, antiderivative size = 198

$$\int (fx)^m (d + ex^2) (a + \operatorname{arccosh}(cx)) dx = -\frac{be(fx)^{2+m}\sqrt{-1+cx}\sqrt{1+cx}}{cf^2(3+m)^2} + \frac{d(fx)^{1+m}(a + \operatorname{arccosh}(cx))}{f(1+m)} + \frac{e(fx)^{3+m}(a + \operatorname{arccosh}(cx))}{f^3(3+m)} - \frac{b(e(1+m)(2+m) + c^2d(3+m)^2)(fx)^{2+m}\sqrt{1-c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{cf^2(1+m)(2+m)(3+m)^2\sqrt{-1+cx}\sqrt{1+cx}}$$

output

```
d*(f*x)^(1+m)*(a+b*arccosh(c*x))/f/(1+m)+e*(f*x)^(3+m)*(a+b*arccosh(c*x))/f^3/(3+m)-b*e*(f*x)^(2+m)*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c/f^2/(3+m)^2-b*(e*(1+m)*(2+m)+c^2*d*(3+m)^2)*(f*x)^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], c^2*x^2)*(-c^2*x^2+1)^(1/2)/c/f^2/(1+m)/(2+m)/(3+m)^2/(c*x-1)^(1/2)/(c*x+1)^(1/2)
```

**3.521.2 Mathematica [A] (verified)**

Time = 0.43 (sec) , antiderivative size = 186, normalized size of antiderivative = 0.94

$$\int (fx)^m (d + ex^2) (a + \operatorname{barccosh}(cx)) dx$$

$$= x(fx)^m \left( -\frac{bcdx\sqrt{1-c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{2}, \frac{4+m}{2}, c^2x^2\right)}{(2+3m+m^2)\sqrt{-1+cx}\sqrt{1+cx}} \right. \\ \left. + \frac{\frac{(d(3+m)+e(1+m)x^2)(a+\operatorname{barccosh}(cx))}{1+m} - \frac{bcex^3\sqrt{1-c^2x^2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4+m}{2}, \frac{6+m}{2}, c^2x^2\right)}{(4+m)\sqrt{-1+cx}\sqrt{1+cx}}}{3+m} \right)$$

input `Integrate[(f*x)^m*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`output `x*(f*x)^m*(-((b*c*d*x*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, c^2*x^2])/((2 + 3*m + m^2)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])) + (((d*(3 + m) + e*(1 + m)*x^2)*(a + b*ArcCosh[c*x]))/(1 + m) - (b*c*e*x^3*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (4 + m)/2, (6 + m)/2, c^2*x^2])/((4 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(3 + m))`**3.521.3 Rubi [A] (verified)**Time = 0.42 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.238$ , Rules used = {6371, 960, 136, 279, 278}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2) (fx)^m (a + \operatorname{barccosh}(cx)) dx$$

$$\downarrow \text{6371}$$

$$-\frac{bc \int \frac{(fx)^{m+1} (e(m+1)x^2 + d(m+3))}{\sqrt{cx-1}\sqrt{cx+1}} dx}{f(m^2 + 4m + 3)} + \frac{d(fx)^{m+1} (a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{e(fx)^{m+3} (a + \operatorname{barccosh}(cx))}{f^3(m+3)}$$

$$\downarrow \text{960}$$

$$\begin{aligned}
& \frac{bc \left( \left( \frac{e^{(m+1)(m+2)}}{c^2(m+3)} + d(m+3) \right) \int \frac{(fx)^{m+1}}{\sqrt{cx-1}\sqrt{cx+1}} dx + \frac{e^{(m+1)\sqrt{cx-1}\sqrt{cx+1}}(fx)^{m+2}}{c^2 f(m+3)} \right)}{f(m^2 + 4m + 3)} + \\
& \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{e(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} \\
& \quad \downarrow 136 \\
& \frac{bc \left( \frac{\sqrt{c^2x^2-1} \left( \frac{e^{(m+1)(m+2)}}{c^2(m+3)} + d(m+3) \right) \int \frac{(fx)^{m+1}}{\sqrt{c^2x^2-1}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{e^{(m+1)\sqrt{cx-1}\sqrt{cx+1}}(fx)^{m+2}}{c^2 f(m+3)} \right)}{f(m^2 + 4m + 3)} + \\
& \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{e(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} \\
& \quad \downarrow 279 \\
& \frac{bc \left( \frac{\sqrt{1-c^2x^2} \left( \frac{e^{(m+1)(m+2)}}{c^2(m+3)} + d(m+3) \right) \int \frac{(fx)^{m+1}}{\sqrt{1-c^2x^2}} dx}{\sqrt{cx-1}\sqrt{cx+1}} + \frac{e^{(m+1)\sqrt{cx-1}\sqrt{cx+1}}(fx)^{m+2}}{c^2 f(m+3)} \right)}{f(m^2 + 4m + 3)} + \\
& \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{e(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} \\
& \quad \downarrow 278 \\
& \frac{d(fx)^{m+1}(a + \operatorname{barccosh}(cx))}{f(m+1)} + \frac{e(fx)^{m+3}(a + \operatorname{barccosh}(cx))}{f^3(m+3)} - \\
& \frac{bc \left( \frac{\sqrt{1-c^2x^2}(fx)^{m+2} \left( \frac{e^{(m+1)(m+2)}}{c^2(m+3)} + d(m+3) \right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+2}{2}, \frac{m+4}{2}, c^2x^2\right)}{f(m+2)\sqrt{cx-1}\sqrt{cx+1}} + \frac{e^{(m+1)\sqrt{cx-1}\sqrt{cx+1}}(fx)^{m+2}}{c^2 f(m+3)} \right)}{f(m^2 + 4m + 3)}
\end{aligned}$$

input `Int[(f*x)^m*(d + e*x^2)*(a + b*ArcCosh[c*x]),x]`

output `(d*(f*x)^(1 + m)*(a + b*ArcCosh[c*x]))/(f*(1 + m)) + (e*(f*x)^(3 + m)*(a + b*ArcCosh[c*x]))/(f^3*(3 + m)) - (b*c*((e*(1 + m)*(f*x)^(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(c^2*f*(3 + m)) + (((e*(1 + m)*(2 + m))/(c^2*(3 + m)) + d*(3 + m))*(f*x)^(2 + m)*Sqrt[1 - c^2*x^2]*Hypergeometric2F1[1/2, (2 + m)/2, (4 + m)/2, c^2*x^2])/(f*(2 + m)*Sqrt[-1 + c*x]*Sqrt[1 + c*x]))/(f*(3 + 4*m + m^2))`

## 3.521.3.1 Defintions of rubi rules used

rule 136 `Int[((f_.)*(x_))^(p_.)*((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_] := Simp[(a + b*x)^FracPart[m]*((c + d*x)^FracPart[m]/(a*c + b*d*x^2)^FracPart[m]) Int[(a*c + b*d*x^2)^m*(f*x)^p, x], x] /; FreeQ[{a, b, c, d, f, m, n, p}, x] && EqQ[b*c + a*d, 0] && EqQ[n, m]`

rule 278 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/2, (m + 1)/2 + 1, (-b)*(x^2/a)], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 279 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[a^IntPart[p]*((a + b*x^2)^FracPart[p]/(1 + b*(x^2/a))^FracPart[p]) Int[(c*x)^m*(1 + b*(x^2/a))^p, x], x] /; FreeQ[{a, b, c, m, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])`

rule 960 `Int[((e_.)*(x_))^(m_.)*((a1_) + (b1_.)*(x_)^(non2_.))^(p_.)*((a2_) + (b2_.)*(x_)^(non2_.))^(p_.)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a1 + b1*x^(n/2))^(p + 1)*((a2 + b2*x^(n/2))^(p + 1)/(b1*b2*e*(m + n*(p + 1) + 1))), x] - Simp[(a1*a2*d*(m + 1) - b1*b2*c*(m + n*(p + 1) + 1))/(b1*b2*(m + n*(p + 1) + 1)) Int[(e*x)^m*(a1 + b1*x^(n/2))^p*(a2 + b2*x^(n/2))^p, x], x] /; FreeQ[{a1, b1, a2, b2, c, d, e, m, n, p}, x] && EqQ[non2, n/2] && EqQ[a2*b1 + a1*b2, 0] && NeQ[m + n*(p + 1) + 1, 0]`

rule 6371 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2), x_Symbol] := Simp[d*(f*x)^(m + 1)*((a + b*ArcCosh[c*x])/(f*(m + 1))), x] + (Simp[e*(f*x)^(m + 3)*((a + b*ArcCosh[c*x])/(f^3*(m + 3))), x] - Simp[b*(c/(f*(m + 1)*(m + 3))) Int[(f*x)^(m + 1)*((d*(m + 3) + e*(m + 1)*x^2)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])], x], x]) /; FreeQ[{a, b, c, d, e, f, m}, x] && NeQ[c^2*d + e, 0] && NeQ[m, -1] && NeQ[m, -3]`

**3.521.4 Maple [F]**

$$\int (fx)^m (ex^2 + d) (a + b \operatorname{arccosh}(cx)) dx$$

input `int((f*x)^m*(e*x^2+d)*(a+b*arccosh(c*x)),x)`

output `int((f*x)^m*(e*x^2+d)*(a+b*arccosh(c*x)),x)`

**3.521.5 Fracas [F]**

$$\int (fx)^m (d + ex^2) (a + b \operatorname{arccosh}(cx)) dx = \int (ex^2 + d) (b \operatorname{arccosh}(cx) + a) (fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="fracas")`

output `integral((a*e*x^2 + a*d + (b*e*x^2 + b*d)*arccosh(c*x))*(f*x)^m, x)`

**3.521.6 Sympy [F]**

$$\int (fx)^m (d + ex^2) (a + b \operatorname{arccosh}(cx)) dx = \int (fx)^m (a + b \operatorname{acosh}(cx)) (d + ex^2) dx$$

input `integrate((f*x)**m*(e*x**2+d)*(a+b*acosh(c*x)),x)`

output `Integral((f*x)**m*(a + b*acosh(c*x))*(d + e*x**2), x)`

**3.521.7 Maxima [F]**

$$\int (fx)^m (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \int (ex^2 + d)(b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `a*e*f^m*x^3*x^m/(m + 3) + (b*e*f^m*(m + 1)*x^3 + b*d*f^m*(m + 3)*x)*x^m*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))/(m^2 + 4*m + 3) + (f*x)^(m + 1)*a*d/(f*(m + 1)) + integrate((b*c*e*f^m*(m + 1)*x^3 + b*c*d*f^m*(m + 3)*x)*x^m/(m^2 + 4*m + 3)*c^3*x^3 - (m^2 + 4*m + 3)*c*x + ((m^2 + 4*m + 3)*c^2*x^2 - m^2 - 4*m - 3)*sqrt(c*x + 1)*sqrt(c*x - 1)), x) - integrate((b*c^2*e*f^m*(m + 1)*x^4 + b*c^2*d*f^m*(m + 3)*x^2)*x^m/((m^2 + 4*m + 3)*c^2*x^2 - m^2 - 4*m - 3), x)`

**3.521.8 Giac [F]**

$$\int (fx)^m (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \int (ex^2 + d)(b \operatorname{arcosh}(cx) + a)(fx)^m dx$$

input `integrate((f*x)^m*(e*x^2+d)*(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)*(b*arccosh(c*x) + a)*(f*x)^m, x)`

**3.521.9 Mupad [F(-1)]**

Timed out.

$$\int (fx)^m (d + ex^2) (a + \operatorname{barccosh}(cx)) dx = \int (a + b \operatorname{acosh}(cx)) (fx)^m (ex^2 + d) dx$$

input `int((a + b*acosh(c*x))*(f*x)^m*(d + e*x^2),x)`

output `int((a + b*acosh(c*x))*(f*x)^m*(d + e*x^2), x)`

**3.522**  $\int \frac{(fx)^m(a+b\text{arccosh}(cx))}{d+ex^2} dx$

3.522.1 Optimal result . . . . . 3854  
 3.522.2 Mathematica [N/A] . . . . . 3854  
 3.522.3 Rubi [N/A] . . . . . 3855  
 3.522.4 Maple [N/A] (verified) . . . . . 3855  
 3.522.5 Fricas [N/A] . . . . . 3856  
 3.522.6 Sympy [N/A] . . . . . 3856  
 3.522.7 Maxima [N/A] . . . . . 3856  
 3.522.8 Giac [N/A] . . . . . 3857  
 3.522.9 Mupad [N/A] . . . . . 3857

**3.522.1 Optimal result**

Integrand size = 23, antiderivative size = 23

$$\int \frac{(fx)^m(a + \text{barccosh}(cx))}{d + ex^2} dx = \text{Int}\left(\frac{(fx)^m(a + \text{barccosh}(cx))}{d + ex^2}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d),x)`

**3.522.2 Mathematica [N/A]**

Not integrable

Time = 2.86 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m(a + \text{barccosh}(cx))}{d + ex^2} dx = \int \frac{(fx)^m(a + \text{barccosh}(cx))}{d + ex^2} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2), x]`

**3.522.3 Rubi [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d + ex^2} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d + ex^2} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2),x]`

output `$Aborted`

**3.522.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_.)]*(b_.))^(n_.)*((f_.)*(x_.))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.522.4 Maple [N/A] (verified)**

Not integrable

Time = 2.47 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{ex^2 + d} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d),x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d),x)`



**3.522.5 Fracas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{ex^2 + d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="fricas")`output `integral((b*arccosh(c*x) + a)*(f*x)^m/(e*x^2 + d), x)`**3.522.6 Sympy [N/A]**

Not integrable

Time = 16.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d + ex^2} dx = \int \frac{(fx)^m (a + b \operatorname{acosh}(cx))}{d + ex^2} dx$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(e*x**2+d),x)`output `Integral((f*x)**m*(a + b*acosh(c*x))/(d + e*x**2), x)`**3.522.7 Maxima [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{ex^2 + d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="maxima")`output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(e*x^2 + d), x)`

---

3.522.  $\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{d + ex^2} dx$

**3.522.8 Giac [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{ex^2 + d} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(e*x^2 + d), x)`**3.522.9 Mupad [N/A]**

Not integrable

Time = 3.55 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{d + ex^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{ex^2 + d} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d + e*x^2),x)`output `int(((a + b*acosh(c*x))*(f*x)^m)/(d + e*x^2), x)`

$$3.523 \quad \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx$$

3.523.1 Optimal result	3858
3.523.2 Mathematica [N/A]	3858
3.523.3 Rubi [N/A]	3859
3.523.4 Maple [N/A] (verified)	3859
3.523.5 Fricas [N/A]	3860
3.523.6 Sympy [F(-1)]	3860
3.523.7 Maxima [N/A]	3860
3.523.8 Giac [N/A]	3861
3.523.9 Mupad [N/A]	3861

### 3.523.1 Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx = \operatorname{Int} \left( \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2}, x \right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^2,x)`

### 3.523.2 Mathematica [N/A]

Not integrable

Time = 2.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2, x]`

---

3.523.  $\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^2} dx$

**3.523.3 Rubi [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2)^2,x]`

output `$Aborted`

**3.523.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.523.4 Maple [N/A] (verified)**

Not integrable

Time = 1.38 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(ex^2 + d)^2} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^2,x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^2,x)`

**3.523.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.57

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(ex^2 + d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="fricas")`output `integral((b*arccosh(c*x) + a)*(f*x)^m/(e^2*x^4 + 2*d*e*x^2 + d^2), x)`**3.523.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(e*x**2+d)**2,x)`output `Timed out`**3.523.7 Maxima [N/A]**

Not integrable

Time = 0.32 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(ex^2 + d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="maxima")`output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(e*x^2 + d)^2, x)`

**3.523.8 Giac [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(ex^2 + d)^2} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^2,x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(e*x^2 + d)^2, x)`**3.523.9 Mupad [N/A]**

Not integrable

Time = 3.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d + ex^2)^2} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{(ex^2 + d)^2} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d + e*x^2)^2,x)`output `int(((a + b*acosh(c*x))*(f*x)^m)/(d + e*x^2)^2, x)`

$$3.524 \quad \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$$

3.524.1 Optimal result	3862
3.524.2 Mathematica [N/A]	3862
3.524.3 Rubi [N/A]	3863
3.524.4 Maple [N/A] (verified)	3863
3.524.5 Fricas [N/A]	3864
3.524.6 Sympy [F(-1)]	3864
3.524.7 Maxima [N/A]	3864
3.524.8 Giac [N/A]	3865
3.524.9 Mupad [N/A]	3865

### 3.524.1 Optimal result

Integrand size = 23, antiderivative size = 23

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx = \operatorname{Int}\left(\frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3}, x\right)$$

output `Unintegrable((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^3,x)`

### 3.524.2 Mathematica [N/A]

Not integrable

Time = 3.60 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$$

input `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

output `Integrate[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3, x]`

**3.524.3 Rubi [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6375}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$$

↓ 6375

$$\int \frac{(fx)^m (a + \operatorname{arccosh}(cx))}{(d + ex^2)^3} dx$$

input `Int[((f*x)^m*(a + b*ArcCosh[c*x]))/(d + e*x^2)^3,x]`

output `$Aborted`

**3.524.3.1 Defintions of rubi rules used**

rule 6375 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((f_.)*(x_))^(m_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(f*x)^m*(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]`

**3.524.4 Maple [N/A] (verified)**

Not integrable

Time = 2.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{(fx)^m (a + b \operatorname{arccosh}(cx))}{(ex^2 + d)^3} dx$$

input `int((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^3,x)`

output `int((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^3,x)`



**3.524.5 Fracas [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.04

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(ex^2 + d)^3} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="fricas")`

output `integral((b*arccosh(c*x) + a)*(f*x)^m/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**3.524.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \text{Timed out}$$

input `integrate((f*x)**m*(a+b*acosh(c*x))/(e*x**2+d)**3,x)`

output `Timed out`

**3.524.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m(a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(ex^2 + d)^3} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(e*x^2 + d)^3, x)`

**3.524.8 Giac [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)(fx)^m}{(ex^2 + d)^3} dx$$

input `integrate((f*x)^m*(a+b*arccosh(c*x))/(e*x^2+d)^3,x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)*(f*x)^m/(e*x^2 + d)^3, x)`**3.524.9 Mupad [N/A]**

Not integrable

Time = 3.31 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \frac{(fx)^m (a + \operatorname{barccosh}(cx))}{(d + ex^2)^3} dx = \int \frac{(a + b \operatorname{acosh}(cx)) (fx)^m}{(ex^2 + d)^3} dx$$

input `int(((a + b*acosh(c*x))*(f*x)^m)/(d + e*x^2)^3,x)`output `int(((a + b*acosh(c*x))*(f*x)^m)/(d + e*x^2)^3, x)`

### 3.525 $\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx))^2 dx$

3.525.1 Optimal result . . . . .	3867
3.525.2 Mathematica [A] (verified) . . . . .	3868
3.525.3 Rubi [A] (verified) . . . . .	3869
3.525.4 Maple [A] (verified) . . . . .	3870
3.525.5 Fricas [A] (verification not implemented) . . . . .	3871
3.525.6 Sympy [F] . . . . .	3872
3.525.7 Maxima [A] (verification not implemented) . . . . .	3873
3.525.8 Giac [F(-2)] . . . . .	3874
3.525.9 Mupad [F(-1)] . . . . .	3874

**3.525.1 Optimal result**

Integrand size = 20, antiderivative size = 609

$$\begin{aligned}
\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx))^2 dx = & 2b^2 d^3 x + \frac{4b^2 d^2 ex}{3c^2} + \frac{16b^2 de^2 x}{25c^4} + \frac{32b^2 e^3 x}{245c^6} \\
& + \frac{2}{9} b^2 d^2 ex^3 + \frac{8b^2 de^2 x^3}{75c^2} + \frac{16b^2 e^3 x^3}{735c^4} \\
& + \frac{6}{125} b^2 de^2 x^5 + \frac{12b^2 e^3 x^5}{1225c^2} + \frac{2}{343} b^2 e^3 x^7 \\
& - \frac{2bd^3 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{c} \\
& - \frac{4bd^2 e \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{3c^3} \\
& - \frac{16bde^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{25c^5} \\
& - \frac{32be^3 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{245c^7} \\
& - \frac{2bd^2 ex^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{3c} \\
& - \frac{8bde^2 x^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{25c^3} \\
& - \frac{16be^3 x^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{245c^5} \\
& - \frac{6bde^2 x^4 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{25c} \\
& - \frac{12be^3 x^4 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{245c^3} \\
& - \frac{2be^3 x^6 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{49c} \\
& + d^3 x (a + \operatorname{barccosh}(cx))^2 + d^2 ex^3 (a + \operatorname{barccosh}(cx))^2 \\
& + \frac{3}{5} de^2 x^5 (a + \operatorname{barccosh}(cx))^2 \\
& + \frac{1}{7} e^3 x^7 (a + \operatorname{barccosh}(cx))^2
\end{aligned}$$

output  $2*b^2*d^3*x+4/3*b^2*d^2*e*x/c^2+16/25*b^2*d*e^2*x/c^4+32/245*b^2*e^3*x/c^6+2/9*b^2*d^2*e*x^3+8/75*b^2*d*e^2*x^3/c^2+16/735*b^2*e^3*x^3/c^4+6/125*b^2*d*e^2*x^5+12/1225*b^2*e^3*x^5/c^2+2/343*b^2*e^3*x^7+d^3*x*(a+b*arccosh(c*x))^2+d^2*e*x^3*(a+b*arccosh(c*x))^2+3/5*d*e^2*x^5*(a+b*arccosh(c*x))^2+1/7*e^3*x^7*(a+b*arccosh(c*x))^2-2*b*d^3*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-4/3*b*d^2*e*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-16/25*b*d*e^2*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5-32/245*b*e^3*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^7-2/3*b*d^2*e*x^2*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-8/25*b*d*e^2*x^2*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-16/245*b*e^3*x^2*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^5-6/25*b*d*e^2*x^4*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-12/245*b*e^3*x^4*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-2/49*b*e^3*x^6*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c$

### 3.525.2 Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 453, normalized size of antiderivative = 0.74

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx))^2 dx$$

$$= \frac{11025a^2c^7x(35d^3 + 35d^2ex^2 + 21de^2x^4 + 5e^3x^6) - 210ab\sqrt{-1 + cx}\sqrt{1 + cx}(240e^3 + 24c^2e^2(49d + 5ex^2))}{c^7}$$

input `Integrate[(d + e*x^2)^3*(a + b*ArcCosh[c*x])^2,x]`

output  $(11025*a^2*c^7*x*(35*d^3 + 35*d^2*e*x^2 + 21*d*e^2*x^4 + 5*e^3*x^6) - 210*a*b*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(240*e^3 + 24*c^2*e^2*(49*d + 5*e*x^2) + 2*c^4*e*(1225*d^2 + 294*d*e*x^2 + 45*e^2*x^4) + c^6*(3675*d^3 + 1225*d^2*e*x^2 + 441*d*e^2*x^4 + 75*e^3*x^6)) + 2*b^2*c*x*(25200*e^3 + 840*c^2*e^2*(147*d + 5*e*x^2) + 210*c^4*e*(1225*d^2 + 98*d*e*x^2 + 9*e^2*x^4) + c^6*(385875*d^3 + 42875*d^2*e*x^2 + 9261*d*e^2*x^4 + 1125*e^3*x^6)) - 210*b*(-105*a*c^7*x*(35*d^3 + 35*d^2*e*x^2 + 21*d*e^2*x^4 + 5*e^3*x^6) + b*\operatorname{Sqrt}[-1 + c*x]*\operatorname{Sqrt}[1 + c*x]*(240*e^3 + 24*c^2*e^2*(49*d + 5*e*x^2) + 2*c^4*e*(1225*d^2 + 294*d*e*x^2 + 45*e^2*x^4) + c^6*(3675*d^3 + 1225*d^2*e*x^2 + 441*d*e^2*x^4 + 75*e^3*x^6)))*\operatorname{ArcCosh}[c*x] + 11025*b^2*c^7*x*(35*d^3 + 35*d^2*e*x^2 + 21*d*e^2*x^4 + 5*e^3*x^6)*\operatorname{ArcCosh}[c*x]^2)/(385875*c^7)$

**3.525.3 Rubi [A] (verified)**

Time = 2.39 (sec) , antiderivative size = 609, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx))^2 dx$$

↓ 6324

$$\int (d^3(a + \operatorname{barccosh}(cx))^2 + 3d^2ex^2(a + \operatorname{barccosh}(cx))^2 + 3de^2x^4(a + \operatorname{barccosh}(cx))^2 + e^3x^6(a + \operatorname{barccosh}(cx))^2)$$

↓ 2009

$$\begin{aligned} & - \frac{32be^3\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{245c^7} - \frac{16bde^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{25c^5} - \\ & \frac{16be^3x^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{245c^5} - \frac{4bd^2e\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{3c^3} - \\ & \frac{8bde^2x^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{25c^3} - \frac{12be^3x^4\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{245c^3} + \\ & d^3x(a + \operatorname{barccosh}(cx))^2 - \frac{2bd^3\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{c} + d^2ex^3(a + \operatorname{barccosh}(cx))^2 - \\ & \frac{2bd^2ex^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{3c} + \frac{3}{5}de^2x^5(a + \operatorname{barccosh}(cx))^2 - \\ & \frac{6bde^2x^4\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{25c} + \frac{1}{7}e^3x^7(a + \operatorname{barccosh}(cx))^2 - \\ & \frac{2be^3x^6\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{49c} + \frac{32b^2e^3x}{245c^6} + \frac{16b^2de^2x}{25c^4} + \frac{16b^2e^3x^3}{735c^4} + \frac{4b^2d^2ex}{3c^2} + \\ & \frac{8b^2de^2x^3}{75c^2} + \frac{12b^2e^3x^5}{1225c^2} + 2b^2d^3x + \frac{2}{9}b^2d^2ex^3 + \frac{6}{125}b^2de^2x^5 + \frac{2}{343}b^2e^3x^7 \end{aligned}$$

input `Int[(d + e*x^2)^3*(a + b*ArcCosh[c*x])^2,x]`

output  $2*b^2*d^3*x + (4*b^2*d^2*e*x)/(3*c^2) + (16*b^2*d*e^2*x)/(25*c^4) + (32*b^2*e^3*x)/(245*c^6) + (2*b^2*d^2*e*x^3)/9 + (8*b^2*d*e^2*x^3)/(75*c^2) + (16*b^2*e^3*x^3)/(735*c^4) + (6*b^2*d*e^2*x^5)/125 + (12*b^2*e^3*x^5)/(1225*c^2) + (2*b^2*e^3*x^7)/343 - (2*b*d^3*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/c - (4*b*d^2*e*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(3*c^3) - (16*b*d*e^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(25*c^5) - (32*b*e^3*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(245*c^7) - (2*b*d^2*e*x^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(3*c) - (8*b*d*e^2*x^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(25*c^3) - (16*b*e^3*x^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(245*c^5) - (6*b*d*e^2*x^4*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(25*c) - (12*b*e^3*x^4*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(245*c^3) - (2*b*e^3*x^6*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(49*c) + d^3*x*(a + b*ArcCosh[c*x])^2 + d^2*e*x^3*(a + b*ArcCosh[c*x])^2 + (3*d*e^2*x^5*(a + b*ArcCosh[c*x])^2)/5 + (e^3*x^7*(a + b*ArcCosh[c*x])^2)/7$

### 3.525.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

### 3.525.4 Maple [A] (verified)

Time = 0.81 (sec) , antiderivative size = 632, normalized size of antiderivative = 1.04

method	result
derivativedivides	$\frac{a^2(d^3c^7x + d^2c^7ex^3 + \frac{3}{5}dc^7e^2x^5 + \frac{1}{7}e^3c^7x^7)}{c^6} + \frac{b^2\left(c^6d^3(\operatorname{arccosh}(cx))^2xc - 2\operatorname{arccosh}(cx)\sqrt{cx-1}\sqrt{cx+1} + 2cx\right) + c^4d^2e(9\operatorname{arccosh}(cx))^2}{c^6}$
default	$\frac{a^2(d^3c^7x + d^2c^7ex^3 + \frac{3}{5}dc^7e^2x^5 + \frac{1}{7}e^3c^7x^7)}{c^6} + \frac{b^2\left(c^6d^3(\operatorname{arccosh}(cx))^2xc - 2\operatorname{arccosh}(cx)\sqrt{cx-1}\sqrt{cx+1} + 2cx\right) + c^4d^2e(9\operatorname{arccosh}(cx))^2}{c^6}$
parts	$a^2\left(\frac{1}{7}e^3x^7 + \frac{3}{5}de^2x^5 + d^2ex^3 + d^3x\right) + \frac{b^2(-15750\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6x^6e^3 - 92610\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6x^5e^3 - 15750\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6x^4e^3 - 92610\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6x^3e^3 - 15750\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6x^2e^3 - 92610\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6xe^3 - 15750\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6e^3 - 92610\sqrt{cx+1}\sqrt{cx-1}\operatorname{arccosh}(cx)c^6)}{c^6}$

3.525.  $\int (d + ex^2)^3 (a + b\operatorname{arccosh}(cx))^2 dx$

input `int((e*x^2+d)^3*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{c} \left( \frac{a^2}{c^6} (d^3 c^7 x + d^2 c^7 e x^3 + \frac{3}{5} d c^7 e^2 x^5 + \frac{1}{7} e^3 c^7 x^7) + \frac{b^2}{c^6} (c^6 d^3 (\operatorname{arccosh}(c x))^2 x c - 2 \operatorname{arccosh}(c x) (c x - 1)^{1/2} (c x + 1)^{1/2} + 2 c x) + \frac{1}{9} c^4 d^2 e (9 \operatorname{arccosh}(c x)^2 x^3 c^3 - 6 (c x + 1)^{1/2} \operatorname{arccosh}(c x) (c x - 1)^{1/2} c^2 x^2 - 12 \operatorname{arccosh}(c x) (c x - 1)^{1/2} (c x + 1)^{1/2} + 2 c^3 x^3 + 12 c x) + \frac{1}{375} c^2 d e^2 (225 \operatorname{arccosh}(c x)^2 c^5 x^5 - 90 (c x - 1)^{1/2} (c x + 1)^{1/2} \operatorname{arccosh}(c x) c^4 x^4 - 120 (c x + 1)^{1/2} \operatorname{arccosh}(c x) (c x - 1)^{1/2} c^2 x^2 + 18 c^5 x^5 - 240 \operatorname{arccosh}(c x) (c x - 1)^{1/2} (c x + 1)^{1/2} + 40 c^3 x^3 + 240 c x) + \frac{1}{25725} e^3 (3675 \operatorname{arccosh}(c x)^2 c^7 x^7 - 1050 \operatorname{arccosh}(c x) (c x - 1)^{1/2} (c x + 1)^{1/2} c^6 x^6 - 1260 (c x - 1)^{1/2} (c x + 1)^{1/2} \operatorname{arccosh}(c x) c^4 x^4 + 150 c^7 x^7 - 1680 (c x + 1)^{1/2} \operatorname{arccosh}(c x) (c x - 1)^{1/2} c^2 x^2 + 252 c^5 x^5 - 3360 \operatorname{arccosh}(c x) (c x - 1)^{1/2} (c x + 1)^{1/2} + 560 c^3 x^3 + 3360 c x) \right) + 2 a b / c^6 (\operatorname{arccosh}(c x) d^3 c^7 x + \operatorname{arccosh}(c x) d^2 c^7 e x^3 + \frac{3}{5} \operatorname{arccosh}(c x) d c^7 e^2 x^5 + \frac{1}{7} \operatorname{arccosh}(c x) e^3 c^7 x^7 - \frac{1}{3675} (c x - 1)^{1/2} (c x + 1)^{1/2} (75 c^6 e^3 x^6 + 441 c^6 d e^2 x^4 + 1225 c^6 d^2 e e x^2 + 90 c^4 e^3 x^4 + 3675 c^6 d^3 + 588 c^4 d e^2 x^2 + 2450 c^4 d^2 e + 120 c^2 e^3 x^2 + 1176 c^2 d e^2 + 240 e^3))$$

### 3.525.5 Fracas [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 586, normalized size of antiderivative = 0.96

$$\int (d + ex^2)^3 (a + b \operatorname{arccosh}(cx))^2 dx$$

$$= \frac{1125 (49 a^2 + 2 b^2) c^7 e^3 x^7 + 189 (49 (25 a^2 + 2 b^2) c^7 d e^2 + 20 b^2 c^5 e^3) x^5 + 35 (1225 (9 a^2 + 2 b^2) c^7 d^2 e + 1176$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))^2,x, algorithm="fracas")`



output  $\frac{1}{385875} \cdot (1125 \cdot (49a^2 + 2b^2) \cdot c^7 \cdot e^3 \cdot x^7 + 189 \cdot (49 \cdot (25a^2 + 2b^2) \cdot c^7 \cdot d \cdot e^2 + 20b^2 \cdot c^5 \cdot e^3) \cdot x^5 + 35 \cdot (1225 \cdot (9a^2 + 2b^2) \cdot c^7 \cdot d^2 \cdot e + 1176b^2 \cdot c^5 \cdot d \cdot e^2 + 240b^2 \cdot c^3 \cdot e^3) \cdot x^3 + 11025 \cdot (5b^2 \cdot c^7 \cdot e^3 \cdot x^7 + 21b^2 \cdot c^7 \cdot d \cdot e^2 \cdot x^5 + 35b^2 \cdot c^7 \cdot d^2 \cdot e \cdot x^3 + 35b^2 \cdot c^7 \cdot d^3 \cdot x) \cdot \log(cx + \sqrt{c^2x^2 - 1})^2 + 105 \cdot (3675 \cdot (a^2 + 2b^2) \cdot c^7 \cdot d^3 + 4900b^2 \cdot c^5 \cdot d^2 \cdot e + 2352b^2 \cdot c^3 \cdot d \cdot e^2 + 480b^2 \cdot c \cdot e^3) \cdot x + 210 \cdot (525 \cdot a \cdot b \cdot c^7 \cdot e^3 \cdot x^7 + 2205 \cdot a \cdot b \cdot c^7 \cdot d \cdot e^2 \cdot x^5 + 3675 \cdot a \cdot b \cdot c^7 \cdot d^2 \cdot e \cdot x^3 + 3675 \cdot a \cdot b \cdot c^7 \cdot d^3 \cdot x - (75b^2 \cdot c^6 \cdot e^3 \cdot x^6 + 3675b^2 \cdot c^6 \cdot d^3 + 2450b^2 \cdot c^4 \cdot d^2 \cdot e + 1176b^2 \cdot c^2 \cdot d \cdot e^2 + 240b^2 \cdot e^3 + 9 \cdot (49b^2 \cdot c^6 \cdot d \cdot e^2 + 10b^2 \cdot c^4 \cdot e^3) \cdot x^4 + (1225b^2 \cdot c^6 \cdot d^2 \cdot e + 588b^2 \cdot c^4 \cdot d \cdot e^2 + 120b^2 \cdot c^2 \cdot e^3) \cdot x^2) \cdot \sqrt{c^2x^2 - 1}) \cdot \log(cx + \sqrt{c^2x^2 - 1}) - 210 \cdot (75 \cdot a \cdot b \cdot c^6 \cdot e^3 \cdot x^6 + 3675 \cdot a \cdot b \cdot c^6 \cdot d^3 + 2450 \cdot a \cdot b \cdot c^4 \cdot d^2 \cdot e + 1176 \cdot a \cdot b \cdot c^2 \cdot d \cdot e^2 + 240 \cdot a \cdot b \cdot e^3 + 9 \cdot (49 \cdot a \cdot b \cdot c^6 \cdot d \cdot e^2 + 10 \cdot a \cdot b \cdot c^4 \cdot e^3) \cdot x^4 + (1225 \cdot a \cdot b \cdot c^6 \cdot d^2 \cdot e + 588 \cdot a \cdot b \cdot c^4 \cdot d \cdot e^2 + 120 \cdot a \cdot b \cdot c^2 \cdot e^3) \cdot x^2) \cdot \sqrt{c^2x^2 - 1}) / c^7$

### 3.525.6 Sympy [F]

$$\int (d + ex^2)^3 (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (d + ex^2)^3 dx$$

input `integrate((e*x**2+d)**3*(a+b*acosh(c*x))**2,x)`

output `Integral((a + b*acosh(c*x))**2*(d + e*x**2)**3, x)`

**3.525.7 Maxima [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 684, normalized size of antiderivative = 1.12

$$\begin{aligned}
\int (d + ex^2)^3 (a + \operatorname{arccosh}(cx))^2 dx &= \frac{1}{7} b^2 e^3 x^7 \operatorname{arccosh}(cx)^2 + \frac{1}{7} a^2 e^3 x^7 \\
&+ \frac{3}{5} b^2 d e^2 x^5 \operatorname{arccosh}(cx)^2 + \frac{3}{5} a^2 d e^2 x^5 + b^2 d^2 e x^3 \operatorname{arccosh}(cx)^2 + a^2 d^2 e x^3 \\
&+ b^2 d^3 x \operatorname{arccosh}(cx)^2 + \frac{2}{3} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \right) a b d^2 e \\
&- \frac{2}{9} \left( 3c \left( \frac{\sqrt{c^2 x^2 - 1} x^2}{c^2} + \frac{2\sqrt{c^2 x^2 - 1}}{c^4} \right) \operatorname{arccosh}(cx) - \frac{c^2 x^3 + 6x}{c^2} \right) b^2 d^2 e \\
&+ \frac{2}{25} \left( 15x^5 \operatorname{arccosh}(cx) - \left( \frac{3\sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4\sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8\sqrt{c^2 x^2 - 1}}{c^6} \right) c \right) a b d e^2 \\
&- \frac{2}{375} \left( 15 \left( \frac{3\sqrt{c^2 x^2 - 1} x^4}{c^2} + \frac{4\sqrt{c^2 x^2 - 1} x^2}{c^4} + \frac{8\sqrt{c^2 x^2 - 1}}{c^6} \right) c \operatorname{arccosh}(cx) - \frac{9c^4 x^5 + 20c^2 x^3 + 120x}{c^4} \right) a b e^2 \\
&+ \frac{2}{245} \left( 35x^7 \operatorname{arccosh}(cx) - \left( \frac{5\sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6\sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8\sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16\sqrt{c^2 x^2 - 1}}{c^8} \right) c \right) a b e^3 \\
&- \frac{2}{25725} \left( 105 \left( \frac{5\sqrt{c^2 x^2 - 1} x^6}{c^2} + \frac{6\sqrt{c^2 x^2 - 1} x^4}{c^4} + \frac{8\sqrt{c^2 x^2 - 1} x^2}{c^6} + \frac{16\sqrt{c^2 x^2 - 1}}{c^8} \right) c \operatorname{arccosh}(cx) - \frac{75c^4 x^7 + 210c^2 x^5 + 120x}{c^4} \right) a b e^3 \\
&+ 2b^2 d^3 \left( x - \frac{\sqrt{c^2 x^2 - 1} \operatorname{arccosh}(cx)}{c} \right) + a^2 d^3 x + \frac{2(cx \operatorname{arccosh}(cx) - \sqrt{c^2 x^2 - 1}) a b d^3}{c}
\end{aligned}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $\frac{1}{7}b^2e^3x^7\operatorname{arccosh}(cx)^2 + \frac{1}{7}a^2e^3x^7 + \frac{3}{5}b^2d^2e^2x^5\operatorname{arccosh}(cx)^2 + \frac{3}{5}a^2d^2e^2x^5 + b^2d^2e^2x^3\operatorname{arccosh}(cx)^2 + a^2d^2e^2x^3 + b^2d^3x\operatorname{arccosh}(cx)^2 + \frac{2}{3}(3x^3\operatorname{arccosh}(cx) - c(\sqrt{c^2x^2 - 1})x^2/c^2 + 2\sqrt{c^2x^2 - 1}/c^4)*a*b*d^2e - \frac{2}{9}(3c(\sqrt{c^2x^2 - 1})x^2/c^2 + 2\sqrt{c^2x^2 - 1}/c^4)*\operatorname{arccosh}(cx) - (c^2x^3 + 6x)/c^2)*b^2d^2e + \frac{2}{25}(15x^5\operatorname{arccosh}(cx) - (3\sqrt{c^2x^2 - 1})x^4/c^2 + 4\sqrt{c^2x^2 - 1})x^2/c^4 + 8\sqrt{c^2x^2 - 1}/c^6)*c)*a*b*d^2e - \frac{2}{3}75(15(3\sqrt{c^2x^2 - 1})x^4/c^2 + 4\sqrt{c^2x^2 - 1})x^2/c^4 + 8\sqrt{c^2x^2 - 1}/c^6)*c*\operatorname{arccosh}(cx) - (9c^4x^5 + 20c^2x^3 + 120x)/c^4)*b^2d^2e + \frac{2}{245}(35x^7\operatorname{arccosh}(cx) - (5\sqrt{c^2x^2 - 1})x^6/c^2 + 6\sqrt{c^2x^2 - 1})x^4/c^4 + 8\sqrt{c^2x^2 - 1})x^2/c^6 + 16\sqrt{c^2x^2 - 1}/c^8)*c)*a*b*e^3 - \frac{2}{25725}(105(5\sqrt{c^2x^2 - 1})x^6/c^2 + 6\sqrt{c^2x^2 - 1})x^4/c^4 + 8\sqrt{c^2x^2 - 1})x^2/c^6 + 16\sqrt{c^2x^2 - 1}/c^8)*c*\operatorname{arccosh}(cx) - (75c^6x^7 + 126c^4x^5 + 280c^2x^3 + 1680x)/c^6)*b^2e^3 + 2*b^2*d^3*(x - \sqrt{c^2x^2 - 1})*\operatorname{arccosh}(cx)/c + a^2*d^3*x + 2*(c*x*\operatorname{arccosh}(cx) - \sqrt{c^2x^2 - 1})*a*b*d^3/c$

### 3.525.8 Giac [F(-2)]

Exception generated.

$$\int (d + ex^2)^3 (a + \operatorname{arccosh}(cx))^2 dx = \text{Exception raised: RuntimeError}$$

input `integrate((e*x^2+d)^3*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const vector & l) Error: Bad Argument Value

### 3.525.9 Mupad [F(-1)]

Timed out.

$$\int (d + ex^2)^3 (a + \operatorname{arccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (ex^2 + d)^3 dx$$

input `int((a + b*acosh(c*x))^2*(d + e*x^2)^3,x)`

output `int((a + b*acosh(c*x))^2*(d + e*x^2)^3, x)`

---

3.525.  $\int (d + ex^2)^3 (a + \operatorname{arccosh}(cx))^2 dx$

### 3.526 $\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx$

3.526.1 Optimal result . . . . .	3875
3.526.2 Mathematica [A] (verified) . . . . .	3876
3.526.3 Rubi [A] (verified) . . . . .	3876
3.526.4 Maple [A] (verified) . . . . .	3878
3.526.5 Fricas [A] (verification not implemented) . . . . .	3878
3.526.6 Sympy [F] . . . . .	3879
3.526.7 Maxima [A] (verification not implemented) . . . . .	3879
3.526.8 Giac [F(-2)] . . . . .	3880
3.526.9 Mupad [F(-1)] . . . . .	3880

#### 3.526.1 Optimal result

Integrand size = 20, antiderivative size = 359

$$\begin{aligned}
 \int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx = & 2b^2 d^2 x + \frac{8b^2 dex}{9c^2} + \frac{16b^2 e^2 x}{75c^4} \\
 & + \frac{4}{27} b^2 dex^3 + \frac{8b^2 e^2 x^3}{225c^2} + \frac{2}{125} b^2 e^2 x^5 \\
 & - \frac{2bd^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{c} \\
 & - \frac{8bde \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{9c^3} \\
 & - \frac{16be^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{75c^5} \\
 & - \frac{4bdex^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{9c} \\
 & - \frac{8be^2 x^2 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{75c^3} \\
 & - \frac{2be^2 x^4 \sqrt{-1 + cx} \sqrt{1 + cx} (a + \operatorname{barccosh}(cx))}{25c} \\
 & + d^2 x (a + \operatorname{barccosh}(cx))^2 + \frac{2}{3} dex^3 (a + \operatorname{barccosh}(cx))^2 \\
 & + \frac{1}{5} e^2 x^5 (a + \operatorname{barccosh}(cx))^2
 \end{aligned}$$

output  $2*b^2*d^2*x+8/9*b^2*d*e*x/c^2+16/75*b^2*e^2*x/c^4+4/27*b^2*d*e*x^3+8/225*b^2*e^2*x^3/c^2+2/125*b^2*e^2*x^5+d^2*x*(a+b*arccosh(c*x))^2+2/3*d*e*x^3*(a+b*arccosh(c*x))^2+1/5*e^2*x^5*(a+b*arccosh(c*x))^2-2*b*d^2*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c-8/9*b*d*e*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3-16/75*b*e^2*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^5-4/9*b*d*e*x^2*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c-8/75*b*e^2*x^2*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c^3-2/25*b*e^2*x^4*(a+b*arccosh(c*x))*(c*x-1)^{(1/2)}*(c*x+1)^{(1/2)}/c$

### 3.526.2 Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 299, normalized size of antiderivative = 0.83

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx$$

$$= \frac{225a^2c^5x(15d^2 + 10dex^2 + 3e^2x^4) - 30ab\sqrt{-1 + cx}\sqrt{1 + cx}(24e^2 + 4c^2e(25d + 3ex^2) + c^4(225d^2 + 50dex^2 + 9e^2x^4)) + 2b^2c^5x(15d^2 + 10dexe^2 + 3e^2x^4) - 30ab\sqrt{-1 + cx}\sqrt{1 + cx}(24e^2 + 4c^2e(25d + 3ex^2) + c^4(225d^2 + 50dex^2 + 9e^2x^4)) + b\sqrt{-1 + cx}\sqrt{1 + cx}(24e^2 + 4c^2e(25d + 3ex^2) + c^4(225d^2 + 50dexe^2 + 9e^2x^4))\operatorname{ArcCosh}[cx] + 225b^2c^5x(15d^2 + 10dexe^2 + 3e^2x^4)\operatorname{ArcCosh}[cx]^2}{(3375c^5)}$$

input `Integrate[(d + e*x^2)^2*(a + b*ArcCosh[c*x])^2,x]`

output  $(225*a^2*c^5*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4) - 30*a*b*\sqrt{-1 + c*x}*\sqrt{1 + c*x}*(24*e^2 + 4*c^2*e*(25*d + 3*e*x^2) + c^4*(225*d^2 + 50*d*e*x^2 + 9*e^2*x^4)) + 2*b^2*c^5*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4) - 30*b*(-15*a*c^5*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4) + b*\sqrt{-1 + c*x}*\sqrt{1 + c*x}*(24*e^2 + 4*c^2*e*(25*d + 3*e*x^2) + c^4*(225*d^2 + 50*d*e*x^2 + 9*e^2*x^4)))*\operatorname{ArcCosh}[c*x] + 225*b^2*c^5*x*(15*d^2 + 10*d*e*x^2 + 3*e^2*x^4)*\operatorname{ArcCosh}[c*x]^2)/(3375*c^5)$

### 3.526.3 Rubi [A] (verified)

Time = 1.47 (sec) , antiderivative size = 359, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx$$

3.526.  $\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx$

$$\begin{aligned}
 & \int (d^2(a + \operatorname{barccosh}(cx))^2 + 2dex^2(a + \operatorname{barccosh}(cx))^2 + e^2x^4(a + \operatorname{barccosh}(cx))^2) dx \\
 & \quad \downarrow \text{6324} \\
 & \quad \downarrow \text{2009} \\
 & \frac{16be^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{75c^5} - \frac{8bde\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{9c^3} - \\
 & \quad \frac{8be^2x^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{75c^3} + d^2x(a + \operatorname{barccosh}(cx))^2 - \\
 & \quad \frac{2bd^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{c} + \frac{2}{3}dex^3(a + \operatorname{barccosh}(cx))^2 - \\
 & \quad \frac{4bdex^2\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{9c} + \frac{1}{5}e^2x^5(a + \operatorname{barccosh}(cx))^2 - \\
 & \quad \frac{2be^2x^4\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{25c} + \frac{16b^2e^2x}{75c^4} + \frac{8b^2dex}{9c^2} + \frac{8b^2e^2x^3}{225c^2} + 2b^2d^2x + \frac{4}{27}b^2dex^3 + \\
 & \quad \frac{2}{125}b^2e^2x^5
 \end{aligned}$$

input `Int[(d + e*x^2)^2*(a + b*ArcCosh[c*x])^2,x]`

output `2*b^2*d^2*x + (8*b^2*d*e*x)/(9*c^2) + (16*b^2*e^2*x)/(75*c^4) + (4*b^2*d*e*x^3)/27 + (8*b^2*e^2*x^3)/(225*c^2) + (2*b^2*e^2*x^5)/125 - (2*b*d^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/c - (8*b*d*e*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(9*c^3) - (16*b*e^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(75*c^5) - (4*b*d*e*x^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(9*c) - (8*b*e^2*x^2*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(75*c^3) - (2*b*e^2*x^4*sqrt[-1 + c*x]*sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/(25*c) + d^2*x*(a + b*ArcCosh[c*x])^2 + (2*d*e*x^3*(a + b*ArcCosh[c*x])^2)/3 + (e^2*x^5*(a + b*ArcCosh[c*x])^2)/5`

### 3.526.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

### 3.526.4 Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 402, normalized size of antiderivative = 1.12

method	result
derivativedivides	$\frac{a^2(d^2c^5x + \frac{2}{3}dc^5ex^3 + \frac{1}{5}e^2c^5x^5)}{c^4} + \frac{b^2(c^4d^2(\operatorname{arccosh}(cx))^2xc - 2\operatorname{arccosh}(cx)\sqrt{cx-1}\sqrt{cx+1} + 2cx) + 2c^2de(9\operatorname{arccosh}(cx)^2x^3c^3 - 6\sqrt{cx-1}\sqrt{cx+1})}{c^4}}$
default	$\frac{a^2(d^2c^5x + \frac{2}{3}dc^5ex^3 + \frac{1}{5}e^2c^5x^5)}{c^4} + \frac{b^2(c^4d^2(\operatorname{arccosh}(cx))^2xc - 2\operatorname{arccosh}(cx)\sqrt{cx-1}\sqrt{cx+1} + 2cx) + 2c^2de(9\operatorname{arccosh}(cx)^2x^3c^3 - 6\sqrt{cx-1}\sqrt{cx+1})}{c^4}}$
parts	$a^2\left(\frac{1}{5}e^2x^5 + \frac{2}{3}dex^3 + d^2x\right) + \frac{b^2(675\operatorname{arccosh}(cx)^2c^5x^5e^2 + 2250\operatorname{arccosh}(cx)^2c^5x^3de + 3375\operatorname{arccosh}(cx)^2c^5x^2d^2e + 2250\operatorname{arccosh}(cx)c^5x^5e^2 + 2250\operatorname{arccosh}(cx)c^5x^3de + 100c^2d^2e + 24e^2)}{c^4}$

input `int((e*x^2+d)^2*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output 
$$\frac{1}{c} \left( \frac{a^2}{c^4} (d^2c^5x + \frac{2}{3}dc^5ex^3 + \frac{1}{5}e^2c^5x^5) + \frac{b^2}{c^4} (c^4d^2(\operatorname{arccosh}(cx))^2xc - 2\operatorname{arccosh}(cx)(cx-1)^{1/2}(cx+1)^{1/2} + 2cx) + \frac{2c^2de(9\operatorname{arccosh}(cx)^2x^3c^3 - 6\sqrt{cx-1}\sqrt{cx+1})}{c^4} \right) + \frac{b^2}{c^4} (c^4d^2(\operatorname{arccosh}(cx))^2xc - 2\operatorname{arccosh}(cx)(cx-1)^{1/2}(cx+1)^{1/2} + 2cx) + \frac{2c^2de(9\operatorname{arccosh}(cx)^2x^3c^3 - 6\sqrt{cx-1}\sqrt{cx+1})}{c^4} + \frac{1}{1125}e^2(225\operatorname{arccosh}(cx)^2c^5x^5 - 90(cx-1)^{1/2}(cx+1)^{1/2}\operatorname{arccosh}(cx)c^4x^4 - 120(cx+1)^{1/2}\operatorname{arccosh}(cx)(cx-1)^{1/2}c^2x^2 + 18c^5x^5 - 240\operatorname{arccosh}(cx)(cx-1)^{1/2}(cx+1)^{1/2} + 40c^3x^3 + 240cx) + 2ab/c^4(\operatorname{arccosh}(cx)d^2c^5x + \frac{2}{3}\operatorname{arccosh}(cx)dc^5ex^3 + \frac{1}{5}\operatorname{arccosh}(cx)e^2c^5x^5 - \frac{1}{225}(cx-1)^{1/2}(cx+1)^{1/2}(9c^4e^2x^4 + 50c^4d^2e^2x^2 + 225c^4d^2 + 12c^2e^2x^2 + 100c^2d^2e + 24e^2))$$

### 3.526.5 Fracas [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 380, normalized size of antiderivative = 1.06

$$\int (d + ex^2)^2 (a + b\operatorname{arccosh}(cx))^2 dx = \frac{27(25a^2 + 2b^2)c^5e^2x^5 + 10(25(9a^2 + 2b^2)c^5de + 12b^2c^3e^2)x^3 + 225(3b^2c^5e^2x^5 + 10b^2c^5dex^3 + 15b^2c^5d^2e^2x^2 + 10b^2c^5d^2e^2x + 15b^2c^5d^2e^2)}{c^4}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output  $\frac{1}{3375} \cdot (27 \cdot (25a^2 + 2b^2) \cdot c^5 \cdot e^{2x^5} + 10 \cdot (25 \cdot (9a^2 + 2b^2) \cdot c^5 \cdot d \cdot e + 12b^2 \cdot c^3 \cdot e^2) \cdot x^3 + 225 \cdot (3b^2 \cdot c^5 \cdot e^{2x^5} + 10b^2 \cdot c^5 \cdot d \cdot e \cdot x^3 + 15b^2 \cdot c^5 \cdot d^2 \cdot x) \cdot \log(cx + \sqrt{c^2x^2 - 1})^2 + 15 \cdot (225 \cdot (a^2 + 2b^2) \cdot c^5 \cdot d^2 + 200b^2 \cdot c^3 \cdot d \cdot e + 48b^2 \cdot c \cdot e^2) \cdot x + 30 \cdot (45 \cdot a \cdot b \cdot c^5 \cdot e^{2x^5} + 150 \cdot a \cdot b \cdot c^5 \cdot d \cdot e \cdot x^3 + 225 \cdot a \cdot b \cdot c^5 \cdot d^2 \cdot x - (9b^2 \cdot c^4 \cdot e^{2x^4} + 225b^2 \cdot c^4 \cdot d^2 + 100b^2 \cdot c^2 \cdot d \cdot e + 24b^2 \cdot e^2 + 2 \cdot (25b^2 \cdot c^4 \cdot d \cdot e + 6b^2 \cdot c^2 \cdot e^2) \cdot x^2) \cdot \sqrt{c^2x^2 - 1}) \cdot \log(cx + \sqrt{c^2x^2 - 1}) - 30 \cdot (9 \cdot a \cdot b \cdot c^4 \cdot e^{2x^4} + 225 \cdot a \cdot b \cdot c^4 \cdot d^2 + 100 \cdot a \cdot b \cdot c^2 \cdot d \cdot e + 24 \cdot a \cdot b \cdot e^2 + 2 \cdot (25 \cdot a \cdot b \cdot c^4 \cdot d \cdot e + 6 \cdot a \cdot b \cdot c^2 \cdot e^2) \cdot x^2) \cdot \sqrt{c^2x^2 - 1}) / c^5$

### 3.526.6 Sympy [F]

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (d + ex^2)^2 dx$$

input `integrate((e*x**2+d)**2*(a+b*acosh(c*x))**2,x)`

output `Integral((a + b*acosh(c*x))**2*(d + e*x**2)**2, x)`

### 3.526.7 Maxima [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 429, normalized size of antiderivative = 1.19

$$\begin{aligned} & \int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx \\ &= \frac{1}{5} b^2 e^2 x^5 \operatorname{arcosh}(cx)^2 + \frac{1}{5} a^2 e^2 x^5 + \frac{2}{3} b^2 d e x^3 \operatorname{arcosh}(cx)^2 + \frac{2}{3} a^2 d e x^3 \\ &+ b^2 d^2 x \operatorname{arcosh}(cx)^2 + \frac{4}{9} \left( 3x^3 \operatorname{arcosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \right) abde \\ &- \frac{4}{27} \left( 3c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \operatorname{arcosh}(cx) - \frac{c^2x^3 + 6x}{c^2} \right) b^2 de \\ &+ \frac{2}{75} \left( 15x^5 \operatorname{arcosh}(cx) - \left( \frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6} \right) c \right) abe^2 \\ &- \frac{2}{1125} \left( 15 \left( \frac{3\sqrt{c^2x^2 - 1}x^4}{c^2} + \frac{4\sqrt{c^2x^2 - 1}x^2}{c^4} + \frac{8\sqrt{c^2x^2 - 1}}{c^6} \right) c \operatorname{arcosh}(cx) - \frac{9c^4x^5 + 20c^2x^3 + 120x}{c^4} \right) \\ &+ 2b^2d^2 \left( x - \frac{\sqrt{c^2x^2 - 1} \operatorname{arcosh}(cx)}{c} \right) + a^2d^2x + \frac{2(cx \operatorname{arcosh}(cx) - \sqrt{c^2x^2 - 1})abd^2}{c} \end{aligned}$$

---

3.526.  $\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx$



input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `1/5*b^2*e^2*x^5*arccosh(c*x)^2 + 1/5*a^2*e^2*x^5 + 2/3*b^2*d*e*x^3*arccosh(c*x)^2 + 2/3*a^2*d*e*x^3 + b^2*d^2*x*arccosh(c*x)^2 + 4/9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4))*a*b*d*e - 4/27*(3*c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/c^4)*arccosh(c*x) - (c^2*x^3 + 6*x)/c^2)*b^2*d*e + 2/75*(15*x^5*arccosh(c*x) - (3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6))*a*b*e^2 - 2/1125*(15*(3*sqrt(c^2*x^2 - 1)*x^4/c^2 + 4*sqrt(c^2*x^2 - 1)*x^2/c^4 + 8*sqrt(c^2*x^2 - 1)/c^6)*c*arccosh(c*x) - (9*c^4*x^5 + 20*c^2*x^3 + 120*x)/c^4)*b^2*e^2 + 2*b^2*d^2*(x - sqrt(c^2*x^2 - 1)*arccosh(c*x)/c) + a^2*d^2*x + 2*(c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))*a*b*d^2/c`

### 3.526.8 Giac [F(-2)]

Exception generated.

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: RuntimeError}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output Exception raised: RuntimeError >> an error occurred running a Giac command :INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index\_m & i,const ve cteur & l) Error: Bad Argument Value

### 3.526.9 Mupad [F(-1)]

Timed out.

$$\int (d + ex^2)^2 (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (ex^2 + d)^2 dx$$

input `int((a + b*acosh(c*x))^2*(d + e*x^2)^2,x)`

output `int((a + b*acosh(c*x))^2*(d + e*x^2)^2, x)`

### 3.527 $\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx$

3.527.1 Optimal result . . . . .	3881
3.527.2 Mathematica [A] (verified) . . . . .	3882
3.527.3 Rubi [A] (verified) . . . . .	3882
3.527.4 Maple [A] (verified) . . . . .	3883
3.527.5 Fracas [A] (verification not implemented) . . . . .	3884
3.527.6 Sympy [F] . . . . .	3884
3.527.7 Maxima [A] (verification not implemented) . . . . .	3885
3.527.8 Giac [F(-2)] . . . . .	3885
3.527.9 Mupad [F(-1)] . . . . .	3886

#### 3.527.1 Optimal result

Integrand size = 18, antiderivative size = 168

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx = 2b^2 dx + \frac{4b^2 ex}{9c^2} + \frac{2}{27} b^2 ex^3$$

$$- \frac{2bd\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx))}{c}$$

$$- \frac{4be\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx))}{9c^3}$$

$$- \frac{2bex^2\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx))}{9c}$$

$$+ dx(a + \operatorname{barccosh}(cx))^2 + \frac{1}{3} ex^3(a + \operatorname{barccosh}(cx))^2$$

```
output 2*b^2*d*x+4/9*b^2*e*x/c^2+2/27*b^2*e*x^3+d*x*(a+b*arccosh(c*x))^2+1/3*e*x^3*(a+b*arccosh(c*x))^2-2*b*d*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c-4/9*b*e*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c^3-2/9*b*e*x^2*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c
```

**3.527.2 Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.04

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx$$

$$= \frac{9a^2c^3x(3d + ex^2) - 6ab\sqrt{-1 + cx}\sqrt{1 + cx}(2e + c^2(9d + ex^2)) + 2b^2cx(6e + c^2(27d + ex^2)) - 6b(-3ac^3x + 27c^3)}{27c^3}$$

input `Integrate[(d + e*x^2)*(a + b*ArcCosh[c*x])^2,x]`output `(9*a^2*c^3*x*(3*d + e*x^2) - 6*a*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*e + c^2*(9*d + e*x^2)) + 2*b^2*c*x*(6*e + c^2*(27*d + e*x^2)) - 6*b*(-3*a*c^3*x*(3*d + e*x^2) + b*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*(2*e + c^2*(9*d + e*x^2)))*ArcCosh[c*x] + 9*b^2*c^3*x*(3*d + e*x^2)*ArcCosh[c*x]^2)/(27*c^3)`**3.527.3 Rubi [A] (verified)**Time = 0.79 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx$$

$$\downarrow \text{6324}$$

$$\int (d(a + \operatorname{barccosh}(cx))^2 + ex^2(a + \operatorname{barccosh}(cx))^2) dx$$

$$\downarrow \text{2009}$$

$$\frac{4be\sqrt{cx - 1}\sqrt{cx + 1}(a + \operatorname{barccosh}(cx))}{9c^3} + dx(a + \operatorname{barccosh}(cx))^2 - \frac{2bd\sqrt{cx - 1}\sqrt{cx + 1}(a + \operatorname{barccosh}(cx))}{c} + \frac{1}{3}ex^3(a + \operatorname{barccosh}(cx))^2 - \frac{2bex^2\sqrt{cx - 1}\sqrt{cx + 1}(a + \operatorname{barccosh}(cx))}{9c} + \frac{4b^2ex}{9c^2} + 2b^2dx + \frac{2}{27}b^2ex^3$$

input `Int[(d + e*x^2)*(a + b*ArcCosh[c*x])^2,x]`

---

 3.527.  $\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx$

```
output 2*b^2*d*x + (4*b^2*e*x)/(9*c^2) + (2*b^2*e*x^3)/27 - (2*b*d*Sqrt[-1 + c*x]
*Sqrt[1 + c*x]*(a + b*ArcCosh[c*x]))/c - (4*b*e*Sqrt[-1 + c*x]*Sqrt[1 + c*
x]*(a + b*ArcCosh[c*x]))/(9*c^3) - (2*b*e*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]
*(a + b*ArcCosh[c*x]))/(9*c) + d*x*(a + b*ArcCosh[c*x])^2 + (e*x^3*(a + b*
ArcCosh[c*x])^2)/3
```

### 3.527.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6324 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n_)*((d_) + (e_.)*(x_)^2)^(p_.),
x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x],
x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] &&
(p > 0 || IGtQ[n, 0])
```

### 3.527.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.21

method	result
parts	$a^2 \left( \frac{1}{3} x^3 e + dx \right) + \frac{b^2 \left( \frac{e \left( 9 \operatorname{arccosh}(cx)^2 x^3 c^3 - 6 \sqrt{cx+1} \operatorname{arccosh}(cx) \sqrt{cx-1} c^2 x^2 - 12 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} + 2c^3 x^3 + 12cx \right)}{27c^2} \right)}{c}$
derivativedivides	$\frac{a^2 \left( dc^3x + \frac{1}{3} e c^3 x^3 \right)}{c^2} + \frac{b^2 \left( dc^2 \left( \operatorname{arccosh}(cx)^2 xc - 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} + 2cx \right) + \frac{e \left( 9 \operatorname{arccosh}(cx)^2 x^3 c^3 - 6 \sqrt{cx+1} \operatorname{arccosh}(cx) \sqrt{cx-1} \right)}{c^2} \right)}{c^2}$
default	$\frac{a^2 \left( dc^3x + \frac{1}{3} e c^3 x^3 \right)}{c^2} + \frac{b^2 \left( dc^2 \left( \operatorname{arccosh}(cx)^2 xc - 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} + 2cx \right) + \frac{e \left( 9 \operatorname{arccosh}(cx)^2 x^3 c^3 - 6 \sqrt{cx+1} \operatorname{arccosh}(cx) \sqrt{cx-1} \right)}{c^2} \right)}{c^2}$

```
input int((e*x^2+d)*(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output a^2*(1/3*x^3*e+d*x)+b^2/c*(1/27*e*(9*arccosh(c*x)^2*x^3*c^3-6*(c*x+1)^(1/2)
)*arccosh(c*x)*(c*x-1)^(1/2)*c^2*x^2-12*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)
^(1/2)+2*c^3*x^3+12*c*x)/c^2+d*(arccosh(c*x)^2*x*c-2*arccosh(c*x)*(c*x-1)^(
1/2)*(c*x+1)^(1/2)+2*c*x))+2*a*b/c*(1/3*c*arccosh(c*x)*x^3*e+arccosh(c*x)
*d*c*x-1/9/c^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(c^2*e*x^2+9*c^2*d+2*e))
```

---

3.527.  $\int (d + ex^2) (a + b \operatorname{arccosh}(cx))^2 dx$

**3.527.5 Fracas [A] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.24

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx$$

$$= \frac{(9a^2 + 2b^2)c^3ex^3 + 9(b^2c^3ex^3 + 3b^2c^3dx) \log(cx + \sqrt{c^2x^2 - 1})^2 + 3(9(a^2 + 2b^2)c^3d + 4b^2ce)x + 6(3a^2c^3d + 2b^2ce)x^2}{c^3}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))^2,x, algorithm="fricas")`output `1/27*((9*a^2 + 2*b^2)*c^3*e*x^3 + 9*(b^2*c^3*e*x^3 + 3*b^2*c^3*d*x)*log(c*x + sqrt(c^2*x^2 - 1))^2 + 3*(9*(a^2 + 2*b^2)*c^3*d + 4*b^2*c^3*e)*x + 6*(3*a*b*c^3*e*x^3 + 9*a*b*c^3*d*x - (b^2*c^2*e*x^2 + 9*b^2*c^2*d + 2*b^2*e)*sqrt(c^2*x^2 - 1))*log(c*x + sqrt(c^2*x^2 - 1)) - 6*(a*b*c^2*e*x^2 + 9*a*b*c^2*d + 2*a*b*e)*sqrt(c^2*x^2 - 1))/c^3`**3.527.6 Sympy [F]**

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (d + ex^2) dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x))**2,x)`output `Integral((a + b*acosh(c*x))**2*(d + e*x**2), x)`

**3.527.7 Maxima [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.30

$$\begin{aligned}
& \int (d + ex^2) (a + \operatorname{arccosh}(cx))^2 dx \\
&= \frac{1}{3} b^2 ex^3 \operatorname{arccosh}(cx)^2 + \frac{1}{3} a^2 ex^3 + b^2 dx \operatorname{arccosh}(cx)^2 \\
&+ \frac{2}{9} \left( 3x^3 \operatorname{arccosh}(cx) - c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \right) abe \\
&- \frac{2}{27} \left( 3c \left( \frac{\sqrt{c^2x^2 - 1}x^2}{c^2} + \frac{2\sqrt{c^2x^2 - 1}}{c^4} \right) \operatorname{arccosh}(cx) - \frac{c^2x^3 + 6x}{c^2} \right) b^2e \\
&+ 2b^2d \left( x - \frac{\sqrt{c^2x^2 - 1} \operatorname{arccosh}(cx)}{c} \right) + a^2dx + \frac{2(cx \operatorname{arccosh}(cx) - \sqrt{c^2x^2 - 1})abd}{c}
\end{aligned}$$

```
input integrate((e*x^2+d)*(a+b*arccosh(c*x))^2,x, algorithm="maxima")
```

```
output 1/3*b^2*e*x^3*arccosh(c*x)^2 + 1/3*a^2*e*x^3 + b^2*d*x*arccosh(c*x)^2 + 2/
9*(3*x^3*arccosh(c*x) - c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)
/c^4))*a*b*e - 2/27*(3*c*(sqrt(c^2*x^2 - 1)*x^2/c^2 + 2*sqrt(c^2*x^2 - 1)/
c^4)*arccosh(c*x) - (c^2*x^3 + 6*x)/c^2)*b^2*e + 2*b^2*d*(x - sqrt(c^2*x^2
- 1)*arccosh(c*x)/c) + a^2*d*x + 2*(c*x*arccosh(c*x) - sqrt(c^2*x^2 - 1))
*a*b*d/c
```

**3.527.8 Giac [F(-2)]**

Exception generated.

$$\int (d + ex^2) (a + \operatorname{arccosh}(cx))^2 dx = \text{Exception raised: RuntimeError}$$

```
input integrate((e*x^2+d)*(a+b*arccosh(c*x))^2,x, algorithm="giac")
```

```
output Exception raised: RuntimeError >> an error occurred running a Giac command
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve
cteur & l) Error: Bad Argument Value
```

**3.527.9 Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 (ex^2 + d) dx$$

input `int((a + b*acosh(c*x))^2*(d + e*x^2),x)`output `int((a + b*acosh(c*x))^2*(d + e*x^2), x)`

### 3.528 $\int (a + \operatorname{barccosh}(cx))^2 dx$

3.528.1 Optimal result . . . . .	3887
3.528.2 Mathematica [A] (verified) . . . . .	3887
3.528.3 Rubi [A] (verified) . . . . .	3888
3.528.4 Maple [A] (verified) . . . . .	3889
3.528.5 Fricas [B] (verification not implemented) . . . . .	3889
3.528.6 Sympy [F] . . . . .	3890
3.528.7 Maxima [A] (verification not implemented) . . . . .	3890
3.528.8 Giac [B] (verification not implemented) . . . . .	3890
3.528.9 Mupad [F(-1)] . . . . .	3891

#### 3.528.1 Optimal result

Integrand size = 10, antiderivative size = 51

$$\int (a + \operatorname{barccosh}(cx))^2 dx = 2b^2x - \frac{2b\sqrt{-1 + cx}\sqrt{1 + cx}(a + \operatorname{barccosh}(cx))}{c} + x(a + \operatorname{barccosh}(cx))^2$$

output `2*b^2*x+x*(a+b*arccosh(c*x))^2-2*b*(a+b*arccosh(c*x))*(c*x-1)^(1/2)*(c*x+1)^(1/2)/c`

#### 3.528.2 Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.65

$$\int (a + \operatorname{barccosh}(cx))^2 dx = (a^2 + 2b^2) x - \frac{2ab\sqrt{-1 + cx}\sqrt{1 + cx}}{c} + \frac{2b(acx - b\sqrt{-1 + cx}\sqrt{1 + cx}) \operatorname{arccosh}(cx)}{c} + b^2x\operatorname{arccosh}(cx)^2$$

input `Integrate[(a + b*ArcCosh[c*x])^2,x]`

output `(a^2 + 2*b^2)*x - (2*a*b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/c + (2*b*(a*c*x - b*Sqrt[-1 + c*x]*Sqrt[1 + c*x])*ArcCosh[c*x])/c + b^2*x*ArcCosh[c*x]^2`



**3.528.3 Rubi [A] (verified)**

Time = 0.39 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.08, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$ , Rules used = {6294, 6330, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + \operatorname{barccosh}(cx))^2 dx$$

$$\downarrow 6294$$

$$x(a + \operatorname{barccosh}(cx))^2 - 2bc \int \frac{x(a + \operatorname{barccosh}(cx))}{\sqrt{cx-1}\sqrt{cx+1}} dx$$

$$\downarrow 6330$$

$$x(a + \operatorname{barccosh}(cx))^2 - 2bc \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{c^2} - \frac{b \int 1 dx}{c} \right)$$

$$\downarrow 24$$

$$x(a + \operatorname{barccosh}(cx))^2 - 2bc \left( \frac{\sqrt{cx-1}\sqrt{cx+1}(a + \operatorname{barccosh}(cx))}{c^2} - \frac{bx}{c} \right)$$

input `Int[(a + b*ArcCosh[c*x])^2,x]`

output `x*(a + b*ArcCosh[c*x])^2 - 2*b*c*(-((b*x)/c) + (Sqrt[-1 + c*x]*Sqrt[1 + c*x])*(a + b*ArcCosh[c*x]))/c^2)`

**3.528.3.1 Defintions of rubi rules used**

rule 24 `Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_., x_Symbol] :> Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]))], x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 6330 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p_) * ((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 + e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && E qQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]`

### 3.528.4 Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.53

method	result	size
derivativedivides	$\frac{cx a^2 + b^2 (\operatorname{arccosh}(cx)^2 xc - 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} + 2cx) + 2ab (cx \operatorname{arccosh}(cx) - \sqrt{cx-1} \sqrt{cx+1})}{c}$	78
default	$\frac{cx a^2 + b^2 (\operatorname{arccosh}(cx)^2 xc - 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} + 2cx) + 2ab (cx \operatorname{arccosh}(cx) - \sqrt{cx-1} \sqrt{cx+1})}{c}$	78
parts	$a^2 x + \frac{b^2 (\operatorname{arccosh}(cx)^2 xc - 2 \operatorname{arccosh}(cx) \sqrt{cx-1} \sqrt{cx+1} + 2cx)}{c} + \frac{2ab (cx \operatorname{arccosh}(cx) - \sqrt{cx-1} \sqrt{cx+1})}{c}$	79

input `int((a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

output `1/c*(c*x*a^2+b^2*(arccosh(c*x)^2*x*c-2*arccosh(c*x)*(c*x-1)^(1/2)*(c*x+1)^(1/2)+2*c*x)+2*a*b*(c*x*arccosh(c*x)-(c*x-1)^(1/2)*(c*x+1)^(1/2)))`

### 3.528.5 Fracas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 96 vs.  $2(47) = 94$ .

Time = 0.26 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.88

$$\int (a + b \operatorname{arccosh}(cx))^2 dx$$

$$= \frac{b^2 cx \log (cx + \sqrt{c^2 x^2 - 1})^2 + (a^2 + 2b^2) cx - 2 \sqrt{c^2 x^2 - 1} ab + 2 (abcx - \sqrt{c^2 x^2 - 1} b^2) \log (cx + \sqrt{c^2 x^2 - 1})}{c}$$

input `integrate((a+b*arccosh(c*x))^2,x, algorithm="fracas")`

output  $(b^2cx \log(cx + \sqrt{c^2x^2 - 1})^2 + (a^2 + 2b^2)cx - 2\sqrt{c^2x^2 - 1}ab + 2(abcx - \sqrt{c^2x^2 - 1}b^2)\log(cx + \sqrt{c^2x^2 - 1}))/c$

### 3.528.6 Sympy [F]

$$\int (a + b \operatorname{arccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 dx$$

input `integrate((a+b*acosh(c*x))**2,x)`

output `Integral((a + b*acosh(c*x))**2, x)`

### 3.528.7 Maxima [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.41

$$\int (a + b \operatorname{arccosh}(cx))^2 dx = b^2x \operatorname{arccosh}(cx)^2 + 2b^2 \left( x - \frac{\sqrt{c^2x^2 - 1} \operatorname{arccosh}(cx)}{c} \right) + a^2x + \frac{2(cx \operatorname{arccosh}(cx) - \sqrt{c^2x^2 - 1})ab}{c}$$

input `integrate((a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $b^2x \operatorname{arccosh}(cx)^2 + 2b^2(x - \sqrt{c^2x^2 - 1} \operatorname{arccosh}(cx)/c) + a^2x + 2(cx \operatorname{arccosh}(cx) - \sqrt{c^2x^2 - 1})ab/c$

### 3.528.8 Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs.  $2(47) = 94$ .

Time = 0.39 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.18

$$\int (a + \operatorname{barccosh}(cx))^2 dx$$

$$= 2 \left( x \log \left( cx + \sqrt{c^2 x^2 - 1} \right) - \frac{\sqrt{c^2 x^2 - 1}}{c} \right) ab$$

$$+ \left( x \log \left( cx + \sqrt{c^2 x^2 - 1} \right)^2 + 2c \left( \frac{x}{c} - \frac{\sqrt{c^2 x^2 - 1} \log \left( cx + \sqrt{c^2 x^2 - 1} \right)}{c^2} \right) \right) b^2 + a^2 x$$

input `integrate((a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `2*(x*log(c*x + sqrt(c^2*x^2 - 1)) - sqrt(c^2*x^2 - 1)/c)*a*b + (x*log(c*x + sqrt(c^2*x^2 - 1))^2 + 2*c*(x/c - sqrt(c^2*x^2 - 1)*log(c*x + sqrt(c^2*x^2 - 1))/c^2))*b^2 + a^2*x`

### 3.528.9 Mupad [F(-1)]

Timed out.

$$\int (a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 dx$$

input `int((a + b*acosh(c*x))^2,x)`

output `int((a + b*acosh(c*x))^2, x)`

$$3.529 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^2}{d+ex^2} dx$$

3.529.1 Optimal result . . . . .	3893
3.529.2 Mathematica [A] (verified) . . . . .	3894
3.529.3 Rubi [A] (verified) . . . . .	3895
3.529.4 Maple [F] . . . . .	3897
3.529.5 Fracas [F] . . . . .	3898
3.529.6 Sympy [F] . . . . .	3898
3.529.7 Maxima [F(-2)] . . . . .	3898
3.529.8 Giac [F] . . . . .	3899
3.529.9 Mupad [F(-1)] . . . . .	3899

## 3.529.1 Optimal result

Integrand size = 20, antiderivative size = 763

$$\begin{aligned}
\int \frac{(a + \operatorname{barccosh}(cx))^2}{d + ex^2} dx = & \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{2\sqrt{-d}\sqrt{e}} \\
& - \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{2\sqrt{-d}\sqrt{e}} \\
& + \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(1 - \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{2\sqrt{-d}\sqrt{e}} \\
& - \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(1 + \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{2\sqrt{-d}\sqrt{e}} \\
& - \frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}} \\
& + \frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}} \\
& - \frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}} \\
& + \frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}} \\
& + \frac{b^2 \operatorname{PolyLog}\left(3, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}} \\
& - \frac{b^2 \operatorname{PolyLog}\left(3, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} - \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}} \\
& + \frac{b^2 \operatorname{PolyLog}\left(3, -\frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}} \\
& - \frac{b^2 \operatorname{PolyLog}\left(3, \frac{\sqrt{ee} \operatorname{arccosh}(cx)}{c\sqrt{-d} + \sqrt{-c^2 d - e}}\right)}{\sqrt{-d}\sqrt{e}}
\end{aligned}$$

output

```

1/2*(a+b*arccosh(c*x))^2*ln(1-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c
*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(1/2)/e^(1/2)-1/2*(a+b*arccosh(c*x))^2
*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(
1/2)))/(-d)^(1/2)/e^(1/2)+1/2*(a+b*arccosh(c*x))^2*ln(1-(c*x+(c*x-1)^(1/2)
*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(1/2)/e^(1/2
)-1/2*(a+b*arccosh(c*x))^2*ln(1+(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/
(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(1/2)/e^(1/2)-b*(a+b*arccosh(c*x))*p
olylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-
e)^(1/2)))/(-d)^(1/2)/e^(1/2)+b*(a+b*arccosh(c*x))*polylog(2,(c*x+(c*x-1)^(
1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(1/2)/e
^(1/2)-b*(a+b*arccosh(c*x))*polylog(2,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e
^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(1/2)/e^(1/2)+b*(a+b*arccosh(
c*x))*polylog(2,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-
c^2*d-e)^(1/2)))/(-d)^(1/2)/e^(1/2)+b^2*polylog(3,-(c*x+(c*x-1)^(1/2)*(c*x
+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d-e)^(1/2)))/(-d)^(1/2)/e^(1/2)-b^2
*polylog(3,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)-(-c^2*d
-e)^(1/2)))/(-d)^(1/2)/e^(1/2)+b^2*polylog(3,-(c*x+(c*x-1)^(1/2)*(c*x+1)^(
1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(1/2)))/(-d)^(1/2)/e^(1/2)-b^2*pol
ylog(3,(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))*e^(1/2)/(c*(-d)^(1/2)+(-c^2*d-e)^(
1/2)))/(-d)^(1/2)/e^(1/2)

```

### 3.529.2 Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 623, normalized size of antiderivative = 0.82

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{d + ex^2} dx$$

$$= \frac{- (a + b \operatorname{arccosh}(cx))^2 \log \left( 1 + \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c \sqrt{-d - \sqrt{-c^2 d - e}}} \right) + (a + b \operatorname{arccosh}(cx))^2 \log \left( 1 + \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{-c \sqrt{-d + \sqrt{-c^2 d - e}}} \right) + (a + b \operatorname{arccosh}(cx))^2 \log \left( 1 + \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{c \sqrt{-d - \sqrt{-c^2 d - e}}} \right) + (a + b \operatorname{arccosh}(cx))^2 \log \left( 1 + \frac{\sqrt{e} e^{\operatorname{arccosh}(cx)}}{-c \sqrt{-d + \sqrt{-c^2 d - e}}} \right)}{2}$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(d + e*x^2),x]`

output  $(-((a + b*\text{ArcCosh}[c*x])^2*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])]) + (a + b*\text{ArcCosh}[c*x])^2*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(-(c*\text{Sqrt}[-d]) + \text{Sqrt}[-(c^2*d) - e])]) + (a + b*\text{ArcCosh}[c*x])^2*\text{Log}[1 - (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])]) - (a + b*\text{ArcCosh}[c*x])^2*\text{Log}[1 + (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])]) + 2*b*(a + b*\text{ArcCosh}[c*x])*PolyLog[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])]) - 2*b*(a + b*\text{ArcCosh}[c*x])*PolyLog[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(-(c*\text{Sqrt}[-d]) + \text{Sqrt}[-(c^2*d) - e])]) - 2*b*(a + b*\text{ArcCosh}[c*x])*PolyLog[2, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e]))] + 2*b*(a + b*\text{ArcCosh}[c*x])*PolyLog[2, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])]) - 2*b^2*PolyLog[3, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] - \text{Sqrt}[-(c^2*d) - e])]) + 2*b^2*PolyLog[3, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(-(c*\text{Sqrt}[-d]) + \text{Sqrt}[-(c^2*d) - e])]) + 2*b^2*PolyLog[3, -((\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e]))] - 2*b^2*PolyLog[3, (\text{Sqrt}[e]*E^{\text{ArcCosh}[c*x]})/(c*\text{Sqrt}[-d] + \text{Sqrt}[-(c^2*d) - e])])/(2*\text{Sqrt}[-d]*\text{Sqrt}[e])$

### 3.529.3 Rubi [A] (verified)

Time = 1.76 (sec) , antiderivative size = 763, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \text{barccosh}(cx))^2}{d + ex^2} dx$$

↓ 6324

$$\int \left( \frac{\sqrt{-d}(a + \text{barccosh}(cx))^2}{2d(\sqrt{-d} - \sqrt{ex})} + \frac{\sqrt{-d}(a + \text{barccosh}(cx))^2}{2d(\sqrt{-d} + \sqrt{ex})} \right) dx$$

↓ 2009



$$\begin{aligned}
& -\frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}} + \\
& \frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}} - \\
& \frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}} + \\
& \frac{b(a + \operatorname{barccosh}(cx)) \operatorname{PolyLog}\left(2, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}} + \\
& \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{c^2(-d)-e}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{c^2(-d)-e}} + 1\right)}{2\sqrt{-d}\sqrt{e}} + \\
& \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(1 - \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d)-e+c\sqrt{-d}}}\right)}{2\sqrt{-d}\sqrt{e}} - \frac{(a + \operatorname{barccosh}(cx))^2 \log\left(\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{c^2(-d)-e+c\sqrt{-d}}} + 1\right)}{2\sqrt{-d}\sqrt{e}} + \\
& \frac{b^2 \operatorname{PolyLog}\left(3, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}} - \frac{b^2 \operatorname{PolyLog}\left(3, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{c\sqrt{-d}-\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}} + \\
& \frac{b^2 \operatorname{PolyLog}\left(3, -\frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}} - \frac{b^2 \operatorname{PolyLog}\left(3, \frac{\sqrt{e}e^{\operatorname{arccosh}(cx)}}{\sqrt{-dc}+\sqrt{-dc^2-e}}\right)}{\sqrt{-d}\sqrt{e}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])^2/(d + e*x^2), x]`

```
output ((a + b*ArcCosh[c*x])^2*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) - ((a + b*ArcCosh[c*x])^2*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) + ((a + b*ArcCosh[c*x])^2*Log[1 - (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) - ((a + b*ArcCosh[c*x])^2*Log[1 + (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(2*Sqrt[-d]*Sqrt[e]) - (b*(a + b*ArcCosh[c*x])*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])]/(Sqrt[-d]*Sqrt[e]) + (b*(a + b*ArcCosh[c*x])*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(Sqrt[-d]*Sqrt[e]) - (b*(a + b*ArcCosh[c*x])*PolyLog[2, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])]/(Sqrt[-d]*Sqrt[e]) + (b*(a + b*ArcCosh[c*x])*PolyLog[2, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(Sqrt[-d]*Sqrt[e]) + (b^2*PolyLog[3, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])]/(Sqrt[-d]*Sqrt[e]) - (b^2*PolyLog[3, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] - Sqrt[-(c^2*d) - e])])/(Sqrt[-d]*Sqrt[e]) + (b^2*PolyLog[3, -((Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])]/(Sqrt[-d]*Sqrt[e]) - (b^2*PolyLog[3, (Sqrt[e]*E^ArcCosh[c*x])/(c*Sqrt[-d] + Sqrt[-(c^2*d) - e])])/(Sqrt[-d]*Sqrt[e]))
```

### 3.529.3.1 Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 6324 Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])
```

### 3.529.4 Maple [F]

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{ex^2 + d} dx$$

```
input int((a+b*arccosh(c*x))^2/(e*x^2+d), x)
```

```
output int((a+b*arccosh(c*x))^2/(e*x^2+d), x)
```

**3.529.5 Fricas [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{d + ex^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{ex^2 + d} dx$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d),x, algorithm="fricas")`

output `integral((b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/(e*x^2 + d), x)`

**3.529.6 Sympy [F]**

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{d + ex^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{d + ex^2} dx$$

input `integrate((a+b*acosh(c*x))**2/(e*x**2+d),x)`

output `Integral((a + b*acosh(c*x))**2/(d + e*x**2), x)`

**3.529.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.529.8 Giac [F]**

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{d + ex^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{ex^2 + d} dx$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^2/(e*x^2 + d), x)`

**3.529.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{d + ex^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{ex^2 + d} dx$$

input `int((a + b*acosh(c*x))^2/(d + e*x^2),x)`

output `int((a + b*acosh(c*x))^2/(d + e*x^2), x)`

### 3.530 $\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx$

3.530.1 Optimal result . . . . .	3900
3.530.2 Mathematica [N/A] . . . . .	3900
3.530.3 Rubi [N/A] . . . . .	3901
3.530.4 Maple [N/A] (verified) . . . . .	3901
3.530.5 Fricas [N/A] . . . . .	3902
3.530.6 Sympy [N/A] . . . . .	3902
3.530.7 Maxima [F(-2)] . . . . .	3902
3.530.8 Giac [N/A] . . . . .	3903
3.530.9 Mupad [N/A] . . . . .	3903

#### 3.530.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx = \operatorname{Int}\left(\sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^2*(e*x^2+d)^(1/2),x)`

#### 3.530.2 Mathematica [N/A]

Not integrable

Time = 15.37 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx = \int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx$$

input `Integrate[Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `Integrate[Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])^2, x]`

**3.530.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \sqrt{d + ex^2}(a + \operatorname{arccosh}(cx))^2 dx$$

↓ 6325

$$\int \sqrt{d + ex^2}(a + \operatorname{arccosh}(cx))^2 dx$$

input `Int[Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

**3.530.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^n_.*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.530.4 Maple [N/A] (verified)**

Not integrable

Time = 0.67 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int (a + b \operatorname{arccosh}(cx))^2 \sqrt{ex^2 + d} dx$$

input `int((a+b*arccosh(c*x))^2*(e*x^2+d)^(1/2),x)`

output `int((a+b*arccosh(c*x))^2*(e*x^2+d)^(1/2),x)`

**3.530.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx = \int \sqrt{ex^2 + d}(b \operatorname{arcosh}(cx) + a)^2 dx$$

input `integrate((a+b*arccosh(c*x))^2*(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral((b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)*sqrt(e*x^2 + d), x)`

**3.530.6 Sympy [N/A]**

Not integrable

Time = 1.64 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 \sqrt{d + ex^2} dx$$

input `integrate((a+b*acosh(c*x))**2*(e*x**2+d)**(1/2),x)`

output `Integral((a + b*acosh(c*x))**2*sqrt(d + e*x**2), x)`

**3.530.7 Maxima [F(-2)]**

Exception generated.

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))^2*(e*x^2+d)^(1/2),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.530.8 Giac [N/A]**

Not integrable

Time = 0.40 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx = \int \sqrt{ex^2 + d}(b \operatorname{arcosh}(cx) + a)^2 dx$$

input `integrate((a+b*arccosh(c*x))^2*(e*x^2+d)^(1/2),x, algorithm="giac")`output `integrate(sqrt(e*x^2 + d)*(b*arccosh(c*x) + a)^2, x)`**3.530.9 Mupad [N/A]**

Not integrable

Time = 3.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2 dx = \int (a + b \operatorname{acosh}(cx))^2 \sqrt{ex^2 + d} dx$$

input `int((a + b*acosh(c*x))^2*(d + e*x^2)^(1/2),x)`output `int((a + b*acosh(c*x))^2*(d + e*x^2)^(1/2), x)`



### 3.531 $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{\sqrt{d+ex^2}} dx$

3.531.1 Optimal result . . . . .	3904
3.531.2 Mathematica [N/A] . . . . .	3904
3.531.3 Rubi [N/A] . . . . .	3905
3.531.4 Maple [N/A] (verified) . . . . .	3905
3.531.5 Fricas [N/A] . . . . .	3906
3.531.6 Sympy [N/A] . . . . .	3906
3.531.7 Maxima [F(-2)] . . . . .	3906
3.531.8 Giac [N/A] . . . . .	3907
3.531.9 Mupad [N/A] . . . . .	3907

#### 3.531.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x)`

#### 3.531.2 Mathematica [N/A]

Not integrable

Time = 8.23 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^2/Sqrt[d + e*x^2],x]`

output `Integrate[(a + b*ArcCosh[c*x])^2/Sqrt[d + e*x^2], x]`

**3.531.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx$$

↓ 6325

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^2/Sqrt[d + e*x^2],x]`

output `$Aborted`

**3.531.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.531.4 Maple [N/A] (verified)**

Not integrable

Time = 0.78 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{ex^2 + d}} dx$$

input `int((a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x)`

output `int((a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x)`

**3.531.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.55

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{\sqrt{ex^2 + d}} dx$$

```
input integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x, algorithm="fricas")
```

```
output integral((b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)/sqrt(e*x^2 + d),
x)
```

**3.531.6 Sympy [N/A]**

Not integrable

Time = 1.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{\sqrt{d + ex^2}} dx$$

```
input integrate((a+b*acosh(c*x))**2/(e*x**2+d)**(1/2),x)
```

```
output Integral((a + b*acosh(c*x))**2/sqrt(d + e*x**2), x)
```

**3.531.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{\sqrt{d + ex^2}} dx = \text{Exception raised: ValueError}$$

```
input integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e>0)', see `assume?` for more de
tails)Is e
```

**3.531.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d + ex^2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{\sqrt{ex^2 + d}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^2/sqrt(e*x^2 + d), x)`**3.531.9 Mupad [N/A]**

Not integrable

Time = 3.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{\sqrt{d + ex^2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{\sqrt{ex^2 + d}} dx$$

input `int((a + b*acosh(c*x))^2/(d + e*x^2)^(1/2),x)`output `int((a + b*acosh(c*x))^2/(d + e*x^2)^(1/2), x)`

$$3.532 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d+ex^2)^{3/2}} dx$$

3.532.1 Optimal result	3908
3.532.2 Mathematica [N/A]	3908
3.532.3 Rubi [N/A]	3909
3.532.4 Maple [N/A] (verified)	3909
3.532.5 Fracas [N/A]	3910
3.532.6 Sympy [N/A]	3910
3.532.7 Maxima [F(-2)]	3910
3.532.8 Giac [N/A]	3911
3.532.9 Mupad [N/A]	3911

### 3.532.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^2/(e*x^2+d)^(3/2),x)`

### 3.532.2 Mathematica [N/A]

Not integrable

Time = 13.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(d + e*x^2)^(3/2),x]`

output `Integrate[(a + b*ArcCosh[c*x])^2/(d + e*x^2)^(3/2), x]`

---

3.532.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d+ex^2)^{3/2}} dx$

**3.532.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx$$

↓ 6325

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^2/(d + e*x^2)^(3/2),x]`

output `$Aborted`

**3.532.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.532.4 Maple [N/A] (verified)**

Not integrable

Time = 1.62 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(ex^2 + d)^{\frac{3}{2}}} dx$$

input `int((a+b*arccosh(c*x))^2/(e*x^2+d)^(3/2),x)`

output `int((a+b*arccosh(c*x))^2/(e*x^2+d)^(3/2),x)`

---

3.532.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d+ex^2)^{3/2}} dx$

**3.532.5 Fricas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.45

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(ex^2 + d)^{3/2}} dx$$

```
input integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(3/2),x, algorithm="fricas")
```

```
output integral((b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)*sqrt(e*x^2 + d)/(
e^2*x^4 + 2*d*e*x^2 + d^2), x)
```

**3.532.6 Sympy [N/A]**

Not integrable

Time = 7.62 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(d + ex^2)^{3/2}} dx$$

```
input integrate((a+b*acosh(c*x))**2/(e*x**2+d)**(3/2),x)
```

```
output Integral((a + b*acosh(c*x))**2/(d + e*x**2)**(3/2), x)
```

**3.532.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d + ex^2)^{3/2}} dx = \text{Exception raised: ValueError}$$

```
input integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(3/2),x, algorithm="maxima")
```

```
output Exception raised: ValueError >> Computation failed since Maxima requested
additional constraints; using the 'assume' command before evaluation *may*
help (example of legal syntax is 'assume(e+c^2*d>0)', see `assume?` for m
ore detail
```

---

3.532.  $\int \frac{(a+b \operatorname{arccosh}(cx))^2}{(d+ex^2)^{3/2}} dx$

**3.532.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d + ex^2)^{3/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(ex^2 + d)^{3/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(3/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^2/(e*x^2 + d)^(3/2), x)`**3.532.9 Mupad [N/A]**

Not integrable

Time = 3.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d + ex^2)^{3/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(ex^2 + d)^{3/2}} dx$$

input `int((a + b*acosh(c*x))^2/(d + e*x^2)^(3/2),x)`output `int((a + b*acosh(c*x))^2/(d + e*x^2)^(3/2), x)`



$$3.533 \quad \int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d+ex^2)^{5/2}} dx$$

3.533.1 Optimal result	3912
3.533.2 Mathematica [N/A]	3912
3.533.3 Rubi [N/A]	3913
3.533.4 Maple [N/A] (verified)	3913
3.533.5 Fricas [N/A]	3914
3.533.6 Sympy [N/A]	3914
3.533.7 Maxima [N/A]	3914
3.533.8 Giac [N/A]	3915
3.533.9 Mupad [N/A]	3915

### 3.533.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^2/(e*x^2+d)^(5/2),x)`

### 3.533.2 Mathematica [N/A]

Not integrable

Time = 26.79 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx = \int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^2/(d + e*x^2)^(5/2),x]`

output `Integrate[(a + b*ArcCosh[c*x])^2/(d + e*x^2)^(5/2), x]`

---

3.533.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d+ex^2)^{5/2}} dx$

**3.533.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx$$

↓ 6325

$$\int \frac{(a + \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx$$

input `Int[(a + b*ArcCosh[c*x])^2/(d + e*x^2)^(5/2),x]`

output `$Aborted`

**3.533.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.533.4 Maple [N/A] (verified)**

Not integrable

Time = 1.66 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(ex^2 + d)^{5/2}} dx$$

input `int((a+b*arccosh(c*x))^2/(e*x^2+d)^(5/2),x)`

output `int((a+b*arccosh(c*x))^2/(e*x^2+d)^(5/2),x)`

---

3.533.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d+ex^2)^{5/2}} dx$

**3.533.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 65, normalized size of antiderivative = 2.95

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(ex^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(5/2),x, algorithm="fricas")`

output `integral((b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2)*sqrt(e*x^2 + d)/(e^3*x^6 + 3*d*e^2*x^4 + 3*d^2*e*x^2 + d^3), x)`

**3.533.6 Sympy [N/A]**

Not integrable

Time = 167.63 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(d + ex^2)^{5/2}} dx$$

input `integrate((a+b*acosh(c*x))**2/(e*x**2+d)**(5/2),x)`

output `Integral((a + b*acosh(c*x))**2/(d + e*x**2)**(5/2), x)`

**3.533.7 Maxima [N/A]**

Not integrable

Time = 0.72 (sec) , antiderivative size = 107, normalized size of antiderivative = 4.86

$$\int \frac{(a + b \operatorname{arccosh}(cx))^2}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^2}{(ex^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(5/2),x, algorithm="maxima")`

output `1/3*a^2*(2*x/(sqrt(e*x^2 + d)*d^2) + x/((e*x^2 + d)^(3/2)*d)) + integrate(b^2*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(e*x^2 + d)^(5/2) + 2*a*b*log(c*x + sqrt(c*x + 1))*sqrt(c*x - 1)/(e*x^2 + d)^(5/2), x)`

---

3.533.  $\int \frac{(a+b\operatorname{arccosh}(cx))^2}{(d+ex^2)^{5/2}} dx$

**3.533.8 Giac [N/A]**

Not integrable

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d + ex^2)^{5/2}} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^2}{(ex^2 + d)^{5/2}} dx$$

input `integrate((a+b*arccosh(c*x))^2/(e*x^2+d)^(5/2),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^2/(e*x^2 + d)^(5/2), x)`**3.533.9 Mupad [N/A]**

Not integrable

Time = 3.36 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^2}{(d + ex^2)^{5/2}} dx = \int \frac{(a + b \operatorname{acosh}(cx))^2}{(ex^2 + d)^{5/2}} dx$$

input `int((a + b*acosh(c*x))^2/(d + e*x^2)^(5/2),x)`output `int((a + b*acosh(c*x))^2/(d + e*x^2)^(5/2), x)`

$$3.534 \quad \int \frac{(d+ex^2)^2}{a+b\operatorname{arccosh}(cx)} dx$$

3.534.1 Optimal result . . . . .	3917
3.534.2 Mathematica [A] (verified) . . . . .	3918
3.534.3 Rubi [A] (verified) . . . . .	3918
3.534.4 Maple [A] (verified) . . . . .	3920
3.534.5 Fricas [F] . . . . .	3921
3.534.6 Sympy [F] . . . . .	3921
3.534.7 Maxima [F] . . . . .	3921
3.534.8 Giac [F] . . . . .	3922
3.534.9 Mupad [F(-1)] . . . . .	3922

**3.534.1 Optimal result**

Integrand size = 20, antiderivative size = 388

$$\begin{aligned}
\int \frac{(d+ex^2)^2}{a+b\operatorname{arccosh}(cx)} dx = & -\frac{d^2 \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right)}{bc} \\
& -\frac{de \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right)}{2bc^3} \\
& -\frac{e^2 \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right)}{8bc^5} \\
& -\frac{de \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{3a}{b}\right)}{2bc^3} \\
& -\frac{3e^2 \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{3a}{b}\right)}{16bc^5} \\
& -\frac{e^2 \operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{5a}{b}\right)}{16bc^5} \\
& +\frac{d^2 \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{bc} \\
& +\frac{de \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2bc^3} \\
& +\frac{e^2 \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8bc^5} \\
& +\frac{de \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{2bc^3} \\
& +\frac{3e^2 \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^5} \\
& +\frac{e^2 \cosh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16bc^5}
\end{aligned}$$

output  $d^2 \cosh(a/b) \operatorname{Shi}((a+b \operatorname{arccosh}(cx))/b) / b/c + 1/2 d e \cosh(a/b) \operatorname{Shi}((a+b \operatorname{arccosh}(cx))/b) / b/c^3 + 1/8 e^2 \cosh(a/b) \operatorname{Shi}((a+b \operatorname{arccosh}(cx))/b) / b/c^5 + 1/2 d e \cosh(3a/b) \operatorname{Shi}(3(a+b \operatorname{arccosh}(cx))/b) / b/c^3 + 3/16 e^2 \cosh(3a/b) \operatorname{Shi}(3(a+b \operatorname{arccosh}(cx))/b) / b/c^5 + 1/16 e^2 \cosh(5a/b) \operatorname{Shi}(5(a+b \operatorname{arccosh}(cx))/b) / b/c^5 - d^2 \operatorname{Chi}((a+b \operatorname{arccosh}(cx))/b) \sinh(a/b) / b/c - 1/2 d e \operatorname{Chi}((a+b \operatorname{arccosh}(cx))/b) \sinh(a/b) / b/c^3 - 1/8 e^2 \operatorname{Chi}((a+b \operatorname{arccosh}(cx))/b) \sinh(a/b) / b/c^5 - 1/2 d e \operatorname{Chi}(3(a+b \operatorname{arccosh}(cx))/b) \sinh(3a/b) / b/c^3 - 3/16 e^2 \operatorname{Chi}(3(a+b \operatorname{arccosh}(cx))/b) \sinh(3a/b) / b/c^5 - 1/16 e^2 \operatorname{Chi}(5(a+b \operatorname{arccosh}(cx))/b) \sinh(5a/b) / b/c^5$

### 3.534.2 Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.65

$$\int \frac{(d + ex^2)^2}{a + b \operatorname{arccosh}(cx)} dx$$

$$= \frac{-2(8c^4 d^2 + 4c^2 d e + e^2) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \sinh\left(\frac{a}{b}\right) - e(8c^2 d + 3e) \operatorname{Chi}\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \sinh\left(\frac{3a}{b}\right)}{16 b^2 c^5}$$

input `Integrate[(d + e*x^2)^2/(a + b*ArcCosh[c*x]),x]`

output  $(-2*(8*c^4*d^2 + 4*c^2*d*e + e^2)*\operatorname{CoshIntegral}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Sinh}[a/b] - e*(8*c^2*d + 3*e)*\operatorname{CoshIntegral}[3*(a/b + \operatorname{ArcCosh}[c*x])]*\operatorname{Sinh}[(3*a)/b] - e^2*\operatorname{CoshIntegral}[5*(a/b + \operatorname{ArcCosh}[c*x])]*\operatorname{Sinh}[(5*a)/b] + 16*c^4*d^2*\operatorname{Cosh}[a/b]*\operatorname{SinhIntegral}[a/b + \operatorname{ArcCosh}[c*x]] + 8*c^2*d*e*\operatorname{Cosh}[a/b]*\operatorname{SinhIntegral}[a/b + \operatorname{ArcCosh}[c*x]] + 2*e^2*\operatorname{Cosh}[a/b]*\operatorname{SinhIntegral}[a/b + \operatorname{ArcCosh}[c*x]] + 8*c^2*d*e*\operatorname{Cosh}[(3*a)/b]*\operatorname{SinhIntegral}[3*(a/b + \operatorname{ArcCosh}[c*x])] + 3*e^2*\operatorname{Cosh}[(3*a)/b]*\operatorname{SinhIntegral}[3*(a/b + \operatorname{ArcCosh}[c*x])] + e^2*\operatorname{Cosh}[(5*a)/b]*\operatorname{SinhIntegral}[5*(a/b + \operatorname{ArcCosh}[c*x])])/(16*b*c^5)$

### 3.534.3 Rubi [A] (verified)

Time = 1.00 (sec) , antiderivative size = 388, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

---

3.534.  $\int \frac{(d+ex^2)^2}{a+b \operatorname{arccosh}(cx)} dx$

$$\begin{aligned}
 & \int \frac{(d + ex^2)^2}{a + \operatorname{barccosh}(cx)} dx \\
 & \quad \downarrow \text{6324} \\
 & \int \left( \frac{d^2}{a + \operatorname{barccosh}(cx)} + \frac{2dex^2}{a + \operatorname{barccosh}(cx)} + \frac{e^2x^4}{a + \operatorname{barccosh}(cx)} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{e^2 \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{8bc^5} - \frac{3e^2 \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{16bc^5} - \\
 & \frac{e^2 \sinh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{16bc^5} + \frac{e^2 \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{8bc^5} + \\
 & \frac{3e^2 \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{16bc^5} + \frac{e^2 \cosh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a + \operatorname{barccosh}(cx))}{b}\right)}{16bc^5} - \\
 & \frac{de \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{2bc^3} - \frac{de \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{2bc^3} + \\
 & \frac{de \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{2bc^3} + \frac{de \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a + \operatorname{barccosh}(cx))}{b}\right)}{2bc^3} - \\
 & \frac{d^2 \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{bc} + \frac{d^2 \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a + \operatorname{barccosh}(cx)}{b}\right)}{bc}
 \end{aligned}$$

input `Int[(d + e*x^2)^2/(a + b*ArcCosh[c*x]),x]`

output `-(d^2*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b])/(b*c) - (d*e*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b])/(2*b*c^3) - (e^2*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b])/(8*b*c^5) - (d*e*CoshIntegral[(3*(a + b*ArcCosh[c*x]))/b]*Sinh[(3*a)/b])/(2*b*c^3) - (3*e^2*CoshIntegral[(3*(a + b*ArcCosh[c*x]))/b]*Sinh[(3*a)/b])/(16*b*c^5) - (e^2*CoshIntegral[(5*(a + b*ArcCosh[c*x]))/b]*Sinh[(5*a)/b])/(16*b*c^5) + (d^2*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(b*c) + (d*e*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(2*b*c^3) + (e^2*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(8*b*c^5) + (d*e*Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x]))/b])/(2*b*c^3) + (3*e^2*Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x]))/b])/(16*b*c^5) + (e^2*Cosh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x]))/b])/(16*b*c^5)`



3.534.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

3.534.4 Maple [A] (verified)

Time = 2.65 (sec) , antiderivative size = 380, normalized size of antiderivative = 0.98

method	result
derivativedivides	$\frac{e^2 e^{\frac{5a}{b}} \operatorname{Ei}_1\left(5 \operatorname{arccosh}(cx) + \frac{5a}{b}\right)}{32c^4 b} - \frac{e^2 e^{-\frac{5a}{b}} \operatorname{Ei}_1\left(-5 \operatorname{arccosh}(cx) - \frac{5a}{b}\right)}{32c^4 b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right) d^2}{2b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right) de}{4c^2 b} + \dots$
default	$\frac{e^2 e^{\frac{5a}{b}} \operatorname{Ei}_1\left(5 \operatorname{arccosh}(cx) + \frac{5a}{b}\right)}{32c^4 b} - \frac{e^2 e^{-\frac{5a}{b}} \operatorname{Ei}_1\left(-5 \operatorname{arccosh}(cx) - \frac{5a}{b}\right)}{32c^4 b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right) d^2}{2b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right) de}{4c^2 b} + \dots$

input `int((e*x^2+d)^2/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `1/c*(1/32/c^4*e^2/b*exp(5*a/b)*Ei(1,5*arccosh(c*x)+5*a/b)-1/32/c^4*e^2/b*exp(-5*a/b)*Ei(1,-5*arccosh(c*x)-5*a/b)+1/2/b*exp(a/b)*Ei(1,arccosh(c*x)+a/b)*d^2+1/4/c^2/b*exp(a/b)*Ei(1,arccosh(c*x)+a/b)*d*e+1/16/c^4/b*exp(a/b)*Ei(1,arccosh(c*x)+a/b)*e^2-1/2/b*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)*d^2-1/4/c^2/b*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)*d*e-1/16/c^4/b*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)*e^2+1/4/c^2*e/b*exp(3*a/b)*Ei(1,3*arccosh(c*x)+3*a/b)*d+3/32/c^4*e^2/b*exp(3*a/b)*Ei(1,3*arccosh(c*x)+3*a/b)-1/4/c^2*e/b*exp(-3*a/b)*Ei(1,-3*arccosh(c*x)-3*a/b)*d-3/32/c^4*e^2/b*exp(-3*a/b)*Ei(1,-3*arccosh(c*x)-3*a/b))`

3.534. 
$$\int \frac{(d+ex^2)^2}{a+b\operatorname{arccosh}(cx)} dx$$

**3.534.5 Fracas [F]**

$$\int \frac{(d + ex^2)^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(ex^2 + d)^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((e^2*x^4 + 2*d*e*x^2 + d^2)/(b*arccosh(c*x) + a), x)`

**3.534.6 Sympy [F]**

$$\int \frac{(d + ex^2)^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(d + ex^2)^2}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate((e*x**2+d)**2/(a+b*acosh(c*x)),x)`

output `Integral((d + e*x**2)**2/(a + b*acosh(c*x)), x)`

**3.534.7 Maxima [F]**

$$\int \frac{(d + ex^2)^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(ex^2 + d)^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((e*x^2 + d)^2/(b*arccosh(c*x) + a), x)`

**3.534.8 Giac [F]**

$$\int \frac{(d + ex^2)^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(ex^2 + d)^2}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)^2/(b*arccosh(c*x) + a), x)`

**3.534.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{(ex^2 + d)^2}{a + b \operatorname{acosh}(cx)} dx$$

input `int((d + e*x^2)^2/(a + b*acosh(c*x)),x)`

output `int((d + e*x^2)^2/(a + b*acosh(c*x)), x)`

### 3.535 $\int \frac{d+ex^2}{a+b\operatorname{arccosh}(cx)} dx$

3.535.1 Optimal result . . . . .	3923
3.535.2 Mathematica [A] (verified) . . . . .	3924
3.535.3 Rubi [A] (verified) . . . . .	3924
3.535.4 Maple [A] (verified) . . . . .	3925
3.535.5 Fricas [F] . . . . .	3926
3.535.6 Sympy [F] . . . . .	3926
3.535.7 Maxima [F] . . . . .	3926
3.535.8 Giac [F] . . . . .	3927
3.535.9 Mupad [F(-1)] . . . . .	3927

#### 3.535.1 Optimal result

Integrand size = 18, antiderivative size = 139

$$\int \frac{d+ex^2}{a+b\operatorname{arccosh}(cx)} dx = -\frac{(4c^2d+e)\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)\sinh\left(\frac{a}{b}\right)}{4bc^3} - \frac{e\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)\sinh\left(\frac{3a}{b}\right)}{4bc^3} + \frac{(4c^2d+e)\cosh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4bc^3} + \frac{e\cosh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4bc^3}$$

```
output 1/4*(4*c^2*d+e)*cosh(a/b)*Shi((a+b*arccosh(c*x))/b)/b/c^3+1/4*e*cosh(3*a/b)*Shi(3*(a+b*arccosh(c*x))/b)/b/c^3-1/4*(4*c^2*d+e)*Chi((a+b*arccosh(c*x))/b)*sinh(a/b)/b/c^3-1/4*e*Chi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)/b/c^3
```

**3.535.2 Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.90

$$\int \frac{d + ex^2}{a + b \operatorname{arccosh}(cx)} dx$$

$$= \frac{-((4c^2d + e) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \sinh\left(\frac{a}{b}\right) - e \operatorname{Chi}\left(3\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)\right) \sinh\left(\frac{3a}{b}\right) + 4c^2d \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b}\right) - 4c^2d \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3a}{b}\right))}{4bc^3}$$

input `Integrate[(d + e*x^2)/(a + b*ArcCosh[c*x]),x]`output `(-((4*c^2*d + e)*CoshIntegral[a/b + ArcCosh[c*x]]*Sinh[a/b]) - e*CoshIntegral[3*(a/b + ArcCosh[c*x]]*Sinh[(3*a)/b] + 4*c^2*d*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + e*Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + e*Cosh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])])/(4*b*c^3)`**3.535.3 Rubi [A] (verified)**Time = 0.56 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.29, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{d + ex^2}{a + b \operatorname{arccosh}(cx)} dx$$

$$\downarrow 6324$$

$$\int \left( \frac{d}{a + b \operatorname{arccosh}(cx)} + \frac{ex^2}{a + b \operatorname{arccosh}(cx)} \right) dx$$

$$\downarrow 2009$$

$$\frac{e \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{4bc^3} - \frac{e \sinh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a + b \operatorname{arccosh}(cx))}{b}\right)}{4bc^3} +$$

$$\frac{e \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{4bc^3} + \frac{e \cosh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a + b \operatorname{arccosh}(cx))}{b}\right)}{4bc^3} -$$

$$\frac{d \sinh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{bc} + \frac{d \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{bc}$$

input `Int[(d + e*x^2)/(a + b*ArcCosh[c*x]),x]`

output `-(d*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b])/(b*c) - (e*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b])/(4*b*c^3) - (e*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b]*Sinh[(3*a)/b])/(4*b*c^3) + (d*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(b*c) + (e*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(4*b*c^3) + (e*Cosh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/(4*b*c^3)`

### 3.535.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

### 3.535.4 Maple [A] (verified)

Time = 0.94 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.28

method	result
derivativedivides	$-\frac{e e^{-\frac{3a}{b}} \operatorname{Ei}_1\left(-3 \operatorname{arccosh}(cx) - \frac{3a}{b}\right)}{8c^2b} + \frac{e e^{\frac{3a}{b}} \operatorname{Ei}_1\left(3 \operatorname{arccosh}(cx) + \frac{3a}{b}\right)}{8c^2b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right)d}{2b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right)e}{8c^2b} - \frac{e^{-\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) - \frac{a}{b}\right)e}{8c^2b} - \frac{e^{-\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) - \frac{a}{b}\right)d}{2b}$
default	$-\frac{e e^{-\frac{3a}{b}} \operatorname{Ei}_1\left(-3 \operatorname{arccosh}(cx) - \frac{3a}{b}\right)}{8c^2b} + \frac{e e^{\frac{3a}{b}} \operatorname{Ei}_1\left(3 \operatorname{arccosh}(cx) + \frac{3a}{b}\right)}{8c^2b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right)d}{2b} + \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right)e}{8c^2b} - \frac{e^{-\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) - \frac{a}{b}\right)e}{8c^2b} - \frac{e^{-\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) - \frac{a}{b}\right)d}{2b}$

input `int((e*x^2+d)/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`

output `1/c*(-1/8*e/c^2/b*exp(-3*a/b)*Ei(1,-3*arccosh(c*x)-3*a/b)+1/8*e/c^2/b*exp(3*a/b)*Ei(1,3*arccosh(c*x)+3*a/b)+1/2/b*exp(a/b)*Ei(1,arccosh(c*x)+a/b)*d+1/8/c^2/b*exp(a/b)*Ei(1,arccosh(c*x)+a/b)*e-1/2/b*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)*d-1/8/c^2/b*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)*e)`

3.535. 
$$\int \frac{d+ex^2}{a+b\operatorname{arccosh}(cx)} dx$$

**3.535.5 Fracas [F]**

$$\int \frac{d + ex^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{ex^2 + d}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral((e*x^2 + d)/(b*arccosh(c*x) + a), x)`

**3.535.6 Sympy [F]**

$$\int \frac{d + ex^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{d + ex^2}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate((e*x**2+d)/(a+b*acosh(c*x)),x)`

output `Integral((d + e*x**2)/(a + b*acosh(c*x)), x)`

**3.535.7 Maxima [F]**

$$\int \frac{d + ex^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{ex^2 + d}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate((e*x^2 + d)/(b*arccosh(c*x) + a), x)`

**3.535.8 Giac [F]**

$$\int \frac{d + ex^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{ex^2 + d}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate((e*x^2 + d)/(b*arccosh(c*x) + a), x)`

**3.535.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{d + ex^2}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{ex^2 + d}{a + b \operatorname{acosh}(cx)} dx$$

input `int((d + e*x^2)/(a + b*acosh(c*x)),x)`

output `int((d + e*x^2)/(a + b*acosh(c*x)), x)`



### 3.536 $\int \frac{1}{a+b\operatorname{arccosh}(cx)} dx$

3.536.1 Optimal result	3928
3.536.2 Mathematica [A] (verified)	3928
3.536.3 Rubi [C] (verified)	3929
3.536.4 Maple [A] (verified)	3932
3.536.5 Fricas [F]	3932
3.536.6 Sympy [F]	3932
3.536.7 Maxima [F]	3933
3.536.8 Giac [F]	3933
3.536.9 Mupad [F(-1)]	3933

#### 3.536.1 Optimal result

Integrand size = 10, antiderivative size = 54

$$\int \frac{1}{a + b\operatorname{arccosh}(cx)} dx = -\frac{\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right)}{bc} + \frac{\cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{bc}$$

output `cosh(a/b)*Shi((a+b*arccosh(c*x))/b)/b/c-Chi((a+b*arccosh(c*x))/b)*sinh(a/b)/b/c`

#### 3.536.2 Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.85

$$\int \frac{1}{a + b\operatorname{arccosh}(cx)} dx = -\frac{\operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) \sinh\left(\frac{a}{b}\right) - \cosh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)}{bc}$$

input `Integrate[(a + b*ArcCosh[c*x])^(-1),x]`

output `-((CoshIntegral[a/b + ArcCosh[c*x]]*Sinh[a/b] - Cosh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]])/(b*c))`

**3.536.3 Rubi [C] (verified)**

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$ , Rules used = {6296, 25, 3042, 26, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + \operatorname{barccosh}(cx)} dx \\
 & \quad \downarrow \text{6296} \\
 & \frac{\int -\frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc} \\
 & \quad \downarrow \text{25} \\
 & \frac{\int \frac{\sinh\left(\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc} \\
 & \quad \downarrow \text{3042} \\
 & \frac{\int -\frac{i \sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc} \\
 & \quad \downarrow \text{26} \\
 & \frac{i \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a + \operatorname{barccosh}(cx))}{bc} \\
 & \quad \downarrow \text{3784} \\
 & \frac{i \left( i \sinh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a + \operatorname{barccosh}(cx)) + \cosh\left(\frac{a}{b}\right) \int -\frac{i \sinh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a + \operatorname{barccosh}(cx)) \right)}{bc} \\
 & \quad \downarrow \text{26}
 \end{aligned}$$

$$i \left( i \sinh \left( \frac{a}{b} \right) \int \frac{\cosh \left( \frac{a+b \operatorname{arccosh}(cx)}{b} \right)}{a+b \operatorname{arccosh}(cx)} d(a + \operatorname{arccosh}(cx)) - i \cosh \left( \frac{a}{b} \right) \int \frac{\sinh \left( \frac{a+b \operatorname{arccosh}(cx)}{b} \right)}{a+b \operatorname{arccosh}(cx)} d(a + \operatorname{arccosh}(cx)) \right)$$

bc

$$\downarrow \text{3042}$$

$$i \left( i \sinh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{i(a+b \operatorname{arccosh}(cx)) + \pi/2}{b} \right)}{a+b \operatorname{arccosh}(cx)} d(a + \operatorname{arccosh}(cx)) - i \cosh \left( \frac{a}{b} \right) \int - \frac{i \sin \left( \frac{i(a+b \operatorname{arccosh}(cx))}{b} \right)}{a+b \operatorname{arccosh}(cx)} d(a + \operatorname{arccosh}(cx)) \right)$$

bc

$$\downarrow \text{26}$$

$$i \left( i \sinh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{i(a+b \operatorname{arccosh}(cx)) + \pi/2}{b} \right)}{a+b \operatorname{arccosh}(cx)} d(a + \operatorname{arccosh}(cx)) - \cosh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{i(a+b \operatorname{arccosh}(cx))}{b} \right)}{a+b \operatorname{arccosh}(cx)} d(a + \operatorname{arccosh}(cx)) \right)$$

bc

$$\downarrow \text{3779}$$

$$i \left( i \sinh \left( \frac{a}{b} \right) \int \frac{\sin \left( \frac{i(a+b \operatorname{arccosh}(cx)) + \pi/2}{b} \right)}{a+b \operatorname{arccosh}(cx)} d(a + \operatorname{arccosh}(cx)) - i \cosh \left( \frac{a}{b} \right) \operatorname{Shi} \left( \frac{a+b \operatorname{arccosh}(cx)}{b} \right) \right)$$

bc

$$\downarrow \text{3782}$$

$$i \left( i \sinh \left( \frac{a}{b} \right) \operatorname{Chi} \left( \frac{a+b \operatorname{arccosh}(cx)}{b} \right) - i \cosh \left( \frac{a}{b} \right) \operatorname{Shi} \left( \frac{a+b \operatorname{arccosh}(cx)}{b} \right) \right)$$

bc

input `Int[(a + b*ArcCosh[c*x])^(-1), x]`

output `(I*(I*CoshIntegral[(a + b*ArcCosh[c*x])/b]*Sinh[a/b] - I*Cosh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b]))/(b*c)`

## 3.536.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`
- rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`
- rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`
- rule 6296 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[1/(b*c) Subst[Int[x^n*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x]`

**3.536.4 Maple [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.04

method	result	size
derivativedivides	$\frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right) - e^{-\frac{a}{b}} \operatorname{Ei}_1\left(-\operatorname{arccosh}(cx) - \frac{a}{b}\right)}{2b} \cdot c$	56
default	$\frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx) + \frac{a}{b}\right) - e^{-\frac{a}{b}} \operatorname{Ei}_1\left(-\operatorname{arccosh}(cx) - \frac{a}{b}\right)}{2b} \cdot c$	56

input `int(1/(a+b*arccosh(c*x)),x,method=_RETURNVERBOSE)`output `1/c*(1/2/b*exp(a/b)*Ei(1,arccosh(c*x)+a/b)-1/2/b*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b))`**3.536.5 Fracas [F]**

$$\int \frac{1}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{1}{b \operatorname{arccosh}(cx) + a} dx$$

input `integrate(1/(a+b*arccosh(c*x)),x, algorithm="fracas")`output `integral(1/(b*arccosh(c*x) + a), x)`**3.536.6 Sympy [F]**

$$\int \frac{1}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{1}{a + b \operatorname{acosh}(cx)} dx$$

input `integrate(1/(a+b*acosh(c*x)),x)`output `Integral(1/(a + b*acosh(c*x)), x)`

**3.536.7 Maxima [F]**

$$\int \frac{1}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{1}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(1/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(1/(b*arccosh(c*x) + a), x)`

**3.536.8 Giac [F]**

$$\int \frac{1}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{1}{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate(1/(a+b*arccosh(c*x)),x, algorithm="giac")`

output `integrate(1/(b*arccosh(c*x) + a), x)`

**3.536.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{a + b \operatorname{arccosh}(cx)} dx = \int \frac{1}{a + b \operatorname{acosh}(cx)} dx$$

input `int(1/(a + b*acosh(c*x)),x)`

output `int(1/(a + b*acosh(c*x)), x)`

**3.537**  $\int \frac{1}{(d+ex^2)(a+b\mathbf{arccosh}(cx))} dx$

3.537.1 Optimal result . . . . .	3934
3.537.2 Mathematica [N/A] . . . . .	3934
3.537.3 Rubi [N/A] . . . . .	3935
3.537.4 Maple [N/A] (verified) . . . . .	3935
3.537.5 Fricas [N/A] . . . . .	3936
3.537.6 Sympy [N/A] . . . . .	3936
3.537.7 Maxima [N/A] . . . . .	3936
3.537.8 Giac [N/A] . . . . .	3937
3.537.9 Mupad [N/A] . . . . .	3937

**3.537.1 Optimal result**

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(d+ex^2)(a+b\mathbf{arccosh}(cx))} dx = \mathbf{Int}\left(\frac{1}{(d+ex^2)(a+b\mathbf{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/(e*x^2+d)/(a+b*arccosh(c*x)),x)`

**3.537.2 Mathematica [N/A]**

Not integrable

Time = 0.57 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d+ex^2)(a+b\mathbf{arccosh}(cx))} dx = \int \frac{1}{(d+ex^2)(a+b\mathbf{arccosh}(cx))} dx$$

input `Integrate[1/((d + e*x^2)*(a + b*ArcCosh[c*x])),x]`

output `Integrate[1/((d + e*x^2)*(a + b*ArcCosh[c*x])), x]`

**3.537.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)(a + \operatorname{arccosh}(cx))} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)(a + \operatorname{arccosh}(cx))} dx$$

input `Int[1/((d + e*x^2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.537.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n_)*((d_) + (e_.)*(x_)^2)^p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.537.4 Maple [N/A] (verified)**

Not integrable

Time = 0.82 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(ex^2 + d)(a + b \operatorname{arccosh}(cx))} dx$$

input `int(1/(e*x^2+d)/(a+b*arccosh(c*x)),x)`

output `int(1/(e*x^2+d)/(a+b*arccosh(c*x)),x)`



**3.537.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.45

$$\int \frac{1}{(d + ex^2)(a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x)),x, algorithm="fricas")`output `integral(1/(a*e*x^2 + a*d + (b*e*x^2 + b*d)*arccosh(c*x)), x)`**3.537.6 Sympy [N/A]**

Not integrable

Time = 9.36 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{1}{(d + ex^2)(a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))(d + ex^2)} dx$$

input `integrate(1/(e*x**2+d)/(a+b*acosh(c*x)),x)`output `Integral(1/((a + b*acosh(c*x))*(d + e*x**2)), x)`**3.537.7 Maxima [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)(a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x)),x, algorithm="maxima")`output `integrate(1/((e*x^2 + d)*(b*arccosh(c*x) + a)), x)`

**3.537.8 Giac [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)(a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)*(b*arccosh(c*x) + a)), x)`**3.537.9 Mupad [N/A]**

Not integrable

Time = 2.87 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)(a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))(ex^2 + d)} dx$$

input `int(1/((a + b*acosh(c*x))*(d + e*x^2)),x)`output `int(1/((a + b*acosh(c*x))*(d + e*x^2)), x)`

**3.538**  $\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))} dx$

3.538.1 Optimal result . . . . .	3938
3.538.2 Mathematica [N/A] . . . . .	3938
3.538.3 Rubi [N/A] . . . . .	3939
3.538.4 Maple [N/A] (verified) . . . . .	3939
3.538.5 Fricas [N/A] . . . . .	3940
3.538.6 Sympy [F(-1)] . . . . .	3940
3.538.7 Maxima [N/A] . . . . .	3940
3.538.8 Giac [N/A] . . . . .	3941
3.538.9 Mupad [N/A] . . . . .	3941

**3.538.1 Optimal result**

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^2/(a+b*arccosh(c*x)),x)`

**3.538.2 Mathematica [N/A]**

Not integrable

Time = 2.61 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])),x]`

output `Integrate[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])), x]`

**3.538.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))} dx$$

input `Int[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.538.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.538.4 Maple [N/A] (verified)**

Not integrable

Time = 0.74 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(ex^2 + d)^2 (a + b \operatorname{arccosh}(cx))} dx$$

input `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x)),x)`

output `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x)),x)`

**3.538.5 Fracas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.65

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(1/(a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arccosh(c*x)), x)`

**3.538.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))} dx = \text{Timed out}$$

input `integrate(1/(e*x**2+d)**2/(a+b*acosh(c*x)),x)`

output `Timed out`

**3.538.7 Maxima [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(1/((e*x^2 + d)^2*(b*arccosh(c*x) + a)), x)`

**3.538.8 Giac [N/A]**

Not integrable

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^2 (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)^2*(b*arccosh(c*x) + a)), x)`**3.538.9 Mupad [N/A]**

Not integrable

Time = 2.84 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^2} dx$$

input `int(1/((a + b*acosh(c*x))*(d + e*x^2)^2),x)`output `int(1/((a + b*acosh(c*x))*(d + e*x^2)^2), x)`

$$3.539 \quad \int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx$$

3.539.1 Optimal result . . . . .	3942
3.539.2 Mathematica [N/A] . . . . .	3942
3.539.3 Rubi [N/A] . . . . .	3943
3.539.4 Maple [N/A] (verified) . . . . .	3943
3.539.5 Fricas [N/A] . . . . .	3944
3.539.6 Sympy [N/A] . . . . .	3944
3.539.7 Maxima [N/A] . . . . .	3944
3.539.8 Giac [N/A] . . . . .	3945
3.539.9 Mupad [N/A] . . . . .	3945

### 3.539.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx = \operatorname{Int}\left(\frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)}, x\right)$$

output `Unintegrable((e*x^2+d)^(1/2)/(a+b*arccosh(c*x)),x)`

### 3.539.2 Mathematica [N/A]

Not integrable

Time = 0.98 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx$$

input `Integrate[Sqrt[d + e*x^2]/(a + b*ArcCosh[c*x]),x]`

output `Integrate[Sqrt[d + e*x^2]/(a + b*ArcCosh[c*x]), x]`

**3.539.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx$$

↓ 6325

$$\int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx$$

input `Int[Sqrt[d + e*x^2]/(a + b*ArcCosh[c*x]),x]`

output `$Aborted`

**3.539.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.539.4 Maple [N/A] (verified)**

Not integrable

Time = 1.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{ex^2+d}}{a+b\operatorname{arccosh}(cx)} dx$$

input `int((e*x^2+d)^(1/2)/(a+b*arccosh(c*x)),x)`

output `int((e*x^2+d)^(1/2)/(a+b*arccosh(c*x)),x)`



**3.539.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}}{a+\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{ex^2+d}}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate((e*x^2+d)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)/(b*arccosh(c*x) + a), x)`

**3.539.6 Sympy [N/A]**

Not integrable

Time = 0.47 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{\sqrt{d+ex^2}}{a+\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{d+ex^2}}{a+b\operatorname{acosh}(cx)} dx$$

input `integrate((e*x**2+d)**(1/2)/(a+b*acosh(c*x)),x)`

output `Integral(sqrt(d + e*x**2)/(a + b*acosh(c*x)), x)`

**3.539.7 Maxima [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}}{a+\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{ex^2+d}}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate((e*x^2+d)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(sqrt(e*x^2 + d)/(b*arccosh(c*x) + a), x)`

**3.539.8 Giac [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{ex^2+d}}{b\operatorname{arcosh}(cx)+a} dx$$

input `integrate((e*x^2+d)^(1/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(sqrt(e*x^2 + d)/(b*arccosh(c*x) + a), x)`**3.539.9 Mupad [N/A]**

Not integrable

Time = 2.79 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d+ex^2}}{a+b\operatorname{arccosh}(cx)} dx = \int \frac{\sqrt{ex^2+d}}{a+b\operatorname{acosh}(cx)} dx$$

input `int((d + e*x^2)^(1/2)/(a + b*acosh(c*x)),x)`output `int((d + e*x^2)^(1/2)/(a + b*acosh(c*x)), x)`

**3.540**      $\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx$

3.540.1 Optimal result . . . . . 3946  
 3.540.2 Mathematica [N/A] . . . . . 3946  
 3.540.3 Rubi [N/A] . . . . . 3947  
 3.540.4 Maple [N/A] (verified) . . . . . 3947  
 3.540.5 Fricas [N/A] . . . . . 3948  
 3.540.6 Sympy [N/A] . . . . . 3948  
 3.540.7 Maxima [N/A] . . . . . 3948  
 3.540.8 Giac [N/A] . . . . . 3949  
 3.540.9 Mupad [N/A] . . . . . 3949

**3.540.1 Optimal result**

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx = \operatorname{Int}\left(\frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/(a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x)`

**3.540.2 Mathematica [N/A]**

Not integrable

Time = 0.81 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx$$

input `Integrate[1/(Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])),x]`

output `Integrate[1/(Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])), x]`

**3.540.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{d + ex^2}(a + \operatorname{arccosh}(cx))} dx$$

↓ 6325

$$\int \frac{1}{\sqrt{d + ex^2}(a + \operatorname{arccosh}(cx))} dx$$

input `Int[1/(Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.540.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.540.4 Maple [N/A] (verified)**

Not integrable

Time = 1.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx)) \sqrt{ex^2 + d}} dx$$

input `int(1/(a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x)`

output `int(1/(a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x)`

**3.540.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.77

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{ex^2+d}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(1/(a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x, algorithm="fricas")`output `integral(sqrt(e*x^2 + d)/(a*e*x^2 + a*d + (b*e*x^2 + b*d)*arccosh(c*x)), x)`**3.540.6 Sympy [N/A]**

Not integrable

Time = 0.73 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{(a+b\operatorname{acosh}(cx))\sqrt{d+ex^2}} dx$$

input `integrate(1/(a+b*acosh(c*x))/(e*x**2+d)**(1/2),x)`output `Integral(1/((a + b*acosh(c*x))*sqrt(d + e*x**2)), x)`**3.540.7 Maxima [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{ex^2+d}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(1/(a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x, algorithm="maxima")`output `integrate(1/(sqrt(e*x^2 + d)*(b*arccosh(c*x) + a)), x)`

---

3.540.  $\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx$

**3.540.8 Giac [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{\sqrt{ex^2+d}(b\operatorname{arcosh}(cx)+a)} dx$$

input `integrate(1/(a+b*arccosh(c*x))/(e*x^2+d)^(1/2),x, algorithm="giac")`output `integrate(1/(sqrt(e*x^2 + d)*(b*arccosh(c*x) + a)), x)`**3.540.9 Mupad [N/A]**

Not integrable

Time = 2.82 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))} dx = \int \frac{1}{(a+b\operatorname{acosh}(cx))\sqrt{ex^2+d}} dx$$

input `int(1/((a + b*acosh(c*x))*(d + e*x^2)^(1/2)),x)`output `int(1/((a + b*acosh(c*x))*(d + e*x^2)^(1/2)), x)`

**3.541**  $\int \frac{1}{(d+ex^2)^{3/2}(a+b\text{arccosh}(cx))} dx$

3.541.1 Optimal result . . . . .	3950
3.541.2 Mathematica [N/A] . . . . .	3950
3.541.3 Rubi [N/A] . . . . .	3951
3.541.4 Maple [N/A] (verified) . . . . .	3951
3.541.5 Fricas [N/A] . . . . .	3952
3.541.6 Sympy [N/A] . . . . .	3952
3.541.7 Maxima [N/A] . . . . .	3952
3.541.8 Giac [N/A] . . . . .	3953
3.541.9 Mupad [N/A] . . . . .	3953

**3.541.1 Optimal result**

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \text{arccosh}(cx))} dx = \text{Int}\left(\frac{1}{(d + ex^2)^{3/2} (a + \text{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x)),x)`

**3.541.2 Mathematica [N/A]**

Not integrable

Time = 1.42 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \text{arccosh}(cx))} dx = \int \frac{1}{(d + ex^2)^{3/2} (a + \text{arccosh}(cx))} dx$$

input `Integrate[1/((d + e*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `Integrate[1/((d + e*x^2)^(3/2)*(a + b*ArcCosh[c*x])), x]`

**3.541.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))} dx$$

input `Int[1/((d + e*x^2)^(3/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.541.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.541.4 Maple [N/A] (verified)**

Not integrable

Time = 1.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x)),x)`

output `int(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x)),x)`



**3.541.5 Fracas [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.86

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)/(a*e^2*x^4 + 2*a*d*e*x^2 + a*d^2 + (b*e^2*x^4 + 2*b*d*e*x^2 + b*d^2)*arccosh(c*x)), x)`

**3.541.6 Sympy [N/A]**

Not integrable

Time = 3.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx)) (d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate(1/(e*x**2+d)**(3/2)/(a+b*acosh(c*x)),x)`

output `Integral(1/((a + b*acosh(c*x))*(d + e*x**2)**(3/2)), x)`

**3.541.7 Maxima [N/A]**

Not integrable

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(1/((e*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)), x)`

---

3.541.  $\int \frac{1}{(d+ex^2)^{3/2}(a+b\operatorname{arccosh}(cx))} dx$

**3.541.8 Giac [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{3/2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)), x)`**3.541.9 Mupad [N/A]**

Not integrable

Time = 2.96 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{3/2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^{3/2}} dx$$

input `int(1/((a + b*acosh(c*x))*(d + e*x^2)^(3/2)),x)`output `int(1/((a + b*acosh(c*x))*(d + e*x^2)^(3/2)), x)`

$$3.542 \quad \int \frac{1}{(d+ex^2)^{5/2}(a+b\text{arccosh}(cx))} dx$$

3.542.1 Optimal result . . . . .	3954
3.542.2 Mathematica [N/A] . . . . .	3954
3.542.3 Rubi [N/A] . . . . .	3955
3.542.4 Maple [N/A] (verified) . . . . .	3955
3.542.5 Fricas [N/A] . . . . .	3956
3.542.6 Sympy [N/A] . . . . .	3956
3.542.7 Maxima [N/A] . . . . .	3956
3.542.8 Giac [N/A] . . . . .	3957
3.542.9 Mupad [N/A] . . . . .	3957

### 3.542.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d+ex^2)^{5/2}(a+b\text{arccosh}(cx))} dx = \text{Int}\left(\frac{1}{(d+ex^2)^{5/2}(a+b\text{arccosh}(cx))}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x)),x)`

### 3.542.2 Mathematica [N/A]

Not integrable

Time = 2.81 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d+ex^2)^{5/2}(a+b\text{arccosh}(cx))} dx = \int \frac{1}{(d+ex^2)^{5/2}(a+b\text{arccosh}(cx))} dx$$

input `Integrate[1/((d + e*x^2)^(5/2)*(a + b*ArcCosh[c*x])),x]`

output `Integrate[1/((d + e*x^2)^(5/2)*(a + b*ArcCosh[c*x])), x]`

**3.542.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))} dx$$

input `Int[1/((d + e*x^2)^(5/2)*(a + b*ArcCosh[c*x])),x]`

output `$Aborted`

**3.542.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.542.4 Maple [N/A] (verified)**

Not integrable

Time = 1.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d)^{5/2} (a + b \operatorname{arccosh}(cx))} dx$$

input `int(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x)),x)`

output `int(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x)),x)`

**3.542.5 Fracas [N/A]**

Not integrable

Time = 0.31 (sec) , antiderivative size = 87, normalized size of antiderivative = 3.95

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)/(a*e^3*x^6 + 3*a*d*e^2*x^4 + 3*a*d^2*e*x^2 + a*d^3 + (b*e^3*x^6 + 3*b*d*e^2*x^4 + 3*b*d^2*e*x^2 + b*d^3)*arccosh(c*x)), x)`

**3.542.6 Sympy [N/A]**

Not integrable

Time = 34.56 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx)) (d + ex^2)^{5/2}} dx$$

input `integrate(1/(e*x**2+d)**(5/2)/(a+b*acosh(c*x)),x)`

output `Integral(1/((a + b*acosh(c*x))*(d + e*x**2)**(5/2)), x)`

**3.542.7 Maxima [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="maxima")`

output `integrate(1/((e*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)), x)`

---

3.542.  $\int \frac{1}{(d+ex^2)^{5/2}(a+b\operatorname{arccosh}(cx))} dx$

**3.542.8 Giac [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{5/2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{(ex^2 + d)^{5/2} (b \operatorname{arcosh}(cx) + a)} dx$$

input `integrate(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x)),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)), x)`**3.542.9 Mupad [N/A]**

Not integrable

Time = 2.96 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{5/2} (a + b \operatorname{arccosh}(cx))} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx)) (ex^2 + d)^{5/2}} dx$$

input `int(1/((a + b*acosh(c*x))*(d + e*x^2)^(5/2)),x)`output `int(1/((a + b*acosh(c*x))*(d + e*x^2)^(5/2)), x)`

$$3.543 \quad \int \frac{(d+ex^2)^2}{(a+b\operatorname{arccosh}(cx))^2} dx$$

3.543.1 Optimal result	3959
3.543.2 Mathematica [A] (warning: unable to verify)	3960
3.543.3 Rubi [A] (verified)	3961
3.543.4 Maple [B] (verified)	3963
3.543.5 Fricas [F]	3964
3.543.6 Sympy [F]	3965
3.543.7 Maxima [F]	3965
3.543.8 Giac [F]	3966
3.543.9 Mupad [F(-1)]	3966

**3.543.1 Optimal result**

Integrand size = 20, antiderivative size = 510

$$\begin{aligned}
\int \frac{(d+ex^2)^2}{(a+b\operatorname{arccosh}(cx))^2} dx = & -\frac{d^2\sqrt{-1+cx}\sqrt{1+cx}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{2dex^2\sqrt{-1+cx}\sqrt{1+cx}}{bc(a+b\operatorname{arccosh}(cx))} \\
& - \frac{e^2x^4\sqrt{-1+cx}\sqrt{1+cx}}{bc(a+b\operatorname{arccosh}(cx))} + \frac{d^2\cosh\left(\frac{a}{b}\right)\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c} \\
& + \frac{de\cosh\left(\frac{a}{b}\right)\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2b^2c^3} \\
& + \frac{e^2\cosh\left(\frac{a}{b}\right)\operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b^2c^5} \\
& + \frac{3de\cosh\left(\frac{3a}{b}\right)\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{2b^2c^3} \\
& + \frac{9e^2\cosh\left(\frac{3a}{b}\right)\operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^5} \\
& + \frac{5e^2\cosh\left(\frac{5a}{b}\right)\operatorname{Chi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^5} \\
& - \frac{d^2\sinh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c} \\
& - \frac{de\sinh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{2b^2c^3} \\
& - \frac{e^2\sinh\left(\frac{a}{b}\right)\operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{8b^2c^5} \\
& - \frac{3de\sinh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{2b^2c^3} \\
& - \frac{9e^2\sinh\left(\frac{3a}{b}\right)\operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^5} \\
& - \frac{5e^2\sinh\left(\frac{5a}{b}\right)\operatorname{Shi}\left(\frac{5(a+b\operatorname{arccosh}(cx))}{b}\right)}{16b^2c^5}
\end{aligned}$$



output  $d^2 \operatorname{Chi}\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \cosh\left(\frac{a}{b}\right) / b^2 / c + 1/2 d e \operatorname{Chi}\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \cosh\left(\frac{a}{b}\right) / b^2 / c^3 + 1/8 e^2 \operatorname{Chi}\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \cosh\left(\frac{a}{b}\right) / b^2 / c^5 + 3/2 d e \operatorname{Chi}\left(\frac{3(a+b \operatorname{arccosh}(cx))}{b}\right) \cosh\left(\frac{3a}{b}\right) / b^2 / c^3 + 9/16 e^2 \operatorname{Chi}\left(\frac{3(a+b \operatorname{arccosh}(cx))}{b}\right) \cosh\left(\frac{3a}{b}\right) / b^2 / c^5 + 5/16 e^2 \operatorname{Chi}\left(\frac{5(a+b \operatorname{arccosh}(cx))}{b}\right) \cosh\left(\frac{5a}{b}\right) / b^2 / c^5 - d^2 \operatorname{Shi}\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right) / b^2 / c - 1/2 d e \operatorname{Shi}\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right) / b^2 / c^3 - 1/8 e^2 \operatorname{Shi}\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \sinh\left(\frac{a}{b}\right) / b^2 / c^5 - 3/2 d e \operatorname{Shi}\left(\frac{3(a+b \operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{3a}{b}\right) / b^2 / c^3 - 9/16 e^2 \operatorname{Shi}\left(\frac{3(a+b \operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{3a}{b}\right) / b^2 / c^5 - 5/16 e^2 \operatorname{Shi}\left(\frac{5(a+b \operatorname{arccosh}(cx))}{b}\right) \sinh\left(\frac{5a}{b}\right) / b^2 / c^5 - d^2 (cx-1)^{1/2} (cx+1)^{1/2} / b / c / (a+b \operatorname{arccosh}(cx)) - 2 d e x^2 (cx-1)^{1/2} (cx+1)^{1/2} / b / c / (a+b \operatorname{arccosh}(cx)) - e^2 x^4 (cx-1)^{1/2} (cx+1)^{1/2} / b / c / (a+b \operatorname{arccosh}(cx))$

### 3.543.2 Mathematica [A] (warning: unable to verify)

Time = 2.58 (sec) , antiderivative size = 663, normalized size of antiderivative = 1.30

$$\int \frac{(d + ex^2)^2}{(a + b \operatorname{arccosh}(cx))^2} dx = \frac{16bc^4 d^2 \sqrt{\frac{-1+cx}{1+cx}} + 16bc^5 d^2 x \sqrt{\frac{-1+cx}{1+cx}} + 32bc^4 dex^2 \sqrt{\frac{-1+cx}{1+cx}} + 32bc^5 dex^3 \sqrt{\frac{-1+cx}{1+cx}} + 16bc^4 e^2 x^4 \sqrt{\frac{-1+cx}{1+cx}} + 16bc^5 e^2 x^5 \sqrt{\frac{-1+cx}{1+cx}}}{(a+b \operatorname{arccosh}(cx))^2}$$

input `Integrate[(d + e*x^2)^2/(a + b*ArcCosh[c*x])^2,x]`

output

```

-1/16*(16*b*c^4*d^2*Sqrt[(-1 + c*x)/(1 + c*x)] + 16*b*c^5*d^2*x*Sqrt[(-1 +
c*x)/(1 + c*x)] + 32*b*c^4*d*e*x^2*Sqrt[(-1 + c*x)/(1 + c*x)] + 32*b*c^5*
d*e*x^3*Sqrt[(-1 + c*x)/(1 + c*x)] + 16*b*c^4*e^2*x^4*Sqrt[(-1 + c*x)/(1 +
c*x)] + 16*b*c^5*e^2*x^5*Sqrt[(-1 + c*x)/(1 + c*x)] - 2*(8*c^4*d^2 + 4*c^
2*d*e + e^2)*(a + b*ArcCosh[c*x])*Cosh[a/b]*CoshIntegral[a/b + ArcCosh[c*x
]] - 3*e*(8*c^2*d + 3*e)*(a + b*ArcCosh[c*x])*Cosh[(3*a)/b]*CoshIntegral[3
*(a/b + ArcCosh[c*x])] - 5*a*e^2*Cosh[(5*a)/b]*CoshIntegral[5*(a/b + ArcCo
sh[c*x])] - 5*b*e^2*ArcCosh[c*x]*Cosh[(5*a)/b]*CoshIntegral[5*(a/b + ArcCo
sh[c*x])] + 16*a*c^4*d^2*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + 8*a*
c^2*d*e*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + 2*a*e^2*Sinh[a/b]*Sin
hIntegral[a/b + ArcCosh[c*x]] + 16*b*c^4*d^2*ArcCosh[c*x]*Sinh[a/b]*SinhIn
tegral[a/b + ArcCosh[c*x]] + 8*b*c^2*d*e*ArcCosh[c*x]*Sinh[a/b]*SinhIntegr
al[a/b + ArcCosh[c*x]] + 2*b*e^2*ArcCosh[c*x]*Sinh[a/b]*SinhIntegral[a/b +
ArcCosh[c*x]] + 24*a*c^2*d*e*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[
c*x])] + 9*a*e^2*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] + 24*b
*c^2*d*e*ArcCosh[c*x]*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] +
9*b*e^2*ArcCosh[c*x]*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] +
5*a*e^2*Sinh[(5*a)/b]*SinhIntegral[5*(a/b + ArcCosh[c*x])] + 5*b*e^2*ArcC
osh[c*x]*Sinh[(5*a)/b]*SinhIntegral[5*(a/b + ArcCosh[c*x])]/(b^2*c^5*(a +
b*ArcCosh[c*x]))

```

### 3.543.3 Rubi [A] (verified)

Time = 1.19 (sec) , antiderivative size = 510, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2}{(a + \operatorname{barccosh}(cx))^2} dx$$

$$\downarrow \text{6324}$$

$$\int \left( \frac{d^2}{(a + \operatorname{barccosh}(cx))^2} + \frac{2dex^2}{(a + \operatorname{barccosh}(cx))^2} + \frac{e^2x^4}{(a + \operatorname{barccosh}(cx))^2} \right) dx$$

$$\downarrow \text{2009}$$

---

3.543.  $\int \frac{(d+ex^2)^2}{(a+\operatorname{barccosh}(cx))^2} dx$

$$\begin{aligned}
& \frac{e^2 \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{8b^2c^5} + \frac{9e^2 \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{16b^2c^5} + \\
& \frac{5e^2 \cosh\left(\frac{5a}{b}\right) \operatorname{Chi}\left(\frac{5(a+\operatorname{barccosh}(cx))}{b}\right)}{16b^2c^5} - \frac{e^2 \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{8b^2c^5} - \\
& \frac{9e^2 \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{16b^2c^5} - \frac{5e^2 \sinh\left(\frac{5a}{b}\right) \operatorname{Shi}\left(\frac{5(a+\operatorname{barccosh}(cx))}{b}\right)}{16b^2c^5} + \\
& \frac{de \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{2b^2c^3} + \frac{3de \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{2b^2c^3} - \\
& \frac{de \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{2b^2c^3} - \frac{3de \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{2b^2c^3} + \\
& \frac{d^2 \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{b^2c} - \frac{d^2 \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{b^2c} - \frac{d^2 \sqrt{cx-1} \sqrt{cx+1}}{bc(a+\operatorname{barccosh}(cx))} - \\
& \frac{2dex^2 \sqrt{cx-1} \sqrt{cx+1}}{bc(a+\operatorname{barccosh}(cx))} - \frac{e^2 x^4 \sqrt{cx-1} \sqrt{cx+1}}{bc(a+\operatorname{barccosh}(cx))}
\end{aligned}$$

input `Int[(d + e*x^2)^2/(a + b*ArcCosh[c*x])^2,x]`

output `-((d^2*sqrt[-1 + c*x]*sqrt[1 + c*x])/(b*c*(a + b*ArcCosh[c*x]))) - (2*d*e*x^2*sqrt[-1 + c*x]*sqrt[1 + c*x])/(b*c*(a + b*ArcCosh[c*x])) - (e^2*x^4*sqrt[-1 + c*x]*sqrt[1 + c*x])/(b*c*(a + b*ArcCosh[c*x])) + (d^2*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/(b^2*c) + (d*e*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/(2*b^2*c^3) + (e^2*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/(8*b^2*c^5) + (3*d*e*Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/(2*b^2*c^3) + (9*e^2*Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/(16*b^2*c^5) + (5*e^2*Cosh[(5*a)/b]*CoshIntegral[(5*(a + b*ArcCosh[c*x])/b])/(16*b^2*c^5) - (d^2*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(b^2*c) - (d*e*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(2*b^2*c^3) - (e^2*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(8*b^2*c^5) - (3*d*e*Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/(2*b^2*c^3) - (9*e^2*Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/(16*b^2*c^5) - (5*e^2*Sinh[(5*a)/b]*SinhIntegral[(5*(a + b*ArcCosh[c*x])/b])/(16*b^2*c^5)`

### 3.543.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x],  
x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] &&  
(p > 0 || IGtQ[n, 0])`

### 3.543.4 Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1101 vs. 2(478) = 956.

Time = 1.98 (sec) , antiderivative size = 1102, normalized size of antiderivative = 2.16

method	result	size
derivativedivides	Expression too large to display	1102
default	Expression too large to display	1102

input `int((e*x^2+d)^2/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`

---

3.543.  $\int \frac{(d+ex^2)^2}{(a+b\operatorname{arccosh}(cx))^2} dx$

output `1/c*(1/32*(-16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+12*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2)+16*c^5*x^5-20*c^3*x^3+5*c*x)*e^2/c^4/b/(a+b*arccosh(c*x))-5/32*e^2/c^4/b^2*exp(5*a/b)*Ei(1,5*arccosh(c*x)+5*a/b)-1/32/b*e^2/c^4*(16*c^5*x^5-20*c^3*x^3+16*(c*x+1)^(1/2)*(c*x-1)^(1/2)*c^4*x^4+5*c*x-12*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))-5/32/b^2*e^2/c^4*exp(-5*a/b)*Ei(1,-5*arccosh(c*x)-5*a/b)+1/2*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*d^2/b/(a+b*arccosh(c*x))-1/2*d^2/b^2*exp(a/b)*Ei(1,arccosh(c*x)+a/b)+1/4*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*d*e/c^2/b/(a+b*arccosh(c*x))-1/4/c^2*d*e/b^2*exp(a/b)*Ei(1,arccosh(c*x)+a/b)+1/16*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*e^2/c^4/b/(a+b*arccosh(c*x))-1/16/c^4*e^2/b^2*exp(a/b)*Ei(1,arccosh(c*x)+a/b)-1/2/b*d^2*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))-1/2/b^2*d^2*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)-1/4/c^2/b*d*e*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))-1/4/c^2/b^2*d*e*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)-1/16/c^4/b*e^2*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))-1/16/c^4/b^2*e^2*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)+1/4*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c^3*x^3-3*c*x)*d*e/c^2/b/(a+b*arccosh(c*x))+3/32*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c^3*x^3-3*c*x)*e^2/c^4/b/(a+b*arccosh(c*x))-3/4*e/c^2/b^2*exp(3*a/b)*Ei(1,3*arccosh(c*x)+3*a/b)*d-9/32*e^2/c^4/b^2*exp(3*a/b)*Ei(1,3*arccos...`

### 3.543.5 Fracas [F]

$$\int \frac{(d + ex^2)^2}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(ex^2 + d)^2}{(b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral((e^2*x^4 + 2*d*e*x^2 + d^2)/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`

## 3.543.6 Sympy [F]

$$\int \frac{(d + ex^2)^2}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(d + ex^2)^2}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((e*x**2+d)**2/(a+b*acosh(c*x))**2,x)`

output `Integral((d + e*x**2)**2/(a + b*acosh(c*x))**2, x)`

## 3.543.7 Maxima [F]

$$\int \frac{(d + ex^2)^2}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{(ex^2 + d)^2}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-(c^3*e^2*x^7 + (2*c^3*d*e - c*e^2)*x^5 - c*d^2*x + (c^3*d^2 - 2*c*d*e)*x^3 + (c^2*e^2*x^6 + (2*c^2*d*e - e^2)*x^4 + (c^2*d^2 - 2*d*e)*x^2 - d^2)*sqrt(c*x + 1)*sqrt(c*x - 1))/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate((5*c^5*e^2*x^8 + 2*(3*c^5*d*e - 5*c^3*e^2)*x^6 + (c^5*d^2 - 12*c^3*d*e + 5*c*e^2)*x^4 + (5*c^3*e^2*x^6 + 3*(2*c^3*d*e - c*e^2)*x^4 + c*d^2 + (c^3*d^2 - 2*c*d*e)*x^2)*(c*x + 1)*(c*x - 1) + c*d^2 - 2*(c^3*d^2 - 3*c*d*e)*x^2 + (10*c^4*e^2*x^7 + (12*c^4*d*e - 13*c^2*e^2)*x^5 + 2*(c^4*d^2 - 7*c^2*d*e + 2*e^2)*x^3 - (c^2*d^2 - 4*d*e)*x)*sqrt(c*x + 1)*sqrt(c*x - 1))/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.543.8 Giac [F]**

$$\int \frac{(d + ex^2)^2}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(ex^2 + d)^2}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((e*x^2 + d)^2/(b*arccosh(c*x) + a)^2, x)`

**3.543.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{(ex^2 + d)^2}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `int((d + e*x^2)^2/(a + b*acosh(c*x))^2,x)`

output `int((d + e*x^2)^2/(a + b*acosh(c*x))^2, x)`

### 3.544 $\int \frac{d+ex^2}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.544.1 Optimal result . . . . .	3967
3.544.2 Mathematica [A] (warning: unable to verify) . . . . .	3968
3.544.3 Rubi [A] (verified) . . . . .	3968
3.544.4 Maple [A] (verified) . . . . .	3970
3.544.5 Fracas [F] . . . . .	3970
3.544.6 Sympy [F] . . . . .	3971
3.544.7 Maxima [F] . . . . .	3971
3.544.8 Giac [F] . . . . .	3971
3.544.9 Mupad [F(-1)] . . . . .	3972

#### 3.544.1 Optimal result

Integrand size = 18, antiderivative size = 257

$$\int \frac{d+ex^2}{(a+b\operatorname{arccosh}(cx))^2} dx = -\frac{d\sqrt{-1+cx}\sqrt{1+cx}}{bc(a+b\operatorname{arccosh}(cx))} - \frac{ex^2\sqrt{-1+cx}\sqrt{1+cx}}{bc(a+b\operatorname{arccosh}(cx))} + \frac{d \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c} + \frac{e \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4b^2c^3} + \frac{3e \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b^2c^3} - \frac{d \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c} - \frac{e \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{4b^2c^3} - \frac{3e \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+b\operatorname{arccosh}(cx))}{b}\right)}{4b^2c^3}$$

output

```
d*Chi((a+b*arccosh(c*x))/b)*cosh(a/b)/b^2/c+1/4*e*Chi((a+b*arccosh(c*x))/b)
*cosh(a/b)/b^2/c^3+3/4*e*Chi(3*(a+b*arccosh(c*x))/b)*cosh(3*a/b)/b^2/c^3-
d*Shi((a+b*arccosh(c*x))/b)*sinh(a/b)/b^2/c-1/4*e*Shi((a+b*arccosh(c*x))/b)
*sinh(a/b)/b^2/c^3-3/4*e*Shi(3*(a+b*arccosh(c*x))/b)*sinh(3*a/b)/b^2/c^3-
d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(a+b*arccosh(c*x))-e*x^2*(c*x-1)^(1/2)*(
c*x+1)^(1/2)/b/c/(a+b*arccosh(c*x))
```



### 3.544.2 Mathematica [A] (warning: unable to verify)

Time = 1.00 (sec) , antiderivative size = 338, normalized size of antiderivative = 1.32

$$\int \frac{d + ex^2}{(a + \operatorname{barccosh}(cx))^2} dx = \frac{4bc^2d\sqrt{\frac{-1+cx}{1+cx}} + 4bc^3dx\sqrt{\frac{-1+cx}{1+cx}} + 4bc^2ex^2\sqrt{\frac{-1+cx}{1+cx}} + 4bc^3ex^3\sqrt{\frac{-1+cx}{1+cx}} - (4c^2d + e)(a + \operatorname{barccosh}(cx))}{\dots}$$

input `Integrate[(d + e*x^2)/(a + b*ArcCosh[c*x])^2,x]`

output `-1/4*(4*b*c^2*d*Sqrt[(-1 + c*x)/(1 + c*x)] + 4*b*c^3*d*x*Sqrt[(-1 + c*x)/(1 + c*x)] + 4*b*c^2*e*x^2*Sqrt[(-1 + c*x)/(1 + c*x)] + 4*b*c^3*e*x^3*Sqrt[(-1 + c*x)/(1 + c*x)] - (4*c^2*d + e)*(a + b*ArcCosh[c*x])*Cosh[a/b]*CoshIntegral[a/b + ArcCosh[c*x]] - 3*e*(a + b*ArcCosh[c*x])*Cosh[(3*a)/b]*CoshIntegral[3*(a/b + ArcCosh[c*x])] + 4*a*c^2*d*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + a*e*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + 4*b*c^2*d*ArcCosh[c*x]*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + b*e*ArcCosh[c*x]*Sinh[a/b]*SinhIntegral[a/b + ArcCosh[c*x]] + 3*a*e*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])] + 3*b*e*ArcCosh[c*x]*Sinh[(3*a)/b]*SinhIntegral[3*(a/b + ArcCosh[c*x])])/(b^2*c^3*(a + b*ArcCosh[c*x]))`

### 3.544.3 Rubi [A] (verified)

Time = 0.80 (sec) , antiderivative size = 257, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{d + ex^2}{(a + \operatorname{barccosh}(cx))^2} dx \xrightarrow{6324} \int \left( \frac{d}{(a + \operatorname{barccosh}(cx))^2} + \frac{ex^2}{(a + \operatorname{barccosh}(cx))^2} \right) dx \xrightarrow{2009}$$

$$\frac{e \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{4b^2c^3} + \frac{3e \cosh\left(\frac{3a}{b}\right) \operatorname{Chi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{4b^2c^3} -$$

$$\frac{e \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{4b^2c^3} - \frac{3e \sinh\left(\frac{3a}{b}\right) \operatorname{Shi}\left(\frac{3(a+\operatorname{barccosh}(cx))}{b}\right)}{4b^2c^3} +$$

$$\frac{d \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{b^2c} - \frac{d \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+\operatorname{barccosh}(cx)}{b}\right)}{b^2c} - \frac{d\sqrt{cx-1}\sqrt{cx+1}}{bc(a+\operatorname{barccosh}(cx))} -$$

$$\frac{ex^2\sqrt{cx-1}\sqrt{cx+1}}{bc(a+\operatorname{barccosh}(cx))}$$

input `Int[(d + e*x^2)/(a + b*ArcCosh[c*x])^2,x]`

output `-((d*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(b*c*(a + b*ArcCosh[c*x]))) - (e*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(b*c*(a + b*ArcCosh[c*x])) + (d*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/(b^2*c) + (e*Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b])/(4*b^2*c^3) + (3*e*Cosh[(3*a)/b]*CoshIntegral[(3*(a + b*ArcCosh[c*x])/b])/(4*b^2*c^3) - (d*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(b^2*c) - (e*Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(4*b^2*c^3) - (3*e*Sinh[(3*a)/b]*SinhIntegral[(3*(a + b*ArcCosh[c*x])/b])/(4*b^2*c^3)`

### 3.544.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

## 3.544.4 Maple [A] (verified)

Time = 1.12 (sec) , antiderivative size = 465, normalized size of antiderivative = 1.81

method	result
derivativedivides	$\frac{(-4\sqrt{cx-1}\sqrt{cx+1}c^2x^2+\sqrt{cx-1}\sqrt{cx+1}+4c^3x^3-3cx)e^{-3e\frac{3a}{b}}\text{Ei}_1\left(3\text{arccosh}(cx)+\frac{3a}{b}\right)-e(4c^3x^3-3cx+4\sqrt{cx-1}\sqrt{cx+1}c^2x^2-\sqrt{cx-1}\sqrt{cx+1})}{8c^2b(a+b\text{arccosh}(cx))} - \frac{3e\frac{3a}{b}\text{Ei}_1\left(3\text{arccosh}(cx)+\frac{3a}{b}\right)-e(4c^3x^3-3cx+4\sqrt{cx-1}\sqrt{cx+1}c^2x^2-\sqrt{cx-1}\sqrt{cx+1})}{8c^2b^2} - \frac{e(4c^3x^3-3cx+4\sqrt{cx-1}\sqrt{cx+1}c^2x^2-\sqrt{cx-1}\sqrt{cx+1})}{8c^2b(a+b\text{arccosh}(cx))}$
default	$\frac{(-4\sqrt{cx-1}\sqrt{cx+1}c^2x^2+\sqrt{cx-1}\sqrt{cx+1}+4c^3x^3-3cx)e^{-3e\frac{3a}{b}}\text{Ei}_1\left(3\text{arccosh}(cx)+\frac{3a}{b}\right)-e(4c^3x^3-3cx+4\sqrt{cx-1}\sqrt{cx+1}c^2x^2-\sqrt{cx-1}\sqrt{cx+1})}{8c^2b(a+b\text{arccosh}(cx))} - \frac{3e\frac{3a}{b}\text{Ei}_1\left(3\text{arccosh}(cx)+\frac{3a}{b}\right)-e(4c^3x^3-3cx+4\sqrt{cx-1}\sqrt{cx+1}c^2x^2-\sqrt{cx-1}\sqrt{cx+1})}{8c^2b^2} - \frac{e(4c^3x^3-3cx+4\sqrt{cx-1}\sqrt{cx+1}c^2x^2-\sqrt{cx-1}\sqrt{cx+1})}{8c^2b(a+b\text{arccosh}(cx))}$

```
input int((e*x^2+d)/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)
```

```
output 1/c*(1/8*(-4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2+(c*x-1)^(1/2)*(c*x+1)^(1/2)+4*c^3*x^3-3*c*x)*e/c^2/b/(a+b*arccosh(c*x))-3/8*e/c^2/b^2*exp(3*a/b)*Ei(1,3*arccosh(c*x)+3*a/b)-1/8*e/c^2/b*(4*c^3*x^3-3*c*x+4*(c*x-1)^(1/2)*(c*x+1)^(1/2)*c^2*x^2-(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))-3/8*e/c^2/b^2*exp(-3*a/b)*Ei(1,-3*arccosh(c*x)-3*a/b)+1/2*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*d/b/(a+b*arccosh(c*x))+1/8*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)*e/c^2/b/(a+b*arccosh(c*x))-1/2/b^2*exp(a/b)*Ei(1,arccosh(c*x)+a/b)*d-1/8/c^2/b^2*exp(a/b)*Ei(1,arccosh(c*x)+a/b)*e-1/2/b*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))*d-1/8/c^2/b*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))*e-1/2/b^2*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)*d-1/8/c^2/b^2*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b)*e)
```

## 3.544.5 Fracas [F]

$$\int \frac{d+ex^2}{(a+b\text{arccosh}(cx))^2} dx = \int \frac{ex^2+d}{(b\text{arccosh}(cx)+a)^2} dx$$

```
input integrate((e*x^2+d)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")
```

```
output integral((e*x^2 + d)/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)
```

**3.544.6 Sympy [F]**

$$\int \frac{d + ex^2}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{d + ex^2}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate((e*x**2+d)/(a+b*acosh(c*x))**2,x)`

output `Integral((d + e*x**2)/(a + b*acosh(c*x))**2, x)`

**3.544.7 Maxima [F]**

$$\int \frac{d + ex^2}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{ex^2 + d}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-(c^3*e*x^5 + (c^3*d - c*e)*x^3 - c*d*x + (c^2*e*x^4 + (c^2*d - e)*x^2 - d)*sqrt(c*x + 1)*sqrt(c*x - 1))/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate(((3*c^5*e*x^6 + (c^5*d - 6*c^3*e)*x^4 + (3*c^3*e*x^4 + (c^3*d - c*e)*x^2 + c*d)*(c*x + 1)*(c*x - 1) - (2*c^3*d - 3*c*e)*x^2 + (6*c^4*e*x^5 + (2*c^4*d - 7*c^2*e)*x^3 - (c^2*d - 2*e)*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + c*d)/(a*b*c^5*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^3*x^2 - 2*a*b*c^3*x^2 + a*b*c + 2*(a*b*c^4*x^3 - a*b*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^3*x^2 - 2*b^2*c^3*x^2 + b^2*c + 2*(b^2*c^4*x^3 - b^2*c^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.544.8 Giac [F]**

$$\int \frac{d + ex^2}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{ex^2 + d}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((e*x^2 + d)/(b*arccosh(c*x) + a)^2, x)`

**3.544.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{d + ex^2}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{ex^2 + d}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `int((d + e*x^2)/(a + b*acosh(c*x))^2, x)`output `int((d + e*x^2)/(a + b*acosh(c*x))^2, x)`

### 3.545 $\int \frac{1}{(a+b\operatorname{arccosh}(cx))^2} dx$

3.545.1 Optimal result . . . . .	3973
3.545.2 Mathematica [A] (verified) . . . . .	3973
3.545.3 Rubi [A] (verified) . . . . .	3974
3.545.4 Maple [A] (verified) . . . . .	3977
3.545.5 Fricas [F] . . . . .	3977
3.545.6 Sympy [F] . . . . .	3977
3.545.7 Maxima [F] . . . . .	3978
3.545.8 Giac [F] . . . . .	3978
3.545.9 Mupad [F(-1)] . . . . .	3978

#### 3.545.1 Optimal result

Integrand size = 10, antiderivative size = 90

$$\int \frac{1}{(a + b\operatorname{arccosh}(cx))^2} dx = -\frac{\sqrt{-1+cx}\sqrt{1+cx}}{bc(a + b\operatorname{arccosh}(cx))} + \frac{\cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c} - \frac{\sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c}$$

output `Chi((a+b*arccosh(c*x))/b)*cosh(a/b)/b^2/c-Shi((a+b*arccosh(c*x))/b)*sinh(a/b)/b^2/c-(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(a+b*arccosh(c*x))`

#### 3.545.2 Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.89

$$\int \frac{1}{(a + b\operatorname{arccosh}(cx))^2} dx = \frac{-\frac{b\sqrt{\frac{-1+cx}{1+cx}}(1+cx)}{a+b\operatorname{arccosh}(cx)} + \cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right) - \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a}{b} + \operatorname{arccosh}(cx)\right)}{b^2c}$$

input `Integrate[(a + b*ArcCosh[c*x])^(-2), x]`

output  $(-((b*\text{Sqrt}[(-1 + c*x)/(1 + c*x)]*(1 + c*x))/(a + b*\text{ArcCosh}[c*x])) + \text{Cosh}[a/b]*\text{CoshIntegral}[a/b + \text{ArcCosh}[c*x]] - \text{Sinh}[a/b]*\text{SinhIntegral}[a/b + \text{ArcCosh}[c*x]])/(b^2*c)$

### 3.545.3 Rubi [A] (verified)

Time = 0.77 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.96, number of steps used = 10, number of rules used = 9,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.900$ , Rules used = {6295, 6368, 3042, 3784, 26, 3042, 26, 3779, 3782}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + b \operatorname{arccosh}(cx))^2} dx \\
 & \quad \downarrow 6295 \\
 & \frac{c \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}(a+b \operatorname{arccosh}(cx))} dx}{b} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a + b \operatorname{arccosh}(cx))} \\
 & \quad \downarrow 6368 \\
 & \frac{\int \frac{\cosh\left(\frac{a}{b} - \frac{a+b \operatorname{arccosh}(cx)}{b}\right)}{a+b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{b^2 c} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a + b \operatorname{arccosh}(cx))} \\
 & \quad \downarrow 3042 \\
 & -\frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a + b \operatorname{arccosh}(cx))} + \frac{\int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+b \operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{b^2 c} \\
 & \quad \downarrow 3784 \\
 & -\frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a + b \operatorname{arccosh}(cx))} + \\
 & \frac{\cosh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right)}{a+b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx)) - i \sinh\left(\frac{a}{b}\right) \int -\frac{i \sinh\left(\frac{a+b \operatorname{arccosh}(cx)}{b}\right)}{a+b \operatorname{arccosh}(cx)} d(a + b \operatorname{arccosh}(cx))}{b^2 c} \\
 & \quad \downarrow 26
 \end{aligned}$$

$$\begin{aligned}
& \frac{\cosh\left(\frac{a}{b}\right) \int \frac{\cosh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - \sinh\left(\frac{a}{b}\right) \int \frac{\sinh\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c} \\
& \quad - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a+b\operatorname{arccosh}(cx))} \\
& \quad \downarrow \text{3042} \\
& \quad - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a+b\operatorname{arccosh}(cx))} + \\
& \quad \frac{\cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) - \sinh\left(\frac{a}{b}\right) \int \frac{i \sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c} \\
& \quad \downarrow \text{26} \\
& \quad - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a+b\operatorname{arccosh}(cx))} + \\
& \quad \frac{i \sinh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx)) + \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c} \\
& \quad \downarrow \text{3779} \\
& \quad - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a+b\operatorname{arccosh}(cx))} + \\
& \quad \frac{-\sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) + \cosh\left(\frac{a}{b}\right) \int \frac{\sin\left(\frac{i(a+b\operatorname{arccosh}(cx))}{b} + \frac{\pi}{2}\right)}{a+b\operatorname{arccosh}(cx)} d(a+b\operatorname{arccosh}(cx))}{b^2c} \\
& \quad \downarrow \text{3782} \\
& \quad \frac{\cosh\left(\frac{a}{b}\right) \operatorname{Chi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right) - \sinh\left(\frac{a}{b}\right) \operatorname{Shi}\left(\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{b^2c} - \frac{\sqrt{cx-1}\sqrt{cx+1}}{bc(a+b\operatorname{arccosh}(cx))}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])^(-2), x]`

output `-((Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(b*c*(a + b*ArcCosh[c*x]))) + (Cosh[a/b]*CoshIntegral[(a + b*ArcCosh[c*x])/b] - Sinh[a/b]*SinhIntegral[(a + b*ArcCosh[c*x])/b])/(b^2*c)`



## 3.545.3.1 Defintions of rubi rules used

- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3779 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]`
- rule 3782 `Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]`
- rule 3784 `Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[Cos[(d*e - c*f)/d] Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Simp[Sin[(d*e - c*f)/d] Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]`
- rule 6295 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_, x_Symbol] := Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*((a + b*ArcCosh[c*x])^(n + 1)/(b*c*(n + 1))), x] - Simp[c/(b*(n + 1)) Int[x*((a + b*ArcCosh[c*x])^(n + 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && LtQ[n, -1]`
- rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*(x_)^m_.*((d1_.) + (e1_.)*(x_))^p_.*((d2_.) + (e2_.)*(x_))^p_.), x_Symbol] := Simp[(1/(b*c^(m + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^p] Subst[Int[x^n*Cosh[-a/b + x/b]^m*Sinh[-a/b + x/b]^(2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

**3.545.4 Maple [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.39

method	result	size
derivativedivides	$\frac{-\sqrt{cx-1}\sqrt{cx+1}+cx}{2b(a+b \operatorname{arccosh}(cx))} - \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx)+\frac{a}{b}\right)}{2b^2} - \frac{cx+\sqrt{cx-1}\sqrt{cx+1}}{2b(a+b \operatorname{arccosh}(cx))} - \frac{e^{-\frac{a}{b}} \operatorname{Ei}_1\left(-\operatorname{arccosh}(cx)-\frac{a}{b}\right)}{2b^2}$	125
default	$\frac{-\sqrt{cx-1}\sqrt{cx+1}+cx}{2b(a+b \operatorname{arccosh}(cx))} - \frac{e^{\frac{a}{b}} \operatorname{Ei}_1\left(\operatorname{arccosh}(cx)+\frac{a}{b}\right)}{2b^2} - \frac{cx+\sqrt{cx-1}\sqrt{cx+1}}{2b(a+b \operatorname{arccosh}(cx))} - \frac{e^{-\frac{a}{b}} \operatorname{Ei}_1\left(-\operatorname{arccosh}(cx)-\frac{a}{b}\right)}{2b^2}$	125

input `int(1/(a+b*arccosh(c*x))^2,x,method=_RETURNVERBOSE)`output `1/c*(1/2*(-(c*x-1)^(1/2)*(c*x+1)^(1/2)+c*x)/b/(a+b*arccosh(c*x))-1/2/b^2*exp(a/b)*Ei(1,arccosh(c*x)+a/b)-1/2/b*(c*x+(c*x-1)^(1/2)*(c*x+1)^(1/2))/(a+b*arccosh(c*x))-1/2/b^2*exp(-a/b)*Ei(1,-arccosh(c*x)-a/b))`**3.545.5 Fracas [F]**

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(1/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`output `integral(1/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`**3.545.6 Sympy [F]**

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `integrate(1/(a+b*acosh(c*x))**2,x)`output `Integral((a + b*acosh(c*x))**(-2), x)`

**3.545.7 Maxima [F]**

$$\int \frac{1}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-(c^3*x^3 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*x)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate((c^4*x^4 - 2*c^2*x^2 + (c^2*x^2 + 1)*(c*x + 1)*(c*x - 1) + (2*c^3*x^3 - c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + 1)/(a*b*c^4*x^4 + (c*x + 1)*(c*x - 1)*a*b*c^2*x^2 - 2*a*b*c^2*x^2 + 2*(a*b*c^3*x^3 - a*b*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + a*b + (b^2*c^4*x^4 + (c*x + 1)*(c*x - 1)*b^2*c^2*x^2 - 2*b^2*c^2*x^2 + 2*(b^2*c^3*x^3 - b^2*c*x)*sqrt(c*x + 1)*sqrt(c*x - 1)) + b^2)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

**3.545.8 Giac [F]**

$$\int \frac{1}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^(-2), x)`

**3.545.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `int(1/(a + b*acosh(c*x))^2,x)`

output `int(1/(a + b*acosh(c*x))^2, x)`

$$3.546 \quad \int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^2} dx$$

3.546.1 Optimal result . . . . .	3979
3.546.2 Mathematica [N/A] . . . . .	3979
3.546.3 Rubi [N/A] . . . . .	3980
3.546.4 Maple [N/A] (verified) . . . . .	3980
3.546.5 Fricas [N/A] . . . . .	3981
3.546.6 Sympy [N/A] . . . . .	3981
3.546.7 Maxima [N/A] . . . . .	3981
3.546.8 Giac [N/A] . . . . .	3982
3.546.9 Mupad [N/A] . . . . .	3983

### 3.546.1 Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/(e*x^2+d)/(a+b*arccosh(c*x))^2,x)`

### 3.546.2 Mathematica [N/A]

Not integrable

Time = 17.95 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/((d + e*x^2)*(a + b*ArcCosh[c*x]))^2, x]`

output `Integrate[1/((d + e*x^2)*(a + b*ArcCosh[c*x]))^2, x]`

**3.546.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)(a + \operatorname{arccosh}(cx))^2} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)(a + \operatorname{arccosh}(cx))^2} dx$$

input `Int[1/((d + e*x^2)*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.546.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.546.4 Maple [N/A] (verified)**

Not integrable

Time = 0.90 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(ex^2 + d)(a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/(e*x^2+d)/(a+b*arccosh(c*x))^2,x)`

output `int(1/(e*x^2+d)/(a+b*arccosh(c*x))^2,x)`

**3.546.5 Fracas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 57, normalized size of antiderivative = 2.85

$$\int \frac{1}{(d + ex^2)(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`output `integral(1/(a^2*e*x^2 + a^2*d + (b^2*e*x^2 + b^2*d)*arccosh(c*x)^2 + 2*(a*b*e*x^2 + a*b*d)*arccosh(c*x)), x)`**3.546.6 Sympy [N/A]**

Not integrable

Time = 89.47 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(d + ex^2)(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (d + ex^2)} dx$$

input `integrate(1/(e*x**2+d)/(a+b*acosh(c*x))**2,x)`output `Integral(1/((a + b*acosh(c*x))**2*(d + e*x**2)), x)`**3.546.7 Maxima [N/A]**

Not integrable

Time = 1.32 (sec) , antiderivative size = 816, normalized size of antiderivative = 40.80

$$\int \frac{1}{(d + ex^2)(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output  $-(c^3x^3 + (c^2x^2 - 1)\sqrt{cx + 1}\sqrt{cx - 1} - cx)/(a^3b^3c^3e^4 + (c^3d - ce)a^2bx^2 - a^2bc^3d + (a^2bc^2e^3x^3 + a^2bc^2d^2x)\sqrt{cx + 1}\sqrt{cx - 1} + (b^2c^3e^4x^4 + (c^3d - ce)b^2x^2 - b^2c^3d + (b^2c^2e^3x^3 + b^2c^2d^2x)\sqrt{cx + 1}\sqrt{cx - 1}))\log(cx + \sqrt{cx + 1}\sqrt{cx - 1})) - \text{integrate}((c^5e^6x^6 - (c^5d + 2c^3e)x^4 + (c^3e^4x^4 - (c^3d + 3ce)x^2 - cd)(cx + 1)(cx - 1) + (2c^3d + ce)x^2 + (2c^4e^5x^5 - (2c^4d + 5c^2e)x^3 + (c^2d + 2e)x)\sqrt{cx + 1}\sqrt{cx - 1} - cd)/(a^5b^5c^5e^2x^8 + 2(c^5d^2e - c^3e^2)a^4bx^6 + (c^5d^2 - 4c^3d^2e + ce^2)a^3bx^4 + a^2bc^5d^2 - 2(c^3d^2 - cd^2e)a^2bx^2 + (a^2bc^3e^2x^6 + 2a^2bc^3d^2e^2x^4 + a^2bc^3d^2x^2)(cx + 1)(cx - 1) + 2(a^2bc^4e^2x^7 + (2c^4d^2e - c^2e^2)a^2bx^5 - a^2bc^2d^2x + (c^4d^2 - 2c^2d^2e)a^2bx^3)\sqrt{cx + 1}\sqrt{cx - 1} + (b^2c^5e^2x^8 + 2(c^5d^2e - c^3e^2)b^2x^6 + (c^5d^2 - 4c^3d^2e + ce^2)b^2x^4 + b^2c^5d^2 - 2(c^3d^2 - cd^2e)b^2x^2 + (b^2c^3e^2x^6 + 2b^2c^3d^2e^2x^4 + b^2c^3d^2x^2)(cx + 1)(cx - 1) + 2(b^2c^4e^2x^7 + (2c^4d^2e - c^2e^2)b^2x^5 - b^2c^2d^2x + (c^4d^2 - 2c^2d^2e)b^2x^3)\sqrt{cx + 1}\sqrt{cx - 1}))\log(cx + \sqrt{cx + 1}\sqrt{cx - 1})), x)$

### 3.546.8 Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(1/((e*x^2 + d)*(b*arccosh(c*x) + a)^2), x)`

**3.546.9 Mupad [N/A]**

Not integrable

Time = 2.96 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (ex^2 + d)} dx$$

input `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)),x)`output `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)), x)`



**3.547**  $\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))^2} dx$

3.547.1 Optimal result . . . . .	3984
3.547.2 Mathematica [N/A] . . . . .	3984
3.547.3 Rubi [N/A] . . . . .	3985
3.547.4 Maple [N/A] (verified) . . . . .	3985
3.547.5 Fricas [N/A] . . . . .	3986
3.547.6 Sympy [F(-1)] . . . . .	3986
3.547.7 Maxima [N/A] . . . . .	3986
3.547.8 Giac [N/A] . . . . .	3987
3.547.9 Mupad [N/A] . . . . .	3988

**3.547.1 Optimal result**

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^2,x)`

**3.547.2 Mathematica [N/A]**

Not integrable

Time = 29.87 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])^2), x]`

**3.547.3 Rubi [N/A]**

Not integrable

Time = 0.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^2} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^2} dx$$

input `Int[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.547.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.547.4 Maple [N/A] (verified)**

Not integrable

Time = 0.66 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(ex^2 + d)^2 (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^2,x)`

output `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^2,x)`

**3.547.5 Fricas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 98, normalized size of antiderivative = 4.90

$$\int \frac{1}{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^2 (b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(1/(a^2*e^2*x^4 + 2*a^2*d*e*x^2 + a^2*d^2 + (b^2*e^2*x^4 + 2*b^2*d*e*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*e^2*x^4 + 2*a*b*d*e*x^2 + a*b*d^2)*arccosh(c*x)), x)`

**3.547.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))^2} dx = \text{Timed out}$$

input `integrate(1/(e*x**2+d)**2/(a+b*acosh(c*x))**2,x)`

output `Timed out`

**3.547.7 Maxima [N/A]**

Not integrable

Time = 1.77 (sec) , antiderivative size = 1078, normalized size of antiderivative = 53.90

$$\int \frac{1}{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^2 (b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`



**3.547.9 Mupad [N/A]**

Not integrable

Time = 2.90 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (ex^2 + d)^2} dx$$

input `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^2), x)`output `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^2), x)`

$$3.548 \quad \int \frac{\sqrt{d+ex^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

3.548.1 Optimal result	3989
3.548.2 Mathematica [N/A]	3989
3.548.3 Rubi [N/A]	3990
3.548.4 Maple [N/A] (verified)	3990
3.548.5 Fricas [N/A]	3991
3.548.6 Sympy [N/A]	3991
3.548.7 Maxima [N/A]	3991
3.548.8 Giac [N/A]	3992
3.548.9 Mupad [N/A]	3992

### 3.548.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{\sqrt{d+ex^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{\sqrt{d+ex^2}}{(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable((e*x^2+d)^(1/2)/(a+b*arccosh(c*x))^2,x)`

### 3.548.2 Mathematica [N/A]

Not integrable

Time = 10.79 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{d+ex^2}}{(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{d+ex^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[Sqrt[d + e*x^2]/(a + b*ArcCosh[c*x])^2,x]`

output `Integrate[Sqrt[d + e*x^2]/(a + b*ArcCosh[c*x])^2, x]`

**3.548.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{d+ex^2}}{(a+\operatorname{arccosh}(cx))^2} dx$$

↓ 6325

$$\int \frac{\sqrt{d+ex^2}}{(a+\operatorname{arccosh}(cx))^2} dx$$

input `Int[Sqrt[d + e*x^2]/(a + b*ArcCosh[c*x])^2,x]`

output `$Aborted`

**3.548.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.548.4 Maple [N/A] (verified)**

Not integrable

Time = 1.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{ex^2+d}}{(a+b \operatorname{arccosh}(cx))^2} dx$$

input `int((e*x^2+d)^(1/2)/(a+b*arccosh(c*x))^2,x)`

output `int((e*x^2+d)^(1/2)/(a+b*arccosh(c*x))^2,x)`

**3.548.5 Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.64

$$\int \frac{\sqrt{d+ex^2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{ex^2+d}}{(b \operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate((e*x^2+d)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`output `integral(sqrt(e*x^2 + d)/(b^2*arccosh(c*x)^2 + 2*a*b*arccosh(c*x) + a^2), x)`**3.548.6 Sympy [N/A]**

Not integrable

Time = 0.87 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{d+ex^2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{d+ex^2}}{(a+b \operatorname{acosh}(cx))^2} dx$$

input `integrate((e*x**2+d)**(1/2)/(a+b*acosh(c*x))**2,x)`output `Integral(sqrt(d + e*x**2)/(a + b*acosh(c*x))**2, x)`**3.548.7 Maxima [N/A]**

Not integrable

Time = 0.81 (sec) , antiderivative size = 596, normalized size of antiderivative = 27.09

$$\int \frac{\sqrt{d+ex^2}}{(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{\sqrt{ex^2+d}}{(b \operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate((e*x^2+d)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`



output `-(c^3*x^3 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*x)*sqrt(e*x^2 + d)/(a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*a*b*c^2*x - a*b*c + (b^2*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))) + integrate((2*c^5*e*x^6 + (c^5*d - 4*c^3*e)*x^4 + (2*c^3*e*x^4 + c^3*d*x^2 + c*d)*(c*x + 1)*(c*x - 1) - 2*(c^3*d - c*e)*x^2 + (4*c^4*e*x^5 + 2*(c^4*d - 2*c^2*e)*x^3 - (c^2*d - e)*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + c*d)*sqrt(e*x^2 + d)/(a*b*c^5*e*x^6 + (c^5*d - 2*c^3*e)*a*b*x^4 - (2*c^3*d - c*e)*a*b*x^2 + a*b*c*d + (a*b*c^3*e*x^4 + a*b*c^3*d*x^2)*(c*x + 1)*(c*x - 1) + 2*(a*b*c^4*e*x^5 - a*b*c^2*d*x + (c^4*d - c^2*e)*a*b*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1) + (b^2*c^5*e*x^6 + (c^5*d - 2*c^3*e)*b^2*x^4 - (2*c^3*d - c*e)*b^2*x^2 + b^2*c*d + (b^2*c^3*e*x^4 + b^2*c^3*d*x^2)*(c*x + 1)*(c*x - 1) + 2*(b^2*c^4*e*x^5 - b^2*c^2*d*x + (c^4*d - c^2*e)*b^2*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1))*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1))), x)`

### 3.548.8 Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d + ex^2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{ex^2 + d}}{(b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate((e*x^2+d)^(1/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(sqrt(e*x^2 + d)/(b*arccosh(c*x) + a)^2, x)`

### 3.548.9 Mupad [N/A]

Not integrable

Time = 3.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{d + ex^2}}{(a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{\sqrt{ex^2 + d}}{(a + b \operatorname{acosh}(cx))^2} dx$$

input `int((d + e*x^2)^(1/2)/(a + b*acosh(c*x))^2,x)`

output `int((d + e*x^2)^(1/2)/(a + b*acosh(c*x))^2, x)`

---

3.548.  $\int \frac{\sqrt{d+ex^2}}{(a+b\operatorname{arccosh}(cx))^2} dx$

$$3.549 \quad \int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

3.549.1 Optimal result	3993
3.549.2 Mathematica [N/A]	3993
3.549.3 Rubi [N/A]	3994
3.549.4 Maple [N/A] (verified)	3994
3.549.5 Fricas [N/A]	3995
3.549.6 Sympy [N/A]	3995
3.549.7 Maxima [N/A]	3995
3.549.8 Giac [N/A]	3996
3.549.9 Mupad [N/A]	3996

### 3.549.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/(a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x)`

### 3.549.2 Mathematica [N/A]

Not integrable

Time = 10.97 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/(Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])^2),x]`

output `Integrate[1/(Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])^2), x]`

**3.549.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{d+ex^2}(a+\operatorname{arccosh}(cx))^2} dx$$

↓ 6325

$$\int \frac{1}{\sqrt{d+ex^2}(a+\operatorname{arccosh}(cx))^2} dx$$

input `Int[1/(Sqrt[d + e*x^2]*(a + b*ArcCosh[c*x])^2), x]`

output `$Aborted`

**3.549.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_.*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.549.4 Maple [N/A] (verified)**

Not integrable

Time = 1.31 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(a+b \operatorname{arccosh}(cx))^2 \sqrt{ex^2+d}} dx$$

input `int(1/(a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2), x)`

output `int(1/(a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2), x)`

**3.549.5 Fricas [N/A]**

Not integrable

Time = 0.30 (sec) , antiderivative size = 67, normalized size of antiderivative = 3.05

$$\int \frac{1}{\sqrt{d+ex^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{\sqrt{ex^2+d}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(1/(a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)/(a^2*e*x^2 + a^2*d + (b^2*e*x^2 + b^2*d)*arccosh(c*x)^2 + 2*(a*b*e*x^2 + a*b*d)*arccosh(c*x)), x)`

**3.549.6 Sympy [N/A]**

Not integrable

Time = 1.98 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d+ex^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a+b\operatorname{acosh}(cx))^2\sqrt{d+ex^2}} dx$$

input `integrate(1/(a+b*acosh(c*x))**2/(e*x**2+d)**(1/2),x)`

output `Integral(1/((a + b*acosh(c*x))**2*sqrt(d + e*x**2)), x)`

**3.549.7 Maxima [N/A]**

Not integrable

Time = 1.10 (sec) , antiderivative size = 580, normalized size of antiderivative = 26.36

$$\int \frac{1}{\sqrt{d+ex^2}(a+\operatorname{barccosh}(cx))^2} dx = \int \frac{1}{\sqrt{ex^2+d}(b\operatorname{arcosh}(cx)+a)^2} dx$$

input `integrate(1/(a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x, algorithm="maxima")`

```
output -(c^3*x^3 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*x)/((b^2*c^3*x^2
+ sqrt(c*x + 1)*sqrt(c*x - 1)*b^2*c^2*x - b^2*c)*sqrt(e*x^2 + d)*log(c*x
+ sqrt(c*x + 1)*sqrt(c*x - 1)) + (a*b*c^3*x^2 + sqrt(c*x + 1)*sqrt(c*x - 1
)*a*b*c^2*x - a*b*c)*sqrt(e*x^2 + d)) + integrate((c^5*d*x^4 - 2*c^3*d*x^2
+ ((c^3*d + 2*c*e)*x^2 + c*d)*(c*x + 1)*(c*x - 1) + (2*(c^4*d + c^2*e)*x^
3 - (c^2*d + e)*x)*sqrt(c*x + 1)*sqrt(c*x - 1) + c*d)/((b^2*c^5*e*x^6 + (c
^5*d - 2*c^3*e)*b^2*x^4 - (2*c^3*d - c*e)*b^2*x^2 + b^2*c*d + (b^2*c^3*e*x
^4 + b^2*c^3*d*x^2)*(c*x + 1)*(c*x - 1) + 2*(b^2*c^4*e*x^5 - b^2*c^2*d*x +
(c^4*d - c^2*e)*b^2*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1))*sqrt(e*x^2 + d)*log
(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + (a*b*c^5*e*x^6 + (c^5*d - 2*c^3*e)*a
*b*x^4 - (2*c^3*d - c*e)*a*b*x^2 + a*b*c*d + (a*b*c^3*e*x^4 + a*b*c^3*d*x^
2)*(c*x + 1)*(c*x - 1) + 2*(a*b*c^4*e*x^5 - a*b*c^2*d*x + (c^4*d - c^2*e)*
a*b*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1))*sqrt(e*x^2 + d)), x)
```

### 3.549.8 Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{\sqrt{ex^2 + d}(b \operatorname{arcosh}(cx) + a)^2} dx$$

```
input integrate(1/(a+b*arccosh(c*x))^2/(e*x^2+d)^(1/2),x, algorithm="giac")
```

```
output integrate(1/(sqrt(e*x^2 + d)*(b*arccosh(c*x) + a)^2), x)
```

### 3.549.9 Mupad [N/A]

Not integrable

Time = 3.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{d + ex^2}(a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 \sqrt{ex^2 + d}} dx$$

```
input int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^(1/2)),x)
```

```
output int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^(1/2)), x)
```

---

3.549.  $\int \frac{1}{\sqrt{d+ex^2}(a+b\operatorname{arccosh}(cx))^2} dx$

**3.550**  $\int \frac{1}{(d+ex^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.550.1 Optimal result	3997
3.550.2 Mathematica [N/A]	3997
3.550.3 Rubi [N/A]	3998
3.550.4 Maple [N/A] (verified)	3998
3.550.5 Fracas [N/A]	3999
3.550.6 Sympy [N/A]	3999
3.550.7 Maxima [N/A]	3999
3.550.8 Giac [N/A]	4000
3.550.9 Mupad [N/A]	4001

**3.550.1 Optimal result**

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d+ex^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x))^2,x)`

**3.550.2 Mathematica [N/A]**

Not integrable

Time = 16.34 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d+ex^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(d+ex^2)^{3/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/((d + e*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/((d + e*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.550.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `Int[1/((d + e*x^2)^(3/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.550.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.550.4 Maple [N/A] (verified)**

Not integrable

Time = 1.52 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x))^2,x)`

output `int(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x))^2,x)`

**3.550.5 Fracas [N/A]**

Not integrable

Time = 0.26 (sec) , antiderivative size = 108, normalized size of antiderivative = 4.91

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)/(a^2*e^2*x^4 + 2*a^2*d*e*x^2 + a^2*d^2 + (b^2*e^2*x^4 + 2*b^2*d*e*x^2 + b^2*d^2)*arccosh(c*x)^2 + 2*(a*b*e^2*x^4 + 2*a*b*d*e*x^2 + a*b*d^2)*arccosh(c*x)), x)`

**3.550.6 Sympy [N/A]**

Not integrable

Time = 15.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (d + ex^2)^{\frac{3}{2}}} dx$$

input `integrate(1/(e*x**2+d)**(3/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(1/((a + b*acosh(c*x))**2*(d + e*x**2)**(3/2)), x)`

**3.550.7 Maxima [N/A]**

Not integrable

Time = 1.99 (sec) , antiderivative size = 857, normalized size of antiderivative = 38.95

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$



input `integrate(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output 
$$-(c^3x^3 + (c^2x^2 - 1)\sqrt{cx + 1}\sqrt{cx - 1} - cx)/((b^2c^3e^x^4 + (c^3d - ce)b^2x^2 - b^2cd + (b^2c^2e^x^3 + b^2c^2dx)\sqrt{cx + 1}\sqrt{cx - 1})\sqrt{e^x^2 + d}\log(cx + \sqrt{cx + 1}\sqrt{cx - 1})) + (a^2b^2c^3e^x^4 + (c^3d - ce)a^2bx^2 - a^2cd + (a^2b^2c^2e^x^3 + a^2b^2c^2dx)\sqrt{cx + 1}\sqrt{cx - 1})\sqrt{e^x^2 + d} - \int (2c^5e^x^6 - (c^5d + 4c^3e)x^4 + (2c^3e^x^4 - (c^3d + 4ce)x^2 - cd)(cx + 1)(cx - 1) + 2(c^3d + ce)x^2 + (4c^4e^x^5 - 2(c^4d + 4c^2e)x^3 + (c^2d + 3e)x)\sqrt{cx + 1}\sqrt{cx - 1} - cd)/((b^2c^5e^2x^8 + 2(c^5de - c^3e^2)b^2x^6 + (c^5d^2 - 4c^3de + ce^2)b^2x^4 + b^2cd^2 - 2(c^3d^2 - cde)b^2x^2 + (b^2c^3e^2x^6 + 2b^2c^3de^x^4 + b^2c^3d^2x^2)(cx + 1)(cx - 1) + 2(b^2c^4e^2x^7 + (2c^4de - c^2e^2)b^2x^5 - b^2c^2d^2x + (c^4d^2 - 2c^2de)b^2x^3)\sqrt{cx + 1}\sqrt{cx - 1})\sqrt{e^x^2 + d}\log(cx + \sqrt{cx + 1}\sqrt{cx - 1})) + (a^2b^2c^5e^2x^8 + 2(c^5de - c^3e^2)a^2bx^6 + (c^5d^2 - 4c^3de + ce^2)a^2bx^4 + a^2cd^2 - 2(c^3d^2 - cde)a^2bx^2 + (a^2b^2c^3e^2x^6 + 2a^2b^2c^3de^x^4 + a^2b^2c^3d^2x^2)(cx + 1)(cx - 1) + 2(a^2b^2c^4e^2x^7 + (2c^4de - c^2e^2)a^2bx^5 - a^2b^2c^2d^2x + (c^4d^2 - 2c^2de)a^2bx^3)\sqrt{cx + 1}\sqrt{cx - 1})\sqrt{e^x^2 + d}), x)$$

### 3.550.8 Giac [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{3/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^{\frac{3}{2}} (b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)^(3/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(1/((e*x^2 + d)^(3/2)*(b*arccosh(c*x) + a)^2), x)`

**3.550.9 Mupad [N/A]**

Not integrable

Time = 3.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{3/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (ex^2 + d)^{3/2}} dx$$

input `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^(3/2)),x)`output `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^(3/2)), x)`

**3.551**  $\int \frac{1}{(d+ex^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$

3.551.1 Optimal result . . . . .	4002
3.551.2 Mathematica [N/A] . . . . .	4002
3.551.3 Rubi [N/A] . . . . .	4003
3.551.4 Maple [N/A] (verified) . . . . .	4003
3.551.5 Fricas [N/A] . . . . .	4004
3.551.6 Sympy [N/A] . . . . .	4004
3.551.7 Maxima [N/A] . . . . .	4004
3.551.8 Giac [N/A] . . . . .	4005
3.551.9 Mupad [N/A] . . . . .	4006

**3.551.1 Optimal result**

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d+ex^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.551.2 Mathematica [N/A]**

Not integrable

Time = 24.97 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d+ex^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(d+ex^2)^{5/2}(a+b\operatorname{arccosh}(cx))^2} dx$$

input `Integrate[1/((d + e*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

output `Integrate[1/((d + e*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2), x]`

**3.551.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `Int[1/((d + e*x^2)^(5/2)*(a + b*ArcCosh[c*x])^2),x]`

output `$Aborted`

**3.551.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.551.4 Maple [N/A] (verified)**

Not integrable

Time = 1.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx$$

input `int(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x))^2,x)`

output `int(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x))^2,x)`

**3.551.5 Fracas [N/A]**

Not integrable

Time = 0.27 (sec) , antiderivative size = 149, normalized size of antiderivative = 6.77

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="fricas")`

output `integral(sqrt(e*x^2 + d)/(a^2*e^3*x^6 + 3*a^2*d*e^2*x^4 + 3*a^2*d^2*e*x^2 + a^2*d^3 + (b^2*e^3*x^6 + 3*b^2*d*e^2*x^4 + 3*b^2*d^2*e*x^2 + b^2*d^3)*arccosh(c*x)^2 + 2*(a*b*e^3*x^6 + 3*a*b*d*e^2*x^4 + 3*a*b*d^2*e*x^2 + a*b*d^3)*arccosh(c*x)), x)`

**3.551.6 Sympy [N/A]**

Not integrable

Time = 170.44 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (d + ex^2)^{\frac{5}{2}}} dx$$

input `integrate(1/(e*x**2+d)**(5/2)/(a+b*acosh(c*x))**2,x)`

output `Integral(1/((a + b*acosh(c*x))**2*(d + e*x**2)**(5/2)), x)`

**3.551.7 Maxima [N/A]**

Not integrable

Time = 2.98 (sec) , antiderivative size = 1117, normalized size of antiderivative = 50.77

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^{\frac{5}{2}} (b \operatorname{arcosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="maxima")`

output `-(c^3*x^3 + (c^2*x^2 - 1)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*x)/((b^2*c^3*e^2*x^6 + (2*c^3*d*e - c*e^2)*b^2*x^4 - b^2*c*d^2 + (c^3*d^2 - 2*c*d*e)*b^2*x^2 + (b^2*c^2*e^2*x^5 + 2*b^2*c^2*d*e*x^3 + b^2*c^2*d^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*sqrt(e*x^2 + d)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + (a*b*c^3*e^2*x^6 + (2*c^3*d*e - c*e^2)*a*b*x^4 - a*b*c*d^2 + (c^3*d^2 - 2*c*d*e)*a*b*x^2 + (a*b*c^2*e^2*x^5 + 2*a*b*c^2*d*e*x^3 + a*b*c^2*d^2*x)*sqrt(c*x + 1)*sqrt(c*x - 1))*sqrt(e*x^2 + d) - integrate((4*c^5*e*x^6 - (c^5*d + 8*c^3*e)*x^4 + (4*c^3*e*x^4 - (c^3*d + 6*c*e)*x^2 - c*d)*(c*x + 1)*(c*x - 1) + 2*(c^3*d + 2*c*e)*x^2 + (8*c^4*e*x^5 - 2*(c^4*d + 7*c^2*e)*x^3 + (c^2*d + 5*e)*x)*sqrt(c*x + 1)*sqrt(c*x - 1) - c*d)/((b^2*c^5*e^3*x^10 + (3*c^5*d*e^2 - 2*c^3*e^3)*b^2*x^8 + (3*c^5*d^2*e - 6*c^3*d*e^2 + c*e^3)*b^2*x^6 + (c^5*d^3 - 6*c^3*d^2*e + 3*c*d*e^2)*b^2*x^4 + b^2*c*d^3 - (2*c^3*d^3 - 3*c*d^2*e)*b^2*x^2 + (b^2*c^3*e^3*x^8 + 3*b^2*c^3*d*e^2*x^6 + 3*b^2*c^3*d^2*e*x^4 + b^2*c^3*d^3*x^2)*(c*x + 1)*(c*x - 1) + 2*(b^2*c^4*e^3*x^9 + (3*c^4*d*e^2 - c^2*e^3)*b^2*x^7 - b^2*c^2*d^3*x + 3*(c^4*d^2*e - c^2*d*e^2)*b^2*x^5 + (c^4*d^3 - 3*c^2*d^2*e)*b^2*x^3)*sqrt(c*x + 1)*sqrt(c*x - 1))*sqrt(e*x^2 + d)*log(c*x + sqrt(c*x + 1)*sqrt(c*x - 1)) + (a*b*c^5*e^3*x^10 + (3*c^5*d*e^2 - 2*c^3*e^3)*a*b*x^8 + (3*c^5*d^2*e - 6*c^3*d*e^2 + c*e^3)*a*b*x^6 + (c^5*d^3 - 6*c^3*d^2*e + 3*c*d*e^2)*a*b*x^4 + a*b*c*d^3 - (2*c^3*d^3 - 3*c*d^2*e)*a*b*x^2 + (a*b*c^3*e^3*x^8 + 3*a*b*c^3*d*e^2*x^6 + 3*a...`

### 3.551.8 Giac [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{5/2} (a + b \operatorname{arccosh}(cx))^2} dx = \int \frac{1}{(ex^2 + d)^{5/2} (b \operatorname{arccosh}(cx) + a)^2} dx$$

input `integrate(1/(e*x^2+d)^(5/2)/(a+b*arccosh(c*x))^2,x, algorithm="giac")`

output `integrate(1/((e*x^2 + d)^(5/2)*(b*arccosh(c*x) + a)^2), x)`

**3.551.9 Mupad [N/A]**

Not integrable

Time = 3.41 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^{5/2} (a + \operatorname{barccosh}(cx))^2} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^2 (ex^2 + d)^{5/2}} dx$$

input `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^(5/2)),x)`output `int(1/((a + b*acosh(c*x))^2*(d + e*x^2)^(5/2)), x)`

### 3.552 $\int (d + ex^2)^2 \sqrt{a + \operatorname{barccosh}(cx)} dx$

3.552.1 Optimal result . . . . .	4008
3.552.2 Mathematica [A] (verified) . . . . .	4009
3.552.3 Rubi [A] (verified) . . . . .	4010
3.552.4 Maple [F] . . . . .	4012
3.552.5 Fricas [F(-2)] . . . . .	4012
3.552.6 Sympy [F] . . . . .	4013
3.552.7 Maxima [F] . . . . .	4013
3.552.8 Giac [F] . . . . .	4013
3.552.9 Mupad [F(-1)] . . . . .	4014



## 3.552.1 Optimal result

Integrand size = 22, antiderivative size = 672

$$\begin{aligned}
\int (d + ex^2)^2 \sqrt{a + \operatorname{barccosh}(cx)} dx &= d^2x \sqrt{a + \operatorname{barccosh}(cx)} + \frac{2}{3} dex^3 \sqrt{a + \operatorname{barccosh}(cx)} \\
&+ \frac{1}{5} e^2 x^5 \sqrt{a + \operatorname{barccosh}(cx)} \\
&\quad - \frac{\sqrt{bd^2} e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{4c} \\
&\quad - \frac{\sqrt{bde} e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c^3} \\
&\quad - \frac{\sqrt{be^2} e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{32c^5} \\
&\quad - \frac{\sqrt{bde} e^{\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erf}\left(\frac{\sqrt{3} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{24c^3} \\
&\quad - \frac{\sqrt{be^2} e^{\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erf}\left(\frac{\sqrt{3} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{64c^5} \\
&\quad - \frac{\sqrt{be^2} e^{\frac{5a}{b}} \sqrt{\frac{\pi}{5}} \operatorname{erf}\left(\frac{\sqrt{5} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{320c^5} \\
&\quad - \frac{\sqrt{bd^2} e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{4c} \\
&\quad - \frac{\sqrt{bde} e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c^3} \\
&\quad - \frac{\sqrt{be^2} e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{32c^5} \\
&\quad - \frac{\sqrt{bde} e^{-\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erfi}\left(\frac{\sqrt{3} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{24c^3} \\
&\quad - \frac{\sqrt{be^2} e^{-\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erfi}\left(\frac{\sqrt{3} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{64c^5} \\
&\quad - \frac{\sqrt{be^2} e^{-\frac{5a}{b}} \sqrt{\frac{\pi}{5}} \operatorname{erfi}\left(\frac{\sqrt{5} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{320c^5}
\end{aligned}$$

output

$$\begin{aligned}
& -1/1600*e^2*\exp(5*a/b)*\operatorname{erf}(5^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2} \\
& *5^{1/2}*Pi^{1/2}/c^5-1/1600*e^2*\operatorname{erfi}(5^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b \\
& ^{1/2})*b^{1/2}*5^{1/2}*Pi^{1/2}/c^5/\exp(5*a/b)-1/72*d*e*\exp(3*a/b)*\operatorname{erf}(3^{1/2} \\
& *(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*3^{1/2}*Pi^{1/2}/c^3-1/192 \\
& *e^2*\exp(3*a/b)*\operatorname{erf}(3^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*3^{1/2} \\
& *Pi^{1/2}/c^5-1/72*d*e*\operatorname{erfi}(3^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b \\
& ^{1/2}*3^{1/2}*Pi^{1/2}/c^3/\exp(3*a/b)-1/192*e^2*\operatorname{erfi}(3^{1/2}*(a+b*\operatorname{arccosh} \\
& (c*x))^{1/2}/b^{1/2})*b^{1/2}*3^{1/2}*Pi^{1/2}/c^5/\exp(3*a/b)-1/4*d^2*\exp \\
& (a/b)*\operatorname{erf}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*Pi^{1/2}/c-1/8*d*e*\exp \\
& (a/b)*\operatorname{erf}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*Pi^{1/2}/c^3-1/32*e^2*e \\
& xp(a/b)*\operatorname{erf}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*Pi^{1/2}/c^5-1/4*d^2 \\
& *\operatorname{erfi}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*Pi^{1/2}/c/\exp(a/b)-1/8*d* \\
& e*\operatorname{erfi}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*Pi^{1/2}/c^3/\exp(a/b)-1/3 \\
& 2*e^2*\operatorname{erfi}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*b^{1/2}*Pi^{1/2}/c^5/\exp(a/b) \\
& +d^2*x*(a+b*\operatorname{arccosh}(c*x))^{1/2}+2/3*d*e*x^3*(a+b*\operatorname{arccosh}(c*x))^{1/2}+1/5*e \\
& ^2*x^5*(a+b*\operatorname{arccosh}(c*x))^{1/2}
\end{aligned}$$

### 3.552.2 Mathematica [A] (verified)

Time = 4.91 (sec) , antiderivative size = 536, normalized size of antiderivative = 0.80

$$\begin{aligned}
& \int (d + ex^2)^2 \sqrt{a + b \operatorname{arccosh}(cx)} dx \\
& = \frac{be^{-\frac{5a}{b}} \left( 450e^{\frac{6a}{b}} \left( 8ac^4d^2 \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} + 8bc^4d^2 \operatorname{arccosh}(cx) \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} - be(4c^2d + e) \sqrt{-\frac{a+b}{b}} \right) \right)}{
\end{aligned}$$

input `Integrate[(d + e*x^2)^2*Sqrt[a + b*ArcCosh[c*x]], x]`

output  $(b*(450*E^{((6*a)/b)}*(8*a*c^4*d^2*\text{Sqrt}[a/b + \text{ArcCosh}[c*x]] + 8*b*c^4*d^2*\text{ArcCosh}[c*x]*\text{Sqrt}[a/b + \text{ArcCosh}[c*x]] - b*e*(4*c^2*d + e)*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])/b])*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])^2/b^2])*\text{Gamma}[3/2, a/b + \text{ArcCosh}[c*x]] - 9*\text{Sqrt}[5]*b*e^2*\text{Sqrt}[a/b + \text{ArcCosh}[c*x]]*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])^2/b^2])*\text{Gamma}[3/2, (-5*(a + b*\text{ArcCosh}[c*x]))/b] - E^{((2*a)/b)}*(25*\text{Sqrt}[3]*b*e*(8*c^2*d + 3*e)*\text{Sqrt}[a/b + \text{ArcCosh}[c*x]]*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])^2/b^2])*\text{Gamma}[3/2, (-3*(a + b*\text{ArcCosh}[c*x]))/b] + 450*E^{((2*a)/b)}*(8*a*c^4*d^2*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])/b] + 8*b*c^4*d^2*\text{ArcCosh}[c*x]*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])/b] + b*e*(4*c^2*d + e)*\text{Sqrt}[a/b + \text{ArcCosh}[c*x]]*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])^2/b^2])*\text{Gamma}[3/2, -(a + b*\text{ArcCosh}[c*x])/b] + b*e*E^{((6*a)/b)}*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])/b]*\text{Sqrt}[-(a + b*\text{ArcCosh}[c*x])^2/b^2])*(25*\text{Sqrt}[3]*(8*c^2*d + 3*e)*\text{Gamma}[3/2, (3*(a + b*\text{ArcCosh}[c*x]))/b] + 9*\text{Sqrt}[5]*e*E^{((2*a)/b)}*\text{Gamma}[3/2, (5*(a + b*\text{ArcCosh}[c*x]))/b])))/(7200*c^5*E^{((5*a)/b)}*(a + b*\text{ArcCosh}[c*x])^{(3/2)})$

### 3.552.3 Rubi [A] (verified)

Time = 2.67 (sec) , antiderivative size = 672, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2)^2 \sqrt{a + \text{barccosh}(cx)} dx$$

$$\downarrow 6324$$

$$\int (d^2 \sqrt{a + \text{barccosh}(cx)} + 2dex^2 \sqrt{a + \text{barccosh}(cx)} + e^2 x^4 \sqrt{a + \text{barccosh}(cx)}) dx$$

$$\downarrow 2009$$

$$\begin{aligned}
& \frac{\sqrt{\pi}\sqrt{b}e^2e^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{32c^5} - \frac{\sqrt{\frac{\pi}{3}}\sqrt{b}e^2e^{\frac{3a}{b}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{64c^5} \\
& \frac{\sqrt{\frac{\pi}{5}}\sqrt{b}e^2e^{\frac{5a}{b}}\operatorname{erfi}\left(\frac{\sqrt{5}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{320c^5} - \frac{\sqrt{\pi}\sqrt{b}e^2e^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{64c^5} \\
& \frac{\sqrt{\frac{\pi}{3}}\sqrt{b}e^2e^{-\frac{3a}{b}}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{64c^5} - \frac{\sqrt{\frac{\pi}{5}}\sqrt{b}e^2e^{-\frac{5a}{b}}\operatorname{erfi}\left(\frac{\sqrt{5}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{320c^5} \\
& \frac{\sqrt{\pi}\sqrt{b}dee^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c^3} - \frac{\sqrt{\frac{\pi}{3}}\sqrt{b}dee^{\frac{3a}{b}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{24c^3} \\
& \frac{\sqrt{\pi}\sqrt{b}dee^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c^3} - \frac{\sqrt{\frac{\pi}{3}}\sqrt{b}dee^{-\frac{3a}{b}}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{24c^3} \\
& \frac{\sqrt{\pi}\sqrt{b}d^2e^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{4c} - \frac{\sqrt{\pi}\sqrt{b}d^2e^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{4c} + \\
& d^2x\sqrt{a+\operatorname{barccosh}(cx)} + \frac{2}{3}dex^3\sqrt{a+\operatorname{barccosh}(cx)} + \frac{1}{5}e^2x^5\sqrt{a+\operatorname{barccosh}(cx)}
\end{aligned}$$

input `Int[(d + e*x^2)^2*Sqrt[a + b*ArcCosh[c*x]],x]`

output `d^2*x*Sqrt[a + b*ArcCosh[c*x]] + (2*d*e*x^3*Sqrt[a + b*ArcCosh[c*x]])/3 + (e^2*x^5*Sqrt[a + b*ArcCosh[c*x]])/5 - (Sqrt[b]*d^2*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(4*c) - (Sqrt[b]*d*e*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(8*c^3) - (Sqrt[b]*e^2*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(32*c^5) - (Sqrt[b]*d*e*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(24*c^3) - (Sqrt[b]*e^2*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*c^5) - (Sqrt[b]*e^2*E^((5*a)/b)*Sqrt[Pi/5]*Erf[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(320*c^5) - (Sqrt[b]*d^2*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(4*c*E^(a/b)) - (Sqrt[b]*d*e*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(8*c^3*E^(a/b)) - (Sqrt[b]*e^2*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(32*c^5*E^(a/b)) - (Sqrt[b]*d*e*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(24*c^3*E^((3*a)/b)) - (Sqrt[b]*e^2*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(64*c^5*E^((3*a)/b)) - (Sqrt[b]*e^2*Sqrt[Pi/5]*Erfi[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(320*c^5*E^((5*a)/b))`

## 3.552.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

## 3.552.4 Maple [F]

$$\int (e x^2 + d)^2 \sqrt{a + b \operatorname{arccosh}(c x)} dx$$

input `int((e*x^2+d)^2*(a+b*arccosh(c*x))^(1/2),x)`

output `int((e*x^2+d)^2*(a+b*arccosh(c*x))^(1/2),x)`

## 3.552.5 Fricas [F(-2)]

Exception generated.

$$\int (d + e x^2)^2 \sqrt{a + b \operatorname{arccosh}(c x)} dx = \text{Exception raised: TypeError}$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.552.6 Sympy [F]**

$$\int (d + ex^2)^2 \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int \sqrt{a + b \operatorname{arccosh}(cx)} (d + ex^2)^2 dx$$

input `integrate((e*x**2+d)**2*(a+b*acosh(c*x))**(1/2),x)`

output `Integral(sqrt(a + b*acosh(c*x))*(d + e*x**2)**2, x)`

**3.552.7 Maxima [F]**

$$\int (d + ex^2)^2 \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int (ex^2 + d)^2 \sqrt{b \operatorname{arccosh}(cx) + a} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate((e*x^2 + d)^2*sqrt(b*arccosh(c*x) + a), x)`

**3.552.8 Giac [F]**

$$\int (d + ex^2)^2 \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int (ex^2 + d)^2 \sqrt{b \operatorname{arccosh}(cx) + a} dx$$

input `integrate((e*x^2+d)^2*(a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`

output `integrate((e*x^2 + d)^2*sqrt(b*arccosh(c*x) + a), x)`

**3.552.9 Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2)^2 \sqrt{a + \operatorname{barccosh}(cx)} dx = \int \sqrt{a + b \operatorname{acosh}(cx)} (ex^2 + d)^2 dx$$

input `int((a + b*acosh(c*x))^(1/2)*(d + e*x^2)^2,x)`output `int((a + b*acosh(c*x))^(1/2)*(d + e*x^2)^2, x)`

### 3.553 $\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx$

3.553.1 Optimal result . . . . .	4015
3.553.2 Mathematica [A] (verified) . . . . .	4016
3.553.3 Rubi [A] (verified) . . . . .	4017
3.553.4 Maple [F] . . . . .	4018
3.553.5 Fricas [F(-2)] . . . . .	4018
3.553.6 Sympy [F] . . . . .	4019
3.553.7 Maxima [F] . . . . .	4019
3.553.8 Giac [F] . . . . .	4019
3.553.9 Mupad [F(-1)] . . . . .	4020

#### 3.553.1 Optimal result

Integrand size = 20, antiderivative size = 322

$$\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx = dx \sqrt{a + b \operatorname{arccosh}(cx)} + \frac{1}{3} ex^3 \sqrt{a + b \operatorname{arccosh}(cx)}$$

$$- \frac{\sqrt{b} d e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4c}$$

$$- \frac{\sqrt{b} e e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16c^3}$$

$$- \frac{\sqrt{b} e e^{\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erf}\left(\frac{\sqrt{3} \sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{48c^3}$$

$$- \frac{\sqrt{b} d e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4c}$$

$$- \frac{\sqrt{b} e e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16c^3}$$

$$- \frac{\sqrt{b} e e^{-\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erfi}\left(\frac{\sqrt{3} \sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{48c^3}$$



output  $-1/144*e*exp(3*a/b)*erf(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*3^(1/2)*Pi^(1/2)/c^3-1/144*e*erfi(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*3^(1/2)*Pi^(1/2)/c^3/exp(3*a/b)-1/4*d*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*Pi^(1/2)/c-1/16*e*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*Pi^(1/2)/c^3-1/4*d*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*Pi^(1/2)/c/exp(a/b)-1/16*e*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*Pi^(1/2)/c^3/exp(a/b)+d*x*(a+b*arccosh(c*x))^(1/2)+1/3*e*x^3*(a+b*arccosh(c*x))^(1/2)$

### 3.553.2 Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 317, normalized size of antiderivative = 0.98

$$\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx$$

$$= \frac{de^{-\frac{a}{b}} \sqrt{a + b \operatorname{arccosh}(cx)} \left( \frac{e^{\frac{2a}{b}} \Gamma\left(\frac{3}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right)}{\sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)}} + \frac{\Gamma\left(\frac{3}{2}, -\frac{a+b \operatorname{arccosh}(cx)}{b}\right)}{\sqrt{-\frac{a+b \operatorname{arccosh}(cx)}{b}}}\right)}{2c} + \frac{ee^{-\frac{3a}{b}} \sqrt{a + b \operatorname{arccosh}(cx)} \left( 9e^{\frac{4a}{b}} \sqrt{-\frac{a+b \operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{3}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) + \sqrt{3} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{3}{2}, -\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \right)}{72c^3}$$

input `Integrate[(d + e*x^2)*Sqrt[a + b*ArcCosh[c*x]], x]`

output  $(d*\operatorname{Sqrt}[a + b*\operatorname{ArcCosh}[c*x]]*((E^((2*a)/b)*\operatorname{Gamma}[3/2, a/b + \operatorname{ArcCosh}[c*x]])/\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]] + \operatorname{Gamma}[3/2, -((a + b*\operatorname{ArcCosh}[c*x])/b)]/\operatorname{Sqrt}[-((a + b*\operatorname{ArcCosh}[c*x])/b)]))/(2*c*E^(a/b)) + (e*\operatorname{Sqrt}[a + b*\operatorname{ArcCosh}[c*x]]*(9*E^((4*a)/b)*\operatorname{Sqrt}[-((a + b*\operatorname{ArcCosh}[c*x])/b)]*\operatorname{Gamma}[3/2, a/b + \operatorname{ArcCosh}[c*x]] + \operatorname{Sqrt}[3]*\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[3/2, (-3*(a + b*\operatorname{ArcCosh}[c*x]))/b] + 9*E^((2*a)/b)*\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[3/2, -((a + b*\operatorname{ArcCosh}[c*x])/b)] + \operatorname{Sqrt}[3]*E^((6*a)/b)*\operatorname{Sqrt}[-((a + b*\operatorname{ArcCosh}[c*x])/b)]*\operatorname{Gamma}[3/2, (3*(a + b*\operatorname{ArcCosh}[c*x])/b)]))/(72*c^3*E^((3*a)/b)*\operatorname{Sqrt}[-((a + b*\operatorname{ArcCosh}[c*x])/b)])$

**3.553.3 Rubi [A] (verified)**

Time = 1.46 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int (d + ex^2) \sqrt{a + \operatorname{barccosh}(cx)} dx \\
 & \quad \downarrow \text{6324} \\
 & \int \left( d\sqrt{a + \operatorname{barccosh}(cx)} + ex^2 \sqrt{a + \operatorname{barccosh}(cx)} \right) dx \\
 & \quad \downarrow \text{2009} \\
 & \frac{\sqrt{\pi} \sqrt{b} e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{16c^3} - \frac{\sqrt{\frac{\pi}{3}} \sqrt{b} e^{\frac{3a}{b}} \operatorname{erf}\left(\frac{\sqrt{3} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{48c^3} \\
 & - \frac{\sqrt{\pi} \sqrt{b} e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{16c^3} - \frac{\sqrt{\frac{\pi}{3}} \sqrt{b} e^{-\frac{3a}{b}} \operatorname{erfi}\left(\frac{\sqrt{3} \sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{48c^3} \\
 & \frac{\sqrt{\pi} \sqrt{b} d e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{4c} - \frac{\sqrt{\pi} \sqrt{b} d e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{4c} + dx \sqrt{a + \operatorname{barccosh}(cx)} + \\
 & \quad \frac{1}{3} ex^3 \sqrt{a + \operatorname{barccosh}(cx)}
 \end{aligned}$$

input `Int[(d + e*x^2)*Sqrt[a + b*ArcCosh[c*x]],x]`

output `d*x*Sqrt[a + b*ArcCosh[c*x]] + (e*x^3*Sqrt[a + b*ArcCosh[c*x]])/3 - (Sqrt[b]*d*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(4*c) - (Sqrt[b]*e*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(16*c^3) - (Sqrt[b]*e*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(48*c^3) - (Sqrt[b]*d*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(4*c*E^(a/b)) - (Sqrt[b]*e*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(16*c^3*E^(a/b)) - (Sqrt[b]*e*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(48*c^3*E^((3*a)/b))`

## 3.553.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.),  
x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x],  
x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] &&  
(p > 0 || IGtQ[n, 0])`

## 3.553.4 Maple [F]

$$\int (ex^2 + d) \sqrt{a + b \operatorname{arccosh}(cx)} dx$$

input `int((e*x^2+d)*(a+b*arccosh(c*x))^(1/2),x)`

output `int((e*x^2+d)*(a+b*arccosh(c*x))^(1/2),x)`

## 3.553.5 Fricas [F(-2)]

Exception generated.

$$\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx = \text{Exception raised: TypeError}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: inte  
grate: implementation incomplete (constant residues)`

**3.553.6 Sympy [F]**

$$\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int \sqrt{a + b \operatorname{acosh}(cx)} (d + ex^2) dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x))**(1/2),x)`

output `Integral(sqrt(a + b*acosh(c*x))*(d + e*x**2), x)`

**3.553.7 Maxima [F]**

$$\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int (ex^2 + d) \sqrt{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate((e*x^2 + d)*sqrt(b*arccosh(c*x) + a), x)`

**3.553.8 Giac [F]**

$$\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int (ex^2 + d) \sqrt{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`

output `integrate((e*x^2 + d)*sqrt(b*arccosh(c*x) + a), x)`

**3.553.9 Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int \sqrt{a + b \operatorname{acosh}(cx)} (ex^2 + d) dx$$

input `int((a + b*acosh(c*x))^(1/2)*(d + e*x^2),x)`output `int((a + b*acosh(c*x))^(1/2)*(d + e*x^2), x)`

### 3.554 $\int \sqrt{a + b \operatorname{arccosh}(cx)} dx$

3.554.1 Optimal result . . . . .	4021
3.554.2 Mathematica [A] (verified) . . . . .	4021
3.554.3 Rubi [A] (verified) . . . . .	4022
3.554.4 Maple [F] . . . . .	4024
3.554.5 Fracas [F(-2)] . . . . .	4025
3.554.6 Sympy [F] . . . . .	4025
3.554.7 Maxima [F] . . . . .	4025
3.554.8 Giac [F] . . . . .	4026
3.554.9 Mupad [F(-1)] . . . . .	4026

#### 3.554.1 Optimal result

Integrand size = 12, antiderivative size = 102

$$\int \sqrt{a + b \operatorname{arccosh}(cx)} dx = x \sqrt{a + b \operatorname{arccosh}(cx)} - \frac{\sqrt{b} e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4c} - \frac{\sqrt{b} e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4c}$$

```
output -1/4*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*Pi^(1/2)/c-1/4
*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*b^(1/2)*Pi^(1/2)/c/exp(a/b)+x*(a+b
*arccosh(c*x))^(1/2)
```

#### 3.554.2 Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.98

$$\int \sqrt{a + b \operatorname{arccosh}(cx)} dx = \frac{e^{-\frac{a}{b}} \sqrt{a + b \operatorname{arccosh}(cx)} \left( \frac{e^{\frac{2a}{b}} \Gamma\left(\frac{3}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right)}{\sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)}} + \frac{\Gamma\left(\frac{3}{2}, -\frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{\sqrt{-\frac{a + b \operatorname{arccosh}(cx)}{b}}}\right)}{2c}$$

input `Integrate[Sqrt[a + b*ArcCosh[c*x]], x]`

output `(Sqrt[a + b*ArcCosh[c*x]]*((E^((2*a)/b)*Gamma[3/2, a/b + ArcCosh[c*x]])/Sqrt[a/b + ArcCosh[c*x]] + Gamma[3/2, -(a + b*ArcCosh[c*x])/b])/Sqrt[-((a + b*ArcCosh[c*x])/b]))/(2*c*E^(a/b))`

### 3.554.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.02, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6294, 6368, 3042, 3788, 26, 2611, 2633, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \sqrt{a + \operatorname{barccosh}(cx)} dx \\
 & \quad \downarrow 6294 \\
 & x\sqrt{a + \operatorname{barccosh}(cx)} - \frac{1}{2}bc \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a + \operatorname{barccosh}(cx)}} dx \\
 & \quad \downarrow 6368 \\
 & x\sqrt{a + \operatorname{barccosh}(cx)} - \frac{\int \frac{\cosh\left(\frac{a}{b} - \frac{a + b\operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{2c} \\
 & \quad \downarrow 3042 \\
 & x\sqrt{a + \operatorname{barccosh}(cx)} - \frac{\int \frac{\sin\left(\frac{ia}{b} - \frac{i(a + b\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right)}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{2c} \\
 & \quad \downarrow 3788 \\
 & \frac{x\sqrt{a + \operatorname{barccosh}(cx)} - \frac{1}{2}i \int -\frac{ie^{-\operatorname{arccosh}(cx)}}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i \int \frac{ie^{\operatorname{arccosh}(cx)}}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{2c} \\
 & \quad \downarrow 26
 \end{aligned}$$

$$\begin{aligned}
& \frac{\frac{1}{2} \int \frac{e^{-\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a+\operatorname{barccosh}(cx)) + \frac{1}{2} \int \frac{e^{\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a+\operatorname{barccosh}(cx))}{2c} \\
& \quad \downarrow \text{2611} \\
& \frac{\int e^{\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}} d\sqrt{a+\operatorname{barccosh}(cx)} + \int e^{\frac{a+\operatorname{barccosh}(cx)}{b} - \frac{a}{b}} d\sqrt{a+\operatorname{barccosh}(cx)}}{2c} \\
& \quad \downarrow \text{2633} \\
& \frac{x\sqrt{a+\operatorname{barccosh}(cx)} - \int e^{\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}} d\sqrt{a+\operatorname{barccosh}(cx)} + \frac{1}{2}\sqrt{\pi}\sqrt{b}e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{2c} \\
& \quad \downarrow \text{2634} \\
& x\sqrt{a+\operatorname{barccosh}(cx)} - \frac{\frac{1}{2}\sqrt{\pi}\sqrt{b}e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right) + \frac{1}{2}\sqrt{\pi}\sqrt{b}e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{2c}
\end{aligned}$$

input `Int[Sqrt[a + b*ArcCosh[c*x]], x]`

output `x*Sqrt[a + b*ArcCosh[c*x]] - ((Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/2 + (Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(2*E^(a/b))/(2*c)`

### 3.554.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2611 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`



rule 2633 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))2), x_Symbol] := Simp[Fa*Sqrt  
[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{  
F, a, b, c, d}, x] && PosQ[b]`

rule 2634 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))2), x_Symbol] := Simp[Fa*Sqrt  
[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; Fr  
eeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3788 `Int[((c_.) + (d_.)*(x_))(m_.)*sin[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol  
] := Simp[I/2 Int[(c + d*x)m/(E(I*k*Pi)*E(I*(e + f*x))), x], x] - Simp  
[I/2 Int[(c + d*x)m*E(I*k*Pi)*E(I*(e + f*x)), x], x] /; FreeQ[{c, d, e,  
f, m}, x] && IntegerQ[2*k]`

rule 6294 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))(n_.), x_Symbol] := Simp[x*(a + b*A  
rcCosh[c*x])n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])(n - 1)/(Sqrt  
[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))(n_.)*(x_)(m_.)*((d1_.) + (e1_.)*(x  
_))(p_.)*((d2_.) + (e2_.)*(x_))(p_.), x_Symbol] := Simp[(1/(b*c(m + 1)))*  
Simp[(d1 + e1*x)p/(1 + c*x)p*Simp[(d2 + e2*x)p/(-1 + c*x)p Subst[In  
t[xn*Cosh[-a/b + x/b]m*Sinh[-a/b + x/b](2*p + 1), x], x, a + b*ArcCosh[c  
*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[  
e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.554.4 Maple [F]

$$\int \sqrt{a + b \operatorname{arccosh}(cx)} dx$$

input `int((a+b*arccosh(c*x))(1/2),x)`

output `int((a+b*arccosh(c*x))(1/2),x)`

**3.554.5 Fracas [F(-2)]**

Exception generated.

$$\int \sqrt{a + \operatorname{barccosh}(cx)} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.554.6 Sympy [F]**

$$\int \sqrt{a + \operatorname{barccosh}(cx)} dx = \int \sqrt{a + b \operatorname{acosh}(cx)} dx$$

input `integrate((a+b*acosh(c*x))**(1/2),x)`

output `Integral(sqrt(a + b*acosh(c*x)), x)`

**3.554.7 Maxima [F]**

$$\int \sqrt{a + \operatorname{barccosh}(cx)} dx = \int \sqrt{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*arccosh(c*x) + a), x)`

**3.554.8 Giac [F]**

$$\int \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int \sqrt{b \operatorname{arcosh}(cx) + a} dx$$

input `integrate((a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`

output `integrate(sqrt(b*arccosh(c*x) + a), x)`

**3.554.9 Mupad [F(-1)]**

Timed out.

$$\int \sqrt{a + b \operatorname{arccosh}(cx)} dx = \int \sqrt{a + b \operatorname{acosh}(cx)} dx$$

input `int((a + b*acosh(c*x))^(1/2),x)`

output `int((a + b*acosh(c*x))^(1/2), x)`

$$3.555 \quad \int \frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{d+ex^2} dx$$

3.555.1 Optimal result	4027
3.555.2 Mathematica [F(-1)]	4027
3.555.3 Rubi [N/A]	4028
3.555.4 Maple [N/A] (verified)	4028
3.555.5 Fricas [F(-2)]	4029
3.555.6 Sympy [N/A]	4029
3.555.7 Maxima [F(-2)]	4029
3.555.8 Giac [N/A]	4030
3.555.9 Mupad [N/A]	4030

### 3.555.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{\sqrt{a + \operatorname{arccosh}(cx)}}{d + ex^2} dx = \operatorname{Int}\left(\frac{\sqrt{a + \operatorname{arccosh}(cx)}}{d + ex^2}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^(1/2)/(e*x^2+d),x)`

### 3.555.2 Mathematica [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + \operatorname{arccosh}(cx)}}{d + ex^2} dx = \$Aborted$$

input `Integrate[Sqrt[a + b*ArcCosh[c*x]]/(d + e*x^2),x]`

output `$Aborted`

**3.555.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{d + ex^2} dx$$

↓ 6325

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{d + ex^2} dx$$

input `Int[Sqrt[a + b*ArcCosh[c*x]]/(d + e*x^2), x]`

output `$Aborted`

**3.555.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.555.4 Maple [N/A] (verified)**

Not integrable

Time = 1.41 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{ex^2 + d} dx$$

input `int((a+b*arccosh(c*x))^(1/2)/(e*x^2+d), x)`

output `int((a+b*arccosh(c*x))^(1/2)/(e*x^2+d), x)`

**3.555.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{d + ex^2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^(1/2)/(e*x^2+d),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.555.6 Sympy [N/A]**

Not integrable

Time = 1.59 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{d + ex^2} dx = \int \frac{\sqrt{a + b \operatorname{acosh}(cx)}}{d + ex^2} dx$$

input `integrate((a+b*acosh(c*x))**(1/2)/(e*x**2+d),x)`

output `Integral(sqrt(a + b*acosh(c*x))/(d + e*x**2), x)`

**3.555.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))^(1/2)/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

---

3.555.  $\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{d + ex^2} dx$

**3.555.8 Giac [N/A]**

Not integrable

Time = 12.58 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{d + ex^2} dx = \int \frac{\sqrt{b \operatorname{arccosh}(cx) + a}}{ex^2 + d} dx$$

input `integrate((a+b*arccosh(c*x))^(1/2)/(e*x^2+d),x, algorithm="giac")`output `integrate(sqrt(b*arccosh(c*x) + a)/(e*x^2 + d), x)`**3.555.9 Mupad [N/A]**

Not integrable

Time = 3.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{d + ex^2} dx = \int \frac{\sqrt{a + b \operatorname{acosh}(cx)}}{ex^2 + d} dx$$

input `int((a + b*acosh(c*x))^(1/2)/(d + e*x^2),x)`output `int((a + b*acosh(c*x))^(1/2)/(d + e*x^2), x)`

$$3.556 \quad \int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx$$

3.556.1 Optimal result	4031
3.556.2 Mathematica [F(-1)]	4031
3.556.3 Rubi [N/A]	4032
3.556.4 Maple [N/A] (verified)	4032
3.556.5 Fricas [F(-2)]	4033
3.556.6 Sympy [N/A]	4033
3.556.7 Maxima [N/A]	4033
3.556.8 Giac [N/A]	4034
3.556.9 Mupad [N/A]	4034

### 3.556.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx = \operatorname{Int} \left( \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2}, x \right)$$

output `Unintegrable((a+b*arccosh(c*x))^(1/2)/(e*x^2+d)^2,x)`

### 3.556.2 Mathematica [F(-1)]

Timed out.

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx = \$Aborted$$

input `Integrate[Sqrt[a + b*ArcCosh[c*x]]/(d + e*x^2)^2,x]`

output `$Aborted`

---


$$3.556. \quad \int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx$$



**3.556.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx$$

↓ 6325

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx$$

input `Int[Sqrt[a + b*ArcCosh[c*x]]/(d + e*x^2)^2,x]`

output `$Aborted`

**3.556.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.556.4 Maple [N/A] (verified)**

Not integrable

Time = 0.93 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(ex^2 + d)^2} dx$$

input `int((a+b*arccosh(c*x))^(1/2)/(e*x^2+d)^2,x)`

output `int((a+b*arccosh(c*x))^(1/2)/(e*x^2+d)^2,x)`

---

3.556.  $\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx$

**3.556.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{(d + ex^2)^2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^(1/2)/(e*x^2+d)^2,x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.556.6 Sympy [N/A]**

Not integrable

Time = 41.82 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{(d + ex^2)^2} dx = \int \frac{\sqrt{a + b \operatorname{acosh}(cx)}}{(d + ex^2)^2} dx$$

input `integrate((a+b*acosh(c*x))**(1/2)/(e*x**2+d)**2,x)`

output `Integral(sqrt(a + b*acosh(c*x))/(d + e*x**2)**2, x)`

**3.556.7 Maxima [N/A]**

Not integrable

Time = 0.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{(d + ex^2)^2} dx = \int \frac{\sqrt{b \operatorname{arcosh}(cx) + a}}{(ex^2 + d)^2} dx$$

input `integrate((a+b*arccosh(c*x))^(1/2)/(e*x^2+d)^2,x, algorithm="maxima")`

output `integrate(sqrt(b*arccosh(c*x) + a)/(e*x^2 + d)^2, x)`

---

3.556.  $\int \frac{\sqrt{a + \operatorname{barccosh}(cx)}}{(d + ex^2)^2} dx$

**3.556.8 Giac [N/A]**

Not integrable

Time = 12.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx = \int \frac{\sqrt{b \operatorname{arccosh}(cx) + a}}{(ex^2 + d)^2} dx$$

input `integrate((a+b*arccosh(c*x))^(1/2)/(e*x^2+d)^2,x, algorithm="giac")`output `integrate(sqrt(b*arccosh(c*x) + a)/(e*x^2 + d)^2, x)`**3.556.9 Mupad [N/A]**

Not integrable

Time = 3.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{a + b \operatorname{arccosh}(cx)}}{(d + ex^2)^2} dx = \int \frac{\sqrt{a + b \operatorname{acosh}(cx)}}{(ex^2 + d)^2} dx$$

input `int((a + b*acosh(c*x))^(1/2)/(d + e*x^2)^2,x)`output `int((a + b*acosh(c*x))^(1/2)/(d + e*x^2)^2, x)`

### 3.557 $\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx$

3.557.1 Optimal result . . . . .	4035
3.557.2 Mathematica [A] (warning: unable to verify) . . . . .	4036
3.557.3 Rubi [A] (verified) . . . . .	4037
3.557.4 Maple [F] . . . . .	4039
3.557.5 Fricas [F(-2)] . . . . .	4039
3.557.6 Sympy [F] . . . . .	4039
3.557.7 Maxima [F] . . . . .	4040
3.557.8 Giac [F(-2)] . . . . .	4040
3.557.9 Mupad [F(-1)] . . . . .	4040

#### 3.557.1 Optimal result

Integrand size = 20, antiderivative size = 442

$$\begin{aligned}
 & \int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx = \\
 & \frac{3bd\sqrt{-1 + cx}\sqrt{1 + cx}\sqrt{a + \operatorname{barccosh}(cx)}}{2c} \\
 & - \frac{be\sqrt{-1 + cx}\sqrt{1 + cx}\sqrt{a + \operatorname{barccosh}(cx)}}{3c^3} \\
 & - \frac{bex^2\sqrt{-1 + cx}\sqrt{1 + cx}\sqrt{a + \operatorname{barccosh}(cx)}}{6c} \\
 & + dx(a + \operatorname{barccosh}(cx))^{3/2} + \frac{1}{3}ex^3(a + \operatorname{barccosh}(cx))^{3/2} \\
 & - \frac{3b^{3/2}de^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c} - \frac{3b^{3/2}ee^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{32c^3} \\
 & - \frac{b^{3/2}ee^{\frac{3a}{b}}\sqrt{\frac{\pi}{3}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{96c^3} + \frac{3b^{3/2}de^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c} \\
 & + \frac{3b^{3/2}ee^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{32c^3} + \frac{b^{3/2}ee^{-\frac{3a}{b}}\sqrt{\frac{\pi}{3}}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{96c^3}
 \end{aligned}$$

```
output d*x*(a+b*arccosh(c*x))^(3/2)+1/3*e*x^3*(a+b*arccosh(c*x))^(3/2)-1/288*b^(3/2)*e*exp(3*a/b)*erf(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/c^3+1/288*b^(3/2)*e*erfi(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/c^3/exp(3*a/b)-3/8*b^(3/2)*d*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c-3/32*b^(3/2)*e*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c^3+3/8*b^(3/2)*d*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c/exp(a/b)+3/32*b^(3/2)*e*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c^3/exp(a/b)-3/2*b*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(a+b*arccosh(c*x))^(1/2)/c-1/3*b*e*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(a+b*arccosh(c*x))^(1/2)/c^3-1/6*b*e*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(a+b*arccosh(c*x))^(1/2)/c
```

### 3.557.2 Mathematica [A] (warning: unable to verify)

Time = 1.99 (sec) , antiderivative size = 812, normalized size of antiderivative = 1.84

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx = \frac{ade^{-\frac{a}{b}} \sqrt{a + \operatorname{barccosh}(cx)} \left( \frac{e^{\frac{2a}{b}} \Gamma\left(\frac{3}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right)}{\sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)}} + \frac{\Gamma\left(\frac{3}{2}, -\frac{a+b\operatorname{arccosh}(cx)}{b}\right)}{\sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}}}\right)}{2c} + \frac{aee^{-\frac{3a}{b}} \sqrt{a + \operatorname{barccosh}(cx)} \left( 9e^{\frac{4a}{b}} \sqrt{-\frac{a+b\operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{3}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) + \sqrt{3} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{3}{2}, -\frac{a+b\operatorname{arccosh}(cx)}{b}\right) \right)}{8c} + \frac{bd \left( -12 \sqrt{\frac{-1+cx}{1+cx}} (1 + cx) \sqrt{a + \operatorname{barccosh}(cx)} + 8cx \operatorname{arccosh}(cx) \sqrt{a + \operatorname{barccosh}(cx)} + \frac{72c^3 \sqrt{a + \operatorname{barccosh}(cx)}}{(2a+3b)\sqrt{\pi}} \operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right) \right)}{8c} + \frac{\sqrt{b}e \left( 9 \left( -12\sqrt{b} \sqrt{\frac{-1+cx}{1+cx}} (1 + cx) \sqrt{a + \operatorname{barccosh}(cx)} + 8\sqrt{b}cx \operatorname{arccosh}(cx) \sqrt{a + \operatorname{barccosh}(cx)} + (2a + 3b) \sqrt{a + \operatorname{barccosh}(cx)} \right) \right)}{8c}$$

```
input Integrate[(d + e*x^2)*(a + b*ArcCosh[c*x])^(3/2), x]
```

output

```
(a*d*Sqrt[a + b*ArcCosh[c*x]]*(E^((2*a)/b)*Gamma[3/2, a/b + ArcCosh[c*x]]
)/Sqrt[a/b + ArcCosh[c*x]] + Gamma[3/2, -((a + b*ArcCosh[c*x])/b)]/Sqrt[-(
(a + b*ArcCosh[c*x])/b)))/(2*c*E^(a/b)) + (a*e*Sqrt[a + b*ArcCosh[c*x]]*(
9*E^((4*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[3/2, a/b + ArcCosh[c*x
]] + Sqrt[3]*Sqrt[a/b + ArcCosh[c*x]]*Gamma[3/2, (-3*(a + b*ArcCosh[c*x]))
/b] + 9*E^((2*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[3/2, -((a + b*ArcCosh[c
*x])/b)] + Sqrt[3]*E^((6*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[3/2,
(3*(a + b*ArcCosh[c*x]))/b]))/(72*c^3*E^((3*a)/b)*Sqrt[-((a + b*ArcCosh[c*
x])^2/b^2)]) + (b*d*(-12*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Sqrt[a + b*A
rcCosh[c*x]] + 8*c*x*ArcCosh[c*x]*Sqrt[a + b*ArcCosh[c*x]] + ((2*a + 3*b)*
Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]*(Cosh[a/b] - Sinh[a/b]))/S
qrt[b] + ((2*a - 3*b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]*(Cosh
[a/b] + Sinh[a/b])/Sqrt[b]))/(8*c) + (Sqrt[b]*e*(9*(-12*Sqrt[b]*Sqrt[(-1
+ c*x)/(1 + c*x)]*(1 + c*x)*Sqrt[a + b*ArcCosh[c*x]] + 8*Sqrt[b]*c*x*ArcCo
sh[c*x]*Sqrt[a + b*ArcCosh[c*x]] + (2*a + 3*b)*Sqrt[Pi]*Erfi[Sqrt[a + b*Ar
cCosh[c*x]]/Sqrt[b]]*(Cosh[a/b] - Sinh[a/b]) + (2*a - 3*b)*Sqrt[Pi]*Erf[Sq
rt[a + b*ArcCosh[c*x]]/Sqrt[b]]*(Cosh[a/b] + Sinh[a/b])) + (2*a + b)*Sqrt[
3*Pi]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]*(Cosh[(3*a)/b] - Si
nh[(3*a)/b]) + (2*a - b)*Sqrt[3*Pi]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])
/Sqrt[b]]*(Cosh[(3*a)/b] + Sinh[(3*a)/b]) + 12*Sqrt[b]*Sqrt[a + b*ArcCo...
```

### 3.557.3 Rubi [A] (verified)

Time = 2.02 (sec) , antiderivative size = 442, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx$$

$$\downarrow \text{6324}$$

$$\int \left( d(a + \operatorname{barccosh}(cx))^{3/2} + ex^2(a + \operatorname{barccosh}(cx))^{3/2} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{3\sqrt{\pi}b^{3/2}ee^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{32c^3} - \frac{\sqrt{\frac{\pi}{3}}b^{3/2}ee^{\frac{3a}{b}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{96c^3} + \\
& \frac{3\sqrt{\pi}b^{3/2}ee^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{32c^3} + \frac{\sqrt{\frac{\pi}{3}}b^{3/2}ee^{-\frac{3a}{b}}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{96c^3} - \\
& \frac{3\sqrt{\pi}b^{3/2}de^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c} + \frac{3\sqrt{\pi}b^{3/2}de^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c} - \\
& \frac{be\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}}{3c^3} + dx(a+\operatorname{barccosh}(cx))^{3/2} - \\
& \frac{3bd\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}}{2c} + \frac{1}{3}ex^3(a+\operatorname{barccosh}(cx))^{3/2} - \\
& \frac{be x^2\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}}{6c}
\end{aligned}$$

input `Int[(d + e*x^2)*(a + b*ArcCosh[c*x])^(3/2),x]`

output `(-3*b*d*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Sqrt[a + b*ArcCosh[c*x]]/(2*c) - (b*e*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Sqrt[a + b*ArcCosh[c*x]]/(3*c^3) - (b*e*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Sqrt[a + b*ArcCosh[c*x]]/(6*c) + d*x*(a + b*ArcCosh[c*x])^(3/2) + (e*x^3*(a + b*ArcCosh[c*x])^(3/2))/3 - (3*b^(3/2)*d*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(8*c) - (3*b^(3/2)*e*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(32*c^3) - (b^(3/2)*e*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(96*c^3) + (3*b^(3/2)*d*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(8*c*E^(a/b)) + (3*b^(3/2)*e*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(32*c^3*E^(a/b)) + (b^(3/2)*e*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(96*c^3*E^((3*a)/b)))`

### 3.557.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

**3.557.4 Maple [F]**

$$\int (e x^2 + d) (a + b \operatorname{arccosh}(cx))^{\frac{3}{2}} dx$$

input `int((e*x^2+d)*(a+b*arccosh(c*x))^(3/2),x)`

output `int((e*x^2+d)*(a+b*arccosh(c*x))^(3/2),x)`

**3.557.5 Fracas [F(-2)]**

Exception generated.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{\frac{3}{2}} dx = \text{Exception raised: TypeError}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.557.6 Sympy [F]**

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{\frac{3}{2}} dx = \int (a + b \operatorname{acosh}(cx))^{\frac{3}{2}} (d + ex^2) dx$$

input `integrate((e*x**2+d)*(a+b*acosh(c*x))**(3/2),x)`

output `Integral((a + b*acosh(c*x))**(3/2)*(d + e*x**2), x)`



**3.557.7 Maxima [F]**

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx = \int (ex^2 + d)(b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}} dx$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((e*x^2 + d)*(b*arccosh(c*x) + a)^(3/2), x)`

**3.557.8 Giac [F(-2)]**

Exception generated.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx = \text{Exception raised: RuntimeError}$$

input `integrate((e*x^2+d)*(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `Exception raised: RuntimeError >> an error occurred running a Giac command  
:INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const ve  
cteur & l) Error: Bad Argument Value`

**3.557.9 Mupad [F(-1)]**

Timed out.

$$\int (d + ex^2) (a + \operatorname{barccosh}(cx))^{3/2} dx = \int (a + b \operatorname{acosh}(cx))^{3/2} (ex^2 + d) dx$$

input `int((a + b*acosh(c*x))^(3/2)*(d + e*x^2),x)`

output `int((a + b*acosh(c*x))^(3/2)*(d + e*x^2), x)`

### 3.558 $\int (a + \operatorname{barccosh}(cx))^{3/2} dx$

3.558.1 Optimal result . . . . .	4041
3.558.2 Mathematica [A] (warning: unable to verify) . . . . .	4042
3.558.3 Rubi [C] (verified) . . . . .	4042
3.558.4 Maple [F] . . . . .	4046
3.558.5 Fracas [F(-2)] . . . . .	4046
3.558.6 Sympy [F] . . . . .	4046
3.558.7 Maxima [F] . . . . .	4047
3.558.8 Giac [F] . . . . .	4047
3.558.9 Mupad [F(-1)] . . . . .	4047

#### 3.558.1 Optimal result

Integrand size = 12, antiderivative size = 140

$$\int (a + \operatorname{barccosh}(cx))^{3/2} dx = -\frac{3b\sqrt{-1 + cx}\sqrt{1 + cx}\sqrt{a + \operatorname{barccosh}(cx)}}{2c}$$

$$+ x(a + \operatorname{barccosh}(cx))^{3/2} - \frac{3b^{3/2}e^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c}$$

$$+ \frac{3b^{3/2}e^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8c}$$

```
output x*(a+b*arccosh(c*x))^(3/2)-3/8*b^(3/2)*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c+3/8*b^(3/2)*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c/exp(a/b)-3/2*b*(c*x-1)^(1/2)*(c*x+1)^(1/2)*(a+b*arccosh(c*x))^(1/2)/c
```

### 3.558.2 Mathematica [A] (warning: unable to verify)

Time = 0.40 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.92

$$\int (a + b \operatorname{arccosh}(cx))^{3/2} dx = \frac{ae^{-\frac{a}{b}} \sqrt{a + b \operatorname{arccosh}(cx)} \left( \frac{e^{\frac{2a}{b}} \Gamma\left(\frac{3}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right)}{\sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)}} + \frac{\Gamma\left(\frac{3}{2}, -\frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{\sqrt{-\frac{a + b \operatorname{arccosh}(cx)}{b}}}\right)}{2c} + \frac{b \left( -12 \sqrt{\frac{-1+cx}{1+cx}} (1+cx) \sqrt{a + b \operatorname{arccosh}(cx)} + 8cx \operatorname{arccosh}(cx) \sqrt{a + b \operatorname{arccosh}(cx)} + \frac{(2a+3b)\sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a+b \operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{\sqrt{b}} \right)}{8c}$$

input `Integrate[(a + b*ArcCosh[c*x])^(3/2), x]`

output `(a*Sqrt[a + b*ArcCosh[c*x]]*(E^((2*a)/b)*Gamma[3/2, a/b + ArcCosh[c*x]])/Sqrt[a/b + ArcCosh[c*x]] + Gamma[3/2, -(a + b*ArcCosh[c*x])/b])/Sqrt[-((a + b*ArcCosh[c*x])/b)))/(2*c*E^(a/b)) + (b*(-12*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x)*Sqrt[a + b*ArcCosh[c*x]] + 8*c*x*ArcCosh[c*x]*Sqrt[a + b*ArcCosh[c*x]] + ((2*a + 3*b)*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]*(Cosh[a/b] - Sinh[a/b]))/Sqrt[b] + ((2*a - 3*b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]*(Cosh[a/b] + Sinh[a/b]))/Sqrt[b]))/(8*c)`

### 3.558.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.87 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$ , Rules used = {6294, 6330, 6296, 25, 3042, 26, 3789, 2611, 2633, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + b \operatorname{arccosh}(cx))^{3/2} dx$$

↓ 6294

$$\begin{aligned}
& x(a + \operatorname{barccosh}(cx))^{3/2} - \frac{3}{2}bc \int \frac{x\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{cx-1}\sqrt{cx+1}} dx \\
& \quad \downarrow \text{6330} \\
& x(a + \operatorname{barccosh}(cx))^{3/2} - \frac{3}{2}bc \left( \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a + \operatorname{barccosh}(cx)}}{c^2} - \frac{b \int \frac{1}{\sqrt{a + \operatorname{barccosh}(cx)}} dx}{2c} \right) \\
& \quad \downarrow \text{6296} \\
& \frac{3}{2}bc \left( \frac{x(a + \operatorname{barccosh}(cx))^{3/2} - \int \frac{\sinh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{c^2} \right) \\
& \quad \downarrow \text{25} \\
& \frac{3}{2}bc \left( \frac{x(a + \operatorname{barccosh}(cx))^{3/2} - \int \frac{\sinh\left(\frac{a}{b} - \frac{a + \operatorname{barccosh}(cx)}{b}\right)}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{2c^2} + \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a + \operatorname{barccosh}(cx)}}{c^2} \right) \\
& \quad \downarrow \text{3042} \\
& \frac{3}{2}bc \left( \frac{x(a + \operatorname{barccosh}(cx))^{3/2} - \int \frac{i \sin\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{c^2} + \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a + \operatorname{barccosh}(cx)}}{2c^2} \right) \\
& \quad \downarrow \text{26} \\
& \frac{3}{2}bc \left( \frac{x(a + \operatorname{barccosh}(cx))^{3/2} - i \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a + \operatorname{barccosh}(cx))}{b}\right)}{\sqrt{a + \operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx))}{c^2} \right) \\
& \quad \downarrow \text{3789}
\end{aligned}$$

$$\frac{3}{2}bc \left( \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}}{c^2} - \frac{i \left( \frac{1}{2}i \int \frac{e^{-\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a+\operatorname{barccosh}(cx)) - \frac{1}{2}i \int \frac{e^{\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a+\operatorname{barccosh}(cx)) \right)}{2c^2} \right)$$

↓ 2611

$$\frac{3}{2}bc \left( \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}}{c^2} - \frac{i \left( i \int e^{\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}} d\sqrt{a+\operatorname{barccosh}(cx)} - i \int e^{\frac{a+\operatorname{barccosh}(cx)}{b} - \frac{a}{b}} d\sqrt{a+\operatorname{barccosh}(cx)} \right)}{2c^2} \right)$$

↓ 2633

$$\frac{3}{2}bc \left( \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}}{c^2} - \frac{i \left( i \int e^{\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}} d\sqrt{a+\operatorname{barccosh}(cx)} - \frac{1}{2}i\sqrt{\pi}\sqrt{b}e^{-\frac{a}{b}} \operatorname{erfi} \left( \frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) \right)}{2c^2} \right)$$

↓ 2634

$$\frac{3}{2}bc \left( \frac{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}}{c^2} - \frac{i \left( \frac{1}{2}i\sqrt{\pi}\sqrt{b}e^{a/b} \operatorname{erf} \left( \frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{2}i\sqrt{\pi}\sqrt{b}e^{-\frac{a}{b}} \operatorname{erfi} \left( \frac{\sqrt{a+\operatorname{barccosh}(cx)}}{\sqrt{b}} \right) \right)}{2c^2} \right)$$

input `Int[(a + b*ArcCosh[c*x])^(3/2), x]`

output `x*(a + b*ArcCosh[c*x])^(3/2) - (3*b*c*((Sqrt[-1 + c*x]*Sqrt[1 + c*x]*Sqrt[a + b*ArcCosh[c*x]])/c^2 - ((I/2)*((I/2)*Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]] - ((I/2)*Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/E^(a/b)))/c^2)/2`

## 3.558.3.1 Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`
- rule 2611 `Int[(F_)^((g_)*(e_) + (f_)*(x_))/Sqrt[(c_) + (d_)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`
- rule 2633 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`
- rule 2634 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`
- rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`
- rule 3789 `Int[((c_) + (d_)*(x_)^m)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)^m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)^m*E^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]`
- rule 6294 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[x*(a + b*ArcCosh[c*x])^n, x] - Simp[b*c*n Int[x*((a + b*ArcCosh[c*x])^(n - 1))/(Sqrt[1 + c*x]*Sqrt[-1 + c*x]), x], x] /; FreeQ[{a, b, c}, x] && GtQ[n, 0]`
- rule 6296 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[1/(b*c) Subst[Int[x^n*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, n}, x]`

```
rule 6330 Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d1_) + (e1_.)*(x_))^(p
_)*((d2_) + (e2_.)*(x_))^(p_), x_Symbol] := Simp[(d1 + e1*x)^(p + 1)*(d2 +
e2*x)^(p + 1)*((a + b*ArcCosh[c*x])^n/(2*e1*e2*(p + 1))), x] - Simp[b*(n/(2
*c*(p + 1)))*Simp[(d1 + e1*x)^p/(1 + c*x)^p]*Simp[(d2 + e2*x)^p/(-1 + c*x)^
p] Int[(1 + c*x)^(p + 1/2)*(-1 + c*x)^(p + 1/2)*(a + b*ArcCosh[c*x])^(n -
1), x], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, p}, x] && EqQ[e1, c*d1] && E
qQ[e2, (-c)*d2] && GtQ[n, 0] && NeQ[p, -1]
```

### 3.558.4 Maple [F]

$$\int (a + b \operatorname{arccosh}(cx))^{\frac{3}{2}} dx$$

```
input int((a+b*arccosh(c*x))^(3/2),x)
```

```
output int((a+b*arccosh(c*x))^(3/2),x)
```

### 3.558.5 Fracas [F(-2)]

Exception generated.

$$\int (a + b \operatorname{arccosh}(cx))^{\frac{3}{2}} dx = \text{Exception raised: TypeError}$$

```
input integrate((a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")
```

```
output Exception raised: TypeError >> Error detected within library code: inte
grate: implementation incomplete (constant residues)
```

### 3.558.6 Sympy [F]

$$\int (a + b \operatorname{arccosh}(cx))^{\frac{3}{2}} dx = \int (a + b \operatorname{acosh}(cx))^{\frac{3}{2}} dx$$

```
input integrate((a+b*acosh(c*x))**(3/2),x)
```

```
output Integral((a + b*acosh(c*x))**(3/2), x)
```

**3.558.7 Maxima [F]**

$$\int (a + b \operatorname{arccosh}(cx))^{3/2} dx = \int (b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}} dx$$

input `integrate((a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^(3/2), x)`

**3.558.8 Giac [F]**

$$\int (a + b \operatorname{arccosh}(cx))^{3/2} dx = \int (b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}} dx$$

input `integrate((a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^(3/2), x)`

**3.558.9 Mupad [F(-1)]**

Timed out.

$$\int (a + b \operatorname{arccosh}(cx))^{3/2} dx = \int (a + b \operatorname{acosh}(cx))^{3/2} dx$$

input `int((a + b*acosh(c*x))^(3/2),x)`

output `int((a + b*acosh(c*x))^(3/2), x)`



**3.559**  $\int \frac{(a+b\operatorname{arccosh}(cx))^{3/2}}{d+ex^2} dx$

3.559.1 Optimal result	4048
3.559.2 Mathematica [F(-1)]	4048
3.559.3 Rubi [N/A]	4049
3.559.4 Maple [N/A] (verified)	4049
3.559.5 Fricas [F(-2)]	4050
3.559.6 Sympy [N/A]	4050
3.559.7 Maxima [F(-2)]	4050
3.559.8 Giac [N/A]	4051
3.559.9 Mupad [N/A]	4051

**3.559.1 Optimal result**

Integrand size = 22, antiderivative size = 22

$$\int \frac{(a + \operatorname{arccosh}(cx))^{3/2}}{d + ex^2} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^{3/2}}{d + ex^2}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^(3/2)/(e*x^2+d),x)`

**3.559.2 Mathematica [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{arccosh}(cx))^{3/2}}{d + ex^2} dx = \$Aborted$$

input `Integrate[(a + b*ArcCosh[c*x])^(3/2)/(d + e*x^2),x]`

output `$Aborted`

**3.559.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{d + ex^2} dx$$

↓ 6325

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{d + ex^2} dx$$

input `Int[(a + b*ArcCosh[c*x])^(3/2)/(d + e*x^2),x]`

output `$Aborted`

**3.559.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.)^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.559.4 Maple [N/A] (verified)**

Not integrable

Time = 1.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{ex^2 + d} dx$$

input `int((a+b*arccosh(c*x))^(3/2)/(e*x^2+d),x)`

output `int((a+b*arccosh(c*x))^(3/2)/(e*x^2+d),x)`

**3.559.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{(a + \operatorname{barccosh}(cx))^{3/2}}{d + ex^2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^(3/2)/(e*x^2+d),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.559.6 Sympy [N/A]**

Not integrable

Time = 37.43 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{(a + \operatorname{barccosh}(cx))^{3/2}}{d + ex^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^{3/2}}{d + ex^2} dx$$

input `integrate((a+b*acosh(c*x))**(3/2)/(e*x**2+d),x)`

output `Integral((a + b*acosh(c*x))**(3/2)/(d + e*x**2), x)`

**3.559.7 Maxima [F(-2)]**

Exception generated.

$$\int \frac{(a + \operatorname{barccosh}(cx))^{3/2}}{d + ex^2} dx = \text{Exception raised: ValueError}$$

input `integrate((a+b*arccosh(c*x))^(3/2)/(e*x^2+d),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(e>0)', see `assume?` for more details)Is e`

**3.559.8 Giac [N/A]**

Not integrable

Time = 19.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{d + ex^2} dx = \int \frac{(b \operatorname{arccosh}(cx) + a)^{3/2}}{ex^2 + d} dx$$

input `integrate((a+b*arccosh(c*x))^(3/2)/(e*x^2+d),x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^(3/2)/(e*x^2 + d), x)`**3.559.9 Mupad [N/A]**

Not integrable

Time = 3.35 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{d + ex^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^{3/2}}{ex^2 + d} dx$$

input `int((a + b*acosh(c*x))^(3/2)/(d + e*x^2),x)`output `int((a + b*acosh(c*x))^(3/2)/(d + e*x^2), x)`

**3.560**  $\int \frac{(a+b\operatorname{arccosh}(cx))^{3/2}}{(d+ex^2)^2} dx$

3.560.1 Optimal result . . . . . 4052  
 3.560.2 Mathematica [F(-1)] . . . . . 4052  
 3.560.3 Rubi [N/A] . . . . . 4053  
 3.560.4 Maple [N/A] (verified) . . . . . 4053  
 3.560.5 Fricas [F(-2)] . . . . . 4054  
 3.560.6 Sympy [F(-1)] . . . . . 4054  
 3.560.7 Maxima [N/A] . . . . . 4054  
 3.560.8 Giac [N/A] . . . . . 4055  
 3.560.9 Mupad [N/A] . . . . . 4055

**3.560.1 Optimal result**

Integrand size = 22, antiderivative size = 22

$$\int \frac{(a + \operatorname{arccosh}(cx))^{3/2}}{(d + ex^2)^2} dx = \operatorname{Int}\left(\frac{(a + \operatorname{arccosh}(cx))^{3/2}}{(d + ex^2)^2}, x\right)$$

output `Unintegrable((a+b*arccosh(c*x))^(3/2)/(e*x^2+d)^2,x)`

**3.560.2 Mathematica [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{arccosh}(cx))^{3/2}}{(d + ex^2)^2} dx = \$Aborted$$

input `Integrate[(a + b*ArcCosh[c*x])^(3/2)/(d + e*x^2)^2,x]`

output `$Aborted`

**3.560.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{(d + ex^2)^2} dx$$

↓ 6325

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{(d + ex^2)^2} dx$$

input `Int[(a + b*ArcCosh[c*x])^(3/2)/(d + e*x^2)^2,x]`

output `$Aborted`

**3.560.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.560.4 Maple [N/A] (verified)**

Not integrable

Time = 1.22 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{\frac{3}{2}}}{(ex^2 + d)^2} dx$$

input `int((a+b*arccosh(c*x))^(3/2)/(e*x^2+d)^2,x)`

output `int((a+b*arccosh(c*x))^(3/2)/(e*x^2+d)^2,x)`

---

3.560.  $\int \frac{(a+b \operatorname{arccosh}(cx))^{3/2}}{(d+ex^2)^2} dx$

**3.560.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{(a + \operatorname{barccosh}(cx))^{3/2}}{(d + ex^2)^2} dx = \text{Exception raised: TypeError}$$

input `integrate((a+b*arccosh(c*x))^(3/2)/(e*x^2+d)^2,x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.560.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{(a + \operatorname{barccosh}(cx))^{3/2}}{(d + ex^2)^2} dx = \text{Timed out}$$

input `integrate((a+b*acosh(c*x))**(3/2)/(e*x**2+d)**2,x)`

output `Timed out`

**3.560.7 Maxima [N/A]**

Not integrable

Time = 0.71 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + \operatorname{barccosh}(cx))^{3/2}}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}}}{(ex^2 + d)^2} dx$$

input `integrate((a+b*arccosh(c*x))^(3/2)/(e*x^2+d)^2,x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^(3/2)/(e*x^2 + d)^2, x)`

**3.560.8 Giac [N/A]**

Not integrable

Time = 19.26 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{(d + ex^2)^2} dx = \int \frac{(b \operatorname{arcosh}(cx) + a)^{3/2}}{(ex^2 + d)^2} dx$$

input `integrate((a+b*arccosh(c*x))^(3/2)/(e*x^2+d)^2,x, algorithm="giac")`output `integrate((b*arccosh(c*x) + a)^(3/2)/(e*x^2 + d)^2, x)`**3.560.9 Mupad [N/A]**

Not integrable

Time = 3.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \operatorname{arccosh}(cx))^{3/2}}{(d + ex^2)^2} dx = \int \frac{(a + b \operatorname{acosh}(cx))^{3/2}}{(ex^2 + d)^2} dx$$

input `int((a + b*acosh(c*x))^(3/2)/(d + e*x^2)^2,x)`output `int((a + b*acosh(c*x))^(3/2)/(d + e*x^2)^2, x)`



$$3.561 \quad \int \frac{(d+ex^2)^2}{\sqrt{a+b \operatorname{arccosh}(cx)}} dx$$

3.561.1 Optimal result	4057
3.561.2 Mathematica [A] (verified)	4058
3.561.3 Rubi [A] (verified)	4059
3.561.4 Maple [F]	4061
3.561.5 Fricas [F(-2)]	4061
3.561.6 Sympy [F]	4062
3.561.7 Maxima [F]	4062
3.561.8 Giac [F]	4062
3.561.9 Mupad [F(-1)]	4063

## 3.561.1 Optimal result

Integrand size = 22, antiderivative size = 608

$$\begin{aligned}
\int \frac{(d+ex^2)^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx = & -\frac{d^2 e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}} \\
& -\frac{dee^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} \\
& -\frac{e^2 e^{a/b} \sqrt{\pi} \operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16\sqrt{bc^5}} \\
& -\frac{dee^{\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} \\
& -\frac{e^2 e^{\frac{3a}{b}} \sqrt{3\pi} \operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}} \\
& -\frac{e^2 e^{\frac{5a}{b}} \sqrt{\frac{\pi}{5}} \operatorname{erf}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}} \\
& +\frac{d^2 e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}} \\
& +\frac{dee^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} \\
& +\frac{e^2 e^{-\frac{a}{b}} \sqrt{\pi} \operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16\sqrt{bc^5}} \\
& +\frac{dee^{-\frac{3a}{b}} \sqrt{\frac{\pi}{3}} \operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} \\
& +\frac{e^2 e^{-\frac{3a}{b}} \sqrt{3\pi} \operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}} \\
& +\frac{e^2 e^{-\frac{5a}{b}} \sqrt{\frac{\pi}{5}} \operatorname{erfi}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}}
\end{aligned}$$

---

3.561.  $\int \frac{(d+ex^2)^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx$

output

```
-1/160*e^2*exp(5*a/b)*erf(5^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*5^(1/2)
)*Pi^(1/2)/c^5/b^(1/2)+1/160*e^2*erfi(5^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(
1/2))*5^(1/2)*Pi^(1/2)/c^5/exp(5*a/b)/b^(1/2)-1/12*d*e*exp(3*a/b)*erf(3^(1
/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/c^3/b^(1/2)+1/12*d*
e*erfi(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/c^3/exp(
3*a/b)/b^(1/2)-1/2*d^2*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(
1/2)/c/b^(1/2)-1/4*d*e*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(
1/2)/c^3/b^(1/2)-1/16*e^2*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*P
i^(1/2)/c^5/b^(1/2)+1/2*d^2*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2
)/c/exp(a/b)/b^(1/2)+1/4*d*e*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/
2)/c^3/exp(a/b)/b^(1/2)+1/16*e^2*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi
^(1/2)/c^5/exp(a/b)/b^(1/2)-1/32*e^2*exp(3*a/b)*erf(3^(1/2)*(a+b*arccosh(c
*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/c^5/b^(1/2)+1/32*e^2*erfi(3^(1/2)*(a+
b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/c^5/exp(3*a/b)/b^(1/2)
```

### 3.561.2 Mathematica [A] (verified)

Time = 0.81 (sec) , antiderivative size = 530, normalized size of antiderivative = 0.87

$$\int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

$$= e^{-\frac{5a}{b}} \left( 30(8c^4d^2 + 4c^2de + e^2) e^{\frac{6a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) + 3\sqrt{5}e^2 \sqrt{-\frac{a+b \operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{a+b \operatorname{arccosh}(cx)}{b}\right) \right)$$

input `Integrate[(d + e*x^2)^2/Sqrt[a + b*ArcCosh[c*x]],x]`

```
output (30*(8*c^4*d^2 + 4*c^2*d*e + e^2)*E^((6*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, a/b + ArcCosh[c*x]] + 3*Sqrt[5]*e^2*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-5*(a + b*ArcCosh[c*x]))/b] + 40*Sqrt[3]*c^2*d*e*E^((2*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-3*(a + b*ArcCosh[c*x]))/b] + 15*Sqrt[3]*e^2*E^((2*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-3*(a + b*ArcCosh[c*x]))/b] + 240*c^4*d^2*E^((4*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, -((a + b*ArcCosh[c*x])/b)] + 120*c^2*d*e*E^((4*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, -((a + b*ArcCosh[c*x])/b)] + 30*e^2*E^((4*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, -((a + b*ArcCosh[c*x])/b)] + 40*Sqrt[3]*c^2*d*e*E^((8*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (3*(a + b*ArcCosh[c*x]))/b] + 15*Sqrt[3]*e^2*E^((8*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (3*(a + b*ArcCosh[c*x]))/b] + 3*Sqrt[5]*e^2*E^((10*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (5*(a + b*ArcCosh[c*x]))/b])/(480*c^5*E^((5*a)/b)*Sqrt[a + b*ArcCosh[c*x]])
```

### 3.561.3 Rubi [A] (verified)

Time = 1.37 (sec) , antiderivative size = 608, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

↓ 6324

$$\int \left( \frac{d^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} + \frac{2dex^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} + \frac{e^2x^4}{\sqrt{a + b \operatorname{arccosh}(cx)}} \right) dx$$

↓ 2009

$$\begin{aligned}
 & \frac{\sqrt{\pi}e^2e^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16\sqrt{bc^5}} - \frac{\sqrt{3\pi}e^2e^{\frac{3a}{b}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}} - \\
 & \frac{\sqrt{\frac{\pi}{5}}e^2e^{\frac{5a}{b}}\operatorname{erf}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}} + \frac{\sqrt{\pi}e^2e^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{16\sqrt{bc^5}} + \\
 & \frac{\sqrt{3\pi}e^2e^{-\frac{3a}{b}}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}} + \frac{\sqrt{\frac{\pi}{5}}e^2e^{-\frac{5a}{b}}\operatorname{erfi}\left(\frac{\sqrt{5}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{32\sqrt{bc^5}} - \\
 & \frac{\sqrt{\pi}dee^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} - \frac{\sqrt{\frac{\pi}{3}}dee^{\frac{3a}{b}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} + \\
 & \frac{\sqrt{\pi}dee^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} + \frac{\sqrt{\frac{\pi}{3}}dee^{-\frac{3a}{b}}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4\sqrt{bc^3}} - \\
 & \frac{\sqrt{\pi}d^2e^{a/b}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}} + \frac{\sqrt{\pi}d^2e^{-\frac{a}{b}}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}}
 \end{aligned}$$

input `Int[(d + e*x^2)^2/Sqrt[a + b*ArcCosh[c*x]],x]`

output `-1/2*(d^2*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(Sqrt[b]*c) - (d*e*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(4*Sqrt[b]*c^3) - (e^2*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(16*Sqrt[b]*c^5) - (d*e*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(4*Sqrt[b]*c^3) - (e^2*E^((3*a)/b)*Sqrt[3*Pi]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(32*Sqrt[b]*c^5) - (e^2*E^((5*a)/b)*Sqrt[Pi/5]*Erf[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(32*Sqrt[b]*c^5) + (d^2*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(2*Sqrt[b]*c*E^(a/b)) + (d*e*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(4*Sqrt[b]*c^3*E^(a/b)) + (e^2*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(16*Sqrt[b]*c^5*E^(a/b)) + (d*e*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(4*Sqrt[b]*c^3*E^((3*a)/b)) + (e^2*Sqrt[3*Pi]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(32*Sqrt[b]*c^5*E^((3*a)/b)) + (e^2*Sqrt[Pi/5]*Erfi[(Sqrt[5]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(32*Sqrt[b]*c^5*E^((5*a)/b))`

3.561.  $\int \frac{(d+ex^2)^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx$

## 3.561.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

## 3.561.4 Maple [F]

$$\int \frac{(ex^2 + d)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

input `int((e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x)`

output `int((e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x)`

## 3.561.5 Fracas [F(-2)]

Exception generated.

$$\int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \text{Exception raised: TypeError}$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.561.6 Sympy [F]**

$$\int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{acosh}(cx)}} dx$$

input `integrate((e*x**2+d)**2/(a+b*acosh(c*x))**(1/2),x)`

output `Integral((d + e*x**2)**2/sqrt(a + b*acosh(c*x)), x)`

**3.561.7 Maxima [F]**

$$\int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{(ex^2 + d)^2}{\sqrt{b \operatorname{arcosh}(cx) + a}} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate((e*x^2 + d)^2/sqrt(b*arccosh(c*x) + a), x)`

**3.561.8 Giac [F]**

$$\int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{(ex^2 + d)^2}{\sqrt{b \operatorname{arcosh}(cx) + a}} dx$$

input `integrate((e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`

output `integrate((e*x^2 + d)^2/sqrt(b*arccosh(c*x) + a), x)`

**3.561.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{(d + ex^2)^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{(ex^2 + d)^2}{\sqrt{a + b \operatorname{acosh}(cx)}} dx$$

input `int((d + e*x^2)^2/(a + b*acosh(c*x))^(1/2),x)`output `int((d + e*x^2)^2/(a + b*acosh(c*x))^(1/2), x)`



$$3.562 \quad \int \frac{d+ex^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx$$

3.562.1 Optimal result . . . . .	4064
3.562.2 Mathematica [A] (verified) . . . . .	4065
3.562.3 Rubi [A] (verified) . . . . .	4065
3.562.4 Maple [F] . . . . .	4067
3.562.5 Fricas [F(-2)] . . . . .	4067
3.562.6 Sympy [F] . . . . .	4067
3.562.7 Maxima [F] . . . . .	4068
3.562.8 Giac [F] . . . . .	4068
3.562.9 Mupad [F(-1)] . . . . .	4068

### 3.562.1 Optimal result

Integrand size = 20, antiderivative size = 287

$$\int \frac{d+ex^2}{\sqrt{a+b\operatorname{arccosh}(cx)}} dx = -\frac{de^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}} - \frac{ee^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}} - \frac{ee^{\frac{3a}{b}}\sqrt{\frac{\pi}{3}}\operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}} + \frac{de^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}} + \frac{ee^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}} + \frac{ee^{-\frac{3a}{b}}\sqrt{\frac{\pi}{3}}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}}$$

output 
$$\begin{aligned} & -1/24*e*\exp(3*a/b)*\operatorname{erf}(3^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})*3^{1/2}*P \\ & i^{1/2}/c^3/b^{1/2}+1/24*e*\operatorname{erfi}(3^{1/2}*(a+b*\operatorname{arccosh}(c*x))^{1/2}/b^{1/2})* \\ & 3^{1/2}*Pi^{1/2}/c^3/\exp(3*a/b)/b^{1/2}-1/2*d*\exp(a/b)*\operatorname{erf}((a+b*\operatorname{arccosh}(c* \\ & x))^{1/2}/b^{1/2})*Pi^{1/2}/c/b^{1/2}-1/8*e*\exp(a/b)*\operatorname{erf}((a+b*\operatorname{arccosh}(c*x) \\ & )^{1/2}/b^{1/2})*Pi^{1/2}/c^3/b^{1/2}+1/2*d*\operatorname{erfi}((a+b*\operatorname{arccosh}(c*x))^{1/2}/ \\ & b^{1/2})*Pi^{1/2}/c/\exp(a/b)/b^{1/2}+1/8*e*\operatorname{erfi}((a+b*\operatorname{arccosh}(c*x))^{1/2}/b \\ & ^{1/2})*Pi^{1/2}/c^3/\exp(a/b)/b^{1/2} \end{aligned}$$

### 3.562.2 Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 213, normalized size of antiderivative = 0.74

$$\int \frac{d + ex^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

$$= \frac{e^{-\frac{3a}{b}} \left( 3(4c^2d + e) e^{\frac{4a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) + \sqrt{3}e \sqrt{-\frac{a+b \operatorname{arccosh}(cx)}{b}} \Gamma\left(\frac{1}{2}, -\frac{3(a+b \operatorname{arccosh}(cx))}{b}\right) \right)}{24c^3}$$

input `Integrate[(d + e*x^2)/Sqrt[a + b*ArcCosh[c*x]],x]`

output 
$$\begin{aligned} & (3*(4*c^2*d + e)*E^{((4*a)/b)*\operatorname{Sqrt}[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[1/2, a/b + \operatorname{Arc} \\ & \operatorname{Cosh}[c*x]] + \operatorname{Sqrt}[3]*e*\operatorname{Sqrt}[-(a + b*\operatorname{ArcCosh}[c*x])/b]*\operatorname{Gamma}[1/2, (-3*(a + \\ & b*\operatorname{ArcCosh}[c*x])/b] + 3*(4*c^2*d + e)*E^{((2*a)/b)*\operatorname{Sqrt}[-(a + b*\operatorname{ArcCosh}[c \\ & *x])/b]*\operatorname{Gamma}[1/2, -(a + b*\operatorname{ArcCosh}[c*x])/b] + \operatorname{Sqrt}[3]*e*E^{((6*a)/b)*\operatorname{Sqr} \\ & t[a/b + \operatorname{ArcCosh}[c*x]]*\operatorname{Gamma}[1/2, (3*(a + b*\operatorname{ArcCosh}[c*x])/b)]/(24*c^3*E^{(( \\ & 3*a)/b)*\operatorname{Sqrt}[a + b*\operatorname{ArcCosh}[c*x]]]) \end{aligned}$$

### 3.562.3 Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 287, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{d + ex^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

---

3.562.  $\int \frac{d+ex^2}{\sqrt{a+b \operatorname{arccosh}(cx)}} dx$

$$\begin{aligned}
 & \int \left( \frac{d}{\sqrt{a + \operatorname{barccosh}(cx)}} + \frac{ex^2}{\sqrt{a + \operatorname{barccosh}(cx)}} \right) dx \\
 & \quad \downarrow \text{6324} \\
 & \quad \downarrow \text{2009} \\
 & -\frac{\sqrt{\pi} e e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}} - \frac{\sqrt{\frac{\pi}{3}} e e^{\frac{3a}{b}} \operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}} + \\
 & \frac{\sqrt{\pi} e e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}} + \frac{\sqrt{\frac{\pi}{3}} e e^{-\frac{3a}{b}} \operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{8\sqrt{bc^3}} - \\
 & \frac{\sqrt{\pi} d e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}} + \frac{\sqrt{\pi} d e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a + \operatorname{barccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}}
 \end{aligned}$$

input `Int[(d + e*x^2)/Sqrt[a + b*ArcCosh[c*x]],x]`

output `-1/2*(d*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(Sqrt[b]*c) - (e*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(8*Sqrt[b]*c^3) - (e*E^((3*a)/b)*Sqrt[Pi/3]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(8*Sqrt[b]*c^3) + (d*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(2*Sqrt[b]*c*E^(a/b)) + (e*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]]/(8*Sqrt[b]*c^3*E^(a/b)) + (e*Sqrt[Pi/3]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]]/(8*Sqrt[b]*c^3*E^((3*a)/b)))`

### 3.562.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

**3.562.4 Maple [F]**

$$\int \frac{e x^2 + d}{\sqrt{a + b \operatorname{arccosh}(c x)}} dx$$

input `int((e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x)`

output `int((e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x)`

**3.562.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{d + e x^2}{\sqrt{a + b \operatorname{arccosh}(c x)}} dx = \text{Exception raised: TypeError}$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.562.6 Sympy [F]**

$$\int \frac{d + e x^2}{\sqrt{a + b \operatorname{arccosh}(c x)}} dx = \int \frac{d + e x^2}{\sqrt{a + b \operatorname{acosh}(c x)}} dx$$

input `integrate((e*x**2+d)/(a+b*acosh(c*x))**(1/2),x)`

output `Integral((d + e*x**2)/sqrt(a + b*acosh(c*x)), x)`

**3.562.7 Maxima [F]**

$$\int \frac{d + ex^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{ex^2 + d}{\sqrt{b \operatorname{arccosh}(cx) + a}} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate((e*x^2 + d)/sqrt(b*arccosh(c*x) + a), x)`

**3.562.8 Giac [F]**

$$\int \frac{d + ex^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{ex^2 + d}{\sqrt{b \operatorname{arccosh}(cx) + a}} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`

output `integrate((e*x^2 + d)/sqrt(b*arccosh(c*x) + a), x)`

**3.562.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{d + ex^2}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{ex^2 + d}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

input `int((d + e*x^2)/(a + b*arccosh(c*x))^(1/2),x)`

output `int((d + e*x^2)/(a + b*arccosh(c*x))^(1/2), x)`

**3.563**  $\int \frac{1}{\sqrt{a+b\text{arccosh}(cx)}} dx$

3.563.1 Optimal result . . . . . 4069  
 3.563.2 Mathematica [A] (verified) . . . . . 4069  
 3.563.3 Rubi [C] (verified) . . . . . 4070  
 3.563.4 Maple [F] . . . . . 4072  
 3.563.5 Fracas [F(-2)] . . . . . 4073  
 3.563.6 Sympy [F] . . . . . 4073  
 3.563.7 Maxima [F] . . . . . 4073  
 3.563.8 Giac [F] . . . . . 4074  
 3.563.9 Mupad [F(-1)] . . . . . 4074

**3.563.1 Optimal result**

Integrand size = 12, antiderivative size = 88

$$\int \frac{1}{\sqrt{a+b\text{arccosh}(cx)}} dx = -\frac{e^{a/b}\sqrt{\pi}\text{erf}\left(\frac{\sqrt{a+b\text{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}} + \frac{e^{-\frac{a}{b}}\sqrt{\pi}\text{erfi}\left(\frac{\sqrt{a+b\text{arccosh}(cx)}}{\sqrt{b}}\right)}{2\sqrt{bc}}$$

output `-1/2*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c/b^(1/2)+1/2*erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/c/exp(a/b)/b^(1/2)`

**3.563.2 Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.14

$$\int \frac{1}{\sqrt{a+b\text{arccosh}(cx)}} dx = \frac{e^{-\frac{a}{b}}\left(e^{\frac{2a}{b}}\sqrt{\frac{a}{b}+\text{arccosh}(cx)}\Gamma\left(\frac{1}{2},\frac{a}{b}+\text{arccosh}(cx)\right)+\sqrt{-\frac{a+b\text{arccosh}(cx)}{b}}\Gamma\left(\frac{1}{2},-\frac{a+b\text{arccosh}(cx)}{b}\right)\right)}{2c\sqrt{a+b\text{arccosh}(cx)}}$$

input `Integrate[1/Sqrt[a + b*ArcCosh[c*x]],x]`

output  $(E^{((2*a)/b)*\text{Sqrt}[a/b + \text{ArcCosh}[c*x]]*\text{Gamma}[1/2, a/b + \text{ArcCosh}[c*x]] + \text{Sqrt}[-((a + b*\text{ArcCosh}[c*x])/b)]*\text{Gamma}[1/2, -((a + b*\text{ArcCosh}[c*x])/b)])/(2*c*E^{(a/b)*\text{Sqrt}[a + b*\text{ArcCosh}[c*x]])}$

### 3.563.3 Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.09, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6296, 25, 3042, 26, 3789, 2611, 2633, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

↓ 6296

$$\frac{\int -\frac{\sinh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a + b \operatorname{arccosh}(cx)}} d(a + b \operatorname{arccosh}(cx))}{bc}$$

↓ 25

$$\frac{\int \frac{\sinh\left(\frac{a}{b} - \frac{a + b \operatorname{arccosh}(cx)}{b}\right)}{\sqrt{a + b \operatorname{arccosh}(cx)}} d(a + b \operatorname{arccosh}(cx))}{bc}$$

↓ 3042

$$\frac{\int -\frac{i \sin\left(\frac{ia}{b} - \frac{i(a + b \operatorname{arccosh}(cx))}{b}\right)}{\sqrt{a + b \operatorname{arccosh}(cx)}} d(a + b \operatorname{arccosh}(cx))}{bc}$$

↓ 26

$$\frac{i \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a + b \operatorname{arccosh}(cx))}{b}\right)}{\sqrt{a + b \operatorname{arccosh}(cx)}} d(a + b \operatorname{arccosh}(cx))}{bc}$$

↓ 3789

---

3.563.  $\int \frac{1}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$

$$\begin{aligned}
& \frac{i \left( \frac{1}{2} i \int \frac{e^{-\operatorname{arccosh}(cx)}}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx)) - \frac{1}{2} i \int \frac{e^{\operatorname{arccosh}(cx)}}{\sqrt{a+b\operatorname{arccosh}(cx)}} d(a+b\operatorname{arccosh}(cx)) \right)}{bc} \\
& \quad \downarrow \text{2611} \\
& \frac{i \left( i \int e^{\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}} d\sqrt{a+b\operatorname{arccosh}(cx)} - i \int e^{\frac{a+b\operatorname{arccosh}(cx)}{b} - \frac{a}{b}} d\sqrt{a+b\operatorname{arccosh}(cx)} \right)}{bc} \\
& \quad \downarrow \text{2633} \\
& \frac{i \left( i \int e^{\frac{a}{b} - \frac{a+b\operatorname{arccosh}(cx)}{b}} d\sqrt{a+b\operatorname{arccosh}(cx)} - \frac{1}{2} i \sqrt{\pi} \sqrt{b} e^{-\frac{a}{b}} \operatorname{erfi} \left( \frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)}{bc} \\
& \quad \downarrow \text{2634} \\
& \frac{i \left( \frac{1}{2} i \sqrt{\pi} \sqrt{b} e^{a/b} \operatorname{erf} \left( \frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) - \frac{1}{2} i \sqrt{\pi} \sqrt{b} e^{-\frac{a}{b}} \operatorname{erfi} \left( \frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)}{bc}
\end{aligned}$$

input `Int[1/Sqrt[a + b*ArcCosh[c*x]], x]`

output `(I*((I/2)*Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]] - (I/2)*Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/E^(a/b)) / (b*c)`

### 3.563.3.1 Defintions of rubi rules used

rule 25 `Int[-(F x_), x_Symbol] :> Simp[Identity[-1] Int[F x, x], x]`

rule 26 `Int[(Complex[0, a_])*(F x_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[F x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2611 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`



rule 2633 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_))2), x_Symbol] := Simp[F^a*Sqrt  
[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{  
F, a, b, c, d}, x] && PosQ[b]`

rule 2634 `Int[(F_)^((a_) + (b_)*((c_) + (d_)*(x_))2), x_Symbol] := Simp[F^a*Sqrt  
[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; Fr  
eeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear  
Q[u, x]`

rule 3789 `Int[((c_) + (d_)*(x_))(m_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[I  
/2 Int[(c + d*x)m/E^(I*(e + f*x)), x], x] - Simp[I/2 Int[(c + d*x)m*E  
^(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x]`

rule 6296 `Int[((a_) + ArcCosh[(c_)*(x_)])*(b_)(n_), x_Symbol] := Simp[1/(b*c) S  
ubst[Int[xn*Sinh[-a/b + x/b], x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a,  
b, c, n}, x]`

### 3.563.4 Maple [F]

$$\int \frac{1}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

input `int(1/(a+b*arccosh(c*x))^(1/2),x)`

output `int(1/(a+b*arccosh(c*x))^(1/2),x)`

**3.563.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{1}{\sqrt{a + \operatorname{barccosh}(cx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.563.6 Sympy [F]**

$$\int \frac{1}{\sqrt{a + \operatorname{barccosh}(cx)}} dx = \int \frac{1}{\sqrt{a + b \operatorname{acosh}(cx)}} dx$$

input `integrate(1/(a+b*acosh(c*x))**(1/2),x)`

output `Integral(1/sqrt(a + b*acosh(c*x)), x)`

**3.563.7 Maxima [F]**

$$\int \frac{1}{\sqrt{a + \operatorname{barccosh}(cx)}} dx = \int \frac{1}{\sqrt{b \operatorname{arcosh}(cx) + a}} dx$$

input `integrate(1/(a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate(1/sqrt(b*arccosh(c*x) + a), x)`

**3.563.8 Giac [F]**

$$\int \frac{1}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{1}{\sqrt{b \operatorname{arcosh}(cx) + a}} dx$$

input `integrate(1/(a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`

output `integrate(1/sqrt(b*arccosh(c*x) + a), x)`

**3.563.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{\sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{1}{\sqrt{a + b \operatorname{acosh}(cx)}} dx$$

input `int(1/(a + b*acosh(c*x))^(1/2),x)`

output `int(1/(a + b*acosh(c*x))^(1/2), x)`

$$3.564 \quad \int \frac{1}{(d+ex^2)\sqrt{a+b\operatorname{arccosh}(cx)}} dx$$

3.564.1 Optimal result	4075
3.564.2 Mathematica [N/A]	4075
3.564.3 Rubi [N/A]	4076
3.564.4 Maple [N/A] (verified)	4076
3.564.5 Fricas [F(-2)]	4077
3.564.6 Sympy [N/A]	4077
3.564.7 Maxima [N/A]	4077
3.564.8 Giac [N/A]	4078
3.564.9 Mupad [N/A]	4078

### 3.564.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d+ex^2)\sqrt{a+b\operatorname{arccosh}(cx)}} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)\sqrt{a+b\operatorname{arccosh}(cx)}}, x\right)$$

output `Unintegrable(1/(e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x)`

### 3.564.2 Mathematica [N/A]

Not integrable

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d+ex^2)\sqrt{a+b\operatorname{arccosh}(cx)}} dx = \int \frac{1}{(d+ex^2)\sqrt{a+b\operatorname{arccosh}(cx)}} dx$$

input `Integrate[1/((d + e*x^2)*Sqrt[a + b*ArcCosh[c*x]]),x]`

output `Integrate[1/((d + e*x^2)*Sqrt[a + b*ArcCosh[c*x]]), x]`

**3.564.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2) \sqrt{a + \operatorname{barccosh}(cx)}} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2) \sqrt{a + \operatorname{barccosh}(cx)}} dx$$

input `Int[1/((d + e*x^2)*Sqrt[a + b*ArcCosh[c*x]]),x]`

output `$Aborted`

**3.564.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.564.4 Maple [N/A] (verified)**

Not integrable

Time = 1.36 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d) \sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

input `int(1/(e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x)`

output `int(1/(e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x)`

**3.564.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{1}{(d + ex^2) \sqrt{a + \operatorname{barccosh}(cx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.564.6 Sympy [N/A]**

Not integrable

Time = 5.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(d + ex^2) \sqrt{a + \operatorname{barccosh}(cx)}} dx = \int \frac{1}{\sqrt{a + b \operatorname{acosh}(cx)} (d + ex^2)} dx$$

input `integrate(1/(e*x**2+d)/(a+b*acosh(c*x))**(1/2),x)`

output `Integral(1/(sqrt(a + b*acosh(c*x))*(d + e*x**2)), x)`

**3.564.7 Maxima [N/A]**

Not integrable

Time = 0.50 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2) \sqrt{a + \operatorname{barccosh}(cx)}} dx = \int \frac{1}{(ex^2 + d) \sqrt{b \operatorname{arcosh}(cx) + a}} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate(1/((e*x^2 + d)*sqrt(b*arccosh(c*x) + a)), x)`

**3.564.8 Giac [N/A]**

Not integrable

Time = 12.92 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{1}{(ex^2 + d) \sqrt{b \operatorname{arcosh}(cx) + a}} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)*sqrt(b*arccosh(c*x) + a)), x)`**3.564.9 Mupad [N/A]**

Not integrable

Time = 3.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2) \sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{1}{\sqrt{a + b \operatorname{acosh}(cx)} (ex^2 + d)} dx$$

input `int(1/((a + b*acosh(c*x))^(1/2)*(d + e*x^2)),x)`output `int(1/((a + b*acosh(c*x))^(1/2)*(d + e*x^2)), x)`

$$3.565 \quad \int \frac{1}{(d+ex^2)^2 \sqrt{a+b\operatorname{arccosh}(cx)}} dx$$

3.565.1 Optimal result	4079
3.565.2 Mathematica [N/A]	4079
3.565.3 Rubi [N/A]	4080
3.565.4 Maple [N/A] (verified)	4080
3.565.5 Fricas [F(-2)]	4081
3.565.6 Sympy [N/A]	4081
3.565.7 Maxima [N/A]	4081
3.565.8 Giac [N/A]	4082
3.565.9 Mupad [N/A]	4082

### 3.565.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d+ex^2)^2 \sqrt{a+b\operatorname{arccosh}(cx)}} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)^2 \sqrt{a+b\operatorname{arccosh}(cx)}}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x)`

### 3.565.2 Mathematica [N/A]

Not integrable

Time = 0.23 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d+ex^2)^2 \sqrt{a+b\operatorname{arccosh}(cx)}} dx = \int \frac{1}{(d+ex^2)^2 \sqrt{a+b\operatorname{arccosh}(cx)}} dx$$

input `Integrate[1/((d + e*x^2)^2*Sqrt[a + b*ArcCosh[c*x]]),x]`

output `Integrate[1/((d + e*x^2)^2*Sqrt[a + b*ArcCosh[c*x]]), x]`



**3.565.3 Rubi [N/A]**

Not integrable

Time = 0.22 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^2 \sqrt{a + \operatorname{arccosh}(cx)}} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^2 \sqrt{a + \operatorname{arccosh}(cx)}} dx$$

input `Int[1/((d + e*x^2)^2*Sqrt[a + b*ArcCosh[c*x]]),x]`

output `$Aborted`

**3.565.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.565.4 Maple [N/A] (verified)**

Not integrable

Time = 1.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d)^2 \sqrt{a + b \operatorname{arccosh}(cx)}} dx$$

input `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x)`

output `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x)`

**3.565.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{1}{(d + ex^2)^2 \sqrt{a + \operatorname{barccosh}(cx)}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.565.6 Sympy [N/A]**

Not integrable

Time = 151.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^2 \sqrt{a + \operatorname{barccosh}(cx)}} dx = \int \frac{1}{\sqrt{a + b \operatorname{acosh}(cx)} (d + ex^2)^2} dx$$

input `integrate(1/(e*x**2+d)**2/(a+b*acosh(c*x))**(1/2),x)`

output `Integral(1/(sqrt(a + b*acosh(c*x))*(d + e*x**2)**2), x)`

**3.565.7 Maxima [N/A]**

Not integrable

Time = 0.53 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^2 \sqrt{a + \operatorname{barccosh}(cx)}} dx = \int \frac{1}{(ex^2 + d)^2 \sqrt{b \operatorname{arcosh}(cx) + a}} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x, algorithm="maxima")`

output `integrate(1/((e*x^2 + d)^2*sqrt(b*arccosh(c*x) + a)), x)`

---

3.565.  $\int \frac{1}{(d+ex^2)^2 \sqrt{a+\operatorname{barccosh}(cx)}} dx$

**3.565.8 Giac [N/A]**

Not integrable

Time = 12.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^2 \sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{1}{(ex^2 + d)^2 \sqrt{b \operatorname{arccosh}(cx) + a}} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(1/2),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)^2*sqrt(b*arccosh(c*x) + a)), x)`**3.565.9 Mupad [N/A]**

Not integrable

Time = 3.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^2 \sqrt{a + b \operatorname{arccosh}(cx)}} dx = \int \frac{1}{\sqrt{a + b \operatorname{acosh}(cx)} (ex^2 + d)^2} dx$$

input `int(1/((a + b*acosh(c*x))^(1/2)*(d + e*x^2)^2),x)`output `int(1/((a + b*acosh(c*x))^(1/2)*(d + e*x^2)^2), x)`

### 3.566 $\int \frac{d+ex^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.566.1 Optimal result . . . . .	4083
3.566.2 Mathematica [A] (warning: unable to verify) . . . . .	4084
3.566.3 Rubi [A] (verified) . . . . .	4084
3.566.4 Maple [F] . . . . .	4086
3.566.5 Fracas [F(-2)] . . . . .	4086
3.566.6 Sympy [F] . . . . .	4086
3.566.7 Maxima [F] . . . . .	4087
3.566.8 Giac [F] . . . . .	4087
3.566.9 Mupad [F(-1)] . . . . .	4087

#### 3.566.1 Optimal result

Integrand size = 20, antiderivative size = 358

$$\int \frac{d+ex^2}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = -\frac{2d\sqrt{-1+cx}\sqrt{1+cx}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} - \frac{2ex^2\sqrt{-1+cx}\sqrt{1+cx}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}}$$

$$+ \frac{de^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}c} + \frac{ee^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3}$$

$$+ \frac{ee^{\frac{3a}{b}}\sqrt{3\pi}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3} + \frac{de^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}c}$$

$$+ \frac{ee^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3} + \frac{ee^{-\frac{3a}{b}}\sqrt{3\pi}\operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3}$$

output

```
d*exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c+1/4*e*
exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^3+d*erfi
((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c/exp(a/b)+1/4*e*erfi(
(a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c^3/exp(a/b)+1/4*e*exp(
3*a/b)*erf(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1/2)/b^(3
/2)/c^3+1/4*e*erfi(3^(1/2)*(a+b*arccosh(c*x))^(1/2)/b^(1/2))*3^(1/2)*Pi^(1
/2)/b^(3/2)/c^3/exp(3*a/b)-2*d*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(a+b*arccos
h(c*x))^(1/2)-2*e*x^2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(a+b*arccosh(c*x))^(
1/2)
```

**3.566.2 Mathematica [A] (warning: unable to verify)**

Time = 1.26 (sec) , antiderivative size = 268, normalized size of antiderivative = 0.75

$$\int \frac{d + ex^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \frac{e^{-\frac{3a}{b}} \left( - \left( (4c^2d + e) e^{\frac{4a}{b}} \sqrt{\frac{a}{b} + \operatorname{arccosh}(cx)} \Gamma\left(\frac{1}{2}, \frac{a}{b} + \operatorname{arccosh}(cx)\right) \right) \right)}{\dots} + \sqrt{3}e\sqrt{\dots}$$

input `Integrate[(d + e*x^2)/(a + b*ArcCosh[c*x])^(3/2),x]`

output `(-((4*c^2*d + e)*E^((4*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, a/b + ArcCosh[c*x]]) + Sqrt[3]*e*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, (-3*(a + b*ArcCosh[c*x])/b)] + (4*c^2*d + e)*E^((2*a)/b)*Sqrt[-((a + b*ArcCosh[c*x])/b)]*Gamma[1/2, -((a + b*ArcCosh[c*x])/b)] - E^((3*a)/b)*(2*(4*c^2*d + e)*Sqrt[(-1 + c*x)/(1 + c*x)]*(1 + c*x) + Sqrt[3]*e*E^((3*a)/b)*Sqrt[a/b + ArcCosh[c*x]]*Gamma[1/2, (3*(a + b*ArcCosh[c*x])/b)] + 2*e*Sinh[3*ArcCosh[c*x]]))/(4*b*c^3*E^((3*a)/b)*Sqrt[a + b*ArcCosh[c*x]])`

**3.566.3 Rubi [A] (verified)**Time = 1.05 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {6324, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{d + ex^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx$$

$$\downarrow \text{6324}$$

$$\int \left( \frac{d}{(a + b \operatorname{arccosh}(cx))^{3/2}} + \frac{ex^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{\sqrt{\pi} e e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a+\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3} + \frac{\sqrt{3\pi} e e^{\frac{3a}{b}} \operatorname{erf}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3} + \\
& \frac{\sqrt{\pi} e e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3} + \frac{\sqrt{3\pi} e e^{-\frac{3a}{b}} \operatorname{erfi}\left(\frac{\sqrt{3}\sqrt{a+\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{4b^{3/2}c^3} + \\
& \frac{\sqrt{\pi} d e^{a/b} \operatorname{erf}\left(\frac{\sqrt{a+\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}c} + \frac{\sqrt{\pi} d e^{-\frac{a}{b}} \operatorname{erfi}\left(\frac{\sqrt{a+\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}c} - \frac{2d\sqrt{cx-1}\sqrt{cx+1}}{bc\sqrt{a+\operatorname{arccosh}(cx)}} - \\
& \frac{2ex^2\sqrt{cx-1}\sqrt{cx+1}}{bc\sqrt{a+\operatorname{arccosh}(cx)}}
\end{aligned}$$

input `Int[(d + e*x^2)/(a + b*ArcCosh[c*x])^(3/2),x]`

output `(-2*d*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(b*c*Sqrt[a + b*ArcCosh[c*x]]) - (2*e*x^2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(b*c*Sqrt[a + b*ArcCosh[c*x]]) + (d*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(b^(3/2)*c) + (e*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(4*b^(3/2)*c^3) + (e*E^((3*a)/b)*Sqrt[3*Pi]*Erf[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(4*b^(3/2)*c^3) + (d*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(b^(3/2)*c*E^(a/b)) + (e*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(4*b^(3/2)*c^3*E^(a/b)) + (e*Sqrt[3*Pi]*Erfi[(Sqrt[3]*Sqrt[a + b*ArcCosh[c*x]])/Sqrt[b]])/(4*b^(3/2)*c^3*E^((3*a)/b))`

### 3.566.3.1 Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 6324 `Int[((a_.) + ArcCosh[(c_.)*(x_)])*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*ArcCosh[c*x])^n, (d + e*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, n}, x] && NeQ[c^2*d + e, 0] && IntegerQ[p] && (p > 0 || IGtQ[n, 0])`

**3.566.4 Maple [F]**

$$\int \frac{e x^2 + d}{(a + b \operatorname{arccosh}(c x))^{3/2}} dx$$

input `int((e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

output `int((e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

**3.566.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{d + e x^2}{(a + b \operatorname{arccosh}(c x))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.566.6 Sympy [F]**

$$\int \frac{d + e x^2}{(a + b \operatorname{arccosh}(c x))^{3/2}} dx = \int \frac{d + e x^2}{(a + b \operatorname{acosh}(c x))^{3/2}} dx$$

input `integrate((e*x**2+d)/(a+b*acosh(c*x))**(3/2),x)`

output `Integral((d + e*x**2)/(a + b*acosh(c*x))**(3/2), x)`

**3.566.7 Maxima [F]**

$$\int \frac{d + ex^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{ex^2 + d}{(b \operatorname{arccosh}(cx) + a)^{3/2}} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((e*x^2 + d)/(b*arccosh(c*x) + a)^(3/2), x)`

**3.566.8 Giac [F]**

$$\int \frac{d + ex^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{ex^2 + d}{(b \operatorname{arccosh}(cx) + a)^{3/2}} dx$$

input `integrate((e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `integrate((e*x^2 + d)/(b*arccosh(c*x) + a)^(3/2), x)`

**3.566.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{d + ex^2}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{ex^2 + d}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int((d + e*x^2)/(a + b*acosh(c*x))^(3/2),x)`

output `int((d + e*x^2)/(a + b*acosh(c*x))^(3/2), x)`



**3.567**  $\int \frac{1}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.567.1 Optimal result . . . . . 4088  
 3.567.2 Mathematica [F] . . . . . 4088  
 3.567.3 Rubi [A] (verified) . . . . . 4089  
 3.567.4 Maple [F] . . . . . 4091  
 3.567.5 Fricas [F(-2)] . . . . . 4092  
 3.567.6 Sympy [F] . . . . . 4092  
 3.567.7 Maxima [F] . . . . . 4092  
 3.567.8 Giac [F] . . . . . 4093  
 3.567.9 Mupad [F(-1)] . . . . . 4093

**3.567.1 Optimal result**

Integrand size = 12, antiderivative size = 120

$$\int \frac{1}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = -\frac{2\sqrt{-1+cx}\sqrt{1+cx}}{bc\sqrt{a+b\operatorname{arccosh}(cx)}} + \frac{e^{a/b}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}c} + \frac{e^{-\frac{a}{b}}\sqrt{\pi}\operatorname{erfi}\left(\frac{\sqrt{a+b\operatorname{arccosh}(cx)}}{\sqrt{b}}\right)}{b^{3/2}c}$$

output `exp(a/b)*erf((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c+erfi((a+b*arccosh(c*x))^(1/2)/b^(1/2))*Pi^(1/2)/b^(3/2)/c/exp(a/b)-2*(c*x-1)^(1/2)*(c*x+1)^(1/2)/b/c/(a+b*arccosh(c*x))^(1/2)`

**3.567.2 Mathematica [F]**

$$\int \frac{1}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

input `Integrate[(a + b*ArcCosh[c*x])^(-3/2), x]`

output `Integrate[(a + b*ArcCosh[c*x])^(-3/2), x]`

**3.567.3 Rubi [A] (verified)**

Time = 0.81 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.08, number of steps used = 9, number of rules used = 8,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$ , Rules used = {6295, 6368, 3042, 3788, 26, 2611, 2633, 2634}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(a + \operatorname{barccosh}(cx))^{3/2}} dx \\
 & \quad \downarrow \text{6295} \\
 & \frac{2c \int \frac{x}{\sqrt{cx-1}\sqrt{cx+1}\sqrt{a+\operatorname{barccosh}(cx)}} dx}{b} - \frac{2\sqrt{cx-1}\sqrt{cx+1}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\
 & \quad \downarrow \text{6368} \\
 & \frac{2 \int \frac{\cosh\left(\frac{a}{b} - \frac{a+\operatorname{barccosh}(cx)}{b}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{a+\operatorname{barccosh}(cx)}}}{b^2c} - \frac{2\sqrt{cx-1}\sqrt{cx+1}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\
 & \quad \downarrow \text{3042} \\
 & -\frac{2\sqrt{cx-1}\sqrt{cx+1}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} + \frac{2 \int \frac{\sin\left(\frac{ia}{b} - \frac{i(a+\operatorname{barccosh}(cx))}{b} + \frac{\pi}{2}\right) d(a + \operatorname{barccosh}(cx))}{\sqrt{a+\operatorname{barccosh}(cx)}}}{b^2c} \\
 & \quad \downarrow \text{3788} \\
 & -\frac{2\sqrt{cx-1}\sqrt{cx+1}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} + \\
 & \frac{2 \left( \frac{1}{2}i \int -\frac{ie^{-\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) - \frac{1}{2}i \int \frac{ie^{\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) \right)}{b^2c} \\
 & \quad \downarrow \text{26} \\
 & \frac{2 \left( \frac{1}{2} \int \frac{e^{-\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) + \frac{1}{2} \int \frac{e^{\operatorname{arccosh}(cx)}}{\sqrt{a+\operatorname{barccosh}(cx)}} d(a + \operatorname{barccosh}(cx)) \right)}{b^2c} - \\
 & \frac{2\sqrt{cx-1}\sqrt{cx+1}}{bc\sqrt{a+\operatorname{barccosh}(cx)}} \\
 & \quad \downarrow \text{2611}
 \end{aligned}$$

---

3.567.  $\int \frac{1}{(a+\operatorname{barccosh}(cx))^{3/2}} dx$

$$\begin{aligned}
& \frac{2 \left( \int e^{\frac{a}{b} - \frac{a + \operatorname{arccosh}(cx)}{b}} d\sqrt{a + \operatorname{arccosh}(cx)} + \int e^{\frac{a + \operatorname{arccosh}(cx)}{b} - \frac{a}{b}} d\sqrt{a + \operatorname{arccosh}(cx)} \right)}{\frac{b^2 c}{2\sqrt{cx-1}\sqrt{cx+1}} bc\sqrt{a + \operatorname{arccosh}(cx)}} \\
& \quad \downarrow \text{2633} \\
& \frac{2 \left( \int e^{\frac{a}{b} - \frac{a + \operatorname{arccosh}(cx)}{b}} d\sqrt{a + \operatorname{arccosh}(cx)} + \frac{1}{2} \sqrt{\pi} \sqrt{b} e^{-\frac{a}{b}} \operatorname{erfi} \left( \frac{\sqrt{a + \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)}{\frac{b^2 c}{2\sqrt{cx-1}\sqrt{cx+1}} bc\sqrt{a + \operatorname{arccosh}(cx)}} \\
& \quad \downarrow \text{2634} \\
& \frac{2 \left( \frac{1}{2} \sqrt{\pi} \sqrt{b} e^{a/b} \operatorname{erf} \left( \frac{\sqrt{a + \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) + \frac{1}{2} \sqrt{\pi} \sqrt{b} e^{-\frac{a}{b}} \operatorname{erfi} \left( \frac{\sqrt{a + \operatorname{arccosh}(cx)}}{\sqrt{b}} \right) \right)}{\frac{b^2 c}{2\sqrt{cx-1}\sqrt{cx+1}} bc\sqrt{a + \operatorname{arccosh}(cx)}}
\end{aligned}$$

input `Int[(a + b*ArcCosh[c*x])^(-3/2), x]`

output `(-2*Sqrt[-1 + c*x]*Sqrt[1 + c*x])/(b*c*Sqrt[a + b*ArcCosh[c*x]]) + (2*((Sqrt[b]*E^(a/b)*Sqrt[Pi]*Erf[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/2 + (Sqrt[b]*Sqrt[Pi]*Erfi[Sqrt[a + b*ArcCosh[c*x]]/Sqrt[b]])/(2*E^(a/b)))/(b^2*c)`

### 3.567.3.1 Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_]*(F_x_), x_Symbol] := Simp[(Complex[Identity[0], a]) Int[F_x, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 2611 `Int[(F_)^((g_.)*((e_.) + (f_.)*(x_)))/Sqrt[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[2/d Subst[Int[F^(g*(e - c*(f/d)) + f*g*(x^2/d)), x], x, Sqrt[c + d*x]], x] /; FreeQ[{F, c, d, e, f, g}, x] && !TrueQ[$UseGamma]`

rule 2633 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erfi[(c + d*x)*Rt[b*Log[F], 2]]/(2*d*Rt[b*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && PosQ[b]`

rule 2634 `Int[(F_)^((a_.) + (b_.)*((c_.) + (d_.)*(x_))2), x_Symbol] := Simp[F^a*Sqrt[Pi]*(Erf[(c + d*x)*Rt[(-b)*Log[F], 2]]/(2*d*Rt[(-b)*Log[F], 2])), x] /; FreeQ[{F, a, b, c, d}, x] && NegQ[b]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3788 `Int[((c_.) + (d_.)*(x_))(m_.)*sin[(e_.) + Pi*(k_.) + (f_.)*(x_)], x_Symbol] := Simp[I/2 Int[(c + d*x)m/(E(I*k*Pi)*E(I*(e + f*x))), x], x] - Simp[I/2 Int[(c + d*x)m*E(I*k*Pi)*E(I*(e + f*x)), x], x] /; FreeQ[{c, d, e, f, m}, x] && IntegerQ[2*k]`

rule 6295 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_), x_Symbol] := Simp[Sqrt[1 + c*x]*Sqrt[-1 + c*x]*((a + b*ArcCosh[c*x])(n + 1)/(b*c*(n + 1))), x] - Simp[c/(b*(n + 1)) Int[x*((a + b*ArcCosh[c*x])(n + 1)/(Sqrt[1 + c*x]*Sqrt[-1 + c*x])), x], x] /; FreeQ[{a, b, c}, x] && LtQ[n, -1]`

rule 6368 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))(n_.)(x_)(m_.)((d1_) + (e1_.)*(x_)(p_.)((d2_) + (e2_.)*(x_)(p_.)), x_Symbol] := Simp[(1/(b*c(m + 1)))*Simp[(d1 + e1*x)p/(1 + c*x)p]*Simp[(d2 + e2*x)p/(-1 + c*x)p Subst[Int[xn*Cosh[-a/b + x/b]m*Sinh[-a/b + x/b](2*p + 1), x], x, a + b*ArcCosh[c*x]], x] /; FreeQ[{a, b, c, d1, e1, d2, e2, n}, x] && EqQ[e1, c*d1] && EqQ[e2, (-c)*d2] && IGtQ[p + 3/2, 0] && IGtQ[m, 0]`

### 3.567.4 Maple [F]

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^{\frac{3}{2}}} dx$$

input `int(1/(a+b*arccosh(c*x))^(3/2),x)`

output `int(1/(a+b*arccosh(c*x))^(3/2),x)`

**3.567.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.567.6 Sympy [F]**

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^{\frac{3}{2}}} dx$$

input `integrate(1/(a+b*acosh(c*x))**(3/2),x)`

output `Integral((a + b*acosh(c*x))**(-3/2), x)`

**3.567.7 Maxima [F]**

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(b \operatorname{arcosh}(cx) + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate((b*arccosh(c*x) + a)^(-3/2), x)`

**3.567.8 Giac [F]**

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(b \operatorname{arcosh}(cx) + a)^{3/2}} dx$$

input `integrate(1/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`

output `integrate((b*arccosh(c*x) + a)^(-3/2), x)`

**3.567.9 Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{(a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^{3/2}} dx$$

input `int(1/(a + b*acosh(c*x))^(3/2),x)`

output `int(1/(a + b*acosh(c*x))^(3/2), x)`

$$3.568 \quad \int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

3.568.1 Optimal result	4094
3.568.2 Mathematica [N/A]	4094
3.568.3 Rubi [N/A]	4095
3.568.4 Maple [N/A] (verified)	4095
3.568.5 Fricas [F(-2)]	4096
3.568.6 Sympy [N/A]	4096
3.568.7 Maxima [N/A]	4096
3.568.8 Giac [N/A]	4097
3.568.9 Mupad [N/A]	4097

### 3.568.1 Optimal result

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^{3/2}} dx = \operatorname{Int}\left(\frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^{3/2}}, x\right)$$

output `Unintegrable(1/(e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

### 3.568.2 Mathematica [N/A]

Not integrable

Time = 0.20 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(d+ex^2)(a+b\operatorname{arccosh}(cx))^{3/2}} dx$$

input `Integrate[1/((d + e*x^2)*(a + b*ArcCosh[c*x]))^(3/2),x]`

output `Integrate[1/((d + e*x^2)*(a + b*ArcCosh[c*x]))^(3/2), x]`

**3.568.3 Rubi [N/A]**

Not integrable

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^{3/2}} dx$$

input `Int[1/((d + e*x^2)*(a + b*ArcCosh[c*x])^(3/2)),x]`

output `$Aborted`

**3.568.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^n_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.568.4 Maple [N/A] (verified)**

Not integrable

Time = 1.41 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d)(a + b \operatorname{arccosh}(cx))^{\frac{3}{2}}} dx$$

input `int(1/(e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`

output `int(1/(e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x)`



**3.568.5 Fricas [F(-2)]**

Exception generated.

$$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.568.6 Sympy [N/A]**

Not integrable

Time = 36.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^{\frac{3}{2}}(d + ex^2)} dx$$

input `integrate(1/(e*x**2+d)/(a+b*acosh(c*x))**(3/2),x)`

output `Integral(1/((a + b*acosh(c*x))**(3/2)*(d + e*x**2)), x)`

**3.568.7 Maxima [N/A]**

Not integrable

Time = 0.51 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arccosh}(cx) + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate(1/((e*x^2 + d)*(b*arccosh(c*x) + a)^(3/2)), x)`

**3.568.8 Giac [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{1}{(ex^2 + d)(b \operatorname{arcosh}(cx) + a)^{3/2}} dx$$

input `integrate(1/(e*x^2+d)/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)*(b*arccosh(c*x) + a)^(3/2)), x)`**3.568.9 Mupad [N/A]**

Not integrable

Time = 3.99 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)(a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^{3/2} (ex^2 + d)} dx$$

input `int(1/((a + b*acosh(c*x))^(3/2)*(d + e*x^2)),x)`output `int(1/((a + b*acosh(c*x))^(3/2)*(d + e*x^2)), x)`

**3.569**  $\int \frac{1}{(d+ex^2)^2(a+b\operatorname{arccosh}(cx))^{3/2}} dx$

3.569.1 Optimal result . . . . .	4098
3.569.2 Mathematica [N/A] . . . . .	4098
3.569.3 Rubi [N/A] . . . . .	4099
3.569.4 Maple [N/A] (verified) . . . . .	4099
3.569.5 Fricas [F(-2)] . . . . .	4100
3.569.6 Sympy [F(-1)] . . . . .	4100
3.569.7 Maxima [N/A] . . . . .	4100
3.569.8 Giac [N/A] . . . . .	4101
3.569.9 Mupad [N/A] . . . . .	4101

**3.569.1 Optimal result**

Integrand size = 22, antiderivative size = 22

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^{3/2}} dx = \operatorname{Int}\left(\frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^{3/2}}, x\right)$$

output `Unintegrable(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

**3.569.2 Mathematica [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^{3/2}} dx$$

input `Integrate[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])^(3/2)),x]`

output `Integrate[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])^(3/2)), x]`

**3.569.3 Rubi [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {6325}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^{3/2}} dx$$

↓ 6325

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{arccosh}(cx))^{3/2}} dx$$

input `Int[1/((d + e*x^2)^2*(a + b*ArcCosh[c*x])^(3/2)),x]`

output `$Aborted`

**3.569.3.1 Defintions of rubi rules used**

rule 6325 `Int[((a_.) + ArcCosh[(c_.)*(x_)]*(b_.))^(n_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] :> Unintegrable[(d + e*x^2)^p*(a + b*ArcCosh[c*x])^n, x] /; FreeQ[{a, b, c, d, e, n, p}, x]`

**3.569.4 Maple [N/A] (verified)**

Not integrable

Time = 1.21 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(ex^2 + d)^2 (a + b \operatorname{arccosh}(cx))^{\frac{3}{2}}} dx$$

input `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

output `int(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x)`

**3.569.5 Fracas [F(-2)]**

Exception generated.

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Exception raised: TypeError}$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)`

**3.569.6 Sympy [F(-1)]**

Timed out.

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))^{3/2}} dx = \text{Timed out}$$

input `integrate(1/(e*x**2+d)**2/(a+b*acosh(c*x))**(3/2),x)`

output `Timed out`

**3.569.7 Maxima [N/A]**

Not integrable

Time = 0.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^2 (a + \operatorname{barccosh}(cx))^{3/2}} dx = \int \frac{1}{(ex^2 + d)^2 (b \operatorname{arccosh}(cx) + a)^{\frac{3}{2}}} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="maxima")`

output `integrate(1/((e*x^2 + d)^2*(b*arccosh(c*x) + a)^(3/2)), x)`

**3.569.8 Giac [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(ex^2 + d)^2 (b \operatorname{arccosh}(cx) + a)^{3/2}} dx$$

input `integrate(1/(e*x^2+d)^2/(a+b*arccosh(c*x))^(3/2),x, algorithm="giac")`output `integrate(1/((e*x^2 + d)^2*(b*arccosh(c*x) + a)^(3/2)), x)`**3.569.9 Mupad [N/A]**

Not integrable

Time = 3.78 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(d + ex^2)^2 (a + b \operatorname{arccosh}(cx))^{3/2}} dx = \int \frac{1}{(a + b \operatorname{acosh}(cx))^{3/2} (ex^2 + d)^2} dx$$

input `int(1/((a + b*acosh(c*x))^(3/2)*(d + e*x^2)^2),x)`output `int(1/((a + b*acosh(c*x))^(3/2)*(d + e*x^2)^2), x)`

## APPENDIX

4.1 Listing of Grading functions . . . . .	4102
--	------

## 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

### 4.1.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
```

```

(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A"," "}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A"," "}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.$
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
      ,
      finalresult={"F","Contains unresolved integral."}
    ]
  ];

  finalresult
]

```



```

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType,expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]],2]],
            Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3,ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
            If[Head[expn]===RootSum,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, CsCh,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

### 4.1.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
            print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
      return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of
                        convert(leaf_count_result,string)," vs. $2 (" ,
                        convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(" ,
                        convert(leaf_count_optimal,string)," )=" ,convert(2*leaf_cou
        fi;
      fi;
    fi;
  fi;

```

```

else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  end if
end if

```

```

elif type(expn, ``~`) then
  if type(op(2,expn), 'integer') then
    ExpnType(op(1,expn))
  elif type(op(2,expn), 'rational') then
    if type(op(1,expn), 'rational') then
      1
    else
      max(2, ExpnType(op(1,expn)))
    end if
  else
    max(3, ExpnType(op(1,expn)), ExpnType(op(2,expn)))
  end if
elif type(expn, ``+`) or type(expn, ``*`) then
  max(ExpnType(op(1,expn)), max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3, ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4, apply(max, map(ExpnType, [op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5, apply(max, map(ExpnType, [op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6, apply(max, map(ExpnType, [op(expn)])))
elif op(0,expn)='int' then
  max(8, apply(max, map(ExpnType, [op(expn)]))) else
  9
end if
end proc:

```

```

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

```

```

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,

```

```

        GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
        EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
    member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u),op(2..nops(u),u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

### 4.1.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):

```

```
if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
        return True
    else:
        return False
else:
    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
```



```

return 1
elif isinstance(expn,list):
    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
        return 1
    else:
        return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

*#main function*

```

def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

    #print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```

#### 4.1.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#      Albert Rich to use with Sagemath. This is used to
#      grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#      'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#      issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
```

```

        return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

```

```

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric', 'hypergeometric_M', 'hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=", expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception, AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):

```

```

    return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
elif is_sqrt(expn):
    if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
elif expn.operator() == operator.pow: #instance(expn,Pow)
    if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)
        return expnType(expn.operands()[0]) #expnType(expn.args[0])
    elif type(expn.operands()[1])==Rational: #instance(expn.args[1],Rational)
        if type(expn.operands()[0])==Rational: #instance(expn.args[0],Rational)
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #instance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

```

```

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)",type(result))
    print("type(optimal)",type(optimal))

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = " "
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal." + str(leaf_
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

```

```
return grade, grade_annotation
```