

Computer Algebra Independent Integration Tests

Summer 2024

360-table-of-integrals

Nasser M. Abbasi

May 17, 2024

Compiled on May 17, 2024 at 9:11pm

Contents

1	Introduction	7
1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26
2	detailed summary tables of results	27
2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	74
3	Listing of integrals	80
3.1	$\int x^n dx$	85
3.2	$\int \frac{1}{x} dx$	90
3.3	$\int e^x dx$	94
3.4	$\int a^x dx$	99
3.5	$\int \sin(x) dx$	104
3.6	$\int \cos(x) dx$	109
3.7	$\int \csc^2(x) dx$	114

3.8	$\int \sec^2(x) dx$	119
3.9	$\int \sec(x) \tan(x) dx$	124
3.10	$\int \cot(x) \csc(x) dx$	129
3.11	$\int \tan(x) dx$	134
3.12	$\int \cot(x) dx$	139
3.13	$\int \csc(x) dx$	144
3.14	$\int \sec(x) dx$	149
3.15	$\int \frac{1}{1+x^2} dx$	154
3.16	$\int \frac{1}{1-x^2} dx$	159
3.17	$\int \frac{1}{\sqrt{1-x^2}} dx$	164
3.18	$\int \frac{1}{\sqrt{1+x^2}} dx$	169
3.19	$\int \frac{1}{\sqrt{-1+x^2}} dx$	174
3.20	$\int \sinh(x) dx$	179
3.21	$\int \cosh(x) dx$	184
3.22	$\int \operatorname{csch}^2(x) dx$	189
3.23	$\int \operatorname{sech}^2(x) dx$	194
3.24	$\int \tanh(x) dx$	199
3.25	$\int \operatorname{coth}(x) dx$	204
3.26	$\int \operatorname{csch}(x) dx$	209
3.27	$\int (a + bx)^m dx$	214
3.28	$\int \frac{1}{a+bx} dx$	219
3.29	$\int \frac{x}{a+bx} dx$	224
3.30	$\int \frac{x^2}{a+bx} dx$	229
3.31	$\int \frac{1}{(a+bx)^2} dx$	234
3.32	$\int \frac{x}{(a+bx)^2} dx$	239
3.33	$\int \frac{x^2}{(a+bx)^2} dx$	244
3.34	$\int \frac{1}{(a+bx)^3} dx$	249
3.35	$\int \frac{x}{(a+bx)^3} dx$	254
3.36	$\int \frac{x^2}{(a+bx)^3} dx$	259
3.37	$\int \frac{x^3}{(a+bx)^3} dx$	264
3.38	$\int \frac{1}{(a+bx)^4} dx$	269
3.39	$\int \frac{x}{(a+bx)^4} dx$	274
3.40	$\int \frac{x^2}{(a+bx)^4} dx$	279
3.41	$\int \frac{x^3}{(a+bx)^4} dx$	284
3.42	$\int \frac{1}{(a+bx)^5} dx$	289
3.43	$\int \frac{x}{(a+bx)^5} dx$	294
3.44	$\int \frac{x^2}{(a+bx)^5} dx$	299

3.45	$\int \frac{x^3}{(a+bx)^5} dx$	304
3.46	$\int \frac{1}{x(a+bx)} dx$	309
3.47	$\int \frac{1}{x^2(a+bx)} dx$	314
3.48	$\int \frac{1}{x^3(a+bx)} dx$	319
3.49	$\int \frac{1}{x^2(a+bx)^2} dx$	324
3.50	$\int \frac{1}{x^3(a+bx)^2} dx$	329
3.51	$\int \frac{1}{x(a+bx)^3} dx$	334
3.52	$\int \frac{1}{x^2(a+bx)^3} dx$	339
3.53	$\int \frac{1}{x^3(a+bx)^3} dx$	344
3.54	$\int \frac{1}{x(a+bx)^4} dx$	349
3.55	$\int \frac{1}{x^2(a+bx)^4} dx$	354
3.56	$\int \frac{1}{x^3(a+bx)^4} dx$	359
3.57	$\int \frac{1}{x(a+bx)^5} dx$	365
3.58	$\int \frac{1}{x^2(a+bx)^5} dx$	371
3.59	$\int \frac{1}{x^3(a+bx)^5} dx$	377
3.60	$\int \frac{1}{a+bx^2} dx$	383
3.61	$\int x(a+bx^2)^{-m} dx$	388
3.62	$\int \frac{1}{a+bx^3} dx$	393
3.63	$\int \frac{x}{a+bx^3} dx$	401
3.64	$\int \frac{x^2}{a+bx^3} dx$	409
3.65	$\int \frac{x^3}{a+bx^3} dx$	414
3.66	$\int \frac{x^4}{a+bx^3} dx$	423
3.67	$\int \frac{1}{(a+bx^3)^2} dx$	432
3.68	$\int \frac{x}{(a+bx^3)^2} dx$	442
3.69	$\int \frac{x^2}{(a+bx^3)^2} dx$	451
3.70	$\int \frac{x^3}{(a+bx^3)^2} dx$	456
3.71	$\int \frac{1}{x(a+bx^3)} dx$	466
3.72	$\int \frac{1}{x^2(a+bx^3)} dx$	471
3.73	$\int \frac{1}{x^3(a+bx^3)} dx$	480
3.74	$\int \frac{1}{x(a+bx^3)^2} dx$	489
3.75	$\int \frac{1}{x^2(a+bx^3)^2} dx$	494
3.76	$\int \frac{1}{x^3(a+bx^3)^2} dx$	505
3.77	$\int \frac{1}{a+bx^4} dx$	517
3.78	$\int \frac{x}{a+bx^4} dx$	526
3.79	$\int \frac{x^2}{a+bx^4} dx$	531
3.80	$\int \frac{x^3}{a+bx^4} dx$	540

3.81	$\int \frac{1}{(a+bx^4)^2} dx$	545
3.82	$\int \frac{x}{(a+bx^4)^2} dx$	555
3.83	$\int \frac{x^2}{(a+bx^4)^2} dx$	561
3.84	$\int \frac{x^3}{(a+bx^4)^2} dx$	570
3.85	$\int \frac{1}{x(a+bx^4)} dx$	575
3.86	$\int \frac{1}{x^2(a+bx^4)} dx$	580
3.87	$\int \frac{1}{1+x} dx$	590
3.88	$\int \frac{1}{1+x^2} dx$	595
3.89	$\int \frac{1}{1+x^3} dx$	600
3.90	$\int \frac{1}{1+x^4} dx$	606
3.91	$\int \frac{1}{1-x} dx$	613
3.92	$\int \frac{1}{1-x^2} dx$	618
3.93	$\int \frac{1}{-1+x^2} dx$	623
3.94	$\int \frac{1}{1-x^3} dx$	628
3.95	$\int \frac{1}{1-x^4} dx$	634
3.96	$\int \frac{x}{1+x} dx$	639
3.97	$\int \frac{x}{1+x^2} dx$	644
3.98	$\int \frac{x}{1+x^3} dx$	649
3.99	$\int \frac{x}{1+x^4} dx$	655
3.100	$\int \frac{x}{1-x} dx$	660
3.101	$\int \frac{x}{1-x^2} dx$	665
3.102	$\int \frac{x}{1-x^3} dx$	670
3.103	$\int \frac{x}{1-x^4} dx$	676
3.104	$\int \frac{1}{x(1+x^2)} dx$	681
3.105	$\int \frac{1}{x(1-x^2)} dx$	686
3.106	$\int \frac{a+bx}{A+Bx} dx$	691
3.107	$\int \frac{1}{(a+bx)(A+Bx)} dx$	696
3.108	$\int \frac{x}{(a+bx)(A+Bx)} dx$	701
3.109	$\int \frac{1}{\sqrt{x}(a+bx)} dx$	706
3.110	$\int \frac{\sqrt{x}}{a+bx} dx$	711
3.111	$\int \frac{x^{3/2}}{a+bx} dx$	716
3.112	$\int \frac{x^{5/2}}{a+bx} dx$	722
3.113	$\int \frac{1}{\sqrt{x}(a+bx)^2} dx$	729
3.114	$\int \frac{\sqrt{x}}{(a+bx)^2} dx$	735
3.115	$\int \frac{x^{3/2}}{(a+bx)^2} dx$	741
3.116	$\int \frac{x^{5/2}}{(a+bx)^2} dx$	747

3.117	$\int \frac{1}{\sqrt{x}(a+bx)^3} dx$	754
3.118	$\int \frac{\sqrt{x}}{(a+bx)^3} dx$	760
3.119	$\int \frac{x^{3/2}}{(a+bx)^3} dx$	767
3.120	$\int \frac{x^{5/2}}{(a+bx)^3} dx$	773
3.121	$\int \frac{1}{\sqrt{x}(a+bx^2)} dx$	780
3.122	$\int \frac{\sqrt{x}}{a+bx^2} dx$	790
3.123	$\int \frac{x^{3/2}}{a+bx^2} dx$	800
3.124	$\int \frac{x^{5/2}}{a+bx^2} dx$	810
3.125	$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx$	820
3.126	$\int \frac{\sqrt{x}}{(a+bx^2)^2} dx$	831
3.127	$\int \frac{x^{3/2}}{(a+bx^2)^2} dx$	841
3.128	$\int \frac{x^{5/2}}{(a+bx^2)^2} dx$	850
3.129	$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx$	860
3.130	$\int \frac{\sqrt{x}}{(a+bx^2)^3} dx$	872
3.131	$\int \frac{x^{3/2}}{(a+bx^2)^3} dx$	884
3.132	$\int \frac{x^{5/2}}{(a+bx^2)^3} dx$	895
3.133	$\int \frac{1}{\sqrt{a+bx}} dx$	906
3.134	$\int \frac{x}{\sqrt{a+bx}} dx$	911
3.135	$\int \frac{x^2}{\sqrt{a+bx}} dx$	916
3.136	$\int \frac{1}{\sqrt{(a+bx)^3}} dx$	922
3.137	$\int \frac{x}{\sqrt{(a+bx)^3}} dx$	927
3.138	$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx$	932
3.139	$\int \frac{1}{x\sqrt{a+bx}} dx$	937
3.140	$\int \frac{\sqrt{a+bx}}{x} dx$	942
3.141	$\int \frac{\sqrt{a+bx}}{x^2} dx$	947
3.142	$\int \frac{\sqrt{a+bx}}{x^3} dx$	953
3.143	$\int \frac{\sqrt{(a+bx)^3}}{x} dx$	959
3.144	$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx$	965
3.145	$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx$	971
3.146	$\int \frac{1}{x^2\sqrt{a+bx}} dx$	977
3.147	$\int \frac{1}{x^3\sqrt{a+bx}} dx$	983
3.148	$\int \frac{1}{x\sqrt{(a+bx)^3}} dx$	989
3.149	$\int \frac{1}{x^2\sqrt{(a+bx)^3}} dx$	995

3.150	$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx$	1002
3.151	$\int \frac{1}{x^3 \sqrt{(a+bx)^2}} dx$	1010
3.152	$\int \frac{\sqrt[3]{a+bx}}{x} dx$	1017
3.153	$\int \frac{\sqrt[3]{a+bx}}{x^2} dx$	1025
3.154	$\int \frac{\sqrt[3]{a+bx}}{x^3} dx$	1033
3.155	$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx$	1043
3.156	$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx$	1051
3.157	$\int \frac{1}{x \sqrt[3]{a+bx}} dx$	1060
3.158	$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx$	1068
3.159	$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx$	1075
3.160	$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx$	1083
3.161	$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx$	1088
3.162	$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx$	1097
3.163	$\int \frac{A+Bx}{\sqrt{a+bx}} dx$	1108
4	Appendix	1113
4.1	Listing of Grading functions	1113
4.2	Links to plain text integration problems used in this report for each CAS	131

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	8
1.2	Results	9
1.3	Time and leaf size Performance	13
1.4	Performance based on number of rules Rubi used	15
1.5	Performance based on number of steps Rubi used	16
1.6	Solved integrals histogram based on leaf size of result	17
1.7	Solved integrals histogram based on CPU time used	18
1.8	Leaf size vs. CPU time used	19
1.9	list of integrals with no known antiderivative	20
1.10	List of integrals solved by CAS but has no known antiderivative	20
1.11	list of integrals solved by CAS but failed verification	20
1.12	Timing	21
1.13	Verification	21
1.14	Important notes about some of the results	22
1.15	Current tree layout of integration tests	25
1.16	Design of the test system	26

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [**163**]. This is test number [360].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (163)	0.00 (0)
Mathematica	100.00 (163)	0.00 (0)
Fricas	100.00 (163)	0.00 (0)
Giac	100.00 (163)	0.00 (0)
Maple	97.55 (159)	2.45 (4)
Reduce	96.32 (157)	3.68 (6)
Mupad	92.64 (151)	7.36 (12)
Maxima	92.64 (151)	7.36 (12)
Sympy	88.34 (144)	11.66 (19)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	93.865	6.135	0.000	0.000
Mathematica	92.638	7.362	0.000	0.000
Maple	87.117	0.000	10.429	2.454
Giac	82.209	17.791	0.000	0.000
Maxima	81.595	11.043	0.000	7.362
Fricas	64.417	25.153	10.429	0.000
Sympy	58.282	26.380	3.681	11.656
Mupad	0.000	92.638	0.000	7.362
Reduce	0.000	96.319	0.000	3.681

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Giac	0	0.00	0.00	0.00
Maple	4	100.00	0.00	0.00
Reduce	6	100.00	0.00	0.00
Mupad	12	0.00	100.00	0.00
Maxima	12	91.67	0.00	8.33
Sympy	19	84.21	15.79	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Mathematica	0.05
Maple	0.07
Maxima	0.07
Fricas	0.10
Mupad	0.11
Giac	0.21
Rubi	0.24
Reduce	0.25
Sympy	2.34

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maple	39.22	0.94	30.00	0.95
Mupad	42.14	1.01	33.00	1.00
Mathematica	53.17	1.28	38.00	1.00
Maxima	58.28	1.27	36.00	1.00
Giac	63.32	1.52	37.00	1.10
Rubi	65.85	1.19	42.00	1.08
Reduce	79.90	1.72	45.00	1.22
Fricas	100.66	2.09	62.00	1.62
Sympy	128.05	2.25	38.00	1.06

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```


For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	28
2.2	Detailed conclusion table per each integral for all CAS systems	33
2.3	Detailed conclusion table specific for Rubi results	74

2.1 List of integrals sorted by grade for each CAS

Rubi	28
Mma	29
Maple	29
Fricas	30
Maxima	30
Giac	31
Mupad	31
Sympy	32
Reduce	32

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 80, 82, 84, 85, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163 }

B grade { 77, 79, 81, 83, 86, 121, 122, 123, 124, 137 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 80, 82, 84, 85, 87, 88, 89, 90, 91, 94, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163 }

B grade { 16, 17, 18, 19, 77, 79, 81, 83, 86, 92, 93, 95 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 64, 69, 71, 74, 78, 80, 82, 84, 85, 87, 88, 89, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 152, 153, 154, 157, 159, 160, 161, 162, 163 }

B grade { }

C grade { 62, 63, 65, 66, 67, 68, 70, 72, 73, 75, 76, 77, 79, 81, 83, 86, 90 }

F normal fail { 151, 155, 156, 158 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 9, 10, 15, 19, 20, 21, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 40, 41, 45, 46, 47, 48, 49, 50, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 80, 82, 84, 85, 87, 88, 89, 90, 91, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 133, 134, 135, 139, 140, 141, 142, 144, 145, 146, 147, 152, 153, 154, 157, 160, 161, 162, 163 }

B grade { 8, 11, 12, 13, 14, 16, 17, 18, 22, 23, 24, 25, 26, 38, 39, 42, 43, 44, 51, 52, 53, 54, 55, 56, 57, 58, 59, 92, 93, 136, 137, 138, 143, 148, 149, 150, 151, 155, 156, 158, 159 }

C grade { 77, 79, 81, 83, 86, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20, 21, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 42, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 80, 82, 84, 85, 87, 88, 89, 90, 91, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 126, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 146, 147, 152, 153, 154, 157, 160, 161, 162, 163 }

B grade { 16, 22, 23, 39, 43, 44, 77, 79, 81, 83, 86, 92, 93, 121, 122, 123, 124, 127 }

C grade { }

F normal fail { 143, 144, 145, 148, 149, 150, 151, 155, 156, 158, 159 }

F(-1) timedout fail { }

F(-2) exception fail { 61 }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 78, 80, 82, 84, 85, 87, 88, 89, 90, 91, 94, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 123, 124, 125, 126, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 152, 153, 154, 157, 160, 161, 162, 163 }
}

B grade { 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 77, 79, 81, 83, 86, 92, 93, 95, 121, 122, 127, 128, 151, 155, 156, 158, 159 }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 146, 147, 152, 153, 154, 157, 161, 162, 163 }
}

C grade { }

F normal fail { }

F(-1) timedout fail { 143, 144, 145, 148, 149, 150, 151, 155, 156, 158, 159, 160 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 3, 4, 5, 6, 9, 10, 11, 12, 15, 17, 18, 19, 20, 21, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 37, 41, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 79, 80, 81, 83, 84, 85, 86, 87, 88, 89, 90, 91, 94, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 121, 122, 123, 124, 133, 139, 140, 141, 142, 146, 147, 163 }

B grade { 7, 8, 13, 14, 16, 24, 25, 34, 38, 39, 40, 42, 43, 44, 60, 61, 78, 82, 92, 93, 95, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 125, 126, 127, 128, 130, 134, 135, 136 }

C grade { 152, 153, 154, 157, 161, 162 }

F normal fail { 22, 23, 137, 138, 143, 144, 145, 148, 149, 150, 151, 155, 156, 158, 159, 160 }

F(-1) timedout fail { 129, 131, 132 }

F(-2) exception fail { }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 152, 153, 154, 157, 160, 161, 162, 163 }

C grade { }

F normal fail { 61, 151, 155, 156, 158, 159 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	11	11	10	12	11	10	20
N.S.	1	1.00	1.00	1.00	1.00	0.91	1.09	1.00	0.91	1.82
time (sec)	N/A	0.132	0.001	0.060	0.031	0.120	0.016	0.107	0.255	0.256

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	3	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.50	1.00	1.00
time (sec)	N/A	0.123	0.000	0.014	0.037	0.071	0.036	0.111	0.245	0.005

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	3	2	2	2	2	3	2
N.S.	1	1.00	1.00	1.00	0.67	0.67	0.67	0.67	1.00	0.67
time (sec)	N/A	0.135	0.000	0.019	0.036	0.074	0.027	0.115	0.233	0.004

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	8	8	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.148	0.001	0.023	0.032	0.088	0.036	0.107	0.250	0.182

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	4	4	3	4	4	4
N.S.	1	1.00	1.00	1.25	1.00	1.00	0.75	1.00	1.00	1.00
time (sec)	N/A	0.152	0.001	0.102	0.027	0.087	0.038	0.106	0.227	0.012

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	2	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.162	0.001	0.044	0.027	0.090	0.035	0.108	0.236	0.179

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	6	8	7	6	8	4
N.S.	1	1.00	1.00	1.25	1.50	2.00	1.75	1.50	2.00	1.00
time (sec)	N/A	0.170	0.007	0.066	0.024	0.080	0.041	0.119	0.267	0.008

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	7	5	2	7	2
N.S.	1	1.00	1.00	1.50	1.00	3.50	2.50	1.00	3.50	1.00
time (sec)	N/A	0.172	0.003	0.056	0.033	0.087	0.041	0.117	0.243	0.014

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	4	4	3	4	2	12
N.S.	1	1.00	1.00	1.50	2.00	2.00	1.50	2.00	1.00	6.00
time (sec)	N/A	0.158	0.004	0.045	0.037	0.084	0.026	0.112	0.247	0.219

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	6	6	5	6	4	6
N.S.	1	1.00	1.00	1.25	1.50	1.50	1.25	1.50	1.00	1.50
time (sec)	N/A	0.166	0.001	0.034	0.032	0.088	0.054	0.110	0.227	0.194

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	5	5	5	6	3	11	5	6	9	5
N.S.	1	1.00	1.00	1.20	0.60	2.20	1.00	1.20	1.80	1.00
time (sec)	N/A	0.156	0.000	0.017	0.030	0.096	0.051	0.115	0.258	0.016

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	11	3	4	17	3
N.S.	1	1.00	1.00	1.33	1.00	3.67	1.00	1.33	5.67	1.00
time (sec)	N/A	0.154	0.001	0.039	0.035	0.089	0.043	0.104	0.230	0.012

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	5	5	6	8	19	15	6	5	5
N.S.	1	0.71	0.71	0.86	1.14	2.71	2.14	0.86	0.71	0.71
time (sec)	N/A	0.158	0.002	0.021	0.023	0.092	0.076	0.117	0.270	0.017

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	6	3	3	7	6	17	15	25	17	11
N.S.	1	0.50	0.50	1.17	1.00	2.83	2.50	4.17	2.83	1.83
time (sec)	N/A	0.155	0.000	0.056	0.032	0.099	0.059	0.112	0.280	0.001

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	2	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.121	0.003	0.030	0.108	0.072	0.040	0.097	0.276	0.024

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	19	3	13	13	12	15	13	2
N.S.	1	1.00	9.50	1.50	6.50	6.50	6.00	7.50	6.50	1.00
time (sec)	N/A	0.123	0.002	0.030	0.024	0.086	0.046	0.115	0.232	0.001

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	20	3	2	18	2	17	2	2
N.S.	1	1.00	10.00	1.50	1.00	9.00	1.00	8.50	1.00	1.00
time (sec)	N/A	0.166	0.001	0.218	0.110	0.093	0.069	0.116	0.244	0.004

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	16	3	2	14	2	25	9	2
N.S.	1	1.00	8.00	1.50	1.00	7.00	1.00	12.50	4.50	1.00
time (sec)	N/A	0.154	0.001	0.066	0.125	0.088	0.068	0.113	0.227	0.002

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	38	11	14	14	10	26	9	10
N.S.	1	1.00	3.17	0.92	1.17	1.17	0.83	2.17	0.75	0.83
time (sec)	N/A	0.174	0.002	0.074	0.032	0.087	0.065	0.121	0.268	0.098

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	11	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	5.50	1.00	1.00
time (sec)	N/A	0.236	0.001	0.039	0.032	0.094	0.073	0.110	0.251	0.010

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	11	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	5.50	1.00	1.00
time (sec)	N/A	0.234	0.001	0.033	0.024	0.077	0.073	0.110	0.236	0.009

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	10	20	0	10	16	4
N.S.	1	1.00	1.00	1.25	2.50	5.00	0.00	2.50	4.00	1.00
time (sec)	N/A	0.287	0.001	0.060	0.024	0.068	0.000	0.108	0.261	0.005

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	10	20	0	10	16	2
N.S.	1	1.00	1.00	1.50	5.00	10.00	0.00	5.00	8.00	1.00
time (sec)	N/A	0.287	0.000	0.067	0.040	0.070	0.000	0.109	0.236	0.001

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	18	7	11	12	3
N.S.	1	1.00	1.00	1.33	1.00	6.00	2.33	3.67	4.00	1.00
time (sec)	N/A	0.206	0.000	0.023	0.023	0.079	0.062	0.121	0.245	0.001

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	3	4	3	18	12	12	16	3
N.S.	1	1.00	1.00	1.33	1.00	6.00	4.00	4.00	5.33	1.00
time (sec)	N/A	0.246	0.001	0.055	0.034	0.094	0.159	0.109	0.241	0.016

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	7	5	5	6	5	17	5	14	15	5
N.S.	1	0.71	0.71	0.86	0.71	2.43	0.71	2.00	2.14	0.71
time (sec)	N/A	0.213	0.000	0.015	0.045	0.086	0.141	0.112	0.275	0.007

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	19	18	20	20	18	21	18
N.S.	1	1.00	1.00	1.06	1.00	1.11	1.11	1.00	1.17	1.00
time (sec)	N/A	0.202	0.004	0.042	0.030	0.089	0.022	0.108	0.249	0.221

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	10	10	7	11	10	10
N.S.	1	1.00	1.00	1.10	1.00	1.00	0.70	1.10	1.00	1.00
time (sec)	N/A	0.215	0.001	0.029	0.031	0.078	0.021	0.124	0.234	0.016

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	19	18	17	14	19	17	18
N.S.	1	1.00	1.00	1.06	1.00	0.94	0.78	1.06	0.94	1.00
time (sec)	N/A	0.263	0.003	0.029	0.023	0.103	0.050	0.112	0.259	0.022

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	30	29	29	26	30	29	29
N.S.	1	1.00	1.00	0.97	0.94	0.94	0.84	0.97	0.94	0.94
time (sec)	N/A	0.226	0.003	0.033	0.032	0.078	0.051	0.107	0.242	0.024

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	12	13	10	12	12	12
N.S.	1	1.00	1.00	1.08	1.00	1.08	0.83	1.00	1.00	1.00
time (sec)	N/A	0.136	0.002	0.029	0.037	0.077	0.068	0.108	0.232	0.016

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	23	20	24	26	28	20	42	33	23
N.S.	1	0.96	0.83	1.00	1.08	1.17	0.83	1.75	1.38	0.96
time (sec)	N/A	0.230	0.006	0.033	0.029	0.082	0.061	0.113	0.244	0.020

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	29	34	36	47	31	50	46	36
N.S.	1	1.00	0.88	1.03	1.09	1.42	0.94	1.52	1.39	1.09
time (sec)	N/A	0.182	0.010	0.035	0.031	0.089	0.083	0.114	0.266	0.173

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	24	26	12	23	26
N.S.	1	1.00	1.00	0.93	0.86	1.71	1.86	0.86	1.64	1.86
time (sec)	N/A	0.138	0.002	0.034	0.026	0.092	0.114	0.110	0.238	0.158

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	17	20	19	32	32	32	18	26	32
N.S.	1	0.74	0.87	0.83	1.39	1.39	1.39	0.78	1.13	1.39
time (sec)	N/A	0.138	0.005	0.036	0.028	0.071	0.090	0.108	0.218	0.018

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	41	33	36	48	61	46	37	71	46
N.S.	1	1.11	0.89	0.97	1.30	1.65	1.24	1.00	1.92	1.24
time (sec)	N/A	0.186	0.010	0.040	0.025	0.082	0.104	0.107	0.239	0.164

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	50	40	45	57	83	58	44	85	43
N.S.	1	0.88	0.70	0.79	1.00	1.46	1.02	0.77	1.49	0.75
time (sec)	N/A	0.199	0.032	0.038	0.036	0.085	0.139	0.114	0.250	0.224

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	35	37	12	34	37
N.S.	1	1.00	1.00	0.93	0.86	2.50	2.64	0.86	2.43	2.64
time (sec)	N/A	0.138	0.002	0.037	0.028	0.079	0.132	0.114	0.265	0.022

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	30	20	19	43	43	44	18	42	44
N.S.	1	1.20	0.80	0.76	1.72	1.72	1.76	0.72	1.68	1.76
time (sec)	N/A	0.166	0.005	0.037	0.027	0.083	0.118	0.110	0.262	0.163

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	17	31	30	54	54	56	29	37	56
N.S.	1	0.50	0.91	0.88	1.59	1.59	1.65	0.85	1.09	1.65
time (sec)	N/A	0.141	0.008	0.039	0.035	0.082	0.125	0.111	0.268	0.165

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	58	44	47	70	94	70	46	108	45
N.S.	1	1.16	0.88	0.94	1.40	1.88	1.40	0.92	2.16	0.90
time (sec)	N/A	0.199	0.013	0.039	0.036	0.099	0.150	0.108	0.252	0.054

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	46	49	12	45	48
N.S.	1	1.00	1.00	0.93	0.86	3.29	3.50	0.86	3.21	3.43
time (sec)	N/A	0.134	0.002	0.042	0.036	0.100	0.166	0.109	0.246	0.188

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	30	20	19	54	54	56	31	53	18
N.S.	1	1.20	0.80	0.76	2.16	2.16	2.24	1.24	2.12	0.72
time (sec)	N/A	0.165	0.005	0.044	0.027	0.080	0.169	0.115	0.223	0.033

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	47	31	30	65	65	68	46	64	22
N.S.	1	1.24	0.82	0.79	1.71	1.71	1.79	1.21	1.68	0.58
time (sec)	N/A	0.182	0.007	0.040	0.031	0.094	0.192	0.110	0.263	0.195

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	17	42	41	76	76	80	61	48	48
N.S.	1	0.36	0.89	0.87	1.62	1.62	1.70	1.30	1.02	1.02
time (sec)	N/A	0.135	0.007	0.046	0.035	0.084	0.212	0.120	0.260	0.037

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	18	18	16	18	16	10	20	15	15
N.S.	1	1.20	1.20	1.07	1.20	1.07	0.67	1.33	1.00	1.00
time (sec)	N/A	0.141	0.004	0.037	0.023	0.099	0.090	0.107	0.253	0.030

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	28	28	26	28	26	19	30	26	25
N.S.	1	1.17	1.17	1.08	1.17	1.08	0.79	1.25	1.08	1.04
time (sec)	N/A	0.170	0.004	0.041	0.036	0.093	0.092	0.113	0.256	0.173

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	42	42	41	40	41	31	45	43	38
N.S.	1	1.14	1.14	1.11	1.08	1.11	0.84	1.22	1.16	1.03
time (sec)	N/A	0.183	0.004	0.047	0.024	0.111	0.113	0.110	0.268	0.040

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	42	35	43	45	63	37	52	70	45
N.S.	1	1.05	0.88	1.08	1.12	1.58	0.92	1.30	1.75	1.12
time (sec)	N/A	0.186	0.030	0.047	0.032	0.093	0.176	0.112	0.248	0.201

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	58	53	57	64	86	54	74	86	57
N.S.	1	1.02	0.93	1.00	1.12	1.51	0.95	1.30	1.51	1.00
time (sec)	N/A	0.208	0.037	0.043	0.025	0.114	0.159	0.115	0.242	0.040

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	43	37	41	51	80	46	43	97	43
N.S.	1	1.13	0.97	1.08	1.34	2.11	1.21	1.13	2.55	1.13
time (sec)	N/A	0.186	0.021	0.043	0.034	0.100	0.150	0.108	0.290	0.243

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	57	53	56	69	109	66	60	119	63
N.S.	1	1.12	1.04	1.10	1.35	2.14	1.29	1.18	2.33	1.24
time (sec)	N/A	0.202	0.034	0.047	0.031	0.096	0.173	0.109	0.243	0.180

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	76	68	72	86	130	78	73	138	79
N.S.	1	1.19	1.06	1.12	1.34	2.03	1.22	1.14	2.16	1.23
time (sec)	N/A	0.221	0.031	0.048	0.025	0.209	0.194	0.111	0.250	0.196

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	57	48	52	73	124	70	54	145	60
N.S.	1	1.12	0.94	1.02	1.43	2.43	1.37	1.06	2.84	1.18
time (sec)	N/A	0.197	0.021	0.054	0.034	0.150	0.247	0.111	0.229	0.073

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	70	64	69	91	153	90	71	171	85
N.S.	1	1.13	1.03	1.11	1.47	2.47	1.45	1.15	2.76	1.37
time (sec)	N/A	0.219	0.039	0.053	0.041	0.097	0.245	0.114	0.248	0.049

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	93	79	83	108	174	104	86	190	101
N.S.	1	1.18	1.00	1.05	1.37	2.20	1.32	1.09	2.41	1.28
time (sec)	N/A	0.240	0.039	0.055	0.042	0.090	0.239	0.106	0.248	0.054

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	71	59	63	95	168	94	89	197	77
N.S.	1	1.11	0.92	0.98	1.48	2.62	1.47	1.39	3.08	1.20
time (sec)	N/A	0.212	0.028	0.049	0.036	0.091	0.229	0.110	0.251	0.215

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	87	75	82	113	197	114	108	223	107
N.S.	1	1.13	0.97	1.06	1.47	2.56	1.48	1.40	2.90	1.39
time (sec)	N/A	0.235	0.044	0.053	0.033	0.086	0.271	0.112	0.253	0.193

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	105	90	94	130	218	128	128	242	123
N.S.	1	1.18	1.01	1.06	1.46	2.45	1.44	1.44	2.72	1.38
time (sec)	N/A	0.259	0.044	0.056	0.040	0.103	0.403	0.116	0.258	0.063

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	24	24	16	15	67	53	15	23	16
N.S.	1	1.20	1.20	0.80	0.75	3.35	2.65	0.75	1.15	0.80
time (sec)	N/A	0.147	0.005	0.037	0.125	0.099	0.071	0.112	0.232	0.032

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	27	24	26	0	29	102	23	15	23
N.S.	1	1.08	0.96	1.04	0.00	1.16	4.08	0.92	0.60	0.92
time (sec)	N/A	0.152	0.004	0.053	0.000	0.103	1.301	0.111	0.240	0.228

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	111	89	27	98	299	20	112	67	99
N.S.	1	1.18	0.95	0.29	1.04	3.18	0.21	1.19	0.71	1.05
time (sec)	N/A	0.278	0.024	0.029	0.123	0.101	0.074	0.112	0.249	0.139

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	116	89	27	98	304	24	112	65	111
N.S.	1	1.15	0.88	0.27	0.97	3.01	0.24	1.11	0.64	1.10
time (sec)	N/A	0.263	0.008	0.031	0.115	0.110	0.058	0.113	0.258	0.291

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	10	14	37	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.67	0.93	2.47	0.87
time (sec)	N/A	0.147	0.002	0.027	0.026	0.094	0.072	0.115	0.279	0.020

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	123	108	34	106	106	22	111	78	114
N.S.	1	1.23	1.08	0.34	1.06	1.06	0.22	1.11	0.78	1.14
time (sec)	N/A	0.285	0.013	0.025	0.109	0.095	0.081	0.120	0.286	0.144

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	133	111	37	109	123	32	114	81	120
N.S.	1	1.18	0.98	0.33	0.96	1.09	0.28	1.01	0.72	1.06
time (sec)	N/A	0.282	0.013	0.030	0.126	0.105	0.091	0.120	0.253	0.268

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	136	118	46	122	389	39	127	171	128
N.S.	1	1.21	1.05	0.41	1.09	3.47	0.35	1.13	1.53	1.14
time (sec)	N/A	0.283	0.048	0.030	0.107	0.101	0.135	0.114	0.249	0.247

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	143	119	48	124	386	44	129	158	138
N.S.	1	1.15	0.96	0.39	1.00	3.11	0.35	1.04	1.27	1.11
time (sec)	N/A	0.289	0.042	0.030	0.107	0.111	0.148	0.115	0.237	0.319

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	15	15	14	17	14
N.S.	1	1.00	1.00	0.94	0.88	0.94	0.94	0.88	1.06	0.88
time (sec)	N/A	0.145	0.003	0.031	0.023	0.081	0.100	0.112	0.292	0.155

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	136	118	43	114	391	39	130	171	108
N.S.	1	1.18	1.03	0.37	0.99	3.40	0.34	1.13	1.49	0.94
time (sec)	N/A	0.297	0.042	0.033	0.111	0.118	0.132	0.117	0.287	0.135

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	26	22	21	23	18	15	22	45	18
N.S.	1	1.24	1.05	1.00	1.10	0.86	0.71	1.05	2.14	0.86
time (sec)	N/A	0.157	0.005	0.030	0.024	0.094	0.136	0.115	0.270	0.191

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	131	114	53	106	103	29	121	84	102
N.S.	1	1.19	1.04	0.48	0.96	0.94	0.26	1.10	0.76	0.93
time (sec)	N/A	0.290	0.014	0.038	0.117	0.098	0.089	0.115	0.250	0.280

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	128	119	54	106	143	32	115	96	128
N.S.	1	1.21	1.12	0.51	1.00	1.35	0.30	1.08	0.91	1.21
time (sec)	N/A	0.281	0.017	0.036	0.120	0.085	0.122	0.118	0.240	0.271

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	39	33	35	37	47	34	45	114	34
N.S.	1	1.03	0.87	0.92	0.97	1.24	0.89	1.18	3.00	0.89
time (sec)	N/A	0.188	0.010	0.042	0.030	0.088	0.239	0.111	0.234	0.037

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	158	131	73	126	146	56	139	183	120
N.S.	1	1.20	0.99	0.55	0.95	1.11	0.42	1.05	1.39	0.91
time (sec)	N/A	0.329	0.060	0.044	0.117	0.104	0.175	0.118	0.271	0.282

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	155	129	74	128	180	58	131	202	146
N.S.	1	1.23	1.02	0.59	1.02	1.43	0.46	1.04	1.60	1.16
time (sec)	N/A	0.313	0.063	0.046	0.124	0.097	0.202	0.119	0.261	0.391

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	C	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	200	134	27	169	112	20	179	112	33
N.S.	1	3.17	2.13	0.43	2.68	1.78	0.32	2.84	1.78	0.52
time (sec)	N/A	0.673	0.034	0.026	0.113	0.080	0.082	0.116	0.244	0.051

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	29	29	19	18	72	56	18	67	19
N.S.	1	1.16	1.16	0.76	0.72	2.88	2.24	0.72	2.68	0.76
time (sec)	N/A	0.262	0.006	0.032	0.109	0.113	0.099	0.113	0.235	0.170

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	C	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	200	134	27	169	124	26	179	112	35
N.S.	1	3.17	2.13	0.43	2.68	1.97	0.41	2.84	1.78	0.56
time (sec)	N/A	0.615	0.013	0.026	0.113	0.119	0.076	0.121	0.265	0.054

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	10	14	47	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.67	0.93	3.13	0.87
time (sec)	N/A	0.215	0.003	0.026	0.033	0.072	0.078	0.109	0.228	0.159

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	C	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	225	183	46	189	183	39	194	305	58
N.S.	1	2.78	2.26	0.57	2.33	2.26	0.48	2.40	3.77	0.72
time (sec)	N/A	0.717	0.075	0.033	0.108	0.123	0.161	0.118	0.238	0.200

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	53	49	40	39	129	83	39	166	37
N.S.	1	1.10	1.02	0.83	0.81	2.69	1.73	0.81	3.46	0.77
time (sec)	N/A	0.282	0.022	0.041	0.108	0.118	0.170	0.115	0.292	0.027

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	C	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	227	184	48	191	196	46	196	305	60
N.S.	1	2.64	2.14	0.56	2.22	2.28	0.53	2.28	3.55	0.70
time (sec)	N/A	0.691	0.082	0.032	0.125	0.089	0.145	0.115	0.256	0.063

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	15	15	14	17	14
N.S.	1	1.00	1.00	0.94	0.88	0.94	0.94	0.88	1.06	0.88
time (sec)	N/A	0.174	0.004	0.031	0.022	0.079	0.153	0.113	0.259	0.162

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	26	22	21	23	18	15	24	55	18
N.S.	1	1.24	1.05	1.00	1.10	0.86	0.71	1.14	2.62	0.86
time (sec)	N/A	0.170	0.005	0.033	0.026	0.080	0.160	0.109	0.237	0.202

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	B	C	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	215	179	50	183	128	29	187	150	51
N.S.	1	2.99	2.49	0.69	2.54	1.78	0.40	2.60	2.08	0.71
time (sec)	N/A	0.517	0.021	0.040	0.121	0.091	0.105	0.114	0.254	0.202

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	4	5	4	4	3	5	4	4
N.S.	1	1.00	1.00	1.25	1.00	1.00	0.75	1.25	1.00	1.00
time (sec)	N/A	0.148	0.000	0.019	0.022	0.066	0.019	0.122	0.246	0.011

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	2	2	2	2	2	2
N.S.	1	1.00	1.00	1.50	1.00	1.00	1.00	1.00	1.00	1.00
time (sec)	N/A	0.139	0.002	0.000	0.118	0.096	0.059	0.119	0.256	0.001

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	45	40	33	34	34	41	35	33	31
N.S.	1	1.05	0.93	0.77	0.79	0.79	0.95	0.81	0.77	0.72
time (sec)	N/A	0.226	0.008	0.037	0.111	0.083	0.070	0.110	0.269	0.051

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	90	64	22	72	60	73	72	56	33
N.S.	1	1.38	0.98	0.34	1.11	0.92	1.12	1.11	0.86	0.51
time (sec)	N/A	0.309	0.014	0.036	0.120	0.117	0.069	0.115	0.252	0.040

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	5	7	6	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.62	0.88	0.75	0.75
time (sec)	N/A	0.148	0.001	0.023	0.035	0.097	0.019	0.111	0.235	0.014

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2	2	19	3	13	13	12	15	13	2
N.S.	1	1.00	9.50	1.50	6.50	6.50	6.00	7.50	6.50	1.00
time (sec)	N/A	0.151	0.002	0.000	0.037	0.073	0.045	0.113	0.235	0.002

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	4	4	19	5	13	13	12	15	13	4
N.S.	1	1.00	4.75	1.25	3.25	3.25	3.00	3.75	3.25	1.00
time (sec)	N/A	0.153	0.002	0.028	0.029	0.080	0.042	0.111	0.266	0.039

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	45	40	31	32	32	41	33	31	46
N.S.	1	1.05	0.93	0.72	0.74	0.74	0.95	0.77	0.72	1.07
time (sec)	N/A	0.229	0.007	0.040	0.115	0.091	0.070	0.111	0.252	0.190

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	A	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	13	25	10	17	17	17	19	17	9
N.S.	1	1.44	2.78	1.11	1.89	1.89	1.89	2.11	1.89	1.00
time (sec)	N/A	0.174	0.004	0.042	0.112	0.087	0.078	0.107	0.256	0.019

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	9	8	8	5	9	8	8
N.S.	1	1.00	1.00	1.12	1.00	1.00	0.62	1.12	1.00	1.00
time (sec)	N/A	0.163	0.001	0.022	0.039	0.078	0.036	0.111	0.261	0.014

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	9	8	8	7	8	8	8
N.S.	1	1.00	1.00	0.90	0.80	0.80	0.70	0.80	0.80	0.80
time (sec)	N/A	0.161	0.001	0.023	0.022	0.078	0.036	0.112	0.241	0.017

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	45	40	35	34	34	41	35	33	46
N.S.	1	1.12	1.00	0.88	0.85	0.85	1.02	0.88	0.82	1.15
time (sec)	N/A	0.233	0.005	0.036	0.114	0.097	0.077	0.117	0.251	0.189

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	8	8	8	7	6	6	5	6	29	6
N.S.	1	1.00	1.00	0.88	0.75	0.75	0.62	0.75	3.62	0.75
time (sec)	N/A	0.166	0.003	0.027	0.110	0.112	0.048	0.115	0.268	0.032

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	7	11	10	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.58	0.92	0.83	0.83
time (sec)	N/A	0.177	0.002	0.026	0.024	0.073	0.031	0.103	0.268	0.016

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	9	8	8	8	9	13	8
N.S.	1	1.00	1.00	0.75	0.67	0.67	0.67	0.75	1.08	0.67
time (sec)	N/A	0.154	0.001	0.029	0.030	0.084	0.030	0.110	0.267	0.022

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	46	41	33	32	32	41	33	31	46
N.S.	1	1.12	1.00	0.80	0.78	0.78	1.00	0.80	0.76	1.12
time (sec)	N/A	0.228	0.005	0.035	0.112	0.081	0.064	0.118	0.255	0.188

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	8	23	7	17	17	15	18	21	6
N.S.	1	0.40	1.15	0.35	0.85	0.85	0.75	0.90	1.05	0.30
time (sec)	N/A	0.159	0.003	0.036	0.028	0.089	0.043	0.111	0.258	0.039

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	17	13	12	15	11	10	15	11	11
N.S.	1	1.42	1.08	1.00	1.25	0.92	0.83	1.25	0.92	0.92
time (sec)	N/A	0.164	0.002	0.028	0.035	0.082	0.046	0.111	0.264	0.161

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	19	15	12	15	11	10	16	15	11
N.S.	1	1.36	1.07	0.86	1.07	0.79	0.71	1.14	1.07	0.79
time (sec)	N/A	0.167	0.003	0.037	0.026	0.076	0.050	0.113	0.228	0.177

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	26	25	26	25	24	20	26	1	26
N.S.	1	1.04	1.00	1.04	1.00	0.96	0.80	1.04	0.04	1.04
time (sec)	N/A	0.187	0.007	0.046	0.024	0.087	0.082	0.109	0.238	0.034

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	36	26	27	36	26	128	46	12	25
N.S.	1	1.44	1.04	1.08	1.44	1.04	5.12	1.84	0.48	1.00
time (sec)	N/A	0.166	0.009	0.056	0.030	0.088	0.189	0.112	0.268	0.210

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	44	38	38	44	38	138	46	33	37
N.S.	1	1.26	1.09	1.09	1.26	1.09	3.94	1.31	0.94	1.06
time (sec)	N/A	0.194	0.012	0.073	0.024	0.127	0.406	0.109	0.240	0.209

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	29	29	19	18	68	73	18	25	19
N.S.	1	1.81	1.81	1.19	1.12	4.25	4.56	1.12	1.56	1.19
time (sec)	N/A	0.147	0.018	0.069	0.110	0.114	0.435	0.110	0.233	0.182

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	40	40	32	31	85	88	31	29	28
N.S.	1	1.29	1.29	1.03	1.00	2.74	2.84	1.00	0.94	0.90
time (sec)	N/A	0.157	0.025	0.066	0.118	0.098	0.298	0.114	0.252	0.178

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	59	49	42	42	103	107	45	39	37
N.S.	1	1.31	1.09	0.93	0.93	2.29	2.38	1.00	0.87	0.82
time (sec)	N/A	0.172	0.040	0.066	0.120	0.093	0.598	0.115	0.228	0.188

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	78	61	53	54	132	122	59	55	48
N.S.	1	1.37	1.07	0.93	0.95	2.32	2.14	1.04	0.96	0.84
time (sec)	N/A	0.178	0.051	0.078	0.112	0.090	2.022	0.111	0.236	0.168

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	45	45	36	35	116	277	35	59	33
N.S.	1	1.50	1.50	1.20	1.17	3.87	9.23	1.17	1.97	1.10
time (sec)	N/A	0.155	0.053	0.066	0.117	0.103	2.498	0.119	0.253	0.166

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	46	46	37	37	115	269	36	60	34
N.S.	1	1.48	1.48	1.19	1.19	3.71	8.68	1.16	1.94	1.10
time (sec)	N/A	0.159	0.058	0.076	0.125	0.107	1.628	0.109	0.259	0.023

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	65	54	47	49	134	332	46	67	46
N.S.	1	1.30	1.08	0.94	0.98	2.68	6.64	0.92	1.34	0.92
time (sec)	N/A	0.166	0.080	0.105	0.113	0.106	3.177	0.113	0.236	0.039

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	84	68	56	63	161	389	65	84	58
N.S.	1	1.22	0.99	0.81	0.91	2.33	5.64	0.94	1.22	0.84
time (sec)	N/A	0.178	0.084	0.110	0.118	0.104	8.302	0.116	0.231	0.180

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	72	59	59	60	186	632	47	113	57
N.S.	1	1.26	1.04	1.04	1.05	3.26	11.09	0.82	1.98	1.00
time (sec)	N/A	0.170	0.077	0.072	0.112	0.118	8.740	0.114	0.219	0.188

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	72	60	52	64	186	627	52	110	56
N.S.	1	1.14	0.95	0.83	1.02	2.95	9.95	0.83	1.75	0.89
time (sec)	N/A	0.174	0.116	0.076	0.116	0.101	5.576	0.112	0.278	0.181

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	73	59	50	61	185	605	47	113	58
N.S.	1	0.84	0.68	0.57	0.70	2.13	6.95	0.54	1.30	0.67
time (sec)	N/A	0.174	0.113	0.079	0.111	0.100	10.681	0.110	0.250	0.180

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	92	70	56	73	200	683	59	120	69
N.S.	1	0.91	0.69	0.55	0.72	1.98	6.76	0.58	1.19	0.68
time (sec)	N/A	0.195	0.117	0.145	0.106	0.120	18.818	0.109	0.239	0.188

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	214	92	106	172	120	104	182	112	37
N.S.	1	2.16	0.93	1.07	1.74	1.21	1.05	1.84	1.13	0.37
time (sec)	N/A	0.380	0.124	0.062	0.127	0.087	1.662	0.116	0.230	0.054

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	B	C	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	214	91	106	172	132	104	182	112	38
N.S.	1	2.12	0.90	1.05	1.70	1.31	1.03	1.80	1.11	0.38
time (sec)	N/A	0.394	0.107	0.056	0.119	0.115	1.030	0.116	0.267	0.044

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	B	C	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	229	118	115	185	117	110	178	145	55
N.S.	1	2.04	1.05	1.03	1.65	1.04	0.98	1.59	1.29	0.49
time (sec)	N/A	0.404	0.137	0.074	0.131	0.097	1.787	0.116	0.229	0.052

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	B	C	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	231	119	116	186	158	124	178	147	54
N.S.	1	2.04	1.05	1.03	1.65	1.40	1.10	1.58	1.30	0.48
time (sec)	N/A	0.417	0.134	0.083	0.118	0.116	4.810	0.119	0.242	0.187

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	241	128	124	194	193	316	199	306	64
N.S.	1	1.90	1.01	0.98	1.53	1.52	2.49	1.57	2.41	0.50
time (sec)	N/A	0.417	0.243	0.079	0.123	0.093	25.081	0.118	0.288	0.205

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	129	241	128	127	194	204	400	199	305	64
N.S.	1	1.87	0.99	0.98	1.50	1.58	3.10	1.54	2.36	0.50
time (sec)	N/A	0.416	0.248	0.079	0.116	0.105	17.878	0.116	0.264	0.050

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	B	C	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	241	127	127	195	191	323	199	304	64
N.S.	1	1.94	1.02	1.02	1.57	1.54	2.60	1.60	2.45	0.52
time (sec)	N/A	0.412	0.269	0.087	0.121	0.097	29.131	0.119	0.243	0.051

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	241	128	124	195	200	393	199	307	64
N.S.	1	1.91	1.02	0.98	1.55	1.59	3.12	1.58	2.44	0.51
time (sec)	N/A	0.411	0.265	0.112	0.130	0.096	53.455	0.124	0.246	0.191

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	270	138	147	217	268	0	209	477	86
N.S.	1	1.86	0.95	1.01	1.50	1.85	0.00	1.44	3.29	0.59
time (sec)	N/A	0.455	0.239	0.092	0.119	0.096	0.000	0.116	0.289	0.061

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	270	138	150	217	281	887	209	478	86
N.S.	1	1.84	0.94	1.02	1.48	1.91	6.03	1.42	3.25	0.59
time (sec)	N/A	0.452	0.244	0.089	0.124	0.108	106.023	0.123	0.292	0.198

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	139	270	137	138	221	279	0	211	477	85
N.S.	1	1.94	0.99	0.99	1.59	2.01	0.00	1.52	3.43	0.61
time (sec)	N/A	0.460	0.352	0.106	0.131	0.104	0.000	0.120	0.263	0.053

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	C	F(-1)	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	177	270	136	138	222	289	0	212	478	85
N.S.	1	1.53	0.77	0.78	1.25	1.63	0.00	1.20	2.70	0.48
time (sec)	N/A	0.470	0.382	0.181	0.140	0.106	0.000	0.119	0.235	0.045

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	13	12	12	10	12	11	12
N.S.	1	1.00	1.00	0.93	0.86	0.86	0.71	0.86	0.79	0.86
time (sec)	N/A	0.180	0.002	0.060	0.032	0.093	0.019	0.116	0.267	0.014

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	32	23	21	26	19	162	23	18	25
N.S.	1	1.19	0.85	0.78	0.96	0.70	6.00	0.85	0.67	0.93
time (sec)	N/A	0.260	0.014	0.060	0.033	0.083	0.639	0.115	0.226	0.017

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	51	35	32	41	31	600	37	30	37
N.S.	1	1.31	0.90	0.82	1.05	0.79	15.38	0.95	0.77	0.95
time (sec)	N/A	0.261	0.022	0.069	0.026	0.104	0.964	0.122	0.239	0.024

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	21	21	20	12	53	27	12	13	21
N.S.	1	1.50	1.50	1.43	0.86	3.79	1.93	0.86	0.93	1.50
time (sec)	N/A	0.230	0.009	0.154	0.045	0.078	0.771	0.114	0.303	0.200

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	51	28	27	30	62	0	29	20	28
N.S.	1	2.43	1.33	1.29	1.43	2.95	0.00	1.38	0.95	1.33
time (sec)	N/A	0.338	0.017	0.165	0.051	0.083	0.000	0.108	0.301	0.223

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	B	F	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	70	41	39	41	73	0	51	31	40
N.S.	1	1.79	1.05	1.00	1.05	1.87	0.00	1.31	0.79	1.03
time (sec)	N/A	0.341	0.027	0.211	0.046	0.094	0.000	0.115	0.260	0.253

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	23	23	18	32	53	24	21	31	17
N.S.	1	0.55	0.55	0.43	0.76	1.26	0.57	0.50	0.74	0.40
time (sec)	N/A	0.239	0.020	0.073	0.118	0.095	0.578	0.112	0.301	0.029

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	35	35	28	42	70	68	32	38	27
N.S.	1	0.65	0.65	0.52	0.78	1.30	1.26	0.59	0.70	0.50
time (sec)	N/A	0.212	0.025	0.066	0.108	0.108	0.804	0.117	0.290	0.165

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	39	39	32	47	90	44	38	51	31
N.S.	1	0.64	0.64	0.52	0.77	1.48	0.72	0.62	0.84	0.51
time (sec)	N/A	0.257	0.055	0.091	0.108	0.126	0.982	0.113	0.292	0.030

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	63	55	44	88	116	97	66	72	48
N.S.	1	0.70	0.61	0.49	0.98	1.29	1.08	0.73	0.80	0.53
time (sec)	N/A	0.284	0.085	0.106	0.122	0.110	2.091	0.115	0.289	0.037

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	72	65	56	0	243	0	44	51	0
N.S.	1	1.11	1.00	0.86	0.00	3.74	0.00	0.68	0.78	0.00
time (sec)	N/A	0.333	0.062	0.166	0.000	0.117	0.000	0.116	0.294	0.000

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	76	67	68	0	260	0	50	59	0
N.S.	1	0.82	0.72	0.73	0.00	2.80	0.00	0.54	0.63	0.00
time (sec)	N/A	0.275	0.071	0.214	0.000	0.106	0.000	0.119	0.262	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	82	80	67	0	286	0	64	73	0
N.S.	1	0.61	0.59	0.50	0.00	2.12	0.00	0.47	0.54	0.00
time (sec)	N/A	0.202	0.115	0.236	0.000	0.108	0.000	0.121	0.282	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	41	41	34	60	90	44	44	51	33
N.S.	1	0.61	0.61	0.51	0.90	1.34	0.66	0.66	0.76	0.49
time (sec)	N/A	0.160	0.054	0.089	0.106	0.143	1.239	0.113	0.266	0.034

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	69	56	45	92	120	102	69	73	51
N.S.	1	0.85	0.69	0.56	1.14	1.48	1.26	0.85	0.90	0.63
time (sec)	N/A	0.174	0.081	0.084	0.112	0.100	2.594	0.113	0.255	0.176

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	59	56	47	0	276	0	37	57	0
N.S.	1	1.04	0.98	0.82	0.00	4.84	0.00	0.65	1.00	0.00
time (sec)	N/A	0.190	0.037	0.142	0.000	0.102	0.000	0.116	0.262	0.000

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	85	68	58	0	317	0	64	73	0
N.S.	1	1.18	0.94	0.81	0.00	4.40	0.00	0.89	1.01	0.00
time (sec)	N/A	0.201	0.077	0.215	0.000	0.109	0.000	0.115	0.247	0.000

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	113	87	74	0	355	0	80	92	0
N.S.	1	1.24	0.96	0.81	0.00	3.90	0.00	0.88	1.01	0.00
time (sec)	N/A	0.222	0.107	0.255	0.000	0.111	0.000	0.118	0.240	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	97	113	0	0	225	0	219	24	0
N.S.	1	1.23	1.43	0.00	0.00	2.85	0.00	2.77	0.30	0.00
time (sec)	N/A	0.223	0.090	0.000	0.000	0.098	0.000	2.970	0.248	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	90	113	87	86	91	180	87	159	107
N.S.	1	0.98	1.23	0.95	0.93	0.99	1.96	0.95	1.73	1.16
time (sec)	N/A	0.205	0.059	0.481	0.110	0.105	1.213	0.328	0.261	0.040

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	96	119	95	93	135	643	96	168	117
N.S.	1	0.79	0.98	0.78	0.76	1.11	5.27	0.79	1.38	0.96
time (sec)	N/A	0.208	0.151	0.137	0.117	0.102	1.320	0.342	0.275	0.040

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	124	129	118	139	184	2266	128	204	196
N.S.	1	0.89	0.92	0.84	0.99	1.31	16.19	0.91	1.46	1.40
time (sec)	N/A	0.223	0.217	0.152	0.122	0.106	1.992	0.328	0.298	0.242

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	104	123	157	0	0	322	0	243	24	0
N.S.	1	1.18	1.51	0.00	0.00	3.10	0.00	2.34	0.23	0.00
time (sec)	N/A	0.317	0.145	0.000	0.000	0.090	0.000	2.999	0.241	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	118	151	183	0	0	322	0	295	24	0
N.S.	1	1.28	1.55	0.00	0.00	2.73	0.00	2.50	0.20	0.00
time (sec)	N/A	0.402	0.131	0.000	0.000	0.091	0.000	2.948	0.254	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	75	95	75	76	213	155	77	141	99
N.S.	1	0.96	1.22	0.96	0.97	2.73	1.99	0.99	1.81	1.27
time (sec)	N/A	0.249	0.052	0.081	0.120	0.138	1.055	0.349	0.264	0.192

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	95	112	136	0	0	238	0	249	55	0
N.S.	1	1.18	1.43	0.00	0.00	2.51	0.00	2.62	0.58	0.00
time (sec)	N/A	0.349	0.084	0.000	0.000	0.134	0.000	2.948	0.289	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	116	139	109	0	560	0	243	62	0
N.S.	1	1.01	1.21	0.95	0.00	4.87	0.00	2.11	0.54	0.00
time (sec)	N/A	0.362	0.170	0.083	0.000	0.115	0.000	2.979	0.288	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	30	31	44	11	13	0	12	13	0
N.S.	1	0.22	0.22	0.32	0.08	0.09	0.00	0.09	0.09	0.00
time (sec)	N/A	0.238	1.013	0.283	0.033	0.080	0.000	0.113	0.265	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	101	120	95	106	306	831	100	166	130
N.S.	1	0.96	1.14	0.90	1.01	2.91	7.91	0.95	1.58	1.24
time (sec)	N/A	0.258	0.152	0.112	0.125	0.105	1.396	0.371	0.248	0.234

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	129	149	109	142	296	2730	130	206	182
N.S.	1	1.10	1.27	0.93	1.21	2.53	23.33	1.11	1.76	1.56
time (sec)	N/A	0.369	0.155	0.134	0.126	0.113	2.311	0.319	0.264	0.255

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	40	29	26	39	25	53	39	16	28
N.S.	1	0.91	0.66	0.59	0.89	0.57	1.20	0.89	0.36	0.64
time (sec)	N/A	0.319	0.072	0.069	0.027	0.090	0.409	0.111	0.244	0.168

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [12] had the largest ratio of [1.5000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	3	0.333
2	A	1	1	1.00	3	0.333
3	A	1	1	1.00	3	0.333
4	A	1	1	1.00	3	0.333
5	A	2	2	1.00	2	1.000
6	A	2	2	1.00	2	1.000
7	A	4	3	1.00	4	0.750
8	A	4	3	1.00	4	0.750
9	A	4	3	1.00	5	0.600
10	A	5	4	1.00	5	0.800
11	A	2	2	1.00	2	1.000
12	A	3	3	1.00	2	1.500
13	A	2	2	0.71	2	1.000
14	A	2	2	0.50	2	1.000
15	A	1	1	1.00	7	0.143
16	A	1	1	1.00	9	0.111
17	A	1	1	1.00	11	0.091
18	A	1	1	1.00	9	0.111
19	A	3	2	1.00	9	0.222
20	A	3	3	1.00	2	1.500
21	A	2	2	1.00	2	1.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	5	4	1.00	4	1.000
23	A	4	3	1.00	4	0.750
24	A	3	3	1.00	2	1.500
25	A	3	3	1.00	2	1.500
26	A	3	3	0.71	2	1.500
27	A	1	1	1.00	7	0.143
28	A	1	1	1.00	7	0.143
29	A	2	2	1.00	9	0.222
30	A	2	2	1.00	11	0.182
31	A	1	1	1.00	7	0.143
32	A	2	2	0.96	9	0.222
33	A	2	2	1.00	11	0.182
34	A	1	1	1.00	7	0.143
35	A	1	1	0.74	9	0.111
36	A	2	2	1.11	11	0.182
37	A	2	2	0.88	11	0.182
38	A	1	1	1.00	7	0.143
39	A	2	2	1.20	9	0.222
40	A	1	1	0.50	11	0.091
41	A	2	2	1.16	11	0.182
42	A	1	1	1.00	7	0.143
43	A	2	2	1.20	9	0.222
44	A	2	2	1.24	11	0.182
45	A	1	1	0.36	11	0.091
46	A	3	3	1.20	11	0.273
47	A	2	2	1.17	11	0.182
48	A	2	2	1.14	11	0.182
49	A	2	2	1.05	11	0.182
50	A	2	2	1.02	11	0.182
51	A	2	2	1.13	11	0.182
52	A	2	2	1.12	11	0.182
53	A	2	2	1.19	11	0.182

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	2	2	1.12	11	0.182
55	A	2	2	1.13	11	0.182
56	A	2	2	1.18	11	0.182
57	A	2	2	1.11	11	0.182
58	A	2	2	1.13	11	0.182
59	A	2	2	1.18	11	0.182
60	A	1	1	1.20	9	0.111
61	A	1	1	1.08	13	0.077
62	A	9	8	1.18	9	0.889
63	A	9	8	1.15	11	0.727
64	A	1	1	1.00	13	0.077
65	A	10	9	1.23	13	0.692
66	A	10	9	1.18	13	0.692
67	A	10	9	1.21	9	1.000
68	A	10	9	1.15	11	0.818
69	A	1	1	1.00	13	0.077
70	A	10	9	1.18	13	0.692
71	A	5	4	1.24	13	0.308
72	A	10	9	1.19	13	0.692
73	A	10	9	1.21	13	0.692
74	A	4	3	1.03	13	0.231
75	A	11	10	1.20	13	0.769
76	A	11	10	1.23	13	0.769
77	B	9	8	3.17	9	0.889
78	A	3	2	1.16	11	0.182
79	B	9	8	3.17	13	0.615
80	A	1	1	1.00	13	0.077
81	B	10	9	2.78	9	1.000
82	A	4	3	1.10	11	0.273
83	B	10	9	2.64	13	0.692
84	A	1	1	1.00	13	0.077
85	A	5	4	1.24	13	0.308

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	B	10	9	2.99	13	0.692
87	A	1	1	1.00	5	0.200
88	A	1	1	1.00	7	0.143
89	A	8	7	1.05	7	1.000
90	A	9	8	1.38	7	1.143
91	A	1	1	1.00	7	0.143
92	A	1	1	1.00	9	0.111
93	A	1	1	1.00	7	0.143
94	A	7	6	1.05	9	0.667
95	A	3	3	1.44	9	0.333
96	A	2	2	1.00	7	0.286
97	A	1	1	1.00	9	0.111
98	A	8	7	1.12	9	0.778
99	A	3	2	1.00	9	0.222
100	A	2	2	1.00	9	0.222
101	A	1	1	1.00	11	0.091
102	A	7	6	1.12	11	0.545
103	A	3	2	0.40	11	0.182
104	A	5	4	1.42	11	0.364
105	A	5	4	1.36	13	0.308
106	A	2	2	1.04	13	0.154
107	A	2	2	1.44	15	0.133
108	A	2	2	1.26	16	0.125
109	A	3	2	1.81	13	0.154
110	A	4	3	1.29	13	0.231
111	A	5	4	1.31	13	0.308
112	A	6	5	1.37	13	0.385
113	A	4	3	1.50	13	0.231
114	A	4	3	1.48	13	0.231
115	A	5	4	1.30	13	0.308
116	A	6	5	1.22	13	0.385
117	A	5	4	1.26	13	0.308

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
118	A	5	4	1.14	13	0.308
119	A	5	4	0.84	13	0.308
120	A	6	5	0.91	13	0.385
121	B	10	9	2.16	15	0.600
122	B	10	9	2.12	15	0.600
123	B	11	10	2.04	15	0.667
124	B	11	10	2.04	15	0.667
125	A	11	10	1.90	15	0.667
126	A	11	10	1.87	15	0.667
127	A	11	10	1.94	15	0.667
128	A	11	10	1.91	15	0.667
129	A	12	11	1.86	15	0.733
130	A	12	11	1.84	15	0.733
131	A	12	11	1.94	15	0.733
132	A	12	11	1.53	15	0.733
133	A	1	1	1.00	9	0.111
134	A	2	2	1.19	11	0.182
135	A	2	2	1.31	13	0.154
136	A	4	3	1.50	11	0.273
137	B	3	3	2.43	13	0.231
138	A	3	3	1.79	15	0.200
139	A	3	2	0.55	13	0.154
140	A	4	3	0.65	13	0.231
141	A	4	3	0.64	13	0.231
142	A	5	4	0.70	13	0.308
143	A	6	5	1.11	15	0.333
144	A	6	5	0.82	15	0.333
145	A	6	5	0.61	15	0.333
146	A	4	3	0.61	13	0.231
147	A	5	4	0.85	13	0.308
148	A	5	4	1.04	15	0.267
149	A	6	5	1.18	15	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
150	A	7	6	1.24	15	0.400
151	A	6	5	1.23	15	0.333
152	A	6	5	0.98	13	0.385
153	A	6	5	0.79	13	0.385
154	A	7	6	0.89	13	0.462
155	A	7	6	1.18	15	0.400
156	A	8	7	1.28	15	0.467
157	A	5	4	0.96	13	0.308
158	A	7	6	1.18	15	0.400
159	A	7	6	1.01	15	0.400
160	A	3	3	0.22	15	0.200
161	A	6	5	0.96	13	0.385
162	A	7	6	1.10	13	0.462
163	A	2	2	0.91	15	0.133

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^n dx$	85
3.2	$\int \frac{1}{x} dx$	90
3.3	$\int e^x dx$	94
3.4	$\int a^x dx$	99
3.5	$\int \sin(x) dx$	104
3.6	$\int \cos(x) dx$	109
3.7	$\int \csc^2(x) dx$	114
3.8	$\int \sec^2(x) dx$	119
3.9	$\int \sec(x) \tan(x) dx$	124
3.10	$\int \cot(x) \csc(x) dx$	129
3.11	$\int \tan(x) dx$	134
3.12	$\int \cot(x) dx$	139
3.13	$\int \csc(x) dx$	144
3.14	$\int \sec(x) dx$	149
3.15	$\int \frac{1}{1+x^2} dx$	154
3.16	$\int \frac{1}{1-x^2} dx$	159
3.17	$\int \frac{1}{\sqrt{1-x^2}} dx$	164
3.18	$\int \frac{1}{\sqrt{1+x^2}} dx$	169
3.19	$\int \frac{1}{\sqrt{-1+x^2}} dx$	174
3.20	$\int \sinh(x) dx$	179
3.21	$\int \cosh(x) dx$	184
3.22	$\int \operatorname{csch}^2(x) dx$	189
3.23	$\int \operatorname{sech}^2(x) dx$	194
3.24	$\int \tanh(x) dx$	199
3.25	$\int \operatorname{coth}(x) dx$	204
3.26	$\int \operatorname{csch}(x) dx$	209

3.27	$\int (a + bx)^m dx$	214
3.28	$\int \frac{1}{a+bx} dx$	219
3.29	$\int \frac{x}{a+bx} dx$	224
3.30	$\int \frac{x^2}{a+bx} dx$	229
3.31	$\int \frac{1}{(a+bx)^2} dx$	234
3.32	$\int \frac{x}{(a+bx)^2} dx$	239
3.33	$\int \frac{x^2}{(a+bx)^2} dx$	244
3.34	$\int \frac{1}{(a+bx)^3} dx$	249
3.35	$\int \frac{x}{(a+bx)^3} dx$	254
3.36	$\int \frac{x^2}{(a+bx)^3} dx$	259
3.37	$\int \frac{x^3}{(a+bx)^3} dx$	264
3.38	$\int \frac{1}{(a+bx)^4} dx$	269
3.39	$\int \frac{x}{(a+bx)^4} dx$	274
3.40	$\int \frac{x^2}{(a+bx)^4} dx$	279
3.41	$\int \frac{x^3}{(a+bx)^4} dx$	284
3.42	$\int \frac{1}{(a+bx)^5} dx$	289
3.43	$\int \frac{x}{(a+bx)^5} dx$	294
3.44	$\int \frac{x^2}{(a+bx)^5} dx$	299
3.45	$\int \frac{x^3}{(a+bx)^5} dx$	304
3.46	$\int \frac{1}{x(a+bx)} dx$	309
3.47	$\int \frac{1}{x^2(a+bx)} dx$	314
3.48	$\int \frac{1}{x^3(a+bx)} dx$	319
3.49	$\int \frac{1}{x^2(a+bx)^2} dx$	324
3.50	$\int \frac{1}{x^3(a+bx)^2} dx$	329
3.51	$\int \frac{1}{x(a+bx)^3} dx$	334
3.52	$\int \frac{1}{x^2(a+bx)^3} dx$	339
3.53	$\int \frac{1}{x^3(a+bx)^3} dx$	344
3.54	$\int \frac{1}{x(a+bx)^4} dx$	349
3.55	$\int \frac{1}{x^2(a+bx)^4} dx$	354
3.56	$\int \frac{1}{x^3(a+bx)^4} dx$	359
3.57	$\int \frac{1}{x(a+bx)^5} dx$	365
3.58	$\int \frac{1}{x^2(a+bx)^5} dx$	371
3.59	$\int \frac{1}{x^3(a+bx)^5} dx$	377
3.60	$\int \frac{1}{a+bx^2} dx$	383
3.61	$\int x(a + bx^2)^{-m} dx$	388
3.62	$\int \frac{1}{a+bx^3} dx$	393

3.63	$\int \frac{x}{a+bx^3} dx$	401
3.64	$\int \frac{x^2}{a+bx^3} dx$	409
3.65	$\int \frac{x^3}{a+bx^3} dx$	414
3.66	$\int \frac{x^4}{a+bx^3} dx$	423
3.67	$\int \frac{1}{(a+bx^3)^2} dx$	432
3.68	$\int \frac{x}{(a+bx^3)^2} dx$	442
3.69	$\int \frac{x^2}{(a+bx^3)^2} dx$	451
3.70	$\int \frac{x^3}{(a+bx^3)^2} dx$	456
3.71	$\int \frac{1}{x(a+bx^3)} dx$	466
3.72	$\int \frac{1}{x^2(a+bx^3)} dx$	471
3.73	$\int \frac{1}{x^3(a+bx^3)} dx$	480
3.74	$\int \frac{1}{x(a+bx^3)^2} dx$	489
3.75	$\int \frac{1}{x^2(a+bx^3)^2} dx$	494
3.76	$\int \frac{1}{x^3(a+bx^3)^2} dx$	505
3.77	$\int \frac{1}{a+bx^4} dx$	517
3.78	$\int \frac{x}{a+bx^4} dx$	526
3.79	$\int \frac{x^2}{a+bx^4} dx$	531
3.80	$\int \frac{x^3}{a+bx^4} dx$	540
3.81	$\int \frac{1}{(a+bx^4)^2} dx$	545
3.82	$\int \frac{x}{(a+bx^4)^2} dx$	555
3.83	$\int \frac{x^2}{(a+bx^4)^2} dx$	561
3.84	$\int \frac{x^3}{(a+bx^4)^2} dx$	570
3.85	$\int \frac{1}{x(a+bx^4)} dx$	575
3.86	$\int \frac{1}{x^2(a+bx^4)} dx$	580
3.87	$\int \frac{1}{1+x} dx$	590
3.88	$\int \frac{1}{1+x^2} dx$	595
3.89	$\int \frac{1}{1+x^3} dx$	600
3.90	$\int \frac{1}{1+x^4} dx$	606
3.91	$\int \frac{1}{1-x} dx$	613
3.92	$\int \frac{1}{1-x^2} dx$	618
3.93	$\int \frac{1}{-1+x^2} dx$	623
3.94	$\int \frac{1}{1-x^3} dx$	628
3.95	$\int \frac{1}{1-x^4} dx$	634
3.96	$\int \frac{x}{1+x} dx$	639
3.97	$\int \frac{x}{1+x^2} dx$	644
3.98	$\int \frac{x}{1+x^3} dx$	649

3.99	$\int \frac{x}{1+x^4} dx$	655
3.100	$\int \frac{x}{1-x} dx$	660
3.101	$\int \frac{x}{1-x^2} dx$	665
3.102	$\int \frac{x}{1-x^3} dx$	670
3.103	$\int \frac{x}{1-x^4} dx$	676
3.104	$\int \frac{1}{x(1+x^2)} dx$	681
3.105	$\int \frac{1}{x(1-x^2)} dx$	686
3.106	$\int \frac{a+bx}{A+Bx} dx$	691
3.107	$\int \frac{1}{(a+bx)(A+Bx)} dx$	696
3.108	$\int \frac{x}{(a+bx)(A+Bx)} dx$	701
3.109	$\int \frac{1}{\sqrt{x}(a+bx)} dx$	706
3.110	$\int \frac{\sqrt{x}}{a+bx} dx$	711
3.111	$\int \frac{x^{3/2}}{a+bx} dx$	716
3.112	$\int \frac{x^{5/2}}{a+bx} dx$	722
3.113	$\int \frac{1}{\sqrt{x}(a+bx)^2} dx$	729
3.114	$\int \frac{\sqrt{x}}{(a+bx)^2} dx$	735
3.115	$\int \frac{x^{3/2}}{(a+bx)^2} dx$	741
3.116	$\int \frac{x^{5/2}}{(a+bx)^2} dx$	747
3.117	$\int \frac{1}{\sqrt{x}(a+bx)^3} dx$	754
3.118	$\int \frac{\sqrt{x}}{(a+bx)^3} dx$	760
3.119	$\int \frac{x^{3/2}}{(a+bx)^3} dx$	767
3.120	$\int \frac{x^{5/2}}{(a+bx)^3} dx$	773
3.121	$\int \frac{1}{\sqrt{x}(a+bx^2)} dx$	780
3.122	$\int \frac{\sqrt{x}}{a+bx^2} dx$	790
3.123	$\int \frac{x^{3/2}}{a+bx^2} dx$	800
3.124	$\int \frac{x^{5/2}}{a+bx^2} dx$	810
3.125	$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx$	820
3.126	$\int \frac{\sqrt{x}}{(a+bx^2)^2} dx$	831
3.127	$\int \frac{x^{3/2}}{(a+bx^2)^2} dx$	841
3.128	$\int \frac{x^{5/2}}{(a+bx^2)^2} dx$	850
3.129	$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx$	860
3.130	$\int \frac{\sqrt{x}}{(a+bx^2)^3} dx$	872
3.131	$\int \frac{x^{3/2}}{(a+bx^2)^3} dx$	884
3.132	$\int \frac{x^{5/2}}{(a+bx^2)^3} dx$	895

3.133	$\int \frac{1}{\sqrt{a+bx}} dx$	906
3.134	$\int \frac{x}{\sqrt{a+bx}} dx$	911
3.135	$\int \frac{x^2}{\sqrt{a+bx}} dx$	916
3.136	$\int \frac{1}{\sqrt{(a+bx)^3}} dx$	922
3.137	$\int \frac{x}{\sqrt{(a+bx)^3}} dx$	927
3.138	$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx$	932
3.139	$\int \frac{1}{x\sqrt{a+bx}} dx$	937
3.140	$\int \frac{\sqrt{a+bx}}{x} dx$	942
3.141	$\int \frac{\sqrt{a+bx}}{x^2} dx$	947
3.142	$\int \frac{\sqrt{a+bx}}{x^3} dx$	953
3.143	$\int \frac{\sqrt{(a+bx)^3}}{x} dx$	959
3.144	$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx$	965
3.145	$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx$	971
3.146	$\int \frac{1}{x^2\sqrt{a+bx}} dx$	977
3.147	$\int \frac{1}{x^3\sqrt{a+bx}} dx$	983
3.148	$\int \frac{1}{x\sqrt{(a+bx)^3}} dx$	989
3.149	$\int \frac{1}{x^2\sqrt{(a+bx)^3}} dx$	995
3.150	$\int \frac{1}{x^3\sqrt{(a+bx)^3}} dx$	1002
3.151	$\int \frac{1}{x\sqrt[3]{(a+bx)^2}} dx$	1010
3.152	$\int \frac{\sqrt[3]{a+bx}}{x} dx$	1017
3.153	$\int \frac{\sqrt[3]{a+bx}}{x^2} dx$	1025
3.154	$\int \frac{\sqrt[3]{a+bx}}{x^3} dx$	1033
3.155	$\int \frac{1}{x^2\sqrt[3]{(a+bx)^2}} dx$	1043
3.156	$\int \frac{1}{x^3\sqrt[3]{(a+bx)^2}} dx$	1051
3.157	$\int \frac{1}{x\sqrt[3]{a+bx}} dx$	1060
3.158	$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx$	1068
3.159	$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx$	1075
3.160	$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx$	1083
3.161	$\int \frac{1}{x^2\sqrt[3]{a+bx}} dx$	1088
3.162	$\int \frac{1}{x^3\sqrt[3]{a+bx}} dx$	1097
3.163	$\int \frac{A+Bx}{\sqrt{a+bx}} dx$	1108

3.1 $\int x^n dx$

Optimal result	85
Mathematica [A] (verified)	85
Rubi [A] (verified)	86
Maple [A] (verified)	87
Fricas [A] (verification not implemented)	87
Sympy [A] (verification not implemented)	88
Maxima [A] (verification not implemented)	88
Giac [A] (verification not implemented)	88
Mupad [B] (verification not implemented)	89
Reduce [B] (verification not implemented)	89

Optimal result

Integrand size = 3, antiderivative size = 11

$$\int x^n dx = \frac{x^{1+n}}{1+n}$$

output `x^(1+n)/(1+n)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int x^n dx = \frac{x^{1+n}}{1+n}$$

input `Integrate[x^n,x]`

output `x^(1 + n)/(1 + n)`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x^n dx$$

$$\downarrow 15$$

$$\frac{x^{n+1}}{n+1}$$

input `Int [x^n, x]`

output `x^(1 + n)/(1 + n)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] :> Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

method	result	size
risch	$\frac{x x^n}{1+n}$	11
parallelrisch	$\frac{x x^n}{1+n}$	11
orering	$\frac{x x^n}{1+n}$	11
gosper	$\frac{x^{1+n}}{1+n}$	12
default	$\frac{x^{1+n}}{1+n}$	12
norman	$\frac{x e^{n \ln(x)}}{1+n}$	13

input `int(x^n,x,method=_RETURNVERBOSE)`

output `x/(1+n)*x^n`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int x^n dx = \frac{xx^n}{n+1}$$

input `integrate(x^n,x, algorithm="fricas")`

output `x*x^n/(n + 1)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.09

$$\int x^n dx = \begin{cases} \frac{x^{n+1}}{n+1} & \text{for } n \neq -1 \\ \log(x) & \text{otherwise} \end{cases}$$

input `integrate(x**n,x)`

output `Piecewise((x**(n + 1)/(n + 1), Ne(n, -1)), (log(x), True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int x^n dx = \frac{x^{n+1}}{n+1}$$

input `integrate(x^n,x, algorithm="maxima")`

output `x^(n + 1)/(n + 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int x^n dx = \frac{x^{n+1}}{n+1}$$

input `integrate(x^n,x, algorithm="giac")`

output `x^(n + 1)/(n + 1)`

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.82

$$\int x^n dx = \begin{cases} \ln(x) & \text{if } n = -1 \\ \frac{x^{n+1}}{n+1} & \text{if } n \neq -1 \end{cases}$$

input `int(x^n,x)`

output `piecewise(n == -1, log(x), n ~= -1, x^(n + 1)/(n + 1))`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.91

$$\int x^n dx = \frac{x^{n+1}}{n+1}$$

input `int(x^n,x)`

output `(x**(n+1))/(n + 1)`

3.2 $\int \frac{1}{x} dx$

Optimal result	90
Mathematica [A] (verified)	90
Rubi [A] (verified)	91
Maple [A] (verified)	91
Fricas [A] (verification not implemented)	92
Sympy [A] (verification not implemented)	92
Maxima [A] (verification not implemented)	92
Giac [A] (verification not implemented)	93
Mupad [B] (verification not implemented)	93
Reduce [B] (verification not implemented)	93

Optimal result

Integrand size = 3, antiderivative size = 2

$$\int \frac{1}{x} dx = \log(x)$$

output `ln(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `Integrate[x^(-1), x]`

output `Log[x]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {14}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x} dx$$

↓ 14

$$\log(x)$$

input `Int [x(-1), x]`

output `Log [x]`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] :> Simp[a*Log[x], x] /; FreeQ[a, x]`

Maple [A] (verified)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\ln(x)$	3
norman	$\ln(x)$	3
risch	$\ln(x)$	3
parallelrisc	$\ln(x)$	3

input `int(1/x,x,method=_RETURNVERBOSE)`

output `ln(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x, algorithm="fricas")`

output `log(x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x)`

output `log(x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `integrate(1/x,x, algorithm="maxima")`

output `log(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \frac{1}{x} dx = \log(|x|)$$

input `integrate(1/x,x, algorithm="giac")`

output `log(abs(x))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \ln(x)$$

input `int(1/x,x)`

output `log(x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{x} dx = \log(x)$$

input `int(1/x,x)`

output `log(x)`

3.3 $\int e^x dx$

Optimal result	94
Mathematica [A] (verified)	94
Rubi [A] (verified)	95
Maple [A] (verified)	96
Fricas [A] (verification not implemented)	96
Sympy [A] (verification not implemented)	97
Maxima [A] (verification not implemented)	97
Giac [A] (verification not implemented)	97
Mupad [B] (verification not implemented)	98
Reduce [B] (verification not implemented)	98

Optimal result

Integrand size = 3, antiderivative size = 3

$$\int e^x dx = e^x$$

output `exp(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e^x dx = e^x$$

input `Integrate[E^x,x]`

output `E^x`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int e^x dx$$

$$\downarrow 2624$$

$$e^x$$

input `Int [E^x,x]`

output `E^x`

Defintions of rubi rules used

rule 2624 `Int[((F_)^(v_))^(n_), x_Symbol] := Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;`
`FreeQ[{F, n}, x] && LinearQ[v, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

method	result	size
gospers	e^x	3
lookup	e^x	3
derivativedivides	e^x	3
default	e^x	3
norman	e^x	3
risch	e^x	3
parallelrisch	e^x	3
orering	e^x	3
meijerg	$-1 + e^x$	5

input `int(exp(x),x,method=_RETURNVERBOSE)`

output `exp(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="fricas")`

output `e^x`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x)`

output `exp(x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="maxima")`

output `e^x`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `integrate(exp(x),x, algorithm="giac")`

output `e^x`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.67

$$\int e^x dx = e^x$$

input `int(exp(x),x)`

output `exp(x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int e^x dx = e^x$$

input `int(exp(x),x)`

output `e**x`

3.4 $\int a^x dx$

Optimal result	99
Mathematica [A] (verified)	99
Rubi [A] (verified)	100
Maple [A] (verified)	101
Fricas [A] (verification not implemented)	101
Sympy [A] (verification not implemented)	102
Maxima [A] (verification not implemented)	102
Giac [A] (verification not implemented)	102
Mupad [B] (verification not implemented)	103
Reduce [B] (verification not implemented)	103

Optimal result

Integrand size = 3, antiderivative size = 8

$$\int a^x dx = \frac{a^x}{\log(x)}$$

output `a^x/ln(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `Integrate[a^x,x]`

output `a^x/Log[a]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2624}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int a^x dx$$

$$\downarrow 2624$$

$$\frac{a^x}{\log(a)}$$

input `Int[a^x, x]`

output `a^x/Log[a]`

Defintions of rubi rules used

rule 2624

```
Int[((F_)^(v_))^(n_), x_Symbol] :> Simp[(F^v)^n/(n*Log[F]*D[v, x]), x] /;
FreeQ[{F, n}, x] && LinearQ[v, x]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
gosper	$\frac{a^x}{\ln(a)}$	9
derivativedivides	$\frac{a^x}{\ln(a)}$	9
default	$\frac{a^x}{\ln(a)}$	9
risch	$\frac{a^x}{\ln(a)}$	9
parallelrisch	$\frac{a^x}{\ln(a)}$	9
orering	$\frac{a^x}{\ln(a)}$	9
norman	$\frac{e^{x \ln(a)}}{\ln(a)}$	11
meijerg	$-\frac{1-e^{x \ln(a)}}{\ln(a)}$	16

input `int(a^x,x,method=_RETURNVERBOSE)`

output `1/ln(a)*a^x`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `integrate(a^x,x, algorithm="fricas")`

output `a^x/log(a)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \begin{cases} \frac{a^x}{\log(a)} & \text{for } \log(a) \neq 0 \\ x & \text{otherwise} \end{cases}$$

input `integrate(a**x,x)`

output `Piecewise((a**x/log(a), Ne(log(a), 0)), (x, True))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `integrate(a^x,x, algorithm="maxima")`

output `a^x/log(a)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `integrate(a^x,x, algorithm="giac")`

output `a^x/log(a)`

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\ln(a)}$$

input `int(a^x,x)`

output `a^x/log(a)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int a^x dx = \frac{a^x}{\log(a)}$$

input `int(a^x,x)`

output `a**x/log(a)`

3.5 $\int \sin(x) dx$

Optimal result	104
Mathematica [A] (verified)	104
Rubi [A] (verified)	105
Maple [A] (verified)	106
Fricas [A] (verification not implemented)	106
Sympy [A] (verification not implemented)	107
Maxima [A] (verification not implemented)	107
Giac [A] (verification not implemented)	107
Mupad [B] (verification not implemented)	108
Reduce [B] (verification not implemented)	108

Optimal result

Integrand size = 2, antiderivative size = 4

$$\int \sin(x) dx = -\cos(x)$$

output `-cos(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `Integrate[Sin[x],x]`

output `-Cos[x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sin(x) dx \\ \downarrow 3042 \\ \int \sin(x) dx \\ \downarrow 3118 \\ -\cos(x) \end{array}$$

input `Int[Sin[x],x]`

output `-Cos[x]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
lookup	$-\cos(x)$	5
default	$-\cos(x)$	5
risch	$-\cos(x)$	5
orering	$-\cos(x)$	5
parallelrisch	$-\cos(x) - 1$	7
norman	$-\frac{2}{1+\tan(\frac{x}{2})^2}$	13
meijerg	$\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(x)}{\sqrt{\pi}} \right)$	16

input `int(sin(x),x,method=_RETURNVERBOSE)`

output `−cos(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x, algorithm="fricas")`

output `−cos(x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x)`

output `-cos(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x, algorithm="maxima")`

output `-cos(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `integrate(sin(x),x, algorithm="giac")`

output `-cos(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `int(sin(x),x)`

output `-cos(x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \sin(x) dx = -\cos(x)$$

input `int(sin(x),x)`

output `- cos(x)`

3.6 $\int \cos(x) dx$

Optimal result	109
Mathematica [A] (verified)	109
Rubi [A] (verified)	110
Maple [A] (verified)	111
Fricas [A] (verification not implemented)	111
Sympy [A] (verification not implemented)	112
Maxima [A] (verification not implemented)	112
Giac [A] (verification not implemented)	112
Mupad [B] (verification not implemented)	113
Reduce [B] (verification not implemented)	113

Optimal result

Integrand size = 2, antiderivative size = 2

$$\int \cos(x) dx = \sin(x)$$

output `sin(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `Integrate[Cos[x], x]`

output `Sin[x]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \cos(x) dx \\ \downarrow 3042 \\ \int \sin\left(x + \frac{\pi}{2}\right) dx \\ \downarrow 3117 \\ \sin(x) \end{array}$$

input `Int[Cos[x], x]`

output `Sin[x]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
lookup	$\sin(x)$	3
default	$\sin(x)$	3
meijerg	$\sin(x)$	3
risch	$\sin(x)$	3
parallelrisc	$\sin(x)$	3
orering	$\sin(x)$	3
norman	$\frac{2 \tan(\frac{x}{2})}{1 + \tan(\frac{x}{2})^2}$	17

input `int(cos(x),x,method=_RETURNVERBOSE)`

output `sin(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="fricas")`

output `sin(x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x)`

output `sin(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="maxima")`

output `sin(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `integrate(cos(x),x, algorithm="giac")`

output `sin(x)`

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `int(cos(x),x)`

output `sin(x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cos(x) dx = \sin(x)$$

input `int(cos(x),x)`

output `sin(x)`

3.7 $\int \csc^2(x) dx$

Optimal result	114
Mathematica [A] (verified)	114
Rubi [A] (verified)	115
Maple [A] (verified)	116
Fricas [A] (verification not implemented)	116
Sympy [B] (verification not implemented)	117
Maxima [A] (verification not implemented)	117
Giac [A] (verification not implemented)	117
Mupad [B] (verification not implemented)	118
Reduce [B] (verification not implemented)	118

Optimal result

Integrand size = 4, antiderivative size = 4

$$\int \csc^2(x) dx = -\cot(x)$$

output `-cot(x)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \csc^2(x) dx = -\cot(x)$$

input `Integrate[Csc[x]^2,x]`

output `-Cot[x]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \csc^2(x) dx \\ & \quad \downarrow \text{3042} \\ & \int \csc(x)^2 dx \\ & \quad \downarrow \text{4254} \\ & - \int 1 d \cot(x) \\ & \quad \downarrow \text{24} \\ & - \cot(x) \end{aligned}$$

input `Int [Csc [x]^2,x]`

output `-Cot [x]`

Defintions of rubi rules used

rule 24 `Int [a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$-\cot(x)$	5
risch	$-\frac{2i}{e^{2ix}-1}$	13
parallelrisch	$\frac{\tan(\frac{x}{2})}{2} - \frac{\cot(\frac{x}{2})}{2}$	14
norman	$\frac{-\frac{1}{2} + \frac{\tan(\frac{x}{2})^2}{2}}{\tan(\frac{x}{2})}$	18

input `int(csc(x)^2,x,method=_RETURNVERBOSE)`

output `-cot(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 2.00

$$\int \csc^2(x) dx = -\frac{\cos(x)}{\sin(x)}$$

input `integrate(csc(x)^2,x, algorithm="fricas")`

output `-cos(x)/sin(x)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.75

$$\int \csc^2(x) dx = -\frac{\cos(x)}{\sin(x)}$$

input `integrate(csc(x)**2,x)`

output `-cos(x)/sin(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \csc^2(x) dx = -\frac{1}{\tan(x)}$$

input `integrate(csc(x)^2,x, algorithm="maxima")`

output `-1/tan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \csc^2(x) dx = -\frac{1}{\tan(x)}$$

input `integrate(csc(x)^2,x, algorithm="giac")`

output `-1/tan(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \csc^2(x) dx = -\cot(x)$$

input `int(1/sin(x)^2,x)`

output `-cot(x)`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 8, normalized size of antiderivative = 2.00

$$\int \csc^2(x) dx = -\frac{\cos(x)}{\sin(x)}$$

input `int(csc(x)^2,x)`

output `(- cos(x))/sin(x)`

3.8 $\int \sec^2(x) dx$

Optimal result	119
Mathematica [A] (verified)	119
Rubi [A] (verified)	120
Maple [A] (verified)	121
Fricas [B] (verification not implemented)	121
Sympy [B] (verification not implemented)	122
Maxima [A] (verification not implemented)	122
Giac [A] (verification not implemented)	122
Mupad [B] (verification not implemented)	123
Reduce [B] (verification not implemented)	123

Optimal result

Integrand size = 4, antiderivative size = 2

$$\int \sec^2(x) dx = \tan(x)$$

output `tan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `Integrate[Sec[x]^2,x]`

output `Tan[x]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \sec^2(x) dx \\
 \downarrow 3042 \\
 \int \csc\left(x + \frac{\pi}{2}\right)^2 dx \\
 \downarrow 4254 \\
 - \int 1d(-\tan(x)) \\
 \downarrow 24 \\
 \tan(x)
 \end{array}$$

input `Int [Sec [x]^2,x]`

output `Tan [x]`

Defintions of rubi rules used

rule 24 `Int [a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinear Q[u, x]`

rule 4254

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\tan(x)$	3
parallelrisch	$\frac{\sin(x)}{\cos(x)}$	8
risch	$\frac{2i}{e^{2ix}+1}$	13
norman	$-\frac{2 \tan(\frac{x}{2})}{\tan(\frac{x}{2})^2-1}$	17

input

```
int(sec(x)^2,x,method=_RETURNVERBOSE)
```

output

```
tan(x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(2) = 4$.

Time = 0.09 (sec) , antiderivative size = 7, normalized size of antiderivative = 3.50

$$\int \sec^2(x) dx = \frac{\sin(x)}{\cos(x)}$$

input

```
integrate(sec(x)^2,x, algorithm="fricas")
```

output

```
sin(x)/cos(x)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5 vs. $2(2) = 4$.

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 2.50

$$\int \sec^2(x) dx = \frac{\sin(x)}{\cos(x)}$$

input `integrate(sec(x)**2,x)`

output `sin(x)/cos(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `integrate(sec(x)^2,x, algorithm="maxima")`

output `tan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `integrate(sec(x)^2,x, algorithm="giac")`

output `tan(x)`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec^2(x) dx = \tan(x)$$

input `int(1/cos(x)^2,x)`

output `tan(x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 7, normalized size of antiderivative = 3.50

$$\int \sec^2(x) dx = \frac{\sin(x)}{\cos(x)}$$

input `int(sec(x)^2,x)`

output `sin(x)/cos(x)`

3.9 $\int \sec(x) \tan(x) dx$

Optimal result	124
Mathematica [A] (verified)	124
Rubi [A] (verified)	125
Maple [A] (verified)	126
Fricas [A] (verification not implemented)	126
Sympy [A] (verification not implemented)	127
Maxima [A] (verification not implemented)	127
Giac [A] (verification not implemented)	127
Mupad [B] (verification not implemented)	128
Reduce [B] (verification not implemented)	128

Optimal result

Integrand size = 5, antiderivative size = 2

$$\int \sec(x) \tan(x) dx = \sec(x)$$

output `sec(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec(x) \tan(x) dx = \sec(x)$$

input `Integrate[Sec[x]*Tan[x],x]`

output `Sec[x]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3042, 3086, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \tan(x) \sec(x) dx \\ \downarrow 3042 \\ \int \tan(x) \sec(x) dx \\ \downarrow 3086 \\ \int 1 d\sec(x) \\ \downarrow 24 \\ \sec(x) \end{array}$$

input `Int [Sec [x] *Tan [x] , x]`

output `Sec [x]`

Defintions of rubi rules used

rule 24 `Int [a_ , x_Symbol] :> Simp [a*x, x] /; FreeQ [a, x]`

rule 3042 `Int [u_ , x_Symbol] :> Int [DeactivateTrig [u, x] , x] /; FunctionOfTrigOfLinear Q [u, x]`

rule 3086

```
Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(
n_.), x_Symbol] :> Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2
), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2
] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
derivativedivides	$\sec(x)$	3
default	$\sec(x)$	3
risch	$\frac{2e^{ix}}{e^{2ix}+1}$	17

input `int(sec(x)*tan(x),x,method=_RETURNVERBOSE)`output `sec(x)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 4, normalized size of antiderivative = 2.00

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x, algorithm="fricas")`output `1/cos(x)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x)`

output `1/cos(x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 4, normalized size of antiderivative = 2.00

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x, algorithm="maxima")`

output `1/cos(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 4, normalized size of antiderivative = 2.00

$$\int \sec(x) \tan(x) dx = \frac{1}{\cos(x)}$$

input `integrate(sec(x)*tan(x),x, algorithm="giac")`

output `1/cos(x)`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 12, normalized size of antiderivative = 6.00

$$\int \sec(x) \tan(x) dx = -\frac{2}{\tan\left(\frac{x}{2}\right)^2 - 1}$$

input `int(tan(x)/cos(x),x)`

output `-2/(tan(x/2)^2 - 1)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sec(x) \tan(x) dx = \sec(x)$$

input `int(sec(x)*tan(x),x)`

output `sec(x)`

3.10 $\int \cot(x) \csc(x) dx$

Optimal result	129
Mathematica [A] (verified)	129
Rubi [A] (verified)	130
Maple [A] (verified)	131
Fricas [A] (verification not implemented)	132
Sympy [A] (verification not implemented)	132
Maxima [A] (verification not implemented)	132
Giac [A] (verification not implemented)	133
Mupad [B] (verification not implemented)	133
Reduce [B] (verification not implemented)	133

Optimal result

Integrand size = 5, antiderivative size = 4

$$\int \cot(x) \csc(x) dx = -\csc(x)$$

output `-csc(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cot(x) \csc(x) dx = -\csc(x)$$

input `Integrate[Cot[x]*Csc[x],x]`

output `-Csc[x]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {3042, 25, 3086, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \cot(x) \csc(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int \tan\left(x - \frac{\pi}{2}\right) \left(-\sec\left(x - \frac{\pi}{2}\right)\right) dx \\
 & \quad \downarrow \text{25} \\
 & - \int \sec\left(x - \frac{\pi}{2}\right) \tan\left(x - \frac{\pi}{2}\right) dx \\
 & \quad \downarrow \text{3086} \\
 & - \int 1 d \csc(x) \\
 & \quad \downarrow \text{24} \\
 & - \csc(x)
 \end{aligned}$$

input `Int [Cot [x] *Csc [x] , x]`

output `-Csc [x]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3086 `Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a/f Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
derivativdivides	$-\csc(x)$	5
default	$-\csc(x)$	5
risch	$-\frac{2ie^{ix}}{e^{2ix}-1}$	18

input `int(cot(x)*csc(x), x, method=_RETURNVERBOSE)`

output `-csc(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `integrate(cot(x)*csc(x),x, algorithm="fricas")`

output `-1/sin(x)`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `integrate(cot(x)*csc(x),x)`

output `-1/sin(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `integrate(cot(x)*csc(x),x, algorithm="maxima")`

output `-1/sin(x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `integrate(cot(x)*csc(x),x, algorithm="giac")`

output `-1/sin(x)`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.50

$$\int \cot(x) \csc(x) dx = -\frac{1}{\sin(x)}$$

input `int(cot(x)/sin(x),x)`

output `-1/sin(x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \cot(x) \csc(x) dx = -\csc(x)$$

input `int(cot(x)*csc(x),x)`

output `- csc(x)`

3.11 $\int \tan(x) dx$

Optimal result	134
Mathematica [A] (verified)	134
Rubi [A] (verified)	135
Maple [A] (verified)	136
Fricas [B] (verification not implemented)	136
Sympy [A] (verification not implemented)	137
Maxima [A] (verification not implemented)	137
Giac [A] (verification not implemented)	137
Mupad [B] (verification not implemented)	138
Reduce [B] (verification not implemented)	138

Optimal result

Integrand size = 2, antiderivative size = 5

$$\int \tan(x) dx = -\log(\cos(x))$$

output `-ln(cos(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \tan(x) dx = -\log(\cos(x))$$

input `Integrate[Tan[x], x]`

output `-Log[Cos[x]]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \tan(x) dx \\ \downarrow 3042 \\ \int \tan(x) dx \\ \downarrow 3956 \\ -\log(\cos(x)) \end{array}$$

input `Int[Tan[x], x]`

output `-Log[Cos[x]]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
lookup	$-\ln(\cos(x))$	6
default	$-\ln(\cos(x))$	6
derivativdivides	$\frac{\ln(1+\tan(x)^2)}{2}$	10
norman	$\frac{\ln(1+\tan(x)^2)}{2}$	10
parallelrisc	$\frac{\ln(1+\tan(x)^2)}{2}$	10
risc	$ix - \ln(e^{2ix} + 1)$	16

input `int(tan(x),x,method=_RETURNVERBOSE)`

output `-ln(cos(x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(5) = 10$.

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 2.20

$$\int \tan(x) dx = -\frac{1}{2} \log\left(\frac{1}{\tan(x)^2 + 1}\right)$$

input `integrate(tan(x),x, algorithm="fricas")`

output `-1/2*log(1/(tan(x)^2 + 1))`

Sympy [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \tan(x) dx = -\log(\cos(x))$$

input `integrate(tan(x),x)`

output `-log(cos(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.60

$$\int \tan(x) dx = \log(\sec(x))$$

input `integrate(tan(x),x, algorithm="maxima")`

output `log(sec(x))`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

$$\int \tan(x) dx = -\log(|\cos(x)|)$$

input `integrate(tan(x),x, algorithm="giac")`

output `-log(abs(cos(x)))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \tan(x) dx = -\ln(\cos(x))$$

input `int(tan(x),x)`

output `-log(cos(x))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.80

$$\int \tan(x) dx = \frac{\log(\tan(x)^2 + 1)}{2}$$

input `int(tan(x),x)`

output `log(tan(x)**2 + 1)/2`

3.12 $\int \cot(x) dx$

Optimal result	139
Mathematica [A] (verified)	139
Rubi [A] (verified)	140
Maple [A] (verified)	141
Fricas [B] (verification not implemented)	141
Sympy [A] (verification not implemented)	142
Maxima [A] (verification not implemented)	142
Giac [A] (verification not implemented)	142
Mupad [B] (verification not implemented)	143
Reduce [B] (verification not implemented)	143

Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \cot(x) dx = \log(\sin(x))$$

output `ln(sin(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `Integrate[Cot[x], x]`

output `Log[Sin[x]]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 25, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \cot(x) dx \\ & \quad \downarrow \text{3042} \\ & \int -\tan\left(x + \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{25} \\ & -\int \tan\left(x + \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{3956} \\ & \log(\sin(x)) \end{aligned}$$

input `Int[Cot[x],x]`

output `Log[Sin[x]]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956 `Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\ln(\sin(x))$	4
default	$\ln(\sin(x))$	4
derivativedivides	$-\frac{\ln(1+\cot(x)^2)}{2}$	10
parallelrisc	$\ln\left(\frac{1}{\sqrt{\sec(x)^2}}\right) + \ln(\tan(x))$	12
norman	$-\frac{\ln(1+\tan(x)^2)}{2} + \ln(\tan(x))$	14
risc	$-ix + \ln(e^{2ix} - 1)$	14

input `int(cot(x),x,method=_RETURNVERBOSE)`

output `ln(sin(x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(3) = 6$.

Time = 0.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 3.67

$$\int \cot(x) dx = \frac{1}{2} \log\left(-\frac{1}{2} \cos(2x) + \frac{1}{2}\right)$$

input `integrate(cot(x),x, algorithm="fricas")`

output `1/2*log(-1/2*cos(2*x) + 1/2)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `integrate(cot(x),x)`

output `log(sin(x))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \log(\sin(x))$$

input `integrate(cot(x),x, algorithm="maxima")`

output `log(sin(x))`

Giac [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

$$\int \cot(x) dx = \log(|\sin(x)|)$$

input `integrate(cot(x),x, algorithm="giac")`

output `log(abs(sin(x)))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \cot(x) dx = \ln(\sin(x))$$

input `int(cot(x),x)`

output `log(sin(x))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 17, normalized size of antiderivative = 5.67

$$\int \cot(x) dx = -\log\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + \log\left(\tan\left(\frac{x}{2}\right)\right)$$

input `int(cot(x),x)`

output `- log(tan(x/2)**2 + 1) + log(tan(x/2))`

3.13 $\int \csc(x) dx$

Optimal result	144
Mathematica [A] (verified)	144
Rubi [A] (verified)	145
Maple [A] (verified)	146
Fricas [B] (verification not implemented)	146
Sympy [B] (verification not implemented)	147
Maxima [A] (verification not implemented)	147
Giac [A] (verification not implemented)	147
Mupad [B] (verification not implemented)	148
Reduce [B] (verification not implemented)	148

Optimal result

Integrand size = 2, antiderivative size = 7

$$\int \csc(x) dx = \log\left(\tan\left(\frac{x}{2}\right)\right)$$

output `ln(tan(1/2*x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \csc(x) dx = -\operatorname{arctanh}(\cos(x))$$

input `Integrate[Csc[x], x]`

output `-ArcTanh[Cos[x]]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \csc(x) dx \\ \downarrow 3042 \\ \int \csc(x) dx \\ \downarrow 4257 \\ -\operatorname{arctanh}(\cos(x)) \end{array}$$

input `Int[Csc[x], x]`

output `-ArcTanh[Cos[x]]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

method	result	size
norman	$\ln\left(\tan\left(\frac{x}{2}\right)\right)$	6
parallelrisc	$\ln\left(\tan\left(\frac{x}{2}\right)\right)$	6
lookup	$-\ln(\csc(x) + \cot(x))$	9
default	$-\ln(\csc(x) + \cot(x))$	9
risch	$\ln(e^{ix} - 1) - \ln(e^{ix} + 1)$	20

input `int(csc(x),x,method=_RETURNVERBOSE)`

output `ln(tan(1/2*x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(5) = 10$.

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.71

$$\int \csc(x) dx = -\frac{1}{2} \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) + \frac{1}{2} \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right)$$

input `integrate(csc(x),x, algorithm="fricas")`

output `-1/2*log(1/2*cos(x) + 1/2) + 1/2*log(-1/2*cos(x) + 1/2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(5) = 10$.

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.14

$$\int \csc(x) dx = \frac{\log(\cos(x) - 1)}{2} - \frac{\log(\cos(x) + 1)}{2}$$

input `integrate(csc(x),x)`

output `log(cos(x) - 1)/2 - log(cos(x) + 1)/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

$$\int \csc(x) dx = -\log(\cot(x) + \csc(x))$$

input `integrate(csc(x),x, algorithm="maxima")`

output `-log(cot(x) + csc(x))`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

$$\int \csc(x) dx = \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)$$

input `integrate(csc(x),x, algorithm="giac")`

output `log(abs(tan(1/2*x)))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \csc(x) dx = \ln\left(\tan\left(\frac{x}{2}\right)\right)$$

input `int(1/sin(x),x)`

output `log(tan(x/2))`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \csc(x) dx = \log\left(\tan\left(\frac{x}{2}\right)\right)$$

input `int(csc(x),x)`

output `log(tan(x/2))`

3.14 $\int \sec(x) dx$

Optimal result	149
Mathematica [A] (verified)	149
Rubi [A] (verified)	150
Maple [A] (verified)	151
Fricas [B] (verification not implemented)	151
Sympy [B] (verification not implemented)	152
Maxima [A] (verification not implemented)	152
Giac [B] (verification not implemented)	152
Mupad [B] (verification not implemented)	153
Reduce [B] (verification not implemented)	153

Optimal result

Integrand size = 2, antiderivative size = 6

$$\int \sec(x) dx = \log(\sec(x) + \tan(x))$$

output `ln(sec(x)+tan(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50

$$\int \sec(x) dx = \operatorname{coth}^{-1}(\sin(x))$$

input `Integrate[Sec[x], x]`

output `ArcCoth[Sin[x]]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sec(x) dx \\ \downarrow 3042 \\ \int \csc\left(x + \frac{\pi}{2}\right) dx \\ \downarrow 4257 \\ \operatorname{arctanh}(\sin(x)) \end{array}$$

input `Int[Sec[x], x]`

output `ArcTanh[Sin[x]]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
lookup	$\ln(\sec(x) + \tan(x))$	7
default	$\ln(\sec(x) + \tan(x))$	7
norman	$-\ln(\tan(\frac{x}{2}) - 1) + \ln(\tan(\frac{x}{2}) + 1)$	18
parallelsch	$-\ln(\tan(\frac{x}{2}) - 1) + \ln(\tan(\frac{x}{2}) + 1)$	18
risch	$\ln(e^{ix} + i) - \ln(e^{ix} - i)$	22

input `int(sec(x),x,method=_RETURNVERBOSE)`

output `ln(sec(x)+tan(x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(6) = 12$.

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.83

$$\int \sec(x) dx = \frac{1}{2} \log(\sin(x) + 1) - \frac{1}{2} \log(-\sin(x) + 1)$$

input `integrate(sec(x),x, algorithm="fricas")`

output `1/2*log(sin(x) + 1) - 1/2*log(-sin(x) + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(7) = 14$.

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \sec(x) dx = -\frac{\log(\sin(x) - 1)}{2} + \frac{\log(\sin(x) + 1)}{2}$$

input `integrate(sec(x),x)`

output `-log(sin(x) - 1)/2 + log(sin(x) + 1)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \sec(x) dx = \log(\sec(x) + \tan(x))$$

input `integrate(sec(x),x, algorithm="maxima")`

output `log(sec(x) + tan(x))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. $2(6) = 12$.

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 4.17

$$\int \sec(x) dx = \frac{1}{4} \log\left(\left|\frac{1}{\sin(x)} + \sin(x) + 2\right|\right) - \frac{1}{4} \log\left(\left|\frac{1}{\sin(x)} + \sin(x) - 2\right|\right)$$

input `integrate(sec(x),x, algorithm="giac")`

output `1/4*log(abs(1/sin(x) + sin(x) + 2)) - 1/4*log(abs(1/sin(x) + sin(x) - 2))`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.83

$$\int \sec(x) dx = \ln\left(\frac{1}{\cos(x)}\right) + \ln(\sin(x) + 1)$$

input `int(1/cos(x),x)`

output `log(1/cos(x)) + log(sin(x) + 1)`

Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.83

$$\int \sec(x) dx = -\log\left(\tan\left(\frac{x}{2}\right) - 1\right) + \log\left(\tan\left(\frac{x}{2}\right) + 1\right)$$

input `int(sec(x),x)`

output `- log(tan(x/2) - 1) + log(tan(x/2) + 1)`

3.15 $\int \frac{1}{1+x^2} dx$

Optimal result	154
Mathematica [A] (verified)	154
Rubi [A] (verified)	155
Maple [A] (verified)	155
Fricas [A] (verification not implemented)	156
Sympy [A] (verification not implemented)	156
Maxima [A] (verification not implemented)	157
Giac [A] (verification not implemented)	157
Mupad [B] (verification not implemented)	157
Reduce [B] (verification not implemented)	158

Optimal result

Integrand size = 7, antiderivative size = 2

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

output `arctan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `Integrate[(1 + x^2)^(-1),x]`

output `ArcTan[x]`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 + 1} dx$$

↓ 216

$$\arctan(x)$$

input `Int[(1 + x^2)^(-1), x]`

output `ArcTan[x]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\arctan(x)$	3
meijerg	$\arctan(x)$	3
risch	$\arctan(x)$	3
parallelrisch	$\frac{i \ln(x+i)}{2} - \frac{i \ln(x-i)}{2}$	18

input `int(1/(x^2+1),x,method=_RETURNVERBOSE)`

output `arctan(x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `integrate(1/(x^2+1),x, algorithm="fricas")`

output `arctan(x)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \operatorname{atan}(x)$$

input `integrate(1/(x**2+1),x)`

output `atan(x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `integrate(1/(x^2+1),x, algorithm="maxima")`

output `arctan(x)`

Giac [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `integrate(1/(x^2+1),x, algorithm="giac")`

output `arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \operatorname{atan}(x)$$

input `int(1/(x^2 + 1),x)`

output `atan(x)`

Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \text{atan}(x)$$

input `int(1/(x^2+1),x)`

output `atan(x)`

3.16 $\int \frac{1}{1-x^2} dx$

Optimal result	159
Mathematica [B] (verified)	159
Rubi [A] (verified)	160
Maple [A] (verified)	161
Fricas [B] (verification not implemented)	161
Sympy [B] (verification not implemented)	162
Maxima [B] (verification not implemented)	162
Giac [B] (verification not implemented)	162
Mupad [B] (verification not implemented)	163
Reduce [B] (verification not implemented)	163

Optimal result

Integrand size = 9, antiderivative size = 2

$$\int \frac{1}{1-x^2} dx = \operatorname{arctanh}(x)$$

output `arctanh(x)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(2) = 4$.

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 9.50

$$\int \frac{1}{1-x^2} dx = -\frac{1}{2} \log(1-x) + \frac{1}{2} \log(1+x)$$

input `Integrate[(1 - x^2)^(-1), x]`

output `-1/2*Log[1 - x] + Log[1 + x]/2`

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1-x^2} dx$$

↓ 219

$$\operatorname{arctanh}(x)$$

input `Int[(1 - x^2)^(-1), x]`

output `ArcTanh[x]`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\operatorname{arctanh}(x)$	3
meijerg	$\operatorname{arctanh}(x)$	3
norman	$-\frac{\ln(x-1)}{2} + \frac{\ln(1+x)}{2}$	14
risch	$-\frac{\ln(x-1)}{2} + \frac{\ln(1+x)}{2}$	14
parallelrisc	$-\frac{\ln(x-1)}{2} + \frac{\ln(1+x)}{2}$	14

input `int(1/(-x^2+1),x,method=_RETURNVERBOSE)`

output `arctanh(x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(2) = 4$.

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(x+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^2+1),x, algorithm="fricas")`

output `1/2*log(x + 1) - 1/2*log(x - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(2) = 4$.

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 6.00

$$\int \frac{1}{1-x^2} dx = -\frac{\log(x-1)}{2} + \frac{\log(x+1)}{2}$$

input `integrate(1/(-x**2+1),x)`

output `-log(x - 1)/2 + log(x + 1)/2`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(2) = 4$.

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(x+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^2+1),x, algorithm="maxima")`

output `1/2*log(x + 1) - 1/2*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(2) = 4$.

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 7.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(|x+1|) - \frac{1}{2} \log(|x-1|)$$

input `integrate(1/(-x^2+1),x, algorithm="giac")`

output $1/2*\log(\text{abs}(x + 1)) - 1/2*\log(\text{abs}(x - 1))$

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-x^2} dx = \text{atanh}(x)$$

input `int(-1/(x^2 - 1),x)`

output `atanh(x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = -\frac{\log(x-1)}{2} + \frac{\log(x+1)}{2}$$

input `int(1/(-x^2+1),x)`

output `(- log(x - 1) + log(x + 1))/2`

3.17 $\int \frac{1}{\sqrt{1-x^2}} dx$

Optimal result	164
Mathematica [B] (verified)	164
Rubi [A] (verified)	165
Maple [A] (verified)	165
Fricas [B] (verification not implemented)	166
Sympy [A] (verification not implemented)	166
Maxima [A] (verification not implemented)	167
Giac [B] (verification not implemented)	167
Mupad [B] (verification not implemented)	167
Reduce [B] (verification not implemented)	168

Optimal result

Integrand size = 11, antiderivative size = 2

$$\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin(x)$$

output `arcsin(x)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 20 vs. 2(2) = 4.

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 10.00

$$\int \frac{1}{\sqrt{1-x^2}} dx = -2 \arctan \left(\frac{\sqrt{1-x^2}}{1+x} \right)$$

input `Integrate[1/Sqrt[1 - x^2],x]`

output `-2*ArcTan[Sqrt[1 - x^2]/(1 + x)]`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {223}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-x^2}} dx$$

↓ 223

$$\arcsin(x)$$

input `Int[1/Sqrt[1 - x^2], x]`

output `ArcSin[x]`

Defintions of rubi rules used

rule 223 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\arcsin(x)$	3
meijerg	$\arcsin(x)$	3
pseudoelliptic	$-\arctan\left(\frac{\sqrt{-x^2+1}}{x}\right)$	17
trager	$\text{RootOf}(_Z^2 + 1) \ln(\text{RootOf}(_Z^2 + 1) \sqrt{-x^2 + 1} + x)$	27

input `int(1/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsin(x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(2) = 4$.

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 9.00

$$\int \frac{1}{\sqrt{1-x^2}} dx = -2 \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right)$$

input `integrate(1/(-x^2+1)^(1/2),x, algorithm="fricas")`

output `-2*arctan((sqrt(-x^2 + 1) - 1)/x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}} dx = \text{asin}(x)$$

input `integrate(1/(-x**2+1)**(1/2),x)`

output `asin(x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin(x)$$

input `integrate(1/(-x^2+1)^(1/2),x, algorithm="maxima")`

output `arcsin(x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(2) = 4$.

Time = 0.12 (sec) , antiderivative size = 17, normalized size of antiderivative = 8.50

$$\int \frac{1}{\sqrt{1-x^2}} dx = \frac{1}{2} \sqrt{-x^2 + 1}x + \frac{1}{2} \arcsin(x)$$

input `integrate(1/(-x^2+1)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(-x^2 + 1)*x + 1/2*arcsin(x)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}} dx = \operatorname{asin}(x)$$

input `int(1/(1 - x^2)^(1/2),x)`

output `asin(x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-x^2}} dx = \text{asin}(x)$$

input `int(1/(-x^2+1)^(1/2),x)`

output `asin(x)`

3.18 $\int \frac{1}{\sqrt{1+x^2}} dx$

Optimal result	169
Mathematica [B] (verified)	169
Rubi [A] (verified)	170
Maple [A] (verified)	170
Fricas [B] (verification not implemented)	171
Sympy [A] (verification not implemented)	171
Maxima [A] (verification not implemented)	172
Giac [B] (verification not implemented)	172
Mupad [B] (verification not implemented)	172
Reduce [B] (verification not implemented)	173

Optimal result

Integrand size = 9, antiderivative size = 2

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{arcsinh}(x)$$

output `arcsinh(x)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 16 vs. 2(2) = 4.

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 8.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = -\log\left(-x + \sqrt{1+x^2}\right)$$

input `Integrate[1/Sqrt[1 + x^2],x]`

output `-Log[-x + Sqrt[1 + x^2]]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 + 1}} dx$$

↓ 222

$$\operatorname{arcsinh}(x)$$

input `Int[1/Sqrt[1 + x^2], x]`

output `ArcSinh[x]`

Defintions of rubi rules used

rule 222 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\operatorname{arcsinh}(x)$	3
meijerg	$\operatorname{arcsinh}(x)$	3
pseudoelliptic	$\operatorname{arctanh}\left(\frac{\sqrt{x^2+1}}{x}\right)$	13
trager	$-\ln(x - \sqrt{x^2 + 1})$	15

input `int(1/(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

output `arcsinh(x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(2) = 4$.

Time = 0.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 7.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = -\log(-x + \sqrt{x^2+1})$$

input `integrate(1/(x^2+1)^(1/2),x, algorithm="fricas")`

output `-log(-x + sqrt(x^2 + 1))`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{asinh}(x)$$

input `integrate(1/(x**2+1)**(1/2),x)`

output `asinh(x)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{arsinh}(x)$$

input `integrate(1/(x^2+1)^(1/2),x, algorithm="maxima")`

output `arcsinh(x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 25 vs. 2(2) = 4.

Time = 0.11 (sec) , antiderivative size = 25, normalized size of antiderivative = 12.50

$$\int \frac{1}{\sqrt{1+x^2}} dx = \frac{1}{2} \sqrt{x^2+1} x - \frac{1}{2} \log(-x + \sqrt{x^2+1})$$

input `integrate(1/(x^2+1)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(x^2 + 1)*x - 1/2*log(-x + sqrt(x^2 + 1))`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1+x^2}} dx = \operatorname{asinh}(x)$$

input `int(1/(x^2 + 1)^(1/2),x)`

output `asinh(x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 9, normalized size of antiderivative = 4.50

$$\int \frac{1}{\sqrt{1+x^2}} dx = \log(\sqrt{x^2+1} + x)$$

input `int(1/(x^2+1)^(1/2),x)`

output `log(sqrt(x**2 + 1) + x)`

3.19 $\int \frac{1}{\sqrt{-1+x^2}} dx$

Optimal result	174
Mathematica [B] (verified)	174
Rubi [A] (verified)	175
Maple [A] (verified)	176
Fricas [A] (verification not implemented)	176
Sympy [A] (verification not implemented)	176
Maxima [A] (verification not implemented)	177
Giac [B] (verification not implemented)	177
Mupad [B] (verification not implemented)	177
Reduce [B] (verification not implemented)	178

Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{1}{\sqrt{-1+x^2}} dx = \log \left(x + \sqrt{-1+x^2} \right)$$

output

```
ln(x+(x^2-1)^(1/2))
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 38 vs. 2(12) = 24.

Time = 0.00 (sec) , antiderivative size = 38, normalized size of antiderivative = 3.17

$$\int \frac{1}{\sqrt{-1+x^2}} dx = -\frac{1}{2} \log \left(1 - \frac{x}{\sqrt{-1+x^2}} \right) + \frac{1}{2} \log \left(1 + \frac{x}{\sqrt{-1+x^2}} \right)$$

input

```
Integrate[1/Sqrt[-1 + x^2],x]
```

output

```
-1/2*Log[1 - x/Sqrt[-1 + x^2]] + Log[1 + x/Sqrt[-1 + x^2]]/2
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {224, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 - 1}} dx$$

↓ 224

$$\int \frac{1}{1 - \frac{x^2}{x^2 - 1}} d \frac{x}{\sqrt{x^2 - 1}}$$

↓ 219

$$\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2 - 1}}\right)$$

input `Int[1/Sqrt[-1 + x^2], x]`

output `ArcTanh[x/Sqrt[-1 + x^2]]`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
default	$\ln(x + \sqrt{x^2 - 1})$	11
pseudoelliptic	$\operatorname{arctanh}\left(\frac{\sqrt{x^2-1}}{x}\right)$	13
trager	$-\ln(-\sqrt{x^2 - 1} + x)$	15
meijerg	$\frac{\sqrt{-\operatorname{signum}(x^2-1)} \arcsin(x)}{\sqrt{\operatorname{signum}(x^2-1)}}$	22

input `int(1/(x^2-1)^(1/2),x,method=_RETURNVERBOSE)`output `ln(x+(x^2-1)^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sqrt{-1+x^2}} dx = -\log(-x + \sqrt{x^2 - 1})$$

input `integrate(1/(x^2-1)^(1/2),x, algorithm="fricas")`output `-log(-x + sqrt(x^2 - 1))`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{-1+x^2}} dx = \log(x + \sqrt{x^2 - 1})$$

input `integrate(1/(x**2-1)**(1/2),x)`

output `log(x + sqrt(x**2 - 1))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sqrt{-1+x^2}} dx = \log\left(2x + 2\sqrt{x^2-1}\right)$$

input `integrate(1/(x^2-1)^(1/2),x, algorithm="maxima")`

output `log(2*x + 2*sqrt(x^2 - 1))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(10) = 20.

Time = 0.12 (sec) , antiderivative size = 26, normalized size of antiderivative = 2.17

$$\int \frac{1}{\sqrt{-1+x^2}} dx = \frac{1}{2} \sqrt{x^2-1}x + \frac{1}{2} \log\left(\left|-x + \sqrt{x^2-1}\right|\right)$$

input `integrate(1/(x^2-1)^(1/2),x, algorithm="giac")`

output `1/2*sqrt(x^2 - 1)*x + 1/2*log(abs(-x + sqrt(x^2 - 1)))`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{-1+x^2}} dx = \ln\left(x + \sqrt{x^2-1}\right)$$

input `int(1/(x^2 - 1)^(1/2),x)`

output `log(x + (x^2 - 1)^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{1}{\sqrt{-1 + x^2}} dx = \log(\sqrt{x^2 - 1} + x)$$

input `int(1/(x^2-1)^(1/2),x)`

output `log(sqrt(x**2 - 1) + x)`

3.20 $\int \sinh(x) dx$

Optimal result	179
Mathematica [A] (verified)	179
Rubi [A] (verified)	180
Maple [A] (verified)	181
Fricas [A] (verification not implemented)	181
Sympy [A] (verification not implemented)	182
Maxima [A] (verification not implemented)	182
Giac [B] (verification not implemented)	182
Mupad [B] (verification not implemented)	183
Reduce [B] (verification not implemented)	183

Optimal result

Integrand size = 2, antiderivative size = 2

$$\int \sinh(x) dx = \cosh(x)$$

output `cosh(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `Integrate[Sinh[x], x]`

output `Cosh[x]`

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 26, 3118}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \sinh(x) dx \\ \downarrow 3042 \\ \int -i \sin(ix) dx \\ \downarrow 26 \\ -i \int \sin(ix) dx \\ \downarrow 3118 \\ \cosh(x) \end{array}$$

input `Int[Sinh[x],x]`

output `Cosh[x]`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3118 `Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
lookup	$\cosh(x)$	3
default	$\cosh(x)$	3
orering	$\cosh(x)$	3
parallelrisc	$1 + \cosh(x)$	5
risc	$\frac{e^x}{2} + \frac{e^{-x}}{2}$	12
meijerg	$-\sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cosh(x)}{\sqrt{\pi}} \right)$	17

input `int(sinh(x),x,method=_RETURNVERBOSE)`

output `cosh(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `integrate(sinh(x),x, algorithm="fricas")`

output `cosh(x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `integrate(sinh(x),x)`

output `cosh(x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `integrate(sinh(x),x, algorithm="maxima")`

output `cosh(x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(2) = 4$.

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 5.50

$$\int \sinh(x) dx = \frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(sinh(x),x, algorithm="giac")`

output `1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `int(sinh(x),x)`

output `cosh(x)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \sinh(x) dx = \cosh(x)$$

input `int(sinh(x),x)`

output `cosh(x)`

3.21 $\int \cosh(x) dx$

Optimal result	184
Mathematica [A] (verified)	184
Rubi [A] (verified)	185
Maple [A] (verified)	186
Fricas [A] (verification not implemented)	186
Sympy [A] (verification not implemented)	187
Maxima [A] (verification not implemented)	187
Giac [B] (verification not implemented)	187
Mupad [B] (verification not implemented)	188
Reduce [B] (verification not implemented)	188

Optimal result

Integrand size = 2, antiderivative size = 2

$$\int \cosh(x) dx = \sinh(x)$$

output `sinh(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `Integrate[Cosh[x], x]`

output `Sinh[x]`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 3117}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \cosh(x) dx \\ \downarrow 3042 \\ \int \sin\left(\frac{\pi}{2} + ix\right) dx \\ \downarrow 3117 \\ \sinh(x) \end{array}$$

input `Int[Cosh[x], x]`

output `Sinh[x]`

Defintions of rubi rules used

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3117 `Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
lookup	$\sinh(x)$	3
default	$\sinh(x)$	3
meijerg	$\sinh(x)$	3
parallelrisch	$\sinh(x)$	3
orering	$\sinh(x)$	3
risch	$\frac{e^x}{2} - \frac{e^{-x}}{2}$	12

input `int(cosh(x),x,method=_RETURNVERBOSE)`

output `sinh(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `integrate(cosh(x),x, algorithm="fricas")`

output `sinh(x)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `integrate(cosh(x),x)`

output `sinh(x)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `integrate(cosh(x),x, algorithm="maxima")`

output `sinh(x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(2) = 4$.

Time = 0.11 (sec) , antiderivative size = 11, normalized size of antiderivative = 5.50

$$\int \cosh(x) dx = -\frac{1}{2} e^{(-x)} + \frac{1}{2} e^x$$

input `integrate(cosh(x),x, algorithm="giac")`

output `-1/2*e^(-x) + 1/2*e^x`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `int(cosh(x),x)`

output `sinh(x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \cosh(x) dx = \sinh(x)$$

input `int(cosh(x),x)`

output `sinh(x)`

3.22 $\int \operatorname{csch}^2(x) dx$

Optimal result	189
Mathematica [A] (verified)	189
Rubi [A] (verified)	190
Maple [A] (verified)	191
Fricas [B] (verification not implemented)	192
Sympy [F]	192
Maxima [B] (verification not implemented)	192
Giac [B] (verification not implemented)	193
Mupad [B] (verification not implemented)	193
Reduce [B] (verification not implemented)	193

Optimal result

Integrand size = 4, antiderivative size = 4

$$\int \operatorname{csch}^2(x) dx = -\operatorname{coth}(x)$$

output `-coth(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}^2(x) dx = -\operatorname{coth}(x)$$

input `Integrate[Csch[x]^2,x]`

output `-Coth[x]`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3042, 25, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \operatorname{csch}^2(x) dx \\
 & \quad \downarrow \text{3042} \\
 & \int -\operatorname{csc}(ix)^2 dx \\
 & \quad \downarrow \text{25} \\
 & -\int \operatorname{csc}(ix)^2 dx \\
 & \quad \downarrow \text{4254} \\
 & -i \int 1d(-i \operatorname{coth}(x)) \\
 & \quad \downarrow \text{24} \\
 & -\operatorname{coth}(x)
 \end{aligned}$$

input `Int [Csch [x]^2, x]`

output `-Coth [x]`

Defintions of rubi rules used

rule 24 `Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 3042 `Int[u_, x_Symbol] := Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 4254 `Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$-\coth(x)$	5
risch	$-\frac{2}{e^{2x}-1}$	11
parallelrisch	$-\frac{\coth(\frac{x}{2})}{2} - \frac{\tanh(\frac{x}{2})}{2}$	14

input `int(csch(x)^2,x,method=_RETURNVERBOSE)`

output `-coth(x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(4) = 8$.

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 5.00

$$\int \operatorname{csch}^2(x) dx = -\frac{2}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1}$$

input `integrate(csch(x)^2,x, algorithm="fricas")`

output `-2/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)`

Sympy [F]

$$\int \operatorname{csch}^2(x) dx = \int \operatorname{csch}^2(x) dx$$

input `integrate(csch(x)**2,x)`

output `Integral(csch(x)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(4) = 8$.

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 2.50

$$\int \operatorname{csch}^2(x) dx = \frac{2}{e^{(-2x)} - 1}$$

input `integrate(csch(x)^2,x, algorithm="maxima")`

output `2/(e^(-2*x) - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(4) = 8$.

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 2.50

$$\int \operatorname{csch}^2(x) dx = -\frac{2}{e^{2x} - 1}$$

input `integrate(csch(x)^2,x, algorithm="giac")`

output `-2/(e^(2*x) - 1)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \operatorname{csch}^2(x) dx = -\operatorname{coth}(x)$$

input `int(1/sinh(x)^2,x)`

output `-coth(x)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 4.00

$$\int \operatorname{csch}^2(x) dx = -\frac{2e^{2x}}{e^{2x} - 1}$$

input `int(csch(x)^2,x)`

output `(- 2*e**(2*x))/(e**(2*x) - 1)`

3.23 $\int \operatorname{sech}^2(x) dx$

Optimal result	194
Mathematica [A] (verified)	194
Rubi [A] (verified)	195
Maple [A] (verified)	196
Fricas [B] (verification not implemented)	196
Sympy [F]	197
Maxima [B] (verification not implemented)	197
Giac [B] (verification not implemented)	197
Mupad [B] (verification not implemented)	198
Reduce [B] (verification not implemented)	198

Optimal result

Integrand size = 4, antiderivative size = 2

$$\int \operatorname{sech}^2(x) dx = \tanh(x)$$

output `tanh(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}^2(x) dx = \tanh(x)$$

input `Integrate[Sech[x]^2,x]`

output `Tanh[x]`

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3042, 4254, 24}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \operatorname{sech}^2(x) dx \\ \downarrow 3042 \\ \int \csc\left(\frac{\pi}{2} + ix\right)^2 dx \\ \downarrow 4254 \\ i \int 1d(-i \tanh(x)) \\ \downarrow 24 \\ \tanh(x) \end{array}$$

input `Int [Sech [x]^2, x]`

output `Tanh [x]`

Defintions of rubi rules used

rule 24 `Int [a_, x_Symbol] :> Simp [a*x, x] /; FreeQ [a, x]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinear Q [u, x]`

rule 4254

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Simp[-d^(-1) Subst[Int[Exp
andIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\tanh(x)$	3
parallelrisc	$\frac{\sinh(x)}{\cosh(x)}$	8
risc	$-\frac{2}{e^{2x}+1}$	11

input

```
int(sech(x)^2,x,method=_RETURNVERBOSE)
```

output

```
tanh(x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(2) = 4$.

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 10.00

$$\int \operatorname{sech}^2(x) dx = -\frac{2}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1}$$

input

```
integrate(sech(x)^2,x, algorithm="fricas")
```

output

```
-2/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1)
```

Sympy [F]

$$\int \operatorname{sech}^2(x) dx = \int \operatorname{sech}^2(x) dx$$

input `integrate(sech(x)**2,x)`

output `Integral(sech(x)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(2) = 4$.

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 5.00

$$\int \operatorname{sech}^2(x) dx = \frac{2}{e^{(-2x)} + 1}$$

input `integrate(sech(x)^2,x, algorithm="maxima")`

output `2/(e^(-2*x) + 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(2) = 4$.

Time = 0.11 (sec) , antiderivative size = 10, normalized size of antiderivative = 5.00

$$\int \operatorname{sech}^2(x) dx = -\frac{2}{e^{(2x)} + 1}$$

input `integrate(sech(x)^2,x, algorithm="giac")`

output `-2/(e^(2*x) + 1)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \operatorname{sech}^2(x) dx = \tanh(x)$$

input `int(1/cosh(x)^2,x)`

output `tanh(x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 8.00

$$\int \operatorname{sech}^2(x) dx = \frac{2e^{2x}}{e^{2x} + 1}$$

input `int(sech(x)^2,x)`

output `(2*e**(2*x))/(e**(2*x) + 1)`

3.24 $\int \tanh(x) dx$

Optimal result	199
Mathematica [A] (verified)	199
Rubi [A] (verified)	200
Maple [A] (verified)	201
Fricas [B] (verification not implemented)	201
Sympy [B] (verification not implemented)	202
Maxima [A] (verification not implemented)	202
Giac [B] (verification not implemented)	202
Mupad [B] (verification not implemented)	203
Reduce [B] (verification not implemented)	203

Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \tanh(x) dx = \log(\cosh(x))$$

output `ln(cosh(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \tanh(x) dx = \log(\cosh(x))$$

input `Integrate[Tanh[x], x]`

output `Log[Cosh[x]]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \tanh(x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \tan(ix) dx \\ & \quad \downarrow \text{26} \\ & -i \int \tan(ix) dx \\ & \quad \downarrow \text{3956} \\ & \log(\cosh(x)) \end{aligned}$$

input `Int [Tanh [x] , x]`

output `Log [Cosh [x]]`

Defintions of rubi rules used

rule 26 `Int [(Complex [0, a_])*(Fx_), x_Symbol] :> Simp [(Complex [Identity [0], a]) Int [Fx, x], x] /; FreeQ [a, x] && EqQ [a^2, 1]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinear Q [u, x]`

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\ln(\cosh(x))$	4
derivativedivides	$\ln(\cosh(x))$	4
default	$\ln(\cosh(x))$	4
risch	$-x + \ln(e^{2x} + 1)$	12
parallelrisc	$-\ln(1 - \tanh(x)) - x$	14

input

```
int(tanh(x),x,method=_RETURNVERBOSE)
```

output

```
ln(cosh(x))
```

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(3) = 6$.

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 6.00

$$\int \tanh(x) dx = -x + \log\left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)}\right)$$

input

```
integrate(tanh(x),x, algorithm="fricas")
```

output

```
-x + log(2*cosh(x)/(cosh(x) - sinh(x)))
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 7 vs. $2(3) = 6$.

Time = 0.06 (sec) , antiderivative size = 7, normalized size of antiderivative = 2.33

$$\int \tanh(x) dx = x - \log(\tanh(x) + 1)$$

input `integrate(tanh(x), x)`

output `x - log(tanh(x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \tanh(x) dx = \log(\cosh(x))$$

input `integrate(tanh(x), x, algorithm="maxima")`

output `log(cosh(x))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11 vs. $2(3) = 6$.

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 3.67

$$\int \tanh(x) dx = -x + \log(e^{2x} + 1)$$

input `integrate(tanh(x), x, algorithm="giac")`

output `-x + log(e^(2*x) + 1)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \tanh(x) dx = \ln(\cosh(x))$$

input `int(tanh(x), x)`

output `log(cosh(x))`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 4.00

$$\int \tanh(x) dx = \log(e^{2x} + 1) - x$$

input `int(tanh(x), x)`

output `log(e**(2*x) + 1) - x`

3.25 $\int \coth(x) dx$

Optimal result	204
Mathematica [A] (verified)	204
Rubi [A] (verified)	205
Maple [A] (verified)	206
Fricas [B] (verification not implemented)	206
Sympy [B] (verification not implemented)	207
Maxima [A] (verification not implemented)	207
Giac [B] (verification not implemented)	207
Mupad [B] (verification not implemented)	208
Reduce [B] (verification not implemented)	208

Optimal result

Integrand size = 2, antiderivative size = 3

$$\int \coth(x) dx = \log(\sinh(x))$$

output `ln(sinh(x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \coth(x) dx = \log(\sinh(x))$$

input `Integrate[Coth[x], x]`

output `Log[Sinh[x]]`

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 26, 3956}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \coth(x) dx \\ & \quad \downarrow \text{3042} \\ & \int -i \tan\left(\frac{\pi}{2} + ix\right) dx \\ & \quad \downarrow \text{26} \\ & -i \int \tan\left(ix + \frac{\pi}{2}\right) dx \\ & \quad \downarrow \text{3956} \\ & \log(\sinh(x)) \end{aligned}$$

input `Int[Coth[x],x]`

output `Log[Sinh[x]]`

Defintions of rubi rules used

rule 26 `Int[(Complex[0, a_])*(Fx_), x_Symbol] :> Simp[(Complex[Identity[0], a]) Int[Fx, x], x] /; FreeQ[a, x] && EqQ[a^2, 1]`

rule 3042 `Int[u_, x_Symbol] :> Int[DeactivateTrig[u, x], x] /; FunctionOfTrigOfLinearQ[u, x]`

rule 3956

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.33

method	result	size
lookup	$\ln(\sinh(x))$	4
derivativedivides	$\ln(\sinh(x))$	4
default	$\ln(\sinh(x))$	4
risch	$-x + \ln(e^{2x} - 1)$	12
parallelrisch	$\ln(\tanh(x)) - \ln(1 - \tanh(x)) - x$	17

input

```
int(coth(x),x,method=_RETURNVERBOSE)
```

output

```
ln(sinh(x))
```

Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 18 vs. $2(3) = 6$.

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 6.00

$$\int \coth(x) dx = -x + \log\left(\frac{2 \sinh(x)}{\cosh(x) - \sinh(x)}\right)$$

input

```
integrate(coth(x),x, algorithm="fricas")
```

output

```
-x + log(2*sinh(x)/(cosh(x) - sinh(x)))
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(3) = 6$.

Time = 0.16 (sec) , antiderivative size = 12, normalized size of antiderivative = 4.00

$$\int \coth(x) dx = x - \log(\tanh(x) + 1) + \log(\tanh(x))$$

input `integrate(coth(x),x)`

output `x - log(tanh(x) + 1) + log(tanh(x))`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \coth(x) dx = \log(\sinh(x))$$

input `integrate(coth(x),x, algorithm="maxima")`

output `log(sinh(x))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(3) = 6$.

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 4.00

$$\int \coth(x) dx = -x + \log(|e^{(2x)} - 1|)$$

input `integrate(coth(x),x, algorithm="giac")`

output `-x + log(abs(e^(2*x) - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.00

$$\int \coth(x) dx = \ln(\sinh(x))$$

input `int(coth(x),x)`

output `log(sinh(x))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 5.33

$$\int \coth(x) dx = \log(e^x - 1) + \log(e^x + 1) - x$$

input `int(coth(x),x)`

output `log(e**x - 1) + log(e**x + 1) - x`

3.26 $\int \operatorname{csch}(x) dx$

Optimal result	209
Mathematica [A] (verified)	209
Rubi [A] (verified)	210
Maple [A] (verified)	211
Fricas [B] (verification not implemented)	211
Sympy [A] (verification not implemented)	212
Maxima [A] (verification not implemented)	212
Giac [B] (verification not implemented)	212
Mupad [B] (verification not implemented)	213
Reduce [B] (verification not implemented)	213

Optimal result

Integrand size = 2, antiderivative size = 7

$$\int \operatorname{csch}(x) dx = \log\left(\tanh\left(\frac{x}{2}\right)\right)$$

output `ln(tanh(1/2*x))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \operatorname{csch}(x) dx = -\operatorname{arctanh}(\operatorname{cosh}(x))$$

input `Integrate[Csch[x], x]`

output `-ArcTanh[Cosh[x]]`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 1.500$, Rules used = {3042, 26, 4257}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c} \int \operatorname{csch}(x) dx \\ \downarrow 3042 \\ \int i \operatorname{csc}(ix) dx \\ \downarrow 26 \\ i \int \operatorname{csc}(ix) dx \\ \downarrow 4257 \\ -\operatorname{arctanh}(\cosh(x)) \end{array}$$

input `Int [Csch [x] , x]`

output `-ArcTanh [Cosh [x]]`

Defintions of rubi rules used

rule 26 `Int [(Complex [0, a_])*(Fx_), x_Symbol] :> Simp [(Complex [Identity [0], a]) Int [Fx, x], x] /; FreeQ [a, x] && EqQ [a^2, 1]`

rule 3042 `Int [u_, x_Symbol] :> Int [DeactivateTrig [u, x], x] /; FunctionOfTrigOfLinear Q [u, x]`

rule 4257 `Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.86

method	result	size
lookup	$\ln\left(\tanh\left(\frac{x}{2}\right)\right)$	6
default	$\ln\left(\tanh\left(\frac{x}{2}\right)\right)$	6
parallelrisch	$\ln\left(\tanh\left(\frac{x}{2}\right)\right)$	6
risch	$\ln(-1 + e^x) - \ln(e^x + 1)$	14

input `int(csch(x),x,method=_RETURNVERBOSE)`

output `ln(tanh(1/2*x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(5) = 10$.

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.43

$$\int \operatorname{csch}(x) dx = -\log(\cosh(x) + \sinh(x) + 1) + \log(\cosh(x) + \sinh(x) - 1)$$

input `integrate(csch(x),x, algorithm="fricas")`

output `-log(cosh(x) + sinh(x) + 1) + log(cosh(x) + sinh(x) - 1)`

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \operatorname{csch}(x) dx = \log\left(\tanh\left(\frac{x}{2}\right)\right)$$

input `integrate(csch(x), x)`

output `log(tanh(x/2))`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \operatorname{csch}(x) dx = \log\left(\tanh\left(\frac{1}{2} x\right)\right)$$

input `integrate(csch(x), x, algorithm="maxima")`

output `log(tanh(1/2*x))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(5) = 10$.

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 2.00

$$\int \operatorname{csch}(x) dx = -\log(e^x + 1) + \log(|e^x - 1|)$$

input `integrate(csch(x), x, algorithm="giac")`

output `-log(e^x + 1) + log(abs(e^x - 1))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.71

$$\int \operatorname{csch}(x) dx = \ln \left(\tanh \left(\frac{x}{2} \right) \right)$$

input `int(1/sinh(x),x)`

output `log(tanh(x/2))`

Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.14

$$\int \operatorname{csch}(x) dx = \log(e^x - 1) - \log(e^x + 1)$$

input `int(csch(x),x)`

output `log(e**x - 1) - log(e**x + 1)`

3.27 $\int (a + bx)^m dx$

Optimal result	214
Mathematica [A] (verified)	214
Rubi [A] (verified)	215
Maple [A] (verified)	216
Fricas [A] (verification not implemented)	216
Sympy [A] (verification not implemented)	217
Maxima [A] (verification not implemented)	217
Giac [A] (verification not implemented)	217
Mupad [B] (verification not implemented)	218
Reduce [B] (verification not implemented)	218

Optimal result

Integrand size = 7, antiderivative size = 18

$$\int (a + bx)^m dx = \frac{(a + bx)^{1+m}}{b(1+m)}$$

output $(b*x+a)^{(1+m)}/b/(1+m)$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (a + bx)^m dx = \frac{(a + bx)^{1+m}}{b(1+m)}$$

input `Integrate[(a + b*x)^m,x]`

output $(a + b*x)^{(1 + m)}/(b*(1 + m))$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (a + bx)^m dx$$

$$\downarrow 17$$

$$\frac{(a + bx)^{m+1}}{b(m + 1)}$$

input `Int[(a + b*x)^m,x]`

output `(a + b*x)^(1 + m)/(b*(1 + m))`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

method	result	size
gospers	$\frac{(bx+a)^{1+m}}{b(1+m)}$	19
default	$\frac{(bx+a)^{1+m}}{b(1+m)}$	19
risch	$\frac{(bx+a)(bx+a)^m}{b(1+m)}$	22
orering	$\frac{(bx+a)(bx+a)^m}{b(1+m)}$	22
parallelrisc	$\frac{x(bx+a)^m ab + (bx+a)^m a^2}{(1+m)ab}$	36
norman	$\frac{x e^{m \ln(bx+a)}}{1+m} + \frac{a e^{m \ln(bx+a)}}{b(1+m)}$	37

input `int((b*x+a)^m,x,method=_RETURNVERBOSE)`output `(b*x+a)^(1+m)/b/(1+m)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (a + bx)^m dx = \frac{(bx + a)(bx + a)^m}{bm + b}$$

input `integrate((b*x+a)^m,x, algorithm="fricas")`output `(b*x + a)*(b*x + a)^m/(b*m + b)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (a + bx)^m dx = \frac{\begin{cases} \frac{(a+bx)^{m+1}}{m+1} & \text{for } m \neq -1 \\ \log(a + bx) & \text{otherwise} \end{cases}}{b}$$

input `integrate((b*x+a)**m,x)`output `Piecewise(((a + b*x)**(m + 1)/(m + 1), Ne(m, -1)), (log(a + b*x), True))/b`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (a + bx)^m dx = \frac{(bx + a)^{m+1}}{b(m + 1)}$$

input `integrate((b*x+a)^m,x, algorithm="maxima")`output `(b*x + a)^(m + 1)/(b*(m + 1))`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (a + bx)^m dx = \frac{(bx + a)^{m+1}}{b(m + 1)}$$

input `integrate((b*x+a)^m,x, algorithm="giac")`output `(b*x + a)^(m + 1)/(b*(m + 1))`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (a + bx)^m dx = \frac{(a + bx)^{m+1}}{b(m+1)}$$

input `int((a + b*x)^m,x)`

output `(a + b*x)^(m + 1)/(b*(m + 1))`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.17

$$\int (a + bx)^m dx = \frac{(bx + a)^m (bx + a)}{b(m+1)}$$

input `int((b*x+a)^m,x)`

output `((a + b*x)**m*(a + b*x))/(b*(m + 1))`

3.28 $\int \frac{1}{a+bx} dx$

Optimal result	219
Mathematica [A] (verified)	219
Rubi [A] (verified)	220
Maple [A] (verified)	220
Fricas [A] (verification not implemented)	221
Sympy [A] (verification not implemented)	221
Maxima [A] (verification not implemented)	222
Giac [A] (verification not implemented)	222
Mupad [B] (verification not implemented)	222
Reduce [B] (verification not implemented)	223

Optimal result

Integrand size = 7, antiderivative size = 10

$$\int \frac{1}{a+bx} dx = \frac{\log(a+bx)}{b}$$

output `ln(b*x+a)/b`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a+bx} dx = \frac{\log(a+bx)}{b}$$

input `Integrate[(a + b*x)^(-1),x]`

output `Log[a + b*x]/b`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a + bx} dx$$

↓ 16

$$\frac{\log(a + bx)}{b}$$

input `Int[(a + b*x)^(-1),x]`

output `Log[a + b*x]/b`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
default	$\frac{\ln(bx+a)}{b}$	11
norman	$\frac{\ln(bx+a)}{b}$	11
risch	$\frac{\ln(bx+a)}{b}$	11
parallelrisk	$\frac{\ln(bx+a)}{b}$	11

input `int(1/(b*x+a),x,method=_RETURNVERBOSE)`

output `ln(b*x+a)/b`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a+bx} dx = \frac{\log(bx+a)}{b}$$

input `integrate(1/(b*x+a),x, algorithm="fricas")`

output `log(b*x + a)/b`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{1}{a+bx} dx = \frac{\log(a+bx)}{b}$$

input `integrate(1/(b*x+a),x)`

output `log(a + b*x)/b`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + bx} dx = \frac{\log(bx + a)}{b}$$

input `integrate(1/(b*x+a),x, algorithm="maxima")`

output `log(b*x + a)/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{1}{a + bx} dx = \frac{\log(|bx + a|)}{b}$$

input `integrate(1/(b*x+a),x, algorithm="giac")`

output `log(abs(b*x + a))/b`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + bx} dx = \frac{\ln(a + bx)}{b}$$

input `int(1/(a + b*x),x)`

output `log(a + b*x)/b`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + bx} dx = \frac{\log(bx + a)}{b}$$

input `int(1/(b*x+a),x)`

output `log(a + b*x)/b`

3.29 $\int \frac{x}{a+bx} dx$

Optimal result	224
Mathematica [A] (verified)	224
Rubi [A] (verified)	225
Maple [A] (verified)	226
Fricas [A] (verification not implemented)	226
Sympy [A] (verification not implemented)	226
Maxima [A] (verification not implemented)	227
Giac [A] (verification not implemented)	227
Mupad [B] (verification not implemented)	227
Reduce [B] (verification not implemented)	228

Optimal result

Integrand size = 9, antiderivative size = 18

$$\int \frac{x}{a+bx} dx = \frac{x}{b} - \frac{a \log(a+bx)}{b^2}$$

output `x/b-a*ln(b*x+a)/b^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x}{a+bx} dx = \frac{x}{b} - \frac{a \log(a+bx)}{b^2}$$

input `Integrate[x/(a + b*x), x]`

output `x/b - (a*Log[a + b*x])/b^2`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{a + bx} dx$$

$$\downarrow 49$$

$$\int \left(\frac{1}{b} - \frac{a}{b(a + bx)} \right) dx$$

$$\downarrow 2009$$

$$\frac{x}{b} - \frac{a \log(a + bx)}{b^2}$$

input

```
Int[x/(a + b*x), x]
```

output

```
x/b - (a*Log[a + b*x])/b^2
```

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

method	result	size
default	$\frac{x}{b} - \frac{a \ln(bx+a)}{b^2}$	19
norman	$\frac{x}{b} - \frac{a \ln(bx+a)}{b^2}$	19
risch	$\frac{x}{b} - \frac{a \ln(bx+a)}{b^2}$	19
parallelrisch	$-\frac{a \ln(bx+a) - bx}{b^2}$	19

input `int(x/(b*x+a),x,method=_RETURNVERBOSE)`output `x/b-a*ln(b*x+a)/b^2`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{x}{a+bx} dx = \frac{bx - a \log(bx+a)}{b^2}$$

input `integrate(x/(b*x+a),x, algorithm="fricas")`output `(b*x - a*log(b*x + a))/b^2`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \frac{x}{a+bx} dx = -\frac{a \log(a+bx)}{b^2} + \frac{x}{b}$$

input `integrate(x/(b*x+a),x)`

output `-a*log(a + b*x)/b**2 + x/b`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x}{a + bx} dx = \frac{x}{b} - \frac{a \log(bx + a)}{b^2}$$

input `integrate(x/(b*x+a),x, algorithm="maxima")`

output `x/b - a*log(b*x + a)/b^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

$$\int \frac{x}{a + bx} dx = \frac{x}{b} - \frac{a \log(|bx + a|)}{b^2}$$

input `integrate(x/(b*x+a),x, algorithm="giac")`

output `x/b - a*log(abs(b*x + a))/b^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{x}{a + bx} dx = -\frac{a \ln(a + bx) - bx}{b^2}$$

input `int(x/(a + b*x),x)`

output `-(a*log(a + b*x) - b*x)/b^2`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{x}{a + bx} dx = \frac{-\log(bx + a) a + bx}{b^2}$$

input `int(x/(b*x+a),x)`

output `(- log(a + b*x)*a + b*x)/b**2`

3.30 $\int \frac{x^2}{a+bx} dx$

Optimal result	229
Mathematica [A] (verified)	229
Rubi [A] (verified)	230
Maple [A] (verified)	231
Fricas [A] (verification not implemented)	231
Sympy [A] (verification not implemented)	231
Maxima [A] (verification not implemented)	232
Giac [A] (verification not implemented)	232
Mupad [B] (verification not implemented)	232
Reduce [B] (verification not implemented)	233

Optimal result

Integrand size = 11, antiderivative size = 31

$$\int \frac{x^2}{a+bx} dx = -\frac{ax}{b^2} + \frac{x^2}{2b} + \frac{a^2 \log(a+bx)}{b^2}$$

output

```
-a*x/b^2+1/2*x^2/b+a^2*ln(b*x+a)/b^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{a+bx} dx = -\frac{ax}{b^2} + \frac{x^2}{2b} + \frac{a^2 \log(a+bx)}{b^3}$$

input

```
Integrate[x^2/(a + b*x),x]
```

output

```
-((a*x)/b^2) + x^2/(2*b) + (a^2*Log[a + b*x])/b^3
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{a + bx} dx$$

↓ 49

$$\int \left(\frac{a^2}{b^2(a + bx)} - \frac{a}{b^2} + \frac{x}{b} \right) dx$$

↓ 2009

$$\frac{a^2 \log(a + bx)}{b^3} - \frac{ax}{b^2} + \frac{x^2}{2b}$$

input `Int[x^2/(a + b*x),x]`

output `-((a*x)/b^2) + x^2/(2*b) + (a^2*Log[a + b*x])/b^3`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

method	result	size
default	$-\frac{\frac{1}{2}x^2b+ax}{b^2} + \frac{a^2 \ln(bx+a)}{b^3}$	30
norman	$\frac{x^2}{2b} - \frac{ax}{b^2} + \frac{a^2 \ln(bx+a)}{b^3}$	30
risch	$\frac{x^2}{2b} - \frac{ax}{b^2} + \frac{a^2 \ln(bx+a)}{b^3}$	30
parallelrisch	$\frac{b^2x^2+2a^2 \ln(bx+a)-2bax}{2b^3}$	30

input `int(x^2/(b*x+a),x,method=_RETURNVERBOSE)`output `-1/b^2*(-1/2*x^2*b+a*x)+a^2/b^3*ln(b*x+a)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{a+bx} dx = \frac{b^2x^2 - 2abx + 2a^2 \log(bx+a)}{2b^3}$$

input `integrate(x^2/(b*x+a),x, algorithm="fricas")`output `1/2*(b^2*x^2 - 2*a*b*x + 2*a^2*log(b*x + a))/b^3`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int \frac{x^2}{a+bx} dx = \frac{a^2 \log(a+bx)}{b^3} - \frac{ax}{b^2} + \frac{x^2}{2b}$$

input `integrate(x**2/(b*x+a),x)`

output `a**2*log(a + b*x)/b**3 - a*x/b**2 + x**2/(2*b)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{a + bx} dx = \frac{a^2 \log (bx + a)}{b^3} + \frac{bx^2 - 2ax}{2b^2}$$

input `integrate(x^2/(b*x+a),x, algorithm="maxima")`

output `a^2*log(b*x + a)/b^3 + 1/2*(b*x^2 - 2*a*x)/b^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

$$\int \frac{x^2}{a + bx} dx = \frac{a^2 \log (|bx + a|)}{b^3} + \frac{bx^2 - 2ax}{2b^2}$$

input `integrate(x^2/(b*x+a),x, algorithm="giac")`

output `a^2*log(abs(b*x + a))/b^3 + 1/2*(b*x^2 - 2*a*x)/b^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{a + bx} dx = \frac{2a^2 \ln (a + bx) + b^2 x^2 - 2abx}{2b^3}$$

input `int(x^2/(a + b*x),x)`

output `(2*a^2*log(a + b*x) + b^2*x^2 - 2*a*b*x)/(2*b^3)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{a+bx} dx = \frac{2 \log(bx+a) a^2 - 2abx + b^2 x^2}{2b^3}$$

input `int(x^2/(b*x+a),x)`

output `(2*log(a + b*x)*a**2 - 2*a*b*x + b**2*x**2)/(2*b**3)`

3.31 $\int \frac{1}{(a+bx)^2} dx$

Optimal result	234
Mathematica [A] (verified)	234
Rubi [A] (verified)	235
Maple [A] (verified)	236
Fricas [A] (verification not implemented)	236
Sympy [A] (verification not implemented)	237
Maxima [A] (verification not implemented)	237
Giac [A] (verification not implemented)	237
Mupad [B] (verification not implemented)	238
Reduce [B] (verification not implemented)	238

Optimal result

Integrand size = 7, antiderivative size = 12

$$\int \frac{1}{(a+bx)^2} dx = -\frac{1}{b(a+bx)}$$

output `-1/b/(b*x+a)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+bx)^2} dx = -\frac{1}{b(a+bx)}$$

input `Integrate[(a + b*x)^(-2),x]`

output `-(1/(b*(a + b*x)))`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx)^2} dx$$

$$\downarrow 17$$

$$-\frac{1}{b(a + bx)}$$

input `Int[(a + b*x)^(-2),x]`

output `-(1/(b*(a + b*x)))`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result	size
gospers	$-\frac{1}{b(bx+a)}$	13
default	$-\frac{1}{b(bx+a)}$	13
norman	$\frac{x}{a(bx+a)}$	13
risch	$-\frac{1}{b(bx+a)}$	13
parallelrisch	$-\frac{1}{b(bx+a)}$	13
orering	$-\frac{1}{b(bx+a)}$	13

input `int(1/(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `-1/b/(b*x+a)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{1}{(a+bx)^2} dx = -\frac{1}{b^2x+ab}$$

input `integrate(1/(b*x+a)^2,x, algorithm="fricas")`

output `-1/(b^2*x + a*b)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{(a+bx)^2} dx = -\frac{1}{ab+b^2x}$$

input `integrate(1/(b*x+a)**2,x)`

output `-1/(a*b + b**2*x)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+bx)^2} dx = -\frac{1}{(bx+a)b}$$

input `integrate(1/(b*x+a)^2,x, algorithm="maxima")`

output `-1/((b*x + a)*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+bx)^2} dx = -\frac{1}{(bx+a)b}$$

input `integrate(1/(b*x+a)^2,x, algorithm="giac")`

output `-1/((b*x + a)*b)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + bx)^2} dx = -\frac{1}{b(a + bx)}$$

input `int(1/(a + b*x)^2,x)`

output `-1/(b*(a + b*x))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + bx)^2} dx = \frac{x}{a(bx + a)}$$

input `int(1/(b*x+a)^2,x)`

output `x/(a*(a + b*x))`

3.32 $\int \frac{x}{(a+bx)^2} dx$

Optimal result	239
Mathematica [A] (verified)	239
Rubi [A] (verified)	240
Maple [A] (verified)	241
Fricas [A] (verification not implemented)	241
Sympy [A] (verification not implemented)	241
Maxima [A] (verification not implemented)	242
Giac [A] (verification not implemented)	242
Mupad [B] (verification not implemented)	242
Reduce [B] (verification not implemented)	243

Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \frac{x}{(a+bx)^2} dx = -\frac{x}{b(a+bx)} + \frac{\log(a+bx)}{b^2}$$

output

```
-x/b/(b*x+a)+ln(b*x+a)/b^2
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{x}{(a+bx)^2} dx = \frac{\frac{a}{a+bx} + \log(a+bx)}{b^2}$$

input

```
Integrate[x/(a + b*x)^2,x]
```

output

```
(a/(a + b*x) + Log[a + b*x])/b^2
```


Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a+bx)^2} dx$$

$$\downarrow 49$$

$$\int \left(\frac{1}{b(a+bx)} - \frac{a}{b(a+bx)^2} \right) dx$$

$$\downarrow 2009$$

$$\frac{a}{b^2(a+bx)} + \frac{\log(a+bx)}{b^2}$$

input `Int[x/(a + b*x)^2,x]`

output `a/(b^2*(a + b*x)) + Log[a + b*x]/b^2`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{a}{b^2(bx+a)} + \frac{\ln(bx+a)}{b^2}$	24
norman	$\frac{a}{b^2(bx+a)} + \frac{\ln(bx+a)}{b^2}$	24
risch	$\frac{a}{b^2(bx+a)} + \frac{\ln(bx+a)}{b^2}$	24
parallelrisc	$\frac{\ln(bx+a)xb+a \ln(bx+a)+a}{b^2(bx+a)}$	31

input `int(x/(b*x+a)^2,x,method=_RETURNVERBOSE)`output `a/b^2/(b*x+a)+ln(b*x+a)/b^2`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{x}{(a+bx)^2} dx = \frac{(bx+a) \log(bx+a) + a}{b^3x + ab^2}$$

input `integrate(x/(b*x+a)^2,x, algorithm="fricas")`output `((b*x + a)*log(b*x + a) + a)/(b^3*x + a*b^2)`**Sympy [A] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{x}{(a+bx)^2} dx = \frac{a}{ab^2 + b^3x} + \frac{\log(a+bx)}{b^2}$$

input `integrate(x/(b*x+a)**2,x)`

output $a/(a*b**2 + b**3*x) + \log(a + b*x)/b**2$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{x}{(a+bx)^2} dx = \frac{a}{b^3x + ab^2} + \frac{\log(bx+a)}{b^2}$$

input `integrate(x/(b*x+a)^2,x, algorithm="maxima")`

output $a/(b^3*x + a*b^2) + \log(b*x + a)/b^2$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.75

$$\int \frac{x}{(a+bx)^2} dx = -\frac{\log\left(\frac{|bx+a|}{(bx+a)^2|b|}\right)}{b} - \frac{a}{(bx+a)b}$$

input `integrate(x/(b*x+a)^2,x, algorithm="giac")`

output $-(\log(\text{abs}(b*x + a)/((b*x + a)^2*\text{abs}(b))))/b - a/((b*x + a)*b)/b$

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.96

$$\int \frac{x}{(a+bx)^2} dx = \frac{\ln(a+bx)}{b^2} + \frac{a}{b^2(a+bx)}$$

input `int(x/(a + b*x)^2,x)`

output `log(a + b*x)/b^2 + a/(b^2*(a + b*x))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.38

$$\int \frac{x}{(a + bx)^2} dx = \frac{\log(bx + a) a + \log(bx + a) bx - bx}{b^2 (bx + a)}$$

input `int(x/(b*x+a)^2,x)`

output `(log(a + b*x)*a + log(a + b*x)*b*x - b*x)/(b**2*(a + b*x))`

3.33 $\int \frac{x^2}{(a+bx)^2} dx$

Optimal result	244
Mathematica [A] (verified)	244
Rubi [A] (verified)	245
Maple [A] (verified)	246
Fricas [A] (verification not implemented)	246
Sympy [A] (verification not implemented)	247
Maxima [A] (verification not implemented)	247
Giac [A] (verification not implemented)	247
Mupad [B] (verification not implemented)	248
Reduce [B] (verification not implemented)	248

Optimal result

Integrand size = 11, antiderivative size = 33

$$\int \frac{x^2}{(a+bx)^2} dx = \frac{x}{b^2} - \frac{a^2}{b^3(a+bx)} - \frac{2a \log(a+bx)}{b^3}$$

output `x/b^2-a^2/b^3/(b*x+a)-2*a*ln(b*x+a)/b^3`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

$$\int \frac{x^2}{(a+bx)^2} dx = \frac{bx - \frac{a^2}{a+bx} - 2a \log(a+bx)}{b^3}$$

input `Integrate[x^2/(a + b*x)^2,x]`

output `(b*x - a^2/(a + b*x) - 2*a*Log[a + b*x])/b^3`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(a+bx)^2} dx$$

$$\downarrow 49$$

$$\int \left(\frac{a^2}{b^2(a+bx)^2} - \frac{2a}{b^2(a+bx)} + \frac{1}{b^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{a^2}{b^3(a+bx)} - \frac{2a \log(a+bx)}{b^3} + \frac{x}{b^2}$$

input `Int[x^2/(a + b*x)^2,x]`

output `x/b^2 - a^2/(b^3*(a + b*x)) - (2*a*Log[a + b*x])/b^3`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.03

method	result	size
default	$\frac{x}{b^2} - \frac{a^2}{b^3(bx+a)} - \frac{2a \ln(bx+a)}{b^3}$	34
risch	$\frac{x}{b^2} - \frac{a^2}{b^3(bx+a)} - \frac{2a \ln(bx+a)}{b^3}$	34
norman	$\frac{\frac{x^2}{b} - \frac{2a^2}{b^3}}{bx+a} - \frac{2a \ln(bx+a)}{b^3}$	38
parallelrisch	$-\frac{2 \ln(bx+a)xab - b^2x^2 + 2a^2 \ln(bx+a) + 2a^2}{b^3(bx+a)}$	49

input `int(x^2/(b*x+a)^2,x,method=_RETURNVERBOSE)`output `x/b^2-a^2/b^3/(b*x+a)-2*a*ln(b*x+a)/b^3`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.42

$$\int \frac{x^2}{(a+bx)^2} dx = \frac{b^2x^2 + abx - a^2 - 2(abx + a^2) \log(bx + a)}{b^4x + ab^3}$$

input `integrate(x^2/(b*x+a)^2,x, algorithm="fricas")`output `(b^2*x^2 + a*b*x - a^2 - 2*(a*b*x + a^2)*log(b*x + a))/(b^4*x + a*b^3)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{(a+bx)^2} dx = -\frac{a^2}{ab^3 + b^4x} - \frac{2a \log(a+bx)}{b^3} + \frac{x}{b^2}$$

input `integrate(x**2/(b*x+a)**2,x)`output `-a**2/(a*b**3 + b**4*x) - 2*a*log(a + b*x)/b**3 + x/b**2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.09

$$\int \frac{x^2}{(a+bx)^2} dx = -\frac{a^2}{b^4x + ab^3} + \frac{x}{b^2} - \frac{2a \log(bx+a)}{b^3}$$

input `integrate(x^2/(b*x+a)^2,x, algorithm="maxima")`output `-a^2/(b^4*x + a*b^3) + x/b^2 - 2*a*log(b*x + a)/b^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.52

$$\int \frac{x^2}{(a+bx)^2} dx = \frac{2a \log\left(\frac{|bx+a|}{(bx+a)^2|b|}\right)}{b^3} + \frac{bx+a}{b^3} - \frac{a^2}{(bx+a)b^3}$$

input `integrate(x^2/(b*x+a)^2,x, algorithm="giac")`output `2*a*log(abs(b*x + a)/((b*x + a)^2*abs(b)))/b^3 + (b*x + a)/b^3 - a^2/((b*x + a)*b^3)`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.09

$$\int \frac{x^2}{(a+bx)^2} dx = \frac{x}{b^2} - \frac{a^2}{x b^4 + a b^3} - \frac{2 a \ln(a+bx)}{b^3}$$

input `int(x^2/(a + b*x)^2,x)`output `x/b^2 - a^2/(a*b^3 + b^4*x) - (2*a*log(a + b*x))/b^3`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

$$\int \frac{x^2}{(a+bx)^2} dx = \frac{-2 \log(bx+a) a^2 - 2 \log(bx+a) abx + 2abx + b^2 x^2}{b^3 (bx+a)}$$

input `int(x^2/(b*x+a)^2,x)`output `(- 2*log(a + b*x)*a**2 - 2*log(a + b*x)*a*b*x + 2*a*b*x + b**2*x**2)/(b**3*(a + b*x))`

3.34 $\int \frac{1}{(a+bx)^3} dx$

Optimal result	249
Mathematica [A] (verified)	249
Rubi [A] (verified)	250
Maple [A] (verified)	251
Fricas [A] (verification not implemented)	251
Sympy [B] (verification not implemented)	252
Maxima [A] (verification not implemented)	252
Giac [A] (verification not implemented)	252
Mupad [B] (verification not implemented)	253
Reduce [B] (verification not implemented)	253

Optimal result

Integrand size = 7, antiderivative size = 14

$$\int \frac{1}{(a+bx)^3} dx = -\frac{1}{2b(a+bx)^2}$$

output

```
-1/2/b/(b*x+a)^2
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+bx)^3} dx = -\frac{1}{2b(a+bx)^2}$$

input

```
Integrate[(a + b*x)^(-3),x]
```

output

```
-1/2*1/(b*(a + b*x)^2)
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a+bx)^3} dx$$

↓ 17

$$-\frac{1}{2b(a+bx)^2}$$

input `Int[(a + b*x)^(-3),x]`

output `-1/2*1/(b*(a + b*x)^2)`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] :> Simp[c*((a + b*x)^(m + 1))/(b*(m + 1))], x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
gospers	$-\frac{1}{2b(bx+a)^2}$	13
default	$-\frac{1}{2b(bx+a)^2}$	13
norman	$-\frac{1}{2b(bx+a)^2}$	13
risch	$-\frac{1}{2b(bx+a)^2}$	13
parallelrisch	$-\frac{1}{2b(bx+a)^2}$	13
orering	$-\frac{1}{2b(bx+a)^2}$	13

input `int(1/(b*x+a)^3,x,method=_RETURNVERBOSE)`output `-1/2/b/(b*x+a)^2`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.71

$$\int \frac{1}{(a+bx)^3} dx = -\frac{1}{2(b^3x^2 + 2ab^2x + a^2b)}$$

input `integrate(1/(b*x+a)^3,x, algorithm="fricas")`output `-1/2/(b^3*x^2 + 2*a*b^2*x + a^2*b)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 26 vs. $2(12) = 24$.

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.86

$$\int \frac{1}{(a + bx)^3} dx = -\frac{1}{2a^2b + 4ab^2x + 2b^3x^2}$$

input `integrate(1/(b*x+a)**3,x)`

output `-1/(2*a**2*b + 4*a*b**2*x + 2*b**3*x**2)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a + bx)^3} dx = -\frac{1}{2(bx + a)^2b}$$

input `integrate(1/(b*x+a)^3,x, algorithm="maxima")`

output `-1/2/((b*x + a)^2*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a + bx)^3} dx = -\frac{1}{2(bx + a)^2b}$$

input `integrate(1/(b*x+a)^3,x, algorithm="giac")`

output `-1/2/((b*x + a)^2*b)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.86

$$\int \frac{1}{(a + bx)^3} dx = -\frac{1}{2a^2b + 4ab^2x + 2b^3x^2}$$

input `int(1/(a + b*x)^3,x)`

output `-1/(2*a^2*b + 2*b^3*x^2 + 4*a*b^2*x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.64

$$\int \frac{1}{(a + bx)^3} dx = -\frac{1}{2b(b^2x^2 + 2abx + a^2)}$$

input `int(1/(b*x+a)^3,x)`

output `(- 1)/(2*b*(a**2 + 2*a*b*x + b**2*x**2))`

3.35 $\int \frac{x}{(a+bx)^3} dx$

Optimal result	254
Mathematica [A] (verified)	254
Rubi [A] (verified)	255
Maple [A] (verified)	256
Fricas [A] (verification not implemented)	256
Sympy [A] (verification not implemented)	257
Maxima [A] (verification not implemented)	257
Giac [A] (verification not implemented)	257
Mupad [B] (verification not implemented)	258
Reduce [B] (verification not implemented)	258

Optimal result

Integrand size = 9, antiderivative size = 23

$$\int \frac{x}{(a+bx)^3} dx = \frac{-\frac{a}{2b^2} - \frac{x}{b}}{(a+bx)^2}$$

output $(-1/2*a/b^2-x/b)/(b*x+a)^2$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \frac{x}{(a+bx)^3} dx = -\frac{a+2bx}{2b^2(a+bx)^2}$$

input `Integrate[x/(a + b*x)^3,x]`

output $-1/2*(a + 2*b*x)/(b^2*(a + b*x)^2)$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a+bx)^3} dx$$

$$\downarrow 48$$

$$\frac{x^2}{2a(a+bx)^2}$$

input `Int[x/(a + b*x)^3,x]`

output `x^2/(2*a*(a + b*x)^2)`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp
 [(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
 a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

method	result	size
gospers	$-\frac{2bx+a}{2b^2(bx+a)^2}$	19
orering	$-\frac{2bx+a}{2b^2(bx+a)^2}$	19
parallelrisch	$\frac{-2bx-a}{2b^2(bx+a)^2}$	21
norman	$\frac{-\frac{a}{2b^2} - \frac{x}{b}}{(bx+a)^2}$	22
risch	$\frac{-\frac{a}{2b^2} - \frac{x}{b}}{(bx+a)^2}$	22
default	$-\frac{1}{b^2(bx+a)} + \frac{a}{2b^2(bx+a)^2}$	27

input `int(x/(b*x+a)^3,x,method=_RETURNVERBOSE)`output `-1/2*(2*b*x+a)/b^2/(b*x+a)^2`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.39

$$\int \frac{x}{(a+bx)^3} dx = -\frac{2bx+a}{2(b^4x^2+2ab^3x+a^2b^2)}$$

input `integrate(x/(b*x+a)^3,x, algorithm="fricas")`output `-1/2*(2*b*x + a)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.39

$$\int \frac{x}{(a+bx)^3} dx = \frac{-a-2bx}{2a^2b^2+4ab^3x+2b^4x^2}$$

input `integrate(x/(b*x+a)**3,x)`output `(-a - 2*b*x)/(2*a**2*b**2 + 4*a*b**3*x + 2*b**4*x**2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.39

$$\int \frac{x}{(a+bx)^3} dx = -\frac{2bx+a}{2(b^4x^2+2ab^3x+a^2b^2)}$$

input `integrate(x/(b*x+a)^3,x, algorithm="maxima")`output `-1/2*(2*b*x + a)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{x}{(a+bx)^3} dx = -\frac{2bx+a}{2(bx+a)^2b^2}$$

input `integrate(x/(b*x+a)^3,x, algorithm="giac")`output `-1/2*(2*b*x + a)/((b*x + a)^2*b^2)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.39

$$\int \frac{x}{(a+bx)^3} dx = -\frac{\frac{a}{2b^2} + \frac{x}{b}}{a^2 + 2abx + b^2x^2}$$

input `int(x/(a + b*x)^3,x)`output `-(a/(2*b^2) + x/b)/(a^2 + b^2*x^2 + 2*a*b*x)`**Reduce [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int \frac{x}{(a+bx)^3} dx = \frac{x^2}{2a(b^2x^2 + 2abx + a^2)}$$

input `int(x/(b*x+a)^3,x)`output `x**2/(2*a*(a**2 + 2*a*b*x + b**2*x**2))`

3.36 $\int \frac{x^2}{(a+bx)^3} dx$

Optimal result	259
Mathematica [A] (verified)	259
Rubi [A] (verified)	260
Maple [A] (verified)	261
Fricas [A] (verification not implemented)	261
Sympy [A] (verification not implemented)	262
Maxima [A] (verification not implemented)	262
Giac [A] (verification not implemented)	262
Mupad [B] (verification not implemented)	263
Reduce [B] (verification not implemented)	263

Optimal result

Integrand size = 11, antiderivative size = 37

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{\frac{3a^2}{2b^3} + \frac{2ax}{b^2}}{(a+bx)^2} + \frac{\log(a+bx)}{b^3}$$

output

```
(3/2*a^2/b^3+2*a*x/b^2)/(b*x+a)^2+ln(b*x+a)/b^3
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{\frac{a(3a+4bx)}{(a+bx)^2} + 2 \log(a+bx)}{2b^3}$$

input

```
Integrate[x^2/(a + b*x)^3,x]
```

output

```
((a*(3*a + 4*b*x))/(a + b*x)^2 + 2*Log[a + b*x])/(2*b^3)
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(a+bx)^3} dx$$

$$\downarrow 49$$

$$\int \left(\frac{a^2}{b^2(a+bx)^3} - \frac{2a}{b^2(a+bx)^2} + \frac{1}{b^2(a+bx)} \right) dx$$

$$\downarrow 2009$$

$$-\frac{a^2}{2b^3(a+bx)^2} + \frac{2a}{b^3(a+bx)} + \frac{\log(a+bx)}{b^3}$$

input `Int[x^2/(a + b*x)^3,x]`

output `-1/2*a^2/(b^3*(a + b*x)^2) + (2*a)/(b^3*(a + b*x)) + Log[a + b*x]/b^3`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.97

method	result	size
norman	$\frac{3a^2 + 2ax}{2b^3 + b^2} + \frac{\ln(bx+a)}{b^3}$	36
risch	$\frac{3a^2 + 2ax}{2b^3 + b^2} + \frac{\ln(bx+a)}{b^3}$	36
default	$\frac{2a}{b^3(bx+a)} - \frac{a^2}{2b^3(bx+a)^2} + \frac{\ln(bx+a)}{b^3}$	40
parallelrisch	$\frac{2 \ln(bx+a)x^2b^2 + 4 \ln(bx+a)xab + 2a^2 \ln(bx+a) + 4bax + 3a^2}{2b^3(bx+a)^2}$	60

input `int(x^2/(b*x+a)^3,x,method=_RETURNVERBOSE)`output `(3/2*a^2/b^3+2*a*x/b^2)/(b*x+a)^2+ln(b*x+a)/b^3`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.65

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{4abx + 3a^2 + 2(b^2x^2 + 2abx + a^2) \log(bx+a)}{2(b^5x^2 + 2ab^4x + a^2b^3)}$$

input `integrate(x^2/(b*x+a)^3,x, algorithm="fricas")`output `1/2*(4*a*b*x + 3*a^2 + 2*(b^2*x^2 + 2*a*b*x + a^2)*log(b*x + a))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.24

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{3a^2 + 4abx}{2a^2b^3 + 4ab^4x + 2b^5x^2} + \frac{\log(a+bx)}{b^3}$$

input `integrate(x**2/(b*x+a)**3,x)`output `(3*a**2 + 4*a*b*x)/(2*a**2*b**3 + 4*a*b**4*x + 2*b**5*x**2) + log(a + b*x)/b**3`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.30

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{4abx + 3a^2}{2(b^5x^2 + 2ab^4x + a^2b^3)} + \frac{\log(bx+a)}{b^3}$$

input `integrate(x^2/(b*x+a)^3,x, algorithm="maxima")`output `1/2*(4*a*b*x + 3*a^2)/(b^5*x^2 + 2*a*b^4*x + a^2*b^3) + log(b*x + a)/b^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{\log(|bx+a|)}{b^3} + \frac{4ax + \frac{3a^2}{b}}{2(bx+a)^2b^2}$$

input `integrate(x^2/(b*x+a)^3,x, algorithm="giac")`output `log(abs(b*x + a))/b^3 + 1/2*(4*a*x + 3*a^2/b)/((b*x + a)^2*b^2)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.24

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{\ln(a+bx)}{b^3} + \frac{\frac{3a^2}{2b^3} + \frac{2ax}{b^2}}{a^2 + 2abx + b^2x^2}$$

input `int(x^2/(a + b*x)^3,x)`output `log(a + b*x)/b^3 + ((3*a^2)/(2*b^3) + (2*a*x)/b^2)/(a^2 + b^2*x^2 + 2*a*b*x)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.92

$$\int \frac{x^2}{(a+bx)^3} dx = \frac{2 \log(bx+a) a^2 + 4 \log(bx+a) abx + 2 \log(bx+a) b^2 x^2 + a^2 - 2b^2 x^2}{2b^3 (b^2 x^2 + 2abx + a^2)}$$

input `int(x^2/(b*x+a)^3,x)`output `(2*log(a + b*x)*a**2 + 4*log(a + b*x)*a*b*x + 2*log(a + b*x)*b**2*x**2 + a**2 - 2*b**2*x**2)/(2*b**3*(a**2 + 2*a*b*x + b**2*x**2))`

3.37 $\int \frac{x^3}{(a+bx)^3} dx$

Optimal result	264
Mathematica [A] (verified)	264
Rubi [A] (verified)	265
Maple [A] (verified)	266
Fricas [A] (verification not implemented)	266
Sympy [A] (verification not implemented)	267
Maxima [A] (verification not implemented)	267
Giac [A] (verification not implemented)	267
Mupad [B] (verification not implemented)	268
Reduce [B] (verification not implemented)	268

Optimal result

Integrand size = 11, antiderivative size = 57

$$\int \frac{x^3}{(a+bx)^3} dx = \frac{-\frac{5a^3}{2b^4} - \frac{2a^2x}{b^3} + \frac{2ax^2}{b^2} + \frac{x^3}{b}}{(a+bx)^2} - \frac{3a \log(a+bx)}{b^4}$$

output

$$\frac{(-5/2*a^3/b^4 - 2*a^2*x/b^3 + 2*a*x^2/b^2 + x^3/b)/(b*x+a)^2 - 3*a*\ln(b*x+a)/b^4}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.70

$$\int \frac{x^3}{(a+bx)^3} dx = -\frac{-2bx + \frac{a^2(5a+6bx)}{(a+bx)^2} + 6a \log(a+bx)}{2b^4}$$

input

`Integrate[x^3/(a + b*x)^3,x]`

output

$$-1/2*(-2*b*x + (a^2*(5*a + 6*b*x))/(a + b*x)^2 + 6*a*\text{Log}[a + b*x])/b^4$$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.88, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(a+bx)^3} dx$$

$$\downarrow 49$$

$$\int \left(-\frac{a^3}{b^3(a+bx)^3} + \frac{3a^2}{b^3(a+bx)^2} - \frac{3a}{b^3(a+bx)} + \frac{1}{b^3} \right) dx$$

$$\downarrow 2009$$

$$\frac{a^3}{2b^4(a+bx)^2} - \frac{3a^2}{b^4(a+bx)} - \frac{3a \log(a+bx)}{b^4} + \frac{x}{b^3}$$

input `Int[x^3/(a + b*x)^3,x]`

output `x/b^3 + a^3/(2*b^4*(a + b*x)^2) - (3*a^2)/(b^4*(a + b*x)) - (3*a*Log[a + b*x])/b^4`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{x}{b^3} + \frac{-3a^2x - \frac{5a^3}{2b}}{b^3(bx+a)^2} - \frac{3a \ln(bx+a)}{b^4}$	45
norman	$\frac{\frac{x^3}{b} - \frac{9a^3}{2b^4} - \frac{6a^2x}{b^3}}{(bx+a)^2} - \frac{3a \ln(bx+a)}{b^4}$	47
default	$\frac{x}{b^3} - \frac{3a^2}{b^4(bx+a)} + \frac{a^3}{2b^4(bx+a)^2} - \frac{3a \ln(bx+a)}{b^4}$	49
parallelrisch	$-\frac{6 \ln(bx+a)x^2 a b^2 - 2b^3 x^3 + 12 \ln(bx+a)x a^2 b + 6 \ln(bx+a)a^3 + 12a^2 bx + 9a^3}{2b^4(bx+a)^2}$	73

input `int(x^3/(b*x+a)^3,x,method=_RETURNVERBOSE)`output `x/b^3+(-3*a^2*x-5/2*a^3/b)/b^3/(b*x+a)^2-3*a*ln(b*x+a)/b^4`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.46

$$\int \frac{x^3}{(a+bx)^3} dx = \frac{2b^3x^3 + 4ab^2x^2 - 4a^2bx - 5a^3 - 6(ab^2x^2 + 2a^2bx + a^3) \log(bx+a)}{2(b^6x^2 + 2ab^5x + a^2b^4)}$$

input `integrate(x^3/(b*x+a)^3,x, algorithm="fricas")`output `1/2*(2*b^3*x^3 + 4*a*b^2*x^2 - 4*a^2*b*x - 5*a^3 - 6*(a*b^2*x^2 + 2*a^2*b*x + a^3)*log(b*x + a))/(b^6*x^2 + 2*a*b^5*x + a^2*b^4)`

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.02

$$\int \frac{x^3}{(a+bx)^3} dx = -\frac{3a \log(a+bx)}{b^4} + \frac{-5a^3 - 6a^2bx}{2a^2b^4 + 4ab^5x + 2b^6x^2} + \frac{x}{b^3}$$

input `integrate(x**3/(b*x+a)**3,x)`output `-3*a*log(a + b*x)/b**4 + (-5*a**3 - 6*a**2*b*x)/(2*a**2*b**4 + 4*a*b**5*x + 2*b**6*x**2) + x/b**3`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{x^3}{(a+bx)^3} dx = -\frac{6a^2bx + 5a^3}{2(b^6x^2 + 2ab^5x + a^2b^4)} + \frac{x}{b^3} - \frac{3a \log(bx+a)}{b^4}$$

input `integrate(x^3/(b*x+a)^3,x, algorithm="maxima")`output `-1/2*(6*a^2*b*x + 5*a^3)/(b^6*x^2 + 2*a*b^5*x + a^2*b^4) + x/b^3 - 3*a*log(b*x + a)/b^4`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.77

$$\int \frac{x^3}{(a+bx)^3} dx = \frac{x}{b^3} - \frac{3a \log(|bx+a|)}{b^4} - \frac{6a^2bx + 5a^3}{2(bx+a)^2b^4}$$

input `integrate(x^3/(b*x+a)^3,x, algorithm="giac")`output `x/b^3 - 3*a*log(abs(b*x + a))/b^4 - 1/2*(6*a^2*b*x + 5*a^3)/((b*x + a)^2*b^4)`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.75

$$\int \frac{x^3}{(a+bx)^3} dx = -\frac{3a \ln(a+bx) - bx + \frac{3a^2}{a+bx} - \frac{a^3}{2(a+bx)^2}}{b^4}$$

input `int(x^3/(a + b*x)^3,x)`output `-(3*a*log(a + b*x) - b*x + (3*a^2)/(a + b*x) - a^3/(2*(a + b*x)^2))/b^4`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.49

$$\int \frac{x^3}{(a+bx)^3} dx = \frac{-6 \log(bx+a) a^3 - 12 \log(bx+a) a^2 bx - 6 \log(bx+a) a b^2 x^2 - 3a^3 + 6a b^2 x^2 + 2b^3 x^3}{2b^4 (b^2 x^2 + 2abx + a^2)}$$

input `int(x^3/(b*x+a)^3,x)`output `(- 6*log(a + b*x)*a**3 - 12*log(a + b*x)*a**2*b*x - 6*log(a + b*x)*a*b**2*x**2 - 3*a**3 + 6*a*b**2*x**2 + 2*b**3*x**3)/(2*b**4*(a**2 + 2*a*b*x + b**2*x**2))`

3.38 $\int \frac{1}{(a+bx)^4} dx$

Optimal result	269
Mathematica [A] (verified)	269
Rubi [A] (verified)	270
Maple [A] (verified)	271
Fricas [B] (verification not implemented)	271
Sympy [B] (verification not implemented)	272
Maxima [A] (verification not implemented)	272
Giac [A] (verification not implemented)	272
Mupad [B] (verification not implemented)	273
Reduce [B] (verification not implemented)	273

Optimal result

Integrand size = 7, antiderivative size = 14

$$\int \frac{1}{(a+bx)^4} dx = -\frac{1}{3b(a+bx)^3}$$

output

```
-1/3/b/(b*x+a)^3
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+bx)^4} dx = -\frac{1}{3b(a+bx)^3}$$

input

```
Integrate[(a + b*x)^(-4),x]
```

output

```
-1/3*1/(b*(a + b*x)^3)
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx)^4} dx$$

↓ 17

$$-\frac{1}{3b(a + bx)^3}$$

input `Int[(a + b*x)^(-4),x]`

output `-1/3*1/(b*(a + b*x)^3)`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
gospers	$-\frac{1}{3b(bx+a)^3}$	13
default	$-\frac{1}{3b(bx+a)^3}$	13
norman	$-\frac{1}{3b(bx+a)^3}$	13
risch	$-\frac{1}{3b(bx+a)^3}$	13
parallelrisch	$-\frac{1}{3b(bx+a)^3}$	13
orering	$-\frac{1}{3b(bx+a)^3}$	13

input `int(1/(b*x+a)^4,x,method=_RETURNVERBOSE)`

output `-1/3/b/(b*x+a)^3`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. 2(12) = 24.

Time = 0.08 (sec) , antiderivative size = 35, normalized size of antiderivative = 2.50

$$\int \frac{1}{(a+bx)^4} dx = -\frac{1}{3(b^4x^3 + 3ab^3x^2 + 3a^2b^2x + a^3b)}$$

input `integrate(1/(b*x+a)^4,x, algorithm="fricas")`

output `-1/3/(b^4*x^3 + 3*a*b^3*x^2 + 3*a^2*b^2*x + a^3*b)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 37 vs. $2(12) = 24$.

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.64

$$\int \frac{1}{(a+bx)^4} dx = -\frac{1}{3a^3b + 9a^2b^2x + 9ab^3x^2 + 3b^4x^3}$$

input `integrate(1/(b*x+a)**4,x)`

output `-1/(3*a**3*b + 9*a**2*b**2*x + 9*a*b**3*x**2 + 3*b**4*x**3)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a+bx)^4} dx = -\frac{1}{3(bx+a)^3b}$$

input `integrate(1/(b*x+a)^4,x, algorithm="maxima")`

output `-1/3/((b*x + a)^3*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a+bx)^4} dx = -\frac{1}{3(bx+a)^3b}$$

input `integrate(1/(b*x+a)^4,x, algorithm="giac")`

output `-1/3/((b*x + a)^3*b)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.64

$$\int \frac{1}{(a + bx)^4} dx = -\frac{1}{3a^3b + 9a^2b^2x + 9ab^3x^2 + 3b^4x^3}$$

input `int(1/(a + b*x)^4,x)`output `-1/(3*a^3*b + 3*b^4*x^3 + 9*a^2*b^2*x + 9*a*b^3*x^2)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.43

$$\int \frac{1}{(a + bx)^4} dx = -\frac{1}{3b(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3)}$$

input `int(1/(b*x+a)^4,x)`output `(- 1)/(3*b*(a**3 + 3*a**2*b*x + 3*a*b**2*x**2 + b**3*x**3))`

3.39 $\int \frac{x}{(a+bx)^4} dx$

Optimal result	274
Mathematica [A] (verified)	274
Rubi [A] (verified)	275
Maple [A] (verified)	276
Fricas [B] (verification not implemented)	276
Sympy [B] (verification not implemented)	277
Maxima [B] (verification not implemented)	277
Giac [A] (verification not implemented)	277
Mupad [B] (verification not implemented)	278
Reduce [B] (verification not implemented)	278

Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \frac{x}{(a+bx)^4} dx = \frac{-\frac{a}{6b^2} - \frac{x}{2b}}{(a+bx)^3}$$

output $(-1/6*a/b^2-1/2*x/b)/(b*x+a)^3$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{x}{(a+bx)^4} dx = -\frac{a+3bx}{6b^2(a+bx)^3}$$

input `Integrate[x/(a + b*x)^4,x]`

output $-1/6*(a + 3*b*x)/(b^2*(a + b*x)^3)$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.20, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a+bx)^4} dx$$

↓ 53

$$\int \left(\frac{1}{b(a+bx)^3} - \frac{a}{b(a+bx)^4} \right) dx$$

↓ 2009

$$\frac{a}{3b^2(a+bx)^3} - \frac{1}{2b^2(a+bx)^2}$$

input `Int[x/(a + b*x)^4,x]`

output `a/(3*b^2*(a + b*x)^3) - 1/(2*b^2*(a + b*x)^2)`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

method	result	size
gospers	$-\frac{3bx+a}{6b^2(bx+a)^3}$	19
orering	$-\frac{3bx+a}{6b^2(bx+a)^3}$	19
norman	$\frac{-\frac{a}{6b^2} - \frac{x}{2b}}{(bx+a)^3}$	22
risch	$\frac{-\frac{a}{6b^2} - \frac{x}{2b}}{(bx+a)^3}$	22
parallelrisch	$\frac{-3b^2x-ab}{6b^3(bx+a)^3}$	24
default	$-\frac{1}{2b^2(bx+a)^2} + \frac{a}{3b^2(bx+a)^3}$	27

input `int(x/(b*x+a)^4,x,method=_RETURNVERBOSE)`

output `-1/6*(3*b*x+a)/b^2/(b*x+a)^3`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(21) = 42.

Time = 0.08 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.72

$$\int \frac{x}{(a+bx)^4} dx = -\frac{3bx+a}{6(b^5x^3+3ab^4x^2+3a^2b^3x+a^3b^2)}$$

input `integrate(x/(b*x+a)^4,x, algorithm="fricas")`

output `-1/6*(3*b*x + a)/(b^5*x^3 + 3*a*b^4*x^2 + 3*a^2*b^3*x + a^3*b^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(19) = 38$.

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.76

$$\int \frac{x}{(a+bx)^4} dx = \frac{-a-3bx}{6a^3b^2 + 18a^2b^3x + 18ab^4x^2 + 6b^5x^3}$$

input `integrate(x/(b*x+a)**4,x)`

output `(-a - 3*b*x)/(6*a**3*b**2 + 18*a**2*b**3*x + 18*a*b**4*x**2 + 6*b**5*x**3)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. $2(21) = 42$.

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.72

$$\int \frac{x}{(a+bx)^4} dx = -\frac{3bx+a}{6(b^5x^3 + 3ab^4x^2 + 3a^2b^3x + a^3b^2)}$$

input `integrate(x/(b*x+a)^4,x, algorithm="maxima")`

output `-1/6*(3*b*x + a)/(b^5*x^3 + 3*a*b^4*x^2 + 3*a^2*b^3*x + a^3*b^2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{x}{(a+bx)^4} dx = -\frac{3bx+a}{6(bx+a)^3b^2}$$

input `integrate(x/(b*x+a)^4,x, algorithm="giac")`

output `-1/6*(3*b*x + a)/((b*x + a)^3*b^2)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.76

$$\int \frac{x}{(a+bx)^4} dx = -\frac{\frac{a}{6b^2} + \frac{x}{2b}}{a^3 + 3a^2bx + 3ab^2x^2 + b^3x^3}$$

input `int(x/(a + b*x)^4,x)`output `-(a/(6*b^2) + x/(2*b))/(a^3 + b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.68

$$\int \frac{x}{(a+bx)^4} dx = \frac{-3bx - a}{6b^2(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3)}$$

input `int(x/(b*x+a)^4,x)`output `(- a - 3*b*x)/(6*b**2*(a**3 + 3*a**2*b*x + 3*a*b**2*x**2 + b**3*x**3))`

3.40 $\int \frac{x^2}{(a+bx)^4} dx$

Optimal result	279
Mathematica [A] (verified)	279
Rubi [A] (verified)	280
Maple [A] (verified)	281
Fricas [A] (verification not implemented)	281
Sympy [B] (verification not implemented)	282
Maxima [A] (verification not implemented)	282
Giac [A] (verification not implemented)	282
Mupad [B] (verification not implemented)	283
Reduce [B] (verification not implemented)	283

Optimal result

Integrand size = 11, antiderivative size = 34

$$\int \frac{x^2}{(a+bx)^4} dx = \frac{-\frac{a^2}{3b^3} - \frac{ax}{b^2} - \frac{x^2}{b}}{(a+bx)^3}$$

output `(-1/3*a^2/b^3-a*x/b^2-x^2/b)/(b*x+a)^3`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

$$\int \frac{x^2}{(a+bx)^4} dx = -\frac{a^2 + 3abx + 3b^2x^2}{3b^3(a+bx)^3}$$

input `Integrate[x^2/(a + b*x)^4,x]`

output `-1/3*(a^2 + 3*a*b*x + 3*b^2*x^2)/(b^3*(a + b*x)^3)`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.50, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(a + bx)^4} dx$$

↓ 48

$$\frac{x^3}{3a(a + bx)^3}$$

input `Int[x^2/(a + b*x)^4,x]`

output `x^3/(3*a*(a + b*x)^3)`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp`
`[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{`
`a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.88

method	result	size
gospers	$-\frac{3b^2x^2+3bax+a^2}{3b^3(bx+a)^3}$	30
orering	$-\frac{3b^2x^2+3bax+a^2}{3b^3(bx+a)^3}$	30
parallelrisc	$-\frac{3b^2x^2-3bax-a^2}{3b^3(bx+a)^3}$	32
norman	$-\frac{\frac{a^2}{3b^3}-\frac{ax}{b^2}-\frac{x^2}{b}}{(bx+a)^3}$	33
risc	$-\frac{\frac{a^2}{3b^3}-\frac{ax}{b^2}-\frac{x^2}{b}}{(bx+a)^3}$	33
default	$-\frac{1}{b^3(bx+a)} + \frac{a}{b^3(bx+a)^2} - \frac{a^2}{3b^3(bx+a)^3}$	41

input `int(x^2/(b*x+a)^4,x,method=_RETURNVERBOSE)`output $-1/3*(3*b^2*x^2+3*a*b*x+a^2)/b^3/(b*x+a)^3$ **Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.59

$$\int \frac{x^2}{(a+bx)^4} dx = -\frac{3b^2x^2+3abx+a^2}{3(b^6x^3+3ab^5x^2+3a^2b^4x+a^3b^3)}$$

input `integrate(x^2/(b*x+a)^4,x, algorithm="fricas")`output $-1/3*(3*b^2*x^2+3*a*b*x+a^2)/(b^6*x^3+3*a*b^5*x^2+3*a^2*b^4*x+a^3*b^3)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(27) = 54$.

Time = 0.12 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.65

$$\int \frac{x^2}{(a+bx)^4} dx = \frac{-a^2 - 3abx - 3b^2x^2}{3a^3b^3 + 9a^2b^4x + 9ab^5x^2 + 3b^6x^3}$$

input `integrate(x**2/(b*x+a)**4,x)`

output `(-a**2 - 3*a*b*x - 3*b**2*x**2)/(3*a**3*b**3 + 9*a**2*b**4*x + 9*a*b**5*x**2 + 3*b**6*x**3)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.59

$$\int \frac{x^2}{(a+bx)^4} dx = -\frac{3b^2x^2 + 3abx + a^2}{3(b^6x^3 + 3ab^5x^2 + 3a^2b^4x + a^3b^3)}$$

input `integrate(x^2/(b*x+a)^4,x, algorithm="maxima")`

output `-1/3*(3*b^2*x^2 + 3*a*b*x + a^2)/(b^6*x^3 + 3*a*b^5*x^2 + 3*a^2*b^4*x + a^3*b^3)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85

$$\int \frac{x^2}{(a+bx)^4} dx = -\frac{3b^2x^2 + 3abx + a^2}{3(bx+a)^3b^3}$$

input `integrate(x^2/(b*x+a)^4,x, algorithm="giac")`

output `-1/3*(3*b^2*x^2 + 3*a*b*x + a^2)/((b*x + a)^3*b^3)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.65

$$\int \frac{x^2}{(a+bx)^4} dx = -\frac{a^2 + 3abx + 3b^2x^2}{3a^3b^3 + 9a^2b^4x + 9ab^5x^2 + 3b^6x^3}$$

input `int(x^2/(a + b*x)^4,x)`output `-(a^2 + 3*b^2*x^2 + 3*a*b*x)/(3*a^3*b^3 + 3*b^6*x^3 + 9*a^2*b^4*x + 9*a*b^5*x^2)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.09

$$\int \frac{x^2}{(a+bx)^4} dx = \frac{x^3}{3a(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3)}$$

input `int(x^2/(b*x+a)^4,x)`output `x**3/(3*a*(a**3 + 3*a**2*b*x + 3*a*b**2*x**2 + b**3*x**3))`

3.41 $\int \frac{x^3}{(a+bx)^4} dx$

Optimal result	284
Mathematica [A] (verified)	284
Rubi [A] (verified)	285
Maple [A] (verified)	286
Fricas [A] (verification not implemented)	286
Sympy [A] (verification not implemented)	287
Maxima [A] (verification not implemented)	287
Giac [A] (verification not implemented)	287
Mupad [B] (verification not implemented)	288
Reduce [B] (verification not implemented)	288

Optimal result

Integrand size = 11, antiderivative size = 50

$$\int \frac{x^3}{(a+bx)^4} dx = \frac{\frac{11a^3}{6b^4} + \frac{9a^2x}{2b^2} + \frac{3ax^2}{b^2}}{(a+bx)^3} + \frac{\log(a+bx)}{b^4}$$

output $(11/6*a^3/b^4+9/2*a^2*x/b^2+3*a*x^2/b^2)/(b*x+a)^3+\ln(b*x+a)/b^4$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.88

$$\int \frac{x^3}{(a+bx)^4} dx = \frac{\frac{a(11a^2+27abx+18b^2x^2)}{(a+bx)^3} + 6\log(a+bx)}{6b^4}$$

input `Integrate[x^3/(a + b*x)^4,x]`

output $((a*(11*a^2 + 27*a*b*x + 18*b^2*x^2))/(a + b*x)^3 + 6*Log[a + b*x])/(6*b^4)$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.16, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(a+bx)^4} dx$$

↓ 49

$$\int \left(-\frac{a^3}{b^3(a+bx)^4} + \frac{3a^2}{b^3(a+bx)^3} - \frac{3a}{b^3(a+bx)^2} + \frac{1}{b^3(a+bx)} \right) dx$$

↓ 2009

$$\frac{a^3}{3b^4(a+bx)^3} - \frac{3a^2}{2b^4(a+bx)^2} + \frac{3a}{b^4(a+bx)} + \frac{\log(a+bx)}{b^4}$$

input `Int[x^3/(a + b*x)^4,x]`

output `a^3/(3*b^4*(a + b*x)^3) - (3*a^2)/(2*b^4*(a + b*x)^2) + (3*a)/(b^4*(a + b*x)) + Log[a + b*x]/b^4`

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.94

method	result	size
norman	$\frac{11a^3 + 3ax^2 + 9a^2x}{6b^4 + b^2 + 2b^3} + \frac{\ln(bx+a)}{b^4}$	47
risch	$\frac{11a^3 + 3ax^2 + 9a^2x}{6b^4 + b^2 + 2b^3} + \frac{\ln(bx+a)}{b^4}$	47
default	$\frac{3a}{b^4(bx+a)} - \frac{3a^2}{2b^4(bx+a)^2} + \frac{\ln(bx+a)}{b^4} + \frac{a^3}{3b^4(bx+a)^3}$	55
parallelrisch	$\frac{6 \ln(bx+a)x^3b^3 + 18 \ln(bx+a)x^2ab^2 + 18 \ln(bx+a)xa^2b + 18ab^2x^2 + 6 \ln(bx+a)a^3 + 27a^2bx + 11a^3}{6b^4(bx+a)^3}$	88

input `int(x^3/(b*x+a)^4,x,method=_RETURNVERBOSE)`output $(11/6*a^3/b^4+3*a*x^2/b^2+9/2*a^2*x/b^3)/(b*x+a)^3+\ln(b*x+a)/b^4$ **Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.88

$$\int \frac{x^3}{(a+bx)^4} dx$$

$$= \frac{18ab^2x^2 + 27a^2bx + 11a^3 + 6(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3)\log(bx+a)}{6(b^7x^3 + 3ab^6x^2 + 3a^2b^5x + a^3b^4)}$$

input `integrate(x^3/(b*x+a)^4,x, algorithm="fricas")`output $1/6*(18*a*b^2*x^2 + 27*a^2*b*x + 11*a^3 + 6*(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*\log(b*x + a))/(b^7*x^3 + 3*a*b^6*x^2 + 3*a^2*b^5*x + a^3*b^4)$

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.40

$$\int \frac{x^3}{(a+bx)^4} dx = \frac{11a^3 + 27a^2bx + 18ab^2x^2}{6a^3b^4 + 18a^2b^5x + 18ab^6x^2 + 6b^7x^3} + \frac{\log(a+bx)}{b^4}$$

input `integrate(x**3/(b*x+a)**4,x)`output `(11*a**3 + 27*a**2*b*x + 18*a*b**2*x**2)/(6*a**3*b**4 + 18*a**2*b**5*x + 18*a*b**6*x**2 + 6*b**7*x**3) + log(a + b*x)/b**4`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.40

$$\int \frac{x^3}{(a+bx)^4} dx = \frac{18ab^2x^2 + 27a^2bx + 11a^3}{6(b^7x^3 + 3ab^6x^2 + 3a^2b^5x + a^3b^4)} + \frac{\log(bx+a)}{b^4}$$

input `integrate(x^3/(b*x+a)^4,x, algorithm="maxima")`output `1/6*(18*a*b^2*x^2 + 27*a^2*b*x + 11*a^3)/(b^7*x^3 + 3*a*b^6*x^2 + 3*a^2*b^5*x + a^3*b^4) + log(b*x + a)/b^4`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.92

$$\int \frac{x^3}{(a+bx)^4} dx = \frac{\log(|bx+a|)}{b^4} + \frac{18abx^2 + 27a^2x + \frac{11a^3}{b}}{6(bx+a)^3b^3}$$

input `integrate(x^3/(b*x+a)^4,x, algorithm="giac")`output `log(abs(b*x + a))/b^4 + 1/6*(18*a*b*x^2 + 27*a^2*x + 11*a^3/b)/((b*x + a)^3*b^3)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.90

$$\int \frac{x^3}{(a+bx)^4} dx = \frac{\ln(a+bx) + \frac{3a}{a+bx} - \frac{3a^2}{2(a+bx)^2} + \frac{a^3}{3(a+bx)^3}}{b^4}$$

input `int(x^3/(a + b*x)^4,x)`output `(log(a + b*x) + (3*a)/(a + b*x) - (3*a^2)/(2*(a + b*x)^2) + a^3/(3*(a + b*x)^3))/b^4`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 108, normalized size of antiderivative = 2.16

$$\int \frac{x^3}{(a+bx)^4} dx = \frac{6 \log(bx+a) a^3 + 18 \log(bx+a) a^2 bx + 18 \log(bx+a) a b^2 x^2 + 6 \log(bx+a) b^3 x^3 + 5a^3 + 9a^2 bx - 6b^3 x^3}{6b^4 (b^3 x^3 + 3a b^2 x^2 + 3a^2 bx + a^3)}$$

input `int(x^3/(b*x+a)^4,x)`output `(6*log(a + b*x)*a**3 + 18*log(a + b*x)*a**2*b*x + 18*log(a + b*x)*a*b**2*x**2 + 6*log(a + b*x)*b**3*x**3 + 5*a**3 + 9*a**2*b*x - 6*b**3*x**3)/(6*b**4*(a**3 + 3*a**2*b*x + 3*a*b**2*x**2 + b**3*x**3))`

3.42 $\int \frac{1}{(a+bx)^5} dx$

Optimal result	289
Mathematica [A] (verified)	289
Rubi [A] (verified)	290
Maple [A] (verified)	291
Fricas [B] (verification not implemented)	291
Sympy [B] (verification not implemented)	292
Maxima [A] (verification not implemented)	292
Giac [A] (verification not implemented)	292
Mupad [B] (verification not implemented)	293
Reduce [B] (verification not implemented)	293

Optimal result

Integrand size = 7, antiderivative size = 14

$$\int \frac{1}{(a+bx)^5} dx = -\frac{1}{256b^4(a+bx)^4}$$

output `-1/256/b^4/(b*x+a)^4`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+bx)^5} dx = -\frac{1}{4b(a+bx)^4}$$

input `Integrate[(a + b*x)^(-5),x]`

output `-1/4*1/(b*(a + b*x)^4)`

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a+bx)^5} dx$$

↓ 17

$$-\frac{1}{4b(a+bx)^4}$$

input `Int[(a + b*x)^(-5),x]`

output `-1/4*1/(b*(a + b*x)^4)`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
gospers	$-\frac{1}{4(bx+a)^4b}$	13
default	$-\frac{1}{4(bx+a)^4b}$	13
norman	$-\frac{1}{4(bx+a)^4b}$	13
risch	$-\frac{1}{4(bx+a)^4b}$	13
parallelrisch	$-\frac{1}{4(bx+a)^4b}$	13
orering	$-\frac{1}{4(bx+a)^4b}$	13

input `int(1/(b*x+a)^5,x,method=_RETURNVERBOSE)`

output `-1/4/(b*x+a)^4/b`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. 2(12) = 24.

Time = 0.10 (sec) , antiderivative size = 46, normalized size of antiderivative = 3.29

$$\int \frac{1}{(a+bx)^5} dx = -\frac{1}{4(b^5x^4 + 4ab^4x^3 + 6a^2b^3x^2 + 4a^3b^2x + a^4b)}$$

input `integrate(1/(b*x+a)^5,x, algorithm="fricas")`

output `-1/4/(b^5*x^4 + 4*a*b^4*x^3 + 6*a^2*b^3*x^2 + 4*a^3*b^2*x + a^4*b)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(14) = 28$.

Time = 0.17 (sec) , antiderivative size = 49, normalized size of antiderivative = 3.50

$$\int \frac{1}{(a+bx)^5} dx = -\frac{1}{4a^4b + 16a^3b^2x + 24a^2b^3x^2 + 16ab^4x^3 + 4b^5x^4}$$

input `integrate(1/(b*x+a)**5,x)`

output `-1/(4*a**4*b + 16*a**3*b**2*x + 24*a**2*b**3*x**2 + 16*a*b**4*x**3 + 4*b**5*x**4)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a+bx)^5} dx = -\frac{1}{4(bx+a)^4b}$$

input `integrate(1/(b*x+a)^5,x, algorithm="maxima")`

output `-1/4/((b*x + a)^4*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a+bx)^5} dx = -\frac{1}{4(bx+a)^4b}$$

input `integrate(1/(b*x+a)^5,x, algorithm="giac")`

output `-1/4/((b*x + a)^4*b)`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 48, normalized size of antiderivative = 3.43

$$\int \frac{1}{(a + bx)^5} dx = -\frac{1}{4a^4b + 16a^3b^2x + 24a^2b^3x^2 + 16ab^4x^3 + 4b^5x^4}$$

input `int(1/(a + b*x)^5,x)`output `-1/(4*a^4*b + 4*b^5*x^4 + 16*a^3*b^2*x + 16*a*b^4*x^3 + 24*a^2*b^3*x^2)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 45, normalized size of antiderivative = 3.21

$$\int \frac{1}{(a + bx)^5} dx = -\frac{1}{4b(b^4x^4 + 4ab^3x^3 + 6a^2b^2x^2 + 4a^3bx + a^4)}$$

input `int(1/(b*x+a)^5,x)`output `(- 1)/(4*b*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2 + 4*a*b**3*x**3 + b**4*x**4))`

3.43 $\int \frac{x}{(a+bx)^5} dx$

Optimal result	294
Mathematica [A] (verified)	294
Rubi [A] (verified)	295
Maple [A] (verified)	296
Fricas [B] (verification not implemented)	296
Sympy [B] (verification not implemented)	297
Maxima [B] (verification not implemented)	297
Giac [A] (verification not implemented)	297
Mupad [B] (verification not implemented)	298
Reduce [B] (verification not implemented)	298

Optimal result

Integrand size = 9, antiderivative size = 25

$$\int \frac{x}{(a+bx)^5} dx = \frac{-\frac{a}{12b^2} - \frac{x}{3b}}{(a+bx)^4}$$

output $(-1/12*a/b^2-1/3*x/b)/(b*x+a)^4$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{x}{(a+bx)^5} dx = -\frac{a+4bx}{12b^2(a+bx)^4}$$

input `Integrate[x/(a + b*x)^5,x]`

output $-1/12*(a + 4*b*x)/(b^2*(a + b*x)^4)$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.20, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a+bx)^5} dx$$

↓ 53

$$\int \left(\frac{1}{b(a+bx)^4} - \frac{a}{b(a+bx)^5} \right) dx$$

↓ 2009

$$\frac{a}{4b^2(a+bx)^4} - \frac{1}{3b^2(a+bx)^3}$$

input `Int[x/(a + b*x)^5,x]`

output `a/(4*b^2*(a + b*x)^4) - 1/(3*b^2*(a + b*x)^3)`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

method	result	size
gospers	$-\frac{4bx+a}{12b^2(bx+a)^4}$	19
orering	$-\frac{4bx+a}{12b^2(bx+a)^4}$	19
norman	$\frac{-\frac{a}{12b^2} - \frac{x}{3b}}{(bx+a)^4}$	22
risch	$\frac{-\frac{a}{12b^2} - \frac{x}{3b}}{(bx+a)^4}$	22
parallelrisch	$\frac{-4b^3x - ab^2}{12b^4(bx+a)^4}$	26
default	$-\frac{1}{3b^2(bx+a)^3} + \frac{a}{4b^2(bx+a)^4}$	27

input `int(x/(b*x+a)^5,x,method=_RETURNVERBOSE)`

output `-1/12*(4*b*x+a)/b^2/(b*x+a)^4`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(21) = 42.

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.16

$$\int \frac{x}{(a+bx)^5} dx = -\frac{4bx+a}{12(b^6x^4 + 4ab^5x^3 + 6a^2b^4x^2 + 4a^3b^3x + a^4b^2)}$$

input `integrate(x/(b*x+a)^5,x, algorithm="fricas")`

output `-1/12*(4*b*x + a)/(b^6*x^4 + 4*a*b^5*x^3 + 6*a^2*b^4*x^2 + 4*a^3*b^3*x + a^4*b^2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(19) = 38$.

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.24

$$\int \frac{x}{(a+bx)^5} dx = \frac{-a-4bx}{12a^4b^2 + 48a^3b^3x + 72a^2b^4x^2 + 48ab^5x^3 + 12b^6x^4}$$

input `integrate(x/(b*x+a)**5,x)`

output `(-a - 4*b*x)/(12*a**4*b**2 + 48*a**3*b**3*x + 72*a**2*b**4*x**2 + 48*a*b**5*x**3 + 12*b**6*x**4)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. $2(21) = 42$.

Time = 0.03 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.16

$$\int \frac{x}{(a+bx)^5} dx = -\frac{4bx+a}{12(b^6x^4 + 4ab^5x^3 + 6a^2b^4x^2 + 4a^3b^3x + a^4b^2)}$$

input `integrate(x/(b*x+a)^5,x, algorithm="maxima")`

output `-1/12*(4*b*x + a)/(b^6*x^4 + 4*a*b^5*x^3 + 6*a^2*b^4*x^2 + 4*a^3*b^3*x + a^4*b^2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.24

$$\int \frac{x}{(a+bx)^5} dx = -\frac{\frac{4}{(bx+a)^3b} - \frac{3a}{(bx+a)^4b}}{12b}$$

input `integrate(x/(b*x+a)^5,x, algorithm="giac")`

output $-1/12*(4/((b*x + a)^3*b) - 3*a/((b*x + a)^4*b))/b$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{x}{(a + bx)^5} dx = -\frac{a + 4bx}{12b^2(a + bx)^4}$$

input `int(x/(a + b*x)^5,x)`

output $-(a + 4*b*x)/(12*b^2*(a + b*x)^4)$

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.12

$$\int \frac{x}{(a + bx)^5} dx = \frac{-4bx - a}{12b^2(b^4x^4 + 4ab^3x^3 + 6a^2b^2x^2 + 4a^3bx + a^4)}$$

input `int(x/(b*x+a)^5,x)`

output $(-a - 4*b*x)/(12*b**2*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2 + 4*a*b**3*x**3 + b**4*x**4))$

3.44 $\int \frac{x^2}{(a+bx)^5} dx$

Optimal result	299
Mathematica [A] (verified)	299
Rubi [A] (verified)	300
Maple [A] (verified)	301
Fricas [B] (verification not implemented)	301
Sympy [B] (verification not implemented)	302
Maxima [B] (verification not implemented)	302
Giac [A] (verification not implemented)	302
Mupad [B] (verification not implemented)	303
Reduce [B] (verification not implemented)	303

Optimal result

Integrand size = 11, antiderivative size = 38

$$\int \frac{x^2}{(a+bx)^5} dx = \frac{-\frac{a^2}{12b^3} - \frac{ax}{3b^2} - \frac{x^2}{2b}}{(a+bx)^4}$$

output $(-1/12*a^2/b^3-1/3*a*x/b^2-1/2*x^2/b)/(b*x+a)^4$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.82

$$\int \frac{x^2}{(a+bx)^5} dx = -\frac{a^2 + 4abx + 6b^2x^2}{12b^3(a+bx)^4}$$

input `Integrate[x^2/(a + b*x)^5,x]`

output $-1/12*(a^2 + 4*a*b*x + 6*b^2*x^2)/(b^3*(a + b*x)^4)$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.24, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(a+bx)^5} dx$$

$$\downarrow \text{53}$$

$$\int \left(\frac{a^2}{b^2(a+bx)^5} - \frac{2a}{b^2(a+bx)^4} + \frac{1}{b^2(a+bx)^3} \right) dx$$

$$\downarrow \text{2009}$$

$$-\frac{a^2}{4b^3(a+bx)^4} + \frac{2a}{3b^3(a+bx)^3} - \frac{1}{2b^3(a+bx)^2}$$

input `Int[x^2/(a + b*x)^5,x]`

output `-1/4*a^2/(b^3*(a + b*x)^4) + (2*a)/(3*b^3*(a + b*x)^3) - 1/(2*b^3*(a + b*x)^2)`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

method	result	size
gospers	$-\frac{6b^2x^2+4bax+a^2}{12b^3(bx+a)^4}$	30
orering	$-\frac{6b^2x^2+4bax+a^2}{12b^3(bx+a)^4}$	30
norman	$-\frac{\frac{a^2}{12b^3} - \frac{ax}{3b^2} - \frac{x^2}{2b}}{(bx+a)^4}$	33
risch	$-\frac{\frac{a^2}{12b^3} - \frac{ax}{3b^2} - \frac{x^2}{2b}}{(bx+a)^4}$	33
parallelrisch	$-\frac{6b^3x^2-4ab^2x-a^2b}{12b^4(bx+a)^4}$	35
default	$-\frac{1}{2b^3(bx+a)^2} + \frac{2a}{3b^3(bx+a)^3} - \frac{a^2}{4b^3(bx+a)^4}$	42

input `int(x^2/(b*x+a)^5,x,method=_RETURNVERBOSE)`

output `-1/12*(6*b^2*x^2+4*a*b*x+a^2)/b^3/(b*x+a)^4`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(32) = 64.

Time = 0.09 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.71

$$\int \frac{x^2}{(a+bx)^5} dx = -\frac{6b^2x^2 + 4abx + a^2}{12(b^7x^4 + 4ab^6x^3 + 6a^2b^5x^2 + 4a^3b^4x + a^4b^3)}$$

input `integrate(x^2/(b*x+a)^5,x, algorithm="fricas")`

output `-1/12*(6*b^2*x^2 + 4*a*b*x + a^2)/(b^7*x^4 + 4*a*b^6*x^3 + 6*a^2*b^5*x^2 + 4*a^3*b^4*x + a^4*b^3)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(31) = 62$.

Time = 0.19 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.79

$$\int \frac{x^2}{(a+bx)^5} dx = \frac{-a^2 - 4abx - 6b^2x^2}{12a^4b^3 + 48a^3b^4x + 72a^2b^5x^2 + 48ab^6x^3 + 12b^7x^4}$$

input `integrate(x**2/(b*x+a)**5,x)`

output `(-a**2 - 4*a*b*x - 6*b**2*x**2)/(12*a**4*b**3 + 48*a**3*b**4*x + 72*a**2*b**5*x**2 + 48*a*b**6*x**3 + 12*b**7*x**4)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(32) = 64$.

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.71

$$\int \frac{x^2}{(a+bx)^5} dx = -\frac{6b^2x^2 + 4abx + a^2}{12(b^7x^4 + 4ab^6x^3 + 6a^2b^5x^2 + 4a^3b^4x + a^4b^3)}$$

input `integrate(x^2/(b*x+a)^5,x, algorithm="maxima")`

output `-1/12*(6*b^2*x^2 + 4*a*b*x + a^2)/(b^7*x^4 + 4*a*b^6*x^3 + 6*a^2*b^5*x^2 + 4*a^3*b^4*x + a^4*b^3)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.21

$$\int \frac{x^2}{(a+bx)^5} dx = -\frac{6}{(bx+a)^2b^2} - \frac{8a}{(bx+a)^3b^2} + \frac{3a^2}{(bx+a)^4b^2}$$

input `integrate(x^2/(b*x+a)^5,x, algorithm="giac")`

output

$$\frac{-1/12*(6/((b*x + a)^2*b^2) - 8*a/((b*x + a)^3*b^2) + 3*a^2/((b*x + a)^4*b^2))/b}$$

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.58

$$\int \frac{x^2}{(a + bx)^5} dx = \frac{x^3(4a + bx)}{12a^2(a + bx)^4}$$

input

int(x^2/(a + b*x)^5,x)

output

(x^3*(4*a + b*x))/(12*a^2*(a + b*x)^4)

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.68

$$\int \frac{x^2}{(a + bx)^5} dx = \frac{-6b^2x^2 - 4abx - a^2}{12b^3(b^4x^4 + 4ab^3x^3 + 6a^2b^2x^2 + 4a^3bx + a^4)}$$

input

int(x^2/(b*x+a)^5,x)

output

(- a**2 - 4*a*b*x - 6*b**2*x**2)/(12*b**3*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2 + 4*a*b**3*x**3 + b**4*x**4))

3.45 $\int \frac{x^3}{(a+bx)^5} dx$

Optimal result	304
Mathematica [A] (verified)	304
Rubi [A] (verified)	305
Maple [A] (verified)	306
Fricas [A] (verification not implemented)	306
Sympy [A] (verification not implemented)	307
Maxima [A] (verification not implemented)	307
Giac [A] (verification not implemented)	307
Mupad [B] (verification not implemented)	308
Reduce [B] (verification not implemented)	308

Optimal result

Integrand size = 11, antiderivative size = 47

$$\int \frac{x^3}{(a+bx)^5} dx = \frac{-\frac{a^3}{4b^4} - \frac{a^2x}{b^3} - \frac{3ax^3}{2b^2} - \frac{x^3}{b}}{(a+bx)^4}$$

output `(-1/4*a^3/b^4-a^2*x/b^3-3/2*a*x^3/b^2-x^3/b)/(b*x+a)^4`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.89

$$\int \frac{x^3}{(a+bx)^5} dx = -\frac{a^3 + 4a^2bx + 6ab^2x^2 + 4b^3x^3}{4b^4(a+bx)^4}$$

input `Integrate[x^3/(a + b*x)^5,x]`

output `-1/4*(a^3 + 4*a^2*b*x + 6*a*b^2*x^2 + 4*b^3*x^3)/(b^4*(a + b*x)^4)`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.36, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {48}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(a+bx)^5} dx$$

↓ 48

$$\frac{x^4}{4a(a+bx)^4}$$

input `Int[x^3/(a + b*x)^5,x]`

output `x^4/(4*a*(a + b*x)^4)`

Defintions of rubi rules used

rule 48 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp`
`[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{`
`a, b, c, d, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.87

method	result	size
gospers	$-\frac{4b^3x^3+6ab^2x^2+4a^2bx+a^3}{4(bx+a)^4b^4}$	41
orering	$-\frac{4b^3x^3+6ab^2x^2+4a^2bx+a^3}{4(bx+a)^4b^4}$	41
parallelrisch	$-\frac{4b^3x^3-6ab^2x^2-4a^2bx-a^3}{4b^4(bx+a)^4}$	43
norman	$-\frac{\frac{x^3}{b}-\frac{3ax^2}{2b^2}-\frac{a^2x}{b^3}-\frac{a^3}{4b^4}}{(bx+a)^4}$	44
risch	$-\frac{\frac{x^3}{b}-\frac{3ax^2}{2b^2}-\frac{a^2x}{b^3}-\frac{a^3}{4b^4}}{(bx+a)^4}$	44
default	$-\frac{1}{b^4(bx+a)} + \frac{3a}{2b^4(bx+a)^2} - \frac{a^2}{b^4(bx+a)^3} + \frac{a^3}{4b^4(bx+a)^4}$	57

input `int(x^3/(b*x+a)^5,x,method=_RETURNVERBOSE)`output $-1/4*(4*b^3*x^3+6*a*b^2*x^2+4*a^2*b*x+a^3)/(b*x+a)^4/b^4$ **Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.62

$$\int \frac{x^3}{(a+bx)^5} dx = -\frac{4b^3x^3+6ab^2x^2+4a^2bx+a^3}{4(b^8x^4+4ab^7x^3+6a^2b^6x^2+4a^3b^5x+a^4b^4)}$$

input `integrate(x^3/(b*x+a)^5,x, algorithm="fricas")`output $-1/4*(4*b^3*x^3+6*a*b^2*x^2+4*a^2*b*x+a^3)/(b^8*x^4+4*a*b^7*x^3+6*a^2*b^6*x^2+4*a^3*b^5*x+a^4*b^4)$

Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.70

$$\int \frac{x^3}{(a+bx)^5} dx = \frac{-a^3 - 4a^2bx - 6ab^2x^2 - 4b^3x^3}{4a^4b^4 + 16a^3b^5x + 24a^2b^6x^2 + 16ab^7x^3 + 4b^8x^4}$$

input `integrate(x**3/(b*x+a)**5,x)`output `(-a**3 - 4*a**2*b*x - 6*a*b**2*x**2 - 4*b**3*x**3)/(4*a**4*b**4 + 16*a**3*b**5*x + 24*a**2*b**6*x**2 + 16*a*b**7*x**3 + 4*b**8*x**4)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.62

$$\int \frac{x^3}{(a+bx)^5} dx = -\frac{4b^3x^3 + 6ab^2x^2 + 4a^2bx + a^3}{4(b^8x^4 + 4ab^7x^3 + 6a^2b^6x^2 + 4a^3b^5x + a^4b^4)}$$

input `integrate(x^3/(b*x+a)^5,x, algorithm="maxima")`output `-1/4*(4*b^3*x^3 + 6*a*b^2*x^2 + 4*a^2*b*x + a^3)/(b^8*x^4 + 4*a*b^7*x^3 + 6*a^2*b^6*x^2 + 4*a^3*b^5*x + a^4*b^4)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.30

$$\int \frac{x^3}{(a+bx)^5} dx = -\frac{\frac{4}{(bx+a)b} - \frac{6a}{(bx+a)^2b} + \frac{4a^2}{(bx+a)^3b} - \frac{a^3}{(bx+a)^4b}}{4b^3}$$

input `integrate(x^3/(b*x+a)^5,x, algorithm="giac")`output `-1/4*(4/((b*x + a)*b) - 6*a/((b*x + a)^2*b) + 4*a^2/((b*x + a)^3*b) - a^3/((b*x + a)^4*b))/b^3`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.02

$$\int \frac{x^3}{(a+bx)^5} dx = \frac{\frac{3a}{2(a+bx)^2} - \frac{1}{a+bx} - \frac{a^2}{(a+bx)^3} + \frac{a^3}{4(a+bx)^4}}{b^4}$$

input `int(x^3/(a + b*x)^5,x)`output `((3*a)/(2*(a + b*x)^2) - 1/(a + b*x) - a^2/(a + b*x)^3 + a^3/(4*(a + b*x)^4))/b^4`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.02

$$\int \frac{x^3}{(a+bx)^5} dx = \frac{x^4}{4a(b^4x^4 + 4ab^3x^3 + 6a^2b^2x^2 + 4a^3bx + a^4)}$$

input `int(x^3/(b*x+a)^5,x)`output `x**4/(4*a*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2 + 4*a*b**3*x**3 + b**4*x**4))`

3.46 $\int \frac{1}{x(a+bx)} dx$

Optimal result	309
Mathematica [A] (verified)	309
Rubi [A] (verified)	310
Maple [A] (verified)	311
Fricas [A] (verification not implemented)	311
Sympy [A] (verification not implemented)	312
Maxima [A] (verification not implemented)	312
Giac [A] (verification not implemented)	312
Mupad [B] (verification not implemented)	313
Reduce [B] (verification not implemented)	313

Optimal result

Integrand size = 11, antiderivative size = 15

$$\int \frac{1}{x(a+bx)} dx = -\frac{\log\left(\frac{a+bx}{x}\right)}{a}$$

output `-ln((b*x+a)/x)/a`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \frac{1}{x(a+bx)} dx = \frac{\log(x)}{a} - \frac{\log(a+bx)}{a}$$

input `Integrate[1/(x*(a + b*x)),x]`

output `Log[x]/a - Log[a + b*x]/a`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x(a+bx)} dx \\ & \quad \downarrow 47 \\ & \frac{\int \frac{1}{x} dx}{a} - \frac{b \int \frac{1}{a+bx} dx}{a} \\ & \quad \downarrow 14 \\ & \frac{\log(x)}{a} - \frac{b \int \frac{1}{a+bx} dx}{a} \\ & \quad \downarrow 16 \\ & \frac{\log(x)}{a} - \frac{\log(a+bx)}{a} \end{aligned}$$

input `Int[1/(x*(a + b*x)),x]`

output `Log[x]/a - Log[a + b*x]/a`

Defintions of rubi rules used

rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
parallelsch	$\frac{\ln(x) - \ln(bx+a)}{a}$	16
default	$-\frac{\ln(bx+a)}{a} + \frac{\ln(x)}{a}$	19
norman	$-\frac{\ln(bx+a)}{a} + \frac{\ln(x)}{a}$	19
risch	$\frac{\ln(-x)}{a} - \frac{\ln(bx+a)}{a}$	21

input `int(1/x/(b*x+a),x,method=_RETURNVERBOSE)`

output `(ln(x)-ln(b*x+a))/a`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(a+bx)} dx = -\frac{\log(bx+a) - \log(x)}{a}$$

input `integrate(1/x/(b*x+a),x, algorithm="fricas")`

output `-(log(b*x + a) - log(x))/a`

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{1}{x(a+bx)} dx = \frac{\log(x) - \log\left(\frac{a}{b} + x\right)}{a}$$

input `integrate(1/x/(b*x+a),x)`output `(log(x) - log(a/b + x))/a`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \frac{1}{x(a+bx)} dx = -\frac{\log(bx+a)}{a} + \frac{\log(x)}{a}$$

input `integrate(1/x/(b*x+a),x, algorithm="maxima")`output `-log(b*x + a)/a + log(x)/a`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.33

$$\int \frac{1}{x(a+bx)} dx = -\frac{\log(|bx+a|)}{a} + \frac{\log(|x|)}{a}$$

input `integrate(1/x/(b*x+a),x, algorithm="giac")`output `-log(abs(b*x + a))/a + log(abs(x))/a`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(a+bx)} dx = -\frac{2 \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a}$$

input `int(1/(x*(a + b*x)),x)`

output `-(2*atanh((2*b*x)/a + 1))/a`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{x(a+bx)} dx = \frac{-\log(bx+a) + \log(x)}{a}$$

input `int(1/x/(b*x+a),x)`

output `(- log(a + b*x) + log(x))/a`

3.47 $\int \frac{1}{x^2(a+bx)} dx$

Optimal result	314
Mathematica [A] (verified)	314
Rubi [A] (verified)	315
Maple [A] (verified)	316
Fricas [A] (verification not implemented)	316
Sympy [A] (verification not implemented)	316
Maxima [A] (verification not implemented)	317
Giac [A] (verification not implemented)	317
Mupad [B] (verification not implemented)	317
Reduce [B] (verification not implemented)	318

Optimal result

Integrand size = 11, antiderivative size = 24

$$\int \frac{1}{x^2(a+bx)} dx = -\frac{1}{ax} + \frac{b \log\left(\frac{a+bx}{x}\right)}{a^2}$$

output `-1/a/x+b*ln((b*x+a)/x)/a^2`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{1}{x^2(a+bx)} dx = -\frac{1}{ax} - \frac{b \log(x)}{a^2} + \frac{b \log(a+bx)}{a^2}$$

input `Integrate[1/(x^2*(a + b*x)),x]`

output `-(1/(a*x)) - (b*Log[x])/a^2 + (b*Log[a + b*x])/a^2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2(a+bx)} dx$$

↓ 54

$$\int \left(\frac{b^2}{a^2(a+bx)} - \frac{b}{a^2x} + \frac{1}{ax^2} \right) dx$$

↓ 2009

$$-\frac{b \log(x)}{a^2} + \frac{b \log(a+bx)}{a^2} - \frac{1}{ax}$$

input `Int[1/(x^2*(a + b*x)),x]`

output `-(1/(a*x)) - (b*Log[x])/a^2 + (b*Log[a + b*x])/a^2`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

method	result	size
parallelrisc	$-\frac{b \ln(x)x - \ln(bx+a)xb+a}{a^2x}$	26
default	$\frac{b \ln(bx+a)}{a^2} - \frac{1}{ax} - \frac{b \ln(x)}{a^2}$	29
norman	$\frac{b \ln(bx+a)}{a^2} - \frac{1}{ax} - \frac{b \ln(x)}{a^2}$	29
risc	$-\frac{1}{ax} + \frac{b \ln(-bx-a)}{a^2} - \frac{b \ln(x)}{a^2}$	32

input `int(1/x^2/(b*x+a),x,method=_RETURNVERBOSE)`output `-(b*ln(x)*x-ln(b*x+a)*x*b+a)/a^2/x`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^2(a+bx)} dx = \frac{bx \log(bx+a) - bx \log(x) - a}{a^2x}$$

input `integrate(1/x^2/(b*x+a),x, algorithm="fricas")`output `(b*x*log(b*x + a) - b*x*log(x) - a)/(a^2*x)`**Sympy [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \frac{1}{x^2(a+bx)} dx = -\frac{1}{ax} + \frac{b(-\log(x) + \log(\frac{a}{b} + x))}{a^2}$$

input `integrate(1/x**2/(b*x+a),x)`

output `-1/(a*x) + b*(-log(x) + log(a/b + x))/a**2`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.17

$$\int \frac{1}{x^2(a+bx)} dx = \frac{b \log(bx+a)}{a^2} - \frac{b \log(x)}{a^2} - \frac{1}{ax}$$

input `integrate(1/x^2/(b*x+a),x, algorithm="maxima")`

output `b*log(b*x + a)/a^2 - b*log(x)/a^2 - 1/(a*x)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.25

$$\int \frac{1}{x^2(a+bx)} dx = \frac{b \log(|bx+a|)}{a^2} - \frac{b \log(|x|)}{a^2} - \frac{1}{ax}$$

input `integrate(1/x^2/(b*x+a),x, algorithm="giac")`

output `b*log(abs(b*x + a))/a^2 - b*log(abs(x))/a^2 - 1/(a*x)`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.04

$$\int \frac{1}{x^2(a+bx)} dx = \frac{2b \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^2} - \frac{1}{ax}$$

input `int(1/(x^2*(a + b*x)),x)`

output `(2*b*atanh((2*b*x)/a + 1))/a^2 - 1/(a*x)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^2(a+bx)} dx = \frac{\log(bx+a)bx - \log(x)bx - a}{a^2x}$$

input `int(1/x^2/(b*x+a),x)`

output `(log(a + b*x)*b*x - log(x)*b*x - a)/(a**2*x)`

3.48 $\int \frac{1}{x^3(a+bx)} dx$

Optimal result	319
Mathematica [A] (verified)	319
Rubi [A] (verified)	320
Maple [A] (verified)	321
Fricas [A] (verification not implemented)	321
Sympy [A] (verification not implemented)	321
Maxima [A] (verification not implemented)	322
Giac [A] (verification not implemented)	322
Mupad [B] (verification not implemented)	322
Reduce [B] (verification not implemented)	323

Optimal result

Integrand size = 11, antiderivative size = 37

$$\int \frac{1}{x^3(a+bx)} dx = -\frac{1}{2ax^2} + \frac{b}{a^2x} - \frac{b^2 \log\left(\frac{a+bx}{x}\right)}{a^3}$$

output `-1/2/a/x^2+b/a^2/x-b^2*ln((b*x+a)/x)/a^3`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14

$$\int \frac{1}{x^3(a+bx)} dx = -\frac{1}{2ax^2} + \frac{b}{a^2x} + \frac{b^2 \log(x)}{a^3} - \frac{b^2 \log(a+bx)}{a^3}$$

input `Integrate[1/(x^3*(a + b*x)),x]`

output `-1/2*1/(a*x^2) + b/(a^2*x) + (b^2*Log[x])/a^3 - (b^2*Log[a + b*x])/a^3`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.14, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3(a+bx)} dx$$

$$\downarrow 54$$

$$\int \left(-\frac{b^3}{a^3(a+bx)} + \frac{b^2}{a^3x} - \frac{b}{a^2x^2} + \frac{1}{ax^3} \right) dx$$

$$\downarrow 2009$$

$$\frac{b^2 \log(x)}{a^3} - \frac{b^2 \log(a+bx)}{a^3} + \frac{b}{a^2x} - \frac{1}{2ax^2}$$

input `Int[1/(x^3*(a + b*x)),x]`

output `-1/2*1/(a*x^2) + b/(a^2*x) + (b^2*Log[x])/a^3 - (b^2*Log[a + b*x])/a^3`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11

method	result	size
default	$-\frac{b^2 \ln(bx+a)}{a^3} - \frac{1}{2ax^2} + \frac{b^2 \ln(x)}{a^3} + \frac{b}{a^2x}$	41
norman	$\frac{\frac{bx}{a^2} - \frac{1}{2a}}{x^2} + \frac{b^2 \ln(x)}{a^3} - \frac{b^2 \ln(bx+a)}{a^3}$	41
risch	$\frac{\frac{bx}{a^2} - \frac{1}{2a}}{x^2} + \frac{b^2 \ln(-x)}{a^3} - \frac{b^2 \ln(bx+a)}{a^3}$	43
parallelrisch	$\frac{2b^2 \ln(x)x^2 - 2 \ln(bx+a)x^2 b^2 + 2bax - a^2}{2a^3x^2}$	44

input `int(1/x^3/(b*x+a),x,method=_RETURNVERBOSE)`output `-b^2/a^3*ln(b*x+a)-1/2/a/x^2+b^2/a^3*ln(x)+b/a^2/x`**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.11

$$\int \frac{1}{x^3(a+bx)} dx = -\frac{2b^2x^2 \log(bx+a) - 2b^2x^2 \log(x) - 2abx + a^2}{2a^3x^2}$$

input `integrate(1/x^3/(b*x+a),x, algorithm="fricas")`output `-1/2*(2*b^2*x^2*log(b*x + a) - 2*b^2*x^2*log(x) - 2*a*b*x + a^2)/(a^3*x^2)`**Sympy [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.84

$$\int \frac{1}{x^3(a+bx)} dx = \frac{-a+2bx}{2a^2x^2} + \frac{b^2(\log(x) - \log(\frac{a}{b} + x))}{a^3}$$

input `integrate(1/x**3/(b*x+a),x)`

output $(-a + 2bx)/(2a^2x^2) + b^2(\log(x) - \log(a/b + x))/a^3$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^3(a+bx)} dx = -\frac{b^2 \log(bx+a)}{a^3} + \frac{b^2 \log(x)}{a^3} + \frac{2bx-a}{2a^2x^2}$$

input `integrate(1/x^3/(b*x+a),x, algorithm="maxima")`

output $-b^2 \log(bx+a)/a^3 + b^2 \log(x)/a^3 + 1/2(2bx-a)/(a^2x^2)$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.22

$$\int \frac{1}{x^3(a+bx)} dx = -\frac{b^2 \log(|bx+a|)}{a^3} + \frac{b^2 \log(|x|)}{a^3} + \frac{2abx-a^2}{2a^3x^2}$$

input `integrate(1/x^3/(b*x+a),x, algorithm="giac")`

output $-b^2 \log(\text{abs}(bx+a))/a^3 + b^2 \log(\text{abs}(x))/a^3 + 1/2(2a*bx - a^2)/(a^3x^2)$

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.03

$$\int \frac{1}{x^3(a+bx)} dx = -\frac{\frac{a^2}{2} - abx}{a^3x^2} - \frac{2b^2 \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^3}$$

input `int(1/(x^3*(a + b*x)),x)`

output $-\frac{a^2/2 - a*b*x}{a^3*x^2} - \frac{(2*b^2*atanh((2*b*x)/a + 1))}{a^3}$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16

$$\int \frac{1}{x^3(a+bx)} dx = \frac{-2\log(bx+a)b^2x^2 + 2\log(x)b^2x^2 - a^2 + 2abx}{2a^3x^2}$$

input `int(1/x^3/(b*x+a),x)`

output `(- 2*log(a + b*x)*b**2*x**2 + 2*log(x)*b**2*x**2 - a**2 + 2*a*b*x)/(2*a**3*x**2)`

3.49 $\int \frac{1}{x^2(a+bx)^2} dx$

Optimal result	324
Mathematica [A] (verified)	324
Rubi [A] (verified)	325
Maple [A] (verified)	326
Fricas [A] (verification not implemented)	326
Sympy [A] (verification not implemented)	327
Maxima [A] (verification not implemented)	327
Giac [A] (verification not implemented)	327
Mupad [B] (verification not implemented)	328
Reduce [B] (verification not implemented)	328

Optimal result

Integrand size = 11, antiderivative size = 40

$$\int \frac{1}{x^2(a+bx)^2} dx = \frac{-\frac{2b}{a^2} - \frac{1}{ax}}{a+bx} + \frac{2b \log\left(\frac{a+bx}{x}\right)}{a^3}$$

output

```
(-2*b/a^2-1/a/x)/(b*x+a)+2*b*ln((b*x+a)/x)/a^3
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.88

$$\int \frac{1}{x^2(a+bx)^2} dx = -\frac{a\left(\frac{1}{x} + \frac{b}{a+bx}\right) + 2b \log(x) - 2b \log(a+bx)}{a^3}$$

input

```
Integrate[1/(x^2*(a + b*x)^2),x]
```

output

```
-((a*(x^(-1) + b/(a + b*x)) + 2*b*Log[x] - 2*b*Log[a + b*x])/a^3)
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.05, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2(a+bx)^2} dx$$

$$\downarrow 54$$

$$\int \left(\frac{2b^2}{a^3(a+bx)} - \frac{2b}{a^3x} + \frac{b^2}{a^2(a+bx)^2} + \frac{1}{a^2x^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2b \log(x)}{a^3} + \frac{2b \log(a+bx)}{a^3} - \frac{b}{a^2(a+bx)} - \frac{1}{a^2x}$$

input `Int[1/(x^2*(a + b*x)^2),x]`

output `-(1/(a^2*x)) - b/(a^2*(a + b*x)) - (2*b*Log[x])/a^3 + (2*b*Log[a + b*x])/a^3`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.08

method	result	size
default	$-\frac{b}{a^2(bx+a)} + \frac{2b \ln(bx+a)}{a^3} - \frac{1}{a^2x} - \frac{2b \ln(x)}{a^3}$	43
risch	$\frac{\frac{2bx}{a^2} - \frac{1}{a}}{x(bx+a)} + \frac{2b \ln(-bx-a)}{a^3} - \frac{2b \ln(x)}{a^3}$	49
norman	$\frac{\frac{2b^2x^2}{a^3} - \frac{1}{a}}{x(bx+a)} - \frac{2b \ln(x)}{a^3} + \frac{2b \ln(bx+a)}{a^3}$	50
parallelrisch	$-\frac{2b^2 \ln(x)x^2 - 2 \ln(bx+a)x^2b^2 + 2 \ln(x)xab - 2 \ln(bx+a)xab - 2b^2x^2 + a^2}{a^3x(bx+a)}$	70

input `int(1/x^2/(b*x+a)^2,x,method=_RETURNVERBOSE)`output `-b/a^2/(b*x+a)+2*b/a^3*ln(b*x+a)-1/a^2/x-2*b/a^3*ln(x)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.58

$$\int \frac{1}{x^2(a+bx)^2} dx = -\frac{2abx + a^2 - 2(b^2x^2 + abx) \log(bx+a) + 2(b^2x^2 + abx) \log(x)}{a^3bx^2 + a^4x}$$

input `integrate(1/x^2/(b*x+a)^2,x, algorithm="fricas")`output `-(2*a*b*x + a^2 - 2*(b^2*x^2 + a*b*x)*log(b*x + a) + 2*(b^2*x^2 + a*b*x)*log(x))/(a^3*b*x^2 + a^4*x)`

Sympy [A] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.92

$$\int \frac{1}{x^2(a+bx)^2} dx = \frac{-a-2bx}{a^3x+a^2bx^2} + \frac{2b(-\log(x) + \log(\frac{a}{b} + x))}{a^3}$$

input `integrate(1/x**2/(b*x+a)**2,x)`output `(-a - 2*b*x)/(a**3*x + a**2*b*x**2) + 2*b*(-log(x) + log(a/b + x))/a**3`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.12

$$\int \frac{1}{x^2(a+bx)^2} dx = -\frac{2bx+a}{a^2bx^2+a^3x} + \frac{2b \log(bx+a)}{a^3} - \frac{2b \log(x)}{a^3}$$

input `integrate(1/x^2/(b*x+a)^2,x, algorithm="maxima")`output `-(2*b*x + a)/(a^2*b*x^2 + a^3*x) + 2*b*log(b*x + a)/a^3 - 2*b*log(x)/a^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.30

$$\int \frac{1}{x^2(a+bx)^2} dx = -\frac{2b \log\left(\left|-\frac{a}{bx+a} + 1\right|\right)}{a^3} - \frac{b}{(bx+a)a^2} + \frac{b}{a^3\left(\frac{a}{bx+a} - 1\right)}$$

input `integrate(1/x^2/(b*x+a)^2,x, algorithm="giac")`output `-2*b*log(abs(-a/(b*x + a) + 1))/a^3 - b/((b*x + a)*a^2) + b/(a^3*(a/(b*x + a) - 1))`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.12

$$\int \frac{1}{x^2(a+bx)^2} dx = \frac{2b \ln\left(\frac{a+bx}{x}\right)}{a^3} - \frac{1}{ax(a+bx)} - \frac{2b}{a^2(a+bx)}$$

input `int(1/(x^2*(a + b*x)^2),x)`output `(2*b*log((a + b*x)/x))/a^3 - 1/(a*x*(a + b*x)) - (2*b)/(a^2*(a + b*x))`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.75

$$\int \frac{1}{x^2(a+bx)^2} dx$$

$$= \frac{2 \log(bx+a) abx + 2 \log(bx+a) b^2 x^2 - 2 \log(x) abx - 2 \log(x) b^2 x^2 - a^2 + 2b^2 x^2}{a^3 x (bx+a)}$$

input `int(1/x^2/(b*x+a)^2,x)`output `(2*log(a + b*x)*a*b*x + 2*log(a + b*x)*b**2*x**2 - 2*log(x)*a*b*x - 2*log(x)*b**2*x**2 - a**2 + 2*b**2*x**2)/(a**3*x*(a + b*x))`

3.50 $\int \frac{1}{x^3(a+bx)^2} dx$

Optimal result	329
Mathematica [A] (verified)	329
Rubi [A] (verified)	330
Maple [A] (verified)	331
Fricas [A] (verification not implemented)	331
Sympy [A] (verification not implemented)	332
Maxima [A] (verification not implemented)	332
Giac [A] (verification not implemented)	332
Mupad [B] (verification not implemented)	333
Reduce [B] (verification not implemented)	333

Optimal result

Integrand size = 11, antiderivative size = 57

$$\int \frac{1}{x^3(a+bx)^2} dx = \frac{\frac{3b^2}{a^3} - \frac{1}{2ax^2} + \frac{3b}{2a^2x}}{a+bx} - \frac{3b^2 \log\left(\frac{a+bx}{x}\right)}{a^4}$$

output `(3*b^2/a^3-1/2/a/x^2+3/2*b/a^2/x)/(b*x+a)-3*b^2*ln((b*x+a)/x)/a^4`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^3(a+bx)^2} dx = \frac{a\left(-\frac{a}{x^2} + \frac{4b}{x} + \frac{2b^2}{a+bx}\right) + 6b^2 \log(x) - 6b^2 \log(a+bx)}{2a^4}$$

input `Integrate[1/(x^3*(a + b*x)^2),x]`

output `(a*(-(a/x^2) + (4*b)/x + (2*b^2)/(a + b*x)) + 6*b^2*Log[x] - 6*b^2*Log[a + b*x])/(2*a^4)`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.02, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3(a+bx)^2} dx$$

$$\downarrow 54$$

$$\int \left(-\frac{3b^3}{a^4(a+bx)} + \frac{3b^2}{a^4x} - \frac{b^3}{a^3(a+bx)^2} - \frac{2b}{a^3x^2} + \frac{1}{a^2x^3} \right) dx$$

$$\downarrow 2009$$

$$\frac{3b^2 \log(x)}{a^4} - \frac{3b^2 \log(a+bx)}{a^4} + \frac{b^2}{a^3(a+bx)} + \frac{2b}{a^3x} - \frac{1}{2a^2x^2}$$

input `Int[1/(x^3*(a + b*x)^2),x]`

output `-1/2*1/(a^2*x^2) + (2*b)/(a^3*x) + b^2/(a^3*(a + b*x)) + (3*b^2*Log[x])/a^4 - (3*b^2*Log[a + b*x])/a^4`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{3b^2 \ln(bx+a)}{a^4} + \frac{b^2}{a^3(bx+a)} - \frac{1}{2a^2x^2} + \frac{2b}{a^3x} + \frac{3b^2 \ln(x)}{a^4}$	57
norman	$\frac{-\frac{3b^3x^3}{a^4} - \frac{1}{2a} + \frac{3bx}{2a^2}}{x^2(bx+a)} + \frac{3b^2 \ln(x)}{a^4} - \frac{3b^2 \ln(bx+a)}{a^4}$	61
risch	$\frac{\frac{3b^2x^2}{a^3} + \frac{3bx}{2a^2} - \frac{1}{2a}}{x^2(bx+a)} + \frac{3b^2 \ln(-x)}{a^4} - \frac{3b^2 \ln(bx+a)}{a^4}$	63
parallelrisch	$\frac{6 \ln(x)x^3b^3 - 6 \ln(bx+a)x^3b^3 + 6 \ln(x)x^2ab^2 - 6 \ln(bx+a)x^2ab^2 - 6b^3x^3 + 3a^2bx - a^3}{2a^4x^2(bx+a)}$	87

input `int(1/x^3/(b*x+a)^2,x,method=_RETURNVERBOSE)`output `-3*b^2/a^4*ln(b*x+a)+b^2/a^3/(b*x+a)-1/2/a^2/x^2+2*b/a^3/x+3*b^2/a^4*ln(x)`**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.51

$$\int \frac{1}{x^3(a+bx)^2} dx$$

$$= \frac{6ab^2x^2 + 3a^2bx - a^3 - 6(b^3x^3 + ab^2x^2) \log(bx+a) + 6(b^3x^3 + ab^2x^2) \log(x)}{2(a^4bx^3 + a^5x^2)}$$

input `integrate(1/x^3/(b*x+a)^2,x, algorithm="fricas")`output `1/2*(6*a*b^2*x^2 + 3*a^2*b*x - a^3 - 6*(b^3*x^3 + a*b^2*x^2)*log(b*x + a) + 6*(b^3*x^3 + a*b^2*x^2)*log(x))/(a^4*b*x^3 + a^5*x^2)`

Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.95

$$\int \frac{1}{x^3(a+bx)^2} dx = \frac{-a^2 + 3abx + 6b^2x^2}{2a^4x^2 + 2a^3bx^3} + \frac{3b^2(\log(x) - \log(\frac{a}{b} + x))}{a^4}$$

input `integrate(1/x**3/(b*x+a)**2,x)`output `(-a**2 + 3*a*b*x + 6*b**2*x**2)/(2*a**4*x**2 + 2*a**3*b*x**3) + 3*b**2*(log(x) - log(a/b + x))/a**4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.12

$$\int \frac{1}{x^3(a+bx)^2} dx = \frac{6b^2x^2 + 3abx - a^2}{2(a^3bx^3 + a^4x^2)} - \frac{3b^2 \log(bx + a)}{a^4} + \frac{3b^2 \log(x)}{a^4}$$

input `integrate(1/x^3/(b*x+a)^2,x, algorithm="maxima")`output `1/2*(6*b^2*x^2 + 3*a*b*x - a^2)/(a^3*b*x^3 + a^4*x^2) - 3*b^2*log(b*x + a)/a^4 + 3*b^2*log(x)/a^4`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.30

$$\int \frac{1}{x^3(a+bx)^2} dx = \frac{3b^2 \log\left(\left|-\frac{a}{bx+a} + 1\right|\right)}{a^4} + \frac{b^2}{(bx+a)a^3} - \frac{\frac{6ab^2}{bx+a} - 5b^2}{2a^4\left(\frac{a}{bx+a} - 1\right)^2}$$

input `integrate(1/x^3/(b*x+a)^2,x, algorithm="giac")`output `3*b^2*log(abs(-a/(b*x + a) + 1))/a^4 + b^2/((b*x + a)*a^3) - 1/2*(6*a*b^2/(b*x + a) - 5*b^2)/(a^4*(a/(b*x + a) - 1)^2)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^3(a+bx)^2} dx = \frac{\frac{3b^2x^2}{a^3} - \frac{1}{2a} + \frac{3bx}{2a^2}}{bx^3 + ax^2} - \frac{6b^2 \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^4}$$

input `int(1/(x^3*(a + b*x)^2),x)`output `((3*b^2*x^2)/a^3 - 1/(2*a) + (3*b*x)/(2*a^2))/(a*x^2 + b*x^3) - (6*b^2*atanh((2*b*x)/a + 1))/a^4`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.51

$$\int \frac{1}{x^3(a+bx)^2} dx = \frac{-6 \log(bx+a) a b^2 x^2 - 6 \log(bx+a) b^3 x^3 + 6 \log(x) a b^2 x^2 + 6 \log(x) b^3 x^3 - a^3 + 3a^2 b x - 6b^3 x^3}{2a^4 x^2 (bx+a)}$$

input `int(1/x^3/(b*x+a)^2,x)`output `(- 6*log(a + b*x)*a*b**2*x**2 - 6*log(a + b*x)*b**3*x**3 + 6*log(x)*a*b**2*x**2 + 6*log(x)*b**3*x**3 - a**3 + 3*a**2*b*x - 6*b**3*x**3)/(2*a**4*x**2*(a + b*x))`

3.51 $\int \frac{1}{x(a+bx)^3} dx$

Optimal result	334
Mathematica [A] (verified)	334
Rubi [A] (verified)	335
Maple [A] (verified)	336
Fricas [B] (verification not implemented)	336
Sympy [A] (verification not implemented)	337
Maxima [A] (verification not implemented)	337
Giac [A] (verification not implemented)	337
Mupad [B] (verification not implemented)	338
Reduce [B] (verification not implemented)	338

Optimal result

Integrand size = 11, antiderivative size = 38

$$\int \frac{1}{x(a+bx)^3} dx = \frac{\frac{3}{2a} + \frac{bx}{a^2}}{(a+bx)^2} - \frac{\log\left(\frac{a+bx}{x}\right)}{a^3}$$

output (3/2/a+b*x/a^2)/(b*x+a)^2-ln((b*x+a)/x)/a^3

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.97

$$\int \frac{1}{x(a+bx)^3} dx = \frac{\frac{a(3a+2bx)}{(a+bx)^2} + 2\log(x) - 2\log(a+bx)}{2a^3}$$

input Integrate[1/(x*(a + b*x)^3),x]

output ((a*(3*a + 2*b*x))/(a + b*x)^2 + 2*Log[x] - 2*Log[a + b*x])/(2*a^3)

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(a+bx)^3} dx$$

$$\downarrow 54$$

$$\int \left(-\frac{b}{a^3(a+bx)} + \frac{1}{a^3x} - \frac{b}{a^2(a+bx)^2} - \frac{b}{a(a+bx)^3} \right) dx$$

$$\downarrow 2009$$

$$-\frac{\log(a+bx)}{a^3} + \frac{\log(x)}{a^3} + \frac{1}{a^2(a+bx)} + \frac{1}{2a(a+bx)^2}$$

input `Int[1/(x*(a + b*x)^3),x]`

output `1/(2*a*(a + b*x)^2) + 1/(a^2*(a + b*x)) + Log[x]/a^3 - Log[a + b*x]/a^3`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.08

method	result	size
risch	$\frac{\frac{3}{2a} + \frac{bx}{a^2}}{(bx+a)^2} + \frac{\ln(-x)}{a^3} - \frac{\ln(bx+a)}{a^3}$	41
default	$-\frac{\ln(bx+a)}{a^3} + \frac{1}{a^2(bx+a)} + \frac{1}{2a(bx+a)^2} + \frac{\ln(x)}{a^3}$	42
norman	$-\frac{2bx - \frac{3b^2x^2}{2a^3}}{(bx+a)^2} + \frac{\ln(x)}{a^3} - \frac{\ln(bx+a)}{a^3}$	46
parallelrisch	$\frac{2b^2 \ln(x)x^2 - 2 \ln(bx+a)x^2b^2 + 4 \ln(x)xab - 4 \ln(bx+a)xab - 3b^2x^2 + 2 \ln(x)a^2 - 2a^2 \ln(bx+a) - 4bax}{2a^3(bx+a)^2}$	87

input `int(1/x/(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `(3/2/a+b*x/a^2)/(b*x+a)^2+1/a^3*ln(-x)-1/a^3*ln(b*x+a)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. 2(38) = 76.

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.11

$$\int \frac{1}{x(a+bx)^3} dx$$

$$= \frac{2abx + 3a^2 - 2(b^2x^2 + 2abx + a^2) \log(bx+a) + 2(b^2x^2 + 2abx + a^2) \log(x)}{2(a^3b^2x^2 + 2a^4bx + a^5)}$$

input `integrate(1/x/(b*x+a)^3,x, algorithm="fricas")`

output `1/2*(2*a*b*x + 3*a^2 - 2*(b^2*x^2 + 2*a*b*x + a^2)*log(b*x + a) + 2*(b^2*x^2 + 2*a*b*x + a^2)*log(x))/(a^3*b^2*x^2 + 2*a^4*b*x + a^5)`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.21

$$\int \frac{1}{x(a+bx)^3} dx = \frac{3a+2bx}{2a^4+4a^3bx+2a^2b^2x^2} + \frac{\log(x) - \log\left(\frac{a}{b} + x\right)}{a^3}$$

input `integrate(1/x/(b*x+a)**3,x)`output `(3*a + 2*b*x)/(2*a**4 + 4*a**3*b*x + 2*a**2*b**2*x**2) + (log(x) - log(a/b + x))/a**3`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int \frac{1}{x(a+bx)^3} dx = \frac{2bx+3a}{2(a^2b^2x^2+2a^3bx+a^4)} - \frac{\log(bx+a)}{a^3} + \frac{\log(x)}{a^3}$$

input `integrate(1/x/(b*x+a)^3,x, algorithm="maxima")`output `1/2*(2*b*x + 3*a)/(a^2*b^2*x^2 + 2*a^3*b*x + a^4) - log(b*x + a)/a^3 + log(x)/a^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13

$$\int \frac{1}{x(a+bx)^3} dx = -\frac{\log(|bx+a|)}{a^3} + \frac{\log(|x|)}{a^3} + \frac{2abx+3a^2}{2(bx+a)^2a^3}$$

input `integrate(1/x/(b*x+a)^3,x, algorithm="giac")`output `-log(abs(b*x + a))/a^3 + log(abs(x))/a^3 + 1/2*(2*a*b*x + 3*a^2)/((b*x + a)^2*a^3)`

Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.13

$$\int \frac{1}{x(a+bx)^3} dx = \frac{1}{a^2+bx a} - \frac{\ln\left(\frac{a+bx}{x}\right)}{a^2} + \frac{1}{2a(a+bx)^2}$$

input `int(1/(x*(a + b*x)^3),x)`output `(1/(a^2 + a*b*x) - log((a + b*x)/x)/a^2)/a + 1/(2*a*(a + b*x)^2)`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.55

$$\int \frac{1}{x(a+bx)^3} dx = \frac{-2\log(bx+a)a^2 - 4\log(bx+a)abx - 2\log(bx+a)b^2x^2 + 2\log(x)a^2 + 4\log(x)abx + 2\log(x)b^2x^2}{2a^3(b^2x^2 + 2abx + a^2)}$$

input `int(1/x/(b*x+a)^3,x)`output `(- 2*log(a + b*x)*a**2 - 4*log(a + b*x)*a*b*x - 2*log(a + b*x)*b**2*x**2 + 2*log(x)*a**2 + 4*log(x)*a*b*x + 2*log(x)*b**2*x**2 + 2*a**2 - b**2*x**2)/(2*a**3*(a**2 + 2*a*b*x + b**2*x**2))`

3.52 $\int \frac{1}{x^2(a+bx)^3} dx$

Optimal result	339
Mathematica [A] (verified)	339
Rubi [A] (verified)	340
Maple [A] (verified)	341
Fricas [B] (verification not implemented)	341
Sympy [A] (verification not implemented)	342
Maxima [A] (verification not implemented)	342
Giac [A] (verification not implemented)	342
Mupad [B] (verification not implemented)	343
Reduce [B] (verification not implemented)	343

Optimal result

Integrand size = 11, antiderivative size = 51

$$\int \frac{1}{x^2(a+bx)^3} dx = \frac{-\frac{9b}{2a^2} - \frac{1}{ax} - \frac{3b^2x}{a^3}}{(a+bx)^2} + \frac{3b \log\left(\frac{a+bx}{x}\right)}{a^4}$$

output $(-9/2*b/a^2-1/a/x-3*b^2*x/a^3)/(b*x+a)^2+3*b*\ln((b*x+a)/x)/a^4$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.04

$$\int \frac{1}{x^2(a+bx)^3} dx = -\frac{\frac{a(2a^2+9abx+6b^2x^2)}{x(a+bx)^2} + 6b \log(x) - 6b \log(a+bx)}{2a^4}$$

input `Integrate[1/(x^2*(a + b*x)^3),x]`

output $-1/2*((a*(2*a^2 + 9*a*b*x + 6*b^2*x^2))/(x*(a + b*x)^2) + 6*b*\text{Log}[x] - 6*b*\text{Log}[a + b*x])/a^4$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.12, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2(a+bx)^3} dx$$

$$\downarrow 54$$

$$\int \left(\frac{3b^2}{a^4(a+bx)} - \frac{3b}{a^4x} + \frac{2b^2}{a^3(a+bx)^2} + \frac{1}{a^3x^2} + \frac{b^2}{a^2(a+bx)^3} \right) dx$$

$$\downarrow 2009$$

$$-\frac{3b \log(x)}{a^4} + \frac{3b \log(a+bx)}{a^4} - \frac{2b}{a^3(a+bx)} - \frac{1}{a^3x} - \frac{b}{2a^2(a+bx)^2}$$

input `Int[1/(x^2*(a + b*x)^3),x]`

output `-(1/(a^3*x)) - b/(2*a^2*(a + b*x)^2) - (2*b)/(a^3*(a + b*x)) - (3*b*Log[x])/a^4 + (3*b*Log[a + b*x])/a^4`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.10

method	result
default	$-\frac{b}{2a^2(bx+a)^2} + \frac{3b \ln(bx+a)}{a^4} - \frac{2b}{a^3(bx+a)} - \frac{1}{a^3x} - \frac{3b \ln(x)}{a^4}$
risch	$\frac{-\frac{3b^2x^2}{a^3} - \frac{9bx}{2a^2} - \frac{1}{a}}{x(bx+a)^2} - \frac{3b \ln(x)}{a^4} + \frac{3b \ln(-bx-a)}{a^4}$
norman	$\frac{-\frac{1}{a} + \frac{6b^2x^2}{a^3} + \frac{9b^3x^3}{2a^4}}{x(bx+a)^2} - \frac{3b \ln(x)}{a^4} + \frac{3b \ln(bx+a)}{a^4}$
parallelrisch	$-\frac{6 \ln(x)x^3b^3 - 6 \ln(bx+a)x^3b^3 + 12 \ln(x)x^2a^2b^2 - 12 \ln(bx+a)x^2a^2b^2 - 9b^3x^3 + 6 \ln(x)a^2b - 6 \ln(bx+a)a^2b - 12a^2b^2x^2 + 2a^3}{2a^4x(bx+a)^2}$

input `int(1/x^2/(b*x+a)^3,x,method=_RETURNVERBOSE)`output `-1/2*b/a^2/(b*x+a)^2+3/a^4*b*ln(b*x+a)-2*b/a^3/(b*x+a)-1/a^3/x-3/a^4*b*ln(x)`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 109 vs. 2(50) = 100.

Time = 0.10 (sec) , antiderivative size = 109, normalized size of antiderivative = 2.14

$$\int \frac{1}{x^2(a+bx)^3} dx = \frac{6ab^2x^2 + 9a^2bx + 2a^3 - 6(b^3x^3 + 2ab^2x^2 + a^2bx) \log(bx+a) + 6(b^3x^3 + 2ab^2x^2 + a^2bx) \log(x)}{2(a^4b^2x^3 + 2a^5bx^2 + a^6x)}$$

input `integrate(1/x^2/(b*x+a)^3,x, algorithm="fricas")`output `-1/2*(6*a*b^2*x^2 + 9*a^2*b*x + 2*a^3 - 6*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)*log(b*x + a) + 6*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)*log(x))/(a^4*b^2*x^3 + 2*a^5*b*x^2 + a^6*x)`

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.29

$$\int \frac{1}{x^2(a+bx)^3} dx = \frac{-2a^2 - 9abx - 6b^2x^2}{2a^5x + 4a^4bx^2 + 2a^3b^2x^3} + \frac{3b(-\log(x) + \log(\frac{a}{b} + x))}{a^4}$$

input `integrate(1/x**2/(b*x+a)**3,x)`output `(-2*a**2 - 9*a*b*x - 6*b**2*x**2)/(2*a**5*x + 4*a**4*b*x**2 + 2*a**3*b**2*x**3) + 3*b*(-log(x) + log(a/b + x))/a**4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.35

$$\int \frac{1}{x^2(a+bx)^3} dx = -\frac{6b^2x^2 + 9abx + 2a^2}{2(a^3b^2x^3 + 2a^4bx^2 + a^5x)} + \frac{3b \log(bx + a)}{a^4} - \frac{3b \log(x)}{a^4}$$

input `integrate(1/x^2/(b*x+a)^3,x, algorithm="maxima")`output `-1/2*(6*b^2*x^2 + 9*a*b*x + 2*a^2)/(a^3*b^2*x^3 + 2*a^4*b*x^2 + a^5*x) + 3*b*log(b*x + a)/a^4 - 3*b*log(x)/a^4`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.18

$$\int \frac{1}{x^2(a+bx)^3} dx = \frac{3b \log(|bx + a|)}{a^4} - \frac{3b \log(|x|)}{a^4} - \frac{6ab^2x^2 + 9a^2bx + 2a^3}{2(bx + a)^2a^4x}$$

input `integrate(1/x^2/(b*x+a)^3,x, algorithm="giac")`output `3*b*log(abs(b*x + a))/a^4 - 3*b*log(abs(x))/a^4 - 1/2*(6*a*b^2*x^2 + 9*a^2*b*x + 2*a^3)/((b*x + a)^2*a^4*x)`

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.24

$$\int \frac{1}{x^2(a+bx)^3} dx = \frac{6b \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^4} - \frac{\frac{1}{a} + \frac{3b^2x^2}{a^3} + \frac{9bx}{2a^2}}{a^2x + 2abx^2 + b^2x^3}$$

input `int(1/(x^2*(a + b*x)^3),x)`output `(6*b*atanh((2*b*x)/a + 1))/a^4 - (1/a + (3*b^2*x^2)/a^3 + (9*b*x)/(2*a^2)) / (a^2*x + b^2*x^3 + 2*a*b*x^2)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.33

$$\int \frac{1}{x^2(a+bx)^3} dx = \frac{6 \log(bx+a) a^2 bx + 12 \log(bx+a) a b^2 x^2 + 6 \log(bx+a) b^3 x^3 - 6 \log(x) a^2 bx - 12 \log(x) a b^2 x^2 - 6 \log(x) a^2}{2a^4 x (b^2 x^2 + 2abx + a^2)}$$

input `int(1/x^2/(b*x+a)^3,x)`output `(6*log(a + b*x)*a**2*b*x + 12*log(a + b*x)*a*b**2*x**2 + 6*log(a + b*x)*b**3*x**3 - 6*log(x)*a**2*b*x - 12*log(x)*a*b**2*x**2 - 6*log(x)*b**3*x**3 - 2*a**3 - 6*a**2*b*x + 3*b**3*x**3)/(2*a**4*x*(a**2 + 2*a*b*x + b**2*x**2))`

3.53 $\int \frac{1}{x^3(a+bx)^3} dx$

Optimal result	344
Mathematica [A] (verified)	344
Rubi [A] (verified)	345
Maple [A] (verified)	346
Fricas [B] (verification not implemented)	346
Sympy [A] (verification not implemented)	347
Maxima [A] (verification not implemented)	347
Giac [A] (verification not implemented)	347
Mupad [B] (verification not implemented)	348
Reduce [B] (verification not implemented)	348

Optimal result

Integrand size = 11, antiderivative size = 64

$$\int \frac{1}{x^3(a+bx)^3} dx = \frac{\frac{9b^2}{a^3} - \frac{1}{2ax^2} + \frac{2b}{a^2x} + \frac{6b^3x}{a^4}}{(a+bx)^2} - \frac{6b^2 \log\left(\frac{a+bx}{x}\right)}{a^5}$$

output `((9*b^2/a^3-1/2/a/x^2+2*b/a^2/x+6*b^3*x/a^4)/(b*x+a)^2-6*b^2*ln((b*x+a)/x)/a^5)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.06

$$\int \frac{1}{x^3(a+bx)^3} dx = \frac{\frac{a(-a^3+4a^2bx+18ab^2x^2+12b^3x^3)}{x^2(a+bx)^2} + 12b^2 \log(x) - 12b^2 \log(a+bx)}{2a^5}$$

input `Integrate[1/(x^3*(a + b*x)^3),x]`

output `((a*(-a^3 + 4*a^2*b*x + 18*a*b^2*x^2 + 12*b^3*x^3))/(x^2*(a + b*x)^2) + 12*b^2*Log[x] - 12*b^2*Log[a + b*x])/(2*a^5)`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.19, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3(a+bx)^3} dx$$

↓ 54

$$\int \left(-\frac{6b^3}{a^5(a+bx)} + \frac{6b^2}{a^5x} - \frac{3b^3}{a^4(a+bx)^2} - \frac{3b}{a^4x^2} - \frac{b^3}{a^3(a+bx)^3} + \frac{1}{a^3x^3} \right) dx$$

↓ 2009

$$\frac{6b^2 \log(x)}{a^5} - \frac{6b^2 \log(a+bx)}{a^5} + \frac{3b^2}{a^4(a+bx)} + \frac{3b}{a^4x} + \frac{b^2}{2a^3(a+bx)^2} - \frac{1}{2a^3x^2}$$

input `Int[1/(x^3*(a + b*x)^3),x]`

output `-1/2*1/(a^3*x^2) + (3*b)/(a^4*x) + b^2/(2*a^3*(a + b*x)^2) + (3*b^2)/(a^4*(a + b*x)) + (6*b^2*Log[x])/a^5 - (6*b^2*Log[a + b*x])/a^5`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.12

method	result
norman	$\frac{-\frac{9b^4x^4}{a^5} - \frac{1}{2a} + \frac{2bx}{a^2} - \frac{12b^3x^3}{a^4}}{x^2(bx+a)^2} + \frac{6b^2 \ln(x)}{a^5} - \frac{6b^2 \ln(bx+a)}{a^5}$
default	$-\frac{6b^2 \ln(bx+a)}{a^5} + \frac{3b^2}{a^4(bx+a)} + \frac{b^2}{2a^3(bx+a)^2} - \frac{1}{2a^3x^2} + \frac{6b^2 \ln(x)}{a^5} + \frac{3b}{a^4x}$
risch	$\frac{\frac{6b^3x^3}{a^4} + \frac{9b^2x^2}{a^3} + \frac{2bx}{a^2} - \frac{1}{2a}}{x^2(bx+a)^2} + \frac{6b^2 \ln(-x)}{a^5} - \frac{6b^2 \ln(bx+a)}{a^5}$
parallelrisch	$\frac{12 \ln(x)x^4b^6 - 12 \ln(bx+a)x^4b^6 + 24 \ln(x)x^3ab^5 - 24 \ln(bx+a)x^3ab^5 + 12 \ln(x)x^2a^2b^4 - 12 \ln(bx+a)x^2a^2b^4 + 12x^3ab^5 + 18x^2a^2b^4}{2a^5b^2x^2(bx+a)^2}$

input `int(1/x^3/(b*x+a)^3,x,method=_RETURNVERBOSE)`output
$$\frac{(-9b^4/a^5x^4 - 1/2/a + 2bx/a^2 - 12b^3x^3/a^4)/x^2/(bx+a)^2 + 6/a^5b^2 \ln(x) - 6/a^5b^2 \ln(bx+a)}$$
Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 130 vs. 2(63) = 126.

Time = 0.21 (sec) , antiderivative size = 130, normalized size of antiderivative = 2.03

$$\int \frac{1}{x^3(a+bx)^3} dx$$

$$= \frac{12ab^3x^3 + 18a^2b^2x^2 + 4a^3bx - a^4 - 12(b^4x^4 + 2ab^3x^3 + a^2b^2x^2) \log(bx+a) + 12(b^4x^4 + 2ab^3x^3 + a^2b^2x^2) \log(x)}{2(a^5b^2x^4 + 2a^6bx^3 + a^7x^2)}$$

input `integrate(1/x^3/(b*x+a)^3,x, algorithm="fricas")`output
$$\frac{1/2*(12*a*b^3*x^3 + 18*a^2*b^2*x^2 + 4*a^3*b*x - a^4 - 12*(b^4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*\log(b*x + a) + 12*(b^4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*\log(x)}{(a^5*b^2*x^4 + 2*a^6*b*x^3 + a^7*x^2)}$$

Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.22

$$\int \frac{1}{x^3(a+bx)^3} dx = \frac{-a^3 + 4a^2bx + 18ab^2x^2 + 12b^3x^3}{2a^6x^2 + 4a^5bx^3 + 2a^4b^2x^4} + \frac{6b^2(\log(x) - \log(\frac{a}{b} + x))}{a^5}$$

input `integrate(1/x**3/(b*x+a)**3,x)`output `(-a**3 + 4*a**2*b*x + 18*a*b**2*x**2 + 12*b**3*x**3)/(2*a**6*x**2 + 4*a**5*b*x**3 + 2*a**4*b**2*x**4) + 6*b**2*(log(x) - log(a/b + x))/a**5`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.34

$$\int \frac{1}{x^3(a+bx)^3} dx = \frac{12b^3x^3 + 18ab^2x^2 + 4a^2bx - a^3}{2(a^4b^2x^4 + 2a^5bx^3 + a^6x^2)} - \frac{6b^2 \log(bx+a)}{a^5} + \frac{6b^2 \log(x)}{a^5}$$

input `integrate(1/x^3/(b*x+a)^3,x, algorithm="maxima")`output `1/2*(12*b^3*x^3 + 18*a*b^2*x^2 + 4*a^2*b*x - a^3)/(a^4*b^2*x^4 + 2*a^5*b*x^3 + a^6*x^2) - 6*b^2*log(b*x + a)/a^5 + 6*b^2*log(x)/a^5`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.14

$$\int \frac{1}{x^3(a+bx)^3} dx = -\frac{6b^2 \log(|bx+a|)}{a^5} + \frac{6b^2 \log(|x|)}{a^5} + \frac{12b^3x^3 + 18ab^2x^2 + 4a^2bx - a^3}{2(bx^2 + ax)^2 a^4}$$

input `integrate(1/x^3/(b*x+a)^3,x, algorithm="giac")`output `-6*b^2*log(abs(b*x + a))/a^5 + 6*b^2*log(abs(x))/a^5 + 1/2*(12*b^3*x^3 + 18*a*b^2*x^2 + 4*a^2*b*x - a^3)/((b*x^2 + a*x)^2*a^4)`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.23

$$\int \frac{1}{x^3(a+bx)^3} dx = \frac{\frac{9b^2x^2}{a^3} - \frac{1}{2a} + \frac{6b^3x^3}{a^4} + \frac{2bx}{a^2}}{a^2x^2 + 2abx^3 + b^2x^4} - \frac{12b^2 \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^5}$$

input `int(1/(x^3*(a + b*x)^3),x)`output `((9*b^2*x^2)/a^3 - 1/(2*a) + (6*b^3*x^3)/a^4 + (2*b*x)/a^2)/(a^2*x^2 + b^2*x^4 + 2*a*b*x^3) - (12*b^2*atanh((2*b*x)/a + 1))/a^5`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 138, normalized size of antiderivative = 2.16

$$\int \frac{1}{x^3(a+bx)^3} dx = \frac{-12 \log(bx+a) a^2 b^2 x^2 - 24 \log(bx+a) a b^3 x^3 - 12 \log(bx+a) b^4 x^4 + 12 \log(x) a^2 b^2 x^2 + 24 \log(x) a b^3 x^3}{2a^5 x^2 (b^2 x^2 + 2abx + a^2)}$$

input `int(1/x^3/(b*x+a)^3,x)`output `(- 12*log(a + b*x)*a**2*b**2*x**2 - 24*log(a + b*x)*a*b**3*x**3 - 12*log(a + b*x)*b**4*x**4 + 12*log(x)*a**2*b**2*x**2 + 24*log(x)*a*b**3*x**3 + 12*log(x)*b**4*x**4 - a**4 + 4*a**3*b*x + 12*a**2*b**2*x**2 - 6*b**4*x**4)/(2*a**5*x**2*(a**2 + 2*a*b*x + b**2*x**2))`

3.54 $\int \frac{1}{x(a+bx)^4} dx$

Optimal result	349
Mathematica [A] (verified)	349
Rubi [A] (verified)	350
Maple [A] (verified)	351
Fricas [B] (verification not implemented)	351
Sympy [A] (verification not implemented)	352
Maxima [A] (verification not implemented)	352
Giac [A] (verification not implemented)	352
Mupad [B] (verification not implemented)	353
Reduce [B] (verification not implemented)	353

Optimal result

Integrand size = 11, antiderivative size = 51

$$\int \frac{1}{x(a+bx)^4} dx = \frac{\frac{11}{6a} + \frac{5bx}{2a^2} + \frac{b^2x^2}{a^3}}{(a+bx)^3} - \frac{\log\left(\frac{a+bx}{x}\right)}{a^4}$$

output $(11/6/a+5/2*b*x/a^2+b^2*x^2/a^3)/(b*x+a)^3-\ln((b*x+a)/x)/a^4$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.94

$$\int \frac{1}{x(a+bx)^4} dx = \frac{\frac{a(11a^2+15abx+6b^2x^2)}{(a+bx)^3} + 6\log(x) - 6\log(a+bx)}{6a^4}$$

input `Integrate[1/(x*(a + b*x)^4),x]`

output $((a*(11*a^2 + 15*a*b*x + 6*b^2*x^2))/(a + b*x)^3 + 6*Log[x] - 6*Log[a + b*x])/ (6*a^4)$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.12, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(a+bx)^4} dx$$

↓ 54

$$\int \left(-\frac{b}{a^4(a+bx)} + \frac{1}{a^4x} - \frac{b}{a^3(a+bx)^2} - \frac{b}{a^2(a+bx)^3} - \frac{b}{a(a+bx)^4} \right) dx$$

↓ 2009

$$-\frac{\log(a+bx)}{a^4} + \frac{\log(x)}{a^4} + \frac{1}{a^3(a+bx)} + \frac{1}{2a^2(a+bx)^2} + \frac{1}{3a(a+bx)^3}$$

input `Int[1/(x*(a + b*x)^4),x]`

output `1/(3*a*(a + b*x)^3) + 1/(2*a^2*(a + b*x)^2) + 1/(a^3*(a + b*x)) + Log[x]/a^4 - Log[a + b*x]/a^4`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.02

method	result
risch	$\frac{\frac{11}{6a} + \frac{5bx}{2a^2} + \frac{b^2x^2}{a^3}}{(bx+a)^3} + \frac{\ln(-x)}{a^4} - \frac{\ln(bx+a)}{a^4}$
default	$-\frac{\ln(bx+a)}{a^4} + \frac{1}{a^3(bx+a)} + \frac{1}{2a^2(bx+a)^2} + \frac{1}{3a(bx+a)^3} + \frac{\ln(x)}{a^4}$
norman	$\frac{-\frac{3bx}{a^2} - \frac{9b^2x^2}{2a^3} - \frac{11b^3x^3}{6a^4}}{(bx+a)^3} + \frac{\ln(x)}{a^4} - \frac{\ln(bx+a)}{a^4}$
parallelrisch	$\frac{6 \ln(x)x^3b^3 - 6 \ln(bx+a)x^3b^3 + 18 \ln(x)x^2ab^2 - 18 \ln(bx+a)x^2ab^2 - 11b^3x^3 + 18 \ln(x)xa^2b - 18 \ln(bx+a)xa^2b - 27ab^2x^2 + 6 \ln(x)a^2b^2}{6a^4(bx+a)^3}$

input `int(1/x/(b*x+a)^4,x,method=_RETURNVERBOSE)`output `(11/6/a+5/2*b*x/a^2+b^2*x^2/a^3)/(b*x+a)^3+1/a^4*ln(-x)-1/a^4*ln(b*x+a)`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(49) = 98.

Time = 0.15 (sec) , antiderivative size = 124, normalized size of antiderivative = 2.43

$$\int \frac{1}{x(a+bx)^4} dx$$

$$= \frac{6ab^2x^2 + 15a^2bx + 11a^3 - 6(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3) \log(bx+a) + 6(b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3) \log(x)}{6(a^4b^3x^3 + 3a^5b^2x^2 + 3a^6bx + a^7)}$$

input `integrate(1/x/(b*x+a)^4,x, algorithm="fricas")`output `1/6*(6*a*b^2*x^2 + 15*a^2*b*x + 11*a^3 - 6*(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*log(b*x + a) + 6*(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*log(x))/(a^4*b^3*x^3 + 3*a^5*b^2*x^2 + 3*a^6*b*x + a^7)`

Sympy [A] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.37

$$\int \frac{1}{x(a+bx)^4} dx = \frac{11a^2 + 15abx + 6b^2x^2}{6a^6 + 18a^5bx + 18a^4b^2x^2 + 6a^3b^3x^3} + \frac{\log(x) - \log\left(\frac{a}{b} + x\right)}{a^4}$$

input `integrate(1/x/(b*x+a)**4,x)`output `(11*a**2 + 15*a*b*x + 6*b**2*x**2)/(6*a**6 + 18*a**5*b*x + 18*a**4*b**2*x**2 + 6*a**3*b**3*x**3) + (log(x) - log(a/b + x))/a**4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.43

$$\int \frac{1}{x(a+bx)^4} dx = \frac{6b^2x^2 + 15abx + 11a^2}{6(a^3b^3x^3 + 3a^4b^2x^2 + 3a^5bx + a^6)} - \frac{\log(bx+a)}{a^4} + \frac{\log(x)}{a^4}$$

input `integrate(1/x/(b*x+a)^4,x, algorithm="maxima")`output `1/6*(6*b^2*x^2 + 15*a*b*x + 11*a^2)/(a^3*b^3*x^3 + 3*a^4*b^2*x^2 + 3*a^5*b*x + a^6) - log(b*x + a)/a^4 + log(x)/a^4`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.06

$$\int \frac{1}{x(a+bx)^4} dx = -\frac{\log(|bx+a|)}{a^4} + \frac{\log(|x|)}{a^4} + \frac{6ab^2x^2 + 15a^2bx + 11a^3}{6(bx+a)^3a^4}$$

input `integrate(1/x/(b*x+a)^4,x, algorithm="giac")`output `-log(abs(b*x + a))/a^4 + log(abs(x))/a^4 + 1/6*(6*a*b^2*x^2 + 15*a^2*b*x + 11*a^3)/((b*x + a)^3*a^4)`

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.18

$$\int \frac{1}{x(a+bx)^4} dx = \frac{\frac{1}{a^2+bx} - \frac{\ln\left(\frac{a+bx}{x}\right)}{a^2}}{a} + \frac{1}{2a(a+bx)^2} + \frac{1}{3a(a+bx)^3}$$

input `int(1/(x*(a + b*x)^4),x)`output `((1/(a^2 + a*b*x) - log((a + b*x)/x)/a^2)/a + 1/(2*a*(a + b*x)^2))/a + 1/(3*a*(a + b*x)^3)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 145, normalized size of antiderivative = 2.84

$$\int \frac{1}{x(a+bx)^4} dx = \frac{-6 \log(bx+a) a^3 - 18 \log(bx+a) a^2 bx - 18 \log(bx+a) a b^2 x^2 - 6 \log(bx+a) b^3 x^3 + 6 \log(x) a^3 + 18 \log(x) a^2 bx + 18 \log(x) a b^2 x^2 + 6 \log(x) b^3 x^3 + 9 a^3 + 9 a^2 b x - 2 b^3 x^3}{6 a^4 (b^3 x^3 + 3 a b^2 x^2 + 3 a^2 b x + a^3)}$$

input `int(1/x/(b*x+a)^4,x)`output `(- 6*log(a + b*x)*a**3 - 18*log(a + b*x)*a**2*b*x - 18*log(a + b*x)*a*b**2*x**2 - 6*log(a + b*x)*b**3*x**3 + 6*log(x)*a**3 + 18*log(x)*a**2*b*x + 18*log(x)*a*b**2*x**2 + 6*log(x)*b**3*x**3 + 9*a**3 + 9*a**2*b*x - 2*b**3*x**3)/(6*a**4*(a**3 + 3*a**2*b*x + 3*a*b**2*x**2 + b**3*x**3))`

3.55 $\int \frac{1}{x^2(a+bx)^4} dx$

Optimal result	354
Mathematica [A] (verified)	354
Rubi [A] (verified)	355
Maple [A] (verified)	356
Fricas [B] (verification not implemented)	356
Sympy [A] (verification not implemented)	357
Maxima [A] (verification not implemented)	357
Giac [A] (verification not implemented)	357
Mupad [B] (verification not implemented)	358
Reduce [B] (verification not implemented)	358

Optimal result

Integrand size = 11, antiderivative size = 62

$$\int \frac{1}{x^2(a+bx)^4} dx = \frac{-\frac{22b}{3a^2} - \frac{1}{ax} - \frac{10b^2x}{a^3} - \frac{4b^3x^2}{a^4}}{(a+bx)^3} + \frac{4b \log\left(\frac{a+bx}{x}\right)}{a^5}$$

output (-22/3*b/a^2-1/a/x-10*b^2*x/a^3-4*b^3*x^2/a^4)/(b*x+a)^3+4*b*ln((b*x+a)/x)/a^5

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.03

$$\int \frac{1}{x^2(a+bx)^4} dx = -\frac{\frac{a(3a^3+22a^2bx+30ab^2x^2+12b^3x^3)}{x(a+bx)^3} + 12b \log(x) - 12b \log(a+bx)}{3a^5}$$

input Integrate[1/(x^2*(a + b*x)^4),x]

output -1/3*((a*(3*a^3 + 22*a^2*b*x + 30*a*b^2*x^2 + 12*b^3*x^3))/(x*(a + b*x)^3) + 12*b*Log[x] - 12*b*Log[a + b*x])/a^5

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.13, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2(a+bx)^4} dx$$

↓ 54

$$\int \left(\frac{4b^2}{a^5(a+bx)} - \frac{4b}{a^5x} + \frac{3b^2}{a^4(a+bx)^2} + \frac{1}{a^4x^2} + \frac{2b^2}{a^3(a+bx)^3} + \frac{b^2}{a^2(a+bx)^4} \right) dx$$

↓ 2009

$$-\frac{4b \log(x)}{a^5} + \frac{4b \log(a+bx)}{a^5} - \frac{3b}{a^4(a+bx)} - \frac{1}{a^4x} - \frac{b}{a^3(a+bx)^2} - \frac{b}{3a^2(a+bx)^3}$$

input `Int[1/(x^2*(a + b*x)^4),x]`

output `-(1/(a^4*x)) - b/(3*a^2*(a + b*x)^3) - b/(a^3*(a + b*x)^2) - (3*b)/(a^4*(a + b*x)) - (4*b*Log[x])/a^5 + (4*b*Log[a + b*x])/a^5`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.11

method	result
default	$-\frac{b}{3a^2(bx+a)^3} + \frac{4b \ln(bx+a)}{a^5} - \frac{3b}{a^4(bx+a)} - \frac{b}{a^3(bx+a)^2} - \frac{1}{a^4x} - \frac{4b \ln(x)}{a^5}$
risch	$-\frac{\frac{4b^3x^3}{a^4} - \frac{10b^2x^2}{a^3} - \frac{22bx}{3a^2} - \frac{1}{a}}{x(bx+a)^3} + \frac{4b \ln(-bx-a)}{a^5} - \frac{4b \ln(x)}{a^5}$
norman	$-\frac{\frac{1}{a} + \frac{12b^2x^2}{a^3} + \frac{18b^3x^3}{a^4} + \frac{22b^4x^4}{3a^5}}{x(bx+a)^3} - \frac{4b \ln(x)}{a^5} + \frac{4b \ln(bx+a)}{a^5}$
parallelrisch	$-\frac{12 \ln(x)x^4b^4 - 12 \ln(bx+a)x^4b^4 + 36 \ln(x)x^3ab^3 - 36 \ln(bx+a)x^3ab^3 - 22b^4x^4 + 36 \ln(x)x^2a^2b^2 - 36 \ln(bx+a)x^2a^2b^2 - 54ab^3}{3a^5x(bx+a)^3}$

input `int(1/x^2/(b*x+a)^4,x,method=_RETURNVERBOSE)`output `-1/3*b/a^2/(b*x+a)^3+4/a^5*b*ln(b*x+a)-3*b/a^4/(b*x+a)-b/a^3/(b*x+a)^2-1/a^4/x-4/a^5*b*ln(x)`**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(61) = 122.

Time = 0.10 (sec) , antiderivative size = 153, normalized size of antiderivative = 2.47

$$\int \frac{1}{x^2(a+bx)^4} dx = \frac{12ab^3x^3 + 30a^2b^2x^2 + 22a^3bx + 3a^4 - 12(b^4x^4 + 3ab^3x^3 + 3a^2b^2x^2 + a^3bx) \log(bx+a) + 12(b^4x^4 - 3(a^5b^3x^4 + 3a^6b^2x^3 + 3a^7bx^2 + a^8x))}{3(a^5b^3x^4 + 3a^6b^2x^3 + 3a^7bx^2 + a^8x)}$$

input `integrate(1/x^2/(b*x+a)^4,x, algorithm="fricas")`output `-1/3*(12*a*b^3*x^3 + 30*a^2*b^2*x^2 + 22*a^3*b*x + 3*a^4 - 12*(b^4*x^4 + 3*a*b^3*x^3 + 3*a^2*b^2*x^2 + a^3*b*x)*log(b*x + a) + 12*(b^4*x^4 + 3*a*b^3*x^3 + 3*a^2*b^2*x^2 + a^3*b*x)*log(x))/(a^5*b^3*x^4 + 3*a^6*b^2*x^3 + 3*a^7*b*x^2 + a^8*x)`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.45

$$\int \frac{1}{x^2(a+bx)^4} dx = \frac{-3a^3 - 22a^2bx - 30ab^2x^2 - 12b^3x^3}{3a^7x + 9a^6bx^2 + 9a^5b^2x^3 + 3a^4b^3x^4} + \frac{4b(-\log(x) + \log(\frac{a}{b} + x))}{a^5}$$

input `integrate(1/x**2/(b*x+a)**4,x)`output `(-3*a**3 - 22*a**2*b*x - 30*a*b**2*x**2 - 12*b**3*x**3)/(3*a**7*x + 9*a**6*b*x**2 + 9*a**5*b**2*x**3 + 3*a**4*b**3*x**4) + 4*b*(-log(x) + log(a/b + x))/a**5`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.47

$$\int \frac{1}{x^2(a+bx)^4} dx = -\frac{12b^3x^3 + 30ab^2x^2 + 22a^2bx + 3a^3}{3(a^4b^3x^4 + 3a^5b^2x^3 + 3a^6bx^2 + a^7x)} + \frac{4b \log(bx+a)}{a^5} - \frac{4b \log(x)}{a^5}$$

input `integrate(1/x^2/(b*x+a)^4,x, algorithm="maxima")`output `-1/3*(12*b^3*x^3 + 30*a*b^2*x^2 + 22*a^2*b*x + 3*a^3)/(a^4*b^3*x^4 + 3*a^5*b^2*x^3 + 3*a^6*b*x^2 + a^7*x) + 4*b*log(b*x + a)/a^5 - 4*b*log(x)/a^5`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.15

$$\int \frac{1}{x^2(a+bx)^4} dx = \frac{4b \log(|bx+a|)}{a^5} - \frac{4b \log(|x|)}{a^5} - \frac{12ab^3x^3 + 30a^2b^2x^2 + 22a^3bx + 3a^4}{3(bx+a)^3a^5x}$$

input `integrate(1/x^2/(b*x+a)^4,x, algorithm="giac")`

output `4*b*log(abs(b*x + a))/a^5 - 4*b*log(abs(x))/a^5 - 1/3*(12*a*b^3*x^3 + 30*a^2*b^2*x^2 + 22*a^3*b*x + 3*a^4)/((b*x + a)^3*a^5*x)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.37

$$\int \frac{1}{x^2(a+bx)^4} dx = \frac{8b \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^5} - \frac{\frac{1}{a} + \frac{10b^2x^2}{a^3} + \frac{4b^3x^3}{a^4} + \frac{22bx}{3a^2}}{a^3x + 3a^2bx^2 + 3ab^2x^3 + b^3x^4}$$

input `int(1/(x^2*(a + b*x)^4),x)`

output `(8*b*atanh((2*b*x)/a + 1))/a^5 - (1/a + (10*b^2*x^2)/a^3 + (4*b^3*x^3)/a^4 + (22*b*x)/(3*a^2))/(a^3*x + b^3*x^4 + 3*a^2*b*x^2 + 3*a*b^2*x^3)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.76

$$\int \frac{1}{x^2(a+bx)^4} dx = \frac{12 \log(bx+a) a^3 bx + 36 \log(bx+a) a^2 b^2 x^2 + 36 \log(bx+a) a b^3 x^3 + 12 \log(bx+a) b^4 x^4 - 12 \log(x) a^3 b}{3a^5x(b^3x^3 + 3ab^2x^2 + \dots)}$$

input `int(1/x^2/(b*x+a)^4,x)`

output `(12*log(a + b*x)*a**3*b*x + 36*log(a + b*x)*a**2*b**2*x**2 + 36*log(a + b*x)*a*b**3*x**3 + 12*log(a + b*x)*b**4*x**4 - 12*log(x)*a**3*b*x - 36*log(x)*a**2*b**2*x**2 - 36*log(x)*a*b**3*x**3 - 12*log(x)*b**4*x**4 - 3*a**4 - 18*a**3*b*x - 18*a**2*b**2*x**2 + 4*b**4*x**4)/(3*a**5*x*(a**3 + 3*a**2*b*x + 3*a*b**2*x**2 + b**3*x**3))`

3.56 $\int \frac{1}{x^3(a+bx)^4} dx$

Optimal result	359
Mathematica [A] (verified)	359
Rubi [A] (verified)	360
Maple [A] (verified)	361
Fricas [B] (verification not implemented)	361
Sympy [A] (verification not implemented)	362
Maxima [A] (verification not implemented)	362
Giac [A] (verification not implemented)	363
Mupad [B] (verification not implemented)	363
Reduce [B] (verification not implemented)	364

Optimal result

Integrand size = 11, antiderivative size = 79

$$\int \frac{1}{x^3(a+bx)^4} dx = \frac{\frac{55b^2}{3a^3} - \frac{1}{2ax^2} + \frac{5b}{2a^2x} + \frac{25b^3x}{a^4} + \frac{10b^4x^2}{a^5}}{(a+bx)^3} - \frac{10b^2 \log\left(\frac{a+bx}{x}\right)}{a^6}$$

output

```
(55/3*b^2/a^3-1/2/a/x^2+5/2*b/a^2/x+25*b^3*x/a^4+10*b^4*x^2/a^5)/(b*x+a)^3-10*b^2*ln((b*x+a)/x)/a^6
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^3(a+bx)^4} dx = \frac{\frac{a(-3a^4+15a^3bx+110a^2b^2x^2+150ab^3x^3+60b^4x^4)}{x^2(a+bx)^3} + 60b^2 \log(x) - 60b^2 \log(a+bx)}{6a^6}$$

input

```
Integrate[1/(x^3*(a + b*x)^4),x]
```

output

```
((a*(-3*a^4 + 15*a^3*b*x + 110*a^2*b^2*x^2 + 150*a*b^3*x^3 + 60*b^4*x^4))/(x^2*(a + b*x)^3) + 60*b^2*Log[x] - 60*b^2*Log[a + b*x])/(6*a^6)
```


Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.18, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3(a+bx)^4} dx$$

↓ 54

$$\int \left(-\frac{10b^3}{a^6(a+bx)} + \frac{10b^2}{a^6x} - \frac{6b^3}{a^5(a+bx)^2} - \frac{4b}{a^5x^2} - \frac{3b^3}{a^4(a+bx)^3} + \frac{1}{a^4x^3} - \frac{b^3}{a^3(a+bx)^4} \right) dx$$

↓ 2009

$$\frac{10b^2 \log(x)}{a^6} - \frac{10b^2 \log(a+bx)}{a^6} + \frac{6b^2}{a^5(a+bx)} + \frac{4b}{a^5x} + \frac{3b^2}{2a^4(a+bx)^2} - \frac{1}{2a^4x^2} + \frac{b^2}{3a^3(a+bx)^3}$$

input `Int[1/(x^3*(a + b*x)^4),x]`

output `-1/2*1/(a^4*x^2) + (4*b)/(a^5*x) + b^2/(3*a^3*(a + b*x)^3) + (3*b^2)/(2*a^4*(a + b*x)^2) + (6*b^2)/(a^5*(a + b*x)) + (10*b^2*Log[x])/a^6 - (10*b^2*Log[a + b*x])/a^6`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.05

method	result
norman	$-\frac{1}{2a} + \frac{5bx}{2a^2} - \frac{30b^3x^3}{a^4} - \frac{45b^4x^4}{a^5} - \frac{55b^5x^5}{3a^6} + \frac{10b^2 \ln(x)}{a^6} - \frac{10b^2 \ln(bx+a)}{a^6}$
risch	$\frac{10b^4x^4}{a^5} + \frac{25b^3x^3}{a^4} + \frac{55b^2x^2}{3a^3} + \frac{5bx}{2a^2} - \frac{1}{2a} + \frac{10b^2 \ln(-x)}{a^6} - \frac{10b^2 \ln(bx+a)}{a^6}$
default	$-\frac{10b^2 \ln(bx+a)}{a^6} + \frac{6b^2}{a^5(bx+a)} + \frac{3b^2}{2a^4(bx+a)^2} + \frac{b^2}{3a^3(bx+a)^3} - \frac{1}{2a^4x^2} + \frac{10b^2 \ln(x)}{a^6} + \frac{4b}{a^5x}$
parallelrisch	$\frac{60 \ln(x)x^5b^5 - 60 \ln(bx+a)x^5b^5 + 180 \ln(x)x^4ab^4 - 180 \ln(bx+a)x^4ab^4 - 110b^5x^5 + 180 \ln(x)x^3a^2b^3 - 180 \ln(bx+a)x^3a^2b^3 - 27}{6a^6x^2(bx+a)^3}$

input `int(1/x^3/(b*x+a)^4,x,method=_RETURNVERBOSE)`output
$$\frac{(-1/2/a+5/2*b*x/a^2-30*b^3*x^3/a^4-45*b^4/a^5*x^4-55/3*b^5/a^6*x^5)/x^2/(b*x+a)^3+10/a^6*b^2*\ln(x)-10/a^6*b^2*\ln(b*x+a)}$$
Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 174 vs. 2(74) = 148.

Time = 0.09 (sec) , antiderivative size = 174, normalized size of antiderivative = 2.20

$$\int \frac{1}{x^3(a+bx)^4} dx = \frac{60ab^4x^4 + 150a^2b^3x^3 + 110a^3b^2x^2 + 15a^4bx - 3a^5 - 60(b^5x^5 + 3ab^4x^4 + 3a^2b^3x^3 + a^3b^2x^2) \log(bx+a)}{6(a^6b^3x^5 + 3a^7b^2x^4 + 3a^8bx^3 + a^9x^2)}$$

input `integrate(1/x^3/(b*x+a)^4,x, algorithm="fricas")`output
$$\frac{1/6*(60*a*b^4*x^4 + 150*a^2*b^3*x^3 + 110*a^3*b^2*x^2 + 15*a^4*b*x - 3*a^5 - 60*(b^5*x^5 + 3*a*b^4*x^4 + 3*a^2*b^3*x^3 + a^3*b^2*x^2)*\log(b*x + a) + 60*(b^5*x^5 + 3*a*b^4*x^4 + 3*a^2*b^3*x^3 + a^3*b^2*x^2)*\log(x)}{(a^6*b^3*x^5 + 3*a^7*b^2*x^4 + 3*a^8*b*x^3 + a^9*x^2)}$$

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.32

$$\int \frac{1}{x^3(a+bx)^4} dx = \frac{-3a^4 + 15a^3bx + 110a^2b^2x^2 + 150ab^3x^3 + 60b^4x^4}{6a^8x^2 + 18a^7bx^3 + 18a^6b^2x^4 + 6a^5b^3x^5} + \frac{10b^2(\log(x) - \log(\frac{a}{b} + x))}{a^6}$$

input `integrate(1/x**3/(b*x+a)**4,x)`output `(-3*a**4 + 15*a**3*b*x + 110*a**2*b**2*x**2 + 150*a*b**3*x**3 + 60*b**4*x**4)/(6*a**8*x**2 + 18*a**7*b*x**3 + 18*a**6*b**2*x**4 + 6*a**5*b**3*x**5) + 10*b**2*(log(x) - log(a/b + x))/a**6`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.37

$$\int \frac{1}{x^3(a+bx)^4} dx = \frac{60b^4x^4 + 150ab^3x^3 + 110a^2b^2x^2 + 15a^3bx - 3a^4}{6(a^5b^3x^5 + 3a^6b^2x^4 + 3a^7bx^3 + a^8x^2)} - \frac{10b^2 \log(bx+a)}{a^6} + \frac{10b^2 \log(x)}{a^6}$$

input `integrate(1/x^3/(b*x+a)^4,x, algorithm="maxima")`output `1/6*(60*b^4*x^4 + 150*a*b^3*x^3 + 110*a^2*b^2*x^2 + 15*a^3*b*x - 3*a^4)/(a^5*b^3*x^5 + 3*a^6*b^2*x^4 + 3*a^7*b*x^3 + a^8*x^2) - 10*b^2*log(b*x + a)/a^6 + 10*b^2*log(x)/a^6`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.09

$$\int \frac{1}{x^3(a+bx)^4} dx = -\frac{10b^2 \log(|bx+a|)}{a^6} + \frac{10b^2 \log(|x|)}{a^6} + \frac{60ab^4x^4 + 150a^2b^3x^3 + 110a^3b^2x^2 + 15a^4bx - 3a^5}{6(bx+a)^3a^6x^2}$$

input `integrate(1/x^3/(b*x+a)^4,x, algorithm="giac")`

output `-10*b^2*log(abs(b*x + a))/a^6 + 10*b^2*log(abs(x))/a^6 + 1/6*(60*a*b^4*x^4 + 150*a^2*b^3*x^3 + 110*a^3*b^2*x^2 + 15*a^4*b*x - 3*a^5)/((b*x + a)^3*a^6*x^2)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.28

$$\int \frac{1}{x^3(a+bx)^4} dx = \frac{55b^2x^2}{3a^3} - \frac{1}{2a} + \frac{25b^3x^3}{a^4} + \frac{10b^4x^4}{a^5} + \frac{5bx}{2a^2} - \frac{20b^2 \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^6}$$

input `int(1/(x^3*(a + b*x)^4),x)`

output `((55*b^2*x^2)/(3*a^3) - 1/(2*a) + (25*b^3*x^3)/a^4 + (10*b^4*x^4)/a^5 + (5*b*x)/(2*a^2))/(a^3*x^2 + b^3*x^5 + 3*a^2*b*x^3 + 3*a*b^2*x^4) - (20*b^2*a*tanh((2*b*x)/a + 1))/a^6`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.41

$$\int \frac{1}{x^3(a+bx)^4} dx$$

$$= \frac{-60 \log(bx+a) a^3 b^2 x^2 - 180 \log(bx+a) a^2 b^3 x^3 - 180 \log(bx+a) a b^4 x^4 - 60 \log(bx+a) b^5 x^5 + 60 \log(x) a^3 b^2 x^2 + 180 \log(x) a^2 b^3 x^3 + 180 \log(x) a b^4 x^4 + 60 \log(x) b^5 x^5 - 3 a^5 + 15 a^4 b x + 90 a^3 b^2 x^2 + 90 a^2 b^3 x^3 - 20 b^5 x^5}{6 a^6 x^2 (b^3 x^3 - \dots)}$$

input `int(1/x^3/(b*x+a)^4,x)`output `(- 60*log(a + b*x)*a**3*b**2*x**2 - 180*log(a + b*x)*a**2*b**3*x**3 - 180*log(a + b*x)*a*b**4*x**4 - 60*log(a + b*x)*b**5*x**5 + 60*log(x)*a**3*b**2*x**2 + 180*log(x)*a**2*b**3*x**3 + 180*log(x)*a*b**4*x**4 + 60*log(x)*b**5*x**5 - 3*a**5 + 15*a**4*b*x + 90*a**3*b**2*x**2 + 90*a**2*b**3*x**3 - 20*b**5*x**5)/(6*a**6*x**2*(a**3 + 3*a**2*b*x + 3*a*b**2*x**2 + b**3*x**3))`

3.57 $\int \frac{1}{x(a+bx)^5} dx$

Optimal result	365
Mathematica [A] (verified)	365
Rubi [A] (verified)	366
Maple [A] (verified)	367
Fricas [B] (verification not implemented)	367
Sympy [A] (verification not implemented)	368
Maxima [A] (verification not implemented)	368
Giac [A] (verification not implemented)	369
Mupad [B] (verification not implemented)	369
Reduce [B] (verification not implemented)	369

Optimal result

Integrand size = 11, antiderivative size = 64

$$\int \frac{1}{x(a+bx)^5} dx = \frac{\frac{25}{12a} + \frac{13bx}{3a^2} + \frac{7b^2x^2}{2a^3} + \frac{b^3x^3}{a^4}}{(a+bx)^4} - \frac{\log\left(\frac{a+bx}{x}\right)}{a^5}$$

output (25/12/a+13/3*b*x/a^2+7/2*b^2*x^2/a^3+b^3*x^3/a^4)/(b*x+a)^4-ln((b*x+a)/x)/a^5

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.92

$$\int \frac{1}{x(a+bx)^5} dx = \frac{\frac{a(25a^3+52a^2bx+42ab^2x^2+12b^3x^3)}{(a+bx)^4} + 12\log(x) - 12\log(a+bx)}{12a^5}$$

input Integrate[1/(x*(a + b*x)^5),x]

output ((a*(25*a^3 + 52*a^2*b*x + 42*a*b^2*x^2 + 12*b^3*x^3))/(a + b*x)^4 + 12*Log[x] - 12*Log[a + b*x])/(12*a^5)

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.11, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(a+bx)^5} dx$$

↓ 54

$$\int \left(-\frac{b}{a^5(a+bx)} + \frac{1}{a^5x} - \frac{b}{a^4(a+bx)^2} - \frac{b}{a^3(a+bx)^3} - \frac{b}{a^2(a+bx)^4} - \frac{b}{a(a+bx)^5} \right) dx$$

↓ 2009

$$-\frac{\log(a+bx)}{a^5} + \frac{\log(x)}{a^5} + \frac{1}{a^4(a+bx)} + \frac{1}{2a^3(a+bx)^2} + \frac{1}{3a^2(a+bx)^3} + \frac{1}{4a(a+bx)^4}$$

input `Int[1/(x*(a + b*x)^5), x]`

output `1/(4*a*(a + b*x)^4) + 1/(3*a^2*(a + b*x)^3) + 1/(2*a^3*(a + b*x)^2) + 1/(a^4*(a + b*x)) + Log[x]/a^5 - Log[a + b*x]/a^5`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.98

method	result
risch	$\frac{25}{12a} + \frac{13bx}{3a^2} + \frac{7b^2x^2}{2a^3} + \frac{b^3x^3}{a^4} + \frac{\ln(-x)}{a^5} - \frac{\ln(bx+a)}{a^5}$
default	$-\frac{\ln(bx+a)}{a^5} + \frac{1}{a^4(bx+a)} + \frac{1}{2a^3(bx+a)^2} + \frac{1}{3a^2(bx+a)^3} + \frac{1}{4a(bx+a)^4} + \frac{\ln(x)}{a^5}$
norman	$-\frac{4bx}{a^2} - \frac{9b^2x^2}{a^3} - \frac{22b^3x^3}{3a^4} - \frac{25b^4x^4}{12a^5} + \frac{\ln(x)}{a^5} - \frac{\ln(bx+a)}{a^5}$
parallelrisch	$\frac{12 \ln(x)x^4b^4 - 12 \ln(bx+a)x^4b^4 + 48 \ln(x)x^3ab^3 - 48 \ln(bx+a)x^3ab^3 - 25b^4x^4 + 72 \ln(x)x^2a^2b^2 - 72 \ln(bx+a)x^2a^2b^2 - 88ab^3x^3}{12a^5(bx+a)^4}$

input `int(1/x/(b*x+a)^5,x,method=_RETURNVERBOSE)`output $(25/12/a+13/3*b*x/a^2+7/2*b^2*x^2/a^3+b^3*x^3/a^4)/(b*x+a)^4+1/a^5*\ln(-x)-1/a^5*\ln(b*x+a)$ **Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(60) = 120.

Time = 0.09 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.62

$$\int \frac{1}{x(a+bx)^5} dx = \frac{12ab^3x^3 + 42a^2b^2x^2 + 52a^3bx + 25a^4 - 12(b^4x^4 + 4ab^3x^3 + 6a^2b^2x^2 + 4a^3bx + a^4) \log(bx+a) + 12(a^5b^4x^4 + 4a^6b^3x^3 + 6a^7b^2x^2 + 4a^8bx + a^9)}{12(a^5b^4x^4 + 4a^6b^3x^3 + 6a^7b^2x^2 + 4a^8bx + a^9)}$$

input `integrate(1/x/(b*x+a)^5,x, algorithm="fricas")`output $1/12*(12*a*b^3*x^3 + 42*a^2*b^2*x^2 + 52*a^3*b*x + 25*a^4 - 12*(b^4*x^4 + 4*a*b^3*x^3 + 6*a^2*b^2*x^2 + 4*a^3*b*x + a^4)*\log(b*x + a) + 12*(b^4*x^4 + 4*a*b^3*x^3 + 6*a^2*b^2*x^2 + 4*a^3*b*x + a^4)*\log(x))/(a^5*b^4*x^4 + 4*a^6*b^3*x^3 + 6*a^7*b^2*x^2 + 4*a^8*b*x + a^9)$

Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.47

$$\int \frac{1}{x(a+bx)^5} dx = \frac{25a^3 + 52a^2bx + 42ab^2x^2 + 12b^3x^3}{12a^8 + 48a^7bx + 72a^6b^2x^2 + 48a^5b^3x^3 + 12a^4b^4x^4} + \frac{\log(x) - \log\left(\frac{a}{b} + x\right)}{a^5}$$

input `integrate(1/x/(b*x+a)**5,x)`output `(25*a**3 + 52*a**2*b*x + 42*a*b**2*x**2 + 12*b**3*x**3)/(12*a**8 + 48*a**7*b*x + 72*a**6*b**2*x**2 + 48*a**5*b**3*x**3 + 12*a**4*b**4*x**4) + (log(x) - log(a/b + x))/a**5`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.48

$$\int \frac{1}{x(a+bx)^5} dx = \frac{12b^3x^3 + 42ab^2x^2 + 52a^2bx + 25a^3}{12(a^4b^4x^4 + 4a^5b^3x^3 + 6a^6b^2x^2 + 4a^7bx + a^8)} - \frac{\log(bx+a)}{a^5} + \frac{\log(x)}{a^5}$$

input `integrate(1/x/(b*x+a)^5,x, algorithm="maxima")`output `1/12*(12*b^3*x^3 + 42*a*b^2*x^2 + 52*a^2*b*x + 25*a^3)/(a^4*b^4*x^4 + 4*a^5*b^3*x^3 + 6*a^6*b^2*x^2 + 4*a^7*b*x + a^8) - log(b*x + a)/a^5 + log(x)/a^5`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.39

$$\int \frac{1}{x(a+bx)^5} dx = \frac{1}{12} b \left(\frac{12 \log \left(\left| -\frac{a}{bx+a} + 1 \right| \right)}{a^5 b} + \frac{\frac{12 b^3}{bx+a} + \frac{6 ab^3}{(bx+a)^2} + \frac{4 a^2 b^3}{(bx+a)^3} + \frac{3 a^3 b^3}{(bx+a)^4}}{a^4 b^4} \right)$$

input `integrate(1/x/(b*x+a)^5,x, algorithm="giac")`

output `1/12*b*(12*log(abs(-a/(b*x + a) + 1))/(a^5*b) + (12*b^3/(b*x + a) + 6*a*b^3/(b*x + a)^2 + 4*a^2*b^3/(b*x + a)^3 + 3*a^3*b^3/(b*x + a)^4)/(a^4*b^4))`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.20

$$\int \frac{1}{x(a+bx)^5} dx = \frac{\frac{\frac{1}{a^2+bx} - \frac{\ln\left(\frac{a+bx}{x}\right)}{a^2}}{a} + \frac{1}{2a(a+bx)^2}}{a} + \frac{1}{3a(a+bx)^3} + \frac{1}{4a(a+bx)^4}$$

input `int(1/(x*(a + b*x)^5),x)`

output `((((1/(a^2 + a*b*x) - log((a + b*x)/x)/a^2)/a + 1/(2*a*(a + b*x)^2))/a + 1/(3*a*(a + b*x)^3))/a + 1/(4*a*(a + b*x)^4)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 197, normalized size of antiderivative = 3.08

$$\int \frac{1}{x(a+bx)^5} dx = \frac{-12 \log(bx+a) a^4 - 48 \log(bx+a) a^3 bx - 72 \log(bx+a) a^2 b^2 x^2 - 48 \log(bx+a) a b^3 x^3 - 12 \log(bx+a) b^4 x^4}{12 a^5 (b^4 x^4)}$$

input `int(1/x/(b*x+a)^5,x)`

output `(- 12*log(a + b*x)*a**4 - 48*log(a + b*x)*a**3*b*x - 72*log(a + b*x)*a**2
*b**2*x**2 - 48*log(a + b*x)*a*b**3*x**3 - 12*log(a + b*x)*b**4*x**4 + 12*
log(x)*a**4 + 48*log(x)*a**3*b*x + 72*log(x)*a**2*b**2*x**2 + 48*log(x)*a*
b**3*x**3 + 12*log(x)*b**4*x**4 + 22*a**4 + 40*a**3*b*x + 24*a**2*b**2*x**
2 - 3*b**4*x**4)/(12*a**5*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2 + 4*a*b**3
*x**3 + b**4*x**4))`

3.58 $\int \frac{1}{x^2(a+bx)^5} dx$

Optimal result	371
Mathematica [A] (verified)	371
Rubi [A] (verified)	372
Maple [A] (verified)	373
Fricas [B] (verification not implemented)	373
Sympy [A] (verification not implemented)	374
Maxima [A] (verification not implemented)	374
Giac [A] (verification not implemented)	375
Mupad [B] (verification not implemented)	375
Reduce [B] (verification not implemented)	376

Optimal result

Integrand size = 11, antiderivative size = 77

$$\int \frac{1}{x^2(a+bx)^5} dx = \frac{-\frac{125b}{12a^2} - \frac{1}{ax} - \frac{65b^2x}{3a^3} - \frac{35b^3x^2}{2a^4} - \frac{5b^4x^3}{a^5}}{(a+bx)^4} + \frac{5b \log\left(\frac{a+bx}{x}\right)}{a^6}$$

output

```
(-125/12*b/a^2-1/a/x-65/3*b^2*x/a^3-35/2*b^3*x^2/a^4-5*b^4*x^3/a^5)/(b*x+a
)^4+5*b*ln((b*x+a)/x)/a^6
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.97

$$\int \frac{1}{x^2(a+bx)^5} dx = -\frac{a(12a^4+125a^3bx+260a^2b^2x^2+210ab^3x^3+60b^4x^4)}{12a^6x(a+bx)^4} + 60b \log(x) - 60b \log(a+bx)$$

input

```
Integrate[1/(x^2*(a + b*x)^5),x]
```

output

```
-1/12*((a*(12*a^4 + 125*a^3*b*x + 260*a^2*b^2*x^2 + 210*a*b^3*x^3 + 60*b^4
*x^4))/(x*(a + b*x)^4) + 60*b*Log[x] - 60*b*Log[a + b*x])/a^6
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.13, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2(a+bx)^5} dx$$

↓ 54

$$\int \left(\frac{5b^2}{a^6(a+bx)} - \frac{5b}{a^6x} + \frac{4b^2}{a^5(a+bx)^2} + \frac{1}{a^5x^2} + \frac{3b^2}{a^4(a+bx)^3} + \frac{2b^2}{a^3(a+bx)^4} + \frac{b^2}{a^2(a+bx)^5} \right) dx$$

↓ 2009

$$-\frac{5b \log(x)}{a^6} + \frac{5b \log(a+bx)}{a^6} - \frac{4b}{a^5(a+bx)} - \frac{1}{a^5x} - \frac{3b}{2a^4(a+bx)^2} - \frac{2b}{3a^3(a+bx)^3} - \frac{b}{4a^2(a+bx)^4}$$

input `Int[1/(x^2*(a + b*x)^5),x]`

output `-(1/(a^5*x)) - b/(4*a^2*(a + b*x)^4) - (2*b)/(3*a^3*(a + b*x)^3) - (3*b)/(2*a^4*(a + b*x)^2) - (4*b)/(a^5*(a + b*x)) - (5*b*Log[x])/a^6 + (5*b*Log[a + b*x])/a^6`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.06

method	result
default	$-\frac{b}{4a^2(bx+a)^4} + \frac{5b \ln(bx+a)}{a^6} - \frac{4b}{a^5(bx+a)} - \frac{3b}{2a^4(bx+a)^2} - \frac{2b}{3a^3(bx+a)^3} - \frac{1}{a^5x} - \frac{5b \ln(x)}{a^6}$
risch	$-\frac{5b^4x^4}{a^5} - \frac{35b^3x^3}{2a^4} - \frac{65b^2x^2}{3a^3} - \frac{125bx}{12a^2} - \frac{1}{a} + \frac{5b \ln(-bx-a)}{a^6} - \frac{5b \ln(x)}{a^6}$
norman	$-\frac{1}{a} + \frac{20b^2x^2}{a^3} + \frac{45b^3x^3}{a^4} + \frac{110b^4x^4}{3a^5} + \frac{125b^5x^5}{12a^6} - \frac{5b \ln(x)}{a^6} + \frac{5b \ln(bx+a)}{a^6}$
parallelrisch	$-\frac{60 \ln(x)x^5b^5 - 60 \ln(bx+a)x^5b^5 + 240 \ln(x)x^4ab^4 - 240 \ln(bx+a)x^4ab^4 - 125b^5x^5 + 360 \ln(x)x^3a^2b^3 - 360 \ln(bx+a)x^3a^2b^3 - 12a^6x(bx+a)}{12a^6x(bx+a)}$

input `int(1/x^2/(b*x+a)^5,x,method=_RETURNVERBOSE)`output
$$-1/4*b/a^2/(b*x+a)^4 + 5/a^6*b*\ln(b*x+a) - 4*b/a^5/(b*x+a) - 3/2/a^4*b/(b*x+a)^2 - 2/3*b/a^3/(b*x+a)^3 - 1/a^5/x - 5/a^6*b*\ln(x)$$
Fricas [B] (verification not implemented)Leaf count of result is larger than twice the leaf count of optimal. 197 vs. $2(72) = 144$.

Time = 0.09 (sec) , antiderivative size = 197, normalized size of antiderivative = 2.56

$$\int \frac{1}{x^2(a+bx)^5} dx = \frac{60ab^4x^4 + 210a^2b^3x^3 + 260a^3b^2x^2 + 125a^4bx + 12a^5 - 60(b^5x^5 + 4ab^4x^4 + 6a^2b^3x^3 + 4a^3b^2x^2 + a^4bx + a^5)}{12(a^6b^4x^5 + 4a^7b^3x^4 + 6a^8b^2x^3 + 4a^9bx^2 + a^{10})} + \frac{60(b^5x^5 + 4ab^4x^4 + 6a^2b^3x^3 + 4a^3b^2x^2 + a^4bx + a^5)}{12(a^6b^4x^5 + 4a^7b^3x^4 + 6a^8b^2x^3 + 4a^9bx^2 + a^{10})} \ln(x)$$

input `integrate(1/x^2/(b*x+a)^5,x, algorithm="fricas")`output
$$-1/12*(60*a*b^4*x^4 + 210*a^2*b^3*x^3 + 260*a^3*b^2*x^2 + 125*a^4*b*x + 12*a^5 - 60*(b^5*x^5 + 4*a*b^4*x^4 + 6*a^2*b^3*x^3 + 4*a^3*b^2*x^2 + a^4*b*x)) * \log(b*x + a) + 60*(b^5*x^5 + 4*a*b^4*x^4 + 6*a^2*b^3*x^3 + 4*a^3*b^2*x^2 + a^4*b*x) * \log(x) / (a^6*b^4*x^5 + 4*a^7*b^3*x^4 + 6*a^8*b^2*x^3 + 4*a^9*b*x^2 + a^{10})$$

Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.48

$$\int \frac{1}{x^2(a+bx)^5} dx = \frac{-12a^4 - 125a^3bx - 260a^2b^2x^2 - 210ab^3x^3 - 60b^4x^4}{12a^9x + 48a^8bx^2 + 72a^7b^2x^3 + 48a^6b^3x^4 + 12a^5b^4x^5} + \frac{5b(-\log(x) + \log(\frac{a}{b} + x))}{a^6}$$

input `integrate(1/x**2/(b*x+a)**5,x)`output `(-12*a**4 - 125*a**3*b*x - 260*a**2*b**2*x**2 - 210*a*b**3*x**3 - 60*b**4*x**4)/(12*a**9*x + 48*a**8*b*x**2 + 72*a**7*b**2*x**3 + 48*a**6*b**3*x**4 + 12*a**5*b**4*x**5) + 5*b*(-log(x) + log(a/b + x))/a**6`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.47

$$\int \frac{1}{x^2(a+bx)^5} dx = -\frac{60b^4x^4 + 210ab^3x^3 + 260a^2b^2x^2 + 125a^3bx + 12a^4}{12(a^5b^4x^5 + 4a^6b^3x^4 + 6a^7b^2x^3 + 4a^8bx^2 + a^9x)} + \frac{5b \log(bx+a)}{a^6} - \frac{5b \log(x)}{a^6}$$

input `integrate(1/x^2/(b*x+a)^5,x, algorithm="maxima")`output `-1/12*(60*b^4*x^4 + 210*a*b^3*x^3 + 260*a^2*b^2*x^2 + 125*a^3*b*x + 12*a^4)/(a^5*b^4*x^5 + 4*a^6*b^3*x^4 + 6*a^7*b^2*x^3 + 4*a^8*b*x^2 + a^9*x) + 5*b*log(b*x + a)/a^6 - 5*b*log(x)/a^6`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.40

$$\int \frac{1}{x^2(a+bx)^5} dx = -\frac{5b \log\left(\left|-\frac{a}{bx+a} + 1\right|\right)}{a^6} + \frac{b}{a^6\left(\frac{a}{bx+a} - 1\right)} - \frac{\frac{48a^3b^9}{bx+a} + \frac{18a^4b^9}{(bx+a)^2} + \frac{8a^5b^9}{(bx+a)^3} + \frac{3a^6b^9}{(bx+a)^4}}{12a^8b^8}$$

input `integrate(1/x^2/(b*x+a)^5,x, algorithm="giac")`output `-5*b*log(abs(-a/(b*x + a) + 1))/a^6 + b/(a^6*(a/(b*x + a) - 1)) - 1/12*(48*a^3*b^9/(b*x + a) + 18*a^4*b^9/(b*x + a)^2 + 8*a^5*b^9/(b*x + a)^3 + 3*a^6*b^9/(b*x + a)^4)/(a^8*b^8)`**Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.39

$$\int \frac{1}{x^2(a+bx)^5} dx = \frac{10b \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^6} - \frac{\frac{1}{a} + \frac{65b^2x^2}{3a^3} + \frac{35b^3x^3}{2a^4} + \frac{5b^4x^4}{a^5} + \frac{125bx}{12a^2}}{a^4x + 4a^3bx^2 + 6a^2b^2x^3 + 4ab^3x^4 + b^4x^5}$$

input `int(1/(x^2*(a + b*x)^5),x)`output `(10*b*atanh((2*b*x)/a + 1))/a^6 - (1/a + (65*b^2*x^2)/(3*a^3) + (35*b^3*x^3)/(2*a^4) + (5*b^4*x^4)/a^5 + (125*b*x)/(12*a^2))/(a^4*x + b^4*x^5 + 4*a^3*b*x^2 + 4*a*b^3*x^4 + 6*a^2*b^2*x^3)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.90

$$\int \frac{1}{x^2(a+bx)^5} dx$$

$$= \frac{60 \log(bx+a) a^4 bx + 240 \log(bx+a) a^3 b^2 x^2 + 360 \log(bx+a) a^2 b^3 x^3 + 240 \log(bx+a) a b^4 x^4 + 60 \log(bx+a) a^5}{(12 a^6 x^5 + 4 a^5 b x^4 + 6 a^4 b^2 x^3 + 4 a^3 b^3 x^2 + b^4 x)$$

input `int(1/x^2/(b*x+a)^5,x)`output `(60*log(a + b*x)*a**4*b*x + 240*log(a + b*x)*a**3*b**2*x**2 + 360*log(a + b*x)*a**2*b**3*x**3 + 240*log(a + b*x)*a*b**4*x**4 + 60*log(a + b*x)*b**5*x**5 - 60*log(x)*a**4*b*x - 240*log(x)*a**3*b**2*x**2 - 360*log(x)*a**2*b**3*x**3 - 240*log(x)*a*b**4*x**4 - 60*log(x)*b**5*x**5 - 12*a**5 - 110*a**4*b*x - 200*a**3*b**2*x**2 - 120*a**2*b**3*x**3 + 15*b**5*x**5)/(12*a**6*x*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2 + 4*a*b**3*x**3 + b**4*x**4))`

3.59 $\int \frac{1}{x^3(a+bx)^5} dx$

Optimal result	377
Mathematica [A] (verified)	377
Rubi [A] (verified)	378
Maple [A] (verified)	379
Fricas [B] (verification not implemented)	379
Sympy [A] (verification not implemented)	380
Maxima [A] (verification not implemented)	380
Giac [A] (verification not implemented)	381
Mupad [B] (verification not implemented)	381
Reduce [B] (verification not implemented)	382

Optimal result

Integrand size = 11, antiderivative size = 89

$$\int \frac{1}{x^3(a+bx)^5} dx = \frac{\frac{65b^2}{a^4} + \frac{125b^2}{4a^3} - \frac{1}{2ax^2} + \frac{3b}{a^2x} + \frac{105b^4x^2}{2a^5} + \frac{15b^5x^3}{a^6}}{(a+bx)^4} - \frac{15b^2 \log\left(\frac{a+bx}{x}\right)}{a^7}$$

output (65*b^2/a^4+125/4*b^2/a^3-1/2/a/x^2+3*b/a^2/x+105/2*b^4*x^2/a^5+15*b^5*x^3/a^6)/(b*x+a)^4-15*b^2*ln((b*x+a)/x)/a^7

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.01

$$\int \frac{1}{x^3(a+bx)^5} dx = \frac{\frac{a(-2a^5+12a^4bx+125a^3b^2x^2+260a^2b^3x^3+210ab^4x^4+60b^5x^5)}{x^2(a+bx)^4} + 60b^2 \log(x) - 60b^2 \log(a+bx)}{4a^7}$$

input Integrate[1/(x^3*(a + b*x)^5),x]

output

```
((a*(-2*a^5 + 12*a^4*b*x + 125*a^3*b^2*x^2 + 260*a^2*b^3*x^3 + 210*a*b^4*x^4 + 60*b^5*x^5))/(x^2*(a + b*x)^4) + 60*b^2*Log[x] - 60*b^2*Log[a + b*x])/
(4*a^7)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.18, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3(a+bx)^5} dx$$

↓ 54

$$\int \left(-\frac{15b^3}{a^7(a+bx)} + \frac{15b^2}{a^7x} - \frac{10b^3}{a^6(a+bx)^2} - \frac{5b}{a^6x^2} - \frac{6b^3}{a^5(a+bx)^3} + \frac{1}{a^5x^3} - \frac{3b^3}{a^4(a+bx)^4} - \frac{b^3}{a^3(a+bx)^5} \right) dx$$

↓ 2009

$$\frac{15b^2 \log(x)}{a^7} - \frac{15b^2 \log(a+bx)}{a^7} + \frac{10b^2}{a^6(a+bx)} + \frac{5b}{a^6x} + \frac{3b^2}{a^5(a+bx)^2} - \frac{1}{2a^5x^2} + \frac{b^2}{a^4(a+bx)^3} + \frac{b^2}{4a^3(a+bx)^4}$$

input

```
Int[1/(x^3*(a + b*x)^5),x]
```

output

```
-1/2*1/(a^5*x^2) + (5*b)/(a^6*x) + b^2/(4*a^3*(a + b*x)^4) + b^2/(a^4*(a + b*x)^3) + (3*b^2)/(a^5*(a + b*x)^2) + (10*b^2)/(a^6*(a + b*x)) + (15*b^2*Log[x])/a^7 - (15*b^2*Log[a + b*x])/a^7
```

Definitions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.06

method	result
norman	$\frac{-\frac{1}{2a} + \frac{3bx}{a^2} - \frac{60b^3x^3}{a^4} - \frac{135b^4x^4}{a^5} - \frac{110b^5x^5}{a^6} - \frac{125b^6x^6}{4a^7}}{x^2(bx+a)^4} + \frac{15b^2 \ln(x)}{a^7} - \frac{15b^2 \ln(bx+a)}{a^7}$
risch	$\frac{\frac{15b^5x^5}{a^6} + \frac{105b^4x^4}{2a^5} + \frac{65b^3x^3}{a^4} + \frac{125b^2x^2}{4a^3} + \frac{3bx}{a^2} - \frac{1}{2a}}{x^2(bx+a)^4} + \frac{15b^2 \ln(-x)}{a^7} - \frac{15b^2 \ln(bx+a)}{a^7}$
default	$-\frac{15b^2 \ln(bx+a)}{a^7} + \frac{10b^2}{a^6(bx+a)} + \frac{3b^2}{a^5(bx+a)^2} + \frac{b^2}{a^4(bx+a)^3} + \frac{b^2}{4a^3(bx+a)^4} - \frac{1}{2a^5x^2} + \frac{15b^2 \ln(x)}{a^7} + \frac{5b}{a^6x}$
parallelrisc	$\frac{60 \ln(x)x^6b^6 - 60 \ln(bx+a)x^6b^6 + 240 \ln(x)x^5ab^5 - 240 \ln(bx+a)x^5ab^5 - 125x^6b^6 + 360 \ln(x)x^4a^2b^4 - 360 \ln(bx+a)x^4a^2b^4 - 44}{4a^7x^2}$

input `int(1/x^3/(b*x+a)^5,x,method=_RETURNVERBOSE)`

output
$$\frac{(-1/2/a+3*b*x/a^2-60*b^3*x^3/a^4-135*b^4/a^5*x^4-110*b^5/a^6*x^5-125/4*b^6/a^7*x^6)/x^2/(b*x+a)^4+15/a^7*b^2*\ln(x)-15/a^7*b^2*\ln(b*x+a)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 218 vs. 2(84) = 168.

Time = 0.10 (sec) , antiderivative size = 218, normalized size of antiderivative = 2.45

$$\int \frac{1}{x^3(a+bx)^5} dx$$

$$= \frac{60ab^5x^5 + 210a^2b^4x^4 + 260a^3b^3x^3 + 125a^4b^2x^2 + 12a^5bx - 2a^6 - 60(b^6x^6 + 4ab^5x^5 + 6a^2b^4x^4 + 4a^3b^3x^3 + 12a^4b^2x^2 + 12a^5bx - 2a^6) \ln(x) - 15a^7b^2 \ln(bx+a)}{4(a^7b^4x^6 + 4a^8b^3x^5 + 6a^9b^2x^4 + \dots)}$$

input `integrate(1/x^3/(b*x+a)^5,x, algorithm="fricas")`

output
$$\frac{1}{4} \cdot (60 \cdot a \cdot b^5 \cdot x^5 + 210 \cdot a^2 \cdot b^4 \cdot x^4 + 260 \cdot a^3 \cdot b^3 \cdot x^3 + 125 \cdot a^4 \cdot b^2 \cdot x^2 + 12 \cdot a^5 \cdot b \cdot x - 2 \cdot a^6 - 60 \cdot (b^6 \cdot x^6 + 4 \cdot a \cdot b^5 \cdot x^5 + 6 \cdot a^2 \cdot b^4 \cdot x^4 + 4 \cdot a^3 \cdot b^3 \cdot x^3 + a^4 \cdot b^2 \cdot x^2) \cdot \log(b \cdot x + a) + 60 \cdot (b^6 \cdot x^6 + 4 \cdot a \cdot b^5 \cdot x^5 + 6 \cdot a^2 \cdot b^4 \cdot x^4 + 4 \cdot a^3 \cdot b^3 \cdot x^3 + a^4 \cdot b^2 \cdot x^2) \cdot \log(x)) / (a^7 \cdot b^4 \cdot x^6 + 4 \cdot a^8 \cdot b^3 \cdot x^5 + 6 \cdot a^9 \cdot b^2 \cdot x^4 + 4 \cdot a^{10} \cdot b \cdot x^3 + a^{11} \cdot x^2)$$

Sympy [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.44

$$\int \frac{1}{x^3(a+bx)^5} dx = \frac{-2a^5 + 12a^4bx + 125a^3b^2x^2 + 260a^2b^3x^3 + 210ab^4x^4 + 60b^5x^5}{4a^{10}x^2 + 16a^9bx^3 + 24a^8b^2x^4 + 16a^7b^3x^5 + 4a^6b^4x^6} + \frac{15b^2(\log(x) - \log(\frac{a}{b} + x))}{a^7}$$

input `integrate(1/x**3/(b*x+a)**5,x)`

output
$$\frac{(-2 \cdot a^{**5} + 12 \cdot a^{**4} \cdot b \cdot x + 125 \cdot a^{**3} \cdot b^{**2} \cdot x^{**2} + 260 \cdot a^{**2} \cdot b^{**3} \cdot x^{**3} + 210 \cdot a \cdot b^{**4} \cdot x^{**4} + 60 \cdot b^{**5} \cdot x^{**5}) / (4 \cdot a^{**10} \cdot x^{**2} + 16 \cdot a^{**9} \cdot b \cdot x^{**3} + 24 \cdot a^{**8} \cdot b^{**2} \cdot x^{**4} + 16 \cdot a^{**7} \cdot b^{**3} \cdot x^{**5} + 4 \cdot a^{**6} \cdot b^{**4} \cdot x^{**6}) + 15 \cdot b^{**2} \cdot (\log(x) - \log(a/b + x))}{a^{**7}}$$

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.46

$$\int \frac{1}{x^3(a+bx)^5} dx = \frac{60b^5x^5 + 210ab^4x^4 + 260a^2b^3x^3 + 125a^3b^2x^2 + 12a^4bx - 2a^5}{4(a^6b^4x^6 + 4a^7b^3x^5 + 6a^8b^2x^4 + 4a^9bx^3 + a^{10}x^2)} - \frac{15b^2 \log(bx+a)}{a^7} + \frac{15b^2 \log(x)}{a^7}$$

input `integrate(1/x^3/(b*x+a)^5,x, algorithm="maxima")`

output

$$\frac{1}{4} \frac{(60b^5x^5 + 210a^2b^4x^4 + 260a^2b^3x^3 + 125a^3b^2x^2 + 12a^4bx - 2a^5)}{(a^6b^4x^6 + 4a^7b^3x^5 + 6a^8b^2x^4 + 4a^9bx^3 + a^{10}x^2)} - 15b^2 \log(bx + a)/a^7 + 15b^2 \log(x)/a^7$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.44

$$\int \frac{1}{x^3(a+bx)^5} dx = \frac{15b^2 \log\left(\left|-\frac{a}{bx+a} + 1\right|\right)}{a^7} - \frac{\frac{12ab^2}{bx+a} - 11b^2}{2a^7\left(\frac{a}{bx+a} - 1\right)^2} + \frac{\frac{40a^6b^{14}}{bx+a} + \frac{12a^7b^{14}}{(bx+a)^2} + \frac{4a^8b^{14}}{(bx+a)^3} + \frac{a^9b^{14}}{(bx+a)^4}}{4a^{12}b^{12}}$$

input

```
integrate(1/x^3/(b*x+a)^5,x, algorithm="giac")
```

output

$$15b^2 \log(\text{abs}(-a/(bx+a) + 1))/a^7 - 1/2 * (12a^2b^2/(bx+a) - 11b^2) / (a^7 * (a/(bx+a) - 1)^2) + 1/4 * (40a^6b^{14}/(bx+a) + 12a^7b^{14}/(bx+a)^2 + 4a^8b^{14}/(bx+a)^3 + a^9b^{14}/(bx+a)^4) / (a^{12}b^{12})$$

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.38

$$\int \frac{1}{x^3(a+bx)^5} dx = \frac{\frac{125b^2x^2}{4a^3} - \frac{1}{2a} + \frac{65b^3x^3}{a^4} + \frac{105b^4x^4}{2a^5} + \frac{15b^5x^5}{a^6} + \frac{3bx}{a^2}}{a^4x^2 + 4a^3bx^3 + 6a^2b^2x^4 + 4ab^3x^5 + b^4x^6} - \frac{30b^2 \operatorname{atanh}\left(\frac{2bx}{a} + 1\right)}{a^7}$$

input

```
int(1/(x^3*(a + b*x)^5),x)
```

output

$$\left(\frac{125b^2x^2}{4a^3} - \frac{1}{2a} + \frac{65b^3x^3}{a^4} + \frac{105b^4x^4}{2a^5} + \frac{15b^5x^5}{a^6} + \frac{3bx}{a^2}\right) / (a^4x^2 + b^4x^6 + 4a^3bx^3 + 4a^2b^2x^4) - (30b^2 \operatorname{atanh}((2bx)/a + 1))/a^7$$

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 242, normalized size of antiderivative = 2.72

$$\int \frac{1}{x^3(a+bx)^5} dx$$

$$= \frac{-60 \log(bx+a) a^4 b^2 x^2 - 240 \log(bx+a) a^3 b^3 x^3 - 360 \log(bx+a) a^2 b^4 x^4 - 240 \log(bx+a) a b^5 x^5 - 60 a^6}{4 a^7 x^2 (a^4 + 4 a^3 b x + 6 a^2 b^2 x^2 + 4 a b^3 x^3 + b^4 x^4)}$$

input `int(1/x^3/(b*x+a)^5,x)`

output

```
( - 60*log(a + b*x)*a**4*b**2*x**2 - 240*log(a + b*x)*a**3*b**3*x**3 - 360
*log(a + b*x)*a**2*b**4*x**4 - 240*log(a + b*x)*a*b**5*x**5 - 60*log(a + b
*x)*b**6*x**6 + 60*log(x)*a**4*b**2*x**2 + 240*log(x)*a**3*b**3*x**3 + 360
*log(x)*a**2*b**4*x**4 + 240*log(x)*a*b**5*x**5 + 60*log(x)*b**6*x**6 - 2*
a**6 + 12*a**5*b*x + 110*a**4*b**2*x**2 + 200*a**3*b**3*x**3 + 120*a**2*b*
*4*x**4 - 15*b**6*x**6)/(4*a**7*x**2*(a**4 + 4*a**3*b*x + 6*a**2*b**2*x**2
+ 4*a*b**3*x**3 + b**4*x**4))
```

3.60 $\int \frac{1}{a+bx^2} dx$

Optimal result	383
Mathematica [A] (verified)	383
Rubi [A] (verified)	384
Maple [A] (verified)	384
Fricas [A] (verification not implemented)	385
Sympy [B] (verification not implemented)	385
Maxima [A] (verification not implemented)	386
Giac [A] (verification not implemented)	386
Mupad [B] (verification not implemented)	386
Reduce [B] (verification not implemented)	387

Optimal result

Integrand size = 9, antiderivative size = 20

$$\int \frac{1}{a+bx^2} dx = \frac{\arctan\left(\sqrt{\frac{b}{a}}x\right)}{\sqrt{ab}}$$

output `arctan((b/a)^(1/2)*x)/(a*b)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20

$$\int \frac{1}{a+bx^2} dx = \frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `Integrate[(a + b*x^2)^(-1),x]`

output `ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(Sqrt[a]*Sqrt[b])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.20, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{a + bx^2} dx$$

↓ 218

$$\frac{\arctan\left(\frac{\sqrt{bx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `Int[(a + b*x^2)^(-1), x]`

output `ArcTan[(Sqrt[b]*x)/Sqrt[a]]/(Sqrt[a]*Sqrt[b])`

Defintions of rubi rules used

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}}$	16
risch	$-\frac{\ln(bx + \sqrt{-ab})}{2\sqrt{-ab}} + \frac{\ln(-bx + \sqrt{-ab})}{2\sqrt{-ab}}$	41

input `int(1/(b*x^2+a),x,method=_RETURNVERBOSE)`

output `1/(a*b)^(1/2)*arctan(b*x/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 67, normalized size of antiderivative = 3.35

$$\int \frac{1}{a + bx^2} dx = \left[-\frac{\sqrt{-ab} \log\left(\frac{bx^2 - 2\sqrt{-ab}x - a}{bx^2 + a}\right)}{2ab}, \frac{\sqrt{ab} \arctan\left(\frac{\sqrt{ab}x}{a}\right)}{ab} \right]$$

input `integrate(1/(b*x^2+a),x, algorithm="fricas")`

output `[-1/2*sqrt(-a*b)*log((b*x^2 - 2*sqrt(-a*b)*x - a)/(b*x^2 + a))/(a*b), sqrt(a*b)*arctan(sqrt(a*b)*x/a)/(a*b)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(15) = 30.

Time = 0.07 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.65

$$\int \frac{1}{a + bx^2} dx = -\frac{\sqrt{-\frac{1}{ab}} \log\left(-a\sqrt{-\frac{1}{ab}} + x\right)}{2} + \frac{\sqrt{-\frac{1}{ab}} \log\left(a\sqrt{-\frac{1}{ab}} + x\right)}{2}$$

input `integrate(1/(b*x**2+a),x)`

output `-sqrt(-1/(a*b))*log(-a*sqrt(-1/(a*b)) + x)/2 + sqrt(-1/(a*b))*log(a*sqrt(-1/(a*b)) + x)/2`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{1}{a + bx^2} dx = \frac{\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/(b*x^2+a),x, algorithm="maxima")`output `arctan(b*x/sqrt(a*b))/sqrt(a*b)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{1}{a + bx^2} dx = \frac{\arctan\left(\frac{bx}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/(b*x^2+a),x, algorithm="giac")`output `arctan(b*x/sqrt(a*b))/sqrt(a*b)`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.80

$$\int \frac{1}{a + bx^2} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b}x}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `int(1/(a + b*x^2),x)`output `atan((b^(1/2)*x)/a^(1/2))/(a^(1/2)*b^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{1}{a + bx^2} dx = \frac{\sqrt{b} \sqrt{a} \operatorname{atan}\left(\frac{bx}{\sqrt{b}\sqrt{a}}\right)}{ab}$$

input `int(1/(b*x^2+a),x)`

output `(sqrt(b)*sqrt(a)*atan((b*x)/(sqrt(b)*sqrt(a))))/(a*b)`

3.61 $\int x(a + bx^2)^{-m} dx$

Optimal result	388
Mathematica [A] (verified)	388
Rubi [A] (verified)	389
Maple [A] (verified)	390
Fricas [A] (verification not implemented)	390
Sympy [B] (verification not implemented)	391
Maxima [F(-2)]	391
Giac [A] (verification not implemented)	392
Mupad [B] (verification not implemented)	392
Reduce [F]	392

Optimal result

Integrand size = 13, antiderivative size = 25

$$\int x(a + bx^2)^{-m} dx = -\frac{(a + bx^2)^{1-m}}{2b(-1 + m)}$$

output

$$-1/2*(b*x^2+a)^(1-m)/b/(-1+m)$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int x(a + bx^2)^{-m} dx = \frac{(a + bx^2)^{1-m}}{2b - 2bm}$$

input

$$\text{Integrate}[x/(a + b*x^2)^m, x]$$

output

$$(a + b*x^2)^(1 - m)/(2*b - 2*b*m)$$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x(a + bx^2)^{-m} dx$$

$$\downarrow \text{241}$$

$$\frac{(a + bx^2)^{1-m}}{2b(1 - m)}$$

input `Int[x/(a + b*x^2)^m,x]`

output `(a + b*x^2)^(1 - m)/(2*b*(1 - m))`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

method	result	size
derivativedivides	$\frac{(x^2b+a)^{1-m}}{2b(1-m)}$	26
default	$\frac{(x^2b+a)^{1-m}}{2b(1-m)}$	26
gosper	$-\frac{(x^2b+a)(x^2b+a)^{-m}}{2b(-1+m)}$	29
risch	$-\frac{(x^2b+a)(x^2b+a)^{-m}}{2b(-1+m)}$	29
orering	$-\frac{(x^2b+a)(x^2b+a)^{-m}}{2b(-1+m)}$	29
norman	$\left(-\frac{x^2}{2(-1+m)} - \frac{a}{2b(-1+m)}\right) e^{-m \ln(x^2b+a)}$	37
parallelrisch	$\frac{(-x^2ab-a^2)(x^2b+a)^{-m}}{2b(-1+m)a}$	38

input `int(x/((b*x^2+a)^m),x,method=_RETURNVERBOSE)`output `1/2/b*(b*x^2+a)^(1-m)/(1-m)`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

$$\int x(a+bx^2)^{-m} dx = -\frac{bx^2+a}{2(bm-b)(bx^2+a)^m}$$

input `integrate(x/((b*x^2+a)^m),x, algorithm="fricas")`output `-1/2*(b*x^2+a)/((b*m-b)*(b*x^2+a)^m)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 102 vs. $2(17) = 34$.

Time = 1.30 (sec) , antiderivative size = 102, normalized size of antiderivative = 4.08

$$\int x(a + bx^2)^{-m} dx = \begin{cases} \frac{x^2}{2a} & \text{for } b = 0 \wedge m = 1 \\ \frac{a^{-m}x^2}{2} & \text{for } b = 0 \\ \frac{\log\left(x - \sqrt{-\frac{a}{b}}\right)}{2b} + \frac{\log\left(x + \sqrt{-\frac{a}{b}}\right)}{2b} & \text{for } m = 1 \\ -\frac{a}{2bm(a+bx^2)^m} - \frac{bx^2}{2bm(a+bx^2)^m - 2b(a+bx^2)^m} & \text{otherwise} \end{cases}$$

input `integrate(x/((b*x**2+a)**m),x)`

output `Piecewise((x**2/(2*a), Eq(b, 0) & Eq(m, 1)), (x**2/(2*a**m), Eq(b, 0)), (log(x - sqrt(-a/b))/(2*b) + log(x + sqrt(-a/b))/(2*b), Eq(m, 1)), (-a/(2*b*m*(a + b*x**2)**m - 2*b*(a + b*x**2)**m) - b*x**2/(2*b*m*(a + b*x**2)**m - 2*b*(a + b*x**2)**m), True))`

Maxima [F(-2)]

Exception generated.

$$\int x(a + bx^2)^{-m} dx = \text{Exception raised: ValueError}$$

input `integrate(x/((b*x^2+a)^m),x, algorithm="maxima")`

output `Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(-m>0)', see `assume?` for more details)Is`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int x(a + bx^2)^{-m} dx = -\frac{(bx^2 + a)^{-m+1}}{2b(m-1)}$$

input `integrate(x/((b*x^2+a)^m),x, algorithm="giac")`

output `-1/2*(b*x^2 + a)^(-m + 1)/(b*(m - 1))`

Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int x(a + bx^2)^{-m} dx = -\frac{(bx^2 + a)^{1-m}}{2b(m-1)}$$

input `int(x/(a + b*x^2)^m,x)`

output `-(a + b*x^2)^(1 - m)/(2*b*(m - 1))`

Reduce [F]

$$\int x(a + bx^2)^{-m} dx = \int \frac{x}{(bx^2 + a)^m} dx$$

input `int(x/((b*x^2+a)^m),x)`

output `int(x/(a + b*x**2)**m,x)`

3.62 $\int \frac{1}{a+bx^3} dx$

Optimal result	393
Mathematica [A] (verified)	393
Rubi [A] (verified)	394
Maple [C] (verified)	397
Fricas [A] (verification not implemented)	397
Sympy [A] (verification not implemented)	398
Maxima [A] (verification not implemented)	398
Giac [A] (verification not implemented)	399
Mupad [B] (verification not implemented)	399
Reduce [B] (verification not implemented)	400

Optimal result

Integrand size = 9, antiderivative size = 94

$$\int \frac{1}{a+bx^3} dx = \frac{\sqrt[3]{\frac{a}{b}} \left(\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{2\sqrt[3]{\frac{a}{b}} - x} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}} + x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}} x + x^2} \right) \right)}{3a}$$

output `1/3*(a/b)^(1/3)*(3^(1/2)*arctan(3^(1/2)*x/(2*(a/b)^(1/3)-x))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2)))/a`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.95

$$\int \frac{1}{a+bx^3} dx = \frac{2\sqrt{3} \arctan \left(\frac{1 - 2\sqrt[3]{bx}}{\sqrt{3}\sqrt[3]{a}} \right) - 2 \log \left(\sqrt[3]{a} + \sqrt[3]{bx} \right) + \log \left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2 \right)}{6a^{2/3}\sqrt[3]{b}}$$

input `Integrate[(a + b*x^3)^(-1),x]`

output `-1/6*(2*Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/Sqrt[3]] - 2*Log[a^(1/3) + b^(1/3)*x] + Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/(a^(2/3)*b^(1/3))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.18, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + bx^3} dx \\
 & \quad \downarrow 750 \\
 & \frac{\int \frac{2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}}} dx}{3a^{2/3}} \\
 & \quad \downarrow 16 \\
 & \frac{\int \frac{2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
 & \quad \downarrow 1142 \\
 & \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\int -\frac{\sqrt[3]{b}(\sqrt[3]{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
 & \quad \downarrow 25
 \end{aligned}$$

$$\begin{aligned}
& \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 27 \\
& \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1082 \\
& \frac{\frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{3 \int \frac{1}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} d\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{-\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^{-3}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 217 \\
& \frac{\frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \\
& \quad \downarrow 1103 \\
& \frac{\frac{\log(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx+b^{2/3}x^2})}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}}
\end{aligned}$$

input `Int[(a + b*x^3)^(-1), x]`

output `Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3))`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.29

method	result	size
risch	$\frac{\sum_{-R=\text{RootOf}(b-Z^3+a)} \frac{\ln(x-R)}{-R^2}}{3b}$	27
default	$\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{3}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$	91

input `int(1/(b*x^3+a),x,method=_RETURNVERBOSE)`

output `1/3/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 299, normalized size of antiderivative = 3.18

$$\int \frac{1}{a + bx^3} dx$$

$$= \frac{3 \sqrt{\frac{1}{3}} ab \sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}} \log\left(\frac{2abx^3 - 3(a^2b)^{\frac{1}{3}}ax - a^2 + 3\sqrt{\frac{1}{3}}\left(2abx^2 + (a^2b)^{\frac{2}{3}}x - (a^2b)^{\frac{1}{3}}a\right)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}}}{bx^3 + a}\right) - (a^2b)^{\frac{2}{3}} \log\left(abx^2 - \dots\right)}{6a^2b}$$

input `integrate(1/(b*x^3+a),x, algorithm="fricas")`

output

```
[1/6*(3*sqrt(1/3)*a*b*sqrt(-(a^2*b)^(1/3)/b)*log((2*a*b*x^3 - 3*(a^2*b)^(1/3)*a*x - a^2 + 3*sqrt(1/3)*(2*a*b*x^2 + (a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt(-(a^2*b)^(1/3)/b))/(b*x^3 + a) - (a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 2*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^2*b), 1/6*(6*sqrt(1/3)*a*b*sqrt((a^2*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*(a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt((a^2*b)^(1/3)/b)/a^2 - (a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 2*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^2*b)]
```

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.21

$$\int \frac{1}{a + bx^3} dx = \text{RootSum}(27t^3a^2b - 1, (t \mapsto t \log(3ta + x)))$$

input

```
integrate(1/(b*x**3+a),x)
```

output

```
RootSum(27*_t**3*a**2*b - 1, Lambda(_t, _t*log(3*_t*a + x)))
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.04

$$\int \frac{1}{a + bx^3} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input

```
integrate(1/(b*x^3+a),x, algorithm="maxima")
```

output

```
1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(b*(a/b)^(2/3)) - 1/6*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(b*(a/b)^(2/3)) + 1/3*log(x + (a/b)^(1/3))/(b*(a/b)^(2/3))
```

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.19

$$\int \frac{1}{a + bx^3} dx = -\frac{\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{3a} + \frac{\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3ab}$$

$$+ \frac{(-ab^2)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6ab}$$

input `integrate(1/(b*x^3+a),x, algorithm="giac")`output `-1/3*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/a + 1/3*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/(a*b) + 1/6*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/(a*b)`**Mupad [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.05

$$\int \frac{1}{a + bx^3} dx = \frac{\ln(b^{1/3}x + a^{1/3})}{3a^{2/3}b^{1/3}} + \frac{\ln\left(3b^2x + \frac{3a^{1/3}b^{5/3}(-1+\sqrt{3}1i)}{2}\right)(-1 + \sqrt{3}1i)}{6a^{2/3}b^{1/3}}$$

$$- \frac{\ln\left(3b^2x - \frac{3a^{1/3}b^{5/3}(1+\sqrt{3}1i)}{2}\right)(1 + \sqrt{3}1i)}{6a^{2/3}b^{1/3}}$$

input `int(1/(a + b*x^3),x)`output `log(b^(1/3)*x + a^(1/3))/(3*a^(2/3)*b^(1/3)) + (log(3*b^2*x + (3*a^(1/3)*b^(5/3)*(3^(1/2)*1i - 1))/2)*(3^(1/2)*1i - 1))/(6*a^(2/3)*b^(1/3)) - (log(3*b^2*x - (3*a^(1/3)*b^(5/3)*(3^(1/2)*1i + 1))/2)*(3^(1/2)*1i + 1))/(6*a^(2/3)*b^(1/3))`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.71

$$\int \frac{1}{a + bx^3} dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) - \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) + 2\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right)}{6a^{\frac{2}{3}}b^{\frac{1}{3}}}$$

input `int(1/(b*x^3+a),x)`output `(a**(1/3)*(-2*sqrt(3)*atan((a**(1/3)-2*b**(1/3)*x)/(a**(1/3)*sqrt(3))) - log(a**(2/3)-b**(1/3)*a**(1/3)*x+b**(2/3)*x**2)+2*log(a**(1/3)+b**(1/3)*x))/(6*b**(1/3)*a)`

3.63 $\int \frac{x}{a+bx^3} dx$

Optimal result	401
Mathematica [A] (verified)	401
Rubi [A] (verified)	402
Maple [C] (verified)	405
Fricas [A] (verification not implemented)	405
Sympy [A] (verification not implemented)	406
Maxima [A] (verification not implemented)	406
Giac [A] (verification not implemented)	407
Mupad [B] (verification not implemented)	407
Reduce [B] (verification not implemented)	408

Optimal result

Integrand size = 11, antiderivative size = 101

$$\int \frac{x}{a+bx^3} dx = \frac{-\sqrt{3} \arctan\left(\frac{-\sqrt[3]{\frac{a}{b}}+2x}{\sqrt{3}\sqrt[3]{\frac{a}{b}}}\right) + \frac{1}{2} \log\left(\frac{\left(\sqrt[3]{\frac{a}{b}}+x\right)^2}{\left(\frac{a}{b}\right)^{2/3}-\sqrt[3]{\frac{a}{b}}x+x^2}\right)}{3\sqrt[3]{\frac{a}{b}}}$$

output

```
-1/3*(-3^(1/2)*arctan(1/3*(-(a/b)^(1/3)+2*x)*3^(1/2)/(a/b)^(1/3))+1/2*ln((
(a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2)))/(a/b)^(1/3)/b
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.88

$$\int \frac{x}{a+bx^3} dx = \frac{-2\sqrt{3} \arctan\left(\frac{1-2\sqrt[3]{\frac{b}{x}}}{\sqrt{3}\sqrt[3]{\frac{a}{b}}}\right) - 2 \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right) + \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6\sqrt[3]{ab^{2/3}}}$$

input `Integrate[x/(a + b*x^3),x]`

output $(-2\sqrt{3}\operatorname{ArcTan}[(1 - (2b^{1/3}x)/a^{1/3})/\sqrt{3}] - 2\operatorname{Log}[a^{1/3} + b^{1/3}x] + \operatorname{Log}[a^{2/3} - a^{1/3}b^{1/3}x + b^{2/3}x^2])/(6a^{1/3}b^{2/3})$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.15, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.727$, Rules used = {821, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{a + bx^3} dx \\
 & \quad \downarrow 821 \\
 & \frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} \\
 & \quad \downarrow 16 \\
 & \frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \\
 & \quad \downarrow 1142 \\
 & \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int -\frac{\sqrt[3]{b}(\sqrt[3]{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \\
 & \quad \downarrow 25
 \end{aligned}$$

$$\begin{aligned}
& \frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx - \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a} - 2 \sqrt[3]{bx})}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx}{2 \sqrt[3]{b}}}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3 \sqrt[3]{ab^{2/3}}} \\
& \quad \downarrow 27 \\
& \frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx - \frac{1}{2} \int \frac{\sqrt[3]{a} - 2 \sqrt[3]{bx}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3 \sqrt[3]{ab^{2/3}}} \\
& \quad \downarrow 1082 \\
& \frac{3 \int \frac{1}{\left(1 - \frac{2 \sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} d\left(1 - \frac{2 \sqrt[3]{bx}}{\sqrt[3]{a}}\right) - \frac{1}{2} \int \frac{\sqrt[3]{a} - 2 \sqrt[3]{bx}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx}{\sqrt[3]{b}}}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3 \sqrt[3]{ab^{2/3}}} \\
& \quad \downarrow 217 \\
& \frac{-\frac{1}{2} \int \frac{\sqrt[3]{a} - 2 \sqrt[3]{bx}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2 \sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3 \sqrt[3]{ab^{2/3}}} \\
& \quad \downarrow 1103 \\
& \frac{\frac{\log(a^{2/3} - \sqrt[3]{a} \sqrt[3]{bx} + b^{2/3} x^2)}{2 \sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2 \sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3 \sqrt[3]{ab^{2/3}}}
\end{aligned}$$

input

Int[x/(a + b*x^3), x]

output

$$\begin{aligned}
& -1/3 * \text{Log}[a^{(1/3)} + b^{(1/3)} * x] / (a^{(1/3)} * b^{(2/3)}) + (-((\text{Sqrt}[3] * \text{ArcTan}[(1 - \\
& (2 * b^{(1/3)} * x) / a^{(1/3)}) / \text{Sqrt}[3]]) / b^{(1/3)}) + \text{Log}[a^{(2/3)} - a^{(1/3)} * b^{(1/3)} * \\
& x + b^{(2/3)} * x^2] / (2 * b^{(1/3)})) / (3 * a^{(1/3)} * b^{(1/3)})
\end{aligned}$$

Defintions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 821 $\text{Int}[(x_)/((a_)+(b_)*(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{(-1)} \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]) \text{ Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.27

method	result	size
risch	$\frac{\sum_{-R=\text{RootOf}(bZ^3+a)} \frac{\ln(x-R)}{-R}}{3b}$	27
default	$-\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{\frac{\left(\frac{a}{b}\right)^{\frac{1}{3}}}{3}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}}$	91

input `int(x/(b*x^3+a),x,method=_RETURNVERBOSE)`

output `1/3/b*sum(1/_R*ln(x-_R),_R=RootOf(_Z^3*b+a))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 304, normalized size of antiderivative = 3.01

$$\int \frac{x}{a + bx^3} dx$$

$$= \frac{3 \sqrt{\frac{1}{3}} ab \sqrt{\frac{(-ab^2)^{\frac{1}{3}}}{a}} \log\left(\frac{2b^2x^3 - ab + 3\sqrt{\frac{1}{3}}(abx + 2(-ab^2)^{\frac{2}{3}}x^2 + (-ab^2)^{\frac{1}{3}}a) \sqrt{\frac{(-ab^2)^{\frac{1}{3}}}{a} - 3(-ab^2)^{\frac{2}{3}}x}}{bx^3 + a}}\right) + (-ab^2)^{\frac{2}{3}} \log(b^2x^3 + a)}{6ab^2}$$

input `integrate(x/(b*x^3+a),x, algorithm="fricas")`

output

```
[1/6*(3*sqrt(1/3)*a*b*sqrt((-a*b^2)^(1/3)/a)*log((2*b^2*x^3 - a*b + 3*sqrt(1/3)*(a*b*x + 2*(-a*b^2)^(2/3)*x^2 + (-a*b^2)^(1/3)*a)*sqrt((-a*b^2)^(1/3)/a) - 3*(-a*b^2)^(2/3)*x)/(b*x^3 + a)) + (-a*b^2)^(2/3)*log(b^2*x^2 + (-a*b^2)^(1/3)*b*x + (-a*b^2)^(2/3)) - 2*(-a*b^2)^(2/3)*log(b*x - (-a*b^2)^(1/3)))/(a*b^2), 1/6*(6*sqrt(1/3)*a*b*sqrt(-(-a*b^2)^(1/3)/a)*arctan(sqrt(1/3)*(2*b*x + (-a*b^2)^(1/3))*sqrt(-(-a*b^2)^(1/3)/a)/b) + (-a*b^2)^(2/3)*log(b^2*x^2 + (-a*b^2)^(1/3)*b*x + (-a*b^2)^(2/3)) - 2*(-a*b^2)^(2/3)*log(b*x - (-a*b^2)^(1/3)))/(a*b^2)]
```

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.24

$$\int \frac{x}{a + bx^3} dx = \text{RootSum}(27t^3ab^2 + 1, (t \mapsto t \log(9t^2ab + x)))$$

input

```
integrate(x/(b*x**3+a),x)
```

output

```
RootSum(27*_t**3*a*b**2 + 1, Lambda(_t, _t*log(9*_t**2*a*b + x)))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.97

$$\int \frac{x}{a + bx^3} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6b\left(\frac{a}{b}\right)^{\frac{1}{3}}} - \frac{\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}}$$

input

```
integrate(x/(b*x^3+a),x, algorithm="maxima")
```

output

```
1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(b*(a/b)^(1/3)) + 1/6*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(b*(a/b)^(1/3)) - 1/3*log(x + (a/b)^(1/3))/(b*(a/b)^(1/3))
```

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.11

$$\int \frac{x}{a + bx^3} dx = -\frac{\left(-\frac{a}{b}\right)^{\frac{2}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{3a} - \frac{\sqrt{3}(-ab^2)^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3ab^2} + \frac{\left(-ab^2\right)^{\frac{2}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6ab^2}$$

input `integrate(x/(b*x^3+a),x, algorithm="giac")`output `-1/3*(-a/b)^(2/3)*log(abs(x - (-a/b)^(1/3)))/a - 1/3*sqrt(3)*(-a*b^2)^(2/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/(a*b^2) + 1/6*(-a*b^2)^(2/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/(a*b^2)`**Mupad [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.10

$$\int \frac{x}{a + bx^3} dx = \frac{\ln\left(b^{1/3}x - (-a)^{1/3}\right)}{3(-a)^{1/3}b^{2/3}} + \frac{\ln\left(bx - \frac{(-a)^{1/3}b^{2/3}(-1+\sqrt{3}i)^2}{4}\right)(-1 + \sqrt{3}i)}{6(-a)^{1/3}b^{2/3}} - \frac{\ln\left(bx - \frac{(-a)^{1/3}b^{2/3}(1+\sqrt{3}i)^2}{4}\right)(1 + \sqrt{3}i)}{6(-a)^{1/3}b^{2/3}}$$

input `int(x/(a + b*x^3),x)`output `log(b^(1/3)*x - (-a)^(1/3))/(3*(-a)^(1/3)*b^(2/3)) + (log(b*x - ((-a)^(1/3)*b^(2/3)*(3^(1/2)*1i - 1)^2)/4)*(3^(1/2)*1i - 1))/(6*(-a)^(1/3)*b^(2/3)) - (log(b*x - ((-a)^(1/3)*b^(2/3)*(3^(1/2)*1i + 1)^2)/4)*(3^(1/2)*1i + 1))/(6*(-a)^(1/3)*b^(2/3))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.64

$$\int \frac{x}{a + bx^3} dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) + \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) - 2\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right)}{6b^{\frac{2}{3}}a^{\frac{1}{3}}}$$

input `int(x/(b*x^3+a),x)`

output `(- 2*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3))) + log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2) - 2*log(a**(1/3) + b**(1/3)*x))/(6*b**(2/3)*a**(1/3))`

3.64 $\int \frac{x^2}{a+bx^3} dx$

Optimal result	409
Mathematica [A] (verified)	409
Rubi [A] (verified)	410
Maple [A] (verified)	411
Fricas [A] (verification not implemented)	411
Sympy [A] (verification not implemented)	412
Maxima [A] (verification not implemented)	412
Giac [A] (verification not implemented)	412
Mupad [B] (verification not implemented)	413
Reduce [B] (verification not implemented)	413

Optimal result

Integrand size = 13, antiderivative size = 15

$$\int \frac{x^2}{a+bx^3} dx = \frac{\log(a+bx^3)}{3b}$$

output `1/3*ln(b*x^3+a)/b`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{a+bx^3} dx = \frac{\log(a+bx^3)}{3b}$$

input `Integrate[x^2/(a + b*x^3),x]`

output `Log[a + b*x^3]/(3*b)`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{a + bx^3} dx$$

↓ 792

$$\frac{\log(a + bx^3)}{3b}$$

input `Int[x^2/(a + b*x^3), x]`

output `Log[a + b*x^3]/(3*b)`

Defintions of rubi rules used

rule 792 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$\frac{\ln(bx^3+a)}{3b}$	14
default	$\frac{\ln(bx^3+a)}{3b}$	14
norman	$\frac{\ln(bx^3+a)}{3b}$	14
risch	$\frac{\ln(bx^3+a)}{3b}$	14
parallelrisch	$\frac{\ln(bx^3+a)}{3b}$	14

input `int(x^2/(b*x^3+a),x,method=_RETURNVERBOSE)`output `1/3*ln(b*x^3+a)/b`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^2}{a + bx^3} dx = \frac{\log(bx^3 + a)}{3b}$$

input `integrate(x^2/(b*x^3+a),x, algorithm="fricas")`output `1/3*log(b*x^3 + a)/b`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{x^2}{a + bx^3} dx = \frac{\log(a + bx^3)}{3b}$$

input `integrate(x**2/(b*x**3+a),x)`

output `log(a + b*x**3)/(3*b)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^2}{a + bx^3} dx = \frac{\log(bx^3 + a)}{3b}$$

input `integrate(x^2/(b*x^3+a),x, algorithm="maxima")`

output `1/3*log(b*x^3 + a)/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{x^2}{a + bx^3} dx = \frac{\log(|bx^3 + a|)}{3b}$$

input `integrate(x^2/(b*x^3+a),x, algorithm="giac")`

output `1/3*log(abs(b*x^3 + a))/b`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^2}{a + bx^3} dx = \frac{\ln(bx^3 + a)}{3b}$$

input `int(x^2/(a + b*x^3),x)`output `log(a + b*x^3)/(3*b)`**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.47

$$\int \frac{x^2}{a + bx^3} dx = \frac{\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) + \log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right)}{3b}$$

input `int(x^2/(b*x^3+a),x)`output `(log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2) + log(a**(1/3) + b**(1/3)*x))/(3*b)`

3.65 $\int \frac{x^3}{a+bx^3} dx$

Optimal result	414
Mathematica [A] (verified)	414
Rubi [A] (verified)	415
Maple [C] (verified)	419
Fricas [A] (verification not implemented)	419
Sympy [A] (verification not implemented)	420
Maxima [A] (verification not implemented)	420
Giac [A] (verification not implemented)	421
Mupad [B] (verification not implemented)	421
Reduce [B] (verification not implemented)	422

Optimal result

Integrand size = 13, antiderivative size = 100

$$\int \frac{x^3}{a+bx^3} dx = \frac{x}{b} - \frac{\sqrt[3]{\frac{a}{b}} \left(\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{2\sqrt[3]{\frac{a}{b}} - x} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}} + x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}} x + x^2} \right) \right)}{3b}$$

output

$$\frac{x}{b} - \frac{1}{3b} \left(\sqrt[3]{\frac{a}{b}} \left(\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{2\sqrt[3]{\frac{a}{b}} - x} \right) + \frac{1}{2} \ln \left(\frac{\left(\sqrt[3]{\frac{a}{b}} + x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}} x + x^2} \right) \right) \right)$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.08

$$\int \frac{x^3}{a+bx^3} dx = \frac{6\sqrt[3]{bx} + 2\sqrt{3}\sqrt[3]{a} \arctan \left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}} \right) - 2\sqrt[3]{a} \log \left(\sqrt[3]{a} + \sqrt[3]{bx} \right) + \sqrt[3]{a} \log \left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2 \right)}{6b^{4/3}}$$

input `Integrate[x^3/(a + b*x^3),x]`

output $(6*b^{(1/3)}*x + 2*\text{Sqrt}[3]*a^{(1/3)}*\text{ArcTan}[(1 - (2*b^{(1/3)}*x)/a^{(1/3)})/\text{Sqrt}[3]] - 2*a^{(1/3)}*\text{Log}[a^{(1/3)} + b^{(1/3)}*x] + a^{(1/3)}*\text{Log}[a^{(2/3)} - a^{(1/3)}*b^{(1/3)}*x + b^{(2/3)}*x^2])/(6*b^{(4/3)})$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.23, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {843, 750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{a + bx^3} dx$$

$$\downarrow 843$$

$$\frac{x}{b} - \frac{a}{b} \int \frac{1}{bx^3 + a} dx$$

$$\downarrow 750$$

$$\frac{x}{b} - \frac{a}{b} \left(\int \frac{2\sqrt[3]{a} - \sqrt[3]{b}x}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{b}x + a^{2/3}} dx + \int \frac{1}{\sqrt[3]{b}x + \sqrt[3]{a}} dx \right)$$

$$\downarrow 16$$

$$\frac{x}{b} - \frac{a}{b} \left(\int \frac{2\sqrt[3]{a} - \sqrt[3]{b}x}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{b}x + a^{2/3}} dx + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b}x)}{3a^{2/3}\sqrt[3]{b}} \right)$$

$$\downarrow 1142$$

$$\frac{x}{b} - \frac{a \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx - \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a} - 2 \sqrt[3]{b_x})}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{2 \sqrt[3]{b}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 a^{2/3} \sqrt[3]{b}} \right)}{b}$$

25

$$\frac{x}{b} - \frac{a \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx + \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a} - 2 \sqrt[3]{b_x})}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{2 \sqrt[3]{b}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 a^{2/3} \sqrt[3]{b}} \right)}{b}$$

27

$$\frac{x}{b} - \frac{a \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a} - 2 \sqrt[3]{b_x}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{3 a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 a^{2/3} \sqrt[3]{b}} \right)}{b}$$

1082

$$\frac{x}{b} - \frac{a \left(\frac{\frac{1}{2} \int \frac{\sqrt[3]{a} - 2 \sqrt[3]{b_x}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx + \frac{\int \frac{1}{\left(1 - 2 \frac{\sqrt[3]{b_x}}{\sqrt[3]{a}}\right)^2} d\left(1 - 2 \frac{\sqrt[3]{b_x}}{\sqrt[3]{a}}\right) - \left(1 - 2 \frac{\sqrt[3]{b_x}}{\sqrt[3]{a}}\right)^{-3}}{\sqrt[3]{b}}}{3 a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 a^{2/3} \sqrt[3]{b}} \right)}{b}$$

217

$$\frac{x}{b} - \frac{a \left(\frac{\frac{1}{2} \int \frac{\sqrt[3]{a-2\sqrt[3]{b}x}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{b}x + a^{2/3}} dx - \frac{\sqrt[3]{a} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}} \right)}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}} \right)}{b}$$

↓ 1103

$$\frac{x}{b} - \frac{a \left(\frac{\frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{2\sqrt[3]{b}} - \frac{\sqrt[3]{a} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}} \right)}{b}$$

input `Int[x^3/(a + b*x^3), x]`

output `x/b - (a*(Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3))))/b`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[((a_) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 750 $\text{Int}[((a_) + (b_*)(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 843 $\text{Int}[((c_*)(x_)^m)*((a_) + (b_*)(x_)^n)^{p_}), x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1})/(b*(m+n*p+1))), x] - \text{Simp}[a*c^n*(m-n+1)/(b*(m+n*p+1)) \text{ Int}[(c*x)^{(m-n)}*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[m, n-1] \ \&\& \ \text{NeQ}[m+n*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$
- rule 1082 $\text{Int}[((a_) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[((d_) + (e_*)(x_))/((a_) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[((d_) + (e_*)(x_))/((a_) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.02 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.34

method	result	size
risch	$\frac{x}{b} - \frac{a \left(\sum_{R=\text{RootOf}(bZ^3+a)} \frac{\ln(x-R)}{-R^2} \right)}{3b^2}$	34
default	$\frac{x}{b} - \frac{\left(\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} \right) a}{b}$	103

input `int(x^3/(b*x^3+a),x,method=_RETURNVERBOSE)`

output `x/b-1/3/b^2*a*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.06

$$\int \frac{x^3}{a + bx^3} dx = \frac{2\sqrt{3}\left(-\frac{a}{b}\right)^{\frac{1}{3}} \arctan\left(\frac{2\sqrt{3}bx\left(-\frac{a}{b}\right)^{\frac{2}{3}}-\sqrt{3}a}{3a}\right) - \left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right) + 2\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{6b}$$

input `integrate(x^3/(b*x^3+a),x, algorithm="fricas")`

output

```
1/6*(2*sqrt(3)*(-a/b)^(1/3)*arctan(1/3*(2*sqrt(3)*b*x*(-a/b)^(2/3) - sqrt(
3)*a)/a) - (-a/b)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3)) + 2*(-a/b
)^(1/3)*log(x - (-a/b)^(1/3)) + 6*x)/b
```

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.22

$$\int \frac{x^3}{a + bx^3} dx = \text{RootSum}(27t^3b^4 + a, (t \mapsto t \log(-3tb + x))) + \frac{x}{b}$$

input

```
integrate(x**3/(b*x**3+a),x)
```

output

```
RootSum(27*_t**3*b**4 + a, Lambda(_t, _t*log(-3*_t*b + x))) + x/b
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.06

$$\int \frac{x^3}{a + bx^3} dx = \frac{x}{b} - \frac{\sqrt{3}a \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b^2\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{a \log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6b^2\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{a \log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3b^2\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input

```
integrate(x^3/(b*x^3+a),x, algorithm="maxima")
```

output

```
x/b - 1/3*sqrt(3)*a*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(b
^2*(a/b)^(2/3)) + 1/6*a*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(b^2*(a/b)
^(2/3)) - 1/3*a*log(x + (a/b)^(1/3))/(b^2*(a/b)^(2/3))
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.11

$$\int \frac{x^3}{a + bx^3} dx = \frac{\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{3b} + \frac{x}{b} - \frac{\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b^2} - \frac{\left(-ab^2\right)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6b^2}$$

input `integrate(x^3/(b*x^3+a),x, algorithm="giac")`output `1/3*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/b + x/b - 1/3*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/b^2 - 1/6*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/b^2`**Mupad [B] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.14

$$\int \frac{x^3}{a + bx^3} dx = \frac{x}{b} + \frac{(-a)^{1/3} \ln\left(\left(-a\right)^{4/3} + a b^{1/3} x\right)}{3 b^{4/3}} - \frac{(-a)^{1/3} \ln\left(3\left(-a\right)^{4/3} b^{2/3} \left(\frac{1}{2} + \frac{\sqrt{3} i i}{2}\right) - 3 a b x\right) \left(\frac{1}{2} + \frac{\sqrt{3} i i}{2}\right)}{3 b^{4/3}} + \frac{(-a)^{1/3} \ln\left(9\left(-a\right)^{4/3} b^{2/3} \left(-\frac{1}{6} + \frac{\sqrt{3} i i}{6}\right) + 3 a b x\right) \left(-\frac{1}{6} + \frac{\sqrt{3} i i}{6}\right)}{b^{4/3}}$$

input `int(x^3/(a + b*x^3),x)`output `x/b + ((-a)^(1/3)*log((-a)^(4/3) + a*b^(1/3)*x))/(3*b^(4/3)) - ((-a)^(1/3)*log(3*(-a)^(4/3)*b^(2/3)*((3^(1/2)*1i)/2 + 1/2) - 3*a*b*x)*((3^(1/2)*1i)/2 + 1/2))/(3*b^(4/3)) + ((-a)^(1/3)*log(9*(-a)^(4/3)*b^(2/3)*((3^(1/2)*1i)/6 - 1/6) + 3*a*b*x)*((3^(1/2)*1i)/6 - 1/6))/b^(4/3)`

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.78

$$\int \frac{x^3}{a + bx^3} dx$$

$$= \frac{2a^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) + a^{\frac{1}{3}}\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) - 2a^{\frac{1}{3}}\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) + 6b^{\frac{1}{3}}x}{6b^{\frac{4}{3}}}$$

input `int(x^3/(b*x^3+a),x)`output `(2*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3))) + a
(1/3)*log(a(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2) - 2*a**(1/3)*l
og(a**(1/3) + b**(1/3)*x) + 6*b**(1/3)*x)/(6*b**(1/3)*b)`

3.66 $\int \frac{x^4}{a+bx^3} dx$

Optimal result	423
Mathematica [A] (verified)	424
Rubi [A] (verified)	424
Maple [C] (verified)	428
Fricas [A] (verification not implemented)	428
Sympy [A] (verification not implemented)	429
Maxima [A] (verification not implemented)	429
Giac [A] (verification not implemented)	430
Mupad [B] (verification not implemented)	430
Reduce [B] (verification not implemented)	431

Optimal result

Integrand size = 13, antiderivative size = 113

$$\int \frac{x^4}{a+bx^3} dx = \frac{x^2}{2b} + \frac{a \left(-\sqrt{3} \arctan \left(\frac{-\sqrt[3]{\frac{a}{b}}+2x}{\sqrt{3}\sqrt[3]{\frac{a}{b}}} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}}+x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}}x+x^2} \right) \right)}{3\sqrt[3]{\frac{a}{b}}b^2}$$

output

```
1/2*x^2/b+1/3*a*(-3^(1/2)*arctan(1/3*(-(a/b)^(1/3)+2*x)*3^(1/2)/(a/b)^(1/3)))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2))/(a/b)^(1/3)/b^2
```


Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 111, normalized size of antiderivative = 0.98

$$\int \frac{x^4}{a + bx^3} dx$$

$$= \frac{3b^{2/3}x^2 + 2\sqrt{3}a^{2/3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right) + 2a^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right) - a^{2/3} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6b^{5/3}}$$

input `Integrate[x^4/(a + b*x^3),x]`

output `(3*b^(2/3)*x^2 + 2*Sqrt[3]*a^(2/3)*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/Sqrt[3]] + 2*a^(2/3)*Log[a^(1/3) + b^(1/3)*x] - a^(2/3)*Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/(6*b^(5/3))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.18, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {843, 821, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^4}{a + bx^3} dx$$

$$\downarrow \text{843}$$

$$\frac{x^2}{2b} - \frac{a}{b} \int \frac{x}{bx^3 + a} dx$$

$$\downarrow \text{821}$$

$$\frac{x^2}{2b} - \frac{a \left(\frac{\int \frac{\sqrt[3]{bx+\sqrt[3]{a}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\int \frac{1}{\sqrt[3]{bx+\sqrt[3]{a}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} \right)}{b}$$

↓ 16

$$\frac{x^2}{2b} - \frac{a \left(\frac{\int \frac{\sqrt[3]{bx+\sqrt[3]{a}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{b}$$

↓ 1142

$$\frac{x^2}{2b} - \frac{a \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{b}$$

↓ 25

$$\frac{x^2}{2b} - \frac{a \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{b}$$

↓ 27

$$\frac{x^2}{2b} - \frac{a \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{b}$$

↓ 1082

$$\frac{x^2}{2b} - \frac{a \left(\frac{\int \frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} dx}{\sqrt[3]{b}} - \frac{\int \frac{\sqrt[3]{a} - 2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx} + a^{2/3}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{b}$$

217

$$\frac{x^2}{2b} - \frac{a \left(\frac{-\frac{1}{2} \int \frac{\sqrt[3]{a} - 2\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx} + a^{2/3}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{b}$$

1103

$$\frac{x^2}{2b} - \frac{a \left(\frac{\frac{\log(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2)}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{b}$$

input `Int[x^4/(a + b*x^3), x]`

output `x^2/(2*b) - (a*(-1/3*Log[a^(1/3) + b^(1/3)*x]/(a^(1/3)*b^(2/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])/b^(1/3)) + Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(1/3)*b^(1/3)))/b`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])]$
- rule 821 $\text{Int}[(x_)/((a_)+(b_)*(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{(-1)} \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]) \text{ Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}\{a, b\}, x]$
- rule 843 $\text{Int}[(c_)*(x_)^m*((a_)+(b_)*(x_)^n)^p, x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*(m+n*p+1))), x] - \text{Simp}[a*c^n*((m-n+1)/(b*(m+n*p+1))) \text{ Int}[(c*x)^{(m-n)}*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, p\}, x \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[m, n-1] \&\& \text{NeQ}[m+n*p+1, 0] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1142

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.33

method	result	size
risch	$\frac{x^2}{2b} - \frac{a \left(\sum_{-R=\text{RootOf}(b-Z^3+a)} \frac{\ln(x-R)}{-R} \right)}{3b^2}$	37
default	$\frac{x^2}{2b} - \frac{\left(\begin{aligned} &-\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x-\left(\frac{a}{b}\right)^{\frac{1}{3}}-1\right)}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}} \end{aligned} \right) a}{b}$	106

input

```
int(x^4/(b*x^3+a),x,method=_RETURNVERBOSE)
```

output

```
1/2*x^2/b-1/3/b^2*a*sum(1/_R*ln(x-_R),_R=RootOf(_Z^3*b+a))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.09

$$\int \frac{x^4}{a + bx^3} dx$$

$$= \frac{3x^2 - 2\sqrt{3}\left(\frac{a^2}{b^2}\right)^{\frac{1}{3}} \arctan\left(\frac{2\sqrt{3}bx\left(\frac{a^2}{b^2}\right)^{\frac{1}{3}} - \sqrt{3}a}{3a}\right) - \left(\frac{a^2}{b^2}\right)^{\frac{1}{3}} \log\left(ax^2 - bx\left(\frac{a^2}{b^2}\right)^{\frac{2}{3}} + a\left(\frac{a^2}{b^2}\right)^{\frac{1}{3}}\right) + 2\left(\frac{a^2}{b^2}\right)^{\frac{1}{3}} \log\left(\dots\right)}{6b}$$

input `integrate(x^4/(b*x^3+a),x, algorithm="fricas")`

output $\frac{1}{6}(3x^2 - 2\sqrt{3}(a^2/b^2)^{1/3}\arctan(1/3(2\sqrt{3}bx(a^2/b^2)^{1/3} - \sqrt{3}a)/a) - (a^2/b^2)^{1/3}\log(ax^2 - bx(a^2/b^2)^{2/3} + a(a^2/b^2)^{1/3}) + 2(a^2/b^2)^{1/3}\log(ax + b(a^2/b^2)^{2/3}))/b$

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.28

$$\int \frac{x^4}{a + bx^3} dx = \text{RootSum} \left(27t^3b^5 - a^2, \left(t \mapsto t \log \left(\frac{9t^2b^3}{a} + x \right) \right) \right) + \frac{x^2}{2b}$$

input `integrate(x**4/(b*x**3+a),x)`

output `RootSum(27*_t**3*b**5 - a**2, Lambda(_t, _t*log(9*_t**2*b**3/a + x))) + x**2/(2*b)`

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.96

$$\int \frac{x^4}{a + bx^3} dx = \frac{x^2}{2b} - \frac{\sqrt{3}a \arctan \left(\frac{\sqrt{3} \left(2x - \left(\frac{a}{b} \right)^{1/3} \right)}{3 \left(\frac{a}{b} \right)^{1/3}} \right)}{3b^2 \left(\frac{a}{b} \right)^{1/3}} - \frac{a \log \left(x^2 - x \left(\frac{a}{b} \right)^{1/3} + \left(\frac{a}{b} \right)^{2/3} \right)}{6b^2 \left(\frac{a}{b} \right)^{1/3}} + \frac{a \log \left(x + \left(\frac{a}{b} \right)^{1/3} \right)}{3b^2 \left(\frac{a}{b} \right)^{1/3}}$$

input `integrate(x^4/(b*x^3+a),x, algorithm="maxima")`

output $\frac{1}{2}x^2/b - \frac{1}{3}\sqrt{3}a\arctan(1/3\sqrt{3}(2x - (a/b)^{1/3})/(a/b)^{1/3})/(b^2(a/b)^{1/3}) - \frac{1}{6}a\log(x^2 - x(a/b)^{1/3} + (a/b)^{2/3})/(b^2(a/b)^{1/3}) + \frac{1}{3}a\log(x + (a/b)^{1/3})/(b^2(a/b)^{1/3})$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.01

$$\int \frac{x^4}{a+bx^3} dx = \frac{x^2}{2b} + \frac{\left(-\frac{a}{b}\right)^{\frac{2}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{3b} + \frac{\sqrt{3}(-ab^2)^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b^3} - \frac{(-ab^2)^{\frac{2}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6b^3}$$

input `integrate(x^4/(b*x^3+a),x, algorithm="giac")`output $\frac{1}{2}x^2/b + \frac{1}{3}(-a/b)^{2/3} \log(\text{abs}(x - (-a/b)^{1/3}))/b + \frac{1}{3}\sqrt{3}*(-a*b^2)^{2/3} \arctan(1/3*\sqrt{3}*(2*x + (-a/b)^{1/3})/(-a/b)^{1/3})/b^3 - 1/6*(-a*b^2)^{2/3} \log(x^2 + x*(-a/b)^{1/3} + (-a/b)^{2/3})/b^3$ **Mupad [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.06

$$\int \frac{x^4}{a+bx^3} dx = \frac{x^2}{2b} + \frac{a^{2/3} \ln\left(\frac{a^{7/3}}{b^{4/3}} + \frac{a^2 x}{b}\right)}{3b^{5/3}} - \frac{a^{2/3} \ln\left(\frac{a^2 x}{b} + \frac{a^{7/3} \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^2}{b^{4/3}}\right)}{3b^{5/3}} \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) + \frac{a^{2/3} \ln\left(\frac{a^2 x}{b} + \frac{9a^{7/3} \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)^2}{b^{4/3}}\right)}{b^{5/3}} \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)$$

input `int(x^4/(a + b*x^3),x)`output $x^2/(2*b) + (a^{2/3} \log(a^{7/3}/b^{4/3} + (a^2*x)/b))/(3*b^{5/3}) - (a^{2/3} \log((a^2*x)/b + (a^{7/3}*((3^{1/2}*1i)/2 + 1/2)^2)/b^{4/3}))*((3^{1/2}*1i)/2 + 1/2))/(3*b^{5/3}) + (a^{2/3} \log((a^2*x)/b + (9*a^{7/3}*((3^{1/2}*1i)/6 - 1/6)^2)/b^{4/3}))*((3^{1/2}*1i)/6 - 1/6))/b^{5/3}$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.72

$$\int \frac{x^4}{a + bx^3} dx$$

$$= \frac{2\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) a + 3b^{\frac{2}{3}}a^{\frac{1}{3}}x^2 - \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) a + 2\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) a}{6b^{\frac{5}{3}}a^{\frac{1}{3}}}$$

input `int(x^4/(b*x^3+a),x)`output `(2*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*a + 3*b**(2/3)*a**(1/3)*x**2 - log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a + 2*log(a**(1/3) + b**(1/3)*x)*a)/(6*b**(2/3)*a**(1/3)*b)`

3.67 $\int \frac{1}{(a+bx^3)^2} dx$

Optimal result	432
Mathematica [A] (verified)	433
Rubi [A] (verified)	433
Maple [C] (verified)	437
Fricas [A] (verification not implemented)	438
Sympy [A] (verification not implemented)	438
Maxima [A] (verification not implemented)	439
Giac [A] (verification not implemented)	439
Mupad [B] (verification not implemented)	440
Reduce [B] (verification not implemented)	440

Optimal result

Integrand size = 9, antiderivative size = 112

$$\int \frac{1}{(a+bx^3)^2} dx = \frac{x}{3a(a+bx^3)} + \frac{2\sqrt[3]{\frac{a}{b}} \left(\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{2\sqrt[3]{\frac{a}{b}} - x} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}} + x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}} x + x^2} \right) \right)}{9a^2}$$

```
output 1/3*x/a/(b*x^3+a)+2/9*(a/b)^(1/3)*(3^(1/2)*arctan(3^(1/2)*x/(2*(a/b)^(1/3)
-x))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2)))/a^2
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.05

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$= \frac{\frac{3a^{2/3}x}{a+bx^3} - \frac{2\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{b}x}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}} + \frac{2 \log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{\sqrt[3]{b}} - \frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{\sqrt[3]{b}}}{9a^{5/3}}$$

input

```
Integrate[(a + b*x^3)^(-2), x]
```

output

```
((3*a^(2/3)*x)/(a + b*x^3) - (2*sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/sqrt[3]])/b^(1/3) + (2*Log[a^(1/3) + b^(1/3)*x])/b^(1/3) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/b^(1/3))/(9*a^(5/3))
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.21, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {749, 750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$\downarrow 749$$

$$\frac{2 \int \frac{1}{bx^3+a} dx}{3a} + \frac{x}{3a(a + bx^3)}$$

$$\downarrow 750$$

$$2 \left(\frac{\int \frac{{}_2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}} dx}{3a^{2/3}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 16

$$2 \left(\frac{\int \frac{{}_2\sqrt[3]{a} - \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 1142

$$2 \left(\frac{\frac{{}_3\sqrt{b}({}_3\sqrt{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 25

$$2 \left(\frac{\frac{{}_3\sqrt{b}({}_3\sqrt{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 27

$$2 \left(\frac{\frac{{}_3\sqrt{b}({}_3\sqrt{a} - 2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{1}{2} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

↓ 1082

$$2 \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{bx}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{1}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} dx \left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^{-3}}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

217

$$2 \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{bx}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

1103

$$2 \left(\frac{\frac{\log(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx+b^{2/3}x^2})}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) + \frac{x}{3a(a+bx^3)}$$

input `Int[(a + b*x^3)^(-2), x]`

output `x/(3*a*(a + b*x^3)) + (2*(Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])]/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3)))/(3*a)`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])]$
- rule 749 $\text{Int}[(a_)+(b_)*(x_)^n)^p, x_Symbol] \rightarrow \text{Simp}[(-x)*((a + b*x^n)^{p+1}/(a*n*(p+1))), x] + \text{Simp}[(n*(p+1)+1)/(a*n*(p+1)) \text{ Int}[(a + b*x^n)^{p+1}, x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (\text{IntegerQ}[2*p] \ || \ \text{Denominator}[p + 1/n] < \text{Denominator}[p])]$
- rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}\{a, b\}, x]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.41

method	result	size
risch	$\frac{x}{3a(bx^3+a)} + \frac{2 \left(\sum_{R=\text{RootOf}(b-Z^3+a)} \frac{\ln(x-R)}{-R^2} \right)}{9ab}$	46
default	$\frac{x}{3a(bx^3+a)} + \frac{\frac{2 \ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right) - \ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}} - \left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}} - \left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}}{a} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{3}\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$	112

input

```
int(1/(b*x^3+a)^2,x,method=_RETURNVERBOSE)
```

output

```
1/3*x/a/(b*x^3+a)+2/9/a/b*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 389, normalized size of antiderivative = 3.47

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$= \frac{3a^2bx + 3\sqrt{\frac{1}{3}}(ab^2x^3 + a^2b)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}} \log\left(\frac{2abx^3 - 3(a^2b)^{\frac{1}{3}}ax - a^2 + 3\sqrt{\frac{1}{3}}(2abx^2 + (a^2b)^{\frac{2}{3}}x - (a^2b)^{\frac{1}{3}}a)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}}}{bx^3 + a}\right)}{9(a^3b^2x^3 + a^4)}$$

input `integrate(1/(b*x^3+a)^2,x, algorithm="fricas")`

output

```
[1/9*(3*a^2*b*x + 3*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt(-(a^2*b)^(1/3)/b)*
log((2*a*b*x^3 - 3*(a^2*b)^(1/3)*a*x - a^2 + 3*sqrt(1/3)*(2*a*b*x^2 + (a^2*
b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt(-(a^2*b)^(1/3)/b))/(b*x^3 + a) - (b*x^
3 + a)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) + 2*
(b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^3*b^2*x^3 + a^4*b
), 1/9*(3*a^2*b*x + 6*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt((a^2*b)^(1/3)/b)*
arctan(sqrt(1/3)*(2*(a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt((a^2*b)^(1/3)/
b)/a^2) - (b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b
)^(1/3)*a) + 2*(b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^3*
b^2*x^3 + a^4*b)]
```

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.35

$$\int \frac{1}{(a + bx^3)^2} dx = \frac{x}{3a^2 + 3abx^3} + \text{RootSum}\left(729t^3a^5b - 8, \left(t \mapsto t \log\left(\frac{9ta^2}{2} + x\right)\right)\right)$$

input `integrate(1/(b*x**3+a)**2,x)`

output `x/(3*a**2 + 3*a*b*x**3) + RootSum(729*_t**3*a**5*b - 8, Lambda(_t, _t*log(9*_t*a**2/2 + x)))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.09

$$\int \frac{1}{(a + bx^3)^2} dx = \frac{x}{3(abx^3 + a^2)} + \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{2\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input `integrate(1/(b*x^3+a)^2,x, algorithm="maxima")`

output `1/3*x/(a*b*x^3 + a^2) + 2/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(a*b*(a/b)^(2/3)) - 1/9*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(a*b*(a/b)^(2/3)) + 2/9*log(x + (a/b)^(1/3))/(a*b*(a/b)^(2/3))`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.13

$$\int \frac{1}{(a + bx^3)^2} dx = -\frac{2\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{9a^2} + \frac{x}{3(bx^3 + a)a} + \frac{2\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9a^2b} + \frac{\left(-ab^2\right)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{9a^2b}$$

input `integrate(1/(b*x^3+a)^2,x, algorithm="giac")`

output

```
-2/9*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/a^2 + 1/3*x/((b*x^3 + a)*a) +
2/9*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)
^(1/3))/(a^2*b) + 1/9*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/
3))/(a^2*b)
```

Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.14

$$\int \frac{1}{(a + bx^3)^2} dx = \frac{x}{3a(bx^3 + a)} + \frac{2 \ln \left(\frac{2b^{5/3}}{a^{2/3}} + \frac{2b^2x}{a} \right)}{9a^{5/3}b^{1/3}}$$

$$+ \frac{\ln \left(\frac{2b^2x}{a} + \frac{b^{5/3}(-1+\sqrt{3}i)}{a^{2/3}} \right) (-1 + \sqrt{3}i)}{9a^{5/3}b^{1/3}}$$

$$- \frac{\ln \left(\frac{2b^2x}{a} - \frac{b^{5/3}(1+\sqrt{3}i)}{a^{2/3}} \right) (1 + \sqrt{3}i)}{9a^{5/3}b^{1/3}}$$

input

```
int(1/(a + b*x^3)^2,x)
```

output

```
x/(3*a*(a + b*x^3)) + (2*log((2*b^(5/3))/a^(2/3) + (2*b^2*x)/a))/(9*a^(5/3)
)*b^(1/3)) + (log((2*b^2*x)/a + (b^(5/3)*(3^(1/2)*1i - 1))/a^(2/3))*(3^(1/
2)*1i - 1))/(9*a^(5/3)*b^(1/3)) - (log((2*b^2*x)/a - (b^(5/3)*(3^(1/2)*1i
+ 1))/a^(2/3))*(3^(1/2)*1i + 1))/(9*a^(5/3)*b^(1/3))
```

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.53

$$\int \frac{1}{(a + bx^3)^2} dx$$

$$= \frac{-2a^{\frac{4}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}}-2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) - 2a^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}}-2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) b x^3 - a^{\frac{4}{3}} \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) - a^{\frac{1}{3}} \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}x\right)}{9b^{\frac{1}{3}}a^2(bx^3 + a)}$$

input

```
int(1/(b*x^3+a)^2,x)
```

output

```
( - 2*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*
a - 2*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*
b*x**3 - a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a -
a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*b*x**3 + 2*a*
*(1/3)*log(a**(1/3) + b**(1/3)*x)*a + 2*a**(1/3)*log(a**(1/3) + b**(1/3)*x
)*b*x**3 + 3*b**(1/3)*a*x)/(9*b**(1/3)*a**2*(a + b*x**3))
```

3.68 $\int \frac{x}{(a+bx^3)^2} dx$

Optimal result	442
Mathematica [A] (verified)	443
Rubi [A] (verified)	443
Maple [C] (verified)	447
Fricas [A] (verification not implemented)	447
Sympy [A] (verification not implemented)	448
Maxima [A] (verification not implemented)	448
Giac [A] (verification not implemented)	449
Mupad [B] (verification not implemented)	449
Reduce [B] (verification not implemented)	450

Optimal result

Integrand size = 11, antiderivative size = 124

$$\int \frac{x}{(a+bx^3)^2} dx = \frac{x^2}{3a(a+bx^3)} - \frac{-\sqrt{3} \arctan\left(\frac{-\sqrt[3]{\frac{a}{b}}+2x}{\sqrt{3}\sqrt[3]{\frac{a}{b}}}\right) + \frac{1}{2} \log\left(\frac{\left(\sqrt[3]{\frac{a}{b}}+x\right)^2}{\left(\frac{a}{b}\right)^{2/3} - \sqrt[3]{\frac{a}{b}}x+x^2}\right)}{9a\sqrt[3]{\frac{a}{b}}b}$$

output

```
1/3*x^2/a/(b*x^3+a)-1/9*(-3^(1/2)*arctan(1/3*(-(a/b)^(1/3)+2*x)*3^(1/2)/(a/b)^(1/3))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2)))/a/(a/b)^(1/3)/b
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.96

$$\int \frac{x}{(a + bx^3)^2} dx$$

$$= \frac{6\sqrt[3]{ax^2}}{a+bx^3} - \frac{2\sqrt{3} \arctan\left(\frac{1 - 2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{b^{2/3}} - \frac{2 \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right)}{b^{2/3}} + \frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{b^{2/3}}$$

$$18a^{4/3}$$

input `Integrate[x/(a + b*x^3)^2,x]`

output $\left(\frac{6a^{1/3}x^2}{a + bx^3} - \frac{2\sqrt{3}\text{ArcTan}\left[\frac{1 - (2b^{1/3})x}{a^{1/3}}\right]}{\sqrt{3}}\right)/b^{2/3} - \frac{2\text{Log}[a^{1/3} + b^{1/3}x]}{b^{2/3}} + \frac{\text{Log}[a^{2/3} - a^{1/3}b^{1/3}x + b^{2/3}x^2]}{b^{2/3}}\bigg)/(18a^{4/3})$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.15, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.818$, Rules used = {819, 821, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a + bx^3)^2} dx$$

$$\downarrow 819$$

$$\frac{\int \frac{x}{bx^3+a} dx}{3a} + \frac{x^2}{3a(a + bx^3)}$$

$$\downarrow 821$$

$$\begin{aligned}
& \frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} + \frac{x^2}{3a(a+bx^3)} \\
& \quad \downarrow 16 \\
& \frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}}}{3a} + \frac{x^2}{3a(a+bx^3)} \\
& \quad \downarrow 1142 \\
& \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a-2}\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}}}{3a} + \frac{x^2}{3a(a+bx^3)} \\
& \quad \downarrow 25 \\
& \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a-2}\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}}}{3a} + \frac{x^2}{3a(a+bx^3)} \\
& \quad \downarrow 27 \\
& \frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{1}{2} \int \frac{\sqrt[3]{a-2}\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}}}{3a} + \frac{x^2}{3a(a+bx^3)} \\
& \quad \downarrow 1082 \\
& \frac{3 \int \frac{1}{\left(1 - 2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} d\left(1 - 2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right) - \frac{1}{2} \int \frac{\sqrt[3]{a-2}\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}}}{3\sqrt[3]{a}\sqrt[3]{b}} + \frac{x^2}{3a(a+bx^3)} \\
& \quad \downarrow 217
\end{aligned}$$

$$\begin{aligned}
 & \frac{-\frac{1}{2} \int \frac{\sqrt[3]{a-2\sqrt[3]{b}x}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{b}x + a^{2/3}} dx - \frac{\sqrt[3]{3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}}}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3\sqrt[3]{ab^{2/3}}} + \frac{x^2}{3a(a+bx^3)} \\
 & \quad \downarrow \text{1103} \\
 & \frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{2\sqrt[3]{b}} - \frac{\sqrt[3]{3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}}}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3\sqrt[3]{ab^{2/3}}} + \frac{x^2}{3a(a+bx^3)}
 \end{aligned}$$

input `Int[x/(a + b*x^3)^2,x]`

output `x^2/(3*a*(a + b*x^3)) + (-1/3*Log[a^(1/3) + b^(1/3)*x]/(a^(1/3)*b^(2/3)) + ((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3]])/b^(1/3)) + Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(1/3)*b^(1/3))/(3*a)`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 $\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 819 $\text{Int}[(c_.)*(x_)^m*((a_.) + (b_.)*(x_)^n)^{p_}], x_Symbol] \rightarrow \text{Simp}[(-c*x)^{(m+1)}*((a + b*x^n)^{(p+1)}/(a*c*n*(p+1))), x] + \text{Simp}[(m + n*(p + 1) + 1)/(a*n*(p + 1)) \ \text{Int}[(c*x)^m*(a + b*x^n)^{(p+1)}, x], x] /;$ $\text{FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 821 $\text{Int}[(x_)/((a_.) + (b_.)*(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{(-1)} \ \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]) \ \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /;$ $\text{FreeQ}[\{a, b\}, x]$

rule 1082 $\text{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S \ \text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c]) /;$ $\text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_.) + (e_.)*(x_)/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}[(d_.) + (e_.)*(x_)/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \ \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \ \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /;$ $\text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.39

method	result	size
risch	$\frac{x^2}{3a(bx^3+a)} + \frac{\sum_{R=\text{RootOf}(bZ^3+a)} \frac{\ln(x-R)}{-R}}{9ab}$	48
default	$\frac{x^2}{3a(bx^3+a)} + \frac{-\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{1}{3}}}{3a} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{\frac{3}{b}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}}$	114

```
input int(x/(b*x^3+a)^2,x,method=_RETURNVERBOSE)
```

```
output 1/3*x^2/a/(b*x^3+a)+1/9/a/b*sum(1/_R*ln(x-_R),_R=RootOf(_Z^3*b+a))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 386, normalized size of antiderivative = 3.11

$$\int \frac{x}{(a+bx^3)^2} dx$$

$$= \frac{6ab^2x^2 + 3\sqrt{\frac{1}{3}}(ab^2x^3 + a^2b)\sqrt{-\frac{(ab^2)^{\frac{1}{3}}}{a}} \log\left(\frac{2b^2x^3 - ab + 3\sqrt{\frac{1}{3}}(abx + 2(ab^2)^{\frac{2}{3}}x^2 - (ab^2)^{\frac{1}{3}}a)\sqrt{-\frac{(ab^2)^{\frac{1}{3}}}{a}} - 3(ab^2)^{\frac{2}{3}}x}{bx^3 + a}}\right) + \dots}{18(a^2b^3x^3 + a^3b)}$$

```
input integrate(x/(b*x^3+a)^2,x, algorithm="fricas")
```


output

```
[1/18*(6*a*b^2*x^2 + 3*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt(-(a*b^2)^(1/3)/a)
)*log((2*b^2*x^3 - a*b + 3*sqrt(1/3)*(a*b*x + 2*(a*b^2)^(2/3)*x^2 - (a*b^2)^(1/3)*a)*sqrt(-(a*b^2)^(1/3)/a) - 3*(a*b^2)^(2/3)*x)/(b*x^3 + a)) + (b*x^3 + a)*(a*b^2)^(2/3)*log(b^2*x^2 - (a*b^2)^(1/3)*b*x + (a*b^2)^(2/3)) - 2*(b*x^3 + a)*(a*b^2)^(2/3)*log(b*x + (a*b^2)^(1/3)))/(a^2*b^3*x^3 + a^3*b^2), 1/18*(6*a*b^2*x^2 - 6*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt((a*b^2)^(1/3)/a)*arctan(-sqrt(1/3)*(2*b*x - (a*b^2)^(1/3))*sqrt((a*b^2)^(1/3)/a)/b) + (b*x^3 + a)*(a*b^2)^(2/3)*log(b^2*x^2 - (a*b^2)^(1/3)*b*x + (a*b^2)^(2/3)) - 2*(b*x^3 + a)*(a*b^2)^(2/3)*log(b*x + (a*b^2)^(1/3)))/(a^2*b^3*x^3 + a^3*b^2)]
```

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.35

$$\int \frac{x}{(a + bx^3)^2} dx = \frac{x^2}{3a^2 + 3abx^3} + \text{RootSum}(729t^3a^4b^2 + 1, (t \mapsto t \log(81t^2a^3b + x)))$$

input

```
integrate(x/(b*x**3+a)**2,x)
```

output

```
x**2/(3*a**2 + 3*a*b*x**3) + RootSum(729*_t**3*a**4*b**2 + 1, Lambda(_t, _t*log(81*_t**2*a**3*b + x)))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.00

$$\int \frac{x}{(a + bx^3)^2} dx = \frac{x^2}{3(abx^3 + a^2)} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{18ab\left(\frac{a}{b}\right)^{\frac{1}{3}}} - \frac{\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{9ab\left(\frac{a}{b}\right)^{\frac{1}{3}}}$$

input

```
integrate(x/(b*x^3+a)^2,x, algorithm="maxima")
```

output

```
1/3*x^2/(a*b*x^3 + a^2) + 1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3)))/(a/b)^(1/3))/(a*b*(a/b)^(1/3)) + 1/18*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(a*b*(a/b)^(1/3)) - 1/9*log(x + (a/b)^(1/3))/(a*b*(a/b)^(1/3))
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.04

$$\int \frac{x}{(a + bx^3)^2} dx = \frac{x^2}{3(bx^3 + a)a} - \frac{\left(-\frac{a}{b}\right)^{\frac{2}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{9a^2}$$

$$- \frac{\sqrt{3}(-ab^2)^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9a^2b^2}$$

$$+ \frac{\left(-ab^2\right)^{\frac{2}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{18a^2b^2}$$

input

```
integrate(x/(b*x^3+a)^2,x, algorithm="giac")
```

output

```
1/3*x^2/((b*x^3 + a)*a) - 1/9*(-a/b)^(2/3)*log(abs(x - (-a/b)^(1/3)))/a^2 - 1/9*sqrt(3)*(-a*b^2)^(2/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/(a^2*b^2) + 1/18*(-a*b^2)^(2/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/(a^2*b^2)
```

Mupad [B] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.11

$$\int \frac{x}{(a + bx^3)^2} dx$$

$$= \frac{x^2}{3a(bx^3 + a)} + \frac{(-1)^{1/3} \ln\left(\frac{(-1)^{2/3} b^{2/3}}{9a^{5/3}} + \frac{bx}{9a^2}\right)}{9a^{4/3} b^{2/3}}$$

$$- \frac{(-1)^{1/3} \ln\left((-1)^{2/3} a^{1/3} - 2b^{1/3} x + (-1)^{1/6} \sqrt{3} a^{1/3}\right) \left(\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{9a^{4/3} b^{2/3}}$$

$$+ \frac{(-1)^{1/3} \ln\left(2b^{1/3} x - (-1)^{2/3} a^{1/3} + (-1)^{1/6} \sqrt{3} a^{1/3}\right) \left(-\frac{1}{2} + \frac{\sqrt{3} \operatorname{li}}{2}\right)}{9a^{4/3} b^{2/3}}$$

input `int(x/(a + b*x^3)^2,x)`

output
$$\frac{x^2/(3*a*(a + b*x^3)) + ((-1)^{(1/3)}*\log((-1)^{(2/3)}*b^{(2/3)})/(9*a^{(5/3)}) + (b*x)/(9*a^2))/(9*a^{(4/3)}*b^{(2/3)}) - ((-1)^{(1/3)}*\log((-1)^{(2/3)}*a^{(1/3)} - 2*b^{(1/3)}*x + (-1)^{(1/6)}*3^{(1/2)}*a^{(1/3)}*((3^{(1/2)}*i)/2 + 1/2))/(9*a^{(4/3)}*b^{(2/3)}) + ((-1)^{(1/3)}*\log(2*b^{(1/3)}*x - (-1)^{(2/3)}*a^{(1/3)} + (-1)^{(1/6)}*3^{(1/2)}*a^{(1/3)}*((3^{(1/2)}*i)/2 - 1/2))/(9*a^{(4/3)}*b^{(2/3)})}$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.27

$$\int \frac{x}{(a + bx^3)^2} dx$$

$$= \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) a - 2\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) b x^3 + 6b^{\frac{2}{3}}a^{\frac{1}{3}}x^2 + \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) a + \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) b x^3 - 2\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) a - 2\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) b x^3}{18b^{\frac{2}{3}}a^{\frac{4}{3}}(bx^3 + a)}$$

input `int(x/(b*x^3+a)^2,x)`

output
$$\left(- 2*\sqrt{3}*\operatorname{atan}\left(\frac{a^{(1/3)} - 2*b^{(1/3)}*x}{a^{(1/3)}*\sqrt{3}}\right)*a - 2*\sqrt{3}*\operatorname{atan}\left(\frac{a^{(1/3)} - 2*b^{(1/3)}*x}{a^{(1/3)}*\sqrt{3}}\right)*b*x**3 + 6*b^{(2/3)}*a^{(1/3)}*x**2 + \log(a^{(2/3)} - b^{(1/3)}*a^{(1/3)}*x + b^{(2/3)}*x**2)*a + \log(a^{(2/3)} - b^{(1/3)}*a^{(1/3)}*x + b^{(2/3)}*x**2)*b*x**3 - 2*\log(a^{(1/3)} + b^{(1/3)}*x)*a - 2*\log(a^{(1/3)} + b^{(1/3)}*x)*b*x**3 \right) / (18*b^{(2/3)}*a^{(4/3)}*(a + b*x**3))$$

$$3.69 \quad \int \frac{x^2}{(a+bx^3)^2} dx$$

Optimal result	451
Mathematica [A] (verified)	451
Rubi [A] (verified)	452
Maple [A] (verified)	453
Fricas [A] (verification not implemented)	453
Sympy [A] (verification not implemented)	454
Maxima [A] (verification not implemented)	454
Giac [A] (verification not implemented)	454
Mupad [B] (verification not implemented)	455
Reduce [B] (verification not implemented)	455

Optimal result

Integrand size = 13, antiderivative size = 16

$$\int \frac{x^2}{(a+bx^3)^2} dx = -\frac{1}{3b(a+bx^3)}$$

output `-1/3/b/(b*x^3+a)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{x^2}{(a+bx^3)^2} dx = -\frac{1}{3b(a+bx^3)}$$

input `Integrate[x^2/(a + b*x^3)^2,x]`

output `-1/3*1/(b*(a + b*x^3))`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {793}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(a + bx^3)^2} dx$$

↓ 793

$$-\frac{1}{3b(a + bx^3)}$$

input `Int[x^2/(a + b*x^3)^2,x]`

output `-1/3*1/(b*(a + b*x^3))`

Defintions of rubi rules used

rule 793

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
gospers	$-\frac{1}{3b(bx^3+a)}$	15
derivativedivides	$-\frac{1}{3b(bx^3+a)}$	15
default	$-\frac{1}{3b(bx^3+a)}$	15
norman	$-\frac{1}{3b(bx^3+a)}$	15
risch	$-\frac{1}{3b(bx^3+a)}$	15
parallelrisch	$-\frac{1}{3b(bx^3+a)}$	15
orering	$-\frac{1}{3b(bx^3+a)}$	15

input `int(x^2/(b*x^3+a)^2,x,method=_RETURNVERBOSE)`output `-1/3/b/(b*x^3+a)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{(a + bx^3)^2} dx = -\frac{1}{3(b^2x^3 + ab)}$$

input `integrate(x^2/(b*x^3+a)^2,x, algorithm="fricas")`output `-1/3/(b^2*x^3 + a*b)`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{x^2}{(a + bx^3)^2} dx = -\frac{1}{3ab + 3b^2x^3}$$

input `integrate(x**2/(b*x**3+a)**2,x)`output `-1/(3*a*b + 3*b**2*x**3)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x^2}{(a + bx^3)^2} dx = -\frac{1}{3(bx^3 + a)b}$$

input `integrate(x^2/(b*x^3+a)^2,x, algorithm="maxima")`output `-1/3/((b*x^3 + a)*b)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x^2}{(a + bx^3)^2} dx = -\frac{1}{3(bx^3 + a)b}$$

input `integrate(x^2/(b*x^3+a)^2,x, algorithm="giac")`output `-1/3/((b*x^3 + a)*b)`

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x^2}{(a + bx^3)^2} dx = -\frac{1}{3b(bx^3 + a)}$$

input `int(x^2/(a + b*x^3)^2,x)`

output `-1/(3*b*(a + b*x^3))`

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{x^2}{(a + bx^3)^2} dx = \frac{x^3}{3a(bx^3 + a)}$$

input `int(x^2/(b*x^3+a)^2,x)`

output `x**3/(3*a*(a + b*x**3))`

3.70 $\int \frac{x^3}{(a+bx^3)^2} dx$

Optimal result	456
Mathematica [A] (verified)	457
Rubi [A] (verified)	457
Maple [C] (verified)	460
Fricas [A] (verification not implemented)	461
Sympy [A] (verification not implemented)	462
Maxima [A] (verification not implemented)	462
Giac [A] (verification not implemented)	463
Mupad [B] (verification not implemented)	464
Reduce [B] (verification not implemented)	464

Optimal result

Integrand size = 13, antiderivative size = 115

$$\int \frac{x^3}{(a+bx^3)^2} dx = -\frac{x}{3b(a+bx^3)} + \frac{\sqrt[3]{\frac{a}{b}} \left(\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{2\sqrt[3]{\frac{a}{b}} - x} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}} + x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}} x + x^2} \right) \right)}{9ab}$$

```
output -1/3*x/b/(b*x^3+a)+1/9*(a/b)^(1/3)*(3^(1/2)*arctan(3^(1/2)*x/(2*(a/b)^(1/3)
)-x))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2))/a/b
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.03

$$\int \frac{x^3}{(a + bx^3)^2} dx$$

$$= \frac{-\frac{6\sqrt[3]{bx^3}}{a+bx^3} - \frac{2\sqrt{3} \arctan\left(\frac{1 - 2\sqrt[3]{bx^3}}{\sqrt[3]{a}}\right)}{a^{2/3}} + \frac{2 \log\left(\sqrt[3]{a} + \sqrt[3]{bx^3}\right)}{a^{2/3}} - \frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx^3} + b^{2/3}x^2\right)}{a^{2/3}}}{18b^{4/3}}$$

input `Integrate[x^3/(a + b*x^3)^2,x]`

output `((-6*b^(1/3)*x)/(a + b*x^3) - (2*Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/Sqrt[3]])/a^(2/3) + (2*Log[a^(1/3) + b^(1/3)*x])/a^(2/3) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/a^(2/3))/(18*b^(4/3))`

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.18, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {817, 750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(a + bx^3)^2} dx$$

$$\downarrow 817$$

$$\frac{\int \frac{1}{bx^3+a} dx}{3b} - \frac{x}{3b(a + bx^3)}$$

$$\downarrow 750$$

$$\frac{\int \frac{{}_2\sqrt[3]{a}-\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{bx+\sqrt[3]{a}}}dx}{3a^{2/3}} - \frac{x}{3b(a+bx^3)}$$

↓ 16

$$\frac{\int \frac{{}_2\sqrt[3]{a}-\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a+\sqrt[3]{bx}}\right)}{3a^{2/3}\sqrt[3]{b}} - \frac{x}{3b(a+bx^3)}$$

↓ 1142

$$\frac{\frac{3}{2}\sqrt[3]{a}\int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx - \frac{\int \frac{\sqrt[3]{b}\left(\sqrt[3]{a}-2\sqrt[3]{bx}\right)}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a+\sqrt[3]{bx}}\right)}{3a^{2/3}\sqrt[3]{b}} - \frac{x}{3b(a+bx^3)}}$$

↓ 25

$$\frac{\frac{3}{2}\sqrt[3]{a}\int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx + \frac{\int \frac{\sqrt[3]{b}\left(\sqrt[3]{a}-2\sqrt[3]{bx}\right)}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a+\sqrt[3]{bx}}\right)}{3a^{2/3}\sqrt[3]{b}} - \frac{x}{3b(a+bx^3)}}$$

↓ 27

$$\frac{\frac{3}{2}\sqrt[3]{a}\int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx + \frac{1}{2}\int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a+\sqrt[3]{bx}}\right)}{3a^{2/3}\sqrt[3]{b}} - \frac{x}{3b(a+bx^3)}}$$

↓ 1082

$$\frac{\frac{1}{2}\int \frac{\sqrt[3]{a}-2\sqrt[3]{bx}}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}}dx + \frac{{}_3f\left(\frac{1}{\left(1-2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2}d\left(1-2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)\right)}{\left(1-2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^{-3}\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a+\sqrt[3]{bx}}\right)}{3a^{2/3}\sqrt[3]{b}} - \frac{x}{3b(a+bx^3)}}$$

↓ 217

$$\frac{\frac{1}{2} \int \frac{\sqrt[3]{a-2\sqrt[3]{bx}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt[3]{b} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt[3]{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} - \frac{x}{3b(a+bx^3)}$$

↓ 1103

$$\frac{\frac{\log(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx+b^{2/3}x^2})}{2\sqrt[3]{b}} - \frac{\sqrt[3]{b} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt[3]{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} - \frac{x}{3b(a+bx^3)}$$

input `Int[x^3/(a + b*x^3)^2,x]`

output `-1/3*x/(b*(a + b*x^3)) + (Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])]/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3)))/(3*b)`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 750 $\text{Int}[(a_ + (b_ \cdot x)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3 \cdot \text{Rt}[a, 3]^2) \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x), x], x] + \text{Simp}[1/(3 \cdot \text{Rt}[a, 3]^2) \text{Int}[(2 \cdot \text{Rt}[a, 3] - \text{Rt}[b, 3] \cdot x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3] \cdot \text{Rt}[b, 3] \cdot x + \text{Rt}[b, 3]^2 \cdot x^2), x], x] /;$
 $\text{FreeQ}\{a, b\}, x]$

rule 817 $\text{Int}[(c_ \cdot x)^{m_} \cdot (a_ + (b_ \cdot x)^n)^{p_}, x_Symbol] \rightarrow \text{Simp}[c^{n-1} \cdot (c \cdot x)^{m-n+1} \cdot (a + b \cdot x^n)^{p+1} / (b \cdot n \cdot (p+1)), x] - \text{Simp}[c^n \cdot ((m-n+1)/(b \cdot n \cdot (p+1))) \text{Int}[(c \cdot x)^{m-n} \cdot (a + b \cdot x^n)^{p+1}, x], x] /;$
 $\text{FreeQ}\{a, b, c\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[m+1, n] \ \&\& \ ! \ \text{ILtQ}[(m+n \cdot (p+1)+1)/n, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

rule 1082 $\text{Int}[(a_ + (b_ \cdot x) + (c_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot S \text{implify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$
 $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$
 $\text{FreeQ}\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_ + (e_ \cdot x))/((a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$
 $\text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142 $\text{Int}[(d_ + (e_ \cdot x))/((a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.37

method	result	size
risch	$-\frac{x}{3b(bx^3+a)} + \frac{\sum_{R=\text{RootOf}(bZ^3+a)} \frac{\ln(x-R)}{-R^2}}{9b^2}$	43
default	$-\frac{x}{3b(bx^3+a)} + \frac{\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}}-\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{2}{3}}}}{3b} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}-1\right)}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}}$	112

```
input int(x^3/(b*x^3+a)^2,x,method=_RETURNVERBOSE)
```

```
output -1/3*x/b/(b*x^3+a)+1/9/b^2*sum(1/_R^2*ln(x-_R),_R=RootOf(_Z^3*b+a))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 391, normalized size of antiderivative = 3.40

$$\int \frac{x^3}{(a+bx^3)^2} dx$$

$$= \frac{6a^2bx - 3\sqrt{\frac{1}{3}(ab^2x^3 + a^2b)}\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}} \log\left(\frac{2abx^3 - 3(a^2b)^{\frac{1}{3}}ax - a^2 + 3\sqrt{\frac{1}{3}}\left(2abx^2 + (a^2b)^{\frac{2}{3}}x - (a^2b)^{\frac{1}{3}}a\right)\sqrt{-\frac{(a^2b)^{\frac{1}{3}}}{b}}}{bx^3+a}}{18(a^2b^3x^3 + a^3b^2)}\right)}{18(a^2b^3x^3 + a^3b^2)} + \frac{6a^2bx - 6\sqrt{\frac{1}{3}(ab^2x^3 + a^2b)}\sqrt{\frac{(a^2b)^{\frac{1}{3}}}{b}} \arctan\left(\frac{\sqrt{\frac{1}{3}}\left(2(a^2b)^{\frac{2}{3}}x - (a^2b)^{\frac{1}{3}}a\right)\sqrt{\frac{(a^2b)^{\frac{1}{3}}}{b}}}{a^2}\right) + (bx^3 + a)(a^2b)^{\frac{2}{3}} \log(a)}{18(a^2b^3x^3 + a^3b^2)}$$

input `integrate(x^3/(b*x^3+a)^2,x, algorithm="fricas")`

output `[-1/18*(6*a^2*b*x - 3*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt(-(a^2*b)^(1/3)/b) *log((2*a*b*x^3 - 3*(a^2*b)^(1/3)*a*x - a^2 + 3*sqrt(1/3)*(2*a*b*x^2 + (a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt(-(a^2*b)^(1/3)/b))/(b*x^3 + a)) + (b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) - 2*(b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^2*b^3*x^3 + a^3*b^2), -1/18*(6*a^2*b*x - 6*sqrt(1/3)*(a*b^2*x^3 + a^2*b)*sqrt((a^2*b)^(1/3)/b)*arctan(sqrt(1/3)*(2*(a^2*b)^(2/3)*x - (a^2*b)^(1/3)*a)*sqrt((a^2*b)^(1/3)/b)/a^2) + (b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x^2 - (a^2*b)^(2/3)*x + (a^2*b)^(1/3)*a) - 2*(b*x^3 + a)*(a^2*b)^(2/3)*log(a*b*x + (a^2*b)^(2/3)))/(a^2*b^3*x^3 + a^3*b^2)]`

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.34

$$\int \frac{x^3}{(a + bx^3)^2} dx = -\frac{x}{3ab + 3b^2x^3} + \text{RootSum}(729t^3a^2b^4 - 1, (t \mapsto t \log(9tab + x)))$$

input `integrate(x**3/(b*x**3+a)**2,x)`

output `-x/(3*a*b + 3*b**2*x**3) + RootSum(729*_t**3*a**2*b**4 - 1, Lambda(_t, _t*log(9*_t*a*b + x)))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.99

$$\int \frac{x^3}{(a + bx^3)^2} dx = -\frac{x}{3(b^2x^3 + ab)} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9b^2\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{18b^2\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{9b^2\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input `integrate(x^3/(b*x^3+a)^2,x, algorithm="maxima")`

output
$$-1/3*x/(b^2*x^3 + a*b) + 1/9*\sqrt{3}*\arctan(1/3*\sqrt{3}*(2*x - (a/b)^{1/3})/(a/b)^{1/3})/(b^2*(a/b)^{2/3}) - 1/18*\log(x^2 - x*(a/b)^{1/3} + (a/b)^{2/3})/(b^2*(a/b)^{2/3}) + 1/9*\log(x + (a/b)^{1/3})/(b^2*(a/b)^{2/3})$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.13

$$\int \frac{x^3}{(a + bx^3)^2} dx = -\frac{\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{9ab} - \frac{x}{3(bx^3 + a)b}$$

$$+ \frac{\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9ab^2}$$

$$+ \frac{\left(-ab^2\right)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{18ab^2}$$

input `integrate(x^3/(b*x^3+a)^2,x, algorithm="giac")`

output
$$-1/9*(-a/b)^{1/3}*\log(\text{abs}(x - (-a/b)^{1/3}))/(*b) - 1/3*x/((b*x^3 + a)*b)$$

$$+ 1/9*\sqrt{3}*(-a*b^2)^{1/3}*\arctan(1/3*\sqrt{3}*(2*x + (-a/b)^{1/3})/(-a/b)^{1/3})/(a*b^2) + 1/18*(-a*b^2)^{1/3}*\log(x^2 + x*(-a/b)^{1/3} + (-a/b)^{2/3})/(a*b^2)$$

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.94

$$\int \frac{x^3}{(a + bx^3)^2} dx = \frac{\ln(b^{1/3}x + a^{1/3})}{9a^{2/3}b^{4/3}} - \frac{x}{3b(bx^3 + a)}$$

$$+ \frac{\ln\left(bx + \frac{a^{1/3}b^{2/3}(-1+\sqrt{3}1i)}{2}\right)(-1 + \sqrt{3}1i)}{18a^{2/3}b^{4/3}}$$

$$- \frac{\ln\left(bx - \frac{a^{1/3}b^{2/3}(1+\sqrt{3}1i)}{2}\right)(1 + \sqrt{3}1i)}{18a^{2/3}b^{4/3}}$$

input `int(x^3/(a + b*x^3)^2,x)`output `log(b^(1/3)*x + a^(1/3))/(9*a^(2/3)*b^(4/3)) - x/(3*b*(a + b*x^3)) + (log(b*x + (a^(1/3)*b^(2/3)*(3^(1/2)*1i - 1))/2)*(3^(1/2)*1i - 1))/(18*a^(2/3)*b^(4/3)) - (log(b*x - (a^(1/3)*b^(2/3)*(3^(1/2)*1i + 1))/2)*(3^(1/2)*1i + 1))/(18*a^(2/3)*b^(4/3))`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.49

$$\int \frac{x^3}{(a + bx^3)^2} dx$$

$$= \frac{-2a^{\frac{4}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}}-2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) - 2a^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}}-2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) b x^3 - a^{\frac{4}{3}}\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) - a^{\frac{1}{3}}\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right)}{18b^{\frac{4}{3}}a(bx^3 + a)}$$

input `int(x^3/(b*x^3+a)^2,x)`

output

```
( - 2*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*
a - 2*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*
b*x**3 - a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a -
a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*b*x**3 + 2*a*
*(1/3)*log(a**(1/3) + b**(1/3)*x)*a + 2*a**(1/3)*log(a**(1/3) + b**(1/3)*x
)*b*x**3 - 6*b**(1/3)*a*x)/(18*b**(1/3)*a*b*(a + b*x**3))
```

3.71 $\int \frac{1}{x(a+bx^3)} dx$

Optimal result	466
Mathematica [A] (verified)	466
Rubi [A] (verified)	467
Maple [A] (verified)	468
Fricas [A] (verification not implemented)	469
Sympy [A] (verification not implemented)	469
Maxima [A] (verification not implemented)	469
Giac [A] (verification not implemented)	470
Mupad [B] (verification not implemented)	470
Reduce [B] (verification not implemented)	470

Optimal result

Integrand size = 13, antiderivative size = 21

$$\int \frac{1}{x(a+bx^3)} dx = \frac{\log\left(\frac{x^3}{a+bx^3}\right)}{3a}$$

output `1/3*ln(x^3/(b*x^3+a))/a`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{1}{x(a+bx^3)} dx = \frac{\log(x)}{a} - \frac{\log(a+bx^3)}{3a}$$

input `Integrate[1/(x*(a + b*x^3)),x]`

output `Log[x]/a - Log[a + b*x^3]/(3*a)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {798, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x(a+bx^3)} dx \\
 & \quad \downarrow 798 \\
 & \frac{1}{3} \int \frac{1}{x^3(bx^3+a)} dx^3 \\
 & \quad \downarrow 47 \\
 & \frac{1}{3} \left(\frac{\int \frac{1}{x^3} dx^3}{a} - \frac{b \int \frac{1}{bx^3+a} dx^3}{a} \right) \\
 & \quad \downarrow 14 \\
 & \frac{1}{3} \left(\frac{\log(x^3)}{a} - \frac{b \int \frac{1}{bx^3+a} dx^3}{a} \right) \\
 & \quad \downarrow 16 \\
 & \frac{1}{3} \left(\frac{\log(x^3)}{a} - \frac{\log(a+bx^3)}{a} \right)
 \end{aligned}$$

input `Int[1/(x*(a + b*x^3)),x]`

output `(Log[x^3]/a - Log[a + b*x^3]/a)/3`

Definitions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{\ln(bx^3+a)}{3a} + \frac{\ln(x)}{a}$	21
norman	$-\frac{\ln(bx^3+a)}{3a} + \frac{\ln(x)}{a}$	21
risch	$-\frac{\ln(bx^3+a)}{3a} + \frac{\ln(x)}{a}$	21
parallelrisc	$\frac{3\ln(x) - \ln(bx^3+a)}{3a}$	21

input `int(1/x/(b*x^3+a),x,method=_RETURNVERBOSE)`

output `-1/3/a*ln(b*x^3+a)+1/a*ln(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \frac{1}{x(a+bx^3)} dx = -\frac{\log(bx^3+a) - 3\log(x)}{3a}$$

input `integrate(1/x/(b*x^3+a),x, algorithm="fricas")`output `-1/3*(log(b*x^3 + a) - 3*log(x))/a`**Sympy [A] (verification not implemented)**

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1}{x(a+bx^3)} dx = \frac{\log(x)}{a} - \frac{\log(\frac{a}{b} + x^3)}{3a}$$

input `integrate(1/x/(b*x**3+a),x)`output `log(x)/a - log(a/b + x**3)/(3*a)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{1}{x(a+bx^3)} dx = -\frac{\log(bx^3+a)}{3a} + \frac{\log(x^3)}{3a}$$

input `integrate(1/x/(b*x^3+a),x, algorithm="maxima")`output `-1/3*log(b*x^3 + a)/a + 1/3*log(x^3)/a`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{1}{x(a+bx^3)} dx = -\frac{\log(|bx^3+a|)}{3a} + \frac{\log(|x|)}{a}$$

input `integrate(1/x/(b*x^3+a),x, algorithm="giac")`output `-1/3*log(abs(b*x^3 + a))/a + log(abs(x))/a`**Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \frac{1}{x(a+bx^3)} dx = -\frac{\ln(bx^3+a) - 3 \ln(x)}{3a}$$

input `int(1/(x*(a + b*x^3)),x)`output `-(log(a + b*x^3) - 3*log(x))/(3*a)`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.14

$$\int \frac{1}{x(a+bx^3)} dx = \frac{-\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) - \log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) + 3 \log(x)}{3a}$$

input `int(1/x/(b*x^3+a),x)`output `(- log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2) - log(a**(1/3) + b**
**(1/3)*x) + 3*log(x))/(3*a)`

3.72 $\int \frac{1}{x^2(a+bx^3)} dx$

Optimal result	471
Mathematica [A] (verified)	471
Rubi [A] (verified)	472
Maple [C] (verified)	476
Fricas [A] (verification not implemented)	476
Sympy [A] (verification not implemented)	477
Maxima [A] (verification not implemented)	477
Giac [A] (verification not implemented)	478
Mupad [B] (verification not implemented)	478
Reduce [B] (verification not implemented)	479

Optimal result

Integrand size = 13, antiderivative size = 110

$$\int \frac{1}{x^2(a+bx^3)} dx = -\frac{1}{ax} + \frac{-\sqrt{3} \arctan\left(\frac{-\sqrt[3]{\frac{a}{b}}+2x}{\sqrt{3}\sqrt[3]{\frac{a}{b}}}\right) + \frac{1}{2} \log\left(\frac{\left(\sqrt[3]{\frac{a}{b}}+x\right)^2}{\left(\frac{a}{b}\right)^{2/3} - \sqrt[3]{\frac{a}{b}}x+x^2}\right)}{3a\sqrt[3]{\frac{a}{b}}}$$

output

```
-1/a/x+1/3*(-3^(1/2)*arctan(1/3*(-(a/b)^(1/3)+2*x)*3^(1/2)/(a/b)^(1/3))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2)))/a/(a/b)^(1/3)
```

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.04

$$\int \frac{1}{x^2(a+bx^3)} dx = \frac{-6\sqrt[3]{a} + 2\sqrt{3}\sqrt[3]{bx} \arctan\left(\frac{1-2\sqrt[3]{bx}}{\sqrt{3}\sqrt[3]{a}}\right) + 2\sqrt[3]{bx} \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right) - \sqrt[3]{bx} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x^2\right)}{6a^{4/3}x}$$

input `Integrate[1/(x^2*(a + b*x^3)),x]`

output $(-6*a^{1/3} + 2*\text{Sqrt}[3]*b^{1/3}*x*\text{ArcTan}[(1 - (2*b^{1/3}*x)/a^{1/3})/\text{Sqrt}[3]] + 2*b^{1/3}*x*\text{Log}[a^{1/3} + b^{1/3}*x] - b^{1/3}*x*\text{Log}[a^{2/3} - a^{1/3}/3*b^{1/3}*x + b^{2/3}*x^2])/(6*a^{4/3}*x)$

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.19, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {847, 821, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2(a + bx^3)} dx$$

$$\downarrow 847$$

$$-\frac{b \int \frac{x}{bx^3+a} dx}{a} - \frac{1}{ax}$$

$$\downarrow 821$$

$$-\frac{b \left(\frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} \right)}{a} - \frac{1}{ax}$$

$$\downarrow 16$$

$$-\frac{b \left(\frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right)}{a} - \frac{1}{ax}$$

$$\downarrow 1142$$

$$b \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx + \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a-2} \sqrt[3]{b_x})}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{2 \sqrt[3]{b}}}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 \sqrt[3]{ab^{2/3}}} \right) \frac{1}{ax}$$

a

25

$$b \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx - \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a-2} \sqrt[3]{b_x})}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{2 \sqrt[3]{b}}}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 \sqrt[3]{ab^{2/3}}} \right) \frac{1}{ax}$$

a

27

$$b \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx - \frac{1}{2} \int \frac{\sqrt[3]{a-2} \sqrt[3]{b_x}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 \sqrt[3]{ab^{2/3}}} \right) \frac{1}{ax}$$

a

1082

$$b \left(\frac{\frac{3 \int \frac{1}{\left(1 - \frac{2 \sqrt[3]{b_x}}{\sqrt[3]{a}}\right)^2} d\left(1 - \frac{2 \sqrt[3]{b_x}}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}} - \frac{1}{2} \int \frac{\sqrt[3]{a-2} \sqrt[3]{b_x}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 \sqrt[3]{ab^{2/3}}} \right) \frac{1}{ax}$$

a

217

$$\begin{array}{c}
 \left(\frac{b \left(\frac{-\frac{1}{2} \int \frac{\sqrt[3]{a-2}\sqrt[3]{b}x}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{b}x + a^{2/3}} dx - \frac{\sqrt[3]{3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}} \right) - \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3\sqrt[3]{ab^{2/3}}}}{3\sqrt[3]{a}\sqrt[3]{b}} \right)}{a} - \frac{1}{ax} \\
 \downarrow \text{1103} \\
 \left(\frac{b \left(\frac{\frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{2\sqrt[3]{b}} - \frac{\sqrt[3]{3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}} \right) - \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3\sqrt[3]{ab^{2/3}}}}{3\sqrt[3]{a}\sqrt[3]{b}} \right)}{a} - \frac{1}{ax}
 \end{array}$$

input `Int[1/(x^2*(a + b*x^3)),x]`

output `-(1/(a*x)) - (b*(-1/3*Log[a^(1/3) + b^(1/3)*x]/(a^(1/3)*b^(2/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])]/b^(1/3)) + Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(1/3)*b^(1/3)))/a`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[((a_) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 821 $\text{Int}[(x_)/((a_) + (b_*)(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{-1} \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]) \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 847 $\text{Int}[((c_*)(x_)^m)*((a_) + (b_*)(x_)^n)^p, x_Symbol] \rightarrow \text{Simp}[(c*x)^{m+1}*((a + b*x^n)^{p+1}/(a*c*(m+1))), x] - \text{Simp}[b*(m+n*(p+1)+1)/(a*c^n*(m+1)) \text{Int}[(c*x)^{m+n}*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$
- rule 1082 $\text{Int}[((a_) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[((d_) + (e_*)(x_))/((a_) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[((d_) + (e_*)(x_))/((a_) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.48

method	result	size
risch	$-\frac{1}{ax} + \frac{\left(\sum_{R=\text{RootOf}(a^4-Z^3-b)} -R \ln((-4-R^3 a^4+3b)x-a^3-R^2) \right)}{3}$	53
default	$-\frac{1}{ax} - \frac{\left(\begin{aligned} & \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3b\left(\frac{a}{b}\right)^{\frac{1}{3}}} \end{aligned} \right)}{a} b$	106

```
input int(1/x^2/(b*x^3+a),x,method=_RETURNVERBOSE)
```

```
output -1/a/x+1/3*sum(_R*ln((-4*_R^3*a^4+3*b)*x-a^3*_R^2),_R=RootOf(_Z^3*a^4-b))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.94

$$\int \frac{1}{x^2(a+bx^3)} dx = \frac{2\sqrt{3}x\left(\frac{b}{a}\right)^{\frac{1}{3}} \arctan\left(\frac{2}{3}\sqrt{3}x\left(\frac{b}{a}\right)^{\frac{1}{3}} - \frac{1}{3}\sqrt{3}\right) + x\left(\frac{b}{a}\right)^{\frac{1}{3}} \log\left(bx^2 - ax\left(\frac{b}{a}\right)^{\frac{2}{3}} + a\left(\frac{b}{a}\right)^{\frac{1}{3}}\right) - 2x\left(\frac{b}{a}\right)^{\frac{1}{3}} \log\left(bx + a\right)}{6ax}$$

```
input integrate(1/x^2/(b*x^3+a),x, algorithm="fricas")
```

output

```
-1/6*(2*sqrt(3)*x*(b/a)^(1/3)*arctan(2/3*sqrt(3)*x*(b/a)^(1/3) - 1/3*sqrt(3)) + x*(b/a)^(1/3)*log(b*x^2 - a*x*(b/a)^(2/3) + a*(b/a)^(1/3)) - 2*x*(b/a)^(1/3)*log(b*x + a*(b/a)^(2/3)) + 6)/(a*x)
```

Sympy [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.26

$$\int \frac{1}{x^2(a+bx^3)} dx = \text{RootSum}\left(27t^3a^4 - b, \left(t \mapsto t \log\left(\frac{9t^2a^3}{b} + x\right)\right)\right) - \frac{1}{ax}$$

input

```
integrate(1/x**2/(b*x**3+a),x)
```

output

```
RootSum(27*_t**3*a**4 - b, Lambda(_t, _t*log(9*_t**2*a**3/b + x))) - 1/(a*x)
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.96

$$\int \frac{1}{x^2(a+bx^3)} dx = -\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3a\left(\frac{a}{b}\right)^{\frac{1}{3}}} - \frac{\log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6a\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{\log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3a\left(\frac{a}{b}\right)^{\frac{1}{3}}} - \frac{1}{ax}$$

input

```
integrate(1/x^2/(b*x^3+a),x, algorithm="maxima")
```

output

```
-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(a*(a/b)^(1/3)) - 1/6*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(a*(a/b)^(1/3)) + 1/3*log(x + (a/b)^(1/3))/(a*(a/b)^(1/3)) - 1/(a*x)
```

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.10

$$\int \frac{1}{x^2(a+bx^3)} dx = \frac{b\left(-\frac{a}{b}\right)^{\frac{2}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{3a^2} + \frac{\sqrt{3}(-ab^2)^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3a^2b}$$

$$- \frac{(-ab^2)^{\frac{2}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6a^2b} - \frac{1}{ax}$$

input `integrate(1/x^2/(b*x^3+a),x, algorithm="giac")`output `1/3*b*(-a/b)^(2/3)*log(abs(x - (-a/b)^(1/3)))/a^2 + 1/3*sqrt(3)*(-a*b^2)^(2/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/(a^2*b) - 1/6*(-a*b^2)^(2/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/(a^2*b) - 1/(a*x)`**Mupad [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.93

$$\int \frac{1}{x^2(a+bx^3)} dx = \frac{b^{1/3} \ln(b^{1/3}x + a^{1/3})}{3a^{4/3}} - \frac{1}{ax}$$

$$- \frac{b^{1/3} \ln(4b^{1/3}x - 2a^{1/3} + \sqrt{3}a^{1/3}2i) \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{3a^{4/3}}$$

$$+ \frac{b^{1/3} \ln(4b^{1/3}x - 2a^{1/3} - \sqrt{3}a^{1/3}2i) \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)}{a^{4/3}}$$

input `int(1/(x^2*(a + b*x^3)),x)`output `(b^(1/3)*log(b^(1/3)*x + a^(1/3)))/(3*a^(4/3)) - 1/(a*x) - (b^(1/3)*log(3^(1/2)*a^(1/3)*2i + 4*b^(1/3)*x - 2*a^(1/3))*((3^(1/2)*1i)/2 + 1/2))/(3*a^(4/3)) + (b^(1/3)*log(4*b^(1/3)*x - 3^(1/2)*a^(1/3)*2i - 2*a^(1/3))*((3^(1/2)*1i)/6 - 1/6))/a^(4/3)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.76

$$\int \frac{1}{x^2 (a + bx^3)} dx$$

$$= \frac{2\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) bx - 6b^{\frac{2}{3}}a^{\frac{1}{3}} - \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) bx + 2\log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right) bx}{6b^{\frac{2}{3}}a^{\frac{4}{3}}x}$$

input `int(1/x^2/(b*x^3+a), x)`output `(2*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*b*x - 6*b**(2/3)*a**(1/3) - log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*b*x + 2*log(a**(1/3) + b**(1/3)*x)*b*x)/(6*b**(2/3)*a**(1/3)*a*x)`

3.73 $\int \frac{1}{x^3(a+bx^3)} dx$

Optimal result	480
Mathematica [A] (verified)	481
Rubi [A] (verified)	481
Maple [C] (verified)	485
Fricas [A] (verification not implemented)	485
Sympy [A] (verification not implemented)	486
Maxima [A] (verification not implemented)	486
Giac [A] (verification not implemented)	487
Mupad [B] (verification not implemented)	487
Reduce [B] (verification not implemented)	488

Optimal result

Integrand size = 13, antiderivative size = 106

$$\int \frac{1}{x^3(a+bx^3)} dx = -\frac{1}{2ax^2} - \frac{\sqrt[3]{\frac{a}{b}}b \left(\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{2\sqrt[3]{\frac{a}{b}}-x} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}}+x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}}x+x^2} \right) \right)}{3a^2}$$

```
output -1/2/a/x^2-1/3*(a/b)^(1/3)*b*(3^(1/2)*arctan(3^(1/2)*x/(2*(a/b)^(1/3)-x))+
1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2))/a^2
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.12

$$\int \frac{1}{x^3(a+bx^3)} dx$$

$$= \frac{-3a^{2/3} + 2\sqrt{3}b^{2/3}x^2 \arctan\left(\frac{1 - 2\sqrt[3]{bx}}{\sqrt[3]{a}}\right) - 2b^{2/3}x^2 \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right) + b^{2/3}x^2 \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + b^{2/3}x\right)}{6a^{5/3}x^2}$$

input `Integrate[1/(x^3*(a + b*x^3)),x]`

output `(-3*a^(2/3) + 2*Sqrt[3]*b^(2/3)*x^2*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/Sqrt[3]] - 2*b^(2/3)*x^2*Log[a^(1/3) + b^(1/3)*x] + b^(2/3)*x^2*Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/(6*a^(5/3)*x^2)`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.21, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {847, 750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3(a+bx^3)} dx$$

$$\downarrow 847$$

$$-\frac{b \int \frac{1}{bx^3+a} dx}{a} - \frac{1}{2ax^2}$$

$$\downarrow 750$$

$$\frac{b \left(\frac{\int \frac{2\sqrt[3]{a}-\sqrt[3]{b}x}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{b}x+\sqrt[3]{a}} dx}{3a^{2/3}} \right)}{a} - \frac{1}{2ax^2}$$

↓ 16

$$\frac{b \left(\frac{\int \frac{2\sqrt[3]{a}-\sqrt[3]{b}x}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a}+\sqrt[3]{b}x)}{3a^{2/3}\sqrt[3]{b}} \right)}{a} - \frac{1}{2ax^2}$$

↓ 1142

$$\frac{b \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx - \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{b}x)}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a}+\sqrt[3]{b}x)}{3a^{2/3}\sqrt[3]{b}} \right)}{a} - \frac{1}{2ax^2}$$

↓ 25

$$\frac{b \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx + \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{b}x)}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{2\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a}+\sqrt[3]{b}x)}{3a^{2/3}\sqrt[3]{b}} \right)}{a} - \frac{1}{2ax^2}$$

↓ 27

$$\frac{b \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a}-2\sqrt[3]{b}x}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a}+\sqrt[3]{b}x)}{3a^{2/3}\sqrt[3]{b}} \right)}{a} - \frac{1}{2ax^2}$$

↓ 1082

$$b \left(\frac{\frac{1}{2} \int \frac{\sqrt[3]{a-2}\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{1}{\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} d\left(1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{-3\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) - \frac{1}{2ax^2}$$

a

217

$$b \left(\frac{\frac{1}{2} \int \frac{\sqrt[3]{a-2}\sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) - \frac{1}{2ax^2}$$

a

1103

$$b \left(\frac{\frac{\log(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx+b^{2/3}x^2})}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3}\sqrt[3]{b}} \right) - \frac{1}{2ax^2}$$

a

input

`Int[1/(x^3*(a + b*x^3)),x]`

output

`-1/2*1/(a*x^2) - (b*(Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3])/b^(1/3)) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(2/3))))/a`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 847 $\text{Int}[(c_)*(x_)^m*((a_)+(b_)*(x_)^n)^p, x_Symbol] \rightarrow \text{Simp}[(c*x)^{m+1}*((a + b*x^n)^{p+1}/(a*c*(m+1))), x] - \text{Simp}[b*((m+n*(p+1)+1)/(a*c^n*(m+1)) \text{ Int}[(c*x)^{m+n}*(a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142

```
Int[((d._) + (e._)*(x_))/((a_) + (b._)*(x_) + (c._)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.51

method	result	size
risch	$-\frac{1}{2ax^2} + \frac{\left(\sum_{-R=\text{RootOf}(a^5-Z^3+b^2)} -R \ln((-4-R^3a^5-3b^2)x-a^2b-R) \right)}{3}$	54
default	$-\frac{1}{2ax^2} - \frac{\left(\frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{\ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{6b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{\left(\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2x}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}-1\right)}{\frac{\left(\frac{a}{b}\right)^{\frac{1}{3}}}{3}} \right) \right)}{3b\left(\frac{a}{b}\right)^{\frac{2}{3}}} \right)}{a} b$	106

input

```
int(1/x^3/(b*x^3+a),x,method=_RETURNVERBOSE)
```

output

```
-1/2/a/x^2+1/3*sum(_R*ln((-4*_R^3*a^5-3*b^2)*x-a^2*b*_R),_R=RootOf(_Z^3*a^5+b^2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.35

$$\int \frac{1}{x^3(a+bx^3)} dx$$

$$= \frac{2\sqrt{3}x^2\left(-\frac{b^2}{a^2}\right)^{\frac{1}{3}} \arctan\left(\frac{2\sqrt{3}ax\left(-\frac{b^2}{a^2}\right)^{\frac{2}{3}}-\sqrt{3}b}{3b}\right) - x^2\left(-\frac{b^2}{a^2}\right)^{\frac{1}{3}} \log\left(b^2x^2+abx\left(-\frac{b^2}{a^2}\right)^{\frac{1}{3}}+a^2\left(-\frac{b^2}{a^2}\right)^{\frac{2}{3}}\right) + 2x^2}{6ax^2}$$

input `integrate(1/x^3/(b*x^3+a),x, algorithm="fricas")`

output
$$\frac{1}{6} \cdot (2 \cdot \sqrt{3}) \cdot x^2 \cdot (-b^2/a^2)^{1/3} \cdot \arctan(1/3 \cdot (2 \cdot \sqrt{3}) \cdot a \cdot x \cdot (-b^2/a^2)^{1/3} - \sqrt{3} \cdot b) / b - x^2 \cdot (-b^2/a^2)^{1/3} \cdot \log(b^2 \cdot x^2 + a \cdot b \cdot x \cdot (-b^2/a^2)^{1/3} + a^2 \cdot (-b^2/a^2)^{2/3}) + 2 \cdot x^2 \cdot (-b^2/a^2)^{1/3} \cdot \log(b \cdot x - a \cdot (-b^2/a^2)^{1/3}) - 3) / (a \cdot x^2)$$

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.30

$$\int \frac{1}{x^3(a+bx^3)} dx = \text{RootSum} \left(27t^3a^5 + b^2, \left(t \mapsto t \log \left(-\frac{3ta^2}{b} + x \right) \right) \right) - \frac{1}{2ax^2}$$

input `integrate(1/x**3/(b*x**3+a),x)`

output `RootSum(27*_t**3*a**5 + b**2, Lambda(_t, _t*log(-3*_t*a**2/b + x))) - 1/(2*a*x**2)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00

$$\int \frac{1}{x^3(a+bx^3)} dx = -\frac{\sqrt{3} \arctan \left(\frac{\sqrt{3} \left(2x - \left(\frac{a}{b} \right)^{1/3} \right)}{3 \left(\frac{a}{b} \right)^{1/3}} \right)}{3a \left(\frac{a}{b} \right)^{2/3}} + \frac{\log \left(x^2 - x \left(\frac{a}{b} \right)^{1/3} + \left(\frac{a}{b} \right)^{2/3} \right)}{6a \left(\frac{a}{b} \right)^{2/3}} - \frac{\log \left(x + \left(\frac{a}{b} \right)^{1/3} \right)}{3a \left(\frac{a}{b} \right)^{2/3}} - \frac{1}{2ax^2}$$

input `integrate(1/x^3/(b*x^3+a),x, algorithm="maxima")`

output

```
-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(a*(a/b)^(2/3)) + 1/6*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(a*(a/b)^(2/3)) - 1/3*log(x + (a/b)^(1/3))/(a*(a/b)^(2/3)) - 1/2/(a*x^2)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.08

$$\int \frac{1}{x^3(a+bx^3)} dx = \frac{b\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{3a^2} - \frac{\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3a^2} - \frac{(-ab^2)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6a^2} - \frac{1}{2ax^2}$$

input

```
integrate(1/x^3/(b*x^3+a),x, algorithm="giac")
```

output

```
1/3*b*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/a^2 - 1/3*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/a^2 - 1/6*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/a^2 - 1/2/(a*x^2)
```

Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.21

$$\int \frac{1}{x^3(a+bx^3)} dx = \frac{b^{2/3} \ln\left((-a)^{7/3} - a^2 b^{1/3} x\right)}{3(-a)^{5/3}} - \frac{1}{2ax^2} - \frac{b^{2/3} \ln\left(3a^2 b^3 x + 3(-a)^{7/3} b^{8/3} \left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)\right) \left(\frac{1}{2} + \frac{\sqrt{3}li}{2}\right)}{3(-a)^{5/3}} + \frac{b^{2/3} \ln\left(3a^2 b^3 x - 9(-a)^{7/3} b^{8/3} \left(-\frac{1}{6} + \frac{\sqrt{3}li}{6}\right)\right) \left(-\frac{1}{6} + \frac{\sqrt{3}li}{6}\right)}{(-a)^{5/3}}$$

input

```
int(1/(x^3*(a + b*x^3)),x)
```


output

$$\begin{aligned} & (b^{2/3} \log((-a)^{7/3} - a^2 b^{1/3} x)) / (3(-a)^{5/3}) - 1/(2a x^2) - (\\ & b^{2/3} \log(3a^2 b^3 x + 3(-a)^{7/3} b^{8/3} ((3^{1/2} i)/2 + 1/2)) * ((3 \\ & ^{1/2} i)/2 + 1/2)) / (3(-a)^{5/3}) + (b^{2/3} \log(3a^2 b^3 x - 9(-a)^{7/3} \\ & b^{8/3} ((3^{1/2} i)/6 - 1/6)) * ((3^{1/2} i)/6 - 1/6)) / (-a)^{5/3} \end{aligned}$$
Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.91

$$\int \frac{1}{x^3 (a + bx^3)} dx$$

$$= \frac{2a^{1/3} \sqrt{3} \operatorname{atan}\left(\frac{a^{1/3} - 2b^{1/3} x}{a^{1/3} \sqrt{3}}\right) b x^2 + a^{1/3} \log\left(a^{2/3} - b^{1/3} a^{1/3} x + b^{2/3} x^2\right) b x^2 - 2a^{1/3} \log\left(a^{1/3} + b^{1/3} x\right) b x^2 - 3b^{1/3} a}{6b^{1/3} a^2 x^2}$$

input

`int(1/x^3/(b*x^3+a),x)`

output

$$\begin{aligned} & (2*a^{1/3}*\sqrt{3}*\operatorname{atan}((a^{1/3} - 2*b^{1/3}*x)/(a^{1/3}*\sqrt{3}))*b*x \\ & **2 + a^{1/3}*\log(a^{2/3} - b^{1/3}*a^{1/3}*x + b^{2/3}*x**2)*b*x**2 \\ & - 2*a^{1/3}*\log(a^{1/3} + b^{1/3}*x)*b*x**2 - 3*b^{1/3}*a)/(6*b^{1/3} \\ & *a**2*x**2) \end{aligned}$$

3.74 $\int \frac{1}{x(a+bx^3)^2} dx$

Optimal result	489
Mathematica [A] (verified)	489
Rubi [A] (verified)	490
Maple [A] (verified)	491
Fricas [A] (verification not implemented)	491
Sympy [A] (verification not implemented)	492
Maxima [A] (verification not implemented)	492
Giac [A] (verification not implemented)	492
Mupad [B] (verification not implemented)	493
Reduce [B] (verification not implemented)	493

Optimal result

Integrand size = 13, antiderivative size = 38

$$\int \frac{1}{x(a+bx^3)^2} dx = \frac{1}{3a(a+bx^3)} + \frac{\log\left(\frac{x^3}{a+bx^3}\right)}{3a^2}$$

output $1/3/a/(b*x^3+a)+1/3*\ln(x^3/(b*x^3+a))/a^2$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{1}{x(a+bx^3)^2} dx = \frac{\frac{a}{a+bx^3} + 3 \log(x) - \log(a+bx^3)}{3a^2}$$

input `Integrate[1/(x*(a + b*x^3)^2),x]`

output $(a/(a + b*x^3) + 3*\text{Log}[x] - \text{Log}[a + b*x^3])/(3*a^2)$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {798, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x(a+bx^3)^2} dx$$

$$\downarrow 798$$

$$\frac{1}{3} \int \frac{1}{x^3(bx^3+a)^2} dx^3$$

$$\downarrow 54$$

$$\frac{1}{3} \int \left(-\frac{b}{a^2(bx^3+a)} - \frac{b}{a(bx^3+a)^2} + \frac{1}{a^2x^3} \right) dx^3$$

$$\downarrow 2009$$

$$\frac{1}{3} \left(-\frac{\log(a+bx^3)}{a^2} + \frac{\log(x^3)}{a^2} + \frac{1}{a(a+bx^3)} \right)$$

input `Int[1/(x*(a + b*x^3)^2),x]`

output `(1/(a*(a + b*x^3)) + Log[x^3]/a^2 - Log[a + b*x^3]/a^2)/3`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst
[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a,
b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

method	result	size
risch	$\frac{1}{3a(bx^3+a)} + \frac{\ln(x)}{a^2} - \frac{\ln(bx^3+a)}{3a^2}$	35
norman	$-\frac{bx^3}{3a^2(bx^3+a)} + \frac{\ln(x)}{a^2} - \frac{\ln(bx^3+a)}{3a^2}$	39
default	$-\frac{b\left(-\frac{a}{b(bx^3+a)} + \frac{\ln(bx^3+a)}{b}\right)}{3a^2} + \frac{\ln(x)}{a^2}$	42
parallelrisc	$\frac{3\ln(x)x^3b - \ln(bx^3+a)x^3b - bx^3 + 3\ln(x)a - \ln(bx^3+a)a}{3a^2(bx^3+a)}$	60

input `int(1/x/(b*x^3+a)^2,x,method=_RETURNVERBOSE)`

output `1/3/a/(b*x^3+a)+1/a^2*ln(x)-1/3/a^2*ln(b*x^3+a)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.24

$$\int \frac{1}{x(a+bx^3)^2} dx = -\frac{(bx^3+a)\log(bx^3+a) - 3(bx^3+a)\log(x) - a}{3(a^2bx^3+a^3)}$$

input `integrate(1/x/(b*x^3+a)^2,x, algorithm="fricas")`

output `-1/3*((b*x^3 + a)*log(b*x^3 + a) - 3*(b*x^3 + a)*log(x) - a)/(a^2*b*x^3 + a^3)`

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \frac{1}{x(a+bx^3)^2} dx = \frac{1}{3a^2 + 3abx^3} + \frac{\log(x)}{a^2} - \frac{\log\left(\frac{a}{b} + x^3\right)}{3a^2}$$

input `integrate(1/x/(b*x**3+a)**2,x)`output `1/(3*a**2 + 3*a*b*x**3) + log(x)/a**2 - log(a/b + x**3)/(3*a**2)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.97

$$\int \frac{1}{x(a+bx^3)^2} dx = \frac{1}{3(abx^3 + a^2)} - \frac{\log(bx^3 + a)}{3a^2} + \frac{\log(x^3)}{3a^2}$$

input `integrate(1/x/(b*x^3+a)^2,x, algorithm="maxima")`output `1/3/(a*b*x^3 + a^2) - 1/3*log(b*x^3 + a)/a^2 + 1/3*log(x^3)/a^2`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.18

$$\int \frac{1}{x(a+bx^3)^2} dx = -\frac{\log(|bx^3 + a|)}{3a^2} + \frac{\log(|x|)}{a^2} + \frac{bx^3 + 2a}{3(bx^3 + a)a^2}$$

input `integrate(1/x/(b*x^3+a)^2,x, algorithm="giac")`output `-1/3*log(abs(b*x^3 + a))/a^2 + log(abs(x))/a^2 + 1/3*(b*x^3 + 2*a)/((b*x^3 + a)*a^2)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \frac{1}{x(a+bx^3)^2} dx = \frac{\ln(x)}{a^2} + \frac{1}{3a(bx^3+a)} - \frac{\ln(bx^3+a)}{3a^2}$$

input `int(1/(x*(a + b*x^3)^2),x)`output `log(x)/a^2 + 1/(3*a*(a + b*x^3)) - log(a + b*x^3)/(3*a^2)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.00

$$\int \frac{1}{x(a+bx^3)^2} dx = \frac{-\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right)a - \log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right)bx^3 - \log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right)a - \log\left(a^{\frac{1}{3}} + b^{\frac{1}{3}}x\right)bx^3}{3a^2(bx^3+a)}$$

input `int(1/x/(b*x^3+a)^2,x)`output `(- log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a - log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*b*x**3 - log(a**(1/3) + b**(1/3)*x)*a - log(a**(1/3) + b**(1/3)*x)*b*x**3 + 3*log(x)*a + 3*log(x)*b*x**3 - b*x**3)/(3*a**2*(a + b*x**3))`

3.75 $\int \frac{1}{x^2(a+bx^3)^2} dx$

Optimal result	494
Mathematica [A] (verified)	495
Rubi [A] (verified)	495
Maple [C] (verified)	501
Fricas [A] (verification not implemented)	501
Sympy [A] (verification not implemented)	502
Maxima [A] (verification not implemented)	502
Giac [A] (verification not implemented)	503
Mupad [B] (verification not implemented)	503
Reduce [B] (verification not implemented)	504

Optimal result

Integrand size = 13, antiderivative size = 132

$$\int \frac{1}{x^2(a+bx^3)^2} dx = \frac{-\frac{1}{ax} - \frac{4bx^2}{3a^2}}{a+bx^3} + \frac{4 \left(-\sqrt{3} \arctan \left(\frac{-\sqrt[3]{\frac{a}{b}} + 2x}{\sqrt{3} \sqrt[3]{\frac{a}{b}}} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}} + x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}} x + x^2} \right) \right)}{9a^2 \sqrt[3]{\frac{a}{b}}}$$

output

```
(-1/a/x-4/3*b*x^2/a^2)/(b*x^3+a)+4/9*(-3^(1/2)*arctan(1/3*(-(a/b)^(1/3)+2*x)*3^(1/2)/(a/b)^(1/3))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2)))/a^2/(a/b)^(1/3)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 131, normalized size of antiderivative = 0.99

$$\int \frac{1}{x^2 (a + bx^3)^2} dx$$

$$= \frac{-\frac{9\sqrt[3]{a}}{x} - \frac{3\sqrt[3]{a}bx^2}{a+bx^3} + 4\sqrt{3}\sqrt[3]{b} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right) + 4\sqrt[3]{b} \log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right) - 2\sqrt[3]{b} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^2\right)}{9a^{7/3}}$$

input `Integrate[1/(x^2*(a + b*x^3)^2),x]`output `((-9*a^(1/3))/x - (3*a^(1/3)*b*x^2)/(a + b*x^3) + 4*Sqrt[3]*b^(1/3)*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/Sqrt[3]] + 4*b^(1/3)*Log[a^(1/3) + b^(1/3)*x] - 2*b^(1/3)*Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/(9*a^(7/3))`**Rubi [A] (verified)**Time = 0.33 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.20, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.769$, Rules used = {819, 847, 821, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 (a + bx^3)^2} dx$$

$$\downarrow \text{819}$$

$$\frac{4 \int \frac{1}{x^2 (bx^3 + a)} dx}{3a} + \frac{1}{3ax (a + bx^3)}$$

$$\downarrow \text{847}$$

$$\frac{4 \left(-\frac{b \int \frac{x}{bx^3 + a} dx}{a} - \frac{1}{ax} \right)}{3a} + \frac{1}{3ax (a + bx^3)}$$

$$\begin{aligned}
 & \downarrow 821 \\
 & \left(\frac{4}{3a} \left(\frac{b}{a} \left(\frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\int \frac{1}{\sqrt[3]{bx} + \sqrt[3]{a}} dx}{3\sqrt[3]{a}\sqrt[3]{b}} \right) - \frac{1}{ax} \right) + \frac{1}{3ax(a+bx^3)} \right) \\
 & \downarrow 16 \\
 & \left(\frac{4}{3a} \left(\frac{b}{a} \left(\frac{\int \frac{\sqrt[3]{bx} + \sqrt[3]{a}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right) - \frac{1}{ax} \right) + \frac{1}{3ax(a+bx^3)} \right) \\
 & \downarrow 1142 \\
 & \left(\frac{4}{3a} \left(\frac{b}{a} \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx}{2\sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3\sqrt[3]{ab^{2/3}}} \right) - \frac{1}{ax} \right) + \frac{1}{3ax(a+bx^3)} \right) \\
 & \downarrow 25
 \end{aligned}$$

$$4 \left(\frac{b \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx - \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a-2} \sqrt[3]{b_x})}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{2 \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 \sqrt[3]{ab^{2/3}}} \right)}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{1}{ax} \right) +$$

$$\frac{3a_1}{3ax(a+bx^3)}$$

↓ 27

$$4 \left(\frac{b \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx - \frac{1}{2} \int \frac{\sqrt[3]{a-2} \sqrt[3]{b_x}}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{b_{x+a^{2/3}}}} dx}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{\log(\sqrt[3]{a} + \sqrt[3]{b_x})}{3 \sqrt[3]{ab^{2/3}}} \right)}{a} - \frac{1}{ax} \right) +$$

$$\frac{3a_1}{3ax(a+bx^3)}$$

↓ 1082

$$\left(\frac{4 \left(\frac{b \left(\frac{3 \int \frac{1}{\left(1 - 2 \frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2} dx \left(1 - 2 \frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}} - \frac{1}{2} \int \frac{\sqrt[3]{a} - 2 \sqrt[3]{bx}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx} + a^{2/3}} dx - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3 \sqrt[3]{ab^{2/3}}} \right)}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{1}{ax} \right)}{a} \right) + \frac{3a}{3ax(a+bx^3)}$$

↓ 217

$$\left(\frac{4 \left(\frac{b \left(\frac{\sqrt{3} \arctan \left(\frac{1 - 2 \frac{\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}} \right)}{\sqrt[3]{b}} - \frac{1}{2} \int \frac{\sqrt[3]{a} - 2 \sqrt[3]{bx}}{b^{2/3} x^2 - \sqrt[3]{a} \sqrt[3]{bx} + a^{2/3}} dx - \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3 \sqrt[3]{ab^{2/3}}} \right)}{3 \sqrt[3]{a} \sqrt[3]{b}} - \frac{1}{ax} \right)}{a} \right) + \frac{1}{3ax(a+bx^3)}$$

↓ 1103

$$\frac{4 \left(\frac{b \left(\frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt{3}} \right)}{3\sqrt[3]{a}\sqrt[3]{b}} - \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3\sqrt[3]{a}b^{2/3}} \right)}{a} - \frac{1}{ax} \right)}{3a} + \frac{1}{3ax(a + bx^3)}$$

input `Int[1/(x^2*(a + b*x^3)^2),x]`

output `1/(3*a*x*(a + b*x^3)) + (4*(-(1/(a*x)) - (b*(-1/3*Log[a^(1/3) + b^(1/3)*x]/(a^(1/3)*b^(2/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3]))/b^(1/3)) + Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3)))/(3*a^(1/3)*b^(1/3))))/a)/(3*a)`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

rule 819 $\text{Int}[(c_ \cdot x)^{m_} \cdot (a_ + (b_ \cdot x)^n)^{p_}, x_Symbol] \rightarrow \text{Simp}[(-c \cdot x)^{m+1} \cdot (a + b \cdot x^n)^{p+1} / (a \cdot c \cdot n \cdot (p+1)), x] + \text{Simp}[(m + n \cdot (p+1) + 1) / (a \cdot n \cdot (p+1)) \text{Int}[(c \cdot x)^m \cdot (a + b \cdot x^n)^{p+1}, x], x] /;$ FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p, x]

rule 821 $\text{Int}[x / (a_ + (b_ \cdot x)^3), x_Symbol] \rightarrow \text{Simp}[-(3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3])^{(-1)} \text{Int}[1 / (\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x), x], x] + \text{Simp}[1 / (3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3]) \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x) / (\text{Rt}[a, 3]^2 - \text{Rt}[a, 3] \cdot \text{Rt}[b, 3] \cdot x + \text{Rt}[b, 3]^2 \cdot x^2), x], x] /;$ FreeQ[{a, b}, x]

rule 847 $\text{Int}[(c_ \cdot x)^{m_} \cdot (a_ + (b_ \cdot x)^n)^{p_}, x_Symbol] \rightarrow \text{Simp}[(c \cdot x)^{m+1} \cdot (a + b \cdot x^n)^{p+1} / (a \cdot c \cdot (m+1)), x] - \text{Simp}[b \cdot (m + n \cdot (p+1) + 1) / (a \cdot c^n \cdot (m+1)) \text{Int}[(c \cdot x)^{m+n} \cdot (a + b \cdot x^n)^p, x], x] /;$ FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]

rule 1082 $\text{Int}[(a_ + (b_ \cdot x) + (c_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$ RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4 \cdot a \cdot c]) /;

 FreeQ[{a, b, c}, x]

rule 1103 $\text{Int}[(d_ + (e_ \cdot x)) / (a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]] / b), x] /;$ FreeQ[{a, b, c, d, e}, x] && EqQ[2 \cdot c \cdot d - b \cdot e, 0]

rule 1142 $\text{Int}[(d_ + (e_ \cdot x)) / (a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e) / (2 \cdot c) \text{Int}[1 / (a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e / (2 \cdot c) \text{Int}[(b + 2 \cdot c \cdot x) / (a + b \cdot x + c \cdot x^2), x], x] /;$ FreeQ[{a, b, c, d, e}, x]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.55

method	result	size
risch	$\frac{-\frac{4bx^3}{3a^2} - \frac{1}{a}}{x(bx^3+a)} + \frac{4 \left(\sum_{-R=\text{RootOf}(a^7-Z^3-b)} -R \ln((-4-R^3 a^7+3b)x-a^5-R^2) \right)}{9}$	73
default	$b \frac{\frac{x^2}{3bx^3+3a} - \frac{4 \ln\left(\left(\frac{b}{a}\right)^{\frac{1}{3}}+x\right)}{9b\left(\frac{b}{a}\right)^{\frac{1}{3}}} + \frac{2 \ln\left(\left(\frac{b}{a}\right)^{\frac{2}{3}}-\left(\frac{b}{a}\right)^{\frac{1}{3}}x+x^2\right)}{9b\left(\frac{b}{a}\right)^{\frac{1}{3}}} + \frac{4\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{-2x}{\left(\frac{b}{a}\right)^{\frac{1}{3}}}-1\right)}{\frac{b}{a}}\right)}{9b\left(\frac{b}{a}\right)^{\frac{1}{3}}}}{a^2} - \frac{1}{a^2x}$	120

```
input int(1/x^2/(b*x^3+a)^2,x,method=_RETURNVERBOSE)
```

```
output (-4/3*b/a^2*x^3-1/a)/x/(b*x^3+a)+4/9*sum(_R*ln((-4*_R^3*a^7+3*b)*x-a^5*_R^2),_R=RootOf(_Z^3*a^7-b))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.11

$$\int \frac{1}{x^2 (a + bx^3)^2} dx = \frac{12bx^3 + 4\sqrt{3}(bx^4 + ax)\left(\frac{b}{a}\right)^{\frac{1}{3}} \arctan\left(\frac{2}{3}\sqrt{3}x\left(\frac{b}{a}\right)^{\frac{1}{3}} - \frac{1}{3}\sqrt{3}\right) + 2(bx^4 + ax)\left(\frac{b}{a}\right)^{\frac{1}{3}} \log\left(bx^2 - ax\left(\frac{b}{a}\right)^{\frac{2}{3}} + a\right)}{9(a^2bx^4 + a^3x)}$$

```
input integrate(1/x^2/(b*x^3+a)^2,x, algorithm="fricas")
```

output

```
-1/9*(12*b*x^3 + 4*sqrt(3)*(b*x^4 + a*x)*(b/a)^(1/3)*arctan(2/3*sqrt(3)*x*(b/a)^(1/3) - 1/3*sqrt(3)) + 2*(b*x^4 + a*x)*(b/a)^(1/3)*log(b*x^2 - a*x*(b/a)^(2/3) + a*(b/a)^(1/3)) - 4*(b*x^4 + a*x)*(b/a)^(1/3)*log(b*x + a*(b/a)^(2/3)) + 9*a)/(a^2*b*x^4 + a^3*x)
```

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.42

$$\int \frac{1}{x^2(a+bx^3)^2} dx = \frac{-3a-4bx^3}{3a^3x+3a^2bx^4} + \text{RootSum}\left(729t^3a^7-64b, \left(t \mapsto t \log\left(\frac{81t^2a^5}{16b}+x\right)\right)\right)$$

input

```
integrate(1/x**2/(b*x**3+a)**2,x)
```

output

```
(-3*a - 4*b*x**3)/(3*a**3*x + 3*a**2*b*x**4) + RootSum(729*_t**3*a**7 - 64*b, Lambda(_t, _t*log(81*_t**2*a**5/(16*b) + x)))
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.95

$$\int \frac{1}{x^2(a+bx^3)^2} dx = -\frac{4bx^3+3a}{3(a^2bx^4+a^3x)} - \frac{4\sqrt{3}\arctan\left(\frac{\sqrt{3}\left(2x-\left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9a^2\left(\frac{a}{b}\right)^{\frac{1}{3}}} - \frac{2\log\left(x^2-x\left(\frac{a}{b}\right)^{\frac{1}{3}}+\left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{9a^2\left(\frac{a}{b}\right)^{\frac{1}{3}}} + \frac{4\log\left(x+\left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{9a^2\left(\frac{a}{b}\right)^{\frac{1}{3}}}$$

input

```
integrate(1/x^2/(b*x^3+a)^2,x, algorithm="maxima")
```

output

$$-1/3*(4*b*x^3 + 3*a)/(a^2*b*x^4 + a^3*x) - 4/9*\sqrt{3}*\arctan(1/3*\sqrt{3}*(2*x - (a/b)^{1/3})/(a/b)^{1/3})/(a^2*(a/b)^{1/3}) - 2/9*\log(x^2 - x*(a/b)^{1/3} + (a/b)^{2/3})/(a^2*(a/b)^{1/3}) + 4/9*\log(x + (a/b)^{1/3})/(a^2*(a/b)^{1/3})$$
Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.05

$$\int \frac{1}{x^2(a+bx^3)^2} dx = \frac{4b\left(-\frac{a}{b}\right)^{\frac{2}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{9a^3} + \frac{4\sqrt{3}(-ab^2)^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9a^3b} - \frac{4bx^3 + 3a}{3(bx^4 + ax)a^2} - \frac{2(-ab^2)^{\frac{2}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{9a^3b}$$

input

```
integrate(1/x^2/(b*x^3+a)^2,x, algorithm="giac")
```

output

$$4/9*b*(-a/b)^{2/3}*\log(\text{abs}(x - (-a/b)^{1/3}))/a^3 + 4/9*\sqrt{3}*(-a*b^2)^{2/3}*\arctan(1/3*\sqrt{3}*(2*x + (-a/b)^{1/3})/(-a/b)^{1/3})/(a^3*b) - 1/3*(4*b*x^3 + 3*a)/((b*x^4 + a*x)*a^2) - 2/9*(-a*b^2)^{2/3}*\log(x^2 + x*(-a/b)^{1/3} + (-a/b)^{2/3})/(a^3*b)$$
Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 120, normalized size of antiderivative = 0.91

$$\int \frac{1}{x^2(a+bx^3)^2} dx = \frac{4b^{1/3} \ln(b^{1/3}x + a^{1/3})}{9a^{7/3}} - \frac{\frac{1}{a} + \frac{4bx^3}{3a^2}}{bx^4 + ax} - \frac{4b^{1/3} \ln(4b^{1/3}x - 2a^{1/3} + \sqrt{3}a^{1/3}2i) \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{9a^{7/3}} + \frac{b^{1/3} \ln(4b^{1/3}x - 2a^{1/3} - \sqrt{3}a^{1/3}2i) \left(-\frac{2}{9} + \frac{\sqrt{3}2i}{9}\right)}{a^{7/3}}$$

input `int(1/(x^2*(a + b*x^3)^2),x)`

output
$$\frac{(4*b^{1/3}*\log(b^{1/3}*x + a^{1/3}))/((9*a^{7/3})) - (1/a + (4*b*x^3)/(3*a^2))/((a*x + b*x^4)) - (4*b^{1/3}*\log(3^{1/2}*a^{1/3}*2i + 4*b^{1/3}*x - 2*a^{1/3}))*((3^{1/2}*1i)/2 + 1/2))/((9*a^{7/3})) + (b^{1/3}*\log(4*b^{1/3}*x - 3^{1/2}*a^{1/3}*2i - 2*a^{1/3}))*((3^{1/2}*2i)/9 - 2/9))/a^{7/3}}$$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.39

$$\int \frac{1}{x^2 (a + bx^3)^2} dx$$

$$= \frac{4\sqrt{3} \operatorname{atan}\left(\frac{a^{1/3} - 2b^{1/3}x}{a^{1/3}\sqrt{3}}\right) abx + 4\sqrt{3} \operatorname{atan}\left(\frac{a^{1/3} - 2b^{1/3}x}{a^{1/3}\sqrt{3}}\right) b^2 x^4 - 9b^{2/3} a^{4/3} - 12b^{5/3} a^{1/3} x^3 - 2 \log\left(a^{2/3} - b^{1/3} a^{1/3} x + b^{2/3} x^2\right)}{9b^{2/3} a^{7/3} x (bx^3 + a)}$$

input `int(1/x^2/(b*x^3+a)^2,x)`

output
$$(4*\sqrt{3}*\operatorname{atan}((a^{1/3} - 2*b^{1/3}*x)/(a^{1/3}*\sqrt{3}))*a*b*x + 4*\sqrt{3}*\operatorname{atan}((a^{1/3} - 2*b^{1/3}*x)/(a^{1/3}*\sqrt{3}))*b**2*x**4 - 9*b^{2/3}*a^{4/3} - 12*b^{5/3}*a^{1/3}*x**3 - 2*\log(a^{2/3} - b^{1/3}*a^{1/3}*x + b^{2/3}*x**2))*a*b*x - 2*\log(a^{2/3} - b^{1/3}*a^{1/3}*x + b^{2/3}*x**2))*b**2*x**4 + 4*\log(a^{1/3} + b^{1/3}*x)*a*b*x + 4*\log(a^{1/3} + b^{1/3}*x))*b**2*x**4)/(9*b^{2/3}*a^{7/3}*x*(a + b*x**3))$$

3.76 $\int \frac{1}{x^3(a+bx^3)^2} dx$

Optimal result	505
Mathematica [A] (verified)	506
Rubi [A] (verified)	506
Maple [C] (verified)	512
Fricas [A] (verification not implemented)	513
Sympy [A] (verification not implemented)	513
Maxima [A] (verification not implemented)	514
Giac [A] (verification not implemented)	514
Mupad [B] (verification not implemented)	515
Reduce [B] (verification not implemented)	516

Optimal result

Integrand size = 13, antiderivative size = 126

$$\int \frac{1}{x^3(a+bx^3)^2} dx = \frac{-\frac{1}{2ax^2} - \frac{5bx}{6a^2}}{a+bx^3} - \frac{5\sqrt[3]{\frac{a}{b}}b \left(\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{2\sqrt[3]{\frac{a}{b}}-x} \right) + \frac{1}{2} \log \left(\frac{\left(\sqrt[3]{\frac{a}{b}}+x \right)^2}{\left(\frac{a}{b} \right)^{2/3} - \sqrt[3]{\frac{a}{b}}x+x^2} \right) \right)}{9a^3}$$

output

```
(-1/2/a/x^2-5/6*b*x/a^2)/(b*x^3+a)-5/9*(a/b)^(1/3)*b*(3^(1/2)*arctan(3^(1/2)*x/(2*(a/b)^(1/3)-x))+1/2*ln(((a/b)^(1/3)+x)^2/((a/b)^(2/3)-(a/b)^(1/3)*x+x^2)))/a^3
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.02

$$\int \frac{1}{x^3 (a + bx^3)^2} dx$$

$$= \frac{-\frac{9a^{2/3}}{x^2} - \frac{6a^{2/3}bx}{a+bx^3} + 10\sqrt{3}b^{2/3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{b}x}{\sqrt[3]{a}}}{\sqrt[3]{a}}\right) - 10b^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right) + 5b^{2/3} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{bx} + \sqrt[3]{bx^2}\right)}{18a^{8/3}}$$

input `Integrate[1/(x^3*(a + b*x^3)^2),x]`

output `((-9*a^(2/3))/x^2 - (6*a^(2/3)*b*x)/(a + b*x^3) + 10*Sqrt[3]*b^(2/3)*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3))/Sqrt[3]] - 10*b^(2/3)*Log[a^(1/3) + b^(1/3)*x] + 5*b^(2/3)*Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2])/(18*a^(8/3))`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.23, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.769$, Rules used = {819, 847, 750, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 (a + bx^3)^2} dx$$

$$\downarrow 819$$

$$\frac{5 \int \frac{1}{x^3 (bx^3 + a)} dx}{3a} + \frac{1}{3ax^2 (a + bx^3)}$$

$$\downarrow 847$$

$$\frac{5 \left(-\frac{b \int \frac{1}{bx^3+a} dx}{a} - \frac{1}{2ax^2} \right)}{3a} + \frac{1}{3ax^2(a+bx^3)}$$

750

$$\frac{5 \left(\frac{b \left(\frac{\int \frac{2\sqrt[3]{a}-\sqrt[3]{b}x}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{b}x+\sqrt[3]{a}} dx}{3a^{2/3}} \right)}{a} - \frac{1}{2ax^2} \right)}{3a} + \frac{1}{3ax^2(a+bx^3)}$$

16

$$\frac{5 \left(\frac{b \left(\frac{\int \frac{2\sqrt[3]{a}-\sqrt[3]{b}x}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{3a^{2/3}} + \frac{\log(\sqrt[3]{a}+\sqrt[3]{b}x)}{3a^{2/3}\sqrt[3]{b}} \right)}{a} - \frac{1}{2ax^2} \right)}{3a} + \frac{1}{3ax^2(a+bx^3)}$$

1142

$$\frac{5 \left(\frac{b \left(\frac{\frac{3}{2}\sqrt[3]{a} \int \frac{1}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx - \frac{\int \frac{\sqrt[3]{b}(\sqrt[3]{a}-2\sqrt[3]{b}x)}{b^{2/3}x^2-\sqrt[3]{a}\sqrt[3]{b}x+a^{2/3}} dx}{2\sqrt[3]{b}} + \frac{\log(\sqrt[3]{a}+\sqrt[3]{b}x)}{3a^{2/3}\sqrt[3]{b}} \right)}{a} - \frac{1}{2ax^2} \right)}{3a} + \frac{1}{3ax^2(a+bx^3)}$$

$$\begin{aligned}
 & \downarrow 25 \\
 & \left(\left(\frac{b \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx + \frac{\int \frac{\sqrt[3]{b} (\sqrt[3]{a-2} \sqrt[3]{bx})}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx}{2 \sqrt[3]{b}} + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3} \sqrt[3]{b}} \right)}{3a^{2/3}} \right)}{a} - \frac{1}{2ax^2} \right) + \\
 & \frac{3a}{3ax^2(a+bx^3)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 27 \\
 & \left(\left(\frac{b \left(\frac{\frac{3}{2} \sqrt[3]{a} \int \frac{1}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx + \frac{1}{2} \int \frac{\sqrt[3]{a-2} \sqrt[3]{bx}}{b^{2/3}x^2 - \sqrt[3]{a} \sqrt[3]{bx+a^{2/3}}} dx + \frac{\log(\sqrt[3]{a} + \sqrt[3]{bx})}{3a^{2/3} \sqrt[3]{b}} \right)}{3a^{2/3}} \right)}{a} - \frac{1}{2ax^2} \right) + \\
 & \frac{3a}{3ax^2(a+bx^3)}
 \end{aligned}$$

1082

$$\left(\frac{5}{b} \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{bx}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx + \frac{3 \int \frac{1}{\left(1 - 2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^2 - d} \left(1 - 2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right) - \left(1 - 2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}\right)^{-3}}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right)}{3a^{2/3}\sqrt[3]{b}} \right) - \frac{1}{2ax^2} \right) + \frac{3a}{3ax^2(a+bx^3)}$$

↓ 217

$$\left(\frac{5}{b} \left(\frac{\int \frac{\sqrt[3]{a-2\sqrt[3]{bx}}}{b^{2/3}x^2 - \sqrt[3]{a}\sqrt[3]{bx+a^{2/3}}} dx - \frac{\sqrt{3} \arctan\left(\frac{1 - 2\frac{\sqrt[3]{bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt[3]{b}}}{3a^{2/3}} + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{bx}\right)}{3a^{2/3}\sqrt[3]{b}} \right) - \frac{1}{2ax^2} \right) + \frac{1}{3ax^2(a+bx^3)}$$

↓ 1103

$$\frac{b \left(\frac{\log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2\right)}{2\sqrt[3]{b}} - \frac{\sqrt{3} \arctan\left(\frac{1 - 2\sqrt[3]{b}x}{\sqrt[3]{a}}\right)}{\sqrt[3]{b}} \right) + \frac{\log\left(\sqrt[3]{a} + \sqrt[3]{b}x\right)}{3a^{2/3}\sqrt[3]{b}}}{a} - \frac{1}{2ax^2} + \frac{3a}{3ax^2(a + bx^3)}$$

input `Int[1/(x^3*(a + b*x^3)^2),x]`

output `1/(3*a*x^2*(a + b*x^3)) + (5*(-1/2*1/(a*x^2) - (b*(Log[a^(1/3) + b^(1/3)*x]/(3*a^(2/3)*b^(1/3)) + (-((Sqrt[3]*ArcTan[(1 - (2*b^(1/3)*x)/a^(1/3)]/Sqrt[3]))/b^(1/3) - Log[a^(2/3) - a^(1/3)*b^(1/3)*x + b^(2/3)*x^2]/(2*b^(1/3))))/(3*a^(2/3))))/a)/(3*a)`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 750 $\text{Int}[(a_*) + (b_*)(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 819 $\text{Int}[(c_*)(x_)^{m_}) * ((a_*) + (b_*)(x_)^{n_})^{p_}, x_Symbol] \rightarrow \text{Simp}[(-(c*x)^{m+1}) * ((a + b*x^n)^{p+1} / (a*c*n*(p+1))), x] + \text{Simp}[(m + n*(p+1) + 1) / (a*n*(p+1)) \text{ Int}[(c*x)^m * (a + b*x^n)^{p+1}, x], x] /; \text{FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$
- rule 847 $\text{Int}[(c_*)(x_)^{m_}) * ((a_*) + (b_*)(x_)^{n_})^{p_}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{m+1} * ((a + b*x^n)^{p+1} / (a*c*(m+1))), x] - \text{Simp}[b * ((m + n*(p+1) + 1) / (a*c^n*(m+1))) \text{ Int}[(c*x)^{m+n} * (a + b*x^n)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$
- rule 1082 $\text{Int}[(a_*) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_*) + (e_*)(x_)/((a_*) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1142

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.05 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.59

method	result	size
risch	$\frac{-\frac{5bx^3}{6a^2} - \frac{1}{2a}}{x^2(bx^3+a)} + \frac{5 \left(\sum_{-R=\text{RootOf}(a^8-Z^3+b^2)} -R \ln((-4-R^3a^8-3b^2)x-a^3b-R) \right)}{9}$	74
default	$b \left(\frac{x}{3bx^3+3a} + \frac{5 \ln\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}+x\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{5 \ln\left(\left(\frac{a}{b}\right)^{\frac{2}{3}}-\left(\frac{a}{b}\right)^{\frac{1}{3}}x+x^2\right)}{18b\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{5\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\left(\frac{a}{b}\right)^{\frac{1}{3}}-1\right)}{\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9b\left(\frac{a}{b}\right)^{\frac{2}{3}}} \right) - \frac{1}{2a^2x^2}$	118

input

```
int(1/x^3/(b*x^3+a)^2,x,method=_RETURNVERBOSE)
```

output

```
(-5/6*b/a^2*x^3-1/2/a)/x^2/(b*x^3+a)+5/9*sum(_R*ln((-4*_R^3*a^8-3*b^2)*x-a
^3*b*_R),_R=RootOf(_Z^3*a^8+b^2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.43

$$\int \frac{1}{x^3 (a + bx^3)^2} dx = \frac{15bx^3 + 10\sqrt{3}(bx^5 + ax^2)\left(\frac{b^2}{a^2}\right)^{\frac{1}{3}} \arctan\left(\frac{2\sqrt{3}ax\left(\frac{b^2}{a^2}\right)^{\frac{2}{3}} - \sqrt{3}b}{3b}\right) - 5(bx^5 + ax^2)\left(\frac{b^2}{a^2}\right)^{\frac{1}{3}} \log\left(b^2x^2 - abx\left(\frac{b^2}{a^2}\right)^{\frac{1}{3}} + a^2\left(\frac{b^2}{a^2}\right)^{\frac{2}{3}}\right) + 10(bx^5 + ax^2)\left(\frac{b^2}{a^2}\right)^{\frac{1}{3}} \log(bx + a\left(\frac{b^2}{a^2}\right)^{\frac{1}{3}}) + 9a}{18(a^2bx^5 + a^3x^2)}$$

input `integrate(1/x^3/(b*x^3+a)^2,x, algorithm="fricas")`output `-1/18*(15*b*x^3 + 10*sqrt(3)*(b*x^5 + a*x^2)*(b^2/a^2)^(1/3)*arctan(1/3*(2*sqrt(3)*a*x*(b^2/a^2)^(2/3) - sqrt(3)*b)/b) - 5*(b*x^5 + a*x^2)*(b^2/a^2)^(1/3)*log(b^2*x^2 - a*b*x*(b^2/a^2)^(1/3) + a^2*(b^2/a^2)^(2/3)) + 10*(b*x^5 + a*x^2)*(b^2/a^2)^(1/3)*log(b*x + a*(b^2/a^2)^(1/3)) + 9*a)/(a^2*b*x^5 + a^3*x^2)`**Sympy [A] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.46

$$\int \frac{1}{x^3 (a + bx^3)^2} dx = \frac{-3a - 5bx^3}{6a^3x^2 + 6a^2bx^5} + \text{RootSum}\left(729t^3a^8 + 125b^2, \left(t \mapsto t \log\left(-\frac{9ta^3}{5b} + x\right)\right)\right)$$

input `integrate(1/x**3/(b*x**3+a)**2,x)`output `(-3*a - 5*b*x**3)/(6*a**3*x**2 + 6*a**2*b*x**5) + RootSum(729*_t**3*a**8 + 125*b**2, Lambda(_t, _t*log(-9*_t*a**3/(5*b) + x))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.02

$$\int \frac{1}{x^3 (a + bx^3)^2} dx = -\frac{5bx^3 + 3a}{6(a^2bx^5 + a^3x^2)} - \frac{5\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2x - \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9a^2\left(\frac{a}{b}\right)^{\frac{2}{3}}} + \frac{5 \log\left(x^2 - x\left(\frac{a}{b}\right)^{\frac{1}{3}} + \left(\frac{a}{b}\right)^{\frac{2}{3}}\right)}{18a^2\left(\frac{a}{b}\right)^{\frac{2}{3}}} - \frac{5 \log\left(x + \left(\frac{a}{b}\right)^{\frac{1}{3}}\right)}{9a^2\left(\frac{a}{b}\right)^{\frac{2}{3}}}$$

input `integrate(1/x^3/(b*x^3+a)^2,x, algorithm="maxima")`output `-1/6*(5*b*x^3 + 3*a)/(a^2*b*x^5 + a^3*x^2) - 5/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - (a/b)^(1/3))/(a/b)^(1/3))/(a^2*(a/b)^(2/3)) + 5/18*log(x^2 - x*(a/b)^(1/3) + (a/b)^(2/3))/(a^2*(a/b)^(2/3)) - 5/9*log(x + (a/b)^(1/3))/(a^2*(a/b)^(2/3))`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.04

$$\int \frac{1}{x^3 (a + bx^3)^2} dx = \frac{5b\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|x - \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right|\right)}{9a^3} - \frac{bx}{3(bx^3 + a)a^2} - \frac{5\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2x + \left(-\frac{a}{b}\right)^{\frac{1}{3}}\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{9a^3} - \frac{5(-ab^2)^{\frac{1}{3}} \log\left(x^2 + x\left(-\frac{a}{b}\right)^{\frac{1}{3}} + \left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{18a^3} - \frac{1}{2a^2x^2}$$

input `integrate(1/x^3/(b*x^3+a)^2,x, algorithm="giac")`

output

```
5/9*b*(-a/b)^(1/3)*log(abs(x - (-a/b)^(1/3)))/a^3 - 1/3*b*x/((b*x^3 + a)*a^2) - 5/9*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*(2*x + (-a/b)^(1/3))/(-a/b)^(1/3))/a^3 - 5/18*(-a*b^2)^(1/3)*log(x^2 + x*(-a/b)^(1/3) + (-a/b)^(2/3))/a^3 - 1/2/(a^2*x^2)
```

Mupad [B] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.16

$$\int \frac{1}{x^3 (a + bx^3)^2} dx$$

$$= \frac{5(-1)^{1/3} b^{2/3} \ln\left((-1)^{1/3} a^{1/3} - b^{1/3} x\right)}{9 a^{8/3}} - \frac{\frac{1}{2a} + \frac{5bx^3}{6a^2}}{bx^5 + ax^2}$$

$$- \frac{5(-1)^{1/3} b^{2/3} \ln\left((-1)^{1/3} a^{1/3} + 2b^{1/3} x + (-1)^{5/6} \sqrt{3} a^{1/3}\right) \left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{9 a^{8/3}}$$

$$+ \frac{5(-1)^{1/3} b^{2/3} \ln\left((-1)^{1/3} a^{1/3} + 2b^{1/3} x - (-1)^{5/6} \sqrt{3} a^{1/3}\right) \left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{9 a^{8/3}}$$

input

```
int(1/(x^3*(a + b*x^3)^2),x)
```

output

```
(5*(-1)^(1/3)*b^(2/3)*log((-1)^(1/3)*a^(1/3) - b^(1/3)*x)/(9*a^(8/3)) - (1/(2*a) + (5*b*x^3)/(6*a^2))/(a*x^2 + b*x^5) - (5*(-1)^(1/3)*b^(2/3)*log((-1)^(1/3)*a^(1/3) + 2*b^(1/3)*x + (-1)^(5/6)*3^(1/2)*a^(1/3))*((3^(1/2)*1i)/2 + 1/2)/(9*a^(8/3)) + (5*(-1)^(1/3)*b^(2/3)*log((-1)^(1/3)*a^(1/3) + 2*b^(1/3)*x - (-1)^(5/6)*3^(1/2)*a^(1/3))*((3^(1/2)*1i)/2 - 1/2)/(9*a^(8/3))
```

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 202, normalized size of antiderivative = 1.60

$$\int \frac{1}{x^3 (a + bx^3)^2} dx$$

$$= \frac{10a^{\frac{4}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) bx^2 + 10a^{\frac{1}{3}}\sqrt{3} \operatorname{atan}\left(\frac{a^{\frac{1}{3}} - 2b^{\frac{1}{3}}x}{a^{\frac{1}{3}}\sqrt{3}}\right) b^2x^5 + 5a^{\frac{4}{3}}\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) bx^2 + 5a^{\frac{1}{3}}\log\left(a^{\frac{2}{3}} - b^{\frac{1}{3}}a^{\frac{1}{3}}x + b^{\frac{2}{3}}x^2\right) b^2x^5}{18b^{\frac{1}{3}}a^3x^2}$$

input `int(1/x^3/(b*x^3+a)^2,x)`output `(10*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*a*b*x**2 + 10*a**(1/3)*sqrt(3)*atan((a**(1/3) - 2*b**(1/3)*x)/(a**(1/3)*sqrt(3)))*b**2*x**5 + 5*a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*a*b*x**2 + 5*a**(1/3)*log(a**(2/3) - b**(1/3)*a**(1/3)*x + b**(2/3)*x**2)*b**2*x**5 - 10*a**(1/3)*log(a**(1/3) + b**(1/3)*x)*a*b*x**2 - 10*a**(1/3)*log(a**(1/3) + b**(1/3)*x)*b**2*x**5 - 9*b**(1/3)*a**2 - 15*b**(1/3)*a*b*x**3)/(18*b**(1/3)*a**3*x**2*(a + b*x**3))`

3.77 $\int \frac{1}{a+bx^4} dx$

Optimal result	517
Mathematica [B] (verified)	517
Rubi [B] (verified)	518
Maple [C] (verified)	521
Fricas [C] (verification not implemented)	522
Sympy [A] (verification not implemented)	522
Maxima [B] (verification not implemented)	523
Giac [B] (verification not implemented)	523
Mupad [B] (verification not implemented)	524
Reduce [B] (verification not implemented)	524

Optimal result

Integrand size = 9, antiderivative size = 63

$$\int \frac{1}{a + bx^4} dx = \frac{\sqrt[4]{-\frac{a}{b}} \left(2 \arctan \left(\frac{x}{\sqrt[4]{-\frac{a}{b}}} \right) + \log \left(\frac{\sqrt[4]{-\frac{a}{b}} + x}{-\sqrt[4]{-\frac{a}{b}} + x} \right) \right)}{4a}$$

output

```
1/4*(-a/b)^(1/4)*(2*arctan(x/(-a/b)^(1/4))+ln(((a/b)^(1/4)+x)/((-a/b)^(1/4)+x)))/a
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 134 vs. 2(63) = 126.

Time = 0.03 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.13

$$\int \frac{1}{a + bx^4} dx = \frac{-2 \arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a}} \right) + 2 \arctan \left(1 + \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a}} \right) - \log \left(\sqrt{a} - \sqrt{2} \sqrt[4]{a} \sqrt[4]{bx} + \sqrt{bx^2} \right) + \log \left(\sqrt{a} + \sqrt{2} \sqrt[4]{a} \sqrt[4]{bx} + \sqrt{bx^2} \right)}{4\sqrt{2}a^{3/4}\sqrt[4]{b}}$$

input `Integrate[(a + b*x^4)^(-1),x]`

output $(-2*\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{(1/4)}*x)/a^{(1/4)}] + 2*\text{ArcTan}[1 + (\text{Sqrt}[2]*b^{(1/4)}*x)/a^{(1/4)}] - \text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*x + \text{Sqrt}[b]*x^2] + \text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*x + \text{Sqrt}[b]*x^2])/(4*\text{Sqrt}[2]*a^{(3/4)}*b^{(1/4)})$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 200 vs. $2(63) = 126$.

Time = 0.67 (sec) , antiderivative size = 200, normalized size of antiderivative = 3.17, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.889$, Rules used = {755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{a + bx^4} dx \\
 & \quad \downarrow 755 \\
 & \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{\sqrt{bx^2}+\sqrt{a}}{bx^4+a} dx}{2\sqrt{a}} \\
 & \quad \downarrow 1476 \\
 & \frac{\int \frac{1}{x^2 - \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt[4]{b}}} dx}{2\sqrt{b}} + \frac{\int \frac{1}{x^2 + \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt[4]{b}}} dx}{2\sqrt{b}} + \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{a}} \\
 & \quad \downarrow 1082 \\
 & \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{1}{\left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)^2} d\left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{\left(\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}} + 1\right)^2} d\left(\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \\
 & \quad \downarrow 217
 \end{aligned}$$

output
$$\begin{aligned} & (-\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{(1/4)*x})/a^{(1/4)}]/(\text{Sqrt}[2]*a^{(1/4)*b^{(1/4)}}) + \text{ArcTan}[1 + (\text{Sqrt}[2]*b^{(1/4)*x})/a^{(1/4)}]/(\text{Sqrt}[2]*a^{(1/4)*b^{(1/4)}}))/(2*\text{Sqrt}[a] \\ &] + (-1/2*\text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{(1/4)*b^{(1/4)*x} + \text{Sqrt}[b]*x^2]/(\text{Sqrt}[2] \\ & *a^{(1/4)*b^{(1/4)}}) + \text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{(1/4)*b^{(1/4)*x} + \text{Sqrt}[b]*x^2] \\ & / (2*\text{Sqrt}[2]*a^{(1/4)*b^{(1/4)}}))/(2*\text{Sqrt}[a]) \end{aligned}$$

Defintions of rubi rules used

rule 25
$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27
$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[a \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ /; } \text{FreeQ}[a, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (b_)*(Gx_)] \text{ /; } \text{FreeQ}[b, \text{x}]$$

rule 217
$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \text{ :> } \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] \text{ /; } \text{FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 755
$$\text{Int}[(a_ + (b_)*(x_)^4)^{-1}, \text{x_Symbol}] \text{ :> } \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*r) \quad \text{Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}] + \text{Simp}[1/(2*r) \quad \text{Int}[(r + s*x^2)/(a + b*x^4), \text{x}], \text{x}]] \text{ /; } \text{FreeQ}[\{a, b\}, \text{x}] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 1082
$$\text{Int}[(a_ + (b_)*(x_) + (c_)*(x_)^2)^{-1}, \text{x_Symbol}] \text{ :> } \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \quad \text{Subst}[\text{Int}[1/(q - x^2), \text{x}], \text{x}, 1 + 2*c*(x/b)], \text{x}] \text{ /; } \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \text{!RationalQ}[b^2 - 4*a*c])] \text{ /; } \text{FreeQ}[\{a, b, c\}, \text{x}]$$

rule 1103
$$\text{Int}[(d_ + (e_)*(x_))/((a_ + (b_)*(x_) + (c_)*(x_)^2), \text{x_Symbol}] \text{ :> } \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, \text{x}]]/b), \text{x}] \text{ /; } \text{FreeQ}[\{a, b, c, d, e\}, \text{x}] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$$

rule 1476

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.43

method	result	size
risch	$\frac{\sum_{-R=\text{RootOf}(bZ^4+a)} \frac{\ln(x-R)}{-R^3}}{4b}$	27
default	$\frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2} \left(\ln\left(\frac{x^2+\left(\frac{a}{b}\right)^{\frac{1}{4}}x\sqrt{2}+\sqrt{\frac{a}{b}}}{x^2-\left(\frac{a}{b}\right)^{\frac{1}{4}}x\sqrt{2}+\sqrt{\frac{a}{b}}}\right) + 2\arctan\left(\frac{\sqrt{2}x}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}+1\right) + 2\arctan\left(\frac{\sqrt{2}x}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}-1\right) \right)}{8a}$	102

input

```
int(1/(b*x^4+a),x,method=_RETURNVERBOSE)
```

output

```
1/4/b*sum(1/_R^3*ln(x-_R),_R=RootOf(_Z^4*b+a))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.08 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.78

$$\begin{aligned} \int \frac{1}{a + bx^4} dx &= \frac{1}{4} \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} \log \left(a \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} + x \right) \\ &\quad + \frac{1}{4} i \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} \log \left(i a \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} + x \right) \\ &\quad - \frac{1}{4} i \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} \log \left(-i a \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} + x \right) \\ &\quad - \frac{1}{4} \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} \log \left(-a \left(-\frac{1}{a^3b} \right)^{\frac{1}{4}} + x \right) \end{aligned}$$

input `integrate(1/(b*x^4+a),x, algorithm="fricas")`

output `1/4*(-1/(a^3*b))^(1/4)*log(a*(-1/(a^3*b))^(1/4) + x) + 1/4*I*(-1/(a^3*b))^(1/4)*log(I*a*(-1/(a^3*b))^(1/4) + x) - 1/4*I*(-1/(a^3*b))^(1/4)*log(-I*a*(-1/(a^3*b))^(1/4) + x) - 1/4*(-1/(a^3*b))^(1/4)*log(-a*(-1/(a^3*b))^(1/4) + x)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.32

$$\int \frac{1}{a + bx^4} dx = \text{RootSum} (256t^4 a^3 b + 1, (t \mapsto t \log(4ta + x)))$$

input `integrate(1/(b*x**4+a),x)`

output `RootSum(256*_t**4*a**3*b + 1, Lambda(_t, _t*log(4*_t*a + x)))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. $2(53) = 106$.

Time = 0.11 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.68

$$\int \frac{1}{a + bx^4} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} + \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{4\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} - \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{4\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} \\ + \frac{\sqrt{2} \log\left(\sqrt{bx^2} + \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}x} + \sqrt{a}\right)}{8a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log\left(\sqrt{bx^2} - \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}x} + \sqrt{a}\right)}{8a^{\frac{3}{4}}b^{\frac{1}{4}}}$$

input `integrate(1/(b*x^4+a),x, algorithm="maxima")`

output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x + sqrt(2)*a^(1/4)*b^(1/4))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(a)*sqrt(sqrt(a)*sqrt(b))) + 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x - sqrt(2)*a^(1/4)*b^(1/4))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(a)*sqrt(sqrt(a)*sqrt(b))) + 1/8*sqrt(2)*log(sqrt(b)*x^2 + sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(3/4)*b^(1/4)) - 1/8*sqrt(2)*log(sqrt(b)*x^2 - sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(3/4)*b^(1/4))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. $2(53) = 106$.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 2.84

$$\int \frac{1}{a + bx^4} dx = \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x + \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{4ab} \\ + \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x - \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{4ab} \\ + \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{8ab} \\ - \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{8ab}$$

input `integrate(1/(b*x^4+a),x, algorithm="giac")`

output `1/4*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x + sqrt(2)*(a/b)^(1/4))/(a/b)^(1/4))/(a*b) + 1/4*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x - sqrt(2)*(a/b)^(1/4))/(a/b)^(1/4))/(a*b) + 1/8*sqrt(2)*(a*b^3)^(1/4)*log(x^2 + sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a*b) - 1/8*sqrt(2)*(a*b^3)^(1/4)*log(x^2 - sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a*b)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.52

$$\int \frac{1}{a + bx^4} dx = -\frac{\operatorname{atan}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right) + \operatorname{atanh}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right)}{2(-a)^{3/4}b^{1/4}}$$

input `int(1/(a + b*x^4),x)`

output `-(atan((b^(1/4)*x)/(-a)^(1/4)) + atanh((b^(1/4)*x)/(-a)^(1/4)))/(2*(-a)^(3/4)*b^(1/4))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.78

$$\int \frac{1}{a + bx^4} dx = \frac{\sqrt{2} \left(-2 \operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{b}x}{b^{1/4}a^{1/4}\sqrt{2}}\right) + 2 \operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}+2\sqrt{b}x}{b^{1/4}a^{1/4}\sqrt{2}}\right) - \log\left(-b^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{b}x^2\right) + \log\left(b^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{b}x^2\right) \right)}{8b^{1/4}a^{3/4}}$$

input `int(1/(b*x^4+a),x)`

output

```
(b**(3/4)*a**(1/4)*sqrt(2)*( - 2*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2))) + 2*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2))) - log( - b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2) + log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)))/(8*a*b)
```

3.78 $\int \frac{x}{a+bx^4} dx$

Optimal result	526
Mathematica [A] (verified)	526
Rubi [A] (verified)	527
Maple [A] (verified)	528
Fricas [A] (verification not implemented)	528
Sympy [B] (verification not implemented)	528
Maxima [A] (verification not implemented)	529
Giac [A] (verification not implemented)	529
Mupad [B] (verification not implemented)	530
Reduce [B] (verification not implemented)	530

Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \frac{x}{a+bx^4} dx = \frac{\arctan\left(\sqrt{\frac{b}{a}}x^2\right)}{2\sqrt{ab}}$$

output `1/2*arctan((b/a)^(1/2)*x^2)/(a*b)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16

$$\int \frac{x}{a+bx^4} dx = \frac{\arctan\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right)}{2\sqrt{a}\sqrt{b}}$$

input `Integrate[x/(a + b*x^4),x]`

output `ArcTan[(Sqrt[b]*x^2)/Sqrt[a]]/(2*Sqrt[a]*Sqrt[b])`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.16, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {807, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{a + bx^4} dx$$

↓ 807

$$\frac{1}{2} \int \frac{1}{bx^4 + a} dx^2$$

↓ 218

$$\frac{\arctan\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right)}{2\sqrt{a}\sqrt{b}}$$

input `Int[x/(a + b*x^4),x]`

output `ArcTan[(Sqrt[b]*x^2)/Sqrt[a]]/(2*Sqrt[a]*Sqrt[b])`

Defintions of rubi rules used

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{\arctan\left(\frac{bx^2}{\sqrt{ab}}\right)}{2\sqrt{ab}}$	19
risch	$-\frac{\ln(x^2\sqrt{-ab}-a)}{4\sqrt{-ab}} + \frac{\ln(x^2\sqrt{-ab}+a)}{4\sqrt{-ab}}$	46

input `int(x/(b*x^4+a),x,method=_RETURNVERBOSE)`

output `1/2/(a*b)^(1/2)*arctan(b*x^2/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.88

$$\int \frac{x}{a + bx^4} dx = \left[-\frac{\sqrt{-ab} \log\left(\frac{bx^4 - 2\sqrt{-ab}x^2 - a}{bx^4 + a}\right)}{4ab}, -\frac{\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{bx^2}\right)}{2ab} \right]$$

input `integrate(x/(b*x^4+a),x, algorithm="fricas")`

output `[-1/4*sqrt(-a*b)*log((b*x^4 - 2*sqrt(-a*b)*x^2 - a)/(b*x^4 + a))/(a*b), -1/2*sqrt(a*b)*arctan(sqrt(a*b)/(b*x^2))/(a*b)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(19) = 38$.

Time = 0.10 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.24

$$\int \frac{x}{a + bx^4} dx = -\frac{\sqrt{-\frac{1}{ab}} \log\left(-a\sqrt{-\frac{1}{ab}} + x^2\right)}{4} + \frac{\sqrt{-\frac{1}{ab}} \log\left(a\sqrt{-\frac{1}{ab}} + x^2\right)}{4}$$

input `integrate(x/(b*x**4+a),x)`

output `-sqrt(-1/(a*b))*log(-a*sqrt(-1/(a*b)) + x**2)/4 + sqrt(-1/(a*b))*log(a*sqrt(-1/(a*b)) + x**2)/4`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{x}{a + bx^4} dx = \frac{\arctan\left(\frac{bx^2}{\sqrt{ab}}\right)}{2\sqrt{ab}}$$

input `integrate(x/(b*x^4+a),x, algorithm="maxima")`

output `1/2*arctan(b*x^2/sqrt(a*b))/sqrt(a*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \frac{x}{a + bx^4} dx = \frac{\arctan\left(\frac{bx^2}{\sqrt{ab}}\right)}{2\sqrt{ab}}$$

input `integrate(x/(b*x^4+a),x, algorithm="giac")`

output `1/2*arctan(b*x^2/sqrt(a*b))/sqrt(a*b)`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

$$\int \frac{x}{a + bx^4} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b}x^2}{\sqrt{a}}\right)}{2\sqrt{a}\sqrt{b}}$$

input `int(x/(a + b*x^4),x)`output `atan((b^(1/2)*x^2)/a^(1/2))/(2*a^(1/2)*b^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.68

$$\int \frac{x}{a + bx^4} dx = -\frac{\sqrt{b}\sqrt{a}\left(\operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{b}x}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) + \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{b}x}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right)\right)}{2ab}$$

input `int(x/(b*x^4+a),x)`output `(- sqrt(b)*sqrt(a)*(atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2))) + atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2))))/(2*a*b)`

3.79 $\int \frac{x^2}{a+bx^4} dx$

Optimal result	531
Mathematica [B] (verified)	531
Rubi [B] (verified)	532
Maple [C] (verified)	535
Fricas [C] (verification not implemented)	536
Sympy [A] (verification not implemented)	536
Maxima [B] (verification not implemented)	537
Giac [B] (verification not implemented)	537
Mupad [B] (verification not implemented)	538
Reduce [B] (verification not implemented)	538

Optimal result

Integrand size = 13, antiderivative size = 63

$$\int \frac{x^2}{a+bx^4} dx = -\frac{-2 \arctan\left(\frac{x}{\sqrt[4]{-\frac{a}{b}}}\right) + \log\left(\frac{\sqrt[4]{-\frac{a}{b}}+x}{-\sqrt[4]{-\frac{a}{b}}+x}\right)}{4\sqrt[4]{-\frac{a}{b}}b}$$

output

```
-1/4*(-2*arctan(x/(-a/b)^(1/4))+ln(((a/b)^(1/4)+x)/(-(a/b)^(1/4)+x)))/(-a/b)^(1/4)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 134 vs. 2(63) = 126.

Time = 0.01 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.13

$$\int \frac{x^2}{a+bx^4} dx = \frac{-2 \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt{a}}\right) + 2 \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt{a}}\right) + \log\left(\sqrt{a} - \sqrt{2}\sqrt[4]{a}\sqrt[4]{bx} + \sqrt{bx^2}\right) - \log\left(\sqrt{a} + \sqrt{2}\sqrt[4]{a}\sqrt[4]{bx} + \sqrt{bx^2}\right)}{4\sqrt{2}\sqrt[4]{ab^3/4}}$$

input `Integrate[x^2/(a + b*x^4),x]`

output $(-2*\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{(1/4)}*x)/a^{(1/4)}] + 2*\text{ArcTan}[1 + (\text{Sqrt}[2]*b^{(1/4)})*x)/a^{(1/4)}] + \text{Log}[\text{Sqrt}[a - \text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*x + \text{Sqrt}[b]*x^2] - \text{Log}[\text{Sqrt}[a + \text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*x + \text{Sqrt}[b]*x^2)]/(4*\text{Sqrt}[2]*a^{(1/4)}*b^{(3/4)})$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 200 vs. $2(63) = 126$.

Time = 0.62 (sec) , antiderivative size = 200, normalized size of antiderivative = 3.17, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$, Rules used = {826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{a + bx^4} dx \\
 & \quad \downarrow 826 \\
 & \frac{\int \frac{\sqrt{bx^2 + \sqrt{a}}}{bx^4 + a} dx}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a - \sqrt{bx^2}}}{bx^4 + a} dx}{2\sqrt{b}} \\
 & \quad \downarrow 1476 \\
 & \frac{\int \frac{1}{x^2 - \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} + \frac{\int \frac{1}{x^2 + \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a - \sqrt{bx^2}}}{bx^4 + a} dx}{2\sqrt{b}} \\
 & \quad \downarrow 1082 \\
 & \frac{\int \frac{1}{-\left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)^2} d\left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{-\left(\frac{\sqrt{2}\sqrt[4]{b}x + 1}{\sqrt[4]{a}}\right)^2} d\left(\frac{\sqrt{2}\sqrt[4]{b}x + 1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a - \sqrt{bx^2}}}{bx^4 + a} dx}{2\sqrt{b}} \\
 & \quad \downarrow 217
 \end{aligned}$$

$$\begin{aligned}
& \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}x+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{b}x^2}{bx^4+a} dx}{2\sqrt{b}} \\
& \quad \downarrow 1479 \\
& \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}x+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \\
& \frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}x}{\sqrt[4]{b}\left(x^2-\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}x+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x^2+\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \\
& \quad \downarrow 25 \\
& \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}x+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}x}{\sqrt[4]{b}\left(x^2-\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}x+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x^2+\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \\
& \quad \downarrow 27 \\
& \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}x+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}x}{\sqrt[4]{b}\left(x^2-\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{b}x+\sqrt[4]{a}}{\sqrt[4]{b}\left(x^2+\frac{\sqrt{2}\sqrt[4]{a}x+\sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \\
& \quad \downarrow 1103 \\
& \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}x+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \\
& \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x+\sqrt{a}+\sqrt{b}x^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x+\sqrt{a}+\sqrt{b}x^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \\
& \quad \downarrow \\
& \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x+\sqrt{a}+\sqrt{b}x^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x+\sqrt{a}+\sqrt{b}x^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}
\end{aligned}$$

input `Int[x^2/(a + b*x^4),x]`

output

$$\begin{aligned} & (-\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{(1/4)}*x)/a^{(1/4)}]/(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)})) + \text{ArcTan}[1 + (\text{Sqrt}[2]*b^{(1/4)}*x)/a^{(1/4)}]/(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}) \\ & - (-1/2*\text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*x + \text{Sqrt}[b]*x^2]/(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}) \\ & + \text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*x + \text{Sqrt}[b]*x^2]/(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)})) \\ & / (2*\text{Sqrt}[b]) \end{aligned}$$
Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), \text{x_Symbol}] \text{ :> } \text{Simp}[a \text{ Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[a, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (b_)*(Gx_)] \text{ ; FreeQ}[b, \text{x}]$$

rule 217

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \text{ :> } \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], \text{x}] \text{ ; FreeQ}[\{a, b\}, \text{x}] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 826

$$\text{Int}[(x_)^2/((a_ + (b_)*(x_)^4), \text{x_Symbol}] \text{ :> } \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \text{ Int}[(r + s*x^2)/(a + b*x^4), \text{x}], \text{x}] - \text{Simp}[1/(2*s) \text{ Int}[(r - s*x^2)/(a + b*x^4), \text{x}], \text{x}]] \text{ ; FreeQ}[\{a, b\}, \text{x}] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$$

rule 1082

$$\text{Int}[(a_ + (b_)*(x_) + (c_)*(x_)^2)^{-1}, \text{x_Symbol}] \text{ :> } \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), \text{x}], \text{x}, 1 + 2*c*(x/b)], \text{x}] \text{ ; RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \text{!RationalQ}[b^2 - 4*a*c])] \text{ ; FreeQ}[\{a, b, c\}, \text{x}]$$

rule 1103

$$\text{Int}[(d_ + (e_)*(x_))/((a_ + (b_)*(x_) + (c_)*(x_)^2), \text{x_Symbol}] \text{ :> } \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, \text{x}]]/b), \text{x}] \text{ ; FreeQ}[\{a, b, c, d, e\}, \text{x}] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$$

rule 1476

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.43

method	result	size
risch	$\frac{\sum_{-R=\text{RootOf}(bZ^4+a)} \frac{\ln(x-R)}{-R}}{4b}$	27
default	$\frac{\sqrt{2} \left(\ln \left(\frac{x^2 - \left(\frac{a}{b}\right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}}{x^2 + \left(\frac{a}{b}\right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}} \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} + 1 \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} - 1 \right) \right)}{8b \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	102

input

```
int(x^2/(b*x^4+a), x, method=_RETURNVERBOSE)
```

output

```
1/4/b*sum(1/_R*ln(x-_R), _R=RootOf(_Z^4*b+a))
```


Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.97

$$\int \frac{x^2}{a + bx^4} dx = \frac{1}{4} \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + x \right) \\ - \frac{1}{4} i \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(i ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + x \right) \\ + \frac{1}{4} i \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(-i ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + x \right) \\ - \frac{1}{4} \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(-ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + x \right)$$

input `integrate(x^2/(b*x^4+a),x, algorithm="fricas")`

output `1/4*(-1/(a*b^3))^(1/4)*log(a*b^2*(-1/(a*b^3))^(3/4) + x) - 1/4*I*(-1/(a*b^3))^(1/4)*log(I*a*b^2*(-1/(a*b^3))^(3/4) + x) + 1/4*I*(-1/(a*b^3))^(1/4)*log(-I*a*b^2*(-1/(a*b^3))^(3/4) + x) - 1/4*(-1/(a*b^3))^(1/4)*log(-a*b^2*(-1/(a*b^3))^(3/4) + x)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.41

$$\int \frac{x^2}{a + bx^4} dx = \text{RootSum} (256t^4 ab^3 + 1, (t \mapsto t \log (64t^3 ab^2 + x)))$$

input `integrate(x**2/(b*x**4+a),x)`

output `RootSum(256*_t**4*a*b**3 + 1, Lambda(_t, _t*log(64*_t**3*a*b**2 + x)))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. $2(55) = 110$.

Time = 0.11 (sec) , antiderivative size = 169, normalized size of antiderivative = 2.68

$$\int \frac{x^2}{a + bx^4} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} + \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{4\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} + \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} - \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{4\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} - \frac{\sqrt{2} \log\left(\sqrt{bx^2} + \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}x} + \sqrt{a}\right)}{8a^{\frac{1}{4}}b^{\frac{3}{4}}} + \frac{\sqrt{2} \log\left(\sqrt{bx^2} - \sqrt{2a^{\frac{1}{4}}b^{\frac{1}{4}}x} + \sqrt{a}\right)}{8a^{\frac{1}{4}}b^{\frac{3}{4}}}$$

input `integrate(x^2/(b*x^4+a),x, algorithm="maxima")`

output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x + sqrt(2)*a^(1/4)*b^(1/4))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(sqrt(a)*sqrt(b))*sqrt(b)) + 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x - sqrt(2)*a^(1/4)*b^(1/4))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(sqrt(a)*sqrt(b))*sqrt(b)) - 1/8*sqrt(2)*log(sqrt(b)*x^2 + sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(1/4)*b^(3/4)) + 1/8*sqrt(2)*log(sqrt(b)*x^2 - sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(1/4)*b^(3/4))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. $2(55) = 110$.

Time = 0.12 (sec) , antiderivative size = 179, normalized size of antiderivative = 2.84

$$\int \frac{x^2}{a + bx^4} dx = \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(2x + \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{4ab^3} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(2x - \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{4ab^3} - \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{8ab^3} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{8ab^3}$$

input `integrate(x^2/(b*x^4+a),x, algorithm="giac")`

output
$$\frac{1}{4}\sqrt{2}\frac{(ab^3)^{3/4}\arctan\left(\frac{1}{2}\sqrt{2}\frac{2x+\sqrt{2}\frac{a}{b}^{1/4}}{\frac{a}{b}^{1/4}}\right)}{(ab^3)^{3/4}} + \frac{1}{4}\sqrt{2}\frac{(ab^3)^{3/4}\arctan\left(\frac{1}{2}\sqrt{2}\frac{2x-\sqrt{2}\frac{a}{b}^{1/4}}{\frac{a}{b}^{1/4}}\right)}{(ab^3)^{3/4}} - \frac{1}{8}\sqrt{2}\frac{(ab^3)^{3/4}\log\left(x^2+\sqrt{2}x\frac{a}{b}^{1/4}+\sqrt{\frac{a}{b}}\right)}{(ab^3)^{3/4}} + \frac{1}{8}\sqrt{2}\frac{(ab^3)^{3/4}\log\left(x^2-\sqrt{2}x\frac{a}{b}^{1/4}+\sqrt{\frac{a}{b}}\right)}{(ab^3)^{3/4}}$$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.56

$$\int \frac{x^2}{a+bx^4} dx = \frac{\operatorname{atan}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right) - \operatorname{atanh}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right)}{2(-a)^{1/4}b^{3/4}}$$

input `int(x^2/(a + b*x^4),x)`

output
$$\frac{(\operatorname{atan}(b^{1/4}x/(-a)^{1/4}) - \operatorname{atanh}(b^{1/4}x/(-a)^{1/4}))}{(2*(-a)^{1/4})b^{3/4}}$$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.78

$$\int \frac{x^2}{a+bx^4} dx = \frac{\sqrt{2}\left(-2\operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{b}x}{b^{1/4}a^{1/4}\sqrt{2}}\right) + 2\operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}+2\sqrt{b}x}{b^{1/4}a^{1/4}\sqrt{2}}\right) + \log\left(-b^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} + \sqrt{b}x^2\right) - \log\left(b^{1/4}a^{1/4}\sqrt{2}x + \sqrt{a} - \sqrt{b}x^2\right)\right)}{8b^{3/4}a^{1/4}}$$

input `int(x^2/(b*x^4+a),x)`

output

```
(b**(1/4)*a**(3/4)*sqrt(2)*( - 2*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2))) + 2*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2))) + log( - b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2) - log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)))/(8*a*b)
```

3.80 $\int \frac{x^3}{a+bx^4} dx$

Optimal result	540
Mathematica [A] (verified)	540
Rubi [A] (verified)	541
Maple [A] (verified)	542
Fricas [A] (verification not implemented)	542
Sympy [A] (verification not implemented)	543
Maxima [A] (verification not implemented)	543
Giac [A] (verification not implemented)	543
Mupad [B] (verification not implemented)	544
Reduce [B] (verification not implemented)	544

Optimal result

Integrand size = 13, antiderivative size = 15

$$\int \frac{x^3}{a+bx^4} dx = \frac{\log(a+bx^4)}{4b}$$

output `1/4*ln(b*x^4+a)/b`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{x^3}{a+bx^4} dx = \frac{\log(a+bx^4)}{4b}$$

input `Integrate[x^3/(a + b*x^4),x]`

output `Log[a + b*x^4]/(4*b)`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {792}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{a + bx^4} dx$$

$$\downarrow 792$$

$$\frac{\log(a + bx^4)}{4b}$$

input `Int[x^3/(a + b*x^4), x]`

output `Log[a + b*x^4]/(4*b)`

Defintions of rubi rules used

rule 792 `Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$\frac{\ln(bx^4+a)}{4b}$	14
default	$\frac{\ln(bx^4+a)}{4b}$	14
norman	$\frac{\ln(bx^4+a)}{4b}$	14
risch	$\frac{\ln(bx^4+a)}{4b}$	14
parallelrisch	$\frac{\ln(bx^4+a)}{4b}$	14

input `int(x^3/(b*x^4+a),x,method=_RETURNVERBOSE)`output `1/4*ln(b*x^4+a)/b`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^3}{a + bx^4} dx = \frac{\log(bx^4 + a)}{4b}$$

input `integrate(x^3/(b*x^4+a),x, algorithm="fricas")`output `1/4*log(b*x^4 + a)/b`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.67

$$\int \frac{x^3}{a + bx^4} dx = \frac{\log(a + bx^4)}{4b}$$

input `integrate(x**3/(b*x**4+a),x)`output `log(a + b*x**4)/(4*b)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^3}{a + bx^4} dx = \frac{\log(bx^4 + a)}{4b}$$

input `integrate(x^3/(b*x^4+a),x, algorithm="maxima")`output `1/4*log(b*x^4 + a)/b`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{x^3}{a + bx^4} dx = \frac{\log(|bx^4 + a|)}{4b}$$

input `integrate(x^3/(b*x^4+a),x, algorithm="giac")`output `1/4*log(abs(b*x^4 + a))/b`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{x^3}{a + bx^4} dx = \frac{\ln(bx^4 + a)}{4b}$$

input `int(x^3/(a + b*x^4),x)`output `log(a + b*x^4)/(4*b)`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 47, normalized size of antiderivative = 3.13

$$\int \frac{x^3}{a + bx^4} dx = \frac{\log\left(-b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}x + \sqrt{a} + \sqrt{b}x^2\right) + \log\left(b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}x + \sqrt{a} + \sqrt{b}x^2\right)}{4b}$$

input `int(x^3/(b*x^4+a),x)`output `(log(-b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2) + log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2))/(4*b)`

3.81 $\int \frac{1}{(a+bx^4)^2} dx$

Optimal result	545
Mathematica [B] (verified)	545
Rubi [B] (verified)	546
Maple [C] (verified)	550
Fricas [C] (verification not implemented)	551
Sympy [A] (verification not implemented)	551
Maxima [B] (verification not implemented)	552
Giac [B] (verification not implemented)	553
Mupad [B] (verification not implemented)	553
Reduce [B] (verification not implemented)	554

Optimal result

Integrand size = 9, antiderivative size = 81

$$\int \frac{1}{(a+bx^4)^2} dx = \frac{x}{4a(a+bx^4)} + \frac{3\sqrt[4]{-\frac{a}{b}} \left(2 \arctan \left(\frac{x}{\sqrt[4]{-\frac{a}{b}}} \right) + \log \left(\frac{\sqrt[4]{-\frac{a}{b}+x}}{-\sqrt[4]{-\frac{a}{b}+x}} \right) \right)}{16a^2}$$

output

`1/4*x/a/(b*x^4+a)+3/16*(-a/b)^(1/4)*(2*arctan(x/(-a/b)^(1/4))+ln(((a/b)^(1/4)+x)/(-(a/b)^(1/4)+x)))/a^2`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 183 vs. 2(81) = 162.

Time = 0.08 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.26

$$\int \frac{1}{(a+bx^4)^2} dx = \frac{8a^{3/4}x}{a+bx^4} - \frac{6\sqrt{2} \arctan \left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}} \right)}{\sqrt[4]{b}} + \frac{6\sqrt{2} \arctan \left(1 + \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}} \right)}{\sqrt[4]{b}} - \frac{3\sqrt{2} \log \left(\sqrt{a} - \sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x + \sqrt{bx^2} \right)}{\sqrt[4]{b}} + \frac{3\sqrt{2} \log \left(\sqrt{a} + \sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x + \sqrt{bx^2} \right)}{\sqrt[4]{b}}$$

$32a^{7/4}$

input `Integrate[(a + b*x^4)^(-2),x]`

output $((8*a^{(3/4)*x})/(a + b*x^4) - (6*\text{Sqrt}[2]*\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{(1/4)*x})/a^{(1/4)}])/b^{(1/4)} + (6*\text{Sqrt}[2]*\text{ArcTan}[1 + (\text{Sqrt}[2]*b^{(1/4)*x})/a^{(1/4)}])/b^{(1/4)} - (3*\text{Sqrt}[2]*\text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{(1/4)*b^{(1/4)*x}} + \text{Sqrt}[b]*x^2])/b^{(1/4)} + (3*\text{Sqrt}[2]*\text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{(1/4)*b^{(1/4)*x}} + \text{Sqrt}[b]*x^2])/b^{(1/4)})/(32*a^{(7/4)})$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 225 vs. 2(81) = 162.

Time = 0.72 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.78, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {749, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx^4)^2} dx$$

$$\downarrow 749$$

$$\frac{3 \int \frac{1}{bx^4+a} dx}{4a} + \frac{x}{4a(a + bx^4)}$$

$$\downarrow 755$$

$$\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{b}x^2}{bx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{\sqrt{b}x^2+\sqrt{a}}{bx^4+a} dx}{2\sqrt{a}} \right)}{4a} + \frac{x}{4a(a + bx^4)}$$

$$\downarrow 1476$$

$$3 \left(\frac{\int \frac{1}{x^2 - \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} + \frac{\int \frac{1}{x^2 + \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} + \frac{\int \frac{\sqrt{a}-\sqrt{b}x^2}{bx^4+a} dx}{2\sqrt{a}} \right) + \frac{x}{4a(a + bx^4)}$$

$$\begin{aligned}
 & \downarrow 1082 \\
 & 3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{a}} + \frac{\int \frac{1}{\left(1-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt{a}}\right)^2} d\left(1-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt{a}}\right)^2} d\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right) \\
 & \frac{4a}{x} \\
 & \frac{4a}{4a(a+bx^4)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 217 \\
 & 3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{a}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right) \\
 & \frac{x}{4a(a+bx^4)}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 1479 \\
 & 3 \left(\frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{bx}}{\sqrt[4]{b}\left(x^2-\frac{\sqrt{2}\sqrt[4]{ax}}{\sqrt{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{bx}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x^2+\frac{\sqrt{2}\sqrt[4]{ax}}{\sqrt{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right) \\
 & \frac{4a}{x} \\
 & \frac{4a}{4a(a+bx^4)}
 \end{aligned}$$

$$\downarrow 25$$

$$\begin{aligned}
 & 3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a-2} \sqrt[4]{b} x}{\sqrt[4]{b} \left(x^2 - \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} x + \sqrt[4]{a}\right)}{\sqrt[4]{b} \left(x^2 + \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} x + 1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} x}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) \\
 & \frac{4a}{x} \\
 & \frac{4a}{4a(a+bx^4)} \\
 & \downarrow 27 \\
 & 3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a-2} \sqrt[4]{b} x}{x^2 - \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{b}}} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{b} x + \sqrt[4]{a}}{x^2 + \frac{\sqrt{2} \sqrt[4]{a} x + \sqrt{a}}{\sqrt[4]{b}}} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} x + 1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} x}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) \\
 & \frac{4a}{x} \\
 & \frac{4a}{4a(a+bx^4)} \\
 & \downarrow 1103 \\
 & 3 \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} x + 1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} x}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\log\left(\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} x + \sqrt{a} + \sqrt{b} x^2\right) - \log\left(-\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} x + \sqrt{a} + \sqrt{b} x^2\right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) \\
 & \frac{4a}{x} \\
 & \frac{4a}{4a(a+bx^4)}
 \end{aligned}$$

input

`Int[(a + b*x^4)^(-2), x]`

output

```
x/(4*a*(a + b*x^4)) + (3*((-ArcTan[1 - (Sqrt[2]*b^(1/4)*x)/a^(1/4)]/(Sqrt
[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a
^(1/4)*b^(1/4)))/(2*Sqrt[a]) + (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)
*x + Sqrt[b]*x^2]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)
)*b^(1/4)*x + Sqrt[b]*x^2]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[a]))/(4*a
)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 217

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

rule 749

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p +
1)/(a*n*(p + 1))), x] + Simp[(n*(p + 1) + 1)/(a*n*(p + 1)) Int[(a + b*x^
n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (Inte
gerQ[2*p] || Denominator[p + 1/n] < Denominator[p])
```

rule 755

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4)
, x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

rule 1082

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.57

method	result	size
risch	$\frac{x}{4a(bx^4+a)} + \frac{3 \left(\sum_{R=\text{RootOf}(bZ^4+a)} \frac{\ln(x-R)}{-R^3} \right)}{16ab}$	46
default	$\frac{x}{4a(bx^4+a)} + \frac{3 \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{x^2 + \left(\frac{a}{b} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}}{x^2 - \left(\frac{a}{b} \right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}} \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} + 1 \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} - 1 \right) \right)}{32a^2}$	118

input `int(1/(b*x^4+a)^2,x,method=_RETURNVERBOSE)`

output `1/4*x/a/(b*x^4+a)+3/16/a/b*sum(1/_R^3*ln(x-_R),_R=RootOf(_Z^4*b+a))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.26

$$\int \frac{1}{(a + bx^4)^2} dx$$

$$= \frac{3(abx^4 + a^2)\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} \log\left(a^2\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} + x\right) - 3(-i abx^4 - i a^2)\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} \log\left(i a^2\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} + x\right) - 3(i abx^4 + i a^2)\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} \log\left(-i a^2\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} + x\right) + 4x}{16(abx^4 + a^2)}$$

input `integrate(1/(b*x^4+a)^2,x, algorithm="fricas")`

output
$$\frac{1}{16} \frac{3(a b x^4 + a^2) \left(-\frac{1}{a^7 b}\right)^{\frac{1}{4}} \log\left(a^2 \left(-\frac{1}{a^7 b}\right)^{\frac{1}{4}} + x\right) - 3(-i a b x^4 - i a^2) \left(-\frac{1}{a^7 b}\right)^{\frac{1}{4}} \log\left(i a^2 \left(-\frac{1}{a^7 b}\right)^{\frac{1}{4}} + x\right) - 3(i a b x^4 + i a^2) \left(-\frac{1}{a^7 b}\right)^{\frac{1}{4}} \log\left(-i a^2 \left(-\frac{1}{a^7 b}\right)^{\frac{1}{4}} + x\right) + 4x}{(a b x^4 + a^2)}$$

Sympy [A] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.48

$$\int \frac{1}{(a + bx^4)^2} dx$$

$$= \frac{x}{4a^2 + 4abx^4} + \text{RootSum}\left(65536t^4 a^7 b + 81, \left(t \mapsto t \log\left(\frac{16ta^2}{3} + x\right)\right)\right)$$

input `integrate(1/(b*x**4+a)**2,x)`

output
$$\frac{x}{4a^2 + 4abx^4} + \text{RootSum}(65536*_t**4*a**7*b + 81, \text{Lambda}(_t, _t*\log(16*_t*a**2/3 + x)))$$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(69) = 138$.

Time = 0.11 (sec) , antiderivative size = 189, normalized size of antiderivative = 2.33

$$\int \frac{1}{(a + bx^4)^2} dx = \frac{x}{4(abx^4 + a^2)} + \frac{3}{32a} \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} + \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} - \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{\sqrt{2} \log(\sqrt{bx^2 + \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}x + \sqrt{a}})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log(\sqrt{bx^2 - \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}x + \sqrt{a}})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} \right)$$

input `integrate(1/(b*x^4+a)^2,x, algorithm="maxima")`

output

```
1/4*x/(a*b*x^4 + a^2) + 3/32*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x +
sqrt(2)*a^(1/4)*b^(1/4))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(a)*sqrt(sqrt(a)*sqrt
(b))) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x - sqrt(2)*a^(1/4)*b^(1/4
))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(a)*sqrt(sqrt(a)*sqrt(b))) + sqrt(2)*log(sq
rt(b)*x^2 + sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(3/4)*b^(1/4)) - sqrt(
2)*log(sqrt(b)*x^2 - sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(3/4)*b^(1/4
))/a
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. $2(69) = 138$.

Time = 0.12 (sec) , antiderivative size = 194, normalized size of antiderivative = 2.40

$$\int \frac{1}{(a + bx^4)^2} dx = \frac{x}{4(bx^4 + a)a} + \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x + \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{16a^2b}$$

$$+ \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(2x - \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{16a^2b}$$

$$+ \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{32a^2b}$$

$$- \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{32a^2b}$$

input `integrate(1/(b*x^4+a)^2,x, algorithm="giac")`

output `1/4*x/((b*x^4 + a)*a) + 3/16*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x + sqrt(2)*(a/b)^(1/4))/(a/b)^(1/4))/(a^2*b) + 3/16*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(2*x - sqrt(2)*(a/b)^(1/4))/(a/b)^(1/4))/(a^2*b) + 3/32*sqrt(2)*(a*b^3)^(1/4)*log(x^2 + sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a^2*b) - 3/32*sqrt(2)*(a*b^3)^(1/4)*log(x^2 - sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a^2*b)`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.72

$$\int \frac{1}{(a + bx^4)^2} dx = \frac{x}{4a(bx^4 + a)} + \frac{3 \operatorname{atan}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{7/4}b^{1/4}} + \frac{3 \operatorname{atanh}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{7/4}b^{1/4}}$$

input `int(1/(a + b*x^4)^2,x)`

output

$$\frac{x}{4*a*(a + b*x^4)} + \frac{(3*atan((b^(1/4)*x)/(-a)^(1/4)))/(8*(-a)^(7/4)*b^(1/4)) + (3*atanh((b^(1/4)*x)/(-a)^(1/4)))/(8*(-a)^(7/4)*b^(1/4))}{1}$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 305, normalized size of antiderivative = 3.77

$$\int \frac{1}{(a + bx^4)^2} dx$$

$$= \frac{-6b^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) - 6b^{\frac{7}{4}}a^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^4 + 6b^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) + 6b^{\frac{7}{4}}a^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^4}{(a + bx^4)^2}$$

input

int(1/(b*x^4+a)^2,x)

output

```
( - 6*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*a - 6*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**4 + 6*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*a + 6*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**4 - 3*b**(3/4)*a**(1/4)*sqrt(2)*log(- b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*a - 3*b**(3/4)*a**(1/4)*sqrt(2)*log(- b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*b*x**4 + 3*b**(3/4)*a**(1/4)*sqrt(2)*log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*a + 3*b**(3/4)*a**(1/4)*sqrt(2)*log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*b*x**4 + 8*a*b*x)/(32*a**2*b*(a + b*x**4))
```

3.82 $\int \frac{x}{(a+bx^4)^2} dx$

Optimal result	555
Mathematica [A] (verified)	555
Rubi [A] (verified)	556
Maple [A] (verified)	557
Fricas [A] (verification not implemented)	558
Sympy [B] (verification not implemented)	558
Maxima [A] (verification not implemented)	559
Giac [A] (verification not implemented)	559
Mupad [B] (verification not implemented)	559
Reduce [B] (verification not implemented)	560

Optimal result

Integrand size = 11, antiderivative size = 48

$$\int \frac{x}{(a+bx^4)^2} dx = \frac{x^2}{4a(a+bx^4)} + \frac{\arctan\left(\sqrt{\frac{b}{a}}x^2\right)}{4a\sqrt{ab}}$$

output $1/4*x^2/a/(b*x^4+a)+1/4*\arctan((b/a)^{(1/2)}*x^2)/a/(a*b)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.02

$$\int \frac{x}{(a+bx^4)^2} dx = \frac{x^2}{4a(a+bx^4)} + \frac{\arctan\left(\frac{\sqrt{bx^2}}{\sqrt{a}}\right)}{4a^{3/2}\sqrt{b}}$$

input `Integrate[x/(a + b*x^4)^2,x]`

output $x^2/(4*a*(a + b*x^4)) + \text{ArcTan}[(\text{Sqrt}[b]*x^2)/\text{Sqrt}[a]]/(4*a^{(3/2)}*\text{Sqrt}[b])$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.10, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {807, 215, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{(a + bx^4)^2} dx \\
 & \quad \downarrow \text{807} \\
 & \frac{1}{2} \int \frac{1}{(bx^4 + a)^2} dx^2 \\
 & \quad \downarrow \text{215} \\
 & \frac{1}{2} \left(\frac{\int \frac{1}{bx^4+a} dx^2}{2a} + \frac{x^2}{2a(a + bx^4)} \right) \\
 & \quad \downarrow \text{218} \\
 & \frac{1}{2} \left(\frac{\arctan\left(\frac{\sqrt{b}x^2}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{b}} + \frac{x^2}{2a(a + bx^4)} \right)
 \end{aligned}$$

input `Int[x/(a + b*x^4)^2,x]`

output `(x^2/(2*a*(a + b*x^4)) + ArcTan[(Sqrt[b]*x^2)/Sqrt[a]]/(2*a^(3/2)*Sqrt[b]))/2`

Definitions of rubi rules used

rule 215 `Int[((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-x)*((a + b*x^2)^(p + 1) / (2*a*(p + 1))), x] + Simp[(2*p + 3)/(2*a*(p + 1)) Int[(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b}, x] && LtQ[p, -1] && (IntegerQ[4*p] || IntegerQ[6*p])`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{x^2}{4a(bx^4+a)} + \frac{\arctan\left(\frac{bx^2}{\sqrt{ab}}\right)}{4a\sqrt{ab}}$	40
risch	$\frac{x^2}{4a(bx^4+a)} - \frac{\ln(x^2\sqrt{-ab}-a)}{8\sqrt{-ab}a} + \frac{\ln(x^2\sqrt{-ab}+a)}{8\sqrt{-ab}a}$	69

input `int(x/(b*x^4+a)^2,x,method=_RETURNVERBOSE)`

output `1/4*x^2/a/(b*x^4+a)+1/4/a/(a*b)^(1/2)*arctan(b*x^2/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 129, normalized size of antiderivative = 2.69

$$\int \frac{x}{(a + bx^4)^2} dx = \left[\frac{2abx^2 - (bx^4 + a)\sqrt{-ab} \log\left(\frac{bx^4 - 2\sqrt{-ab}x^2 - a}{bx^4 + a}\right)}{8(a^2b^2x^4 + a^3b)}, \frac{abx^2 - (bx^4 + a)\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{bx^2}\right)}{4(a^2b^2x^4 + a^3b)} \right]$$

input `integrate(x/(b*x^4+a)^2,x, algorithm="fricas")`

output `[1/8*(2*a*b*x^2 - (b*x^4 + a)*sqrt(-a*b)*log((b*x^4 - 2*sqrt(-a*b)*x^2 - a)/(b*x^4 + a)))/(a^2*b^2*x^4 + a^3*b), 1/4*(a*b*x^2 - (b*x^4 + a)*sqrt(a*b)*arctan(sqrt(a*b)/(b*x^2)))/(a^2*b^2*x^4 + a^3*b)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. 2(34) = 68.

Time = 0.17 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.73

$$\int \frac{x}{(a + bx^4)^2} dx = \frac{x^2}{4a^2 + 4abx^4} - \frac{\sqrt{-\frac{1}{a^3b}} \log\left(-a^2 \sqrt{-\frac{1}{a^3b}} + x^2\right)}{8} + \frac{\sqrt{-\frac{1}{a^3b}} \log\left(a^2 \sqrt{-\frac{1}{a^3b}} + x^2\right)}{8}$$

input `integrate(x/(b*x**4+a)**2,x)`

output `x**2/(4*a**2 + 4*a*b*x**4) - sqrt(-1/(a**3*b))*log(-a**2*sqrt(-1/(a**3*b)) + x**2)/8 + sqrt(-1/(a**3*b))*log(a**2*sqrt(-1/(a**3*b)) + x**2)/8`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.81

$$\int \frac{x}{(a + bx^4)^2} dx = \frac{x^2}{4(abx^4 + a^2)} + \frac{\arctan\left(\frac{bx^2}{\sqrt{ab}}\right)}{4\sqrt{aba}}$$

input `integrate(x/(b*x^4+a)^2,x, algorithm="maxima")`output `1/4*x^2/(a*b*x^4 + a^2) + 1/4*arctan(b*x^2/sqrt(a*b))/(sqrt(a*b)*a)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.81

$$\int \frac{x}{(a + bx^4)^2} dx = \frac{x^2}{4(bx^4 + a)a} + \frac{\arctan\left(\frac{bx^2}{\sqrt{ab}}\right)}{4\sqrt{aba}}$$

input `integrate(x/(b*x^4+a)^2,x, algorithm="giac")`output `1/4*x^2/((b*x^4 + a)*a) + 1/4*arctan(b*x^2/sqrt(a*b))/(sqrt(a*b)*a)`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.77

$$\int \frac{x}{(a + bx^4)^2} dx = \frac{x^2}{4a(bx^4 + a)} + \frac{\operatorname{atan}\left(\frac{\sqrt{b}x^2}{\sqrt{a}}\right)}{4a^{3/2}\sqrt{b}}$$

input `int(x/(a + b*x^4)^2,x)`output `x^2/(4*a*(a + b*x^4)) + atan((b^(1/2)*x^2)/a^(1/2))/(4*a^(3/2)*b^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 166, normalized size of antiderivative = 3.46

$$\int \frac{x}{(a + bx^4)^2} dx$$

$$= \frac{-\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{b}x}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) a - \sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{b}x}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) bx^4 - \sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{b}x}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) a - \sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{b}x}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) bx^4}{4a^2b(bx^4 + a)}$$

input `int(x/(b*x^4+a)^2,x)`output `(- sqrt(b)*sqrt(a)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*a - sqrt(b)*sqrt(a)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**4 - sqrt(b)*sqrt(a)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*a - sqrt(b)*sqrt(a)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**4 + a*b*x**2)/(4*a**2*b*(a + b*x**4))`

3.83 $\int \frac{x^2}{(a+bx^4)^2} dx$

Optimal result	561
Mathematica [B] (verified)	561
Rubi [B] (verified)	562
Maple [C] (verified)	566
Fricas [C] (verification not implemented)	566
Sympy [A] (verification not implemented)	567
Maxima [B] (verification not implemented)	567
Giac [B] (verification not implemented)	568
Mupad [B] (verification not implemented)	568
Reduce [B] (verification not implemented)	569

Optimal result

Integrand size = 13, antiderivative size = 86

$$\int \frac{x^2}{(a+bx^4)^2} dx = \frac{x^3}{4a(a+bx^4)} - \frac{-2 \arctan\left(\frac{x}{\sqrt[4]{-\frac{a}{b}}}\right) + \log\left(\frac{\sqrt[4]{-\frac{a}{b}}+x}{-\sqrt[4]{-\frac{a}{b}}+x}\right)}{16a\sqrt[4]{-\frac{a}{b}}}$$

output

```
1/4*x^3/a/(b*x^4+a)-1/16*(-2*arctan(x/(-a/b)^(1/4))+ln(((a/b)^(1/4)+x)/(-a/b)^(1/4)+x)))/a/(-a/b)^(1/4)/b
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 184 vs. 2(86) = 172.

Time = 0.08 (sec) , antiderivative size = 184, normalized size of antiderivative = 2.14

$$\int \frac{x^2}{(a+bx^4)^2} dx = \frac{8\sqrt[4]{ax^3}}{a+bx^4} - \frac{2\sqrt{2} \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{b^{3/4}} + \frac{2\sqrt{2} \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{b^{3/4}} + \frac{\sqrt{2} \log\left(\sqrt{a} - \sqrt{2}\sqrt[4]{a}\sqrt[4]{bx} + \sqrt{bx^2}\right)}{b^{3/4}} - \frac{\sqrt{2} \log\left(\sqrt{a} + \sqrt{2}\sqrt[4]{a}\sqrt[4]{bx} + \sqrt{bx^2}\right)}{b^{3/4}}$$

$32a^{5/4}$

input `Integrate[x^2/(a + b*x^4)^2,x]`

output $((8*a^{1/4}*x^3)/(a + b*x^4) - (2*\text{Sqrt}[2]*\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{1/4}*x)/a^{1/4}])/b^{3/4} + (2*\text{Sqrt}[2]*\text{ArcTan}[1 + (\text{Sqrt}[2]*b^{1/4}*x)/a^{1/4}])/b^{3/4} + (\text{Sqrt}[2]*\text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{1/4}*b^{1/4}*x + \text{Sqrt}[b]*x^2])/b^{3/4} - (\text{Sqrt}[2]*\text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{1/4}*b^{1/4}*x + \text{Sqrt}[b]*x^2])/b^{3/4}))/ (32*a^{5/4})$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 227 vs. $2(86) = 172$.

Time = 0.69 (sec) , antiderivative size = 227, normalized size of antiderivative = 2.64, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {819, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^2}{(a + bx^4)^2} dx \\
 & \quad \downarrow 819 \\
 & \int \frac{x^2}{bx^4+a} dx + \frac{x^3}{4a(a + bx^4)} \\
 & \quad \downarrow 826 \\
 & \frac{\int \frac{\sqrt{bx^2+\sqrt{a}}}{bx^4+a} dx}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{b}} + \frac{x^3}{4a(a + bx^4)} \\
 & \quad \downarrow 1476 \\
 & \frac{\int \frac{1}{x^2 - \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{b}} + \frac{\sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} + \frac{\int \frac{1}{x^2 + \frac{\sqrt{2}\sqrt[4]{a}x + \sqrt{a}}{\sqrt{b}} + \frac{\sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{b}} + \frac{x^3}{4a(a + bx^4)} \\
 & \quad \downarrow 1082
 \end{aligned}$$

$$\frac{\int \frac{1}{\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)^2} d\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right) - \int \frac{1}{\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right)^2} d\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{b}} + \frac{x^3}{4a(a+bx^4)}$$

217

$$\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{b}} + \frac{x^3}{4a(a+bx^4)}$$

1479

$$\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{2\sqrt{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{bx}}{\sqrt[4]{b}\left(x^2 - \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{bx} + \sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x^2 + \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{4a^3 x^3}{4a(a+bx^4)}$$

25

$$\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{2\sqrt{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{bx}}{\sqrt[4]{b}\left(x^2 - \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{bx} + \sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x^2 + \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{4a^3 x^3}{4a(a+bx^4)}$$

27

$$\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{2\sqrt{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{bx}}{x^2 - \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{bx} + \sqrt[4]{a}}{x^2 + \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}} dx}{2\sqrt[4]{a}\sqrt[4]{b}} + \frac{x^3}{4a(a+bx^4)}$$

1103

$$\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}x+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x+\sqrt{a}+\sqrt{b}x^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x+\sqrt{a}+\sqrt{b}x^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{\frac{4a}{x^3}} + \frac{4a}{4a(a+bx^4)}$$

input `Int[x^2/(a + b*x^4)^2,x]`

output `x^3/(4*a*(a + b*x^4)) + ((-(ArcTan[1 - (Sqrt[2]*b^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]) - (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*x + Sqrt[b]*x^2]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*x + Sqrt[b]*x^2]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]))/(4*a)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 819 `Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^n)^(p + 1)/(a*c*n*(p + 1))), x] + Simp[(m + n*(p + 1) + 1)/(a*n*(p + 1)) Int[(c*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.56

method	result	size
risch	$\frac{x^3}{4a(bx^4+a)} + \frac{\sum_{R=\text{RootOf}(bZ^4+a)} \frac{\ln(x-R)}{-R}}{16ab}$	48
default	$\frac{x^3}{4a(bx^4+a)} + \frac{\sqrt{2} \left(\ln \left(\frac{x^2 - \left(\frac{a}{b}\right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}}{x^2 + \left(\frac{a}{b}\right)^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}} \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{b}\right)^{\frac{1}{4}} + 1} \right) + 2 \arctan \left(\frac{\sqrt{2} x}{\left(\frac{a}{b}\right)^{\frac{1}{4}} - 1} \right) \right)}{32ab \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	123

input `int(x^2/(b*x^4+a)^2,x,method=_RETURNVERBOSE)`

output `1/4*x^3/a/(b*x^4+a)+1/16/a/b*sum(1/_R*ln(x-_R),_R=RootOf(_Z^4*b+a))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.28

$$\int \frac{x^2}{(a+bx^4)^2} dx$$

$$= \frac{4x^3 + (abx^4 + a^2) \left(-\frac{1}{a^5b^3}\right)^{\frac{1}{4}} \log\left(a^4b^2 \left(-\frac{1}{a^5b^3}\right)^{\frac{3}{4}} + x\right) - (i abx^4 + i a^2) \left(-\frac{1}{a^5b^3}\right)^{\frac{1}{4}} \log\left(i a^4b^2 \left(-\frac{1}{a^5b^3}\right)^{\frac{3}{4}} + x\right)}{16(ab)}$$

input `integrate(x^2/(b*x^4+a)^2,x, algorithm="fricas")`

output `1/16*(4*x^3 + (a*b*x^4 + a^2)*(-1/(a^5*b^3))^(1/4)*log(a^4*b^2*(-1/(a^5*b^3))^(3/4) + x) - (I*a*b*x^4 + I*a^2)*(-1/(a^5*b^3))^(1/4)*log(I*a^4*b^2*(-1/(a^5*b^3))^(3/4) + x) - (-I*a*b*x^4 - I*a^2)*(-1/(a^5*b^3))^(1/4)*log(-I*a^4*b^2*(-1/(a^5*b^3))^(3/4) + x) - (a*b*x^4 + a^2)*(-1/(a^5*b^3))^(1/4)*log(-a^4*b^2*(-1/(a^5*b^3))^(3/4) + x))/(a*b*x^4 + a^2)`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.53

$$\int \frac{x^2}{(a + bx^4)^2} dx = \frac{x^3}{4a^2 + 4abx^4} + \text{RootSum}(65536t^4a^5b^3 + 1, (t \mapsto t \log(4096t^3a^4b^2 + x)))$$

input `integrate(x**2/(b*x**4+a)**2,x)`

output `x**3/(4*a**2 + 4*a*b*x**4) + RootSum(65536*_t**4*a**5*b**3 + 1, Lambda(_t, _t*log(4096*_t**3*a**4*b**2 + x)))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 191 vs. $2(76) = 152$.

Time = 0.12 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.22

$$\int \frac{x^2}{(a + bx^4)^2} dx = \frac{x^3}{4(abx^4 + a^2)} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} + \sqrt{2a^{1/4}b^{1/4}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} - \sqrt{2a^{1/4}b^{1/4}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} - \frac{\sqrt{2} \log(\sqrt{bx^2 + \sqrt{2a^{1/4}b^{1/4}}x + \sqrt{a}})}{a^{1/4}b^{3/4}} + \frac{\sqrt{2} \log(\sqrt{bx^2 - \sqrt{2a^{1/4}b^{1/4}}x + \sqrt{a}})}{a^{1/4}b^{3/4}}$$

$32a$

input `integrate(x^2/(b*x^4+a)^2,x, algorithm="maxima")`

output `1/4*x^3/(a*b*x^4 + a^2) + 1/32*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x + sqrt(2)*a^(1/4)*b^(1/4))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x - sqrt(2)*a^(1/4)*b^(1/4))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) - sqrt(2)*log(sqrt(b)*x^2 + sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(1/4)*b^(3/4)) + sqrt(2)*log(sqrt(b)*x^2 - sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(1/4)*b^(3/4)))/a`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. $2(76) = 152$.

Time = 0.12 (sec) , antiderivative size = 196, normalized size of antiderivative = 2.28

$$\int \frac{x^2}{(a + bx^4)^2} dx = \frac{x^3}{4(bx^4 + a)a} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(2x + \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{16 a^2 b^3}$$

$$+ \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(2x - \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{16 a^2 b^3}$$

$$- \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{32 a^2 b^3}$$

$$+ \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{32 a^2 b^3}$$

input

```
integrate(x^2/(b*x^4+a)^2,x, algorithm="giac")
```

output

```
1/4*x^3/((b*x^4 + a)*a) + 1/16*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*(2*x + sqrt(2)*(a/b)^(1/4))/(a/b)^(1/4))/(a^2*b^3) + 1/16*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*(2*x - sqrt(2)*(a/b)^(1/4))/(a/b)^(1/4))/(a^2*b^3) - 1/32*sqrt(2)*(a*b^3)^(3/4)*log(x^2 + sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a^2*b^3) + 1/32*sqrt(2)*(a*b^3)^(3/4)*log(x^2 - sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a^2*b^3)
```

Mupad [B] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.70

$$\int \frac{x^2}{(a + bx^4)^2} dx = \frac{\operatorname{atanh}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{5/4}b^{3/4}} - \frac{\operatorname{atan}\left(\frac{b^{1/4}x}{(-a)^{1/4}}\right)}{8(-a)^{5/4}b^{3/4}} + \frac{x^3}{4a(bx^4 + a)}$$

input

```
int(x^2/(a + b*x^4)^2,x)
```

output

```
atanh((b^(1/4)*x)/(-a)^(1/4))/(8*(-a)^(5/4)*b^(3/4)) - atan((b^(1/4)*x)/(-a)^(1/4))/(8*(-a)^(5/4)*b^(3/4)) + x^3/(4*a*(a + b*x^4))
```

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 305, normalized size of antiderivative = 3.55

$$\int \frac{x^2}{(a + bx^4)^2} dx$$

$$= \frac{-2b^{\frac{1}{4}}a^{\frac{7}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) - 2b^{\frac{5}{4}}a^{\frac{3}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^4 + 2b^{\frac{1}{4}}a^{\frac{7}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) + 2b^{\frac{5}{4}}a^{\frac{3}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{bx}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^4}{(a + bx^4)^2}$$

input

```
int(x^2/(b*x^4+a)^2,x)
```

output

```
( - 2*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*a - 2*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**4 + 2*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*a + 2*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(b)*x)/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**4 + b**(1/4)*a**(3/4)*sqrt(2)*log(- b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*a + b**(1/4)*a**(3/4)*sqrt(2)*log(- b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*b*x**4 - b**(1/4)*a**(3/4)*sqrt(2)*log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*a - b**(1/4)*a**(3/4)*sqrt(2)*log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*b*x**4 + 8*a*b*x**3)/(32*a**2*b*(a + b*x**4))
```

3.84 $\int \frac{x^3}{(a+bx^4)^2} dx$

Optimal result	570
Mathematica [A] (verified)	570
Rubi [A] (verified)	571
Maple [A] (verified)	572
Fricas [A] (verification not implemented)	572
Sympy [A] (verification not implemented)	573
Maxima [A] (verification not implemented)	573
Giac [A] (verification not implemented)	573
Mupad [B] (verification not implemented)	574
Reduce [B] (verification not implemented)	574

Optimal result

Integrand size = 13, antiderivative size = 16

$$\int \frac{x^3}{(a+bx^4)^2} dx = -\frac{1}{4b(a+bx^4)}$$

output `-1/4/b/(b*x^4+a)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{x^3}{(a+bx^4)^2} dx = -\frac{1}{4b(a+bx^4)}$$

input `Integrate[x^3/(a + b*x^4)^2,x]`

output `-1/4*1/(b*(a + b*x^4))`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {793}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^3}{(a + bx^4)^2} dx$$

↓ 793

$$-\frac{1}{4b(a + bx^4)}$$

input `Int[x^3/(a + b*x^4)^2,x]`

output `-1/4*1/(b*(a + b*x^4))`

Defintions of rubi rules used

rule 793

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
gospers	$-\frac{1}{4b(bx^4+a)}$	15
derivativedivides	$-\frac{1}{4b(bx^4+a)}$	15
default	$-\frac{1}{4b(bx^4+a)}$	15
norman	$-\frac{1}{4b(bx^4+a)}$	15
risch	$-\frac{1}{4b(bx^4+a)}$	15
parallelrisch	$-\frac{1}{4b(bx^4+a)}$	15
orering	$-\frac{1}{4b(bx^4+a)}$	15

input `int(x^3/(b*x^4+a)^2,x,method=_RETURNVERBOSE)`output `-1/4/b/(b*x^4+a)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{x^3}{(a + bx^4)^2} dx = -\frac{1}{4(b^2x^4 + ab)}$$

input `integrate(x^3/(b*x^4+a)^2,x, algorithm="fricas")`output `-1/4/(b^2*x^4 + a*b)`

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{x^3}{(a + bx^4)^2} dx = -\frac{1}{4ab + 4b^2x^4}$$

input `integrate(x**3/(b*x**4+a)**2,x)`output `-1/(4*a*b + 4*b**2*x**4)`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x^3}{(a + bx^4)^2} dx = -\frac{1}{4(bx^4 + a)b}$$

input `integrate(x^3/(b*x^4+a)^2,x, algorithm="maxima")`output `-1/4/((b*x^4 + a)*b)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x^3}{(a + bx^4)^2} dx = -\frac{1}{4(bx^4 + a)b}$$

input `integrate(x^3/(b*x^4+a)^2,x, algorithm="giac")`output `-1/4/((b*x^4 + a)*b)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{x^3}{(a + bx^4)^2} dx = -\frac{1}{4b(bx^4 + a)}$$

input `int(x^3/(a + b*x^4)^2,x)`

output `-1/(4*b*(a + b*x^4))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{x^3}{(a + bx^4)^2} dx = \frac{x^4}{4a(bx^4 + a)}$$

input `int(x^3/(b*x^4+a)^2,x)`

output `x**4/(4*a*(a + b*x**4))`

3.85 $\int \frac{1}{x(a+bx^4)} dx$

Optimal result	575
Mathematica [A] (verified)	575
Rubi [A] (verified)	576
Maple [A] (verified)	577
Fricas [A] (verification not implemented)	578
Sympy [A] (verification not implemented)	578
Maxima [A] (verification not implemented)	578
Giac [A] (verification not implemented)	579
Mupad [B] (verification not implemented)	579
Reduce [B] (verification not implemented)	579

Optimal result

Integrand size = 13, antiderivative size = 21

$$\int \frac{1}{x(a+bx^4)} dx = \frac{\log\left(\frac{x^4}{a+bx^4}\right)}{4a}$$

output `1/4*ln(x^4/(b*x^4+a))/a`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{1}{x(a+bx^4)} dx = \frac{\log(x)}{a} - \frac{\log(a+bx^4)}{4a}$$

input `Integrate[1/(x*(a + b*x^4)),x]`

output `Log[x]/a - Log[a + b*x^4]/(4*a)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.24, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {798, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x(a+bx^4)} dx \\
 & \quad \downarrow 798 \\
 & \frac{1}{4} \int \frac{1}{x^4(bx^4+a)} dx^4 \\
 & \quad \downarrow 47 \\
 & \frac{1}{4} \left(\frac{\int \frac{1}{x^4} dx^4}{a} - \frac{b \int \frac{1}{bx^4+a} dx^4}{a} \right) \\
 & \quad \downarrow 14 \\
 & \frac{1}{4} \left(\frac{\log(x^4)}{a} - \frac{b \int \frac{1}{bx^4+a} dx^4}{a} \right) \\
 & \quad \downarrow 16 \\
 & \frac{1}{4} \left(\frac{\log(x^4)}{a} - \frac{\log(a+bx^4)}{a} \right)
 \end{aligned}$$

input `Int[1/(x*(a + b*x^4)),x]`

output `(Log[x^4]/a - Log[a + b*x^4]/a)/4`

Definitions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{\ln(bx^4+a)}{4a} + \frac{\ln(x)}{a}$	21
norman	$-\frac{\ln(bx^4+a)}{4a} + \frac{\ln(x)}{a}$	21
risch	$-\frac{\ln(bx^4+a)}{4a} + \frac{\ln(x)}{a}$	21
parallelrisc	$\frac{4\ln(x) - \ln(bx^4+a)}{4a}$	21

input `int(1/x/(b*x^4+a),x,method=_RETURNVERBOSE)`

output `-1/4/a*ln(b*x^4+a)+1/a*ln(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \frac{1}{x(a+bx^4)} dx = -\frac{\log(bx^4+a) - 4\log(x)}{4a}$$

input `integrate(1/x/(b*x^4+a),x, algorithm="fricas")`output `-1/4*(log(b*x^4 + a) - 4*log(x))/a`**Sympy [A] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1}{x(a+bx^4)} dx = \frac{\log(x)}{a} - \frac{\log(\frac{a}{b} + x^4)}{4a}$$

input `integrate(1/x/(b*x**4+a),x)`output `log(x)/a - log(a/b + x**4)/(4*a)`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{1}{x(a+bx^4)} dx = -\frac{\log(bx^4+a)}{4a} + \frac{\log(x^4)}{4a}$$

input `integrate(1/x/(b*x^4+a),x, algorithm="maxima")`output `-1/4*log(b*x^4 + a)/a + 1/4*log(x^4)/a`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.14

$$\int \frac{1}{x(a+bx^4)} dx = \frac{\log(x^4)}{4a} - \frac{\log(|bx^4+a|)}{4a}$$

input `integrate(1/x/(b*x^4+a),x, algorithm="giac")`

output `1/4*log(x^4)/a - 1/4*log(abs(b*x^4 + a))/a`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \frac{1}{x(a+bx^4)} dx = -\frac{\ln(bx^4+a) - 4 \ln(x)}{4a}$$

input `int(1/(x*(a + b*x^4)),x)`

output `-(log(a + b*x^4) - 4*log(x))/(4*a)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.62

$$\int \frac{1}{x(a+bx^4)} dx = \frac{-\log\left(-b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}x + \sqrt{a} + \sqrt{b}x^2\right) - \log\left(b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}x + \sqrt{a} + \sqrt{b}x^2\right) + 4\log(x)}{4a}$$

input `int(1/x/(b*x^4+a),x)`

output `(- log(- b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2) - log(b**(1/4)*a**(1/4)*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2) + 4*log(x))/(4*a)`

3.86 $\int \frac{1}{x^2(a+bx^4)} dx$

Optimal result	580
Mathematica [B] (verified)	580
Rubi [B] (verified)	581
Maple [C] (verified)	585
Fricas [C] (verification not implemented)	586
Sympy [A] (verification not implemented)	586
Maxima [B] (verification not implemented)	587
Giac [B] (verification not implemented)	588
Mupad [B] (verification not implemented)	588
Reduce [B] (verification not implemented)	589

Optimal result

Integrand size = 13, antiderivative size = 72

$$\int \frac{1}{x^2(a+bx^4)} dx = -\frac{1}{ax} + \frac{-2 \arctan\left(\frac{x}{\sqrt[4]{-a/b}}\right) + \log\left(\frac{\sqrt[4]{-a/b}+x}{-\sqrt[4]{-a/b}+x}\right)}{4a\sqrt[4]{-a/b}}$$

```
output -1/a/x+1/4*(-2*arctan(x/(-a/b)^(1/4))+ln(((a/b)^(1/4)+x)/(-(a/b)^(1/4)+x)))/a/(-a/b)^(1/4)
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 179 vs. 2(72) = 144.

Time = 0.02 (sec) , antiderivative size = 179, normalized size of antiderivative = 2.49

$$\int \frac{1}{x^2(a+bx^4)} dx = \frac{-8\sqrt[4]{a} + 2\sqrt{2}\sqrt[4]{b}x \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right) - 2\sqrt{2}\sqrt[4]{b}x \arctan\left(1 + \frac{\sqrt{2}\sqrt[4]{b}x}{\sqrt[4]{a}}\right) - \sqrt{2}\sqrt[4]{b}x \log\left(\sqrt{a} - \sqrt{2}\sqrt[4]{a}\sqrt[4]{b}x\right)}{8a^{5/4}x}$$

input `Integrate[1/(x^2*(a + b*x^4)),x]`

output $(-8a^{1/4} + 2\sqrt{2}b^{1/4}x\text{ArcTan}[1 - (\sqrt{2}b^{1/4}x)/a^{1/4}] - 2\sqrt{2}b^{1/4}x\text{ArcTan}[1 + (\sqrt{2}b^{1/4}x)/a^{1/4}] - \sqrt{2}b^{1/4}x\text{Log}[\sqrt{a} - \sqrt{2}a^{1/4}b^{1/4}x + \sqrt{bx^2} + \sqrt{2}b^{1/4}x\text{Log}[\sqrt{a} + \sqrt{2}a^{1/4}b^{1/4}x + \sqrt{bx^2}])/(8a^{5/4}x)$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 215 vs. $2(72) = 144$.

Time = 0.52 (sec) , antiderivative size = 215, normalized size of antiderivative = 2.99, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {847, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^2(a + bx^4)} dx \\
 & \quad \downarrow \text{847} \\
 & \frac{b \int \frac{x^2}{bx^4+a} dx}{a} - \frac{1}{ax} \\
 & \quad \downarrow \text{826} \\
 & \frac{b \left(\frac{\int \frac{\sqrt{bx^2+\sqrt{a}}}{bx^4+a} dx}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{b}} \right)}{a} - \frac{1}{ax} \\
 & \quad \downarrow \text{1476} \\
 & \frac{b \left(\frac{\int \frac{1}{x^2 - \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} + \frac{\int \frac{1}{x^2 + \frac{\sqrt{2}\sqrt[4]{ax} + \sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx^2}}{bx^4+a} dx}{2\sqrt{b}} \right)}{a} - \frac{1}{ax}
 \end{aligned}$$

$$\begin{array}{c} \downarrow 1082 \\ b \left(\frac{\int \frac{1}{\left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a}}\right)^2} d\left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{1}{\left(\frac{\sqrt{2} \sqrt[4]{bx} + 1}{\sqrt[4]{a}}\right)^2} d\left(\frac{\sqrt{2} \sqrt[4]{bx} + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx^2}}{bx^4 + a} dx}{2\sqrt{b}} \right) \\ \hline a \qquad \qquad \qquad \frac{1}{ax} \end{array}$$

$$\begin{array}{c} \downarrow 217 \\ b \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{bx} + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx^2}}{bx^4 + a} dx}{2\sqrt{b}} \right) \\ \hline a \qquad \qquad \qquad \frac{1}{ax} \end{array}$$

$$\begin{array}{c} \downarrow 1479 \\ b \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{bx} + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{bx}}{\sqrt[4]{b} \left(x^2 - \frac{\sqrt{2} \sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{bx} + \sqrt[4]{a}\right)}{\sqrt[4]{b} \left(x^2 + \frac{\sqrt{2} \sqrt[4]{ax} + \sqrt{a}}{\sqrt[4]{b}}\right)} dx}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) \\ \hline \frac{1}{ax} \qquad \qquad \qquad a \\ \downarrow 25 \end{array}$$

$$\begin{aligned}
 & b \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{bx}}{\sqrt[4]{b}\left(x^2-\frac{\sqrt{2}\sqrt[4]{ax}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{bx}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x^2+\frac{\sqrt{2}\sqrt[4]{ax}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right) \\
 & \qquad \qquad \qquad \frac{a}{ax} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & b \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{bx}}{x^2-\frac{\sqrt{2}\sqrt[4]{ax}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{bx}+\sqrt[4]{a}}{x^2+\frac{\sqrt{2}\sqrt[4]{ax}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}} dx}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right) \\
 & \qquad \qquad \qquad \frac{a}{ax} \\
 & \qquad \qquad \qquad \downarrow 1103 \\
 & b \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{bx}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{bx}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{bx}+\sqrt{a}+\sqrt{bx}^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{bx}+\sqrt{a}+\sqrt{bx}^2\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right) \\
 & \qquad \qquad \qquad \frac{a}{ax}
 \end{aligned}$$

input `Int [1/(x^2*(a + b*x^4)),x]`

output `-(1/(a*x)) - (b*((-(ArcTan[1 - (Sqrt[2]*b^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*x)/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]) - (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*x + Sqrt[b]*x^2]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*x + Sqrt[b]*x^2]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]))/a`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \text{ Int}[\text{Fx}, \text{x}], \text{x}] /; \text{FreeQ}[\text{a}, \text{x}] \ \&\& \ !\text{MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] /; \text{FreeQ}[\text{b}, \text{x}]$
- rule 217 $\text{Int}[(\text{a}_) + (\text{b}_.)*(\text{x}_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-\text{a}, 2]*\text{Rt}[-\text{b}, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Rt}[-\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{LtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$
- rule 826 $\text{Int}[(\text{x}_)^2/((\text{a}_) + (\text{b}_.)*(\text{x}_)^4), \text{x_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[\text{a}/\text{b}, 2]]\}, \text{Simp}[1/(2*\text{s}) \text{ Int}[(\text{r} + \text{s}*\text{x}^2)/(\text{a} + \text{b}*\text{x}^4), \text{x}], \text{x}] - \text{Simp}[1/(2*\text{s}) \text{ Int}[(\text{r} - \text{s}*\text{x}^2)/(\text{a} + \text{b}*\text{x}^4), \text{x}], \text{x}]] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ (\text{GtQ}[\text{a}/\text{b}, 0] \ || \ (\text{PosQ}[\text{a}/\text{b}] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{a}]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{b}]])$
- rule 847 $\text{Int}[(\text{c}_.)*(\text{x}_))^{(\text{m}_)}*((\text{a}_) + (\text{b}_.)*(\text{x}_)^{\text{n}_})^{(\text{p}_)}, \text{x_Symbol}] \rightarrow \text{Simp}[(\text{c}*\text{x})^{(\text{m} + 1)}*((\text{a} + \text{b}*\text{x}^{\text{n}})^{(\text{p} + 1)}/(\text{a}*\text{c}^{(\text{m} + 1)})), \text{x}] - \text{Simp}[\text{b}*((\text{m} + \text{n}*(\text{p} + 1) + 1)/(\text{a}*\text{c}^{\text{n}}*(\text{m} + 1))) \text{ Int}[(\text{c}*\text{x})^{(\text{m} + \text{n})}*(\text{a} + \text{b}*\text{x}^{\text{n}})^{\text{p}}, \text{x}], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \ \&\& \ \text{IGtQ}[\text{n}, 0] \ \&\& \ \text{LtQ}[\text{m}, -1] \ \&\& \ \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, \text{n}, \text{m}, \text{p}, \text{x}]$
- rule 1082 $\text{Int}[(\text{a}_) + (\text{b}_.)*(\text{x}_) + (\text{c}_.)*(\text{x}_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = 1 - 4*\text{Simplify}[\text{a}*(\text{c}/\text{b}^2)]\}, \text{Simp}[-2/\text{b} \text{ Subst}[\text{Int}[1/(\text{q} - \text{x}^2), \text{x}], \text{x}, 1 + 2*\text{c}*(\text{x}/\text{b})], \text{x}] /; \text{RationalQ}[\text{q}] \ \&\& \ (\text{EqQ}[\text{q}^2, 1] \ || \ !\text{RationalQ}[\text{b}^2 - 4*\text{a}*\text{c}]) /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$
- rule 1103 $\text{Int}[(\text{d}_) + (\text{e}_.)*(\text{x}_))/((\text{a}_.) + (\text{b}_.)*(\text{x}_) + (\text{c}_.)*(\text{x}_)^2), \text{x_Symbol}] \rightarrow \text{Simp}[\text{d}*(\text{Log}[\text{RemoveContent}[\text{a} + \text{b}*\text{x} + \text{c}*\text{x}^2, \text{x}]]/\text{b}), \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{d}, \text{e}\}, \text{x}] \ \&\& \ \text{EqQ}[2*\text{c}*\text{d} - \text{b}*\text{e}, 0]$

rule 1476

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.69

method	result	size
risch	$-\frac{1}{ax} + \frac{\sum_{R=\text{RootOf}(a^5-Z^4+b)} -R \ln((5-R^4 a^5+4b)x+R^3 a^4)}{4}$	50
default	$-\frac{1}{ax} - \frac{\sqrt{2} \left(\ln \left(\frac{x^2 - (\frac{a}{b})^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}}{x^2 + (\frac{a}{b})^{\frac{1}{4}} x \sqrt{2} + \sqrt{\frac{a}{b}}} \right) + 2 \arctan \left(\frac{\sqrt{2} x}{(\frac{a}{b})^{\frac{1}{4}}} + 1 \right) + 2 \arctan \left(\frac{\sqrt{2} x}{(\frac{a}{b})^{\frac{1}{4}}} - 1 \right) \right)}{8a(\frac{a}{b})^{\frac{1}{4}}}$	111

input

```
int(1/x^2/(b*x^4+a), x, method=_RETURNVERBOSE)
```

output

```
-1/a/x+1/4*sum(_R*ln((5*_R^4*a^5+4*b)*x+_R^3*a^4), _R=RootOf(_Z^4*a^5+b))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.78

$$\int \frac{1}{x^2(a+bx^4)} dx = \frac{ax\left(-\frac{b}{a^5}\right)^{\frac{1}{4}} \log\left(a^4\left(-\frac{b}{a^5}\right)^{\frac{3}{4}} + bx\right) - iax\left(-\frac{b}{a^5}\right)^{\frac{1}{4}} \log\left(ia^4\left(-\frac{b}{a^5}\right)^{\frac{3}{4}} + bx\right) + iax\left(-\frac{b}{a^5}\right)^{\frac{1}{4}} \log\left(-ia^4\left(-\frac{b}{a^5}\right)^{\frac{3}{4}} + bx\right)}{4ax}$$

input `integrate(1/x^2/(b*x^4+a),x, algorithm="fricas")`

output `-1/4*(a*x*(-b/a^5)^(1/4)*log(a^4*(-b/a^5)^(3/4) + b*x) - I*a*x*(-b/a^5)^(1/4)*log(I*a^4*(-b/a^5)^(3/4) + b*x) + I*a*x*(-b/a^5)^(1/4)*log(-I*a^4*(-b/a^5)^(3/4) + b*x) - a*x*(-b/a^5)^(1/4)*log(-a^4*(-b/a^5)^(3/4) + b*x) + 4)/(a*x)`

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.40

$$\int \frac{1}{x^2(a+bx^4)} dx = \text{RootSum}\left(256t^4a^5 + b, \left(t \mapsto t \log\left(-\frac{64t^3a^4}{b} + x\right)\right)\right) - \frac{1}{ax}$$

input `integrate(1/x**2/(b*x**4+a),x)`

output `RootSum(256*_t**4*a**5 + b, Lambda(_t, _t*log(-64*_t**3*a**4/b + x))) - 1/(a*x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 183 vs. $2(64) = 128$.

Time = 0.12 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.54

$$\int \frac{1}{x^2(a+bx^4)} dx =$$

$$\frac{b \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} + \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{bx} - \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} - \frac{\sqrt{2} \log(\sqrt{bx^2 + \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}x + \sqrt{a}})}{a^{\frac{1}{4}}b^{\frac{3}{4}}} + \frac{\sqrt{2} \log(\sqrt{bx^2 - \sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}x + \sqrt{a}})}{a^{\frac{1}{4}}b^{\frac{3}{4}}} \right)}{8a} - \frac{1}{ax}$$

input `integrate(1/x^2/(b*x^4+a),x, algorithm="maxima")`

output `-1/8*b*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x + sqrt(2)*a^(1/4)*b^(1/4)))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(sqrt(a)*sqrt(b))*sqrt(b)) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*sqrt(b)*x - sqrt(2)*a^(1/4)*b^(1/4)))/sqrt(sqrt(a)*sqrt(b)))/(sqrt(sqrt(a)*sqrt(b))*sqrt(b)) - sqrt(2)*log(sqrt(b)*x^2 + sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(1/4)*b^(3/4)) + sqrt(2)*log(sqrt(b)*x^2 - sqrt(2)*a^(1/4)*b^(1/4)*x + sqrt(a))/(a^(1/4)*b^(3/4))/a - 1/(a*x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(64) = 128$.

Time = 0.11 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.60

$$\int \frac{1}{x^2(a+bx^4)} dx = -\frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(2x+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{4a^2b^2} - \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(2x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{4a^2b^2} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(x^2 + \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{8a^2b^2} - \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(x^2 - \sqrt{2}x\left(\frac{a}{b}\right)^{\frac{1}{4}} + \sqrt{\frac{a}{b}}\right)}{8a^2b^2} - \frac{1}{ax}$$

input `integrate(1/x^2/(b*x^4+a),x, algorithm="giac")`

output `-1/4*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*(2*x + sqrt(2)*(a/b)^(1/4)))/(a/b)^(1/4))/(a^2*b^2) - 1/4*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*(2*x - sqrt(2)*(a/b)^(1/4))/(a/b)^(1/4))/(a^2*b^2) + 1/8*sqrt(2)*(a*b^3)^(3/4)*log(x^2 + sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a^2*b^2) - 1/8*sqrt(2)*(a*b^3)^(3/4)*log(x^2 - sqrt(2)*x*(a/b)^(1/4) + sqrt(a/b))/(a^2*b^2) - 1/(a*x)`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\int \frac{1}{x^2(a+bx^4)} dx = \frac{(-b)^{1/4} \operatorname{atanh}\left(\frac{(-b)^{1/4}x}{a^{1/4}}\right)}{2a^{5/4}} - \frac{(-b)^{1/4} \operatorname{atan}\left(\frac{(-b)^{1/4}x}{a^{1/4}}\right)}{2a^{5/4}} - \frac{1}{ax}$$

input `int(1/(x^2*(a + b*x^4)),x)`

output $((-b)^{1/4} \operatorname{atanh}((-b)^{1/4} x / a^{1/4}) / (2a^{5/4}) - (-b)^{1/4} \operatorname{atan}((-b)^{1/4} x / a^{1/4}) / (2a^{5/4}) - 1/(ax))$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 150, normalized size of antiderivative = 2.08

$$\int \frac{1}{x^2(a+bx^4)} dx$$

$$= \frac{2b^{1/4} a^{3/4} \sqrt{2} \operatorname{atan}\left(\frac{b^{1/4} a^{1/4} \sqrt{2} - 2\sqrt{b}x}{b^{1/4} a^{1/4} \sqrt{2}}\right) x - 2b^{1/4} a^{3/4} \sqrt{2} \operatorname{atan}\left(\frac{b^{1/4} a^{1/4} \sqrt{2} + 2\sqrt{b}x}{b^{1/4} a^{1/4} \sqrt{2}}\right) x - b^{1/4} a^{3/4} \sqrt{2} \log\left(-b^{1/4} a^{1/4} \sqrt{2} x + \sqrt{a} + \sqrt{b} x^2\right) x + b^{1/4} a^{3/4} \sqrt{2} \log\left(b^{1/4} a^{1/4} \sqrt{2} x + \sqrt{a} + \sqrt{b} x^2\right) x - 8a}{8a^2 x}$$

input `int(1/x^2/(b*x^4+a),x)`

output $(2*b^{1/4}*a^{3/4}*sqrt(2)*atan((b^{1/4}*a^{1/4}*sqrt(2) - 2*sqrt(b)*x)/(b^{1/4}*a^{1/4}*sqrt(2)))*x - 2*b^{1/4}*a^{3/4}*sqrt(2)*atan((b^{1/4}*a^{1/4}*sqrt(2) + 2*sqrt(b)*x)/(b^{1/4}*a^{1/4}*sqrt(2)))*x - b^{1/4}*a^{3/4}*sqrt(2)*log(-b^{1/4}*a^{1/4}*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*x + b^{1/4}*a^{3/4}*sqrt(2)*log(b^{1/4}*a^{1/4}*sqrt(2)*x + sqrt(a) + sqrt(b)*x**2)*x - 8*a)/(8*a**2*x)$

3.87 $\int \frac{1}{1+x} dx$

Optimal result	590
Mathematica [A] (verified)	590
Rubi [A] (verified)	591
Maple [A] (verified)	591
Fricas [A] (verification not implemented)	592
Sympy [A] (verification not implemented)	592
Maxima [A] (verification not implemented)	593
Giac [A] (verification not implemented)	593
Mupad [B] (verification not implemented)	593
Reduce [B] (verification not implemented)	594

Optimal result

Integrand size = 5, antiderivative size = 4

$$\int \frac{1}{1+x} dx = \log(1+x)$$

output `ln(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x} dx = \log(1+x)$$

input `Integrate[(1 + x)^(-1), x]`

output `Log[1 + x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x+1} dx$$

↓ 16

$$\log(x+1)$$

input `Int[(1 + x)^(-1), x]`

output `Log[1 + x]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$\ln(1+x)$	5
norman	$\ln(1+x)$	5
meijerg	$\ln(1+x)$	5
risch	$\ln(1+x)$	5
parallelrisch	$\ln(1+x)$	5

input `int(1/(1+x),x,method=_RETURNVERBOSE)`

output `ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x} dx = \log(x+1)$$

input `integrate(1/(1+x),x, algorithm="fricas")`

output `log(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.75

$$\int \frac{1}{1+x} dx = \log(x+1)$$

input `integrate(1/(1+x),x)`

output `log(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x} dx = \log(x+1)$$

input `integrate(1/(1+x),x, algorithm="maxima")`

output `log(x + 1)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

$$\int \frac{1}{1+x} dx = \log(|x+1|)$$

input `integrate(1/(1+x),x, algorithm="giac")`

output `log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x} dx = \ln(x+1)$$

input `int(1/(x + 1),x)`

output `log(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x} dx = \log(x+1)$$

input `int(1/(1+x),x)`

output `log(x + 1)`

3.88 $\int \frac{1}{1+x^2} dx$

Optimal result	595
Mathematica [A] (verified)	595
Rubi [A] (verified)	596
Maple [A] (verified)	596
Fricas [A] (verification not implemented)	597
Sympy [A] (verification not implemented)	597
Maxima [A] (verification not implemented)	598
Giac [A] (verification not implemented)	598
Mupad [B] (verification not implemented)	598
Reduce [B] (verification not implemented)	599

Optimal result

Integrand size = 7, antiderivative size = 2

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

output `arctan(x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `Integrate[(1 + x^2)^(-1),x]`

output `ArcTan[x]`

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 + 1} dx$$

↓ 216

$$\arctan(x)$$

input `Int[(1 + x^2)^(-1), x]`

output `ArcTan[x]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Maple [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\arctan(x)$	3
meijerg	$\arctan(x)$	3
risch	$\arctan(x)$	3
parallelrisch	$\frac{i \ln(x+i)}{2} - \frac{i \ln(x-i)}{2}$	18

input `int(1/(x^2+1),x,method=_RETURNVERBOSE)`

output `arctan(x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `integrate(1/(x^2+1),x, algorithm="fricas")`

output `arctan(x)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \operatorname{atan}(x)$$

input `integrate(1/(x**2+1),x)`

output `atan(x)`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `integrate(1/(x^2+1),x, algorithm="maxima")`

output `arctan(x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \arctan(x)$$

input `integrate(1/(x^2+1),x, algorithm="giac")`

output `arctan(x)`

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \operatorname{atan}(x)$$

input `int(1/(x^2 + 1),x)`

output `atan(x)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1+x^2} dx = \text{atan}(x)$$

input `int(1/(x^2+1),x)`

output `atan(x)`

3.89 $\int \frac{1}{1+x^3} dx$

Optimal result	600
Mathematica [A] (verified)	600
Rubi [A] (verified)	601
Maple [A] (verified)	603
Fricas [A] (verification not implemented)	603
Sympy [A] (verification not implemented)	604
Maxima [A] (verification not implemented)	604
Giac [A] (verification not implemented)	604
Mupad [B] (verification not implemented)	605
Reduce [B] (verification not implemented)	605

Optimal result

Integrand size = 7, antiderivative size = 43

$$\int \frac{1}{1+x^3} dx = \frac{\arctan\left(\frac{\sqrt{3}x}{2-x}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(\frac{1+x}{\sqrt{1-x+x^2}}\right)$$

output `1/3*arctan(3^(1/2)*x/(2-x))*3^(1/2)+1/3*ln((1+x)/(x^2-x+1)^(1/2))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

$$\int \frac{1}{1+x^3} dx = \frac{\arctan\left(\frac{-1+2x}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3} \log(1+x) - \frac{1}{6} \log(1-x+x^2)$$

input `Integrate[(1 + x^3)^(-1), x]`

output `ArcTan[(-1 + 2*x)/Sqrt[3]]/Sqrt[3] + Log[1 + x]/3 - Log[1 - x + x^2]/6`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.05, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {750, 16, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 + 1} dx \\
 & \quad \downarrow \text{750} \\
 & \frac{1}{3} \int \frac{2-x}{x^2-x+1} dx + \frac{1}{3} \int \frac{1}{x+1} dx \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{3} \int \frac{2-x}{x^2-x+1} dx + \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^2-x+1} dx - \frac{1}{2} \int -\frac{1-2x}{x^2-x+1} dx \right) + \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^2-x+1} dx + \frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx \right) + \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx - 3 \int \frac{1}{-(2x-1)^2-3} d(2x-1) \right) + \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx + \sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) \right) + \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) - \frac{1}{2} \log(x^2-x+1) \right) + \frac{1}{3} \log(x+1)
 \end{aligned}$$

input

```
Int[(1 + x^3)^(-1), x]
```

output $\text{Log}[1 + x]/3 + (\text{Sqrt}[3]*\text{ArcTan}[(-1 + 2*x)/\text{Sqrt}[3]] - \text{Log}[1 - x + x^2]/2)/3$

Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{Int}[Fx, x], x]$

rule 217 $\text{Int}(((a_) + (b_)*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

rule 750 $\text{Int}(((a_) + (b_)*(x_)^3)^{-1}), x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$

rule 1083 $\text{Int}(((a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[-2 \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}(((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

rule 1142 $\text{Int}(((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

method	result	size
risch	$\frac{\ln(1+x)}{3} - \frac{\ln(x^2-x+1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{2(-\frac{1}{2}+x)\sqrt{3}}{3}\right)}{3}$	33
default	$\frac{\ln(1+x)}{3} - \frac{\ln(x^2-x+1)}{6} + \frac{\arctan\left(\frac{(-1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	35
meijerg	$\frac{x \ln\left(1+(x^3)^{\frac{1}{3}}\right)}{3(x^3)^{\frac{1}{3}}} - \frac{x \ln\left(1-(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{6(x^3)^{\frac{1}{3}}} + \frac{x\sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2-(x^3)^{\frac{1}{3}}}\right)}{3(x^3)^{\frac{1}{3}}}$	74

input `int(1/(x^3+1),x,method=_RETURNVERBOSE)`output `1/3*ln(1+x)-1/6*ln(x^2-x+1)+1/3*3^(1/2)*arctan(2/3*(-1/2+x)*3^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \frac{1}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) - \frac{1}{6} \log(x^2-x+1) + \frac{1}{3} \log(x+1)$$

input `integrate(1/(x^3+1),x, algorithm="fricas")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) - 1/6*log(x^2 - x + 1) + 1/3*log(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \frac{1}{1+x^3} dx = \frac{\log(x+1)}{3} - \frac{\log(x^2-x+1)}{6} + \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} - \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(1/(x**3+1),x)`output `log(x + 1)/3 - log(x**2 - x + 1)/6 + sqrt(3)*atan(2*sqrt(3)*x/3 - sqrt(3)/3)/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \frac{1}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) - \frac{1}{6} \log(x^2-x+1) + \frac{1}{3} \log(x+1)$$

input `integrate(1/(x^3+1),x, algorithm="maxima")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) - 1/6*log(x^2 - x + 1) + 1/3*log(x + 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.81

$$\int \frac{1}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) - \frac{1}{6} \log(x^2-x+1) + \frac{1}{3} \log(|x+1|)$$

input `integrate(1/(x^3+1),x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) - 1/6*log(x^2 - x + 1) + 1/3*log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.72

$$\int \frac{1}{1+x^3} dx = \frac{\ln(x+1)}{3} - \frac{\ln\left(\left(x-\frac{1}{2}\right)^2 + \frac{3}{4}\right)}{6} + \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}\left(x-\frac{1}{2}\right)}{3}\right)}{3}$$

input `int(1/(x^3 + 1),x)`output `log(x + 1)/3 - log((x - 1/2)^2 + 3/4)/6 + (3^(1/2)*atan((2*3^(1/2)*(x - 1/2))/3))/3`**Reduce [B] (verification not implemented)**

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

$$\int \frac{1}{1+x^3} dx = \frac{\sqrt{3} \operatorname{atan}\left(\frac{2x-1}{\sqrt{3}}\right)}{3} - \frac{\log(x^2 - x + 1)}{6} + \frac{\log(x + 1)}{3}$$

input `int(1/(x^3+1),x)`output `(2*sqrt(3)*atan((2*x - 1)/sqrt(3)) - log(x**2 - x + 1) + 2*log(x + 1))/6`

3.90 $\int \frac{1}{1+x^4} dx$

Optimal result	606
Mathematica [A] (verified)	606
Rubi [A] (verified)	607
Maple [C] (verified)	609
Fricas [A] (verification not implemented)	610
Sympy [A] (verification not implemented)	610
Maxima [A] (verification not implemented)	611
Giac [A] (verification not implemented)	611
Mupad [B] (verification not implemented)	612
Reduce [B] (verification not implemented)	612

Optimal result

Integrand size = 7, antiderivative size = 65

$$\int \frac{1}{1+x^4} dx = \frac{\arctan\left(\frac{\sqrt{2}x}{1-x^2}\right)}{2\sqrt{2}} + \frac{\log\left(\frac{1+\sqrt{2}x+x^2}{1-\sqrt{2}x+x^2}\right)}{4\sqrt{2}}$$

output $1/4*\arctan(2^{(1/2)*x}/(-x^2+1))*2^{(1/2)}+1/8*\ln((1+x*2^{(1/2)}+x^2)/(1-x*2^{(1/2)}+x^2))*2^{(1/2)}$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.98

$$\int \frac{1}{1+x^4} dx = \frac{-2 \arctan(1 - \sqrt{2}x) + 2 \arctan(1 + \sqrt{2}x) - \log(1 - \sqrt{2}x + x^2) + \log(1 + \sqrt{2}x + x^2)}{4\sqrt{2}}$$

input `Integrate[(1 + x^4)^(-1), x]`

output $(-2*\text{ArcTan}[1 - \text{Sqrt}[2]*x] + 2*\text{ArcTan}[1 + \text{Sqrt}[2]*x] - \text{Log}[1 - \text{Sqrt}[2]*x + x^2] + \text{Log}[1 + \text{Sqrt}[2]*x + x^2])/(4*\text{Sqrt}[2])$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.38, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 1.143$, Rules used = {755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^4 + 1} dx \\
 & \quad \downarrow \text{755} \\
 & \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx + \frac{1}{2} \int \frac{x^2 + 1}{x^4 + 1} dx \\
 & \quad \downarrow \text{1476} \\
 & \frac{1}{2} \left(\frac{1}{2} \int \frac{1}{x^2 - \sqrt{2}x + 1} dx + \frac{1}{2} \int \frac{1}{x^2 + \sqrt{2}x + 1} dx \right) + \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx \\
 & \quad \downarrow \text{1082} \\
 & \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx + \frac{1}{2} \left(\frac{\int \frac{1}{-(1 - \sqrt{2}x)^2 - 1} d(1 - \sqrt{2}x)}{\sqrt{2}} - \frac{\int \frac{1}{-(\sqrt{2}x + 1)^2 - 1} d(\sqrt{2}x + 1)}{\sqrt{2}} \right) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{2} \int \frac{1 - x^2}{x^4 + 1} dx + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}x)}{\sqrt{2}} \right) \\
 & \quad \downarrow \text{1479} \\
 & \frac{1}{2} \left(-\frac{\int -\frac{\sqrt{2} - 2x}{x^2 - \sqrt{2}x + 1} dx}{2\sqrt{2}} - \frac{\int -\frac{\sqrt{2}(\sqrt{2}x + 1)}{x^2 + \sqrt{2}x + 1} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}x)}{\sqrt{2}} \right) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{2} \left(\frac{\int \frac{\sqrt{2} - 2x}{x^2 - \sqrt{2}x + 1} dx}{2\sqrt{2}} + \frac{\int \frac{\sqrt{2}(\sqrt{2}x + 1)}{x^2 + \sqrt{2}x + 1} dx}{2\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x + 1)}{\sqrt{2}} - \frac{\arctan(1 - \sqrt{2}x)}{\sqrt{2}} \right) \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\frac{1}{2} \left(\int \frac{\sqrt{2}-2x}{x^2-\sqrt{2}x+1} dx + \frac{1}{2} \int \frac{\sqrt{2}x+1}{x^2+\sqrt{2}x+1} dx \right) + \frac{1}{2} \left(\frac{\arctan(\sqrt{2}x+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}x)}{\sqrt{2}} \right)$$

↓ 1103

$$\frac{1}{2} \left(\frac{\arctan(\sqrt{2}x+1)}{\sqrt{2}} - \frac{\arctan(1-\sqrt{2}x)}{\sqrt{2}} \right) + \frac{1}{2} \left(\frac{\log(x^2+\sqrt{2}x+1)}{2\sqrt{2}} - \frac{\log(x^2-\sqrt{2}x+1)}{2\sqrt{2}} \right)$$

input `Int[(1 + x^4)^(-1), x]`

output `(-(ArcTan[1 - Sqrt[2]*x]/Sqrt[2]) + ArcTan[1 + Sqrt[2]*x]/Sqrt[2])/2 + (-1/2*Log[1 - Sqrt[2]*x + x^2]/Sqrt[2] + Log[1 + Sqrt[2]*x + x^2]/(2*Sqrt[2]))/2`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.34

method	result
risch	$\frac{\left(\sum_{R=\text{RootOf}(_Z^4+1)} \frac{\ln(x-R)}{-R^3} \right)}{4}$
default	$\frac{\sqrt{2} \left(\ln\left(\frac{1+\sqrt{2}x+x^2}{1-\sqrt{2}x+x^2}\right) + 2 \arctan(\sqrt{2}x+1) + 2 \arctan(\sqrt{2}x-1) \right)}{8}$
meijerg	$-\frac{x\sqrt{2} \ln\left(1-\sqrt{2}(x^4)^{\frac{1}{4}}+\sqrt{x^4}\right)}{8(x^4)^{\frac{1}{4}}} + \frac{x\sqrt{2} \arctan\left(\frac{\sqrt{2}(x^4)^{\frac{1}{4}}}{2-\sqrt{2}(x^4)^{\frac{1}{4}}}\right)}{4(x^4)^{\frac{1}{4}}} + \frac{x\sqrt{2} \ln\left(1+\sqrt{2}(x^4)^{\frac{1}{4}}+\sqrt{x^4}\right)}{8(x^4)^{\frac{1}{4}}} + \frac{x\sqrt{2} \arctan\left(\frac{\sqrt{2}(x^4)^{\frac{1}{4}}}{2+\sqrt{2}(x^4)^{\frac{1}{4}}}\right)}{4(x^4)^{\frac{1}{4}}}$

input `int(1/(x^4+1),x,method=_RETURNVERBOSE)`

output `1/4*sum(1/_R^3*ln(x-_R),_R=RootOf(_Z^4+1))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.92

$$\int \frac{1}{1+x^4} dx = \frac{1}{4} \sqrt{2} \arctan(\sqrt{2}x+1) + \frac{1}{4} \sqrt{2} \arctan(\sqrt{2}x-1) \\ + \frac{1}{8} \sqrt{2} \log(x^2 + \sqrt{2}x + 1) - \frac{1}{8} \sqrt{2} \log(x^2 - \sqrt{2}x + 1)$$

input `integrate(1/(x^4+1),x, algorithm="fricas")`

output `1/4*sqrt(2)*arctan(sqrt(2)*x + 1) + 1/4*sqrt(2)*arctan(sqrt(2)*x - 1) + 1/8*sqrt(2)*log(x^2 + sqrt(2)*x + 1) - 1/8*sqrt(2)*log(x^2 - sqrt(2)*x + 1)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.12

$$\int \frac{1}{1+x^4} dx = -\frac{\sqrt{2} \log(x^2 - \sqrt{2}x + 1)}{8} + \frac{\sqrt{2} \log(x^2 + \sqrt{2}x + 1)}{8} \\ + \frac{\sqrt{2} \operatorname{atan}(\sqrt{2}x - 1)}{4} + \frac{\sqrt{2} \operatorname{atan}(\sqrt{2}x + 1)}{4}$$

input `integrate(1/(x**4+1),x)`

output `-sqrt(2)*log(x**2 - sqrt(2)*x + 1)/8 + sqrt(2)*log(x**2 + sqrt(2)*x + 1)/8 + sqrt(2)*atan(sqrt(2)*x - 1)/4 + sqrt(2)*atan(sqrt(2)*x + 1)/4`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11

$$\int \frac{1}{1+x^4} dx = \frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (2x + \sqrt{2}) \right) + \frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (2x - \sqrt{2}) \right) \\ + \frac{1}{8} \sqrt{2} \log (x^2 + \sqrt{2}x + 1) - \frac{1}{8} \sqrt{2} \log (x^2 - \sqrt{2}x + 1)$$

input `integrate(1/(x^4+1),x, algorithm="maxima")`output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x + sqrt(2))) + 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x - sqrt(2))) + 1/8*sqrt(2)*log(x^2 + sqrt(2)*x + 1) - 1/8*sqrt(2)*log(x^2 - sqrt(2)*x + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11

$$\int \frac{1}{1+x^4} dx = \frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (2x + \sqrt{2}) \right) + \frac{1}{4} \sqrt{2} \arctan \left(\frac{1}{2} \sqrt{2} (2x - \sqrt{2}) \right) \\ + \frac{1}{8} \sqrt{2} \log (x^2 + \sqrt{2}x + 1) - \frac{1}{8} \sqrt{2} \log (x^2 - \sqrt{2}x + 1)$$

input `integrate(1/(x^4+1),x, algorithm="giac")`output `1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x + sqrt(2))) + 1/4*sqrt(2)*arctan(1/2*sqrt(2)*(2*x - sqrt(2))) + 1/8*sqrt(2)*log(x^2 + sqrt(2)*x + 1) - 1/8*sqrt(2)*log(x^2 - sqrt(2)*x + 1)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.51

$$\int \frac{1}{1+x^4} dx = \sqrt{2} \operatorname{atan}\left(\sqrt{2}x\left(\frac{1}{2} - \frac{1}{2}i\right)\right) \left(\frac{1}{4} + \frac{1}{4}i\right) + \sqrt{2} \operatorname{atan}\left(\sqrt{2}x\left(\frac{1}{2} + \frac{1}{2}i\right)\right) \left(\frac{1}{4} - \frac{1}{4}i\right)$$

input `int(1/(x^4 + 1),x)`output `2^(1/2)*atan(2^(1/2)*x*(1/2 - 1i/2))*(1/4 + 1i/4) + 2^(1/2)*atan(2^(1/2)*x*(1/2 + 1i/2))*(1/4 - 1i/4)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.86

$$\int \frac{1}{1+x^4} dx = \frac{\sqrt{2} \left(-2 \operatorname{atan}\left(\frac{\sqrt{2}-2x}{\sqrt{2}}\right) + 2 \operatorname{atan}\left(\frac{\sqrt{2}+2x}{\sqrt{2}}\right) - \log(-\sqrt{2}x + x^2 + 1) + \log(\sqrt{2}x + x^2 + 1) \right)}{8}$$

input `int(1/(x^4+1),x)`output `(sqrt(2)*(- 2*atan((sqrt(2) - 2*x)/sqrt(2)) + 2*atan((sqrt(2) + 2*x)/sqrt(2)) - log(- sqrt(2)*x + x**2 + 1) + log(sqrt(2)*x + x**2 + 1)))/8`

3.91 $\int \frac{1}{1-x} dx$

Optimal result	613
Mathematica [A] (verified)	613
Rubi [A] (verified)	614
Maple [A] (verified)	614
Fricas [A] (verification not implemented)	615
Sympy [A] (verification not implemented)	615
Maxima [A] (verification not implemented)	616
Giac [A] (verification not implemented)	616
Mupad [B] (verification not implemented)	616
Reduce [B] (verification not implemented)	617

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \frac{1}{1-x} dx = -\log(1-x)$$

output `-ln(1-x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-x} dx = -\log(1-x)$$

input `Integrate[(1 - x)^(-1),x]`

output `-Log[1 - x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1-x} dx$$

↓ 16

$$-\log(1-x)$$

input `Int[(1 - x)^(-1), x]`

output `-Log[1 - x]`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
norman	$-\ln(x-1)$	7
risch	$-\ln(x-1)$	7
parallelrisch	$-\ln(x-1)$	7
default	$-\ln(1-x)$	9
meijerg	$-\ln(1-x)$	9

input `int(1/(1-x),x,method=_RETURNVERBOSE)`

output `-ln(x-1)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{1-x} dx = -\log(x-1)$$

input `integrate(1/(1-x),x, algorithm="fricas")`

output `-log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{1}{1-x} dx = -\log(x-1)$$

input `integrate(1/(1-x),x)`

output `-log(x - 1)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{1-x} dx = -\log(x-1)$$

input `integrate(1/(1-x),x, algorithm="maxima")`

output `-log(x - 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

$$\int \frac{1}{1-x} dx = -\log(|x-1|)$$

input `integrate(1/(1-x),x, algorithm="giac")`

output `-log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{1-x} dx = -\ln(x-1)$$

input `int(-1/(x - 1),x)`

output `-log(x - 1)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{1}{1-x} dx = -\log(x-1)$$

input `int(1/(1-x),x)`

output `- log(x - 1)`

3.92 $\int \frac{1}{1-x^2} dx$

Optimal result	618
Mathematica [B] (verified)	618
Rubi [A] (verified)	619
Maple [A] (verified)	620
Fricas [B] (verification not implemented)	620
Sympy [B] (verification not implemented)	621
Maxima [B] (verification not implemented)	621
Giac [B] (verification not implemented)	621
Mupad [B] (verification not implemented)	622
Reduce [B] (verification not implemented)	622

Optimal result

Integrand size = 9, antiderivative size = 2

$$\int \frac{1}{1-x^2} dx = \operatorname{arctanh}(x)$$

output `arctanh(x)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(2) = 4$.

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 9.50

$$\int \frac{1}{1-x^2} dx = -\frac{1}{2} \log(1-x) + \frac{1}{2} \log(1+x)$$

input `Integrate[(1 - x^2)^(-1), x]`

output `-1/2*Log[1 - x] + Log[1 + x]/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{1-x^2} dx$$

↓ 219

$$\operatorname{arctanh}(x)$$

input `Int[(1 - x^2)^(-1), x]`

output `ArcTanh[x]`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Maple [A] (verified)

Time = 0.00 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
default	$\operatorname{arctanh}(x)$	3
meijerg	$\operatorname{arctanh}(x)$	3
norman	$-\frac{\ln(x-1)}{2} + \frac{\ln(1+x)}{2}$	14
risch	$-\frac{\ln(x-1)}{2} + \frac{\ln(1+x)}{2}$	14
parallelrisc	$-\frac{\ln(x-1)}{2} + \frac{\ln(1+x)}{2}$	14

input `int(1/(-x^2+1),x,method=_RETURNVERBOSE)`

output `arctanh(x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(2) = 4$.

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(x+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^2+1),x, algorithm="fricas")`

output `1/2*log(x + 1) - 1/2*log(x - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(2) = 4$.

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 6.00

$$\int \frac{1}{1-x^2} dx = -\frac{\log(x-1)}{2} + \frac{\log(x+1)}{2}$$

input `integrate(1/(-x**2+1),x)`

output `-log(x - 1)/2 + log(x + 1)/2`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(2) = 4$.

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(x+1) - \frac{1}{2} \log(x-1)$$

input `integrate(1/(-x^2+1),x, algorithm="maxima")`

output `1/2*log(x + 1) - 1/2*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(2) = 4$.

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 7.50

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \log(|x+1|) - \frac{1}{2} \log(|x-1|)$$

input `integrate(1/(-x^2+1),x, algorithm="giac")`

output $1/2*\log(\text{abs}(x + 1)) - 1/2*\log(\text{abs}(x - 1))$

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-x^2} dx = \text{atanh}(x)$$

input `int(-1/(x^2 - 1),x)`

output `atanh(x)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 13, normalized size of antiderivative = 6.50

$$\int \frac{1}{1-x^2} dx = -\frac{\log(x-1)}{2} + \frac{\log(x+1)}{2}$$

input `int(1/(-x^2+1),x)`

output `(- log(x - 1) + log(x + 1))/2`

3.93 $\int \frac{1}{-1+x^2} dx$

Optimal result	623
Mathematica [B] (verified)	623
Rubi [A] (verified)	624
Maple [A] (verified)	625
Fricas [B] (verification not implemented)	625
Sympy [B] (verification not implemented)	626
Maxima [B] (verification not implemented)	626
Giac [B] (verification not implemented)	626
Mupad [B] (verification not implemented)	627
Reduce [B] (verification not implemented)	627

Optimal result

Integrand size = 7, antiderivative size = 4

$$\int \frac{1}{-1+x^2} dx = -\operatorname{coth}^{-1}(x)$$

output `-arccoth(x)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 19 vs. $2(4) = 8$.

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 4.75

$$\int \frac{1}{-1+x^2} dx = \frac{1}{2} \log(1-x) - \frac{1}{2} \log(1+x)$$

input `Integrate[(-1 + x^2)^(-1), x]`

output `Log[1 - x]/2 - Log[1 + x]/2`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 - 1} dx$$

↓ 220

$$-\operatorname{arctanh}(x)$$

input `Int[(-1 + x^2)^(-1), x]`

output `-ArcTanh[x]`

Defintions of rubi rules used

rule 220 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.25

method	result	size
default	$-\operatorname{arctanh}(x)$	5
meijerg	$-\operatorname{arctanh}(x)$	5
norman	$\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}$	14
risch	$\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}$	14
parallelrisch	$\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}$	14

input `int(1/(x^2-1),x,method=_RETURNVERBOSE)`

output `-arctanh(x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(4) = 8$.

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 3.25

$$\int \frac{1}{-1+x^2} dx = -\frac{1}{2} \log(x+1) + \frac{1}{2} \log(x-1)$$

input `integrate(1/(x^2-1),x, algorithm="fricas")`

output `-1/2*log(x + 1) + 1/2*log(x - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12 vs. $2(3) = 6$.

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 3.00

$$\int \frac{1}{-1+x^2} dx = \frac{\log(x-1)}{2} - \frac{\log(x+1)}{2}$$

input `integrate(1/(x**2-1),x)`

output `log(x - 1)/2 - log(x + 1)/2`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. $2(4) = 8$.

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 3.25

$$\int \frac{1}{-1+x^2} dx = -\frac{1}{2} \log(x+1) + \frac{1}{2} \log(x-1)$$

input `integrate(1/(x^2-1),x, algorithm="maxima")`

output `-1/2*log(x + 1) + 1/2*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 15 vs. $2(4) = 8$.

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 3.75

$$\int \frac{1}{-1+x^2} dx = -\frac{1}{2} \log(|x+1|) + \frac{1}{2} \log(|x-1|)$$

input `integrate(1/(x^2-1),x, algorithm="giac")`

output $-1/2*\log(\text{abs}(x + 1)) + 1/2*\log(\text{abs}(x - 1))$

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 4, normalized size of antiderivative = 1.00

$$\int \frac{1}{-1 + x^2} dx = -\text{atanh}(x)$$

input $\text{int}(1/(x^2 - 1), x)$

output $-\text{atanh}(x)$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 3.25

$$\int \frac{1}{-1 + x^2} dx = \frac{\log(x - 1)}{2} - \frac{\log(x + 1)}{2}$$

input $\text{int}(1/(x^2-1), x)$

output $(\log(x - 1) - \log(x + 1))/2$

3.94 $\int \frac{1}{1-x^3} dx$

Optimal result	628
Mathematica [A] (verified)	628
Rubi [A] (verified)	629
Maple [A] (verified)	631
Fricas [A] (verification not implemented)	631
Sympy [A] (verification not implemented)	632
Maxima [A] (verification not implemented)	632
Giac [A] (verification not implemented)	632
Mupad [B] (verification not implemented)	633
Reduce [B] (verification not implemented)	633

Optimal result

Integrand size = 9, antiderivative size = 43

$$\int \frac{1}{1-x^3} dx = \frac{\arctan\left(\frac{\sqrt{3}x}{2+x}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(\frac{\sqrt{1+x+x^2}}{1-x}\right)$$

output `1/3*arctan(3^(1/2)*x/(2+x))*3^(1/2)+1/3*ln((x^2+x+1)^(1/2)/(1-x))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.93

$$\int \frac{1}{1-x^3} dx = \frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3} \log(1-x) + \frac{1}{6} \log(1+x+x^2)$$

input `Integrate[(1 - x^3)^(-1),x]`

output `ArcTan[(1 + 2*x)/Sqrt[3]]/Sqrt[3] - Log[1 - x]/3 + Log[1 + x + x^2]/6`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.05, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {750, 16, 1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{1-x^3} dx \\
 & \quad \downarrow \text{750} \\
 & \frac{1}{3} \int \frac{x+2}{x^2+x+1} dx + \frac{1}{3} \int \frac{1}{1-x} dx \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{3} \int \frac{x+2}{x^2+x+1} dx - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^2+x+1} dx + \frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx \right) - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx - 3 \int \frac{1}{-(2x+1)^2-3} d(2x+1) \right) - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx + \sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) \right) - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) + \frac{1}{2} \log(x^2+x+1) \right) - \frac{1}{3} \log(1-x)
 \end{aligned}$$

input `Int[(1 - x^3)^(-1), x]`

output `-1/3*Log[1 - x] + (Sqrt[3]*ArcTan[(1 + 2*x)/Sqrt[3]] + Log[1 + x + x^2]/2)/3`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$
- rule 750 $\text{Int}[(a_)+(b_)*(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1083 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.72

method	result	size
risch	$-\frac{\ln(x-1)}{3} + \frac{\ln(x^2+x+1)}{6} + \frac{\sqrt{3} \arctan\left(\frac{2\left(\frac{1}{2}+x\right)\sqrt{3}}{3}\right)}{3}$	31
default	$\frac{\ln(x^2+x+1)}{6} + \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} - \frac{\ln(x-1)}{3}$	33
meijerg	$-\frac{x \left(\ln\left(1-(x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1+(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2+(x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{1}{3}}}$	62

input `int(1/(-x^3+1),x,method=_RETURNVERBOSE)`output `-1/3*ln(x-1)+1/6*ln(x^2+x+1)+1/3*3^(1/2)*arctan(2/3*(1/2+x)*3^(1/2))`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

$$\int \frac{1}{1-x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{6} \log(x^2+x+1) - \frac{1}{3} \log(x-1)$$

input `integrate(1/(-x^3+1),x, algorithm="fricas")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/6*log(x^2 + x + 1) - 1/3*log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \frac{1}{1-x^3} dx = -\frac{\log(x-1)}{3} + \frac{\log(x^2+x+1)}{6} + \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(1/(-x**3+1),x)`output `-log(x - 1)/3 + log(x**2 + x + 1)/6 + sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

$$\int \frac{1}{1-x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{6} \log(x^2+x+1) - \frac{1}{3} \log(x-1)$$

input `integrate(1/(-x^3+1),x, algorithm="maxima")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/6*log(x^2 + x + 1) - 1/3*log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

$$\int \frac{1}{1-x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{6} \log(x^2+x+1) - \frac{1}{3} \log(|x-1|)$$

input `integrate(1/(-x^3+1),x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/6*log(x^2 + x + 1) - 1/3*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.07

$$\int \frac{1}{1-x^3} dx = -\frac{\ln(x-1)}{3} - \ln\left(x + \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right) \\ + \ln\left(x + \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)$$

input `int(-1/(x^3 - 1),x)`output `log(x + (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 + 1/6) - log(x - (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 - 1/6) - log(x - 1)/3`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.72

$$\int \frac{1}{1-x^3} dx = \frac{\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right)}{3} + \frac{\log(x^2 + x + 1)}{6} - \frac{\log(x-1)}{3}$$

input `int(1/(-x^3+1),x)`output `(2*sqrt(3)*atan((2*x + 1)/sqrt(3)) + log(x**2 + x + 1) - 2*log(x - 1))/6`

3.95 $\int \frac{1}{1-x^4} dx$

Optimal result	634
Mathematica [B] (verified)	634
Rubi [A] (verified)	635
Maple [A] (verified)	636
Fricas [A] (verification not implemented)	636
Sympy [B] (verification not implemented)	637
Maxima [A] (verification not implemented)	637
Giac [B] (verification not implemented)	637
Mupad [B] (verification not implemented)	638
Reduce [B] (verification not implemented)	638

Optimal result

Integrand size = 9, antiderivative size = 9

$$\int \frac{1}{1-x^4} dx = \frac{1}{2}(\arctan(x) + \operatorname{arctanh}(x))$$

output `1/2*arctan(x)+1/2*arctanh(x)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. $2(9) = 18$.

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.78

$$\int \frac{1}{1-x^4} dx = \frac{\arctan(x)}{2} - \frac{1}{4} \log(1-x) + \frac{1}{4} \log(1+x)$$

input `Integrate[(1 - x^4)^(-1),x]`

output `ArcTan[x]/2 - Log[1 - x]/4 + Log[1 + x]/4`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.44, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{1-x^4} dx \\ & \quad \downarrow \text{756} \\ & \frac{1}{2} \int \frac{1}{1-x^2} dx + \frac{1}{2} \int \frac{1}{x^2+1} dx \\ & \quad \downarrow \text{216} \\ & \frac{1}{2} \int \frac{1}{1-x^2} dx + \frac{\arctan(x)}{2} \\ & \quad \downarrow \text{219} \\ & \frac{\arctan(x)}{2} + \frac{\operatorname{arctanh}(x)}{2} \end{aligned}$$

input `Int[(1 - x^4)^(-1), x]`

output `ArcTan[x]/2 + ArcTanh[x]/2`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 756

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x]
+ Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.11

method	result	size
default	$\frac{\arctan(x)}{2} + \frac{\operatorname{arctanh}(x)}{2}$	10
risch	$\frac{\arctan(x)}{2} + \frac{\ln(1+x)}{4} - \frac{\ln(x-1)}{4}$	18
parallelrisch	$\frac{i \ln(x+i)}{4} - \frac{i \ln(x-i)}{4} + \frac{\ln(1+x)}{4} - \frac{\ln(x-1)}{4}$	30
meijerg	$-\frac{x \left(\ln \left(1 - (x^4)^{\frac{1}{4}} \right) - \ln \left(1 + (x^4)^{\frac{1}{4}} \right) - 2 \arctan \left((x^4)^{\frac{1}{4}} \right) \right)}{4(x^4)^{\frac{1}{4}}}$	38

input `int(1/(-x^4+1),x,method=_RETURNVERBOSE)`

output `1/2*arctan(x)+1/2*arctanh(x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int \frac{1}{1-x^4} dx = \frac{1}{2} \arctan(x) + \frac{1}{4} \log(x+1) - \frac{1}{4} \log(x-1)$$

input `integrate(1/(-x^4+1),x, algorithm="fricas")`

output `1/2*arctan(x) + 1/4*log(x + 1) - 1/4*log(x - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(8) = 16$.

Time = 0.08 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int \frac{1}{1-x^4} dx = -\frac{\log(x-1)}{4} + \frac{\log(x+1)}{4} + \frac{\operatorname{atan}(x)}{2}$$

input `integrate(1/(-x**4+1),x)`

output `-log(x - 1)/4 + log(x + 1)/4 + atan(x)/2`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int \frac{1}{1-x^4} dx = \frac{1}{2} \arctan(x) + \frac{1}{4} \log(x+1) - \frac{1}{4} \log(x-1)$$

input `integrate(1/(-x^4+1),x, algorithm="maxima")`

output `1/2*arctan(x) + 1/4*log(x + 1) - 1/4*log(x - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. $2(9) = 18$.

Time = 0.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 2.11

$$\int \frac{1}{1-x^4} dx = \frac{1}{2} \arctan(x) + \frac{1}{4} \log(|x+1|) - \frac{1}{4} \log(|x-1|)$$

input `integrate(1/(-x^4+1),x, algorithm="giac")`

output `1/2*arctan(x) + 1/4*log(abs(x + 1)) - 1/4*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.00

$$\int \frac{1}{1-x^4} dx = \frac{\operatorname{atan}(x)}{2} + \frac{\operatorname{atanh}(x)}{2}$$

input `int(-1/(x^4 - 1),x)`

output `atan(x)/2 + atanh(x)/2`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.89

$$\int \frac{1}{1-x^4} dx = \frac{\operatorname{atan}(x)}{2} - \frac{\log(x-1)}{4} + \frac{\log(x+1)}{4}$$

input `int(1/(-x^4+1),x)`

output `(2*atan(x) - log(x - 1) + log(x + 1))/4`

3.96 $\int \frac{x}{1+x} dx$

Optimal result	639
Mathematica [A] (verified)	639
Rubi [A] (verified)	640
Maple [A] (verified)	641
Fricas [A] (verification not implemented)	641
Sympy [A] (verification not implemented)	641
Maxima [A] (verification not implemented)	642
Giac [A] (verification not implemented)	642
Mupad [B] (verification not implemented)	642
Reduce [B] (verification not implemented)	643

Optimal result

Integrand size = 7, antiderivative size = 8

$$\int \frac{x}{1+x} dx = x - \log(1+x)$$

output `x-ln(1+x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x} dx = x - \log(1+x)$$

input `Integrate[x/(1 + x), x]`

output `x - Log[1 + x]`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x+1} dx$$

$$\downarrow 49$$

$$\int \left(\frac{1}{-x-1} + 1 \right) dx$$

$$\downarrow 2009$$

$$x - \log(x+1)$$

input

```
Int[x/(1 + x),x]
```

output

```
x - Log[1 + x]
```

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

method	result	size
default	$x - \ln(1 + x)$	9
norman	$x - \ln(1 + x)$	9
meijerg	$x - \ln(1 + x)$	9
risch	$x - \ln(1 + x)$	9
parallelsch	$x - \ln(1 + x)$	9

input `int(x/(1+x),x,method=_RETURNVERBOSE)`

output `x-ln(1+x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x} dx = x - \log(x+1)$$

input `integrate(x/(1+x),x, algorithm="fricas")`

output `x - log(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{x}{1+x} dx = x - \log(x+1)$$

input `integrate(x/(1+x),x)`

output `x - log(x + 1)`

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x} dx = x - \log(x+1)$$

input `integrate(x/(1+x),x, algorithm="maxima")`

output `x - log(x + 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 1.12

$$\int \frac{x}{1+x} dx = x - \log(|x+1|)$$

input `integrate(x/(1+x),x, algorithm="giac")`

output `x - log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x} dx = x - \ln(x+1)$$

input `int(x/(x + 1),x)`

output `x - log(x + 1)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x} dx = -\log(x+1) + x$$

input `int(x/(1+x),x)`

output `- log(x + 1) + x`

3.97 $\int \frac{x}{1+x^2} dx$

Optimal result	644
Mathematica [A] (verified)	644
Rubi [A] (verified)	645
Maple [A] (verified)	646
Fricas [A] (verification not implemented)	646
Sympy [A] (verification not implemented)	647
Maxima [A] (verification not implemented)	647
Giac [A] (verification not implemented)	647
Mupad [B] (verification not implemented)	648
Reduce [B] (verification not implemented)	648

Optimal result

Integrand size = 9, antiderivative size = 10

$$\int \frac{x}{1+x^2} dx = \frac{1}{2} \log(1+x^2)$$

output `1/2*ln(x^2+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x^2} dx = \frac{1}{2} \log(1+x^2)$$

input `Integrate[x/(1 + x^2), x]`

output `Log[1 + x^2]/2`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^2 + 1} dx$$

$$\downarrow 240$$

$$\frac{1}{2} \log(x^2 + 1)$$

input

```
Int[x/(1 + x^2), x]
```

output

```
Log[1 + x^2]/2
```

Defintions of rubi rules used

rule 240

```
Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]
```

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.90

method	result	size
derivativedivides	$\frac{\ln(x^2+1)}{2}$	9
default	$\frac{\ln(x^2+1)}{2}$	9
norman	$\frac{\ln(x^2+1)}{2}$	9
meijerg	$\frac{\ln(x^2+1)}{2}$	9
risch	$\frac{\ln(x^2+1)}{2}$	9
parallelrisch	$\frac{\ln(x^2+1)}{2}$	9

input `int(x/(x^2+1),x,method=_RETURNVERBOSE)`

output `1/2*ln(x^2+1)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{1+x^2} dx = \frac{1}{2} \log(x^2 + 1)$$

input `integrate(x/(x^2+1),x, algorithm="fricas")`

output `1/2*log(x^2 + 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{x}{1+x^2} dx = \frac{\log(x^2+1)}{2}$$

input `integrate(x/(x**2+1),x)`

output `log(x**2 + 1)/2`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{1+x^2} dx = \frac{1}{2} \log(x^2+1)$$

input `integrate(x/(x^2+1),x, algorithm="maxima")`

output `1/2*log(x^2 + 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{1+x^2} dx = \frac{1}{2} \log(x^2+1)$$

input `integrate(x/(x^2+1),x, algorithm="giac")`

output `1/2*log(x^2 + 1)`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{1+x^2} dx = \frac{\ln(x^2+1)}{2}$$

input `int(x/(x^2 + 1),x)`

output `log(x^2 + 1)/2`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{x}{1+x^2} dx = \frac{\log(x^2+1)}{2}$$

input `int(x/(x^2+1),x)`

output `log(x**2 + 1)/2`

3.98 $\int \frac{x}{1+x^3} dx$

Optimal result	649
Mathematica [A] (verified)	649
Rubi [A] (verified)	650
Maple [A] (verified)	652
Fricas [A] (verification not implemented)	652
Sympy [A] (verification not implemented)	653
Maxima [A] (verification not implemented)	653
Giac [A] (verification not implemented)	653
Mupad [B] (verification not implemented)	654
Reduce [B] (verification not implemented)	654

Optimal result

Integrand size = 9, antiderivative size = 40

$$\int \frac{x}{1+x^3} dx = \frac{\arctan\left(\frac{-1+2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{6} \log\left(\frac{(1+x)^2}{1-x+x^2}\right)$$

output `1/3*arctan(1/3*(-1+2*x)*3^(1/2))*3^(1/2)-1/6*ln((1+x)^2/(x^2-x+1))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x^3} dx = \frac{\arctan\left(\frac{-1+2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3} \log(1+x) + \frac{1}{6} \log(1-x+x^2)$$

input `Integrate[x/(1 + x^3), x]`

output `ArcTan[(-1 + 2*x)/Sqrt[3]]/Sqrt[3] - Log[1 + x]/3 + Log[1 - x + x^2]/6`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {821, 16, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{x^3 + 1} dx \\
 & \quad \downarrow \text{821} \\
 & \frac{1}{3} \int \frac{x+1}{x^2-x+1} dx - \frac{1}{3} \int \frac{1}{x+1} dx \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{3} \int \frac{x+1}{x^2-x+1} dx - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^2-x+1} dx + \frac{1}{2} \int -\frac{1-2x}{x^2-x+1} dx \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{25} \\
 & \frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^2-x+1} dx - \frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(-\frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx - 3 \int \frac{1}{-(2x-1)^2-3} d(2x-1) \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) - \frac{1}{2} \int \frac{1-2x}{x^2-x+1} dx \right) - \frac{1}{3} \log(x+1) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2x-1}{\sqrt{3}} \right) + \frac{1}{2} \log(x^2-x+1) \right) - \frac{1}{3} \log(x+1)
 \end{aligned}$$

input

Int[x/(1 + x^3), x]

output
$$-1/3 \cdot \text{Log}[1 + x] + (\text{Sqrt}[3] \cdot \text{ArcTan}[(-1 + 2x)/\text{Sqrt}[3]] + \text{Log}[1 - x + x^2]/2)/3$$

Defintions of rubi rules used

rule 16
$$\text{Int}[(c_)/((a_)+(b_)(x_)), x_Symbol] \rightarrow \text{Simp}[c \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$$

rule 25
$$\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[Fx, x], x]$$

rule 217
$$\text{Int}(((a_)+(b_)(x_)^2)^{-1}, x_Symbol) \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1}) \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 821
$$\text{Int}[(x_)/((a_)+(b_)(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3])^{-1} \quad \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x), x], x] + \text{Simp}[1/(3 \cdot \text{Rt}[a, 3] \cdot \text{Rt}[b, 3]) \quad \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3] \cdot \text{Rt}[b, 3] \cdot x + \text{Rt}[b, 3]^2 \cdot x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$$

rule 1083
$$\text{Int}(((a_)+(b_)(x_)+(c_)(x_)^2)^{-1}, x_Symbol) \rightarrow \text{Simp}[-2 \quad \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /; \text{FreeQ}[\{a, b, c\}, x]$$

rule 1103
$$\text{Int}(((d_)+(e_)(x_))/((a_)+(b_)(x_)+(c_)(x_)^2), x_Symbol) \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$$

rule 1142
$$\text{Int}(((d_)+(e_)(x_))/((a_)+(b_)(x_)+(c_)(x_)^2), x_Symbol) \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \quad \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \quad \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.88

method	result	size
default	$-\frac{\ln(1+x)}{3} + \frac{\ln(x^2-x+1)}{6} + \frac{\arctan\left(\frac{(-1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3}$	35
risch	$\frac{\ln(4x^2-4x+4)}{6} + \frac{\arctan\left(\frac{(-1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} - \frac{\ln(1+x)}{3}$	37
meijerg	$-\frac{x^2 \ln\left(1+(x^3)^{\frac{1}{3}}\right)}{3(x^3)^{\frac{2}{3}}} + \frac{x^2 \ln\left(1-(x^3)^{\frac{1}{3}}+(x^3)^{\frac{2}{3}}\right)}{6(x^3)^{\frac{2}{3}}} + \frac{x^2\sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2-(x^3)^{\frac{1}{3}}}\right)}{3(x^3)^{\frac{2}{3}}}$	80

input `int(x/(x^3+1),x,method=_RETURNVERBOSE)`output `-1/3*ln(1+x)+1/6*ln(x^2-x+1)+1/3*arctan(1/3*(-1+2*x)*3^(1/2))*3^(1/2)`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

$$\int \frac{x}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(x+1)$$

input `integrate(x/(x^3+1),x, algorithm="fricas")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + 1/6*log(x^2 - x + 1) - 1/3*log(x + 1)`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.02

$$\int \frac{x}{1+x^3} dx = -\frac{\log(x+1)}{3} + \frac{\log(x^2-x+1)}{6} + \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x - \sqrt{3}}{3}\right)}{3}$$

input `integrate(x/(x**3+1),x)`output `-log(x + 1)/3 + log(x**2 - x + 1)/6 + sqrt(3)*atan(2*sqrt(3)*x/3 - sqrt(3)/3)/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.85

$$\int \frac{x}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(x+1)$$

input `integrate(x/(x^3+1),x, algorithm="maxima")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + 1/6*log(x^2 - x + 1) - 1/3*log(x + 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.88

$$\int \frac{x}{1+x^3} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x-1)\right) + \frac{1}{6} \log(x^2-x+1) - \frac{1}{3} \log(|x+1|)$$

input `integrate(x/(x^3+1),x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x - 1)) + 1/6*log(x^2 - x + 1) - 1/3*log(abs(x + 1))`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.15

$$\int \frac{x}{1+x^3} dx = -\frac{\ln(x+1)}{3} - \ln\left(x - \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right) \\ + \ln\left(x - \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) \left(\frac{1}{6} + \frac{\sqrt{3}1i}{6}\right)$$

input `int(x/(x^3 + 1),x)`output `log(x + (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/6 + 1/6) - log(x - (3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/6 - 1/6) - log(x + 1)/3`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.82

$$\int \frac{x}{1+x^3} dx = \frac{\sqrt{3} \operatorname{atan}\left(\frac{2x-1}{\sqrt{3}}\right)}{3} + \frac{\log(x^2 - x + 1)}{6} - \frac{\log(x + 1)}{3}$$

input `int(x/(x^3+1),x)`output `(2*sqrt(3)*atan((2*x - 1)/sqrt(3)) + log(x**2 - x + 1) - 2*log(x + 1))/6`

3.99 $\int \frac{x}{1+x^4} dx$

Optimal result	655
Mathematica [A] (verified)	655
Rubi [A] (verified)	656
Maple [A] (verified)	657
Fricas [A] (verification not implemented)	657
Sympy [A] (verification not implemented)	657
Maxima [A] (verification not implemented)	658
Giac [A] (verification not implemented)	658
Mupad [B] (verification not implemented)	658
Reduce [B] (verification not implemented)	659

Optimal result

Integrand size = 9, antiderivative size = 8

$$\int \frac{x}{1+x^4} dx = \frac{\arctan(x^2)}{2}$$

output

```
1/2*arctan(x^2)
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00

$$\int \frac{x}{1+x^4} dx = \frac{\arctan(x^2)}{2}$$

input

```
Integrate[x/(1 + x^4), x]
```

output

```
ArcTan[x^2]/2
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {807, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{x^4 + 1} dx$$

$$\downarrow 807$$

$$\frac{1}{2} \int \frac{1}{x^4 + 1} dx^2$$

$$\downarrow 216$$

$$\frac{\arctan(x^2)}{2}$$

input

```
Int[x/(1 + x^4), x]
```

output

```
ArcTan[x^2]/2
```

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

rule 807

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.88

method	result	size
default	$\frac{\arctan(x^2)}{2}$	7
meijerg	$\frac{\arctan(x^2)}{2}$	7
risch	$\frac{\arctan(x^2)}{2}$	7
parallelrisc	$\frac{i \ln(x^2+i)}{4} - \frac{i \ln(x^2-i)}{4}$	22

input `int(x/(x^4+1),x,method=_RETURNVERBOSE)`output `1/2*arctan(x^2)`**Fricas [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{1}{2} \arctan(x^2)$$

input `integrate(x/(x^4+1),x, algorithm="fricas")`output `1/2*arctan(x^2)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.62

$$\int \frac{x}{1+x^4} dx = \frac{\text{atan}(x^2)}{2}$$

input `integrate(x/(x**4+1),x)`

output `atan(x**2)/2`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{1}{2} \arctan(x^2)$$

input `integrate(x/(x^4+1),x, algorithm="maxima")`

output `1/2*arctan(x^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{1}{2} \arctan(x^2)$$

input `integrate(x/(x^4+1),x, algorithm="giac")`

output `1/2*arctan(x^2)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.75

$$\int \frac{x}{1+x^4} dx = \frac{\operatorname{atan}(x^2)}{2}$$

input `int(x/(x^4 + 1),x)`

output `atan(x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 3.62

$$\int \frac{x}{1+x^4} dx = -\frac{\operatorname{atan}\left(\frac{\sqrt{2}-2x}{\sqrt{2}}\right)}{2} - \frac{\operatorname{atan}\left(\frac{\sqrt{2}+2x}{\sqrt{2}}\right)}{2}$$

input `int(x/(x^4+1),x)`output `(- (atan((sqrt(2) - 2*x)/sqrt(2)) + atan((sqrt(2) + 2*x)/sqrt(2))))/2`

3.100 $\int \frac{x}{1-x} dx$

Optimal result	660
Mathematica [A] (verified)	660
Rubi [A] (verified)	661
Maple [A] (verified)	662
Fricas [A] (verification not implemented)	662
Sympy [A] (verification not implemented)	662
Maxima [A] (verification not implemented)	663
Giac [A] (verification not implemented)	663
Mupad [B] (verification not implemented)	663
Reduce [B] (verification not implemented)	664

Optimal result

Integrand size = 9, antiderivative size = 12

$$\int \frac{x}{1-x} dx = -x - \log(1-x)$$

output `-x-ln(1-x)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{1-x} dx = -x - \log(1-x)$$

input `Integrate[x/(1 - x), x]`

output `-x - Log[1 - x]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{1-x} dx$$

$$\downarrow 49$$

$$\int \left(\frac{1}{1-x} - 1 \right) dx$$

$$\downarrow 2009$$

$$-x - \log(1-x)$$

input

```
Int[x/(1 - x),x]
```

output

```
-x - Log[1 - x]
```

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_)^(m_.))*((c_.) + (d_.)*(x_)^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
default	$-x - \ln(x - 1)$	11
norman	$-x - \ln(x - 1)$	11
risch	$-x - \ln(x - 1)$	11
parallelrisc	$-x - \ln(x - 1)$	11
meijerg	$-x - \ln(1 - x)$	13

input `int(x/(1-x),x,method=_RETURNVERBOSE)`

output `-x-ln(x-1)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{x}{1-x} dx = -x - \log(x - 1)$$

input `integrate(x/(1-x),x, algorithm="fricas")`

output `-x - log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{x}{1-x} dx = -x - \log(x - 1)$$

input `integrate(x/(1-x),x)`

output `-x - log(x - 1)`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{x}{1-x} dx = -x - \log(x-1)$$

input `integrate(x/(1-x),x, algorithm="maxima")`

output `-x - log(x - 1)`

Giac [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{x}{1-x} dx = -x - \log(|x-1|)$$

input `integrate(x/(1-x),x, algorithm="giac")`

output `-x - log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{x}{1-x} dx = -x - \ln(x-1)$$

input `int(-x/(x - 1),x)`

output `- x - log(x - 1)`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{x}{1-x} dx = -\log(x-1) - x$$

input `int(x/(1-x),x)`

output `- (log(x - 1) + x)`

3.101 $\int \frac{x}{1-x^2} dx$

Optimal result	665
Mathematica [A] (verified)	665
Rubi [A] (verified)	666
Maple [A] (verified)	667
Fricas [A] (verification not implemented)	667
Sympy [A] (verification not implemented)	668
Maxima [A] (verification not implemented)	668
Giac [A] (verification not implemented)	668
Mupad [B] (verification not implemented)	669
Reduce [B] (verification not implemented)	669

Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{x}{1-x^2} dx = -\frac{1}{2} \log(1-x^2)$$

output `-1/2*ln(-x^2+1)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{x}{1-x^2} dx = -\frac{1}{2} \log(1-x^2)$$

input `Integrate[x/(1 - x^2), x]`

output `-1/2*Log[1 - x^2]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {240}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{1-x^2} dx$$

↓ 240

$$-\frac{1}{2} \log(1-x^2)$$

input

```
Int[x/(1 - x^2), x]
```

output

```
-1/2*Log[1 - x^2]
```

Defintions of rubi rules used

rule 240

```
Int[(x_)/((a_) + (b_.)*(x_)^2), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^2, x]]/(2*b), x] /; FreeQ[{a, b}, x]
```

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
risch	$-\frac{\ln(x^2-1)}{2}$	9
derivativedivides	$-\frac{\ln(-x^2+1)}{2}$	11
meijerg	$-\frac{\ln(-x^2+1)}{2}$	11
default	$-\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}$	14
norman	$-\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}$	14
parallelrisch	$-\frac{\ln(x-1)}{2} - \frac{\ln(1+x)}{2}$	14

input `int(x/(-x^2+1),x,method=_RETURNVERBOSE)`output `-1/2*ln(x^2-1)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{1-x^2} dx = -\frac{1}{2} \log(x^2 - 1)$$

input `integrate(x/(-x^2+1),x, algorithm="fricas")`output `-1/2*log(x^2 - 1)`

Sympy [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{1-x^2} dx = -\frac{\log(x^2-1)}{2}$$

input `integrate(x/(-x**2+1),x)`

output `-log(x**2 - 1)/2`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{1-x^2} dx = -\frac{1}{2} \log(x^2-1)$$

input `integrate(x/(-x^2+1),x, algorithm="maxima")`

output `-1/2*log(x^2 - 1)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

$$\int \frac{x}{1-x^2} dx = -\frac{1}{2} \log(|x^2-1|)$$

input `integrate(x/(-x^2+1),x, algorithm="giac")`

output `-1/2*log(abs(x^2 - 1))`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{x}{1-x^2} dx = -\frac{\ln(x^2-1)}{2}$$

input `int(-x/(x^2 - 1),x)`

output `-log(x^2 - 1)/2`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{x}{1-x^2} dx = -\frac{\log(x-1)}{2} - \frac{\log(x+1)}{2}$$

input `int(x/(-x^2+1),x)`

output `(- (log(x - 1) + log(x + 1)))/2`

3.102 $\int \frac{x}{1-x^3} dx$

Optimal result	670
Mathematica [A] (verified)	670
Rubi [A] (verified)	671
Maple [A] (verified)	673
Fricas [A] (verification not implemented)	673
Sympy [A] (verification not implemented)	674
Maxima [A] (verification not implemented)	674
Giac [A] (verification not implemented)	674
Mupad [B] (verification not implemented)	675
Reduce [B] (verification not implemented)	675

Optimal result

Integrand size = 11, antiderivative size = 41

$$\int \frac{x}{1-x^3} dx = -\frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{6} \log\left(\frac{(1-x)^2}{1+x+x^2}\right)$$

output `-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)-1/6*ln((1-x)^2/(x^2+x+1))`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{x}{1-x^3} dx = -\frac{\arctan\left(\frac{1+2x}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{3} \log(1-x) + \frac{1}{6} \log(1+x+x^2)$$

input `Integrate[x/(1 - x^3), x]`

output `-(ArcTan[(1 + 2*x)/Sqrt[3]]/Sqrt[3]) - Log[1 - x]/3 + Log[1 + x + x^2]/6`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.545$, Rules used = {821, 16, 1142, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{1-x^3} dx \\
 & \quad \downarrow \text{821} \\
 & \frac{1}{3} \int \frac{1}{1-x} dx - \frac{1}{3} \int \frac{1-x}{x^2+x+1} dx \\
 & \quad \downarrow \text{16} \\
 & -\frac{1}{3} \int \frac{1-x}{x^2+x+1} dx - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1142} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx - \frac{3}{2} \int \frac{1}{x^2+x+1} dx \right) - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1083} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx + 3 \int \frac{1}{-(2x+1)^2-3} d(2x+1) \right) - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{217} \\
 & \frac{1}{3} \left(\frac{1}{2} \int \frac{2x+1}{x^2+x+1} dx - \sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) \right) - \frac{1}{3} \log(1-x) \\
 & \quad \downarrow \text{1103} \\
 & \frac{1}{3} \left(\frac{1}{2} \log(x^2+x+1) - \sqrt{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) \right) - \frac{1}{3} \log(1-x)
 \end{aligned}$$

input `Int[x/(1 - x^3), x]`

output `-1/3*Log[1 - x] + (-(Sqrt[3]*ArcTan[(1 + 2*x)/Sqrt[3]]) + Log[1 + x + x^2])/3`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$
- rule 821 $\text{Int}(x_)/((a_)+(b_)*(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{(-1)} \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]) \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 1083 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[-2 \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_)+(e_)*(x_)/((a_)+(b_)*(x_)+(c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{\ln(x^2+x+1)}{6} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} - \frac{\ln(x-1)}{3}$	33
risch	$\frac{\ln(4x^2+4x+4)}{6} - \frac{\arctan\left(\frac{(1+2x)\sqrt{3}}{3}\right)\sqrt{3}}{3} - \frac{\ln(x-1)}{3}$	37
meijerg	$-\frac{x^2 \left(\ln\left(1 - (x^3)^{\frac{1}{3}}\right) - \frac{\ln\left(1 + (x^3)^{\frac{1}{3}} + (x^3)^{\frac{2}{3}}\right)}{2} + \sqrt{3} \arctan\left(\frac{\sqrt{3}(x^3)^{\frac{1}{3}}}{2 + (x^3)^{\frac{1}{3}}}\right) \right)}{3(x^3)^{\frac{2}{3}}}$	63

input `int(x/(-x^3+1),x,method=_RETURNVERBOSE)`output `1/6*ln(x^2+x+1)-1/3*arctan(1/3*(1+2*x)*3^(1/2))*3^(1/2)-1/3*ln(x-1)`**Fricas [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \frac{x}{1-x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{6} \log(x^2+x+1) - \frac{1}{3} \log(x-1)$$

input `integrate(x/(-x^3+1),x, algorithm="fricas")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/6*log(x^2 + x + 1) - 1/3*log(x - 1)`

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{x}{1-x^3} dx = -\frac{\log(x-1)}{3} + \frac{\log(x^2+x+1)}{6} - \frac{\sqrt{3} \operatorname{atan}\left(\frac{2\sqrt{3}x}{3} + \frac{\sqrt{3}}{3}\right)}{3}$$

input `integrate(x/(-x**3+1),x)`output `-log(x - 1)/3 + log(x**2 + x + 1)/6 - sqrt(3)*atan(2*sqrt(3)*x/3 + sqrt(3)/3)/3`**Maxima [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \frac{x}{1-x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{6} \log(x^2+x+1) - \frac{1}{3} \log(x-1)$$

input `integrate(x/(-x^3+1),x, algorithm="maxima")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/6*log(x^2 + x + 1) - 1/3*log(x - 1)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \frac{x}{1-x^3} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x+1)\right) + \frac{1}{6} \log(x^2+x+1) - \frac{1}{3} \log(|x-1|)$$

input `integrate(x/(-x^3+1),x, algorithm="giac")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*x + 1)) + 1/6*log(x^2 + x + 1) - 1/3*log(abs(x - 1))`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.12

$$\int \frac{x}{1-x^3} dx = -\frac{\ln(x-1)}{3} + \ln\left(x + \frac{1}{2} - \frac{\sqrt{3} \text{li}}{2}\right) \left(\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6}\right) - \ln\left(x + \frac{1}{2} + \frac{\sqrt{3} \text{li}}{2}\right) \left(-\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6}\right)$$

input `int(-x/(x^3 - 1),x)`output `log(x - (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 + 1/6) - log(x - 1)/3 - log(x + (3^(1/2)*1i)/2 + 1/2)*((3^(1/2)*1i)/6 - 1/6)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \frac{x}{1-x^3} dx = -\frac{\sqrt{3} \operatorname{atan}\left(\frac{2x+1}{\sqrt{3}}\right)}{3} + \frac{\log(x^2 + x + 1)}{6} - \frac{\log(x-1)}{3}$$

input `int(x/(-x^3+1),x)`output `(- 2*sqrt(3)*atan((2*x + 1)/sqrt(3)) + log(x**2 + x + 1) - 2*log(x - 1))/6`

3.103 $\int \frac{x}{1-x^4} dx$

Optimal result	676
Mathematica [A] (verified)	676
Rubi [A] (verified)	677
Maple [A] (verified)	678
Fricas [A] (verification not implemented)	678
Sympy [A] (verification not implemented)	679
Maxima [A] (verification not implemented)	679
Giac [A] (verification not implemented)	679
Mupad [B] (verification not implemented)	680
Reduce [B] (verification not implemented)	680

Optimal result

Integrand size = 11, antiderivative size = 20

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log \left(\frac{1+x^2}{1-x^2} \right)$$

output `1/4*ln((x^2+1)/(-x^2+1))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{x}{1-x^4} dx = -\frac{1}{4} \log(1-x^2) + \frac{1}{4} \log(1+x^2)$$

input `Integrate[x/(1 - x^4),x]`

output `-1/4*Log[1 - x^2] + Log[1 + x^2]/4`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.40, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {807, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{1-x^4} dx$$

$$\downarrow 807$$

$$\frac{1}{2} \int \frac{1}{1-x^4} dx^2$$

$$\downarrow 219$$

$$\frac{\operatorname{arctanh}(x^2)}{2}$$

input `Int[x/(1 - x^4), x]`

output `ArcTanh[x^2]/2`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.35

method	result	size
meijerg	$\frac{\operatorname{arctanh}(x^2)}{2}$	7
risch	$\frac{\ln(x^2+1)}{4} - \frac{\ln(x^2-1)}{4}$	18
default	$-\frac{\ln(x-1)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22
norman	$-\frac{\ln(x-1)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22
parallelrisch	$-\frac{\ln(x-1)}{4} - \frac{\ln(1+x)}{4} + \frac{\ln(x^2+1)}{4}$	22

input `int(x/(-x^4+1),x,method=_RETURNVERBOSE)`output `1/2*arctanh(x^2)`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2+1) - \frac{1}{4} \log(x^2-1)$$

input `integrate(x/(-x^4+1),x, algorithm="fricas")`output `1/4*log(x^2 + 1) - 1/4*log(x^2 - 1)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.75

$$\int \frac{x}{1-x^4} dx = -\frac{\log(x^2-1)}{4} + \frac{\log(x^2+1)}{4}$$

input `integrate(x/(-x**4+1),x)`output `-log(x**2 - 1)/4 + log(x**2 + 1)/4`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2+1) - \frac{1}{4} \log(x^2-1)$$

input `integrate(x/(-x^4+1),x, algorithm="maxima")`output `1/4*log(x^2 + 1) - 1/4*log(x^2 - 1)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{x}{1-x^4} dx = \frac{1}{4} \log(x^2+1) - \frac{1}{4} \log(|x^2-1|)$$

input `integrate(x/(-x^4+1),x, algorithm="giac")`output `1/4*log(x^2 + 1) - 1/4*log(abs(x^2 - 1))`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.30

$$\int \frac{x}{1-x^4} dx = \frac{\operatorname{atanh}(x^2)}{2}$$

input `int(-x/(x^4 - 1),x)`

output `atanh(x^2)/2`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.05

$$\int \frac{x}{1-x^4} dx = \frac{\log(x^2+1)}{4} - \frac{\log(x-1)}{4} - \frac{\log(x+1)}{4}$$

input `int(x/(-x^4+1),x)`

output `(log(x**2 + 1) - log(x - 1) - log(x + 1))/4`

3.104 $\int \frac{1}{x(1+x^2)} dx$

Optimal result	681
Mathematica [A] (verified)	681
Rubi [A] (verified)	682
Maple [A] (verified)	683
Fricas [A] (verification not implemented)	684
Sympy [A] (verification not implemented)	684
Maxima [A] (verification not implemented)	684
Giac [A] (verification not implemented)	685
Mupad [B] (verification not implemented)	685
Reduce [B] (verification not implemented)	685

Optimal result

Integrand size = 11, antiderivative size = 12

$$\int \frac{1}{x(1+x^2)} dx = \log\left(\frac{x}{\sqrt{1+x^2}}\right)$$

output `ln(x/(x^2+1)^(1/2))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

$$\int \frac{1}{x(1+x^2)} dx = \log(x) - \frac{1}{2} \log(1+x^2)$$

input `Integrate[1/(x*(1 + x^2)),x]`

output `Log[x] - Log[1 + x^2]/2`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.42, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x(x^2+1)} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{1}{x^2(x^2+1)} dx^2 \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(\int \frac{1}{x^2} dx^2 - \int \frac{1}{x^2+1} dx^2 \right) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(\log(x^2) - \int \frac{1}{x^2+1} dx^2 \right) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(x^2) - \log(x^2+1))
 \end{aligned}$$

input `Int[1/(x*(1 + x^2)),x]`

output `(Log[x^2] - Log[1 + x^2])/2`

Defintions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{\ln(x^2+1)}{2} + \ln(x)$	12
norman	$-\frac{\ln(x^2+1)}{2} + \ln(x)$	12
meijerg	$-\frac{\ln(x^2+1)}{2} + \ln(x)$	12
risch	$-\frac{\ln(x^2+1)}{2} + \ln(x)$	12
parallelrisc	$-\frac{\ln(x^2+1)}{2} + \ln(x)$	12

input `int(1/x/(x^2+1),x,method=_RETURNVERBOSE)`

output `-1/2*ln(x^2+1)+ln(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{x(1+x^2)} dx = -\frac{1}{2} \log(x^2 + 1) + \log(x)$$

input `integrate(1/x/(x^2+1),x, algorithm="fricas")`output `-1/2*log(x^2 + 1) + log(x)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{x(1+x^2)} dx = \log(x) - \frac{\log(x^2 + 1)}{2}$$

input `integrate(1/x/(x**2+1),x)`output `log(x) - log(x**2 + 1)/2`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

$$\int \frac{1}{x(1+x^2)} dx = -\frac{1}{2} \log(x^2 + 1) + \frac{1}{2} \log(x^2)$$

input `integrate(1/x/(x^2+1),x, algorithm="maxima")`output `-1/2*log(x^2 + 1) + 1/2*log(x^2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.25

$$\int \frac{1}{x(1+x^2)} dx = -\frac{1}{2} \log(x^2 + 1) + \frac{1}{2} \log(x^2)$$

input `integrate(1/x/(x^2+1),x, algorithm="giac")`

output `-1/2*log(x^2 + 1) + 1/2*log(x^2)`

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{x(1+x^2)} dx = \ln(x) - \frac{\ln(x^2 + 1)}{2}$$

input `int(1/(x*(x^2 + 1)),x)`

output `log(x) - log(x^2 + 1)/2`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

$$\int \frac{1}{x(1+x^2)} dx = -\frac{\log(x^2 + 1)}{2} + \log(x)$$

input `int(1/x/(x^2+1),x)`

output `(- log(x**2 + 1) + 2*log(x))/2`

3.105 $\int \frac{1}{x(1-x^2)} dx$

Optimal result	686
Mathematica [A] (verified)	686
Rubi [A] (verified)	687
Maple [A] (verified)	688
Fricas [A] (verification not implemented)	689
Sympy [A] (verification not implemented)	689
Maxima [A] (verification not implemented)	689
Giac [A] (verification not implemented)	690
Mupad [B] (verification not implemented)	690
Reduce [B] (verification not implemented)	690

Optimal result

Integrand size = 13, antiderivative size = 14

$$\int \frac{1}{x(1-x^2)} dx = \log\left(\frac{x}{\sqrt{1-x^2}}\right)$$

output `ln(x/(-x^2+1)^(1/2))`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(1-x^2)} dx = \log(x) - \frac{1}{2} \log(1-x^2)$$

input `Integrate[1/(x*(1-x^2)),x]`

output `Log[x] - Log[1-x^2]/2`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {243, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x(1-x^2)} dx \\
 & \quad \downarrow \text{243} \\
 & \frac{1}{2} \int \frac{1}{x^2(1-x^2)} dx^2 \\
 & \quad \downarrow \text{47} \\
 & \frac{1}{2} \left(\int \frac{1}{x^2} dx^2 + \int \frac{1}{1-x^2} dx^2 \right) \\
 & \quad \downarrow \text{14} \\
 & \frac{1}{2} \left(\int \frac{1}{1-x^2} dx^2 + \log(x^2) \right) \\
 & \quad \downarrow \text{16} \\
 & \frac{1}{2} (\log(x^2) - \log(1-x^2))
 \end{aligned}$$

input `Int[1/(x*(1 - x^2)),x]`

output `(Log[x^2] - Log[1 - x^2])/2`

Definitions of rubi rules used

- rule 14 `Int[(a_.)/(x_), x_Symbol] := Simp[a*Log[x], x] /; FreeQ[a, x]`
- rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`
- rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`
- rule 243 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[1/2 Subst[Int[x^((m - 1)/2)*(a + b*x)^p, x], x, x^2], x] /; FreeQ[{a, b, m, p}, x] && IntegerQ[(m - 1)/2]`

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

method	result	size
risch	$\ln(x) - \frac{\ln(x^2-1)}{2}$	12
default	$-\frac{\ln(1+x)}{2} + \ln(x) - \frac{\ln(x-1)}{2}$	16
norman	$-\frac{\ln(1+x)}{2} + \ln(x) - \frac{\ln(x-1)}{2}$	16
parallelrisc	$-\frac{\ln(1+x)}{2} + \ln(x) - \frac{\ln(x-1)}{2}$	16
meijerg	$\ln(x) + \frac{i\pi}{2} - \frac{\ln(-x^2+1)}{2}$	18

input `int(1/x/(-x^2+1),x,method=_RETURNVERBOSE)`

output `ln(x)-1/2*ln(x^2-1)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \frac{1}{x(1-x^2)} dx = -\frac{1}{2} \log(x^2 - 1) + \log(x)$$

input `integrate(1/x/(-x^2+1),x, algorithm="fricas")`output `-1/2*log(x^2 - 1) + log(x)`**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{x(1-x^2)} dx = \log(x) - \frac{\log(x^2 - 1)}{2}$$

input `integrate(1/x/(-x**2+1),x)`output `log(x) - log(x**2 - 1)/2`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(1-x^2)} dx = -\frac{1}{2} \log(x^2 - 1) + \frac{1}{2} \log(x^2)$$

input `integrate(1/x/(-x^2+1),x, algorithm="maxima")`output `-1/2*log(x^2 - 1) + 1/2*log(x^2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{1}{x(1-x^2)} dx = \frac{1}{2} \log(x^2) - \frac{1}{2} \log(|x^2 - 1|)$$

input `integrate(1/x/(-x^2+1),x, algorithm="giac")`

output `1/2*log(x^2) - 1/2*log(abs(x^2 - 1))`

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \frac{1}{x(1-x^2)} dx = \ln(x) - \frac{\ln(x^2 - 1)}{2}$$

input `int(-1/(x*(x^2 - 1)),x)`

output `log(x) - log(x^2 - 1)/2`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(1-x^2)} dx = -\frac{\log(x-1)}{2} - \frac{\log(x+1)}{2} + \log(x)$$

input `int(1/x/(-x^2+1),x)`

output `(- log(x - 1) - log(x + 1) + 2*log(x))/2`

3.106 $\int \frac{a+bx}{A+Bx} dx$

Optimal result	691
Mathematica [A] (verified)	691
Rubi [A] (verified)	692
Maple [A] (verified)	693
Fricas [A] (verification not implemented)	693
Sympy [A] (verification not implemented)	693
Maxima [A] (verification not implemented)	694
Giac [A] (verification not implemented)	694
Mupad [B] (verification not implemented)	694
Reduce [B] (verification not implemented)	695

Optimal result

Integrand size = 13, antiderivative size = 25

$$\int \frac{a+bx}{A+Bx} dx = \frac{bx}{B} + \frac{(-Ab+aB)\log(A+Bx)}{B^2}$$

output `b*x/B+(-A*b+B*a)*ln(B*x+A)/B^2`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{a+bx}{A+Bx} dx = \frac{bx}{B} + \frac{(-Ab+aB)\log(A+Bx)}{B^2}$$

input `Integrate[(a + b*x)/(A + B*x),x]`

output `(b*x)/B + ((-A*b) + a*B)*Log[A + B*x])/B^2`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + bx}{A + Bx} dx$$

$$\downarrow 49$$

$$\int \left(\frac{aB - Ab}{B(A + Bx)} + \frac{b}{B} \right) dx$$

$$\downarrow 2009$$

$$\frac{bx}{B} - \frac{(Ab - aB) \log(A + Bx)}{B^2}$$

input

```
Int[(a + b*x)/(A + B*x),x]
```

output

```
(b*x)/B - ((A*b - a*B)*Log[A + B*x])/B^2
```

Defintions of rubi rules used

rule 49

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x]
&& IGtQ[m, 0] && IGtQ[m + n + 2, 0]
```

rule 2009

```
Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

method	result	size
default	$\frac{bx}{B} + \frac{(-Ab+Ba)\ln(Bx+A)}{B^2}$	26
norman	$\frac{bx}{B} - \frac{(Ab-Ba)\ln(Bx+A)}{B^2}$	27
parallelsch	$-\frac{A\ln(Bx+A)b-B\ln(Bx+A)a-xbB}{B^2}$	31
risch	$\frac{bx}{B} - \frac{\ln(Bx+A)Ab}{B^2} + \frac{\ln(Bx+A)a}{B}$	32

input `int((b*x+a)/(B*x+A),x,method=_RETURNVERBOSE)`output `b*x/B+(-A*b+B*a)*ln(B*x+A)/B^2`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{a + bx}{A + Bx} dx = \frac{Bbx + (Ba - Ab) \log(Bx + A)}{B^2}$$

input `integrate((b*x+a)/(B*x+A),x, algorithm="fricas")`output `(B*b*x + (B*a - A*b)*log(B*x + A))/B^2`**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{a + bx}{A + Bx} dx = \frac{bx}{B} + \frac{(-Ab + Ba) \log(A + Bx)}{B^2}$$

input `integrate((b*x+a)/(B*x+A),x)`

output $b*x/B + (-A*b + B*a)*\log(A + B*x)/B**2$

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{a + bx}{A + Bx} dx = \frac{bx}{B} + \frac{(Ba - Ab) \log(Bx + A)}{B^2}$$

input `integrate((b*x+a)/(B*x+A),x, algorithm="maxima")`

output $b*x/B + (B*a - A*b)*\log(B*x + A)/B^2$

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{a + bx}{A + Bx} dx = \frac{bx}{B} + \frac{(Ba - Ab) \log(|Bx + A|)}{B^2}$$

input `integrate((b*x+a)/(B*x+A),x, algorithm="giac")`

output $b*x/B + (B*a - A*b)*\log(\text{abs}(B*x + A))/B^2$

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{a + bx}{A + Bx} dx = \frac{bx}{B} - \frac{\ln(A + Bx) (Ab - Ba)}{B^2}$$

input `int((a + b*x)/(A + B*x),x)`

output $(b*x)/B - (\log(A + B*x)*(A*b - B*a))/B^2$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 1, normalized size of antiderivative = 0.04

$$\int \frac{a + bx}{A + Bx} dx = x$$

input `int((b*x+a)/(B*x+A), x)`

output `x`

3.107 $\int \frac{1}{(a+bx)(A+Bx)} dx$

Optimal result	696
Mathematica [A] (verified)	696
Rubi [A] (verified)	697
Maple [A] (verified)	698
Fricas [A] (verification not implemented)	698
Sympy [B] (verification not implemented)	698
Maxima [A] (verification not implemented)	699
Giac [A] (verification not implemented)	699
Mupad [B] (verification not implemented)	700
Reduce [B] (verification not implemented)	700

Optimal result

Integrand size = 15, antiderivative size = 25

$$\int \frac{1}{(a+bx)(A+Bx)} dx = \frac{\log\left(\frac{A+Bx}{a+bx}\right)}{-Ab+aB}$$

output `ln((B*x+A)/(b*x+a))/(-A*b+B*a)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{1}{(a+bx)(A+Bx)} dx = \frac{\log(a+bx) - \log(A+Bx)}{Ab - aB}$$

input `Integrate[1/((a + b*x)*(A + B*x)),x]`

output `(Log[a + b*x] - Log[A + B*x])/(A*b - a*B)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.44, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {47, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(a + bx)(A + Bx)} dx$$

$$\downarrow 47$$

$$\frac{b \int \frac{1}{a+bx} dx}{Ab - aB} - \frac{B \int \frac{1}{A+Bx} dx}{Ab - aB}$$

$$\downarrow 16$$

$$\frac{\log(a + bx)}{Ab - aB} - \frac{\log(A + Bx)}{Ab - aB}$$

input `Int[1/((a + b*x)*(A + B*x)),x]`

output `Log[a + b*x]/(A*b - a*B) - Log[A + B*x]/(A*b - a*B)`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] :> Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 47 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] :> Simp[b/(b*c - a*d) Int[1/(a + b*x), x], x] - Simp[d/(b*c - a*d) Int[1/(c + d*x), x], x] /; FreeQ[{a, b, c, d}, x]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.08

method	result	size
parallelrisc	$\frac{-\ln(Bx+A)+\ln(bx+a)}{Ab-Ba}$	27
default	$-\frac{\ln(Bx+A)}{Ab-Ba} + \frac{\ln(bx+a)}{Ab-Ba}$	37
norman	$-\frac{\ln(Bx+A)}{Ab-Ba} + \frac{\ln(bx+a)}{Ab-Ba}$	37
risc	$\frac{\ln(-bx-a)}{Ab-Ba} - \frac{\ln(Bx+A)}{Ab-Ba}$	40

input `int(1/(b*x+a)/(B*x+A),x,method=_RETURNVERBOSE)`

output `(-ln(B*x+A)+ln(b*x+a))/(A*b-B*a)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{1}{(a+bx)(A+Bx)} dx = \frac{\log(Bx+A) - \log(bx+a)}{Ba - Ab}$$

input `integrate(1/(b*x+a)/(B*x+A),x, algorithm="fricas")`

output `(log(B*x + A) - log(b*x + a))/(B*a - A*b)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 128 vs. $2(17) = 34$.

Time = 0.19 (sec) , antiderivative size = 128, normalized size of antiderivative = 5.12

$$\int \frac{1}{(a+bx)(A+Bx)} dx = \frac{\log\left(x + \frac{-\frac{A^2b^2}{-Ab+Ba} + \frac{2ABab}{-Ab+Ba} + Ab - \frac{B^2a^2}{-Ab+Ba} + Ba}{2Bb}\right)}{-Ab+Ba} - \frac{\log\left(x + \frac{\frac{A^2b^2}{-Ab+Ba} - \frac{2ABab}{-Ab+Ba} + Ab + \frac{B^2a^2}{-Ab+Ba} + Ba}{2Bb}\right)}{-Ab+Ba}$$

input `integrate(1/(b*x+a)/(B*x+A),x)`

output `log(x + (-A**2*b**2/(-A*b + B*a) + 2*A*B*a*b/(-A*b + B*a) + A*b - B**2*a**2/(-A*b + B*a) + B*a)/(2*B*b))/(-A*b + B*a) - log(x + (A**2*b**2/(-A*b + B*a) - 2*A*B*a*b/(-A*b + B*a) + A*b + B**2*a**2/(-A*b + B*a) + B*a)/(2*B*b))/(-A*b + B*a)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.44

$$\int \frac{1}{(a+bx)(A+Bx)} dx = \frac{\log(Bx+A)}{Ba-Ab} - \frac{\log(bx+a)}{Ba-Ab}$$

input `integrate(1/(b*x+a)/(B*x+A),x, algorithm="maxima")`

output `log(B*x + A)/(B*a - A*b) - log(b*x + a)/(B*a - A*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.84

$$\int \frac{1}{(a+bx)(A+Bx)} dx = \frac{B \log(|Bx+A|)}{B^2a-ABb} - \frac{b \log(|bx+a|)}{Bab-Ab^2}$$

input `integrate(1/(b*x+a)/(B*x+A),x, algorithm="giac")`

output $B \cdot \log(\text{abs}(B \cdot x + A)) / (B^2 \cdot a - A \cdot B \cdot b) - b \cdot \log(\text{abs}(b \cdot x + a)) / (B \cdot a \cdot b - A \cdot b^2)$

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + bx)(A + Bx)} dx = \frac{\ln\left(\frac{a+bx}{A+Bx}\right)}{Ab - Ba}$$

input `int(1/((A + B*x)*(a + b*x)),x)`

output $\log((a + b \cdot x) / (A + B \cdot x)) / (A \cdot b - B \cdot a)$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.48

$$\int \frac{1}{(a + bx)(A + Bx)} dx = \frac{x}{a(bx + a)}$$

input `int(1/(b*x+a)/(B*x+A),x)`

output $x / (a \cdot (a + b \cdot x))$

3.108 $\int \frac{x}{(a+bx)(A+Bx)} dx$

Optimal result	701
Mathematica [A] (verified)	701
Rubi [A] (verified)	702
Maple [A] (verified)	703
Fricas [A] (verification not implemented)	703
Sympy [B] (verification not implemented)	703
Maxima [A] (verification not implemented)	704
Giac [A] (verification not implemented)	705
Mupad [B] (verification not implemented)	705
Reduce [B] (verification not implemented)	705

Optimal result

Integrand size = 16, antiderivative size = 35

$$\int \frac{x}{(a+bx)(A+Bx)} dx = \frac{\frac{a \log(a+bx)}{b} - \frac{A \log(A+Bx)}{B}}{-Ab + aB}$$

output $(a*\ln(b*x+a)/b-A*\ln(B*x+A)/B)/(-A*b+B*a)$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int \frac{x}{(a+bx)(A+Bx)} dx = -\frac{aB \log(a+bx) - Ab \log(A+Bx)}{Ab^2B - abB^2}$$

input `Integrate[x/((a + b*x)*(A + B*x)),x]`

output $-((a*B*\text{Log}[a + b*x] - A*b*\text{Log}[A + B*x])/(A*b^2*B - a*b*B^2))$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {86, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(a+bx)(A+Bx)} dx$$

↓ 86

$$\int \left(\frac{A}{(A+Bx)(Ab-aB)} - \frac{a}{(a+bx)(Ab-aB)} \right) dx$$

↓ 2009

$$\frac{A \log(A+Bx)}{B(Ab-aB)} - \frac{a \log(a+bx)}{b(Ab-aB)}$$

input `Int[x/((a + b*x)*(A + B*x)),x]`

output `-((a*Log[a + b*x])/(b*(A*b - a*B))) + (A*Log[A + B*x])/(B*(A*b - a*B))`

Defintions of rubi rules used

rule 86 `Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_)^(n_.))*((e_.) + (f_.)*(x_)^(p_.)), x_] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

method	result	size
parallelrisch	$\frac{A \ln(Bx+A)b - a \ln(bx+a)B}{B(Ab-Ba)b}$	38
default	$\frac{A \ln(Bx+A)}{(Ab-Ba)B} - \frac{a \ln(bx+a)}{(Ab-Ba)b}$	45
norman	$\frac{A \ln(Bx+A)}{(Ab-Ba)B} - \frac{a \ln(bx+a)}{(Ab-Ba)b}$	45
risch	$-\frac{a \ln(bx+a)}{(Ab-Ba)b} + \frac{A \ln(-Bx-A)}{(Ab-Ba)B}$	48

input `int(x/(b*x+a)/(B*x+A),x,method=_RETURNVERBOSE)`output `(A*ln(B*x+A)*b-a*ln(b*x+a)*B)/B/(A*b-B*a)/b`**Fricas [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int \frac{x}{(a+bx)(A+Bx)} dx = -\frac{Ab \log(Bx+A) - Ba \log(bx+a)}{B^2 ab - ABb^2}$$

input `integrate(x/(b*x+a)/(B*x+A),x, algorithm="fricas")`output `-(A*b*log(B*x + A) - B*a*log(b*x + a))/(B^2*a*b - A*B*b^2)`**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 138 vs. 2(26) = 52.

Time = 0.41 (sec) , antiderivative size = 138, normalized size of antiderivative = 3.94

$$\int \frac{x}{(a+bx)(A+Bx)} dx = -\frac{A \log \left(x + \frac{-\frac{A^3 b^2}{B(-Ab+Ba)} + \frac{2A^2 ab}{-Ab+Ba} - \frac{ABa^2}{-Ab+Ba} + 2Aa}{Ab+Ba} \right)}{B(-Ab+Ba)} + \frac{a \log \left(x + \frac{\frac{A^2 ab}{-Ab+Ba} - \frac{2ABa^2}{-Ab+Ba} + 2Aa + \frac{B^2 a^3}{b(-Ab+Ba)}}{Ab+Ba} \right)}{b(-Ab+Ba)}$$

input `integrate(x/(b*x+a)/(B*x+A),x)`

output `-A*log(x + (-A**3*b**2/(B*(-A*b + B*a)) + 2*A**2*a*b/(-A*b + B*a) - A*B*a**2/(-A*b + B*a) + 2*A*a)/(A*b + B*a))/(B*(-A*b + B*a)) + a*log(x + (A**2*a*b/(-A*b + B*a) - 2*A*B*a**2/(-A*b + B*a) + 2*A*a + B**2*a**3/(b*(-A*b + B*a)))/(A*b + B*a))/(b*(-A*b + B*a))`

Maxima [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.26

$$\int \frac{x}{(a+bx)(A+Bx)} dx = -\frac{A \log(Bx + A)}{B^2 a - ABb} + \frac{a \log(bx + a)}{Bab - Ab^2}$$

input `integrate(x/(b*x+a)/(B*x+A),x, algorithm="maxima")`

output `-A*log(B*x + A)/(B^2*a - A*B*b) + a*log(b*x + a)/(B*a*b - A*b^2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.31

$$\int \frac{x}{(a+bx)(A+Bx)} dx = -\frac{A \log(|Bx+A|)}{B^2a - ABb} + \frac{a \log(|bx+a|)}{Bab - Ab^2}$$

input `integrate(x/(b*x+a)/(B*x+A),x, algorithm="giac")`

output `-A*log(abs(B*x + A))/(B^2*a - A*B*b) + a*log(abs(b*x + a))/(B*a*b - A*b^2)`

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.06

$$\int \frac{x}{(a+bx)(A+Bx)} dx = \frac{Ab \ln(A+Bx) - Ba \ln(a+bx)}{Bb(Ab - Ba)}$$

input `int(x/((A + B*x)*(a + b*x)),x)`

output `(A*b*log(A + B*x) - B*a*log(a + b*x))/(B*b*(A*b - B*a))`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{x}{(a+bx)(A+Bx)} dx = \frac{\log(bx+a)a + \log(bx+a)bx - bx}{b^2(bx+a)}$$

input `int(x/(b*x+a)/(B*x+A),x)`

output `(log(a + b*x)*a + log(a + b*x)*b*x - b*x)/(b**2*(a + b*x))`

3.109 $\int \frac{1}{\sqrt{x}(a+bx)} dx$

Optimal result	706
Mathematica [A] (verified)	706
Rubi [A] (verified)	707
Maple [A] (verified)	708
Fricas [A] (verification not implemented)	708
Sympy [B] (verification not implemented)	708
Maxima [A] (verification not implemented)	709
Giac [A] (verification not implemented)	710
Mupad [B] (verification not implemented)	710
Reduce [B] (verification not implemented)	710

Optimal result

Integrand size = 13, antiderivative size = 16

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \frac{2 \arctan\left(\frac{bx}{a}\right)}{\sqrt{ab}}$$

output `2*arctan(b*x/a)/(a*b)^(1/2)`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \frac{2 \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `Integrate[1/(Sqrt[x]*(a + b*x)),x]`

output `(2*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/(Sqrt[a]*Sqrt[b])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x}(a+bx)} dx$$

$$\downarrow \text{73}$$

$$2 \int \frac{1}{a+bx} d\sqrt{x}$$

$$\downarrow \text{218}$$

$$\frac{2 \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `Int[1/(Sqrt[x]*(a + b*x)),x]`

output `(2*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/(Sqrt[a]*Sqrt[b])`

Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

method	result	size
derivativedivides	$\frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}}$	19
default	$\frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}}$	19

input `int(1/x^(1/2)/(b*x+a),x,method=_RETURNVERBOSE)`

output `2/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 4.25

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \left[-\frac{\sqrt{-ab} \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right)}{ab}, -\frac{2\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right)}{ab} \right]$$

input `integrate(1/x^(1/2)/(b*x+a),x, algorithm="fricas")`

output `[-sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a))/(a*b), -2*sqrt(a*b)*arctan(sqrt(a*b)/(b*sqrt(x)))/(a*b)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(14) = 28$.

Time = 0.44 (sec) , antiderivative size = 73, normalized size of antiderivative = 4.56

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \begin{cases} \frac{\infty}{\sqrt{x}} & \text{for } a = 0 \wedge b = 0 \\ \frac{2\sqrt{x}}{a} & \text{for } b = 0 \\ -\frac{2}{b\sqrt{x}} & \text{for } a = 0 \\ \frac{\log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{b\sqrt{-\frac{a}{b}}} - \frac{\log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{b\sqrt{-\frac{a}{b}}} & \text{otherwise} \end{cases}$$

input `integrate(1/x**(1/2)/(b*x+a),x)`

output `Piecewise((zoo/sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*sqrt(x)/a, Eq(b, 0)), (-2/(b*sqrt(x)), Eq(a, 0)), (log(sqrt(x) - sqrt(-a/b))/(b*sqrt(-a/b)) - log(sqrt(x) + sqrt(-a/b))/(b*sqrt(-a/b)), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/x^(1/2)/(b*x+a),x, algorithm="maxima")`

output `2*arctan(b*sqrt(x)/sqrt(a*b))/sqrt(a*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \frac{2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

input `integrate(1/x^(1/2)/(b*x+a),x, algorithm="giac")`output `2*arctan(b*sqrt(x)/sqrt(a*b))/sqrt(a*b)`**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \frac{2 \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

input `int(1/(x^(1/2)*(a + b*x)),x)`output `(2*atan((b^(1/2)*x^(1/2))/a^(1/2)))/(a^(1/2)*b^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.56

$$\int \frac{1}{\sqrt{x}(a+bx)} dx = \frac{2\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)}{ab}$$

input `int(1/x^(1/2)/(b*x+a),x)`output `(2*sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))/(a*b)`

3.110 $\int \frac{\sqrt{x}}{a+bx} dx$

Optimal result	711
Mathematica [A] (verified)	711
Rubi [A] (verified)	712
Maple [A] (verified)	713
Fricas [A] (verification not implemented)	713
Sympy [B] (verification not implemented)	714
Maxima [A] (verification not implemented)	714
Giac [A] (verification not implemented)	715
Mupad [B] (verification not implemented)	715
Reduce [B] (verification not implemented)	715

Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{\sqrt{x}}{a+bx} dx = \frac{2\sqrt{x}}{b} - \frac{2a \arctan\left(\frac{bx}{a}\right)}{b\sqrt{ab}}$$

output $2*x^{(1/2)}/b-2*a*\arctan(b*x/a)/b/(a*b)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29

$$\int \frac{\sqrt{x}}{a+bx} dx = \frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}}$$

input `Integrate[Sqrt[x]/(a + b*x), x]`

output $(2*\text{Sqrt}[x])/b - (2*\text{Sqrt}[a]*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[x])/(\text{Sqrt}[a])])/b^{(3/2)}$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {60, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{a + bx} dx$$

$$\downarrow 60$$

$$\frac{2\sqrt{x}}{b} - \frac{a \int \frac{1}{\sqrt{x}(a+bx)} dx}{b}$$

$$\downarrow 73$$

$$\frac{2\sqrt{x}}{b} - \frac{2a \int \frac{1}{a+bx} d\sqrt{x}}{b}$$

$$\downarrow 218$$

$$\frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}}$$

input `Int[Sqrt[x]/(a + b*x), x]`

output `(2*Sqrt[x])/b - (2*Sqrt[a]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/b^(3/2)`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt
[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$\frac{2\sqrt{x}}{b} - \frac{2a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b\sqrt{ab}}$	32
default	$\frac{2\sqrt{x}}{b} - \frac{2a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b\sqrt{ab}}$	32
risch	$\frac{2\sqrt{x}}{b} - \frac{2a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b\sqrt{ab}}$	32

input

```
int(x^(1/2)/(b*x+a), x, method=_RETURNVERBOSE)
```

output

```
2*x^(1/2)/b-2*a/b/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.74

$$\int \frac{\sqrt{x}}{a + bx} dx = \left[\frac{\sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx + a}\right) + 2\sqrt{x}}{b}, -\frac{2\left(\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - \sqrt{x}\right)}{b} \right]$$

input `integrate(x^(1/2)/(b*x+a),x, algorithm="fricas")`

output `[(sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*sqrt(x))/b, -2*(sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) - sqrt(x))/b]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 88 vs. $2(26) = 52$.

Time = 0.30 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.84

$$\int \frac{\sqrt{x}}{a+bx} dx = \begin{cases} \infty \sqrt{x} & \text{for } a = 0 \wedge b = 0 \\ \frac{2x^{\frac{3}{2}}}{3a} & \text{for } b = 0 \\ \frac{2\sqrt{x}}{b} & \text{for } a = 0 \\ -\frac{a \log\left(\sqrt{x} - \sqrt{-\frac{a}{b}}\right)}{b^2 \sqrt{-\frac{a}{b}}} + \frac{a \log\left(\sqrt{x} + \sqrt{-\frac{a}{b}}\right)}{b^2 \sqrt{-\frac{a}{b}}} + \frac{2\sqrt{x}}{b} & \text{otherwise} \end{cases}$$

input `integrate(x**(1/2)/(b*x+a),x)`

output `Piecewise((zoo*sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(3/2)/(3*a), Eq(b, 0)), (2*sqrt(x)/b, Eq(a, 0)), (-a*log(sqrt(x) - sqrt(-a/b))/(b**2*sqrt(-a/b)) + a*log(sqrt(x) + sqrt(-a/b))/(b**2*sqrt(-a/b)) + 2*sqrt(x)/b, True))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{x}}{a+bx} dx = -\frac{2a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}} + \frac{2\sqrt{x}}{b}$$

input `integrate(x^(1/2)/(b*x+a),x, algorithm="maxima")`

output `-2*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b) + 2*sqrt(x)/b`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{x}}{a+bx} dx = -\frac{2a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}} + \frac{2\sqrt{x}}{b}$$

input `integrate(x^(1/2)/(b*x+a),x, algorithm="giac")`output `-2*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b) + 2*sqrt(x)/b`**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

$$\int \frac{\sqrt{x}}{a+bx} dx = \frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}}$$

input `int(x^(1/2)/(a + b*x),x)`output `(2*x^(1/2))/b - (2*a^(1/2)*atan((b^(1/2)*x^(1/2))/a^(1/2)))/b^(3/2)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{x}}{a+bx} dx = \frac{-2\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) + 2\sqrt{x}b}{b^2}$$

input `int(x^(1/2)/(b*x+a),x)`output `(2*(- sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a))) + sqrt(x)*b))/b**2`

3.111 $\int \frac{x^{3/2}}{a+bx} dx$

Optimal result	716
Mathematica [A] (verified)	716
Rubi [A] (verified)	717
Maple [A] (verified)	718
Fricas [A] (verification not implemented)	719
Sympy [B] (verification not implemented)	719
Maxima [A] (verification not implemented)	720
Giac [A] (verification not implemented)	720
Mupad [B] (verification not implemented)	720
Reduce [B] (verification not implemented)	721

Optimal result

Integrand size = 13, antiderivative size = 45

$$\int \frac{x^{3/2}}{a+bx} dx = 2\sqrt{x} \left(-\frac{a}{b^2} + \frac{x}{3b} \right) + \frac{2a^2 \arctan\left(\frac{bx}{a}\right)}{b^2\sqrt{ab}}$$

output `2*x^(1/2)*(-a/b^2+1/3*x/b)+2*a^2*arctan(b*x/a)/b^2/(a*b)^(1/2)`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.09

$$\int \frac{x^{3/2}}{a+bx} dx = \frac{2\sqrt{x}(-3a+bx)}{3b^2} + \frac{2a^{3/2} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{5/2}}$$

input `Integrate[x^(3/2)/(a + b*x),x]`

output `(2*Sqrt[x]*(-3*a + b*x))/(3*b^2) + (2*a^(3/2)*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/b^(5/2)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.31, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {60, 60, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{3/2}}{a+bx} dx \\
 & \quad \downarrow 60 \\
 & \frac{2x^{3/2}}{3b} - \frac{a \int \frac{\sqrt{x}}{a+bx} dx}{b} \\
 & \quad \downarrow 60 \\
 & \frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{a \int \frac{1}{\sqrt{x}(a+bx)} dx}{b} \right)}{b} \\
 & \quad \downarrow 73 \\
 & \frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{2a \int \frac{1}{a+bx} d\sqrt{x}}{b} \right)}{b} \\
 & \quad \downarrow 218 \\
 & \frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}} \right)}{b}
 \end{aligned}$$

input `Int[x^(3/2)/(a + b*x),x]`

output `(2*x^(3/2))/(3*b) - (a*((2*sqrt[x])/b - (2*sqrt[a]*ArcTan[(sqrt[b]*sqrt[x])/sqrt[a]])/b^(3/2)))/b`

Definitions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

method	result	size
risch	$-\frac{2(-bx+3a)\sqrt{x}}{3b^2} + \frac{2a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b^2\sqrt{ab}}$	42
derivativedivides	$-\frac{2\left(-\frac{x^{\frac{3}{2}}b}{3} + \sqrt{x}a\right)}{b^2} + \frac{2a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b^2\sqrt{ab}}$	43
default	$-\frac{2\left(-\frac{x^{\frac{3}{2}}b}{3} + \sqrt{x}a\right)}{b^2} + \frac{2a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b^2\sqrt{ab}}$	43

input `int(x^(3/2)/(b*x+a), x, method=_RETURNVERBOSE)`

output
$$-2/3*(-b*x+3*a)*x^{(1/2)}/b^2+2*a^2/b^2/(a*b)^{(1/2)}*\arctan(b*x^{(1/2)}/(a*b)^{(1/2)})$$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 103, normalized size of antiderivative = 2.29

$$\int \frac{x^{3/2}}{a+bx} dx = \left[\frac{3a\sqrt{-\frac{a}{b}} \log\left(\frac{bx+2b\sqrt{x}\sqrt{-\frac{a}{b}}-a}{bx+a}\right) + 2(bx-3a)\sqrt{x}}{3b^2}, \frac{2\left(3a\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) + (bx-3a)\sqrt{x}\right)}{3b^2} \right]$$

input `integrate(x^(3/2)/(b*x+a),x, algorithm="fricas")`

output `[1/3*(3*a*sqrt(-a/b)*log((b*x + 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(b*x - 3*a)*sqrt(x))/b^2, 2/3*(3*a*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) + (b*x - 3*a)*sqrt(x))/b^2]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. $2(37) = 74$.

Time = 0.60 (sec) , antiderivative size = 107, normalized size of antiderivative = 2.38

$$\int \frac{x^{3/2}}{a+bx} dx = \begin{cases} \tilde{\infty}x^{\frac{3}{2}} & \text{for } a = 0 \wedge b = 0 \\ \frac{2x^{\frac{5}{2}}}{5a} & \text{for } b = 0 \\ \frac{2x^{\frac{3}{2}}}{3b} & \text{for } a = 0 \\ \frac{a^2 \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{b^3\sqrt{-\frac{a}{b}}} - \frac{a^2 \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{b^3\sqrt{-\frac{a}{b}}} - \frac{2a\sqrt{x}}{b^2} + \frac{2x^{\frac{3}{2}}}{3b} & \text{otherwise} \end{cases}$$

input `integrate(x**(3/2)/(b*x+a),x)`

output `Piecewise((zoo*x**(3/2), Eq(a, 0) & Eq(b, 0)), (2*x**(5/2)/(5*a), Eq(b, 0)), (2*x**(3/2)/(3*b), Eq(a, 0)), (a**2*log(sqrt(x) - sqrt(-a/b))/(b**3*sqrt(-a/b)) - a**2*log(sqrt(x) + sqrt(-a/b))/(b**3*sqrt(-a/b)) - 2*a*sqrt(x)/b**2 + 2*x**(3/2)/(3*b), True))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

$$\int \frac{x^{3/2}}{a+bx} dx = \frac{2a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}} + \frac{2\left(bx^{\frac{3}{2}} - 3a\sqrt{x}\right)}{3b^2}$$

input `integrate(x^(3/2)/(b*x+a),x, algorithm="maxima")`output `2*a^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2) + 2/3*(b*x^(3/2) - 3*a*sqrt(x))/b^2`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \frac{x^{3/2}}{a+bx} dx = \frac{2a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}} + \frac{2\left(b^2x^{\frac{3}{2}} - 3ab\sqrt{x}\right)}{3b^3}$$

input `integrate(x^(3/2)/(b*x+a),x, algorithm="giac")`output `2*a^2*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2) + 2/3*(b^2*x^(3/2) - 3*a*b*sqrt(x))/b^3`**Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.82

$$\int \frac{x^{3/2}}{a+bx} dx = \frac{2x^{3/2}}{3b} - \frac{2a\sqrt{x}}{b^2} + \frac{2a^{3/2} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{5/2}}$$

input `int(x^(3/2)/(a + b*x),x)`

output

$$\frac{(2x^{3/2})/(3b) - (2ax^{1/2})/b^2 + (2a^{3/2} \operatorname{atan}((b^{1/2}x^{1/2})/a^{1/2}))/b^{5/2}}$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int \frac{x^{3/2}}{a+bx} dx = \frac{2\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a - 2\sqrt{x}ab + \frac{2\sqrt{x}b^2x}{3}}{b^3}$$

input

```
int(x^(3/2)/(b*x+a),x)
```

output

```
(2*(3*sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a - 3*sqrt(x)*a*
b + sqrt(x)*b**2*x))/(3*b**3)
```

3.112 $\int \frac{x^{5/2}}{a+bx} dx$

Optimal result	722
Mathematica [A] (verified)	722
Rubi [A] (verified)	723
Maple [A] (verified)	724
Fricas [A] (verification not implemented)	725
Sympy [B] (verification not implemented)	725
Maxima [A] (verification not implemented)	726
Giac [A] (verification not implemented)	727
Mupad [B] (verification not implemented)	727
Reduce [B] (verification not implemented)	727

Optimal result

Integrand size = 13, antiderivative size = 57

$$\int \frac{x^{5/2}}{a+bx} dx = 2\sqrt{x} \left(\frac{a^2}{b^2} - \frac{ax}{3b^2} + \frac{x^2}{5b} \right) - \frac{2a^3 \arctan\left(\frac{bx}{a}\right)}{b^3\sqrt{ab}}$$

output $2*x^{(1/2)}*(a^2/b^2-1/3*a*x/b^2+1/5*x^2/b)-2*a^3*\arctan(b*x/a)/b^3/(a*b)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.07

$$\int \frac{x^{5/2}}{a+bx} dx = \frac{2\sqrt{x}(15a^2 - 5abx + 3b^2x^2)}{15b^3} - \frac{2a^{5/2} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{7/2}}$$

input `Integrate[x^(5/2)/(a + b*x),x]`

output $(2*\text{Sqrt}[x]*(15*a^2 - 5*a*b*x + 3*b^2*x^2))/(15*b^3) - (2*a^{(5/2)}*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[x])/ \text{Sqrt}[a]])/b^{(7/2)}$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.37, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {60, 60, 60, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{5/2}}{a+bx} dx \\
 & \quad \downarrow 60 \\
 & \frac{2x^{5/2}}{5b} - \frac{a \int \frac{x^{3/2}}{a+bx} dx}{b} \\
 & \quad \downarrow 60 \\
 & \frac{2x^{5/2}}{5b} - \frac{a \left(\frac{2x^{3/2}}{3b} - \frac{a \int \frac{\sqrt{x}}{a+bx} dx}{b} \right)}{b} \\
 & \quad \downarrow 60 \\
 & \frac{2x^{5/2}}{5b} - \frac{a \left(\frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{a \int \frac{1}{\sqrt{x}(a+bx)} dx}{b} \right)}{b} \right)}{b} \\
 & \quad \downarrow 73 \\
 & \frac{2x^{5/2}}{5b} - \frac{a \left(\frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{2a \int \frac{1}{a+bx} d\sqrt{x}}{b} \right)}{b} \right)}{b} \\
 & \quad \downarrow 218 \\
 & \frac{2x^{5/2}}{5b} - \frac{a \left(\frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}} \right)}{b} \right)}{b}
 \end{aligned}$$

input `Int[x^(5/2)/(a + b*x), x]`

output `(2*x^(5/2))/(5*b) - (a*((2*x^(3/2))/(3*b) - (a*((2*Sqrt[x])/b - (2*Sqrt[a]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/b^(3/2)))/b)/b`

Defintions of rubi rules used

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.93

method	result	size
risch	$\frac{2(3b^2x^2 - 5bax + 15a^2)\sqrt{x}}{15b^3} - \frac{2a^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b^3\sqrt{ab}}$	53
derivativedivides	$\frac{\frac{2x}{5}b^2 - \frac{2ba}{3}x^{\frac{3}{2}} + 2\sqrt{x}a^2}{b^3} - \frac{2a^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b^3\sqrt{ab}}$	54
default	$\frac{\frac{2x}{5}b^2 - \frac{2ba}{3}x^{\frac{3}{2}} + 2\sqrt{x}a^2}{b^3} - \frac{2a^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b^3\sqrt{ab}}$	54

input `int(x^(5/2)/(b*x+a),x,method=_RETURNVERBOSE)`

output $\frac{2}{15} \cdot (3b^2x^2 - 5abx + 15a^2) \cdot x^{1/2} / b^3 - 2a^3 / b^3 / (ab)^{1/2} \cdot \arctan(b \cdot x^{1/2} / (ab)^{1/2})$

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.32

$$\int \frac{x^{5/2}}{a+bx} dx = \left[\frac{15a^2 \sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx+a}\right) + 2(3b^2x^2 - 5abx + 15a^2)\sqrt{x}}{15b^3}, \right. \\ \left. - \frac{2\left(15a^2 \sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - (3b^2x^2 - 5abx + 15a^2)\sqrt{x}\right)}{15b^3} \right]$$

input `integrate(x^(5/2)/(b*x+a),x, algorithm="fricas")`

output `[1/15*(15*a^2*sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(3*b^2*x^2 - 5*a*b*x + 15*a^2)*sqrt(x))/b^3, -2/15*(15*a^2*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) - (3*b^2*x^2 - 5*a*b*x + 15*a^2)*sqrt(x))/b^3]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 122 vs. $2(49) = 98$.

Time = 2.02 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.14

$$\int \frac{x^{5/2}}{a+bx} dx = \begin{cases} \tilde{\infty} x^{5/2} & \text{for } a = 0 \wedge b = 0 \\ \frac{2x^{7/2}}{7a} & \text{for } b = 0 \\ \frac{2x^{5/2}}{5b} & \text{for } a = 0 \\ -\frac{a^3 \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{b^4 \sqrt{-\frac{a}{b}}} + \frac{a^3 \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{b^4 \sqrt{-\frac{a}{b}}} + \frac{2a^2 \sqrt{x}}{b^3} - \frac{2ax^{3/2}}{3b^2} + \frac{2x^{5/2}}{5b} & \text{otherwise} \end{cases}$$

input `integrate(x**(5/2)/(b*x+a),x)`

output `Piecewise((zoo*x**(5/2), Eq(a, 0) & Eq(b, 0)), (2*x**(7/2)/(7*a), Eq(b, 0)), (2*x**(5/2)/(5*b), Eq(a, 0)), (-a**3*log(sqrt(x) - sqrt(-a/b))/(b**4*sqrt(-a/b)) + a**3*log(sqrt(x) + sqrt(-a/b))/(b**4*sqrt(-a/b)) + 2*a**2*sqrt(x)/b**3 - 2*a*x**(3/2)/(3*b**2) + 2*x**(5/2)/(5*b), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.95

$$\int \frac{x^{5/2}}{a+bx} dx = -\frac{2a^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^3}} + \frac{2\left(3b^2x^{5/2} - 5abx^{3/2} + 15a^2\sqrt{x}\right)}{15b^3}$$

input `integrate(x^(5/2)/(b*x+a),x, algorithm="maxima")`

output `-2*a^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 2/15*(3*b^2*x^(5/2) - 5*a*b*x^(3/2) + 15*a^2*sqrt(x))/b^3`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.04

$$\int \frac{x^{5/2}}{a+bx} dx = -\frac{2a^3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^3}} + \frac{2\left(3b^4x^{5/2} - 5ab^3x^{3/2} + 15a^2b^2\sqrt{x}\right)}{15b^5}$$

input `integrate(x^(5/2)/(b*x+a),x, algorithm="giac")`

output `-2*a^3*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 2/15*(3*b^4*x^(5/2) - 5*a*b^3*x^(3/2) + 15*a^2*b^2*sqrt(x))/b^5`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.84

$$\int \frac{x^{5/2}}{a+bx} dx = \frac{2x^{5/2}}{5b} - \frac{2ax^{3/2}}{3b^2} + \frac{2a^2\sqrt{x}}{b^3} - \frac{2a^{5/2} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{7/2}}$$

input `int(x^(5/2)/(a + b*x),x)`

output `(2*x^(5/2))/(5*b) - (2*a*x^(3/2))/(3*b^2) + (2*a^2*x^(1/2))/b^3 - (2*a^(5/2)*atan((b^(1/2)*x^(1/2))/a^(1/2)))/b^(7/2)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.96

$$\int \frac{x^{5/2}}{a+bx} dx = \frac{-2\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a^2 + 2\sqrt{x} a^2 b - \frac{2\sqrt{x} a b^2 x}{3} + \frac{2\sqrt{x} b^3 x^2}{5}}{b^4}$$

input `int(x^(5/2)/(b*x+a),x)`

output

```
(2*( - 15*sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a**2 + 15*sqrt(x)*a**2*b - 5*sqrt(x)*a*b**2*x + 3*sqrt(x)*b**3*x**2))/(15*b**4)
```

3.113 $\int \frac{1}{\sqrt{x}(a+bx)^2} dx$

Optimal result	729
Mathematica [A] (verified)	729
Rubi [A] (verified)	730
Maple [A] (verified)	731
Fricas [A] (verification not implemented)	731
Sympy [B] (verification not implemented)	732
Maxima [A] (verification not implemented)	733
Giac [A] (verification not implemented)	733
Mupad [B] (verification not implemented)	733
Reduce [B] (verification not implemented)	734

Optimal result

Integrand size = 13, antiderivative size = 30

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx = \frac{\sqrt{x} \arctan\left(\frac{bx}{a}\right)}{a^2 \sqrt{ab}(a+bx)}$$

output $x^{(1/2)}*\arctan(b*x/a)/a^2/(a*b)^{(1/2)/(b*x+a)}$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx = \frac{\sqrt{x}}{a(a+bx)} + \frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{a^{3/2}\sqrt{b}}$$

input `Integrate[1/(Sqrt[x]*(a + b*x)^2),x]`

output `Sqrt[x]/(a*(a + b*x)) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]]/(a^(3/2)*Sqrt[b])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.50, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {52, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx$$

$$\downarrow 52$$

$$\frac{\int \frac{1}{\sqrt{x}(a+bx)} dx}{2a} + \frac{\sqrt{x}}{a(a+bx)}$$

$$\downarrow 73$$

$$\frac{\int \frac{1}{a+bx} d\sqrt{x}}{a} + \frac{\sqrt{x}}{a(a+bx)}$$

$$\downarrow 218$$

$$\frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{a^{3/2}\sqrt{b}} + \frac{\sqrt{x}}{a(a+bx)}$$

input `Int[1/(Sqrt[x]*(a + b*x)^2), x]`

output `Sqrt[x]/(a*(a + b*x)) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]]/(a^(3/2)*Sqrt[b])`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt
[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$\frac{\sqrt{x}}{a(bx+a)} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{a\sqrt{ab}}$	36
default	$\frac{\sqrt{x}}{a(bx+a)} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{a\sqrt{ab}}$	36

input

```
int(1/x^(1/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)
```

output

```
x^(1/2)/a/(b*x+a)+1/a/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.87

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx$$

$$= \left[\frac{2ab\sqrt{x} - \sqrt{-ab}(bx+a) \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right)}{2(a^2b^2x+a^3b)}, \frac{ab\sqrt{x} - \sqrt{ab}(bx+a) \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right)}{a^2b^2x+a^3b} \right]$$

input

```
integrate(1/x^(1/2)/(b*x+a)^2,x, algorithm="fricas")
```

output

```
[1/2*(2*a*b*sqrt(x) - sqrt(-a*b)*(b*x + a)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a)))/(a^2*b^2*x + a^3*b), (a*b*sqrt(x) - sqrt(a*b)*(b*x + a)*arctan(sqrt(a*b)/(b*sqrt(x))))/(a^2*b^2*x + a^3*b)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 277 vs. $2(26) = 52$.

Time = 2.50 (sec) , antiderivative size = 277, normalized size of antiderivative = 9.23

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx$$

$$= \begin{cases} \frac{\infty}{x^{\frac{3}{2}}} \\ \frac{2\sqrt{x}}{a^2} \\ -\frac{2}{3b^2x^{\frac{3}{2}}} \\ \frac{a \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{2a^2b\sqrt{-\frac{a}{b}}+2ab^2x\sqrt{-\frac{a}{b}}} - \frac{a \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{2a^2b\sqrt{-\frac{a}{b}}+2ab^2x\sqrt{-\frac{a}{b}}} + \frac{2b\sqrt{x}\sqrt{-\frac{a}{b}}}{2a^2b\sqrt{-\frac{a}{b}}+2ab^2x\sqrt{-\frac{a}{b}}} + \frac{bx \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{2a^2b\sqrt{-\frac{a}{b}}+2ab^2x\sqrt{-\frac{a}{b}}} - \frac{bx \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{2a^2b\sqrt{-\frac{a}{b}}+2ab^2x\sqrt{-\frac{a}{b}}} \end{cases}$$

input

```
integrate(1/x**(1/2)/(b*x+a)**2,x)
```

output

```
Piecewise((zoo/x**(3/2), Eq(a, 0) & Eq(b, 0)), (2*sqrt(x)/a**2, Eq(b, 0)), (-2/(3*b**2*x**(3/2)), Eq(a, 0)), (a*log(sqrt(x) - sqrt(-a/b))/(2*a**2*b*sqrt(-a/b) + 2*a*b**2*x*sqrt(-a/b)) - a*log(sqrt(x) + sqrt(-a/b))/(2*a**2*b*sqrt(-a/b) + 2*a*b**2*x*sqrt(-a/b)) + 2*b*sqrt(x)*sqrt(-a/b)/(2*a**2*b*sqrt(-a/b) + 2*a*b**2*x*sqrt(-a/b)) + b*x*log(sqrt(x) - sqrt(-a/b))/(2*a**2*b*sqrt(-a/b) + 2*a*b**2*x*sqrt(-a/b)) - b*x*log(sqrt(x) + sqrt(-a/b))/(2*a**2*b*sqrt(-a/b) + 2*a*b**2*x*sqrt(-a/b)), True))
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx = \frac{\sqrt{x}}{abx+a^2} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba}}$$

input `integrate(1/x^(1/2)/(b*x+a)^2,x, algorithm="maxima")`output `sqrt(x)/(a*b*x + a^2) + arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx = \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{aba}} + \frac{\sqrt{x}}{(bx+a)a}$$

input `integrate(1/x^(1/2)/(b*x+a)^2,x, algorithm="giac")`output `arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a) + sqrt(x)/((b*x + a)*a)`**Mupad [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.10

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx = \frac{\sqrt{x}}{a(a+bx)} + \frac{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{a^{3/2}\sqrt{b}}$$

input `int(1/(x^(1/2)*(a + b*x)^2),x)`output `x^(1/2)/(a*(a + b*x)) + atan((b^(1/2)*x^(1/2))/a^(1/2))/(a^(3/2)*b^(1/2))`

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.97

$$\int \frac{1}{\sqrt{x}(a+bx)^2} dx = \frac{\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a + \sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) bx + \sqrt{x} ab}{a^2 b (bx + a)}$$

input `int(1/x^(1/2)/(b*x+a)^2,x)`output `(sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a + sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*b*x + sqrt(x)*a*b)/(a**2*b*(a + b*x))`

3.114 $\int \frac{\sqrt{x}}{(a+bx)^2} dx$

Optimal result	735
Mathematica [A] (verified)	735
Rubi [A] (verified)	736
Maple [A] (verified)	737
Fricas [A] (verification not implemented)	737
Sympy [B] (verification not implemented)	738
Maxima [A] (verification not implemented)	739
Giac [A] (verification not implemented)	739
Mupad [B] (verification not implemented)	739
Reduce [B] (verification not implemented)	740

Optimal result

Integrand size = 13, antiderivative size = 31

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx = -\frac{\sqrt{x} \arctan\left(\frac{bx}{a}\right)}{b^2 \sqrt{ab}(a+bx)}$$

output `-x^(1/2)*arctan(b*x/a)/b^2/(a*b)^(1/2)/(b*x+a)`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.48

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx = -\frac{\sqrt{x}}{b(a+bx)} + \frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{ab}^{3/2}}$$

input `Integrate[Sqrt[x]/(a + b*x)^2,x]`

output `-(Sqrt[x]/(b*(a + b*x))) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]]/(Sqrt[a]*b^(3/2))`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.48, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {51, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx$$

$$\downarrow \text{51}$$

$$\frac{\int \frac{1}{\sqrt{x}(a+bx)} dx}{2b} - \frac{\sqrt{x}}{b(a+bx)}$$

$$\downarrow \text{73}$$

$$\frac{\int \frac{1}{a+bx} d\sqrt{x}}{b} - \frac{\sqrt{x}}{b(a+bx)}$$

$$\downarrow \text{218}$$

$$\frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{a}b^{3/2}} - \frac{\sqrt{x}}{b(a+bx)}$$

input `Int[Sqrt[x]/(a + b*x)^2,x]`

output `-(Sqrt[x]/(b*(a + b*x))) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]]/(Sqrt[a]*b^(3/2))`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
 t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

method	result	size
derivativedivides	$-\frac{\sqrt{x}}{b(bx+a)} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b\sqrt{ab}}$	37
default	$-\frac{\sqrt{x}}{b(bx+a)} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b\sqrt{ab}}$	37

input `int(x^(1/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `-x^(1/2)/b/(b*x+a)+1/b/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 115, normalized size of antiderivative = 3.71

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx = \left[-\frac{2ab\sqrt{x} + \sqrt{-ab}(bx+a) \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right)}{2(ab^3x+a^2b^2)}, \right. \\ \left. -\frac{ab\sqrt{x} + \sqrt{ab}(bx+a) \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right)}{ab^3x+a^2b^2} \right]$$

input `integrate(x^(1/2)/(b*x+a)^2,x, algorithm="fricas")`

output

```
[-1/2*(2*a*b*sqrt(x) + sqrt(-a*b)*(b*x + a)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a)))/(a*b^3*x + a^2*b^2), -(a*b*sqrt(x) + sqrt(a*b)*(b*x + a)*arctan(sqrt(a*b)/(b*sqrt(x))))/(a*b^3*x + a^2*b^2)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 269 vs. $2(27) = 54$.

Time = 1.63 (sec) , antiderivative size = 269, normalized size of antiderivative = 8.68

$$\int \frac{\sqrt{x}}{(a + bx)^2} dx$$

$$= \begin{cases} \frac{\infty}{\sqrt{x}} \\ \frac{2x^{\frac{3}{2}}}{3a^2} \\ -\frac{2}{b^2\sqrt{x}} \\ \frac{a \log\left(\sqrt{x} - \sqrt{-\frac{a}{b}}\right)}{2ab^2\sqrt{-\frac{a}{b}} + 2b^3x\sqrt{-\frac{a}{b}}} - \frac{a \log\left(\sqrt{x} + \sqrt{-\frac{a}{b}}\right)}{2ab^2\sqrt{-\frac{a}{b}} + 2b^3x\sqrt{-\frac{a}{b}}} - \frac{2b\sqrt{x}\sqrt{-\frac{a}{b}}}{2ab^2\sqrt{-\frac{a}{b}} + 2b^3x\sqrt{-\frac{a}{b}}} + \frac{bx \log\left(\sqrt{x} - \sqrt{-\frac{a}{b}}\right)}{2ab^2\sqrt{-\frac{a}{b}} + 2b^3x\sqrt{-\frac{a}{b}}} - \frac{bx \log\left(\sqrt{x} + \sqrt{-\frac{a}{b}}\right)}{2ab^2\sqrt{-\frac{a}{b}} + 2b^3x\sqrt{-\frac{a}{b}}} \end{cases}$$

input

```
integrate(x**(1/2)/(b*x+a)**2,x)
```

output

```
Piecewise((zoo/sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(3/2)/(3*a**2), Eq(b, 0)), (-2/(b**2*sqrt(x)), Eq(a, 0)), (a*log(sqrt(x) - sqrt(-a/b))/(2*a*b**2*sqrt(-a/b) + 2*b**3*x*sqrt(-a/b)) - a*log(sqrt(x) + sqrt(-a/b))/(2*a*b**2*sqrt(-a/b) + 2*b**3*x*sqrt(-a/b)) - 2*b*sqrt(x)*sqrt(-a/b)/(2*a*b**2*sqrt(-a/b) + 2*b**3*x*sqrt(-a/b)) + b*x*log(sqrt(x) - sqrt(-a/b))/(2*a*b**2*sqrt(-a/b) + 2*b**3*x*sqrt(-a/b)) - b*x*log(sqrt(x) + sqrt(-a/b))/(2*a*b**2*sqrt(-a/b) + 2*b**3*x*sqrt(-a/b)), True))
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.19

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx = -\frac{\sqrt{x}}{b^2x+ab} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}}$$

input `integrate(x^(1/2)/(b*x+a)^2,x, algorithm="maxima")`output `-sqrt(x)/(b^2*x + a*b) + arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.16

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx = \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb}} - \frac{\sqrt{x}}{(bx+a)b}$$

input `integrate(x^(1/2)/(b*x+a)^2,x, algorithm="giac")`output `arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b) - sqrt(x)/((b*x + a)*b)`**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.10

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{a}b^{3/2}} - \frac{\sqrt{x}}{b(a+bx)}$$

input `int(x^(1/2)/(a + b*x)^2,x)`output `atan((b^(1/2)*x^(1/2))/a^(1/2))/(a^(1/2)*b^(3/2)) - x^(1/2)/(b*(a + b*x))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.94

$$\int \frac{\sqrt{x}}{(a+bx)^2} dx = \frac{\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a + \sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) bx - \sqrt{x} ab}{a b^2 (bx + a)}$$

input `int(x^(1/2)/(b*x+a)^2,x)`

output `(sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a + sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*b*x - sqrt(x)*a*b)/(a*b**2*(a + b*x))`

3.115 $\int \frac{x^{3/2}}{(a+bx)^2} dx$

Optimal result	741
Mathematica [A] (verified)	741
Rubi [A] (verified)	742
Maple [A] (verified)	743
Fricas [A] (verification not implemented)	744
Sympy [B] (verification not implemented)	744
Maxima [A] (verification not implemented)	745
Giac [A] (verification not implemented)	745
Mupad [B] (verification not implemented)	746
Reduce [B] (verification not implemented)	746

Optimal result

Integrand size = 13, antiderivative size = 50

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = \frac{2x^{3/2}}{b(a+bx)} + \frac{3a\sqrt{x} \arctan\left(\frac{bx}{a}\right)}{b^3\sqrt{ab}(a+bx)}$$

output `2*x^(3/2)/b/(b*x+a)+3*a*x^(1/2)*arctan(b*x/a)/b^3/(a*b)^(1/2)/(b*x+a)`

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.08

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = \frac{\sqrt{x}(3a+2bx)}{b^2(a+bx)} - \frac{3\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{5/2}}$$

input `Integrate[x^(3/2)/(a + b*x)^2,x]`

output `(Sqrt[x]*(3*a + 2*b*x))/(b^2*(a + b*x)) - (3*Sqrt[a]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/b^(5/2)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.30, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {51, 60, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{3/2}}{(a+bx)^2} dx \\
 & \quad \downarrow \text{51} \\
 & \frac{3 \int \frac{\sqrt{x}}{a+bx} dx}{2b} - \frac{x^{3/2}}{b(a+bx)} \\
 & \quad \downarrow \text{60} \\
 & \frac{3 \left(\frac{2\sqrt{x}}{b} - \frac{a \int \frac{1}{\sqrt{x}(a+bx)} dx \right)}{2b} - \frac{x^{3/2}}{b(a+bx)} \\
 & \quad \downarrow \text{73} \\
 & \frac{3 \left(\frac{2\sqrt{x}}{b} - \frac{2a \int \frac{1}{a+bx} d\sqrt{x} \right)}{2b} - \frac{x^{3/2}}{b(a+bx)} \\
 & \quad \downarrow \text{218} \\
 & \frac{3 \left(\frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}} \right)}{2b} - \frac{x^{3/2}}{b(a+bx)}
 \end{aligned}$$

input `Int[x^(3/2)/(a + b*x)^2,x]`

output `-(x^(3/2)/(b*(a + b*x))) + (3*((2*sqrt[x])/b - (2*sqrt[a]*ArcTan[(sqrt[b]*sqrt[x])/sqrt[a]])/b^(3/2)))/(2*b)`

Defintions of rubi rules used

```
rule 51 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]
```

```
rule 60 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 218 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{2\sqrt{x}}{b^2} - \frac{2a \left(-\frac{\sqrt{x}}{2(bx+a)} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{b^2}$	47
default	$\frac{2\sqrt{x}}{b^2} - \frac{2a \left(-\frac{\sqrt{x}}{2(bx+a)} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{b^2}$	47
risch	$\frac{2\sqrt{x}}{b^2} + \frac{a\sqrt{x}}{b^2(bx+a)} - \frac{3a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{b^2\sqrt{ab}}$	47

input `int(x^(3/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `2*x^(1/2)/b^2-2*a/b^2*(-1/2*x^(1/2)/(b*x+a)+3/2/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.68

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = \left[\frac{3(bx+a)\sqrt{-\frac{a}{b}} \log\left(\frac{bx-2b\sqrt{x}\sqrt{-\frac{a}{b}}-a}{bx+a}\right) + 2(2bx+3a)\sqrt{x}}{2(b^3x+ab^2)}, \right. \\ \left. - \frac{3(bx+a)\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - (2bx+3a)\sqrt{x}}{b^3x+ab^2} \right]$$

input `integrate(x^(3/2)/(b*x+a)^2,x, algorithm="fricas")`

output `[1/2*(3*(b*x + a)*sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(2*b*x + 3*a)*sqrt(x))/(b^3*x + a*b^2), -(3*(b*x + a)*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) - (2*b*x + 3*a)*sqrt(x))/(b^3*x + a*b^2)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. 2(42) = 84.

Time = 3.18 (sec) , antiderivative size = 332, normalized size of antiderivative = 6.64

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = \begin{cases} \infty\sqrt{x} \\ \frac{2x^{\frac{5}{2}}}{5a^2} \\ \frac{2\sqrt{x}}{b^2} \\ -\frac{3a^2 \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{2ab^3\sqrt{-\frac{a}{b}}+2b^4x\sqrt{-\frac{a}{b}}} + \frac{3a^2 \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{2ab^3\sqrt{-\frac{a}{b}}+2b^4x\sqrt{-\frac{a}{b}}} + \frac{6ab\sqrt{x}\sqrt{-\frac{a}{b}}}{2ab^3\sqrt{-\frac{a}{b}}+2b^4x\sqrt{-\frac{a}{b}}} - \frac{3abx \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{2ab^3\sqrt{-\frac{a}{b}}+2b^4x\sqrt{-\frac{a}{b}}} + \frac{3abx \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{2ab^3\sqrt{-\frac{a}{b}}+2b^4x\sqrt{-\frac{a}{b}}} \end{cases}$$

input `integrate(x**(3/2)/(b*x+a)**2,x)`

output `Piecewise((zoo*sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(5/2)/(5*a**2), Eq(b, 0)), (2*sqrt(x)/b**2, Eq(a, 0)), (-3*a**2*log(sqrt(x) - sqrt(-a/b))/(2*a*b**3*sqrt(-a/b) + 2*b**4*x*sqrt(-a/b)) + 3*a**2*log(sqrt(x) + sqrt(-a/b))/(2*a*b**3*sqrt(-a/b) + 2*b**4*x*sqrt(-a/b)) + 6*a*b*sqrt(x)*sqrt(-a/b)/(2*a*b**3*sqrt(-a/b) + 2*b**4*x*sqrt(-a/b)) - 3*a*b*x*log(sqrt(x) - sqrt(-a/b))/(2*a*b**3*sqrt(-a/b) + 2*b**4*x*sqrt(-a/b)) + 3*a*b*x*log(sqrt(x) + sqrt(-a/b))/(2*a*b**3*sqrt(-a/b) + 2*b**4*x*sqrt(-a/b)) + 4*b**2*x**(3/2)*sqrt(-a/b)/(2*a*b**3*sqrt(-a/b) + 2*b**4*x*sqrt(-a/b)), True))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.98

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = \frac{a\sqrt{x}}{b^3x+ab^2} - \frac{3a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}} + \frac{2\sqrt{x}}{b^2}$$

input `integrate(x^(3/2)/(b*x+a)^2,x, algorithm="maxima")`

output `a*sqrt(x)/(b^3*x + a*b^2) - 3*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2) + 2*sqrt(x)/b^2`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.92

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = -\frac{3a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^2}} + \frac{a\sqrt{x}}{(bx+a)b^2} + \frac{2\sqrt{x}}{b^2}$$

input `integrate(x^(3/2)/(b*x+a)^2,x, algorithm="giac")`

output `-3*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2) + a*sqrt(x)/((b*x + a)*b^2) + 2*sqrt(x)/b^2`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.92

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = \frac{2\sqrt{x}}{b^2} + \frac{a\sqrt{x}}{xb^3+ab^2} - \frac{3\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{5/2}}$$

input `int(x^(3/2)/(a + b*x)^2,x)`output `(2*x^(1/2))/b^2 + (a*x^(1/2))/(a*b^2 + b^3*x) - (3*a^(1/2)*atan((b^(1/2)*x^(1/2))/a^(1/2)))/b^(5/2)`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.34

$$\int \frac{x^{3/2}}{(a+bx)^2} dx = \frac{-3\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)a - 3\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)bx + 3\sqrt{x}ab + 2\sqrt{x}b^2x}{b^3(bx+a)}$$

input `int(x^(3/2)/(b*x+a)^2,x)`output `(- 3*sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a - 3*sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*b*x + 3*sqrt(x)*a*b + 2*sqrt(x)*b**2*x)/(b**3*(a + b*x))`

3.116 $\int \frac{x^{5/2}}{(a+bx)^2} dx$

Optimal result	747
Mathematica [A] (verified)	747
Rubi [A] (verified)	748
Maple [A] (verified)	750
Fricas [A] (verification not implemented)	750
Sympy [B] (verification not implemented)	751
Maxima [A] (verification not implemented)	751
Giac [A] (verification not implemented)	752
Mupad [B] (verification not implemented)	752
Reduce [B] (verification not implemented)	753

Optimal result

Integrand size = 13, antiderivative size = 69

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = \frac{2\sqrt{x}\left(-\frac{5ax}{3b^2} + \frac{x^2}{3b}\right)}{a+bx} - \frac{5a^2\sqrt{x} \arctan\left(\frac{bx}{a}\right)}{b^4\sqrt{ab}(a+bx)}$$

output

$$\frac{2x^{1/2}(-5/3ax/b^2+1/3x^2/b)/(bx+a)-5a^2x^{1/2}\arctan(bx/a)/b^4}{(ab)^{1/2}/(bx+a)}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.99

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = \frac{\sqrt{x}(-15a^2 - 10abx + 2b^2x^2)}{3b^3(a+bx)} + \frac{5a^{3/2} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{7/2}}$$

input

`Integrate[x^(5/2)/(a + b*x)^2,x]`

output

`(Sqrt[x]*(-15*a^2 - 10*a*b*x + 2*b^2*x^2))/(3*b^3*(a + b*x)) + (5*a^(3/2)*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/b^(7/2)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.22, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {51, 60, 60, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{5/2}}{(a+bx)^2} dx \\
 & \quad \downarrow \text{51} \\
 & \frac{5 \int \frac{x^{3/2}}{a+bx} dx}{2b} - \frac{x^{5/2}}{b(a+bx)} \\
 & \quad \downarrow \text{60} \\
 & \frac{5 \left(\frac{2x^{3/2}}{3b} - \frac{a \int \frac{\sqrt{x}}{a+bx} dx}{b} \right)}{2b} - \frac{x^{5/2}}{b(a+bx)} \\
 & \quad \downarrow \text{60} \\
 & \frac{5 \left(\frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{a \int \frac{1}{\sqrt{x}(a+bx)} dx}{b} \right)}{b} \right)}{2b} - \frac{x^{5/2}}{b(a+bx)} \\
 & \quad \downarrow \text{73} \\
 & \frac{5 \left(\frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{2a \int \frac{1}{a+bx} d\sqrt{x}}{b} \right)}{b} \right)}{2b} - \frac{x^{5/2}}{b(a+bx)} \\
 & \quad \downarrow \text{218} \\
 & \frac{5 \left(\frac{2x^{3/2}}{3b} - \frac{a \left(\frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}} \right)}{b} \right)}{2b} - \frac{x^{5/2}}{b(a+bx)}
 \end{aligned}$$

input `Int[x^(5/2)/(a + b*x)^2,x]`

output `-(x^(5/2)/(b*(a + b*x))) + (5*((2*x^(3/2))/(3*b) - (a*((2*Sqrt[x])/b - (2*Sqrt[a]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/b^(3/2)))/b)/(2*b)`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(b*(m + n + 1)))] Int[(a + b*x)^(m)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.81

method	result	size
risch	$-\frac{2(-bx+6a)\sqrt{x}}{3b^3} + \frac{a^2 \left(-\frac{\sqrt{x}}{bx+a} + \frac{5 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}} \right)}{b^3}$	56
derivativedivides	$-\frac{2 \left(-\frac{x}{3} \frac{2}{b} + 2\sqrt{x} a \right)}{b^3} + \frac{2a^2 \left(-\frac{\sqrt{x}}{2(bx+a)} + \frac{5 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{b^3}$	59
default	$-\frac{2 \left(-\frac{x}{3} \frac{2}{b} + 2\sqrt{x} a \right)}{b^3} + \frac{2a^2 \left(-\frac{\sqrt{x}}{2(bx+a)} + \frac{5 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{2\sqrt{ab}} \right)}{b^3}$	59

input `int(x^(5/2)/(b*x+a)^2,x,method=_RETURNVERBOSE)`

output `-2/3*(-b*x+6*a)*x^(1/2)/b^3+a^2/b^3*(-x^(1/2)/(b*x+a)+5/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 161, normalized size of antiderivative = 2.33

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = \left[\frac{15(abx+a^2)\sqrt{-\frac{a}{b}} \log\left(\frac{bx+2b\sqrt{x}\sqrt{-\frac{a}{b}}-a}{bx+a}\right) + 2(2b^2x^2-10abx-15a^2)\sqrt{x}}{6(b^4x+ab^3)}, \frac{15(abx+a^2)\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) + (2b^2x^2-10abx-15a^2)\sqrt{x}}{6(b^4x+ab^3)} \right]$$

input `integrate(x^(5/2)/(b*x+a)^2,x, algorithm="fricas")`

output `[1/6*(15*(a*b*x + a^2)*sqrt(-a/b)*log((b*x + 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(2*b^2*x^2 - 10*a*b*x - 15*a^2)*sqrt(x))/(b^4*x + a*b^3), 1/3*(15*(a*b*x + a^2)*sqrt(a/b)*arctan(b*sqrt(x)*sqrt(a/b)/a) + (2*b^2*x^2 - 10*a*b*x - 15*a^2)*sqrt(x))/(b^4*x + a*b^3)]`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 389 vs. $2(60) = 120$.

Time = 8.30 (sec) , antiderivative size = 389, normalized size of antiderivative = 5.64

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = \begin{cases} \tilde{\infty} x^{\frac{3}{2}} \\ \frac{2x^{\frac{7}{2}}}{7a^2} \\ \frac{2x^{\frac{3}{2}}}{3b^2} \\ \frac{15a^3 \log(\sqrt{x} - \sqrt{-\frac{a}{b}})}{6ab^4 \sqrt{-\frac{a}{b}} + 6b^5 x \sqrt{-\frac{a}{b}}} - \frac{15a^3 \log(\sqrt{x} + \sqrt{-\frac{a}{b}})}{6ab^4 \sqrt{-\frac{a}{b}} + 6b^5 x \sqrt{-\frac{a}{b}}} - \frac{30a^2 b \sqrt{x} \sqrt{-\frac{a}{b}}}{6ab^4 \sqrt{-\frac{a}{b}} + 6b^5 x \sqrt{-\frac{a}{b}}} + \frac{15a^2 b x \log(\sqrt{x} - \sqrt{-\frac{a}{b}})}{6ab^4 \sqrt{-\frac{a}{b}} + 6b^5 x \sqrt{-\frac{a}{b}}} - \frac{15a^2 b x \log(\sqrt{x} + \sqrt{-\frac{a}{b}})}{6ab^4 \sqrt{-\frac{a}{b}} + 6b^5 x \sqrt{-\frac{a}{b}}} \end{cases}$$

input `integrate(x**(5/2)/(b*x+a)**2,x)`

output `Piecewise((zoo*x**(3/2), Eq(a, 0) & Eq(b, 0)), (2*x**(7/2)/(7*a**2), Eq(b, 0)), (2*x**(3/2)/(3*b**2), Eq(a, 0)), (15*a**3*log(sqrt(x) - sqrt(-a/b))/(6*a*b**4*sqrt(-a/b) + 6*b**5*x*sqrt(-a/b)) - 15*a**3*log(sqrt(x) + sqrt(-a/b))/(6*a*b**4*sqrt(-a/b) + 6*b**5*x*sqrt(-a/b)) - 30*a**2*b*sqrt(x)*sqrt(-a/b)/(6*a*b**4*sqrt(-a/b) + 6*b**5*x*sqrt(-a/b)) + 15*a**2*b*x*log(sqrt(x) - sqrt(-a/b))/(6*a*b**4*sqrt(-a/b) + 6*b**5*x*sqrt(-a/b)) - 15*a**2*b*x*log(sqrt(x) + sqrt(-a/b))/(6*a*b**4*sqrt(-a/b) + 6*b**5*x*sqrt(-a/b)) - 20*a*b**2*x**(3/2)*sqrt(-a/b)/(6*a*b**4*sqrt(-a/b) + 6*b**5*x*sqrt(-a/b)) + 4*b**3*x**(5/2)*sqrt(-a/b)/(6*a*b**4*sqrt(-a/b) + 6*b**5*x*sqrt(-a/b)), True))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.91

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = -\frac{a^2 \sqrt{x}}{b^4 x + ab^3} + \frac{5a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{abb^3}} + \frac{2\left(bx^{\frac{3}{2}} - 6a\sqrt{x}\right)}{3b^3}$$

input `integrate(x^(5/2)/(b*x+a)^2,x, algorithm="maxima")`

output
$$-a^2\sqrt{x}/(b^4x + a^3) + 5a^2\arctan(b\sqrt{x}/\sqrt{ab})/(\sqrt{ab})^3 + 2/3*(b^4x^{3/2} - 6a^3\sqrt{x})/b^3$$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.94

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = \frac{5a^2 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{\sqrt{ab}b^3} - \frac{a^2\sqrt{x}}{(bx+a)b^3} + \frac{2(b^4x^{3/2} - 6ab^3\sqrt{x})}{3b^6}$$

input `integrate(x^(5/2)/(b*x+a)^2,x, algorithm="giac")`

output
$$5a^2\arctan(b\sqrt{x}/\sqrt{ab})/(\sqrt{ab})^3 - a^2\sqrt{x}/((b^4x + a^3)^3) + 2/3*(b^4x^{3/2} - 6a^3\sqrt{x})/b^6$$

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.84

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = \frac{2x^{3/2}}{3b^2} - \frac{4a\sqrt{x}}{b^3} - \frac{a^2\sqrt{x}}{xb^4 + ab^3} + \frac{5a^{3/2} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{7/2}}$$

input `int(x^(5/2)/(a + b*x)^2,x)`

output
$$(2*x^{3/2})/(3*b^2) - (4*a*x^{1/2})/b^3 - (a^2*x^{1/2})/(a*b^3 + b^4*x) + (5*a^{3/2}*atan((b^{1/2}*x^{1/2})/a^{1/2}))/b^{7/2}$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.22

$$\int \frac{x^{5/2}}{(a+bx)^2} dx = \frac{15\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)a^2 + 15\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)abx - 15\sqrt{x}a^2b - 10\sqrt{x}ab^2x + 2\sqrt{x}}{3b^4(bx+a)}$$

input `int(x^(5/2)/(b*x+a)^2,x)`output `(15*sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a**2 + 15*sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a*b*x - 15*sqrt(x)*a**2*b - 10*sqrt(x)*a*b**2*x + 2*sqrt(x)*b**3*x**2)/(3*b**4*(a + b*x))`

3.117 $\int \frac{1}{\sqrt{x}(a+bx)^3} dx$

Optimal result	754
Mathematica [A] (verified)	754
Rubi [A] (verified)	755
Maple [A] (verified)	756
Fricas [A] (verification not implemented)	757
Sympy [B] (verification not implemented)	757
Maxima [A] (verification not implemented)	758
Giac [A] (verification not implemented)	758
Mupad [B] (verification not implemented)	759
Reduce [B] (verification not implemented)	759

Optimal result

Integrand size = 13, antiderivative size = 57

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx = \sqrt{x} \left(\frac{1}{2a(a+bx)^2} + \frac{1}{4a^2(a+bx)} \right) + \frac{3 \arctan\left(\frac{bx}{a}\right)}{4a^2\sqrt{ab}}$$

output

```
x^(1/2)*(1/2/a/(b*x+a)^2+1/4/a^2/(b*x+a))+3/4*arctan(b*x/a)/a^2/(a*b)^(1/2)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.04

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx = \frac{\sqrt{x}(5a+3bx)}{4a^2(a+bx)^2} + \frac{3 \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4a^{5/2}\sqrt{b}}$$

input

```
Integrate[1/(Sqrt[x]*(a + b*x)^3),x]
```

output

```
(Sqrt[x]*(5*a + 3*b*x))/(4*a^2*(a + b*x)^2) + (3*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/(4*a^(5/2)*Sqrt[b])
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.26, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {52, 52, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{x}(a+bx)^3} dx \\
 & \quad \downarrow 52 \\
 & \frac{3 \int \frac{1}{\sqrt{x}(a+bx)^2} dx}{4a} + \frac{\sqrt{x}}{2a(a+bx)^2} \\
 & \quad \downarrow 52 \\
 & \frac{3 \left(\frac{\int \frac{1}{\sqrt{x}(a+bx)} dx}{2a} + \frac{\sqrt{x}}{a(a+bx)} \right)}{4a} + \frac{\sqrt{x}}{2a(a+bx)^2} \\
 & \quad \downarrow 73 \\
 & \frac{3 \left(\frac{\int \frac{1}{a+bx} d\sqrt{x}}{a} + \frac{\sqrt{x}}{a(a+bx)} \right)}{4a} + \frac{\sqrt{x}}{2a(a+bx)^2} \\
 & \quad \downarrow 218 \\
 & \frac{3 \left(\frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{a^{3/2}\sqrt{b}} + \frac{\sqrt{x}}{a(a+bx)} \right)}{4a} + \frac{\sqrt{x}}{2a(a+bx)^2}
 \end{aligned}$$

input `Int[1/(Sqrt[x]*(a + b*x)^3),x]`

output `Sqrt[x]/(2*a*(a + b*x)^2) + (3*(Sqrt[x]/(a*(a + b*x)) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]]/(a^(3/2)*Sqrt[b]))/(4*a)`

Definitions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.04

method	result	size
derivativedivides	$\frac{\sqrt{x}}{2a(bx+a)^2} + \frac{3\sqrt{x}}{4a(bx+a)} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4a\sqrt{ab}}$	59
default	$\frac{\sqrt{x}}{2a(bx+a)^2} + \frac{3\sqrt{x}}{4a(bx+a)} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4a\sqrt{ab}}$	59

input `int(1/x^(1/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `1/2*x^(1/2)/a/(b*x+a)^2+3/2/a*(1/2*x^(1/2)/a/(b*x+a)+1/2/a/(a*b)^(1/2)*arc tan(b*x^(1/2)/(a*b)^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 186, normalized size of antiderivative = 3.26

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx$$

$$= \left[\frac{3(b^2x^2 + 2abx + a^2)\sqrt{-ab} \log\left(\frac{bx - a - 2\sqrt{-ab}\sqrt{x}}{bx+a}\right) - 2(3ab^2x + 5a^2b)\sqrt{x}}{8(a^3b^3x^2 + 2a^4b^2x + a^5b)}, \right. \\ \left. - \frac{3(b^2x^2 + 2abx + a^2)\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right) - (3ab^2x + 5a^2b)\sqrt{x}}{4(a^3b^3x^2 + 2a^4b^2x + a^5b)} \right]$$

input `integrate(1/x^(1/2)/(b*x+a)^3,x, algorithm="fricas")`

output

```
[-1/8*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*
sqrt(x))/(b*x + a)) - 2*(3*a*b^2*x + 5*a^2*b)*sqrt(x))/(a^3*b^3*x^2 + 2*a^
4*b^2*x + a^5*b), -1/4*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(a*b)*arctan(sqrt(
a*b)/(b*sqrt(x))) - (3*a*b^2*x + 5*a^2*b)*sqrt(x))/(a^3*b^3*x^2 + 2*a^4*b^
2*x + a^5*b)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 632 vs. 2(48) = 96.

Time = 8.74 (sec) , antiderivative size = 632, normalized size of antiderivative = 11.09

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx$$

$$= \left\{ \begin{array}{l} \frac{\infty}{x^{\frac{5}{2}}} \\ \frac{2\sqrt{x}}{a^3} \\ -\frac{2}{5b^3x^{\frac{5}{2}}} \\ \frac{3a^2 \log\left(\sqrt{x} - \sqrt{-\frac{a}{b}}\right)}{8a^4b\sqrt{-\frac{a}{b}} + 16a^3b^2x\sqrt{-\frac{a}{b}} + 8a^2b^3x^2\sqrt{-\frac{a}{b}}} - \frac{3a^2 \log\left(\sqrt{x} + \sqrt{-\frac{a}{b}}\right)}{8a^4b\sqrt{-\frac{a}{b}} + 16a^3b^2x\sqrt{-\frac{a}{b}} + 8a^2b^3x^2\sqrt{-\frac{a}{b}}} + \frac{10ab\sqrt{x}\sqrt{-\frac{a}{b}}}{8a^4b\sqrt{-\frac{a}{b}} + 16a^3b^2x\sqrt{-\frac{a}{b}} + 8a^2b^3x^2\sqrt{-\frac{a}{b}}} \end{array} \right.$$

input `integrate(1/x**(1/2)/(b*x+a)**3,x)`

output

```
Piecewise((zoo/x**(5/2), Eq(a, 0) & Eq(b, 0)), (2*sqrt(x)/a**3, Eq(b, 0)),
(-2/(5*b**3*x**(5/2)), Eq(a, 0)), (3*a**2*log(sqrt(x) - sqrt(-a/b))/(8*a**
4*b*sqrt(-a/b) + 16*a**3*b**2*x*sqrt(-a/b) + 8*a**2*b**3*x**2*sqrt(-a/b))
- 3*a**2*log(sqrt(x) + sqrt(-a/b))/(8*a**4*b*sqrt(-a/b) + 16*a**3*b**2*x*
sqrt(-a/b) + 8*a**2*b**3*x**2*sqrt(-a/b)) + 10*a*b*sqrt(x)*sqrt(-a/b)/(8*a
**4*b*sqrt(-a/b) + 16*a**3*b**2*x*sqrt(-a/b) + 8*a**2*b**3*x**2*sqrt(-a/b)
) + 6*a*b*x*log(sqrt(x) - sqrt(-a/b))/(8*a**4*b*sqrt(-a/b) + 16*a**3*b**2*
x*sqrt(-a/b) + 8*a**2*b**3*x**2*sqrt(-a/b)) - 6*a*b*x*log(sqrt(x) + sqrt(-
a/b))/(8*a**4*b*sqrt(-a/b) + 16*a**3*b**2*x*sqrt(-a/b) + 8*a**2*b**3*x**2*
sqrt(-a/b)) + 6*b**2*x**(3/2)*sqrt(-a/b)/(8*a**4*b*sqrt(-a/b) + 16*a**3*b*
**2*x*sqrt(-a/b) + 8*a**2*b**3*x**2*sqrt(-a/b)) + 3*b**2*x**2*log(sqrt(x) -
sqrt(-a/b))/(8*a**4*b*sqrt(-a/b) + 16*a**3*b**2*x*sqrt(-a/b) + 8*a**2*b**
3*x**2*sqrt(-a/b)) - 3*b**2*x**2*log(sqrt(x) + sqrt(-a/b))/(8*a**4*b*sqrt(-
a/b) + 16*a**3*b**2*x*sqrt(-a/b) + 8*a**2*b**3*x**2*sqrt(-a/b)), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.05

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx = \frac{3bx^{\frac{3}{2}} + 5a\sqrt{x}}{4(a^2b^2x^2 + 2a^3bx + a^4)} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{aba^2}}$$

input

```
integrate(1/x^(1/2)/(b*x+a)^3,x, algorithm="maxima")
```

output

```
1/4*(3*b*x^(3/2) + 5*a*sqrt(x))/(a^2*b^2*x^2 + 2*a^3*b*x + a^4) + 3/4*arct
an(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a^2)
```

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.82

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx = \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{aba^2}} + \frac{3bx^{\frac{3}{2}} + 5a\sqrt{x}}{4(bx+a)^2a^2}$$

input

```
integrate(1/x^(1/2)/(b*x+a)^3,x, algorithm="giac")
```

output $\frac{3}{4} \arctan(b\sqrt{x}/\sqrt{a*b})/(\sqrt{a*b}*a^2) + 1/4*(3*b*x^{(3/2)} + 5*a*\sqrt{x})/((b*x + a)^2*a^2)$

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx = \frac{\frac{5\sqrt{x}}{4a} + \frac{3bx^{3/2}}{4a^2}}{a^2 + 2abx + b^2x^2} + \frac{3 \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4a^{5/2}\sqrt{b}}$$

input `int(1/(x^(1/2)*(a + b*x)^3),x)`

output $((5*x^{(1/2)})/(4*a) + (3*b*x^{(3/2)})/(4*a^2))/(a^2 + b^2*x^2 + 2*a*b*x) + (3*\operatorname{atan}((b^{(1/2)}*x^{(1/2)})/a^{(1/2)}))/(4*a^{(5/2)}*b^{(1/2)})$

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.98

$$\int \frac{1}{\sqrt{x}(a+bx)^3} dx = \frac{3\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a^2 + 6\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) abx + 3\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) b^2x^2 + 5\sqrt{x}a^2b + 3\sqrt{x}ab^2x}{4a^3b(b^2x^2 + 2abx + a^2)}$$

input `int(1/x^(1/2)/(b*x+a)^3,x)`

output $(3*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{x}*b)/(\sqrt{b}*\sqrt{a}))*a**2 + 6*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{x}*b)/(\sqrt{b}*\sqrt{a}))*a*b*x + 3*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{x}*b)/(\sqrt{b}*\sqrt{a}))*b**2*x**2 + 5*\sqrt{x}*a**2*b + 3*\sqrt{x}*a*b**2*x)/(4*a**3*b*(a**2 + 2*a*b*x + b**2*x**2))$

3.118 $\int \frac{\sqrt{x}}{(a+bx)^3} dx$

Optimal result	760
Mathematica [A] (verified)	760
Rubi [A] (verified)	761
Maple [A] (verified)	762
Fricas [A] (verification not implemented)	763
Sympy [B] (verification not implemented)	763
Maxima [A] (verification not implemented)	764
Giac [A] (verification not implemented)	765
Mupad [B] (verification not implemented)	765
Reduce [B] (verification not implemented)	765

Optimal result

Integrand size = 13, antiderivative size = 63

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = \sqrt{x} \left(-\frac{1}{2b(a+bx)^2} + \frac{1}{4ab(a+bx)} \right) + \frac{\arctan\left(\frac{bx}{a}\right)}{4ab\sqrt{ab}}$$

output

```
x^(1/2)*(-1/2/b/(b*x+a)^2+1/4/a/b/(b*x+a))+1/4*arctan(b*x/a)/a/b/(a*b)^(1/2)
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.95

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = -\frac{\sqrt{x}(a-bx)}{4ab(a+bx)^2} + \frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4a^{3/2}b^{3/2}}$$

input

```
Integrate[Sqrt[x]/(a + b*x)^3,x]
```

output

```
-1/4*(Sqrt[x]*(a - b*x))/(a*b*(a + b*x)^2) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]]/(4*a^(3/2)*b^(3/2))
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.14, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {51, 52, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x}}{(a+bx)^3} dx \\
 & \quad \downarrow \text{51} \\
 & \frac{\int \frac{1}{\sqrt{x}(a+bx)^2} dx}{4b} - \frac{\sqrt{x}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{52} \\
 & \frac{\int \frac{1}{\sqrt{x}(a+bx)} dx}{4b} + \frac{\sqrt{x}}{a(a+bx)} - \frac{\sqrt{x}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{\int \frac{1}{a+bx} d\sqrt{x} + \frac{\sqrt{x}}{a(a+bx)}}{4b} - \frac{\sqrt{x}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{218} \\
 & \frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{a^{3/2}\sqrt{b}} + \frac{\sqrt{x}}{a(a+bx)} - \frac{\sqrt{x}}{2b(a+bx)^2}
 \end{aligned}$$

input `Int[Sqrt[x]/(a + b*x)^3,x]`

output `-1/2*Sqrt[x]/(b*(a + b*x)^2) + (Sqrt[x]/(a*(a + b*x)) + ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]]/Sqrt[a])/(a^(3/2)*Sqrt[b])/(4*b)`

Definitions of rubi rules used

rule 51 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^n/(b*(m + 1))), x] - \text{Simp}[d*(n/(b*(m + 1)))]$
 $\text{Int}[(a + b*x)^{(m + 1)}(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d, n}, x]
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]

rule 52 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^{(n + 1)})/((b*c - a*d)*(m + 1)), x] - \text{Simp}[d*((m + n + 2)/((b*c - a*d)*(m + 1)))]$
 $\text{Int}[(a + b*x)^{(m + 1)}(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]

rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_))^{(n_)}, x_Symbol] \rightarrow \text{With}[p = \text{Denominator}[m], \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p*(m + 1) - 1)}(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}, x]]] /;$ FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

rule 218 $\text{Int}[(a_.) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b]

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$\frac{x^{\frac{3}{2}} - \frac{\sqrt{x}}{4b}}{(bx+a)^2} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4ba\sqrt{ab}}$	52
default	$\frac{x^{\frac{3}{2}} - \frac{\sqrt{x}}{4b}}{(bx+a)^2} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4ba\sqrt{ab}}$	52

input `int(x^(1/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)`

output

```
2*(1/8/a*x^(3/2)-1/8*x^(1/2)/b)/(b*x+a)^2+1/4/b/a/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 186, normalized size of antiderivative = 2.95

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = \left[-\frac{(b^2x^2 + 2abx + a^2)\sqrt{-ab} \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right) - 2(ab^2x - a^2b)\sqrt{x}}{8(a^2b^4x^2 + 2a^3b^3x + a^4b^2)}, \right. \\ \left. -\frac{(b^2x^2 + 2abx + a^2)\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right) - (ab^2x - a^2b)\sqrt{x}}{4(a^2b^4x^2 + 2a^3b^3x + a^4b^2)} \right]$$

input

```
integrate(x^(1/2)/(b*x+a)^3,x, algorithm="fricas")
```

output

```
[-1/8*((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*sqrt(x))/(b*x + a)) - 2*(a*b^2*x - a^2*b)*sqrt(x))/(a^2*b^4*x^2 + 2*a^3*b^3*x + a^4*b^2), -1/4*((b^2*x^2 + 2*a*b*x + a^2)*sqrt(a*b)*arctan(sqrt(a*b)/(b*sqrt(x))) - (a*b^2*x - a^2*b)*sqrt(x))/(a^2*b^4*x^2 + 2*a^3*b^3*x + a^4*b^2)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 627 vs. 2(46) = 92.

Time = 5.58 (sec) , antiderivative size = 627, normalized size of antiderivative = 9.95

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = \left\{ \begin{array}{l} \frac{\infty}{x^{\frac{3}{2}}} \\ \frac{2x^{\frac{3}{2}}}{3a^3} \\ -\frac{2}{3b^3x^{\frac{3}{2}}} \\ \frac{a^2 \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{8a^3b^2\sqrt{-\frac{a}{b}}+16a^2b^3x\sqrt{-\frac{a}{b}}+8ab^4x^2\sqrt{-\frac{a}{b}}} - \frac{a^2 \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{8a^3b^2\sqrt{-\frac{a}{b}}+16a^2b^3x\sqrt{-\frac{a}{b}}+8ab^4x^2\sqrt{-\frac{a}{b}}} - \frac{2ab\sqrt{x}\sqrt{-\frac{a}{b}}}{8a^3b^2\sqrt{-\frac{a}{b}}+16a^2b^3x\sqrt{-\frac{a}{b}}+8ab^4x^2\sqrt{-\frac{a}{b}}} \end{array} \right.$$

input `integrate(x**(1/2)/(b*x+a)**3,x)`

output `Piecewise((zoo/x**(3/2), Eq(a, 0) & Eq(b, 0)), (2*x**(3/2)/(3*a**3), Eq(b, 0)), (-2/(3*b**3*x**(3/2)), Eq(a, 0)), (a**2*log(sqrt(x) - sqrt(-a/b))/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)) - a**2*log(sqrt(x) + sqrt(-a/b))/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)) - 2*a*b*sqrt(x)*sqrt(-a/b)/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)) + 2*a*b*x*log(sqrt(x) - sqrt(-a/b))/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)) - 2*a*b*x*log(sqrt(x) + sqrt(-a/b))/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)) + 2*b**2*x**(3/2)*sqrt(-a/b)/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)) + b**2*x**2*log(sqrt(x) - sqrt(-a/b))/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)) - b**2*x**2*log(sqrt(x) + sqrt(-a/b))/(8*a**3*b**2*sqrt(-a/b) + 16*a**2*b**3*x*sqrt(-a/b) + 8*a*b**4*x**2*sqrt(-a/b)), True))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.02

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = \frac{bx^{\frac{3}{2}} - a\sqrt{x}}{4(ab^3x^2 + 2a^2b^2x + a^3b)} + \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{ab}}$$

input `integrate(x^(1/2)/(b*x+a)^3,x, algorithm="maxima")`

output `1/4*(b*x^(3/2) - a*sqrt(x))/(a*b^3*x^2 + 2*a^2*b^2*x + a^3*b) + 1/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = \frac{\arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abab}} + \frac{bx^{\frac{3}{2}} - a\sqrt{x}}{4(bx+a)^2 ab}$$

input `integrate(x^(1/2)/(b*x+a)^3,x, algorithm="giac")`output `1/4*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*a*b) + 1/4*(b*x^(3/2) - a*sqrt(x))/((b*x + a)^2*a*b)`**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.89

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = \frac{\frac{x^{3/2}}{4a} - \frac{\sqrt{x}}{4b}}{a^2 + 2abx + b^2x^2} + \frac{\operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4a^{3/2}b^{3/2}}$$

input `int(x^(1/2)/(a + b*x)^3,x)`output `(x^(3/2)/(4*a) - x^(1/2)/(4*b))/(a^2 + b^2*x^2 + 2*a*b*x) + atan((b^(1/2)*x^(1/2))/a^(1/2))/(4*a^(3/2)*b^(3/2))`**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.75

$$\int \frac{\sqrt{x}}{(a+bx)^3} dx = \frac{\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)a^2 + 2\sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)abx + \sqrt{b}\sqrt{a}\operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right)b^2x^2 - \sqrt{x}a^2b + \sqrt{x}ab^2x}{4a^2b^2(b^2x^2 + 2abx + a^2)}$$

input `int(x^(1/2)/(b*x+a)^3,x)`

output

```
(sqrt(b)*sqrt(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a**2 + 2*sqrt(b)*sqrt
(a)*atan((sqrt(x)*b)/(sqrt(b)*sqrt(a)))*a*b*x + sqrt(b)*sqrt(a)*atan((sqrt
(x)*b)/(sqrt(b)*sqrt(a)))*b**2*x**2 - sqrt(x)*a**2*b + sqrt(x)*a*b**2*x)/(
4*a**2*b**2*(a**2 + 2*a*b*x + b**2*x**2))
```

3.119 $\int \frac{x^{3/2}}{(a+bx)^3} dx$

Optimal result	767
Mathematica [A] (verified)	767
Rubi [A] (verified)	768
Maple [A] (verified)	769
Fricas [A] (verification not implemented)	770
Sympy [B] (verification not implemented)	770
Maxima [A] (verification not implemented)	771
Giac [A] (verification not implemented)	771
Mupad [B] (verification not implemented)	772
Reduce [B] (verification not implemented)	772

Optimal result

Integrand size = 13, antiderivative size = 87

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = -\frac{2x^{3/2}}{b(a+bx)^2} + \frac{3a \left(\sqrt{x} \left(-\frac{1}{2b(a+bx)^2} + \frac{1}{4ab(a+bx)} \right) + \frac{\arctan\left(\frac{bx}{a}\right)}{4ab\sqrt{ab}} \right)}{b}$$

output

```
-2*x^(3/2)/b/(b*x+a)^2+3*a*(x^(1/2)*(-1/2/b/(b*x+a)^2+1/4/a/b/(b*x+a))+1/4*arctan(b*x/a)/a/b/(a*b)^(1/2))/b
```

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.68

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = -\frac{\sqrt{x}(3a+5bx)}{4b^2(a+bx)^2} + \frac{3 \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4\sqrt{ab}^{5/2}}$$

input

```
Integrate[x^(3/2)/(a + b*x)^3,x]
```

output

```
-1/4*(Sqrt[x]*(3*a + 5*b*x))/(b^2*(a + b*x)^2) + (3*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/(4*Sqrt[a]*b^(5/2))
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.84, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {51, 51, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{3/2}}{(a+bx)^3} dx \\
 & \quad \downarrow \text{51} \\
 & \frac{3 \int \frac{\sqrt{x}}{(a+bx)^2} dx}{4b} - \frac{x^{3/2}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{51} \\
 & \frac{3 \left(\frac{\int \frac{1}{\sqrt{x}(a+bx)} dx}{2b} - \frac{\sqrt{x}}{b(a+bx)} \right)}{4b} - \frac{x^{3/2}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{3 \left(\frac{\int \frac{1}{a+bx} d\sqrt{x}}{b} - \frac{\sqrt{x}}{b(a+bx)} \right)}{4b} - \frac{x^{3/2}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{218} \\
 & \frac{3 \left(\frac{\arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{ab^{3/2}}} - \frac{\sqrt{x}}{b(a+bx)} \right)}{4b} - \frac{x^{3/2}}{2b(a+bx)^2}
 \end{aligned}$$

input

 $\text{Int}[x^{(3/2)}/(a + b*x)^3, x]$

output

$$-1/2*x^{(3/2)}/(b*(a + b*x)^2) + (3*(-(\text{Sqrt}[x]/(b*(a + b*x)))) + \text{ArcTan}[(\text{Sqrt}[b]*\text{Sqrt}[x])/(\text{Sqrt}[a])]/(\text{Sqrt}[a]*b^{(3/2)})))/(4*b)$$

Definitions of rubi rules used

rule 51

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 218

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.57

method	result	size
derivativedivides	$\frac{-\frac{5x^{\frac{3}{2}}}{4b} - \frac{3a\sqrt{x}}{4b^2}}{(bx+a)^2} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4b^2\sqrt{ab}}$	50
default	$\frac{-\frac{5x^{\frac{3}{2}}}{4b} - \frac{3a\sqrt{x}}{4b^2}}{(bx+a)^2} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4b^2\sqrt{ab}}$	50

input

```
int(x^(3/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)
```

output

```
2*(-5/8*x^(3/2)/b-3/8*a/b^2*x^(1/2))/(b*x+a)^2+3/4/b^2/(a*b)^(1/2)*arctan(
b*x^(1/2)/(a*b)^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.13

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = \left[-\frac{3(b^2x^2 + 2abx + a^2)\sqrt{-ab} \log\left(\frac{bx-a-2\sqrt{-ab}\sqrt{x}}{bx+a}\right) + 2(5ab^2x + 3a^2b)\sqrt{x}}{8(ab^5x^2 + 2a^2b^4x + a^3b^3)}, \right. \\ \left. -\frac{3(b^2x^2 + 2abx + a^2)\sqrt{ab} \arctan\left(\frac{\sqrt{ab}}{b\sqrt{x}}\right) + (5ab^2x + 3a^2b)\sqrt{x}}{4(ab^5x^2 + 2a^2b^4x + a^3b^3)} \right]$$

input `integrate(x^(3/2)/(b*x+a)^3,x, algorithm="fricas")`

output

```
[-1/8*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a*b)*log((b*x - a - 2*sqrt(-a*b)*
sqrt(x))/(b*x + a)) + 2*(5*a*b^2*x + 3*a^2*b)*sqrt(x))/(a*b^5*x^2 + 2*a^2*b
b^4*x + a^3*b^3), -1/4*(3*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(a*b)*arctan(sqrt(
a*b)/(b*sqrt(x))) + (5*a*b^2*x + 3*a^2*b)*sqrt(x))/(a*b^5*x^2 + 2*a^2*b^4*
x + a^3*b^3)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 605 vs. 2(66) = 132.

Time = 10.68 (sec) , antiderivative size = 605, normalized size of antiderivative = 6.95

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = \left\{ \begin{array}{l} \frac{\infty}{\sqrt{x}} \\ \frac{2x^{\frac{5}{2}}}{5a^3} \\ -\frac{2}{b^3\sqrt{x}} \\ \frac{3a^2 \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{8a^2b^3\sqrt{-\frac{a}{b}+16ab^4x\sqrt{-\frac{a}{b}}+8b^5x^2\sqrt{-\frac{a}{b}}}} - \frac{3a^2 \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{8a^2b^3\sqrt{-\frac{a}{b}+16ab^4x\sqrt{-\frac{a}{b}}+8b^5x^2\sqrt{-\frac{a}{b}}}} - \frac{6ab\sqrt{x}\sqrt{-\frac{a}{b}}}{8a^2b^3\sqrt{-\frac{a}{b}+16ab^4x\sqrt{-\frac{a}{b}}+8b^5x^2\sqrt{-\frac{a}{b}}}} \end{array} \right.$$

input `integrate(x**(3/2)/(b*x+a)**3,x)`

output

```
Piecewise((zoo/sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(5/2)/(5*a**3), Eq(b,
0)), (-2/(b**3*sqrt(x)), Eq(a, 0)), (3*a**2*log(sqrt(x) - sqrt(-a/b))/(8*a
**2*b**3*sqrt(-a/b) + 16*a*b**4*x*sqrt(-a/b) + 8*b**5*x**2*sqrt(-a/b)) - 3
*a**2*log(sqrt(x) + sqrt(-a/b))/(8*a**2*b**3*sqrt(-a/b) + 16*a*b**4*x*sqrt
(-a/b) + 8*b**5*x**2*sqrt(-a/b)) - 6*a*b*sqrt(x)*sqrt(-a/b)/(8*a**2*b**3*s
qrt(-a/b) + 16*a*b**4*x*sqrt(-a/b) + 8*b**5*x**2*sqrt(-a/b)) + 6*a*b*x*log
(sqrt(x) - sqrt(-a/b))/(8*a**2*b**3*sqrt(-a/b) + 16*a*b**4*x*sqrt(-a/b) +
8*b**5*x**2*sqrt(-a/b)) - 6*a*b*x*log(sqrt(x) + sqrt(-a/b))/(8*a**2*b**3*s
qrt(-a/b) + 16*a*b**4*x*sqrt(-a/b) + 8*b**5*x**2*sqrt(-a/b)) - 10*b**2*x**
(3/2)*sqrt(-a/b)/(8*a**2*b**3*sqrt(-a/b) + 16*a*b**4*x*sqrt(-a/b) + 8*b**5
*x**2*sqrt(-a/b)) + 3*b**2*x**2*log(sqrt(x) - sqrt(-a/b))/(8*a**2*b**3*sqr
t(-a/b) + 16*a*b**4*x*sqrt(-a/b) + 8*b**5*x**2*sqrt(-a/b)) - 3*b**2*x**2*1
og(sqrt(x) + sqrt(-a/b))/(8*a**2*b**3*sqrt(-a/b) + 16*a*b**4*x*sqrt(-a/b)
+ 8*b**5*x**2*sqrt(-a/b)), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.70

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = -\frac{5bx^{3/2} + 3a\sqrt{x}}{4(b^4x^2 + 2ab^3x + a^2b^2)} + \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^2}}$$

input

```
integrate(x^(3/2)/(b*x+a)^3,x, algorithm="maxima")
```

output

```
-1/4*(5*b*x^(3/2) + 3*a*sqrt(x))/(b^4*x^2 + 2*a*b^3*x + a^2*b^2) + 3/4*arc
tan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^2)
```

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.54

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = \frac{3 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^2}} - \frac{5bx^{3/2} + 3a\sqrt{x}}{4(bx+a)^2b^2}$$

input

```
integrate(x^(3/2)/(b*x+a)^3,x, algorithm="giac")
```

output $\frac{3}{4} \arctan(b\sqrt{x}/\sqrt{a*b})/(\sqrt{a*b}*b^2) - \frac{1}{4}*(5*b*x^{(3/2)} + 3*a*sqr(x))/((b*x + a)^2*b^2)$

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.67

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = \frac{3 \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4\sqrt{a}b^{5/2}} - \frac{\frac{5x^{3/2}}{4b} + \frac{3a\sqrt{x}}{4b^2}}{a^2 + 2abx + b^2x^2}$$

input `int(x^(3/2)/(a + b*x)^3,x)`

output $(3*\operatorname{atan}((b^{(1/2)}*x^{(1/2)})/a^{(1/2)}))/(4*a^{(1/2)}*b^{(5/2)}) - ((5*x^{(3/2)})/(4*b) + (3*a*x^{(1/2)})/(4*b^2))/(a^2 + b^2*x^2 + 2*a*b*x)$

Reduce [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.30

$$\int \frac{x^{3/2}}{(a+bx)^3} dx = \frac{3\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a^2 + 6\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) abx + 3\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) b^2x^2 - 3\sqrt{x}}{4ab^3(b^2x^2 + 2abx + a^2)}$$

input `int(x^(3/2)/(b*x+a)^3,x)`

output $(3*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{x}*b)/(\sqrt{b}*\sqrt{a}))*a**2 + 6*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{x}*b)/(\sqrt{b}*\sqrt{a}))*a*b*x + 3*\sqrt{b}*\sqrt{a}*\operatorname{atan}((\sqrt{x}*b)/(\sqrt{b}*\sqrt{a}))*b**2*x**2 - 3*\sqrt{x})*a**2*b - 5*\sqrt{x})*a*b**2*x)/(4*a*b**3*(a**2 + 2*a*b*x + b**2*x**2))$

3.120 $\int \frac{x^{5/2}}{(a+bx)^3} dx$

Optimal result	773
Mathematica [A] (verified)	773
Rubi [A] (verified)	774
Maple [A] (verified)	776
Fricas [A] (verification not implemented)	776
Sympy [B] (verification not implemented)	777
Maxima [A] (verification not implemented)	778
Giac [A] (verification not implemented)	778
Mupad [B] (verification not implemented)	778
Reduce [B] (verification not implemented)	779

Optimal result

Integrand size = 13, antiderivative size = 101

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = \frac{2\sqrt{x}\left(\frac{5ax}{b^2} + \frac{x^2}{b}\right)}{(a+bx)^2} - \frac{15a^2\left(\sqrt{x}\left(-\frac{1}{2b(a+bx)^2} + \frac{1}{4ab(a+bx)}\right) + \frac{\arctan\left(\frac{bx}{a}\right)}{4ab\sqrt{ab}}\right)}{b^2}$$

output

```
2*x^(1/2)*(5*a*x/b^2+x^2/b)/(b*x+a)^2-15*a^2*(x^(1/2)*(-1/2/b/(b*x+a)^2+1/4/a/b/(b*x+a))+1/4*arctan(b*x/a)/a/b/(a*b)^(1/2))/b^2
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.69

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = \frac{\sqrt{x}(15a^2 + 25abx + 8b^2x^2)}{4b^3(a+bx)^2} - \frac{15\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4b^{7/2}}$$

input

```
Integrate[x^(5/2)/(a + b*x)^3,x]
```

output

```
(Sqrt[x]*(15*a^2 + 25*a*b*x + 8*b^2*x^2))/(4*b^3*(a + b*x)^2) - (15*Sqrt[a]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/(4*b^(7/2))
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.91, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {51, 51, 60, 73, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{5/2}}{(a+bx)^3} dx \\
 & \quad \downarrow \text{51} \\
 & \frac{5 \int \frac{x^{3/2}}{(a+bx)^2} dx}{4b} - \frac{x^{5/2}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{51} \\
 & \frac{5 \left(\frac{3 \int \frac{\sqrt{x}}{a+bx} dx}{2b} - \frac{x^{3/2}}{b(a+bx)} \right)}{4b} - \frac{x^{5/2}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{60} \\
 & \frac{5 \left(\frac{3 \left(\frac{2\sqrt{x}}{b} - \frac{a \int \frac{1}{\sqrt{x}(a+bx)} dx}{b} \right)}{2b} - \frac{x^{3/2}}{b(a+bx)} \right)}{4b} - \frac{x^{5/2}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{5 \left(\frac{3 \left(\frac{2\sqrt{x}}{b} - \frac{2a \int \frac{1}{a+bx} d\sqrt{x}}{b} \right)}{2b} - \frac{x^{3/2}}{b(a+bx)} \right)}{4b} - \frac{x^{5/2}}{2b(a+bx)^2} \\
 & \quad \downarrow \text{218} \\
 & \frac{5 \left(\frac{3 \left(\frac{2\sqrt{x}}{b} - \frac{2\sqrt{a} \arctan\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{b^{3/2}} \right)}{2b} - \frac{x^{3/2}}{b(a+bx)} \right)}{4b} - \frac{x^{5/2}}{2b(a+bx)^2}
 \end{aligned}$$

input `Int[x^(5/2)/(a + b*x)^3,x]`

output `-1/2*x^(5/2)/(b*(a + b*x)^2) + (5*(-x^(3/2)/(b*(a + b*x))) + (3*((2*Sqrt[x])/b - (2*Sqrt[a]*ArcTan[(Sqrt[b]*Sqrt[x])/Sqrt[a]])/b^(3/2)))/(2*b))/(4*b)`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 218 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.55

method	result	size
derivativedivides	$\frac{2\sqrt{x}}{b^3} - \frac{2a \left(\frac{-9x^{\frac{3}{2}}b - 7\sqrt{x}a}{(bx+a)^2} + \frac{15 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{8\sqrt{ab}} \right)}{b^3}$	56
default	$\frac{2\sqrt{x}}{b^3} - \frac{2a \left(\frac{-9x^{\frac{3}{2}}b - 7\sqrt{x}a}{(bx+a)^2} + \frac{15 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{8\sqrt{ab}} \right)}{b^3}$	56
risch	$\frac{2\sqrt{x}}{b^3} - \frac{a \left(\frac{-9x^{\frac{3}{2}}b - 7\sqrt{x}a}{(bx+a)^2} + \frac{15 \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{ab}} \right)}{b^3}$	57

input `int(x^(5/2)/(b*x+a)^3,x,method=_RETURNVERBOSE)`

output `2*x^(1/2)/b^3-2*a/b^3*((-9/8*x^(3/2)*b-7/8*x^(1/2)*a)/(b*x+a)^2+15/8/(a*b)^(1/2)*arctan(b*x^(1/2)/(a*b)^(1/2)))`

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.98

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = \left[\frac{15(b^2x^2 + 2abx + a^2)\sqrt{-\frac{a}{b}} \log\left(\frac{bx - 2b\sqrt{x}\sqrt{-\frac{a}{b}} - a}{bx+a}\right) + 2(8b^2x^2 + 25abx + 15a^2)\sqrt{x}}{8(b^5x^2 + 2ab^4x + a^2b^3)}, \right. \\ \left. - \frac{15(b^2x^2 + 2abx + a^2)\sqrt{\frac{a}{b}} \arctan\left(\frac{b\sqrt{x}\sqrt{\frac{a}{b}}}{a}\right) - (8b^2x^2 + 25abx + 15a^2)\sqrt{x}}{4(b^5x^2 + 2ab^4x + a^2b^3)} \right]$$

input `integrate(x^(5/2)/(b*x+a)^3,x, algorithm="fricas")`

output

```
[1/8*(15*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a/b)*log((b*x - 2*b*sqrt(x)*sqrt(-a/b) - a)/(b*x + a)) + 2*(8*b^2*x^2 + 25*a*b*x + 15*a^2)*sqrt(x))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3), -1/4*(15*(b^2*x^2 + 2*a*b*x + a^2)*sqrt(a/b)*arc tan(b*sqrt(x)*sqrt(a/b)/a) - (8*b^2*x^2 + 25*a*b*x + 15*a^2)*sqrt(x))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 683 vs. $2(82) = 164$.

Time = 18.82 (sec) , antiderivative size = 683, normalized size of antiderivative = 6.76

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = \begin{cases} \tilde{\infty}\sqrt{x} \\ \frac{2x^{7/2}}{7a^3} \\ \frac{2\sqrt{x}}{b^3} \\ -\frac{15a^3 \log\left(\sqrt{x}-\sqrt{-\frac{a}{b}}\right)}{8a^2b^4\sqrt{-\frac{a}{b}}+16ab^5x\sqrt{-\frac{a}{b}}+8b^6x^2\sqrt{-\frac{a}{b}}} + \frac{15a^3 \log\left(\sqrt{x}+\sqrt{-\frac{a}{b}}\right)}{8a^2b^4\sqrt{-\frac{a}{b}}+16ab^5x\sqrt{-\frac{a}{b}}+8b^6x^2\sqrt{-\frac{a}{b}}} + \frac{30a^2b\sqrt{x}\sqrt{-\frac{a}{b}}}{8a^2b^4\sqrt{-\frac{a}{b}}+16ab^5x\sqrt{-\frac{a}{b}}+8b^6x^2\sqrt{-\frac{a}{b}}} \end{cases}$$

input

```
integrate(x**(5/2)/(b*x+a)**3,x)
```

output

```
Piecewise((zoo*sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(7/2)/(7*a**3), Eq(b, 0)), (2*sqrt(x)/b**3, Eq(a, 0)), (-15*a**3*log(sqrt(x) - sqrt(-a/b))/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) + 15*a**3*log(sqrt(x) + sqrt(-a/b))/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) + 30*a**2*b*sqrt(x)*sqrt(-a/b)/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) - 30*a**2*b*x*log(sqrt(x) - sqrt(-a/b))/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) + 30*a**2*b*x*log(sqrt(x) + sqrt(-a/b))/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) + 50*a*b**2*x**(3/2)*sqrt(-a/b)/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) - 15*a*b**2*x**2*log(sqrt(x) - sqrt(-a/b))/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) + 15*a*b**2*x**2*log(sqrt(x) + sqrt(-a/b))/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b)) + 16*b**3*x**(5/2)*sqrt(-a/b)/(8*a**2*b**4*sqrt(-a/b) + 16*a*b**5*x*sqrt(-a/b) + 8*b**6*x**2*sqrt(-a/b))), True))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.72

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = \frac{9abx^{3/2} + 7a^2\sqrt{x}}{4(b^5x^2 + 2ab^4x + a^2b^3)} - \frac{15a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^3}} + \frac{2\sqrt{x}}{b^3}$$

input `integrate(x^(5/2)/(b*x+a)^3,x, algorithm="maxima")`output `1/4*(9*a*b*x^(3/2) + 7*a^2*sqrt(x))/(b^5*x^2 + 2*a*b^4*x + a^2*b^3) - 15/4*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 2*sqrt(x)/b^3`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.58

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = -\frac{15a \arctan\left(\frac{b\sqrt{x}}{\sqrt{ab}}\right)}{4\sqrt{abb^3}} + \frac{2\sqrt{x}}{b^3} + \frac{9abx^{3/2} + 7a^2\sqrt{x}}{4(bx+a)^2b^3}$$

input `integrate(x^(5/2)/(b*x+a)^3,x, algorithm="giac")`output `-15/4*a*arctan(b*sqrt(x)/sqrt(a*b))/(sqrt(a*b)*b^3) + 2*sqrt(x)/b^3 + 1/4*(9*a*b*x^(3/2) + 7*a^2*sqrt(x))/((b*x + a)^2*b^3)`**Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.68

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = \frac{\frac{7a^2\sqrt{x}}{4} + \frac{9abx^{3/2}}{4}}{a^2b^3 + 2ab^4x + b^5x^2} + \frac{2\sqrt{x}}{b^3} - \frac{15\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{b}\sqrt{x}}{\sqrt{a}}\right)}{4b^{7/2}}$$

input `int(x^(5/2)/(a + b*x)^3,x)`

output
$$\left(\frac{7a^2x^{1/2}}{4} + \frac{9abx^{3/2}}{4}\right) / (a^2b^3 + b^5x^2 + 2ab^4x) + \frac{2x^{1/2}}{b^3} - \frac{15a^{1/2} \operatorname{atan}\left(\frac{b^{1/2}x^{1/2}}{a^{1/2}}\right)}{4b^{7/2}}$$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.19

$$\int \frac{x^{5/2}}{(a+bx)^3} dx = \frac{-15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a^2 - 30\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) abx - 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) b^2x^2 + 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a^2 + 30\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) abx + 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) b^2x^2}{4b^4(b^2x^2 + 2abx + a^2)}$$

input `int(x^(5/2)/(b*x+a)^3,x)`

output
$$\left(-15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a^2 - 30\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) abx - 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) b^2x^2 + 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) a^2 + 30\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) abx + 15\sqrt{b}\sqrt{a} \operatorname{atan}\left(\frac{\sqrt{x}b}{\sqrt{b}\sqrt{a}}\right) b^2x^2\right) / (4b^4(b^2x^2 + 2abx + a^2))$$

3.121 $\int \frac{1}{\sqrt{x}(a+bx^2)} dx$

Optimal result	780
Mathematica [A] (verified)	780
Rubi [B] (verified)	781
Maple [A] (verified)	785
Fricas [C] (verification not implemented)	785
Sympy [A] (verification not implemented)	786
Maxima [B] (verification not implemented)	786
Giac [B] (verification not implemented)	788
Mupad [B] (verification not implemented)	788
Reduce [B] (verification not implemented)	789

Optimal result

Integrand size = 15, antiderivative size = 99

$$\int \frac{1}{\sqrt{x}(a+bx^2)} dx = \frac{\arctan\left(\frac{\sqrt[4]{2}\sqrt{\frac{a}{b}}\sqrt{x}}{\sqrt{\frac{a}{b}}-x}\right) + \log\left(\frac{\sqrt{\frac{a}{b}}+\sqrt[4]{2}\sqrt{\frac{a}{b}}\sqrt{x+x}}{\sqrt{a+bx^2}}\right)}{\sqrt{2}\left(\frac{a}{b}\right)^{3/4}b}$$

output

```
1/2*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))+ln(((a/b)^(1/2)+
^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/(a/b)^(3/4)/b
```

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.93

$$\int \frac{1}{\sqrt{x}(a+bx^2)} dx = \frac{-\arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right) + \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}+\sqrt{bx}}\right)}{\sqrt{2}a^{3/4}\sqrt[4]{b}}$$

input

```
Integrate[1/(Sqrt[x]*(a + b*x^2)),x]
```

output

```
(-ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])] + ArcTan
h[(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])/(Sqrt[a] + Sqrt[b]*x)]/(Sqrt[2]*a^(3/
4)*b^(1/4))
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 214 vs. 2(99) = 198.

Time = 0.38 (sec) , antiderivative size = 214, normalized size of antiderivative = 2.16, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt{x}(a+bx^2)} dx \\
 & \quad \downarrow \text{266} \\
 & 2 \int \frac{1}{bx^2+a} d\sqrt{x} \\
 & \quad \downarrow \text{755} \\
 & 2 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} \right) \\
 & \quad \downarrow \text{1476} \\
 & 2 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \frac{\sqrt{a}}{\sqrt{b}}}}{2\sqrt{b}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \frac{\sqrt{a}}{\sqrt{b}}}}{2\sqrt{b}} d\sqrt{x}}{2\sqrt{b}}}{2\sqrt{a}} \right) \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

$$2 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{-x-1} d\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

217

$$2 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

1479

$$2 \left(\frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

25

$$2 \left(\frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

27

$$2 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{x - \sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}} d\sqrt{x}}{\frac{\sqrt[4]{b}}{2\sqrt{2} \sqrt[4]{a} \sqrt{b}}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}}{x + \sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}} d\sqrt{x}}{\frac{\sqrt[4]{b}}{2\sqrt{2} \sqrt[4]{a} \sqrt{b}}} + \frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right)$$

↓ 1103

$$2 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}}}{2\sqrt{a}} + \frac{\frac{\log\left(\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx}\right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\log\left(-\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx}\right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}}}{2\sqrt{a}} \right)$$

input `Int[1/(Sqrt[x]*(a + b*x^2)),x]`

output `2*((-(ArcTan[1 - (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[a]) + (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[a])`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.07

method	result	size
derivativedivides	$\frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\frac{a}{b}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1}\right)\right)}{4a}$	106
default	$\frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\frac{a}{b}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1}\right)\right)}{4a}$	106

input `int(1/x^(1/2)/(b*x^2+a),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{4}\frac{(a/b)^{1/4}}{a^{1/2}}\left(\ln\left(\frac{(a/b)^{1/2}+2^{1/2}(a/b)^{1/4}x^{1/2}+x}{(x-2^{1/2})(a/b)^{1/4}x^{1/2}+(a/b)^{1/2}}\right)+2\arctan\left(\frac{2^{1/2}}{(a/b)^{1/4}x^{1/2}+1}\right)+2\arctan\left(\frac{2^{1/2}}{(a/b)^{1/4}x^{1/2}-1}\right)\right)$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.21

$$\begin{aligned} \int \frac{1}{\sqrt{x}(a+bx^2)} dx &= \frac{1}{2} \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} \log \left(a \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} + \sqrt{x}\right) \\ &\quad + \frac{1}{2} i \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} \log \left(i a \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} + \sqrt{x}\right) \\ &\quad - \frac{1}{2} i \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} \log \left(-i a \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} + \sqrt{x}\right) \\ &\quad - \frac{1}{2} \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} \log \left(-a \left(-\frac{1}{a^3b}\right)^{\frac{1}{4}} + \sqrt{x}\right) \end{aligned}$$

input `integrate(1/x^(1/2)/(b*x^2+a),x, algorithm="fricas")`

output

```
1/2*(-1/(a^3*b))^(1/4)*log(a*(-1/(a^3*b))^(1/4) + sqrt(x)) + 1/2*I*(-1/(a^
3*b))^(1/4)*log(I*a*(-1/(a^3*b))^(1/4) + sqrt(x)) - 1/2*I*(-1/(a^3*b))^(1/
4)*log(-I*a*(-1/(a^3*b))^(1/4) + sqrt(x)) - 1/2*(-1/(a^3*b))^(1/4)*log(-a*
(-1/(a^3*b))^(1/4) + sqrt(x))
```

Sympy [A] (verification not implemented)

Time = 1.66 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.05

$$\int \frac{1}{\sqrt{x}(a+bx^2)} dx$$

$$= \begin{cases} \frac{\infty}{x^{\frac{3}{2}}} & \text{for } a = 0 \wedge b = 0 \\ -\frac{2}{3bx^{\frac{3}{2}}} & \text{for } a = 0 \\ \frac{2\sqrt{x}}{a} & \text{for } b = 0 \\ -\frac{\sqrt[4]{-\frac{a}{b}} \log\left(\sqrt{x} - \sqrt[4]{-\frac{a}{b}}\right)}{2a} + \frac{\sqrt[4]{-\frac{a}{b}} \log\left(\sqrt{x} + \sqrt[4]{-\frac{a}{b}}\right)}{2a} + \frac{\sqrt[4]{-\frac{a}{b}} \operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{a} & \text{otherwise} \end{cases}$$

input

```
integrate(1/x**(1/2)/(b*x**2+a),x)
```

output

```
Piecewise((zoo/x**(3/2), Eq(a, 0) & Eq(b, 0)), (-2/(3*b*x**(3/2)), Eq(a, 0
)), (2*sqrt(x)/a, Eq(b, 0)), (-(-a/b)**(1/4)*log(sqrt(x) - (-a/b)**(1/4))/
(2*a) + (-a/b)**(1/4)*log(sqrt(x) + (-a/b)**(1/4))/(2*a) + (-a/b)**(1/4)*a
tan(sqrt(x)/(-a/b)**(1/4))/a, True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 172 vs. 2(79) = 158.

Time = 0.13 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.74

$$\int \frac{1}{\sqrt{x}(a+bx^2)} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}+2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{2\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}-2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{2\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{\sqrt{2} \log\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{bx} + \sqrt{a}\right)}{4a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log\left(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{bx} + \sqrt{a}\right)}{4a^{\frac{3}{4}}b^{\frac{1}{4}}}$$

input `integrate(1/x^(1/2)/(b*x^2+a),x, algorithm="maxima")`

output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + 1/4*sqrt(2)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4)) - 1/4*sqrt(2)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. $2(79) = 158$.

Time = 0.12 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.84

$$\int \frac{1}{\sqrt{x}(a+bx^2)} dx = \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}+2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{2ab} + \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}-2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{2ab} + \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{4ab} - \frac{\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{4ab}$$

input `integrate(1/x^(1/2)/(b*x^2+a),x, algorithm="giac")`

output `1/2*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/(a*b) + 1/2*sqrt(2)*(a*b^3)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/(a*b) + 1/4*sqrt(2)*(a*b^3)^(1/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a*b) - 1/4*sqrt(2)*(a*b^3)^(1/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a*b)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.37

$$\int \frac{1}{\sqrt{x}(a+bx^2)} dx = -\frac{\operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right) + \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{(-a)^{3/4}b^{1/4}}$$

input `int(1/(x^(1/2)*(a + b*x^2)),x)`

output

$$\frac{-(\operatorname{atan}((b^{1/4}x^{1/2})/(-a)^{1/4})) + \operatorname{atanh}((b^{1/4}x^{1/2})/(-a)^{1/4}))}{((-a)^{3/4}b^{1/4})}$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.13

$$\int \frac{1}{\sqrt{x}(a+bx^2)} dx$$

$$= \frac{\sqrt{2} \left(-2\operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{1/4}a^{1/4}\sqrt{2}}\right) + 2\operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}+2\sqrt{x}\sqrt{b}}{b^{1/4}a^{1/4}\sqrt{2}}\right) - \log\left(-\sqrt{x}b^{1/4}a^{1/4}\sqrt{2} + \sqrt{a} + \sqrt{b}x\right) + \log\left(\sqrt{x}b^{1/4}a^{1/4}\sqrt{2} + \sqrt{a} + \sqrt{b}x\right) \right)}{4b^{1/4}a^{3/4}}$$

input

$$\operatorname{int}(1/x^{1/2}/(b*x^2+a), x)$$

output

$$\frac{(b^{3/4}a^{1/4}\sqrt{2} * (-2\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2} - 2\sqrt{x}\sqrt{b})/b^{1/4}a^{1/4}\sqrt{2})) + 2\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2} + 2\sqrt{x}\sqrt{b})/b^{1/4}a^{1/4}\sqrt{2}) - \log(-\sqrt{x}b^{1/4}a^{1/4}\sqrt{2} + \sqrt{a} + \sqrt{b}x) + \log(\sqrt{x}b^{1/4}a^{1/4}\sqrt{2} + \sqrt{a} + \sqrt{b}x)}{4a^{3/4}b^{1/4}}$$

3.122 $\int \frac{\sqrt{x}}{a+bx^2} dx$

Optimal result	790
Mathematica [A] (verified)	790
Rubi [B] (verified)	791
Maple [A] (verified)	795
Fricas [C] (verification not implemented)	795
Sympy [A] (verification not implemented)	796
Maxima [B] (verification not implemented)	797
Giac [B] (verification not implemented)	798
Mupad [B] (verification not implemented)	798
Reduce [B] (verification not implemented)	799

Optimal result

Integrand size = 15, antiderivative size = 101

$$\int \frac{\sqrt{x}}{a + bx^2} dx = \frac{\arctan\left(\frac{\sqrt[4]{\frac{a}{b}}\sqrt{x}}{\sqrt{\frac{a}{b}-x}}\right) - \log\left(\frac{\sqrt{\frac{a}{b}+\sqrt[4]{\frac{a}{b}}\sqrt{x+x}}}{\sqrt{a+bx^2}}\right)}{\sqrt[4]{\frac{a}{b}}b}$$

output

$1/2*(\arctan(2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)}/((a/b)^{(1/2)}-x))-\ln(((a/b)^{(1/2)}+2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)+x}/(b*x^2+a)^{(1/2)}))*2^{(1/2)}/(a/b)^{(1/4)}/b$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.90

$$\int \frac{\sqrt{x}}{a + bx^2} dx = -\frac{\arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right) + \operatorname{arctanh}\left(\frac{\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a+\sqrt{bx}}}\right)}{\sqrt[4]{ab^3/4}}$$

input

`Integrate[Sqrt[x]/(a + b*x^2), x]`

output

```

-((ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])] + ArcTan
h[(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])/(Sqrt[a] + Sqrt[b]*x)])/(Sqrt[2]*a^(1
/4)*b^(3/4))

```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 214 vs. $2(101) = 202$.

Time = 0.39 (sec) , antiderivative size = 214, normalized size of antiderivative = 2.12, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x}}{a + bx^2} dx \\
 & \quad \downarrow \text{266} \\
 & 2 \int \frac{x}{bx^2 + a} d\sqrt{x} \\
 & \quad \downarrow \text{826} \\
 & 2 \left(\frac{\int \frac{\sqrt{bx} + \sqrt{a}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right) \\
 & \quad \downarrow \text{1476} \\
 & 2 \left(\frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt{b}} + \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt{b}} + \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right) \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

$$2 \left(\frac{\int \frac{1}{-x-1} d\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right)$$

217

$$2 \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right)$$

1479

$$2 \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right)$$

25

$$2 \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right)$$

27

$$2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{x-\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}}{x+\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}} d\sqrt{x}}{2\sqrt[4]{a}\sqrt[4]{b}} \right)$$

↓ 1103

$$2 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

input `Int[Sqrt[x]/(a + b*x^2),x]`

output `2*((-ArcTan[1 - (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]) - (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b])`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.05

method	result	size
derivativedivides	$\frac{\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + \sqrt{\frac{a}{b}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + x} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} + 1 \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} - 1 \right) \right)}{4b \left(\frac{a}{b} \right)^{\frac{1}{4}}}$	106
default	$\frac{\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + \sqrt{\frac{a}{b}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + x} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} + 1 \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} - 1 \right) \right)}{4b \left(\frac{a}{b} \right)^{\frac{1}{4}}}$	106

input `int(x^(1/2)/(b*x^2+a),x,method=_RETURNVERBOSE)`output `1/4/b/(a/b)^(1/4)*2^(1/2)*(ln((x-2^(1/2)*(a/b)^(1/4)*x^(1/2)+(a/b)^(1/2))/
((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x))+2*arctan(2^(1/2)/(a/b)^(1/4)*
x^(1/2)+1)+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)-1))`**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.31

$$\begin{aligned} \int \frac{\sqrt{x}}{a + bx^2} dx &= \frac{1}{2} \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + \sqrt{x} \right) \\ &\quad - \frac{1}{2} i \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(i ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + \sqrt{x} \right) \\ &\quad + \frac{1}{2} i \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(-i ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + \sqrt{x} \right) \\ &\quad - \frac{1}{2} \left(-\frac{1}{ab^3} \right)^{\frac{1}{4}} \log \left(-ab^2 \left(-\frac{1}{ab^3} \right)^{\frac{3}{4}} + \sqrt{x} \right) \end{aligned}$$

input `integrate(x^(1/2)/(b*x^2+a),x, algorithm="fricas")`

output

```
1/2*(-1/(a*b^3))^(1/4)*log(a*b^2*(-1/(a*b^3))^(3/4) + sqrt(x)) - 1/2*I*(-1/(a*b^3))^(1/4)*log(I*a*b^2*(-1/(a*b^3))^(3/4) + sqrt(x)) + 1/2*I*(-1/(a*b^3))^(1/4)*log(-I*a*b^2*(-1/(a*b^3))^(3/4) + sqrt(x)) - 1/2*(-1/(a*b^3))^(1/4)*log(-a*b^2*(-1/(a*b^3))^(3/4) + sqrt(x))
```

Sympy [A] (verification not implemented)

Time = 1.03 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.03

$$\int \frac{\sqrt{x}}{a + bx^2} dx = \begin{cases} \frac{\infty}{\sqrt{x}} & \text{for } a = 0 \wedge b = 0 \\ \frac{2x^{3/2}}{3a} & \text{for } b = 0 \\ -\frac{2}{b\sqrt{x}} & \text{for } a = 0 \\ \frac{\log\left(\sqrt{x} - \sqrt[4]{-\frac{a}{b}}\right)}{2b\sqrt[4]{-\frac{a}{b}}} - \frac{\log\left(\sqrt{x} + \sqrt[4]{-\frac{a}{b}}\right)}{2b\sqrt[4]{-\frac{a}{b}}} + \frac{\operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{b\sqrt[4]{-\frac{a}{b}}} & \text{otherwise} \end{cases}$$

input

```
integrate(x**(1/2)/(b*x**2+a),x)
```

output

```
Piecewise((zoo/sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(3/2)/(3*a), Eq(b, 0)), (-2/(b*sqrt(x)), Eq(a, 0)), (log(sqrt(x) - (-a/b)**(1/4))/(2*b*(-a/b)**(1/4)) - log(sqrt(x) + (-a/b)**(1/4))/(2*b*(-a/b)**(1/4)) + atan(sqrt(x)/(-a/b)**(1/4))/(b*(-a/b)**(1/4)), True))
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 172 vs. $2(81) = 162$.

Time = 0.12 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.70

$$\int \frac{\sqrt{x}}{a + bx^2} dx = \frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} + 2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{2\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} + \frac{\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} - 2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{2\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} - \frac{\sqrt{2} \log\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a}\right)}{4a^{\frac{1}{4}}b^{\frac{3}{4}}} + \frac{\sqrt{2} \log\left(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a}\right)}{4a^{\frac{1}{4}}b^{\frac{3}{4}}}$$

input `integrate(x^(1/2)/(b*x^2+a),x, algorithm="maxima")`

output `1/2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) + 1/2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) - 1/4*sqrt(2)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4)) + 1/4*sqrt(2)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4))`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. $2(81) = 162$.

Time = 0.12 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.80

$$\int \frac{\sqrt{x}}{a + bx^2} dx = \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}} + 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{2ab^3} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}} - 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{2ab^3} - \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}} + x + \sqrt{\frac{a}{b}}\right)}{4ab^3} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}} + x + \sqrt{\frac{a}{b}}\right)}{4ab^3}$$

input `integrate(x^(1/2)/(b*x^2+a),x, algorithm="giac")`

output `1/2*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/(a*b^3) + 1/2*sqrt(2)*(a*b^3)^(3/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/(a*b^3) - 1/4*sqrt(2)*(a*b^3)^(3/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a*b^3) + 1/4*sqrt(2)*(a*b^3)^(3/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a*b^3)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.38

$$\int \frac{\sqrt{x}}{a + bx^2} dx = \frac{\operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right) - \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{(-a)^{1/4} b^{3/4}}$$

input `int(x^(1/2)/(a + b*x^2),x)`

output $(\operatorname{atan}((b^{1/4}x^{1/2})/(-a)^{1/4}) - \operatorname{atanh}((b^{1/4}x^{1/2})/(-a)^{1/4}))/((-a)^{1/4}b^{3/4})$

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.11

$$\int \frac{\sqrt{x}}{a + bx^2} dx$$

$$= \frac{\sqrt{2} \left(-2 \operatorname{atan} \left(\frac{b^{1/4} a^{1/4} \sqrt{2} - 2\sqrt{x} \sqrt{b}}{b^{1/4} a^{1/4} \sqrt{2}} \right) + 2 \operatorname{atan} \left(\frac{b^{1/4} a^{1/4} \sqrt{2} + 2\sqrt{x} \sqrt{b}}{b^{1/4} a^{1/4} \sqrt{2}} \right) + \log \left(-\sqrt{x} b^{1/4} a^{1/4} \sqrt{2} + \sqrt{a} + \sqrt{b} x \right) - \log \left(\sqrt{x} b^{1/4} a^{1/4} \sqrt{2} + \sqrt{a} + \sqrt{b} x \right) \right)}{4b^{3/4} a^{1/4}}$$

input $\operatorname{int}(x^{1/2}/(b*x^2+a), x)$

output $(b^{1/4}a^{3/4}\sqrt{2}*(-2*\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2}-2*\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2}))+2*\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2}+2*\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2}))+\log(-\sqrt{x}b^{1/4}a^{1/4}\sqrt{2}+\sqrt{a}+\sqrt{b}x)-\log(\sqrt{x}b^{1/4}a^{1/4}\sqrt{2}+\sqrt{a}+\sqrt{b}x))/(4*a*b)$

3.123 $\int \frac{x^{3/2}}{a+bx^2} dx$

Optimal result	800
Mathematica [A] (verified)	800
Rubi [B] (verified)	801
Maple [A] (verified)	805
Fricas [C] (verification not implemented)	805
Sympy [A] (verification not implemented)	806
Maxima [B] (verification not implemented)	806
Giac [A] (verification not implemented)	807
Mupad [B] (verification not implemented)	808
Reduce [B] (verification not implemented)	808

Optimal result

Integrand size = 15, antiderivative size = 112

$$\int \frac{x^{3/2}}{a+bx^2} dx = \frac{2\sqrt{x}}{b} - \frac{a \left(\arctan \left(\frac{\sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}-x}} \right) + \log \left(\frac{\sqrt{\frac{a}{b}+\sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x}+x}}{\sqrt{a+bx^2}} \right) \right)}{\sqrt{2} \left(\frac{a}{b}\right)^{3/4} b^2}$$

output

```
2*x^(1/2)/b-1/2*a*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))+ln(
((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/(a/b)^(3/4)/b^2
```

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.05

$$\int \frac{x^{3/2}}{a+bx^2} dx = \frac{4\sqrt[4]{b}\sqrt{x} + \sqrt{2}\sqrt[4]{a} \arctan \left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}} \right) - \sqrt{2}\sqrt[4]{a} \operatorname{arctanh} \left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a+\sqrt{bx}}} \right)}{2b^{5/4}}$$

input

```
Integrate[x^(3/2)/(a + b*x^2),x]
```

output

```
(4*b^(1/4)*Sqrt[x] + Sqrt[2]*a^(1/4)*ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x]]) - Sqrt[2]*a^(1/4)*ArcTanh[(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])/(Sqrt[a] + Sqrt[b]*x)]/(2*b^(5/4))
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 229 vs. 2(112) = 224.

Time = 0.40 (sec) , antiderivative size = 229, normalized size of antiderivative = 2.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {262, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x^{3/2}}{a + bx^2} dx \\
 & \quad \downarrow \text{262} \\
 & \frac{2\sqrt{x}}{b} - \frac{a \int \frac{1}{\sqrt{x}(bx^2+a)} dx}{b} \\
 & \quad \downarrow \text{266} \\
 & \frac{2\sqrt{x}}{b} - \frac{2a \int \frac{1}{bx^2+a} d\sqrt{x}}{b} \\
 & \quad \downarrow \text{755} \\
 & \frac{2\sqrt{x}}{b} - \frac{2a \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} \right)}{b} \\
 & \quad \downarrow \text{1476} \\
 & \frac{2\sqrt{x}}{b} - \frac{2a \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{a}} \right)}{b} \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

$$\frac{2\sqrt{x}}{b} - \frac{2a \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{-x-1} d\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{2\sqrt{a}} \right)}{b}$$

217

$$\frac{2\sqrt{x}}{b} - \frac{2a \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right)}{b}$$

1479

$$\frac{2\sqrt{x}}{b} - \frac{2a \left(\frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right)}{b}$$

25

$$\frac{2\sqrt{x}}{b} - \frac{2a \left(\frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right)}{b}$$

27

rule 217 $\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{(-1)} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 262 $\text{Int}[(c_ \cdot x)^m \cdot (a_ + (b_ \cdot x)^2)^p, x_Symbol] \rightarrow \text{Simp}[c \cdot (c \cdot x)^{m-1} \cdot ((a + b \cdot x^2)^{p+1} / (b \cdot (m + 2 \cdot p + 1))), x] - \text{Simp}[a \cdot c^2 \cdot ((m - 1) / (b \cdot (m + 2 \cdot p + 1))) \ \text{Int}[(c \cdot x)^{m-2} \cdot (a + b \cdot x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{GtQ}[m, 2 - 1] \ \&\& \ \text{NeQ}[m + 2 \cdot p + 1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 266 $\text{Int}[(c_ \cdot x)^m \cdot (a_ + (b_ \cdot x)^2)^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k \cdot (m + 1) - 1} \cdot (a + b \cdot (x^{2 \cdot k}/c^2))^p, x], x, (c \cdot x)^{1/k}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 755 $\text{Int}[(a_ + (b_ \cdot x)^4)^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2 \cdot r) \ \text{Int}[(r - s \cdot x^2)/(a + b \cdot x^4), x], x] + \text{Simp}[1/(2 \cdot r) \ \text{Int}[(r + s \cdot x^2)/(a + b \cdot x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 1082 $\text{Int}[(a_ + (b_ \cdot x) + (c_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \text{!RationalQ}[b^2 - 4 \cdot a \cdot c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_ + (e_ \cdot x))/(a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1476 $\text{Int}[(d_ + (e_ \cdot x)^2)/(a_ + (c_ \cdot x)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2 \cdot (d/e), 2]\}, \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e + q \cdot x + x^2, x], x], x] + \text{Simp}[e/(2 \cdot c) \ \text{Int}[1/\text{Simp}[d/e - q \cdot x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[d \cdot e]$

rule 1479

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$\frac{2\sqrt{x}}{b} - \frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2} \left(\ln\left(\frac{\sqrt{\frac{a}{b}+\sqrt{2}}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\sqrt{\frac{a}{b}}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}+1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}-1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) \right)}{4b}$	115
default	$\frac{2\sqrt{x}}{b} - \frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2} \left(\ln\left(\frac{\sqrt{\frac{a}{b}+\sqrt{2}}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\sqrt{\frac{a}{b}}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}+1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}-1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) \right)}{4b}$	115
risch	$\frac{2\sqrt{x}}{b} - \frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2} \left(\ln\left(\frac{\sqrt{\frac{a}{b}+\sqrt{2}}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\sqrt{\frac{a}{b}}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}+1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}-1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) \right)}{4b}$	115

input

```
int(x^(3/2)/(b*x^2+a), x, method=_RETURNVERBOSE)
```

output

```
2*x^(1/2)/b-1/4/b*(a/b)^(1/4)*2^(1/2)*(ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(x-2^(1/2)*(a/b)^(1/4)*x^(1/2)+(a/b)^(1/2))))+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)+1)+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)-1))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.04

$$\int \frac{x^{3/2}}{a + bx^2} dx = \frac{b\left(-\frac{a}{b^5}\right)^{\frac{1}{4}} \log\left(b\left(-\frac{a}{b^5}\right)^{\frac{1}{4}} + \sqrt{x}\right) + ib\left(-\frac{a}{b^5}\right)^{\frac{1}{4}} \log\left(ib\left(-\frac{a}{b^5}\right)^{\frac{1}{4}} + \sqrt{x}\right) - ib\left(-\frac{a}{b^5}\right)^{\frac{1}{4}} \log\left(-ib\left(-\frac{a}{b^5}\right)^{\frac{1}{4}} + \sqrt{x}\right)}{2b}$$

input `integrate(x^(3/2)/(b*x^2+a),x, algorithm="fricas")`

output
$$-1/2*(b*(-a/b^5)^{1/4}*\log(b*(-a/b^5)^{1/4} + \text{sqrt}(x)) + I*b*(-a/b^5)^{1/4})*\log(I*b*(-a/b^5)^{1/4} + \text{sqrt}(x)) - I*b*(-a/b^5)^{1/4}*\log(-I*b*(-a/b^5)^{1/4} + \text{sqrt}(x)) - b*(-a/b^5)^{1/4}*\log(-b*(-a/b^5)^{1/4} + \text{sqrt}(x)) - 4*\text{sqrt}(x))/b$$

Sympy [A] (verification not implemented)

Time = 1.79 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.98

$$\int \frac{x^{3/2}}{a + bx^2} dx = \begin{cases} \tilde{\infty}\sqrt{x} & \text{for } a = 0 \\ \frac{2x^{5/2}}{5a} & \text{for } b = 0 \\ \frac{2\sqrt{x}}{b} & \text{for } a = 0 \\ \frac{2\sqrt{x}}{b} + \frac{\sqrt[4]{-\frac{a}{b}} \log\left(\sqrt{x} - \sqrt[4]{-\frac{a}{b}}\right)}{2b} - \frac{\sqrt[4]{-\frac{a}{b}} \log\left(\sqrt{x} + \sqrt[4]{-\frac{a}{b}}\right)}{2b} - \frac{\sqrt[4]{-\frac{a}{b}} \operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{b} & \text{otherwise} \end{cases}$$

input `integrate(x**(3/2)/(b*x**2+a),x)`

output `Piecewise((zoo*sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(5/2)/(5*a), Eq(b, 0)), (2*sqrt(x)/b, Eq(a, 0)), (2*sqrt(x)/b + (-a/b)**(1/4)*log(sqrt(x) - (-a/b)**(1/4))/(2*b) - (-a/b)**(1/4)*log(sqrt(x) + (-a/b)**(1/4))/(2*b) - (-a/b)**(1/4)*atan(sqrt(x)/(-a/b)**(1/4))/b, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs. $2(89) = 178$.

Time = 0.13 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.65

$$\int \frac{x^{3/2}}{a + bx^2} dx = \frac{2\sqrt{2}\sqrt{a} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}a^{1/4}b^{1/4} + 2\sqrt{b}\sqrt{x}\right)}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{\sqrt{a}\sqrt{b}}} + \frac{2\sqrt{2}\sqrt{a} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}a^{1/4}b^{1/4} - 2\sqrt{b}\sqrt{x}\right)}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{\sqrt{a}\sqrt{b}}} + \frac{\sqrt{2}a^{1/4} \log\left(\sqrt{2}a^{1/4}b^{1/4}\sqrt{x} + \sqrt{bx} + \sqrt{a}\right)}{b^{1/4}} - \frac{\sqrt{2}a^{1/4}}{4b} + \frac{2\sqrt{x}}{b}$$

input `integrate(x^(3/2)/(b*x^2+a),x, algorithm="maxima")`

output `-1/4*(2*sqrt(2)*sqrt(a)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b)) + 2*sqrt(2)*sqrt(a)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b)) + sqrt(2)*a^(1/4)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/b^(1/4) - sqrt(2)*a^(1/4)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/b + 2*sqrt(x)/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.59

$$\int \frac{x^{3/2}}{a + bx^2} dx = -\frac{\sqrt{2}(ab^3)^{1/4} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} + 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{2b^2} - \frac{\sqrt{2}(ab^3)^{1/4} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} - 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{2b^2} - \frac{\sqrt{2}(ab^3)^{1/4} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{4b^2} + \frac{\sqrt{2}(ab^3)^{1/4} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{4b^2} + \frac{2\sqrt{x}}{b}$$

input `integrate(x^(3/2)/(b*x^2+a),x, algorithm="giac")`

output
$$-1/2*\sqrt{2}*(a*b^3)^{1/4}*\arctan(1/2*\sqrt{2}*(\sqrt{2}*(a/b)^{1/4} + 2*\sqrt{x})/(a/b)^{1/4})/b^2 - 1/2*\sqrt{2}*(a*b^3)^{1/4}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*(a/b)^{1/4} - 2*\sqrt{x})/(a/b)^{1/4})/b^2 - 1/4*\sqrt{2}*(a*b^3)^{1/4}*\log(\sqrt{2}*\sqrt{x}*(a/b)^{1/4} + x + \sqrt{a/b})/b^2 + 1/4*\sqrt{2}*(a*b^3)^{1/4}*\log(-\sqrt{2}*\sqrt{x}*(a/b)^{1/4} + x + \sqrt{a/b})/b^2 + 2*\sqrt{x}/b$$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.49

$$\int \frac{x^{3/2}}{a + bx^2} dx = \frac{2\sqrt{x}}{b} - \frac{(-a)^{1/4} \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{b^{5/4}} - \frac{(-a)^{1/4} \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{b^{5/4}}$$

input `int(x^(3/2)/(a + b*x^2),x)`

output
$$(2*x^{1/2})/b - ((-a)^{1/4}*\operatorname{atan}((b^{1/4}*x^{1/2})/(-a)^{1/4}))/b^{5/4} - ((-a)^{1/4}*\operatorname{atanh}((b^{1/4}*x^{1/2})/(-a)^{1/4}))/b^{5/4}$$

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.29

$$\int \frac{x^{3/2}}{a + bx^2} dx = \frac{2b^{\frac{3}{4}}a^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) - 2b^{\frac{3}{4}}a^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) + b^{\frac{3}{4}}a^{\frac{1}{4}}\sqrt{2} \log(-\sqrt{x}b)}{4b^2}$$

input `int(x^(3/2)/(b*x^2+a),x)`

output

```
(2*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))) - 2*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))) + b**(3/4)*a**(1/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x) - b**(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x) + 8*sqrt(x)*b)/(4*b**2)
```

3.124 $\int \frac{x^{5/2}}{a+bx^2} dx$

Optimal result	810
Mathematica [A] (verified)	810
Rubi [B] (verified)	811
Maple [A] (verified)	815
Fricas [C] (verification not implemented)	815
Sympy [A] (verification not implemented)	816
Maxima [B] (verification not implemented)	817
Giac [A] (verification not implemented)	818
Mupad [B] (verification not implemented)	818
Reduce [B] (verification not implemented)	819

Optimal result

Integrand size = 15, antiderivative size = 113

$$\int \frac{x^{5/2}}{a+bx^2} dx = \frac{x^{3/2}}{b} - \frac{a \left(\arctan \left(\frac{\sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}} - x} \right) - \log \left(\frac{\sqrt{\frac{a}{b}} + \sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x} + x}{\sqrt{a+bx^2}} \right) \right)}{\sqrt{2} \sqrt[4]{\frac{a}{b}} b^2}$$

output

```
x^(3/2)/b-1/2*a*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))-ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/(a/b)^(1/4)/b^2
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.05

$$\int \frac{x^{5/2}}{a+bx^2} dx = \frac{4b^{3/4}x^{3/2} + 3\sqrt{2}a^{3/4} \arctan \left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}} \right) + 3\sqrt{2}a^{3/4} \operatorname{arctanh} \left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a+\sqrt{bx}}} \right)}{6b^{7/4}}$$

input

```
Integrate[x^(5/2)/(a + b*x^2),x]
```

output

$$(4*b^{(3/4)}*x^{(3/2)} + 3*\text{Sqrt}[2]*a^{(3/4)}*\text{ArcTan}[(\text{Sqrt}[a] - \text{Sqrt}[b]*x)/(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*\text{Sqrt}[x])] + 3*\text{Sqrt}[2]*a^{(3/4)}*\text{ArcTanh}[(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*\text{Sqrt}[x])/(\text{Sqrt}[a] + \text{Sqrt}[b]*x))]/(6*b^{(7/4)})$$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 231 vs. $2(113) = 226$.

Time = 0.42 (sec) , antiderivative size = 231, normalized size of antiderivative = 2.04, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {262, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^{5/2}}{a + bx^2} dx \\ & \quad \downarrow 262 \\ & \frac{2x^{3/2}}{3b} - \frac{a \int \frac{\sqrt{x}}{bx^2+a} dx}{b} \\ & \quad \downarrow 266 \\ & \frac{2x^{3/2}}{3b} - \frac{2a \int \frac{x}{bx^2+a} d\sqrt{x}}{b} \\ & \quad \downarrow 826 \\ & \frac{2x^{3/2}}{3b} - \frac{2a \left(\frac{\int \frac{\sqrt{bx} + \sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} \right)}{b} \\ & \quad \downarrow 1476 \\ & \frac{2x^{3/2}}{3b} - \frac{2a \left(\frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}} + \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} \right)}{b} \\ & \quad \downarrow 1082 \end{aligned}$$

$$\begin{aligned}
 & \frac{2x^{3/2}}{3b} - \frac{2a \left(\frac{\int \frac{1}{-x-1} d \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1 \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right)}{b} \\
 & \quad \downarrow \text{217} \\
 & \frac{2x^{3/2}}{3b} - \frac{2a \left(\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1 \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right)}{b} \\
 & \quad \downarrow \text{1479} \\
 & \frac{2x^{3/2}}{3b} - \frac{2a \left(\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1 \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a} \right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right)}{b} \\
 & \quad \downarrow \text{25} \\
 & \frac{2x^{3/2}}{3b} - \frac{2a \left(\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1 \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a} \right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right)}{b} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\begin{array}{c}
 \frac{2x^{3/2}}{3b} - \\
 2a \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}{2\sqrt{b}}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt[4]{b}}+ \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x} - \int \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}}{x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt[4]{b}}+ \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}}{x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt[4]{b}}+ \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt{b}} \right) \\
 \hline
 b \\
 \downarrow \text{1103} \\
 \frac{2x^{3/2}}{3b} - \\
 2a \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right) - \arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}{2\sqrt{b}}} - \frac{\log\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}\right) - \log\left(\frac{-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}\right)}{2\sqrt{b}} \right) \\
 \hline
 b
 \end{array}$$

input `Int[x^(5/2)/(a + b*x^2),x]`

output $(2x^{3/2})/(3b) - (2a*((-(\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{1/4}*\text{Sqrt}[x])/a^{1/4}]/(\text{Sqrt}[2]*a^{1/4}*b^{1/4}))) + \text{ArcTan}[1 + (\text{Sqrt}[2]*b^{1/4}*\text{Sqrt}[x])/a^{1/4}]/(\text{Sqrt}[2]*a^{1/4}*b^{1/4})))/(2*\text{Sqrt}[b]) - (-1/2*\text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{1/4}*b^{1/4}*\text{Sqrt}[x] + \text{Sqrt}[b]*x]/(\text{Sqrt}[2]*a^{1/4}*b^{1/4}) + \text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{1/4}*b^{1/4}*\text{Sqrt}[x] + \text{Sqrt}[b]*x]/(2*\text{Sqrt}[2]*a^{1/4}*b^{1/4}))/ (2*\text{Sqrt}[b]))/b$

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 $\text{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 262 $\text{Int}[(c_.)*(x_.)^m*((a_.) + (b_.)*(x_.)^2)^p], x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a + b*x^2)^{(p+1})/(b*(m+2*p+1))), x] - \text{Simp}[a*c^2*((m-1)/(b*(m+2*p+1))) \ \text{Int}[(c*x)^{(m-2)}*(a + b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{GtQ}[m, 2-1] \ \&\& \ \text{NeQ}[m+2*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 266 $\text{Int}[(c_.)*(x_.)^m*((a_.) + (b_.)*(x_.)^2)^p], x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \ \text{Subst}[\text{Int}[x^{k*(m+1)-1}*(a + b*(x^{2*k}/c^2))^p, x], x, (c*x)^{(1/k)}, x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$

rule 826 $\text{Int}[(x_.)^2/((a_.) + (b_.)*(x_.)^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \ \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Simp}[1/(2*s) \ \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

rule 1082 $\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_.) + (e_.)*(x_.)]/((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$

rule 1476 $\text{Int}[(d_.) + (e_.)*(x_.)^2]/((a_.) + (c_.)*(x_.)^4), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Simp}[e/(2*c) \ \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Simp}[e/(2*c) \ \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[d*e]$

rule 1479

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.03

method	result	size
derivativedivides	$\frac{2x^{\frac{3}{2}}}{3b} - \frac{a\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x + \sqrt{\frac{a}{b}}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x + x}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} + 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} - 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) \right)}{4b^2 \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	116
default	$\frac{2x^{\frac{3}{2}}}{3b} - \frac{a\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x + \sqrt{\frac{a}{b}}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x + x}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} + 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} - 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) \right)}{4b^2 \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	116
risch	$\frac{2x^{\frac{3}{2}}}{3b} - \frac{a\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x + \sqrt{\frac{a}{b}}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x + x}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} + 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} - 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) \right)}{4b^2 \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	116

input

```
int(x^(5/2)/(b*x^2+a), x, method=_RETURNVERBOSE)
```

output

```
2/3*x^(3/2)/b-1/4*a/b^2/(a/b)^(1/4)*2^(1/2)*(ln((x-2^(1/2)*(a/b)^(1/4)*x^(1/2)+(a/b)^(1/2))/(a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x))+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)+1)+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)-1))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.12 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.40

$$\int \frac{x^{5/2}}{a + bx^2} dx =$$

$$3b \left(-\frac{a^3}{b^7}\right)^{\frac{1}{4}} \log \left(b^5 \left(-\frac{a^3}{b^7}\right)^{\frac{3}{4}} + a^2 \sqrt{x} \right) - 3ib \left(-\frac{a^3}{b^7}\right)^{\frac{1}{4}} \log \left(ib^5 \left(-\frac{a^3}{b^7}\right)^{\frac{3}{4}} + a^2 \sqrt{x} \right) + 3ib \left(-\frac{a^3}{b^7}\right)^{\frac{1}{4}} \log \left(-ib^5 \right)$$

6b

input `integrate(x^(5/2)/(b*x^2+a),x, algorithm="fricas")`

output `-1/6*(3*b*(-a^3/b^7)^(1/4)*log(b^5*(-a^3/b^7)^(3/4) + a^2*sqrt(x)) - 3*I*b*(-a^3/b^7)^(1/4)*log(I*b^5*(-a^3/b^7)^(3/4) + a^2*sqrt(x)) + 3*I*b*(-a^3/b^7)^(1/4)*log(-I*b^5*(-a^3/b^7)^(3/4) + a^2*sqrt(x)) - 3*b*(-a^3/b^7)^(1/4)*log(-b^5*(-a^3/b^7)^(3/4) + a^2*sqrt(x)) - 4*x^(3/2))/b`

Sympy [A] (verification not implemented)

Time = 4.81 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.10

$$\int \frac{x^{5/2}}{a + bx^2} dx = \begin{cases} \infty x^{\frac{3}{2}} & \text{for } a = 0 \wedge b = 0 \\ \frac{2x^{\frac{7}{2}}}{7a} & \text{for } b = 0 \\ \frac{2x^{\frac{3}{2}}}{3b} & \text{for } a = 0 \\ -\frac{a \log\left(\sqrt{x} - \sqrt[4]{-\frac{a}{b}}\right)}{2b^2 \sqrt[4]{-\frac{a}{b}}} + \frac{a \log\left(\sqrt{x} + \sqrt[4]{-\frac{a}{b}}\right)}{2b^2 \sqrt[4]{-\frac{a}{b}}} - \frac{a \operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{b^2 \sqrt[4]{-\frac{a}{b}}} + \frac{2x^{\frac{3}{2}}}{3b} & \text{otherwise} \end{cases}$$

input `integrate(x**(5/2)/(b*x**2+a),x)`

output `Piecewise((zoo*x**(3/2), Eq(a, 0) & Eq(b, 0)), (2*x**(7/2)/(7*a), Eq(b, 0)), (2*x**(3/2)/(3*b), Eq(a, 0)), (-a*log(sqrt(x) - (-a/b)**(1/4))/(2*b**2*(-a/b)**(1/4)) + a*log(sqrt(x) + (-a/b)**(1/4))/(2*b**2*(-a/b)**(1/4)) - a*atan(sqrt(x)/(-a/b)**(1/4))/(b**2*(-a/b)**(1/4)) + 2*x**(3/2)/(3*b), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(90) = 180$.

Time = 0.12 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.65

$$\int \frac{x^{5/2}}{a + bx^2} dx =$$

$$\frac{a \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}a^{1/4}b^{1/4} + 2\sqrt{b}\sqrt{x}\right)}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} \right) + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}a^{1/4}b^{1/4} - 2\sqrt{b}\sqrt{x}\right)}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} - \frac{\sqrt{2} \log\left(\sqrt{2}a^{1/4}b^{1/4}\sqrt{x} + \sqrt{b}x + \sqrt{a}\right)}{a^{1/4}b^{3/4}} + \frac{\sqrt{2} \log\left(-\sqrt{2}a^{1/4}b^{1/4}\sqrt{x} + \sqrt{b}x + \sqrt{a}\right)}{a^{1/4}b^{3/4}}}{4b} + \frac{2x^{3/2}}{3b}$$

input `integrate(x^(5/2)/(b*x^2+a),x, algorithm="maxima")`

output `-1/4*a*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b)) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b)) - sqrt(2)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4)) + sqrt(2)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4))/b + 2/3*x^(3/2)/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.58

$$\int \frac{x^{5/2}}{a + bx^2} dx = \frac{2x^{3/2}}{3b} - \frac{\sqrt{2}(ab^3)^{3/4} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} + 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{2b^4}$$

$$- \frac{\sqrt{2}(ab^3)^{3/4} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} - 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{2b^4}$$

$$+ \frac{\sqrt{2}(ab^3)^{3/4} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{4b^4}$$

$$- \frac{\sqrt{2}(ab^3)^{3/4} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{4b^4}$$

input `integrate(x^(5/2)/(b*x^2+a),x, algorithm="giac")`output `2/3*x^(3/2)/b - 1/2*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/b^4 - 1/2*sqrt(2)*(a*b^3)^(3/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/b^4 + 1/4*sqrt(2)*(a*b^3)^(3/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/b^4 - 1/4*sqrt(2)*(a*b^3)^(3/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/b^4`**Mupad [B] (verification not implemented)**

Time = 0.19 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.48

$$\int \frac{x^{5/2}}{a + bx^2} dx = \frac{2x^{3/2}}{3b} + \frac{(-a)^{3/4} \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{b^{7/4}} - \frac{(-a)^{3/4} \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{b^{7/4}}$$

input `int(x^(5/2)/(a + b*x^2),x)`output `(2*x^(3/2))/(3*b) + ((-a)^(3/4)*atan((b^(1/4)*x^(1/2))/(-a)^(1/4)))/b^(7/4) - ((-a)^(3/4)*atanh((b^(1/4)*x^(1/2))/(-a)^(1/4)))/b^(7/4)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.30

$$\int \frac{x^{5/2}}{a + bx^2} dx = \frac{6b^{1/4}a^{3/4}\sqrt{2} \operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{1/4}a^{1/4}\sqrt{2}}\right) - 6b^{1/4}a^{3/4}\sqrt{2} \operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}+2\sqrt{x}\sqrt{b}}{b^{1/4}a^{1/4}\sqrt{2}}\right) - 3b^{1/4}a^{3/4}\sqrt{2} \log\left(-\sqrt{x}\right)}{12b^2}$$

input `int(x^(5/2)/(b*x^2+a),x)`

output

```
(6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))) - 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))) - 3*b**(1/4)*a**(3/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x) + 3*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x) + 8*sqrt(x)*b*x)/(12*b**2)
```

3.125 $\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx$

Optimal result	820
Mathematica [A] (verified)	821
Rubi [A] (verified)	821
Maple [A] (verified)	825
Fricas [C] (verification not implemented)	826
Sympy [B] (verification not implemented)	827
Maxima [A] (verification not implemented)	828
Giac [A] (verification not implemented)	829
Mupad [B] (verification not implemented)	829
Reduce [B] (verification not implemented)	830

Optimal result

Integrand size = 15, antiderivative size = 127

$$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx = \frac{\sqrt{x}}{2a(a+bx^2)} + \frac{3 \left(\arctan \left(\frac{\sqrt[4]{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}-x}} \right) + \log \left(\frac{\sqrt{\frac{a}{b}+\sqrt{2}} \sqrt[4]{\frac{a}{b}} \sqrt{x+x}}{\sqrt{a+bx^2}} \right) \right)}{4\sqrt{2}a \left(\frac{a}{b}\right)^{3/4} b}$$

output

```
1/2*x^(1/2)/a/(b*x^2+a)+3/8*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))+ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/a/(a/b)^(3/4)/b
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.01

$$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx = \frac{\frac{4a^{3/4}\sqrt{x}}{a+bx^2} - \frac{3\sqrt{2} \arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right)}{\sqrt[4]{b}} + \frac{3\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}+\sqrt{bx}}\right)}{\sqrt[4]{b}}}{8a^{7/4}}$$

input `Integrate[1/(Sqrt[x]*(a + b*x^2)^2), x]`output `((4*a^(3/4)*Sqrt[x])/(a + b*x^2) - (3*Sqrt[2]*ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x]])/b^(1/4) + (3*Sqrt[2]*ArcTanh[(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])/(Sqrt[a] + Sqrt[b]*x)]/b^(1/4))/(8*a^(7/4))`**Rubi [A] (verified)**Time = 0.42 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.90, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {253, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{x}(a+bx^2)^2} dx \\ & \quad \downarrow \text{253} \\ & \frac{3 \int \frac{1}{\sqrt{x}(bx^2+a)} dx}{4a} + \frac{\sqrt{x}}{2a(a+bx^2)} \\ & \quad \downarrow \text{266} \\ & \frac{3 \int \frac{1}{bx^2+a} d\sqrt{x}}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} \\ & \quad \downarrow \text{755} \end{aligned}$$

$$3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

1476

$$3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\int \frac{1}{x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}}}{2\sqrt{a}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

1082

$$3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\int \frac{1}{x-1} d\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{x-1} d\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

217

$$3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

1479

$$3 \left(\frac{\frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

↓ 25

$$3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a} \right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\frac{2a}{\sqrt{x}}}{2a(a+bx^2)}$$

↓ 27

$$3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}}{x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2 \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\frac{2a}{\sqrt{x}}}{2a(a+bx^2)}$$

↓ 1103

$$3 \left(\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\log \left(\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx} \right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\log \left(-\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx} \right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\frac{2a}{\sqrt{x}}}{2a(a+bx^2)}$$

input `Int [1/(Sqrt [x]*(a + b*x^2)^2), x]`

output

$$\frac{\sqrt{x}}{2a(a + bx^2)} + \frac{3\left(-\operatorname{ArcTan}\left[\frac{1 - (\sqrt{2})^{1/4}\sqrt{x}}{a^{1/4}}\right]/(\sqrt{2})^{1/4}b^{1/4}\right) + \operatorname{ArcTan}\left[\frac{1 + (\sqrt{2})^{1/4}\sqrt{x}}{a^{1/4}}\right]/(\sqrt{2})^{1/4}b^{1/4}}{2\sqrt{a}} + \frac{-1/2\log[\sqrt{a} - \sqrt{2}a^{1/4}b^{1/4}\sqrt{x}] + \sqrt{bx}}{(\sqrt{2})^{1/4}b^{1/4}} + \frac{\log[\sqrt{a} + \sqrt{2}a^{1/4}b^{1/4}\sqrt{x}] + \sqrt{bx}}{2\sqrt{2}a^{1/4}b^{1/4}}}{2a}$$

Defintions of rubi rules used

rule 25

$$\operatorname{Int}[-(Fx_), x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[Fx, x], x]$$

rule 27

$$\operatorname{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[Fx, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ \text{!MatchQ}[Fx, (b_*)(Gx_)] \text{ ; FreeQ}[b, x]$$

rule 217

$$\operatorname{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[-b, 2])^{-1})*\operatorname{ArcTan}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[-a, 2])], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{PosQ}[a/b] \ \&\& \ (\operatorname{LtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$$

rule 253

$$\operatorname{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \operatorname{Simp}[(-c*x)^{(m+1)}((a + b*x^2)^{(p+1})/(2*a*c*(p+1))), x] + \operatorname{Simp}[(m + 2*p + 3)/(2*a*(p + 1)) \operatorname{Int}[(c*x)^m*(a + b*x^2)^{(p+1)}, x], x] \text{ ; FreeQ}[\{a, b, c, m\}, x] \ \&\& \ \operatorname{LtQ}[p, -1] \ \&\& \ \operatorname{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 266

$$\operatorname{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^2)^{(p_*)}, x_Symbol] \rightarrow \operatorname{With}[\{k = \operatorname{Denominator}[m]\}, \operatorname{Simp}[k/c \operatorname{Subst}[\operatorname{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{2*k}/c^2))^p, x], x, (c*x)^{(1/k)}], x]] \text{ ; FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \operatorname{FractionQ}[m] \ \&\& \ \operatorname{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 755

$$\operatorname{Int}[(a_*) + (b_*)(x_)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numerator}[\operatorname{Rt}[a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[a/b, 2]]\}, \operatorname{Simp}[1/(2*r) \operatorname{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \operatorname{Simp}[1/(2*r) \operatorname{Int}[(r + s*x^2)/(a + b*x^4), x], x]] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ (\operatorname{GtQ}[a/b, 0] \ || \ (\operatorname{PosQ}[a/b] \ \&\& \ \operatorname{AtomQ}[\operatorname{SplitProduct}[\operatorname{SumBaseQ}, a]] \ \&\& \ \operatorname{AtomQ}[\operatorname{SplitProduct}[\operatorname{SumBaseQ}, b]]))$$

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.98

method	result	size
derivativedivides	$\frac{\sqrt{x}}{2a(x^2b+a)} + \frac{3\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\sqrt{\frac{a}{b}}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1}\right)\right)}{16a^2}$	124
default	$\frac{\sqrt{x}}{2a(x^2b+a)} + \frac{3\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\sqrt{\frac{a}{b}}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1}\right)\right)}{16a^2}$	124

input `int(1/x^(1/2)/(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output

$$\frac{1}{2}x^{1/2}/a/(b*x^2+a)+3/16/a^2*(a/b)^{1/4}*2^{1/2}*(\ln(((a/b)^{1/2}+2^{1/2}*(1/2)*(a/b)^{1/4}*x^{1/2}+x)/(x-2^{1/2)*(a/b)^{1/4}*x^{1/2}+(a/b)^{1/2}))) + 2*\arctan(2^{1/2)/(a/b)^{1/4}*x^{1/2}+1)+2*\arctan(2^{1/2)/(a/b)^{1/4}*x^{1/2}-1))$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.52

$$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx$$

$$= \frac{3(abx^2+a^2)\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} \log\left(a^2\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} + \sqrt{x}\right) - 3(-i abx^2 - i a^2)\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} \log\left(i a^2\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} + \sqrt{x}\right) - 3(i abx^2 + i a^2)\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} \log\left(-i a^2\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} + \sqrt{x}\right) - 3(-i abx^2 - i a^2)\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} \log\left(-i a^2\left(-\frac{1}{a^7b}\right)^{\frac{1}{4}} + \sqrt{x}\right)}{8(abx^2+a^2)^2}$$

input

```
integrate(1/x^(1/2)/(b*x^2+a)^2,x, algorithm="fricas")
```

output

$$\frac{1}{8}*(3*(a*b*x^2 + a^2)*(-1/(a^7*b))^{1/4}*\log(a^2*(-1/(a^7*b))^{1/4} + \sqrt{x}) - 3*(-I*a*b*x^2 - I*a^2)*(-1/(a^7*b))^{1/4}*\log(I*a^2*(-1/(a^7*b))^{1/4} + \sqrt{x}) - 3*(I*a*b*x^2 + I*a^2)*(-1/(a^7*b))^{1/4}*\log(-I*a^2*(-1/(a^7*b))^{1/4} + \sqrt{x}) - 3*(a*b*x^2 + a^2)*(-1/(a^7*b))^{1/4}*\log(-a^2*(-1/(a^7*b))^{1/4} + \sqrt{x}) + 4*\sqrt{x})/(a*b*x^2 + a^2)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 316 vs. $2(100) = 200$.

Time = 25.08 (sec) , antiderivative size = 316, normalized size of antiderivative = 2.49

$$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx$$

$$= \begin{cases} \frac{\infty}{x^{\frac{7}{2}}} \\ \frac{2\sqrt{x}}{a^2} \\ -\frac{2}{7b^2x^{\frac{7}{2}}} \\ \frac{4a\sqrt{x}}{8a^3+8a^2bx^2} - \frac{3a\sqrt[4]{-\frac{a}{b}}\log\left(\sqrt{x}-\sqrt[4]{-\frac{a}{b}}\right)}{8a^3+8a^2bx^2} + \frac{3a\sqrt[4]{-\frac{a}{b}}\log\left(\sqrt{x}+\sqrt[4]{-\frac{a}{b}}\right)}{8a^3+8a^2bx^2} + \frac{6a\sqrt[4]{-\frac{a}{b}}\operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{8a^3+8a^2bx^2} - \frac{3bx^2\sqrt[4]{-\frac{a}{b}}\log}{8a^3+8a^2bx^2} \end{cases}$$

input `integrate(1/x**(1/2)/(b*x**2+a)**2,x)`

output `Piecewise((zoo/x**(7/2), Eq(a, 0) & Eq(b, 0)), (2*sqrt(x)/a**2, Eq(b, 0)), (-2/(7*b**2*x**(7/2)), Eq(a, 0)), (4*a*sqrt(x)/(8*a**3 + 8*a**2*b*x**2) - 3*a*(-a/b)**(1/4)*log(sqrt(x) - (-a/b)**(1/4))/(8*a**3 + 8*a**2*b*x**2) + 3*a*(-a/b)**(1/4)*log(sqrt(x) + (-a/b)**(1/4))/(8*a**3 + 8*a**2*b*x**2) + 6*a*(-a/b)**(1/4)*atan(sqrt(x)/(-a/b)**(1/4))/(8*a**3 + 8*a**2*b*x**2) - 3*b*x**2*(-a/b)**(1/4)*log(sqrt(x) - (-a/b)**(1/4))/(8*a**3 + 8*a**2*b*x**2) + 3*b*x**2*(-a/b)**(1/4)*log(sqrt(x) + (-a/b)**(1/4))/(8*a**3 + 8*a**2*b*x**2) + 6*b*x**2*(-a/b)**(1/4)*atan(sqrt(x)/(-a/b)**(1/4))/(8*a**3 + 8*a**2*b*x**2), True))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.53

$$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx$$

$$= \frac{3 \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}+2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} \right) + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}-2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{\sqrt{2} \log(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x}+\sqrt{bx}+\sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x}+\sqrt{bx}+\sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}}}{16a} + \frac{\sqrt{x}}{2(abx^2+a^2)}$$

input `integrate(1/x^(1/2)/(b*x^2+a)^2,x, algorithm="maxima")`output `3/16*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + sqrt(2)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4)) - sqrt(2)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4)) / a + 1/2*sqrt(x)/(a*b*x^2 + a^2)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.57

$$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx = \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}+2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{8a^2b} + \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}-2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{8a^2b} + \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{16a^2b} - \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{16a^2b} + \frac{\sqrt{x}}{2(bx^2+a)a}$$

input `integrate(1/x^(1/2)/(b*x^2+a)^2,x, algorithm="giac")`output `3/8*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/(a^2*b) + 3/8*sqrt(2)*(a*b^3)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/(a^2*b) + 3/16*sqrt(2)*(a*b^3)^(1/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^2*b) - 3/16*sqrt(2)*(a*b^3)^(1/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^2*b) + 1/2*sqrt(x)/((b*x^2 + a)*a)`**Mupad [B] (verification not implemented)**

Time = 0.20 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.50

$$\int \frac{1}{\sqrt{x}(a+bx^2)^2} dx = \frac{\sqrt{x}}{2a(bx^2+a)} + \frac{3 \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{7/4}b^{1/4}} + \frac{3 \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{7/4}b^{1/4}}$$

input `int(1/(x^(1/2)*(a + b*x^2)^2),x)`

output

$$x^{1/2}/(2*a*(a + b*x^2)) + (3*atan((b^{1/4}*x^{1/2})/(-a)^{1/4}))/(-4*(-a)^{7/4}*b^{1/4}) + (3*atanh((b^{1/4}*x^{1/2})/(-a)^{1/4}))/(-4*(-a)^{7/4}*b^{1/4})$$

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.41

$$\int \frac{1}{\sqrt{x}(a + bx^2)^2} dx$$

$$= \frac{-6b^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) - 6b^{\frac{7}{4}}a^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^2 + 6b^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right)}{\dots}$$

input

int(1/x^(1/2)/(b*x^2+a)^2,x)

output

$$\begin{aligned} & (-6b^{3/4}a^{5/4}\sqrt{2}\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{x}\sqrt{b})/\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2})^2 - 6b^{7/4}a^{1/4}\sqrt{2}\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2})^2 \\ & + 6b^{3/4}a^{5/4}\sqrt{2}\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2}+2\sqrt{x}\sqrt{b})/\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2})^2 + 6b^{3/4}a^{5/4}\sqrt{2}\operatorname{atan}(b^{1/4}a^{1/4}\sqrt{2}+2\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2})^2 \\ & - 3b^{3/4}a^{1/4}\sqrt{2}\log(-\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2}) + \sqrt{a} + \sqrt{b}\sqrt{x} - 3b^{3/4}a^{1/4}\sqrt{2}\log(-\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2}) + \sqrt{a} + \sqrt{b}\sqrt{x} \\ & + 3b^{3/4}a^{1/4}\sqrt{2}\log(\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2}) + \sqrt{a} + \sqrt{b}\sqrt{x} + 3b^{3/4}a^{1/4}\sqrt{2}\log(\sqrt{x}\sqrt{b})/(b^{1/4}a^{1/4}\sqrt{2}) + \sqrt{a} + \sqrt{b}\sqrt{x} \\ & + 8\sqrt{x}\sqrt{a}\sqrt{b}/(16a^2b(a + b^2x^2)) \end{aligned}$$

3.126 $\int \frac{\sqrt{x}}{(a+bx^2)^2} dx$

Optimal result	831
Mathematica [A] (verified)	831
Rubi [A] (verified)	832
Maple [A] (verified)	836
Fricas [C] (verification not implemented)	836
Sympy [B] (verification not implemented)	837
Maxima [A] (verification not implemented)	838
Giac [A] (verification not implemented)	838
Mupad [B] (verification not implemented)	839
Reduce [B] (verification not implemented)	839

Optimal result

Integrand size = 15, antiderivative size = 129

$$\int \frac{\sqrt{x}}{(a+bx^2)^2} dx = \frac{x^{3/2}}{2a(a+bx^2)} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{\frac{a}{b}}\sqrt{x}}{\sqrt{\frac{a}{b}-x}}\right) - \log\left(\frac{\sqrt{\frac{a}{b}+\sqrt{2}\sqrt[4]{\frac{a}{b}}\sqrt{x}+x}}{\sqrt{a+bx^2}}\right)}{4\sqrt{2}a\sqrt[4]{\frac{a}{b}}b}$$

output

```
1/2*x^(3/2)/a/(b*x^2+a)+1/8*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))-ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/a/(a/b)^(1/4)/b
```

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.99

$$\int \frac{\sqrt{x}}{(a+bx^2)^2} dx = \frac{4\sqrt[4]{a}x^{3/2}}{a+bx^2} - \frac{\sqrt{2}\arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right)}{b^{3/4}} - \frac{\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a+\sqrt{bx}}}\right)}{b^{3/4}}}{8a^{5/4}}$$

input `Integrate[Sqrt[x]/(a + b*x^2)^2,x]`

output $((4*a^{(1/4)}*x^{(3/2)})/(a + b*x^2) - (\text{Sqrt}[2]*\text{ArcTan}[(\text{Sqrt}[a] - \text{Sqrt}[b]*x)/(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*\text{Sqrt}[x])])/b^{(3/4)} - (\text{Sqrt}[2]*\text{ArcTanh}[(\text{Sqrt}[2]*a^{(1/4)}*b^{(1/4)}*\text{Sqrt}[x])/(\text{Sqrt}[a] + \text{Sqrt}[b]*x])/b^{(3/4)})/(8*a^{(5/4)})$

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.87, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {253, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x}}{(a + bx^2)^2} dx \\
 & \quad \downarrow \text{253} \\
 & \frac{\int \frac{\sqrt{x}}{bx^2+a} dx}{4a} + \frac{x^{3/2}}{2a(a + bx^2)} \\
 & \quad \downarrow \text{266} \\
 & \frac{\int \frac{x}{bx^2+a} d\sqrt{x}}{2a} + \frac{x^{3/2}}{2a(a + bx^2)} \\
 & \quad \downarrow \text{826} \\
 & \frac{\int \frac{\sqrt{bx} + \sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a + bx^2)} \\
 & \quad \downarrow \text{1476} \\
 & \frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \frac{\sqrt{a}}{\sqrt{b}}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \frac{\sqrt{a}}{\sqrt{b}}}} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a + bx^2)} \\
 & \quad \downarrow \text{1082}
 \end{aligned}$$

$$\frac{\int \frac{1}{-x-1} d\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a + bx^2)}$$

217

$$\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right) - \arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a + bx^2)}$$

1479

$$\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right) - \arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{2a x^{3/2}}{2a(a + bx^2)}$$

25

$$\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right) - \arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{2a x^{3/2}}{2a(a + bx^2)}$$

27

$$\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1\right) - \arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}}{x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2 \sqrt[4]{a} \sqrt[4]{b}} + \frac{2a x^{3/2}}{2a(a + bx^2)}$$

1103

$$\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} - \frac{\frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} + \frac{2a}{x^{3/2}}}{2a(a+bx^2)}$$

input `Int[Sqrt[x]/(a + b*x^2)^2,x]`

output `x^(3/2)/(2*a*(a + b*x^2)) + ((-(ArcTan[1 - (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]) - (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*a)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 253 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Simp[(m + 2*p + 3)/(2*a*(p + 1) Int[(c*x)^m*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && LtQ[p, -1] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.98

method	result	size
derivativedivides	$\frac{x^{\frac{3}{2}}}{2a(x^2b+a)} + \frac{\sqrt{2} \left(\ln \left(\frac{x-\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+\sqrt{\frac{a}{b}}}}{\sqrt{\frac{a}{b}+\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+x}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}} + 1} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}} - 1} \right) \right)}{16ab \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	127
default	$\frac{x^{\frac{3}{2}}}{2a(x^2b+a)} + \frac{\sqrt{2} \left(\ln \left(\frac{x-\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+\sqrt{\frac{a}{b}}}}{\sqrt{\frac{a}{b}+\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+x}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}} + 1} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}} - 1} \right) \right)}{16ab \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	127

input `int(x^(1/2)/(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output `1/2*x^(3/2)/a/(b*x^2+a)+1/16/a/b/(a/b)^(1/4)*2^(1/2)*(ln((x-2^(1/2)*(a/b)^(1/4)*x^(1/2)+(a/b)^(1/2))/((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x))+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)+1)+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)-1))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.58

$$\int \frac{\sqrt{x}}{(a+bx^2)^2} dx$$

$$= \frac{(abx^2+a^2)\left(-\frac{1}{a^5b^3}\right)^{\frac{1}{4}} \log\left(a^4b^2\left(-\frac{1}{a^5b^3}\right)^{\frac{3}{4}}+\sqrt{x}\right) - (i abx^2 + i a^2)\left(-\frac{1}{a^5b^3}\right)^{\frac{1}{4}} \log\left(i a^4b^2\left(-\frac{1}{a^5b^3}\right)^{\frac{3}{4}}+\sqrt{x}\right) - (i abx^2 - i a^2)\left(-\frac{1}{a^5b^3}\right)^{\frac{1}{4}} \log\left(-i a^4b^2\left(-\frac{1}{a^5b^3}\right)^{\frac{3}{4}}+\sqrt{x}\right) + (i abx^2 + i a^2)\left(-\frac{1}{a^5b^3}\right)^{\frac{1}{4}} \log\left(-i a^4b^2\left(-\frac{1}{a^5b^3}\right)^{\frac{3}{4}}+\sqrt{x}\right)}{8}$$

input `integrate(x^(1/2)/(b*x^2+a)^2,x, algorithm="fricas")`

output

```
1/8*((a*b*x^2 + a^2)*(-1/(a^5*b^3))^(1/4)*log(a^4*b^2*(-1/(a^5*b^3))^(3/4)
+ sqrt(x)) - (I*a*b*x^2 + I*a^2)*(-1/(a^5*b^3))^(1/4)*log(I*a^4*b^2*(-1/(
a^5*b^3))^(3/4) + sqrt(x)) - (-I*a*b*x^2 - I*a^2)*(-1/(a^5*b^3))^(1/4)*log
(-I*a^4*b^2*(-1/(a^5*b^3))^(3/4) + sqrt(x)) - (a*b*x^2 + a^2)*(-1/(a^5*b^3
))^(1/4)*log(-a^4*b^2*(-1/(a^5*b^3))^(3/4) + sqrt(x)) + 4*x^(3/2)/(a*b*x^
2 + a^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 400 vs. $2(97) = 194$.

Time = 17.88 (sec) , antiderivative size = 400, normalized size of antiderivative = 3.10

$$\int \frac{\sqrt{x}}{(a + bx^2)^2} dx$$

$$= \begin{cases} \frac{\infty}{x^{\frac{5}{2}}} \\ \frac{2x^{\frac{3}{2}}}{3a^2} \\ -\frac{2}{5b^2x^{\frac{5}{2}}} \\ \frac{a \log\left(\sqrt{x} - \sqrt[4]{-\frac{a}{b}}\right)}{8a^2b^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}} + 8ab^2x^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}}} - \frac{a \log\left(\sqrt{x} + \sqrt[4]{-\frac{a}{b}}\right)}{8a^2b^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}} + 8ab^2x^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}}} + \frac{2a \operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{8a^2b^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}} + 8ab^2x^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}}} + \frac{4bx^{\frac{3}{2}}\sqrt[4]{-\frac{a}{b}}}{8a^2b^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}} + 8ab^2x^{\frac{4}{3}}\sqrt[4]{-\frac{a}{b}}} + \dots \end{cases}$$

input

```
integrate(x**(1/2)/(b*x**2+a)**2,x)
```

output

```
Piecewise((zoo/x**(5/2), Eq(a, 0) & Eq(b, 0)), (2*x**(3/2)/(3*a**2), Eq(b,
0)), (-2/(5*b**2*x**(5/2)), Eq(a, 0)), (a*log(sqrt(x) - (-a/b)**(1/4))/(8
*a**2*b*(-a/b)**(1/4) + 8*a*b**2*x**2*(-a/b)**(1/4)) - a*log(sqrt(x) + (-a
/b)**(1/4))/(8*a**2*b*(-a/b)**(1/4) + 8*a*b**2*x**2*(-a/b)**(1/4)) + 2*a*a
tan(sqrt(x)/(-a/b)**(1/4))/(8*a**2*b*(-a/b)**(1/4) + 8*a*b**2*x**2*(-a/b)*
*(1/4)) + 4*b*x**(3/2)*(-a/b)**(1/4)/(8*a**2*b*(-a/b)**(1/4) + 8*a*b**2*x
**2*(-a/b)**(1/4)) + b*x**2*log(sqrt(x) - (-a/b)**(1/4))/(8*a**2*b*(-a/b)**
(1/4) + 8*a*b**2*x**2*(-a/b)**(1/4)) - b*x**2*log(sqrt(x) + (-a/b)**(1/4))
/(8*a**2*b*(-a/b)**(1/4) + 8*a*b**2*x**2*(-a/b)**(1/4)) + 2*b*x**2*atan(sq
rt(x)/(-a/b)**(1/4))/(8*a**2*b*(-a/b)**(1/4) + 8*a*b**2*x**2*(-a/b)**(1/4
)), True))
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 194, normalized size of antiderivative = 1.50

$$\int \frac{\sqrt{x}}{(a+bx^2)^2} dx = \frac{x^{\frac{3}{2}}}{2(abx^2+a^2)} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}+2\sqrt{b}\sqrt{x}\right)}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}-2\sqrt{b}\sqrt{x}\right)}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} - \frac{\sqrt{2} \log\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x}+\sqrt{bx}+\sqrt{a}\right)}{a^{\frac{1}{4}}b^{\frac{3}{4}}} + \frac{\sqrt{2} \log\left(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x}+\sqrt{bx}+\sqrt{a}\right)}{a^{\frac{1}{4}}b^{\frac{3}{4}}}$$

input `integrate(x^(1/2)/(b*x^2+a)^2,x, algorithm="maxima")`

output

```
1/2*x^(3/2)/(a*b*x^2 + a^2) + 1/16*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*
a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*
sqrt(b))*sqrt(b) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4)
- 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b))*sqrt(b)
) - sqrt(2)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a
^(1/4)*b^(3/4)) + sqrt(2)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x
+ sqrt(a))/(a^(1/4)*b^(3/4))/a
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.54

$$\int \frac{\sqrt{x}}{(a+bx^2)^2} dx = \frac{x^{\frac{3}{2}}}{2(bx^2+a)a} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}+2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{8a^2b^3} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}-2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{8a^2b^3} - \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{16a^2b^3} + \frac{\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{16a^2b^3}$$

input `integrate(x^(1/2)/(b*x^2+a)^2,x, algorithm="giac")`

output `1/2*x^(3/2)/((b*x^2 + a)*a) + 1/8*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*
(sqrt(2)(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/(a^2*b^3) + 1/8*sqrt(2)*(a
*b^3)^(3/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1
/4))/(a^2*b^3) - 1/16*sqrt(2)*(a*b^3)^(3/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4
) + x + sqrt(a/b))/(a^2*b^3) + 1/16*sqrt(2)*(a*b^3)^(3/4)*log(-sqrt(2)*sqr
t(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^2*b^3)`

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.50

$$\int \frac{\sqrt{x}}{(a + bx^2)^2} dx = \frac{x^{3/2}}{2a(bx^2 + a)} - \frac{\operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{5/4}b^{3/4}} + \frac{\operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{5/4}b^{3/4}}$$

input `int(x^(1/2)/(a + b*x^2)^2,x)`

output `x^(3/2)/(2*a*(a + b*x^2)) - atan((b^(1/4)*x^(1/2))/(-a)^(1/4))/(4*(-a)^(5/
4)*b^(3/4)) + atanh((b^(1/4)*x^(1/2))/(-a)^(1/4))/(4*(-a)^(5/4)*b^(3/4))`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 305, normalized size of antiderivative = 2.36

$$\int \frac{\sqrt{x}}{(a + bx^2)^2} dx$$

$$= \frac{-2b^{\frac{1}{4}}a^{\frac{7}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) - 2b^{\frac{5}{4}}a^{\frac{3}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^2 + 2b^{\frac{1}{4}}a^{\frac{7}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}+2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right)}{\dots}$$

input `int(x^(1/2)/(b*x^2+a)^2,x)`

output

```
( - 2*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)
)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))*a - 2*b**(1/4)*a**(3/4)*sqrt(2)*at
an((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt
(2))*b*x**2 + 2*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2)
+ 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))*a + 2*b**(1/4)*a**(3/4)
*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a*
*(1/4)*sqrt(2))*b*x**2 + b**(1/4)*a**(3/4)*sqrt(2)*log( - sqrt(x)*b**(1/4)
)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a + b**(1/4)*a**(3/4)*sqrt(2)*lo
g( - sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b*x**2 - b**
(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + s
qrt(b)*x)*a - b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt
(2) + sqrt(a) + sqrt(b)*x)*b*x**2 + 8*sqrt(x)*a*b*x)/(16*a**2*b*(a + b*x**
2))
```

3.127 $\int \frac{x^{3/2}}{(a+bx^2)^2} dx$

Optimal result	841
Mathematica [A] (verified)	841
Rubi [A] (verified)	842
Maple [A] (verified)	846
Fricas [C] (verification not implemented)	846
Sympy [B] (verification not implemented)	847
Maxima [B] (verification not implemented)	847
Giac [B] (verification not implemented)	848
Mupad [B] (verification not implemented)	849
Reduce [B] (verification not implemented)	849

Optimal result

Integrand size = 15, antiderivative size = 124

$$\int \frac{x^{3/2}}{(a+bx^2)^2} dx = -\frac{\sqrt{x}}{2b(a+bx^2)} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{\frac{a}{b}}\sqrt{x}}{\sqrt{\frac{a}{b}}-x}\right) + \log\left(\frac{\sqrt{\frac{a}{b}}+\sqrt{2}\sqrt[4]{\frac{a}{b}}\sqrt{x+x}}{\sqrt{a+bx^2}}\right)}{4\sqrt{2}\left(\frac{a}{b}\right)^{3/4}b^2}$$

output

```
-1/2*x^(1/2)/b/(b*x^2+a)+1/8*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))+ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*^(1/2)/(a/b)^(3/4)/b^2
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.02

$$\int \frac{x^{3/2}}{(a+bx^2)^2} dx = \frac{-\frac{4\sqrt[4]{b}\sqrt{x}}{a+bx^2} - \frac{\sqrt{2}\arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right)}{a^{3/4}} + \frac{\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a+\sqrt{bx}}}\right)}{a^{3/4}}}{8b^{5/4}}$$

input

```
Integrate[x^(3/2)/(a + b*x^2)^2,x]
```

output

$$\left((-4b^{1/4} \sqrt{x}) / (a + bx^2) - (\sqrt{2} \operatorname{ArcTan}[(\sqrt{a} - \sqrt{b})x] / (\sqrt{2} a^{1/4} b^{1/4} \sqrt{x})) \right) / a^{3/4} + (\sqrt{2} \operatorname{ArcTanh}[(\sqrt{2} a^{1/4} b^{1/4} \sqrt{x}) / (\sqrt{a} + \sqrt{b})x]) / a^{3/4} / (8b^{5/4})$$

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.94, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {252, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^{3/2}}{(a + bx^2)^2} dx$$

$$\downarrow 252$$

$$\frac{\int \frac{1}{\sqrt{x}(bx^2+a)} dx}{4b} - \frac{\sqrt{x}}{2b(a + bx^2)}$$

$$\downarrow 266$$

$$\frac{\int \frac{1}{bx^2+a} d\sqrt{x}}{2b} - \frac{\sqrt{x}}{2b(a + bx^2)}$$

$$\downarrow 755$$

$$\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} - \frac{\sqrt{x}}{2b(a + bx^2)}$$

$$\downarrow 1476$$

$$\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{a}} - \frac{\sqrt{x}}{2b(a + bx^2)}$$

$$\downarrow 1082$$

$$\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{-x-1} d\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} - \frac{\sqrt{x}}{2b(a+bx^2)}$$

217

$$\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} - \frac{\sqrt{x}}{2b(a+bx^2)}$$

1479

$$\frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}}}{\frac{2b}{\sqrt{x}}}{2b(a+bx^2)}$$

25

$$\frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}}}{\frac{2b}{\sqrt{x}}}{2b(a+bx^2)}$$

27

$$\frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}}{x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}}+\frac{\sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt[4]{a}\sqrt[4]{b}} + \frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}}}{\frac{2b}{\sqrt{x}}}{2b(a+bx^2)}$$

1103

$$\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} + \frac{\frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} - \frac{\frac{2b}{\sqrt{x}}}{2b(a+bx^2)}$$

input `Int[x^(3/2)/(a + b*x^2)^2,x]`

output `-1/2*Sqrt[x]/(b*(a + b*x^2)) + ((-ArcTan[1 - (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[a]) + (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*b)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] :> Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 252 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[c*(c*x)^(m-1)*((a + b*x^2)^(p+1)/(2*b*(p+1))), x] - Simp[c^2*((m-1)/(2*b*(p+1))) Int[(c*x)^(m-2)*(a + b*x^2)^(p+1), x], x] /; FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m+2*p+3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.02

method	result	size
derivativedivides	$-\frac{\sqrt{x}}{2b(x^2b+a)} + \frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\frac{a}{b}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}-1\right)\right)}{16ba}$	127
default	$-\frac{\sqrt{x}}{2b(x^2b+a)} + \frac{\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\frac{a}{b}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}-1\right)\right)}{16ba}$	127

input `int(x^(3/2)/(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output
$$-1/2*x^{(1/2)}/b/(b*x^2+a)+1/16/b*(a/b)^{(1/4)}/a*2^{(1/2)}*(\ln(((a/b)^{(1/2)}+2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)+x})/(x-2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)}+(a/b)^{(1/2)}))+2*\arctan(2^{(1/2)}/(a/b)^{(1/4)}*x^{(1/2)}+1)+2*\arctan(2^{(1/2)}/(a/b)^{(1/4)}*x^{(1/2)}-1))$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.54

$$\int \frac{x^{3/2}}{(a+bx^2)^2} dx = \frac{(b^2x^2+ab)\left(-\frac{1}{a^3b^5}\right)^{\frac{1}{4}}\log\left(ab\left(-\frac{1}{a^3b^5}\right)^{\frac{1}{4}}+\sqrt{x}\right)-(-ib^2x^2-iab)\left(-\frac{1}{a^3b^5}\right)^{\frac{1}{4}}\log\left(iab\left(-\frac{1}{a^3b^5}\right)^{\frac{1}{4}}+\sqrt{x}\right)}{16ba}$$

input `integrate(x^(3/2)/(b*x^2+a)^2,x, algorithm="fricas")`

output
$$1/8*((b^2*x^2+a*b)*(-1/(a^3*b^5))^{(1/4)}*\log(a*b*(-1/(a^3*b^5))^{(1/4)}+sqrt(x))-(-I*b^2*x^2-I*a*b)*(-1/(a^3*b^5))^{(1/4)}*\log(I*a*b*(-1/(a^3*b^5))^{(1/4)}+sqrt(x))-(I*b^2*x^2+I*a*b)*(-1/(a^3*b^5))^{(1/4)}*\log(-I*a*b*(-1/(a^3*b^5))^{(1/4)}+sqrt(x))-(b^2*x^2+a*b)*(-1/(a^3*b^5))^{(1/4)}*\log(-a*b*(-1/(a^3*b^5))^{(1/4)}+sqrt(x))-4*sqrt(x))/(b^2*x^2+a*b)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 323 vs. $2(97) = 194$.

Time = 29.13 (sec) , antiderivative size = 323, normalized size of antiderivative = 2.60

$$\int \frac{x^{3/2}}{(a + bx^2)^2} dx = \begin{cases} \frac{\infty}{x^{3/2}}, \\ \frac{2x^{5/2}}{5a^2}, \\ -\frac{2}{3b^2x^{3/2}}, \\ -\frac{4a\sqrt{x}}{8a^2b+8ab^2x^2} - \frac{a\sqrt[4]{-\frac{a}{b}}\log\left(\sqrt{x}-\sqrt[4]{-\frac{a}{b}}\right)}{8a^2b+8ab^2x^2} + \frac{a\sqrt[4]{-\frac{a}{b}}\log\left(\sqrt{x}+\sqrt[4]{-\frac{a}{b}}\right)}{8a^2b+8ab^2x^2} + \frac{2a\sqrt[4]{-\frac{a}{b}}\operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{8a^2b+8ab^2x^2} \end{cases}$$

input `integrate(x**(3/2)/(b*x**2+a)**2,x)`

output `Piecewise((zoo/x**(3/2), Eq(a, 0) & Eq(b, 0)), (2*x**(5/2)/(5*a**2), Eq(b, 0)), (-2/(3*b**2*x**(3/2)), Eq(a, 0)), (-4*a*sqrt(x)/(8*a**2*b + 8*a*b**2*x**2) - a*(-a/b)**(1/4)*log(sqrt(x) - (-a/b)**(1/4))/(8*a**2*b + 8*a*b**2*x**2) + a*(-a/b)**(1/4)*log(sqrt(x) + (-a/b)**(1/4))/(8*a**2*b + 8*a*b**2*x**2) + 2*a*(-a/b)**(1/4)*atan(sqrt(x)/(-a/b)**(1/4))/(8*a**2*b + 8*a*b**2*x**2) - b*x**2*(-a/b)**(1/4)*log(sqrt(x) - (-a/b)**(1/4))/(8*a**2*b + 8*a*b**2*x**2) + b*x**2*(-a/b)**(1/4)*log(sqrt(x) + (-a/b)**(1/4))/(8*a**2*b + 8*a*b**2*x**2) + 2*b*x**2*(-a/b)**(1/4)*atan(sqrt(x)/(-a/b)**(1/4))/(8*a**2*b + 8*a*b**2*x**2), True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 195 vs. $2(97) = 194$.

Time = 0.12 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.57

$$\int \frac{x^{3/2}}{(a + bx^2)^2} dx = \frac{2\sqrt{2}\arctan\left(\frac{\sqrt{2}\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}+2\sqrt{b}\sqrt{x}\right)}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{2\sqrt{2}\arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}-2\sqrt{b}\sqrt{x}\right)}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{a}\sqrt{\sqrt{a}\sqrt{b}}} + \frac{\sqrt{2}\log\left(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x}+\sqrt{bx}+\sqrt{a}\right)}{a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{x}}{2(b^2x^2 + ab)}$$

input `integrate(x^(3/2)/(b*x^2+a)^2,x, algorithm="maxima")`

output
$$\frac{1}{16} \cdot (2 \cdot \sqrt{2}) \cdot \arctan\left(\frac{1}{2} \sqrt{2} \cdot (\sqrt{2} \cdot a^{1/4} \cdot b^{1/4} + 2 \cdot \sqrt{b} \cdot \sqrt{x}) / \sqrt{a} \cdot \sqrt{b}\right) / (\sqrt{a} \cdot \sqrt{b}) + 2 \cdot \sqrt{2} \cdot \arctan\left(-\frac{1}{2} \sqrt{2} \cdot (\sqrt{2} \cdot a^{1/4} \cdot b^{1/4} - 2 \cdot \sqrt{b} \cdot \sqrt{x}) / \sqrt{a} \cdot \sqrt{b}\right) / (\sqrt{a} \cdot \sqrt{b}) + \sqrt{2} \cdot \log(\sqrt{2} \cdot a^{1/4} \cdot b^{1/4} \cdot \sqrt{x} + \sqrt{b} \cdot x + \sqrt{a}) / (a^{3/4} \cdot b^{1/4}) - \sqrt{2} \cdot \log(-\sqrt{2} \cdot a^{1/4} \cdot b^{1/4} \cdot \sqrt{x} + \sqrt{b} \cdot x + \sqrt{a}) / (a^{3/4} \cdot b^{1/4}) / b - 1/2 \cdot \sqrt{x} / (b^2 \cdot x^2 + a \cdot b)$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 199 vs. $2(97) = 194$.

Time = 0.12 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.60

$$\int \frac{x^{3/2}}{(a + bx^2)^2} dx = \frac{\sqrt{2}(ab^3)^{1/4} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} + 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{8ab^2} + \frac{\sqrt{2}(ab^3)^{1/4} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} - 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{8ab^2} + \frac{\sqrt{2}(ab^3)^{1/4} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{16ab^2} - \frac{\sqrt{2}(ab^3)^{1/4} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{16ab^2} - \frac{\sqrt{x}}{2(bx^2 + a)b}$$

input `integrate(x^(3/2)/(b*x^2+a)^2,x, algorithm="giac")`

output
$$\frac{1}{8} \sqrt{2} \cdot (a \cdot b^3)^{1/4} \cdot \arctan\left(\frac{1}{2} \sqrt{2} \cdot (\sqrt{2} \cdot (a/b)^{1/4} + 2 \cdot \sqrt{b} \cdot \sqrt{x}) / (a/b)^{1/4}\right) / (a \cdot b^2) + \frac{1}{8} \sqrt{2} \cdot (a \cdot b^3)^{1/4} \cdot \arctan\left(-\frac{1}{2} \sqrt{2} \cdot (\sqrt{2} \cdot (a/b)^{1/4} - 2 \cdot \sqrt{b} \cdot \sqrt{x}) / (a/b)^{1/4}\right) / (a \cdot b^2) + \frac{1}{16} \sqrt{2} \cdot (a \cdot b^3)^{1/4} \cdot \log(\sqrt{2} \cdot \sqrt{x} \cdot (a/b)^{1/4} + x + \sqrt{a/b}) / (a \cdot b^2) - \frac{1}{16} \sqrt{2} \cdot (a \cdot b^3)^{1/4} \cdot \log(-\sqrt{2} \cdot \sqrt{x} \cdot (a/b)^{1/4} + x + \sqrt{a/b}) / (a \cdot b^2) - 1/2 \cdot \sqrt{x} / ((b \cdot x^2 + a) \cdot b)$$

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.52

$$\int \frac{x^{3/2}}{(a + bx^2)^2} dx = -\frac{\sqrt{x}}{2b(bx^2 + a)} - \frac{\operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{3/4}b^{5/4}} - \frac{\operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{3/4}b^{5/4}}$$

input `int(x^(3/2)/(a + b*x^2)^2,x)`output `- x^(1/2)/(2*b*(a + b*x^2)) - atan((b^(1/4)*x^(1/2))/(-a)^(1/4))/(4*(-a)^(3/4)*b^(5/4)) - atanh((b^(1/4)*x^(1/2))/(-a)^(1/4))/(4*(-a)^(3/4)*b^(5/4))`**Reduce [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.45

$$\int \frac{x^{3/2}}{(a + bx^2)^2} dx = \frac{-2b^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) - 2b^{\frac{7}{4}}a^{\frac{1}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right) x^2 + 2b^{\frac{3}{4}}a^{\frac{5}{4}}\sqrt{2} \operatorname{atan}\left(\frac{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{\frac{1}{4}}a^{\frac{1}{4}}\sqrt{2}}\right)}{(a + bx^2)^2}$$

input `int(x^(3/2)/(b*x^2+a)^2,x)`output `(- 2*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a - 2*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**2 + 2*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a + 2*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**2 - b**(3/4)*a**(1/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a - b**(3/4)*a**(1/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b*x**2 + b**(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a + b**(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b*x**2 - 8*sqrt(x)*a*b)/(16*a*b**2*(a + b*x**2))`

3.128 $\int \frac{x^{5/2}}{(a+bx^2)^2} dx$

Optimal result	850
Mathematica [A] (verified)	850
Rubi [A] (verified)	851
Maple [A] (verified)	856
Fricas [C] (verification not implemented)	856
Sympy [B] (verification not implemented)	857
Maxima [A] (verification not implemented)	857
Giac [B] (verification not implemented)	858
Mupad [B] (verification not implemented)	859
Reduce [B] (verification not implemented)	859

Optimal result

Integrand size = 15, antiderivative size = 126

$$\int \frac{x^{5/2}}{(a+bx^2)^2} dx = -\frac{x^{3/2}}{2b(a+bx^2)} + \frac{3 \left(\arctan \left(\frac{\sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}} - x} \right) - \log \left(\frac{\sqrt{\frac{a}{b}} + \sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x+x}}{\sqrt{a+bx^2}} \right) \right)}{4\sqrt{2} \sqrt[4]{\frac{a}{b}} b^2}$$

output

```

-1/2*x^(3/2)/b/(b*x^2+a)+3/8*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))-ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/(a/b)^(1/4)/b^2
    
```

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.02

$$\int \frac{x^{5/2}}{(a+bx^2)^2} dx = \frac{-\frac{4b^{3/4}x^{3/2}}{a+bx^2} - \frac{3\sqrt{2} \arctan \left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}\sqrt{x}} \right)}{\sqrt[4]{a}} - \frac{3\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}\sqrt{x}}{\sqrt{a}+\sqrt{bx}} \right)}{\sqrt[4]{a}}}{8b^{7/4}}$$

input `Integrate[x^(5/2)/(a + b*x^2)^2,x]`

output
$$\frac{((-4*b^{3/4}*x^{3/2})/(a + b*x^2) - (3*Sqrt[2]*ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^{1/4}*b^{1/4}*Sqrt[x])])/a^{1/4} - (3*Sqrt[2]*ArcTanh[(Sqrt[2]*a^{1/4}*b^{1/4}*Sqrt[x])/(Sqrt[a] + Sqrt[b]*x)])/a^{1/4})/(8*b^{7/4})$$

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.91, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {252, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^{5/2}}{(a + bx^2)^2} dx \\ & \quad \downarrow \text{252} \\ & \frac{3 \int \frac{\sqrt{x}}{bx^2+a} dx}{4b} - \frac{x^{3/2}}{2b(a + bx^2)} \\ & \quad \downarrow \text{266} \\ & \frac{3 \int \frac{x}{bx^2+a} d\sqrt{x}}{2b} - \frac{x^{3/2}}{2b(a + bx^2)} \\ & \quad \downarrow \text{826} \\ & \frac{3 \left(\frac{\int \frac{\sqrt{bx} + \sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} \right)}{2b} - \frac{x^{3/2}}{2b(a + bx^2)} \\ & \quad \downarrow \text{1476} \end{aligned}$$

$$3 \left(\frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{b}x}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right) - \frac{x^{3/2}}{2b(a + bx^2)}$$

1082

$$3 \left(\frac{\int \frac{1}{-x-1} d\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{b}x}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right) - \frac{x^{3/2}}{2b(a + bx^2)}$$

217

$$3 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a} - \sqrt{b}x}{bx^2 + a} d\sqrt{x}}{2\sqrt{b}} \right) - \frac{x^{3/2}}{2b(a + bx^2)}$$

1479

$$3 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}} + 1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a} - 2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

$$\frac{x^{3/2}}{2b(a + bx^2)}$$

25

$$3 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

$$\frac{2b}{x^{3/2} (2b(a+bx^2))}$$

27

$$3 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}}{x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2\sqrt[4]{a}\sqrt{b}} \right)$$

$$\frac{2b}{x^{3/2} (2b(a+bx^2))}$$

1103

$$3 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} - \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)$$

$$\frac{2b}{x^{3/2} (2b(a+bx^2))}$$

input `Int [x^(5/2)/(a + b*x^2)^2,x]`

output

$$\begin{aligned}
& -1/2*x^{(3/2)}/(b*(a + b*x^2)) + (3*((-ArcTan[1 - (Sqrt[2]*b^{(1/4)}*Sqrt[x]) \\
& /a^{(1/4)}]/(Sqrt[2]*a^{(1/4)*b^{(1/4)}})) + ArcTan[1 + (Sqrt[2]*b^{(1/4)}*Sqrt[x] \\
&)/a^{(1/4)}]/(Sqrt[2]*a^{(1/4)*b^{(1/4)}}))/(2*Sqrt[b]) - (-1/2*Log[Sqrt[a] - Sqrt[2]*a^{(1/4)*b^{(1/4)}*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^{(1/4)*b^{(1/4)}}) + Log[Sqrt[a] + Sqrt[2]*a^{(1/4)*b^{(1/4)}*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^{(1/4)*b^{(1/4)}}))/(2*Sqrt[b]))/(2*b)
\end{aligned}$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(F_x), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[F_x, x], x]$$

rule 27

$$\text{Int}[(a_)*(F_x), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[F_x, x], x] \text{ ; FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[F_x, (b_)*(G_x) \text{ ; FreeQ}[b, x]$$

rule 217

$$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \text{ ; FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$

rule 252

$$\text{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a + b*x^2)^{(p+1)}/(2*b*(p+1))), x] - \text{Simp}[c^2*((m-1)/(2*b*(p+1))) \quad \text{Int}[(c*x)^{(m-2)}*(a + b*x^2)^{(p+1)}, x], x] \text{ ; FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ !\text{LtQ}[(m + 2*p + 3)/2, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 266

$$\text{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \quad \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(2*k)}/c^2))^p, x], x, (c*x)^{(1/k)}], x]] \text{ ; FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{FractionQ}[m] \ \&\& \ \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 826 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*s) Int[(r + s*x^2)/(a + b*x^4), x], x] - Simp[1/(2*s) Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.98

method	result	size
derivativedivides	$-\frac{x^{\frac{3}{2}}}{2b(x^2b+a)} + \frac{3\sqrt{2} \left(\ln \left(\frac{x-\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+\sqrt{\frac{a}{b}}} \right)}{\sqrt{\frac{a}{b}}+\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+x}} \right) + 2 \arctan \left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1} \right) + 2 \arctan \left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1} \right)}{16b^2 \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	124
default	$-\frac{x^{\frac{3}{2}}}{2b(x^2b+a)} + \frac{3\sqrt{2} \left(\ln \left(\frac{x-\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+\sqrt{\frac{a}{b}}} \right)}{\sqrt{\frac{a}{b}}+\sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x+x}} \right) + 2 \arctan \left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1} \right) + 2 \arctan \left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1} \right)}{16b^2 \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	124

input `int(x^(5/2)/(b*x^2+a)^2,x,method=_RETURNVERBOSE)`

output
$$-1/2*x^{(3/2)}/b/(b*x^2+a)+3/16/b^2/(a/b)^{(1/4)}*2^{(1/2)}*(\ln((x-2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)}+(a/b)^{(1/2))}/((a/b)^{(1/2)}+2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)+x}))+2*\arctan(2^{(1/2)}/(a/b)^{(1/4)}*x^{(1/2)+1}))+2*\arctan(2^{(1/2)}/(a/b)^{(1/4)}*x^{(1/2)}-1))$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.59

$$\int \frac{x^{5/2}}{(a+bx^2)^2} dx = \frac{3(b^2x^2+ab)\left(-\frac{1}{ab^7}\right)^{\frac{1}{4}} \log\left(ab^5\left(-\frac{1}{ab^7}\right)^{\frac{3}{4}}+\sqrt{x}\right) - 3(ib^2x^2+iab)\left(-\frac{1}{ab^7}\right)^{\frac{1}{4}} \log\left(iab^5\left(-\frac{1}{ab^7}\right)^{\frac{3}{4}}+\sqrt{x}\right)}{16b^2\left(\frac{a}{b}\right)^{\frac{1}{4}}}$$

input `integrate(x^(5/2)/(b*x^2+a)^2,x, algorithm="fricas")`

output
$$1/8*(3*(b^2*x^2+a*b)*(-1/(a*b^7))^{(1/4)}*\log(a*b^5*(-1/(a*b^7))^{(3/4)}+sqrt(x))-3*(I*b^2*x^2+I*a*b)*(-1/(a*b^7))^{(1/4)}*\log(I*a*b^5*(-1/(a*b^7))^{(3/4)}+sqrt(x))-3*(-I*b^2*x^2-I*a*b)*(-1/(a*b^7))^{(1/4)}*\log(-I*a*b^5*(-1/(a*b^7))^{(3/4)}+sqrt(x))-3*(b^2*x^2+a*b)*(-1/(a*b^7))^{(1/4)}*\log(-a*b^5*(-1/(a*b^7))^{(3/4)}+sqrt(x))-4*x^{(3/2)})/(b^2*x^2+a*b)$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 393 vs. $2(100) = 200$.

Time = 53.45 (sec) , antiderivative size = 393, normalized size of antiderivative = 3.12

$$\int \frac{x^{5/2}}{(a + bx^2)^2} dx = \begin{cases} \frac{\infty}{\sqrt{x}} \\ \frac{2x^{7/2}}{7a^2} \\ -\frac{2}{b^2\sqrt{x}} \\ \frac{3a \log\left(\sqrt{x} - \sqrt[4]{-\frac{a}{b}}\right)}{8ab^2 \sqrt[4]{-\frac{a}{b} + 8b^3x^2} \sqrt[4]{-\frac{a}{b}}} - \frac{3a \log\left(\sqrt{x} + \sqrt[4]{-\frac{a}{b}}\right)}{8ab^2 \sqrt[4]{-\frac{a}{b} + 8b^3x^2} \sqrt[4]{-\frac{a}{b}}} + \frac{6a \operatorname{atan}\left(\frac{\sqrt{x}}{\sqrt[4]{-\frac{a}{b}}}\right)}{8ab^2 \sqrt[4]{-\frac{a}{b} + 8b^3x^2} \sqrt[4]{-\frac{a}{b}}} - \frac{4bx^{3/2} \sqrt[4]{-\frac{a}{b}}}{8ab^2 \sqrt[4]{-\frac{a}{b} + 8b^3x^2}} \end{cases}$$

input

```
integrate(x**(5/2)/(b*x**2+a)**2,x)
```

output

```
Piecewise((zoo/sqrt(x), Eq(a, 0) & Eq(b, 0)), (2*x**(7/2)/(7*a**2), Eq(b, 0)), (-2/(b**2*sqrt(x)), Eq(a, 0)), (3*a*log(sqrt(x) - (-a/b)**(1/4))/(8*a*b**2*(-a/b)**(1/4) + 8*b**3*x**2*(-a/b)**(1/4)) - 3*a*log(sqrt(x) + (-a/b)**(1/4))/(8*a*b**2*(-a/b)**(1/4) + 8*b**3*x**2*(-a/b)**(1/4)) + 6*a*atan(sqrt(x)/(-a/b)**(1/4))/(8*a*b**2*(-a/b)**(1/4) + 8*b**3*x**2*(-a/b)**(1/4)) - 4*b*x**(3/2)*(-a/b)**(1/4)/(8*a*b**2*(-a/b)**(1/4) + 8*b**3*x**2*(-a/b)**(1/4)) + 3*b*x**2*log(sqrt(x) - (-a/b)**(1/4))/(8*a*b**2*(-a/b)**(1/4) + 8*b**3*x**2*(-a/b)**(1/4)) - 3*b*x**2*log(sqrt(x) + (-a/b)**(1/4))/(8*a*b**2*(-a/b)**(1/4) + 8*b**3*x**2*(-a/b)**(1/4)) + 6*b*x**2*atan(sqrt(x)/(-a/b)**(1/4))/(8*a*b**2*(-a/b)**(1/4) + 8*b**3*x**2*(-a/b)**(1/4)), True))
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.55

$$\int \frac{x^{5/2}}{(a + bx^2)^2} dx = -\frac{x^{3/2}}{2(b^2x^2 + ab)} + 3 \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}a^{1/4}b^{1/4} + 2\sqrt{b}\sqrt{x}\right)}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{b}\sqrt{b}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}a^{1/4}b^{1/4} - 2\sqrt{b}\sqrt{x}\right)}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{b}\sqrt{b}} - \frac{\sqrt{2} \log\left(\sqrt{2}a^{1/4}b^{1/4}\sqrt{x} + \sqrt{bx} + \sqrt{a}\right)}{a^{1/4}b^{3/4}} + \frac{\sqrt{2} \log\left(-\sqrt{2}a^{1/4}b^{1/4}\sqrt{x} + \sqrt{bx} + \sqrt{a}\right)}{a^{1/4}b^{3/4}} \right)$$

input `integrate(x^(5/2)/(b*x^2+a)^2,x, algorithm="maxima")`

output
$$\begin{aligned} & -1/2*x^{(3/2)}/(b^2*x^2 + a*b) + 3/16*(2*\sqrt{2}*\arctan(1/2*\sqrt{2}*(\sqrt{2} \\ & *a^{(1/4)}*b^{(1/4)} + 2*\sqrt{b}*\sqrt{x}))/\sqrt{(\sqrt{a}*\sqrt{b}))/(\sqrt{(\sqrt{a} \\ & *\sqrt{b}))*\sqrt{b}} + 2*\sqrt{2}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*a^{(1/4)}*b^{(1/4)} \\ &) - 2*\sqrt{b}*\sqrt{x}))/\sqrt{(\sqrt{a}*\sqrt{b}))/(\sqrt{(\sqrt{a}*\sqrt{b}))*\sqrt{b}} \\ & - \sqrt{2}*\log(\sqrt{2}*a^{(1/4)}*b^{(1/4)}*\sqrt{x} + \sqrt{b}*x + \sqrt{a}))/ \\ & (a^{(1/4)}*b^{(3/4)}) + \sqrt{2}*\log(-\sqrt{2}*a^{(1/4)}*b^{(1/4)}*\sqrt{x} + \sqrt{b}* \\ & x + \sqrt{a}))/(a^{(1/4)}*b^{(3/4)})/b \end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 199 vs. $2(99) = 198$.

Time = 0.12 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.58

$$\begin{aligned} \int \frac{x^{5/2}}{(a + bx^2)^2} dx &= -\frac{x^{3/2}}{2(bx^2 + a)b} + \frac{3\sqrt{2}(ab^3)^{3/4} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} + 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{8ab^4} \\ &+ \frac{3\sqrt{2}(ab^3)^{3/4} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} - 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{8ab^4} \\ &- \frac{3\sqrt{2}(ab^3)^{3/4} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{16ab^4} \\ &+ \frac{3\sqrt{2}(ab^3)^{3/4} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{16ab^4} \end{aligned}$$

input `integrate(x^(5/2)/(b*x^2+a)^2,x, algorithm="giac")`

output
$$\begin{aligned} & -1/2*x^{(3/2)}/((b*x^2 + a)*b) + 3/8*\sqrt{2}*(a*b^3)^{(3/4)}*\arctan(1/2*\sqrt{2} \\ &)*(\sqrt{2}*(a/b)^{(1/4)} + 2*\sqrt{x}))/((a/b)^{(1/4)})/(a*b^4) + 3/8*\sqrt{2}*(a* \\ & b^3)^{(3/4)}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*(a/b)^{(1/4)} - 2*\sqrt{x}))/((a/b)^{(1/4)} \\ &)/(a*b^4) - 3/16*\sqrt{2}*(a*b^3)^{(3/4)}*\log(\sqrt{2}*\sqrt{x}*(a/b)^{(1/4)} + \\ & x + \sqrt{a/b}))/((a*b^4) + 3/16*\sqrt{2}*(a*b^3)^{(3/4)}*\log(-\sqrt{2}*\sqrt{x}* \\ & (a/b)^{(1/4)} + x + \sqrt{a/b}))/((a*b^4) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.51

$$\int \frac{x^{5/2}}{(a + bx^2)^2} dx = \frac{3 \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{1/4}b^{7/4}} - \frac{x^{3/2}}{2b(bx^2 + a)} - \frac{3 \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{4(-a)^{1/4}b^{7/4}}$$

input `int(x^(5/2)/(a + b*x^2)^2,x)`output `(3*atan((b^(1/4)*x^(1/2))/(-a)^(1/4)))/(4*(-a)^(1/4)*b^(7/4)) - x^(3/2)/(2*b*(a + b*x^2)) - (3*atanh((b^(1/4)*x^(1/2))/(-a)^(1/4)))/(4*(-a)^(1/4)*b^(7/4))`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 307, normalized size of antiderivative = 2.44

$$\int \frac{x^{5/2}}{(a + bx^2)^2} dx = \frac{-6b^{1/4}a^{7/4}\sqrt{2} \operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{1/4}a^{1/4}\sqrt{2}}\right) - 6b^{5/4}a^{3/4}\sqrt{2} \operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{1/4}a^{1/4}\sqrt{2}}\right) x^2 + 6b^{1/4}a^{7/4}\sqrt{2} \operatorname{atan}\left(\frac{b^{1/4}a^{1/4}\sqrt{2}-2\sqrt{x}\sqrt{b}}{b^{1/4}a^{1/4}\sqrt{2}}\right)}{(a + bx^2)^2}$$

input `int(x^(5/2)/(b*x^2+a)^2,x)`output `(- 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a - 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**2 + 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a + 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b*x**2 + 3*b**(1/4)*a**(3/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a + 3*b**(1/4)*a**(3/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b*x**2 - 3*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a - 3*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b*x**2 - 8*sqrt(x)*a*b*x)/(16*a*b**2*(a + b*x**2))`

3.129 $\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx$

Optimal result	860
Mathematica [A] (verified)	861
Rubi [A] (verified)	861
Maple [A] (verified)	867
Fricas [C] (verification not implemented)	868
Sympy [F(-1)]	868
Maxima [A] (verification not implemented)	869
Giac [A] (verification not implemented)	870
Mupad [B] (verification not implemented)	870
Reduce [B] (verification not implemented)	871

Optimal result

Integrand size = 15, antiderivative size = 145

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx = \sqrt{x} \left(\frac{1}{4a(a+bx^2)^2} + \frac{7}{16a^2(a+bx^2)} \right) + \frac{21 \left(\arctan \left(\frac{\sqrt{2} \sqrt{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}-x}} \right) + \log \left(\frac{\sqrt{\frac{a}{b}+\sqrt{2} \sqrt{\frac{a}{b}} \sqrt{x+x}}}{\sqrt{a+bx^2}} \right) \right)}{32\sqrt{2}a^2 \left(\frac{a}{b}\right)^{3/4} b}$$

output

```
x^(1/2)*(1/4/a/(b*x^2+a)^2+7/16/a^2/(b*x^2+a))+21/64*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))+ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/a^2/(a/b)^(3/4)/b
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.95

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx$$

$$= \frac{\frac{4a^{3/4}\sqrt{x}(11a+7bx^2)}{(a+bx^2)^2} - \frac{21\sqrt{2}\arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right)}{\sqrt[4]{b}} + \frac{21\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}+\sqrt{bx}}\right)}{\sqrt[4]{b}}}{64a^{11/4}}$$

input

```
Integrate[1/(Sqrt[x]*(a + b*x^2)^3), x]
```

output

```
((4*a^(3/4)*Sqrt[x]*(11*a + 7*b*x^2))/(a + b*x^2)^2 - (21*Sqrt[2]*ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x]])/b^(1/4) + (21*Sqrt[2]*ArcTanh[(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x]]/(Sqrt[a] + Sqrt[b]*x)]/b^(1/4))/(64*a^(11/4))
```

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.86, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.733$, Rules used = {253, 253, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx$$

$$\downarrow 253$$

$$\frac{7 \int \frac{1}{\sqrt{x}(bx^2+a)^2} dx}{8a} + \frac{\sqrt{x}}{4a(a+bx^2)^2}$$

$$\downarrow 253$$

$$7 \left(\frac{3 \int \frac{1}{\sqrt{x}(bx^2+a)} dx}{8a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right) + \frac{\sqrt{x}}{4a(a+bx^2)^2}$$

266

$$7 \left(\frac{3 \int \frac{1}{bx^2+a} d\sqrt{x}}{8a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right) + \frac{\sqrt{x}}{4a(a+bx^2)^2}$$

755

$$7 \left(\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} \right)}{8a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right) + \frac{\sqrt{x}}{4a(a+bx^2)^2}$$

1476

$$7 \left(\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{x - \sqrt{2} \sqrt[4]{a}\sqrt{x} + \sqrt{a}} \frac{d\sqrt{x}}{\sqrt{b}} + \frac{\int \frac{1}{x + \sqrt{2} \sqrt[4]{a}\sqrt{x} + \sqrt{a}} \frac{d\sqrt{x}}{\sqrt{b}}}{2\sqrt{a}} \right)}{8a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right) + \frac{\sqrt{x}}{4a(a+bx^2)^2}$$

1082

$$\left(\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\int \frac{1}{-x-1} d \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1 \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right)}{8a} + \frac{\sqrt{x}}{4a(a+bx^2)^2}$$

217

$$\left(\frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} + 1 \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right)}{8a} + \frac{\sqrt{x}}{4a(a+bx^2)^2}$$

1479

$$\left(\frac{3}{7} \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt[4]{b}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a} \right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt[4]{b}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\sqrt{x}}{2a(a+b)} \right)$$

8a

$$\frac{\sqrt{x}}{4a(a+bx^2)^2}$$

↓ 25

$$\left(\frac{3}{7} \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt[4]{b}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a} \right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt[4]{b}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)} \right)$$

8a

$$\frac{\sqrt{x}}{4a(a+bx^2)^2}$$

↓ 27

$$\left(\frac{3 \left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a-2} \sqrt[4]{b} \sqrt{x}}{x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt{b}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}}{x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2 \sqrt[4]{a} \sqrt{b}} + \frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right)$$

$$\frac{\sqrt{x} \cdot 8a}{4a(a+bx^2)^2}$$

↓ 1103

$$\left(\frac{3 \left(\frac{\arctan\left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\log\left(\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx}\right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\log\left(-\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx}\right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} \right)$$

$$\frac{\sqrt{x} \cdot 8a}{4a(a+bx^2)^2}$$

input

```
Int [1/(Sqrt [x]*(a + b*x^2)^3), x]
```

output

```
Sqrt[x]/(4*a*(a + b*x^2)^2) + (7*(Sqrt[x]/(2*a*(a + b*x^2)) + (3*((-ArcTan[1 - (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[a]) + (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[a]))/(2*a))/(8*a)
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]
```

rule 217

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

rule 253

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^2)^(p + 1)/(2*a*c*(p + 1))), x] + Simp[(m + 2*p + 3)/(2*a*(p + 1)) Int[(c*x)^m*(a + b*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && LtQ[p, -1] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 266

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]
```

rule 755

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

rule 1082 `Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])]; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.01

method	result
derivativedivides	$\frac{\sqrt{x}}{4a(x^2b+a)^2} + \frac{7\sqrt{x}}{16a(x^2b+a)} + \frac{21\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x}+x}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x}+\sqrt{\frac{a}{b}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1}\right)\right)}{128a^2}$
default	$\frac{\sqrt{x}}{4a(x^2b+a)^2} + \frac{7\sqrt{x}}{16a(x^2b+a)} + \frac{21\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2}\left(\ln\left(\frac{\sqrt{\frac{a}{b}}+\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x}+x}{x-\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x}+\sqrt{\frac{a}{b}}}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}+1}\right)+2\arctan\left(\frac{\sqrt{2}\sqrt{x}}{\left(\frac{a}{b}\right)^{\frac{1}{4}}-1}\right)\right)}{128a^2}$

input `int(1/x^(1/2)/(b*x^2+a)^3,x,method=_RETURNVERBOSE)`

output

```
1/4*x^(1/2)/a/(b*x^2+a)^2+7/4/a*(1/4*x^(1/2)/a/(b*x^2+a)+3/32/a^2*(a/b)^(1/4)*2^(1/2)*(ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(x-2^(1/2)*(a/b)^(1/4)*x^(1/2)+(a/b)^(1/2)))+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)+1)+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)-1)))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 268, normalized size of antiderivative = 1.85

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx$$

$$= \frac{21(a^2b^2x^4 + 2a^3bx^2 + a^4)\left(-\frac{1}{a^{11}b}\right)^{\frac{1}{4}} \log\left(a^3\left(-\frac{1}{a^{11}b}\right)^{\frac{1}{4}} + \sqrt{x}\right) - 21(-ia^2b^2x^4 - 2ia^3bx^2 - ia^4)\left(-\frac{1}{a^{11}b}\right)^{\frac{1}{4}}}{\dots}$$

input

```
integrate(1/x^(1/2)/(b*x^2+a)^3,x, algorithm="fricas")
```

output

```
1/64*(21*(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)*(-1/(a^11*b))^(1/4)*log(a^3*(-1/(a^11*b))^(1/4) + sqrt(x)) - 21*(-I*a^2*b^2*x^4 - 2*I*a^3*b*x^2 - I*a^4)*(-1/(a^11*b))^(1/4)*log(I*a^3*(-1/(a^11*b))^(1/4) + sqrt(x)) - 21*(I*a^2*b^2*x^4 + 2*I*a^3*b*x^2 + I*a^4)*(-1/(a^11*b))^(1/4)*log(-I*a^3*(-1/(a^11*b))^(1/4) + sqrt(x)) - 21*(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)*(-1/(a^11*b))^(1/4)*log(-a^3*(-1/(a^11*b))^(1/4) + sqrt(x)) + 4*(7*b*x^2 + 11*a)*sqrt(x))/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx = \text{Timed out}$$

input

```
integrate(1/x**(1/2)/(b*x**2+a)**3,x)
```

output

```
Timed out
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx = \frac{7bx^{\frac{5}{2}} + 11a\sqrt{x}}{16(a^2b^2x^4 + 2a^3bx^2 + a^4)} + 21 \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} + 2\sqrt{b}\sqrt{x})}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{a}\sqrt{b}} + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} - 2\sqrt{b}\sqrt{x})}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{a}\sqrt{b}} + \frac{\sqrt{2} \log(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} \right) + \frac{\sqrt{2} \log(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} + \frac{\sqrt{2} \log(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}}$$

input `integrate(1/x^(1/2)/(b*x^2+a)^3,x, algorithm="maxima")`output `1/16*(7*b*x^(5/2) + 11*a*sqrt(x))/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4) + 21/128*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + sqrt(2)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4)) - sqrt(2)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4))/a^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.44

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx = \frac{21\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}+2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{64a^3b}$$

$$+ \frac{21\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}-2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{64a^3b}$$

$$+ \frac{21\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{128a^3b}$$

$$- \frac{21\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{128a^3b} + \frac{7bx^{\frac{5}{2}}+11a\sqrt{x}}{16(bx^2+a)^2a^2}$$

input `integrate(1/x^(1/2)/(b*x^2+a)^3,x, algorithm="giac")`output `21/64*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/(a^3*b) + 21/64*sqrt(2)*(a*b^3)^(1/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/(a^3*b) + 21/128*sqrt(2)*(a*b^3)^(1/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^3*b) - 21/128*sqrt(2)*(a*b^3)^(1/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^3*b) + 1/16*(7*b*x^(5/2) + 11*a*sqrt(x))/((b*x^2 + a)^2*a^2)`**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.59

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx = \frac{\frac{11\sqrt{x}}{16a} + \frac{7bx^{5/2}}{16a^2}}{a^2 + 2abx^2 + b^2x^4} - \frac{21 \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{11/4}b^{1/4}} - \frac{21 \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{11/4}b^{1/4}}$$

input `int(1/(x^(1/2)*(a + b*x^2)^3),x)`

output

```
((11*x^(1/2))/(16*a) + (7*b*x^(5/2))/(16*a^2))/(a^2 + b^2*x^4 + 2*a*b*x^2)
- (21*atan((b^(1/4)*x^(1/2))/(-a)^(1/4)))/(32*(-a)^(11/4)*b^(1/4)) - (21*
atanh((b^(1/4)*x^(1/2))/(-a)^(1/4)))/(32*(-a)^(11/4)*b^(1/4))
```

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 477, normalized size of antiderivative = 3.29

$$\int \frac{1}{\sqrt{x}(a+bx^2)^3} dx = \text{Too large to display}$$

input

```
int(1/x^(1/2)/(b*x^2+a)^3,x)
```

output

```
( - 42*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a**2 - 84*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a*b*x**2 - 42*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b**2*x**4 + 42*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a**2 + 84*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a*b*x**2 + 42*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b**2*x**4 - 21*b**(3/4)*a**(1/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a**2 - 42*b**(3/4)*a**(1/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 - 21*b**(3/4)*a**(1/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 + 21*b**(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a**2 + 42*b**(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 + 21*b**(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 + 88*sqrt(x)*a**2*b + 56*sqrt(x)*a*b**2*x**2)/(128*a**3*b*(a**2 + 2*a*b*x**2 + b**2*x**4))
```


3.130 $\int \frac{\sqrt{x}}{(a+bx^2)^3} dx$

Optimal result	872
Mathematica [A] (verified)	873
Rubi [A] (verified)	873
Maple [A] (verified)	878
Fricas [C] (verification not implemented)	879
Sympy [B] (verification not implemented)	879
Maxima [A] (verification not implemented)	880
Giac [A] (verification not implemented)	881
Mupad [B] (verification not implemented)	882
Reduce [B] (verification not implemented)	882

Optimal result

Integrand size = 15, antiderivative size = 147

$$\int \frac{\sqrt{x}}{(a+bx^2)^3} dx = x^{3/2} \left(\frac{1}{4a(a+bx^2)^2} + \frac{5}{16a^2(a+bx^2)} \right) + \frac{5 \left(\arctan \left(\frac{\sqrt{2} \sqrt{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}} - x} \right) - \log \left(\frac{\sqrt{\frac{a}{b}} + \sqrt{2} \sqrt{\frac{a}{b}} \sqrt{x} + x}{\sqrt{a+bx^2}} \right) \right)}{32\sqrt{2}a^2 \sqrt[4]{\frac{a}{b}} b}$$

output

```
x^(3/2)*(1/4/a/(b*x^2+a)^2+5/16/a^2/(b*x^2+a))+5/64*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))-ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/a^2/(a/b)^(1/4)/b
```

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.94

$$\int \frac{\sqrt{x}}{(a+bx^2)^3} dx$$

$$= \frac{4\sqrt[4]{ax^{3/2}}(9a+5bx^2)}{(a+bx^2)^2} - \frac{5\sqrt{2} \arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right)}{b^{3/4}} - \frac{5\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}+\sqrt{bx}}\right)}{b^{3/4}}$$

$$64a^{9/4}$$

input `Integrate[Sqrt[x]/(a + b*x^2)^3,x]`

output `((4*a^(1/4)*x^(3/2)*(9*a + 5*b*x^2))/(a + b*x^2)^2 - (5*Sqrt[2]*ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x]])/b^(3/4) - (5*Sqrt[2]*ArcTanh[(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x]/(Sqrt[a] + Sqrt[b]*x)]/b^(3/4)))/(64*a^(9/4))`

Rubi [A] (verified)

Time = 0.45 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.84, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.733$, Rules used = {253, 253, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{x}}{(a+bx^2)^3} dx$$

$$\downarrow 253$$

$$\frac{5 \int \frac{\sqrt{x}}{(bx^2+a)^2} dx}{8a} + \frac{x^{3/2}}{4a(a+bx^2)^2}$$

$$\downarrow 253$$

$$\begin{aligned}
 & \frac{5 \left(\frac{\int \frac{\sqrt{x}}{bx^2+a} dx}{4a} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8a} + \frac{x^{3/2}}{4a(a+bx^2)^2} \\
 & \quad \downarrow \text{266} \\
 & \frac{5 \left(\frac{\int \frac{x}{bx^2+a} d\sqrt{x}}{2a} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8a} + \frac{x^{3/2}}{4a(a+bx^2)^2} \\
 & \quad \downarrow \text{826} \\
 & \frac{5 \left(\frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8a} + \frac{x^{3/2}}{4a(a+bx^2)^2} \\
 & \quad \downarrow \text{1476} \\
 & \frac{5 \left(\frac{\int \frac{1}{x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{1}{x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x} + \sqrt{a}}{\sqrt{b}}} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8a} + \frac{x^{3/2}}{4a(a+bx^2)^2} \\
 & \quad \downarrow \text{1082} \\
 & \frac{5 \left(\frac{\int \frac{1}{-x-1} d \left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}} \right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{1}{-x-1} d \left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}} + 1 \right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8a} + \frac{x^{3/2}}{4a(a+bx^2)^2} \\
 & \quad \downarrow \text{217}
 \end{aligned}$$

$$5 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right) + \frac{x^{3/2}}{4a(a+bx^2)^2}$$

8a

1479

$$5 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)^2} \right)$$

8a

$$\frac{x^{3/2}}{4a(a+bx^2)^2}$$

25

$$5 \left(\frac{\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x - \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x} + \sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x + \frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt[4]{b}} + \frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)^2} \right) +$$

8a

$$\frac{x^{3/2}}{4a(a+bx^2)^2}$$

27

$$5 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{x-\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt{a}}{x+\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}} d\sqrt{x}}{2\sqrt[4]{a}\sqrt[4]{b}} \right) + \frac{x^{3/2}}{2a(a+bx^2)}$$

$$\frac{8a}{x^{3/2}} \frac{1}{4a(a+bx^2)^2}$$

1103

$$5 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}\right)}{2a} - \frac{\log\left(\frac{-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}\right)}{2\sqrt{b}} \right) + \frac{x^{3/2}}{2a(a+bx^2)}$$

$$\frac{8a}{x^{3/2}} \frac{1}{4a(a+bx^2)^2}$$

input `Int[Sqrt[x]/(a + b*x^2)^3,x]`

output `x^(3/2)/(4*a*(a + b*x^2)^2) + (5*(x^(3/2)/(2*a*(a + b*x^2))) + ((- (ArcTan[1 - (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]) - (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]))/(8*a)`

Definitions of rubi rules used

- rule 25 $\text{Int}[-(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, \text{x}], \text{x}]$
- rule 27 $\text{Int}[(\text{a}_)*(\text{Fx}_), \text{x_Symbol}] \rightarrow \text{Simp}[\text{a} \quad \text{Int}[\text{Fx}, \text{x}], \text{x}] \text{ ; FreeQ}[\text{a}, \text{x}] \ \&\& \ \text{!MatchQ}[\text{Fx}, (\text{b}_)*(\text{Gx}_)] \text{ ; FreeQ}[\text{b}, \text{x}]$
- rule 217 $\text{Int}[(\text{a}_) + (\text{b}_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{Rt}[-\text{a}, 2]*\text{Rt}[-\text{b}, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-\text{b}, 2]*(\text{x}/\text{Rt}[-\text{a}, 2])], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ \text{PosQ}[\text{a}/\text{b}] \ \&\& \ (\text{LtQ}[\text{a}, 0] \ || \ \text{LtQ}[\text{b}, 0])$
- rule 253 $\text{Int}[(\text{c}_)*(x_)^{\text{m}_}*(\text{a}_) + (\text{b}_)*(x_)^2)^{\text{p}_}, \text{x_Symbol}] \rightarrow \text{Simp}[(-\text{c}*x)^{\text{m} + 1}*(\text{a} + \text{b}*x^2)^{\text{p} + 1}/(2*\text{a}*c*(\text{p} + 1)), \text{x}] + \text{Simp}[(\text{m} + 2*\text{p} + 3)/(2*\text{a}*(\text{p} + 1)) \quad \text{Int}[(\text{c}*x)^{\text{m}}*(\text{a} + \text{b}*x^2)^{\text{p} + 1}, \text{x}], \text{x}] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{m}\}, \text{x}] \ \&\& \ \text{LtQ}[\text{p}, -1] \ \&\& \ \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$
- rule 266 $\text{Int}[(\text{c}_)*(x_)^{\text{m}_}*(\text{a}_) + (\text{b}_)*(x_)^2)^{\text{p}_}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{k} = \text{Denominator}[\text{m}]\}, \text{Simp}[\text{k}/\text{c} \quad \text{Subst}[\text{Int}[x^{(\text{k}*(\text{m} + 1) - 1)*(a + b*(x^{2*\text{k}}/c^2)})^{\text{p}}, \text{x}], \text{x}, (\text{c}*x)^{1/\text{k}}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}, \text{p}\}, \text{x}] \ \&\& \ \text{FractionQ}[\text{m}] \ \&\& \ \text{IntBinomialQ}[\text{a}, \text{b}, \text{c}, 2, \text{m}, \text{p}, \text{x}]$
- rule 826 $\text{Int}[(x_)^2/((\text{a}_) + (\text{b}_)*(x_)^4), \text{x_Symbol}] \rightarrow \text{With}[\{\text{r} = \text{Numerator}[\text{Rt}[\text{a}/\text{b}, 2]], \text{s} = \text{Denominator}[\text{Rt}[\text{a}/\text{b}, 2]]\}, \text{Simp}[1/(2*\text{s}) \quad \text{Int}[(\text{r} + \text{s}*x^2)/(\text{a} + \text{b}*x^4), \text{x}], \text{x}] - \text{Simp}[1/(2*\text{s}) \quad \text{Int}[(\text{r} - \text{s}*x^2)/(\text{a} + \text{b}*x^4), \text{x}], \text{x}]] \text{ ; FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \ \&\& \ (\text{GtQ}[\text{a}/\text{b}, 0] \ || \ (\text{PosQ}[\text{a}/\text{b}] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{a}]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, \text{b}]])$
- rule 1082 $\text{Int}[(\text{a}_) + (\text{b}_)*(x_) + (\text{c}_)*(x_)^2)^{-1}, \text{x_Symbol}] \rightarrow \text{With}[\{\text{q} = 1 - 4*\text{Simplify}[\text{a}*(\text{c}/\text{b}^2)]\}, \text{Simp}[-2/\text{b} \quad \text{Subst}[\text{Int}[1/(\text{q} - x^2), \text{x}], \text{x}, 1 + 2*\text{c}*(\text{x}/\text{b})], \text{x}] \text{ ; RationalQ}[\text{q}] \ \&\& \ (\text{EqQ}[\text{q}^2, 1] \ || \ \text{!RationalQ}[\text{b}^2 - 4*\text{a}*c]) \text{ ; FreeQ}[\{\text{a}, \text{b}, \text{c}\}, \text{x}]$

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.02

method	result	size
derivativedivides	$\frac{x^{\frac{3}{2}}}{4a(x^2b+a)^2} + \frac{\frac{5x^{\frac{3}{2}}}{16a(x^2b+a)} + \frac{5\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + \sqrt{\frac{a}{b}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + x} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} + 1}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} - 1}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} \right) \right)}{128ab \left(\frac{a}{b} \right)^{\frac{1}{4}}}{a}$	150
default	$\frac{x^{\frac{3}{2}}}{4a(x^2b+a)^2} + \frac{\frac{5x^{\frac{3}{2}}}{16a(x^2b+a)} + \frac{5\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + \sqrt{\frac{a}{b}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b} \right)^{\frac{1}{4}} \sqrt{x} + x} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} + 1}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} - 1}{\left(\frac{a}{b} \right)^{\frac{1}{4}}} \right) \right)}{128ab \left(\frac{a}{b} \right)^{\frac{1}{4}}}{a}$	150

input `int(x^(1/2)/(b*x^2+a)^3,x,method=_RETURNVERBOSE)`

output `1/4*x^(3/2)/a/(b*x^2+a)^2+5/4/a*(1/4*x^(3/2)/a/(b*x^2+a)+1/32/a/b/(a/b)^(1/4)*2^(1/2)*(ln((x-2^(1/2)*(a/b)^(1/4)*x^(1/2)+(a/b)^(1/2))/((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x))+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)+1)+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)-1))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.91

$$\int \frac{\sqrt{x}}{(a + bx^2)^3} dx$$

$$= \frac{5(a^2b^2x^4 + 2a^3bx^2 + a^4)\left(-\frac{1}{a^9b^3}\right)^{\frac{1}{4}} \log\left(a^7b^2\left(-\frac{1}{a^9b^3}\right)^{\frac{3}{4}} + \sqrt{x}\right) - 5\left(ia^2b^2x^4 + 2ia^3bx^2 + ia^4\right)\left(-\frac{1}{a^9b^3}\right)^{\frac{1}{4}} \log\left(ia^7b^2\left(-\frac{1}{a^9b^3}\right)^{\frac{3}{4}} + \sqrt{x}\right)}{1}$$

input `integrate(x^(1/2)/(b*x^2+a)^3,x, algorithm="fricas")`

output `1/64*(5*(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)*(-1/(a^9*b^3))^(1/4)*log(a^7*b^2*(-1/(a^9*b^3))^(3/4) + sqrt(x)) - 5*(I*a^2*b^2*x^4 + 2*I*a^3*b*x^2 + I*a^4)*(-1/(a^9*b^3))^(1/4)*log(I*a^7*b^2*(-1/(a^9*b^3))^(3/4) + sqrt(x)) - 5*(-I*a^2*b^2*x^4 - 2*I*a^3*b*x^2 - I*a^4)*(-1/(a^9*b^3))^(1/4)*log(-I*a^7*b^2*(-1/(a^9*b^3))^(3/4) + sqrt(x)) - 5*(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)*(-1/(a^9*b^3))^(1/4)*log(-a^7*b^2*(-1/(a^9*b^3))^(3/4) + sqrt(x)) + 4*(5*b*x^3 + 9*a*x)*sqrt(x))/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 887 vs. 2(119) = 238.

Time = 106.02 (sec) , antiderivative size = 887, normalized size of antiderivative = 6.03

$$\int \frac{\sqrt{x}}{(a + bx^2)^3} dx = \text{Too large to display}$$

input `integrate(x**(1/2)/(b*x**2+a)**3,x)`

output

```
Piecewise((zoo/x**(9/2), Eq(a, 0) & Eq(b, 0)), (2*x**(3/2)/(3*a**3), Eq(b,
0)), (-2/(9*b**3*x**(9/2)), Eq(a, 0)), (5*a**2*log(sqrt(x) - (-a/b)**(1/4
)))/(64*a**4*b*(-a/b)**(1/4) + 128*a**3*b**2*x**2*(-a/b)**(1/4) + 64*a**2*b
**3*x**4*(-a/b)**(1/4)) - 5*a**2*log(sqrt(x) + (-a/b)**(1/4))/(64*a**4*b*(
-a/b)**(1/4) + 128*a**3*b**2*x**2*(-a/b)**(1/4) + 64*a**2*b**3*x**4*(-a/b)
**2*(1/4)) + 10*a**2*atan(sqrt(x)/(-a/b)**(1/4))/(64*a**4*b*(-a/b)**(1/4) +
128*a**3*b**2*x**2*(-a/b)**(1/4) + 64*a**2*b**3*x**4*(-a/b)**(1/4)) + 36*a
*b*x**(3/2)*(-a/b)**(1/4)/(64*a**4*b*(-a/b)**(1/4) + 128*a**3*b**2*x**2*(-
a/b)**(1/4) + 64*a**2*b**3*x**4*(-a/b)**(1/4)) + 10*a*b*x**2*log(sqrt(x) -
(-a/b)**(1/4))/(64*a**4*b*(-a/b)**(1/4) + 128*a**3*b**2*x**2*(-a/b)**(1/4
)) + 64*a**2*b**3*x**4*(-a/b)**(1/4)) - 10*a*b*x**2*log(sqrt(x) + (-a/b)**(
1/4))/(64*a**4*b*(-a/b)**(1/4) + 128*a**3*b**2*x**2*(-a/b)**(1/4) + 64*a**
2*b**3*x**4*(-a/b)**(1/4)) + 20*a*b*x**2*atan(sqrt(x)/(-a/b)**(1/4))/(64*a
**4*b*(-a/b)**(1/4) + 128*a**3*b**2*x**2*(-a/b)**(1/4) + 64*a**2*b**3*x**4
*(-a/b)**(1/4)) + 20*b**2*x**(7/2)*(-a/b)**(1/4)/(64*a**4*b*(-a/b)**(1/4)
+ 128*a**3*b**2*x**2*(-a/b)**(1/4) + 64*a**2*b**3*x**4*(-a/b)**(1/4)) + 5*
b**2*x**4*log(sqrt(x) - (-a/b)**(1/4))/(64*a**4*b*(-a/b)**(1/4) + 128*a**3
*b**2*x**2*(-a/b)**(1/4) + 64*a**2*b**3*x**4*(-a/b)**(1/4)) - 5*b**2*x**4*
log(sqrt(x) + (-a/b)**(1/4))/(64*a**4*b*(-a/b)**(1/4) + 128*a**3*b**2*x**2
*(-a/b)**(1/4) + 64*a**2*b**3*x**4*(-a/b)**(1/4)) + 10*b**2*x**4*atan(s...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.48

$$\int \frac{\sqrt{x}}{(a+bx^2)^3} dx = \frac{5bx^{\frac{7}{2}} + 9ax^{\frac{3}{2}}}{16(a^2b^2x^4 + 2a^3bx^2 + a^4)}$$

$$+ \frac{5 \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} + 2\sqrt{b}\sqrt{x})}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{b}\sqrt{b}} \right) + \frac{2\sqrt{2} \arctan\left(-\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} - 2\sqrt{b}\sqrt{x})}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{b}\sqrt{b}}}{128a^2} - \frac{\sqrt{2} \log(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{bx} + \sqrt{a})}{a^{\frac{1}{4}}b^{\frac{3}{4}}} + \frac{\sqrt{2} \log(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} - \sqrt{bx} + \sqrt{a})}{a^{\frac{1}{4}}b^{\frac{3}{4}}}$$

input

```
integrate(x^(1/2)/(b*x^2+a)^3,x, algorithm="maxima")
```

output

```
1/16*(5*b*x^(7/2) + 9*a*x^(3/2))/(a^2*b^2*x^4 + 2*a^3*b*x^2 + a^4) + 5/128
*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x)
))/sqrt(sqrt(a)*sqrt(b))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) + 2*sqrt(2)*arct
an(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)
*sqrt(b))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) - sqrt(2)*log(sqrt(2)*a^(1/4)*b
^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4)) + sqrt(2)*log(-sqr
t(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4))/a^2
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.42

$$\int \frac{\sqrt{x}}{(a+bx^2)^3} dx = \frac{5bx^{\frac{7}{2}} + 9ax^{\frac{3}{2}}}{16(bx^2+a)^2a^2} + \frac{5\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{\frac{a}{b}}^{\frac{1}{4}} + 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{64a^3b^3}$$

$$+ \frac{5\sqrt{2}(ab^3)^{\frac{3}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{\frac{a}{b}}^{\frac{1}{4}} - 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{64a^3b^3}$$

$$- \frac{5\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}} + x + \sqrt{\frac{a}{b}}\right)}{128a^3b^3}$$

$$+ \frac{5\sqrt{2}(ab^3)^{\frac{3}{4}} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}} + x + \sqrt{\frac{a}{b}}\right)}{128a^3b^3}$$

input

```
integrate(x^(1/2)/(b*x^2+a)^3,x, algorithm="giac")
```

output

```
1/16*(5*b*x^(7/2) + 9*a*x^(3/2))/((b*x^2 + a)^2*a^2) + 5/64*sqrt(2)*(a*b^3
)^(3/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/
(a^3*b^3) + 5/64*sqrt(2)*(a*b^3)^(3/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)
^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/(a^3*b^3) - 5/128*sqrt(2)*(a*b^3)^(3/4)*l
og(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^3*b^3) + 5/128*sqrt(2)*
(a*b^3)^(3/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^3*b^3)
```

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{x}}{(a + bx^2)^3} dx = \frac{\frac{9x^{3/2}}{16a} + \frac{5bx^{7/2}}{16a^2}}{a^2 + 2abx^2 + b^2x^4} + \frac{5 \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{9/4}b^{3/4}} - \frac{5 \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{9/4}b^{3/4}}$$

input `int(x^(1/2)/(a + b*x^2)^3,x)`output `((9*x^(3/2))/(16*a) + (5*b*x^(7/2))/(16*a^2))/(a^2 + b^2*x^4 + 2*a*b*x^2) + (5*atan((b^(1/4)*x^(1/2))/(-a)^(1/4)))/(32*(-a)^(9/4)*b^(3/4)) - (5*atanh((b^(1/4)*x^(1/2))/(-a)^(1/4)))/(32*(-a)^(9/4)*b^(3/4))`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 478, normalized size of antiderivative = 3.25

$$\int \frac{\sqrt{x}}{(a + bx^2)^3} dx = \text{Too large to display}$$

input `int(x^(1/2)/(b*x^2+a)^3,x)`

output

```
( - 10*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a**2 - 20*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a*b*x**2 - 10*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b**2*x**4 + 10*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a**2 + 20*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a*b*x**2 + 10*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b**2*x**4 + 5*b**(1/4)*a**(3/4)*sqrt(2)*log( - sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a**2 + 10*b**(1/4)*a**(3/4)*sqrt(2)*log( - sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 + 5*b**(1/4)*a**(3/4)*sqrt(2)*log( - sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 - 5*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a**2 - 10*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 - 5*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 + 72*sqrt(x)*a**2*b*x + 40*sqrt(x)*a*b**2*x**3)/(128*a**3*b*(a**2 + 2*a*b*x**2 + b**2*x**4))
```

3.131 $\int \frac{x^{3/2}}{(a+bx^2)^3} dx$

Optimal result	884
Mathematica [A] (verified)	884
Rubi [A] (verified)	885
Maple [A] (verified)	890
Fricas [C] (verification not implemented)	890
Sympy [F(-1)]	891
Maxima [A] (verification not implemented)	891
Giac [A] (verification not implemented)	892
Mupad [B] (verification not implemented)	893
Reduce [B] (verification not implemented)	893

Optimal result

Integrand size = 15, antiderivative size = 139

$$\int \frac{x^{3/2}}{(a+bx^2)^3} dx = \frac{\sqrt{x}(-3a+bx^2)}{16ab(a+bx^2)^2} + \frac{3 \left(\arctan \left(\frac{\sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}} - x} \right) + \log \left(\frac{\sqrt{\frac{a}{b}} + \sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x+x}}{\sqrt{a+bx^2}} \right) \right)}{32\sqrt{2}a \left(\frac{a}{b}\right)^{3/4} b^2}$$

output `1/16*x^(1/2)*(b*x^2-3*a)/a/b/(b*x^2+a)^2+3/64*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))+ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/a/(a/b)^(3/4)/b^2`

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.99

$$\int \frac{x^{3/2}}{(a+bx^2)^3} dx = \frac{4a^{3/4} \sqrt[4]{b} \sqrt{x} (-3a+bx^2)}{(a+bx^2)^2} - 3\sqrt{2} \arctan \left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x}} \right) + 3\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x}}{\sqrt{a+\sqrt{bx}}} \right)$$

input `Integrate[x^(3/2)/(a + b*x^2)^3,x]`

output

$$\left((4a^{3/4}b^{1/4}\sqrt{x}(-3a + bx^2))/(a + bx^2)^2 - 3\sqrt{2}\operatorname{ArcTan}[(\sqrt{a} - \sqrt{b}x)/(\sqrt{2}a^{1/4}b^{1/4}\sqrt{x})] + 3\sqrt{2}\operatorname{ArcTanh}[(\sqrt{2}a^{1/4}b^{1/4}\sqrt{x})/(\sqrt{a} + \sqrt{b}x)] \right) / (64a^{7/4}b^{5/4})$$
Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.94, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.733$, Rules used = {252, 253, 266, 755, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^{3/2}}{(a + bx^2)^3} dx \\ & \quad \downarrow \text{252} \\ & \frac{\int \frac{1}{\sqrt{x}(bx^2+a)^2} dx}{8b} - \frac{\sqrt{x}}{4b(a + bx^2)^2} \\ & \quad \downarrow \text{253} \\ & \frac{3 \int \frac{1}{\sqrt{x}(bx^2+a)} dx}{8b} + \frac{\sqrt{x}}{2a(a+bx^2)} - \frac{\sqrt{x}}{4b(a + bx^2)^2} \\ & \quad \downarrow \text{266} \\ & \frac{3 \int \frac{1}{bx^2+a} d\sqrt{x}}{8b} + \frac{\sqrt{x}}{2a(a+bx^2)} - \frac{\sqrt{x}}{4b(a + bx^2)^2} \\ & \quad \downarrow \text{755} \\ & \frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} \right)}{8b} + \frac{\sqrt{x}}{2a(a+bx^2)} - \frac{\sqrt{x}}{4b(a + bx^2)^2} \\ & \quad \downarrow \text{1476} \end{aligned}$$

$$\begin{aligned}
 & \frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{x-\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}} \frac{d\sqrt{x}}{\sqrt{b}}}{2\sqrt{b}} + \frac{\int \frac{1}{x+\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt{a}} \frac{d\sqrt{x}}{\sqrt{b}}}{2\sqrt{b}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} - \frac{\sqrt{x}}{4b(a+bx^2)^2} \\
 & \quad \downarrow 1082 \\
 & \frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\int \frac{1}{-x-1} d\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} - \frac{\int \frac{1}{-x-1} d\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{a}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} - \frac{\sqrt{x}}{4b(a+bx^2)^2} \\
 & \quad \downarrow 217 \\
 & \frac{3 \left(\frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{a}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} - \frac{\sqrt{x}}{4b(a+bx^2)^2} \\
 & \quad \downarrow 1479 \\
 & \frac{3 \left(\frac{\int -\frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int -\frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}}{\sqrt{b}}+\frac{\sqrt{a}}{\sqrt{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right)}{2a} + \frac{\sqrt{x}}{2a(a+bx^2)} \\
 & \quad \frac{\sqrt{x}}{4b(a+bx^2)^2} \\
 & \quad \downarrow 25
 \end{aligned}$$

$$\left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{b} \left(x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \left(\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a} \right)}{\sqrt[4]{b} \left(x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}} \right)} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

$$\frac{8b}{4b(a+bx^2)^2}$$

↓ 27

$$\left(\frac{\int \frac{\sqrt{2} \sqrt[4]{a} - 2 \sqrt[4]{b} \sqrt{x}}{x - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\int \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + \sqrt[4]{a}}{x + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{x} + \sqrt{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2 \sqrt[4]{a} \sqrt[4]{b}} + \frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

$$\frac{8b}{4b(a+bx^2)^2}$$

↓ 1103

$$\left(\frac{\arctan \left(\frac{\sqrt{2} \sqrt[4]{b} \sqrt{x} + 1}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2} \sqrt[4]{b} \sqrt{x}}{\sqrt[4]{a}} \right)}{\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} + \frac{\log \left(\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx} \right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} - \frac{\log \left(-\sqrt{2} \sqrt[4]{a} \sqrt[4]{b} \sqrt{x} + \sqrt{a} + \sqrt{bx} \right)}{2\sqrt{2} \sqrt[4]{a} \sqrt[4]{b}} \right) + \frac{\sqrt{x}}{2a(a+bx^2)}$$

$$\frac{8b}{4b(a+bx^2)^2}$$

input

Int [x^(3/2)/(a + b*x^2)^3, x]

output

$$-1/4*\text{Sqrt}[x]/(b*(a + b*x^2)^2) + (\text{Sqrt}[x]/(2*a*(a + b*x^2)) + (3*((-\text{ArcTan}[1 - (\text{Sqrt}[2]*b^{1/4}*\text{Sqrt}[x])/a^{1/4}]/(\text{Sqrt}[2]*a^{1/4}*b^{1/4})) + \text{ArcTan}[1 + (\text{Sqrt}[2]*b^{1/4}*\text{Sqrt}[x])/a^{1/4}]/(\text{Sqrt}[2]*a^{1/4}*b^{1/4})))/(2*\text{Sqrt}[a]) + (-1/2*\text{Log}[\text{Sqrt}[a] - \text{Sqrt}[2]*a^{1/4}*b^{1/4}*\text{Sqrt}[x] + \text{Sqrt}[b]*x]/(\text{Sqrt}[2]*a^{1/4}*b^{1/4}) + \text{Log}[\text{Sqrt}[a] + \text{Sqrt}[2]*a^{1/4}*b^{1/4}*\text{Sqrt}[x] + \text{Sqrt}[b]*x]/(2*\text{Sqrt}[2]*a^{1/4}*b^{1/4}))/2*\text{Sqrt}[a]))/(2*a))/(8*b)$$

Defintions of rubi rules used

rule 25

$$\text{Int}[-(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \quad \text{Int}[\text{Fx}, x], x]$$

rule 27

$$\text{Int}[(a_)*(\text{Fx}_), x_Symbol] \rightarrow \text{Simp}[a \quad \text{Int}[\text{Fx}, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[\text{Fx}, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$$

rule 217

$$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$$

rule 252

$$\text{Int}[((c_)*(x_))^{(m_)*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a + b*x^2)^{(p+1)}/(2*b*(p+1))), x] - \text{Simp}[c^2*((m-1)/(2*b*(p+1))) \quad \text{Int}[(c*x)^{(m-2)}*(a + b*x^2)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{LtQ}[p, -1] \&\& \text{GtQ}[m, 1] \&\& \text{!ILtQ}[(m + 2*p + 3)/2, 0] \&\& \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 253

$$\text{Int}[((c_)*(x_))^{(m_)*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(-(c*x)^{(m+1)}*((a + b*x^2)^{(p+1)}/(2*a*c*(p+1))), x] + \text{Simp}[(m + 2*p + 3)/(2*a*(p+1)) \quad \text{Int}[(c*x)^m*(a + b*x^2)^{(p+1)}, x], x] /; \text{FreeQ}[\{a, b, c, m\}, x] \&\& \text{LtQ}[p, -1] \&\& \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 266

$$\text{Int}[((c_)*(x_))^{(m_)*((a_) + (b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \quad \text{Subst}[\text{Int}[x^{(k*(m+1)-1)}*(a + b*(x^{(2*k)}/c^2))^p, x], x, (c*x)^{(1/k)}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, 2, m, p, x]$$

rule 755 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Simp[1/(2*r) Int[(r - s*x^2)/(a + b*x^4), x], x] + Simp[1/(2*r) Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`

rule 1476 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x], x] + Simp[e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]`

rule 1479 `Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.99

method	result	size
derivativedivides	$\frac{x^{\frac{5}{2}} - 3\sqrt{x}}{16a - 16b} + \frac{3\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2} \left(\ln\left(\frac{\sqrt{\frac{a}{b} + \sqrt{2}}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}{x - \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\sqrt{\frac{a}{b}}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}+1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}-1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) \right)}{128ba^2}$	138
default	$\frac{x^{\frac{5}{2}} - 3\sqrt{x}}{16a - 16b} + \frac{3\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{2} \left(\ln\left(\frac{\sqrt{\frac{a}{b} + \sqrt{2}}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+x}}{x - \sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}\sqrt{x+\sqrt{\frac{a}{b}}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}+1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) + 2\arctan\left(\frac{\sqrt{2}\sqrt{x}-1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right) \right)}{128ba^2}$	138

input `int(x^(3/2)/(b*x^2+a)^3,x,method=_RETURNVERBOSE)`

output
$$2*(1/32/a*x^{(5/2)}-3/32*x^{(1/2)}/b)/(b*x^2+a)^2+3/128/b/a^2*(a/b)^{(1/4)}*2^{(1/2)}*(\ln(((a/b)^{(1/2)}+2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)+x}/(x-2^{(1/2)}*(a/b)^{(1/4)}*x^{(1/2)}+(a/b)^{(1/2)}))+2*\arctan(2^{(1/2)}/(a/b)^{(1/4)}*x^{(1/2)}+1)+2*\arctan(2^{(1/2)}/(a/b)^{(1/4)}*x^{(1/2)}-1))$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 279, normalized size of antiderivative = 2.01

$$\int \frac{x^{3/2}}{(a+bx^2)^3} dx = \frac{3(ab^3x^4 + 2a^2b^2x^2 + a^3b)\left(-\frac{1}{a^7b^5}\right)^{\frac{1}{4}} \log\left(a^2b\left(-\frac{1}{a^7b^5}\right)^{\frac{1}{4}} + \sqrt{x}\right) - 3(-iab^3x^4 - 2ia^2b^2x^2}$$

input `integrate(x^(3/2)/(b*x^2+a)^3,x, algorithm="fricas")`

```
output 1/64*(3*(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b)*(-1/(a^7*b^5))^(1/4)*log(a^2*b
*(-1/(a^7*b^5))^(1/4) + sqrt(x)) - 3*(-I*a*b^3*x^4 - 2*I*a^2*b^2*x^2 - I*a
^3*b)*(-1/(a^7*b^5))^(1/4)*log(I*a^2*b*(-1/(a^7*b^5))^(1/4) + sqrt(x)) - 3
*(I*a*b^3*x^4 + 2*I*a^2*b^2*x^2 + I*a^3*b)*(-1/(a^7*b^5))^(1/4)*log(-I*a^2
*b*(-1/(a^7*b^5))^(1/4) + sqrt(x)) - 3*(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b)
*(-1/(a^7*b^5))^(1/4)*log(-a^2*b*(-1/(a^7*b^5))^(1/4) + sqrt(x)) + 4*(b*x^
2 - 3*a)*sqrt(x))/(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{x^{3/2}}{(a + bx^2)^3} dx = \text{Timed out}$$

```
input integrate(x**(3/2)/(b*x**2+a)**3,x)
```

```
output Timed out
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.59

$$\int \frac{x^{3/2}}{(a + bx^2)^3} dx = \frac{bx^{\frac{5}{2}} - 3a\sqrt{x}}{16(ab^3x^4 + 2a^2b^2x^2 + a^3b)} + \frac{3 \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} + 2\sqrt{b}\sqrt{x})}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{a}\sqrt{b}} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}} - 2\sqrt{b}\sqrt{x})}{2\sqrt{a}\sqrt{b}}\right)}{\sqrt{a}\sqrt{a}\sqrt{b}} + \frac{\sqrt{2} \log(\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} - \frac{\sqrt{2} \log(-\sqrt{2}a^{\frac{1}{4}}b^{\frac{1}{4}}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{\frac{3}{4}}b^{\frac{1}{4}}} \right)}{128ab}$$

```
input integrate(x^(3/2)/(b*x^2+a)^3,x, algorithm="maxima")
```

output

```
1/16*(b*x^(5/2) - 3*a*sqrt(x))/(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b) + 3/128
*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x)
))/sqrt(sqrt(a)*sqrt(b))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + 2*sqrt(2)*arct
an(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)
*sqrt(b))/sqrt(a)*sqrt(sqrt(a)*sqrt(b)) + sqrt(2)*log(sqrt(2)*a^(1/4)*b
^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4)) - sqrt(2)*log(-sqr
t(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(3/4)*b^(1/4))/(a*
b)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.52

$$\int \frac{x^{3/2}}{(a+bx^2)^3} dx = \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}+2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{64a^2b^2}$$

$$+ \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{\frac{1}{4}}-2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{\frac{1}{4}}}\right)}{64a^2b^2}$$

$$+ \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{128a^2b^2}$$

$$- \frac{3\sqrt{2}(ab^3)^{\frac{1}{4}} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{\frac{1}{4}}+x+\sqrt{\frac{a}{b}}\right)}{128a^2b^2} + \frac{bx^{\frac{5}{2}}-3a\sqrt{x}}{16(bx^2+a)^2ab}$$

input

```
integrate(x^(3/2)/(b*x^2+a)^3,x, algorithm="giac")
```

output

```
3/64*sqrt(2)*(a*b^3)^(1/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt
t(x))/(a/b)^(1/4))/(a^2*b^2) + 3/64*sqrt(2)*(a*b^3)^(1/4)*arctan(-1/2*sqrt
(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/(a^2*b^2) + 3/128*sqrt(
2)*(a*b^3)^(1/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^2*b^2
) - 3/128*sqrt(2)*(a*b^3)^(1/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqr
t(a/b))/(a^2*b^2) + 1/16*(b*x^(5/2) - 3*a*sqrt(x))/((b*x^2 + a)^2*a*b)
```

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.61

$$\int \frac{x^{3/2}}{(a + bx^2)^3} dx = \frac{\frac{x^{5/2}}{16a} - \frac{3\sqrt{x}}{16b}}{a^2 + 2abx^2 + b^2x^4} + \frac{3 \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{7/4}b^{5/4}} + \frac{3 \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{7/4}b^{5/4}}$$

input `int(x^(3/2)/(a + b*x^2)^3,x)`output `(x^(5/2)/(16*a) - (3*x^(1/2))/(16*b))/(a^2 + b^2*x^4 + 2*a*b*x^2) + (3*atan(b^(1/4)*x^(1/2)/(-a)^(1/4))/(32*(-a)^(7/4)*b^(5/4)) + (3*atanh(b^(1/4)*x^(1/2)/(-a)^(1/4))/(32*(-a)^(7/4)*b^(5/4))`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 477, normalized size of antiderivative = 3.43

$$\int \frac{x^{3/2}}{(a + bx^2)^3} dx = \text{Too large to display}$$

input `int(x^(3/2)/(b*x^2+a)^3,x)`

output

```
( - 6*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)
)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))*a**2 - 12*b**(3/4)*a**(1/4)*sqrt(2)
)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*
sqrt(2))*a*b*x**2 - 6*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*s
qrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))*b**2*x**4 + 6*b**
(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b)
)/(b**(1/4)*a**(1/4)*sqrt(2))*a**2 + 12*b**(3/4)*a**(1/4)*sqrt(2)*atan((b
**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))
)*a*b*x**2 + 6*b**(3/4)*a**(1/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) +
2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2))*b**2*x**4 - 3*b**(3/4)*a**
(1/4)*sqrt(2)*log( - sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)
*x)*a**2 - 6*b**(3/4)*a**(1/4)*sqrt(2)*log( - sqrt(x)*b**(1/4)*a**(1/4)*sq
rt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 - 3*b**(3/4)*a**(1/4)*sqrt(2)*log( -
sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 + 3*b**
*(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) +
sqrt(b)*x)*a**2 + 6*b**(3/4)*a**(1/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)
)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 + 3*b**(3/4)*a**(1/4)*sqrt(2)*lo
g(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 - 24*
sqrt(x)*a**2*b + 8*sqrt(x)*a*b**2*x**2)/(128*a**2*b**2*(a**2 + 2*a*b*x**2
+ b**2*x**4))
```

3.132 $\int \frac{x^{5/2}}{(a+bx^2)^3} dx$

Optimal result	895
Mathematica [A] (verified)	896
Rubi [A] (verified)	896
Maple [A] (verified)	901
Fricas [C] (verification not implemented)	902
Sympy [F(-1)]	902
Maxima [A] (verification not implemented)	903
Giac [A] (verification not implemented)	904
Mupad [B] (verification not implemented)	904
Reduce [B] (verification not implemented)	905

Optimal result

Integrand size = 15, antiderivative size = 177

$$\int \frac{x^{5/2}}{(a+bx^2)^3} dx = -\frac{2x^{3/2}}{5b(a+bx^2)^2} + \frac{3a \left(x^{3/2} \left(\frac{1}{4a(a+bx^2)^2} + \frac{5}{16a^2(a+bx^2)} \right) + \frac{5 \left(\arctan \left(\frac{\sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x}}{\sqrt{\frac{a}{b}} - x} \right) - \log \left(\frac{\sqrt{\frac{a}{b}} + \sqrt{2} \sqrt[4]{\frac{a}{b}} \sqrt{x+x}}{\sqrt{a+bx^2}} \right) \right)}{32\sqrt{2}a^2 \sqrt[4]{\frac{a}{b}} b} \right)}{5b}$$

output

```
-2/5*x^(3/2)/b/(b*x^2+a)^2+3/5*a*(x^(3/2)*(1/4/a/(b*x^2+a)^2+5/16/a^2/(b*x^2+a))+5/64*(arctan(2^(1/2)*(a/b)^(1/4)*x^(1/2)/((a/b)^(1/2)-x))-ln(((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x)/(b*x^2+a)^(1/2)))*2^(1/2)/a^2/(a/b)^(1/4)/b)/b
```


Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.77

$$\int \frac{x^{5/2}}{(a + bx^2)^3} dx = \frac{-\frac{4\sqrt[4]{ab^{3/4}x^{3/2}(a-3bx^2)}}{(a+bx^2)^2} - 3\sqrt{2} \arctan\left(\frac{\sqrt{a}-\sqrt{bx}}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}\right) - 3\sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}}{\sqrt{a}+\sqrt{bx}}\right)}{64a^{5/4}b^{7/4}}$$

input `Integrate[x^(5/2)/(a + b*x^2)^3,x]`

output `((-4*a^(1/4)*b^(3/4)*x^(3/2)*(a - 3*b*x^2))/(a + b*x^2)^2 - 3*Sqrt[2]*ArcTan[(Sqrt[a] - Sqrt[b]*x)/(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])] - 3*Sqrt[2]*ArcTanh[(Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x])/(Sqrt[a] + Sqrt[b]*x)])/(64*a^(5/4)*b^(7/4))`

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.53, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.733$, Rules used = {252, 253, 266, 826, 1476, 1082, 217, 1479, 25, 27, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x^{5/2}}{(a + bx^2)^3} dx \\ & \quad \downarrow \text{252} \\ & \frac{3 \int \frac{\sqrt{x}}{(bx^2+a)^2} dx}{8b} - \frac{x^{3/2}}{4b(a + bx^2)^2} \\ & \quad \downarrow \text{253} \\ & \frac{3 \left(\frac{\int \frac{\sqrt{x}}{bx^2+a} dx}{4a} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8b} - \frac{x^{3/2}}{4b(a + bx^2)^2} \\ & \quad \downarrow \text{266} \end{aligned}$$

$$\begin{aligned}
 & \frac{3 \left(\frac{\int \frac{x}{bx^2+a} d\sqrt{x}}{2a} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8b} - \frac{x^{3/2}}{4b(a+bx^2)^2} \\
 & \quad \downarrow \text{826} \\
 & \frac{3 \left(\frac{\int \frac{\sqrt{bx}+\sqrt{a}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8b} - \frac{x^{3/2}}{4b(a+bx^2)^2} \\
 & \quad \downarrow \text{1476} \\
 & \frac{3 \left(\frac{\int \frac{\frac{1}{x-\sqrt{2}\sqrt[4]{a}\sqrt{x}+\frac{\sqrt{a}}{\sqrt{b}}}}{2\sqrt{b}} d\sqrt{x}}{2\sqrt{b}} + \frac{\int \frac{\frac{1}{x+\sqrt{2}\sqrt[4]{a}\sqrt{x}+\frac{\sqrt{a}}{\sqrt{b}}}}{2\sqrt{b}} d\sqrt{x}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8b} - \frac{x^{3/2}}{4b(a+bx^2)^2} \\
 & \quad \downarrow \text{1082} \\
 & \frac{3 \left(\frac{\int \frac{\frac{1}{-x-1} d \left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{4\sqrt{a}} \right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} - \frac{\int \frac{\frac{1}{-x-1} d \left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{4\sqrt{a}} + 1 \right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8b} - \frac{x^{3/2}}{4b(a+bx^2)^2} \\
 & \quad \downarrow \text{217} \\
 & \frac{3 \left(\frac{\frac{\arctan \left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{4\sqrt{a}} + 1 \right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan \left(1 - \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{4\sqrt{a}} \right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}}}{2\sqrt{b}} - \frac{\int \frac{\sqrt{a}-\sqrt{bx}}{bx^2+a} d\sqrt{x}}{2\sqrt{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right)}{8b} - \frac{x^{3/2}}{4b(a+bx^2)^2} \\
 & \quad \downarrow \text{1479}
 \end{aligned}$$

$$3 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right) + \frac{x^{3/2}}{2a(a+bx^2)}$$

$$\frac{x^{3/2}}{4b(a+bx^2)^2} \quad 8b$$

↓ 25

$$3 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{b}\left(x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\left(\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}\right)}{\sqrt[4]{b}\left(x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}\right)} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} \right) + \frac{x^{3/2}}{2a(a+bx^2)}$$

$$\frac{x^{3/2}}{4b(a+bx^2)^2} \quad 8b$$

↓ 27

$$3 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+1}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\int \frac{\sqrt{2}\sqrt[4]{a}-2\sqrt[4]{b}\sqrt{x}}{x-\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{\int \frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}+\sqrt[4]{a}}{x+\frac{\sqrt{2}\sqrt[4]{a}\sqrt{x}+\sqrt[4]{a}}{\sqrt[4]{b}}} d\sqrt{x}}{2\sqrt[4]{a}\sqrt[4]{b}} \right) + \frac{x^{3/2}}{2a(a+bx^2)}$$

$$\frac{x^{3/2}}{4b(a+bx^2)^2} \quad 8b$$

↓ 1103

$$3 \left(\frac{\arctan\left(\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}+1\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\arctan\left(1-\frac{\sqrt{2}\sqrt[4]{b}\sqrt{x}}{\sqrt[4]{a}}\right)}{\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} - \frac{\log\left(-\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}\sqrt{x}+\sqrt{a}+\sqrt{bx}\right)}{2\sqrt{2}\sqrt[4]{a}\sqrt[4]{b}} + \frac{x^{3/2}}{2a(a+bx^2)} \right) \\ \frac{x^{3/2}}{4b(a+bx^2)^2} \quad \frac{8b}{4b(a+bx^2)^2}$$

input `Int[x^(5/2)/(a + b*x^2)^3,x]`

output `-1/4*x^(3/2)/(b*(a + b*x^2)^2) + (3*(x^(3/2)/(2*a*(a + b*x^2))) + ((-(ArcTan[1 - (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4))) + ArcTan[1 + (Sqrt[2]*b^(1/4)*Sqrt[x])/a^(1/4)]/(Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]) - (-1/2*Log[Sqrt[a] - Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(Sqrt[2]*a^(1/4)*b^(1/4)) + Log[Sqrt[a] + Sqrt[2]*a^(1/4)*b^(1/4)*Sqrt[x] + Sqrt[b]*x]/(2*Sqrt[2]*a^(1/4)*b^(1/4)))/(2*Sqrt[b]))/(2*a)))/(8*b)`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 252 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[c*(c*x)^{(m-1)}*((a+b*x^2)^{(p+1)}/(2*b*(p+1))), x] - \text{Simp}[c^2*(m-1)/(2*b*(p+1)) \text{Int}[(c*x)^{(m-2)}*(a+b*x^2)^{(p+1)}, x], x] /;$ FreeQ[{a, b, c}, x] && LtQ[p, -1] && GtQ[m, 1] && !ILtQ[(m+2*p+3)/2, 0] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 253 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(-c*x)^{(m+1)}*((a+b*x^2)^{(p+1)}/(2*a*c*(p+1))), x] + \text{Simp}[(m+2*p+3)/(2*a*(p+1)) \text{Int}[(c*x)^m*(a+b*x^2)^{(p+1)}, x], x] /;$ FreeQ[{a, b, c, m}, x] && LtQ[p, -1] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 266 $\text{Int}[\{(c_)*(x_)\}^{(m_)}*((a_)+(b_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Simp}[k/c \text{Subst}[\text{Int}[x^{k*(m+1)-1}*(a+b*(x^{2*k}/c^2))^p, x], x, (c*x)^{1/k}], x]] /;$ FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]

rule 826 $\text{Int}[(x_)^2/((a_)+(b_)*(x_)^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Simp}[1/(2*s) \text{Int}[(r+s*x^2)/(a+b*x^4), x], x] - \text{Simp}[1/(2*s) \text{Int}[(r-s*x^2)/(a+b*x^4), x], x]] /;$ FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

rule 1082 $\text{Int}[\{(a_)+(b_)*(x_)+(c_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q-x^2), x], x, 1+2*c*(x/b)], x] /;$ RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2-4*a*c]) /;

rule 1103 $\text{Int}[\{(d_)+(e_)*(x_)\} / \{(a_)+(b_)*(x_)+(c_)*(x_)^2\}, x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a+b*x+c*x^2, x]]/b), x] /;$ FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d-b*e, 0]

rule 1476

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Simp[e/(2*c) Int[1/Simp[d/e + q*x + x^2, x], x] + Simp[
e/(2*c) Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x]
&& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

rule 1479

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Simp[e/(2*c*q) Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Simp[e/(2*c*q) Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; F
reeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{\frac{3x^{\frac{7}{2}} - x^{\frac{3}{2}}}{16a - 16b}}{(x^2b+a)^2} + \frac{3\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x} + \sqrt{\frac{a}{b}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x} + x} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} + 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} - 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) \right)}{128b^2a \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	138
default	$\frac{\frac{3x^{\frac{7}{2}} - x^{\frac{3}{2}}}{16a - 16b}}{(x^2b+a)^2} + \frac{3\sqrt{2} \left(\ln \left(\frac{x - \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x} + \sqrt{\frac{a}{b}}}{\sqrt{\frac{a}{b}} + \sqrt{2} \left(\frac{a}{b}\right)^{\frac{1}{4}} \sqrt{x} + x} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} + 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{x} - 1}{\left(\frac{a}{b}\right)^{\frac{1}{4}}} \right) \right)}{128b^2a \left(\frac{a}{b}\right)^{\frac{1}{4}}}$	138

input

```
int(x^(5/2)/(b*x^2+a)^3,x,method=_RETURNVERBOSE)
```

output

```
2*(3/32/a*x^(7/2)-1/32*x^(3/2)/b)/(b*x^2+a)^2+3/128/b^2/a/(a/b)^(1/4)*2^(1/2)*
(ln((x-2^(1/2)*(a/b)^(1/4)*x^(1/2)+(a/b)^(1/2))/((a/b)^(1/2)+2^(1/2)*(a/b)^(1/4)*x^(1/2)+x))+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)+1)+2*arctan(2^(1/2)/(a/b)^(1/4)*x^(1/2)-1))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.11 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.63

$$\int \frac{x^{5/2}}{(a + bx^2)^3} dx = \frac{3(ab^3x^4 + 2a^2b^2x^2 + a^3b)\left(-\frac{1}{a^5b^7}\right)^{\frac{1}{4}} \log\left(a^4b^5\left(-\frac{1}{a^5b^7}\right)^{\frac{3}{4}} + \sqrt{x}\right) - 3(ia^3b^3x^4 + 2ia^2b^2x^2 - 3ia^3b)}{(a + bx^2)^3}$$

input `integrate(x^(5/2)/(b*x^2+a)^3,x, algorithm="fricas")`

output `1/64*(3*(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b)*(-1/(a^5*b^7))^(1/4)*log(a^4*b^5*(-1/(a^5*b^7))^(3/4) + sqrt(x)) - 3*(I*a*b^3*x^4 + 2*I*a^2*b^2*x^2 + I*a^3*b)*(-1/(a^5*b^7))^(1/4)*log(I*a^4*b^5*(-1/(a^5*b^7))^(3/4) + sqrt(x)) - 3*(-I*a*b^3*x^4 - 2*I*a^2*b^2*x^2 - I*a^3*b)*(-1/(a^5*b^7))^(1/4)*log(-I*a^4*b^5*(-1/(a^5*b^7))^(3/4) + sqrt(x)) - 3*(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b)*(-1/(a^5*b^7))^(1/4)*log(-a^4*b^5*(-1/(a^5*b^7))^(3/4) + sqrt(x)) + 4*(3*b*x^3 - a*x)*sqrt(x))/(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b)`

Sympy [F(-1)]

Timed out.

$$\int \frac{x^{5/2}}{(a + bx^2)^3} dx = \text{Timed out}$$

input `integrate(x**(5/2)/(b*x**2+a)**3,x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.25

$$\int \frac{x^{5/2}}{(a+bx^2)^3} dx = \frac{3bx^{7/2} - ax^{3/2}}{16(ab^3x^4 + 2a^2b^2x^2 + a^3b)} + \frac{3}{128ab} \left(\frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{1/4}b^{1/4} + 2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} + \frac{2\sqrt{2} \arctan\left(\frac{\sqrt{2}(\sqrt{2}a^{1/4}b^{1/4} - 2\sqrt{b}\sqrt{x})}{2\sqrt{\sqrt{a}\sqrt{b}}}\right)}{\sqrt{\sqrt{a}\sqrt{b}\sqrt{b}}} - \frac{\sqrt{2} \log(\sqrt{2}a^{1/4}b^{1/4}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{1/4}b^{3/4}} + \frac{\sqrt{2} \log(-\sqrt{2}a^{1/4}b^{1/4}\sqrt{x} + \sqrt{b}x + \sqrt{a})}{a^{1/4}b^{3/4}} \right)$$

input `integrate(x^(5/2)/(b*x^2+a)^3,x, algorithm="maxima")`

output `1/16*(3*b*x^(7/2) - a*x^(3/2))/(a*b^3*x^4 + 2*a^2*b^2*x^2 + a^3*b) + 3/128*(2*sqrt(2)*arctan(1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) + 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) + 2*sqrt(2)*arctan(-1/2*sqrt(2)*(sqrt(2)*a^(1/4)*b^(1/4) - 2*sqrt(b)*sqrt(x))/sqrt(sqrt(a)*sqrt(b)))/sqrt(sqrt(a)*sqrt(b))*sqrt(b) - sqrt(2)*log(sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4)) + sqrt(2)*log(-sqrt(2)*a^(1/4)*b^(1/4)*sqrt(x) + sqrt(b)*x + sqrt(a))/(a^(1/4)*b^(3/4))/(a*b)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.20

$$\int \frac{x^{5/2}}{(a+bx^2)^3} dx = \frac{3bx^{7/2} - ax^{3/2}}{16(bx^2+a)^2ab} + \frac{3\sqrt{2}(ab^3)^{3/4} \arctan\left(\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} + 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{64a^2b^4}$$

$$+ \frac{3\sqrt{2}(ab^3)^{3/4} \arctan\left(-\frac{\sqrt{2}\left(\sqrt{2}\left(\frac{a}{b}\right)^{1/4} - 2\sqrt{x}\right)}{2\left(\frac{a}{b}\right)^{1/4}}\right)}{64a^2b^4}$$

$$- \frac{3\sqrt{2}(ab^3)^{3/4} \log\left(\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{128a^2b^4}$$

$$+ \frac{3\sqrt{2}(ab^3)^{3/4} \log\left(-\sqrt{2}\sqrt{x}\left(\frac{a}{b}\right)^{1/4} + x + \sqrt{\frac{a}{b}}\right)}{128a^2b^4}$$

input `integrate(x^(5/2)/(b*x^2+a)^3,x, algorithm="giac")`output `1/16*(3*b*x^(7/2) - a*x^(3/2))/((b*x^2 + a)^2*a*b) + 3/64*sqrt(2)*(a*b^3)^(3/4)*arctan(1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) + 2*sqrt(x))/(a/b)^(1/4))/(a^2*b^4) + 3/64*sqrt(2)*(a*b^3)^(3/4)*arctan(-1/2*sqrt(2)*(sqrt(2)*(a/b)^(1/4) - 2*sqrt(x))/(a/b)^(1/4))/(a^2*b^4) - 3/128*sqrt(2)*(a*b^3)^(3/4)*log(sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^2*b^4) + 3/128*sqrt(2)*(a*b^3)^(3/4)*log(-sqrt(2)*sqrt(x)*(a/b)^(1/4) + x + sqrt(a/b))/(a^2*b^4)`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.48

$$\int \frac{x^{5/2}}{(a+bx^2)^3} dx = \frac{\frac{3x^{7/2}}{16a} - \frac{x^{3/2}}{16b}}{a^2 + 2abx^2 + b^2x^4} - \frac{3 \operatorname{atan}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{5/4}b^{7/4}} + \frac{3 \operatorname{atanh}\left(\frac{b^{1/4}\sqrt{x}}{(-a)^{1/4}}\right)}{32(-a)^{5/4}b^{7/4}}$$

input `int(x^(5/2)/(a + b*x^2)^3,x)`

output

```
((3*x^(7/2))/(16*a) - x^(3/2)/(16*b))/(a^2 + b^2*x^4 + 2*a*b*x^2) - (3*atan(b^(1/4)*x^(1/2))/(-a)^(1/4))/(32*(-a)^(5/4)*b^(7/4)) + (3*atanh(b^(1/4)*x^(1/2))/(-a)^(1/4))/(32*(-a)^(5/4)*b^(7/4))
```

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 478, normalized size of antiderivative = 2.70

$$\int \frac{x^{5/2}}{(a + bx^2)^3} dx = \text{Too large to display}$$

input

```
int(x^(5/2)/(b*x^2+a)^3,x)
```

output

```
( - 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a**2 - 12*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a*b*x**2 - 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) - 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b**2*x**4 + 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a**2 + 12*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*a*b*x**2 + 6*b**(1/4)*a**(3/4)*sqrt(2)*atan((b**(1/4)*a**(1/4)*sqrt(2) + 2*sqrt(x)*sqrt(b))/(b**(1/4)*a**(1/4)*sqrt(2)))*b**2*x**4 + 3*b**(1/4)*a**(3/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a**2 + 6*b**(1/4)*a**(3/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 + 3*b**(1/4)*a**(3/4)*sqrt(2)*log(-sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 - 3*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a**2 - 6*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*a*b*x**2 - 3*b**(1/4)*a**(3/4)*sqrt(2)*log(sqrt(x)*b**(1/4)*a**(1/4)*sqrt(2) + sqrt(a) + sqrt(b)*x)*b**2*x**4 - 8*sqrt(x)*a**2*b*x + 24*sqrt(x)*a*b**2*x**3)/(128*a**2*b**2*(a**2 + 2*a*b*x**2 + b**2*x**4))
```

3.133 $\int \frac{1}{\sqrt{a+bx}} dx$

Optimal result	906
Mathematica [A] (verified)	906
Rubi [A] (verified)	907
Maple [A] (verified)	908
Fricas [A] (verification not implemented)	908
Sympy [A] (verification not implemented)	909
Maxima [A] (verification not implemented)	909
Giac [A] (verification not implemented)	909
Mupad [B] (verification not implemented)	910
Reduce [B] (verification not implemented)	910

Optimal result

Integrand size = 9, antiderivative size = 14

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}}{b}$$

output `2*(b*x+a)^(1/2)/b`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}}{b}$$

input `Integrate[1/Sqrt[a + b*x],x]`

output `(2*Sqrt[a + b*x])/b`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{a+bx}} dx$$

↓ 17

$$\frac{2\sqrt{a+bx}}{b}$$

input `Int[1/Sqrt[a + b*x],x]`

output `(2*Sqrt[a + b*x])/b`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1))], x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

method	result	size
gosper	$\frac{2\sqrt{bx+a}}{b}$	13
derivativedivides	$\frac{2\sqrt{bx+a}}{b}$	13
default	$\frac{2\sqrt{bx+a}}{b}$	13
trager	$\frac{2\sqrt{bx+a}}{b}$	13
risch	$\frac{2\sqrt{bx+a}}{b}$	13
pseudoelliptic	$\frac{2\sqrt{bx+a}}{b}$	13
orering	$\frac{2\sqrt{bx+a}}{b}$	13

input `int(1/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`output `2*(b*x+a)^(1/2)/b`**Fricas [A] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{bx+a}}{b}$$

input `integrate(1/(b*x+a)^(1/2),x, algorithm="fricas")`output `2*sqrt(b*x + a)/b`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}}{b}$$

input `integrate(1/(b*x+a)**(1/2),x)`

output `2*sqrt(a + b*x)/b`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{bx+a}}{b}$$

input `integrate(1/(b*x+a)^(1/2),x, algorithm="maxima")`

output `2*sqrt(b*x + a)/b`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{bx+a}}{b}$$

input `integrate(1/(b*x+a)^(1/2),x, algorithm="giac")`

output `2*sqrt(b*x + a)/b`

Mupad [B] (verification not implemented)

Time = 0.01 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}}{b}$$

input `int(1/(a + b*x)^(1/2),x)`

output `(2*(a + b*x)^(1/2))/b`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sqrt{a+bx}} dx = \frac{2\sqrt{bx+a}}{b}$$

input `int(1/(b*x+a)^(1/2),x)`

output `(2*sqrt(a + b*x))/b`

3.134 $\int \frac{x}{\sqrt{a+bx}} dx$

Optimal result	911
Mathematica [A] (verified)	911
Rubi [A] (verified)	912
Maple [A] (verified)	913
Fricas [A] (verification not implemented)	913
Sympy [B] (verification not implemented)	914
Maxima [A] (verification not implemented)	914
Giac [A] (verification not implemented)	915
Mupad [B] (verification not implemented)	915
Reduce [B] (verification not implemented)	915

Optimal result

Integrand size = 11, antiderivative size = 27

$$\int \frac{x}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}(-a + \frac{1}{3}(a+bx))}{b^2}$$

output `2*(b*x+a)^(1/2)*(-2/3*a+1/3*b*x)/b^2`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{a+bx}} dx = \frac{2(-2a+bx)\sqrt{a+bx}}{3b^2}$$

input `Integrate[x/Sqrt[a + b*x],x]`

output `(2*(-2*a + b*x)*Sqrt[a + b*x])/(3*b^2)`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.19, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{a+bx}} dx$$

↓ 53

$$\int \left(\frac{\sqrt{a+bx}}{b} - \frac{a}{b\sqrt{a+bx}} \right) dx$$

↓ 2009

$$\frac{2(a+bx)^{3/2}}{3b^2} - \frac{2a\sqrt{a+bx}}{b^2}$$

input `Int[x/Sqrt[a + b*x],x]`

output `(-2*a*Sqrt[a + b*x])/b^2 + (2*(a + b*x)^(3/2))/(3*b^2)`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.78

method	result	size
gospers	$-\frac{2\sqrt{bx+a}(-bx+2a)}{3b^2}$	21
trager	$-\frac{2\sqrt{bx+a}(-bx+2a)}{3b^2}$	21
risch	$-\frac{2\sqrt{bx+a}(-bx+2a)}{3b^2}$	21
pseudoelliptic	$-\frac{2\sqrt{bx+a}(-bx+2a)}{3b^2}$	21
orering	$-\frac{2\sqrt{bx+a}(-bx+2a)}{3b^2}$	21
derivativedivides	$\frac{2(bx+a)^{\frac{3}{2}}}{3} - \frac{2\sqrt{bx+a}a}{b^2}$	26
default	$\frac{2(bx+a)^{\frac{3}{2}}}{3} - \frac{2\sqrt{bx+a}a}{b^2}$	26

input `int(x/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/3*(b*x+a)^(1/2)*(-b*x+2*a)/b^2`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \frac{x}{\sqrt{a+bx}} dx = \frac{2\sqrt{bx+a}(bx-2a)}{3b^2}$$

input `integrate(x/(b*x+a)^(1/2),x, algorithm="fricas")`

output `2/3*sqrt(b*x + a)*(b*x - 2*a)/b^2`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(22) = 44$.

Time = 0.64 (sec) , antiderivative size = 162, normalized size of antiderivative = 6.00

$$\int \frac{x}{\sqrt{a+bx}} dx = -\frac{4a^{\frac{7}{2}}\sqrt{1+\frac{bx}{a}}}{3a^2b^2+3ab^3x} + \frac{4a^{\frac{7}{2}}}{3a^2b^2+3ab^3x} - \frac{2a^{\frac{5}{2}}bx\sqrt{1+\frac{bx}{a}}}{3a^2b^2+3ab^3x} \\ + \frac{4a^{\frac{5}{2}}bx}{3a^2b^2+3ab^3x} + \frac{2a^{\frac{3}{2}}b^2x^2\sqrt{1+\frac{bx}{a}}}{3a^2b^2+3ab^3x}$$

input `integrate(x/(b*x+a)**(1/2),x)`

output `-4*a**(7/2)*sqrt(1 + b*x/a)/(3*a**2*b**2 + 3*a*b**3*x) + 4*a**(7/2)/(3*a**2*b**2 + 3*a*b**3*x) - 2*a**(5/2)*b*x*sqrt(1 + b*x/a)/(3*a**2*b**2 + 3*a*b**3*x) + 4*a**(5/2)*b*x/(3*a**2*b**2 + 3*a*b**3*x) + 2*a**(3/2)*b**2*x**2*sqrt(1 + b*x/a)/(3*a**2*b**2 + 3*a*b**3*x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{x}{\sqrt{a+bx}} dx = \frac{2(bx+a)^{\frac{3}{2}}}{3b^2} - \frac{2\sqrt{bx+aa}}{b^2}$$

input `integrate(x/(b*x+a)^(1/2),x, algorithm="maxima")`

output `2/3*(b*x + a)^(3/2)/b^2 - 2*sqrt(b*x + a)*a/b^2`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{x}{\sqrt{a+bx}} dx = \frac{2 \left((bx+a)^{\frac{3}{2}} - 3\sqrt{bx+aa} \right)}{3b^2}$$

input `integrate(x/(b*x+a)^(1/2),x, algorithm="giac")`

output `2/3*((b*x + a)^(3/2) - 3*sqrt(b*x + a)*a)/b^2`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.93

$$\int \frac{x}{\sqrt{a+bx}} dx = -\frac{6a\sqrt{a+bx} - 2(a+bx)^{3/2}}{3b^2}$$

input `int(x/(a + b*x)^(1/2),x)`

output `-(6*a*(a + b*x)^(1/2) - 2*(a + b*x)^(3/2))/(3*b^2)`

Reduce [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.67

$$\int \frac{x}{\sqrt{a+bx}} dx = \frac{2\sqrt{bx+a}(bx-2a)}{3b^2}$$

input `int(x/(b*x+a)^(1/2),x)`

output `(2*sqrt(a + b*x)*(- 2*a + b*x))/(3*b**2)`

3.135 $\int \frac{x^2}{\sqrt{a+bx}} dx$

Optimal result	916
Mathematica [A] (verified)	916
Rubi [A] (verified)	917
Maple [A] (verified)	918
Fricas [A] (verification not implemented)	918
Sympy [B] (verification not implemented)	919
Maxima [A] (verification not implemented)	920
Giac [A] (verification not implemented)	920
Mupad [B] (verification not implemented)	921
Reduce [B] (verification not implemented)	921

Optimal result

Integrand size = 13, antiderivative size = 39

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}(a^2 - \frac{2}{3}a(a+bx) + \frac{1}{5}(a+bx)^2)}{b^3}$$

output $2*(b*x+a)^{(1/2)}*(a^2-2/3*a*(b*x+a)+1/5*(b*x+a)^2)/b^3$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}(8a^2 - 4abx + 3b^2x^2)}{15b^3}$$

input `Integrate[x^2/Sqrt[a + b*x],x]`

output $(2*\text{Sqrt}[a + b*x]*(8*a^2 - 4*a*b*x + 3*b^2*x^2))/(15*b^3)$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.31, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\sqrt{a+bx}} dx$$

↓ 53

$$\int \left(\frac{a^2}{b^2\sqrt{a+bx}} - \frac{2a\sqrt{a+bx}}{b^2} + \frac{(a+bx)^{3/2}}{b^2} \right) dx$$

↓ 2009

$$\frac{2a^2\sqrt{a+bx}}{b^3} + \frac{2(a+bx)^{5/2}}{5b^3} - \frac{4a(a+bx)^{3/2}}{3b^3}$$

input `Int[x^2/Sqrt[a + b*x],x]`

output `(2*a^2*Sqrt[a + b*x])/b^3 - (4*a*(a + b*x)^(3/2))/(3*b^3) + (2*(a + b*x)^(5/2))/(5*b^3)`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

method	result	size
gospers	$\frac{2\sqrt{bx+a}(3b^2x^2-4bax+8a^2)}{15b^3}$	32
trager	$\frac{2\sqrt{bx+a}(3b^2x^2-4bax+8a^2)}{15b^3}$	32
risch	$\frac{2\sqrt{bx+a}(3b^2x^2-4bax+8a^2)}{15b^3}$	32
pseudoelliptic	$\frac{2\sqrt{bx+a}(3b^2x^2-4bax+8a^2)}{15b^3}$	32
orering	$\frac{2\sqrt{bx+a}(3b^2x^2-4bax+8a^2)}{15b^3}$	32
derivativedivides	$\frac{\frac{2(bx+a)^{\frac{5}{2}}}{5} - \frac{4a(bx+a)^{\frac{3}{2}}}{b^3} + 2\sqrt{bx+a}a^2}{b^3}$	37
default	$\frac{\frac{2(bx+a)^{\frac{5}{2}}}{5} - \frac{4a(bx+a)^{\frac{3}{2}}}{b^3} + 2\sqrt{bx+a}a^2}{b^3}$	37

input `int(x^2/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`output `2/15*(b*x+a)^(1/2)*(3*b^2*x^2-4*a*b*x+8*a^2)/b^3`**Fricas [A] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{2(3b^2x^2 - 4abx + 8a^2)\sqrt{bx+a}}{15b^3}$$

input `integrate(x^2/(b*x+a)^(1/2),x, algorithm="fricas")`output `2/15*(3*b^2*x^2 - 4*a*b*x + 8*a^2)*sqrt(b*x + a)/b^3`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 600 vs. $2(34) = 68$.

Time = 0.96 (sec) , antiderivative size = 600, normalized size of antiderivative = 15.38

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{16a^{\frac{21}{2}} \sqrt{1 + \frac{bx}{a}}}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} - \frac{16a^{\frac{21}{2}}}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} + \frac{40a^{\frac{19}{2}} bx \sqrt{1 + \frac{bx}{a}}}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} - \frac{48a^{\frac{19}{2}} bx}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} + \frac{30a^{\frac{17}{2}} b^2x^2 \sqrt{1 + \frac{bx}{a}}}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} - \frac{48a^{\frac{17}{2}} b^2x^2}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} + \frac{10a^{\frac{15}{2}} b^3x^3 \sqrt{1 + \frac{bx}{a}}}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} - \frac{16a^{\frac{15}{2}} b^3x^3}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} + \frac{10a^{\frac{13}{2}} b^4x^4 \sqrt{1 + \frac{bx}{a}}}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} - \frac{10a^{\frac{13}{2}} b^4x^4}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} + \frac{6a^{\frac{11}{2}} b^5x^5 \sqrt{1 + \frac{bx}{a}}}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3} - \frac{6a^{\frac{11}{2}} b^5x^5}{15a^8b^3 + 45a^7b^4x + 45a^6b^5x^2 + 15a^5b^6x^3}$$

input `integrate(x**2/(b*x+a)**(1/2),x)`

output

```
16*a**(21/2)*sqrt(1 + b*x/a)/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) - 16*a**(21/2)/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) + 40*a**(19/2)*b*x*sqrt(1 + b*x/a)/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) - 48*a**(19/2)*b*x/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) + 30*a**(17/2)*b**2*x**2*sqrt(1 + b*x/a)/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) - 48*a**(17/2)*b**2*x**2/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) + 10*a**(15/2)*b**3*x**3*sqrt(1 + b*x/a)/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) - 16*a**(15/2)*b**3*x**3/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) + 10*a**(13/2)*b**4*x**4*sqrt(1 + b*x/a)/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3) + 6*a**(11/2)*b**5*x**5*sqrt(1 + b*x/a)/(15*a**8*b**3 + 45*a**7*b**4*x + 45*a**6*b**5*x**2 + 15*a**5*b**6*x**3)
```

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{2(bx+a)^{\frac{5}{2}}}{5b^3} - \frac{4(bx+a)^{\frac{3}{2}}a}{3b^3} + \frac{2\sqrt{bx+aa^2}}{b^3}$$

input

```
integrate(x^2/(b*x+a)^(1/2),x, algorithm="maxima")
```

output

```
2/5*(b*x + a)^(5/2)/b^3 - 4/3*(b*x + a)^(3/2)*a/b^3 + 2*sqrt(b*x + a)*a^2/b^3
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.95

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{2\left(3(bx+a)^{\frac{5}{2}} - 10(bx+a)^{\frac{3}{2}}a + 15\sqrt{bx+aa^2}\right)}{15b^3}$$

input

```
integrate(x^2/(b*x+a)^(1/2),x, algorithm="giac")
```

output $2/15*(3*(b*x + a)^{(5/2)} - 10*(b*x + a)^{(3/2)}*a + 15*\text{sqrt}(b*x + a)*a^2)/b^3$

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.95

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{6(a+bx)^{5/2} - 20a(a+bx)^{3/2} + 30a^2\sqrt{a+bx}}{15b^3}$$

input $\text{int}(x^2/(a + b*x)^{(1/2)}, x)$

output $(6*(a + b*x)^{(5/2)} - 20*a*(a + b*x)^{(3/2)} + 30*a^2*(a + b*x)^{(1/2)})/(15*b^3)$

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.77

$$\int \frac{x^2}{\sqrt{a+bx}} dx = \frac{2\sqrt{bx+a}(3b^2x^2 - 4abx + 8a^2)}{15b^3}$$

input $\text{int}(x^2/(b*x+a)^{(1/2)}, x)$

output $(2*\text{sqrt}(a + b*x)*(8*a**2 - 4*a*b*x + 3*b**2*x**2))/(15*b**3)$

3.136 $\int \frac{1}{\sqrt{(a+bx)^3}} dx$

Optimal result	922
Mathematica [A] (verified)	922
Rubi [A] (verified)	923
Maple [A] (verified)	924
Fricas [B] (verification not implemented)	924
Sympy [B] (verification not implemented)	925
Maxima [A] (verification not implemented)	925
Giac [A] (verification not implemented)	925
Mupad [B] (verification not implemented)	926
Reduce [B] (verification not implemented)	926

Optimal result

Integrand size = 11, antiderivative size = 14

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = -\frac{2}{b\sqrt{a+bx}}$$

output `-2/b/(b*x+a)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = -\frac{2(a+bx)}{b\sqrt{(a+bx)^3}}$$

input `Integrate[1/Sqrt[(a + b*x)^3],x]`

output `(-2*(a + b*x))/(b*Sqrt[(a + b*x)^3])`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.50, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {239, 20, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt{(a+bx)^3}} dx \\ & \quad \downarrow \text{239} \\ & \frac{\int \frac{1}{\sqrt{(a+bx)^3}} d(a+bx)}{b} \\ & \quad \downarrow \text{20} \\ & \frac{(a+bx)^{3/2} \int \frac{1}{(a+bx)^{3/2}} d(a+bx)}{b\sqrt{(a+bx)^3}} \\ & \quad \downarrow \text{15} \\ & -\frac{2(a+bx)}{b\sqrt{(a+bx)^3}} \end{aligned}$$

input `Int[1/Sqrt[(a + b*x)^3],x]`

output `(-2*(a + b*x))/(b*Sqrt[(a + b*x)^3])`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 20 `Int[((a_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a*x^n)^p/x^(n*p) Int[x^(n*p), x], x] /; FreeQ[{a, n, p}, x] && !IntegerQ[p]`

rule 239 `Int[((a_.) + (b_.)*(v_)^(n_))^(p_), x_Symbol] := Simp[1/Coefficient[v, x, 1] Subst[Int[(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, n, p}, x] && LinearQ[v, x] && NeQ[v, x]`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

method	result	size
gospers	$-\frac{2(bx+a)}{b\sqrt{(bx+a)^3}}$	20
default	$-\frac{2(bx+a)}{b\sqrt{(bx+a)^3}}$	20
orering	$-\frac{2(bx+a)}{b\sqrt{(bx+a)^3}}$	20
trager	$-\frac{2\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}}{(bx+a)^2b}$	42

input `int(1/((b*x+a)^3)^(1/2),x,method=_RETURNVERBOSE)`

output `-2*(b*x+a)/b/((b*x+a)^3)^(1/2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. $2(12) = 24$.

Time = 0.08 (sec) , antiderivative size = 53, normalized size of antiderivative = 3.79

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = -\frac{2\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}}{b^3x^2+2ab^2x+a^2b}$$

input `integrate(1/((b*x+a)^3)^(1/2),x, algorithm="fricas")`

output `-2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)/(b^3*x^2 + 2*a*b^2*x + a^2*b)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(12) = 24$.

Time = 0.77 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.93

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = \begin{cases} -\frac{2(\frac{a}{b}+x)}{\sqrt{(a+bx)^3}} & \text{for } b \neq 0 \\ \frac{x}{\sqrt{a^3}} & \text{otherwise} \end{cases}$$

input `integrate(1/((b*x+a)**3)**(1/2),x)`

output `Piecewise((-2*(a/b + x)/sqrt((a + b*x)**3), Ne(b, 0)), (x/sqrt(a**3), True))`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = -\frac{2}{\sqrt{bx+ab}}$$

input `integrate(1/((b*x+a)^3)^(1/2),x, algorithm="maxima")`

output `-2/(sqrt(b*x + a)*b)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = -\frac{2}{\sqrt{bx+ab}}$$

input `integrate(1/((b*x+a)^3)^(1/2),x, algorithm="giac")`

output `-2/(sqrt(b*x + a)*b)`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = -\frac{2\sqrt{(a+bx)^3}}{b(a+bx)^2}$$

input `int(1/((a + b*x)^3)^(1/2),x)`

output `-(2*((a + b*x)^3)^(1/2))/(b*(a + b*x)^2)`

Reduce [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

$$\int \frac{1}{\sqrt{(a+bx)^3}} dx = -\frac{2}{\sqrt{bx+a}b}$$

input `int(1/((b*x+a)^3)^(1/2),x)`

output `(- 2)/(sqrt(a + b*x)*b)`

3.137 $\int \frac{x}{\sqrt{(a+bx)^3}} dx$

Optimal result	927
Mathematica [A] (verified)	927
Rubi [B] (verified)	928
Maple [A] (verified)	929
Fricas [B] (verification not implemented)	930
Sympy [F]	930
Maxima [A] (verification not implemented)	930
Giac [A] (verification not implemented)	931
Mupad [B] (verification not implemented)	931
Reduce [B] (verification not implemented)	931

Optimal result

Integrand size = 13, antiderivative size = 21

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \frac{2(2a+bx)}{b^2\sqrt{a+bx}}$$

output

```
2*(b*x+2*a)/b^2/(b*x+a)^(1/2)
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.33

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \frac{2(a+bx)(2a+bx)}{b^2\sqrt{(a+bx)^3}}$$

input

```
Integrate[x/Sqrt[(a + b*x)^3],x]
```

output

```
(2*(a + b*x)*(2*a + b*x))/(b^2*Sqrt[(a + b*x)^3])
```


Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs. $2(21) = 42$.

Time = 0.34 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.43, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2008, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{\sqrt{(a+bx)^3}} dx \\
 & \quad \downarrow \text{2008} \\
 & \frac{(a+bx)^{3/2} \int \frac{x}{(a+bx)^{3/2}} dx}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{53} \\
 & \frac{(a+bx)^{3/2} \int \left(\frac{1}{b\sqrt{a+bx}} - \frac{a}{b(a+bx)^{3/2}} \right) dx}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{2009} \\
 & \frac{(a+bx)^{3/2} \left(\frac{2a}{b^2\sqrt{a+bx}} + \frac{2\sqrt{a+bx}}{b^2} \right)}{\sqrt{(a+bx)^3}}
 \end{aligned}$$

input `Int[x/Sqrt[(a + b*x)^3],x]`

output `((a + b*x)^(3/2)*((2*a)/(b^2*Sqrt[a + b*x]) + (2*Sqrt[a + b*x])/b^2))/Sqrt[(a + b*x)^3]`

Definitions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

rule 2008 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Expon[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p), x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.29

method	result	size
gospers	$\frac{2(bx+a)(bx+2a)}{b^2\sqrt{(bx+a)^3}}$	27
default	$\frac{2(bx+a)(bx+2a)}{b^2\sqrt{(bx+a)^3}}$	27
orering	$\frac{2(bx+a)(bx+2a)}{b^2\sqrt{(bx+a)^3}}$	27
risch	$\frac{2(bx+a)^2}{b^2\sqrt{(bx+a)^3}} + \frac{2a(bx+a)}{b^2\sqrt{(bx+a)^3}}$	43
trager	$\frac{2(bx+2a)\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}}{(bx+a)^2b^2}$	49

input `int(x/((b*x+a)^3)^(1/2),x,method=_RETURNVERBOSE)`

output `2*(b*x+a)*(b*x+2*a)/b^2/((b*x+a)^3)^(1/2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 62 vs. $2(19) = 38$.

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 2.95

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \frac{2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}(bx + 2a)}{b^4x^2 + 2ab^3x + a^2b^2}$$

input `integrate(x/((b*x+a)^3)^(1/2),x, algorithm="fricas")`

output `2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*(b*x + 2*a)/(b^4*x^2 + 2*a*b^3*x + a^2*b^2)`

Sympy [F]

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \int \frac{x}{\sqrt{(a+bx)^3}} dx$$

input `integrate(x/((b*x+a)**3)**(1/2),x)`

output `Integral(x/sqrt((a + b*x)**3), x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.43

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \frac{2(b^2x^2 + 3abx + 2a^2)}{(bx + a)^{\frac{3}{2}}b^2}$$

input `integrate(x/((b*x+a)^3)^(1/2),x, algorithm="maxima")`

output `2*(b^2*x^2 + 3*a*b*x + 2*a^2)/((b*x + a)^(3/2)*b^2)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.38

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \frac{2 \left(\frac{\sqrt{bx+a}}{b} + \frac{a}{\sqrt{bx+ab}} \right)}{b}$$

input `integrate(x/((b*x+a)^3)^(1/2),x, algorithm="giac")`output `2*(sqrt(b*x + a)/b + a/(sqrt(b*x + a)*b))/b`**Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.33

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \frac{2(2a+bx)\sqrt{(a+bx)^3}}{b^2(a+bx)^2}$$

input `int(x/((a + b*x)^3)^(1/2),x)`output `(2*(2*a + b*x)*((a + b*x)^3)^(1/2))/(b^2*(a + b*x)^2)`**Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

$$\int \frac{x}{\sqrt{(a+bx)^3}} dx = \frac{2bx+4a}{\sqrt{bx+ab^2}}$$

input `int(x/((b*x+a)^3)^(1/2),x)`output `(2*(2*a + b*x))/(sqrt(a + b*x)*b**2)`

3.138 $\int \frac{x^2}{\sqrt{(a+bx)^3}} dx$

Optimal result	932
Mathematica [A] (verified)	932
Rubi [A] (verified)	933
Maple [A] (verified)	934
Fricas [B] (verification not implemented)	935
Sympy [F]	935
Maxima [A] (verification not implemented)	935
Giac [A] (verification not implemented)	936
Mupad [B] (verification not implemented)	936
Reduce [B] (verification not implemented)	936

Optimal result

Integrand size = 15, antiderivative size = 39

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = \frac{2(-a^2 - 2a(a+bx) + \frac{1}{3}(a+bx)^2)}{b^3\sqrt{a+bx}}$$

output `2*(-a^2-2*a*(b*x+a)+1/3*(b*x+a)^2)/b^3/(b*x+a)^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = \frac{2(a+bx)(-8a^2 - 4abx + b^2x^2)}{3b^3\sqrt{(a+bx)^3}}$$

input `Integrate[x^2/Sqrt[(a + b*x)^3], x]`

output `(2*(a + b*x)*(-8*a^2 - 4*a*b*x + b^2*x^2))/(3*b^3*Sqrt[(a + b*x)^3])`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.79, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2008, 53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx$$

$$\downarrow \text{2008}$$

$$\frac{(a+bx)^{3/2} \int \frac{x^2}{(a+bx)^{3/2}} dx}{\sqrt{(a+bx)^3}}$$

$$\downarrow \text{53}$$

$$\frac{(a+bx)^{3/2} \int \left(\frac{a^2}{b^2(a+bx)^{3/2}} - \frac{2a}{b^2\sqrt{a+bx}} + \frac{\sqrt{a+bx}}{b^2} \right) dx}{\sqrt{(a+bx)^3}}$$

$$\downarrow \text{2009}$$

$$\frac{(a+bx)^{3/2} \left(-\frac{2a^2}{b^3\sqrt{a+bx}} - \frac{4a\sqrt{a+bx}}{b^3} + \frac{2(a+bx)^{3/2}}{3b^3} \right)}{\sqrt{(a+bx)^3}}$$

input `Int[x^2/Sqrt[(a + b*x)^3],x]`

output `((a + b*x)^(3/2)*((-2*a^2)/(b^3*Sqrt[a + b*x]) - (4*a*Sqrt[a + b*x])/b^3 + (2*(a + b*x)^(3/2))/(3*b^3)))/Sqrt[(a + b*x)^3]`

Defintions of rubi rules used

rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n},
x] && IGtQ[m, 0] && ( !IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0])
|| LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

rule 2008

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Exp
on[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p),
x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x
] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

method	result	size
gosper	$-\frac{2(bx+a)(-b^2x^2+4bax+8a^2)}{3b^3\sqrt{(bx+a)^3}}$	39
default	$-\frac{2(bx+a)(-b^2x^2+4bax+8a^2)}{3b^3\sqrt{(bx+a)^3}}$	39
orering	$-\frac{2(bx+a)(-b^2x^2+4bax+8a^2)}{3b^3\sqrt{(bx+a)^3}}$	39
risch	$-\frac{2(-bx+5a)(bx+a)^2}{3b^3\sqrt{(bx+a)^3}} - \frac{2a^2(bx+a)}{b^3\sqrt{(bx+a)^3}}$	53
trager	$-\frac{2(-b^2x^2+4bax+8a^2)\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}}{3(bx+a)^2b^3}$	61

input

```
int(x^2/((b*x+a)^3)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-2/3*(b*x+a)*(-b^2*x^2+4*a*b*x+8*a^2)/b^3/((b*x+a)^3)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. $2(33) = 66$.

Time = 0.09 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.87

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = \frac{2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}(b^2x^2 - 4abx - 8a^2)}{3(b^5x^2 + 2ab^4x + a^2b^3)}$$

input `integrate(x^2/((b*x+a)^3)^(1/2),x, algorithm="fricas")`

output `2/3*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*(b^2*x^2 - 4*a*b*x - 8*a^2)/(b^5*x^2 + 2*a*b^4*x + a^2*b^3)`

Sympy [F]

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = \int \frac{x^2}{\sqrt{(a+bx)^3}} dx$$

input `integrate(x**2/((b*x+a)**3)**(1/2),x)`

output `Integral(x**2/sqrt((a + b*x)**3), x)`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = \frac{2(b^3x^3 - 3ab^2x^2 - 12a^2bx - 8a^3)}{3(bx+a)^{\frac{3}{2}}b^3}$$

input `integrate(x^2/((b*x+a)^3)^(1/2),x, algorithm="maxima")`

output `2/3*(b^3*x^3 - 3*a*b^2*x^2 - 12*a^2*b*x - 8*a^3)/((b*x + a)^(3/2)*b^3)`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.31

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = -\frac{2 \left(\frac{3a^2}{\sqrt{bx+ab}} - \frac{(bx+a)^{\frac{3}{2}} b^2 - 6\sqrt{bx+ab} b^2}{b^3} \right)}{3b^2}$$

input `integrate(x^2/((b*x+a)^3)^(1/2),x, algorithm="giac")`output `-2/3*(3*a^2/(sqrt(b*x + a)*b) - ((b*x + a)^(3/2)*b^2 - 6*sqrt(b*x + a)*a*b^2)/b^3)/b^2`**Mupad [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.03

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = -\frac{2\sqrt{(a+bx)^3}(8a^2+4abx-b^2x^2)}{3b^3(a+bx)^2}$$

input `int(x^2/((a + b*x)^3)^(1/2),x)`output `-(2*((a + b*x)^3)^(1/2)*(8*a^2 - b^2*x^2 + 4*a*b*x))/(3*b^3*(a + b*x)^2)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int \frac{x^2}{\sqrt{(a+bx)^3}} dx = \frac{\frac{2}{3}b^2x^2 - \frac{8}{3}abx - \frac{16}{3}a^2}{\sqrt{bx+a}b^3}$$

input `int(x^2/((b*x+a)^3)^(1/2),x)`output `(2*(- 8*a**2 - 4*a*b*x + b**2*x**2))/(3*sqrt(a + b*x)*b**3)`

3.139 $\int \frac{1}{x\sqrt{a+bx}} dx$

Optimal result	937
Mathematica [A] (verified)	937
Rubi [A] (verified)	938
Maple [A] (verified)	939
Fricas [A] (verification not implemented)	939
Sympy [A] (verification not implemented)	940
Maxima [A] (verification not implemented)	940
Giac [A] (verification not implemented)	940
Mupad [B] (verification not implemented)	941
Reduce [B] (verification not implemented)	941

Optimal result

Integrand size = 13, antiderivative size = 42

$$\int \frac{1}{x\sqrt{a+bx}} dx = \frac{\log\left(\frac{-\sqrt{a}+\sqrt{a+bx}}{\sqrt{a}+\sqrt{a+bx}}\right)}{\sqrt{a}}$$

output $\ln((-a^{(1/2)}+(b*x+a)^{(1/2)})/(a^{(1/2)}+(b*x+a)^{(1/2)))/a^{(1/2)}$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.55

$$\int \frac{1}{x\sqrt{a+bx}} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{\sqrt{a}}$$

input `Integrate[1/(x*Sqrt[a + b*x]),x]`

output $(-2*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*x]/\operatorname{Sqrt}[a]])/\operatorname{Sqrt}[a]$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.55, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt{a+bx}} dx$$

$$\downarrow \text{73}$$

$$\frac{2 \int \frac{1}{\frac{a+bx}{b} - \frac{a}{b}} d\sqrt{a+bx}}{b}$$

$$\downarrow \text{221}$$

$$\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{\sqrt{a}}$$

input `Int[1/(x*Sqrt[a + b*x]),x]`

output `(-2*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/Sqrt[a]`

Defintions of rubi rules used

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.43

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{\sqrt{a}}$	18
default	$-\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{\sqrt{a}}$	18
pseudoelliptic	$-\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{\sqrt{a}}$	18

input `int(1/x/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output `-2/a^(1/2)*arctanh((b*x+a)^(1/2)/a^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.26

$$\int \frac{1}{x\sqrt{a+bx}} dx = \left[\frac{\log\left(\frac{bx-2\sqrt{bx+a}\sqrt{a}+2a}{x}\right)}{\sqrt{a}}, \frac{2\sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right)}{a} \right]$$

input `integrate(1/x/(b*x+a)^(1/2),x, algorithm="fricas")`

output `[log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x)/sqrt(a), 2*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a))/a]`

Sympy [A] (verification not implemented)

Time = 0.58 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.57

$$\int \frac{1}{x\sqrt{a+bx}} dx = -\frac{2 \operatorname{asinh}\left(\frac{\sqrt{a}}{\sqrt{b}\sqrt{x}}\right)}{\sqrt{a}}$$

input `integrate(1/x/(b*x+a)**(1/2),x)`output `-2*asinh(sqrt(a)/(sqrt(b)*sqrt(x)))/sqrt(a)`**Maxima [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.76

$$\int \frac{1}{x\sqrt{a+bx}} dx = \frac{\log\left(\frac{\sqrt{bx+a}-\sqrt{a}}{\sqrt{bx+a}+\sqrt{a}}\right)}{\sqrt{a}}$$

input `integrate(1/x/(b*x+a)^(1/2),x, algorithm="maxima")`output `log((sqrt(b*x + a) - sqrt(a))/(sqrt(b*x + a) + sqrt(a)))/sqrt(a)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.50

$$\int \frac{1}{x\sqrt{a+bx}} dx = \frac{2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}}$$

input `integrate(1/x/(b*x+a)^(1/2),x, algorithm="giac")`output `2*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a)`

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.40

$$\int \frac{1}{x\sqrt{a+bx}} dx = -\frac{2 \operatorname{atanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{\sqrt{a}}$$

input `int(1/(x*(a + b*x)^(1/2)),x)`output `-(2*atanh((a + b*x)^(1/2)/a^(1/2)))/a^(1/2)`**Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.74

$$\int \frac{1}{x\sqrt{a+bx}} dx = \frac{\sqrt{a} (\log(\sqrt{bx+a} - \sqrt{a}) - \log(\sqrt{bx+a} + \sqrt{a}))}{a}$$

input `int(1/x/(b*x+a)^(1/2),x)`output `(sqrt(a)*(log(sqrt(a + b*x) - sqrt(a)) - log(sqrt(a + b*x) + sqrt(a))))/a`

3.140 $\int \frac{\sqrt{a+bx}}{x} dx$

Optimal result	942
Mathematica [A] (verified)	942
Rubi [A] (verified)	943
Maple [A] (verified)	944
Fricas [A] (verification not implemented)	944
Sympy [A] (verification not implemented)	945
Maxima [A] (verification not implemented)	945
Giac [A] (verification not implemented)	946
Mupad [B] (verification not implemented)	946
Reduce [B] (verification not implemented)	946

Optimal result

Integrand size = 13, antiderivative size = 54

$$\int \frac{\sqrt{a+bx}}{x} dx = 2\sqrt{a+bx} + \sqrt{a} \log \left(\frac{-\sqrt{a} + \sqrt{a+bx}}{\sqrt{a} + \sqrt{a+bx}} \right)$$

output

```
2*(b*x+a)^(1/2)+a^(1/2)*ln((-a^(1/2)+(b*x+a)^(1/2))/(a^(1/2)+(b*x+a)^(1/2)))
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.65

$$\int \frac{\sqrt{a+bx}}{x} dx = 2\sqrt{a+bx} - 2\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a+bx}}{\sqrt{a}} \right)$$

input

```
Integrate[Sqrt[a + b*x]/x,x]
```

output

```
2*Sqrt[a + b*x] - 2*Sqrt[a]*ArcTanh[Sqrt[a + b*x]/Sqrt[a]]
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.65, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx}}{x} dx$$

$$\downarrow 60$$

$$a \int \frac{1}{x\sqrt{a+bx}} dx + 2\sqrt{a+bx}$$

$$\downarrow 73$$

$$\frac{2a \int \frac{1}{\frac{a+bx}{b} - \frac{a}{b}} d\sqrt{a+bx}}{b} + 2\sqrt{a+bx}$$

$$\downarrow 221$$

$$2\sqrt{a+bx} - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)$$

input `Int[Sqrt[a + b*x]/x,x]`

output `2*Sqrt[a + b*x] - 2*Sqrt[a]*ArcTanh[Sqrt[a + b*x]/Sqrt[a]]`

Defintions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(
b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```


rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.52

method	result	size
derivativedivides	$2\sqrt{bx+a} - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)$	28
default	$2\sqrt{bx+a} - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)$	28
pseudoelliptic	$2\sqrt{bx+a} - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)$	28

input

```
int((b*x+a)^(1/2)/x,x,method=_RETURNVERBOSE)
```

output

```
2*(b*x+a)^(1/2)-2*a^(1/2)*arctanh((b*x+a)^(1/2)/a^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.30

$$\int \frac{\sqrt{a+bx}}{x} dx = \left[\sqrt{a} \log\left(\frac{bx - 2\sqrt{bx+a}\sqrt{a} + 2a}{x}\right) + 2\sqrt{bx+a}, 2\sqrt{-a} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + 2\sqrt{bx+a} \right]$$

input

```
integrate((b*x+a)^(1/2)/x,x, algorithm="fricas")
```

output `[sqrt(a)*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*sqrt(b*x + a), 2*sqrt(-a)*arctan(sqrt(-a)/sqrt(b*x + a)) + 2*sqrt(b*x + a)]`

Sympy [A] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.26

$$\int \frac{\sqrt{a+bx}}{x} dx = -2\sqrt{a} \operatorname{asinh}\left(\frac{\sqrt{a}}{\sqrt{b}\sqrt{x}}\right) + \frac{2a}{\sqrt{b}\sqrt{x}\sqrt{\frac{a}{bx}+1}} + \frac{2\sqrt{b}\sqrt{x}}{\sqrt{\frac{a}{bx}+1}}$$

input `integrate((b*x+a)**(1/2)/x,x)`

output `-2*sqrt(a)*asinh(sqrt(a)/(sqrt(b)*sqrt(x))) + 2*a/(sqrt(b)*sqrt(x)*sqrt(a/(b*x) + 1)) + 2*sqrt(b)*sqrt(x)/sqrt(a/(b*x) + 1)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.78

$$\int \frac{\sqrt{a+bx}}{x} dx = \sqrt{a} \log\left(\frac{\sqrt{bx+a}-\sqrt{a}}{\sqrt{bx+a}+\sqrt{a}}\right) + 2\sqrt{bx+a}$$

input `integrate((b*x+a)^(1/2)/x,x, algorithm="maxima")`

output `sqrt(a)*log((sqrt(b*x + a) - sqrt(a))/(sqrt(b*x + a) + sqrt(a))) + 2*sqrt(b*x + a)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{a+bx}}{x} dx = \frac{2a \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + 2\sqrt{bx+a}$$

input `integrate((b*x+a)^(1/2)/x,x, algorithm="giac")`output `2*a*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + 2*sqrt(b*x + a)`**Mupad [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.50

$$\int \frac{\sqrt{a+bx}}{x} dx = 2\sqrt{a+bx} - 2\sqrt{a} \operatorname{atanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)$$

input `int((a + b*x)^(1/2)/x,x)`output `2*(a + b*x)^(1/2) - 2*a^(1/2)*atanh((a + b*x)^(1/2)/a^(1/2))`**Reduce [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.70

$$\int \frac{\sqrt{a+bx}}{x} dx = 2\sqrt{bx+a} + \sqrt{a} \log(\sqrt{bx+a} - \sqrt{a}) - \sqrt{a} \log(\sqrt{bx+a} + \sqrt{a})$$

input `int((b*x+a)^(1/2)/x,x)`output `2*sqrt(a + b*x) + sqrt(a)*log(sqrt(a + b*x) - sqrt(a)) - sqrt(a)*log(sqrt(a + b*x) + sqrt(a))`

3.141 $\int \frac{\sqrt{a+bx}}{x^2} dx$

Optimal result	947
Mathematica [A] (verified)	947
Rubi [A] (verified)	948
Maple [A] (verified)	949
Fricas [A] (verification not implemented)	950
Sympy [A] (verification not implemented)	950
Maxima [A] (verification not implemented)	951
Giac [A] (verification not implemented)	951
Mupad [B] (verification not implemented)	951
Reduce [B] (verification not implemented)	952

Optimal result

Integrand size = 13, antiderivative size = 61

$$\int \frac{\sqrt{a+bx}}{x^2} dx = -\frac{\sqrt{a+bx}}{x} + \frac{b \log\left(\frac{-\sqrt{a}+\sqrt{a+bx}}{\sqrt{a}+\sqrt{a+bx}}\right)}{2\sqrt{a}}$$

output

```
-(b*x+a)^(1/2)/x+1/2*b*ln((-a^(1/2)+(b*x+a)^(1/2))/(a^(1/2)+(b*x+a)^(1/2)))/a^(1/2)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.64

$$\int \frac{\sqrt{a+bx}}{x^2} dx = -\frac{\sqrt{a+bx}}{x} - \frac{b \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{\sqrt{a}}$$

input

```
Integrate[Sqrt[a + b*x]/x^2,x]
```

output

```
-(Sqrt[a + b*x]/x) - (b*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/Sqrt[a]
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.64, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {51, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{a+bx}}{x^2} dx$$

↓ 51

$$\frac{1}{2}b \int \frac{1}{x\sqrt{a+bx}} dx - \frac{\sqrt{a+bx}}{x}$$

↓ 73

$$\int \frac{1}{\frac{a+bx}{b} - \frac{a}{b}} d\sqrt{a+bx} - \frac{\sqrt{a+bx}}{x}$$

↓ 221

$$-\frac{\text{barctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\sqrt{a+bx}}{x}$$

input `Int[Sqrt[a + b*x]/x^2,x]`

output `-(Sqrt[a + b*x]/x) - (b*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/Sqrt[a]`

Defintions of rubi rules used

rule 51

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]
```

rule 73 `Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.52

method	result	size
risch	$-\frac{\sqrt{bx+a}}{x} - \frac{b \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{\sqrt{a}}$	32
pseudoelliptic	$b\left(-\frac{\sqrt{bx+a}}{bx} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{\sqrt{a}}\right)$	36
derivativedivides	$2b\left(-\frac{\sqrt{bx+a}}{2bx} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{2\sqrt{a}}\right)$	37
default	$2b\left(-\frac{\sqrt{bx+a}}{2bx} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{2\sqrt{a}}\right)$	37

input `int((b*x+a)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

output `-(b*x+a)^(1/2)/x-b/a^(1/2)*arctanh((b*x+a)^(1/2)/a^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.48

$$\int \frac{\sqrt{a+bx}}{x^2} dx = \left[\frac{\sqrt{abx} \log\left(\frac{bx-2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2\sqrt{bx+aa}}{2ax}, \frac{\sqrt{-abx} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) - \sqrt{bx+aa}}{ax} \right]$$

input `integrate((b*x+a)^(1/2)/x^2,x, algorithm="fricas")`output `[1/2*(sqrt(a)*b*x*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*sqrt(b*x + a)*a)/(a*x), (sqrt(-a)*b*x*arctan(sqrt(-a)/sqrt(b*x + a)) - sqrt(b*x + a)*a)/(a*x)]`**Sympy [A] (verification not implemented)**

Time = 0.98 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.72

$$\int \frac{\sqrt{a+bx}}{x^2} dx = -\frac{\sqrt{b}\sqrt{\frac{a}{bx}+1}}{\sqrt{x}} - \frac{b \operatorname{asinh}\left(\frac{\sqrt{a}}{\sqrt{b}\sqrt{x}}\right)}{\sqrt{a}}$$

input `integrate((b*x+a)**(1/2)/x**2,x)`output `-sqrt(b)*sqrt(a/(b*x) + 1)/sqrt(x) - b*asinh(sqrt(a)/(sqrt(b)*sqrt(x)))/sqrt(a)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{a+bx}}{x^2} dx = \frac{b \log\left(\frac{\sqrt{bx+a}-\sqrt{a}}{\sqrt{bx+a}+\sqrt{a}}\right)}{2\sqrt{a}} - \frac{\sqrt{bx+a}}{x}$$

input `integrate((b*x+a)^(1/2)/x^2,x, algorithm="maxima")`output `1/2*b*log((sqrt(b*x + a) - sqrt(a))/(sqrt(b*x + a) + sqrt(a)))/sqrt(a) - sqrt(b*x + a)/x`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.62

$$\int \frac{\sqrt{a+bx}}{x^2} dx = b \left(\frac{\arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} - \frac{\sqrt{bx+a}}{bx} \right)$$

input `integrate((b*x+a)^(1/2)/x^2,x, algorithm="giac")`output `b*(arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) - sqrt(b*x + a)/(b*x))`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.51

$$\int \frac{\sqrt{a+bx}}{x^2} dx = -\frac{\sqrt{a+bx}}{x} - \frac{b \operatorname{atanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{\sqrt{a}}$$

input `int((a + b*x)^(1/2)/x^2,x)`output `-(a + b*x)^(1/2)/x - (b*atanh((a + b*x)^(1/2)/a^(1/2)))/a^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.84

$$\int \frac{\sqrt{a+bx}}{x^2} dx$$

$$= \frac{-2\sqrt{bx+a}a + \sqrt{a}\log(\sqrt{bx+a} - \sqrt{a})bx - \sqrt{a}\log(\sqrt{bx+a} + \sqrt{a})bx}{2ax}$$

input `int((b*x+a)^(1/2)/x^2,x)`output `(- 2*sqrt(a + b*x)*a + sqrt(a)*log(sqrt(a + b*x) - sqrt(a))*b*x - sqrt(a)
*log(sqrt(a + b*x) + sqrt(a))*b*x)/(2*a*x)`

3.142 $\int \frac{\sqrt{a+bx}}{x^3} dx$

Optimal result	953
Mathematica [A] (verified)	953
Rubi [A] (verified)	954
Maple [A] (verified)	955
Fricas [A] (verification not implemented)	956
Sympy [A] (verification not implemented)	956
Maxima [A] (verification not implemented)	957
Giac [A] (verification not implemented)	957
Mupad [B] (verification not implemented)	957
Reduce [B] (verification not implemented)	958

Optimal result

Integrand size = 13, antiderivative size = 90

$$\int \frac{\sqrt{a+bx}}{x^3} dx = \frac{b\sqrt{a+bx}}{4ax} - \frac{\sqrt{(a+bx)^3}}{2ax^2} - \frac{b^2 \log\left(\frac{-\sqrt{a}+\sqrt{a+bx}}{\sqrt{a}+\sqrt{a+bx}}\right)}{8a^{3/2}}$$

output $1/4*b*(b*x+a)^{(1/2)}/a/x-1/2*((b*x+a)^3)^{(1/2)}/a/x^2-1/8*b^2*\ln((-a^{(1/2)}+b*x+a)^{(1/2)})/(a^{(1/2)}+(b*x+a)^{(1/2)})/a^{(3/2)}$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.61

$$\int \frac{\sqrt{a+bx}}{x^3} dx = -\frac{\sqrt{a+bx}(2a+bx)}{4ax^2} + \frac{b^2 \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{4a^{3/2}}$$

input `Integrate[Sqrt[a + b*x]/x^3,x]`

output $-1/4*(\operatorname{Sqrt}[a + b*x]*(2*a + b*x))/(a*x^2) + (b^2*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*x]/\operatorname{Sqrt}[a]])/(4*a^{(3/2)})$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.70, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {51, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{a+bx}}{x^3} dx \\
 & \quad \downarrow \text{51} \\
 & \frac{1}{4}b \int \frac{1}{x^2\sqrt{a+bx}} dx - \frac{\sqrt{a+bx}}{2x^2} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{4}b \left(-\frac{b \int \frac{1}{x\sqrt{a+bx}} dx}{2a} - \frac{\sqrt{a+bx}}{ax} \right) - \frac{\sqrt{a+bx}}{2x^2} \\
 & \quad \downarrow \text{73} \\
 & \frac{1}{4}b \left(-\frac{\int \frac{\frac{1}{a+bx} - \frac{a}{b}}{a} d\sqrt{a+bx}}{a} - \frac{\sqrt{a+bx}}{ax} \right) - \frac{\sqrt{a+bx}}{2x^2} \\
 & \quad \downarrow \text{221} \\
 & \frac{1}{4}b \left(\frac{\text{barctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{\sqrt{a+bx}}{ax} \right) - \frac{\sqrt{a+bx}}{2x^2}
 \end{aligned}$$

input

```
Int[Sqrt[a + b*x]/x^3,x]
```

output

```
-1/2*Sqrt[a + b*x]/x^2 + (b*(-(Sqrt[a + b*x]/(a*x)) + (b*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/a^(3/2)))/4
```

Definitions of rubi rules used

rule 51

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x
] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]
```

rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((
m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x],
x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 221

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.49

method	result	size
risch	$-\frac{\sqrt{bx+a}(bx+2a)}{4x^2a} + \frac{b^2 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{4a^{\frac{3}{2}}}$	44
pseudoelliptic	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)b^2x^2 - (2a^{\frac{3}{2}} + bx\sqrt{a})\sqrt{bx+a}}{4a^{\frac{3}{2}}x^2}$	50
derivativedivides	$2b^2 \left(-\frac{(bx+a)^{\frac{3}{2}}}{8a} + \frac{\sqrt{bx+a}}{8} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{8a^{\frac{3}{2}}} \right)$	54
default	$2b^2 \left(-\frac{(bx+a)^{\frac{3}{2}}}{8a} + \frac{\sqrt{bx+a}}{8} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{8a^{\frac{3}{2}}} \right)$	54

input `int((b*x+a)^(1/2)/x^3,x,method=_RETURNVERBOSE)`

output
$$-1/4*(b*x+a)^{(1/2)}*(b*x+2*a)/x^2/a+1/4*b^2/a^{(3/2)}*\operatorname{arctanh}((b*x+a)^{(1/2)}/a^{(1/2)})$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.29

$$\int \frac{\sqrt{a+bx}}{x^3} dx = \left[\frac{\sqrt{ab^2x^2} \log\left(\frac{bx+2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2(abx+2a^2)\sqrt{bx+a}}{8a^2x^2}, \right. \\ \left. - \frac{\sqrt{-ab^2x^2} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (abx+2a^2)\sqrt{bx+a}}{4a^2x^2} \right]$$

input `integrate((b*x+a)^(1/2)/x^3,x, algorithm="fricas")`

output
$$\left[\frac{1}{8}*(\operatorname{sqrt}(a)*b^2*x^2*\log((b*x+2*\operatorname{sqrt}(b*x+a))*\operatorname{sqrt}(a)+2*a)/x) - 2*(a*b*x+2*a^2)*\operatorname{sqrt}(b*x+a)/(a^2*x^2), -1/4*(\operatorname{sqrt}(-a)*b^2*x^2*\operatorname{arctan}(\operatorname{sqrt}(-a)/\operatorname{sqrt}(b*x+a)) + (a*b*x+2*a^2)*\operatorname{sqrt}(b*x+a))/(a^2*x^2) \right]$$

Sympy [A] (verification not implemented)

Time = 2.09 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{a+bx}}{x^3} dx = -\frac{a}{2\sqrt{bx^2}\sqrt{\frac{a}{bx}+1}} - \frac{3\sqrt{b}}{4x^{\frac{3}{2}}\sqrt{\frac{a}{bx}+1}} - \frac{b^{\frac{3}{2}}}{4a\sqrt{x}\sqrt{\frac{a}{bx}+1}} + \frac{b^2 \operatorname{asinh}\left(\frac{\sqrt{a}}{\sqrt{b}\sqrt{x}}\right)}{4a^{\frac{3}{2}}}$$

input `integrate((b*x+a)**(1/2)/x**3,x)`

output
$$-a/(2*\operatorname{sqrt}(b)*x^{(5/2)}*\operatorname{sqrt}(a/(b*x)+1)) - 3*\operatorname{sqrt}(b)/(4*x^{(3/2)}*\operatorname{sqrt}(a/(b*x)+1)) - b^{(3/2)}/(4*a*\operatorname{sqrt}(x)*\operatorname{sqrt}(a/(b*x)+1)) + b^{(2)}*\operatorname{asinh}(\operatorname{sqrt}(a)/(\operatorname{sqrt}(b)*\operatorname{sqrt}(x)))/(4*a^{(3/2)})$$

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.98

$$\int \frac{\sqrt{a+bx}}{x^3} dx = -\frac{b^2 \log\left(\frac{\sqrt{bx+a}-\sqrt{a}}{\sqrt{bx+a}+\sqrt{a}}\right)}{8a^{\frac{3}{2}}} - \frac{(bx+a)^{\frac{3}{2}}b^2 + \sqrt{bx+a}aab^2}{4((bx+a)^2a - 2(bx+a)a^2 + a^3)}$$

input `integrate((b*x+a)^(1/2)/x^3,x, algorithm="maxima")`output `-1/8*b^2*log((sqrt(b*x + a) - sqrt(a))/(sqrt(b*x + a) + sqrt(a)))/a^(3/2)
- 1/4*((b*x + a)^(3/2)*b^2 + sqrt(b*x + a)*a*b^2)/((b*x + a)^2*a - 2*(b*x + a)*a^2 + a^3)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.73

$$\int \frac{\sqrt{a+bx}}{x^3} dx = -\frac{b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa}} + \frac{(bx+a)^{\frac{3}{2}}b^3 + \sqrt{bx+a}aab^3}{4b}$$

input `integrate((b*x+a)^(1/2)/x^3,x, algorithm="giac")`output `-1/4*(b^3*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a) + ((b*x + a)^(3/2)*b^3 + sqrt(b*x + a)*a*b^3)/(a*b^2*x^2))/b`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.53

$$\int \frac{\sqrt{a+bx}}{x^3} dx = \frac{b^2 \operatorname{atanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{4a^{3/2}} - \frac{(a+bx)^{3/2}}{4ax^2} - \frac{\sqrt{a+bx}}{4x^2}$$

input `int((a + b*x)^(1/2)/x^3,x)`

output

$$(b^2 \operatorname{atanh}((a + b*x)^{1/2}/a^{1/2}))/ (4*a^{3/2}) - (a + b*x)^{3/2}/(4*a*x^2) - (a + b*x)^{1/2}/(4*x^2)$$

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.80

$$\int \frac{\sqrt{a+bx}}{x^3} dx$$

$$= \frac{-4\sqrt{bx+a}a^2 - 2\sqrt{bx+a}abx - \sqrt{a}\log(\sqrt{bx+a} - \sqrt{a})b^2x^2 + \sqrt{a}\log(\sqrt{bx+a} + \sqrt{a})b^2x^2}{8a^2x^2}$$

input

```
int((b*x+a)^(1/2)/x^3,x)
```

output

```
( - 4*sqrt(a + b*x)*a**2 - 2*sqrt(a + b*x)*a*b*x - sqrt(a)*log(sqrt(a + b*x) - sqrt(a))*b**2*x**2 + sqrt(a)*log(sqrt(a + b*x) + sqrt(a))*b**2*x**2)/
(8*a**2*x**2)
```

3.143 $\int \frac{\sqrt{(a+bx)^3}}{x} dx$

Optimal result	959
Mathematica [A] (verified)	959
Rubi [A] (verified)	960
Maple [A] (verified)	961
Fricas [B] (verification not implemented)	962
Sympy [F]	962
Maxima [F]	963
Giac [A] (verification not implemented)	963
Mupad [F(-1)]	963
Reduce [B] (verification not implemented)	964

Optimal result

Integrand size = 15, antiderivative size = 65

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = 2\sqrt{a+bx} \left(a + \frac{1}{3}(a+bx) \right) + a^{3/2} \log \left(\frac{-\sqrt{a} + \sqrt{a+bx}}{\sqrt{a} + \sqrt{a+bx}} \right)$$

output `2*(b*x+a)^(1/2)*(4/3*a+1/3*b*x)+a^(3/2)*ln((-a^(1/2)+(b*x+a)^(1/2))/(a^(1/2)+(b*x+a)^(1/2)))`

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = \frac{2\sqrt{(a+bx)^3} \left(\sqrt{a+bx}(4a+bx) - 3a^{3/2} \operatorname{arctanh} \left(\frac{\sqrt{a+bx}}{\sqrt{a}} \right) \right)}{3(a+bx)^{3/2}}$$

input `Integrate[Sqrt[(a + b*x)^3]/x,x]`

output `(2*Sqrt[(a + b*x)^3]*(Sqrt[a + b*x]*(4*a + b*x) - 3*a^(3/2)*ArcTanh[Sqrt[a + b*x]/Sqrt[a]]))/(3*(a + b*x)^(3/2))`

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2008, 60, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{(a+bx)^3}}{x} dx \\
 & \quad \downarrow \text{2008} \\
 & \frac{\sqrt{(a+bx)^3} \int \frac{(a+bx)^{3/2}}{x} dx}{(a+bx)^{3/2}} \\
 & \quad \downarrow \text{60} \\
 & \frac{\sqrt{(a+bx)^3} \left(a \int \frac{\sqrt{a+bx}}{x} dx + \frac{2}{3} (a+bx)^{3/2} \right)}{(a+bx)^{3/2}} \\
 & \quad \downarrow \text{60} \\
 & \frac{\sqrt{(a+bx)^3} \left(a \left(a \int \frac{1}{x\sqrt{a+bx}} dx + 2\sqrt{a+bx} \right) + \frac{2}{3} (a+bx)^{3/2} \right)}{(a+bx)^{3/2}} \\
 & \quad \downarrow \text{73} \\
 & \frac{\sqrt{(a+bx)^3} \left(a \left(\frac{2a \int \frac{\frac{1}{a+bx} - \frac{a}{b}}{b} d\sqrt{a+bx}}{b} + 2\sqrt{a+bx} \right) + \frac{2}{3} (a+bx)^{3/2} \right)}{(a+bx)^{3/2}} \\
 & \quad \downarrow \text{221} \\
 & \frac{\sqrt{(a+bx)^3} \left(a \left(2\sqrt{a+bx} - 2\sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a+bx}}{\sqrt{a}} \right) \right) + \frac{2}{3} (a+bx)^{3/2} \right)}{(a+bx)^{3/2}}
 \end{aligned}$$

input `Int[Sqrt[(a + b*x)^3]/x,x]`

output `(Sqrt[(a + b*x)^3]*((2*(a + b*x)^(3/2))/3 + a*(2*Sqrt[a + b*x] - 2*Sqrt[a]*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])))/(a + b*x)^(3/2)`

Definitions of rubi rules used

rule 60

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*((b*c - a*d)/(
b*(m + n + 1))) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!Integer
Q[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinear
Q[a, b, c, d, m, n, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

rule 2008

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Exp
on[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p),
x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x
] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.86

method	result	size
default	$-\frac{2\sqrt{(bx+a)^3} \left(3a^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right) - (bx+a)^{\frac{3}{2}} - 3\sqrt{bx+a} \right)}{3(bx+a)^{\frac{3}{2}}}$	56

input

```
int(((b*x+a)^3)^(1/2)/x,x,method=_RETURNVERBOSE)
```

output
$$-2/3*((b*x+a)^3)^{(1/2)}*(3*a^{(3/2)}*\operatorname{arctanh}((b*x+a)^{(1/2)}/a^{(1/2)})-(b*x+a)^{(3/2)}-3*(b*x+a)^{(1/2)}*a)/(b*x+a)^{(3/2)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. $2(49) = 98$.

Time = 0.12 (sec) , antiderivative size = 243, normalized size of antiderivative = 3.74

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = \left[\frac{3(abx + a^2)\sqrt{a} \log\left(\frac{b^2x^2 + 3abx + 2a^2 - 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}\sqrt{a}}{bx^2 + ax}\right) + 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}(bx + a)}{3(bx + a)} \right]$$

input `integrate(((b*x+a)^3)^(1/2)/x,x, algorithm="fricas")`

output
$$\left[\frac{1}{3} * (3 * (a * b * x + a^2) * \sqrt{a} * \log((b^2 * x^2 + 3 * a * b * x + 2 * a^2 - 2 * \sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3}) * \sqrt{a}) / (b * x^2 + a * x)) + 2 * \sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3} * (b * x + 4 * a) / (b * x + a), \frac{2}{3} * (3 * (a * b * x + a^2) * \sqrt{-a} * \arctan(\sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3}) * \sqrt{-a} / (b^2 * x^2 + 2 * a * b * x + a^2)) + \sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3} * (b * x + 4 * a) / (b * x + a) \right]$$

Sympy [F]

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = \int \frac{\sqrt{(a+bx)^3}}{x} dx$$

input `integrate(((b*x+a)**3)**(1/2)/x,x)`

output `Integral(sqrt((a + b*x)**3)/x, x)`

Maxima [F]

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = \int \frac{\sqrt{(bx+a)^3}}{x} dx$$

input `integrate(((b*x+a)^3)^(1/2)/x,x, algorithm="maxima")`

output `integrate(sqrt((b*x + a)^3)/x, x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.68

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = \frac{2a^2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + \frac{2}{3}(bx+a)^{\frac{3}{2}} + 2\sqrt{bx+aa}$$

input `integrate(((b*x+a)^3)^(1/2)/x,x, algorithm="giac")`

output `2*a^2*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + 2/3*(b*x + a)^(3/2) + 2*sqrt(b*x + a)*a`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = \int \frac{\sqrt{(a+bx)^3}}{x} dx$$

input `int(((a + b*x)^3)^(1/2)/x,x)`

output `int(((a + b*x)^3)^(1/2)/x, x)`

Reduce [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.78

$$\int \frac{\sqrt{(a+bx)^3}}{x} dx = \frac{8\sqrt{bx+a}a}{3} + \frac{2\sqrt{bx+a}bx}{3} + \sqrt{a} \log(\sqrt{bx+a} - \sqrt{a})a - \sqrt{a} \log(\sqrt{bx+a} + \sqrt{a})a$$

input `int(((b*x+a)^3)^(1/2)/x,x)`output `(8*sqrt(a + b*x)*a + 2*sqrt(a + b*x)*b*x + 3*sqrt(a)*log(sqrt(a + b*x) - sqrt(a))*a - 3*sqrt(a)*log(sqrt(a + b*x) + sqrt(a))*a)/3`

3.144 $\int \frac{\sqrt{(a+bx)^3}}{x^2} dx$

Optimal result	965
Mathematica [A] (verified)	965
Rubi [A] (verified)	966
Maple [A] (verified)	968
Fricas [A] (verification not implemented)	968
Sympy [F]	969
Maxima [F]	969
Giac [A] (verification not implemented)	969
Mupad [F(-1)]	970
Reduce [B] (verification not implemented)	970

Optimal result

Integrand size = 15, antiderivative size = 93

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx = -\frac{\sqrt{(a+bx)^5}}{ax} + \frac{3b\left(2\sqrt{a+bx}\left(a + \frac{1}{3}(a+bx)\right) + a^{3/2} \log\left(\frac{-\sqrt{a} + \sqrt{a+bx}}{\sqrt{a} + \sqrt{a+bx}}\right)\right)}{2a}$$

output `-((b*x+a)^5)^(1/2)/a/x+3/2*b*(2*(b*x+a)^(1/2)*(4/3*a+1/3*b*x)+a^(3/2)*ln((-a^(1/2)+(b*x+a)^(1/2))/(a^(1/2)+(b*x+a)^(1/2))))/a`

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.72

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx = -\frac{\sqrt{(a+bx)^3}\left((a-2bx)\sqrt{a+bx} + 3\sqrt{abx} \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)\right)}{x(a+bx)^{3/2}}$$

input `Integrate[Sqrt[(a + b*x)^3]/x^2,x]`

output

$$-\left(\frac{\sqrt{(a+bx)^3} \left((a-2bx)\sqrt{a+bx} + 3\sqrt{a}bx \operatorname{ArcTanh}\left[\frac{\sqrt{a+bx}}{\sqrt{a}}\right] \right)}{x(a+bx)^{3/2}}\right)$$
Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.82, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2008, 51, 60, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{(a+bx)^3}}{x^2} dx \\ & \quad \downarrow \text{2008} \\ & \frac{\sqrt{(a+bx)^3} \int \frac{(a+bx)^{3/2}}{x^2} dx}{(a+bx)^{3/2}} \\ & \quad \downarrow \text{51} \\ & \frac{\sqrt{(a+bx)^3} \left(\frac{3}{2}b \int \frac{\sqrt{a+bx}}{x} dx - \frac{(a+bx)^{3/2}}{x} \right)}{(a+bx)^{3/2}} \\ & \quad \downarrow \text{60} \\ & \frac{\sqrt{(a+bx)^3} \left(\frac{3}{2}b \left(a \int \frac{1}{x\sqrt{a+bx}} dx + 2\sqrt{a+bx} \right) - \frac{(a+bx)^{3/2}}{x} \right)}{(a+bx)^{3/2}} \\ & \quad \downarrow \text{73} \\ & \frac{\sqrt{(a+bx)^3} \left(\frac{3}{2}b \left(\frac{2a \int \frac{1}{\frac{a+bx}{b} - \frac{a}{b}} d\sqrt{a+bx}}{b} + 2\sqrt{a+bx} \right) - \frac{(a+bx)^{3/2}}{x} \right)}{(a+bx)^{3/2}} \\ & \quad \downarrow \text{221} \\ & \frac{\sqrt{(a+bx)^3} \left(\frac{3}{2}b \left(2\sqrt{a+bx} - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right) \right) - \frac{(a+bx)^{3/2}}{x} \right)}{(a+bx)^{3/2}} \end{aligned}$$

input `Int[Sqrt[(a + b*x)^3]/x^2,x]`

output `(Sqrt[(a + b*x)^3]*(-((a + b*x)^(3/2)/x) + (3*b*(2*Sqrt[a + b*x] - 2*Sqrt[a])*ArcTanh[Sqrt[a + b*x]/Sqrt[a]]))/2)/(a + b*x)^(3/2)`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1))] Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 2008 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Expon[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p), x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]`

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.73

method	result	size
default	$-\frac{\sqrt{(bx+a)^3} \left(-2bx\sqrt{bx+a}\sqrt{a} + 3 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right) abx + a^{\frac{3}{2}}\sqrt{bx+a} \right)}{(bx+a)^{\frac{3}{2}}x\sqrt{a}}$	68
risch	$-\frac{a\sqrt{(bx+a)^3}}{(bx+a)x} + \frac{b\left(4\sqrt{bx+a} - 6\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)\right)\sqrt{(bx+a)^3}}{2(bx+a)^{\frac{3}{2}}}$	70

input `int(((b*x+a)^3)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

output
$$-\left((b*x+a)^3\right)^{1/2} * \left(-2*b*x*(b*x+a)^{1/2} * a^{1/2} + 3*\operatorname{arctanh}\left((b*x+a)^{1/2}/a^{1/2}\right) * a*b*x + a^{3/2} * (b*x+a)^{1/2}\right) / (b*x+a)^{3/2} / x / a^{1/2}$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 260, normalized size of antiderivative = 2.80

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx$$

$$= \left[\frac{3(b^2x^2 + abx)\sqrt{a} \log\left(\frac{b^2x^2 + 3abx + 2a^2 - 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}\sqrt{a}}{bx^2 + ax}\right) + 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}(2bx + a)}{2(bx^2 + ax)} \right]$$

input `integrate(((b*x+a)^3)^(1/2)/x^2,x, algorithm="fricas")`

output
$$\left[\frac{1}{2} * \left(3 * (b^2 * x^2 + a * b * x) * \sqrt{a} * \log\left(\frac{b^2 * x^2 + 3 * a * b * x + 2 * a^2 - 2 * \sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3} * \sqrt{a}}{b * x^2 + a * x}\right) + 2 * \sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3} * (2 * b * x - a) / (b * x^2 + a * x), \right. \right. \\ \left. \left. 3 * (b^2 * x^2 + a * b * x) * \sqrt{-a} * \arctan\left(\frac{\sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3} * \sqrt{-a}}{b^2 * x^2 + 2 * a * b * x + a^2}\right) + \sqrt{b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3} * (2 * b * x - a) / (b * x^2 + a * x) \right]$$

Sympy [F]

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx = \int \frac{\sqrt{(a+bx)^3}}{x^2} dx$$

input `integrate(((b*x+a)**3)**(1/2)/x**2,x)`

output `Integral(sqrt((a + b*x)**3)/x**2, x)`

Maxima [F]

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx = \int \frac{\sqrt{(bx+a)^3}}{x^2} dx$$

input `integrate(((b*x+a)^3)^(1/2)/x^2,x, algorithm="maxima")`

output `integrate(sqrt((b*x + a)^3)/x^2, x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.54

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx = \left(\frac{3a \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} + 2\sqrt{bx+a} - \frac{\sqrt{bx+aa}}{bx} \right) b$$

input `integrate(((b*x+a)^3)^(1/2)/x^2,x, algorithm="giac")`

output `(3*a*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) + 2*sqrt(b*x + a) - sqrt(b*x + a)*a/(b*x))*b`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx = \int \frac{\sqrt{(a+bx)^3}}{x^2} dx$$

input `int(((a + b*x)^3)^(1/2)/x^2,x)`output `int(((a + b*x)^3)^(1/2)/x^2, x)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.63

$$\int \frac{\sqrt{(a+bx)^3}}{x^2} dx$$

$$= \frac{-2\sqrt{bx+a}a + 4\sqrt{bx+a}bx + 3\sqrt{a}\log(\sqrt{bx+a} - \sqrt{a})bx - 3\sqrt{a}\log(\sqrt{bx+a} + \sqrt{a})bx}{2x}$$

input `int(((b*x+a)^3)^(1/2)/x^2,x)`output `(- 2*sqrt(a + b*x)*a + 4*sqrt(a + b*x)*b*x + 3*sqrt(a)*log(sqrt(a + b*x) - sqrt(a))*b*x - 3*sqrt(a)*log(sqrt(a + b*x) + sqrt(a))*b*x)/(2*x)`

3.145 $\int \frac{\sqrt{(a+bx)^3}}{x^3} dx$

Optimal result	971
Mathematica [A] (verified)	971
Rubi [A] (verified)	972
Maple [A] (verified)	974
Fricas [A] (verification not implemented)	974
Sympy [F]	975
Maxima [F]	975
Giac [A] (verification not implemented)	975
Mupad [F(-1)]	976
Reduce [B] (verification not implemented)	976

Optimal result

Integrand size = 15, antiderivative size = 135

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx = \left(-\frac{1}{2ax^2} - \frac{b}{4a^2x} \right) \sqrt{(a+bx)^5} + \frac{3b^2 \left(\frac{b\sqrt{a+bx}}{4ax} - \frac{\sqrt{(a+bx)^3}}{2ax^2} - \frac{b^2 \log\left(\frac{-\sqrt{a}+\sqrt{a+bx}}{\sqrt{a}+\sqrt{a+bx}}\right)}{8a^{3/2}} \right)}{8a^2}$$

output

$$\left(-\frac{1}{2} \frac{1}{a} x^{-2} - \frac{1}{4} \frac{b}{a^2} \frac{1}{x}\right) \sqrt{(bx+a)^5} + \frac{3}{8} \frac{b^2}{a^2} \left(\frac{1}{4} \frac{b \sqrt{bx+a}}{ax} + \frac{1}{2} \frac{1}{a} \frac{1}{x} - \frac{1}{2} \frac{1}{a} \sqrt{(bx+a)^3} \sqrt{\frac{1}{a} x^{-2}} - \frac{1}{8} \frac{b^2}{a^2} \ln\left(\frac{-\sqrt{a} + \sqrt{bx+a}}{\sqrt{a} + \sqrt{bx+a}}\right) \sqrt{\frac{1}{a} x^{-2}} + \frac{1}{a} \sqrt{(bx+a)^3} \sqrt{\frac{1}{a} x^{-2}}\right) \frac{1}{a^2}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx = -\frac{\sqrt{(a+bx)^3} \left(\sqrt{a} \sqrt{a+bx} (2a+5bx) + 3b^2 x^2 \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right) \right)}{4\sqrt{a} x^2 (a+bx)^{3/2}}$$

input

```
Integrate[Sqrt[(a + b*x)^3]/x^3,x]
```

output

```
-1/4*(Sqrt[(a + b*x)^3]*(Sqrt[a]*Sqrt[a + b*x]*(2*a + 5*b*x) + 3*b^2*x^2*ArcTanh[Sqrt[a + b*x]/Sqrt[a]]))/(Sqrt[a]*x^2*(a + b*x)^(3/2))
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.61, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2008, 51, 51, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx$$

↓ 2008

$$\frac{\sqrt{(a+bx)^3} \int \frac{(a+bx)^{3/2}}{x^3} dx}{(a+bx)^{3/2}}$$

↓ 51

$$\frac{\sqrt{(a+bx)^3} \left(\frac{3}{4} b \int \frac{\sqrt{a+bx}}{x^2} dx - \frac{(a+bx)^{3/2}}{2x^2} \right)}{(a+bx)^{3/2}}$$

↓ 51

$$\frac{\sqrt{(a+bx)^3} \left(\frac{3}{4} b \left(\frac{1}{2} b \int \frac{1}{x\sqrt{a+bx}} dx - \frac{\sqrt{a+bx}}{x} \right) - \frac{(a+bx)^{3/2}}{2x^2} \right)}{(a+bx)^{3/2}}$$

↓ 73

$$\frac{\sqrt{(a+bx)^3} \left(\frac{3}{4} b \left(\int \frac{1}{\frac{a+bx}{b} - \frac{a}{b}} d\sqrt{a+bx} - \frac{\sqrt{a+bx}}{x} \right) - \frac{(a+bx)^{3/2}}{2x^2} \right)}{(a+bx)^{3/2}}$$

↓ 221

$$\frac{\sqrt{(a+bx)^3} \left(\frac{3}{4} b \left(-\frac{\text{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\sqrt{a+bx}}{x} \right) - \frac{(a+bx)^{3/2}}{2x^2} \right)}{(a+bx)^{3/2}}$$

input `Int[Sqrt[(a + b*x)^3]/x^3,x]`

output `(Sqrt[(a + b*x)^3]*(-1/2*(a + b*x)^(3/2)/x^2 + (3*b*(-(Sqrt[a + b*x]/x) - (b*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/Sqrt[a])/4))/(a + b*x)^(3/2)`

Defintions of rubi rules used

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))]
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x]
&& ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 2008 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Expon[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p), x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]`

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.50

method	result	size
risch	$-\frac{(5bx+2a)\sqrt{(bx+a)^3}}{4(bx+a)x^2} - \frac{3b^2 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)\sqrt{(bx+a)^3}}{4\sqrt{a}(bx+a)^{\frac{3}{2}}}$	67
default	$-\frac{\sqrt{(bx+a)^3} \left(3 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right) b^2 x^2 + 5(bx+a)^{\frac{3}{2}} \sqrt{a} - 3a^{\frac{3}{2}} \sqrt{bx+a} \right)}{4(bx+a)^{\frac{3}{2}} x^2 \sqrt{a}}$	70

input `int(((b*x+a)^3)^(1/2)/x^3,x,method=_RETURNVERBOSE)`

output
$$-1/4/(b*x+a)*(5*b*x+2*a)/x^2*((b*x+a)^3)^(1/2)-3/4*b^2/a^(1/2)*\operatorname{arctanh}((b*x+a)^(1/2)/a^(1/2))*((b*x+a)^3)^(1/2)/(b*x+a)^(3/2)$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.12

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx = \left[\frac{3(b^3x^3 + ab^2x^2)\sqrt{a} \log\left(\frac{b^2x^2+3abx+2a^2-2\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}\sqrt{a}}{bx^2+ax}\right) - 2\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}(5}{8(abx^3+a^2x^2)} \right]$$

input `integrate(((b*x+a)^3)^(1/2)/x^3,x, algorithm="fricas")`

output
$$\left[\frac{1}{8} * (3 * (b^3 * x^3 + a * b^2 * x^2) * \operatorname{sqrt}(a) * \log((b^2 * x^2 + 3 * a * b * x + 2 * a^2 - 2 * \operatorname{sqrt}(b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3)) * \operatorname{sqrt}(a)) / (b * x^2 + a * x)) - 2 * \operatorname{sqrt}(b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3) * (5 * a * b * x + 2 * a^2)) / (a * b * x^3 + a^2 * x^2), \frac{1}{4} * (3 * (b^3 * x^3 + a * b^2 * x^2) * \operatorname{sqrt}(-a) * \operatorname{arctan}(\operatorname{sqrt}(b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3)) * \operatorname{sqrt}(-a) / (b^2 * x^2 + 2 * a * b * x + a^2)) - \operatorname{sqrt}(b^3 * x^3 + 3 * a * b^2 * x^2 + 3 * a^2 * b * x + a^3) * (5 * a * b * x + 2 * a^2)) / (a * b * x^3 + a^2 * x^2) \right]$$

Sympy [F]

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx = \int \frac{\sqrt{(a+bx)^3}}{x^3} dx$$

input `integrate(((b*x+a)**3)**(1/2)/x**3,x)`

output `Integral(sqrt((a + b*x)**3)/x**3, x)`

Maxima [F]

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx = \int \frac{\sqrt{(bx+a)^3}}{x^3} dx$$

input `integrate(((b*x+a)^3)^(1/2)/x^3,x, algorithm="maxima")`

output `integrate(sqrt((b*x + a)^3)/x^3, x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.47

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx = \frac{3b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-a}} - \frac{5(bx+a)^{\frac{3}{2}}b^3 - 3\sqrt{bx+a}ab^3}{4b}$$

input `integrate(((b*x+a)^3)^(1/2)/x^3,x, algorithm="giac")`

output `1/4*(3*b^3*arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a) - (5*(b*x + a)^(3/2)*b^3 - 3*sqrt(b*x + a)*a*b^3)/(b^2*x^2))/b`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx = \int \frac{\sqrt{(a+bx)^3}}{x^3} dx$$

input `int(((a + b*x)^3)^(1/2)/x^3,x)`output `int(((a + b*x)^3)^(1/2)/x^3, x)`**Reduce [B] (verification not implemented)**

Time = 0.28 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.54

$$\int \frac{\sqrt{(a+bx)^3}}{x^3} dx$$

$$= \frac{-4\sqrt{bx+a}a^2 - 10\sqrt{bx+a}abx + 3\sqrt{a}\log(\sqrt{bx+a} - \sqrt{a})b^2x^2 - 3\sqrt{a}\log(\sqrt{bx+a} + \sqrt{a})b^2x^2}{8ax^2}$$

input `int(((b*x+a)^3)^(1/2)/x^3,x)`output `(- 4*sqrt(a + b*x)*a**2 - 10*sqrt(a + b*x)*a*b*x + 3*sqrt(a)*log(sqrt(a + b*x) - sqrt(a))*b**2*x**2 - 3*sqrt(a)*log(sqrt(a + b*x) + sqrt(a))*b**2*x**2)/(8*a*x**2)`

3.146 $\int \frac{1}{x^2\sqrt{a+bx}} dx$

Optimal result	977
Mathematica [A] (verified)	977
Rubi [A] (verified)	978
Maple [A] (verified)	979
Fricas [A] (verification not implemented)	980
Sympy [A] (verification not implemented)	980
Maxima [A] (verification not implemented)	981
Giac [A] (verification not implemented)	981
Mupad [B] (verification not implemented)	981
Reduce [B] (verification not implemented)	982

Optimal result

Integrand size = 13, antiderivative size = 67

$$\int \frac{1}{x^2\sqrt{a+bx}} dx = -\frac{\sqrt{a+bx}}{ax} - \frac{b \log\left(\frac{-\sqrt{a}+\sqrt{a+bx}}{\sqrt{a}+\sqrt{a+bx}}\right)}{2\sqrt{ax}}$$

output `-(b*x+a)^(1/2)/a/x-1/2*b*ln((-a^(1/2)+(b*x+a)^(1/2))/(a^(1/2)+(b*x+a)^(1/2)))/a^(1/2)/x`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.61

$$\int \frac{1}{x^2\sqrt{a+bx}} dx = -\frac{\sqrt{a+bx}}{ax} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{a^{3/2}}$$

input `Integrate[1/(x^2*Sqrt[a + b*x]),x]`

output `-(Sqrt[a + b*x]/(a*x)) + (b*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/a^(3/2)`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.61, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^2 \sqrt{a+bx}} dx \\
 & \quad \downarrow \text{52} \\
 & -\frac{b \int \frac{1}{x \sqrt{a+bx}} dx}{2a} - \frac{\sqrt{a+bx}}{ax} \\
 & \quad \downarrow \text{73} \\
 & -\frac{\int \frac{1}{\frac{a+bx}{b} - \frac{a}{b}} d\sqrt{a+bx}}{a} - \frac{\sqrt{a+bx}}{ax} \\
 & \quad \downarrow \text{221} \\
 & \frac{\text{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{\sqrt{a+bx}}{ax}
 \end{aligned}$$

input `Int[1/(x^2*Sqrt[a + b*x]),x]`

output `-(Sqrt[a + b*x]/(a*x)) + (b*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/a^(3/2)`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
 {p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
 d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
 Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
 inearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
 /Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.51

method	result	size
risch	$-\frac{\sqrt{bx+a}}{ax} + \frac{b \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{a^{\frac{3}{2}}}$	34
pseudoelliptic	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)bx - \sqrt{bx+a}\sqrt{a}}{a^{\frac{3}{2}}x}$	36
derivativedivides	$2b\left(-\frac{\sqrt{bx+a}}{2abx} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{2a^{\frac{3}{2}}}\right)$	40
default	$2b\left(-\frac{\sqrt{bx+a}}{2abx} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{2a^{\frac{3}{2}}}\right)$	40

input `int(1/x^2/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output `-(b*x+a)^(1/2)/a/x+b/a^(3/2)*arctanh((b*x+a)^(1/2)/a^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.34

$$\int \frac{1}{x^2 \sqrt{a+bx}} dx = \left[\frac{\sqrt{abx} \log\left(\frac{bx+2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) - 2\sqrt{bx+aa}}{2a^2x}, \right. \\ \left. - \frac{\sqrt{-abx} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + \sqrt{bx+aa}}{a^2x} \right]$$

input `integrate(1/x^2/(b*x+a)^(1/2),x, algorithm="fricas")`output `[1/2*(sqrt(a)*b*x*log((b*x + 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) - 2*sqrt(b*x + a)*a)/(a^2*x), -(sqrt(-a)*b*x*arctan(sqrt(-a)/sqrt(b*x + a)) + sqrt(b*x + a)*a)/(a^2*x)]`**Sympy [A] (verification not implemented)**

Time = 1.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.66

$$\int \frac{1}{x^2 \sqrt{a+bx}} dx = -\frac{\sqrt{b}\sqrt{\frac{a}{bx}+1}}{a\sqrt{x}} + \frac{b \operatorname{asinh}\left(\frac{\sqrt{a}}{\sqrt{b}\sqrt{x}}\right)}{a^{\frac{3}{2}}}$$

input `integrate(1/x**2/(b*x+a)**(1/2),x)`output `-sqrt(b)*sqrt(a/(b*x) + 1)/(a*sqrt(x)) + b*asinh(sqrt(a)/(sqrt(b)*sqrt(x)))/a**(3/2)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.90

$$\int \frac{1}{x^2 \sqrt{a+bx}} dx = -\frac{\sqrt{bx+a}}{(bx+a)a-a^2} - \frac{b \log\left(\frac{\sqrt{bx+a}-\sqrt{a}}{\sqrt{bx+a}+\sqrt{a}}\right)}{2a^{\frac{3}{2}}}$$

input `integrate(1/x^2/(b*x+a)^(1/2),x, algorithm="maxima")`output `-sqrt(b*x + a)*b/((b*x + a)*a - a^2) - 1/2*b*log((sqrt(b*x + a) - sqrt(a)) / (sqrt(b*x + a) + sqrt(a)))/a^(3/2)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.66

$$\int \frac{1}{x^2 \sqrt{a+bx}} dx = -b \left(\frac{\arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa}} + \frac{\sqrt{bx+a}}{abx} \right)$$

input `integrate(1/x^2/(b*x+a)^(1/2),x, algorithm="giac")`output `-b*(arctan(sqrt(b*x + a)/sqrt(-a))/sqrt(-a)*a) + sqrt(b*x + a)/(a*b*x)`**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.49

$$\int \frac{1}{x^2 \sqrt{a+bx}} dx = \frac{b \operatorname{atanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{\sqrt{a+bx}}{ax}$$

input `int(1/(x^2*(a + b*x)^(1/2)),x)`output `(b*atanh((a + b*x)^(1/2)/a^(1/2)))/a^(3/2) - (a + b*x)^(1/2)/(a*x)`

Reduce [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.76

$$\int \frac{1}{x^2 \sqrt{a+bx}} dx$$

$$= \frac{-2\sqrt{bx+a}a - \sqrt{a} \log(\sqrt{bx+a} - \sqrt{a})bx + \sqrt{a} \log(\sqrt{bx+a} + \sqrt{a})bx}{2a^2x}$$

input `int(1/x^2/(b*x+a)^(1/2),x)`output `(- 2*sqrt(a + b*x)*a - sqrt(a)*log(sqrt(a + b*x) - sqrt(a))*b*x + sqrt(a)
*log(sqrt(a + b*x) + sqrt(a))*b*x)/(2*a**2*x)`

3.147 $\int \frac{1}{x^3 \sqrt{a+bx}} dx$

Optimal result	983
Mathematica [A] (verified)	983
Rubi [A] (verified)	984
Maple [A] (verified)	985
Fricas [A] (verification not implemented)	986
Sympy [A] (verification not implemented)	986
Maxima [A] (verification not implemented)	987
Giac [A] (verification not implemented)	987
Mupad [B] (verification not implemented)	987
Reduce [B] (verification not implemented)	988

Optimal result

Integrand size = 13, antiderivative size = 81

$$\int \frac{1}{x^3 \sqrt{a+bx}} dx = \left(-\frac{1}{2ax^2} + \frac{3b}{4a^2x} \right) \sqrt{a+bx} + \frac{3b^2 \log\left(\frac{-\sqrt{a} + \sqrt{a+bx}}{\sqrt{a} + \sqrt{a+bx}}\right)}{8a^{5/2}}$$

output

```
(-1/2/a/x^2+3/4*b/a^2/x)*(b*x+a)^(1/2)+3/8*b^2*ln((-a^(1/2)+(b*x+a)^(1/2))
/(a^(1/2)+(b*x+a)^(1/2)))/a^(5/2)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.69

$$\int \frac{1}{x^3 \sqrt{a+bx}} dx = \frac{\sqrt{a+bx}(-2a+3bx)}{4a^2x^2} - \frac{3b^2 \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{4a^{5/2}}$$

input

```
Integrate[1/(x^3*Sqrt[a + b*x]),x]
```

output

```
(Sqrt[a + b*x]*(-2*a + 3*b*x))/(4*a^2*x^2) - (3*b^2*ArcTanh[Sqrt[a + b*x]/
Sqrt[a]])/(4*a^(5/2))
```


Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.85, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {52, 52, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^3 \sqrt{a+bx}} dx \\
 & \quad \downarrow 52 \\
 & -\frac{3b \int \frac{1}{x^2 \sqrt{a+bx}} dx}{4a} - \frac{\sqrt{a+bx}}{2ax^2} \\
 & \quad \downarrow 52 \\
 & -\frac{3b \left(-\frac{b \int \frac{1}{x \sqrt{a+bx}} dx}{2a} - \frac{\sqrt{a+bx}}{ax} \right)}{4a} - \frac{\sqrt{a+bx}}{2ax^2} \\
 & \quad \downarrow 73 \\
 & -\frac{3b \left(-\frac{\int \frac{\frac{1}{a+bx} - \frac{a}{b}}{a} d\sqrt{a+bx}}{4a} - \frac{\sqrt{a+bx}}{ax} \right)}{4a} - \frac{\sqrt{a+bx}}{2ax^2} \\
 & \quad \downarrow 221 \\
 & -\frac{3b \left(\frac{\text{barctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{\sqrt{a+bx}}{ax} \right)}{4a} - \frac{\sqrt{a+bx}}{2ax^2}
 \end{aligned}$$

input

```
Int[1/(x^3*Sqrt[a + b*x]),x]
```

output

```
-1/2*Sqrt[a + b*x]/(a*x^2) - (3*b*(-(Sqrt[a + b*x]/(a*x)) + (b*ArcTanh[Sqr
t[a + b*x]/Sqrt[a]]/a^(3/2)))/(4*a)
```

Defintions of rubi rules used

```
rule 52 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((
m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x],
x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]
```

```
rule 73 Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

```
rule 221 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.56

method	result	size
risch	$-\frac{\sqrt{bx+a}(-3bx+2a)}{4a^2x^2} - \frac{3b^2 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{4a^{\frac{5}{2}}}$	45
pseudoelliptic	$\frac{-3 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)b^2x^2+3bx\sqrt{bx+a}\sqrt{a}-2a^{\frac{3}{2}}\sqrt{bx+a}}{4a^{\frac{5}{2}}x^2}$	56
derivativedivides	$2b^2 \left(-\frac{\sqrt{bx+a}}{4a b^2x^2} - \frac{3 \left(-\frac{\sqrt{bx+a}}{2abx} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{2a^{\frac{3}{2}}} \right)}{4a} \right)$	66
default	$2b^2 \left(-\frac{\sqrt{bx+a}}{4a b^2x^2} - \frac{3 \left(-\frac{\sqrt{bx+a}}{2abx} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{2a^{\frac{3}{2}}} \right)}{4a} \right)$	66

```
input int(1/x^3/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-1/4*(b*x+a)^(1/2)*(-3*b*x+2*a)/a^2/x^2-3/4*b^2/a^(5/2)*arctanh((b*x+a)^(1/2)/a^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.48

$$\int \frac{1}{x^3 \sqrt{a+bx}} dx$$

$$= \left[\frac{3 \sqrt{ab^2 x^2} \log\left(\frac{bx-2\sqrt{bx+a}\sqrt{a+2a}}{x}\right) + 2(3abx - 2a^2)\sqrt{bx+a}}{8a^3 x^2}, \frac{3\sqrt{-ab^2 x^2} \arctan\left(\frac{\sqrt{-a}}{\sqrt{bx+a}}\right) + (3abx - 2a^2)\sqrt{bx+a}}{4a^3 x^2} \right]$$

input

```
integrate(1/x^3/(b*x+a)^(1/2),x, algorithm="fricas")
```

output

```
[1/8*(3*sqrt(a)*b^2*x^2*log((b*x - 2*sqrt(b*x + a)*sqrt(a) + 2*a)/x) + 2*(3*a*b*x - 2*a^2)*sqrt(b*x + a))/(a^3*x^2), 1/4*(3*sqrt(-a)*b^2*x^2*arctan(sqrt(-a)/sqrt(b*x + a)) + (3*a*b*x - 2*a^2)*sqrt(b*x + a))/(a^3*x^2)]
```

Sympy [A] (verification not implemented)

Time = 2.59 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.26

$$\int \frac{1}{x^3 \sqrt{a+bx}} dx = -\frac{1}{2\sqrt{b}x^{\frac{5}{2}}\sqrt{\frac{a}{bx}+1}} + \frac{\sqrt{b}}{4ax^{\frac{3}{2}}\sqrt{\frac{a}{bx}+1}}$$

$$+ \frac{3b^{\frac{3}{2}}}{4a^2\sqrt{x}\sqrt{\frac{a}{bx}+1}} - \frac{3b^2 \operatorname{asinh}\left(\frac{\sqrt{a}}{\sqrt{b}\sqrt{x}}\right)}{4a^{\frac{5}{2}}}$$

input

```
integrate(1/x**3/(b*x+a)**(1/2),x)
```

output

```
-1/(2*sqrt(b)*x**(5/2)*sqrt(a/(b*x) + 1)) + sqrt(b)/(4*a*x**(3/2)*sqrt(a/(b*x) + 1)) + 3*b**(3/2)/(4*a**2*sqrt(x)*sqrt(a/(b*x) + 1)) - 3*b**2*asinh(sqrt(a)/(sqrt(b)*sqrt(x)))/(4*a**(5/2))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.14

$$\int \frac{1}{x^3 \sqrt{a+bx}} dx = \frac{3b^2 \log\left(\frac{\sqrt{bx+a}-\sqrt{a}}{\sqrt{bx+a}+\sqrt{a}}\right)}{8a^{\frac{5}{2}}} + \frac{3(bx+a)^{\frac{3}{2}}b^2 - 5\sqrt{bx+a}ab^2}{4((bx+a)^2a^2 - 2(bx+a)a^3 + a^4)}$$

input `integrate(1/x^3/(b*x+a)^(1/2),x, algorithm="maxima")`output `3/8*b^2*log((sqrt(b*x + a) - sqrt(a))/(sqrt(b*x + a) + sqrt(a)))/a^(5/2) + 1/4*(3*(b*x + a)^(3/2)*b^2 - 5*sqrt(b*x + a)*a*b^2)/((b*x + a)^2*a^2 - 2*(b*x + a)*a^3 + a^4)`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.85

$$\int \frac{1}{x^3 \sqrt{a+bx}} dx = \frac{3b^3 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa^2}} + \frac{3(bx+a)^{\frac{3}{2}}b^3 - 5\sqrt{bx+a}ab^3}{4ba^2b^2x^2}$$

input `integrate(1/x^3/(b*x+a)^(1/2),x, algorithm="giac")`output `1/4*(3*b^3*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^2) + (3*(b*x + a)^(3/2)*b^3 - 5*sqrt(b*x + a)*a*b^3)/(a^2*b^2*x^2))/b`**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.63

$$\int \frac{1}{x^3 \sqrt{a+bx}} dx = \frac{3(a+bx)^{3/2}}{4a^2x^2} - \frac{5\sqrt{a+bx}}{4ax^2} - \frac{3b^2 \operatorname{atanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{4a^{5/2}}$$

input `int(1/(x^3*(a + b*x)^(1/2)),x)`

output

$$\frac{(3*(a + b*x)^{(3/2)})/(4*a^2*x^2) - (5*(a + b*x)^{(1/2)})/(4*a*x^2) - (3*b^2*a \tanh((a + b*x)^{(1/2)}/a^{(1/2)}))/(4*a^{(5/2)})}{8a^3x^2}$$

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.90

$$\int \frac{1}{x^3 \sqrt{a + bx}} dx$$

$$= \frac{-4\sqrt{bx + a} a^2 + 6\sqrt{bx + a} abx + 3\sqrt{a} \log(\sqrt{bx + a} - \sqrt{a}) b^2 x^2 - 3\sqrt{a} \log(\sqrt{bx + a} + \sqrt{a}) b^2 x^2}{8a^3x^2}$$

input

```
int(1/x^3/(b*x+a)^(1/2),x)
```

output

```
( - 4*sqrt(a + b*x)*a**2 + 6*sqrt(a + b*x)*a*b*x + 3*sqrt(a)*log(sqrt(a +
b*x) - sqrt(a))*b**2*x**2 - 3*sqrt(a)*log(sqrt(a + b*x) + sqrt(a))*b**2*x*
*2)/(8*a**3*x**2)
```

3.148 $\int \frac{1}{x\sqrt{(a+bx)^3}} dx$

Optimal result	989
Mathematica [A] (verified)	989
Rubi [A] (verified)	990
Maple [A] (verified)	991
Fricas [B] (verification not implemented)	992
Sympy [F]	992
Maxima [F]	993
Giac [A] (verification not implemented)	993
Mupad [F(-1)]	993
Reduce [B] (verification not implemented)	994

Optimal result

Integrand size = 15, antiderivative size = 57

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx = \frac{2}{a\sqrt{a+bx}} + \frac{\log\left(\frac{-\sqrt{a}+\sqrt{a+bx}}{\sqrt{a}+\sqrt{a+bx}}\right)}{a^{3/2}}$$

output $2/a/(b*x+a)^{(1/2)}+\ln((-a^{(1/2)}+(b*x+a)^{(1/2)})/(a^{(1/2)}+(b*x+a)^{(1/2)}))/a^{(3/2)}$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.98

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx = \frac{2(a+bx)\left(\sqrt{a}-\sqrt{a+bx}\operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)\right)}{a^{3/2}\sqrt{(a+bx)^3}}$$

input `Integrate[1/(x*Sqrt[(a + b*x)^3]),x]`

output $(2*(a + b*x)*(Sqrt[a] - Sqrt[a + b*x]*ArcTanh[Sqrt[a + b*x]/Sqrt[a]]))/(a^{(3/2)}*Sqrt[(a + b*x)^3])$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.04, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2008, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x\sqrt{(a+bx)^3}} dx \\
 & \quad \downarrow \text{2008} \\
 & \frac{(a+bx)^{3/2} \int \frac{1}{x(a+bx)^{3/2}} dx}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{61} \\
 & \frac{(a+bx)^{3/2} \left(\frac{\int \frac{1}{x\sqrt{a+bx}} dx}{a} + \frac{2}{a\sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{73} \\
 & \frac{(a+bx)^{3/2} \left(\frac{2 \int \frac{\frac{1}{a+bx} - \frac{a}{b}}{ab} d\sqrt{a+bx}}{ab} + \frac{2}{a\sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{221} \\
 & \frac{(a+bx)^{3/2} \left(\frac{2}{a\sqrt{a+bx}} - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{a^{3/2}} \right)}{\sqrt{(a+bx)^3}}
 \end{aligned}$$

input `Int[1/(x*Sqrt[(a + b*x)^3]),x]`

output `((a + b*x)^(3/2)*(2/(a*Sqrt[a + b*x]) - (2*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/a^(3/2)))/Sqrt[(a + b*x)^3]`

Definitions of rubi rules used

rule 61

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((
m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x],
x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0]
) || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n])) && IntLinearQ[a, b, c, d
, m, n, x]
```

rule 73

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

rule 2008

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Exp
on[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p),
x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x]
&& GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.82

method	result	size
default	$\frac{2(bx+a)\left(\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)a\sqrt{bx+a}-a^{\frac{3}{2}}\right)}{\sqrt{(bx+a)^3}a^{\frac{5}{2}}}$	47

input

```
int(1/x/((b*x+a)^3)^(1/2),x,method=_RETURNVERBOSE)
```


output

```
-2*(b*x+a)*(arctanh((b*x+a)^(1/2)/a^(1/2))*a*(b*x+a)^(1/2)-a^(3/2))/((b*x+a)^3)^(1/2)/a^(5/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. $2(45) = 90$.

Time = 0.10 (sec) , antiderivative size = 276, normalized size of antiderivative = 4.84

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx$$

$$= \left[\frac{(b^2x^2 + 2abx + a^2)\sqrt{a} \log\left(\frac{b^2x^2 + 3abx + 2a^2 - 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}\sqrt{a}}{bx^2 + ax}\right) + 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}}{a^2b^2x^2 + 2a^3bx + a^4} \right]$$

input

```
integrate(1/x/((b*x+a)^3)^(1/2),x, algorithm="fricas")
```

output

```
[((b^2*x^2 + 2*a*b*x + a^2)*sqrt(a)*log((b^2*x^2 + 3*a*b*x + 2*a^2 - 2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*sqrt(a))/(b*x^2 + a*x)) + 2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*a)/(a^2*b^2*x^2 + 2*a^3*b*x + a^4), 2*((b^2*x^2 + 2*a*b*x + a^2)*sqrt(-a)*arctan(sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*sqrt(-a)/(b^2*x^2 + 2*a*b*x + a^2)) + sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*a)/(a^2*b^2*x^2 + 2*a^3*b*x + a^4)]
```

Sympy [F]

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx = \int \frac{1}{x\sqrt{(a+bx)^3}} dx$$

input

```
integrate(1/x/((b*x+a)**3)**(1/2),x)
```

output

```
Integral(1/(x*sqrt((a + b*x)**3)), x)
```

Maxima [F]

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx = \int \frac{1}{\sqrt{(bx+a)^3}x} dx$$

input `integrate(1/x/((b*x+a)^3)^(1/2),x, algorithm="maxima")`

output `integrate(1/(sqrt((b*x + a)^3)*x), x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.65

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx = \frac{2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa}} + \frac{2}{\sqrt{bx+aa}}$$

input `integrate(1/x/((b*x+a)^3)^(1/2),x, algorithm="giac")`

output `2*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a) + 2/(sqrt(b*x + a)*a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx = \int \frac{1}{x\sqrt{(a+bx)^3}} dx$$

input `int(1/(x*((a + b*x)^3)^(1/2)),x)`

output `int(1/(x*((a + b*x)^3)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00

$$\int \frac{1}{x\sqrt{(a+bx)^3}} dx$$

$$= \frac{\sqrt{a}\sqrt{bx+a}\log(\sqrt{bx+a}-\sqrt{a}) - \sqrt{a}\sqrt{bx+a}\log(\sqrt{bx+a}+\sqrt{a}) + 2a}{\sqrt{bx+a}a^2}$$

input

```
int(1/x/((b*x+a)^3)^(1/2),x)
```

output

```
(sqrt(a)*sqrt(a + b*x)*log(sqrt(a + b*x) - sqrt(a)) - sqrt(a)*sqrt(a + b*x)
)*log(sqrt(a + b*x) + sqrt(a)) + 2*a)/(sqrt(a + b*x)*a**2)
```

3.149 $\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx$

Optimal result	995
Mathematica [A] (verified)	995
Rubi [A] (verified)	996
Maple [A] (verified)	998
Fricas [B] (verification not implemented)	998
Sympy [F]	999
Maxima [F]	999
Giac [A] (verification not implemented)	1000
Mupad [F(-1)]	1000
Reduce [B] (verification not implemented)	1000

Optimal result

Integrand size = 15, antiderivative size = 72

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx = \frac{-\frac{3b}{a^2} - \frac{1}{ax}}{\sqrt{a+bx}} - \frac{3b \log\left(\frac{-\sqrt{a} + \sqrt{a+bx}}{\sqrt{a} + \sqrt{a+bx}}\right)}{2a^{5/2}}$$

output $(-3*b/a^2-1/a/x)/(b*x+a)^{(1/2)}-3/2*b*\ln((-a^{(1/2)}+(b*x+a)^{(1/2)})/(a^{(1/2)}+(b*x+a)^{(1/2)}))/a^{(5/2)}$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.94

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx = -\frac{(a+bx)\left(\sqrt{a}(a+3bx) - 3bx\sqrt{a+bx}\operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)\right)}{a^{5/2}x\sqrt{(a+bx)^3}}$$

input `Integrate[1/(x^2*Sqrt[(a + b*x)^3]),x]`

output $-(((a + b*x)*(Sqrt[a]*(a + 3*b*x) - 3*b*x*Sqrt[a + b*x]*ArcTanh[Sqrt[a + b*x]/Sqrt[a]]))/(a^{(5/2)}*x*Sqrt[(a + b*x)^3]))$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.18, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2008, 52, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx \\
 & \quad \downarrow \text{2008} \\
 & \frac{(a+bx)^{3/2} \int \frac{1}{x^2 (a+bx)^{3/2}} dx}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{52} \\
 & \frac{(a+bx)^{3/2} \left(-\frac{3b \int \frac{1}{x(a+bx)^{3/2}} dx}{2a} - \frac{1}{ax\sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{61} \\
 & \frac{(a+bx)^{3/2} \left(-\frac{3b \left(\frac{\int \frac{1}{x\sqrt{a+bx}} dx}{a} + \frac{2}{a\sqrt{a+bx}} \right)}{2a} - \frac{1}{ax\sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{73} \\
 & \frac{(a+bx)^{3/2} \left(-\frac{3b \left(\frac{2 \int \frac{1}{\frac{a+bx}{b} - \frac{a}{b}} d\sqrt{a+bx}}{ab} + \frac{2}{a\sqrt{a+bx}} \right)}{2a} - \frac{1}{ax\sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}} \\
 & \quad \downarrow \text{221} \\
 & \frac{(a+bx)^{3/2} \left(-\frac{3b \left(\frac{2}{a\sqrt{a+bx}} - \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{a+bx}}{\sqrt{a}} \right)}{a^{3/2}} \right)}{2a} - \frac{1}{ax\sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}}
 \end{aligned}$$

input `Int[1/(x^2*Sqrt[(a + b*x)^3]),x]`

output `((a + b*x)^(3/2)*(-1/(a*x*Sqrt[a + b*x])) - (3*b*(2/(a*Sqrt[a + b*x]) - (2*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/a^(3/2)))/(2*a))/Sqrt[(a + b*x)^3]`

Defintions of rubi rules used

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 61 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 221 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

rule 2008 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Expon[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p), x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]`

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{(bx+a) \left(3\sqrt{bx+a} \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right) bx - 3bx\sqrt{a-a^{\frac{3}{2}}}\right)}{\sqrt{(bx+a)^3} a^{\frac{5}{2}} x}$	58
risch	$-\frac{(bx+a)^2}{a^2 x \sqrt{(bx+a)^3}} - \frac{b \left(\frac{4}{\sqrt{bx+a}} - \frac{6 \operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{\sqrt{a}} \right) (bx+a)^{\frac{3}{2}}}{2a^2 \sqrt{(bx+a)^3}}$	75

input `int(1/x^2/((b*x+a)^3)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{(b*x+a)*(3*(b*x+a)^(1/2)*\operatorname{arctanh}((b*x+a)^(1/2)/a^(1/2))*b*x-3*b*x*a^(1/2)-a^(3/2))}{((b*x+a)^3)^(1/2)/a^(5/2)/x}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(58) = 116.

Time = 0.11 (sec) , antiderivative size = 317, normalized size of antiderivative = 4.40

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx$$

$$= \frac{\left[3(b^3x^3 + 2ab^2x^2 + a^2bx)\sqrt{a} \log\left(\frac{b^2x^2 + 3abx + 2a^2 + 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}\sqrt{a}}{bx^2 + ax}\right) - 2\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3} \right]}{2(a^3b^2x^3 + 2a^4bx^2 + a^5x)}$$

$$- \frac{3(b^3x^3 + 2ab^2x^2 + a^2bx)\sqrt{-a} \arctan\left(\frac{\sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}\sqrt{-a}}{b^2x^2 + 2abx + a^2}\right) + \sqrt{b^3x^3 + 3ab^2x^2 + 3a^2bx + a^3}(3)}{a^3b^2x^3 + 2a^4bx^2 + a^5x}$$

input `integrate(1/x^2/((b*x+a)^3)^(1/2),x, algorithm="fricas")`

output

```
[1/2*(3*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)*sqrt(a)*log((b^2*x^2 + 3*a*b*x +
2*a^2 + 2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*sqrt(a))/(b*x^2 +
a*x)) - 2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*(3*a*b*x + a^2))/
(a^3*b^2*x^3 + 2*a^4*b*x^2 + a^5*x), -(3*(b^3*x^3 + 2*a*b^2*x^2 + a^2*b*x)
*sqrt(-a)*arctan(sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*sqrt(-a)/(b
^2*x^2 + 2*a*b*x + a^2)) + sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*(
3*a*b*x + a^2))/(a^3*b^2*x^3 + 2*a^4*b*x^2 + a^5*x)]
```

Sympy [F]

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx = \int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx$$

input

```
integrate(1/x**2/((b*x+a)**3)**(1/2), x)
```

output

```
Integral(1/(x**2*sqrt((a + b*x)**3)), x)
```

Maxima [F]

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx = \int \frac{1}{\sqrt{(bx+a)^3} x^2} dx$$

input

```
integrate(1/x^2/((b*x+a)^3)^(1/2), x, algorithm="maxima")
```

output

```
integrate(1/(sqrt((b*x + a)^3)*x^2), x)
```


Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.89

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx = -\frac{3b \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{\sqrt{-aa^2}} - \frac{3(bx+a)b - 2ab}{\left((bx+a)^{\frac{3}{2}} - \sqrt{bx+aa}\right)a^2}$$

input `integrate(1/x^2/((b*x+a)^3)^(1/2),x, algorithm="giac")`output `-3*b*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^2) - (3*(b*x + a)*b - 2*a*b)/(((b*x + a)^(3/2) - sqrt(b*x + a))*a^2)`**Mupad [F(-1)]**

Timed out.

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx = \int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx$$

input `int(1/(x^2*((a + b*x)^3)^(1/2)),x)`output `int(1/(x^2*((a + b*x)^3)^(1/2)), x)`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.01

$$\int \frac{1}{x^2 \sqrt{(a+bx)^3}} dx = \frac{-3\sqrt{a} \sqrt{bx+a} \log(\sqrt{bx+a} - \sqrt{a}) bx + 3\sqrt{a} \sqrt{bx+a} \log(\sqrt{bx+a} + \sqrt{a}) bx - 2a^2 - 6abx}{2\sqrt{bx+a} a^3 x}$$

input `int(1/x^2/((b*x+a)^3)^(1/2),x)`

output

```
( - 3*sqrt(a)*sqrt(a + b*x)*log(sqrt(a + b*x) - sqrt(a))*b*x + 3*sqrt(a)*s  
qrt(a + b*x)*log(sqrt(a + b*x) + sqrt(a))*b*x - 2*a**2 - 6*a*b*x)/(2*sqrt(  
a + b*x)*a**3*x)
```

3.150 $\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx$

Optimal result	1002
Mathematica [A] (verified)	1002
Rubi [A] (verified)	1003
Maple [A] (verified)	1006
Fricas [B] (verification not implemented)	1006
Sympy [F]	1007
Maxima [F]	1007
Giac [A] (verification not implemented)	1008
Mupad [F(-1)]	1008
Reduce [B] (verification not implemented)	1008

Optimal result

Integrand size = 15, antiderivative size = 91

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx = \frac{15b^2}{4a^3} - \frac{1}{2ax^2} + \frac{5b}{4a^2x} + \frac{15b^2 \log\left(\frac{-\sqrt{a} + \sqrt{a+bx}}{\sqrt{a} + \sqrt{a+bx}}\right)}{8a^{5/2}}$$

output

$$\frac{(15/4*b^2/a^3-1/2/a/x^2+5/4*b/a^2/x)/(b*x+a)^{(1/2)}+15/8*b^2*\ln((-a^{(1/2)}+(b*x+a)^{(1/2)})/(a^{(1/2)}+(b*x+a)^{(1/2)}))/a^{(5/2)}}{}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.96

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx = -\frac{(a+bx)\left(\sqrt{a}(2a^2-5abx-15b^2x^2)+15b^2x^2\sqrt{a+bx}\operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)\right)}{4a^{7/2}x^2\sqrt{(a+bx)^3}}$$

input

```
Integrate[1/(x^3*Sqrt[(a + b*x)^3]),x]
```

output

$$-1/4*((a + b*x)*(Sqrt[a]*(2*a^2 - 5*a*b*x - 15*b^2*x^2) + 15*b^2*x^2*Sqrt[a + b*x])*ArcTanh[Sqrt[a + b*x]/Sqrt[a]])/(a^(7/2)*x^2*Sqrt[(a + b*x)^3])$$
Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.24, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2008, 52, 52, 61, 73, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx$$

$$\downarrow \text{2008}$$

$$\frac{(a+bx)^{3/2} \int \frac{1}{x^3 (a+bx)^{3/2}} dx}{\sqrt{(a+bx)^3}}$$

$$\downarrow \text{52}$$

$$\frac{(a+bx)^{3/2} \left(-\frac{5b \int \frac{1}{x^2 (a+bx)^{3/2}} dx}{4a} - \frac{1}{2ax^2 \sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}}$$

$$\downarrow \text{52}$$

$$\frac{(a+bx)^{3/2} \left(-\frac{5b \left(-\frac{3b \int \frac{1}{x (a+bx)^{3/2}} dx}{2a} - \frac{1}{ax \sqrt{a+bx}} \right)}{4a} - \frac{1}{2ax^2 \sqrt{a+bx}} \right)}{\sqrt{(a+bx)^3}}$$

$$\downarrow \text{61}$$

$$\begin{array}{c}
 (a + bx)^{3/2} \left(-\frac{5b \left(\frac{3b \left(\frac{\int \frac{1}{x\sqrt{a+bx}} dx}{a} + \frac{2}{a\sqrt{a+bx}} \right)}{2a} - \frac{1}{ax\sqrt{a+bx}} \right)}{4a} - \frac{1}{2ax^2\sqrt{a+bx}} \right) \\
 \hline
 \sqrt{(a + bx)^3} \\
 \downarrow \text{73} \\
 (a + bx)^{3/2} \left(-\frac{5b \left(\frac{3b \left(\frac{2 \int \frac{1}{a+bx} \frac{d\sqrt{a+bx}}{b} - \frac{a}{ab} + \frac{2}{a\sqrt{a+bx}} \right)}{2a} - \frac{1}{ax\sqrt{a+bx}} \right)}{4a} - \frac{1}{2ax^2\sqrt{a+bx}} \right) \\
 \hline
 \sqrt{(a + bx)^3} \\
 \downarrow \text{221} \\
 (a + bx)^{3/2} \left(-\frac{5b \left(\frac{3b \left(\frac{2}{a\sqrt{a+bx}} - \frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+bx}}{\sqrt{a}}\right)}{a^{3/2}} \right)}{2a} - \frac{1}{ax\sqrt{a+bx}} \right)}{4a} - \frac{1}{2ax^2\sqrt{a+bx}} \right) \\
 \hline
 \sqrt{(a + bx)^3}
 \end{array}$$

input `Int[1/(x^3*sqrt[(a + b*x)^3]),x]`

output `((a + b*x)^(3/2)*(-1/2*1/(a*x^2*sqrt[a + b*x]) - (5*b*(-1/(a*x*sqrt[a + b*x])) - (3*b*(2/(a*sqrt[a + b*x]) - (2*ArcTanh[Sqrt[a + b*x]/sqrt[a]))/a^(3/2)))/(2*a)))/(4*a))/sqrt[(a + b*x)^3]`

Definitions of rubi rules used

- rule 52 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)} / ((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*((m + n + 2) / ((b*c - a*d)*(m + 1))) \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ \text{FractionQ}[n] \ \&\& \ \text{LtQ}[n, 0]$
- rule 61 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)} / ((b*c - a*d)*(m + 1))), x] - \text{Simp}[d*((m + n + 2) / ((b*c - a*d)*(m + 1))) \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ !(\text{LtQ}[n, -1] \ \&\& \ (\text{EqQ}[a, 0] \ || \ (\text{NeQ}[c, 0] \ \&\& \ \text{LtQ}[m - n, 0] \ \&\& \ \text{IntegerQ}[n]))) \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 73 $\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \ \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}, x]] /;$ $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 221 $\text{Int}[(a_.) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /;$ $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b]$
- rule 2008 $\text{Int}[(u_.)(Px_)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{a = \text{Rt}[\text{Coeff}[Px, x, 0], \text{Expon}[Px, x]], b = \text{Rt}[\text{Coeff}[Px, x, \text{Expon}[Px, x]], \text{Expon}[Px, x]]\}, \text{Simp}[(a + b*x)^{\text{Expon}[Px, x]} / (a + b*x)^{(\text{Expon}[Px, x]*p)} \text{Int}[u*(a + b*x)^{(\text{Expon}[Px, x]*p)}, x], x] /;$ $\text{EqQ}[Px, (a + b*x)^{\text{Expon}[Px, x]}] /;$ $!\text{IntegerQ}[p] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ \text{GtQ}[\text{Expon}[Px, x], 1] \ \&\& \ \text{NeQ}[\text{Coeff}[Px, x, 0], 0]$

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.81

method	result	size
default	$-\frac{(bx+a)\left(15\sqrt{bx+a}\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)b^2x^2-5a^{\frac{3}{2}}bx-15b^2x^2\sqrt{a+2a^{\frac{5}{2}}}\right)}{4\sqrt{(bx+a)^3}a^{\frac{7}{2}}x^2}$	74
risch	$-\frac{(bx+a)^2(-7bx+2a)}{4a^3x^2\sqrt{(bx+a)^3}} + \frac{b^2\left(\frac{16}{\sqrt{bx+a}} - \frac{30\operatorname{arctanh}\left(\frac{\sqrt{bx+a}}{\sqrt{a}}\right)}{\sqrt{a}}\right)(bx+a)^{\frac{3}{2}}}{8a^3\sqrt{(bx+a)^3}}$	85

input `int(1/x^3/((b*x+a)^3)^(1/2),x,method=_RETURNVERBOSE)`output
$$-1/4*(b*x+a)*(15*(b*x+a)^(1/2)*\operatorname{arctanh}((b*x+a)^(1/2)/a^(1/2))*b^2*x^2-5*a^(3/2)*b*x-15*b^2*x^2*a^(1/2)+2*a^(5/2))/((b*x+a)^3)^(1/2)/a^(7/2)/x^2$$
Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 172 vs. 2(72) = 144.

Time = 0.11 (sec) , antiderivative size = 355, normalized size of antiderivative = 3.90

$$\int \frac{1}{x^3\sqrt{(a+bx)^3}} dx$$

$$= \left[\frac{15(b^4x^4 + 2ab^3x^3 + a^2b^2x^2)\sqrt{a}\log\left(\frac{b^2x^2+3abx+2a^2-2\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}\sqrt{a}}{bx^2+ax}\right) + 2\sqrt{b^3x^3+3ab^2x^2+3a^2bx+a^3}}{8(a^4b^2x^4 + 2a^5bx^3 + a^6x^2)} \right]$$

input `integrate(1/x^3/((b*x+a)^3)^(1/2),x, algorithm="fricas")`

output

```
[1/8*(15*(b^4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*sqrt(a)*log((b^2*x^2 + 3*a*
b*x + 2*a^2 - 2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*sqrt(a))/(b*
x^2 + a*x)) + 2*sqrt(b^3*x^3 + 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*(15*a*b^2*x^
2 + 5*a^2*b*x - 2*a^3))/(a^4*b^2*x^4 + 2*a^5*b*x^3 + a^6*x^2), 1/4*(15*(b^
4*x^4 + 2*a*b^3*x^3 + a^2*b^2*x^2)*sqrt(-a)*arctan(sqrt(b^3*x^3 + 3*a*b^2*
x^2 + 3*a^2*b*x + a^3)*sqrt(-a)/(b^2*x^2 + 2*a*b*x + a^2)) + sqrt(b^3*x^3
+ 3*a*b^2*x^2 + 3*a^2*b*x + a^3)*(15*a*b^2*x^2 + 5*a^2*b*x - 2*a^3))/(a^4*
b^2*x^4 + 2*a^5*b*x^3 + a^6*x^2)]
```

Sympy [F]

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx = \int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx$$

input

```
integrate(1/x**3/((b*x+a)**3)**(1/2), x)
```

output

```
Integral(1/(x**3*sqrt((a + b*x)**3)), x)
```

Maxima [F]

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx = \int \frac{1}{\sqrt{(bx+a)^3} x^3} dx$$

input

```
integrate(1/x^3/((b*x+a)^3)^(1/2), x, algorithm="maxima")
```

output

```
integrate(1/(sqrt((b*x + a)^3)*x^3), x)
```


Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.88

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx = \frac{15b^2 \arctan\left(\frac{\sqrt{bx+a}}{\sqrt{-a}}\right)}{4\sqrt{-aa^3}} + \frac{2b^2}{\sqrt{bx+aa^3}} + \frac{7(bx+a)^{\frac{3}{2}}b^2 - 9\sqrt{bx+aa^3}b^2}{4a^3b^2x^2}$$

input `integrate(1/x^3/((b*x+a)^3)^(1/2),x, algorithm="giac")`

output `15/4*b^2*arctan(sqrt(b*x + a)/sqrt(-a))/(sqrt(-a)*a^3) + 2*b^2/(sqrt(b*x + a)*a^3) + 1/4*(7*(b*x + a)^(3/2)*b^2 - 9*sqrt(b*x + a)*a*b^2)/(a^3*b^2*x^2)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx = \int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx$$

input `int(1/(x^3*((a + b*x)^3)^(1/2)),x)`

output `int(1/(x^3*((a + b*x)^3)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.01

$$\int \frac{1}{x^3 \sqrt{(a+bx)^3}} dx = \frac{15\sqrt{a}\sqrt{bx+a}\log(\sqrt{bx+a}-\sqrt{a})b^2x^2 - 15\sqrt{a}\sqrt{bx+a}\log(\sqrt{bx+a}+\sqrt{a})b^2x^2 - 4a^3 + 10a^2bx + 30a^2}{8\sqrt{bx+a}a^4x^2}$$

input `int(1/x^3/((b*x+a)^3)^(1/2),x)`

output

```
(15*sqrt(a)*sqrt(a + b*x)*log(sqrt(a + b*x) - sqrt(a))*b**2*x**2 - 15*sqrt
(a)*sqrt(a + b*x)*log(sqrt(a + b*x) + sqrt(a))*b**2*x**2 - 4*a**3 + 10*a**
2*b*x + 30*a*b**2*x**2)/(8*sqrt(a + b*x)*a**4*x**2)
```

3.151
$$\int \frac{1}{x \sqrt[3]{(a + bx)^2}} dx$$

Optimal result	1010
Mathematica [A] (verified)	1010
Rubi [A] (verified)	1011
Maple [F]	1013
Fricas [B] (verification not implemented)	1013
Sympy [F]	1014
Maxima [F]	1014
Giac [B] (verification not implemented)	1015
Mupad [F(-1)]	1015
Reduce [F]	1016

Optimal result

Integrand size = 15, antiderivative size = 79

$$\int \frac{1}{x \sqrt[3]{(a + bx)^2}} dx = \frac{-\sqrt{3} \arctan\left(\frac{\sqrt{3} \sqrt[3]{a + bx}}{2 \sqrt[3]{a} + \sqrt[3]{a + bx}}\right) + \frac{3}{2} \log\left(\frac{-\sqrt[3]{a} + \sqrt[3]{a + bx}}{\sqrt[3]{x}}\right)}{\sqrt[3]{a^2}}$$

output

```
(-3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/(a^2)^(1/3)
```

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.43

$$\int \frac{1}{x \sqrt[3]{(a + bx)^2}} dx = \frac{(a + bx)^{2/3} \left(2\sqrt{3} \arctan\left(\frac{1 + 2 \sqrt[3]{a + bx}}{\sqrt[3]{a}}\right) - 2 \log\left(\sqrt[3]{a} - \sqrt[3]{a + bx}\right) + \log\left(a^{2/3} + \sqrt[3]{a} \sqrt[3]{a + bx} + (a + bx)\right) \right)}{2a^{2/3} \sqrt[3]{(a + bx)^2}}$$

input `Integrate[1/(x*((a + b*x)^2)^(1/3)),x]`

output
$$-1/2*((a + b*x)^(2/3)*(2*\text{Sqrt}[3]*\text{ArcTan}[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/\text{Sqrt}[3]) - 2*\text{Log}[a^(1/3) - (a + b*x)^(1/3)] + \text{Log}[a^(2/3) + a^(1/3)*(a + b*x)^(1/3) + (a + b*x)^(2/3)])/(a^(2/3)*((a + b*x)^2)^(1/3))$$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.23, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2008, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \sqrt[3]{(a+bx)^2}} dx$$

$$\downarrow \text{2008}$$

$$\frac{(a+bx)^{2/3} \int \frac{1}{x(a+bx)^{2/3}} dx}{\sqrt[3]{(a+bx)^2}}$$

$$\downarrow \text{69}$$

$$\frac{(a+bx)^{2/3} \left(-\frac{3 \int \frac{1}{\sqrt[3]{a}-\sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2a^{2/3}} - \frac{3 \int \frac{1}{a^{2/3}+\sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2a^{2/3}} \right)}{\sqrt[3]{(a+bx)^2}}$$

$$\downarrow \text{16}$$

$$\frac{(a+bx)^{2/3} \left(-\frac{3 \int \frac{1}{a^{2/3}+\sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} + \frac{3 \log\left(\sqrt[3]{a}-\sqrt[3]{a+bx}\right)}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right)}{\sqrt[3]{(a+bx)^2}}$$

$$\downarrow \text{1082}$$

$$\frac{(a+bx)^{2/3} \left(\frac{3 \int \frac{1}{-(a+bx)^{2/3}-3} d\left(\frac{2\sqrt[3]{a+bx}+1}{\sqrt[3]{a}}\right)}{a^{2/3}} + \frac{3 \log\left(\sqrt[3]{a}-\sqrt[3]{a+bx}\right)}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right)}{\sqrt[3]{(a+bx)^2}}$$

↓ 217

$$\frac{(a+bx)^{2/3} \left(-\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{a+bx}+1}{\sqrt[3]{a}}\right)}{a^{2/3}} + \frac{3 \log\left(\sqrt[3]{a}-\sqrt[3]{a+bx}\right)}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right)}{\sqrt[3]{(a+bx)^2}}$$

input `Int[1/(x*((a + b*x)^2)^(1/3)),x]`

output `((a + b*x)^(2/3)*(-(Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]))/a^(2/3)) - Log[x]/(2*a^(2/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)])/(2*a^(2/3)))/((a + b*x)^2)^(1/3)`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 69 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2), x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Fre
eQ[{a, b, c}, x]
```

rule 2008

```
Int[(u_)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Exp
on[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p),
x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x
] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

Maple [F]

$$\int \frac{1}{x ((bx + a)^2)^{\frac{1}{3}}} dx$$

input

```
int(1/x/((b*x+a)^2)^(1/3),x)
```

output

```
int(1/x/((b*x+a)^2)^(1/3),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 225 vs. $2(60) = 120$.

Time = 0.10 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.85

$$\int \frac{1}{x \sqrt[3]{(a+bx)^2}} dx$$

$$= \frac{2\sqrt{3}(a^2)^{\frac{1}{6}} a \arctan\left(\frac{\sqrt{3}(a^2)^{\frac{1}{6}}\left((a^2)^{\frac{1}{3}}(bx+a)+2(b^2x^2+2abx+a^2)^{\frac{1}{3}}a\right)}{3(abx+a^2)}\right) - (a^2)^{\frac{2}{3}} \log\left(\frac{(b^2x^2+2abx+a^2)^{\frac{2}{3}}a^2+(b^2x^2+2abx+a^2)^{\frac{1}{3}}a^2}{b^2x^2}\right)}{2a^2}$$

input

```
integrate(1/x/((b*x+a)^2)^(1/3),x, algorithm="fricas")
```

output

```
1/2*(2*sqrt(3)*(a^2)^(1/6)*a*arctan(1/3*sqrt(3)*(a^2)^(1/6)*((a^2)^(1/3)*(
b*x + a) + 2*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(a*b*x + a^2)) - (a^2)^(2/
3)*log(((b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a^2 + (b^2*x^2 + 2*a*b*x + a^2)^(1
/3)*(a*b*x + a^2)*(a^2)^(1/3) + (b^2*x^2 + 2*a*b*x + a^2)*(a^2)^(2/3))/(b^
2*x^2 + 2*a*b*x + a^2)) + 2*(a^2)^(2/3)*log(-((a^2)^(1/3)*(b*x + a) - (b^2
*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(b*x + a)))/a^2
```

Sympy [F]

$$\int \frac{1}{x \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{x \sqrt[3]{(a+bx)^2}} dx$$

input

```
integrate(1/x/((b*x+a)**2)**(1/3),x)
```

output

```
Integral(1/(x*((a + b*x)**2)**(1/3)), x)
```

Maxima [F]

$$\int \frac{1}{x \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{((bx+a)^2)^{\frac{1}{3}} x} dx$$

input

```
integrate(1/x/((b*x+a)^2)^(1/3),x, algorithm="maxima")
```

output

```
integrate(1/(((b*x + a)^2)^(1/3)*x), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. $2(60) = 120$.

Time = 2.97 (sec) , antiderivative size = 219, normalized size of antiderivative = 2.77

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx$$

$$= -\frac{\sqrt{3}(\operatorname{asgn}(bx+a))^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}(2(bx\operatorname{sgn}(bx+a)+\operatorname{asgn}(bx+a))^{\frac{1}{3}}+(\operatorname{asgn}(bx+a))^{\frac{1}{3}})}{3(\operatorname{asgn}(bx+a))^{\frac{1}{3}}}\right)}{\operatorname{asgn}(bx+a)}$$

$$- \frac{(\operatorname{asgn}(bx+a))^{\frac{1}{3}} \log\left((bx\operatorname{sgn}(bx+a)+\operatorname{asgn}(bx+a))^{\frac{2}{3}}+(bx\operatorname{sgn}(bx+a)+\operatorname{asgn}(bx+a))^{\frac{1}{3}}(\operatorname{asgn}(bx+a))^{\frac{1}{3}}\right)}{2\operatorname{asgn}(bx+a)}$$

$$+ \frac{(\operatorname{asgn}(bx+a))^{\frac{1}{3}} \log\left(\left|(bx\operatorname{sgn}(bx+a)+\operatorname{asgn}(bx+a))^{\frac{1}{3}}-(\operatorname{asgn}(bx+a))^{\frac{1}{3}}\right|\right)}{\operatorname{asgn}(bx+a)}$$

input `integrate(1/x/((b*x+a)^2)^(1/3),x, algorithm="giac")`

output `-sqrt(3)*(a*sgn(b*x + a))^(1/3)*arctan(1/3*sqrt(3)*(2*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(1/3))/(a*sgn(b*x + a))^(1/3))/(a*sgn(b*x + a)) - 1/2*(a*sgn(b*x + a))^(1/3)*log((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(2/3) + (b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3)*(a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(2/3))/(a*sgn(b*x + a)) + (a*sgn(b*x + a))^(1/3)*log(abs((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) - (a*sgn(b*x + a))^(1/3)))/(a*sgn(b*x + a))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{x ((a+bx)^2)^{1/3}} dx$$

input `int(1/(x*((a + b*x)^2)^(1/3)),x)`

output `int(1/(x*((a + b*x)^2)^(1/3)), x)`

Reduce [F]

$$\int \frac{1}{x \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{(b^2x^2 + 2abx + a^2)^{\frac{1}{3}} x} dx$$

input `int(1/x/((b*x+a)^2)^(1/3),x)`

output `int(1/((a**2 + 2*a*b*x + b**2*x**2)**(1/3)*x),x)`

3.152 $\int \frac{\sqrt[3]{a+bx}}{x} dx$

Optimal result	1017
Mathematica [A] (verified)	1017
Rubi [A] (verified)	1018
Maple [A] (verified)	1020
Fricas [A] (verification not implemented)	1021
Sympy [C] (verification not implemented)	1022
Maxima [A] (verification not implemented)	1023
Giac [A] (verification not implemented)	1023
Mupad [B] (verification not implemented)	1024
Reduce [B] (verification not implemented)	1024

Optimal result

Integrand size = 13, antiderivative size = 92

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = 3\sqrt[3]{a+bx} + \frac{a \left(-\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{a+bx}}{2\sqrt[3]{a} + \sqrt[3]{a+bx}} \right) + \frac{3}{2} \log \left(\frac{-\sqrt[3]{a} + \sqrt[3]{a+bx}}{\sqrt[3]{x}} \right) \right)}{\sqrt[3]{a^2}}$$

output

```
3*(b*x+a)^(1/3)+a*(-3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/(a^2)^(1/3)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.23

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = 3\sqrt[3]{a+bx} - \sqrt{3}\sqrt[3]{a} \arctan \left(\frac{1 + \frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}}}{\sqrt{3}} \right) + \sqrt[3]{a} \log \left(\sqrt[3]{a} - \sqrt[3]{a+bx} \right) - \frac{1}{2}\sqrt[3]{a} \log \left(a^{2/3} + \sqrt[3]{a}\sqrt[3]{a+bx} + (a+bx)^{2/3} \right)$$

input `Integrate[(a + b*x)^(1/3)/x,x]`

output $3*(a + b*x)^{(1/3)} - \text{Sqrt}[3]*a^{(1/3)}*\text{ArcTan}[(1 + (2*(a + b*x)^{(1/3}))/a^{(1/3)})/\text{Sqrt}[3]] + a^{(1/3)}*\text{Log}[a^{(1/3)} - (a + b*x)^{(1/3)}] - (a^{(1/3)}*\text{Log}[a^{(2/3)} + a^{(1/3)}*(a + b*x)^{(1/3)} + (a + b*x)^{(2/3)}])/2$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {60, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[3]{a+bx}}{x} dx$$

$$\downarrow 60$$

$$a \int \frac{1}{x(a+bx)^{2/3}} dx + 3\sqrt[3]{a+bx}$$

$$\downarrow 69$$

$$a \left(-\frac{3 \int \frac{1}{\sqrt[3]{a}-\sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2a^{2/3}} - \frac{3 \int \frac{1}{a^{2/3}+\sqrt[3]{a+bx}\sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2a^{2/3}} \right) + 3\sqrt[3]{a+bx}$$

$$\downarrow 16$$

$$a \left(-\frac{3 \int \frac{1}{a^{2/3}+\sqrt[3]{a+bx}\sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a}-\sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) + 3\sqrt[3]{a+bx}$$

$$\downarrow 1082$$

$$a \left(\frac{3 \int \frac{1}{-(a+bx)^{2/3}-3} d \left(\frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}} + 1 \right)}{a^{2/3}} + \frac{3 \log \left(\sqrt[3]{a} - \sqrt[3]{a+bx} \right)}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) + 3\sqrt[3]{a+bx}$$

↓ 217

$$a \left(-\frac{\sqrt{3} \arctan \left(\frac{2\sqrt[3]{a+bx}+1}{\sqrt[3]{a}} \right)}{a^{2/3}} + \frac{3 \log \left(\sqrt[3]{a} - \sqrt[3]{a+bx} \right)}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) + 3\sqrt[3]{a+bx}$$

input `Int[(a + b*x)^(1/3)/x,x]`

output `3*(a + b*x)^(1/3) + a*(-((Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]])/a^(2/3)) - Log[x]/(2*a^(2/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)])/(2*a^(2/3)))`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

```
rule 69 Int[1/((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(2/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2),
x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1
/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)],
x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.95

method	result
pseudoelliptic	$3(bx + a)^{\frac{1}{3}} + a^{\frac{1}{3}} \ln \left(-a^{\frac{1}{3}} + (bx + a)^{\frac{1}{3}} \right) - \frac{a^{\frac{1}{3}} \ln \left((bx+a)^{\frac{2}{3}} + a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}} + a^{\frac{2}{3}} \right)}{2} - a^{\frac{1}{3}} \sqrt{3} \arctan$
derivativedivides	$3(bx + a)^{\frac{1}{3}} + 3 \left(\frac{\ln \left(-a^{\frac{1}{3}} + (bx+a)^{\frac{1}{3}} \right)}{3a^{\frac{2}{3}}} - \frac{\ln \left((bx+a)^{\frac{2}{3}} + a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}} + a^{\frac{2}{3}} \right)}{6a^{\frac{2}{3}}} - \frac{\sqrt{3} \arctan \left(\frac{\sqrt{3} \left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}} + 1 \right)}{3} \right)}{3a^{\frac{2}{3}}} \right)$
default	$3(bx + a)^{\frac{1}{3}} + 3 \left(\frac{\ln \left(-a^{\frac{1}{3}} + (bx+a)^{\frac{1}{3}} \right)}{3a^{\frac{2}{3}}} - \frac{\ln \left((bx+a)^{\frac{2}{3}} + a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}} + a^{\frac{2}{3}} \right)}{6a^{\frac{2}{3}}} - \frac{\sqrt{3} \arctan \left(\frac{\sqrt{3} \left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}} + 1 \right)}{3} \right)}{3a^{\frac{2}{3}}} \right)$

input `int((b*x+a)^(1/3)/x,x,method=_RETURNVERBOSE)`

output `3*(b*x+a)^(1/3)+a^(1/3)*ln(-a^(1/3)+(b*x+a)^(1/3))-1/2*a^(1/3)*ln((b*x+a)^(2/3)+a^(1/3)*(b*x+a)^(1/3)+a^(2/3))-a^(1/3)*3^(1/2)*arctan(2/3*3^(1/2)/a^(1/3)*(b*x+a)^(1/3)+1/3*3^(1/2))`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.99

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = -\sqrt{3}a^{\frac{1}{3}} \arctan\left(\frac{2\sqrt{3}(bx+a)^{\frac{1}{3}}a^{\frac{2}{3}} + \sqrt{3}a}{3a}\right) - \frac{1}{2}a^{\frac{1}{3}} \log\left((bx+a)^{\frac{2}{3}} + (bx+a)^{\frac{1}{3}}a^{\frac{1}{3}} + a^{\frac{2}{3}}\right) + a^{\frac{1}{3}} \log\left((bx+a)^{\frac{1}{3}} - a^{\frac{1}{3}}\right) + 3(bx+a)^{\frac{1}{3}}$$

input `integrate((b*x+a)^(1/3)/x,x, algorithm="fricas")`

output `-sqrt(3)*a^(1/3)*arctan(1/3*(2*sqrt(3)*(b*x + a)^(1/3)*a^(2/3) + sqrt(3)*a)/a) - 1/2*a^(1/3)*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3)) + a^(1/3)*log((b*x + a)^(1/3) - a^(1/3)) + 3*(b*x + a)^(1/3)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.21 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.96

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = \frac{4\sqrt[3]{a} \log\left(1 - \frac{\sqrt[3]{b} \sqrt[3]{\frac{a}{b} + x}}{\sqrt[3]{a}}\right) \Gamma\left(\frac{4}{3}\right)}{3\Gamma\left(\frac{7}{3}\right)} + \frac{4\sqrt[3]{a} e^{-\frac{2i\pi}{3}} \log\left(1 - \frac{\sqrt[3]{b} \sqrt[3]{\frac{a}{b} + x} e^{\frac{2i\pi}{3}}}{\sqrt[3]{a}}\right) \Gamma\left(\frac{4}{3}\right)}{3\Gamma\left(\frac{7}{3}\right)} + \frac{4\sqrt[3]{a} e^{\frac{2i\pi}{3}} \log\left(1 - \frac{\sqrt[3]{b} \sqrt[3]{\frac{a}{b} + x} e^{\frac{4i\pi}{3}}}{\sqrt[3]{a}}\right) \Gamma\left(\frac{4}{3}\right)}{3\Gamma\left(\frac{7}{3}\right)} + \frac{4\sqrt[3]{b} \sqrt[3]{\frac{a}{b} + x} \Gamma\left(\frac{4}{3}\right)}{\Gamma\left(\frac{7}{3}\right)}$$

input `integrate((b*x+a)**(1/3)/x,x)`

output `4*a**(1/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)/a**(1/3))*gamma(4/3)/(3*gamma(7/3)) + 4*a**(1/3)*exp(-2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)*exp_polar(2*I*pi/3)/a**(1/3))*gamma(4/3)/(3*gamma(7/3)) + 4*a**(1/3)*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)*exp_polar(4*I*pi/3)/a**(1/3))*gamma(4/3)/(3*gamma(7/3)) + 4*b**(1/3)*(a/b + x)**(1/3)*gamma(4/3)/gamma(7/3)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.93

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = -\sqrt{3}a^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}}\right) - \frac{1}{2}a^{\frac{1}{3}} \log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}\right) + a^{\frac{1}{3}} \log\left((bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}\right) + 3(bx+a)^{\frac{1}{3}}$$

input `integrate((b*x+a)^(1/3)/x,x, algorithm="maxima")`output `-sqrt(3)*a^(1/3)*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3)) - 1/2*a^(1/3)*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3)) + a^(1/3)*log((b*x + a)^(1/3) - a^(1/3)) + 3*(b*x + a)^(1/3)`**Giac [A] (verification not implemented)**

Time = 0.33 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.95

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = -\sqrt{3}a^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}}\right) - \frac{1}{2}a^{\frac{1}{3}} \log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}\right) + a^{\frac{1}{3}} \log\left(\left|(bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}\right|\right) + 3(bx+a)^{\frac{1}{3}}$$

input `integrate((b*x+a)^(1/3)/x,x, algorithm="giac")`output `-sqrt(3)*a^(1/3)*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3)) - 1/2*a^(1/3)*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3)) + a^(1/3)*log(abs((b*x + a)^(1/3) - a^(1/3))) + 3*(b*x + a)^(1/3)`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.16

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = a^{1/3} \ln \left(9a(a+bx)^{1/3} - 9a^{4/3} \right) + 3(a+bx)^{1/3} + \frac{a^{1/3} \ln \left(9a(a+bx)^{1/3} - \frac{9a^{4/3}(-1+\sqrt{3}i)}{2} \right) (-1+\sqrt{3}i) - a^{1/3} \ln \left(9a(a+bx)^{1/3} - \frac{9a^{4/3}(1+\sqrt{3}i)}{2} \right) (1+\sqrt{3}i)}{2}$$

input `int((a + b*x)^(1/3)/x,x)`output `a^(1/3)*log(9*a*(a + b*x)^(1/3) - 9*a^(4/3)) + 3*(a + b*x)^(1/3) + (a^(1/3))*log(9*a*(a + b*x)^(1/3) - (9*a^(4/3)*(3^(1/2)*1i - 1))/2)*(3^(1/2)*1i - 1)/2 - (a^(1/3)*log(9*a*(a + b*x)^(1/3) + (9*a^(4/3)*(3^(1/2)*1i + 1))/2)*(3^(1/2)*1i + 1)/2`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.73

$$\int \frac{\sqrt[3]{a+bx}}{x} dx = \frac{2\sqrt{3} \operatorname{atan} \left(\frac{2(bx+a)^{1/6} + a^{1/6}}{a^{1/6}\sqrt{3}} \right) a - 2\sqrt{3} \operatorname{atan} \left(\frac{2(bx+a)^{1/6} - a^{1/6}}{a^{1/6}\sqrt{3}} \right) a + 6a^{2/3}(bx+a)^{1/3} + 2 \log \left((bx+a)^{1/6} + a^{1/6} \right) a + 2 \log \left((bx+a)^{1/6} - a^{1/6} \right) a}{2}$$

input `int((b*x+a)^(1/3)/x,x)`output `(2*sqrt(3)*atan((2*(a + b*x)**(1/6) + a**(1/6))/(a**(1/6)*sqrt(3)))*a - 2*sqrt(3)*atan((2*(a + b*x)**(1/6) - a**(1/6))/(a**(1/6)*sqrt(3)))*a + 6*a**(2/3)*(a + b*x)**(1/3) + 2*log((a + b*x)**(1/6) + a**(1/6))*a + 2*log((a + b*x)**(1/6) - a**(1/6))*a - log(-a**(1/6)*(a + b*x)**(1/6) + (a + b*x)**(1/3) + a**(1/3))*a - log(a**(1/6)*(a + b*x)**(1/6) + (a + b*x)**(1/3) + a**(1/3))*a)/(2*a**(2/3))`

3.153 $\int \frac{\sqrt[3]{a+bx}}{x^2} dx$

Optimal result	1025
Mathematica [A] (verified)	1025
Rubi [A] (verified)	1026
Maple [A] (verified)	1028
Fricas [A] (verification not implemented)	1029
Sympy [C] (verification not implemented)	1029
Maxima [A] (verification not implemented)	1030
Giac [A] (verification not implemented)	1031
Mupad [B] (verification not implemented)	1031
Reduce [B] (verification not implemented)	1032

Optimal result

Integrand size = 13, antiderivative size = 122

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx = \frac{b\sqrt[3]{a+bx}}{a} + \frac{(-a-bx)\sqrt[3]{a+bx}}{ax} + \frac{b\left(-\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{a+bx}}{2\sqrt[3]{a}+\sqrt[3]{a+bx}}\right) + \frac{3}{2} \log\left(\frac{-\sqrt[3]{a}+\sqrt[3]{a+bx}}{\sqrt[3]{x}}\right)\right)}{3\sqrt[3]{a^2}}$$

```
output b*(b*x+a)^(1/3)/a+(-b*x-a)*(b*x+a)^(1/3)/a/x+1/3*b*(-3^(1/2)*arctan(3^(1/2)
)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))
/x^(1/3))/(a^2)^(1/3)
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.98

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx = \frac{6a^{2/3}\sqrt[3]{a+bx} + 2\sqrt{3}bx \arctan\left(\frac{1+2\sqrt[3]{a+bx}}{\sqrt[3]{a}}\right) - 2bx \log\left(\sqrt[3]{a} - \sqrt[3]{a+bx}\right) + bx \log\left(a^{2/3} + \sqrt[3]{a}\sqrt[3]{a+bx}\right)}{6a^{2/3}x}$$

input `Integrate[(a + b*x)^(1/3)/x^2,x]`

output
$$-1/6*(6*a^{2/3}*(a + b*x)^{1/3} + 2*\text{Sqrt}[3]*b*x*\text{ArcTan}[(1 + (2*(a + b*x)^{1/3}))/a^{1/3}]/\text{Sqrt}[3]] - 2*b*x*\text{Log}[a^{1/3} - (a + b*x)^{1/3}] + b*x*\text{Log}[a^{2/3} + a^{1/3}*(a + b*x)^{1/3} + (a + b*x)^{2/3}]/(a^{2/3}*x)$$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.79, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {51, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx$$

$$\downarrow 51$$

$$\frac{1}{3}b \int \frac{1}{x(a+bx)^{2/3}} dx - \frac{\sqrt[3]{a+bx}}{x}$$

$$\downarrow 69$$

$$\frac{1}{3}b \left(-\frac{3 \int \frac{1}{\sqrt[3]{a}-\sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2a^{2/3}} - \frac{3 \int \frac{1}{a^{2/3}+\sqrt[3]{a+bx}\sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2a^{2/3}} \right) -$$

$$\frac{\sqrt[3]{a+bx}}{x}$$

$$\downarrow 16$$

$$\frac{1}{3}b \left(-\frac{3 \int \frac{1}{a^{2/3}+\sqrt[3]{a+bx}\sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a}-\sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) -$$

$$\frac{\sqrt[3]{a+bx}}{x}$$

$$\downarrow 1082$$

$$\frac{1}{3}b \left(\frac{3 \int \frac{1}{-(a+bx)^{2/3}-3} d\left(\frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}} + 1\right)}{a^{2/3}} + \frac{3 \log\left(\sqrt[3]{a} - \sqrt[3]{a+bx}\right)}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) - \frac{\sqrt[3]{a+bx}}{x}$$

↓ 217

$$\frac{1}{3}b \left(-\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{a+bx} + 1}{\sqrt[3]{a}}\right)}{a^{2/3}} + \frac{3 \log\left(\sqrt[3]{a} - \sqrt[3]{a+bx}\right)}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) - \frac{\sqrt[3]{a+bx}}{x}$$

input `Int[(a + b*x)^(1/3)/x^2,x]`

output `-((a + b*x)^(1/3)/x) + (b*(-((Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))]/a^(1/3))/Sqrt[3]))/a^(2/3)) - Log[x]/(2*a^(2/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)]/(2*a^(2/3))))/3`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1)))] Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 69 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2), x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

```
rule 217 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.78

method	result
derivativedivides	$3b \left(-\frac{(bx+a)^{\frac{1}{3}}}{3bx} + \frac{\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{9a^{\frac{2}{3}}} - \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{18a^{\frac{2}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{3}\right)}{9a^{\frac{2}{3}}}$
default	$3b \left(-\frac{(bx+a)^{\frac{1}{3}}}{3bx} + \frac{\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{9a^{\frac{2}{3}}} - \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{18a^{\frac{2}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{3}\right)}{9a^{\frac{2}{3}}}$
pseudoelliptic	$\frac{-\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{3}\right) bx + \ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right) bx - \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right) bx}{2} - 3(bx+a)^{\frac{1}{3}} a^{\frac{2}{3}}}{3a^{\frac{2}{3}} x}$

```
input int((b*x+a)^(1/3)/x^2,x,method=_RETURNVERBOSE)
```

```
output 3*b*(-1/3*(b*x+a)^(1/3)/b/x+1/9/a^(2/3)*ln(-a^(1/3)+(b*x+a)^(1/3))-1/18/a^(2/3)*ln((b*x+a)^(2/3)+a^(1/3)*(b*x+a)^(1/3)+a^(2/3))-1/9/a^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/a^(1/3)*(b*x+a)^(1/3)+1)))
```

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.11

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx = \frac{6 \sqrt{\frac{1}{3}} (a^2)^{\frac{1}{6}} abx \arctan \left(\frac{\sqrt{\frac{1}{3}} (a^2)^{\frac{1}{6}} \left((a^2)^{\frac{1}{3}} a + 2 (a^2)^{\frac{2}{3}} (bx+a)^{\frac{1}{3}} \right)}{a^2} \right) + (a^2)^{\frac{2}{3}} bx \log \left((bx+a)^{\frac{2}{3}} a + (a^2)^{\frac{1}{3}} a + (a^2)^{\frac{2}{3}} (bx+a)^{\frac{1}{3}} \right)}{6 a^2 x}$$

input `integrate((b*x+a)^(1/3)/x^2,x, algorithm="fricas")`

output `-1/6*(6*sqrt(1/3)*(a^2)^(1/6)*a*b*x*arctan(sqrt(1/3)*(a^2)^(1/6)*((a^2)^(1/3)*a + 2*(a^2)^(2/3)*(b*x + a)^(1/3))/a^2) + (a^2)^(2/3)*b*x*log((b*x + a)^(2/3)*a + (a^2)^(1/3)*a + (a^2)^(2/3)*(b*x + a)^(1/3)) - 2*(a^2)^(2/3)*b*x*log((b*x + a)^(1/3)*a - (a^2)^(2/3)) + 6*(b*x + a)^(1/3)*a^2/(a^2*x)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.32 (sec) , antiderivative size = 643, normalized size of antiderivative = 5.27

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx = \text{Too large to display}$$

input `integrate((b*x+a)**(1/3)/x**2,x)`

output

```

4*a**(7/3)*b*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)/a**(1/3))*gam
ma(4/3)/(9*a**3*exp(2*I*pi/3)*gamma(7/3) - 9*a**2*b*(a/b + x)*exp(2*I*pi/3
)*gamma(7/3)) + 4*a**(7/3)*b*log(1 - b**(1/3)*(a/b + x)**(1/3)*exp_polar(2
*I*pi/3)/a**(1/3))*gamma(4/3)/(9*a**3*exp(2*I*pi/3)*gamma(7/3) - 9*a**2*b*
(a/b + x)*exp(2*I*pi/3)*gamma(7/3)) + 4*a**(7/3)*b*exp(-2*I*pi/3)*log(1 -
b**(1/3)*(a/b + x)**(1/3)*exp_polar(4*I*pi/3)/a**(1/3))*gamma(4/3)/(9*a**3
*exp(2*I*pi/3)*gamma(7/3) - 9*a**2*b*(a/b + x)*exp(2*I*pi/3)*gamma(7/3)) -
4*a**(4/3)*b**2*(a/b + x)*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)
/a**(1/3))*gamma(4/3)/(9*a**3*exp(2*I*pi/3)*gamma(7/3) - 9*a**2*b*(a/b + x
)*exp(2*I*pi/3)*gamma(7/3)) - 4*a**(4/3)*b**2*(a/b + x)*log(1 - b**(1/3)*(
a/b + x)**(1/3)*exp_polar(2*I*pi/3)/a**(1/3))*gamma(4/3)/(9*a**3*exp(2*I*p
i/3)*gamma(7/3) - 9*a**2*b*(a/b + x)*exp(2*I*pi/3)*gamma(7/3)) - 4*a**(4/3
)*b**2*(a/b + x)*exp(-2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)*exp_pola
r(4*I*pi/3)/a**(1/3))*gamma(4/3)/(9*a**3*exp(2*I*pi/3)*gamma(7/3) - 9*a**2
*b*(a/b + x)*exp(2*I*pi/3)*gamma(7/3)) + 12*a**2*b**(4/3)*(a/b + x)**(1/3)
*exp(2*I*pi/3)*gamma(4/3)/(9*a**3*exp(2*I*pi/3)*gamma(7/3) - 9*a**2*b*(a/b
+ x)*exp(2*I*pi/3)*gamma(7/3))

```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.76

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx = -\frac{\sqrt{3}b \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{3a^{\frac{2}{3}}}
- \frac{b \log\left((bx+a)^{\frac{2}{3}} + (bx+a)^{\frac{1}{3}}a^{\frac{1}{3}} + a^{\frac{2}{3}}\right)}{6a^{\frac{2}{3}}}
+ \frac{b \log\left((bx+a)^{\frac{1}{3}} - a^{\frac{1}{3}}\right)}{3a^{\frac{2}{3}}} - \frac{(bx+a)^{\frac{1}{3}}}{x}$$

input

```
integrate((b*x+a)^(1/3)/x^2,x, algorithm="maxima")
```

output

```

-1/3*sqrt(3)*b*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3))/a
^(2/3) - 1/6*b*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3))/a
^(2/3) + 1/3*b*log((b*x + a)^(1/3) - a^(1/3))/a^(2/3) - (b*x + a)^(1/3)/x

```

Giac [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx = -\frac{1}{6} b \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{a^{\frac{2}{3}}} + \frac{\log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{a^{\frac{2}{3}}} - \frac{2\log\left(\left|(bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}\right|\right)}{a^{\frac{2}{3}}}\right)$$

input `integrate((b*x+a)^(1/3)/x^2,x, algorithm="giac")`output `-1/6*b*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3))/a^(2/3) + log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3))/a^(2/3) - 2*log(abs((b*x + a)^(1/3) - a^(1/3)))/a^(2/3) + 6*(b*x + a)^(1/3)/(b*x))`**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.96

$$\int \frac{\sqrt[3]{a+bx}}{x^2} dx = \frac{b \ln\left(3b(a+bx)^{1/3} - 3a^{1/3}b\right)}{3a^{2/3}} - \frac{(a+bx)^{1/3}}{x} - \frac{\ln\left(\frac{3a^{1/3}(b-\sqrt{3}bi)}{2} + 3b(a+bx)^{1/3}\right)(b-\sqrt{3}bi)}{6a^{2/3}} - \frac{\ln\left(\frac{3a^{1/3}(b+\sqrt{3}bi)}{2} + 3b(a+bx)^{1/3}\right)(b+\sqrt{3}bi)}{6a^{2/3}}$$

input `int((a + b*x)^(1/3)/x^2,x)`

output

$$\frac{(b \log(3b(a + bx)^{1/3} - 3a^{1/3}b)) / (3a^{2/3}) - (a + bx)^{1/3} / x - (\log((3a^{1/3})(b - 3^{1/2}b^{1/2})) / 2 + 3b(a + bx)^{1/3})(b - 3^{1/2}b^{1/2}) / (6a^{2/3}) - (\log((3a^{1/3})(b + 3^{1/2}b^{1/2})) / 2 + 3b(a + bx)^{1/3})(b + 3^{1/2}b^{1/2}) / (6a^{2/3})}{1}$$
Reduce [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.38

$$\int \frac{\sqrt[3]{a + bx}}{x^2} dx$$

$$= \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}+a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) bx - 2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}-a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) bx - 6a^{\frac{2}{3}}(bx+a)^{\frac{1}{3}} + 2\log\left((bx+a)^{\frac{1}{6}}+a^{\frac{1}{6}}\right) bx + \dots}{1}$$

input

$$\operatorname{int}((b*x+a)^{(1/3)}/x^2,x)$$

output

$$\frac{(2\sqrt{3})\operatorname{atan}\left(\frac{2(a + bx)^{1/6} + a^{1/6}}{a^{1/6}\sqrt{3}}\right)*bx - 2\sqrt{3}\operatorname{atan}\left(\frac{2(a + bx)^{1/6} - a^{1/6}}{a^{1/6}\sqrt{3}}\right)*bx - 6a^{2/3}(a + bx)^{1/3} + 2\log((a + bx)^{1/6} + a^{1/6})*bx + 2\log((a + bx)^{1/6} - a^{1/6})*bx - \log(-a^{1/6}(a + bx)^{1/6} + (a + bx)^{1/3} + a^{1/3})*bx - \log(a^{1/6}(a + bx)^{1/6} + (a + bx)^{1/3} + a^{1/3})*bx}{6a^{2/3}x}$$

3.154 $\int \frac{\sqrt[3]{a+bx}}{x^3} dx$

Optimal result	1033
Mathematica [A] (verified)	1033
Rubi [A] (verified)	1034
Maple [A] (verified)	1037
Fricas [A] (verification not implemented)	1038
Sympy [C] (verification not implemented)	1038
Maxima [A] (verification not implemented)	1039
Giac [A] (verification not implemented)	1040
Mupad [B] (verification not implemented)	1041
Reduce [B] (verification not implemented)	1041

Optimal result

Integrand size = 13, antiderivative size = 140

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx = -\frac{b^2 \sqrt[3]{a+bx}}{3a^2} + \left(-\frac{1}{2ax^2} + \frac{b}{3a^2x}\right) (a+bx)^{4/3} - \frac{b^2 \left(-\sqrt{3} \arctan\left(\frac{\sqrt{3} \sqrt[3]{a+bx}}{2\sqrt[3]{a} + \sqrt[3]{a+bx}}\right) + \frac{3}{2} \log\left(\frac{-\sqrt[3]{a} + \sqrt[3]{a+bx}}{\sqrt[3]{x}}\right)\right)}{9a\sqrt[3]{a^2}}$$

```
output -1/3*b^2*(b*x+a)^(1/3)/a^2+(-1/2/a/x^2+1/3*b/a^2/x)*(b*x+a)^(4/3)-1/9*b^2*
(-3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/a/(a^2)^(1/3)
```

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx = -\frac{3a^{2/3} \sqrt[3]{a+bx} (3a+bx)}{x^2} + 2\sqrt{3}b^2 \arctan\left(\frac{1+2\sqrt[3]{a+bx}}{\sqrt[3]{a}}\right) - 2b^2 \log\left(\sqrt[3]{a} - \sqrt[3]{a+bx}\right) + b^2 \log\left(a^{2/3} + \sqrt[3]{a}\right)$$

$18a^{5/3}$

input `Integrate[(a + b*x)^(1/3)/x^3,x]`

output `((-3*a^(2/3)*(a + b*x)^(1/3)*(3*a + b*x))/x^2 + 2*sqrt[3]*b^2*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3))/sqrt[3]] - 2*b^2*Log[a^(1/3) - (a + b*x)^(1/3)] + b^2*Log[a^(2/3) + a^(1/3)*(a + b*x)^(1/3) + (a + b*x)^(2/3)])/(18*a^(5/3))`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.89, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {51, 52, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt[3]{a+bx}}{x^3} dx \\
 & \quad \downarrow \text{51} \\
 & \frac{1}{6}b \int \frac{1}{x^2(a+bx)^{2/3}} dx - \frac{\sqrt[3]{a+bx}}{2x^2} \\
 & \quad \downarrow \text{52} \\
 & \frac{1}{6}b \left(-\frac{2b \int \frac{1}{x(a+bx)^{2/3}} dx}{3a} - \frac{\sqrt[3]{a+bx}}{ax} \right) - \frac{\sqrt[3]{a+bx}}{2x^2} \\
 & \quad \downarrow \text{69} \\
 & \frac{1}{6}b \left(\frac{2b \left(-\frac{3 \int \frac{1}{\sqrt[3]{a}-\sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2a^{2/3}} - \frac{3 \int \frac{1}{a^{2/3}+\sqrt[3]{a+bx}} \sqrt[3]{a+(a+bx)^{2/3}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2a^{2/3}} \right)}{3a} - \frac{\sqrt[3]{a+bx}}{ax} \right) - \frac{\sqrt[3]{a+bx}}{2x^2}
 \end{aligned}$$

$$\downarrow 16$$

$$\frac{1}{6}b \left(\frac{2b \left(\frac{{}_3F_1 \left(\frac{1}{a^{2/3} + \sqrt[3]{a+bx}} \right)}{\sqrt[3]{a} + (a+bx)^{2/3}} d \sqrt[3]{a+bx} \right)}{2\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right)}{3a} - \frac{\sqrt[3]{a+bx}}{ax} \right)$$

$$\frac{\sqrt[3]{a+bx}}{2x^2}$$

$$\downarrow 1082$$

$$\frac{1}{6}b \left(\frac{2b \left(\frac{{}_3F_1 \left(\frac{1}{-(a+bx)^{2/3} - 3} \right) d \left(\frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}} + 1 \right)}{a^{2/3}} \right)}{3a} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) - \frac{\sqrt[3]{a+bx}}{ax}$$

$$\frac{\sqrt[3]{a+bx}}{2x^2}$$

$$\downarrow 217$$

$$\frac{1}{6}b \left(\frac{2b \left(\frac{\sqrt{3} \arctan \left(\frac{2\sqrt[3]{a+bx} + 1}{\sqrt[3]{a}} \right)}{a^{2/3}} \right)}{3a} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) - \frac{\sqrt[3]{a+bx}}{ax}$$

$$\frac{\sqrt[3]{a+bx}}{2x^2}$$

input `Int[(a + b*x)^(1/3)/x^3,x]`

output `-1/2*(a + b*x)^(1/3)/x^2 + (b*(-((a + b*x)^(1/3)/(a*x)) - (2*b*(-((Sqrt[3] *ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3))/Sqrt[3]])/a^(2/3)) - Log[x]/(2*a^(2/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)]/(2*a^(2/3))))/(3*a)))/6`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 51 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Simp[d*(n/(b*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && GtQ[n, 0]`

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1))) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 69 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2), x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 1082

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])]; Fre
eQ[{a, b, c}, x]
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 118, normalized size of antiderivative = 0.84

method	result
derivativedivides	$3b^2 \left(-\frac{\frac{(bx+a)^{\frac{4}{3}}}{18a} + \frac{(bx+a)^{\frac{1}{3}}}{9}}{b^2x^2} - \frac{\ln\left(-a^{\frac{1}{3}} + (bx+a)^{\frac{1}{3}}\right)}{3a^{\frac{2}{3}}} - \frac{\ln\left((bx+a)^{\frac{2}{3}} + a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}} + a^{\frac{2}{3}}\right)}{6a^{\frac{2}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}} + 1\right)}{\frac{1}{3}}\right)}{3a^{\frac{2}{3}}} \right)$
default	$3b^2 \left(-\frac{\frac{(bx+a)^{\frac{4}{3}}}{18a} + \frac{(bx+a)^{\frac{1}{3}}}{9}}{b^2x^2} - \frac{\ln\left(-a^{\frac{1}{3}} + (bx+a)^{\frac{1}{3}}\right)}{3a^{\frac{2}{3}}} - \frac{\ln\left((bx+a)^{\frac{2}{3}} + a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}} + a^{\frac{2}{3}}\right)}{6a^{\frac{2}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}} + 1\right)}{\frac{1}{3}}\right)}{3a^{\frac{2}{3}}} \right)$
pseudoelliptic	$\frac{2b^2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2(bx+a)^{\frac{1}{3}} + a^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}}\right) x^2 - 2b^2 \ln\left(-a^{\frac{1}{3}} + (bx+a)^{\frac{1}{3}}\right) x^2 + b^2 \ln\left((bx+a)^{\frac{2}{3}} + a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}} + a^{\frac{2}{3}}\right) x^2 - 3bx(bx+a)^{\frac{1}{3}}}{18a^{\frac{5}{3}}x^2}$

input

```
int((b*x+a)^(1/3)/x^3,x,method=_RETURNVERBOSE)
```

output

```
3*b^2*(-(1/18/a*(b*x+a)^(4/3)+1/9*(b*x+a)^(1/3))/b^2/x^2-1/9/a*(1/3/a^(2/3)
)*ln(-a^(1/3)+(b*x+a)^(1/3))-1/6/a^(2/3)*ln((b*x+a)^(2/3)+a^(1/3)*(b*x+a)^(
1/3)+a^(2/3))-1/3/a^(2/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/a^(1/3)*(b*x+a)^(
1/3)+1))))
```

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx$$

$$= \frac{6\sqrt{\frac{1}{3}}ab^2x^2\sqrt{-(-a^2)^{\frac{1}{3}}}\arctan\left(-\frac{\sqrt{\frac{1}{3}}\left((-a^2)^{\frac{1}{3}}a-2(-a^2)^{\frac{2}{3}}(bx+a)^{\frac{1}{3}}\right)\sqrt{-(-a^2)^{\frac{1}{3}}}}{a^2}\right)+(-a^2)^{\frac{2}{3}}b^2x^2\log\left((bx+a)^{\frac{2}{3}}a\right)}{18}$$

input `integrate((b*x+a)^(1/3)/x^3,x, algorithm="fricas")`

output `1/18*(6*sqrt(1/3)*a*b^2*x^2*sqrt(-(-a^2)^(1/3))*arctan(-sqrt(1/3)*((-a^2)^(1/3)*a - 2*(-a^2)^(2/3)*(b*x + a)^(1/3))*sqrt(-(-a^2)^(1/3))/a^2) + (-a^2)^(2/3)*b^2*x^2*log((b*x + a)^(2/3)*a - (-a^2)^(1/3)*a + (-a^2)^(2/3)*(b*x + a)^(1/3)) - 2*(-a^2)^(2/3)*b^2*x^2*log((b*x + a)^(1/3)*a - (-a^2)^(2/3)) - 3*(a^2*b*x + 3*a^3)*(b*x + a)^(1/3)/(a^3*x^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.99 (sec) , antiderivative size = 2266, normalized size of antiderivative = 16.19

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx = \text{Too large to display}$$

input `integrate((b*x+a)**(1/3)/x**3,x)`

output

```

-4*a**(16/3)*b**2*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)/a**(1/3)
)*gamma(4/3)/(27*a**7*exp(2*I*pi/3)*gamma(7/3) - 81*a**6*b*(a/b + x)*exp(2
*I*pi/3)*gamma(7/3) + 81*a**5*b**2*(a/b + x)**2*exp(2*I*pi/3)*gamma(7/3) -
27*a**4*b**3*(a/b + x)**3*exp(2*I*pi/3)*gamma(7/3)) - 4*a**(16/3)*b**2*lo
g(1 - b**(1/3)*(a/b + x)**(1/3)*exp_polar(2*I*pi/3)/a**(1/3))*gamma(4/3)/(
27*a**7*exp(2*I*pi/3)*gamma(7/3) - 81*a**6*b*(a/b + x)*exp(2*I*pi/3)*gamma
(7/3) + 81*a**5*b**2*(a/b + x)**2*exp(2*I*pi/3)*gamma(7/3) - 27*a**4*b**3*
(a/b + x)**3*exp(2*I*pi/3)*gamma(7/3) - 4*a**(16/3)*b**2*exp(-2*I*pi/3)*l
og(1 - b**(1/3)*(a/b + x)**(1/3)*exp_polar(4*I*pi/3)/a**(1/3))*gamma(4/3)/
(27*a**7*exp(2*I*pi/3)*gamma(7/3) - 81*a**6*b*(a/b + x)*exp(2*I*pi/3)*gamm
a(7/3) + 81*a**5*b**2*(a/b + x)**2*exp(2*I*pi/3)*gamma(7/3) - 27*a**4*b**3
*(a/b + x)**3*exp(2*I*pi/3)*gamma(7/3)) + 12*a**(13/3)*b**3*(a/b + x)*exp(
2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)/a**(1/3))*gamma(4/3)/(27*a**7*
exp(2*I*pi/3)*gamma(7/3) - 81*a**6*b*(a/b + x)*exp(2*I*pi/3)*gamma(7/3) +
81*a**5*b**2*(a/b + x)**2*exp(2*I*pi/3)*gamma(7/3) - 27*a**4*b**3*(a/b + x
)**3*exp(2*I*pi/3)*gamma(7/3)) + 12*a**(13/3)*b**3*(a/b + x)*log(1 - b**(1
/3)*(a/b + x)**(1/3)*exp_polar(2*I*pi/3)/a**(1/3))*gamma(4/3)/(27*a**7*exp
(2*I*pi/3)*gamma(7/3) - 81*a**6*b*(a/b + x)*exp(2*I*pi/3)*gamma(7/3) + 81*
a**5*b**2*(a/b + x)**2*exp(2*I*pi/3)*gamma(7/3) - 27*a**4*b**3*(a/b + x)**
3*exp(2*I*pi/3)*gamma(7/3)) + 12*a**(13/3)*b**3*(a/b + x)*exp(-2*I*pi/3...

```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.99

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx = \frac{\sqrt{3}b^2 \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{9a^{\frac{5}{3}}}$$

$$+ \frac{b^2 \log\left((bx+a)^{\frac{2}{3}} + (bx+a)^{\frac{1}{3}}a^{\frac{1}{3}} + a^{\frac{2}{3}}\right)}{18a^{\frac{5}{3}}}$$

$$- \frac{b^2 \log\left((bx+a)^{\frac{1}{3}} - a^{\frac{1}{3}}\right)}{9a^{\frac{5}{3}}} - \frac{(bx+a)^{\frac{4}{3}}b^2 + 2(bx+a)^{\frac{1}{3}}ab^2}{6((bx+a)^2a - 2(bx+a)a^2 + a^3)}$$

input

```
integrate((b*x+a)^(1/3)/x^3,x, algorithm="maxima")
```


output

$$\frac{1}{9}\sqrt{3}b^2\arctan\left(\frac{1}{3}\sqrt{3}\frac{2(bx+a)^{1/3}+a^{1/3}}{a^{1/3}}\right)/a^{5/3} + \frac{1}{18}b^2\log\left(\frac{(bx+a)^{2/3}+(bx+a)^{1/3}a^{1/3}+a^{2/3}}{a^{5/3}}\right) - \frac{1}{9}b^2\log\left(\frac{(bx+a)^{1/3}-a^{1/3}}{a^{5/3}}\right) - \frac{1}{6}\frac{(bx+a)^{4/3}b^2+2(bx+a)^{1/3}ab^2}{(bx+a)^2a-2(bx+a)a^2+a^3}$$
Giac [A] (verification not implemented)

Time = 0.33 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx$$

$$= \frac{2\sqrt{3}b^3 \arctan\left(\frac{\sqrt{3}\left(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}}\right)}{a^{\frac{5}{3}}} + \frac{b^3 \log\left(\frac{(bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}}{a^{\frac{5}{3}}}\right)}{a^{\frac{5}{3}}} - \frac{2b^3 \log\left(\left|\frac{(bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}}{a^{\frac{5}{3}}}\right|\right)}{a^{\frac{5}{3}}} - \frac{3\left((bx+a)^{\frac{4}{3}}b^3+2(bx+a)^{\frac{1}{3}}ab^3\right)}{ab^2x^2}$$

$18b$

input

`integrate((b*x+a)^(1/3)/x^3,x, algorithm="giac")`

output

$$\frac{1}{18}\frac{2\sqrt{3}b^3\arctan\left(\frac{1}{3}\sqrt{3}\frac{2(bx+a)^{1/3}+a^{1/3}}{a^{1/3}}\right)}{a^{5/3}} + \frac{b^3\log\left(\frac{(bx+a)^{2/3}+(bx+a)^{1/3}a^{1/3}+a^{2/3}}{a^{5/3}}\right)}{a^{5/3}} - \frac{2b^3\log\left(\frac{\text{abs}\left((bx+a)^{1/3}-a^{1/3}\right)}{a^{5/3}}\right)}{a^{5/3}} - \frac{3\left((bx+a)^{4/3}b^3+2(bx+a)^{1/3}ab^3\right)}{(ab^2x^2)/b}$$

Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.40

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx = \frac{b^2 \ln\left(\frac{b^2}{(-a)^{2/3}} - \frac{b^2(a+bx)^{1/3}}{a}\right)}{9(-a)^{5/3}} - \frac{\ln\left(\frac{b^2 + \sqrt{3}b^2 1i}{2(-a)^{2/3}} + \frac{b^2(a+bx)^{1/3}}{a}\right) (b^2 + \sqrt{3}b^2 1i)}{18(-a)^{5/3}} - \frac{\frac{b^2(a+bx)^{1/3}}{3} + \frac{b^2(a+bx)^{4/3}}{6a}}{(a+bx)^2 - 2a(a+bx) + a^2} + \frac{b^2 \ln\left(\frac{b^2(a+bx)^{1/3}}{a} - \frac{b^2\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{(-a)^{2/3}}\right) \left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{9(-a)^{5/3}}$$

input `int((a + b*x)^(1/3)/x^3,x)`output $(b^2 \cdot \log(b^2 / (-a)^{2/3} - (b^2 \cdot (a + b \cdot x)^{1/3}) / a)) / (9 \cdot (-a)^{5/3}) - (\log((3^{1/2} \cdot b^2 \cdot 1i + b^2) / (2 \cdot (-a)^{2/3}) + (b^2 \cdot (a + b \cdot x)^{1/3}) / a) \cdot (3^{1/2} \cdot b^2 \cdot 1i + b^2)) / (18 \cdot (-a)^{5/3}) - ((b^2 \cdot (a + b \cdot x)^{1/3}) / 3 + (b^2 \cdot (a + b \cdot x)^{4/3}) / (6 \cdot a)) / ((a + b \cdot x)^2 - 2 \cdot a \cdot (a + b \cdot x) + a^2) + (b^2 \cdot \log((b^2 \cdot (a + b \cdot x)^{1/3}) / a - (b^2 \cdot ((3^{1/2} \cdot 1i) / 2 - 1/2)) / (-a)^{2/3})) \cdot ((3^{1/2} \cdot 1i) / 2 - 1/2)) / (9 \cdot (-a)^{5/3})$ **Reduce [B] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.46

$$\int \frac{\sqrt[3]{a+bx}}{x^3} dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}} + a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) b^2 x^2 + 2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}} - a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) b^2 x^2 - 9a^{\frac{5}{3}}(bx+a)^{\frac{1}{3}} - 3a^{\frac{2}{3}}(bx+a)^{\frac{1}{3}} bx - 21}{=}$$

input `int((b*x+a)^(1/3)/x^3,x)`

output

```
( - 2*sqrt(3)*atan((2*(a + b*x)**(1/6) + a**(1/6))/(a**(1/6)*sqrt(3)))*b**  
2*x**2 + 2*sqrt(3)*atan((2*(a + b*x)**(1/6) - a**(1/6))/(a**(1/6)*sqrt(3))  
) *b**2*x**2 - 9*a**(2/3)*(a + b*x)**(1/3)*a - 3*a**(2/3)*(a + b*x)**(1/3)*  
b*x - 2*log((a + b*x)**(1/6) + a**(1/6))*b**2*x**2 - 2*log((a + b*x)**(1/6  
) - a**(1/6))*b**2*x**2 + log( - a**(1/6)*(a + b*x)**(1/6) + (a + b*x)**(1  
/3) + a**(1/3))*b**2*x**2 + log(a**(1/6)*(a + b*x)**(1/6) + (a + b*x)**(1/  
3) + a**(1/3))*b**2*x**2)/(18*a**(2/3)*a*x**2)
```

3.155 $\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx$

Optimal result	1043
Mathematica [A] (verified)	1043
Rubi [A] (verified)	1044
Maple [F]	1047
Fricas [B] (verification not implemented)	1047
Sympy [F]	1048
Maxima [F]	1048
Giac [B] (verification not implemented)	1049
Mupad [F(-1)]	1049
Reduce [F]	1050

Optimal result

Integrand size = 15, antiderivative size = 104

$$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx = -\frac{\sqrt[3]{a+bx}}{ax} - \frac{2b \left(-\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{a+bx}}{2 \sqrt[3]{a} + \sqrt[3]{a+bx}} \right) + \frac{3}{2} \log \left(\frac{-\sqrt[3]{a} + \sqrt[3]{a+bx}}{\sqrt[3]{x}} \right) \right)}{3a \sqrt[3]{a^2}}$$

output

$$-(b*x+a)^{(1/3)}/a/x-2/3*b*(-3^{(1/2)}*\arctan(3^{(1/2)}*(b*x+a)^{(1/3)}/(2*a^{(1/3)}+(b*x+a)^{(1/3)}))+3/2*\ln((-a^{(1/3)}+(b*x+a)^{(1/3)})/x^{(1/3)})/a/(a^2)^{(1/3)}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.51

$$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx = \frac{-3a^{5/3} - 3a^{2/3}bx + 2\sqrt{3}bx(a+bx)^{2/3} \arctan \left(\frac{1+2\sqrt[3]{a+bx}}{\sqrt[3]{a}} \right) - 2bx(a+bx)^{2/3} \log \left(\sqrt[3]{a} - \sqrt[3]{a+bx} \right)}{3a^{5/3}x \sqrt[3]{(a+bx)^2}}$$

input `Integrate[1/(x^2*((a + b*x)^2)^(1/3)),x]`

output $(-3a^{5/3} - 3a^{2/3}bx + 2\sqrt[3]{3}bx(a + bx)^{2/3}\text{ArcTan}[\frac{1 + (a + bx)^{1/3}}{a^{1/3}}/\sqrt[3]{3}] - 2bx(a + bx)^{2/3}\text{Log}[a^{1/3} - (a + bx)^{1/3}] + bx(a + bx)^{2/3}\text{Log}[a^{2/3} + a^{1/3}(a + bx)^{1/3} + (a + bx)^{2/3}]) / (3a^{5/3}x((a + bx)^2)^{1/3})$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.18, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2008, 52, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx \\
 & \quad \downarrow \text{2008} \\
 & \frac{(a+bx)^{2/3} \int \frac{1}{x^2 (a+bx)^{2/3}} dx}{\sqrt[3]{(a+bx)^2}} \\
 & \quad \downarrow \text{52} \\
 & \frac{(a+bx)^{2/3} \left(-\frac{2b \int \frac{1}{x(a+bx)^{2/3}} dx}{3a} - \frac{\sqrt[3]{a+bx}}{ax} \right)}{\sqrt[3]{(a+bx)^2}} \\
 & \quad \downarrow \text{69} \\
 & \frac{(a+bx)^{2/3} \left(\frac{2b \left(-\frac{\int \frac{1}{\sqrt[3]{a}-\sqrt[3]{a+bx}} dx \sqrt[3]{a+bx}}{2a^{2/3}} - \frac{\int \frac{1}{a^{2/3}+\sqrt[3]{a+bx}} dx \sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2a^{2/3}} \right)}{3a} - \frac{\sqrt[3]{a+bx}}{ax} \right)}{\sqrt[3]{(a+bx)^2}}
 \end{aligned}$$

↓ 16

$$(a + bx)^{2/3} \left(\frac{2b \left(\frac{{}^3f \frac{1}{a^{2/3} + \sqrt[3]{a+bx}} \sqrt[3]{a+(a+bx)^{2/3}} d \sqrt[3]{a+bx}}{2 \sqrt[3]{a}} + \frac{{}^3\log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}}}{{}^3a} \right) - \frac{\sqrt[3]{a+bx}}{ax}}{\sqrt[3]{(a+bx)^2}} \right)$$

$\sqrt[3]{(a+bx)^2}$

↓ 1082

$$(a + bx)^{2/3} \left(\frac{2b \left(\frac{{}^3f \frac{1}{-(a+bx)^{2/3} - 3} d \left(\frac{2 \sqrt[3]{a+bx} + 1}{\sqrt[3]{a}} \right)}{a^{2/3}} + \frac{{}^3\log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}}}{{}^3a} \right) - \frac{\sqrt[3]{a+bx}}{ax}}{\sqrt[3]{(a+bx)^2}} \right)$$

$\sqrt[3]{(a+bx)^2}$

↓ 217

$$(a + bx)^{2/3} \left(\frac{2b \left(\frac{\sqrt{3} \arctan \left(\frac{2 \sqrt[3]{a+bx} + 1}{\sqrt[3]{a}} \right)}{a^{2/3}} + \frac{{}^3\log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}}}{{}^3a} \right) - \frac{\sqrt[3]{a+bx}}{ax}}{\sqrt[3]{(a+bx)^2}} \right)$$

$\sqrt[3]{(a+bx)^2}$

input `Int [1/(x^2*((a + b*x)^2)^(1/3)),x]`

output $((a + b*x)^{2/3} * (-((a + b*x)^{1/3} / (a*x)) - (2*b * (-((\text{Sqrt}[3] * \text{ArcTan}[(1 + (2*(a + b*x)^{1/3}) / a^{1/3}) / \text{Sqrt}[3]]) / a^{2/3}) - \text{Log}[x] / (2*a^{2/3})) + (3 * \text{Log}[a^{1/3} - (a + b*x)^{1/3}]) / (2*a^{2/3}))) / (3*a)) / ((a + b*x)^2)^{1/3}$

Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_.) + (b_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 52 $\text{Int}[(a_.) + (b_.)*(x_)]^{(m_)} * ((c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)} * ((c + d*x)^{(n + 1)} / ((b*c - a*d)*(m + 1))), x] - \text{Simp}[d * ((m + n + 2) / ((b*c - a*d)*(m + 1))) \text{Int}[(a + b*x)^{(m + 1)} * (c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{ILtQ}[m, -1] \&\& \text{FractionQ}[n] \&\& \text{LtQ}[n, 0]$

rule 69 $\text{Int}[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^{2/3}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]] / (2*b*q^2), x] + (-\text{Simp}[3 / (2*b*q) \text{Subst}[\text{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{1/3}], x] - \text{Simp}[3 / (2*b*q^2) \text{Subst}[\text{Int}[1/(q - x), x], x, (c + d*x)^{1/3}], x]]) /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{PosQ}[(b*c - a*d)/b]$

rule 217 $\text{Int}[(a_.) + (b_.)*(x_)]^{2} \wedge (-1), x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2] * \text{Rt}[-b, 2]) \wedge (-1)) * \text{ArcTan}[\text{Rt}[-b, 2] * (x / \text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

rule 1082 $\text{Int}[(a_.) + (b_.)*(x_)] + (c_.)*(x_)]^{2} \wedge (-1), x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 2008

```
Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px,
x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Exp
on[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p),
x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x
] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]
```

Maple [F]

$$\int \frac{1}{x^2 ((bx + a)^2)^{\frac{1}{3}}} dx$$

input

```
int(1/x^2/((b*x+a)^2)^(1/3),x)
```

output

```
int(1/x^2/((b*x+a)^2)^(1/3),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 322 vs. $2(80) = 160$.

Time = 0.09 (sec) , antiderivative size = 322, normalized size of antiderivative = 3.10

$$\int \frac{1}{x^2 \sqrt[3]{(a + bx)^2}} dx =$$

$$6 \sqrt{\frac{1}{3}} (ab^2 x^2 + a^2 bx) \sqrt{-(-a^2)^{\frac{1}{3}}} \arctan \left(-\frac{\sqrt{\frac{1}{3}} \left((-a^2)^{\frac{1}{3}} (bx+a) - 2 (b^2 x^2 + 2 abx + a^2)^{\frac{1}{3}} a \right) \sqrt{-(-a^2)^{\frac{1}{3}}}}{abx + a^2} \right) + 3 (b^2 x^2 +$$

input

```
integrate(1/x^2/((b*x+a)^2)^(1/3),x, algorithm="fricas")
```


output

```
-1/3*(6*sqrt(1/3)*(a*b^2*x^2 + a^2*b*x)*sqrt(-(-a^2)^(1/3))*arctan(-sqrt(1/3)*((-a^2)^(1/3)*(b*x + a) - 2*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)*sqrt(-(-a^2)^(1/3)))/(a*b*x + a^2)) + 3*(b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a^2 - (b^2*x^2 + a*b*x)*(-a^2)^(2/3)*log(((b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a^2 - (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(a*b*x + a^2)*(-a^2)^(1/3) + (b^2*x^2 + 2*a*b*x + a^2)*(-a^2)^(2/3))/(b^2*x^2 + 2*a*b*x + a^2)) + 2*(b^2*x^2 + a*b*x)*(-a^2)^(2/3)*log(((a^2)^(1/3)*(b*x + a) + (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(b*x + a)))/(a^3*b*x^2 + a^4*x)
```

Sympy [F]

$$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx$$

input

```
integrate(1/x**2/((b*x+a)**2)**(1/3), x)
```

output

```
Integral(1/(x**2*((a + b*x)**2)**(1/3)), x)
```

Maxima [F]

$$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{((bx+a)^2)^{\frac{1}{3}} x^2} dx$$

input

```
integrate(1/x^2/((b*x+a)^2)^(1/3), x, algorithm="maxima")
```

output

```
integrate(1/(((b*x + a)^2)^(1/3)*x^2), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. $2(80) = 160$.

Time = 3.00 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.34

$$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx$$

$$= \frac{1}{3} b \left(\frac{2\sqrt{3}(\operatorname{asgn}(bx+a))^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}(2(bx\operatorname{asgn}(bx+a)+\operatorname{asgn}(bx+a))^{\frac{1}{3}}+(\operatorname{asgn}(bx+a))^{\frac{1}{3}})}{3(\operatorname{asgn}(bx+a))^{\frac{1}{3}}}\right)}{a^2} \right) + \frac{(\operatorname{asgn}(bx+a))^{\frac{1}{3}} \log\left(\frac{(bx\operatorname{asgn}(bx+a)+\operatorname{asgn}(bx+a))^{\frac{1}{3}}+(\operatorname{asgn}(bx+a))^{\frac{1}{3}}}{\operatorname{asgn}(bx+a)}\right)}{+ a}$$

input `integrate(1/x^2/((b*x+a)^2)^(1/3),x, algorithm="giac")`

output `1/3*b*(2*sqrt(3)*(a*sgn(b*x + a))^(1/3)*arctan(1/3*sqrt(3)*(2*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(1/3))/(a*sgn(b*x + a))^(1/3))/a^2 + (a*sgn(b*x + a))^(1/3)*log((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(2/3) + (b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3)*(a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(2/3))/a^2 - 2*(a*sgn(b*x + a))^(1/3)*log(abs((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) - (a*sgn(b*x + a))^(1/3)))/a^2 - 3*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3)/(a*b*x*sgn(b*x + a)^2)*sgn(b*x + a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{x^2 ((a+bx)^2)^{1/3}} dx$$

input `int(1/(x^2*((a + b*x)^2)^(1/3)),x)`

output `int(1/(x^2*((a + b*x)^2)^(1/3)), x)`

Reduce [F]

$$\int \frac{1}{x^2 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{(b^2x^2 + 2abx + a^2)^{\frac{1}{3}} x^2} dx$$

input `int(1/x^2/((b*x+a)^2)^(1/3),x)`

output `int(1/((a**2 + 2*a*b*x + b**2*x**2)**(1/3)*x**2),x)`

$$3.156 \quad \int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx$$

Optimal result	1051
Mathematica [A] (verified)	1052
Rubi [A] (verified)	1052
Maple [F]	1057
Fricas [B] (verification not implemented)	1057
Sympy [F]	1058
Maxima [F]	1058
Giac [B] (verification not implemented)	1058
Mupad [F(-1)]	1059
Reduce [F]	1059

Optimal result

Integrand size = 15, antiderivative size = 118

$$\begin{aligned} & \int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx \\ &= \left(-\frac{1}{2ax^2} + \frac{5b}{6a^2x} \right) \sqrt[3]{a+bx} \\ & \quad + \frac{5b^2 \left(-\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{a+bx}}{2 \sqrt[3]{a} + \sqrt[3]{a+bx}} \right) + \frac{3}{2} \log \left(\frac{-\sqrt[3]{a} + \sqrt[3]{a+bx}}{\sqrt[3]{x}} \right) \right)}{9(a^2)^{4/3}} \end{aligned}$$

output

```
(-1/2/a/x^2+5/6*b/a^2/x)*(b*x+a)^(1/3)+5/9*b^2*(-3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/(a^2)^(4/3)
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.55

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx$$

$$= \frac{-9a^{8/3} + 6a^{5/3}bx + 15a^{2/3}b^2x^2 - 10\sqrt{3}b^2x^2(a+bx)^{2/3} \arctan\left(\frac{1 + \frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}}}{\sqrt{3}}\right) + 10b^2x^2(a+bx)^{2/3} \log}{18a^{8/3}x^2\sqrt[3]{(a+bx)^2}}$$

input `Integrate[1/(x^3*((a + b*x)^2)^(1/3)), x]`

output `(-9*a^(8/3) + 6*a^(5/3)*b*x + 15*a^(2/3)*b^2*x^2 - 10*Sqrt[3]*b^2*x^2*(a + b*x)^(2/3)*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3))/Sqrt[3]] + 10*b^2*x^2*(a + b*x)^(2/3)*Log[a^(1/3) - (a + b*x)^(1/3)] - 5*b^2*x^2*(a + b*x)^(2/3)*Log[a^(2/3) + a^(1/3)*(a + b*x)^(1/3) + (a + b*x)^(2/3)]/(18*a^(8/3)*x^2*((a + b*x)^2)^(1/3))`

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.28, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {2008, 52, 52, 69, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx$$

$$\downarrow \text{2008}$$

$$\frac{(a+bx)^{2/3} \int \frac{1}{x^3 (a+bx)^{2/3}} dx}{\sqrt[3]{(a+bx)^2}}$$

$$\downarrow \text{52}$$

$$\frac{(a + bx)^{2/3} \left(-\frac{5b \int \frac{1}{x^2(a+bx)^{2/3}} dx}{6a} - \frac{\sqrt[3]{a + bx}}{2ax^2} \right)}{\sqrt[3]{(a + bx)^2}}$$

↓ 52

$$\frac{(a + bx)^{2/3} \left(-\frac{5b \left(-\frac{2b \int \frac{1}{x(a+bx)^{2/3}} dx}{3a} - \frac{\sqrt[3]{a + bx}}{ax} \right)}{6a} - \frac{\sqrt[3]{a + bx}}{2ax^2} \right)}{\sqrt[3]{(a + bx)^2}}$$

↓ 69

$$\frac{(a + bx)^{2/3} \left(5b \left(\frac{2b \left(\frac{3 \int \frac{1}{\sqrt[3]{a} - \sqrt[3]{a + bx}} dx \sqrt[3]{a + bx}}{2a^{2/3}} - \frac{3 \int \frac{1}{a^{2/3} + \sqrt[3]{a} + bx} dx \sqrt[3]{a + bx}}{2 \sqrt[3]{a}} - \frac{\log(x)}{2a^{2/3}} \right)}{3a} - \frac{\sqrt[3]{a + bx}}{ax} \right)}{6a} \right)}{\sqrt[3]{(a + bx)^2}}$$

↓ 16

$$\left(\begin{array}{l} 5b \left(\frac{3 \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx}} \sqrt[3]{a+(a+bx)^{2/3}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) - \frac{\sqrt[3]{a+bx}}{ax} \\ (a+bx)^{2/3} - \frac{\quad}{6a} \end{array} \right) - \frac{\sqrt[3]{a+bx}}{2ax}$$

$$\sqrt[3]{(a+bx)^2}$$

↓ 1082

$$\left(\begin{array}{l} 5b \left(\frac{3 \int \frac{1}{-(a+bx)^{2/3} - 3} d\left(\frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}} + 1\right)}{a^{2/3}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2a^{2/3}} - \frac{\log(x)}{2a^{2/3}} \right) - \frac{\sqrt[3]{a+bx}}{ax} \\ (a+bx)^{2/3} - \frac{\quad}{6a} \end{array} \right) - \frac{\sqrt[3]{a+bx}}{2ax^2}$$

$$\sqrt[3]{(a+bx)^2}$$

↓ 217

$$\frac{(a + bx)^{2/3} \left(\frac{2b \left(\frac{\sqrt{3} \arctan \left(\frac{2\sqrt[3]{a+bx} + 1}{\sqrt[3]{a}} \right)}{a^{2/3}} \right) + \frac{3 \log \left(\frac{\sqrt[3]{a} - \sqrt[3]{a+bx}}{2a^{2/3}} \right) - \frac{\log(x)}{2a^{2/3}}}{3a} \right) - \frac{5b \sqrt[3]{a+bx}}{3a} - \frac{\sqrt[3]{a+bx}}{2ax^2} \right)}{6a} - \frac{\sqrt[3]{a+bx}}{2ax^2}$$

$$\frac{\sqrt[3]{(a + bx)^2}}{\sqrt[3]{(a + bx)^2}}$$

input `Int [1/(x^3*((a + b*x)^2)^(1/3)),x]`

output `((a + b*x)^(2/3)*(-1/2*(a + b*x)^(1/3)/(a*x^2) - (5*b*(-((a + b*x)^(1/3)/(a*x)) - (2*b*(-((Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]))/a^(2/3)) - Log[x]/(2*a^(2/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)])/(2*a^(2/3)))))/(3*a)))/(6*a)))/((a + b*x)^2)^(1/3)`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 52 $\text{Int}[(a_)+(b_)*(x_)]^{(m_)}*((c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^{(n+1)}/((b*c - a*d)*(m+1))), x] - \text{Simp}[d*((m + n + 2)/((b*c - a*d)*(m+1))) \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&\& \text{ILtQ}[m, -1] \&\& \text{FractionQ}[n] \&\& \text{LtQ}[n, 0]$
- rule 69 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_)]^{(2/3)}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]]/(2*b*q^2), x] + (-\text{Simp}[3/(2*b*q) \text{Subst}[\text{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{(1/3)}], x] - \text{Simp}[3/(2*b*q^2) \text{Subst}[\text{Int}[1/(q - x), x], x, (c + d*x)^{(1/3)}], x])] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{PosQ}[(b*c - a*d)/b]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2]^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2]^{(-1)}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 2008 $\text{Int}[(u_)*(P_x)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{a = \text{Rt}[\text{Coeff}[P_x, x, 0], \text{Expon}[P_x, x]], b = \text{Rt}[\text{Coeff}[P_x, x, \text{Expon}[P_x, x]], \text{Expon}[P_x, x]]\}, \text{Simp}[(a + b*x)^{\text{Expon}[P_x, x]}*u*(a + b*x)^{(\text{Expon}[P_x, x]*p)} \text{Int}[u*(a + b*x)^{(\text{Expon}[P_x, x]*p)}, x], x] /; \text{EqQ}[P_x, (a + b*x)^{\text{Expon}[P_x, x]}] /; \text{!IntegerQ}[p] \&\& \text{PolyQ}[P_x, x] \&\& \text{GtQ}[\text{Expon}[P_x, x], 1] \&\& \text{NeQ}[\text{Coeff}[P_x, x, 0], 0]$

Maple [F]

$$\int \frac{1}{x^3 ((bx + a)^2)^{\frac{1}{3}}} dx$$

input `int(1/x^3/((b*x+a)^2)^(1/3),x)`

output `int(1/x^3/((b*x+a)^2)^(1/3),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 322 vs. $2(91) = 182$.

Time = 0.09 (sec) , antiderivative size = 322, normalized size of antiderivative = 2.73

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx$$

$$= \frac{30 \sqrt{\frac{1}{3}} (ab^3 x^3 + a^2 b^2 x^2) (a^2)^{\frac{1}{6}} \arctan \left(\frac{\sqrt{\frac{1}{3}} (a^2)^{\frac{1}{6}} \left((a^2)^{\frac{1}{3}} (bx+a) + 2(b^2 x^2 + 2abx + a^2)^{\frac{1}{3}} a \right)}{abx + a^2} \right) - 5(b^3 x^3 + ab^2 x^2) (a^2)^{\frac{2}{3}}}{1}$$

input `integrate(1/x^3/((b*x+a)^2)^(1/3),x, algorithm="fricas")`

output `1/18*(30*sqrt(1/3)*(a*b^3*x^3 + a^2*b^2*x^2)*(a^2)^(1/6)*arctan(sqrt(1/3)*(a^2)^(1/6)*((a^2)^(1/3)*(b*x + a) + 2*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(a*b*x + a^2)) - 5*(b^3*x^3 + a*b^2*x^2)*(a^2)^(2/3)*log(((b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a^2 + (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(a*b*x + a^2)*(a^2)^(1/3) + (b^2*x^2 + 2*a*b*x + a^2)*(a^2)^(2/3))/(b^2*x^2 + 2*a*b*x + a^2)) + 10*(b^3*x^3 + a*b^2*x^2)*(a^2)^(2/3)*log(-((a^2)^(1/3)*(b*x + a) - (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(b*x + a)) + 3*(5*a^2*b*x - 3*a^3)*(b^2*x^2 + 2*a*b*x + a^2)^(2/3)/(a^4*b*x^3 + a^5*x^2)`

Sympy [F]

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx$$

input `integrate(1/x**3/((b*x+a)**2)**(1/3), x)`

output `Integral(1/(x**3*((a + b*x)**2)**(1/3)), x)`

Maxima [F]

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{((bx+a)^2)^{\frac{1}{3}} x^3} dx$$

input `integrate(1/x^3/((b*x+a)^2)^(1/3), x, algorithm="maxima")`

output `integrate(1/(((b*x + a)^2)^(1/3)*x^3), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 295 vs. $2(91) = 182$.

Time = 2.95 (sec) , antiderivative size = 295, normalized size of antiderivative = 2.50

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx = \frac{10 \sqrt{3} (\operatorname{asgn}(bx+a))^{\frac{1}{3}} b^3 \arctan \left(\frac{\sqrt{3} \left(2 (bx \operatorname{sgn}(bx+a) + \operatorname{asgn}(bx+a))^{\frac{1}{3}} + (\operatorname{asgn}(bx+a))^{\frac{1}{3}} \right)}{3 (\operatorname{asgn}(bx+a))^{\frac{1}{3}}} \right)}{a^3} + \frac{5 (\operatorname{asgn}(bx+a))^{\frac{1}{3}} b^3 \log \left((bx \operatorname{sgn}(bx+a) + \operatorname{asgn}(bx+a))^{\frac{1}{3}} \right)}{a^3}$$

input `integrate(1/x^3/((b*x+a)^2)^(1/3), x, algorithm="giac")`

output

```
-1/18*(10*sqrt(3)*(a*sgn(b*x + a))^(1/3)*b^3*arctan(1/3*sqrt(3)*(2*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(1/3))/(a*sgn(b*x + a))^(1/3))/a^3 + 5*(a*sgn(b*x + a))^(1/3)*b^3*log((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(2/3) + (b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3)*(a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(2/3))/a^3 - 10*(a*sgn(b*x + a))^(1/3)*b^3*log(abs((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) - (a*sgn(b*x + a))^(1/3)))/a^3 - 3*(5*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(4/3)*b^3*sgn(b*x + a) - 8*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3)*a*b^3)/(a^2*b^2*x^2*sgn(b*x + a)^2))/(b*sgn(b*x + a))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{x^3 ((a+bx)^2)^{1/3}} dx$$

input

```
int(1/(x^3*((a + b*x)^2)^(1/3)),x)
```

output

```
int(1/(x^3*((a + b*x)^2)^(1/3)), x)
```

Reduce [F]

$$\int \frac{1}{x^3 \sqrt[3]{(a+bx)^2}} dx = \int \frac{1}{(b^2x^2 + 2abx + a^2)^{\frac{1}{3}} x^3} dx$$

input

```
int(1/x^3/((b*x+a)^2)^(1/3),x)
```

output

```
int(1/((a**2 + 2*a*b*x + b**2*x**2)**(1/3)*x**3),x)
```

3.157 $\int \frac{1}{x \sqrt[3]{a + bx}} dx$

Optimal result	1060
Mathematica [A] (verified)	1060
Rubi [A] (verified)	1061
Maple [A] (verified)	1063
Fricas [A] (verification not implemented)	1063
Sympy [C] (verification not implemented)	1064
Maxima [A] (verification not implemented)	1065
Giac [A] (verification not implemented)	1065
Mupad [B] (verification not implemented)	1066
Reduce [B] (verification not implemented)	1066

Optimal result

Integrand size = 13, antiderivative size = 78

$$\int \frac{1}{x \sqrt[3]{a + bx}} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3} \sqrt[3]{a + bx}}{2 \sqrt[3]{a} + \sqrt[3]{a + bx}}\right) + \frac{3}{2} \log\left(\frac{-\sqrt[3]{a} + \sqrt[3]{a + bx}}{\sqrt[3]{x}}\right)}{\sqrt[3]{a^2}}$$

output `(3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/(a^2)^(1/3)`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.22

$$\int \frac{1}{x \sqrt[3]{a + bx}} dx = \frac{2\sqrt{3} \arctan\left(\frac{1 + 2 \sqrt[3]{a + bx}}{\sqrt[3]{a}}\right) + 2 \log\left(\sqrt[3]{a} - \sqrt[3]{a + bx}\right) - \log\left(a^{2/3} + \sqrt[3]{a} \sqrt[3]{a + bx} + (a + bx)^{2/3}\right)}{2 \sqrt[3]{a}}$$

input `Integrate[1/(x*(a + b*x)^(1/3)),x]`

output

```
(2*Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3))/Sqrt[3]] + 2*Log[a^(1/3) - (a + b*x)^(1/3)] - Log[a^(2/3) + a^(1/3)*(a + b*x)^(1/3) + (a + b*x)^(2/3)])/(2*a^(1/3))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.96, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x\sqrt[3]{a+bx}} dx \\
 & \quad \downarrow 67 \\
 & \frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx}\sqrt[3]{a} + (a+bx)^{2/3}} d\sqrt[3]{a+bx} - \frac{3 \int \frac{1}{\sqrt[3]{a}-\sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \\
 & \quad \downarrow 16 \\
 & \frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx}\sqrt[3]{a} + (a+bx)^{2/3}} d\sqrt[3]{a+bx} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \\
 & \quad \downarrow 1082 \\
 & -\frac{3 \int \frac{1}{-(a+bx)^{2/3}-3} d\left(\frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}} + 1\right)}{\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \\
 & \quad \downarrow 217 \\
 & \frac{\sqrt{3} \arctan\left(\frac{\frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}} + 1}{\sqrt{3}}\right)}{\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}}
 \end{aligned}$$

input

```
Int[1/(x*(a + b*x)^(1/3)),x]
```

output $(\sqrt{3} \operatorname{ArcTan}[(1 + (2(a + bx)^{1/3})/a^{1/3})/\sqrt{3}])/a^{1/3} - \operatorname{Log}[x/(2a^{1/3}) + (3 \operatorname{Log}[a^{1/3} - (a + bx)^{1/3}])/(2a^{1/3})]$

Defintions of rubi rules used

rule 16 $\operatorname{Int}[(c_.) / ((a_.) + (b_.) * (x_)), x_Symbol] \rightarrow \operatorname{Simp}[c * (\operatorname{Log}[\operatorname{RemoveContent}[a + b * x, x]] / b), x] / ; \operatorname{FreeQ}[\{a, b, c\}, x]$

rule 67 $\operatorname{Int}[1 / (((a_.) + (b_.) * (x_)) * ((c_.) + (d_.) * (x_))^{1/3}), x_Symbol] \rightarrow \operatorname{With}[\{q = \operatorname{Rt}[(b * c - a * d) / b, 3]\}, \operatorname{Simp}[-\operatorname{Log}[\operatorname{RemoveContent}[a + b * x, x]] / (2 * b * q), x] + (\operatorname{Simp}[3 / (2 * b) \operatorname{Subst}[\operatorname{Int}[1 / (q^2 + q * x + x^2), x], x, (c + d * x)^{1/3}], x] - \operatorname{Simp}[3 / (2 * b * q) \operatorname{Subst}[\operatorname{Int}[1 / (q - x), x], x, (c + d * x)^{1/3}], x])] / ; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{PosQ}[(b * c - a * d) / b]$

rule 217 $\operatorname{Int}[(a_.) + (b_.) * (x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2] * \operatorname{Rt}[-b, 2])^{-1} * \operatorname{ArcTan}[\operatorname{Rt}[-b, 2] * (x / \operatorname{Rt}[-a, 2])], x] / ; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{PosQ}[a / b] \& \& (\operatorname{LtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])]$

rule 1082 $\operatorname{Int}[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2)^{-1}, x_Symbol] \rightarrow \operatorname{With}[\{q = 1 - 4 * \operatorname{Simplify}[a * (c / b^2)]\}, \operatorname{Simp}[-2 / b \operatorname{Subst}[\operatorname{Int}[1 / (q - x^2), x], x, 1 + 2 * c * (x / b)], x] / ; \operatorname{RationalQ}[q] \&\& (\operatorname{EqQ}[q^2, 1] \parallel \operatorname{!RationalQ}[b^2 - 4 * a * c])] / ; \operatorname{FreeQ}[\{a, b, c\}, x]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$\frac{\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{a^{\frac{1}{3}}} - \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{2a^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{3}\right)}{a^{\frac{1}{3}}}$	75
default	$\frac{\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{a^{\frac{1}{3}}} - \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{2a^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{3}\right)}{a^{\frac{1}{3}}}$	75
pseudoelliptic	$\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}}\right) + 2\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right) - \ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{2a^{\frac{1}{3}}}$	75

input `int(1/x/(b*x+a)^(1/3), x, method=_RETURNVERBOSE)`

output `1/a^(1/3)*ln(-a^(1/3)+(b*x+a)^(1/3))-1/2/a^(1/3)*ln((b*x+a)^(2/3)+a^(1/3)*(b*x+a)^(1/3)+a^(2/3))+3^(1/2)/a^(1/3)*arctan(1/3*3^(1/2)*(2/a^(1/3)*(b*x+a)^(1/3)+1))`

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 213, normalized size of antiderivative = 2.73

$$\int \frac{1}{x\sqrt[3]{a+bx}} dx$$

$$= \frac{\sqrt{3}a\sqrt{-\frac{1}{a^{\frac{2}{3}}}} \log\left(\frac{2bx+\sqrt{3}\left(2(bx+a)^{\frac{2}{3}}a^{\frac{2}{3}}-(bx+a)^{\frac{1}{3}}a-a^{\frac{4}{3}}\right)\sqrt{-\frac{1}{a^{\frac{2}{3}}}-3(bx+a)^{\frac{1}{3}}a^{\frac{2}{3}}+3a}}{x}}\right) - a^{\frac{2}{3}} \log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}\right)}{2a}$$

input `integrate(1/x/(b*x+a)^(1/3), x, algorithm="fricas")`

output

```
[1/2*(sqrt(3)*a*sqrt(-1/a^(2/3))*log((2*b*x + sqrt(3)*(2*(b*x + a)^(2/3)*a
^(2/3) - (b*x + a)^(1/3)*a - a^(4/3))*sqrt(-1/a^(2/3)) - 3*(b*x + a)^(1/3)
*a^(2/3) + 3*a)/x) - a^(2/3)*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3)
+ a^(2/3)) + 2*a^(2/3)*log((b*x + a)^(1/3) - a^(1/3)))/a, 1/2*(2*sqrt(3)*
a^(2/3)*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3)) - a^(2/3)
)*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3)) + 2*a^(2/3)*log
((b*x + a)^(1/3) - a^(1/3)))/a]
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.05 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.99

$$\int \frac{1}{x\sqrt[3]{a+bx}} dx = \frac{2 \log \left(1 - \frac{\sqrt[3]{b} \sqrt[3]{\frac{a}{b} + x}}{\sqrt[3]{a}} \right) \Gamma\left(\frac{2}{3}\right)}{3\sqrt[3]{a} \Gamma\left(\frac{5}{3}\right)} + \frac{2e^{\frac{2i\pi}{3}} \log \left(1 - \frac{\sqrt[3]{b} \sqrt[3]{\frac{a}{b} + x} e^{\frac{2i\pi}{3}}}{\sqrt[3]{a}} \right) \Gamma\left(\frac{2}{3}\right)}{3\sqrt[3]{a} \Gamma\left(\frac{5}{3}\right)} + \frac{2e^{-\frac{2i\pi}{3}} \log \left(1 - \frac{\sqrt[3]{b} \sqrt[3]{\frac{a}{b} + x} e^{\frac{4i\pi}{3}}}{\sqrt[3]{a}} \right) \Gamma\left(\frac{2}{3}\right)}{3\sqrt[3]{a} \Gamma\left(\frac{5}{3}\right)}$$

input

```
integrate(1/x/(b*x+a)**(1/3),x)
```

output

```
2*log(1 - b**(1/3)*(a/b + x)**(1/3)/a**(1/3))*gamma(2/3)/(3*a**(1/3)*gamma
(5/3)) + 2*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)*exp_polar(2*I*pi
i/3)/a**(1/3))*gamma(2/3)/(3*a**(1/3)*gamma(5/3)) + 2*exp(-2*I*pi/3)*log(1
- b**(1/3)*(a/b + x)**(1/3)*exp_polar(4*I*pi/3)/a**(1/3))*gamma(2/3)/(3*a
**(1/3)*gamma(5/3))
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.97

$$\int \frac{1}{x\sqrt[3]{a+bx}} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{a^{\frac{1}{3}}} - \frac{\log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{2a^{\frac{1}{3}}} + \frac{\log\left((bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}\right)}{a^{\frac{1}{3}}}$$

input `integrate(1/x/(b*x+a)^(1/3),x, algorithm="maxima")`output `sqrt(3)*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3))/a^(1/3) - 1/2*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3))/a^(1/3) + log((b*x + a)^(1/3) - a^(1/3))/a^(1/3)`**Giac [A] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.99

$$\int \frac{1}{x\sqrt[3]{a+bx}} dx = \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{a^{\frac{1}{3}}} - \frac{\log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{2a^{\frac{1}{3}}} + \frac{\log\left(\left|(bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}\right|\right)}{a^{\frac{1}{3}}}$$

input `integrate(1/x/(b*x+a)^(1/3),x, algorithm="giac")`output `sqrt(3)*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3))/a^(1/3) - 1/2*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3))/a^(1/3) + log(abs((b*x + a)^(1/3) - a^(1/3)))/a^(1/3)`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.27

$$\int \frac{1}{x\sqrt[3]{a+bx}} dx = \frac{\ln\left(9(a+bx)^{1/3} - 9a^{1/3}\right)}{a^{1/3}} + \frac{\ln\left(9(a+bx)^{1/3} - \frac{9a^{1/3}(-1+\sqrt{3}1i)^2}{4}\right)(-1+\sqrt{3}1i)}{2a^{1/3}} - \frac{\ln\left(9(a+bx)^{1/3} - \frac{9a^{1/3}(1+\sqrt{3}1i)^2}{4}\right)(1+\sqrt{3}1i)}{2a^{1/3}}$$

input `int(1/(x*(a + b*x)^(1/3)),x)`output `log(9*(a + b*x)^(1/3) - 9*a^(1/3))/a^(1/3) + (log(9*(a + b*x)^(1/3) - (9*a^(1/3)*(3^(1/2)*1i - 1)^2)/4)*(3^(1/2)*1i - 1))/(2*a^(1/3)) - (log(9*(a + b*x)^(1/3) - (9*a^(1/3)*(3^(1/2)*1i + 1)^2)/4)*(3^(1/2)*1i + 1))/(2*a^(1/3))`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.81

$$\int \frac{1}{x\sqrt[3]{a+bx}} dx = \frac{-2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}+a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) + 2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}-a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) + 2\log\left((bx+a)^{\frac{1}{6}}+a^{\frac{1}{6}}\right) + 2\log\left((bx+a)^{\frac{1}{6}}-a^{\frac{1}{6}}\right)}{2a^{\frac{1}{3}}}$$

input `int(1/x/(b*x+a)^(1/3),x)`

output

```
( - 2*sqrt(3)*atan((2*(a + b*x)**(1/6) + a**(1/6))/(a**(1/6)*sqrt(3))) + 2
*sqrt(3)*atan((2*(a + b*x)**(1/6) - a**(1/6))/(a**(1/6)*sqrt(3))) + 2*log(
(a + b*x)**(1/6) + a**(1/6)) + 2*log((a + b*x)**(1/6) - a**(1/6)) - log( -
a**(1/6)*(a + b*x)**(1/6) + (a + b*x)**(1/3) + a**(1/3)) - log(a**(1/6)*(
a + b*x)**(1/6) + (a + b*x)**(1/3) + a**(1/3)))/(2*a**(1/3))
```

3.158 $\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx$

Optimal result	1068
Mathematica [A] (verified)	1068
Rubi [A] (verified)	1069
Maple [F]	1072
Fricas [B] (verification not implemented)	1072
Sympy [F]	1073
Maxima [F]	1073
Giac [B] (verification not implemented)	1073
Mupad [F(-1)]	1074
Reduce [F]	1074

Optimal result

Integrand size = 15, antiderivative size = 95

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx = \frac{3}{2} \sqrt[3]{(a+bx)^2} + \frac{a \left(\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{a+bx}}{2 \sqrt[3]{a} + \sqrt[3]{a+bx}} \right) + \frac{3}{2} \log \left(\frac{-\sqrt[3]{a} + \sqrt[3]{a+bx}}{\sqrt[3]{x}} \right) \right)}{\sqrt[3]{a^2}}$$

output

$$\frac{3/2*((b*x+a)^2)^(1/3)+a*(3^(1/2)*\arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*\ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3))}{(a^2)^(1/3)}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.43

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx = \frac{\sqrt[3]{(a+bx)^2} \left(3(a+bx)^{2/3} + 2\sqrt{3}a^{2/3} \arctan \left(\frac{1+2\sqrt[3]{a+bx}}{\sqrt[3]{a}} \right) + 2a^{2/3} \log \left(\sqrt[3]{a} - \sqrt[3]{a+bx} \right) - a^{2/3} \log \left(\sqrt[3]{a} + \sqrt[3]{a+bx} \right) \right)}{2(a+bx)^{2/3}}$$

input `Integrate[((a + b*x)^2)^(1/3)/x,x]`

output `((((a + b*x)^2)^(1/3)*(3*(a + b*x)^(2/3) + 2*Sqrt[3]*a^(2/3)*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]] + 2*a^(2/3)*Log[a^(1/3) - (a + b*x)^(1/3)] - a^(2/3)*Log[a^(2/3) + a^(1/3)*(a + b*x)^(1/3) + (a + b*x)^(2/3)]))/ (2*(a + b*x)^(2/3))`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.18, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2008, 60, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx$$

↓ 2008

$$\frac{\sqrt[3]{(a+bx)^2} \int \frac{(a+bx)^{2/3}}{x} dx}{(a+bx)^{2/3}}$$

↓ 60

$$\frac{\sqrt[3]{(a+bx)^2} \left(a \int \frac{1}{x \sqrt[3]{a+bx}} dx + \frac{3}{2} (a+bx)^{2/3} \right)}{(a+bx)^{2/3}}$$

↓ 67

$$\frac{\sqrt[3]{(a+bx)^2} \left(a \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} - \frac{3 \int \frac{1}{\sqrt[3]{a} - \sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2 \sqrt[3]{a}} - \frac{\log(x)}{2 \sqrt[3]{a}} \right) + \frac{3}{2} (a+bx)^{2/3} \right)}{(a+bx)^{2/3}}$$

↓ 16

$$\frac{\sqrt[3]{(a+bx)^2} \left(a \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right) + \frac{3}{2}(a+bx)^{2/3} \right)}{(a+bx)^{2/3}}$$

↓ 1082

$$\frac{\sqrt[3]{(a+bx)^2} \left(a \left(-\frac{3 \int \frac{1}{-(a+bx)^{2/3} - 3} d\left(\frac{2\sqrt[3]{a+bx} + 1}{\sqrt[3]{a}}\right)}{\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right) + \frac{3}{2}(a+bx)^{2/3} \right)}{(a+bx)^{2/3}}$$

↓ 217

$$\frac{\sqrt[3]{(a+bx)^2} \left(a \left(\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{a+bx} + 1}{\sqrt[3]{a}}\right)}{\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right) + \frac{3}{2}(a+bx)^{2/3} \right)}{(a+bx)^{2/3}}$$

input `Int[((a + b*x)^2)^(1/3)/x,x]`

output `((((a + b*x)^2)^(1/3)*((3*(a + b*x)^(2/3))/2 + a*((Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]))/a^(1/3) - Log[x]/(2*a^(1/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)]/(2*a^(1/3))))) / (a + b*x)^(2/3)`

Defintions of rubi rules used

- rule 16 $\text{Int}[(c_.)/((a_.) + (b_.)*(x_.)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 60 $\text{Int}[(a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Simp}[n*((b*c - a*d)/(b*(m + n + 1)) \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$
- rule 67 $\text{Int}[1/(((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^{(1/3)}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]]/(2*b*q), x] + (\text{Simp}[3/(2*b) \text{Subst}[\text{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{(1/3)}], x] - \text{Simp}[3/(2*b*q) \text{Subst}[\text{Int}[1/(q - x), x], x, (c + d*x)^{(1/3)}], x])] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{PosQ}[(b*c - a*d)/b]$
- rule 217 $\text{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] || \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] || !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 2008 $\text{Int}[(u_.)*(P_x)^{(p_.)}, x_Symbol] \rightarrow \text{With}[\{a = \text{Rt}[\text{Coeff}[P_x, x, 0], \text{Expon}[P_x, x]], b = \text{Rt}[\text{Coeff}[P_x, x, \text{Expon}[P_x, x]], \text{Expon}[P_x, x]]\}, \text{Simp}[(a + b*x)^{\text{Expon}[P_x, x]} / (a + b*x)^{(\text{Expon}[P_x, x]*p)} \text{Int}[u*(a + b*x)^{(\text{Expon}[P_x, x]*p)}, x], x] /; \text{EqQ}[P_x, (a + b*x)^{\text{Expon}[P_x, x]}] /; !\text{IntegerQ}[p] \&\& \text{PolyQ}[P_x, x] \&\& \text{GtQ}[\text{Expon}[P_x, x], 1] \&\& \text{NeQ}[\text{Coeff}[P_x, x, 0], 0]$

Maple [F]

$$\int \frac{((bx+a)^2)^{\frac{1}{3}}}{x} dx$$

input `int(((b*x+a)^2)^(1/3)/x,x)`

output `int(((b*x+a)^2)^(1/3)/x,x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. $2(73) = 146$.

Time = 0.13 (sec) , antiderivative size = 238, normalized size of antiderivative = 2.51

$$\begin{aligned} & \int \frac{\sqrt[3]{(a+bx)^2}}{x} dx \\ &= -\sqrt{3}(a^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}(abx+a^2) + 2\sqrt{3}(b^2x^2+2abx+a^2)^{\frac{1}{3}}(a^2)^{\frac{2}{3}}}{3(abx+a^2)}}\right) \\ & \quad - \frac{1}{2}(a^2)^{\frac{1}{3}} \log\left(\frac{(b^2x^2+2abx+a^2)^{\frac{2}{3}}a^2 + (b^2x^2+2abx+a^2)^{\frac{1}{3}}(abx+a^2)(a^2)^{\frac{1}{3}} + (b^2x^2+2abx+a^2)(a^2)^{\frac{2}{3}}}{b^2x^2+2abx+a^2}}\right) \\ & \quad + (a^2)^{\frac{1}{3}} \log\left(-\frac{(a^2)^{\frac{1}{3}}(bx+a) - (b^2x^2+2abx+a^2)^{\frac{1}{3}}a}{bx+a}\right) + \frac{3}{2}(b^2x^2+2abx+a^2)^{\frac{1}{3}} \end{aligned}$$

input `integrate(((b*x+a)^2)^(1/3)/x,x, algorithm="fricas")`

output `-sqrt(3)*(a^2)^(1/3)*arctan(1/3*(sqrt(3)*(a*b*x + a^2) + 2*sqrt(3)*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(a^2)^(2/3))/(a*b*x + a^2)) - 1/2*(a^2)^(1/3)*log(((b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a^2 + (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(a*b*x + a^2)*(a^2)^(1/3) + (b^2*x^2 + 2*a*b*x + a^2)*(a^2)^(2/3))/(b^2*x^2 + 2*a*b*x + a^2)) + (a^2)^(1/3)*log(-((a^2)^(1/3)*(b*x + a) - (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(b*x + a)) + 3/2*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)`

Sympy [F]

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx = \int \frac{\sqrt[3]{(a+bx)^2}}{x} dx$$

input `integrate(((b*x+a)**2)**(1/3)/x,x)`

output `Integral(((a + b*x)**2)**(1/3)/x, x)`

Maxima [F]

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx = \int \frac{((bx+a)^2)^{\frac{1}{3}}}{x} dx$$

input `integrate(((b*x+a)^2)^(1/3)/x,x, algorithm="maxima")`

output `integrate(((b*x + a)^2)^(1/3)/x, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 249 vs. $2(73) = 146$.

Time = 2.95 (sec) , antiderivative size = 249, normalized size of antiderivative = 2.62

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx = \frac{1}{2} \left(\frac{2\sqrt{3}(a\operatorname{sgn}(bx+a))^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}(2(bx\operatorname{sgn}(bx+a)+a\operatorname{sgn}(bx+a))^{\frac{1}{3}}+(a\operatorname{sgn}(bx+a))^{\frac{1}{3}})}{3(a\operatorname{sgn}(bx+a))^{\frac{1}{3}}}\right)}{\operatorname{sgn}(bx+a)} - \frac{(a\operatorname{sgn}(bx+a))^{\frac{2}{3}} \log\left((b\right.}{+ a)} \right)$$

input `integrate(((b*x+a)^2)^(1/3)/x,x, algorithm="giac")`

output `1/2*(2*sqrt(3)*(a*sgn(b*x + a))^(2/3)*arctan(1/3*sqrt(3)*(2*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(1/3))/(a*sgn(b*x + a))^(1/3))/sgn(b*x + a) - (a*sgn(b*x + a))^(2/3)*log((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(2/3) + (b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3)*(a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(2/3))/sgn(b*x + a) + 2*(a*sgn(b*x + a))^(2/3)*log(abs((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) - (a*sgn(b*x + a))^(1/3)))/sgn(b*x + a) + 3*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(2/3)/sgn(b*x + a)*sgn(b*x + a)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx = \int \frac{((a+bx)^2)^{1/3}}{x} dx$$

input `int(((a + b*x)^2)^(1/3)/x,x)`

output `int(((a + b*x)^2)^(1/3)/x, x)`

Reduce [F]

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x} dx = \frac{3(b^2x^2 + 2abx + a^2)^{\frac{1}{3}}}{2} + \left(\int \frac{(b^2x^2 + 2abx + a^2)^{\frac{1}{3}}}{bx^2 + ax} dx \right) a$$

input `int(((b*x+a)^2)^(1/3)/x,x)`

output `(3*(a**2 + 2*a*b*x + b**2*x**2)**(1/3) + 2*int((a**2 + 2*a*b*x + b**2*x**2)**(1/3)/(a*x + b*x**2),x)*a)/2`

3.159 $\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx$

Optimal result	1075
Mathematica [A] (verified)	1075
Rubi [A] (verified)	1076
Maple [A] (verified)	1079
Fricas [B] (verification not implemented)	1079
Sympy [F]	1080
Maxima [F]	1081
Giac [B] (verification not implemented)	1081
Mupad [F(-1)]	1082
Reduce [F]	1082

Optimal result

Integrand size = 15, antiderivative size = 115

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx = \frac{b\sqrt[3]{(a+bx)^2}}{a} - \frac{\sqrt[3]{(a+bx)^5}}{ax} + \frac{b\left(\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{a+bx}}{2\sqrt[3]{a}+\sqrt[3]{a+bx}}\right) + \frac{3}{2} \log\left(\frac{-\sqrt[3]{a}+\sqrt[3]{a+bx}}{\sqrt[3]{x}}\right)\right)}{\sqrt[3]{a^2}}$$

output

```
b*((b*x+a)^2)^(1/3)/a-((b*x+a)^5)^(1/3)/a/x+b*(3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/(a^2)^(1/3)
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx = \frac{\sqrt[3]{(a+bx)^2} \left(3\sqrt[3]{a}(a+bx)^{2/3} - 2\sqrt{3}bx \arctan\left(\frac{1+2\sqrt[3]{a+bx}}{\sqrt[3]{a}}\right) - 2bx \log\left(\sqrt[3]{a} - \sqrt[3]{a+bx}\right) + bx \log\left(\sqrt[3]{a} + \sqrt[3]{a+bx}\right) \right)}{3\sqrt[3]{ax}(a+bx)^{2/3}}$$

input `Integrate[((a + b*x)^2)^(1/3)/x^2,x]`

output `-1/3*(((a + b*x)^2)^(1/3)*(3*a^(1/3)*(a + b*x)^(2/3) - 2*Sqrt[3]*b*x*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]] - 2*b*x*Log[a^(1/3) - (a + b*x)^(1/3)] + b*x*Log[a^(2/3) + a^(1/3)*(a + b*x)^(1/3) + (a + b*x)^(2/3)])))/(a^(1/3)*x*(a + b*x)^(2/3))`

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.01, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2008, 51, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx \\
 & \quad \downarrow \text{2008} \\
 & \frac{\sqrt[3]{(a+bx)^2} \int \frac{(a+bx)^{2/3}}{x^2} dx}{(a+bx)^{2/3}} \\
 & \quad \downarrow \text{51} \\
 & \frac{\sqrt[3]{(a+bx)^2} \left(\frac{2}{3} b \int \frac{1}{x \sqrt[3]{a+bx}} dx - \frac{(a+bx)^{2/3}}{x} \right)}{(a+bx)^{2/3}} \\
 & \quad \downarrow \text{67} \\
 & \frac{\sqrt[3]{(a+bx)^2} \left(\frac{2}{3} b \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} - \frac{3 \int \frac{1}{\sqrt[3]{a} - \sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2 \sqrt[3]{a}} - \frac{\log(x)}{2 \sqrt[3]{a}} \right) - \frac{(a+bx)^{2/3}}{x} \right)}{(a+bx)^{2/3}} \\
 & \quad \downarrow \text{16}
 \end{aligned}$$

$$\frac{\sqrt[3]{(a+bx)^2} \left(\frac{2}{3}b \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right) - \frac{(a+bx)^{2/3}}{x} \right)}{(a+bx)^{2/3}}$$

↓ 1082

$$\frac{\sqrt[3]{(a+bx)^2} \left(\frac{2}{3}b \left(-\frac{3 \int \frac{1}{-(a+bx)^{2/3} - 3} d\left(\frac{2\sqrt[3]{a+bx}}{\sqrt[3]{a}} + 1\right)}{\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right) - \frac{(a+bx)^{2/3}}{x} \right)}{(a+bx)^{2/3}}$$

↓ 217

$$\frac{\sqrt[3]{(a+bx)^2} \left(\frac{2}{3}b \left(\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{a+bx} + 1}{\sqrt[3]{a}}\right)}{\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a} - \sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right) - \frac{(a+bx)^{2/3}}{x} \right)}{(a+bx)^{2/3}}$$

input `Int[((a + b*x)^2)^(1/3)/x^2,x]`

output `((a + b*x)^2)^(1/3)*(-(a + b*x)^(2/3)/x + (2*b*((Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]))/a^(1/3) - Log[x]/(2*a^(1/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)]/(2*a^(1/3))))/3)/(a + b*x)^(2/3)`

Definitions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 51 $\text{Int}[(a_)+(b_)*(x_)]^{(m_)}*((c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n/(b*(m+1)), x] - \text{Simp}[d*(n/(b*(m+1))) \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \ \&\& \ \text{ILtQ}[m, -1] \ \&\& \ \text{FractionQ}[n] \ \&\& \ \text{GtQ}[n, 0]$
- rule 67 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_)]^{(1/3)}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]]/(2*b*q), x] + (\text{Simp}[3/(2*b) \ \text{Subst}[\text{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{(1/3)}], x] - \text{Simp}[3/(2*b*q) \ \text{Subst}[\text{Int}[1/(q - x), x], x, (c + d*x)^{(1/3)}], x])] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{PosQ}[(b*c - a*d)/b]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2]^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2]^{(-1)}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[a*(c/b^2)]\}, \text{Simp}[-2/b \ \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 2008 $\text{Int}[(u_)*(P_x)^{(p_)}, x_Symbol] \rightarrow \text{With}[\{a = \text{Rt}[\text{Coeff}[P_x, x, 0], \text{Expon}[P_x, x]], b = \text{Rt}[\text{Coeff}[P_x, x, \text{Expon}[P_x, x]], \text{Expon}[P_x, x]]\}, \text{Simp}[(a + b*x)^{\text{Expon}[P_x, x]}*p/(a + b*x)^{(\text{Expon}[P_x, x]*p)} \ \text{Int}[u*(a + b*x)^{(\text{Expon}[P_x, x]*p)}, x], x] /; \text{EqQ}[P_x, (a + b*x)^{\text{Expon}[P_x, x]}] /; !\text{IntegerQ}[p] \ \&\& \ \text{PolyQ}[P_x, x] \ \&\& \ \text{GtQ}[\text{Expon}[P_x, x], 1] \ \&\& \ \text{NeQ}[\text{Coeff}[P_x, x, 0], 0]$

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.95

method	result
risch	$-\frac{((bx+a)^2)^{\frac{1}{3}}}{x} + \frac{2b \left(\frac{\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{a^{\frac{1}{3}}} - \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{2a^{\frac{1}{3}}} + \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{\frac{a^{\frac{1}{3}}}{3}}\right)}{a^{\frac{1}{3}}} \right)}{3(bx+a)^{\frac{2}{3}}} ((bx+a)^2)^{\frac{1}{3}}$

```
input int(((b*x+a)^2)^(1/3)/x^2,x,method=_RETURNVERBOSE)
```

```
output -((b*x+a)^2)^(1/3)/x+2/3*b*(1/a^(1/3)*ln(-a^(1/3)+(b*x+a)^(1/3))-1/2/a^(1/3)*ln((b*x+a)^(2/3)+a^(1/3)*(b*x+a)^(1/3)+a^(2/3))+3^(1/2)/a^(1/3)*arctan(1/3*3^(1/2)*(2/a^(1/3)*(b*x+a)^(1/3)+1)))/(b*x+a)^(2/3)*((b*x+a)^2)^(1/3)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 230 vs. 2(93) = 186.

Time = 0.11 (sec) , antiderivative size = 560, normalized size of antiderivative = 4.87

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx$$

$$= \left[\frac{3 \sqrt{\frac{1}{3} abx} \sqrt{-\frac{1}{a^{\frac{2}{3}}}} \log \left(-\frac{b^2 x^2 + 4 abx + 3 a^2 + 3 \sqrt{\frac{1}{3}} \left((b^2 x^2 + 2 abx + a^2)^{\frac{1}{3}} (bx+a) a^{\frac{2}{3}} - 2 (b^2 x^2 + 2 abx + a^2)^{\frac{2}{3}} a + (b^2 x^2 + 2 abx + a^2) a^{\frac{1}{3}} \right)}{bx^2 + ax}} \right)}{6 \sqrt{\frac{1}{3}} a^{\frac{2}{3}} bx \arctan \left(\frac{\sqrt{\frac{1}{3}} \left((bx+a) a^{\frac{1}{3}} + 2 (b^2 x^2 + 2 abx + a^2)^{\frac{1}{3}} a^{\frac{2}{3}} \right)}{(bx+a) a^{\frac{1}{3}}} \right)} + a^{\frac{2}{3}} bx \log \left(\frac{(b^2 x^2 + 2 abx + a^2)^{\frac{1}{3}} (bx+a) a^{\frac{2}{3}} + (b^2 x^2 + 2 abx + a^2)^{\frac{2}{3}} a + (b^2 x^2 + 2 abx + a^2) a^{\frac{1}{3}}}{b^2 x^2 + 2 abx + a^2} \right)} \right]$$

input `integrate(((b*x+a)^2)^(1/3)/x^2,x, algorithm="fricas")`

output `[1/3*(3*sqrt(1/3)*a*b*x*sqrt(-1/a^(2/3))*log(-(b^2*x^2 + 4*a*b*x + 3*a^2 + 3*sqrt(1/3)*((b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(b*x + a)*a^(2/3) - 2*(b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a + (b^2*x^2 + 2*a*b*x + a^2)*a^(1/3))*sqrt(-1/a^(2/3)) - 3*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(b*x + a)*a^(1/3))/(b*x^2 + a*x)) - a^(2/3)*b*x*log(((b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(b*x + a)*a^(2/3) + (b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a + (b^2*x^2 + 2*a*b*x + a^2)*a^(1/3))/(b^2*x^2 + 2*a*b*x + a^2)) + 2*a^(2/3)*b*x*log(-((b*x + a)*a^(2/3) - (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(b*x + a)) - 3*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(a*x), -1/3*(6*sqrt(1/3)*a^(2/3)*b*x*arctan(sqrt(1/3)*((b*x + a)*a^(1/3) + 2*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a^(2/3))/(b*x + a)*a^(1/3))) + a^(2/3)*b*x*log(((b^2*x^2 + 2*a*b*x + a^2)^(1/3)*(b*x + a)*a^(2/3) + (b^2*x^2 + 2*a*b*x + a^2)^(2/3)*a + (b^2*x^2 + 2*a*b*x + a^2)*a^(1/3))/(b^2*x^2 + 2*a*b*x + a^2)) - 2*a^(2/3)*b*x*log(-((b*x + a)*a^(2/3) - (b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(b*x + a)) + 3*(b^2*x^2 + 2*a*b*x + a^2)^(1/3)*a)/(a*x)]`

Sympy [F]

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx = \int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx$$

input `integrate(((b*x+a)**2)**(1/3)/x**2,x)`

output `Integral(((a + b*x)**2)**(1/3)/x**2, x)`

Maxima [F]

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx = \int \frac{((bx+a)^2)^{\frac{1}{3}}}{x^2} dx$$

input `integrate(((b*x+a)^2)^(1/3)/x^2,x, algorithm="maxima")`

output `integrate(((b*x + a)^2)^(1/3)/x^2, x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 243 vs. 2(93) = 186.

Time = 2.98 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.11

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx = \frac{1}{3} \left(\frac{2\sqrt{3}(a\operatorname{sgn}(bx+a))^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}(2(bx\operatorname{sgn}(bx+a)+a\operatorname{sgn}(bx+a))^{\frac{1}{3}}+(a\operatorname{sgn}(bx+a))^{\frac{1}{3}})}{3(a\operatorname{sgn}(bx+a))^{\frac{1}{3}}}\right)}{a} - \frac{(a\operatorname{sgn}(bx+a))^{\frac{2}{3}} \log\left((b+bx+a)^2\right)}{a} \right)$$

input `integrate(((b*x+a)^2)^(1/3)/x^2,x, algorithm="giac")`

output `1/3*(2*sqrt(3)*(a*sgn(b*x + a))^(2/3)*arctan(1/3*sqrt(3)*(2*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(1/3))/(a*sgn(b*x + a))^(1/3))/a - (a*sgn(b*x + a))^(2/3)*log((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(2/3) + (b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3)*(a*sgn(b*x + a))^(1/3) + (a*sgn(b*x + a))^(2/3))/a + 2*(a*sgn(b*x + a))^(2/3)*log(abs((b*x*sgn(b*x + a) + a*sgn(b*x + a))^(1/3) - (a*sgn(b*x + a))^(1/3)))/a - 3*(b*x*sgn(b*x + a) + a*sgn(b*x + a))^(2/3)/(b*x*sgn(b*x + a)^2)*b*sgn(b*x + a)^2`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx = \int \frac{((a+bx)^2)^{1/3}}{x^2} dx$$

input `int(((a + b*x)^2)^(1/3)/x^2,x)`output `int(((a + b*x)^2)^(1/3)/x^2, x)`**Reduce [F]**

$$\int \frac{\sqrt[3]{(a+bx)^2}}{x^2} dx = \frac{-3(b^2x^2 + 2abx + a^2)^{\frac{1}{3}} + 2 \left(\int \frac{(b^2x^2 + 2abx + a^2)^{\frac{1}{3}}}{bx^2 + ax} dx \right) bx}{3x}$$

input `int(((b*x+a)^2)^(1/3)/x^2,x)`output `(- 3*(a**2 + 2*a*b*x + b**2*x**2)**(1/3) + 2*int((a**2 + 2*a*b*x + b**2*x**2)**(1/3)/(a*x + b*x**2),x)*b*x)/(3*x)`

3.160 $\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx$

Optimal result	1083
Mathematica [A] (verified)	1083
Rubi [A] (verified)	1084
Maple [A] (verified)	1085
Fricas [A] (verification not implemented)	1085
Sympy [F]	1086
Maxima [A] (verification not implemented)	1086
Giac [A] (verification not implemented)	1086
Mupad [F(-1)]	1087
Reduce [B] (verification not implemented)	1087

Optimal result

Integrand size = 15, antiderivative size = 138

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = \left(\frac{b}{6a^3} - \frac{1}{2ax^2} \right) (a+bx)^{5/3} - \frac{b^2 \sqrt[3]{(a+bx)^2}}{6a^2} - \frac{b^2 \left(\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{a+bx}}{2 \sqrt[3]{a} + \sqrt[3]{a+bx}} \right) + \frac{3}{2} \log \left(\frac{-\sqrt[3]{a} + \sqrt[3]{a+bx}}{\sqrt[3]{x}} \right) \right)}{9a \sqrt[3]{a^2}}$$

output

```
(1/6*b/a^3-1/2/a/x^2)*(b*x+a)^(5/3)-1/6*b^2*((b*x+a)^2)^(1/3)/a^2-1/9*b^2*(3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/a/(a^2)^(1/3)
```

Mathematica [A] (verified)

Time = 1.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.22

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = \frac{\sqrt[3]{(a+bx)^3}(-a+bx \log(x))}{x(a+bx)}$$

input

```
Integrate[((a + b*x)^3)^(1/3)/x^2,x]
```

output $((a + bx)^3)^{1/3}(-a + b \cdot \text{Log}[x]) / (x(a + bx))$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.22, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2008, 49, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx \\ & \quad \downarrow \text{2008} \\ & \frac{\sqrt[3]{(a+bx)^3} \int \frac{a+bx}{x^2} dx}{a+bx} \\ & \quad \downarrow \text{49} \\ & \frac{\sqrt[3]{(a+bx)^3} \int \left(\frac{a}{x^2} + \frac{b}{x}\right) dx}{a+bx} \\ & \quad \downarrow \text{2009} \\ & \frac{\sqrt[3]{(a+bx)^3} \left(b \log(x) - \frac{a}{x}\right)}{a+bx} \end{aligned}$$

input $\text{Int}[(a + bx)^3)^{1/3}/x^2, x]$

output $((a + bx)^3)^{1/3}(-(a/x) + b \cdot \text{Log}[x]) / (a + bx)$

Defintions of rubi rules used

rule 49 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[m, 0] && IGtQ[m + n + 2, 0]`

rule 2008 `Int[(u_.)*(Px_)^(p_), x_Symbol] := With[{a = Rt[Coeff[Px, x, 0], Expon[Px, x]], b = Rt[Coeff[Px, x, Expon[Px, x]], Expon[Px, x]]}, Simp[((a + b*x)^Expon[Px, x])^p/(a + b*x)^(Expon[Px, x]*p) Int[u*(a + b*x)^(Expon[Px, x]*p), x], x] /; EqQ[Px, (a + b*x)^Expon[Px, x]] /; !IntegerQ[p] && PolyQ[Px, x] && GtQ[Expon[Px, x], 1] && NeQ[Coeff[Px, x, 0], 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.32

method	result	size
risch	$-\frac{((bx+a)^3)^{\frac{1}{3}} a}{(bx+a)x} + \frac{((bx+a)^3)^{\frac{1}{3}} b \ln(x)}{bx+a}$	44

input `int(((b*x+a)^3)^(1/3)/x^2,x,method=_RETURNVERBOSE)`

output `-((b*x+a)^3)^(1/3)/(b*x+a)*a/x+((b*x+a)^3)^(1/3)/(b*x+a)*b*ln(x)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.09

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = \frac{bx \log(x) - a}{x}$$

input `integrate(((b*x+a)^3)^(1/3)/x^2,x, algorithm="fricas")`

output `(b*x*log(x) - a)/x`

Sympy [F]

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = \int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx$$

input `integrate(((b*x+a)**3)**(1/3)/x**2,x)`

output `Integral(((a + b*x)**3)**(1/3)/x**2, x)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.08

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = b \log(x) - \frac{a}{x}$$

input `integrate(((b*x+a)^3)^(1/3)/x^2,x, algorithm="maxima")`

output `b*log(x) - a/x`

Giac [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.09

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = b \log(|x|) - \frac{a}{x}$$

input `integrate(((b*x+a)^3)^(1/3)/x^2,x, algorithm="giac")`

output `b*log(abs(x)) - a/x`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = \int \frac{((a+bx)^3)^{1/3}}{x^2} dx$$

input `int(((a + b*x)^3)^(1/3)/x^2,x)`output `int(((a + b*x)^3)^(1/3)/x^2, x)`**Reduce [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.09

$$\int \frac{\sqrt[3]{(a+bx)^3}}{x^2} dx = \frac{\log(x) bx - a}{x}$$

input `int(((b*x+a)^3)^(1/3)/x^2,x)`output `(log(x)*b*x - a)/x`

3.161 $\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx$

Optimal result	1088
Mathematica [A] (verified)	1088
Rubi [A] (verified)	1089
Maple [A] (verified)	1091
Fricas [A] (verification not implemented)	1092
Sympy [C] (verification not implemented)	1093
Maxima [A] (verification not implemented)	1094
Giac [A] (verification not implemented)	1094
Mupad [B] (verification not implemented)	1095
Reduce [B] (verification not implemented)	1095

Optimal result

Integrand size = 13, antiderivative size = 105

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx = -\frac{\sqrt[3]{(a+bx)^2}}{ax} - \frac{b \left(\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{a+bx}}{2 \sqrt[3]{a} + \sqrt[3]{a+bx}} \right) + \frac{3}{2} \log \left(\frac{-\sqrt[3]{a} + \sqrt[3]{a+bx}}{\sqrt[3]{x}} \right) \right)}{3a \sqrt[3]{a^2}}$$

output

```

-((b*x+a)^2)^(1/3)/a/x-1/3*b*(3^(1/2)*arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/a/(a^2)^(1/3)
    
```

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.14

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx = \frac{6 \sqrt[3]{a}(a+bx)^{2/3} + 2\sqrt{3}bx \arctan \left(\frac{1 + 2 \sqrt[3]{a+bx}}{\sqrt[3]{a}} \right) + 2bx \log \left(\sqrt[3]{a} - \sqrt[3]{a+bx} \right) - bx \log \left(a^{2/3} + \sqrt[3]{a} \sqrt[3]{a+bx} \right)}{6a^{4/3}x}$$

input `Integrate[1/(x^2*(a + b*x)^(1/3)),x]`

output
$$-1/6*(6*a^{1/3}*(a + b*x)^{2/3} + 2*\text{Sqrt}[3]*b*x*\text{ArcTan}[(1 + (2*(a + b*x)^{1/3}))/a^{1/3}]/\text{Sqrt}[3]] + 2*b*x*\text{Log}[a^{1/3} - (a + b*x)^{1/3}] - b*x*\text{Log}[a^{2/3} + a^{1/3}*(a + b*x)^{1/3} + (a + b*x)^{2/3}]/(a^{4/3}*x)$$

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.96, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {52, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx$$

$$\downarrow 52$$

$$-\frac{b \int \frac{1}{x \sqrt[3]{a+bx}} dx}{3a} - \frac{(a+bx)^{2/3}}{ax}$$

$$\downarrow 67$$

$$-\frac{b \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} - \frac{3 \int \frac{1}{\sqrt[3]{a}-\sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right)}{\frac{3a}{(a+bx)^{2/3}} - ax}$$

$$\downarrow 16$$

$$-\frac{b \left(\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} + \frac{3 \log(\sqrt[3]{a}-\sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right)}{\frac{3a}{(a+bx)^{2/3}} - ax}$$

$$\downarrow 1082$$

$$\begin{array}{c}
 \frac{b \left(-\frac{3 \int \frac{1}{-(a+bx)^{2/3}-3} dx \left(\frac{2\sqrt[3]{a+bx}+1}{\sqrt[3]{a}} \right) + \frac{3 \log(\sqrt[3]{a}-\sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right)}{3a} - \frac{(a+bx)^{2/3}}{ax} \\
 \downarrow 217 \\
 \frac{b \left(\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{a+bx}+1}{\sqrt[3]{a}}\right)}{\sqrt[3]{a}} + \frac{3 \log(\sqrt[3]{a}-\sqrt[3]{a+bx})}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right)}{3a} - \frac{(a+bx)^{2/3}}{ax}
 \end{array}$$

input `Int[1/(x^2*(a + b*x)^(1/3)),x]`

output `-((a + b*x)^(2/3)/(a*x)) - (b*((Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))]/a^(1/3))/Sqrt[3]])/a^(1/3) - Log[x]/(2*a^(1/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)]/(2*a^(1/3))))/(3*a)`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

```
rule 217 Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

```
rule 1082 Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.90

method	result
risch	$-\frac{(bx+a)^{\frac{2}{3}}}{ax} - \frac{b \ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{3a^{\frac{4}{3}}} + \frac{b \ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{6a^{\frac{4}{3}}} - \frac{b\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{\frac{a^{\frac{1}{3}}}{3}}\right)}{3a^{\frac{4}{3}}}$
pseudoelliptic	$-2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{\frac{a^{\frac{1}{3}}}{3}}\right)bx - 2 \ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)bx + \ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)bx - 6(bx+a)^{\frac{2}{3}}a$ <hr/> $6a^{\frac{4}{3}}x$
derivativedivides	$3b \left(-\frac{(bx+a)^{\frac{2}{3}}}{3abx} + \frac{\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}} + \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{\frac{a^{\frac{1}{3}}}{3}}\right)}{3a^{\frac{1}{3}}} \right)$
default	$3b \left(-\frac{(bx+a)^{\frac{2}{3}}}{3abx} + \frac{\ln\left(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}} + \frac{\ln\left((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{6a^{\frac{1}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}+1\right)}{\frac{a^{\frac{1}{3}}}{3}}\right)}{3a^{\frac{1}{3}}} \right)$

input `int(1/x^2/(b*x+a)^(1/3),x,method=_RETURNVERBOSE)`

output `-1/a*(b*x+a)^(2/3)/x-1/3*b/a^(4/3)*ln(-a^(1/3)+(b*x+a)^(1/3))+1/6*b/a^(4/3)*ln((b*x+a)^(2/3)+a^(1/3)*(b*x+a)^(1/3)+a^(2/3))-1/3*b/a^(4/3)*3^(1/2)*arctan(1/3*3^(1/2)*(2/a^(1/3)*(b*x+a)^(1/3)+1))`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.91

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx$$

$$= \frac{3 \sqrt[3]{\frac{1}{3} abx} \sqrt{\frac{(-a)^{\frac{1}{3}}}{a}} \log \left(\frac{2bx - 3 \sqrt[3]{\frac{1}{3}} (2(bx+a)^{\frac{2}{3}} (-a)^{\frac{2}{3}} - (bx+a)^{\frac{1}{3}} a + (-a)^{\frac{1}{3}} a) \sqrt{\frac{(-a)^{\frac{1}{3}}}{a} - 3(bx+a)^{\frac{1}{3}} (-a)^{\frac{2}{3}} + 3a}}{x}} \right) + (-a)^{\frac{2}{3}} bx \log \left(\frac{2(bx+a)^{\frac{2}{3}} (-a)^{\frac{2}{3}} - (bx+a)^{\frac{1}{3}} a + (-a)^{\frac{1}{3}} a}{x} \right)}{6a^2x} + \frac{6 \sqrt[3]{\frac{1}{3} abx} \sqrt{-\frac{(-a)^{\frac{1}{3}}}{a}} \arctan \left(\sqrt{\frac{1}{3}} (2(bx+a)^{\frac{1}{3}} - (-a)^{\frac{1}{3}}) \sqrt{-\frac{(-a)^{\frac{1}{3}}}{a}} \right) - (-a)^{\frac{2}{3}} bx \log \left(\frac{(bx+a)^{\frac{2}{3}} - (bx+a)^{\frac{1}{3}} a + (-a)^{\frac{1}{3}} a}{x} \right)}{6a^2x}$$

input `integrate(1/x^2/(b*x+a)^(1/3),x, algorithm="fricas")`

output `[1/6*(3*sqrt(1/3)*a*b*x*sqrt((-a)^(1/3)/a)*log((2*b*x - 3*sqrt(1/3)*(2*(b*x + a)^(2/3)*(-a)^(2/3) - (b*x + a)^(1/3)*a + (-a)^(1/3)*a)*sqrt((-a)^(1/3)/a) - 3*(b*x + a)^(1/3)*(-a)^(2/3) + 3*a)/x) + (-a)^(2/3)*b*x*log((b*x + a)^(2/3) - (b*x + a)^(1/3)*(-a)^(1/3) + (-a)^(2/3)) - 2*(-a)^(2/3)*b*x*log((b*x + a)^(1/3) + (-a)^(1/3)) - 6*(b*x + a)^(2/3)*a/(a^2*x), -1/6*(6*sqrt(1/3)*a*b*x*sqrt(-(-a)^(1/3)/a)*arctan(sqrt(1/3)*(2*(b*x + a)^(1/3) - (-a)^(1/3))*sqrt(-(-a)^(1/3)/a)) - (-a)^(2/3)*b*x*log((b*x + a)^(2/3) - (b*x + a)^(1/3)*(-a)^(1/3) + (-a)^(2/3)) + 2*(-a)^(2/3)*b*x*log((b*x + a)^(1/3) + (-a)^(1/3)) + 6*(b*x + a)^(2/3)*a/(a^2*x)]`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.40 (sec) , antiderivative size = 831, normalized size of antiderivative = 7.91

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx = \text{Too large to display}$$

input `integrate(1/x**2/(b*x+a)**(1/3),x)`

output

```
-2*a**(5/3)*b**(7/3)*(a/b + x)**(4/3)*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b
+ x)**(1/3)/a**(1/3))*gamma(2/3)/(9*a**3*b**(4/3)*(a/b + x)**(4/3)*exp(2*I
*pi/3)*gamma(5/3) - 9*a**2*b**(7/3)*(a/b + x)**(7/3)*exp(2*I*pi/3)*gamma(5
/3)) - 2*a**(5/3)*b**(7/3)*(a/b + x)**(4/3)*exp(-2*I*pi/3)*log(1 - b**(1/3)
)*(a/b + x)**(1/3)*exp_polar(2*I*pi/3)/a**(1/3))*gamma(2/3)/(9*a**3*b**(4/
3)*(a/b + x)**(4/3)*exp(2*I*pi/3)*gamma(5/3) - 9*a**2*b**(7/3)*(a/b + x)**
(7/3)*exp(2*I*pi/3)*gamma(5/3)) - 2*a**(5/3)*b**(7/3)*(a/b + x)**(4/3)*log
(1 - b**(1/3)*(a/b + x)**(1/3)*exp_polar(4*I*pi/3)/a**(1/3))*gamma(2/3)/(9
*a**3*b**(4/3)*(a/b + x)**(4/3)*exp(2*I*pi/3)*gamma(5/3) - 9*a**2*b**(7/3)
*(a/b + x)**(7/3)*exp(2*I*pi/3)*gamma(5/3)) + 2*a**(2/3)*b**(10/3)*(a/b +
x)**(7/3)*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)/a**(1/3))*gamma(
2/3)/(9*a**3*b**(4/3)*(a/b + x)**(4/3)*exp(2*I*pi/3)*gamma(5/3) - 9*a**2*b
**(7/3)*(a/b + x)**(7/3)*exp(2*I*pi/3)*gamma(5/3)) + 2*a**(2/3)*b**(10/3)*
(a/b + x)**(7/3)*exp(-2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)*exp_pola
r(2*I*pi/3)/a**(1/3))*gamma(2/3)/(9*a**3*b**(4/3)*(a/b + x)**(4/3)*exp(2*I
*pi/3)*gamma(5/3) - 9*a**2*b**(7/3)*(a/b + x)**(7/3)*exp(2*I*pi/3)*gamma(5
/3)) + 2*a**(2/3)*b**(10/3)*(a/b + x)**(7/3)*log(1 - b**(1/3)*(a/b + x)**(
1/3)*exp_polar(4*I*pi/3)/a**(1/3))*gamma(2/3)/(9*a**3*b**(4/3)*(a/b + x)**
(4/3)*exp(2*I*pi/3)*gamma(5/3) - 9*a**2*b**(7/3)*(a/b + x)**(7/3)*exp(2*I*
pi/3)*gamma(5/3)) + 6*a*b**3*(a/b + x)**2*exp(2*I*pi/3)*gamma(2/3)/(9*a...
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.01

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx = -\frac{\sqrt{3}b \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{3a^{\frac{4}{3}}} - \frac{(bx+a)^{\frac{2}{3}}b}{(bx+a)a-a^2}$$

$$+ \frac{b \log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{6a^{\frac{4}{3}}} - \frac{b \log\left((bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}\right)}{3a^{\frac{4}{3}}}$$

input `integrate(1/x^2/(b*x+a)^(1/3),x, algorithm="maxima")`output `-1/3*sqrt(3)*b*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3))/a^(4/3) - (b*x + a)^(2/3)*b/((b*x + a)*a - a^2) + 1/6*b*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3))/a^(4/3) - 1/3*b*log((b*x + a)^(1/3) - a^(1/3))/a^(4/3)`**Giac [A] (verification not implemented)**

Time = 0.37 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.95

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx =$$

$$-\frac{1}{6}b \left(\frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{a^{\frac{4}{3}}} - \frac{\log\left((bx+a)^{\frac{2}{3}}+(bx+a)^{\frac{1}{3}}a^{\frac{1}{3}}+a^{\frac{2}{3}}\right)}{a^{\frac{4}{3}}} + \frac{2 \log\left(\left|(bx+a)^{\frac{1}{3}}-a^{\frac{1}{3}}\right|\right)}{a^{\frac{4}{3}}}\right)$$

input `integrate(1/x^2/(b*x+a)^(1/3),x, algorithm="giac")`output `-1/6*b*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3))/a^(4/3) - log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3))/a^(4/3) + 2*log(abs((b*x + a)^(1/3) - a^(1/3)))/a^(4/3) + 6*(b*x + a)^(2/3)/(a*b*x)`

Mupad [B] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.24

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx = -\frac{(a+bx)^{2/3}}{ax} + \frac{\ln\left(\frac{(b-\sqrt{3}bi)^2}{4a^{5/3}} - \frac{b^2(a+bx)^{1/3}}{a^2}\right)(b-\sqrt{3}bi)}{6a^{4/3}} + \frac{\ln\left(\frac{(b+\sqrt{3}bi)^2}{4a^{5/3}} - \frac{b^2(a+bx)^{1/3}}{a^2}\right)(b+\sqrt{3}bi)}{6a^{4/3}} - \frac{b \ln\left((a+bx)^{1/3} - a^{1/3}\right)}{3a^{4/3}}$$

input `int(1/(x^2*(a + b*x)^(1/3)),x)`output `(log((b - 3^(1/2)*b*1i)^2/(4*a^(5/3)) - (b^2*(a + b*x)^(1/3))/a^2)*(b - 3^(1/2)*b*1i))/(6*a^(4/3)) - (a + b*x)^(2/3)/(a*x) + (log((b + 3^(1/2)*b*1i)^2/(4*a^(5/3)) - (b^2*(a + b*x)^(1/3))/a^2)*(b + 3^(1/2)*b*1i))/(6*a^(4/3)) - (b*log((a + b*x)^(1/3) - a^(1/3)))/(3*a^(4/3))`**Reduce [B] (verification not implemented)**

Time = 0.25 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.58

$$\int \frac{1}{x^2 \sqrt[3]{a+bx}} dx = \frac{2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}+a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) bx - 2\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}-a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) bx - 6a^{\frac{1}{3}}(bx+a)^{\frac{2}{3}} - 2\log\left((bx+a)^{\frac{1}{6}}+a^{\frac{1}{6}}\right) bx - \dots}{\dots}$$

input `int(1/x^2/(b*x+a)^(1/3),x)`

output

```
(2*sqrt(3)*atan((2*(a + b*x)**(1/6) + a**(1/6))/(a**(1/6)*sqrt(3)))*b*x -  
2*sqrt(3)*atan((2*(a + b*x)**(1/6) - a**(1/6))/(a**(1/6)*sqrt(3)))*b*x - 6  
*a**(1/3)*(a + b*x)**(2/3) - 2*log((a + b*x)**(1/6) + a**(1/6))*b*x - 2*lo  
g((a + b*x)**(1/6) - a**(1/6))*b*x + log(- a**(1/6)*(a + b*x)**(1/6) + (a  
+ b*x)**(1/3) + a**(1/3))*b*x + log(a**(1/6)*(a + b*x)**(1/6) + (a + b*x)  
**(1/3) + a**(1/3))*b*x)/(6*a**(1/3)*a*x)
```

3.162 $\int \frac{1}{x^3 \sqrt[3]{a + bx}} dx$

Optimal result	1097
Mathematica [A] (verified)	1098
Rubi [A] (verified)	1098
Maple [A] (verified)	1101
Fricas [A] (verification not implemented)	1103
Sympy [C] (verification not implemented)	1104
Maxima [A] (verification not implemented)	1105
Giac [A] (verification not implemented)	1105
Mupad [B] (verification not implemented)	1106
Reduce [B] (verification not implemented)	1106

Optimal result

Integrand size = 13, antiderivative size = 117

$$\int \frac{1}{x^3 \sqrt[3]{a + bx}} dx = \left(-\frac{1}{2ax^2} + \frac{2b}{3a^2x} \right) \sqrt[3]{a + bx} + \frac{2b^2 \left(\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{a + bx}}{2 \sqrt[3]{a} + \sqrt[3]{a + bx}} \right) + \frac{3}{2} \log \left(\frac{-\sqrt[3]{a} + \sqrt[3]{a + bx}}{\sqrt[3]{x}} \right) \right)}{9(a^2)^{4/3}}$$

output

$(-1/2/a/x^2+2/3*b/a^2/x)*(b*x+a)^(1/3)+2/9*b^2*(3^(1/2)*\arctan(3^(1/2)*(b*x+a)^(1/3)/(2*a^(1/3)+(b*x+a)^(1/3)))+3/2*\ln((-a^(1/3)+(b*x+a)^(1/3))/x^(1/3)))/(a^2)^(4/3)$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.27

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx = -\frac{(a+bx)^{2/3}(7a-4(a+bx))}{6a^2x^2} + \frac{2b^2 \arctan\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{a+bx}}{\sqrt{3}\sqrt[3]{a}}\right)}{3\sqrt{3}a^{7/3}} + \frac{2b^2 \log\left(\sqrt[3]{a} - \sqrt[3]{a+bx}\right)}{9a^{7/3}} - \frac{b^2 \log\left(a^{2/3} + \sqrt[3]{a}\sqrt[3]{a+bx} + (a+bx)^{2/3}\right)}{9a^{7/3}}$$

input `Integrate[1/(x^3*(a + b*x)^(1/3)),x]`

output `-1/6*((a + b*x)^(2/3)*(7*a - 4*(a + b*x)))/(a^2*x^2) + (2*b^2*ArcTan[1/Sqrt[3] + (2*(a + b*x)^(1/3))/(Sqrt[3]*a^(1/3))]/(3*Sqrt[3]*a^(7/3)) + (2*b^2*Log[a^(1/3) - (a + b*x)^(1/3)])/(9*a^(7/3)) - (b^2*Log[a^(2/3) + a^(1/3)*(a + b*x)^(1/3) + (a + b*x)^(2/3)])/(9*a^(7/3))`

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.10, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {52, 52, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx$$

↓ 52

$$-\frac{2b \int \frac{1}{x^2 \sqrt[3]{a+bx}} dx}{3a} - \frac{(a+bx)^{2/3}}{2ax^2}$$

↓ 52

$$\frac{2b \left(-\frac{b \int \frac{1}{x \sqrt[3]{a+bx}} dx}{3a} - \frac{(a+bx)^{2/3}}{ax} \right)}{3a} - \frac{(a+bx)^{2/3}}{2ax^2}$$

↓ 67

$$2b \left(\frac{b \left(\frac{\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} - \frac{\int \frac{1}{\sqrt[3]{a} - \sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right)}{3a} - \frac{(a+bx)^{2/3}}{ax} \right)$$

$$\frac{3a}{(a+bx)^{2/3}} - \frac{3a}{2ax^2}$$

↓ 16

$$2b \left(\frac{b \left(\frac{\frac{3}{2} \int \frac{1}{a^{2/3} + \sqrt[3]{a+bx} \sqrt[3]{a+(a+bx)^{2/3}}} d\sqrt[3]{a+bx} + \frac{\int \frac{1}{\sqrt[3]{a} - \sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right)}{3a} - \frac{(a+bx)^{2/3}}{ax} \right)$$

$$\frac{3a}{(a+bx)^{2/3}} - \frac{3a}{2ax^2}$$

↓ 1082

$$2b \left(\frac{b \left(-\frac{\int \frac{1}{-(a+bx)^{2/3}-3} d\left(\frac{2\sqrt[3]{a+bx}+1}{\sqrt[3]{a}}\right)}{\sqrt[3]{a}} + \frac{\int \frac{1}{\sqrt[3]{a} - \sqrt[3]{a+bx}} d\sqrt[3]{a+bx}}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}} \right)}{3a} - \frac{(a+bx)^{2/3}}{ax} \right)$$

$$\frac{3a}{(a+bx)^{2/3}} - \frac{3a}{2ax^2}$$

↓ 217

$$\frac{2b \left(\frac{b \left(\frac{\sqrt{3} \arctan \left(\frac{2\sqrt[3]{a+bx} + 1}{\sqrt[3]{a}} \right)}{\sqrt{3}} \right) + \frac{3 \log \left(\sqrt[3]{a} - \sqrt[3]{a+bx} \right)}{2\sqrt[3]{a}} - \frac{\log(x)}{2\sqrt[3]{a}}}{\sqrt[3]{a}} \right) - \frac{(a+bx)^{2/3}}{ax}}{3a} - \frac{(a+bx)^{2/3}}{2ax^2}}{3a} - \frac{(a+bx)^{2/3}}{2ax^2}$$

input `Int[1/(x^3*(a + b*x)^(1/3)),x]`

output `-1/2*(a + b*x)^(2/3)/(a*x^2) - (2*b*(-((a + b*x)^(2/3)/(a*x)) - (b*((Sqrt[3]*ArcTan[(1 + (2*(a + b*x)^(1/3))/a^(1/3)]/Sqrt[3]))/a^(1/3) - Log[x]/(2*a^(1/3)) + (3*Log[a^(1/3) - (a + b*x)^(1/3)]/(2*a^(1/3)))))/(3*a)))/(3*a)`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 52 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Simp[d*((m + n + 2)/((b*c - a*d)*(m + 1)) Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && ILtQ[m, -1] && FractionQ[n] && LtQ[n, 0]`

rule 67

```
Int[1/((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(1/3)), x_Symbol] := With[
{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x
] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)],
x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /
; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]
```

rule 217

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

rule 1082

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Fre
eQ[{a, b, c}, x]
```

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 109, normalized size of antiderivative = 0.93

method	result
risch	$-\frac{(bx+a)^{\frac{2}{3}}(-4bx+3a)}{6a^2x^2} + \frac{2b^2 \ln(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}})}{9a^{\frac{7}{3}}} - \frac{b^2 \ln((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}})}{9a^{\frac{7}{3}}} + \frac{2b^2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}\right)}{3a^{\frac{1}{3}}}\right)}{9a^{\frac{7}{3}}}$
pseudoelliptic	$\frac{4b^2\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}}\right)}{3a^{\frac{1}{3}}}\right)}{18a^{\frac{7}{3}}x^2} + \frac{x^2+4b^2 \ln(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}})x^2-2b^2 \ln((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}})x^2+12b^2 \ln\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}\right)}{18a^{\frac{7}{3}}x^2}$
derivativedivides	$3b^2 \left(-\frac{(bx+a)^{\frac{2}{3}}}{6ab^2x^2} - \frac{2 \left(-\frac{(bx+a)^{\frac{2}{3}}}{3abx} + \frac{\ln(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}})}{3a^{\frac{1}{3}}} + \frac{\ln((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}})}{6a^{\frac{1}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}\right)}{3a^{\frac{1}{3}}}\right)}{3a} \right)}{3a} \right)$
default	$3b^2 \left(-\frac{(bx+a)^{\frac{2}{3}}}{6ab^2x^2} - \frac{2 \left(-\frac{(bx+a)^{\frac{2}{3}}}{3abx} + \frac{\ln(-a^{\frac{1}{3}}+(bx+a)^{\frac{1}{3}})}{3a^{\frac{1}{3}}} + \frac{\ln((bx+a)^{\frac{2}{3}}+a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}}+a^{\frac{2}{3}})}{6a^{\frac{1}{3}}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(\frac{2(bx+a)^{\frac{1}{3}}}{a^{\frac{1}{3}}}\right)}{3a^{\frac{1}{3}}}\right)}{3a} \right)}{3a} \right)$

input `int(1/x^3/(b*x+a)^(1/3),x,method=_RETURNVERBOSE)`

output
$$-1/6*(b*x+a)^{(2/3)}*(-4*b*x+3*a)/a^2/x^2+2/9*b^2/a^{(7/3)}*\ln(-a^{(1/3)}+(b*x+a)^{(1/3)})-1/9*b^2/a^{(7/3)}*\ln((b*x+a)^{(2/3)}+a^{(1/3)}*(b*x+a)^{(1/3)}+a^{(2/3)})+2/9*b^2/a^{(7/3)}*3^{(1/2)}*\arctan(1/3*3^{(1/2)}*(2/a^{(1/3)}*(b*x+a)^{(1/3)}+1))$$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 296, normalized size of antiderivative = 2.53

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx$$

$$= \frac{6 \sqrt{\frac{1}{3}} ab^2 x^2 \sqrt{-\frac{1}{a^{\frac{2}{3}}}} \log \left(\frac{2bx+3 \sqrt{\frac{1}{3}} (2(bx+a)^{\frac{2}{3}} a^{\frac{2}{3}} - (bx+a)^{\frac{1}{3}} a - a^{\frac{4}{3}}) \sqrt{-\frac{1}{a^{\frac{2}{3}}}} - 3(bx+a)^{\frac{1}{3}} a^{\frac{2}{3}} + 3a}{x} \right) - 2a^{\frac{2}{3}} b^2 x^2 \log((bx + a)^{\frac{2}{3}} + a^{\frac{1}{3}}(bx+a)^{\frac{1}{3}} + a^{\frac{2}{3}})}{18 a^3 x^2}$$

input `integrate(1/x^3/(b*x+a)^(1/3),x, algorithm="fricas")`

output
$$\left[\frac{1}{18} * (6 * \sqrt{1/3} * a * b^2 * x^2 * \sqrt{-1/a^{(2/3)}}) * \log((2 * b * x + 3 * \sqrt{1/3}) * (2 * (b * x + a)^{(2/3)} * a^{(2/3)} - (b * x + a)^{(1/3)} * a - a^{(4/3)}) * \sqrt{-1/a^{(2/3)}} - 3 * (b * x + a)^{(1/3)} * a^{(2/3)} + 3 * a) / x) - 2 * a^{(2/3)} * b^2 * x^2 * \log((b * x + a)^{(2/3)} + (b * x + a)^{(1/3)} * a^{(1/3)} + a^{(2/3)}) + 4 * a^{(2/3)} * b^2 * x^2 * \log((b * x + a)^{(1/3)} - a^{(1/3)} + 3 * (4 * a * b * x - 3 * a^2) * (b * x + a)^{(2/3)}) / (a^3 * x^2), \frac{1}{18} * (12 * \sqrt{1/3} * a^{(2/3)} * b^2 * x^2 * \arctan(\sqrt{1/3} * (2 * (b * x + a)^{(1/3)} + a^{(1/3)}) / a^{(1/3)}) - 2 * a^{(2/3)} * b^2 * x^2 * \log((b * x + a)^{(2/3)} + (b * x + a)^{(1/3)} * a^{(1/3)} + a^{(2/3)}) + 4 * a^{(2/3)} * b^2 * x^2 * \log((b * x + a)^{(1/3)} - a^{(1/3)} + 3 * (4 * a * b * x - 3 * a^2) * (b * x + a)^{(2/3)}) / (a^3 * x^2)) \right]$$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 2.31 (sec) , antiderivative size = 2730, normalized size of antiderivative = 23.33

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx = \text{Too large to display}$$

input `integrate(1/x**3/(b*x+a)**(1/3),x)`

output

```
4*a**(14/3)*b**(10/3)*(a/b + x)**(4/3)*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b
+ x)**(1/3)/a**(1/3))*gamma(2/3)/(27*a**7*b**(4/3)*(a/b + x)**(4/3)*exp(2
*I*pi/3)*gamma(5/3) - 81*a**6*b**(7/3)*(a/b + x)**(7/3)*exp(2*I*pi/3)*gamm
a(5/3) + 81*a**5*b**(10/3)*(a/b + x)**(10/3)*exp(2*I*pi/3)*gamma(5/3) - 27
*a**4*b**(13/3)*(a/b + x)**(13/3)*exp(2*I*pi/3)*gamma(5/3)) + 4*a**(14/3)*
b**(10/3)*(a/b + x)**(4/3)*exp(-2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(1/3
))*exp_polar(2*I*pi/3)/a**(1/3))*gamma(2/3)/(27*a**7*b**(4/3)*(a/b + x)**(4
/3)*exp(2*I*pi/3)*gamma(5/3) - 81*a**6*b**(7/3)*(a/b + x)**(7/3)*exp(2*I*
pi/3)*gamma(5/3) + 81*a**5*b**(10/3)*(a/b + x)**(10/3)*exp(2*I*pi/3)*gamma(
5/3) - 27*a**4*b**(13/3)*(a/b + x)**(13/3)*exp(2*I*pi/3)*gamma(5/3)) + 4*a
**(14/3)*b**(10/3)*(a/b + x)**(4/3)*log(1 - b**(1/3)*(a/b + x)**(1/3)*exp_
polar(4*I*pi/3)/a**(1/3))*gamma(2/3)/(27*a**7*b**(4/3)*(a/b + x)**(4/3)*ex
p(2*I*pi/3)*gamma(5/3) - 81*a**6*b**(7/3)*(a/b + x)**(7/3)*exp(2*I*pi/3)*g
amma(5/3) + 81*a**5*b**(10/3)*(a/b + x)**(10/3)*exp(2*I*pi/3)*gamma(5/3) -
27*a**4*b**(13/3)*(a/b + x)**(13/3)*exp(2*I*pi/3)*gamma(5/3)) - 12*a**(11
/3)*b**(13/3)*(a/b + x)**(7/3)*exp(2*I*pi/3)*log(1 - b**(1/3)*(a/b + x)**(
1/3)/a**(1/3))*gamma(2/3)/(27*a**7*b**(4/3)*(a/b + x)**(4/3)*exp(2*I*pi/3)
*gamma(5/3) - 81*a**6*b**(7/3)*(a/b + x)**(7/3)*exp(2*I*pi/3)*gamma(5/3) +
81*a**5*b**(10/3)*(a/b + x)**(10/3)*exp(2*I*pi/3)*gamma(5/3) - 27*a**4*b*
*(13/3)*(a/b + x)**(13/3)*exp(2*I*pi/3)*gamma(5/3)) - 12*a**(11/3)*b**(...
```

Maxima [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.21

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx = \frac{2\sqrt{3}b^2 \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{9a^{\frac{7}{3}}} - \frac{b^2 \log\left((bx+a)^{\frac{2}{3}} + (bx+a)^{\frac{1}{3}}a^{\frac{1}{3}} + a^{\frac{2}{3}}\right)}{9a^{\frac{7}{3}}} + \frac{2b^2 \log\left((bx+a)^{\frac{1}{3}} - a^{\frac{1}{3}}\right)}{9a^{\frac{7}{3}}} + \frac{4(bx+a)^{\frac{5}{3}}b^2 - 7(bx+a)^{\frac{2}{3}}ab^2}{6((bx+a)^2a^2 - 2(bx+a)a^3 + a^4)}$$

input `integrate(1/x^3/(b*x+a)^(1/3),x, algorithm="maxima")`output
$$\frac{2/9*\sqrt{3}*b^2*\arctan(1/3*\sqrt{3}*(2*(b*x + a)^{(1/3)} + a^{(1/3)})/a^{(1/3)})/a^{(7/3)} - 1/9*b^2*\log((b*x + a)^{(2/3)} + (b*x + a)^{(1/3)}*a^{(1/3)} + a^{(2/3)})/a^{(7/3)} + 2/9*b^2*\log((b*x + a)^{(1/3)} - a^{(1/3)})/a^{(7/3)} + 1/6*(4*(b*x + a)^{(5/3)}*b^2 - 7*(b*x + a)^{(2/3)}*a*b^2)/((b*x + a)^2*a^2 - 2*(b*x + a)*a^3 + a^4)}$$
Giac [A] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.11

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx = \frac{4\sqrt{3}b^3 \arctan\left(\frac{\sqrt{3}(2(bx+a)^{\frac{1}{3}}+a^{\frac{1}{3}})}{3a^{\frac{1}{3}}}\right)}{a^{\frac{7}{3}}} - \frac{2b^3 \log\left((bx+a)^{\frac{2}{3}} + (bx+a)^{\frac{1}{3}}a^{\frac{1}{3}} + a^{\frac{2}{3}}\right)}{a^{\frac{7}{3}}} + \frac{4b^3 \log\left(\left|(bx+a)^{\frac{1}{3}} - a^{\frac{1}{3}}\right|\right)}{a^{\frac{7}{3}}} + \frac{3(4(bx+a)^{\frac{5}{3}}b^3 - 7(bx+a)a^2b^2x^2)}{18b}$$

input `integrate(1/x^3/(b*x+a)^(1/3),x, algorithm="giac")`

output

```
1/18*(4*sqrt(3)*b^3*arctan(1/3*sqrt(3)*(2*(b*x + a)^(1/3) + a^(1/3))/a^(1/3)))/a^(7/3) - 2*b^3*log((b*x + a)^(2/3) + (b*x + a)^(1/3)*a^(1/3) + a^(2/3))/a^(7/3) + 4*b^3*log(abs((b*x + a)^(1/3) - a^(1/3)))/a^(7/3) + 3*(4*(b*x + a)^(5/3)*b^3 - 7*(b*x + a)^(2/3)*a*b^3)/(a^2*b^2*x^2)/b
```

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.56

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx = \frac{2b^2 \ln\left(\frac{(a+bx)^{1/3} - a^{1/3}}{9a^{7/3}}\right) - \frac{7b^2(a+bx)^{2/3}}{6a} - \frac{2b^2(a+bx)^{5/3}}{3a^2}}{\ln\left(\frac{4b^4(a+bx)^{1/3}}{9a^4} - \frac{(b^2 + \sqrt{3}b^2 i)^2}{9a^{11/3}}\right) (b^2 + \sqrt{3}b^2 i)} - \frac{b^2 \ln\left(\frac{4b^4(a+bx)^{1/3}}{9a^4} - \frac{9b^4\left(-\frac{1}{9} + \frac{\sqrt{3}i}{9}\right)^2}{a^{11/3}}\right) \left(-\frac{1}{9} + \frac{\sqrt{3}i}{9}\right)}{a^{7/3}}$$

input

```
int(1/(x^3*(a + b*x)^(1/3)),x)
```

output

```
(2*b^2*log((a + b*x)^(1/3) - a^(1/3)))/(9*a^(7/3)) - ((7*b^2*(a + b*x)^(2/3))/(6*a) - (2*b^2*(a + b*x)^(5/3))/(3*a^2))/((a + b*x)^2 - 2*a*(a + b*x) + a^2) - (log((4*b^4*(a + b*x)^(1/3))/(9*a^4) - (3^(1/2)*b^2*i + b^2)^2/(9*a^(11/3))))*(3^(1/2)*b^2*i + b^2)/(9*a^(7/3)) + (b^2*log((4*b^4*(a + b*x)^(1/3))/(9*a^4) - (9*b^4*((3^(1/2)*i)/9 - 1/9)^2)/a^(11/3))*((3^(1/2)*i)/9 - 1/9))/a^(7/3)
```

Reduce [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.76

$$\int \frac{1}{x^3 \sqrt[3]{a+bx}} dx = \frac{-4\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}+a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) b^2 x^2 + 4\sqrt{3} \operatorname{atan}\left(\frac{2(bx+a)^{\frac{1}{6}}-a^{\frac{1}{6}}}{a^{\frac{1}{6}}\sqrt{3}}\right) b^2 x^2 - 9a^{\frac{4}{3}}(bx+a)^{\frac{2}{3}} + 12a^{\frac{1}{3}}(bx+a)^{\frac{2}{3}} bx + 4}{=}$$

input `int(1/x^3/(b*x+a)^(1/3),x)`

output
$$\begin{aligned} & (-4\sqrt{3}\operatorname{atan}((2(a+bx)^{1/6}+a^{1/6})/(a^{1/6}\sqrt{3}))b^2x^2 \\ & + 4\sqrt{3}\operatorname{atan}((2(a+bx)^{1/6}-a^{1/6})/(a^{1/6}\sqrt{3})) \\ &)b^2x^2 - 9a^{1/3}(a+bx)^{2/3}a + 12a^{1/3}(a+bx)^{2/3} \\ & *bx + 4\log((a+bx)^{1/6}+a^{1/6})b^2x^2 + 4\log((a+bx)^{1/6} \\ & - a^{1/6})b^2x^2 - 2\log(-a^{1/6}(a+bx)^{1/6} + (a+bx)^{1/3} \\ & + a^{1/3})b^2x^2 - 2\log(a^{1/6}(a+bx)^{1/6} + (a+bx)^{1/3} \\ & + a^{1/3})b^2x^2)/(18a^{1/3}a^2x^2) \end{aligned}$$

3.163 $\int \frac{A+Bx}{\sqrt{a+bx}} dx$

Optimal result	1108
Mathematica [A] (verified)	1108
Rubi [A] (verified)	1109
Maple [A] (verified)	1110
Fricas [A] (verification not implemented)	1110
Sympy [A] (verification not implemented)	1111
Maxima [A] (verification not implemented)	1111
Giac [A] (verification not implemented)	1111
Mupad [B] (verification not implemented)	1112
Reduce [B] (verification not implemented)	1112

Optimal result

Integrand size = 15, antiderivative size = 44

$$\int \frac{A+Bx}{\sqrt{a+bx}} dx = \frac{2A\sqrt{a+bx}}{b} + \frac{2B\sqrt{a+bx}(-a + \frac{1}{3}(a+bx))}{b^2}$$

output `2*(b*x+a)^(1/2)*A/b+2*(b*x+a)^(1/2)*(-2/3*a+1/3*b*x)*B/b^2`

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.66

$$\int \frac{A+Bx}{\sqrt{a+bx}} dx = \frac{2\sqrt{a+bx}(3Ab - 2aB + bBx)}{3b^2}$$

input `Integrate[(A + B*x)/Sqrt[a + b*x],x]`

output `(2*Sqrt[a + b*x]*(3*A*b - 2*a*B + b*B*x))/(3*b^2)`

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.91, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {53, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{A + Bx}{\sqrt{a + bx}} dx$$

↓ 53

$$\int \left(\frac{Ab - aB}{b\sqrt{a + bx}} + \frac{B\sqrt{a + bx}}{b} \right) dx$$

↓ 2009

$$\frac{2\sqrt{a + bx}(Ab - aB)}{b^2} + \frac{2B(a + bx)^{3/2}}{3b^2}$$

input `Int[(A + B*x)/Sqrt[a + b*x],x]`

output `(2*(A*b - a*B)*Sqrt[a + b*x])/b^2 + (2*B*(a + b*x)^(3/2))/(3*b^2)`

Defintions of rubi rules used

rule 53 `Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]`

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.59

method	result	size
gospers	$\frac{2\sqrt{bx+a}(xbB+3Ab-2Ba)}{3b^2}$	26
trager	$\frac{2\sqrt{bx+a}(xbB+3Ab-2Ba)}{3b^2}$	26
risch	$\frac{2\sqrt{bx+a}(xbB+3Ab-2Ba)}{3b^2}$	26
pseudoelliptic	$\frac{2\left(\left(\frac{Bx}{3}+A\right)b-\frac{2Ba}{3}\right)\sqrt{bx+a}}{b^2}$	26
orering	$\frac{2\sqrt{bx+a}(xbB+3Ab-2Ba)}{3b^2}$	26
derivativdivides	$\frac{\frac{2B(bx+a)^{\frac{3}{2}}}{3}+2\sqrt{bx+a}Ab-2\sqrt{bx+a}Ba}{b^2}$	38
default	$\frac{2B(bx+a)^{\frac{3}{2}}+2\sqrt{bx+a}Ab-2\sqrt{bx+a}Ba}{b^2}$	38

input `int((B*x+A)/(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

output `2/3*(b*x+a)^(1/2)*(B*b*x+3*A*b-2*B*a)/b^2`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.57

$$\int \frac{A+Bx}{\sqrt{a+bx}} dx = \frac{2(Bbx-2Ba+3Ab)\sqrt{bx+a}}{3b^2}$$

input `integrate((B*x+A)/(b*x+a)^(1/2),x, algorithm="fricas")`

output `2/3*(B*b*x - 2*B*a + 3*A*b)*sqrt(b*x + a)/b^2`

Sympy [A] (verification not implemented)

Time = 0.41 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.20

$$\int \frac{A + Bx}{\sqrt{a + bx}} dx = \begin{cases} \frac{2A\sqrt{a+bx} + \frac{2B\left(-a\sqrt{a+bx} + \frac{(a+bx)^{\frac{3}{2}}}{3}\right)}{b}}{b} & \text{for } b \neq 0 \\ \frac{Ax + \frac{Bx^2}{2}}{\sqrt{a}} & \text{otherwise} \end{cases}$$

input `integrate((B*x+A)/(b*x+a)**(1/2),x)`output `Piecewise(((2*A*sqrt(a + b*x) + 2*B*(-a*sqrt(a + b*x) + (a + b*x)**(3/2)/3)/b)/b, Ne(b, 0)), ((A*x + B*x**2/2)/sqrt(a), True))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.89

$$\int \frac{A + Bx}{\sqrt{a + bx}} dx = \frac{2 \left(3\sqrt{bx + a}A + \frac{(bx+a)^{\frac{3}{2}} - 3\sqrt{bx+aa}B}{b} \right)}{3b}$$

input `integrate((B*x+A)/(b*x+a)^(1/2),x, algorithm="maxima")`output `2/3*(3*sqrt(b*x + a)*A + ((b*x + a)^(3/2) - 3*sqrt(b*x + a)*a)*B/b)/b`**Giac [A] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.89

$$\int \frac{A + Bx}{\sqrt{a + bx}} dx = \frac{2 \left(3\sqrt{bx + a}A + \frac{(bx+a)^{\frac{3}{2}} - 3\sqrt{bx+aa}B}{b} \right)}{3b}$$

input `integrate((B*x+A)/(b*x+a)^(1/2),x, algorithm="giac")`

output `2/3*(3*sqrt(b*x + a)*A + ((b*x + a)^(3/2) - 3*sqrt(b*x + a)*a)*B/b)/b`

Mupad [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.64

$$\int \frac{A + Bx}{\sqrt{a + bx}} dx = \frac{2\sqrt{a + bx}(3Ab - 3Ba + B(a + bx))}{3b^2}$$

input `int((A + B*x)/(a + b*x)^(1/2),x)`

output `(2*(a + b*x)^(1/2)*(3*A*b - 3*B*a + B*(a + b*x)))/(3*b^2)`

Reduce [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.36

$$\int \frac{A + Bx}{\sqrt{a + bx}} dx = \frac{2\sqrt{bx + a}(bx + a)}{3b}$$

input `int((B*x+A)/(b*x+a)^(1/2),x)`

output `(2*sqrt(a + b*x)*(a + b*x))/(3*b)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 1113
4.2 Links to plain text integration problems used in this report for each CAS . 1131

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leaf
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
      ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A",""}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal.
      ]
    ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "
    ,

```

```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]===Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]===Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]===Plus || Head[expn]===Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```


Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result    := ExpnType(result);
      ExpnType_optimal   := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#     is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well
    fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co
        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```



```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```

```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```



```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file