

Computer Algebra Independent Integration Tests

Summer 2024

0-Independent-test-suites/12-Welz-Problems

Nasser M. Abbasi

May 18, 2024

Compiled on May 18, 2024 at 6:40am

Contents

1	Introduction	6
1.1	Listing of CAS systems tested	7
1.2	Results	8
1.3	Time and leaf size Performance	12
1.4	Performance based on number of rules Rubi used	14
1.5	Performance based on number of steps Rubi used	15
1.6	Solved integrals histogram based on leaf size of result	16
1.7	Solved integrals histogram based on CPU time used	17
1.8	Leaf size vs. CPU time used	18
1.9	list of integrals with no known antiderivative	19
1.10	List of integrals solved by CAS but has no known antiderivative	19
1.11	list of integrals solved by CAS but failed verification	19
1.12	Timing	20
1.13	Verification	20
1.14	Important notes about some of the results	21
1.15	Current tree layout of integration tests	24
1.16	Design of the test system	25
2	detailed summary tables of results	26
2.1	List of integrals sorted by grade for each CAS	27
2.2	Detailed conclusion table per each integral for all CAS systems	32
2.3	Detailed conclusion table specific for Rubi results	62
3	Listing of integrals	66
3.1	$\int \frac{1}{\sqrt{1-ax}} dx$	71
3.2	$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$	76
3.3	$\int \frac{1}{(2x+\sqrt{1+x^2})^2} dx$	82
3.4	$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx$	89
3.5	$\int \frac{1}{(2\sqrt{x}+\sqrt{1+x})^2} dx$	96

3.6	$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx$	102
3.7	$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx$	109
3.8	$\int \frac{1}{(\sqrt{-1+x}+\sqrt{x})^2\sqrt{-1+x}} dx$	115
3.9	$\int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx$	120
3.10	$\int \frac{(\sqrt{x}-\sqrt{-1+x^2})^2}{(1+x-x^2)^2\sqrt{-1+x^2}} dx$	129
3.11	$\int \left(\frac{1}{\sqrt{2(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2(1+x)^2\sqrt{i+x^2}}} \right) dx$	138
3.12	$\int \frac{\sqrt{x^2+\sqrt{1+x^4}}}{(1+x)^2\sqrt{1+x^4}} dx$	144
3.13	$\int \frac{\sqrt{x^2+\sqrt{1+x^4}}}{(1+x)\sqrt{1+x^4}} dx$	151
3.14	$\int \frac{\sqrt{x^2+\sqrt{1+x^4}}}{\sqrt{1+x^4}} dx$	157
3.15	$\int \frac{\sqrt{-x^2+\sqrt{1+x^4}}}{\sqrt{1+x^4}} dx$	162
3.16	$\int \frac{(-1+x)^{3/2}+(1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx$	167
3.17	$\int (x + \sqrt{a+x^2})^b dx$	172
3.18	$\int (x - \sqrt{a+x^2})^b dx$	178
3.19	$\int \frac{(x+\sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx$	183
3.20	$\int \frac{(x-\sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx$	188
3.21	$\int \frac{1}{(a+be^{px})^2} dx$	193
3.22	$\int \frac{1}{(be^{-px}+ae^{px})^2} dx$	198
3.23	$\int \frac{x}{(be^{-px}+ae^{px})^2} dx$	203
3.24	$\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx$	209
3.25	$\int \frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a^2+x^2}} dx$	218
3.26	$\int \frac{\sqrt{bx+\sqrt{a+b^2x^2}}}{\sqrt{a+b^2x^2}} dx$	223
3.27	$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx$	228
3.28	$\int \frac{x}{\sqrt{x+\sqrt{a^2+x^2}}} dx$	235
3.29	$\int x^3 \log^3(2+x) \log(3+x) dx$	242
3.30	$\int \frac{(x+\sqrt{b+x^2})^a}{\sqrt{b+x^2}} dx$	252
3.31	$\int (x + \sqrt{b+x^2})^a dx$	257
3.32	$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx$	263
3.33	$\int \frac{1}{x\sqrt[3]{1-x^2}} dx$	268
3.34	$\int \frac{1}{x(1-x^2)^{2/3}} dx$	275
3.35	$\int \frac{1}{\sqrt[3]{1-x^3}} dx$	282

3.36	$\int \frac{1}{x \sqrt[3]{1-x^3}} dx$	287
3.37	$\int \frac{1}{(1+x) \sqrt[3]{1-x^3}} dx$	294
3.38	$\int \frac{x}{(1+x) \sqrt[3]{1-x^3}} dx$	300
3.39	$\int \frac{1}{x \sqrt[3]{2-3x+x^2}} dx$	306
3.40	$\int \frac{1}{\sqrt[3]{-5+7x-3x^2+x^3}} dx$	312
3.41	$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx$	319
3.42	$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx$	326
3.43	$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx$	333
3.44	$\int \frac{2-(1+k)x}{\sqrt[3]{(1-x)x(1-kx)(1-(1+k)x)}} dx$	339
3.45	$\int \frac{1-kx}{(1+(-2+k)x)((1-x)x(1-kx))^{2/3}} dx$	345
3.46	$\int \frac{a+bx+cx^2}{(1-x+x^2) \sqrt[3]{1-x^3}} dx$	353
3.47	$\int \frac{1}{(3-2x)^{11/2}(1+x+2x^2)^5} dx$	361
3.48	$\int \frac{1}{(3-2x)^{21/2}(1+x+2x^2)^{10}} dx$	376
3.49	$\int \frac{1}{(3-2x)^{41/2}(1+x+2x^2)^{20}} dx$	392
3.50	$\int \frac{1}{(3-2x+x^2)^{11/2}(1+x+2x^2)^5} dx$	407
3.51	$\int \frac{1}{(3-2x+x^2)^{21/2}(1+x+2x^2)^{10}} dx$	422
3.52	$\int \frac{-a-\sqrt{1+a^2}+x}{(-a+\sqrt{1+a^2}+x)\sqrt{(-a+x)(1+x^2)}} dx$	436
3.53	$\int \frac{a+bx}{\sqrt[3]{1-x^2(3+x^2)}} dx$	446
3.54	$\int \frac{a+bx}{(3-x^2) \sqrt[3]{1+x^2}} dx$	454
3.55	$\int \frac{1}{x \sqrt[3]{4-6x+3x^2}} dx$	462
3.56	$\int x \sqrt[3]{1-x^3} dx$	468
3.57	$\int \frac{\sqrt[3]{1-x^3}}{x} dx$	474
3.58	$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx$	481
3.59	$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx$	487
3.60	$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx$	494
3.61	$\int \frac{2+x}{(1+x+x^2) \sqrt[3]{2+x^3}} dx$	499
3.62	$\int \frac{3-3x+30x^2+160x^3}{9+24x-12x^2+80x^3+320x^4} dx$	504
3.63	$\int \frac{3+12x+20x^2}{9+24x-12x^2+80x^3+320x^4} dx$	509
3.64	$\int \frac{-84-576x-400x^2+2560x^3}{9+24x-12x^2+80x^3+320x^4} dx$	515
3.65	$\int \frac{\sqrt{1-x^4}}{1+x^4} dx$	522

3.66	$\int \frac{\sqrt{1+x^4}}{1-x^4} dx$	527
3.67	$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx$	533
3.68	$\int \frac{\sqrt{1+px^2-x^4}}{1+x^4} dx$	539
3.69	$\int \frac{a+bx}{(2-x^2)\sqrt[4]{-1+x^2}} dx$	545
3.70	$\int \frac{a+bx}{\sqrt[4]{-1-x^2(2+x^2)}} dx$	552
3.71	$\int \frac{a+bx}{\sqrt[4]{1-x^2(2-x^2)}} dx$	559
3.72	$\int \frac{a+bx}{\sqrt[4]{1+x^2(2+x^2)}} dx$	565
3.73	$\int \frac{x}{\sqrt{1-x^3(4-x^3)}} dx$	571
3.74	$\int \frac{x}{(4-dx^3)\sqrt{-1+dx^3}} dx$	578
3.75	$\int \frac{x}{\sqrt{-1+x^3(8+x^3)}} dx$	585
3.76	$\int \frac{x}{(8-dx^3)\sqrt{1+dx^3}} dx$	593
3.77	$\int \frac{1}{\sqrt[3]{1-3x^2(3-x^2)}} dx$	602
3.78	$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx$	608
3.79	$\int \frac{1}{\sqrt[3]{1-x^2(3+x^2)}} dx$	614
3.80	$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx$	621
3.81	$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx$	627
3.82	$\int \frac{-2+a+x}{(-a+x)\sqrt{(2-a)ax+(-1-2a+a^2)x^2+x^3}} dx$	634
3.83	$\int \frac{-a+(-1+2a)x}{(-a+x)\sqrt{a^2x-(-1+2a+a^2)x^2+(-1+2a)x^3}} dx$	642
3.84	$\int \frac{1-\sqrt[3]{2}x}{(2^{2/3}+x)\sqrt{1+x^3}} dx$	650
3.85	$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx$	656
3.86	$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx$	661
3.87	$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx$	668
3.88	$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx$	675
3.89	$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx$	682
3.90	$\int \frac{1-\sqrt{3}+x}{(1+\sqrt{3}+x)\sqrt{-4+4\sqrt{3}x^2+x^4}} dx$	689
3.91	$\int \frac{1+\sqrt{3}+x}{(1-\sqrt{3}+x)\sqrt{-4-4\sqrt{3}x^2+x^4}} dx$	695
3.92	$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx$	702
3.93	$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx$	707
3.94	$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx$	714

3.95	$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx$	720
3.96	$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx$	725
3.97	$\int \frac{1}{\sqrt[3]{1-x^3(1+x^3)}} dx$	733
3.98	$\int \frac{x}{\sqrt[3]{1-x^3(1+x^3)}} dx$	739
3.99	$\int \frac{x^2}{\sqrt[3]{1-x^3(1+x^3)}} dx$	749
3.100	$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx$	756
3.101	$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3(1+x^3)}} dx$	763
3.102	$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx$	770
3.103	$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx$	777
3.104	$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx$	782
3.105	$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx$	787
3.106	$\int (1-x^3)^{2/3} dx$	792
3.107	$\int \frac{(1-x^3)^{2/3}}{x} dx$	798
3.108	$\int \frac{(1-x^3)^{2/3}}{a+bx} dx$	805
3.109	$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$	812
3.110	$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$	817
3.111	$\int \frac{(1-x^3)^{2/3}}{1+x} dx$	824
3.112	$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx$	830
3.113	$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx$	837
3.114	$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx$	844
3.115	$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx$	854
3.116	$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx$	860
4	Appendix	870
4.1	Listing of Grading functions	870
4.2	Links to plain text integration problems used in this report for each CAS888	

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	7
1.2	Results	8
1.3	Time and leaf size Performance	12
1.4	Performance based on number of rules Rubi used	14
1.5	Performance based on number of steps Rubi used	15
1.6	Solved integrals histogram based on leaf size of result	16
1.7	Solved integrals histogram based on CPU time used	17
1.8	Leaf size vs. CPU time used	18
1.9	list of integrals with no known antiderivative	19
1.10	List of integrals solved by CAS but has no known antiderivative	19
1.11	list of integrals solved by CAS but failed verification	19
1.12	Timing	20
1.13	Verification	20
1.14	Important notes about some of the results	21
1.15	Current tree layout of integration tests	24
1.16	Design of the test system	25

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [116]. This is test number [12].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 14 (January 9, 2024) on windows 10 pro.
2. Rubi 4.17.3 (Sept 25, 2023) on Mathematica 14 on windows 10m pro.
3. Maple 2024 (March 1, 2024) on windows 10 pro.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.4.0 on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
5. FriCAS 1.3.10 built with sbcl 2.3.11 (January 10, 2024) on Linux Manjaro 23.1.2 KDE via sagemath 10.3.
6. Giac/Xcas 1.9.0-99 on Linux via sagemath 10.3.
7. Sympy 1.12 using Python 3.11.6 (Nov 14 2023, 09:36:21) [GCC 13.2.1 20230801] on Linux Manjaro 23.1.2 KDE.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.
9. Reduce CSL rev 6687 (January 9, 2024) on Linux Manjaro 23.1.2 KDE.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

Reduce was called directly.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	94.83 (110)	5.17 (6)
Mathematica	87.93 (102)	12.07 (14)
Fricas	77.59 (90)	22.41 (26)
Maple	75.86 (88)	24.14 (28)
Mupad	31.90 (37)	68.10 (79)
Giac	31.03 (36)	68.97 (80)
Sympy	25.00 (29)	75.00 (87)
Reduce	23.28 (27)	76.72 (89)
Maxima	17.24 (20)	82.76 (96)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	87.931	3.448	3.448	5.172
Mathematica	55.172	2.586	30.172	12.069
Fricas	36.207	39.655	1.724	22.414
Maple	32.759	6.034	37.069	24.138
Giac	20.690	9.483	0.862	68.966
Maxima	15.517	1.724	0.000	82.759
Sympy	10.345	6.034	8.621	75.000
Mupad	0.000	31.897	0.000	68.103
Reduce	0.000	23.276	0.000	76.724

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	6	100.00	0.00	0.00
Mathematica	14	100.00	0.00	0.00
Fricas	26	42.31	26.92	30.77
Maple	28	100.00	0.00	0.00
Mupad	79	0.00	100.00	0.00
Giac	80	98.75	1.25	0.00
Sympy	87	90.80	8.05	1.15
Reduce	89	100.00	0.00	0.00
Maxima	96	98.96	0.00	1.04

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.08
Reduce	0.17
Rubi	0.37
Mupad	0.51
Sympy	0.67
Giac	1.29
Fricas	2.40
Mathematica	3.79
Maple	5.74

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Maxima	81.80	1.16	62.00	1.05
Mathematica	139.19	1.20	83.50	1.08
Rubi	151.02	1.29	90.00	1.00
Mupad	151.70	1.71	76.00	1.09
Sympy	208.41	5.09	37.00	0.82
Giac	247.72	1.54	69.50	1.10
Maple	321.58	2.97	111.00	1.18
Fricas	398.66	3.00	210.00	1.67
Reduce	1378.59	2.91	37.00	1.29

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Reduce {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {40, 41, 42, 82}

Mathematica {53, 69, 70, 71, 72, 74, 75, 77, 78, 79, 80, 88, 89, 106, 115}

Maple {37, 38, 39, 55, 74, 76, 77, 100, 101, 102, 110, 116}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Reduce Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'  
'display2d : false'  
'domain : complex'  
'keepfloat : true'  
'load(to_poly_solve)'  
'load(simplify_sum)'  
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib  
maxima_lib.set('extra_definite_integration_methods', '[]')  
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:  
    # 1.7 is a fudge factor since it is low side from actual leaf count  
    leafCount = round(1.7*count_ops(anti))  
  
except Exception as ee:  
    leafCount = 1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')  
the_variable = evalin(symengine, 'x')  
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Current tree layout of integration tests





Figure 1.6: CAS integration tests tree

1.16 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "E"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
January 13, 2024
Design note

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	27
2.2	Detailed conclusion table per each integral for all CAS systems	32
2.3	Detailed conclusion table specific for Rubi results	62

2.1 List of integrals sorted by grade for each CAS

Rubi	27
Mma	27
Maple	28
Fricas	28
Maxima	29
Giac	29
Mupad	30
Sympy	30
Reduce	30

Rubi

A grade { 1, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 46, 47, 50, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116 }

B grade { 10, 100, 101, 102 }

C grade { 2, 52, 82, 83 }

F normal fail { 43, 44, 45, 48, 49, 51 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 39, 41, 42, 43, 44, 52, 55, 56, 57, 62, 66, 67, 81, 82, 83, 84, 85, 90, 91, 92, 97, 98, 99, 100, 101, 102, 103, 104, 105, 107, 113, 116 }

B grade { 12, 13, 54 }

C grade { 2, 15, 24, 40, 47, 48, 49, 50, 51, 53, 63, 64, 65, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 86, 87, 88, 89, 94, 96, 106, 114, 115 }

F normal fail { 38, 45, 46, 58, 59, 60, 61, 93, 95, 108, 109, 110, 111, 112 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 4, 6, 7, 8, 11, 16, 19, 20, 21, 22, 23, 25, 26, 30, 32, 33, 36, 41, 44, 47, 48, 49, 50, 51, 62, 63, 66, 67, 81, 94, 96, 97, 99, 103, 104, 105, 113 }

B grade { 5, 9, 10, 17, 24, 31, 85 }

C grade { 2, 3, 15, 28, 34, 35, 37, 38, 39, 40, 52, 55, 56, 57, 59, 64, 65, 68, 73, 74, 75, 76, 77, 78, 79, 82, 83, 84, 86, 87, 88, 89, 90, 91, 92, 93, 100, 101, 102, 106, 107, 110, 116 }

F normal fail { 12, 13, 14, 18, 27, 29, 42, 43, 45, 46, 53, 54, 58, 60, 61, 69, 70, 71, 72, 80, 95, 98, 108, 109, 111, 112, 114, 115 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 3, 5, 6, 8, 11, 15, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 28, 30, 31, 32, 33, 34, 36, 40, 52, 56, 57, 62, 63, 64, 65, 66, 67, 81, 83, 98, 99, 106, 107, 113, 116 }

B grade { 4, 7, 9, 10, 12, 13, 14, 24, 27, 35, 37, 39, 41, 42, 43, 45, 47, 48, 49, 50, 51, 55, 59, 73, 74, 75, 76, 77, 78, 79, 80, 85, 86, 87, 88, 89, 90, 91, 93, 94, 96, 97, 100, 101, 102, 110 }

C grade { 68, 82 }

F normal fail { 2, 29, 61, 103, 104, 105, 109, 111, 112, 114, 115 }

F(-1) timedout fail { 44, 69, 70, 71, 72, 95, 108 }

F(-2) exception fail { 38, 46, 53, 54, 58, 60, 84, 92 }

Maxima

A grade { 1, 6, 16, 21, 22, 23, 29, 32, 33, 34, 35, 36, 56, 57, 62, 96, 99, 107 }

B grade { 2, 106 }

C grade { }

F normal fail { 3, 4, 5, 7, 8, 9, 10, 12, 13, 14, 15, 17, 18, 19, 20, 24, 25, 26, 27, 28, 30, 31, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 58, 59, 60, 61, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 97, 98, 100, 101, 102, 103, 104, 105, 108, 109, 110, 111, 112, 113, 114, 115, 116 }

F(-1) timedout fail { }

F(-2) exception fail { 11 }

Giac

A grade { 1, 6, 8, 16, 19, 20, 21, 22, 23, 25, 26, 30, 33, 34, 36, 41, 49, 57, 62, 63, 64, 96, 99, 107 }

B grade { 2, 3, 4, 5, 7, 9, 10, 11, 24, 47, 48 }

C grade { 50 }

F normal fail { 12, 13, 14, 15, 17, 18, 27, 28, 29, 31, 32, 35, 37, 38, 39, 40, 42, 43, 44, 45, 46, 52, 53, 54, 55, 56, 58, 59, 60, 61, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 97, 98, 100, 101, 102, 103, 104, 105, 106, 108, 109, 110, 111, 112, 113, 114, 115, 116 }

F(-1) timedout fail { 51 }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 5, 8, 16, 19, 20, 21, 22, 23, 25, 26, 30, 33, 34, 35, 36, 41, 47, 48, 49, 57, 62, 63, 64, 73, 74, 75, 81, 82, 84, 85, 96, 99, 106, 107 }

C grade { }

F normal fail { }

F(-1) timedout fail { 4, 6, 7, 9, 10, 11, 12, 13, 14, 15, 17, 18, 24, 27, 28, 29, 31, 32, 37, 38, 39, 40, 42, 43, 44, 45, 46, 50, 51, 52, 53, 54, 55, 56, 58, 59, 60, 61, 65, 66, 67, 68, 69, 70, 71, 72, 76, 77, 78, 79, 80, 83, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 97, 98, 100, 101, 102, 103, 104, 105, 108, 109, 110, 111, 112, 113, 114, 115, 116 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 14, 15, 21, 22, 23, 25, 26, 62, 63, 64 }

B grade { 8, 16, 17, 19, 20, 30, 31 }

C grade { 27, 28, 33, 34, 35, 36, 56, 57, 106, 107 }

F normal fail { 3, 4, 5, 6, 7, 9, 12, 13, 18, 24, 29, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 50, 53, 54, 55, 58, 59, 60, 61, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 108, 109, 110, 111, 112, 113, 114, 115, 116 }

F(-1) timedout fail { 10, 32, 47, 48, 49, 51, 52 }

F(-2) exception fail { 11 }

Reduce

A grade { }

B grade { 1, 2, 3, 4, 5, 7, 8, 14, 16, 17, 18, 19, 20, 21, 22, 23, 25, 26, 27, 28, 30, 31, 32, 47, 48, 49, 62 }

C grade { }

F normal fail { 6, 9, 10, 11, 12, 13, 15, 24, 29, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74,

75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99,
100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	12	13	12	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.80	0.87	0.80	0.87
time (sec)	N/A	0.141	0.002	0.105	0.024	0.067	0.017	0.120	0.154	0.016

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	C	C	B	F	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	49	37	34	41	0	42	41	31	43
N.S.	1	3.27	2.47	2.27	2.73	0.00	2.80	2.73	2.07	2.87
time (sec)	N/A	0.271	0.020	0.214	0.025	0.000	3.959	0.128	0.155	0.289

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	60	73	0	100	0	177	107	204
N.S.	1	1.00	0.73	0.89	0.00	1.22	0.00	2.16	1.30	2.49
time (sec)	N/A	0.291	0.191	0.217	0.000	0.068	0.000	0.129	0.148	0.277

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	71	52	0	80	0	94	160	0
N.S.	1	1.00	1.65	1.21	0.00	1.86	0.00	2.19	3.72	0.00
time (sec)	N/A	0.181	0.076	0.445	0.000	0.066	0.000	0.123	0.149	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	F	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	73	115	0	105	0	133	113	82
N.S.	1	1.00	0.99	1.55	0.00	1.42	0.00	1.80	1.53	1.11
time (sec)	N/A	0.281	0.066	0.030	0.000	0.074	0.000	0.150	0.153	1.145

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	70	65	53	89	0	84	24	0
N.S.	1	1.00	1.09	1.02	0.83	1.39	0.00	1.31	0.38	0.00
time (sec)	N/A	0.210	0.182	0.284	0.105	0.074	0.000	0.132	0.158	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	B	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	55	37	0	83	0	101	157	0
N.S.	1	1.00	1.15	0.77	0.00	1.73	0.00	2.10	3.27	0.00
time (sec)	N/A	0.168	0.100	0.246	0.000	0.069	0.000	0.128	0.156	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	26	21	0	18	53	20	19	21
N.S.	1	1.00	0.87	0.70	0.00	0.60	1.77	0.67	0.63	0.70
time (sec)	N/A	0.285	0.176	0.084	0.000	0.064	0.224	0.127	0.156	0.364

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	220	365	195	1206	0	341	0	367	605	0
N.S.	1	1.66	0.89	5.48	0.00	1.55	0.00	1.67	2.75	0.00
time (sec)	N/A	0.776	6.190	0.207	0.000	0.089	0.000	1.314	0.237	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	B	F	B	F(-1)	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	220	541	195	1636	0	341	0	358	977	0
N.S.	1	2.46	0.89	7.44	0.00	1.55	0.00	1.63	4.44	0.00
time (sec)	N/A	1.128	6.324	0.809	0.000	0.082	0.000	1.417	0.183	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-2)	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	138	126	146	0	161	0	547	39	0
N.S.	1	1.00	0.91	1.06	0.00	1.17	0.00	3.96	0.28	0.00
time (sec)	N/A	0.250	3.630	0.602	0.000	0.093	0.000	0.144	180.017	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	125	151	272	0	0	342	0	0	41	0
N.S.	1	1.21	2.18	0.00	0.00	2.74	0.00	0.00	0.33	0.00
time (sec)	N/A	0.388	2.338	0.000	0.000	1.071	0.000	0.000	0.166	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	205	0	0	332	0	0	31	0
N.S.	1	1.00	2.53	0.00	0.00	4.10	0.00	0.00	0.38	0.00
time (sec)	N/A	0.347	1.086	0.000	0.000	3.270	0.000	0.000	0.160	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	A	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	44	0	0	60	15	0	42	0
N.S.	1	1.00	1.42	0.00	0.00	1.94	0.48	0.00	1.35	0.00
time (sec)	N/A	0.225	0.132	0.000	0.000	0.207	0.906	0.000	0.159	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	57	22	0	42	15	0	29	0
N.S.	1	1.00	1.73	0.67	0.00	1.27	0.45	0.00	0.88	0.00
time (sec)	N/A	0.232	0.159	0.120	0.000	0.216	0.416	0.000	0.159	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	16	15	28	56	15	26	15
N.S.	1	1.00	1.00	0.84	0.79	1.47	2.95	0.79	1.37	0.79
time (sec)	N/A	0.421	0.009	0.243	0.027	0.076	0.658	0.136	0.148	0.281

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	51	43	120	0	32	2236	0	30	0
N.S.	1	0.98	0.83	2.31	0.00	0.62	43.00	0.00	0.58	0.00
time (sec)	N/A	0.204	0.128	0.042	0.000	0.094	1.506	0.000	0.143	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	56	55	50	0	0	33	0	0	37	0
N.S.	1	0.98	0.89	0.00	0.00	0.59	0.00	0.00	0.66	0.00
time (sec)	N/A	0.208	0.137	0.000	0.000	0.091	0.000	0.000	0.169	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	0	15	311	15	14	15
N.S.	1	1.00	1.00	0.94	0.00	0.88	18.29	0.88	0.82	0.88
time (sec)	N/A	0.215	0.025	0.093	0.000	0.074	1.309	0.120	0.155	0.180

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	0	18	41	18	23	18
N.S.	1	1.00	1.00	0.95	0.00	0.90	2.05	0.90	1.15	0.90
time (sec)	N/A	0.230	0.024	0.153	0.000	0.079	0.846	0.123	0.148	0.188

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	45	39	43	40	52	36	41	71	58
N.S.	1	1.07	0.93	1.02	0.95	1.24	0.86	0.98	1.69	1.38
time (sec)	N/A	0.211	0.040	0.056	0.042	0.075	0.056	0.120	0.154	0.243

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	20	20	19	20	19	26	24
N.S.	1	1.00	1.00	0.91	0.91	0.86	0.91	0.86	1.18	1.09
time (sec)	N/A	0.181	0.021	0.060	0.034	0.067	0.043	0.122	0.157	0.223

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	49	50	51	58	51	74	69	47
N.S.	1	1.00	0.79	0.81	0.82	0.94	0.82	1.19	1.11	0.76
time (sec)	N/A	0.308	0.070	0.119	0.042	0.074	0.072	0.123	0.155	0.235

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	B	F	B	F	B	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	94	239	158	0	245	0	304	283	0
N.S.	1	1.09	2.78	1.84	0.00	2.85	0.00	3.53	3.29	0.00
time (sec)	N/A	0.308	0.300	1.258	0.000	0.081	0.000	0.143	0.203	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	19	16	0	15	15	15	13	15
N.S.	1	1.00	1.00	0.84	0.00	0.79	0.79	0.79	0.68	0.79
time (sec)	N/A	0.205	0.031	0.111	0.000	0.065	0.082	0.126	0.157	0.256

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	23	0	22	27	22	20	22
N.S.	1	1.00	1.00	0.88	0.00	0.85	1.04	0.85	0.77	0.85
time (sec)	N/A	0.226	0.042	0.144	0.000	0.072	0.396	0.119	0.162	0.276

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	C	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	63	69	56	0	0	237	46	0	102	0
N.S.	1	1.10	0.89	0.00	0.00	3.76	0.73	0.00	1.62	0.00
time (sec)	N/A	0.344	0.066	0.000	0.000	0.089	1.032	0.000	0.162	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	C	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	92	82	25	0	257	51	0	117	0
N.S.	1	1.12	1.00	0.30	0.00	3.13	0.62	0.00	1.43	0.00
time (sec)	N/A	0.229	0.062	0.029	0.000	0.088	1.654	0.000	0.153	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	606	741	412	0	518	0	0	0	425	0
N.S.	1	1.22	0.68	0.00	0.85	0.00	0.00	0.00	0.70	0.00
time (sec)	N/A	4.787	0.184	0.000	0.054	0.000	0.000	0.000	0.158	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	0	15	311	15	14	15
N.S.	1	1.00	1.00	0.94	0.00	0.88	18.29	0.88	0.82	0.88
time (sec)	N/A	0.190	0.026	0.118	0.000	0.069	1.223	0.118	0.147	0.178

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	A	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	51	46	120	0	32	2236	0	30	0
N.S.	1	0.98	0.88	2.31	0.00	0.62	43.00	0.00	0.58	0.00
time (sec)	N/A	0.191	0.133	0.039	0.000	0.094	1.164	0.000	0.154	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	33	44	48	48	0	0	44	0
N.S.	1	1.00	0.97	1.29	1.41	1.41	0.00	0.00	1.29	0.00
time (sec)	N/A	0.224	0.124	0.878	0.098	0.080	0.000	0.000	0.160	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	61	77	63	62	64	36	64	15	86
N.S.	1	1.05	1.33	1.09	1.07	1.10	0.62	1.10	0.26	1.48
time (sec)	N/A	0.178	0.044	1.589	0.118	0.071	0.399	0.127	0.157	0.311

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	62	77	48	62	64	37	64	15	76
N.S.	1	1.07	1.33	0.83	1.07	1.10	0.64	1.10	0.26	1.31
time (sec)	N/A	0.177	0.035	1.627	0.113	0.069	0.432	0.131	0.161	0.260

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	B	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	86	12	78	82	29	0	11	10
N.S.	1	1.00	1.76	0.24	1.59	1.67	0.59	0.00	0.22	0.20
time (sec)	N/A	0.148	0.028	2.067	0.111	0.082	0.392	0.000	0.156	0.186

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	61	79	63	62	64	32	63	15	80
N.S.	1	1.11	1.44	1.15	1.13	1.16	0.58	1.15	0.27	1.45
time (sec)	N/A	0.180	0.034	2.253	0.111	0.068	0.405	0.132	0.158	0.286

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	148	1143	0	298	0	0	25	0
N.S.	1	1.00	1.53	11.78	0.00	3.07	0.00	0.00	0.26	0.00
time (sec)	N/A	0.208	0.938	7.403	0.000	1.759	0.000	0.000	0.154	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	N/A	No	TBD	TBD	TBD	TBD	TBD	TBD
size	145	145	0	1421	0	0	0	0	27	0
N.S.	1	1.00	0.00	9.80	0.00	0.00	0.00	0.00	0.19	0.00
time (sec)	N/A	0.312	0.000	16.865	0.000	0.000	0.000	0.000	0.161	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	110	120	162	1592	0	274	0	0	16	0
N.S.	1	1.09	1.47	14.47	0.00	2.49	0.00	0.00	0.15	0.00
time (sec)	N/A	0.193	0.172	7.765	0.000	0.998	0.000	0.000	0.159	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	F	F	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	86	85	653	0	120	0	0	17	0
N.S.	1	1.06	1.05	8.06	0.00	1.48	0.00	0.00	0.21	0.00
time (sec)	N/A	0.217	0.010	1.166	0.000	0.230	0.000	0.000	0.157	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	A	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	86	130	92	0	415	0	67	15	37
N.S.	1	1.30	1.97	1.39	0.00	6.29	0.00	1.02	0.23	0.56
time (sec)	N/A	0.227	0.571	0.752	0.000	0.768	0.000	0.127	0.169	0.229

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	B	F	F	F	F(-1)
verified	N/A	No	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	94	145	0	0	665	0	0	22	0
N.S.	1	1.19	1.84	0.00	0.00	8.42	0.00	0.00	0.28	0.00
time (sec)	N/A	0.233	0.647	0.000	0.000	0.628	0.000	0.000	0.150	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	F	F	B	F	F	F	F(-1)
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	118	0	189	0	0	1496	0	0	30	0
N.S.	1	0.00	1.60	0.00	0.00	12.68	0.00	0.00	0.25	0.00
time (sec)	N/A	0.000	2.089	0.000	0.000	13.548	0.000	0.000	0.151	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	A	A	F	F(-1)	F	F	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	0	128	111	0	0	0	0	210	0
N.S.	1	0.00	1.15	1.00	0.00	0.00	0.00	0.00	1.89	0.00
time (sec)	N/A	0.000	15.185	1.243	0.000	0.000	0.000	0.000	0.280	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	F	F	F	B	F	F	F	F(-1)
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	176	0	0	0	0	932	0	0	140	0
N.S.	1	0.00	0.00	0.00	0.00	5.30	0.00	0.00	0.80	0.00
time (sec)	N/A	0.000	0.000	0.000	0.000	52.261	0.000	0.000	0.205	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	493	570	0	0	0	0	0	0	130	0
N.S.	1	1.16	0.00	0.00	0.00	0.00	0.00	0.00	0.26	0.00
time (sec)	N/A	0.777	0.000	0.000	0.000	0.000	0.000	0.000	0.161	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	B	F(-1)	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	407	506	178	277	0	636	0	795	4694	343
N.S.	1	1.24	0.44	0.68	0.00	1.56	0.00	1.95	11.53	0.84
time (sec)	N/A	0.878	4.255	2.543	0.000	0.095	0.000	0.447	0.167	0.273

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	A	F	B	F(-1)	B	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	648	0	253	352	0	1086	0	1000	10154	567
N.S.	1	0.00	0.39	0.54	0.00	1.68	0.00	1.54	15.67	0.88
time (sec)	N/A	0.000	6.725	3.135	0.000	0.121	0.000	0.598	0.220	0.318

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	A	F	B	F(-1)	A	B	B
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1058	0	1242	502	0	1986	0	1410	21074	1017
N.S.	1	0.00	1.17	0.47	0.00	1.88	0.00	1.33	19.92	0.96
time (sec)	N/A	0.000	36.087	8.081	0.000	0.317	0.000	0.933	0.593	0.683

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	B	F	C	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	378	424	733	452	0	1014	0	2509	124	0
N.S.	1	1.12	1.94	1.20	0.00	2.68	0.00	6.64	0.33	0.00
time (sec)	N/A	1.327	2.582	2.503	0.000	0.101	0.000	34.719	0.222	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	F	C	A	F	B	F(-1)	F(-1)	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	638	0	1431	552	0	1714	0	0	224	0
N.S.	1	0.00	2.24	0.87	0.00	2.69	0.00	0.00	0.35	0.00
time (sec)	N/A	0.000	21.416	5.678	0.000	0.182	0.000	0.000	0.437	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	C	F	A	F(-1)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	429	99	787	0	546	0	0	337	0
N.S.	1	6.50	1.50	11.92	0.00	8.27	0.00	0.00	5.11	0.00
time (sec)	N/A	1.204	1.313	1.842	0.000	0.122	0.000	0.000	0.987	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	198	203	145	0	0	0	0	0	65	0
N.S.	1	1.03	0.73	0.00	0.00	0.00	0.00	0.00	0.33	0.00
time (sec)	N/A	0.284	10.154	0.000	0.000	0.000	0.000	0.000	0.164	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	B	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	198	198	1072	0	0	0	0	0	59	0
N.S.	1	1.00	5.41	0.00	0.00	0.00	0.00	0.00	0.30	0.00
time (sec)	N/A	0.268	9.752	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	168	1593	0	171	0	0	18	0
N.S.	1	1.00	1.73	16.42	0.00	1.76	0.00	0.00	0.19	0.00
time (sec)	N/A	0.172	0.184	12.414	0.000	1.066	0.000	0.000	0.170	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	76	99	15	105	96	32	0	37	0
N.S.	1	1.04	1.36	0.21	1.44	1.32	0.44	0.00	0.51	0.00
time (sec)	N/A	0.167	0.146	1.512	0.116	0.070	0.497	0.000	0.154	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	75	88	49	71	73	37	72	33	83
N.S.	1	1.12	1.31	0.73	1.06	1.09	0.55	1.07	0.49	1.24
time (sec)	N/A	0.187	0.048	1.952	0.105	0.069	0.518	0.123	0.155	0.210

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	482	482	0	0	0	0	0	0	67	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.665	0.000	0.000	0.000	0.000	0.000	0.000	0.159	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	280	408	0	925	0	3744	0	0	22	0
N.S.	1	1.46	0.00	3.30	0.00	13.37	0.00	0.00	0.08	0.00
time (sec)	N/A	0.557	0.000	9.348	0.000	5.063	0.000	0.000	0.161	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	232	232	0	0	0	0	0	0	71	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.31	0.00
time (sec)	N/A	0.395	0.000	0.000	0.000	0.000	0.000	0.000	0.154	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	168	168	0	0	0	0	0	0	69	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.41	0.00
time (sec)	N/A	0.344	0.000	0.000	0.000	0.000	0.000	0.000	0.180	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	23	23	22	23	23	23
N.S.	1	1.00	1.00	0.88	0.92	0.92	0.88	0.92	0.92	0.92
time (sec)	N/A	0.171	0.005	0.065	0.026	0.062	0.035	0.122	0.154	0.039

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	86	52	0	43	73	40	85	53
N.S.	1	1.00	1.46	0.88	0.00	0.73	1.24	0.68	1.44	0.90
time (sec)	N/A	0.184	0.014	0.099	0.000	0.068	0.066	0.129	0.159	0.200

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	85	99	70	0	66	100	64	108	76
N.S.	1	1.09	1.27	0.90	0.00	0.85	1.28	0.82	1.38	0.97
time (sec)	N/A	0.255	0.013	0.093	0.000	0.068	0.074	0.133	0.151	0.211

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	57	56	0	56	0	0	18	0
N.S.	1	1.00	1.16	1.14	0.00	1.14	0.00	0.00	0.37	0.00
time (sec)	N/A	0.149	0.143	1.790	0.000	0.107	0.000	0.000	0.145	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	44	62	0	61	0	0	18	0
N.S.	1	1.00	0.83	1.17	0.00	1.15	0.00	0.00	0.34	0.00
time (sec)	N/A	0.154	0.172	1.550	0.000	0.107	0.000	0.000	0.144	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	101	79	89	0	345	0	0	23	0
N.S.	1	1.35	1.05	1.19	0.00	4.60	0.00	0.00	0.31	0.00
time (sec)	N/A	0.258	0.250	2.428	0.000	0.143	0.000	0.000	0.179	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	C	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	92	107	0	229	0	0	23	0
N.S.	1	1.00	0.54	0.63	0.00	1.34	0.00	0.00	0.13	0.00
time (sec)	N/A	0.283	0.241	1.343	0.000	0.154	0.000	0.000	0.155	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	80	88	157	0	0	0	0	0	59	0
N.S.	1	1.10	1.96	0.00	0.00	0.00	0.00	0.00	0.74	0.00
time (sec)	N/A	0.211	10.192	0.000	0.000	0.000	0.000	0.000	0.159	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	88	96	162	0	0	0	0	0	62	0
N.S.	1	1.09	1.84	0.00	0.00	0.00	0.00	0.00	0.70	0.00
time (sec)	N/A	0.210	10.178	0.000	0.000	0.000	0.000	0.000	0.160	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	149	151	144	0	0	0	0	0	67	0
N.S.	1	1.01	0.97	0.00	0.00	0.00	0.00	0.00	0.45	0.00
time (sec)	N/A	0.221	10.188	0.000	0.000	0.000	0.000	0.000	0.173	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	135	137	152	0	0	0	0	0	57	0
N.S.	1	1.01	1.13	0.00	0.00	0.00	0.00	0.00	0.42	0.00
time (sec)	N/A	0.210	10.142	0.000	0.000	0.000	0.000	0.000	0.156	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	127	127	28	164	0	1019	0	0	24	653
N.S.	1	1.00	0.22	1.29	0.00	8.02	0.00	0.00	0.19	5.14
time (sec)	N/A	0.182	10.019	11.852	0.000	0.145	0.000	0.000	0.162	0.249

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	B
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	157	157	54	240	0	1669	0	0	31	331
N.S.	1	1.00	0.34	1.53	0.00	10.63	0.00	0.00	0.20	2.11
time (sec)	N/A	0.203	10.024	0.883	0.000	0.212	0.000	0.000	0.162	8.403

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	48	421	0	261	0	0	22	533
N.S.	1	1.00	0.65	5.69	0.00	3.53	0.00	0.00	0.30	7.20
time (sec)	N/A	0.422	10.021	8.427	0.000	0.129	0.000	0.000	0.151	0.124

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	32	383	0	491	0	0	31	0
N.S.	1	1.00	0.31	3.72	0.00	4.77	0.00	0.00	0.30	0.00
time (sec)	N/A	0.642	10.023	1.681	0.000	0.237	0.000	0.000	0.143	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	126	438	0	956	0	0	31	0
N.S.	1	1.00	1.56	5.41	0.00	11.80	0.00	0.00	0.38	0.00
time (sec)	N/A	0.157	4.158	8.730	0.000	1.100	0.000	0.000	0.176	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	126	443	0	345	0	0	29	0
N.S.	1	1.00	1.56	5.47	0.00	4.26	0.00	0.00	0.36	0.00
time (sec)	N/A	0.162	3.923	2.066	0.000	0.859	0.000	0.000	0.173	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	118	938	0	1132	0	0	29	0
N.S.	1	1.00	1.04	8.30	0.00	10.02	0.00	0.00	0.26	0.00
time (sec)	N/A	0.170	3.925	14.201	0.000	0.501	0.000	0.000	0.163	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	109	109	124	0	0	1103	0	0	27	0
N.S.	1	1.00	1.14	0.00	0.00	10.12	0.00	0.00	0.25	0.00
time (sec)	N/A	0.169	4.052	0.000	0.000	0.492	0.000	0.000	0.161	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	29	33	0	85	0	0	130	217
N.S.	1	1.00	0.33	0.38	0.00	0.98	0.00	0.00	1.49	2.49
time (sec)	N/A	0.495	10.357	1.063	0.000	0.097	0.000	0.000	1.371	0.089

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	C	F	C	F	F	F	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	546	46	317	0	70	0	0	228	207
N.S.	1	7.69	0.65	4.46	0.00	0.99	0.00	0.00	3.21	2.92
time (sec)	N/A	1.170	10.360	0.759	0.000	0.085	0.000	0.000	5.843	0.443

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	C	A	C	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	233	52	367	0	63	0	0	51	0
N.S.	1	5.07	1.13	7.98	0.00	1.37	0.00	0.00	1.11	0.00
time (sec)	N/A	0.833	10.851	0.920	0.000	0.088	0.000	0.000	180.023	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F(-2)	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	34	112	0	0	0	0	54	67
N.S.	1	1.00	1.06	3.50	0.00	0.00	0.00	0.00	1.69	2.09
time (sec)	N/A	0.234	1.209	3.578	0.000	0.000	0.000	0.000	0.164	0.976

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	B	F	B	F	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	31	44	0	44	0	0	24	204
N.S.	1	1.00	1.35	1.91	0.00	1.91	0.00	0.00	1.04	8.87
time (sec)	N/A	0.192	0.868	0.542	0.000	0.079	0.000	0.000	0.160	0.123

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	218	218	47	351	0	511	0	0	90	0
N.S.	1	1.00	0.22	1.61	0.00	2.34	0.00	0.00	0.41	0.00
time (sec)	N/A	0.237	10.044	70.818	0.000	0.205	0.000	0.000	0.227	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	210	210	50	350	0	657	0	0	90	0
N.S.	1	1.00	0.24	1.67	0.00	3.13	0.00	0.00	0.43	0.00
time (sec)	N/A	0.233	10.053	72.366	0.000	0.212	0.000	0.000	0.231	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	222	222	65	347	0	659	0	0	90	0
N.S.	1	1.00	0.29	1.56	0.00	2.97	0.00	0.00	0.41	0.00
time (sec)	N/A	0.218	10.053	61.048	0.000	0.215	0.000	0.000	0.289	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	214	214	68	350	0	516	0	0	90	0
N.S.	1	1.00	0.32	1.64	0.00	2.41	0.00	0.00	0.42	0.00
time (sec)	N/A	0.222	10.049	61.290	0.000	0.199	0.000	0.000	0.285	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	73	77	327	0	323	0	0	261	0
N.S.	1	1.12	1.18	5.03	0.00	4.97	0.00	0.00	4.02	0.00
time (sec)	N/A	0.267	8.545	2.500	0.000	0.161	0.000	0.000	0.267	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	70	77	311	0	112	0	0	261	0
N.S.	1	1.11	1.22	4.94	0.00	1.78	0.00	0.00	4.14	0.00
time (sec)	N/A	0.276	8.471	2.625	0.000	0.156	0.000	0.000	0.262	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	F(-2)	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	92	818	0	0	0	0	47	0
N.S.	1	1.00	1.74	15.43	0.00	0.00	0.00	0.00	0.89	0.00
time (sec)	N/A	0.206	0.747	2.490	0.000	0.000	0.000	0.000	0.175	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	111	0	1421	0	267	0	0	21	0
N.S.	1	1.03	0.00	13.16	0.00	2.47	0.00	0.00	0.19	0.00
time (sec)	N/A	0.278	0.000	5.063	0.000	0.771	0.000	0.000	0.152	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	189	111	0	1252	0	0	31	0
N.S.	1	1.00	1.93	1.13	0.00	12.78	0.00	0.00	0.32	0.00
time (sec)	N/A	0.174	0.883	1.144	0.000	101.790	0.000	0.000	0.168	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	154	154	0	0	0	0	0	0	79	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.51	0.00
time (sec)	N/A	0.408	0.000	0.000	0.000	0.000	0.000	0.000	0.204	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	A	A	B	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	100	182	92	110	387	0	113	35	157
N.S.	1	1.04	1.90	0.96	1.15	4.03	0.00	1.18	0.36	1.64
time (sec)	N/A	0.214	0.339	0.832	0.125	0.093	0.000	3.443	0.179	0.359

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	114	95	0	246	0	0	27	0
N.S.	1	1.00	1.30	1.08	0.00	2.80	0.00	0.00	0.31	0.00
time (sec)	N/A	0.168	0.235	3.100	0.000	1.544	0.000	0.000	0.150	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	F	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	233	249	283	0	0	336	0	0	29	0
N.S.	1	1.07	1.21	0.00	0.00	1.44	0.00	0.00	0.12	0.00
time (sec)	N/A	0.390	0.720	0.000	0.000	1.565	0.000	0.000	0.169	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	A	A	F	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	86	104	80	86	84	0	87	31	100
N.S.	1	1.05	1.27	0.98	1.05	1.02	0.00	1.06	0.38	1.22
time (sec)	N/A	0.199	0.092	3.595	0.108	0.082	0.000	0.138	0.149	0.349

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	135	383	145	720	0	287	0	0	81	0
N.S.	1	2.84	1.07	5.33	0.00	2.13	0.00	0.00	0.60	0.00
time (sec)	N/A	0.574	1.098	7.268	0.000	5.654	0.000	0.000	0.160	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	135	383	145	720	0	287	0	0	81	0
N.S.	1	2.84	1.07	5.33	0.00	2.13	0.00	0.00	0.60	0.00
time (sec)	N/A	0.597	1.057	6.322	0.000	5.607	0.000	0.000	0.155	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	B	A	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	119	357	139	687	0	265	0	0	69	0
N.S.	1	3.00	1.17	5.77	0.00	2.23	0.00	0.00	0.58	0.00
time (sec)	N/A	0.570	1.030	7.411	0.000	5.235	0.000	0.000	0.169	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	34	0	0	0	0	133	0
N.S.	1	1.00	1.00	0.79	0.00	0.00	0.00	0.00	3.09	0.00
time (sec)	N/A	0.201	10.175	0.406	0.000	0.000	0.000	0.000	0.168	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	34	0	0	0	0	81	0
N.S.	1	1.00	1.00	0.79	0.00	0.00	0.00	0.00	1.88	0.00
time (sec)	N/A	0.225	10.138	0.358	0.000	0.000	0.000	0.000	0.157	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	43	34	0	0	0	0	81	0
N.S.	1	1.00	1.10	0.87	0.00	0.00	0.00	0.00	2.08	0.00
time (sec)	N/A	0.178	10.017	0.214	0.000	0.000	0.000	0.000	0.155	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	C	B	A	C	F	F	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	70	101	12	105	94	31	0	34	10
N.S.	1	1.04	1.51	0.18	1.57	1.40	0.46	0.00	0.51	0.15
time (sec)	N/A	0.161	0.085	1.471	0.136	0.084	0.531	0.000	0.156	0.222

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	A	A	C	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	76	90	66	73	75	41	74	35	91
N.S.	1	1.09	1.29	0.94	1.04	1.07	0.59	1.06	0.50	1.30
time (sec)	N/A	0.184	0.049	2.071	0.110	0.077	0.521	0.121	0.164	0.240

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F(-1)	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	384	375	0	0	0	0	0	0	88	0
N.S.	1	0.98	0.00	0.00	0.00	0.00	0.00	0.00	0.23	0.00
time (sec)	N/A	0.737	0.000	0.000	0.000	0.000	0.000	0.000	0.171	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	234	234	0	0	0	0	0	0	32	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.14	0.00
time (sec)	N/A	0.406	0.000	0.000	0.000	0.000	0.000	0.000	0.177	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	C	F	B	F	F	F	F(-1)
verified	N/A	Yes	N/A	No	TBD	TBD	TBD	TBD	TBD	TBD
size	199	250	0	842	0	2155	0	0	648	0
N.S.	1	1.26	0.00	4.23	0.00	10.83	0.00	0.00	3.26	0.00
time (sec)	N/A	0.474	0.000	24.925	0.000	1.684	0.000	0.000	0.203	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	177	182	0	0	0	0	0	0	69	0
N.S.	1	1.03	0.00	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.382	0.000	0.000	0.000	0.000	0.000	0.000	0.162	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	F	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	177	182	0	0	0	0	0	0	69	0
N.S.	1	1.03	0.00	0.00	0.00	0.00	0.00	0.00	0.39	0.00
time (sec)	N/A	0.407	0.000	0.000	0.000	0.000	0.000	0.000	0.156	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	A	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	138	204	179	0	191	0	0	19	0
N.S.	1	1.05	1.55	1.36	0.00	1.45	0.00	0.00	0.14	0.00
time (sec)	N/A	0.205	0.330	1.069	0.000	0.096	0.000	0.000	0.153	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	250	272	26	0	0	0	0	0	20	0
N.S.	1	1.09	0.10	0.00	0.00	0.00	0.00	0.00	0.08	0.00
time (sec)	N/A	0.425	10.014	0.000	0.000	0.000	0.000	0.000	0.154	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	C	F	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	383	655	138	0	0	0	0	0	42	0
N.S.	1	1.71	0.36	0.00	0.00	0.00	0.00	0.00	0.11	0.00
time (sec)	N/A	1.143	15.090	0.000	0.000	0.000	0.000	0.000	0.161	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Reduce	Mupad
grade	N/A	A	A	C	F	A	F	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD	TBD
size	272	295	283	1151	0	341	0	0	19	0
N.S.	1	1.08	1.04	4.23	0.00	1.25	0.00	0.00	0.07	0.00
time (sec)	N/A	0.352	1.878	6.398	0.000	1.612	0.000	0.000	0.144	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [47] had the largest ratio of [1.30000000000000004]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	10	0.100
2	C	5	4	3.27	37	0.108
3	A	2	2	1.00	15	0.133
4	A	5	4	1.00	19	0.211
5	A	2	2	1.00	17	0.118
6	A	8	7	1.00	17	0.412
7	A	4	3	1.00	17	0.176
8	A	2	2	1.00	23	0.087
9	A	2	2	1.66	27	0.074
10	B	2	2	2.46	39	0.051
11	A	1	1	1.00	45	0.022
12	A	5	4	1.21	32	0.125
13	A	4	3	1.00	32	0.094
14	A	3	2	1.00	27	0.074
15	A	3	2	1.00	29	0.069
16	A	2	2	1.00	30	0.067
17	A	4	3	0.98	13	0.231
18	A	4	3	0.98	15	0.200
19	A	3	2	1.00	23	0.087
20	A	3	2	1.00	25	0.080
21	A	4	3	1.07	11	0.273

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
22	A	3	2	1.00	18	0.111
23	A	7	6	1.00	20	0.300
24	A	8	7	1.09	31	0.226
25	A	3	2	1.00	29	0.069
26	A	3	2	1.00	35	0.057
27	A	7	6	1.10	32	0.188
28	A	8	7	1.12	21	0.333
29	A	5	5	1.22	14	0.357
30	A	3	2	1.00	23	0.087
31	A	4	3	0.98	13	0.231
32	A	2	2	1.00	33	0.061
33	A	6	5	1.05	15	0.333
34	A	6	5	1.07	15	0.333
35	A	1	1	1.00	11	0.091
36	A	6	5	1.11	15	0.333
37	A	1	1	1.00	17	0.059
38	A	3	3	1.00	18	0.167
39	A	3	3	1.09	16	0.188
40	A	6	5	1.06	17	0.294
41	A	6	5	1.30	13	0.385
42	A	6	5	1.19	16	0.312
43	F	0	0	N/A	0.000	N/A
44	F	0	0	N/A	0.000	N/A
45	F	0	0	N/A	0.000	N/A
46	A	2	2	1.16	32	0.062
47	A	27	26	1.24	20	1.300
48	F	0	0	N/A	0.000	N/A
49	F	0	0	N/A	0.000	N/A
50	A	24	23	1.12	23	1.000
51	F	0	0	N/A	0.000	N/A
52	C	11	10	6.50	48	0.208
53	A	8	7	1.03	24	0.292

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
54	A	8	7	1.00	24	0.292
55	A	1	1	1.00	18	0.056
56	A	2	2	1.04	13	0.154
57	A	7	6	1.12	15	0.400
58	A	2	2	1.00	17	0.118
59	A	2	2	1.46	22	0.091
60	A	2	2	1.00	17	0.118
61	A	2	2	1.00	21	0.095
62	A	1	1	1.00	38	0.026
63	A	1	1	1.00	33	0.030
64	A	3	3	1.09	38	0.079
65	A	1	1	1.00	19	0.053
66	A	5	4	1.00	19	0.211
67	A	5	4	1.35	24	0.167
68	A	1	1	1.00	24	0.042
69	A	8	7	1.10	24	0.292
70	A	8	7	1.09	24	0.292
71	A	3	3	1.01	26	0.115
72	A	3	3	1.01	22	0.136
73	A	1	1	1.00	22	0.045
74	A	1	1	1.00	23	0.043
75	A	10	9	1.00	18	0.500
76	A	9	8	1.00	23	0.348
77	A	1	1	1.00	21	0.048
78	A	1	1	1.00	19	0.053
79	A	1	1	1.00	19	0.053
80	A	1	1	1.00	19	0.053
81	A	6	5	1.00	34	0.147
82	C	6	5	7.69	40	0.125
83	C	8	7	5.07	51	0.137
84	A	3	2	1.00	29	0.069
85	A	3	2	1.00	18	0.111

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
86	A	1	1	1.00	25	0.040
87	A	1	1	1.00	25	0.040
88	A	1	1	1.00	25	0.040
89	A	1	1	1.00	25	0.040
90	A	3	2	1.12	40	0.050
91	A	3	2	1.11	40	0.050
92	A	1	1	1.00	18	0.056
93	A	3	3	1.03	15	0.200
94	A	1	1	1.00	21	0.048
95	A	2	2	1.00	23	0.087
96	A	6	5	1.04	24	0.208
97	A	1	1	1.00	19	0.053
98	A	12	11	1.07	20	0.550
99	A	6	5	1.05	22	0.227
100	B	2	2	2.84	25	0.080
101	B	3	3	2.84	24	0.125
102	B	2	2	3.00	23	0.087
103	A	2	2	1.00	20	0.100
104	A	2	2	1.00	25	0.080
105	A	3	3	1.00	19	0.158
106	A	2	2	1.04	11	0.182
107	A	7	6	1.09	15	0.400
108	A	6	6	0.98	19	0.316
109	A	2	2	1.00	22	0.091
110	A	2	2	1.26	27	0.074
111	A	5	5	1.03	17	0.294
112	A	6	6	1.03	27	0.222
113	A	3	3	1.05	19	0.158
114	A	14	13	1.09	20	0.650
115	A	2	2	1.71	24	0.083
116	A	11	10	1.08	19	0.526

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{1}{\sqrt{1-ax}} dx$	71
3.2	$\int \frac{-2\log(-\sqrt{-1+ax})+\log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$	76
3.3	$\int \frac{1}{(2x+\sqrt{1+x^2})^2} dx$	82
3.4	$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx$	89
3.5	$\int \frac{1}{(2\sqrt{x}+\sqrt{1+x})^2} dx$	96
3.6	$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx$	102
3.7	$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx$	109
3.8	$\int \frac{1}{(\sqrt{-1+x}+\sqrt{x})^2\sqrt{-1+x}} dx$	115
3.9	$\int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx$	120
3.10	$\int \frac{(\sqrt{x}-\sqrt{-1+x^2})^2}{(1+x-x^2)^2\sqrt{-1+x^2}} dx$	129
3.11	$\int \left(\frac{1}{\sqrt{2(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2(1+x)^2\sqrt{i+x^2}}} \right) dx$	138
3.12	$\int \frac{\sqrt{x^2+\sqrt{1+x^4}}}{(1+x)^2\sqrt{1+x^4}} dx$	144
3.13	$\int \frac{\sqrt{x^2+\sqrt{1+x^4}}}{(1+x)\sqrt{1+x^4}} dx$	151
3.14	$\int \frac{\sqrt{x^2+\sqrt{1+x^4}}}{\sqrt{1+x^4}} dx$	157
3.15	$\int \frac{\sqrt{-x^2+\sqrt{1+x^4}}}{\sqrt{1+x^4}} dx$	162
3.16	$\int \frac{(-1+x)^{3/2}+(1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx$	167
3.17	$\int (x + \sqrt{a+x^2})^b dx$	172
3.18	$\int (x - \sqrt{a+x^2})^b dx$	178
3.19	$\int \frac{(x+\sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx$	183
3.20	$\int \frac{(x-\sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx$	188

3.21	$\int \frac{1}{(a+be^{px})^2} dx$	193
3.22	$\int \frac{1}{(be^{-px}+ae^{px})^2} dx$	198
3.23	$\int \frac{x}{(be^{-px}+ae^{px})^2} dx$	203
3.24	$\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx$	209
3.25	$\int \frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a^2+x^2}} dx$	218
3.26	$\int \frac{\sqrt{bx+\sqrt{a+b^2x^2}}}{\sqrt{a+b^2x^2}} dx$	223
3.27	$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx$	228
3.28	$\int \frac{\sqrt{x+\sqrt{a^2+x^2}}}{x} dx$	235
3.29	$\int x^3 \log^3(2+x) \log(3+x) dx$	242
3.30	$\int \frac{(x+\sqrt{b+x^2})^a}{\sqrt{b+x^2}} dx$	252
3.31	$\int (x+\sqrt{b+x^2})^a dx$	257
3.32	$\int (6+3x^a+2x^{2a})^{\frac{1}{a}} (x^a+x^{2a}+x^{3a}) dx$	263
3.33	$\int \frac{1}{x\sqrt[3]{1-x^2}} dx$	268
3.34	$\int \frac{1}{x(1-x^2)^{2/3}} dx$	275
3.35	$\int \frac{1}{\sqrt[3]{1-x^3}} dx$	282
3.36	$\int \frac{1}{x\sqrt[3]{1-x^3}} dx$	287
3.37	$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx$	294
3.38	$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx$	300
3.39	$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx$	306
3.40	$\int \frac{1}{\sqrt[3]{-5+7x-3x^2+x^3}} dx$	312
3.41	$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx$	319
3.42	$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx$	326
3.43	$\int \frac{1}{x\sqrt[3]{(-1+x)(q-2qx+x^2)}} dx$	333
3.44	$\int \frac{2-(1+k)x}{\sqrt[3]{(1-x)x(1-kx)(1-(1+k)x)}} dx$	339
3.45	$\int \frac{1-kx}{(1+(-2+k)x)((1-x)x(1-kx))^{2/3}} dx$	345
3.46	$\int \frac{a+bx+cx^2}{(1-x+x^2)\sqrt[3]{1-x^3}} dx$	353
3.47	$\int \frac{1}{(3-2x)^{11/2}(1+x+2x^2)^5} dx$	361
3.48	$\int \frac{1}{(3-2x)^{21/2}(1+x+2x^2)^{10}} dx$	376
3.49	$\int \frac{1}{(3-2x)^{41/2}(1+x+2x^2)^{20}} dx$	392
3.50	$\int \frac{1}{(3-2x+x^2)^{11/2}(1+x+2x^2)^5} dx$	407
3.51	$\int \frac{1}{(3-2x+x^2)^{21/2}(1+x+2x^2)^{10}} dx$	422

3.52	$\int \frac{-a-\sqrt{1+a^2+x}}{(-a+\sqrt{1+a^2+x})\sqrt{(-a+x)(1+x^2)}} dx$	436
3.53	$\int \frac{a+bx}{\sqrt[3]{1-x^2(3+x^2)}} dx$	446
3.54	$\int \frac{a+bx}{(3-x^2)\sqrt[3]{1+x^2}} dx$	454
3.55	$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx$	462
3.56	$\int x\sqrt[3]{1-x^3} dx$	468
3.57	$\int \frac{\sqrt[3]{1-x^3}}{x} dx$	474
3.58	$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx$	481
3.59	$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx$	487
3.60	$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx$	494
3.61	$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx$	499
3.62	$\int \frac{3-3x+30x^2+160x^3}{9+24x-12x^2+80x^3+320x^4} dx$	504
3.63	$\int \frac{3+12x+20x^2}{9+24x-12x^2+80x^3+320x^4} dx$	509
3.64	$\int \frac{-84-576x-400x^2+2560x^3}{9+24x-12x^2+80x^3+320x^4} dx$	515
3.65	$\int \frac{\sqrt{1-x^4}}{1+x^4} dx$	522
3.66	$\int \frac{\sqrt{1+x^4}}{1-x^4} dx$	527
3.67	$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx$	533
3.68	$\int \frac{\sqrt{1+px^2-x^4}}{1+x^4} dx$	539
3.69	$\int \frac{a+bx}{(2-x^2)\sqrt[4]{-1+x^2}} dx$	545
3.70	$\int \frac{a+bx}{\sqrt[4]{-1-x^2(2+x^2)}} dx$	552
3.71	$\int \frac{a+bx}{\sqrt[4]{1-x^2(2-x^2)}} dx$	559
3.72	$\int \frac{a+bx}{\sqrt[4]{1+x^2(2+x^2)}} dx$	565
3.73	$\int \frac{x}{\sqrt{1-x^3(4-x^3)}} dx$	571
3.74	$\int \frac{x}{(4-dx^3)\sqrt{-1+dx^3}} dx$	578
3.75	$\int \frac{x}{\sqrt{-1+x^3(8+x^3)}} dx$	585
3.76	$\int \frac{x}{(8-dx^3)\sqrt{1+dx^3}} dx$	593
3.77	$\int \frac{1}{\sqrt[3]{1-3x^2(3-x^2)}} dx$	602
3.78	$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx$	608
3.79	$\int \frac{1}{\sqrt[3]{1-x^2(3+x^2)}} dx$	614
3.80	$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx$	621
3.81	$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx$	627
3.82	$\int \frac{-2+a+x}{(-a+x)\sqrt{(2-a)ax+(-1-2a+a^2)x^2+x^3}} dx$	634

3.83	$\int \frac{-a+(-1+2a)x}{(-a+x)\sqrt{a^2x-(-1+2a+a^2)x^2+(-1+2a)x^3}} dx$	642
3.84	$\int \frac{1-\sqrt[3]{2x}}{(2^{2/3}+x)\sqrt{1+x^3}} dx$	650
3.85	$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx$	656
3.86	$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3+x^3})} dx$	661
3.87	$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3+x^3})} dx$	668
3.88	$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3+x^3})} dx$	675
3.89	$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3+x^3})} dx$	682
3.90	$\int \frac{1-\sqrt{3+x}}{(1+\sqrt{3+x})\sqrt{-4+4\sqrt{3}x^2+x^4}} dx$	689
3.91	$\int \frac{1+\sqrt{3+x}}{(1-\sqrt{3+x})\sqrt{-4-4\sqrt{3}x^2+x^4}} dx$	695
3.92	$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx$	702
3.93	$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx$	707
3.94	$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx$	714
3.95	$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx$	720
3.96	$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx$	725
3.97	$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx$	733
3.98	$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx$	739
3.99	$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx$	749
3.100	$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx$	756
3.101	$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx$	763
3.102	$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx$	770
3.103	$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx$	777
3.104	$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx$	782
3.105	$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx$	787
3.106	$\int (1-x^3)^{2/3} dx$	792
3.107	$\int \frac{(1-x^3)^{2/3}}{x} dx$	798
3.108	$\int \frac{(1-x^3)^{2/3}}{a+bx} dx$	805
3.109	$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$	812
3.110	$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$	817

3.111	$\int \frac{(1-x^3)^{2/3}}{1+x} dx$	824
3.112	$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx$	830
3.113	$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx$	837
3.114	$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx$	844
3.115	$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx$	854
3.116	$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx$	860

3.1 $\int \frac{1}{\sqrt{1-ax}} dx$

Optimal result	71
Mathematica [A] (verified)	71
Rubi [A] (verified)	72
Maple [A] (verified)	73
Fricas [A] (verification not implemented)	73
Sympy [A] (verification not implemented)	74
Maxima [A] (verification not implemented)	74
Giac [A] (verification not implemented)	74
Mupad [B] (verification not implemented)	75
Reduce [B] (verification not implemented)	75

Optimal result

Integrand size = 10, antiderivative size = 15

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{1-ax}}{a}$$

output

```
-2*(-a*x+1)^(1/2)/a
```

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{1-ax}}{a}$$

input

```
Integrate[1/Sqrt[1 - a*x],x]
```

output

```
(-2*Sqrt[1 - a*x])/a
```


Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {17}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{1-ax}} dx$$

$$\downarrow 17$$

$$-\frac{2\sqrt{1-ax}}{a}$$

input `Int[1/Sqrt[1 - a*x],x]`

output `(-2*Sqrt[1 - a*x])/a`

Defintions of rubi rules used

rule 17 `Int[(c_.)*((a_.) + (b_.)*(x_))^(m_.), x_Symbol] := Simp[c*((a + b*x)^(m + 1))/(b*(m + 1)), x] /; FreeQ[{a, b, c, m}, x] && NeQ[m, -1]`

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
gospers	$-\frac{2\sqrt{-ax+1}}{a}$	14
derivativdivides	$-\frac{2\sqrt{-ax+1}}{a}$	14
default	$-\frac{2\sqrt{-ax+1}}{a}$	14
trager	$-\frac{2\sqrt{-ax+1}}{a}$	14
pseudoelliptic	$-\frac{2\sqrt{-ax+1}}{a}$	14
risch	$\frac{2ax-2}{a\sqrt{-ax+1}}$	19
orering	$\frac{2ax-2}{a\sqrt{-ax+1}}$	19
meijerg	$-\frac{2\sqrt{\pi}+2\sqrt{\pi}\sqrt{-ax+1}}{\sqrt{\pi}a}$	28

input `int(1/(-a*x+1)^(1/2),x,method=_RETURNVERBOSE)`output `-2*(-a*x+1)^(1/2)/a`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{-ax+1}}{a}$$

input `integrate(1/(-a*x+1)^(1/2),x, algorithm="fricas")`output `-2*sqrt(-a*x + 1)/a`

Sympy [A] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{-ax+1}}{a}$$

input `integrate(1/(-a*x+1)**(1/2),x)`output `-2*sqrt(-a*x + 1)/a`**Maxima [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{-ax+1}}{a}$$

input `integrate(1/(-a*x+1)^(1/2),x, algorithm="maxima")`output `-2*sqrt(-a*x + 1)/a`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{-ax+1}}{a}$$

input `integrate(1/(-a*x+1)^(1/2),x, algorithm="giac")`output `-2*sqrt(-a*x + 1)/a`

Mupad [B] (verification not implemented)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{1-ax}}{a}$$

input `int(1/(1 - a*x)^(1/2),x)`

output `-(2*(1 - a*x)^(1/2))/a`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{1-ax}} dx = -\frac{2\sqrt{-ax+1}}{a}$$

input `int(1/(-a*x+1)^(1/2),x)`

output `(- 2*sqrt(- a*x + 1))/a`

3.2
$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$$

Optimal result	76
Mathematica [C] (verified)	76
Rubi [C] (verified)	77
Maple [C] (verified)	78
Fricas [F]	79
Sympy [A] (verification not implemented)	79
Maxima [B] (verification not implemented)	79
Giac [B] (verification not implemented)	80
Mupad [B] (verification not implemented)	80
Reduce [B] (verification not implemented)	81

Optimal result

Integrand size = 37, antiderivative size = 15

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx = -\frac{2\sqrt{1-ax}}{a}$$

output `-2*(-a*x+1)^(1/2)/a`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 2 in optimal.

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 2.47

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx = \frac{\sqrt{-1+ax}(-2 \log(-\sqrt{-1+ax}) + \log(-1+ax))}{a\pi}$$

input `Integrate[(-2*Log[-Sqrt[-1 + a*x]] + Log[-1 + a*x])/(2*Pi*Sqrt[-1 + a*x]), x]`

output $(\text{Sqrt}[-1 + a*x]*(-2*\text{Log}[-\text{Sqrt}[-1 + a*x]] + \text{Log}[-1 + a*x]))/(a*\text{Pi})$

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 2 in optimal.

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 3.27, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.108$, Rules used = {27, 25, 7267, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\log(ax-1) - 2\log(-\sqrt{ax-1})}{2\pi\sqrt{ax-1}} dx \\
 & \quad \downarrow 27 \\
 & \int \frac{-2\log(-\sqrt{ax-1}) - \log(ax-1)}{2\pi\sqrt{ax-1}} dx \\
 & \quad \downarrow 25 \\
 & \int \frac{2\log(-\sqrt{ax-1}) - \log(ax-1)}{2\pi\sqrt{ax-1}} dx \\
 & \quad \downarrow 7267 \\
 & \int \frac{(2\log(-\sqrt{ax-1}) - \log(ax-1)) d\sqrt{ax-1}}{\pi a} \\
 & \quad \downarrow 2009 \\
 & \frac{2\sqrt{ax-1}\log(-\sqrt{ax-1}) - \sqrt{ax-1}\log(ax-1)}{\pi a}
 \end{aligned}$$

input $\text{Int}[(-2*\text{Log}[-\text{Sqrt}[-1 + a*x]] + \text{Log}[-1 + a*x])/(2*\text{Pi}*\text{Sqrt}[-1 + a*x]),x]$

output $-((2*\text{Sqrt}[-1 + a*x]*\text{Log}[-\text{Sqrt}[-1 + a*x]] - \text{Sqrt}[-1 + a*x]*\text{Log}[-1 + a*x])/(a*\text{Pi}))$

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7267 `Int[u_, x_Symbol] := With[{lst = SubstForFractionalPowerOfLinear[u, x]}, Simp[lst[[2]]*lst[[4]] Subst[Int[lst[[1]], x], x, lst[[3]]^(1/lst[[2])]], x] /; !FalseQ[lst] && SubstForFractionalPowerQ[u, lst[[3]], x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 3 vs. order 2.

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.27

method	result	size
gospers	$\frac{\sqrt{ax-1} (\ln(ax-1) - 2\ln(-\sqrt{ax-1}))}{a\pi}$	34
orering	$\frac{\sqrt{ax-1} (\ln(ax-1) - 2\ln(-\sqrt{ax-1}))}{a\pi}$	34
derivativedivides	$\frac{-2\ln(-\sqrt{ax-1})\sqrt{ax-1} + \sqrt{ax-1} \ln(ax-1)}{\pi a}$	42
default	$\frac{-2\ln(-\sqrt{ax-1})\sqrt{ax-1} + \sqrt{ax-1} \ln(ax-1)}{\pi a}$	42
meijerg	$\frac{i\sqrt{-\text{signum}(ax-1)} (-2\sqrt{\pi} + 2\sqrt{\pi}\sqrt{-ax+1})}{\sqrt{\pi}\sqrt{\text{signum}(ax-1)} a}$	47
parts	$\frac{-2\ln(-\sqrt{ax-1})\sqrt{ax-1} + 2\sqrt{ax-1}}{a\pi} + \frac{\sqrt{ax-1} \ln(ax-1) - 2\sqrt{ax-1}}{\pi a}$	68

input `int(1/2*(ln(a*x-1)-2*ln(-(a*x-1)^(1/2)))/Pi/(a*x-1)^(1/2),x,method=_RETURN
VERBOSE)`

output `(a*x-1)^(1/2)*(ln(a*x-1)-2*ln(-(a*x-1)^(1/2)))/a/Pi`

Fricas [F]

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx = \int \frac{\log(ax-1) - 2 \log(-\sqrt{ax-1})}{2\pi\sqrt{ax-1}} dx$$

input `integrate(1/2*(log(a*x-1)-2*log(-(a*x-1)^(1/2)))/pi/(a*x-1)^(1/2),x, algorithm="fricas")`

output 0

Sympy [A] (verification not implemented)

Time = 3.96 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.80

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$$

$$= \frac{\begin{cases} \frac{-2\sqrt{ax-1} \log(-\sqrt{ax-1}) + \sqrt{ax-1} \log(ax-1)}{a} & \text{for } a \neq 0 \\ \pi x & \text{otherwise} \end{cases}}{\pi}$$

input `integrate(1/2*(ln(a*x-1)-2*ln(-(a*x-1)**(1/2)))/pi/(a*x-1)**(1/2),x)`

output `Piecewise((((-2*sqrt(a*x - 1)*log(-sqrt(a*x - 1)) + sqrt(a*x - 1)*log(a*x - 1))/a, Ne(a, 0)), (pi*x, True))/pi`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(13) = 26.

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.73

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$$

$$= \frac{\sqrt{ax-1} \log(ax-1) - 2 \sqrt{ax-1} \log(-\sqrt{ax-1})}{\pi a}$$

input `integrate(1/2*(log(a*x-1)-2*log(-(a*x-1)^(1/2)))/pi/(a*x-1)^(1/2),x, algo
ithm="maxima")`

output `(sqrt(a*x - 1)*log(a*x - 1) - 2*sqrt(a*x - 1)*log(-sqrt(a*x - 1)))/(pi*a)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(13) = 26.

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.73

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$$

$$= \frac{\sqrt{ax-1} \log(ax-1) - 2\sqrt{ax-1} \log(-\sqrt{ax-1})}{\pi a}$$

input `integrate(1/2*(log(a*x-1)-2*log(-(a*x-1)^(1/2)))/pi/(a*x-1)^(1/2),x, algo
ithm="giac")`

output `(sqrt(a*x - 1)*log(a*x - 1) - 2*sqrt(a*x - 1)*log(-sqrt(a*x - 1)))/(pi*a)`

Mupad [B] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 43, normalized size of antiderivative = 2.87

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$$

$$= -\frac{2 \ln(-\sqrt{ax-1}) \sqrt{ax-1} - \ln(ax-1) \sqrt{ax-1}}{\Pi a}$$

input `int((log(a*x - 1)/2 - log(-(a*x - 1)^(1/2)))/(Pi*(a*x - 1)^(1/2)),x)`

output `-(2*log(-(a*x - 1)^(1/2))*(a*x - 1)^(1/2) - log(a*x - 1)*(a*x - 1)^(1/2))/
(Pi*a)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.07

$$\int \frac{-2 \log(-\sqrt{-1+ax}) + \log(-1+ax)}{2\pi\sqrt{-1+ax}} dx$$

$$= \frac{\sqrt{ax-1}(-2 \log(-\sqrt{ax-1}) + \log(ax-1))}{a\pi}$$

input `int(1/2*(log(a*x-1)-2*log(-sqrt(a*x-1)))/Pi/sqrt(a*x-1),x)`output `(sqrt(a*x - 1)*(- 2*log(- sqrt(a*x - 1)) + log(a*x - 1)))/(a*pi)`

3.3 $\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx$

Optimal result	82
Mathematica [A] (verified)	82
Rubi [A] (verified)	83
Maple [C] (verified)	84
Fricas [A] (verification not implemented)	85
Sympy [F]	85
Maxima [F]	86
Giac [B] (verification not implemented)	86
Mupad [B] (verification not implemented)	87
Reduce [B] (verification not implemented)	87

Optimal result

Integrand size = 15, antiderivative size = 82

$$\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx = \frac{4x}{3(1-3x^2)} - \frac{2\sqrt{1+x^2}}{3(1-3x^2)} - \frac{\operatorname{arctanh}(\sqrt{3}x)}{3\sqrt{3}} + \frac{\operatorname{arctanh}(\frac{1}{2}\sqrt{3}\sqrt{1+x^2})}{3\sqrt{3}}$$

output $4/3*x/(-3*x^2+1)-1/9*\operatorname{arctanh}(x*3^{(1/2)})*3^{(1/2)}+1/9*\operatorname{arctanh}(1/2*3^{(1/2)}*(x^2+1)^{(1/2)})*3^{(1/2)}-2/3*(x^2+1)^{(1/2)/(-3*x^2+1)}$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.73

$$\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx = \frac{6(-2x + \sqrt{1+x^2}) - 2\sqrt{3}(-1 + 3x^2) \operatorname{arctanh}\left(\frac{x-\sqrt{1+x^2}}{\sqrt{3}}\right)}{-9 + 27x^2}$$

input `Integrate[(2*x + Sqrt[1 + x^2])^(-2), x]`

output

```
(6*(-2*x + Sqrt[1 + x^2]) - 2*Sqrt[3]*(-1 + 3*x^2)*ArcTanh[(x - Sqrt[1 + x^2])/Sqrt[3]])/(-9 + 27*x^2)
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{x^2+1}+2x)^2} dx$$

↓ 7293

$$\int \left(-\frac{4\sqrt{x^2+1}x}{(3x^2-1)^2} + \frac{5}{3(3x^2-1)} + \frac{8}{3(3x^2-1)^2} \right) dx$$

↓ 2009

$$\frac{\operatorname{arctanh}\left(\frac{1}{2}\sqrt{3}\sqrt{x^2+1}\right)}{3\sqrt{3}} - \frac{\operatorname{arctanh}(\sqrt{3}x)}{3\sqrt{3}} + \frac{4x}{3(1-3x^2)} - \frac{2\sqrt{x^2+1}}{3(1-3x^2)}$$

input

```
Int[(2*x + Sqrt[1 + x^2])^(-2), x]
```

output

```
(4*x)/(3*(1 - 3*x^2)) - (2*Sqrt[1 + x^2])/(3*(1 - 3*x^2)) - ArcTanh[Sqrt[3]*x]/(3*Sqrt[3]) + ArcTanh[(Sqrt[3]*Sqrt[1 + x^2])/2]/(3*Sqrt[3])
```

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

```
rule 7293 Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.22 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.89

method	result
trager	$-\frac{4x}{3(3x^2-1)} + \frac{2\sqrt{x^2+1}}{3(3x^2-1)} + \frac{\text{RootOf}(-Z^2-3) \ln\left(-\frac{2\text{RootOf}(-Z^2-3)+3\sqrt{x^2+1}}{\text{RootOf}(-Z^2-3)x+1}\right)}{9}$
default	$-\frac{x}{2(3x^2-1)} - \frac{\text{arctanh}(x\sqrt{3})\sqrt{3}}{9} - \frac{5x}{18(x^2-\frac{1}{3})} - \sqrt{3} \left(-\frac{\left(x-\frac{\sqrt{3}}{3}\right)^2 + \frac{2\sqrt{3}\left(x-\frac{\sqrt{3}}{3}\right)}{3} + \frac{4}{3}}{12\left(x-\frac{\sqrt{3}}{3}\right)} + \sqrt{3} \frac{\sqrt{9\left(x-\frac{\sqrt{3}}{3}\right)^2 + 6\sqrt{3}}}{3} \right)^{\frac{3}{2}}$

```
input int(1/(2*x+(x^2+1)^(1/2))^2,x,method=_RETURNVERBOSE)
```

output

```
-4/3*x/(3*x^2-1)+2/3/(3*x^2-1)*(x^2+1)^(1/2)+1/9*RootOf(_Z^2-3)*ln(-(2*RootOf(_Z^2-3)+3*(x^2+1)^(1/2))/(RootOf(_Z^2-3)*x+1))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.22

$$\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx$$

$$= \frac{\sqrt{3}(3x^2 - 1) \log\left(\frac{3x^2 - 2\sqrt{3}x + 1}{3x^2 - 1}\right) + \sqrt{3}(3x^2 - 1) \log\left(\frac{3x^2 + 4\sqrt{3}\sqrt{x^2 + 1} + 7}{3x^2 - 1}\right) - 24x + 12\sqrt{x^2 + 1}}{18(3x^2 - 1)}$$

input

```
integrate(1/(2*x+(x^2+1)^(1/2))^2,x, algorithm="fricas")
```

output

```
1/18*(sqrt(3)*(3*x^2 - 1)*log((3*x^2 - 2*sqrt(3)*x + 1)/(3*x^2 - 1)) + sqrt(3)*(3*x^2 - 1)*log((3*x^2 + 4*sqrt(3)*sqrt(x^2 + 1) + 7)/(3*x^2 - 1)) - 24*x + 12*sqrt(x^2 + 1))/(3*x^2 - 1)
```

Sympy [F]

$$\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx = \int \frac{1}{(2x + \sqrt{x^2+1})^2} dx$$

input

```
integrate(1/(2*x+(x**2+1)**(1/2))**2,x)
```

output

```
Integral((2*x + sqrt(x**2 + 1))**(-2), x)
```

Maxima [F]

$$\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx = \int \frac{1}{(2x + \sqrt{x^2+1})^2} dx$$

input `integrate(1/(2*x+(x^2+1)^(1/2))^2,x, algorithm="maxima")`

output `integrate((2*x + sqrt(x^2 + 1))^(-2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 177 vs. 2(60) = 120.

Time = 0.13 (sec) , antiderivative size = 177, normalized size of antiderivative = 2.16

$$\begin{aligned} \int \frac{1}{(2x + \sqrt{1+x^2})^2} dx &= \frac{1}{18} \sqrt{3} \log \left(\frac{|6x - 2\sqrt{3}|}{|6x + 2\sqrt{3}|} \right) \\ &\quad - \frac{1}{18} \sqrt{3} \log \left(-\frac{|-6x - 8\sqrt{3} + 6\sqrt{x^2+1} - \frac{6}{x-\sqrt{x^2+1}}|}{2 \left(3x - 4\sqrt{3} - 3\sqrt{x^2+1} + \frac{3}{x-\sqrt{x^2+1}} \right)} \right) \\ &\quad - \frac{4 \left(x - \sqrt{x^2+1} + \frac{1}{x-\sqrt{x^2+1}} \right)}{3 \left(3 \left(x - \sqrt{x^2+1} + \frac{1}{x-\sqrt{x^2+1}} \right)^2 - 16 \right)} - \frac{4x}{3(3x^2 - 1)} \end{aligned}$$

input `integrate(1/(2*x+(x^2+1)^(1/2))^2,x, algorithm="giac")`

output `1/18*sqrt(3)*log(abs(6*x - 2*sqrt(3))/abs(6*x + 2*sqrt(3))) - 1/18*sqrt(3)*log(-1/2*abs(-6*x - 8*sqrt(3) + 6*sqrt(x^2 + 1) - 6/(x - sqrt(x^2 + 1)))/(3*x - 4*sqrt(3) - 3*sqrt(x^2 + 1) + 3/(x - sqrt(x^2 + 1)))) - 4/3*(x - sqrt(x^2 + 1) + 1/(x - sqrt(x^2 + 1)))/(3*(x - sqrt(x^2 + 1) + 1/(x - sqrt(x^2 + 1)))^2 - 16) - 4/3*x/(3*x^2 - 1)`

Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.49

$$\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx = \frac{\sqrt{3} \left(\ln \left(x - \frac{\sqrt{3}}{3} \right) - \ln \left(x + \sqrt{3} + 2\sqrt{x^2+1} \right) \right)}{18} - \frac{4x}{9 \left(x^2 - \frac{1}{3} \right)}$$

$$+ \frac{\sqrt{3} \left(\ln \left(x + \frac{\sqrt{3}}{3} \right) - \ln \left(x - \sqrt{3} - 2\sqrt{x^2+1} \right) \right)}{18}$$

$$- \frac{\sqrt{3} \left(6 \ln \left(x - \frac{\sqrt{3}}{3} \right) - 6 \ln \left(x + \sqrt{3} + 2\sqrt{x^2+1} \right) \right)}{54}$$

$$- \frac{\sqrt{3} \left(6 \ln \left(x + \frac{\sqrt{3}}{3} \right) - 6 \ln \left(x - \sqrt{3} - 2\sqrt{x^2+1} \right) \right)}{54}$$

$$+ \frac{\sqrt{3} \sqrt{x^2+1}}{9 \left(x - \frac{\sqrt{3}}{3} \right)} - \frac{\sqrt{3} \sqrt{x^2+1}}{9 \left(x + \frac{\sqrt{3}}{3} \right)} + \frac{\sqrt{3} \operatorname{atan}(\sqrt{3} x \operatorname{li}) \operatorname{li}}{9}$$

input

```
int(1/(2*x + (x^2 + 1)^(1/2))^2,x)
```

output

```
(3^(1/2)*(log(x - 3^(1/2)/3) - log(x + 3^(1/2) + 2*(x^2 + 1)^(1/2)))/18 +
(3^(1/2)*atan(3^(1/2)*x*1i)*1i)/9 - (4*x)/(9*(x^2 - 1/3)) + (3^(1/2)*(log
(x + 3^(1/2)/3) - log(x - 3^(1/2) - 2*(x^2 + 1)^(1/2)))/18 - (3^(1/2)*(6*
log(x - 3^(1/2)/3) - 6*log(x + 3^(1/2) + 2*(x^2 + 1)^(1/2)))/54 - (3^(1/2)
)*(6*log(x + 3^(1/2)/3) - 6*log(x - 3^(1/2) - 2*(x^2 + 1)^(1/2)))/54 + (3
^(1/2)*(x^2 + 1)^(1/2))/(9*(x - 3^(1/2)/3)) - (3^(1/2)*(x^2 + 1)^(1/2))/(9
*(x + 3^(1/2)/3))
```

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.30

$$\int \frac{1}{(2x + \sqrt{1+x^2})^2} dx$$

$$= \frac{6\sqrt{x^2+1} - 3\sqrt{3} \log(3\sqrt{x^2+1} - \sqrt{3} + 3x) x^2 + \sqrt{3} \log(3\sqrt{x^2+1} - \sqrt{3} + 3x) + 3\sqrt{3} \log(3\sqrt{x^2+1} + \sqrt{3} + 3x)}{27x^2 - 9}$$

input

```
int(1/(2*x+(x^2+1)^(1/2))^2,x)
```


output

```
(6*sqrt(x**2 + 1) - 3*sqrt(3)*log(3*sqrt(x**2 + 1) - sqrt(3) + 3*x)*x**2 +  
sqrt(3)*log(3*sqrt(x**2 + 1) - sqrt(3) + 3*x) + 3*sqrt(3)*log(3*sqrt(x**2  
+ 1) + sqrt(3) + 3*x)*x**2 - sqrt(3)*log(3*sqrt(x**2 + 1) + sqrt(3) + 3*x  
) - 12*x)/(9*(3*x**2 - 1))
```

3.4 $\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx$

Optimal result	89
Mathematica [A] (verified)	89
Rubi [A] (verified)	90
Maple [A] (verified)	91
Fricas [B] (verification not implemented)	92
Sympy [F]	93
Maxima [F]	93
Giac [B] (verification not implemented)	93
Mupad [F(-1)]	94
Reduce [B] (verification not implemented)	94

Optimal result

Integrand size = 19, antiderivative size = 43

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = \frac{3x\sqrt{-1+x^2}}{8(4-3x^2)} + \frac{5}{16} \operatorname{arctanh}\left(\frac{x}{2\sqrt{-1+x^2}}\right)$$

output

```
5/16*arctanh(1/2*x/(x^2-1)^(1/2))+3/8*x*(x^2-1)^(1/2)/(-3*x^2+4)
```

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.65

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = -\frac{3x\sqrt{-1+x^2}}{8(-4+3x^2)} + \frac{5}{32} \log\left(2-x^2+x\sqrt{-1+x^2}\right) - \frac{5}{32} \log\left(2-3x^2+3x\sqrt{-1+x^2}\right)$$

input

```
Integrate[1/(Sqrt[-1+x^2]*(-4+3*x^2)^2),x]
```

output

```
(-3*x*Sqrt[-1+x^2])/(8*(-4+3*x^2)) + (5*Log[2-x^2+x*Sqrt[-1+x^2]])/32 - (5*Log[2-3*x^2+3*x*Sqrt[-1+x^2]])/32
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {296, 25, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2 - 1} (3x^2 - 4)^2} dx$$

$$\downarrow \text{296}$$

$$\frac{3x\sqrt{x^2 - 1}}{8(4 - 3x^2)} - \frac{5}{8} \int -\frac{1}{(4 - 3x^2)\sqrt{x^2 - 1}} dx$$

$$\downarrow \text{25}$$

$$\frac{5}{8} \int \frac{1}{(4 - 3x^2)\sqrt{x^2 - 1}} dx + \frac{3\sqrt{x^2 - 1}x}{8(4 - 3x^2)}$$

$$\downarrow \text{291}$$

$$\frac{5}{8} \int \frac{1}{4 - \frac{x^2}{x^2 - 1}} d\frac{x}{\sqrt{x^2 - 1}} + \frac{3\sqrt{x^2 - 1}x}{8(4 - 3x^2)}$$

$$\downarrow \text{219}$$

$$\frac{5}{16} \operatorname{arctanh}\left(\frac{x}{2\sqrt{x^2 - 1}}\right) + \frac{3\sqrt{x^2 - 1}x}{8(4 - 3x^2)}$$

input `Int[1/(Sqrt[-1 + x^2]*(-4 + 3*x^2)^2),x]`

output `(3*x*Sqrt[-1 + x^2])/(8*(4 - 3*x^2)) + (5*ArcTanh[x/(2*Sqrt[-1 + x^2])])/16`

Defintions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`
- rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`
- rule 296 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))], x] + Simp[(b*c + 2*(p + 1)*(b*c - a*d))/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && EqQ[2*(p + q + 2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.21

method	result
trager	$-\frac{3x\sqrt{x^2-1}}{8(3x^2-4)} + \frac{5 \ln\left(-\frac{4\sqrt{x^2-1}x+5x^2-4}{3x^2-4}\right)}{32}$
pseudoelliptic	$\frac{(-15x^2+20) \ln\left(\frac{2\sqrt{x^2-1}-x}{x}\right) + 15 \ln\left(\frac{x+2\sqrt{x^2-1}}{x}\right) x^2 - 12\sqrt{x^2-1}x - 20 \ln\left(\frac{x+2\sqrt{x^2-1}}{x}\right)}{96x^2-128}$
risch	$-\frac{3x\sqrt{x^2-1}}{8(3x^2-4)} + \frac{5 \operatorname{arctanh}\left(\frac{3\left(\frac{2}{3} + \frac{4\sqrt{3}\left(x-\frac{2\sqrt{3}}{3}\right)}{3}\right)\sqrt{3}}{2\sqrt{9\left(x-\frac{2\sqrt{3}}{3}\right)^2+12\sqrt{3}\left(x-\frac{2\sqrt{3}}{3}\right)+3}}\right)}{32} - \frac{5 \operatorname{arctanh}\left(\frac{3\left(\frac{2}{3} - \frac{4\sqrt{3}\left(x+\frac{2\sqrt{3}}{3}\right)}{3}\right)\sqrt{3}}{2\sqrt{9\left(x+\frac{2\sqrt{3}}{3}\right)^2-12\sqrt{3}\left(x+\frac{2\sqrt{3}}{3}\right)+3}}\right)}{32}$
default	$-\frac{\sqrt{\left(x-\frac{2\sqrt{3}}{3}\right)^2 + \frac{4\sqrt{3}\left(x-\frac{2\sqrt{3}}{3}\right)}{3}} + \frac{1}{3}}{16\left(x-\frac{2\sqrt{3}}{3}\right)} + \frac{5 \operatorname{arctanh}\left(\frac{3\left(\frac{2}{3} + \frac{4\sqrt{3}\left(x-\frac{2\sqrt{3}}{3}\right)}{3}\right)\sqrt{3}}{2\sqrt{9\left(x-\frac{2\sqrt{3}}{3}\right)^2+12\sqrt{3}\left(x-\frac{2\sqrt{3}}{3}\right)+3}}\right)}{32} - \frac{\sqrt{\left(x+\frac{2\sqrt{3}}{3}\right)^2 - \frac{4\sqrt{3}\left(x+\frac{2\sqrt{3}}{3}\right)}{3}}}{16\left(x+\frac{2\sqrt{3}}{3}\right)}$

input `int(1/(3*x^2-4)^2/(x^2-1)^(1/2),x,method=_RETURNVERBOSE)`

output `-3/8*x/(3*x^2-4)*(x^2-1)^(1/2)+5/32*ln(-(4*(x^2-1)^(1/2)*x+5*x^2-4)/(3*x^2-4))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. $2(33) = 66$.

Time = 0.07 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.86

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = \frac{12x^2 + 5(3x^2 - 4) \log(3x^2 - 3\sqrt{x^2-1}x - 2) - 5(3x^2 - 4) \log(x^2 - \sqrt{x^2-1}x - 2) + 12\sqrt{x^2-1}}{32(3x^2 - 4)}$$

input `integrate(1/(3*x^2-4)^2/(x^2-1)^(1/2),x, algorithm="fricas")`

output

```
-1/32*(12*x^2 + 5*(3*x^2 - 4)*log(3*x^2 - 3*sqrt(x^2 - 1)*x - 2) - 5*(3*x^2 - 4)*log(x^2 - sqrt(x^2 - 1)*x - 2) + 12*sqrt(x^2 - 1)*x - 16)/(3*x^2 - 4)
```

Sympy [F]

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = \int \frac{1}{\sqrt{(x-1)(x+1)}(3x^2-4)^2} dx$$

input

```
integrate(1/(3*x**2-4)**2/(x**2-1)**(1/2), x)
```

output

```
Integral(1/(sqrt((x - 1)*(x + 1))*(3*x**2 - 4)**2), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = \int \frac{1}{(3x^2-4)^2\sqrt{x^2-1}} dx$$

input

```
integrate(1/(3*x^2-4)^2/(x^2-1)^(1/2), x, algorithm="maxima")
```

output

```
integrate(1/((3*x^2 - 4)^2*sqrt(x^2 - 1)), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(33) = 66$.

Time = 0.12 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.19

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = \frac{5(x-\sqrt{x^2-1})^2-3}{4\left(3(x-\sqrt{x^2-1})^4-10(x-\sqrt{x^2-1})^2+3\right)} - \frac{5}{32} \log\left(\left|3(x-\sqrt{x^2-1})^2-1\right|\right) + \frac{5}{32} \log\left(\left|(x-\sqrt{x^2-1})^2-3\right|\right)$$

input `integrate(1/(3*x^2-4)^2/(x^2-1)^(1/2),x, algorithm="giac")`

output `1/4*(5*(x - sqrt(x^2 - 1))^2 - 3)/(3*(x - sqrt(x^2 - 1))^4 - 10*(x - sqrt(x^2 - 1))^2 + 3) - 5/32*log(abs(3*(x - sqrt(x^2 - 1))^2 - 1)) + 5/32*log(abs((x - sqrt(x^2 - 1))^2 - 3))`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = \int \frac{1}{\sqrt{x^2-1}(3x^2-4)^2} dx$$

input `int(1/((x^2 - 1)^(1/2)*(3*x^2 - 4)^2),x)`

output `int(1/((x^2 - 1)^(1/2)*(3*x^2 - 4)^2), x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 160, normalized size of antiderivative = 3.72

$$\int \frac{1}{\sqrt{-1+x^2}(-4+3x^2)^2} dx = \frac{-12\sqrt{x^2-1}x - 15\log(\sqrt{x^2-1} - \sqrt{3} + x)x^2 + 20\log(\sqrt{x^2-1} - \sqrt{3} + x) - 15\log(\sqrt{x^2-1} + \sqrt{3} + x)}{}$$

input `int(1/(3*x^2-4)^2/(x^2-1)^(1/2),x)`

output `(- 12*sqrt(x**2 - 1)*x - 15*log(sqrt(x**2 - 1) - sqrt(3) + x)*x**2 + 20*log(sqrt(x**2 - 1) - sqrt(3) + x) - 15*log(sqrt(x**2 - 1) + sqrt(3) + x)*x**2 + 20*log(sqrt(x**2 - 1) + sqrt(3) + x) + 15*log(3*sqrt(x**2 - 1) - sqrt(3) + 3*x)*x**2 - 20*log(3*sqrt(x**2 - 1) - sqrt(3) + 3*x) + 15*log(3*sqrt(x**2 - 1) + sqrt(3) + 3*x)*x**2 - 20*log(3*sqrt(x**2 - 1) + sqrt(3) + 3*x))/ (32*(3*x**2 - 4))`

3.5 $\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx$

Optimal result	96
Mathematica [A] (verified)	96
Rubi [A] (verified)	97
Maple [B] (verified)	98
Fricas [A] (verification not implemented)	98
Sympy [F]	99
Maxima [F]	99
Giac [B] (verification not implemented)	99
Mupad [B] (verification not implemented)	100
Reduce [B] (verification not implemented)	101

Optimal result

Integrand size = 17, antiderivative size = 74

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx = \frac{8}{9(1-3x)} - \frac{4\sqrt{x}\sqrt{1+x}}{3(1-3x)} - \frac{8\operatorname{arcsinh}(\sqrt{x})}{9} + \frac{10}{9}\operatorname{arctanh}\left(\frac{2\sqrt{x}}{\sqrt{1+x}}\right) + \frac{5}{9}\log(1-3x)$$

output `8/9/(1-3*x)-8/9*arcsinh(x^(1/2))+10/9*arctanh(2*x^(1/2)/(1+x)^(1/2))+5/9*ln(1-3*x)-4/3*x^(1/2)*(1+x)^(1/2)/(1-3*x)`

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.99

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx = \frac{2(-4 + 6\sqrt{x}\sqrt{1+x} + (1-3x)\log(-\sqrt{x} + \sqrt{1+x}) + 5(-1+3x)\log(1-x + \sqrt{x}\sqrt{1+x}))}{-9 + 27x}$$

input `Integrate[(2*Sqrt[x] + Sqrt[1 + x])^(-2), x]`

output

```
(2*(-4 + 6*Sqrt[x]*Sqrt[1 + x] + (1 - 3*x)*Log[-Sqrt[x] + Sqrt[1 + x]] + 5
*(-1 + 3*x)*Log[1 - x + Sqrt[x]*Sqrt[1 + x]])/(-9 + 27*x)
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(2\sqrt{x} + \sqrt{x+1})^2} dx$$

↓ 7293

$$\int \left(-\frac{4\sqrt{x}\sqrt{x+1}}{(3x-1)^2} + \frac{5}{3(3x-1)} + \frac{8}{3(3x-1)^2} \right) dx$$

↓ 2009

$$-\frac{8}{9}\operatorname{arcsinh}(\sqrt{x}) + \frac{10}{9}\operatorname{arctanh}\left(\frac{2\sqrt{x}}{\sqrt{x+1}}\right) - \frac{4\sqrt{x}\sqrt{x+1}}{3(1-3x)} + \frac{8}{9(1-3x)} + \frac{5}{9}\log(1-3x)$$

input

```
Int[(2*Sqrt[x] + Sqrt[1 + x])^(-2), x]
```

output

```
8/(9*(1 - 3*x)) - (4*Sqrt[x]*Sqrt[1 + x])/(3*(1 - 3*x)) - (8*ArcSinh[Sqrt[
x]])/9 + (10*ArcTanh[(2*Sqrt[x])/Sqrt[1 + x]])/9 + (5*Log[1 - 3*x])/9
```

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(54) = 108$.

Time = 0.03 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.55

method	result
default	$-\frac{8}{9(3x-1)} + \frac{5 \ln(3x-1)}{9} - \frac{\sqrt{x} \sqrt{1+x} \left(12 \ln\left(\frac{1}{2}+x+\sqrt{x(1+x)}\right) x - 15 \operatorname{arctanh}\left(\frac{5x+1}{4\sqrt{x(1+x)}}\right) x - 4 \ln\left(\frac{1}{2}+x+\sqrt{x(1+x)}\right) + 5 \operatorname{arctanh}\left(\frac{5x+1}{4\sqrt{x(1+x)}}\right) \right)}{9\sqrt{x(1+x)}(3x-1)}$

input `int(1/(2*x^(1/2)+(1+x)^(1/2))^2,x,method=_RETURNVERBOSE)`

output
$$-\frac{8}{9(3x-1)} + \frac{5 \ln(3x-1)}{9} - \frac{1}{9} x^{1/2} (1+x)^{1/2} \left(12 \ln\left(\frac{1}{2}+x+(x(1+x))^{1/2}\right) x - 15 \operatorname{arctanh}\left(\frac{1}{4} \frac{5x+1}{(x(1+x))^{1/2}}\right) x - 4 \ln\left(\frac{1}{2}+x+(x(1+x))^{1/2}\right) + 5 \operatorname{arctanh}\left(\frac{1}{4} \frac{5x+1}{(x(1+x))^{1/2}}\right) - 12 \frac{(x(1+x))^{1/2}}{(x(1+x))^{1/2}} \right) / (3x-1)$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.42

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx = \frac{5(3x-1) \log(3\sqrt{x+1}\sqrt{x} - 3x - 1) - 4(3x-1) \log(2\sqrt{x+1}\sqrt{x} - 2x - 1) - 5(3x-1) \log(\sqrt{x+1})}{9(3x-1)}$$

input `integrate(1/(2*x^(1/2)+(1+x)^(1/2))^2,x, algorithm="fricas")`

output

```
-1/9*(5*(3*x - 1)*log(3*sqrt(x + 1)*sqrt(x) - 3*x - 1) - 4*(3*x - 1)*log(2
*sqrt(x + 1)*sqrt(x) - 2*x - 1) - 5*(3*x - 1)*log(sqrt(x + 1)*sqrt(x) - x
+ 1) - 5*(3*x - 1)*log(3*x - 1) - 12*sqrt(x + 1)*sqrt(x) - 12*x + 12)/(3*x
- 1)
```

Sympy [F]

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx = \int \frac{1}{(2\sqrt{x} + \sqrt{x+1})^2} dx$$

input

```
integrate(1/(2*x**(1/2)+(1+x)**(1/2))**2,x)
```

output

```
Integral((2*sqrt(x) + sqrt(x + 1))**(-2), x)
```

Maxima [F]

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx = \int \frac{1}{(\sqrt{x+1} + 2\sqrt{x})^2} dx$$

input

```
integrate(1/(2*x^(1/2)+(1+x)^(1/2))^2,x, algorithm="maxima")
```

output

```
integrate((sqrt(x + 1) + 2*sqrt(x))^-2, x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 133 vs. $2(54) = 108$.

Time = 0.15 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.80

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx = -\frac{8 \left(5(\sqrt{x+1} - \sqrt{x})^2 - 3 \right)}{9 \left(3(\sqrt{x+1} - \sqrt{x})^4 - 10(\sqrt{x+1} - \sqrt{x})^2 + 3 \right)} - \frac{5x+1}{3(3x-1)} + \frac{4}{9} \log \left(\left(\sqrt{x+1} - \sqrt{x} \right)^2 \right) - \frac{5}{9} \log \left(\left| 3(\sqrt{x+1} - \sqrt{x})^2 - 1 \right| \right) + \frac{5}{9} \log \left(\left| (\sqrt{x+1} - \sqrt{x})^2 - 3 \right| \right) + \frac{5}{9} \log(|3x-1|)$$

input `integrate(1/(2*x^(1/2)+(1+x)^(1/2))^2,x, algorithm="giac")`

output `-8/9*(5*(sqrt(x + 1) - sqrt(x))^2 - 3)/(3*(sqrt(x + 1) - sqrt(x))^4 - 10*(sqrt(x + 1) - sqrt(x))^2 + 3) - 1/3*(5*x + 1)/(3*x - 1) + 4/9*log((sqrt(x + 1) - sqrt(x))^2) - 5/9*log(abs(3*(sqrt(x + 1) - sqrt(x))^2 - 1)) + 5/9*log(abs((sqrt(x + 1) - sqrt(x))^2 - 3)) + 5/9*log(abs(3*x - 1))`

Mupad [B] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.11

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx = \frac{10 \operatorname{atanh} \left(\frac{2662400 \sqrt{x}}{81 \left(\frac{665600 x}{81 (\sqrt{x+1}-1)^2} + \frac{665600}{81} \right) (\sqrt{x+1}-1)} \right)}{9} + \frac{5 \ln \left(x - \frac{1}{3} \right)}{9} - \frac{16 \operatorname{atanh} \left(\frac{\sqrt{x}}{\sqrt{x+1}-1} \right)}{9} - \frac{8}{27 \left(x - \frac{1}{3} \right)} + \frac{4 \sqrt{x} \sqrt{x+1}}{3(3x-1)}$$

input `int(1/((x + 1)^(1/2) + 2*x^(1/2))^2,x)`

output `(10*atanh((2662400*x^(1/2))/(81*((665600*x)/(81*((x + 1)^(1/2) - 1)^2) + 665600/81))*((x + 1)^(1/2) - 1)))/9 + (5*log(x - 1/3))/9 - (16*atanh(x^(1/2)/((x + 1)^(1/2) - 1)))/9 - 8/(27*(x - 1/3)) + (4*x^(1/2)*(x + 1)^(1/2))/(3*(3*x - 1))`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.53

$$\int \frac{1}{(2\sqrt{x} + \sqrt{1+x})^2} dx$$

$$= \frac{12\sqrt{x}\sqrt{x+1} - 54\log(\sqrt{x+1} + \sqrt{x})x + 18\log(\sqrt{x+1} + \sqrt{x}) + 30\log(3\sqrt{x+1} + 3\sqrt{x} - \sqrt{3})x - 1}{2}$$

input

```
int(1/(2*x^(1/2)+(1+x)^(1/2))^2,x)
```

output

```
(2*(6*sqrt(x)*sqrt(x + 1) - 27*log(sqrt(x + 1) + sqrt(x))*x + 9*log(sqrt(x
+ 1) + sqrt(x)) + 15*log(3*sqrt(x + 1) + 3*sqrt(x) - sqrt(3))*x - 5*log(3
*sqrt(x + 1) + 3*sqrt(x) - sqrt(3)) + 15*log(3*sqrt(x + 1) + 3*sqrt(x) + s
qrt(3))*x - 5*log(3*sqrt(x + 1) + 3*sqrt(x) + sqrt(3)) + 6*x - 6))/(9*(3*x
- 1))
```

3.6 $\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx$

Optimal result	102
Mathematica [A] (verified)	102
Rubi [A] (verified)	103
Maple [A] (verified)	105
Fricas [A] (verification not implemented)	105
Sympy [F]	106
Maxima [A] (verification not implemented)	106
Giac [A] (verification not implemented)	107
Mupad [F(-1)]	107
Reduce [F]	107

Optimal result

Integrand size = 17, antiderivative size = 64

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = \frac{\sqrt{-1+x^2}}{i-x} - \frac{i \arctan\left(\frac{1-ix}{\sqrt{2}\sqrt{-1+x^2}}\right)}{\sqrt{2}} + \operatorname{arctanh}\left(\frac{x}{\sqrt{-1+x^2}}\right)$$

output

```
arctanh(x/(x^2-1)^(1/2))-1/2*I*arctan(1/2*(1-I*x)*2^(1/2)/(x^2-1)^(1/2))*2
^(1/2)+(x^2-1)^(1/2)/(I-x)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = \frac{\sqrt{-1+x^2}}{i-x} - \sqrt{2} \operatorname{arctanh}\left(\frac{1+ix-i\sqrt{-1+x^2}}{\sqrt{2}}\right) - \log\left(-x + \sqrt{-1+x^2}\right)$$

input

```
Integrate[Sqrt[-1 + x^2]/(-I + x)^2,x]
```

output

$$\text{Sqrt}[-1 + x^2]/(1 - x) - \text{Sqrt}[2]*\text{ArcTanh}[(1 + I*x - I*\text{Sqrt}[-1 + x^2])/ \text{Sqrt}[2]] - \text{Log}[-x + \text{Sqrt}[-1 + x^2]]$$
Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {492, 25, 605, 224, 219, 488, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt{x^2 - 1}}{(x - i)^2} dx \\ & \quad \downarrow 492 \\ & \int -\frac{x}{(i - x)\sqrt{x^2 - 1}} dx + \frac{\sqrt{x^2 - 1}}{-x + i} \\ & \quad \downarrow 25 \\ & \frac{\sqrt{x^2 - 1}}{-x + i} - \int \frac{x}{(i - x)\sqrt{x^2 - 1}} dx \\ & \quad \downarrow 605 \\ & \int \frac{1}{\sqrt{x^2 - 1}} dx - i \int \frac{1}{(i - x)\sqrt{x^2 - 1}} dx + \frac{\sqrt{x^2 - 1}}{-x + i} \\ & \quad \downarrow 224 \\ & -i \int \frac{1}{(i - x)\sqrt{x^2 - 1}} dx + \int \frac{1}{1 - \frac{x^2}{x^2 - 1}} d \frac{x}{\sqrt{x^2 - 1}} + \frac{\sqrt{x^2 - 1}}{-x + i} \\ & \quad \downarrow 219 \\ & -i \int \frac{1}{(i - x)\sqrt{x^2 - 1}} dx + \text{arctanh}\left(\frac{x}{\sqrt{x^2 - 1}}\right) + \frac{\sqrt{x^2 - 1}}{-x + i} \\ & \quad \downarrow 488 \\ & i \int \frac{1}{-\frac{(1 - ix)^2}{x^2 - 1} - 2} d \frac{1 - ix}{\sqrt{x^2 - 1}} + \text{arctanh}\left(\frac{x}{\sqrt{x^2 - 1}}\right) + \frac{\sqrt{x^2 - 1}}{-x + i} \end{aligned}$$

$$\downarrow 217$$

$$-\frac{i \arctan\left(\frac{1-ix}{\sqrt{2}\sqrt{x^2-1}}\right)}{\sqrt{2}} + \operatorname{arctanh}\left(\frac{x}{\sqrt{x^2-1}}\right) + \frac{\sqrt{x^2-1}}{-x+i}$$

input `Int[Sqrt[-1 + x^2]/(-1 + x)^2,x]`

output `Sqrt[-1 + x^2]/(1 - x) - (1*ArcTan[(1 - 1*x)/(Sqrt[2]*Sqrt[-1 + x^2]))/Sqrt[2] + ArcTanh[x/Sqrt[-1 + x^2]]`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 224 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

rule 488 `Int[1/(((c_) + (d_.)*(x_))*Sqrt[(a_) + (b_.)*(x_)^2]), x_Symbol] := -Subst[Int[1/(b*c^2 + a*d^2 - x^2), x], x, (a*d - b*c*x)/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x]`

rule 492 `Int[((c_) + (d_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[(c + d*x)^(n + 1)*((a + b*x^2)^p/(d*(n + 1))), x] - Simp[2*b*(p/(d*(n + 1)))*Int[x*(c + d*x)^(n + 1)*(a + b*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, n}, x] && GtQ[p, 0] && (IntegerQ[p] || LtQ[n, -1]) && NeQ[n, -1] && !LtQ[n + 2*p + 1, 0] && IntQuadraticQ[a, 0, b, c, d, n, p, x]`

rule 605 `Int[((x_)^(m_))*((a_) + (b_)*(x_)^2)^(p_)/((c_) + (d_)*(x_)), x_Symbol] := Simp[1/d Int[x^(m - 1)*(a + b*x^2)^p, x], x] - Simp[c/d Int[x^(m - 1)*((a + b*x^2)^p/(c + d*x)), x], x] /; FreeQ[{a, b, c, d, p}, x] && IGtQ[m, 0] && LtQ[-1, p, 0]`

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.02

method	result
risch	$-\frac{\sqrt{x^2-1}}{x-i} + \frac{i\sqrt{2} \arctan\left(\frac{(-4+2i(x-i))\sqrt{2}}{4\sqrt{(x-i)^2+2i(x-i)-2}}\right)}{2} + \ln(x + \sqrt{x^2-1})$
default	$\frac{((x-i)^2+2i(x-i)-2)^{\frac{3}{2}}}{2x-2i} - \frac{i\left(\sqrt{(x-i)^2+2i(x-i)-2} + i \ln\left(x + \sqrt{(x-i)^2+2i(x-i)-2}\right) - \sqrt{2} \arctan\left(\frac{(-4+2i(x-i))\sqrt{2}}{4\sqrt{(x-i)^2+2i(x-i)-2}}\right)\right)}{2}$

input `int((x^2-1)^(1/2)/(x-I)^2,x,method=_RETURNVERBOSE)`

output
$$-(x^2-1)^{(1/2)}/(x-I)+1/2*I*2^{(1/2)}*\arctan(1/4*(-4+2*I*(x-I))*2^{(1/2)}/((x-I)^2+2*I*(x-I)-2)^{(1/2)})+\ln(x+(x^2-1)^{(1/2)})$$

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.39

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = \frac{-\sqrt{2}(x-i) \log(-x+i\sqrt{2}+\sqrt{x^2-1}+i) - \sqrt{2}(x-i) \log(-x-i\sqrt{2}+\sqrt{x^2-1}+i) + 2(x-i) \log(x-i)}{2(x-i)}$$

input `integrate((x^2-1)^(1/2)/(-I+x)^2,x, algorithm="fricas")`

output `-1/2*(sqrt(2)*(x - I)*log(-x + I*sqrt(2) + sqrt(x^2 - 1) + I) - sqrt(2)*(x - I)*log(-x - I*sqrt(2) + sqrt(x^2 - 1) + I) + 2*(x - I)*log(-x + sqrt(x^2 - 1)) + 2*x + 2*sqrt(x^2 - 1) - 2*I)/(x - I)`

Sympy [F]

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = \int \frac{\sqrt{(x-1)(x+1)}}{(x-i)^2} dx$$

input `integrate((x**2-1)**(1/2)/(-I+x)**2,x)`

output `Integral(sqrt((x - 1)*(x + 1))/(x - I)**2, x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = \frac{1}{2}i\sqrt{2}\arcsin\left(\frac{ix}{|x-i|} - \frac{1}{|x-i|}\right) - \frac{\sqrt{x^2-1}}{x-i} + \log\left(2x + 2\sqrt{x^2-1}\right)$$

input `integrate((x^2-1)^(1/2)/(-I+x)^2,x, algorithm="maxima")`

output `1/2*I*sqrt(2)*arcsin(I*x/abs(x - I) - 1/abs(x - I)) - sqrt(x^2 - 1)/(x - I) + log(2*x + 2*sqrt(x^2 - 1))`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = i\sqrt{2} \arctan\left(-\frac{1}{2}\sqrt{2}(x - \sqrt{x^2-1} - i)\right) + \frac{2(ix - i\sqrt{x^2-1} - 1)}{(x - \sqrt{x^2-1})^2 - 2ix + 2i\sqrt{x^2-1} + 1} - \log\left(|-x + \sqrt{x^2-1}|\right)$$

input `integrate((x^2-1)^(1/2)/(-I+x)^2,x, algorithm="giac")`

output `I*sqrt(2)*arctan(-1/2*sqrt(2)*(x - sqrt(x^2 - 1) - I)) + 2*(I*x - I*sqrt(x^2 - 1) - 1)/((x - sqrt(x^2 - 1))^2 - 2*I*x + 2*I*sqrt(x^2 - 1) + 1) - log(abs(-x + sqrt(x^2 - 1)))`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = \int \frac{\sqrt{x^2-1}}{(x-i)^2} dx$$

input `int((x^2 - 1)^(1/2)/(x - 1i)^2,x)`

output `int((x^2 - 1)^(1/2)/(x - 1i)^2, x)`

Reduce [F]

$$\int \frac{\sqrt{-1+x^2}}{(-i+x)^2} dx = -\left(\int \frac{\sqrt{x^2-1}}{2ix - x^2 + 1} dx\right)$$

input `int((x^2-1)^(1/2)/(-I+x)^2,x)`

output `- int(sqrt(x**2 - 1)/(2*i*x - x**2 + 1),x)`

$$3.7 \quad \int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx$$

Optimal result	109
Mathematica [A] (verified)	109
Rubi [A] (verified)	110
Maple [A] (verified)	111
Fricas [B] (verification not implemented)	112
Sympy [F]	112
Maxima [F]	113
Giac [B] (verification not implemented)	113
Mupad [F(-1)]	114
Reduce [B] (verification not implemented)	114

Optimal result

Integrand size = 17, antiderivative size = 48

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx = -\frac{x\sqrt{-1+x^2}}{4(1+x^2)} + \frac{3\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{-1+x^2}}\right)}{4\sqrt{2}}$$

output `3/8*arctanh(x*2^(1/2)/(x^2-1)^(1/2))*2^(1/2)-1/4*x*(x^2-1)^(1/2)/(x^2+1)`

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.15

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx = \frac{1}{8} \left(-\frac{2x\sqrt{-1+x^2}}{1+x^2} + 3\sqrt{2}\operatorname{arctanh}\left(\frac{1+x^2-x\sqrt{-1+x^2}}{\sqrt{2}}\right) \right)$$

input `Integrate[1/(Sqrt[-1 + x^2]*(1 + x^2)^2), x]`

output `((-2*x*Sqrt[-1 + x^2])/(1 + x^2) + 3*Sqrt[2]*ArcTanh[(1 + x^2 - x*Sqrt[-1 + x^2])/Sqrt[2]])/8`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {296, 291, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt{x^2-1}(x^2+1)^2} dx$$

$$\downarrow \text{296}$$

$$\frac{3}{4} \int \frac{1}{\sqrt{x^2-1}(x^2+1)} dx - \frac{x\sqrt{x^2-1}}{4(x^2+1)}$$

$$\downarrow \text{291}$$

$$\frac{3}{4} \int \frac{1}{1-\frac{2x^2}{x^2-1}} d\frac{x}{\sqrt{x^2-1}} - \frac{x\sqrt{x^2-1}}{4(x^2+1)}$$

$$\downarrow \text{219}$$

$$\frac{3\text{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{x^2-1}}\right)}{4\sqrt{2}} - \frac{x\sqrt{x^2-1}}{4(x^2+1)}$$

input `Int [1/(Sqrt [-1 + x^2] *(1 + x^2)^2), x]`

output `-1/4*(x*Sqrt [-1 + x^2])/(1 + x^2) + (3*ArcTanh [(Sqrt [2]*x)/Sqrt [-1 + x^2]])/(4*Sqrt [2])`

Defintions of rubi rules used

- rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

- rule 291 `Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*((c_) + (d_.)*(x_)^2)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

- rule 296 `Int[((a_) + (b_.)*(x_)^2)^(p_)*((c_) + (d_.)*(x_)^2)^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^2)^(p + 1)*((c + d*x^2)^(q + 1)/(2*a*(p + 1)*(b*c - a*d))], x] + Simp[(b*c + 2*(p + 1)*(b*c - a*d))/(2*a*(p + 1)*(b*c - a*d)) Int[(a + b*x^2)^(p + 1)*(c + d*x^2)^q, x], x] /; FreeQ[{a, b, c, d, q}, x] && NeQ[b*c - a*d, 0] && EqQ[2*(p + q + 2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]`

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.77

method	result	size
risch	$\frac{3 \operatorname{arctanh}\left(\frac{x\sqrt{2}}{\sqrt{x^2-1}}\right)\sqrt{2}}{8} - \frac{x\sqrt{x^2-1}}{4(x^2+1)}$	37
default	$-\frac{x}{8\sqrt{x^2-1}\left(\frac{x^2}{x^2-1} - \frac{1}{2}\right)} + \frac{3 \operatorname{arctanh}\left(\frac{x\sqrt{2}}{\sqrt{x^2-1}}\right)\sqrt{2}}{8}$	45
pseudoelliptic	$\frac{(3x^2+3)\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{x^2-1}}{2x}\right) - 2\sqrt{x^2-1}x}{8x^2+8}$	49
trager	$-\frac{x\sqrt{x^2-1}}{4(x^2+1)} + \frac{3 \operatorname{RootOf}\left(_Z^2 - 2\right) \ln\left(\frac{3 \operatorname{RootOf}\left(_Z^2 - 2\right) x^2 + 4\sqrt{x^2-1}x - \operatorname{RootOf}\left(_Z^2 - 2\right)}{x^2+1}\right)}{16}$	66

input `int(1/(x^2+1)^2/(x^2-1)^(1/2), x, method=_RETURNVERBOSE)`

output `3/8*arctanh(x*2^(1/2)/(x^2-1)^(1/2))*2^(1/2)-1/4*x*(x^2-1)^(1/2)/(x^2+1)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 83 vs. $2(36) = 72$.

Time = 0.07 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.73

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx$$

$$= \frac{3\sqrt{2}(x^2+1)\log\left(\frac{9x^2+2\sqrt{2}(3x^2-1)+2\sqrt{x^2-1}(3\sqrt{2}x+4x)-3}{x^2+1}\right) - 4x^2 - 4\sqrt{x^2-1}x - 4}{16(x^2+1)}$$

input `integrate(1/(x^2+1)^2/(x^2-1)^(1/2),x, algorithm="fricas")`

output `1/16*(3*sqrt(2)*(x^2 + 1)*log((9*x^2 + 2*sqrt(2)*(3*x^2 - 1) + 2*sqrt(x^2 - 1)*(3*sqrt(2)*x + 4*x) - 3)/(x^2 + 1)) - 4*x^2 - 4*sqrt(x^2 - 1)*x - 4)/(x^2 + 1)`

Sympy [F]

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx = \int \frac{1}{\sqrt{(x-1)(x+1)}(x^2+1)^2} dx$$

input `integrate(1/(x**2+1)**2/(x**2-1)**(1/2),x)`

output `Integral(1/(sqrt((x - 1)*(x + 1))*(x**2 + 1)**2), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx = \int \frac{1}{(x^2+1)^2\sqrt{x^2-1}} dx$$

input `integrate(1/(x^2+1)^2/(x^2-1)^(1/2),x, algorithm="maxima")`

output `integrate(1/((x^2 + 1)^2*sqrt(x^2 - 1)), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 101 vs. $2(36) = 72$.

Time = 0.13 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.10

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx = -\frac{3}{16}\sqrt{2}\log\left(\frac{(x-\sqrt{x^2-1})^2-2\sqrt{2}+3}{(x-\sqrt{x^2-1})^2+2\sqrt{2}+3}\right) - \frac{3(x-\sqrt{x^2-1})^2+1}{2((x-\sqrt{x^2-1})^4+6(x-\sqrt{x^2-1})^2+1)}$$

input `integrate(1/(x^2+1)^2/(x^2-1)^(1/2),x, algorithm="giac")`

output `-3/16*sqrt(2)*log(((x - sqrt(x^2 - 1))^2 - 2*sqrt(2) + 3)/((x - sqrt(x^2 - 1))^2 + 2*sqrt(2) + 3)) - 1/2*(3*(x - sqrt(x^2 - 1))^2 + 1)/((x - sqrt(x^2 - 1))^4 + 6*(x - sqrt(x^2 - 1))^2 + 1)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx = \int \frac{1}{\sqrt{x^2-1}(x^2+1)^2} dx$$

input `int(1/((x^2 - 1)^(1/2)*(x^2 + 1)^2),x)`output `int(1/((x^2 - 1)^(1/2)*(x^2 + 1)^2), x)`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 157, normalized size of antiderivative = 3.27

$$\int \frac{1}{\sqrt{-1+x^2}(1+x^2)^2} dx$$

$$= \frac{-4\sqrt{x^2-1}x + 3\sqrt{2}\log(\sqrt{x^2-1} - \sqrt{2}i + i + x)x^2 + 3\sqrt{2}\log(\sqrt{x^2-1} - \sqrt{2}i + i + x) + 3\sqrt{2}\log(\sqrt{x^2-1} + \sqrt{2}i - i + x)x^2 + 3\sqrt{2}\log(\sqrt{x^2-1} + \sqrt{2}i - i + x) - 3\sqrt{2}\log(2\sqrt{x^2-1}x + 2\sqrt{2} + 2x^2 + 2)x^2 - 3\sqrt{2}\log(2\sqrt{x^2-1}x + 2\sqrt{2} + 2x^2 + 2))/(16(x^2 + 1))$$

input `int(1/(x^2+1)^2/(x^2-1)^(1/2),x)`output `(- 4*sqrt(x**2 - 1)*x + 3*sqrt(2)*log(sqrt(x**2 - 1) - sqrt(2)*i + i + x) *x**2 + 3*sqrt(2)*log(sqrt(x**2 - 1) - sqrt(2)*i + i + x) + 3*sqrt(2)*log(sqrt(x**2 - 1) + sqrt(2)*i - i + x)*x**2 + 3*sqrt(2)*log(sqrt(x**2 - 1) + sqrt(2)*i - i + x) - 3*sqrt(2)*log(2*sqrt(x**2 - 1)*x + 2*sqrt(2) + 2*x**2 + 2)*x**2 - 3*sqrt(2)*log(2*sqrt(x**2 - 1)*x + 2*sqrt(2) + 2*x**2 + 2))/(16*(x**2 + 1))`

$$3.8 \quad \int \frac{1}{(\sqrt{-1+x}+\sqrt{x})^2 \sqrt{-1+x}} dx$$

Optimal result	115
Mathematica [A] (verified)	115
Rubi [A] (verified)	116
Maple [A] (verified)	117
Fricas [A] (verification not implemented)	117
Sympy [B] (verification not implemented)	117
Maxima [F]	118
Giac [A] (verification not implemented)	118
Mupad [B] (verification not implemented)	118
Reduce [B] (verification not implemented)	119

Optimal result

Integrand size = 23, antiderivative size = 30

$$\int \frac{1}{(\sqrt{-1+x}+\sqrt{x})^2 \sqrt{-1+x}} dx = 2\sqrt{-1+x} + \frac{4}{3}(-1+x)^{3/2} - \frac{4x^{3/2}}{3}$$

output $4/3*(-1+x)^{(3/2)}-4/3*x^{(3/2)}+2*(-1+x)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

$$\int \frac{1}{(\sqrt{-1+x}+\sqrt{x})^2 \sqrt{-1+x}} dx = -\frac{4x^{3/2}}{3} + \frac{2}{3}\sqrt{-1+x}(1+2x)$$

input `Integrate[1/((Sqrt[-1 + x] + Sqrt[x])^2*Sqrt[-1 + x]),x]`

output $(-4*x^{(3/2)})/3 + (2*Sqrt[-1 + x]*(1 + 2*x))/3$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {7240, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(\sqrt{x-1} + \sqrt{x})^2 \sqrt{x-1}} dx$$

↓ 7240

$$\int \left(\frac{2x}{\sqrt{x-1}} - 2\sqrt{x} - \frac{1}{\sqrt{x-1}} \right) dx$$

↓ 2009

$$-\frac{4x^{3/2}}{3} + \frac{4}{3}(x-1)^{3/2} + 2\sqrt{x-1}$$

input `Int[1/((Sqrt[-1 + x] + Sqrt[x])^2*Sqrt[-1 + x]),x]`

output `2*Sqrt[-1 + x] + (4*(-1 + x)^(3/2))/3 - (4*x^(3/2))/3`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7240 `Int[(u_.)*((e_.)*Sqrt[(a_.) + (b_.)*(x_)^(n_.)] + (f_.)*Sqrt[(c_.) + (d_.)*(x_)^(n_.)])^(m_), x_Symbol] := Simp[(a*e^2 - c*f^2)^m Int[ExpandIntegrand[u/(e*Sqrt[a + b*x^n] - f*Sqrt[c + d*x^n])^m, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && ILtQ[m, 0] && EqQ[b*e^2 - d*f^2, 0]`

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{4(-1+x)^{\frac{3}{2}}}{3} - \frac{4x^{\frac{3}{2}}}{3} + 2\sqrt{-1+x}$	21

input `int(1/(-1+x)^(1/2)/((-1+x)^(1/2)+x^(1/2))^2,x,method=_RETURNVERBOSE)`

output `4/3*(-1+x)^(3/2)-4/3*x^(3/2)+2*(-1+x)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.60

$$\int \frac{1}{(\sqrt{-1+x} + \sqrt{x})^2 \sqrt{-1+x}} dx = \frac{2}{3} (2x+1)\sqrt{x-1} - \frac{4}{3}x^{\frac{3}{2}}$$

input `integrate(1/(-1+x)^(1/2)/((-1+x)^(1/2)+x^(1/2))^2,x, algorithm="fricas")`

output `2/3*(2*x + 1)*sqrt(x - 1) - 4/3*x^(3/2)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(26) = 52.

Time = 0.22 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.77

$$\int \frac{1}{(\sqrt{-1+x} + \sqrt{x})^2 \sqrt{-1+x}} dx = -\frac{4\sqrt{x}}{6\sqrt{x}\sqrt{x-1} + 6x - 3} - \frac{2\sqrt{x-1}}{6\sqrt{x}\sqrt{x-1} + 6x - 3}$$

input `integrate(1/(-1+x)**(1/2)/((-1+x)**(1/2)+x**(1/2))**2,x)`

output
$$\frac{-4\sqrt{x}}{(6\sqrt{x})\sqrt{x-1} + 6x - 3} - \frac{2\sqrt{x-1}}{(6\sqrt{x})\sqrt{x-1} + 6x - 3}$$

Maxima [F]

$$\int \frac{1}{(\sqrt{-1+x} + \sqrt{x})^2 \sqrt{-1+x}} dx = \int \frac{1}{\sqrt{x-1}(\sqrt{x-1} + \sqrt{x})^2} dx$$

input `integrate(1/(-1+x)^(1/2)/((-1+x)^(1/2)+x^(1/2))^2,x, algorithm="maxima")`

output `integrate(1/(sqrt(x - 1)*(sqrt(x - 1) + sqrt(x))^2), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.67

$$\int \frac{1}{(\sqrt{-1+x} + \sqrt{x})^2 \sqrt{-1+x}} dx = \frac{4}{3} (x-1)^{\frac{3}{2}} - \frac{4}{3} x^{\frac{3}{2}} + 2\sqrt{x-1}$$

input `integrate(1/(-1+x)^(1/2)/((-1+x)^(1/2)+x^(1/2))^2,x, algorithm="giac")`

output
$$\frac{4}{3}(x-1)^{3/2} - \frac{4}{3}x^{3/2} + 2\sqrt{x-1}$$

Mupad [B] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.70

$$\int \frac{1}{(\sqrt{-1+x} + \sqrt{x})^2 \sqrt{-1+x}} dx = \frac{4x\sqrt{x-1}}{3} + \frac{2\sqrt{x-1}}{3} - \frac{4x^{3/2}}{3}$$

input `int(1/(((x - 1)^(1/2) + x^(1/2))^2*(x - 1)^(1/2)),x)`

output $(4*x*(x - 1)^{(1/2)})/3 + (2*(x - 1)^{(1/2)})/3 - (4*x^{(3/2)})/3$

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.63

$$\int \frac{1}{(\sqrt{-1+x} + \sqrt{x})^2 \sqrt{-1+x}} dx = \frac{4\sqrt{x-1}x}{3} + \frac{2\sqrt{x-1}}{3} - \frac{4\sqrt{x}x}{3}$$

input `int(1/(-1+x)^(1/2)/((-1+x)^(1/2)+x^(1/2))^2,x)`

output $(2*(2*\text{sqrt}(x - 1)*x + \text{sqrt}(x - 1) - 2*\text{sqrt}(x)*x))/3$

$$3.9 \quad \int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx$$

Optimal result	120
Mathematica [A] (verified)	121
Rubi [A] (verified)	121
Maple [B] (verified)	123
Fricas [B] (verification not implemented)	124
Sympy [F]	125
Maxima [F]	125
Giac [B] (verification not implemented)	126
Mupad [F(-1)]	127
Reduce [F]	127

Optimal result

Integrand size = 27, antiderivative size = 220

$$\begin{aligned} & \int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx \\ &= \frac{2-4x}{5(\sqrt{x}+\sqrt{-1+x^2})} + \frac{1}{25}\sqrt{-110+50\sqrt{5}} \arctan\left(\frac{1}{2}\sqrt{2+2\sqrt{5}\sqrt{x}}\right) \\ & \quad - \frac{1}{50}\sqrt{-110+50\sqrt{5}} \arctan\left(\frac{\sqrt{-2+2\sqrt{5}\sqrt{-1+x^2}}}{2-(1-\sqrt{5})x}\right) \\ & \quad - \frac{1}{25}\sqrt{110+50\sqrt{5}} \operatorname{arctanh}\left(\frac{1}{2}\sqrt{-2+2\sqrt{5}\sqrt{x}}\right) \\ & \quad - \frac{1}{50}\sqrt{110+50\sqrt{5}} \operatorname{arctanh}\left(\frac{\sqrt{2+2\sqrt{5}\sqrt{-1+x^2}}}{2-x-\sqrt{5}x}\right) \end{aligned}$$

output

```
1/5*(2-4*x)/(x^(1/2)+(x^2-1)^(1/2))-1/50*arctan((x^2-1)^(1/2)*(-2+2*5^(1/2))^(1/2)/(2-x*(-5^(1/2)+1)))*(-110+50*5^(1/2))^(1/2)+1/25*arctan(1/2*x^(1/2)*(2+2*5^(1/2))^(1/2))*(-110+50*5^(1/2))^(1/2)-1/25*arctanh(1/2*x^(1/2)*(-2+2*5^(1/2))^(1/2))*(110+50*5^(1/2))^(1/2)-1/50*arctanh((x^2-1)^(1/2)*(2+2*5^(1/2))^(1/2)/(2-x-x*5^(1/2)))*(110+50*5^(1/2))^(1/2)
```

Mathematica [A] (verified)

Time = 6.19 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx$$

$$= \frac{1}{25} \left(-\frac{10(-1+2x)(-\sqrt{x}+\sqrt{-1+x^2})}{-1-x+x^2} \right.$$

$$+ \sqrt{-110+50\sqrt{5}} \arctan \left(\sqrt{\frac{1}{2}(1+\sqrt{5})} \sqrt{x} \right)$$

$$- \sqrt{-110+50\sqrt{5}} \arctan \left(\frac{\sqrt{-2+\sqrt{5}\sqrt{-1+x^2}}}{1+x} \right)$$

$$- \sqrt{110+50\sqrt{5}} \operatorname{arctanh} \left(\sqrt{\frac{1}{2}(-1+\sqrt{5})} \sqrt{x} \right)$$

$$\left. + \sqrt{110+50\sqrt{5}} \operatorname{arctanh} \left(\frac{\sqrt{2+\sqrt{5}\sqrt{-1+x^2}}}{1+x} \right) \right)$$

input `Integrate[1/(Sqrt[-1 + x^2]*(Sqrt[x] + Sqrt[-1 + x^2])^2),x]`

output `((-10*(-1 + 2*x)*(-Sqrt[x] + Sqrt[-1 + x^2]))/(-1 - x + x^2) + Sqrt[-110 + 50*Sqrt[5]]*ArcTan[Sqrt[(1 + Sqrt[5])/2]*Sqrt[x]] - Sqrt[-110 + 50*Sqrt[5]]*ArcTan[(Sqrt[-2 + Sqrt[5]]*Sqrt[-1 + x^2])/(1 + x)] - Sqrt[110 + 50*Sqrt[5]]*ArcTanh[Sqrt[(-1 + Sqrt[5])/2]*Sqrt[x]] + Sqrt[110 + 50*Sqrt[5]]*ArcTanh[(Sqrt[2 + Sqrt[5]]*Sqrt[-1 + x^2])/(1 + x)])/25`

Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 365, normalized size of antiderivative = 1.66, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{1}{\sqrt{x^2-1}(\sqrt{x^2-1}+\sqrt{x})^2} dx \\
& \quad \downarrow \text{7293} \\
& \int \left(\frac{2x}{\sqrt{x^2-1}(x^2-x-1)^2} - \frac{2\sqrt{x}}{(x^2-x-1)^2} + \frac{1}{\sqrt{x^2-1}(x^2-x-1)} \right) dx \\
& \quad \downarrow \text{2009} \\
& -\frac{2}{5}\sqrt{\frac{1}{5}(5\sqrt{5}-2)} \arctan\left(\frac{2-(1-\sqrt{5})x}{\sqrt{2(\sqrt{5}-1)\sqrt{x^2-1}}}\right) + \\
& \sqrt{\frac{2}{5(\sqrt{5}-1)}} \arctan\left(\frac{2-(1-\sqrt{5})x}{\sqrt{2(\sqrt{5}-1)\sqrt{x^2-1}}}\right) + \frac{1}{5}\sqrt{\frac{2}{5}(5\sqrt{5}-11)} \arctan\left(\sqrt{\frac{2}{\sqrt{5}-1}}\sqrt{x}\right) - \\
& \frac{2}{5}\sqrt{\frac{1}{5}(2+5\sqrt{5})} \operatorname{arctanh}\left(\frac{2-(1+\sqrt{5})x}{\sqrt{2(1+\sqrt{5})\sqrt{x^2-1}}}\right) + \\
& \sqrt{\frac{2}{5(1+\sqrt{5})}} \operatorname{arctanh}\left(\frac{2-(1+\sqrt{5})x}{\sqrt{2(1+\sqrt{5})\sqrt{x^2-1}}}\right) - \\
& \frac{1}{5}\sqrt{\frac{2}{5}(11+5\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{1+\sqrt{5}}}\sqrt{x}\right) - \frac{2\sqrt{x^2-1}(1-2x)}{5(-x^2+x+1)} + \frac{2\sqrt{x}(1-2x)}{5(-x^2+x+1)}
\end{aligned}$$

input `Int[1/(Sqrt[-1 + x^2]*(Sqrt[x] + Sqrt[-1 + x^2])^2),x]`

output `(2*(1 - 2*x)*Sqrt[x])/(5*(1 + x - x^2)) - (2*(1 - 2*x)*Sqrt[-1 + x^2])/(5*(1 + x - x^2)) + (Sqrt[(2*(-11 + 5*Sqrt[5]))/5]*ArcTan[Sqrt[2/(-1 + Sqrt[5]])]*Sqrt[x])/5 + Sqrt[2/(5*(-1 + Sqrt[5]))]*ArcTan[(2 - (1 - Sqrt[5])*x)/(Sqrt[2*(-1 + Sqrt[5]])*Sqrt[-1 + x^2])]) - (2*Sqrt[(-2 + 5*Sqrt[5])/5]*ArcTan[(2 - (1 - Sqrt[5])*x)/(Sqrt[2*(-1 + Sqrt[5]])*Sqrt[-1 + x^2])])/5 - (Sqrt[(2*(11 + 5*Sqrt[5]))/5]*ArcTanh[Sqrt[2/(1 + Sqrt[5]])]*Sqrt[x])/5 + Sqrt[2/(5*(1 + Sqrt[5]))]*ArcTanh[(2 - (1 + Sqrt[5])*x)/(Sqrt[2*(1 + Sqrt[5]])*Sqrt[-1 + x^2])]) - (2*Sqrt[(2 + 5*Sqrt[5])/5]*ArcTanh[(2 - (1 + Sqrt[5])*x)/(Sqrt[2*(1 + Sqrt[5]])*Sqrt[-1 + x^2])])/5`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :=> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7293 `Int[u_, x_Symbol] :=> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1205 vs. $2(158) = 316$.

Time = 0.21 (sec) , antiderivative size = 1206, normalized size of antiderivative = 5.48

method	result	size
default	Expression too large to display	1206

input `int(1/(x^2-1)^(1/2)/(x^(1/2)+(x^2-1)^(1/2))^2,x,method=_RETURNVERBOSE)`

output

```

-6/25*5^(1/2)/(2+2*5^(1/2))^(1/2)*arctanh(2*(1+5^(1/2)+(5^(1/2)+1)*(x-1/2*
5^(1/2)-1/2))/(2+2*5^(1/2))^(1/2)/(4*(x-1/2*5^(1/2)-1/2)^2+4*(5^(1/2)+1)*(
x-1/2*5^(1/2)-1/2)+2+2*5^(1/2))^(1/2))-6/25*5^(1/2)/(-2+2*5^(1/2))^(1/2)*a
rctan(2*(1-5^(1/2)+(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2))/(-2+2*5^(1/2))^(1/2)/
(4*(x+1/2*5^(1/2)-1/2)^2+4*(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2)+2-2*5^(1/2))^(
1/2))+2/5-2/5*5^(1/2))*(-1/4/(1/2-1/2*5^(1/2)))/(x+1/2*5^(1/2)-1/2)*((x+1/
2*5^(1/2)-1/2)^2+(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2)+1/2-1/2*5^(1/2))^(1/2)-1
/4*(-5^(1/2)+1)/(1/2-1/2*5^(1/2)))/(-2+2*5^(1/2))^(1/2)*arctan(2*(1-5^(1/2)
+(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2))/(-2+2*5^(1/2))^(1/2)/(4*(x+1/2*5^(1/2)-
1/2)^2+4*(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2)+2-2*5^(1/2))^(1/2)))+(2/5+2/5*5^
(1/2))*(-1/4/(1/2+1/2*5^(1/2)))/(x-1/2*5^(1/2)-1/2)*((x-1/2*5^(1/2)-1/2)^2+
(5^(1/2)+1)*(x-1/2*5^(1/2)-1/2)+1/2+1/2*5^(1/2))^(1/2)+1/4*(5^(1/2)+1)/(1/
2+1/2*5^(1/2))/(2+2*5^(1/2))^(1/2)*arctanh(2*(1+5^(1/2)+(5^(1/2)+1)*(x-1/2
*5^(1/2)-1/2))/(2+2*5^(1/2))^(1/2)/(4*(x-1/2*5^(1/2)-1/2)^2+4*(5^(1/2)+1)*
(x-1/2*5^(1/2)-1/2)+2+2*5^(1/2))^(1/2))+1/5*(5^(1/2)-1)^2*(-1/4/(1/2-1/2*
5^(1/2)))/(x+1/2*5^(1/2)-1/2)*((x+1/2*5^(1/2)-1/2)^2+(-5^(1/2)+1)*(x+1/2*5^
(1/2)-1/2)+1/2-1/2*5^(1/2))^(1/2)-1/4*(-5^(1/2)+1)/(1/2-1/2*5^(1/2)))/(-2+2
*5^(1/2))^(1/2)*arctan(2*(1-5^(1/2)+(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2))/(-2+
2*5^(1/2))^(1/2)/(4*(x+1/2*5^(1/2)-1/2)^2+4*(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/
2)+2-2*5^(1/2))^(1/2))+1/5*(5^(1/2)+1)^2*(-1/4/(1/2+1/2*5^(1/2)))/(x-1/...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 341 vs. $2(153) = 306$.

Time = 0.09 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.55

$$\int \frac{1}{\sqrt{-1+x^2} (\sqrt{x} + \sqrt{-1+x^2})^2} dx$$

$$= \frac{2(x^2 - x - 1)\sqrt{\frac{1}{2}\sqrt{5} - \frac{11}{10}} \arctan\left(\sqrt{x}(2\sqrt{5} + 5)\sqrt{\frac{1}{2}\sqrt{5} - \frac{11}{10}}\right) - 2(x^2 - x - 1)\sqrt{\frac{1}{2}\sqrt{5} - \frac{11}{10}} \arctan\left(\dots\right)}{\dots}$$

input

```
integrate(1/(x^2-1)^(1/2)/(x^(1/2)+(x^2-1)^(1/2))^2,x, algorithm="fricas")
```

output

```
1/5*(2*(x^2 - x - 1)*sqrt(1/2*sqrt(5) - 11/10)*arctan(sqrt(x)*(2*sqrt(5) +
5)*sqrt(1/2*sqrt(5) - 11/10)) - 2*(x^2 - x - 1)*sqrt(1/2*sqrt(5) - 11/10)
*arctan(-1/2*(sqrt(5)*(4*x + 3) - 2*sqrt(x^2 - 1)*(2*sqrt(5) + 5) + 10*x +
5)*sqrt(1/2*sqrt(5) - 11/10)) + (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11/10)*l
og((3*sqrt(5) - 5)*sqrt(1/2*sqrt(5) + 11/10) - 2*x + sqrt(5) + 2*sqrt(x^2
- 1) + 1) - (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11/10)*log((3*sqrt(5) - 5)*sq
rt(1/2*sqrt(5) + 11/10) + 2*sqrt(x)) - (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11
/10)*log(-(3*sqrt(5) - 5)*sqrt(1/2*sqrt(5) + 11/10) - 2*x + sqrt(5) + 2*sq
rt(x^2 - 1) + 1) + (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11/10)*log(-(3*sqrt(5)
- 5)*sqrt(1/2*sqrt(5) + 11/10) + 2*sqrt(x)) - 4*x^2 - 2*sqrt(x^2 - 1)*(2*
x - 1) + 2*(2*x - 1)*sqrt(x) + 4*x + 4)/(x^2 - x - 1)
```

Sympy [F]

$$\int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx = \int \frac{1}{\sqrt{(x-1)(x+1)}(\sqrt{x}+\sqrt{x^2-1})^2} dx$$

input

```
integrate(1/(x**2-1)**(1/2)/(x**(1/2)+(x**2-1)**(1/2))**2,x)
```

output

```
Integral(1/(sqrt((x - 1)*(x + 1))*(sqrt(x) + sqrt(x**2 - 1))**2), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx = \int \frac{1}{\sqrt{x^2-1}(\sqrt{x^2-1}+\sqrt{x})^2} dx$$

input

```
integrate(1/(x^2-1)^(1/2)/(x^(1/2)+(x^2-1)^(1/2))^2,x, algorithm="maxima")
```

output

```
integrate(1/(sqrt(x^2 - 1)*(sqrt(x^2 - 1) + sqrt(x))^2), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 367 vs. $2(153) = 306$.

Time = 1.31 (sec) , antiderivative size = 367, normalized size of antiderivative = 1.67

$$\begin{aligned}
& \int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx \\
&= \frac{2}{5} \sqrt{\frac{1}{10}} \sqrt{5\sqrt{5}-11} \arctan\left(\frac{2x+\sqrt{5}-2\sqrt{x^2-1}-1}{\sqrt{2\sqrt{5}-2}}\right) \\
&+ \frac{1}{5} \sqrt{\frac{1}{10}} \sqrt{5\sqrt{5}+11} \log\left(\left|-153040x+22956\sqrt{5}\sqrt{50\sqrt{5}+110}+76520\sqrt{5}+153040\sqrt{x^2-1}-382\right|\right) \\
&- \frac{1}{5} \sqrt{\frac{1}{10}} \sqrt{5\sqrt{5}+11} \log\left(\left|-153040x-22956\sqrt{5}\sqrt{50\sqrt{5}+110}+76520\sqrt{5}+153040\sqrt{x^2-1}+382\right|\right) \\
&+ \frac{1}{25} \sqrt{50\sqrt{5}-110} \arctan\left(\frac{\sqrt{x}}{\sqrt{\frac{1}{2}\sqrt{5}-\frac{1}{2}}}\right) \\
&- \frac{1}{50} \sqrt{50\sqrt{5}+110} \log\left(\sqrt{x}+\sqrt{\frac{1}{2}\sqrt{5}+\frac{1}{2}}\right) \\
&+ \frac{1}{50} \sqrt{50\sqrt{5}+110} \log\left(\left|\sqrt{x}-\sqrt{\frac{1}{2}\sqrt{5}+\frac{1}{2}}\right|\right) \\
&+ \frac{4\left((x-\sqrt{x^2-1})^3+2(x-\sqrt{x^2-1})^2+3x-3\sqrt{x^2-1}-2\right)}{5\left((x-\sqrt{x^2-1})^4-2(x-\sqrt{x^2-1})^3-2(x-\sqrt{x^2-1})^2-2x+2\sqrt{x^2-1}+1\right)} \\
&+ \frac{2\left(2x^{\frac{3}{2}}-\sqrt{x}\right)}{5(x^2-x-1)}
\end{aligned}$$

input `integrate(1/(x^2-1)^(1/2)/(x^(1/2)+(x^2-1)^(1/2))^2,x, algorithm="giac")`

output

```
2/5*sqrt(1/10)*sqrt(5*sqrt(5) - 11)*arctan((2*x + sqrt(5) - 2*sqrt(x^2 - 1)
) - 1)/sqrt(2*sqrt(5) - 2)) + 1/5*sqrt(1/10)*sqrt(5*sqrt(5) + 11)*log(abs(
-153040*x + 22956*sqrt(5)*sqrt(50*sqrt(5) + 110) + 76520*sqrt(5) + 153040*
sqrt(x^2 - 1) - 38260*sqrt(50*sqrt(5) + 110) + 76520)) - 1/5*sqrt(1/10)*sq
rt(5*sqrt(5) + 11)*log(abs(-153040*x - 22956*sqrt(5)*sqrt(50*sqrt(5) + 110
) + 76520*sqrt(5) + 153040*sqrt(x^2 - 1) + 38260*sqrt(50*sqrt(5) + 110) +
76520)) + 1/25*sqrt(50*sqrt(5) - 110)*arctan(sqrt(x)/sqrt(1/2*sqrt(5) - 1/
2)) - 1/50*sqrt(50*sqrt(5) + 110)*log(sqrt(x) + sqrt(1/2*sqrt(5) + 1/2)) +
1/50*sqrt(50*sqrt(5) + 110)*log(abs(sqrt(x) - sqrt(1/2*sqrt(5) + 1/2))) +
4/5*((x - sqrt(x^2 - 1))^3 + 2*(x - sqrt(x^2 - 1))^2 + 3*x - 3*sqrt(x^2 -
1) - 2)/((x - sqrt(x^2 - 1))^4 - 2*(x - sqrt(x^2 - 1))^3 - 2*(x - sqrt(x^
2 - 1))^2 - 2*x + 2*sqrt(x^2 - 1) + 1) + 2/5*(2*x^(3/2) - sqrt(x))/(x^2 -
x - 1)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx = \int \frac{1}{\sqrt{x^2-1}(\sqrt{x^2-1}+\sqrt{x})^2} dx$$

input

```
int(1/((x^2 - 1)^(1/2)*((x^2 - 1)^(1/2) + x^(1/2))^2), x)
```

output

```
int(1/((x^2 - 1)^(1/2)*((x^2 - 1)^(1/2) + x^(1/2))^2), x)
```

Reduce [F]

$$\int \frac{1}{\sqrt{-1+x^2}(\sqrt{x}+\sqrt{-1+x^2})^2} dx = \text{Too large to display}$$

input

```
int(1/(x^2-1)^(1/2)/(x^(1/2)+(x^2-1)^(1/2))^2, x)
```


output

```
(6*sqrt(sqrt(5) - 1)*sqrt(10)*atan((2*sqrt(x))/(sqrt(sqrt(5) - 1)*sqrt(2))
)*x**2 - 6*sqrt(sqrt(5) - 1)*sqrt(10)*atan((2*sqrt(x))/(sqrt(sqrt(5) - 1)*
sqrt(2)))*x - 6*sqrt(sqrt(5) - 1)*sqrt(10)*atan((2*sqrt(x))/(sqrt(sqrt(5)
- 1)*sqrt(2))) - 10*sqrt(sqrt(5) - 1)*sqrt(2)*atan((2*sqrt(x))/(sqrt(sqrt(
5) - 1)*sqrt(2)))*x**2 + 10*sqrt(sqrt(5) - 1)*sqrt(2)*atan((2*sqrt(x))/(sq
rt(sqrt(5) - 1)*sqrt(2)))*x + 10*sqrt(sqrt(5) - 1)*sqrt(2)*atan((2*sqrt(x)
)/(sqrt(sqrt(5) - 1)*sqrt(2))) - 100*sqrt(x**2 - 1)*x + 100*sqrt(x**2 - 1)
+ 3*sqrt(sqrt(5) + 1)*sqrt(10)*log(- sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2)
)*x**2 - 3*sqrt(sqrt(5) + 1)*sqrt(10)*log(- sqrt(sqrt(5) + 1) + sqrt(x)*s
qrt(2))*x - 3*sqrt(sqrt(5) + 1)*sqrt(10)*log(- sqrt(sqrt(5) + 1) + sqrt(x)
)*sqrt(2)) - 3*sqrt(sqrt(5) + 1)*sqrt(10)*log(sqrt(sqrt(5) + 1) + sqrt(x)*
sqrt(2))*x**2 + 3*sqrt(sqrt(5) + 1)*sqrt(10)*log(sqrt(sqrt(5) + 1) + sqrt(
x)*sqrt(2))*x + 3*sqrt(sqrt(5) + 1)*sqrt(10)*log(sqrt(sqrt(5) + 1) + sqrt(
x)*sqrt(2)) + 5*sqrt(sqrt(5) + 1)*sqrt(2)*log(- sqrt(sqrt(5) + 1) + sqrt(
x)*sqrt(2))*x**2 - 5*sqrt(sqrt(5) + 1)*sqrt(2)*log(- sqrt(sqrt(5) + 1) +
sqrt(x)*sqrt(2))*x - 5*sqrt(sqrt(5) + 1)*sqrt(2)*log(- sqrt(sqrt(5) + 1)
+ sqrt(x)*sqrt(2)) - 5*sqrt(sqrt(5) + 1)*sqrt(2)*log(sqrt(sqrt(5) + 1) + s
qrt(x)*sqrt(2))*x**2 + 5*sqrt(sqrt(5) + 1)*sqrt(2)*log(sqrt(sqrt(5) + 1) +
sqrt(x)*sqrt(2))*x + 5*sqrt(sqrt(5) + 1)*sqrt(2)*log(sqrt(sqrt(5) + 1) +
sqrt(x)*sqrt(2)) + 80*sqrt(x)*x - 40*sqrt(x) + 100*int(sqrt(x**2 - 1)/(...
```

3.10
$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx$$

Optimal result	129
Mathematica [A] (verified)	130
Rubi [B] (verified)	130
Maple [B] (verified)	132
Fricas [B] (verification not implemented)	133
Sympy [F(-1)]	134
Maxima [F]	134
Giac [B] (verification not implemented)	135
Mupad [F(-1)]	136
Reduce [F]	136

Optimal result

Integrand size = 39, antiderivative size = 220

$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx = \frac{2-4x}{5(\sqrt{x} + \sqrt{-1+x^2})} + \frac{1}{25} \sqrt{-110+50\sqrt{5}} \arctan\left(\frac{1}{2} \sqrt{2+2\sqrt{5}\sqrt{x}}\right) - \frac{1}{50} \sqrt{-110+50\sqrt{5}} \arctan\left(\frac{\sqrt{-2+2\sqrt{5}\sqrt{-1+x^2}}}{2-(1-\sqrt{5})x}\right) - \frac{1}{25} \sqrt{110+50\sqrt{5}} \operatorname{arctanh}\left(\frac{1}{2} \sqrt{-2+2\sqrt{5}\sqrt{x}}\right) - \frac{1}{50} \sqrt{110+50\sqrt{5}} \operatorname{arctanh}\left(\frac{\sqrt{2+2\sqrt{5}\sqrt{-1+x^2}}}{2-x-\sqrt{5}x}\right)$$

output

```
1/5*(2-4*x)/(x^(1/2)+(x^2-1)^(1/2))-1/50*arctan((x^2-1)^(1/2)*(-2+2*5^(1/2))^(1/2)/(2-x*(-5^(1/2)+1)))*(-110+50*5^(1/2))^(1/2)+1/25*arctan(1/2*x^(1/2)*(2+2*5^(1/2))^(1/2))*(-110+50*5^(1/2))^(1/2)-1/25*arctanh(1/2*x^(1/2)*(-2+2*5^(1/2))^(1/2))*(110+50*5^(1/2))^(1/2)-1/50*arctanh((x^2-1)^(1/2)*(2+2*5^(1/2))^(1/2)/(2-x-x*5^(1/2)))*(110+50*5^(1/2))^(1/2)
```

Mathematica [A] (verified)

Time = 6.32 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.89

$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx = \frac{1}{25} \left(-\frac{10(-1+2x)(-\sqrt{x} + \sqrt{-1+x^2})}{-1-x+x^2} \right. \\ \left. + \sqrt{-110+50\sqrt{5}} \arctan \left(\sqrt{\frac{1}{2}}(1+\sqrt{5})\sqrt{x} \right) \right. \\ \left. - \sqrt{-110+50\sqrt{5}} \arctan \left(\frac{\sqrt{-2+\sqrt{5}\sqrt{-1+x^2}}}{1+x} \right) \right. \\ \left. - \sqrt{110+50\sqrt{5}} \operatorname{arctanh} \left(\sqrt{\frac{1}{2}}(-1+\sqrt{5})\sqrt{x} \right) \right. \\ \left. + \sqrt{110+50\sqrt{5}} \operatorname{arctanh} \left(\frac{\sqrt{2+\sqrt{5}\sqrt{-1+x^2}}}{1+x} \right) \right)$$

input `Integrate[(Sqrt[x] - Sqrt[-1 + x^2])^2/((1 + x - x^2)^2*Sqrt[-1 + x^2]),x]`

output `((-10*(-1 + 2*x)*(-Sqrt[x] + Sqrt[-1 + x^2]))/(-1 - x + x^2) + Sqrt[-110 + 50*Sqrt[5]]*ArcTan[Sqrt[(1 + Sqrt[5])/2]*Sqrt[x]] - Sqrt[-110 + 50*Sqrt[5]]*ArcTan[(Sqrt[-2 + Sqrt[5]]*Sqrt[-1 + x^2])/(1 + x)] - Sqrt[110 + 50*Sqrt[5]]*ArcTanh[Sqrt[(-1 + Sqrt[5])/2]*Sqrt[x]] + Sqrt[110 + 50*Sqrt[5]]*ArcTanh[(Sqrt[2 + Sqrt[5]]*Sqrt[-1 + x^2])/(1 + x))]/25`

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 541 vs. 2(220) = 440.

Time = 1.13 (sec) , antiderivative size = 541, normalized size of antiderivative = 2.46, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.051$, Rules used = {7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{x} - \sqrt{x^2 - 1})^2}{(-x^2 + x + 1)^2 \sqrt{x^2 - 1}} dx$$

↓ 7293

$$\int \left(\frac{x^2}{\sqrt{x^2 - 1} (x^2 - x - 1)^2} + \frac{x}{\sqrt{x^2 - 1} (x^2 - x - 1)^2} - \frac{2\sqrt{x}}{(x^2 - x - 1)^2} - \frac{1}{\sqrt{x^2 - 1} (x^2 - x - 1)^2} \right) dx$$

↓ 2009

$$\begin{aligned} & \frac{1}{5} \sqrt{\frac{1}{5} (2 + 5\sqrt{5})} \arctan \left(\frac{2 - (1 - \sqrt{5})x}{\sqrt{2(\sqrt{5} - 1)} \sqrt{x^2 - 1}} \right) - \\ & \frac{1}{5} \sqrt{\frac{1}{5} (5\sqrt{5} - 2)} \arctan \left(\frac{2 - (1 - \sqrt{5})x}{\sqrt{2(\sqrt{5} - 1)} \sqrt{x^2 - 1}} \right) - \\ & \frac{1}{5} \sqrt{\frac{1}{10} (5\sqrt{5} - 11)} \arctan \left(\frac{2 - (1 - \sqrt{5})x}{\sqrt{2(\sqrt{5} - 1)} \sqrt{x^2 - 1}} \right) + \\ & \frac{1}{5} \sqrt{\frac{2}{5} (5\sqrt{5} - 11)} \arctan \left(\sqrt{\frac{2}{\sqrt{5} - 1}} \sqrt{x} \right) + \\ & \frac{1}{5} \sqrt{\frac{1}{10} (11 + 5\sqrt{5})} \operatorname{arctanh} \left(\frac{2 - (1 + \sqrt{5})x}{\sqrt{2(1 + \sqrt{5})} \sqrt{x^2 - 1}} \right) - \\ & \frac{1}{5} \sqrt{\frac{1}{5} (2 + 5\sqrt{5})} \operatorname{arctanh} \left(\frac{2 - (1 + \sqrt{5})x}{\sqrt{2(1 + \sqrt{5})} \sqrt{x^2 - 1}} \right) - \\ & \frac{1}{5} \sqrt{\frac{1}{5} (5\sqrt{5} - 2)} \operatorname{arctanh} \left(\frac{2 - (1 + \sqrt{5})x}{\sqrt{2(1 + \sqrt{5})} \sqrt{x^2 - 1}} \right) - \\ & \frac{1}{5} \sqrt{\frac{2}{5} (11 + 5\sqrt{5})} \operatorname{arctanh} \left(\sqrt{\frac{2}{1 + \sqrt{5}}} \sqrt{x} \right) - \frac{\sqrt{x^2 - 1}(1 - 2x)}{5(-x^2 + x + 1)} + \frac{2\sqrt{x}(1 - 2x)}{5(-x^2 + x + 1)} - \\ & \frac{(3 - x)\sqrt{x^2 - 1}}{5(-x^2 + x + 1)} + \frac{(x + 2)\sqrt{x^2 - 1}}{5(-x^2 + x + 1)} \end{aligned}$$

input

```
Int[(Sqrt[x] - Sqrt[-1 + x^2])^2/((1 + x - x^2)^2*Sqrt[-1 + x^2]),x]
```

output

$$\begin{aligned} & (2*(1 - 2*x)*\text{Sqrt}[x])/(5*(1 + x - x^2)) - ((1 - 2*x)*\text{Sqrt}[-1 + x^2])/(5*(1 \\ & + x - x^2)) - ((3 - x)*\text{Sqrt}[-1 + x^2])/(5*(1 + x - x^2)) + ((2 + x)*\text{Sqrt} \\ & [-1 + x^2])/(5*(1 + x - x^2)) + (\text{Sqrt}[(2*(-11 + 5*\text{Sqrt}[5]))/5]*\text{ArcTan}[\text{Sqrt} \\ & [2/(-1 + \text{Sqrt}[5])]*\text{Sqrt}[x]])/5 - (\text{Sqrt}[(-11 + 5*\text{Sqrt}[5])/10]*\text{ArcTan}[(2 - (1 \\ & - \text{Sqrt}[5])*x)/(\text{Sqrt}[2*(-1 + \text{Sqrt}[5])]*\text{Sqrt}[-1 + x^2])])/5 - (\text{Sqrt}[(-2 + 5 \\ & * \text{Sqrt}[5])/5]*\text{ArcTan}[(2 - (1 - \text{Sqrt}[5])*x)/(\text{Sqrt}[2*(-1 + \text{Sqrt}[5])]*\text{Sqrt}[-1 \\ & + x^2])])/5 + (\text{Sqrt}[(2 + 5*\text{Sqrt}[5])/5]*\text{ArcTan}[(2 - (1 - \text{Sqrt}[5])*x)/(\text{Sqrt} \\ & [2*(-1 + \text{Sqrt}[5])]*\text{Sqrt}[-1 + x^2])])/5 - (\text{Sqrt}[(2*(11 + 5*\text{Sqrt}[5]))/5]*\text{ArcT} \\ & \text{anh}[\text{Sqrt}[2/(1 + \text{Sqrt}[5])]*\text{Sqrt}[x]])/5 - (\text{Sqrt}[(-2 + 5*\text{Sqrt}[5])/5]*\text{ArcTanh} \\ & [(2 - (1 + \text{Sqrt}[5])*x)/(\text{Sqrt}[2*(1 + \text{Sqrt}[5])]*\text{Sqrt}[-1 + x^2])])/5 - (\text{Sqrt}[(\\ & 2 + 5*\text{Sqrt}[5])/5]*\text{ArcTanh}[(2 - (1 + \text{Sqrt}[5])*x)/(\text{Sqrt}[2*(1 + \text{Sqrt}[5])]*\text{Sqr} \\ & \text{t}[-1 + x^2])])/5 + (\text{Sqrt}[(11 + 5*\text{Sqrt}[5])/10]*\text{ArcTanh}[(2 - (1 + \text{Sqrt}[5])*x \\ &)/(\text{Sqrt}[2*(1 + \text{Sqrt}[5])]*\text{Sqrt}[-1 + x^2])])/5 \end{aligned}$$
Defintions of rubi rules used

rule 2009

$$\text{Int}[u_, x_Symbol] \text{ :> Simp}[\text{IntSum}[u, x], x] \text{ /; SumQ}[u]$$

rule 7293

$$\text{Int}[u_, x_Symbol] \text{ :> With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] \text{ /; SumQ}[v]$$

$$\text{]}$$
Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1635 vs. $2(158) = 316$.

Time = 0.81 (sec) , antiderivative size = 1636, normalized size of antiderivative = 7.44

method	result	size
default	Expression too large to display	1636

input

$$\text{int}((x^{(1/2)} - (x^2 - 1)^{(1/2)})^2 / (-x^2 + x + 1)^2 / (x^2 - 1)^{(1/2)}, x, \text{method} = \text{_RETURNV}$$

$$\text{ERBOSE})$$

output

```

2/25*5^(1/2)/(2+2*5^(1/2))^(1/2)*arctanh(2*(1+5^(1/2)+(5^(1/2)+1)*(x-1/2*5
^(1/2)-1/2))/(2+2*5^(1/2))^(1/2)/(4*(x-1/2*5^(1/2)-1/2)^2+4*(5^(1/2)+1)*(x
-1/2*5^(1/2)-1/2)+2+2*5^(1/2))^(1/2))+2/25*5^(1/2)/(-2+2*5^(1/2))^(1/2)*ar
ctan(2*(1-5^(1/2)+(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2))/(-2+2*5^(1/2))^(1/2)/(
4*(x+1/2*5^(1/2)-1/2)^2+4*(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2)+2-2*5^(1/2))^(1
/2))+2/5-2/5*5^(1/2))*(-1/4/(1/2-1/2*5^(1/2)))/(x+1/2*5^(1/2)-1/2)*((x+1/2
*5^(1/2)-1/2)^2+(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2)+1/2-1/2*5^(1/2))^(1/2)-1/
4*(-5^(1/2)+1)/(1/2-1/2*5^(1/2))/(-2+2*5^(1/2))^(1/2)*arctan(2*(1-5^(1/2)+
(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2))/(-2+2*5^(1/2))^(1/2)/(4*(x+1/2*5^(1/2)-1
/2)^2+4*(-5^(1/2)+1)*(x+1/2*5^(1/2)-1/2)+2-2*5^(1/2))^(1/2)))+2/5+2/5*5^(
1/2))*(-1/4/(1/2+1/2*5^(1/2)))/(x-1/2*5^(1/2)-1/2)*((x-1/2*5^(1/2)-1/2)^2+(
5^(1/2)+1)*(x-1/2*5^(1/2)-1/2)+1/2+1/2*5^(1/2))^(1/2)+1/4*(5^(1/2)+1)/(1/2
+1/2*5^(1/2))/(2+2*5^(1/2))^(1/2)*arctanh(2*(1+5^(1/2)+(5^(1/2)+1)*(x-1/2*
5^(1/2)-1/2))/(2+2*5^(1/2))^(1/2)/(4*(x-1/2*5^(1/2)-1/2)^2+4*(5^(1/2)+1)*(
x-1/2*5^(1/2)-1/2)+2+2*5^(1/2))^(1/2))+2/5*x^(1/2)/(x-1/2*5^(1/2)-1/2)-8/
25*(5/2+5^(1/2))/(2+2*5^(1/2))^(1/2)*arctanh(2*x^(1/2)/(2+2*5^(1/2))^(1/2)
)+2/5*x^(1/2)/(x+1/2*5^(1/2)-1/2)-8/25*(-5/2+5^(1/2))/(-2+2*5^(1/2))^(1/2)
*arctan(2*x^(1/2)/(-2+2*5^(1/2))^(1/2))-1/5/(1/2+1/2*5^(1/2))/(x-1/2*5^(1/
2)-1/2)*((x-1/2*5^(1/2)-1/2)^2+(5^(1/2)+1)*(x-1/2*5^(1/2)-1/2)+1/2+1/2*5^(
1/2))^(3/2)+1/10*(5^(1/2)+1)/(1/2+1/2*5^(1/2))*1/2*(4*(x-1/2*5^(1/2)-1...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 341 vs. $2(153) = 306$.

Time = 0.08 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.55

$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx$$

$$= \frac{2(x^2 - x - 1)\sqrt{\frac{1}{2}\sqrt{5} - \frac{11}{10}} \arctan\left(\sqrt{x}(2\sqrt{5} + 5)\sqrt{\frac{1}{2}\sqrt{5} - \frac{11}{10}}\right) - 2(x^2 - x - 1)\sqrt{\frac{1}{2}\sqrt{5} - \frac{11}{10}} \arctan\left(\dots\right)}{\dots}$$

input

```

integrate((x^(1/2)-(x^2-1)^(1/2))^2/(-x^2+x+1)^2/(x^2-1)^(1/2),x, algorithm
m="fricas")

```

output

```
1/5*(2*(x^2 - x - 1)*sqrt(1/2*sqrt(5) - 11/10)*arctan(sqrt(x)*(2*sqrt(5) +
5)*sqrt(1/2*sqrt(5) - 11/10)) - 2*(x^2 - x - 1)*sqrt(1/2*sqrt(5) - 11/10)
*arctan(-1/2*(sqrt(5)*(4*x + 3) - 2*sqrt(x^2 - 1)*(2*sqrt(5) + 5) + 10*x +
5)*sqrt(1/2*sqrt(5) - 11/10)) + (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11/10)*l
og((3*sqrt(5) - 5)*sqrt(1/2*sqrt(5) + 11/10) - 2*x + sqrt(5) + 2*sqrt(x^2
- 1) + 1) - (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11/10)*log((3*sqrt(5) - 5)*sq
rt(1/2*sqrt(5) + 11/10) + 2*sqrt(x)) - (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11
/10)*log(-(3*sqrt(5) - 5)*sqrt(1/2*sqrt(5) + 11/10) - 2*x + sqrt(5) + 2*sq
rt(x^2 - 1) + 1) + (x^2 - x - 1)*sqrt(1/2*sqrt(5) + 11/10)*log(-(3*sqrt(5)
- 5)*sqrt(1/2*sqrt(5) + 11/10) + 2*sqrt(x)) - 4*x^2 - 2*sqrt(x^2 - 1)*(2*
x - 1) + 2*(2*x - 1)*sqrt(x) + 4*x + 4)/(x^2 - x - 1)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx = \text{Timed out}$$

input

```
integrate((x**(1/2)-(x**2-1)**(1/2))**2/(-x**2+x+1)**2/(x**2-1)**(1/2),x)
```

output

Timed out

Maxima [F]

$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx = \int \frac{(\sqrt{x^2-1} - \sqrt{x})^2}{(x^2-x-1)^2 \sqrt{x^2-1}} dx$$

input

```
integrate((x^(1/2)-(x^2-1)^(1/2))^2/(-x^2+x+1)^2/(x^2-1)^(1/2),x, algorith
m="maxima")
```

output

```
-2/5*(x^(5/2) - 3*x^(3/2))/(x^2 - x - 1) + integrate(1/5*(x^(3/2) + sqrt(x
))/(x^2 - x - 1), x) + integrate((x^2 + x - 1)/((x^4 - 2*x^3 - x^2 + 2*x +
1)*sqrt(x + 1)*sqrt(x - 1)), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 358 vs. $2(153) = 306$.

Time = 1.42 (sec) , antiderivative size = 358, normalized size of antiderivative = 1.63

$$\begin{aligned}
\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx &= \frac{2}{5} \sqrt{\frac{1}{2} \sqrt{5} - \frac{11}{10}} \arctan \left(\frac{2x + \sqrt{5} - 2\sqrt{x^2-1} - 1}{\sqrt{2}\sqrt{5} - 2} \right) \\
&+ \frac{1}{25} \sqrt{50\sqrt{5} - 110} \arctan \left(\frac{\sqrt{x}}{\sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}}} \right) \\
&- \frac{1}{50} \sqrt{50\sqrt{5} + 110} \log \left(\sqrt{x} + \sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}} \right) \\
&- \frac{1}{5} \sqrt{\frac{1}{2}\sqrt{5} + \frac{11}{10}} \log \left(\left| -520x - 78\sqrt{5}\sqrt{50\sqrt{5} + 110} + 260\sqrt{5} + 520\sqrt{x^2-1} + 130\sqrt{50\sqrt{5} + 110} \right| \right) \\
&+ \frac{1}{5} \sqrt{\frac{1}{2}\sqrt{5} + \frac{11}{10}} \log \left(\left| -1040x + 156\sqrt{5}\sqrt{50\sqrt{5} + 110} + 520\sqrt{5} + 1040\sqrt{x^2-1} - 260\sqrt{50\sqrt{5} + 110} \right| \right) \\
&+ \frac{1}{50} \sqrt{50\sqrt{5} + 110} \log \left(\left| \sqrt{x} - \sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}} \right| \right) \\
&+ \frac{4 \left((x - \sqrt{x^2-1})^3 + 2(x - \sqrt{x^2-1})^2 + 3x - 3\sqrt{x^2-1} - 2 \right)}{5 \left((x - \sqrt{x^2-1})^4 - 2(x - \sqrt{x^2-1})^3 - 2(x - \sqrt{x^2-1})^2 - 2x + 2\sqrt{x^2-1} + 1 \right)} \\
&+ \frac{2 \left(2x^{\frac{3}{2}} - \sqrt{x} \right)}{5(x^2 - x - 1)}
\end{aligned}$$

input

```
integrate((x^(1/2)-(x^2-1)^(1/2))^2/(-x^2+x+1)^2/(x^2-1)^(1/2),x, algorithm
m="giac")
```


output

```
2/5*sqrt(1/2*sqrt(5) - 11/10)*arctan((2*x + sqrt(5) - 2*sqrt(x^2 - 1) - 1)
/sqrt(2*sqrt(5) - 2)) + 1/25*sqrt(50*sqrt(5) - 110)*arctan(sqrt(x)/sqrt(1/
2*sqrt(5) - 1/2)) - 1/50*sqrt(50*sqrt(5) + 110)*log(sqrt(x) + sqrt(1/2*sq
rt(5) + 1/2)) - 1/5*sqrt(1/2*sqrt(5) + 11/10)*log(abs(-520*x - 78*sqrt(5)*s
qrt(50*sqrt(5) + 110) + 260*sqrt(5) + 520*sqrt(x^2 - 1) + 130*sqrt(50*sqrt
(5) + 110) + 260)) + 1/5*sqrt(1/2*sqrt(5) + 11/10)*log(abs(-1040*x + 156*s
qrt(5)*sqrt(50*sqrt(5) + 110) + 520*sqrt(5) + 1040*sqrt(x^2 - 1) - 260*sq
rt(50*sqrt(5) + 110) + 520)) + 1/50*sqrt(50*sqrt(5) + 110)*log(abs(sqrt(x)
- sqrt(1/2*sqrt(5) + 1/2))) + 4/5*((x - sqrt(x^2 - 1))^3 + 2*(x - sqrt(x^2
- 1))^2 + 3*x - 3*sqrt(x^2 - 1) - 2)/((x - sqrt(x^2 - 1))^4 - 2*(x - sqrt
(x^2 - 1))^3 - 2*(x - sqrt(x^2 - 1))^2 - 2*x + 2*sqrt(x^2 - 1) + 1) + 2/5*
(2*x^(3/2) - sqrt(x))/(x^2 - x - 1)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx = \int \frac{(\sqrt{x^2-1} - \sqrt{x})^2}{\sqrt{x^2-1} (-x^2+x+1)^2} dx$$

input

```
int(((x^2 - 1)^(1/2) - x^(1/2))^2/((x^2 - 1)^(1/2)*(x - x^2 + 1)^2), x)
```

output

```
int(((x^2 - 1)^(1/2) - x^(1/2))^2/((x^2 - 1)^(1/2)*(x - x^2 + 1)^2), x)
```

Reduce [F]

$$\int \frac{(\sqrt{x} - \sqrt{-1+x^2})^2}{(1+x-x^2)^2 \sqrt{-1+x^2}} dx = \text{Too large to display}$$

input

```
int((x^(1/2)-(x^2-1)^(1/2))^2/(-x^2+x+1)^2/(x^2-1)^(1/2), x)
```

output

```
(6*sqrt(sqrt(5) - 1)*sqrt(10)*atan((2*sqrt(x))/(sqrt(sqrt(5) - 1)*sqrt(2))
)*x**2 - 6*sqrt(sqrt(5) - 1)*sqrt(10)*atan((2*sqrt(x))/(sqrt(sqrt(5) - 1)*
sqrt(2)))*x - 6*sqrt(sqrt(5) - 1)*sqrt(10)*atan((2*sqrt(x))/(sqrt(sqrt(5)
- 1)*sqrt(2))) - 10*sqrt(sqrt(5) - 1)*sqrt(2)*atan((2*sqrt(x))/(sqrt(sqrt(
5) - 1)*sqrt(2)))*x**2 + 10*sqrt(sqrt(5) - 1)*sqrt(2)*atan((2*sqrt(x))/(sq
rt(sqrt(5) - 1)*sqrt(2)))*x + 10*sqrt(sqrt(5) - 1)*sqrt(2)*atan((2*sqrt(x)
)/(sqrt(sqrt(5) - 1)*sqrt(2))) + 3*sqrt(sqrt(5) + 1)*sqrt(10)*log( - sqrt(
sqrt(5) + 1) + sqrt(x)*sqrt(2))*x**2 - 3*sqrt(sqrt(5) + 1)*sqrt(10)*log( -
sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2))*x - 3*sqrt(sqrt(5) + 1)*sqrt(10)*log
( - sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2)) - 3*sqrt(sqrt(5) + 1)*sqrt(10)*lo
g(sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2))*x**2 + 3*sqrt(sqrt(5) + 1)*sqrt(10)
*log(sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2))*x + 3*sqrt(sqrt(5) + 1)*sqrt(10)
*log(sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2)) + 5*sqrt(sqrt(5) + 1)*sqrt(2)*lo
g( - sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2))*x**2 - 5*sqrt(sqrt(5) + 1)*sqrt(
2)*log( - sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2))*x - 5*sqrt(sqrt(5) + 1)*sqr
t(2)*log(sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2))*x**2 + 5*sqrt(sqrt(5) + 1)*sqr
t(2)*log(sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2))*x + 5*sqrt(sqrt(5) + 1)*sqr
t(2)*log(sqrt(sqrt(5) + 1) + sqrt(x)*sqrt(2)) + 80*sqrt(x)*x - 40*sqrt(x)
+ 100*int(x**2/(sqrt(x**2 - 1))*x**4 - 2*sqrt(x**2 - 1)*x**3 - sqrt(x**...
```

3.11 $\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx$

Optimal result	138
Mathematica [A] (verified)	139
Rubi [A] (verified)	139
Maple [A] (verified)	140
Fricas [A] (verification not implemented)	140
Sympy [F(-2)]	141
Maxima [F(-2)]	141
Giac [B] (verification not implemented)	142
Mupad [F(-1)]	143
Reduce [F]	143

Optimal result

Integrand size = 45, antiderivative size = 138

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx$$

$$= -\frac{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{-i+x^2}}{\sqrt{2}(1+x)} - \frac{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{i+x^2}}{\sqrt{2}(1+x)}$$

$$+ \frac{\operatorname{arctanh}\left(\frac{i+x}{\sqrt{1-i}\sqrt{-i+x^2}}\right)}{(1-i)^{3/2}\sqrt{2}} - \frac{\operatorname{arctanh}\left(\frac{i-x}{\sqrt{1+i}\sqrt{i+x^2}}\right)}{(1+i)^{3/2}\sqrt{2}}$$

output

```
1/2*arctanh((I+x)/(1-I)^(1/2)/(-I+x^2)^(1/2))/(1-I)^(3/2)*2^(1/2)-1/2*arctanh((I-x)/(1+I)^(1/2)/(I+x^2)^(1/2))/(1+I)^(3/2)*2^(1/2)-(1/4+1/4*I)*(-I+x^2)^(1/2)/(1+x)*2^(1/2)+(-1/4+1/4*I)*(I+x^2)^(1/2)/(1+x)*2^(1/2)
```

Mathematica [A] (verified)

Time = 3.63 (sec) , antiderivative size = 126, normalized size of antiderivative = 0.91

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx =$$

$$\frac{\left(\frac{1}{2} - \frac{i}{2}\right) \left(i\sqrt{-i+x^2} + \sqrt{i+x^2} + \frac{2(1+x) \arctan\left(\sqrt{-\frac{1}{2}-\frac{i}{2}}(1+x-\sqrt{-i+x^2})\right)}{\sqrt{1-i}} \right) + (1+i)^{3/2}(1+x) \arctan\left(\sqrt{-\frac{1}{2}-\frac{i}{2}}(1+x-\sqrt{-i+x^2})\right)}{\sqrt{2}(1+x)}$$

input

```
Integrate[1/(Sqrt[2]*(1+x)^2*Sqrt[-I+x^2]) + 1/(Sqrt[2]*(1+x)^2*Sqrt[I+x^2]),x]
```

output

```
((-1/2 + I/2)*(I*Sqrt[-I + x^2] + Sqrt[I + x^2] + (2*(1 + x)*ArcTan[Sqrt[-1/2 - I/2]*(1 + x - Sqrt[-I + x^2])]))/Sqrt[1 - I] + (1 + I)^(3/2)*(1 + x)*ArcTan[Sqrt[-1/2 + I/2]*(1 + x - Sqrt[I + x^2])])/(Sqrt[2]*(1 + x))
```

Rubi [A] (verified)

Time = 0.25 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.022$, Rules used = {2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \left(\frac{1}{\sqrt{2}(x+1)^2\sqrt{x^2+i}} + \frac{1}{\sqrt{2}(x+1)^2\sqrt{x^2-i}} \right) dx$$

$$\downarrow \text{2009}$$

$$\frac{\operatorname{arctanh}\left(\frac{x+i}{\sqrt{1-i}\sqrt{x^2-i}}\right)}{(1-i)^{3/2}\sqrt{2}} - \frac{\operatorname{arctanh}\left(\frac{-x+i}{\sqrt{1+i}\sqrt{x^2+i}}\right)}{(1+i)^{3/2}\sqrt{2}} - \frac{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{x^2-i}}{\sqrt{2}(x+1)} - \frac{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{x^2+i}}{\sqrt{2}(x+1)}$$

input

```
Int[1/(Sqrt[2]*(1+x)^2*Sqrt[-I+x^2]) + 1/(Sqrt[2]*(1+x)^2*Sqrt[I+x^2]),x]
```

```
output ((-1/2 - I/2)*Sqrt[-I + x^2])/(Sqrt[2]*(1 + x)) - ((1/2 - I/2)*Sqrt[I + x^2])/(Sqrt[2]*(1 + x)) + ArcTanh[(I + x)/(Sqrt[1 - I]*Sqrt[-I + x^2])]/((1 - I)^(3/2)*Sqrt[2]) - ArcTanh[(I - x)/(Sqrt[1 + I]*Sqrt[I + x^2])]/((1 + I)^(3/2)*Sqrt[2])
```

Defintions of rubi rules used

```
rule 2009 Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.06

method	result
default	$\frac{\sqrt{2} \left(\frac{(-\frac{1}{2} - \frac{i}{2})\sqrt{(1+x)^2 - 2x - 1 - i}}{1+x} + \frac{(-\frac{1}{2} - \frac{i}{2}) \ln\left(\frac{-2i - 2x + 2\sqrt{1-i}\sqrt{(1+x)^2 - 2x - 1 - i}}{1+x}\right)}{\sqrt{1-i}} \right)}{2} + \frac{\sqrt{2} \left(\frac{(-\frac{1}{2} + \frac{i}{2})\sqrt{(1+x)^2 - 2x - 1 + i}}{1+x} + \frac{(-\frac{1}{2} + \frac{i}{2}) \ln\left(\frac{-2i - 2x + 2\sqrt{1+i}\sqrt{(1+x)^2 - 2x - 1 + i}}{1+x}\right)}{\sqrt{1+i}} \right)}{2}$

```
input int(1/2/(1+x)^2*2^(1/2)/(x^2-I)^(1/2)+1/2/(1+x)^2*2^(1/2)/(x^2+I)^(1/2),x, method=_RETURNVERBOSE)
```

```
output 1/2*2^(1/2)*((-1/2-1/2*I)/(1+x)*((1+x)^2-2*x-1-I)^(1/2)-(1/2+1/2*I)/(1-I)^(1/2)*ln((-2*I-2*x+2*(1-I)^(1/2)*((1+x)^2-2*x-1-I)^(1/2))/(1+x)))+1/2*2^(1/2)*((-1/2+1/2*I)/(1+x)*((1+x)^2-2*x-1+I)^(1/2)+(-1/2+1/2*I)/(1+I)^(1/2)*ln((2*I-2*x+2*(1+I)^(1/2)*((1+x)^2-2*x-1+I)^(1/2))/(1+x))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.17

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx$$

$$= \frac{\sqrt{-\frac{1}{2}i + \frac{1}{2}}(-i-1)x - i + 1 \log\left(\sqrt{2}\sqrt{-\frac{1}{2}i + \frac{1}{2}} - x + \sqrt{x^2 - i} - 1\right) + \sqrt{-\frac{1}{2}i + \frac{1}{2}}((i-1)x + i - 1)}{2}$$

input `integrate(1/2/(1+x)^2*2^(1/2)/(-I+x^2)^(1/2)+1/2/(1+x)^2*2^(1/2)/(I+x^2)^(1/2),x, algorithm="fricas")`

output `(sqrt(-1/2*I + 1/2)*(-I - 1)*x - I + 1)*log(sqrt(2)*sqrt(-1/2*I + 1/2) - x + sqrt(x^2 - I) - 1) + sqrt(-1/2*I + 1/2)*((I - 1)*x + I - 1)*log(-sqrt(2)*sqrt(-1/2*I + 1/2) - x + sqrt(x^2 - I) - 1) + sqrt(-1/2*I - 1/2)*(-I + 1)*x - I - 1)*log(I*sqrt(2)*sqrt(-1/2*I - 1/2) - x + sqrt(x^2 + I) - 1) + sqrt(-1/2*I - 1/2)*((I + 1)*x + I + 1)*log(-I*sqrt(2)*sqrt(-1/2*I - 1/2) - x + sqrt(x^2 + I) - 1) + sqrt(2)*(-I + 1)*x - I - 1) - sqrt(2)*sqrt(x^2 + I) - I*sqrt(2)*sqrt(x^2 - I))/((2*I + 2)*x + 2*I + 2)`

Sympy [F(-2)]

Exception generated.

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx = \text{Exception raised: TypeError}$$

input `integrate(1/2/(1+x)**2*2**(1/2)/(-I+x**2)**(1/2)+1/2/(1+x)**2*2**(1/2)/(I+x**2)**(1/2),x)`

output `Exception raised: TypeError >> Invalid comparison of non-real I`

Maxima [F(-2)]

Exception generated.

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx = \text{Exception raised: RuntimeError}$$

input `integrate(1/2/(1+x)^2*2^(1/2)/(-I+x^2)^(1/2)+1/2/(1+x)^2*2^(1/2)/(I+x^2)^(1/2),x, algorithm="maxima")`

output `Exception raised: RuntimeError >> ECL says: In function CAR, the value of the first argument is 1which is not of the expected type LIST`

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 547 vs. $2(82) = 164$.

Time = 0.14 (sec) , antiderivative size = 547, normalized size of antiderivative = 3.96

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx$$

$$= \sqrt{2} \left(\frac{-(i-1)\sqrt{2x^2+2\sqrt{x^4+1}}\left(\frac{i}{x^2+\sqrt{x^4+1}}+1\right) + (2i-2)x + 2i + 2}{\left(\sqrt{2x^2+2\sqrt{x^4+1}}\left(\frac{i}{x^2+\sqrt{x^4+1}}+1\right) - 2x\right)^2 - 4\sqrt{2x^2+2\sqrt{x^4+1}}\left(\frac{i}{x^2+\sqrt{x^4+1}}+1\right) + 8x - 4i} \right.$$

$$\left. + \sqrt{2} \left(\frac{(i+1)\sqrt{2x^2+2\sqrt{x^4+1}}\left(-\frac{i}{x^2+\sqrt{x^4+1}}+1\right) - (2i+2)x - 2i + 2}{\left(\sqrt{2x^2+2\sqrt{x^4+1}}\left(-\frac{i}{x^2+\sqrt{x^4+1}}+1\right) - 2x\right)^2 - 4\sqrt{2x^2+2\sqrt{x^4+1}}\left(-\frac{i}{x^2+\sqrt{x^4+1}}+1\right) + 8x + 4i} \right) \right)$$

input

```
integrate(1/2/(1+x)^2*2^(1/2)/(-I+x^2)^(1/2)+1/2/(1+x)^2*2^(1/2)/(I+x^2)^(1/2),x, algorithm="giac")
```

output

```
sqrt(2)*((-I - 1)*sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(I/(x^2 + sqrt(x^4 + 1)) + 1) + (2*I - 2)*x + 2*I + 2)/((sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(I/(x^2 + sqrt(x^4 + 1)) + 1) - 2*x)^2 - 4*sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(I/(x^2 + sqrt(x^4 + 1)) + 1) + 8*x - 4*I) - (I - 1)*arctan((sqrt(2)*(sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(I/(x^2 + sqrt(x^4 + 1)) + 1) - 2*x) - sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(I/(x^2 + sqrt(x^4 + 1)) + 1) + 2*x - 2*sqrt(2) + 2)/(sqrt(2)*sqrt(2*sqrt(2) - 2) - (I + 1)*sqrt(2*sqrt(2) - 2)))/(sqrt(2*sqrt(2) - 2)*(-I/(sqrt(2) - 1) + 1))) + sqrt(2)*(((I + 1)*sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(-I/(x^2 + sqrt(x^4 + 1)) + 1) - (2*I + 2)*x - 2*I + 2)/((sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(-I/(x^2 + sqrt(x^4 + 1)) + 1) - 2*x)^2 - 4*sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(-I/(x^2 + sqrt(x^4 + 1)) + 1) + 8*x + 4*I) + (I + 1)*arctan((sqrt(2)*(sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(-I/(x^2 + sqrt(x^4 + 1)) + 1) - 2*x) - sqrt(2*x^2 + 2*sqrt(x^4 + 1))*(-I/(x^2 + sqrt(x^4 + 1)) + 1) + 2*x - 2*sqrt(2) + 2)/(sqrt(2)*sqrt(2*sqrt(2) - 2) + (I - 1)*sqrt(2*sqrt(2) - 2)))/(sqrt(2*sqrt(2) - 2)*(I/(sqrt(2) - 1) + 1)))
```

Mupad [F(-1)]

Timed out.

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx$$

$$= \int \frac{\sqrt{2}}{2\sqrt{x^2-i}(x+1)^2} + \frac{\sqrt{2}}{2\sqrt{x^2+1i}(x+1)^2} dx$$

input `int(2^(1/2)/(2*(x^2 - 1i)^(1/2)*(x + 1)^2) + 2^(1/2)/(2*(x^2 + 1i)^(1/2)*(x + 1)^2), x)`

output `int(2^(1/2)/(2*(x^2 - 1i)^(1/2)*(x + 1)^2) + 2^(1/2)/(2*(x^2 + 1i)^(1/2)*(x + 1)^2), x)`

Reduce [F]

$$\int \left(\frac{1}{\sqrt{2}(1+x)^2\sqrt{-i+x^2}} + \frac{1}{\sqrt{2}(1+x)^2\sqrt{i+x^2}} \right) dx$$

$$= \int \left(\frac{\sqrt{2}}{2(x+1)^2\sqrt{x^2-i}} + \frac{\sqrt{2}}{2(x+1)^2\sqrt{x^2+i}} \right) dx$$

input `int(1/2/(1+x)^2*2^(1/2)/(-I+x^2)^(1/2)+1/2/(1+x)^2*2^(1/2)/(I+x^2)^(1/2), x)`

output `int(1/2/(1+x)^2*2^(1/2)/(-I+x^2)^(1/2)+1/2/(1+x)^2*2^(1/2)/(I+x^2)^(1/2), x)`

3.12 $\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{(1+x)^2 \sqrt{1+x^4}} dx$

Optimal result	144
Mathematica [B] (verified)	145
Rubi [A] (verified)	146
Maple [F]	147
Fricas [B] (verification not implemented)	148
Sympy [F]	148
Maxima [F]	149
Giac [F]	149
Mupad [F(-1)]	149
Reduce [F]	150

Optimal result

Integrand size = 32, antiderivative size = 125

$$\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{(1+x)^2 \sqrt{1+x^4}} dx = -\frac{\sqrt{1-ix^2}}{2(1+x)} - \frac{\sqrt{1+ix^2}}{2(1+x)} - \frac{1}{4}(1-i)^{3/2} \operatorname{arctanh}\left(\frac{1+ix}{\sqrt{1-i}\sqrt{1-ix^2}}\right) - \frac{1}{4}(1+i)^{3/2} \operatorname{arctanh}\left(\frac{1-ix}{\sqrt{1+i}\sqrt{1+ix^2}}\right)$$

output

```
-1/4*(1-I)^(3/2)*arctanh((1+I*x)/(1-I)^(1/2)/(1-I*x^2)^(1/2))-1/4*(1+I)^(3/2)*arctanh((1-I*x)/(1+I)^(1/2)/(1+I*x^2)^(1/2))-1/2*(1-I*x^2)^(1/2)/(1+x)-1/2*(1+I*x^2)^(1/2)/(1+x)
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 272 vs. $2(125) = 250$.

Time = 2.34 (sec) , antiderivative size = 272, normalized size of antiderivative = 2.18

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)^2 \sqrt{1 + x^4}} dx = \frac{1}{2} \left(\frac{-1 - 2x^4 - \sqrt{1 + x^4} - x^2(1 + 2\sqrt{1 + x^4})}{(1 + x)(x^2 + \sqrt{1 + x^4})^{3/2}} \right. \\ \left. + \frac{\arctan\left(\sqrt{1 + \sqrt{2}}\sqrt{x^2 + \sqrt{1 + x^4}}\right)}{\sqrt{-1 + \sqrt{2}}} \right. \\ \left. - \sqrt{1 + \sqrt{2}} \arctan\left(\frac{\sqrt{2}(-1 + \sqrt{2})x\sqrt{x^2 + \sqrt{1 + x^4}}}{1 + x^2 + \sqrt{1 + x^4}}\right) \right. \\ \left. - \frac{\operatorname{arctanh}\left(\sqrt{-1 + \sqrt{2}}\sqrt{x^2 + \sqrt{1 + x^4}}\right)}{\sqrt{1 + \sqrt{2}}} \right. \\ \left. + \sqrt{-1 + \sqrt{2}} \operatorname{arctanh}\left(\frac{\sqrt{2}(1 + \sqrt{2})x\sqrt{x^2 + \sqrt{1 + x^4}}}{1 + x^2 + \sqrt{1 + x^4}}\right) \right)$$

input

```
Integrate[Sqrt[x^2 + Sqrt[1 + x^4]]/((1 + x)^2*Sqrt[1 + x^4]),x]
```

output

```
((-1 - 2*x^4 - Sqrt[1 + x^4] - x^2*(1 + 2*Sqrt[1 + x^4]))/((1 + x)*(x^2 + Sqrt[1 + x^4])^(3/2)) + ArcTan[Sqrt[1 + Sqrt[2]]*Sqrt[x^2 + Sqrt[1 + x^4]]/Sqrt[-1 + Sqrt[2]] - Sqrt[1 + Sqrt[2]]*ArcTan[(Sqrt[2*(-1 + Sqrt[2]])*x*Sqrt[x^2 + Sqrt[1 + x^4]])/(1 + x^2 + Sqrt[1 + x^4])] - ArcTanh[Sqrt[-1 + Sqrt[2]]*Sqrt[x^2 + Sqrt[1 + x^4]]]/Sqrt[1 + Sqrt[2]] + Sqrt[-1 + Sqrt[2]]*ArcTanh[(Sqrt[2*(1 + Sqrt[2]])*x*Sqrt[x^2 + Sqrt[1 + x^4]])/(1 + x^2 + Sqrt[1 + x^4])])/2
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.21, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2558, 491, 488, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{\sqrt{x^4+1}+x^2}}{(x+1)^2\sqrt{x^4+1}} dx \\
 & \quad \downarrow \text{2558} \\
 & \left(\frac{1}{2} - \frac{i}{2}\right) \int \frac{1}{(x+1)^2\sqrt{1-ix^2}} dx + \left(\frac{1}{2} + \frac{i}{2}\right) \int \frac{1}{(x+1)^2\sqrt{ix^2+1}} dx \\
 & \quad \downarrow \text{491} \\
 & \left(\frac{1}{2} - \frac{i}{2}\right) \left(\left(\frac{1}{2} - \frac{i}{2}\right) \int \frac{1}{(x+1)\sqrt{1-ix^2}} dx - \frac{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{1-ix^2}}{x+1} \right) + \\
 & \left(\frac{1}{2} + \frac{i}{2}\right) \left(\left(\frac{1}{2} + \frac{i}{2}\right) \int \frac{1}{(x+1)\sqrt{ix^2+1}} dx - \frac{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{1+ix^2}}{x+1} \right) \\
 & \quad \downarrow \text{488} \\
 & \left(\frac{1}{2} - \frac{i}{2}\right) \left(\left(-\frac{1}{2} + \frac{i}{2}\right) \int \frac{1}{(1-i) - \frac{(ix+1)^2}{1-ix^2}} d \frac{ix+1}{\sqrt{1-ix^2}} - \frac{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{1-ix^2}}{x+1} \right) + \\
 & \left(\frac{1}{2} + \frac{i}{2}\right) \left(\left(-\frac{1}{2} - \frac{i}{2}\right) \int \frac{1}{(1+i) - \frac{(1-ix)^2}{ix^2+1}} d \frac{1-ix}{\sqrt{ix^2+1}} - \frac{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{1+ix^2}}{x+1} \right) \\
 & \quad \downarrow \text{219} \\
 & \left(\frac{1}{2} - \frac{i}{2}\right) \left(-\frac{1}{2}\sqrt{1-i} \operatorname{arctanh}\left(\frac{1+ix}{\sqrt{1-i}\sqrt{1-ix^2}}\right) - \frac{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{1-ix^2}}{x+1} \right) + \\
 & \left(\frac{1}{2} + \frac{i}{2}\right) \left(-\frac{1}{2}\sqrt{1+i} \operatorname{arctanh}\left(\frac{1-ix}{\sqrt{1+i}\sqrt{1+ix^2}}\right) - \frac{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{1+ix^2}}{x+1} \right)
 \end{aligned}$$

input

```
Int[Sqrt[x^2 + Sqrt[1 + x^4]]/((1 + x)^2*Sqrt[1 + x^4]),x]
```

output

```
(1/2 - I/2)*((( -1/2 - I/2)*Sqrt[1 - I*x^2])/(1 + x) - (Sqrt[1 - I]*ArcTanh
[(1 + I*x)/(Sqrt[1 - I]*Sqrt[1 - I*x^2])])/2) + (1/2 + I/2)*((( -1/2 + I/2)
*Sqrt[1 + I*x^2])/(1 + x) - (Sqrt[1 + I]*ArcTanh[(1 - I*x)/(Sqrt[1 + I]*Sqr
t[1 + I*x^2])])/2)
```

Defintions of rubi rules used

rule 219

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 488

```
Int[1/(((c_) + (d_)*(x_))*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := -Subst[
Int[1/(b*c^2 + a*d^2 - x^2), x], x, (a*d - b*c*x)/Sqrt[a + b*x^2]] /; FreeQ
[{a, b, c, d}, x]
```

rule 491

```
Int[((c_) + (d_)*(x_)^(n_))*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := Simp[
d*(c + d*x)^(n + 1)*((a + b*x^2)^(p + 1)/((n + 1)*(b*c^2 + a*d^2))), x] + S
imp[b*(c/(b*c^2 + a*d^2)) Int[(c + d*x)^(n + 1)*(a + b*x^2)^p, x], x] /;
FreeQ[{a, b, c, d, n, p}, x] && EqQ[n + 2*p + 3, 0]
```

rule 2558

```
Int((((c_) + (d_)*(x_)^(m_))*Sqrt[(b_)*(x_)^2 + Sqrt[(a_) + (e_)*(x_)^
4]])/Sqrt[(a_) + (e_)*(x_)^4], x_Symbol] := Simp[(1 - I)/2 Int[(c + d*x)
^m/Sqrt[Sqrt[a] - I*b*x^2], x], x] + Simp[(1 + I)/2 Int[(c + d*x)^m/Sqrt[
Sqrt[a] + I*b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[e, b^2] && G
tQ[a, 0]
```

Maple [F]

$$\int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{(1+x)^2 \sqrt{x^4 + 1}} dx$$

input

```
int((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)^2/(x^4+1)^(1/2),x)
```

output `int((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)^2/(x^4+1)^(1/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 342 vs. $2(81) = 162$.

Time = 1.07 (sec) , antiderivative size = 342, normalized size of antiderivative = 2.74

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)^2 \sqrt{1 + x^4}} dx$$

$$= \frac{2(x+1)\sqrt{\sqrt{2}+1} \arctan\left(\frac{(x^3+x^2-\sqrt{2}(x^3+1)+\sqrt{x^4+1})(\sqrt{2}x-x-1)-x+1}{x^2-2x+1}\sqrt{x^2+\sqrt{x^4+1}\sqrt{\sqrt{2}+1}}\right) + (x+1)\sqrt{\sqrt{2}-1}}{\dots}$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)^2/(x^4+1)^(1/2),x, algorithm="fricas")`

output `1/8*(2*(x + 1)*sqrt(sqrt(2) + 1)*arctan((x^3 + x^2 - sqrt(2)*(x^3 + 1) + sqrt(x^4 + 1)*(sqrt(2)*x - x - 1) - x + 1)*sqrt(x^2 + sqrt(x^4 + 1))*sqrt(sqrt(2) + 1)/(x^2 - 2*x + 1)) + (x + 1)*sqrt(sqrt(2) - 1)*log(-((2*x^3 - sqrt(2)*(x^3 - x^2 - x - 1) + sqrt(x^4 + 1)*(sqrt(2)*(x - 1) - 2*x) - 2)*sqrt(x^2 + sqrt(x^4 + 1)) + (sqrt(2)*(x^2 + 1) + 2*sqrt(x^4 + 1))*sqrt(sqrt(2) - 1))/(x^2 + 2*x + 1)) - (x + 1)*sqrt(sqrt(2) - 1)*log(-((2*x^3 - sqrt(2)*(x^3 - x^2 - x - 1) + sqrt(x^4 + 1)*(sqrt(2)*(x - 1) - 2*x) - 2)*sqrt(x^2 + sqrt(x^4 + 1)) - (sqrt(2)*(x^2 + 1) + 2*sqrt(x^4 + 1))*sqrt(sqrt(2) - 1))/(x^2 + 2*x + 1)) + 4*sqrt(x^2 + sqrt(x^4 + 1))*(x^2 - sqrt(x^4 + 1) - 1))/(x + 1)`

Sympy [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)^2 \sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{(x + 1)^2 \sqrt{x^4 + 1}} dx$$

input `integrate((x**2+(x**4+1)**(1/2))**(1/2)/(1+x)**2/(x**4+1)**(1/2),x)`

output `Integral(sqrt(x**2 + sqrt(x**4 + 1))/((x + 1)**2*sqrt(x**4 + 1)), x)`

Maxima [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)^2 \sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1} (x + 1)^2} dx$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)^2/(x^4+1)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x^2 + sqrt(x^4 + 1))/(sqrt(x^4 + 1)*(x + 1)^2), x)`

Giac [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)^2 \sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1} (x + 1)^2} dx$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)^2/(x^4+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(x^2 + sqrt(x^4 + 1))/(sqrt(x^4 + 1)*(x + 1)^2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)^2 \sqrt{1 + x^4}} dx = \int \frac{\sqrt{\sqrt{x^4 + 1} + x^2}}{\sqrt{x^4 + 1} (x + 1)^2} dx$$

input `int(((x^4 + 1)^(1/2) + x^2)^(1/2)/((x^4 + 1)^(1/2)*(x + 1)^2),x)`

output `int(((x^4 + 1)^(1/2) + x^2)^(1/2)/((x^4 + 1)^(1/2)*(x + 1)^2), x)`

Reduce [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)^2 \sqrt{1 + x^4}} dx = \int \frac{\sqrt{\sqrt{x^4 + 1} + x^2} \sqrt{x^4 + 1}}{x^6 + 2x^5 + x^4 + x^2 + 2x + 1} dx$$

input `int((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)^2/(x^4+1)^(1/2),x)`

output `int((sqrt(sqrt(x**4 + 1) + x**2))*sqrt(x**4 + 1))/(x**6 + 2*x**5 + x**4 + x**2 + 2*x + 1),x)`

3.13 $\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{(1+x)\sqrt{1+x^4}} dx$

Optimal result	151
Mathematica [B] (verified)	151
Rubi [A] (verified)	152
Maple [F]	153
Fricas [B] (verification not implemented)	154
Sympy [F]	155
Maxima [F]	155
Giac [F]	155
Mupad [F(-1)]	156
Reduce [F]	156

Optimal result

Integrand size = 32, antiderivative size = 81

$$\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{(1+x)\sqrt{1+x^4}} dx = -\frac{1}{2}\sqrt{1-i}\operatorname{arctanh}\left(\frac{1+ix}{\sqrt{1-i}\sqrt{1-ix^2}}\right) - \frac{1}{2}\sqrt{1+i}\operatorname{arctanh}\left(\frac{1-ix}{\sqrt{1+i}\sqrt{1+ix^2}}\right)$$

output `-1/2*arctanh((1+I*x)/(1-I)^(1/2)/(1-I*x^2)^(1/2))*(1-I)^(1/2)-1/2*arctanh((1-I*x)/(1+I)^(1/2)/(1+I*x^2)^(1/2))*(1+I)^(1/2)`

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 205 vs. 2(81) = 162.

Time = 1.09 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.53

$$\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{(1+x)\sqrt{1+x^4}} dx = \frac{\sqrt{-1+\sqrt{2}} \left(\arctan\left(\sqrt{1+\sqrt{2}}\sqrt{x^2+\sqrt{1+x^4}}\right) - \arctan\left(\frac{\sqrt{2(-1+\sqrt{2})}x\sqrt{x^2+\sqrt{1+x^4}}}{1+x^2+\sqrt{1+x^4}}\right) \right) - \sqrt{1+\sqrt{2}}\operatorname{arctan}\left(\frac{\sqrt{1+x^2}}{1+x}\right)}{\sqrt{2}}$$

input `Integrate[Sqrt[x^2 + Sqrt[1 + x^4]]/((1 + x)*Sqrt[1 + x^4]),x]`

output `(Sqrt[-1 + Sqrt[2]]*(ArcTan[Sqrt[1 + Sqrt[2]]*Sqrt[x^2 + Sqrt[1 + x^4]]] - ArcTan[(Sqrt[2*(-1 + Sqrt[2]))*x*Sqrt[x^2 + Sqrt[1 + x^4]]]/(1 + x^2 + Sqrt[1 + x^4]))] - Sqrt[1 + Sqrt[2]]*ArcTanh[Sqrt[-1 + Sqrt[2]]*Sqrt[x^2 + Sqrt[1 + x^4]]] + Sqrt[1 + Sqrt[2]]*ArcTanh[(Sqrt[2*(1 + Sqrt[2]))*x*Sqrt[x^2 + Sqrt[1 + x^4]]]/(1 + x^2 + Sqrt[1 + x^4]))]/Sqrt[2]`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.094$, Rules used = {2558, 488, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sqrt{x^4 + 1} + x^2}}{(x + 1)\sqrt{x^4 + 1}} dx$$

$$\downarrow 2558$$

$$\left(\frac{1}{2} - \frac{i}{2}\right) \int \frac{1}{(x + 1)\sqrt{1 - ix^2}} dx + \left(\frac{1}{2} + \frac{i}{2}\right) \int \frac{1}{(x + 1)\sqrt{ix^2 + 1}} dx$$

$$\downarrow 488$$

$$\left(-\frac{1}{2} + \frac{i}{2}\right) \int \frac{1}{(1 - i) - \frac{(ix+1)^2}{1-ix^2}} d \frac{ix + 1}{\sqrt{1 - ix^2}} - \left(\frac{1}{2} + \frac{i}{2}\right) \int \frac{1}{(1 + i) - \frac{(1-ix)^2}{ix^2+1}} d \frac{1 - ix}{\sqrt{ix^2 + 1}}$$

$$\downarrow 219$$

$$-\frac{1}{2}\sqrt{1 - i}\operatorname{arctanh}\left(\frac{1 + ix}{\sqrt{1 - i}\sqrt{1 - ix^2}}\right) - \frac{1}{2}\sqrt{1 + i}\operatorname{arctanh}\left(\frac{1 - ix}{\sqrt{1 + i}\sqrt{1 + ix^2}}\right)$$

input `Int[Sqrt[x^2 + Sqrt[1 + x^4]]/((1 + x)*Sqrt[1 + x^4]),x]`

output

```
-1/2*(Sqrt[1 - I]*ArcTanh[(1 + I*x)/(Sqrt[1 - I]*Sqrt[1 - I*x^2])]) - (Sqrt[1 + I]*ArcTanh[(1 - I*x)/(Sqrt[1 + I]*Sqrt[1 + I*x^2])])/2
```

Defintions of rubi rules used

rule 219

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

rule 488

```
Int[1/(((c_) + (d_)*(x_))*Sqrt[(a_) + (b_)*(x_)^2]), x_Symbol] := -Subst[Int[1/(b*c^2 + a*d^2 - x^2), x], x, (a*d - b*c*x)/Sqrt[a + b*x^2]] /; FreeQ[{a, b, c, d}, x]
```

rule 2558

```
Int[(((c_) + (d_)*(x_)^(m_))*Sqrt[(b_)*(x_)^2 + Sqrt[(a_) + (e_)*(x_)^4]])/Sqrt[(a_) + (e_)*(x_)^4], x_Symbol] := Simp[(1 - I)/2 Int[(c + d*x)^m/Sqrt[Sqrt[a] - I*b*x^2], x], x] + Simp[(1 + I)/2 Int[(c + d*x)^m/Sqrt[Sqrt[a] + I*b*x^2], x], x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[e, b^2] && GtQ[a, 0]
```

Maple [F]

$$\int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{(1+x)\sqrt{x^4 + 1}} dx$$

input

```
int((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)/(x^4+1)^(1/2),x)
```

output

```
int((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)/(x^4+1)^(1/2),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(49) = 98$.

Time = 3.27 (sec) , antiderivative size = 332, normalized size of antiderivative = 4.10

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)\sqrt{1 + x^4}} dx$$

$$= \frac{1}{2} \sqrt{\frac{1}{2}} \sqrt{2} - \frac{1}{2} \arctan \left(\frac{(2x^2 - \sqrt{2}(x^3 - x^2 + x + 1) + \sqrt{x^4 + 1})(\sqrt{2}(x - 1) - 2) - 2x}{x^2 - 2x + 1}} \sqrt{x^2 + \sqrt{x^4 + 1}} \right)$$

$$- \frac{1}{4} \sqrt{\frac{1}{2}} \sqrt{2} + \frac{1}{2} \log \left(- \frac{(2x^3 - \sqrt{2}(x^3 - x^2 - x - 1) + \sqrt{x^4 + 1})(\sqrt{2}(x - 1) - 2) - 2}{x^2 + 2x + 1}} \sqrt{x^2 + \sqrt{x^4 + 1}} \right)$$

$$+ \frac{1}{4} \sqrt{\frac{1}{2}} \sqrt{2} + \frac{1}{2} \log \left(- \frac{(2x^3 - \sqrt{2}(x^3 - x^2 - x - 1) + \sqrt{x^4 + 1})(\sqrt{2}(x - 1) - 2) - 2}{x^2 + 2x + 1}} \sqrt{x^2 + \sqrt{x^4 + 1}} \right)$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)/(x^4+1)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(1/2*sqrt(2) - 1/2)*arctan((2*x^2 - sqrt(2)*(x^3 - x^2 + x + 1) + sqrt(x^4 + 1)*(sqrt(2)*(x - 1) - 2) - 2*x)*sqrt(x^2 + sqrt(x^4 + 1))*sqrt(1/2*sqrt(2) - 1/2)/(x^2 - 2*x + 1)) - 1/4*sqrt(1/2*sqrt(2) + 1/2)*log(-((2*x^3 - sqrt(2)*(x^3 - x^2 - x - 1) + sqrt(x^4 + 1)*(sqrt(2)*(x - 1) - 2*x) - 2)*sqrt(x^2 + sqrt(x^4 + 1)) + 2*(x^2 - sqrt(2)*(x^2 + 1) + sqrt(x^4 + 1)*(sqrt(2) - 2) + 1)*sqrt(1/2*sqrt(2) + 1/2))/(x^2 + 2*x + 1)) + 1/4*sqrt(1/2*sqrt(2) + 1/2)*log(-((2*x^3 - sqrt(2)*(x^3 - x^2 - x - 1) + sqrt(x^4 + 1)*(sqrt(2)*(x - 1) - 2*x) - 2)*sqrt(x^2 + sqrt(x^4 + 1)) - 2*(x^2 - sqrt(2)*(x^2 + 1) + sqrt(x^4 + 1)*(sqrt(2) - 2) + 1)*sqrt(1/2*sqrt(2) + 1/2))/(x^2 + 2*x + 1))`

Sympy [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)\sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{(x + 1)\sqrt{x^4 + 1}} dx$$

input `integrate((x**2+(x**4+1)**(1/2))**(1/2)/(1+x)/(x**4+1)**(1/2),x)`

output `Integral(sqrt(x**2 + sqrt(x**4 + 1))/((x + 1)*sqrt(x**4 + 1)), x)`

Maxima [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)\sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1}(x + 1)} dx$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)/(x^4+1)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x^2 + sqrt(x^4 + 1))/(sqrt(x^4 + 1)*(x + 1)), x)`

Giac [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)\sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1}(x + 1)} dx$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)/(x^4+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(x^2 + sqrt(x^4 + 1))/(sqrt(x^4 + 1)*(x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)\sqrt{1 + x^4}} dx = \int \frac{\sqrt{\sqrt{x^4 + 1} + x^2}}{\sqrt{x^4 + 1} (x + 1)} dx$$

input `int(((x^4 + 1)^(1/2) + x^2)^(1/2)/((x^4 + 1)^(1/2)*(x + 1)), x)`

output `int(((x^4 + 1)^(1/2) + x^2)^(1/2)/((x^4 + 1)^(1/2)*(x + 1)), x)`

Reduce [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{(1 + x)\sqrt{1 + x^4}} dx = \int \frac{\sqrt{\sqrt{x^4 + 1} + x^2} \sqrt{x^4 + 1}}{x^5 + x^4 + x + 1} dx$$

input `int((x^2+(x^4+1)^(1/2))^(1/2)/(1+x)/(x^4+1)^(1/2), x)`

output `int((sqrt(sqrt(x**4 + 1) + x**2))*sqrt(x**4 + 1))/(x**5 + x**4 + x + 1), x)`

3.14 $\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx$

Optimal result	157
Mathematica [A] (verified)	157
Rubi [A] (verified)	158
Maple [F]	159
Fricas [B] (verification not implemented)	159
Sympy [A] (verification not implemented)	159
Maxima [F]	160
Giac [F]	160
Mupad [F(-1)]	160
Reduce [B] (verification not implemented)	161

Optimal result

Integrand size = 27, antiderivative size = 31

$$\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{x^2 + \sqrt{1+x^4}}}\right)}{\sqrt{2}}$$

output `1/2*arctanh(x*2^(1/2)/(x^2+(x^4+1)^(1/2))^(1/2))*2^(1/2)`

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.42

$$\int \frac{\sqrt{x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx = \frac{\log\left(x^2 + \sqrt{1+x^4} + \sqrt{2}x\sqrt{x^2 + \sqrt{1+x^4}}\right)}{\sqrt{2}}$$

input `Integrate[Sqrt[x^2 + Sqrt[1 + x^4]]/Sqrt[1 + x^4],x]`

output `Log[x^2 + Sqrt[1 + x^4] + Sqrt[2]*x*Sqrt[x^2 + Sqrt[1 + x^4]]]/Sqrt[2]`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {2557, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sqrt{x^4+1}+x^2}}{\sqrt{x^4+1}} dx$$

↓ 2557

$$\int \frac{1}{1 - \frac{2x^2}{\sqrt{x^4+1+x^2}}} d \frac{x}{\sqrt{\sqrt{x^4+1}+x^2}}$$

↓ 219

$$\frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{\sqrt{x^4+1}+x^2}}\right)}{\sqrt{2}}$$

input `Int[Sqrt[x^2 + Sqrt[1 + x^4]]/Sqrt[1 + x^4], x]`

output `ArcTanh[(Sqrt[2]*x)/Sqrt[x^2 + Sqrt[1 + x^4]]]/Sqrt[2]`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 2557 `Int[Sqrt[(c_.)*(x_)^2 + (d_.)*Sqrt[(a_) + (b_.)*(x_)^4]]/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := Simp[d Subst[Int[1/(1 - 2*c*x^2), x], x, x/Sqrt[c*x^2 + d*Sqrt[a + b*x^4]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[c^2 - b*d^2, 0]`

Maple [F]

$$\int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1}} dx$$

input `int((x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x)`

output `int((x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(24) = 48$.

Time = 0.21 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.94

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \frac{1}{4} \sqrt{2} \log \left(4x^4 + 4\sqrt{x^4 + 1}x^2 + 2 \left(\sqrt{2}x^3 + \sqrt{2}\sqrt{x^4 + 1}x \right) \sqrt{x^2 + \sqrt{x^4 + 1} + 1} \right)$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x, algorithm="fricas")`

output `1/4*sqrt(2)*log(4*x^4 + 4*sqrt(x^4 + 1)*x^2 + 2*(sqrt(2)*x^3 + sqrt(2)*sqrt(x^4 + 1)*x)*sqrt(x^2 + sqrt(x^4 + 1)) + 1)`

Sympy [A] (verification not implemented)

Time = 0.91 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.48

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \frac{G_{3,3}^{2,2} \left(\begin{matrix} 1, 1 & \frac{1}{2} \\ \frac{1}{4}, \frac{3}{4} & 0 \end{matrix} \middle| x^4 \right)}{4\sqrt{\pi}}$$

input `integrate((x**2+(x**4+1)**(1/2))**(1/2)/(x**4+1)**(1/2),x)`

output `meijerg(((1, 1), (1/2,)), ((1/4, 3/4), (0,)), x**4)/(4*sqrt(pi))`

Maxima [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1}} dx$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x^2 + sqrt(x^4 + 1))/sqrt(x^4 + 1), x)`

Giac [F]

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \int \frac{\sqrt{x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1}} dx$$

input `integrate((x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(x^2 + sqrt(x^4 + 1))/sqrt(x^4 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \int \frac{\sqrt{\sqrt{x^4 + 1} + x^2}}{\sqrt{x^4 + 1}} dx$$

input `int(((x^4 + 1)^(1/2) + x^2)^(1/2)/(x^4 + 1)^(1/2),x)`

output `int(((x^4 + 1)^(1/2) + x^2)^(1/2)/(x^4 + 1)^(1/2), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

$$\int \frac{\sqrt{x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx$$

$$= \frac{\sqrt{2} \left(-\log\left(\sqrt{\sqrt{x^4 + 1} + x^2} - \sqrt{2}x\right) + \log\left(\sqrt{\sqrt{x^4 + 1} + x^2} + \sqrt{2}x\right) \right)}{4}$$

input `int((x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x)`output `(sqrt(2)*(-log(sqrt(sqrt(x**4 + 1) + x**2) - sqrt(2)*x) + log(sqrt(sqrt(x**4 + 1) + x**2) + sqrt(2)*x)))/4`

3.15 $\int \frac{\sqrt{-x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx$

Optimal result	162
Mathematica [C] (verified)	162
Rubi [A] (verified)	163
Maple [C] (verified)	164
Fricas [A] (verification not implemented)	164
Sympy [A] (verification not implemented)	164
Maxima [F]	165
Giac [F]	165
Mupad [F(-1)]	165
Reduce [F]	166

Optimal result

Integrand size = 29, antiderivative size = 33

$$\int \frac{\sqrt{-x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx = \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{-x^2 + \sqrt{1+x^4}}}\right)}{\sqrt{2}}$$

output `1/2*arctan(x*2^(1/2)/(-x^2+(x^4+1)^(1/2))^(1/2))*2^(1/2)`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.73

$$\int \frac{\sqrt{-x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx = -\frac{i \log\left(ix^2 - i\sqrt{1+x^4} + \sqrt{2}x\sqrt{-x^2 + \sqrt{1+x^4}}\right)}{\sqrt{2}}$$

input `Integrate[Sqrt[-x^2 + Sqrt[1 + x^4]]/Sqrt[1 + x^4],x]`

output `((-I)*Log[I*x^2 - I*Sqrt[1 + x^4] + Sqrt[2]*x*Sqrt[-x^2 + Sqrt[1 + x^4]]]) /Sqrt[2]`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {2557, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sqrt{x^4+1}-x^2}}{\sqrt{x^4+1}} dx$$

↓ 2557

$$\int \frac{1}{\frac{2x^2}{\sqrt{x^4+1-x^2}} + 1} d \frac{x}{\sqrt{\sqrt{x^4+1}-x^2}}$$

↓ 216

$$\frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{\sqrt{x^4+1}-x^2}}\right)}{\sqrt{2}}$$

input `Int[Sqrt[-x^2 + Sqrt[1 + x^4]]/Sqrt[1 + x^4],x]`

output `ArcTan[(Sqrt[2]*x)/Sqrt[-x^2 + Sqrt[1 + x^4]]]/Sqrt[2]`

Defintions of rubi rules used

rule 216 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 2557 `Int[Sqrt[(c_)*(x_)^2 + (d_)*Sqrt[(a_) + (b_)*(x_)^4]]/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := Simp[d Subst[Int[1/(1 - 2*c*x^2), x], x, x/Sqrt[c*x^2 + d*Sqrt[a + b*x^4]]], x] /; FreeQ[{a, b, c, d}, x] && EqQ[c^2 - b*d^2, 0]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.67

method	result	size
meijerg	$-\frac{\sqrt{2} \operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{3}{4}, \frac{5}{4}\right], \left[\frac{3}{2}, \frac{3}{2}\right], -\frac{1}{x^4}\right)}{4x^2}$	22

input `int((-x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/4*2^(1/2)/x^2*hypergeom([1/2,3/4,5/4],[3/2,3/2],-1/x^4)`

Fricas [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.27

$$\int \frac{\sqrt{-x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx = \frac{1}{2} \sqrt{2} \arctan \left(\left(\sqrt{2}x^3 + \sqrt{2}\sqrt{x^4+1}x \right) \sqrt{-x^2 + \sqrt{x^4+1}} \right)$$

input `integrate((-x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x, algorithm="fricas")`

output `1/2*sqrt(2)*arctan((sqrt(2)*x^3 + sqrt(2)*sqrt(x^4 + 1)*x)*sqrt(-x^2 + sqrt(x^4 + 1)))`

Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.45

$$\int \frac{\sqrt{-x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx = \frac{G_{3,3}^{2,2} \left(\begin{matrix} \frac{1}{2}, 1 \\ \frac{1}{4}, \frac{3}{4} \end{matrix} \middle| x^4 \right)}{4\sqrt{\pi}}$$

input `integrate((-x**2+(x**4+1)**(1/2))**(1/2)/(x**4+1)**(1/2),x)`

output `meijerg(((1/2, 1), (1,)), ((1/4, 3/4), (0,)), x**4)/(4*sqrt(pi))`

Maxima [F]

$$\int \frac{\sqrt{-x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \int \frac{\sqrt{-x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1}} dx$$

input `integrate((-x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(-x^2 + sqrt(x^4 + 1))/sqrt(x^4 + 1), x)`

Giac [F]

$$\int \frac{\sqrt{-x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \int \frac{\sqrt{-x^2 + \sqrt{x^4 + 1}}}{\sqrt{x^4 + 1}} dx$$

input `integrate((-x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x, algorithm="giac")`

output `integrate(sqrt(-x^2 + sqrt(x^4 + 1))/sqrt(x^4 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{-x^2 + \sqrt{1 + x^4}}}{\sqrt{1 + x^4}} dx = \int \frac{\sqrt{\sqrt{x^4 + 1} - x^2}}{\sqrt{x^4 + 1}} dx$$

input `int(((x^4 + 1)^(1/2) - x^2)^(1/2)/(x^4 + 1)^(1/2),x)`

output `int(((x^4 + 1)^(1/2) - x^2)^(1/2)/(x^4 + 1)^(1/2), x)`

Reduce [F]

$$\int \frac{\sqrt{-x^2 + \sqrt{1+x^4}}}{\sqrt{1+x^4}} dx = \int \frac{\sqrt{\sqrt{x^4+1} - x^2} \sqrt{x^4+1}}{x^4+1} dx$$

input `int((-x^2+(x^4+1)^(1/2))^(1/2)/(x^4+1)^(1/2),x)`

output `int((sqrt(sqrt(x**4 + 1) - x**2)*sqrt(x**4 + 1))/(x**4 + 1),x)`

$$3.16 \quad \int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx$$

Optimal result	167
Mathematica [A] (verified)	167
Rubi [A] (verified)	168
Maple [A] (verified)	169
Fricas [A] (verification not implemented)	169
Sympy [B] (verification not implemented)	170
Maxima [A] (verification not implemented)	170
Giac [A] (verification not implemented)	170
Mupad [B] (verification not implemented)	171
Reduce [B] (verification not implemented)	171

Optimal result

Integrand size = 30, antiderivative size = 19

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = -\frac{2}{\sqrt{-1+x}} - \frac{2}{\sqrt{1+x}}$$

output `-2/(-1+x)^(1/2)-2/(1+x)^(1/2)`

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = -\frac{2}{\sqrt{-1+x}} - \frac{2}{\sqrt{1+x}}$$

input `Integrate[((-1 + x)^(3/2) + (1 + x)^(3/2))/((-1 + x)^(3/2)*(1 + x)^(3/2)), x]`

output `-2/Sqrt[-1 + x] - 2/Sqrt[1 + x]`

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {7239, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x-1)^{3/2} + (x+1)^{3/2}}{(x-1)^{3/2}(x+1)^{3/2}} dx$$

↓ 7239

$$\int \left(\frac{1}{(x+1)^{3/2}} + \frac{1}{(x-1)^{3/2}} \right) dx$$

↓ 2009

$$-\frac{2}{\sqrt{x+1}} - \frac{2}{\sqrt{x-1}}$$

input `Int[((-1 + x)^(3/2) + (1 + x)^(3/2))/((-1 + x)^(3/2)*(1 + x)^(3/2)),x]`

output `-2/Sqrt[-1 + x] - 2/Sqrt[1 + x]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 7239 `Int[u_, x_Symbol] :> With[{v = SimplifyIntegrand[u, x]}, Int[v, x] /; SimplifierIntegrandQ[v, u, x]]`

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
default	$-\frac{2}{\sqrt{-1+x}} - \frac{2}{\sqrt{1+x}}$	16
meijerg	$\frac{2\sqrt{\pi} - \frac{2\sqrt{\pi}}{\sqrt{1+x}}}{\sqrt{\pi}} - \frac{2(-\text{signum}(-1+x))^{\frac{3}{2}} \left(\sqrt{\pi} - \frac{\sqrt{\pi}}{\sqrt{1-x}} \right)}{\sqrt{\pi} \text{signum}(-1+x)^{\frac{3}{2}}}$	56
orering	$-\frac{4x \left((-1+x)^{\frac{3}{2}} + (1+x)^{\frac{3}{2}} \right)}{(-1+x)^{\frac{3}{2}} (1+x)^{\frac{3}{2}}} - \frac{4(-1+x)(1+x) \left(\frac{\frac{3\sqrt{-1+x}}{2} + \frac{3\sqrt{1+x}}{2}}{(-1+x)^{\frac{3}{2}} (1+x)^{\frac{3}{2}}} - \frac{3 \left((-1+x)^{\frac{3}{2}} + (1+x)^{\frac{3}{2}} \right)}{2(-1+x)^{\frac{5}{2}} (1+x)^{\frac{3}{2}}} - \frac{3 \left((-1+x)^{\frac{3}{2}} + (1+x)^{\frac{3}{2}} \right)}{2(-1+x)^{\frac{3}{2}} (1+x)^{\frac{5}{2}}} \right)}{3}$	107

input `int(((−1+x)^(3/2)+(1+x)^(3/2))/(−1+x)^(3/2)/(1+x)^(3/2),x,method=_RETURNVE
RBOSE)`

output `−2/(−1+x)^(1/2)−2/(1+x)^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.47

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = -\frac{2((x+1)\sqrt{x-1} + \sqrt{x+1}(x-1))}{x^2-1}$$

input `integrate(((−1+x)^(3/2)+(1+x)^(3/2))/(−1+x)^(3/2)/(1+x)^(3/2),x, algorithm
="fricas")`

output `−2*((x + 1)*sqrt(x − 1) + sqrt(x + 1)*(x − 1))/(x^2 − 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(17) = 34$.

Time = 0.66 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.95

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = -\frac{2x\sqrt{x-1}}{x^2-1} - \frac{2x\sqrt{x+1}}{x^2-1} - \frac{2\sqrt{x-1}}{x^2-1} + \frac{2\sqrt{x+1}}{x^2-1}$$

input `integrate(((−1+x)**(3/2)+(1+x)**(3/2))/(−1+x)**(3/2)/(1+x)**(3/2),x)`

output `−2*x*sqrt(x − 1)/(x**2 − 1) − 2*x*sqrt(x + 1)/(x**2 − 1) − 2*sqrt(x − 1)/(x**2 − 1) + 2*sqrt(x + 1)/(x**2 − 1)`

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = -\frac{2}{\sqrt{x+1}} - \frac{2}{\sqrt{x-1}}$$

input `integrate(((−1+x)^(3/2)+(1+x)^(3/2))/(−1+x)^(3/2)/(1+x)^(3/2),x, algorithm="maxima")`

output `−2/sqrt(x + 1) − 2/sqrt(x − 1)`

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = -\frac{2}{\sqrt{x+1}} - \frac{2}{\sqrt{x-1}}$$

input `integrate(((−1+x)^(3/2)+(1+x)^(3/2))/(−1+x)^(3/2)/(1+x)^(3/2),x, algorithm="giac")`

output $-2/\sqrt{x + 1} - 2/\sqrt{x - 1}$

Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = -\frac{2}{\sqrt{x-1}} - \frac{2}{\sqrt{x+1}}$$

input `int(((x - 1)^(3/2) + (x + 1)^(3/2))/((x - 1)^(3/2)*(x + 1)^(3/2)),x)`

output $-2/(x - 1)^{1/2} - 2/(x + 1)^{1/2}$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.37

$$\int \frac{(-1+x)^{3/2} + (1+x)^{3/2}}{(-1+x)^{3/2}(1+x)^{3/2}} dx = \frac{-2\sqrt{x-1} - 2\sqrt{x+1}}{\sqrt{x+1}\sqrt{x-1}}$$

input `int(((-1+x)^(3/2)+(1+x)^(3/2))/((-1+x)^(3/2)/(1+x)^(3/2)),x)`

output $(-2*(\sqrt{x - 1} + \sqrt{x + 1}))/(\sqrt{x + 1}*\sqrt{x - 1})$

3.17 $\int \left(x + \sqrt{a + x^2}\right)^b dx$

Optimal result	172
Mathematica [A] (verified)	172
Rubi [A] (verified)	173
Maple [B] (verified)	174
Fricas [A] (verification not implemented)	175
Sympy [B] (verification not implemented)	175
Maxima [F]	176
Giac [F]	177
Mupad [F(-1)]	177
Reduce [B] (verification not implemented)	177

Optimal result

Integrand size = 13, antiderivative size = 52

$$\int \left(x + \sqrt{a + x^2}\right)^b dx = -\frac{a(x + \sqrt{a + x^2})^{-1+b}}{2(1-b)} + \frac{(x + \sqrt{a + x^2})^{1+b}}{2(1+b)}$$

output `-1/2*a*(x+(x^2+a)^(1/2))^(1+b)/(1+b)+1/2*(x+(x^2+a)^(1/2))^(1+b)/(1+b)`

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.83

$$\int \left(x + \sqrt{a + x^2}\right)^b dx = \frac{(x + \sqrt{a + x^2})^{-1+b} (ab + (-1 + b)x(x + \sqrt{a + x^2}))}{-1 + b^2}$$

input `Integrate[(x + Sqrt[a + x^2])^b,x]`

output `((x + Sqrt[a + x^2])^(1+b)*(a*b + (-1 + b)*x*(x + Sqrt[a + x^2])))/(-1 + b^2)`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2542, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{a+x^2} + x)^b dx$$

$$\downarrow \text{2542}$$

$$\frac{1}{2} \int (x + \sqrt{x^2 + a})^{b-2} \left((x + \sqrt{x^2 + a})^2 + a \right) d(x + \sqrt{x^2 + a})$$

$$\downarrow \text{244}$$

$$\frac{1}{2} \int \left(a(x + \sqrt{x^2 + a})^{b-2} + (x + \sqrt{x^2 + a})^b \right) d(x + \sqrt{x^2 + a})$$

$$\downarrow \text{2009}$$

$$\frac{1}{2} \left(\frac{(\sqrt{a+x^2} + x)^{b+1}}{b+1} - \frac{a(\sqrt{a+x^2} + x)^{b-1}}{1-b} \right)$$

input `Int[(x + Sqrt[a + x^2])^b,x]`

output `((-(a*(x + Sqrt[a + x^2])^(-1 + b))/(1 - b)) + (x + Sqrt[a + x^2])^(1 + b)/(1 + b))/2`

Definitions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2]))^(n_)^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(44) = 88.

Time = 0.04 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.31

method	result	size
meijerg	$\frac{a^{\frac{b}{2} + \frac{1}{2}b} \left(\frac{8\sqrt{\pi} x^{1+b} a^{-\frac{b}{2} - \frac{1}{2}} \left(\frac{ab}{x^2} + b - 1 \right) \left(\sqrt{1 + \frac{a}{x^2}} + 1 \right)^{-1+b}}{(1+b)b(2b-2)} + \frac{4\sqrt{\pi} x^{1+b} a^{-\frac{b}{2} - \frac{1}{2}} \sqrt{1 + \frac{a}{x^2}} \left(\sqrt{1 + \frac{a}{x^2}} + 1 \right)^{-1+b}}{(1+b)b} \right)}{4\sqrt{\pi}}$	120

input `int((x+(x^2+a)^(1/2))^b,x,method=_RETURNVERBOSE)`

output `1/4*a^(1/2*b+1/2)/Pi^(1/2)*b*(8*Pi^(1/2)/(1+b)/b*x^(1+b)*a^(-1/2*b-1/2)*(a*b/x^2+b-1)/(2*b-2)*((1+1/x^2*a)^(1/2)+1)^(-1+b)+4*Pi^(1/2)/(1+b)/b*x^(1+b)*a^(-1/2*b-1/2)*(1+1/x^2*a)^(1/2)*((1+1/x^2*a)^(1/2)+1)^(-1+b))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.62

$$\int \left(x + \sqrt{a + x^2}\right)^b dx = \frac{(\sqrt{x^2 + ab} - x)(x + \sqrt{x^2 + a})^b}{b^2 - 1}$$

input `integrate((x+(x^2+a)^(1/2))^b,x, algorithm="fricas")`

output `(sqrt(x^2 + a)*b - x)*(x + sqrt(x^2 + a))^b/(b^2 - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2236 vs. 2(37) = 74.

Time = 1.51 (sec) , antiderivative size = 2236, normalized size of antiderivative = 43.00

$$\int \left(x + \sqrt{a + x^2}\right)^b dx = \text{Too large to display}$$

input `integrate((x+(x**2+a)**(1/2))**b,x)`

output

```
Piecewise((2*a**(9/2)*a**(b/2 + 1/2)*b*cosh(b*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - b/2)/(2*a**(9/2)*b**2*gamma(1 - b/2) - 2*a**(9/2)*gamma(1 - b/2) + 2*a**(7/2)*b**2*x**2*gamma(1 - b/2) - 2*a**(7/2)*x**2*gamma(1 - b/2)) - 2*a**(9/2)*a**(b/2 + 1/2)*b*gamma(1 - b/2)/(2*a**(9/2)*b**2*gamma(1 - b/2) - 2*a**(9/2)*gamma(1 - b/2) + 2*a**(7/2)*b**2*x**2*gamma(1 - b/2) - 2*a**(7/2)*x**2*gamma(1 - b/2)) - 2*a**(7/2)*a**(b/2 + 1/2)*b*x**2*sqrt(a/x**2 + 1)*sinh(b*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - b/2)/(2*a**(9/2)*b**2*gamma(1 - b/2) - 2*a**(9/2)*gamma(1 - b/2) + 2*a**(7/2)*b**2*x**2*gamma(1 - b/2) - 2*a**(7/2)*x**2*gamma(1 - b/2)) + 4*a**(7/2)*a**(b/2 + 1/2)*b*x**2*cosh(b*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - b/2)/(2*a**(9/2)*b**2*gamma(1 - b/2) - 2*a**(9/2)*gamma(1 - b/2) + 2*a**(7/2)*b**2*x**2*gamma(1 - b/2) - 2*a**(7/2)*x**2*gamma(1 - b/2)) - 2*a**(7/2)*a**(b/2 + 1/2)*b*x**2*gamma(1 - b/2)/(2*a**(9/2)*b**2*gamma(1 - b/2) - 2*a**(9/2)*gamma(1 - b/2) + 2*a**(7/2)*b**2*x**2*gamma(1 - b/2) - 2*a**(7/2)*x**2*gamma(1 - b/2)) - 2*a**(7/2)*a**(b/2 + 1/2)*x**2*sqrt(a/x**2 + 1)*sinh(b*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - b/2)/(2*a**(9/2)*b**2*gamma(1 - b/2) - 2*a**(9/2)*gamma(1 - b/2) + 2*a**(7/2)*b**2*x**2*gamma(1 - b/2) - 2*a**(7/2)*x**2*gamma(1 - b/2)) + 2*a**(7/2)*a**(b/2 + 1/2)*x**2*cosh(b*asinh(x/sqrt(a)) + asinh(x/sqrt(a)))*gamma(1 - b/2)/(2*a**(9/2)*b**2*gamma(1 - b/2) - 2*a**(9/2)*gamma(1 - b/2) + 2*a**(7/2)*b**2*x**2*gam...
```

Maxima [F]

$$\int (x + \sqrt{a + x^2})^b dx = \int (x + \sqrt{x^2 + a})^b dx$$

input

```
integrate((x+(x^2+a)^(1/2))^b,x, algorithm="maxima")
```

output

```
integrate((x + sqrt(x^2 + a))^b, x)
```

Giac [F]

$$\int (x + \sqrt{a + x^2})^b dx = \int (x + \sqrt{x^2 + a})^b dx$$

input `integrate((x+(x^2+a)^(1/2))^b,x, algorithm="giac")`

output `integrate((x + sqrt(x^2 + a))^b, x)`

Mupad [F(-1)]

Timed out.

$$\int (x + \sqrt{a + x^2})^b dx = \int (x + \sqrt{x^2 + a})^b dx$$

input `int((x + (a + x^2)^(1/2))^b,x)`

output `int((x + (a + x^2)^(1/2))^b, x)`

Reduce [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.58

$$\int (x + \sqrt{a + x^2})^b dx = \frac{(\sqrt{x^2 + a} + x)^b (\sqrt{x^2 + a} b - x)}{b^2 - 1}$$

input `int((x+(x^2+a)^(1/2))^b,x)`

output `((sqrt(a + x**2) + x)**b*(sqrt(a + x**2)*b - x))/(b**2 - 1)`

3.18 $\int \left(x - \sqrt{a + x^2}\right)^b dx$

Optimal result	178
Mathematica [A] (verified)	178
Rubi [A] (verified)	179
Maple [F]	180
Fricas [A] (verification not implemented)	180
Sympy [F]	181
Maxima [F]	181
Giac [F]	181
Mupad [F(-1)]	182
Reduce [B] (verification not implemented)	182

Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \left(x - \sqrt{a + x^2}\right)^b dx = -\frac{a(x - \sqrt{a + x^2})^{-1+b}}{2(1 - b)} + \frac{(x - \sqrt{a + x^2})^{1+b}}{2(1 + b)}$$

output `-1/2*a*(x-(x^2+a)^(1/2))^(1+b)/(1+b)+1/2*(x-(x^2+a)^(1/2))^(1+b)/(1+b)`

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.89

$$\int \left(x - \sqrt{a + x^2}\right)^b dx = \frac{1}{2} \left(x - \sqrt{a + x^2}\right)^{-1+b} \left(\frac{a}{-1 + b} + \frac{(x - \sqrt{a + x^2})^2}{1 + b} \right)$$

input `Integrate[(x - Sqrt[a + x^2])^b,x]`

output `((x - Sqrt[a + x^2])^(-1 + b)*(a/(-1 + b) + (x - Sqrt[a + x^2])^2/(1 + b)))/2`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2542, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (x - \sqrt{a + x^2})^b dx$$

$$\downarrow \text{2542}$$

$$\frac{1}{2} \int (x - \sqrt{x^2 + a})^{b-2} \left((x - \sqrt{x^2 + a})^2 + a \right) d(x - \sqrt{x^2 + a})$$

$$\downarrow \text{244}$$

$$\frac{1}{2} \int \left(a(x - \sqrt{x^2 + a})^{b-2} + (x - \sqrt{x^2 + a})^b \right) d(x - \sqrt{x^2 + a})$$

$$\downarrow \text{2009}$$

$$\frac{1}{2} \left(\frac{(x - \sqrt{a + x^2})^{b+1}}{b+1} - \frac{a(x - \sqrt{a + x^2})^{b-1}}{1-b} \right)$$

input `Int[(x - Sqrt[a + x^2])^b,x]`

output `((-(a*(x - Sqrt[a + x^2])^(-1 + b))/(1 - b)) + (x - Sqrt[a + x^2])^(1 + b)/(1 + b))/2`

Defintions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.))^(p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [F]

$$\int (x - \sqrt{x^2 + a})^b dx$$

input `int((x-(x^2+a)^(1/2))^b,x)`

output `int((x-(x^2+a)^(1/2))^b,x)`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.59

$$\int (x - \sqrt{a + x^2})^b dx = -\frac{(\sqrt{x^2 + ab} + x)(x - \sqrt{x^2 + a})^b}{b^2 - 1}$$

input `integrate((x-(x^2+a)^(1/2))^b,x, algorithm="fricas")`

output `-(sqrt(x^2 + a)*b + x)*(x - sqrt(x^2 + a))^b/(b^2 - 1)`

Sympy [F]

$$\int (x - \sqrt{a + x^2})^b dx = \int (x - \sqrt{a + x^2})^b dx$$

input `integrate((x-(x**2+a)**(1/2))**b,x)`

output `Integral((x - sqrt(a + x**2))**b, x)`

Maxima [F]

$$\int (x - \sqrt{a + x^2})^b dx = \int (x - \sqrt{x^2 + a})^b dx$$

input `integrate((x-(x^2+a)^(1/2))^b,x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^b, x)`

Giac [F]

$$\int (x - \sqrt{a + x^2})^b dx = \int (x - \sqrt{x^2 + a})^b dx$$

input `integrate((x-(x^2+a)^(1/2))^b,x, algorithm="giac")`

output `integrate((x - sqrt(x^2 + a))^b, x)`

Mupad [F(-1)]

Timed out.

$$\int (x - \sqrt{a + x^2})^b dx = \int (x - \sqrt{x^2 + a})^b dx$$

input `int((x - (a + x^2)^(1/2))^b, x)`output `int((x - (a + x^2)^(1/2))^b, x)`**Reduce [B] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.66

$$\int (x - \sqrt{a + x^2})^b dx = -\frac{a^b(-1)^b(\sqrt{x^2 + a}b + x)}{(\sqrt{x^2 + a} + x)^b(b^2 - 1)}$$

input `int((x-(x^2+a)^(1/2))^b,x)`output `(- a**b*(- 1)**b*(sqrt(a + x**2)*b + x))/((sqrt(a + x**2) + x)**b*(b**2 - 1))`

3.19
$$\int \frac{(x + \sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx$$

Optimal result	183
Mathematica [A] (verified)	183
Rubi [A] (verified)	184
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	185
Sympy [B] (verification not implemented)	185
Maxima [F]	186
Giac [A] (verification not implemented)	187
Mupad [B] (verification not implemented)	187
Reduce [B] (verification not implemented)	187

Optimal result

Integrand size = 23, antiderivative size = 17

$$\int \frac{(x + \sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx = \frac{(x + \sqrt{a+x^2})^b}{b}$$

output

```
(x+(x^2+a)^(1/2))^b/b
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{(x + \sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx = \frac{(x + \sqrt{a+x^2})^b}{b}$$

input

```
Integrate[(x + Sqrt[a + x^2])^b/Sqrt[a + x^2],x]
```

output

```
(x + Sqrt[a + x^2])^b/b
```


Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2547, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{a+x^2}+x)^b}{\sqrt{a+x^2}} dx$$

↓ 2547

$$\int (\sqrt{a+x^2}+x)^{b-1} d(\sqrt{a+x^2}+x)$$

↓ 15

$$\frac{(\sqrt{a+x^2}+x)^b}{b}$$

input `Int[(x + Sqrt[a + x^2])^b/Sqrt[a + x^2],x]`

output `(x + Sqrt[a + x^2])^b/b`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{(x+\sqrt{x^2+a})^b}{b}$	16
default	$\frac{(x+\sqrt{x^2+a})^b}{b}$	16

input `int((x+(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

output `(x+(x^2+a)^(1/2))^b/b`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{x^2 + a})^b}{b}$$

input `integrate((x+(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x, algorithm="fricas")`

output `(x + sqrt(x^2 + a))^b/b`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. $2(12) = 24$.

Time = 1.31 (sec) , antiderivative size = 311, normalized size of antiderivative = 18.29

$$\int \frac{(x + \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx$$

$$= \begin{cases} \frac{\sqrt{a} a^{\frac{b}{2}} \sinh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{bx \sqrt{\frac{a}{x^2} + 1}} - \frac{2a^{\frac{b}{2}} \cosh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right) \Gamma\left(1 - \frac{b}{2}\right)}{b^2 \Gamma\left(-\frac{b}{2}\right)} + \frac{a^{\frac{b}{2}} x \cosh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{\sqrt{ab}} + \frac{a^{\frac{b}{2}} x \sinh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{\sqrt{ab}} \\ \frac{a^{\frac{b}{2}} \sinh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{b \sqrt{1 + \frac{x^2}{a}}} - \frac{2a^{\frac{b}{2}} \cosh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right) \Gamma\left(1 - \frac{b}{2}\right)}{b^2 \Gamma\left(-\frac{b}{2}\right)} + \frac{a^{\frac{b}{2}} x^2 \sinh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{ab \sqrt{1 + \frac{x^2}{a}}} + \frac{a^{\frac{b}{2}} x \cosh\left(b \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{a}}\right)\right)}{ab \sqrt{1 + \frac{x^2}{a}}} \end{cases}$$

input `integrate((x+(x**2+a)**(1/2))**b/(x**2+a)**(1/2),x)`

output `Piecewise((sqrt(a)*a**(b/2)*sinh(b*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(b*x*sqrt(a/x**2 + 1)) - 2*a**(b/2)*cosh(b*asinh(x/sqrt(a)))*gamma(1 - b/2)/(b**2*gamma(-b/2)) + a**(b/2)*x*cosh(b*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(sqrt(a)*b) + a**(b/2)*x*sinh(b*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(sqrt(a)*b*sqrt(a/x**2 + 1)), Abs(x**2/a) > 1), (a**(b/2)*sinh(b*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(b*sqrt(1 + x**2/a)) - 2*a**(b/2)*cosh(b*asinh(x/sqrt(a)))*gamma(1 - b/2)/(b**2*gamma(-b/2)) + a**(b/2)*x**2*sinh(b*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(a*b*sqrt(1 + x**2/a)) + a**(b/2)*x*cosh(b*asinh(x/sqrt(a)) - asinh(x/sqrt(a)))/(sqrt(a)*b), True))`

Maxima [F]

$$\int \frac{(x + \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = \int \frac{(x + \sqrt{x^2 + a})^b}{\sqrt{x^2 + a}} dx$$

input `integrate((x+(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 + a))^b/sqrt(x^2 + a), x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{x^2 + a})^b}{b}$$

input `integrate((x+(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x, algorithm="giac")`output `(x + sqrt(x^2 + a))^b/b`**Mupad [B] (verification not implemented)**

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = \frac{(x + \sqrt{x^2 + a})^b}{b}$$

input `int((x + (a + x^2)^(1/2))^b/(a + x^2)^(1/2),x)`output `(x + (a + x^2)^(1/2))^b/b`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{(x + \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = \frac{(\sqrt{x^2 + a} + x)^b}{b}$$

input `int((x+(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x)`output `(sqrt(a + x**2) + x)**b/b`

$$3.20 \quad \int \frac{(x - \sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx$$

Optimal result	188
Mathematica [A] (verified)	188
Rubi [A] (verified)	189
Maple [A] (verified)	190
Fricas [A] (verification not implemented)	190
Sympy [B] (verification not implemented)	190
Maxima [F]	191
Giac [A] (verification not implemented)	191
Mupad [B] (verification not implemented)	192
Reduce [B] (verification not implemented)	192

Optimal result

Integrand size = 25, antiderivative size = 20

$$\int \frac{(x - \sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx = -\frac{(x - \sqrt{a+x^2})^b}{b}$$

output

```
-(x-(x^2+a)^(1/2))^b/b
```

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(x - \sqrt{a+x^2})^b}{\sqrt{a+x^2}} dx = -\frac{(x - \sqrt{a+x^2})^b}{b}$$

input

```
Integrate[(x - Sqrt[a + x^2])^b/Sqrt[a + x^2], x]
```

output

```
-((x - Sqrt[a + x^2])^b/b)
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2547, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x - \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx$$

↓ 2547

$$- \int (x - \sqrt{x^2 + a})^{b-1} d(x - \sqrt{x^2 + a})$$

↓ 15

$$-\frac{(x - \sqrt{a + x^2})^b}{b}$$

input `Int[(x - Sqrt[a + x^2])^b/Sqrt[a + x^2],x]`

output `-((x - Sqrt[a + x^2])^b/b)`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

method	result	size
derivativedivides	$-\frac{(x-\sqrt{x^2+a})^b}{b}$	19
default	$-\frac{(x-\sqrt{x^2+a})^b}{b}$	19

input `int((x-(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

output `-(x-(x^2+a)^(1/2))^b/b`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(x - \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = -\frac{(x - \sqrt{x^2 + a})^b}{b}$$

input `integrate((x-(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x, algorithm="fricas")`

output `-(x - sqrt(x^2 + a))^b/b`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. $2(14) = 28$.

Time = 0.85 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.05

$$\int \frac{(x - \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = \begin{cases} -\frac{(x - \sqrt{a + x^2})^b}{b} & \text{for } b \neq 0 \\ \begin{cases} \log(2x + 2\sqrt{a + x^2}) & \text{for } a \neq 0 \\ \frac{x \log(x)}{\sqrt{x^2}} & \text{otherwise} \end{cases} & \text{otherwise} \end{cases}$$

input `integrate((x-(x**2+a)**(1/2))**b/(x**2+a)**(1/2),x)`

output `Piecewise((-x - sqrt(a + x**2))**b/b, Ne(b, 0)), (Piecewise((log(2*x + 2*sqrt(a + x**2)), Ne(a, 0)), (x*log(x)/sqrt(x**2), True)), True))`

Maxima [F]

$$\int \frac{(x - \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = \int \frac{(x - \sqrt{x^2 + a})^b}{\sqrt{x^2 + a}} dx$$

input `integrate((x-(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate((x - sqrt(x^2 + a))^b/sqrt(x^2 + a), x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(x - \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = -\frac{(x - \sqrt{x^2 + a})^b}{b}$$

input `integrate((x-(x^2+a)^(1/2))^b/(x^2+a)^(1/2),x, algorithm="giac")`

output `-(x - sqrt(x^2 + a))^b/b`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(x - \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = -\frac{(x - \sqrt{x^2 + a})^b}{b}$$

input `int((x - (a + x^2)^(1/2))^b/(a + x^2)^(1/2), x)`output `-(x - (a + x^2)^(1/2))^b/b`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int \frac{(x - \sqrt{a + x^2})^b}{\sqrt{a + x^2}} dx = -\frac{a^b(-1)^b}{(\sqrt{x^2 + a} + x)^b b}$$

input `int((x-(x^2+a)^(1/2))^b/(x^2+a)^(1/2), x)`output `(- a**b*(- 1)**b)/((sqrt(a + x**2) + x)**b*b)`

3.21 $\int \frac{1}{(a+be^{px})^2} dx$

Optimal result	193
Mathematica [A] (verified)	193
Rubi [A] (verified)	194
Maple [A] (verified)	195
Fricas [A] (verification not implemented)	195
Sympy [A] (verification not implemented)	196
Maxima [A] (verification not implemented)	196
Giac [A] (verification not implemented)	196
Mupad [B] (verification not implemented)	197
Reduce [B] (verification not implemented)	197

Optimal result

Integrand size = 11, antiderivative size = 42

$$\int \frac{1}{(a + be^{px})^2} dx = \frac{1}{a(a + be^{px})p} + \frac{x}{a^2} - \frac{\log(a + be^{px})}{a^2p}$$

output 1/a/(a+b*exp(p*x))/p+x/a^2-ln(a+b*exp(p*x))/a^2/p

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.93

$$\int \frac{1}{(a + be^{px})^2} dx = \frac{\frac{a}{a+be^{px}} + \log(e^{px}) - \log(a + be^{px})}{a^2p}$$

input Integrate[(a + b*E^(p*x))^(-2), x]

output (a/(a + b*E^(p*x)) + Log[E^(p*x)] - Log[a + b*E^(p*x)])/(a^2*p)

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.07, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {2720, 54, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(a + be^{px})^2} dx \\ & \quad \downarrow \text{2720} \\ & \int \frac{e^{-px}}{(a+be^{px})^2} de^{px} \\ & \quad \downarrow \text{54} \\ & \int \left(-\frac{b}{a^2(a+be^{px})} - \frac{b}{a(a+be^{px})^2} + \frac{e^{-px}}{a^2} \right) de^{px} \\ & \quad \downarrow \text{2009} \\ & \frac{-\frac{\log(a+be^{px})}{a^2} + \frac{\log(e^{px})}{a^2} + \frac{1}{a(a+be^{px})}}{p} \end{aligned}$$

input `Int[(a + b*E^(p*x))^-2], x]`

output `(1/(a*(a + b*E^(p*x))) + Log[E^(p*x)]/a^2 - Log[a + b*E^(p*x)]/a^2)/p`

Defintions of rubi rules used

rule 54 `Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d}, x] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2720

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x]
  Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Funct
ionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ
[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))
*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.02

method	result	size
derivativedivides	$\frac{\frac{\ln(e^{px})}{a^2} - \frac{\ln(a+be^{px})}{a^2} + \frac{1}{a(a+be^{px})}}{p}$	43
default	$\frac{\frac{\ln(e^{px})}{a^2} - \frac{\ln(a+be^{px})}{a^2} + \frac{1}{a(a+be^{px})}}{p}$	43
risch	$\frac{x}{a^2} + \frac{1}{a(a+be^{px})p} - \frac{\ln(e^{px} + \frac{a}{b})}{a^2p}$	43
norman	$\frac{\frac{x}{a} + \frac{bx e^{px}}{a^2} - \frac{b e^{px}}{a^2 p}}{a+be^{px}} - \frac{\ln(a+be^{px})}{a^2 p}$	59
parallelrisch	$-\frac{-b^2 e^{px} xp + \ln(a+be^{px}) e^{px} b^2 - xabp + \ln(a+be^{px}) ab - ab}{a^2 bp(a+be^{px})}$	73

input

```
int(1/(a+b*exp(p*x))^2,x,method=_RETURNVERBOSE)
```

output

```
1/p*(1/a^2*ln(exp(p*x))-1/a^2*ln(a+b*exp(p*x))+1/a/(a+b*exp(p*x)))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.24

$$\int \frac{1}{(a + be^{px})^2} dx = \frac{bp x e^{(px)} + apx - (be^{(px)} + a) \log (be^{(px)} + a) + a}{a^2 b p e^{(px)} + a^3 p}$$

input

```
integrate(1/(a+b*exp(p*x))^2,x, algorithm="fricas")
```

output $(b*p*x*e^{(p*x)} + a*p*x - (b*e^{(p*x)} + a)*\log(b*e^{(p*x)} + a) + a)/(a^2*b*p*e^{(p*x)} + a^3*p)$

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{1}{(a + be^{px})^2} dx = \frac{1}{a^2p + abpe^{px}} + \frac{x}{a^2} - \frac{\log\left(\frac{a}{b} + e^{px}\right)}{a^2p}$$

input `integrate(1/(a+b*exp(p*x))**2,x)`

output $1/(a**2*p + a*b*p*exp(p*x)) + x/a**2 - \log(a/b + exp(p*x))/(a**2*p)$

Maxima [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.95

$$\int \frac{1}{(a + be^{px})^2} dx = \frac{x}{a^2} + \frac{1}{(abe^{px} + a^2)p} - \frac{\log(be^{px} + a)}{a^2p}$$

input `integrate(1/(a+b*exp(p*x))^2,x, algorithm="maxima")`

output $x/a^2 + 1/((a*b*e^{(p*x)} + a^2)*p) - \log(b*e^{(p*x)} + a)/(a^2*p)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.98

$$\int \frac{1}{(a + be^{px})^2} dx = \frac{x}{a^2} - \frac{\log(|be^{px} + a|)}{a^2p} + \frac{1}{(be^{px} + a)ap}$$

input `integrate(1/(a+b*exp(p*x))^2,x, algorithm="giac")`

output $x/a^2 - \log(\text{abs}(b \cdot e^{(p \cdot x)} + a))/(a^2 \cdot p) + 1/((b \cdot e^{(p \cdot x)} + a) \cdot a \cdot p)$

Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.38

$$\int \frac{1}{(a + b e^{p x})^2} dx = \frac{\frac{x}{a} + \frac{b x e^{p x}}{a^2} - \frac{b e^{p x}}{a^2 p}}{a + b e^{p x}} - \frac{\ln(a + b e^{p x})}{a^2 p}$$

input `int(1/(a + b*exp(p*x))^2,x)`

output $(x/a + (b \cdot x \cdot \exp(p \cdot x))/a^2 - (b \cdot \exp(p \cdot x))/(a^2 \cdot p))/(a + b \cdot \exp(p \cdot x)) - \log(a + b \cdot \exp(p \cdot x))/(a^2 \cdot p)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.69

$$\int \frac{1}{(a + b e^{p x})^2} dx = \frac{-e^{p x} \log(e^{p x} b + a) b + e^{p x} b p x - e^{p x} b - \log(e^{p x} b + a) a + a p x}{a^2 p (e^{p x} b + a)}$$

input `int(1/(a+b*exp(p*x))^2,x)`

output $(-e^{(p \cdot x)} \cdot \log(e^{(p \cdot x)} \cdot b + a) \cdot b + e^{(p \cdot x)} \cdot b \cdot p \cdot x - e^{(p \cdot x)} \cdot b - \log(e^{(p \cdot x)} \cdot b + a) \cdot a + a \cdot p \cdot x)/(a^2 \cdot p \cdot (e^{(p \cdot x)} \cdot b + a))$

$$3.22 \quad \int \frac{1}{(be^{-px} + ae^{px})^2} dx$$

Optimal result	198
Mathematica [A] (verified)	198
Rubi [A] (verified)	199
Maple [A] (verified)	200
Fricas [A] (verification not implemented)	200
Sympy [A] (verification not implemented)	201
Maxima [A] (verification not implemented)	201
Giac [A] (verification not implemented)	201
Mupad [B] (verification not implemented)	202
Reduce [B] (verification not implemented)	202

Optimal result

Integrand size = 18, antiderivative size = 22

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = -\frac{1}{2a(b + ae^{2px})p}$$

output `-1/2/a/(b+a*exp(2*p*x))/p`

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = -\frac{1}{2a(b + ae^{2px})p}$$

input `Integrate[(b/E^(p*x) + a*E^(p*x))^(-2), x]`

output `-1/2*1/(a*(b + a*E^(2*p*x))*p)`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2720, 241}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(ae^{px} + be^{-px})^2} dx$$

↓ 2720

$$\int \frac{e^{px}}{(e^{2px}a+b)^2} de^{px}$$

↓ 241

$$\frac{1}{2ap(ae^{2px} + b)}$$

input `Int[(b/E^(p*x) + a*E^(p*x))^(-2), x]`

output `-1/2*1/(a*(b + a*E^(2*p*x))*p)`

Defintions of rubi rules used

rule 241 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(a + b*x^2)^(p + 1)/(2*b*(p + 1)), x] /; FreeQ[{a, b, p}, x] && NeQ[p, -1]`

rule 2720 `Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Simp[v/D[v, x] Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x)) *(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

method	result	size
risch	$-\frac{1}{2a(b+ae^{2px})p}$	20
derivativedivides	$-\frac{1}{2a(b+ae^{2px})p}$	21
default	$-\frac{1}{2a(b+ae^{2px})p}$	21
norman	$-\frac{1}{2a(b+ae^{2px})p}$	21
parallelrisch	$-\frac{1}{2a(b+ae^{2px})p}$	21

input `int(1/(b/exp(p*x)+a*exp(p*x))^2,x,method=_RETURNVERBOSE)`

output `-1/2/a/(b+a*exp(2*p*x))/p`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = -\frac{1}{2(a^2pe^{(2px)} + abp)}$$

input `integrate(1/(b/exp(p*x)+a*exp(p*x))^2,x, algorithm="fricas")`

output `-1/2/(a^2*p*e^(2*p*x) + a*b*p)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = \frac{1}{2abp + 2b^2pe^{-2px}}$$

input `integrate(1/(b/exp(p*x)+a*exp(p*x))**2,x)`output `1/(2*a*b*p + 2*b**2*p*exp(-2*p*x))`**Maxima [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.91

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = \frac{1}{2(b^2e^{(-2px)} + ab)p}$$

input `integrate(1/(b/exp(p*x)+a*exp(p*x))^2,x, algorithm="maxima")`output `1/2/((b^2*e^(-2*p*x) + a*b)*p)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = -\frac{1}{2(ae^{(2px)} + b)ap}$$

input `integrate(1/(b/exp(p*x)+a*exp(p*x))^2,x, algorithm="giac")`output `-1/2/((a*e^(2*p*x) + b)*a*p)`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = \frac{e^{2px}}{2bp(b + ae^{2px})}$$

input `int(1/(a*exp(p*x) + b*exp(-p*x))^2,x)`output `exp(2*p*x)/(2*b*p*(b + a*exp(2*p*x)))`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.18

$$\int \frac{1}{(be^{-px} + ae^{px})^2} dx = \frac{e^{2px}}{2bp(e^{2px}a + b)}$$

input `int(1/(b/exp(p*x)+a*exp(p*x))^2,x)`output `e**(2*p*x)/(2*b*p*(e**(2*p*x)*a + b))`

3.23 $\int \frac{x}{(be^{-px} + ae^{px})^2} dx$

Optimal result	203
Mathematica [A] (verified)	203
Rubi [A] (verified)	204
Maple [A] (verified)	206
Fricas [A] (verification not implemented)	206
Sympy [A] (verification not implemented)	207
Maxima [A] (verification not implemented)	207
Giac [A] (verification not implemented)	207
Mupad [B] (verification not implemented)	208
Reduce [B] (verification not implemented)	208

Optimal result

Integrand size = 20, antiderivative size = 62

$$\int \frac{x}{(be^{-px} + ae^{px})^2} dx = \frac{x}{2abp} - \frac{x}{2a(b + ae^{2px})p} - \frac{\log(b + ae^{2px})}{4abp^2}$$

output `1/2*x/a/b/p-1/2*x/a/(b+a*exp(2*p*x))/p-1/4*ln(b+a*exp(2*p*x))/a/b/p^2`

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.79

$$\int \frac{x}{(be^{-px} + ae^{px})^2} dx = \frac{\frac{2e^{2px} px}{b+ae^{2px}} - \frac{\log(b+ae^{2px})}{a}}{4bp^2}$$

input `Integrate[x/(b/E^(p*x) + a*E^(p*x))^2,x]`

output `((2*E^(2*p*x)*p*x)/(b + a*E^(2*p*x)) - Log[b + a*E^(2*p*x)])/a/(4*b*p^2)`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {2721, 2621, 2720, 47, 14, 16}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{(ae^{px} + be^{-px})^2} dx \\
 & \quad \downarrow \text{2721} \\
 & \int \frac{xe^{2px}}{(ae^{2px} + b)^2} dx \\
 & \quad \downarrow \text{2621} \\
 & \frac{\int \frac{1}{e^{2px}a+b} dx}{2ap} - \frac{x}{2ap(ae^{2px} + b)} \\
 & \quad \downarrow \text{2720} \\
 & \frac{\int \frac{e^{-2px}}{e^{2px}a+b} de^{2px}}{4ap^2} - \frac{x}{2ap(ae^{2px} + b)} \\
 & \quad \downarrow \text{47} \\
 & \frac{\int e^{-2px} de^{2px}}{b} - \frac{a \int \frac{1}{e^{2px}a+b} de^{2px}}{b} - \frac{x}{2ap(ae^{2px} + b)} \\
 & \quad \downarrow \text{14} \\
 & \frac{\log(e^{2px})}{b} - \frac{a \int \frac{1}{e^{2px}a+b} de^{2px}}{b} - \frac{x}{2ap(ae^{2px} + b)} \\
 & \quad \downarrow \text{16} \\
 & \frac{\log(e^{2px})}{b} - \frac{\log(ae^{2px}+b)}{b} - \frac{x}{2ap(ae^{2px} + b)}
 \end{aligned}$$

input `Int [x/(b/E^(p*x) + a*E^(p*x))^2, x]`

output
$$-1/2*x/(a*(b + a*E^{(2*p*x)})^p) + (\text{Log}[E^{(2*p*x)}]/b - \text{Log}[b + a*E^{(2*p*x)}])/b)/(4*a*p^2)$$

Defintions of rubi rules used

rule 14
$$\text{Int}[(a_)/(x_), x_Symbol] \rightarrow \text{Simp}[a*\text{Log}[x], x] \text{ ; FreeQ}[a, x]$$

rule 16
$$\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] \text{ ; FreeQ}[\{a, b, c\}, x]$$

rule 47
$$\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[1/(a + b*x), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[1/(c + d*x), x], x] \text{ ; FreeQ}[\{a, b, c, d\}, x]$$

rule 2621
$$\text{Int}[(F_)^{(g_)*((e_)+(f_)*(x_)))^{(n_)*((a_)+(b_)*(F_)^{(g_)*((e_)+(f_)*(x_)))^{(n_)}}^{(p_)*((c_)+(d_)*(x_))^{(m_)}}, x_Symbol] \rightarrow \text{Simp}[(c + d*x)^m*((a + b*(F^{(g*(e + f*x)))^n)^{(p + 1)})/(b*f*g*n*(p + 1)*\text{Log}[F])), x] - \text{Simp}[d*(m/(b*f*g*n*(p + 1)*\text{Log}[F])) \text{ Int}[(c + d*x)^{(m - 1)}*(a + b*(F^{(g*(e + f*x)))^n})^{(p + 1)}, x], x] \text{ ; FreeQ}[\{F, a, b, c, d, e, f, g, m, n, p\}, x] \ \&\& \ \text{NeQ}[p, -1]$$

rule 2720
$$\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Simp}[v/D[v, x] \text{ Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] \text{ ; FunctionOfExponentialQ}[u, x] \ \&\& \ \text{!MatchQ}[u, (w_)*((a_)*(v_)^{(n_)})^{(m_)} \text{ ; FreeQ}[\{a, m, n\}, x] \ \&\& \ \text{IntegerQ}[m*n] \ \&\& \ \text{!MatchQ}[u, E^{(c_)*((a_)+(b_)*x)}*(F_)[v_] \text{ ; FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{InverseFunctionQ}[F[x]]]$$

rule 2721
$$\text{Int}[(u_)*((a_)*(F_)^{(v_)+(b_)*(F_)^{(w_)})^{(n_)}}, x_Symbol] \rightarrow \text{Int}[u*F^{(n*v)}*(a + b*F^{\text{ExpandToSum}[w - v, x]})^n, x] \text{ ; FreeQ}[\{F, a, b, n\}, x] \ \&\& \ \text{ILtQ}[n, 0] \ \&\& \ \text{LinearQ}[\{v, w\}, x]$$

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{-\frac{\ln(b+a e^{2px})}{4ba} + \frac{px e^{2px}}{2b(b+a e^{2px})}}{p^2}$	50
default	$\frac{-\frac{\ln(b+a e^{2px})}{4ba} + \frac{px e^{2px}}{2b(b+a e^{2px})}}{p^2}$	50
norman	$\frac{x e^{2px}}{2bp(b+a e^{2px})} - \frac{\ln(b+a e^{2px})}{4ab p^2}$	51
risch	$\frac{x}{2abp} - \frac{x}{2a(b+a e^{2px})p} - \frac{\ln(e^{2px} + \frac{b}{a})}{4ab p^2}$	57
parallelrisc	$-\frac{2 e^{2px} apx + \ln(b+a e^{2px})e^{2px} a + \ln(b+a e^{2px})b}{4ab p^2(b+a e^{2px})}$	68

input `int(x/(b/exp(p*x)+a*exp(p*x))^2,x,method=_RETURNVERBOSE)`

output `1/p^2*(-1/4/b/a*ln(a*exp(p*x)^2+b)+1/2*p*x*exp(p*x)^2/b/(a*exp(p*x)^2+b))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \frac{x}{(be^{-px} + ae^{px})^2} dx = \frac{2apxe^{(2px)} - (ae^{(2px)} + b) \log(ae^{(2px)} + b)}{4(a^2bp^2e^{(2px)} + ab^2p^2)}$$

input `integrate(x/(b/exp(p*x)+a*exp(p*x))^2,x, algorithm="fricas")`

output `1/4*(2*a*p*x*e^(2*p*x) - (a*e^(2*p*x) + b)*log(a*e^(2*p*x) + b))/(a^2*b*p^2*e^(2*p*x) + a*b^2*p^2)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.82

$$\int \frac{x}{(be^{-px} + ae^{px})^2} dx = \frac{x}{2abp + 2b^2pe^{-2px}} - \frac{x}{2abp} - \frac{\log\left(\frac{a}{b} + e^{-2px}\right)}{4abp^2}$$

input `integrate(x/(b/exp(p*x)+a*exp(p*x))**2,x)`output `x/(2*a*b*p + 2*b**2*p*exp(-2*p*x)) - x/(2*a*b*p) - log(a/b + exp(-2*p*x))/(4*a*b*p**2)`**Maxima [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.82

$$\int \frac{x}{(be^{-px} + ae^{px})^2} dx = \frac{xe^{(2px)}}{2(abpe^{(2px)} + b^2p)} - \frac{\log\left(\frac{ae^{(2px)}+b}{a}\right)}{4abp^2}$$

input `integrate(x/(b/exp(p*x)+a*exp(p*x))^2,x, algorithm="maxima")`output `1/2*x*e^(2*p*x)/(a*b*p*e^(2*p*x) + b^2*p) - 1/4*log((a*e^(2*p*x) + b)/a)/(a*b*p^2)`**Giac [A] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.19

$$\int \frac{x}{(be^{-px} + ae^{px})^2} dx = \frac{2apxe^{(2px)} - ae^{(2px)} \log(-ae^{(2px)} - b) - b \log(-ae^{(2px)} - b)}{4(a^2bp^2e^{(2px)} + ab^2p^2)}$$

input `integrate(x/(b/exp(p*x)+a*exp(p*x))^2,x, algorithm="giac")`

output $\frac{1}{4} \cdot (2 \cdot a \cdot p \cdot x \cdot e^{(2 \cdot p \cdot x)} - a \cdot e^{(2 \cdot p \cdot x)} \cdot \log(-a \cdot e^{(2 \cdot p \cdot x)} - b) - b \cdot \log(-a \cdot e^{(2 \cdot p \cdot x)} - b)) / (a^2 \cdot b \cdot p^2 \cdot e^{(2 \cdot p \cdot x)} + a \cdot b^2 \cdot p^2)$

Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.76

$$\int \frac{x}{(b e^{-p x} + a e^{p x})^2} dx = \frac{x e^{2 p x}}{2 b p (b + a e^{2 p x})} - \frac{\ln(b + a e^{2 p x})}{4 a b p^2}$$

input `int(x/(a*exp(p*x) + b*exp(-p*x))^2,x)`

output $(x \cdot \exp(2 \cdot p \cdot x)) / (2 \cdot b \cdot p \cdot (b + a \cdot \exp(2 \cdot p \cdot x))) - \log(b + a \cdot \exp(2 \cdot p \cdot x)) / (4 \cdot a \cdot b \cdot p^2)$

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.11

$$\int \frac{x}{(b e^{-p x} + a e^{p x})^2} dx = \frac{-e^{2 p x} \log(e^{2 p x} a + b) a + 2 e^{2 p x} a p x - \log(e^{2 p x} a + b) b}{4 a b p^2 (e^{2 p x} a + b)}$$

input `int(x/(b/exp(p*x)+a*exp(p*x))^2,x)`

output $(-e^{(2 \cdot p \cdot x)} \cdot \log(e^{(2 \cdot p \cdot x)} \cdot a + b) \cdot a + 2 \cdot e^{(2 \cdot p \cdot x)} \cdot a \cdot p \cdot x - \log(e^{(2 \cdot p \cdot x)} \cdot a + b) \cdot b) / (4 \cdot a \cdot b \cdot p^2 \cdot (e^{(2 \cdot p \cdot x)} \cdot a + b))$

3.24 $\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx$

Optimal result	209
Mathematica [C] (verified)	210
Rubi [A] (verified)	210
Maple [B] (verified)	213
Fricas [B] (verification not implemented)	214
Sympy [F]	215
Maxima [F]	215
Giac [B] (verification not implemented)	216
Mupad [F(-1)]	217
Reduce [F]	217

Optimal result

Integrand size = 31, antiderivative size = 86

$$\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx = \frac{(1+x)\sqrt{1-x+x^2}}{1+x+x^2} + \sqrt{2} \arctan\left(\frac{\sqrt{2}(1+x)}{\sqrt{1-x+x^2}}\right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{\frac{2}{3}}(1-x)}{\sqrt{1-x+x^2}}\right)}{\sqrt{6}}$$

output

```
arctan((1+x)*2^(1/2)/(x^2-x+1)^(1/2))*2^(1/2)-1/6*arctanh(1/3*(1-x)*6^(1/2)
)/(x^2-x+1)^(1/2))*6^(1/2)+(1+x)*(x^2-x+1)^(1/2)/(x^2+x+1)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.30 (sec) , antiderivative size = 239, normalized size of antiderivative = 2.78

$$\int \frac{1 - x + 3x^2}{\sqrt{1 - x + x^2} (1 + x + x^2)^2} dx$$

$$= \frac{(1+x)\sqrt{1-x+x^2}}{1+x+x^2} - \text{RootSum} \left[3 + 6\#1 + \#1^2 - 2\#1^3 \right. \\ \left. + \#1^4 \&, \frac{19 \log(-x + \sqrt{1-x+x^2} - \#1) + 6 \log(-x + \sqrt{1-x+x^2} - \#1) \#1}{3 + \#1 - 3\#1^2 + 2\#1^3} \& \right] \\ - \frac{1}{2} \text{RootSum} \left[3 + 6\#1 + \#1^2 - 2\#1^3 \right. \\ \left. + \#1^4 \&, \frac{-36 \log(-x + \sqrt{1-x+x^2} - \#1) - 6 \log(-x + \sqrt{1-x+x^2} - \#1) \#1 + \log(-x + \sqrt{1-x+x^2} - \#1) \#1^2}{3 + \#1 - 3\#1^2 + 2\#1^3} \& \right]$$

input `Integrate[(1 - x + 3*x^2)/(Sqrt[1 - x + x^2]*(1 + x + x^2)^2), x]`

output `((1 + x)*Sqrt[1 - x + x^2])/(1 + x + x^2) - RootSum[3 + 6*#1 + #1^2 - 2*#1^3 + #1^4 & , (19*Log[-x + Sqrt[1 - x + x^2] - #1] + 6*Log[-x + Sqrt[1 - x + x^2] - #1]*#1)/(3 + #1 - 3*#1^2 + 2*#1^3) &] - RootSum[3 + 6*#1 + #1^2 - 2*#1^3 + #1^4 & , (-36*Log[-x + Sqrt[1 - x + x^2] - #1] - 6*Log[-x + Sqrt[1 - x + x^2] - #1]*#1 + Log[-x + Sqrt[1 - x + x^2] - #1]*#1^2)/(3 + #1 - 3*#1^2 + 2*#1^3) &]/2`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.09, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {2135, 27, 1368, 27, 1362, 217, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{3x^2 - x + 1}{\sqrt{x^2 - x + 1}(x^2 + x + 1)^2} dx \\
& \quad \downarrow \text{2135} \\
& \frac{1}{12} \int \frac{6(3-x)}{\sqrt{x^2 - x + 1}(x^2 + x + 1)} dx + \frac{\sqrt{x^2 - x + 1}(x+1)}{x^2 + x + 1} \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \int \frac{3-x}{\sqrt{x^2 - x + 1}(x^2 + x + 1)} dx + \frac{\sqrt{x^2 - x + 1}(x+1)}{x^2 + x + 1} \\
& \quad \downarrow \text{1368} \\
& \frac{1}{2} \left(\frac{1}{4} \int \frac{4(x+1)}{\sqrt{x^2 - x + 1}(x^2 + x + 1)} dx - \frac{1}{4} \int -\frac{8(1-x)}{\sqrt{x^2 - x + 1}(x^2 + x + 1)} dx \right) + \\
& \quad \frac{\sqrt{x^2 - x + 1}(x+1)}{x^2 + x + 1} \\
& \quad \downarrow \text{27} \\
& \frac{1}{2} \left(2 \int \frac{1-x}{\sqrt{x^2 - x + 1}(x^2 + x + 1)} dx + \int \frac{x+1}{\sqrt{x^2 - x + 1}(x^2 + x + 1)} dx \right) + \frac{\sqrt{x^2 - x + 1}(x+1)}{x^2 + x + 1} \\
& \quad \downarrow \text{1362} \\
& \frac{1}{2} \left(2 \int \frac{1}{3 - \frac{2(1-x)^2}{x^2 - x + 1}} d\left(-\frac{1-x}{\sqrt{x^2 - x + 1}}\right) - 12 \int \frac{1}{-\frac{18(x+1)^2}{x^2 - x + 1} - 9} d\frac{3(x+1)}{\sqrt{x^2 - x + 1}} \right) + \\
& \quad \frac{\sqrt{x^2 - x + 1}(x+1)}{x^2 + x + 1} \\
& \quad \downarrow \text{217} \\
& \frac{1}{2} \left(2 \int \frac{1}{3 - \frac{2(1-x)^2}{x^2 - x + 1}} d\left(-\frac{1-x}{\sqrt{x^2 - x + 1}}\right) + 2\sqrt{2} \arctan\left(\frac{\sqrt{2}(x+1)}{\sqrt{x^2 - x + 1}}\right) \right) + \frac{\sqrt{x^2 - x + 1}(x+1)}{x^2 + x + 1} \\
& \quad \downarrow \text{219} \\
& \frac{1}{2} \left(2\sqrt{2} \arctan\left(\frac{\sqrt{2}(x+1)}{\sqrt{x^2 - x + 1}}\right) - \sqrt{\frac{2}{3}} \operatorname{arctanh}\left(\frac{\sqrt{\frac{2}{3}}(1-x)}{\sqrt{x^2 - x + 1}}\right) \right) + \frac{\sqrt{x^2 - x + 1}(x+1)}{x^2 + x + 1}
\end{aligned}$$

input `Int[(1 - x + 3*x^2)/(Sqrt[1 - x + x^2]*(1 + x + x^2)^2),x]`

output
$$\frac{((1+x)\sqrt{1-x+x^2})/(1+x+x^2) + (2\sqrt{2}\operatorname{ArcTan}[(\sqrt{2}(1+x))/\sqrt{1-x+x^2}]) - \sqrt{2/3}\operatorname{ArcTanh}[(\sqrt{2/3}(1-x))/\sqrt{1-x+x^2}])}{2}$$

Defintions of rubi rules used

rule 27
$$\operatorname{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \operatorname{Simp}[a \operatorname{Int}[Fx, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[Fx, (b_*)(Gx_)] /; \operatorname{FreeQ}[b, x]$$

rule 217
$$\operatorname{Int}[((a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[-b, 2])^{(-1)}*\operatorname{ArcTan}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{PosQ}[a/b] \ \&\& \ (\operatorname{LtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$$

rule 219
$$\operatorname{Int}[((a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$$

rule 1362
$$\operatorname{Int}[((g_*) + (h_*)(x_))/(((a_*) + (b_*)(x_) + (c_*)(x_)^2)*\sqrt{(d_*) + (e_*)(x_) + (f_*)(x_)^2})], x_Symbol] \rightarrow \operatorname{Simp}[-2*g*(g*b - 2*a*h) \operatorname{Subst}[\operatorname{Int}[1/\operatorname{Simp}[g*(g*b - 2*a*h)*(b^2 - 4*a*c) - (b*d - a*e)*x^2, x], x], x, \operatorname{Simp}[g*b - 2*a*h - (b*h - 2*g*c)*x, x]/\sqrt{d + e*x + f*x^2}], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, f, g, h\}, x] \ \&\& \ \operatorname{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \operatorname{NeQ}[e^2 - 4*d*f, 0] \ \&\& \ \operatorname{NeQ}[b*d - a*e, 0] \ \&\& \ \operatorname{EqQ}[h^2*(b*d - a*e) - 2*g*h*(c*d - a*f) + g^2*(c*e - b*f), 0]$$

rule 1368
$$\operatorname{Int}[((g_*) + (h_*)(x_))/(((a_*) + (b_*)(x_) + (c_*)(x_)^2)*\sqrt{(d_*) + (e_*)(x_) + (f_*)(x_)^2})], x_Symbol] \rightarrow \operatorname{With}[\{q = \operatorname{Rt}[(c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f)], 2\}, \operatorname{Simp}[1/(2*q) \operatorname{Int}[\operatorname{Simp}[h*(b*d - a*e) - g*(c*d - a*f - q) - (g*(c*e - b*f) - h*(c*d - a*f + q))*x, x]/((a + b*x + c*x^2)*\sqrt{d + e*x + f*x^2}), x], x] - \operatorname{Simp}[1/(2*q) \operatorname{Int}[\operatorname{Simp}[h*(b*d - a*e) - g*(c*d - a*f + q) - (g*(c*e - b*f) - h*(c*d - a*f - q))*x, x]/((a + b*x + c*x^2)*\sqrt{d + e*x + f*x^2}), x], x]] /; \operatorname{FreeQ}[\{a, b, c, d, e, f, g, h\}, x] \ \&\& \ \operatorname{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \operatorname{NeQ}[e^2 - 4*d*f, 0] \ \&\& \ \operatorname{NeQ}[b*d - a*e, 0] \ \&\& \ \operatorname{NegQ}[b^2 - 4*a*c]$$

rule 2135

```

Int[(Px_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_)*((d_) + (e_)*(x_) + (f_
)*(x_)^2)^(q_), x_Symbol] :=> With[{A = Coeff[Px, x, 0], B = Coeff[Px, x, 1]
, C = Coeff[Px, x, 2]}, Simp[(a + b*x + c*x^2)^(p + 1)*((d + e*x + f*x^2)^(
q + 1)/((b^2 - 4*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1))*(
(A*c - a*C)*(2*a*c*e - b*(c*d + a*f)) + (A*b - a*B)*(2*c^2*d + b^2*f - c*(b
*e + 2*a*f)) + c*(A*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)) - B*(b*c*d - 2*a*c*
e + a*b*f) + C*(b^2*d - a*b*e - 2*a*(c*d - a*f)))*x), x] + Simp[1/((b^2 - 4
*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1)) Int[(a + b*x + c
*x^2)^(p + 1)*(d + e*x + f*x^2)^q*Simp[(b*B - 2*A*c - 2*a*C)*((c*d - a*f)^2
- (b*d - a*e)*(c*e - b*f))*(p + 1) + (b^2*(C*d + A*f) - b*(B*c*d + A*c*e +
a*C*e + a*B*f) + 2*(A*c*(c*d - a*f) - a*(c*C*d - B*c*e - a*C*f)))*(a*f*(p
+ 1) - c*d*(p + 2)) - e*((A*c - a*C)*(2*a*c*e - b*(c*d + a*f)) + (A*b - a*B
)*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)))*(p + q + 2) - (2*f*((A*c - a*C)*(2*a
*c*e - b*(c*d + a*f)) + (A*b - a*B)*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)))*(p
+ q + 2) - (b^2*(C*d + A*f) - b*(B*c*d + A*c*e + a*C*e + a*B*f) + 2*(A*c*(
c*d - a*f) - a*(c*C*d - B*c*e - a*C*f)))*(b*f*(p + 1) - c*e*(2*p + q + 4))
*x - c*f*(b^2*(C*d + A*f) - b*(B*c*d + A*c*e + a*C*e + a*B*f) + 2*(A*c*(c*d
- a*f) - a*(c*C*d - B*c*e - a*C*f)))*(2*p + 2*q + 5)*x^2, x], x]] /; F
reeQ[{a, b, c, d, e, f, q}, x] && PolyQ[Px, x, 2] && LtQ[p, -1] && NeQ[(c*d
- a*f)^2 - (b*d - a*e)*(c*e - b*f), 0] && !( !IntegerQ[p] && !LtQ[q, -1])
&& !IGtQ[q, 0]

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 157 vs. $2(71) = 142$.

Time = 1.26 (sec) , antiderivative size = 158, normalized size of antiderivative = 1.84

method	result
risch	$\frac{(1+x)\sqrt{x^2-x+1}}{x^2+x+1} + \frac{\sqrt{\frac{(1+x)^2}{(1-x)^2}+3} \left(6\sqrt{2} \arctan\left(\frac{2(1+x)\sqrt{2}}{(1-x)\sqrt{\frac{(1+x)^2}{(1-x)^2}+3}}\right) - \sqrt{6} \operatorname{arctanh}\left(\frac{\sqrt{\frac{(1+x)^2}{(1-x)^2}+3}\sqrt{6}}{4}\right) \right)}{6\sqrt{\frac{(1+x)^2}{(1-x)^2}+3} \left(\frac{1+x}{1-x}+1\right)}$
default	$\frac{9\sqrt{2} \arctan\left(\frac{2(1+x)\sqrt{2}}{(1-x)\sqrt{\frac{(1+x)^2}{(1-x)^2}+3}}\right) \sqrt{\frac{(1+x)^2}{(1-x)^2}+3} (1+x)^2}{(1-x)^2} - \frac{6 \operatorname{arctanh}\left(\frac{\sqrt{\frac{(1+x)^2}{(1-x)^2}+3}\sqrt{6}}{4}\right) \sqrt{\frac{(1+x)^2}{(1-x)^2}+3}\sqrt{6} (1+x)^2}{(1-x)^2} + 3\sqrt{2} \arctan\left(\frac{2(1+x)}{(1-x)\sqrt{\frac{(1+x)^2}{(1-x)^2}+3}}\right)$
trager	$\frac{(1+x)\sqrt{x^2-x+1}}{x^2+x+1} - \operatorname{RootOf}(576_Z^4 + 528_Z^2 + 169) \ln\left(\frac{3456x \operatorname{RootOf}(576_Z^4 + 528_Z^2 + 169)^5 + 6312x R}{\dots}\right)$

```
input int((3*x^2-x+1)/(x^2+x+1)^2/(x^2-x+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
output (1+x)*(x^2-x+1)^(1/2)/(x^2+x+1)+1/6*((1+x)^2/(1-x)^2+3)^(1/2)*(6*2^(1/2)*arctan(2*(1+x)/(1-x)/((1+x)^2/(1-x)^2+3)^(1/2)*2^(1/2))-6^(1/2)*arctanh(1/4*((1+x)^2/(1-x)^2+3)^(1/2)*6^(1/2)))/(((1+x)^2/(1-x)^2+3)/((1+x)/(1-x)+1))^2)^(1/2)/((1+x)/(1-x)+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 245 vs. 2(69) = 138.

Time = 0.08 (sec) , antiderivative size = 245, normalized size of antiderivative = 2.85

$$\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx$$

$$= \frac{12\sqrt{2}(x^2+x+1) \arctan\left(-\frac{1}{2}\sqrt{6}\sqrt{2}(2x-1) + \sqrt{x^2-x+1}(\sqrt{6}\sqrt{2}+2\sqrt{2}) - 2\sqrt{2}(x-1)\right) - 12\sqrt{2}}{\dots}$$

```
input integrate((3*x^2-x+1)/(x^2+x+1)^2/(x^2-x+1)^(1/2),x, algorithm="fricas")
```

output

```
1/12*(12*sqrt(2)*(x^2 + x + 1)*arctan(-1/2*sqrt(6)*sqrt(2)*(2*x - 1) + sqrt(x^2 - x + 1)*(sqrt(6)*sqrt(2) + 2*sqrt(2)) - 2*sqrt(2)*(x - 1)) - 12*sqrt(2)*(x^2 + x + 1)*arctan(-1/2*sqrt(6)*sqrt(2)*(2*x - 1) + sqrt(x^2 - x + 1)*(sqrt(6)*sqrt(2) - 2*sqrt(2)) + 2*sqrt(2)*(x - 1)) + sqrt(6)*(x^2 + x + 1)*log(2*x^2 - sqrt(x^2 - x + 1)*(2*x + sqrt(6) + 1) + sqrt(6)*(x + 1) + 4) - sqrt(6)*(x^2 + x + 1)*log(2*x^2 - sqrt(x^2 - x + 1)*(2*x - sqrt(6) + 1) - sqrt(6)*(x + 1) + 4) + 12*x^2 + 12*sqrt(x^2 - x + 1)*(x + 1) + 12*x + 12)/(x^2 + x + 1)
```

Sympy [F]

$$\int \frac{1 - x + 3x^2}{\sqrt{1 - x + x^2} (1 + x + x^2)^2} dx = \int \frac{3x^2 - x + 1}{\sqrt{x^2 - x + 1} (x^2 + x + 1)^2} dx$$

input

```
integrate((3*x**2-x+1)/(x**2+x+1)**2/(x**2-x+1)**(1/2),x)
```

output

```
Integral((3*x**2 - x + 1)/(sqrt(x**2 - x + 1)*(x**2 + x + 1)**2), x)
```

Maxima [F]

$$\int \frac{1 - x + 3x^2}{\sqrt{1 - x + x^2} (1 + x + x^2)^2} dx = \int \frac{3x^2 - x + 1}{(x^2 + x + 1)^2 \sqrt{x^2 - x + 1}} dx$$

input

```
integrate((3*x^2-x+1)/(x^2+x+1)^2/(x^2-x+1)^(1/2),x, algorithm="maxima")
```

output

```
integrate((3*x^2 - x + 1)/((x^2 + x + 1)^2*sqrt(x^2 - x + 1)), x)
```


Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(69) = 138$.

Time = 0.14 (sec) , antiderivative size = 304, normalized size of antiderivative = 3.53

$$\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx$$

$$= -\frac{1}{3}\sqrt{6}\sqrt{3}\arctan\left(-\frac{2x+\sqrt{6}-2\sqrt{x^2-x+1}+1}{\sqrt{3}+\sqrt{2}}\right)$$

$$+ \frac{1}{3}\sqrt{6}\sqrt{3}\arctan\left(-\frac{2x-\sqrt{6}-2\sqrt{x^2-x+1}+1}{\sqrt{3}-\sqrt{2}}\right)$$

$$+ \frac{1}{12}\sqrt{6}\log\left(4\left(\sqrt{6}\sqrt{3}+3\sqrt{3}\right)^2+36\left(2x+\sqrt{6}-2\sqrt{x^2-x+1}+1\right)^2\right)$$

$$- \frac{1}{12}\sqrt{6}\log\left(4\left(\sqrt{6}\sqrt{3}-3\sqrt{3}\right)^2+36\left(2x-\sqrt{6}-2\sqrt{x^2-x+1}+1\right)^2\right)$$

$$+ \frac{(x-\sqrt{x^2-x+1})^3+4(x-\sqrt{x^2-x+1})^2-10x+10\sqrt{x^2-x+1}+5}{(x-\sqrt{x^2-x+1})^4+2(x-\sqrt{x^2-x+1})^3+(x-\sqrt{x^2-x+1})^2-6x+6\sqrt{x^2-x+1}+3}$$

input `integrate((3*x^2-x+1)/(x^2+x+1)^2/(x^2-x+1)^(1/2),x, algorithm="giac")`

output `-1/3*sqrt(6)*sqrt(3)*arctan(-(2*x + sqrt(6) - 2*sqrt(x^2 - x + 1) + 1)/(sqrt(3) + sqrt(2))) + 1/3*sqrt(6)*sqrt(3)*arctan(-(2*x - sqrt(6) - 2*sqrt(x^2 - x + 1) + 1)/(sqrt(3) - sqrt(2))) + 1/12*sqrt(6)*log(4*(sqrt(6)*sqrt(3) + 3*sqrt(3))^2 + 36*(2*x + sqrt(6) - 2*sqrt(x^2 - x + 1) + 1)^2) - 1/12*sqrt(6)*log(4*(sqrt(6)*sqrt(3) - 3*sqrt(3))^2 + 36*(2*x - sqrt(6) - 2*sqrt(x^2 - x + 1) + 1)^2) + ((x - sqrt(x^2 - x + 1))^3 + 4*(x - sqrt(x^2 - x + 1))^2 - 10*x + 10*sqrt(x^2 - x + 1) + 5)/((x - sqrt(x^2 - x + 1))^4 + 2*(x - sqrt(x^2 - x + 1))^3 + (x - sqrt(x^2 - x + 1))^2 - 6*x + 6*sqrt(x^2 - x + 1) + 3)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx = \int \frac{3x^2-x+1}{\sqrt{x^2-x+1}(x^2+x+1)^2} dx$$

input `int((3*x^2 - x + 1)/((x^2 - x + 1)^(1/2)*(x + x^2 + 1)^2), x)`

output `int((3*x^2 - x + 1)/((x^2 - x + 1)^(1/2)*(x + x^2 + 1)^2), x)`

Reduce [F]

$$\int \frac{1-x+3x^2}{\sqrt{1-x+x^2}(1+x+x^2)^2} dx$$

$$= \frac{4\sqrt{x^2-x+1}x + 6\sqrt{x^2-x+1} + 14\left(\int \frac{\sqrt{x^2-x+1}}{x^6+x^5+2x^4+x^3+2x^2+x+1} dx\right) x^2 + 14\left(\int \frac{\sqrt{x^2-x+1}}{x^6+x^5+2x^4+x^3+2x^2+x+1} dx\right) x}{1}$$

input `int((3*x^2-x+1)/(x^2+x+1)^2/(x^2-x+1)^(1/2), x)`

output `(2*(2*sqrt(x**2 - x + 1))*x + 3*sqrt(x**2 - x + 1) + 7*int(sqrt(x**2 - x + 1)/(x**6 + x**5 + 2*x**4 + x**3 + 2*x**2 + x + 1),x)*x**2 + 7*int(sqrt(x**2 - x + 1)/(x**6 + x**5 + 2*x**4 + x**3 + 2*x**2 + x + 1),x)*x + 7*int(sqrt(x**2 - x + 1)/(x**6 + x**5 + 2*x**4 + x**3 + 2*x**2 + x + 1),x) + 8*int((sqrt(x**2 - x + 1)*x**2)/(x**6 + x**5 + 2*x**4 + x**3 + 2*x**2 + x + 1),x)*x**2 + 8*int((sqrt(x**2 - x + 1)*x**2)/(x**6 + x**5 + 2*x**4 + x**3 + 2*x**2 + x + 1),x)*x + 8*int((sqrt(x**2 - x + 1)*x**2)/(x**6 + x**5 + 2*x**4 + x**3 + 2*x**2 + x + 1),x))/ (9*(x**2 + x + 1))`

3.25 $\int \frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a^2+x^2}} dx$

Optimal result	218
Mathematica [A] (verified)	218
Rubi [A] (verified)	219
Maple [A] (verified)	220
Fricas [A] (verification not implemented)	220
Sympy [A] (verification not implemented)	220
Maxima [F]	221
Giac [A] (verification not implemented)	221
Mupad [B] (verification not implemented)	221
Reduce [B] (verification not implemented)	222

Optimal result

Integrand size = 29, antiderivative size = 19

$$\int \frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a^2+x^2}} dx = 2\sqrt{x+\sqrt{a^2+x^2}}$$

output `2*(x+(a^2+x^2)^(1/2))^(1/2)`

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a^2+x^2}} dx = 2\sqrt{x+\sqrt{a^2+x^2}}$$

input `Integrate[Sqrt[x + Sqrt[a^2 + x^2]]/Sqrt[a^2 + x^2],x]`

output `2*Sqrt[x + Sqrt[a^2 + x^2]]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {2547, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sqrt{a^2 + x^2} + x}}{\sqrt{a^2 + x^2}} dx$$

↓ 2547

$$\int \frac{1}{\sqrt{\sqrt{a^2 + x^2} + x}} d(\sqrt{a^2 + x^2} + x)$$

↓ 15

$$2\sqrt{\sqrt{a^2 + x^2} + x}$$

input `Int[Sqrt[x + Sqrt[a^2 + x^2]]/Sqrt[a^2 + x^2],x]`

output `2*Sqrt[x + Sqrt[a^2 + x^2]]`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

method	result	size
derivativedivides	$2\sqrt{x + \sqrt{a^2 + x^2}}$	16
default	$2\sqrt{x + \sqrt{a^2 + x^2}}$	16

input `int((x+(a^2+x^2)^(1/2))^(1/2)/(a^2+x^2)^(1/2),x,method=_RETURNVERBOSE)`

output `2*(x+(a^2+x^2)^(1/2))^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a^2 + x^2}} dx = 2\sqrt{x + \sqrt{a^2 + x^2}}$$

input `integrate((x+(a^2+x^2)^(1/2))^(1/2)/(a^2+x^2)^(1/2),x, algorithm="fricas")`

output `2*sqrt(x + sqrt(a^2 + x^2))`

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a^2 + x^2}} dx = 2\sqrt{x + \sqrt{a^2 + x^2}}$$

input `integrate((x+(a**2+x**2)**(1/2))**(1/2)/(a**2+x**2)**(1/2),x)`

output `2*sqrt(x + sqrt(a**2 + x**2))`

Maxima [F]

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a^2 + x^2}} dx = \int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a^2 + x^2}} dx$$

input `integrate((x+(a^2+x^2)^(1/2))^(1/2)/(a^2+x^2)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(x + sqrt(a^2 + x^2))/sqrt(a^2 + x^2), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a^2 + x^2}} dx = 2 \sqrt{x + \sqrt{a^2 + x^2}}$$

input `integrate((x+(a^2+x^2)^(1/2))^(1/2)/(a^2+x^2)^(1/2),x, algorithm="giac")`

output `2*sqrt(x + sqrt(a^2 + x^2))`

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a^2 + x^2}} dx = 2 \sqrt{x + \sqrt{a^2 + x^2}}$$

input `int((x + (a^2 + x^2)^(1/2))^(1/2)/(a^2 + x^2)^(1/2),x)`

output `2*(x + (a^2 + x^2)^(1/2))^(1/2)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.68

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a^2 + x^2}} dx = 2\sqrt{\sqrt{a^2 + x^2} + x}$$

input `int((x+(a^2+x^2)^(1/2))^(1/2)/(a^2+x^2)^(1/2),x)`

output `2*sqrt(sqrt(a**2 + x**2) + x)`

$$3.26 \quad \int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx$$

Optimal result	223
Mathematica [A] (verified)	223
Rubi [A] (verified)	224
Maple [A] (verified)	225
Fricas [A] (verification not implemented)	225
Sympy [A] (verification not implemented)	225
Maxima [F]	226
Giac [A] (verification not implemented)	226
Mupad [B] (verification not implemented)	227
Reduce [B] (verification not implemented)	227

Optimal result

Integrand size = 35, antiderivative size = 26

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \frac{2\sqrt{bx + \sqrt{a + b^2x^2}}}{b}$$

output `2*(b*x+(b^2*x^2+a)^(1/2))^(1/2)/b`

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \frac{2\sqrt{bx + \sqrt{a + b^2x^2}}}{b}$$

input `Integrate[Sqrt[b*x + Sqrt[a + b^2*x^2]]/Sqrt[a + b^2*x^2], x]`

output `(2*Sqrt[b*x + Sqrt[a + b^2*x^2]])/b`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.057$, Rules used = {2547, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sqrt{a + b^2x^2} + bx}}{\sqrt{a + b^2x^2}} dx$$

↓ 2547

$$\int \frac{1}{\sqrt{bx + \sqrt{b^2x^2 + a}}} d\left(\frac{bx + \sqrt{b^2x^2 + a}}{b}\right)$$

↓ 15

$$\frac{2\sqrt{\sqrt{a + b^2x^2} + bx}}{b}$$

input `Int[Sqrt[b*x + Sqrt[a + b^2*x^2]]/Sqrt[a + b^2*x^2],x]`

output `(2*Sqrt[b*x + Sqrt[a + b^2*x^2]])/b`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))], x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$\frac{2\sqrt{bx+\sqrt{b^2x^2+a}}}{b}$	23
default	$\frac{2\sqrt{bx+\sqrt{b^2x^2+a}}}{b}$	23

input `int((b*x+(b^2*x^2+a)^(1/2))^(1/2)/(b^2*x^2+a)^(1/2),x,method=_RETURNVERBOS E)`

output `2*(b*x+(b^2*x^2+a)^(1/2))^(1/2)/b`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \frac{2\sqrt{bx + \sqrt{b^2x^2 + a}}}{b}$$

input `integrate((b*x+(b^2*x^2+a)^(1/2))^(1/2)/(b^2*x^2+a)^(1/2),x, algorithm="fr icas")`

output `2*sqrt(b*x + sqrt(b^2*x^2 + a))/b`

Sympy [A] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \begin{cases} \frac{2\sqrt{bx+\sqrt{a+b^2x^2}}}{b} & \text{for } b \neq 0 \\ \frac{x}{\sqrt[4]{a}} & \text{otherwise} \end{cases}$$

input `integrate((b*x+(b**2*x**2+a)**(1/2))**(1/2)/(b**2*x**2+a)**(1/2),x)`

output `Piecewise((2*sqrt(b*x + sqrt(a + b**2*x**2))/b, Ne(b, 0)), (x/a**(1/4), True))`

Maxima [F]

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \int \frac{\sqrt{bx + \sqrt{b^2x^2 + a}}}{\sqrt{b^2x^2 + a}} dx$$

input `integrate((b*x+(b^2*x^2+a)^(1/2))^(1/2)/(b^2*x^2+a)^(1/2),x, algorithm="maxima")`

output `integrate(sqrt(b*x + sqrt(b^2*x^2 + a))/sqrt(b^2*x^2 + a), x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \frac{2\sqrt{bx + \sqrt{b^2x^2 + a}}}{b}$$

input `integrate((b*x+(b^2*x^2+a)^(1/2))^(1/2)/(b^2*x^2+a)^(1/2),x, algorithm="giac")`

output `2*sqrt(b*x + sqrt(b^2*x^2 + a))/b`

Mupad [B] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \frac{2\sqrt{\sqrt{b^2x^2 + a} + bx}}{b}$$

input `int(((a + b^2*x^2)^(1/2) + b*x)^(1/2)/(a + b^2*x^2)^(1/2),x)`output `(2*((a + b^2*x^2)^(1/2) + b*x)^(1/2))/b`**Reduce [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{\sqrt{bx + \sqrt{a + b^2x^2}}}{\sqrt{a + b^2x^2}} dx = \frac{2\sqrt{\sqrt{b^2x^2 + a} + bx}}{b}$$

input `int((b*x+(b^2*x^2+a)^(1/2))^(1/2)/(b^2*x^2+a)^(1/2),x)`output `(2*sqrt(sqrt(a + b**2*x**2) + b*x))/b`

3.27 $\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx$

Optimal result	228
Mathematica [A] (verified)	228
Rubi [A] (verified)	229
Maple [F]	231
Fricas [B] (verification not implemented)	231
Sympy [C] (verification not implemented)	232
Maxima [F]	232
Giac [F]	233
Mupad [F(-1)]	233
Reduce [B] (verification not implemented)	233

Optimal result

Integrand size = 32, antiderivative size = 63

$$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx = -\frac{2 \arctan\left(\frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a}}\right)}{a^{3/2}}$$

output

`-2*arctan((x+(a^2+x^2)^(1/2))^(1/2)/a^(1/2))/a^(3/2)-2*arctanh((x+(a^2+x^2)^(1/2))^(1/2)/a^(1/2))/a^(3/2)`

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.89

$$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx = -\frac{2\left(\arctan\left(\frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a}}\right) + \operatorname{arctanh}\left(\frac{\sqrt{x+\sqrt{a^2+x^2}}}{\sqrt{a}}\right)\right)}{a^{3/2}}$$

input

`Integrate[1/(x*Sqrt[a^2+x^2]*Sqrt[x+Sqrt[a^2+x^2]]),x]`

output

`(-2*(ArcTan[Sqrt[x+Sqrt[a^2+x^2]]/Sqrt[a]]+ArcTanh[Sqrt[x+Sqrt[a^2+x^2]]/Sqrt[a]]))/a^(3/2)`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.10, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {2545, 25, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{x\sqrt{a^2+x^2}\sqrt{\sqrt{a^2+x^2}+x}} dx \\
 & \quad \downarrow \text{2545} \\
 & 2 \int -\frac{1}{\sqrt{x+\sqrt{a^2+x^2}} \left(a^2 - (x+\sqrt{a^2+x^2})^2\right)} d(x+\sqrt{a^2+x^2}) \\
 & \quad \downarrow \text{25} \\
 & -2 \int \frac{1}{\sqrt{x+\sqrt{a^2+x^2}} \left(a^2 - (x+\sqrt{a^2+x^2})^2\right)} d(x+\sqrt{a^2+x^2}) \\
 & \quad \downarrow \text{266} \\
 & -4 \int \frac{1}{a^2 - (x+\sqrt{a^2+x^2})^2} d\sqrt{x+\sqrt{a^2+x^2}} \\
 & \quad \downarrow \text{756} \\
 & -4 \left(\frac{\int \frac{1}{a-x-\sqrt{a^2+x^2}} d\sqrt{x+\sqrt{a^2+x^2}}}{2a} + \frac{\int \frac{1}{a+x+\sqrt{a^2+x^2}} d\sqrt{x+\sqrt{a^2+x^2}}}{2a} \right) \\
 & \quad \downarrow \text{216} \\
 & -4 \left(\frac{\int \frac{1}{a-x-\sqrt{a^2+x^2}} d\sqrt{x+\sqrt{a^2+x^2}}}{2a} + \frac{\arctan\left(\frac{\sqrt{\sqrt{a^2+x^2}+x}}{\sqrt{a}}\right)}{2a^{3/2}} \right) \\
 & \quad \downarrow \text{219} \\
 & -4 \left(\frac{\arctan\left(\frac{\sqrt{\sqrt{a^2+x^2}+x}}{\sqrt{a}}\right)}{2a^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{\sqrt{a^2+x^2}+x}}{\sqrt{a}}\right)}{2a^{3/2}} \right)
 \end{aligned}$$

input `Int[1/(x*Sqrt[a^2 + x^2]*Sqrt[x + Sqrt[a^2 + x^2]]),x]`

output `-4*(ArcTan[Sqrt[x + Sqrt[a^2 + x^2]]/Sqrt[a]]/(2*a^(3/2)) + ArcTanh[Sqrt[x + Sqrt[a^2 + x^2]]/Sqrt[a]]/(2*a^(3/2)))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] :> Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] :> With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 756 `Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] :> With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 2545

```
Int[(x_)^(p_.)*((g_) + (i_.)*(x_)^2)^(m_.)*((e_.)*(x_) + (f_.)*Sqrt[(a_) +
(c_.)*(x_)^2])^(n_.), x_Symbol] :> Simp[(1/(2^(2*m + p + 1)*e^(p + 1)*f^(2*
m)))*(i/c)^m Subst[Int[x^(n - 2*m - p - 2)*((-a)*f^2 + x^2)^p*(a*f^2 + x^
2)^(2*m + 1), x], x, e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, e, f, g,
i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[p, 2*m]
&& (IntegerQ[m] || GtQ[i/c, 0])
```

Maple [F]

$$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx$$

input

```
int(1/x/(a^2+x^2)^(1/2)/(x+(a^2+x^2)^(1/2))^(1/2),x)
```

output

```
int(1/x/(a^2+x^2)^(1/2)/(x+(a^2+x^2)^(1/2))^(1/2),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 125 vs. $2(47) = 94$.

Time = 0.09 (sec) , antiderivative size = 237, normalized size of antiderivative = 3.76

$$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx$$

$$= \frac{2\sqrt{a} \arctan\left(-\frac{(\sqrt{ax}-\sqrt{a^2+x^2}\sqrt{a})\sqrt{x+\sqrt{a^2+x^2}}}{a^2}\right) + \sqrt{a} \log\left(\frac{a^2+\sqrt{a^2+x^2}a - ((a-x)\sqrt{a+\sqrt{a^2+x^2}\sqrt{a}})\sqrt{x+\sqrt{a^2+x^2}}}{x}\right)}{a^2},$$

input

```
integrate(1/x/(a^2+x^2)^(1/2)/(x+(a^2+x^2)^(1/2))^(1/2),x, algorithm="fric
as")
```


output

```
[(2*sqrt(a)*arctan(-(sqrt(a)*x - sqrt(a^2 + x^2))*sqrt(a))*sqrt(x + sqrt(a^2 + x^2))/a^2) + sqrt(a)*log((a^2 + sqrt(a^2 + x^2)*a - ((a - x)*sqrt(a) + sqrt(a^2 + x^2))*sqrt(a))*sqrt(x + sqrt(a^2 + x^2)))/x)/a^2, (2*sqrt(-a)*arctan(-(sqrt(-a)*x - sqrt(a^2 + x^2))*sqrt(-a))*sqrt(x + sqrt(a^2 + x^2))/a^2) - sqrt(-a)*log(-(a^2 - sqrt(a^2 + x^2)*a - (sqrt(-a)*(a + x) - sqrt(a^2 + x^2))*sqrt(-a))*sqrt(x + sqrt(a^2 + x^2)))/x)/a^2]
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.03 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.73

$$\int \frac{1}{x\sqrt{a^2 + x^2}\sqrt{x + \sqrt{a^2 + x^2}}} dx = -\frac{\Gamma^2\left(\frac{3}{4}\right)\Gamma\left(\frac{5}{4}\right) {}_3F_2\left(\begin{matrix} \frac{3}{4}, \frac{3}{4}, \frac{5}{4} \\ \frac{3}{2}, \frac{7}{4} \end{matrix} \middle| \frac{a^2 e^{i\pi}}{x^2}\right)}{\pi x^{\frac{3}{2}}\Gamma\left(\frac{7}{4}\right)}$$

input

```
integrate(1/x/(a**2+x**2)**(1/2)/(x+(a**2+x**2)**(1/2))**(1/2),x)
```

output

```
-gamma(3/4)**2*gamma(5/4)*hyper((3/4, 3/4, 5/4), (3/2, 7/4), a**2*exp_polar(I*pi)/x**2)/(pi*x**(3/2)*gamma(7/4))
```

Maxima [F]

$$\int \frac{1}{x\sqrt{a^2 + x^2}\sqrt{x + \sqrt{a^2 + x^2}}} dx = \int \frac{1}{\sqrt{a^2 + x^2}\sqrt{x + \sqrt{a^2 + x^2}}} dx$$

input

```
integrate(1/x/(a^2+x^2)^(1/2)/(x+(a^2+x^2)^(1/2))^(1/2),x, algorithm="maxima")
```

output

```
integrate(1/(sqrt(a^2 + x^2)*sqrt(x + sqrt(a^2 + x^2)))*x, x)
```

Giac [F]

$$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx = \int \frac{1}{\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}x} dx$$

input `integrate(1/x/(a^2+x^2)^(1/2)/(x+(a^2+x^2)^(1/2))^(1/2),x, algorithm="giac")`

output `integrate(1/(sqrt(a^2 + x^2)*sqrt(x + sqrt(a^2 + x^2))*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx = \int \frac{1}{x\sqrt{x+\sqrt{a^2+x^2}}\sqrt{a^2+x^2}} dx$$

input `int(1/(x*(x + (a^2 + x^2)^(1/2))^(1/2)*(a^2 + x^2)^(1/2)),x)`

output `int(1/(x*(x + (a^2 + x^2)^(1/2))^(1/2)*(a^2 + x^2)^(1/2)), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.62

$$\int \frac{1}{x\sqrt{a^2+x^2}\sqrt{x+\sqrt{a^2+x^2}}} dx$$

$$= \frac{\sqrt{a} \left(\operatorname{atan} \left(\frac{\sqrt{a}\sqrt{\sqrt{a^2+x^2}+x}\sqrt{a^2+x^2}-\sqrt{a}\sqrt{\sqrt{a^2+x^2}+x}a-\sqrt{a}\sqrt{\sqrt{a^2+x^2}+x}x}{2a^2} \right) + \log \left(\sqrt{\sqrt{a^2+x^2}+x} - \sqrt{a} \right) - \log \left(\sqrt{\sqrt{a^2+x^2}+x} + \sqrt{a} \right) \right)}{a^2}$$

input `int(1/x/(a^2+x^2)^(1/2)/(x+(a^2+x^2)^(1/2))^(1/2),x)`

output

```
(sqrt(a)*(atan((sqrt(a)*sqrt(sqrt(a**2 + x**2) + x)*sqrt(a**2 + x**2) - sqrt(a)*sqrt(sqrt(a**2 + x**2) + x)*a - sqrt(a)*sqrt(sqrt(a**2 + x**2) + x)*x)/(2*a**2)) + log(sqrt(sqrt(a**2 + x**2) + x) - sqrt(a)) - log(sqrt(sqrt(a**2 + x**2) + x) + sqrt(a))))/a**2
```

3.28 $\int \frac{\sqrt{x+\sqrt{a^2+x^2}}}{x} dx$

Optimal result	235
Mathematica [A] (verified)	235
Rubi [A] (verified)	236
Maple [C] (verified)	238
Fricas [A] (verification not implemented)	239
Sympy [C] (verification not implemented)	240
Maxima [F]	240
Giac [F]	240
Mupad [F(-1)]	241
Reduce [B] (verification not implemented)	241

Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx = 2\sqrt{x + \sqrt{a^2 + x^2}} - 2\sqrt{a} \arctan\left(\frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a}}\right) - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a}}\right)$$

output

```
-2*arctan((x+(a^2+x^2)^(1/2))^(1/2)/a^(1/2))*a^(1/2)-2*arctanh((x+(a^2+x^2)^(1/2))^(1/2)/a^(1/2))*a^(1/2)+2*(x+(a^2+x^2)^(1/2))^(1/2)
```

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx = 2\sqrt{x + \sqrt{a^2 + x^2}} - 2\sqrt{a} \arctan\left(\frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a}}\right) - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{x + \sqrt{a^2 + x^2}}}{\sqrt{a}}\right)$$

input

```
Integrate[Sqrt[x + Sqrt[a^2 + x^2]]/x,x]
```

output

$$2\sqrt{x + \sqrt{a^2 + x^2}} - 2\sqrt{a} \operatorname{ArcTan}[\sqrt{x + \sqrt{a^2 + x^2}}] / \sqrt{a} - 2\sqrt{a} \operatorname{ArcTanh}[\sqrt{x + \sqrt{a^2 + x^2}}] / \sqrt{a}$$
Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.12, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2544, 25, 363, 266, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{\sqrt{a^2 + x^2} + x}}{x} dx$$

$$\downarrow 2544$$

$$\int -\frac{(\sqrt{a^2 + x^2} + x)^2 + a^2}{\sqrt{\sqrt{a^2 + x^2} + x} (a^2 - (\sqrt{a^2 + x^2} + x)^2)} d(\sqrt{a^2 + x^2} + x)$$

$$\downarrow 25$$

$$-\int \frac{a^2 + (x + \sqrt{a^2 + x^2})^2}{\sqrt{x + \sqrt{a^2 + x^2}} (a^2 - (x + \sqrt{a^2 + x^2})^2)} d(x + \sqrt{a^2 + x^2})$$

$$\downarrow 363$$

$$2\sqrt{\sqrt{a^2 + x^2} + x} - 2a^2 \int \frac{1}{\sqrt{x + \sqrt{a^2 + x^2}} (a^2 - (x + \sqrt{a^2 + x^2})^2)} d(x + \sqrt{a^2 + x^2})$$

$$\downarrow 266$$

$$2\sqrt{\sqrt{a^2 + x^2} + x} - 4a^2 \int \frac{1}{a^2 - (x + \sqrt{a^2 + x^2})^2} d\sqrt{x + \sqrt{a^2 + x^2}}$$

$$\downarrow 756$$

$$2\sqrt{\sqrt{a^2 + x^2} + x} - 4a^2 \left(\frac{\int \frac{1}{a-x-\sqrt{a^2+x^2}} d\sqrt{x + \sqrt{a^2 + x^2}}}{2a} + \frac{\int \frac{1}{a+x+\sqrt{a^2+x^2}} d\sqrt{x + \sqrt{a^2 + x^2}}}{2a} \right)$$

$$\begin{aligned}
 & \downarrow 216 \\
 & 2\sqrt{\sqrt{a^2+x^2}+x} - 4a^2 \left(\frac{\int \frac{1}{a-x-\sqrt{a^2+x^2}} d\sqrt{x+\sqrt{a^2+x^2}}}{2a} + \frac{\arctan\left(\frac{\sqrt{\sqrt{a^2+x^2}+x}}{\sqrt{a}}\right)}{2a^{3/2}} \right) \\
 & \downarrow 219 \\
 & 2\sqrt{\sqrt{a^2+x^2}+x} - 4a^2 \left(\frac{\arctan\left(\frac{\sqrt{\sqrt{a^2+x^2}+x}}{\sqrt{a}}\right)}{2a^{3/2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{\sqrt{a^2+x^2}+x}}{\sqrt{a}}\right)}{2a^{3/2}} \right)
 \end{aligned}$$

input `Int[Sqrt[x + Sqrt[a^2 + x^2]]/x,x]`

output `2*Sqrt[x + Sqrt[a^2 + x^2]] - 4*a^2*(ArcTan[Sqrt[x + Sqrt[a^2 + x^2]]/Sqrt[a]]/(2*a^(3/2))) + ArcTanh[Sqrt[x + Sqrt[a^2 + x^2]]/Sqrt[a]]/(2*a^(3/2))`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 266 `Int[((c_.)*(x_))^(m)*((a_) + (b_.)*(x_)^2)^(p), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m+1)-1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 363 `Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^2)^(p_)*((c_) + (d_)*(x_)^2), x_Symbol] := Simp[d*(e*x)^(m + 1)*((a + b*x^2)^(p + 1)/(b*e*(m + 2*p + 3))), x] - Simp[(a*d*(m + 1) - b*c*(m + 2*p + 3))/(b*(m + 2*p + 3)) Int[(e*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && NeQ[m + 2*p + 3, 0]`

rule 756 `Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[r/(2*a) Int[1/(r - s*x^2), x], x] + Simp[r/(2*a) Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 2544 `Int[((g_) + (h_)*(x_))^(m_)*((e_)*(x_) + (f_)*Sqrt[(a_) + (c_)*(x_)^2])^(n_), x_Symbol] := Simp[1/(2^(m + 1)*e^(m + 1)) Subst[Int[x^(n - m - 2)*(a*f^2 + x^2)*((-a)*f^2*h + 2*e*g*x + h*x^2)^m, x], x, e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[m]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.30

method	result	size
meijerg	$2\sqrt{2}\sqrt{x}\operatorname{hypergeom}\left(\left[-\frac{1}{4}, -\frac{1}{4}, \frac{1}{4}\right], \left[\frac{1}{2}, \frac{3}{4}\right], -\frac{a^2}{x^2}\right)$	25

input `int((x+(a^2+x^2)^(1/2))^(1/2)/x,x,method=_RETURNVERBOSE)`

output `2*2^(1/2)*x^(1/2)*hypergeom([-1/4,-1/4,1/4],[1/2,3/4],-a^2/x^2)`

Fricas [A] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. $2(62) = 124$.

Time = 0.09 (sec) , antiderivative size = 257, normalized size of antiderivative = 3.13

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx$$

$$= \left[2\sqrt{a} \arctan \left(-\frac{(\sqrt{ax} - \sqrt{a^2 + x^2}\sqrt{a})\sqrt{x + \sqrt{a^2 + x^2}}}{a^2} \right) \right. \\ \left. + \sqrt{a} \log \left(\frac{a^2 + \sqrt{a^2 + x^2}a - ((a - x)\sqrt{a} + \sqrt{a^2 + x^2}\sqrt{a})\sqrt{x + \sqrt{a^2 + x^2}}}{x} \right) \right. \\ \left. + 2\sqrt{x + \sqrt{a^2 + x^2}}, 2\sqrt{-a} \arctan \left(-\frac{(\sqrt{-ax} - \sqrt{a^2 + x^2}\sqrt{-a})\sqrt{x + \sqrt{a^2 + x^2}}}{a^2} \right) \right. \\ \left. + \sqrt{-a} \log \left(-\frac{a^2 - \sqrt{a^2 + x^2}a + (\sqrt{-a}(a + x) - \sqrt{a^2 + x^2}\sqrt{-a})\sqrt{x + \sqrt{a^2 + x^2}}}{x} \right) \right. \\ \left. + 2\sqrt{x + \sqrt{a^2 + x^2}} \right]$$

input `integrate((x+(a^2+x^2)^(1/2))^(1/2)/x,x, algorithm="fricas")`

output `[2*sqrt(a)*arctan(-(sqrt(a)*x - sqrt(a^2 + x^2)*sqrt(a))*sqrt(x + sqrt(a^2 + x^2)))/a^2) + sqrt(a)*log((a^2 + sqrt(a^2 + x^2)*a - ((a - x)*sqrt(a) + sqrt(a^2 + x^2)*sqrt(a))*sqrt(x + sqrt(a^2 + x^2)))/x) + 2*sqrt(x + sqrt(a^2 + x^2)), 2*sqrt(-a)*arctan(-(sqrt(-a)*x - sqrt(a^2 + x^2)*sqrt(-a))*sqrt(x + sqrt(a^2 + x^2)))/a^2) + sqrt(-a)*log(-(a^2 - sqrt(a^2 + x^2)*a + (sqrt(-a)*(a + x) - sqrt(a^2 + x^2)*sqrt(-a))*sqrt(x + sqrt(a^2 + x^2)))/x) + 2*sqrt(x + sqrt(a^2 + x^2))]`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.65 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.62

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx = \frac{\sqrt{x}\Gamma^2\left(-\frac{1}{4}\right)\Gamma\left(\frac{1}{4}\right) {}_3F_2\left(\begin{matrix} -\frac{1}{4}, -\frac{1}{4}, \frac{1}{4} \\ \frac{1}{2}, \frac{3}{4} \end{matrix} \middle| \frac{a^2 e^{i\pi}}{x^2}\right)}{8\pi\Gamma\left(\frac{3}{4}\right)}$$

input `integrate((x+(a**2+x**2)**(1/2))**(1/2)/x,x)`

output `sqrt(x)*gamma(-1/4)**2*gamma(1/4)*hyper((-1/4, -1/4, 1/4), (1/2, 3/4), a**2*exp_polar(I*pi)/x**2)/(8*pi*gamma(3/4))`

Maxima [F]

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx = \int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx$$

input `integrate((x+(a^2+x^2)^(1/2))^(1/2)/x,x, algorithm="maxima")`

output `integrate(sqrt(x + sqrt(a^2 + x^2))/x, x)`

Giac [F]

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx = \int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx$$

input `integrate((x+(a^2+x^2)^(1/2))^(1/2)/x,x, algorithm="giac")`

output `integrate(sqrt(x + sqrt(a^2 + x^2))/x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx = \int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx$$

input `int((x + (a^2 + x^2)^(1/2))^(1/2)/x,x)`output `int((x + (a^2 + x^2)^(1/2))^(1/2)/x, x)`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.43

$$\begin{aligned} & \int \frac{\sqrt{x + \sqrt{a^2 + x^2}}}{x} dx \\ &= \sqrt{a} \operatorname{atan} \left(\frac{\sqrt{a} \sqrt{\sqrt{a^2 + x^2} + x} \sqrt{a^2 + x^2} - \sqrt{a} \sqrt{\sqrt{a^2 + x^2} + x} a - \sqrt{a} \sqrt{\sqrt{a^2 + x^2} + x} x}{2a^2} \right) \\ & \quad + 2\sqrt{\sqrt{a^2 + x^2} + x} + \sqrt{a} \log \left(\sqrt{\sqrt{a^2 + x^2} + x} - \sqrt{a} \right) \\ & \quad - \sqrt{a} \log \left(\sqrt{\sqrt{a^2 + x^2} + x} + \sqrt{a} \right) \end{aligned}$$

input `int((x+(a^2+x^2)^(1/2))^(1/2)/x,x)`output `sqrt(a)*atan((sqrt(a)*sqrt(sqrt(a**2 + x**2) + x)*sqrt(a**2 + x**2) - sqrt(a)*sqrt(sqrt(a**2 + x**2) + x)*a - sqrt(a)*sqrt(sqrt(a**2 + x**2) + x)*x)/(2*a**2)) + 2*sqrt(sqrt(a**2 + x**2) + x) + sqrt(a)*log(sqrt(sqrt(a**2 + x**2) + x) - sqrt(a)) - sqrt(a)*log(sqrt(sqrt(a**2 + x**2) + x) + sqrt(a))`

3.29 $\int x^3 \log^3(2+x) \log(3+x) dx$

Optimal result	243
Mathematica [A] (verified)	244
Rubi [A] (verified)	245
Maple [F]	248
Fricas [F]	248
Sympy [F]	249
Maxima [A] (verification not implemented)	250
Giac [F]	250
Mupad [F(-1)]	251
Reduce [F]	251

Optimal result

Integrand size = 14, antiderivative size = 606

$$\begin{aligned}
\int x^3 \log^3(2+x) \log(3+x) dx = & -\frac{302177x}{1152} + \frac{8029x^2}{2304} - \frac{763x^3}{3456} + \frac{3x^4}{256} + \frac{377}{64}(2+x)^2 \\
& - \frac{71}{216}(2+x)^3 + \frac{3}{256}(2+x)^4 + \frac{2069}{144} \log(2+x) \\
& - \frac{187}{64}x^2 \log(2+x) + \frac{83}{288}x^3 \log(2+x) - \frac{3}{128}x^4 \log(2+x) \\
& + \frac{6733}{32}(2+x) \log(2+x) - \frac{377}{32}(2+x)^2 \log(2+x) \\
& + \frac{71}{72}(2+x)^3 \log(2+x) - \frac{3}{64}(2+x)^4 \log(2+x) \\
& - \frac{43}{12} \log^2(2+x) - \frac{17}{48}x^3 \log^2(2+x) + \frac{3}{64}x^4 \log^2(2+x) \\
& - \frac{1251}{16}(2+x) \log^2(2+x) + \frac{273}{32}(2+x)^2 \log^2(2+x) \\
& - \frac{3}{4}(2+x)^3 \log^2(2+x) + \frac{3}{64}(2+x)^4 \log^2(2+x) \\
& + \frac{65}{4}(2+x) \log^3(2+x) - \frac{33}{8}(2+x)^2 \log^3(2+x) \\
& + \frac{3}{4}(2+x)^3 \log^3(2+x) - \frac{1}{16}(2+x)^4 \log^3(2+x) \\
& + \frac{3891}{128} \log(3+x) - \frac{115}{48}x^2 \log(3+x) \\
& + \frac{37}{144}x^3 \log(3+x) - \frac{3}{128}x^4 \log(3+x) \\
& + \frac{415}{12}(3+x) \log(3+x) - \frac{4083}{32} \log(2+x) \log(3+x) \\
& - 25x \log(2+x) \log(3+x) + \frac{13}{4}x^2 \log(2+x) \log(3+x) \\
& - \frac{7}{12}x^3 \log(2+x) \log(3+x) + \frac{3}{32}x^4 \log(2+x) \log(3+x) \\
& + \frac{963}{16} \log^2(2+x) \log(3+x) + 6x \log^2(2+x) \log(3+x) \\
& - \frac{3}{2}x^2 \log^2(2+x) \log(3+x) + \frac{1}{2}x^3 \log^2(2+x) \log(3+x) \\
& - \frac{3}{16}x^4 \log^2(2+x) \log(3+x) - \frac{81}{4} \log^3(2+x) \log(3+x) \\
& + \frac{1}{4}x^4 \log^3(2+x) \log(3+x) - \frac{5609 \operatorname{PolyLog}(2, -2-x)}{96} \\
& + \frac{563}{8} \log(2+x) \operatorname{PolyLog}(2, -2-x) \\
& - \frac{195}{4} \log^2(2+x) \operatorname{PolyLog}(2, -2-x) \\
& - \frac{563 \operatorname{PolyLog}(3, -2-x)}{8} \\
& + \frac{195}{2} \log(2+x) \operatorname{PolyLog}(3, -2-x) \\
& - \frac{195 \operatorname{PolyLog}(4, -2-x)}{2}
\end{aligned}$$

output

```

-302177/1152*x+3/256*x^4-3/64*(2+x)^4*ln(2+x)-17/48*x^3*ln(2+x)^2+3/64*x^4
*ln(2+x)^2-1251/16*(2+x)*ln(2+x)^2+273/32*(2+x)^2*ln(2+x)^2-3/4*(2+x)^3*ln
(2+x)^2+3/64*(2+x)^4*ln(2+x)^2+65/4*(2+x)*ln(2+x)^3-33/8*(2+x)^2*ln(2+x)^3
+3/4*(2+x)^3*ln(2+x)^3-1/16*(2+x)^4*ln(2+x)^3-115/48*x^2*ln(3+x)+37/144*x^
3*ln(3+x)-3/128*x^4*ln(3+x)+415/12*(3+x)*ln(3+x)-4083/32*ln(2+x)*ln(3+x)+9
63/16*ln(2+x)^2*ln(3+x)-81/4*ln(2+x)^3*ln(3+x)+563/8*ln(2+x)*polylog(2,-2-
x)-195/4*ln(2+x)^2*polylog(2,-2-x)+195/2*ln(2+x)*polylog(3,-2-x)-187/64*x^
2*ln(2+x)+83/288*x^3*ln(2+x)-3/128*x^4*ln(2+x)+6733/32*(2+x)*ln(2+x)-377/3
2*(2+x)^2*ln(2+x)+71/72*(2+x)^3*ln(2+x)-763/3456*x^3-43/12*ln(2+x)^2-5609/
96*polylog(2,-2-x)-563/8*polylog(3,-2-x)-195/2*polylog(4,-2-x)+377/64*(2+x
)^2-71/216*(2+x)^3+3/256*(2+x)^4+3891/128*ln(3+x)+2069/144*ln(2+x)+8029/23
04*x^2-25*x*ln(2+x)*ln(3+x)+13/4*x^2*ln(2+x)*ln(3+x)-7/12*x^3*ln(2+x)*ln(3
+x)+3/32*x^4*ln(2+x)*ln(3+x)+6*x*ln(2+x)^2*ln(3+x)-3/2*x^2*ln(2+x)^2*ln(3+
x)+1/2*x^3*ln(2+x)^2*ln(3+x)-3/16*x^4*ln(2+x)^2*ln(3+x)+1/4*x^4*ln(2+x)^3*
ln(3+x)

```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 412, normalized size of antiderivative = 0.68

$$\int x^3 \log^3(2+x) \log(3+x) dx$$

$$= \frac{-195984 - 558290x + 17705x^2 - 1050x^3 + 54x^4 + 910528 \log(2+x) + 400008x \log(2+x) - 22836x^2 \log(2+x) - 145000x^2 \log(3+x) + 910528x \log(3+x) - 22836x^2 \log(3+x) + 145000x^2 \log(3+x)^2 - 33000x^2 \log(3+x)^3}{(2+x)^4}$$

input

```
Integrate[x^3*Log[2 + x]^3*Log[3 + x],x]
```

output

```
(-195984 - 558290*x + 17705*x^2 - 1050*x^3 + 54*x^4 + 910528*Log[2 + x] +
400008*x*Log[2 + x] - 22836*x^2*Log[2 + x] + 2072*x^3*Log[2 + x] - 162*x^4
*Log[2 + x] - 302016*Log[2 + x]^2 - 118800*x*Log[2 + x]^2 + 11880*x^2*Log[
2 + x]^2 - 1680*x^3*Log[2 + x]^2 + 216*x^4*Log[2 + x]^2 + 48384*Log[2 + x]
^3 + 15552*x*Log[2 + x]^3 - 2592*x^2*Log[2 + x]^3 + 576*x^3*Log[2 + x]^3 -
144*x^4*Log[2 + x]^3 + 309078*Log[3 + x] + 79680*x*Log[3 + x] - 5520*x^2*
Log[3 + x] + 592*x^3*Log[3 + x] - 54*x^4*Log[3 + x] - 293976*Log[2 + x]*Lo
g[3 + x] - 57600*x*Log[2 + x]*Log[3 + x] + 7488*x^2*Log[2 + x]*Log[3 + x]
- 1344*x^3*Log[2 + x]*Log[3 + x] + 216*x^4*Log[2 + x]*Log[3 + x] + 138672*
Log[2 + x]^2*Log[3 + x] + 13824*x*Log[2 + x]^2*Log[3 + x] - 3456*x^2*Log[2
+ x]^2*Log[3 + x] + 1152*x^3*Log[2 + x]^2*Log[3 + x] - 432*x^4*Log[2 + x]
^2*Log[3 + x] - 46656*Log[2 + x]^3*Log[3 + x] + 576*x^4*Log[2 + x]^3*Log[3
+ x] - 24*(5609 - 6756*Log[2 + x] + 4680*Log[2 + x]^2)*PolyLog[2, -2 - x]
+ 288*(-563 + 780*Log[2 + x])*PolyLog[3, -2 - x] - 224640*PolyLog[4, -2 -
x])/2304
```

Rubi [A] (verified)

Time = 4.79 (sec) , antiderivative size = 741, normalized size of antiderivative = 1.22, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {2889, 2863, 2009, 7293, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int x^3 \log^3(x+2) \log(x+3) dx \\
 & \quad \downarrow \text{2889} \\
 & -\frac{1}{4} \int \frac{x^4 \log^3(x+2)}{x+3} dx - \frac{3}{4} \int \frac{x^4 \log^2(x+2) \log(x+3)}{x+2} dx + \frac{1}{4} x^4 \log^3(x+2) \log(x+3) \\
 & \quad \downarrow \text{2863} \\
 & -\frac{3}{4} \int \frac{x^4 \log^2(x+2) \log(x+3)}{x+2} dx - \\
 & \frac{1}{4} \int \left(x^3 \log^3(x+2) - 3x^2 \log^3(x+2) + 9x \log^3(x+2) + \frac{81 \log^3(x+2)}{x+3} - 27 \log^3(x+2) \right) dx + \\
 & \quad \frac{1}{4} x^4 \log^3(x+2) \log(x+3)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 2009 \\
 & -\frac{3}{4} \int \frac{x^4 \log^2(x+2) \log(x+3)}{x+2} dx + \\
 & \frac{1}{4} \left(-486 \operatorname{PolyLog}(4, -x-2) - 243 \operatorname{PolyLog}(2, -x-2) \log^2(x+2) + 486 \operatorname{PolyLog}(3, -x-2) \log(x+2) + \frac{3}{128} \right. \\
 & \quad \left. \frac{1}{4} x^4 \log^3(x+2) \log(x+3) \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 7293 \\
 & -\frac{3}{4} \int \left(\log^2(x+2) \log(x+3) x^3 - 2 \log^2(x+2) \log(x+3) x^2 + 4 \log^2(x+2) \log(x+3) x + \frac{16 \log^2(x+2) \log(x+3)}{x+2} \right) dx \\
 & \frac{1}{4} \left(-486 \operatorname{PolyLog}(4, -x-2) - 243 \operatorname{PolyLog}(2, -x-2) \log^2(x+2) + 486 \operatorname{PolyLog}(3, -x-2) \log(x+2) + \frac{3}{128} \right. \\
 & \quad \left. \frac{1}{4} x^4 \log^3(x+2) \log(x+3) \right)
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 2009 \\
 & -\frac{3}{4} \left(\frac{5609 \operatorname{PolyLog}(2, -x-2)}{72} + \frac{563 \operatorname{PolyLog}(3, -x-2)}{6} - 32 \operatorname{PolyLog}(4, -x-2) - 16 \operatorname{PolyLog}(2, -x-2) \log^2(x+2) \right) \\
 & \frac{1}{4} \left(-486 \operatorname{PolyLog}(4, -x-2) - 243 \operatorname{PolyLog}(2, -x-2) \log^2(x+2) + 486 \operatorname{PolyLog}(3, -x-2) \log(x+2) + \frac{3}{128} \right. \\
 & \quad \left. \frac{1}{4} x^4 \log^3(x+2) \log(x+3) \right)
 \end{aligned}$$

input `Int[x^3*Log[2 + x]^3*Log[3 + x],x]`

output

```
(x^4*Log[2 + x]^3*Log[3 + x])/4 + (-390*x + (99*(2 + x)^2)/8 - (2*(2 + x)^3)/3 + (3*(2 + x)^4)/128 + 390*(2 + x)*Log[2 + x] - (99*(2 + x)^2*Log[2 + x])/4 + 2*(2 + x)^3*Log[2 + x] - (3*(2 + x)^4*Log[2 + x])/32 - 195*(2 + x)*Log[2 + x]^2 + (99*(2 + x)^2*Log[2 + x]^2)/4 - 3*(2 + x)^3*Log[2 + x]^2 + (3*(2 + x)^4*Log[2 + x]^2)/16 + 65*(2 + x)*Log[2 + x]^3 - (33*(2 + x)^2*Log[2 + x]^3)/2 + 3*(2 + x)^3*Log[2 + x]^3 - ((2 + x)^4*Log[2 + x]^3)/4 - 81*Log[2 + x]^3*Log[3 + x] - 243*Log[2 + x]^2*PolyLog[2, -2 - x] + 486*Log[2 + x]*PolyLog[3, -2 - x] - 486*PolyLog[4, -2 - x])/4 - (3*((189857*x)/864 - (8029*x^2)/1728 + (763*x^3)/2592 - x^4/64 - (179*(2 + x)^2)/48 + (35*(2 + x)^3)/162 - (2 + x)^4/128 - (2069*Log[2 + x])/108 + (187*x^2*Log[2 + x])/48 - (83*x^3*Log[2 + x])/216 + (x^4*Log[2 + x])/32 - (3613*(2 + x)*Log[2 + x])/24 + (179*(2 + x)^2*Log[2 + x])/24 - (35*(2 + x)^3*Log[2 + x])/54 + ((2 + x)^4*Log[2 + x])/32 + (43*Log[2 + x]^2)/9 + (17*x^3*Log[2 + x]^2)/36 - (x^4*Log[2 + x]^2)/16 + (157*(2 + x)*Log[2 + x]^2)/4 - (25*(2 + x)^2*Log[2 + x]^2)/8 - (1297*Log[3 + x])/32 + (115*x^2*Log[3 + x])/36 - (37*x^3*Log[3 + x])/108 + (x^4*Log[3 + x])/32 - (415*(3 + x)*Log[3 + x])/9 + (1361*Log[2 + x]*Log[3 + x])/8 + (100*x*Log[2 + x]*Log[3 + x])/3 - (13*x^2*Log[2 + x]*Log[3 + x])/3 + (7*x^3*Log[2 + x]*Log[3 + x])/9 - (x^4*Log[2 + x]*Log[3 + x])/8 - (321*Log[2 + x]^2*Log[3 + x])/4 - 8*x*Log[2 + x]^2*Log[3 + x] + 2*x^2*Log[2 + x]^2*Log[3 + x] - (2*x^3*Log[2 + x]^2*Log[3 + x])/3 ...
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2863

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]*(b_.))^(p_.)*((h_.)*(x_))^(m_.)*((f_) + (g_.)*(x_)^(r_.))^(q_.), x_Symbol] := Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n]]^p, (h*x)^m*(f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q, r}, x] && IntegerQ[m] && IntegerQ[q]
```


rule 2889

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.)*(x_))^(n_.)]*(b_.))^(p_.)*((f_.) + Log
[(h_.)*((i_.) + (j_.)*(x_))^(m_.)]*(g_.))*(x_)^(r_.), x_Symbol] := Simp[x^(
r + 1)*(a + b*Log[c*(d + e*x)^n])^p*((f + g*Log[h*(i + j*x)^m])/(r + 1)), x
] + (-Simp[g*j*(m/(r + 1)) Int[x^(r + 1)*((a + b*Log[c*(d + e*x)^n])^p/(i
+ j*x)), x], x] - Simp[b*e*n*(p/(r + 1)) Int[x^(r + 1)*(a + b*Log[c*(d +
e*x)^n])^(p - 1)*((f + g*Log[h*(i + j*x)^m])/(d + e*x)), x], x] /; FreeQ[
{a, b, c, d, e, f, g, h, i, j, m, n}, x] && IGtQ[p, 0] && IntegerQ[r] && (E
qQ[p, 1] || GtQ[r, 0]) && NeQ[r, -1]
```

rule 7293

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Maple [F]

$$\int x^3 \ln(2+x)^3 \ln(3+x) dx$$

input

```
int(x^3*ln(2+x)^3*ln(3+x),x)
```

output

```
int(x^3*ln(2+x)^3*ln(3+x),x)
```

Fricas [F]

$$\int x^3 \log^3(2+x) \log(3+x) dx = \int x^3 \log(x+3) \log(x+2)^3 dx$$

input

```
integrate(x^3*log(2+x)^3*log(3+x),x, algorithm="fricas")
```

output

```
integral(x^3*log(x + 3)*log(x + 2)^3, x)
```

SymPy [F]

$$\int x^3 \log^3(2+x) \log(3+x) dx = \left(\frac{x^4 \log(x+2)^3}{4} - \frac{3x^4 \log(x+2)^2}{16} + \frac{3x^4 \log(x+2)}{32} - \frac{3x^4}{128} + \frac{x^3 \log(x+2)^2}{2} - \frac{7x^3 \log(x+2)}{12} + \frac{37x^3}{144} - \frac{3x^2 \log(x+2)^2}{2} + \frac{13x^2 \log(x+2)}{4} - \frac{115x^2}{48} + 6x \log(x+2)^2 - 25x \log(x+2) + \frac{415x}{12} - 4 \log(x+2)^3 + 25 \log(x+2)^2 - \frac{415 \log(x+2)}{6} + \frac{10955281}{240000} \right) \log(x+3) \\ - \int \frac{24900000x}{x+3} dx + \int \left(-\frac{1725000x^2}{x+3} \right) dx + \int \frac{185000x^3}{x+3} dx + \int \left(-\frac{16875x^4}{x+3} \right) dx + \int \left(-\frac{49800000 \log(x+2)}{x+3} \right) dx + \int$$

input `integrate(x**3*ln(2+x)**3*ln(3+x), x)`

output `(x**4*log(x + 2)**3/4 - 3*x**4*log(x + 2)**2/16 + 3*x**4*log(x + 2)/32 - 3*x**4/128 + x**3*log(x + 2)**2/2 - 7*x**3*log(x + 2)/12 + 37*x**3/144 - 3*x**2*log(x + 2)**2/2 + 13*x**2*log(x + 2)/4 - 115*x**2/48 + 6*x*log(x + 2)**2 - 25*x*log(x + 2) + 415*x/12 - 4*log(x + 2)**3 + 25*log(x + 2)**2 - 415*log(x + 2)/6 + 10955281/240000)*log(x + 3) - (Integral(24900000*x/(x + 3), x) + Integral(-1725000*x**2/(x + 3), x) + Integral(185000*x**3/(x + 3), x) + Integral(-16875*x**4/(x + 3), x) + Integral(-49800000*log(x + 2)/(x + 3), x) + Integral(18000000*log(x + 2)**2/(x + 3), x) + Integral(-2880000*log(x + 2)**3/(x + 3), x) + Integral(-18000000*x*log(x + 2)/(x + 3), x) + Integral(4320000*x*log(x + 2)**2/(x + 3), x) + Integral(2340000*x**2*log(x + 2)/(x + 3), x) + Integral(-1080000*x**2*log(x + 2)**2/(x + 3), x) + Integral(-420000*x**3*log(x + 2)/(x + 3), x) + Integral(360000*x**3*log(x + 2)**2/(x + 3), x) + Integral(67500*x**4*log(x + 2)/(x + 3), x) + Integral(-135000*x**4*log(x + 2)**2/(x + 3), x) + Integral(180000*x**4*log(x + 2)**3/(x + 3), x) + Integral(32865843/(x + 3), x))/720000`

Maxima [A] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 518, normalized size of antiderivative = 0.85

$$\int x^3 \log^3(2+x) \log(3+x) dx = \text{Too large to display}$$

input `integrate(x^3*log(2+x)^3*log(3+x),x, algorithm="maxima")`

output

```

3/128*x^4 + 1/16*(4*x^4*log(x + 3) - x^4 + 4*x^3 - 18*x^2 + 108*x - 324*log(x + 3))*log(x + 2)^3 - 65/4*log(x + 3)*log(x + 2)^3 + 195/4*log(x + 3)*log(x + 2)^2*log(-x - 2) - 175/384*x^3 + 1/96*(9*x^4 - 70*x^3 + 495*x^2 - 6*(3*x^4 - 8*x^3 + 24*x^2 - 96*x)*log(x + 3) + 4680*log(x + 3)*log(-x - 2) - 4950*x + 4680*dilog(x + 3) + 5778*log(x + 3) + 6048*log(x + 2))*log(x + 2)^2 + 195/4*dilog(x + 3)*log(x + 2)^2 - 195/4*dilog(-x - 2)*log(x + 2)^2 + 563/16*log(x + 3)*log(x + 2)^2 + 21*log(x + 2)^3 + 17705/2304*x^2 + 1/8*(780*log(x + 2)^2 - 563*log(x + 2))*dilog(-x - 2) - 1/1152*(27*x^4 - 296*x^3 - 18720*log(x + 2)^3 + 2760*x^2 + 40536*log(x + 2)^2 - 39840*x - 67308*log(x + 2))*log(x + 3) - 1/1152*(81*x^4 - 1036*x^3 + 56160*log(x + 3)*log(x + 2)^2 + 112320*log(x + 3)*log(x + 2)*log(-x - 2) + 11418*x^2 - 12*(9*x^4 - 56*x^3 + 312*x^2 + 4680*log(x + 2)^2 - 2400*x - 6756*log(x + 2))*log(x + 3) + 112320*dilog(x + 3)*log(x + 2) + 112320*dilog(-x - 2)*log(x + 2) - 81072*log(x + 3)*log(x + 2) + 72576*log(x + 2)^2 - 200004*x - 81072*dilog(-x - 2) + 146988*log(x + 3) + 302016*log(x + 2) - 112320*polylog(3, -x - 2))*log(x + 2) + 563/8*dilog(-x - 2)*log(x + 2) - 5609/96*log(x + 3)*log(x + 2) + 1573/12*log(x + 2)^2 - 279145/1152*x - 5609/96*dilog(-x - 2) + 17171/128*log(x + 3) + 14227/36*log(x + 2) - 195/2*polylog(4, -x - 2) - 563/8*polylog(3, -x - 2)

```

Giac [F]

$$\int x^3 \log^3(2+x) \log(3+x) dx = \int x^3 \log(x+3) \log(x+2)^3 dx$$

input `integrate(x^3*log(2+x)^3*log(3+x),x, algorithm="giac")`

output `integrate(x^3*log(x + 3)*log(x + 2)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \log^3(2+x) \log(3+x) dx = \int x^3 \ln(x+2)^3 \ln(x+3) dx$$

input `int(x^3*log(x + 2)^3*log(x + 3),x)`output `int(x^3*log(x + 2)^3*log(x + 3), x)`**Reduce [F]**

$$\int x^3 \log^3(2+x) \log(3+x) dx = \text{Too large to display}$$

input `int(x^3*log(2+x)^3*log(3+x),x)`

output

```
(37440*int(log(x + 2)**3/(x**2 + 5*x + 6),x) - 81072*int(log(x + 2)**2/(x**2 + 5*x + 6),x) + 134616*int(log(x + 2)/(x**2 + 5*x + 6),x) + 576*log(x + 3)*log(x + 2)**3*x**4 - 9216*log(x + 3)*log(x + 2)**3 - 432*log(x + 3)*log(x + 2)**2*x**4 + 1152*log(x + 3)*log(x + 2)**2*x**3 - 3456*log(x + 3)*log(x + 2)**2*x**2 + 13824*log(x + 3)*log(x + 2)**2*x + 57600*log(x + 3)*log(x + 2)**2 + 216*log(x + 3)*log(x + 2)*x**4 - 1344*log(x + 3)*log(x + 2)*x**3 + 7488*log(x + 3)*log(x + 2)*x**2 - 57600*log(x + 3)*log(x + 2)*x - 159360*log(x + 3)*log(x + 2) - 54*log(x + 3)*x**4 + 592*log(x + 3)*x**3 - 5520*log(x + 3)*x**2 + 79680*log(x + 3)*x + 309078*log(x + 3) - 9360*log(x + 2)**4 - 144*log(x + 2)**3*x**4 + 576*log(x + 2)**3*x**3 - 2592*log(x + 2)**3*x**2 + 15552*log(x + 2)**3*x + 75408*log(x + 2)**3 + 216*log(x + 2)**2*x**4 - 1680*log(x + 2)**2*x**3 + 11880*log(x + 2)**2*x**2 - 118800*log(x + 2)**2*x - 369324*log(x + 2)**2 - 162*log(x + 2)*x**4 + 2072*log(x + 2)*x**3 - 22836*log(x + 2)*x**2 + 400008*log(x + 2)*x + 910528*log(x + 2) + 54*x**4 - 1050*x**3 + 17705*x**2 - 558290*x)/2304
```

$$3.30 \quad \int \frac{(x + \sqrt{b+x^2})^a}{\sqrt{b+x^2}} dx$$

Optimal result	252
Mathematica [A] (verified)	252
Rubi [A] (verified)	253
Maple [A] (verified)	254
Fricas [A] (verification not implemented)	254
Sympy [B] (verification not implemented)	254
Maxima [F]	255
Giac [A] (verification not implemented)	255
Mupad [B] (verification not implemented)	256
Reduce [B] (verification not implemented)	256

Optimal result

Integrand size = 23, antiderivative size = 17

$$\int \frac{(x + \sqrt{b+x^2})^a}{\sqrt{b+x^2}} dx = \frac{(x + \sqrt{b+x^2})^a}{a}$$

output $(x + \sqrt{b+x^2})^a / a$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{(x + \sqrt{b+x^2})^a}{\sqrt{b+x^2}} dx = \frac{(x + \sqrt{b+x^2})^a}{a}$$

input `Integrate[(x + Sqrt[b + x^2])^a/Sqrt[b + x^2], x]`

output $(x + \sqrt{b+x^2})^a / a$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2547, 15}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(\sqrt{b+x^2}+x)^a}{\sqrt{b+x^2}} dx$$

↓ 2547

$$\int (\sqrt{b+x^2}+x)^{a-1} d(\sqrt{b+x^2}+x)$$

↓ 15

$$\frac{(\sqrt{b+x^2}+x)^a}{a}$$

input `Int[(x + Sqrt[b + x^2])^a/Sqrt[b + x^2],x]`

output `(x + Sqrt[b + x^2])^a/a`

Defintions of rubi rules used

rule 15 `Int[(a_.)*(x_)^(m_.), x_Symbol] := Simp[a*(x^(m + 1)/(m + 1)), x] /; FreeQ[{a, m}, x] && NeQ[m, -1]`

rule 2547 `Int[((g_) + (i_.)*(x_)^2)^(m_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.), x_Symbol] := Simp[(1/(2^(2*m + 1)*e*f^(2*m)))*(i/c)^m Subst[Int[x^n*((d^2 + a*f^2 - 2*d*x + x^2)^(2*m + 1)/(-d + x)^(2*(m + 1))), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, i, n}, x] && EqQ[e^2 - c*f^2, 0] && EqQ[c*g - a*i, 0] && IntegerQ[2*m] && (IntegerQ[m] || GtQ[i/c, 0])`

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{(x+\sqrt{x^2+b})^a}{a}$	16
default	$\frac{(x+\sqrt{x^2+b})^a}{a}$	16

input `int((x+(x^2+b)^(1/2))^a/(x^2+b)^(1/2),x,method=_RETURNVERBOSE)`

output `(x+(x^2+b)^(1/2))^a/a`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{b + x^2})^a}{\sqrt{b + x^2}} dx = \frac{(x + \sqrt{x^2 + b})^a}{a}$$

input `integrate((x+(x^2+b)^(1/2))^a/(x^2+b)^(1/2),x, algorithm="fricas")`

output `(x + sqrt(x^2 + b))^a/a`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. 2(12) = 24.

Time = 1.22 (sec) , antiderivative size = 311, normalized size of antiderivative = 18.29

$$\int \frac{(x + \sqrt{b + x^2})^a}{\sqrt{b + x^2}} dx = \begin{cases} \frac{\sqrt{b}^{\frac{a}{2}} \sinh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{ax\sqrt{\frac{b}{x^2}+1}} + \frac{b^{\frac{a}{2}}x \cosh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{a\sqrt{b}} + \frac{b^{\frac{a}{2}}x \sinh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{a\sqrt{b}\sqrt{\frac{b}{x^2}+1}} - \frac{2b^{\frac{a}{2}} \cosh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{a\sqrt{b}} \\ \frac{b^{\frac{a}{2}} \sinh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{a\sqrt{1+\frac{x^2}{b}}} + \frac{b^{\frac{a}{2}}x^2 \sinh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{ab\sqrt{1+\frac{x^2}{b}}} + \frac{b^{\frac{a}{2}}x \cosh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{a\sqrt{b}} - \frac{2b^{\frac{a}{2}} \cosh\left(a \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right) - \operatorname{asinh}\left(\frac{x}{\sqrt{b}}\right)\right)}{a\sqrt{b}} \end{cases}$$

input `integrate((x+(x**2+b)**(1/2))**a/(x**2+b)**(1/2),x)`

output `Piecewise((sqrt(b)*b**(a/2)*sinh(a*asinh(x/sqrt(b)) - asinh(x/sqrt(b)))/(a*x*sqrt(b/x**2 + 1)) + b**(a/2)*x*cosh(a*asinh(x/sqrt(b)) - asinh(x/sqrt(b)))/(a*sqrt(b)) + b**(a/2)*x*sinh(a*asinh(x/sqrt(b)) - asinh(x/sqrt(b)))/(a*sqrt(b)*sqrt(b/x**2 + 1)) - 2*b**(a/2)*cosh(a*asinh(x/sqrt(b)))*gamma(1 - a/2)/(a**2*gamma(-a/2)), Abs(x**2/b) > 1), (b**(a/2)*sinh(a*asinh(x/sqrt(b)) - asinh(x/sqrt(b)))/(a*sqrt(1 + x**2/b)) + b**(a/2)*x**2*sinh(a*asinh(x/sqrt(b)) - asinh(x/sqrt(b)))/(a*b*sqrt(1 + x**2/b)) + b**(a/2)*x*cosh(a*asinh(x/sqrt(b)) - asinh(x/sqrt(b)))/(a*sqrt(b)) - 2*b**(a/2)*cosh(a*asinh(x/sqrt(b)))*gamma(1 - a/2)/(a**2*gamma(-a/2)), True))`

Maxima [F]

$$\int \frac{(x + \sqrt{b + x^2})^a}{\sqrt{b + x^2}} dx = \int \frac{(x + \sqrt{x^2 + b})^a}{\sqrt{x^2 + b}} dx$$

input `integrate((x+(x^2+b)^(1/2))^a/(x^2+b)^(1/2),x, algorithm="maxima")`

output `integrate((x + sqrt(x^2 + b))^a/sqrt(x^2 + b), x)`

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{b + x^2})^a}{\sqrt{b + x^2}} dx = \frac{(x + \sqrt{x^2 + b})^a}{a}$$

input `integrate((x+(x^2+b)^(1/2))^a/(x^2+b)^(1/2),x, algorithm="giac")`

output `(x + sqrt(x^2 + b))^a/a`

Mupad [B] (verification not implemented)

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{(x + \sqrt{b + x^2})^a}{\sqrt{b + x^2}} dx = \frac{(x + \sqrt{x^2 + b})^a}{a}$$

input `int((x + (b + x^2)^(1/2))^a/(b + x^2)^(1/2), x)`output `(x + (b + x^2)^(1/2))^a/a`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

$$\int \frac{(x + \sqrt{b + x^2})^a}{\sqrt{b + x^2}} dx = \frac{(\sqrt{x^2 + b} + x)^a}{a}$$

input `int((x+(x^2+b)^(1/2))^a/(x^2+b)^(1/2), x)`output `(sqrt(b + x**2) + x)**a/a`

3.31 $\int \left(x + \sqrt{b + x^2}\right)^a dx$

Optimal result	257
Mathematica [A] (verified)	257
Rubi [A] (verified)	258
Maple [B] (verified)	259
Fricas [A] (verification not implemented)	260
Sympy [B] (verification not implemented)	260
Maxima [F]	261
Giac [F]	262
Mupad [F(-1)]	262
Reduce [B] (verification not implemented)	262

Optimal result

Integrand size = 13, antiderivative size = 52

$$\int \left(x + \sqrt{b + x^2}\right)^a dx = -\frac{b(x + \sqrt{b + x^2})^{-1+a}}{2(1-a)} + \frac{(x + \sqrt{b + x^2})^{1+a}}{2(1+a)}$$

output

```
-1/2*b*(x+(x^2+b)^(1/2))^(1+a)/(1+a)+1/2*(x+(x^2+b)^(1/2))^(1+a)/(1+a)
```

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.88

$$\int \left(x + \sqrt{b + x^2}\right)^a dx = \frac{1}{2} \left(x + \sqrt{b + x^2}\right)^{-1+a} \left(\frac{b}{-1+a} + \frac{(x + \sqrt{b + x^2})^2}{1+a} \right)$$

input

```
Integrate[(x + Sqrt[b + x^2])^a,x]
```

output

```
((x + Sqrt[b + x^2])^(-1 + a)*(b/(-1 + a) + (x + Sqrt[b + x^2])^2/(1 + a)))/2
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.98, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2542, 244, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (\sqrt{b+x^2} + x)^a dx$$

$$\downarrow \text{2542}$$

$$\frac{1}{2} \int (x + \sqrt{x^2 + b})^{a-2} \left((x + \sqrt{x^2 + b})^2 + b \right) d(x + \sqrt{x^2 + b})$$

$$\downarrow \text{244}$$

$$\frac{1}{2} \int \left(b(x + \sqrt{x^2 + b})^{a-2} + (x + \sqrt{x^2 + b})^a \right) d(x + \sqrt{x^2 + b})$$

$$\downarrow \text{2009}$$

$$\frac{1}{2} \left(\frac{(\sqrt{b+x^2} + x)^{a+1}}{a+1} - \frac{b(\sqrt{b+x^2} + x)^{a-1}}{1-a} \right)$$

input `Int[(x + Sqrt[b + x^2])^a,x]`

output `((-(b*(x + Sqrt[b + x^2])^(-1 + a))/(1 - a)) + (x + Sqrt[b + x^2])^(1 + a)/(1 + a))/2`

Definitions of rubi rules used

rule 244 `Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^2)^(p_.), x_Symbol] := Int[Expand Integrand[(c*x)^m*(a + b*x^2)^p, x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[p, 0]`

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2542 `Int[((g_.) + (h_.)*((d_.) + (e_.)*(x_) + (f_.)*Sqrt[(a_) + (c_.)*(x_)^2])^(n_.))^p_.), x_Symbol] := Simp[1/(2*e) Subst[Int[(g + h*x^n)^p*((d^2 + a*f^2 - 2*d*x + x^2)/(d - x)^2), x], x, d + e*x + f*Sqrt[a + c*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h, n}, x] && EqQ[e^2 - c*f^2, 0] && IntegerQ[p]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(44) = 88.

Time = 0.04 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.31

method	result	size
meijerg	$\frac{b^{\frac{a}{2} + \frac{1}{2}} a \left(\frac{8\sqrt{\pi} x^{1+a} b^{-\frac{a}{2} - \frac{1}{2}} \left(\frac{ab}{x^2} + a - 1 \right) \left(\sqrt{1 + \frac{b}{x^2}} + 1 \right)^{a-1}}{(1+a)a(2a-2)} + \frac{4\sqrt{\pi} x^{1+a} b^{-\frac{a}{2} - \frac{1}{2}} \sqrt{1 + \frac{b}{x^2}} \left(\sqrt{1 + \frac{b}{x^2}} + 1 \right)^{a-1}}{(1+a)a} \right)}{4\sqrt{\pi}}$	120

input `int((x+(x^2+b)^(1/2))^a,x,method=_RETURNVERBOSE)`

output `1/4*b^(1/2*a+1/2)/Pi^(1/2)*a*(8*Pi^(1/2)/(1+a)/a*x^(1+a)*b^(-1/2*a-1/2)*(a*b/x^2+a-1)/(2*a-2)*((1+b/x^2)^(1/2)+1)^(a-1)+4*Pi^(1/2)/(1+a)/a*x^(1+a)*b^(-1/2*a-1/2)*(1+b/x^2)^(1/2)*((1+b/x^2)^(1/2)+1)^(a-1))`

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.62

$$\int \left(x + \sqrt{b + x^2}\right)^a dx = \frac{(\sqrt{x^2 + ba} - x)(x + \sqrt{x^2 + b})^a}{a^2 - 1}$$

input `integrate((x+(x^2+b)^(1/2))^a,x, algorithm="fricas")`

output `(sqrt(x^2 + b)*a - x)*(x + sqrt(x^2 + b))^a/(a^2 - 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2236 vs. 2(37) = 74.

Time = 1.16 (sec) , antiderivative size = 2236, normalized size of antiderivative = 43.00

$$\int \left(x + \sqrt{b + x^2}\right)^a dx = \text{Too large to display}$$

input `integrate((x+(x**2+b)**(1/2))**a,x)`

output

```
Piecewise((-a**2*b**4*b**(a/2 + 1/2)*x*sqrt(b/x**2 + 1)*sinh(a*asinh(x/sqr
t(b)))*gamma(-a/2)/(2*a**2*b**(9/2)*gamma(1 - a/2) + 2*a**2*b**(7/2)*x**2*
gamma(1 - a/2) - 2*b**(9/2)*gamma(1 - a/2) - 2*b**(7/2)*x**2*gamma(1 - a/2
)) - a**2*b**3*b**(a/2 + 1/2)*x**3*sqrt(b/x**2 + 1)*sinh(a*asinh(x/sqrt(b)
))*gamma(-a/2)/(2*a**2*b**(9/2)*gamma(1 - a/2) + 2*a**2*b**(7/2)*x**2*gamma
a(1 - a/2) - 2*b**(9/2)*gamma(1 - a/2) - 2*b**(7/2)*x**2*gamma(1 - a/2)) +
2*a*b**(9/2)*b**(a/2 + 1/2)*cosh(a*asinh(x/sqrt(b)) + asinh(x/sqrt(b)))*g
amma(1 - a/2)/(2*a**2*b**(9/2)*gamma(1 - a/2) + 2*a**2*b**(7/2)*x**2*gamma
(1 - a/2) - 2*b**(9/2)*gamma(1 - a/2) - 2*b**(7/2)*x**2*gamma(1 - a/2)) -
2*a*b**(9/2)*b**(a/2 + 1/2)*gamma(1 - a/2)/(2*a**2*b**(9/2)*gamma(1 - a/2)
+ 2*a**2*b**(7/2)*x**2*gamma(1 - a/2) - 2*b**(9/2)*gamma(1 - a/2) - 2*b**
(7/2)*x**2*gamma(1 - a/2)) - 2*a*b**(7/2)*b**(a/2 + 1/2)*x**2*sqrt(b/x**2
+ 1)*sinh(a*asinh(x/sqrt(b)) + asinh(x/sqrt(b)))*gamma(1 - a/2)/(2*a**2*b*
*(9/2)*gamma(1 - a/2) + 2*a**2*b**(7/2)*x**2*gamma(1 - a/2) - 2*b**(9/2)*g
amma(1 - a/2) - 2*b**(7/2)*x**2*gamma(1 - a/2)) + 4*a*b**(7/2)*b**(a/2 + 1
/2)*x**2*cosh(a*asinh(x/sqrt(b)) + asinh(x/sqrt(b)))*gamma(1 - a/2)/(2*a**
2*b**(9/2)*gamma(1 - a/2) + 2*a**2*b**(7/2)*x**2*gamma(1 - a/2) - 2*b**(9/
2)*gamma(1 - a/2) - 2*b**(7/2)*x**2*gamma(1 - a/2)) - 2*a*b**(7/2)*b**(a/2
+ 1/2)*x**2*gamma(1 - a/2)/(2*a**2*b**(9/2)*gamma(1 - a/2) + 2*a**2*b**(7
/2)*x**2*gamma(1 - a/2) - 2*b**(9/2)*gamma(1 - a/2) - 2*b**(7/2)*x**2*g...
```

Maxima [F]

$$\int (x + \sqrt{b + x^2})^a dx = \int (x + \sqrt{x^2 + b})^a dx$$

input

```
integrate((x+(x^2+b)^(1/2))^a,x, algorithm="maxima")
```

output

```
integrate((x + sqrt(x^2 + b))^a, x)
```

Giac [F]

$$\int (x + \sqrt{b + x^2})^a dx = \int (x + \sqrt{x^2 + b})^a dx$$

input `integrate((x+(x^2+b)^(1/2))^a,x, algorithm="giac")`

output `integrate((x + sqrt(x^2 + b))^a, x)`

Mupad [F(-1)]

Timed out.

$$\int (x + \sqrt{b + x^2})^a dx = \int (x + \sqrt{x^2 + b})^a dx$$

input `int((x + (b + x^2)^(1/2))^a,x)`

output `int((x + (b + x^2)^(1/2))^a, x)`

Reduce [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.58

$$\int (x + \sqrt{b + x^2})^a dx = \frac{(\sqrt{x^2 + b} + x)^a (\sqrt{x^2 + b} a - x)}{a^2 - 1}$$

input `int((x+(x^2+b)^(1/2))^a,x)`

output `((sqrt(b + x**2) + x)**a*(sqrt(b + x**2)*a - x))/(a**2 - 1)`

3.32 $\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx$

Optimal result	263
Mathematica [A] (verified)	263
Rubi [A] (verified)	264
Maple [A] (verified)	265
Fricas [A] (verification not implemented)	265
Sympy [F(-1)]	266
Maxima [A] (verification not implemented)	266
Giac [F]	266
Mupad [F(-1)]	267
Reduce [B] (verification not implemented)	267

Optimal result

Integrand size = 33, antiderivative size = 34

$$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx = \frac{x^{1+a}(6 + 3x^a + 2x^{2a})^{1+\frac{1}{a}}}{6(1+a)}$$

output $1/6*x^{(1+a)}*(6+3*x^a+2*x^{(2*a)})^{(1+1/a)/(1+a)}$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.97

$$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx = \frac{x^{1+a}(6 + 3x^a + 2x^{2a})^{1+\frac{1}{a}}}{6 + 6a}$$

input $\text{Integrate}[(6 + 3*x^a + 2*x^{(2*a)})^a*(x^a + x^{(2*a)} + x^{(3*a)}), x]$

output $(x^{(1+a)}*(6 + 3*x^a + 2*x^{(2*a)})^{(1+a^{-1})})/(6 + 6*a)$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.061$, Rules used = {2028, 2285}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (2x^{2a} + 3x^a + 6)^{\frac{1}{a}} (x^{2a} + x^{3a} + x^a) dx$$

$$\downarrow \text{2028}$$

$$\int x^a (x^{2a} + x^a + 1) (2x^{2a} + 3x^a + 6)^{\frac{1}{a}} dx$$

$$\downarrow \text{2285}$$

$$\frac{x^{a+1} (2x^{2a} + 3x^a + 6)^{\frac{1}{a}+1}}{6(a+1)}$$

input `Int[(6 + 3*x^a + 2*x^(2*a))^a^(-1)*(x^a + x^(2*a) + x^(3*a)),x]`

output `(x^(1 + a)*(6 + 3*x^a + 2*x^(2*a))^(1 + a^(-1)))/(6*(1 + a))`

Defintions of rubi rules used

rule 2028 `Int[(Fx_.)*((a_.)*(x_)^(r_.) + (b_.)*(x_)^(s_.) + (c_.)*(x_)^(t_.))^(p_.), x_Symbol] := Int[x^(p*r)*(a + b*x^(s - r) + c*x^(t - r))^p*Fx, x] /; FreeQ[{a, b, c, r, s, t}, x] && IntegerQ[p] && PosQ[s - r] && PosQ[t - r] && !(EqQ[p, 1] && EqQ[u, 1])`

rule 2285

```
Int[((g_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.))^(p_.)*
((d_) + (e_.)*(x_)^(n_.) + (f_.)*(x_)^(n2_.)), x_Symbol] :> Simp[d*(g*x)^(m
+ 1)*((a + b*x^n + c*x^(2*n))^(p + 1)/(a*g*(m + 1))), x] /; FreeQ[{a, b, c
, d, e, f, g, m, n, p}, x] && EqQ[n2, 2*n] && EqQ[a*e*(m + 1) - b*d*(m + n*
(p + 1) + 1), 0] && EqQ[a*f*(m + 1) - c*d*(m + 2*n*(p + 1) + 1), 0] && NeQ[
m, -1]
```

Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29

method	result	size
risch	$\frac{x x^a (6+3x^a+2x^{2a})(6+3x^a+2x^{2a})^{\frac{1}{a}}}{6+6a}$	44

input

```
int((6+3*x^a+2*x^(2*a))^(1/a)*(x^a+x^(2*a)+x^(3*a)),x,method=_RETURNVERBOS
E)
```

output

```
1/6*x*x^a*(6+3*x^a+2*(x^a)^2)/(1+a)*(6+3*x^a+2*(x^a)^2)^(1/a)
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.41

$$\int (6+3x^a+2x^{2a})^{\frac{1}{a}} (x^a+x^{2a}+x^{3a}) dx = \frac{(2xx^{3a} + 3xx^{2a} + 6xx^a)(2x^{2a} + 3x^a + 6)^{\left(\frac{1}{a}\right)}}{6(a+1)}$$

input

```
integrate((6+3*x^a+2*x^(2*a))^(1/a)*(x^a+x^(2*a)+x^(3*a)),x, algorithm="fr
icas")
```

output

```
1/6*(2*x*x^(3*a) + 3*x*x^(2*a) + 6*x*x^a)*(2*x^(2*a) + 3*x^a + 6)^(1/a)/(a
+ 1)
```

Sympy [F(-1)]

Timed out.

$$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx = \text{Timed out}$$

input `integrate((6+3*x**a+2*x**(2*a))**(1/a)*(x**a+x**(2*a)+x**(3*a)),x)`

output `Timed out`

Maxima [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.41

$$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx = \frac{(2xx^{3a} + 3xx^{2a} + 6xx^a)(2x^{2a} + 3x^a + 6)^{\left(\frac{1}{a}\right)}}{6(a+1)}$$

input `integrate((6+3*x^a+2*x^(2*a))^(1/a)*(x^a+x^(2*a)+x^(3*a)),x, algorithm="maxima")`

output `1/6*(2*x*x^(3*a) + 3*x*x^(2*a) + 6*x*x^a)*(2*x^(2*a) + 3*x^a + 6)^(1/a)/(a + 1)`

Giac [F]

$$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx = \int (2x^{2a} + 3x^a + 6)^{\left(\frac{1}{a}\right)} (x^{3a} + x^{2a} + x^a) dx$$

input `integrate((6+3*x^a+2*x^(2*a))^(1/a)*(x^a+x^(2*a)+x^(3*a)),x, algorithm="giac")`

output `integrate((2*x^(2*a) + 3*x^a + 6)^(1/a)*(x^(3*a) + x^(2*a) + x^a), x)`

Mupad [F(-1)]

Timed out.

$$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx = \int (x^a + x^{2a} + x^{3a}) (3x^a + 2x^{2a} + 6)^{1/a} dx$$

input `int((x^a + x^(2*a) + x^(3*a))*(3*x^a + 2*x^(2*a) + 6)^(1/a), x)`

output `int((x^a + x^(2*a) + x^(3*a))*(3*x^a + 2*x^(2*a) + 6)^(1/a), x)`

Reduce [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29

$$\int (6 + 3x^a + 2x^{2a})^{\frac{1}{a}} (x^a + x^{2a} + x^{3a}) dx = \frac{x^a(2x^{2a} + 3x^a + 6)^{\frac{1}{a}} x(2x^{2a} + 3x^a + 6)}{6a + 6}$$

input `int((6+3*x^a+2*x^(2*a))^(1/a)*(x^a+x^(2*a)+x^(3*a)), x)`

output `(x**a*(2*x**(2*a) + 3*x**a + 6)**(1/a)*x*(2*x**(2*a) + 3*x**a + 6))/(6*(a + 1))`

3.33 $\int \frac{1}{x \sqrt[3]{1-x^2}} dx$

Optimal result	268
Mathematica [A] (verified)	268
Rubi [A] (verified)	269
Maple [A] (verified)	271
Fricas [A] (verification not implemented)	271
Sympy [C] (verification not implemented)	272
Maxima [A] (verification not implemented)	272
Giac [A] (verification not implemented)	273
Mupad [B] (verification not implemented)	273
Reduce [F]	274

Optimal result

Integrand size = 15, antiderivative size = 58

$$\int \frac{1}{x \sqrt[3]{1-x^2}} dx = \frac{1}{2} \sqrt{3} \arctan \left(\frac{1 + 2\sqrt[3]{1-x^2}}{\sqrt{3}} \right) - \frac{\log(x)}{2} + \frac{3}{4} \log \left(1 - \sqrt[3]{1-x^2} \right)$$

output

```
-1/2*ln(x)+3/4*ln(1-(-x^2+1)^(1/3))+1/2*arctan(1/3*(1+2*(-x^2+1)^(1/3))*3^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.33

$$\int \frac{1}{x \sqrt[3]{1-x^2}} dx = \frac{1}{4} \left(2\sqrt{3} \arctan \left(\frac{1 + 2\sqrt[3]{1-x^2}}{\sqrt{3}} \right) + 2 \log \left(-1 + \sqrt[3]{1-x^2} \right) - \log \left(1 + \sqrt[3]{1-x^2} + (1-x^2)^{2/3} \right) \right)$$

input

```
Integrate[1/(x*(1-x^2)^(1/3)),x]
```

output

```
(2*Sqrt[3]*ArcTan[(1 + 2*(1 - x^2)^(1/3))/Sqrt[3]] + 2*Log[-1 + (1 - x^2)^(1/3)] - Log[1 + (1 - x^2)^(1/3) + (1 - x^2)^(2/3)])/4
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {243, 67, 16, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt[3]{1-x^2}} dx$$

$$\downarrow 243$$

$$\frac{1}{2} \int \frac{1}{x^2\sqrt[3]{1-x^2}} dx^2$$

$$\downarrow 67$$

$$\frac{1}{2} \left(-\frac{3}{2} \int \frac{1}{1-\sqrt[3]{1-x^2}} d\sqrt[3]{1-x^2} + \frac{3}{2} \int \frac{1}{x^4 + \sqrt[3]{1-x^2} + 1} d\sqrt[3]{1-x^2} - \frac{1}{2} \log(x^2) \right)$$

$$\downarrow 16$$

$$\frac{1}{2} \left(\frac{3}{2} \int \frac{1}{x^4 + \sqrt[3]{1-x^2} + 1} d\sqrt[3]{1-x^2} - \frac{1}{2} \log(x^2) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^2}) \right)$$

$$\downarrow 1083$$

$$\frac{1}{2} \left(-3 \int \frac{1}{-x^4 - 3} d(2\sqrt[3]{1-x^2} + 1) - \frac{1}{2} \log(x^2) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^2}) \right)$$

$$\downarrow 217$$

$$\frac{1}{2} \left(\sqrt{3} \arctan \left(\frac{2\sqrt[3]{1-x^2} + 1}{\sqrt{3}} \right) - \frac{\log(x^2)}{2} + \frac{3}{2} \log(1 - \sqrt[3]{1-x^2}) \right)$$

input

```
Int[1/(x*(1 - x^2)^(1/3)),x]
```

output $(\sqrt{3} \operatorname{ArcTan}[(1 + 2(1 - x^2)^{1/3})/\sqrt{3}] - \operatorname{Log}[x^2]/2 + (3 \operatorname{Log}[1 - (1 - x^2)^{1/3}])/2)/2$

Defintions of rubi rules used

rule 16 $\operatorname{Int}[(c_./((a_.) + (b_.)*(x_))), x_Symbol] \rightarrow \operatorname{Simp}[c*(\operatorname{Log}[\operatorname{RemoveContent}[a + b*x, x]]/b), x] /; \operatorname{FreeQ}[\{a, b, c\}, x]$

rule 67 $\operatorname{Int}[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^{1/3}), x_Symbol] \rightarrow \operatorname{With}[\{q = \operatorname{Rt}[(b*c - a*d)/b, 3]\}, \operatorname{Simp}[-\operatorname{Log}[\operatorname{RemoveContent}[a + b*x, x]]/(2*b*q), x] + (\operatorname{Simp}[3/(2*b) \operatorname{Subst}[\operatorname{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{1/3}], x] - \operatorname{Simp}[3/(2*b*q) \operatorname{Subst}[\operatorname{Int}[1/(q - x), x], x, (c + d*x)^{1/3}], x]) /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{PosQ}[(b*c - a*d)/b]$

rule 217 $\operatorname{Int}(((a_.) + (b_.)*(x_)^2)^{-1}), x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[-b, 2])^{-1})*\operatorname{ArcTan}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \& \& (\operatorname{LtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$

rule 243 $\operatorname{Int}[(x_)^{(m_.)}*((a_.) + (b_.)*(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[1/2 \operatorname{Subst}[\operatorname{Int}[x^{(m-1)/2}*(a + b*x)^p, x], x, x^2], x] /; \operatorname{FreeQ}[\{a, b, m, p\}, x] \&\& \operatorname{IntegerQ}[(m-1)/2]$

rule 1083 $\operatorname{Int}(((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^{-1}), x_Symbol] \rightarrow \operatorname{Simp}[-2 \operatorname{Subst}[\operatorname{Int}[1/\operatorname{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \operatorname{FreeQ}[\{a, b, c\}, x]$

Maple [A] (verified)

Time = 1.59 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

method	result
pseudoelliptic	$-\frac{\ln\left((-x^2+1)^{\frac{2}{3}}+(-x^2+1)^{\frac{1}{3}}+1\right)}{4} + \frac{\arctan\left(\frac{(1+2(-x^2+1)^{\frac{1}{3}})\sqrt{3}}{3}\right)\sqrt{3}}{2} + \frac{\ln\left((-x^2+1)^{\frac{1}{3}}-1\right)}{2}$
meijerg	$\frac{\sqrt{3}\Gamma\left(\frac{2}{3}\right)\left(\frac{2\left(-\frac{\pi\sqrt{3}}{6}-\frac{3\ln(3)}{2}+2\ln(x)+i\pi\right)\pi\sqrt{3}}{3\Gamma\left(\frac{2}{3}\right)} + \frac{2\pi\sqrt{3}x^2\operatorname{hypergeom}\left(\left[1,1,\frac{4}{3}\right],[2,2],x^2\right)}{9\Gamma\left(\frac{2}{3}\right)}\right)}{4\pi}$
trager	$\operatorname{RootOf}\left(-Z^2+Z+1\right)\ln\left(-\frac{\operatorname{RootOf}\left(-Z^2+Z+1\right)^2x^2+15\operatorname{RootOf}\left(-Z^2+Z+1\right)\left(-x^2+1\right)^{\frac{2}{3}}-\operatorname{RootOf}\left(-Z^2+Z+1\right)}{\dots}\right)$

input `int(1/x/(-x^2+1)^(1/3),x,method=_RETURNVERBOSE)`output `-1/4*ln((-x^2+1)^(2/3)+(-x^2+1)^(1/3)+1)+1/2*arctan(1/3*(1+2*(-x^2+1)^(1/3)))*3^(1/2)*3^(1/2)+1/2*ln((-x^2+1)^(1/3)-1)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

$$\int \frac{1}{x\sqrt[3]{1-x^2}} dx = \frac{1}{2} \sqrt{3} \arctan\left(\frac{2}{3} \sqrt{3}(-x^2+1)^{\frac{1}{3}} + \frac{1}{3} \sqrt{3}\right) - \frac{1}{4} \log\left(\left(-x^2+1\right)^{\frac{2}{3}} + \left(-x^2+1\right)^{\frac{1}{3}} + 1\right) + \frac{1}{2} \log\left(\left(-x^2+1\right)^{\frac{1}{3}} - 1\right)$$

input `integrate(1/x/(-x^2+1)^(1/3),x, algorithm="fricas")`output `1/2*sqrt(3)*arctan(2/3*sqrt(3)*(-x^2 + 1)^(1/3) + 1/3*sqrt(3)) - 1/4*log((-x^2 + 1)^(2/3) + (-x^2 + 1)^(1/3) + 1) + 1/2*log((-x^2 + 1)^(1/3) - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.62

$$\int \frac{1}{x\sqrt{1-x^2}} dx = -\frac{e^{-\frac{i\pi}{3}}\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{4}{3} \middle| \frac{1}{x^2}\right)}{2x^{\frac{2}{3}}\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/x/(-x**2+1)**(1/3),x)`

output `-exp(-I*pi/3)*gamma(1/3)*hyper((1/3, 1/3), (4/3,), x**(-2))/(2*x**(2/3)*gamma(4/3))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.07

$$\int \frac{1}{x\sqrt{1-x^2}} dx = \frac{1}{2}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}\left(2(-x^2+1)^{\frac{1}{3}}+1\right)\right) - \frac{1}{4}\log\left(\left(-x^2+1\right)^{\frac{2}{3}}+\left(-x^2+1\right)^{\frac{1}{3}}+1\right) + \frac{1}{2}\log\left(\left(-x^2+1\right)^{\frac{1}{3}}-1\right)$$

input `integrate(1/x/(-x^2+1)^(1/3),x, algorithm="maxima")`

output `1/2*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^2 + 1)^(1/3) + 1)) - 1/4*log((-x^2 + 1)^(2/3) + (-x^2 + 1)^(1/3) + 1) + 1/2*log((-x^2 + 1)^(1/3) - 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

$$\int \frac{1}{x\sqrt[3]{1-x^2}} dx = \frac{1}{2} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \left(2(-x^2+1)^{\frac{1}{3}} + 1 \right) \right) - \frac{1}{4} \log \left((-x^2+1)^{\frac{2}{3}} + (-x^2+1)^{\frac{1}{3}} + 1 \right) + \frac{1}{2} \log \left(-(-x^2+1)^{\frac{1}{3}} + 1 \right)$$

input `integrate(1/x/(-x^2+1)^(1/3),x, algorithm="giac")`output `1/2*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^2 + 1)^(1/3) + 1)) - 1/4*log((-x^2 + 1)^(2/3) + (-x^2 + 1)^(1/3) + 1) + 1/2*log(-(-x^2 + 1)^(1/3) + 1)`**Mupad [B] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.48

$$\int \frac{1}{x\sqrt[3]{1-x^2}} dx = \frac{\ln \left(\frac{9(1-x^2)^{1/3}}{4} - \frac{9}{4} \right)}{2} + \ln \left(\frac{9(1-x^2)^{1/3}}{4} - 9 \left(-\frac{1}{4} + \frac{\sqrt{3} \text{li}}{4} \right)^2 \right) \left(-\frac{1}{4} + \frac{\sqrt{3} \text{li}}{4} \right) - \ln \left(\frac{9(1-x^2)^{1/3}}{4} - 9 \left(\frac{1}{4} + \frac{\sqrt{3} \text{li}}{4} \right)^2 \right) \left(\frac{1}{4} + \frac{\sqrt{3} \text{li}}{4} \right)$$

input `int(1/(x*(1 - x^2)^(1/3)),x)`output `log((9*(1 - x^2)^(1/3))/4 - 9/4)/2 + log((9*(1 - x^2)^(1/3))/4 - 9*((3^(1/2)*1i)/4 - 1/4)^2)*((3^(1/2)*1i)/4 - 1/4) - log((9*(1 - x^2)^(1/3))/4 - 9*((3^(1/2)*1i)/4 + 1/4)^2)*((3^(1/2)*1i)/4 + 1/4)`

Reduce [F]

$$\int \frac{1}{x\sqrt[3]{1-x^2}} dx = \int \frac{1}{(-x^2+1)^{\frac{1}{3}}x} dx$$

input `int(1/x/(-x^2+1)^(1/3),x)`

output `int(1/((-x**2+1)**(1/3)*x),x)`

3.34 $\int \frac{1}{x(1-x^2)^{2/3}} dx$

Optimal result	275
Mathematica [A] (verified)	275
Rubi [A] (verified)	276
Maple [C] (verified)	278
Fricas [A] (verification not implemented)	278
Sympy [C] (verification not implemented)	279
Maxima [A] (verification not implemented)	279
Giac [A] (verification not implemented)	280
Mupad [B] (verification not implemented)	280
Reduce [F]	281

Optimal result

Integrand size = 15, antiderivative size = 58

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = -\frac{1}{2}\sqrt{3} \arctan\left(\frac{1+2\sqrt[3]{1-x^2}}{\sqrt{3}}\right) - \frac{\log(x)}{2} + \frac{3}{4} \log\left(1 - \sqrt[3]{1-x^2}\right)$$

output

$-1/2*\ln(x)+3/4*\ln(1-(-x^2+1)^{(1/3)})-1/2*\arctan(1/3*(1+2*(-x^2+1)^{(1/3)))*3^{(1/2)})*3^{(1/2)}$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.33

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = \frac{1}{4} \left(-2\sqrt{3} \arctan\left(\frac{1+2\sqrt[3]{1-x^2}}{\sqrt{3}}\right) + 2 \log\left(-1 + \sqrt[3]{1-x^2}\right) - \log\left(1 + \sqrt[3]{1-x^2} + (1-x^2)^{2/3}\right) \right)$$

input

`Integrate[1/(x*(1 - x^2)^(2/3)),x]`

output

$$\frac{(-2\sqrt{3}\operatorname{ArcTan}[(1 + 2(1 - x^2)^{1/3})/\sqrt{3}] + 2\operatorname{Log}[-1 + (1 - x^2)^{1/3}] - \operatorname{Log}[1 + (1 - x^2)^{1/3} + (1 - x^2)^{2/3}])}{4}$$
Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.07, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {243, 69, 16, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{x(1-x^2)^{2/3}} dx \\ & \quad \downarrow \text{243} \\ & \frac{1}{2} \int \frac{1}{x^2(1-x^2)^{2/3}} dx^2 \\ & \quad \downarrow \text{69} \\ & \frac{1}{2} \left(-\frac{3}{2} \int \frac{1}{1-\sqrt[3]{1-x^2}} d\sqrt[3]{1-x^2} - \frac{3}{2} \int \frac{1}{x^4 + \sqrt[3]{1-x^2} + 1} d\sqrt[3]{1-x^2} - \frac{1}{2} \log(x^2) \right) \\ & \quad \downarrow \text{16} \\ & \frac{1}{2} \left(-\frac{3}{2} \int \frac{1}{x^4 + \sqrt[3]{1-x^2} + 1} d\sqrt[3]{1-x^2} - \frac{1}{2} \log(x^2) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^2}) \right) \\ & \quad \downarrow \text{1083} \\ & \frac{1}{2} \left(3 \int \frac{1}{-x^4 - 3} d(2\sqrt[3]{1-x^2} + 1) - \frac{1}{2} \log(x^2) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^2}) \right) \\ & \quad \downarrow \text{217} \\ & \frac{1}{2} \left(-\sqrt{3} \arctan\left(\frac{2\sqrt[3]{1-x^2} + 1}{\sqrt{3}}\right) - \frac{\log(x^2)}{2} + \frac{3}{2} \log(1 - \sqrt[3]{1-x^2}) \right) \end{aligned}$$

input

$$\operatorname{Int}[1/(x*(1-x^2)^{2/3}), x]$$

output $(-\sqrt{3} \operatorname{ArcTan}[(1 + 2(1 - x^2)^{1/3})/\sqrt{3}]) - \operatorname{Log}[x^2]/2 + (3 \operatorname{Log}[1 - (1 - x^2)^{1/3}])/2)/2$

Defintions of rubi rules used

rule 16 $\operatorname{Int}[(c_./((a_.) + (b_.)(x_)), x_Symbol] \rightarrow \operatorname{Simp}[c*(\operatorname{Log}[\operatorname{RemoveContent}[a + b*x, x]]/b), x] /; \operatorname{FreeQ}[\{a, b, c\}, x]$

rule 69 $\operatorname{Int}[1/(((a_.) + (b_.)(x_))*((c_.) + (d_.)(x_))^{2/3}), x_Symbol] \rightarrow \operatorname{With}[\{q = \operatorname{Rt}[(b*c - a*d)/b, 3]\}, \operatorname{Simp}[-\operatorname{Log}[\operatorname{RemoveContent}[a + b*x, x]]/(2*b*q^2), x] + (-\operatorname{Simp}[3/(2*b*q) \operatorname{Subst}[\operatorname{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{1/3}], x] - \operatorname{Simp}[3/(2*b*q^2) \operatorname{Subst}[\operatorname{Int}[1/(q - x), x], x, (c + d*x)^{1/3}], x])]/; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{PosQ}[(b*c - a*d)/b]$

rule 217 $\operatorname{Int}(((a_.) + (b_.)(x_)^2)^{-1}), x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[-b, 2])^{-1})*\operatorname{ArcTan}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \& \& (\operatorname{LtQ}[a, 0] \mid \mid \operatorname{LtQ}[b, 0])]$

rule 243 $\operatorname{Int}[(x_)^{(m_.)*((a_.) + (b_.)(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[1/2 \operatorname{Subst}[\operatorname{Int}[x^{((m - 1)/2)*(a + b*x)^p}, x], x, x^2], x] /; \operatorname{FreeQ}[\{a, b, m, p\}, x] \&\& \operatorname{IntegerQ}[(m - 1)/2]$

rule 1083 $\operatorname{Int}(((a_.) + (b_.)(x_) + (c_.)(x_)^2)^{-1}), x_Symbol] \rightarrow \operatorname{Simp}[-2 \operatorname{Subst}[\operatorname{Int}[1/\operatorname{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \operatorname{FreeQ}[\{a, b, c\}, x]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 1.63 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.83

method	result
meijerg	$\frac{\left(\frac{\pi\sqrt{3}}{6} - \frac{3\ln(3)}{2} + 2\ln(x) + i\pi\right)\Gamma\left(\frac{2}{3}\right) + \frac{2\Gamma\left(\frac{2}{3}\right)x^2 \operatorname{hypergeom}\left(\left[1, 1, \frac{5}{3}\right], [2, 2], x^2\right)}{3}}{2\Gamma\left(\frac{2}{3}\right)}$
pseudoelliptic	$-\frac{\ln\left((-x^2+1)^{\frac{2}{3}} + (-x^2+1)^{\frac{1}{3}} + 1\right)}{4} - \frac{\arctan\left(\frac{(1+2(-x^2+1)^{\frac{1}{3}})\sqrt{3}}{3}\right)\sqrt{3}}{2} + \frac{\ln\left((-x^2+1)^{\frac{1}{3}} - 1\right)}{2}$
trager	$\ln\left(-\frac{4\operatorname{RootOf}\left(-Z^2+Z+1\right)^2x^2 - 15\operatorname{RootOf}\left(-Z^2+Z+1\right)(-x^2+1)^{\frac{2}{3}} - 9\operatorname{RootOf}\left(-Z^2+Z+1\right)x^2 - 16\operatorname{RootOf}\left(-Z^2+Z+1\right)}{x^2}\right)$

input `int(1/x/(-x^2+1)^(2/3), x, method=_RETURNVERBOSE)`

output `1/2/GAMMA(2/3)*((1/6*Pi*3^(1/2)-3/2*ln(3)+2*ln(x)+I*Pi)*GAMMA(2/3)+2/3*GAMMA(2/3)*x^2*hypergeom([1,1,5/3],[2,2],x^2))`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = -\frac{1}{2}\sqrt{3} \arctan\left(\frac{2}{3}\sqrt{3}(-x^2+1)^{\frac{1}{3}} + \frac{1}{3}\sqrt{3}\right) - \frac{1}{4} \log\left((-x^2+1)^{\frac{2}{3}} + (-x^2+1)^{\frac{1}{3}} + 1\right) + \frac{1}{2} \log\left((-x^2+1)^{\frac{1}{3}} - 1\right)$$

input `integrate(1/x/(-x^2+1)^(2/3), x, algorithm="fricas")`

output `-1/2*sqrt(3)*arctan(2/3*sqrt(3)*(-x^2+1)^(1/3)+1/3*sqrt(3))-1/4*log((-x^2+1)^(2/3)+(-x^2+1)^(1/3)+1)+1/2*log((-x^2+1)^(1/3)-1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.64

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = -\frac{e^{-\frac{2i\pi}{3}} \Gamma\left(\frac{2}{3}\right) {}_2F_1\left(\frac{2}{3}, \frac{2}{3} \middle| \frac{1}{x^2}\right)}{2x^{4/3} \Gamma\left(\frac{5}{3}\right)}$$

input `integrate(1/x/(-x**2+1)**(2/3),x)`

output `-exp(-2*I*pi/3)*gamma(2/3)*hyper((2/3, 2/3), (5/3,), x**(-2))/(2*x**(4/3)*gamma(5/3))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.07

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = -\frac{1}{2} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(2(-x^2+1)^{1/3} + 1\right)\right) - \frac{1}{4} \log\left(\left(-x^2+1\right)^{2/3} + \left(-x^2+1\right)^{1/3} + 1\right) + \frac{1}{2} \log\left(\left(-x^2+1\right)^{1/3} - 1\right)$$

input `integrate(1/x/(-x^2+1)^(2/3),x, algorithm="maxima")`

output `-1/2*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^2 + 1)^(1/3) + 1)) - 1/4*log((-x^2 + 1)^(2/3) + (-x^2 + 1)^(1/3) + 1) + 1/2*log((-x^2 + 1)^(1/3) - 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = -\frac{1}{2}\sqrt{3} \arctan\left(\frac{1}{3}\sqrt{3}\left(2(-x^2+1)^{1/3}+1\right)\right) - \frac{1}{4} \log\left(\left(-x^2+1\right)^{2/3} + \left(-x^2+1\right)^{1/3} + 1\right) + \frac{1}{2} \log\left(-\left(-x^2+1\right)^{1/3} + 1\right)$$

input `integrate(1/x/(-x^2+1)^(2/3),x, algorithm="giac")`output `-1/2*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^2 + 1)^(1/3) + 1)) - 1/4*log((-x^2 + 1)^(2/3) + (-x^2 + 1)^(1/3) + 1) + 1/2*log(-(-x^2 + 1)^(1/3) + 1)`**Mupad [B] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.31

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = \frac{\ln\left(\frac{9(1-x^2)^{1/3}}{4} - \frac{9}{4}\right)}{2} + \ln\left(\frac{9(1-x^2)^{1/3}}{2} + \frac{9}{4} - \frac{\sqrt{3}9i}{4}\right) \left(-\frac{1}{4} + \frac{\sqrt{3}1i}{4}\right) - \ln\left(\frac{9(1-x^2)^{1/3}}{2} + \frac{9}{4} + \frac{\sqrt{3}9i}{4}\right) \left(\frac{1}{4} + \frac{\sqrt{3}1i}{4}\right)$$

input `int(1/(x*(1 - x^2)^(2/3)),x)`output `log((9*(1 - x^2)^(1/3))/4 - 9/4)/2 + log((9*(1 - x^2)^(1/3))/2 - (3^(1/2))*9i)/4 + 9/4)*((3^(1/2)*1i)/4 - 1/4) - log((3^(1/2)*9i)/4 + (9*(1 - x^2)^(1/3))/2 + 9/4)*((3^(1/2)*1i)/4 + 1/4)`

Reduce **[F]**

$$\int \frac{1}{x(1-x^2)^{2/3}} dx = \int \frac{1}{(-x^2+1)^{2/3} x} dx$$

input `int(1/x/(-x^2+1)^(2/3),x)`

output `int(1/((-x**2+1)**(2/3)*x),x)`

3.35 $\int \frac{1}{\sqrt[3]{1-x^3}} dx$

Optimal result	282
Mathematica [A] (verified)	282
Rubi [A] (verified)	283
Maple [C] (verified)	284
Fricas [B] (verification not implemented)	284
Sympy [C] (verification not implemented)	285
Maxima [A] (verification not implemented)	285
Giac [F]	286
Mupad [B] (verification not implemented)	286
Reduce [F]	286

Optimal result

Integrand size = 11, antiderivative size = 49

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = -\frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2} \log\left(x + \sqrt[3]{1-x^3}\right)$$

output

$1/2*\ln(x+(-x^3+1)^{(1/3)})-1/3*\arctan(1/3*(1-2*x/(-x^3+1)^{(1/3}))*3^{(1/2)})*3^{(1/2)}$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.76

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = \frac{\arctan\left(\frac{-1+\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{6} \log\left(1 + \frac{x^2}{(1-x^3)^{2/3}} - \frac{x}{\sqrt[3]{1-x^3}}\right) + \frac{1}{3} \log\left(1 + \frac{x}{\sqrt[3]{1-x^3}}\right)$$

input

`Integrate[(1 - x^3)^(-1/3), x]`

output

```
ArcTan[(-1 + (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - Log[1 + x^2/(1 - x^3)^(2/3) - x/(1 - x^3)^(1/3)]/6 + Log[1 + x/(1 - x^3)^(1/3)]/3
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx$$

↓ 769

$$\frac{1}{2} \log\left(\sqrt[3]{1-x^3} + x\right) - \frac{\arctan\left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}}$$

input

```
Int[(1 - x^3)^(-1/3), x]
```

output

```
-(ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3]) + Log[x + (1 - x^3)^(1/3)]/2
```

Defintions of rubi rules used

rule 769

```
Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] :=> Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 2.07 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.24

method	result
meijerg	$x \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{1}{3}\right], \left[\frac{4}{3}\right], x^3\right)$
pseudoelliptic	$-\frac{\ln\left(\frac{(-x^3+1)^{\frac{2}{3}}-x(-x^3+1)^{\frac{1}{3}}+x^2}{x^2}\right)}{6} + \frac{\sqrt{3} \arctan\left(\frac{(-2(-x^3+1)^{\frac{1}{3}}+x)\sqrt{3}}{3x}\right)}{3} + \frac{\ln\left(\frac{x+(-x^3+1)^{\frac{1}{3}}}{x}\right)}{3}$
trager	$\frac{\operatorname{RootOf}(_Z^2 + _Z + 1) \ln\left(-\operatorname{RootOf}(_Z^2 + _Z + 1)^2 x^3 - \operatorname{RootOf}(_Z^2 + _Z + 1) x^3 + 3x(-x^3+1)^{\frac{2}{3}} - 3x^2(-x^3+1)\right)}{3}$

input `int(1/(-x^3+1)^(1/3), x, method=_RETURNVERBOSE)`

output `x*hypergeom([1/3, 1/3], [4/3], x^3)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(40) = 80$.

Time = 0.08 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.67

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = -\frac{1}{3} \sqrt{3} \arctan\left(-\frac{\sqrt{3}x - 2\sqrt{3}(-x^3+1)^{\frac{1}{3}}}{3x}\right) + \frac{1}{3} \log\left(\frac{x + (-x^3+1)^{\frac{1}{3}}}{x}\right) - \frac{1}{6} \log\left(\frac{x^2 - (-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{2}{3}}}{x^2}\right)$$

input `integrate(1/(-x^3+1)^(1/3), x, algorithm="fricas")`

output

```
-1/3*sqrt(3)*arctan(-1/3*(sqrt(3)*x - 2*sqrt(3)*(-x^3 + 1)^(1/3))/x) + 1/3
*log((x + (-x^3 + 1)^(1/3))/x) - 1/6*log((x^2 - (-x^3 + 1)^(1/3)*x + (-x^3
+ 1)^(2/3))/x^2)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.39 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.59

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{4}{3} \middle| x^3 e^{2i\pi}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input

```
integrate(1/(-x**3+1)**(1/3),x)
```

output

```
x*gamma(1/3)*hyper((1/3, 1/3), (4/3,), x**3*exp_polar(2*I*pi))/(3*gamma(4/
3))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.59

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(\frac{2(-x^3+1)^{\frac{1}{3}}}{x} - 1\right)\right) + \frac{1}{3} \log\left(\frac{(-x^3+1)^{\frac{1}{3}}}{x} + 1\right) - \frac{1}{6} \log\left(-\frac{(-x^3+1)^{\frac{1}{3}}}{x} + \frac{(-x^3+1)^{\frac{2}{3}}}{x^2} + 1\right)$$

input

```
integrate(1/(-x^3+1)^(1/3),x, algorithm="maxima")
```

output

```
-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3)/x - 1)) + 1/3*log((-x^
3 + 1)^(1/3)/x + 1) - 1/6*log(-(-x^3 + 1)^(1/3)/x + (-x^3 + 1)^(2/3)/x^2 +
1)
```

Giac [F]

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = \int \frac{1}{(-x^3+1)^{\frac{1}{3}}} dx$$

input `integrate(1/(-x^3+1)^(1/3),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(-1/3), x)`

Mupad [B] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.20

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = x {}_2F_1\left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; x^3\right)$$

input `int(1/(1 - x^3)^(1/3),x)`

output `x*hypergeom([1/3, 1/3], 4/3, x^3)`

Reduce [F]

$$\int \frac{1}{\sqrt[3]{1-x^3}} dx = \int \frac{1}{(-x^3+1)^{\frac{1}{3}}} dx$$

input `int(1/(-x^3+1)^(1/3),x)`

output `int(1/(- x**3 + 1)**(1/3),x)`

$$3.36 \quad \int \frac{1}{x \sqrt[3]{1-x^3}} dx$$

Optimal result	287
Mathematica [A] (verified)	287
Rubi [A] (verified)	288
Maple [A] (verified)	290
Fricas [A] (verification not implemented)	290
Sympy [C] (verification not implemented)	291
Maxima [A] (verification not implemented)	291
Giac [A] (verification not implemented)	292
Mupad [B] (verification not implemented)	292
Reduce [F]	293

Optimal result

Integrand size = 15, antiderivative size = 55

$$\int \frac{1}{x \sqrt[3]{1-x^3}} dx = \frac{\arctan\left(\frac{1+2\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\log(x)}{2} + \frac{1}{2} \log\left(1 - \sqrt[3]{1-x^3}\right)$$

output

```
-1/2*ln(x)+1/2*ln(1-(-x^3+1)^(1/3))+1/3*arctan(1/3*(1+2*(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.44

$$\int \frac{1}{x \sqrt[3]{1-x^3}} dx = \frac{\arctan\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(-1 + \sqrt[3]{1-x^3}\right) - \frac{1}{6} \log\left(1 + \sqrt[3]{1-x^3} + (1-x^3)^{2/3}\right)$$

input

```
Integrate[1/(x*(1 - x^3)^(1/3)),x]
```


output

```
ArcTan[1/Sqrt[3] + (2*(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + Log[-1 + (1 - x^3)^(1/3)]/3 - Log[1 + (1 - x^3)^(1/3) + (1 - x^3)^(2/3)]/6
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.11, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {798, 67, 16, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x^3 \sqrt[3]{1-x^3}} dx$$

$$\downarrow 798$$

$$\frac{1}{3} \int \frac{1}{x^3 \sqrt[3]{1-x^3}} dx^3$$

$$\downarrow 67$$

$$\frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{1 - \sqrt[3]{1-x^3}} d\sqrt[3]{1-x^3} + \frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{1-x^3} + 1} d\sqrt[3]{1-x^3} - \frac{1}{2} \log(x^3) \right)$$

$$\downarrow 16$$

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{1-x^3} + 1} d\sqrt[3]{1-x^3} - \frac{1}{2} \log(x^3) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^3}) \right)$$

$$\downarrow 1083$$

$$\frac{1}{3} \left(-3 \int \frac{1}{-x^6 - 3} d(2\sqrt[3]{1-x^3} + 1) - \frac{1}{2} \log(x^3) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^3}) \right)$$

$$\downarrow 217$$

$$\frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2\sqrt[3]{1-x^3} + 1}{\sqrt{3}} \right) - \frac{\log(x^3)}{2} + \frac{3}{2} \log(1 - \sqrt[3]{1-x^3}) \right)$$

input

```
Int[1/(x*(1 - x^3)^(1/3)),x]
```

output $(\sqrt{3} \cdot \text{ArcTan}[(1 + 2(1 - x^3)^{1/3})/\sqrt{3}] - \text{Log}[x^3]/2 + (3 \cdot \text{Log}[1 - (1 - x^3)^{1/3}])/2)/3$

Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)(x_)), x_Symbol] \rightarrow \text{Simp}[c \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 67 $\text{Int}[1/(((a_)+(b_)(x_))*((c_)+(d_)(x_))^{1/3}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b \cdot c - a \cdot d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b \cdot x, x]]/(2 \cdot b \cdot q), x] + (\text{Simp}[3/(2 \cdot b) \text{ Subst}[\text{Int}[1/(q^2 + q \cdot x + x^2), x], x, (c + d \cdot x)^{1/3}], x] - \text{Simp}[3/(2 \cdot b \cdot q) \text{ Subst}[\text{Int}[1/(q - x), x], x, (c + d \cdot x)^{1/3}], x])] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{PosQ}[(b \cdot c - a \cdot d)/b]$

rule 217 $\text{Int}(((a_)+(b_)(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1}) \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \& \& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

rule 798 $\text{Int}[(x_)^{(m_)} \cdot ((a_)+(b_)(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n) - 1}) \cdot (a + b \cdot x)^p, x], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&\& \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

rule 1083 $\text{Int}(((a_)+(b_)(x_)+(c_)(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[-2 \text{ Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] /; \text{FreeQ}[\{a, b, c\}, x]$

Maple [A] (verified)

Time = 2.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.15

method	result
pseudoelliptic	$-\frac{\ln\left((-x^3+1)^{\frac{2}{3}}+(-x^3+1)^{\frac{1}{3}}+1\right)}{6} + \frac{\arctan\left(\frac{(1+2(-x^3+1)^{\frac{1}{3}})\sqrt{3}}{3}\right)\sqrt{3}}{3} + \frac{\ln\left((-x^3+1)^{\frac{1}{3}}-1\right)}{3}$
meijerg	$\frac{\sqrt{3}\Gamma\left(\frac{2}{3}\right)\left(\frac{2\left(-\frac{\pi\sqrt{3}}{6}-\frac{3\ln(3)}{2}+3\ln(x)+i\pi\right)\pi\sqrt{3}}{3\Gamma\left(\frac{2}{3}\right)} + \frac{2\pi\sqrt{3}x^3\operatorname{hypergeom}\left(\left[1,1,\frac{4}{3}\right],[2,2],x^3\right)}{9\Gamma\left(\frac{2}{3}\right)}\right)}{6\pi}$
trager	$\operatorname{RootOf}\left(-Z^2+_Z+1\right)\ln\left(-\frac{1438\operatorname{RootOf}\left(-Z^2+_Z+1\right)^2x^3+9855\operatorname{RootOf}\left(-Z^2+_Z+1\right)x^3-5502\operatorname{RootOf}\left(-Z^2+_Z+1\right)}{\dots}\right)$

input `int(1/x/(-x^3+1)^(1/3),x,method=_RETURNVERBOSE)`output `-1/6*ln((-x^3+1)^(2/3)+(-x^3+1)^(1/3)+1)+1/3*arctan(1/3*(1+2*(-x^3+1)^(1/3)))*3^(1/2)+1/3*ln((-x^3+1)^(1/3)-1)`**Fricas [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.16

$$\int \frac{1}{x\sqrt[3]{1-x^3}} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{2}{3} \sqrt{3}(-x^3+1)^{\frac{1}{3}} + \frac{1}{3} \sqrt{3}\right) - \frac{1}{6} \log\left(\left(-x^3+1\right)^{\frac{2}{3}} + \left(-x^3+1\right)^{\frac{1}{3}} + 1\right) + \frac{1}{3} \log\left(\left(-x^3+1\right)^{\frac{1}{3}} - 1\right)$$

input `integrate(1/x/(-x^3+1)^(1/3),x, algorithm="fricas")`output `1/3*sqrt(3)*arctan(2/3*sqrt(3)*(-x^3 + 1)^(1/3) + 1/3*sqrt(3)) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log((-x^3 + 1)^(1/3) - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.41 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.58

$$\int \frac{1}{x\sqrt[3]{1-x^3}} dx = -\frac{e^{-\frac{i\pi}{3}}\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{4}{3} \middle| \frac{1}{x^3}\right)}{3x\Gamma\left(\frac{4}{3}\right)}$$

input `integrate(1/x/(-x**3+1)**(1/3),x)`

output `-exp(-I*pi/3)*gamma(1/3)*hyper((1/3, 1/3), (4/3,), x**(-3))/(3*x*gamma(4/3))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.13

$$\int \frac{1}{x\sqrt[3]{1-x^3}} dx = \frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}\left(2(-x^3+1)^{\frac{1}{3}}+1\right)\right) - \frac{1}{6}\log\left(\left(-x^3+1\right)^{\frac{2}{3}}+\left(-x^3+1\right)^{\frac{1}{3}}+1\right) + \frac{1}{3}\log\left(\left(-x^3+1\right)^{\frac{1}{3}}-1\right)$$

input `integrate(1/x/(-x^3+1)^(1/3),x, algorithm="maxima")`

output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3) + 1)) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log((-x^3 + 1)^(1/3) - 1)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.15

$$\int \frac{1}{x\sqrt[3]{1-x^3}} dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \left(2(-x^3+1)^{\frac{1}{3}} + 1 \right) \right) - \frac{1}{6} \log \left((-x^3+1)^{\frac{2}{3}} + (-x^3+1)^{\frac{1}{3}} + 1 \right) + \frac{1}{3} \log \left(\left| (-x^3+1)^{\frac{1}{3}} - 1 \right| \right)$$

input `integrate(1/x/(-x^3+1)^(1/3),x, algorithm="giac")`output `1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3) + 1)) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log(abs((-x^3 + 1)^(1/3) - 1))`**Mupad [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.45

$$\int \frac{1}{x\sqrt[3]{1-x^3}} dx = \frac{\ln \left((1-x^3)^{1/3} - 1 \right)}{3} + \ln \left((1-x^3)^{1/3} - 9 \left(-\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6} \right)^2 \right) \left(-\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6} \right) - \ln \left((1-x^3)^{1/3} - 9 \left(\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6} \right)^2 \right) \left(\frac{1}{6} + \frac{\sqrt{3} \text{li}}{6} \right)$$

input `int(1/(x*(1 - x^3)^(1/3)),x)`output `log((1 - x^3)^(1/3) - 1)/3 + log((1 - x^3)^(1/3) - 9*((3^(1/2)*1i)/6 - 1/6)^2)*((3^(1/2)*1i)/6 - 1/6) - log((1 - x^3)^(1/3) - 9*((3^(1/2)*1i)/6 + 1/6)^2)*((3^(1/2)*1i)/6 + 1/6)`

Reduce [F]

$$\int \frac{1}{x\sqrt[3]{1-x^3}} dx = \int \frac{1}{(-x^3+1)^{\frac{1}{3}}x} dx$$

input `int(1/x/(-x^3+1)^(1/3),x)`

output `int(1/((-x**3+1)**(1/3)*x),x)`

3.37 $\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx$

Optimal result	294
Mathematica [A] (verified)	294
Rubi [A] (verified)	295
Maple [C] (warning: unable to verify)	296
Fricas [B] (verification not implemented)	297
Sympy [F]	298
Maxima [F]	298
Giac [F]	299
Mupad [F(-1)]	299
Reduce [F]	299

Optimal result

Integrand size = 17, antiderivative size = 97

$$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = -\frac{\sqrt{3} \arctan\left(\frac{1+\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{2\sqrt[3]{2}} - \frac{\log((1-x)(1+x)^2)}{4\sqrt[3]{2}} + \frac{3 \log\left(-1+x+2^{2/3}\sqrt[3]{1-x^3}\right)}{4\sqrt[3]{2}}$$

output -1/8*ln((1-x)*(1+x)^2)*2^(2/3)+3/8*ln(-1+x+2^(2/3)*(-x^3+1)^(1/3))*2^(2/3)
-1/4*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)*2^(2/3)

Mathematica [A] (verified)

Time = 0.94 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.53

$$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1-x^3}}{\sqrt[3]{2}-\sqrt[3]{2}x+\sqrt[3]{1-x^3}}\right) + 2 \log\left(-\sqrt[3]{2} + \sqrt[3]{2}x + 2\sqrt[3]{1-x^3}\right) - \log\left(2^{2/3} - 2 \cdot 2^{2/3}x + 2^{2/3}x^2 - \dots\right)}{4\sqrt[3]{2}}$$

input `Integrate[1/((1 + x)*(1 - x^3)^(1/3)),x]`

output `(2*Sqrt[3]*ArcTan[(Sqrt[3]*(1 - x^3)^(1/3))/(2^(1/3) - 2^(1/3)*x + (1 - x^3)^(1/3))] + 2*Log[-2^(1/3) + 2^(1/3)*x + 2*(1 - x^3)^(1/3)] - Log[2^(2/3) - 2*2^(2/3)*x + 2^(2/3)*x^2 - 2*(-1 + x)*(2 - 2*x^3)^(1/3) + 4*(1 - x^3)^(2/3)])/(4*2^(1/3))`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {2574}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx$$

↓ 2574

$$-\frac{\sqrt{3} \arctan\left(\frac{\sqrt[3]{2(1-x)}+1}{\frac{\sqrt[3]{1-x^3}}{\sqrt{3}}}\right)}{2\sqrt[3]{2}} + \frac{3 \log\left(2^{2/3}\sqrt[3]{1-x^3} + x - 1\right)}{4\sqrt[3]{2}} - \frac{\log\left((1-x)(x+1)^2\right)}{4\sqrt[3]{2}}$$

input `Int[1/((1 + x)*(1 - x^3)^(1/3)),x]`

output `-1/2*(Sqrt[3]*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]])/2^(1/3) - Log[(1 - x)*(1 + x)^2]/(4*2^(1/3)) + (3*Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3)])/(4*2^(1/3))`

Defintions of rubi rules used

rule 2574

```
Int[1/(((c_) + (d_.)*(x_))*((a_) + (b_.)*(x_)^3)^(1/3)), x_Symbol] :> Simp[
Sqrt[3]*(ArcTan[(1 - 2^(1/3)*Rt[b, 3]*((c - d*x)/(d*(a + b*x^3)^(1/3)))))/Sqrt[3]]/(2^(4/3)*Rt[b, 3]*c), x] + (Simp[Log[(c + d*x)^2*(c - d*x)]/(2^(7/3)*Rt[b, 3]*c), x] - Simp[(3*Log[Rt[b, 3]*(c - d*x) + 2^(2/3)*d*(a + b*x^3)^(1/3)])/(2^(7/3)*Rt[b, 3]*c), x]) /; FreeQ[{a, b, c, d}, x] && EqQ[b*c^3 + a*d^3, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 7.40 (sec) , antiderivative size = 1143, normalized size of antiderivative = 11.78

method	result	size
trager	Expression too large to display	1143

input

```
int(1/(1+x)/(-x^3+1)^(1/3),x,method=_RETURNVERBOSE)
```

output

```

-1/4*ln((20*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)^2*RootOf(_
Z^3-4)^2*x+6*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z
^3-4)^3*x+8*(-x^3+1)^(2/3)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_
Z^2)*RootOf(_Z^3-4)^2-18*(-x^3+1)^(1/3)*RootOf(_Z^3-4)*RootOf(RootOf(_Z^3-
4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*x-13*(-x^3+1)^(1/3)*RootOf(_Z^3-4)^2*x+18
*(-x^3+1)^(1/3)*RootOf(_Z^3-4)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)
+4*_Z^2)-70*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*x^2+13*(-x
^3+1)^(1/3)*RootOf(_Z^3-4)^2-21*RootOf(_Z^3-4)*x^2-20*RootOf(RootOf(_Z^3-4
)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*x-6*RootOf(_Z^3-4)*x-36*(-x^3+1)^(2/3)-70*
RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)-21*RootOf(_Z^3-4))/(1+
x)^2)*RootOf(_Z^3-4)-1/2*ln((20*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4
)+4*_Z^2)^2*RootOf(_Z^3-4)^2*x+6*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-
4)+4*_Z^2)*RootOf(_Z^3-4)^3*x+8*(-x^3+1)^(2/3)*RootOf(RootOf(_Z^3-4)^2+2*_
Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^2-18*(-x^3+1)^(1/3)*RootOf(_Z^3-4)
*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*x-13*(-x^3+1)^(1/3)*R
ootOf(_Z^3-4)^2*x+18*(-x^3+1)^(1/3)*RootOf(_Z^3-4)*RootOf(RootOf(_Z^3-4)^2
+2*_Z*RootOf(_Z^3-4)+4*_Z^2)-70*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4
)+4*_Z^2)*x^2+13*(-x^3+1)^(1/3)*RootOf(_Z^3-4)^2-21*RootOf(_Z^3-4)*x^2-20*
RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*x-6*RootOf(_Z^3-4)*x-3
6*(-x^3+1)^(2/3)-70*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 298 vs. $2(71) = 142$.

Time = 1.76 (sec) , antiderivative size = 298, normalized size of antiderivative = 3.07

$$\begin{aligned}
& \int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = \frac{1}{6} \\
& \cdot 2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \arctan \left(\frac{2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \left(8 \cdot 2^{\frac{2}{3}} (x^4 + 2x^3 + 2x^2 + 2x + 1) (-x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}} (13x^6 + 2x^5 + 19x^4 - 4x^3 - 3(3x^6 - 18x^5 - 3x^4 - 28x^3 - \dots \right)}{3(3x^6 - 18x^5 - 3x^4 - 28x^3 - \dots} \right)}{24} \\
& \cdot 2^{\frac{2}{3}} \log \left(\frac{4 \cdot 2^{\frac{2}{3}} (-x^3 + 1)^{\frac{2}{3}} (x^2 + 1) + 2^{\frac{1}{3}} (5x^4 + 6x^2 + 5) - 2(3x^3 - x^2 + x - 3) (-x^3 + 1)^{\frac{1}{3}}}{x^4 + 4x^3 + 6x^2 + 4x + 1} \right) \\
& + \frac{1}{12} \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} (x^2 + 2x + 1) - 2 \cdot 2^{\frac{1}{3}} (-x^3 + 1)^{\frac{1}{3}} (x - 1) - 4(-x^3 + 1)^{\frac{2}{3}}}{x^2 + 2x + 1} \right)
\end{aligned}$$

input `integrate(1/(1+x)/(-x^3+1)^(1/3),x, algorithm="fricas")`

output
$$\frac{1}{6}2^{1/6}\sqrt{3/2}\arctan\left(\frac{1}{3}2^{1/6}\sqrt{3/2}\right)(8\cdot 2^{2/3})(x^4 + 2x^3 + 2x^2 + 2x + 1)(-x^3 + 1)^{2/3} + 2^{1/3}(13x^6 + 2x^5 + 19x^4 - 4x^3 + 19x^2 + 2x + 13) - 4(5x^5 - 5x^4 + 6x^3 - 6x^2 + 5x - 5)(-x^3 + 1)^{1/3}) / (3x^6 - 18x^5 - 3x^4 - 28x^3 - 3x^2 - 18x + 3) - 1/24\cdot 2^{2/3}\log((4\cdot 2^{2/3})(-x^3 + 1)^{2/3}(x^2 + 1) + 2^{1/3}(5x^4 + 6x^2 + 5) - 2(3x^3 - x^2 + x - 3)(-x^3 + 1)^{1/3}) / (x^4 + 4x^3 + 6x^2 + 4x + 1)) + 1/12\cdot 2^{2/3}\log((2^{2/3}(x^2 + 2x + 1) - 2\cdot 2^{1/3})(-x^3 + 1)^{1/3}(x - 1) - 4(-x^3 + 1)^{2/3}) / (x^2 + 2x + 1))$$

Sympy [F]

$$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{1}{\sqrt[3]{-(x-1)(x^2+x+1)}(x+1)} dx$$

input `integrate(1/(1+x)/(-x**3+1)**(1/3),x)`

output `Integral(1/((-x - 1)*(x**2 + x + 1))**(1/3)*(x + 1)), x)`

Maxima [F]

$$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{1}{(-x^3+1)^{1/3}(x+1)} dx$$

input `integrate(1/(1+x)/(-x^3+1)^(1/3),x, algorithm="maxima")`

output `integrate(1/((-x^3 + 1)^(1/3)*(x + 1)), x)`

Giac [F]

$$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{1}{(-x^3+1)^{\frac{1}{3}}(x+1)} dx$$

input `integrate(1/(1+x)/(-x^3+1)^(1/3),x, algorithm="giac")`

output `integrate(1/((-x^3 + 1)^(1/3)*(x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{1}{(1-x^3)^{1/3}(x+1)} dx$$

input `int(1/((1 - x^3)^(1/3)*(x + 1)),x)`

output `int(1/((1 - x^3)^(1/3)*(x + 1)), x)`

Reduce [F]

$$\int \frac{1}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{1}{(-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int(1/(1+x)/(-x^3+1)^(1/3),x)`

output `int(1/((- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)`

3.38 $\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx$

Optimal result	300
Mathematica [F]	301
Rubi [A] (verified)	301
Maple [C] (warning: unable to verify)	303
Fricas [F(-2)]	304
Sympy [F]	304
Maxima [F]	304
Giac [F]	305
Mupad [F(-1)]	305
Reduce [F]	305

Optimal result

Integrand size = 18, antiderivative size = 145

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \frac{\sqrt{3} \arctan\left(\frac{1+\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{2\sqrt[3]{2}} - \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\log((1-x)(1+x)^2)}{4\sqrt[3]{2}} + \frac{1}{2} \log\left(x + \sqrt[3]{1-x^3}\right) - \frac{3 \log\left(-1+x+2^{2/3}\sqrt[3]{1-x^3}\right)}{4\sqrt[3]{2}}$$

output

```
1/8*ln((1-x)*(1+x)^2)*2^(2/3)+1/2*ln(x+(-x^3+1)^(1/3))-3/8*ln(-1+x+2^(2/3)
*(-x^3+1)^(1/3))*2^(2/3)-1/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(1/2))*3^(
1/2)+1/4*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)*2^(
2/3)
```

Mathematica [F]

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx$$

input `Integrate[x/((1 + x)*(1 - x^3)^(1/3)), x]`

output `Integrate[x/((1 + x)*(1 - x^3)^(1/3)), x]`

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2577, 769, 2574}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{(x+1)\sqrt[3]{1-x^3}} dx \\ & \quad \downarrow \text{2577} \\ & \int \frac{1}{\sqrt[3]{1-x^3}} dx - \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx \\ & \quad \downarrow \text{769} \\ & - \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{\arctan\left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2} \log(\sqrt[3]{1-x^3} + x) \\ & \quad \downarrow \text{2574} \end{aligned}$$

$$\frac{\sqrt{3} \arctan\left(\frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}\right)}{2\sqrt[3]{2}} - \frac{\arctan\left(\frac{1-\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2} \log\left(\sqrt[3]{1-x^3}+x\right) - \frac{3 \log\left(2^{2/3}\sqrt[3]{1-x^3}+x-1\right)}{4\sqrt[3]{2}} + \frac{\log((1-x)(x+1)^2)}{4\sqrt[3]{2}}$$

input `Int[x/((1 + x)*(1 - x^3)^(1/3)),x]`

output `(Sqrt[3]*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/(2*2^(1/3)) - ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + Log[(1 - x)*(1 + x)^2]/(4*2^(1/3)) + Log[x + (1 - x^3)^(1/3)]/2 - (3*Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3)]/(4*2^(1/3)))`

Defintions of rubi rules used

rule 769 `Int[((a_) + (b_)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

rule 2574 `Int[1/(((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^3)^(1/3)), x_Symbol] := Simp[Sqrt[3]*(ArcTan[(1 - 2^(1/3)*Rt[b, 3]*((c - d*x)/(d*(a + b*x^3)^(1/3))))/Sqrt[3]]/(2^(4/3)*Rt[b, 3]*c), x] + (Simp[Log[(c + d*x)^2*(c - d*x)]/(2^(7/3)*Rt[b, 3]*c), x] - Simp[(3*Log[Rt[b, 3]*(c - d*x) + 2^(2/3)*d*(a + b*x^3)^(1/3)]/(2^(7/3)*Rt[b, 3]*c), x]) /; FreeQ[{a, b, c, d}, x] && EqQ[b*c^3 + a*d^3, 0]`

rule 2577 `Int[((e_) + (f_)*(x_))/(((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^3)^(1/3)), x_Symbol] := Simp[f/d Int[1/(a + b*x^3)^(1/3), x], x] + Simp[(d*e - c*f)/d Int[1/((c + d*x)*(a + b*x^3)^(1/3)), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 16.86 (sec) , antiderivative size = 1421, normalized size of antiderivative = 9.80

method	result	size
trager	Expression too large to display	1421

input `int(x/(1+x)/(-x^3+1)^(1/3),x,method=_RETURNVERBOSE)`

output

```

1/3*ln(-RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)^2*RootOf(_Z^3+4)^4*x^3+3*(-x^3+1)^(2/3)*RootOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*x-RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*x-RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*x^3+2*RootOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)+6*x^2*(-x^3+1)^(1/3)+2*x^3-2)+1/2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*ln((10*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)^2*RootOf(_Z^3+4)^2*x+2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)^3*x+4*(-x^3+1)^(2/3)*RootOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)+9*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)*x-2*(-x^3+1)^(1/3)*RootOf(_Z^3+4)^2*x-9*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)-35*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*x^2+2*(-x^3+1)^(1/3)*RootOf(_Z^3+4)^2-7*x^2*RootOf(_Z^3+4)-30*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*x-6*x*RootOf(_Z^3+4)-26*(-x^3+1)^(2/3)-35*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)-7*RootOf(_Z^3+4))/(1+x)^2)+1/4*RootOf(_Z^3+4)*ln(-(8*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)^2*RootOf(_Z^3+4)^2*x+10*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)^3*x+8*(-x^3+1)^(2/3)*RootOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)-26*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+...

```


Fricas [F(-2)]

Exception generated.

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \text{Exception raised: TypeError}$$

input `integrate(x/(1+x)/(-x^3+1)^(1/3),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (residue poly has multiple non-linear factors)`

Sympy [F]

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{x}{\sqrt[3]{-(x-1)(x^2+x+1)}(x+1)} dx$$

input `integrate(x/(1+x)/(-x**3+1)**(1/3),x)`

output `Integral(x/((-x - 1)*(x**2 + x + 1))**(1/3)*(x + 1)), x)`

Maxima [F]

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{x}{(-x^3+1)^{\frac{1}{3}}(x+1)} dx$$

input `integrate(x/(1+x)/(-x^3+1)^(1/3),x, algorithm="maxima")`

output `integrate(x/((-x^3 + 1)^(1/3)*(x + 1)), x)`

Giac [F]

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{x}{(-x^3+1)^{\frac{1}{3}}(x+1)} dx$$

input `integrate(x/(1+x)/(-x^3+1)^(1/3),x, algorithm="giac")`

output `integrate(x/((-x^3 + 1)^(1/3)*(x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{x}{(1-x^3)^{1/3}(x+1)} dx$$

input `int(x/((1 - x^3)^(1/3)*(x + 1)),x)`

output `int(x/((1 - x^3)^(1/3)*(x + 1)), x)`

Reduce [F]

$$\int \frac{x}{(1+x)\sqrt[3]{1-x^3}} dx = \int \frac{x}{(-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int(x/(1+x)/(-x^3+1)^(1/3),x)`

output `int(x/((- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)`

3.39 $\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx$

Optimal result	306
Mathematica [A] (verified)	306
Rubi [A] (verified)	307
Maple [C] (warning: unable to verify)	308
Fricas [B] (verification not implemented)	309
Sympy [F]	310
Maxima [F]	310
Giac [F]	311
Mupad [F(-1)]	311
Reduce [F]	311

Optimal result

Integrand size = 16, antiderivative size = 110

$$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx = -\frac{\sqrt{3} \arctan\left(\frac{1}{\sqrt{3}} + \frac{\sqrt[3]{2(2-x)}}{\sqrt{3}\sqrt[3]{2-3x+x^2}}\right) - \frac{\log(2-x)}{4\sqrt[3]{2}}}{2\sqrt[3]{2}} - \frac{\log(x)}{2\sqrt[3]{2}} + \frac{3 \log\left(2-x-2^{2/3}\sqrt[3]{2-3x+x^2}\right)}{4\sqrt[3]{2}}$$

output

```
-1/8*ln(2-x)*2^(2/3)-1/4*ln(x)*2^(2/3)+3/8*ln(2-x-2^(2/3)*(x^2-3*x+2)^(1/3))
)*2^(2/3)+1/4*arctan(-1/3*3^(1/2)-1/3*2^(1/3)*(2-x)/(x^2-3*x+2)^(1/3)*3^(
1/2))*3^(1/2)*2^(2/3)
```

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.47

$$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx = \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{2-3x+x^2}}{2\sqrt[3]{2}-\sqrt[3]{2x+\sqrt[3]{2-3x+x^2}}}\right) + 2 \log\left(-2\sqrt[3]{2} + \sqrt[3]{2}x + 2\sqrt[3]{2-3x+x^2}\right) - \log\left(4 \cdot 2^{2/3} - 4 \cdot 2\right)}{4\sqrt[3]{2}}$$

input `Integrate[1/(x*(2 - 3*x + x^2)^(1/3)),x]`

output $(2\sqrt[3]{3}\text{ArcTan}[\sqrt[3]{3}(2 - 3x + x^2)^{1/3}]/(2\sqrt[3]{2} - 2^{1/3})x + (2 - 3x + x^2)^{1/3}] + 2\text{Log}[-2\sqrt[3]{2} + 2^{1/3})x + 2(2 - 3x + x^2)^{1/3}] - \text{Log}[4\sqrt[3]{2} - 4\sqrt[3]{2}x + 2^{2/3}x^2 - 2\sqrt[3]{2}(-2 + x)(2 - 3x + x^2)^{1/3} + 4(2 - 3x + x^2)^{2/3}]/(4\sqrt[3]{2})$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.09, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {1177, 27, 133}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x\sqrt[3]{x^2 - 3x + 2}} dx$$

$$\downarrow 1177$$

$$\frac{2^{2/3}\sqrt[3]{x-2}\sqrt[3]{x-1} \int \frac{1}{2^{2/3}\sqrt[3]{x-2}\sqrt[3]{x-1x}} dx}{\sqrt[3]{x^2 - 3x + 2}}$$

$$\downarrow 27$$

$$\frac{\sqrt[3]{x-2}\sqrt[3]{x-1} \int \frac{1}{\sqrt[3]{x-2}\sqrt[3]{x-1x}} dx}{\sqrt[3]{x^2 - 3x + 2}}$$

$$\downarrow 133$$

$$\frac{\sqrt[3]{x-2}\sqrt[3]{x-1} \left(-\frac{\sqrt{3}\arctan\left(\frac{1}{\sqrt{3}} - \frac{\sqrt{2}(x-2)^{2/3}}{\sqrt{3}\sqrt[3]{x-1}}\right)}{2\sqrt[3]{2}} + \frac{3\log\left(-\frac{(x-2)^{2/3}}{2^{2/3}} - \sqrt[3]{x-1}\right)}{4\sqrt[3]{2}} - \frac{\log(x)}{2\sqrt[3]{2}} \right)}{\sqrt[3]{x^2 - 3x + 2}}$$

input `Int[1/(x*(2 - 3*x + x^2)^(1/3)),x]`

output

$$\frac{((-2 + x)^{1/3}*(-1 + x)^{1/3}*(-1/2*\sqrt[3]{3}*\text{ArcTan}[1/\sqrt[3]{3}] - (2^{1/3})*(-2 + x)^{2/3})/(\sqrt[3]{3}*(-1 + x)^{1/3})]/2^{1/3} + (3*\text{Log}[-((-2 + x)^{2/3}/2^{2/3}) - (-1 + x)^{1/3}])/(4*2^{1/3}) - \text{Log}[x]/(2*2^{1/3})))/(2 - 3*x + x^2)^{1/3}}$$

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_) /; FreeQ[b, x]]
```

rule 133

```
Int[1/(((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(1/3))*((e_.) + (f_.)*(x_.))^(1/3)), x_] := With[{q = Rt[b*((b*e - a*f)/(b*c - a*d)^2), 3]}, Simp[-Log[a + b*x]/(2*q*(b*c - a*d)), x] + (-Simp[Sqrt[3]*(ArcTan[1/Sqrt[3] + 2*q*((c + d*x)^(2/3)/(Sqrt[3]*(e + f*x)^(1/3)))]/(2*q*(b*c - a*d))), x] + Simp[3*(Log[q*(c + d*x)^(2/3) - (e + f*x)^(1/3)]/(4*q*(b*c - a*d))), x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[2*b*d*e - b*c*f - a*d*f, 0]
```

rule 1177

```
Int[1/(((d_.) + (e_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(1/3)), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(b + q + 2*c*x)^(1/3)*((b - q + 2*c*x)^(1/3)/(a + b*x + c*x^2)^(1/3)) Int[1/((d + e*x)*(b + q + 2*c*x)^(1/3)*(b - q + 2*c*x)^(1/3)), x], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d^2 - b*c*d*e - 2*b^2*e^2 + 9*a*c*e^2, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 7.76 (sec) , antiderivative size = 1592, normalized size of antiderivative = 14.47

method	result	size
trager	Expression too large to display	1592

input

```
int(1/x/(x^2-3*x+2)^(1/3),x,method=_RETURNVERBOSE)
```

output

```

-1/4*ln((12*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^
3-4)^3*x^2+136*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)^2*RootO
f(_Z^3-4)^2*x^2-54*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*Roo
tOf(_Z^3-4)^3*x-612*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)^2*
RootOf(_Z^3-4)^2*x+216*(x^2-3*x+2)^(2/3)*RootOf(_Z^3-4)^2*RootOf(RootOf(_Z
^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)+54*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf
(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^3+612*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(
_Z^3-4)+4*_Z^2)^2*RootOf(_Z^3-4)^2+129*(x^2-3*x+2)^(1/3)*RootOf(_Z^3-4)^2*
x+474*(x^2-3*x+2)^(1/3)*RootOf(_Z^3-4)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf
(_Z^3-4)+4*_Z^2)*x-258*(x^2-3*x+2)^(1/3)*RootOf(_Z^3-4)^2-948*(x^2-3*x+2)^(
1/3)*RootOf(_Z^3-4)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)+2
1*RootOf(_Z^3-4)*x^2+238*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^
2)*x^2-180*RootOf(_Z^3-4)*x-2040*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-
4)+4*_Z^2)*x+948*(x^2-3*x+2)^(2/3)+180*RootOf(_Z^3-4)+2040*RootOf(RootOf(
_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2))/x^2)*RootOf(_Z^3-4)-1/2*ln((12*RootO
f(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^3*x^2+136*Ro
otOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)^2*RootOf(_Z^3-4)^2*x^2-5
4*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^3*x-6
12*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)^2*RootOf(_Z^3-4)^2*
x+216*(x^2-3*x+2)^(2/3)*RootOf(_Z^3-4)^2*RootOf(RootOf(_Z^3-4)^2+2*_Z*R...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 274 vs. $2(81) = 162$.

Time = 1.00 (sec) , antiderivative size = 274, normalized size of antiderivative = 2.49

$$\begin{aligned}
& \int \frac{1}{x\sqrt{2-3x+x^2}} dx = -\frac{1}{6} \\
& \cdot 2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \arctan \left(\frac{2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \left(24 \cdot 2^{\frac{2}{3}} (x^4 - 6x^3 + 6x^2) (x^2 - 3x + 2)^{\frac{2}{3}} + 2^{\frac{1}{3}} (x^6 + 36x^5 - 612x^4 + 2880x^3 - 3168x^2 + 1296x - 216) \right)}{3(x^6 - 108x^5 + 972x^4 - 3168x^3 + 3168x^2 - 1296x + 216)}} \right) \\
& + \frac{1}{12} \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} x^2 + 6 \cdot 2^{\frac{1}{3}} (x^2 - 3x + 2)^{\frac{1}{3}} (x - 2) + 12 (x^2 - 3x + 2)^{\frac{2}{3}}}{x^2} \right) - \frac{1}{24} \\
& \cdot 2^{\frac{2}{3}} \log \left(\frac{12 \cdot 2^{\frac{2}{3}} (x^2 - 3x + 2)^{\frac{2}{3}} (x^2 - 6x + 6) + 2^{\frac{1}{3}} (x^4 - 36x^3 + 180x^2 - 288x + 144) - 6(x^3 - 14x^2 + 14x - 6)}{x^4} \right)
\end{aligned}$$

input

```
integrate(1/x/(x^2-3*x+2)^(1/3),x, algorithm="fricas")
```

output

```
-1/6*2^(1/6)*sqrt(3/2)*arctan(1/3*2^(1/6)*sqrt(3/2)*(24*2^(2/3)*(x^4 - 6*x^3 + 6*x^2)*(x^2 - 3*x + 2)^(2/3) + 2^(1/3)*(x^6 + 36*x^5 - 612*x^4 + 2880*x^3 - 5760*x^2 + 5184*x - 1728) + 12*(x^5 - 38*x^4 + 252*x^3 - 648*x^2 + 720*x - 288)*(x^2 - 3*x + 2)^(1/3))/(x^6 - 108*x^5 + 972*x^4 - 3456*x^3 + 6048*x^2 - 5184*x + 1728)) + 1/12*2^(2/3)*log((2^(2/3)*x^2 + 6*2^(1/3)*(x^2 - 3*x + 2)^(1/3)*(x - 2) + 12*(x^2 - 3*x + 2)^(2/3))/x^2) - 1/24*2^(2/3)*log((12*2^(2/3)*(x^2 - 3*x + 2)^(2/3)*(x^2 - 6*x + 6) + 2^(1/3)*(x^4 - 36*x^3 + 180*x^2 - 288*x + 144) - 6*(x^3 - 14*x^2 + 36*x - 24)*(x^2 - 3*x + 2)^(1/3))/x^4)
```

Sympy [F]

$$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx = \int \frac{1}{x\sqrt[3]{(x-2)(x-1)}} dx$$

input

```
integrate(1/x/(x**2-3*x+2)**(1/3),x)
```

output

```
Integral(1/(x*((x - 2)*(x - 1))**(1/3)), x)
```

Maxima [F]

$$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx = \int \frac{1}{(x^2-3x+2)^{\frac{1}{3}}x} dx$$

input

```
integrate(1/x/(x^2-3*x+2)^(1/3),x, algorithm="maxima")
```

output

```
integrate(1/((x^2 - 3*x + 2)^(1/3)*x), x)
```

Giac [F]

$$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx = \int \frac{1}{(x^2-3x+2)^{\frac{1}{3}}x} dx$$

input `integrate(1/x/(x^2-3*x+2)^(1/3),x, algorithm="giac")`

output `integrate(1/((x^2 - 3*x + 2)^(1/3)*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx = \int \frac{1}{x(x^2-3x+2)^{\frac{1}{3}}} dx$$

input `int(1/(x*(x^2 - 3*x + 2)^(1/3)),x)`

output `int(1/(x*(x^2 - 3*x + 2)^(1/3)), x)`

Reduce [F]

$$\int \frac{1}{x\sqrt[3]{2-3x+x^2}} dx = \int \frac{1}{(x^2-3x+2)^{\frac{1}{3}}x} dx$$

input `int(1/x/(x^2-3*x+2)^(1/3),x)`

output `int(1/((x**2 - 3*x + 2)**(1/3)*x),x)`

3.40 $\int \frac{1}{\sqrt[3]{-5 + 7x - 3x^2 + x^3}} dx$

Optimal result	312
Mathematica [C] (verified)	312
Rubi [A] (warning: unable to verify)	313
Maple [C] (verified)	315
Fricas [A] (verification not implemented)	316
Sympy [F]	316
Maxima [F]	317
Giac [F]	317
Mupad [F(-1)]	317
Reduce [F]	318

Optimal result

Integrand size = 17, antiderivative size = 81

$$\int \frac{1}{\sqrt[3]{-5 + 7x - 3x^2 + x^3}} dx = \frac{1}{2}\sqrt{3} \arctan\left(\frac{1}{\sqrt{3}} + \frac{2(-1+x)}{\sqrt{3}\sqrt[3]{-5 + 7x - 3x^2 + x^3}}\right) + \frac{1}{4} \log(1-x) - \frac{3}{4} \log\left(1-x + \sqrt[3]{-5 + 7x - 3x^2 + x^3}\right)$$

output `1/4*ln(1-x)-3/4*ln(1-x+(x^3-3*x^2+7*x-5)^(1/3))+1/2*arctan(1/3*3^(1/2)+2/3*(-1+x)/(x^3-3*x^2+7*x-5)^(1/3)*3^(1/2))*3^(1/2)`

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.05

$$\int \frac{1}{\sqrt[3]{-5 + 7x - 3x^2 + x^3}} dx = \frac{3\sqrt[3]{(2-i) + ix}\sqrt[3]{i(-1+x)}((-1+2i) + x) \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{3}, \frac{1}{3}, \frac{5}{3}, -\frac{1}{4}i((-1+2i) + x), -\frac{1}{2}i((-1+2i) + x)\right)}{4\sqrt[3]{-5 + 7x - 3x^2 + x^3}}$$

input `Integrate[(-5 + 7*x - 3*x^2 + x^3)^(-1/3), x]`

output

```
(3*((2 - I) + I*x)^(1/3)*(I*(-1 + x))^(1/3)*((-1 + 2*I) + x)*AppellF1[2/3,
1/3, 1/3, 5/3, (-1/4*I)*((-1 + 2*I) + x), (-1/2*I)*((-1 + 2*I) + x)]/(4*
(-5 + 7*x - 3*x^2 + x^3)^(1/3))
```

Rubi [A] (warning: unable to verify)

Time = 0.22 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.06, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2481, 1917, 266, 807, 769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{\sqrt[3]{x^3 - 3x^2 + 7x - 5}} dx \\
 & \quad \downarrow \text{2481} \\
 & \int \frac{1}{\sqrt[3]{(x-1)^3 + 4(x-1)}} d(x-1) \\
 & \quad \downarrow \text{1917} \\
 & \frac{\sqrt[3]{(x-1)^2 + 4\sqrt[3]{x-1}} \int \frac{1}{\sqrt[3]{(x-1)^2 + 4\sqrt[3]{x-1}}} d(x-1)}{\sqrt[3]{(x-1)^3 + 4(x-1)}} \\
 & \quad \downarrow \text{266} \\
 & \frac{3\sqrt[3]{(x-1)^2 + 4\sqrt[3]{x-1}} \int \frac{\sqrt[3]{x-1}}{\sqrt[3]{(x-1)^2 + 4}} d\sqrt[3]{x-1}}{\sqrt[3]{(x-1)^3 + 4(x-1)}} \\
 & \quad \downarrow \text{807} \\
 & \frac{3\sqrt[3]{(x-1)^2 + 4\sqrt[3]{x-1}} \int \frac{1}{\sqrt[3]{x+3}} d(x-1)^{2/3}}{2\sqrt[3]{(x-1)^3 + 4(x-1)}} \\
 & \quad \downarrow \text{769}
 \end{aligned}$$

$$\frac{3\sqrt[3]{(x-1)^2+4}\sqrt[3]{x-1} \left(\frac{\arctan\left(\frac{\sqrt[3]{x+3}+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(-x + \sqrt[3]{x+3} + 1\right) \right)}{2\sqrt[3]{(x-1)^3+4(x-1)}}$$

input `Int[(-5 + 7*x - 3*x^2 + x^3)^(-1/3), x]`

output `(3*(4 + (-1 + x)^2)^(1/3)*(-1 + x)^(1/3)*(ArcTan[(1 + (2*(-1 + x)^(2/3)))/(3 + x)^(1/3)]/Sqrt[3])/Sqrt[3] - Log[1 - x + (3 + x)^(1/3)]/2)/(2*(4*(-1 + x) + (-1 + x)^3)^(1/3))`

Defintions of rubi rules used

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntegerBinomialQ[a, b, c, 2, m, p, x]`

rule 769 `Int[((a_) + (b_.)*(x_)^3)^(1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^p_, x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1 /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 1917 `Int[((a_.)*(x_)^(j_.) + (b_.)*(x_)^(n_.))^p_, x_Symbol] := Simp[(a*x^j + b*x^n)^FracPart[p]/(x^(j*FracPart[p])*(a + b*x^(n - j))^FracPart[p]) Int[x^(j*p)*(a + b*x^(n - j))^p, x], x] /; FreeQ[{a, b, j, n, p}, x] && !IntegerQ[p] && NeQ[n, j] && PosQ[n - j]`

rule 2481

```
Int[(Px_)^(p_), x_Symbol] := With[{a = Coeff[Px, x, 0], b = Coeff[Px, x, 1],
c = Coeff[Px, x, 2], d = Coeff[Px, x, 3]}, Subst[Int[Simp[(2*c^3 - 9*b*c*d + 27*a*d^2)/(27*d^2) - (c^2 - 3*b*d)*(x/(3*d)) + d*x^3, x]^p, x], x, c/(3*d) + x]] /; FreeQ[p, x] && PolyQ[Px, x, 3]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.17 (sec) , antiderivative size = 653, normalized size of antiderivative = 8.06

method	result
trager	$\frac{\text{RootOf}(_Z^2 - _Z + 1) \ln\left(-304 \text{RootOf}(_Z^2 - _Z + 1)^2 x^2 + 624 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - 3x^2 + 7x - 5)^{\frac{2}{3}} + 624 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - 3x^2 + 7x - 5)^{\frac{1}{3}}\right)}{\dots}$

input

```
int(1/(x^3-3*x^2+7*x-5)^(1/3),x,method=_RETURNVERBOSE)
```

output

```
1/2*RootOf(_Z^2-_Z+1)*ln(-304*RootOf(_Z^2-_Z+1)^2*x^2+624*RootOf(_Z^2-_Z+1)
)*(x^3-3*x^2+7*x-5)^(2/3)+624*RootOf(_Z^2-_Z+1)*(x^3-3*x^2+7*x-5)^(1/3)*x+
608*RootOf(_Z^2-_Z+1)^2*x+928*RootOf(_Z^2-_Z+1)*x^2+51*(x^3-3*x^2+7*x-5)^(
2/3)-624*RootOf(_Z^2-_Z+1)*(x^3-3*x^2+7*x-5)^(1/3)+51*(x^3-3*x^2+7*x-5)^(1
/3)*x-1856*RootOf(_Z^2-_Z+1)*x-253*x^2-51*(x^3-3*x^2+7*x-5)^(1/3)+2356*Ro
otOf(_Z^2-_Z+1)+506*x-713)-1/2*ln(-304*RootOf(_Z^2-_Z+1)^2*x^2-624*RootOf(_
Z^2-_Z+1)*(x^3-3*x^2+7*x-5)^(2/3)-624*RootOf(_Z^2-_Z+1)*(x^3-3*x^2+7*x-5)^(
1/3)*x+608*RootOf(_Z^2-_Z+1)^2*x-320*RootOf(_Z^2-_Z+1)*x^2+675*(x^3-3*x^2
+7*x-5)^(2/3)+624*RootOf(_Z^2-_Z+1)*(x^3-3*x^2+7*x-5)^(1/3)+675*(x^3-3*x^2
+7*x-5)^(1/3)*x+640*RootOf(_Z^2-_Z+1)*x+371*x^2-675*(x^3-3*x^2+7*x-5)^(1/3
)-2356*RootOf(_Z^2-_Z+1)-742*x+1643)*RootOf(_Z^2-_Z+1)+1/2*ln(-304*RootOf(
_Z^2-_Z+1)^2*x^2-624*RootOf(_Z^2-_Z+1)*(x^3-3*x^2+7*x-5)^(2/3)-624*RootOf(
_Z^2-_Z+1)*(x^3-3*x^2+7*x-5)^(1/3)*x+608*RootOf(_Z^2-_Z+1)^2*x-320*RootOf(
_Z^2-_Z+1)*x^2+675*(x^3-3*x^2+7*x-5)^(2/3)+624*RootOf(_Z^2-_Z+1)*(x^3-3*x^
2+7*x-5)^(1/3)+675*(x^3-3*x^2+7*x-5)^(1/3)*x+640*RootOf(_Z^2-_Z+1)*x+371*x
^2-675*(x^3-3*x^2+7*x-5)^(1/3)-2356*RootOf(_Z^2-_Z+1)-742*x+1643)
```

Fricas [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.48

$$\int \frac{1}{\sqrt[3]{-5 + 7x - 3x^2 + x^3}} dx =$$

$$-\frac{1}{2} \sqrt{3} \arctan \left(\frac{22791076 \sqrt{3} (x^3 - 3x^2 + 7x - 5)^{\frac{1}{3}} (x - 1) + \sqrt{3} (20389537 x^2 - 40779074 x + 53222437)}{7204617 x^2 - 14409234 x - 20666867} \right)$$

$$-\frac{1}{4} \log \left(3 (x^3 - 3x^2 + 7x - 5)^{\frac{1}{3}} (x - 1) - 3 (x^3 - 3x^2 + 7x - 5)^{\frac{2}{3}} + 4 \right)$$

input `integrate(1/(x^3-3*x^2+7*x-5)^(1/3),x, algorithm="fricas")`

output `-1/2*sqrt(3)*arctan((22791076*sqrt(3)*(x^3 - 3*x^2 + 7*x - 5)^(1/3)*(x - 1) + sqrt(3)*(20389537*x^2 - 40779074*x + 53222437) + 17987998*sqrt(3)*(x^3 - 3*x^2 + 7*x - 5)^(2/3))/(7204617*x^2 - 14409234*x - 20666867)) - 1/4*log(3*(x^3 - 3*x^2 + 7*x - 5)^(1/3)*(x - 1) - 3*(x^3 - 3*x^2 + 7*x - 5)^(2/3) + 4)`

Sympy [F]

$$\int \frac{1}{\sqrt[3]{-5 + 7x - 3x^2 + x^3}} dx = \int \frac{1}{\sqrt[3]{x^3 - 3x^2 + 7x - 5}} dx$$

input `integrate(1/(x**3-3*x**2+7*x-5)**(1/3),x)`

output `Integral((x**3 - 3*x**2 + 7*x - 5)**(-1/3), x)`

Maxima [F]

$$\int \frac{1}{\sqrt[3]{-5+7x-3x^2+x^3}} dx = \int \frac{1}{(x^3-3x^2+7x-5)^{\frac{1}{3}}} dx$$

input `integrate(1/(x^3-3*x^2+7*x-5)^(1/3),x, algorithm="maxima")`

output `integrate((x^3 - 3*x^2 + 7*x - 5)^(-1/3), x)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{-5+7x-3x^2+x^3}} dx = \int \frac{1}{(x^3-3x^2+7x-5)^{\frac{1}{3}}} dx$$

input `integrate(1/(x^3-3*x^2+7*x-5)^(1/3),x, algorithm="giac")`

output `integrate((x^3 - 3*x^2 + 7*x - 5)^(-1/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{-5+7x-3x^2+x^3}} dx = \int \frac{1}{(x^3-3x^2+7x-5)^{\frac{1}{3}}} dx$$

input `int(1/(7*x - 3*x^2 + x^3 - 5)^(1/3),x)`

output `int(1/(7*x - 3*x^2 + x^3 - 5)^(1/3), x)`

Reduce [F]

$$\int \frac{1}{\sqrt[3]{-5 + 7x - 3x^2 + x^3}} dx = \int \frac{1}{(x^3 - 3x^2 + 7x - 5)^{\frac{1}{3}}} dx$$

input `int(1/(x^3-3*x^2+7*x-5)^(1/3),x)`

output `int(1/(x**3 - 3*x**2 + 7*x - 5)**(1/3),x)`

3.41
$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx$$

Optimal result	319
Mathematica [A] (verified)	319
Rubi [A] (warning: unable to verify)	320
Maple [A] (verified)	322
Fricas [B] (verification not implemented)	322
Sympy [F]	323
Maxima [F]	323
Giac [A] (verification not implemented)	324
Mupad [B] (verification not implemented)	324
Reduce [F]	325

Optimal result

Integrand size = 13, antiderivative size = 66

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx = \frac{1}{2}\sqrt{3} \arctan\left(\frac{1}{\sqrt{3}} + \frac{2x}{\sqrt{3}\sqrt[3]{x(-q+x^2)}}\right) + \frac{\log(x)}{4} - \frac{3}{4} \log\left(-x + \sqrt[3]{x(-q+x^2)}\right)$$

output

```
1/4*ln(x)-3/4*ln(-x+(x*(x^2-q))^(1/3))+1/2*arctan(1/3*3^(1/2)+2/3*x/(x*(x^2-q))^(1/3)*3^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.97

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx = \frac{\sqrt[3]{x}\sqrt[3]{-q+x^2} \left(2\sqrt{3} \arctan\left(\frac{\sqrt{3}x^{2/3}}{x^{2/3}+2\sqrt[3]{-q+x^2}}\right) - 2 \log\left(-x^{2/3} + \sqrt[3]{-q+x^2}\right) + \log\left(x^{4/3} + x^{2/3}\sqrt[3]{-q+x^2}\right) \right)}{4\sqrt[3]{-qx+x^3}}$$

input

```
Integrate[(x*(-q + x^2))^(1/3),x]
```


output

$$\frac{(x^{1/3}*(-q + x^2)^{1/3}*(2*\sqrt{3}*\text{ArcTan}[(\sqrt{3}*x^{2/3})/(x^{2/3} + 2*(-q + x^2)^{1/3})]) - 2*\text{Log}[-x^{2/3} + (-q + x^2)^{1/3}] + \text{Log}[x^{4/3} + x^{2/3}*(-q + x^2)^{1/3} + (-q + x^2)^{2/3}])}{4*(-(q*x) + x^3)^{1/3}}$$

Rubi [A] (warning: unable to verify)

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.30, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {2078, 1917, 266, 807, 769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt[3]{x(x^2 - q)}} dx \\ & \quad \downarrow \text{2078} \\ & \int \frac{1}{\sqrt[3]{x^3 - qx}} dx \\ & \quad \downarrow \text{1917} \\ & \frac{\sqrt[3]{x} \sqrt[3]{x^2 - q} \int \frac{1}{\sqrt[3]{x} \sqrt[3]{x^2 - q}} dx}{\sqrt[3]{x^3 - qx}} \\ & \quad \downarrow \text{266} \\ & \frac{3 \sqrt[3]{x} \sqrt[3]{x^2 - q} \int \frac{\sqrt[3]{x}}{\sqrt[3]{x^2 - q}} d\sqrt[3]{x}}{\sqrt[3]{x^3 - qx}} \\ & \quad \downarrow \text{807} \\ & \frac{3 \sqrt[3]{x} \sqrt[3]{x^2 - q} \int \frac{1}{\sqrt[3]{x - q}} dx^{2/3}}{2 \sqrt[3]{x^3 - qx}} \\ & \quad \downarrow \text{769} \end{aligned}$$

$$\frac{3\sqrt[3]{x}\sqrt[3]{x^2 - q} \left(\frac{\arctan\left(\frac{\sqrt[3]{x^2/3} + 1}{\sqrt[3]{x - q}}\right)}{\sqrt{3}} - \frac{1}{2} \log(\sqrt[3]{x - q} - x^{2/3}) \right)}{2\sqrt[3]{x^3 - qx}}$$

input `Int[(x*(-q + x^2))^(1/3),x]`

output `(3*x^(1/3)*(-q + x^2)^(1/3)*(ArcTan[(1 + (2*x^(2/3)))/(-q + x)^(1/3)]/Sqrt[3])/Sqrt[3] - Log[-x^(2/3) + (-q + x)^(1/3)]/2)/(2*(-q*x) + x^3)^(1/3)`

Definitions of rubi rules used

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntegerBinomialQ[a, b, c, 2, m, p, x]`

rule 769 `Int[((a_) + (b_.)*(x_)^3)^(1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^n)^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1 /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 1917 `Int[((a_.)*(x_)^(j_.) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Simp[(a*x^j + b*x^n)^FracPart[p]/(x^(j*FracPart[p])*(a + b*x^(n - j))^FracPart[p]) Int[x^(j*p)*(a + b*x^(n - j))^p, x], x] /; FreeQ[{a, b, j, n, p}, x] && !IntegerQ[p] && NeQ[n, j] && PosQ[n - j]`

rule 2078

```
Int[(u_)^(p_), x_Symbol] := Int[ExpandToSum[u, x]^p, x] /; FreeQ[p, x] && GeneralizedBinomialQ[u, x] && !GeneralizedBinomialMatchQ[u, x]
```

Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.39

method	result
pseudoelliptic	$\frac{\ln\left(\frac{(-x(-x^2+q))^{\frac{2}{3}} + (-x(-x^2+q))^{\frac{1}{3}}x+x^2}{x^2}\right)}{4} - \frac{\sqrt{3} \arctan\left(\frac{(x+2(-x(-x^2+q))^{\frac{1}{3}})\sqrt{3}}{3x}\right)}{2} - \frac{\ln\left(\frac{(-x(-x^2+q))^{\frac{1}{3}}-x}{x}\right)}{2}$

input

```
int(1/(x*(x^2-q))^(1/3),x,method=_RETURNVERBOSE)
```

output

```
1/4*ln(((x*(-x^2+q))^(2/3)+(-x*(-x^2+q))^(1/3)*x+x^2)/x^2)-1/2*3^(1/2)*arctan(1/3*(x+2*(-x*(-x^2+q))^(1/3))*3^(1/2)/x)-1/2*ln(((x*(-x^2+q))^(1/3)-x)/x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 415 vs. 2(52) = 104.

Time = 0.77 (sec) , antiderivative size = 415, normalized size of antiderivative = 6.29

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx$$

$$= \frac{1}{2} \sqrt{3} \arctan\left(\frac{4\sqrt{3}(q^{12} - 15q^{10} + 90q^8 - 351q^6 + 810q^4 - 1215q^2 + 729)(x^3 - qx)^{\frac{1}{3}}x - 2\sqrt{3}(q^{12} + 6q^9 - 12q^6 + 8q^3 - q)}{3(x^3 - qx)^{\frac{1}{3}}x + q + 3(x^3 - qx)^{\frac{2}{3}}}\right) - \frac{1}{4} \log\left(-3(x^3 - qx)^{\frac{1}{3}}x + q + 3(x^3 - qx)^{\frac{2}{3}}\right)$$

input

```
integrate(1/(x*(x^2-q))^(1/3),x, algorithm="fricas")
```

output

```

1/2*sqrt(3)*arctan((4*sqrt(3)*(q^12 - 15*q^10 + 90*q^8 - 351*q^6 + 810*q^4
- 1215*q^2 + 729)*(x^3 - q*x)^(1/3)*x - 2*sqrt(3)*(q^12 + 6*q^11 - 15*q^1
0 - 54*q^9 + 90*q^8 + 270*q^7 - 351*q^6 - 810*q^5 + 810*q^4 + 1458*q^3 - 1
215*q^2 - 1458*q + 729)*(x^3 - q*x)^(2/3) - sqrt(3)*(q^13 + 10*q^12 - 15*q
^11 - 282*q^10 + 90*q^9 + 2178*q^8 - 351*q^7 - 6534*q^6 + 810*q^5 + 7614*q
^4 - 1215*q^3 - (q^12 - 6*q^11 - 15*q^10 + 54*q^9 + 90*q^8 - 270*q^7 - 351
*q^6 + 810*q^5 + 810*q^4 - 1458*q^3 - 1215*q^2 + 1458*q + 729)*x^2 - 2430*
q^2 + 729*q))/(q^13 + 18*q^12 + 81*q^11 - 162*q^10 - 1350*q^9 + 810*q^8 +
6561*q^7 - 2430*q^6 - 12150*q^5 + 4374*q^4 + 6561*q^3 - 9*(q^12 + 2*q^11 -
15*q^10 - 18*q^9 + 90*q^8 + 90*q^7 - 351*q^6 - 270*q^5 + 810*q^4 + 486*q^
3 - 1215*q^2 - 486*q + 729)*x^2 - 4374*q^2 + 729*q)) - 1/4*log(-3*(x^3 - q
*x)^(1/3)*x + q + 3*(x^3 - q*x)^(2/3))

```

Sympy [F]

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx = \int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx$$

input

```
integrate(1/(x*(x**2-q))**(1/3),x)
```

output

```
Integral((x*(-q + x**2))**(-1/3), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx = \int \frac{1}{((x^2 - q)x)^{\frac{1}{3}}} dx$$

input

```
integrate(1/(x*(x^2-q))^(1/3),x, algorithm="maxima")
```

output

```
integrate(((x^2 - q)*x)^(-1/3), x)
```

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.02

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx = -\frac{1}{2} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \left(2 \left(-\frac{q}{x^2} + 1 \right)^{\frac{1}{3}} + 1 \right) \right) \\ + \frac{1}{4} \log \left(\left(-\frac{q}{x^2} + 1 \right)^{\frac{2}{3}} + \left(-\frac{q}{x^2} + 1 \right)^{\frac{1}{3}} + 1 \right) \\ - \frac{1}{2} \log \left(\left| \left(-\frac{q}{x^2} + 1 \right)^{\frac{1}{3}} - 1 \right| \right)$$

input `integrate(1/(x*(x^2-q))^(1/3),x, algorithm="giac")`output `-1/2*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-q/x^2 + 1)^(1/3) + 1)) + 1/4*log((-q/x^2 + 1)^(2/3) + (-q/x^2 + 1)^(1/3) + 1) - 1/2*log(abs((-q/x^2 + 1)^(1/3) - 1))`**Mupad [B] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.56

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx = \frac{3x \left(1 - \frac{x^2}{q}\right)^{1/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; \frac{x^2}{q}\right)}{2(x^3 - qx)^{1/3}}$$

input `int(1/(-x*(q - x^2))^(1/3),x)`output `(3*x*(1 - x^2/q)^(1/3)*hypergeom([1/3, 1/3], 4/3, x^2/q))/(2*(x^3 - q*x)^(1/3))`

Reduce [F]

$$\int \frac{1}{\sqrt[3]{x(-q+x^2)}} dx = \int \frac{1}{x^{\frac{1}{3}}(x^2-q)^{\frac{1}{3}}} dx$$

input `int(1/(x*(x^2-q))^(1/3),x)`

output `int(1/(x**(1/3)*(-q+x**2)**(1/3)),x)`

3.42 $\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx$

Optimal result	326
Mathematica [A] (verified)	326
Rubi [A] (warning: unable to verify)	327
Maple [F]	329
Fricas [B] (verification not implemented)	329
Sympy [F]	330
Maxima [F]	331
Giac [F]	331
Mupad [F(-1)]	331
Reduce [F]	332

Optimal result

Integrand size = 16, antiderivative size = 79

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \frac{1}{2}\sqrt{3} \arctan\left(\frac{1}{\sqrt{3}} + \frac{2(-1+x)}{\sqrt{3}\sqrt[3]{(-1+x)(q-2x+x^2)}}\right) + \frac{1}{4} \log(1-x) - \frac{3}{4} \log\left(1-x + \sqrt[3]{(-1+x)(q-2x+x^2)}\right)$$

output

```
1/4*ln(1-x)-3/4*ln(1-x+((-1+x)*(x^2+q-2*x))^(1/3))+1/2*arctan(1/3*3^(1/2)+2/3*(-1+x)/((-1+x)*(x^2+q-2*x))^(1/3)*3^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.84

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \frac{\sqrt[3]{-1+x}\sqrt[3]{q+(-2+x)x}\left(2\sqrt{3} \arctan\left(\frac{\sqrt{3}(-1+x)^{2/3}}{(-1+x)^{2/3}+2\sqrt[3]{q+(-2+x)x}}\right) - 2 \log\left(-(-1+x)^{2/3} + \sqrt[3]{q+(-2+x)x}\right)\right)}{4\sqrt[3]{(-1+x)(q+(-2+x)x)}}$$

input `Integrate[((-1 + x)*(q - 2*x + x^2))^(1/3), x]`

output
$$\frac{((-1 + x)^{1/3}*(q + (-2 + x)*x)^{1/3}*(2*\sqrt{3}*\text{ArcTan}[\sqrt{3}*(-1 + x)^{2/3}])/((-1 + x)^{2/3} + 2*(q + (-2 + x)*x)^{1/3}) - 2*\text{Log}[(-1 + x)^{2/3} + (q + (-2 + x)*x)^{1/3}] + \text{Log}[(-1 + x)^{4/3} + (-1 + x)^{2/3}*(q + (-2 + x)*x)^{1/3} + (q + (-2 + x)*x)^{2/3}]}{4*((-1 + x)*(q + (-2 + x)*x))^{1/3}}$$

Rubi [A] (warning: unable to verify)

Time = 0.23 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.19, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {2481, 1917, 266, 807, 769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{\sqrt[3]{(x-1)(q+x^2-2x)}} dx \\ & \quad \downarrow \text{2481} \\ & \int \frac{1}{\sqrt[3]{(x-1)^3 - (1-q)(x-1)}} d(x-1) \\ & \quad \downarrow \text{1917} \\ & \frac{\sqrt[3]{x-1} \sqrt[3]{q+(x-1)^2-1} \int \frac{1}{\sqrt[3]{(x-1)^2+q-1} \sqrt[3]{x-1}} d(x-1)}{\sqrt[3]{(x-1)^3 - (1-q)(x-1)}} \\ & \quad \downarrow \text{266} \\ & \frac{3\sqrt[3]{x-1} \sqrt[3]{q+(x-1)^2-1} \int \frac{\sqrt[3]{x-1}}{\sqrt[3]{(x-1)^2+q-1}} d\sqrt[3]{x-1}}{\sqrt[3]{(x-1)^3 - (1-q)(x-1)}} \\ & \quad \downarrow \text{807} \\ & \frac{3\sqrt[3]{x-1} \sqrt[3]{q+(x-1)^2-1} \int \frac{1}{\sqrt[3]{q+x-2}} d(x-1)^{2/3}}{2\sqrt[3]{(x-1)^3 - (1-q)(x-1)}} \end{aligned}$$

$$\begin{array}{c}
 \downarrow 769 \\
 \frac{3\sqrt[3]{x-1}\sqrt[3]{q+(x-1)^2-1} \left(\frac{\arctan\left(\frac{\frac{2(x-1)^{2/3}}{\sqrt[3]{q+x-2}}+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(\sqrt[3]{q+x-2}-x+1\right) \right)}{2\sqrt[3]{(x-1)^3-(1-q)(x-1)}}
 \end{array}$$

input `Int[((-1 + x)*(q - 2*x + x^2))^(1/3), x]`

output `(3*(-1 + q + (-1 + x)^2)^(1/3)*(-1 + x)^(1/3)*(ArcTan[(1 + (2*(-1 + x)^(2/3)))/(-2 + q + x)^(1/3)]/Sqrt[3])/Sqrt[3] - Log[1 - x + (-2 + q + x)^(1/3)]/2))/(2*(-((1 - q)*(-1 + x)) + (-1 + x)^3)^(1/3))`

Defintions of rubi rules used

rule 266 `Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^2)^(p_), x_Symbol] := With[{k = Denominator[m]}, Simp[k/c Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(2*k)/c^2))^(p), x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && FractionQ[m] && IntBinomialQ[a, b, c, 2, m, p, x]`

rule 769 `Int[((a_) + (b_.)*(x_)^3)^(1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

rule 807 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^n)^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Simp[1/k Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^(p), x], x, x^k], x] /; k != 1 /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]`

rule 1917

```
Int[((a_.)*(x_)^(j_.) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Simp[(a*x^j +
b*x^n)^FracPart[p]/(x^(j*FracPart[p])*(a + b*x^(n - j))^FracPart[p]) Int[
x^(j*p)*(a + b*x^(n - j))^p, x], x] /; FreeQ[{a, b, j, n, p}, x] && !Integ
erQ[p] && NeQ[n, j] && PosQ[n - j]
```

rule 2481

```
Int[(Px_)^(p_), x_Symbol] := With[{a = Coeff[Px, x, 0], b = Coeff[Px, x, 1]
, c = Coeff[Px, x, 2], d = Coeff[Px, x, 3]}, Subst[Int[Simp[(2*c^3 - 9*b*c*
d + 27*a*d^2)/(27*d^2) - (c^2 - 3*b*d)*(x/(3*d)) + d*x^3, x]^p, x], x, c/(3
*d) + x]] /; FreeQ[p, x] && PolyQ[Px, x, 3]
```

Maple [F]

$$\int \frac{1}{((-1+x)(x^2+q-2x))^{\frac{1}{3}}} dx$$

input

```
int(1/((-1+x)*(x^2+q-2*x))^(1/3),x)
```

output

```
int(1/((-1+x)*(x^2+q-2*x))^(1/3),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 665 vs. $2(65) = 130$.

Time = 0.63 (sec) , antiderivative size = 665, normalized size of antiderivative = 8.42

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \text{Too large to display}$$

input

```
integrate(1/((-1+x)*(x^2+q-2*x))^(1/3),x, algorithm="fricas")
```

output

```

1/2*sqrt(3)*arctan((2*sqrt(3)*(q^12 - 18*q^11 + 117*q^10 - 346*q^9 + 414*q
^8 - 18*q^7 + 69*q^6 - 774*q^5 - 234*q^4 + 1058*q^3 + 621*q^2 + 378*q - 53
9)*(x^3 + (q + 2)*x - 3*x^2 - q)^(2/3) + 4*sqrt(3)*(q^12 - 12*q^11 + 51*q^
10 - 70*q^9 - 90*q^8 + 288*q^7 - 57*q^6 + 54*q^5 - 810*q^4 + 320*q^3 + 291
*q^2 - (q^12 - 12*q^11 + 51*q^10 - 70*q^9 - 90*q^8 + 288*q^7 - 57*q^6 + 54
*q^5 - 810*q^4 + 320*q^3 + 291*q^2 + 714*q + 49)*x + 714*q + 49)*(x^3 + (q
+ 2)*x - 3*x^2 - q)^(1/3) - sqrt(3)*(q^13 - 22*q^12 + 177*q^11 - 514*q^10
- 434*q^9 + 5346*q^8 - 8247*q^7 - 4542*q^6 + 19638*q^5 - 8050*q^4 - 10343
*q^3 + (q^12 - 6*q^11 - 15*q^10 + 206*q^9 - 594*q^8 + 594*q^7 - 183*q^6 +
882*q^5 - 1386*q^4 - 418*q^3 - 39*q^2 + 1050*q + 637)*x^2 + 6186*q^2 - 2*(
q^12 - 6*q^11 - 15*q^10 + 206*q^9 - 594*q^8 + 594*q^7 - 183*q^6 + 882*q^5
- 1386*q^4 - 418*q^3 - 39*q^2 + 1050*q + 637)*x + 1501*q + 32))/(q^13 - 22
*q^12 + 249*q^11 - 1546*q^10 + 4702*q^9 - 4230*q^8 - 10623*q^7 + 25338*q^6
- 3546*q^5 - 31306*q^4 + 18817*q^3 + 9*(q^12 - 14*q^11 + 73*q^10 - 162*q^
9 + 78*q^8 + 186*q^7 - 15*q^6 - 222*q^5 - 618*q^4 + 566*q^3 + 401*q^2 + 60
2*q - 147)*x^2 + 9714*q^2 - 18*(q^12 - 14*q^11 + 73*q^10 - 162*q^9 + 78*q^
8 + 186*q^7 - 15*q^6 - 222*q^5 - 618*q^4 + 566*q^3 + 401*q^2 + 602*q - 147
)*x - 995*q + 8)) - 1/4*log(3*(x^3 + (q + 2)*x - 3*x^2 - q)^(1/3)*(x - 1)
+ q - 3*(x^3 + (q + 2)*x - 3*x^2 - q)^(2/3) - 1)

```

Sympy [F]

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \int \frac{1}{\sqrt[3]{(x-1)(q+x^2-2x)}} dx$$

input

```
integrate(1/((-1+x)*(x**2+q-2*x))**(1/3),x)
```

output

```
Integral(((x - 1)*(q + x**2 - 2*x))**(-1/3), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \int \frac{1}{((x^2+q-2x)(x-1))^{\frac{1}{3}}} dx$$

input `integrate(1/((-1+x)*(x^2+q-2*x))^(1/3),x, algorithm="maxima")`

output `integrate(((x^2 + q - 2*x)*(x - 1))^(-1/3), x)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \int \frac{1}{((x^2+q-2x)(x-1))^{\frac{1}{3}}} dx$$

input `integrate(1/((-1+x)*(x^2+q-2*x))^(1/3),x, algorithm="giac")`

output `integrate(((x^2 + q - 2*x)*(x - 1))^(-1/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \int \frac{1}{((x-1)(x^2-2x+q))^{1/3}} dx$$

input `int(1/((x - 1)*(q - 2*x + x^2))^(1/3),x)`

output `int(1/((x - 1)*(q - 2*x + x^2))^(1/3), x)`

Reduce **[F]**

$$\int \frac{1}{\sqrt[3]{(-1+x)(q-2x+x^2)}} dx = \int \frac{1}{(x^3 + qx - 3x^2 - q + 2x)^{\frac{1}{3}}} dx$$

input `int(1/((-1+x)*(x^2+q-2*x))^(1/3),x)`

output `int(1/(q*x - q + x**3 - 3*x**2 + 2*x)**(1/3),x)`

3.43
$$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx$$

Optimal result	333
Mathematica [A] (verified)	334
Rubi [F]	334
Maple [F]	335
Fricas [B] (verification not implemented)	335
Sympy [F]	336
Maxima [F]	337
Giac [F]	337
Mupad [F(-1)]	337
Reduce [F]	338

Optimal result

Integrand size = 21, antiderivative size = 118

$$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx$$

$$= \frac{\sqrt{3} \arctan\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{q}(-1+x)}{\sqrt{3}\sqrt[3]{(-1+x)(q-2qx+x^2)}}\right)}{2\sqrt[3]{q}} + \frac{\log(1-x)}{4\sqrt[3]{q}}$$

$$+ \frac{\log(x)}{2\sqrt[3]{q}} - \frac{3 \log\left(-\sqrt[3]{q}(-1+x) + \sqrt[3]{(-1+x)(q-2qx+x^2)}\right)}{4\sqrt[3]{q}}$$

output

```
1/4*ln(1-x)/q^(1/3)+1/2*ln(x)/q^(1/3)-3/4*ln(-q^(1/3)*(-1+x)+((-1+x)*(-2*q
*x+x^2+q))^(1/3))/q^(1/3)+1/2*arctan(1/3*3^(1/2)+2/3*q^(1/3)*(-1+x)/((-1+x
)*(-2*q*x+x^2+q))^(1/3)*3^(1/2))/q^(1/3)
```

Mathematica [A] (verified)

Time = 2.09 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.60

$$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx$$

$$= \frac{\sqrt[3]{-1+x} \sqrt[3]{q-2qx+x^2} \left(2\sqrt{3} \arctan \left(\frac{\sqrt{3} \sqrt[3]{q} (-1+x)^{2/3}}{\sqrt[3]{q} (-1+x)^{2/3} + 2 \sqrt[3]{q-2qx+x^2}} \right) - 2 \log \left(-\sqrt[3]{q} (-1+x)^{2/3} + \sqrt[3]{q-2qx+x^2} \right) \right)}{4 \sqrt[3]{q} \sqrt[3]{(-1+x)(q-2qx+x^2)}}$$

input `Integrate[1/(x*((-1 + x)*(q - 2*q*x + x^2))^(1/3)),x]`

output `((-1 + x)^(1/3)*(q - 2*q*x + x^2)^(1/3)*(2*Sqrt[3]*ArcTan[(Sqrt[3]*q^(1/3)*(-1 + x)^(2/3))/(q^(1/3)*(-1 + x)^(2/3) + 2*(q - 2*q*x + x^2)^(1/3)]) - 2*Log[-(q^(1/3)*(-1 + x)^(2/3)) + (q - 2*q*x + x^2)^(1/3)] + Log[q^(2/3)*(-1 + x)^(4/3) + q^(1/3)*(-1 + x)^(2/3)*(q - 2*q*x + x^2)^(1/3) + (q - 2*q*x + x^2)^(2/3)]))/(4*q^(1/3)*((-1 + x)*(q - 2*q*x + x^2))^(1/3))`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \sqrt[3]{(x-1)(-2qx+q+x^2)}} dx$$

↓ 2490

$$\int \frac{1}{\left(\frac{1}{3}(-2q-1) + \frac{1}{3}(2q+1) + x\right) \sqrt[3]{\left(\frac{1}{3}(-2q-1) + x\right)^3 - \frac{1}{3}(1-4q)(1-q) \left(\frac{1}{3}(-2q-1) + x\right) - \frac{2}{27}(1-q)^2}}$$

↓ 7299

$$\int \frac{1}{\left(\frac{1}{3}(-2q-1) + \frac{1}{3}(2q+1) + x\right) \sqrt[3]{\left(\frac{1}{3}(-2q-1) + x\right)^3 - \frac{1}{3}(1-4q)(1-q) \left(\frac{1}{3}(-2q-1) + x\right) - \frac{2}{27}(1-q)^2}}$$

input `Int[1/(x*((-1 + x)*(q - 2*q*x + x^2))^(1/3)),x]`

output `$Aborted`

Defintions of rubi rules used

rule 2490 `Int[(P3_)^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := With[{a = Coeff[P3, x, 0], b = Coeff[P3, x, 1], c = Coeff[P3, x, 2], d = Coeff[P3, x, 3]}, Subst[Int[((3*d*e - c*f)/(3*d) + f*x)^m*SimP[(2*c^3 - 9*b*c*d + 27*a*d^2)/(27*d^2) - (c^2 - 3*b*d)*(x/(3*d)) + d*x^3, x]^p, x], x, x + c/(3*d)] /; NeQ[c, 0]] /; FreeQ[{e, f, m, p}, x] && PolyQ[P3, x, 3]`

rule 7299 `Int[u_, x_] := CannotIntegrate[u, x]`

Maple [F]

$$\int \frac{1}{x((-1+x)(-2qx+x^2+q))^{\frac{1}{3}}} dx$$

input `int(1/x/((-1+x)*(-2*q*x+x^2+q))^(1/3),x)`

output `int(1/x/((-1+x)*(-2*q*x+x^2+q))^(1/3),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 661 vs. $2(100) = 200$.

Time = 13.55 (sec) , antiderivative size = 1496, normalized size of antiderivative = 12.68

$$\int \frac{1}{x^3 \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx = \text{Too large to display}$$

input `integrate(1/x/((-1+x)*(-2*q*x+x^2+q))^(1/3),x, algorithm="fricas")`

output

```
[1/12*(sqrt(3)*q*sqrt((-q)^(1/3)/q)*log(-((q^3 - 30*q^2 - 51*q - 1)*x^6 +
54*(q^3 + 6*q^2 + 2*q)*x^5 - 27*(17*q^3 + 26*q^2 + 2*q)*x^4 + 486*q^3*x +
540*(2*q^3 + q^2)*x^3 - 81*q^3 - 135*(8*q^3 + q^2)*x^2 + 9*((2*q^2 - q - 1)
)*x^4 - 6*(q^2 - q)*x^3 + 3*(q^2 - q)*x^2)*(-(2*q + 1)*x^2 + x^3 + 3*q*x -
q)^(2/3)*(-q)^(1/3) + 9*((q^2 + 7*q + 1)*x^5 - (19*q^2 + 25*q + 1)*x^4 +
9*(7*q^2 + 3*q)*x^3 + 45*q^2*x - 9*(9*q^2 + q)*x^2 - 9*q^2)*(-(2*q + 1)*x^
2 + x^3 + 3*q*x - q)^(1/3)*(-q)^(2/3) + sqrt(3)*(3*((4*q^2 + 13*q + 1)*x^4
- 6*(7*q^2 + 5*q)*x^3 - 72*q^2*x + 3*(31*q^2 + 5*q)*x^2 + 18*q^2)*(-(2*q
+ 1)*x^2 + x^3 + 3*q*x - q)^(2/3)*(-q)^(2/3) + 3*((q^3 - 5*q^2 - 5*q)*x^5
+ 5*(q^3 + 7*q^2 + q)*x^4 - 45*q^3*x - 45*(q^3 + q^2)*x^3 + 9*q^3 + 15*(5*
q^3 + q^2)*x^2)*(-(2*q + 1)*x^2 + x^3 + 3*q*x - q)^(1/3) + ((q^3 + 24*q^2
+ 3*q - 1)*x^6 - 54*(q^3 + 2*q^2)*x^5 + 81*(3*q^3 + 2*q^2)*x^4 - 162*q^3*x
- 108*(4*q^3 + q^2)*x^3 + 27*q^3 + 27*(14*q^3 + q^2)*x^2)*(-q)^(1/3))*sqrt
t((-q)^(1/3)/q)/x^6) - 2*(-q)^(2/3)*log(((q)^(2/3)*(q - 1)*x^2 + 3*(-(2*
q + 1)*x^2 + x^3 + 3*q*x - q)^(1/3)*(q*x - q)*(-q)^(1/3) + 3*(-(2*q + 1)*x
^2 + x^3 + 3*q*x - q)^(2/3)*q)/x^2) + (-q)^(2/3)*log((3*((2*q + 1)*x^2 - 6
*q*x + 3*q)*(-(2*q + 1)*x^2 + x^3 + 3*q*x - q)^(2/3)*(-q)^(2/3) + 3*((q^2
+ 2*q)*x^3 + 9*q^2*x - (7*q^2 + 2*q)*x^2 - 3*q^2)*(-(2*q + 1)*x^2 + x^3 +
3*q*x - q)^(1/3) - ((q^2 + 7*q + 1)*x^4 - 18*(q^2 + q)*x^3 - 36*q^2*x + 9*
(5*q^2 + q)*x^2 + 9*q^2)*(-q)^(1/3))/x^4))/q, 1/12*(2*sqrt(3)*q*sqrt(-(...
```

Sympy [F]

$$\int \frac{1}{x^3 \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx = \int \frac{1}{x^3 \sqrt[3]{(x-1)(-2qx+q+x^2)}} dx$$

input

```
integrate(1/x/((-1+x)*(-2*q*x+x**2+q))**(1/3),x)
```

output

```
Integral(1/(x*((x - 1)*(-2*q*x + q + x**2))**(1/3)), x)
```

Maxima [F]

$$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx = \int \frac{1}{(-(2qx-x^2-q)(x-1))^{\frac{1}{3}} x} dx$$

input `integrate(1/x/((-1+x)*(-2*q*x+x^2+q))^(1/3),x, algorithm="maxima")`

output `integrate(1/((-2*q*x - x^2 - q)*(x - 1))^(1/3)*x, x)`

Giac [F]

$$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx = \int \frac{1}{(-(2qx-x^2-q)(x-1))^{\frac{1}{3}} x} dx$$

input `integrate(1/x/((-1+x)*(-2*q*x+x^2+q))^(1/3),x, algorithm="giac")`

output `integrate(1/((-2*q*x - x^2 - q)*(x - 1))^(1/3)*x, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx = \int \frac{1}{x((x-1)(x^2-2qx+q))^{1/3}} dx$$

input `int(1/(x*((x - 1)*(q - 2*q*x + x^2))^(1/3)),x)`

output `int(1/(x*((x - 1)*(q - 2*q*x + x^2))^(1/3)), x)`

Reduce [F]

$$\int \frac{1}{x \sqrt[3]{(-1+x)(q-2qx+x^2)}} dx = \int \frac{1}{(-2qx^2+x^3+3qx-x^2-q)^{\frac{1}{3}} x} dx$$

input `int(1/x/((-1+x)*(-2*q*x+x^2+q))^(1/3),x)`

output `int(1/((-2*q*x**2 + 3*q*x - q + x**3 - x**2)**(1/3)*x),x)`

$$3.44 \quad \int \frac{2-(1+k)x}{\sqrt[3]{(1-x)x(1-kx)}(1-(1+k)x)} dx$$

Optimal result	339
Mathematica [A] (verified)	340
Rubi [F]	340
Maple [A] (verified)	342
Fricas [F(-1)]	342
Sympy [F]	343
Maxima [F]	343
Giac [F]	343
Mupad [F(-1)]	344
Reduce [F]	344

Optimal result

Integrand size = 36, antiderivative size = 111

$$\int \frac{2-(1+k)x}{\sqrt[3]{(1-x)x(1-kx)}(1-(1+k)x)} dx = \frac{\sqrt{3} \arctan\left(\frac{1+\frac{2\sqrt[3]{k}x}{\sqrt[3]{(1-x)x(1-kx)}}}{\sqrt{3}}\right)}{\sqrt[3]{k}} + \frac{\log(x)}{2\sqrt[3]{k}} + \frac{\log(1-(1+k)x)}{2\sqrt[3]{k}} - \frac{3 \log\left(-\sqrt[3]{k}x + \sqrt[3]{(1-x)x(1-kx)}\right)}{2\sqrt[3]{k}}$$

output

```
1/2*ln(x)/k^(1/3)+1/2*ln(1-(1+k)*x)/k^(1/3)-3/2*ln(-k^(1/3)*x+((1-x)*x*(-k*x+1))^(1/3))/k^(1/3)+arctan(1/3*(1+2*k^(1/3)*x/((1-x)*x*(-k*x+1))^(1/3))*3^(1/2))*3^(1/2)/k^(1/3)
```

Mathematica [A] (verified)

Time = 15.18 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.15

$$\int \frac{2 - (1+k)x}{\sqrt[3]{(1-x)x(1-kx)}(1 - (1+k)x)} dx$$

$$= \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{kx}}{\sqrt[3]{kx+2}\sqrt[3]{(-1+x)x(-1+kx)}}\right) - 2\log\left(-\sqrt[3]{kx} + \sqrt[3]{(-1+x)x(-1+kx)}\right) + \log\left(k^{2/3}x^2\right)}{2\sqrt[3]{k}}$$

input

```
Integrate[(2 - (1 + k)*x)/(((1 - x)*x*(1 - k*x))^(1/3)*(1 - (1 + k)*x)),x]
```

output

```
(2*Sqrt[3]*ArcTan[(Sqrt[3]*k^(1/3)*x)/(k^(1/3)*x + 2*((-1 + x)*x*(-1 + k*x))^(1/3)]) - 2*Log[-(k^(1/3)*x) + ((-1 + x)*x*(-1 + k*x))^(1/3)] + Log[k^(2/3)*x^2 + k^(1/3)*x*((-1 + x)*x*(-1 + k*x))^(1/3) + ((-1 + x)*x*(-1 + k*x))^(2/3)])/(2*k^(1/3))
```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2 - (k+1)x}{\sqrt[3]{(1-x)x(1-kx)}(1 - (k+1)x)} dx$$

$$\downarrow 2467$$

$$\frac{\sqrt[3]{x}\sqrt[3]{kx^2 - (k+1)x + 1} \int \frac{2 - (k+1)x}{\sqrt[3]{x(1-(k+1)x)}\sqrt[3]{kx^2 - (k+1)x + 1}} dx}{\sqrt[3]{(1-x)x(1-kx)}}$$

$$\downarrow 2035$$

$$\frac{3\sqrt[3]{x}\sqrt[3]{kx^2 - (k+1)x + 1} \int \frac{\sqrt[3]{x(2-(k+1)x)}}{(1-(k+1)x)\sqrt[3]{kx^2 - (k+1)x + 1}} d\sqrt[3]{x}}{\sqrt[3]{(1-x)x(1-kx)}}$$

$$\downarrow 7276$$

$$\frac{3\sqrt[3]{x}\sqrt[3]{kx^2 - (k+1)x + 1} \int \left(\frac{\sqrt[3]{x}}{(1-(k+1)x)\sqrt[3]{kx^2 - (k+1)x + 1}} + \frac{\sqrt[3]{x}}{\sqrt[3]{kx^2 - (k+1)x + 1}} \right) d\sqrt[3]{x}}{\sqrt[3]{(1-x)x(1-kx)}}$$

↓ 2009

$$\frac{3\sqrt[3]{x}\sqrt[3]{kx^2 - (k+1)x + 1} \left(\int \frac{\sqrt[3]{x}}{((-k-1)x+1)\sqrt[3]{kx^2 + (-k-1)x + 1}} d\sqrt[3]{x} + \frac{\sqrt[3]{1-xx^{2/3}}\sqrt[3]{1-kx} \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{3}, \frac{1}{3}, \frac{5}{3}, \frac{\sqrt[3]{x}}{\sqrt[3]{kx^2 - (k+1)x + 1}}\right)}{2\sqrt[3]{kx^2 - (k+1)x + 1}} \right)}{\sqrt[3]{(1-x)x(1-kx)}}$$

input

```
Int[(2 - (1 + k)*x)/(((1 - x)*x*(1 - k*x))^(1/3)*(1 - (1 + k)*x)), x]
```

output

```
$Aborted
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2035

```
Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

rule 2467

```
Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[Px^F
racPart[p]/(x^(r*FracPart[p])*ExpandToSum[Px/x^r, x]^FracPart[p]) Int[x^(
p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x], x] /; IGtQ[r, 0] /; FreeQ[p, x] && P
olyQ[Px, x] && !IntegerQ[p] && !MonomialQ[Px, x] && !PolyQ[Fx, x]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [A] (verified)

Time = 1.24 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00

method	result
pseudoelliptic	$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(k^{\frac{1}{3}}x + 2((-1+x)x(kx-1))^{\frac{1}{3}}\right)}{3k^{\frac{1}{3}}x}\right) - \ln\left(\frac{k^{\frac{2}{3}}x^2 + k^{\frac{1}{3}}((-1+x)x(kx-1))^{\frac{1}{3}}x + ((-1+x)x(kx-1))^{\frac{2}{3}}}{x^2}\right) + \ln\left(-k^{\frac{1}{3}}x + ((-1+x)x(kx-1))^{\frac{1}{3}}\right)}{k^{\frac{1}{3}}}$

input

```
int((2-(1+k)*x)/((1-x)*x*(-k*x+1))^(1/3)/(1-(1+k)*x),x,method=_RETURNVERBOSE)
```

output

```
-1/k^(1/3)*(3^(1/2)*arctan(1/3*3^(1/2)*(k^(1/3)*x+2*((-1+x)*x*(k*x-1))^(1/3)))/k^(1/3)/x-1/2*ln((k^(2/3)*x^2+k^(1/3)*((-1+x)*x*(k*x-1))^(1/3)*x+((-1+x)*x*(k*x-1))^(2/3))/x^2)+ln((-k^(1/3)*x+((-1+x)*x*(k*x-1))^(1/3))/x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{2 - (1 + k)x}{\sqrt[3]{(1 - x)x(1 - kx)}(1 - (1 + k)x)} dx = \text{Timed out}$$

input

```
integrate((2-(1+k)*x)/((1-x)*x*(-k*x+1))^(1/3)/(1-(1+k)*x),x, algorithm="fricas")
```

output

Timed out

Sympy [F]

$$\int \frac{2 - (1+k)x}{\sqrt[3]{(1-x)x(1-kx)}(1 - (1+k)x)} dx = \int \frac{kx + x - 2}{\sqrt[3]{x(x-1)(kx-1)}(kx + x - 1)} dx$$

input `integrate((2-(1+k)*x)/((1-x)*x*(-k*x+1))**(1/3)/(1-(1+k)*x), x)`

output `Integral((k*x + x - 2)/((x*(x - 1)*(k*x - 1))**(1/3)*(k*x + x - 1)), x)`

Maxima [F]

$$\int \frac{2 - (1+k)x}{\sqrt[3]{(1-x)x(1-kx)}(1 - (1+k)x)} dx = \int \frac{(k+1)x - 2}{((kx-1)(x-1)x)^{\frac{1}{3}}((k+1)x-1)} dx$$

input `integrate((2-(1+k)*x)/((1-x)*x*(-k*x+1))^(1/3)/(1-(1+k)*x), x, algorithm="maxima")`

output `integrate(((k + 1)*x - 2)/(((k*x - 1)*(x - 1)*x)^(1/3)*((k + 1)*x - 1)), x)`

Giac [F]

$$\int \frac{2 - (1+k)x}{\sqrt[3]{(1-x)x(1-kx)}(1 - (1+k)x)} dx = \int \frac{(k+1)x - 2}{((kx-1)(x-1)x)^{\frac{1}{3}}((k+1)x-1)} dx$$

input `integrate((2-(1+k)*x)/((1-x)*x*(-k*x+1))^(1/3)/(1-(1+k)*x), x, algorithm="giac")`

output `integrate(((k + 1)*x - 2)/(((k*x - 1)*(x - 1)*x)^(1/3)*((k + 1)*x - 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{2 - (1+k)x}{\sqrt[3]{(1-x)x(1-kx)(1-(1+k)x)}} dx = \int \frac{x(k+1) - 2}{(x(k+1) - 1)(x(kx - 1)(x - 1))^{1/3}} dx$$

input `int((x*(k + 1) - 2)/((x*(k + 1) - 1)*(x*(k*x - 1)*(x - 1))^(1/3)), x)`

output `int((x*(k + 1) - 2)/((x*(k + 1) - 1)*(x*(k*x - 1)*(x - 1))^(1/3)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{2 - (1+k)x}{\sqrt[3]{(1-x)x(1-kx)(1-(1+k)x)}} dx \\ &= \left(\int \frac{x}{x^{\frac{4}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}}k + x^{\frac{4}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}} - x^{\frac{1}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}}} dx \right) k \\ & \quad + \int \frac{x}{x^{\frac{4}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}}k + x^{\frac{4}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}} - x^{\frac{1}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}}} dx \\ & \quad - 2 \left(\int \frac{1}{x^{\frac{4}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}}k + x^{\frac{4}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}} - x^{\frac{1}{3}}(kx^2 - kx - x + 1)^{\frac{1}{3}}} dx \right) \end{aligned}$$

input `int((2-(1+k)*x)/((1-x)*x*(-k*x+1))^(1/3)/(1-(1+k)*x), x)`

output `int(x/(x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)*k*x + x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)*x - x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)), x)*k + int(x/(x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)*k*x + x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)*x - x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)), x) - 2*int(1/(x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)*k*x + x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)*x - x**(1/3)*(k*x**2 - k*x - x + 1)**(1/3)), x)`

3.45 $\int \frac{1-kx}{(1+(-2+k)x)((1-x)x(1-kx))^{2/3}} dx$

Optimal result	345
Mathematica [F]	346
Rubi [F]	346
Maple [F]	349
Fricas [B] (verification not implemented)	349
Sympy [F]	350
Maxima [F]	351
Giac [F]	351
Mupad [F(-1)]	351
Reduce [F]	352

Optimal result

Integrand size = 33, antiderivative size = 176

$$\int \frac{1-kx}{(1+(-2+k)x)((1-x)x(1-kx))^{2/3}} dx =$$

$$\frac{\sqrt{3} \arctan\left(\frac{1+\frac{\sqrt[3]{2(1-kx)}}{\sqrt[3]{1-k}\sqrt{(1-x)x(1-kx)}}}{\sqrt{3}}\right)}{2^{2/3}\sqrt[3]{1-k}} + \frac{\log(1-(2-k)x)}{2^{2/3}\sqrt[3]{1-k}}$$

$$+ \frac{\log(1-kx)}{2 \cdot 2^{2/3}\sqrt[3]{1-k}} - \frac{3 \log\left(-1+kx+2^{2/3}\sqrt[3]{1-k}\sqrt{(1-x)x(1-kx)}\right)}{2 \cdot 2^{2/3}\sqrt[3]{1-k}}$$

output

```
1/2*ln(1-(2-k)*x)*2^(1/3)/(1-k)^(1/3)+1/4*ln(-k*x+1)*2^(1/3)/(1-k)^(1/3)-3
/4*ln(-1+k*x+2^(2/3)*(1-k)^(1/3)*((1-x)*x*(-k*x+1))^(1/3))*2^(1/3)/(1-k)^(
1/3)-1/2*arctan(1/3*(1+2^(1/3)*(-k*x+1)/(1-k)^(1/3)/((1-x)*x*(-k*x+1))^(1/
3))*3^(1/2))*3^(1/2)*2^(1/3)/(1-k)^(1/3)
```

Mathematica [F]

$$\int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx = \int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx$$

input `Integrate[(1 - k*x)/((1 + (-2 + k)*x)*((1 - x)*x*(1 - k*x))^(2/3)), x]`

output `Integrate[(1 - k*x)/((1 + (-2 + k)*x)*((1 - x)*x*(1 - k*x))^(2/3)), x]`

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1 - kx}{((k - 2)x + 1)((1 - x)x(1 - kx))^{2/3}} dx \\ & \quad \downarrow \text{2467} \\ & \frac{x^{2/3}(kx^2 - (k + 1)x + 1)^{2/3} \int \frac{1 - kx}{x^{2/3}(1 - (2 - k)x)(kx^2 - (k + 1)x + 1)^{2/3}} dx}{((1 - x)x(1 - kx))^{2/3}} \\ & \quad \downarrow \text{2035} \\ & \frac{3x^{2/3}(kx^2 - (k + 1)x + 1)^{2/3} \int \frac{1 - kx}{(1 - (2 - k)x)(kx^2 - (k + 1)x + 1)^{2/3}} d\sqrt[3]{x}}{((1 - x)x(1 - kx))^{2/3}} \\ & \quad \downarrow \text{1395} \\ & \frac{3(1 - x)^{2/3}x^{2/3}(1 - kx)^{2/3} \int \frac{\sqrt[3]{1 - kx}}{(1 - x)^{2/3}(1 - (2 - k)x)} d\sqrt[3]{x}}{((1 - x)x(1 - kx))^{2/3}} \\ & \quad \downarrow \text{1028} \\ & \frac{3(1 - x)^{2/3}x^{2/3}(1 - kx)^{2/3} \left(\frac{k \int \frac{1}{(1 - x)^{2/3}(1 - kx)^{2/3}} d\sqrt[3]{x}}{2 - k} + \frac{2(1 - k) \int \frac{1}{(1 - x)^{2/3}(1 - (2 - k)x)(1 - kx)^{2/3}} d\sqrt[3]{x}}{2 - k} \right)}{((1 - x)x(1 - kx))^{2/3}} \\ & \quad \downarrow \text{905} \end{aligned}$$

$$\frac{3(1-x)^{2/3}x^{2/3}(1-kx)^{2/3} \left(\frac{2(1-k) \int \frac{1}{(1-x)^{2/3}(1-(2-k)x)(1-kx)^{2/3}} d\sqrt[3]{x}}{2-k} + \frac{k \sqrt[3]{x} \left(\frac{1-x}{1-kx}\right)^{2/3} \sqrt[3]{1-kx} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \frac{1-x}{1-kx}\right)}{(2-k)(1-x)^{2/3}} \right)}{((1-x)x(1-kx))^{2/3}}$$

↓ 1030

$$\frac{3(1-x)^{2/3}x^{2/3}(1-kx)^{2/3} \left(\frac{2(1-k) \int \left(\frac{1}{3(1-\sqrt[3]{2-k}\sqrt[3]{x})} \frac{1}{(1-x)^{2/3}(1-kx)^{2/3}} + \frac{1}{3(\sqrt[3]{-1}\sqrt[3]{2-k}\sqrt[3]{x+1})} \frac{1}{(1-x)^{2/3}(1-kx)^{2/3}} + \frac{1}{3(1-\sqrt[3]{2-k}\sqrt[3]{x})} \frac{1}{(1-x)^{2/3}(1-kx)^{2/3}} \right)}{2-k} \right)}{((1-x)x(1-kx))^{2/3}}$$

↓ 2009

$$\frac{3(1-x)^{2/3}x^{2/3}(1-kx)^{2/3} \left(\frac{2(1-k) \left(\frac{1}{3} \int \frac{1}{(1-\sqrt[3]{2-k}\sqrt[3]{x})} \frac{1}{(1-x)^{2/3}(1-kx)^{2/3}} d\sqrt[3]{x} + \frac{1}{3} \int \frac{1}{(\sqrt[3]{-1}\sqrt[3]{2-k}\sqrt[3]{x+1})} \frac{1}{(1-x)^{2/3}(1-kx)^{2/3}} d\sqrt[3]{x} \right)}{2-k} \right)}{((1-x)x(1-kx))^{2/3}}$$

input

```
Int[(1 - k*x)/((1 + (-2 + k)*x)*((1 - x)*x*(1 - k*x))^(2/3)), x]
```

output

```
$Aborted
```

Defintions of rubi rules used

rule 905

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[x*((a + b*x^n)^p/(c*(c*((a + b*x^n)/(a*(c + d*x^n))))^p*(c + d*x^n)^(1/n + p)))*Hypergeometric2F1[1/n, -p, 1 + 1/n, -(b*c - a*d)*(x^n/(a*(c + d*x^n)))]], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 1) + 1, 0]
```

rule 1028

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_
.)*(x_)^(n_))^(r_), x_Symbol] := Simp[d/b Int[(a + b*x^n)^(p + 1)*(c + d*
x^n)^(q - 1)*(e + f*x^n)^r, x], x] + Simp[(b*c - a*d)/b Int[(a + b*x^n)^p
*(c + d*x^n)^(q - 1)*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n, r
}, x] && ILtQ[p, 0] && GtQ[q, 0]
```

rule 1030

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_
.)*(x_)^(n_))^(r_), x_Symbol] := With[{u = ExpandIntegrand[(a + b*x^n)^p*(c
+ d*x^n)^q*(e + f*x^n)^r, x]}, Int[u, x] /; SumQ[u]] /; FreeQ[{a, b, c, d,
e, f, p, q, r}, x] && IGtQ[n, 0]
```

rule 1395

```
Int[(u_.)*((a_) + (c_.)*(x_)^(n2_.) + (b_.)*(x_)^(n_))^(p_)*((d_) + (e_.)*(
x_)^(n_))^(q_.), x_Symbol] := Simp[(a + b*x^n + c*x^(2*n))^FracPart[p]/((d
+ e*x^n)^FracPart[p]*(a/d + c*(x^n/e))^FracPart[p]) Int[u*(d + e*x^n)^(p
+ q)*(a/d + (c/e)*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, n, p, q}, x] && E
qQ[n2, 2*n] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[p] && !(EqQ[q,
1] && EqQ[n, 2])
```

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2035

```
Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x]] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

rule 2467

```
Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[Px^F
racPart[p]/(x^(r*FracPart[p]))*ExpandToSum[Px/x^r, x]^FracPart[p]) Int[x^(
p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x], x] /; IGtQ[r, 0] /; FreeQ[p, x] && P
olyQ[Px, x] && !IntegerQ[p] && !MonomialQ[Px, x] && !PolyQ[Fx, x]
```

Maple [F]

$$\int \frac{-kx + 1}{(1 + (-2 + k)x)((1 - x)x(-kx + 1))^{2/3}} dx$$

input `int((-k*x+1)/(1+(-2+k)*x)/((1-x)*x*(-k*x+1))^(2/3),x)`

output `int((-k*x+1)/(1+(-2+k)*x)/((1-x)*x*(-k*x+1))^(2/3),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 932 vs. 2(132) = 264.

Time = 52.26 (sec) , antiderivative size = 932, normalized size of antiderivative = 5.30

$$\int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx = \text{Too large to display}$$

input `integrate((-k*x+1)/(1+(-2+k)*x)/((1-x)*x*(-k*x+1))^(2/3),x, algorithm="fricas")`

output

```

1/6*sqrt(3)*2^(1/3)*arctan(1/3*(24*sqrt(3)*2^(1/3)*((k^5 - 3*k^4 - 4*k^3 +
22*k^2 - 24*k + 8)*x^4 - 2*(k^4 - 10*k^3 + 27*k^2 - 26*k + 8)*x^3 - 6*(k^
3 - 4*k^2 + 4*k - 1)*x^2 - 2*(k^2 - 1)*x + k - 1)*(k*x^3 - (k + 1)*x^2 + x
)^(2/3)/(k - 1)^(1/3) - 6*sqrt(3)*2^(2/3)*((k^6 + 27*k^5 - 40*k^4 - 20*k^3
+ 48*k^2 - 16*k)*x^5 - (33*k^5 + 55*k^4 - 220*k^3 + 132*k^2 + 16*k - 16)*
x^4 + 2*(55*k^4 - 55*k^3 - 66*k^2 + 82*k - 16)*x^3 - 2*(55*k^3 - 99*k^2 +
38*k + 6)*x^2 + (33*k^2 - 61*k + 28)*x - k + 1)*(k*x^3 - (k + 1)*x^2 + x)^(
1/3)/(k - 1)^(2/3) + sqrt(3)*((k^6 - 48*k^5 - 192*k^4 + 416*k^3 - 48*k^2
- 192*k + 64)*x^6 + 6*(7*k^5 + 104*k^4 - 80*k^3 - 176*k^2 + 176*k - 32)*x^
5 - 3*(139*k^4 + 256*k^3 - 768*k^2 + 352*k + 16)*x^4 + 4*(203*k^3 - 192*k^
2 - 120*k + 104)*x^3 - 3*(139*k^2 - 208*k + 64)*x^2 + 6*(7*k - 8)*x + 1))/
((k^6 + 96*k^5 - 48*k^4 - 160*k^3 + 240*k^2 - 192*k + 64)*x^6 - 6*(17*k^5
+ 64*k^4 - 112*k^3 + 80*k^2 - 80*k + 32)*x^5 + 3*(149*k^4 + 32*k^3 - 96*k^
2 - 160*k + 80)*x^4 - 4*(157*k^3 - 24*k^2 - 168*k + 40)*x^3 + 3*(149*k^2 -
128*k - 16)*x^2 - 6*(17*k - 16)*x + 1))/(k - 1)^(1/3) - 1/12*2^(1/3)*log(
(12*2^(2/3)*(k*x^3 - (k + 1)*x^2 + x)^(2/3)*((k^3 + k^2 - 4*k + 2)*x^2 - 2
*(2*k^2 - 3*k + 1)*x + k - 1)/(k - 1)^(2/3) + 6*((k^3 + 8*k^2 - 8*k)*x^3 -
(11*k^2 - 8)*x^2 + (11*k - 8)*x - 1)*(k*x^3 - (k + 1)*x^2 + x)^(1/3) + 2^(
1/3)*((k^4 + 28*k^3 - 12*k^2 - 32*k + 16)*x^4 - 4*(8*k^3 + 15*k^2 - 30*k
+ 8)*x^3 + 6*(13*k^2 - 10*k - 2)*x^2 - 4*(8*k - 7)*x + 1)/(k - 1)^(1/3)...

```

Sympy [F]

$$\int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx =$$

$$- \int \frac{kx}{kx(kx^3 - kx^2 - x^2 + x)^{2/3} - 2x(kx^3 - kx^2 - x^2 + x)^{2/3} + (kx^3 - kx^2 - x^2 + x)^{2/3}} dx$$

$$- \int \left(-\frac{1}{kx(kx^3 - kx^2 - x^2 + x)^{2/3} - 2x(kx^3 - kx^2 - x^2 + x)^{2/3} + (kx^3 - kx^2 - x^2 + x)^{2/3}} \right) dx$$

input

```
integrate((-k*x+1)/(1+(-2+k)*x)/((1-x)*x*(-k*x+1))**(2/3), x)
```

output

```

-Integral(k*x/(k*x*(k*x**3 - k*x**2 - x**2 + x)**(2/3) - 2*x*(k*x**3 - k*x
**2 - x**2 + x)**(2/3) + (k*x**3 - k*x**2 - x**2 + x)**(2/3)), x) - Integr
al(-1/(k*x*(k*x**3 - k*x**2 - x**2 + x)**(2/3) - 2*x*(k*x**3 - k*x**2 - x
**2 + x)**(2/3) + (k*x**3 - k*x**2 - x**2 + x)**(2/3)), x)

```

Maxima [F]

$$\int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx = \int -\frac{kx - 1}{((kx - 1)(x - 1)x)^{2/3} ((k - 2)x + 1)} dx$$

input `integrate((-k*x+1)/(1+(-2+k)*x)/((1-x)*x*(-k*x+1))^(2/3),x, algorithm="maxima")`

output `-integrate((k*x - 1)/(((k*x - 1)*(x - 1)*x)^(2/3))*((k - 2)*x + 1), x)`

Giac [F]

$$\int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx = \int -\frac{kx - 1}{((kx - 1)(x - 1)x)^{2/3} ((k - 2)x + 1)} dx$$

input `integrate((-k*x+1)/(1+(-2+k)*x)/((1-x)*x*(-k*x+1))^(2/3),x, algorithm="giac")`

output `integrate(-(k*x - 1)/(((k*x - 1)*(x - 1)*x)^(2/3))*((k - 2)*x + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx = -\int \frac{kx - 1}{(x(k - 2) + 1)(x(kx - 1)(x - 1))^{2/3}} dx$$

input `int(-(k*x - 1)/((x*(k - 2) + 1)*(x*(k*x - 1)*(x - 1))^(2/3)),x)`

output `-int((k*x - 1)/((x*(k - 2) + 1)*(x*(k*x - 1)*(x - 1))^(2/3)), x)`

Reduce [F]

$$\int \frac{1 - kx}{(1 + (-2 + k)x)((1 - x)x(1 - kx))^{2/3}} dx =$$

$$- \left(\int \frac{x}{x^{5/3} (kx^2 - kx - x + 1)^{2/3} k - 2x^{5/3} (kx^2 - kx - x + 1)^{2/3} + x^{2/3} (kx^2 - kx - x + 1)^{2/3}} dx \right) k$$

$$+ \int \frac{1}{x^{5/3} (kx^2 - kx - x + 1)^{2/3} k - 2x^{5/3} (kx^2 - kx - x + 1)^{2/3} + x^{2/3} (kx^2 - kx - x + 1)^{2/3}} dx$$

input

```
int((-k*x+1)/(1+(-2+k)*x)/((1-x)*x*(-k*x+1))^(2/3),x)
```

output

```
- int(x/(x**(2/3)*(k*x**2 - k*x - x + 1)**(2/3)*k*x - 2*x**(2/3)*(k*x**2 - k*x - x + 1)**(2/3)*x + x**(2/3)*(k*x**2 - k*x - x + 1)**(2/3)),x)*k + int(1/(x**(2/3)*(k*x**2 - k*x - x + 1)**(2/3)*k*x - 2*x**(2/3)*(k*x**2 - k*x - x + 1)**(2/3)*x + x**(2/3)*(k*x**2 - k*x - x + 1)**(2/3)),x)
```

$$3.46 \quad \int \frac{a+bx+cx^2}{(1-x+x^2)\sqrt[3]{1-x^3}} dx$$

Optimal result	354
Mathematica [F]	355
Rubi [A] (verified)	355
Maple [F]	357
Fricas [F(-2)]	358
Sympy [F]	358
Maxima [F]	358
Giac [F]	359
Mupad [F(-1)]	359
Reduce [F]	359

Optimal result

Integrand size = 32, antiderivative size = 493

$$\begin{aligned}
& \int \frac{a + bx + cx^2}{(1 - x + x^2) \sqrt[3]{1 - x^3}} dx \\
&= \frac{(a + b) \arctan \left(\frac{1 - \frac{2 \sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt[3]{2}\sqrt{3}} + \frac{(a + b) \arctan \left(\frac{1 + \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{2\sqrt[3]{2}\sqrt{3}} \\
&\quad - \frac{c \arctan \left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} - \frac{(a - c) \arctan \left(\frac{1 - \frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt[3]{2}\sqrt{3}} \\
&\quad + \frac{(b + c) \arctan \left(\frac{1 + 2^{2/3} \sqrt[3]{1-x^3}}{\sqrt{3}} \right)}{\sqrt[3]{2}\sqrt{3}} + \frac{(a + b) \log((1-x)(1+x)^2)}{12\sqrt[3]{2}} \\
&\quad - \frac{(a - c) \log(1+x^3)}{6\sqrt[3]{2}} - \frac{(b + c) \log(1+x^3)}{6\sqrt[3]{2}} \\
&\quad + \frac{(a + b) \log \left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} \right)}{6\sqrt[3]{2}} - \frac{(a + b) \log \left(1 + \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} \right)}{3\sqrt[3]{2}} \\
&\quad + \frac{(b + c) \log(\sqrt[3]{2} - \sqrt[3]{1-x^3})}{2\sqrt[3]{2}} + \frac{(a - c) \log(-\sqrt[3]{2}x - \sqrt[3]{1-x^3})}{2\sqrt[3]{2}} \\
&\quad + \frac{1}{2}c \log(x + \sqrt[3]{1-x^3}) - \frac{(a + b) \log(-1 + x + 2^{2/3} \sqrt[3]{1-x^3})}{4\sqrt[3]{2}}
\end{aligned}$$

output

```

1/24*(a+b)*ln((1-x)*(1+x)^2)*2^(2/3)-1/12*(a-c)*ln(x^3+1)*2^(2/3)-1/12*(b+
c)*ln(x^3+1)*2^(2/3)+1/12*(a+b)*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3
))*(1-x)/(-x^3+1)^(1/3))*2^(2/3)-1/6*(a+b)*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3
))*2^(2/3)+1/4*(b+c)*ln(2^(1/3)-(-x^3+1)^(1/3))*2^(2/3)+1/4*(a-c)*ln(-2^(1
/3)*x-(-x^3+1)^(1/3))*2^(2/3)+1/2*c*ln(x+(-x^3+1)^(1/3))-1/8*(a+b)*ln(-1+x
+2^(2/3)*(-x^3+1)^(1/3))*2^(2/3)+1/6*(a+b)*arctan(1/3*(1-2*2^(1/3)*(1-x)/(
-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)+1/12*(a+b)*arctan(1/3*(1+2^(1/3)*(
1-x)/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)-1/3*c*arctan(1/3*(1-2*x/(-x^
3+1)^(1/3))*3^(1/2))*3^(1/2)-1/6*(a-c)*arctan(1/3*(1-2*2^(1/3)*x/(-x^3+1)^(
1/3))*3^(1/2))*2^(2/3)*3^(1/2)+1/6*(b+c)*arctan(1/3*(1+2^(2/3)*(-x^3+1)^(
1/3))*3^(1/2))*2^(2/3)*3^(1/2)

```

Mathematica [F]

$$\int \frac{a + bx + cx^2}{(1 - x + x^2)\sqrt[3]{1 - x^3}} dx = \int \frac{a + bx + cx^2}{(1 - x + x^2)\sqrt[3]{1 - x^3}} dx$$

input

```
Integrate[(a + b*x + c*x^2)/((1 - x + x^2)*(1 - x^3)^(1/3)), x]
```

output

```
Integrate[(a + b*x + c*x^2)/((1 - x + x^2)*(1 - x^3)^(1/3)), x]
```

Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 570, normalized size of antiderivative = 1.16, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + bx + cx^2}{(x^2 - x + 1)\sqrt[3]{1 - x^3}} dx$$

↓ 2583

$$\begin{aligned}
& \int \left(\frac{x(a+b)}{\sqrt[3]{1-x^3}(x^3+1)} + \frac{a}{\sqrt[3]{1-x^3}(x^3+1)} + \frac{x^2(b+c)}{\sqrt[3]{1-x^3}(x^3+1)} + \frac{cx^3}{\sqrt[3]{1-x^3}(x^3+1)} \right) dx \\
& \quad \downarrow \text{2009} \\
& \frac{(a+b) \arctan \left(\frac{1 - \frac{2}{3} \sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} \right)}{\sqrt[3]{2}\sqrt{3}} + \frac{(a+b) \arctan \left(\frac{\frac{\sqrt[3]{2(1-x)}}{3} + 1}{\sqrt[3]{1-x^3}} \right)}{2\sqrt[3]{2}\sqrt{3}} - \frac{a \arctan \left(\frac{1 - \frac{2}{3} \sqrt[3]{2x}}{\sqrt[3]{1-x^3}} \right)}{\sqrt[3]{2}\sqrt{3}} + \\
& \frac{(a+b) \log \left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1 \right)}{6\sqrt[3]{2}} - \frac{(a+b) \log \left(\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} - \\
& \frac{(a+b) \log \left(2^{2/3} \sqrt[3]{1-x^3} + x - 1 \right)}{4\sqrt[3]{2}} + \frac{(a+b) \log \left((1-x)(x+1)^2 \right)}{12\sqrt[3]{2}} - \frac{a \log(x^3+1)}{6\sqrt[3]{2}} + \\
& \frac{a \log \left(-\sqrt[3]{1-x^3} - \sqrt[3]{2x} \right)}{2\sqrt[3]{2}} + \frac{\arctan \left(\frac{2^{2/3} \sqrt[3]{1-x^3} + 1}{\sqrt{3}} \right) (b+c)}{\sqrt[3]{2}\sqrt{3}} - \frac{c \arctan \left(\frac{1 - \frac{2}{3} \sqrt[3]{2x}}{\sqrt[3]{1-x^3}} \right)}{\sqrt{3}} + \\
& \frac{c \arctan \left(\frac{1 - \frac{2}{3} \sqrt[3]{2x}}{\sqrt[3]{1-x^3}} \right)}{\sqrt[3]{2}\sqrt{3}} - \frac{(b+c) \log(x^3+1)}{6\sqrt[3]{2}} + \frac{(b+c) \log \left(\sqrt[3]{2} - \sqrt[3]{1-x^3} \right)}{2\sqrt[3]{2}} + \\
& \frac{c \log(x^3+1)}{6\sqrt[3]{2}} - \frac{c \log \left(-\sqrt[3]{1-x^3} - \sqrt[3]{2x} \right)}{2\sqrt[3]{2}} + \frac{1}{2} c \log \left(\sqrt[3]{1-x^3} + x \right)
\end{aligned}$$

input

```
Int[(a + b*x + c*x^2)/((1 - x + x^2)*(1 - x^3)^(1/3)),x]
```

output

```
((a + b)*ArcTan[(1 - (2*2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + ((a + b)*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/(2*2^(1/3)*Sqrt[3]) - (c*ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - (a*ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + (c*ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + ((b + c)*ArcTan[(1 + 2^(2/3)*(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + ((a + b)*Log[(1 - x)*(1 + x)^2])/(12*2^(1/3)) - (a*Log[1 + x^3])/(6*2^(1/3)) + (c*Log[1 + x^3])/(6*2^(1/3)) - ((b + c)*Log[1 + x^3])/((6*2^(1/3)) + ((a + b)*Log[1 + (2^(2/3)*(1 - x)^2)/(1 - x^3)^(2/3) - (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)])/(6*2^(1/3)) - ((a + b)*Log[1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)])/(3*2^(1/3)) + ((b + c)*Log[2^(1/3) - (1 - x^3)^(1/3)])/(2*2^(1/3)) + (a*Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)])/(2*2^(1/3)) - (c*Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)])/(2*2^(1/3)) + (c*Log[x + (1 - x^3)^(1/3)])/2 - ((a + b)*Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3)])/(4*2^(1/3))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2583

```
Int[(Px_.)*((c_) + (d_.)*(x_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^3)^(p_.), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]
```

Maple [F]

$$\int \frac{cx^2 + bx + a}{(x^2 - x + 1)(-x^3 + 1)^{\frac{1}{3}}} dx$$

input

```
int((c*x^2+b*x+a)/(x^2-x+1)/(-x^3+1)^(1/3),x)
```

output

```
int((c*x^2+b*x+a)/(x^2-x+1)/(-x^3+1)^(1/3),x)
```

Fricas [F(-2)]

Exception generated.

$$\int \frac{a + bx + cx^2}{(1 - x + x^2) \sqrt[3]{1 - x^3}} dx = \text{Exception raised: TypeError}$$

input `integrate((c*x^2+b*x+a)/(x^2-x+1)/(-x^3+1)^(1/3),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (residue poly has multiple non-linear factors)`

Sympy [F]

$$\int \frac{a + bx + cx^2}{(1 - x + x^2) \sqrt[3]{1 - x^3}} dx = \int \frac{a + bx + cx^2}{\sqrt[3]{-(x - 1)(x^2 + x + 1)(x^2 - x + 1)}} dx$$

input `integrate((c*x**2+b*x+a)/(x**2-x+1)/(-x**3+1)**(1/3),x)`

output `Integral((a + b*x + c*x**2)/((-x - 1)*(x**2 + x + 1))**(1/3)*(x**2 - x + 1)), x)`

Maxima [F]

$$\int \frac{a + bx + cx^2}{(1 - x + x^2) \sqrt[3]{1 - x^3}} dx = \int \frac{cx^2 + bx + a}{(-x^3 + 1)^{\frac{1}{3}}(x^2 - x + 1)} dx$$

input `integrate((c*x^2+b*x+a)/(x^2-x+1)/(-x^3+1)^(1/3),x, algorithm="maxima")`

output `integrate((c*x^2 + b*x + a)/((-x^3 + 1)^(1/3)*(x^2 - x + 1)), x)`

Giac [F]

$$\int \frac{a + bx + cx^2}{(1 - x + x^2) \sqrt[3]{1 - x^3}} dx = \int \frac{cx^2 + bx + a}{(-x^3 + 1)^{\frac{1}{3}}(x^2 - x + 1)} dx$$

input `integrate((c*x^2+b*x+a)/(x^2-x+1)/(-x^3+1)^(1/3),x, algorithm="giac")`

output `integrate((c*x^2 + b*x + a)/((-x^3 + 1)^(1/3)*(x^2 - x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + bx + cx^2}{(1 - x + x^2) \sqrt[3]{1 - x^3}} dx = \int \frac{cx^2 + bx + a}{(1 - x^3)^{\frac{1}{3}}(x^2 - x + 1)} dx$$

input `int((a + b*x + c*x^2)/((1 - x^3)^(1/3)*(x^2 - x + 1)),x)`

output `int((a + b*x + c*x^2)/((1 - x^3)^(1/3)*(x^2 - x + 1)), x)`

Reduce [F]

$$\begin{aligned} \int \frac{a + bx + cx^2}{(1 - x + x^2) \sqrt[3]{1 - x^3}} dx &= \left(\int \frac{x^2}{(-x^3 + 1)^{\frac{1}{3}} x^2 - (-x^3 + 1)^{\frac{1}{3}} x + (-x^3 + 1)^{\frac{1}{3}}} dx \right) c \\ &+ \left(\int \frac{x}{(-x^3 + 1)^{\frac{1}{3}} x^2 - (-x^3 + 1)^{\frac{1}{3}} x + (-x^3 + 1)^{\frac{1}{3}}} dx \right) b \\ &+ \left(\int \frac{1}{(-x^3 + 1)^{\frac{1}{3}} x^2 - (-x^3 + 1)^{\frac{1}{3}} x + (-x^3 + 1)^{\frac{1}{3}}} dx \right) a \end{aligned}$$

input `int((c*x^2+b*x+a)/(x^2-x+1)/(-x^3+1)^(1/3),x)`

output

```
int(x**2/((- x**3 + 1)**(1/3)*x**2 - (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)*c + int(x/((- x**3 + 1)**(1/3)*x**2 - (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)*b + int(1/((- x**3 + 1)**(1/3)*x**2 - (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)*a
```

$$3.47 \quad \int \frac{1}{(3-2x)^{11/2}(1+x+2x^2)^5} dx$$

Optimal result	361
Mathematica [C] (verified)	362
Rubi [A] (verified)	363
Maple [A] (verified)	370
Fricas [B] (verification not implemented)	371
Sympy [F(-1)]	372
Maxima [F]	372
Giac [B] (verification not implemented)	372
Mupad [B] (verification not implemented)	373
Reduce [B] (verification not implemented)	374

Optimal result

Integrand size = 20, antiderivative size = 407

$$\int \frac{1}{(3-2x)^{11/2}(1+x+2x^2)^5} dx = -\frac{19255}{395136(3-2x)^{9/2}} - \frac{462025}{30118144(3-2x)^{7/2}}$$

$$- \frac{38491}{8605184(3-2x)^{5/2}} - \frac{141045}{120472576(3-2x)^{3/2}} - \frac{38225}{240945152\sqrt{3-2x}}$$

$$+ \frac{x}{28(3-2x)^{9/2}(1+x+2x^2)^4} + \frac{23+73x}{1176(3-2x)^{9/2}(1+x+2x^2)^3}$$

$$+ \frac{1387+3049x}{32928(3-2x)^{9/2}(1+x+2x^2)^2} + \frac{5(3049+4377x)}{153664(3-2x)^{9/2}(1+x+2x^2)}$$

$$+ \frac{5\sqrt{\frac{1}{2}}(149046503977+40815066112\sqrt{14}) \arctan\left(\frac{\sqrt{7+2\sqrt{14}-2\sqrt{3-2x}}}{\sqrt{-7+2\sqrt{14}}}\right)}{3373232128}$$

$$- \frac{5\sqrt{\frac{1}{2}}(149046503977+40815066112\sqrt{14}) \arctan\left(\frac{\sqrt{7+2\sqrt{14}+2\sqrt{3-2x}}}{\sqrt{-7+2\sqrt{14}}}\right)}{3373232128}$$

$$+ \frac{5\sqrt{\frac{1}{2}}(-149046503977+40815066112\sqrt{14}) \log\left(3+\sqrt{14}-\sqrt{7+2\sqrt{14}}\sqrt{3-2x}-2x\right)}{6746464256}$$

$$- \frac{5\sqrt{\frac{1}{2}}(-149046503977+40815066112\sqrt{14}) \log\left(3+\sqrt{14}+\sqrt{7+2\sqrt{14}}\sqrt{3-2x}-2x\right)}{6746464256}$$

output

```
-19255/395136/(3-2*x)^(9/2)-462025/30118144/(3-2*x)^(7/2)-38491/8605184/(3-2*x)^(5/2)-141045/120472576/(3-2*x)^(3/2)+1/28*x/(3-2*x)^(9/2)/(2*x^2+x+1)^4+1/1176*(23+73*x)/(3-2*x)^(9/2)/(2*x^2+x+1)^3+1/32928*(1387+3049*x)/(3-2*x)^(9/2)/(2*x^2+x+1)^2+5/153664*(3049+4377*x)/(3-2*x)^(9/2)/(2*x^2+x+1)-38225/240945152/(3-2*x)^(1/2)+5/13492928512*ln(3-2*x+14^(1/2)-(3-2*x)^(1/2))*(7+2*14^(1/2))^(1/2))*(-298093007954+81630132224*14^(1/2))^(1/2)-5/13492928512*ln(3-2*x+14^(1/2)+(3-2*x)^(1/2))*(7+2*14^(1/2))^(1/2))*(-298093007954+81630132224*14^(1/2))^(1/2)+5/6746464256*arctan((-2*(3-2*x)^(1/2)+(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2))*(298093007954+81630132224*14^(1/2))^(1/2)-5/6746464256*arctan((2*(3-2*x)^(1/2)+(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2))*(298093007954+81630132224*14^(1/2))^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.25 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.44

$$\int \frac{1}{(3-2x)^{11/2}(1+x+2x^2)^5} dx = \frac{-14(40289347-429812744x+135202154x^2-1073855156x^3+1627773523x^4-1470758860x^5+2888625656x^6-3106712560x^7+2343370048x^8-2443779648x^9+1873554048x^{10}-677249280x^{11}+88070400x^{12})}{(3-2x)^{9/2}(1+x+2x^2)^4} - 45\sqrt{149046503977+(12577271771I)*\text{Sqrt}[7]}*\text{ArcTan}[(\text{Sqrt}[-1-I/\text{Sqrt}[7]]*\text{Sqrt}[3-2*x])/2] - 45\sqrt{149046503977-(12577271771I)*\text{Sqrt}[7]}*\text{ArcTan}[(\text{Sqrt}[-1+I/\text{Sqrt}[7]]*\text{Sqrt}[3-2*x])/2]}/30359089152$$

input

```
Integrate[1/((3 - 2*x)^(11/2)*(1 + x + 2*x^2)^5),x]
```

output

```
((-14*(40289347 - 429812744*x + 135202154*x^2 - 1073855156*x^3 + 1627773523*x^4 - 1470758860*x^5 + 2888625656*x^6 - 3106712560*x^7 + 2343370048*x^8 - 2443779648*x^9 + 1873554048*x^10 - 677249280*x^11 + 88070400*x^12))/((3 - 2*x)^(9/2)*(1 + x + 2*x^2)^4) - 45*Sqrt[149046503977 + (12577271771*I)*Sqrt[7]]*ArcTan[(Sqrt[-1 - I/Sqrt[7]]*Sqrt[3 - 2*x])/2] - 45*Sqrt[149046503977 - (12577271771*I)*Sqrt[7]]*ArcTan[(Sqrt[-1 + I/Sqrt[7]]*Sqrt[3 - 2*x])/2])/30359089152
```

Rubi [A] (verified)

Time = 0.88 (sec) , antiderivative size = 506, normalized size of antiderivative = 1.24, number of steps used = 27, number of rules used = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 1.300$, Rules used = {1165, 27, 1235, 27, 1235, 27, 1235, 27, 1198, 27, 1198, 27, 1198, 27, 1198, 27, 1198, 27, 1197, 27, 1483, 1142, 25, 1083, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3-2x)^{11/2} (2x^2+x+1)^5} dx \\
 & \quad \downarrow \text{1165} \\
 & \frac{1}{784} \int \frac{28(25-23x)}{(3-2x)^{11/2} (2x^2+x+1)^4} dx + \frac{x}{28(3-2x)^{9/2} (2x^2+x+1)^4} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{28} \int \frac{25-23x}{(3-2x)^{11/2} (2x^2+x+1)^4} dx + \frac{x}{28(3-2x)^{9/2} (2x^2+x+1)^4} \\
 & \quad \downarrow \text{1235} \\
 & \frac{1}{28} \left(\frac{1}{588} \int \frac{14(831-1387x)}{(3-2x)^{11/2} (2x^2+x+1)^3} dx + \frac{73x+23}{42(3-2x)^{9/2} (2x^2+x+1)^3} \right) + \\
 & \quad \frac{1}{28(3-2x)^{9/2} (2x^2+x+1)^4} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{28} \left(\frac{1}{42} \int \frac{831-1387x}{(3-2x)^{11/2} (2x^2+x+1)^3} dx + \frac{73x+23}{42(3-2x)^{9/2} (2x^2+x+1)^3} \right) + \\
 & \quad \frac{1}{28(3-2x)^{9/2} (2x^2+x+1)^4} \\
 & \quad \downarrow \text{1235} \\
 & \frac{1}{28} \left(\frac{1}{392} \int \frac{210(664-3049x)}{(3-2x)^{11/2} (2x^2+x+1)^2} dx + \frac{3049x+1387}{28(3-2x)^{9/2} (2x^2+x+1)^2} \right) + \frac{73x+23}{42(3-2x)^{9/2} (2x^2+x+1)^3} \\
 & \quad \frac{1}{28(3-2x)^{9/2} (2x^2+x+1)^4} \\
 & \quad \downarrow \text{27}
 \end{aligned}$$

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \int \frac{664 - 3049x}{(3 - 2x)^{11/2} (2x^2 + x + 1)^2} dx + \frac{3049x + 1387}{28(3 - 2x)^{9/2} (2x^2 + x + 1)^2} \right) + \frac{73x + 23}{42(3 - 2x)^{9/2} (2x^2 + x + 1)^3} \right) \frac{1}{28(3 - 2x)^{9/2} (2x^2 + x + 1)^4}$$

↓ 1235

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{196} \int -\frac{14(48147x + 22129)}{(3 - 2x)^{11/2} (2x^2 + x + 1)} dx + \frac{4377x + 3049}{14(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) + \frac{3049x + 1387}{28(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) \frac{1}{28(3 - 2x)^{9/2} (2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{4377x + 3049}{14(3 - 2x)^{9/2} (2x^2 + x + 1)} - \frac{1}{14} \int \frac{48147x + 22129}{(3 - 2x)^{11/2} (2x^2 + x + 1)} dx \right) + \frac{3049x + 1387}{28(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) \frac{1}{28(3 - 2x)^{9/2} (2x^2 + x + 1)^4}$$

↓ 1198

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(-\frac{1}{28} \int \frac{14(26957x + 5767)}{(3 - 2x)^{9/2} (2x^2 + x + 1)} dx - \frac{26957}{18(3 - 2x)^{9/2}} \right) + \frac{4377x + 3049}{14(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) + \frac{3049x + 1387}{28(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) \frac{1}{28(3 - 2x)^{9/2} (2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(-\frac{1}{2} \int \frac{26957x + 5767}{(3 - 2x)^{9/2} (2x^2 + x + 1)} dx - \frac{26957}{18(3 - 2x)^{9/2}} \right) + \frac{4377x + 3049}{14(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) + \frac{3049x + 1387}{28(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) \frac{1}{28(3 - 2x)^{9/2} (2x^2 + x + 1)^4}$$

↓ 1198

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(-\frac{1}{28} \int -\frac{2(3889 - 92405x)}{(3 - 2x)^{7/2} (2x^2 + x + 1)} dx - \frac{92405}{98(3 - 2x)^{7/2}} \right) - \frac{26957}{18(3 - 2x)^{9/2}} \right) + \frac{4377x + 3049}{14(3 - 2x)^{9/2} (2x^2 + x + 1)} \right) \right) \frac{1}{28(3 - 2x)^{9/2} (2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \int \frac{3889 - 92405x}{(3-2x)^{7/2}(2x^2+x+1)} dx - \frac{92405}{98(3-2x)^{7/2}} \right) - \frac{26957}{18(3-2x)^{9/2}} \right) + \frac{4377x + 28(3-2x)^{9/2}(2x^2+x+1)^4}{28(3-2x)^{9/2}(2x^2+x+1)^4} \right) \right) \right)$$

↓ 1198

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{28} \int \frac{14(15423 - 38491x)}{(3-2x)^{5/2}(2x^2+x+1)} dx - \frac{38491}{10(3-2x)^{5/2}} \right) - \frac{92405}{98(3-2x)^{7/2}} \right) - \frac{26957}{18(3-2x)^{9/2}} \right) \right) \right) \right)$$

↓ 27

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{2} \int \frac{15423 - 38491x}{(3-2x)^{5/2}(2x^2+x+1)} dx - \frac{38491}{10(3-2x)^{5/2}} \right) - \frac{92405}{98(3-2x)^{7/2}} \right) - \frac{26957}{18(3-2x)^{9/2}} \right) \right) \right) \right)$$

↓ 1198

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{28} \int \frac{2(100183 - 84627x)}{(3-2x)^{3/2}(2x^2+x+1)} dx - \frac{28209}{14(3-2x)^{3/2}} \right) - \frac{38491}{10(3-2x)^{5/2}} \right) - \frac{92405}{98(3-2x)^{7/2}} \right) \right) \right) \right)$$

↓ 27

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{14} \int \frac{100183 - 84627x}{(3-2x)^{3/2}(2x^2+x+1)} dx - \frac{28209}{14(3-2x)^{3/2}} \right) - \frac{38491}{10(3-2x)^{5/2}} \right) - \frac{92405}{98(3-2x)^{7/2}} \right) \right) \right) \right)$$

↓ 1198

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{28} \int \frac{14(69337 - 7645x)}{\sqrt{3-2x}(2x^2+x+1)} dx - \frac{7645}{2\sqrt{3-2x}} \right) - \frac{28209}{14(3-2x)^{3/2}} \right) - \frac{38491}{10(3-2x)^{5/2}} \right) \right) \right) \right)$$

↓ 27

↓ 1083

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{2} \left(-\frac{\frac{1}{2}(115739 - 7645\sqrt{14}) \int \frac{\sqrt{7+2\sqrt{14}}-2\sqrt{3-2x}}{-2x-\sqrt{7+2\sqrt{14}}\sqrt{3-2x}+\sqrt{14}+3} d\sqrt{3-2x} - \sqrt{7+2\sqrt{14}}}{2\sqrt{14}(7+2\sqrt{14})} \right. \frac{x}{28(3-2x)^{9/2}(2x^2+x+1)^4}$$

↓ 217

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{2} \left(-\frac{\frac{1}{2}(115739 - 7645\sqrt{14}) \int \frac{\sqrt{7+2\sqrt{14}}-2\sqrt{3-2x}}{-2x-\sqrt{7+2\sqrt{14}}\sqrt{3-2x}+\sqrt{14}+3} d\sqrt{3-2x} + \sqrt{\frac{7+2\sqrt{14}}{2\sqrt{14}}}}{2\sqrt{14}(7+2\sqrt{14})} \right. \frac{x}{28(3-2x)^{9/2}(2x^2+x+1)^4}$$

↓ 1103

$$\frac{1}{28} \left(\frac{1}{42} \left(\frac{15}{28} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{2} \left(\frac{1}{14} \left(\frac{1}{2} \left(-\frac{\sqrt{\frac{7+2\sqrt{14}}{2\sqrt{14}-7}}(115739 + 7645\sqrt{14}) \arctan\left(\frac{2\sqrt{3-2x}-\sqrt{7+2\sqrt{14}}}{\sqrt{2\sqrt{14}-7}}\right) - \frac{1}{2}(115739 - 7645\sqrt{14})}{2\sqrt{14}(7+2\sqrt{14})} \right. \frac{x}{28(3-2x)^{9/2}(2x^2+x+1)^4}$$

input

```
Int[1/((3 - 2*x)^(11/2)*(1 + x + 2*x^2)^5),x]
```

output

```
x/(28*(3 - 2*x)^(9/2)*(1 + x + 2*x^2)^4) + ((23 + 73*x)/(42*(3 - 2*x)^(9/2)*
(1 + x + 2*x^2)^3) + ((1387 + 3049*x)/(28*(3 - 2*x)^(9/2)*(1 + x + 2*x^2)^2) + (15*((3049 + 4377*x)/(14*(3 - 2*x)^(9/2)*(1 + x + 2*x^2)) + (-26957/(18*(3 - 2*x)^(9/2)) + (-92405/(98*(3 - 2*x)^(7/2)) + (-38491/(10*(3 - 2*x)^(5/2)) + (-28209/(14*(3 - 2*x)^(3/2)) + (-7645/(2*Sqrt[3 - 2*x])) + (-1/2*(Sqrt[(7 + 2*Sqrt[14])/(-7 + 2*Sqrt[14]])*(115739 + 7645*Sqrt[14])*ArcTan[(-Sqrt[7 + 2*Sqrt[14]] + 2*Sqrt[3 - 2*x])/Sqrt[-7 + 2*Sqrt[14]]] - ((115739 - 7645*Sqrt[14])*Log[3 + Sqrt[14] - Sqrt[7 + 2*Sqrt[14]]*Sqrt[3 - 2*x] - 2*x])/2)/Sqrt[14*(7 + 2*Sqrt[14]]) - (Sqrt[(7 + 2*Sqrt[14])/(-7 + 2*Sqrt[14]])*(115739 + 7645*Sqrt[14])*ArcTan[(Sqrt[7 + 2*Sqrt[14]] + 2*Sqrt[3 - 2*x])/Sqrt[-7 + 2*Sqrt[14]]] + ((115739 - 7645*Sqrt[14])*Log[3 + Sqrt[14] + Sqrt[7 + 2*Sqrt[14]]*Sqrt[3 - 2*x] - 2*x])/2)/(2*Sqrt[14*(7 + 2*Sqrt[14])])]/2)/14)/2)/14)/2)/14)/28)/42)/28
```


Definitions of rubi rules used

- rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`
- rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`
- rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`
- rule 1083 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]`
- rule 1103 `Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]`
- rule 1142 `Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c) Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]`
- rule 1165 `Int[((d_.) + (e_.)*(x_)^m)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^p, x_Symbol] := Simp[(d + e*x)^(m + 1)*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^m*Simp[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && LtQ[p, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]`

rule 1197

```
Int[((f_.) + (g_.)*(x_))/(Sqrt[(d_.) + (e_.)*(x_)]*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)), x_Symbol] := Simp[2 Subst[Int[(e*f - d*g + g*x^2)/(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4), x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x]
```

rule 1198

```
Int[((d_.) + (e_.)*(x_)^m)*((f_.) + (g_.)*(x_)))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(e*f - d*g)*((d + e*x)^(m + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Simp[1/(c*d^2 - b*d*e + a*e^2) Int[(d + e*x)^(m + 1)*(Simp[c*d*f - f*b*e + a*e*g - c*(e*f - d*g)*x, x]/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && FractionQ[m] && LtQ[m, -1]
```

rule 1235

```
Int[((d_.) + (e_.)*(x_)^m)*((f_.) + (g_.)*(x_)))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(f*(b*c*d - b^2*e + 2*a*c*e) - a*g*(2*c*d - b*e) + c*(f*(2*c*d - b*e) - g*(b*d - 2*a*e))*x)/((a + b*x + c*x^2)^(p + 1))/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^m*(a + b*x + c*x^2)^(p + 1)*Simp[f*(b*c*d*e*(2*p - m + 2) + b^2*e^2*(p + m + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3)) - g*(a*e*(b*e - 2*c*d*m + b*e*m) - b*d*(3*c*d - b*e + 2*c*d*p - b*e*p)) + c*e*(g*(b*d - 2*a*e) - f*(2*c*d - b*e))*(m + 2*p + 4)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && LtQ[p, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

rule 1483

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Simp[1/(2*c*q*r) Int[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Simp[1/(2*c*q*r) Int[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]
```

Maple [A] (verified)

Time = 2.54 (sec) , antiderivative size = 277, normalized size of antiderivative = 0.68

method	result
pseudoelliptic	$5 \left(\frac{\left(\frac{\sqrt{7+2\sqrt{14}} (146319\sqrt{14}-569986) (\ln(3-2x+\sqrt{14}-\sqrt{3-2x}\sqrt{7+2\sqrt{14}}) - \ln(3-2x+\sqrt{14}+\sqrt{3-2x}\sqrt{7+2\sqrt{14}})) \sqrt{-7+2\sqrt{14}}}{2} \right)}{\dots} \right) + \dots$
derivativedivides	$\frac{\frac{567651623\sqrt{3-2x}}{32} - \frac{6194606411(3-2x)^{\frac{3}{2}}}{192} + \frac{9801432515(3-2x)^{\frac{5}{2}}}{384} - \frac{8763772549(3-2x)^{\frac{7}{2}}}{768} + \frac{149630663(3-2x)^{\frac{9}{2}}}{48} - \frac{200063633(3-2x)}{384}}{6588344((3-2x)^2-7+14x)^4}$
default	$\frac{\frac{567651623\sqrt{3-2x}}{32} - \frac{6194606411(3-2x)^{\frac{3}{2}}}{192} + \frac{9801432515(3-2x)^{\frac{5}{2}}}{384} - \frac{8763772549(3-2x)^{\frac{7}{2}}}{768} + \frac{149630663(3-2x)^{\frac{9}{2}}}{48} - \frac{200063633(3-2x)}{384}}{6588344((3-2x)^2-7+14x)^4}$
trager	Expression too large to display
risch	$\frac{-88070400x^{12} - 677249280x^{11} + 1873554048x^{10} - 2443779648x^9 + 2343370048x^8 - 3106712560x^7 + 2888625656x^6 - 142168506368(2x^2+x+1)^4\sqrt{3-2x}}{(2x^2+x+1)^4\sqrt{3-2x}}$

```
input int(1/(3-2*x)^(11/2)/(2*x^2+x+1)^5,x,method=_RETURNVERBOSE)
```

```
output -5/26353376/(3-2*x)^(9/2)/(-7+2*14^(1/2))^(1/2)*((1/2*(7+2*14^(1/2))^(1/2)
*(146319*14^(1/2)-569986)*(ln(3-2*x+14^(1/2)-(3-2*x)^(1/2)*(7+2*14^(1/2))^(1/2))-ln(3-2*x+14^(1/2)+(3-2*x)^(1/2)*(7+2*14^(1/2))^(1/2)))*(-7+2*14^(1/2))^(1/2)+(115739*14^(1/2)+107030)*(arctan((2*(3-2*x)^(1/2)-(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2))+arctan((2*(3-2*x)^(1/2)+(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2))))*(x-3/2)^4*(x^2+1/2*x+1/2)^4*(3-2*x)^(1/2)+214060*(-73537943/4403520*x^5-24405799/2001600*x^3+1627773523/88070400*x^4+361078207/11008800*x^6+67601077/44035200*x^2+36615157/1376100*x^8-53726593/11008800*x+x^12-38833907/1100880*x^7+1626349/76450*x^10+40289347/88070400-4242673/152900*x^9-58789/7645*x^11)*(-7+2*14^(1/2))^(1/2))/(2*x^2+x+1)^4
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 636 vs. $2(298) = 596$.

Time = 0.09 (sec) , antiderivative size = 636, normalized size of antiderivative = 1.56

$$\int \frac{1}{(3-2x)^{11/2} (1+x+2x^2)^5} dx = \text{Too large to display}$$

input `integrate(1/(3-2*x)^(11/2)/(2*x^2+x+1)^5,x, algorithm="fricas")`

output

```
-1/60718178304*(90*sqrt(1/2)*(512*x^13 - 2816*x^12 + 5632*x^11 - 5888*x^10
+ 6848*x^9 - 8992*x^8 + 6112*x^7 - 4240*x^6 + 4994*x^5 - 1707*x^4 + 936*x
^3 - 1242*x^2 - 162*x - 243)*sqrt(40815066112*sqrt(14) + 149046503977)*arc
tan(2/88040902397*sqrt(1/2)*(sqrt(1/2)*sqrt(40815066112*sqrt(14) - 1490465
03977)*(2*sqrt(14) + 7) + sqrt(-2*x + 3)*(115739*sqrt(14) - 107030))*sqrt(
40815066112*sqrt(14) + 149046503977)) - 90*sqrt(1/2)*(512*x^13 - 2816*x^12
+ 5632*x^11 - 5888*x^10 + 6848*x^9 - 8992*x^8 + 6112*x^7 - 4240*x^6 + 499
4*x^5 - 1707*x^4 + 936*x^3 - 1242*x^2 - 162*x - 243)*sqrt(40815066112*sqrt
(14) + 149046503977)*arctan(2/88040902397*sqrt(1/2)*(sqrt(1/2)*sqrt(408150
66112*sqrt(14) - 149046503977)*(2*sqrt(14) + 7) - sqrt(-2*x + 3)*(115739*s
qrt(14) - 107030))*sqrt(40815066112*sqrt(14) + 149046503977)) + 45*sqrt(1/
2)*(512*x^13 - 2816*x^12 + 5632*x^11 - 5888*x^10 + 6848*x^9 - 8992*x^8 + 6
112*x^7 - 4240*x^6 + 4994*x^5 - 1707*x^4 + 936*x^3 - 1242*x^2 - 162*x - 24
3)*sqrt(40815066112*sqrt(14) - 149046503977)*log(5*sqrt(1/2)*sqrt(-2*x + 3
))*sqrt(40815066112*sqrt(14) - 149046503977)*(146319*sqrt(14) + 569986) - 1
25772717710*x + 62886358855*sqrt(14) + 188659076565) - 45*sqrt(1/2)*(512*x
^13 - 2816*x^12 + 5632*x^11 - 5888*x^10 + 6848*x^9 - 8992*x^8 + 6112*x^7 -
4240*x^6 + 4994*x^5 - 1707*x^4 + 936*x^3 - 1242*x^2 - 162*x - 243)*sqrt(4
0815066112*sqrt(14) - 149046503977)*log(-5*sqrt(1/2)*sqrt(-2*x + 3))*sqrt(4
0815066112*sqrt(14) - 149046503977)*(146319*sqrt(14) + 569986) - 125772...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(3-2x)^{11/2} (1+x+2x^2)^5} dx = \text{Timed out}$$

input `integrate(1/(3-2*x)**(11/2)/(2*x**2+x+1)**5,x)`output `Timed out`**Maxima [F]**

$$\int \frac{1}{(3-2x)^{11/2} (1+x+2x^2)^5} dx = \int \frac{1}{(2x^2+x+1)^5 (-2x+3)^{\frac{11}{2}}} dx$$

input `integrate(1/(3-2*x)^(11/2)/(2*x^2+x+1)^5,x, algorithm="maxima")`output `integrate(1/((2*x^2 + x + 1)^5*(-2*x + 3)^(11/2)), x)`**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 795 vs. 2(298) = 596.

Time = 0.45 (sec) , antiderivative size = 795, normalized size of antiderivative = 1.95

$$\int \frac{1}{(3-2x)^{11/2} (1+x+2x^2)^5} dx = \text{Too large to display}$$

input `integrate(1/(3-2*x)^(11/2)/(2*x^2+x+1)^5,x, algorithm="giac")`

output

```

-5/1511207993344*sqrt(7)*(22935*14^(3/4)*sqrt(7)*(sqrt(14) + 4)*sqrt(-2*sqrt(14) + 8) + 7645*14^(3/4)*sqrt(7)*(sqrt(14) - 4)*sqrt(-2*sqrt(14) + 8) + 53515*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) + 160545*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) + 925912*14^(1/4)*sqrt(7)*sqrt(-2*sqrt(14) + 8) + 6481384*14^(1/4)*sqrt(2*sqrt(14) + 8))*arctan(1/28*14^(3/4)*(14^(1/4)*sqrt(1/2)*sqrt(sqrt(14) + 4) + 2*sqrt(-2*x + 3))/sqrt(-1/8*sqrt(14) + 1/2)) - 5/1511207993344*sqrt(7)*(22935*14^(3/4)*sqrt(7)*(sqrt(14) + 4)*sqrt(-2*sqrt(14) + 8) + 7645*14^(3/4)*sqrt(7)*(sqrt(14) - 4)*sqrt(-2*sqrt(14) + 8) + 53515*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) + 160545*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) + 925912*14^(1/4)*sqrt(7)*sqrt(-2*sqrt(14) + 8) + 6481384*14^(1/4)*sqrt(2*sqrt(14) + 8))*arctan(-1/28*14^(3/4)*(14^(1/4)*sqrt(1/2)*sqrt(sqrt(14) + 4) - 2*sqrt(-2*x + 3))/sqrt(-1/8*sqrt(14) + 1/2)) - 5/3022415986688*sqrt(7)*(7645*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) + 22935*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) - 160545*14^(3/4)*(sqrt(14) + 4)*sqrt(-2*sqrt(14) + 8) - 53515*14^(3/4)*(sqrt(14) - 4)*sqrt(-2*sqrt(14) + 8) + 925912*14^(1/4)*sqrt(7)*sqrt(2*sqrt(14) + 8) - 6481384*14^(1/4)*sqrt(-2*sqrt(14) + 8))*log(14^(1/4)*sqrt(1/2)*sqrt(-2*x + 3)*sqrt(sqrt(14) + 4) - 2*x + sqrt(14) + 3) + 5/3022415986688*sqrt(7)*(7645*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) + 22935*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) - 1605...

```

Mupad [B] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 343, normalized size of antiderivative = 0.84

$$\int \frac{1}{(3 - 2x)^{11/2} (1 + x + 2x^2)^5} dx = \text{Too large to display}$$

input

```
int(1/((3 - 2*x)^(11/2)*(x + 2*x^2 + 1)^5),x)
```

output

```
(atan(((3 - 2*x)^(1/2)*(7^(1/2)*12577271771i - 149046503977)^(1/2)*1572158
971375i)/(391663056253676053933850624*((7^(1/2)*181960107187971125i)/19583
1528126838026966925312 - 230036728532618625/27975932589548289566703616)) -
(1572158971375*7^(1/2)*(3 - 2*x)^(1/2)*(7^(1/2)*12577271771i - 1490465039
77)^(1/2))/(391663056253676053933850624*((7^(1/2)*181960107187971125i)/195
831528126838026966925312 - 230036728532618625/27975932589548289566703616))
)*(7^(1/2)*12577271771i - 149046503977)^(1/2)*5i)/3373232128 - ((272*x)/44
1 - (164*(2*x - 3)^2)/441 + (1966*(2*x - 3)^3)/3087 - (9091*(2*x - 3)^4)/3
087 - (32070727*(2*x - 3)^5)/5531904 - (41014777*(2*x - 3)^6)/11063808 - (
141921511*(2*x - 3)^7)/154893312 + (23262655*(2*x - 3)^8)/309786624 + (157
1659*(2*x - 3)^9)/15059072 + (468427*(2*x - 3)^10)/17210368 + (394105*(2*x
- 3)^11)/120472576 + (38225*(2*x - 3)^12)/240945152 - 520/441)/(38416*(3
- 2*x)^(9/2) - 76832*(3 - 2*x)^(11/2) + 68600*(3 - 2*x)^(13/2) - 35672*(3
- 2*x)^(15/2) + 11809*(3 - 2*x)^(17/2) - 2548*(3 - 2*x)^(19/2) + 350*(3 -
2*x)^(21/2) - 28*(3 - 2*x)^(23/2) + (3 - 2*x)^(25/2)) - atan(((3 - 2*x)^(
1/2)*(- 7^(1/2)*12577271771i - 149046503977)^(1/2)*1572158971375i)/(391663
056253676053933850624*((7^(1/2)*181960107187971125i)/195831528126838026966
925312 + 230036728532618625/27975932589548289566703616)) + (1572158971375*
7^(1/2)*(3 - 2*x)^(1/2)*(- 7^(1/2)*12577271771i - 149046503977)^(1/2))/(39
1663056253676053933850624*((7^(1/2)*181960107187971125i)/19583152812683...
```

Reduce [B] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 4694, normalized size of antiderivative = 11.53

$$\int \frac{1}{(3-2x)^{11/2} (1+x+2x^2)^5} dx = \text{Too large to display}$$

input

```
int(1/(3-2*x)^(11/2)/(2*x^2+x+1)^5,x)
```

output

```
( - 3371189760*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt
(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**12 + 1348475
9040*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x +
3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**11 - 16855948800*sqrt(
2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(
2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**10 + 13484759040*sqrt(2*sqrt(14)
- 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14)
+ 7))/sqrt(2*sqrt(14) - 7))*x**9 - 24862524480*sqrt(2*sqrt(14) - 7)*sqrt(
- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt
(2*sqrt(14) - 7))*x**8 + 21912733440*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)
*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14)
- 7))*x**7 - 7374477600*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*at
an((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**6
+ 16855948800*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(
- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**5 - 759834567
0*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3)
- sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**4 - 158024520*sqrt(2*sqrt
(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt
(14) + 7))/sqrt(2*sqrt(14) - 7))*x**3 - 6399993060*sqrt(2*sqrt(14) - 7)*sq
rt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7)...
```


$$3.48 \quad \int \frac{1}{(3-2x)^{21/2}(1+x+2x^2)^{10}} dx$$

Optimal result	376
Mathematica [C] (verified)	377
Rubi [F]	378
Maple [A] (verified)	385
Fricas [B] (verification not implemented)	386
Sympy [F(-1)]	387
Maxima [F]	388
Giac [B] (verification not implemented)	388
Mupad [B] (verification not implemented)	389
Reduce [B] (verification not implemented)	390

Optimal result

Integrand size = 20, antiderivative size = 648

$$\int \frac{1}{(3-2x)^{21/2}(1+x+2x^2)^{10}} dx = \text{Too large to display}$$

output

```

4718120139975/351733660450816/(3-2*x)^(19/2)-815900548375/629418129227776/
(3-2*x)^(17/2)-3029508823715/1555033025150976/(3-2*x)^(15/2)-1351574302182
5/13476952884641792/(3-2*x)^(13/2)-5846828446875/14513641568075776/(3-2*x)
^(11/2)-37283626871975/261245548225363968/(3-2*x)^(9/2)-132355162272575/28
44673747342852096/(3-2*x)^(7/2)-11557581705725/812763927812243456/(3-2*x)
^(5/2)-46601678385075/11378694989371408384/(3-2*x)^(3/2)+1/63*x/(3-2*x)^(19
/2)/(2*x^2+x+1)^9+1/7056*(53+173*x)/(3-2*x)^(19/2)/(2*x^2+x+1)^8+1/691488*
(8477+21409*x)/(3-2*x)^(19/2)/(2*x^2+x+1)^7+5/6453888*(21409+47471*x)/(3-2
*x)^(19/2)/(2*x^2+x+1)^6+41/90354432*(47471+92875*x)/(3-2*x)^(19/2)/(2*x^2
+x+1)^5+41/5059848192*(3436375+5677637*x)/(3-2*x)^(19/2)/(2*x^2+x+1)^4+451
/10119696384*(811091+998691*x)/(3-2*x)^(19/2)/(2*x^2+x+1)^3+451/2833514987
52*(28962039+14627273*x)/(3-2*x)^(19/2)/(2*x^2+x+1)^2+11275/3966920982528*
(14627273-35058731*x)/(3-2*x)^(19/2)/(2*x^2+x+1)-24229218097975/2275738997
8742816768/(3-2*x)^(1/2)+11275/1274413838809597739008*ln(3-2*x+14^(1/2)-(3
-2*x)^(1/2)*(7+2*14^(1/2))^(1/2))*(9756589235-2148932869*14^(1/2))*(-14+4*
14^(1/2))^(1/2)-11275/1274413838809597739008*ln(3-2*x+14^(1/2)+(3-2*x)^(1/
2)*(7+2*14^(1/2))^(1/2))*(9756589235-2148932869*14^(1/2))*(-14+4*14^(1/2))
^(1/2)+11275/637206919404798869504*arctan((-2*(3-2*x)^(1/2)+(7+2*14^(1/2))
^(1/2))/(-7+2*14^(1/2))^(1/2))*(9756589235+2148932869*14^(1/2))*(14+4*14^(
1/2))^(1/2)-11275/637206919404798869504*arctan((2*(3-2*x)^(1/2)+(7+2*14...

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.73 (sec) , antiderivative size = 253, normalized size of antiderivative = 0.39

$$\int \frac{1}{(3-2x)^{21/2}(1+x+2x^2)^{10}} dx = \frac{14(-4884417100172357749737+205702452014540322797289x+111926768697602999806116x^2}{\dots}$$

input

```
Integrate[1/((3 - 2*x)^(21/2)*(1 + x + 2*x^2)^10), x]
```

output

```

((14*(-4884417100172357749737 + 205702452014540322797289*x + 1119267686976
02999806116*x^2 + 1362587089603925431664856*x^3 - 809990362095044210054958
*x^4 + 3673303058277822225386926*x^5 - 8685973988079840377705700*x^6 + 107
18131725916893151555068*x^7 - 27246604251076689552043953*x^8 + 41613884937
255303086792337*x^9 - 59791102681494117572149176*x^10 + 102031573634317834
547976132*x^11 - 133312541377246386115890240*x^12 + 1726496922946149692741
68896*x^13 - 229408132984166521977166336*x^14 + 25881925681516324984544793
6*x^15 - 282644664539994827031006720*x^16 + 304010591010966811155955200*x^
17 - 287279159180291305208156160*x^18 + 253788172995391086570485760*x^19 -
216634228326470609547509760*x^20 + 162290307223249502039654400*x^21 - 106
701725825102321939251200*x^22 + 65360120291258796757811200*x^23 - 33969890
064381284111155200*x^24 + 12365045055896811105484800*x^25 - 26219489415962
37063782400*x^26 + 240031204937714427494400*x^27))/((3 - 2*x)^(19/2)*(1 +
x + 2*x^2)^9) - 426093525*Sqrt[2293002953699236822393 + (30540258843957888
971*I)*Sqrt[7]]*ArcTan[(Sqrt[-1 - I/Sqrt[7]]*Sqrt[3 - 2*x])/2] - 426093525
*Sqrt[2293002953699236822393 - (30540258843957888971*I)*Sqrt[7]]*ArcTan[(S
qrt[-1 + I/Sqrt[7]]*Sqrt[3 - 2*x])/2])/12040343345613377038712832

```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3-2x)^{21/2} (2x^2+x+1)^{10}} dx \\
 & \quad \downarrow \text{1165} \\
 & \int \frac{28(60-53x)}{(3-2x)^{21/2} (2x^2+x+1)^9} dx + \frac{x}{63(3-2x)^{19/2} (2x^2+x+1)^9} \\
 & \quad \downarrow \text{27} \\
 & \frac{1}{63} \int \frac{60-53x}{(3-2x)^{21/2} (2x^2+x+1)^9} dx + \frac{x}{63(3-2x)^{19/2} (2x^2+x+1)^9} \\
 & \quad \downarrow \text{1235}
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{63} \left(\frac{\int \frac{14(6466-8477x)}{(3-2x)^{21/2}(2x^2+x+1)^8} dx}{1568} + \frac{173x+53}{112(3-2x)^{19/2}(2x^2+x+1)^8} \right) + \\
& \qquad \qquad \qquad \frac{x}{63(3-2x)^{19/2}(2x^2+x+1)^9} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{1}{63} \left(\frac{1}{112} \int \frac{6466-8477x}{(3-2x)^{21/2}(2x^2+x+1)^8} dx + \frac{173x+53}{112(3-2x)^{19/2}(2x^2+x+1)^8} \right) + \\
& \qquad \qquad \qquad \frac{x}{63(3-2x)^{19/2}(2x^2+x+1)^9} \\
& \qquad \qquad \qquad \downarrow 1235 \\
& \frac{1}{63} \left(\frac{1}{112} \left(\frac{\int \frac{630(13031-21409x)}{(3-2x)^{21/2}(2x^2+x+1)^7} dx}{1372} + \frac{21409x+8477}{98(3-2x)^{19/2}(2x^2+x+1)^7} \right) + \frac{173x+53}{112(3-2x)^{19/2}(2x^2+x+1)^8} \right) + \\
& \qquad \qquad \qquad \frac{x}{63(3-2x)^{19/2}(2x^2+x+1)^9} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \int \frac{13031-21409x}{(3-2x)^{21/2}(2x^2+x+1)^7} dx + \frac{21409x+8477}{98(3-2x)^{19/2}(2x^2+x+1)^7} \right) + \frac{173x+53}{112(3-2x)^{19/2}(2x^2+x+1)^8} \right) + \\
& \qquad \qquad \qquad \frac{x}{63(3-2x)^{19/2}(2x^2+x+1)^9} \\
& \qquad \qquad \qquad \downarrow 1235 \\
& \frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{\int \frac{574(22702-47471x)}{(3-2x)^{21/2}(2x^2+x+1)^6} dx}{1176} + \frac{47471x+21409}{84(3-2x)^{19/2}(2x^2+x+1)^6} \right) + \frac{21409x+8477}{98(3-2x)^{19/2}(2x^2+x+1)^7} \right) + \right) + \\
& \qquad \qquad \qquad \frac{x}{63(3-2x)^{19/2}(2x^2+x+1)^9} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \int \frac{22702-47471x}{(3-2x)^{21/2}(2x^2+x+1)^6} dx + \frac{47471x+21409}{84(3-2x)^{19/2}(2x^2+x+1)^6} \right) + \frac{21409x+8477}{98(3-2x)^{19/2}(2x^2+x+1)^7} \right) + \right) + \\
& \qquad \qquad \qquad \frac{x}{63(3-2x)^{19/2}(2x^2+x+1)^9} \\
& \qquad \qquad \qquad \downarrow 1235
\end{aligned}$$

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{980} \int \frac{14(1120631 - 3436375x)}{x(3-2x)^{21/2}(2x^2+x+1)^5} dx + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5} \right) + \frac{47471x+2}{84(3-2x)^{19/2}(2x^2+x+1)^5} \right) \right) \right) + \frac{47471x+2}{84(3-2x)^{19/2}(2x^2+x+1)^5}$$

$$\frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 27

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \int \frac{1120631 - 3436375x}{x(3-2x)^{21/2}(2x^2+x+1)^5} dx + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5} \right) + \frac{47471x+2}{84(3-2x)^{19/2}(2x^2+x+1)^5} \right) \right) \right) + \frac{47471x+2}{84(3-2x)^{19/2}(2x^2+x+1)^5}$$

$$\frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 1235

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{1}{784} \int \frac{3234(93800 - 811091x)}{x(3-2x)^{21/2}(2x^2+x+1)^4} dx + \frac{5677637x+3436375}{56(3-2x)^{19/2}(2x^2+x+1)^4} \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5} \right) \right) \right) \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5}$$

$$\frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 27

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \int \frac{93800 - 811091x}{x(3-2x)^{21/2}(2x^2+x+1)^4} dx + \frac{5677637x+3436375}{56(3-2x)^{19/2}(2x^2+x+1)^4} \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5} \right) \right) \right) \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5}$$

$$\frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 1235

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{1}{588} \int -\frac{14(28962039x+7167383)}{x(3-2x)^{21/2}(2x^2+x+1)^3} dx + \frac{998691x+811091}{42(3-2x)^{19/2}(2x^2+x+1)^3} \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5} \right) \right) \right) \right) \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5}$$

$$\frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 27

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{998691x+811091}{42(3-2x)^{19/2}(2x^2+x+1)^3} - \frac{1}{42} \int \frac{28962039x+7167383}{x(3-2x)^{21/2}(2x^2+x+1)^3} dx \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5} \right) \right) \right) \right) \right) + \frac{92875x+47471}{70(3-2x)^{19/2}(2x^2+x+1)^5}$$

$$\frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 1235

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{1}{42} \left(\frac{14627273x + 28962039}{28(3-2x)^{19/2}(2x^2+x+1)^2} - \frac{1}{392} \int \frac{350(14627273x + 24843002)}{(3-2x)^{21/2}(2x^2+x+1)^2} dx \right) + \frac{1}{42} \right) \right) \right) \right) \right) \right) \frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 27

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{1}{42} \left(\frac{14627273x + 28962039}{28(3-2x)^{19/2}(2x^2+x+1)^2} - \frac{25}{28} \int \frac{14627273x + 24843002}{(3-2x)^{21/2}(2x^2+x+1)^2} dx \right) + \frac{1}{42} \right) \right) \right) \right) \right) \right) \frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 1235

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{1}{42} \left(\frac{14627273x + 28962039}{28(3-2x)^{19/2}(2x^2+x+1)^2} - \frac{25}{28} \left(\frac{1}{196} \int \frac{42(158887401 - 245411117x)}{(3-2x)^{21/2}(2x^2+x+1)} dx - \frac{1}{42} \right) \right) \right) \right) \right) \right) \right) \frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 27

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{1}{42} \left(\frac{14627273x + 28962039}{28(3-2x)^{19/2}(2x^2+x+1)^2} - \frac{25}{28} \left(\frac{3}{14} \int \frac{158887401 - 245411117x}{(3-2x)^{21/2}(2x^2+x+1)} dx - \frac{1}{14} \right) \right) \right) \right) \right) \right) \right) \frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 1198

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{1}{42} \left(\frac{14627273x + 28962039}{28(3-2x)^{19/2}(2x^2+x+1)^2} - \frac{25}{28} \left(\frac{3}{14} \left(\frac{1}{28} \int \frac{2(880960721 - 418458549x)}{(3-2x)^{19/2}(2x^2+x+1)} dx - \frac{1}{28} \right) \right) \right) \right) \right) \right) \right) \right) \frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 27

$$\frac{1}{63} \left(\frac{1}{112} \left(\frac{45}{98} \left(\frac{41}{84} \left(\frac{1}{70} \left(\frac{33}{8} \left(\frac{1}{42} \left(\frac{14627273x + 28962039}{28(3-2x)^{19/2}(2x^2+x+1)^2} - \frac{25}{28} \left(\frac{3}{14} \left(\frac{1}{14} \int \frac{880960721 - 418458549x}{(3-2x)^{19/2}(2x^2+x+1)} dx - \frac{1}{14} \right) \right) \right) \right) \right) \right) \right) \right) \frac{1}{63(3-2x)^{19/2}(2x^2+x+1)^9}$$

↓ 1198

Definitions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 1165 `Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^m*Simp[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && LtQ[p, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]`

rule 1198 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[(e*f - d*g)*((d + e*x)^(m + 1)/((m + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Simp[1/(c*d^2 - b*d*e + a*e^2) Int[(d + e*x)^(m + 1)*(Simp[c*d*f - f*b*e + a*e*g - c*(e*f - d*g)*x, x]/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && FractionQ[m] && LtQ[m, -1]`

rule 1235 `Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(f*(b*c*d - b^2*e + 2*a*c*e) - a*g*(2*c*d - b*e) + c*(f*(2*c*d - b*e) - g*(b*d - 2*a*e))*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x] + Simp[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^m*(a + b*x + c*x^2)^(p + 1)*Simp[f*(b*c*d*e*(2*p - m + 2) + b^2*e^2*(p + m + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3)) - g*(a*e*(b*e - 2*c*d*m + b*e*m) - b*d*(3*c*d - b*e + 2*c*d*p - b*e*p)) + c*e*(g*(b*d - 2*a*e) - f*(2*c*d - b*e))*(m + 2*p + 4)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && LtQ[p, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])`

Maple [A] (verified)

Time = 3.14 (sec) , antiderivative size = 352, normalized size of antiderivative = 0.54

method	result
pseudoelliptic	$\frac{11275 \left(x - \frac{3}{2}\right)^9 \left(x^2 + \frac{1}{2}x + \frac{1}{2}\right)^9 \left(\frac{\sqrt{7+2\sqrt{14}} \left(18352320711\sqrt{14} - 69111417106\right) \left(\ln\left(3-2x+\sqrt{14}-\sqrt{3-2x}\sqrt{7+2\sqrt{14}}\right) - \ln\left(3-2x+\sqrt{14}+\sqrt{3-2x}\sqrt{7+2\sqrt{14}}\right)\right)}{2}\right)}{245}$
derivativedivides	$\frac{1}{5367029731(3-2x)^{\frac{19}{2}}} + \frac{5}{4802079233(3-2x)^{\frac{17}{2}}} + \frac{73}{23727920916(3-2x)^{\frac{15}{2}}} + \frac{165}{25705247659(3-2x)^{\frac{13}{2}}} + \frac{1}{2214605(3-2x)^{\frac{11}{2}}}$
default	$\frac{1}{5367029731(3-2x)^{\frac{19}{2}}} + \frac{5}{4802079233(3-2x)^{\frac{17}{2}}} + \frac{73}{23727920916(3-2x)^{\frac{15}{2}}} + \frac{165}{25705247659(3-2x)^{\frac{13}{2}}} + \frac{1}{2214605(3-2x)^{\frac{11}{2}}}$
trager	Expression too large to display
risch	$-\frac{240031204937714427494400x^{27} - 2621948941596237063782400x^{26} + 12365045055896811105484800x^{25} - 3396989000000000000000000x^{24} + \dots}{\dots}$

input

```
int(1/(3-2*x)^(21/2)/(2*x^2+x+1)^10,x,method=_RETURNVERBOSE)
```

output

```

11275/2430751493090816/(-7+2*14^(1/2))^(1/2)*((x-3/2)^9*(x^2+1/2*x+1/2)^9*
(1/2*(7+2*14^(1/2))^(1/2)*(18352320711*14^(1/2)-69111417106)*(ln(3-2*x+14^
(1/2)-(3-2*x)^(1/2)*(7+2*14^(1/2))^(1/2))-ln(3-2*x+14^(1/2)+(3-2*x)^(1/2)*
(7+2*14^(1/2))^(1/2)))*(-7+2*14^(1/2))^(1/2)+(9756589235*14^(1/2)+30085060
166)*(arctan((2*(3-2*x)^(1/2)-(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2))
+arctan((2*(3-2*x)^(1/2)+(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2))))*(3
-2*x)^(1/2)+60170120332*(-7+2*14^(1/2))^(1/2)*(22855828001615591421921/266
70133881968269721600*x+193096157908388472533/152458844599666176*x^17-46547
1892878599/515743888560*x^20+271924352600651293/257184285761920*x^19+33243
27068969447/4916758404272*x^21-63047885074067/141829569354*x^22+2106543768
2057/77361583284*x^23-1216492052933/8595731476*x^24-38014370445393391293/3
1762259291597120*x^18-1129001874807303405453/2032784594662215680*x^12-1092
8359993529274103333/11434413344974963200*x^14-299208867441559564523/254098
074332776960*x^16-20254438577741909746663/81311383786488627200*x^10+112774
755927521146576673/650491070291909017600*x^9+12198896895542543585363/28698
135454054809600*x^11+21932125545555763373243/30491768919933235200*x^13-221
517107732330809366211/1951473210875727052800*x^8+297725881275469254209863/
6667533470492067430400*x^7-9651082208977600419673/266701338819682697216*x^
6+442803288917/8595731476*x^25+144203782903185201071/133735828596198400*x^
15-134998393682507368342493/40005200822952404582400*x^4+847930065890931...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1086 vs. $2(491) = 982$.

Time = 0.12 (sec) , antiderivative size = 1086, normalized size of antiderivative = 1.68

$$\int \frac{1}{(3-2x)^{21/2}(1+x+2x^2)^{10}} dx = \text{Too large to display}$$

input

```
integrate(1/(3-2*x)^(21/2)/(2*x^2+x+1)^10,x, algorithm="fricas")
```

output

```

-1/24080686691226754077425664*(852187050*sqrt(1/2)*(524288*x^28 - 5505024*
x^27 + 24772608*x^26 - 64684032*x^25 + 119734272*x^24 - 194052096*x^23 + 2
95206912*x^22 - 386777088*x^21 + 449261568*x^20 - 515594240*x^19 + 5405030
40*x^18 - 496581120*x^17 + 467712000*x^16 - 411828480*x^15 + 303534720*x^1
4 - 248434368*x^13 + 186495624*x^12 - 105219828*x^11 + 83621482*x^10 - 497
93667*x^9 + 19105065*x^8 - 20036484*x^7 + 5497632*x^6 - 2235114*x^5 + 3276
126*x^4 + 734832*x^3 + 826686*x^2 + 137781*x + 59049)*sqrt(613211190464078
675968*sqrt(14) + 2293002953699236822393)*arctan(2/213781811907705222797*s
qrt(1/2)*(sqrt(1/2)*sqrt(613211190464078675968*sqrt(14) - 2293002953699236
822393)*(2*sqrt(14) + 7) + sqrt(-2*x + 3)*(9756589235*sqrt(14) - 300850601
66))*sqrt(613211190464078675968*sqrt(14) + 2293002953699236822393)) - 8521
87050*sqrt(1/2)*(524288*x^28 - 5505024*x^27 + 24772608*x^26 - 64684032*x^2
5 + 119734272*x^24 - 194052096*x^23 + 295206912*x^22 - 386777088*x^21 + 44
9261568*x^20 - 515594240*x^19 + 540503040*x^18 - 496581120*x^17 + 46771200
0*x^16 - 411828480*x^15 + 303534720*x^14 - 248434368*x^13 + 186495624*x^12
- 105219828*x^11 + 83621482*x^10 - 49793667*x^9 + 19105065*x^8 - 20036484
*x^7 + 5497632*x^6 - 2235114*x^5 + 3276126*x^4 + 734832*x^3 + 826686*x^2 +
137781*x + 59049)*sqrt(613211190464078675968*sqrt(14) + 22930029536992368
22393)*arctan(2/213781811907705222797*sqrt(1/2)*(sqrt(1/2)*sqrt(6132111904
64078675968*sqrt(14) - 2293002953699236822393)*(2*sqrt(14) + 7) - sqrt(...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(3-2x)^{21/2} (1+x+2x^2)^{10}} dx = \text{Timed out}$$

input

```
integrate(1/(3-2*x)**(21/2)/(2*x**2+x+1)**10,x)
```

output

Timed out

Maxima [F]

$$\int \frac{1}{(3-2x)^{21/2} (1+x+2x^2)^{10}} dx = \int \frac{1}{(2x^2+x+1)^{10} (-2x+3)^{\frac{21}{2}}} dx$$

input `integrate(1/(3-2*x)^(21/2)/(2*x^2+x+1)^10,x, algorithm="maxima")`

output `integrate(1/((2*x^2 + x + 1)^10*(-2*x + 3)^(21/2)), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1000 vs. $2(491) = 982$.

Time = 0.60 (sec) , antiderivative size = 1000, normalized size of antiderivative = 1.54

$$\int \frac{1}{(3-2x)^{21/2} (1+x+2x^2)^{10}} dx = \text{Too large to display}$$

input `integrate(1/(3-2*x)^(21/2)/(2*x^2+x+1)^10,x, algorithm="giac")`

output

```
-11275/142734349946674946768896*sqrt(7)*(6446798607*14^(3/4)*sqrt(7)*(sqrt(14) + 4)*sqrt(-2*sqrt(14) + 8) + 2148932869*14^(3/4)*sqrt(7)*(sqrt(14) - 4)*sqrt(-2*sqrt(14) + 8) - 15042530083*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) - 45127590249*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) + 78052713880*14^(1/4)*sqrt(7)*sqrt(-2*sqrt(14) + 8) - 546368997160*14^(1/4)*sqrt(2*sqrt(14) + 8))*arctan(1/28*14^(3/4)*(14^(1/4)*sqrt(1/2)*sqrt(sqrt(14) + 4) + 2*sqrt(-2*x + 3))/sqrt(-1/8*sqrt(14) + 1/2)) - 11275/142734349946674946768896*sqrt(7)*(6446798607*14^(3/4)*sqrt(7)*(sqrt(14) + 4)*sqrt(-2*sqrt(14) + 8) + 2148932869*14^(3/4)*sqrt(7)*(sqrt(14) - 4)*sqrt(-2*sqrt(14) + 8) - 15042530083*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) - 45127590249*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) + 78052713880*14^(1/4)*sqrt(7)*sqrt(-2*sqrt(14) + 8) - 546368997160*14^(1/4)*sqrt(2*sqrt(14) + 8))*arctan(-1/28*14^(3/4)*(14^(1/4)*sqrt(1/2)*sqrt(sqrt(14) + 4) - 2*sqrt(-2*x + 3))/sqrt(-1/8*sqrt(14) + 1/2)) - 11275/285468699893349893537792*sqrt(7)*(2148932869*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) + 6446798607*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) + 45127590249*14^(3/4)*(sqrt(14) + 4)*sqrt(-2*sqrt(14) + 8) + 15042530083*14^(3/4)*(sqrt(14) - 4)*sqrt(-2*sqrt(14) + 8) + 78052713880*14^(1/4)*sqrt(7)*sqrt(2*sqrt(14) + 8) + 546368997160*14^(1/4)*sqrt(-2*sqrt(14) + 8))*log(14^(1/4)*sqrt(1/2)*sqrt(-2*x + 3)*sqrt(sqrt(14) + 4) - 2*x + sqrt(14) + 3) + 11275/285...
```

Mupad [B] (verification not implemented)

Time = 0.32 (sec) , antiderivative size = 567, normalized size of antiderivative = 0.88

$$\int \frac{1}{(3 - 2x)^{21/2} (1 + x + 2x^2)^{10}} dx = \text{Too large to display}$$

input

```
int(1/((3 - 2*x)^(21/2)*(x + 2*x^2 + 1)^10),x)
```

output

```

((184192*(2*x - 3)^2)/47481 - (18944*x)/2261 - (15552*(2*x - 3)^3)/4199 +
(5666272*(2*x - 3)^4)/1440257 - (63490768*(2*x - 3)^5)/12962313 + (5334956
72*(2*x - 3)^6)/70572593 - (1111521492*(2*x - 3)^7)/70572593 + (7800732315
8*(2*x - 3)^8)/1482024453 - (250239440467*(2*x - 3)^9)/494008151 + (111869
3654785651073*(2*x - 3)^10)/453254454575104 + (1624300450152249301*(2*x -
3)^11)/97125954551808 + (35048653520674948897*(2*x - 3)^12)/90650890915020
8 + (95527511967437577915*(2*x - 3)^13)/1813017818300416 + (56406629997314
15610547*(2*x - 3)^14)/114220122552926208 + (1737142288764447500149*(2*x -
3)^15)/50764498912411648 + (12971210667229097601055*(2*x - 3)^16)/7107029
84773763072 + (32723441206946795665235*(2*x - 3)^17)/4264217908642578432 +
(102645797034777710681325*(2*x - 3)^18)/39799367147330732032 + (146093178
7430200665315*(2*x - 3)^19)/2094703534070038528 + (687618468821894139745*(
2*x - 3)^20)/4528256169239642112 + (39968995676603847725*(2*x - 3)^21)/150
9418723079880704 + (5940132943613849875*(2*x - 3)^22)/1625527855624486912
+ (5717978503620010375*(2*x - 3)^23)/14629750700620382208 + (1780569958183
25525*(2*x - 3)^24)/5689347494685704192 + (179665281323275*(2*x - 3)^25)/1
01595490976530432 + (1433237383402275*(2*x - 3)^26)/22757389978742816768 +
(24229218097975*(2*x - 3)^27)/22757389978742816768 + 37120/2261)/(2066104
6784*(3 - 2*x)^(19/2) - 92974710528*(3 - 2*x)^(21/2) + 199231522560*(3 - 2
*x)^(23/2) - 270069397248*(3 - 2*x)^(25/2) + 259475340096*(3 - 2*x)^(27...

```

Reduce [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 10154, normalized size of antiderivative = 15.67

$$\int \frac{1}{(3-2x)^{21/2} (1+x+2x^2)^{10}} dx = \text{Too large to display}$$

input

```
int(1/(3-2*x)^(21/2)/(2*x^2+x+1)^10,x)
```

output

```
( - 4099829936255400031027200*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**27 + 36898469426298600279244800*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**26 - 138369260348619751047168000*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**25 + 298262627862580352257228800*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**24 - 488904719898456453699993600*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**23 + 784092475308845255933952000*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**22 - 1132321780519538296069324800*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**21 + 1326038745007605947535360000*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**20 - 1524079748881567966221619200*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**19 + 1745726804810375530399027200*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqr...
```


$$3.49 \quad \int \frac{1}{(3-2x)^{41/2} (1+x+2x^2)^{20}} dx$$

Optimal result	392
Mathematica [C] (verified)	393
Rubi [F]	394
Maple [A] (verified)	400
Fricas [B] (verification not implemented)	401
Sympy [F(-1)]	402
Maxima [F]	403
Giac [A] (verification not implemented)	403
Mupad [B] (verification not implemented)	404
Reduce [B] (verification not implemented)	405

Optimal result

Integrand size = 20, antiderivative size = 1058

$$\int \frac{1}{(3-2x)^{41/2} (1+x+2x^2)^{20}} dx = \text{Too large to display}$$

output

```
-143401467550777247627940437025/73985542663511997461099839851260280832/(3-
2*x)^(9/2)-4611053278117143010907562317585/7250583181024175751187784305423
507521536/(3-2*x)^(7/2)-405965372440630510720926890227/2071595194578335928
910795515835287863296/(3-2*x)^(5/2)-4986681479187781853417316522775/870069
98172290109014253411665082090258432/(3-2*x)^(3/2)-927027754781476746208047
620505/58004665448193406009502274443388060172288/(3-2*x)^(1/2)-39481943432
91401740321996415/202881463139404195937734623232/(3-2*x)^(37/2)-3046882292
62620222736480811/537361713180043545997243056128/(3-2*x)^(35/2)+2124315846
756567455653862925/1688851098565851144562763890688/(3-2*x)^(33/2)+47657515
074514118796095929535/66632852434325399703658138959872/(3-2*x)^(31/2)+3491
1619993974714062172751985/124667917457770102671360389021696/(3-2*x)^(29/2)
+149066309808794760843017404825/1624981820656451683095663001731072/(3-2*x)
^(27/2)+15848613964169066543734380171/601845118761648771516912222863360/(3
-2*x)^(25/2)+11155168222970774232376891145/1685166332532616560247354224017
408/(3-2*x)^(23/2)+14011818498091020272474956375/1011099799519569936148412
5344104448/(3-2*x)^(21/2)-13056959628363355534285785425/106924014357253562
723941220352/(3-2*x)^(39/2)+5/595601664*(751303+1831285*x)/(3-2*x)^(39/2)/
(2*x^2+x+1)^16+1/25015269888*(184959785+429411497*x)/(3-2*x)^(39/2)/(2*x^2
+x+1)^15+1/4902992898048*(41652915209+92630823167*x)/(3-2*x)^(39/2)/(2*x^2
+x+1)^14+1/297448235814912*(2871555518177+6100156355517*x)/(3-2*x)^(39/2)...
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 36.09 (sec) , antiderivative size = 1242, normalized size of antiderivative = 1.17

$$\int \frac{1}{(3-2x)^{41/2}(1+x+2x^2)^{20}} dx = \text{Too large to display}$$

input

```
Integrate[1/((3 - 2*x)^(41/2)*(1 + x + 2*x^2)^20), x]
```

output

```

-1/150276832468*(393*sqrt[3 - 2*x] + 287*(3 - 2*x)^(3/2))/(14 - 7*(3 - 2*x)
) + (3 - 2*x)^2^19 - (-4226921*sqrt[3 - 2*x] + 1313129*(3 - 2*x)^(3/2))/(
75739523563872*(14 - 7*(3 - 2*x) + (3 - 2*x)^2)^18) - (-3401932701*sqrt[3
- 2*x] + 760755809*(3 - 2*x)^(3/2))/(36052013216403072*(14 - 7*(3 - 2*x) +
(3 - 2*x)^2)^17) - (5*(-146490500023*sqrt[3 - 2*x] + 16144709919*(3 - 2*x)
)^(3/2)))/(16151301920948576256*(14 - 7*(3 - 2*x) + (3 - 2*x)^2)^16) - (97
45709632283*sqrt[3 - 2*x] - 4557912048927*(3 - 2*x)^(3/2))/(45223645378656
0135168*(14 - 7*(3 - 2*x) + (3 - 2*x)^2)^15) - (435856117815771*sqrt[3 - 2
*x] - 123609208162571*(3 - 2*x)^(3/2))/(9330352099175345946624*(14 - 7*(3
- 2*x) + (3 - 2*x)^2)^14) - (127435522656997631*sqrt[3 - 2*x] - 3127030241
4674811*(3 - 2*x)^(3/2))/(3396248164099825924571136*(14 - 7*(3 - 2*x) + (3
- 2*x)^2)^13) + (5*(-1540359167602841319*sqrt[3 - 2*x] + 3420265577570880
31*(3 - 2*x)^(3/2)))/(380379794379180503551967232*(14 - 7*(3 - 2*x) + (3 -
2*x)^2)^12) + (5*(-21084628139481190687*sqrt[3 - 2*x] + 41586699245502578
27*(3 - 2*x)^(3/2)))/(13017441852087510566000656384*(14 - 7*(3 - 2*x) + (3
- 2*x)^2)^11) - (1633293973597342712581*sqrt[3 - 2*x] - 23708074415419338
4005*(3 - 2*x)^(3/2))/(728976743716900591696036757504*(14 - 7*(3 - 2*x) +
(3 - 2*x)^2)^10) - (7350432513431022017155*sqrt[3 - 2*x] + 513156431847137
6538977*(3 - 2*x)^(3/2))/(61234046472219649702467087630336*(14 - 7*(3 - 2*
x) + (3 - 2*x)^2)^9) - (-113207386492327172550771*sqrt[3 - 2*x] + 43421...

```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(3-2x)^{41/2} (2x^2+x+1)^{20}} dx \\
 & \quad \downarrow 1165 \\
 & \int \frac{28(130-113x)}{(3-2x)^{41/2} (2x^2+x+1)^{19}} dx + \frac{x}{133(3-2x)^{39/2} (2x^2+x+1)^{19}} \\
 & \quad \downarrow 27 \\
 & \frac{1}{133} \int \frac{130-113x}{(3-2x)^{41/2} (2x^2+x+1)^{19}} dx + \frac{x}{133(3-2x)^{39/2} (2x^2+x+1)^{19}} \\
 & \quad \downarrow 1235
 \end{aligned}$$

$$\begin{aligned}
& \frac{1}{133} \left(\frac{\int \frac{14(33336-40657x)}{(3-2x)^{41/2}(2x^2+x+1)^{18}} dx}{3528} + \frac{373x+113}{252(3-2x)^{39/2}(2x^2+x+1)^{18}} \right) + \\
& \qquad \qquad \qquad \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{1}{133} \left(\frac{1}{252} \int \frac{33336-40657x}{(3-2x)^{41/2}(2x^2+x+1)^{18}} dx + \frac{373x+113}{252(3-2x)^{39/2}(2x^2+x+1)^{18}} \right) + \\
& \qquad \qquad \qquad \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}} \\
& \qquad \qquad \qquad \downarrow 1235 \\
& \frac{1}{133} \left(\frac{1}{252} \left(\frac{\int \frac{210(539991-751303x)}{(3-2x)^{41/2}(2x^2+x+1)^{17}} dx}{3332} + \frac{107329x+40657}{238(3-2x)^{39/2}(2x^2+x+1)^{17}} \right) + \frac{373x+113}{252(3-2x)^{39/2}(2x^2+x+1)^{18}} \right) \\
& \qquad \qquad \qquad \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \int \frac{539991-751303x}{(3-2x)^{41/2}(2x^2+x+1)^{17}} dx + \frac{107329x+40657}{238(3-2x)^{39/2}(2x^2+x+1)^{17}} \right) + \frac{373x+113}{252(3-2x)^{39/2}(2x^2+x+1)^{18}} \right) \\
& \qquad \qquad \qquad \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}} \\
& \qquad \qquad \qquad \downarrow 1235 \\
& \frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{\int \frac{14(122225856-184959785x)}{(3-2x)^{41/2}(2x^2+x+1)^{16}} dx}{3136} + \frac{1831285x+751303}{224(3-2x)^{39/2}(2x^2+x+1)^{16}} \right) + \frac{107329x+40657}{238(3-2x)^{39/2}(2x^2+x+1)^{17}} \right) \right) \\
& \qquad \qquad \qquad \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}} \\
& \qquad \qquad \qquad \downarrow 27 \\
& \frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \int \frac{122225856-184959785x}{(3-2x)^{41/2}(2x^2+x+1)^{16}} dx + \frac{1831285x+751303}{224(3-2x)^{39/2}(2x^2+x+1)^{16}} \right) + \frac{107329x+40657}{238(3-2x)^{39/2}(2x^2+x+1)^{17}} \right) \right) \\
& \qquad \qquad \qquad \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}} \\
& \qquad \qquad \qquad \downarrow 1235
\end{aligned}$$

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{\int \frac{14(25488953979-41652915209x)}{(3-2x)^{41/2}(2x^2+x+1)^{15}} dx}{2940} + \frac{429411497x + 184959785}{210(3-2x)^{39/2}(2x^2+x+1)^{15}} \right) + \frac{1831285x}{224(3-2x)^{39/2}} \right) \right) \right) + \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}}$$

↓ 27

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \int \frac{25488953979 - 41652915209x}{(3-2x)^{41/2}(2x^2+x+1)^{15}} dx + \frac{429411497x + 184959785}{210(3-2x)^{39/2}(2x^2+x+1)^{15}} \right) + \frac{1831285x}{224(3-2x)^{39/2}} \right) \right) \right) + \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}}$$

↓ 1235

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{\int \frac{42(1614300418670-2871555518177x)}{(3-2x)^{41/2}(2x^2+x+1)^{14}} dx}{2744} + \frac{92630823167x + 41652915209}{196(3-2x)^{39/2}(2x^2+x+1)^{14}} \right) + \frac{429411497x + 184959785}{210(3-2x)^{39/2}} \right) \right) \right) \right) + \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}}$$

↓ 27

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \int \frac{1614300418670 - 2871555518177x}{(3-2x)^{41/2}(2x^2+x+1)^{14}} dx + \frac{92630823167x + 41652915209}{196(3-2x)^{39/2}(2x^2+x+1)^{14}} \right) + \frac{429411497x + 184959785}{210(3-2x)^{39/2}} \right) \right) \right) \right) + \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}}$$

↓ 1235

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{\int \frac{98(39357458161627-77559130805859x)}{(3-2x)^{41/2}(2x^2+x+1)^{13}} dx}{2548} + \frac{6100156355517x + 2871555518177}{182(3-2x)^{39/2}(2x^2+x+1)^{13}} \right) + \frac{429411497x + 184959785}{210(3-2x)^{39/2}} \right) \right) \right) \right) \right) + \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}}$$

↓ 27

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{1}{26} \int \frac{39357458161627 - 77559130805859x}{(3-2x)^{41/2}(2x^2+x+1)^{13}} dx + \frac{6100156355517x + 2871555518177}{182(3-2x)^{39/2}(2x^2+x+1)^{13}} \right) + \frac{429411497x + 184959785}{210(3-2x)^{39/2}} \right) \right) \right) \right) \right) + \frac{x}{133(3-2x)^{39/2}(2x^2+x+1)^{19}}$$

↓ 1235

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{1}{26} \left(\int \frac{70(1182110687469684 - 2656658801194921x)}{(3-2x)^{41/2}(2x^2+x+1)^{12}} dx \right. \right. \right. \right. \right. \right. \right. \right. + \frac{156274047129113x + 7755}{168(3-2x)^{39/2}(2x^2+x+1)^{19}} \right. \right. \right. \right. \right. \right. \right. \right. \right.$$

↓ 27

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{1}{26} \left(\frac{5}{168} \int \frac{1182110687469684 - 2656658801194921x}{(3-2x)^{41/2}(2x^2+x+1)^{12}} dx \right. \right. \right. \right. \right. \right. \right. \right. + \frac{156274047129113}{168(3-2x)^{39/2}(2x^2+x+1)^{19}} \right. \right. \right. \right. \right. \right. \right. \right. \right.$$

↓ 1235

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{1}{26} \left(\frac{5}{168} \left(\int \frac{126(16782494726084327 - 45187921585208601x)}{(3-2x)^{41/2}(2x^2+x+1)^{11}} dx \right. \right. \right. \right. \right. \right. \right. \right. \right. + \frac{502088017613428}{154(3-2x)^{39/2}(2x^2+x+1)^{19}} \right. \right. \right. \right. \right. \right. \right. \right. \right.$$

↓ 27

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{1}{26} \left(\frac{5}{168} \left(\frac{9}{154} \int \frac{16782494726084327 - 45187921585208601x}{(3-2x)^{41/2}(2x^2+x+1)^{11}} dx \right. \right. \right. \right. \right. \right. \right. \right. \right. + \frac{50208801}{154(3-2x)^{39/2}(2x^2+x+1)^{19}} \right. \right. \right. \right. \right. \right. \right. \right. \right.$$

↓ 1235

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{1}{26} \left(\frac{5}{168} \left(\frac{9}{154} \left(\int \frac{14(1706599234272796606 - 6063974149878048635x)}{(3-2x)^{41/2}(2x^2+x+1)^{10}} dx \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. + \frac{78752911}{140(3-2x)^{39/2}(2x^2+x+1)^{19}} \right. \right. \right. \right. \right. \right. \right. \right. \right.$$

↓ 27

$$\frac{1}{133} \left(\frac{1}{252} \left(\frac{15}{238} \left(\frac{1}{224} \left(\frac{1}{210} \left(\frac{3}{196} \left(\frac{1}{26} \left(\frac{5}{168} \left(\frac{9}{154} \left(\frac{1}{140} \int \frac{1706599234272796606 - 6063974149878048635x}{(3-2x)^{41/2}(2x^2+x+1)^{10}} dx \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. + \frac{78752911}{140(3-2x)^{39/2}(2x^2+x+1)^{19}} \right. \right. \right. \right. \right. \right. \right. \right. \right.$$

↓ 1235

rule 1165

```
Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Simp[(d + e*x)^(m + 1)*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x]
+ Simp[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^m*Simp[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && LtQ[p, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]
```

rule 1235

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Simp[(d + e*x)^(m + 1)*(f*(b*c*d - b^2*e + 2*a*c*e) - a*g*(2*c*d - b*e) + c*(f*(2*c*d - b*e) - g*(b*d - 2*a*e))*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x]
+ Simp[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)) Int[(d + e*x)^m*(a + b*x + c*x^2)^(p + 1)*Simp[f*(b*c*d*e*(2*p - m + 2) + b^2*e^2*(p + m + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3)) - g*(a*e*(b*e - 2*c*d*m + b*e*m) - b*d*(3*c*d - b*e + 2*c*d*p - b*e*p)) + c*e*(g*(b*d - 2*a*e) - f*(2*c*d - b*e))*(m + 2*p + 4)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && LtQ[p, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

Maple [A] (verified)

Time = 8.08 (sec) , antiderivative size = 502, normalized size of antiderivative = 0.47

method	result	size
pseudoelliptic	Expression too large to display	502
trager	Expression too large to display	733
risch	Expression too large to display	761
derivativdivides	Expression too large to display	820
default	Expression too large to display	820

input

```
int(1/(3-2*x)^(41/2)/(2*x^2+x+1)^20,x,method=_RETURNVERBOSE)
```

output

```

115/5908552821163231304184823545856*((x-3/2)^19*(x^2+1/2*x+1/2)^19*(1/2*(7
+2*14^(1/2))^(1/2)*(62541562556792464940960784209*14^(1/2)-234044028404883
307655877091262)*(ln(3-2*x+14^(1/2)-(3-2*x)^(1/2)*(7+2*14^(1/2))^(1/2))-ln
(3-2*x+14^(1/2)+(3-2*x)^(1/2)*(7+2*14^(1/2))^(1/2)))*(-7+2*14^(1/2))^(1/2)
+(30297118912219360725028693061*14^(1/2)+112855552756005864755762319018)*(
arctan((2*(3-2*x)^(1/2)-(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2))+arctan
((2*(3-2*x)^(1/2)+(7+2*14^(1/2))^(1/2))/(-7+2*14^(1/2))^(1/2)))*(-7+2*14^(1/2))^(1/2)
+225711105512011729511524638036*(-7+2*14^(1/2))^(1/2)*(35128571782045
484630026117801687570244083874053171/4517315819499287681645418200730380459
26298694451200*x+183212802026248860514288501929804345458085916920873773/57
83192372047452224012626430522535438601263513600*x^17-339692530351150840696
302021012775910383197972648611749049/2202191462005902744802141372856060473
890706135449600*x^20+16088554099632166662170276395247605557702557286910897
24699/17617531696047221958417130982848483791125649083596800*x^19+330991985
1041151721938329104326989773026441469925012281/135519474584978630449362546
02191141377788960833536*x^21-410719388838590997785308221509211987369317097
087246119213/1101095731002951372401070686428030236945353067724800*x^22+625
318899982989425383877400685785463778432122219115569133/1101095731002951372
401070686428030236945353067724800*x^23-27406709657651230963424691464485550
30280367060674320773/3329926605855699714922592801697672087536350003200*...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1986 vs. $2(821) = 1642$.

Time = 0.32 (sec) , antiderivative size = 1986, normalized size of antiderivative = 1.88

$$\int \frac{1}{(3-2x)^{41/2}(1+x+2x^2)^{20}} dx = \text{Too large to display}$$

input

```
integrate(1/(3-2*x)^(41/2)/(2*x^2+x+1)^20,x, algorithm="fricas")
```

output

```
-1/336864077912586356135291702496114974019074478550548480*(47704745152350*
sqrt(1/2)*(549755813888*x^58 - 11269994184704*x^57 + 107064944754688*x^56
- 630638638006272*x^55 + 2618521301286912*x^54 - 8342252417974272*x^53 + 2
1849572376576000*x^52 - 49684091485814784*x^51 + 101394501297242112*x^50 -
188583312363618304*x^49 + 323261995581177856*x^48 - 517079841212727296*x^
47 + 778117896260812800*x^46 - 1105641165387988992*x^45 + 1491287028233404
416*x^44 - 1919929663119949824*x^43 + 2363050939901804544*x^42 - 278627402
0645928960*x^41 + 3161145685194047488*x^40 - 3453753931369283584*x^39 + 36
34098467102523392*x^38 - 3697893960325791744*x^37 + 3640651752731836416*x^
36 - 3461798212247617536*x^35 + 3194540251789393920*x^34 - 286154457949529
7024*x^33 + 2477632938217930752*x^32 - 2088430257127768064*x^31 + 17127610
05459316736*x^30 - 1355447485390974976*x^29 + 1048940886155151360*x^28 - 7
90511024135089152*x^27 + 571750925528393856*x^26 - 408374103192240192*x^25
+ 282845069599813728*x^24 - 186113897194906128*x^23 + 123982890381352520*
x^22 - 78116367732251996*x^21 + 46488580159296898*x^20 - 29591055660829971
*x^19 + 16200795673453545*x^18 - 8941894120163277*x^17 + 5578893209169441*
x^16 - 2296849711499532*x^15 + 1448289882400788*x^14 - 756896247319212*x^1
3 + 182213447974992*x^12 - 240797810407770*x^11 + 25549234281774*x^10 - 26
500281727302*x^9 + 25520701332582*x^8 + 9965507230260*x^7 + 10389354811164
*x^6 + 3755740313808*x^5 + 1820618017974*x^4 + 463742325333*x^3 + 13985...
```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(3-2x)^{41/2} (1+x+2x^2)^{20}} dx = \text{Timed out}$$

input

```
integrate(1/(3-2*x)**(41/2)/(2*x**2+x+1)**20,x)
```

output

Timed out

Maxima [F]

$$\int \frac{1}{(3-2x)^{41/2} (1+x+2x^2)^{20}} dx = \int \frac{1}{(2x^2+x+1)^{20} (-2x+3)^{\frac{41}{2}}} dx$$

input `integrate(1/(3-2*x)^(41/2)/(2*x^2+x+1)^20,x, algorithm="maxima")`

output `integrate(1/((2*x^2 + x + 1)^20*(-2*x + 3)^(41/2)), x)`

Giac [A] (verification not implemented)

Time = 0.93 (sec) , antiderivative size = 1410, normalized size of antiderivative = 1.33

$$\int \frac{1}{(3-2x)^{41/2} (1+x+2x^2)^{20}} dx = \text{Too large to display}$$

input `integrate(1/(3-2*x)^(41/2)/(2*x^2+x+1)^20,x, algorithm="giac")`

output

```
-115/363805261691069042491598265308929913400590336*sqrt(7)*(24183332733429
828161949068361*14^(3/4)*sqrt(7)*(sqrt(14) + 4)*sqrt(-2*sqrt(14) + 8) + 80
61110911143276053983022787*14^(3/4)*sqrt(7)*(sqrt(14) - 4)*sqrt(-2*sqrt(14)
) + 8) - 56427776378002932377881159509*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt
(14) + 4) - 169283329134008797133643478527*14^(3/4)*sqrt(2*sqrt(14) + 8)*(
sqrt(14) - 4) + 242376951297754885800229544488*14^(1/4)*sqrt(7)*sqrt(-2*sq
rt(14) + 8) - 1696638659084284200601606811416*14^(1/4)*sqrt(2*sqrt(14) + 8
))*arctan(1/28*14^(3/4)*(14^(1/4)*sqrt(1/2)*sqrt(sqrt(14) + 4) + 2*sqrt(-2
*x + 3))/sqrt(-1/8*sqrt(14) + 1/2)) - 115/36380526169106904249159826530892
9913400590336*sqrt(7)*(24183332733429828161949068361*14^(3/4)*sqrt(7)*(sq
rt(14) + 4)*sqrt(-2*sqrt(14) + 8) + 8061110911143276053983022787*14^(3/4)*s
qrt(7)*(sqrt(14) - 4)*sqrt(-2*sqrt(14) + 8) - 5642777637800293237788115950
9*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) - 1692833291340087971336434
78527*14^(3/4)*sqrt(2*sqrt(14) + 8)*(sqrt(14) - 4) + 242376951297754885800
229544488*14^(1/4)*sqrt(7)*sqrt(-2*sqrt(14) + 8) - 16966386590842842006016
06811416*14^(1/4)*sqrt(2*sqrt(14) + 8))*arctan(-1/28*14^(3/4)*(14^(1/4)*sq
rt(1/2)*sqrt(sqrt(14) + 4) - 2*sqrt(-2*x + 3))/sqrt(-1/8*sqrt(14) + 1/2))
- 115/727610523382138084983196530617859826801180672*sqrt(7)*(8061110911143
276053983022787*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(14) + 4) + 241
83332733429828161949068361*14^(3/4)*sqrt(7)*sqrt(2*sqrt(14) + 8)*(sqrt(...
```

Mupad [B] (verification not implemented)

Time = 0.68 (sec) , antiderivative size = 1017, normalized size of antiderivative = 0.96

$$\int \frac{1}{(3 - 2x)^{41/2} (1 + x + 2x^2)^{20}} dx = \text{Too large to display}$$

input

```
int(1/((3 - 2*x)^(41/2)*(x + 2*x^2 + 1)^20),x)
```

output

```

((64356352*(2*x - 3)^2)/38073 - (5767168*x)/1443 - (7517962240*(2*x - 3)^3
)/5444439 + (1357449428992*(2*x - 3)^4)/1181443263 - (34130408095744*(2*x
- 3)^5)/34261854627 + (1965832636456960*(2*x - 3)^6)/2158496841501 - (9552
588571922432*(2*x - 3)^7)/10792484207505 + (69571472879183872*(2*x - 3)^8)
/75547389452535 - (5204838729946112*(2*x - 3)^9)/5036492630169 + (32508205
2781755904*(2*x - 3)^10)/257635969158645 - (461538785202937088*(2*x - 3)^1
1)/272428464995505 + (17726678744562203264*(2*x - 3)^12)/6992330601551295
- (1432471149647610304*(2*x - 3)^13)/332968123883395 + (204346360124338870
4*(2*x - 3)^14)/241114848329355 - (96972768477343976816*(2*x - 3)^15)/4840
844262612435 + (10833870670122545927656*(2*x - 3)^16)/181389282075536535 -
(44340157049832305729324*(2*x - 3)^17)/181389282075536535 + (691509778132
186261807282*(2*x - 3)^18)/423241658176251915 - (1357735833153708223970340
7*(2*x - 3)^19)/423241658176251915 + (509495943858959939640753039465067261
4981*(2*x - 3)^20)/203594616979243053623625646080 + (474753402737241482257
49886260884632526403*(2*x - 3)^21)/203594616979243053623625646080 + (54736
2406727667345868176230754600752341499*(2*x - 3)^22)/5182408432198914092237
74371840 + (1363217399168846741803250531443496167647559*(2*x - 3)^23)/4385
11482724523500112424468480 + (40035704814224807138997531075224002020138815
9*(2*x - 3)^24)/59856817391897457765345939947520 + (1678035321867106187517
78512174316524508553291*(2*x - 3)^25)/14964204347974364441336484986880 ...

```

Reduce [B] (verification not implemented)

Time = 0.59 (sec) , antiderivative size = 21074, normalized size of antiderivative = 19.92

$$\int \frac{1}{(3-2x)^{41/2} (1+x+2x^2)^{20}} dx = \text{Too large to display}$$

input

```
int(1/(3-2*x)^(41/2)/(2*x^2+x+1)^20,x)
```

output

```
( - 820106290170129967145447171095953091911999318235545600*sqrt(2*sqrt(14)
- 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14)
+ 7))/sqrt(2*sqrt(14) - 7))*x**57 + 1558201951323246937576349625082310874
6327987046475366400*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2
*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**56 - 13
6342670740784107037930592194702201530369886656659456000*sqrt(2*sqrt(14) -
7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) +
7))/sqrt(2*sqrt(14) - 7))*x**55 + 7362504220002341780048251978513918882639
97387945961062400*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*s
qrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**54 - 2801
841883723113399629473669601130241449101170794351820800*sqrt(2*sqrt(14) - 7
)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7
))/sqrt(2*sqrt(14) - 7))*x**53 + 82419144462803992704429042981697480825108
59648395064115200*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*s
qrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**52 - 2023
1548054548101683090925749291372929587761556265055027200*sqrt(2*sqrt(14) -
7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) +
7))/sqrt(2*sqrt(14) - 7))*x**51 + 4376964934361511221918566466393319335736
1154238751703040000*sqrt(2*sqrt(14) - 7)*sqrt(- 2*x + 3)*sqrt(14)*atan((2
*sqrt(- 2*x + 3) - sqrt(2*sqrt(14) + 7))/sqrt(2*sqrt(14) - 7))*x**50 - ...
```

3.50 $\int \frac{1}{(3-2x+x^2)^{11/2}(1+x+2x^2)^5} dx$

Optimal result	407
Mathematica [C] (verified)	408
Rubi [A] (verified)	409
Maple [A] (verified)	416
Fricas [B] (verification not implemented)	417
Sympy [F]	418
Maxima [F]	419
Giac [C] (verification not implemented)	419
Mupad [F(-1)]	420
Reduce [F]	421

Optimal result

Integrand size = 23, antiderivative size = 378

$$\int \frac{1}{(3-2x+x^2)^{11/2}(1+x+2x^2)^5} dx = -\frac{3450497-2004270x}{123480000(3-2x+x^2)^{9/2}} - \frac{4878869-2578034x}{411600000(3-2x+x^2)^{7/2}} - \frac{30316369-15043110x}{6860000000(3-2x+x^2)^{5/2}} - \frac{63043297-29625922x}{41160000000(3-2x+x^2)^{3/2}} - \frac{31(7434109-3088870x)}{41160000000\sqrt{3-2x+x^2}} - \frac{1-10x}{280(3-2x+x^2)^{9/2}(1+x+2x^2)^4} + \frac{28+67x}{1050(3-2x+x^2)^{9/2}(1+x+2x^2)^3} + \frac{5485+8878x}{117600(3-2x+x^2)^{9/2}(1+x+2x^2)^2} + \frac{3(8822+8233x)}{343000(3-2x+x^2)^{9/2}(1+x+2x^2)} + \frac{\sqrt{\frac{1}{70}(151363871237318045+110320475741093888\sqrt{2})} \arctan\left(\frac{\sqrt{\frac{5}{7(151363871237318045+110320475741093888\sqrt{2})}}(3081)}}{\sqrt{3-2x+x^2}}\right)}{13720000000} + \frac{\sqrt{\frac{1}{70}(-151363871237318045+110320475741093888\sqrt{2})} \operatorname{arctanh}\left(\frac{\sqrt{\frac{5}{7(-151363871237318045+110320475741093888\sqrt{2})}}}}{\sqrt{3-2x+x^2}}\right)}{13720000000}$$

output

```

1/123480000*(-3450497+2004270*x)/(x^2-2*x+3)^(9/2)+1/411600000*(-4878869+2
578034*x)/(x^2-2*x+3)^(7/2)+1/6860000000*(-30316369+15043110*x)/(x^2-2*x+3
)^(5/2)+1/41160000000*(-63043297+29625922*x)/(x^2-2*x+3)^(3/2)+1/280*(-1+1
0*x)/(x^2-2*x+3)^(9/2)/(2*x^2+x+1)^4+1/1050*(28+67*x)/(x^2-2*x+3)^(9/2)/(2
*x^2+x+1)^3+1/117600*(5485+8878*x)/(x^2-2*x+3)^(9/2)/(2*x^2+x+1)^2+3/34300
0*(8822+8233*x)/(x^2-2*x+3)^(9/2)/(2*x^2+x+1)-31/411600000000*(7434109-308
8870*x)/(x^2-2*x+3)^(1/2)-1/9604000000000*arctanh(1/7*(308108167+x*(932587
773-620347970*2^(1/2))-312239803*2^(1/2))*35^(1/2)/(-151363871237318045+11
0320475741093888*2^(1/2))^(1/2)/(x^2-2*x+3)^(1/2))*(-10595470986612263150+
7722433301876572160*2^(1/2))^(1/2)+1/9604000000000*arctan(1/7*(308108167+x
12239803*2^(1/2)+x*(932587773+620347970*2^(1/2)))*35^(1/2)/(15136387123731
8045+110320475741093888*2^(1/2))^(1/2)/(x^2-2*x+3)^(1/2))*(105954709866122
63150+7722433301876572160*2^(1/2))^(1/2)

```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 2.58 (sec) , antiderivative size = 733, normalized size of antiderivative = 1.94

$$\int \frac{1}{(3 - 2x + x^2)^{11/2} (1 + x + 2x^2)^5} dx = \text{Too large to display}$$

input

```
Integrate[1/((3 - 2*x + x^2)^(11/2)*(1 + x + 2*x^2)^5),x]
```

output

```

((-53205422447 + 261702502714*x - 266966654968*x^2 + 1002897791524*x^3 - 1
409335257371*x^4 + 25034272226914*x^5 - 3359813871472*x^6 + 4591320676952*x
^7 - 5134334619701*x^8 + 5380603084494*x^9 - 4915797913008*x^10 + 39996561
32532*x^11 - 2679143870481*x^12 + 1459208021718*x^13 - 606785954952*x^14 +
188603773872*x^15 - 38639385552*x^16 + 4596238560*x^17)/((3 - 2*x + x^2)^
(9/2)*(1 + x + 2*x^2)^4) - 49392*RootSum[14 + 7*#1 - 5*#1^2 - #1^3 + #1^4
& , (-6014*Log[-x + Sqrt[3 - 2*x + x^2] - #1] - 10727*Log[-x + Sqrt[3 - 2*
x + x^2] - #1]*#1 + 3229*Log[-x + Sqrt[3 - 2*x + x^2] - #1]*#1^2)/(7 - 10*
#1 - 3*#1^2 + 4*#1^3) & ] - 56448*RootSum[14 + 7*#1 - 5*#1^2 - #1^3 + #1^4
& , (73781*Log[-x + Sqrt[3 - 2*x + x^2] - #1] - 60407*Log[-x + Sqrt[3 - 2
*x + x^2] - #1]*#1 + 13104*Log[-x + Sqrt[3 - 2*x + x^2] - #1]*#1^2)/(7 - 1
0*#1 - 3*#1^2 + 4*#1^3) & ] - 504*RootSum[14 + 7*#1 - 5*#1^2 - #1^3 + #1^4
& , (275935046*Log[-x + Sqrt[3 - 2*x + x^2] - #1] - 208696097*Log[-x + Sq
rt[3 - 2*x + x^2] - #1]*#1 + 50007219*Log[-x + Sqrt[3 - 2*x + x^2] - #1]*#
1^2)/(7 - 10*#1 - 3*#1^2 + 4*#1^3) & ] + 1440*RootSum[14 + 7*#1 - 5*#1^2 -
#1^3 + #1^4 & , (3276009822*Log[-x + Sqrt[3 - 2*x + x^2] - #1] - 24478316
21*Log[-x + Sqrt[3 - 2*x + x^2] - #1]*#1 + 590084719*Log[-x + Sqrt[3 - 2*x
+ x^2] - #1]*#1^2)/(7 - 10*#1 - 3*#1^2 + 4*#1^3) & ] - 18*RootSum[14 + 7*
#1 - 5*#1^2 - #1^3 + #1^4 & , (254137663854*Log[-x + Sqrt[3 - 2*x + x^2] -
#1] - 189631531133*Log[-x + Sqrt[3 - 2*x + x^2] - #1]*#1 + 45801521671...

```

Rubi [A] (verified)

Time = 1.33 (sec) , antiderivative size = 424, normalized size of antiderivative = 1.12, number of steps used = 24, number of rules used = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {1305, 27, 2135, 27, 2135, 27, 2135, 27, 2135, 27, 2135, 27, 2135, 27, 2135, 27, 2135, 27, 1368, 27, 1362, 217, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)^5} dx$$

$$\downarrow 1305$$

$$\int -\frac{5(160x^2 - 267x + 247)}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)^4} dx - \frac{1 - 10x}{280 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^4}$$

$$\downarrow 27$$

$$\begin{aligned}
& \frac{1}{280} \int \frac{160x^2 - 267x + 247}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)^4} dx - \frac{1 - 10x}{280 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^4} \\
& \quad \downarrow 2135 \\
& \frac{1}{280} \left(\frac{\int \frac{70(3752x^2 - 3814x + 2901)}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)^3} dx}{1050} + \frac{4(67x + 28)}{15 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^3} \right) - \\
& \quad \frac{1 - 10x}{280 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^4} \\
& \quad \downarrow 27 \\
& \frac{1}{280} \left(\frac{1}{15} \int \frac{3752x^2 - 3814x + 2901}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)^3} dx + \frac{4(67x + 28)}{15 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^3} \right) - \\
& \quad \frac{1 - 10x}{280 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^4} \\
& \quad \downarrow 2135 \\
& \frac{1}{280} \left(\frac{1}{15} \left(\frac{1}{700} \int \frac{75(35512x^2 - 23581x + 12713)}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)^2} dx + \frac{8878x + 5485}{28 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^2} \right) + \frac{4(67x + 28)}{15 (x^2 - 2x + 3)^9} \right) - \\
& \quad \frac{1 - 10x}{280 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^4} \\
& \quad \downarrow 27 \\
& \frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \int \frac{35512x^2 - 23581x + 12713}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)^2} dx + \frac{8878x + 5485}{28 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^2} \right) + \frac{4(67x + 28)}{15 (x^2 - 2x + 3)^9} \right) - \\
& \quad \frac{1 - 10x}{280 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^4} \\
& \quad \downarrow 2135 \\
& \frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{350} \int -\frac{10(-987960x^2 - 28350x + 486617)}{(x^2 - 2x + 3)^{11/2} (2x^2 + x + 1)} dx + \frac{12(8233x + 8822)}{35 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)} \right) + \frac{4(67x + 28)}{15 (x^2 - 2x + 3)^9} \right) - \right. \\
& \quad \left. \frac{1 - 10x}{280 (x^2 - 2x + 3)^{9/2} (2x^2 + x + 1)^4} \right) \\
& \quad \downarrow 27
\end{aligned}$$

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{12(8233x + 8822)}{35(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)} - \frac{1}{35} \int \frac{-987960x^2 - 28350x + 486617}{(x^2 - 2x + 3)^{11/2}(2x^2 + x + 1)} dx \right) + \frac{8878}{28(x^2 - 2x + 3)} \right) \right)$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 2135

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(- \frac{\int \frac{60(-10689440x^2 + 3332642x + 2083763)}{(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)} dx}{1800} - \frac{3450497 - 2004270x}{90(x^2 - 2x + 3)^{9/2}} \right) + \frac{12(8233x + 8822)}{35(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)} \right) \right)$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(- \frac{1}{30} \int \frac{-10689440x^2 + 3332642x + 2083763}{(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)} dx - \frac{3450497 - 2004270x}{90(x^2 - 2x + 3)^{9/2}} \right) + \frac{12(8233x + 8822)}{35(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)} \right) \right)$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 2135

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(- \frac{\int -\frac{420(10312136x^2 - 5581162x + 620039)}{(x^2 - 2x + 3)^{7/2}(2x^2 + x + 1)} dx}{1400} - \frac{4878869 - 2578034x}{10(x^2 - 2x + 3)^{7/2}} \right) - \frac{3450497 - 2004270x}{90(x^2 - 2x + 3)^{9/2}} \right) \right) \right)$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \int \frac{10312136x^2 - 5581162x + 620039}{(x^2 - 2x + 3)^{7/2}(2x^2 + x + 1)} dx - \frac{4878869 - 2578034x}{10(x^2 - 2x + 3)^{7/2}} \right) - \frac{3450497 - 2004270x}{90(x^2 - 2x + 3)^{9/2}} \right) \right) \right)$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 2135

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\int \frac{100(24068976x^2 - 18512030x + 9163631)}{(x^2 - 2x + 3)^{5/2}(2x^2 + x + 1)} dx - \frac{30316369 - 15043110x}{50(x^2 - 2x + 3)^{5/2}} \right) - \frac{4878869 - 29625922x}{10(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3}$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \int \frac{24068976x^2 - 18512030x + 9163631}{(x^2 - 2x + 3)^{5/2}(2x^2 + x + 1)} dx - \frac{30316369 - 15043110x}{50(x^2 - 2x + 3)^{5/2}} \right) - \frac{4878869 - 29625922x}{10(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3}$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 2135

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{600} \int \frac{20(118503688x^2 - 141252406x + 125053685)}{(x^2 - 2x + 3)^{3/2}(2x^2 + x + 1)} dx - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3}$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{30} \int \frac{118503688x^2 - 141252406x + 125053685}{(x^2 - 2x + 3)^{3/2}(2x^2 + x + 1)} dx - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3}$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 2135

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{30} \left(\frac{1}{200} \int \frac{60(132636591 - 89801606x)}{\sqrt{x^2 - 2x + 3}(2x^2 + x + 1)} dx - \frac{31(7434109 - 3088870x)}{10\sqrt{x^2 - 2x + 3}} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3}$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 27

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{3}{10} \int \frac{132636591 - 89801606x}{\sqrt{x^2 - 2x + 3}(2x^2 + x + 1)} dx - \frac{31(7434109 - 3088870x)}{10\sqrt{x^2 - 2x + 3}} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3} \right) - \frac{63043297 - 29625922x}{30(x^2 - 2x + 3)^3}$$

$$\frac{1 - 10x}{280(x^2 - 2x + 3)^{9/2}(2x^2 + x + 1)^4}$$

↓ 1368

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{30} \left(\frac{3}{10} \left(\int -\frac{5(-((42834985-89801606\sqrt{2})x)-132636591\sqrt{2}+222438197)}{\sqrt{x^2-2x+3}(2x^2+x+1)} dx - \int -\frac{5(-((42834985+89801606\sqrt{2})x)+132636591\sqrt{2}+222438197)}{\sqrt{x^2-2x+3}(2x^2+x+1)} dx \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \frac{1-10x}{280(x^2-2x+3)^{9/2}(2x^2+x+1)^4}$$

↓ 27

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{30} \left(\frac{3}{10} \left(\int \frac{-((42834985+89801606\sqrt{2})x)+132636591\sqrt{2}+222438197}{\sqrt{x^2-2x+3}(2x^2+x+1)} dx - \int \frac{-((42834985-89801606\sqrt{2})x)-132636591\sqrt{2}+222438197}{\sqrt{x^2-2x+3}(2x^2+x+1)} dx \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \frac{1-10x}{280(x^2-2x+3)^{9/2}(2x^2+x+1)^4}$$

↓ 1362

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{(151363871237318045 - 110320475741093888\sqrt{2}) \int \frac{-5((932587773-6...)}{\sqrt{x^2-2x+3}(2x^2+x+1)} \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \frac{1-10x}{280(x^2-2x+3)^{9/2}(2x^2+x+1)^4}$$

↓ 217

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{(151363871237318045 - 110320475741093888\sqrt{2}) \int \frac{-5((932587773-6...)}{\sqrt{x^2-2x+3}(2x^2+x+1)} \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \frac{1-10x}{280(x^2-2x+3)^{9/2}(2x^2+x+1)^4}$$

↓ 219

$$\frac{1}{280} \left(\frac{1}{15} \left(\frac{3}{28} \left(\frac{1}{35} \left(\frac{1}{30} \left(\frac{3}{10} \left(\frac{1}{10} \left(\frac{1}{30} \left(\frac{3}{10} \left(\sqrt{\frac{1}{70} (151363871237318045 + 110320475741093888\sqrt{2})} \right) \arctan \left(\frac{1-10x}{280(x^2-2x+3)^{9/2}(2x^2+x+1)^4} \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \right)$$

input `Int[1/((3 - 2*x + x^2)^(11/2)*(1 + x + 2*x^2)^5),x]`

output

```
-1/280*(1 - 10*x)/((3 - 2*x + x^2)^(9/2)*(1 + x + 2*x^2)^4) + ((4*(28 + 67
*x))/(15*(3 - 2*x + x^2)^(9/2)*(1 + x + 2*x^2)^3) + ((5485 + 8878*x)/(28*(
3 - 2*x + x^2)^(9/2)*(1 + x + 2*x^2)^2) + (3*((12*(8822 + 8233*x))/(35*(3
- 2*x + x^2)^(9/2)*(1 + x + 2*x^2)) + (-1/90*(3450497 - 2004270*x)/(3 - 2*
x + x^2)^(9/2) + (-1/10*(4878869 - 2578034*x)/(3 - 2*x + x^2)^(7/2) + (3*(
-1/50*(30316369 - 15043110*x)/(3 - 2*x + x^2)^(5/2) + (-1/30*(63043297 - 2
9625922*x)/(3 - 2*x + x^2)^(3/2) + ((-31*(7434109 - 3088870*x))/(10*sqrt[3
- 2*x + x^2]) + (3*(sqrt[(151363871237318045 + 110320475741093888*sqrt[2]
)/70]*ArcTan[(sqrt[5/(7*(151363871237318045 + 110320475741093888*sqrt[2]
)]*(308108167 + 312239803*sqrt[2] + (932587773 + 620347970*sqrt[2])*x))/sqrt
[3 - 2*x + x^2]) + ((151363871237318045 - 110320475741093888*sqrt[2])*Arc
Tanh[(sqrt[5/(7*(-151363871237318045 + 110320475741093888*sqrt[2]
)]*(308108167 - 312239803*sqrt[2] + (932587773 - 620347970*sqrt[2])*x))/sqrt[3 - 2
*x + x^2]])/sqrt[70*(-151363871237318045 + 110320475741093888*sqrt[2]
]])))/10)/30)/10)/10)/30)/35)/28)/15)/280
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !Ma
tchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]
```

rule 217

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

rule 219

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 1305

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_)*((d_) + (e_)*(x_) + (f_)*(x
_)^2)^(q_), x_Symbol] := Simp[(2*a*c^2*e - b^2*c*e + b^3*f + b*c*(c*d - 3*a
*f) + c*(2*c^2*d + b^2*f - c*(b*e + 2*a*f))*x*(a + b*x + c*x^2)^(p + 1)*((
d + e*x + f*x^2)^(q + 1)/((b^2 - 4*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e -
b*f))*(p + 1))), x] - Simp[1/((b^2 - 4*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(
c*e - b*f))*(p + 1)) Int[(a + b*x + c*x^2)^(p + 1)*(d + e*x + f*x^2)^q*Si
mp[2*c*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1) - (2*c^2*d + b^2*f
- c*(b*e + 2*a*f))*(a*f*(p + 1) - c*d*(p + 2)) - e*(b^2*c*e - 2*a*c^2*e -
b^3*f - b*c*(c*d - 3*a*f))*(p + q + 2) + (2*f*(2*a*c^2*e - b^2*c*e + b^3*f
+ b*c*(c*d - 3*a*f))*(p + q + 2) - (2*c^2*d + b^2*f - c*(b*e + 2*a*f))*(b*f
*(p + 1) - c*e*(2*p + q + 4))*x + c*f*(2*c^2*d + b^2*f - c*(b*e + 2*a*f))*
(2*p + 2*q + 5)*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && NeQ[b
^2 - 4*a*c, 0] && NeQ[e^2 - 4*d*f, 0] && LtQ[p, -1] && NeQ[(c*d - a*f)^2 -
(b*d - a*e)*(c*e - b*f), 0] && !(IntegerQ[p] && ILtQ[q, -1]) && !IGtQ[q
, 0]
```

rule 1362

```
Int[((g_) + (h_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2)*Sqrt[(d_) +
(e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := Simp[-2*g*(g*b - 2*a*h) Subst[I
nt[1/Simp[g*(g*b - 2*a*h)*(b^2 - 4*a*c) - (b*d - a*e)*x^2, x], x], x, Simp[
g*b - 2*a*h - (b*h - 2*g*c)*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, b
, c, d, e, f, g, h}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[e^2 - 4*d*f, 0] && Ne
Q[b*d - a*e, 0] && EqQ[h^2*(b*d - a*e) - 2*g*h*(c*d - a*f) + g^2*(c*e - b*f
), 0]
```

rule 1368

```
Int[((g_) + (h_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2)*Sqrt[(d_) +
(e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 - (b*d
- a*e)*(c*e - b*f), 2]}, Simp[1/(2*q) Int[Simp[h*(b*d - a*e) - g*(c*d -
a*f - q) - (g*(c*e - b*f) - h*(c*d - a*f + q))*x, x]/((a + b*x + c*x^2)*Sqr
t[d + e*x + f*x^2]), x], x] - Simp[1/(2*q) Int[Simp[h*(b*d - a*e) - g*(c*
d - a*f + q) - (g*(c*e - b*f) - h*(c*d - a*f - q))*x, x]/((a + b*x + c*x^2)
*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, b, c, d, e, f, g, h}, x] && N
eQ[b^2 - 4*a*c, 0] && NeQ[e^2 - 4*d*f, 0] && NeQ[b*d - a*e, 0] && NegQ[b^2
- 4*a*c]
```


rule 2135

```

Int[(Px_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_)*((d_) + (e_)*(x_) + (f_
)*(x_)^2)^(q_), x_Symbol] :=> With[{A = Coeff[Px, x, 0], B = Coeff[Px, x, 1]
, C = Coeff[Px, x, 2]}, Simp[(a + b*x + c*x^2)^(p + 1)*((d + e*x + f*x^2)^(
q + 1)/((b^2 - 4*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1))*(
(A*c - a*C)*(2*a*c*e - b*(c*d + a*f)) + (A*b - a*B)*(2*c^2*d + b^2*f - c*(b
*e + 2*a*f)) + c*(A*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)) - B*(b*c*d - 2*a*c*
e + a*b*f) + C*(b^2*d - a*b*e - 2*a*(c*d - a*f)))*x), x] + Simp[1/((b^2 - 4
*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1)) Int[(a + b*x + c
*x^2)^(p + 1)*(d + e*x + f*x^2)^q*Simp[(b*B - 2*A*c - 2*a*C)*((c*d - a*f)^2
- (b*d - a*e)*(c*e - b*f))*(p + 1) + (b^2*(C*d + A*f) - b*(B*c*d + A*c*e +
a*C*e + a*B*f) + 2*(A*c*(c*d - a*f) - a*(c*C*d - B*c*e - a*C*f)))*(a*f*(p
+ 1) - c*d*(p + 2)) - e*((A*c - a*C)*(2*a*c*e - b*(c*d + a*f)) + (A*b - a*B
)*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)))*(p + q + 2) - (2*f*((A*c - a*C)*(2*a
*c*e - b*(c*d + a*f)) + (A*b - a*B)*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)))*(p
+ q + 2) - (b^2*(C*d + A*f) - b*(B*c*d + A*c*e + a*C*e + a*B*f) + 2*(A*c*(
c*d - a*f) - a*(c*C*d - B*c*e - a*C*f)))*(b*f*(p + 1) - c*e*(2*p + q + 4))
*x - c*f*(b^2*(C*d + A*f) - b*(B*c*d + A*c*e + a*C*e + a*B*f) + 2*(A*c*(c*d
- a*f) - a*(c*C*d - B*c*e - a*C*f)))*(2*p + 2*q + 5)*x^2, x], x]] /; F
reeQ[{a, b, c, d, e, f, q}, x] && PolyQ[Px, x, 2] && LtQ[p, -1] && NeQ[(c*d
- a*f)^2 - (b*d - a*e)*(c*e - b*f), 0] && !( !IntegerQ[p] && !LtQ[q, -1])
&& !IGtQ[q, 0]

```

Maple [A] (verified)

Time = 2.50 (sec) , antiderivative size = 452, normalized size of antiderivative = 1.20

method	result
risch	$\frac{4596238560x^{17} - 38639385552x^{16} + 188603773872x^{15} - 606785954952x^{14} + 1459208021718x^{13} - 2679143870481x^{12} + 399965613253x^{11} - 4596238560x^{10} + 38639385552x^9 - 188603773872x^8 + 606785954952x^7 - 1459208021718x^6 + 2679143870481x^5 - 399965613253x^4 + 4596238560x^3 - 38639385552x^2 + 188603773872x - 606785954952}{(x^2 - 2x + 3)^{11/2}(2x^2 + x + 1)^5}$
trager	Expression too large to display
default	Expression too large to display

input

```
int(1/(x^2-2*x+3)^(11/2)/(2*x^2+x+1)^5,x,method=_RETURNVERBOSE)
```

output

```

1/1234800000000*(4596238560*x^17-38639385552*x^16+188603773872*x^15-606785
954952*x^14+1459208021718*x^13-2679143870481*x^12+3999656132532*x^11-49157
97913008*x^10+5380603084494*x^9-5134334619701*x^8+4591320676952*x^7-335981
3871472*x^6+2503427226914*x^5-1409335257371*x^4+1002897791524*x^3-26696665
4968*x^2+261702502714*x-53205422447)/(x^2-2*x+3)^(9/2)/(2*x^2+x+1)^4+1/268
912000000000*4^(1/2)*((2^(1/2)-1+x)^2/(2^(1/2)+1-x)^2+1)^(1/2)*2^(1/2)*(96
25722625*(-6050+4280*2^(1/2))^(1/2)*arctan(1/49*(-6050+4280*2^(1/2))^(1/2)
/((2^(1/2)-1+x)^2/(2^(1/2)+1-x)^2+1)^(1/2)*(57+40*2^(1/2))*(2^(1/2)-1+x)/(
2^(1/2)+1-x))*(-350+280*2^(1/2))^(1/2)*2^(1/2)+13664181884*(-6050+4280*2^(
1/2))^(1/2)*arctan(1/49*(-6050+4280*2^(1/2))^(1/2)/((2^(1/2)-1+x)^2/(2^(1/
2)+1-x)^2+1)^(1/2)*(57+40*2^(1/2))*(2^(1/2)-1+x)/(2^(1/2)+1-x))*(-350+280*
2^(1/2))^(1/2)+456968008770*2^(1/2)*arctanh(7*((2^(1/2)-1+x)^2/(2^(1/2)+1-
x)^2+1)^(1/2)/(-350+280*2^(1/2))^(1/2))-607941010600*arctanh(7*((2^(1/2)-1
+x)^2/(2^(1/2)+1-x)^2+1)^(1/2)/(-350+280*2^(1/2))^(1/2)))/(((2^(1/2)-1+x)^
2/(2^(1/2)+1-x)^2+1)/((2^(1/2)-1+x)/(2^(1/2)+1-x)+1)^2)^(1/2)/((2^(1/2)-1+
x)/(2^(1/2)+1-x)+1)/(-350+280*2^(1/2))^(1/2)

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1014 vs. $2(298) = 596$.

Time = 0.10 (sec) , antiderivative size = 1014, normalized size of antiderivative = 2.68

$$\int \frac{1}{(3-2x+x^2)^{11/2}(1+x+2x^2)^5} dx = \text{Too large to display}$$

input

```
integrate(1/(x^2-2*x+3)^(11/2)/(2*x^2+x+1)^5,x, algorithm="fricas")
```

output

```

1/2469600000000*(9192477120*x^18 - 73539816960*x^17 + 353910369120*x^16 -
1116885970080*x^15 + 2670989133180*x^14 - 4857075098280*x^13 + 72879107667
00*x^12 - 8932789641360*x^11 + 9990499039980*x^10 - 9478592970360*x^9 + 88
80507427740*x^8 - 6269269395840*x^7 + 5282801694900*x^6 - 2524484029080*x^
5 + 2531952916740*x^4 - 227513808720*x^3 + 18*(16*x^18 - 128*x^17 + 616*x^
16 - 1944*x^15 + 4649*x^14 - 8454*x^13 + 12685*x^12 - 15548*x^11 + 17389*x
^10 - 16498*x^9 + 15457*x^8 - 10912*x^7 + 9195*x^6 - 4394*x^5 + 4407*x^4 -
396*x^3 + 1647*x^2 + 162*x + 243)*sqrt(55160237870546944/35*sqrt(2) + 302
72774247463609/14)*arctan(-5/14293820940408247*(sqrt(2)*(89801606*x - 4696
6621) + (5*sqrt(2)*(x - 1) - sqrt(x^2 - 2*x + 3))*(5*sqrt(2) + 8) + 8*x - 8
)*sqrt(55160237870546944/35*sqrt(2) - 30272774247463609/14) - sqrt(x^2 - 2
*x + 3)*(89801606*sqrt(2) - 42834985) - 42834985*x - 136768227)*sqrt(55160
237870546944/35*sqrt(2) + 30272774247463609/14)) - 18*(16*x^18 - 128*x^17
+ 616*x^16 - 1944*x^15 + 4649*x^14 - 8454*x^13 + 12685*x^12 - 15548*x^11 +
17389*x^10 - 16498*x^9 + 15457*x^8 - 10912*x^7 + 9195*x^6 - 4394*x^5 + 44
07*x^4 - 396*x^3 + 1647*x^2 + 162*x + 243)*sqrt(55160237870546944/35*sqrt(
2) + 30272774247463609/14)*arctan(5/14293820940408247*(sqrt(2)*(89801606*x
- 46966621) - (5*sqrt(2)*(x - 1) - sqrt(x^2 - 2*x + 3))*(5*sqrt(2) + 8) +
8*x - 8)*sqrt(55160237870546944/35*sqrt(2) - 30272774247463609/14) - sqrt(
x^2 - 2*x + 3)*(89801606*sqrt(2) - 42834985) - 42834985*x - 136768227)*...

```

Sympy [F]

$$\int \frac{1}{(3 - 2x + x^2)^{11/2} (1 + x + 2x^2)^5} dx = \int \frac{1}{(x^2 - 2x + 3)^{\frac{11}{2}} (2x^2 + x + 1)^5} dx$$

input

```
integrate(1/(x**2-2*x+3)**(11/2)/(2*x**2+x+1)**5,x)
```

output

```
Integral(1/((x**2 - 2*x + 3)**(11/2)*(2*x**2 + x + 1)**5), x)
```

Maxima [F]

$$\int \frac{1}{(3 - 2x + x^2)^{11/2} (1 + x + 2x^2)^5} dx = \int \frac{1}{(2x^2 + x + 1)^5 (x^2 - 2x + 3)^{\frac{11}{2}}} dx$$

input `integrate(1/(x^2-2*x+3)^(11/2)/(2*x^2+x+1)^5,x, algorithm="maxima")`

output `integrate(1/((2*x^2 + x + 1)^5*(x^2 - 2*x + 3)^(11/2)), x)`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 34.72 (sec) , antiderivative size = 2509, normalized size of antiderivative = 6.64

$$\int \frac{1}{(3 - 2x + x^2)^{11/2} (1 + x + 2x^2)^5} dx = \text{Too large to display}$$

input `integrate(1/(x^2-2*x+3)^(11/2)/(2*x^2+x+1)^5,x, algorithm="giac")`

output

```

1/19208000000000*sqrt(7722433301876572160*sqrt(2) - 10595470986612263150)*
log(3136*(2474301535988301451359142266380914651280177790712513272161012361
81293485559300330785024470114864584026604284622700*sqrt(7)*sqrt(2)*sqrt(77
22433301876572160*sqrt(2) - 10595470986612263150)*(110320475741093888*sqrt
(2) - 151363871237318045)^2 + 14433425626598425132928329887222002132467703
77915632742093923877724211999095918596245976075670043406821858326965750*sq
rt(7)*(110320475741093888*sqrt(2) - 151363871237318045)^3 + 28866851253196
85026585665977444400426493540755831265484187847755448423998191837192491952
151340086813643716653931500*sqrt(2)*(110320475741093888*sqrt(2) - 15136387
1237318045)^3 + 2061917946656917876132618555317428876066814825593761060134
17696817744571299416942320853725095720486688836903852250*sqrt(772243330187
6572160*sqrt(2) - 10595470986612263150)*(110320475741093888*sqrt(2) - 1513
63871237318045)^3 - 104913854112296962573522080729623041436733404265622592
321084093289259251415027575686933144355006438004151420024881229481000*sqrt
(7)*sqrt(2)*(110320475741093888*sqrt(2) - 151363871237318045)^2 - 10491385
41122969625735220807296230414367334042656225923210840932892592514150275756
8693314435500643800415142002488122948100*sqrt(7)*sqrt(7722433301876572160*
sqrt(2) - 10595470986612263150)*(110320475741093888*sqrt(2) - 151363871237
318045)^2 - 20982770822459392514704416145924608287346680853124518464216818
657851850283005515137386628871001287600830284004976245896200*sqrt(2)*sq...

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(3 - 2x + x^2)^{11/2} (1 + x + 2x^2)^5} dx = \int \frac{1}{(2x^2 + x + 1)^5 (x^2 - 2x + 3)^{11/2}} dx$$

input

```
int(1/((x + 2*x^2 + 1)^5*(x^2 - 2*x + 3)^(11/2)),x)
```

output

```
int(1/((x + 2*x^2 + 1)^5*(x^2 - 2*x + 3)^(11/2)), x)
```

Reduce [F]

$$\int \frac{1}{(3 - 2x + x^2)^{11/2} (1 + x + 2x^2)^5} dx = \int \frac{1}{32x^{22} - 304x^{21} + 1696x^{20} - 6360x^{19} + 18130x^{18} - 40343x^{17} -$$

input `int(1/(x^2-2*x+3)^(11/2)/(2*x^2+x+1)^5,x)`

output `int(sqrt(x**2 - 2*x + 3)/(32*x**22 - 304*x**21 + 1696*x**20 - 6360*x**19 + 18130*x**18 - 40343*x**17 + 73881*x**16 - 112604*x**15 + 150340*x**14 - 175580*x**13 + 189860*x**12 - 179940*x**11 + 164080*x**10 - 124970*x**9 + 103430*x**8 - 59524*x**7 + 49708*x**6 - 15372*x**5 + 21060*x**4 + 540*x**3 + 6318*x**2 + 729*x + 729),x)`

$$3.51 \quad \int \frac{1}{(3-2x+x^2)^{21/2} (1+x+2x^2)^{10}} dx$$

Optimal result	422
Mathematica [C] (verified)	423
Rubi [F]	424
Maple [A] (verified)	431
Fricas [B] (verification not implemented)	432
Sympy [F(-1)]	433
Maxima [F]	434
Giac [F(-1)]	434
Mupad [F(-1)]	434
Reduce [F]	435

Optimal result

Integrand size = 23, antiderivative size = 638

$$\int \frac{1}{(3-2x+x^2)^{21/2} (1+x+2x^2)^{10}} dx = \text{Too large to display}$$

output

```

1/1840124479200000000*(37358055634422583-14024622879097678*x)/(x^2-2*x+3)^
(19/2)+1/10427372048800000000*(476849951294984711-125181871472148210*x)/(
x^2-2*x+3)^(17/2)+1/1564105807320000000000*(785175837548333511+194216499
6204584234*x)/(x^2-2*x+3)^(15/2)-11/40666750990320000000000*(750232510630
8201089-7813986379726516886*x)/(x^2-2*x+3)^(13/2)-3/114701092536800000000
000*(69053268515296359011-44840736195018286006*x)/(x^2-2*x+3)^(11/2)+1/938
4634843920000000000000*(-838519439380295335657+466189390555853643870*x)/(x
^2-2*x+3)^(9/2)+1/312821161464000000000000*(-1117646664729238460189+5688
39749685437871554*x)/(x^2-2*x+3)^(7/2)+1/521368602440000000000000*(-655
1405511565449301689+3127298559983309301910*x)/(x^2-2*x+3)^(5/2)+1/10427372
04880000000000000000*(-4179039782398459850819+1886993445589652402694*x)/(x
^2-2*x+3)^(3/2)+1/630*(-1+10*x)/(x^2-2*x+3)^(19/2)/(2*x^2+x+1)^9+1/88200*(
887+2218*x)/(x^2-2*x+3)^(19/2)/(2*x^2+x+1)^8+1/1080450*(14453+29371*x)/(x^
2-2*x+3)^(19/2)/(2*x^2+x+1)^7+1/605052000*(8837931+17459234*x)/(x^2-2*x+3)
^(19/2)/(2*x^2+x+1)^6+1/26471025000*(447940041+813432205*x)/(x^2-2*x+3)^(1
9/2)/(2*x^2+x+1)^5+1/29647548000000*(592729157441+911061463974*x)/(x^2-2*x
+3)^(19/2)/(2*x^2+x+1)^4+1/12353145000000*(277010166219+310705340015*x)/(x
^2-2*x+3)^(19/2)/(2*x^2+x+1)^3+1/276710448000000*(5488221294349+1384103301
166*x)/(x^2-2*x+3)^(19/2)/(2*x^2+x+1)^2+1/2421216420000000*(-3785719779211
7-146548895467025*x)/(x^2-2*x+3)^(19/2)/(2*x^2+x+1)+1/10427372048800000...

```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 21.42 (sec) , antiderivative size = 1431, normalized size of antiderivative = 2.24

$$\int \frac{1}{(3-2x+x^2)^{21/2}(1+x+2x^2)^{10}} dx = \text{Too large to display}$$

input

```
Integrate[1/((3 - 2*x + x^2)^(21/2)*(1 + x + 2*x^2)^10), x]
```


output

```

Sqrt[3 - 2*x + x^2]*((1 - x)/(11875000000*(3 - 2*x + x^2)^10) + (265 - 113
*x)/(403750000000*(3 - 2*x + x^2)^9) + (82361 - 4841*x)/(60562500000000*(3
- 2*x + x^2)^8) + (1062937 + 1642511*x)/(15746250000000000*(3 - 2*x + x^2)
^7) + (7*(-678331 + 833371*x))/(22206250000000000*(3 - 2*x + x^2)^6) + (7*(
-73161291 + 43964675*x))/(90843750000000000*(3 - 2*x + x^2)^5) + (-1340879
383 + 430593031*x)/(1816875000000000000*(3 - 2*x + x^2)^4) - (11*(162612572
3 + 112950205*x))/(30281250000000000000*(3 - 2*x + x^2)^3) - (11*(331157064
7 + 15286717673*x))/(363375000000000000000*(3 - 2*x + x^2)^2) - (11*(-41152
1923277 + 484788625685*x))/(3633750000000000000000*(3 - 2*x + x^2)) + (2519
43 + 221770*x)/(63000000000000*(1 + x + 2*x^2)^9) - (73*(-888423 + 1604678*
x))/(8820000000000000*(1 + x + 2*x^2)^8) + (-2596903794 - 4965311863*x)/(10
8045000000000000*(1 + x + 2*x^2)^7) + (-539608494637 - 334647150510*x)/(121
01040000000000000*(1 + x + 2*x^2)^6) + (-40800462989458 + 56711874696335*x)
/(2647102500000000000000*(1 + x + 2*x^2)^5) + (42018358198215561 + 12919659
7088670934*x)/(29647548000000000000000000*(1 + x + 2*x^2)^4) + (628195598643
14747 + 169630389653846945*x)/(370594350000000000000000000*(1 + x + 2*x^2)^3)
+ (1082422109196374795 + 4797048907791526114*x)/(8301313440000000000000000
0*(1 + x + 2*x^2)^2) + (65571203144429922747 + 367152793968978953465*x)/(3
6318246300000000000000000000000000*(1 + x + 2*x^2))) + ((232442807954946745795*I +
21634177831191924841*Sqrt[7])*ArcTan[(-1350637388604350168995865589487...

```

Rubi [F]

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{1}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^{10}} dx \\
 & \quad \downarrow 1305 \\
 & -\frac{\int -\frac{20(90x^2 - 153x + 148)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^9} dx}{3150} - \frac{1 - 10x}{630 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \\
 & \quad \downarrow 27 \\
 & \frac{2}{315} \int \frac{90x^2 - 153x + 148}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^9} dx - \frac{1 - 10x}{630 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \\
 & \quad \downarrow 2135
 \end{aligned}$$

$$\begin{aligned}
& \frac{2}{315} \left(\frac{\int \frac{5(75412x^2 - 86509x + 80661)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^8} dx}{2800} + \frac{2218x + 887}{560(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^8} \right) - \\
& \quad \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \\
& \quad \downarrow 27 \\
& \frac{2}{315} \left(\frac{1}{560} \int \frac{75412x^2 - 86509x + 80661}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^8} dx + \frac{2218x + 887}{560(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^8} \right) - \\
& \quad \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \\
& \quad \downarrow 2135 \\
& \frac{2}{315} \left(\frac{1}{560} \left(\frac{\int \frac{50(3759488x^2 - 3790178x + 3715561)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^7} dx}{2450} + \frac{4(29371x + 14453)}{49(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^7} \right) + \frac{2218x + 887}{560(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^8} \right) - \\
& \quad \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \\
& \quad \downarrow 27 \\
& \frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \int \frac{3759488x^2 - 3790178x + 3715561}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^7} dx + \frac{4(29371x + 14453)}{49(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^7} \right) + \frac{2218x + 887}{560(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^8} \right) - \\
& \quad \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \\
& \quad \downarrow 2135 \\
& \frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{\int \frac{15(523777020x^2 - 494230435x + 458962907)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^6} dx}{2100} + \frac{17459234x + 8837931}{140(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^6} \right) + \frac{4(29371x + 14453)}{49(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^7} \right) + \frac{2218x + 887}{560(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^8} \right) - \\
& \quad \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \\
& \quad \downarrow 27 \\
& \frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \int \frac{523777020x^2 - 494230435x + 458962907}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^6} dx + \frac{17459234x + 8837931}{140(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^6} \right) + \frac{4(29371x + 14453)}{49(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^7} \right) + \frac{2218x + 887}{560(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^8} \right) - \\
& \quad \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9}
\end{aligned}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\int \frac{10(91104406960x^2 - 76561243634x + 63390281609)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^5} dx + \frac{4(813432205x + 447940041)}{175 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^5} \right) + \frac{1 - 10x}{630 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \right) \right) \right)$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \int \frac{91104406960x^2 - 76561243634x + 63390281609}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^5} dx + \frac{4(813432205x + 447940041)}{175 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^5} \right) + \frac{1 - 10x}{630 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \right) \right) \right)$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\int \frac{15(7895866021108x^2 - 5294487996061x + 3622330118837)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^4} dx + \frac{911061463974x + 59272915}{280 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^5} \right) + \frac{1 - 10x}{630 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \right) \right) \right) \right)$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \int \frac{7895866021108x^2 - 5294487996061x + 3622330118837}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^4} dx + \frac{911061463974x + 59272915}{280 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^5} \right) + \frac{1 - 10x}{630 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \right) \right) \right) \right)$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\int \frac{1050(5965542528288x^2 - 2041006971986x + 660555973049)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^3} dx + \frac{4(310705340015x + 27}{5 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^5} \right) + \frac{1 - 10x}{630 (x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9} \right) \right) \right) \right) \right)$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\int \frac{5965542528288x^2 - 2041006971986x + 660555973049}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^3} dx + \frac{4(31070534001)}{5(x^2 - 2x + 3)} \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{700} \int -\frac{25(-30450272625652x^2 - 90242403939711x + 57003619484663)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^2} dx \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(-\frac{1}{28} \int \frac{-30450272625652x^2 - 90242403939711x + 57003619484663}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)^2} dx + \frac{4(3)}{5} \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(-\frac{1}{350} \int -\frac{10(-11723911637362000x^2 + 9423200395626322x + 2186320)}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)} \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \int \frac{-11723911637362000x^2 + 9423200395626322x + 21863207223365}{(x^2 - 2x + 3)^{21/2} (2x^2 + x + 1)} \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\int \frac{60(168295474549172136x^2 - 211409077626196062x + 28036472352531697)}{(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)} dx + \frac{1}{3800} \right) \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2} (2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \int \frac{16829547454}{\dots} \right. \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\int \frac{20(400581988)}{\dots} \right. \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \int \frac{400581}{\dots} \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\int \frac{20(-5)}{\dots} \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\frac{1}{150} \int \dots \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \right. \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\frac{1}{150} \left(\frac{3}{2} \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\frac{1}{150} \left(\frac{3}{2} \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\frac{1}{150} \left(\frac{3}{2} \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\frac{1}{150} \left(\frac{3}{2} \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 2135

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\frac{1}{150} \left(\frac{3}{2} \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

↓ 27

$$\frac{2}{315} \left(\frac{1}{560} \left(\frac{1}{49} \left(\frac{1}{140} \left(\frac{1}{175} \left(\frac{3}{280} \left(\frac{1}{28} \left(\frac{1}{35} \left(\frac{37358055634422583 - 14024622879097678x}{190(x^2 - 2x + 3)^{19/2}} - \frac{3}{190} \left(\frac{1}{170} \left(\frac{1}{150} \left(\frac{3}{2} \right) \right) \right) \right) \right) \right) \right) \right) \right) \right) \frac{1 - 10x}{630(x^2 - 2x + 3)^{19/2}(2x^2 + x + 1)^9}$$

input `Int[1/((3 - 2*x + x^2)^(21/2)*(1 + x + 2*x^2)^10),x]`

output `$Aborted`

Defintions of rubi rules used

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_) /; FreeQ[b, x]]`

rule 1305 `Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)*((d_.) + (e_.)*(x_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[(2*a*c^2*e - b^2*c*e + b^3*f + b*c*(c*d - 3*a*f) + c*(2*c^2*d + b^2*f - c*(b*e + 2*a*f))*x*(a + b*x + c*x^2)^(p + 1)*((d + e*x + f*x^2)^(q + 1)/((b^2 - 4*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1))), x] - Simp[1/((b^2 - 4*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1)) Int[(a + b*x + c*x^2)^(p + 1)*(d + e*x + f*x^2)^q*Simp[2*c*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1) - (2*c^2*d + b^2*f - c*(b*e + 2*a*f))*(a*f*(p + 1) - c*d*(p + 2)) - e*(b^2*c*e - 2*a*c^2*e - b^3*f - b*c*(c*d - 3*a*f))*(p + q + 2) + (2*f*(2*a*c^2*e - b^2*c*e + b^3*f + b*c*(c*d - 3*a*f))*(p + q + 2) - (2*c^2*d + b^2*f - c*(b*e + 2*a*f))*(b*f*(p + 1) - c*e*(2*p + q + 4)))*x + c*f*(2*c^2*d + b^2*f - c*(b*e + 2*a*f))*(2*p + 2*q + 5)*x^2, x], x] /; FreeQ[{a, b, c, d, e, f, q}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[e^2 - 4*d*f, 0] && LtQ[p, -1] && NeQ[(c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f), 0] && !(IntegerQ[p] && ILtQ[q, -1]) && !IGtQ[q, 0]`

rule 2135

```

Int[(Px_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_)*((d_) + (e_)*(x_) + (f_
)*(x_)^2)^(q_), x_Symbol] :> With[{A = Coeff[Px, x, 0], B = Coeff[Px, x, 1]
, C = Coeff[Px, x, 2]}, Simp[(a + b*x + c*x^2)^(p + 1)*((d + e*x + f*x^2)^(
q + 1)/((b^2 - 4*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1))*(
(A*c - a*C)*(2*a*c*e - b*(c*d + a*f)) + (A*b - a*B)*(2*c^2*d + b^2*f - c*(b
*e + 2*a*f)) + c*(A*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)) - B*(b*c*d - 2*a*c*
e + a*b*f) + C*(b^2*d - a*b*e - 2*a*(c*d - a*f)))*x), x] + Simp[1/((b^2 - 4
*a*c)*((c*d - a*f)^2 - (b*d - a*e)*(c*e - b*f))*(p + 1)) Int[(a + b*x + c
*x^2)^(p + 1)*(d + e*x + f*x^2)^q*Simp[(b*B - 2*A*c - 2*a*C)*((c*d - a*f)^2
- (b*d - a*e)*(c*e - b*f))*(p + 1) + (b^2*(C*d + A*f) - b*(B*c*d + A*c*e +
a*C*e + a*B*f) + 2*(A*c*(c*d - a*f) - a*(c*C*d - B*c*e - a*C*f)))*(a*f*(p
+ 1) - c*d*(p + 2)) - e*((A*c - a*C)*(2*a*c*e - b*(c*d + a*f)) + (A*b - a*B
)*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)))*(p + q + 2) - (2*f*((A*c - a*C)*(2*a
*c*e - b*(c*d + a*f)) + (A*b - a*B)*(2*c^2*d + b^2*f - c*(b*e + 2*a*f)))*(p
+ q + 2) - (b^2*(C*d + A*f) - b*(B*c*d + A*c*e + a*C*e + a*B*f) + 2*(A*c*(
c*d - a*f) - a*(c*C*d - B*c*e - a*C*f)))*(b*f*(p + 1) - c*e*(2*p + q + 4))
*x - c*f*(b^2*(C*d + A*f) - b*(B*c*d + A*c*e + a*C*e + a*B*f) + 2*(A*c*(c*d
- a*f) - a*(c*C*d - B*c*e - a*C*f)))*(2*p + 2*q + 5)*x^2, x], x]] /; F
reeQ[{a, b, c, d, e, f, q}, x] && PolyQ[Px, x, 2] && LtQ[p, -1] && NeQ[(c*d
- a*f)^2 - (b*d - a*e)*(c*e - b*f), 0] && !( !IntegerQ[p] && !LtQ[q, -1])
&& !IGtQ[q, 0]

```

Maple [A] (verified)

Time = 5.68 (sec) , antiderivative size = 552, normalized size of antiderivative = 0.87

method	result
risch	$\frac{3372249001933422237824271360x^{37} - 53502205399640031394796147712x^{36} + 469149394082989701729494575872x^{35} - 2847499220...}{...}$
trager	Expression too large to display
default	Expression too large to display

input

```
int(1/(x^2-2*x+3)^(21/2)/(2*x^2+x+1)^10,x,method=_RETURNVERBOSE)
```


output

```

1/13420027826805600000000000000000*(3372249001933422237824271360*x^37-5350
2205399640031394796147712*x^36+469149394082989701729494575872*x^35-2847499
220912667753383035299072*x^34+13254252261100740556512388253568*x^33-497700
80058525077628064229832576*x^32+156010734937008739388220889457760*x^31-417
516398850754397130111919794336*x^30+971538171913365251873706873353652*x^29
-1993653213575521837888601204380228*x^28+365555347185295760625734541414003
1*x^27-6054769996581738503753686155104785*x^26+915549415851386923027152974
6307221*x^25-12740106677685048178693605103009787*x^24+16442770202470076313
197215936814318*x^23-19772569734288744720189854470201506*x^22+222864376176
21909921609206629636086*x^21-23584986647560742443188031208946882*x^20+2357
9397211179175240196614296051673*x^19-22218747553941794885903840542461607*x
^18+19912295454080246583636391613811979*x^17-16801760806053390242995145349
148613*x^16+13613407965006475288139078599341572*x^15-102793056507331786692
23634020962076*x^14+7606288378303449524327938977040824*x^13-50698382349927
51929471190426115248*x^12+3507425970596197680016078213030977*x^11-19748144
83061344405275851094534735*x^10+1357002388430055881833293557852283*x^9-566
969010759169461615951049236597*x^8+458426000073846882432457044306894*x^7-9
4704557665253489332536549937026*x^6+135183920426913231415208872303230*x^5-
1023095318901774638403186272874*x^4+29398041153524973343917601742151*x^3+1
933957195570062708781629134823*x^2+3397462350398947848063583843461*x-80...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1714 vs. $2(518) = 1036$.

Time = 0.18 (sec) , antiderivative size = 1714, normalized size of antiderivative = 2.69

$$\int \frac{1}{(3-2x+x^2)^{21/2}(1+x+2x^2)^{10}} dx = \text{Too large to display}$$

input

```
integrate(1/(x^2-2*x+3)^(21/2)/(2*x^2+x+1)^10,x, algorithm="fricas")
```

output

```

1/26840055653611200000000000000000*(6744498003866844475648542720*x^38 - 10
4539719059936089372552412160*x^37 + 900390483516223737499080453120*x^36 -
5371992660079941624854064276480*x^35 + 24606457904857698713844252046080*x^
34 - 90963466109277373120851623698560*x^33 + 28089653898954730543292941933
8240*x^32 - 740971416495510997144845440898240*x^31 + 170108003010485593246
3967828449800*x^30 - 3447426219612332424711322428901980*x^29 + 62518630215
92248151838414654875460*x^28 - 10256627551906425809034568826985240*x^27 +
15388905077240219856331650258935040*x^26 - 2127848738855476045716060344160
5700*x^25 + 27334008812742537113204165617656780*x^24 - 3274901989653683403
3895800624267120*x^23 + 36839244026922246796455050563687320*x^22 - 3893137
5096491957607948047018948700*x^21 + 38950602632818864204958532927160740*x^
20 - 36715802061835614047293656120490920*x^19 + 33031150678661088971867773
726923120*x^18 - 27904629463817966302108782777675300*x^17 + 22806781599657
490046875160302230540*x^16 - 17210816451136000844652573117508320*x^15 + 12
969937518120337685296303583114040*x^14 - 855890766495650026437468870542130
0*x^13 + 6174005722191899041430283362811180*x^12 - 33249299464323317158463
17213304040*x^11 + 2534236799713243288588790298370080*x^10 - 9006943415932
84668166111283071020*x^9 + 952602184923756094454184807020100*x^8 - 8685812
1789158018014357835347440*x^7 + 331412539235861366902063441287720*x^6 + 44
740404706662877172620981695660*x^5 + 88530111824581469125218465390060*x...

```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(3 - 2x + x^2)^{21/2} (1 + x + 2x^2)^{10}} dx = \text{Timed out}$$

input

```
integrate(1/(x**2-2*x+3)**(21/2)/(2*x**2+x+1)**10,x)
```

output

Timed out

Maxima [F]

$$\int \frac{1}{(3 - 2x + x^2)^{21/2} (1 + x + 2x^2)^{10}} dx = \int \frac{1}{(2x^2 + x + 1)^{10} (x^2 - 2x + 3)^{\frac{21}{2}}} dx$$

input `integrate(1/(x^2-2*x+3)^(21/2)/(2*x^2+x+1)^10,x, algorithm="maxima")`

output `integrate(1/((2*x^2 + x + 1)^10*(x^2 - 2*x + 3)^(21/2)), x)`

Giac [F(-1)]

Timed out.

$$\int \frac{1}{(3 - 2x + x^2)^{21/2} (1 + x + 2x^2)^{10}} dx = \text{Timed out}$$

input `integrate(1/(x^2-2*x+3)^(21/2)/(2*x^2+x+1)^10,x, algorithm="giac")`

output `Timed out`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(3 - 2x + x^2)^{21/2} (1 + x + 2x^2)^{10}} dx = \int \frac{1}{(2x^2 + x + 1)^{10} (x^2 - 2x + 3)^{21/2}} dx$$

input `int(1/((x + 2*x^2 + 1)^10*(x^2 - 2*x + 3)^(21/2)),x)`

output `int(1/((x + 2*x^2 + 1)^10*(x^2 - 2*x + 3)^(21/2)), x)`

Reduce [F]

$$\int \frac{1}{(3 - 2x + x^2)^{21/2} (1 + x + 2x^2)^{10}} dx = \int \frac{1}{1024x^{42} - 17408x^{41} + 163072x^{40} - 1059840x^{39} + 5294720x^{38} - 21409152x^{37} + 72501024x^{36} - 210353856x^{35} + 532338420x^{34} - 1191461700x^{33} + 2387723889x^{32} - 4329193728x^{31} + 7171025142x^{30} - 10938959220x^{29} + 15482374950x^{28} - 20441554296x^{27} + 25316000622x^{26} - 29502127788x^{25} + 32502050290x^{24} - 33883949360x^{23} + 33595551166x^{22} - 31604802132x^{21} + 28438741598x^{20} - 24253023240x^{19} + 19949752830x^{18} - 15409638228x^{17} + 11697087396x^{16} - 8094599584x^{15} + 5789499650x^{14} - 3456463420x^{13} + 2465340786x^{12} - 1136849672x^{11} + 940451658x^{10} - 234461700x^9 + 339398910x^8 + 4848336x^7 + 110547018x^6 + 20741508x^5 + 26309610x^4 + 5511240x^3 + 3503574x^2 + 472392x + 177147), x)$$

input `int(1/(x^2-2*x+3)^(21/2)/(2*x^2+x+1)^10,x)`

output

```
int(sqrt(x**2 - 2*x + 3)/(1024*x**42 - 17408*x**41 + 163072*x**40 - 1059840*x**39 + 5294720*x**38 - 21409152*x**37 + 72501024*x**36 - 210353856*x**35 + 532338420*x**34 - 1191461700*x**33 + 2387723889*x**32 - 4329193728*x**31 + 7171025142*x**30 - 10938959220*x**29 + 15482374950*x**28 - 20441554296*x**27 + 25316000622*x**26 - 29502127788*x**25 + 32502050290*x**24 - 33883949360*x**23 + 33595551166*x**22 - 31604802132*x**21 + 28438741598*x**20 - 24253023240*x**19 + 19949752830*x**18 - 15409638228*x**17 + 11697087396*x**16 - 8094599584*x**15 + 5789499650*x**14 - 3456463420*x**13 + 2465340786*x**12 - 1136849672*x**11 + 940451658*x**10 - 234461700*x**9 + 339398910*x**8 + 4848336*x**7 + 110547018*x**6 + 20741508*x**5 + 26309610*x**4 + 5511240*x**3 + 3503574*x**2 + 472392*x + 177147),x)
```

$$3.52 \quad \int \frac{-a - \sqrt{1+a^2} + x}{(-a + \sqrt{1+a^2} + x) \sqrt{(-a+x)(1+x^2)}} dx$$

Optimal result	436
Mathematica [A] (verified)	436
Rubi [C] (verified)	437
Maple [C] (verified)	441
Fricas [A] (verification not implemented)	442
Sympy [F(-1)]	442
Maxima [F]	443
Giac [F]	443
Mupad [F(-1)]	444
Reduce [F]	444

Optimal result

Integrand size = 48, antiderivative size = 66

$$\int \frac{-a - \sqrt{1+a^2} + x}{(-a + \sqrt{1+a^2} + x) \sqrt{(-a+x)(1+x^2)}} dx$$

$$= -\sqrt{2} \sqrt{a + \sqrt{1+a^2}} \arctan \left(\frac{\sqrt{2} \sqrt{-a + \sqrt{1+a^2}} (-a+x)}{\sqrt{(-a+x)(1+x^2)}} \right)$$

output

```
-arctan((-a+x)*2^(1/2)*(-a+(a^2+1)^(1/2))^(1/2)/((-a+x)*(x^2+1))^(1/2))*2^(1/2)*(a+(a^2+1)^(1/2))^(1/2)
```

Mathematica [A] (verified)

Time = 1.31 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.50

$$\int \frac{-a - \sqrt{1+a^2} + x}{(-a + \sqrt{1+a^2} + x) \sqrt{(-a+x)(1+x^2)}} dx$$

$$= -\frac{\sqrt{2} \sqrt{-a + \sqrt{1+a^2}} \sqrt{1+x^2} \arctan \left(\frac{\sqrt{2} \sqrt{-a + \sqrt{1+a^2}} \sqrt{-a+x}}{\sqrt{1+x^2}} \right)}{\sqrt{-a + \sqrt{1+a^2}} \sqrt{(-a+x)(1+x^2)}}$$

input

```
Integrate[(-a - Sqrt[1 + a^2] + x)/((-a + Sqrt[1 + a^2] + x)*Sqrt[(-a + x)
*(1 + x^2)]),x]
```

output

```
-((Sqrt[2]*Sqrt[-a + x]*Sqrt[1 + x^2]*ArcTan[(Sqrt[2]*Sqrt[-a + Sqrt[1 + a
^2]]*Sqrt[-a + x])/Sqrt[1 + x^2]])/(Sqrt[-a + Sqrt[1 + a^2]]*Sqrt[(-a + x)
*(1 + x^2)]))
```

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 1.20 (sec) , antiderivative size = 429, normalized size of antiderivative = 6.50, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {7270, 2349, 510, 729, 25, 1416, 1534, 1416, 2212, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{-\sqrt{a^2+1}-a+x}{(\sqrt{a^2+1}-a+x)\sqrt{(x^2+1)(x-a)}} dx$$

$$\downarrow 7270$$

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \int \frac{a-x+\sqrt{a^2+1}}{(a-x-\sqrt{a^2+1})\sqrt{x-a}\sqrt{x^2+1}} dx}{\sqrt{-((x^2+1)(a-x))}}$$

$$\downarrow 2349$$

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \left(2\sqrt{a^2+1} \int \frac{1}{(a-x-\sqrt{a^2+1})\sqrt{x-a}\sqrt{x^2+1}} dx + \int \frac{1}{\sqrt{x-a}\sqrt{x^2+1}} dx \right)}{\sqrt{-((x^2+1)(a-x))}}$$

$$\downarrow 510$$

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \left(2\sqrt{a^2+1} \int \frac{1}{(a-x-\sqrt{a^2+1})\sqrt{x-a}\sqrt{x^2+1}} dx + 2 \int \frac{1}{\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a} \right)}{\sqrt{-((x^2+1)(a-x))}}$$

$$\downarrow 729$$

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \left(2 \int \frac{1}{\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a} + 4\sqrt{a^2+1} \int \frac{1}{(-a+x+\sqrt{a^2+1})\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a} \right)}{\sqrt{-((x^2+1)(a-x))}}$$

↓ 25

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \left(2 \int \frac{1}{\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a} - 4\sqrt{a^2+1} \int \frac{1}{(-a+x+\sqrt{a^2+1})\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a} \right)}{\sqrt{-((x^2+1)(a-x))}}$$

↓ 1416

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \left(\frac{\sqrt[4]{a^2+1} \left(\frac{x-a}{\sqrt{a^2+1}} + 1 \right) \sqrt{\frac{a^2+2a(x-a)+(x-a)^2+1}{(a^2+1) \left(\frac{x-a}{\sqrt{a^2+1}} + 1 \right)^2}} \text{EllipticF} \left(2 \arctan \left(\frac{\sqrt{x-a}}{\sqrt[4]{a^2+1}} \right), \frac{1}{2} \left(1 - \frac{a}{\sqrt{a^2+1}} \right) \right)}{\sqrt{a^2+2a(x-a)+(x-a)^2+1}} - 4\sqrt{a^2+1} \right)}{\sqrt{-((x^2+1)(a-x))}}$$

↓ 1534

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \left(4\sqrt{a^2+1} \left(- \frac{\int \frac{1}{\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a}}{2\sqrt{a^2+1}} - \frac{\int \frac{a-x+\sqrt{a^2+1}}{(-a+x+\sqrt{a^2+1})\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a}}{2\sqrt{a^2+1}} \right) + \frac{\sqrt[4]{a^2+1}}{\sqrt{-((x^2+1)(a-x))}} \right)}{\sqrt{-((x^2+1)(a-x))}}$$

↓ 1416

$$\frac{\sqrt{x^2+1}\sqrt{x-a} \left(4\sqrt{a^2+1} \left(- \frac{\int \frac{a-x+\sqrt{a^2+1}}{(-a+x+\sqrt{a^2+1})\sqrt{a^2+2(x-a)a+(x-a)^2+1}} d\sqrt{x-a}}{2\sqrt{a^2+1}} - \frac{\left(\frac{x-a}{\sqrt{a^2+1}} + 1 \right) \sqrt{\frac{a^2+2a(x-a)+(x-a)^2+1}{(a^2+1) \left(\frac{x-a}{\sqrt{a^2+1}} + 1 \right)^2}} \text{EllipticF} \left(2 \arctan \left(\frac{\sqrt{x-a}}{\sqrt[4]{a^2+1}} \right), \frac{1}{2} \left(1 - \frac{a}{\sqrt{a^2+1}} \right) \right)}{4\sqrt[4]{a^2+1}\sqrt{a^2+2a(x-a)+(x-a)^2+1}} \right) + \frac{\sqrt[4]{a^2+1}}{\sqrt{-((x^2+1)(a-x))}} \right)}{\sqrt{-((x^2+1)(a-x))}}$$

↓ 2212

$$\sqrt{x^2+1}\sqrt{x-a} \left(4\sqrt{a^2+1} \left(-\frac{1}{2} \int \frac{1}{\sqrt{a^2+1}-2\sqrt{a^2+1}(a-\sqrt{a^2+1})(x-a)} d \frac{\sqrt{x-a}}{\sqrt{a^2+2(x-a)a+(x-a)^2+1}} - \frac{\left(\frac{x-a}{\sqrt{a^2+1}}+1\right) \sqrt{\frac{a^2+2a(x-a)}{(a^2+1)\left(\frac{x-a}{\sqrt{a^2+1}}+1\right)}}}{4} \right) \right)$$

↓ 221

$$\sqrt{x^2+1}\sqrt{x-a} \left(4\sqrt{a^2+1} \left(-\frac{\left(\frac{x-a}{\sqrt{a^2+1}}+1\right) \sqrt{\frac{a^2+2a(x-a)+(x-a)^2+1}{(a^2+1)\left(\frac{x-a}{\sqrt{a^2+1}}+1\right)^2}} \operatorname{EllipticF}\left(2 \arctan\left(\frac{\sqrt{x-a}}{\sqrt{a^2+1}}\right), \frac{1}{2}\left(1-\frac{a}{\sqrt{a^2+1}}\right)\right)}{4 \sqrt{a^2+1} \sqrt{a^2+2a(x-a)+(x-a)^2+1}} - \frac{\arctan\left(\frac{\sqrt{x-a}}{\sqrt{a^2+1}}\right)}{\sqrt{-(x^2+1)}}$$

input

```
Int[(-a - Sqrt[1 + a^2] + x)/((-a + Sqrt[1 + a^2] + x)*Sqrt[(-a + x)*(1 + x^2)]), x]
```

output

```
(Sqrt[-a + x]*Sqrt[1 + x^2]*(((1 + a^2)^(1/4)*(1 + (-a + x)/Sqrt[1 + a^2]) *Sqrt[(1 + a^2 + 2*a*(-a + x) + (-a + x)^2]/((1 + a^2)*(1 + (-a + x)/Sqrt[1 + a^2])^2)]*EllipticF[2*ArcTan[Sqrt[-a + x]/(1 + a^2)^(1/4)], (1 - a/Sqrt[1 + a^2])/2])/Sqrt[1 + a^2 + 2*a*(-a + x) + (-a + x)^2] + 4*Sqrt[1 + a^2]*(-1/2*ArcTanh[(Sqrt[2]*Sqrt[a - Sqrt[1 + a^2]])*Sqrt[-a + x])/Sqrt[1 + a^2 + 2*a*(-a + x) + (-a + x)^2])/(Sqrt[2]*Sqrt[1 + a^2]*Sqrt[a - Sqrt[1 + a^2]]) - ((1 + (-a + x)/Sqrt[1 + a^2])*Sqrt[(1 + a^2 + 2*a*(-a + x) + (-a + x)^2]/((1 + a^2)*(1 + (-a + x)/Sqrt[1 + a^2])^2)]*EllipticF[2*ArcTan[Sqrt[-a + x]/(1 + a^2)^(1/4)], (1 - a/Sqrt[1 + a^2])/2])/(4*(1 + a^2)^(1/4)*Sqrt[1 + a^2 + 2*a*(-a + x) + (-a + x)^2])))/Sqrt[-((a - x)*(1 + x^2))]
```

Defintions of rubi rules used

rule 25

```
Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]
```

rule 221

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```


rule 510 $\text{Int}[1/(\text{Sqrt}[(c_)+(d_)(x_)]*\text{Sqrt}[(a_)+(b_)(x_)^2]), x_Symbol] \rightarrow \text{Simp}[2/d \text{ Subst}[\text{Int}[1/\text{Sqrt}[(b*c^2+a*d^2)/d^2-2*b*c*(x^2/d^2)+b*(x^4/d^2)], x], x, \text{Sqrt}[c+d*x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{PosQ}[b/a]$

rule 729 $\text{Int}[1/(\text{Sqrt}[(c_)+(d_)(x_)]*((e_)+(f_)(x_))*\text{Sqrt}[(a_)+(b_)(x_)^2]), x_Symbol] \rightarrow \text{Simp}[2 \text{ Subst}[\text{Int}[1/((d*e-c*f+f*x^2)*\text{Sqrt}[(b*c^2+a*d^2)/d^2-2*b*c*(x^2/d^2)+b*(x^4/d^2)]), x], x, \text{Sqrt}[c+d*x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{PosQ}[b/a]$

rule 1416 $\text{Int}[1/\text{Sqrt}[(a_)+(b_)(x_)^2+(c_)(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(1+q^2*x^2)*(\text{Sqrt}[(a+b*x^2+c*x^4)/(a*(1+q^2*x^2)^2])/(2*q*\text{Sqrt}[a+b*x^2+c*x^4]))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2-b*(q^2/(4*c))], x]] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2-4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

rule 1534 $\text{Int}[1/(((d_)+(e_)(x_)^2)*\text{Sqrt}[(a_)+(b_)(x_)^2+(c_)(x_)^4]), x_Symbol] \rightarrow \text{Simp}[1/(2*d) \text{ Int}[1/\text{Sqrt}[a+b*x^2+c*x^4], x], x] + \text{Simp}[1/(2*d) \text{ Int}[(d-e*x^2)/((d+e*x^2)*\text{Sqrt}[a+b*x^2+c*x^4]), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2-4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2-b*d*e+a*e^2, 0] \ \&\& \ \text{EqQ}[c*d^2-a*e^2, 0]$

rule 2212 $\text{Int}[((A_)+(B_)(x_)^2)/(((d_)+(e_)(x_)^2)*\text{Sqrt}[(a_)+(b_)(x_)^2+(c_)(x_)^4]), x_Symbol] \rightarrow \text{Simp}[A \text{ Subst}[\text{Int}[1/(d-(b*d-2*a*e)*x^2), x], x, x/\text{Sqrt}[a+b*x^2+c*x^4]], x] /; \text{FreeQ}[\{a, b, c, d, e, A, B\}, x] \ \&\& \ \text{EqQ}[c*d^2-a*e^2, 0] \ \&\& \ \text{EqQ}[B*d+A*e, 0]$

rule 2349 $\text{Int}[(P_x)*((c_)+(d_)(x_))^(m_)*((e_)+(f_)(x_))^(n_)*((a_)+(b_)(x_)^2)^(p_), x_Symbol] \rightarrow \text{Int}[\text{PolynomialQuotient}[P_x, c+d*x, x]*(c+d*x)^(m+1)*(e+f*x)^n*(a+b*x^2)^p, x] + \text{Simp}[\text{PolynomialRemainder}[P_x, c+d*x, x] \text{ Int}[(c+d*x)^m*(e+f*x)^n*(a+b*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{PolynomialQ}[P_x, x] \ \&\& \ \text{LtQ}[m, 0] \ \&\& \ !\text{IntegerQ}[n] \ \&\& \ \text{IntegersQ}[2*m, 2*n, 2*p]$

rule 7270

```
Int[(u_.)*((a_.)*(v_)^(m_.)*(w_)^(n_.))^(p_), x_Symbol] := Simp[a^IntPart[p]
]*((a*v^m*w^n)^FracPart[p]/(v^(m*FracPart[p])*w^(n*FracPart[p]))) Int[u*v
^(m*p)*w^(n*p), x], x] /; FreeQ[{a, m, n, p}, x] && !IntegerQ[p] && !Free
Q[v, x] && !FreeQ[w, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 1.84 (sec) , antiderivative size = 787, normalized size of antiderivative = 11.92

method	result
default	$\frac{2i\sqrt{-i(x+i)}\sqrt{\frac{-a+x}{-i-a}}\sqrt{i(x-i)}\operatorname{EllipticF}\left(\frac{\sqrt{2}\sqrt{-i(x+i)}}{2},\sqrt{2}\sqrt{\frac{-i}{-i-a}}\right)}{\sqrt{-ax^2+x^3-ax}} - \frac{2\sqrt{a^2+1}(2ax-x^2+1)\sqrt{-(a-x)(x^2+1)(a^2+1)}}{\left(\frac{i\sqrt{-}}{\dots}\right)}$
elliptic	Expression too large to display

input

```
int((-a+x-(a^2+1)^(1/2))/(-a+x+(a^2+1)^(1/2))/((-a+x)*(x^2+1))^(1/2),x,met
hod=_RETURNVERBOSE)
```

output

```
2*I*(-I*(x+I))^(1/2)*((-a+x)/(-I-a))^(1/2)*(I*(x-I))^(1/2)/(-a*x^2+x^3-a*x
)^(1/2)*EllipticF(1/2*2^(1/2)*(-I*(x+I))^(1/2),2^(1/2)*(-I/(-I-a))^(1/2))-
2*(a^2+1)^(1/2)*(2*a*x-x^2+1)*(-(a-x)*(x^2+1)*(a^2+1))^(1/2)/(-a+x+(a^2+1)
^(1/2))/((-a-x)*(x^2+1))^(1/2)*a^2+(-(a-x)*(x^2+1)*(a^2+1))^(1/2)*a-((a-
x)*(x^2+1)*(a^2+1))^(1/2)*x+(-(a-x)*(x^2+1))^(1/2)*(I*(1-I*x))^(1/2)*(-1/(
-I-a)*a+1/(-I-a)*x)^(1/2)*(I*x+1)^(1/2)/(-a*x^2+x^3-a*x)^(1/2)/(-I-a-(a^2+
1)^(1/2))*EllipticPi(1/2*2^(1/2)*(-I*(x+I))^(1/2),-2*I/(-I-a-(a^2+1)^(1/2)
),2^(1/2)*(-I/(-I-a))^(1/2))+I*(1-I*x)^(1/2)*(-1/(-I-a)*a+1/(-I-a)*x)^(1/2
)*(I*x+1)^(1/2)/(-a*x^2+x^3-a*x)^(1/2)/(-I-a+(a^2+1)^(1/2))*EllipticPi(1/2
*2^(1/2)*(-I*(x+I))^(1/2),-2*I/(-I-a+(a^2+1)^(1/2)),2^(1/2)*(-I/(-I-a))^(1
/2))-I*(a^2+1)^(1/2)*(1-I*x)^(1/2)*(-1/(-I-a)*a+1/(-I-a)*x)^(1/2)*(I*x+1)^(
1/2)/(-a^3*x^2+a^2*x^3-a^3+a^2*x-a*x^2+x^3-a*x)^(1/2)/(-I-a-(a^2+1)^(1/2)
)*EllipticPi(1/2*2^(1/2)*(-I*(x+I))^(1/2),-2*I/(-I-a-(a^2+1)^(1/2)),2^(1/2
)*(-I/(-I-a))^(1/2))+I*(a^2+1)^(1/2)*(1-I*x)^(1/2)*(-1/(-I-a)*a+1/(-I-a)*x
)^(1/2)*(I*x+1)^(1/2)/(-a^3*x^2+a^2*x^3-a^3+a^2*x-a*x^2+x^3-a*x)^(1/2)/(-I
-a+(a^2+1)^(1/2))*EllipticPi(1/2*2^(1/2)*(-I*(x+I))^(1/2),-2*I/(-I-a+(a^2+
1)^(1/2)),2^(1/2)*(-I/(-I-a))^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 546, normalized size of antiderivative = 8.27

$$\int \frac{-a - \sqrt{1 + a^2} + x}{(-a + \sqrt{1 + a^2} + x) \sqrt{(-a + x)(1 + x^2)}} dx$$

$$= \left[\frac{1}{4} \sqrt{-2a - 2\sqrt{a^2 + 1}} \log \left(-\frac{8ax^7 + x^8 + 4(2a^2 + 15)x^6 - 8(4a^3 + 15a)x^5 + 2(8a^4 + 80a^2 + 67)x^4}{4(ax^2 - x^3 + a - x)} \right) \right.$$

$$\left. - \frac{1}{2} \sqrt{2a + 2\sqrt{a^2 + 1}} \arctan \left(-\frac{\sqrt{-ax^2 + x^3 - a + x}(2a^2 - 2ax - x^2 - 2\sqrt{a^2 + 1}(a - x) - 1)\sqrt{2a}}{4(ax^2 - x^3 + a - x)} \right) \right]$$

input `integrate((-a+x-(a^2+1)^(1/2))/(-a+x+(a^2+1)^(1/2))/((-a+x)*(x^2+1))^(1/2), x, algorithm="fricas")`

output `[1/4*sqrt(-2*a - 2*sqrt(a^2 + 1))*log(-(8*a*x^7 + x^8 + 4*(2*a^2 + 15)*x^6 - 8*(4*a^3 + 15*a)*x^5 + 2*(8*a^4 + 80*a^2 + 67)*x^4 + 64*a^4 - 8*(20*a^3 + 37*a)*x^3 + 4*(16*a^4 + 74*a^2 + 15)*x^2 + 48*a^2 - 4*(a*x^6 + 2*(2*a^2 + 3)*x^5 - (4*a^3 - a)*x^4 - 8*a^3 - (4*a^3 + 29*a)*x^2 + 20*x^3 + 2*(10*a^2 + 3)*x - (4*a*x^5 + x^6 - (4*a^2 - 15)*x^4 - 16*a*x^3 + (4*a^2 + 15)*x^2 + 8*a^2 - 20*a*x + 1)*sqrt(a^2 + 1) - 5*a)*sqrt(-a*x^2 + x^3 - a + x)*sqrt(-2*a - 2*sqrt(a^2 + 1)) - 8*(24*a^3 + 13*a)*x + 16*(a*x^6 - x^7 + 15*a*x^4 - 7*x^5 - (12*a^2 + 7)*x^3 + 4*a^3 + (4*a^3 + 15*a)*x^2 - (12*a^2 + 1)*x + a)*sqrt(a^2 + 1) + 1)/(8*a*x^7 - x^8 - 4*(6*a^2 - 1)*x^6 + 8*(4*a^3 - 3*a)*x^5 - 2*(8*a^4 - 24*a^2 + 3)*x^4 - 8*(4*a^3 - 3*a)*x^3 - 4*(6*a^2 - 1)*x^2 - 8*a*x - 1), -1/2*sqrt(2*a + 2*sqrt(a^2 + 1))*arctan(-1/4*sqrt(-a*x^2 + x^3 - a + x)*(2*a^2 - 2*a*x - x^2 - 2*sqrt(a^2 + 1)*(a - x) - 1)*sqrt(2*a + 2*sqrt(a^2 + 1))/(a*x^2 - x^3 + a - x))]`

Sympy [F(-1)]

Timed out.

$$\int \frac{-a - \sqrt{1 + a^2} + x}{(-a + \sqrt{1 + a^2} + x) \sqrt{(-a + x)(1 + x^2)}} dx = \text{Timed out}$$

input `integrate((-a+x-(a**2+1)**(1/2))/(-a+x+(a**2+1)**(1/2))/((-a+x)*(x**2+1))** (1/2), x)`

output Timed out

Maxima [F]

$$\int \frac{-a - \sqrt{1 + a^2} + x}{(-a + \sqrt{1 + a^2} + x) \sqrt{(-a + x)(1 + x^2)}} dx$$

$$= \int \frac{a - x + \sqrt{a^2 + 1}}{\sqrt{-(x^2 + 1)(a - x)(a - x - \sqrt{a^2 + 1})}} dx$$

input `integrate((-a+x-(a^2+1)^(1/2))/(-a+x+(a^2+1)^(1/2))/((-a+x)*(x^2+1))^(1/2),x, algorithm="maxima")`

output `integrate((a - x + sqrt(a^2 + 1))/(sqrt(-(x^2 + 1)*(a - x))*(a - x - sqrt(a^2 + 1))), x)`

Giac [F]

$$\int \frac{-a - \sqrt{1 + a^2} + x}{(-a + \sqrt{1 + a^2} + x) \sqrt{(-a + x)(1 + x^2)}} dx$$

$$= \int \frac{a - x + \sqrt{a^2 + 1}}{\sqrt{-(x^2 + 1)(a - x)(a - x - \sqrt{a^2 + 1})}} dx$$

input `integrate((-a+x-(a^2+1)^(1/2))/(-a+x+(a^2+1)^(1/2))/((-a+x)*(x^2+1))^(1/2),x, algorithm="giac")`

output `integrate((a - x + sqrt(a^2 + 1))/(sqrt(-(x^2 + 1)*(a - x))*(a - x - sqrt(a^2 + 1))), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{-a - \sqrt{1+a^2} + x}{(-a + \sqrt{1+a^2} + x) \sqrt{(-a+x)(1+x^2)}} dx$$

$$= \int -\frac{a-x+\sqrt{a^2+1}}{\sqrt{-(x^2+1)(a-x)}(x-a+\sqrt{a^2+1})} dx$$

input `int(-(a - x + (a^2 + 1)^(1/2))/((-x^2 + 1)*(a - x))^(1/2)*(x - a + (a^2 + 1)^(1/2))), x)`

output `int(-(a - x + (a^2 + 1)^(1/2))/((-x^2 + 1)*(a - x))^(1/2)*(x - a + (a^2 + 1)^(1/2))), x)`

Reduce [F]

$$\int \frac{-a - \sqrt{1+a^2} + x}{(-a + \sqrt{1+a^2} + x) \sqrt{(-a+x)(1+x^2)}} dx$$

$$= -2\sqrt{a^2+1} \left(\int \frac{\sqrt{-a+x} \sqrt{x^2+1} x}{2a^2x^3 - 3ax^4 + x^5 + 2a^2x - 2ax^2 + a - x} dx \right)$$

$$+ 2\sqrt{a^2+1} \left(\int \frac{\sqrt{-a+x} \sqrt{x^2+1}}{2a^2x^3 - 3ax^4 + x^5 + 2a^2x - 2ax^2 + a - x} dx \right) a$$

$$+ \int \frac{\sqrt{-a+x} \sqrt{x^2+1} x^2}{2a^2x^3 - 3ax^4 + x^5 + 2a^2x - 2ax^2 + a - x} dx$$

$$- 2 \left(\int \frac{\sqrt{-a+x} \sqrt{x^2+1} x}{2a^2x^3 - 3ax^4 + x^5 + 2a^2x - 2ax^2 + a - x} dx \right) a$$

$$+ 2 \left(\int \frac{\sqrt{-a+x} \sqrt{x^2+1}}{2a^2x^3 - 3ax^4 + x^5 + 2a^2x - 2ax^2 + a - x} dx \right) a^2$$

$$+ \int \frac{\sqrt{-a+x} \sqrt{x^2+1}}{2a^2x^3 - 3ax^4 + x^5 + 2a^2x - 2ax^2 + a - x} dx$$

input `int((-a+x-(a^2+1)^(1/2))/(-a+x+(a^2+1)^(1/2))/((-a+x)*(x^2+1))^(1/2), x)`

output

```

- 2*sqrt(a**2 + 1)*int((sqrt(- a + x)*sqrt(x**2 + 1)*x)/(2*a**2*x**3 + 2
*a**2*x - 3*a*x**4 - 2*a*x**2 + a + x**5 - x),x) + 2*sqrt(a**2 + 1)*int((s
qrt(- a + x)*sqrt(x**2 + 1))/(2*a**2*x**3 + 2*a**2*x - 3*a*x**4 - 2*a*x**
2 + a + x**5 - x),x)*a + int((sqrt(- a + x)*sqrt(x**2 + 1)*x**2)/(2*a**2*
x**3 + 2*a**2*x - 3*a*x**4 - 2*a*x**2 + a + x**5 - x),x) - 2*int((sqrt(-
a + x)*sqrt(x**2 + 1)*x)/(2*a**2*x**3 + 2*a**2*x - 3*a*x**4 - 2*a*x**2 + a
+ x**5 - x),x)*a + 2*int((sqrt(- a + x)*sqrt(x**2 + 1))/(2*a**2*x**3 + 2
*a**2*x - 3*a*x**4 - 2*a*x**2 + a + x**5 - x),x)*a**2 + int((sqrt(- a + x
)*sqrt(x**2 + 1))/(2*a**2*x**3 + 2*a**2*x - 3*a*x**4 - 2*a*x**2 + a + x**5
- x),x)

```

3.53
$$\int \frac{a+bx}{\sqrt[3]{1-x^2}(3+x^2)} dx$$

Optimal result	446
Mathematica [C] (warning: unable to verify)	447
Rubi [A] (verified)	447
Maple [F]	451
Fricas [F(-2)]	451
Sympy [F]	451
Maxima [F]	452
Giac [F]	452
Mupad [F(-1)]	452
Reduce [F]	453

Optimal result

Integrand size = 24, antiderivative size = 198

$$\int \frac{a+bx}{\sqrt[3]{1-x^2}(3+x^2)} dx = \frac{a \arctan\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\sqrt{3}b \arctan\left(\frac{1+\sqrt[3]{2-2x^2}}{\sqrt{3}}\right)}{2 \cdot 2^{2/3}}$$

$$+ \frac{a \arctan\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}}$$

$$- \frac{a \operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} + \frac{a \operatorname{arctanh}\left(\frac{x}{1+\sqrt[3]{2}\sqrt[3]{1-x^2}}\right)}{2 \cdot 2^{2/3}}$$

$$- \frac{b \log(3+x^2)}{4 \cdot 2^{2/3}} + \frac{3b \log\left(2^{2/3} - \sqrt[3]{1-x^2}\right)}{4 \cdot 2^{2/3}}$$

output

```
-1/12*a*arctanh(x)*2^(1/3)+1/4*a*arctanh(x/(1+2^(1/3)*(-x^2+1)^(1/3)))*2^(1/3)-1/8*b*ln(x^2+3)*2^(1/3)+3/8*b*ln(2^(2/3)-(-x^2+1)^(1/3))*2^(1/3)+1/12*a*arctan(3^(1/2)/x)*2^(1/3)*3^(1/2)+1/12*a*arctan((1-2^(1/3)*(-x^2+1)^(1/3))*3^(1/2)/x)*2^(1/3)*3^(1/2)+1/4*b*arctan(1/3*(1+(-2*x^2+2)^(1/3))*3^(1/2))*3^(1/2)*2^(1/3)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.15 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.73

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2} (3 + x^2)} dx = \frac{1}{6} bx^2 \operatorname{AppellF1} \left(1, \frac{1}{3}, 1, 2, x^2, -\frac{x^2}{3} \right) - \frac{9ax \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, x^2, -\frac{x^2}{3} \right) + 2x^2 \left(\operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{3}, 2, \frac{5}{2}, x^2, -\frac{x^2}{3} \right) - \operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{3}, 1, \frac{5}{2}, x^2, -\frac{x^2}{3} \right) \right)}{\sqrt[3]{1 - x^2} (3 + x^2)}$$

input `Integrate[(a + b*x)/((1 - x^2)^(1/3)*(3 + x^2)),x]`

output `(b*x^2*AppellF1[1, 1/3, 1, 2, x^2, -1/3*x^2])/6 - (9*a*x*AppellF1[1/2, 1/3, 1, 3/2, x^2, -1/3*x^2])/((1 - x^2)^(1/3)*(3 + x^2)*(-9*AppellF1[1/2, 1/3, 1, 3/2, x^2, -1/3*x^2] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5/2, x^2, -1/3*x^2] - AppellF1[3/2, 4/3, 1, 5/2, x^2, -1/3*x^2])))`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {1343, 305, 353, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2} (x^2 + 3)} dx$$

↓ 1343

$$a \int \frac{1}{\sqrt[3]{1 - x^2} (x^2 + 3)} dx + b \int \frac{x}{\sqrt[3]{1 - x^2} (x^2 + 3)} dx$$

↓ 305

$$b \int \frac{x}{\sqrt[3]{1-x^2}(x^2+3)} dx +$$

$$a \left(\frac{\arctan \left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan \left(\frac{\sqrt{3}}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\operatorname{arctanh} \left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2+1}} \right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} \right)$$

↓ 353

$$\frac{1}{2} b \int \frac{1}{\sqrt[3]{1-x^2}(x^2+3)} dx^2 +$$

$$a \left(\frac{\arctan \left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan \left(\frac{\sqrt{3}}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\operatorname{arctanh} \left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2+1}} \right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} \right)$$

↓ 67

$$\frac{1}{2} b \left(-\frac{3 \int \frac{1}{2^{2/3}-\sqrt[3]{1-x^2}} d\sqrt[3]{1-x^2}}{2 \cdot 2^{2/3}} + \frac{3}{2} \int \frac{1}{x^4 + 2^{2/3}\sqrt[3]{1-x^2} + 2\sqrt[3]{2}} d\sqrt[3]{1-x^2} - \frac{\log(x^2+3)}{2 \cdot 2^{2/3}} \right) +$$

$$a \left(\frac{\arctan \left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan \left(\frac{\sqrt{3}}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\operatorname{arctanh} \left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2+1}} \right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} \right)$$

↓ 16

$$\frac{1}{2} b \left(\frac{3}{2} \int \frac{1}{x^4 + 2^{2/3}\sqrt[3]{1-x^2} + 2\sqrt[3]{2}} d\sqrt[3]{1-x^2} - \frac{\log(x^2+3)}{2 \cdot 2^{2/3}} + \frac{3 \log(2^{2/3} - \sqrt[3]{1-x^2})}{2 \cdot 2^{2/3}} \right) +$$

$$a \left(\frac{\arctan \left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1-x^2})}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan \left(\frac{\sqrt{3}}{x} \right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\operatorname{arctanh} \left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2+1}} \right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} \right)$$

↓ 1082

$$\begin{aligned}
& \frac{1}{2}b \left(-\frac{3 \int \frac{1}{-x^4-3} d\left(\sqrt[3]{2}\sqrt[3]{1-x^2}+1\right)}{2^{2/3}} - \frac{\log(x^2+3)}{2 \cdot 2^{2/3}} + \frac{3 \log\left(2^{2/3} - \sqrt[3]{1-x^2}\right)}{2 \cdot 2^{2/3}} \right) + \\
& a \left(\frac{\arctan\left(\frac{\sqrt{3}\left(1-\sqrt[3]{2}\sqrt[3]{1-x^2}\right)}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} + \frac{\arctan\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2}+1}\right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} \right) \\
& \quad \downarrow 217 \\
& a \left(\frac{\arctan\left(\frac{\sqrt{3}\left(1-\sqrt[3]{2}\sqrt[3]{1-x^2}\right)}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} + \frac{\arctan\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2}+1}\right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} \right) + \\
& \frac{1}{2}b \left(\frac{\sqrt{3} \arctan\left(\frac{\sqrt[3]{2}\sqrt[3]{1-x^2}+1}{\sqrt{3}}\right)}{2^{2/3}} - \frac{\log(x^2+3)}{2 \cdot 2^{2/3}} + \frac{3 \log\left(2^{2/3} - \sqrt[3]{1-x^2}\right)}{2 \cdot 2^{2/3}} \right)
\end{aligned}$$

input `Int[(a + b*x)/((1 - x^2)^(1/3)*(3 + x^2)), x]`

output `a*(ArcTan[Sqrt[3]/x]/(2*2^(2/3)*Sqrt[3]) + ArcTan[(Sqrt[3]*(1 - 2^(1/3)*(1 - x^2)^(1/3)))/x]/(2*2^(2/3)*Sqrt[3]) - ArcTanh[x]/(6*2^(2/3)) + ArcTanh[x/(1 + 2^(1/3)*(1 - x^2)^(1/3))]/(2*2^(2/3))] + (b*((Sqrt[3]*ArcTan[(1 + 2^(1/3)*(1 - x^2)^(1/3))/Sqrt[3]])/2^(2/3) - Log[3 + x^2]/(2*2^(2/3)) + (3*Log[2^(2/3) - (1 - x^2)^(1/3)]/(2*2^(2/3))))/2`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^(1/3)), x_Symbol] := With[
 {q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x
] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)],
 x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x]) /
 ; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
 -1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])`

rule 305 `Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Wit
 h[{q = Rt[-b/a, 2]}, Simp[q*(ArcTan[Sqrt[3]/(q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/
 3)*d)), x] + (Simp[q*(ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(
 1/3))]/(2*2^(2/3)*a^(1/3)*d)), x] - Simp[q*(ArcTanh[q*x]/(6*2^(2/3)*a^(1/3
)*d)), x] + Simp[q*(ArcTan[Sqrt[3]*((a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3))]/(
 a^(1/3)*q*x))]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d)), x]) /; FreeQ[{a, b, c, d},
 x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]`

rule 353 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol]
 := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[
 {a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 imply[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Fre
 eQ[{a, b, c}, x]`

rule 1343 `Int[((g_) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q
 _), x_Symbol] := Simp[g Int[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + Simp[h
 Int[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; FreeQ[{a, c, d, f, g, h, p,
 q}, x]`

Maple [F]

$$\int \frac{bx + a}{(-x^2 + 1)^{\frac{1}{3}}(x^2 + 3)} dx$$

input `int((b*x+a)/(-x^2+1)^(1/3)/(x^2+3),x)`

output `int((b*x+a)/(-x^2+1)^(1/3)/(x^2+3),x)`

Fricas [F(-2)]

Exception generated.

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2}(3 + x^2)} dx = \text{Exception raised: TypeError}$$

input `integrate((b*x+a)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (trace 0)`

Sympy [F]

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2}(3 + x^2)} dx = \int \frac{a + bx}{\sqrt[3]{-(x - 1)(x + 1)}(x^2 + 3)} dx$$

input `integrate((b*x+a)/(-x**2+1)**(1/3)/(x**2+3),x)`

output `Integral((a + b*x)/((-x - 1)*(x + 1))**(1/3)*(x**2 + 3)), x)`

Maxima [F]

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2} (3 + x^2)} dx = \int \frac{bx + a}{(x^2 + 3)(-x^2 + 1)^{\frac{1}{3}}} dx$$

input `integrate((b*x+a)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="maxima")`

output `integrate((b*x + a)/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)`

Giac [F]

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2} (3 + x^2)} dx = \int \frac{bx + a}{(x^2 + 3)(-x^2 + 1)^{\frac{1}{3}}} dx$$

input `integrate((b*x+a)/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="giac")`

output `integrate((b*x + a)/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2} (3 + x^2)} dx = \int \frac{a + bx}{(1 - x^2)^{1/3} (x^2 + 3)} dx$$

input `int((a + b*x)/((1 - x^2)^(1/3)*(x^2 + 3)),x)`

output `int((a + b*x)/((1 - x^2)^(1/3)*(x^2 + 3)), x)`

Reduce [F]

$$\int \frac{a + bx}{\sqrt[3]{1 - x^2} (3 + x^2)} dx = \left(\int \frac{x}{(-x^2 + 1)^{\frac{1}{3}} x^2 + 3(-x^2 + 1)^{\frac{1}{3}}} dx \right) b + \left(\int \frac{1}{(-x^2 + 1)^{\frac{1}{3}} x^2 + 3(-x^2 + 1)^{\frac{1}{3}}} dx \right) a$$

input `int((b*x+a)/(-x^2+1)^(1/3)/(x^2+3),x)`

output `int(x/((-x**2+1)**(1/3)*x**2+3*(-x**2+1)**(1/3)),x)*b + int(1/((-x**2+1)**(1/3)*x**2+3*(-x**2+1)**(1/3)),x)*a`

3.54 $\int \frac{a+bx}{(3-x^2)\sqrt[3]{1+x^2}} dx$

Optimal result	454
Mathematica [B] (verified)	455
Rubi [A] (verified)	456
Maple [F]	459
Fricas [F(-2)]	460
Sympy [F]	460
Maxima [F]	460
Giac [F]	461
Mupad [F(-1)]	461
Reduce [F]	461

Optimal result

Integrand size = 24, antiderivative size = 198

$$\int \frac{a+bx}{(3-x^2)\sqrt[3]{1+x^2}} dx = -\frac{a \arctan(x)}{6 \cdot 2^{2/3}} + \frac{a \arctan\left(\frac{x}{1+\sqrt[3]{2}\sqrt[3]{1+x^2}}\right)}{2 \cdot 2^{2/3}} - \frac{\sqrt{3}b \arctan\left(\frac{1+\sqrt[3]{2}\sqrt[3]{1+x^2}}{\sqrt{3}}\right)}{2 \cdot 2^{2/3}} - \frac{a \operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} - \frac{a \operatorname{arctanh}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1+x^2})}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} + \frac{b \log(3-x^2)}{4 \cdot 2^{2/3}} - \frac{3b \log\left(2^{2/3}-\sqrt[3]{1+x^2}\right)}{4 \cdot 2^{2/3}}$$

output

```
-1/12*a*arctan(x)*2^(1/3)+1/4*a*arctan(x/(1+2^(1/3)*(x^2+1)^(1/3)))*2^(1/3)
)+1/8*b*ln(-x^2+3)*2^(1/3)-3/8*b*ln(2^(2/3)-(x^2+1)^(1/3))*2^(1/3)-1/12*a*
arctanh(3^(1/2)/x)*2^(1/3)*3^(1/2)-1/12*a*arctanh((1-2^(1/3)*(x^2+1)^(1/3)
))*3^(1/2)/x)*2^(1/3)*3^(1/2)-1/4*b*arctan(1/3*(1+2^(1/3)*(x^2+1)^(1/3))*3^
(1/2))*3^(1/2)*2^(1/3)
```

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1072 vs. $2(198) = 396$.

Time = 9.75 (sec) , antiderivative size = 1072, normalized size of antiderivative = 5.41

$$\int \frac{a + bx}{(3 - x^2) \sqrt[3]{1 + x^2}} dx = \text{Too large to display}$$

input `Integrate[(a + b*x)/((3 - x^2)*(1 + x^2)^(1/3)),x]`

output

```
((2*(Sqrt[3]*a^4 - 12*a^3*b + 18*Sqrt[3]*a^2*b^2 - 36*a*b^3 + 9*Sqrt[3]*b^4)*ArcTan[(3*b + a*x + 6*2^(1/3)*b*(1 + x^2)^(1/3) - Sqrt[3]*(a + b*x + 2*2^(1/3)*a*(1 + x^2)^(1/3))]/(3*b*(Sqrt[3] - x) + a*(-3 + Sqrt[3]*x)))]/(Sqrt[3]*a^3 - 9*a^2*b + 9*Sqrt[3]*a*b^2 - 9*b^3) - (2*(Sqrt[3]*a^4 + 12*a^3*b + 18*Sqrt[3]*a^2*b^2 + 36*a*b^3 + 9*Sqrt[3]*b^4)*ArcTan[(3*b + a*x + 6*2^(1/3)*b*(1 + x^2)^(1/3) + Sqrt[3]*(a + b*x + 2*2^(1/3)*a*(1 + x^2)^(1/3))]/(3*b*(Sqrt[3] + x) + a*(3 + Sqrt[3]*x)))]/(Sqrt[3]*a^3 + 9*a^2*b + 9*Sqrt[3]*a*b^2 + 9*b^3) + (2*(a^4 - 4*Sqrt[3]*a^3*b + 18*a^2*b^2 - 12*Sqrt[3]*a*b^3 + 9*b^4)*Log[3*b + a*x - 3*2^(1/3)*b*(1 + x^2)^(1/3) - Sqrt[3]*(a + b*x - 2^(1/3)*a*(1 + x^2)^(1/3))]/(Sqrt[3]*a^3 - 9*a^2*b + 9*Sqrt[3]*a*b^2 - 9*b^3) - (2*(a^4 + 4*Sqrt[3]*a^3*b + 18*a^2*b^2 + 12*Sqrt[3]*a*b^3 + 9*b^4)*Log[-3*b - a*x + 3*2^(1/3)*b*(1 + x^2)^(1/3) - Sqrt[3]*(a + b*x - 2^(1/3)*a*(1 + x^2)^(1/3))]/(Sqrt[3]*a^3 + 9*a^2*b + 9*Sqrt[3]*a*b^2 + 9*b^3) - ((a^4 - 4*Sqrt[3]*a^3*b + 18*a^2*b^2 - 12*Sqrt[3]*a*b^3 + 9*b^4)*Log[3*a^2 + 9*b^2 + 12*a*b*x + a^2*x^2 + 3*b^2*x^2 + 3*2^(1/3)*(a^2 + 3*b^2 + 2*a*b*x)*(1 + x^2)^(1/3) + 3*2^(2/3)*(a^2 + 3*b^2)*(1 + x^2)^(2/3) + Sqrt[3]*(-6*a*b - 2*a^2*x - 6*b^2*x - 2*a*b*x^2 - 2^(1/3)*(6*a*b + a^2*x + 3*b^2*x)*(1 + x^2)^(1/3) - 6*2^(2/3)*a*b*(1 + x^2)^(2/3))]/(Sqrt[3]*a^3 - 9*a^2*b + 9*Sqrt[3]*a*b^2 - 9*b^3) + ((a^4 + 4*Sqrt[3]*a^3*b + 18*a^2*b^2 + 12*Sqrt[3]*a*b^3 + 9*b^4)*Log[3*a^2 + 9*b^2 + 12*a*b*x + a^2*x^2 + 3*b^2...
```


Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {1343, 304, 353, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{a + bx}{(3 - x^2) \sqrt[3]{x^2 + 1}} dx \\
 & \quad \downarrow \text{1343} \\
 & a \int \frac{1}{(3 - x^2) \sqrt[3]{x^2 + 1}} dx + b \int \frac{x}{(3 - x^2) \sqrt[3]{x^2 + 1}} dx \\
 & \quad \downarrow \text{304} \\
 & b \int \frac{x}{(3 - x^2) \sqrt[3]{x^2 + 1}} dx + \\
 & a \left(\frac{\arctan\left(\frac{x}{\sqrt[3]{2} \sqrt[3]{x^2 + 1} + 1}\right)}{2 \cdot 2^{2/3}} - \frac{\arctan(x)}{6 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1 - \sqrt[3]{2} \sqrt[3]{x^2 + 1})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} \right) \\
 & \quad \downarrow \text{353} \\
 & \frac{1}{2} b \int \frac{1}{(3 - x^2) \sqrt[3]{x^2 + 1}} dx^2 + \\
 & a \left(\frac{\arctan\left(\frac{x}{\sqrt[3]{2} \sqrt[3]{x^2 + 1} + 1}\right)}{2 \cdot 2^{2/3}} - \frac{\arctan(x)}{6 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1 - \sqrt[3]{2} \sqrt[3]{x^2 + 1})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} \right) \\
 & \quad \downarrow \text{67}
 \end{aligned}$$

$$\frac{1}{2}b \left(\frac{3 \int \frac{1}{2^{2/3} - \sqrt[3]{x^2+1}} d\sqrt[3]{x^2+1}}{2 \cdot 2^{2/3}} - \frac{3}{2} \int \frac{1}{x^4 + 2^{2/3} \sqrt[3]{x^2+1} + 2\sqrt[3]{2}} d\sqrt[3]{x^2+1} + \frac{\log(3-x^2)}{2 \cdot 2^{2/3}} \right) +$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt[3]{2} \sqrt[3]{x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\arctan(x)}{6 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-\sqrt[3]{2} \sqrt[3]{x^2+1})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} \right)$$

↓ 16

$$\frac{1}{2}b \left(-\frac{3}{2} \int \frac{1}{x^4 + 2^{2/3} \sqrt[3]{x^2+1} + 2\sqrt[3]{2}} d\sqrt[3]{x^2+1} + \frac{\log(3-x^2)}{2 \cdot 2^{2/3}} - \frac{3 \log(2^{2/3} - \sqrt[3]{x^2+1})}{2 \cdot 2^{2/3}} \right) +$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt[3]{2} \sqrt[3]{x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\arctan(x)}{6 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-\sqrt[3]{2} \sqrt[3]{x^2+1})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} \right)$$

↓ 1082

$$\frac{1}{2}b \left(\frac{3 \int \frac{1}{-x^4-3} d(\sqrt[3]{2} \sqrt[3]{x^2+1} + 1)}{2^{2/3}} + \frac{\log(3-x^2)}{2 \cdot 2^{2/3}} - \frac{3 \log(2^{2/3} - \sqrt[3]{x^2+1})}{2 \cdot 2^{2/3}} \right) +$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt[3]{2} \sqrt[3]{x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\arctan(x)}{6 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-\sqrt[3]{2} \sqrt[3]{x^2+1})}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} \right)$$

↓ 217

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\arctan(x)}{6 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{x^2+1})}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} \right) +$$

$$\frac{1}{2}b \left(-\frac{\sqrt{3}\arctan\left(\frac{\sqrt[3]{2}\sqrt[3]{x^2+1}}{\sqrt{3}}\right)}{2^{2/3}} + \frac{\log(3-x^2)}{2 \cdot 2^{2/3}} - \frac{3\log(2^{2/3}-\sqrt[3]{x^2+1})}{2 \cdot 2^{2/3}} \right)$$

input `Int[(a + b*x)/((3 - x^2)*(1 + x^2)^(1/3)), x]`

output `a*(-1/6*ArcTan[x]/2^(2/3) + ArcTan[x/(1 + 2^(1/3)*(1 + x^2)^(1/3))]/(2*2^(2/3)) - ArcTanh[Sqrt[3]/x]/(2*2^(2/3)*Sqrt[3]) - ArcTanh[(Sqrt[3]*(1 - 2^(1/3)*(1 + x^2)^(1/3)))/x]/(2*2^(2/3)*Sqrt[3])) + (b*(-((Sqrt[3]*ArcTan[(1 + 2^(1/3)*(1 + x^2)^(1/3)]/Sqrt[3])]/2^(2/3)) + Log[3 - x^2]/(2*2^(2/3)) - (3*Log[2^(2/3) - (1 + x^2)^(1/3)]/(2*2^(2/3)))))/2`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])]/; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 304 `Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[q*(ArcTanh[Sqrt[3]/(q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d)), x] + (-Simp[q*(ArcTan[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3))]/(2*2^(2/3)*a^(1/3)*d)), x] + Simp[q*(ArcTan[q*x]/(6*2^(2/3)*a^(1/3)*d)), x] + Simp[q*(ArcTanh[Sqrt[3]*((a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d)), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && PosQ[b/a]`

rule 353 `Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol] := Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]`

rule 1082 `Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]`

rule 1343 `Int[((g_) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_.)*((d_) + (f_.)*(x_)^2)^(q_.), x_Symbol] := Simp[g Int[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + Simp[h Int[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; FreeQ[{a, c, d, f, g, h, p, q}, x]`

Maple [F]

$$\int \frac{bx + a}{(-x^2 + 3)(x^2 + 1)^{\frac{1}{3}}} dx$$

input `int((b*x+a)/(-x^2+3)/(x^2+1)^(1/3),x)`

output `int((b*x+a)/(-x^2+3)/(x^2+1)^(1/3),x)`

Fricas [F(-2)]

Exception generated.

$$\int \frac{a + bx}{(3 - x^2)\sqrt[3]{1 + x^2}} dx = \text{Exception raised: TypeError}$$

input `integrate((b*x+a)/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (trace 0)`

Sympy [F]

$$\begin{aligned} & \int \frac{a + bx}{(3 - x^2)\sqrt[3]{1 + x^2}} dx \\ &= - \int \frac{a}{x^2\sqrt[3]{x^2 + 1} - 3\sqrt[3]{x^2 + 1}} dx - \int \frac{bx}{x^2\sqrt[3]{x^2 + 1} - 3\sqrt[3]{x^2 + 1}} dx \end{aligned}$$

input `integrate((b*x+a)/(-x**2+3)/(x**2+1)**(1/3),x)`

output `-Integral(a/(x**2*(x**2 + 1)**(1/3) - 3*(x**2 + 1)**(1/3)), x) - Integral(b*x/(x**2*(x**2 + 1)**(1/3) - 3*(x**2 + 1)**(1/3)), x)`

Maxima [F]

$$\int \frac{a + bx}{(3 - x^2)\sqrt[3]{1 + x^2}} dx = \int -\frac{bx + a}{(x^2 + 1)^{\frac{1}{3}}(x^2 - 3)} dx$$

input `integrate((b*x+a)/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="maxima")`

output `-integrate((b*x + a)/((x^2 + 1)^(1/3)*(x^2 - 3)), x)`

Giac [F]

$$\int \frac{a + bx}{(3 - x^2)\sqrt[3]{1 + x^2}} dx = \int -\frac{bx + a}{(x^2 + 1)^{\frac{1}{3}}(x^2 - 3)} dx$$

input `integrate((b*x+a)/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="giac")`

output `integrate(-(b*x + a)/((x^2 + 1)^(1/3)*(x^2 - 3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + bx}{(3 - x^2)\sqrt[3]{1 + x^2}} dx = \int -\frac{a + bx}{(x^2 + 1)^{\frac{1}{3}}(x^2 - 3)} dx$$

input `int(-(a + b*x)/((x^2 + 1)^(1/3)*(x^2 - 3)),x)`

output `int(-(a + b*x)/((x^2 + 1)^(1/3)*(x^2 - 3)), x)`

Reduce [F]

$$\int \frac{a + bx}{(3 - x^2)\sqrt[3]{1 + x^2}} dx = -\left(\int \frac{x}{(x^2 + 1)^{\frac{1}{3}}x^2 - 3(x^2 + 1)^{\frac{1}{3}}} dx\right) b - \left(\int \frac{1}{(x^2 + 1)^{\frac{1}{3}}x^2 - 3(x^2 + 1)^{\frac{1}{3}}} dx\right) a$$

input `int((b*x+a)/(-x^2+3)/(x^2+1)^(1/3),x)`

output `- (int(x/((x**2 + 1)**(1/3)*x**2 - 3*(x**2 + 1)**(1/3)),x)*b + int(1/((x**2 + 1)**(1/3)*x**2 - 3*(x**2 + 1)**(1/3)),x)*a)`

3.55 $\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx$

Optimal result	462
Mathematica [A] (verified)	462
Rubi [A] (verified)	463
Maple [C] (warning: unable to verify)	464
Fricas [B] (verification not implemented)	465
Sympy [F]	466
Maxima [F]	466
Giac [F]	467
Mupad [F(-1)]	467
Reduce [F]	467

Optimal result

Integrand size = 18, antiderivative size = 97

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = -\frac{\arctan\left(\frac{1}{\sqrt{3}} + \frac{2^{2/3}(2-x)}{\sqrt{3}\sqrt[3]{4-6x+3x^2}}\right)}{2^{2/3}\sqrt{3}} - \frac{\log(x)}{2 \cdot 2^{2/3}} + \frac{\log\left(6-3x-3\sqrt[3]{2}\sqrt[3]{4-6x+3x^2}\right)}{2 \cdot 2^{2/3}}$$

output

```
-1/4*ln(x)*2^(1/3)+1/4*ln(6-3*x-3*2^(1/3)*(3*x^2-6*x+4)^(1/3))*2^(1/3)+1/6
*arctan(-1/3*3^(1/2)-1/3*2^(2/3)*(2-x)/(3*x^2-6*x+4)^(1/3)*3^(1/2))*2^(1/3
)*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.73

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = \frac{2\sqrt{3} \arctan\left(\frac{2 \cdot 2^{2/3} - 2^{2/3}x + \sqrt[3]{4-6x+3x^2}}{\sqrt{3}\sqrt[3]{4-6x+3x^2}}\right) - 2 \log\left(-2 \cdot 2^{2/3} + 2^{2/3}x + 2\sqrt[3]{4-6x+3x^2}\right) + \log\left(-4\sqrt[3]{4-6x+3x^2}\right)}{6 \cdot 2^{2/3}}$$

input `Integrate[1/(x*(4 - 6*x + 3*x^2)^(1/3)),x]`

output
$$-1/6*(2*\text{Sqrt}[3]*\text{ArcTan}[(2*2^{(2/3)} - 2^{(2/3)}*x + (4 - 6*x + 3*x^2)^{(1/3)})/(\text{Sqrt}[3]*(4 - 6*x + 3*x^2)^{(1/3)})] - 2*\text{Log}[-2*2^{(2/3)} + 2^{(2/3)}*x + 2*(4 - 6*x + 3*x^2)^{(1/3)}] + \text{Log}[-4*2^{(1/3)} + 4*2^{(1/3)}*x - 2^{(1/3)}*x^2 + 2^{(2/3)}*(-2 + x)*(4 - 6*x + 3*x^2)^{(1/3)} - 2*(4 - 6*x + 3*x^2)^{(2/3)}])/2^{(2/3)}$$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {1175}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{x \sqrt[3]{3x^2 - 6x + 4}} dx$$

↓ 1175

$$-\frac{\arctan\left(\frac{2^{2/3}(2-x)}{\sqrt{3}\sqrt[3]{3x^2 - 6x + 4}} + \frac{1}{\sqrt{3}}\right)}{2^{2/3}\sqrt{3}} + \frac{\log\left(-3\sqrt[3]{2}\sqrt[3]{3x^2 - 6x + 4} - 3x + 6\right)}{2 \cdot 2^{2/3}} - \frac{\log(x)}{2 \cdot 2^{2/3}}$$

input `Int[1/(x*(4 - 6*x + 3*x^2)^(1/3)),x]`

output
$$-(\text{ArcTan}[1/\text{Sqrt}[3] + (2^{(2/3)}*(2 - x))/(\text{Sqrt}[3]*(4 - 6*x + 3*x^2)^{(1/3)})]/(2^{(2/3)}*\text{Sqrt}[3])) - \text{Log}[x]/(2*2^{(2/3)}) + \text{Log}[6 - 3*x - 3*2^{(1/3)}*(4 - 6*x + 3*x^2)^{(1/3)}]/(2*2^{(2/3)})$$

Defintions of rubi rules used

rule 1175

```
Int[1/(((d_.) + (e_.)*(x_))*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(1/3)), x_Sy
mbol] := With[{q = Rt[3*c*e^2*(2*c*d - b*e), 3]}, Simp[(-Sqrt[3])*c*e*(ArcT
an[1/Sqrt[3] + 2*((c*d - b*e - c*e*x)/(Sqrt[3]*q*(a + b*x + c*x^2)^(1/3)))]
/q^2), x] + (-Simp[3*c*e*(Log[d + e*x]/(2*q^2)), x] + Simp[3*c*e*(Log[c*d -
b*e - c*e*x - q*(a + b*x + c*x^2)^(1/3)]/(2*q^2)), x]]) /; FreeQ[{a, b, c,
d, e}, x] && EqQ[c^2*d^2 - b*c*d*e + b^2*e^2 - 3*a*c*e^2, 0] && PosQ[c*e^2
*(2*c*d - b*e)]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 12.41 (sec) , antiderivative size = 1593, normalized size of antiderivative = 16.42

method	result	size
trager	Expression too large to display	1593

input

```
int(1/x/(3*x^2-6*x+4)^(1/3),x,method=_RETURNVERBOSE)
```

output

```

1/6*RootOf(_Z^3-2)*ln(-(48*(3*x^2-6*x+4)^(2/3)*RootOf(RootOf(_Z^3-2)^2+2*_
Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2*x+18*(3*x^2-6*x+4)^(1/3)*RootOf(
RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)*x^2-72*(3*x^2-
6*x+4)^(1/3)*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z
^3-2)*x-40*RootOf(_Z^3-2)+320*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+
4*_Z^2)+30*(3*x^2-6*x+4)^(2/3)*x+96*(3*x^2-6*x+4)^(1/3)*RootOf(_Z^3-2)^2-8
*RootOf(_Z^3-2)^3*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)+3*Ro
otOf(_Z^3-2)*x^3-30*RootOf(_Z^3-2)*x^2+60*RootOf(_Z^3-2)*x-24*RootOf(RootO
f(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*x^3+240*RootOf(RootOf(_Z^3-2)^2+2*_
_Z*RootOf(_Z^3-2)+4*_Z^2)*x^2-480*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3
-2)+4*_Z^2)*x+64*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)^2*Ro
otOf(_Z^3-2)^2-16*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)^2*Ro
otOf(_Z^3-2)^2*x^3+2*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)*Ro
otOf(_Z^3-2)^3*x^3+48*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z^2)^
2*RootOf(_Z^3-2)^2*x^2-6*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_Z
^2)*RootOf(_Z^3-2)^3*x^2-96*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4*_
Z^2)^2*RootOf(_Z^3-2)^2*x+12*RootOf(RootOf(_Z^3-2)^2+2*_Z*RootOf(_Z^3-2)+4
*_Z^2)*RootOf(_Z^3-2)^3*x-96*(3*x^2-6*x+4)^(2/3)*RootOf(RootOf(_Z^3-2)^2+2
*_Z*RootOf(_Z^3-2)+4*_Z^2)*RootOf(_Z^3-2)^2+24*(3*x^2-6*x+4)^(1/3)*RootOf(
_Z^3-2)^2*x^2-96*(3*x^2-6*x+4)^(1/3)*RootOf(_Z^3-2)^2*x+72*(3*x^2-6*x+4...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 171 vs. $2(74) = 148$.

Time = 1.07 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.76

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = -\frac{1}{6}$$

$$\cdot 4^{\frac{1}{6}}\sqrt{3} \arctan \left(\frac{4^{\frac{1}{6}}\sqrt{3} \left(4^{\frac{1}{3}}x^3 + 2 \cdot 4^{\frac{2}{3}}(3x^2 - 6x + 4)^{\frac{2}{3}}(x - 2) + 4(3x^2 - 6x + 4)^{\frac{1}{3}}(x^2 - 4x + 4) \right)}{6(x^3 - 12x^2 + 24x - 16)} \right)$$

$$+ \frac{1}{12} \cdot 4^{\frac{1}{3}} \log \left(\frac{4^{\frac{1}{3}}(x - 2) + 2(3x^2 - 6x + 4)^{\frac{1}{3}}}{x} \right) - \frac{1}{24}$$

$$\cdot 4^{\frac{2}{3}} \log \left(\frac{4^{\frac{2}{3}}(3x^2 - 6x + 4)^{\frac{2}{3}} + 4^{\frac{1}{3}}(x^2 - 4x + 4) - 2(3x^2 - 6x + 4)^{\frac{1}{3}}(x - 2)}{x^2} \right)$$

input

```
integrate(1/x/(3*x^2-6*x+4)^(1/3),x, algorithm="fricas")
```

output

```
-1/6*4^(1/6)*sqrt(3)*arctan(1/6*4^(1/6)*sqrt(3)*(4^(1/3)*x^3 + 2*4^(2/3)*(
3*x^2 - 6*x + 4)^(2/3)*(x - 2) + 4*(3*x^2 - 6*x + 4)^(1/3)*(x^2 - 4*x + 4)
)/(x^3 - 12*x^2 + 24*x - 16)) + 1/12*4^(2/3)*log((4^(1/3)*(x - 2) + 2*(3*x
^2 - 6*x + 4)^(1/3))/x) - 1/24*4^(2/3)*log((4^(2/3)*(3*x^2 - 6*x + 4)^(2/3
) + 4^(1/3)*(x^2 - 4*x + 4) - 2*(3*x^2 - 6*x + 4)^(1/3)*(x - 2))/x^2)
```

Sympy [F]

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = \int \frac{1}{x\sqrt[3]{3x^2-6x+4}} dx$$

input

```
integrate(1/x/(3*x**2-6*x+4)**(1/3),x)
```

output

```
Integral(1/(x*(3*x**2 - 6*x + 4)**(1/3)), x)
```

Maxima [F]

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = \int \frac{1}{(3x^2-6x+4)^{\frac{1}{3}}x} dx$$

input

```
integrate(1/x/(3*x^2-6*x+4)^(1/3),x, algorithm="maxima")
```

output

```
integrate(1/((3*x^2 - 6*x + 4)^(1/3)*x), x)
```

Giac [F]

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = \int \frac{1}{(3x^2-6x+4)^{\frac{1}{3}}x} dx$$

input `integrate(1/x/(3*x^2-6*x+4)^(1/3),x, algorithm="giac")`

output `integrate(1/((3*x^2 - 6*x + 4)^(1/3)*x), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = \int \frac{1}{x(3x^2-6x+4)^{1/3}} dx$$

input `int(1/(x*(3*x^2 - 6*x + 4)^(1/3)),x)`

output `int(1/(x*(3*x^2 - 6*x + 4)^(1/3)), x)`

Reduce [F]

$$\int \frac{1}{x\sqrt[3]{4-6x+3x^2}} dx = \int \frac{1}{(3x^2-6x+4)^{\frac{1}{3}}x} dx$$

input `int(1/x/(3*x^2-6*x+4)^(1/3),x)`

output `int(1/((3*x**2 - 6*x + 4)**(1/3)*x),x)`

3.56 $\int x\sqrt[3]{1-x^3} dx$

Optimal result	468
Mathematica [A] (verified)	468
Rubi [A] (verified)	469
Maple [C] (verified)	470
Fricas [A] (verification not implemented)	471
Sympy [C] (verification not implemented)	471
Maxima [A] (verification not implemented)	472
Giac [F]	472
Mupad [F(-1)]	473
Reduce [F]	473

Optimal result

Integrand size = 13, antiderivative size = 73

$$\int x\sqrt[3]{1-x^3} dx = \frac{1}{3}x^2\sqrt[3]{1-x^3} - \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{1}{6}\log\left(-x - \sqrt[3]{1-x^3}\right)$$

output

$1/3*x^2*(-x^3+1)^{(1/3)}-1/6*\ln(-x-(-x^3+1)^{(1/3)})-1/9*\arctan(1/3*(1-2*x/(-x^3+1)^{(1/3)})*3^{(1/2)})*3^{(1/2)}$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.36

$$\int x\sqrt[3]{1-x^3} dx = \frac{1}{18}\left(6x^2\sqrt[3]{1-x^3} - 2\sqrt{3}\arctan\left(\frac{\sqrt{3}x}{x-2\sqrt[3]{1-x^3}}\right) - 2\log\left(x + \sqrt[3]{1-x^3}\right) + \log\left(x^2 - x\sqrt[3]{1-x^3} + (1-x^3)^{2/3}\right)\right)$$

input

$\text{Integrate}[x*(1-x^3)^{(1/3)},x]$

output

$$(6x^2(1-x^3)^{1/3} - 2\sqrt{3}\operatorname{ArcTan}[(\sqrt{3}x)/(x - 2(1-x^3)^{1/3})] - 2\operatorname{Log}[x + (1-x^3)^{1/3}] + \operatorname{Log}[x^2 - x(1-x^3)^{1/3} + (1-x^3)^{2/3}])/18$$
Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {811, 853}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int x \sqrt[3]{1-x^3} dx$$

$$\downarrow 811$$

$$\frac{1}{3} \int \frac{x}{(1-x^3)^{2/3}} dx + \frac{1}{3} \sqrt[3]{1-x^3} x^2$$

$$\downarrow 853$$

$$\frac{1}{3} \left(-\frac{\arctan\left(\frac{1-\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(-\sqrt[3]{1-x^3} - x\right) \right) + \frac{1}{3} \sqrt[3]{1-x^3} x^2$$

input

$$\text{Int}[x*(1-x^3)^{1/3},x]$$

output

$$(x^2*(1-x^3)^{1/3})/3 + (-(\operatorname{ArcTan}[(1-(2*x))/(1-x^3)^{1/3}]/\sqrt{3}]/\sqrt{3}) - \operatorname{Log}[-x - (1-x^3)^{1/3}])/2/3$$

Defintions of rubi rules used

```
rule 811 Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c*(m + n*p + 1))), x] + Simp[a*n*(p/(m + n*p + 1)) Int[(c*x)^m*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c, m}, x] && I GtQ[n, 0] && GtQ[p, 0] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

```
rule 853 Int[(x_)/((a_) + (b_.)*(x_)^3)^(2/3), x_Symbol] := With[{q = Rt[b, 3]}, Simp[-ArcTan[(1 + 2*q*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*q^2), x] - Simp[Log[q*x - (a + b*x^3)^(1/3)]/(2*q^2), x]] /; FreeQ[{a, b}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 1.51 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.21

method	result
meijerg	$\frac{x^2 \operatorname{hypergeom}\left(\left[-\frac{1}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right)}{2}$
risch	$-\frac{x^2(x^3-1)}{3(-x^3+1)^{\frac{2}{3}}} + \frac{(x^3-1)^{\frac{2}{3}}(-\operatorname{signum}(x^3-1))^{\frac{2}{3}}x^2 \operatorname{hypergeom}\left(\left[\frac{2}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right)}{6 \operatorname{signum}(x^3-1)^{\frac{2}{3}}(-x^3+1)^{\frac{2}{3}}}$
pseudoelliptic	$\frac{6x^2(-x^3+1)^{\frac{1}{3}}+2\sqrt{3} \arctan\left(\frac{(-2(-x^3+1)^{\frac{1}{3}}+x)\sqrt{3}}{3x}\right)+\ln\left(\frac{(-x^3+1)^{\frac{2}{3}}-x(-x^3+1)^{\frac{1}{3}}+x^2}{x^2}\right)-2\ln\left(\frac{x+(-x^3+1)^{\frac{1}{3}}}{x}\right)}{18\left((-x^3+1)^{\frac{2}{3}}-x(-x^3+1)^{\frac{1}{3}}+x^2\right)\left(x+(-x^3+1)^{\frac{1}{3}}\right)}$
trager	$\frac{x^2(-x^3+1)^{\frac{1}{3}}}{3} - \frac{\ln\left(-2\operatorname{RootOf}\left(_Z^2-_Z+1\right)^2x^3+3\operatorname{RootOf}\left(_Z^2-_Z+1\right)(-x^3+1)^{\frac{2}{3}}x-\operatorname{RootOf}\left(_Z^2-_Z+1\right)\right)}{9}$

```
input int(x*(-x^3+1)^(1/3), x, method=_RETURNVERBOSE)
```

```
output 1/2*x^2*hypergeom([-1/3, 2/3], [5/3], x^3)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.32

$$\int x\sqrt[3]{1-x^3} dx = \frac{1}{3}(-x^3+1)^{\frac{1}{3}}x^2 - \frac{1}{9}\sqrt{3}\arctan\left(-\frac{\sqrt{3}x-2\sqrt{3}(-x^3+1)^{\frac{1}{3}}}{3x}\right) - \frac{1}{9}\log\left(\frac{x+(-x^3+1)^{\frac{1}{3}}}{x}\right) + \frac{1}{18}\log\left(\frac{x^2-(-x^3+1)^{\frac{1}{3}}x+(-x^3+1)^{\frac{2}{3}}}{x^2}\right)$$

input `integrate(x*(-x^3+1)^(1/3),x, algorithm="fricas")`

output `1/3*(-x^3 + 1)^(1/3)*x^2 - 1/9*sqrt(3)*arctan(-1/3*(sqrt(3)*x - 2*sqrt(3)*(-x^3 + 1)^(1/3))/x) - 1/9*log((x + (-x^3 + 1)^(1/3))/x) + 1/18*log((x^2 - (-x^3 + 1)^(1/3)*x + (-x^3 + 1)^(2/3))/x^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.50 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.44

$$\int x\sqrt[3]{1-x^3} dx = \frac{x^2\Gamma\left(\frac{2}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{1}{3}, \frac{2}{3} \\ \frac{5}{3} \end{matrix} \middle| x^3 e^{2i\pi}\right)}{3\Gamma\left(\frac{5}{3}\right)}$$

input `integrate(x*(-x**3+1)**(1/3),x)`

output `x**2*gamma(2/3)*hyper((-1/3, 2/3), (5/3,), x**3*exp_polar(2*I*pi))/(3*gamma(5/3))`

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.44

$$\int x\sqrt[3]{1-x^3} dx = -\frac{1}{9}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}\left(\frac{2(-x^3+1)^{\frac{1}{3}}}{x}-1\right)\right) - \frac{(-x^3+1)^{\frac{1}{3}}}{3x\left(\frac{x^3-1}{x^3}-1\right)} - \frac{1}{9}\log\left(\frac{(-x^3+1)^{\frac{1}{3}}}{x}+1\right) + \frac{1}{18}\log\left(-\frac{(-x^3+1)^{\frac{1}{3}}}{x}+\frac{(-x^3+1)^{\frac{2}{3}}}{x^2}+1\right)$$

input `integrate(x*(-x^3+1)^(1/3),x, algorithm="maxima")`output `-1/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3)/x - 1)) - 1/3*(-x^3 + 1)^(1/3)/(x*((x^3 - 1)/x^3 - 1)) - 1/9*log((-x^3 + 1)^(1/3)/x + 1) + 1/18*log(-(-x^3 + 1)^(1/3)/x + (-x^3 + 1)^(2/3)/x^2 + 1)`**Giac [F]**

$$\int x\sqrt[3]{1-x^3} dx = \int (-x^3+1)^{\frac{1}{3}}x dx$$

input `integrate(x*(-x^3+1)^(1/3),x, algorithm="giac")`output `integrate((-x^3 + 1)^(1/3)*x, x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \sqrt[3]{1-x^3} dx = \int x (1-x^3)^{1/3} dx$$

input `int(x*(1 - x^3)^(1/3), x)`output `int(x*(1 - x^3)^(1/3), x)`**Reduce [F]**

$$\int x^3 \sqrt[3]{1-x^3} dx = \frac{(-x^3+1)^{\frac{1}{3}} x^2}{3} - \frac{\left(\int \frac{(-x^3+1)^{\frac{1}{3}} x}{x^3-1} dx \right)}{3}$$

input `int(x*(-x^3+1)^(1/3), x)`output `((- x**3 + 1)**(1/3)*x**2 - int(((- x**3 + 1)**(1/3)*x)/(x**3 - 1), x))/3`

3.57 $\int \frac{\sqrt[3]{1-x^3}}{x} dx$

Optimal result	474
Mathematica [A] (verified)	474
Rubi [A] (verified)	475
Maple [C] (verified)	477
Fricas [A] (verification not implemented)	477
Sympy [C] (verification not implemented)	478
Maxima [A] (verification not implemented)	478
Giac [A] (verification not implemented)	479
Mupad [B] (verification not implemented)	479
Reduce [F]	480

Optimal result

Integrand size = 15, antiderivative size = 67

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = \sqrt[3]{1-x^3} - \frac{\arctan\left(\frac{1+2\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\log(x)}{2} + \frac{1}{2} \log\left(1 - \sqrt[3]{1-x^3}\right)$$

output

```
(-x^3+1)^(1/3)-1/2*ln(x)+1/2*ln(1-(-x^3+1)^(1/3))-1/3*arctan(1/3*(1+2*(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.31

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = \sqrt[3]{1-x^3} - \frac{\arctan\left(\frac{1+2\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(-1 + \sqrt[3]{1-x^3}\right) - \frac{1}{6} \log\left(1 + \sqrt[3]{1-x^3} + (1-x^3)^{2/3}\right)$$

input

```
Integrate[(1 - x^3)^(1/3)/x,x]
```

output

$$(1 - x^3)^{1/3} - \text{ArcTan}[(1 + 2(1 - x^3)^{1/3})/\text{Sqrt}[3]]/\text{Sqrt}[3] + \text{Log}[-1 + (1 - x^3)^{1/3}]/3 - \text{Log}[1 + (1 - x^3)^{1/3} + (1 - x^3)^{2/3}]/6$$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.12, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {798, 60, 69, 16, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx$$

$$\downarrow 798$$

$$\frac{1}{3} \int \frac{\sqrt[3]{1-x^3}}{x^3} dx^3$$

$$\downarrow 60$$

$$\frac{1}{3} \left(\int \frac{1}{x^3(1-x^3)^{2/3}} dx^3 + 3\sqrt[3]{1-x^3} \right)$$

$$\downarrow 69$$

$$\frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{1-\sqrt[3]{1-x^3}} d\sqrt[3]{1-x^3} - \frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{1-x^3} + 1} d\sqrt[3]{1-x^3} + 3\sqrt[3]{1-x^3} - \frac{1}{2} \log(x^3) \right)$$

$$\downarrow 16$$

$$\frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{1-x^3} + 1} d\sqrt[3]{1-x^3} + 3\sqrt[3]{1-x^3} - \frac{1}{2} \log(x^3) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^3}) \right)$$

$$\downarrow 1083$$

$$\frac{1}{3} \left(3 \int \frac{1}{-x^6 - 3} d(2\sqrt[3]{1-x^3} + 1) + 3\sqrt[3]{1-x^3} - \frac{1}{2} \log(x^3) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^3}) \right)$$

$$\downarrow 217$$

$$\frac{1}{3} \left(-\sqrt{3} \arctan \left(\frac{2\sqrt[3]{1-x^3}+1}{\sqrt{3}} \right) + 3\sqrt[3]{1-x^3} - \frac{\log(x^3)}{2} + \frac{3}{2} \log \left(1 - \sqrt[3]{1-x^3} \right) \right)$$

input `Int[(1 - x^3)^(1/3)/x,x]`

output `(3*(1 - x^3)^(1/3) - Sqrt[3]*ArcTan[(1 + 2*(1 - x^3)^(1/3))/Sqrt[3]] - Log[x^3]/2 + (3*Log[1 - (1 - x^3)^(1/3)])/2)/3`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 69 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2), x] + (-Simp[3/(2*b*q) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q^2) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 1083

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 1.95 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.73

method	result
meijerg	$-\frac{-3\left(3 + \frac{\pi\sqrt{3}}{6} - \frac{3\ln(3)}{2} + 3\ln(x) + i\pi\right)\Gamma\left(\frac{2}{3}\right) + \Gamma\left(\frac{2}{3}\right)x^3 \operatorname{hypergeom}\left(\left[\frac{2}{3}, 1, 1\right], [2, 2], x^3\right)}{9\Gamma\left(\frac{2}{3}\right)}$
pseudoelliptic	$(-x^3 + 1)^{\frac{1}{3}} - \frac{\ln\left((-x^3 + 1)^{\frac{2}{3}} + (-x^3 + 1)^{\frac{1}{3}} + 1\right)}{6} - \frac{\arctan\left(\frac{\left(1 + 2(-x^3 + 1)^{\frac{1}{3}}\right)\sqrt{3}}{3}\right)\sqrt{3}}{3} + \frac{\ln\left((-x^3 + 1)^{\frac{1}{3}} - 1\right)}{3}$
trager	$(-x^3 + 1)^{\frac{1}{3}} + \frac{\operatorname{RootOf}(_Z^2 + _Z + 1) \ln\left(-\frac{1438 \operatorname{RootOf}(_Z^2 + _Z + 1)^2 x^3 - 6979 \operatorname{RootOf}(_Z^2 + _Z + 1) x^3 + 5502}{\dots}\right)}{\dots}$

input

```
int((-x^3+1)^(1/3)/x,x,method=_RETURNVERBOSE)
```

output

```
-1/9/GAMMA(2/3)*(-3*(3+1/6*Pi*3^(1/2)-3/2*ln(3)+3*ln(x)+I*Pi)*GAMMA(2/3)+GAMMA(2/3)*x^3*hypergeom([2/3,1,1],[2,2],x^3))
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.09

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = -\frac{1}{3}\sqrt{3}\arctan\left(\frac{2}{3}\sqrt{3}(-x^3+1)^{\frac{1}{3}} + \frac{1}{3}\sqrt{3}\right) + (-x^3+1)^{\frac{1}{3}} - \frac{1}{6}\log\left((-x^3+1)^{\frac{2}{3}} + (-x^3+1)^{\frac{1}{3}} + 1\right) + \frac{1}{3}\log\left((-x^3+1)^{\frac{1}{3}} - 1\right)$$

input

```
integrate((-x^3+1)^(1/3)/x,x, algorithm="fricas")
```

output

```
-1/3*sqrt(3)*arctan(2/3*sqrt(3)*(-x^3 + 1)^(1/3) + 1/3*sqrt(3)) + (-x^3 +
1)^(1/3) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log((-x^
3 + 1)^(1/3) - 1)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.55

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = -\frac{x e^{\frac{i\pi}{3}} \Gamma(-\frac{1}{3}) {}_2F_1\left(-\frac{1}{3}, -\frac{1}{3} \middle| \frac{1}{x^3}\right)}{3\Gamma(\frac{2}{3})}$$

input

```
integrate((-x**3+1)**(1/3)/x,x)
```

output

```
-x*exp(I*pi/3)*gamma(-1/3)*hyper((-1/3, -1/3), (2/3,), x**(-3))/(3*gamma(2
/3))
```

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.06

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = -\frac{1}{3}\sqrt{3}\arctan\left(\frac{1}{3}\sqrt{3}\left(2(-x^3+1)^{\frac{1}{3}}+1\right)\right) + (-x^3+1)^{\frac{1}{3}} \\ -\frac{1}{6}\log\left((-x^3+1)^{\frac{2}{3}}+(-x^3+1)^{\frac{1}{3}}+1\right) + \frac{1}{3}\log\left((-x^3+1)^{\frac{1}{3}}-1\right)$$

input

```
integrate((-x^3+1)^(1/3)/x,x, algorithm="maxima")
```

output

```
-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3) + 1)) + (-x^3 + 1)^(1/
3) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log((-x^3 + 1)
^(1/3) - 1)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.07

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = -\frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \left(2(-x^3+1)^{\frac{1}{3}} + 1 \right) \right) + (-x^3+1)^{\frac{1}{3}} \\ - \frac{1}{6} \log \left((-x^3+1)^{\frac{2}{3}} + (-x^3+1)^{\frac{1}{3}} + 1 \right) + \frac{1}{3} \log \left(\left| (-x^3+1)^{\frac{1}{3}} - 1 \right| \right)$$

input `integrate((-x^3+1)^(1/3)/x,x, algorithm="giac")`output `-1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3) + 1)) + (-x^3 + 1)^(1/3) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log(abs((-x^3 + 1)^(1/3) - 1))`**Mupad [B] (verification not implemented)**

Time = 0.21 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.24

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = \frac{\ln \left((1-x^3)^{1/3} - 1 \right)}{3} + \ln \left(3(1-x^3)^{1/3} + \frac{3}{2} - \frac{\sqrt{3}3i}{2} \right) \left(-\frac{1}{6} + \frac{\sqrt{3}1i}{6} \right) \\ - \ln \left(3(1-x^3)^{1/3} + \frac{3}{2} + \frac{\sqrt{3}3i}{2} \right) \left(\frac{1}{6} + \frac{\sqrt{3}1i}{6} \right) + (1-x^3)^{1/3}$$

input `int((1 - x^3)^(1/3)/x,x)`output `log((1 - x^3)^(1/3) - 1)/3 + log(3*(1 - x^3)^(1/3) - (3^(1/2)*3i)/2 + 3/2) * ((3^(1/2)*1i)/6 - 1/6) - log((3^(1/2)*3i)/2 + 3*(1 - x^3)^(1/3) + 3/2) * ((3^(1/2)*1i)/6 + 1/6) + (1 - x^3)^(1/3)`

Reduce [F]

$$\int \frac{\sqrt[3]{1-x^3}}{x} dx = (-x^3+1)^{\frac{1}{3}} - \left(\int \frac{(-x^3+1)^{\frac{1}{3}}}{x^4-x} dx \right)$$

input `int((-x^3+1)^(1/3)/x,x)`

output `(- x**3 + 1)**(1/3) - int((- x**3 + 1)**(1/3)/(x**4 - x),x)`

3.58 $\int \frac{\sqrt[3]{1-x^3}}{1+x} dx$

Optimal result	481
Mathematica [F]	482
Rubi [A] (verified)	482
Maple [F]	484
Fricas [F(-2)]	484
Sympy [F]	485
Maxima [F]	485
Giac [F]	485
Mupad [F(-1)]	486
Reduce [F]	486

Optimal result

Integrand size = 17, antiderivative size = 482

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = \sqrt[3]{1-x^3} + \frac{\sqrt[3]{2} \arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{1+\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{2^{2/3}\sqrt{3}}$$

$$- \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\sqrt[3]{2} \arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2x}}{\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} - \frac{\sqrt[3]{2} \arctan\left(\frac{1+2^{2/3}\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}}$$

$$- \frac{1}{3}\sqrt[3]{2} \log(1+x^3) + \frac{\log\left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}\right)}{3 \cdot 2^{2/3}} - \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{3 \cdot 2^{2/3}}$$

$$+ \frac{1}{3}\sqrt[3]{2} \log\left(1 + \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right) - \frac{\log\left(2\sqrt[3]{2} + \frac{(1-x)^2}{(1-x^3)^{2/3}} + \frac{2^{2/3}(1-x)}{\sqrt[3]{1-x^3}}\right)}{6 \cdot 2^{2/3}} + \frac{\log\left(\sqrt[3]{2} - \sqrt[3]{1-x^3}\right)}{2^{2/3}} - \frac{1}{2} \log(-x)$$

output

```
(-x^3+1)^(1/3)-1/3*2^(1/3)*ln(x^3+1)+1/6*ln(2^(2/3)+(-1+x)/(-x^3+1)^(1/3))
*2^(1/3)-1/6*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3)*(1-x)/(-x^3+1)^(1
/3))*2^(1/3)+1/3*2^(1/3)*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))-1/12*ln(2*2^(1
/3)+(1-x)^2/(-x^3+1)^(2/3)+2^(2/3)*(1-x)/(-x^3+1)^(1/3))*2^(1/3)+1/2*ln(2^
(1/3)-(-x^3+1)^(1/3))*2^(1/3)-1/2*ln(-x-(-x^3+1)^(1/3))+1/2*ln(-2^(1/3)*x-
(-x^3+1)^(1/3))*2^(1/3)+1/3*2^(1/3)*arctan(1/3*(1-2*2^(1/3)*(1-x)/(-x^3+1)
^(1/3))*3^(1/2))*3^(1/2)+1/6*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3
^(1/2))*2^(1/3)*3^(1/2)-1/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(1/2))*3^(
1/2)+1/3*2^(1/3)*arctan(1/3*(1-2*2^(1/3)*x/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2
)-1/3*2^(1/3)*arctan(1/3*(1+2^(2/3)*(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)
```

Mathematica [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = \int \frac{\sqrt[3]{1-x^3}}{1+x} dx$$

input

```
Integrate[(1 - x^3)^(1/3)/(1 + x), x]
```

output

```
Integrate[(1 - x^3)^(1/3)/(1 + x), x]
```

Rubi [A] (verified)

Time = 0.66 (sec) , antiderivative size = 482, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2581, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[3]{1-x^3}}{x+1} dx$$

↓ 2581

$$\int \left(-\frac{\sqrt[3]{1-x^3}x}{x^3+1} + \frac{\sqrt[3]{1-x^3}}{x^3+1} + \frac{\sqrt[3]{1-x^3}x^2}{x^3+1} \right) dx$$

$$\begin{aligned}
& \begin{array}{c} \downarrow \\ \text{2009} \end{array} \\
& \frac{\sqrt[3]{2} \arctan\left(\frac{1 - 2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{\sqrt[3]{2(1-x)} + 1}{\sqrt[3]{1-x^3}}\right)}{2^{2/3}\sqrt{3}} - \frac{\arctan\left(\frac{1 - \sqrt[3]{\frac{2x}{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \\
& \frac{\sqrt[3]{2} \arctan\left(\frac{1 - \frac{2\sqrt[3]{2x}}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\sqrt[3]{2} \arctan\left(\frac{2^{2/3}\sqrt[3]{1-x^3+1}}{\sqrt{3}}\right)}{\sqrt{3}} + \sqrt[3]{1-x^3} - \frac{1}{3}\sqrt[3]{2} \log(x^3 + 1) + \\
& \frac{\log\left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}\right)}{3 \cdot 2^{2/3}} - \frac{\log\left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1\right)}{3 \cdot 2^{2/3}} + \frac{1}{3}\sqrt[3]{2} \log\left(\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1\right) - \\
& \frac{\log\left(\frac{(1-x)^2}{(1-x^3)^{2/3}} + \frac{2^{2/3}(1-x)}{\sqrt[3]{1-x^3}} + 2\sqrt[3]{2}\right)}{6 \cdot 2^{2/3}} + \frac{\log\left(\sqrt[3]{2} - \sqrt[3]{1-x^3}\right)}{2^{2/3}} - \frac{1}{2} \log\left(-\sqrt[3]{1-x^3} - x\right) + \\
& \frac{\log\left(-\sqrt[3]{1-x^3} - \sqrt[3]{2x}\right)}{2^{2/3}}
\end{aligned}$$

input `Int[(1 - x^3)^(1/3)/(1 + x),x]`

output

$$\begin{aligned}
& (1 - x^3)^{1/3} + (2^{1/3} \operatorname{ArcTan}[(1 - (2 \cdot 2^{1/3})(1 - x))/(1 - x^3)^{1/3}]) / \sqrt{3} \\
& + \operatorname{ArcTan}[(1 + (2^{1/3})(1 - x))/(1 - x^3)^{1/3}] / \sqrt{3} - \operatorname{ArcTan}[(1 - (2x)/(1 - x^3)^{1/3}) / \sqrt{3}] / \sqrt{3} \\
& + (2^{1/3} \operatorname{ArcTan}[(1 - (2 \cdot 2^{1/3}x)/(1 - x^3)^{1/3}) / \sqrt{3}]) / \sqrt{3} \\
& - (2^{1/3} \operatorname{ArcTan}[(1 + 2^{2/3}(1 - x^3)^{1/3}) / \sqrt{3}]) / \sqrt{3} - (2^{1/3} \operatorname{Log}[1 + x^3]) / 3 \\
& + \operatorname{Log}[2^{2/3} - (1 - x)/(1 - x^3)^{1/3}] / (3 \cdot 2^{2/3}) - \operatorname{Log}[1 + (2^{2/3}(1 - x)^2) / (1 - x^3)^{2/3} - (2^{1/3}(1 - x)) / (1 - x^3)^{1/3}] / (3 \cdot 2^{2/3}) \\
& + (2^{1/3} \operatorname{Log}[1 + (2^{1/3}(1 - x)) / (1 - x^3)^{1/3}]) / 3 - \operatorname{Log}[2 \cdot 2^{1/3} + (1 - x)^2 / (1 - x^3)^{2/3} + (2^{2/3}(1 - x)) / (1 - x^3)^{1/3}] / (6 \cdot 2^{2/3}) \\
& + \operatorname{Log}[2^{1/3} - (1 - x^3)^{1/3}] / 2^{2/3} - \operatorname{Log}[-x - (1 - x^3)^{1/3}] / 2 + \operatorname{Log}[-(2^{1/3}x) - (1 - x^3)^{1/3}] / 2^{2/3}
\end{aligned}$$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] :> Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2581 `Int[(Px_)*((c_) + (d_)*(x_))^(q_)*((a_) + (b_)*(x_)^3)^(p_), x_Symbol] :> Int[ExpandIntegrand[(c^3 + d^3*x^3)^q*(a + b*x^3)^p, Px/(c^2 - c*d*x + d^2*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p}, x] && PolyQ[Px, x] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [F]

$$\int \frac{(-x^3 + 1)^{\frac{1}{3}}}{1 + x} dx$$

input `int((-x^3+1)^(1/3)/(1+x),x)`

output `int((-x^3+1)^(1/3)/(1+x),x)`

Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = \text{Exception raised: TypeError}$$

input `integrate((-x^3+1)^(1/3)/(1+x),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (residue poly has multiple non-linear factors)`

Sympy [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = \int \frac{\sqrt[3]{-(x-1)(x^2+x+1)}}{x+1} dx$$

input `integrate((-x**3+1)**(1/3)/(1+x),x)`

output `Integral((-x - 1)*(x**2 + x + 1)**(1/3)/(x + 1), x)`

Maxima [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x+1} dx$$

input `integrate((-x^3+1)^(1/3)/(1+x),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(1/3)/(x + 1), x)`

Giac [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x+1} dx$$

input `integrate((-x^3+1)^(1/3)/(1+x),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(1/3)/(x + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = \int \frac{(1-x^3)^{1/3}}{x+1} dx$$

input `int((1 - x^3)^(1/3)/(x + 1),x)`output `int((1 - x^3)^(1/3)/(x + 1), x)`**Reduce [F]**

$$\int \frac{\sqrt[3]{1-x^3}}{1+x} dx = (-x^3 + 1)^{\frac{1}{3}} - \left(\int \frac{(-x^3 + 1)^{\frac{1}{3}}}{x^4 + x^3 - x - 1} dx \right) - \left(\int \frac{(-x^3 + 1)^{\frac{1}{3}} x^2}{x^4 + x^3 - x - 1} dx \right)$$

input `int((-x^3+1)^(1/3)/(1+x),x)`output `(- x**3 + 1)**(1/3) - int((- x**3 + 1)**(1/3)/(x**4 + x**3 - x - 1),x) -
int(((- x**3 + 1)**(1/3)*x**2)/(x**4 + x**3 - x - 1),x)`

3.59 $\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx$

Optimal result	487
Mathematica [F]	488
Rubi [A] (verified)	488
Maple [C] (verified)	490
Fricas [B] (verification not implemented)	491
Sympy [F]	492
Maxima [F]	492
Giac [F]	492
Mupad [F(-1)]	493
Reduce [F]	493

Optimal result

Integrand size = 22, antiderivative size = 280

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \frac{\sqrt{3} \arctan\left(\frac{1+\frac{2\sqrt[3]{2}(-1+x)}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{2^{2/3}} + \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{1-\frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{2^{2/3}\sqrt{3}} - \frac{\arctan\left(\frac{1+2^{2/3}\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{2^{2/3}\sqrt{3}} - \frac{\log(-3(-1+x)(1-x+x^2))}{2 \cdot 2^{2/3}} + \frac{\log(\sqrt[3]{2}-\sqrt[3]{1-x^3})}{2 \cdot 2^{2/3}} + \frac{3 \log(-\sqrt[3]{2}(-1+x)+\sqrt[3]{1-x^3})}{2 \cdot 2^{2/3}} + \frac{1}{2} \log(x+\sqrt[3]{1-x^3}) - \frac{\log(\sqrt[3]{2}x+\sqrt[3]{1-x^3})}{2 \cdot 2^{2/3}}$$

output

```
-1/4*ln(-3*(-1+x)*(x^2-x+1))*2^(1/3)+1/4*ln(2^(1/3)-(-x^3+1)^(1/3))*2^(1/3)
)+3/4*ln(-2^(1/3)*(-1+x)+(-x^3+1)^(1/3))*2^(1/3)+1/2*ln(x+(-x^3+1)^(1/3))-
1/4*ln(2^(1/3)*x+(-x^3+1)^(1/3))*2^(1/3)+1/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))
)*3^(1/2))*3^(1/2)-1/6*2^(1/3)*arctan(1/3*(1-2*2^(1/3)*x/(-x^3+1)^(1/3))
)*3^(1/2))*3^(1/2)-1/6*2^(1/3)*arctan(1/3*(1+2^(2/3)*(-x^3+1)^(1/3))*3^(1/2)
)*3^(1/2)+1/2*arctan(1/3*(1+2*2^(1/3)*(-1+x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)
)*2^(1/3)
```

Mathematica [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx$$

input

```
Integrate[(1 - x^3)^(1/3)/(1 - x + x^2), x]
```

output

```
Integrate[(1 - x^3)^(1/3)/(1 - x + x^2), x]
```

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 408, normalized size of antiderivative = 1.46, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[3]{1-x^3}}{x^2-x+1} dx$$

↓ 2583

$$\int \left(\frac{\sqrt[3]{1-x^3}x}{x^3+1} + \frac{\sqrt[3]{1-x^3}}{x^3+1} \right) dx$$

↓ 2009

$$\begin{aligned}
& \frac{\sqrt[3]{2} \arctan\left(\frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1}{\sqrt{3}}\right)}{2^{2/3}\sqrt{3}} + \frac{\arctan\left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \\
& \frac{\sqrt[3]{2} \arctan\left(\frac{1 - \frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\log(x^3 + 1)}{3 \cdot 2^{2/3}} + \frac{\log\left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}\right)}{3 \cdot 2^{2/3}} - \\
& \frac{\log\left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1\right)}{3 \cdot 2^{2/3}} + \frac{1}{3} \sqrt[3]{2} \log\left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1\right) - \\
& \frac{\log\left(\frac{(1-x)^2}{(1-x^3)^{2/3}} + \frac{2^{2/3}(1-x)}{\sqrt[3]{1-x^3}} + 2\sqrt[3]{2}\right)}{6 \cdot 2^{2/3}} + \frac{1}{2} \log\left(-\sqrt[3]{1-x^3} - x\right) - \frac{\log\left(-\sqrt[3]{1-x^3} - \sqrt[3]{2}x\right)}{2^{2/3}}
\end{aligned}$$

input `Int[(1 - x^3)^(1/3)/(1 - x + x^2), x]`

output `(2^(1/3)*ArcTan[(1 - (2*2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(2/3)*Sqrt[3]) + ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - (2^(1/3)*ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + Log[1 + x^3]/(3*2^(2/3)) + Log[2^(2/3) - (1 - x)/(1 - x^3)^(1/3)]/(3*2^(2/3)) - Log[1 + (2^(2/3)*(1 - x)^2)/(1 - x^3)^(2/3) - (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(3*2^(2/3)) + (2^(1/3)*Log[1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)])/3 - Log[2*2^(1/3) + (1 - x)^2/(1 - x^3)^(2/3) + (2^(2/3)*(1 - x))/(1 - x^3)^(1/3)]/(6*2^(2/3)) + Log[-x - (1 - x^3)^(1/3)]/2 - Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)]/2^(2/3)`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2583 `Int[(Px_.)*((c_) + (d_)*(x_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^3)^(p_.), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 9.35 (sec) , antiderivative size = 925, normalized size of antiderivative = 3.30

method	result	size
trager	Expression too large to display	925

input `int((-x^3+1)^(1/3)/(x^2-x+1),x,method=_RETURNVERBOSE)`

output

```

-1/6*RootOf(_Z^6+108)*ln(-(RootOf(_Z^6+108)^5*x^4-2*RootOf(_Z^6+108)^4*(-x
^3+1)^(1/3)*x^3+2*RootOf(_Z^6+108)^5*x^3+6*RootOf(_Z^6+108)^4*(-x^3+1)^(1/
3)*x^2-x^2*RootOf(_Z^6+108)^5-2*RootOf(_Z^6+108)^4*(-x^3+1)^(1/3)*x-2*Root
Of(_Z^6+108)^5*x-6*RootOf(_Z^6+108)^2*x^4+36*RootOf(_Z^6+108)*(-x^3+1)^(1/
3)*x^3+RootOf(_Z^6+108)^5-12*RootOf(_Z^6+108)^2*x^3+144*(-x^3+1)^(2/3)*x^2
-108*RootOf(_Z^6+108)*(-x^3+1)^(1/3)*x^2+6*x^2*RootOf(_Z^6+108)^2-144*x*(-
x^3+1)^(2/3)+36*RootOf(_Z^6+108)*(-x^3+1)^(1/3)*x+12*RootOf(_Z^6+108)^2*x-
6*RootOf(_Z^6+108)^2)/(x^2-x+1)^2)+1/72*ln(-(-3*RootOf(_Z^6+108)^4*(-x^3+1
)^(1/3)*x^3+RootOf(_Z^6+108)^4*(-x^3+1)^(1/3)*x^2+RootOf(_Z^6+108)^4*(-x^3
+1)^(1/3)*x-15*RootOf(_Z^6+108)^2*x^4+6*RootOf(_Z^6+108)^2*x^3+72*(-x^3+1)
^(2/3)*x^2+3*x^2*RootOf(_Z^6+108)^2-36*x*(-x^3+1)^(2/3)+6*RootOf(_Z^6+108)
^2*x-3*RootOf(_Z^6+108)^2)/(x^2-x+1)^2)*RootOf(_Z^6+108)^4+1/12*ln(-(-3*Ro
otOf(_Z^6+108)^4*(-x^3+1)^(1/3)*x^3+RootOf(_Z^6+108)^4*(-x^3+1)^(1/3)*x^2+
RootOf(_Z^6+108)^4*(-x^3+1)^(1/3)*x-15*RootOf(_Z^6+108)^2*x^4+6*RootOf(_Z^
6+108)^2*x^3+72*(-x^3+1)^(2/3)*x^2+3*x^2*RootOf(_Z^6+108)^2-36*x*(-x^3+1)^(
2/3)+6*RootOf(_Z^6+108)^2*x-3*RootOf(_Z^6+108)^2)/(x^2-x+1)^2)*RootOf(_Z^
6+108)+1/3*ln(-RootOf(_Z^6+108)^6*x^3-18*(-x^3+1)^(2/3)*RootOf(_Z^6+108)^3
*x+18*RootOf(_Z^6+108)^3*x^3+108*x*(-x^3+1)^(2/3)+216*x^2*(-x^3+1)^(1/3)-1
2*RootOf(_Z^6+108)^3)+1/36*ln(RootOf(_Z^6+108)^6*x^3+36*(-x^3+1)^(2/3)*Roo
tOf(_Z^6+108)^3*x+36*RootOf(_Z^6+108)^3*(-x^3+1)^(1/3)*x^2+216*x*(-x^3+...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3744 vs. $2(218) = 436$.

Time = 5.06 (sec) , antiderivative size = 3744, normalized size of antiderivative = 13.37

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \text{Too large to display}$$

input

```
integrate((-x^3+1)^(1/3)/(x^2-x+1),x, algorithm="fricas")
```

output

```
Too large to include
```

Sympy [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \int \frac{\sqrt[3]{-(x-1)(x^2+x+1)}}{x^2-x+1} dx$$

input `integrate((-x**3+1)**(1/3)/(x**2-x+1),x)`

output `Integral((-x - 1)*(x**2 + x + 1)**(1/3)/(x**2 - x + 1), x)`

Maxima [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x^2-x+1} dx$$

input `integrate((-x^3+1)^(1/3)/(x^2-x+1),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(1/3)/(x^2 - x + 1), x)`

Giac [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x^2-x+1} dx$$

input `integrate((-x^3+1)^(1/3)/(x^2-x+1),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(1/3)/(x^2 - x + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \int \frac{(1-x^3)^{1/3}}{x^2-x+1} dx$$

input `int((1 - x^3)^(1/3)/(x^2 - x + 1),x)`output `int((1 - x^3)^(1/3)/(x^2 - x + 1), x)`**Reduce [F]**

$$\int \frac{\sqrt[3]{1-x^3}}{1-x+x^2} dx = \int \frac{(-x^3+1)^{1/3}}{x^2-x+1} dx$$

input `int((-x^3+1)^(1/3)/(x^2-x+1),x)`output `int((-x**3 + 1)**(1/3)/(x**2 - x + 1),x)`

3.60 $\int \frac{\sqrt[3]{1-x^3}}{2+x} dx$

Optimal result	494
Mathematica [F]	495
Rubi [A] (verified)	495
Maple [F]	496
Fricas [F(-2)]	497
Sympy [F]	497
Maxima [F]	497
Giac [F]	498
Mupad [F(-1)]	498
Reduce [F]	498

Optimal result

Integrand size = 17, antiderivative size = 232

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx$$

$$= \sqrt[3]{1-x^3} + \frac{1}{2}x \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{1}{3}, 1, \frac{4}{3}, x^3, -\frac{x^3}{8}\right) - \frac{2 \arctan\left(\frac{1-\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}}$$

$$+ \sqrt[6]{3} \arctan\left(\frac{1-\frac{3^{2/3}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right) - \sqrt[6]{3} \arctan\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-x^3}}{3\sqrt[6]{3}}\right) - \frac{\log(8+x^3)}{\sqrt[3]{3}}$$

$$+ \frac{1}{2}3^{2/3} \log\left(3^{2/3}-\sqrt[3]{1-x^3}\right) - \log\left(-x-\sqrt[3]{1-x^3}\right) + \frac{1}{2}3^{2/3} \log\left(-\frac{1}{2}3^{2/3}x-\sqrt[3]{1-x^3}\right)$$

output

```
(-x^3+1)^(1/3)+1/2*x*AppellF1(1/3,-1/3,1,4/3,x^3,-1/8*x^3)-3^(1/6)*arctan(
2/9*(-x^3+1)^(1/3)*3^(5/6)+1/3*3^(1/2))+3^(1/6)*arctan(1/3*(1-3^(2/3)*x/(-
x^3+1)^(1/3))*3^(1/2))-1/3*ln(x^3+8)*3^(2/3)+1/2*3^(2/3)*ln(3^(2/3)-(-x^3+
1)^(1/3))-ln(-x-(-x^3+1)^(1/3))+1/2*3^(2/3)*ln(-1/2*3^(2/3)*x-(-x^3+1)^(1/
3))-2/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)
```

Mathematica [F]

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx = \int \frac{\sqrt[3]{1-x^3}}{2+x} dx$$

input `Integrate[(1 - x^3)^(1/3)/(2 + x), x]`

output `Integrate[(1 - x^3)^(1/3)/(2 + x), x]`

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2581, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{\sqrt[3]{1-x^3}}{x+2} dx \\ & \quad \downarrow \text{2581} \\ & \int \left(-\frac{2\sqrt[3]{1-x^3}x}{x^3+8} + \frac{4\sqrt[3]{1-x^3}}{x^3+8} + \frac{\sqrt[3]{1-x^3}x^2}{x^3+8} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{1}{2}x \operatorname{AppellF1} \left(\frac{1}{3}, -\frac{1}{3}, 1, \frac{4}{3}, x^3, -\frac{x^3}{8} \right) - \frac{2 \arctan \left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \\ & \sqrt[6]{3} \arctan \left(\frac{1 - \frac{3^{2/3}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right) - \sqrt[6]{3} \arctan \left(\frac{2\sqrt[3]{1-x^3}}{3\sqrt[6]{3}} + \frac{1}{\sqrt{3}} \right) + \sqrt[3]{1-x^3} - \frac{\log(x^3+8)}{\sqrt[3]{3}} + \\ & \frac{1}{2}3^{2/3} \log \left(3^{2/3} - \sqrt[3]{1-x^3} \right) - \log \left(-\sqrt[3]{1-x^3} - x \right) + \frac{1}{2}3^{2/3} \log \left(-\sqrt[3]{1-x^3} - \frac{1}{2}3^{2/3}x \right) \end{aligned}$$

input `Int[(1 - x^3)^(1/3)/(2 + x),x]`

output `(1 - x^3)^(1/3) + (x*AppellF1[1/3, -1/3, 1, 4/3, x^3, -1/8*x^3])/2 - (2*ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + 3^(1/6)*ArcTan[(1 - (3^(2/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]] - 3^(1/6)*ArcTan[1/Sqrt[3] + (2*(1 - x^3)^(1/3))/(3*3^(1/6))]) - Log[8 + x^3]/3^(1/3) + (3^(2/3)*Log[3^(2/3) - (1 - x^3)^(1/3)])/2 - Log[-x - (1 - x^3)^(1/3)] + (3^(2/3)*Log[-1/2*(3^(2/3)*x) - (1 - x^3)^(1/3)])/2`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2581 `Int[(Px_)*((c_) + (d_)*(x_))^(q_)*((a_) + (b_)*(x_)^3)^(p_), x_Symbol] := Int[ExpandIntegrand[(c^3 + d^3*x^3)^q*(a + b*x^3)^p, Px/(c^2 - c*d*x + d^2*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p}, x] && PolyQ[Px, x] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [F]

$$\int \frac{(-x^3 + 1)^{\frac{1}{3}}}{2 + x} dx$$

input `int((-x^3+1)^(1/3)/(2+x),x)`

output `int((-x^3+1)^(1/3)/(2+x),x)`

Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx = \text{Exception raised: TypeError}$$

input `integrate((-x^3+1)^(1/3)/(2+x),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (residue poly has multiple non-linear factors)`

Sympy [F]

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx = \int \frac{\sqrt[3]{-(x-1)(x^2+x+1)}}{x+2} dx$$

input `integrate((-x**3+1)**(1/3)/(2+x),x)`

output `Integral((-x - 1)*(x**2 + x + 1)**(1/3)/(x + 2), x)`

Maxima [F]

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x+2} dx$$

input `integrate((-x^3+1)^(1/3)/(2+x),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(1/3)/(x + 2), x)`

Giac [F]

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x+2} dx$$

input `integrate((-x^3+1)^(1/3)/(2+x),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(1/3)/(x + 2), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx = \int \frac{(1-x^3)^{1/3}}{x+2} dx$$

input `int((1 - x^3)^(1/3)/(x + 2),x)`

output `int((1 - x^3)^(1/3)/(x + 2), x)`

Reduce [F]

$$\int \frac{\sqrt[3]{1-x^3}}{2+x} dx = (-x^3+1)^{\frac{1}{3}} - \left(\int \frac{(-x^3+1)^{\frac{1}{3}}}{x^4+2x^3-x-2} dx \right) - 2 \left(\int \frac{(-x^3+1)^{\frac{1}{3}} x^2}{x^4+2x^3-x-2} dx \right)$$

input `int((-x^3+1)^(1/3)/(2+x),x)`

output `(- x**3 + 1)**(1/3) - int((- x**3 + 1)**(1/3)/(x**4 + 2*x**3 - x - 2),x) - 2*int(((- x**3 + 1)**(1/3)*x**2)/(x**4 + 2*x**3 - x - 2),x)`

3.61 $\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx$

Optimal result	499
Mathematica [F]	500
Rubi [A] (verified)	500
Maple [F]	501
Fricas [F]	501
Sympy [F]	502
Maxima [F]	502
Giac [F]	502
Mupad [F(-1)]	503
Reduce [F]	503

Optimal result

Integrand size = 21, antiderivative size = 168

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = -\frac{x^2 \operatorname{AppellF1}\left(\frac{2}{3}, 1, \frac{1}{3}, \frac{5}{3}, x^3, -\frac{x^3}{2}\right)}{2\sqrt[3]{2}}$$

$$+ \frac{2 \arctan\left(\frac{1+\frac{2\sqrt[3]{3}x}{\sqrt[3]{2+x^3}}}{\sqrt{3}}\right)}{3^{5/6}}$$

$$+ \frac{\arctan\left(\frac{\sqrt[3]{3}+2\sqrt[3]{2+x^3}}{3^{5/6}}\right)}{3^{5/6}} + \frac{\log(1-x^3)}{6\sqrt[3]{3}}$$

$$+ \frac{\log(\sqrt[3]{3}-\sqrt[3]{2+x^3})}{2\sqrt[3]{3}} - \frac{\log(\sqrt[3]{3}x-\sqrt[3]{2+x^3})}{\sqrt[3]{3}}$$

output

```
-1/4*x^2*AppellF1(2/3,1,1/3,5/3,x^3,-1/2*x^3)*2^(2/3)+1/3*arctan(1/3*(3^(1/3)+2*(x^3+2)^(1/3))*3^(1/6))*3^(1/6)+2/3*arctan(1/3*(1+2*3^(1/3)*x/(x^3+2)^(1/3))*3^(1/2))*3^(1/6)+1/18*ln(-x^3+1)*3^(2/3)+1/6*ln(3^(1/3)-(x^3+2)^(1/3))*3^(2/3)-1/3*ln(3^(1/3)*x-(x^3+2)^(1/3))*3^(2/3)
```

Mathematica [F]

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = \int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx$$

input `Integrate[(2 + x)/((1 + x + x^2)*(2 + x^3)^(1/3)), x]`

output `Integrate[(2 + x)/((1 + x + x^2)*(2 + x^3)^(1/3)), x]`

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+2}{(x^2+x+1)\sqrt[3]{x^3+2}} dx$$

↓ 2583

$$\int \left(-\frac{x}{(1-x^3)\sqrt[3]{x^3+2}} + \frac{2}{(1-x^3)\sqrt[3]{x^3+2}} - \frac{x^2}{(1-x^3)\sqrt[3]{x^3+2}} \right) dx$$

↓ 2009

$$-\frac{x^2 \operatorname{AppellF1}\left(\frac{2}{3}, 1, \frac{1}{3}, \frac{5}{3}, x^3, -\frac{x^3}{2}\right)}{2\sqrt[3]{2}} + \frac{2 \arctan\left(\frac{\frac{2\sqrt[3]{3}x}{\sqrt[3]{x^3+2}}+1}{\sqrt{3}}\right)}{3^{5/6}} + \frac{\arctan\left(\frac{2\sqrt[3]{x^3+2}+\sqrt[3]{3}}{3^{5/6}}\right)}{3^{5/6}} + \frac{\log(1-x^3)}{6\sqrt[3]{3}} + \frac{\log\left(\sqrt[3]{3}-\sqrt[3]{x^3+2}\right)}{2\sqrt[3]{3}} - \frac{\log\left(\sqrt[3]{3}x-\sqrt[3]{x^3+2}\right)}{\sqrt[3]{3}}$$

input `Int[(2 + x)/((1 + x + x^2)*(2 + x^3)^(1/3)), x]`

output

```
-1/2*(x^2*AppellF1[2/3, 1, 1/3, 5/3, x^3, -1/2*x^3])/2^(1/3) + (2*ArcTan[(1 + (2*3^(1/3)*x)/(2 + x^3)^(1/3))/Sqrt[3]])/3^(5/6) + ArcTan[(3^(1/3) + 2*(2 + x^3)^(1/3))/3^(5/6)]/3^(5/6) + Log[1 - x^3]/(6*3^(1/3)) + Log[3^(1/3) - (2 + x^3)^(1/3)]/(2*3^(1/3)) - Log[3^(1/3)*x - (2 + x^3)^(1/3)]/3^(1/3)
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2583

```
Int[(Px_)*((c_) + (d_)*(x_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^3)^(p_), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]
```

Maple [F]

$$\int \frac{2+x}{(x^2+x+1)(x^3+2)^{\frac{1}{3}}} dx$$

input

```
int((2+x)/(x^2+x+1)/(x^3+2)^(1/3),x)
```

output

```
int((2+x)/(x^2+x+1)/(x^3+2)^(1/3),x)
```

Fricas [F]

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = \int \frac{x+2}{(x^3+2)^{\frac{1}{3}}(x^2+x+1)} dx$$

input

```
integrate((2+x)/(x^2+x+1)/(x^3+2)^(1/3),x, algorithm="fricas")
```

output `integral((x^3 + 2)^(2/3)*(x + 2)/(x^5 + x^4 + x^3 + 2*x^2 + 2*x + 2), x)`

Sympy [F]

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = \int \frac{x+2}{\sqrt[3]{x^3+2}(x^2+x+1)} dx$$

input `integrate((2+x)/(x**2+x+1)/(x**3+2)**(1/3), x)`

output `Integral((x + 2)/((x**3 + 2)**(1/3)*(x**2 + x + 1)), x)`

Maxima [F]

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = \int \frac{x+2}{(x^3+2)^{\frac{1}{3}}(x^2+x+1)} dx$$

input `integrate((2+x)/(x^2+x+1)/(x^3+2)^(1/3), x, algorithm="maxima")`

output `integrate((x + 2)/((x^3 + 2)^(1/3)*(x^2 + x + 1)), x)`

Giac [F]

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = \int \frac{x+2}{(x^3+2)^{\frac{1}{3}}(x^2+x+1)} dx$$

input `integrate((2+x)/(x^2+x+1)/(x^3+2)^(1/3), x, algorithm="giac")`

output `integrate((x + 2)/((x^3 + 2)^(1/3)*(x^2 + x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = \int \frac{x+2}{(x^3+2)^{1/3}(x^2+x+1)} dx$$

input `int((x + 2)/((x^3 + 2)^(1/3)*(x + x^2 + 1)), x)`output `int((x + 2)/((x^3 + 2)^(1/3)*(x + x^2 + 1)), x)`**Reduce [F]**

$$\int \frac{2+x}{(1+x+x^2)\sqrt[3]{2+x^3}} dx = \int \frac{x}{(x^3+2)^{\frac{1}{3}}x^2 + (x^3+2)^{\frac{1}{3}}x + (x^3+2)^{\frac{1}{3}}} dx + 2 \left(\int \frac{1}{(x^3+2)^{\frac{1}{3}}x^2 + (x^3+2)^{\frac{1}{3}}x + (x^3+2)^{\frac{1}{3}}} dx \right)$$

input `int((2+x)/(x^2+x+1)/(x^3+2)^(1/3), x)`output `int(x/((x**3 + 2)**(1/3)*x**2 + (x**3 + 2)**(1/3)*x + (x**3 + 2)**(1/3)), x) + 2*int(1/((x**3 + 2)**(1/3)*x**2 + (x**3 + 2)**(1/3)*x + (x**3 + 2)**(1/3)), x)`

$$3.62 \quad \int \frac{3-3x+30x^2+160x^3}{9+24x-12x^2+80x^3+320x^4} dx$$

Optimal result	504
Mathematica [A] (verified)	504
Rubi [A] (verified)	505
Maple [A] (verified)	505
Fricas [A] (verification not implemented)	506
Sympy [A] (verification not implemented)	506
Maxima [A] (verification not implemented)	507
Giac [A] (verification not implemented)	507
Mupad [B] (verification not implemented)	508
Reduce [B] (verification not implemented)	508

Optimal result

Integrand size = 38, antiderivative size = 25

$$\int \frac{3-3x+30x^2+160x^3}{9+24x-12x^2+80x^3+320x^4} dx = \frac{1}{8} \log(9+24x-12x^2+80x^3+320x^4)$$

output `1/8*ln(320*x^4+80*x^3-12*x^2+24*x+9)`

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{3-3x+30x^2+160x^3}{9+24x-12x^2+80x^3+320x^4} dx = \frac{1}{8} \log(9+24x-12x^2+80x^3+320x^4)$$

input `Integrate[(3 - 3*x + 30*x^2 + 160*x^3)/(9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4), x]`

output `Log[9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4]/8`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.026$, Rules used = {2020}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{160x^3 + 30x^2 - 3x + 3}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx$$

↓ 2020

$$\frac{1}{8} \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

input `Int[(3 - 3*x + 30*x^2 + 160*x^3)/(9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4),x]`

output `Log[9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4]/8`

Defintions of rubi rules used

rule 2020 `Int[(Pp_)/(Qq_), x_Symbol] :> With[{p = Expon[Pp, x], q = Expon[Qq, x]}, Simp[Coeff[Pp, x, p]*(Log[RemoveContent[Qq, x]]/(q*Coeff[Qq, x, q])), x] /; EqQ[p, q - 1] && EqQ[Pp, Simplify[(Coeff[Pp, x, p]/(q*Coeff[Qq, x, q]))*D[Qq, x]]] /; PolyQ[Pp, x] && PolyQ[Qq, x]`

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
parallelrisch	$\frac{\ln(x^4 + \frac{1}{4}x^3 - \frac{3}{80}x^2 + \frac{3}{40}x + \frac{9}{320})}{8}$	22
default	$\frac{\ln(320x^4 + 80x^3 - 12x^2 + 24x + 9)}{8}$	24
norman	$\frac{\ln(320x^4 + 80x^3 - 12x^2 + 24x + 9)}{8}$	24
risch	$\frac{\ln(320x^4 + 80x^3 - 12x^2 + 24x + 9)}{8}$	24

input `int((160*x^3+30*x^2-3*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x,method=_RETURN
VERBOSE)`

output `1/8*ln(x^4+1/4*x^3-3/80*x^2+3/40*x+9/320)`

Fricas [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{3 - 3x + 30x^2 + 160x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{1}{8} \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

input `integrate((160*x^3+30*x^2-3*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x, algorit
hm="fricas")`

output `1/8*log(320*x^4 + 80*x^3 - 12*x^2 + 24*x + 9)`

Sympy [A] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{3 - 3x + 30x^2 + 160x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{\log(320x^4 + 80x^3 - 12x^2 + 24x + 9)}{8}$$

input `integrate((160*x**3+30*x**2-3*x+3)/(320*x**4+80*x**3-12*x**2+24*x+9),x)`

output $\log(320x^4 + 80x^3 - 12x^2 + 24x + 9)/8$

Maxima [A] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{3 - 3x + 30x^2 + 160x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{1}{8} \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

input `integrate((160*x^3+30*x^2-3*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x, algorithm="maxima")`

output $1/8*\log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{3 - 3x + 30x^2 + 160x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{1}{8} \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

input `integrate((160*x^3+30*x^2-3*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x, algorithm="giac")`

output $1/8*\log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{3 - 3x + 30x^2 + 160x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{\ln(320x^4 + 80x^3 - 12x^2 + 24x + 9)}{8}$$

input `int((30*x^2 - 3*x + 160*x^3 + 3)/(24*x - 12*x^2 + 80*x^3 + 320*x^4 + 9),x)`output `log(24*x - 12*x^2 + 80*x^3 + 320*x^4 + 9)/8`**Reduce [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \frac{3 - 3x + 30x^2 + 160x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{\log(320x^4 + 80x^3 - 12x^2 + 24x + 9)}{8}$$

input `int((160*x^3+30*x^2-3*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x)`output `log(320*x**4 + 80*x**3 - 12*x**2 + 24*x + 9)/8`

3.63 $\int \frac{3+12x+20x^2}{9+24x-12x^2+80x^3+320x^4} dx$

Optimal result	509
Mathematica [C] (verified)	509
Rubi [A] (verified)	510
Maple [A] (verified)	511
Fricas [A] (verification not implemented)	511
Sympy [A] (verification not implemented)	512
Maxima [F]	512
Giac [A] (verification not implemented)	512
Mupad [B] (verification not implemented)	513
Reduce [F]	513

Optimal result

Integrand size = 33, antiderivative size = 59

$$\int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = -\frac{\arctan\left(\frac{7-40x}{5\sqrt{11}}\right)}{2\sqrt{11}} + \frac{\arctan\left(\frac{57+30x-40x^2+800x^3}{6\sqrt{11}}\right)}{2\sqrt{11}}$$

output

```
-1/22*arctan(1/55*(7-40*x)*11^(1/2))*11^(1/2)+1/22*arctan(1/66*(800*x^3-40*x^2+30*x+57)*11^(1/2))*11^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.46

$$\int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{1}{8} \text{RootSum} \left[9 + 24\#1 - 12\#1^2 + 80\#1^3 + 320\#1^4 \&, \frac{3 \log(x - \#1) + 12 \log(x - \#1)\#1 + 20 \log(x - \#1)\#1^2}{3 - 3\#1 + 30\#1^2 + 160\#1^3} \& \right]$$

input

```
Integrate[(3 + 12*x + 20*x^2)/(9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4),x]
```

output

```
RootSum[9 + 24*#1 - 12*#1^2 + 80*#1^3 + 320*#1^4 & , (3*Log[x - #1] + 12*Log[x - #1]*#1 + 20*Log[x - #1]*#1^2)/(3 - 3*#1 + 30*#1^2 + 160*#1^3) & ]/8
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.030$, Rules used = {2502}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{20x^2 + 12x + 3}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx$$

↓ 2502

$$\frac{\arctan\left(\frac{800x^3 - 40x^2 + 30x + 57}{6\sqrt{11}}\right)}{2\sqrt{11}} - \frac{\arctan\left(\frac{7 - 40x}{5\sqrt{11}}\right)}{2\sqrt{11}}$$

input

```
Int[(3 + 12*x + 20*x^2)/(9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4), x]
```

output

```
-1/2*ArcTan[(7 - 40*x)/(5*Sqrt[11])]/Sqrt[11] + ArcTan[(57 + 30*x - 40*x^2 + 800*x^3)/(6*Sqrt[11])]/(2*Sqrt[11])
```

Defintions of rubi rules used

rule 2502

```
Int[((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)/((a_) + (b_.)*(x_) + (c_.)*(x_)^2 + (d_.)*(x_)^3 + (e_.)*(x_)^4), x_Symbol] :> With[{q = Rt[(-C)*(2*e*(B*d - 4*A*e) + C*(d^2 - 4*c*e)), 2]}, Simp[2*(C^2/q)*ArcTan[(C*d - B*e + 2*C*e*x)/q], x] - Simp[2*(C^2/q)*ArcTan[C*((4*B*c*C - 3*B^2*d - 4*A*C*d + 12*A*B*e + 4*C*(2*c*C - B*d + 2*A*e))*x + 4*C*(2*C*d - B*e))*x^2 + 8*C^2*e*x^3]/(q*(B^2 - 4*A*C))], x] /; FreeQ[{a, b, c, d, e, A, B, C}, x] && EqQ[B^2*d + 2*C*(b*C + A*d) - 2*B*(c*C + 2*A*e), 0] && EqQ[2*B^2*c*C - 8*a*C^3 - B^3*d - 4*A*B*C*d + 4*A*(B^2 + 2*A*C)*e, 0] && NegQ[C*(2*e*(B*d - 4*A*e) + C*(d^2 - 4*c*e))]
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.88

method	result	size
risch	$\frac{\sqrt{11} \arctan\left(\frac{(40x-7)\sqrt{11}}{55}\right)}{22} + \frac{\sqrt{11} \arctan\left(-\frac{20\sqrt{11}x^2 + 5\sqrt{11}x + 19\sqrt{11} + 400\sqrt{11}x^3}{33}\right)}{22}$	52
default	$\frac{i\sqrt{11} \ln\left(80x^2 + (10i\sqrt{11}+10)x + 3i\sqrt{11}-9\right)}{44} - \frac{i\sqrt{11} \ln\left(80x^2 + (-10i\sqrt{11}+10)x - 3i\sqrt{11}-9\right)}{44}$	62

input

```
int((20*x^2+12*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x,method=_RETURNVERBOSE)
```

output

```
1/22*11^(1/2)*arctan(1/55*(40*x-7)*11^(1/2))+1/22*11^(1/2)*arctan(-20/33*11^(1/2)*x^2+5/11*11^(1/2)*x+19/22*11^(1/2)+400/33*11^(1/2)*x^3)
```

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.73

$$\begin{aligned} & \int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx \\ &= \frac{1}{22} \sqrt{11} \arctan\left(\frac{1}{66} \sqrt{11}(800x^3 - 40x^2 + 30x + 57)\right) \\ &+ \frac{1}{22} \sqrt{11} \arctan\left(\frac{1}{55} \sqrt{11}(40x - 7)\right) \end{aligned}$$

input

```
integrate((20*x^2+12*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x, algorithm="fricas")
```

output

```
1/22*sqrt(11)*arctan(1/66*sqrt(11)*(800*x^3 - 40*x^2 + 30*x + 57)) + 1/22*sqrt(11)*arctan(1/55*sqrt(11)*(40*x - 7))
```


Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.24

$$\int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx$$

$$= \frac{\sqrt{11} \cdot \left(2 \operatorname{atan} \left(\frac{8\sqrt{11}x}{11} - \frac{7\sqrt{11}}{55} \right) + 2 \operatorname{atan} \left(\frac{400\sqrt{11}x^3}{33} - \frac{20\sqrt{11}x^2}{33} + \frac{5\sqrt{11}x}{11} + \frac{19\sqrt{11}}{22} \right) \right)}{44}$$

input `integrate((20*x**2+12*x+3)/(320*x**4+80*x**3-12*x**2+24*x+9), x)`output `sqrt(11)*(2*atan(8*sqrt(11)*x/11 - 7*sqrt(11)/55) + 2*atan(400*sqrt(11)*x**3/33 - 20*sqrt(11)*x**2/33 + 5*sqrt(11)*x/11 + 19*sqrt(11)/22))/44`**Maxima [F]**

$$\int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \int \frac{20x^2 + 12x + 3}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx$$

input `integrate((20*x^2+12*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9), x, algorithm="maxima")`output `integrate((20*x^2 + 12*x + 3)/(320*x^4 + 80*x^3 - 12*x^2 + 24*x + 9), x)`**Giac [A] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.68

$$\int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx$$

$$= \frac{1}{22} \sqrt{11} \left(\arctan \left(\frac{1}{66} \sqrt{11} (800x^3 - 40x^2 + 30x + 57) \right) - \arctan \left(-\frac{1}{55} \sqrt{11} (40x - 7) \right) \right)$$

input `integrate((20*x^2+12*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x, algorithm="giac")`

output `1/22*sqrt(11)*(arctan(1/66*sqrt(11)*(800*x^3 - 40*x^2 + 30*x + 57)) - arctan(-1/55*sqrt(11)*(40*x - 7)))`

Mupad [B] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.90

$$\int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx$$

$$= \frac{\sqrt{11} \operatorname{atan}\left(\frac{8\sqrt{11}x}{11} - \frac{7\sqrt{11}}{55}\right)}{22} + \frac{\sqrt{11} \operatorname{atan}\left(\frac{400\sqrt{11}x^3}{33} - \frac{20\sqrt{11}x^2}{33} + \frac{5\sqrt{11}x}{11} + \frac{19\sqrt{11}}{22}\right)}{22}$$

input `int((12*x + 20*x^2 + 3)/(24*x - 12*x^2 + 80*x^3 + 320*x^4 + 9),x)`

output `(11^(1/2)*atan((8*11^(1/2)*x)/11 - (7*11^(1/2))/55))/22 + (11^(1/2)*atan((5*11^(1/2)*x)/11 + (19*11^(1/2))/22 - (20*11^(1/2)*x^2)/33 + (400*11^(1/2)*x^3)/33))/22`

Reduce [F]

$$\int \frac{3 + 12x + 20x^2}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = 20 \left(\int \frac{x^2}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx \right)$$

$$+ 12 \left(\int \frac{x}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx \right)$$

$$+ 3 \left(\int \frac{1}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx \right)$$

input `int((20*x^2+12*x+3)/(320*x^4+80*x^3-12*x^2+24*x+9),x)`

output

```
20*int(x**2/(320*x**4 + 80*x**3 - 12*x**2 + 24*x + 9),x) + 12*int(x/(320*x
**4 + 80*x**3 - 12*x**2 + 24*x + 9),x) + 3*int(1/(320*x**4 + 80*x**3 - 12*
x**2 + 24*x + 9),x)
```

3.64 $\int \frac{-84-576x-400x^2+2560x^3}{9+24x-12x^2+80x^3+320x^4} dx$

Optimal result	515
Mathematica [C] (verified)	515
Rubi [A] (verified)	516
Maple [C] (verified)	517
Fricas [A] (verification not implemented)	518
Sympy [A] (verification not implemented)	518
Maxima [F]	519
Giac [A] (verification not implemented)	519
Mupad [B] (verification not implemented)	520
Reduce [F]	520

Optimal result

Integrand size = 38, antiderivative size = 78

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = 2\sqrt{11} \arctan\left(\frac{7 - 40x}{5\sqrt{11}}\right) - 2\sqrt{11} \arctan\left(\frac{57 + 30x - 40x^2 + 800x^3}{6\sqrt{11}}\right) + 2 \log(9 + 24x - 12x^2 + 80x^3 + 320x^4)$$

output

```
2*ln(320*x^4+80*x^3-12*x^2+24*x+9)+2*arctan(1/55*(7-40*x)*11^(1/2))*11^(1/2)-2*arctan(1/66*(800*x^3-40*x^2+30*x+57)*11^(1/2))*11^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.01 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.27

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \frac{1}{2} \text{RootSum}\left[9 + 24\#1 - 12\#1^2 + 80\#1^3 + 320\#1^4 \&, \frac{-21 \log(x - \#1) - 144 \log(x - \#1)\#1 - 100 \log(x - \#1)\#1^2 + 640 \log(x - \#1)\#1^3}{3 - 3\#1 + 30\#1^2 + 160\#1^3} \&\right]$$

input `Integrate[(-84 - 576*x - 400*x^2 + 2560*x^3)/(9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4), x]`

output `RootSum[9 + 24*#1 - 12*#1^2 + 80*#1^3 + 320*#1^4 & , (-21*Log[x - #1] - 14 4*Log[x - #1]*#1 - 100*Log[x - #1]*#1^2 + 640*Log[x - #1]*#1^3)/(3 - 3*#1 + 30*#1^2 + 160*#1^3) &]/2`

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.09, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.079$, Rules used = {2525, 27, 2502}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{2560x^3 - 400x^2 - 576x - 84}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx$$

$$\downarrow \text{2525}$$

$$\int -\frac{56320(20x^2+12x+3)}{320x^4+80x^3-12x^2+24x+9} dx + 2 \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

$$\downarrow \text{27}$$

$$2 \log(320x^4 + 80x^3 - 12x^2 + 24x + 9) - 44 \int \frac{20x^2 + 12x + 3}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx$$

$$\downarrow \text{2502}$$

$$2 \log(320x^4 + 80x^3 - 12x^2 + 24x + 9) - 44 \left(\frac{\arctan\left(\frac{800x^3 - 40x^2 + 30x + 57}{6\sqrt{11}}\right)}{2\sqrt{11}} - \frac{\arctan\left(\frac{7 - 40x}{5\sqrt{11}}\right)}{2\sqrt{11}} \right)$$

input `Int[(-84 - 576*x - 400*x^2 + 2560*x^3)/(9 + 24*x - 12*x^2 + 80*x^3 + 320*x^4), x]`

output

```
-44*(-1/2*ArcTan[(7 - 40*x)/(5*sqrt[11])]/sqrt[11] + ArcTan[(57 + 30*x - 4
0*x^2 + 800*x^3)/(6*sqrt[11])]/(2*sqrt[11])) + 2*Log[9 + 24*x - 12*x^2 + 8
0*x^3 + 320*x^4]
```

Defintions of rubi rules used

rule 27

```
Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !Ma
tchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]
```

rule 2502

```
Int[((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)/((a_) + (b_.)*(x_) + (c_.)*(x_)^2 +
(d_.)*(x_)^3 + (e_.)*(x_)^4), x_Symbol] := With[{q = Rt[(-C)*(2*e*(B*d - 4
*A*e) + C*(d^2 - 4*c*e)), 2]}, Simp[2*(C^2/q)*ArcTan[(C*d - B*e + 2*C*e*x)/
q], x] - Simp[2*(C^2/q)*ArcTan[C*((4*B*c*C - 3*B^2*d - 4*A*C*d + 12*A*B*e +
4*C*(2*c*C - B*d + 2*A*e)*x + 4*C*(2*C*d - B*e)*x^2 + 8*C^2*e*x^3)/(q*(B^2
- 4*A*C))], x]] /; FreeQ[{a, b, c, d, e, A, B, C}, x] && EqQ[B^2*d + 2*C*
(b*C + A*d) - 2*B*(c*C + 2*A*e), 0] && EqQ[2*B^2*c*C - 8*a*C^3 - B^3*d - 4*
A*B*C*d + 4*A*(B^2 + 2*A*C)*e, 0] && NegQ[C*(2*e*(B*d - 4*A*e) + C*(d^2 - 4
*c*e))]
```

rule 2525

```
Int[(Pm_)/(Qn_), x_Symbol] := With[{m = Expon[Pm, x], n = Expon[Qn, x]}, Si
mp[Coeff[Pm, x, m]*(Log[Qn]/(n*Coeff[Qn, x, n])), x] + Simp[1/(n*Coeff[Qn,
x, n]) Int[ExpandToSum[n*Coeff[Qn, x, n]*Pm - Coeff[Pm, x, m]*D[Qn, x], x
]/Qn, x], x] /; EqQ[m, n - 1] /; PolyQ[Pm, x] && PolyQ[Qn, x]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.90

method	result
default	$4\left(\frac{i\sqrt{11}}{4} + \frac{1}{2}\right) \ln(80x^2 + (-10i\sqrt{11} + 10)x - 3i\sqrt{11} - 9) + 4\left(\frac{1}{2} - \frac{i\sqrt{11}}{4}\right) \ln(80x^2 + (10i\sqrt{11} + 10)x - 3i\sqrt{11} - 9)$
risch	$2 \ln(6400x^4 + 1600x^3 - 240x^2 + 480x + 180) - 2\sqrt{11} \arctan\left(-\frac{20\sqrt{11}x^2}{33} + \frac{5\sqrt{11}x}{11} + \frac{19\sqrt{11}}{22} + \frac{40}{11}\right)$

input `int((2560*x^3-400*x^2-576*x-84)/(320*x^4+80*x^3-12*x^2+24*x+9),x,method=_RETURVERBOSE)`

output `4*(1/4*I*11^(1/2)+1/2)*ln(80*x^2+(-10*I*11^(1/2)+10)*x-3*I*11^(1/2)-9)+4*(1/2-1/4*I*11^(1/2))*ln(80*x^2+(10*I*11^(1/2)+10)*x+3*I*11^(1/2)-9)`

Fricas [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.85

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx$$

$$= -2\sqrt{11} \arctan\left(\frac{1}{66}\sqrt{11}(800x^3 - 40x^2 + 30x + 57)\right)$$

$$- 2\sqrt{11} \arctan\left(\frac{1}{55}\sqrt{11}(40x - 7)\right) + 2 \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

input `integrate((2560*x^3-400*x^2-576*x-84)/(320*x^4+80*x^3-12*x^2+24*x+9),x, algorithm="fricas")`

output `-2*sqrt(11)*arctan(1/66*sqrt(11)*(800*x^3 - 40*x^2 + 30*x + 57)) - 2*sqrt(11)*arctan(1/55*sqrt(11)*(40*x - 7)) + 2*log(320*x^4 + 80*x^3 - 12*x^2 + 24*x + 9)`

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.28

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx$$

$$= \sqrt{11} \left(-2 \operatorname{atan}\left(\frac{8\sqrt{11}x}{11} - \frac{7\sqrt{11}}{55}\right) \right.$$

$$\left. - 2 \operatorname{atan}\left(\frac{400\sqrt{11}x^3}{33} - \frac{20\sqrt{11}x^2}{33} + \frac{5\sqrt{11}x}{11} + \frac{19\sqrt{11}}{22}\right) \right)$$

$$+ 2 \log\left(x^4 + \frac{x^3}{4} - \frac{3x^2}{80} + \frac{3x}{40} + \frac{9}{320}\right)$$

input `integrate((2560*x**3-400*x**2-576*x-84)/(320*x**4+80*x**3-12*x**2+24*x+9), x)`

output `sqrt(11)*(-2*atan(8*sqrt(11)*x/11 - 7*sqrt(11)/55) - 2*atan(400*sqrt(11)*x**3/33 - 20*sqrt(11)*x**2/33 + 5*sqrt(11)*x/11 + 19*sqrt(11)/22)) + 2*log(x**4 + x**3/4 - 3*x**2/80 + 3*x/40 + 9/320)`

Maxima [F]

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = \int \frac{4(640x^3 - 100x^2 - 144x - 21)}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx$$

input `integrate((2560*x^3-400*x^2-576*x-84)/(320*x^4+80*x^3-12*x^2+24*x+9), x, algorithm="maxima")`

output `4*integrate((640*x^3 - 100*x^2 - 144*x - 21)/(320*x^4 + 80*x^3 - 12*x^2 + 24*x + 9), x)`

Giac [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.82

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx =$$

$$-2\sqrt{11} \left(\arctan \left(\frac{1}{66} \sqrt{11} (800x^3 - 40x^2 + 30x + 57) \right) - \arctan \left(-\frac{1}{55} \sqrt{11} (40x - 7) \right) \right)$$

$$+ 2 \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

input `integrate((2560*x^3-400*x^2-576*x-84)/(320*x^4+80*x^3-12*x^2+24*x+9), x, algorithm="giac")`

output `-2*sqrt(11)*(arctan(1/66*sqrt(11)*(800*x^3 - 40*x^2 + 30*x + 57)) - arctan(-1/55*sqrt(11)*(40*x - 7))) + 2*log(320*x^4 + 80*x^3 - 12*x^2 + 24*x + 9)`

Mupad [B] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.97

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = 2 \ln(320x^4 + 80x^3 - 12x^2 + 24x + 9) - 2\sqrt{11} \operatorname{atan}\left(\frac{8\sqrt{11}x}{11} - \frac{7\sqrt{11}}{55}\right) - 2\sqrt{11} \operatorname{atan}\left(\frac{400\sqrt{11}x^3}{33} - \frac{20\sqrt{11}x^2}{33} + \frac{5\sqrt{11}x}{11} + \frac{19\sqrt{11}}{22}\right)$$

input

```
int(-(576*x + 400*x^2 - 2560*x^3 + 84)/(24*x - 12*x^2 + 80*x^3 + 320*x^4 + 9), x)
```

output

```
2*log(24*x - 12*x^2 + 80*x^3 + 320*x^4 + 9) - 2*11^(1/2)*atan((8*11^(1/2)*x)/11 - (7*11^(1/2))/55) - 2*11^(1/2)*atan((5*11^(1/2)*x)/11 + (19*11^(1/2))/22 - (20*11^(1/2)*x^2)/33 + (400*11^(1/2)*x^3)/33)
```

Reduce [F]

$$\int \frac{-84 - 576x - 400x^2 + 2560x^3}{9 + 24x - 12x^2 + 80x^3 + 320x^4} dx = -880 \left(\int \frac{x^2}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx \right) - 528 \left(\int \frac{x}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx \right) - 132 \left(\int \frac{1}{320x^4 + 80x^3 - 12x^2 + 24x + 9} dx \right) + 2 \log(320x^4 + 80x^3 - 12x^2 + 24x + 9)$$

input

```
int((2560*x^3-400*x^2-576*x-84)/(320*x^4+80*x^3-12*x^2+24*x+9), x)
```

output

```
2*( - 440*int(x**2/(320*x**4 + 80*x**3 - 12*x**2 + 24*x + 9),x) - 264*int(x/(320*x**4 + 80*x**3 - 12*x**2 + 24*x + 9),x) - 66*int(1/(320*x**4 + 80*x**3 - 12*x**2 + 24*x + 9),x) + log(320*x**4 + 80*x**3 - 12*x**2 + 24*x + 9))
```

3.65 $\int \frac{\sqrt{1-x^4}}{1+x^4} dx$

Optimal result	522
Mathematica [C] (verified)	522
Rubi [A] (verified)	523
Maple [C] (verified)	524
Fricas [A] (verification not implemented)	524
Sympy [F]	525
Maxima [F]	525
Giac [F]	525
Mupad [F(-1)]	526
Reduce [F]	526

Optimal result

Integrand size = 19, antiderivative size = 49

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = \frac{1}{2} \arctan\left(\frac{x(1+x^2)}{\sqrt{1-x^4}}\right) + \frac{1}{2} \operatorname{arctanh}\left(\frac{x(1-x^2)}{\sqrt{1-x^4}}\right)$$

output `1/2*arctan(x*(x^2+1)/(-x^4+1)^(1/2))+1/2*arctanh(x*(-x^2+1)/(-x^4+1)^(1/2))`

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.14 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.16

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = \left(\frac{1}{4} - \frac{i}{4}\right) \arctan\left(\frac{(1+i)x}{\sqrt{1-x^4}}\right) - \left(\frac{1}{4} + \frac{i}{4}\right) \arctan\left(\frac{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{1-x^4}}{x}\right)$$

input `Integrate[Sqrt[1 - x^4]/(1 + x^4),x]`

output `(1/4 - I/4)*ArcTan[((1 + I)*x)/Sqrt[1 - x^4]] - (1/4 + I/4)*ArcTan[((1/2 + I/2)*Sqrt[1 - x^4])/x]`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {921}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{1-x^4}}{x^4+1} dx$$

↓ 921

$$\frac{1}{2} \arctan\left(\frac{x(x^2+1)}{\sqrt{1-x^4}}\right) + \frac{1}{2} \operatorname{arctanh}\left(\frac{x(1-x^2)}{\sqrt{1-x^4}}\right)$$

input `Int[Sqrt[1 - x^4]/(1 + x^4),x]`

output `ArcTan[(x*(1 + x^2))/Sqrt[1 - x^4]]/2 + ArcTanh[(x*(1 - x^2))/Sqrt[1 - x^4]]/2`

Defintions of rubi rules used

rule 921 `Int[Sqrt[(a_) + (b_.)*(x_)^4]/((c_) + (d_.)*(x_)^4), x_Symbol] :> With[{q = Rt[(-a)*b, 4]}, Simp[(a/(2*c*q))*ArcTan[q*x*((a + q^2*x^2)/(a*Sqrt[a + b*x^4]))], x] + Simp[(a/(2*c*q))*ArcTanh[q*x*((a - q^2*x^2)/(a*Sqrt[a + b*x^4]))], x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c + a*d, 0] && NegQ[a*b]`

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.79 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.14

method	result
pseudoelliptic	$\left(\frac{1}{4} - \frac{i}{4}\right) \left(\ln \left(\frac{(-1+i)\sqrt{-x^4+1}-2x}{ix^2-1} \right) - \arctan \left(\frac{(\frac{1}{2}-\frac{i}{2})\sqrt{-x^4+1}}{x} \right) + \ln(2) \right)$
default	$-\frac{\ln \left(\frac{1+\frac{-x^4+1}{2x^2} - \frac{\sqrt{-x^4+1}}{x}}{1+\frac{-x^4+1}{2x^2} + \frac{\sqrt{-x^4+1}}{x}} \right)}{8} - \frac{\arctan \left(1 + \frac{\sqrt{-x^4+1}}{x} \right)}{4} - \frac{\arctan \left(-1 + \frac{\sqrt{-x^4+1}}{x} \right)}{4}$
elliptic	$-\frac{\ln \left(\frac{1+\frac{-x^4+1}{2x^2} - \frac{\sqrt{-x^4+1}}{x}}{1+\frac{-x^4+1}{2x^2} + \frac{\sqrt{-x^4+1}}{x}} \right)}{8} - \frac{\arctan \left(1 + \frac{\sqrt{-x^4+1}}{x} \right)}{4} - \frac{\arctan \left(-1 + \frac{\sqrt{-x^4+1}}{x} \right)}{4}$
trager	$\text{RootOf}(8_Z^2 - 4_Z + 1) \ln \left(\frac{4 \text{RootOf}(8_Z^2 - 4_Z + 1)x - \sqrt{-x^4+1} - 2x}{4x^2 \text{RootOf}(8_Z^2 - 4_Z + 1) - x^2 + 1} \right) + \frac{\ln \left(-\frac{4 \text{RootOf}(8_Z^2 - 4_Z + 1)}{4x^2 \text{RootOf}(8_Z^2 - 4_Z + 1)} \right)}{2}$

input `int((-x^4+1)^(1/2)/(x^4+1),x,method=_RETURNVERBOSE)`

output $(1/4-1/4*I)*(\ln(((1+I)*(-x^4+1)^(1/2)-2*x)/(I*x^2-1))-\arctan((1/2-1/2*I)*(-x^4+1)^(1/2)/x)+\ln(2))$

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.14

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = -\frac{1}{2} \arctan \left(\frac{\sqrt{-x^4+1}x}{x^2-1} \right) + \frac{1}{4} \log \left(-\frac{x^4-2x^2-2\sqrt{-x^4+1}x-1}{x^4+1} \right)$$

input `integrate((-x^4+1)^(1/2)/(x^4+1),x, algorithm="fricas")`

output $-1/2*\arctan(\text{sqrt}(-x^4+1)*x/(x^2-1))+1/4*\log(-x^4-2*x^2-2*\text{sqrt}(-x^4+1)*x-1)/(x^4+1))$

Sympy [F]

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = \int \frac{\sqrt{-(x-1)(x+1)(x^2+1)}}{x^4+1} dx$$

input `integrate((-x**4+1)**(1/2)/(x**4+1),x)`

output `Integral(sqrt(-(x - 1)*(x + 1)*(x**2 + 1))/(x**4 + 1), x)`

Maxima [F]

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = \int \frac{\sqrt{-x^4+1}}{x^4+1} dx$$

input `integrate((-x^4+1)^(1/2)/(x^4+1),x, algorithm="maxima")`

output `integrate(sqrt(-x^4 + 1)/(x^4 + 1), x)`

Giac [F]

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = \int \frac{\sqrt{-x^4+1}}{x^4+1} dx$$

input `integrate((-x^4+1)^(1/2)/(x^4+1),x, algorithm="giac")`

output `integrate(sqrt(-x^4 + 1)/(x^4 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = \int \frac{\sqrt{1-x^4}}{x^4+1} dx$$

input `int((1 - x^4)^(1/2)/(x^4 + 1),x)`output `int((1 - x^4)^(1/2)/(x^4 + 1), x)`**Reduce [F]**

$$\int \frac{\sqrt{1-x^4}}{1+x^4} dx = \int \frac{\sqrt{-x^4+1}}{x^4+1} dx$$

input `int((-x^4+1)^(1/2)/(x^4+1),x)`output `int(sqrt(-x**4 + 1)/(x**4 + 1),x)`

3.66 $\int \frac{\sqrt{1+x^4}}{1-x^4} dx$

Optimal result	527
Mathematica [A] (verified)	527
Rubi [A] (verified)	528
Maple [A] (verified)	529
Fricas [A] (verification not implemented)	530
Sympy [F]	530
Maxima [F]	531
Giac [F]	531
Mupad [F(-1)]	531
Reduce [F]	532

Optimal result

Integrand size = 19, antiderivative size = 53

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{2\sqrt{2}}$$

output `1/4*arctan(x*2^(1/2)/(x^4+1)^(1/2))*2^(1/2)+1/4*arctanh(x*2^(1/2)/(x^4+1)^(1/2))*2^(1/2)`

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right) + \operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{2\sqrt{2}}$$

input `Integrate[Sqrt[1 + x^4]/(1 - x^4), x]`

output `(ArcTan[(Sqrt[2]*x)/Sqrt[1 + x^4]] + ArcTanh[(Sqrt[2]*x)/Sqrt[1 + x^4]])/(2*Sqrt[2])`

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {920, 756, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{x^4+1}}{1-x^4} dx \\
 & \quad \downarrow \text{920} \\
 & \int \frac{1}{1-\frac{4x^4}{(x^4+1)^2}} d\frac{x}{\sqrt{x^4+1}} \\
 & \quad \downarrow \text{756} \\
 & \frac{1}{2} \int \frac{1}{1-\frac{2x^2}{x^4+1}} d\frac{x}{\sqrt{x^4+1}} + \frac{1}{2} \int \frac{1}{\frac{2x^2}{x^4+1}+1} d\frac{x}{\sqrt{x^4+1}} \\
 & \quad \downarrow \text{216} \\
 & \frac{1}{2} \int \frac{1}{1-\frac{2x^2}{x^4+1}} d\frac{x}{\sqrt{x^4+1}} + \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{x^4+1}}\right)}{2\sqrt{2}} \\
 & \quad \downarrow \text{219} \\
 & \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{x^4+1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{x^4+1}}\right)}{2\sqrt{2}}
 \end{aligned}$$

input `Int[Sqrt[1 + x^4]/(1 - x^4),x]`

output `ArcTan[(Sqrt[2]*x)/Sqrt[1 + x^4]]/(2*Sqrt[2]) + ArcTanh[(Sqrt[2]*x)/Sqrt[1 + x^4]]/(2*Sqrt[2])`

Defintions of rubi rules used

rule 216 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

rule 219 $\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

rule 756 $\text{Int}[(a_ + (b_ \cdot)(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Simp}[r/(2 \cdot a) \text{Int}[1/(r - s \cdot x^2), x], x] + \text{Simp}[r/(2 \cdot a) \text{Int}[1/(r + s \cdot x^2), x], x]] /;$ FreeQ[{a, b}, x] && !GtQ[a/b, 0]

rule 920 $\text{Int}[\text{Sqrt}[(a_ + (b_ \cdot)(x_)^4)/((c_ + (d_ \cdot)(x_)^4)], x_Symbol] \rightarrow \text{Simp}[a/c \text{Subst}[\text{Int}[1/(1 - 4 \cdot a \cdot b \cdot x^4), x], x, x/\text{Sqrt}[a + b \cdot x^4]], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[b \cdot c + a \cdot d, 0] && PosQ[a \cdot b]

Maple [A] (verified)

Time = 1.55 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.17

method	result
default	$\frac{\left(2 \arctan\left(\frac{x\sqrt{2}}{\sqrt{x^4+1}}\right) + \operatorname{arctanh}\left(\frac{(x^2-x+1)\sqrt{2}}{\sqrt{x^4+1}}\right) - \operatorname{arctanh}\left(\frac{(x^2+x+1)\sqrt{2}}{\sqrt{x^4+1}}\right)\right)\sqrt{2}}{8}$
pseudoelliptic	$\frac{\left(2 \arctan\left(\frac{x\sqrt{2}}{\sqrt{x^4+1}}\right) + \operatorname{arctanh}\left(\frac{(x^2-x+1)\sqrt{2}}{\sqrt{x^4+1}}\right) - \operatorname{arctanh}\left(\frac{(x^2+x+1)\sqrt{2}}{\sqrt{x^4+1}}\right)\right)\sqrt{2}}{8}$
elliptic	$\frac{\left(-\frac{\ln\left(-1 + \frac{\sqrt{2}\sqrt{x^4+1}}{2x}\right)}{4} + \frac{\ln\left(1 + \frac{\sqrt{2}\sqrt{x^4+1}}{2x}\right)}{4} - \frac{\arctan\left(\frac{\sqrt{2}\sqrt{x^4+1}}{2x}\right)}{2}\right)\sqrt{2}}{2}$
trager	$-\frac{\operatorname{RootOf}\left(-Z^2+2\right) \ln\left(\frac{\operatorname{RootOf}\left(-Z^2+2\right)x+\sqrt{x^4+1}}{x^2+1}\right)}{4} - \frac{\operatorname{RootOf}\left(-Z^2-2\right) \ln\left(\frac{-\operatorname{RootOf}\left(-Z^2-2\right)x+\sqrt{x^4+1}}{(-1+x)(1+x)}\right)}{4}$

input `int((x^4+1)^(1/2)/(-x^4+1),x,method=_RETURNVERBOSE)`

output `1/8*(2*arctan(x*2^(1/2)/(x^4+1)^(1/2))+arctanh((x^2-x+1)*2^(1/2)/(x^4+1)^(1/2))-arctanh((x^2+x+1)*2^(1/2)/(x^4+1)^(1/2)))*2^(1/2)`

Fricas [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.15

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = \frac{1}{4} \sqrt{2} \arctan\left(\frac{\sqrt{2}x}{\sqrt{x^4+1}}\right) + \frac{1}{8} \sqrt{2} \log\left(\frac{x^4 + 2\sqrt{2}\sqrt{x^4+1}x + 2x^2 + 1}{x^4 - 2x^2 + 1}\right)$$

input `integrate((x^4+1)^(1/2)/(-x^4+1),x, algorithm="fricas")`

output `1/4*sqrt(2)*arctan(sqrt(2)*x/sqrt(x^4 + 1)) + 1/8*sqrt(2)*log((x^4 + 2*sqrt(2)*sqrt(x^4 + 1)*x + 2*x^2 + 1)/(x^4 - 2*x^2 + 1))`

Sympy [F]

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = - \int \frac{\sqrt{x^4+1}}{x^4-1} dx$$

input `integrate((x**4+1)**(1/2)/(-x**4+1),x)`

output `-Integral(sqrt(x**4 + 1)/(x**4 - 1), x)`

Maxima [F]

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = \int -\frac{\sqrt{x^4+1}}{x^4-1} dx$$

input `integrate((x^4+1)^(1/2)/(-x^4+1),x, algorithm="maxima")`

output `-integrate(sqrt(x^4 + 1)/(x^4 - 1), x)`

Giac [F]

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = \int -\frac{\sqrt{x^4+1}}{x^4-1} dx$$

input `integrate((x^4+1)^(1/2)/(-x^4+1),x, algorithm="giac")`

output `integrate(-sqrt(x^4 + 1)/(x^4 - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = -\int \frac{\sqrt{x^4+1}}{x^4-1} dx$$

input `int(-(x^4 + 1)^(1/2)/(x^4 - 1),x)`

output `-int((x^4 + 1)^(1/2)/(x^4 - 1), x)`

Reduce [F]

$$\int \frac{\sqrt{1+x^4}}{1-x^4} dx = - \left(\int \frac{\sqrt{x^4+1}}{x^4-1} dx \right)$$

input `int((x^4+1)^(1/2)/(-x^4+1),x)`

output `- int(sqrt(x**4 + 1)/(x**4 - 1),x)`

3.67 $\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx$

Optimal result	533
Mathematica [A] (verified)	533
Rubi [A] (verified)	534
Maple [A] (verified)	535
Fricas [A] (verification not implemented)	536
Sympy [F]	537
Maxima [F]	537
Giac [F]	538
Mupad [F(-1)]	538
Reduce [F]	538

Optimal result

Integrand size = 24, antiderivative size = 75

$$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx = \frac{1}{4} \sqrt{2-p} \arctan\left(\frac{\sqrt{2-px}}{\sqrt{1+px^2+x^4}}\right) + \frac{1}{4} \sqrt{2+p} \operatorname{arctanh}\left(\frac{\sqrt{2+px}}{\sqrt{1+px^2+x^4}}\right)$$

output

$1/4*\arctan(x*(2-p)^{(1/2)/(x^4+p*x^2+1)^{(1/2))}*(2-p)^{(1/2)}+1/4*\operatorname{arctanh}(x*(2+p)^{(1/2)/(x^4+p*x^2+1)^{(1/2))}*(2+p)^{(1/2)}$

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.05

$$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx = -\frac{1}{4} \sqrt{-2-p} \arctan\left(\frac{\sqrt{-2-px}}{\sqrt{1+px^2+x^4}}\right) + \frac{1}{4} \sqrt{2-p} \arctan\left(\frac{\sqrt{2-px}}{\sqrt{1+px^2+x^4}}\right)$$

input

`Integrate[Sqrt[1 + p*x^2 + x^4]/(1 - x^4), x]`

output

```
-1/4*(Sqrt[-2 - p]*ArcTan[(Sqrt[-2 - p]*x)/Sqrt[1 + p*x^2 + x^4]]) + (Sqrt
[2 - p]*ArcTan[(Sqrt[2 - p]*x)/Sqrt[1 + p*x^2 + x^4]])/4
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.35, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2517, 1406, 218, 221}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{\sqrt{px^2 + x^4 + 1}}{1 - x^4} dx \\
 & \quad \downarrow \text{2517} \\
 & \int \frac{1}{-\frac{(4-p^2)x^4}{(px^2+x^4+1)^2} - \frac{2px^2}{px^2+x^4+1} + 1} d \frac{x}{\sqrt{px^2 + x^4 + 1}} \\
 & \quad \downarrow \text{1406} \\
 & \frac{1}{4}(4-p^2) \int \frac{1}{-\frac{(4-p^2)x^2}{x^4+px^2+1} - p + 2} d \frac{x}{\sqrt{x^4 + px^2 + 1}} - \\
 & \frac{1}{4}(4-p^2) \int \frac{1}{-\frac{(4-p^2)x^2}{x^4+px^2+1} - p - 2} d \frac{x}{\sqrt{x^4 + px^2 + 1}} \\
 & \quad \downarrow \text{218} \\
 & \frac{1}{4}(4-p^2) \int \frac{1}{-\frac{(4-p^2)x^2}{x^4+px^2+1} - p + 2} d \frac{x}{\sqrt{x^4 + px^2 + 1}} + \frac{(4-p^2) \arctan\left(\frac{\sqrt{2-px}}{\sqrt{px^2+x^4+1}}\right)}{4\sqrt{2-p}(p+2)} \\
 & \quad \downarrow \text{221} \\
 & \frac{(4-p^2) \arctan\left(\frac{\sqrt{2-px}}{\sqrt{px^2+x^4+1}}\right)}{4\sqrt{2-p}(p+2)} + \frac{(4-p^2) \operatorname{arctanh}\left(\frac{\sqrt{p+2x}}{\sqrt{px^2+x^4+1}}\right)}{4(2-p)\sqrt{p+2}}
 \end{aligned}$$

input

```
Int[Sqrt[1 + p*x^2 + x^4]/(1 - x^4), x]
```

output
$$\left((4 - p^2) \operatorname{ArcTan}\left[\frac{\sqrt{2 - p}x}{\sqrt{1 + px^2 + x^4}}\right] / (4\sqrt{2 - p}(2 + p)) + (4 - p^2) \operatorname{ArcTanh}\left[\frac{\sqrt{2 + p}x}{\sqrt{1 + px^2 + x^4}}\right] / (4(2 - p)\sqrt{2 + p}) \right)$$

Defintions of rubi rules used

rule 218
$$\operatorname{Int}[(a_ + (b_.)x^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2]/a) \operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \operatorname{PosQ}[a/b]$$

rule 221
$$\operatorname{Int}[(a_ + (b_.)x^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a) \operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b]$$

rule 1406
$$\operatorname{Int}[(a_ + (b_.)x^2 + (c_.)x^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{q = \operatorname{Rt}[b^2 - 4ac, 2]\}, \operatorname{Simp}[c/q \operatorname{Int}[1/(b/2 - q/2 + cx^2), x], x] - \operatorname{Simp}[c/q \operatorname{Int}[1/(b/2 + q/2 + cx^2), x], x]] \text{ ; FreeQ}\{a, b, c, x\} \ \&\& \ \operatorname{NeQ}[b^2 - 4ac, 0] \ \&\& \ \operatorname{PosQ}[b^2 - 4ac]$$

rule 2517
$$\operatorname{Int}[\sqrt{v_}/((d_ + (e_.)x^4), x_Symbol] \rightarrow \operatorname{With}\{a = \operatorname{Coeff}[v, x, 0], b = \operatorname{Coeff}[v, x, 2], c = \operatorname{Coeff}[v, x, 4]\}, \operatorname{Simp}[a/d \operatorname{Subst}[\operatorname{Int}[1/(1 - 2bx^2 + (b^2 - 4ac)x^4), x], x, x/\sqrt{v}], x] \text{ ; EqQ}[c*d + a*e, 0] \ \&\& \ \operatorname{PosQ}[a*c]] \text{ ; FreeQ}\{d, e, x\} \ \&\& \ \operatorname{PolyQ}[v, x^2, 2]$$

Maple [A] (verified)

Time = 2.43 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.19

method	result
elliptic	$\frac{\left(\frac{4\left(\frac{1}{4} + \frac{p}{8}\right) \operatorname{arctanh}\left(\frac{\sqrt{x^4+px^2+1}\sqrt{2}}{x\sqrt{4+2p}}\right)}{\sqrt{4+2p}} + \frac{4\left(\frac{1}{4} - \frac{p}{8}\right) \operatorname{arctanh}\left(\frac{\sqrt{x^4+px^2+1}\sqrt{2}}{x\sqrt{2p-4}}\right)}{\sqrt{2p-4}} \right) \sqrt{2}}{2}$
pseudoelliptic	$\frac{\left(\ln\left(\frac{\sqrt{2+p}\sqrt{x^4+px^2+1}-2-2x^2+x(p-2)}{(1+x)^2}\right) + \ln\left(\frac{\sqrt{2+p}\sqrt{x^4+px^2+1}+2+2x^2+x(p-2)}{(-1+x)^2}\right) + 2\ln(2) \right) \sqrt{2+p}}{8} - \frac{\sqrt{p-2} \left(\ln\left(\frac{\sqrt{p-2}}{\sqrt{p-2}}\right) \right)}{8}$
default	$- \frac{\left(-\ln\left(\frac{\sqrt{2+p}\sqrt{x^4+px^2+1}-2-2x^2+x(p-2)}{(1+x)^2}\right) - \ln\left(\frac{\sqrt{2+p}\sqrt{x^4+px^2+1}+2+2x^2+x(p-2)}{(-1+x)^2}\right) - 2\ln(2) \right) \sqrt{2+p}}{8} - \frac{\sqrt{p-2} \left(\ln\left(\frac{\sqrt{p-2}}{\sqrt{p-2}}\right) \right)}{8}$

```
input int((x^4+p*x^2+1)^(1/2)/(-x^4+1),x,method=_RETURNVERBOSE)
```

```
output 1/2*(4*(1/4+1/8*p)/(4+2*p)^(1/2)*arctanh((x^4+p*x^2+1)^(1/2)*2^(1/2)/x/(4+2*p)^(1/2))+4*(1/4-1/8*p)/(2*p-4)^(1/2)*arctanh((x^4+p*x^2+1)^(1/2)*2^(1/2)/x/(2*p-4)^(1/2)))*2^(1/2)
```

Fricas [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 345, normalized size of antiderivative = 4.60

$$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx$$

$$= \left[\frac{1}{8} \sqrt{p-2} \log \left(\frac{x^4 + 2(p-1)x^2 - 2\sqrt{x^4+px^2+1}\sqrt{p-2}x + 1}{x^4 + 2x^2 + 1} \right) + \frac{1}{8} \sqrt{p+2} \log \left(\frac{x^4 + 2(p+1)x^2 + 2\sqrt{x^4+px^2+1}\sqrt{p+2}x + 1}{x^4 - 2x^2 + 1} \right), \frac{1}{4} \sqrt{-p+2} \arctan \left(\frac{\sqrt{-p+2}x}{\sqrt{x^4+px^2+1}} \right) + \frac{1}{8} \sqrt{p+2} \log \left(\frac{x^4 + 2(p+1)x^2 + 2\sqrt{x^4+px^2+1}\sqrt{p+2}x + 1}{x^4 - 2x^2 + 1} \right), -\frac{1}{4} \sqrt{-p-2} \arctan \left(\frac{\sqrt{-p-2}x}{\sqrt{x^4+px^2+1}} \right) + \frac{1}{8} \sqrt{p-2} \log \left(\frac{x^4 + 2(p-1)x^2 - 2\sqrt{x^4+px^2+1}\sqrt{p-2}x + 1}{x^4 + 2x^2 + 1} \right), \frac{1}{4} \sqrt{-p+2} \arctan \left(\frac{\sqrt{-p+2}x}{\sqrt{x^4+px^2+1}} \right) - \frac{1}{4} \sqrt{-p-2} \arctan \left(\frac{\sqrt{-p-2}x}{\sqrt{x^4+px^2+1}} \right) \right]$$

input `integrate((x^4+p*x^2+1)^(1/2)/(-x^4+1),x, algorithm="fricas")`

output `[1/8*sqrt(p - 2)*log((x^4 + 2*(p - 1)*x^2 - 2*sqrt(x^4 + p*x^2 + 1)*sqrt(p - 2)*x + 1)/(x^4 + 2*x^2 + 1)) + 1/8*sqrt(p + 2)*log((x^4 + 2*(p + 1)*x^2 + 2*sqrt(x^4 + p*x^2 + 1)*sqrt(p + 2)*x + 1)/(x^4 - 2*x^2 + 1)), 1/4*sqrt(-p + 2)*arctan(sqrt(-p + 2)*x/sqrt(x^4 + p*x^2 + 1)) + 1/8*sqrt(p + 2)*log((x^4 + 2*(p + 1)*x^2 + 2*sqrt(x^4 + p*x^2 + 1)*sqrt(p + 2)*x + 1)/(x^4 - 2*x^2 + 1)), -1/4*sqrt(-p - 2)*arctan(sqrt(-p - 2)*x/sqrt(x^4 + p*x^2 + 1)) + 1/8*sqrt(p - 2)*log((x^4 + 2*(p - 1)*x^2 - 2*sqrt(x^4 + p*x^2 + 1)*sqrt(p - 2)*x + 1)/(x^4 + 2*x^2 + 1)), 1/4*sqrt(-p + 2)*arctan(sqrt(-p + 2)*x/sqrt(x^4 + p*x^2 + 1)) - 1/4*sqrt(-p - 2)*arctan(sqrt(-p - 2)*x/sqrt(x^4 + p*x^2 + 1))]`

Sympy [F]

$$\int \frac{\sqrt{1 + px^2 + x^4}}{1 - x^4} dx = - \int \frac{\sqrt{px^2 + x^4 + 1}}{x^4 - 1} dx$$

input `integrate((x**4+p*x**2+1)**(1/2)/(-x**4+1),x)`

output `-Integral(sqrt(p*x**2 + x**4 + 1)/(x**4 - 1), x)`

Maxima [F]

$$\int \frac{\sqrt{1 + px^2 + x^4}}{1 - x^4} dx = \int -\frac{\sqrt{x^4 + px^2 + 1}}{x^4 - 1} dx$$

input `integrate((x^4+p*x^2+1)^(1/2)/(-x^4+1),x, algorithm="maxima")`

output `-integrate(sqrt(x^4 + p*x^2 + 1)/(x^4 - 1), x)`

Giac [F]

$$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx = \int -\frac{\sqrt{x^4+px^2+1}}{x^4-1} dx$$

input `integrate((x^4+p*x^2+1)^(1/2)/(-x^4+1),x, algorithm="giac")`

output `integrate(-sqrt(x^4 + p*x^2 + 1)/(x^4 - 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx = - \int \frac{\sqrt{x^4+px^2+1}}{x^4-1} dx$$

input `int(-(p*x^2 + x^4 + 1)^(1/2)/(x^4 - 1),x)`

output `-int((p*x^2 + x^4 + 1)^(1/2)/(x^4 - 1), x)`

Reduce [F]

$$\int \frac{\sqrt{1+px^2+x^4}}{1-x^4} dx = - \left(\int \frac{\sqrt{x^4+px^2+1}}{x^4-1} dx \right)$$

input `int((x^4+p*x^2+1)^(1/2)/(-x^4+1),x)`

output `- int(sqrt(p*x**2 + x**4 + 1)/(x**4 - 1),x)`

3.68 $\int \frac{\sqrt{1+px^2-x^4}}{1+x^4} dx$

Optimal result	539
Mathematica [C] (verified)	540
Rubi [A] (verified)	540
Maple [C] (verified)	541
Fricas [C] (verification not implemented)	542
Sympy [F]	543
Maxima [F]	543
Giac [F]	543
Mupad [F(-1)]	544
Reduce [F]	544

Optimal result

Integrand size = 24, antiderivative size = 171

$$\int \frac{\sqrt{1+px^2-x^4}}{1+x^4} dx = -\frac{\sqrt{p+\sqrt{4+p^2}} \arctan\left(\frac{\sqrt{p+\sqrt{4+p^2}}x(p-\sqrt{4+p^2}-2x^2)}{2\sqrt{2}\sqrt{1+px^2-x^4}}\right)}{2\sqrt{2}} + \frac{\sqrt{-p+\sqrt{4+p^2}} \operatorname{arctanh}\left(\frac{\sqrt{-p+\sqrt{4+p^2}}x(p+\sqrt{4+p^2}-2x^2)}{2\sqrt{2}\sqrt{1+px^2-x^4}}\right)}{2\sqrt{2}}$$

output

```
1/4*arctanh(1/4*x*(p-2*x^2+(p^2+4)^(1/2))*(-p+(p^2+4)^(1/2))^(1/2)*2^(1/2)
/(-x^4+p*x^2+1)^(1/2))*(-p+(p^2+4)^(1/2))^(1/2)*2^(1/2)-1/4*arctan(1/4*x*(
p-2*x^2-(p^2+4)^(1/2))*(p+(p^2+4)^(1/2))^(1/2)*2^(1/2)/(-x^4+p*x^2+1)^(1/2
))* (p+(p^2+4)^(1/2))^(1/2)*2^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.54

$$\int \frac{\sqrt{1+px^2-x^4}}{1+x^4} dx = \frac{1}{4}i \left(\sqrt{-2i-p} \arctan \left(\frac{\sqrt{-2i-px}}{\sqrt{1+px^2-x^4}} \right) - \sqrt{2i-p} \arctan \left(\frac{\sqrt{2i-px}}{\sqrt{1+px^2-x^4}} \right) \right)$$

input `Integrate[Sqrt[1 + p*x^2 - x^4]/(1 + x^4), x]`

output `(I/4)*(Sqrt[-2*I - p]*ArcTan[(Sqrt[-2*I - p]*x)/Sqrt[1 + p*x^2 - x^4]] - Sqrt[2*I - p]*ArcTan[(Sqrt[2*I - p]*x)/Sqrt[1 + p*x^2 - x^4]])`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {2518}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt{px^2 - x^4 + 1}}{x^4 + 1} dx$$

↓ 2518

$$\frac{\sqrt{\sqrt{p^2+4}-p} \operatorname{arctanh} \left(\frac{\sqrt{\sqrt{p^2+4}-p} (\sqrt{p^2+4+p-2x^2})}{2\sqrt{2}\sqrt{px^2-x^4+1}} \right)}{2\sqrt{2}} - \frac{\sqrt{\sqrt{p^2+4}+p} \operatorname{arctan} \left(\frac{\sqrt{\sqrt{p^2+4}+p} (-\sqrt{p^2+4+p-2x^2})}{2\sqrt{2}\sqrt{px^2-x^4+1}} \right)}{2\sqrt{2}}$$

input `Int[Sqrt[1 + p*x^2 - x^4]/(1 + x^4), x]`

output

```
-1/2*(Sqrt[p + Sqrt[4 + p^2]]*ArcTan[(Sqrt[p + Sqrt[4 + p^2]]*x*(p - Sqrt[4 + p^2] - 2*x^2))/(2*Sqrt[2]*Sqrt[1 + p*x^2 - x^4])])/Sqrt[2] + (Sqrt[-p + Sqrt[4 + p^2]]*ArcTanh[(Sqrt[-p + Sqrt[4 + p^2]]*x*(p + Sqrt[4 + p^2] - 2*x^2))/(2*Sqrt[2]*Sqrt[1 + p*x^2 - x^4])])/(2*Sqrt[2])
```

Defintions of rubi rules used

rule 2518

```
Int[Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]/((d_) + (e_)*(x_)^4), x_Symbol] := With[{q = Sqrt[b^2 - 4*a*c]}, Simp[(-a)*(Sqrt[b + q]/(2*Sqrt[2]*Rt[(-a)*c, 2]*d))*ArcTan[Sqrt[b + q]*x*((b - q + 2*c*x^2)/(2*Sqrt[2]*Rt[(-a)*c, 2]*Sqrt[a + b*x^2 + c*x^4]))], x] + Simp[a*(Sqrt[-b + q]/(2*Sqrt[2]*Rt[(-a)*c, 2]*d))*ArcTanh[Sqrt[-b + q]*x*((b + q + 2*c*x^2)/(2*Sqrt[2]*Rt[(-a)*c, 2]*Sqrt[a + b*x^2 + c*x^4]))], x]] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c*d + a*e, 0] && NegQ[a*c]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.34 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.63

method	result
pseudoelliptic	$-\frac{i \left(\ln(2) + \ln \left(\frac{\sqrt{2i+p} \sqrt{-x^4+px^2+1+2x(i+\frac{p}{2})}}{x^2+i} \right) \right) \sqrt{2i-p} \sqrt{2i+p} + (ip+2) \arctan \left(\frac{\sqrt{-x^4+px^2+1}}{x\sqrt{2i-p}} \right)}{4\sqrt{2i-p}}$
default	$\left(\frac{\sqrt{p+\sqrt{p^2+4}} \sqrt{p^2+4} \ln \left(\frac{-x^4+px^2+1}{x^2} + \frac{\sqrt{-x^4+px^2+1} \sqrt{2} \sqrt{p+\sqrt{p^2+4}}}{x \sqrt{p^2+4}} + \sqrt{p^2+4} \right)}{16} - \frac{\sqrt{p^2+4} (p+\sqrt{p^2+4}) \arctan \left(\frac{2\sqrt{-x^4+px^2+1}}{x} \right)}{8\sqrt{-p+\sqrt{p^2+4}}} \right)$
elliptic	$\left(\frac{\sqrt{p+\sqrt{p^2+4}} \sqrt{p^2+4} \ln \left(\frac{-x^4+px^2+1}{x^2} + \frac{\sqrt{-x^4+px^2+1} \sqrt{2} \sqrt{p+\sqrt{p^2+4}}}{x \sqrt{p^2+4}} + \sqrt{p^2+4} \right)}{16} - \frac{\sqrt{p^2+4} (p+\sqrt{p^2+4}) \arctan \left(\frac{2\sqrt{-x^4+px^2+1}}{x} \right)}{8\sqrt{-p+\sqrt{p^2+4}}} \right)$

input

```
int((-x^4+p*x^2+1)^(1/2)/(x^4+1),x,method=_RETURNVERBOSE)
```

output

```
-1/4/(2*I-p)^(1/2)*(I*(ln(2)+ln(((2*I+p)^(1/2)*(-x^4+p*x^2+1)^(1/2)+2*x*(I
+1/2*p)))/(x^2+I)))*(2*I-p)^(1/2)*(2*I+p)^(1/2)+(I*p+2)*arctan((-x^4+p*x^2+
1)^(1/2)/x/(2*I-p)^(1/2)))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.15 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.34

$$\int \frac{\sqrt{1+px^2-x^4}}{1+x^4} dx$$

$$= \frac{1}{8} \sqrt{-p+2i} \log \left(-\frac{\sqrt{-x^4+px^2+1}(x^2+i) - (ix^3-x)\sqrt{-p+2i}}{x^4+1} \right)$$

$$- \frac{1}{8} \sqrt{-p+2i} \log \left(-\frac{\sqrt{-x^4+px^2+1}(x^2+i) - (-ix^3+x)\sqrt{-p+2i}}{x^4+1} \right)$$

$$- \frac{1}{8} \sqrt{-p-2i} \log \left(-\frac{\sqrt{-x^4+px^2+1}(x^2-i) - (ix^3+x)\sqrt{-p-2i}}{x^4+1} \right)$$

$$+ \frac{1}{8} \sqrt{-p-2i} \log \left(-\frac{\sqrt{-x^4+px^2+1}(x^2-i) - (-ix^3-x)\sqrt{-p-2i}}{x^4+1} \right)$$

input

```
integrate((-x^4+p*x^2+1)^(1/2)/(x^4+1),x, algorithm="fricas")
```

output

```
1/8*sqrt(-p + 2*I)*log(-(sqrt(-x^4 + p*x^2 + 1)*(x^2 + I) - (I*x^3 - x)*sq
rt(-p + 2*I))/(x^4 + 1)) - 1/8*sqrt(-p + 2*I)*log(-(sqrt(-x^4 + p*x^2 + 1)
*(x^2 + I) - (-I*x^3 + x)*sqrt(-p + 2*I))/(x^4 + 1)) - 1/8*sqrt(-p - 2*I)*
log(-(sqrt(-x^4 + p*x^2 + 1)*(x^2 - I) - (I*x^3 + x)*sqrt(-p - 2*I))/(x^4
+ 1)) + 1/8*sqrt(-p - 2*I)*log(-(sqrt(-x^4 + p*x^2 + 1)*(x^2 - I) - (-I*x^
3 - x)*sqrt(-p - 2*I))/(x^4 + 1))
```

Sympy [F]

$$\int \frac{\sqrt{1 + px^2 - x^4}}{1 + x^4} dx = \int \frac{\sqrt{px^2 - x^4 + 1}}{x^4 + 1} dx$$

input `integrate((-x**4+p*x**2+1)**(1/2)/(x**4+1),x)`

output `Integral(sqrt(p*x**2 - x**4 + 1)/(x**4 + 1), x)`

Maxima [F]

$$\int \frac{\sqrt{1 + px^2 - x^4}}{1 + x^4} dx = \int \frac{\sqrt{-x^4 + px^2 + 1}}{x^4 + 1} dx$$

input `integrate((-x^4+p*x^2+1)^(1/2)/(x^4+1),x, algorithm="maxima")`

output `integrate(sqrt(-x^4 + p*x^2 + 1)/(x^4 + 1), x)`

Giac [F]

$$\int \frac{\sqrt{1 + px^2 - x^4}}{1 + x^4} dx = \int \frac{\sqrt{-x^4 + px^2 + 1}}{x^4 + 1} dx$$

input `integrate((-x^4+p*x^2+1)^(1/2)/(x^4+1),x, algorithm="giac")`

output `integrate(sqrt(-x^4 + p*x^2 + 1)/(x^4 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{1 + px^2 - x^4}}{1 + x^4} dx = \int \frac{\sqrt{-x^4 + px^2 + 1}}{x^4 + 1} dx$$

input `int((p*x^2 - x^4 + 1)^(1/2)/(x^4 + 1), x)`output `int((p*x^2 - x^4 + 1)^(1/2)/(x^4 + 1), x)`**Reduce [F]**

$$\int \frac{\sqrt{1 + px^2 - x^4}}{1 + x^4} dx = \int \frac{\sqrt{-x^4 + px^2 + 1}}{x^4 + 1} dx$$

input `int((-x^4+p*x^2+1)^(1/2)/(x^4+1), x)`output `int(sqrt(p*x**2 - x**4 + 1)/(x**4 + 1), x)`

3.69 $\int \frac{a+bx}{(2-x^2)\sqrt[4]{-1+x^2}} dx$

Optimal result	545
Mathematica [C] (warning: unable to verify)	545
Rubi [A] (verified)	546
Maple [F]	549
Fricas [F(-1)]	549
Sympy [F]	550
Maxima [F]	550
Giac [F]	550
Mupad [F(-1)]	551
Reduce [F]	551

Optimal result

Integrand size = 24, antiderivative size = 80

$$\int \frac{a+bx}{(2-x^2)\sqrt[4]{-1+x^2}} dx = \frac{a \arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{-1+x^2}}\right)}{2\sqrt{2}} - b \arctan\left(\sqrt[4]{-1+x^2}\right) + \frac{a \operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{-1+x^2}}\right)}{2\sqrt{2}} + b \operatorname{arctanh}\left(\sqrt[4]{-1+x^2}\right)$$

output -b*arctan((x^2-1)^(1/4))+b*arctanh((x^2-1)^(1/4))+1/4*a*arctan(1/2*x/(x^2-1)^(1/4)*2^(1/2))*2^(1/2)+1/4*a*arctanh(1/2*x/(x^2-1)^(1/4)*2^(1/2))*2^(1/2)

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.19 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.96

$$\int \frac{a + bx}{(2 - x^2) \sqrt[4]{-1 + x^2}} dx$$

$$= \frac{x \left(bx \sqrt[4]{1 - x^2} (-2 + x^2) \operatorname{AppellF1} \left(1, \frac{1}{4}, 1, 2, x^2, \frac{x^2}{2} \right) - \frac{24a \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, x^2, \frac{x^2}{2} \right)}{6 \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, x^2, \frac{x^2}{2} \right) + x^2 \left(2 \operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{4}, 2, \frac{5}{2}, x^2, \frac{x^2}{2} \right) + \operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{4}, 1, \frac{5}{2}, x^2, \frac{x^2}{2} \right) \right)} \right)}{4(-2 + x^2) \sqrt[4]{-1 + x^2}}$$

input `Integrate[(a + b*x)/((2 - x^2)*(-1 + x^2)^(1/4)),x]`

output `(x*(b*x*(1 - x^2)^(1/4)*(-2 + x^2)*AppellF1[1, 1/4, 1, 2, x^2, x^2/2] - (4*a*AppellF1[1/2, 1/4, 1, 3/2, x^2, x^2/2])/(6*AppellF1[1/2, 1/4, 1, 3/2, x^2, x^2/2] + x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, x^2, x^2/2] + AppellF1[3/2, 5/4, 1, 5/2, x^2, x^2/2]))))/(4*(-2 + x^2)*(-1 + x^2)^(1/4))`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.10, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {1343, 309, 353, 73, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + bx}{(2 - x^2) \sqrt[4]{x^2 - 1}} dx$$

$$\downarrow 1343$$

$$a \int \frac{1}{(2 - x^2) \sqrt[4]{x^2 - 1}} dx + b \int \frac{x}{(2 - x^2) \sqrt[4]{x^2 - 1}} dx$$

$$\downarrow 309$$

$$b \int \frac{x}{(2 - x^2) \sqrt[4]{x^2 - 1}} dx + a \left(\frac{\arctan \left(\frac{x}{\sqrt{2} \sqrt[4]{x^2 - 1}} \right)}{2\sqrt{2}} + \frac{\operatorname{arctanh} \left(\frac{x}{\sqrt{2} \sqrt[4]{x^2 - 1}} \right)}{2\sqrt{2}} \right)$$

$$\frac{1}{2}b \int \frac{1}{(2-x^2)\sqrt[4]{x^2-1}} dx^2 + a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} \right)$$

$$2b \int \frac{x^4}{1-x^8} d\sqrt[4]{x^2-1} + a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} \right)$$

$$2b \left(\frac{1}{2} \int \frac{1}{1-x^4} d\sqrt[4]{x^2-1} - \frac{1}{2} \int \frac{1}{x^4+1} d\sqrt[4]{x^2-1} \right) + a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} \right)$$

$$2b \left(\frac{1}{2} \int \frac{1}{1-x^4} d\sqrt[4]{x^2-1} - \frac{1}{2} \arctan\left(\sqrt[4]{x^2-1}\right) \right) + a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} \right)$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{x^2-1}}\right)}{2\sqrt{2}} \right) + 2b \left(\frac{1}{2} \operatorname{arctanh}\left(\sqrt[4]{x^2-1}\right) - \frac{1}{2} \arctan\left(\sqrt[4]{x^2-1}\right) \right)$$

input `Int[(a + b*x)/((2 - x^2)*(-1 + x^2)^(1/4)),x]`

output

```
a*(ArcTan[x/(Sqrt[2]*(-1 + x^2)^(1/4))]/(2*Sqrt[2]) + ArcTanh[x/(Sqrt[2]*(-1 + x^2)^(1/4))]/(2*Sqrt[2])) + 2*b*(-1/2*ArcTan[(-1 + x^2)^(1/4)] + ArcTanh[(-1 + x^2)^(1/4)]/2)
```

Defintions of rubi rules used

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 219

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 309

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/4))*((c_) + (d_.)*(x_)^2), x_Symbol] := Wit
h[{q = Rt[-b^2/a, 4]}, Simp[(b/(2*Sqrt[2]*a*d*q))*ArcTan[q*(x/(Sqrt[2]*(a +
b*x^2)^(1/4)))]], x] + Simp[(b/(2*Sqrt[2]*a*d*q))*ArcTanh[q*(x/(Sqrt[2]*(a
+ b*x^2)^(1/4)))]], x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] &&
NegQ[b^2/a]
```

rule 353

```
Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol]
:= Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[
{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]
```

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 1343 `Int[((g_) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[g Int[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + Simp[h Int[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; FreeQ[{a, c, d, f, g, h, p, q}, x]`

Maple [F]

$$\int \frac{bx + a}{(-x^2 + 2)(x^2 - 1)^{\frac{1}{4}}} dx$$

input `int((b*x+a)/(-x^2+2)/(x^2-1)^(1/4),x)`

output `int((b*x+a)/(-x^2+2)/(x^2-1)^(1/4),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{a + bx}{(2 - x^2) \sqrt[4]{-1 + x^2}} dx = \text{Timed out}$$

input `integrate((b*x+a)/(-x^2+2)/(x^2-1)^(1/4),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{a + bx}{(2 - x^2)\sqrt[4]{-1 + x^2}} dx$$

$$= - \int \frac{a}{x^2\sqrt[4]{x^2 - 1} - 2\sqrt[4]{x^2 - 1}} dx - \int \frac{bx}{x^2\sqrt[4]{x^2 - 1} - 2\sqrt[4]{x^2 - 1}} dx$$

input `integrate((b*x+a)/(-x**2+2)/(x**2-1)**(1/4), x)`

output `-Integral(a/(x**2*(x**2 - 1)**(1/4) - 2*(x**2 - 1)**(1/4)), x) - Integral(b*x/(x**2*(x**2 - 1)**(1/4) - 2*(x**2 - 1)**(1/4)), x)`

Maxima [F]

$$\int \frac{a + bx}{(2 - x^2)\sqrt[4]{-1 + x^2}} dx = \int -\frac{bx + a}{(x^2 - 1)^{\frac{1}{4}}(x^2 - 2)} dx$$

input `integrate((b*x+a)/(-x^2+2)/(x^2-1)^(1/4), x, algorithm="maxima")`

output `-integrate((b*x + a)/((x^2 - 1)^(1/4)*(x^2 - 2)), x)`

Giac [F]

$$\int \frac{a + bx}{(2 - x^2)\sqrt[4]{-1 + x^2}} dx = \int -\frac{bx + a}{(x^2 - 1)^{\frac{1}{4}}(x^2 - 2)} dx$$

input `integrate((b*x+a)/(-x^2+2)/(x^2-1)^(1/4), x, algorithm="giac")`

output `integrate(-(b*x + a)/((x^2 - 1)^(1/4)*(x^2 - 2)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + bx}{(2 - x^2) \sqrt[4]{-1 + x^2}} dx = \int -\frac{a + bx}{(x^2 - 1)^{1/4} (x^2 - 2)} dx$$

input `int(-(a + b*x)/((x^2 - 1)^(1/4)*(x^2 - 2)), x)`output `int(-(a + b*x)/((x^2 - 1)^(1/4)*(x^2 - 2)), x)`**Reduce [F]**

$$\int \frac{a + bx}{(2 - x^2) \sqrt[4]{-1 + x^2}} dx = - \left(\int \frac{x}{(x^2 - 1)^{1/4} x^2 - 2 (x^2 - 1)^{1/4}} dx \right) b$$

$$- \left(\int \frac{1}{(x^2 - 1)^{1/4} x^2 - 2 (x^2 - 1)^{1/4}} dx \right) a$$

input `int((b*x+a)/(-x^2+2)/(x^2-1)^(1/4), x)`output `- (int(x/((x**2 - 1)**(1/4)*x**2 - 2*(x**2 - 1)**(1/4)), x)*b + int(1/((x**2 - 1)**(1/4)*x**2 - 2*(x**2 - 1)**(1/4)), x)*a)`

3.70 $\int \frac{a+bx}{\sqrt[4]{-1-x^2}(2+x^2)} dx$

Optimal result	552
Mathematica [C] (warning: unable to verify)	552
Rubi [A] (verified)	553
Maple [F]	556
Fricas [F(-1)]	556
Sympy [F]	557
Maxima [F]	557
Giac [F]	557
Mupad [F(-1)]	558
Reduce [F]	558

Optimal result

Integrand size = 24, antiderivative size = 88

$$\int \frac{a+bx}{\sqrt[4]{-1-x^2}(2+x^2)} dx = \frac{a \arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{-1-x^2}}\right)}{2\sqrt{2}} + b \arctan\left(\sqrt[4]{-1-x^2}\right) + \frac{a \operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{-1-x^2}}\right)}{2\sqrt{2}} - b \operatorname{arctanh}\left(\sqrt[4]{-1-x^2}\right)$$

output `b*arctan((-x^2-1)^(1/4))-b*arctanh((-x^2-1)^(1/4))+1/4*a*arctan(1/2*x/(-x^2-1)^(1/4)*2^(1/2))*2^(1/2)+1/4*a*arctanh(1/2*x/(-x^2-1)^(1/4)*2^(1/2))*2^(1/2)`

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.18 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.84

$$\int \frac{a + bx}{\sqrt[4]{-1 - x^2} (2 + x^2)} dx$$

$$= \frac{x \left(bx \sqrt[4]{1 + x^2} \operatorname{AppellF1} \left(1, \frac{1}{4}, 1, 2, -x^2, -\frac{x^2}{2} \right) - \frac{24a \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, -x^2, -\frac{x^2}{2} \right)}{(2+x^2) \left(-6 \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, -x^2, -\frac{x^2}{2} \right) + x^2 \left(2 \operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{4}, 2, \frac{5}{2}, -x^2, -\frac{x^2}{2} \right) + \operatorname{AppellF1} \left(\frac{3}{2}, \frac{5}{4}, 1, \frac{5}{2}, -x^2, -\frac{1}{2}x^2 \right) \right) \right)}{4 \sqrt[4]{-1 - x^2}} \right)}{4 \sqrt[4]{-1 - x^2}}$$

input `Integrate[(a + b*x)/((-1 - x^2)^(1/4)*(2 + x^2)),x]`

output `(x*(b*x*(1 + x^2)^(1/4)*AppellF1[1, 1/4, 1, 2, -x^2, -1/2*x^2] - (24*a*AppellF1[1/2, 1/4, 1, 3/2, -x^2, -1/2*x^2])/((2 + x^2)*(-6*AppellF1[1/2, 1/4, 1, 3/2, -x^2, -1/2*x^2] + x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, -x^2, -1/2*x^2] + AppellF1[3/2, 5/4, 1, 5/2, -x^2, -1/2*x^2])))))/(4*(-1 - x^2)^(1/4))`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.09, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.292$, Rules used = {1343, 309, 353, 73, 827, 216, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + bx}{\sqrt[4]{-x^2 - 1} (x^2 + 2)} dx$$

$$\downarrow 1343$$

$$a \int \frac{1}{\sqrt[4]{-x^2 - 1} (x^2 + 2)} dx + b \int \frac{x}{\sqrt[4]{-x^2 - 1} (x^2 + 2)} dx$$

$$\downarrow 309$$

$$b \int \frac{x}{\sqrt[4]{-x^2 - 1} (x^2 + 2)} dx + a \left(\frac{\arctan \left(\frac{x}{\sqrt{2} \sqrt[4]{-x^2 - 1}} \right)}{2\sqrt{2}} + \frac{\operatorname{arctanh} \left(\frac{x}{\sqrt{2} \sqrt[4]{-x^2 - 1}} \right)}{2\sqrt{2}} \right)$$

$$\frac{1}{2}b \int \frac{1}{\sqrt[4]{-x^2-1}(x^2+2)} dx^2 + a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} \right)$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} \right) - 2b \int \frac{x^4}{1-x^8} d\sqrt[4]{-x^2-1}$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} \right) - 2b \left(\frac{1}{2} \int \frac{1}{1-x^4} d\sqrt[4]{-x^2-1} - \frac{1}{2} \int \frac{1}{x^4+1} d\sqrt[4]{-x^2-1} \right)$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} \right) - 2b \left(\frac{1}{2} \int \frac{1}{1-x^4} d\sqrt[4]{-x^2-1} - \frac{1}{2} \arctan\left(\sqrt[4]{-x^2-1}\right) \right)$$

$$a \left(\frac{\arctan\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{2}\sqrt[4]{-x^2-1}}\right)}{2\sqrt{2}} \right) - 2b \left(\frac{1}{2} \operatorname{arctanh}\left(\sqrt[4]{-x^2-1}\right) - \frac{1}{2} \arctan\left(\sqrt[4]{-x^2-1}\right) \right)$$

input `Int[(a + b*x)/((-1 - x^2)^(1/4)*(2 + x^2)),x]`

output

```
a*(ArcTan[x/(Sqrt[2]*(-1 - x^2)^(1/4))]/(2*Sqrt[2]) + ArcTanh[x/(Sqrt[2]*(-1 - x^2)^(1/4))]/(2*Sqrt[2])) - 2*b*(-1/2*ArcTan[(-1 - x^2)^(1/4)] + ArcTanh[(-1 - x^2)^(1/4)]/2)
```

Definitions of rubi rules used

rule 73

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && Lt
Q[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntL
inearQ[a, b, c, d, m, n, x]
```

rule 216

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 219

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

rule 309

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/4))*((c_) + (d_.)*(x_)^2), x_Symbol] := Wit
h[{q = Rt[-b^2/a, 4]}, Simp[(b/(2*Sqrt[2]*a*d*q))*ArcTan[q*(x/(Sqrt[2]*(a +
b*x^2)^(1/4)))]], x] + Simp[(b/(2*Sqrt[2]*a*d*q))*ArcTanh[q*(x/(Sqrt[2]*(a
+ b*x^2)^(1/4)))]], x]] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] &&
NegQ[b^2/a]
```

rule 353

```
Int[(x_)*((a_) + (b_.)*(x_)^2)^(p_.)*((c_) + (d_.)*(x_)^2)^(q_.), x_Symbol]
:= Simp[1/2 Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^2], x] /; FreeQ[
{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0]
```

rule 827 `Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Simp[s/(2*b) Int[1/(r + s*x^2), x], x] - Simp[s/(2*b) Int[1/(r - s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

rule 1343 `Int[((g_) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[g Int[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + Simp[h Int[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; FreeQ[{a, c, d, f, g, h, p, q}, x]`

Maple [F]

$$\int \frac{bx + a}{(-x^2 - 1)^{\frac{1}{4}}(x^2 + 2)} dx$$

input `int((b*x+a)/(-x^2-1)^(1/4)/(x^2+2),x)`

output `int((b*x+a)/(-x^2-1)^(1/4)/(x^2+2),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{a + bx}{\sqrt[4]{-1 - x^2}(2 + x^2)} dx = \text{Timed out}$$

input `integrate((b*x+a)/(-x^2-1)^(1/4)/(x^2+2),x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{a + bx}{\sqrt[4]{-1 - x^2} (2 + x^2)} dx = \int \frac{a + bx}{\sqrt[4]{-x^2 - 1} (x^2 + 2)} dx$$

input `integrate((b*x+a)/(-x**2-1)**(1/4)/(x**2+2),x)`

output `Integral((a + b*x)/((-x**2 - 1)**(1/4)*(x**2 + 2)), x)`

Maxima [F]

$$\int \frac{a + bx}{\sqrt[4]{-1 - x^2} (2 + x^2)} dx = \int \frac{bx + a}{(x^2 + 2)(-x^2 - 1)^{\frac{1}{4}}} dx$$

input `integrate((b*x+a)/(-x^2-1)^(1/4)/(x^2+2),x, algorithm="maxima")`

output `integrate((b*x + a)/((x^2 + 2)*(-x^2 - 1)^(1/4)), x)`

Giac [F]

$$\int \frac{a + bx}{\sqrt[4]{-1 - x^2} (2 + x^2)} dx = \int \frac{bx + a}{(x^2 + 2)(-x^2 - 1)^{\frac{1}{4}}} dx$$

input `integrate((b*x+a)/(-x^2-1)^(1/4)/(x^2+2),x, algorithm="giac")`

output `integrate((b*x + a)/((x^2 + 2)*(-x^2 - 1)^(1/4)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + bx}{\sqrt[4]{-1 - x^2} (2 + x^2)} dx = \int \frac{a + bx}{(-x^2 - 1)^{1/4} (x^2 + 2)} dx$$

input `int((a + b*x)/((- x^2 - 1)^(1/4)*(x^2 + 2)), x)`output `int((a + b*x)/((- x^2 - 1)^(1/4)*(x^2 + 2)), x)`**Reduce [F]**

$$\int \frac{a + bx}{\sqrt[4]{-1 - x^2} (2 + x^2)} dx = - \left(\left(\int \frac{x}{(x^2 + 1)^{1/4} x^2 + 2 (x^2 + 1)^{1/4}} dx \right) b + \left(\int \frac{1}{(x^2 + 1)^{1/4} x^2 + 2 (x^2 + 1)^{1/4}} dx \right) a \right) (-1)^{3/4}$$

input `int((b*x+a)/(-x^2-1)^(1/4)/(x^2+2), x)`output `(int(x/((x**2 + 1)**(1/4)*x**2 + 2*(x**2 + 1)**(1/4)), x)*b + int(1/((x**2 + 1)**(1/4)*x**2 + 2*(x**2 + 1)**(1/4)), x)*a)/(- 1)**(1/4)`

3.71 $\int \frac{a+bx}{\sqrt[4]{1-x^2}(2-x^2)} dx$

Optimal result	559
Mathematica [C] (warning: unable to verify)	560
Rubi [A] (verified)	560
Maple [F]	562
Fricas [F(-1)]	562
Sympy [F]	562
Maxima [F]	563
Giac [F]	563
Mupad [F(-1)]	563
Reduce [F]	564

Optimal result

Integrand size = 26, antiderivative size = 149

$$\int \frac{a+bx}{\sqrt[4]{1-x^2}(2-x^2)} dx = \frac{b \arctan\left(\frac{1-\sqrt{1-x^2}}{\sqrt{2}\sqrt[4]{1-x^2}}\right)}{\sqrt{2}} + \frac{1}{2}a \arctan\left(\frac{1-\sqrt{1-x^2}}{x\sqrt[4]{1-x^2}}\right) + \frac{b \operatorname{arctanh}\left(\frac{1+\sqrt{1-x^2}}{\sqrt{2}\sqrt[4]{1-x^2}}\right)}{\sqrt{2}} + \frac{1}{2}a \operatorname{arctanh}\left(\frac{1+\sqrt{1-x^2}}{x\sqrt[4]{1-x^2}}\right)$$

output

```
1/2*a*arctan((1-(-x^2+1)^(1/2))/x/(-x^2+1)^(1/4))+1/2*a*arctanh((1+(-x^2+1)^(1/2))/x/(-x^2+1)^(1/4))+1/2*b*arctan(1/2*(1-(-x^2+1)^(1/2))/(-x^2+1)^(1/4)*2^(1/2))*2^(1/2)+1/2*b*arctanh(1/2*(1+(-x^2+1)^(1/2))/(-x^2+1)^(1/4)*2^(1/2))*2^(1/2)
```


Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.19 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.97

$$\int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx = \frac{1}{4} bx^2 \operatorname{AppellF1} \left(1, \frac{1}{4}, 1, 2, x^2, \frac{x^2}{2} \right) - \frac{6ax \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, x^2, \frac{x^2}{2} \right)}{\sqrt[4]{1 - x^2} (-2 + x^2) (6 \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, x^2, \frac{x^2}{2} \right) + x^2 (2 \operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{4}, 2, \frac{5}{2}, x^2, \frac{x^2}{2} \right) + \operatorname{AppellF1} \left(\frac{3}{2}, \frac{5}{4}, 1, \frac{5}{2}, x^2, \frac{x^2}{2} \right)))}$$

input `Integrate[(a + b*x)/((1 - x^2)^(1/4)*(2 - x^2)),x]`

output `(b*x^2*AppellF1[1, 1/4, 1, 2, x^2, x^2/2])/4 - (6*a*x*AppellF1[1/2, 1/4, 1, 3/2, x^2, x^2/2])/((1 - x^2)^(1/4)*(-2 + x^2)*(6*AppellF1[1/2, 1/4, 1, 3/2, x^2, x^2/2] + x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, x^2, x^2/2] + AppellF1[3/2, 5/4, 1, 5/2, x^2, x^2/2])))`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {1343, 308, 348}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx \\ & \quad \downarrow 1343 \\ & a \int \frac{1}{\sqrt[4]{1 - x^2} (2 - x^2)} dx + b \int \frac{x}{\sqrt[4]{1 - x^2} (2 - x^2)} dx \\ & \quad \downarrow 308 \\ & b \int \frac{x}{\sqrt[4]{1 - x^2} (2 - x^2)} dx + a \left(\frac{1}{2} \arctan \left(\frac{1 - \sqrt{1 - x^2}}{x \sqrt[4]{1 - x^2}} \right) + \frac{1}{2} \operatorname{arctanh} \left(\frac{\sqrt{1 - x^2} + 1}{x \sqrt[4]{1 - x^2}} \right) \right) \end{aligned}$$

$$\begin{array}{c}
 \downarrow \text{348} \\
 a \left(\frac{1}{2} \arctan \left(\frac{1 - \sqrt{1-x^2}}{x^4 \sqrt{1-x^2}} \right) + \frac{1}{2} \operatorname{arctanh} \left(\frac{\sqrt{1-x^2} + 1}{x^4 \sqrt{1-x^2}} \right) \right) + \\
 b \left(\frac{\arctan \left(\frac{1 - \sqrt{1-x^2}}{\sqrt{2}^4 \sqrt{1-x^2}} \right)}{\sqrt{2}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt{1-x^2} + 1}{\sqrt{2}^4 \sqrt{1-x^2}} \right)}{\sqrt{2}} \right)
 \end{array}$$

input `Int[(a + b*x)/((1 - x^2)^(1/4)*(2 - x^2)),x]`

output `b*(ArcTan[(1 - Sqrt[1 - x^2])/(Sqrt[2]*(1 - x^2)^(1/4))]/Sqrt[2] + ArcTanh[(1 + Sqrt[1 - x^2])/(Sqrt[2]*(1 - x^2)^(1/4))]/Sqrt[2]) + a*(ArcTan[(1 - Sqrt[1 - x^2])/(x*(1 - x^2)^(1/4))]/2 + ArcTanh[(1 + Sqrt[1 - x^2])/(x*(1 - x^2)^(1/4))]/2)`

Defintions of rubi rules used

rule 308 `Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b^2/a, 4]}, Simp[(-b/(2*a*d*q))*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))], x] - Simp[(b/(2*a*d*q))*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]`

rule 348 `Int[(x_)/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Simp[(-(Sqrt[2]*Rt[a, 4]*d)^(-1))*ArcTan[(Rt[a, 4]^2 - Sqrt[a + b*x^2])/(Sqrt[2]*Rt[a, 4]*(a + b*x^2)^(1/4))], x] - Simp[(1/(Sqrt[2]*Rt[a, 4]*d))*ArcTanh[(Rt[a, 4]^2 + Sqrt[a + b*x^2])/(Sqrt[2]*Rt[a, 4]*(a + b*x^2)^(1/4))], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[a]`

rule 1343 `Int[((g_) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[g Int[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + Simp[h Int[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; FreeQ[{a, c, d, f, g, h, p, q}, x]`

Maple [F]

$$\int \frac{bx + a}{(-x^2 + 1)^{\frac{1}{4}} (-x^2 + 2)} dx$$

input `int((b*x+a)/(-x^2+1)^(1/4)/(-x^2+2), x)`

output `int((b*x+a)/(-x^2+1)^(1/4)/(-x^2+2), x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx = \text{Timed out}$$

input `integrate((b*x+a)/(-x^2+1)^(1/4)/(-x^2+2), x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\begin{aligned} & \int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx \\ &= - \int \frac{a}{x^2 \sqrt[4]{1 - x^2} - 2 \sqrt[4]{1 - x^2}} dx - \int \frac{bx}{x^2 \sqrt[4]{1 - x^2} - 2 \sqrt[4]{1 - x^2}} dx \end{aligned}$$

input `integrate((b*x+a)/(-x**2+1)**(1/4)/(-x**2+2), x)`

output `-Integral(a/(x**2*(1 - x**2)**(1/4) - 2*(1 - x**2)**(1/4)), x) - Integral(b*x/(x**2*(1 - x**2)**(1/4) - 2*(1 - x**2)**(1/4)), x)`

Maxima [F]

$$\int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx = \int -\frac{bx + a}{(x^2 - 2)(-x^2 + 1)^{\frac{1}{4}}} dx$$

input `integrate((b*x+a)/(-x^2+1)^(1/4)/(-x^2+2),x, algorithm="maxima")`

output `-integrate((b*x + a)/((x^2 - 2)*(-x^2 + 1)^(1/4)), x)`

Giac [F]

$$\int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx = \int -\frac{bx + a}{(x^2 - 2)(-x^2 + 1)^{\frac{1}{4}}} dx$$

input `integrate((b*x+a)/(-x^2+1)^(1/4)/(-x^2+2),x, algorithm="giac")`

output `integrate(-(b*x + a)/((x^2 - 2)*(-x^2 + 1)^(1/4)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx = \int -\frac{a + bx}{(1 - x^2)^{1/4} (x^2 - 2)} dx$$

input `int(-(a + b*x)/((1 - x^2)^(1/4)*(x^2 - 2)),x)`

output `int(-(a + b*x)/((1 - x^2)^(1/4)*(x^2 - 2)), x)`

Reduce [F]

$$\int \frac{a + bx}{\sqrt[4]{1 - x^2} (2 - x^2)} dx = - \left(\int \frac{x}{(-x^2 + 1)^{\frac{1}{4}} x^2 - 2(-x^2 + 1)^{\frac{1}{4}}} dx \right) b - \left(\int \frac{1}{(-x^2 + 1)^{\frac{1}{4}} x^2 - 2(-x^2 + 1)^{\frac{1}{4}}} dx \right) a$$

input `int((b*x+a)/(-x^2+1)^(1/4)/(-x^2+2),x)`

output `- (int(x/((- x**2 + 1)**(1/4)*x**2 - 2*(- x**2 + 1)**(1/4)),x)*b + int(1/((- x**2 + 1)**(1/4)*x**2 - 2*(- x**2 + 1)**(1/4)),x)*a)`

3.72
$$\int \frac{a+bx}{\sqrt[4]{1+x^2}(2+x^2)} dx$$

Optimal result	565
Mathematica [C] (warning: unable to verify)	566
Rubi [A] (verified)	566
Maple [F]	568
Fricas [F(-1)]	568
Sympy [F]	568
Maxima [F]	569
Giac [F]	569
Mupad [F(-1)]	569
Reduce [F]	570

Optimal result

Integrand size = 22, antiderivative size = 135

$$\int \frac{a+bx}{\sqrt[4]{1+x^2}(2+x^2)} dx = -\frac{b \arctan\left(\frac{1-\sqrt{1+x^2}}{\sqrt{2}\sqrt[4]{1+x^2}}\right)}{\sqrt{2}} - \frac{1}{2}a \arctan\left(\frac{1+\sqrt{1+x^2}}{x\sqrt[4]{1+x^2}}\right) - \frac{1}{2}a \operatorname{arctanh}\left(\frac{1-\sqrt{1+x^2}}{x\sqrt[4]{1+x^2}}\right) - \frac{b \operatorname{arctanh}\left(\frac{1+\sqrt{1+x^2}}{\sqrt{2}\sqrt[4]{1+x^2}}\right)}{\sqrt{2}}$$

output

```
-1/2*a*arctan((1+(x^2+1)^(1/2))/x/(x^2+1)^(1/4))-1/2*a*arctanh((1-(x^2+1)^(1/2))/x/(x^2+1)^(1/4))-1/2*b*arctan(1/2*(1-(x^2+1)^(1/2))/(x^2+1)^(1/4)*2^(1/2))-1/2*b*arctanh(1/2*(1+(x^2+1)^(1/2))/(x^2+1)^(1/4)*2^(1/2))
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.14 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.13

$$\int \frac{a + bx}{\sqrt[4]{1 + x^2} (2 + x^2)} dx = \frac{1}{4} bx^2 \operatorname{AppellF1} \left(1, \frac{1}{4}, 1, 2, -x^2, -\frac{x^2}{2} \right) - \frac{6ax \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, -x^2, -\frac{x^2}{2} \right) - \sqrt[4]{1 + x^2} (2 + x^2) \left(-6 \operatorname{AppellF1} \left(\frac{1}{2}, \frac{1}{4}, 1, \frac{3}{2}, -x^2, -\frac{x^2}{2} \right) + x^2 \left(2 \operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{4}, 2, \frac{5}{2}, -x^2, -\frac{x^2}{2} \right) + \operatorname{AppellF1} \left(\frac{3}{2}, \frac{1}{4}, 2, \frac{5}{2}, -x^2, -\frac{x^2}{2} \right) \right) \right)}{\sqrt[4]{1 + x^2} (2 + x^2)}$$

input `Integrate[(a + b*x)/((1 + x^2)^(1/4)*(2 + x^2)),x]`

output `(b*x^2*AppellF1[1, 1/4, 1, 2, -x^2, -1/2*x^2])/4 - (6*a*x*AppellF1[1/2, 1/4, 1, 3/2, -x^2, -1/2*x^2])/((1 + x^2)^(1/4)*(2 + x^2))*(-6*AppellF1[1/2, 1/4, 1, 3/2, -x^2, -1/2*x^2] + x^2*(2*AppellF1[3/2, 1/4, 2, 5/2, -x^2, -1/2*x^2] + AppellF1[3/2, 5/4, 1, 5/2, -x^2, -1/2*x^2]))`

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.01, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {1343, 308, 348}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + bx}{\sqrt[4]{x^2 + 1} (x^2 + 2)} dx$$

↓ 1343

$$a \int \frac{1}{\sqrt[4]{x^2 + 1} (x^2 + 2)} dx + b \int \frac{x}{\sqrt[4]{x^2 + 1} (x^2 + 2)} dx$$

↓ 308

$$b \int \frac{x}{\sqrt[4]{x^2 + 1} (x^2 + 2)} dx + a \left(-\frac{1}{2} \arctan \left(\frac{\sqrt{x^2 + 1} + 1}{x \sqrt[4]{x^2 + 1}} \right) - \frac{1}{2} \operatorname{arctanh} \left(\frac{1 - \sqrt{x^2 + 1}}{x \sqrt[4]{x^2 + 1}} \right) \right)$$

$$\begin{array}{c}
 \downarrow \text{348} \\
 a \left(-\frac{1}{2} \arctan \left(\frac{\sqrt{x^2+1}+1}{x\sqrt{x^2+1}} \right) - \frac{1}{2} \operatorname{arctanh} \left(\frac{1-\sqrt{x^2+1}}{x\sqrt{x^2+1}} \right) \right) + \\
 b \left(-\frac{\arctan \left(\frac{1-\sqrt{x^2+1}}{\sqrt{2}\sqrt{x^2+1}} \right)}{\sqrt{2}} - \frac{\operatorname{arctanh} \left(\frac{\sqrt{x^2+1}+1}{\sqrt{2}\sqrt{x^2+1}} \right)}{\sqrt{2}} \right)
 \end{array}$$

input `Int[(a + b*x)/((1 + x^2)^(1/4)*(2 + x^2)),x]`

output `a*(-1/2*ArcTan[(1 + Sqrt[1 + x^2])/(x*(1 + x^2)^(1/4))] - ArcTanh[(1 - Sqrt[1 + x^2])/(x*(1 + x^2)^(1/4))]/2) + b*(-(ArcTan[(1 - Sqrt[1 + x^2])/(Sqrt[2]*(1 + x^2)^(1/4))]/Sqrt[2]) - ArcTanh[(1 + Sqrt[1 + x^2])/(Sqrt[2]*(1 + x^2)^(1/4))]/Sqrt[2])`

Defintions of rubi rules used

rule 308 `Int[1/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b^2/a, 4]}, Simp[(-b/(2*a*d*q))*ArcTan[(b + q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))], x] - Simp[(b/(2*a*d*q))*ArcTanh[(b - q^2*Sqrt[a + b*x^2])/(q^3*x*(a + b*x^2)^(1/4))], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[b^2/a]`

rule 348 `Int[(x_)/(((a_) + (b_.)*(x_)^2)^(1/4)*((c_) + (d_.)*(x_)^2)), x_Symbol] := Simp[(-(Sqrt[2]*Rt[a, 4]*d)^(-1))*ArcTan[(Rt[a, 4]^2 - Sqrt[a + b*x^2])/(Sqrt[2]*Rt[a, 4]*(a + b*x^2)^(1/4))], x] - Simp[(1/(Sqrt[2]*Rt[a, 4]*d))*ArcTanh[(Rt[a, 4]^2 + Sqrt[a + b*x^2])/(Sqrt[2]*Rt[a, 4]*(a + b*x^2)^(1/4))], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b*c - 2*a*d, 0] && PosQ[a]`

rule 1343 `Int[((g_) + (h_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[g Int[(a + c*x^2)^p*(d + f*x^2)^q, x], x] + Simp[h Int[x*(a + c*x^2)^p*(d + f*x^2)^q, x], x] /; FreeQ[{a, c, d, f, g, h, p, q}, x]`

Maple [F]

$$\int \frac{bx + a}{(x^2 + 1)^{\frac{1}{4}}(x^2 + 2)} dx$$

input `int((b*x+a)/(x^2+1)^(1/4)/(x^2+2), x)`

output `int((b*x+a)/(x^2+1)^(1/4)/(x^2+2), x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{a + bx}{\sqrt[4]{1 + x^2}(2 + x^2)} dx = \text{Timed out}$$

input `integrate((b*x+a)/(x^2+1)^(1/4)/(x^2+2), x, algorithm="fricas")`

output `Timed out`

Sympy [F]

$$\int \frac{a + bx}{\sqrt[4]{1 + x^2}(2 + x^2)} dx = \int \frac{a + bx}{\sqrt[4]{x^2 + 1}(x^2 + 2)} dx$$

input `integrate((b*x+a)/(x**2+1)**(1/4)/(x**2+2), x)`

output `Integral((a + b*x)/((x**2 + 1)**(1/4)*(x**2 + 2)), x)`

Maxima [F]

$$\int \frac{a + bx}{\sqrt[4]{1 + x^2} (2 + x^2)} dx = \int \frac{bx + a}{(x^2 + 2)(x^2 + 1)^{\frac{1}{4}}} dx$$

input `integrate((b*x+a)/(x^2+1)^(1/4)/(x^2+2),x, algorithm="maxima")`

output `integrate((b*x + a)/((x^2 + 2)*(x^2 + 1)^(1/4)), x)`

Giac [F]

$$\int \frac{a + bx}{\sqrt[4]{1 + x^2} (2 + x^2)} dx = \int \frac{bx + a}{(x^2 + 2)(x^2 + 1)^{\frac{1}{4}}} dx$$

input `integrate((b*x+a)/(x^2+1)^(1/4)/(x^2+2),x, algorithm="giac")`

output `integrate((b*x + a)/((x^2 + 2)*(x^2 + 1)^(1/4)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{a + bx}{\sqrt[4]{1 + x^2} (2 + x^2)} dx = \int \frac{a + bx}{(x^2 + 1)^{1/4} (x^2 + 2)} dx$$

input `int((a + b*x)/((x^2 + 1)^(1/4)*(x^2 + 2)),x)`

output `int((a + b*x)/((x^2 + 1)^(1/4)*(x^2 + 2)), x)`

Reduce [F]

$$\int \frac{a + bx}{\sqrt[4]{1 + x^2} (2 + x^2)} dx = \left(\int \frac{x}{(x^2 + 1)^{\frac{1}{4}} x^2 + 2 (x^2 + 1)^{\frac{1}{4}}} dx \right) b + \left(\int \frac{1}{(x^2 + 1)^{\frac{1}{4}} x^2 + 2 (x^2 + 1)^{\frac{1}{4}}} dx \right) a$$

input `int((b*x+a)/(x^2+1)^(1/4)/(x^2+2),x)`

output `int(x/((x**2 + 1)**(1/4)*x**2 + 2*(x**2 + 1)**(1/4)),x)*b + int(1/((x**2 + 1)**(1/4)*x**2 + 2*(x**2 + 1)**(1/4)),x)*a`

3.73 $\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx$

Optimal result	571
Mathematica [C] (verified)	571
Rubi [A] (verified)	572
Maple [C] (verified)	573
Fricas [B] (verification not implemented)	574
Sympy [F]	575
Maxima [F]	575
Giac [F]	575
Mupad [B] (verification not implemented)	576
Reduce [F]	576

Optimal result

Integrand size = 22, antiderivative size = 127

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = -\frac{\arctan\left(\frac{\sqrt{3}(1-\sqrt[3]{2x})}{\sqrt{1-x^3}}\right)}{3 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan\left(\frac{\sqrt{1-x^3}}{\sqrt{3}}\right)}{3 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{1+\sqrt[3]{2x}}{\sqrt{1-x^3}}\right)}{3 \cdot 2^{2/3}} + \frac{\operatorname{arctanh}(\sqrt{1-x^3})}{9 \cdot 2^{2/3}}$$

output

```
-1/6*arctanh((1+2^(1/3)*x)/(-x^3+1)^(1/2))*2^(1/3)+1/18*arctanh((-x^3+1)^(1/2))*2^(1/3)-1/18*arctan((1-2^(1/3)*x)*3^(1/2)/(-x^3+1)^(1/2))*2^(1/3)*3^(1/2)+1/18*arctan(1/3*(-x^3+1)^(1/2)*3^(1/2))*2^(1/3)*3^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.22

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = \frac{1}{8}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, x^3, \frac{x^3}{4}\right)$$

input `Integrate[x/(Sqrt[1 - x^3]*(4 - x^3)),x]`

output `(x^2*AppellF1[2/3, 1/2, 1, 5/3, x^3, x^3/4])/8`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {986}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx$$

↓ 986

$$-\frac{\arctan\left(\frac{\sqrt{3}(1-\sqrt[3]{2x})}{\sqrt{1-x^3}}\right)}{3 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan\left(\frac{\sqrt{1-x^3}}{\sqrt{3}}\right)}{3 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt[3]{2x+1}}{\sqrt{1-x^3}}\right)}{3 \cdot 2^{2/3}} + \frac{\operatorname{arctanh}\left(\sqrt{1-x^3}\right)}{9 \cdot 2^{2/3}}$$

input `Int[x/(Sqrt[1 - x^3]*(4 - x^3)),x]`

output `-1/3*ArcTan[(Sqrt[3]*(1 - 2^(1/3)*x))/Sqrt[1 - x^3]]/(2^(2/3)*Sqrt[3]) + ArcTan[Sqrt[1 - x^3]/Sqrt[3]]/(3*2^(2/3)*Sqrt[3]) - ArcTanh[(1 + 2^(1/3)*x)/Sqrt[1 - x^3]]/(3*2^(2/3)) + ArcTanh[Sqrt[1 - x^3]]/(9*2^(2/3))`

Defintions of rubi rules used

rule 986

```
Int[(x_)/(((a_) + (b_.)*(x_)^3)*Sqrt[(c_) + (d_.)*(x_)^3]), x_Symbol] :> With[{q = Rt[d/c, 3]}, Simp[q*(ArcTanh[Sqrt[c + d*x^3]/Rt[c, 2]]/(9*2^(2/3)*b*Rt[c, 2])), x] + (-Simp[q*(ArcTanh[Rt[c, 2]*((1 - 2^(1/3)*q*x)/Sqrt[c + d*x^3]))/(3*2^(2/3)*b*Rt[c, 2])), x] + Simp[q*(ArcTan[Sqrt[c + d*x^3]/(Sqrt[3]*Rt[c, 2])]/(3*2^(2/3)*Sqrt[3]*b*Rt[c, 2])), x] - Simp[q*(ArcTan[Sqrt[3]*Rt[c, 2]*((1 + 2^(1/3)*q*x)/Sqrt[c + d*x^3])]/(3*2^(2/3)*Sqrt[3]*b*Rt[c, 2])), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[4*b*c - a*d, 0] && PosQ[c]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 11.85 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.29

method	result
default	$i\sqrt{2} \left(\sum_{-\alpha=\text{RootOf}(_Z^3-4)} \frac{-\alpha^2\sqrt{2}\sqrt{i(-i\sqrt{3}+2x+1)}\sqrt{\frac{-1+x}{i\sqrt{3}-3}}\sqrt{-\frac{i(i\sqrt{3}+2x+1)}{2}}(-2-\alpha^2+\alpha+1+i\sqrt{3}(1-\alpha))\text{EllipticPi}\left(\frac{\sqrt{3}\sqrt{i\left(x+\frac{1}{2}\right)}}{3}\right)}{2\sqrt{-x^3+1}} \right)$
elliptic	$i\sqrt{2} \left(\sum_{-\alpha=\text{RootOf}(_Z^3-4)} \frac{-\alpha^2\sqrt{2}\sqrt{i(-i\sqrt{3}+2x+1)}\sqrt{\frac{-1+x}{i\sqrt{3}-3}}\sqrt{-\frac{i(i\sqrt{3}+2x+1)}{2}}(-2-\alpha^2+\alpha+1+i\sqrt{3}(1-\alpha))\text{EllipticPi}\left(\frac{\sqrt{3}\sqrt{i\left(x+\frac{1}{2}\right)}}{3}\right)}{2\sqrt{-x^3+1}} \right)$
trager	Expression too large to display

input

```
int(x/(-x^3+4)/(-x^3+1)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
1/36*I*2^(1/2)*sum(_alpha^2*(1/2*I*(-I*3^(1/2)+2*x+1))^(1/2)*((-1+x)/(I*3^(1/2)-3))^(1/2)*(-1/2*I*(I*3^(1/2)+2*x+1))^(1/2)/(-x^3+1)^(1/2)*(-2*_alpha^2+_alpha+1+I*3^(1/2)*(1-_alpha))*EllipticPi(1/3*3^(1/2)*(I*(x+1/2-1/2*I*3^(1/2))*3^(1/2))^(1/2),1/2*_alpha-1/3*I*_alpha^2*3^(1/2)-1/2+1/6*I*_alpha*3^(1/2)+1/6*I*3^(1/2),(I*3^(1/2)/(-3/2+1/2*I*3^(1/2)))^(1/2)),_alpha=RootOf(_Z^3-4))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1019 vs. $2(92) = 184$.

Time = 0.14 (sec) , antiderivative size = 1019, normalized size of antiderivative = 8.02

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = \text{Too large to display}$$

input

```
integrate(x/(-x^3+4)/(-x^3+1)^(1/2),x, algorithm="fricas")
```

output

```
-1/36*(-1/432)^(1/6)*(sqrt(-3) + 1)*log(-(x^9 + 66*x^6 - 72*x^3 - 24*(-1/2)^(2/3)*(x^7 + x^4 + sqrt(-3)*(x^7 + x^4 - 2*x) - 2*x) + 6*sqrt(-x^3 + 1)*(648*(-1/432)^(5/6)*(sqrt(-3)*x^5 - x^5) - sqrt(-1/3)*(5*x^6 + 20*x^3 - 16) - (-1/432)^(1/6)*(x^7 + 16*x^4 + sqrt(-3)*(x^7 + 16*x^4 - 8*x) - 8*x)) + 6*(-1/2)^(1/3)*(x^8 + 7*x^5 - 8*x^2 - sqrt(-3)*(x^8 + 7*x^5 - 8*x^2)) + 32)/(x^9 - 12*x^6 + 48*x^3 - 64)) + 1/36*(-1/432)^(1/6)*(sqrt(-3) + 1)*log(-(x^9 + 66*x^6 - 72*x^3 - 24*(-1/2)^(2/3)*(x^7 + x^4 + sqrt(-3)*(x^7 + x^4 - 2*x) - 2*x) - 6*sqrt(-x^3 + 1)*(648*(-1/432)^(5/6)*(sqrt(-3)*x^5 - x^5) - sqrt(-1/3)*(5*x^6 + 20*x^3 - 16) - (-1/432)^(1/6)*(x^7 + 16*x^4 + sqrt(-3)*(x^7 + 16*x^4 - 8*x) - 8*x)) + 6*(-1/2)^(1/3)*(x^8 + 7*x^5 - 8*x^2 - sqrt(-3)*(x^8 + 7*x^5 - 8*x^2)) + 32)/(x^9 - 12*x^6 + 48*x^3 - 64)) - 1/36*(-1/432)^(1/6)*(sqrt(-3) - 1)*log(-(x^9 + 66*x^6 - 72*x^3 - 24*(-1/2)^(2/3)*(x^7 + x^4 - sqrt(-3)*(x^7 + x^4 - 2*x) - 2*x) + 6*sqrt(-x^3 + 1)*(648*(-1/432)^(5/6)*(sqrt(-3)*x^5 + x^5) + sqrt(-1/3)*(5*x^6 + 20*x^3 - 16) + (-1/432)^(1/6)*(x^7 + 16*x^4 - sqrt(-3)*(x^7 + 16*x^4 - 8*x) - 8*x)) + 6*(-1/2)^(1/3)*(x^8 + 7*x^5 - 8*x^2 + sqrt(-3)*(x^8 + 7*x^5 - 8*x^2)) + 32)/(x^9 - 12*x^6 + 48*x^3 - 64)) + 1/36*(-1/432)^(1/6)*(sqrt(-3) - 1)*log(-(x^9 + 66*x^6 - 72*x^3 - 24*(-1/2)^(2/3)*(x^7 + x^4 - sqrt(-3)*(x^7 + x^4 - 2*x) - 2*x) - 6*sqrt(-x^3 + 1)*(648*(-1/432)^(5/6)*(sqrt(-3)*x^5 + x^5) + sqrt(-1/3)*(5*x^6 + 20*x^3 - 16) + (-1/432)^(1/6)*(x^7 + 16*x^4 - sqrt(-3)...
```

Sympy [F]

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = - \int \frac{x}{x^3\sqrt{1-x^3} - 4\sqrt{1-x^3}} dx$$

input `integrate(x/(-x**3+4)/(-x**3+1)**(1/2),x)`

output `-Integral(x/(x**3*sqrt(1 - x**3) - 4*sqrt(1 - x**3)), x)`

Maxima [F]

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = \int -\frac{x}{(x^3-4)\sqrt{-x^3+1}} dx$$

input `integrate(x/(-x^3+4)/(-x^3+1)^(1/2),x, algorithm="maxima")`

output `-integrate(x/((x^3 - 4)*sqrt(-x^3 + 1)), x)`

Giac [F]

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = \int -\frac{x}{(x^3-4)\sqrt{-x^3+1}} dx$$

input `integrate(x/(-x^3+4)/(-x^3+1)^(1/2),x, algorithm="giac")`

output `integrate(-x/((x^3 - 4)*sqrt(-x^3 + 1)), x)`

Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 653, normalized size of antiderivative = 5.14

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = \text{Too large to display}$$

input `int(-x/((1-x^3)^(1/2)*(x^3-4)),x)`

output

```
- (2^(1/3)*((3^(1/2)*1i)/2 + 3/2)*(x^3 - 1)^(1/2)*(-(x - (3^(1/2)*1i)/2 +
1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2)*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1
i)/2 + 3/2))^(1/2)*(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*ellipticPi(-(3
^(1/2)*1i)/2 + 3/2)/(2^(2/3) - 1), asin(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^(
1/2)), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/((3*(1 - x^3)^(1/2)
)*(2^(2/3) - 1)*(((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) - x*((3^(1
/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) + x^3)^(1/2)) - (2^(1/3)*((3^
(1/2)*1i)/2 + 3/2)*(x^3 - 1)^(1/2)*(-(x - (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*
1i)/2 - 3/2))^(1/2)*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1
/2)*(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*ellipticPi(((3^(1/2)*1i)/2 + 3
/2)/(2^(2/3)*((3^(1/2)*1i)/2 + 1/2) + 1), asin(-(x - 1)/((3^(1/2)*1i)/2 +
3/2))^(1/2)), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/((3*((3^(1/
2)*1i)/2 + 1/2)*(1 - x^3)^(1/2)*(2^(2/3)*((3^(1/2)*1i)/2 + 1/2) + 1)*(((3^
(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) - x*((3^(1/2)*1i)/2 - 1/2)*((3^
(1/2)*1i)/2 + 1/2) + 1) + x^3)^(1/2)) - (2^(1/3)*((3^(1/2)*1i)/2 + 3/2)*(x
^3 - 1)^(1/2)*(-(x - (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2)*
(x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*(-(x - 1)/((3^(1/
2)*1i)/2 + 3/2))^(1/2)*ellipticPi(-(3^(1/2)*1i)/2 + 3/2)/(2^(2/3)*((3^(1/
2)*1i)/2 - 1/2) - 1), asin(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)), -((3^
(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/((3*((3^(1/2)*1i)/2 - 1/2)*...
```

Reduce [F]

$$\int \frac{x}{\sqrt{1-x^3}(4-x^3)} dx = \int \frac{\sqrt{-x^3+1}x}{x^6-5x^3+4} dx$$

input `int(x/(-x^3+4)/(-x^3+1)^(1/2),x)`

output `int((sqrt(-x**3 + 1)*x)/(x**6 - 5*x**3 + 4),x)`

3.74 $\int \frac{x}{(4-dx^3)\sqrt{-1+dx^3}} dx$

Optimal result	578
Mathematica [C] (warning: unable to verify)	579
Rubi [A] (verified)	579
Maple [C] (warning: unable to verify)	580
Fricas [B] (verification not implemented)	581
Sympy [F]	582
Maxima [F]	583
Giac [F]	583
Mupad [B] (verification not implemented)	583
Reduce [F]	584

Optimal result

Integrand size = 23, antiderivative size = 157

$$\int \frac{x}{(4-dx^3)\sqrt{-1+dx^3}} dx = -\frac{\arctan\left(\frac{1+\sqrt[3]{2}\sqrt[3]{dx}}{\sqrt{-1+dx^3}}\right)}{3 \cdot 2^{2/3} d^{2/3}} - \frac{\arctan(\sqrt{-1+dx^3})}{9 \cdot 2^{2/3} d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{dx})}{\sqrt{-1+dx^3}}\right)}{3 \cdot 2^{2/3} \sqrt{3} d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{-1+dx^3}}{\sqrt{3}}\right)}{3 \cdot 2^{2/3} \sqrt{3} d^{2/3}}$$

output

```
-1/6*arctan((1+2^(1/3)*d^(1/3)*x)/(d*x^3-1)^(1/2))*2^(1/3)/d^(2/3)-1/18*arctan((d*x^3-1)^(1/2))*2^(1/3)/d^(2/3)-1/18*arctanh((1-2^(1/3)*d^(1/3)*x)*3^(1/2)/(d*x^3-1)^(1/2))*2^(1/3)/d^(2/3)*3^(1/2)-1/18*arctanh(1/3*(d*x^3-1)^(1/2)*3^(1/2))*2^(1/3)/d^(2/3)*3^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.34

$$\int \frac{x}{(4 - dx^3)\sqrt{-1 + dx^3}} dx = \frac{x^2\sqrt{1 - dx^3} \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, dx^3, \frac{dx^3}{4}\right)}{8\sqrt{-1 + dx^3}}$$

input `Integrate[x/((4 - d*x^3)*Sqrt[-1 + d*x^3]),x]`

output `(x^2*Sqrt[1 - d*x^3]*AppellF1[2/3, 1/2, 1, 5/3, d*x^3, (d*x^3)/4])/(8*Sqrt[-1 + d*x^3])`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {987}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{(4 - dx^3)\sqrt{dx^3 - 1}} dx$$

↓ 987

$$\frac{\arctan\left(\frac{\sqrt[3]{2}\sqrt[3]{dx+1}}{\sqrt{dx^3-1}}\right)}{3 \cdot 2^{2/3} d^{2/3}} - \frac{\arctan\left(\sqrt{dx^3 - 1}\right)}{9 \cdot 2^{2/3} d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}\left(1 - \sqrt[3]{2}\sqrt[3]{dx}\right)}{\sqrt{dx^3-1}}\right)}{3 \cdot 2^{2/3} \sqrt{3} d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{dx^3-1}}{\sqrt{3}}\right)}{3 \cdot 2^{2/3} \sqrt{3} d^{2/3}}$$

input `Int[x/((4 - d*x^3)*Sqrt[-1 + d*x^3]),x]`

output

```
-1/3*ArcTan[(1 + 2^(1/3)*d^(1/3)*x)/Sqrt[-1 + d*x^3]]/(2^(2/3)*d^(2/3)) -
ArcTan[Sqrt[-1 + d*x^3]]/(9*2^(2/3)*d^(2/3)) - ArcTanh[(Sqrt[3]*(1 - 2^(1/3)
3)*d^(1/3)*x)/Sqrt[-1 + d*x^3]]/(3*2^(2/3)*Sqrt[3]*d^(2/3)) - ArcTanh[Sqr
t[-1 + d*x^3]/Sqrt[3]]/(3*2^(2/3)*Sqrt[3]*d^(2/3))
```

Defintions of rubi rules used

rule 987

```
Int[(x_)/(((a_) + (b_.)*(x_)^3)*Sqrt[(c_) + (d_.)*(x_)^3]), x_Symbol] := Wi
th[{q = Rt[d/c, 3]}, Simp[(-q)*(ArcTan[Sqrt[c + d*x^3]/Rt[-c, 2]]/(9*2^(2/3)
)*b*Rt[-c, 2]), x] + (-Simp[q*(ArcTan[Rt[-c, 2]*((1 - 2^(1/3)*q*x)/Sqrt[c
+ d*x^3]])/(3*2^(2/3)*b*Rt[-c, 2]), x] - Simp[q*(ArcTanh[Sqrt[c + d*x^3]]/(
Sqrt[3]*Rt[-c, 2]))/(3*2^(2/3)*Sqrt[3]*b*Rt[-c, 2]), x] - Simp[q*(ArcTanh[
Sqrt[3]*Rt[-c, 2]*((1 + 2^(1/3)*q*x)/Sqrt[c + d*x^3]))/(3*2^(2/3)*Sqrt[3]*b
*Rt[-c, 2]), x]]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[4*
b*c - a*d, 0] && NegQ[c]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.88 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.53

method	result
default	$i\sqrt{2} \sum_{-\alpha = \text{RootOf}(dZ^3 - 4)} \frac{\sqrt{-\frac{i\left(2x + \frac{1}{d^{1/3}} + \frac{i\sqrt{3}}{d^{1/3}}\right)d^{1/3}}{2}}}{\sqrt{-\frac{3}{d^{1/3}} - \frac{i\sqrt{3}}{d^{1/3}}}} \sqrt{2} \sqrt{i\left(2x + \frac{1}{d^{1/3}} - \frac{i\sqrt{3}}{d^{1/3}}\right)d^{1/3}} \left(-2_{-\alpha}^2 d + i\sqrt{3}_{-\alpha} d^{2/3} - i\sqrt{3} d^{1/3} +_{-\alpha}\right)$
elliptic	$i\sqrt{2} \sum_{-\alpha = \text{RootOf}(dZ^3 - 4)} \frac{\sqrt{-\frac{i\left(2x + \frac{1}{d^{1/3}} + \frac{i\sqrt{3}}{d^{1/3}}\right)d^{1/3}}{2}}}{\sqrt{-\frac{3}{d^{1/3}} - \frac{i\sqrt{3}}{d^{1/3}}}} \sqrt{2} \sqrt{i\left(2x + \frac{1}{d^{1/3}} - \frac{i\sqrt{3}}{d^{1/3}}\right)d^{1/3}} \left(-2_{-\alpha}^2 d + i\sqrt{3}_{-\alpha} d^{2/3} - i\sqrt{3} d^{1/3} +_{-\alpha}\right)$

input `int(x/(-d*x^3+4)/(d*x^3-1)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/9*I*2^(1/2)*sum(1/_alpha/d^(4/3)*(-1/2*I*(2*x+1/d^(1/3)+I*3^(1/2)/d^(1/3))*d^(1/3))^(1/2)*((x-1/d^(1/3))/(-3/d^(1/3)-I*3^(1/2)/d^(1/3)))^(1/2)*(1/2*I*(2*x+1/d^(1/3)-I*3^(1/2)/d^(1/3))*d^(1/3))^(1/2)/(d*x^3-1)^(1/2)*(-2*_alpha^2*d+I*3^(1/2)*_alpha*d^(2/3)-I*3^(1/2)*d^(1/3)+_alpha*d^(2/3)+d^(1/3))*EllipticPi(1/3*3^(1/2)*(-I*(x+1/2/d^(1/3)+1/2*I*3^(1/2)/d^(1/3))*3^(1/2)*d^(1/3))^(1/2),1/3*I*3^(1/2)*d^(2/3)*_alpha^2-1/6*I*3^(1/2)*d^(1/3)*_alpha-1/6*I*3^(1/2)+1/2*d^(1/3)*_alpha-1/2,(-I*3^(1/2)/d^(1/3)/(-3/2/d^(1/3)-1/2*I*3^(1/2)/d^(1/3)))^(1/2)),_alpha=RootOf(_Z^3*d-4))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1669 vs. $2(110) = 220$.

Time = 0.21 (sec) , antiderivative size = 1669, normalized size of antiderivative = 10.63

$$\int \frac{x}{(4 - dx^3)\sqrt{-1 + dx^3}} dx = \text{Too large to display}$$

input `integrate(x/(-d*x^3+4)/(d*x^3-1)^(1/2),x, algorithm="fricas")`

output

```

-1/36*(1/432)^(1/6)*(sqrt(-3) + 1)*(d^(-4))^(1/6)*log((d^3*x^9 + 66*d^2*x^
6 - 72*d*x^3 - 24*(1/2)^(2/3)*(d^5*x^7 + d^4*x^4 - 2*d^3*x + sqrt(-3)*(d^5
*x^7 + d^4*x^4 - 2*d^3*x))*(d^(-4))^(2/3) - 6*(1/2)^(1/3)*(d^4*x^8 + 7*d^3
*x^5 - 8*d^2*x^2 - sqrt(-3)*(d^4*x^8 + 7*d^3*x^5 - 8*d^2*x^2))*(d^(-4))^(1
/3) + 6*sqrt(d*x^3 - 1)*(648*(1/432)^(5/6)*(sqrt(-3)*d^5*x^5 - d^5*x^5)*(d
^(-4))^(5/6) + sqrt(1/3)*(5*d^4*x^6 + 20*d^3*x^3 - 16*d^2)*sqrt(d^(-4)) -
(1/432)^(1/6)*(d^3*x^7 + 16*d^2*x^4 - 8*d*x + sqrt(-3)*(d^3*x^7 + 16*d^2*x
^4 - 8*d*x))*(d^(-4))^(1/6)) + 32)/(d^3*x^9 - 12*d^2*x^6 + 48*d*x^3 - 64))
+ 1/36*(1/432)^(1/6)*(sqrt(-3) + 1)*(d^(-4))^(1/6)*log((d^3*x^9 + 66*d^2*x
^6 - 72*d*x^3 - 24*(1/2)^(2/3)*(d^5*x^7 + d^4*x^4 - 2*d^3*x + sqrt(-3)*(d
^5*x^7 + d^4*x^4 - 2*d^3*x))*(d^(-4))^(2/3) - 6*(1/2)^(1/3)*(d^4*x^8 + 7*d
^3*x^5 - 8*d^2*x^2 - sqrt(-3)*(d^4*x^8 + 7*d^3*x^5 - 8*d^2*x^2))*(d^(-4))^(
1/3) - 6*sqrt(d*x^3 - 1)*(648*(1/432)^(5/6)*(sqrt(-3)*d^5*x^5 - d^5*x^5)*
(d^(-4))^(5/6) + sqrt(1/3)*(5*d^4*x^6 + 20*d^3*x^3 - 16*d^2)*sqrt(d^(-4))
- (1/432)^(1/6)*(d^3*x^7 + 16*d^2*x^4 - 8*d*x + sqrt(-3)*(d^3*x^7 + 16*d^2
*x^4 - 8*d*x))*(d^(-4))^(1/6)) + 32)/(d^3*x^9 - 12*d^2*x^6 + 48*d*x^3 - 64
)) - 1/36*(1/432)^(1/6)*(sqrt(-3) - 1)*(d^(-4))^(1/6)*log((d^3*x^9 + 66*d^
2*x^6 - 72*d*x^3 - 24*(1/2)^(2/3)*(d^5*x^7 + d^4*x^4 - 2*d^3*x - sqrt(-3)*
(d^5*x^7 + d^4*x^4 - 2*d^3*x))*(d^(-4))^(2/3) - 6*(1/2)^(1/3)*(d^4*x^8 + 7
*d^3*x^5 - 8*d^2*x^2 + sqrt(-3)*(d^4*x^8 + 7*d^3*x^5 - 8*d^2*x^2))*(d^(...

```

Sympy [F]

$$\int \frac{x}{(4 - dx^3)\sqrt{-1 + dx^3}} dx = - \int \frac{x}{dx^3\sqrt{dx^3 - 1} - 4\sqrt{dx^3 - 1}} dx$$

input

```
integrate(x/(-d*x**3+4)/(d*x**3-1)**(1/2), x)
```

output

```
-Integral(x/(d*x**3*sqrt(d*x**3 - 1) - 4*sqrt(d*x**3 - 1)), x)
```

Maxima [F]

$$\int \frac{x}{(4 - dx^3) \sqrt{-1 + dx^3}} dx = \int -\frac{x}{\sqrt{dx^3 - 1}(dx^3 - 4)} dx$$

input `integrate(x/(-d*x^3+4)/(d*x^3-1)^(1/2),x, algorithm="maxima")`

output `-integrate(x/(sqrt(d*x^3 - 1)*(d*x^3 - 4)), x)`

Giac [F]

$$\int \frac{x}{(4 - dx^3) \sqrt{-1 + dx^3}} dx = \int -\frac{x}{\sqrt{dx^3 - 1}(dx^3 - 4)} dx$$

input `integrate(x/(-d*x^3+4)/(d*x^3-1)^(1/2),x, algorithm="giac")`

output `integrate(-x/(sqrt(d*x^3 - 1)*(d*x^3 - 4)), x)`

Mupad [B] (verification not implemented)

Time = 8.40 (sec) , antiderivative size = 331, normalized size of antiderivative = 2.11

$$\begin{aligned} & \int \frac{x}{(4 - dx^3) \sqrt{-1 + dx^3}} dx \\ &= \frac{\sqrt{3} 314928^{1/3} \ln \left(\frac{(54 \sqrt{dx^3 - 1} + 54 \sqrt{3} - 54 2^{1/3} \sqrt{3} d^{1/3} x) (\sqrt{dx^3 - 1} - \sqrt{3} + 2^{1/3} \sqrt{3} d^{1/3} x)^3}{(2^{2/3} - d^{1/3} x)^6} \right)}{2916 d^{2/3}} \\ &+ \frac{\sqrt{3} 314928^{1/3} \ln \left(\frac{(2 \sqrt{dx^3 - 1} + 2 \sqrt{3} + 2^{1/3} \sqrt{3} d^{1/3} x + 2^{1/3} d^{1/3} x 3i)^3 (108 \sqrt{3} - 108 \sqrt{dx^3 - 1} + 54 2^{1/3} \sqrt{3} d^{1/3} x + 2^{1/3} d^{1/3} x 162i)}{(2^{2/3} + 2 d^{1/3} x - 2^{2/3} \sqrt{3} 1i)^6} \right)}{2916 d^{2/3}} \\ &+ \frac{\sqrt{3} 314928^{1/3} \ln \left(\frac{(2 \sqrt{dx^3 - 1} - 2 \sqrt{3} - 2^{1/3} \sqrt{3} d^{1/3} x + 2^{1/3} d^{1/3} x 3i)^3 (108 \sqrt{dx^3 - 1} + 108 \sqrt{3} + 54 2^{1/3} \sqrt{3} d^{1/3} x - 2^{1/3} d^{1/3} x 162i)}{(2^{2/3} + 2 d^{1/3} x + 2^{2/3} \sqrt{3} 1i)^6} \right)}{2916 d^{2/3}} \end{aligned}$$

input `int(-x/((d*x^3 - 1)^(1/2)*(d*x^3 - 4)),x)`

output `(3^(1/2)*314928^(1/3)*log(((54*(d*x^3 - 1)^(1/2) + 54*3^(1/2) - 54*2^(1/3)
*3^(1/2)*d^(1/3)*x)*((d*x^3 - 1)^(1/2) - 3^(1/2) + 2^(1/3)*3^(1/2)*d^(1/3)
*x)^3)/(2^(2/3) - d^(1/3)*x)^6)/(2916*d^(2/3)) + (3^(1/2)*314928^(1/3)*lo
g(((2*(d*x^3 - 1)^(1/2) + 2*3^(1/2) + 2^(1/3)*d^(1/3)*x*3i + 2^(1/3)*3^(1/
2)*d^(1/3)*x)^3*(108*3^(1/2) - 108*(d*x^3 - 1)^(1/2) + 2^(1/3)*d^(1/3)*x*1
62i + 54*2^(1/3)*3^(1/2)*d^(1/3)*x))/(2^(2/3) - 2^(2/3)*3^(1/2)*1i + 2*d^(
1/3)*x)^6)*((3^(1/2)*1i)/2 - 1/2)^(1/2))/(2916*d^(2/3)) + (3^(1/2)*314928^(
1/3)*log(((2*(d*x^3 - 1)^(1/2) - 2*3^(1/2) + 2^(1/3)*d^(1/3)*x*3i - 2^(1/
3)*3^(1/2)*d^(1/3)*x)^3*(108*(d*x^3 - 1)^(1/2) + 108*3^(1/2) - 2^(1/3)*d^(
1/3)*x*162i + 54*2^(1/3)*3^(1/2)*d^(1/3)*x))/(2^(2/3)*3^(1/2)*1i + 2^(2/3)
+ 2*d^(1/3)*x)^6)*((3^(1/2)*1i)/2 + 1/2)^(1/2)*1i)/(2916*d^(2/3))`

Reduce [F]

$$\int \frac{x}{(4 - dx^3)\sqrt{-1 + dx^3}} dx = - \left(\int \frac{\sqrt{dx^3 - 1} x}{d^2x^6 - 5dx^3 + 4} dx \right)$$

input `int(x/(-d*x^3+4)/(d*x^3-1)^(1/2),x)`

output `- int((sqrt(d*x**3 - 1)*x)/(d**2*x**6 - 5*d*x**3 + 4),x)`

3.75 $\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx$

Optimal result	585
Mathematica [C] (warning: unable to verify)	585
Rubi [A] (verified)	586
Maple [C] (verified)	588
Fricas [B] (verification not implemented)	589
Sympy [F]	590
Maxima [F]	591
Giac [F]	591
Mupad [B] (verification not implemented)	591
Reduce [F]	592

Optimal result

Integrand size = 18, antiderivative size = 74

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx = \frac{1}{18} \arctan\left(\frac{(1-x)^2}{3\sqrt{-1+x^3}}\right) + \frac{1}{18} \arctan\left(\frac{1}{3}\sqrt{-1+x^3}\right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-x)}{\sqrt{-1+x^3}}\right)}{6\sqrt{3}}$$

output

```
1/18*arctan(1/3*(1-x)^2/(x^3-1)^(1/2))+1/18*arctan(1/3*(x^3-1)^(1/2))-1/18*arctanh((1-x)*3^(1/2)/(x^3-1)^(1/2))*3^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.65

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx = \frac{x^2\sqrt{1-x^3} \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, x^3, -\frac{x^3}{8}\right)}{16\sqrt{-1+x^3}}$$

input

```
Integrate[x/(Sqrt[-1 + x^3]*(8 + x^3)), x]
```

output

```
(x^2*sqrt[1 - x^3]*AppellF1[2/3, 1/2, 1, 5/3, x^3, -1/8*x^3])/(16*sqrt[-1 + x^3])
```

Rubi [A] (verified)

Time = 0.42 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {988, 25, 946, 73, 216, 2563, 216, 2570, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
 & \int \frac{x}{\sqrt{x^3-1}(x^3+8)} dx \\
 & \quad \downarrow 988 \\
 & -\frac{1}{12} \int \frac{1-x}{(x+2)\sqrt{x^3-1}} dx - \frac{1}{12} \int -\frac{-x^2+2x+2}{(x^2-2x+4)\sqrt{x^3-1}} dx - \frac{1}{4} \int -\frac{x^2}{\sqrt{x^3-1}(x^3+8)} dx \\
 & \quad \downarrow 25 \\
 & -\frac{1}{12} \int \frac{1-x}{(x+2)\sqrt{x^3-1}} dx + \frac{1}{12} \int \frac{-x^2+2x+2}{(x^2-2x+4)\sqrt{x^3-1}} dx + \frac{1}{4} \int \frac{x^2}{\sqrt{x^3-1}(x^3+8)} dx \\
 & \quad \downarrow 946 \\
 & -\frac{1}{12} \int \frac{1-x}{(x+2)\sqrt{x^3-1}} dx + \frac{1}{12} \int \frac{1}{\sqrt{x^3-1}(x^3+8)} dx^3 + \frac{1}{12} \int \frac{-x^2+2x+2}{(x^2-2x+4)\sqrt{x^3-1}} dx \\
 & \quad \downarrow 73 \\
 & -\frac{1}{12} \int \frac{1-x}{(x+2)\sqrt{x^3-1}} dx + \frac{1}{6} \int \frac{1}{x^6+9} d\sqrt{x^3-1} + \frac{1}{12} \int \frac{-x^2+2x+2}{(x^2-2x+4)\sqrt{x^3-1}} dx \\
 & \quad \downarrow 216 \\
 & -\frac{1}{12} \int \frac{1-x}{(x+2)\sqrt{x^3-1}} dx + \frac{1}{12} \int \frac{-x^2+2x+2}{(x^2-2x+4)\sqrt{x^3-1}} dx + \frac{1}{18} \arctan\left(\frac{\sqrt{x^3-1}}{3}\right) \\
 & \quad \downarrow 2563 \\
 & \frac{1}{6} \int \frac{1}{\frac{(1-x)^4}{x^3-1}+9} d\sqrt{x^3-1} + \frac{1}{12} \int \frac{-x^2+2x+2}{(x^2-2x+4)\sqrt{x^3-1}} dx + \frac{1}{18} \arctan\left(\frac{\sqrt{x^3-1}}{3}\right)
 \end{aligned}$$

$$\begin{aligned} & \downarrow 216 \\ & \frac{1}{12} \int \frac{-x^2 + 2x + 2}{(x^2 - 2x + 4)\sqrt{x^3 - 1}} dx + \frac{1}{18} \arctan\left(\frac{(1-x)^2}{3\sqrt{x^3 - 1}}\right) + \frac{1}{18} \arctan\left(\frac{\sqrt{x^3 - 1}}{3}\right) \\ & \downarrow 2570 \\ & \frac{1}{3} \int \frac{1}{\frac{6(1-x)^2}{x^3-1} - 2} d\frac{1-x}{\sqrt{x^3-1}} + \frac{1}{18} \arctan\left(\frac{(1-x)^2}{3\sqrt{x^3-1}}\right) + \frac{1}{18} \arctan\left(\frac{\sqrt{x^3-1}}{3}\right) \\ & \downarrow 220 \\ & \frac{1}{18} \arctan\left(\frac{(1-x)^2}{3\sqrt{x^3-1}}\right) + \frac{1}{18} \arctan\left(\frac{\sqrt{x^3-1}}{3}\right) - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-x)}{\sqrt{x^3-1}}\right)}{6\sqrt{3}} \end{aligned}$$

input `Int[x/(Sqrt[-1 + x^3]*(8 + x^3)),x]`

output `ArcTan[(1 - x)^2/(3*Sqrt[-1 + x^3])]/18 + ArcTan[Sqrt[-1 + x^3]/3]/18 - ArcTanh[(Sqrt[3]*(1 - x))/Sqrt[-1 + x^3]]/(6*Sqrt[3])`

Defintions of rubi rules used

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 73 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Simp[p/b Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

rule 216 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

rule 220 $\text{Int}[(a_+) + (b_+)(x_+)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[b, 2])^{-1}) \cdot \text{ArcTanh}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

rule 946 $\text{Int}[(x_+)^{m_+} \cdot ((a_+) + (b_+)(x_+)^n)^{p_+} \cdot ((c_+) + (d_+)(x_+)^n)^{q_+}, x_Symbol] \rightarrow \text{Simp}[1/n \ \text{Subst}[\text{Int}[(a + b \cdot x)^p \cdot (c + d \cdot x)^q, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, d, m, n, p, q, x\} \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[m - n + 1, 0]$

rule 988 $\text{Int}[(x_+)/((a_+) + (b_+)(x_+)^3) \cdot \text{Sqrt}[(c_+) + (d_+)(x_+)^3], x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[d/c, 3]\}, \text{Simp}[d \cdot (q/(4 \cdot b)) \ \text{Int}[x^2/((8 \cdot c - d \cdot x^3) \cdot \text{Sqrt}[c + d \cdot x^3]), x], x] + (-\text{Simp}[q^2/(12 \cdot b) \ \text{Int}[(1 + q \cdot x)/((2 - q \cdot x) \cdot \text{Sqrt}[c + d \cdot x^3]), x], x] + \text{Simp}[1/(12 \cdot b \cdot c) \ \text{Int}[(2 \cdot c \cdot q^2 - 2 \cdot d \cdot x - d \cdot q \cdot x^2)/((4 + 2 \cdot q \cdot x + q^2 \cdot x^2) \cdot \text{Sqrt}[c + d \cdot x^3]), x], x)])] /;$ $\text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NeQ}[b \cdot c - a \cdot d, 0] \ \&\& \ \text{EqQ}[8 \cdot b \cdot c + a \cdot d, 0]$

rule 2563 $\text{Int}[(e_+) + (f_+)(x_+)/((c_+) + (d_+)(x_+) \cdot \text{Sqrt}[(a_+) + (b_+)(x_+)^3]), x_Symbol] \rightarrow \text{Simp}[-2 \cdot (e/d) \ \text{Subst}[\text{Int}[1/(9 - a \cdot x^2), x], x, (1 + f \cdot (x/e))^2/\text{Sqrt}[a + b \cdot x^3]], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{NeQ}[d \cdot e - c \cdot f, 0] \ \&\& \ \text{EqQ}[b \cdot c^3 + 8 \cdot a \cdot d^3, 0] \ \&\& \ \text{EqQ}[2 \cdot d \cdot e + c \cdot f, 0]$

rule 2570 $\text{Int}[(f_+) + (g_+)(x_+) + (h_+)(x_+)^2)/((c_+) + (d_+)(x_+) + (e_+)(x_+)^2) \cdot \text{Sqrt}[(a_+) + (b_+)(x_+)^3], x_Symbol] \rightarrow \text{Simp}[-2 \cdot g \cdot h \ \text{Subst}[\text{Int}[1/(2 \cdot e \cdot h - (b \cdot d \cdot f - 2 \cdot a \cdot e \cdot h) \cdot x^2), x], x, (1 + 2 \cdot h \cdot (x/g))/\text{Sqrt}[a + b \cdot x^3]], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f, g, h, x\} \ \&\& \ \text{NeQ}[b \cdot d \cdot f - 2 \cdot a \cdot e \cdot h, 0] \ \&\& \ \text{EqQ}[b \cdot g^3 - 8 \cdot a \cdot h^3, 0] \ \&\& \ \text{EqQ}[g^2 + 2 \cdot f \cdot h, 0] \ \&\& \ \text{EqQ}[b \cdot d \cdot f + b \cdot c \cdot g - 4 \cdot a \cdot e \cdot h, 0]$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 8.43 (sec) , antiderivative size = 421, normalized size of antiderivative = 5.69

method	result
default	$\frac{\left(-\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}-\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\operatorname{EllipticPi}\left(\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\frac{i\sqrt{3}}{6}+\frac{1}{2},\sqrt{\frac{\frac{3}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)+\frac{i\left(\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)\left(-\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}}{9\sqrt{x^3-1}}$
trager	$\operatorname{RootOf}\left(_Z^2+1\right)\ln\left(-\frac{-9\operatorname{RootOf}\left(-18\operatorname{RootOf}\left(_Z^2+1\right)_Z+324\right)_Z^2-1}{\operatorname{RootOf}\left(_Z^2+1\right)x^2-162\operatorname{RootOf}\left(-18\operatorname{RootOf}\left(_Z^2+1\right)_Z+324\right)_Z^2-1}\right)$
elliptic	Expression too large to display

input

```
int(x/(x^3+8)/(x^3-1)^(1/2),x,method=_RETURNVERBOSE)
```

output

```
-1/9*(-3/2-1/2*I*3^(1/2))*((-1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2-1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2+1/2*I*3^(1/2))/(3/2+1/2*I*3^(1/2)))^(1/2)/(x^3-1)^(1/2)*EllipticPi(((1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2),1/6*I*3^(1/2)+1/2,((3/2+1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2))+1/9*I*(1/2-1/2*I*3^(1/2))*(-3/2-1/2*I*3^(1/2))*((-1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2-1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2+1/2*I*3^(1/2))/(3/2+1/2*I*3^(1/2)))^(1/2)/(x^3-1)^(1/2)*3^(1/2)*EllipticPi(((1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2),1/6*I*(1+I*3^(1/2))*3^(1/2)+1/3*I*3^(1/2),((3/2+1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2))-1/9*I*(1/2+1/2*I*3^(1/2))*(-3/2-1/2*I*3^(1/2))*((-1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2-1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2+1/2*I*3^(1/2))/(3/2+1/2*I*3^(1/2)))^(1/2)/(x^3-1)^(1/2)*3^(1/2)*EllipticPi(((1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2),1/6*I*(1-I*3^(1/2))*3^(1/2)-2/3*I*3^(1/2),((3/2+1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2)))^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 261 vs. $2(51) = 102$.

Time = 0.13 (sec) , antiderivative size = 261, normalized size of antiderivative = 3.53

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx$$

$$= \frac{1}{216} \sqrt{3} \log \left(\frac{x^6 + 48x^5 + 186x^4 - 56x^3 + 6\sqrt{3}(x^4 + 12x^3 + 12x^2 - 16x)\sqrt{x^3-1} - 120x^2 - 96x + 64}{x^6 - 6x^5 + 24x^4 - 56x^3 + 96x^2 - 96x + 64} \right)$$

$$- \frac{1}{216} \sqrt{3} \log \left(\frac{x^6 + 48x^5 + 186x^4 - 56x^3 - 6\sqrt{3}(x^4 + 12x^3 + 12x^2 - 16x)\sqrt{x^3-1} - 120x^2 - 96x + 64}{x^6 - 6x^5 + 24x^4 - 56x^3 + 96x^2 - 96x + 64} \right)$$

$$+ \frac{1}{54} \arctan \left(\frac{(x^3 - 12x^2 - 6x - 10)\sqrt{x^3-1}}{6(x^4 - x^3 - x + 1)} \right)$$

$$+ \frac{1}{54} \arctan \left(\frac{\sqrt{x^3-1}(x^2 - 8x + 10)}{3(x^3 - 3x^2 + 2)} \right)$$

input `integrate(x/(x^3+8)/(x^3-1)^(1/2),x, algorithm="fricas")`

output `1/216*sqrt(3)*log((x^6 + 48*x^5 + 186*x^4 - 56*x^3 + 6*sqrt(3)*(x^4 + 12*x^3 + 12*x^2 - 16*x)*sqrt(x^3 - 1) - 120*x^2 - 96*x + 64)/(x^6 - 6*x^5 + 24*x^4 - 56*x^3 + 96*x^2 - 96*x + 64)) - 1/216*sqrt(3)*log((x^6 + 48*x^5 + 186*x^4 - 56*x^3 - 6*sqrt(3)*(x^4 + 12*x^3 + 12*x^2 - 16*x)*sqrt(x^3 - 1) - 120*x^2 - 96*x + 64)/(x^6 - 6*x^5 + 24*x^4 - 56*x^3 + 96*x^2 - 96*x + 64)) + 1/54*arctan(1/6*(x^3 - 12*x^2 - 6*x - 10)*sqrt(x^3 - 1)/(x^4 - x^3 - x + 1)) + 1/54*arctan(1/3*sqrt(x^3 - 1)*(x^2 - 8*x + 10)/(x^3 - 3*x^2 + 2))`

Sympy [F]

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx = \int \frac{x}{\sqrt{(x-1)(x^2+x+1)}(x+2)(x^2-2x+4)} dx$$

input `integrate(x/(x**3+8)/(x**3-1)**(1/2),x)`

output `Integral(x/(sqrt((x - 1)*(x**2 + x + 1))*(x + 2)*(x**2 - 2*x + 4)), x)`

Maxima [F]

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx = \int \frac{x}{(x^3+8)\sqrt{x^3-1}} dx$$

input `integrate(x/(x^3+8)/(x^3-1)^(1/2),x, algorithm="maxima")`

output `integrate(x/((x^3 + 8)*sqrt(x^3 - 1)), x)`

Giac [F]

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx = \int \frac{x}{(x^3+8)\sqrt{x^3-1}} dx$$

input `integrate(x/(x^3+8)/(x^3-1)^(1/2),x, algorithm="giac")`

output `integrate(x/((x^3 + 8)*sqrt(x^3 - 1)), x)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 533, normalized size of antiderivative = 7.20

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx = \text{Too large to display}$$

input `int(x/((x^3 - 1)^(1/2)*(x^3 + 8)),x)`

output

```

(((3^(1/2)*1i)/2 + 3/2)*(-(x - (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 - 3/2
))^1/2*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 + 3/2))^1/2*(-(x -
1)/((3^(1/2)*1i)/2 + 3/2))^1/2*ellipticPi((3^(1/2)*1i)/6 + 1/2, asin((-
x - 1)/((3^(1/2)*1i)/2 + 3/2))^1/2), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1
i)/2 - 3/2))/9*(((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) - x*(((3^(
1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) + x^3)^1/2) - (3^(1/2)*((3
^(1/2)*1i)/2 + 3/2)*(-(x - (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 - 3/2))^
1/2*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/2)*1i)/2 + 3/2))^1/2*(-(x - 1)/
(3^(1/2)*1i)/2 + 3/2))^1/2*ellipticPi(-(3^(1/2)*((3^(1/2)*1i)/2 + 3/2)*1
i)/3, asin((-x - 1)/((3^(1/2)*1i)/2 + 3/2))^1/2), -((3^(1/2)*1i)/2 + 3/
2)/((3^(1/2)*1i)/2 - 3/2))*2i)/9*(3^(1/2)*1i - 1)*(((3^(1/2)*1i)/2 - 1/2)
*(((3^(1/2)*1i)/2 + 1/2) - x*(((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2)
+ 1) + x^3)^1/2) - (3^(1/2)*((3^(1/2)*1i)/2 + 3/2)*(-(x - (3^(1/2)*1i)/
2 + 1/2)/((3^(1/2)*1i)/2 - 3/2))^1/2*((x + (3^(1/2)*1i)/2 + 1/2)/((3^(1/
2)*1i)/2 + 3/2))^1/2*(-(x - 1)/((3^(1/2)*1i)/2 + 3/2))^1/2*ellipticPi(
(3^(1/2)*((3^(1/2)*1i)/2 + 3/2)*1i)/3, asin((-x - 1)/((3^(1/2)*1i)/2 + 3/
2))^1/2), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))*2i)/9*(3^(1/2)
)*1i + 1)*(((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) - x*(((3^(1/2)*1i
)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) + x^3)^1/2))

```

Reduce [F]

$$\int \frac{x}{\sqrt{-1+x^3}(8+x^3)} dx = \int \frac{\sqrt{x^3-1}x}{x^6+7x^3-8} dx$$

input

```
int(x/(x^3+8)/(x^3-1)^(1/2),x)
```

output

```
int((sqrt(x**3 - 1)*x)/(x**6 + 7*x**3 - 8),x)
```

3.76 $\int \frac{x}{(8-dx^3)\sqrt{1+dx^3}} dx$

Optimal result	593
Mathematica [C] (verified)	594
Rubi [A] (verified)	594
Maple [C] (warning: unable to verify)	597
Fricas [B] (verification not implemented)	599
Sympy [F]	600
Maxima [F]	600
Giac [F]	600
Mupad [F(-1)]	601
Reduce [F]	601

Optimal result

Integrand size = 23, antiderivative size = 103

$$\int \frac{x}{(8-dx^3)\sqrt{1+dx^3}} dx = -\frac{\arctan\left(\frac{\sqrt{3}(1+\sqrt[3]{dx})}{\sqrt{1+dx^3}}\right)}{6\sqrt{3}d^{2/3}} + \frac{\operatorname{arctanh}\left(\frac{(1+\sqrt[3]{dx})^2}{3\sqrt{1+dx^3}}\right)}{18d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{1}{3}\sqrt{1+dx^3}\right)}{18d^{2/3}}$$

output

```
1/18*arctanh(1/3*(1+d^(1/3)*x)^2/(d*x^3+1)^(1/2))/d^(2/3)-1/18*arctanh(1/3
*(d*x^3+1)^(1/2))/d^(2/3)-1/18*arctan((1+d^(1/3)*x)*3^(1/2)/(d*x^3+1)^(1/2
))/d^(2/3)*3^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.31

$$\int \frac{x}{(8 - dx^3)\sqrt{1 + dx^3}} dx = \frac{1}{16}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, -dx^3, \frac{dx^3}{8}\right)$$

input `Integrate[x/((8 - d*x^3)*Sqrt[1 + d*x^3]),x]`

output `(x^2*AppellF1[2/3, 1/2, 1, 5/3, -(d*x^3), (d*x^3)/8])/16`

Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$, Rules used = {988, 946, 73, 219, 2563, 219, 2570, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{x}{(8 - dx^3)\sqrt{dx^3 + 1}} dx \\ & \quad \downarrow \text{988} \\ & -\frac{\int \frac{-d^{4/3}x^2 - 2dx + 2d^{2/3}}{(d^{2/3}x^2 + 2\sqrt[3]{d}x + 4)\sqrt{dx^3 + 1}} dx}{12d} + \frac{\int \frac{\sqrt[3]{d}x + 1}{(2 - \sqrt[3]{d}x)\sqrt{dx^3 + 1}} dx}{12\sqrt[3]{d}} - \frac{1}{4}\sqrt[3]{d} \int \frac{x^2}{(8 - dx^3)\sqrt{dx^3 + 1}} dx \\ & \quad \downarrow \text{946} \\ & -\frac{\int \frac{-d^{4/3}x^2 - 2dx + 2d^{2/3}}{(d^{2/3}x^2 + 2\sqrt[3]{d}x + 4)\sqrt{dx^3 + 1}} dx}{12d} + \frac{\int \frac{\sqrt[3]{d}x + 1}{(2 - \sqrt[3]{d}x)\sqrt{dx^3 + 1}} dx}{12\sqrt[3]{d}} - \frac{1}{12}\sqrt[3]{d} \int \frac{1}{(8 - dx^3)\sqrt{dx^3 + 1}} dx^3 \\ & \quad \downarrow \text{73} \end{aligned}$$

$$\begin{aligned}
& -\frac{\int \frac{1}{9-x^6} d\sqrt{dx^3+1}}{6d^{2/3}} - \frac{\int \frac{-d^{4/3}x^2-2dx+2d^{2/3}}{(d^{2/3}x^2+2\sqrt[3]{d}x+4)\sqrt{dx^3+1}} dx}{12d} + \frac{\int \frac{\sqrt[3]{d}x+1}{(2-\sqrt[3]{d}x)\sqrt{dx^3+1}} dx}{12\sqrt[3]{d}} \\
& \quad \downarrow \text{219} \\
& -\frac{\int \frac{-d^{4/3}x^2-2dx+2d^{2/3}}{(d^{2/3}x^2+2\sqrt[3]{d}x+4)\sqrt{dx^3+1}} dx}{12d} + \frac{\int \frac{\sqrt[3]{d}x+1}{(2-\sqrt[3]{d}x)\sqrt{dx^3+1}} dx}{12\sqrt[3]{d}} - \frac{\operatorname{arctanh}\left(\frac{1}{3}\sqrt{dx^3+1}\right)}{18d^{2/3}} \\
& \quad \downarrow \text{2563} \\
& \frac{\int \frac{1}{9-\frac{dx^3+1}{d}} d \frac{(\sqrt[3]{d}x+1)^2}{\sqrt{dx^3+1}}}{6d^{2/3}} - \frac{\int \frac{-d^{4/3}x^2-2dx+2d^{2/3}}{(d^{2/3}x^2+2\sqrt[3]{d}x+4)\sqrt{dx^3+1}} dx}{12d} - \frac{\operatorname{arctanh}\left(\frac{1}{3}\sqrt{dx^3+1}\right)}{18d^{2/3}} \\
& \quad \downarrow \text{219} \\
& -\frac{\int \frac{-d^{4/3}x^2-2dx+2d^{2/3}}{(d^{2/3}x^2+2\sqrt[3]{d}x+4)\sqrt{dx^3+1}} dx}{12d} + \frac{\operatorname{arctanh}\left(\frac{(\sqrt[3]{d}x+1)^2}{3\sqrt{dx^3+1}}\right)}{18d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{1}{3}\sqrt{dx^3+1}\right)}{18d^{2/3}} \\
& \quad \downarrow \text{2570} \\
& \frac{1}{3}d^{4/3} \int \frac{1}{6(\sqrt[3]{d}x+1)^2 d^2} d \frac{\sqrt[3]{d}x+1}{\sqrt{dx^3+1}} + \frac{\operatorname{arctanh}\left(\frac{(\sqrt[3]{d}x+1)^2}{3\sqrt{dx^3+1}}\right)}{18d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{1}{3}\sqrt{dx^3+1}\right)}{18d^{2/3}} \\
& \quad \downarrow \text{218} \\
& -\frac{\operatorname{arctan}\left(\frac{\sqrt{3}(\sqrt[3]{d}x+1)}{\sqrt{dx^3+1}}\right)}{6\sqrt{3}d^{2/3}} + \frac{\operatorname{arctanh}\left(\frac{(\sqrt[3]{d}x+1)^2}{3\sqrt{dx^3+1}}\right)}{18d^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{1}{3}\sqrt{dx^3+1}\right)}{18d^{2/3}}
\end{aligned}$$

input `Int[x/((8 - d*x^3)*Sqrt[1 + d*x^3]),x]`

output

$$-1/6*\text{ArcTan}[\sqrt[3]{1+d^{1/3}x}]/\sqrt{1+dx^3}/(\sqrt[3]{d^{2/3}}) + \text{ArcTanh}[(1+d^{1/3}x)^2/(3\sqrt{1+dx^3})]/(18d^{2/3}) - \text{ArcTanh}[\sqrt{1+dx^3}/3]/(18d^{2/3})$$
Defintions of rubi rules used

rule 73

$$\text{Int}[(a_.) + (b_.)*(x_)^{(m_)}*((c_.) + (d_.)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Simp}[p/b \text{ Subst}[\text{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}, x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$$

rule 218

$$\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b]$$

rule 219

$$\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{LtQ}[b, 0])$$

rule 946

$$\text{Int}[(x_)^{(m_)}*((a_.) + (b_.)*(x_)^{(n_)})^{(p_)}*((c_.) + (d_.)*(x_)^{(n_)})^{(q_)}], x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; \text{FreeQ}[\{a, b, c, d, m, n, p, q\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[m - n + 1, 0]$$

rule 988

$$\text{Int}[(x_)/(((a_.) + (b_.)*(x_)^3)*\sqrt{(c_.) + (d_.)*(x_)^3}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[d/c, 3]\}, \text{Simp}[d*(q/(4*b)) \text{ Int}[x^2/((8*c - d*x^3)*\sqrt{c + d*x^3}), x], x] + (-\text{Simp}[q^2/(12*b) \text{ Int}[(1 + q*x)/((2 - q*x)*\sqrt{c + d*x^3}), x], x] + \text{Simp}[1/(12*b*c) \text{ Int}[(2*c*q^2 - 2*d*x - d*q*x^2)/((4 + 2*q*x + q^2*x^2)*\sqrt{c + d*x^3}), x], x)]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[8*b*c + a*d, 0]$$

rule 2563

```
Int[((e_) + (f_)*(x_))/(((c_) + (d_)*(x_))*Sqrt[(a_) + (b_)*(x_)^3]), x_
Symbol] := Simp[-2*(e/d) Subst[Int[1/(9 - a*x^2), x], x, (1 + f*(x/e))^2/
Sqrt[a + b*x^3]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[d*e - c*f, 0] &
& EqQ[b*c^3 + 8*a*d^3, 0] && EqQ[2*d*e + c*f, 0]
```

rule 2570

```
Int[((f_) + (g_)*(x_) + (h_)*(x_)^2)/(((c_) + (d_)*(x_) + (e_)*(x_)^2)*
Sqrt[(a_) + (b_)*(x_)^3]), x_Symbol] := Simp[-2*g*h Subst[Int[1/(2*e*h -
(b*d*f - 2*a*e*h)*x^2), x], x, (1 + 2*h*(x/g))/Sqrt[a + b*x^3]], x] /; Fre
eQ[{a, b, c, d, e, f, g, h}, x] && NeQ[b*d*f - 2*a*e*h, 0] && EqQ[b*g^3 - 8
*a*h^3, 0] && EqQ[g^2 + 2*f*h, 0] && EqQ[b*d*f + b*c*g - 4*a*e*h, 0]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.68 (sec) , antiderivative size = 383, normalized size of antiderivative = 3.72

method	result
default	$i\sqrt{2} \sum_{-\alpha=\text{RootOf}(dZ^3-8)} \frac{(-d^2)^{\frac{1}{3}}\sqrt{2} \sqrt{\frac{id\left(2x+\frac{-i\sqrt{3}(-d^2)^{\frac{1}{3}}+(-d^2)^{\frac{1}{3}}\right)}{(-d^2)^{\frac{1}{3}}}}{\sqrt{-3(-d^2)^{\frac{1}{3}}+i\sqrt{3}(-d^2)^{\frac{1}{3}}}} \sqrt{\frac{d\left(x-\frac{(-d^2)^{\frac{1}{3}}}{d}\right)}{2(-d^2)^{\frac{1}{3}}}}}{2(-d^2)^{\frac{1}{3}}}}$
elliptic	$i\sqrt{2} \sum_{-\alpha=\text{RootOf}(dZ^3-8)} \frac{(-d^2)^{\frac{1}{3}}\sqrt{2} \sqrt{\frac{id\left(2x+\frac{-i\sqrt{3}(-d^2)^{\frac{1}{3}}+(-d^2)^{\frac{1}{3}}\right)}{(-d^2)^{\frac{1}{3}}}}{\sqrt{-3(-d^2)^{\frac{1}{3}}+i\sqrt{3}(-d^2)^{\frac{1}{3}}}} \sqrt{\frac{d\left(x-\frac{(-d^2)^{\frac{1}{3}}}{d}\right)}{2(-d^2)^{\frac{1}{3}}}}}{2(-d^2)^{\frac{1}{3}}}}$

input `int(x/(-d*x^3+8)/(d*x^3+1)^(1/2),x,method=_RETURNVERBOSE)`

output

```
-1/27*I/d^3*2^(1/2)*sum(1/_alpha*(-d^2)^(1/3)*(1/2*I*d*(2*x+1/d*(-I*3^(1/2)
)*(-d^2)^(1/3)+(-d^2)^(1/3)))/(-d^2)^(1/3))^(1/2)*(d*(x-1/d*(-d^2)^(1/3))/
(-3*(-d^2)^(1/3)+I*3^(1/2)*(-d^2)^(1/3)))^(1/2)*(-1/2*I*d*(2*x+1/d*(I*3^(1
/2)*(-d^2)^(1/3)+(-d^2)^(1/3)))/(-d^2)^(1/3))^(1/2)/(d*x^3+1)^(1/2)*(I*(-d
^2)^(1/3)*_alpha*3^(1/2)*d-I*3^(1/2)*(-d^2)^(2/3)+2*_alpha^2*d^2-(-d^2)^(1
/3)*_alpha*d-(-d^2)^(2/3))*EllipticPi(1/3*3^(1/2)*(I*(x+1/2*d*(-d^2)^(1/3)
-1/2*I*3^(1/2)/d*(-d^2)^(1/3))*3^(1/2)*d/(-d^2)^(1/3))^(1/2),-1/18/d*(2*I*
(-d^2)^(1/3)*_alpha^2*3^(1/2)*d-I*(-d^2)^(2/3)*_alpha*3^(1/2)+I*3^(1/2)*d-
3*(-d^2)^(2/3)*_alpha-3*d), (I*3^(1/2)/d*(-d^2)^(1/3)/(-3/2/d*(-d^2)^(1/3)+
1/2*I*3^(1/2)/d*(-d^2)^(1/3)))^(1/2)), _alpha=RootOf(_Z^3*d-8))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 491 vs. $2(73) = 146$.

Time = 0.24 (sec) , antiderivative size = 491, normalized size of antiderivative = 4.77

$$\int \frac{x}{(8 - dx^3)\sqrt{1 + dx^3}} dx$$

$$= \frac{6 \sqrt{\frac{1}{3}} (d^2)^{\frac{1}{6}} d \arctan \left(-\frac{\sqrt{\frac{1}{3}} \left(9 d^3 x^5 - (d^2 x^6 - 40 dx^3 - 32) (d^2)^{\frac{2}{3}} + 3 (5 d^2 x^4 + 8 dx) (d^2)^{\frac{1}{3}} \right) \sqrt{dx^3 + 1} (d^2)^{\frac{1}{6}}}{3 (d^4 x^7 - 7 d^3 x^4 - 8 d^2 x)} \right)}{1} + 2 (d^2)^{\frac{2}{3}} \log \left(\frac{d^4 x^9}{\dots} \right)$$

input

```
integrate(x/(-d*x^3+8)/(d*x^3+1)^(1/2),x, algorithm="fricas")
```

output

```
1/108*(6*sqrt(1/3)*(d^2)^(1/6)*d*arctan(-1/3*sqrt(1/3)*(9*d^3*x^5 - (d^2*x
^6 - 40*d*x^3 - 32)*(d^2)^(2/3) + 3*(5*d^2*x^4 + 8*d*x)*(d^2)^(1/3))*sqrt(
d*x^3 + 1)*(d^2)^(1/6)/(d^4*x^7 - 7*d^3*x^4 - 8*d^2*x)) + 2*(d^2)^(2/3)*lo
g((d^4*x^9 + 318*d^3*x^6 + 1200*d^2*x^3 + 18*(5*d^2*x^7 + 64*d*x^4 + 32*x)
*(d^2)^(2/3) + 6*(7*d^3*x^6 + 152*d^2*x^3 + (d^2*x^7 + 80*d*x^4 + 160*x)*(
d^2)^(2/3) + 6*(5*d^2*x^5 + 32*d*x^2)*(d^2)^(1/3) + 64*d)*sqrt(d*x^3 + 1)
+ 18*(d^3*x^8 + 38*d^2*x^5 + 64*d*x^2)*(d^2)^(1/3) + 640*d)/(d^3*x^9 - 24*
d^2*x^6 + 192*d*x^3 - 512)) - (d^2)^(2/3)*log((d^4*x^9 - 276*d^3*x^6 - 160
8*d^2*x^3 - 18*(d^2*x^7 - 52*d*x^4 - 80*x)*(d^2)^(2/3) - 6*(4*d^3*x^6 + 16
4*d^2*x^3 + (d^2*x^7 - 28*d*x^4 - 272*x)*(d^2)^(2/3) - 24*(d^2*x^5 + d*x^2
)*(d^2)^(1/3) + 160*d)*sqrt(d*x^3 + 1) + 18*(d^3*x^8 + 20*d^2*x^5 - 8*d*x^
2)*(d^2)^(1/3) - 1088*d)/(d^3*x^9 - 24*d^2*x^6 + 192*d*x^3 - 512)))/d^2
```


Sympy [F]

$$\int \frac{x}{(8 - dx^3)\sqrt{1 + dx^3}} dx = - \int \frac{x}{dx^3\sqrt{dx^3 + 1} - 8\sqrt{dx^3 + 1}} dx$$

input `integrate(x/(-d*x**3+8)/(d*x**3+1)**(1/2),x)`

output `-Integral(x/(d*x**3*sqrt(d*x**3 + 1) - 8*sqrt(d*x**3 + 1)), x)`

Maxima [F]

$$\int \frac{x}{(8 - dx^3)\sqrt{1 + dx^3}} dx = \int -\frac{x}{\sqrt{dx^3 + 1}(dx^3 - 8)} dx$$

input `integrate(x/(-d*x^3+8)/(d*x^3+1)^(1/2),x, algorithm="maxima")`

output `-integrate(x/(sqrt(d*x^3 + 1)*(d*x^3 - 8)), x)`

Giac [F]

$$\int \frac{x}{(8 - dx^3)\sqrt{1 + dx^3}} dx = \int -\frac{x}{\sqrt{dx^3 + 1}(dx^3 - 8)} dx$$

input `integrate(x/(-d*x^3+8)/(d*x^3+1)^(1/2),x, algorithm="giac")`

output `integrate(-x/(sqrt(d*x^3 + 1)*(d*x^3 - 8)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{(8 - dx^3) \sqrt{1 + dx^3}} dx = - \int \frac{x}{\sqrt{dx^3 + 1} (dx^3 - 8)} dx$$

input `int(-x/((d*x^3 + 1)^(1/2)*(d*x^3 - 8)),x)`output `-int(x/((d*x^3 + 1)^(1/2)*(d*x^3 - 8)), x)`**Reduce [F]**

$$\int \frac{x}{(8 - dx^3) \sqrt{1 + dx^3}} dx = - \left(\int \frac{\sqrt{dx^3 + 1} x}{d^2x^6 - 7dx^3 - 8} dx \right)$$

input `int(x/(-d*x^3+8)/(d*x^3+1)^(1/2),x)`output `- int((sqrt(d*x**3 + 1)*x)/(d**2*x**6 - 7*d*x**3 - 8),x)`

$$3.77 \quad \int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx$$

Optimal result	602
Mathematica [C] (warning: unable to verify)	602
Rubi [A] (verified)	603
Maple [C] (warning: unable to verify)	604
Fricas [B] (verification not implemented)	605
Sympy [F]	606
Maxima [F]	606
Giac [F]	606
Mupad [F(-1)]	607
Reduce [F]	607

Optimal result

Integrand size = 21, antiderivative size = 81

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx = \frac{1}{4} \arctan\left(\frac{1-\sqrt[3]{1-3x^2}}{x}\right) + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{3}}\right)}{4\sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{(1-\sqrt[3]{1-3x^2})^2}{3\sqrt{3}x}\right)}{4\sqrt{3}}$$

output

```
1/4*arctan((1-(-3*x^2+1)^(1/3))/x)+1/12*arctanh(1/3*x*3^(1/2))*3^(1/2)-1/12*arctanh(1/9*(1-(-3*x^2+1)^(1/3))^2/x*3^(1/2))*3^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 4.16 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.56

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx =$$

$$\frac{9x \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, 3x^2, \frac{x^2}{3}\right)}{\sqrt[3]{1-3x^2}(-3+x^2)} + \frac{9 \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, 3x^2, \frac{x^2}{3}\right)}{(-3+x^2)} + \frac{2x^2 \operatorname{AppellF1}\left(\frac{3}{2}, \frac{1}{3}, 2, \frac{5}{2}, 3x^2, \frac{x^2}{3}\right)}{(-3+x^2)} + 3 \operatorname{AppellF1}\left(\frac{3}{2}, \frac{1}{3}, 2, \frac{5}{2}, 3x^2, \frac{x^2}{3}\right)$$

input `Integrate[1/((1 - 3*x^2)^(1/3)*(3 - x^2)),x]`

output `(-9*x*AppellF1[1/2, 1/3, 1, 3/2, 3*x^2, x^2/3])/((1 - 3*x^2)^(1/3)*(-3 + x^2)*(9*AppellF1[1/2, 1/3, 1, 3/2, 3*x^2, x^2/3] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5/2, 3*x^2, x^2/3] + 3*AppellF1[3/2, 4/3, 1, 5/2, 3*x^2, x^2/3])))`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {307}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx$$

↓ 307

$$\frac{1}{4} \arctan\left(\frac{1 - \sqrt[3]{1-3x^2}}{x}\right) - \frac{\operatorname{arctanh}\left(\frac{(1 - \sqrt[3]{1-3x^2})^2}{3\sqrt[3]{3}x}\right)}{4\sqrt{3}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{3}}\right)}{4\sqrt{3}}$$

input `Int[1/((1 - 3*x^2)^(1/3)*(3 - x^2)),x]`

output `ArcTan[(1 - (1 - 3*x^2)^(1/3))/x]/4 + ArcTanh[x/Sqrt[3]]/(4*Sqrt[3]) - ArcTanh[(1 - (1 - 3*x^2)^(1/3))^2/(3*Sqrt[3]*x)]/(4*Sqrt[3])`

Definitions of rubi rules used

rule 307

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> Wit
h[{q = Rt[-b/a, 2]}, Simp[(-q)*(ArcTanh[q*(x/3)]/(12*Rt[a, 3]*d)), x] + (Si
mp[q*(ArcTanh[(Rt[a, 3] - (a + b*x^2)^(1/3))^2/(3*Rt[a, 3]^2*q*x)]/(12*Rt[a
, 3]*d)), x] - Simp[q*(ArcTan[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3)))/(Rt[
a, 3]*q*x)]/(4*Sqrt[3]*Rt[a, 3]*d)), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[
b*c - a*d, 0] && EqQ[b*c - 9*a*d, 0] && NegQ[b/a]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 8.73 (sec) , antiderivative size = 438, normalized size of antiderivative = 5.41

method	result
trager	$\frac{\text{RootOf}(_Z^2-3) \ln\left(\frac{-96(-3x^2+1)^{\frac{1}{3}} \text{RootOf}(-4_Z \text{RootOf}(_Z^2-3)+48_Z^2+1)^2 \text{RootOf}(_Z^2-3)_{x+4}(-3x^2+1)^{\frac{1}{3}} \text{RootOf}(_Z^2-3)}{\dots}\right)}{\dots}$

input

```
int(1/(-3*x^2+1)^(1/3)/(-x^2+3),x,method=_RETURNVERBOSE)
```

output

```
1/12*RootOf(_Z^2-3)*ln((-96*(-3*x^2+1)^(1/3)*RootOf(-4*_Z*RootOf(_Z^2-3)+4
8*_Z^2+1)^2*RootOf(_Z^2-3)*x+4*(-3*x^2+1)^(1/3)*RootOf(-4*_Z*RootOf(_Z^2-3
)+48*_Z^2+1)*RootOf(_Z^2-3)^2*x+192*RootOf(-4*_Z*RootOf(_Z^2-3)+48*_Z^2+1)
^2*RootOf(_Z^2-3)*x-8*RootOf(-4*_Z*RootOf(_Z^2-3)+48*_Z^2+1)*RootOf(_Z^2-3
)^2*x+6*RootOf(-4*_Z*RootOf(_Z^2-3)+48*_Z^2+1)*RootOf(_Z^2-3)*x^2+12*(-3*x
^2+1)^(1/3)*RootOf(-4*_Z*RootOf(_Z^2-3)+48*_Z^2+1)*RootOf(_Z^2-3)+24*(-3*x
^2+1)^(1/3)*RootOf(-4*_Z*RootOf(_Z^2-3)+48*_Z^2+1)*x-(-3*x^2+1)^(1/3)*Root
Of(_Z^2-3)*x+3*(-3*x^2+1)^(2/3)+6*RootOf(-4*_Z*RootOf(_Z^2-3)+48*_Z^2+1)*R
ootOf(_Z^2-3)-3*(-3*x^2+1)^(1/3))/(x^2-3))+RootOf(-4*_Z*RootOf(_Z^2-3)+48*
_Z^2+1)*ln((-48*(-3*x^2+1)^(1/3)*RootOf(-4*_Z*RootOf(_Z^2-3)+48*_Z^2+1)*x+
2*(-3*x^2+1)^(1/3)*RootOf(_Z^2-3)*x+6*(-3*x^2+1)^(2/3)-96*RootOf(-4*_Z*Ro
otOf(_Z^2-3)+48*_Z^2+1)*x+4*RootOf(_Z^2-3)*x-3*x^2+6*(-3*x^2+1)^(1/3)-3)/(x
^2-3))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 956 vs. $2(59) = 118$.

Time = 1.10 (sec) , antiderivative size = 956, normalized size of antiderivative = 11.80

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx = \text{Too large to display}$$

input `integrate(1/(-3*x^2+1)^(1/3)/(-x^2+3),x, algorithm="fricas")`

output

```
1/72*sqrt(3)*log(-(x^12 + 2598*x^10 + 55143*x^8 + 114228*x^6 - 22113*x^4 -
7290*x^2 + 8*(3*x^10 + 576*x^8 + 5598*x^6 + 5832*x^4 - 729*x^2 - sqrt(3)*
(41*x^9 + 1368*x^7 + 4482*x^5 + 864*x^3 - 243*x))*(-3*x^2 + 1)^(2/3) - 4*sqrt(3)*(25*x^11 + 2359*x^9 + 15426*x^7 + 6966*x^5 - 4347*x^3 + 243*x) - 4*(84*x^10 + 4536*x^8 + 20880*x^6 + 5832*x^4 - 2916*x^2 - sqrt(3)*(x^11 + 521*x^9 + 7362*x^7 + 10746*x^5 - 1971*x^3 - 243*x))*(-3*x^2 + 1)^(1/3) + 729)/(x^12 - 18*x^10 + 135*x^8 - 540*x^6 + 1215*x^4 - 1458*x^2 + 729)) + 1/144*sqrt(3)*log((x^6 - 93*x^4 - 117*x^2 + 2*(3*x^4 + 8*sqrt(3)*x^3 + 18*x^2 - 9))*(-3*x^2 + 1)^(2/3) - 8*sqrt(3)*(x^5 + 13*x^3) + 2*(3*x^4 - 54*x^2 + sqrt(3)*(x^5 - 10*x^3 - 27*x) - 9))*(-3*x^2 + 1)^(1/3) + 9)/(x^6 - 9*x^4 + 27*x^2 - 27)) - 1/144*sqrt(3)*log((x^6 - 93*x^4 - 117*x^2 + 2*(3*x^4 - 8*sqrt(3)*x^3 + 18*x^2 - 9))*(-3*x^2 + 1)^(2/3) + 8*sqrt(3)*(x^5 + 13*x^3) + 2*(3*x^4 - 54*x^2 - sqrt(3)*(x^5 - 10*x^3 - 27*x) - 9))*(-3*x^2 + 1)^(1/3) + 9)/(x^6 - 9*x^4 + 27*x^2 - 27)) + 1/24*arctan(1/3*(216*x^11 - 80424*x^9 + 282096*x^7 - 134352*x^5 + 18360*x^3 + 12*(140*x^9 - 19440*x^7 + 8424*x^5 - 1584*x^3 + sqrt(3)*(x^10 + 589*x^8 + 3946*x^6 - 774*x^4 - 27*x^2 + 9) + 108*x))*(-3*x^2 + 1)^(2/3) + sqrt(3)*(x^12 + 3150*x^10 + 77991*x^8 + 4260*x^6 - 14337*x^4 + 2862*x^2 - 135) - 12*(x^11 - 1591*x^9 + 42426*x^7 - 15102*x^5 + 1269*x^3 + sqrt(3)*(27*x^10 + 2307*x^8 + 4574*x^6 - 2538*x^4 + 279*x^2 - 9) - 27*x))*(-3*x^2 + 1)^(1/3) - 648*x)/(x^12 - 4986*x^10 + 327519*...
```

Sympy [F]

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx = - \int \frac{1}{x^2 \sqrt[3]{1-3x^2} - 3\sqrt[3]{1-3x^2}} dx$$

input `integrate(1/(-3*x**2+1)**(1/3)/(-x**2+3),x)`

output `-Integral(1/(x**2*(1 - 3*x**2)**(1/3) - 3*(1 - 3*x**2)**(1/3)), x)`

Maxima [F]

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx = \int -\frac{1}{(x^2-3)(-3x^2+1)^{\frac{1}{3}}} dx$$

input `integrate(1/(-3*x^2+1)^(1/3)/(-x^2+3),x, algorithm="maxima")`

output `-integrate(1/((x^2 - 3)*(-3*x^2 + 1)^(1/3)), x)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx = \int -\frac{1}{(x^2-3)(-3x^2+1)^{\frac{1}{3}}} dx$$

input `integrate(1/(-3*x^2+1)^(1/3)/(-x^2+3),x, algorithm="giac")`

output `integrate(-1/((x^2 - 3)*(-3*x^2 + 1)^(1/3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx = - \int \frac{1}{(x^2-3)(1-3x^2)^{1/3}} dx$$

input `int(-1/((x^2 - 3)*(1 - 3*x^2)^(1/3)),x)`output `-int(1/((x^2 - 3)*(1 - 3*x^2)^(1/3)), x)`**Reduce [F]**

$$\int \frac{1}{\sqrt[3]{1-3x^2}(3-x^2)} dx = - \left(\int \frac{1}{(-3x^2+1)^{1/3} x^2 - 3(-3x^2+1)^{1/3}} dx \right)$$

input `int(1/(-3*x^2+1)^(1/3)/(-x^2+3),x)`output `- int(1/((- 3*x**2 + 1)**(1/3)*x**2 - 3*(- 3*x**2 + 1)**(1/3)),x)`

3.78 $\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx$

Optimal result	608
Mathematica [C] (warning: unable to verify)	608
Rubi [A] (verified)	609
Maple [C] (verified)	610
Fricas [B] (verification not implemented)	611
Sympy [F]	611
Maxima [F]	612
Giac [F]	612
Mupad [F(-1)]	612
Reduce [F]	613

Optimal result

Integrand size = 19, antiderivative size = 81

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx = \frac{\arctan\left(\frac{x}{\sqrt{3}}\right)}{4\sqrt{3}} + \frac{\arctan\left(\frac{(1-\sqrt[3]{1+3x^2})^2}{3\sqrt{3}x}\right)}{4\sqrt{3}} - \frac{1}{4}\operatorname{arctanh}\left(\frac{1-\sqrt[3]{1+3x^2}}{x}\right)$$

output

```
-1/4*arctanh((1-(3*x^2+1)^(1/3))/x)+1/12*arctan(1/3*x*3^(1/2))*3^(1/2)+1/12*arctan(1/9*(1-(3*x^2+1)^(1/3))^2/x*3^(1/2))*3^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 3.92 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.56

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx =$$

$$\frac{9x \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, -3x^2, -\frac{x^2}{3}\right)}{(3+x^2)\sqrt[3]{1+3x^2}} - \frac{9 \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, -3x^2, -\frac{x^2}{3}\right) + 2x^2 \left(\operatorname{AppellF1}\left(\frac{3}{2}, \frac{1}{3}, 2, \frac{5}{2}, -3x^2, -\frac{x^2}{3}\right) + 3A\right)}{(3+x^2)\sqrt[3]{1+3x^2}}$$

input `Integrate[1/((3 + x^2)*(1 + 3*x^2)^(1/3)),x]`

output `(-9*x*AppellF1[1/2, 1/3, 1, 3/2, -3*x^2, -1/3*x^2])/((3 + x^2)*(1 + 3*x^2)^(1/3)*(-9*AppellF1[1/2, 1/3, 1, 3/2, -3*x^2, -1/3*x^2] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -3*x^2, -1/3*x^2] + 3*AppellF1[3/2, 4/3, 1, 5/2, -3*x^2, -1/3*x^2])))`

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {306}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(x^2 + 3)\sqrt[3]{3x^2 + 1}} dx$$

↓ 306

$$\frac{\arctan\left(\frac{(1 - \sqrt[3]{3x^2 + 1})^2}{3\sqrt[3]{3x^2 + 1}}\right)}{4\sqrt{3}} + \frac{\arctan\left(\frac{x}{\sqrt{3}}\right)}{4\sqrt{3}} - \frac{1}{4}\operatorname{arctanh}\left(\frac{1 - \sqrt[3]{3x^2 + 1}}{x}\right)$$

input `Int[1/((3 + x^2)*(1 + 3*x^2)^(1/3)),x]`

output `ArcTan[x/Sqrt[3]]/(4*Sqrt[3]) + ArcTan[(1 - (1 + 3*x^2)^(1/3))^2/(3*Sqrt[3]*x)]/(4*Sqrt[3]) - ArcTanh[(1 - (1 + 3*x^2)^(1/3))/x]/4`

Definitions of rubi rules used

rule 306

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With
h[{q = Rt[b/a, 2]}, Simp[q*(ArcTan[q*(x/3)]/(12*Rt[a, 3]*d)), x] + (Simp[q*
(ArcTan[(Rt[a, 3] - (a + b*x^2)^(1/3)]/(3*Rt[a, 3]^2*q*x)]/(12*Rt[a, 3]*d
)), x] - Simp[q*(ArcTanh[(Sqrt[3]*(Rt[a, 3] - (a + b*x^2)^(1/3))]/(Rt[a, 3]
*q*x)]/(4*Sqrt[3]*Rt[a, 3]*d)), x)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0] && EqQ[b*c - 9*a*d, 0] && PosQ[b/a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 2.07 (sec) , antiderivative size = 443, normalized size of antiderivative = 5.47

method	result
trager	$\text{RootOf}(48_Z^2 + 12_Z + 1) \ln \left(-\frac{-24\text{RootOf}(48_Z^2 + 12_Z + 1)(3x^2 + 1)^{\frac{1}{3}}x + 12\text{RootOf}(48_Z^2 + 12_Z + 1)}{\dots} \right)$

input

```
int(1/(x^2+3)/(3*x^2+1)^(1/3),x,method=_RETURNVERBOSE)
```

output

```
RootOf(48*_Z^2+12*_Z+1)*ln(-(-24*RootOf(48*_Z^2+12*_Z+1)*(3*x^2+1)^(1/3)*x
+12*RootOf(48*_Z^2+12*_Z+1)*x^2+2*(3*x^2+1)^(2/3)-24*RootOf(48*_Z^2+12*_Z+
1)*(3*x^2+1)^(1/3)-4*(3*x^2+1)^(1/3)*x+48*RootOf(48*_Z^2+12*_Z+1)*x+x^2-4*
(3*x^2+1)^(1/3)-12*RootOf(48*_Z^2+12*_Z+1)+4*x-1)/(x^2+3))-1/4*ln(-((12*Ro
otOf(48*_Z^2+12*_Z+1)*(3*x^2+1)^(1/3)*x-6*RootOf(48*_Z^2+12*_Z+1)*x^2+(3*x^
2+1)^(2/3)+12*RootOf(48*_Z^2+12*_Z+1)*(3*x^2+1)^(1/3)+(3*x^2+1)^(1/3)*x-24
*RootOf(48*_Z^2+12*_Z+1)*x-x^2+(3*x^2+1)^(1/3)+6*RootOf(48*_Z^2+12*_Z+1)-4
*x+1)/(x^2+3))-ln(-((12*RootOf(48*_Z^2+12*_Z+1)*(3*x^2+1)^(1/3)*x-6*RootOf(
48*_Z^2+12*_Z+1)*x^2+(3*x^2+1)^(2/3)+12*RootOf(48*_Z^2+12*_Z+1)*(3*x^2+1)^(
1/3)+(3*x^2+1)^(1/3)*x-24*RootOf(48*_Z^2+12*_Z+1)*x-x^2+(3*x^2+1)^(1/3)+6
*RootOf(48*_Z^2+12*_Z+1)-4*x+1)/(x^2+3))*RootOf(48*_Z^2+12*_Z+1)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 345 vs. $2(59) = 118$.

Time = 0.86 (sec) , antiderivative size = 345, normalized size of antiderivative = 4.26

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx$$

$$= \frac{1}{36} \sqrt{3} \arctan \left(\frac{4\sqrt{3}(3x^4 - 10x^3 - 36x^2 + 18x + 9)(3x^2 + 1)^{\frac{2}{3}} - 4\sqrt{3}(x^5 + 15x^4 - 26x^3 - 54x^2 + 9x - 9)(3x^2 + 1)^{\frac{1}{3}}}{x^6 + 126x^5 - 225x^4 - 828x^3 - 81x^2 - 162x + 81} \right)$$

$$- \frac{1}{36} \sqrt{3} \arctan \left(\frac{2(2\sqrt{3}(23x^3 + 9x)(3x^2 + 1)^{\frac{2}{3}} + \sqrt{3}(x^5 - 80x^3 - 9x)(3x^2 + 1)^{\frac{1}{3}} + \sqrt{3}(11x^5 + 10x^3 - 9x)(3x^2 + 1)^{\frac{1}{3}})}{x^6 - 657x^4 - 189x^2 - 27} \right)$$

$$+ \frac{1}{24} \log \left(\frac{x^6 + 108x^5 + 549x^4 + 360x^3 + 99x^2 + 6(3x^4 + 32x^3 + 42x^2 + 3)(3x^2 + 1)^{\frac{2}{3}} + 6(x^5 + 27x^4 + 70x^3 + 18x^2 + 9x + 3)(3x^2 + 1)^{\frac{1}{3}} + 108x - 9}{x^6 + 9x^4 + 27x^2 + 27} \right)$$

input `integrate(1/(x^2+3)/(3*x^2+1)^(1/3),x, algorithm="fricas")`

output

```
1/36*sqrt(3)*arctan((4*sqrt(3)*(3*x^4 - 10*x^3 - 36*x^2 + 18*x + 9)*(3*x^2 + 1)^(2/3) - 4*sqrt(3)*(x^5 + 15*x^4 - 26*x^3 - 54*x^2 + 9*x - 9)*(3*x^2 + 1)^(1/3) + sqrt(3)*(x^6 - 2*x^5 - 105*x^4 - 28*x^3 + 63*x^2 + 126*x + 9))/(x^6 + 126*x^5 - 225*x^4 - 828*x^3 - 81*x^2 - 162*x + 81)) - 1/36*sqrt(3)*arctan(2*(2*sqrt(3)*(23*x^3 + 9*x)*(3*x^2 + 1)^(2/3) + sqrt(3)*(x^5 - 80*x^3 - 9*x)*(3*x^2 + 1)^(1/3) + sqrt(3)*(11*x^5 + 10*x^3 - 9*x))/(x^6 - 657*x^4 - 189*x^2 - 27)) + 1/24*log((x^6 + 108*x^5 + 549*x^4 + 360*x^3 + 99*x^2 + 6*(3*x^4 + 32*x^3 + 42*x^2 + 3)*(3*x^2 + 1)^(2/3) + 6*(x^5 + 27*x^4 + 70*x^3 + 18*x^2 + 9*x + 3)*(3*x^2 + 1)^(1/3) + 108*x - 9)/(x^6 + 9*x^4 + 27*x^2 + 27))
```

Sympy [F]

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx = \int \frac{1}{(x^2+3)\sqrt[3]{3x^2+1}} dx$$

input `integrate(1/(x**2+3)/(3*x**2+1)**(1/3),x)`

output `Integral(1/((x**2 + 3)*(3*x**2 + 1)**(1/3)), x)`

Maxima [F]

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx = \int \frac{1}{(3x^2+1)^{\frac{1}{3}}(x^2+3)} dx$$

input `integrate(1/(x^2+3)/(3*x^2+1)^(1/3),x, algorithm="maxima")`

output `integrate(1/((3*x^2 + 1)^(1/3)*(x^2 + 3)), x)`

Giac [F]

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx = \int \frac{1}{(3x^2+1)^{\frac{1}{3}}(x^2+3)} dx$$

input `integrate(1/(x^2+3)/(3*x^2+1)^(1/3),x, algorithm="giac")`

output `integrate(1/((3*x^2 + 1)^(1/3)*(x^2 + 3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx = \int \frac{1}{(x^2+3)(3x^2+1)^{\frac{1}{3}}} dx$$

input `int(1/((x^2 + 3)*(3*x^2 + 1)^(1/3)),x)`

output `int(1/((x^2 + 3)*(3*x^2 + 1)^(1/3)), x)`

Reduce [F]

$$\int \frac{1}{(3+x^2)\sqrt[3]{1+3x^2}} dx = \int \frac{1}{(3x^2+1)^{\frac{1}{3}} x^2 + 3(3x^2+1)^{\frac{1}{3}}} dx$$

input `int(1/(x^2+3)/(3*x^2+1)^(1/3),x)`

output `int(1/((3*x**2 + 1)**(1/3)*x**2 + 3*(3*x**2 + 1)**(1/3)),x)`

3.79 $\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx$

Optimal result	614
Mathematica [C] (warning: unable to verify)	615
Rubi [A] (verified)	615
Maple [C] (verified)	616
Fricas [B] (verification not implemented)	617
Sympy [F]	618
Maxima [F]	619
Giac [F]	619
Mupad [F(-1)]	619
Reduce [F]	620

Optimal result

Integrand size = 19, antiderivative size = 113

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx = \frac{\arctan\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan\left(\frac{\sqrt{3}\left(1-\sqrt[3]{2}\sqrt[3]{1-x^2}\right)}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}} + \frac{\operatorname{arctanh}\left(\frac{x}{1+\sqrt[3]{2}\sqrt[3]{1-x^2}}\right)}{2 \cdot 2^{2/3}}$$

output

```
-1/12*arctanh(x)*2^(1/3)+1/4*arctanh(x/(1+2^(1/3)*(-x^2+1)^(1/3)))*2^(1/3)
+1/12*arctan(3^(1/2)/x)*2^(1/3)*3^(1/2)+1/12*arctan((1-2^(1/3)*(-x^2+1)^(1/3))*3^(1/2)/x)*2^(1/3)*3^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 3.93 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.04

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx =$$

$$\frac{9x \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, x^2, -\frac{x^2}{3}\right)}{\sqrt[3]{1-x^2}(3+x^2) \left(-9 \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, x^2, -\frac{x^2}{3}\right) + 2x^2 \left(\operatorname{AppellF1}\left(\frac{3}{2}, \frac{1}{3}, 2, \frac{5}{2}, x^2, -\frac{x^2}{3}\right) - \operatorname{AppellF1}\left(\frac{3}{2}, \frac{1}{3}, 1, \frac{5}{2}, x^2, -\frac{x^2}{3}\right)\right)\right)}$$

input `Integrate[1/((1 - x^2)^(1/3)*(3 + x^2)),x]`

output `(-9*x*AppellF1[1/2, 1/3, 1, 3/2, x^2, -1/3*x^2])/((1 - x^2)^(1/3)*(3 + x^2))*(-9*AppellF1[1/2, 1/3, 1, 3/2, x^2, -1/3*x^2] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5/2, x^2, -1/3*x^2] - AppellF1[3/2, 4/3, 1, 5/2, x^2, -1/3*x^2]))`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {305}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{1-x^2}(x^2+3)} dx$$

↓ 305

$$\frac{\arctan\left(\frac{\sqrt{3}\left(1-\sqrt[3]{2}\sqrt[3]{1-x^2}\right)}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\arctan\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3} \sqrt{3}} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{1-x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}(x)}{6 \cdot 2^{2/3}}$$

input `Int[1/((1 - x^2)^(1/3)*(3 + x^2)),x]`

output

$$\text{ArcTan}[\text{Sqrt}[3]/x]/(2*2^{(2/3)}*\text{Sqrt}[3]) + \text{ArcTan}[(\text{Sqrt}[3]*(1 - 2^{(1/3)}*(1 - x^2)^{(1/3)}))/x]/(2*2^{(2/3)}*\text{Sqrt}[3]) - \text{ArcTanh}[x]/(6*2^{(2/3)}) + \text{ArcTanh}[x/(1 + 2^{(1/3)}*(1 - x^2)^{(1/3)})]/(2*2^{(2/3)})$$

Defintions of rubi rules used

rule 305

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] :> With
h[{q = Rt[-b/a, 2]}, Simp[q*(ArcTan[Sqrt[3]/(q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d)), x] + (Simp[q*(ArcTanh[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3))]/(2*2^(2/3)*a^(1/3)*d)), x] - Simp[q*(ArcTanh[q*x]/(6*2^(2/3)*a^(1/3)*d)), x] + Simp[q*(ArcTan[Sqrt[3]*((a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d)), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && NegQ[b/a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 14.20 (sec) , antiderivative size = 938, normalized size of antiderivative = 8.30

method	result	size
trager	Expression too large to display	938

input

```
int(1/(-x^2+1)^(1/3)/(x^2+3),x,method=_RETURNVERBOSE)
```

output

```

1/36*RootOf(_Z^6+108)*ln((-486*RootOf(_Z^6+108)+189*x^2*RootOf(_Z^6+108)^4
-RootOf(_Z^6+108)^4*x^6-18*RootOf(_Z^6+108)*x^6-72*RootOf(_Z^6+108)^4*x^5-
225*RootOf(_Z^6+108)^4*x^4+72*RootOf(_Z^6+108)^4*x^3-1296*RootOf(_Z^6+108)
*x^5-4050*RootOf(_Z^6+108)*x^4+1296*RootOf(_Z^6+108)*x^3+3402*RootOf(_Z^6+
108)*x^2-27*RootOf(_Z^6+108)^4+1296*(-x^2+1)^(2/3)*x^4+9072*(-x^2+1)^(2/3)
*x^3+3888*(-x^2+1)^(2/3)*x^2-3888*(-x^2+1)^(2/3)*x-6*(-x^2+1)^(1/3)*RootOf
(_Z^6+108)^5*x^5-108*(-x^2+1)^(1/3)*RootOf(_Z^6+108)^5*x^4-144*(-x^2+1)^(1
/3)*RootOf(_Z^6+108)^5*x^3+108*(-x^2+1)^(1/3)*RootOf(_Z^6+108)^5*x^2+54*(-
x^2+1)^(1/3)*RootOf(_Z^6+108)^5*x-36*RootOf(_Z^6+108)^2*(-x^2+1)^(1/3)*x^5
-648*RootOf(_Z^6+108)^2*(-x^2+1)^(1/3)*x^4-864*RootOf(_Z^6+108)^2*(-x^2+1)
^(1/3)*x^3+648*RootOf(_Z^6+108)^2*(-x^2+1)^(1/3)*x^2+324*RootOf(_Z^6+108)^
2*(-x^2+1)^(1/3)*x)/(x^2+3)^3)-1/432*ln((RootOf(_Z^6+108)^4*x^6+72*RootOf(
_Z^6+108)^4*x^5+36*RootOf(_Z^6+108)^2*(-x^2+1)^(1/3)*x^5+225*RootOf(_Z^6+1
08)^4*x^4+648*RootOf(_Z^6+108)^2*(-x^2+1)^(1/3)*x^4-72*RootOf(_Z^6+108)^4*
x^3+648*(-x^2+1)^(2/3)*x^4+864*RootOf(_Z^6+108)^2*(-x^2+1)^(1/3)*x^3-189*x
^2*RootOf(_Z^6+108)^4+4536*(-x^2+1)^(2/3)*x^3-648*RootOf(_Z^6+108)^2*(-x^2
+1)^(1/3)*x^2+1944*(-x^2+1)^(2/3)*x^2-324*RootOf(_Z^6+108)^2*(-x^2+1)^(1/3
)*x+27*RootOf(_Z^6+108)^4-1944*(-x^2+1)^(2/3)*x)/(x^2+3)^3)*RootOf(_Z^6+10
8)^4+1/72*ln((RootOf(_Z^6+108)^4*x^6+72*RootOf(_Z^6+108)^4*x^5+36*RootOf(
_Z^6+108)^2*(-x^2+1)^(1/3)*x^5+225*RootOf(_Z^6+108)^4*x^4+648*RootOf(_Z^...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1132 vs. $2(81) = 162$.

Time = 0.50 (sec) , antiderivative size = 1132, normalized size of antiderivative = 10.02

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx = \text{Too large to display}$$

input

```
integrate(1/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="fricas")
```

output

```

-1/24*(-1/432)^(1/6)*(sqrt(-3) + 1)*log((2*(-1/2)^(2/3)*(5*x^5 - 30*x^3 +
sqrt(-3)*(5*x^5 - 30*x^3 + 9*x) + 9*x) + 12*(3*x^3 + sqrt(-1/3)*(x^4 - 9*x
^2) - 3*x)*(-x^2 + 1)^(2/3) + 2*(-x^2 + 1)^(1/3)*(864*(-1/432)^(5/6)*(x^4
- 3*x^2 - sqrt(-3)*(x^4 - 3*x^2)) - (-1/2)^(1/3)*(x^5 - 18*x^3 - sqrt(-3)*
(x^5 - 18*x^3 + 9*x) + 9*x)) - (-1/432)^(1/6)*(x^6 - 69*x^4 + 63*x^2 + sqr
t(-3)*(x^6 - 69*x^4 + 63*x^2 - 27) - 27))/(x^6 + 9*x^4 + 27*x^2 + 27)) + 1
/24*(-1/432)^(1/6)*(sqrt(-3) + 1)*log((2*(-1/2)^(2/3)*(5*x^5 - 30*x^3 + sq
rt(-3)*(5*x^5 - 30*x^3 + 9*x) + 9*x) + 12*(3*x^3 - sqrt(-1/3)*(x^4 - 9*x^2
) - 3*x)*(-x^2 + 1)^(2/3) - 2*(-x^2 + 1)^(1/3)*(864*(-1/432)^(5/6)*(x^4 -
3*x^2 - sqrt(-3)*(x^4 - 3*x^2)) + (-1/2)^(1/3)*(x^5 - 18*x^3 - sqrt(-3)*(x
^5 - 18*x^3 + 9*x) + 9*x)) + (-1/432)^(1/6)*(x^6 - 69*x^4 + 63*x^2 + sqrt(
-3)*(x^6 - 69*x^4 + 63*x^2 - 27) - 27))/(x^6 + 9*x^4 + 27*x^2 + 27)) + 1/2
4*(-1/432)^(1/6)*(sqrt(-3) - 1)*log((2*(-1/2)^(2/3)*(5*x^5 - 30*x^3 - sqrt
(-3)*(5*x^5 - 30*x^3 + 9*x) + 9*x) + 12*(3*x^3 + sqrt(-1/3)*(x^4 - 9*x^2)
- 3*x)*(-x^2 + 1)^(2/3) + 2*(-x^2 + 1)^(1/3)*(864*(-1/432)^(5/6)*(x^4 - 3*
x^2 + sqrt(-3)*(x^4 - 3*x^2)) - (-1/2)^(1/3)*(x^5 - 18*x^3 + sqrt(-3)*(x^5
- 18*x^3 + 9*x) + 9*x)) - (-1/432)^(1/6)*(x^6 - 69*x^4 + 63*x^2 - sqrt(-3
)*(x^6 - 69*x^4 + 63*x^2 - 27) - 27))/(x^6 + 9*x^4 + 27*x^2 + 27)) - 1/24*
(-1/432)^(1/6)*(sqrt(-3) - 1)*log((2*(-1/2)^(2/3)*(5*x^5 - 30*x^3 - sqrt(-
3)*(5*x^5 - 30*x^3 + 9*x) + 9*x) + 12*(3*x^3 - sqrt(-1/3)*(x^4 - 9*x^2)...

```

Sympy [F]

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx = \int \frac{1}{\sqrt[3]{-(x-1)(x+1)(x^2+3)}} dx$$

input

```
integrate(1/((-x**2+1)**(1/3)/(x**2+3), x)
```

output

```
Integral(1/((-x - 1)*(x + 1)**(1/3)*(x**2 + 3)), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx = \int \frac{1}{(x^2+3)(-x^2+1)^{\frac{1}{3}}} dx$$

input `integrate(1/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="maxima")`

output `integrate(1/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx = \int \frac{1}{(x^2+3)(-x^2+1)^{\frac{1}{3}}} dx$$

input `integrate(1/(-x^2+1)^(1/3)/(x^2+3),x, algorithm="giac")`

output `integrate(1/((x^2 + 3)*(-x^2 + 1)^(1/3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx = \int \frac{1}{(1-x^2)^{1/3}(x^2+3)} dx$$

input `int(1/((1 - x^2)^(1/3)*(x^2 + 3)),x)`

output `int(1/((1 - x^2)^(1/3)*(x^2 + 3)), x)`

Reduce [F]

$$\int \frac{1}{\sqrt[3]{1-x^2}(3+x^2)} dx = \int \frac{1}{(-x^2+1)^{\frac{1}{3}}x^2+3(-x^2+1)^{\frac{1}{3}}} dx$$

input `int(1/(-x^2+1)^(1/3)/(x^2+3),x)`

output `int(1/((-x**2+1)**(1/3)*x**2+3*(-x**2+1)**(1/3)),x)`

3.80 $\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx$

Optimal result	621
Mathematica [C] (warning: unable to verify)	622
Rubi [A] (verified)	622
Maple [F]	623
Fricas [B] (verification not implemented)	623
Sympy [F]	624
Maxima [F]	625
Giac [F]	625
Mupad [F(-1)]	625
Reduce [F]	626

Optimal result

Integrand size = 19, antiderivative size = 109

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = -\frac{\arctan(x)}{6 \cdot 2^{2/3}} + \frac{\arctan\left(\frac{x}{1+\sqrt[3]{2}\sqrt[3]{1+x^2}}\right)}{2 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}(1-\sqrt[3]{2}\sqrt[3]{1+x^2})}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}}$$

output

```
-1/12*arctan(x)*2^(1/3)+1/4*arctan(x/(1+2^(1/3)*(x^2+1)^(1/3)))*2^(1/3)-1/12*arctanh(3^(1/2)/x)*2^(1/3)*3^(1/2)-1/12*arctanh((1-2^(1/3)*(x^2+1)^(1/3))*3^(1/2)/x)*2^(1/3)*3^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 4.05 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.14

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = \frac{9x \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, -x^2, \frac{x^2}{3}\right)}{(-3+x^2)\sqrt[3]{1+x^2} \left(9 \operatorname{AppellF1}\left(\frac{1}{2}, \frac{1}{3}, 1, \frac{3}{2}, -x^2, \frac{x^2}{3}\right) + 2x^2 \left(\operatorname{AppellF1}\left(\frac{3}{2}, \frac{1}{3}, 2, \frac{5}{2}, -x^2, \frac{x^2}{3}\right) - \operatorname{AppellF1}\left(\frac{3}{2}, \frac{1}{3}, 2, \frac{5}{2}, -x^2, \frac{x^2}{3}\right)\right)\right)}$$

input `Integrate[1/((3 - x^2)*(1 + x^2)^(1/3)),x]`

output `(-9*x*AppellF1[1/2, 1/3, 1, 3/2, -x^2, x^2/3])/((-3 + x^2)*(1 + x^2)^(1/3) * (9*AppellF1[1/2, 1/3, 1, 3/2, -x^2, x^2/3] + 2*x^2*(AppellF1[3/2, 1/3, 2, 5/2, -x^2, x^2/3] - AppellF1[3/2, 4/3, 1, 5/2, -x^2, x^2/3])))`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {304}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(3-x^2)\sqrt[3]{x^2+1}} dx \xrightarrow{304} \frac{\arctan\left(\frac{x}{\sqrt[3]{2}\sqrt[3]{x^2+1}}\right)}{2 \cdot 2^{2/3}} - \frac{\arctan(x)}{6 \cdot 2^{2/3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}\left(1-\sqrt[3]{2}\sqrt[3]{x^2+1}\right)}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}}{x}\right)}{2 \cdot 2^{2/3}\sqrt{3}}$$

input `Int[1/((3 - x^2)*(1 + x^2)^(1/3)),x]`

output

```
-1/6*ArcTan[x]/2^(2/3) + ArcTan[x/(1 + 2^(1/3)*(1 + x^2)^(1/3))]/(2*2^(2/3)) - ArcTanh[Sqrt[3]/x]/(2*2^(2/3)*Sqrt[3]) - ArcTanh[(Sqrt[3]*(1 - 2^(1/3)*(1 + x^2)^(1/3)))/x]/(2*2^(2/3)*Sqrt[3])
```

Defintions of rubi rules used

rule 304

```
Int[1/(((a_) + (b_.)*(x_)^2)^(1/3)*((c_) + (d_.)*(x_)^2)), x_Symbol] := With[{q = Rt[b/a, 2]}, Simp[q*(ArcTanh[Sqrt[3]/(q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d)), x] + (-Simp[q*(ArcTan[(a^(1/3)*q*x)/(a^(1/3) + 2^(1/3)*(a + b*x^2)^(1/3))]/(2*2^(2/3)*a^(1/3)*d)), x] + Simp[q*(ArcTan[q*x]/(6*2^(2/3)*a^(1/3)*d)), x] + Simp[q*(ArcTanh[Sqrt[3]*((a^(1/3) - 2^(1/3)*(a + b*x^2)^(1/3)))/(a^(1/3)*q*x)]/(2*2^(2/3)*Sqrt[3]*a^(1/3)*d)), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b*c + 3*a*d, 0] && PosQ[b/a]
```

Maple [F]

$$\int \frac{1}{(-x^2 + 3)(x^2 + 1)^{\frac{1}{3}}} dx$$

input

```
int(1/(-x^2+3)/(x^2+1)^(1/3),x)
```

output

```
int(1/(-x^2+3)/(x^2+1)^(1/3),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1103 vs. $2(77) = 154$.

Time = 0.49 (sec) , antiderivative size = 1103, normalized size of antiderivative = 10.12

$$\int \frac{1}{(3 - x^2) \sqrt[3]{1 + x^2}} dx = \text{Too large to display}$$

input

```
integrate(1/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="fricas")
```


output

```
-1/10368*432^(5/6)*(sqrt(-3) + 1)*log((432^(5/6)*(x^6 + 69*x^4 + 63*x^2 +
sqrt(-3)*(x^6 + 69*x^4 + 63*x^2 + 27) + 27) - 1728*(9*x^3 + sqrt(3)*(x^4 +
9*x^2) + 9*x)*(x^2 + 1)^(2/3) + 432*2^(1/3)*(5*x^5 + 30*x^3 + sqrt(-3)*(5
*x^5 + 30*x^3 + 9*x) + 9*x) + 432*(x^2 + 1)^(1/3)*(2^(2/3)*(x^5 + 18*x^3 -
sqrt(-3)*(x^5 + 18*x^3 + 9*x) + 9*x) + 4*432^(1/6)*(x^4 + 3*x^2 - sqrt(-3
)*(x^4 + 3*x^2)))))/(x^6 - 9*x^4 + 27*x^2 - 27)) + 1/10368*432^(5/6)*(sqrt(
-3) + 1)*log(-(432^(5/6)*(x^6 + 69*x^4 + 63*x^2 + sqrt(-3)*(x^6 + 69*x^4 +
63*x^2 + 27) + 27) + 1728*(9*x^3 - sqrt(3)*(x^4 + 9*x^2) + 9*x)*(x^2 + 1)
^(2/3) - 432*2^(1/3)*(5*x^5 + 30*x^3 + sqrt(-3)*(5*x^5 + 30*x^3 + 9*x) + 9
*x) - 432*(x^2 + 1)^(1/3)*(2^(2/3)*(x^5 + 18*x^3 - sqrt(-3)*(x^5 + 18*x^3
+ 9*x) + 9*x) - 4*432^(1/6)*(x^4 + 3*x^2 - sqrt(-3)*(x^4 + 3*x^2)))))/(x^6
- 9*x^4 + 27*x^2 - 27)) + 1/10368*432^(5/6)*(sqrt(-3) - 1)*log((432^(5/6)*
(x^6 + 69*x^4 + 63*x^2 - sqrt(-3)*(x^6 + 69*x^4 + 63*x^2 + 27) + 27) - 172
8*(9*x^3 + sqrt(3)*(x^4 + 9*x^2) + 9*x)*(x^2 + 1)^(2/3) + 432*2^(1/3)*(5*x
^5 + 30*x^3 - sqrt(-3)*(5*x^5 + 30*x^3 + 9*x) + 9*x) + 432*(x^2 + 1)^(1/3)
*(2^(2/3)*(x^5 + 18*x^3 + sqrt(-3)*(x^5 + 18*x^3 + 9*x) + 9*x) + 4*432^(1/
6)*(x^4 + 3*x^2 + sqrt(-3)*(x^4 + 3*x^2)))))/(x^6 - 9*x^4 + 27*x^2 - 27)) -
1/10368*432^(5/6)*(sqrt(-3) - 1)*log(-(432^(5/6)*(x^6 + 69*x^4 + 63*x^2 -
sqrt(-3)*(x^6 + 69*x^4 + 63*x^2 + 27) + 27) + 1728*(9*x^3 - sqrt(3)*(x^4
+ 9*x^2) + 9*x)*(x^2 + 1)^(2/3) - 432*2^(1/3)*(5*x^5 + 30*x^3 - sqrt(-3...
```

Sympy [F]

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = -\int \frac{1}{x^2\sqrt[3]{x^2+1} - 3\sqrt[3]{x^2+1}} dx$$

input

```
integrate(1/(-x**2+3)/(x**2+1)**(1/3),x)
```

output

```
-Integral(1/(x**2*(x**2 + 1)**(1/3) - 3*(x**2 + 1)**(1/3)), x)
```

Maxima [F]

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = \int -\frac{1}{(x^2+1)^{\frac{1}{3}}(x^2-3)} dx$$

input `integrate(1/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="maxima")`

output `-integrate(1/((x^2 + 1)^(1/3)*(x^2 - 3)), x)`

Giac [F]

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = \int -\frac{1}{(x^2+1)^{\frac{1}{3}}(x^2-3)} dx$$

input `integrate(1/(-x^2+3)/(x^2+1)^(1/3),x, algorithm="giac")`

output `integrate(-1/((x^2 + 1)^(1/3)*(x^2 - 3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = -\int \frac{1}{(x^2+1)^{1/3}(x^2-3)} dx$$

input `int(-1/((x^2 + 1)^(1/3)*(x^2 - 3)),x)`

output `-int(1/((x^2 + 1)^(1/3)*(x^2 - 3)), x)`

Reduce [F]

$$\int \frac{1}{(3-x^2)\sqrt[3]{1+x^2}} dx = - \left(\int \frac{1}{(x^2+1)^{\frac{1}{3}} x^2 - 3(x^2+1)^{\frac{1}{3}}} dx \right)$$

input `int(1/(-x^2+3)/(x^2+1)^(1/3),x)`

output `- int(1/((x**2 + 1)**(1/3)*x**2 - 3*(x**2 + 1)**(1/3)),x)`

3.81
$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx$$

Optimal result	627
Mathematica [A] (verified)	627
Rubi [A] (verified)	628
Maple [A] (verified)	630
Fricas [A] (verification not implemented)	630
Sympy [F]	631
Maxima [F]	631
Giac [F]	631
Mupad [B] (verification not implemented)	632
Reduce [F]	632

Optimal result

Integrand size = 34, antiderivative size = 87

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx = -\frac{2\sqrt{x}\sqrt{a^2-(1+a^2)x+x^2} \arctan\left(\frac{(1-a)\sqrt{x}}{\sqrt{a^2-(1+a^2)x+x^2}}\right)}{(1-a)\sqrt{a^2x-(1+a^2)x^2+x^3}}$$

output

$$-2*\arctan((1-a)*x^{(1/2)}/(a^2-(a^2+1)*x+x^2)^{(1/2)})*x^{(1/2)}*(a^2-(a^2+1)*x+x^2)^{(1/2)}/(1-a)/(a^2*x-(a^2+1)*x^2+x^3)^{(1/2)}$$

Mathematica [A] (verified)

Time = 10.36 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.33

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx = -\frac{2 \arctan\left(\frac{(-1+a)x}{\sqrt{(-1+x)x(-a^2+x)}}\right)}{-1+a}$$

input

`Integrate[(a + x)/((-a + x)*Sqrt[a^2*x - (1 + a^2)*x^2 + x^3]),x]`

output $(-2*\text{ArcTan}[((-1 + a)*x)/\text{Sqrt}[(-1 + x)*x*(-a^2 + x)]]/(-1 + a)$

Rubi [A] (verified)

Time = 0.50 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.147$, Rules used = {2467, 25, 2035, 2212, 218}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a+x}{(x-a)\sqrt{-(a^2+1)x^2+a^2x+x^3}} dx$$

$$\downarrow 2467$$

$$\frac{\sqrt{x}\sqrt{-(a^2+1)x+a^2+x^2} \int -\frac{a+x}{(a-x)\sqrt{x}\sqrt{a^2+x^2-(a^2+1)x}} dx}{\sqrt{-(a^2+1)x^2+a^2x+x^3}}$$

$$\downarrow 25$$

$$-\frac{\sqrt{x}\sqrt{-(a^2+1)x+a^2+x^2} \int \frac{a+x}{(a-x)\sqrt{x}\sqrt{a^2+x^2-(a^2+1)x}} dx}{\sqrt{-(a^2+1)x^2+a^2x+x^3}}$$

$$\downarrow 2035$$

$$-\frac{2\sqrt{x}\sqrt{-(a^2+1)x+a^2+x^2} \int \frac{a+x}{(a-x)\sqrt{a^2+x^2-(a^2+1)x}} d\sqrt{x}}{\sqrt{-(a^2+1)x^2+a^2x+x^3}}$$

$$\downarrow 2212$$

$$-\frac{2a\sqrt{x}\sqrt{-(a^2+1)x+a^2+x^2} \int \frac{1}{ax(1-a)^2+a} d\frac{\sqrt{x}}{\sqrt{a^2+x^2-(a^2+1)x}}}{\sqrt{-(a^2+1)x^2+a^2x+x^3}}$$

$$\downarrow 218$$

$$-\frac{2\sqrt{x}\sqrt{-(a^2+1)x+a^2+x^2} \arctan\left(\frac{(1-a)\sqrt{x}}{\sqrt{-(a^2+1)x+a^2+x^2}}\right)}{(1-a)\sqrt{-(a^2+1)x^2+a^2x+x^3}}$$

input $\text{Int}[(a+x)/((-a+x)*\text{Sqrt}[a^2*x - (1+a^2)*x^2 + x^3]),x]$

output
$$\frac{(-2\sqrt{x}\sqrt{a^2 - (1 + a^2)x + x^2})\operatorname{ArcTan}\left[\frac{(1 - a)\sqrt{x}}{\sqrt{a^2 - (1 + a^2)x + x^2}}\right]}{(1 - a)\sqrt{a^2x - (1 + a^2)x^2 + x^3}}$$

Defintions of rubi rules used

rule 25 $\operatorname{Int}[-(Fx_), x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Identity}[-1] \operatorname{Int}[Fx, x], x]$

rule 218 $\operatorname{Int}[(a_ + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2]/a)\operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b]$

rule 2035 $\operatorname{Int}[(Fx_)(x_)^m, x_Symbol] \rightarrow \operatorname{With}\{k = \operatorname{Denominator}[m]\}, \operatorname{Simp}[k \operatorname{Subst}[\operatorname{Int}[x^{k(m+1)} - 1] \operatorname{SubstPower}[Fx, x, k], x], x, x^{1/k}], x] /; \operatorname{FractionQ}[m] \ \&\& \operatorname{AlgebraicFunctionQ}[Fx, x]$

rule 2212 $\operatorname{Int}[(A_ + (B_)(x_)^2)/((d_ + (e_)(x_)^2)\sqrt{(a_ + (b_)(x_)^2 + (c_)(x_)^4})], x_Symbol] \rightarrow \operatorname{Simp}[A \operatorname{Subst}[\operatorname{Int}[1/(d - (b*d - 2*a*e)x^2), x], x, x/\sqrt{a + b*x^2 + c*x^4}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, A, B\}, x \ \&\& \operatorname{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \operatorname{EqQ}[B*d + A*e, 0]$

rule 2467 $\operatorname{Int}[(Fx_)(Px_)^p, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Expon}[Px, x, \operatorname{Min}]\}, \operatorname{Simp}[Px^{\operatorname{FracPart}[p]}/(x^{(r*\operatorname{FracPart}[p])})\operatorname{ExpandToSum}[Px/x^r, x]^{\operatorname{FracPart}[p]} \operatorname{Int}[x^{(p*r)}\operatorname{ExpandToSum}[Px/x^r, x]^p Fx, x], x] /; \operatorname{IGtQ}[r, 0] /; \operatorname{FreeQ}[p, x] \ \&\& \operatorname{PolyQ}[Px, x] \ \&\& \operatorname{IntegerQ}[p] \ \&\& \operatorname{MonomialQ}[Px, x] \ \&\& \operatorname{PolyQ}[Fx, x]$

Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.38

method	result
default	$\frac{2 \arctan\left(\frac{\sqrt{-(a^2-x)x(-1+x)}}{x(a-1)}\right)}{a-1}$
pseudoelliptic	$\frac{2 \arctan\left(\frac{\sqrt{-(a^2-x)x(-1+x)}}{x(a-1)}\right)}{a-1}$
elliptic	$-\frac{2a^2 \sqrt{-\frac{-a^2+x}{a^2}} \sqrt{\frac{-1+x}{a^2-1}} \sqrt{\frac{x}{a^2}} \operatorname{EllipticF}\left(\sqrt{-\frac{-a^2+x}{a^2}}, \sqrt{\frac{a^2}{a^2-1}}\right)}{\sqrt{-a^2x^2+a^2x+x^3-x^2}} - \frac{4a^3 \sqrt{-\frac{-a^2+x}{a^2}} \sqrt{\frac{-1+x}{a^2-1}} \sqrt{\frac{x}{a^2}} \operatorname{EllipticPi}\left(\sqrt{-\frac{-a^2+x}{a^2}}, \sqrt{\frac{a^2}{a^2-1}}\right)}{\sqrt{-a^2x^2+a^2x+x^3-x^2} (a^2-a)}$

input `int((a+x)/(-a+x)/(a^2*x-(a^2+1)*x^2+x^3)^(1/2),x,method=_RETURNVERBOSE)`

output `2*arctan((-a^2-x)*x*(-1+x))^(1/2)/x/(a-1)/(a-1)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.98

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx = \frac{\arctan\left(\frac{\sqrt{a^2x-(a^2+1)x^2+x^3}(a^2-2(a^2-a+1)x+x^2)}{2((a-1)x^3-(a^3-a^2+a-1)x^2+(a^3-a^2)x)}\right)}{a-1}$$

input `integrate((a+x)/(-a+x)/(a^2*x-(a^2+1)*x^2+x^3)^(1/2),x, algorithm="fricas")`

output `arctan(1/2*sqrt(a^2*x - (a^2 + 1)*x^2 + x^3)*(a^2 - 2*(a^2 - a + 1)*x + x^2)/((a - 1)*x^3 - (a^3 - a^2 + a - 1)*x^2 + (a^3 - a^2)*x)/(a - 1)`

Sympy [F]

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx = \int \frac{a+x}{\sqrt{x(-a^2+x)(x-1)}(-a+x)} dx$$

input `integrate((a+x)/(-a+x)/(a**2*x-(a**2+1)*x**2+x**3)**(1/2),x)`

output `Integral((a + x)/(sqrt(x*(-a**2 + x)*(x - 1))*(-a + x)), x)`

Maxima [F]

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx = \int -\frac{a+x}{\sqrt{a^2x-(a^2+1)x^2+x^3}(a-x)} dx$$

input `integrate((a+x)/(-a+x)/(a^2*x-(a^2+1)*x^2+x^3)^(1/2),x, algorithm="maxima")`

output `-integrate((a + x)/(sqrt(a^2*x - (a^2 + 1)*x^2 + x^3)*(a - x)), x)`

Giac [F]

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx = \int -\frac{a+x}{\sqrt{a^2x-(a^2+1)x^2+x^3}(a-x)} dx$$

input `integrate((a+x)/(-a+x)/(a^2*x-(a^2+1)*x^2+x^3)^(1/2),x, algorithm="giac")`

output `integrate(-(a + x)/(sqrt(a^2*x - (a^2 + 1)*x^2 + x^3)*(a - x)), x)`

Mupad [B] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 217, normalized size of antiderivative = 2.49

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx$$

$$= \frac{4a(a^2-1)\sqrt{\frac{x}{a^2}}\sqrt{\frac{x-1}{a^2-1}}\sqrt{-\frac{x-a^2}{a^2-1}}\Pi\left(-\frac{a^2-1}{a-a^2}; \operatorname{asin}\left(\sqrt{-\frac{x-a^2}{a^2-1}}\right)\middle|\frac{a^2-1}{a^2}\right)}{(a-a^2)\sqrt{a^2x-x^2(a^2+1)+x^3}}$$

$$- \frac{2(a^2-1)\operatorname{F}\left(\operatorname{asin}\left(\sqrt{-\frac{x-a^2}{a^2-1}}\right)\middle|\frac{a^2-1}{a^2}\right)\sqrt{\frac{x}{a^2}}\sqrt{\frac{x-1}{a^2-1}}\sqrt{-\frac{x-a^2}{a^2-1}}}{\sqrt{a^2x-x^2(a^2+1)+x^3}}$$

input `int(-(a + x)/((a - x)*(a^2*x - x^2*(a^2 + 1) + x^3)^(1/2)),x)`output `(4*a*(a^2 - 1)*(x/a^2)^(1/2)*((x - 1)/(a^2 - 1))^(1/2)*(-(x - a^2)/(a^2 - 1))^(1/2)*ellipticPi(-(a^2 - 1)/(a - a^2), asin((-x - a^2)/(a^2 - 1))^(1/2), (a^2 - 1)/a^2))/((a - a^2)*(a^2*x - x^2*(a^2 + 1) + x^3)^(1/2)) - (2*(a^2 - 1)*ellipticF(asin((-x - a^2)/(a^2 - 1))^(1/2), (a^2 - 1)/a^2)*(x/a^2)^(1/2)*((x - 1)/(a^2 - 1))^(1/2)*(-(x - a^2)/(a^2 - 1))^(1/2))/(a^2*x - x^2*(a^2 + 1) + x^3)^(1/2)`**Reduce [F]**

$$\int \frac{a+x}{(-a+x)\sqrt{a^2x-(1+a^2)x^2+x^3}} dx$$

$$= \left(\int \frac{\sqrt{x}\sqrt{x-1}\sqrt{-a^2+x}}{a^3x^2-a^2x^3-a^3x+a^2x^2-ax^3+x^4+ax^2-x^3} dx \right) a$$

$$+ \int \frac{\sqrt{x}\sqrt{x-1}\sqrt{-a^2+x}}{a^3x-a^2x^2-a^3+a^2x-ax^2+x^3+ax-x^2} dx$$

input `int((a+x)/(-a+x)/(a^2*x-(a^2+1)*x^2+x^3)^(1/2),x)`

output

```
int((sqrt(x)*sqrt(x - 1)*sqrt(- a**2 + x))/(a**3*x**2 - a**3*x - a**2*x**
3 + a**2*x**2 - a*x**3 + a*x**2 + x**4 - x**3),x)*a + int((sqrt(x)*sqrt(x
- 1)*sqrt(- a**2 + x))/(a**3*x - a**3 - a**2*x**2 + a**2*x - a*x**2 + a*x
+ x**3 - x**2),x)
```

3.82
$$\int \frac{-2+a+x}{(-a+x)\sqrt{(2-a)ax+(-1-2a+a^2)x^2+x^3}} dx$$

Optimal result	634
Mathematica [A] (verified)	634
Rubi [C] (warning: unable to verify)	635
Maple [C] (verified)	638
Fricas [C] (verification not implemented)	638
Sympy [F]	639
Maxima [F]	639
Giac [F]	640
Mupad [B] (verification not implemented)	640
Reduce [F]	641

Optimal result

Integrand size = 40, antiderivative size = 71

$$\int \frac{-2+a+x}{(-a+x)\sqrt{(2-a)ax+(-1-2a+a^2)x^2+x^3}} dx$$

$$= \frac{\log\left(\frac{-a^2+2(-a+a^2)x+x^2-2a\sqrt{-((-2a+a^2)x+(-1-2a+a^2)x^2+x^3)}}{a^2-2ax+x^2}\right)}{a}$$

output 0

Mathematica [A] (verified)

Time = 10.36 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.65

$$\int \frac{-2+a+x}{(-a+x)\sqrt{(2-a)ax+(-1-2a+a^2)x^2+x^3}} dx$$

$$= -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{(2a-a^2)x+(-1-2a+a^2)x^2+x^3}}{a(-1+x)}\right)}{a}$$

input `Integrate[(-2 + a + x)/((-a + x)*Sqrt[(2 - a)*a*x + (-1 - 2*a + a^2)*x^2 + x^3]), x]`

```
output (-2*ArcTanh[Sqrt[(2*a - a^2)*x + (-1 - 2*a + a^2)*x^2 + x^3]/(a*(-1 + x))]
)/a
```

Rubi [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 1.17 (sec) , antiderivative size = 546, normalized size of antiderivative = 7.69, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2467, 2035, 2226, 1416, 2222}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{a + x - 2}{(x - a)\sqrt{(a^2 - 2a - 1)x^2 + (2 - a)ax + x^3}} dx$$

↓ 2467

$$\frac{\sqrt{x}\sqrt{-(-a^2 + 2a + 1)x + (2 - a)a + x^2} \int \frac{-a-x+2}{(a-x)\sqrt{x}\sqrt{x^2 - (-a^2 + 2a + 1)x + (2-a)a}} dx}{\sqrt{-(-a^2 + 2a + 1)x^2 + (2 - a)ax + x^3}}$$

↓ 2035

$$\frac{2\sqrt{x}\sqrt{-(-a^2 + 2a + 1)x + (2 - a)a + x^2} \int \frac{-a-x+2}{(a-x)\sqrt{x^2 - (-a^2 + 2a + 1)x + (2-a)a}} d\sqrt{x}}{\sqrt{-(-a^2 + 2a + 1)x^2 + (2 - a)ax + x^3}}$$

↓ 2226

$$\frac{2\sqrt{x}\sqrt{-(-a^2 + 2a + 1)x + (2 - a)a + x^2} \left(\frac{\sqrt{(2-a)a} \int \frac{1}{\sqrt{x^2 - (-a^2 + 2a + 1)x + (2-a)a}} d\sqrt{x}}{a} + (\sqrt{2-a} - \sqrt{a}) \sqrt{2-a} \int \frac{1}{\sqrt{x^2 - (-a^2 + 2a + 1)x + (2-a)a}} d\sqrt{x}}{\sqrt{-(-a^2 + 2a + 1)x^2 + (2 - a)ax + x^3}} \right)}{\sqrt{-(-a^2 + 2a + 1)x^2 + (2 - a)ax + x^3}}$$

↓ 1416

$$2\sqrt{x}\sqrt{-(-a^2 + 2a + 1)x + (2 - a)a + x^2} \left((\sqrt{2 - a} - \sqrt{a}) \sqrt{2 - a} \int \frac{\frac{x}{\sqrt{(2-a)a}} + 1}{(a-x)\sqrt{x^2 - (-a^2 + 2a + 1)x + (2-a)a}} d\sqrt{x} + \frac{((2-a))}{\sqrt{-(-a^2 + 2a + 1)x^2 + (2 - a)}}$$

↓ 2222

$$2\sqrt{x}\sqrt{-(-a^2 + 2a + 1)x + (2 - a)a + x^2} \left((\sqrt{2 - a} - \sqrt{a}) \sqrt{2 - a} \left(\frac{\sqrt[4]{(2 - a)a} \left(1 - \frac{a}{\sqrt{(2-a)a}}\right) \left(\frac{x}{\sqrt{(2-a)a}} + 1\right)}{\sqrt{\frac{-(-a^2 + 2a + 1)x^2 + (2 - a)}{(2 - a)}}} \right) \right)$$

input

```
Int[(-2 + a + x)/((-a + x)*Sqrt[(2 - a)*a*x + (-1 - 2*a + a^2)*x^2 + x^3]], x]
```

output

```
(2*Sqrt[x]*Sqrt[(2 - a)*a - (1 + 2*a - a^2)*x + x^2]*(((2 - a)*a)^(3/4)*(1 + x/Sqrt[(2 - a)*a])*Sqrt[((2 - a)*a - (1 + 2*a - a^2)*x + x^2]/((2 - a)*a*(1 + x/Sqrt[(2 - a)*a])^2)]*EllipticF[2*ArcTan[Sqrt[x]/((2 - a)*a)^(1/4)], (2 + (1 + 2*a - a^2)/Sqrt[(2 - a)*a])/4]/(2*a*Sqrt[(2 - a)*a - (1 + 2*a - a^2)*x + x^2]) + (Sqrt[2 - a] - Sqrt[a])*Sqrt[2 - a]*(((1 + a/Sqrt[(2 - a)*a])*ArcTanh[((1 - a)*Sqrt[x])/Sqrt[(2 - a)*a - (1 + 2*a - a^2)*x + x^2]])/(2*(1 - a)*a) + (((2 - a)*a)^(1/4)*(1 - a/Sqrt[(2 - a)*a])*(1 + x/Sqrt[(2 - a)*a])*Sqrt[((2 - a)*a - (1 + 2*a - a^2)*x + x^2]/((2 - a)*a*(1 + x/Sqrt[(2 - a)*a])^2)]*EllipticPi[(2*a - a^2 + Sqrt[(2 - a)*a])/((2*(2 - a)*a), 2*ArcTan[Sqrt[x]/((2 - a)*a)^(1/4)], (2 + (1 + 2*a - a^2)/Sqrt[(2 - a)*a])/4])/((4*a*Sqrt[(2 - a)*a - (1 + 2*a - a^2)*x + x^2])))/Sqrt[(2 - a)*a*x - (1 + 2*a - a^2)*x^2 + x^3]
```

Definitions of rubi rules used

rule 1416 `Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]`

rule 2035 `Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x]] /; FractionQ[m] && AlgebraicFunctionQ[Fx, x]`

rule 2222 `Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(- (B*d - A*e)) * (ArcTanh[Rt[b - c*(d/e) - a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[b - c*(d/e) - a*(e/d), 2])), x] + Simp[(B*d + A*e)*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(4*d*e*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[-(e - d*q^2)^2/(4*d*e*q^2), 2*ArcTan[q*x], 1/2 - b/(4*a*q^2)], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0] && PosQ[B/A] && NegQ[-b + c*(d/e) + a*(e/d)]`

rule 2226 `Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Simp[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2) Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] + Simp[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)) Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]`

rule 2467 `Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[Px^FracPart[p]/(x^(r*FracPart[p]))*ExpandToSum[Px/x^r, x]^FracPart[p] Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x], x] /; IGtQ[r, 0]] /; FreeQ[p, x] && PolyQ[Px, x] && !IntegerQ[p] && !MonomialQ[Px, x] && !PolyQ[Fx, x]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 1.

Time = 0.76 (sec) , antiderivative size = 317, normalized size of antiderivative = 4.46

method	result
default	$\frac{2(2a-2)(a^2-2a)\sqrt{\frac{a^2-2a+x}{a^2-2a}}\sqrt{\frac{-1+x}{-a^2+2a-1}}\sqrt{\frac{x}{-a^2+2a}}\text{EllipticPi}\left(\sqrt{\frac{a^2-2a+x}{a^2-2a}},\frac{-a^2+2a}{-a^2+2a},\sqrt{\frac{-a^2+2a}{-a^2+2a-1}}\right)}{\sqrt{a^2x^2-a^2x-2ax^2+x^3+2ax-x^2}(-a^2+a)} + \frac{2(a^2-2a)\sqrt{\frac{a^2-2a+x}{a^2-2a}}}{\sqrt{a^2x^2-a^2x-2ax^2+x^3+2ax-x^2}(-a^2+a)}$
elliptic	$\frac{2(2a-2)(a^2-2a)\sqrt{\frac{a^2-2a+x}{a^2-2a}}\sqrt{\frac{-1+x}{-a^2+2a-1}}\sqrt{\frac{x}{-a^2+2a}}\text{EllipticPi}\left(\sqrt{\frac{a^2-2a+x}{a^2-2a}},\frac{-a^2+2a}{-a^2+2a},\sqrt{\frac{-a^2+2a}{-a^2+2a-1}}\right)}{\sqrt{a^2x^2-a^2x-2ax^2+x^3+2ax-x^2}(-a^2+a)} + \frac{2(a^2-2a)\sqrt{\frac{a^2-2a+x}{a^2-2a}}}{\sqrt{a^2x^2-a^2x-2ax^2+x^3+2ax-x^2}(-a^2+a)}$

input `int((-2+a+x)/(-a+x)/((2-a)*a*x+(a^2-2*a-1)*x^2+x^3)^(1/2),x,method=_RETURN
VERBOSE)`

output `2*(2*a-2)*(a^2-2*a)*((a^2-2*a+x)/(a^2-2*a))^(1/2)*((-1+x)/(-a^2+2*a-1))^(1/2)*
(x/(-a^2+2*a))^(1/2)/(a^2*x^2-a^2*x-2*a*x^2+x^3+2*a*x-x^2)^(1/2)/(-a^2+
a)*EllipticPi(((a^2-2*a+x)/(a^2-2*a))^(1/2),(-a^2+2*a)/(-a^2+a),((-a^2+2*
a)/(-a^2+2*a-1))^(1/2))+2*(a^2-2*a)*((a^2-2*a+x)/(a^2-2*a))^(1/2)*((-1+x)/
(-a^2+2*a-1))^(1/2)*(x/(-a^2+2*a))^(1/2)/(a^2*x^2-a^2*x-2*a*x^2+x^3+2*a*x-
x^2)^(1/2)*EllipticF(((a^2-2*a+x)/(a^2-2*a))^(1/2),((-a^2+2*a)/(-a^2+2*a-1
))^(1/2))`

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 3 vs. order 1.

Time = 0.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.99

$$\int \frac{-2+a+x}{(-a+x)\sqrt{(2-a)ax+(-1-2a+a^2)x^2+x^3}} dx$$

$$= \frac{\log\left(-\frac{a^2-2(a^2-a)x-x^2+2\sqrt{(a^2-2a-1)x^2+x^3-(a^2-2a)xa}}{a^2-2ax+x^2}\right)}{a}$$

input `integrate((-2+a+x)/(-a+x)/((2-a)*a*x+(a^2-2*a-1)*x^2+x^3)^(1/2),x, algorit
hm="fricas")`

output $\log(-a^2 - 2(a^2 - a)x - x^2 + 2\sqrt{(a^2 - 2a - 1)x^2 + x^3 - (a^2 - 2a)x})a / (a^2 - 2ax + x^2) / a$

Sympy [F]

$$\int \frac{-2 + a + x}{(-a + x)\sqrt{(2 - a)ax + (-1 - 2a + a^2)x^2 + x^3}} dx$$

$$= \int \frac{a + x - 2}{\sqrt{x(x - 1)(a^2 - 2a + x)}(-a + x)} dx$$

input `integrate((-2+a+x)/(-a+x)/((2-a)*a*x+(a**2-2*a-1)*x**2+x**3)**(1/2),x)`

output `Integral((a + x - 2)/(sqrt(x*(x - 1)*(a**2 - 2*a + x))*(-a + x)), x)`

Maxima [F]

$$\int \frac{-2 + a + x}{(-a + x)\sqrt{(2 - a)ax + (-1 - 2a + a^2)x^2 + x^3}} dx$$

$$= \int -\frac{a + x - 2}{\sqrt{-(a - 2)ax + (a^2 - 2a - 1)x^2 + x^3}(a - x)} dx$$

input `integrate((-2+a+x)/(-a+x)/((2-a)*a*x+(a^2-2*a-1)*x^2+x^3)^(1/2),x, algorithm="maxima")`

output `-integrate((a + x - 2)/(sqrt(-(a - 2)*a*x + (a^2 - 2*a - 1)*x^2 + x^3)*(a - x)), x)`

Giac [F]

$$\int \frac{-2 + a + x}{(-a + x)\sqrt{(2 - a)ax + (-1 - 2a + a^2)x^2 + x^3}} dx$$

$$= \int -\frac{a + x - 2}{\sqrt{-(a - 2)ax + (a^2 - 2a - 1)x^2 + x^3}(a - x)} dx$$

input `integrate((-2+a+x)/(-a+x)/((2-a)*a*x+(a^2-2*a-1)*x^2+x^3)^(1/2),x, algorithm="giac")`

output `integrate(-(a + x - 2)/(sqrt(-(a - 2)*a*x + (a^2 - 2*a - 1)*x^2 + x^3)*(a - x)), x)`

Mupad [B] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.92

$$\int \frac{-2 + a + x}{(-a + x)\sqrt{(2 - a)ax + (-1 - 2a + a^2)x^2 + x^3}} dx$$

$$= \frac{2 \sqrt{\frac{x}{2a - a^2}} \sqrt{-\frac{x-1}{a^2 - 2a + 1}} (a - 1)^2 \sqrt{\frac{a^2 - 2a + x}{a^2 - 2a + 1}} \left(a F\left(\operatorname{asin}\left(\sqrt{\frac{a^2 - 2a + x}{a^2 - 2a + 1}}\right) \middle| -\frac{a^2 - 2a + 1}{2a - a^2}\right) - 2 \Pi\left(-\frac{a^2 - 2a + 1}{a - a^2}; \operatorname{asin}\left(\sqrt{\frac{a^2 - 2a + x}{a^2 - 2a + 1}}\right)\right) \right)}{a \sqrt{x^3 + (a^2 - 2a - 1)x^2 + (2a - a^2)x}}$$

input `int(-(a + x - 2)/((a - x)*(x^3 - x^2*(2*a - a^2 + 1) - a*x*(a - 2))^(1/2)),x)`

output `(2*(x/(2*a - a^2))^(1/2)*(-(x - 1)/(a^2 - 2*a + 1))^(1/2)*(a - 1)^2*((x - 2*a + a^2)/(a^2 - 2*a + 1))^(1/2)*(a*ellipticF(asin(((x - 2*a + a^2)/(a^2 - 2*a + 1))^(1/2))), -(a^2 - 2*a + 1)/(2*a - a^2)) - 2*ellipticPi(-(a^2 - 2*a + 1)/(a - a^2), asin(((x - 2*a + a^2)/(a^2 - 2*a + 1))^(1/2))), -(a^2 - 2*a + 1)/(2*a - a^2)))/(a*(x*(2*a - a^2) - x^2*(2*a - a^2 + 1) + x^3)^(1/2))`

Reduce [F]

$$\int \frac{-2 + a + x}{(-a + x)\sqrt{(2 - a)ax + (-1 - 2a + a^2)x^2 + x^3}} dx$$

$$= - \left(\int \frac{\sqrt{x} \sqrt{x - 1} \sqrt{a^2 - 2a + x}}{a^3 x^2 - a^2 x^3 - a^3 x - a^2 x^2 + 3a x^3 - x^4 + 2a^2 x - 3a x^2 + x^3} dx \right) a$$

$$+ 2 \left(\int \frac{\sqrt{x} \sqrt{x - 1} \sqrt{a^2 - 2a + x}}{a^3 x^2 - a^2 x^3 - a^3 x - a^2 x^2 + 3a x^3 - x^4 + 2a^2 x - 3a x^2 + x^3} dx \right)$$

$$- \left(\int \frac{\sqrt{x} \sqrt{x - 1} \sqrt{a^2 - 2a + x}}{a^3 x - a^2 x^2 - a^3 - a^2 x + 3a x^2 - x^3 + 2a^2 - 3a x + x^2} dx \right)$$

input `int((-2+a+x)/(-a+x)/((2-a)*a*x+(a^2-2*a-1)*x^2+x^3)^(1/2),x)`

output `- int((sqrt(x)*sqrt(x - 1)*sqrt(a**2 - 2*a + x))/(a**3*x**2 - a**3*x - a**2*x**3 - a**2*x**2 + 2*a**2*x + 3*a*x**3 - 3*a*x**2 - x**4 + x**3),x)*a + 2*int((sqrt(x)*sqrt(x - 1)*sqrt(a**2 - 2*a + x))/(a**3*x**2 - a**3*x - a**2*x**3 - a**2*x**2 + 2*a**2*x + 3*a*x**3 - 3*a*x**2 - x**4 + x**3),x) - int((sqrt(x)*sqrt(x - 1)*sqrt(a**2 - 2*a + x))/(a**3*x - a**3 - a**2*x**2 - a**2*x + 2*a**2 + 3*a*x**2 - 3*a*x - x**3 + x**2),x)`

3.83
$$\int \frac{-a+(-1+2a)x}{(-a+x)\sqrt{a^2x - (-1+2a+a^2)x^2 + (-1+2a)x^3}} dx$$

Optimal result	642
Mathematica [A] (verified)	642
Rubi [C] (verified)	643
Maple [C] (verified)	646
Fricas [A] (verification not implemented)	647
Sympy [F]	647
Maxima [F]	648
Giac [F]	648
Mupad [F(-1)]	649
Reduce [F]	649

Optimal result

Integrand size = 51, antiderivative size = 46

$$\int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx$$

$$= \log \left(\frac{-a^2 + 2ax + x^2 - 2\left(x + \sqrt{(1-x)x(a^2 + x - 2ax)}\right)}{(a-x)^2} \right)$$

output `ln((-a^2+2*a*x+x^2-2*x-2*((1-x)*x*(a^2-2*a*x+x))^(1/2))/(a-x)^2)`

Mathematica [A] (verified)

Time = 10.85 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.13

$$\int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx$$

$$= -2\operatorname{arctanh} \left(\frac{\sqrt{a^2x + (1 - 2a - a^2)x^2 + (-1 + 2a)x^3}}{-a^2 + (-1 + 2a)x} \right)$$

input

```
Integrate[(-a + (-1 + 2*a)*x)/((-a + x)*Sqrt[a^2*x - (-1 + 2*a + a^2)*x^2 + (-1 + 2*a)*x^3]),x]
```

output

```
-2*ArcTanh[Sqrt[a^2*x + (1 - 2*a - a^2)*x^2 + (-1 + 2*a)*x^3]/(-a^2 + (-1 + 2*a)*x)]
```

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 0.83 (sec) , antiderivative size = 233, normalized size of antiderivative = 5.07, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.137$, Rules used = {2467, 2035, 2228, 1417, 321, 1544, 412}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(2a-1)x-a}{(x-a)\sqrt{-(a^2+2a-1)x^2+a^2x+(2a-1)x^3}} dx$$

$$\downarrow 2467$$

$$\frac{\sqrt{x}\sqrt{(-a^2-2a+1)x+a^2-(1-2a)x^2} \int \frac{a+(1-2a)x}{(a-x)\sqrt{x}\sqrt{a^2-(1-2a)x^2+(-a^2-2a+1)x}} dx}{\sqrt{(-a^2-2a+1)x^2+a^2x-((1-2a)x^3)}}$$

$$\downarrow 2035$$

$$\frac{2\sqrt{x}\sqrt{(-a^2-2a+1)x+a^2-(1-2a)x^2} \int \frac{a+(1-2a)x}{(a-x)\sqrt{a^2-(1-2a)x^2+(-a^2-2a+1)x}} d\sqrt{x}}{\sqrt{(-a^2-2a+1)x^2+a^2x-((1-2a)x^3)}}$$

$$\downarrow 2228$$

$$\frac{2\sqrt{x}\sqrt{(-a^2-2a+1)x+a^2-(1-2a)x^2} \left(2(1-a)a \int \frac{1}{(a-x)\sqrt{a^2-(1-2a)x^2+(-a^2-2a+1)x}} d\sqrt{x} - (1-2a) \int \frac{1}{\sqrt{a^2-(1-2a)x^2+(-a^2-2a+1)x}} d\sqrt{x} \right)}{\sqrt{(-a^2-2a+1)x^2+a^2x-((1-2a)x^3)}}$$

$$\downarrow 1417$$

$$\frac{2\sqrt{x}\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2} \left(2(1 - a)a \int \frac{1}{(a-x)\sqrt{a^2 - (1-2a)x^2 + (-a^2 - 2a + 1)x}} d\sqrt{x} - \frac{(1-2a)\sqrt{1-x}\sqrt{\frac{(1-2a)}{a^2}}}{\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2}} \right)}{\sqrt{(-a^2 - 2a + 1)x^2 + a^2x - ((1 - 2a)x^3)}}$$

↓ 321

$$\frac{2\sqrt{x}\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2} \left(2(1 - a)a \int \frac{1}{(a-x)\sqrt{a^2 - (1-2a)x^2 + (-a^2 - 2a + 1)x}} d\sqrt{x} - \frac{(1-2a)\sqrt{1-x}\sqrt{\frac{(1-2a)}{a^2}}}{\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2}} \right)}{\sqrt{(-a^2 - 2a + 1)x^2 + a^2x - ((1 - 2a)x^3)}}$$

↓ 1544

$$\frac{2\sqrt{x}\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2} \left(\frac{2(1-a)a\sqrt{1-x}\sqrt{\frac{(1-2a)x}{a^2} + 1} \int \frac{1}{\sqrt{1-x(a-x)}\sqrt{\frac{(1-2a)x}{a^2} + 1}} d\sqrt{x}}{\sqrt{(-a^2 - 2a + 1)x + a^2 - (1-2a)x^2}} - \frac{(1-2a)\sqrt{1-x}\sqrt{\frac{(1-2a)}{a^2}}}{\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2}} \right)}{\sqrt{(-a^2 - 2a + 1)x^2 + a^2x - ((1 - 2a)x^3)}}$$

↓ 412

$$\frac{2\sqrt{x}\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2} \left(\frac{2(1-a)\sqrt{1-x}\sqrt{\frac{(1-2a)x}{a^2} + 1} \operatorname{EllipticPi}\left(\frac{1}{a}, \arcsin(\sqrt{x}), -\frac{1-2a}{a^2}\right)}{\sqrt{(-a^2 - 2a + 1)x + a^2 - (1-2a)x^2}} - \frac{(1-2a)\sqrt{1-x}\sqrt{\frac{(1-2a)}{a^2}}}{\sqrt{(-a^2 - 2a + 1)x + a^2 - (1 - 2a)x^2}} \right)}{\sqrt{(-a^2 - 2a + 1)x^2 + a^2x - ((1 - 2a)x^3)}}$$

input `Int[(-a + (-1 + 2*a)*x)/((-a + x)*Sqrt[a^2*x - (-1 + 2*a + a^2)*x^2 + (-1 + 2*a)*x^3], x]`

output `(2*Sqrt[x]*Sqrt[a^2 + (1 - 2*a - a^2)*x - (1 - 2*a)*x^2]*(-(((1 - 2*a)*Sqrt[1 - x]*Sqrt[1 + ((1 - 2*a)*x)/a^2]*EllipticF[ArcSin[Sqrt[x]], -((1 - 2*a)/a^2)]))/Sqrt[a^2 + (1 - 2*a - a^2)*x - (1 - 2*a)*x^2]) + (2*(1 - a)*Sqrt[1 - x]*Sqrt[1 + ((1 - 2*a)*x)/a^2]*EllipticPi[a^(-1), ArcSin[Sqrt[x]], -((1 - 2*a)/a^2)]))/Sqrt[a^2 + (1 - 2*a - a^2)*x - (1 - 2*a)*x^2])/Sqrt[a^2*x + (1 - 2*a - a^2)*x^2 - (1 - 2*a)*x^3]`

Definitions of rubi rules used

rule 321

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)^2]*Sqrt[(c_) + (d_.)*(x_)^2]), x_Symbol] := S
imp[(1/(Sqrt[a]*Sqrt[c]*Rt[-d/c, 2]))*EllipticF[ArcSin[Rt[-d/c, 2]*x], b*(c
/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && NegQ[d/c] && GtQ[c, 0] && GtQ[a,
0] && !(NegQ[b/a] && SimplerSqrtQ[-b/a, -d/c])
```

rule 412

```
Int[1/(((a_) + (b_.)*(x_)^2)*Sqrt[(c_) + (d_.)*(x_)^2]*Sqrt[(e_) + (f_.)*(x
_)^2]), x_Symbol] := Simp[(1/(a*Sqrt[c]*Sqrt[e]*Rt[-d/c, 2]))*EllipticPi[b*
(c/(a*d)), ArcSin[Rt[-d/c, 2]*x], c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e,
f}, x] && !GtQ[d/c, 0] && GtQ[c, 0] && GtQ[e, 0] && !( !GtQ[f/e, 0] && S
implerSqrtQ[-f/e, -d/c])
```

rule 1417

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[b
^2 - 4*a*c, 2]}, Simp[Sqrt[1 + 2*c*(x^2/(b - q))]*(Sqrt[1 + 2*c*(x^2/(b + q
))]/Sqrt[a + b*x^2 + c*x^4]) Int[1/(Sqrt[1 + 2*c*(x^2/(b - q))]*Sqrt[1 +
2*c*(x^2/(b + q))]), x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
&& NegQ[c/a]
```

rule 1544

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_S
ymbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[Sqrt[1 + 2*c*(x^2/(b - q))]*(
Sqrt[1 + 2*c*(x^2/(b + q))]/Sqrt[a + b*x^2 + c*x^4]) Int[1/((d + e*x^2)*S
qrt[1 + 2*c*(x^2/(b - q))]*Sqrt[1 + 2*c*(x^2/(b + q))]), x], x]] /; FreeQ[{
a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0]
&& NegQ[c/a]
```

rule 2035

```
Int[(Fx_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Simp[k Subst
[Int[x^(k*(m + 1) - 1)*SubstPower[Fx, x, k], x], x, x^(1/k)], x]] /; Fracti
onQ[m] && AlgebraicFunctionQ[Fx, x]
```

rule 2228

```
Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 +
(c_.)*(x_)^4]), x_Symbol] := Simp[B/e Int[1/Sqrt[a + b*x^2 + c*x^4], x],
x] + Simp[(e*A - d*B)/e Int[1/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x],
x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[c*d^2 - a*e^2, 0] && NegQ[c/a
]
```

rule 2467

```
Int[(Fx_.)*(Px_)^(p_), x_Symbol] := With[{r = Expon[Px, x, Min]}, Simp[Px^FracPart[p]/(x^(r*FracPart[p]))*ExpandToSum[Px/x^r, x]^FracPart[p]] Int[x^(p*r)*ExpandToSum[Px/x^r, x]^p*Fx, x], x] /; IGtQ[r, 0]] /; FreeQ[p, x] && PolyQ[Px, x] && !IntegerQ[p] && !MonomialQ[Px, x] && !PolyQ[Fx, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 0.92 (sec) , antiderivative size = 367, normalized size of antiderivative = 7.98

method	result
elliptic	$\frac{2a^2 \sqrt{-\frac{(x - \frac{a^2}{-1+2a})(-1+2a)}{a^2}} \sqrt{\frac{-1+x}{\frac{a^2}{-1+2a}-1}} \sqrt{\frac{(-1+2a)x}{a^2}} \operatorname{EllipticF}\left(\sqrt{-\frac{(x - \frac{a^2}{-1+2a})(-1+2a)}{a^2}}, \sqrt{\frac{a^2}{(-1+2a)(\frac{a^2}{-1+2a}-1)}}\right)}{\sqrt{-a^2x^2+2ax^3+a^2x-2ax^2-x^3+x^2}} - \frac{4a^3}{\dots}$
default	$\frac{2a^2 \sqrt{-\frac{(x - \frac{a^2}{-1+2a})(-1+2a)}{a^2}} \sqrt{\frac{-1+x}{\frac{a^2}{-1+2a}-1}} \sqrt{\frac{(-1+2a)x}{a^2}} \operatorname{EllipticF}\left(\sqrt{-\frac{(x - \frac{a^2}{-1+2a})(-1+2a)}{a^2}}, \sqrt{\frac{a^2}{(-1+2a)(\frac{a^2}{-1+2a}-1)}}\right)}{(-1+2a)\sqrt{-a^2x^2+2ax^3+a^2x-2ax^2-x^3+x^2}} - \frac{4a^3}{\dots}$

input

```
int((-a+(-1+2*a)*x)/(-a+x)/(a^2*x-(a^2+2*a-1)*x^2+(-1+2*a)*x^3)^(1/2), x, method=_RETURNVERBOSE)
```

output

```
-2*a^2*(-(x-a^2/(-1+2*a))/a^2*(-1+2*a))^(1/2)*((-1+x)/(a^2/(-1+2*a)-1))^(1/2)*(1/a^2*(-1+2*a)*x)^(1/2)/(-a^2*x^2+2*a*x^3+a^2*x-2*a*x^2-x^3+x^2)^(1/2)*EllipticF(-(x-a^2/(-1+2*a))/a^2*(-1+2*a))^(1/2),(a^2/(-1+2*a)/(a^2/(-1+2*a)-1))^(1/2))-4*a^3*(a-1)/(-1+2*a)*(-(x-a^2/(-1+2*a))/a^2*(-1+2*a))^(1/2)*((-1+x)/(a^2/(-1+2*a)-1))^(1/2)*(1/a^2*(-1+2*a)*x)^(1/2)/(-a^2*x^2+2*a*x^3+a^2*x-2*a*x^2-x^3+x^2)^(1/2)/(a^2/(-1+2*a)-a)*EllipticPi(-(x-a^2/(-1+2*a))/a^2*(-1+2*a))^(1/2),a^2/(-1+2*a)/(a^2/(-1+2*a)-a),(a^2/(-1+2*a)/(a^2/(-1+2*a)-1))^(1/2))
```

Fricas [A] (verification not implemented)

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.37

$$\int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx$$

$$= \log\left(-\frac{a^2 - 2(a - 1)x - x^2 + 2\sqrt{(2a - 1)x^3 + a^2x - (a^2 + 2a - 1)x^2}}{a^2 - 2ax + x^2}\right)$$

input `integrate((-a+(-1+2*a)*x)/(-a+x)/(a^2*x-(a^2+2*a-1)*x^2+(-1+2*a)*x^3)^(1/2),x, algorithm="fricas")`

output `log(-(a^2 - 2*(a - 1)*x - x^2 + 2*sqrt((2*a - 1)*x^3 + a^2*x - (a^2 + 2*a - 1)*x^2))/(a^2 - 2*a*x + x^2))`

Sympy [F]

$$\int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx$$

$$= \int \frac{2ax - a - x}{\sqrt{x(x - 1)}(-a^2 + 2ax - x)(-a + x)} dx$$

input `integrate((-a+(-1+2*a)*x)/(-a+x)/(a**2*x-(a**2+2*a-1)*x**2+(-1+2*a)*x**3)*(1/2),x)`

output `Integral((2*a*x - a - x)/(sqrt(x*(x - 1))*(-a**2 + 2*a*x - x))*(-a + x), x)`

Maxima [F]

$$\int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx$$

$$= \int -\frac{(2a - 1)x - a}{\sqrt{(2a - 1)x^3 + a^2x - (a^2 + 2a - 1)x^2(a - x)}} dx$$

input `integrate((-a+(-1+2*a)*x)/(-a+x)/(a^2*x-(a^2+2*a-1)*x^2+(-1+2*a)*x^3)^(1/2),x, algorithm="maxima")`

output `-integrate(((2*a - 1)*x - a)/(sqrt((2*a - 1)*x^3 + a^2*x - (a^2 + 2*a - 1)*x^2)*(a - x)), x)`

Giac [F]

$$\int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx$$

$$= \int -\frac{(2a - 1)x - a}{\sqrt{(2a - 1)x^3 + a^2x - (a^2 + 2a - 1)x^2(a - x)}} dx$$

input `integrate((-a+(-1+2*a)*x)/(-a+x)/(a^2*x-(a^2+2*a-1)*x^2+(-1+2*a)*x^3)^(1/2),x, algorithm="giac")`

output `integrate(-((2*a - 1)*x - a)/(sqrt((2*a - 1)*x^3 + a^2*x - (a^2 + 2*a - 1)*x^2)*(a - x)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx = \text{Hanged}$$

input

```
int((a - x*(2*a - 1))/((a - x)*(x^3*(2*a - 1) - x^2*(2*a + a^2 - 1) + a^2*x)^(1/2)),x)
```

output

```
\text{Hanged}
```

Reduce [F]

$$\begin{aligned} & \int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (-1 + 2a + a^2)x^2 + (-1 + 2a)x^3}} dx \\ &= \int \frac{-a + (-1 + 2a)x}{(-a + x)\sqrt{a^2x - (a^2 + 2a - 1)x^2 + (-1 + 2a)x^3}} dx \end{aligned}$$

input

```
int((-a+(-1+2*a)*x)/(-a+x)/(a^2*x-(a^2+2*a-1)*x^2+(-1+2*a)*x^3)^(1/2),x)
```

output

```
int((-a+(-1+2*a)*x)/(-a+x)/(a^2*x-(a^2+2*a-1)*x^2+(-1+2*a)*x^3)^(1/2),x)
```

3.84
$$\int \frac{1 - \sqrt[3]{2x}}{(2^{2/3} + x)\sqrt{1+x^3}} dx$$

Optimal result	650
Mathematica [A] (verified)	650
Rubi [A] (verified)	651
Maple [C] (verified)	652
Fricas [F(-2)]	653
Sympy [F]	653
Maxima [F]	653
Giac [F]	654
Mupad [B] (verification not implemented)	654
Reduce [F]	654

Optimal result

Integrand size = 29, antiderivative size = 32

$$\int \frac{1 - \sqrt[3]{2x}}{(2^{2/3} + x)\sqrt{1+x^3}} dx = \frac{2 \arctan\left(\frac{\sqrt{3}(1 + \sqrt[3]{2x})}{\sqrt{1+x^3}}\right)}{\sqrt{3}}$$

output

```
2/3*arctan((1+2^(1/3)*x)*3^(1/2)/(x^3+1)^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 1.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.06

$$\int \frac{1 - \sqrt[3]{2x}}{(2^{2/3} + x)\sqrt{1+x^3}} dx = -\frac{2 \arctan\left(\frac{\sqrt{1+x^3}}{\sqrt{3}(1 + \sqrt[3]{2x})}\right)}{\sqrt{3}}$$

input

```
Integrate[(1 - 2^(1/3)*x)/((2^(2/3) + x)*Sqrt[1 + x^3]),x]
```

output $(-2*\text{ArcTan}[\text{Sqrt}[1 + x^3]/(\text{Sqrt}[3]*(1 + 2^{(1/3)*x}))])/\text{Sqrt}[3]$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.069$, Rules used = {2562, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1 - \sqrt[3]{2}x}{(x + 2^{2/3})\sqrt{x^3 + 1}} dx$$

↓ 2562

$$2 \int \frac{1}{\frac{3(\sqrt[3]{2}x+1)^2}{x^3+1} + 1} d\frac{\sqrt[3]{2}x+1}{\sqrt{x^3+1}}$$

↓ 216

$$\frac{2 \arctan\left(\frac{\sqrt{3}(\sqrt[3]{2}x+1)}{\sqrt{x^3+1}}\right)}{\sqrt{3}}$$

input $\text{Int}[(1 - 2^{(1/3)*x})/((2^{(2/3)} + x)*\text{Sqrt}[1 + x^3]),x]$

output $(2*\text{ArcTan}[(\text{Sqrt}[3]*(1 + 2^{(1/3)*x}))/\text{Sqrt}[1 + x^3]])/\text{Sqrt}[3]$

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 2562

```
Int(((e_) + (f_)*(x_))/(((c_) + (d_)*(x_))*Sqrt[(a_) + (b_)*(x_)^3]), x_
Symbol] := Simp[2*(e/d) Subst[Int[1/(1 + 3*a*x^2), x], x, (1 + 2*d*(x/c))
/Sqrt[a + b*x^3]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[d*e - c*f, 0]
&& EqQ[b*c^3 - 4*a*d^3, 0] && EqQ[2*d*e + c*f, 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 3.58 (sec) , antiderivative size = 112, normalized size of antiderivative = 3.50

method	result
trager	$2^{\frac{1}{3}} \text{RootOf}(-Z^2 + 6 \cdot 2^{\frac{1}{3}}) \ln \left(\frac{3 \cdot 2^{\frac{2}{3}} x^2 \text{RootOf}(-Z^2 + 6 \cdot 2^{\frac{1}{3}}) - 12\sqrt{x^3+1}x - \text{RootOf}(-Z^2 + 6 \cdot 2^{\frac{1}{3}})x^3 + 6 \text{RootOf}(-Z^2 + 6 \cdot 2^{\frac{1}{3}})2^{\frac{1}{3}}x - 6\sqrt{x^3+1}}{(2^{\frac{1}{3}}x+2)^3} \right)$
default	$- \frac{2 \cdot 2^{\frac{1}{3}} \left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}}}{\sqrt{x^3+1}} \text{EllipticF}\left(\sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right) + \frac{6 \left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}}{\sqrt{x^3+1}}$
elliptic	$- \frac{2 \cdot 2^{\frac{1}{3}} \left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}}}{\sqrt{x^3+1}} \text{EllipticF}\left(\sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}}, \sqrt{\frac{-\frac{3}{2} + \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}\right) + \frac{6 \left(\frac{3}{2} - \frac{i\sqrt{3}}{2}\right) \sqrt{\frac{1+x}{\frac{3}{2} - \frac{i\sqrt{3}}{2}}} \sqrt{\frac{x - \frac{1}{2} - \frac{i\sqrt{3}}{2}}{-\frac{3}{2} - \frac{i\sqrt{3}}{2}}}}{\sqrt{x^3+1}}$

input

```
int((1-2^(1/3)*x)/(2^(2/3)+x)/(x^3+1)^(1/2), x, method=_RETURNVERBOSE)
```

output

```
1/6*2^(1/3)*RootOf(-Z^2+6*2^(1/3))*ln(-(3*2^(2/3)*x^2*RootOf(-Z^2+6*2^(1/3)
))-12*(x^3+1)^(1/2)*x-RootOf(-Z^2+6*2^(1/3))*x^3+6*RootOf(-Z^2+6*2^(1/3))
2^(1/3)*x-6*(x^3+1)^(1/2)*2^(2/3)+2*RootOf(-Z^2+6*2^(1/3)))/(2^(1/3)*x+2)
^3)
```

Fricas [F(-2)]

Exception generated.

$$\int \frac{1 - \sqrt[3]{2}x}{(2^{2/3} + x)\sqrt{1 + x^3}} dx = \text{Exception raised: TypeError}$$

input `integrate((1-2^(1/3)*x)/(2^(2/3)+x)/(x^3+1)^(1/2),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: catd ef: division by zero`

Sympy [F]

$$\int \frac{1 - \sqrt[3]{2}x}{(2^{2/3} + x)\sqrt{1 + x^3}} dx = - \int \frac{\sqrt[3]{2}x}{x\sqrt{x^3 + 1} + 2^{2/3}\sqrt{x^3 + 1}} dx - \int \left(-\frac{1}{x\sqrt{x^3 + 1} + 2^{2/3}\sqrt{x^3 + 1}} \right) dx$$

input `integrate((1-2**(1/3)*x)/(2**(2/3)+x)/(x**3+1)**(1/2),x)`

output `-Integral(2**(1/3)*x/(x*sqrt(x**3 + 1) + 2**(2/3)*sqrt(x**3 + 1)), x) - Integral(-1/(x*sqrt(x**3 + 1) + 2**(2/3)*sqrt(x**3 + 1)), x)`

Maxima [F]

$$\int \frac{1 - \sqrt[3]{2}x}{(2^{2/3} + x)\sqrt{1 + x^3}} dx = \int -\frac{2^{1/3}x - 1}{\sqrt{x^3 + 1}(x + 2^{2/3})} dx$$

input `integrate((1-2^(1/3)*x)/(2^(2/3)+x)/(x^3+1)^(1/2),x, algorithm="maxima")`

output `-integrate((2^(1/3)*x - 1)/(sqrt(x^3 + 1)*(x + 2^(2/3))), x)`

Giac [F]

$$\int \frac{1 - \sqrt[3]{2}x}{(2^{2/3} + x)\sqrt{1+x^3}} dx = \int -\frac{2^{1/3}x - 1}{\sqrt{x^3+1}(x+2^{2/3})} dx$$

input `integrate((1-2^(1/3)*x)/(2^(2/3)+x)/(x^3+1)^(1/2),x, algorithm="giac")`

output `integrate(-(2^(1/3)*x - 1)/(sqrt(x^3 + 1)*(x + 2^(2/3))), x)`

Mupad [B] (verification not implemented)

Time = 0.98 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.09

$$\int \frac{1 - \sqrt[3]{2}x}{(2^{2/3} + x)\sqrt{1+x^3}} dx = \frac{\sqrt{3} \ln \left(\frac{(\sqrt{3} \operatorname{li} + \sqrt{x^3+1} + 2^{1/3} \sqrt{3} x \operatorname{li}) (\sqrt{3} \operatorname{li} - \sqrt{x^3+1} + 2^{1/3} \sqrt{3} x \operatorname{li})^3}{(x+2^{2/3})^6} \right) \operatorname{li}}{3}$$

input `int(-(2^(1/3)*x - 1)/((x^3 + 1)^(1/2)*(x + 2^(2/3))),x)`

output `(3^(1/2)*log(((3^(1/2)*li + (x^3 + 1)^(1/2) + 2^(1/3)*3^(1/2)*x*li)*(3^(1/2)*li - (x^3 + 1)^(1/2) + 2^(1/3)*3^(1/2)*x*li)^3)/(x + 2^(2/3))^6)*li)/3`

Reduce [F]

$$\int \frac{1 - \sqrt[3]{2}x}{(2^{2/3} + x)\sqrt{1+x^3}} dx = -2^{1/3} \left(\int \frac{x}{\sqrt{x^3+1} 2^{2/3} + \sqrt{x^3+1} x} dx \right) + \int \frac{1}{\sqrt{x^3+1} 2^{2/3} + \sqrt{x^3+1} x} dx$$

input `int((1-2^(1/3)*x)/(2^(2/3)+x)/(x^3+1)^(1/2),x)`

output

```
- 2**(1/3)*int(x/(sqrt(x**3 + 1)*2**(2/3) + sqrt(x**3 + 1)*x),x) + int(1/  
(sqrt(x**3 + 1)*2**(2/3) + sqrt(x**3 + 1)*x),x)
```


$$3.85 \quad \int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx$$

Optimal result	656
Mathematica [A] (verified)	656
Rubi [A] (verified)	657
Maple [B] (verified)	658
Fricas [B] (verification not implemented)	658
Sympy [F]	659
Maxima [F]	659
Giac [F]	659
Mupad [B] (verification not implemented)	660
Reduce [F]	660

Optimal result

Integrand size = 18, antiderivative size = 23

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx = -\frac{2}{3} \operatorname{arctanh}\left(\frac{(1+x)^2}{3\sqrt{1+x^3}}\right)$$

output `-2/3*arctanh(1/3*(1+x)^2/(x^3+1)^(1/2))`

Mathematica [A] (verified)

Time = 0.87 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.35

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx = -\frac{2}{3} \operatorname{arctanh}\left(\frac{\frac{1}{3} + \frac{2x}{3} + \frac{x^2}{3}}{\sqrt{1+x^3}}\right)$$

input `Integrate[(1 + x)/((-2 + x)*Sqrt[1 + x^3]),x]`

output `(-2*ArcTanh[(1/3 + (2*x)/3 + x^2/3)/Sqrt[1 + x^3]])/3`

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$, Rules used = {2563, 219}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+1}{(x-2)\sqrt{x^3+1}} dx$$

↓ 2563

$$-2 \int \frac{1}{9 - \frac{(x+1)^4}{x^3+1}} d \frac{(x+1)^2}{\sqrt{x^3+1}}$$

↓ 219

$$-\frac{2}{3} \operatorname{arctanh} \left(\frac{(x+1)^2}{3\sqrt{x^3+1}} \right)$$

input `Int[(1 + x)/((-2 + x)*Sqrt[1 + x^3]),x]`

output `(-2*ArcTanh[(1 + x)^2/(3*Sqrt[1 + x^3])])/3`

Defintions of rubi rules used

rule 219 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

rule 2563 `Int[((e_) + (f_.)*(x_))/(((c_) + (d_.)*(x_))*Sqrt[(a_) + (b_.)*(x_)^3]), x_Symbol] :> Simp[-2*(e/d) Subst[Int[1/(9 - a*x^2), x], x, (1 + f*(x/e))^2/Sqrt[a + b*x^3]], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[d*e - c*f, 0] && EqQ[b*c^3 + 8*a*d^3, 0] && EqQ[2*d*e + c*f, 0]`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(17) = 34.

Time = 0.54 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.91

method	result
trager	$\frac{\ln\left(\frac{-x^3+6\sqrt{x^3+1}x-12x^2+6\sqrt{x^3+1}+6x-10}{(-2+x)^3}\right)}{3}$
default	$\frac{2\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\operatorname{EllipticF}\left(\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\sqrt{\frac{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)}{\sqrt{x^3+1}} - 2\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}}$
elliptic	$\frac{2\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\operatorname{EllipticF}\left(\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\sqrt{\frac{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)}{\sqrt{x^3+1}} - 2\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}}$

input `int((1+x)/(-2+x)/(x^3+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/3*ln((-x^3+6*(x^3+1)^(1/2)*x-12*x^2+6*(x^3+1)^(1/2)+6*x-10)/(-2+x)^3)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44 vs. 2(17) = 34.

Time = 0.08 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.91

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx = \frac{1}{3} \log \left(\frac{x^3 + 12x^2 - 6\sqrt{x^3+1}(x+1) - 6x + 10}{x^3 - 6x^2 + 12x - 8} \right)$$

input `integrate((1+x)/(-2+x)/(x^3+1)^(1/2),x, algorithm="fricas")`

output `1/3*log((x^3 + 12*x^2 - 6*sqrt(x^3 + 1)*(x + 1) - 6*x + 10)/(x^3 - 6*x^2 + 12*x - 8))`

Sympy [F]

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx = \int \frac{x+1}{\sqrt{(x+1)(x^2-x+1)}(x-2)} dx$$

input `integrate((1+x)/(-2+x)/(x**3+1)**(1/2),x)`

output `Integral((x + 1)/(sqrt((x + 1)*(x**2 - x + 1))*(x - 2)), x)`

Maxima [F]

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx = \int \frac{x+1}{\sqrt{x^3+1}(x-2)} dx$$

input `integrate((1+x)/(-2+x)/(x^3+1)^(1/2),x, algorithm="maxima")`

output `integrate((x + 1)/(sqrt(x^3 + 1)*(x - 2)), x)`

Giac [F]

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx = \int \frac{x+1}{\sqrt{x^3+1}(x-2)} dx$$

input `integrate((1+x)/(-2+x)/(x^3+1)^(1/2),x, algorithm="giac")`

output `integrate((x + 1)/(sqrt(x^3 + 1)*(x - 2)), x)`

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 204, normalized size of antiderivative = 8.87

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx$$

$$= \frac{(3 + \sqrt{3} i) \sqrt{\frac{x - \frac{1}{2} + \frac{\sqrt{3} i}{2}}{-\frac{3}{2} + \frac{\sqrt{3} i}{2}}} \left(F \left(\operatorname{asin} \left(\sqrt{\frac{x+1}{\frac{3}{2} + \frac{\sqrt{3} i}{2}}} \right) \middle| -\frac{\frac{3}{2} + \frac{\sqrt{3} i}{2}}{-\frac{3}{2} + \frac{\sqrt{3} i}{2}} \right) - \Pi \left(\frac{1}{2} + \frac{\sqrt{3} i}{6}; \operatorname{asin} \left(\sqrt{\frac{x+1}{\frac{3}{2} + \frac{\sqrt{3} i}{2}}} \right) \middle| -\frac{\frac{3}{2} + \frac{\sqrt{3} i}{2}}{-\frac{3}{2} + \frac{\sqrt{3} i}{2}} \right) \right)}{\sqrt{x^3 + \left(-\left(-\frac{1}{2} + \frac{\sqrt{3} i}{2} \right) \left(\frac{1}{2} + \frac{\sqrt{3} i}{2} \right) - 1 \right) x - \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2} \right) \left(\frac{1}{2} + \frac{\sqrt{3} i}{2} \right)}}$$

input `int((x + 1)/((x^3 + 1)^(1/2)*(x - 2)),x)`output `((3^(1/2)*1i + 3)*((x + (3^(1/2)*1i)/2 - 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2) * (ellipticF(asin(((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2)) - ellipticPi((3^(1/2)*1i)/6 + 1/2, asin(((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)), -((3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))) * ((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2) * (((3^(1/2)*1i)/2 - x + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1/2) / (x^3 - x*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) - ((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2))^(1/2)`**Reduce [F]**

$$\int \frac{1+x}{(-2+x)\sqrt{1+x^3}} dx = \int \frac{\sqrt{x^3+1}}{x^3-3x^2+3x-2} dx$$

input `int((1+x)/(-2+x)/(x^3+1)^(1/2),x)`output `int(sqrt(x**3 + 1)/(x**3 - 3*x**2 + 3*x - 2),x)`

3.86 $\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx$

Optimal result	661
Mathematica [C] (verified)	662
Rubi [A] (verified)	662
Maple [C] (verified)	663
Fricas [B] (verification not implemented)	664
Sympy [F]	665
Maxima [F]	665
Giac [F]	666
Mupad [F(-1)]	666
Reduce [F]	666

Optimal result

Integrand size = 25, antiderivative size = 218

$$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx = -\frac{(2-\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(1+x)}{\sqrt{2}\sqrt{1+x^3}}\right)}{2\sqrt{2}3^{3/4}} - \frac{(2-\sqrt{3}) \arctan\left(\frac{(1-\sqrt{3})\sqrt{1+x^3}}{\sqrt{2}3^{3/4}}\right)}{3\sqrt{2}3^{3/4}} - \frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1+\sqrt{3}-2x)}{\sqrt{2}\sqrt{1+x^3}}\right)}{3\sqrt{2}\sqrt[4]{3}} - \frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(1+x)}{\sqrt{2}\sqrt{1+x^3}}\right)}{6\sqrt{2}\sqrt[4]{3}}$$

output

```
-1/12*arctan(1/2*3^(1/4)*(1+x)*(1+3^(1/2))*2^(1/2)/(x^3+1)^(1/2))*(2-3^(1/2))*3^(1/4)*2^(1/2)-1/18*arctan(1/6*(1-3^(1/2))*(x^3+1)^(1/2)*3^(1/4)*2^(1/2))*(2-3^(1/2))*3^(1/4)*2^(1/2)-1/36*arctanh(1/2*3^(1/4)*(1+x)*(1-3^(1/2))*2^(1/2)/(x^3+1)^(1/2))*(2-3^(1/2))*3^(3/4)*2^(1/2)-1/18*arctanh(1/2*3^(1/4)*(1-2*x+3^(1/2))*2^(1/2)/(x^3+1)^(1/2))*(2-3^(1/2))*3^(3/4)*2^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.04 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.22

$$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx = \frac{x^2 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, -x^3, -\frac{x^3}{10+6\sqrt{3}}\right)}{20+12\sqrt{3}}$$

input

```
Integrate[x/(Sqrt[1 + x^3]*(10 + 6*Sqrt[3] + x^3)),x]
```

output

```
(x^2*AppellF1[2/3, 1/2, 1, 5/3, -x^3, -(x^3/(10 + 6*Sqrt[3]))])/(20 + 12*Sqrt[3])
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {989}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{x^3+1}(x^3+6\sqrt{3}+10)} dx$$

↓ 989

$$\frac{(2-\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(x+1)}{\sqrt{2}\sqrt{x^3+1}}\right)}{2\sqrt{2}3^{3/4}} - \frac{(2-\sqrt{3}) \arctan\left(\frac{(1-\sqrt{3})\sqrt{x^3+1}}{\sqrt{2}3^{3/4}}\right)}{3\sqrt{2}3^{3/4}}$$

$$\frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(-2x+\sqrt{3}+1)}{\sqrt{2}\sqrt{x^3+1}}\right)}{3\sqrt{2}\sqrt[4]{3}} - \frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(x+1)}{\sqrt{2}\sqrt{x^3+1}}\right)}{6\sqrt{2}\sqrt[4]{3}}$$

input

```
Int[x/(Sqrt[1 + x^3]*(10 + 6*Sqrt[3] + x^3)),x]
```

output

```
-1/2*((2 - Sqrt[3])*ArcTan[(3^(1/4)*(1 + Sqrt[3])*(1 + x))/(Sqrt[2]*Sqrt[1 + x^3]])/(Sqrt[2]*3^(3/4)) - ((2 - Sqrt[3])*ArcTan[((1 - Sqrt[3])*Sqrt[1 + x^3])/(Sqrt[2]*3^(3/4))])/(3*Sqrt[2]*3^(3/4)) - ((2 - Sqrt[3])*ArcTanh[(3^(1/4)*(1 + Sqrt[3] - 2*x))/(Sqrt[2]*Sqrt[1 + x^3])])/(3*Sqrt[2]*3^(1/4)) - ((2 - Sqrt[3])*ArcTanh[(3^(1/4)*(1 - Sqrt[3])*(1 + x))/(Sqrt[2]*Sqrt[1 + x^3])])/(6*Sqrt[2]*3^(1/4))
```

Defintions of rubi rules used

rule 989

```
Int[(x_)/(Sqrt[(a_) + (b_.)*(x_)^3]*((c_) + (d_.)*(x_)^3)), x_Symbol] := With[{q = Rt[b/a, 3], r = Simplify[(b*c - 10*a*d)/(6*a*d)]}, Simp[(-q)*(2 - r)*(ArcTan[(1 - r)*(Sqrt[a + b*x^3]/(Sqrt[2]*Rt[a, 2]*r^(3/2)))]/(3*Sqrt[2]*Rt[a, 2]*d*r^(3/2))), x] + (-Simp[q*(2 - r)*(ArcTan[Rt[a, 2]*Sqrt[r]*(1 + r)*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])]/(2*Sqrt[2]*Rt[a, 2]*d*r^(3/2))), x] - Simp[q*(2 - r)*(ArcTanh[Rt[a, 2]*Sqrt[r]*((1 + r - 2*q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])]/(3*Sqrt[2]*Rt[a, 2]*d*Sqrt[r])), x] - Simp[q*(2 - r)*(ArcTanh[Rt[a, 2]*(1 - r)*Sqrt[r]*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])]/(6*Sqrt[2]*Rt[a, 2]*d*Sqrt[r])), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b^2*c^2 - 20*a*b*c*d - 8*a^2*d^2, 0] && PosQ[a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 70.82 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.61

method	result
default	$\frac{(1+\sqrt{3})\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\sqrt{3}\operatorname{EllipticPi}\left(\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\frac{\left(-\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{3}}{3},\sqrt{\frac{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)}{9(2+\sqrt{3})\sqrt{x^3+1}}-\frac{\sqrt{2}}{-\alpha}$
elliptic	$\frac{2\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\left(-\frac{(-\sqrt{3}-1)^2}{3}+\frac{2(-\sqrt{3}-1)^2\sqrt{3}}{9}-\frac{\sqrt{3}}{9}-\frac{2(-\sqrt{3}-1)\sqrt{3}}{9}\right)\operatorname{EllipticPi}\left(\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\frac{\sqrt{2}}{3(-\sqrt{3}-1)}\right)}{3(-\sqrt{3}-1)}$
trager	Expression too large to display

input `int(x/(10+x^3+6*3^(1/2))/(x^3+1)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\begin{aligned} & -1/9*(1+3^{(1/2)})/(2+3^{(1/2)})*(3/2-1/2*I*3^{(1/2)})*((1+x)/(3/2-1/2*I*3^{(1/2)}))^{(1/2)} \\ & *((x-1/2-1/2*I*3^{(1/2)})/(-3/2-1/2*I*3^{(1/2)}))^{(1/2)}*((x-1/2+1/2*I*3^{(1/2)})/(-3/2+1/2*I*3^{(1/2)}))^{(1/2)} \\ & /((x^3+1)^{(1/2)}*3^{(1/2)}*EllipticPi(((1+x)/(3/2-1/2*I*3^{(1/2)}))^{(1/2)},1/3*(-3/2+1/2*I*3^{(1/2)})*3^{(1/2)},((-3/2+1/2*I*3^{(1/2)})/(-3/2-1/2*I*3^{(1/2)}))^{(1/2)}-1/18*2^{(1/2)}*sum((-alpha*3^{(1/2)}+_alpha-2)/(-1+2*_alpha-3^{(1/2)})*(3-I*3^{(1/2)})*((1+x)/(3-I*3^{(1/2)}))^{(1/2)}*((-I*3^{(1/2)}+2*x-1)/(-3-I*3^{(1/2)}))^{(1/2)}*((I*3^{(1/2)}+2*x-1)/(I*3^{(1/2)}-3))^{(1/2)})/((x^3+1)^{(1/2)}*(-1+2*_alpha-_alpha*3^{(1/2)})*EllipticPi(((1+x)/(3/2-1/2*I*3^{(1/2)}))^{(1/2)},-1/2*I*_alpha+1/3*I*_alpha*3^{(1/2)}+1/2*_alpha*3^{(1/2)}-_alpha-1/6*I*3^{(1/2)}+1/2,((-3/2+1/2*I*3^{(1/2)})/(-3/2-1/2*I*3^{(1/2)}))^{(1/2)}),_alpha=RootOf(_Z^2+(-3^{(1/2)}-1)*_Z+2*3^{(1/2)}+4)) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 511 vs. $2(148) = 296$.

Time = 0.20 (sec) , antiderivative size = 511, normalized size of antiderivative = 2.34

$$\begin{aligned} & \int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx \\ & = \frac{1}{36} \sqrt{14\sqrt{3}-24} \arctan \left(\frac{(3x^2 + \sqrt{3}(x^2 - 10x - 8) - 18x - 12) \sqrt{14\sqrt{3}-24}}{12\sqrt{x^3+1}} \right) \\ & + \frac{1}{18} \sqrt{\frac{7}{2}\sqrt{3}-6} \arctan \left(\frac{\sqrt{x^3+1}(\sqrt{3}(x-8) + 3x - 12) \sqrt{\frac{7}{2}\sqrt{3}-6}}{3(x^2-x-2)} \right) \\ & + \frac{1}{24} \sqrt{\frac{7}{6}\sqrt{3}-2} \log \left(\frac{x^8 + 14x^7 + 70x^6 + 20x^5 + 100x^4 + 32x^3 - 8x^2 + 6(5x^6 + 46x^5 + 62x^4 + 68x^3 + 32x^2 + 12x + 6)}{x^8 + 14x^7 + 70x^6 + 20x^5 + 100x^4 + 32x^3 - 8x^2 - 6(5x^6 + 46x^5 + 62x^4 + 68x^3 + 32x^2 + 12x + 6)} \right) \\ & - \frac{1}{24} \sqrt{\frac{7}{6}\sqrt{3}-2} \log \left(\frac{x^8 + 14x^7 + 70x^6 + 20x^5 + 100x^4 + 32x^3 - 8x^2 - 6(5x^6 + 46x^5 + 62x^4 + 68x^3 + 32x^2 + 12x + 6)}{x^8 + 14x^7 + 70x^6 + 20x^5 + 100x^4 + 32x^3 - 8x^2 + 6(5x^6 + 46x^5 + 62x^4 + 68x^3 + 32x^2 + 12x + 6)} \right) \end{aligned}$$

input `integrate(x/(10+x^3+6*3^(1/2))/(x^3+1)^(1/2),x, algorithm="fricas")`

output

```

1/36*sqrt(14*sqrt(3) - 24)*arctan(1/12*(3*x^2 + sqrt(3)*(x^2 - 10*x - 8) -
18*x - 12)*sqrt(14*sqrt(3) - 24)/sqrt(x^3 + 1)) + 1/18*sqrt(7/2*sqrt(3) -
6)*arctan(1/3*sqrt(x^3 + 1)*(sqrt(3)*(x - 8) + 3*x - 12)*sqrt(7/2*sqrt(3)
- 6)/(x^2 - x - 2)) + 1/24*sqrt(7/6*sqrt(3) - 2)*log((x^8 + 14*x^7 + 70*x
^6 + 20*x^5 + 100*x^4 + 32*x^3 - 8*x^2 + 6*(5*x^6 + 46*x^5 + 62*x^4 + 68*x
^3 + 28*x^2 + sqrt(3)*(3*x^6 + 26*x^5 + 38*x^4 + 36*x^3 + 20*x^2 + 8*x) +
8*x)*sqrt(x^3 + 1)*sqrt(7/6*sqrt(3) - 2) + 12*sqrt(3)*(x^7 + 2*x^6 + 4*x^5
+ x^4 + 2*x^3 + 4*x^2) + 32*x + 16)/(x^8 - 4*x^7 + 16*x^6 - 16*x^5 + 28*x
^4 + 32*x^3 + 64*x^2 + 32*x + 16)) - 1/24*sqrt(7/6*sqrt(3) - 2)*log((x^8 +
14*x^7 + 70*x^6 + 20*x^5 + 100*x^4 + 32*x^3 - 8*x^2 - 6*(5*x^6 + 46*x^5 +
62*x^4 + 68*x^3 + 28*x^2 + sqrt(3)*(3*x^6 + 26*x^5 + 38*x^4 + 36*x^3 + 20
*x^2 + 8*x) + 8*x)*sqrt(x^3 + 1)*sqrt(7/6*sqrt(3) - 2) + 12*sqrt(3)*(x^7 +
2*x^6 + 4*x^5 + x^4 + 2*x^3 + 4*x^2) + 32*x + 16)/(x^8 - 4*x^7 + 16*x^6 -
16*x^5 + 28*x^4 + 32*x^3 + 64*x^2 + 32*x + 16))

```

Sympy [F]

$$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx = \int \frac{x}{\sqrt{(x+1)(x^2-x+1)}(x^3+10+6\sqrt{3})} dx$$

input

```
integrate(x/(10+x**3+6*3**(1/2))/(x**3+1)**(1/2),x)
```

output

```
Integral(x/(sqrt((x + 1)*(x**2 - x + 1))*(x**3 + 10 + 6*sqrt(3))), x)
```

Maxima [F]

$$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3+6\sqrt{3}+10)\sqrt{x^3+1}} dx$$

input

```
integrate(x/(10+x^3+6*3^(1/2))/(x^3+1)^(1/2),x, algorithm="maxima")
```

output

```
integrate(x/((x^3 + 6*sqrt(3) + 10)*sqrt(x^3 + 1)), x)
```

Giac [F]

$$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3+6\sqrt{3}+10)\sqrt{x^3+1}} dx$$

input `integrate(x/(10+x^3+6*3^(1/2))/(x^3+1)^(1/2),x, algorithm="giac")`

output `integrate(x/((x^3 + 6*sqrt(3) + 10)*sqrt(x^3 + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx = \int \frac{x}{\sqrt{x^3+1}(x^3+6\sqrt{3}+10)} dx$$

input `int(x/((x^3 + 1)^(1/2)*(6*3^(1/2) + x^3 + 10)),x)`

output `int(x/((x^3 + 1)^(1/2)*(6*3^(1/2) + x^3 + 10)), x)`

Reduce [F]

$$\begin{aligned} \int \frac{x}{\sqrt{1+x^3}(10+6\sqrt{3}+x^3)} dx &= -6\sqrt{3} \left(\int \frac{\sqrt{x^3+1} x}{x^9+21x^6+12x^3-8} dx \right) \\ &\quad + \int \frac{\sqrt{x^3+1} x^4}{x^9+21x^6+12x^3-8} dx \\ &\quad + 10 \left(\int \frac{\sqrt{x^3+1} x}{x^9+21x^6+12x^3-8} dx \right) \end{aligned}$$

input `int(x/(10+x^3+6*3^(1/2))/(x^3+1)^(1/2),x)`

output

```
- 6*sqrt(3)*int((sqrt(x**3 + 1)*x)/(x**9 + 21*x**6 + 12*x**3 - 8),x) + in  
t((sqrt(x**3 + 1)*x**4)/(x**9 + 21*x**6 + 12*x**3 - 8),x) + 10*int((sqrt(x  
**3 + 1)*x)/(x**9 + 21*x**6 + 12*x**3 - 8),x)
```

$$3.87 \quad \int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx$$

Optimal result	668
Mathematica [C] (verified)	669
Rubi [A] (verified)	669
Maple [C] (verified)	670
Fricas [B] (verification not implemented)	671
Sympy [F]	672
Maxima [F]	673
Giac [F]	673
Mupad [F(-1)]	673
Reduce [F]	674

Optimal result

Integrand size = 25, antiderivative size = 210

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = -\frac{(2+\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1-\sqrt{3}-2x)}{\sqrt{2}\sqrt{1+x^3}}\right)}{3\sqrt{2}\sqrt[4]{3}} - \frac{(2+\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(1+x)}{\sqrt{2}\sqrt{1+x^3}}\right)}{6\sqrt{2}\sqrt[4]{3}} + \frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(1+x)}{\sqrt{2}\sqrt{1+x^3}}\right)}{2\sqrt{2}3^{3/4}} + \frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{(1+\sqrt{3})\sqrt{1+x^3}}{\sqrt{2}3^{3/4}}\right)}{3\sqrt{2}3^{3/4}}$$

output

```
-1/18*arctan(1/2*3^(1/4)*(1-2*x-3^(1/2))*2^(1/2)/(x^3+1)^(1/2))*(2+3^(1/2))
*3^(3/4)*2^(1/2)-1/36*arctan(1/2*3^(1/4)*(1+x)*(1+3^(1/2))*2^(1/2)/(x^3+1)
^(1/2))*(2+3^(1/2))*3^(3/4)*2^(1/2)+1/12*arctanh(1/2*3^(1/4)*(1+x)*(1-3^(
1/2))*2^(1/2)/(x^3+1)^(1/2))*(2+3^(1/2))*3^(1/4)*2^(1/2)+1/18*arctanh(1/6*
(1+3^(1/2))*(x^3+1)^(1/2)*3^(1/4)*2^(1/2))*(2+3^(1/2))*3^(1/4)*2^(1/2)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.05 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.24

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = -\frac{x^2 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, -x^3, \frac{1}{4}(5+3\sqrt{3})x^3\right)}{4(-5+3\sqrt{3})}$$

input `Integrate[x/(Sqrt[1 + x^3]*(10 - 6*Sqrt[3] + x^3)),x]`

output `-1/4*(x^2*AppellF1[2/3, 1/2, 1, 5/3, -x^3, ((5 + 3*Sqrt[3])*x^3)/4])/(-5 + 3*Sqrt[3])`

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {989}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{x^3+1}(x^3-6\sqrt{3}+10)} dx$$

↓ 989

$$-\frac{(2+\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(-2x-\sqrt{3}+1)}{\sqrt{2}\sqrt{x^3+1}}\right)}{3\sqrt{2}\sqrt[4]{3}} - \frac{(2+\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(x+1)}{\sqrt{2}\sqrt{x^3+1}}\right)}{6\sqrt{2}\sqrt[4]{3}} +$$

$$\frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(x+1)}{\sqrt{2}\sqrt{x^3+1}}\right)}{2\sqrt{2}2^{3/4}} + \frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{(1+\sqrt{3})\sqrt{x^3+1}}{\sqrt{2}2^{3/4}}\right)}{3\sqrt{2}2^{3/4}}$$

input `Int[x/(Sqrt[1 + x^3]*(10 - 6*Sqrt[3] + x^3)),x]`

output

```
-1/3*((2 + Sqrt[3])*ArcTan[(3^(1/4)*(1 - Sqrt[3] - 2*x))/(Sqrt[2]*Sqrt[1 + x^3]])/(Sqrt[2]*3^(1/4)) - ((2 + Sqrt[3])*ArcTan[(3^(1/4)*(1 + Sqrt[3])*(1 + x))/(Sqrt[2]*Sqrt[1 + x^3]])/(6*Sqrt[2]*3^(1/4)) + ((2 + Sqrt[3])*ArcTanh[(3^(1/4)*(1 - Sqrt[3])*(1 + x))/(Sqrt[2]*Sqrt[1 + x^3]])/(2*Sqrt[2]*3^(3/4)) + ((2 + Sqrt[3])*ArcTanh[((1 + Sqrt[3])*Sqrt[1 + x^3])/(Sqrt[2]*3^(3/4))])/(3*Sqrt[2]*3^(3/4)))
```

Defintions of rubi rules used

rule 989

```
Int[(x_)/(Sqrt[(a_) + (b_.)*(x_)^3]*((c_) + (d_.)*(x_)^3)), x_Symbol] := With[{q = Rt[b/a, 3], r = Simplify[(b*c - 10*a*d)/(6*a*d)]}, Simp[(-q)*(2 - r)*(ArcTan[(1 - r)*(Sqrt[a + b*x^3]/(Sqrt[2]*Rt[a, 2]*r^(3/2)))]/(3*Sqrt[2]*Rt[a, 2]*d*r^(3/2))), x] + (-Simp[q*(2 - r)*(ArcTan[Rt[a, 2]*Sqrt[r]*(1 + r)*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3]))]/(2*Sqrt[2]*Rt[a, 2]*d*r^(3/2))), x] - Simp[q*(2 - r)*(ArcTanh[Rt[a, 2]*Sqrt[r]*((1 + r - 2*q*x)/(Sqrt[2]*Sqrt[a + b*x^3]))]/(3*Sqrt[2]*Rt[a, 2]*d*Sqrt[r])), x] - Simp[q*(2 - r)*(ArcTanh[Rt[a, 2]*(1 - r)*Sqrt[r]*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3]))]/(6*Sqrt[2]*Rt[a, 2]*d*Sqrt[r])), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b^2*c^2 - 20*a*b*c*d - 8*a^2*d^2, 0] && PosQ[a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 72.37 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.67

method	result
default	$\frac{(\sqrt{3}-1)\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\sqrt{3}\operatorname{EllipticPi}\left(\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}},-\frac{\left(-\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{3}}{3},\sqrt{\frac{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)}{9(-2+\sqrt{3})\sqrt{x^3+1}}-\frac{\sqrt{2}}{-\alpha}$
elliptic	$\frac{2\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}-\frac{i\sqrt{3}}{2}}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x-\frac{1}{2}+\frac{i\sqrt{3}}{2}}{-\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\left(-\frac{2(\sqrt{3}-1)^2\sqrt{3}}{9}-\frac{(\sqrt{3}-1)^2}{3}+\frac{\sqrt{3}}{9}-\frac{2}{3}+\frac{2(\sqrt{3}-1)\sqrt{3}}{9}\right)\operatorname{EllipticPi}\left(\sqrt{\frac{1+x}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}},-\frac{i\left(\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{3}}{3(\sqrt{3}-1)}\right)}{3(\sqrt{3}-1)\sqrt{x^3+1}}$
trager	Expression too large to display

input `int(x/(10+x^3-6*3^(1/2))/(x^3+1)^(1/2),x,method=_RETURNVERBOSE)`

output `1/9*(3^(1/2)-1)/(-2+3^(1/2))*(3/2-1/2*I*3^(1/2))*((1+x)/(3/2-1/2*I*3^(1/2)))^(1/2)*((x-1/2-1/2*I*3^(1/2))/(-3/2-1/2*I*3^(1/2)))^(1/2)*((x-1/2+1/2*I*3^(1/2))/(-3/2+1/2*I*3^(1/2)))^(1/2)/(x^3+1)^(1/2)*3^(1/2)*EllipticPi(((1+x)/(3/2-1/2*I*3^(1/2)))^(1/2),-1/3*(-3/2+1/2*I*3^(1/2))*3^(1/2),((-3/2+1/2*I*3^(1/2))/(-3/2-1/2*I*3^(1/2)))^(1/2))-1/18*2^(1/2)*sum((-_alpha*3^(1/2)-_alpha+2)/(1-2*_alpha-3^(1/2))*(3-I*3^(1/2))*((1+x)/(3-I*3^(1/2)))^(1/2)*((-I*3^(1/2)+2*x-1)/(-3-I*3^(1/2)))^(1/2)*((I*3^(1/2)+2*x-1)/(I*3^(1/2)-3))^(1/2)/(x^3+1)^(1/2)*(-1+2*_alpha+_alpha*3^(1/2))*EllipticPi(((1+x)/(3/2-1/2*I*3^(1/2)))^(1/2),1/3*I*_alpha*3^(1/2)+1/2*I*_alpha-1/2*_alpha*3^(1/2)-_alpha-1/6*I*3^(1/2)+1/2,((-3/2+1/2*I*3^(1/2))/(-3/2-1/2*I*3^(1/2)))^(1/2)),_alpha=RootOf(_Z^2+(3^(1/2)-1)*_Z-2*3^(1/2)+4))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 657 vs. $2(146) = 292$.

Time = 0.21 (sec) , antiderivative size = 657, normalized size of antiderivative = 3.13

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = \text{Too large to display}$$

input `integrate(x/(10+x^3-6*3^(1/2))/(x^3+1)^(1/2),x, algorithm="fricas")`

output

```

1/6*sqrt(7/6*sqrt(3) + 2)*arctan(1/3*sqrt(x^3 + 1)*(sqrt(3)*(x + 10) - 3*x
- 18)*sqrt(7/6*sqrt(3) + 2)/(x^2 + x)) + 1/72*sqrt(7/2*sqrt(3) + 6)*log((
x^8 - 10*x^7 + 22*x^6 + 20*x^5 - 20*x^4 + 80*x^3 - 8*x^2 + 2*(5*x^6 - 30*x
^5 + 6*x^4 + 28*x^3 + 60*x^2 - 3*sqrt(3)*(x^6 - 6*x^5 + 2*x^4 + 4*x^3 + 12
*x^2 + 8*x) + 24*x + 32)*sqrt(x^3 + 1)*sqrt(7/2*sqrt(3) + 6) + 4*sqrt(3)*(
x^7 - 2*x^6 - 7*x^4 + 14*x^3 - 8*x + 16) - 64*x + 112)/(x^8 - 4*x^7 + 16*x
^6 - 16*x^5 + 28*x^4 + 32*x^3 + 64*x^2 + 32*x + 16)) - 1/72*sqrt(7/2*sqrt(
3) + 6)*log((x^8 - 10*x^7 + 22*x^6 + 20*x^5 - 20*x^4 + 80*x^3 - 8*x^2 - 2*
(5*x^6 - 30*x^5 + 6*x^4 + 28*x^3 + 60*x^2 - 3*sqrt(3)*(x^6 - 6*x^5 + 2*x^4
+ 4*x^3 + 12*x^2 + 8*x) + 24*x + 32)*sqrt(x^3 + 1)*sqrt(7/2*sqrt(3) + 6)
+ 4*sqrt(3)*(x^7 - 2*x^6 - 7*x^4 + 14*x^3 - 8*x + 16) - 64*x + 112)/(x^8 -
4*x^7 + 16*x^6 - 16*x^5 + 28*x^4 + 32*x^3 + 64*x^2 + 32*x + 16)) + 1/72*sq
qrt(14*sqrt(3) + 24)*log((x^8 - 16*x^7 + 112*x^6 - 16*x^5 + 112*x^4 + 224*
x^3 + 64*x^2 + 2*(5*x^6 - 54*x^5 + 96*x^4 - 56*x^3 - 36*x^2 - 3*sqrt(3)*(x
^6 - 10*x^5 + 20*x^4 - 8*x^3 - 4*x^2 + 8*x) + 24*x - 16)*sqrt(x^3 + 1)*sq
rt(14*sqrt(3) + 24) + 16*sqrt(3)*(x^7 - 2*x^6 + 6*x^5 + 5*x^4 + 2*x^3 + 6*x
^2 + 4*x + 4) + 128*x + 112)/(x^8 + 8*x^7 + 16*x^6 - 16*x^5 - 56*x^4 + 32*
x^3 + 64*x^2 - 64*x + 16))

```

SymPy [F]

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = \int \frac{x}{\sqrt{(x+1)(x^2-x+1)}(x^3-6\sqrt{3}+10)} dx$$

input

```
integrate(x/(10+x**3-6*3**(1/2))/(x**3+1)**(1/2),x)
```

output

```
Integral(x/(sqrt((x + 1)*(x**2 - x + 1))*(x**3 - 6*sqrt(3) + 10)), x)
```

Maxima [F]

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3-6\sqrt{3}+10)\sqrt{x^3+1}} dx$$

input `integrate(x/(10+x^3-6*3^(1/2))/(x^3+1)^(1/2),x, algorithm="maxima")`

output `integrate(x/((x^3 - 6*sqrt(3) + 10)*sqrt(x^3 + 1)), x)`

Giac [F]

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3-6\sqrt{3}+10)\sqrt{x^3+1}} dx$$

input `integrate(x/(10+x^3-6*3^(1/2))/(x^3+1)^(1/2),x, algorithm="giac")`

output `integrate(x/((x^3 - 6*sqrt(3) + 10)*sqrt(x^3 + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = \int \frac{x}{\sqrt{x^3+1}(x^3-6\sqrt{3}+10)} dx$$

input `int(x/((x^3 + 1)^(1/2)*(x^3 - 6*3^(1/2) + 10)),x)`

output `int(x/((x^3 + 1)^(1/2)*(x^3 - 6*3^(1/2) + 10)), x)`

Reduce [F]

$$\int \frac{x}{\sqrt{1+x^3}(10-6\sqrt{3}+x^3)} dx = 6\sqrt{3} \left(\int \frac{\sqrt{x^3+1}x}{x^9+21x^6+12x^3-8} dx \right) \\ + \int \frac{\sqrt{x^3+1}x^4}{x^9+21x^6+12x^3-8} dx \\ + 10 \left(\int \frac{\sqrt{x^3+1}x}{x^9+21x^6+12x^3-8} dx \right)$$

input `int(x/(10+x^3-6*3^(1/2))/(x^3+1)^(1/2),x)`

output `6*sqrt(3)*int((sqrt(x**3 + 1)*x)/(x**9 + 21*x**6 + 12*x**3 - 8),x) + int((sqrt(x**3 + 1)*x**4)/(x**9 + 21*x**6 + 12*x**3 - 8),x) + 10*int((sqrt(x**3 + 1)*x)/(x**9 + 21*x**6 + 12*x**3 - 8),x)`

3.88 $\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx$

Optimal result	675
Mathematica [C] (warning: unable to verify)	676
Rubi [A] (verified)	676
Maple [C] (verified)	677
Fricas [B] (verification not implemented)	678
Sympy [F]	679
Maxima [F]	680
Giac [F]	680
Mupad [F(-1)]	680
Reduce [F]	681

Optimal result

Integrand size = 25, antiderivative size = 222

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = \frac{(2-\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(1-x)}{\sqrt{2}\sqrt{-1+x^3}}\right)}{6\sqrt{2}\sqrt[4]{3}} + \frac{(2-\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1+\sqrt{3}+2x)}{\sqrt{2}\sqrt{-1+x^3}}\right)}{3\sqrt{2}\sqrt[4]{3}} + \frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(1-x)}{\sqrt{2}\sqrt{-1+x^3}}\right)}{2\sqrt{2}3^{3/4}} - \frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{(1-\sqrt{3})\sqrt{-1+x^3}}{\sqrt{2}3^{3/4}}\right)}{3\sqrt{2}3^{3/4}}$$

output

```
1/36*arctan(1/2*3^(1/4)*(1-x)*(1-3^(1/2))*2^(1/2)/(x^3-1)^(1/2))*(2-3^(1/2))
)*3^(3/4)*2^(1/2)+1/18*arctan(1/2*3^(1/4)*(1+2*x+3^(1/2))*2^(1/2)/(x^3-1)
)^(1/2))*(2-3^(1/2))*3^(3/4)*2^(1/2)+1/12*arctanh(1/2*3^(1/4)*(1-x)*(1+3^(1
/2))*2^(1/2)/(x^3-1)^(1/2))*(2-3^(1/2))*3^(1/4)*2^(1/2)-1/18*arctanh(1/6*(
1-3^(1/2))*(x^3-1)^(1/2)*3^(1/4)*2^(1/2))*(2-3^(1/2))*3^(1/4)*2^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.05 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.29

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = -\frac{x^2\sqrt{1-x^3} \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, x^3, \frac{x^3}{10+6\sqrt{3}}\right)}{(20+12\sqrt{3})\sqrt{-1+x^3}}$$

input `Integrate[x/(Sqrt[-1 + x^3]*(-10 - 6*Sqrt[3] + x^3)),x]`

output `-((x^2*Sqrt[1 - x^3]*AppellF1[2/3, 1/2, 1, 5/3, x^3, x^3/(10 + 6*Sqrt[3])])/(20 + 12*Sqrt[3])*Sqrt[-1 + x^3])`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 222, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {990}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{x^3-1}(x^3-6\sqrt{3}-10)} dx$$

↓ 990

$$\frac{(2-\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(1-x)}{\sqrt{2}\sqrt{x^3-1}}\right)}{6\sqrt{2}\sqrt[4]{3}} + \frac{(2-\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(2x+\sqrt{3}+1)}{\sqrt{2}\sqrt{x^3-1}}\right)}{3\sqrt{2}\sqrt[4]{3}} +$$

$$\frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(1-x)}{\sqrt{2}\sqrt{x^3-1}}\right)}{2\sqrt{2}3^{3/4}} - \frac{(2-\sqrt{3}) \operatorname{arctanh}\left(\frac{(1-\sqrt{3})\sqrt{x^3-1}}{\sqrt{2}3^{3/4}}\right)}{3\sqrt{2}3^{3/4}}$$

input `Int[x/(Sqrt[-1 + x^3]*(-10 - 6*Sqrt[3] + x^3)),x]`

output

```
((2 - Sqrt[3])*ArcTan[(3^(1/4)*(1 - Sqrt[3])*(1 - x))/(Sqrt[2]*Sqrt[-1 + x^3])])/(6*Sqrt[2]*3^(1/4)) + ((2 - Sqrt[3])*ArcTan[(3^(1/4)*(1 + Sqrt[3] + 2*x))/(Sqrt[2]*Sqrt[-1 + x^3])])/(3*Sqrt[2]*3^(1/4)) + ((2 - Sqrt[3])*ArcTanh[(3^(1/4)*(1 + Sqrt[3])*(1 - x))/(Sqrt[2]*Sqrt[-1 + x^3])])/(2*Sqrt[2]*3^(3/4)) - ((2 - Sqrt[3])*ArcTanh[((1 - Sqrt[3])*Sqrt[-1 + x^3])/(Sqrt[2]*3^(3/4))])/(3*Sqrt[2]*3^(3/4))
```

Defintions of rubi rules used

rule 990

```
Int[(x_)/(Sqrt[(a_) + (b_.)*(x_)^3]*((c_) + (d_.)*(x_)^3)), x_Symbol] := With[{q = Rt[b/a, 3], r = Simplify[(b*c - 10*a*d)/(6*a*d)]}, Simp[q*(2 - r)*(ArcTanh[(1 - r)*(Sqrt[a + b*x^3]/(Sqrt[2]*Rt[-a, 2]*r^(3/2))])/(3*Sqrt[2]*Rt[-a, 2]*d*r^(3/2))], x] + (-Simp[q*(2 - r)*(ArcTanh[Rt[-a, 2]*Sqrt[r]*(1 + r)*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])/(2*Sqrt[2]*Rt[-a, 2]*d*r^(3/2))], x] - Simp[q*(2 - r)*(ArcTan[Rt[-a, 2]*Sqrt[r]*((1 + r - 2*q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])/(3*Sqrt[2]*Rt[-a, 2]*d*Sqrt[r])], x] - Simp[q*(2 - r)*(ArcTan[Rt[-a, 2]*(1 - r)*Sqrt[r]*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])/(6*Sqrt[2]*Rt[-a, 2]*d*Sqrt[r])], x))] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b^2*c^2 - 20*a*b*c*d - 8*a^2*d^2, 0] && NegQ[a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 61.05 (sec) , antiderivative size = 347, normalized size of antiderivative = 1.56

method	result
default	$\frac{(1+\sqrt{3})\left(-\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}-\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\sqrt{3}\operatorname{EllipticPi}\left(\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}},-\frac{\left(\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{3}}{3},\sqrt{\frac{\frac{3}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)}{9(2+\sqrt{3})\sqrt{x^3-1}}-\frac{\sqrt{2}}{\dots}$
elliptic	$\frac{2\left(-\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}-\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\left(\frac{(1+\sqrt{3})^2}{3}-\frac{2(1+\sqrt{3})^2\sqrt{3}}{9}+\frac{2}{3}+\frac{\sqrt{3}}{9}-\frac{2(1+\sqrt{3})\sqrt{3}}{9}\right)\operatorname{EllipticPi}\left(\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\frac{i}{\dots}}{3(1+\sqrt{3})\sqrt{x^3-1}}$
trager	Expression too large to display

input `int(x/(-10+x^3-6*3^(1/2))/(x^3-1)^(1/2),x,method=_RETURNVERBOSE)`

output `-1/9*(1+3^(1/2))/(2+3^(1/2))*(-3/2-1/2*I*3^(1/2))*((-1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2-1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2)*((x+1/2+1/2*I*3^(1/2))/(3/2+1/2*I*3^(1/2)))^(1/2)/(x^3-1)^(1/2)*3^(1/2)*EllipticPi(((1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2),-1/3*(3/2+1/2*I*3^(1/2))*3^(1/2),((3/2+1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2))-1/18*2^(1/2)*sum((-_alpha*3^(1/2)+_alpha+2)/(-1-2*_alpha-3^(1/2))*(-3-I*3^(1/2))*((-1+x)/(-3-I*3^(1/2)))^(1/2)*((-I*3^(1/2)+2*x+1)/(3-I*3^(1/2)))^(1/2)*((I*3^(1/2)+2*x+1)/(I*3^(1/2)+3))^(1/2)/(x^3-1)^(1/2)*(1+2*_alpha-_alpha*3^(1/2))*EllipticPi(((1+x)/(-3/2-1/2*I*3^(1/2)))^(1/2),-1/2*I*_alpha+1/3*I*_alpha*3^(1/2)-1/2*_alpha*3^(1/2)+_alpha+1/6*I*3^(1/2)+1/2,((3/2+1/2*I*3^(1/2))/(3/2-1/2*I*3^(1/2)))^(1/2)),_alpha=RootOf(_Z^2+(1+3^(1/2))*_Z+2*3^(1/2)+4))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 659 vs. $2(146) = 292$.

Time = 0.21 (sec) , antiderivative size = 659, normalized size of antiderivative = 2.97

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = \text{Too large to display}$$

input `integrate(x/(-10+x^3-6*3^(1/2))/(x^3-1)^(1/2),x, algorithm="fricas")`

output

```
-1/6*sqrt(7/6*sqrt(3) - 2)*arctan(1/3*sqrt(x^3 - 1)*(sqrt(3)*(x - 10) + 3*x - 18)*sqrt(7/6*sqrt(3) - 2)/(x^2 - x)) - 1/72*sqrt(7/2*sqrt(3) - 6)*log((x^8 + 10*x^7 + 22*x^6 - 20*x^5 - 20*x^4 - 80*x^3 - 8*x^2 + 2*(5*x^6 + 30*x^5 + 6*x^4 - 28*x^3 + 60*x^2 + 3*sqrt(3)*(x^6 + 6*x^5 + 2*x^4 - 4*x^3 + 12*x^2 - 8*x) - 24*x + 32)*sqrt(x^3 - 1)*sqrt(7/2*sqrt(3) - 6) + 4*sqrt(3)*(x^7 + 2*x^6 + 7*x^4 + 14*x^3 - 8*x - 16) + 64*x + 112)/(x^8 + 4*x^7 + 16*x^6 + 16*x^5 + 28*x^4 - 32*x^3 + 64*x^2 - 32*x + 16)) + 1/72*sqrt(7/2*sqrt(3) - 6)*log((x^8 + 10*x^7 + 22*x^6 - 20*x^5 - 20*x^4 - 80*x^3 - 8*x^2 - 2*(5*x^6 + 30*x^5 + 6*x^4 - 28*x^3 + 60*x^2 + 3*sqrt(3)*(x^6 + 6*x^5 + 2*x^4 - 4*x^3 + 12*x^2 - 8*x) - 24*x + 32)*sqrt(x^3 - 1)*sqrt(7/2*sqrt(3) - 6) + 4*sqrt(3)*(x^7 + 2*x^6 + 7*x^4 + 14*x^3 - 8*x - 16) + 64*x + 112)/(x^8 + 4*x^7 + 16*x^6 + 16*x^5 + 28*x^4 - 32*x^3 + 64*x^2 - 32*x + 16)) + 1/72*sqrt(14*sqrt(3) - 24)*log((x^8 + 16*x^7 + 112*x^6 + 16*x^5 + 112*x^4 - 224*x^3 + 64*x^2 - 2*(5*x^6 + 54*x^5 + 96*x^4 + 56*x^3 - 36*x^2 + 3*sqrt(3)*(x^6 + 10*x^5 + 20*x^4 + 8*x^3 - 4*x^2 - 8*x) - 24*x - 16)*sqrt(x^3 - 1)*sqrt(14*sqrt(3) - 24) + 16*sqrt(3)*(x^7 + 2*x^6 + 6*x^5 - 5*x^4 + 2*x^3 - 6*x^2 + 4*x - 4) - 128*x + 112)/(x^8 - 8*x^7 + 16*x^6 + 16*x^5 - 56*x^4 - 32*x^3 + 64*x^2 + 64*x + 16))
```

Sympy [F]

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = \int \frac{x}{\sqrt{(x-1)(x^2+x+1)}(x^3-6\sqrt{3}-10)} dx$$

input

```
integrate(x/(-10+x**3-6*3**(1/2))/(x**3-1)**(1/2),x)
```

output

```
Integral(x/(sqrt((x - 1)*(x**2 + x + 1))*(x**3 - 6*sqrt(3) - 10)), x)
```


Maxima [F]

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3-6\sqrt{3}-10)\sqrt{x^3-1}} dx$$

input `integrate(x/(-10+x^3-6*3^(1/2))/(x^3-1)^(1/2),x, algorithm="maxima")`

output `integrate(x/((x^3 - 6*sqrt(3) - 10)*sqrt(x^3 - 1)), x)`

Giac [F]

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3-6\sqrt{3}-10)\sqrt{x^3-1}} dx$$

input `integrate(x/(-10+x^3-6*3^(1/2))/(x^3-1)^(1/2),x, algorithm="giac")`

output `integrate(x/((x^3 - 6*sqrt(3) - 10)*sqrt(x^3 - 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = \int -\frac{x}{\sqrt{x^3-1}(-x^3+6\sqrt{3}+10)} dx$$

input `int(-x/((x^3 - 1)^(1/2)*(6*3^(1/2) - x^3 + 10)),x)`

output `int(-x/((x^3 - 1)^(1/2)*(6*3^(1/2) - x^3 + 10)), x)`

Reduce [F]

$$\int \frac{x}{\sqrt{-1+x^3}(-10-6\sqrt{3}+x^3)} dx = 6\sqrt{3} \left(\int \frac{\sqrt{x^3-1}x}{x^9-21x^6+12x^3+8} dx \right) \\ + \int \frac{\sqrt{x^3-1}x^4}{x^9-21x^6+12x^3+8} dx \\ - 10 \left(\int \frac{\sqrt{x^3-1}x}{x^9-21x^6+12x^3+8} dx \right)$$

input `int(x/(-10+x^3-6*3^(1/2))/(x^3-1)^(1/2),x)`

output `6*sqrt(3)*int((sqrt(x**3 - 1)*x)/(x**9 - 21*x**6 + 12*x**3 + 8),x) + int((sqrt(x**3 - 1)*x**4)/(x**9 - 21*x**6 + 12*x**3 + 8),x) - 10*int((sqrt(x**3 - 1)*x)/(x**9 - 21*x**6 + 12*x**3 + 8),x)`

3.89 $\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx$

Optimal result	682
Mathematica [C] (warning: unable to verify)	683
Rubi [A] (verified)	683
Maple [C] (verified)	684
Fricas [B] (verification not implemented)	685
Sympy [F]	686
Maxima [F]	686
Giac [F]	687
Mupad [F(-1)]	687
Reduce [F]	687

Optimal result

Integrand size = 25, antiderivative size = 214

$$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx = -\frac{(2+\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(1-x)}{\sqrt{2}\sqrt{-1+x^3}}\right)}{2\sqrt{2}3^{3/4}} + \frac{(2+\sqrt{3}) \arctan\left(\frac{(1+\sqrt{3})\sqrt{-1+x^3}}{\sqrt{2}3^{3/4}}\right)}{3\sqrt{2}3^{3/4}} + \frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(1-x)}{\sqrt{2}\sqrt{-1+x^3}}\right)}{6\sqrt{2}\sqrt[4]{3}} + \frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1-\sqrt{3}+2x)}{\sqrt{2}\sqrt{-1+x^3}}\right)}{3\sqrt{2}\sqrt[4]{3}}$$

output

```
-1/12*arctan(1/2*3^(1/4)*(1-x)*(1-3^(1/2))*2^(1/2)/(x^3-1)^(1/2))*(2+3^(1/2))*3^(1/4)*2^(1/2)+1/18*arctan(1/6*(1+3^(1/2))*(x^3-1)^(1/2)*3^(1/4)*2^(1/2))*(2+3^(1/2))*3^(1/4)*2^(1/2)+1/18*arctanh(1/2*3^(1/4)*(1+2*x-3^(1/2))*2^(1/2)/(x^3-1)^(1/2))*(2+3^(1/2))*3^(3/4)*2^(1/2)+1/36*arctanh(1/2*3^(1/4)*(1-x)*(1+3^(1/2))*2^(1/2)/(x^3-1)^(1/2))*(2+3^(1/2))*3^(3/4)*2^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 10.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.32

$$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx = \frac{x^2\sqrt{1-x^3} \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{2}, 1, \frac{5}{3}, x^3, -\frac{x^3}{-10+6\sqrt{3}}\right)}{4(-5+3\sqrt{3})\sqrt{-1+x^3}}$$

input `Integrate[x/(Sqrt[-1 + x^3]*(-10 + 6*Sqrt[3] + x^3)),x]`

output `(x^2*Sqrt[1 - x^3]*AppellF1[2/3, 1/2, 1, 5/3, x^3, -(x^3/(-10 + 6*Sqrt[3]))]/(4*(-5 + 3*Sqrt[3])*Sqrt[-1 + x^3])`

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {990}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt{x^3-1}(x^3+6\sqrt{3}-10)} dx$$

↓ 990

$$\frac{(2+\sqrt{3}) \arctan\left(\frac{\sqrt[4]{3}(1-\sqrt{3})(1-x)}{\sqrt{2}\sqrt{x^3-1}}\right)}{2\sqrt{2}3^{3/4}} + \frac{(2+\sqrt{3}) \arctan\left(\frac{(1+\sqrt{3})\sqrt{x^3-1}}{\sqrt{2}3^{3/4}}\right)}{3\sqrt{2}3^{3/4}} +$$

$$\frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(1+\sqrt{3})(1-x)}{\sqrt{2}\sqrt{x^3-1}}\right)}{6\sqrt{2}\sqrt[4]{3}} + \frac{(2+\sqrt{3}) \operatorname{arctanh}\left(\frac{\sqrt[4]{3}(2x-\sqrt{3}+1)}{\sqrt{2}\sqrt{x^3-1}}\right)}{3\sqrt{2}\sqrt[4]{3}}$$

input `Int[x/(Sqrt[-1 + x^3]*(-10 + 6*Sqrt[3] + x^3)),x]`

output

```
-1/2*((2 + Sqrt[3])*ArcTan[(3^(1/4)*(1 - Sqrt[3])*(1 - x))/(Sqrt[2]*Sqrt[-1 + x^3]])/(Sqrt[2]*3^(3/4)) + ((2 + Sqrt[3])*ArcTan[((1 + Sqrt[3])*Sqrt[-1 + x^3])/(Sqrt[2]*3^(3/4))])/(3*Sqrt[2]*3^(3/4)) + ((2 + Sqrt[3])*ArcTanh[(3^(1/4)*(1 + Sqrt[3])*(1 - x))/(Sqrt[2]*Sqrt[-1 + x^3])])/(6*Sqrt[2]*3^(1/4)) + ((2 + Sqrt[3])*ArcTanh[(3^(1/4)*(1 - Sqrt[3] + 2*x))/(Sqrt[2]*Sqrt[-1 + x^3])])/(3*Sqrt[2]*3^(1/4))
```

Defintions of rubi rules used

rule 990

```
Int[(x_)/(Sqrt[(a_) + (b_.)*(x_)^3]*((c_) + (d_.)*(x_)^3)), x_Symbol] := With[{q = Rt[b/a, 3], r = Simplify[(b*c - 10*a*d)/(6*a*d)]}, Simp[q*(2 - r)*(ArcTanh[(1 - r)*(Sqrt[a + b*x^3]/(Sqrt[2]*Rt[-a, 2]*r^(3/2)))]/(3*Sqrt[2]*Rt[-a, 2]*d*r^(3/2))), x] + (-Simp[q*(2 - r)*(ArcTanh[Rt[-a, 2]*Sqrt[r]*(1 + r)*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])]/(2*Sqrt[2]*Rt[-a, 2]*d*r^(3/2))), x] - Simp[q*(2 - r)*(ArcTan[Rt[-a, 2]*Sqrt[r]*((1 + r - 2*q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])]/(3*Sqrt[2]*Rt[-a, 2]*d*Sqrt[r])), x] - Simp[q*(2 - r)*(ArcTan[Rt[-a, 2]*(1 - r)*Sqrt[r]*((1 + q*x)/(Sqrt[2]*Sqrt[a + b*x^3])])]/(6*Sqrt[2]*Rt[-a, 2]*d*Sqrt[r])), x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[b^2*c^2 - 20*a*b*c*d - 8*a^2*d^2, 0] && NegQ[a]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 61.29 (sec) , antiderivative size = 350, normalized size of antiderivative = 1.64

method	result
default	$\frac{(\sqrt{3}-1)\left(-\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}-\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\sqrt{3}\operatorname{EllipticPi}\left(\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\frac{\left(\frac{3}{2}+\frac{i\sqrt{3}}{2}\right)\sqrt{3}}{3},\sqrt{\frac{\frac{3}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\right)}{9(-2+\sqrt{3})\sqrt{x^3-1}}-\frac{\sqrt{2}}{-\alpha=}$
elliptic	$\frac{2\left(-\frac{3}{2}-\frac{i\sqrt{3}}{2}\right)\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}-\frac{i\sqrt{3}}{2}}{\frac{3}{2}-\frac{i\sqrt{3}}{2}}}\sqrt{\frac{x+\frac{1}{2}+\frac{i\sqrt{3}}{2}}{\frac{3}{2}+\frac{i\sqrt{3}}{2}}}\left(\frac{2(1-\sqrt{3})^2\sqrt{3}}{9}+\frac{(1-\sqrt{3})^2}{3}+\frac{2(1-\sqrt{3})\sqrt{3}}{9}+\frac{2}{3}-\frac{\sqrt{3}}{9}\right)\operatorname{EllipticPi}\left(\sqrt{\frac{-1+x}{-\frac{3}{2}-\frac{i\sqrt{3}}{2}}},\frac{i}{3(1-\sqrt{3})}\right)}{3(1-\sqrt{3})\sqrt{x^3-1}}$
trager	Expression too large to display

input `int(x/(-10+x^3+6*3^(1/2))/(x^3-1)^(1/2),x,method=_RETURNVERBOSE)`

output
$$\frac{1}{9} \frac{(3^{1/2}-1)}{(-2+3^{1/2})} \frac{(-3/2-1/2 I 3^{1/2})}{(-3/2-1/2 I 3^{1/2})} \frac{((-1+x)/(-3/2-1/2 I 3^{1/2}))^{1/2}}{((x+1/2-1/2 I 3^{1/2})/(3/2-1/2 I 3^{1/2}))^{1/2}} \frac{((x+1/2+1/2 I 3^{1/2})/(3/2+1/2 I 3^{1/2}))^{1/2}}{(x^3-1)^{1/2} 3^{1/2}} \text{EllipticPi}\left(\frac{(-1+x)/(-3/2-1/2 I 3^{1/2})}{(3/2-1/2 I 3^{1/2})}, \frac{1}{3} \frac{(3/2+1/2 I 3^{1/2}) 3^{1/2}}{(3/2+1/2 I 3^{1/2})/(3/2-1/2 I 3^{1/2})} - \frac{1}{18} 2^{1/2} \sum_{\alpha} \frac{(-\alpha 3^{1/2} - \alpha^2)}{(-3^{1/2} + 2\alpha + 1) (-3 - I 3^{1/2})} \frac{((-1+x)/(-3 - I 3^{1/2}))^{1/2}}{((-I 3^{1/2} + 2x + 1)/(3 - I 3^{1/2}))^{1/2}} \frac{(I 3^{1/2} + 2x + 1)/(I 3^{1/2} + 3)}{(x^3-1)^{1/2}} \frac{(1+2\alpha + \alpha 3^{1/2}) \text{EllipticPi}\left(\frac{(-1+x)/(-3/2-1/2 I 3^{1/2})}{(3/2-1/2 I 3^{1/2})} \right)^{1/2}}{1/3 I \alpha 3^{1/2} + 1/2 I \alpha + 1/2 \alpha 3^{1/2} + \alpha + 1/6 I 3^{1/2} + 1/2}, \frac{(3/2+1/2 I 3^{1/2})/(3/2-1/2 I 3^{1/2})}{(3/2-1/2 I 3^{1/2})} \right)^{1/2}, \alpha = \text{RootOf}(_Z^2 + (1-3^{1/2}) _Z - 2 \cdot 3^{1/2} + 4)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 516 vs. $2(148) = 296$.

Time = 0.20 (sec) , antiderivative size = 516, normalized size of antiderivative = 2.41

$$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx$$

$$= -\frac{1}{36} \sqrt{14\sqrt{3}+24} \arctan\left(-\frac{(3x^2-\sqrt{3}(x^2+10x-8)+18x-12)\sqrt{14\sqrt{3}+24}}{12\sqrt{x^3-1}}\right)$$

$$- \frac{1}{18} \sqrt{\frac{7}{2}\sqrt{3}+6} \arctan\left(\frac{\sqrt{x^3-1}(\sqrt{3}(x+8)-3x-12)\sqrt{\frac{7}{2}\sqrt{3}+6}}{3(x^2+x-2)}\right)$$

$$- \frac{1}{24} \sqrt{\frac{7}{6}\sqrt{3}+2} \log\left(\frac{x^8-14x^7+70x^6-20x^5+100x^4-32x^3-8x^2+6(5x^6-46x^5+62x^4-68x^3+32x^2-4x+1)}{x^8-14x^7+70x^6-20x^5+100x^4-32x^3-8x^2-6(5x^6-46x^5+62x^4-68x^3+32x^2-4x+1)}\right)$$

$$+ \frac{1}{24} \sqrt{\frac{7}{6}\sqrt{3}+2} \log\left(\frac{x^8-14x^7+70x^6-20x^5+100x^4-32x^3-8x^2-6(5x^6-46x^5+62x^4-68x^3+32x^2-4x+1)}{x^8-14x^7+70x^6-20x^5+100x^4-32x^3-8x^2+6(5x^6-46x^5+62x^4-68x^3+32x^2-4x+1)}\right)$$

input `integrate(x/(-10+x^3+6*3^(1/2))/(x^3-1)^(1/2),x, algorithm="fricas")`

output

```
-1/36*sqrt(14*sqrt(3) + 24)*arctan(-1/12*(3*x^2 - sqrt(3)*(x^2 + 10*x - 8)
+ 18*x - 12)*sqrt(14*sqrt(3) + 24)/sqrt(x^3 - 1)) - 1/18*sqrt(7/2*sqrt(3)
+ 6)*arctan(1/3*sqrt(x^3 - 1)*(sqrt(3)*(x + 8) - 3*x - 12)*sqrt(7/2*sqrt(
3) + 6)/(x^2 + x - 2)) - 1/24*sqrt(7/6*sqrt(3) + 2)*log((x^8 - 14*x^7 + 70
*x^6 - 20*x^5 + 100*x^4 - 32*x^3 - 8*x^2 + 6*(5*x^6 - 46*x^5 + 62*x^4 - 68
*x^3 + 28*x^2 - sqrt(3)*(3*x^6 - 26*x^5 + 38*x^4 - 36*x^3 + 20*x^2 - 8*x)
- 8*x)*sqrt(x^3 - 1)*sqrt(7/6*sqrt(3) + 2) + 12*sqrt(3)*(x^7 - 2*x^6 + 4*x
^5 - x^4 + 2*x^3 - 4*x^2) - 32*x + 16)/(x^8 + 4*x^7 + 16*x^6 + 16*x^5 + 28
*x^4 - 32*x^3 + 64*x^2 - 32*x + 16)) + 1/24*sqrt(7/6*sqrt(3) + 2)*log((x^8
- 14*x^7 + 70*x^6 - 20*x^5 + 100*x^4 - 32*x^3 - 8*x^2 - 6*(5*x^6 - 46*x^5
+ 62*x^4 - 68*x^3 + 28*x^2 - sqrt(3)*(3*x^6 - 26*x^5 + 38*x^4 - 36*x^3 +
20*x^2 - 8*x) - 8*x)*sqrt(x^3 - 1)*sqrt(7/6*sqrt(3) + 2) + 12*sqrt(3)*(x^7
- 2*x^6 + 4*x^5 - x^4 + 2*x^3 - 4*x^2) - 32*x + 16)/(x^8 + 4*x^7 + 16*x^6
+ 16*x^5 + 28*x^4 - 32*x^3 + 64*x^2 - 32*x + 16))
```

Sympy [F]

$$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx = \int \frac{x}{\sqrt{(x-1)(x^2+x+1)}(x^3-10+6\sqrt{3})} dx$$

input

```
integrate(x/(-10+x**3+6*3**(1/2))/(x**3-1)**(1/2),x)
```

output

```
Integral(x/(sqrt((x - 1)*(x**2 + x + 1))*(x**3 - 10 + 6*sqrt(3))), x)
```

Maxima [F]

$$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3+6\sqrt{3}-10)\sqrt{x^3-1}} dx$$

input

```
integrate(x/(-10+x^3+6*3^(1/2))/(x^3-1)^(1/2),x, algorithm="maxima")
```

output

```
integrate(x/((x^3 + 6*sqrt(3) - 10)*sqrt(x^3 - 1)), x)
```

Giac [F]

$$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx = \int \frac{x}{(x^3+6\sqrt{3}-10)\sqrt{x^3-1}} dx$$

input `integrate(x/(-10+x^3+6*3^(1/2))/(x^3-1)^(1/2),x, algorithm="giac")`

output `integrate(x/((x^3 + 6*sqrt(3) - 10)*sqrt(x^3 - 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx = \int \frac{x}{\sqrt{x^3-1}(x^3+6\sqrt{3}-10)} dx$$

input `int(x/((x^3 - 1)^(1/2)*(6*3^(1/2) + x^3 - 10)),x)`

output `int(x/((x^3 - 1)^(1/2)*(6*3^(1/2) + x^3 - 10)), x)`

Reduce [F]

$$\begin{aligned} \int \frac{x}{\sqrt{-1+x^3}(-10+6\sqrt{3}+x^3)} dx &= -6\sqrt{3} \left(\int \frac{\sqrt{x^3-1}x}{x^9-21x^6+12x^3+8} dx \right) \\ &\quad + \int \frac{\sqrt{x^3-1}x^4}{x^9-21x^6+12x^3+8} dx \\ &\quad - 10 \left(\int \frac{\sqrt{x^3-1}x}{x^9-21x^6+12x^3+8} dx \right) \end{aligned}$$

input `int(x/(-10+x^3+6*3^(1/2))/(x^3-1)^(1/2),x)`

output

```
- 6*sqrt(3)*int((sqrt(x**3 - 1)*x)/(x**9 - 21*x**6 + 12*x**3 + 8),x) + in  
t((sqrt(x**3 - 1)*x**4)/(x**9 - 21*x**6 + 12*x**3 + 8),x) - 10*int((sqrt(x  
**3 - 1)*x)/(x**9 - 21*x**6 + 12*x**3 + 8),x)
```

3.90
$$\int \frac{1-\sqrt{3}+x}{(1+\sqrt{3}+x)\sqrt{-4+4\sqrt{3}x^2+x^4}} dx$$

Optimal result	689
Mathematica [A] (verified)	689
Rubi [A] (verified)	690
Maple [C] (verified)	691
Fricas [B] (verification not implemented)	692
Sympy [F]	692
Maxima [F]	693
Giac [F]	693
Mupad [F(-1)]	693
Reduce [F]	694

Optimal result

Integrand size = 40, antiderivative size = 65

$$\int \frac{1-\sqrt{3}+x}{(1+\sqrt{3}+x)\sqrt{-4+4\sqrt{3}x^2+x^4}} dx$$

$$= \frac{1}{3}\sqrt{-3+2\sqrt{3}}\operatorname{arctanh}\left(\frac{(1-\sqrt{3}+x)^2}{\sqrt{3}(-3+2\sqrt{3})\sqrt{-4+4\sqrt{3}x^2+x^4}}\right)$$

output

`1/3*arctanh(((1+x-3^(1/2))^2/(-9+6*3^(1/2))^(1/2)/(-4+x^4+4*3^(1/2)*x^2)^(1/2))*(-3+2*3^(1/2))^(1/2))`

Mathematica [A] (verified)

Time = 8.54 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.18

$$\int \frac{1-\sqrt{3}+x}{(1+\sqrt{3}+x)\sqrt{-4+4\sqrt{3}x^2+x^4}} dx$$

$$= \frac{1}{3}\sqrt{-3+2\sqrt{3}}\operatorname{arctanh}\left(\frac{\sqrt{9+6\sqrt{3}}\sqrt{-4+4\sqrt{3}x^2+x^4}}{2+(-2-2\sqrt{3})x+(2+\sqrt{3})x^2}\right)$$

input `Integrate[(1 - Sqrt[3] + x)/((1 + Sqrt[3] + x)*Sqrt[-4 + 4*Sqrt[3]*x^2 + x^4]),x]`

output `(Sqrt[-3 + 2*Sqrt[3]]*ArcTanh[(Sqrt[9 + 6*Sqrt[3]]*Sqrt[-4 + 4*Sqrt[3]*x^2 + x^4])/(2 + (-2 - 2*Sqrt[3])*x + (2 + Sqrt[3])*x^2))]/3`

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.12, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2278, 220}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x - \sqrt{3} + 1}{(x + \sqrt{3} + 1) \sqrt{x^4 + 4\sqrt{3}x^2 - 4}} dx$$

↓ 2278

$$-4(2 - \sqrt{3}) \int \frac{1}{\frac{4(x - \sqrt{3} + 1)^4}{x^4 + 4\sqrt{3}x^2 - 4} + 12(3 - 2\sqrt{3})} d \frac{(x - \sqrt{3} + 1)^2}{\sqrt{x^4 + 4\sqrt{3}x^2 - 4}}$$

↓ 220

$$\frac{(2 - \sqrt{3}) \operatorname{arctanh}\left(\frac{(x - \sqrt{3} + 1)^2}{\sqrt{3(2\sqrt{3} - 3)} \sqrt{x^4 + 4\sqrt{3}x^2 - 4}}\right)}{\sqrt{3(2\sqrt{3} - 3)}}$$

input `Int[(1 - Sqrt[3] + x)/((1 + Sqrt[3] + x)*Sqrt[-4 + 4*Sqrt[3]*x^2 + x^4]),x]`

output `((2 - Sqrt[3])*ArcTanh[(1 - Sqrt[3] + x)^2/(Sqrt[3*(-3 + 2*Sqrt[3]])*Sqrt[-4 + 4*Sqrt[3]*x^2 + x^4])])/Sqrt[3*(-3 + 2*Sqrt[3])]`

Defintions of rubi rules used

```
rule 220 Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])
```

```
rule 2278 Int[((A_) + (B_.)*(x_))/(((d_) + (e_.)*(x_))*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_Symbol] := Simp[(-A^2)*(B*d + A*e)/e Subst[Int[1/(6*A^3*B*d + 3*A^4*e - a*e*x^2), x], x, (A + B*x)^2/Sqrt[a + b*x^2 + c*x^4]], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[B*d - A*e, 0] && EqQ[c^2*d^6 + a*e^4*(13*c*d^2 + b*e^2), 0] && EqQ[b^2*e^4 - 12*c*d^2*(c*d^2 - b*e^2), 0] && EqQ[4*A*c*d*e + B*(2*c*d^2 - b*e^2), 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 2.50 (sec) , antiderivative size = 327, normalized size of antiderivative = 5.03

method	result
elliptic	$\frac{\sqrt{1 - \left(-1 + \frac{\sqrt{3}}{2}\right)x^2} \sqrt{1 - \left(1 + \frac{\sqrt{3}}{2}\right)x^2} \operatorname{EllipticF}\left(x\left(\frac{i\sqrt{3}}{2} - \frac{i}{2}\right), i\sqrt{1 + 4\sqrt{3}\left(1 + \frac{\sqrt{3}}{2}\right)}\right)}{\left(\frac{i\sqrt{3}}{2} - \frac{i}{2}\right)\sqrt{-4 + x^4 + 4\sqrt{3}x^2}} - 2\sqrt{3} \left(-\frac{\operatorname{arctanh}\left(\frac{4(-\sqrt{3}-1)^2\sqrt{3}-8}{2\sqrt{(-\sqrt{3}-1)^4+4(-\sqrt{3}-1)^2+4}}\right)}{2\sqrt{(-\sqrt{3}-1)^4+4}} \right)$

```
input int((1+x-3^(1/2))/(1+x+3^(1/2)))/(-4+x^4+4*3^(1/2)*x^2)^(1/2), x, method=_RET URNVERBOSE)
```

```
output 1/(1/2*I*3^(1/2)-1/2*I)*(1-(-1+1/2*3^(1/2))*x^2)^(1/2)*(1-(1+1/2*3^(1/2))*x^2)^(1/2)/(-4+x^4+4*3^(1/2)*x^2)^(1/2)*EllipticF(x*(1/2*I*3^(1/2)-1/2*I), I*(1+4*3^(1/2)*(1+1/2*3^(1/2))))^(1/2))-2*3^(1/2)*(-1/2/((-3^(1/2)-1)^4+4*(-3^(1/2)-1)^2*3^(1/2)-4)^(1/2)*arctanh(1/2*(4*(-3^(1/2)-1)^2*3^(1/2)-8+4*3^(1/2))*x^2+2*x^2*(-3^(1/2)-1)^2)/((-3^(1/2)-1)^4+4*(-3^(1/2)-1)^2*3^(1/2)-4)^(1/2)/(-4+x^4+4*3^(1/2)*x^2)^(1/2))-1/(-1+1/2*3^(1/2))^(1/2)/(-3^(1/2)-1)*(1-(-1+1/2*3^(1/2))*x^2)^(1/2)*(1-(1+1/2*3^(1/2))*x^2)^(1/2)/(-4+x^4+4*3^(1/2)*x^2)^(1/2)*EllipticPi((-1+1/2*3^(1/2))^(1/2)*x, 1/(-1+1/2*3^(1/2)))/(-3^(1/2)-1)^2, (1+1/2*3^(1/2))^(1/2)/(-1+1/2*3^(1/2))^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 323 vs. $2(47) = 94$.

Time = 0.16 (sec) , antiderivative size = 323, normalized size of antiderivative = 4.97

$$\int \frac{1 - \sqrt{3} + x}{(1 + \sqrt{3} + x) \sqrt{-4 + 4\sqrt{3}x^2 + x^4}} dx$$

$$= \frac{1}{12} \sqrt{2\sqrt{3} - 3} \log \left(-\frac{37x^{12} - 204x^{11} + 804x^{10} - 2408x^9 + 3708x^8 - 5472x^7 + 6432x^6 + 10944x^5 + 14832x^4 + 19264x^3 + 12864x^2 + (54x^{10} - 300x^9 + 1026x^8 - 2232x^7 + 3024x^6 - 3024x^5 - 1008x^4 - 2016x^3 - 2592x^2 + \sqrt{3})(31x^{10} - 176x^9 + 576x^8 - 1320x^7 + 1848x^6 - 1008x^5 + 1344x^4 + 1632x^3 + 1008x^2 + 832x + 256) - 1152x - 480}{\sqrt{x^4 + 4\sqrt{3}x^2 - 4}} \sqrt{2\sqrt{3} - 3} + 3\sqrt{3} \log(3) (7x^{12} - 40x^{11} + 160x^{10} - 400x^9 + 924x^8 - 960x^7 - 1920x^5 - 3696x^4 - 3200x^3 - 2560x^2 - 1280x - 448) + 6528x + 2368 \right) / (x^{12} + 12x^{11} + 48x^{10} + 40x^9 - 180x^8 - 288x^7 + 384x^6 + 576x^5 - 720x^4 - 320x^3 + 768x^2 - 384x + 64)$$

input `integrate((1+x-3^(1/2))/(1+x+3^(1/2))/(-4+x^4+4*3^(1/2)*x^2)^(1/2),x, algo
rithm="fricas")`

output `1/12*sqrt(2*sqrt(3) - 3)*log(-(37*x^12 - 204*x^11 + 804*x^10 - 2408*x^9 +
3708*x^8 - 5472*x^7 + 6432*x^6 + 10944*x^5 + 14832*x^4 + 19264*x^3 + 12864
*x^2 + (54*x^10 - 300*x^9 + 1026*x^8 - 2232*x^7 + 3024*x^6 - 3024*x^5 - 10
08*x^4 - 2016*x^3 - 2592*x^2 + sqrt(3)*(31*x^10 - 176*x^9 + 576*x^8 - 1320
*x^7 + 1848*x^6 - 1008*x^5 + 1344*x^4 + 1632*x^3 + 1008*x^2 + 832*x + 256)
- 1152*x - 480)*sqrt(x^4 + 4*sqrt(3)*x^2 - 4)*sqrt(2*sqrt(3) - 3) + 3*sqrt
t(3)*(7*x^12 - 40*x^11 + 160*x^10 - 400*x^9 + 924*x^8 - 960*x^7 - 1920*x^5
- 3696*x^4 - 3200*x^3 - 2560*x^2 - 1280*x - 448) + 6528*x + 2368)/(x^12 +
12*x^11 + 48*x^10 + 40*x^9 - 180*x^8 - 288*x^7 + 384*x^6 + 576*x^5 - 720*
x^4 - 320*x^3 + 768*x^2 - 384*x + 64))`

Sympy [F]

$$\int \frac{1 - \sqrt{3} + x}{(1 + \sqrt{3} + x) \sqrt{-4 + 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x - \sqrt{3} + 1}{(x + 1 + \sqrt{3}) \sqrt{x^4 + 4\sqrt{3}x^2 - 4}} dx$$

input `integrate((1+x-3**(1/2))/(1+x+3**(1/2))/(-4+x**4+4*3**(1/2)*x**2)**(1/2),x
)`

output `Integral((x - sqrt(3) + 1)/((x + 1 + sqrt(3))*sqrt(x**4 + 4*sqrt(3)*x**2 -
4)), x)`

Maxima [F]

$$\int \frac{1 - \sqrt{3} + x}{(1 + \sqrt{3} + x) \sqrt{-4 + 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x - \sqrt{3} + 1}{\sqrt{x^4 + 4\sqrt{3}x^2 - 4}(x + \sqrt{3} + 1)} dx$$

input `integrate((1+x-3^(1/2))/(1+x+3^(1/2))/(-4+x^4+4*3^(1/2)*x^2)^(1/2),x, algorith="maxima")`

output `integrate((x - sqrt(3) + 1)/(sqrt(x^4 + 4*sqrt(3)*x^2 - 4)*(x + sqrt(3) + 1)), x)`

Giac [F]

$$\int \frac{1 - \sqrt{3} + x}{(1 + \sqrt{3} + x) \sqrt{-4 + 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x - \sqrt{3} + 1}{\sqrt{x^4 + 4\sqrt{3}x^2 - 4}(x + \sqrt{3} + 1)} dx$$

input `integrate((1+x-3^(1/2))/(1+x+3^(1/2))/(-4+x^4+4*3^(1/2)*x^2)^(1/2),x, algorith="giac")`

output `integrate((x - sqrt(3) + 1)/(sqrt(x^4 + 4*sqrt(3)*x^2 - 4)*(x + sqrt(3) + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1 - \sqrt{3} + x}{(1 + \sqrt{3} + x) \sqrt{-4 + 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x - \sqrt{3} + 1}{(x + \sqrt{3} + 1) \sqrt{x^4 + 4\sqrt{3}x^2 - 4}} dx$$

input `int((x - 3^(1/2) + 1)/((x + 3^(1/2) + 1)*(4*3^(1/2)*x^2 + x^4 - 4)^(1/2)), x)`

output `int((x - 3^(1/2) + 1)/((x + 3^(1/2) + 1)*(4*3^(1/2)*x^2 + x^4 - 4)^(1/2)), x)`

Reduce [F]

$$\int \frac{1 - \sqrt{3} + x}{(1 + \sqrt{3} + x) \sqrt{-4 + 4\sqrt{3}x^2 + x^4}} dx = -4\sqrt{3} \left(\int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}}{x^8 - 56x^4 + 16} dx \right) \\ - 2\sqrt{3} \left(\int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}x^3}{x^8 - 56x^4 + 16} dx \right) \\ - 2\sqrt{3} \left(\int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}x^2}{x^8 - 56x^4 + 16} dx \right) \\ - 8\sqrt{3} \left(\int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}x}{x^8 - 56x^4 + 16} dx \right) \\ + 8 \left(\int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}}{x^8 - 56x^4 + 16} dx \right) \\ + \int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}x^4}{x^8 - 56x^4 + 16} dx \\ + 6 \left(\int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}x^2}{x^8 - 56x^4 + 16} dx \right) \\ + 12 \left(\int \frac{\sqrt{4\sqrt{3}x^2 + x^4 - 4}x}{x^8 - 56x^4 + 16} dx \right)$$

input `int((1+x-3^(1/2))/(1+x+3^(1/2))/(-4+x^4+4*3^(1/2)*x^2)^(1/2), x)`

output `- 4*sqrt(3)*int(sqrt(4*sqrt(3)*x**2 + x**4 - 4)/(x**8 - 56*x**4 + 16), x) \\ - 2*sqrt(3)*int((sqrt(4*sqrt(3)*x**2 + x**4 - 4)*x**3)/(x**8 - 56*x**4 + 16), x) \\ - 2*sqrt(3)*int((sqrt(4*sqrt(3)*x**2 + x**4 - 4)*x**2)/(x**8 - 56*x**4 + 16), x) \\ - 8*sqrt(3)*int((sqrt(4*sqrt(3)*x**2 + x**4 - 4)*x)/(x**8 - 56*x**4 + 16), x) \\ + 8*int(sqrt(4*sqrt(3)*x**2 + x**4 - 4)/(x**8 - 56*x**4 + 16), x) \\ + int((sqrt(4*sqrt(3)*x**2 + x**4 - 4)*x**4)/(x**8 - 56*x**4 + 16), x) \\ + 6*int((sqrt(4*sqrt(3)*x**2 + x**4 - 4)*x**2)/(x**8 - 56*x**4 + 16), x) \\ + 12*int((sqrt(4*sqrt(3)*x**2 + x**4 - 4)*x)/(x**8 - 56*x**4 + 16), x)`

3.91
$$\int \frac{1+\sqrt{3}+x}{(1-\sqrt{3}+x)\sqrt{-4-4\sqrt{3}x^2+x^4}} dx$$

Optimal result	695
Mathematica [A] (verified)	695
Rubi [A] (verified)	696
Maple [C] (verified)	697
Fricas [B] (verification not implemented)	698
Sympy [F]	698
Maxima [F]	699
Giac [F]	699
Mupad [F(-1)]	699
Reduce [F]	700

Optimal result

Integrand size = 40, antiderivative size = 63

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx$$

$$= -\frac{1}{3} \sqrt{3 + 2\sqrt{3}} \arctan \left(\frac{(1 + \sqrt{3} + x)^2}{\sqrt{3(3 + 2\sqrt{3})} \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} \right)$$

output `-1/3*arctan(((1+x+3^(1/2))^2/(9+6*3^(1/2))^(1/2)/(-4+x^4-4*3^(1/2)*x^2)^(1/2))*(3+2*3^(1/2))^(1/2)`

Mathematica [A] (verified)

Time = 8.47 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.22

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx$$

$$= -\frac{1}{3} \sqrt{3 + 2\sqrt{3}} \arctan \left(\frac{\sqrt{-9 + 6\sqrt{3}} \sqrt{-4 - 4\sqrt{3}x^2 + x^4}}{-2 + (2 - 2\sqrt{3})x + (-2 + \sqrt{3})x^2} \right)$$

input `Integrate[(1 + Sqrt[3] + x)/((1 - Sqrt[3] + x)*Sqrt[-4 - 4*Sqrt[3]*x^2 + x^4]),x]`

output `-1/3*(Sqrt[3 + 2*Sqrt[3]]*ArcTan[(Sqrt[-9 + 6*Sqrt[3]]*Sqrt[-4 - 4*Sqrt[3]*x^2 + x^4])/(-2 + (2 - 2*Sqrt[3])*x + (-2 + Sqrt[3])*x^2)])`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.11, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$, Rules used = {2278, 216}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x + \sqrt{3} + 1}{(x - \sqrt{3} + 1) \sqrt{x^4 - 4\sqrt{3}x^2 - 4}} dx$$

↓ 2278

$$-4(2 + \sqrt{3}) \int \frac{1}{\frac{4(x + \sqrt{3} + 1)^4}{x^4 - 4\sqrt{3}x^2 - 4} + 12(3 + 2\sqrt{3})} d \frac{(x + \sqrt{3} + 1)^2}{\sqrt{x^4 - 4\sqrt{3}x^2 - 4}}$$

↓ 216

$$-\frac{(2 + \sqrt{3}) \arctan \left(\frac{(x + \sqrt{3} + 1)^2}{\sqrt{3(3 + 2\sqrt{3})} \sqrt{x^4 - 4\sqrt{3}x^2 - 4}} \right)}{\sqrt{3(3 + 2\sqrt{3})}}$$

input `Int[(1 + Sqrt[3] + x)/((1 - Sqrt[3] + x)*Sqrt[-4 - 4*Sqrt[3]*x^2 + x^4]),x]`

output `-(((2 + Sqrt[3])*ArcTan[(1 + Sqrt[3] + x)^2/(Sqrt[3*(3 + 2*Sqrt[3]])*Sqrt[-4 - 4*Sqrt[3]*x^2 + x^4]]))/Sqrt[3*(3 + 2*Sqrt[3])])`

Defintions of rubi rules used

rule 216

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

rule 2278

```
Int[((A_) + (B_)*(x_))/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_)^2 + (c_
.)*(x_)^4]), x_Symbol] := Simp[(-A^2)*(B*d + A*e)/e Subst[Int[1/(6*A^3*
B*d + 3*A^4*e - a*e*x^2), x], x, (A + B*x)^2/Sqrt[a + b*x^2 + c*x^4]], x] /
; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[B*d - A*e, 0] && EqQ[c^2*d^6 + a*e
^4*(13*c*d^2 + b*e^2), 0] && EqQ[b^2*e^4 - 12*c*d^2*(c*d^2 - b*e^2), 0] &&
EqQ[4*A*c*d*e + B*(2*c*d^2 - b*e^2), 0]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 2.62 (sec) , antiderivative size = 311, normalized size of antiderivative = 4.94

method	result
elliptic	$\frac{\sqrt{1 - \left(-1 - \frac{\sqrt{3}}{2}\right)x^2} \sqrt{1 - \left(1 - \frac{\sqrt{3}}{2}\right)x^2} \operatorname{EllipticF}\left(x\left(\frac{i}{2} + \frac{i\sqrt{3}}{2}\right), i\sqrt{1 - 4\sqrt{3}\left(1 - \frac{\sqrt{3}}{2}\right)}\right)}{\left(\frac{i}{2} + \frac{i\sqrt{3}}{2}\right)\sqrt{-4 + x^4 - 4\sqrt{3}x^2}} + 2\sqrt{3} \left(-\frac{\operatorname{arctanh}\left(\frac{-4(\sqrt{3}-1)^2\sqrt{3}-8}{2\sqrt{(\sqrt{3}-1)^4-4(\sqrt{3}-1)}}\right)}{2\sqrt{(\sqrt{3}-1)^4-4(\sqrt{3}-1)}} \right)$

input

```
int((1+x+3^(1/2))/(1+x-3^(1/2))/(-4+x^4-4*3^(1/2)*x^2)^(1/2), x, method=_RET
URNVERBOSE)
```

output

```
1/(1/2*I+1/2*I*3^(1/2))*(1-(-1-1/2*3^(1/2))*x^2)^(1/2)*(1-(1-1/2*3^(1/2))*
x^2)^(1/2)/(-4+x^4-4*3^(1/2)*x^2)^(1/2)*EllipticF(x*(1/2*I+1/2*I*3^(1/2)),
I*(1-4*3^(1/2)*(1-1/2*3^(1/2))))^(1/2)+2*3^(1/2)*(-1/2/((3^(1/2)-1)^4-4*(3
^(1/2)-1)^2*3^(1/2)-4)^(1/2)*arctanh(1/2*(-4*(3^(1/2)-1)^2*3^(1/2)-8-4*3^(
1/2)*x^2+2*x^2*(3^(1/2)-1)^2)/((3^(1/2)-1)^4-4*(3^(1/2)-1)^2*3^(1/2)-4)^(1
/2)/(-4+x^4-4*3^(1/2)*x^2)^(1/2))-1/(-1-1/2*3^(1/2))^(1/2)/(3^(1/2)-1)*(1-
(-1-1/2*3^(1/2))*x^2)^(1/2)*(1-(1-1/2*3^(1/2))*x^2)^(1/2)/(-4+x^4-4*3^(1/2
)*x^2)^(1/2)*EllipticPi((-1-1/2*3^(1/2))^(1/2)*x, 1/(-1-1/2*3^(1/2))/(3^(1/
2)-1)^2, (1-1/2*3^(1/2))^(1/2)/(-1-1/2*3^(1/2))^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. $2(45) = 90$.

Time = 0.16 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.78

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx$$

$$= \frac{1}{6} \sqrt{2\sqrt{3} + 3} \arctan \left(-\frac{(9x^4 - 30x^3 + 18x^2 - 2\sqrt{3}(2x^4 - 10x^3 + 3x^2 - 10x + 2) + 24)\sqrt{x^4 - 4\sqrt{3}x^2 - 4}}{11x^6 - 42x^5 + 66x^4 - 176x^3 - 132x^2 - 168x - 88} \right)$$

input `integrate((1+x+3^(1/2))/(1+x-3^(1/2))/(-4+x^4-4*3^(1/2)*x^2)^(1/2),x, algo
rithm="fricas")`

output `1/6*sqrt(2*sqrt(3) + 3)*arctan(-(9*x^4 - 30*x^3 + 18*x^2 - 2*sqrt(3)*(2*x^4
4 - 10*x^3 + 3*x^2 - 10*x + 2) + 24)*sqrt(x^4 - 4*sqrt(3)*x^2 - 4)*sqrt(2*
sqrt(3) + 3)/(11*x^6 - 42*x^5 + 66*x^4 - 176*x^3 - 132*x^2 - 168*x - 88))`

Sympy [F]

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x + 1 + \sqrt{3}}{(x - \sqrt{3} + 1) \sqrt{x^4 - 4\sqrt{3}x^2 - 4}} dx$$

input `integrate((1+x+3**(1/2))/(1+x-3**(1/2))/(-4+x**4-4*3**(1/2)*x**2)**(1/2),x
)`

output `Integral((x + 1 + sqrt(3))/((x - sqrt(3) + 1)*sqrt(x**4 - 4*sqrt(3)*x**2 -
4)), x)`

Maxima [F]

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x + \sqrt{3} + 1}{\sqrt{x^4 - 4\sqrt{3}x^2 - 4}(x - \sqrt{3} + 1)} dx$$

input `integrate((1+x+3^(1/2))/(1+x-3^(1/2))/(-4+x^4-4*3^(1/2)*x^2)^(1/2),x, algorith="maxima")`

output `integrate((x + sqrt(3) + 1)/(sqrt(x^4 - 4*sqrt(3)*x^2 - 4)*(x - sqrt(3) + 1)), x)`

Giac [F]

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x + \sqrt{3} + 1}{\sqrt{x^4 - 4\sqrt{3}x^2 - 4}(x - \sqrt{3} + 1)} dx$$

input `integrate((1+x+3^(1/2))/(1+x-3^(1/2))/(-4+x^4-4*3^(1/2)*x^2)^(1/2),x, algorith="giac")`

output `integrate((x + sqrt(3) + 1)/(sqrt(x^4 - 4*sqrt(3)*x^2 - 4)*(x - sqrt(3) + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx = \int \frac{x + \sqrt{3} + 1}{\sqrt{x^4 - 4\sqrt{3}x^2 - 4}(x - \sqrt{3} + 1)} dx$$

input `int((x + 3^(1/2) + 1)/((x^4 - 4*3^(1/2)*x^2 - 4)^(1/2)*(x - 3^(1/2) + 1)), x)`

output `int((x + 3^(1/2) + 1)/((x^4 - 4*3^(1/2)*x^2 - 4)^(1/2)*(x - 3^(1/2) + 1)), x)`

Reduce [F]

$$\int \frac{1 + \sqrt{3} + x}{(1 - \sqrt{3} + x) \sqrt{-4 - 4\sqrt{3}x^2 + x^4}} dx = 4\sqrt{3} \left(\int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}}{x^8 - 56x^4 + 16} dx \right) \\ + 2\sqrt{3} \left(\int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}x^3}{x^8 - 56x^4 + 16} dx \right) \\ + 2\sqrt{3} \left(\int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}x^2}{x^8 - 56x^4 + 16} dx \right) \\ + 8\sqrt{3} \left(\int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}x}{x^8 - 56x^4 + 16} dx \right) \\ + 8 \left(\int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}}{x^8 - 56x^4 + 16} dx \right) \\ + \int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}x^4}{x^8 - 56x^4 + 16} dx \\ + 6 \left(\int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}x^2}{x^8 - 56x^4 + 16} dx \right) \\ + 12 \left(\int \frac{\sqrt{-4\sqrt{3}x^2 + x^4 - 4}x}{x^8 - 56x^4 + 16} dx \right)$$

input `int((1+x+3^(1/2))/(1+x-3^(1/2))/(-4+x^4-4*3^(1/2)*x^2)^(1/2), x)`

output

```
4*sqrt(3)*int(sqrt(-4*sqrt(3)*x**2 + x**4 - 4)/(x**8 - 56*x**4 + 16),x)
+ 2*sqrt(3)*int((sqrt(-4*sqrt(3)*x**2 + x**4 - 4)*x**3)/(x**8 - 56*x**4
+ 16),x) + 2*sqrt(3)*int((sqrt(-4*sqrt(3)*x**2 + x**4 - 4)*x**2)/(x**8 -
56*x**4 + 16),x) + 8*sqrt(3)*int((sqrt(-4*sqrt(3)*x**2 + x**4 - 4)*x)/(
x**8 - 56*x**4 + 16),x) + 8*int(sqrt(-4*sqrt(3)*x**2 + x**4 - 4)/(x**8 -
56*x**4 + 16),x) + int((sqrt(-4*sqrt(3)*x**2 + x**4 - 4)*x**4)/(x**8 -
56*x**4 + 16),x) + 6*int((sqrt(-4*sqrt(3)*x**2 + x**4 - 4)*x**2)/(x**8 -
56*x**4 + 16),x) + 12*int((sqrt(-4*sqrt(3)*x**2 + x**4 - 4)*x)/(x**8 -
56*x**4 + 16),x)
```

3.92 $\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx$

Optimal result	702
Mathematica [A] (verified)	702
Rubi [A] (verified)	703
Maple [C] (verified)	704
Fricas [F(-2)]	705
Sympy [F]	705
Maxima [F]	705
Giac [F]	706
Mupad [F(-1)]	706
Reduce [F]	706

Optimal result

Integrand size = 18, antiderivative size = 53

$$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx = \sqrt{3} \arctan\left(\frac{1 + \frac{2(2+x)}{\sqrt[3]{2+x^3}}}{\sqrt{3}}\right) + \log(1+x) - \frac{3}{2} \log\left(2+x-\sqrt[3]{2+x^3}\right)$$

output

```
ln(1+x)-3/2*ln(2+x-(x^3+2)^(1/3))+arctan(1/3*(1+2*(2+x)/(x^3+2)^(1/3))*3^(1/2))*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.75 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.74

$$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx = -\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{2+x^3}}{4+2x+\sqrt[3]{2+x^3}}\right) - \log\left(-2-x+\sqrt[3]{2+x^3}\right) + \frac{1}{2} \log\left(4+4x+x^2+(2+x)\sqrt[3]{2+x^3}+(2+x^3)^{2/3}\right)$$

input

```
Integrate[(-1 + x)/((1 + x)*(2 + x^3)^(1/3)), x]
```

output

$$-(\text{Sqrt}[3]*\text{ArcTan}[(\text{Sqrt}[3]*(2 + x^3)^{(1/3)})/(4 + 2*x + (2 + x^3)^{(1/3)})]) - \text{Log}[-2 - x + (2 + x^3)^{(1/3)}] + \text{Log}[4 + 4*x + x^2 + (2 + x)*(2 + x^3)^{(1/3)} + (2 + x^3)^{(2/3)}]/2$$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.056$, Rules used = {2576}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x-1}{(x+1)\sqrt[3]{x^3+2}} dx$$

↓ 2576

$$\sqrt{3} \arctan\left(\frac{\frac{2(x+2)}{\sqrt[3]{x^3+2}} + 1}{\sqrt{3}}\right) - \frac{3}{2} \log\left(-\sqrt[3]{x^3+2} + x + 2\right) + \log(x+1)$$

input

$$\text{Int}[(-1 + x)/((1 + x)*(2 + x^3)^{(1/3)}), x]$$

output

$$\text{Sqrt}[3]*\text{ArcTan}[(1 + (2*(2 + x))/(2 + x^3)^{(1/3)})/\text{Sqrt}[3]] + \text{Log}[1 + x] - (3*\text{Log}[2 + x - (2 + x^3)^{(1/3)}])/2$$

Defintions of rubi rules used

rule 2576

$$\text{Int}[(e_ + (f_)*(x_))/((c_ + (d_)*(x_))*((a_ + (b_)*(x_)^3)^{(1/3)}), x_Symbol] := \text{Simp}[\text{Sqrt}[3]*f*(\text{ArcTan}[(1 + 2*\text{Rt}[b, 3]*((2*c + d*x)/(d*(a + b*x^3)^{(1/3)})))/\text{Sqrt}[3])/(\text{Rt}[b, 3]*d), x] + (\text{Simp}[(f*\text{Log}[c + d*x])/(\text{Rt}[b, 3]*d), x] - \text{Simp}[(3*f*\text{Log}[\text{Rt}[b, 3]*(2*c + d*x) - d*(a + b*x^3)^{(1/3)}])/(2*\text{Rt}[b, 3]*d), x]) /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{EqQ}[d*e + c*f, 0] \&\& \text{EqQ}[2*b*c^3 - a*d^3, 0]$$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 2.49 (sec) , antiderivative size = 818, normalized size of antiderivative = 15.43

method	result	size
trager	Expression too large to display	818

input `int((-1+x)/(1+x)/(x^3+2)^(1/3),x,method=_RETURNVERBOSE)`

output

```
RootOf(_Z^2-_Z+1)*ln(-(1239*RootOf(_Z^2-_Z+1)^2*x^3-2478*RootOf(_Z^2-_Z+1)
^2*x^2+4504*RootOf(_Z^2-_Z+1)*(x^3+2)^(2/3)*x+4504*(x^3+2)^(1/3)*RootOf(_Z
^2-_Z+1)*x^2+3265*RootOf(_Z^2-_Z+1)*x^3-4956*RootOf(_Z^2-_Z+1)^2*x+9008*Ro
otOf(_Z^2-_Z+1)*(x^3+2)^(2/3)+18016*(x^3+2)^(1/3)*RootOf(_Z^2-_Z+1)*x+1081
6*RootOf(_Z^2-_Z+1)*x^2+335*x*(x^3+2)^(2/3)+335*(x^3+2)^(1/3)*x^2+1574*x^3
+18016*(x^3+2)^(1/3)*RootOf(_Z^2-_Z+1)+21632*RootOf(_Z^2-_Z+1)*x+670*(x^3+
2)^(2/3)+1340*x*(x^3+2)^(1/3)+7870*x^2+17346*RootOf(_Z^2-_Z+1)+1340*(x^3+2
)^(1/3)+15740*x+11018)/(1+x)^2)-ln(-(1239*RootOf(_Z^2-_Z+1)^2*x^3-2478*Ro
otOf(_Z^2-_Z+1)^2*x^2-4504*RootOf(_Z^2-_Z+1)*(x^3+2)^(2/3)*x-4504*(x^3+2)^(
1/3)*RootOf(_Z^2-_Z+1)*x^2-5743*RootOf(_Z^2-_Z+1)*x^3-4956*RootOf(_Z^2-_Z+
1)^2*x-9008*RootOf(_Z^2-_Z+1)*(x^3+2)^(2/3)-18016*(x^3+2)^(1/3)*RootOf(_Z
^2-_Z+1)*x-5860*RootOf(_Z^2-_Z+1)*x^2+4839*x*(x^3+2)^(2/3)+4839*(x^3+2)^(1/
3)*x^2+6078*x^3-18016*(x^3+2)^(1/3)*RootOf(_Z^2-_Z+1)-11720*RootOf(_Z^2-_Z
+1)*x+9678*(x^3+2)^(2/3)+19356*x*(x^3+2)^(1/3)+16208*x^2-17346*RootOf(_Z^2
-_Z+1)+19356*(x^3+2)^(1/3)+32416*x+28364)/(1+x)^2)*RootOf(_Z^2-_Z+1)+ln(-(
1239*RootOf(_Z^2-_Z+1)^2*x^3-2478*RootOf(_Z^2-_Z+1)^2*x^2-4504*RootOf(_Z^2
-_Z+1)*(x^3+2)^(2/3)*x-4504*(x^3+2)^(1/3)*RootOf(_Z^2-_Z+1)*x^2-5743*RootO
f(_Z^2-_Z+1)*x^3-4956*RootOf(_Z^2-_Z+1)^2*x-9008*RootOf(_Z^2-_Z+1)*(x^3+2)
^(2/3)-18016*(x^3+2)^(1/3)*RootOf(_Z^2-_Z+1)*x-5860*RootOf(_Z^2-_Z+1)*x^2+
4839*x*(x^3+2)^(2/3)+4839*(x^3+2)^(1/3)*x^2+6078*x^3-18016*(x^3+2)^(1/3)...
```

Fricas [F(-2)]

Exception generated.

$$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx = \text{Exception raised: TypeError}$$

input `integrate((-1+x)/(1+x)/(x^3+2)^(1/3),x, algorithm="fricas")`

output `Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (residue poly has multiple non-linear factors)`

Sympy [F]

$$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{x-1}{(x+1)\sqrt[3]{x^3+2}} dx$$

input `integrate((-1+x)/(1+x)/(x**3+2)**(1/3),x)`

output `Integral((x - 1)/((x + 1)*(x**3 + 2)**(1/3)), x)`

Maxima [F]

$$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{x-1}{(x^3+2)^{\frac{1}{3}}(x+1)} dx$$

input `integrate((-1+x)/(1+x)/(x^3+2)^(1/3),x, algorithm="maxima")`

output `integrate((x - 1)/((x^3 + 2)^(1/3)*(x + 1)), x)`

Giac [F]

$$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{x-1}{(x^3+2)^{\frac{1}{3}}(x+1)} dx$$

input `integrate((-1+x)/(1+x)/(x^3+2)^(1/3),x, algorithm="giac")`

output `integrate((x - 1)/((x^3 + 2)^(1/3)*(x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{x-1}{(x^3+2)^{\frac{1}{3}}(x+1)} dx$$

input `int((x - 1)/((x^3 + 2)^(1/3)*(x + 1)),x)`

output `int((x - 1)/((x^3 + 2)^(1/3)*(x + 1)), x)`

Reduce [F]

$$\begin{aligned} & \int \frac{-1+x}{(1+x)\sqrt[3]{2+x^3}} dx \\ &= \int \frac{x}{(x^3+2)^{\frac{1}{3}}x + (x^3+2)^{\frac{1}{3}}} dx - \left(\int \frac{1}{(x^3+2)^{\frac{1}{3}}x + (x^3+2)^{\frac{1}{3}}} dx \right) \end{aligned}$$

input `int((-1+x)/(1+x)/(x^3+2)^(1/3),x)`

output `int(x/((x**3 + 2)**(1/3)*x + (x**3 + 2)**(1/3)),x) - int(1/((x**3 + 2)**(1/3)*x + (x**3 + 2)**(1/3)),x)`

3.93 $\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx$

Optimal result	707
Mathematica [F]	708
Rubi [A] (verified)	708
Maple [C] (verified)	710
Fricas [B] (verification not implemented)	711
Sympy [F]	712
Maxima [F]	712
Giac [F]	712
Mupad [F(-1)]	713
Reduce [F]	713

Optimal result

Integrand size = 15, antiderivative size = 108

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx = \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{2+x^3}}}{\sqrt{3}}\right)}{2\sqrt{3}} - \frac{1}{2}\sqrt{3}\arctan\left(\frac{1+\frac{2(2+x)}{\sqrt[3]{2+x^3}}}{\sqrt{3}}\right) - \frac{1}{2}\log(1+x) + \frac{3}{4}\log\left(2+x-\sqrt[3]{2+x^3}\right) - \frac{1}{4}\log\left(-x+\sqrt[3]{2+x^3}\right)$$

output

```
-1/2*ln(1+x)+3/4*ln(2+x-(x^3+2)^(1/3))-1/4*ln(-x+(x^3+2)^(1/3))+1/6*arctan
(1/3*(1+2*x/(x^3+2)^(1/3))*3^(1/2))*3^(1/2)-1/2*arctan(1/3*(1+2*(2+x)/(x^3
+2)^(1/3))*3^(1/2))*3^(1/2)
```

Mathematica [F]

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx$$

input `Integrate[1/((1 + x)*(2 + x^3)^(1/3)), x]`

output `Integrate[1/((1 + x)*(2 + x^3)^(1/3)), x]`

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.03, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2575, 769, 2576}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{1}{(x+1)\sqrt[3]{x^3+2}} dx \\ & \quad \downarrow \text{2575} \\ & \frac{1}{2} \int \frac{1}{\sqrt[3]{x^3+2}} dx + \frac{1}{2} \int \frac{1-x}{(x+1)\sqrt[3]{x^3+2}} dx \\ & \quad \downarrow \text{769} \\ & \frac{1}{2} \int \frac{1-x}{(x+1)\sqrt[3]{x^3+2}} dx + \frac{1}{2} \left(\frac{\arctan\left(\frac{\sqrt[3]{x^3+2}+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(\sqrt[3]{x^3+2}-x\right) \right) \\ & \quad \downarrow \text{2576} \end{aligned}$$

$$\frac{1}{2} \left(-\sqrt{3} \arctan \left(\frac{\frac{2(x+2)}{\sqrt[3]{x^3+2}} + 1}{\sqrt{3}} \right) + \frac{3}{2} \log \left(-\sqrt[3]{x^3+2} + x + 2 \right) - \log(x+1) \right) + \frac{1}{2} \left(\frac{\arctan \left(\frac{\frac{2x}{\sqrt[3]{x^3+2}} + 1}{\sqrt{3}} \right)}{\sqrt{3}} - \frac{1}{2} \log \left(\sqrt[3]{x^3+2} - x \right) \right)$$

input `Int[1/((1 + x)*(2 + x^3)^(1/3)),x]`

output `(- (Sqrt[3]*ArcTan[(1 + (2*(2 + x))/(2 + x^3)^(1/3)]/Sqrt[3]]) - Log[1 + x] + (3*Log[2 + x - (2 + x^3)^(1/3)]/2)/2 + (ArcTan[(1 + (2*x)/(2 + x^3)^(1/3)]/Sqrt[3])/Sqrt[3] - Log[-x + (2 + x^3)^(1/3)]/2)/2`

Defintions of rubi rules used

rule 769 `Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

rule 2575 `Int[1/(((c_) + (d_.)*(x_))*((a_) + (b_.)*(x_)^3)^(1/3)), x_Symbol] := Simp[1/(2*c) Int[1/(a + b*x^3)^(1/3), x], x] + Simp[1/(2*c) Int[(c - d*x)/((c + d*x)*(a + b*x^3)^(1/3)), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[2*b*c^3 - a*d^3, 0]`

rule 2576 `Int[((e_) + (f_.)*(x_))/(((c_) + (d_.)*(x_))*((a_) + (b_.)*(x_)^3)^(1/3)), x_Symbol] := Simp[Sqrt[3]*f*(ArcTan[(1 + 2*Rt[b, 3]*((2*c + d*x)/(d*(a + b*x^3)^(1/3))))/Sqrt[3]]/(Rt[b, 3]*d), x] + (Simp[(f*Log[c + d*x])/(Rt[b, 3]*d), x] - Simp[(3*f*Log[Rt[b, 3]*(2*c + d*x) - d*(a + b*x^3)^(1/3)]/(2*Rt[b, 3]*d), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[d*e + c*f, 0] && EqQ[2*b*c^3 - a*d^3, 0]`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 5.06 (sec) , antiderivative size = 1421, normalized size of antiderivative = 13.16

method	result	size
trager	Expression too large to display	1421

input `int(1/(1+x)/(x^3+2)^(1/3),x,method=_RETURNVERBOSE)`

output

```
1/6*ln((-4550781346817636-6728375859478224*x-68457312523761*x^6+1629509985
3018372*x*(x^3+2)^(2/3)-9125490357912936*(x^3+2)^(1/3)-234710785795752*x^4
-469421571591504*x^2+2151515536461060*x^3+4993190285176576*RootOf(_Z^2+_Z+
1)^2*x^3-15559137585059152*RootOf(_Z^2+_Z+1)-625895428788672*x^5+132631616
9500028*RootOf(_Z^2+_Z+1)^2*x^6+4346750471470680*RootOf(_Z^2+_Z+1)^2*x^5+1
53868976350327*RootOf(_Z^2+_Z+1)*x^6-928201890361806*(x^3+2)^(2/3)*x^4-540
325086981687*(x^3+2)^(1/3)*x^5-1959537324097146*(x^3+2)^(2/3)*x^3-48028896
6205944*(x^3+2)^(1/3)*x^4+5569211342170836*(x^3+2)^(2/3)*x^2+1200722415514
860*(x^3+2)^(1/3)*x^3+7115580883942020*RootOf(_Z^2+_Z+1)*(x^3+2)^(2/3)+337
2637147591320*RootOf(_Z^2+_Z+1)*(x^3+2)^(1/3)+4547369724000096*RootOf(_Z^2
+_Z+1)^2*x^4-868588603920114*RootOf(_Z^2+_Z+1)*x^5+527550776058264*RootOf(
_Z^2+_Z+1)*x^4+10107087250606332*(x^3+2)^(2/3)+8816461926585488*RootOf(_Z^
2+_Z+1)*x^3+1055101552116528*RootOf(_Z^2+_Z+1)*x^2-21283128527537520*RootO
f(_Z^2+_Z+1)*x-14648813469281292*x*(x^3+2)^(1/3)+9094739448000192*RootOf(_
Z^2+_Z+1)^2*x^2+5884831407529536*RootOf(_Z^2+_Z+1)^2*x+36303984101745*Root
Of(_Z^2+_Z+1)^2*(x^3+2)^(2/3)*x^4+288449229728205*RootOf(_Z^2+_Z+1)^2*(x^3
+2)^(1/3)*x^5-1306943427662820*RootOf(_Z^2+_Z+1)^2*(x^3+2)^(2/3)*x^3+12869
27332633530*RootOf(_Z^2+_Z+1)^2*(x^3+2)^(1/3)*x^4-601904942144643*RootOf(_
Z^2+_Z+1)*(x^3+2)^(2/3)*x^4-580773949503594*RootOf(_Z^2+_Z+1)*(x^3+2)^(1/3
)*x^5-3775614346581480*RootOf(_Z^2+_Z+1)^2*(x^3+2)^(2/3)*x^2+1420057746...
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 267 vs. $2(84) = 168$.

Time = 0.77 (sec) , antiderivative size = 267, normalized size of antiderivative = 2.47

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx$$

$$= \frac{1}{6} \sqrt{3} \arctan \left(\frac{13910019318573948542 \sqrt{3}(7114781247 x^4 + 13663058416 x^3 - 46178206896 x^2 - 126842559344 x - 77084338088)(x^3 + 2)^{2/3} - 27820038637147897084 \sqrt{3}(1625757424 x^5 + 16302821713 x^4 + 26102613730 x^3 - 26431113242 x^2 - 80188343316 x - 42779182428)(x^3 + 2)^{1/3} + \sqrt{3}(93292570833559435663132301885 x^6 + 382151535711085278859235047618 x^5 + 673924074224408772959625384792 x^4 + 889426563183087468015580290048 x^3 + 888876515195959220955879945824 x^2 + 351260598258508240019971964880 x - 47674000995597211057816884304)}{(78905434814564721745708464883 x^6 + 337746705836458222863347934450 x^5 + 15598952776058587894336070976 x^4 - 895430525315100108684787964824 x^3 + 361667862240477028869533375352 x^2 + 2541802301011632510645972090336 x + 1554815286823334092314485968880)} \right) + \frac{1}{12} \log \left(\frac{(22 x^6 + 6 x^5 - 48 x^4 + 44 x^3 + 24 x^2 + 3(7 x^4 - 2 x^3 - 32 x^2 - 20 x + 4)(x^3 + 2)^{2/3} + 3(7 x^5 - 16 x^3 + 34 x^2 + 76 x + 32)(x^3 + 2)^{1/3} - 192 x - 140)(x^6 + 6 x^5 + 15 x^4 + 20 x^3 + 15 x^2 + 6 x + 1)}{x^6 + 6 x^5 + 15 x^4 + 20 x^3 + 15 x^2 + 6 x + 1} \right)$$

input `integrate(1/(1+x)/(x^3+2)^(1/3),x, algorithm="fricas")`

output

```
1/6*sqrt(3)*arctan(1/3*(13910019318573948542*sqrt(3)*(7114781247*x^4 + 136
63058416*x^3 - 46178206896*x^2 - 126842559344*x - 77084338088)*(x^3 + 2)^(
2/3) - 27820038637147897084*sqrt(3)*(1625757424*x^5 + 16302821713*x^4 + 26
102613730*x^3 - 26431113242*x^2 - 80188343316*x - 42779182428)*(x^3 + 2)^(
1/3) + sqrt(3)*(93292570833559435663132301885*x^6 + 3821515357110852788592
35047618*x^5 + 673924074224408772959625384792*x^4 + 8894265631830874680155
80290048*x^3 + 888876515195959220955879945824*x^2 + 3512605982585082400199
71964880*x - 47674000995597211057816884304))/(7890543481456472174570846488
3*x^6 + 337746705836458222863347934450*x^5 + 15598952776058587894336070976
*x^4 - 895430525315100108684787964824*x^3 + 361667862240477028869533375352
*x^2 + 2541802301011632510645972090336*x + 1554815286823334092314485968880
)) + 1/12*log((22*x^6 + 6*x^5 - 48*x^4 + 44*x^3 + 24*x^2 + 3*(7*x^4 - 2*x^
3 - 32*x^2 - 20*x + 4)*(x^3 + 2)^(2/3) + 3*(7*x^5 - 16*x^3 + 34*x^2 + 76*x
+ 32)*(x^3 + 2)^(1/3) - 192*x - 140)/(x^6 + 6*x^5 + 15*x^4 + 20*x^3 + 15*
x^2 + 6*x + 1))
```


Sympy [F]

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{1}{(x+1)\sqrt[3]{x^3+2}} dx$$

input `integrate(1/(1+x)/(x**3+2)**(1/3),x)`

output `Integral(1/((x + 1)*(x**3 + 2)**(1/3)), x)`

Maxima [F]

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{1}{(x^3+2)^{\frac{1}{3}}(x+1)} dx$$

input `integrate(1/(1+x)/(x^3+2)^(1/3),x, algorithm="maxima")`

output `integrate(1/((x^3 + 2)^(1/3)*(x + 1)), x)`

Giac [F]

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{1}{(x^3+2)^{\frac{1}{3}}(x+1)} dx$$

input `integrate(1/(1+x)/(x^3+2)^(1/3),x, algorithm="giac")`

output `integrate(1/((x^3 + 2)^(1/3)*(x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{1}{(x^3+2)^{1/3}(x+1)} dx$$

input `int(1/((x^3 + 2)^(1/3)*(x + 1)),x)`output `int(1/((x^3 + 2)^(1/3)*(x + 1)), x)`**Reduce [F]**

$$\int \frac{1}{(1+x)\sqrt[3]{2+x^3}} dx = \int \frac{1}{(x^3+2)^{1/3}x + (x^3+2)^{1/3}} dx$$

input `int(1/(1+x)/(x^3+2)^(1/3),x)`output `int(1/((x**3 + 2)**(1/3)*x + (x**3 + 2)**(1/3)),x)`

3.94 $\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx$

Optimal result	714
Mathematica [C] (verified)	715
Rubi [A] (verified)	715
Maple [A] (verified)	716
Fricas [B] (verification not implemented)	717
Sympy [F]	718
Maxima [F]	718
Giac [F]	718
Mupad [F(-1)]	719
Reduce [F]	719

Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx = \frac{\arctan\left(\frac{1+\frac{2\sqrt[3]{a+bx^3}}{\sqrt[3]{a+b}}}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{a+b}} + \frac{\log(1-x^3)}{6\sqrt[3]{a+b}} - \frac{\log\left(\sqrt[3]{a+bx} - \sqrt[3]{a+bx^3}\right)}{2\sqrt[3]{a+b}}$$

output

```
1/6*ln(-x^3+1)/(a+b)^(1/3)-1/2*ln((a+b)^(1/3)*x-(b*x^3+a)^(1/3))/(a+b)^(1/3)+1/3*arctan(1/3*(1+2*(a+b)^(1/3)*x/(b*x^3+a)^(1/3))*3^(1/2))/(a+b)^(1/3)*3^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.88 (sec) , antiderivative size = 189, normalized size of antiderivative = 1.93

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx$$

$$= \frac{-2\sqrt{-6+6i\sqrt{3}} \arctan\left(\frac{3\sqrt[3]{a+bx}}{\sqrt{3}\sqrt[3]{a+bx}-(3i+\sqrt{3})\sqrt[3]{a+bx^3}}\right) + (1+i\sqrt{3})\left(2\log\left(2\sqrt[3]{a+bx} + (1+i\sqrt{3})\sqrt[3]{a+bx^3}\right)\right)}{12\sqrt[3]{a+b}}$$

input `Integrate[1/((1 - x^3)*(a + b*x^3)^(1/3)),x]`

output `(-2*Sqrt[-6 + (6*I)*Sqrt[3]]*ArcTan[(3*(a + b)^(1/3)*x)/(Sqrt[3]*(a + b)^(1/3)*x - (3*I + Sqrt[3])*(a + b*x^3)^(1/3)]) + (1 + I*Sqrt[3])*(2*Log[2*(a + b)^(1/3)*x + (1 + I*Sqrt[3])*(a + b*x^3)^(1/3)] - Log[-((a + b)^(1/3)*x) + (a + b*x^3)^(1/3)]*(2*I)*(a + b)^(1/3)*x + (I + Sqrt[3])*(a + b*x^3)^(1/3)))/(12*(a + b)^(1/3))`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {901}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx$$

↓ 901

$$\frac{\arctan\left(\frac{\frac{2x\sqrt[3]{a+b}+1}{\sqrt[3]{a+bx^3}}}{\sqrt{3}}\right)}{\sqrt{3}\sqrt[3]{a+b}} + \frac{\log(1-x^3)}{6\sqrt[3]{a+b}} - \frac{\log\left(x\sqrt[3]{a+b} - \sqrt[3]{a+bx^3}\right)}{2\sqrt[3]{a+b}}$$

input `Int[1/((1 - x^3)*(a + b*x^3)^(1/3)),x]`

output `ArcTan[(1 + (2*(a + b)^(1/3)*x)/(a + b*x^3)^(1/3))/Sqrt[3]]/(Sqrt[3]*(a + b)^(1/3)) + Log[1 - x^3]/(6*(a + b)^(1/3)) - Log[(a + b)^(1/3)*x - (a + b*x^3)^(1/3)]/(2*(a + b)^(1/3))`

Defintions of rubi rules used

rule 901 `Int[1/(((a_) + (b_.)*(x_)^3)^(1/3)*((c_) + (d_.)*(x_)^3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/c, 3]}, Simp[ArcTan[(1 + (2*q*x)/(a + b*x^3)^(1/3))/Sqrt[3]]/(Sqrt[3]*c*q), x] + (-Simp[Log[q*x - (a + b*x^3)^(1/3)]/(2*c*q), x] + Simp[Log[c + d*x^3]/(6*c*q), x])] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

Maple [A] (verified)

Time = 1.14 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.13

method	result
pseudoelliptic	$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left((a+b)^{\frac{1}{3}}x+2(bx^3+a)^{\frac{1}{3}}\right)}{3(a+b)^{\frac{1}{3}}x}\right) + \ln\left(\frac{-(a+b)^{\frac{1}{3}}x+(bx^3+a)^{\frac{1}{3}}}{x}\right) - \frac{\ln\left(\frac{(a+b)^{\frac{2}{3}}x^2+(a+b)^{\frac{1}{3}}(bx^3+a)^{\frac{1}{3}}x+(bx^3+a)}{x^2}\right)}{2}}{3(a+b)^{\frac{1}{3}}}$

input `int(1/(-x^3+1)/(b*x^3+a)^(1/3),x,method=_RETURNVERBOSE)`

output `-1/3/(a+b)^(1/3)*(3^(1/2)*arctan(1/3*3^(1/2)*((a+b)^(1/3)*x+2*(b*x^3+a)^(1/3))/(a+b)^(1/3)/x)+ln((-a+b)^(1/3)*x+(b*x^3+a)^(1/3))/x-1/2*ln(((a+b)^(2/3)*x^2+(a+b)^(1/3)*(b*x^3+a)^(1/3)*x+(b*x^3+a)^(2/3))/x^2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 559 vs. $2(78) = 156$.

Time = 101.79 (sec) , antiderivative size = 1252, normalized size of antiderivative = 12.78

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx = \text{Too large to display}$$

input `integrate(1/(-x^3+1)/(b*x^3+a)^(1/3),x, algorithm="fricas")`

output

```
[1/18*(3*sqrt(1/3)*(a + b)*sqrt((-a - b)^(1/3)/(a + b))*log(-((a^3 - 27*a^2*b - 108*a*b^2 - 81*b^3)*x^9 - 3*(10*a^3 + 54*a^2*b + 45*a*b^2)*x^6 - 3*(17*a^3 + 18*a^2*b)*x^3 - a^3 + 9*((2*a^2 + 3*a*b)*x^7 - (a^2 + 3*a*b)*x^4 - a^2*x)*(b*x^3 + a)^(2/3))*(-a - b)^(1/3) + 9*((a^2 + 9*a*b + 9*b^2)*x^8 + (7*a^2 + 9*a*b)*x^5 + a^2*x^2)*(b*x^3 + a)^(1/3))*(-a - b)^(2/3) + 3*sqrt(1/3)*(3*((4*a^2 + 21*a*b + 18*b^2)*x^7 + (13*a^2 + 15*a*b)*x^4 + a^2*x)*(b*x^3 + a)^(2/3))*(-a - b)^(2/3) + 3*((a^3 - 2*a^2*b - 12*a*b^2 - 9*b^3)*x^8 - 5*(a^3 + 4*a^2*b + 3*a*b^2)*x^5 - 5*(a^3 + a^2*b)*x^2)*(b*x^3 + a)^(1/3) + ((a^3 + 27*a^2*b + 54*a*b^2 + 27*b^3)*x^9 + 3*(8*a^3 + 18*a^2*b + 9*a*b^2)*x^6 + 3*a^3*x^3 - a^3))*(-a - b)^(1/3))*sqrt((-a - b)^(1/3)/(a + b)))/(x^9 - 3*x^6 + 3*x^3 - 1) - 2*(-a - b)^(2/3)*log(-3*(b*x^3 + a)^(1/3)*(a + b))*(-a - b)^(1/3)*x^2 + 3*(b*x^3 + a)^(2/3)*(a + b)*x + (a*x^3 - a))*(-a - b)^(2/3))/(x^3 - 1) + (-a - b)^(2/3)*log((3*((2*a + 3*b)*x^4 + a*x)*(b*x^3 + a)^(2/3))*(-a - b)^(2/3) + 3*((a^2 + 4*a*b + 3*b^2)*x^5 + 2*(a^2 + a*b)*x^2)*(b*x^3 + a)^(1/3) - ((a^2 + 9*a*b + 9*b^2)*x^6 + (7*a^2 + 9*a*b)*x^3 + a^2))*(-a - b)^(1/3))/(x^6 - 2*x^3 + 1)))/(a + b), 1/18*(6*sqrt(1/3)*(a + b)*sqrt(-(-a - b)^(1/3)/(a + b))*arctan(sqrt(1/3)*(6*((2*a^2 + 3*a*b)*x^7 - (a^2 + 3*a*b)*x^4 - a^2*x)*(b*x^3 + a)^(2/3))*(-a - b)^(2/3) - 6*((a^3 + 10*a^2*b + 18*a*b^2 + 9*b^3)*x^8 + (7*a^3 + 16*a^2*b + 9*a*b^2)*x^5 + (a^3 + a^2*b)*x^2)*(b*x^3 + a)^(1/3) - ((a^3 - 9*a^2*b - 36*a*b^2 - 27...
```

Sympy [F]

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx = - \int \frac{1}{x^3\sqrt[3]{a+bx^3} - \sqrt[3]{a+bx^3}} dx$$

input `integrate(1/(-x**3+1)/(b*x**3+a)**(1/3),x)`

output `-Integral(1/(x**3*(a + b*x**3)**(1/3) - (a + b*x**3)**(1/3)), x)`

Maxima [F]

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx = \int -\frac{1}{(bx^3+a)^{\frac{1}{3}}(x^3-1)} dx$$

input `integrate(1/(-x^3+1)/(b*x^3+a)^(1/3),x, algorithm="maxima")`

output `-integrate(1/((b*x^3 + a)^(1/3)*(x^3 - 1)), x)`

Giac [F]

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx = \int -\frac{1}{(bx^3+a)^{\frac{1}{3}}(x^3-1)} dx$$

input `integrate(1/(-x^3+1)/(b*x^3+a)^(1/3),x, algorithm="giac")`

output `integrate(-1/((b*x^3 + a)^(1/3)*(x^3 - 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx = - \int \frac{1}{(x^3-1)(bx^3+a)^{1/3}} dx$$

input `int(-1/((x^3 - 1)*(a + b*x^3)^(1/3)),x)`output `-int(1/((x^3 - 1)*(a + b*x^3)^(1/3)), x)`**Reduce [F]**

$$\int \frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} dx = - \left(\int \frac{1}{(bx^3+a)^{1/3} x^3 - (bx^3+a)^{1/3}} dx \right)$$

input `int(1/(-x^3+1)/(b*x^3+a)^(1/3),x)`output `- int(1/((a + b*x**3)**(1/3)*x**3 - (a + b*x**3)**(1/3)),x)`

3.95 $\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx$

Optimal result	720
Mathematica [F]	721
Rubi [A] (verified)	721
Maple [F]	722
Fricas [F(-1)]	722
Sympy [F]	723
Maxima [F]	723
Giac [F]	723
Mupad [F(-1)]	724
Reduce [F]	724

Optimal result

Integrand size = 23, antiderivative size = 154

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \frac{\arctan\left(\frac{1+\frac{2\sqrt[3]{a+bx^3}}{\sqrt[3]{a+b}}}{\sqrt[3]{a+b}}\right)}{\sqrt[3]{a+b}} + \frac{\arctan\left(\frac{1+\frac{2\sqrt[3]{a+bx^3}}{\sqrt[3]{a+b}}}{\sqrt[3]{a+b}}\right)}{\sqrt[3]{a+b}}$$

$$+ \frac{\log\left(\frac{\sqrt[3]{a+b}-\sqrt[3]{a+bx^3}}{2\sqrt[3]{a+b}}\right)}{\log\left(\frac{\sqrt[3]{a+b}-\sqrt[3]{a+bx^3}}{2\sqrt[3]{a+b}}\right)}$$

output

```
1/2*ln((a+b)^(1/3)-(b*x^3+a)^(1/3))/(a+b)^(1/3)-1/2*ln((a+b)^(1/3)*x-(b*x^3+a)^(1/3))/(a+b)^(1/3)+1/3*arctan(1/3*(1+2*(a+b)^(1/3)*x/(b*x^3+a)^(1/3))*3^(1/2))/(a+b)^(1/3)*3^(1/2)+1/3*arctan(1/3*(1+2*(b*x^3+a)^(1/3)/(a+b)^(1/3))*3^(1/2))/(a+b)^(1/3)*3^(1/2)
```

Mathematica [F]

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx$$

input `Integrate[(1 + x)/((1 + x + x^2)*(a + b*x^3)^(1/3)),x]`

output `Integrate[(1 + x)/((1 + x + x^2)*(a + b*x^3)^(1/3)), x]`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+1}{(x^2+x+1)\sqrt[3]{a+bx^3}} dx$$

↓ 2583

$$\int \left(\frac{1}{(1-x^3)\sqrt[3]{a+bx^3}} - \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} \right) dx$$

↓ 2009

$$\frac{\arctan\left(\frac{\frac{2x\sqrt[3]{a+b}}{\sqrt[3]{a+bx^3}}+1}{\sqrt{3}\sqrt[3]{a+b}}\right)}{\sqrt{3}\sqrt[3]{a+b}} + \frac{\arctan\left(\frac{\frac{2\sqrt[3]{a+bx^3}}{\sqrt[3]{a+b}}+1}{\sqrt{3}\sqrt[3]{a+b}}\right)}{\log\left(x\sqrt[3]{a+b}-\sqrt[3]{a+bx^3}\right)} + \frac{\log\left(\sqrt[3]{a+b}-\sqrt[3]{a+bx^3}\right)}{2\sqrt[3]{a+b}}$$

input `Int[(1 + x)/((1 + x + x^2)*(a + b*x^3)^(1/3)),x]`

output

```
ArcTan[(1 + (2*(a + b)^(1/3)*x)/(a + b*x^3)^(1/3))/Sqrt[3]]/(Sqrt[3]*(a +
b)^(1/3)) + ArcTan[(1 + (2*(a + b*x^3)^(1/3))/(a + b)^(1/3))/Sqrt[3]]/(Sqr
t[3]*(a + b)^(1/3)) + Log[(a + b)^(1/3) - (a + b*x^3)^(1/3)]/(2*(a + b)^(1
/3)) - Log[(a + b)^(1/3)*x - (a + b*x^3)^(1/3)]/(2*(a + b)^(1/3))
```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 2583

```
Int[(Px_)*((c_) + (d_)*(x_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^3)^(p
_), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b
*x^3)^p, Px/(c - d*x)^q, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && Poly
Q[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denomina
tor[p], 3]
```

Maple [F]

$$\int \frac{1+x}{(x^2+x+1)(bx^3+a)^{\frac{1}{3}}} dx$$

input

```
int((1+x)/(x^2+x+1)/(b*x^3+a)^(1/3),x)
```

output

```
int((1+x)/(x^2+x+1)/(b*x^3+a)^(1/3),x)
```

Fricas [F(-1)]

Timed out.

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \text{Timed out}$$

input

```
integrate((1+x)/(x^2+x+1)/(b*x^3+a)^(1/3),x, algorithm="fricas")
```

output

```
Timed out
```

Sympy [F]

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \int \frac{x+1}{\sqrt[3]{a+bx^3}(x^2+x+1)} dx$$

input `integrate((1+x)/(x**2+x+1)/(b*x**3+a)**(1/3),x)`

output `Integral((x + 1)/((a + b*x**3)**(1/3)*(x**2 + x + 1)), x)`

Maxima [F]

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \int \frac{x+1}{(bx^3+a)^{\frac{1}{3}}(x^2+x+1)} dx$$

input `integrate((1+x)/(x^2+x+1)/(b*x^3+a)^(1/3),x, algorithm="maxima")`

output `integrate((x + 1)/((b*x^3 + a)^(1/3)*(x^2 + x + 1)), x)`

Giac [F]

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \int \frac{x+1}{(bx^3+a)^{\frac{1}{3}}(x^2+x+1)} dx$$

input `integrate((1+x)/(x^2+x+1)/(b*x^3+a)^(1/3),x, algorithm="giac")`

output `integrate((x + 1)/((b*x^3 + a)^(1/3)*(x^2 + x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \int \frac{x+1}{(bx^3+a)^{1/3}(x^2+x+1)} dx$$

input `int((x + 1)/((a + b*x^3)^(1/3)*(x + x^2 + 1)),x)`output `int((x + 1)/((a + b*x^3)^(1/3)*(x + x^2 + 1)), x)`**Reduce [F]**

$$\int \frac{1+x}{(1+x+x^2)\sqrt[3]{a+bx^3}} dx = \int \frac{x}{(bx^3+a)^{1/3}x^2 + (bx^3+a)^{1/3}x + (bx^3+a)^{1/3}} dx + \int \frac{1}{(bx^3+a)^{1/3}x^2 + (bx^3+a)^{1/3}x + (bx^3+a)^{1/3}} dx$$

input `int((1+x)/(x^2+x+1)/(b*x^3+a)^(1/3),x)`output `int(x/((a + b*x**3)**(1/3)*x**2 + (a + b*x**3)**(1/3)*x + (a + b*x**3)**(1/3)),x) + int(1/((a + b*x**3)**(1/3)*x**2 + (a + b*x**3)**(1/3)*x + (a + b*x**3)**(1/3)),x)`

3.96 $\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx$

Optimal result	725
Mathematica [C] (verified)	726
Rubi [A] (verified)	726
Maple [A] (verified)	728
Fricas [B] (verification not implemented)	729
Sympy [F]	730
Maxima [A] (verification not implemented)	730
Giac [A] (verification not implemented)	731
Mupad [B] (verification not implemented)	732
Reduce [F]	732

Optimal result

Integrand size = 24, antiderivative size = 96

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx = -\frac{\arctan\left(\frac{1+2\sqrt[3]{a+bx^3}}{\sqrt[3]{a+b}}\right)}{\sqrt{3}\sqrt[3]{a+b}} + \frac{\log(1-x^3)}{6\sqrt[3]{a+b}} - \frac{\log\left(\sqrt[3]{a+b}-\sqrt[3]{a+bx^3}\right)}{2\sqrt[3]{a+b}}$$

output

```
1/6*ln(-x^3+1)/(a+b)^(1/3)-1/2*ln((a+b)^(1/3)-(b*x^3+a)^(1/3))/(a+b)^(1/3)
-1/3*arctan(1/3*(1+2*(b*x^3+a)^(1/3)/(a+b)^(1/3))*3^(1/2))/(a+b)^(1/3)*3^(1/2)
```

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.90

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx$$

$$= \frac{2\sqrt{-6+6i\sqrt{3}} \arctan\left(\frac{1+\frac{(-1-i\sqrt{3})\sqrt[3]{a+bx^3}}{\sqrt[3]{a+b}}}{\sqrt[3]{a+b}}\right) - i(-i+\sqrt{3}) \left(\log\left(\left(\sqrt[3]{a+b}-\sqrt[3]{a+bx^3}\right)\left(2\sqrt[3]{a+b}+\sqrt[3]{a+bx^3}\right)\right)\right)}{12\sqrt[3]{a+b}}$$

input

```
Integrate[x^2/((1-x^3)*(a+b*x^3)^(1/3)),x]
```

output

```
(2*Sqrt[-6+(6*I)*Sqrt[3]]*ArcTan[(1+((-1-I*Sqrt[3])*(a+b*x^3)^(1/3)))/(a+b)^(1/3)]/Sqrt[3]-I*(-I+Sqrt[3])*(Log[((a+b)^(1/3)-(a+b*x^3)^(1/3))*(2*(a+b)^(1/3)+(a+b*x^3)^(1/3)-I*Sqrt[3]*(a+b*x^3)^(1/3))]-2*Log[2*(a+b)^(1/3)+(1+I*Sqrt[3])*(a+b*x^3)^(1/3)]))/(12*(a+b)^(1/3))
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.04, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {946, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx$$

$$\downarrow \text{946}$$

$$\frac{1}{3} \int \frac{1}{(1-x^3)\sqrt[3]{bx^3+a}} dx^3$$

↓ 67

$$\frac{1}{3} \left(\frac{3 \int \frac{1}{\sqrt[3]{a+b} - \sqrt[3]{bx^3+a}} d\sqrt[3]{bx^3+a}}{2\sqrt[3]{a+b}} - \frac{3}{2} \int \frac{1}{x^6 + (a+b)^{2/3} + \sqrt[3]{a+b}\sqrt[3]{bx^3+a}} d\sqrt[3]{bx^3+a} + \frac{\log(1-x^3)}{2\sqrt[3]{a+b}} \right)$$

↓ 16

$$\frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{x^6 + (a+b)^{2/3} + \sqrt[3]{a+b}\sqrt[3]{bx^3+a}} d\sqrt[3]{bx^3+a} + \frac{\log(1-x^3)}{2\sqrt[3]{a+b}} - \frac{3 \log(\sqrt[3]{a+b} - \sqrt[3]{a+bx^3})}{2\sqrt[3]{a+b}} \right)$$

↓ 1082

$$\frac{1}{3} \left(\frac{3 \int \frac{1}{-x^6-3} d\left(\frac{2\sqrt[3]{bx^3+a}}{\sqrt[3]{a+b}} + 1\right)}{\sqrt[3]{a+b}} + \frac{\log(1-x^3)}{2\sqrt[3]{a+b}} - \frac{3 \log(\sqrt[3]{a+b} - \sqrt[3]{a+bx^3})}{2\sqrt[3]{a+b}} \right)$$

↓ 217

$$\frac{1}{3} \left(-\frac{\sqrt{3} \arctan\left(\frac{2\sqrt[3]{a+bx^3} + 1}{\sqrt[3]{a+b}}\right)}{\sqrt[3]{a+b}} + \frac{\log(1-x^3)}{2\sqrt[3]{a+b}} - \frac{3 \log(\sqrt[3]{a+b} - \sqrt[3]{a+bx^3})}{2\sqrt[3]{a+b}} \right)$$

input

```
Int[x^2/((1 - x^3)*(a + b*x^3)^(1/3)),x]
```

output

```
(-((Sqrt[3]*ArcTan[(1 + (2*(a + b*x^3)^(1/3))/(a + b)^(1/3)]/Sqrt[3]))/(a + b)^(1/3)) + Log[1 - x^3]/(2*(a + b)^(1/3)) - (3*Log[(a + b)^(1/3) - (a + b*x^3)^(1/3)])/(2*(a + b)^(1/3)))/3
```


Defintions of rubi rules used

rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /;$ $\text{FreeQ}\{a, b, c, x\}$

rule 67 $\text{Int}[1/(((a_)+(b_)*(x_))*((c_)+(d_)*(x_))^{(1/3)}), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]]/(2*b*q), x] + (\text{Simp}[3/(2*b) \text{ Subst}[\text{Int}[1/(q^2 + q*x + x^2), x], x, (c + d*x)^{(1/3)}], x] - \text{Simp}[3/(2*b*q) \text{ Subst}[\text{Int}[1/(q - x), x], x, (c + d*x)^{(1/3)}], x]) /;$ $\text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{PosQ}[(b*c - a*d)/b]$

rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

rule 946 $\text{Int}[(x_)^{(m_)}*((a_)+(b_)*(x_)^{(n_)})^{(p_)}*((c_)+(d_)*(x_)^{(n_)})^{(q_)}, x_Symbol] \rightarrow \text{Simp}[1/n \text{ Subst}[\text{Int}[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, c, d, m, n, p, q, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[m - n + 1, 0]$

rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /;$ $\text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c]) /;$ $\text{FreeQ}\{a, b, c, x\}$

Maple [A] (verified)

Time = 0.83 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.96

method	result	si
pseudoelliptic	$\frac{\arctan\left(\frac{\left((a+b)^{\frac{1}{3}}+2(bx^3+a)^{\frac{1}{3}}\right)\sqrt{3}}{3(a+b)^{\frac{1}{3}}}\right)\sqrt{3}+\ln\left((bx^3+a)^{\frac{1}{3}}-(a+b)^{\frac{1}{3}}\right)-\frac{\ln\left((bx^3+a)^{\frac{2}{3}}+(a+b)^{\frac{1}{3}}(bx^3+a)^{\frac{1}{3}}+(a+b)^{\frac{2}{3}}\right)}{2}}{3(a+b)^{\frac{1}{3}}}$	9

input `int(x^2/(-x^3+1)/(b*x^3+a)^(1/3),x,method=_RETURNVERBOSE)`

output `-1/3/(a+b)^(1/3)*(arctan(1/3*((a+b)^(1/3)+2*(b*x^3+a)^(1/3))*3^(1/2)/(a+b)^(1/3))*3^(1/2)+ln((b*x^3+a)^(1/3)-(a+b)^(1/3))-1/2*ln((b*x^3+a)^(2/3)+(a+b)^(1/3)*(b*x^3+a)^(1/3)+(a+b)^(2/3)))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(75) = 150$.

Time = 0.09 (sec) , antiderivative size = 387, normalized size of antiderivative = 4.03

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx$$

$$= \frac{3\sqrt{\frac{1}{3}}(a+b)\sqrt{\frac{(-a-b)^{\frac{1}{3}}}{a+b}} \log\left(\frac{2bx^3+3\sqrt{\frac{1}{3}}\left((bx^3+a)^{\frac{1}{3}}(a+b)-(a+b)(-a-b)^{\frac{1}{3}}-2(bx^3+a)^{\frac{2}{3}}(-a-b)^{\frac{2}{3}}\right)\sqrt{\frac{(-a-b)^{\frac{1}{3}}}{a+b}}+3a-3(bx^3+a)}{x^3-1}}\right)}{6\sqrt{\frac{1}{3}}(a+b)\sqrt{-\frac{(-a-b)^{\frac{1}{3}}}{a+b}} \arctan\left(\sqrt{\frac{1}{3}}\left(2(bx^3+a)^{\frac{1}{3}}-(-a-b)^{\frac{1}{3}}\right)\sqrt{-\frac{(-a-b)^{\frac{1}{3}}}{a+b}}\right)-(-a-b)^{\frac{2}{3}} \log\left((bx^3+a)^{\frac{1}{3}}-(-a-b)^{\frac{1}{3}}\right)}{6(a+b)}}$$

input `integrate(x^2/(-x^3+1)/(b*x^3+a)^(1/3),x, algorithm="fricas")`

output

```
[1/6*(3*sqrt(1/3)*(a + b)*sqrt((-a - b)^(1/3)/(a + b))*log((2*b*x^3 + 3*sqrt(1/3)*((b*x^3 + a)^(1/3)*(a + b) - (a + b)*(-a - b)^(1/3) - 2*(b*x^3 + a)^(2/3)*(-a - b)^(2/3))*sqrt((-a - b)^(1/3)/(a + b)) + 3*a - 3*(b*x^3 + a)^(1/3)*(-a - b)^(2/3) + b)/(x^3 - 1)) + (-a - b)^(2/3)*log((b*x^3 + a)^(2/3) - (b*x^3 + a)^(1/3)*(-a - b)^(1/3) + (-a - b)^(2/3)) - 2*(-a - b)^(2/3)*log((b*x^3 + a)^(1/3) + (-a - b)^(1/3)))/(a + b), -1/6*(6*sqrt(1/3)*(a + b)*sqrt(-(-a - b)^(1/3)/(a + b))*arctan(sqrt(1/3)*(2*(b*x^3 + a)^(1/3) - (-a - b)^(1/3))*sqrt(-(-a - b)^(1/3)/(a + b))) - (-a - b)^(2/3)*log((b*x^3 + a)^(2/3) - (b*x^3 + a)^(1/3)*(-a - b)^(1/3) + (-a - b)^(2/3)) + 2*(-a - b)^(2/3)*log((b*x^3 + a)^(1/3) + (-a - b)^(1/3)))/(a + b)]
```

Sympy [F]

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx = - \int \frac{x^2}{x^3\sqrt[3]{a+bx^3} - \sqrt[3]{a+bx^3}} dx$$

input

```
integrate(x**2/(-x**3+1)/(b*x**3+a)**(1/3), x)
```

output

```
-Integral(x**2/(x**3*(a + b*x**3)**(1/3) - (a + b*x**3)**(1/3)), x)
```

Maxima [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.15

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx = \frac{2\sqrt{3}b \arctan\left(\frac{\sqrt{3}\left(2(bx^3+a)^{\frac{1}{3}}+(a+b)^{\frac{1}{3}}\right)}{3(a+b)^{\frac{1}{3}}}\right)}{(a+b)^{\frac{1}{3}}} - \frac{b \log\left(\frac{(bx^3+a)^{\frac{2}{3}}+(bx^3+a)^{\frac{1}{3}}(a+b)^{\frac{1}{3}}+(a+b)^{\frac{2}{3}}}{(a+b)^{\frac{1}{3}}}\right)}{(a+b)^{\frac{1}{3}}} + \frac{2b \log\left(\frac{(bx^3+a)^{\frac{1}{3}}-(a+b)^{\frac{1}{3}}}{(a+b)^{\frac{1}{3}}}\right)}{(a+b)^{\frac{1}{3}}}$$

$6b$

input

```
integrate(x^2/(-x^3+1)/(b*x^3+a)^(1/3), x, algorithm="maxima")
```

output

```
-1/6*(2*sqrt(3)*b*arctan(1/3*sqrt(3)*(2*(b*x^3 + a)^(1/3) + (a + b)^(1/3))
/(a + b)^(1/3))/(a + b)^(1/3) - b*log((b*x^3 + a)^(2/3) + (b*x^3 + a)^(1/3)
)*(a + b)^(1/3) + (a + b)^(2/3))/(a + b)^(1/3) + 2*b*log((b*x^3 + a)^(1/3)
- (a + b)^(1/3))/(a + b)^(1/3))/b
```

Giac [A] (verification not implemented)

Time = 3.44 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.18

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx = -\frac{(a+b)^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}(2(bx^3+a)^{\frac{1}{3}}+(a+b)^{\frac{1}{3}})}{3(a+b)^{\frac{1}{3}}}\right)}{\sqrt{3}a + \sqrt{3}b} + \frac{\log\left((bx^3+a)^{\frac{2}{3}} + (bx^3+a)^{\frac{1}{3}}(a+b)^{\frac{1}{3}} + (a+b)^{\frac{2}{3}}\right)}{6(a+b)^{\frac{1}{3}}} - \frac{\log\left(\left|(bx^3+a)^{\frac{1}{3}} - (a+b)^{\frac{1}{3}}\right|\right)}{3(a+b)^{\frac{1}{3}}}$$

input

```
integrate(x^2/(-x^3+1)/(b*x^3+a)^(1/3),x, algorithm="giac")
```

output

```
-(a + b)^(2/3)*arctan(1/3*sqrt(3)*(2*(b*x^3 + a)^(1/3) + (a + b)^(1/3))/(a
+ b)^(1/3))/(sqrt(3)*a + sqrt(3)*b) + 1/6*log((b*x^3 + a)^(2/3) + (b*x^3
+ a)^(1/3)*(a + b)^(1/3) + (a + b)^(2/3))/(a + b)^(1/3) - 1/3*log(abs((b*x
^3 + a)^(1/3) - (a + b)^(1/3)))/(a + b)^(1/3)
```

Mupad [B] (verification not implemented)

Time = 0.36 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.64

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx = \frac{\ln\left((bx^3+a)^{1/3} - \frac{9a+9b}{9(-a-b)^{2/3}}\right)}{3(-a-b)^{1/3}} + \frac{\ln\left((bx^3+a)^{1/3} - \frac{(-1+\sqrt{3}i)^2(9a+9b)}{36(-a-b)^{2/3}}\right)(-1+\sqrt{3}i)}{6(-a-b)^{1/3}} - \frac{\ln\left((bx^3+a)^{1/3} - \frac{(1+\sqrt{3}i)^2(9a+9b)}{36(-a-b)^{2/3}}\right)(1+\sqrt{3}i)}{6(-a-b)^{1/3}}$$

input `int(-x^2/((x^3 - 1)*(a + b*x^3)^(1/3)),x)`output `log((a + b*x^3)^(1/3) - (9*a + 9*b)/(9*(- a - b)^(2/3)))/(3*(- a - b)^(1/3)) + (log((a + b*x^3)^(1/3) - ((3^(1/2)*1i - 1)^2*(9*a + 9*b))/(36*(- a - b)^(2/3)))*(3^(1/2)*1i - 1))/(6*(- a - b)^(1/3)) - (log((a + b*x^3)^(1/3) - ((3^(1/2)*1i + 1)^2*(9*a + 9*b))/(36*(- a - b)^(2/3)))*(3^(1/2)*1i + 1))/(6*(- a - b)^(1/3))`**Reduce [F]**

$$\int \frac{x^2}{(1-x^3)\sqrt[3]{a+bx^3}} dx = -\left(\int \frac{x^2}{(bx^3+a)^{\frac{1}{3}}x^3 - (bx^3+a)^{\frac{1}{3}}} dx\right)$$

input `int(x^2/(-x^3+1)/(b*x^3+a)^(1/3),x)`output `- int(x**2/((a + b*x**3)**(1/3)*x**3 - (a + b*x**3)**(1/3)),x)`

$$3.97 \quad \int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx$$

Optimal result	733
Mathematica [A] (verified)	733
Rubi [A] (verified)	734
Maple [A] (verified)	735
Fricas [B] (verification not implemented)	736
Sympy [F]	736
Maxima [F]	737
Giac [F]	737
Mupad [F(-1)]	737
Reduce [F]	738

Optimal result

Integrand size = 19, antiderivative size = 88

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = -\frac{\arctan\left(\frac{1-\frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt[3]{2}\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt{3}} - \frac{\log(1+x^3)}{6\sqrt[3]{2}} + \frac{\log\left(-\sqrt[3]{2}x - \sqrt[3]{1-x^3}\right)}{2\sqrt[3]{2}}$$

output

$$-1/12*\ln(x^3+1)*2^{(2/3)}+1/4*\ln(-2^{(1/3)}*x-(-x^3+1)^{(1/3)})*2^{(2/3)}-1/6*\arctan(1/3*(1-2*2^{(1/3)}*x/(-x^3+1)^{(1/3)})*3^{(1/2)})*2^{(2/3)}*3^{(1/2)}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.30

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}x}{x-2^{2/3}\sqrt[3]{1-x^3}}\right) - 2 \log\left(2x + 2^{2/3}\sqrt[3]{1-x^3}\right) + \log\left(-2x^2 + 2^{2/3}x\sqrt[3]{1-x^3} - \sqrt[3]{2}(1-x^3)\right)}{6\sqrt[3]{2}}$$

input `Integrate[1/((1 - x^3)^(1/3)*(1 + x^3)),x]`

output `-1/6*(2*Sqrt[3]*ArcTan[(Sqrt[3]*x)/(x - 2^(2/3)*(1 - x^3)^(1/3))] - 2*Log[2*x + 2^(2/3)*(1 - x^3)^(1/3)] + Log[-2*x^2 + 2^(2/3)*x*(1 - x^3)^(1/3) - 2^(1/3)*(1 - x^3)^(2/3)])/2^(1/3)`

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {901}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1}{\sqrt[3]{1-x^3}(x^3+1)} dx$$

↓ 901

$$-\frac{\arctan\left(\frac{1-\frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt{3}} - \frac{\log(x^3+1)}{6\sqrt[3]{2}} + \frac{\log\left(-\sqrt[3]{1-x^3}-\sqrt[3]{2}x\right)}{2\sqrt[3]{2}}$$

input `Int[1/((1 - x^3)^(1/3)*(1 + x^3)),x]`

output `-(ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3])) - Log[1 + x^3]/(6*2^(1/3)) + Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)]/(2*2^(1/3))`

Defintions of rubi rules used

rule 901

```
Int[1/(((a_) + (b_.)*(x_)^3)^(1/3)*((c_) + (d_.)*(x_)^3)), x_Symbol] := With
h[{q = Rt[(b*c - a*d)/c, 3]}, Simp[ArcTan[(1 + (2*q*x)/(a + b*x^3)^(1/3))/S
qrt[3]]/(Sqrt[3]*c*q), x] + (-Simp[Log[q*x - (a + b*x^3)^(1/3)]/(2*c*q), x]
+ Simp[Log[c + d*x^3]/(6*c*q), x])] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0]
```

Maple [A] (verified)

Time = 3.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.08

method	result	size
pseudoelliptic	$2^{\frac{2}{3}} \left(\sqrt{3} \arctan \left(\frac{\sqrt{3} \left(-2^{\frac{2}{3}} (-x^3+1)^{\frac{1}{3}} + x \right)}{3x} \right) + \ln \left(\frac{2^{\frac{1}{3}} x + (-x^3+1)^{\frac{1}{3}}}{x} \right) - \frac{\ln \left(\frac{2^{\frac{2}{3}} x^2 - 2^{\frac{1}{3}} (-x^3+1)^{\frac{1}{3}} x + (-x^3+1)^{\frac{2}{3}}}{x^2} \right)}{2} \right)$	95
trager	Expression too large to display	780

input

```
int(1/(-x^3+1)^(1/3)/(x^3+1),x,method=_RETURNVERBOSE)
```

output

```
1/6*2^(2/3)*(3^(1/2)*arctan(1/3*3^(1/2)*(-2^(2/3)*(-x^3+1)^(1/3)+x)/x)+ln(
(2^(1/3)*x+(-x^3+1)^(1/3))/x)-1/2*ln((2^(2/3)*x^2-2^(1/3)*(-x^3+1)^(1/3)*x
+(-x^3+1)^(2/3))/x^2)
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 246 vs. $2(67) = 134$.

Time = 1.54 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.80

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = -\frac{1}{3} \cdot 2^{\frac{1}{6}} \sqrt{\frac{1}{6}} \arctan \left(\frac{2^{\frac{1}{6}} \sqrt{\frac{1}{6}} \left(6 \cdot 2^{\frac{2}{3}} (5x^7 + 4x^4 - x)(-x^3 + 1)^{\frac{2}{3}} - 2^{\frac{1}{3}} (71x^9 - 111x^6 + 33x^3 - 1) + 12(19x^8 - 16x^5 + x^2)(-x^3 + 1)^{\frac{1}{3}} \right)}{109x^9 - 105x^6 + 3x^3 + 1} \right) + \frac{1}{18} \cdot 2^{\frac{2}{3}} \log \left(\frac{6 \cdot 2^{\frac{1}{3}} (-x^3 + 1)^{\frac{1}{3}} x^2 + 2^{\frac{2}{3}} (x^3 + 1) + 6(-x^3 + 1)^{\frac{2}{3}} x}{x^3 + 1} \right) - \frac{1}{36} \cdot 2^{\frac{2}{3}} \log \left(\frac{3 \cdot 2^{\frac{2}{3}} (5x^4 - x)(-x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}} (19x^6 - 16x^3 + 1) - 12(2x^5 - x^2)(-x^3 + 1)^{\frac{1}{3}}}{x^6 + 2x^3 + 1} \right)$$

input `integrate(1/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="fricas")`

output `-1/3*2^(1/6)*sqrt(1/6)*arctan(2^(1/6)*sqrt(1/6)*(6*2^(2/3)*(5*x^7 + 4*x^4 - x)*(-x^3 + 1)^(2/3) - 2^(1/3)*(71*x^9 - 111*x^6 + 33*x^3 - 1) + 12*(19*x^8 - 16*x^5 + x^2)*(-x^3 + 1)^(1/3))/(109*x^9 - 105*x^6 + 3*x^3 + 1)) + 1/18*2^(2/3)*log((6*2^(1/3)*(-x^3 + 1)^(1/3)*x^2 + 2^(2/3)*(x^3 + 1) + 6*(-x^3 + 1)^(2/3)*x)/(x^3 + 1)) - 1/36*2^(2/3)*log((3*2^(2/3)*(5*x^4 - x)*(-x^3 + 1)^(2/3) + 2^(1/3)*(19*x^6 - 16*x^3 + 1) - 12*(2*x^5 - x^2)*(-x^3 + 1)^(1/3))/(x^6 + 2*x^3 + 1))`

Sympy [F]

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{1}{\sqrt[3]{-(x-1)(x^2+x+1)(x+1)(x^2-x+1)}} dx$$

input `integrate(1/(-x**3+1)**(1/3)/(x**3+1),x)`

output `Integral(1/((-x - 1)*(x**2 + x + 1))**(1/3)*(x + 1)*(x**2 - x + 1)), x)`

Maxima [F]

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{1}{(x^3+1)(-x^3+1)^{\frac{1}{3}}} dx$$

input `integrate(1/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="maxima")`

output `integrate(1/((x^3 + 1)*(-x^3 + 1)^(1/3)), x)`

Giac [F]

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{1}{(x^3+1)(-x^3+1)^{\frac{1}{3}}} dx$$

input `integrate(1/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="giac")`

output `integrate(1/((x^3 + 1)*(-x^3 + 1)^(1/3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{1}{(1-x^3)^{1/3}(x^3+1)} dx$$

input `int(1/((1 - x^3)^(1/3)*(x^3 + 1)),x)`

output `int(1/((1 - x^3)^(1/3)*(x^3 + 1)), x)`

Reduce [F]

$$\int \frac{1}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{1}{(-x^3+1)^{\frac{1}{3}}x^3 + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int(1/(-x^3+1)^(1/3)/(x^3+1),x)`

output `int(1/((-x**3+1)**(1/3)*x**3+(-x**3+1)**(1/3)),x)`

3.98
$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx$$

Optimal result	739
Mathematica [A] (verified)	740
Rubi [A] (verified)	740
Maple [F]	746
Fricas [A] (verification not implemented)	746
Sympy [F]	747
Maxima [F]	747
Giac [F]	748
Mupad [F(-1)]	748
Reduce [F]	748

Optimal result

Integrand size = 20, antiderivative size = 233

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{\arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt{3}} + \frac{\arctan\left(\frac{1+\frac{3}{3}\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}\sqrt{3}}\right)}{2\sqrt[3]{2}\sqrt{3}}$$

$$+ \frac{\log((1-x)(1+x)^2)}{12\sqrt[3]{2}} + \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{6\sqrt[3]{2}}$$

$$- \frac{\log\left(1 + \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{3\sqrt[3]{2}} - \frac{\log\left(-1+x+2^{2/3}\sqrt[3]{1-x^3}\right)}{4\sqrt[3]{2}}$$

output

```
1/24*ln((1-x)*(1+x)^2)*2^(2/3)+1/12*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)-1/6*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)-1/8*ln(-1+x+2^(2/3)*(-x^3+1)^(1/3))*2^(2/3)+1/6*arctan(1/3*(1-2*2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)+1/12*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)*2^(2/3)
```

Mathematica [A] (verified)

Time = 0.72 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.21

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx$$

$$= \frac{-2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1-x^3}}{\sqrt[3]{2}-\sqrt[3]{2x+\sqrt[3]{1-x^3}}}\right) - 4\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1-x^3}}{-2\sqrt[3]{2}+2\sqrt[3]{2x+\sqrt[3]{1-x^3}}}\right) - 4 \log\left(-\sqrt[3]{2} + \sqrt[3]{2x} - \sqrt[3]{1-x^3}\right)}{12\sqrt{3}}$$

input `Integrate[x/((1 - x^3)^(1/3)*(1 + x^3)),x]`

output

```
(-2*Sqrt[3]*ArcTan[(Sqrt[3]*(1 - x^3)^(1/3))/(2^(1/3) - 2^(1/3)*x + (1 - x^3)^(1/3))] - 4*Sqrt[3]*ArcTan[(Sqrt[3]*(1 - x^3)^(1/3))/(-2*2^(1/3) + 2*2^(1/3)*x + (1 - x^3)^(1/3))] - 4*Log[-2^(1/3) + 2^(1/3)*x - (1 - x^3)^(1/3)]) - 2*Log[-2^(1/3) + 2^(1/3)*x + 2*(1 - x^3)^(1/3)] + 2*Log[2^(2/3) - 2*2^(2/3)*x + 2^(2/3)*x^2 + (-1 + x)*(2 - 2*x^3)^(1/3) + (1 - x^3)^(2/3)] + Log[2^(2/3) - 2*2^(2/3)*x + 2^(2/3)*x^2 - 2*(-1 + x)*(2 - 2*x^3)^(1/3) + 4*(1 - x^3)^(2/3)]/(12*2^(1/3))
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.07, number of steps used = 12, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules used = {991, 750, 16, 27, 1142, 25, 27, 1082, 217, 1103, 2574}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x}{\sqrt[3]{1-x^3}(x^3+1)} dx$$

$$\downarrow 991$$

$$-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \int \frac{1}{\frac{2(1-x)^3}{1-x^3} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}$$

$$\downarrow 750$$

$$\begin{aligned}
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \int \frac{\sqrt[3]{2} \left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \\
 & \qquad \qquad \qquad \frac{1}{3} \int \frac{1}{\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} \\
 & \qquad \qquad \qquad \downarrow 16 \\
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \int \frac{\sqrt[3]{2} \left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \\
 & \qquad \qquad \qquad \downarrow 27 \\
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \sqrt[3]{2} \int \frac{2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \\
 & \qquad \qquad \qquad \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \\
 & \qquad \qquad \qquad \downarrow 1142 \\
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \\
 & \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} - \frac{\int -\frac{\sqrt[3]{2} \left(1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2 \cdot 2^{2/3}} \right) - \\
 & \qquad \qquad \qquad \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \\
 & \qquad \qquad \qquad \downarrow 25
 \end{aligned}$$

$$\begin{aligned}
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \\
 & \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} + \frac{\int \frac{\sqrt[3]{2} \left(1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}\right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2 \cdot 2^{2/3}} \right) - \\
 & \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \\
 & \quad \downarrow \text{27} \\
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \\
 & \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} + \frac{\int \frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} \right) - \\
 & \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \\
 & \quad \downarrow \text{1082} \\
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \\
 & \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{-\frac{(1-x)^2}{(1-x^3)^{2/3}} - 3} d \left(1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}\right)}{2^{2/3}} + \frac{\int \frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} \right) - \\
 & \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}}
 \end{aligned}$$

$$\begin{aligned}
 & \downarrow 217 \\
 & \frac{1}{3} \sqrt[3]{2} \left(\frac{\int \frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} - \frac{\sqrt{3} \arctan \left(\frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{2^{2/3}} \right) - \\
 & \frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \\
 & \downarrow 1103 \\
 & -\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \\
 & \frac{1}{3} \sqrt[3]{2} \left(\frac{\sqrt{3} \arctan \left(\frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{2^{2/3}} - \frac{\log \left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{2 \cdot 2^{2/3}} \right) - \\
 & \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \\
 & \downarrow 2574
 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{3} \sqrt[3]{2} \left(\frac{\sqrt{3} \arctan \left(\frac{1 - 2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} \right)}{2^{2/3}} - \frac{\log \left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1 \right)}{2 \cdot 2^{2/3}} \right) + \\
& \frac{1}{3} \left(\frac{\sqrt{3} \arctan \left(\frac{\sqrt[3]{2(1-x)} + 1}{\sqrt[3]{1-x^3}} \right)}{2\sqrt[3]{2}} - \frac{3 \log \left(2^{2/3} \sqrt[3]{1-x^3} + x - 1 \right)}{4\sqrt[3]{2}} + \frac{\log \left((1-x)(x+1)^2 \right)}{4\sqrt[3]{2}} \right) - \\
& \frac{\log \left(\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}}
\end{aligned}$$

input `Int[x/((1 - x^3)^(1/3)*(1 + x^3)),x]`

output `-1/3*(2^(1/3)*(-(Sqrt[3]*ArcTan[(1 - (2*2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/Sqrt[3]))/2^(2/3)) - Log[1 + (2^(2/3)*(1 - x)^2/(1 - x^3)^(2/3) - (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(2*2^(2/3))) - Log[1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(3*2^(1/3)) + ((Sqrt[3]*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/Sqrt[3]))/(2*2^(1/3)) + Log[(1 - x)*(1 + x)^2/(4*2^(1/3)) - (3*Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3])/(4*2^(1/3)))/3`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

- rule 27 $\text{Int}[(a_*)(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_*)(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 750 $\text{Int}[(a_*) + (b_*)(x_)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]^2) \text{ Int}[(2*\text{Rt}[a, 3] - \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 991 $\text{Int}[(x_)/((a_*) + (b_*)(x_)^3)^{1/3}*((c_*) + (d_*)(x_)^3)], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 3]\}, \text{Simp}[-q^2/(3*d) \text{ Int}[1/((1 - q*x)*(a + b*x^3)^{1/3}), x], x] + \text{Simp}[q/d \text{ Subst}[\text{Int}[1/(1 + 2*a*x^3), x], x, (1 + q*x)/(a + b*x^3)^{1/3}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[b*c + a*d, 0]$
- rule 1082 $\text{Int}[(a_*) + (b_*)(x_) + (c_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 1103 $\text{Int}[(d_*) + (e_*)(x_)/((a_*) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{EqQ}[2*c*d - b*e, 0]$
- rule 1142 $\text{Int}[(d_*) + (e_*)(x_)/((a_*) + (b_*)(x_) + (c_*)(x_)^2), x_Symbol] \rightarrow \text{Simp}[(2*c*d - b*e)/(2*c) \text{ Int}[1/(a + b*x + c*x^2), x], x] + \text{Simp}[e/(2*c) \text{ Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x]$

rule 2574

```
Int[1/(((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^3)^(1/3)), x_Symbol] := Simp[
  Sqrt[3]*(ArcTan[(1 - 2^(1/3)*Rt[b, 3]*((c - d*x)/(d*(a + b*x^3)^(1/3))))/Sqrt[3]]/(2^(4/3)*Rt[b, 3]*c)), x] + (Simp[Log[(c + d*x)^2*(c - d*x)]/(2^(7/3)*Rt[b, 3]*c), x] - Simp[(3*Log[Rt[b, 3]*(c - d*x) + 2^(2/3)*d*(a + b*x^3)^(1/3)])/(2^(7/3)*Rt[b, 3]*c), x]) /; FreeQ[{a, b, c, d}, x] && EqQ[b*c^3 + a*d^3, 0]
```

Maple [F]

$$\int \frac{x}{(-x^3 + 1)^{\frac{1}{3}} (x^3 + 1)} dx$$

input

```
int(x/(-x^3+1)^(1/3)/(x^3+1),x)
```

output

```
int(x/(-x^3+1)^(1/3)/(x^3+1),x)
```

Fricas [A] (verification not implemented)

Time = 1.57 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.44

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{1}{6} \cdot 2^{\frac{1}{6}} \sqrt{\frac{1}{6}} \arctan \left(\frac{2^{\frac{1}{6}} \sqrt{\frac{1}{6}} \left(24 \cdot 2^{\frac{2}{3}} (x^{14} - 2x^{11} - 6x^8 - 2x^5 + x^2) (-x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}} (x^{18} + 42x^{15} - 417x^{12} - 102x^9 + 447x^6 - 102x^3 + 1) \right)}{x^{18} - 102x^{15} + 447x^{12} - 102x^9 + 447x^6 - 102x^3 + 1}} \right) - \frac{1}{36} \cdot 2^{\frac{2}{3}} \log \left(-\frac{12(-x^3 + 1)^{\frac{2}{3}} x^2 + 2^{\frac{2}{3}} (x^6 + 2x^3 + 1) - 6 \cdot 2^{\frac{1}{3}} (x^4 - x) (-x^3 + 1)^{\frac{1}{3}}}{x^6 + 2x^3 + 1} \right) + \frac{1}{72} \cdot 2^{\frac{2}{3}} \log \left(\frac{12 \cdot 2^{\frac{2}{3}} (x^8 - 4x^5 + x^2) (-x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}} (x^{12} - 32x^9 + 78x^6 - 32x^3 + 1) + 6(x^{10} - 11x^7 + 11x^4 - 11x + 1)}{x^{12} + 4x^9 + 6x^6 + 4x^3 + 1} \right)$$

input

```
integrate(x/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="fricas")
```

output

```
1/6*2^(1/6)*sqrt(1/6)*arctan(2^(1/6)*sqrt(1/6)*(24*2^(2/3)*(x^14 - 2*x^11
- 6*x^8 - 2*x^5 + x^2)*(-x^3 + 1)^(2/3) + 2^(1/3)*(x^18 + 42*x^15 - 417*x^
12 + 812*x^9 - 417*x^6 + 42*x^3 + 1) - 12*(x^16 - 33*x^13 + 110*x^10 - 110
*x^7 + 33*x^4 - x)*(-x^3 + 1)^(1/3))/(x^18 - 102*x^15 + 447*x^12 - 628*x^9
+ 447*x^6 - 102*x^3 + 1)) - 1/36*2^(2/3)*log(-(12*(-x^3 + 1)^(2/3)*x^2 +
2^(2/3)*(x^6 + 2*x^3 + 1) - 6*2^(1/3)*(x^4 - x)*(-x^3 + 1)^(1/3))/(x^6 + 2
*x^3 + 1)) + 1/72*2^(2/3)*log((12*2^(2/3)*(x^8 - 4*x^5 + x^2)*(-x^3 + 1)^(
2/3) + 2^(1/3)*(x^12 - 32*x^9 + 78*x^6 - 32*x^3 + 1) + 6*(x^10 - 11*x^7 +
11*x^4 - x)*(-x^3 + 1)^(1/3))/(x^12 + 4*x^9 + 6*x^6 + 4*x^3 + 1))
```

Sympy [F]

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x}{\sqrt[3]{-(x-1)(x^2+x+1)(x+1)(x^2-x+1)}} dx$$

input

```
integrate(x/(-x**3+1)**(1/3)/(x**3+1),x)
```

output

```
Integral(x/((-x - 1)*(x**2 + x + 1))**(1/3)*(x + 1)*(x**2 - x + 1)), x)
```

Maxima [F]

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x}{(x^3+1)(-x^3+1)^{\frac{1}{3}}} dx$$

input

```
integrate(x/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="maxima")
```

output

```
integrate(x/((x^3 + 1)*(-x^3 + 1)^(1/3)), x)
```

Giac [F]

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x}{(x^3+1)(-x^3+1)^{\frac{1}{3}}} dx$$

input `integrate(x/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="giac")`

output `integrate(x/((x^3 + 1)*(-x^3 + 1)^(1/3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x}{(1-x^3)^{1/3}(x^3+1)} dx$$

input `int(x/((1 - x^3)^(1/3)*(x^3 + 1)),x)`

output `int(x/((1 - x^3)^(1/3)*(x^3 + 1)), x)`

Reduce [F]

$$\int \frac{x}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x}{(-x^3+1)^{\frac{1}{3}}x^3 + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int(x/(-x^3+1)^(1/3)/(x^3+1),x)`

output `int(x/((- x**3 + 1)**(1/3)*x**3 + (- x**3 + 1)**(1/3)),x)`

3.99 $\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx$

Optimal result	749
Mathematica [A] (verified)	749
Rubi [A] (verified)	750
Maple [A] (verified)	752
Fricas [A] (verification not implemented)	752
Sympy [F]	753
Maxima [A] (verification not implemented)	753
Giac [A] (verification not implemented)	754
Mupad [B] (verification not implemented)	754
Reduce [F]	755

Optimal result

Integrand size = 22, antiderivative size = 82

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{\arctan\left(\frac{1+2^{2/3}\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt{3}} - \frac{\log(1+x^3)}{6\sqrt[3]{2}} + \frac{\log\left(\sqrt[3]{2}-\sqrt[3]{1-x^3}\right)}{2\sqrt[3]{2}}$$

output -1/12*ln(x^3+1)*2^(2/3)+1/4*ln(2^(1/3)-(-x^3+1)^(1/3))*2^(2/3)+1/6*arctan(1/3*(1+2^(2/3)*(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.27

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{2\sqrt{3} \arctan\left(\frac{1+2^{2/3}\sqrt[3]{1-x^3}}{\sqrt{3}}\right) + 2 \log\left(-2 + 2^{2/3}\sqrt[3]{1-x^3}\right) - \log\left(2 + 2^{2/3}\sqrt[3]{1-x^3} + \sqrt[3]{2}(1-x^3)^{2/3}\right)}{6\sqrt[3]{2}}$$

input Integrate[x^2/((1 - x^3)^(1/3)*(1 + x^3)),x]

output

$$\frac{(2\sqrt{3}\operatorname{ArcTan}[(1 + 2^{2/3})(1 - x^3)^{1/3}]/\sqrt{3}] + 2\operatorname{Log}[-2 + 2^{2/3}/3*(1 - x^3)^{1/3}] - \operatorname{Log}[2 + 2^{2/3}/3*(1 - x^3)^{1/3}] + 2^{1/3}*(1 - x^3)^{2/3})/(6*2^{1/3})$$
Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.05, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.227$, Rules used = {946, 67, 16, 1082, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(x^3+1)} dx$$

$$\downarrow 946$$

$$\frac{1}{3} \int \frac{1}{\sqrt[3]{1-x^3}(x^3+1)} dx^3$$

$$\downarrow 67$$

$$\frac{1}{3} \left(-\frac{3 \int \frac{1}{\sqrt[3]{2}-\sqrt[3]{1-x^3}} d\sqrt[3]{1-x^3}}{2\sqrt[3]{2}} + \frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{2}\sqrt[3]{1-x^3} + 2^{2/3}} d\sqrt[3]{1-x^3} - \frac{\log(x^3+1)}{2\sqrt[3]{2}} \right)$$

$$\downarrow 16$$

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{2}\sqrt[3]{1-x^3} + 2^{2/3}} d\sqrt[3]{1-x^3} - \frac{\log(x^3+1)}{2\sqrt[3]{2}} + \frac{3 \log(\sqrt[3]{2} - \sqrt[3]{1-x^3})}{2\sqrt[3]{2}} \right)$$

$$\downarrow 1082$$

$$\frac{1}{3} \left(-\frac{3 \int \frac{1}{-x^6-3} d(2^{2/3}\sqrt[3]{1-x^3}+1)}{\sqrt[3]{2}} - \frac{\log(x^3+1)}{2\sqrt[3]{2}} + \frac{3 \log(\sqrt[3]{2} - \sqrt[3]{1-x^3})}{2\sqrt[3]{2}} \right)$$

$$\downarrow 217$$

$$\frac{1}{3} \left(\frac{\sqrt{3} \arctan\left(\frac{2^{2/3} \sqrt[3]{1-x^3+1}}{\sqrt{3}}\right)}{\sqrt[3]{2}} - \frac{\log(x^3+1)}{2\sqrt[3]{2}} + \frac{3 \log\left(\sqrt[3]{2} - \sqrt[3]{1-x^3}\right)}{2\sqrt[3]{2}} \right)$$

input `Int[x^2/((1 - x^3)^(1/3)*(1 + x^3)),x]`

output `((Sqrt[3]*ArcTan[(1 + 2^(2/3)*(1 - x^3)^(1/3))/Sqrt[3]])/2^(1/3) - Log[1 + x^3]/(2*2^(1/3)) + (3*Log[2^(1/3) - (1 - x^3)^(1/3)])/(2*2^(1/3)))/3`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x]) /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 946 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Simp[1/n Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]`

rule 1082

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Simp[-2/b Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x]
```

Maple [A] (verified)

Time = 3.60 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.98

method	result
pseudoelliptic	$\frac{2^{\frac{2}{3}} \left(2 \arctan \left(\frac{(1+2^{\frac{2}{3}}(-x^3+1)^{\frac{1}{3}})\sqrt{3}}{3} \right) \sqrt{3} + 2 \ln \left((-x^3+1)^{\frac{1}{3}} - 2^{\frac{1}{3}} \right) - \ln \left((-x^3+1)^{\frac{2}{3}} + 2^{\frac{1}{3}}(-x^3+1)^{\frac{1}{3}} + 2^{\frac{2}{3}} \right) \right)}{12}$
trager	$\frac{\text{RootOf}(-Z^3-4) \ln \left(\frac{6 \text{RootOf}(\text{RootOf}(-Z^3-4))^2 + 6_Z \text{RootOf}(-Z^3-4) + 36_Z^2}{\text{RootOf}(-Z^3-4)} \right) x^3 + 45 \text{RootOf}(-Z^3-4)}{\dots}$

input

```
int(x^2/(-x^3+1)^(1/3)/(x^3+1),x,method=_RETURNVERBOSE)
```

output

```
1/12*2^(2/3)*(2*arctan(1/3*(1+2^(2/3)*(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)+2*ln((-x^3+1)^(1/3)-2^(1/3))-ln((-x^3+1)^(2/3)+2^(1/3)*(-x^3+1)^(1/3)+2^(2/3))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.02

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = 2^{\frac{1}{6}} \sqrt{\frac{1}{6}} \arctan \left(2^{\frac{1}{6}} \sqrt{\frac{1}{6}} \left(2^{\frac{1}{3}} + 2(-x^3+1)^{\frac{1}{3}} \right) \right) - \frac{1}{12} \cdot 2^{\frac{2}{3}} \log \left(2^{\frac{2}{3}} + 2^{\frac{1}{3}}(-x^3+1)^{\frac{1}{3}} + (-x^3+1)^{\frac{2}{3}} \right) + \frac{1}{6} \cdot 2^{\frac{2}{3}} \log \left(-2^{\frac{1}{3}} + (-x^3+1)^{\frac{1}{3}} \right)$$

input `integrate(x^2/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="fricas")`

output $2^{1/6} \sqrt{1/6} \arctan(2^{1/6} \sqrt{1/6} (2^{1/3} + 2(-x^3 + 1)^{1/3})) - 1/12 \cdot 2^{2/3} \log(2^{2/3} + 2^{1/3}(-x^3 + 1)^{1/3} + (-x^3 + 1)^{2/3}) + 1/6 \cdot 2^{2/3} \log(-2^{1/3} + (-x^3 + 1)^{1/3})$

Sympy [F]

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x^2}{\sqrt[3]{-(x-1)(x^2+x+1)(x+1)(x^2-x+1)}} dx$$

input `integrate(x**2/(-x**3+1)**(1/3)/(x**3+1),x)`

output `Integral(x**2/((-x - 1)*(x**2 + x + 1))**(1/3)*(x + 1)*(x**2 - x + 1)), x)`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.05

$$\begin{aligned} \int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx &= \frac{1}{6} \sqrt[3]{32} \arctan \left(\frac{1}{6} \sqrt[3]{32} \left(2^{1/3} + 2(-x^3 + 1)^{1/3} \right) \right) \\ &\quad - \frac{1}{12} \cdot 2^{2/3} \log \left(2^{2/3} + 2^{1/3}(-x^3 + 1)^{1/3} + (-x^3 + 1)^{2/3} \right) \\ &\quad + \frac{1}{6} \cdot 2^{2/3} \log \left(-2^{1/3} + (-x^3 + 1)^{1/3} \right) \end{aligned}$$

input `integrate(x^2/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="maxima")`

output $1/6 \sqrt{3} \cdot 2^{2/3} \arctan(1/6 \sqrt{3} \cdot 2^{2/3} (2^{1/3} + 2(-x^3 + 1)^{1/3})) - 1/12 \cdot 2^{2/3} \log(2^{2/3} + 2^{1/3}(-x^3 + 1)^{1/3} + (-x^3 + 1)^{2/3}) + 1/6 \cdot 2^{2/3} \log(-2^{1/3} + (-x^3 + 1)^{1/3})$

Giac [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.06

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{1}{6} \sqrt{3} 2^{\frac{2}{3}} \arctan \left(\frac{1}{6} \sqrt{3} 2^{\frac{2}{3}} \left(2^{\frac{1}{3}} + 2(-x^3+1)^{\frac{1}{3}} \right) \right) - \frac{1}{12} \cdot 2^{\frac{2}{3}} \log \left(2^{\frac{2}{3}} + 2^{\frac{1}{3}}(-x^3+1)^{\frac{1}{3}} + (-x^3+1)^{\frac{2}{3}} \right) + \frac{1}{6} \cdot 2^{\frac{2}{3}} \log \left(\left| -2^{\frac{1}{3}} + (-x^3+1)^{\frac{1}{3}} \right| \right)$$

input `integrate(x^2/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="giac")`output `1/6*sqrt(3)*2^(2/3)*arctan(1/6*sqrt(3)*2^(2/3)*(2^(1/3) + 2*(-x^3 + 1)^(1/3))) - 1/12*2^(2/3)*log(2^(2/3) + 2^(1/3)*(-x^3 + 1)^(1/3) + (-x^3 + 1)^(2/3)) + 1/6*2^(2/3)*log(abs(-2^(1/3) + (-x^3 + 1)^(1/3)))`**Mupad [B] (verification not implemented)**

Time = 0.35 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.22

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{2^{2/3} \ln \left((1-x^3)^{1/3} - 2^{1/3} \right)}{6} + \frac{2^{2/3} \ln \left((1-x^3)^{1/3} - \frac{2^{1/3}(-1+\sqrt{3}i)^2}{4} \right) (-1 + \sqrt{3}i)}{12} - \frac{2^{2/3} \ln \left((1-x^3)^{1/3} - \frac{2^{1/3}(1+\sqrt{3}i)^2}{4} \right) (1 + \sqrt{3}i)}{12}$$

input `int(x^2/((1 - x^3)^(1/3)*(x^3 + 1)),x)`output `(2^(2/3)*log((1 - x^3)^(1/3) - 2^(1/3)))/6 + (2^(2/3)*log((1 - x^3)^(1/3) - (2^(1/3)*(3^(1/2)*1i - 1)^2)/4)*(3^(1/2)*1i - 1)/12 - (2^(2/3)*log((1 - x^3)^(1/3) - (2^(1/3)*(3^(1/2)*1i + 1)^2)/4)*(3^(1/2)*1i + 1)/12`

Reduce [F]

$$\int \frac{x^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x^2}{(-x^3+1)^{\frac{1}{3}}x^3 + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int(x^2/(-x^3+1)^(1/3)/(x^3+1),x)`

output `int(x**2/((- x**3 + 1)**(1/3)*x**3 + (- x**3 + 1)**(1/3)),x)`

3.100 $\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx$

Optimal result	756
Mathematica [A] (verified)	757
Rubi [B] (verified)	757
Maple [C] (warning: unable to verify)	759
Fricas [B] (verification not implemented)	760
Sympy [F]	761
Maxima [F]	761
Giac [F]	761
Mupad [F(-1)]	762
Reduce [F]	762

Optimal result

Integrand size = 25, antiderivative size = 135

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx = \frac{\sqrt{3} \arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2(1-x)}}{\frac{\sqrt[3]{1-x^3}}{\sqrt{3}}}\right)}{\sqrt[3]{2}} + \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{2\sqrt[3]{2}} - \frac{\log\left(1 + \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{\sqrt[3]{2}}$$

output

```
1/4*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)-1/2*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)+1/2*arctan(1/3*(1-2*2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)
```

Mathematica [A] (verified)

Time = 1.10 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.07

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx$$

$$= \frac{-2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1-x^3}}{-2\sqrt[3]{2+2\sqrt[3]{2x+\sqrt[3]{1-x^3}}}}\right) - 2 \log\left(-\sqrt[3]{2} + \sqrt[3]{2}x - \sqrt[3]{1-x^3}\right) + \log\left(2^{2/3} - 2 \cdot 2^{2/3}x + 2^{2/3}\right)}{2\sqrt[3]{2}}$$

input

```
Integrate[(1 + x)/((1 - x + x^2)*(1 - x^3)^(1/3)),x]
```

output

```
(-2*Sqrt[3]*ArcTan[(Sqrt[3]*(1 - x^3)^(1/3))/(-2*2^(1/3) + 2*2^(1/3)*x + (1 - x^3)^(1/3))] - 2*Log[-2^(1/3) + 2^(1/3)*x - (1 - x^3)^(1/3)] + Log[2^(2/3) - 2*2^(2/3)*x + 2^(2/3)*x^2 + (-1 + x)*(2 - 2*x^3)^(1/3) + (1 - x^3)^(2/3)])/(2*2^(1/3))
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 383 vs. $2(135) = 270$.

Time = 0.57 (sec) , antiderivative size = 383, normalized size of antiderivative = 2.84, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x+1}{(x^2-x+1)\sqrt[3]{1-x^3}} dx$$

$$\downarrow \text{2583}$$

$$\int \left(\frac{2x}{\sqrt[3]{1-x^3}(x^3+1)} + \frac{1}{\sqrt[3]{1-x^3}(x^3+1)} + \frac{x^2}{\sqrt[3]{1-x^3}(x^3+1)} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{2^{2/3} \arctan\left(\frac{1 - \frac{2}{3}\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{\sqrt[3]{2(1-x)} + 1}{\sqrt[3]{1-x^3}}\right)}{\sqrt[3]{2}\sqrt{3}} - \frac{\arctan\left(\frac{1 - \frac{2}{3}\sqrt[3]{2x}}{\sqrt[3]{1-x^3}}\right)}{\sqrt[3]{2}\sqrt{3}} + \\
& \frac{\arctan\left(\frac{2^{2/3}\sqrt[3]{1-x^3+1}}{\sqrt[3]{2}\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt{3}} - \frac{\log(x^3+1)}{3\sqrt[3]{2}} + \frac{\log\left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1\right)}{3\sqrt[3]{2}} - \\
& \frac{1}{3}2^{2/3}\log\left(\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1\right) + \frac{\log\left(\sqrt[3]{2} - \sqrt[3]{1-x^3}\right)}{2\sqrt[3]{2}} + \frac{\log\left(-\sqrt[3]{1-x^3} - \sqrt[3]{2x}\right)}{2\sqrt[3]{2}} - \\
& \frac{\log\left(2^{2/3}\sqrt[3]{1-x^3} + x - 1\right)}{2\sqrt[3]{2}} + \frac{\log\left((1-x)(x+1)^2\right)}{6\sqrt[3]{2}}
\end{aligned}$$

input `Int[(1 + x)/((1 - x + x^2)*(1 - x^3)^(1/3)), x]`

output `(2^(2/3)*ArcTan[(1 - (2*2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) - ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + ArcTan[(1 + 2^(2/3)*(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + Log[(1 - x)*(1 + x)^2]/(6*2^(1/3)) - Log[1 + x^3]/(3*2^(1/3)) + Log[1 + (2^(2/3)*(1 - x)^2)/(1 - x^3)^(2/3) - (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(3*2^(1/3)) - (2^(2/3)*Log[1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)])/3 + Log[2^(1/3) - (1 - x^3)^(1/3)]/(2*2^(1/3)) + Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)]/(2*2^(1/3)) - Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3)]/(2*2^(1/3))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2583 `Int[(Px_.)*((c_) + (d_.)*(x_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^3)^(p_.), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 7.27 (sec) , antiderivative size = 720, normalized size of antiderivative = 5.33

method	result	size
trager	Expression too large to display	720

input `int((1+x)/(x^2-x+1)/(-x^3+1)^(1/3),x,method=_RETURNVERBOSE)`

output

```
-1/2*ln(-(RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+
4)^3*x+(-x^3+1)^(2/3)*RootOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf
(_Z^3+4)+4*_Z^2)-(-x^3+1)^(1/3)*RootOf(_Z^3+4)^2*x-2*(-x^3+1)^(1/3)*RootOf
(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)*x+(-x^3+1)^(1
/3)*RootOf(_Z^3+4)^2+2*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(
_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)+x^2*RootOf(_Z^3+4)-3*x*RootOf(_Z^3+4)-2*(-x
^3+1)^(2/3)+RootOf(_Z^3+4)))/(x^2-x+1))*RootOf(_Z^3+4)-ln(-(RootOf(RootOf(_
Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)^3*x+(-x^3+1)^(2/3)*Roo
tOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)-(-x^3+1)
^(1/3)*RootOf(_Z^3+4)^2*x-2*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*Ro
otOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)*x+(-x^3+1)^(1/3)*RootOf(_Z^3+4)^2+2*(-
x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z
^3+4)+x^2*RootOf(_Z^3+4)-3*x*RootOf(_Z^3+4)-2*(-x^3+1)^(2/3)+RootOf(_Z^3+4
)))/(x^2-x+1))*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)+RootOf(Ro
otOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*ln((RootOf(RootOf(_Z^3+4)^2+2*_
_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)^2*x+RootOf(RootOf(_Z^3+4)^2+2*_Z*
RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)*(-x^3+1)^(2/3)-2*(-x^3+1)^(1/3)*Root
Of(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*x+2*(-x^3+1)^(1/3)*RootOf(
RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)-x^2+x-1)/(x^2-x+1))
```


Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(101) = 202$.

Time = 5.65 (sec) , antiderivative size = 287, normalized size of antiderivative = 2.13

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx = -\frac{1}{3} \cdot 2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \arctan \left(\frac{2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \left(2 \cdot 2^{\frac{2}{3}} (x^4 - 4x^3 + 5x^2 - 4x + 1) (-x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}} (x^6 - 7x^5 + 10x^4 - 7x^3 + 10x^2 - 7x + 1) \right)}{3(3x^6 - 9x^5 + 6x^4 - x^3 + 6x^2 - 9x + 3)}} \right) + \frac{1}{12} \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} (-x^3 + 1)^{\frac{2}{3}} (x^2 - 3x + 1) - 2^{\frac{1}{3}} (x^4 - 3x^2 + 1) - 4(-x^3 + 1)^{\frac{1}{3}} (x^2 - x)}{x^4 - 2x^3 + 3x^2 - 2x + 1} \right) - \frac{1}{6} \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} (x^2 - x + 1) - 2 \cdot 2^{\frac{1}{3}} (-x^3 + 1)^{\frac{1}{3}} (x - 1) + 2(-x^3 + 1)^{\frac{2}{3}}}{x^2 - x + 1} \right)$$

input `integrate((1+x)/(x^2-x+1)/(-x^3+1)^(1/3),x, algorithm="fricas")`

output `-1/3*2^(1/6)*sqrt(3/2)*arctan(1/3*2^(1/6)*sqrt(3/2)*(2*2^(2/3)*(x^4 - 4*x^3 + 5*x^2 - 4*x + 1)*(-x^3 + 1)^(2/3) + 2^(1/3)*(x^6 - 7*x^5 + 10*x^4 - 7*x^3 + 10*x^2 - 7*x + 1) + 4*(x^5 - x^4 - 3*x^3 + 3*x^2 + x - 1)*(-x^3 + 1)^(1/3))/(3*x^6 - 9*x^5 + 6*x^4 - x^3 + 6*x^2 - 9*x + 3)) + 1/12*2^(2/3)*log((2^(2/3)*(-x^3 + 1)^(2/3)*(x^2 - 3*x + 1) - 2^(1/3)*(x^4 - 3*x^2 + 1) - 4*(-x^3 + 1)^(1/3)*(x^2 - x))/(x^4 - 2*x^3 + 3*x^2 - 2*x + 1)) - 1/6*2^(2/3)*log((2^(2/3)*(x^2 - x + 1) - 2*2^(1/3)*(-x^3 + 1)^(1/3)*(x - 1) + 2*(-x^3 + 1)^(2/3))/(x^2 - x + 1))`

Sympy [F]

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx = \int \frac{x+1}{\sqrt[3]{-(x-1)(x^2+x+1)}(x^2-x+1)} dx$$

input `integrate((1+x)/(x**2-x+1)/(-x**3+1)**(1/3),x)`

output `Integral((x + 1)/((-x - 1)*(x**2 + x + 1))**(1/3)*(x**2 - x + 1)), x)`

Maxima [F]

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx = \int \frac{x+1}{(-x^3+1)^{\frac{1}{3}}(x^2-x+1)} dx$$

input `integrate((1+x)/(x^2-x+1)/(-x^3+1)^(1/3),x, algorithm="maxima")`

output `integrate((x + 1)/((-x^3 + 1)^(1/3)*(x^2 - x + 1)), x)`

Giac [F]

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx = \int \frac{x+1}{(-x^3+1)^{\frac{1}{3}}(x^2-x+1)} dx$$

input `integrate((1+x)/(x^2-x+1)/(-x^3+1)^(1/3),x, algorithm="giac")`

output `integrate((x + 1)/((-x^3 + 1)^(1/3)*(x^2 - x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx = \int \frac{x+1}{(1-x^3)^{1/3}(x^2-x+1)} dx$$

input `int((x + 1)/((1 - x^3)^(1/3)*(x^2 - x + 1)), x)`output `int((x + 1)/((1 - x^3)^(1/3)*(x^2 - x + 1)), x)`**Reduce [F]**

$$\int \frac{1+x}{(1-x+x^2)\sqrt[3]{1-x^3}} dx = \int \frac{x}{(-x^3+1)^{\frac{1}{3}}x^2 - (-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx + \int \frac{1}{(-x^3+1)^{\frac{1}{3}}x^2 - (-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int((1+x)/(x^2-x+1)/(-x^3+1)^(1/3), x)`output `int(x/((- x**3 + 1)**(1/3)*x**2 - (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)), x) + int(1/((- x**3 + 1)**(1/3)*x**2 - (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)), x)`

3.101
$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx$$

Optimal result	763
Mathematica [A] (verified)	764
Rubi [B] (verified)	764
Maple [C] (warning: unable to verify)	766
Fricas [B] (verification not implemented)	767
Sympy [F]	768
Maxima [F]	768
Giac [F]	769
Mupad [F(-1)]	769
Reduce [F]	769

Optimal result

Integrand size = 24, antiderivative size = 135

$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \frac{\sqrt{3} \arctan\left(\frac{1 - \frac{2}{3}\sqrt[3]{2(1-x)}}{\frac{\sqrt[3]{1-x^3}}{\sqrt{3}}}\right)}{\sqrt[3]{2}} + \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{2\sqrt[3]{2}} - \frac{\log\left(1 + \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{\sqrt[3]{2}}$$

```
output 1/4*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)-1/2*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)+1/2*arctan(1/3*(1-2*2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)
```

Mathematica [A] (verified)

Time = 1.06 (sec) , antiderivative size = 145, normalized size of antiderivative = 1.07

$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx$$

$$= \frac{-2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1-x^3}}{-2\sqrt[3]{2+2\sqrt[3]{2x+\sqrt[3]{1-x^3}}}}\right) - 2\log\left(-\sqrt[3]{2} + \sqrt[3]{2}x - \sqrt[3]{1-x^3}\right) + \log\left(2^{2/3} - 2 \cdot 2^{2/3}x + 2^{2/3}\right)}{2\sqrt[3]{2}}$$

input

```
Integrate[(1 + x)^2/((1 - x^3)^(1/3)*(1 + x^3)),x]
```

output

```
(-2*Sqrt[3]*ArcTan[(Sqrt[3]*(1 - x^3)^(1/3))/(-2*2^(1/3) + 2*2^(1/3)*x + (1 - x^3)^(1/3))] - 2*Log[-2^(1/3) + 2^(1/3)*x - (1 - x^3)^(1/3)] + Log[2^(2/3) - 2*2^(2/3)*x + 2^(2/3)*x^2 + (-1 + x)*(2 - 2*x^3)^(1/3) + (1 - x^3)^(2/3)])/(2*2^(1/3))
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 383 vs. $2(135) = 270$.

Time = 0.60 (sec) , antiderivative size = 383, normalized size of antiderivative = 2.84, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2019, 2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(x+1)^2}{\sqrt[3]{1-x^3}(x^3+1)} dx$$

$$\downarrow \text{2019}$$

$$\int \frac{x+1}{(x^2-x+1)\sqrt[3]{1-x^3}} dx$$

$$\downarrow \text{2583}$$

$$\int \left(\frac{2x}{\sqrt[3]{1-x^3}(x^3+1)} + \frac{1}{\sqrt[3]{1-x^3}(x^3+1)} + \frac{x^2}{\sqrt[3]{1-x^3}(x^3+1)} \right) dx$$

↓ 2009

$$\frac{2^{2/3} \arctan \left(\frac{1 - \frac{2}{3} \sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} \right)}{\sqrt{3}} + \frac{\arctan \left(\frac{\sqrt[3]{2(1-x)} + 1}{\sqrt[3]{1-x^3}} \right)}{\sqrt[3]{2}\sqrt{3}} - \frac{\arctan \left(\frac{1 - \frac{2}{3} \sqrt[3]{2x}}{\sqrt[3]{1-x^3}} \right)}{\sqrt[3]{2}\sqrt{3}} +$$

$$\frac{\arctan \left(\frac{2^{2/3} \sqrt[3]{1-x^3+1}}{\sqrt{3}} \right)}{\sqrt[3]{2}\sqrt{3}} - \frac{\log(x^3+1)}{3\sqrt[3]{2}} + \frac{\log \left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)} + 1}{\sqrt[3]{1-x^3}} \right)}{3\sqrt[3]{2}} -$$

$$\frac{1}{3} 2^{2/3} \log \left(\frac{\sqrt[3]{2(1-x)} + 1}{\sqrt[3]{1-x^3}} \right) + \frac{\log(\sqrt[3]{2} - \sqrt[3]{1-x^3})}{2\sqrt[3]{2}} + \frac{\log(-\sqrt[3]{1-x^3} - \sqrt[3]{2}x)}{2\sqrt[3]{2}} -$$

$$\frac{\log(2^{2/3} \sqrt[3]{1-x^3} + x - 1)}{2\sqrt[3]{2}} + \frac{\log((1-x)(x+1)^2)}{6\sqrt[3]{2}}$$

input `Int[(1 + x)^2/((1 - x^3)^(1/3)*(1 + x^3)),x]`

output `(2^(2/3)*ArcTan[(1 - (2*2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) - ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + ArcTan[(1 + 2^(2/3)*(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) + Log[(1 - x)*(1 + x)^2/(6*2^(1/3)) - Log[1 + x^3]/(3*2^(1/3)) + Log[1 + (2^(2/3)*(1 - x)^2)/(1 - x^3)^(2/3) - (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(3*2^(1/3)) - (2^(2/3)*Log[1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)])/3 + Log[2^(1/3) - (1 - x^3)^(1/3)]/(2*2^(1/3)) + Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)]/(2*2^(1/3)) - Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3)]/(2*2^(1/3))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2019 `Int[(u_.)*(Px_)^(p_.)*(Qx_)^(q_.), x_Symbol] := Int[u*PolynomialQuotient[Px, Qx, x]^p*Qx^(p + q), x] /; FreeQ[q, x] && PolyQ[Px, x] && PolyQ[Qx, x] && EqQ[PolynomialRemainder[Px, Qx, x], 0] && IntegerQ[p] && LtQ[p*q, 0]`

rule 2583 `Int[(Px_.)*((c_) + (d_.)*(x_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^3)^(p_.), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 6.32 (sec) , antiderivative size = 720, normalized size of antiderivative = 5.33

method	result	size
trager	Expression too large to display	720

input `int((1+x)^2/(-x^3+1)^(1/3)/(x^3+1),x,method=_RETURNVERBOSE)`

output

```

-1/2*ln(-(RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+
4)^3*x+(-x^3+1)^(2/3)*RootOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf
(_Z^3+4)+4*_Z^2)-(-x^3+1)^(1/3)*RootOf(_Z^3+4)^2*x-2*(-x^3+1)^(1/3)*RootOf
(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)*x+(-x^3+1)^(1
/3)*RootOf(_Z^3+4)^2+2*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(
_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)+x^2*RootOf(_Z^3+4)-3*x*RootOf(_Z^3+4)-2*(-x
^3+1)^(2/3)+RootOf(_Z^3+4))/(x^2-x+1)*RootOf(_Z^3+4)-ln(-(RootOf(RootOf(_
Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)^3*x+(-x^3+1)^(2/3)*Ro
otOf(_Z^3+4)^2*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)-(-x^3+1)
^(1/3)*RootOf(_Z^3+4)^2*x-2*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*Ro
otOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)*x+(-x^3+1)^(1/3)*RootOf(_Z^3+4)^2+2*(-
x^3+1)^(1/3)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z
^3+4)+x^2*RootOf(_Z^3+4)-3*x*RootOf(_Z^3+4)-2*(-x^3+1)^(2/3)+RootOf(_Z^3+4
))/ (x^2-x+1)*RootOf(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)+RootOf(R
ootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*ln((RootOf(RootOf(_Z^3+4)^2+2*_
_Z*RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)^2*x+RootOf(RootOf(_Z^3+4)^2+2*_Z*
RootOf(_Z^3+4)+4*_Z^2)*RootOf(_Z^3+4)*(-x^3+1)^(2/3)-2*(-x^3+1)^(1/3)*Root
Of(RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)*x+2*(-x^3+1)^(1/3)*RootOf(
RootOf(_Z^3+4)^2+2*_Z*RootOf(_Z^3+4)+4*_Z^2)-x^2+x-1)/(x^2-x+1))

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 287 vs. $2(101) = 202$.

Time = 5.61 (sec) , antiderivative size = 287, normalized size of antiderivative = 2.13

$$\begin{aligned}
& \int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = -\frac{1}{3} \\
& \cdot 2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \arctan \left(\frac{2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \left(2 \cdot 2^{\frac{2}{3}} (x^4 - 4x^3 + 5x^2 - 4x + 1) (-x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}} (x^6 - 7x^5 + 10x^4 - 7x^3 + 1) \right)}{3(3x^6 - 9x^5 + 6x^4 - x^3 + 6x^2)}} \right) \\
& + \frac{1}{12} \\
& \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} (-x^3 + 1)^{\frac{2}{3}} (x^2 - 3x + 1) - 2^{\frac{1}{3}} (x^4 - 3x^2 + 1) - 4(-x^3 + 1)^{\frac{1}{3}} (x^2 - x)}{x^4 - 2x^3 + 3x^2 - 2x + 1}} \right) \\
& - \frac{1}{6} \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} (x^2 - x + 1) - 2 \cdot 2^{\frac{1}{3}} (-x^3 + 1)^{\frac{1}{3}} (x - 1) + 2(-x^3 + 1)^{\frac{2}{3}}}{x^2 - x + 1}} \right)
\end{aligned}$$

input `integrate((1+x)^2/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="fricas")`

output `-1/3*2^(1/6)*sqrt(3/2)*arctan(1/3*2^(1/6)*sqrt(3/2)*(2*2^(2/3)*(x^4 - 4*x^3 + 5*x^2 - 4*x + 1)*(-x^3 + 1)^(2/3) + 2^(1/3)*(x^6 - 7*x^5 + 10*x^4 - 7*x^3 + 10*x^2 - 7*x + 1) + 4*(x^5 - x^4 - 3*x^3 + 3*x^2 + x - 1)*(-x^3 + 1)^(1/3))/(3*x^6 - 9*x^5 + 6*x^4 - x^3 + 6*x^2 - 9*x + 3)) + 1/12*2^(2/3)*log((2^(2/3)*(-x^3 + 1)^(2/3)*(x^2 - 3*x + 1) - 2^(1/3)*(x^4 - 3*x^2 + 1) - 4*(-x^3 + 1)^(1/3)*(x^2 - x))/(x^4 - 2*x^3 + 3*x^2 - 2*x + 1)) - 1/6*2^(2/3)*log((2^(2/3)*(x^2 - x + 1) - 2*2^(1/3)*(-x^3 + 1)^(1/3)*(x - 1) + 2*(-x^3 + 1)^(2/3))/(x^2 - x + 1))`

Sympy [F]

$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x+1}{\sqrt[3]{-(x-1)(x^2+x+1)(x^2-x+1)}} dx$$

input `integrate((1+x)**2/(-x**3+1)**(1/3)/(x**3+1),x)`

output `Integral((x + 1)/((-x - 1)*(x**2 + x + 1))**(1/3)*(x**2 - x + 1)), x)`

Maxima [F]

$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{(x+1)^2}{(x^3+1)(-x^3+1)^{\frac{1}{3}}} dx$$

input `integrate((1+x)^2/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="maxima")`

output `integrate((x + 1)^2/((x^3 + 1)*(-x^3 + 1)^(1/3)), x)`

Giac [F]

$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{(x+1)^2}{(x^3+1)(-x^3+1)^{\frac{1}{3}}} dx$$

input `integrate((1+x)^2/(-x^3+1)^(1/3)/(x^3+1),x, algorithm="giac")`

output `integrate((x + 1)^2/((x^3 + 1)*(-x^3 + 1)^(1/3)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{(x+1)^2}{(1-x^3)^{1/3}(x^3+1)} dx$$

input `int((x + 1)^2/((1 - x^3)^(1/3)*(x^3 + 1)),x)`

output `int((x + 1)^2/((1 - x^3)^(1/3)*(x^3 + 1)), x)`

Reduce [F]

$$\int \frac{(1+x)^2}{\sqrt[3]{1-x^3}(1+x^3)} dx = \int \frac{x}{(-x^3+1)^{\frac{1}{3}}x^2 - (-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx + \int \frac{1}{(-x^3+1)^{\frac{1}{3}}x^2 - (-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int((1+x)^2/(-x^3+1)^(1/3)/(x^3+1),x)`

output `int(x/((- x**3 + 1)**(1/3)*x**2 - (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x) + int(1/((- x**3 + 1)**(1/3)*x**2 - (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)`

3.102
$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx$$

Optimal result	770
Mathematica [A] (verified)	771
Rubi [B] (verified)	771
Maple [C] (warning: unable to verify)	773
Fricas [B] (verification not implemented)	774
Sympy [F]	774
Maxima [F]	775
Giac [F]	775
Mupad [F(-1)]	775
Reduce [F]	776

Optimal result

Integrand size = 23, antiderivative size = 119

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx = -\frac{\sqrt{3} \arctan\left(\frac{1-\frac{2\sqrt[3]{2}(1+x)}{\sqrt[3]{1+x^3}}}{\sqrt{3}}\right)}{\sqrt[3]{2}} - \frac{\log\left(1 + \frac{2^{2/3}(1+x)^2}{(1+x^3)^{2/3}} - \frac{\sqrt[3]{2}(1+x)}{\sqrt[3]{1+x^3}}\right)}{2\sqrt[3]{2}} + \frac{\log\left(1 + \frac{\sqrt[3]{2}(1+x)}{\sqrt[3]{1+x^3}}\right)}{\sqrt[3]{2}}$$

output

```
-1/4*ln(1+2^(2/3)*(1+x)^2/(x^3+1)^(2/3)-2^(1/3)*(1+x)/(x^3+1)^(1/3))*2^(2/3)+1/2*ln(1+2^(1/3)*(1+x)/(x^3+1)^(1/3))*2^(2/3)-1/2*arctan(1/3*(1-2*2^(1/3)*(1+x)/(x^3+1)^(1/3))*3^(1/2))*3^(1/2)*2^(2/3)
```

Mathematica [A] (verified)

Time = 1.03 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.17

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx$$

$$= \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1+x^3}}{-2\sqrt[3]{2}-2\sqrt[3]{2x+\sqrt[3]{1+x^3}}}\right) + 2\log\left(\sqrt[3]{2} + \sqrt[3]{2x} + \sqrt[3]{1+x^3}\right) - \log\left(2^{2/3} + 2 \cdot 2^{2/3}x + 2^{2/3}x^2 - \dots\right)}{2\sqrt[3]{2}}$$

input

```
Integrate[(1 - x)/((1 + x + x^2)*(1 + x^3)^(1/3)),x]
```

output

```
(2*Sqrt[3]*ArcTan[(Sqrt[3]*(1 + x^3)^(1/3))/(-2*2^(1/3) - 2*2^(1/3)*x + (1 + x^3)^(1/3))] + 2*Log[2^(1/3) + 2^(1/3)*x + (1 + x^3)^(1/3)] - Log[2^(2/3) + 2*2^(2/3)*x + 2^(2/3)*x^2 - 2^(1/3)*(1 + x)*(1 + x^3)^(1/3) + (1 + x^3)^(2/3)])/(2*2^(1/3))
```

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 357 vs. $2(119) = 238$.

Time = 0.57 (sec) , antiderivative size = 357, normalized size of antiderivative = 3.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1-x}{(x^2+x+1)\sqrt[3]{x^3+1}} dx$$

$$\downarrow \text{2583}$$

$$\int \left(-\frac{2x}{(1-x^3)\sqrt[3]{x^3+1}} + \frac{1}{(1-x^3)\sqrt[3]{x^3+1}} + \frac{x^2}{(1-x^3)\sqrt[3]{x^3+1}} \right) dx$$

$$\downarrow \text{2009}$$

$$\begin{aligned}
& \frac{\arctan\left(\frac{\sqrt[3]{2}\sqrt[3]{2x}+1}{\sqrt[3]{x^3+1}}\right)}{\sqrt[3]{2}\sqrt{3}} - \frac{2^{2/3}\arctan\left(\frac{1-\sqrt[3]{2}\sqrt[3]{2(x+1)}}{\sqrt[3]{x^3+1}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{\sqrt[3]{2}\sqrt[3]{2(x+1)}+1}{\sqrt[3]{x^3+1}}\right)}{\sqrt[3]{2}\sqrt{3}} \\
& \frac{\arctan\left(\frac{2^{2/3}\sqrt[3]{x^3+1+1}}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt{3}} + \frac{\log(1-x^3)}{3\sqrt[3]{2}} - \frac{\log\left(\frac{2^{2/3}(x+1)^2}{(x^3+1)^{2/3}} - \frac{\sqrt[3]{2}\sqrt[3]{2(x+1)}}{\sqrt[3]{x^3+1}} + 1\right)}{3\sqrt[3]{2}} + \\
& \frac{1}{3}2^{2/3}\log\left(\frac{\sqrt[3]{2}\sqrt[3]{2(x+1)}}{\sqrt[3]{x^3+1}} + 1\right) - \frac{\log\left(\sqrt[3]{2} - \sqrt[3]{x^3+1}\right)}{2\sqrt[3]{2}} - \frac{\log\left(\sqrt[3]{2}x - \sqrt[3]{x^3+1}\right)}{2\sqrt[3]{2}} + \\
& \frac{\log\left(-2^{2/3}\sqrt[3]{x^3+1} + x + 1\right)}{2\sqrt[3]{2}} - \frac{\log\left((1-x)^2(x+1)\right)}{6\sqrt[3]{2}}
\end{aligned}$$

input `Int[(1 - x)/((1 + x + x^2)*(1 + x^3)^(1/3)), x]`

output `ArcTan[(1 + (2*2^(1/3)*x)/(1 + x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) - (2^(2/3)*ArcTan[(1 - (2*2^(1/3)*(1 + x))/(1 + x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - ArcTan[(1 + (2^(1/3)*(1 + x))/(1 + x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) - ArcTan[(1 + 2^(2/3)*(1 + x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3]) - Log[(1 - x)^2*(1 + x)]/(6*2^(1/3)) + Log[1 - x^3]/(3*2^(1/3)) - Log[1 + (2^(2/3)*(1 + x)^2)/(1 + x^3)^(2/3) - (2^(1/3)*(1 + x))/(1 + x^3)^(1/3)]/(3*2^(1/3)) + (2^(2/3)*Log[1 + (2^(1/3)*(1 + x))/(1 + x^3)^(1/3)]/3 - Log[2^(1/3) - (1 + x^3)^(1/3)]/(2*2^(1/3)) - Log[2^(1/3)*x - (1 + x^3)^(1/3)]/(2*2^(1/3)) + Log[1 + x - 2^(2/3)*(1 + x^3)^(1/3)]/(2*2^(1/3))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2583 `Int[(Px_.)*((c_) + (d_.)*(x_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^3)^(p_.), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 7.41 (sec) , antiderivative size = 687, normalized size of antiderivative = 5.77

method	result	size
trager	Expression too large to display	687

input `int((1-x)/(x^2+x+1)/(x^3+1)^(1/3),x,method=_RETURNVERBOSE)`

output

```
RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*ln(-(RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^2*x+RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)*(x^3+1)^(2/3)-2*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*(x^3+1)^(1/3)*x-2*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*(x^3+1)^(1/3)-x^2-x-1)/(x^2+x+1))-1/2*ln((RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^3*x+(x^3+1)^(2/3)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^2-(x^3+1)^(1/3)*RootOf(_Z^3-4)^2*x-2*(x^3+1)^(1/3)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)*x-(x^3+1)^(1/3)*RootOf(_Z^3-4)^2-2*(x^3+1)^(1/3)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)+RootOf(_Z^3-4)*x^2+3*RootOf(_Z^3-4)*x+2*(x^3+1)^(2/3)+RootOf(_Z^3-4))/(x^2+x+1))*RootOf(_Z^3-4)-ln((RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^3*x+(x^3+1)^(2/3)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)^2-(x^3+1)^(1/3)*RootOf(_Z^3-4)^2*x-2*(x^3+1)^(1/3)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)*x-(x^3+1)^(1/3)*RootOf(_Z^3-4)^2-2*(x^3+1)^(1/3)*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)*RootOf(_Z^3-4)+RootOf(_Z^3-4)*x^2+3*RootOf(_Z^3-4)*x+2*(x^3+1)^(2/3)+RootOf(_Z^3-4))/(x^2+x+1))*RootOf(RootOf(_Z^3-4)^2+2*_Z*RootOf(_Z^3-4)+4*_Z^2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 265 vs. $2(93) = 186$.

Time = 5.24 (sec) , antiderivative size = 265, normalized size of antiderivative = 2.23

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx = \frac{1}{3} \cdot 2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \arctan \left(\frac{2^{\frac{1}{6}} \sqrt{\frac{3}{2}} \left(2 \cdot 2^{\frac{2}{3}} (x^4 + 4x^3 + 5x^2 + 4x + 1)(x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}} (x^6 + 7x^5 + 10x^4 + 7x^3 + 10x^2 + 7x + 1) \right)}{3(3x^6 + 9x^5 + 6x^4 + x^3 + 6x^2 + 9x + 3)}} \right) - \frac{1}{12} \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} (x^3 + 1)^{\frac{2}{3}} (x^2 + 3x + 1) - 2^{\frac{1}{3}} (x^4 - 3x^2 + 1) - 4(x^3 + 1)^{\frac{1}{3}} (x^2 + x)}{x^4 + 2x^3 + 3x^2 + 2x + 1}} \right) + \frac{1}{6} \cdot 2^{\frac{2}{3}} \log \left(\frac{2^{\frac{2}{3}} (x^2 + x + 1) + 2 \cdot 2^{\frac{1}{3}} (x^3 + 1)^{\frac{1}{3}} (x + 1) + 2(x^3 + 1)^{\frac{2}{3}}}{x^2 + x + 1}} \right)$$

input `integrate((1-x)/(x^2+x+1)/(x^3+1)^(1/3),x, algorithm="fricas")`

output `1/3*2^(1/6)*sqrt(3/2)*arctan(1/3*2^(1/6)*sqrt(3/2)*(2*2^(2/3)*(x^4 + 4*x^3 + 5*x^2 + 4*x + 1)*(x^3 + 1)^(2/3) + 2^(1/3)*(x^6 + 7*x^5 + 10*x^4 + 7*x^3 + 10*x^2 + 7*x + 1) - 4*(x^5 + x^4 - 3*x^3 - 3*x^2 + x + 1)*(x^3 + 1)^(1/3))/(3*x^6 + 9*x^5 + 6*x^4 + x^3 + 6*x^2 + 9*x + 3)) - 1/12*2^(2/3)*log((2^(2/3)*(x^3 + 1)^(2/3)*(x^2 + 3*x + 1) - 2^(1/3)*(x^4 - 3*x^2 + 1) - 4*(x^3 + 1)^(1/3)*(x^2 + x))/(x^4 + 2*x^3 + 3*x^2 + 2*x + 1)) + 1/6*2^(2/3)*log((2^(2/3)*(x^2 + x + 1) + 2*2^(1/3)*(x^3 + 1)^(1/3)*(x + 1) + 2*(x^3 + 1)^(2/3))/(x^2 + x + 1))`

Sympy [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx = - \int \frac{x}{x^2\sqrt[3]{x^3+1} + x\sqrt[3]{x^3+1} + \sqrt[3]{x^3+1}} dx - \int \left(-\frac{1}{x^2\sqrt[3]{x^3+1} + x\sqrt[3]{x^3+1} + \sqrt[3]{x^3+1}} \right) dx$$

input `integrate((1-x)/(x**2+x+1)/(x**3+1)**(1/3),x)`

output

```
-Integral(x/(x**2*(x**3 + 1)**(1/3) + x*(x**3 + 1)**(1/3) + (x**3 + 1)**(1/3)), x) - Integral(-1/(x**2*(x**3 + 1)**(1/3) + x*(x**3 + 1)**(1/3) + (x**3 + 1)**(1/3)), x)
```

Maxima [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx = \int -\frac{x-1}{(x^3+1)^{\frac{1}{3}}(x^2+x+1)} dx$$

input

```
integrate((1-x)/(x^2+x+1)/(x^3+1)^(1/3),x, algorithm="maxima")
```

output

```
-integrate((x - 1)/((x^3 + 1)^(1/3)*(x^2 + x + 1)), x)
```

Giac [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx = \int -\frac{x-1}{(x^3+1)^{\frac{1}{3}}(x^2+x+1)} dx$$

input

```
integrate((1-x)/(x^2+x+1)/(x^3+1)^(1/3),x, algorithm="giac")
```

output

```
integrate(-(x - 1)/((x^3 + 1)^(1/3)*(x^2 + x + 1)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx = -\int \frac{x-1}{(x^3+1)^{1/3}(x^2+x+1)} dx$$

input

```
int(-(x - 1)/((x^3 + 1)^(1/3)*(x + x^2 + 1)),x)
```


output `-int((x - 1)/((x^3 + 1)^(1/3)*(x + x^2 + 1)), x)`

Reduce [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1+x^3}} dx = -\left(\int \frac{x}{(x^3+1)^{\frac{1}{3}}x^2 + (x^3+1)^{\frac{1}{3}}x + (x^3+1)^{\frac{1}{3}}} dx \right) + \int \frac{1}{(x^3+1)^{\frac{1}{3}}x^2 + (x^3+1)^{\frac{1}{3}}x + (x^3+1)^{\frac{1}{3}}} dx$$

input `int((1-x)/(x^2+x+1)/(x^3+1)^(1/3),x)`

output `- int(x/((x**3 + 1)**(1/3)*x**2 + (x**3 + 1)**(1/3)*x + (x**3 + 1)**(1/3)),x) + int(1/((x**3 + 1)**(1/3)*x**2 + (x**3 + 1)**(1/3)*x + (x**3 + 1)**(1/3)),x)`

3.103 $\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx$

Optimal result	777
Mathematica [A] (verified)	777
Rubi [A] (verified)	778
Maple [A] (verified)	779
Fricas [F]	779
Sympy [F]	779
Maxima [F]	780
Giac [F]	780
Mupad [F(-1)]	780
Reduce [F]	781

Optimal result

Integrand size = 20, antiderivative size = 43

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \frac{1}{\sqrt[3]{1-x^3}} + \frac{x}{\sqrt[3]{1-x^3}} - x^2 \operatorname{Hypergeometric2F1}\left(\frac{2}{3}, \frac{4}{3}, \frac{5}{3}, x^3\right)$$

output `1/(-x^3+1)^(1/3)+x/(-x^3+1)^(1/3)-x^2*hypergeom([2/3, 4/3],[5/3],x^3)`

Mathematica [A] (verified)

Time = 10.17 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \frac{(1+2x)(1-x^3)^{2/3}}{1+x+x^2} + x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)$$

input `Integrate[(1 - x^3)^(2/3)/(1 + x + x^2)^2,x]`

output `((1 + 2*x)*(1 - x^3)^(2/3))/(1 + x + x^2) + x^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3]`

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-x^3)^{2/3}}{(x^2+x+1)^2} dx$$

↓ 2583

$$\int \left(-\frac{2x}{(1-x^3)^{4/3}} + \frac{1}{(1-x^3)^{4/3}} + \frac{x^2}{(1-x^3)^{4/3}} \right) dx$$

↓ 2009

$$x^2 \left(-\text{Hypergeometric2F1} \left(\frac{2}{3}, \frac{4}{3}, \frac{5}{3}, x^3 \right) \right) + \frac{x}{\sqrt[3]{1-x^3}} + \frac{1}{\sqrt[3]{1-x^3}}$$

input `Int[(1 - x^3)^(2/3)/(1 + x + x^2)^2,x]`

output `(1 - x^3)^(-1/3) + x/(1 - x^3)^(1/3) - x^2*Hypergeometric2F1[2/3, 4/3, 5/3, x^3]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2583 `Int[(Px_.)*((c_) + (d_.)*(x_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^3)^(p_.), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [A] (verified)

Time = 0.41 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

method	result	size
risch	$-\frac{(1+2x)(-1+x)}{(-x^3+1)^{\frac{1}{3}}} + x^2 \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right)$	34

input `int((-x^3+1)^(2/3)/(x^2+x+1)^2,x,method=_RETURNVERBOSE)`output `-(1+2*x)*(-1+x)/(-x^3+1)^(1/3)+x^2*hypergeom([1/3,2/3],[5/3],x^3)`**Fricas [F]**

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \int \frac{(-x^3+1)^{\frac{2}{3}}}{(x^2+x+1)^2} dx$$

input `integrate((-x^3+1)^(2/3)/(x^2+x+1)^2,x, algorithm="fricas")`output `integral((-x^3 + 1)^(2/3)/(x^4 + 2*x^3 + 3*x^2 + 2*x + 1), x)`**Sympy [F]**

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \int \frac{(-(x-1)(x^2+x+1))^{\frac{2}{3}}}{(x^2+x+1)^2} dx$$

input `integrate((-x**3+1)**(2/3)/(x**2+x+1)**2,x)`output `Integral(-(x - 1)*(x**2 + x + 1)**(2/3)/(x**2 + x + 1)**2, x)`

Maxima [F]

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \int \frac{(-x^3+1)^{2/3}}{(x^2+x+1)^2} dx$$

input `integrate((-x^3+1)^(2/3)/(x^2+x+1)^2,x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(2/3)/(x^2 + x + 1)^2, x)`

Giac [F]

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \int \frac{(-x^3+1)^{2/3}}{(x^2+x+1)^2} dx$$

input `integrate((-x^3+1)^(2/3)/(x^2+x+1)^2,x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3)/(x^2 + x + 1)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \int \frac{(1-x^3)^{2/3}}{(x^2+x+1)^2} dx$$

input `int((1 - x^3)^(2/3)/(x + x^2 + 1)^2,x)`

output `int((1 - x^3)^(2/3)/(x + x^2 + 1)^2, x)`

Reduce [F]

$$\int \frac{(1-x^3)^{2/3}}{(1+x+x^2)^2} dx = \frac{-(-x^3+1)^{2/3} + 2\left(\int \frac{(-x^3+1)^{2/3}x}{x^5+x^4+x^3-x^2-x-1} dx\right) x^2 + 2\left(\int \frac{(-x^3+1)^{2/3}x}{x^5+x^4+x^3-x^2-x-1} dx\right) x + 2\left(\int \frac{(-x^3+1)^{2/3}}{x^5+x^4+x^3-x^2-x-1} dx\right)}{x^2+x+1}$$

input `int((-x^3+1)^(2/3)/(x^2+x+1)^2,x)`

output `(- (- x**3 + 1)**(2/3) + 2*int(((- x**3 + 1)**(2/3)*x)/(x**5 + x**4 + x**3 - x**2 - x - 1),x)*x**2 + 2*int(((- x**3 + 1)**(2/3)*x)/(x**5 + x**4 + x**3 - x**2 - x - 1),x)*x + 2*int(((- x**3 + 1)**(2/3)*x)/(x**5 + x**4 + x**3 - x**2 - x - 1),x))/(x**2 + x + 1)`

3.104
$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx$$

Optimal result	782
Mathematica [A] (verified)	782
Rubi [A] (verified)	783
Maple [A] (verified)	784
Fricas [F]	784
Sympy [F]	785
Maxima [F]	785
Giac [F]	785
Mupad [F(-1)]	786
Reduce [F]	786

Optimal result

Integrand size = 25, antiderivative size = 43

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = \frac{1}{\sqrt[3]{1-x^3}} + \frac{x}{\sqrt[3]{1-x^3}} - x^2 \text{Hypergeometric2F1}\left(\frac{2}{3}, \frac{4}{3}, \frac{5}{3}, x^3\right)$$

output `1/(-x^3+1)^(1/3)+x/(-x^3+1)^(1/3)-x^2*hypergeom([2/3, 4/3],[5/3],x^3)`

Mathematica [A] (verified)

Time = 10.14 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = \frac{(1+2x)(1-x^3)^{2/3}}{1+x+x^2} + x^2 \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)$$

input `Integrate[(1-x)/((1+x+x^2)*(1-x^3)^(1/3)),x]`

output $((1 + 2x)(1 - x^3)^{2/3})/(1 + x + x^2) + x^2 \text{Hypergeometric2F1}[1/3, 2/3, 5/3, x^3]$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{1-x}{(x^2+x+1)\sqrt[3]{1-x^3}} dx$$

↓ 2583

$$\int \left(-\frac{2x}{(1-x^3)^{4/3}} + \frac{1}{(1-x^3)^{4/3}} + \frac{x^2}{(1-x^3)^{4/3}} \right) dx$$

↓ 2009

$$x^2 \left(-\text{Hypergeometric2F1} \left(\frac{2}{3}, \frac{4}{3}, \frac{5}{3}, x^3 \right) \right) + \frac{x}{\sqrt[3]{1-x^3}} + \frac{1}{\sqrt[3]{1-x^3}}$$

input $\text{Int}[(1-x)/((1+x+x^2)*(1-x^3)^{1/3}),x]$

output $(1-x^3)^{-1/3} + x/(1-x^3)^{1/3} - x^2 \text{Hypergeometric2F1}[2/3, 4/3, 5/3, x^3]$

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2583 `Int[(Px_)*((c_) + (d_)*(x_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^3)^(p_), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

method	result	size
risch	$-\frac{(1+2x)(-1+x)}{(-x^3+1)^{\frac{1}{3}}} + x^2 \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right)$	34

input `int((1-x)/(x^2+x+1)/(-x^3+1)^(1/3),x,method=_RETURNVERBOSE)`

output `-(1+2*x)*(-1+x)/(-x^3+1)^(1/3)+x^2*hypergeom([1/3,2/3],[5/3],x^3)`

Fricas [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = \int -\frac{x-1}{(-x^3+1)^{\frac{1}{3}}(x^2+x+1)} dx$$

input `integrate((1-x)/(x^2+x+1)/(-x^3+1)^(1/3),x, algorithm="fricas")`

output `integral((-x^3 + 1)^(2/3)/(x^4 + 2*x^3 + 3*x^2 + 2*x + 1), x)`

Sympy [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = - \int \frac{x}{x^2\sqrt[3]{1-x^3} + x\sqrt[3]{1-x^3} + \sqrt[3]{1-x^3}} dx - \int \left(-\frac{1}{x^2\sqrt[3]{1-x^3} + x\sqrt[3]{1-x^3} + \sqrt[3]{1-x^3}} \right) dx$$

input `integrate((1-x)/(x**2+x+1)/(-x**3+1)**(1/3), x)`

output `-Integral(x/(x**2*(1 - x**3)**(1/3) + x*(1 - x**3)**(1/3) + (1 - x**3)**(1/3)), x) - Integral(-1/(x**2*(1 - x**3)**(1/3) + x*(1 - x**3)**(1/3) + (1 - x**3)**(1/3)), x)`

Maxima [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = \int -\frac{x-1}{(-x^3+1)^{\frac{1}{3}}(x^2+x+1)} dx$$

input `integrate((1-x)/(x^2+x+1)/(-x^3+1)^(1/3), x, algorithm="maxima")`

output `-integrate((x - 1)/((-x^3 + 1)^(1/3)*(x^2 + x + 1)), x)`

Giac [F]

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = \int -\frac{x-1}{(-x^3+1)^{\frac{1}{3}}(x^2+x+1)} dx$$

input `integrate((1-x)/(x^2+x+1)/(-x^3+1)^(1/3), x, algorithm="giac")`

output `integrate(-(x - 1)/((-x^3 + 1)^(1/3)*(x^2 + x + 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = - \int \frac{x-1}{(1-x^3)^{1/3}(x^2+x+1)} dx$$

input `int(-(x - 1)/((1 - x^3)^(1/3)*(x + x^2 + 1)),x)`output `-int((x - 1)/((1 - x^3)^(1/3)*(x + x^2 + 1)), x)`**Reduce [F]**

$$\int \frac{1-x}{(1+x+x^2)\sqrt[3]{1-x^3}} dx = - \left(\int \frac{x}{(-x^3+1)^{\frac{1}{3}}x^2 + (-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx \right) + \int \frac{1}{(-x^3+1)^{\frac{1}{3}}x^2 + (-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}} dx$$

input `int((1-x)/(x^2+x+1)/(-x^3+1)^(1/3),x)`output `- int(x/((- x**3 + 1)**(1/3)*x**2 + (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x) + int(1/((- x**3 + 1)**(1/3)*x**2 + (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)`

3.105

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx$$

Optimal result	787
Mathematica [A] (verified)	787
Rubi [A] (verified)	788
Maple [A] (verified)	789
Fricas [F]	789
Sympy [F]	790
Maxima [F]	790
Giac [F]	790
Mupad [F(-1)]	791
Reduce [F]	791

Optimal result

Integrand size = 19, antiderivative size = 39

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = \frac{1 + (1-2x)x}{\sqrt[3]{1-x^3}} + x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)$$

output `(1+(1-2*x)*x)/(-x^3+1)^(1/3)+x^2*hypergeom([1/3, 2/3],[5/3],x^3)`

Mathematica [A] (verified)

Time = 10.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = \frac{1}{\sqrt[3]{1-x^3}} + \frac{x}{\sqrt[3]{1-x^3}} - x^2 \operatorname{Hypergeometric2F1}\left(\frac{2}{3}, \frac{4}{3}, \frac{5}{3}, x^3\right)$$

input `Integrate[(1 - x)^2/(1 - x^3)^(4/3), x]`

output `(1 - x^3)^(-1/3) + x/(1 - x^3)^(1/3) - x^2*Hypergeometric2F1[2/3, 4/3, 5/3, x^3]`

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2393, 27, 888}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx$$

↓ 2393

$$\frac{(1-2x)x+1}{\sqrt[3]{1-x^3}} - \int -\frac{2x}{\sqrt[3]{1-x^3}} dx$$

↓ 27

$$2 \int \frac{x}{\sqrt[3]{1-x^3}} dx + \frac{(1-2x)x+1}{\sqrt[3]{1-x^3}}$$

↓ 888

$$x^2 \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) + \frac{(1-2x)x+1}{\sqrt[3]{1-x^3}}$$

input `Int[(1 - x)^2/(1 - x^3)^(4/3), x]`

output `(1 + (1 - 2*x)*x)/(1 - x^3)^(1/3) + x^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3]`

Defintions of rubi rules used

rule 27 `Int[(a_)*(F_x_), x_Symbol] := Simp[a Int[F_x, x], x] /; FreeQ[a, x] && !MatchQ[F_x, (b_)*(G_x_)] /; FreeQ[b, x]`

rule 888

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p
*((c*x)^(m + 1)/(c*(m + 1))]*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1
, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt
Q[p, 0] || GtQ[a, 0])
```

rule 2393

```
Int[(Pq_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Module[{q = Expon[Pq,
x], i}, Simp[(a*Coeff[Pq, x, q] - b*x*ExpandToSum[Pq - Coeff[Pq, x, q]*x^q
, x])*(a + b*x^n)^(p + 1)/(a*b*n*(p + 1)), x] + Simp[1/(a*n*(p + 1)) In
t[Sum[(n*(p + 1) + i + 1)*Coeff[Pq, x, i]*x^i, {i, 0, q - 1}]*a + b*x^n)^(
p + 1), x], x] /; q == n - 1] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[n
, 0] && LtQ[p, -1]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.87

method	result	size
risch	$-\frac{(1+2x)(-1+x)}{(-x^3+1)^{\frac{1}{3}}} + x^2 \operatorname{hypergeom}\left(\left[\frac{1}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right)$	34
meijerg	$\frac{x}{(-x^3+1)^{\frac{1}{3}}} - x^2 \operatorname{hypergeom}\left(\left[\frac{2}{3}, \frac{4}{3}\right], \left[\frac{5}{3}\right], x^3\right) + \frac{x^3 \operatorname{hypergeom}\left(\left[1, \frac{4}{3}\right], [2], x^3\right)}{3}$	41

input

```
int((1-x)^2/(-x^3+1)^(4/3),x,method=_RETURNVERBOSE)
```

output

```
-(1+2*x)*(-1+x)/(-x^3+1)^(1/3)+x^2*hypergeom([1/3,2/3],[5/3],x^3)
```

Fricas [F]

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = \int \frac{(x-1)^2}{(-x^3+1)^{\frac{4}{3}}} dx$$

input

```
integrate((1-x)^2/(-x^3+1)^(4/3),x, algorithm="fricas")
```

output `integral((-x^3 + 1)^(2/3)/(x^4 + 2*x^3 + 3*x^2 + 2*x + 1), x)`

Sympy [F]

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = \int \frac{(x-1)^2}{(-(x-1)(x^2+x+1))^{4/3}} dx$$

input `integrate((1-x)**2/(-x**3+1)**(4/3),x)`

output `Integral((x - 1)**2/(-(x - 1)*(x**2 + x + 1))**(4/3), x)`

Maxima [F]

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = \int \frac{(x-1)^2}{(-x^3+1)^{4/3}} dx$$

input `integrate((1-x)^2/(-x^3+1)^(4/3),x, algorithm="maxima")`

output `x/(-x^3 + 1)^(1/3) - integrate((x^2 - 2*x)/((x^3 - 1)*(x^2 + x + 1)^(1/3)*(-x + 1)^(1/3)), x)`

Giac [F]

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = \int \frac{(x-1)^2}{(-x^3+1)^{4/3}} dx$$

input `integrate((1-x)^2/(-x^3+1)^(4/3),x, algorithm="giac")`

output `integrate((x - 1)^2/(-x^3 + 1)^(4/3), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = \int \frac{(x-1)^2}{(1-x^3)^{4/3}} dx$$

input `int((x - 1)^2/(1 - x^3)^(4/3),x)`output `int((x - 1)^2/(1 - x^3)^(4/3), x)`**Reduce [F]**

$$\int \frac{(1-x)^2}{(1-x^3)^{4/3}} dx = - \left(\int \frac{x}{(-x^3+1)^{1/3} x^2 + (-x^3+1)^{1/3} x + (-x^3+1)^{1/3}} dx \right) + \int \frac{1}{(-x^3+1)^{1/3} x^2 + (-x^3+1)^{1/3} x + (-x^3+1)^{1/3}} dx$$

input `int((1-x)^2/(-x^3+1)^(4/3),x)`output `- int(x/((- x**3 + 1)**(1/3)*x**2 + (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x) + int(1/((- x**3 + 1)**(1/3)*x**2 + (- x**3 + 1)**(1/3)*x + (- x**3 + 1)**(1/3)),x)`

3.106 $\int (1 - x^3)^{2/3} dx$

Optimal result	792
Mathematica [C] (warning: unable to verify)	792
Rubi [A] (verified)	793
Maple [C] (verified)	794
Fricas [A] (verification not implemented)	795
Sympy [C] (verification not implemented)	795
Maxima [B] (verification not implemented)	796
Giac [F]	796
Mupad [B] (verification not implemented)	797
Reduce [F]	797

Optimal result

Integrand size = 11, antiderivative size = 67

$$\int (1 - x^3)^{2/3} dx = \frac{1}{3}x(1 - x^3)^{2/3} - \frac{2 \arctan\left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{1}{3} \log\left(x + \sqrt[3]{1-x^3}\right)$$

output

$1/3*x*(-x^3+1)^{(2/3)}+1/3*\ln(x+(-x^3+1)^{(1/3)})-2/9*\arctan(1/3*(1-2*x/(-x^3+1)^{(1/3}))*3^{(1/2)})*3^{(1/2)}$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.08 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.51

$$\int (1-x^3)^{2/3} dx = \frac{3(-1+x)(1-x^3)^{2/3} \operatorname{AppellF1}\left(\frac{5}{3}, -\frac{2}{3}, -\frac{2}{3}, \frac{8}{3}, -\frac{-1+x}{1-(-1)^{2/3}}, -\frac{-1+x}{1+\sqrt[3]{-1}}\right)}{5\left(1+\frac{-1+x}{1+\sqrt[3]{-1}}\right)^{2/3}\left(1+\frac{-1+x}{1-(-1)^{2/3}}\right)^{2/3}}$$

input

`Integrate[(1 - x^3)^(2/3), x]`

output

```
(3*(-1 + x)*(1 - x^3)^(2/3)*AppellF1[5/3, -2/3, -2/3, 8/3, -((-1 + x)/(1 - (-1)^(2/3))), -((-1 + x)/(1 + (-1)^(1/3)))])/(5*(1 + (-1 + x)/(1 + (-1)^(1/3)))^(2/3)*(1 + (-1 + x)/(1 - (-1)^(2/3)))^(2/3))
```

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.04, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {748, 769}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int (1 - x^3)^{2/3} dx$$

$$\downarrow 748$$

$$\frac{2}{3} \int \frac{1}{\sqrt[3]{1 - x^3}} dx + \frac{1}{3} (1 - x^3)^{2/3} x$$

$$\downarrow 769$$

$$\frac{2}{3} \left(\frac{1}{2} \log \left(\sqrt[3]{1 - x^3} + x \right) - \frac{\arctan \left(\frac{1 - \frac{2x}{\sqrt[3]{1 - x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} \right) + \frac{1}{3} (1 - x^3)^{2/3} x$$

input

```
Int[(1 - x^3)^(2/3), x]
```

output

```
(x*(1 - x^3)^(2/3))/3 + (2*(-(ArcTan[(1 - (2*x))/(1 - x^3)^(1/3)]/Sqrt[3]]/Sqrt[3]) + Log[x + (1 - x^3)^(1/3)]/2))/3
```

Definitions of rubi rules used

rule 748 $\text{Int}[(a_ + (b_ \cdot x_)^{n_})^{p_}, x_Symbol] \rightarrow \text{Simp}[x \cdot (a + b \cdot x^n)^p / (n \cdot p + 1), x] + \text{Simp}[a \cdot n \cdot (p / (n \cdot p + 1)) \text{Int}[(a + b \cdot x^n)^{p-1}, x], x] /;$ FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || LtQ[Denominator[p + 1/n], Denominator[p]])

rule 769 $\text{Int}[(a_ + (b_ \cdot x_)^3)^{-1/3}, x_Symbol] \rightarrow \text{Simp}[\text{ArcTan}[(1 + 2 \cdot \text{Rt}[b, 3] \cdot (x / (a + b \cdot x^3)^{1/3})) / \text{Sqrt}[3]] / (\text{Sqrt}[3] \cdot \text{Rt}[b, 3]), x] - \text{Simp}[\text{Log}[(a + b \cdot x^3)^{1/3} - \text{Rt}[b, 3] \cdot x] / (2 \cdot \text{Rt}[b, 3]), x] /;$ FreeQ[{a, b}, x]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 1.47 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.18

method	result
meijerg	$x \text{ hypergeom} \left(\left[-\frac{2}{3}, \frac{1}{3} \right], \left[\frac{4}{3} \right], x^3 \right)$
risch	$-\frac{x(x^3-1)}{3(-x^3+1)^{\frac{1}{3}}} + \frac{2x \text{ hypergeom} \left(\left[\frac{1}{3}, \frac{1}{3} \right], \left[\frac{4}{3} \right], x^3 \right)}{3}$
pseudoelliptic	$\frac{3x(-x^3+1)^{\frac{2}{3}} + 2\sqrt{3} \arctan \left(\frac{(-2(-x^3+1)^{\frac{1}{3}}+x)\sqrt{3}}{3x} \right) - \ln \left(\frac{(-x^3+1)^{\frac{2}{3}} - x(-x^3+1)^{\frac{1}{3}} + x^2}{x^2} \right) + 2 \ln \left(\frac{x + (-x^3+1)^{\frac{1}{3}}}{x} \right)}{9 \left((-x^3+1)^{\frac{2}{3}} - x(-x^3+1)^{\frac{1}{3}} + x^2 \right) (x + (-x^3+1)^{\frac{1}{3}})}$
trager	$\frac{x(-x^3+1)^{\frac{2}{3}}}{3} + \frac{2 \text{RootOf}(_Z^2 + _Z + 1) \ln \left(\text{RootOf}(_Z^2 + _Z + 1)^2 x^3 + 3 \text{RootOf}(_Z^2 + _Z + 1) (-x^3+1)^{\frac{2}{3}} x - \dots \right)}{3}$

input `int((-x^3+1)^(2/3),x,method=_RETURNVERBOSE)`

output `x*hypergeom([-2/3,1/3],[4/3],x^3)`

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.40

$$\int (1 - x^3)^{2/3} dx = \frac{1}{3} (-x^3 + 1)^{\frac{2}{3}} x - \frac{2}{9} \sqrt{3} \arctan \left(-\frac{\sqrt{3}x - 2\sqrt{3}(-x^3 + 1)^{\frac{1}{3}}}{3x} \right) + \frac{2}{9} \log \left(\frac{x + (-x^3 + 1)^{\frac{1}{3}}}{x} \right) - \frac{1}{9} \log \left(\frac{x^2 - (-x^3 + 1)^{\frac{1}{3}}x + (-x^3 + 1)^{\frac{2}{3}}}{x^2} \right)$$

input `integrate((-x^3+1)^(2/3),x, algorithm="fricas")`

output `1/3*(-x^3 + 1)^(2/3)*x - 2/9*sqrt(3)*arctan(-1/3*(sqrt(3)*x - 2*sqrt(3)*(-x^3 + 1)^(1/3))/x) + 2/9*log((x + (-x^3 + 1)^(1/3))/x) - 1/9*log((x^2 - (-x^3 + 1)^(1/3)*x + (-x^3 + 1)^(2/3))/x^2)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.53 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.46

$$\int (1 - x^3)^{2/3} dx = \frac{x\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(-\frac{2}{3}, \frac{1}{3} \middle| \frac{4}{3} \middle| x^3 e^{2i\pi}\right)}{3\Gamma\left(\frac{4}{3}\right)}$$

input `integrate((-x**3+1)**(2/3),x)`

output `x*gamma(1/3)*hyper((-2/3, 1/3), (4/3,), x**3*exp_polar(2*I*pi))/(3*gamma(4/3))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. $2(52) = 104$.

Time = 0.14 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.57

$$\int (1 - x^3)^{2/3} dx = -\frac{2}{9} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \left(\frac{2(-x^3 + 1)^{1/3}}{x} - 1 \right) \right) - \frac{(-x^3 + 1)^{2/3}}{3x^2 \left(\frac{x^3 - 1}{x^3} - 1 \right)}$$

$$+ \frac{2}{9} \log \left(\frac{(-x^3 + 1)^{1/3}}{x} + 1 \right) - \frac{1}{9} \log \left(-\frac{(-x^3 + 1)^{1/3}}{x} + \frac{(-x^3 + 1)^{2/3}}{x^2} + 1 \right)$$

input `integrate((-x^3+1)^(2/3),x, algorithm="maxima")`

output `-2/9*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3)/x - 1)) - 1/3*(-x^3 + 1)^(2/3)/(x^2*((x^3 - 1)/x^3 - 1)) + 2/9*log((-x^3 + 1)^(1/3)/x + 1) - 1/9*log(-(-x^3 + 1)^(1/3)/x + (-x^3 + 1)^(2/3)/x^2 + 1)`

Giac [F]

$$\int (1 - x^3)^{2/3} dx = \int (-x^3 + 1)^{2/3} dx$$

input `integrate((-x^3+1)^(2/3),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3), x)`

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.15

$$\int (1 - x^3)^{2/3} dx = x {}_2F_1\left(-\frac{2}{3}, \frac{1}{3}; \frac{4}{3}; x^3\right)$$

input `int((1 - x^3)^(2/3),x)`output `x*hypergeom([-2/3, 1/3], 4/3, x^3)`**Reduce [F]**

$$\int (1 - x^3)^{2/3} dx = \frac{(-x^3 + 1)^{\frac{2}{3}} x}{3} - \frac{2 \left(\int \frac{(-x^3 + 1)^{\frac{2}{3}}}{x^3 - 1} dx \right)}{3}$$

input `int((-x^3+1)^(2/3),x)`output `((- x**3 + 1)**(2/3)*x - 2*int((- x**3 + 1)**(2/3)/(x**3 - 1),x))/3`

3.107 $\int \frac{(1-x^3)^{2/3}}{x} dx$

Optimal result	798
Mathematica [A] (verified)	798
Rubi [A] (verified)	799
Maple [C] (verified)	801
Fricas [A] (verification not implemented)	801
Sympy [C] (verification not implemented)	802
Maxima [A] (verification not implemented)	802
Giac [A] (verification not implemented)	803
Mupad [B] (verification not implemented)	803
Reduce [F]	804

Optimal result

Integrand size = 15, antiderivative size = 70

$$\int \frac{(1-x^3)^{2/3}}{x} dx = \frac{1}{2}(1-x^3)^{2/3} + \frac{\arctan\left(\frac{1+2\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\log(x)}{2} + \frac{1}{2}\log\left(1-\sqrt[3]{1-x^3}\right)$$

output `1/2*(-x^3+1)^(2/3)-1/2*ln(x)+1/2*ln(1-(-x^3+1)^(1/3))+1/3*arctan(1/3*(1+2*(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)`

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.29

$$\int \frac{(1-x^3)^{2/3}}{x} dx = \frac{1}{6}\left(3(1-x^3)^{2/3} + 2\sqrt{3}\arctan\left(\frac{1+2\sqrt[3]{1-x^3}}{\sqrt{3}}\right) + 2\log\left(-1+\sqrt[3]{1-x^3}\right) - \log\left(1+\sqrt[3]{1-x^3}+(1-x^3)^{2/3}\right)\right)$$

input `Integrate[(1 - x^3)^(2/3)/x,x]`

output

$$(3*(1 - x^3)^{(2/3)} + 2*\text{Sqrt}[3]*\text{ArcTan}[(1 + 2*(1 - x^3)^{(1/3)})/\text{Sqrt}[3]] + 2*\text{Log}[-1 + (1 - x^3)^{(1/3)}] - \text{Log}[1 + (1 - x^3)^{(1/3)} + (1 - x^3)^{(2/3)}])/6$$
Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.09, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {798, 60, 67, 16, 1083, 217}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-x^3)^{2/3}}{x} dx$$

$$\downarrow 798$$

$$\frac{1}{3} \int \frac{(1-x^3)^{2/3}}{x^3} dx^3$$

$$\downarrow 60$$

$$\frac{1}{3} \left(\int \frac{1}{x^3 \sqrt[3]{1-x^3}} dx^3 + \frac{3}{2} (1-x^3)^{2/3} \right)$$

$$\downarrow 67$$

$$\frac{1}{3} \left(-\frac{3}{2} \int \frac{1}{1-\sqrt[3]{1-x^3}} d\sqrt[3]{1-x^3} + \frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{1-x^3} + 1} d\sqrt[3]{1-x^3} + \frac{3}{2} (1-x^3)^{2/3} - \frac{1}{2} \log(x^3) \right)$$

$$\downarrow 16$$

$$\frac{1}{3} \left(\frac{3}{2} \int \frac{1}{x^6 + \sqrt[3]{1-x^3} + 1} d\sqrt[3]{1-x^3} + \frac{3}{2} (1-x^3)^{2/3} - \frac{1}{2} \log(x^3) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^3}) \right)$$

$$\downarrow 1083$$

$$\frac{1}{3} \left(-3 \int \frac{1}{-x^6 - 3} d(2\sqrt[3]{1-x^3} + 1) + \frac{3}{2} (1-x^3)^{2/3} - \frac{1}{2} \log(x^3) + \frac{3}{2} \log(1 - \sqrt[3]{1-x^3}) \right)$$

$$\downarrow 217$$

$$\frac{1}{3} \left(\sqrt{3} \arctan \left(\frac{2\sqrt[3]{1-x^3}+1}{\sqrt{3}} \right) + \frac{3}{2} (1-x^3)^{2/3} - \frac{\log(x^3)}{2} + \frac{3}{2} \log \left(1 - \sqrt[3]{1-x^3} \right) \right)$$

input `Int[(1 - x^3)^(2/3)/x,x]`

output `((3*(1 - x^3)^(2/3))/2 + Sqrt[3]*ArcTan[(1 + 2*(1 - x^3)^(1/3))/Sqrt[3]] - Log[x^3]/2 + (3*Log[1 - (1 - x^3)^(1/3)])/2)/3`

Defintions of rubi rules used

rule 16 `Int[(c_.)/((a_.) + (b_.)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 60 `Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Simp[n*(b*c - a*d)/(b*(m + n + 1)) Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

rule 67 `Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Simp[3/(2*b) Subst[Int[1/(q^2 + q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Simp[3/(2*b*q) Subst[Int[1/(q - x), x], x, (c + d*x)^(1/3)], x]) /; FreeQ[{a, b, c, d}, x] && PosQ[(b*c - a*d)/b]`

rule 217 `Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 798 `Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[1/n Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

rule 1083

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Simp[-2 Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3.

Time = 2.07 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.94

method	result
meijerg	$-\frac{\sqrt{3}\Gamma\left(\frac{2}{3}\right)\left(-\frac{\left(\frac{3}{2}-\frac{\pi\sqrt{3}}{6}-\frac{3\ln(3)}{2}+3\ln(x)+i\pi\right)\pi\sqrt{3}}{\Gamma\left(\frac{2}{3}\right)}+\frac{2\pi\sqrt{3}x^3\operatorname{hypergeom}\left(\left[\frac{1}{3},1,1\right],[2,2],x^3\right)}{3\Gamma\left(\frac{2}{3}\right)}\right)}{9\pi}$
pseudoelliptic	$\frac{(-x^3+1)^{\frac{2}{3}}}{2}-\frac{\ln\left((-x^3+1)^{\frac{2}{3}}+(-x^3+1)^{\frac{1}{3}}+1\right)}{6}+\frac{\arctan\left(\frac{\left(1+2(-x^3+1)^{\frac{1}{3}}\right)\sqrt{3}}{3}\right)\sqrt{3}}{3}+\frac{\ln\left((-x^3+1)^{\frac{1}{3}}-1\right)}{3}$
trager	$\frac{(-x^3+1)^{\frac{2}{3}}}{2}+\frac{\ln\left(-\frac{1438\operatorname{RootOf}\left(-Z^2+_Z+1\right)^2x^3+9855\operatorname{RootOf}\left(-Z^2+_Z+1\right)x^3-5502\operatorname{RootOf}\left(-Z^2+_Z+1\right)\left(-x^3\right)}{\dots}\right)}{\dots}$

input

```
int((-x^3+1)^(2/3)/x,x,method=_RETURNVERBOSE)
```

output

```
-1/9/Pi*3^(1/2)*GAMMA(2/3)*(-3/2-1/6*Pi*3^(1/2)-3/2*ln(3)+3*ln(x)+I*Pi)*Pi*3^(1/2)/GAMMA(2/3)+2/3*Pi*3^(1/2)/GAMMA(2/3)*x^3*hypergeom([1/3,1,1],[2,2],x^3))
```

Fricas [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.07

$$\int \frac{(1-x^3)^{2/3}}{x} dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{2}{3} \sqrt{3} (-x^3+1)^{\frac{1}{3}} + \frac{1}{3} \sqrt{3} \right) + \frac{1}{2} (-x^3+1)^{\frac{2}{3}} - \frac{1}{6} \log \left((-x^3+1)^{\frac{2}{3}} + (-x^3+1)^{\frac{1}{3}} + 1 \right) + \frac{1}{3} \log \left((-x^3+1)^{\frac{1}{3}} - 1 \right)$$

input `integrate((-x^3+1)^(2/3)/x,x, algorithm="fricas")`

output `1/3*sqrt(3)*arctan(2/3*sqrt(3)*(-x^3 + 1)^(1/3) + 1/3*sqrt(3)) + 1/2*(-x^3 + 1)^(2/3) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log((-x^3 + 1)^(1/3) - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.52 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.59

$$\int \frac{(1-x^3)^{2/3}}{x} dx = -\frac{x^2 e^{\frac{2i\pi}{3}} \Gamma\left(-\frac{2}{3}\right) {}_2F_1\left(\begin{matrix} -\frac{2}{3}, -\frac{2}{3} \\ \frac{1}{3} \end{matrix} \middle| \frac{1}{x^3}\right)}{3\Gamma\left(\frac{1}{3}\right)}$$

input `integrate((-x**3+1)**(2/3)/x,x)`

output `-x**2*exp(2*I*pi/3)*gamma(-2/3)*hyper((-2/3, -2/3), (1/3,), x**(-3))/(3*gamma(1/3))`

Maxima [A] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.04

$$\int \frac{(1-x^3)^{2/3}}{x} dx = \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(2(-x^3+1)^{\frac{1}{3}}+1\right)\right) + \frac{1}{2} (-x^3+1)^{\frac{2}{3}} - \frac{1}{6} \log\left(\left(-x^3+1\right)^{\frac{2}{3}} + \left(-x^3+1\right)^{\frac{1}{3}} + 1\right) + \frac{1}{3} \log\left(\left(-x^3+1\right)^{\frac{1}{3}} - 1\right)$$

input `integrate((-x^3+1)^(2/3)/x,x, algorithm="maxima")`

output

```
1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3) + 1)) + 1/2*(-x^3 + 1)^(2/3) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log((-x^3 + 1)^(1/3) - 1)
```

Giac [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.06

$$\int \frac{(1-x^3)^{2/3}}{x} dx = \frac{1}{3} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3} \left(2(-x^3+1)^{1/3} + 1 \right) \right) + \frac{1}{2} (-x^3+1)^{2/3} - \frac{1}{6} \log \left((-x^3+1)^{2/3} + (-x^3+1)^{1/3} + 1 \right) + \frac{1}{3} \log \left(\left| (-x^3+1)^{1/3} - 1 \right| \right)$$

input

```
integrate((-x^3+1)^(2/3)/x,x, algorithm="giac")
```

output

```
1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(-x^3 + 1)^(1/3) + 1)) + 1/2*(-x^3 + 1)^(2/3) - 1/6*log((-x^3 + 1)^(2/3) + (-x^3 + 1)^(1/3) + 1) + 1/3*log(abs((-x^3 + 1)^(1/3) - 1))
```

Mupad [B] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.30

$$\int \frac{(1-x^3)^{2/3}}{x} dx = \frac{\ln \left((1-x^3)^{1/3} - 1 \right)}{3} + \ln \left((1-x^3)^{1/3} - 9 \left(-\frac{1}{6} + \frac{\sqrt{3} \operatorname{li}}{6} \right)^2 \right) \left(-\frac{1}{6} + \frac{\sqrt{3} \operatorname{li}}{6} \right) - \ln \left((1-x^3)^{1/3} - 9 \left(\frac{1}{6} + \frac{\sqrt{3} \operatorname{li}}{6} \right)^2 \right) \left(\frac{1}{6} + \frac{\sqrt{3} \operatorname{li}}{6} \right)$$

input

```
int((1 - x^3)^(2/3)/x,x)
```

output

```
log((1 - x^3)^(1/3) - 1)/3 + log((1 - x^3)^(1/3) - 9*((3^(1/2)*1i)/6 - 1/6)^2)*((3^(1/2)*1i)/6 - 1/6) - log((1 - x^3)^(1/3) - 9*((3^(1/2)*1i)/6 + 1/6)^2)*((3^(1/2)*1i)/6 + 1/6) + (1 - x^3)^(2/3)/2
```

Reduce [F]

$$\int \frac{(1-x^3)^{2/3}}{x} dx = \frac{(-x^3+1)^{2/3}}{2} - \left(\int \frac{(-x^3+1)^{2/3}}{x^4-x} dx \right)$$

input `int((-x^3+1)^(2/3)/x,x)`

output `((-x**3 + 1)**(2/3) - 2*int((-x**3 + 1)**(2/3)/(x**4 - x),x))/2`

3.108 $\int \frac{(1-x^3)^{2/3}}{a+bx} dx$

Optimal result	805
Mathematica [F]	806
Rubi [A] (verified)	806
Maple [F]	809
Fricas [F(-1)]	810
Sympy [F]	810
Maxima [F]	810
Giac [F]	811
Mupad [F(-1)]	811
Reduce [F]	811

Optimal result

Integrand size = 19, antiderivative size = 384

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \frac{(1-x^3)^{2/3}}{2b} - \frac{(a^3+b^3)x^2 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{3}, 1, \frac{5}{3}, x^3, -\frac{b^3x^3}{a^3}\right)}{2a^2b^2}$$

$$+ \frac{a^2 \arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}b^3} - \frac{(a^3+b^3)^{2/3} \arctan\left(\frac{1-2\sqrt[3]{a^3+b^3x}}{a\sqrt[3]{1-x^3}}\right)}{\sqrt{3}b^3}$$

$$+ \frac{(a^3+b^3)^{2/3} \arctan\left(\frac{1+\frac{2b\sqrt[3]{1-x^3}}{\sqrt[3]{a^3+b^3}}}{\sqrt{3}}\right)}{\sqrt{3}b^3} + \frac{ax^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)}{2b^2}$$

$$- \frac{(a^3+b^3)^{2/3} \log(a^3+b^3x^3)}{3b^3} + \frac{(a^3+b^3)^{2/3} \log\left(-\frac{\sqrt[3]{a^3+b^3x}}{a} - \sqrt[3]{1-x^3}\right)}{2b^3}$$

$$- \frac{a^2 \log\left(x + \sqrt[3]{1-x^3}\right)}{2b^3} + \frac{(a^3+b^3)^{2/3} \log\left(\sqrt[3]{a^3+b^3} - b\sqrt[3]{1-x^3}\right)}{2b^3}$$

output

```

1/2*(-x^3+1)^(2/3)/b-1/2*(a^3+b^3)*x^2*AppellF1(2/3,1/3,1,5/3,x^3,-b^3*x^3
/a^3)/a^2/b^2+1/2*a*x^2*hypergeom([1/3, 2/3],[5/3],x^3)/b^2-1/3*(a^3+b^3)^(
2/3)*ln(b^3*x^3+a^3)/b^3+1/2*(a^3+b^3)^(2/3)*ln(-(a^3+b^3)^(1/3)*x/a-(-x^
3+1)^(1/3))/b^3-1/2*a^2*ln(x+(-x^3+1)^(1/3))/b^3+1/2*(a^3+b^3)^(2/3)*ln((a
^3+b^3)^(1/3)-b*(-x^3+1)^(1/3))/b^3+1/3*a^2*arctan(1/3*(1-2*x/(-x^3+1)^(1/
3))*3^(1/2))/b^3*3^(1/2)-1/3*(a^3+b^3)^(2/3)*arctan(1/3*(1-2*(a^3+b^3)^(1/
3)*x/a/(-x^3+1)^(1/3))*3^(1/2))/b^3*3^(1/2)+1/3*(a^3+b^3)^(2/3)*arctan(1/3
*(1+2*b*(-x^3+1)^(1/3)/(a^3+b^3)^(1/3))*3^(1/2))/b^3*3^(1/2)

```

Mathematica [F]

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \int \frac{(1-x^3)^{2/3}}{a+bx} dx$$

input

```
Integrate[(1 - x^3)^(2/3)/(a + b*x), x]
```

output

```
Integrate[(1 - x^3)^(2/3)/(a + b*x), x]
```

Rubi [A] (verified)

Time = 0.74 (sec) , antiderivative size = 375, normalized size of antiderivative = 0.98, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2578, 888, 2577, 769, 2581, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{array}{c}
 \int \frac{(1-x^3)^{2/3}}{a+bx} dx \\
 \downarrow \text{2578} \\
 \int \frac{b^2-a^2x}{(a+bx)\sqrt[3]{1-x^3}} dx + a \int \frac{x}{\sqrt[3]{1-x^3}} dx + \frac{(1-x^3)^{2/3}}{2b} \\
 \downarrow \text{888}
 \end{array}$$

$$\begin{aligned}
 & \frac{\int \frac{b^2 - a^2 x}{(a+bx) \sqrt[3]{1-x^3}} dx}{b^2} + \frac{ax^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)}{2b^2} + \frac{(1-x^3)^{2/3}}{2b} \\
 & \quad \downarrow \text{2577} \\
 & \frac{(a^3+b^3) \int \frac{1}{(a+bx) \sqrt[3]{1-x^3}} dx}{b^2} - \frac{a^2 \int \frac{1}{\sqrt[3]{1-x^3}} dx}{b} + \frac{ax^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)}{2b^2} + \frac{(1-x^3)^{2/3}}{2b} \\
 & \quad \downarrow \text{769} \\
 & \frac{(a^3+b^3) \int \frac{1}{(a+bx) \sqrt[3]{1-x^3}} dx}{b^2} - \frac{a^2 \left(\frac{1}{2} \log\left(\sqrt[3]{1-x^3}+x\right) - \frac{\arctan\left(\frac{1-\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} \right)}{b} \\
 & \quad \downarrow \text{2581} \\
 & \frac{(a^3+b^3) \int \left(\frac{a^2}{\sqrt[3]{1-x^3}(a^3+b^3x^3)} - \frac{bxa}{\sqrt[3]{1-x^3}(a^3+b^3x^3)} + \frac{b^2x^2}{\sqrt[3]{1-x^3}(a^3+b^3x^3)} \right) dx}{b^2} - \frac{a^2 \left(\frac{1}{2} \log\left(\sqrt[3]{1-x^3}+x\right) - \frac{\arctan\left(\frac{1-\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} \right)}{b} \\
 & \quad \downarrow \text{2009} \\
 & \frac{ax^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)}{2b^2} + \frac{(1-x^3)^{2/3}}{2b}
 \end{aligned}$$

$$\frac{(a^3+b^3) \left(\frac{\arctan\left(\frac{1-2x\sqrt[3]{a^3+b^3}}{a\sqrt[3]{1-x^3}}\right)}{\sqrt[3]{a^3+b^3}} + \frac{\arctan\left(\frac{2b\sqrt[3]{1-x^3}+1}{\sqrt[3]{a^3+b^3}}\right)}{\sqrt[3]{a^3+b^3}} - \frac{\log(a^3+b^3x^3)}{3\sqrt[3]{a^3+b^3}} + \frac{\log\left(-\frac{x\sqrt[3]{a^3+b^3}}{a} - \sqrt[3]{1-x^3}\right)}{2\sqrt[3]{a^3+b^3}} + \frac{\log\left(\sqrt[3]{a^3+b^3}\right)}{2\sqrt[3]{a^3+b^3}} \right)}{b^2} + \frac{(1-x^3)^{2/3}}{2b}$$

input

```
Int[(1 - x^3)^(2/3)/(a + b*x),x]
```

output

```
(1 - x^3)^(2/3)/(2*b) + (a*x^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3])/(2*b^2) + (-((a^2*(-ArcTan[(1 - (2*x)/(1 - x^3)^(1/3)]/Sqrt[3])/Sqrt[3]] + Log[x + (1 - x^3)^(1/3)]/2))/b) + ((a^3 + b^3)*(-1/2*(b*x^2*AppellF1[2/3, 1/3, 1, 5/3, x^3, -(b^3*x^3)/a^3]))/a^2 - ArcTan[(1 - (2*(a^3 + b^3)^(1/3)*x)/(a*(1 - x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*(a^3 + b^3)^(1/3)) + ArcTan[(1 + (2*b*(1 - x^3)^(1/3))/(a^3 + b^3)^(1/3))/Sqrt[3]]/(Sqrt[3]*(a^3 + b^3)^(1/3)) - Log[a^3 + b^3*x^3]/(3*(a^3 + b^3)^(1/3)) + Log[-((a^3 + b^3)^(1/3)*x)/a] - (1 - x^3)^(1/3)]/(2*(a^3 + b^3)^(1/3)) + Log[(a^3 + b^3)^(1/3) - b*(1 - x^3)^(1/3)]/(2*(a^3 + b^3)^(1/3)))/b/b^2
```

Defintions of rubi rules used

rule 769

```
Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]
```

rule 888

```
Int[((c_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])
```

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2577 `Int[((e_.) + (f_.)*(x_))/(((c_.) + (d_.)*(x_))*((a_) + (b_.)*(x_)^3)^(1/3)), x_Symbol] := Simp[f/d Int[1/(a + b*x^3)^(1/3), x], x] + Simp[(d*e - c*f)/d Int[1/((c + d*x)*(a + b*x^3)^(1/3)), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`

rule 2578 `Int[((a_) + (b_.)*(x_)^3)^(2/3)/((c_) + (d_.)*(x_)), x_Symbol] := Simp[(a + b*x^3)^(2/3)/(2*d), x] + (Simp[1/d^2 Int[(a*d^2 + b*c^2*x)/((c + d*x)*(a + b*x^3)^(1/3)), x], x] - Simp[b*(c/d^2 Int[x/(a + b*x^3)^(1/3), x], x) /; FreeQ[{a, b, c, d}, x]`

rule 2581 `Int[(Px_.)*((c_) + (d_.)*(x_))^(q_)*((a_) + (b_.)*(x_)^3)^(p_), x_Symbol] := Int[ExpandIntegrand[(c^3 + d^3*x^3)^q*(a + b*x^3)^p, Px/(c^2 - c*d*x + d^2*x^2)^q, x], x] /; FreeQ[{a, b, c, d, p}, x] && PolyQ[Px, x] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [F]

$$\int \frac{(-x^3 + 1)^{\frac{2}{3}}}{bx + a} dx$$

input `int((-x^3+1)^(2/3)/(b*x+a),x)`

output `int((-x^3+1)^(2/3)/(b*x+a),x)`

Fricas [F(-1)]

Timed out.

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \text{Timed out}$$

input `integrate((-x^3+1)^(2/3)/(b*x+a),x, algorithm="fricas")`

output Timed out

Sympy [F]

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \int \frac{-(x-1)(x^2+x+1)^{2/3}}{a+bx} dx$$

input `integrate((-x**3+1)**(2/3)/(b*x+a),x)`

output `Integral((-x - 1)*(x**2 + x + 1)**(2/3)/(a + b*x), x)`

Maxima [F]

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \int \frac{(-x^3+1)^{2/3}}{bx+a} dx$$

input `integrate((-x^3+1)^(2/3)/(b*x+a),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(2/3)/(b*x + a), x)`

Giac [F]

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \int \frac{(-x^3+1)^{2/3}}{bx+a} dx$$

input `integrate((-x^3+1)^(2/3)/(b*x+a),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3)/(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \int \frac{(1-x^3)^{2/3}}{a+bx} dx$$

input `int((1 - x^3)^(2/3)/(a + b*x),x)`

output `int((1 - x^3)^(2/3)/(a + b*x), x)`

Reduce [F]

$$\int \frac{(1-x^3)^{2/3}}{a+bx} dx = \frac{(-x^3+1)^{2/3} - 2\left(\int \frac{(-x^3+1)^{2/3}}{bx^4+ax^3-bx-a} dx\right)b - 2\left(\int \frac{(-x^3+1)^{2/3}x^2}{bx^4+ax^3-bx-a} dx\right)a}{2b}$$

input `int((-x^3+1)^(2/3)/(b*x+a),x)`

output `((- x**3 + 1)**(2/3) - 2*int((- x**3 + 1)**(2/3)/(a*x**3 - a + b*x**4 - b*x),x)*b - 2*int(((- x**3 + 1)**(2/3)*x**2)/(a*x**3 - a + b*x**4 - b*x), x)*a)/(2*b)`

3.109 $\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$

Optimal result	812
Mathematica [F]	813
Rubi [A] (verified)	813
Maple [F]	814
Fricas [F]	815
Sympy [F]	815
Maxima [F]	815
Giac [F]	816
Mupad [F(-1)]	816
Reduce [F]	816

Optimal result

Integrand size = 22, antiderivative size = 234

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = -\frac{(1-x^3)^{2/3}}{3(1+x^3)} + \frac{x(1-x^3)^{2/3}}{3(1+x^3)} + \frac{2x^2(1-x^3)^{2/3}}{3(1+x^3)}$$

$$-\frac{2^{2/3} \arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2x}}{\sqrt[3]{1-x^3}}\right)}{3\sqrt{3}} - \frac{2^{2/3} \arctan\left(\frac{1+2^{2/3}\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{3\sqrt{3}}$$

$$+ \frac{1}{3}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) - \frac{\log\left(\sqrt[3]{2} - \sqrt[3]{1-x^3}\right)}{3\sqrt[3]{2}} + \frac{\log\left(-\sqrt[3]{2}x - \sqrt[3]{1-x^3}\right)}{3\sqrt[3]{2}}$$

output

```
-1/3*(-x^3+1)^(2/3)/(x^3+1)+1/3*x*(-x^3+1)^(2/3)/(x^3+1)+2/3*x^2*(-x^3+1)^(2/3)/(x^3+1)+1/3*x^2*hypergeom([1/3, 2/3], [5/3], x^3)-1/6*ln(2^(1/3)-(-x^3+1)^(1/3))*2^(2/3)+1/6*ln(-2^(1/3)*x-(-x^3+1)^(1/3))*2^(2/3)-1/9*arctan(1/3*(1-2*2^(1/3)*x/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)-1/9*arctan(1/3*(1+2^(2/3)*(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)
```

Mathematica [F]

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$$

input `Integrate[(1 - x^3)^(2/3)/(1 - x + x^2)^2, x]`

output `Integrate[(1 - x^3)^(2/3)/(1 - x + x^2)^2, x]`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(1-x^3)^{2/3}}{(x^2-x+1)^2} dx \\ & \quad \downarrow \text{2583} \\ & \int \left(\frac{2(1-x^3)^{2/3} x}{(x^3+1)^2} + \frac{(1-x^3)^{2/3}}{(x^3+1)^2} + \frac{(1-x^3)^{2/3} x^2}{(x^3+1)^2} \right) dx \\ & \quad \downarrow \text{2009} \\ & \frac{2^{2/3} \arctan\left(\frac{1-\frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{2^{2/3} \arctan\left(\frac{2^{2/3}\sqrt[3]{1-x^3+1}}{\sqrt{3}}\right)}{3\sqrt{3}} + \\ & \frac{1}{3}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) + \frac{(1-x^3)^{2/3} x}{3(x^3+1)} - \frac{(1-x^3)^{2/3}}{3(x^3+1)} - \\ & \frac{\log\left(\sqrt[3]{2}-\sqrt[3]{1-x^3}\right)}{3\sqrt[3]{2}} + \frac{\log\left(-\sqrt[3]{1-x^3}-\sqrt[3]{2}x\right)}{3\sqrt[3]{2}} + \frac{2(1-x^3)^{2/3} x^2}{3(x^3+1)} \end{aligned}$$

input `Int[(1 - x^3)^(2/3)/(1 - x + x^2)^2,x]`

output `-1/3*(1 - x^3)^(2/3)/(1 + x^3) + (x*(1 - x^3)^(2/3))/(3*(1 + x^3)) + (2*x^2*(1 - x^3)^(2/3))/(3*(1 + x^3)) - (2^(2/3)*ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]])/(3*Sqrt[3]) - (2^(2/3)*ArcTan[(1 + 2^(2/3)*(1 - x^3)^(1/3))/Sqrt[3]])/(3*Sqrt[3]) + (x^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3])/3 - Log[2^(1/3) - (1 - x^3)^(1/3)]/(3*2^(1/3)) + Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)]/(3*2^(1/3))`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2583 `Int[(Px_)*((c_) + (d_)*(x_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^3)^(p_), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [F]

$$\int \frac{(-x^3 + 1)^{\frac{2}{3}}}{(x^2 - x + 1)^2} dx$$

input `int((-x^3+1)^(2/3)/(x^2-x+1)^2,x)`

output `int((-x^3+1)^(2/3)/(x^2-x+1)^2,x)`

Fricas [F]

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(-x^3+1)^{2/3}}{(x^2-x+1)^2} dx$$

input `integrate((-x^3+1)^(2/3)/(x^2-x+1)^2,x, algorithm="fricas")`

output `integral((-x^3 + 1)^(2/3)/(x^4 - 2*x^3 + 3*x^2 - 2*x + 1), x)`

Sympy [F]

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(-(x-1)(x^2+x+1))^{2/3}}{(x^2-x+1)^2} dx$$

input `integrate((-x**3+1)**(2/3)/(x**2-x+1)**2,x)`

output `Integral(-(x - 1)*(x**2 + x + 1)**(2/3)/(x**2 - x + 1)**2, x)`

Maxima [F]

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(-x^3+1)^{2/3}}{(x^2-x+1)^2} dx$$

input `integrate((-x^3+1)^(2/3)/(x^2-x+1)^2,x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(2/3)/(x^2 - x + 1)^2, x)`

Giac [F]

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(-x^3+1)^{\frac{2}{3}}}{(x^2-x+1)^2} dx$$

input `integrate((-x^3+1)^(2/3)/(x^2-x+1)^2,x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3)/(x^2 - x + 1)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(1-x^3)^{2/3}}{(x^2-x+1)^2} dx$$

input `int((1 - x^3)^(2/3)/(x^2 - x + 1)^2,x)`

output `int((1 - x^3)^(2/3)/(x^2 - x + 1)^2, x)`

Reduce [F]

$$\int \frac{(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(-x^3+1)^{\frac{2}{3}}}{x^4-2x^3+3x^2-2x+1} dx$$

input `int((-x^3+1)^(2/3)/(x^2-x+1)^2,x)`

output `int((- x**3 + 1)**(2/3)/(x**4 - 2*x**3 + 3*x**2 - 2*x + 1),x)`

3.110
$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$$

Optimal result	817
Mathematica [F]	818
Rubi [A] (verified)	818
Maple [C] (warning: unable to verify)	819
Fricas [B] (verification not implemented)	820
Sympy [F]	821
Maxima [F]	822
Giac [F]	822
Mupad [F(-1)]	822
Reduce [F]	823

Optimal result

Integrand size = 27, antiderivative size = 199

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \frac{(1-x^3)^{2/3}}{1-x+x^2} - \frac{2 \arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}}$$

$$+ \frac{2^{2/3} \arctan\left(\frac{1-\frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{2^{2/3} \arctan\left(\frac{1+2^{2/3}\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}}$$

$$+ \frac{\log\left(\sqrt[3]{2}-\sqrt[3]{1-x^3}\right)}{\sqrt[3]{2}} - \frac{\log\left(-\sqrt[3]{2}x-\sqrt[3]{1-x^3}\right)}{\sqrt[3]{2}} + \log\left(x+\sqrt[3]{1-x^3}\right)$$

output

```
(-x^3+1)^(2/3)/(x^2-x+1)+1/2*ln(2^(1/3)-(-x^3+1)^(1/3))-1/2*ln(-2^(1/3)*x-(-x^3+1)^(1/3))*2^(2/3)+ln(x+(-x^3+1)^(1/3))-2/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)+1/3*arctan(1/3*(1-2*2^(1/3)*x/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)+1/3*arctan(1/3*(1+2^(2/3)*(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)
```

Mathematica [F]

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx$$

input `Integrate[((1 - 2*x)*(1 - x^3)^(2/3))/(1 - x + x^2)^2,x]`

output `Integrate[((1 - 2*x)*(1 - x^3)^(2/3))/(1 - x + x^2)^2, x]`

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 250, normalized size of antiderivative = 1.26, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {2583, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(x^2-x+1)^2} dx$$

$$\downarrow 2583$$

$$\int \left(-\frac{2(1-x^3)^{2/3}x^3}{(x^3+1)^2} + \frac{(1-x^3)^{2/3}}{(x^3+1)^2} - \frac{3(1-x^3)^{2/3}x^2}{(x^3+1)^2} \right) dx$$

$$\downarrow 2009$$

$$-\frac{2 \arctan \left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{2^{2/3} \arctan \left(\frac{1 - \frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{2^{2/3} \arctan \left(\frac{2^{2/3} \sqrt[3]{1-x^3+1}}{\sqrt{3}} \right)}{\sqrt{3}} +$$

$$\frac{(1-x^3)^{2/3}x}{x^3+1} + \frac{(1-x^3)^{2/3}}{x^3+1} + \frac{\log \left(\sqrt[3]{2} - \sqrt[3]{1-x^3} \right)}{\sqrt[3]{2}} - \frac{2}{3} 2^{2/3} \log \left(-\sqrt[3]{1-x^3} - \sqrt[3]{2}x \right) +$$

$$\frac{\log \left(-\sqrt[3]{1-x^3} - \sqrt[3]{2}x \right)}{3\sqrt[3]{2}} + \log \left(\sqrt[3]{1-x^3} + x \right)$$

input `Int[((1 - 2*x)*(1 - x^3)^(2/3))/(1 - x + x^2)^2,x]`

output `(1 - x^3)^(2/3)/(1 + x^3) + (x*(1 - x^3)^(2/3))/(1 + x^3) - (2*ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]])/Sqrt[3] + (2^(2/3)*ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]])/Sqrt[3] + (2^(2/3)*ArcTan[(1 + 2^(2/3)*(1 - x^3)^(1/3))/Sqrt[3]])/Sqrt[3] + Log[2^(1/3) - (1 - x^3)^(1/3)]/2^(1/3) + Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)]/(3*2^(1/3)) - (2*2^(2/3)*Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)])/3 + Log[x + (1 - x^3)^(1/3)]`

Defintions of rubi rules used

rule 2009 `Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]`

rule 2583 `Int[(Px_.)*((c_) + (d_.)*(x_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*(x_)^3)^(p_.), x_Symbol] := Simp[1/c^q Int[ExpandIntegrand[(c^3 - d^3*x^3)^q*(a + b*x^3)^p, Px/(c - d*x)^q, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && RationalQ[p] && EqQ[Denominator[p], 3]`

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 24.92 (sec) , antiderivative size = 842, normalized size of antiderivative = 4.23

Expression too large to display

input `int((1-2*x)*(-x^3+1)^(2/3)/(x^2-x+1)^2,x)`

output

```

-(x^3-1)/(x^2-x+1)/(-x^3+1)^(1/3)-2/3*RootOf(_Z^2-_Z+1)*ln(RootOf(_Z^2-_Z+1)^2*x^3-3*RootOf(_Z^2-_Z+1)*(-x^3+1)^(2/3)*x+3*RootOf(_Z^2-_Z+1)*(-x^3+1)^(1/3)*x^2-4*RootOf(_Z^2-_Z+1)*x^3+3*x*(-x^3+1)^(2/3)-3*x^2*(-x^3+1)^(1/3)+4*x^3+RootOf(_Z^2-_Z+1)-2)+2/3*ln(RootOf(_Z^2-_Z+1)^2*x^3+3*RootOf(_Z^2-_Z+1)*(-x^3+1)^(2/3)*x-3*RootOf(_Z^2-_Z+1)*(-x^3+1)^(1/3)*x^2+2*RootOf(_Z^2-_Z+1)*x^3+x^3-RootOf(_Z^2-_Z+1)-1)*RootOf(_Z^2-_Z+1)-2/3*ln(RootOf(_Z^2-_Z+1)^2*x^3+3*RootOf(_Z^2-_Z+1)*(-x^3+1)^(2/3)*x-3*RootOf(_Z^2-_Z+1)*(-x^3+1)^(1/3)*x^2+2*RootOf(_Z^2-_Z+1)*x^3+x^3-RootOf(_Z^2-_Z+1)-1)-1/9*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*ln(-((-x^3+1)^(1/3)*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))^2*RootOf(_Z^2-_Z+1)*x-(-x^3+1)^(1/3)*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))^2*x+3*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*RootOf(_Z^2-_Z+1)*x^2+3*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*RootOf(_Z^2-_Z+1)*x-6*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*x^2+54*(-x^3+1)^(2/3)-3*RootOf(_Z^2-_Z+1)*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))-6*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*x+6*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1)))/((x^2-x+1)^2)+1/9*RootOf(_Z^2-_Z+1)*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*ln((6*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*RootOf(_Z^2-_Z+1)*x^2-(-x^3+1)^(1/3)*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))^2-6*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*RootOf(_Z^2-_Z+1)*x-3*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1))*x^2+54*(-x^3+1)^(2/3)-6*RootOf(_Z^2-_Z+1)*RootOf(_Z^3-324+648*RootOf(_Z^2-_Z+1)))+3*Root...

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2155 vs. $2(163) = 326$.

Time = 1.68 (sec) , antiderivative size = 2155, normalized size of antiderivative = 10.83

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \text{Too large to display}$$

input

```
integrate((1-2*x)*(-x^3+1)^(2/3)/(x^2-x+1)^2,x, algorithm="fricas")
```

output

```

-1/12*(8*sqrt(3)*(x^2 - x + 1)*arctan((4*sqrt(3)*(-x^3 + 1)^(1/3)*x^2 + 2*
sqrt(3)*(-x^3 + 1)^(2/3)*x - sqrt(3)*(x^3 - 1))/(9*x^3 - 1)) + (-16/27)^(1
/6)*(x^2 + sqrt(-3)*(x^2 - x + 1) - x + 1)*log(-(4*2^(2/3)*(15*x^6 - 200*x
^5 + 5*x^4 + 216*x^3 + 157*x^2 + sqrt(-3)*(15*x^6 - 200*x^5 + 5*x^4 + 216*
x^3 + 157*x^2 - 124*x - 42) - 124*x - 42) - 16*(31*x^4 + 107*x^3 - 243*x^2
+ 3*sqrt(-1/3)*(23*x^4 - 85*x^3 - 57*x^2 + 104*x - 4) - 26*x + 50)*(-x^3
+ 1)^(2/3) + (-x^3 + 1)^(1/3)*(27*(-16/27)^(5/6)*(4*x^5 + 69*x^4 - 58*x^3
- 77*x^2 - sqrt(-3)*(4*x^5 + 69*x^4 - 58*x^3 - 77*x^2 + 12*x + 23) + 12*x
+ 23) - 8*(-2)^(1/3)*(50*x^5 - 93*x^4 - 88*x^3 - 7*x^2 - sqrt(-3)*(50*x^5
- 93*x^4 - 88*x^3 - 7*x^2 + 150*x - 31) + 150*x - 31)) + 4*(-16/27)^(1/6)*
(131*x^6 - 48*x^5 - 381*x^4 - 152*x^3 + 267*x^2 + sqrt(-3)*(131*x^6 - 48*x
^5 - 381*x^4 - 152*x^3 + 267*x^2 + 276*x - 112) + 276*x - 112))/(x^6 - 3*x
^5 + 6*x^4 - 7*x^3 + 6*x^2 - 3*x + 1)) - (-16/27)^(1/6)*(x^2 + sqrt(-3)*(x
^2 - x + 1) - x + 1)*log(-(4*2^(2/3)*(15*x^6 - 200*x^5 + 5*x^4 + 216*x^3 +
157*x^2 + sqrt(-3)*(15*x^6 - 200*x^5 + 5*x^4 + 216*x^3 + 157*x^2 - 124*x
- 42) - 124*x - 42) - 16*(31*x^4 + 107*x^3 - 243*x^2 - 3*sqrt(-1/3)*(23*x^
4 - 85*x^3 - 57*x^2 + 104*x - 4) - 26*x + 50)*(-x^3 + 1)^(2/3) - (-x^3 + 1
)^(1/3)*(27*(-16/27)^(5/6)*(4*x^5 + 69*x^4 - 58*x^3 - 77*x^2 - sqrt(-3)*(4
*x^5 + 69*x^4 - 58*x^3 - 77*x^2 + 12*x + 23) + 12*x + 23) + 8*(-2)^(1/3)*(
50*x^5 - 93*x^4 - 88*x^3 - 7*x^2 - sqrt(-3)*(50*x^5 - 93*x^4 - 88*x^3 - ...

```

Sympy [F]

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx =$$

$$-\int \left(\frac{(1-x^3)^{2/3}}{x^4-2x^3+3x^2-2x+1} \right) dx - \int \frac{2x(1-x^3)^{2/3}}{x^4-2x^3+3x^2-2x+1} dx$$

input

```
integrate((1-2*x)*(-x**3+1)**(2/3)/(x**2-x+1)**2,x)
```

output

```
-Integral(-(1 - x**3)**(2/3)/(x**4 - 2*x**3 + 3*x**2 - 2*x + 1), x) - Inte
gral(2*x*(1 - x**3)**(2/3)/(x**4 - 2*x**3 + 3*x**2 - 2*x + 1), x)
```

Maxima [F]

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int -\frac{(-x^3+1)^{2/3}(2x-1)}{(x^2-x+1)^2} dx$$

input `integrate((1-2*x)*(-x^3+1)^(2/3)/(x^2-x+1)^2,x, algorithm="maxima")`

output `-integrate((-x^3 + 1)^(2/3)*(2*x - 1)/(x^2 - x + 1)^2, x)`

Giac [F]

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \int -\frac{(-x^3+1)^{2/3}(2x-1)}{(x^2-x+1)^2} dx$$

input `integrate((1-2*x)*(-x^3+1)^(2/3)/(x^2-x+1)^2,x, algorithm="giac")`

output `integrate(-(-x^3 + 1)^(2/3)*(2*x - 1)/(x^2 - x + 1)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = -\int \frac{(2x-1)(1-x^3)^{2/3}}{(x^2-x+1)^2} dx$$

input `int(-((2*x - 1)*(1 - x^3)^(2/3))/(x^2 - x + 1)^2,x)`

output `-int(((2*x - 1)*(1 - x^3)^(2/3))/(x^2 - x + 1)^2, x)`

Reduce [F]

$$\int \frac{(1-2x)(1-x^3)^{2/3}}{(1-x+x^2)^2} dx = \frac{-(-x^3+1)^{2/3} - 2\left(\int \frac{(-x^3+1)^{2/3}}{x^7-2x^6+3x^5-3x^4+3x^3-3x^2+2x-1} dx\right) x^2 + 2\left(\int \frac{(-x^3}{x^7-2x^6+3x^5-3x^4+3x^3-3x^2+2x-1} dx\right)}{1-x+x^2}$$

input `int((1-2*x)*(-x^3+1)^(2/3)/(x^2-x+1)^2,x)`

output

```
( - ( - x**3 + 1)**(2/3) - 2*int(( - x**3 + 1)**(2/3)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x**2 + 2*int(( - x**3 + 1)**(2/3)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x - 2*int(( - x**3 + 1)**(2/3)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x) - 2*int((( - x**3 + 1)**(2/3)*x**4)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x**2 + 2*int((( - x**3 + 1)**(2/3)*x**4)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x - 2*int((( - x**3 + 1)**(2/3)*x**4)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x) + 2*int((( - x**3 + 1)**(2/3)*x**2)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x**2 - 2*int((( - x**3 + 1)**(2/3)*x**2)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x + 2*int((( - x**3 + 1)**(2/3)*x**2)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x) + 4*int((( - x**3 + 1)**(2/3)*x)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x**2 - 4*int((( - x**3 + 1)**(2/3)*x)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x)*x + 4*int((( - x**3 + 1)**(2/3)*x)/(x**7 - 2*x**6 + 3*x**5 - 3*x**4 + 3*x**3 - 3*x**2 + 2*x - 1),x))/(x**2 - x + 1)
```


3.111 $\int \frac{(1-x^3)^{2/3}}{1+x} dx$

Optimal result	824
Mathematica [F]	825
Rubi [A] (verified)	825
Maple [F]	827
Fricas [F]	828
Sympy [F]	828
Maxima [F]	828
Giac [F]	829
Mupad [F(-1)]	829
Reduce [F]	829

Optimal result

Integrand size = 17, antiderivative size = 177

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \frac{1}{2}(1-x^3)^{2/3} - \frac{\sqrt{3} \arctan\left(\frac{1+\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{\sqrt[3]{2}}$$

$$+ \frac{\arctan\left(\frac{1-\sqrt[3]{1-x^3}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)$$

$$- \frac{\log((1-x)(1+x)^2)}{2\sqrt[3]{2}} - \frac{1}{2} \log\left(x + \sqrt[3]{1-x^3}\right) + \frac{3 \log\left(-1+x+2^{2/3}\sqrt[3]{1-x^3}\right)}{2\sqrt[3]{2}}$$

output

```
1/2*(-x^3+1)^(2/3)+1/2*x^2*hypergeom([1/3, 2/3], [5/3], x^3)-1/4*ln((1-x)*(1+x)^2)*2^(2/3)-1/2*ln(x+(-x^3+1)^(1/3))+3/4*ln(-1+x+2^(2/3)*(-x^3+1)^(1/3))*2^(2/3)+1/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)-1/2*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)*2^(2/3)
```

Mathematica [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \int \frac{(1-x^3)^{2/3}}{1+x} dx$$

input `Integrate[(1 - x^3)^(2/3)/(1 + x), x]`

output `Integrate[(1 - x^3)^(2/3)/(1 + x), x]`

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.03, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2578, 888, 2577, 769, 2574}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(1-x^3)^{2/3}}{x+1} dx \\ & \quad \downarrow \text{2578} \\ & \int \frac{x}{\sqrt[3]{1-x^3}} dx + \int \frac{1-x}{(x+1)\sqrt[3]{1-x^3}} dx + \frac{1}{2}(1-x^3)^{2/3} \\ & \quad \downarrow \text{888} \\ & \int \frac{1-x}{(x+1)\sqrt[3]{1-x^3}} dx + \frac{1}{2}x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) + \frac{1}{2}(1-x^3)^{2/3} \\ & \quad \downarrow \text{2577} \\ & - \int \frac{1}{\sqrt[3]{1-x^3}} dx + 2 \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx + \frac{1}{2}x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) + \\ & \quad \frac{1}{2}(1-x^3)^{2/3} \\ & \quad \downarrow \text{769} \end{aligned}$$

$$\begin{aligned}
& 2 \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx + \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) + \\
& \frac{1}{2}(1-x^3)^{2/3} - \frac{1}{2}\log\left(\sqrt[3]{1-x^3} + x\right) \\
& \quad \downarrow \text{2574} \\
& \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \\
& 2 \left(\frac{\sqrt{3} \arctan\left(\frac{\frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{2\sqrt[3]{2}} + \frac{3 \log\left(2^{2/3}\sqrt[3]{1-x^3} + x - 1\right)}{4\sqrt[3]{2}} - \frac{\log\left((1-x)(x+1)^2\right)}{4\sqrt[3]{2}} \right) + \\
& \frac{1}{2}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) + \frac{1}{2}(1-x^3)^{2/3} - \frac{1}{2}\log\left(\sqrt[3]{1-x^3} + x\right)
\end{aligned}$$

input `Int[(1 - x^3)^(2/3)/(1 + x), x]`

output `(1 - x^3)^(2/3)/2 + ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + (x^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3])/2 - Log[x + (1 - x^3)^(1/3)]/2 + 2*(-1/2*(Sqrt[3]*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]])/2^(1/3) - Log[(1 - x)*(1 + x)^2]/(4*2^(1/3)) + (3*Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3)]/(4*2^(1/3)))`

Defintions of rubi rules used

rule 769

`Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

rule 888 `Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

rule 2574 `Int[1/(((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^3)^(1/3)), x_Symbol] := Simp[Sqrt[3]*(ArcTan[(1 - 2^(1/3)*Rt[b, 3]*((c - d*x)/(d*(a + b*x^3)^(1/3)))))/Sqrt[3]]/(2^(4/3)*Rt[b, 3]*c), x] + (Simp[Log[(c + d*x)^2*(c - d*x)]/(2^(7/3)*Rt[b, 3]*c), x] - Simp[(3*Log[Rt[b, 3]*(c - d*x) + 2^(2/3)*d*(a + b*x^3)^(1/3)])/(2^(7/3)*Rt[b, 3]*c), x]) /; FreeQ[{a, b, c, d}, x] && EqQ[b*c^3 + a*d^3, 0]`

rule 2577 `Int[((e_) + (f_)*(x_))/(((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^3)^(1/3)), x_Symbol] := Simp[f/d Int[1/(a + b*x^3)^(1/3), x], x] + Simp[(d*e - c*f)/d Int[1/((c + d*x)*(a + b*x^3)^(1/3)), x], x] /; FreeQ[{a, b, c, d, e, f}, x]`

rule 2578 `Int[((a_) + (b_)*(x_)^3)^(2/3)/((c_) + (d_)*(x_)), x_Symbol] := Simp[(a + b*x^3)^(2/3)/(2*d), x] + (Simp[1/d^2 Int[(a*d^2 + b*c^2*x)/((c + d*x)*(a + b*x^3)^(1/3)), x], x] - Simp[b*(c/d^2) Int[x/(a + b*x^3)^(1/3), x], x]) /; FreeQ[{a, b, c, d}, x]`

Maple [F]

$$\int \frac{(-x^3 + 1)^{\frac{2}{3}}}{1 + x} dx$$

input `int((-x^3+1)^(2/3)/(1+x),x)`

output `int((-x^3+1)^(2/3)/(1+x),x)`

Fricas [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \int \frac{(-x^3+1)^{2/3}}{x+1} dx$$

input `integrate((-x^3+1)^(2/3)/(1+x),x, algorithm="fricas")`

output `integral((-x^3 + 1)^(2/3)/(x + 1), x)`

Sympy [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \int \frac{-(x-1)(x^2+x+1)^{2/3}}{x+1} dx$$

input `integrate((-x**3+1)**(2/3)/(1+x),x)`

output `Integral((-x - 1)*(x**2 + x + 1)**(2/3)/(x + 1), x)`

Maxima [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \int \frac{(-x^3+1)^{2/3}}{x+1} dx$$

input `integrate((-x^3+1)^(2/3)/(1+x),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(2/3)/(x + 1), x)`

Giac [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \int \frac{(-x^3+1)^{2/3}}{x+1} dx$$

input `integrate((-x^3+1)^(2/3)/(1+x),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3)/(x + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \int \frac{(1-x^3)^{2/3}}{x+1} dx$$

input `int((1 - x^3)^(2/3)/(x + 1),x)`

output `int((1 - x^3)^(2/3)/(x + 1), x)`

Reduce [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x} dx = \frac{(-x^3+1)^{2/3}}{2} - \left(\int \frac{(-x^3+1)^{2/3}}{x^4+x^3-x-1} dx \right) - \left(\int \frac{(-x^3+1)^{2/3} x^2}{x^4+x^3-x-1} dx \right)$$

input `int((-x^3+1)^(2/3)/(1+x),x)`

output `((- x**3 + 1)**(2/3) - 2*int((- x**3 + 1)**(2/3)/(x**4 + x**3 - x - 1),x) - 2*int(((- x**3 + 1)**(2/3)*x**2)/(x**4 + x**3 - x - 1),x))/2`

3.112
$$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx$$

Optimal result	830
Mathematica [F]	831
Rubi [A] (verified)	831
Maple [F]	834
Fricas [F]	834
Sympy [F]	834
Maxima [F]	835
Giac [F]	835
Mupad [F(-1)]	835
Reduce [F]	836

Optimal result

Integrand size = 27, antiderivative size = 177

$$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx = \frac{1}{2}(1-x^3)^{2/3} - \frac{\sqrt{3} \arctan\left(\frac{1+\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt[3]{2}}$$

$$+ \frac{\arctan\left(\frac{1-\frac{\sqrt[3]{2x}}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{1}{2}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right)$$

$$- \frac{\log((1-x)(1+x)^2)}{2\sqrt[3]{2}} - \frac{1}{2} \log\left(x + \sqrt[3]{1-x^3}\right) + \frac{3 \log\left(-1+x+2^{2/3}\sqrt[3]{1-x^3}\right)}{2\sqrt[3]{2}}$$

output

```
1/2*(-x^3+1)^(2/3)+1/2*x^2*hypergeom([1/3, 2/3], [5/3], x^3)-1/4*ln((1-x)*(1+x)^2)*2^(2/3)-1/2*ln(x+(-x^3+1)^(1/3))+3/4*ln(-1+x+2^(2/3)*(-x^3+1)^(1/3))*2^(2/3)+1/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)-1/2*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)*2^(2/3)
```

Mathematica [F]

$$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx$$

input `Integrate[((1 - x + x^2)*(1 - x^3)^(2/3))/(1 + x^3), x]`

output `Integrate[((1 - x + x^2)*(1 - x^3)^(2/3))/(1 + x^3), x]`

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.03, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2019, 2578, 888, 2577, 769, 2574}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned} & \int \frac{(x^2 - x + 1)(1 - x^3)^{2/3}}{x^3 + 1} dx \\ & \quad \downarrow \text{2019} \\ & \int \frac{(1 - x^3)^{2/3}}{x + 1} dx \\ & \quad \downarrow \text{2578} \\ & \int \frac{x}{\sqrt[3]{1 - x^3}} dx + \int \frac{1 - x}{(x + 1)\sqrt[3]{1 - x^3}} dx + \frac{1}{2}(1 - x^3)^{2/3} \\ & \quad \downarrow \text{888} \\ & \int \frac{1 - x}{(x + 1)\sqrt[3]{1 - x^3}} dx + \frac{1}{2}x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) + \frac{1}{2}(1 - x^3)^{2/3} \\ & \quad \downarrow \text{2577} \end{aligned}$$

$$\begin{aligned}
& - \int \frac{1}{\sqrt[3]{1-x^3}} dx + 2 \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx + \frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) + \\
& \qquad \qquad \qquad \frac{1}{2} (1-x^3)^{2/3} \\
& \qquad \qquad \qquad \downarrow \text{769} \\
& 2 \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx + \frac{\arctan \left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) + \\
& \qquad \qquad \qquad \frac{1}{2} (1-x^3)^{2/3} - \frac{1}{2} \log \left(\sqrt[3]{1-x^3} + x \right) \\
& \qquad \qquad \qquad \downarrow \text{2574} \\
& \qquad \qquad \qquad \frac{\arctan \left(\frac{1 - \frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \\
& 2 \left(- \frac{\sqrt{3} \arctan \left(\frac{\frac{\sqrt[3]{2(1-x)} + 1}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{2\sqrt[3]{2}} + \frac{3 \log \left(2^{2/3} \sqrt[3]{1-x^3} + x - 1 \right)}{4\sqrt[3]{2}} - \frac{\log \left((1-x)(x+1)^2 \right)}{4\sqrt[3]{2}} \right) + \\
& \qquad \qquad \qquad \frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) + \frac{1}{2} (1-x^3)^{2/3} - \frac{1}{2} \log \left(\sqrt[3]{1-x^3} + x \right)
\end{aligned}$$

input `Int[((1 - x + x^2)*(1 - x^3)^(2/3))/(1 + x^3),x]`

output `(1 - x^3)^(2/3)/2 + ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + (x^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3])/2 - Log[x + (1 - x^3)^(1/3)]/2 + 2*(-1/2*(Sqrt[3]*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]])/2^(1/3) - Log[(1 - x)*(1 + x)^2]/(4*2^(1/3)) + (3*Log[-1 + x + 2^(2/3)*(1 - x^3)^(1/3)]/(4*2^(1/3)))`

Definitions of rubi rules used

rule 769 $\text{Int}[(a + b \cdot x^3)^{-1/3}, x_Symbol] \rightarrow \text{Simp}[\text{ArcTan}[(1 + 2 \cdot \text{Rt}[b, 3] \cdot (x/(a + b \cdot x^3)^{1/3}))/\text{Sqrt}[3]]/(\text{Sqrt}[3] \cdot \text{Rt}[b, 3]), x] - \text{Simp}[\text{Log}[(a + b \cdot x^3)^{1/3} - \text{Rt}[b, 3] \cdot x]/(2 \cdot \text{Rt}[b, 3]), x] /; \text{FreeQ}\{a, b\}, x]$

rule 888 $\text{Int}[(c \cdot x)^m \cdot (a + b \cdot x^n)^p, x_Symbol] \rightarrow \text{Simp}[a^p \cdot ((c \cdot x)^{m+1}/(c \cdot (m+1))) \cdot \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b) \cdot (x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])]$

rule 2019 $\text{Int}[u \cdot (Px)^p \cdot (Qx)^q, x_Symbol] \rightarrow \text{Int}[u \cdot \text{PolynomialQuotient}[Px, Qx, x]^p \cdot Qx^{p+q}, x] /; \text{FreeQ}[q, x] \ \&\& \ \text{PolyQ}[Px, x] \ \&\& \ \text{PolyQ}[Qx, x] \ \&\& \ \text{EqQ}[\text{PolynomialRemainder}[Px, Qx, x], 0] \ \&\& \ \text{IntegerQ}[p] \ \&\& \ \text{LtQ}[p \cdot q, 0]$

rule 2574 $\text{Int}[1/((c + d \cdot x) \cdot (a + b \cdot x^3)^{1/3}), x_Symbol] \rightarrow \text{Simp}[\text{Sqrt}[3] \cdot (\text{ArcTan}[(1 - 2^{1/3} \cdot \text{Rt}[b, 3] \cdot ((c - d \cdot x)/(d \cdot (a + b \cdot x^3)^{1/3})))/\text{Sqrt}[3]]/(2^{4/3} \cdot \text{Rt}[b, 3] \cdot c), x] + (\text{Simp}[\text{Log}[(c + d \cdot x)^2 \cdot (c - d \cdot x)]/(2^{7/3} \cdot \text{Rt}[b, 3] \cdot c), x] - \text{Simp}[(3 \cdot \text{Log}[\text{Rt}[b, 3] \cdot (c - d \cdot x) + 2^{2/3} \cdot d \cdot (a + b \cdot x^3)^{1/3}]/(2^{7/3} \cdot \text{Rt}[b, 3] \cdot c), x)] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[b \cdot c^3 + a \cdot d^3, 0]$

rule 2577 $\text{Int}[(e + f \cdot x)/((c + d \cdot x) \cdot (a + b \cdot x^3)^{1/3}), x_Symbol] \rightarrow \text{Simp}[f/d \ \text{Int}[1/(a + b \cdot x^3)^{1/3}, x], x] + \text{Simp}[(d \cdot e - c \cdot f)/d \ \text{Int}[1/((c + d \cdot x) \cdot (a + b \cdot x^3)^{1/3}), x], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x]$

rule 2578 $\text{Int}[(a + b \cdot x^3)^{2/3}/((c + d \cdot x)), x_Symbol] \rightarrow \text{Simp}[(a + b \cdot x^3)^{2/3}/(2 \cdot d), x] + (\text{Simp}[1/d^2 \ \text{Int}[(a \cdot d^2 + b \cdot c^2 \cdot x)/((c + d \cdot x) \cdot (a + b \cdot x^3)^{1/3}), x], x] - \text{Simp}[b \cdot (c/d^2) \ \text{Int}[x/(a + b \cdot x^3)^{1/3}, x], x]) /; \text{FreeQ}\{a, b, c, d\}, x]$

Maple [F]

$$\int \frac{(x^2 - x + 1)(-x^3 + 1)^{\frac{2}{3}}}{x^3 + 1} dx$$

input `int((x^2-x+1)*(-x^3+1)^(2/3)/(x^3+1),x)`

output `int((x^2-x+1)*(-x^3+1)^(2/3)/(x^3+1),x)`

Fricas [F]

$$\int \frac{(1 - x + x^2)(1 - x^3)^{2/3}}{1 + x^3} dx = \int \frac{(-x^3 + 1)^{\frac{2}{3}}(x^2 - x + 1)}{x^3 + 1} dx$$

input `integrate((x^2-x+1)*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="fricas")`

output `integral((-x^3 + 1)^(2/3)/(x + 1), x)`

Sympy [F]

$$\int \frac{(1 - x + x^2)(1 - x^3)^{2/3}}{1 + x^3} dx = \int \frac{(-(x - 1)(x^2 + x + 1))^{\frac{2}{3}}}{x + 1} dx$$

input `integrate((x**2-x+1)*(-x**3+1)**(2/3)/(x**3+1),x)`

output `Integral((-x - 1)*(x**2 + x + 1)**(2/3)/(x + 1), x)`

Maxima [F]

$$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}(x^2-x+1)}{x^3+1} dx$$

input `integrate((x^2-x+1)*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(2/3)*(x^2 - x + 1)/(x^3 + 1), x)`

Giac [F]

$$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}(x^2-x+1)}{x^3+1} dx$$

input `integrate((x^2-x+1)*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3)*(x^2 - x + 1)/(x^3 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(1-x^3)^{2/3}(x^2-x+1)}{x^3+1} dx$$

input `int(((1 - x^3)^(2/3)*(x^2 - x + 1))/(x^3 + 1),x)`

output `int(((1 - x^3)^(2/3)*(x^2 - x + 1))/(x^3 + 1), x)`

Reduce [F]

$$\int \frac{(1-x+x^2)(1-x^3)^{2/3}}{1+x^3} dx = \frac{(-x^3+1)^{2/3}}{2} - \left(\int \frac{(-x^3+1)^{2/3}}{x^4+x^3-x-1} dx \right) - \left(\int \frac{(-x^3+1)^{2/3} x^2}{x^4+x^3-x-1} dx \right)$$

input `int((x^2-x+1)*(-x^3+1)^(2/3)/(x^3+1),x)`

output `((-x**3+1)**(2/3) - 2*int((-x**3+1)**(2/3)/(x**4+x**3-x-1),x) - 2*int((-x**3+1)**(2/3)*x**2/(x**4+x**3-x-1),x))/2`

3.113 $\int \frac{(1-x^3)^{2/3}}{1+x^3} dx$

Optimal result	837
Mathematica [A] (verified)	838
Rubi [A] (verified)	838
Maple [A] (verified)	840
Fricas [A] (verification not implemented)	841
Sympy [F]	841
Maxima [F]	842
Giac [F]	842
Mupad [F(-1)]	842
Reduce [F]	843

Optimal result

Integrand size = 19, antiderivative size = 132

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{2^{2/3} \arctan\left(\frac{1-\frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\log(1+x^3)}{3\sqrt[3]{2}} + \frac{\log\left(-\sqrt[3]{2}x - \sqrt[3]{1-x^3}\right)}{\sqrt[3]{2}} - \frac{1}{2} \log\left(x + \sqrt[3]{1-x^3}\right)$$

output

```
-1/6*ln(x^3+1)*2^(2/3)+1/2*ln(-2^(1/3)*x-(-x^3+1)^(1/3))*2^(2/3)-1/2*ln(x+
(-x^3+1)^(1/3))+1/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)-1/3
*arctan(1/3*(1-2*2^(1/3)*x/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)
```

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.55

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = \frac{1}{6} \left(2\sqrt{3} \arctan \left(\frac{\sqrt{3}x}{x - 2\sqrt[3]{1-x^3}} \right) - 2 \cdot 2^{2/3} \sqrt{3} \arctan \left(\frac{\sqrt{3}x}{x - 2^{2/3} \sqrt[3]{1-x^3}} \right) - 2 \log \left(x + \sqrt[3]{1-x^3} \right) + 2 \cdot 2^{2/3} \log \left(2x + 2^{2/3} \sqrt[3]{1-x^3} \right) + \log \left(x^2 - x \sqrt[3]{1-x^3} \right) \right)$$

input

```
Integrate[(1 - x^3)^(2/3)/(1 + x^3), x]
```

output

```
(2*Sqrt[3]*ArcTan[(Sqrt[3]*x)/(x - 2*(1 - x^3)^(1/3))] - 2*2^(2/3)*Sqrt[3]*ArcTan[(Sqrt[3]*x)/(x - 2^(2/3)*(1 - x^3)^(1/3))] - 2*Log[x + (1 - x^3)^(1/3)] + 2*2^(2/3)*Log[2*x + 2^(2/3)*(1 - x^3)^(1/3)] + Log[x^2 - x*(1 - x^3)^(1/3) + (1 - x^3)^(2/3)] - 2^(2/3)*Log[-2*x^2 + 2^(2/3)*x*(1 - x^3)^(1/3) - 2^(1/3)*(1 - x^3)^(2/3)])/6
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.05, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {916, 769, 901}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{(1-x^3)^{2/3}}{x^3+1} dx$$

↓ 916

$$2 \int \frac{1}{\sqrt[3]{1-x^3}(x^3+1)} dx - \int \frac{1}{\sqrt[3]{1-x^3}} dx$$

↓ 769

$$\begin{aligned}
& 2 \int \frac{1}{\sqrt[3]{1-x^3}(x^3+1)} dx + \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(\sqrt[3]{1-x^3}+x\right) \\
& \quad \downarrow \text{901} \\
& \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \\
& 2 \left(\frac{\arctan\left(\frac{1-\frac{2\sqrt[3]{2}x}{\sqrt[3]{1-x^3}}}{\sqrt{2}\sqrt{3}}\right)}{\sqrt{2}\sqrt{3}} - \frac{\log(x^3+1)}{6\sqrt[3]{2}} + \frac{\log\left(-\sqrt[3]{1-x^3}-\sqrt[3]{2}x\right)}{2\sqrt[3]{2}} \right) - \\
& \quad \frac{1}{2} \log\left(\sqrt[3]{1-x^3}+x\right)
\end{aligned}$$

input `Int[(1 - x^3)^(2/3)/(1 + x^3),x]`

output `ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + 2*(-(ArcTan[(1 - (2*2^(1/3)*x)/(1 - x^3)^(1/3))/Sqrt[3]]/(2^(1/3)*Sqrt[3])) - Log[1 + x^3]/(6*2^(1/3)) + Log[-(2^(1/3)*x) - (1 - x^3)^(1/3)]/(2*2^(1/3))) - Log[x + (1 - x^3)^(1/3)]/2`

Defintions of rubi rules used

rule 769 `Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]`

rule 901 `Int[1/((a_) + (b_.)*(x_)^3)^(1/3)*((c_) + (d_.)*(x_)^3), x_Symbol] := With[{q = Rt[(b*c - a*d)/c, 3]}, Simp[ArcTan[(1 + (2*q*x)/(a + b*x^3)^(1/3))/Sqrt[3]]/(Sqrt[3]*c*q), x] + (-Simp[Log[q*x - (a + b*x^3)^(1/3)]/(2*c*q), x] + Simp[Log[c + d*x^3]/(6*c*q), x])] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]`

rule 916 `Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[b/d Int[(a + b*x^n)^(p - 1), x], x] - Simp[(b*c - a*d)/d Int[(a + b*x^n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]`

Maple [A] (verified)

Time = 1.07 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.36

method	result
pseudoelliptic	$\frac{2^{\frac{2}{3}} \ln\left(\frac{2^{\frac{1}{3}}x + (-x^3+1)^{\frac{1}{3}}}{x}\right)}{3} - \frac{2^{\frac{2}{3}} \ln\left(\frac{2^{\frac{2}{3}}x^2 - 2^{\frac{1}{3}}(-x^3+1)^{\frac{1}{3}}x + (-x^3+1)^{\frac{2}{3}}}{x^2}\right)}{6} + \frac{\sqrt{3}2^{\frac{2}{3}} \arctan\left(\frac{\sqrt{3}\left(-2^{\frac{2}{3}}(-x^3+1)^{\frac{1}{3}}+x\right)}{3x}\right)}{3}$

input `int((-x^3+1)^(2/3)/(x^3+1),x,method=_RETURNVERBOSE)`

output `1/3*2^(2/3)*ln((2^(1/3)*x+(-x^3+1)^(1/3))/x)-1/6*2^(2/3)*ln((2^(2/3)*x^2-2^(1/3)*(-x^3+1)^(1/3)*x+(-x^3+1)^(2/3))/x^2)+1/3*3^(1/2)*2^(2/3)*arctan(1/3*3^(1/2)*(-2^(2/3)*(-x^3+1)^(1/3)+x)/x)+1/6*ln(((x^3+1)^(2/3)-x*(-x^3+1)^(1/3)+x^2)/x^2)-1/3*3^(1/2)*arctan(1/3*(-2*(-x^3+1)^(1/3)+x)*3^(1/2)/x)-1/3*ln((x+(-x^3+1)^(1/3))/x)`

Fricas [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.45

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = -\frac{1}{3} \cdot 4^{1/3} \sqrt{3} \arctan\left(-\frac{\sqrt{3}x - 4^{1/3}\sqrt{3}(-x^3+1)^{1/3}}{3x}\right) + \frac{1}{3} \sqrt{3} \arctan\left(-\frac{\sqrt{3}x - 2\sqrt{3}(-x^3+1)^{1/3}}{3x}\right) + \frac{1}{3} \cdot 4^{1/3} \log\left(\frac{4^{2/3}x + 2(-x^3+1)^{1/3}}{x}\right) - \frac{1}{6} \cdot 4^{1/3} \log\left(\frac{2 \cdot 4^{1/3}x^2 - 4^{2/3}(-x^3+1)^{1/3}x + 2(-x^3+1)^{2/3}}{x^2}\right) - \frac{1}{3} \log\left(\frac{x + (-x^3+1)^{1/3}}{x}\right) + \frac{1}{6} \log\left(\frac{x^2 - (-x^3+1)^{1/3}x + (-x^3+1)^{2/3}}{x^2}\right)$$

input `integrate((-x^3+1)^(2/3)/(x^3+1),x, algorithm="fricas")`output `-1/3*4^(1/3)*sqrt(3)*arctan(-1/3*(sqrt(3)*x - 4^(1/3)*sqrt(3)*(-x^3 + 1)^(1/3))/x) + 1/3*sqrt(3)*arctan(-1/3*(sqrt(3)*x - 2*sqrt(3)*(-x^3 + 1)^(1/3))/x) + 1/3*4^(1/3)*log((4^(2/3)*x + 2*(-x^3 + 1)^(1/3))/x) - 1/6*4^(1/3)*log((2*4^(1/3)*x^2 - 4^(2/3)*(-x^3 + 1)^(1/3)*x + 2*(-x^3 + 1)^(2/3))/x^2) - 1/3*log((x + (-x^3 + 1)^(1/3))/x) + 1/6*log((x^2 - (-x^3 + 1)^(1/3)*x + (-x^3 + 1)^(2/3))/x^2)`**Sympy [F]**

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{-(x-1)(x^2+x+1)^{2/3}}{(x+1)(x^2-x+1)} dx$$

input `integrate((-x**3+1)**(2/3)/(x**3+1),x)`output `Integral((-x - 1)*(x**2 + x + 1)**(2/3)/((x + 1)*(x**2 - x + 1)), x)`

Maxima [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}}{x^3+1} dx$$

input `integrate((-x^3+1)^(2/3)/(x^3+1),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(2/3)/(x^3 + 1), x)`

Giac [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}}{x^3+1} dx$$

input `integrate((-x^3+1)^(2/3)/(x^3+1),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3)/(x^3 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(1-x^3)^{2/3}}{x^3+1} dx$$

input `int((1 - x^3)^(2/3)/(x^3 + 1),x)`

output `int((1 - x^3)^(2/3)/(x^3 + 1), x)`

Reduce [F]

$$\int \frac{(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}}{x^3+1} dx$$

input `int((-x^3+1)^(2/3)/(x^3+1),x)`

output `int((-x**3+1)**(2/3)/(x**3+1),x)`

3.114 $\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx$

Optimal result	844
Mathematica [C] (verified)	845
Rubi [A] (verified)	845
Maple [F]	851
Fricas [F]	851
Sympy [F]	852
Maxima [F]	852
Giac [F]	852
Mupad [F(-1)]	853
Reduce [F]	853

Optimal result

Integrand size = 20, antiderivative size = 250

$$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx = \frac{2^{2/3} \arctan\left(\frac{1 - \frac{2}{3}\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{1 + \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt[3]{2}\sqrt{3}} - \frac{1}{2}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) + \frac{\log((1-x)(1+x)^2)}{6\sqrt[3]{2}} + \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{3\sqrt[3]{2}} - \frac{1}{3}2^{2/3} \log\left(1 + \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right) - \frac{\log(-1+x+2^{2/3}\sqrt[3]{1-x^3})}{2\sqrt[3]{2}}$$

output

```
-1/2*x^2*hypergeom([1/3, 2/3], [5/3], x^3)+1/12*ln((1-x)*(1+x)^2)*2^(2/3)+1/6*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)-1/3*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)-1/4*ln(-1+x+2^(2/3)*(-x^3+1)^(1/3))*2^(2/3)+1/3*arctan(1/3*(1-2*2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)+1/6*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)*2^(2/3)
```

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 10.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.10

$$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx = \frac{1}{2}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, -\frac{2}{3}, 1, \frac{5}{3}, x^3, -x^3\right)$$

input

```
Integrate[(x*(1 - x^3)^(2/3))/(1 + x^3), x]
```

output

```
(x^2*AppellF1[2/3, -2/3, 1, 5/3, x^3, -x^3])/2
```

Rubi [A] (verified)

Time = 0.43 (sec) , antiderivative size = 272, normalized size of antiderivative = 1.09, number of steps used = 14, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.650$, Rules used = {984, 888, 991, 750, 16, 27, 1142, 25, 27, 1082, 217, 1103, 2574}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{x(1-x^3)^{2/3}}{x^3+1} dx$$

↓ 984

$$2 \int \frac{x}{\sqrt[3]{1-x^3}(x^3+1)} dx - \int \frac{x}{\sqrt[3]{1-x^3}} dx$$

$$\begin{aligned}
& \downarrow 888 \\
& 2 \int \frac{x}{\sqrt[3]{1-x^3}(x^3+1)} dx - \frac{1}{2} x^2 \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) \\
& \downarrow 991 \\
& 2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \int \frac{1}{\frac{2(1-x)^3}{1-x^3} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} \right) - \\
& \quad \frac{1}{2} x^2 \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) \\
& \downarrow 750 \\
& 2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \int \frac{\sqrt[3]{2} \left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{1}{3} \int \frac{1}{\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} \right) - \\
& \quad \frac{1}{2} x^2 \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) \\
& \downarrow 16 \\
& 2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \int \frac{\sqrt[3]{2} \left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \right) - \\
& \quad \frac{1}{2} x^2 \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) \\
& \downarrow 27 \\
& 2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \sqrt[3]{2} \int \frac{2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3\sqrt[3]{2}} \right) - \\
& \quad \frac{1}{2} x^2 \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right) \\
& \downarrow 1142
\end{aligned}$$

$$2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} - \frac{\int -\frac{\sqrt[3]{2} \left(1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2 \cdot 2^{2/3}} \right) \right)$$

$$\frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right)$$

↓ 25

$$2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} + \frac{\int \frac{\sqrt[3]{2} \left(1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2 \cdot 2^{2/3}} \right) \right)$$

$$\frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right)$$

↓ 27

$$2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} + \frac{\int \frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} \right) \right)$$

$$\frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right)$$

↓ 1082

$$2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \sqrt[3]{2} \left(\frac{3 \int \frac{1}{-\frac{(1-x)^2}{(1-x^3)^{2/3}} - 3} d\left(1 - \frac{2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{2^{2/3}} + \frac{\int \frac{1 - \frac{2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1} d\frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} \right) \right)$$

$$\frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right)$$

↓ 217

$$2 \left(-\frac{1}{3} \sqrt[3]{2} \left(\frac{\int \frac{1 - \frac{2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1} d\frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} - \frac{\sqrt{3} \arctan \left(\frac{1 - \frac{2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{2^{2/3}} \right) - \frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \dots \right)$$

$$\frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right)$$

↓ 1103

$$2 \left(-\frac{1}{3} \int \frac{1}{(x+1)\sqrt[3]{1-x^3}} dx - \frac{1}{3} \sqrt[3]{2} \left(-\frac{\sqrt{3} \arctan \left(\frac{1 - \frac{2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\sqrt{3}} \right)}{2^{2/3}} - \frac{\log \left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1 \right)}{2 \cdot 2^{2/3}} \right) - \dots \right)$$

$$\frac{1}{2} x^2 \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right)$$

↓ 2574

$$2 \left(-\frac{1}{3} \sqrt[3]{2} \left(\frac{\sqrt{3} \arctan \left(\frac{1 - \frac{2}{3} \sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} \right)}{2^{2/3}} - \frac{\log \left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1 \right)}{2 \cdot 2^{2/3}} \right) + \frac{1}{3} \frac{\sqrt{3} \arctan \left(\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} + 1 \right)}{2 \sqrt[3]{2}} \right) + \frac{1}{2} x^2 \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3 \right)$$

input `Int[(x*(1 - x^3)^(2/3))/(1 + x^3),x]`

output `-1/2*(x^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3]) + 2*(-1/3*(2^(1/3)*(-(Sqrt[3]*ArcTan[(1 - (2*2^(1/3))*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3])/2^(2/3)) - Log[1 + (2^(2/3)*(1 - x)^2)/(1 - x^3)^(2/3) - (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(2*2^(2/3)))) - Log[1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(3*2^(1/3)) + ((Sqrt[3]*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]])/(2*2^(1/3)) + Log[(1 - x)*(1 + x)^2]/(4*2^(1/3)) - (3*Log[-1 + x + 2^(2/3)]*(1 - x^3)^(1/3)]/(4*2^(1/3)))/3)`

Defintions of rubi rules used

rule 16 `Int[(c_)/((a_) + (b_)*(x_)), x_Symbol] := Simp[c*(Log[RemoveContent[a + b*x, x]]/b), x] /; FreeQ[{a, b, c}, x]`

rule 25 `Int[-(Fx_), x_Symbol] := Simp[Identity[-1] Int[Fx, x], x]`

rule 27 `Int[(a_)*(Fx_), x_Symbol] := Simp[a Int[Fx, x], x] /; FreeQ[a, x] && !MatchQ[Fx, (b_)*(Gx_)] /; FreeQ[b, x]`

rule 217 `Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

rule 750 $\text{Int}[(a_ + (b_ \cdot x)^3)^{-1}, x_Symbol] \rightarrow \text{Simp}[1/(3 \cdot \text{Rt}[a, 3]^2) \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3] \cdot x), x], x] + \text{Simp}[1/(3 \cdot \text{Rt}[a, 3]^2) \text{Int}[(2 \cdot \text{Rt}[a, 3] - \text{Rt}[b, 3] \cdot x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3] \cdot \text{Rt}[b, 3] \cdot x + \text{Rt}[b, 3]^2 \cdot x^2), x], x] /;$
 $\text{FreeQ}\{a, b\}, x]$

rule 888 $\text{Int}[(c_ \cdot x)^{m_} \cdot (a_ + (b_ \cdot x)^n)^{p_}, x_Symbol] \rightarrow \text{Simp}[a^p \cdot ((c \cdot x)^{m+1}/(c \cdot (m+1))) \cdot \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b) \cdot (x^n/a)], x] /;$
 $\text{FreeQ}\{a, b, c, m, n, p\}, x] \&\& !\text{IGtQ}[p, 0] \&\& (\text{ILtQ}[p, 0] \parallel \text{GtQ}[a, 0])$

rule 984 $\text{Int}[(x_ \cdot (a_ + (b_ \cdot x)^n)^{p_}) / (c_ + (d_ \cdot x)^n), x_Symbol] \rightarrow \text{Simp}[b/d \text{Int}[x \cdot (a + b \cdot x^n)^{p-1}, x], x] - \text{Simp}[(b \cdot c - a \cdot d)/d \text{Int}[x \cdot (a + b \cdot x^n)^{p-1}/(c + d \cdot x^n), x], x] /;$
 $\text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& \text{IntBinomialQ}[a, b, c, d, 1, 1, n, p, -1, x]$

rule 991 $\text{Int}[x_ / ((a_ + (b_ \cdot x)^3)^{1/3} \cdot (c_ + (d_ \cdot x)^3)), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b/a, 3]\}, \text{Simp}[-q^2/(3 \cdot d) \text{Int}[1/((1 - q \cdot x) \cdot (a + b \cdot x^3)^{1/3}), x], x] + \text{Simp}[q/d \text{Subst}[\text{Int}[1/(1 + 2 \cdot a \cdot x^3), x], x, (1 + q \cdot x)/(a + b \cdot x^3)^{1/3}], x] /;$
 $\text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b \cdot c - a \cdot d, 0] \&\& \text{EqQ}[b \cdot c + a \cdot d, 0]$

rule 1082 $\text{Int}[(a_ + (b_ \cdot x) + (c_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4 \cdot \text{Simplify}[a \cdot (c/b^2)]\}, \text{Simp}[-2/b \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2 \cdot c \cdot (x/b)], x] /;$
 $\text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \parallel !\text{RationalQ}[b^2 - 4 \cdot a \cdot c]) /;$
 $\text{FreeQ}\{a, b, c\}, x]$

rule 1103 $\text{Int}[(d_ + (e_ \cdot x)) / (a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[d \cdot (\text{Log}[\text{RemoveContent}[a + b \cdot x + c \cdot x^2, x]]/b), x] /;$
 $\text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2 \cdot c \cdot d - b \cdot e, 0]$

rule 1142 $\text{Int}[(d_ + (e_ \cdot x)) / (a_ + (b_ \cdot x) + (c_ \cdot x)^2), x_Symbol] \rightarrow \text{Simp}[(2 \cdot c \cdot d - b \cdot e)/(2 \cdot c) \text{Int}[1/(a + b \cdot x + c \cdot x^2), x], x] + \text{Simp}[e/(2 \cdot c) \text{Int}[(b + 2 \cdot c \cdot x)/(a + b \cdot x + c \cdot x^2), x], x] /;$
 $\text{FreeQ}\{a, b, c, d, e\}, x]$

rule 2574

```
Int[1/(((c_) + (d_)*(x_))*((a_) + (b_)*(x_)^3)^(1/3)), x_Symbol] := Simp[
  Sqrt[3]*(ArcTan[(1 - 2^(1/3)*Rt[b, 3]*((c - d*x)/(d*(a + b*x^3)^(1/3)))))/Sqrt[3]]/(2^(4/3)*Rt[b, 3]*c), x] + (Simp[Log[(c + d*x)^2*(c - d*x)]/(2^(7/3)*Rt[b, 3]*c), x] - Simp[(3*Log[Rt[b, 3]*(c - d*x) + 2^(2/3)*d*(a + b*x^3)^(1/3)])/(2^(7/3)*Rt[b, 3]*c), x]) /; FreeQ[{a, b, c, d}, x] && EqQ[b*c^3 + a*d^3, 0]
```

Maple [F]

$$\int \frac{x(-x^3 + 1)^{\frac{2}{3}}}{x^3 + 1} dx$$

input `int(x*(-x^3+1)^(2/3)/(x^3+1),x)`

output `int(x*(-x^3+1)^(2/3)/(x^3+1),x)`

Fricas [F]

$$\int \frac{x(1 - x^3)^{2/3}}{1 + x^3} dx = \int \frac{(-x^3 + 1)^{\frac{2}{3}}x}{x^3 + 1} dx$$

input `integrate(x*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="fricas")`

output `integral((-x^3 + 1)^(2/3)*x/(x^3 + 1), x)`

Sympy [F]

$$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{x(-(x-1)(x^2+x+1))^{2/3}}{(x+1)(x^2-x+1)} dx$$

input `integrate(x*(-x**3+1)**(2/3)/(x**3+1),x)`

output `Integral(x*(-(x - 1)*(x**2 + x + 1))**(2/3)/((x + 1)*(x**2 - x + 1)), x)`

Maxima [F]

$$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}x}{x^3+1} dx$$

input `integrate(x*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="maxima")`

output `integrate((-x^3 + 1)^(2/3)*x/(x^3 + 1), x)`

Giac [F]

$$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}x}{x^3+1} dx$$

input `integrate(x*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(2/3)*x/(x^3 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{x(1-x^3)^{2/3}}{x^3+1} dx$$

input `int((x*(1 - x^3)^(2/3))/(x^3 + 1),x)`output `int((x*(1 - x^3)^(2/3))/(x^3 + 1), x)`**Reduce [F]**

$$\int \frac{x(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{\frac{2}{3}}x}{x^3+1} dx$$

input `int(x*(-x^3+1)^(2/3)/(x^3+1),x)`output `int(((- x**3 + 1)**(2/3)*x)/(x**3 + 1),x)`

3.115 $\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx$

Optimal result	854
Mathematica [C] (warning: unable to verify)	855
Rubi [A] (verified)	855
Maple [F]	857
Fricas [F]	858
Sympy [F]	858
Maxima [F]	858
Giac [F]	859
Mupad [F(-1)]	859
Reduce [F]	859

Optimal result

Integrand size = 24, antiderivative size = 383

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = -\frac{2^{2/3} \arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2(1-x)}}{\frac{\sqrt[3]{1-x^3}}{\sqrt{3}}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{1+\frac{\sqrt[3]{2(1-x)}}{\sqrt{3}}}{\frac{\sqrt[3]{1-x^3}}{\sqrt{3}}}\right)}{\sqrt[3]{2}\sqrt{3}} + \frac{\arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{2^{2/3} \arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2x}}{\frac{\sqrt[3]{1-x^3}}{\sqrt{3}}}\right)}{\sqrt{3}} + \frac{1}{2}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) - \frac{\log((1-x)(1+x)^2)}{6\sqrt[3]{2}} - \frac{\log(1+x^3)}{3\sqrt[3]{2}} - \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}}{\sqrt[3]{1-x^3}}\right)}{3\sqrt[3]{2}}$$

output

```
1/2*x^2*hypergeom([1/3, 2/3], [5/3], x^3)-1/12*ln((1-x)*(1+x)^2)*2^(2/3)-1/6
*ln(x^3+1)*2^(2/3)-1/6*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3)*(1-x)/
(-x^3+1)^(1/3))*2^(2/3)+1/3*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(2/3)+1/2*
ln(-2^(1/3)*x-(-x^3+1)^(1/3))*2^(2/3)-1/2*ln(x+(-x^3+1)^(1/3))+1/4*ln(-1+x
+2^(2/3)*(-x^3+1)^(1/3))*2^(2/3)-1/3*arctan(1/3*(1-2*2^(1/3)*(1-x)/(-x^3+1
)^(1/3))*3^(1/2))*2^(2/3)*3^(1/2)-1/6*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1
)^(1/3))*3^(1/2))*3^(1/2)*2^(2/3)+1/3*arctan(1/3*(1-2*x/(-x^3+1)^(1/3))*3^(
1/2))*3^(1/2)-1/3*arctan(1/3*(1-2*2^(1/3)*x/(-x^3+1)^(1/3))*3^(1/2))*2^(2/
3)*3^(1/2)
```

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 15.09 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.36

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = -\frac{1}{2}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, -\frac{2}{3}, 1, \frac{5}{3}, x^3, -x^3\right) - \frac{4x(1-x^3)^{2/3} \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{2}{3}, 1, \frac{4}{3}, x^3, -x^3\right)}{(1+x^3)\left(-4 \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{2}{3}, 1, \frac{4}{3}, x^3, -x^3\right) + x^3\left(3 \operatorname{AppellF1}\left(\frac{4}{3}, -\frac{2}{3}, 2, \frac{7}{3}, x^3, -x^3\right) + 2 \operatorname{AppellF1}\left(\frac{4}{3}, \frac{1}{3}, 1, \frac{7}{3}, x^3, -x^3\right)\right)\right)}$$

input

```
Integrate[((1-x)*(1-x^3)^(2/3))/(1+x^3),x]
```

output

```
-1/2*(x^2*AppellF1[2/3, -2/3, 1, 5/3, x^3, -x^3]) - (4*x*(1-x^3)^(2/3)*A
ppellF1[1/3, -2/3, 1, 4/3, x^3, -x^3])/((1+x^3)*(-4*AppellF1[1/3, -2/3,
1, 4/3, x^3, -x^3] + x^3*(3*AppellF1[4/3, -2/3, 2, 7/3, x^3, -x^3] + 2*App
ellF1[4/3, 1/3, 1, 7/3, x^3, -x^3])))
```

Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 655, normalized size of antiderivative = 1.71, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {7276, 2009}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\begin{aligned}
& \int \frac{(1-x)(1-x^3)^{2/3}}{x^3+1} dx \\
& \quad \downarrow 7276 \\
& \int \left(-\frac{2(1-x^3)^{2/3}}{3(-x-1)} + \frac{(-1-(-1)^{2/3})(1-x^3)^{2/3}}{3(\sqrt[3]{-1}x-1)} + \frac{(\sqrt[3]{-1}-1)(1-x^3)^{2/3}}{3(-(-1)^{2/3}x-1)} \right) dx \\
& \quad \downarrow 2009 \\
& -\frac{2^{2/3} \arctan\left(\frac{\sqrt[3]{2}(1-x)+1}{\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} + \frac{(1+(-1)^{2/3}) \arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{3\sqrt{3}} + \\
& \frac{(1-\sqrt[3]{-1}) \arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{2 \arctan\left(\frac{1-\frac{2x}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{3\sqrt{3}} - \\
& \frac{(1-\sqrt[3]{-1}) \arctan\left(\frac{\sqrt[3]{2}(x+\sqrt[3]{-1})}{\sqrt[3]{1-x^3}}\right)}{3\sqrt{3}} + \frac{(1+(-1)^{2/3}) \arctan\left(\frac{(-1)^{2/3}\sqrt[3]{2}(\sqrt[3]{-1}x+1)}{\sqrt[3]{1-x^3}}+1\right)}{3\sqrt{3}} + \\
& \frac{\sqrt[3]{2}\sqrt{3}}{6}(-1)^{2/3}(1-\sqrt[3]{-1})x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) + \\
& \frac{1}{6}(1-\sqrt[3]{-1})x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) + \frac{1}{3}x^2 \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, x^3\right) - \\
& \frac{1}{6}(1+(-1)^{2/3}) \log\left(\sqrt[3]{1-x^3}+x\right) - \frac{1}{6}(1-\sqrt[3]{-1}) \log\left(\sqrt[3]{1-x^3}+x\right) - \\
& \frac{1}{3} \log\left(\sqrt[3]{1-x^3}+x\right) + \frac{(1-\sqrt[3]{-1}) \log\left(-(-2)^{2/3}\sqrt[3]{1-x^3}-(-1)^{2/3}x+1\right)}{2\sqrt[3]{2}} + \\
& \frac{\log\left(-2^{2/3}\sqrt[3]{1-x^3}-x+1\right)}{\sqrt[3]{2}} + \frac{(1+(-1)^{2/3}) \log\left(\sqrt[3]{-12}^{2/3}\sqrt[3]{1-x^3}+\sqrt[3]{-1}x+1\right)}{2\sqrt[3]{2}} - \\
& \frac{\log\left(-((1-x)(x+1)^2)\right)}{3\sqrt[3]{2}} - \frac{(1+(-1)^{2/3}) \log\left(-(-1)^{2/3}(x+(-1)^{2/3})^2(\sqrt[3]{-1}x+1)\right)}{6\sqrt[3]{2}} - \\
& \frac{(1-\sqrt[3]{-1}) \log\left((-1)^{2/3}(x+\sqrt[3]{-1})((-1)^{2/3}x+1)^2\right)}{6\sqrt[3]{2}}
\end{aligned}$$

input `Int[((1-x)*(1-x^3)^(2/3))/(1+x^3),x]`

output

```

-((2^(2/3)*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]])/Sqrt[3
]) + (2*ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]])/(3*Sqrt[3]) + ((1 - (-
-1)^(1/3))*ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]])/(3*Sqrt[3]) + ((1
+ (-1)^(2/3))*ArcTan[(1 - (2*x)/(1 - x^3)^(1/3))/Sqrt[3]])/(3*Sqrt[3]) - (
(1 - (-1)^(1/3))*ArcTan[(1 - (2^(1/3)*((-1)^(1/3) + x))/(1 - x^3)^(1/3))/S
qrt[3]])/(2^(1/3)*Sqrt[3]) - ((1 + (-1)^(2/3))*ArcTan[(1 + ((-1)^(2/3)*2^(
1/3)*(1 + (-1)^(1/3)*x))/(1 - x^3)^(1/3))/Sqrt[3]])/(2^(1/3)*Sqrt[3]) + (x
^2*Hypergeometric2F1[1/3, 2/3, 5/3, x^3])/3 + ((1 - (-1)^(1/3))*x^2*Hyperg
eometric2F1[1/3, 2/3, 5/3, x^3])/6 + ((-1)^(2/3)*(1 - (-1)^(1/3))*x^2*Hype
rgeometric2F1[1/3, 2/3, 5/3, x^3])/6 - Log[-((1 - x)*(1 + x)^2)]/(3*2^(1/3
)) - ((1 + (-1)^(2/3))*Log[-((-1)^(2/3)*((-1)^(2/3) + x)^2*(1 + (-1)^(1/3
)*x)])]/(6*2^(1/3)) - ((1 - (-1)^(1/3))*Log[(-1)^(2/3)*((-1)^(1/3) + x)*(1
+ (-1)^(2/3)*x)^2]/(6*2^(1/3)) - Log[x + (1 - x^3)^(1/3)]/3 - ((1 - (-1)^(
1/3))*Log[x + (1 - x^3)^(1/3)])/6 - ((1 + (-1)^(2/3))*Log[x + (1 - x^3)^(
1/3)])/6 + ((1 - (-1)^(1/3))*Log[1 - (-1)^(2/3)*x - (-2)^(2/3)*(1 - x^3)^(
1/3)])/2*2^(1/3) + Log[1 - x - 2^(2/3)*(1 - x^3)^(1/3)]/2^(1/3) + ((1 +
(-1)^(2/3))*Log[1 + (-1)^(1/3)*x + (-1)^(1/3)*2^(2/3)*(1 - x^3)^(1/3)])/(2
*2^(1/3))

```

Defintions of rubi rules used

rule 2009

```
Int[u_, x_Symbol] := Simp[IntSum[u, x], x] /; SumQ[u]
```

rule 7276

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Maple [F]

$$\int \frac{(1-x)(-x^3+1)^{\frac{2}{3}}}{x^3+1} dx$$

input

```
int((1-x)*(-x^3+1)^(2/3)/(x^3+1),x)
```

output

```
int((1-x)*(-x^3+1)^(2/3)/(x^3+1),x)
```

Fricas [F]

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = \int -\frac{(-x^3+1)^{2/3}(x-1)}{x^3+1} dx$$

input `integrate((1-x)*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="fricas")`

output `integral(-(-x^3 + 1)^(2/3)*(x - 1)/(x^3 + 1), x)`

Sympy [F]

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = -\int \left(-\frac{(1-x^3)^{2/3}}{x^3+1} \right) dx - \int \frac{x(1-x^3)^{2/3}}{x^3+1} dx$$

input `integrate((1-x)*(-x**3+1)**(2/3)/(x**3+1),x)`

output `-Integral(-(1 - x**3)**(2/3)/(x**3 + 1), x) - Integral(x*(1 - x**3)**(2/3)/(x**3 + 1), x)`

Maxima [F]

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = \int -\frac{(-x^3+1)^{2/3}(x-1)}{x^3+1} dx$$

input `integrate((1-x)*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="maxima")`

output `-integrate((-x^3 + 1)^(2/3)*(x - 1)/(x^3 + 1), x)`

Giac [F]

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = \int -\frac{(-x^3+1)^{2/3}(x-1)}{x^3+1} dx$$

input `integrate((1-x)*(-x^3+1)^(2/3)/(x^3+1),x, algorithm="giac")`

output `integrate(-(-x^3 + 1)^(2/3)*(x - 1)/(x^3 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = -\int \frac{(1-x^3)^{2/3}(x-1)}{x^3+1} dx$$

input `int(-((1 - x^3)^(2/3)*(x - 1))/(x^3 + 1),x)`

output `-int(((1 - x^3)^(2/3)*(x - 1))/(x^3 + 1), x)`

Reduce [F]

$$\int \frac{(1-x)(1-x^3)^{2/3}}{1+x^3} dx = \int \frac{(-x^3+1)^{2/3}}{x^3+1} dx - \left(\int \frac{(-x^3+1)^{2/3}x}{x^3+1} dx \right)$$

input `int((1-x)*(-x^3+1)^(2/3)/(x^3+1),x)`

output `int((-x**3 + 1)**(2/3)/(x**3 + 1),x) - int(((- x**3 + 1)**(2/3)*x)/(x**3 + 1),x)`

3.116 $\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx$

Optimal result	860
Mathematica [A] (verified)	861
Rubi [A] (verified)	861
Maple [C] (warning: unable to verify)	866
Fricas [A] (verification not implemented)	867
Sympy [F]	868
Maxima [F]	868
Giac [F]	869
Mupad [F(-1)]	869
Reduce [F]	869

Optimal result

Integrand size = 19, antiderivative size = 272

$$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx = \frac{\sqrt[3]{2} \arctan\left(\frac{1-\frac{2}{3}\sqrt[3]{2(1-x)}}{\frac{\sqrt[3]{1-x^3}}{\sqrt{3}}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{1+\frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}}{\sqrt{3}}\right)}{2^{2/3}\sqrt{3}}$$

$$+ \frac{\log\left(2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}\right)}{3 \cdot 2^{2/3}} - \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right)}{3 \cdot 2^{2/3}}$$

$$+ \frac{1}{3} \sqrt[3]{2} \log\left(1 + \frac{\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}}\right) - \frac{\log\left(2\sqrt[3]{2} + \frac{(1-x)^2}{(1-x^3)^{2/3}} + \frac{2^{2/3}(1-x)}{\sqrt[3]{1-x^3}}\right)}{6 \cdot 2^{2/3}}$$

output

```
1/6*ln(2^(2/3)+(-1+x)/(-x^3+1)^(1/3))*2^(1/3)-1/6*ln(1+2^(2/3)*(1-x)^2/(-x^3+1)^(2/3)-2^(1/3)*(1-x)/(-x^3+1)^(1/3))*2^(1/3)+1/3*2^(1/3)*ln(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))-1/12*ln(2*2^(1/3)+(1-x)^2/(-x^3+1)^(2/3)+2^(2/3)*(1-x)/(-x^3+1)^(1/3))*2^(1/3)+1/3*2^(1/3)*arctan(1/3*(1-2*2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*3^(1/2)+1/6*arctan(1/3*(1+2^(1/3)*(1-x)/(-x^3+1)^(1/3))*3^(1/2))*2^(1/3)*3^(1/2)
```

Mathematica [A] (verified)

Time = 1.88 (sec) , antiderivative size = 283, normalized size of antiderivative = 1.04

$$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx = \frac{2\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1-x^3}}{\sqrt[3]{2}-\sqrt[3]{2x+\sqrt[3]{1-x^3}}}\right) + 4\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{1-x^3}}{-2\sqrt[3]{2}+2\sqrt[3]{2x+\sqrt[3]{1-x^3}}}\right) - 4 \log\left(-\sqrt[3]{2} + \sqrt[3]{2x} - \dots\right)}{\dots}$$

input `Integrate[(1 - x^3)^(1/3)/(1 + x^3), x]`

output `-1/6*(2*Sqrt[3]*ArcTan[(Sqrt[3]*(1 - x^3)^(1/3))/(2^(1/3) - 2^(1/3)*x + (1 - x^3)^(1/3))] + 4*Sqrt[3]*ArcTan[(Sqrt[3]*(1 - x^3)^(1/3))/(-2*2^(1/3) + 2*2^(1/3)*x + (1 - x^3)^(1/3))] - 4*Log[-2^(1/3) + 2^(1/3)*x - (1 - x^3)^(1/3)] - 2*Log[-2^(1/3) + 2^(1/3)*x + 2*(1 - x^3)^(1/3)] + 2*Log[2^(2/3) - 2*2^(2/3)*x + 2^(2/3)*x^2 + (-1 + x)*(2 - 2*x^3)^(1/3) + (1 - x^3)^(2/3)] + Log[2^(2/3) - 2*2^(2/3)*x + 2^(2/3)*x^2 - 2*(-1 + x)*(2 - 2*x^3)^(1/3) + 4*(1 - x^3)^(2/3)]/2^(2/3)`

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 295, normalized size of antiderivative = 1.08, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.526$, Rules used = {927, 982, 821, 16, 1142, 25, 27, 1082, 217, 1103}

Below are the steps used by Rubi to obtain the solution. The rule number used for the transformation is given above next to the arrow. The rules definitions used are listed below.

$$\int \frac{\sqrt[3]{1-x^3}}{x^3+1} dx$$

↓ 927

$$-9 \int \frac{1-x}{\sqrt[3]{1-x^3} \left(4 - \frac{(1-x)^3}{1-x^3}\right) \left(\frac{2(1-x)^3}{1-x^3} + 1\right)} d \frac{1-x}{\sqrt[3]{1-x^3}}$$

↓ 982

$$-9 \left(\frac{1}{9} \int \frac{1-x}{\sqrt[3]{1-x^3} \left(4 - \frac{(1-x)^3}{1-x^3} \right)} d \frac{1-x}{\sqrt[3]{1-x^3}} + \frac{2}{9} \int \frac{1-x}{\sqrt[3]{1-x^3} \left(\frac{2(1-x)^3}{1-x^3} + 1 \right)} d \frac{1-x}{\sqrt[3]{1-x^3}} \right)$$

821

$$-9 \left(\frac{2}{9} \left(\frac{\int \frac{\frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\int \frac{1}{\frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}} d \frac{1-x}{\sqrt[3]{1-x^3}}}{3\sqrt[3]{2}} \right) + \frac{1}{9} \left(\frac{\int \frac{1}{2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}} d \frac{1-x}{\sqrt[3]{1-x^3}}}{3 \cdot 2^{2/3}} \right) \right)$$

16

$$-9 \left(\frac{2}{9} \left(\frac{\int \frac{\frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\log \left(\frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}} \right)}{3 \cdot 2^{2/3}} \right) + \frac{1}{9} \left(\frac{\int \frac{1}{\frac{2^{2/3} - \frac{1-x}{\sqrt[3]{1-x^3}}}{(1-x^3)^{2/3} + \frac{2^{2/3}(1-x)}{\sqrt[3]{1-x^3}} + 2\sqrt[3]{2}}} d \frac{1-x}{\sqrt[3]{1-x^3}}}{3 \cdot 2^{2/3}} \right) \right)$$

1142

$$-9 \left(\frac{2}{9} \left(\frac{\frac{3}{2} \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}} d \frac{1-x}{\sqrt[3]{1-x^3}} + \frac{\int - \frac{\sqrt[3]{2} \left(1 - \frac{2\sqrt[3]{2(1-x)}}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}}} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} - \frac{\log \left(\frac{\sqrt[3]{2(1-x)}+1}{\sqrt[3]{1-x^3}} \right)}{3 \cdot 2^{2/3}} \right) \right)$$

25

$$-9 \left(\frac{2}{9} \left(\frac{\frac{3}{2} \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\int \frac{\sqrt[3]{2} \left(1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} \right)}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}}}{2\sqrt[3]{2}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3 \cdot 2^{2/3}} \right) \right) +$$

↓ 27

$$-9 \left(\frac{2}{9} \left(\frac{\frac{3}{2} \int \frac{1}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{1}{2} \int \frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3 \cdot 2^{2/3}} \right) \right) +$$

↓ 1082

$$-9 \left(\frac{2}{9} \left(\frac{3 \int \frac{1}{-\frac{(1-x)^2}{(1-x^3)^{2/3}} - 3} d \left(1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} \right)}{\sqrt[3]{2}} - \frac{1}{2} \int \frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3 \cdot 2^{2/3}} \right) \right) + \frac{1}{9}$$

↓ 217

$$-9 \left(\frac{2}{9} \left(\frac{-\frac{1}{2} \int \frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1} d \frac{1-x}{\sqrt[3]{1-x^3}} - \frac{\sqrt[3]{3} \arctan \left(\frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt[3]{3}} \right)}{\sqrt[3]{2}}}{3\sqrt[3]{2}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3 \cdot 2^{2/3}} \right) + \frac{1}{9} \left(\dots \right) \right)$$

1103

$$-9 \left(\frac{2}{9} \left(\frac{\frac{\log \left(\frac{2^{2/3}(1-x)^2}{(1-x^3)^{2/3}} - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{2\sqrt[3]{2}} - \frac{\sqrt[3]{3} \arctan \left(\frac{1 - \frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}}}{\sqrt[3]{3}} \right)}{\sqrt[3]{2}}}{3\sqrt[3]{2}} - \frac{\log \left(\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{1-x^3}} + 1 \right)}{3 \cdot 2^{2/3}} \right) + \frac{1}{9} \left(\frac{\sqrt[3]{3} \arctan \left(\dots \right)}{\dots} \right) \right)$$

input `Int[(1 - x^3)^(1/3)/(1 + x^3),x]`

output `-9*((2*((-((Sqrt[3]*ArcTan[(1 - (2*2^(1/3))*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]])/2^(1/3)) + Log[1 + (2^(2/3)*(1 - x)^2/(1 - x^3)^(2/3) - (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(2*2^(1/3)))/(3*2^(1/3)) - Log[1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3)]/(3*2^(2/3)))/9 + (-1/3*Log[2^(2/3) - (1 - x)/(1 - x^3)^(1/3)]/2^(2/3) - (Sqrt[3]*ArcTan[(1 + (2^(1/3)*(1 - x))/(1 - x^3)^(1/3))/Sqrt[3]] - Log[2*2^(1/3) + (1 - x)^2/(1 - x^3)^(2/3) + (2^(2/3)*(1 - x))/(1 - x^3)^(1/3)]/2)/(3*2^(2/3)))/9)`

Defintions of rubi rules used

- rule 16 $\text{Int}[(c_)/((a_)+(b_)*(x_)), x_Symbol] \rightarrow \text{Simp}[c*(\text{Log}[\text{RemoveContent}[a + b*x, x]]/b), x] /; \text{FreeQ}[\{a, b, c\}, x]$
- rule 25 $\text{Int}[-(Fx_), x_Symbol] \rightarrow \text{Simp}[\text{Identity}[-1] \text{ Int}[Fx, x], x]$
- rule 27 $\text{Int}[(a_)*(Fx_), x_Symbol] \rightarrow \text{Simp}[a \text{ Int}[Fx, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[Fx, (b_)*(Gx_)] /; \text{FreeQ}[b, x]$
- rule 217 $\text{Int}[(a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$
- rule 821 $\text{Int}[(x_)/((a_)+(b_)*(x_)^3), x_Symbol] \rightarrow \text{Simp}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{(-1)} \text{ Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Simp}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]) \text{ Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$
- rule 927 $\text{Int}[(a_)+(b_)*(x_)^3)^{1/3}/((c_)+(d_)*(x_)^3), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 3]\}, \text{Simp}[9*(a/(c*q)) \text{ Subst}[\text{Int}[x/((4 - a*x^3)*(1 + 2*a*x^3)), x], x, (1 + q*x)/(a + b*x^3)^{1/3}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[b*c + a*d, 0]$
- rule 982 $\text{Int}[(e_)*(x_)^m/(((a_)+(b_)*(x_)^n)*((c_)+(d_)*(x_)^n))), x_Symbol] \rightarrow \text{Simp}[b/(b*c - a*d) \text{ Int}[(e*x)^m/(a + b*x^n), x], x] - \text{Simp}[d/(b*c - a*d) \text{ Int}[(e*x)^m/(c + d*x^n), x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0]$
- rule 1082 $\text{Int}[(a_)+(b_)*(x_)+(c_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Simp}[-2/b \text{ Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ !\text{RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x]$

rule 1103

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

rule 1142

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> S
imp[(2*c*d - b*e)/(2*c) Int[1/(a + b*x + c*x^2), x], x] + Simp[e/(2*c)
Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x]
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 6.40 (sec) , antiderivative size = 1151, normalized size of antiderivative = 4.23

method	result	size
trager	Expression too large to display	1151

input

```
int((-x^3+1)^(1/3)/(x^3+1),x,method=_RETURNVERBOSE)
```

output

```

1/6*RootOf(_Z^3-2)*ln((6*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2)^4*x^3-18*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)^2*RootOf(_Z^3-2)^3*x^3+x^6*RootOf(_Z^3-2)^2-3*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2)*x^6+18*RootOf(_Z^3-2)^2*(-x^3+1)^(2/3)*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*x^2-6*RootOf(_Z^3-2)*(-x^3+1)^(1/3)*x^4-2*RootOf(_Z^3-2)^2*x^3+6*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2)*x^3+12*(-x^3+1)^(2/3)*x^2+6*RootOf(_Z^3-2)*(-x^3+1)^(1/3)*x+RootOf(_Z^3-2)^2-3*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2))/(1+x)^2/(x^2-x+1)^2)-1/6*ln((6*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2)^4*x^3-18*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)^2*RootOf(_Z^3-2)^3*x^3-x^6*RootOf(_Z^3-2)^2+3*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2)*x^6-18*RootOf(_Z^3-2)^2*(-x^3+1)^(2/3)*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*x^2+6*RootOf(_Z^3-2)*(-x^3+1)^(1/3)*x^4+18*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*x^2+6*RootOf(_Z^3-2)^2*x^3-18*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2)*x^3-6*RootOf(_Z^3-2)*(-x^3+1)^(1/3)*x-18*(-x^3+1)^(1/3)*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*x-RootOf(_Z^3-2)^2+3*RootOf(RootOf(_Z^3-2)^2+3*_Z*RootOf(_Z^3-2)+9*_Z^2)*RootOf(_Z^3-2))/(1+x)^2/(x^2-x+1)^2)*RootOf(_Z^3-2)-1/2*ln((6*RootOf(RootOf(_Z^3-2)^2+3*_Z...

```

Fricas [A] (verification not implemented)

Time = 1.61 (sec) , antiderivative size = 341, normalized size of antiderivative = 1.25

$$\begin{aligned}
& \int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx \\
&= \frac{1}{18} \sqrt{32}^{\frac{1}{3}} \arctan \left(-\frac{6\sqrt{32}^{\frac{2}{3}}(x^{16} - 33x^{13} + 110x^{10} - 110x^7 + 33x^4 - x)(-x^3 + 1)^{\frac{1}{3}} - 24\sqrt{32}^{\frac{1}{3}}(x^{14} - 2x^{11} + x^8)}{3(x^{18} - 102x^{15} + 447x^{12} - 102x^9 + x^6)} \right) \\
&+ \frac{1}{18} \\
&\cdot 2^{\frac{1}{3}} \log \left(-\frac{12(-x^3 + 1)^{\frac{2}{3}}x^2 + 2^{\frac{2}{3}}(x^6 + 2x^3 + 1) - 6 \cdot 2^{\frac{1}{3}}(x^4 - x)(-x^3 + 1)^{\frac{1}{3}}}{x^6 + 2x^3 + 1} \right) - \frac{1}{36} \\
&\cdot 2^{\frac{1}{3}} \log \left(\frac{12 \cdot 2^{\frac{2}{3}}(x^8 - 4x^5 + x^2)(-x^3 + 1)^{\frac{2}{3}} + 2^{\frac{1}{3}}(x^{12} - 32x^9 + 78x^6 - 32x^3 + 1) + 6(x^{10} - 11x^7 + 11x^4 - x)}{x^{12} + 4x^9 + 6x^6 + 4x^3 + 1} \right)
\end{aligned}$$

input

```
integrate((-x^3+1)^(1/3)/(x^3+1),x, algorithm="fricas")
```

output

```
1/18*sqrt(3)*2^(1/3)*arctan(-1/3*(6*sqrt(3)*2^(2/3)*(x^16 - 33*x^13 + 110*
x^10 - 110*x^7 + 33*x^4 - x)*(-x^3 + 1)^(1/3) - 24*sqrt(3)*2^(1/3)*(x^14 -
2*x^11 - 6*x^8 - 2*x^5 + x^2)*(-x^3 + 1)^(2/3) - sqrt(3)*(x^18 + 42*x^15
- 417*x^12 + 812*x^9 - 417*x^6 + 42*x^3 + 1))/(x^18 - 102*x^15 + 447*x^12
- 628*x^9 + 447*x^6 - 102*x^3 + 1)) + 1/18*2^(1/3)*log(-(12*(-x^3 + 1)^(2/
3)*x^2 + 2^(2/3)*(x^6 + 2*x^3 + 1) - 6*2^(1/3)*(x^4 - x)*(-x^3 + 1)^(1/3))
/(x^6 + 2*x^3 + 1)) - 1/36*2^(1/3)*log((12*2^(2/3)*(x^8 - 4*x^5 + x^2)*(-x
^3 + 1)^(2/3) + 2^(1/3)*(x^12 - 32*x^9 + 78*x^6 - 32*x^3 + 1) + 6*(x^10 -
11*x^7 + 11*x^4 - x)*(-x^3 + 1)^(1/3))/(x^12 + 4*x^9 + 6*x^6 + 4*x^3 + 1))
```

Sympy [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx = \int \frac{\sqrt[3]{-(x-1)(x^2+x+1)}}{(x+1)(x^2-x+1)} dx$$

input

```
integrate((-x**3+1)**(1/3)/(x**3+1),x)
```

output

```
Integral((-x - 1)*(x**2 + x + 1)**(1/3)/((x + 1)*(x**2 - x + 1)), x)
```

Maxima [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x^3+1} dx$$

input

```
integrate((-x^3+1)^(1/3)/(x^3+1),x, algorithm="maxima")
```

output

```
integrate((-x^3 + 1)^(1/3)/(x^3 + 1), x)
```

Giac [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x^3+1} dx$$

input `integrate((-x^3+1)^(1/3)/(x^3+1),x, algorithm="giac")`

output `integrate((-x^3 + 1)^(1/3)/(x^3 + 1), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx = \int \frac{(1-x^3)^{1/3}}{x^3+1} dx$$

input `int((1 - x^3)^(1/3)/(x^3 + 1),x)`

output `int((1 - x^3)^(1/3)/(x^3 + 1), x)`

Reduce [F]

$$\int \frac{\sqrt[3]{1-x^3}}{1+x^3} dx = \int \frac{(-x^3+1)^{\frac{1}{3}}}{x^3+1} dx$$

input `int((-x^3+1)^(1/3)/(x^3+1),x)`

output `int((- x**3 + 1)**(1/3)/(x**3 + 1),x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	870
4.2 Links to plain text integration problems used in this report for each CAS .	888

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal."}
        ]
      ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
    ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A",""}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal."}
      ]
    ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order of result is higher than in optimal."}
    ]
  ]

```



```

        finalresult={"F","Contains unresolved integral."}
    ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType,expn]],
  If[Head[expn]==Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]]==Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]],2]],
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
    If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3,ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
    If[HypergeometricFunctionQ[Head[expn]],

```

```

    Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 5]],
    If [AppellFunctionQ [Head [expn]],
      Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 6]],
      If [Head [expn] === RootSum,
        Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 7]],
        If [Head [expn] === Integrate || Head [expn] === Int,
          Apply [Max, Append [Map [ExpnType, Apply [List, expn]], 8]],
          9]]]]]]]]]]]]

```

```

ElementaryFunctionQ [func_] :=
  MemberQ [{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

```

```

SpecialFunctionQ [func_] :=
  MemberQ [{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

```

```

HypergeometricFunctionQ [func_] :=
  MemberQ [{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ [func_] :=
  MemberQ [{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
#                   if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
#                   see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issue";
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal

```

```

#   antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (" ,
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf

            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex
        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well
        fi;

```

```

        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A"," ";
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of
                            convert(leaf_count_result,string)," $ vs. $2(",
                            convert(leaf_count_optimal,string),")=",convert(2*leaf_co

        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function

```

```

# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+'') or type(expn,'*') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9

```

```

    end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.

```

```
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc;
```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
    asinh,acosh,atanh,acoth,asech,acsch
  ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
    gamma,loggamma,digamma,zeta,polylog,LambertW,
    elliptic_f,elliptic_e,elliptic_pi,exp_polar
  ]
```



```

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')

```

```

    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""

```

```

else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is lar
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

```

```

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arcsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:

```

```

    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi','zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral',
        'weierstrassPInverse','weierstrass','weierstrassP','weierstrassZeta',
        'weierstrassPPrime','weierstrassSigma']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

def is_atom(expn):

    #debug=False

```

```

if debug:
    print ("Enter is_atom, expn=",expn)

if not hasattr(expn, 'parent'):
    return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print ("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isins
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

if debug:
    print ("Enter grade_antiderivative for sagemath")
    print ("Enter grade_antiderivative, result=",result)
    print ("Enter grade_antiderivative, optimal=",optimal)
    print ("type(anti)=", type(result))
    print ("type(optimal)=", type(optimal))

```

```

leaf_count_result = tree_size(result) #leaf_count(result)
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result - 2*leaf_count_optimal)
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result - expnType_optimal)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```


4.2 Links to plain text integration problems used in this report for each CAS

1. Mathematica integration problems as .m file
2. Maple integration problems as .txt file
3. Sagemath integration problems as .sage file
4. Reduce integration problems as .txt file
5. Mupad integration problems as .txt file
6. Sympy integration problems as .py file